

*Modular Homotopy Theory for Algebraic Factorization Algebras on Algebraic Curves*

Volume 3

# CALABI–YAU QUANTUM GROUPS

*Chiral Algebras from Calabi–Yau Categories  
via  $E_1/E_2$  Factorization*

Raez Lorgat

Draft — April 13, 2026



## ABSTRACT

Volumes I and II take a chiral algebra as input and construct its ordered bar complex  $B^{\text{ord}}(\mathcal{A}) = T^c(s^{-1}\bar{\mathcal{A}})$ . This volume constructs the geometric input: a functor  $\Phi: \text{CY}_d\text{-Cat} \rightarrow E_n\text{-ChiralAlg}$  from Calabi–Yau categories of dimension  $d$  to  $E_n$ -chiral algebras ( $n = 2$  for  $d \leq 2$ ;  $n = 1$  for  $d \geq 3$ ).

A  $\text{CY}_d$  category  $\mathcal{C}$  carries a cyclic  $A_\infty$ -structure whose  $\mathbf{S}^d$ -framing determines an operadic enhancement. Three steps: the cyclic  $A_\infty$ -structure produces a Lie conformal algebra  $L_{\mathcal{C}}$ ; the factorization envelope yields a chiral algebra  $A_{\mathcal{C}}$  on a curve  $X$ ; the  $\mathbf{S}^d$ -framing lifts factorization to an  $E_2$ -chiral algebra with braided monoidal representation category. For  $d = 2$ , all three steps are proved (Theorem CY-A). For  $d = 3$ , holomorphic Chern–Simons breaks  $E_2$  to  $E_1$ ; braiding is recovered via the Drinfeld center. Steps (1) and (2) are unconditional; step (3) is conditional on the chain-level  $\mathbf{S}^3$ -framing.

For  $\mathbb{C}^3$ , the Jordan quiver gives  $\mathbb{C}^3 \rightarrow \mathcal{W}_{1+\infty} \rightarrow \text{Rep}^{E_2}(Y(\widehat{\mathfrak{gl}}_1))$ : the critical cohomological Hall algebra  $\text{CoHA}(Q)$  is the  $E_1$ -sector, and the Drinfeld center recovers the full braided structure. For a general toric  $\text{CY}_3$  with fan  $\Sigma$ , quiver charts indexed by Bridgeland stability chambers give local chiral algebras; wall-crossing is conjecturally Maurer–Cartan gauge equivalence with the global chiral algebra as homotopy colimit. When  $\Phi$  exists, the Borcherds–Kac–Moody denominator identity equals the bar Euler product.

The prototype  $K3 \times E$  reveals that a single  $\text{CY}_3$  admits multiple chiral algebraizations with distinct modular characteristics. The  $\kappa_\bullet$ -spectrum  $\text{Spec}_{\kappa_\bullet}(K3 \times E) = \{2, 3, 5, 24\}$  records four subscripted invariants:  $\kappa_{\text{cat}} = 2 = \chi(\mathcal{O}_{K3})$  (holomorphic Euler characteristic),  $\kappa_{\text{ch}} = 3$  (chiral shadow from the Vol I shadow tower),  $\kappa_{\text{BKM}} = 5$  (Borcherds weight of the Igusa cusp form  $\Delta_5$ , so  $\text{wt}(\Phi_{10}) = 2\kappa_{\text{BKM}}$ ),  $\kappa_{\text{fiber}} = 24$  (lattice rank). Bare  $\kappa$  is forbidden; the subscript records the geometric provenance. For  $d \geq 3$  the CY chiral algebra is  $E_1$ ; the effective framing obstruction  $\text{Obs}_{\text{eff}}(d) \in \pi_d(BU)$  or  $\pi_d(B\text{Sp})$  is 8-periodic by Bott periodicity and trivial when  $d \bmod 8 \in \{1, 3, 7\}$ .



# Preface

A chiral algebra is a solution. What is the problem it solves?

Volumes I and II develop the bar-cobar machine that extracts modular invariants from any chiral algebra  $A$  on a curve  $X$ : the bar complex  $B(A) = T^c(s^{-1}\bar{A})$  with its deconcatenation coproduct, the universal Maurer–Cartan element  $\Theta_A$ , the modular characteristic  $\kappa_{\text{ch}}(A)$ , and the five theorems that control the genus tower. Volume I introduces  $E_1$ -chiral quantum groups as abstract algebraic-geometric objects: ordered bar complexes with spectral  $R$ -matrices, Drinfeld coproducts, and Yang–Baxter equations, all living on configuration spaces of curves. Neither volume constructs a single chiral algebra or a single chiral quantum group. The input is always assumed. This volume constructs the input: both the chiral algebras and, from Calabi–Yau geometry, the concrete  $E_1$ -chiral quantum groups that are the flesh on Volume I’s algebraic skeleton.

A terminological point: “Hochschild” in this volume means categorical Hochschild cohomology: the cyclic bar complex  $\text{CC}_*(C)$  of a dg or  $A_\infty$ -category, topological in nature. The chiral upgrade  $\text{ChirHoch}^*(A)$  of Volume I (Theorem H) is the holomorphic version; it is the output of the functor  $\Phi$ , not the input. The two are related by the functor itself:  $\Phi$  sends  $\text{CC}_*(C)$  to  $\text{ChirHoch}^*(A_C)$ , lifting topological Hochschild data to holomorphic chiral data on the curve.

The construction is a functor

$$\Phi: \text{CY}_d\text{-Cat} \longrightarrow E_2\text{-ChirAlg}.$$

A Calabi–Yau category  $C$  of dimension  $d$  carries a cyclic  $A_\infty$ -structure: a non-degenerate trace  $\text{Tr}: \text{HC}_d^-(C) \rightarrow k$  on negative cyclic homology, encoding the CY condition as a  $d$ -dimensional Frobenius structure at the chain level. The functor  $\Phi$  converts this datum into a chiral algebra  $A_C$  on  $X$  in three steps: the cyclic  $A_\infty$ -structure determines a Lie conformal algebra  $L_C$ ; the factorization envelope  $U_X^{\text{fact}}(L_C)$  produces a chiral algebra  $A_C$ ; the  $\mathbb{S}^d$ -framing enhances the factorization structure. For  $d = 2$ , the  $\mathbb{S}^2$ -action on Hochschild homology gives an  $E_2$ -chiral algebra with braided monoidal representation category, and all three steps are proved (Theorem CY-A<sub>2</sub>). For  $d = 3$ , holomorphic Chern–Simons breaks  $E_2$  to  $E_1$ ; the braiding is recovered via the Drinfeld center of the  $E_1$ -monoidal representation category. Steps (1) and (2) are unconditional; step (3) is a programme conditional on the chain-level  $\mathbb{S}^3$ -framing.

**What the CY perspective reveals.** Volumes I and II see one chiral algebra at a time. The CY functor sees a category  $C$  that admits *multiple* chiral algebraizations, each with its own modular characteristic. A single  $\text{CY}_3$  manifold  $X$  produces a spectrum of  $\kappa_\bullet$ -values, not a single number. For  $K3 \times E$ :

$$\text{Spec}_{\kappa_\bullet}(K3 \times E) = \{\kappa_{\text{cat}}, \kappa_{\text{ch}}, \kappa_{\text{BKM}}, \kappa_{\text{fiber}}\} = \{2, 3, 5, 24\}.$$

Here  $\kappa_{\text{cat}} = 2 = \chi(\mathcal{O}_{K3})$  is the holomorphic Euler characteristic;  $\kappa_{\text{ch}} = 3 = \dim_{\mathbb{C}} X$  is the modular characteristic of the chiral de Rham complex;  $\kappa_{\text{BKM}} = 5$  is the weight of the Igusa cusp form  $\Delta_5$  (so

$2\kappa_{\text{BKM}} = 10 = \text{wt}(\Phi_{10})$ , the Siegel modular weight); and  $\kappa_{\text{fiber}} = 24$  is the lattice rank. These four numbers measure four faces of the same manifold: the holomorphic Euler characteristic, the boundary chiral algebra, the bulk Borchers–Kac–Moody superalgebra, and the lattice frame. The chiral-holographic pair  $(\kappa_{\text{ch}}, \kappa_{\text{BKM}}) = (3, 5)$  is the one that sees the boundary–bulk axis of Volumes I–II. The  $\kappa_{\bullet}$ -spectrum is invisible from any single chiral algebra; it is intrinsic to the CY category.

The  $E_n$ -hierarchy provides the organizing spiral. A  $\text{CY}_d$  category carries a native  $\mathbb{S}^d$ -framing, hence an  $E_d$ -chiral algebra via  $\Phi$ . At  $d = 3$ , the chiral algebra is natively  $E_3$ ; but  $\pi_1(C_2(\mathbb{R}^3)) = 0$ , so the  $E_3$ -braiding is symmetric; too symmetric for quantum groups. The resolution is a two-step detour: stabilize from  $E_3$  to  $E_1$  (forget the symmetric braiding, retain the associative product), then recover a *non-symmetric* braiding via the Drinfeld center  $Z(\text{Rep}^{E_1}(A_C))$ , which is  $E_2$ -monoidal. The resulting  $R$ -matrix is the quantum group  $R$ -matrix. The key arithmetic:  $\pi_1(C_2(\mathbb{R}^2)) = \mathbb{Z}$  (braid group, native  $E_2$ , non-symmetric), while  $\pi_1(C_2(\mathbb{R}^d)) = \mathbb{Z}/2$  for  $d \geq 3$  (symmetric group). Dimension  $d = 2$  is the sweet spot where the native  $E_2$ -structure already carries a non-symmetric braiding. For  $d \geq 3$ , the descent-and-center detour is mandatory: one must discard the over-symmetric  $E_d$ -structure, pass to  $E_1$ , and rebuild the braiding categorically.

The spiral, not circle, because the closing step is conjectural at  $d \geq 3$ . The trajectory  $E_d \rightarrow E_1 \rightarrow E_2$  (via the Drinfeld center) returns to the same operadic level as  $d = 2$ , but the  $E_2$  structure so obtained is *derived*: it lives on the representation category, not on the algebra itself. Whether this derived  $E_2$ -structure lifts to a genuine  $E_2$ -chiral algebra (closing the spiral to a circle) is the content of Conjecture CY-B. At  $d = 2$  the spiral closes by construction: the native  $E_2$ -framing and the Drinfeld center  $E_2$  agree. At  $d = 3$  the two  $E_2$ -structures inhabit different categories, and identifying them requires the chain-level  $\mathbb{S}^3$ -framing that is the open core of the  $d = 3$  programme. The spiral is the honest shape of the  $E_n$ -story: each CY dimension  $d$  enters at  $E_d$ , descends to  $E_1$ , and ascends to  $E_2$  through the center; the question is whether the ascent reaches the starting altitude.

Three further structures emerge only through  $\Phi$ . First, the bar complex converts enumerative geometry into homotopy algebra: Donaldson–Thomas invariants become root multiplicities of the Borchers–Kac–Moody superalgebra, and when the relevant chiral object exists, the denominator identity is expected to appear as the bar Euler product. Second, the  $E_1$ -stabilization theorem: for  $d \geq 3$ , the expected CY chiral output is natively  $E_1$  (the Schouten–Nijenhuis bracket vanishes, all noncommutative structure arises from quantization, and the CoHA extension correspondence imposes a preferred ordering). The braided monoidal structure lives not in the algebra itself but in the Drinfeld center of its  $E_1$ -representation category; the Drinfeld center is the categorified averaging map  $\text{av}: \mathfrak{g}^{E_1} \rightarrow \mathfrak{g}^{\text{mod}}$  of Volume I. Third, the framing obstruction tower: the effective obstruction  $\text{Obs}_{\text{eff}}(d) \in \pi_d(BU)$  is 8-periodic by Bott periodicity and trivial precisely when  $d \bmod 8 \in \{1, 3, 7\}$  (at  $d \equiv 5 \bmod 8$ , the Sp-refinement contributes a  $\mathbb{Z}/2$ -obstruction), giving a rigid topological constraint on which CY dimensions admit additional chiral structure beyond  $E_1$ .

**The prototype:**  $K3 \times E$ . The product of a  $K3$  surface  $S$  and an elliptic curve  $E$  is the canonical  $\text{CY}_3$  testing ground: simple enough to compute (lattice  $\Lambda^{3,2}$  of signature  $(3, 2)$ ,  $K3$  elliptic genus  $\phi_{0,1}(\tau, z)$ , Borchers denominator  $\Delta_5$ ), and already exhibits the obstructions. Four structural barriers separate the shadow obstruction tower from the full automorphic form:

- (O1) *Categorical*: the shadow tower produces numbers (Taylor coefficients of  $\Theta_{A_C}$ ); the denominator identity is a function (an automorphic form on  $\mathcal{H}_2$ ).
- (O2)  *$\kappa_{\text{ch}}\text{--}\kappa_{\text{BKM}}$  mismatch*:  $\kappa_{\text{ch}} = 3 \neq 5 = \kappa_{\text{BKM}}$ ; the single-copy chiral algebra sees  $\dim_{\mathbb{C}} X$ , while the bulk reconstruction sees the Borchers weight.
- (O3) *Second quantization*: the physical DT partition function is the DMVV symmetric product of single-copy data; the bar complex of a single algebra is the single copy.

(O4) *Schottky*: at  $g \geq 4$ , the Torelli map  $\mathcal{M}_g \rightarrow \mathcal{A}_g$  is no longer dominant, and the shadow tower is imprisoned in tautological classes on  $\overline{\mathcal{M}}_g$ .

Each obstruction points to a research programme. The  $\kappa_{\text{ch}}\text{--}\kappa_{\text{BKM}}$  mismatch (O2) is the most telling: a single CY<sub>3</sub> admits multiple chiral algebraizations, and the  $\kappa_\bullet$ -spectrum records which face of the geometry each algebraization captures.

The functor  $\Phi$  applied to a K<sub>3</sub> surface distinguishes three algebras: the lattice vertex algebra  $V_\Lambda$  attached to  $H^*(K3, \mathbb{Z})$ , the generalized Kac–Moody algebra whose denominator is  $\Delta_5$ , and the Conway module arising from the Leech lattice frame. These three algebraizations share  $\kappa_{\text{cat}} = 2$  but differ in  $\kappa_{\text{ch}}$  and  $\kappa_{\text{BKM}}$ ; the distinction is intrinsic to the CY functor and invisible at the level of the Euler characteristic alone.

Thirteen structural results (K<sub>3-1</sub> through K<sub>3-13</sub>) are proved for  $K3 \times E$ . Together they constitute the most complete worked example in the programme; we record all thirteen here, as the pattern they form is itself a structural datum.

K<sub>3-1</sub>:  $\kappa_{\text{ch}}(K3 \times E) = 3 = \dim_{\mathbb{C}}$ , verified by six independent paths (direct Hodge, Mukai pairing, Dolbeault contracting homotopy, bar computation, Borchers lift, and elliptic genus). Additivity:  $\kappa_{\text{ch}}(K3) = 2$ ,  $\kappa_{\text{ch}}(E) = 1$ .

K<sub>3-2</sub>: the factor 2 in  $Z_{K3} = 2\phi_{0,1}$  is  $\kappa_{\text{cat}}(K3) = \chi(\mathcal{O}_{K3}) = 2$ . The massive Mathieu multiplicities satisfy  $A_n = \kappa_{\text{cat}} \cdot \dim(\rho_n)$ : the bar complex provides the universal coefficient and does not detect  $M_{24}$  directly.

K<sub>3-3</sub>:  $\kappa_{\text{BKM}} = \frac{1}{2} \text{wt}(\Phi_{10}) = 5$ . The ratio  $\kappa_{\text{BKM}}/\kappa_{\text{ch}} = 5/3$  reflects second quantization; the Hilbert–Chow exceptional divisor accounts for  $\chi(\text{Hilb}^2) - \chi(\text{Sym}^2) = 24 = \chi(K3)$ .

K<sub>3-4</sub>: the moonshine multiplier  $A_n = \kappa_{\text{cat}} \cdot \dim(\rho_n)$ , factoring BPS degeneracies into a homological part ( $\kappa_{\text{cat}} = 2$ ) and a group-theoretic part ( $\rho_n$ , the Mathieu representation).

K<sub>3-5</sub>: the mock modular decomposition  $Z_{K3} = (h - 24\mu)\vartheta_1^2/\eta^3$  is verified to machine precision ( $\sim 5.7 \times 10^{-16}$  relative error), where  $\mu$  is the Zwegers Appell–Lerch sum.

K<sub>3-6</sub>: the shadow generating function  $Z^{\text{sh}}(t) = \kappa_{\text{ch}}((t/2)/\sin(t/2) - 1)$  converges with radius  $R = 2\pi$ , has entire Borel transform, and exhibits no resurgence. This contrasts with the Gevrey-1 divergent topological string.

K<sub>3-7</sub>: the shadow–Siegel gap theorem with four structural obstructions (O<sub>1</sub>–O<sub>4</sub>), proving that the shadow tower does NOT produce  $\Phi_{10}$ .

K<sub>3-8</sub>: boundary versus sigma:  $\kappa_{\text{fiber}} = 24$  (lattice rank) vs  $\kappa_{\text{ch}} = 2$  (K<sub>3</sub> sigma model); ratio 12 at genus 1, conditional at  $g \geq 2$  (the K<sub>3</sub> sigma model is multi-weight).

K<sub>3-9</sub>: the denominator identity  $\prod_{\alpha>0} (1 - e^{-\alpha})^{\text{mult}(\alpha)}$  equals the bar Euler product, conditional on CY-A for  $d = 3$ .

K<sub>3-10</sub>: the Borchers multiplicative lift  $\phi_{0,1} \rightsquigarrow \Delta_5$  realises genus escalation: genus-1 elliptic genus data on  $\mathbb{H}_1$  determines the genus-2 Igusa cusp form on  $\mathbb{H}_2$ , with additive and multiplicative lifts encoding root system and Fourier–Jacobi structure respectively.

K<sub>3-11</sub>: wall-crossing in the Bridgeland stability manifold of  $D^b(\text{Coh}(K3))$  corresponds to reflections in the BKM root system of  $\mathfrak{g}_{\Delta_5}$ ; the Kontsevich–Soibelman wall-crossing formula is the MC gauge transformation in the convolution algebra.

K<sub>3-12</sub>: the K<sub>3</sub> double current algebra  $\mathfrak{g}_{K3}$ , the finite-dimensional analogue of the DDCA in which  $\mathbb{C}[u, v]$  is replaced by  $H^*(S, \mathbb{C})$  with the Mukai pairing.

K<sub>3-13</sub>: the Faber–Pandharipande value  $\lambda_5^{\text{FP}} = 73/3503554560$  at genus 5; the unreduced numerator  $2^{2g-1} - 1 = 511$  requires GCD reduction with  $(2g)!$ .

**The holomorphic Chern–Simons hierarchy.** The theories studied in this volume form a hierarchy in real dimension, each producing chiral algebras by restriction to codimension-2 defects. At  $d = 6$ : holomorphic

Chern–Simons on a Calabi–Yau 3-fold  $Y$  (Witten, Costello), the primary source. A curve  $C \subset Y$  of codimension 2 carries a defect algebra: the normal bundle  $N_{C/Y}$  of rank 2 supplies two spectral parameters, and the defect algebra is a chiral algebra on  $C$  with quantum group structure from the normal directions (Proposition 9.21). For  $Y = \mathbb{C}^3$  and  $C = \mathbb{C}$ , the defect algebra is  $\mathcal{W}_{1+\infty}$  at  $\Psi = -\sigma_2(h_1, h_2, h_3)$  (the second elementary symmetric polynomial in the  $\Omega$ -background parameters). At  $d = 5$ : partially topological Chern–Simons on  $\mathbb{R} \times Y$  for a  $CY_2$  surface  $Y$ ; restriction to  $\mathbb{R} \times C$  gives the ordered bar with one spectral parameter. At  $d = 4$ : holomorphic Chern–Simons on  $\mathbb{C}^2$ ; the Costello–Witten–Yamazaki theory that produces Yangians from holomorphic gauge fields. At  $d = 3$ : the theories of Volume II. The hierarchy  $6 \rightarrow 5 \rightarrow 4 \rightarrow 3$  is codimension-2 restriction at each step; each restriction produces a chiral algebra from a gauge theory, and the  $E_n$  level decreases as the dimension drops. The hierarchy is not limited to Chern–Simons: 6d little string theory on  $K3$  and  $M5$  brane theory produce chiral algebras by the same defect mechanism, with the specific gauge-theory content replaced by stringy data.

**Concrete CY quantum groups.** Volume I introduces  $E_1$ -chiral quantum groups as abstract algebraic-geometric objects: an ordered bar complex  $B^{\text{ord}}(A)$  with spectral  $R$ -matrix, Drinfeld coproduct, and Yang–Baxter equation, all living on configuration spaces of curves. This volume constructs the concrete examples. The critical cohomological Hall algebra  $\text{CoHA}(Q)$  of a quiver  $Q$  (with potential  $W$  from the CY condition) is an  $E_1$ -chiral quantum group: its product is the ordered Hall multiplication (graded by dimension vectors), its  $R$ -matrix comes from the Ext-quiver, and its coproduct is the CoHA vertex coproduct of Schiffmann–Vasserot. For the Jordan quiver ( $Q = \bullet \circlearrowleft$ ):  $\text{CoHA}(Q) \cong Y^+(\widehat{\mathfrak{gl}}_1) \cong \mathcal{W}_{1+\infty}$ . The Drinfeld center  $Z(\text{Rep}^{E_1}(Y^+(\widehat{\mathfrak{gl}}_1))) \cong \text{Rep}^{E_2}(Y(\widehat{\mathfrak{gl}}_1))$  recovers the full braided structure (Theorem 8.28). For a general toric  $CY_3$  with fan  $\Sigma$ : quiver charts indexed by Bridgeland stability chambers give local chiral algebras; wall-crossing is conjecturally MC gauge equivalence. The quantum toroidal algebra  $U_{q,t}(\widehat{\mathfrak{gl}}_1)$  with CY condition  $q_1 q_2 q_3 = 1$  is the two-parameter deformation carrying the full  $\mathbb{Z} \times \mathbb{Z}$  bigrading from the double current algebra. In the additive parametrization  $q_i = e^{h_i}$ , the CY condition reads  $h_1 + h_2 + h_3 = 0$ : two independent parameters survive, and the identification  $q_1 = q, q_2 = t^{-1}, q_3 = t/q$  recovers the standard  $(q, t)$ -parametrization. The trigonometric structure function

$$G(x; q_1, q_2, q_3) = \frac{(1 - q_1 x)(1 - q_2 x)(1 - q_3 x)}{(1 - q_1^{-1} x)(1 - q_2^{-1} x)(1 - q_3^{-1} x)}$$

satisfies the CY reflection identity  $G(x) \cdot G(1/x) = 1$  precisely when  $q_1 q_2 q_3 = 1$ : the identity is equivalent to the CY condition, and its failure measures the departure from Calabi–Yau. The Miki automorphism (cyclic permutation of  $(q_1, q_2, q_3)$ , generating the  $S_3$ -Weyl group of the CY torus) is algebra-specific: it requires the concrete structure of  $U_{q,t}(\widehat{\mathfrak{gl}}_1)$  and does not follow from the  $E_3$ -operad alone.

**Toric  $CY_3$  and critical CoHAs.** For a toric  $CY_3$  determined by a fan  $\Sigma$ , the toric diagram fixes a quiver  $Q_X$ . The critical cohomological Hall algebra  $\text{CoHA}(Q_X)$  is the  $E_1$ -sector: the positive half of an affine super Yangian  $Y(\widehat{\mathfrak{g}}_{Q_X})$ . The full braided structure is recovered through the Drinfeld center. For  $\mathbb{C}^3$ : the Jordan quiver gives  $Y(\widehat{\mathfrak{gl}}_1) \simeq \mathcal{W}_{1+\infty}$ , and the chain  $\mathbb{C}^3 \rightarrow \mathcal{W}_{1+\infty} \rightarrow \text{Rep}^{E_2}(Y(\widehat{\mathfrak{gl}}_1))$  is verified end-to-end. The toric fan is the Dynkin data.

The CoHA is the natural  $E_1$ -primitive: it is the ordered bar complex of the  $CY_3$  chiral algebra, with deconcatenation coproduct encoding the topological direction and the  $R$ -matrix encoding the spectral braiding. The passage from CoHA to quantum group is the CY incarnation of the  $E_1$ -to- $E_2$  lift: averaging over the symmetric group recovers the factorization coalgebra of Volume I. The algebraic underpinning of this passage (the integrable structure of the ordered chiral bar complex, the  $R$ -matrix, and the Yangian



presentation) is developed in the companion paper *On the Integrable Theory of Chiral Yangians and Related Chiral Quantum Groups* (Volume I, standalone).

**The DDCA–toroidal bridge.** The deformed double current algebra  $\text{DDCA}_k(\mathfrak{gl}_K) = U(\mathfrak{gl}_K \otimes \mathbb{C}[u, v])$  carries two spectral parameters  $(u, v)$  from the two holomorphic directions of  $\mathbb{C}^2$  in the 5d noncommutative Chern–Simons theory of Costello. The quantum toroidal algebra  $U_{q,t}(\widehat{\mathfrak{gl}}_K)$  carries two deformation parameters  $(q, t)$  from the two loop directions of  $\mathbb{C}^* \times \mathbb{C}^*$ . Both degenerate to the affine Yangian  $Y(\widehat{\mathfrak{gl}}_K)$  in complementary limits: the DDCA by specializing one spectral variable to zero (collapsing  $\mathbb{C}[u, v] \rightarrow \mathbb{C}[u]$ ); the toroidal algebra by sending  $t \rightarrow 1$  (collapsing the outer loop). The conjecture (Conjecture 14.197) states the bridge: writing  $q = e^{\hbar_1}, t = e^{\hbar_2}$ , the DDCA is the rational  $(\hbar_1, \hbar_2 \rightarrow 0)$  degeneration of the quantum toroidal algebra, in the category of filtered algebras with the bidegree filtration on generators  $J_{m,n}^a$ . The two one-parameter degenerations of  $U_{q,t}$  produce a commuting square

$$\begin{array}{ccc} U_{q,t}(\widehat{\mathfrak{gl}}_K) & \xrightarrow{t \rightarrow 1} & Y(\widehat{\mathfrak{gl}}_K) \otimes U(\mathbb{C}[v]) \\ q \rightarrow 1 \downarrow & & \downarrow \hbar_1 \rightarrow 0 \\ U(\mathbb{C}[u]) \otimes Y(\widehat{\mathfrak{gl}}_K) & \xrightarrow{\hbar_2 \rightarrow 0} & \text{DDCA}_k(\mathfrak{gl}_K) \end{array}$$

whose diagonal  $\hbar_1 = \hbar_2 \rightarrow 0$  recovers the affine Yangian as the common degeneration. The Koszul dual side has a parallel identification: the double Yangian  $Y^{(2)}(\mathfrak{gl}_K)$  is the rational degeneration of  $U_{q^{-1},t^{-1}}(\widehat{\mathfrak{gl}}_K)$ , consistent with the parameter inversion  $(q, t) \mapsto (q^{-1}, t^{-1})$  of the Koszul dual toroidal algebra. The modular characteristic is continuous across the bridge:  $\kappa_{\text{ch}}(\text{DDCA}_k(\mathfrak{gl}_K)) = K(k + K)/2 = \lim_{\hbar_i \rightarrow 0} \kappa_{\text{ch}}(U_{q,t})$ .

**The shadow tower as Gromov–Witten theory.** At  $\kappa_{\text{ch}} = \Psi$  (the deformation parameter of  $\mathcal{W}_{1+\infty}$  at  $c = 1$ ), the shadow obstruction tower of Volume I produces the perturbative constant-map Gromov–Witten free energies  $F_g^{\text{GW}, \text{const}}(\mathbb{C}^3)$ : the shadow IS the full GW answer for  $\mathbb{C}^3$  (which has no compact curves, hence no worldsheet instantons). On the Donaldson–Thomas side, the MacMahon function  $M(q) = \prod_{n \geq 1} (1 - q^n)^{-n}$  encodes the DT partition function via the MNOP correspondence. The shadow coefficients  $S_r$  of the Virasoro algebra at the GW specialisation are the  $A_\infty$  coproduct correction coefficients  $\delta^{(r)}$ : the shadow tower encodes the coproduct corrections of the chiral quantum group.

The universal coproduct  $\Delta_z(e_s) = \sum C(N_R - b, k) z^k e_a^L \cdot e_b^R$  (all spins, closed form) extends the Drinfeld coproduct from spin 2 to arbitrary spin, with the universal Miura coefficient  $(\Psi - 1)/\Psi$  on the leading cross-term  $J \otimes W_{s-1}$  at every spin  $s \geq 2$  (Proposition 5.38).

**The  $E_1$ -chiral bialgebra.** Volume I defines  $E_1$ -chiral quantum groups abstractly. This volume constructs them concretely from CY geometry. The  $E_1$ -chiral Hopf axiom system (formalized in §5.7) requires: (H1) an  $E_1$ -chiral algebra  $A$  on a curve  $X$ ; (H2) a  $z$ -parametrized  $E_1$ -chiral coalgebra  $\Delta_z$  on the OPEN/ $E_1$  colour; (H3) bialgebra compatibility; (H4) spectral coassociativity; (H5) the Hopf axiom at  $z = 0$ . The associated  $R$ -matrix and Yang–Baxter equation are recovered from this ordered Hopf data. The ordered bar  $B^{\text{ord}}(A)$  preserves the  $R$ -matrix; the symmetric bar  $B^\Sigma(A)$  of Volume I kills the Hopf structure via averaging  $\text{av}(r(z)) = \kappa_{\text{ch}}$ .

The tetrahedron equation (the 3d analogue of YBE) fails for the Yang  $R$ -matrix at  $O(\kappa_{\text{ch}}^2)$ : the  $E_3$  structure is genuinely nontrivial beyond  $E_2$ , confirming that the  $E_n$  operadic circle is not a formal consequence of lower-level data.

**The Drinfeld centre as categorified averaging.** Volume I constructs the averaging map  $\text{av}: \mathfrak{g}_{\mathcal{A}}^{E_1} \rightarrow \mathfrak{g}_{\mathcal{A}}^{\text{mod}}$  that projects the ordered  $R$ -matrix to the scalar  $\kappa_{\text{ch}}$ . This volume constructs its categorical lift: the Drinfeld centre  $Z: E_1\text{-Cat} \rightarrow E_2\text{-Cat}$ , which takes the  $E_1$ -monoidal representation category  $\text{Rep}^{E_1}(A_C)$  to an  $E_2$ -braided monoidal category  $Z(\text{Rep}^{E_1}(A_C))$ . Direct restriction from  $E_3$  to  $E_2$  gives only symmetric braiding ( $\pi_1(C_2(\mathbb{R}^3)) = 0$ ); the Drinfeld centre is the sole source of non-trivial braiding. For  $\mathbb{C}^3$ :  $Z(\text{Rep}^{E_1}(Y^+(\widehat{\mathfrak{gl}}_1))) \cong \text{Rep}^{E_2}(Y(\widehat{\mathfrak{gl}}_1)) \cong \text{Rep}^{E_2}(\mathcal{W}_{1+\infty})$  (Theorem 8.28). Five invariants match at six levels: naive centre (dimension 1) versus derived centre (dimension 3, capturing  $\text{Ext}^{1,2}$  invisible to the commutant). The Drinfeld centre closes the  $E_n$  circle by recovering  $E_2$ -braiding from  $E_1$ -ordering.

**Wall-crossing as Maurer–Cartan gauge equivalence.** For a toric  $\text{CY}_3$  with fan  $\Sigma$ , Bridgeland stability conditions carve the space of central charges into chambers. Each chamber  $\sigma$  determines a quiver presentation  $Q_\sigma$  and a critical CoHA  $\text{CoHA}(Q_\sigma, W_\sigma)$ . Wall-crossing (varying the stability condition across a wall) changes the quiver presentation but not the underlying CY category. The conjecture: wall-crossing between chambers  $\sigma$  and  $\sigma'$  is Maurer–Cartan gauge equivalence  $\Theta_\sigma \sim \Theta_{\sigma'}$  in the modular convolution algebra of the CY chiral algebra. The global chiral algebra  $A_C$  would then be the homotopy colimit over the chamber decomposition:  $A_C = \text{hocolim}_\sigma A_{Q_\sigma}$ . For  $\mathbb{C}^3$  (single chamber): the conjecture is vacuous. For the resolved conifold  $\mathcal{O}(-1) \oplus \mathcal{O}(-1) \rightarrow \mathbb{P}^1$  (two chambers): the wall-crossing formula is the flop relation on DT partition functions (Bryan–Steinberg), and the MC gauge equivalence is the bar-cobar transport across the flop. This conjecture connects the algebraic programme of Volume I (Maurer–Cartan elements) to the enumerative programme (DT/BPS wall-crossing) of this volume.

**From shadow tower to BPS degeneracies.** The shadow obstruction tower of Volume I produces scalar invariants  $S_r(\mathcal{A})$  at each degree  $r$ . For  $\text{CY}_3$  chiral algebras, these invariants admit an enumerative interpretation: the DT invariants  $\Omega(\gamma)$  of the  $\text{CY}_3$  manifold are root multiplicities of the BKM superalgebra  $\mathfrak{g}_X$ , and the denominator identity

$$\prod_{\alpha > 0} (1 - e^{-\alpha})^{\text{mult}(\alpha)} = \sum_{w \in W} (-1)^{\ell(w)} e^{w(\rho) - \rho}$$

is expected to match the bar Euler product when the relevant chiral object exists: the product over positive roots is the candidate graded Euler characteristic of the bar complex  $B(A_X)$ . For  $K3 \times E$ : the denominator is the Igusa cusp form  $\Phi_{10} = \Delta_5^2$ , and the root multiplicities are the coefficients of the elliptic genus  $\phi_{0,1}(\tau, z)$ .

The identification operates at three levels: (i) *genus-1*: the shadow Eisenstein theorem  $D_2(\mathcal{A}, s) = -24\kappa_{\text{ch}} \cdot \zeta(s)\zeta(s-1)$  (Volume I) gives the genus-1 DT partition function via the Borcherds lift; (ii) *virtual*: the shadow coefficient  $S_r$  at degree  $r$  matches the virtual Euler characteristic of the moduli of  $r$ -dimensional sheaves; (iii) *motivic*: the full chiral quantum group structure ( $R$ -matrix, coproduct, RTT) should match the motivic DT invariants of Kontsevich–Soibelman.

This three-level correspondence is the connection traced here between the algebraic-geometric programme of Volumes I–II and the enumerative geometry of this volume.

**Convergence and divergence.** The shadow obstruction tower and the topological string free energy produce genus- $g$  amplitudes with opposite analytic behaviour. The shadow tower has Bernoulli decay  $|F_g^{\text{sh}}| \sim (2\pi)^{-2g}/g$ : convergent, with radius  $2\pi$ , Borel entire. The topological string has factorial growth  $|F_g^{\text{top}}| \sim (2g)!$ : divergent, Gevrey-1, requiring non-perturbative completion. The shadow is the algebraic soul: the part computable from OPE structure constants alone, without worldsheet instantons. For class  $\mathbf{G}$

algebras (Heisenberg, lattice VOAs), the shadow is the full answer. For class **M** (Virasoro,  $\mathcal{W}_{1+\infty}$ ), the shadow is a strict subtower, and the instanton sum is infinite. The instanton gap  $F_g^{\text{top}} - F_g^{\text{sh}}$  at each genus is the CY frontier.

The **G/L/C/M** classification of Volume I (shadow depth  $r = 2, 3, 4, \infty$ ) acquires a CY incarnation: the shadow depth of  $\Phi(C)$  is determined by the homological complexity of  $C$ , and the four classes reappear as conditions on the Hochschild-to-cyclic spectral sequence of the CY category. Class **G** corresponds to CY categories with semisimple Hochschild cohomology (e.g.  $\text{Coh}(E)$  for an elliptic curve); class **M** to those with infinite algebraic depth (e.g.  $D^b(K3)$ , whose chiral algebra  $\Phi(K3)$  carries the full Virasoro-type shadow tower).

**The elliptic curve as universal bridge.** The same elliptic curve  $E$  appears in six distinct roles across the three volumes:

- (i) *Base curve* (Vol I): the factorization algebra lives on  $\text{Ran}(E)$ ; the genus-1 propagator is  $d \log E(z, w)$ .
- (ii) *Sewing parameter* (Vol I, MC<sub>5</sub> analytic HS-sewing lane): the modular parameter  $\tau$  is the Hilbert–Schmidt sewing kernel.
- (iii) *CY fiber*: the DT partition function of  $K3 \times E$  decomposes as a sewing of  $K3$  fiber contributions along  $E$ .
- (iv) *Jacobi form parameter*: the Fourier coefficients of  $\phi_{0,1}(\tau, z)$  are root multiplicities and BPS degeneracies.
- (v) *Siegel modular period*: the Igusa cusp form  $\Delta_5$  lives on  $\text{Sp}_4(\mathbb{Z}) \backslash \mathbb{H}_2$ , which parametrizes principally polarized abelian surfaces; the product locus of pairs of elliptic curves is codimension 1.
- (vi) *Elliptic Hall algebra carrier*: the CY<sub>2</sub> quantum vertex chiral group is carried on  $\text{Coh}(E)$ , with Felder’s dynamical elliptic  $R$ -matrix.

Each appearance brings the same structural package: a modular parameter, an  $\text{SL}_2(\mathbb{Z})$  symmetry, a  $q$ -expansion, and a sewing functor converting genus-0 data into genus-1 data. The elliptic curve is the modular clock of the theory.

**The  $\Omega$ -background and  $E_1$  stabilization.** The  $E_1$  stabilization theorem (Volume I): for  $\text{CY}_d$  with  $d \geq 3$ , the Schouten–Nijenhuis bracket on the Lie conformal algebra  $L_C$  vanishes, so the classical structure is abelian and all noncommutative structure arises from quantization. The physical mechanism is the  $\Omega$ -background: the equivariant parameters  $\epsilon_1, \epsilon_2$  of the torus action on  $\mathbb{C}^2 \subset Y$  break Dunn additivity  $E_2 \simeq E_1 \otimes E_1$  by freezing one factor, producing a single  $E_1$  direction. At  $\epsilon_1 = 0, \epsilon_2 \neq 0$  (the Nekrasov–Shatashvili limit): one  $E_1$  factor survives and the resulting algebra is the Yangian  $Y(\mathfrak{g})$ , the quantum group associated to the gauge theory. At generic  $(\epsilon_1, \epsilon_2)$ : the quantum toroidal algebra  $U_{q,t}(\widehat{\mathfrak{g}})$  with  $q = e^{R\epsilon_1}$ ,  $t = e^{-R\epsilon_2}$ , and CY condition  $qt^{-1}(qt)^{1/2} = 1$  (for CY<sub>3</sub>:  $q_1q_2q_3 = 1$ ). The passage from  $E_2$  to  $E_1$  is not a loss of structure but a specialization of parameters: the  $\Omega$ -background IS the  $E_1$  stabilization.

This mechanism connects all three volumes: Volume I introduces  $E_1$ -chiral quantum groups as abstract algebraic-geometric objects; this volume constructs them concretely from the  $\Omega$ -background of gauge theories on CY manifolds; Volume II realizes their derived centres as 3d gauge theories. The  $\Omega$ -background is the physical face of the  $E_1$  stabilization theorem.

**Ten research programmes from  $K3 \times E$ .** The  $K3 \times E$  prototype generates ten formal research programmes, each grounded in 90+ compute engines and detailed in Chapter 14: (A) constructive CY gluing via Bridgeland charts; (B)  $\kappa_{\text{cat}}$  as universal moonshine multiplier ( $A_n = \kappa_{\text{cat}} \cdot \dim(\rho_n)$ ); (C) the second-quantization bridge ( $\kappa_{\text{BKM}}/\kappa_{\text{ch}} = 5/3$ ); (D) Schottky shadow ( $g \geq 4$  Torelli descent); (E) denominator

= bar Euler product; (F) quantum toroidal  $U_{q,t}(\widehat{\mathfrak{gl}}_1)$  as the two-parameter chiral quantum group with CY condition  $q_1 q_2 q_3 = 1$ ; (G) BPS/DT correspondence (three-level: genus-1, virtual, motivic); (H)  $\Omega$ -background as  $E_1$  stabilization; (I) wall-crossing as MC gauge equivalence; (J) the DDCA–toroidal bridge. Each programme connects the abstract algebraic-geometric framework of Volume I to concrete enumerative geometry.

**Maturity and scope.** This volume is the least mature of the three. The CY-to-chiral functor  $\Phi$  is proved for  $d = 2$  (Theorem CY-A<sub>2</sub>) and is a programme for  $d = 3$ , conditional on the chain-level  $\mathbb{S}^3$ -framing. The quantum group realization (CY-C) is conjectural: the object  $C(\mathfrak{g}, q)$  is not constructed in general. The modular CY characteristic (CY-D) is well-defined only when  $A_C$  exists. What is proved is stated as proved; what is conjectural is stated as conjecture; what is a programme is described as such.

The following chapter-and-major-sector assessment records the honest state of each component.


- *CY categories* (~ 200 lines): definitions and basic properties only; a stub awaiting expansion of the cyclic  $A_\infty$ -structure and the CY trace.
- *Cyclic  $A_\infty$ -structures* (~ 200 lines): stub; the higher homotopy Frobenius relations are stated but not developed beyond degree 3.
- *Hochschild calculus* (~ 390 lines): the HKR theorem and basic Hochschild-to-cyclic spectral sequence; sufficient for the downstream applications but does not develop the full calculus (cap product, Connes operator, Tsygan formality).
- *$E_n$ -factorization* (~ 840 lines): the  $E_1$  stabilization theorem is proved; the framing obstruction tower is complete. The Bott-periodic table through  $d = 8$  is verified.
- *$E_1$ -chiral algebras* (~ 1,590 lines): the  $E_1$ -chiral Hopf axiom system (H<sub>1</sub>–H<sub>5</sub>) is formalized and verified computationally for Heisenberg and  $\mathcal{W}_{1+\infty}$ ; the spectral coassociativity theorem is proved from factorization. The most developed theoretical chapter.
- *$E_2$ -chiral algebras* (~ 540 lines): definition, braided factorization, and basic structural results; the passage from  $E_1$  to  $E_2$  via the Drinfeld center is stated but the full proof is deferred to the Drinfeld center chapter.
- *CY-to-chiral functor* (~ 2,220 lines): the three-step construction (cyclic  $A_\infty \rightarrow$  Lie conformal  $\rightarrow$  factorization envelope  $\rightarrow E_d$ -enhancement), proved for  $d = 2$ ; the Čech resolution for compact  $CY_3$  (quintic). Conditional on the  $\mathbb{S}^3$ -framing for  $d = 3$ .
- *Drinfeld center* (~ 820 lines): the categorical lift of the averaging map; the equivalence  $Z(\text{Rep}^{E_1}(A)) \simeq \text{Rep}^{E_2}(Z^{\text{der}}(A))$  with five invariant checks at six levels for  $\mathbb{C}^3$ .
- *Toric  $CY_3$  CoHA* (~ 930 lines): the Jordan quiver computation end-to-end; quiver-chart gluing for general toric  $CY_3$ ; shuffle algebra presentation. Proved.
- *Toroidal and elliptic algebras* (~ 6,160 lines): the crown jewel. The  $K3 \times E$  programme (K<sub>3-1</sub> through K<sub>3-13</sub>), all ten research programmes (A through J), the DDCA–toroidal bridge, the Fay trisecant identity, the M2-brane model, and the double-loop holographic dual. The most developed example chapter.
- *Fukaya categories* (~ 710 lines): symplectic input; elliptic curve and K<sub>3</sub> surface fully developed; compact  $CY_3$  conditional on the chain-level  $\mathbb{S}^3$ -framing.
- *Derived categories* (~ 200 lines): stub.
- *Matrix factorizations* (~ 210 lines): stub; ADE singularities and the LG/CY correspondence stated but not developed.
- *Quantum group representations* (~ 580 lines): Kazhdan–Lusztig equivalence, affine Yangian realization, critical CoHA; partially developed.

- *Geometric Langlands* ( $\sim 200$  lines): stub; the Langlands = Koszul duality conjecture is stated but the connection to the CY programme is not developed.
- *Bar-cobar bridge, modular Koszul bridge, holographic datum* ( $\sim 530 + 380 + 1,010$  lines): the cross-volume connections; partially developed.

The  $K3 \times E$  sector inside the toroidal-and-elliptic chapter, with its thirteen structural results and ten research directions, is the most developed component; the remaining landscape (toric  $CY_3$ , Fukaya categories, matrix factorizations, geometric Langlands) consists of varying mixtures of theorems, programmes, and conjectures.

**The three volumes.** Volume I constructs  $E_n$ -chiral algebras as algebraic-geometric objects on curves and their configuration spaces: the bar complex on every curve geometry,  $R$ -matrices, Yangian structures,  $E_1$ -chiral quantum groups as abstract algebraic-geometric objects, derived chiral centres with  $E_2/E_3$  structure, the  $SC^{\text{ch},\text{top}}$  datum on the pair  $(\mathcal{Z}_{\text{ch}}^{\text{der}}(A), A)$ , and the five theorems as structural properties of the categorical logarithm. Volume II interprets these constructions physically: the derived chiral centre IS the bulk of a real 3d holomorphic-topological gauge theory; the  $SC^{\text{ch},\text{top}}$  pair IS a boundary-bulk system; 3d quantum gravity is the climax. This volume constructs the concrete examples: the functor  $\Phi: CY_d\text{-Cat} \rightarrow E_2\text{-ChirAlg}$  produces chiral algebras from Calabi–Yau categories, and the CY quantum groups (CoHA, quantum toroidal algebras, BPS algebras) are concrete  $E_1$ -chiral quantum groups instantiating Volume I’s abstract framework. The theories studied include 6d holomorphic Chern–Simons on  $CY_3$  (the primary focus), 4d and 5d HT Chern–Simons, 6d little string theory, and  $M_5$  brane theory.

The three volumes form a triangle:



Volume III closes the triangle from below: the CY category  $C$  sits beneath all three vertices, supplying the chiral algebra via  $\Phi$ , the modular trace via the CY Euler characteristic, and the quantum group via the CoHA.

At a higher altitude, the entire programme is factorization (co)homology at varying geometric inputs. Volume I computes factorization homology on the curve  $X$ : the chiral algebra  $A$  is the input, the bar complex  $B(A)$  on  $\text{Ran}(X)$  is the output, and the five theorems extract genus-by-genus invariants. Volume II computes factorization homology on the slab  $\mathbb{C} \times \mathbb{R}$ : the bar complex  $B(A)$  is an  $E_1$  dg coalgebra whose differential encodes holomorphic data along  $\mathbb{C}$  and whose coproduct encodes topological data along  $\mathbb{R}$ , and the output is the 3d holomorphic-topological field theory. This volume computes factorization homology on the CY category  $C$ : the cyclic bar complex  $CC_*(C)$  is the input to  $\Phi$ , and the output is the chiral algebra that feeds Volumes I–II. The geometric input changes; the machine is the same.

**Organisation.** Part I establishes the categorical foundations: CY categories, cyclic  $A_\infty$ -structures, the Hochschild calculus, and the  $E_1/E_2/E_n$  chiral hierarchy with the framing obstruction tower. Part II constructs the bridge functor  $\Phi$  from CY categories to quantum chiral algebras and proves Theorem CY-A<sub>2</sub> for  $d = 2$ , with the identification of automorphic correction and shadow obstruction tower as the main structural comparison. In Part II, the Drinfeld center chapter gives the categorical mechanism that lifts  $E_1$ -monoidal representation categories to  $E_2$ -braided monoidal categories, realizing the quantum group  $R$ -matrix as the half-braiding of the center construction. Part III works through the standard CY landscape,

centred on the  $K3 \times E$  programme with its thirteen structural results and ten research programmes. Part IV develops the seven faces of  $r_{CY}(z)$ : the bar-cobar bridge to Volume I, the CY holographic datum, and the modular Koszul bridge. Part V connects to the geometric Langlands programme and identifies the open frontiers.

Volume I asks what the logarithm does. Volume II asks how it composes. Volume III asks where it comes from.

# Contents

|                                                                  |           |
|------------------------------------------------------------------|-----------|
| <b>Preface</b>                                                   | <b>3</b>  |
| <b>Contents</b>                                                  | <b>i</b>  |
| <b>I The CY Engine</b>                                           | <b>I</b>  |
| <b>1 Introduction</b>                                            | <b>3</b>  |
| 1.1 The question . . . . .                                       | 3         |
| 1.2 The $E_1/E_2$ chiral hierarchy . . . . .                     | 4         |
| The $E_1$ stabilization theorem . . . . .                        | 4         |
| 1.3 Relation to Volumes I and II . . . . .                       | 5         |
| 1.4 The analytic gap and the Čech resolution . . . . .           | 6         |
| 1.5 $CY_3$ combinatorics as generalized root data . . . . .      | 7         |
| 1.6 Automorphic correction as shadow obstruction tower . . . . . | 8         |
| 1.7 Main results . . . . .                                       | 8         |
| 1.8 What is proved versus what is conjectural . . . . .          | 9         |
| 1.9 The $E_1$ -chiral bialgebra and Hopf axioms . . . . .        | 11        |
| 1.10 The ten research programmes . . . . .                       | 11        |
| 1.11 The landscape of examples . . . . .                         | 13        |
| 1.12 Guide for the reader . . . . .                              | 14        |
| 1.13 Conventions and notation . . . . .                          | 14        |
| <b>2 Calabi–Yau Categories</b>                                   | <b>17</b> |
| 2.1 Smooth and proper dg categories . . . . .                    | 17        |
| 2.2 The Calabi–Yau condition . . . . .                           | 17        |
| 2.3 Hochschild cohomology and the CY structure . . . . .         | 18        |
| 2.4 Cyclic $A_\infty$ structure . . . . .                        | 19        |
| 2.5 Interface with the CY-to-chiral functor $\Phi$ . . . . .     | 20        |
| <b>3 Cyclic <math>A_\infty</math>-Structures</b>                 | <b>23</b> |
| 3.1 $A_\infty$ -algebras . . . . .                               | 23        |
| 3.2 Cyclic structures and CY dimension . . . . .                 | 24        |
| 3.3 Examples . . . . .                                           | 25        |
| 3.4 The bridge to the Vol III functor . . . . .                  | 26        |

|          |                                                                    |           |
|----------|--------------------------------------------------------------------|-----------|
| <b>4</b> | <b>Hochschild Calculus for CY Categories</b>                       | <b>29</b> |
| 4.1      | Hochschild homology and cohomology . . . . .                       | 29        |
| 4.2      | The Gerstenhaber bracket . . . . .                                 | 30        |
| 4.3      | The Connes $B$ -operator and cyclic homology . . . . .             | 31        |
| 4.4      | The Hochschild-to-cyclic spectral sequence . . . . .               | 31        |
| 4.5      | CY Hochschild duality . . . . .                                    | 32        |
| 4.6      | The $(2 - d)$ -shifted Poisson structure . . . . .                 | 32        |
| 4.7      | Categorical Hodge theory . . . . .                                 | 33        |
| <b>5</b> | <b><math>E_1</math>-Chiral Algebras</b>                            | <b>35</b> |
| 5.1      | Factorization algebras and the $E_n$ hierarchy . . . . .           | 35        |
| 5.2      | The ordered bar as the $E_1$ primitive . . . . .                   | 36        |
| 5.3      | $E_1$ -chiral algebras from CY categories . . . . .                | 37        |
| 5.4      | Connection to Volume II . . . . .                                  | 39        |
| 5.5      | Operadic dictionary for the CY reader . . . . .                    | 40        |
| 5.6      | $E_1$ -chiral Koszul duality: three standard families . . . . .    | 41        |
|          | Family I: Heisenberg $H_k$ (class G) . . . . .                     | 42        |
|          | Family II: Kac–Moody $V_k(\mathfrak{sl}_2)$ (class L) . . . . .    | 43        |
|          | Family III: Virasoro $\mathcal{C}$ (class M) . . . . .             | 44        |
| 5.7      | $E_1$ -chiral bialgebras . . . . .                                 | 46        |
|          | The $E_1$ -chiral monoidal category . . . . .                      | 46        |
|          | Algebra and coalgebra objects . . . . .                            | 47        |
|          | The $E_1$ -chiral bialgebra axiom . . . . .                        | 47        |
|          | Why $E_1$ and not $E_\infty$ . . . . .                             | 50        |
|          | The Drinfeld center recovers $E_2$ . . . . .                       | 51        |
|          | Examples . . . . .                                                 | 52        |
|          | The factorization-homology coproduct . . . . .                     | 53        |
| 5.8      | Closing: the ordered bar as the primitive of Vol III . . . . .     | 56        |
| <b>6</b> | <b><math>E_2</math>-Chiral Algebras</b>                            | <b>57</b> |
| 6.1      | The $E_2$ operad and its categorical content . . . . .             | 57        |
| 6.2      | $E_2$ -chiral algebras from cyclic $A_\infty$ data . . . . .       | 57        |
| 6.3      | Braided tensor categories from $E_2$ -chiral structure . . . . .   | 58        |
| 6.4      | Connection to Volume II: the Drinfeld center . . . . .             | 59        |
| 6.5      | $E_2$ -chiral Koszul duality and the three bar complexes . . . . . | 59        |
| <b>7</b> | <b><math>E_n</math>-Factorization and Higher Chiral Structure</b>  | <b>65</b> |
| 7.1      | Dunn additivity and the $E_n$ hierarchy . . . . .                  | 65        |
| 7.2      | Factorization algebras on $\mathbb{C}^n$ . . . . .                 | 66        |
| 7.3      | The $E_n$ -chiral bar complex . . . . .                            | 66        |
| 7.4      | The framing obstruction tower . . . . .                            | 66        |
| 7.5      | The $E_n$ landscape: the $E_1$ stabilization theorem . . . . .     | 67        |
| 7.6      | Explicit computations for $d = 4$ varieties . . . . .              | 69        |
|          | $\mathbb{C}^4$ . . . . .                                           | 69        |
|          | $K3 \times K3$ . . . . .                                           | 70        |
|          | The sextic 4-fold in $\mathbb{P}^5$ . . . . .                      | 70        |



|      |                                                                                   |    |
|------|-----------------------------------------------------------------------------------|----|
| 7.7  | The $d = 5$ functor: $\mathbb{Z}_2$ obstruction from Sp . . . . .                 | 70 |
| 7.8  | The $d = 6$ case and higher . . . . .                                             | 71 |
| 7.9  | Relation to topological field theory . . . . .                                    | 71 |
| 7.10 | $E_3$ -chiral structure from holomorphic Chern–Simons on $\mathbb{C}^3$ . . . . . | 72 |
|      | The $E_n$ hierarchy from theory dimension . . . . .                               | 72 |
|      | $E_3$ -chiral factorization on $\mathbb{C}^3$ : precise conditions . . . . .      | 72 |
|      | $E_3$ -chiral Koszul duality . . . . .                                            | 73 |
|      | The Miki automorphism from the CY torus Weyl group on $\mathbb{C}^3$ . . . . .    | 78 |
|      | The $S_3$ -embedding in $SO(3)$ . . . . .                                         | 78 |
|      | From holomorphic torus action to Miki triality . . . . .                          | 79 |
|      | Rational degeneration and $S_3$ -breaking . . . . .                               | 80 |
|      | Factorization homology on $\mathbb{C}^3$ and the chiral quantum group . . . . .   | 80 |
|      | The Nekrasov partition function as $E_3$ factorization homology . . . . .         | 81 |
| 7.11 | The $E_n$ -chiral Koszul duality cascade . . . . .                                | 83 |
|      | The pattern . . . . .                                                             | 84 |
|      | Comparison table . . . . .                                                        | 84 |
|      | Obstructions at each level . . . . .                                              | 85 |

## II The CY Characteristic Datum 87

|          |                                                                         |           |
|----------|-------------------------------------------------------------------------|-----------|
| <b>8</b> | <b>From CY Categories to Chiral Algebras</b>                            | <b>89</b> |
| 8.1      | The cyclic-to-chiral passage . . . . .                                  | 89        |
| 8.2      | The CY-to-chiral functor . . . . .                                      | 90        |
| 8.3      | The $d = 3$ functor chain for $\mathbb{C}^3$ . . . . .                  | 96        |
|          | The five-step chain . . . . .                                           | 96        |
|          | Abelianity of the classical bracket . . . . .                           | 98        |
|          | Hochschild cohomology and the unique deformation . . . . .              | 98        |
|          | Necessity of the $\Omega$ -deformation for noncompact $CY_3$ . . . . .  | 99        |
| 8.4      | The $\mathbb{S}^3$ -framing obstruction: universal triviality . . . . . | 100       |
|          | The Čech contracting homotopy for compact $CY_3$ . . . . .              | 101       |
| 8.5      | The Drinfeld center and $E_2$ enhancement . . . . .                     | 103       |
|          | The CY condition and the universal $R$ -matrix . . . . .                | 103       |
| 8.6      | The bar complex and shadow tower of $\mathbb{C}^3$ . . . . .            | 104       |
| 8.7      | $E_1$ universality for $CY_3$ chiral algebras . . . . .                 | 106       |
| 8.8      | The quiver-chart gluing construction . . . . .                          | 109       |
|          | Quiver chart atlases . . . . .                                          | 109       |
|          | The $E_1$ -chiral hocolim . . . . .                                     | III       |
|          | $E_1$ descent and the Čech spectral sequence . . . . .                  | 116       |
|          | The Čech nerve of a chart cover . . . . .                               | 116       |
|          | The $E_1$ descent spectral sequence . . . . .                           | 117       |
|          | Explicit degeneration for standard $CY_3$ atlases . . . . .             | 119       |
|          | The punchline: why $d = 3$ gives $E_1$ . . . . .                        | 120       |
|          | The three-component $E_1 \rightarrow E_2$ obstruction . . . . .         | 121       |
|          | The center-hocolim obstruction . . . . .                                | 122       |
|          | Bar-hocolim commutation . . . . .                                       | 123       |

|           |                                                                            |            |
|-----------|----------------------------------------------------------------------------|------------|
|           | DT = hocolim shadow . . . . .                                              | 123        |
|           | Connection to the factorization envelope . . . . .                         | 126        |
|           | Standard $CY_3$ atlas data . . . . .                                       | 127        |
| 8.9       | The CY-to-chiral programme at $d = 3$ . . . . .                            | 127        |
|           | Status summary . . . . .                                                   | 130        |
| 8.10      | Compatibility with Koszul duality . . . . .                                | 131        |
| 8.11      | The CY modular characteristic . . . . .                                    | 131        |
| 8.12      | Cross-volume bridges . . . . .                                             | 133        |
|           | Shadow tower identification . . . . .                                      | 133        |
|           | $E_1$ -chiral coalgebra and the derived center . . . . .                   | 133        |
|           | The BCOV holomorphic anomaly equation as genus spectral sequence . . . . . | 134        |
|           | The completed modular Koszul datum for $CY_3$ . . . . .                    | 134        |
| 8.13      | Beyond $CY_3$ : chiral algebras from non-CY geometries . . . . .           | 135        |
| <b>9</b>  | <b>Quantum Chiral Algebras</b> . . . . .                                   | <b>137</b> |
| 9.1       | Definition and structure . . . . .                                         | 137        |
| 9.2       | The CY $R$ -matrix . . . . .                                               | 138        |
| 9.3       | Drinfeld center and the $E_1 \rightarrow E_2$ passage . . . . .            | 138        |
| 9.4       | Comparison with the Maulik–Okounkov $R$ -matrix . . . . .                  | 140        |
| 9.5       | Quantum chiral algebras from holomorphic Chern–Simons theory . . . . .     | 141        |
|           | The chiral Chevalley–Eilenberg complex . . . . .                           | 143        |
|           | The universal defect and holomorphic Wilson lines . . . . .                | 144        |
|           | Codimension-2 defect OPE from the 6d action . . . . .                      | 146        |
| 9.6       | The quantum chiral Koszul dual . . . . .                                   | 148        |
|           | Chiral Koszul duality: boundary and defect . . . . .                       | 148        |
|           | The $E_n$ level of the Koszul dual . . . . .                               | 149        |
|           | Deformation of chiral CE cochains to $E_3$ . . . . .                       | 150        |
|           | Chiral quantum groups from defect Wilson lines . . . . .                   | 151        |
| <b>10</b> | <b>The Modular Trace</b> . . . . .                                         | <b>153</b> |
| 10.1      | CY trace as modular characteristic . . . . .                               | 153        |
| 10.2      | CY Euler characteristic versus modular characteristic . . . . .            | 154        |
| 10.3      | Modularity constraints on the shadow tower . . . . .                       | 154        |
| 10.4      | Genus expansion and Gromov–Witten invariants . . . . .                     | 156        |
| 10.5      | The CY shadow obstruction tower . . . . .                                  | 156        |
| 10.6      | Complementarity for CY pairs . . . . .                                     | 156        |
| <b>11</b> | <b>Quantum Groups: Foundations</b> . . . . .                               | <b>157</b> |
| 11.1      | Quantized enveloping algebras . . . . .                                    | 157        |
| 11.2      | The universal $R$ -matrix and Yang–Baxter equation . . . . .               | 158        |
| 11.3      | Representation categories . . . . .                                        | 159        |
| 11.4      | The four regimes . . . . .                                                 | 161        |
| 11.5      | Connection to Conjecture CY-C . . . . .                                    | 161        |
| <b>12</b> | <b>Braided Factorization</b> . . . . .                                     | <b>163</b> |
| 12.1      | The categorical $R$ -matrix and Yang–Baxter equation . . . . .             | 163        |

|            |                                                                            |            |
|------------|----------------------------------------------------------------------------|------------|
| 12.2       | Braided monoidal categories and $E_2$ -algebras . . . . .                  | 164        |
| 12.3       | Factorization homology with $E_2$ -coefficients . . . . .                  | 164        |
| 12.4       | The $E_2$ -equivariant bar complex . . . . .                               | 165        |
| 12.5       | The braided bar-cobar adjunction . . . . .                                 | 166        |
| 12.6       | Quantum group structure from braided Koszul duality . . . . .              | 167        |
| 12.7       | The braided shadow obstruction tower . . . . .                             | 168        |
| 12.8       | Braided complementarity . . . . .                                          | 169        |
| 12.9       | The categorical $S$ -matrix and $E_3$ scattering . . . . .                 | 169        |
| <b>13</b>  | <b>The Drinfeld Center and Bulk Algebras</b>                               | <b>173</b> |
| 13.1       | The Drinfeld center of a monoidal category . . . . .                       | 173        |
| 13.2       | Drinfeld center as chiral derived center . . . . .                         | 174        |
|            | The $\mathbb{C}^3$ theorem: Yangian and $\mathcal{W}_{1+\infty}$ . . . . . | 175        |
|            | The Ben-Zvi–Francis–Nadler loop space identification . . . . .             | 176        |
| 13.3       | Bulk-boundary correspondence . . . . .                                     | 176        |
| 13.4       | CY enhancement of the Drinfeld center . . . . .                            | 177        |
| 13.5       | Categorical Theorem H . . . . .                                            | 177        |
| 13.6       | Quantum group reconstruction from the center . . . . .                     | 177        |
| 13.7       | The Drinfeld double and the slab . . . . .                                 | 178        |
| 13.8       | The $E_n$ -circle and the categorified averaging map . . . . .             | 179        |
| 13.9       | The $E_3 \rightarrow E_2$ restriction and symmetric braiding . . . . .     | 180        |
| <b>III</b> | <b>The CY Landscape</b>                                                    | <b>183</b> |
| <b>14</b>  | <b>Toroidal and elliptic algebras</b>                                      | <b>185</b> |
| 14.1       | Double affine setup . . . . .                                              | 186        |
|            | Definition . . . . .                                                       | 186        |
|            | Koszul dual of the toroidal algebra . . . . .                              | 188        |
|            | Connection to the three main theorems of Vol I . . . . .                   | 189        |
| 14.2       | Elliptic quantum groups . . . . .                                          | 189        |
|            | Felder’s elliptic quantum group . . . . .                                  | 189        |
| 14.3       | Elliptic $R$ -matrices and theta functions . . . . .                       | 190        |
|            | Theta function identities . . . . .                                        | 190        |
|            | Fay identity and bar nilpotency . . . . .                                  | 190        |
|            | Bar complex with theta functions . . . . .                                 | 191        |
| 14.4       | Explicit elliptic bar complex for the Heisenberg algebra . . . . .         | 193        |
|            | Bar elements and the elliptic propagator . . . . .                         | 193        |
|            | Comparison table: elliptic versus rational . . . . .                       | 195        |
| 14.5       | Dynamical Yang–Baxter equation for $\mathfrak{sl}_2$ . . . . .             | 196        |
|            | The dynamical Yang–Baxter equation . . . . .                               | 196        |
|            | Explicit verification for $\mathfrak{sl}_2$ . . . . .                      | 196        |
| 14.6       | Shuffle algebra presentation . . . . .                                     | 198        |
|            | Definition of the shuffle algebra . . . . .                                | 198        |
|            | Explicit shuffle products . . . . .                                        | 199        |
| 14.7       | Comparison: rational, trigonometric, elliptic . . . . .                    | 199        |

|       |                                                                              |     |
|-------|------------------------------------------------------------------------------|-----|
|       | The elliptic curve as shared geometry . . . . .                              | 199 |
| 14.8  | Calabi–Yau local-to-global gluing . . . . .                                  | 200 |
|       | McKay correspondence and ADE quiver charts . . . . .                         | 200 |
|       | Ginzburg dg algebras and Jacobian algebras . . . . .                         | 201 |
|       | $A_\infty$ -bimodule transition data . . . . .                               | 202 |
|       | Čech descent and the global category . . . . .                               | 202 |
|       | Wall-crossing during chart transitions . . . . .                             | 202 |
| 14.9  | The $K_3$ chiral algebra . . . . .                                           | 203 |
|       | The $\mathcal{N} = 4$ superconformal algebra at $c = 6$ . . . . .            | 203 |
|       | Bar complex of the $\mathcal{N} = 4$ SCA . . . . .                           | 204 |
|       | Shadow depth classification . . . . .                                        | 205 |
|       | The $K_3$ elliptic genus and Mathieu moonshine . . . . .                     | 205 |
|       | Lattice VOA and Gepner model . . . . .                                       | 207 |
|       | Chiral de Rham complex on $K_3$ . . . . .                                    | 207 |
|       | Koszul dual of the $K_3$ chiral algebra . . . . .                            | 207 |
|       | The abelian $(2, 0)$ pushforward . . . . .                                   | 207 |
| 14.10 | The Calabi–Yau threefold $K3 \times E$ : geometry and invariants . . . . .   | 209 |
|       | Hodge diamond and topological invariants . . . . .                           | 209 |
|       | HKR decomposition and deformation space . . . . .                            | 209 |
|       | Noncommutative deformations and Brauer group . . . . .                       | 210 |
|       | Autoequivalences and semiorthogonal structure . . . . .                      | 210 |
|       | Factorization homology . . . . .                                             | 210 |
| 14.11 | Enumerative geometry of $K3 \times E$ . . . . .                              | 210 |
|       | Gromov–Witten theory and BPS invariants . . . . .                            | 210 |
|       | Bridgeland stability on $K_3$ . . . . .                                      | 211 |
| 14.12 | BPS spectrum and black hole entropy . . . . .                                | 211 |
|       | Charge lattice and BPS states . . . . .                                      | 211 |
|       | Black hole entropy from $1/\Phi_{10}$ . . . . .                              | 211 |
|       | Fourier–Jacobi as genus escalation . . . . .                                 | 212 |
| 14.13 | Twisted holography on $K3 \times E$ . . . . .                                | 212 |
| 14.14 | Mock modular forms and the BKM root system . . . . .                         | 213 |
|       | BKM root multiplicities . . . . .                                            | 213 |
|       | The Hilbert scheme bridge . . . . .                                          | 214 |
|       | Fourier–Jacobi expansion . . . . .                                           | 215 |
| 14.15 | Grand synthesis and open problems . . . . .                                  | 215 |
|       | Research programmes . . . . .                                                | 218 |
| 14.16 | The Borcherds–Kac–Moody algebra of $K3 \times E$ . . . . .                   | 226 |
|       | The Igusa cusp form and the DVV formula . . . . .                            | 226 |
|       | The Borcherds multiplicative lift . . . . .                                  | 227 |
|       | Shadow tower data for $K3 \times E$ . . . . .                                | 227 |
|       | The shadow–Siegel gap . . . . .                                              | 228 |
|       | Euler characteristic and Donaldson–Thomas structure . . . . .                | 229 |
| 14.17 | The $CY_3$ geometry . . . . .                                                | 231 |
| 14.18 | The lattice $\Lambda^{3,2}$ and the isomorphism $Sp_4 \simeq SO_+$ . . . . . | 231 |
| 14.19 | The hyperbolic sublattice and reflection groups . . . . .                    | 232 |

|           |                                                                                     |            |
|-----------|-------------------------------------------------------------------------------------|------------|
| 14.20     | The generalized BKM superalgebra $\mathfrak{g}_{\Delta_5}$                          | 232        |
| 14.21     | The denominator identity: $\Delta_5 = \Phi$                                         | 233        |
| 14.22     | The weak Jacobi form $\phi_{0,1}$ and root multiplicities                           | 233        |
| 14.23     | The quantum vertex chiral group $G(K3 \times E)$                                    | 233        |
| 14.24     | CoHA structure and the BKM superalgebra                                             | 234        |
| 14.25     | The relative chiral algebra                                                         | 235        |
| 14.26     | From fiber $\kappa_{\text{ch}}$ to global $\kappa_{\text{ch}}$ : the Borcherds lift | 237        |
| 14.27     | The Maulik–Okounkov $R$ -matrix                                                     | 237        |
| 14.28     | Diophantine gaps and the $D$ -stratification                                        | 238        |
| 14.29     | Euler characteristics and the modular characteristic                                | 238        |
| 14.30     | Five routes to $G(K3 \times E)$                                                     | 239        |
| 14.31     | DT and GW invariants of $K3 \times E$                                               | 240        |
|           | Degree-zero DT and the MacMahon function                                            | 240        |
|           | DT/PT correspondence                                                                | 240        |
|           | Yau–Zaslow BPS counts                                                               | 240        |
| 14.32     | Bridgeland stability and wall-crossing                                              | 241        |
|           | Stability conditions on $K3$                                                        | 241        |
|           | Walls of marginal stability                                                         | 242        |
|           | Kontsevich–Soibelman wall-crossing formula                                          | 242        |
| 14.33     | The Borcherds lift: genus-1 to genus-2 bridge                                       | 243        |
|           | Multiplicative lift (Borcherds product)                                             | 243        |
|           | Additive lift (Saito–Kurokawa / Maass)                                              | 243        |
|           | The genus-1 to genus-2 bridge                                                       | 244        |
| 14.34     | Scattering diagrams and the shadow MC element                                       | 244        |
|           | The scattering diagram of $K3 \times E$                                             | 244        |
|           | Shadow tower and scattering: two MC elements                                        | 245        |
| 14.35     | The complete enumerative–algebraic dictionary                                       | 246        |
| 14.36     | Cross-volume structural results                                                     | 246        |
| 14.37     | Costello’s M2-brane model and the double-loop holographic dual                      | 247        |
|           | The 5d noncommutative Chern–Simons theory                                           | 247        |
|           | The classical double-current algebra                                                | 247        |
|           | The Koszul dual bulk $L_\infty$ model                                               | 248        |
|           | Quantum double-loop deformation                                                     | 248        |
|           | Sector/level/genus matching                                                         | 248        |
|           | The DDCA–toroidal bridge                                                            | 249        |
| 14.38     | The elliptic Sklyanin bracket and the three-parameter $\hbar$                       | 252        |
| <b>15</b> | <b>Toric CY<sub>3</sub> and Critical CoHAs</b>                                      | <b>255</b> |
| 15.1      | Toric CY <sub>3</sub> threefolds and their quivers                                  | 255        |
| 15.2      | The critical cohomological Hall algebra                                             | 256        |
| 15.3      | $\mathbb{C}^3$ and the affine Yangian $Y(\widehat{\mathfrak{gl}}_1)$                | 256        |
| 15.4      | General toric CY <sub>3</sub> and affine super Yangians                             | 256        |
| 15.5      | The CoHA as $E_1$ -sector                                                           | 256        |
| 15.6      | Root datum from toric geometry                                                      | 257        |
| 15.7      | The conifold bar complex                                                            | 257        |

|           |                                                                           |            |
|-----------|---------------------------------------------------------------------------|------------|
| 15.8      | Local $\mathbb{P}^2$ . . . . .                                            | 258        |
| 15.9      | Local $\mathbb{P}^1 \times \mathbb{P}^1$ . . . . .                        | 259        |
| 15.10     | 5d Chern–Simons theory . . . . .                                          | 260        |
| 15.11     | Wall-crossing as MC gauge equivalence . . . . .                           | 260        |
| 15.12     | The $\kappa_{\text{ch}}$ -table for toric $\text{CY}_3$ . . . . .         | 261        |
| 15.13     | Gaudin Hamiltonians from toric $\text{CY}_3$ collision residues . . . . . | 262        |
| 15.14     | The chiral quantum group of a toric $\text{CY}_3$ . . . . .               | 263        |
| 15.15     | The chiral quantum group equivalence for toric $\text{CY}_3$ . . . . .    | 265        |
| <b>16</b> | <b>Fukaya Categories</b> . . . . .                                        | <b>269</b> |
| 16.1      | The Fukaya category: definition and $A_\infty$ -structure . . . . .       | 269        |
| 16.2      | CY 1-folds: elliptic curves . . . . .                                     | 270        |
| 16.3      | CY 2-folds: $K_3$ and abelian surfaces . . . . .                          | 270        |
| 16.4      | CY 3-folds . . . . .                                                      | 271        |
| 16.5      | Wrapped Fukaya categories . . . . .                                       | 272        |
| 16.6      | Homological mirror symmetry and the CY-to-chiral functor . . . . .        | 273        |
| 16.7      | The quantum chiral algebra of a Fukaya category . . . . .                 | 274        |
| 16.8      | Homological mirror symmetry and chiral algebra equivalence . . . . .      | 275        |
| 16.9      | Open-string sector and the Swiss-cheese structure . . . . .               | 276        |
| <b>17</b> | <b>Derived Categories of CY Manifolds</b> . . . . .                       | <b>279</b> |
| 17.1      | $D^b(\text{Coh}(X))$ as a $\text{CY}_d$ category . . . . .                | 279        |
| 17.2      | Homological mirror symmetry as chiral equivalence . . . . .               | 281        |
| 17.3      | Exceptional collections, quivers, and local CY . . . . .                  | 282        |
| 17.4      | Bridgeland stability conditions and wall-crossing . . . . .               | 283        |
| <b>18</b> | <b>Matrix Factorizations</b> . . . . .                                    | <b>285</b> |
| 18.1      | $\text{MF}(W)$ as a CY category . . . . .                                 | 285        |
| 18.2      | LG/CY correspondence . . . . .                                            | 287        |
| 18.3      | ADE singularities and $\mathcal{W}$ -algebras . . . . .                   | 288        |
| <b>19</b> | <b>Quantum Group Representations</b> . . . . .                            | <b>291</b> |
| 19.1      | $\text{Rep}_q(\mathfrak{g})$ as a braided monoidal category . . . . .     | 291        |
| 19.2      | The $R$ -matrix as categorical $r(z)$ . . . . .                           | 292        |
| 19.3      | Kazhdan–Lusztig equivalence . . . . .                                     | 293        |
| 19.4      | Conjecture CY-C: quantum group realization . . . . .                      | 293        |
| 19.5      | Yangian and RTT realizations . . . . .                                    | 294        |
| 19.6      | BPS algebras and quantum groups . . . . .                                 | 294        |
| 19.7      | Wall-crossing and Kontsevich–Soibelman . . . . .                          | 295        |
| 19.8      | The modular characteristic $\kappa_{\text{cat}}$ . . . . .                | 296        |
| 19.9      | Shadow obstruction tower for quantum group categories . . . . .           | 297        |
| <b>IV</b> | <b>The Seven Faces of <math>r_{\text{CY}}(z)</math></b> . . . . .         | <b>299</b> |
| <b>20</b> | <b>The Bar-Cobar Bridge to Volume I</b> . . . . .                         | <b>301</b> |
| 20.1      | CY bar complex vs chiral bar complex . . . . .                            | 301        |

|           |                                                                                                   |            |
|-----------|---------------------------------------------------------------------------------------------------|------------|
| 20.2      | The shadow tower of $\mathbb{C}^3$ : MacMahon meets Koszul . . . . .                              | 302        |
| 20.3      | Open string field theory: Koszul duality at chain level . . . . .                                 | 303        |
| 20.4      | From shuffle product to $\lambda$ -bracket: the seven-step chain . . . . .                        | 303        |
| 20.5      | Koszul duality: CY vs chiral . . . . .                                                            | 304        |
| 20.6      | The five theorems in the CY setting . . . . .                                                     | 305        |
| 20.7      | Compact $\text{CY}_3$ examples . . . . .                                                          | 305        |
|           | The banana manifold: $\kappa_{\text{ch}} = 0$ with nontrivial tower . . . . .                     | 305        |
|           | Enriques $\times E$ : the $\mathbb{Z}_2$ quotient of $K3 \times E$ . . . . .                      | 306        |
|           | The quintic threefold: non-integer $\kappa_{\text{ch}}$ and BKM obstruction . . . . .             | 307        |
|           | Hochschild homology of compact $\text{CY}_3$ examples . . . . .                                   | 307        |
| 20.8      | Integrable systems and lattice models . . . . .                                                   | 308        |
|           | The Calogero–Moser/Bethe dictionary . . . . .                                                     | 308        |
|           | Lattice finite-size corrections . . . . .                                                         | 308        |
|           | Crystal melting: combinatorial vs bar-algebraic degree . . . . .                                  | 309        |
| 20.9      | Grand atlas of $\text{CY}_3$ shadow invariants . . . . .                                          | 309        |
| <b>21</b> | <b>The seven faces of <math>r_{\text{CY}}(z)</math> for Calabi–Yau chiral algebras</b>            | <b>311</b> |
| 21.1      | The $\text{CY}_3$ collision residue . . . . .                                                     | 311        |
| 21.2      | Face 1: the CY bar-cobar twisting morphism . . . . .                                              | 312        |
| 21.3      | Face 2: CoHA and perverse coherent sheaves . . . . .                                              | 314        |
| 21.4      | Face 3: classical CY Poisson coisson (genus 0 only) . . . . .                                     | 314        |
| 21.5      | Face 4: the Maulik–Okounkov $R$ -matrix (for $K3 \times E$ ) . . . . .                            | 315        |
| 21.6      | Face 5: the affine super Yangian $Y(\widehat{\mathfrak{gl}}_1)$ for toric $\text{CY}_3$ . . . . . | 316        |
|           | Gaudin Hamiltonians for toric $\text{CY}_3$ from collision residues . . . . .                     | 317        |
| 21.7      | Face 6: elliptic Sklyanin bracket (for toroidal CY) . . . . .                                     | 317        |
|           | The three-parameter $\hbar$ identification . . . . .                                              | 318        |
| 21.8      | Face 7: Gaudin model from $\text{CY}_3$ collision residues . . . . .                              | 318        |
| 21.9      | The seven-face master theorem in the CY setting . . . . .                                         | 319        |
| 21.10     | Cross-volume bridge . . . . .                                                                     | 320        |
| <b>22</b> | <b>Modular Koszul Duality and CY Geometry</b>                                                     | <b>323</b> |
| 22.1      | The modular convolution algebra for CY categories . . . . .                                       | 323        |
| 22.2      | CY complementarity . . . . .                                                                      | 325        |
|           | The $d = 2$ theorem . . . . .                                                                     | 326        |
|           | The $d = 3$ conjecture . . . . .                                                                  | 327        |
| 22.3      | The CY shadow CohFT . . . . .                                                                     | 327        |
|           | CohFT structure . . . . .                                                                         | 327        |
|           | The $K3 \times E$ CohFT and the Igusa cusp form . . . . .                                         | 328        |
|           | Toric $\text{CY}_3$ and the DT CohFT . . . . .                                                    | 329        |
| 22.4      | The Hochschild bridge . . . . .                                                                   | 329        |
| 22.5      | Principal examples . . . . .                                                                      | 330        |
|           | $K3 \times E$ . . . . .                                                                           | 330        |
|           | $\mathbb{C}^3$ and $\mathcal{W}_{1+\infty}$ . . . . .                                             | 331        |

|                                                                     |            |
|---------------------------------------------------------------------|------------|
| <b>V The CY Frontier</b>                                            | <b>333</b> |
| <b>23 Geometric Langlands and CY Quantum Groups</b>                 | <b>335</b> |
| 23.1 The Feigin–Frenkel center at the critical level . . . . .      | 335        |
| 23.2 CY Langlands for $d = 1, 2, 3$ . . . . .                       | 337        |
| 23.3 Quantum geometric Langlands and the shadow parameter . . . . . | 339        |
| 23.4 Open problems . . . . .                                        | 340        |
| <b>A Conventions</b>                                                | <b>343</b> |
| A.1 Grading conventions . . . . .                                   | 343        |
| A.2 Sign conventions . . . . .                                      | 343        |
| A.3 Categorical conventions . . . . .                               | 343        |
| A.4 Operadic conventions . . . . .                                  | 343        |
| A.5 Bar complex and Koszul duality conventions . . . . .            | 343        |
| <b>Bibliography</b>                                                 | <b>345</b> |



## **Part I**

# **The CY Engine**



# Chapter I

## Introduction

### I.1 The question

A Calabi–Yau category  $C$  of dimension  $d$  carries a canonical cyclic  $A_\infty$ -structure: a non-degenerate trace

$$\mathrm{Tr}: \mathrm{HH}_\bullet(C) \longrightarrow k[-d]$$

on its Hochschild homology, encoding the CY condition as a  $d$ -dimensional Frobenius structure at the chain level. The cyclic bar complex  $\mathrm{CC}_\bullet(C)$  is the primary invariant; it carries an  $\mathbb{S}^d$ -action from the  $d$ -sphere framing of the trace, and its  $\mathbb{S}^d$ -equivariant structure governs higher-genus amplitudes.

A chiral algebra  $A$  on a curve  $X$  carries a bar complex  $B(A)$ , a factorization coalgebra on  $\mathrm{Ran}(X)$  whose differential encodes holomorphic OPE residues and whose coproduct encodes topological interval-cutting. At genus  $g \geq 1$ , the bar complex acquires curvature  $\kappa_{\mathrm{ch}}(A) \cdot \omega_g$  from the Hodge bundle, and the full modular structure is controlled by the universal Maurer–Cartan element  $\Theta_A := D_A - d_0$  (Volume I). Together with the  $\mathrm{SC}^{\mathrm{ch}, \mathrm{top}}$  structure on the derived center pair  $(C_{\mathrm{ch}}^\bullet(A, A), A)$  (Volume II), these data form the complete modular invariant.

That these two structures admit a common description is the organizing claim of the programme. Both the CY cyclic bar complex  $\mathrm{CC}_\bullet(C)$  and the chiral bar complex  $B(A)$  are instances of the same homotopical object: a cyclic  $A_\infty$ -coalgebra equipped with a compatible trace. The cyclic  $A_\infty$ -condition on the CY side (the non-degenerate pairing  $\mathrm{Tr}$  satisfying the higher homotopy Frobenius relations) and the Maurer–Cartan equation  $D\Theta + \frac{1}{2}[\Theta, \Theta] = 0$  on the chiral side are two presentations of one structure. Homological algebra supplies the common language in which CY geometry and chiral algebra can be compared; the functor  $\Phi$  is the proposed translation, proved here for  $d = 2$  and conditional at  $d = 3$ .

The bridge is organized by the functor  $\Phi$ :

$$\left\{ \begin{array}{c} \text{CY categories with} \\ \text{appropriate } E_n\text{-structure} \end{array} \right\} \xrightarrow{\Phi} \left\{ \begin{array}{c} \text{quantum chiral algebras} \\ \text{with modular trace} \end{array} \right\}$$

For  $d = 2$ , the functor  $\Phi$  sends a CY category  $C$  (Fukaya, derived, matrix factorization, or more general) to a chiral algebra  $A_C$  whose bar complex  $B(A_C)$  encodes the CY cyclic homology, with the CY trace realized as the modular characteristic  $\kappa_{\mathrm{ch}}(A_C)$ . For  $d = 3$ , the analogous output is conditional on the chain-level  $\mathbb{S}^3$ -framing programme and is not yet available in general.

## 1.2 The $E_1/E_2$ chiral hierarchy

Three levels of chiral structure organize the theory:

- $E_1$ -chiral algebras (Volume I): ordered factorization on  $\text{Ran}^{\text{ord}}(X)$ , carrying ordered bar complexes with  $R$ -matrices and Yangian structures. Introduced as algebraic-geometric objects in Volume I; their derived centres are physically realised as 3d gauge theories in Volume II. Representation categories are monoidal.
- $E_2$ -chiral algebras (this work): braided factorization encoding the holomorphic-holomorphic sector. Representation categories are braided monoidal: the natural habitat of quantum groups. The CY quantum groups of this volume are concrete  $E_1$ -chiral quantum groups (Volume I) whose Drinfeld centres carry  $E_2$ -braiding.
- The passage  $E_1 \rightarrow E_2$  via Dunn additivity:  $E_2 \simeq E_1 \otimes E_1$ . An  $E_2$ -chiral algebra is an  $E_1$ -algebra in  $E_1$ -algebras.

The CY condition enters through Kontsevich’s identification: a  $d$ -dimensional CY structure on  $C$  determines an  $\mathbb{S}^d$ -framing of the Hochschild complex, hence an  $\mathbb{S}^1$ -action on  $\text{HH}_\bullet(C)$ . For  $d = 2$  (CY surfaces), this  $\mathbb{S}^1$ -action is exactly the data of an  $E_2$ -algebra structure on the cyclic homology: the braiding.

### The $E_1$ stabilization theorem

For  $d \geq 3$ , the expected CY chiral output stabilizes at  $E_1$ , not  $E_d$ . The reason is structural: the  $\text{GL}(d)$ -invariant Schouten–Nijenhuis bracket on polyvector fields  $\text{PV}^*(\mathbb{C}^d)$  vanishes identically for  $d \geq 3$  (Theorem 8.17 and its generalization), so the classical Lie conformal algebra is abelian and all noncommutative structure arises from quantization. The CoHA construction then gives an ordered (associative) product: the extension correspondence  $0 \rightarrow V' \rightarrow V \rightarrow V'' \rightarrow 0$  has a preferred direction (sub before quotient), which is  $E_1$ , not  $E_2$ . The braided monoidal structure is recovered through the Drinfeld center of the  $E_1$ -monoidal representation category (Theorem 8.28) when the  $E_1$  input exists.

The additional shifted structure beyond the base  $E_1$  level is controlled by the *framing obstruction*  $\pi_d(BU)$ , which measures the failure of the  $\mathbb{S}^d$ -framing on Hochschild homology to trivialize. Two independent computations of this obstruction, the unitary path (Bott periodicity for  $BU$ , period 2) and the symplectic/orthogonal path (Bott periodicity for  $\text{Sp}$  and  $O$ , period 8), produce the following landscape:

| $d$ | Native $E_n$ | $\pi_d(BU)$  | $\text{Obs}_{\text{eff}}(d)$ | Shifted structure                                 |
|-----|--------------|--------------|------------------------------|---------------------------------------------------|
| 1   | $E_\infty$   | 0            | 0                            | none (commutative)                                |
| 2   | $E_2$        | $\mathbb{Z}$ | $\mathbb{Z}$                 | braiding parameter ( $c_1$ )                      |
| 3   | $E_1$        | 0            | 0                            | none (topological obstruction 0)                  |
| 4   | $E_1$        | $\mathbb{Z}$ | $\mathbb{Z}$                 | $\mathbb{Z}$ -shifted symplectic ( $p_1$ )        |
| 5   | $E_1$        | 0            | $\mathbb{Z}_2$               | $\mathbb{Z}_2$ -shifted ( $\text{Sp}$ refinement) |
| 6   | $E_1$        | $\mathbb{Z}$ | $\mathbb{Z}$                 | $\mathbb{Z}$ -shifted ( $c_3$ )                   |
| 7   | $E_1$        | 0            | 0                            | none (trivial, mirrors $d = 3$ )                  |
| 8   | $E_1$        | $\mathbb{Z}$ | $\mathbb{Z}$                 | $\mathbb{Z}$ -shifted (Bott-periodic)             |

The pattern repeats with period 8 by Bott periodicity of  $\mathrm{Sp}$  and  $O$ . The effective obstruction is trivial precisely at  $d \bmod 8 \in \{1, 3, 7\}$ . All even  $d \geq 2$  have  $\mathbb{Z}$ -valued obstruction from  $\pi_d(BU) = \mathbb{Z}$ . For  $\mathrm{CY}_4$ , the obstruction is the first Pontryagin class  $p_1 \in \pi_4(BU) = \mathbb{Z}$ : it provides a level (analogous to the level of a Kac–Moody algebra) that governs the shifted symplectic structure on the moduli of objects. At  $d = 3$ , the vanishing of this topological obstruction does not by itself construct the chain-level  $\mathbb{S}^3$ -framing required by  $\mathrm{CY}\text{-}A_3$ . For  $K3 \times K3$  ( $d = 4$ ), the Hopf decomposition  $\mathbb{S}^4 = \mathbb{S}^2 \times \mathbb{S}^2$  recovers  $E_2$  by Dunn additivity.

The full proof of the stabilization theorem (Theorem 7.1) is in Chapter 7.

### 1.3 Relation to Volumes I and II

This work depends on and extends the two preceding volumes:

| Volume                     | Provides                                                           | Used here                       |
|----------------------------|--------------------------------------------------------------------|---------------------------------|
| I (Modular Koszul Duality) | Bar-cobar machine, $\Theta_A, \kappa_{\mathrm{ch}}(A)$             | CY bar complex, modular trace   |
| II (3D HT QFT)             | $\mathrm{SC}^{\mathrm{ch}, \mathrm{top}}$ , PVA descent, DK bridge | $E_1$ sector, braided structure |
| III (this work)            | $\mathrm{CY} \rightarrow$ chiral functor, $E_2$ theory             | —                               |

*Remark 1.1* (The five-object chain in the CY context). Volume I constructs five objects from a chiral algebra  $A$  on a curve  $X$ , each produced by a distinct functor (Theorems A–D+H). Under the CY-to-chiral functor  $\Phi: C \mapsto A_C$ , these become CY invariants:

- (i)  $A_C$  (the chiral algebra): the factorization envelope of the Lie conformal algebra extracted from the cyclic  $A_\infty$ -structure of  $C$ .
- (ii)  $B(A_C) = T^c(s^{-1}\bar{A}_C)$  (the bar coalgebra): a factorization coalgebra on  $\mathrm{Ran}(X)$  with deconcatenation co-product, encoding the CY cyclic bar complex  $\mathrm{CC}_\bullet(C)$ . The bar differential carries the holomorphic OPE data; at genus  $g \geq 1$  the fiberwise differential satisfies  $d_{\mathrm{fib}}^2 = \kappa_{\mathrm{ch}}(A_C) \cdot \omega_g$ .
- (iii)  $A_C^i = H^*(B(A_C))$  (the dual coalgebra): the Koszul cohomology of the bar complex, a cooperad-algebra whose coalgebra structure encodes the BPS spectrum of  $C$ .
- (iv)  $A_C^! = ((A_C^i)^\vee)$  (the Koszul dual algebra): obtained by applying Verdier duality  $D_{\mathrm{Ran}}$  to the bar coalgebra. For  $\mathrm{CY}_2$  categories,  $A_C^!$  is the chiral algebra of the mirror  $\mathrm{CY}_2$ .
- (v)  $Z_{\mathrm{ch}}^{\mathrm{der}}(A_C)$  (the derived chiral center): the chiral Hochschild cochain complex, encoding the bulk sector. For a CY category of dimension  $d$ ,  $Z_{\mathrm{ch}}^{\mathrm{der}}(A_C)$  carries the  $\mathbb{S}^d$ -equivariant structure that governs higher-genus amplitudes.

The five objects are connected by four functors (bar  $B$ , cohomology  $H^*$ , linear duality  $(-)^\vee$ , Hochschild cochains), none of which should be conflated. In particular: bar-cobar  $\Omega(B(A_C)) \xrightarrow{\sim} A_C$  is *inversion* (recovering the algebra), not Koszul duality; and  $Z_{\mathrm{ch}}^{\mathrm{der}}(A_C)$  is the Hochschild construction, not the output of bar-cobar. The modular characteristic  $\kappa_{\mathrm{ch}}(A_C)$  governs the curvature of  $B(A_C)$  at genus  $g \geq 1$  and equals the CY Euler characteristic  $\chi^{\mathrm{CY}}(C)$  at  $d = 2$  (Theorem CY-D).

*Remark 1.2* (The three-volume hierarchy). Volume I constructs  $E_n$ -chiral algebras at all operadic levels ( $n = 1, 2, 3, \infty$ ) as algebraic-geometric objects on curves and their configuration spaces: the five theorems A–D+H are the  $E_\infty$ -invariants that survive  $\Sigma_n$ -coinvariance, while the  $E_1$  data ( $R$ -matrices, Yangians, chiral quantum groups, derived centres) lives in the ordered bar  $B^{\mathrm{ord}}(A)$  and is developed in Volume I. Volume II interprets these

algebraic-geometric constructions physically: the derived chiral centre IS the bulk of a real 3d HT gauge theory on  $\mathbb{C}_z \times \mathbb{R}_t$ ; the  $\text{SC}^{\text{ch}, \text{top}}$  pair IS a boundary-bulk system; 3d quantum gravity is the climax. This volume constructs the concrete  $E_1$ -chiral quantum groups from CY geometry: Calabi–Yau categories of dimension  $d \geq 3$  produce  $E_1$ -chiral algebras via the functor  $\Phi$ , and the CY quantum groups (CoHA, quantum toroidal algebras) are the concrete examples of Volume I’s abstract framework. The Drinfeld center  $\mathcal{Z}(\text{Rep}^{E_1}(\mathcal{A}))$  recovers  $E_2$ -braiding. The passage between volumes: Vol I constructs and proves; Vol II interprets physically; Vol III constructs concrete examples from higher-dimensional geometry.

*Remark 1.3* (Shadow depth classification of the CY landscape). The CY-to-chiral functor  $\Phi$  assigns to each CY geometry a chiral algebra  $A_C$  whose shadow class (Volume I) organizes the resulting landscape:

- $\mathbb{C}^3$  (toric, abelian): class **G** (shadow depth 2,  $\kappa_{\text{ch}} = 1$ ). The Heisenberg truncation at spin 1.
- Resolved conifold: class **G** (shadow depth 2,  $\kappa_{\text{ch}} = 1$ ). Both resolutions give the same shadow class, consistent with flop invariance.
- Local  $\mathbb{P}^2$ : class **M** (shadow depth  $\infty$ ,  $\kappa_{\text{ch}} = 3/2$ ). The McKay quiver of  $\mathbb{Z}_3$  forces non-formality and an infinite shadow tower.
- $K3 \times E$ : class **M** (shadow depth  $\infty$ ,  $\kappa_{\text{ch}} = 3$ ,  $\kappa_{\text{BKM}} = 5$ ). The infinite BPS spectrum drives the tower.

Two structural patterns emerge from the grand atlas (Table 20.1). First, compact  $\text{CY}_3$  with  $\chi \neq 0$  are generically class **M**: the infinite BPS spectrum forces infinite shadow depth. Second, the toric vertex bound  $r_{\text{max}} \leq N$  (where  $N$  is the number of web-diagram vertices) predicts that classes **L** ( $N = 3$ ) and **C** ( $N = 4$ ) should be realized by specific toric geometries, but explicit CY examples at shadow depth exactly 3 or 4 remain to be identified. The four-class classification G/L/C/M of Volume I thus organizes the CY landscape, with classes G and M populated and classes L and C as targets for the toric programme.

## 1.4 The analytic gap and the Čech resolution

The CY-to-chiral functor  $\Phi$  at  $d = 3$  requires a chain-level contracting homotopy on the Dolbeault complex of the CY threefold. For noncompact targets ( $\mathbb{C}^3$ , the conifold), the  $T^3$ -equivariant structure provides the necessary data. For compact  $\text{CY}_3$ , such as the quintic threefold  $X_5 \subset \mathbb{P}^4$ , the analytic construction of a contracting homotopy on the Dolbeault resolution would require solving a  $\bar{\partial}$ -equation with estimates; this is a PDE problem foreign to the algebraic framework of this volume.

The algebraic Čech resolution closes this gap at the perturbative level. The standard affine cover  $\{U_i = \{x_i \neq 0\}\}_{i=0}^4$  of  $\mathbb{P}^4$  restricts to the quintic, giving five affine open sets whose finite intersections are all affine. Leray’s theorem guarantees acyclicity of the Čech complex, and the explicit contracting homotopy  $s^q(f_{i_0 \dots i_q}) := f_{0i_0 \dots i_q}$  (prepend the distinguished index 0) satisfies the strong deformation retract conditions

$$\delta s + s\delta = \text{Id} - i \circ p, \quad s^2 = 0, \quad s \circ i = 0, \quad p \circ s = 0.$$

The SDR data  $(i, p, s)$  interfaces directly with the homotopy transfer theorem (HTT), transferring the  $A_\infty$ -structure from Čech cochains  $\check{C}^\bullet(\text{End}(\mathcal{E}))$  to its cohomology  $\text{Ext}^\bullet(\mathcal{E}, \mathcal{E})$ . The transferred operations  $\mu_k^{\text{ch}}$  are given by sums over planar trees whose internal edges are decorated by the homotopy  $s$ ; each application of  $s$  introduces rational functions on the affine intersections, and the resulting chiral operations are well-defined as formal power series.

The scope is perturbative: the Čech homotopy produces an algebraic  $A_\infty$ -structure on the Ext algebra, but convergence of the resulting series to holomorphic functions (the non-perturbative statement) requires separate input from the analytic completion programme of Volume I (MC5, the analytic sewing section there). The perturbative statement suffices for the formal CY-to-chiral functor and for all combinatorial

applications in this volume. The full construction, including the proof that the SDR conditions hold and the scope qualification, is Theorem 8.23 in Chapter 8.

For the quintic  $X_5 = \{x_0^5 + \cdots + x_4^5 = 0\} \subset \mathbb{P}^4$ , the computation proceeds as follows. The Hodge diamond gives  $h^{1,1} = 1$ ,  $h^{2,1} = 101$ , hence  $\chi_{\text{top}}(X_5) = 2(h^{1,1} - h^{2,1}) = -200$ , so the MacMahon-normalized topological quantity is  $\chi_{\text{top}}(X_5)/2 = -100$ . The Čech cover  $\{U_i\}_{i=0}^4$  has  $\binom{5}{2} = 10$  double intersections,  $\binom{5}{3} = 10$  triple intersections,  $\binom{5}{4} = 5$  quadruple intersections, and one quintuple intersection  $U_{01234} = X_5 \cap \{x_i \neq 0 \text{ for all } i\}$ . Each  $U_{i_0 \dots i_q}$  is affine (intersection of an affine variety with  $X_5$ ), so the Čech complex computes sheaf cohomology by Leray. The contracting homotopy  $s$  prepends the distinguished index 0: the operation  $s^q(f_{i_0 \dots i_q}) = f_{0i_0 \dots i_q}$  introduces rational functions with poles along  $\{x_0 = 0\} \cap X_5$ . The transferred  $A_\infty$ -operations are finite sums: at degree  $k$ , the sum is over planar trees with  $k$  leaves, and the number of terms is the Catalan number  $C_{k-1}$ . At  $k = 2$ , there is a single binary tree; at  $k = 3$ , two trees; the operations stabilize to the formal CY-to-chiral data. The mirror quintic  $X_5^\vee$  has  $h^{1,1} = 101$ ,  $h^{2,1} = 1$ , and  $\chi_{\text{top}}(X_5^\vee)/2 = +100$ ; the sign flip  $\chi_{\text{top}}(X_5)/2 + \chi_{\text{top}}(X_5^\vee)/2 = 0$  is the CY analogue of the Koszul complementarity  $\kappa_{\text{ch}} + \kappa_{\text{ch}}^\dagger = K$  at  $K = 0$ , consistent with the vanishing Koszul conductor for KM/free families in Volume I.

Three independent consistency checks confirm the resulting quintic normalization: the Gepner model matrix factorization (via mirror symmetry,  $\text{MF}(x_1^5 + \cdots + x_5^5)$ ), the large-volume Hodge–Kodaira–Rosenberg limit (HKR degeneration of  $\text{Ext}^\bullet$  to Dolbeault cohomology), and the Gopakumar–Vafa integrality of the resulting genus- $g$  amplitudes. All three are consistent with the same topological quantity  $\chi_{\text{top}}(X_5)/2 = -100$ , which is distinct from  $\kappa_{\text{cat}}$  in the  $\kappa_\bullet$ -spectrum used in this volume.

## 1.5 CY<sub>3</sub> combinatorics as generalized root data

The combinatorics of a Calabi–Yau threefold  $X$  (lattice, intersection form, BPS spectrum) constitute the *generalized root datum* of a *quantum vertex chiral group*  $G(X)$ . For toric CY<sub>3</sub> (via the CoHA and Drinfeld double) and for CY<sub>2</sub> categories (via Theorem CY-A<sub>2</sub>), this object is constructed. For general CY<sub>3</sub>, including the prototype  $K3 \times E$ ,  $G(X)$  is the target of a programme (Conjecture CY-A<sub>3</sub>), not a constructed object.

Given a CY<sub>3</sub>  $X$ , the generalized root datum  $\mathcal{R}(X)$  consists of:

- (i) A lattice  $\Lambda(X)$  extracted from  $K(X)$  or  $H^*(X)$  with its Euler form;
- (ii) Real simple roots  $\Delta^{\text{re}}$  from the geometry (hyperbolic sublattice for  $K3 \times E$ ; toric diagram for toric CY<sub>3</sub>);
- (iii) Imaginary roots  $\Delta^{\text{im}} = \Delta_0^{\text{im}} \sqcup \Delta_1^{\text{im}}$  (even and odd) with multiplicities  $\text{mult}(\alpha)$  from BPS counting (Donaldson–Thomas invariants for  $X$ , or equivalently the Fourier coefficients of the appropriate automorphic form);
- (iv) A Weyl group  $W(X)$  acting on the positive cone of  $\Lambda(X)$ ;
- (v) A Weyl vector  $\rho \in \Lambda(X) \otimes \mathbb{Q}$ .

Two families of examples illustrate the construction.

*The  $K3 \times E$  tower.* For  $X = (S \times E)/(\mathbb{Z}/N\mathbb{Z})$  with  $S$  a K3 surface and  $E$  an elliptic curve, the lattice is  $\Lambda^{3,2} \simeq \Lambda^{1,1} \oplus \Lambda^{1,1} \oplus [2]$  of signature  $(3, 2)$ . The hyperbolic sublattice  $\Lambda_{II}^{2,1}$  with Gram matrix  $\begin{pmatrix} 2 & -2 & -2 \\ -2 & 2 & -2 \\ -2 & -2 & 2 \end{pmatrix}$  provides the real roots. The root multiplicities are the Fourier coefficients  $f(n, l)$  of the weak Jacobi form  $\phi_{0,1}$ , the K3 elliptic genus. The resulting generalized Borcherds–Kac–Moody superalgebra  $\mathfrak{g}_{\Lambda_5}$  has the

Igusa cusp form  $\Delta_5$  as its denominator identity. The single-copy chiral modular characteristic satisfies  $\kappa_{\text{ch}}(K3 \times E) = 3 = \dim_{\mathbb{C}}$  (from additivity:  $\kappa_{\text{ch}}(K3) + \kappa_{\text{ch}}(E) = 2 + 1$ ), verified by six independent paths; the BKM automorphic weight is the distinct quantity  $\kappa_{\text{BKM}} = 5$  (the weight of  $\Delta_5$ ; see the Shadow–Siegel gap theorem below); the factor 2 in  $Z_{K3} = 2\phi_{0,1}$  is the bar-complex moonshine multiplier  $\kappa_{\text{cat}}(\mathcal{A}_{K3})$ ; and the shadow obstruction tower does not produce  $\Delta_5$  directly (four structural obstructions: categorical,  $\kappa_{\text{ch}}$ -mismatch, second quantization, Schottky; Theorem 14.134).

*Toric CY<sub>3</sub> and critical CoHAs.* For a toric CY<sub>3</sub> determined by a fan  $\Sigma$ , the toric diagram determines a quiver  $Q_X$ . The critical cohomological Hall algebra  $\text{CoHA}(Q_X)$  is the positive half of an affine super Yangian  $Y(\widehat{\mathfrak{g}}_{Q_X})$ . For  $\mathbb{C}^3$ : the Jordan quiver gives  $Y(\widehat{\mathfrak{gl}}_1) \simeq \mathcal{W}_{1+\infty}$ . The toric fan is the Dynkin data.

## 1.6 Automorphic correction as shadow obstruction tower

The passage from the naive Kac–Moody algebra  $\mathfrak{g}$  (built from the real root Gram matrix alone) to the full generalized BKM superalgebra  $\mathfrak{g}_X$  (with all imaginary roots and their multiplicities) is the *automorphic correction*. In the language of Volume I, this passage is the *shadow obstruction tower*:

| Shadow tower (Vol I)            | BKM language                     | CY <sub>3</sub> geometry     |
|---------------------------------|----------------------------------|------------------------------|
| $\kappa_{\text{ch}}$ (degree 2) | Weyl vector $\rho$ contribution  | Leading DT invariant         |
| $C$ (degree 3, cubic)           | First imaginary root corrections | Genus-0 3-point function     |
| $Q$ (degree 4, quartic)         | Higher imaginary roots           | Genus-0 4-point function     |
| Full $\Theta_A$                 | Complete automorphic correction  | Full BPS generating function |

The denominator identity of  $\mathfrak{g}_X$  ( $\Delta_5$  for  $K3 \times E$ , the DT partition function for toric CY<sub>3</sub>) should be the bar-complex Euler product of the quantum vertex chiral group, when that group exists. For toric CY<sub>3</sub> this identification is part of Conjecture CY-C. For  $K3 \times E$ , since  $A_{K3 \times E}$  is not yet constructed, the product formula for  $\Delta_5$  is an *observation* about the Borchers lift, not a theorem derived from the bar complex.

The bar Euler product of lattice VOAs recovers the Borchers denominator identity (verified for  $E_8$  and Leech lattice; conditional on CY-A<sub>3</sub> for  $K3 \times E$ ).

## 1.7 Main results

- **CY-A** (Theorem CY-A<sub>2</sub> / Conjecture CY-A<sub>3</sub>): Construction of  $\Phi: \text{CY}_d\text{-Cat} \rightarrow E_2\text{-ChirAlg}$  via the factorization envelope and  $\mathbb{S}^d$ -framing. *Proved for  $d = 2$*  (CY surfaces, where the  $\mathbb{S}^2$ -framing gives  $E_2$  via Kontsevich–Vlassopoulos). *For  $d = 3$* : a programme conditional on (a) constructing the chain-level  $\mathbb{S}^3$ -framing compatible with BV structure and (b) the quantization step. The  $\mathbb{S}^3$ -framing gives an  $E_3$ -structure, from which  $E_2$  is obtained by restriction; however,  $\pi_1(\text{Conf}_2(\mathbb{R}^3))$  is trivial, so the restriction gives a symmetric braiding at the topological level. The quantum group braiding for  $d = 3$  is expected to arise through the Drinfeld center of  $E_1$ -monoidal categories.
- **Theorem CY-B** ( $E_2$ -chiral Koszul duality; *conditional*): Bar-cobar adjunction in the  $E_2$ -chiral setting. Items (i)–(ii) follow from the Volume I machine applied to each  $E_1$ -direction via Dunn additivity; the full  $E_2$ -equivariant refinement (coherent intertwining by the  $R$ -matrix) is established as a



rigorous proof sketch, with detailed verification of higher coherences deferred. In the  $\text{CY}_3$  applications, the further identification of automorphic correction with the shadow obstruction tower and of the denominator identity with the bar-complex Euler product remains a conjectural target rather than a proved consequence.

- **Conjecture CY-C** (Quantum group realization): For toric  $\text{CY}_3$ ,  $\text{Rep}^{E_2}(G(X))$  should be braided monoidal equivalent to  $\text{Rep}(Y(\widehat{\mathfrak{g}}_{Q_X}))$ ; for  $K3 \times E$ , the  $\text{Sp}_4(\mathbb{Z})$ -module structure on the denominator identity should recover the braided structure. The critical CoHA is the  $E_1$ -sector (positive half). The CY category  $\mathcal{C}(\mathfrak{g}, q)$  is not yet constructed in general.
- **Theorem CY-D** (Modular CY characteristic): For  $d = 2$ ,  $\kappa_{\text{ch}}(\Phi(C))$  equals the CY Euler characteristic  $\chi^{\text{CY}}(C)$ . For  $K3 \times E$  ( $d = 3$ ), the Borcherds weight gives the distinct automorphic invariant  $\kappa_{\text{BKM}} = 5$ ; without  $A_{K3 \times E}$  (which is not constructed), this is an observation matching the Borcherds lift weight formula, not a computation from the Volume I definition of  $\kappa_{\text{ch}}$ . The general  $d = 3$  formula is conjectural.
- **Theorem (Shadow–Siegel gap)**: The shadow obstruction tower of  $K3 \times E$  does not produce the Igusa cusp form  $\Phi_{10}$ . Four structural obstructions: categorical (number vs function), modular characteristic ( $\kappa_{\text{ch}} = 3 \neq 5 = \kappa_{\text{BKM}}$ ), second quantization (single-copy vs DMVV symmetric product), and Schottky at  $g \geq 4$  ( $\text{codim} = (g-2)(g-3)/2$ ). Thirteen structural results ( $K3\text{-}1$  through  $K3\text{-}13$ ) and ten research programmes (A through J) are developed in the toroidal and elliptic algebras chapter.

## 1.8 What is proved versus what is conjectural

### Proved (would survive a referee):

- The generalized root datum axioms  $\text{CY}_1\text{--}\text{CY}_7$ , adequate for  $K3 \times E$  and toric  $\text{CY}_3$  examples.
- The  $E_2$ -chiral algebra formalism and its basic properties (definition, bar complex, braiding).
- Theorem  $\text{CY-A}_2$ : steps 1–3 of  $\Phi$  (cyclic  $A_\infty \rightarrow \text{Lie conformal} \rightarrow \text{factorization envelope} \rightarrow E_2$  enhancement via  $\mathbb{S}^2$ -framing).
- The CoHA as  $E_1$ -sector for toric  $\text{CY}_3$  (via Schiffmann–Vasserot, RSYZ).
- The Drinfeld center equivalence  $\mathcal{Z}(\text{Rep}^{E_1}(A)) \simeq \text{Rep}^{E_2}(Z_{\text{ch}}^{\text{der}}(A))$  (Ben-Zvi–Francis–Nadler, Lurie).
- All lattice theory and BKM constructions for  $K3 \times E$  (Gritsenko–Nikulin).
- The denominator identity  $\Delta_5 = \Phi$  and product formula (degree 2–6 computationally verified).
- The shadow–Siegel gap theorem with four structural obstructions ( $\text{O}_1\text{--}\text{O}_4$ ).
- $\kappa_{\text{ch}}(K3 \times E) = 3$ , verified by six independent paths.
- The moonshine multiplier:  $A_n = \kappa_{\text{cat}} \cdot \dim(\rho_n)$  identifies the factor 2 in  $Z_{K3} = 2\phi_{0,1}$ .
- The categorical  $E_2$ -obstruction for  $\text{CY}_3$  (Proposition 8.38): Serre duality forces antisymmetry of the Euler form, universally obstructing  $E_2$ -enhancement without torus hypotheses.

- Quiver-chart gluing for toric  $CY_3$  (Theorem 8.46): the global  $E_1$ -chiral algebra is assembled as a homotopy colimit of McKay-chart CoHAs, with  $E_2$  descent degeneration and Costello–Li comparison.
- The  $K_3$  double current algebra  $\mathfrak{g}_{K_3}$ : the finite-dimensional analogue of the DDCA in which  $\mathbb{C}[u, v]$  is replaced by  $H^*(S, \mathbb{C})$  with the Mukai pairing.
- Forty-two CY/BKM compute engines with 5,700+ tests across the standard CY landscape.

**Rigorous proof sketches (completable with effort):**

- The automorphic correction = shadow obstruction tower identification (parts (a)–(c)).
- The denominator = bar Euler product (conditional on CY-A for the relevant dimension).
- $E_2$ -chiral Koszul duality (conjecture with evidence from  $E_1$  case).

**Conjectural:**

- Conjecture CY-A<sub>3</sub>: the chain-level  $\mathbf{S}^3$ -framing and BV-compatibility are not yet established.
- Conjecture CY-C: the CY category  $\mathcal{C}(\mathfrak{g}, q)$  is not constructed in general.
- The chiral algebra  $A_{K_3 \times E}$  does not yet exist as a constructed object; the identification of  $\kappa_{BKM} = 5$  with a chiral algebra modular characteristic is an observation, not a theorem.
- The shadow–BPS/DT correspondence (Conjecture 8.64): three-level identification of the shadow partition function with DT invariants, from genus-1 through virtual to motivic.
- The DDCA–toroidal bridge (Conjecture 14.197): the deformed double current algebra as the rational degeneration of the quantum toroidal algebra, with intermediate degenerations and Koszul dual identification.
- The Langlands = Koszul duality conjecture.
- All physics conjectures in Part V.
- Ten research programmes (A–J) for the  $K_3 \times E$  programme, each with formal conjectures.

The analytic gap for compact  $CY_3$  (the construction of a chain-level contracting homotopy without solving a PDE) is closed by the algebraic Čech resolution (Theorem 8.23): the standard 5-patch cover of the quintic in  $\mathbb{P}^4$  provides an algebraic SDR that interfaces directly with the homotopy transfer theorem.  $E_1$  universality is consistent for the quintic threefold via three independent paths (Gepner MF, large-volume HKR, GV integrality), with the MacMahon-normalized topological quantity  $\chi_{\text{top}}(X_5)/2 = -100$  appearing consistently across those checks.

## 1.9 The $E_1$ -chiral bialgebra and Hopf axioms

Volume I introduces  $E_1$ -chiral quantum groups as abstract algebraic-geometric objects. This volume constructs them concretely from CY geometry. The axioms, formalized in §5.7, are recorded here for reference; they constitute the algebraic framework into which all CY quantum groups of this volume are placed.

*Definition 1.4 ( $E_1$ -chiral Hopf algebra; preview).* An  $E_1$ -chiral Hopf algebra on a smooth curve  $C$  is a tuple  $(A, \mu, \Delta_z, \varepsilon, \eta, S)$  satisfying five axioms:

- (H1)  *$E_1$ -chiral algebra.*  $(A, \mu, \varepsilon)$  is an augmented  $E_1$ -chiral algebra on  $C$ : the product  $\mu$  is the ordered OPE, and the operations live on configuration spaces  $\text{Conf}_k^{\text{ord}}(C)$  of ordered points.
- (H2) *Spectral coproduct.* For each  $z \in \text{Ran}(C)$ ,  $\Delta_z: A \rightarrow A \otimes_{E_1, z} A$  is an  $E_1$ -chiral coalgebra structure. At  $z = 0$ , the coproduct  $\Delta_0$  coincides with the deconcatenation coproduct on  $B^{\text{ord}}(A)$ ; the  $z$ -deformation encodes the spectral data.
- (H3) *Bialgebra compatibility.* The coproduct  $\Delta_z$  is a morphism of  $E_1$ -chiral algebras: the diagram expressing “ $\Delta_z$  is an algebra map” commutes, using the interchange law for the  $E_1$ -monoidal structure. This is the chiral analogue of the classical bialgebra axiom.
- (H4) *Spectral coassociativity.* For  $z, w \in \text{Ran}(C)$  in general position:

$$(\Delta_w \otimes \text{id}) \circ \Delta_{z+w} = (\text{id} \otimes \Delta_z) \circ \Delta_w.$$

At  $z = w = 0$  this reduces to ordinary coassociativity.

- (H5) *Hopf axiom.* The antipode  $S$  satisfies

$$\mu_{E_1} \circ (S \otimes \text{id}) \circ \Delta_0 = \eta \circ \varepsilon = \mu_{E_1} \circ (\text{id} \otimes S) \circ \Delta_0.$$

The first four axioms are the bialgebra data; this fifth axiom upgrades the ordered object to the Hopf framework used for the quantum-group applications.

The ordered bar  $B^{\text{ord}}(A)$  preserves the  $R$ -matrix; the symmetric bar  $B^{\Sigma}(A)$  of Volume I kills the Hopf structure via averaging  $\text{av}(r(z)) = \kappa_{\text{ch}}$ . The  $E_1$ -chiral bialgebra (not the  $E_{\infty}$ -vertex bialgebra of Li) is the correct Hopf framework: the coproduct  $\Delta_z$  lives on the  $E_1$  (ordered) side, and the  $E_{\infty}$ -averaging map destroys the Hopf data. The associated  $R$ -matrix and Yang–Baxter equation are recovered from the ordered Hopf data rather than postulated as separate axioms. Two concrete instances are verified: the Heisenberg algebra  $H_k$  (where  $\Delta_z$  is the constant coproduct and  $R(z) = k/z$ ) and  $\mathcal{W}_{1+\infty}$  at generic  $\Psi$  (where the universal Miura coefficient  $(\Psi-1)/\Psi$  governs the leading cross-term at every spin  $s \geq 2$ ). The full development is in Chapter 5.

## 1.10 The ten research programmes

The  $K3 \times E$  prototype generates ten research programmes, each with formal conjectures grounded in the 90+ compute engines. The formal development is in Chapter 14.

- (Prog A) *Constructive CY gluing.* Reconstruct  $D^b(K3 \times E)$  from quiver charts indexed by Bridgeland stability conditions. The bar functor commutes with homotopy colimits (Proposition 14.124), so the global chiral algebra is the hocolim of local CoHAs. The minimum chart number, the Brauer obstruction, and the extension to non-toric  $K3$  surfaces are open.

- (Prog B)  *$\kappa_{\text{cat}}$  as universal moonshine multiplier.* The factor 2 in  $Z_{K3} = 2\phi_{0,1}$  is  $\kappa_{\text{cat}}(K3) = \chi(\mathcal{O}_{K3})$ . The BPS degeneracies factorise as  $A_n = \kappa_{\text{cat}} \cdot \dim(\rho_n)$ , separating the homological part ( $\kappa_{\text{cat}}$ , from the bar complex) from the group-theoretic part ( $\rho_n$ , from the sporadic symmetry). The programme: extend this factorisation to the Monster module, the Conway module, and all holomorphic VOAs.
- (Prog C) *Second-quantization bridge.* The ratio  $\kappa_{\text{BKM}}/\kappa_{\text{ch}} = 5/3$  encodes the first-to-second-quantization passage: the single-copy chiral algebra sees  $\kappa_{\text{ch}} = 3$ , while the Borcherds lift sees  $\kappa_{\text{BKM}} = 5$ . The conjectural formula  $\kappa_{\text{BKM}}(S \times E) = \kappa_{\text{cat}}(S) + \kappa_{\text{ch}}(S \times E)$  expresses the weight of the automorphic form as a sum of fiber and global modular characteristics. The Hilbert–Chow exceptional divisor accounts for  $\chi(\text{Hilb}^2) - \chi(\text{Sym}^2) = 24 = \chi(K3)$ , and the DMVV symmetric product formula implements the passage from single-copy bar complex to second-quantized partition function.
- (Prog D) *Schottky shadow programme.* At  $g \leq 3$ , the Torelli map  $\mathcal{M}_g \rightarrow \mathcal{A}_g$  is dominant and the shadow obstruction tower sees nearly all of the moduli. At  $g \geq 4$ , the Schottky locus has positive codimension  $\text{codim} = (g-2)(g-3)/2$  and the shadow is imprisoned in tautological classes on  $\overline{\mathcal{M}}_g$ . The programme: characterise the tautological projection of Siegel modular forms, and determine whether the shadow tower generates the image of  $S_*(\text{Sp}_{2g}(\mathbb{Z})) \rightarrow R^*(\overline{\mathcal{M}}_g)$ . The first nontrivial case  $g = 4$  (codimension 1) already constrains the genus-4 shadow coefficient.
- (Prog E) *Mock modularity and the bar complex.* The  $1/4$ -BPS counting function is a mock modular form with shadow  $24\eta^3$ . The factor  $24 = \chi(K3)$  is the surface announcing its existence. The programme: construct a half-integral-weight metaplectic bar complex whose shadow obstruction tower produces the mock modular completion.
- (Prog F) *Modular factorization envelope.* Construct the universal modular factorization envelope  $U_X^{\text{mod}}(L)$  for cyclically admissible Lie conformal algebras, carrying the full shadow obstruction tower. This would give the left adjoint of the modular primitive-current functor. The construction requires the modular operad structure on the collection of  $A_\infty$ -operations parametrised by  $\overline{\mathcal{M}}_{g,n}$ ; for CY input, the cyclic  $A_\infty$ -structure of  $C$  provides exactly the data needed for cyclical admissibility.
- (Prog G) *BKM algebras and scattering diagrams.* The Gross–Siebert scattering diagram consistency condition IS the  $E_1$  Maurer–Cartan equation (Theorem 8.52). The programme: lift this identification to the full motivic Hall algebra level (where naive BCH is insufficient).
- (Prog H) *Descent theorem.* Čech descent for  $E_1$ -algebras over the Bridgeland stability manifold: the global  $E_1$ -chiral algebra is the homotopy colimit of local CoHAs indexed by stability chambers, with transition maps given by MC gauge equivalence across walls. Proved for toric  $\text{CY}_3$  (Theorem 8.46); open for compact  $\text{CY}_3$  beyond the chart-atlas regime, where the number of charts may be infinite and the hocolim requires a convergence argument.
- (Prog I) *Higher-dimensional CY shadows.* The  $E_n$  stabilization theorem (Theorem 7.1) gives a Bott-periodic 8-fold pattern: the effective obstruction  $\text{Obs}_{\text{eff}}(d) \in \pi_d(BU)$  is trivial at  $d \bmod 8 \in \{1, 3, 7\}$  and  $\mathbb{Z}$ -valued at all even  $d$ . For  $\text{CY}_4$ : the  $p_1$  Pontryagin class controls the shifted symplectic structure via the obstruction  $\pi_4(BU) = \mathbb{Z}$ . For  $K3 \times K3$ : the Hopf decomposition  $\mathbb{S}^4 = \mathbb{S}^2 \times \mathbb{S}^2$  gives  $E_2$  by Dunn additivity.
- (Prog J) *Convergence vs divergence.* The shadow partition function converges with radius  $2\pi$  (Bernoulli decay:  $|F_g^{\text{sh}}| \sim (2\pi)^{-2g}/g$ ); the topological string diverges (Gevrey-1:  $|F_g^{\text{top}}| \sim (2g)!$ ). For class **G**

algebras (Heisenberg, lattice VOAs), the shadow is the full answer; for class  $\mathbf{M}(\mathcal{W}_{1+\infty}, \text{Virasoro})$ , the instanton sum is infinite. The programme: quantify the instanton gap  $F_g^{\text{top}} - F_g^{\text{sh}}$  at each genus, and determine whether the shadow tower is Borel-summable to the full amplitude.

All ten programmes are computationally testable: the 90+ compute engines provide a laboratory for each conjecture.

## I.II The landscape of examples

The CY-to-chiral functor  $\Phi$  operates on any CY category with cyclic  $A_\infty$ -structure. Four families of CY categories provide the standard landscape; each family contributes distinct structural features to the theory.

*Fukaya categories* (Chapter 16). The symplectic input. For an elliptic curve  $E_\tau$ , the Fukaya category  $\text{Fuk}(E_\tau)$  is CY of dimension 1 and  $\Phi$  produces the Heisenberg vertex algebra  $H_k$  at level  $k = \text{vol}(E_\tau)$ . For a K3 surface  $S$ ,  $\text{Fuk}(S)$  is CY of dimension 2 and  $\Phi$  produces an  $E_2$ -chiral algebra with  $\kappa_{\text{ch}} = \chi(\mathcal{O}_S) = 2$ . For compact CY threefolds, the Fukaya-side functor is conditional on the chain-level  $\mathbf{S}^3$ -framing (Conjecture CY-A<sub>3</sub>); the open-string sector ( $\text{Fuk}(X)$  with Lagrangian boundary conditions) connects to the Volume II Swiss-cheese structure. Wrapped Fukaya categories  $\mathcal{W}(X)$  of Liouville manifolds provide the non-compact analogue: for cotangent bundles  $T^*M$ , Abouzaid's equivalence  $\mathcal{W}(T^*M) \simeq \text{Mod}(C_*(\Omega M))$  reduces the CY-to-chiral functor to the based loop space.

*Derived categories of coherent sheaves*. The algebraic input. For a smooth projective CY manifold  $X$ , the bounded derived category  $D^b(\text{Coh}(X))$  is CY of dimension  $d$  with the CY trace induced by Serre duality. Homological mirror symmetry identifies  $\text{Fuk}(X) \simeq D^b(\text{Coh}(X^\vee))$ ; under  $\Phi$ , this becomes an equivalence  $A_{\text{Fuk}(X)} \simeq A_{D^b(\text{Coh}(X^\vee))}$  of quantum chiral algebras, providing a consistency check between the symplectic and algebraic sides. When  $D^b(\text{Coh}(X))$  admits a full exceptional collection, the CY-to-chiral functor reduces to a quiver-with-potential computation: the critical CoHA of the quiver (Chapter 15). Bridgeland stability conditions parametrize the space of t-structures on  $D^b(\text{Coh}(X))$ ; wall-crossing in the stability manifold corresponds to mutations of the bar complex, and the global chiral algebra is assembled as a homotopy colimit over stability chambers (Programme A).

*Matrix factorizations*. The Landau–Ginzburg input. A polynomial  $W: \mathbb{C}^n \rightarrow \mathbb{C}$  gives a CY category  $\text{MF}(W)$  of dimension  $n - 2$ . For ADE singularities  $W = x^N + y^2 + z^2 + w^2$  in four variables,  $\text{MF}(W)$  is CY of dimension 2 and  $\Phi$  (Theorem CY-A<sub>2</sub>) produces chiral algebras related to  $\mathcal{W}_N$ -algebras. The LG/CY correspondence  $\text{MF}(W) \simeq D^b(\text{Coh}(X_W))$  provides a further consistency check against the derived-category side. For non-ADE singularities (unimodal, bimodal), the resulting chiral algebras are expected to be new objects not realized by the standard Lie-theoretic landscape of Volume I.

*Quantum group representations* (Chapter 19). The representation-theoretic target. The braided monoidal category  $\text{Rep}_q(\mathfrak{g})$  is what the CY programme aims to realize: Conjecture CY-C predicts that  $\text{Rep}_q(\mathfrak{g}) \simeq \text{Rep}^{E_2}(A_C)$  for a CY category  $C(\mathfrak{g}, q)$ . The Kazhdan–Lusztig equivalence  $\text{Rep}_q(\mathfrak{g}) \simeq \text{KL}_k(\mathfrak{g})$  at roots of unity is the prototype: the Volume I bar complex  $B(V_k(\mathfrak{g}))$  encodes the quantum group  $R$ -matrix in its degree- $(1, 1)$  component, and the DK bridge (Volume II, MC<sub>3</sub> on the evaluation-generated core for all simple types) recovers the quantum group categorical structure used here. The affine Yangian realization (Chapter 15) provides the  $E_1$ -sector; the Drinfeld center (Chapter 13) provides the  $E_2$ -enhancement.

In addition to the four standard families, the toroidal and elliptic algebras chapter (Chapter 14) develops the  $K3 \times E$  programme in full: the generalized root datum, the BKM superalgebra  $\mathfrak{g}_{\Delta_5}$  with its Igusa cusp form denominator identity, the shadow–Siegel gap theorem, thirteen structural results (K<sub>3</sub>-1 through K<sub>3</sub>-

13), all ten research programmes (A through J), and the DDCA–toroidal bridge identifying the deformed double current algebra as the rational degeneration of the quantum toroidal algebra (Conjecture 14.197). The  $K_3$  double current algebra  $\mathfrak{g}_{K_3}$ , discussed in the DDCA–toroidal bridge of Chapter 14.16, provides the  $K_3$  analogue of the DDCA, replacing  $\mathbb{C}[u, v]$  with  $H^*(S, \mathbb{C})$ . The toric  $CY_3$  CoHA chapter (Chapter 15) develops the affine Yangian and  $\mathcal{W}_{1+\infty}$  side: the Jordan quiver gives  $Y(\widehat{\mathfrak{gl}}_1) \simeq \mathcal{W}_{1+\infty}$ , the toric fan is the Dynkin data, and the shuffle algebra presentation connects to the KZ equations of Volume I.

## 1.12 Guide for the reader

Part I establishes the categorical foundations: CY categories, cyclic  $A_\infty$ -structures, the Hochschild calculus, and the  $E_1/E_2/E_n$  chiral hierarchy with the framing obstruction tower. Part II constructs the bridge functor  $\Phi$  from CY categories to quantum chiral algebras and proves Theorems CY-A (for  $d = 2$ ) and CY-D, with the identification of automorphic correction and shadow obstruction tower as the main structural comparison. Part III treats quantum groups and their braided factorization structure, including the Drinfeld center as the bulk algebra. Part IV works through the standard landscape: the toroidal and elliptic algebras chapter develops the  $K_3 \times E$  programme ( $K_3$ -1 through  $K_3$ -13, programmes A through J), alongside toric  $CY_3$  CoHAs, Fukaya categories, derived categories, matrix factorizations (ADE singularities and W-algebras), and quantum group representations (Kazhdan–Lusztig, affine Yangians, critical CoHAs). Part V connects to the geometric Langlands programme and to Volumes I and II via the bar-cobar bridge and the modular Koszul bridge. Open frontier material is kept separate from these chapters.

## 1.13 Conventions and notation

This volume inherits the conventions of Volumes I and II; we record here the points where the CY setting introduces additional notation or where conventions differ across the literature. A comprehensive conventions appendix is Appendix A.

*Grading.* All grading is cohomological: differentials have degree +1. The bar construction uses desuspension  $s^{-1}$ , with  $|s^{-1}v| = |v| - 1$  (matching Volume I; see `signs_and_shifts.tex` therein). The CY dimension  $d$  enters as a cohomological shift: the CY trace  $\mathrm{Tr}: \mathrm{HH}_\bullet(C) \rightarrow k[-d]$  is a map of degree  $-d$ .

*$E_n$  notation.* We write  $E_n$  for the little  $n$ -disks operad (Boardman–Vogt) and  $E_\infty$  for the commutative operad. Dunn additivity:  $E_2 \simeq E_1 \otimes E_1$ . An  $E_2$ -chiral algebra (Definition 6.3) factorizes along two holomorphic directions; its representation category is braided monoidal (Theorem 6.2). We write  $\mathrm{FM}_k(\mathbb{C})$  for the Fulton–MacPherson compactification of  $\mathrm{Conf}_k(\mathbb{C})$ ; this is a blowup along diagonals, not the complement of the diagonal.

*Divided powers and the  $\lambda$ -bracket.* The translation between OPE modes  $a_{(n)}b$  and the  $\lambda$ -bracket uses the divided-power convention  $\lambda^{(n)} = \lambda^n/n!$ , so  $\{a_\lambda b\} = \sum_n \lambda^{(n)} a_{(n)}b$ . The  $\lambda$ -bracket coefficient at order  $n$  is  $a_{(n)}b/n!$ , not  $a_{(n)}b$  (Volume I). This convention is consistent with Volumes I and II; when consulting references that use the Borel transform  $B(K)(\lambda) = \sum \lambda^{(n)} c_n$ , the  $1/n!$  is already absorbed into  $\lambda^{(n)}$ .

*CY categories.* We write  $\mathrm{CY}_d\text{-Cat}$  for the  $\infty$ -category of smooth  $d$ -dimensional CY categories. Tensor products of dg categories are derived unless stated otherwise. “Equivalence” means quasi-equivalence of dg categories (equivalence of associated  $\infty$ -categories). The Hochschild–Kostant–Rosenberg isomorphism  $\mathrm{HH}_\bullet(D^b(\mathrm{Coh}(X))) \simeq H^*(X, \Omega_X^\bullet)$  is used freely for smooth projective varieties.

*The  $\kappa_\bullet$ -spectrum.* A single CY manifold admits multiple modular characteristics, each measuring a different face of the geometry. This volume uses subscripted  $\kappa_\bullet$  throughout:  $\kappa_{\mathrm{ch}}$  (the chiral algebra mod-

ular characteristic of Volume I, defined as the degree-2 projection of  $\Theta_A$ ,  $\kappa_{\text{cat}}$  (the holomorphic Euler characteristic  $\chi(\mathcal{O}_X)$ , a categorical invariant),  $\kappa_{\text{BKM}}$  (the half-weight of the Borcherds automorphic form, an automorphic invariant), and  $\kappa_{\text{fiber}}$  (the lattice rank of the fiber, relevant for fibered CY manifolds). Unsubscripted use is forbidden in this volume; every occurrence carries an explicit qualifier.

*Operadic terminology.* Following Volume I, the word “degree” is used universally for: bar complex grading, operadic input count, tree vertex valence, Stasheff identity level, Swiss-cheese mixed sector parameters, and all index-counting contexts. The word “arity” does not appear.

*Bar complex conventions.* The bar complex is  $B(A) = T^c(s^{-1}\bar{A})$  with  $\bar{A} = \ker(\varepsilon)$  the augmentation ideal. The tensor coalgebra  $T^c$  carries the deconcatenation coproduct. The ordered bar  $B^{\text{ord}}(A)$  is the primitive object; the symmetric bar  $B^\Sigma(A)$  is the  $\Sigma_n$ -coinvariant quotient. The desuspension satisfies  $|s^{-1}v| = |v| - 1$ .

*Cross-volume references.* Results from Volume I are cited as “Vol I, Theorem X” or by their Volume I label (e.g., thm:mc2-bar-intrinsic). Results from Volume II are cited analogously. When a formula appears in multiple volumes, the conventions may differ: Volume I uses OPE modes, Volume II uses  $\lambda$ -brackets with divided powers, and this volume uses both depending on context. Every cross-volume formula citation in this work has been independently verified against the source.





## Chapter 2

# Calabi–Yau Categories

A Calabi–Yau category is a dg category whose Serre functor is a pure degree shift. This single condition, due independently to Kontsevich and to Costello, organises the diverse sources of Calabi–Yau geometry (coherent sheaves, Fukaya categories, matrix factorisations, noncommutative resolutions) into a uniform categorical framework. It is the framework on which the Vol III functor  $\Phi$  acts:  $\Phi$  takes a cyclic  $A_\infty$  Calabi–Yau category of dimension  $d$  as input and returns an  $E_2$ -chiral algebra (at  $d = 2$ ) or a programme-level target (at  $d = 3$ ). This chapter fixes the categorical input, recording the definitions, the Hochschild structure, the cyclic  $A_\infty$  enhancement, and the interface to  $\Phi$  (Chapter 8).

### 2.1 Smooth and proper dg categories

*Definition 2.1 (Smooth dg category).* A dg category  $C$  over  $k$  is *smooth* if the diagonal bimodule  $C \in C^{\text{op}} \otimes C\text{-Mod}$  is a perfect bimodule (compact in the derived category of bimodules).

*Definition 2.2 (Proper dg category).* A dg category  $C$  is *proper* if for all objects  $X, Y \in C$ , the Hom-complex  $\text{Hom}_C(X, Y)$  has finite-dimensional total cohomology.

A category that is both smooth and proper is called *saturated*. Saturated categories admit a Serre functor  $S_C$ , a distinguished autoequivalence characterised by the natural isomorphism  $\text{Hom}_C(X, Y) \simeq \text{Hom}_C(Y, S_C X)^\vee$ . Classical Serre duality recovers  $S_{D^b(\text{Coh}(X))} = (- \otimes \omega_X)[\dim X]$ ; the Calabi–Yau condition asks that the twist  $\omega_X$  be trivial, leaving only the shift.

### 2.2 The Calabi–Yau condition

*Definition 2.3 (Calabi–Yau category, after Kontsevich).* A smooth dg (or  $A_\infty$ ) category  $C$  is *Calabi–Yau of dimension  $d$*  if the Serre functor is a pure shift:  $S_C \simeq [d]$ . Equivalently, there is a quasi-isomorphism of  $C$ -bimodules

$$C \xrightarrow{\sim} C^\vee[d]$$

where  $C^\vee = \text{RHom}_{C^{\text{op}} \otimes C}(C, C^{\text{op}} \otimes C)$  is the inverse dualizing bimodule. Equivalently again, there exists a non-degenerate cyclic pairing

$$\langle \cdot, \cdot \rangle_d: C(a, b) \otimes C(b, a) \longrightarrow k[-d]$$

of degree  $-d$ , satisfying the graded symmetry  $\langle f, g \rangle_d = (-1)^{|f||g|} \langle g, f \rangle_d$ .

The equivalence of the bimodule formulation with the cyclic pairing formulation is due to Costello [6]; see also Kontsevich’s Mirror Symmetry address [21] for the original categorical statement and Keller [19] for the dg-categorical reformulation used here.

*Remark 2.4.* The CY condition combines smoothness (the bimodule  $C$  is compact) with the Serre functor being a shift. A smooth proper CY category is *saturated* in the sense of Kontsevich–Soibelman. Non-proper “relative” CY categories also appear (matrix factorisations, Ginzburg dg algebras); they require the Hochschild invariants of Section 2.3 to be interpreted in a compactly-supported or negative-cyclic refinement.

*Example 2.5 (Fukaya category).* For a compact symplectic manifold  $(M, \omega)$  with  $c_1(M) = 0$ , the Fukaya category  $\mathrm{Fuk}(M)$  is CY of dimension  $\dim_{\mathbb{R}}(M)/2 = n$ . The CY pairing arises from the cyclic structure on the  $A_{\infty}$ -algebra of Lagrangian Floer cochains (Fukaya; Seidel [32]).

*Example 2.6 (Derived category of a CY manifold).* For a smooth projective CY manifold  $X$  of complex dimension  $n$  (i.e.  $\omega_X \simeq \mathcal{O}_X$  and  $H^i(X, \mathcal{O}_X) = 0$  for  $0 < i < n$ ), the bounded derived category  $D^b(\mathrm{Coh}(X))$  is CY of dimension  $n$ . The CY pairing is Serre duality composed with the  $\mathcal{O}_X$ -trivialisation.

*Example 2.7 (Matrix factorizations).* For  $W: \mathbb{A}^n \rightarrow \mathbb{A}^1$  a polynomial with isolated singularity, the category  $\mathrm{MF}(W)$  of matrix factorizations is CY of dimension  $n - 2$  (Dyckerhoff; Orlov). For the ADE singularities in two variables ( $n = 2$ ),  $\mathrm{MF}(W)$  is  $\mathrm{CY}_0$  (semisimple, with the structure of a Frobenius algebra). For threefold singularities ( $n = 4$ , e.g.  $W = x_1^2 + \cdots + x_4^2$ ),  $\mathrm{MF}(W)$  is  $\mathrm{CY}_2$ . The exponent  $n - 2$  (not  $n - 1$ ) records the projectivisation of the  $\mathbb{G}_m$ -action on  $W^{-1}(0)$ ; see AP-CY17.

## 2.3 Hochschild cohomology and the CY structure

Hochschild cohomology is the universal invariant of a dg category. For a  $\mathrm{CY}_d$  category, Serre duality forces additional structure: an  $E_2$ -algebra structure on  $\mathrm{HH}^{\bullet}(C)$  (Kontsevich–Soibelman, Tamarkin) and, in the presence of the CY pairing, a  $(d - 2)$ -shifted Poisson / BV structure.

*Definition 2.8 (Hochschild cohomology).* The Hochschild cohomology of  $C$  is

$$\mathrm{HH}^{\bullet}(C) = \mathrm{RHom}_{C^{\mathrm{op}} \otimes C}(C, C),$$

the derived endomorphisms of the diagonal bimodule. It is a graded-commutative algebra under Yoneda product and carries the Gerstenhaber bracket of degree  $-1$ .

**THEOREM 2.9** (*Deligne conjecture; Kontsevich–Soibelman, Tamarkin*). <sup>[PE]</sup> The pair (Yoneda product, Gerstenhaber bracket) on  $\mathrm{HH}^{\bullet}(C)$  is the homology of an  $E_2$ -algebra structure, canonical up to contractible choice.

For a  $\mathrm{CY}_d$  category, the cyclic pairing relates  $\mathrm{HH}^{\bullet}(C)$  to  $\mathrm{HH}_{\bullet}(C)[d]$ . Transporting structure through this duality produces the higher Calabi–Yau brackets.

**PROPOSITION 2.10** (*CY bracket spectrum*). <sup>[PE]</sup> For a  $\mathrm{CY}_d$  category  $C$ , the Hochschild invariants carry the following additional structure:

- $d = 1$ : a BV operator  $\Delta: \mathrm{HH}^{\bullet}(C) \rightarrow \mathrm{HH}^{\bullet-1}(C)$  whose deviation from being a derivation recovers the Gerstenhaber bracket;

- $d = 2$ : a degree  $-2$  Poisson bracket  $\{-, -\}_2$  on  $\mathrm{HH}^\bullet(C)$ , part of a 2-shifted symplectic structure on the moduli of objects;
- $d \geq 3$ : a  $(d-2)$ -shifted Poisson structure /  $P_{d-1}$ -algebra, refining to the Lie-cobar level in the cyclic setting.

THEOREM 2.II (*HKR for smooth varieties*). <sup>[PE]</sup> For  $X$  a smooth projective variety over  $k$  of characteristic zero, the Hochschild–Kostant–Rosenberg isomorphism gives

$$\mathrm{HH}^p(D^b(\mathrm{Coh}(X))) \simeq \bigoplus_{q+r=p} H^q(X, \wedge^r T_X).$$

When  $X$  is CY $_n$ , the trivialisation  $\omega_X \simeq \mathcal{O}_X$  identifies  $\wedge^r T_X \simeq \Omega_X^{n-r}$ , so

$$\mathrm{HH}^p(D^b(\mathrm{Coh}(X))) \simeq \bigoplus_{q+r=p} H^q(X, \Omega_X^{n-r}).$$

The sign and compatibility with the Mukai pairing are recorded in Căldăraru [3].

EXAMPLE 2.12 (*Hochschild cohomology of K3*). For a K3 surface  $X$ , the Hodge diamond is

$$h^{0,0} = 1, \quad h^{2,0} = h^{0,2} = 1, \quad h^{1,1} = 20, \quad h^{2,2} = 1,$$

with all other entries zero. Applied to the HKR sum  $\mathrm{HH}^p(D^b(\mathrm{Coh}(X))) = \bigoplus_{q+r=p} H^q(X, \Omega_X^{2-r})$ , this gives

$$\mathrm{HH}^0 \simeq H^0(\mathcal{O}) \oplus H^2(T_X) \simeq k \oplus k, \quad \mathrm{HH}^1 \simeq H^1(T_X) \simeq k^{20}, \quad \mathrm{HH}^2 \simeq H^0(\wedge^2 T_X) \oplus H^2(\mathcal{O}) \simeq k \oplus k.$$

The total dimension is

$$\dim_k \mathrm{HH}^\bullet(D^b(\mathrm{Coh}(K3))) = 2 + 20 + 2 = 24,$$

matching the topological Euler characteristic  $\chi(K3) = 24$ . The Mukai pairing on  $\mathrm{HH}^\bullet$  is the non-degenerate pairing  $\langle -, - \rangle_{\mathrm{Mukai}}$  of Mukai [28]; it coincides with the CY pairing produced by Definition 2.3 at  $d = 2$ .

REMARK 2.13 ( $\kappa_{\mathrm{cat}}$  from  $\chi$ ). For  $X$  a CY $_2$ , the categorical kappa of the Vol III spectrum is

$$\kappa_{\mathrm{cat}}(X) = \chi(\mathcal{O}_X),$$

the holomorphic Euler characteristic. For K3:  $\chi(\mathcal{O}_{K3}) = 1 - 0 + 1 = 2$ , giving  $\kappa_{\mathrm{cat}}(K3) = 2$ . Do not confuse this with the topological Euler characteristic 24 (which is  $\dim \mathrm{HH}^\bullet$ ), with  $\kappa_{\mathrm{ch}}(K3 \times E) = 3$  (chiral-algebra kappa via  $\Phi$ ), or with  $\kappa_{\mathrm{BKM}}(K3 \times E) = 5$  (Borcherds weight).

## 2.4 Cyclic $A_\infty$ structure

The CY pairing in Definition 2.3 is cohomological. The categorical input to  $\Phi$  requires a chain-level enhancement: a cyclic  $A_\infty$ -structure for which every higher operation  $m_n$  is compatible with the pairing.

*Definition 2.14 (Cyclic  $A_\infty$  category).* A cyclic  $A_\infty$ -structure of degree  $-d$  on an  $A_\infty$ -category  $C$  is a collection of pairings

$$\langle -, - \rangle_d : C(a_0, a_1) \otimes C(a_1, a_0) \longrightarrow k[-d]$$

together with the cyclic symmetry condition

$$\langle m_n(f_1, \dots, f_n), f_0 \rangle_d = (-1)^\epsilon \langle m_n(f_0, f_1, \dots, f_{n-1}), f_n \rangle_d$$

for all  $n \geq 2$ , where  $\epsilon$  is the Koszul sign determined by the degrees of the  $f_i$ .

*Definition 2.15 (Negative cyclic CY class).* A  $d$ -dimensional CY structure on a smooth dg category  $C$  is a class

$$[\sigma] \in \mathrm{HC}_d^-(C)$$

in the negative cyclic homology, whose image under the canonical map  $\mathrm{HC}^- \rightarrow \mathrm{HH}$  is a non-degenerate trace  $\mathrm{Tr} : \mathrm{HH}_d(C) \rightarrow k$ . The lift to  $\mathrm{HC}^-$  is essential for the  $\mathbb{S}^d$ -framing used in Chapter 8; see AP-CY2.

**THEOREM 2.16 (Cyclic  $A_\infty$  enhancement).** <sup>[PE]</sup> Every smooth proper  $\mathrm{CY}_d$  category admits a cyclic  $A_\infty$ -enhancement, unique up to  $A_\infty$ -quasi-isomorphism preserving the pairing. The proof is due to Kontsevich–Soibelman in the formal case, Costello [6] for the TCFT enhancement, and Seidel [32] for the Fukaya case.

At  $d = 2$  the enhancement is explicit enough to compute directly; it drives the  $K3$  computations feeding the Borchers denominator (Vol III Chapter 14.16). At  $d = 3$  the enhancement was constructed by Sheridan [33] for the quintic threefold, as part of the HMS proof; the general  $d = 3$  construction remains programme-level (conditional on  $\mathbb{S}^3$ -framing, see AP-CY6).

## 2.5 Interface with the CY-to-chiral functor $\Phi$

The Vol III functor

$$\Phi : \mathrm{CY}_d\text{-}\mathbf{Cat}^{\mathrm{cyc}} \longrightarrow E_2\text{-}\mathbf{ChirAlg}$$

takes as input the data of this chapter: a smooth proper cyclic  $A_\infty$  category of dimension  $d$ , with its negative cyclic CY class  $[\sigma] \in \mathrm{HC}_d^-(C)$ . The construction (Chapter 8) proceeds by (i) extracting the Gerstenhaber algebra  $\mathrm{HH}^\bullet(C)$ , (ii) promoting it to a Lie conformal algebra using the CY pairing, (iii) taking the factorization envelope on a curve, (iv) and enhancing to  $E_2$  using the  $\mathbb{S}^2$ -framing at  $d = 2$ .

**THEOREM 2.17 (Categorical kappa equals Euler characteristic:  $\mathrm{CY}\text{-}D_2$ ).** <sup>[PH]</sup> For  $C$  a smooth proper  $\mathrm{CY}_2$  category such that  $\Phi(C)$  exists, the modular characteristic of the chiral algebra satisfies

$$\kappa_{\mathrm{ch}}(\Phi(C)) = \chi^{\mathrm{CY}}(C),$$

the categorical Euler characteristic defined via the Mukai pairing. When  $C = D^b(\mathrm{Coh}(X))$  for  $X$  a  $K3$  surface, the right-hand side is  $\kappa_{\mathrm{cat}}(X) = \chi(\mathcal{O}_{K3}) = 2$ , which implies a contribution to  $\kappa_{\mathrm{ch}}(\Phi(D^b(\mathrm{Coh}(K3))))$ ; the full  $K3 \times E$  chiral kappa is 3, combining the  $K3$  and  $E$  factors.

*Remark 2.18 (The  $d = 3$  case).* For  $d = 3$ , the functor  $\Phi$  is not yet unconditional: it depends on the chain-level  $\mathbb{S}^3$ -framing programme (CY-A3, AP-CY6). Any downstream statement inheriting the  $d = 3$   $\Phi$  must carry <sup>[CD]</sup> and name CY-A3 as its dependency. The  $\mathbb{C}^3$  case is verified end-to-end in Chapter 8, Theorem 8.15, providing a Rosetta Stone for the general  $d = 3$  functor.

*Remark 2.19* (What  $\Phi$  does not see). Not every invariant of  $C$  survives the passage through  $\Phi$ . The functor retains the cyclic  $A_\infty$ -structure and the Gerstenhaber bracket; it loses the fine structure of the triangulated /  $A_\infty$  category (individual objects, t-structures, stability conditions) that is visible only via the fuller categorical CoHA construction of Chapter 15. The surviving numerical data are organized by the kappa-spectrum:  $\kappa_{\text{cat}}$  is always present,  $\kappa_{\text{ch}}$  exists only when  $\Phi$  is defined, and  $\kappa_{\text{BKM}}$  records the Borchers-side lift for  $K3 \times E$ .

*Remark 2.20* (Cross-volume conventions). Vol III uses lambda-bracket conventions for the Lie conformal algebras produced by  $\Phi$ : an OPE of the form  $T(z)T(w) \sim (c/2)(z-w)^{-4} + \dots$  is rewritten as  $\{T_\lambda T\} = (c/12)\lambda^3 + \dots$ , absorbing the combinatorial factor  $3! = 6$  from the divided-power  $\lambda^{(3)} = \lambda^3/3!$ . Vol I uses OPE modes directly; care is required when transporting formulas between volumes (see the concordance). The Hochschild / cyclic invariants of this chapter are convention-independent: they depend only on the chain-level  $A_\infty$ -structure.



## Chapter 3

# Cyclic $A_\infty$ -Structures

A Calabi–Yau category enters this volume through a single structural datum: a cyclic  $A_\infty$ -algebra of dimension  $d$ . Everything that follows, the functor  $\Phi$  to chiral algebras, the modular characteristic  $\kappa_{\text{cat}}$ , the four subscripted kappas of the CY kappa-spectrum, depends on this input. This chapter fixes the definitions, records the standard examples (elliptic curve,  $K_3$ , quintic), and states the bridge to  $\Pi$  precisely. The content is classical (Stasheff, Kontsevich, Keller, Costello); the Vol III role is the specific identification of  $d$  with the CY dimension appearing in the CY-A programme.

### 3.1 $A_\infty$ -algebras

*Definition 3.1 ( $A_\infty$ -algebra).* An  $A_\infty$ -algebra over a field  $k$  is a  $\mathbb{Z}$ -graded  $k$ -vector space  $A$  equipped with operations

$$\mu_n: A^{\otimes n} \longrightarrow A, \quad \deg(\mu_n) = 2 - n, \quad n \geq 1,$$

satisfying, for every  $n \geq 1$ , the  $A_\infty$ -relation

$$\sum_{\substack{p+q=n+1 \\ 1 \leq i \leq p}} (-1)^{\dagger(i,p,q)} \mu_p(a_1, \dots, a_{i-1}, \mu_q(a_i, \dots, a_{i+q-1}), a_{i+q}, \dots, a_n) = 0,$$

where  $\dagger(i, p, q) = (i-1)(q-1) + q \sum_{j=1}^{i-1} |a_j|$  follows the Koszul sign rule (see Appendix A).

Specializing to small  $n$  yields the familiar relations:

- $n = 1$ :  $\mu_1^2 = 0$ , so  $(A, \mu_1)$  is a cochain complex.
- $n = 2$ :  $\mu_1(\mu_2(a, b)) = \mu_2(\mu_1 a, b) + (-1)^{|a|} \mu_2(a, \mu_1 b)$ , so  $\mu_1$  is a derivation of  $\mu_2$ .
- $n = 3$ :  $\mu_2$  is associative up to the explicit homotopy  $\mu_3$ , giving the Stasheff associahedron relation.

An  $A_\infty$ -algebra with  $\mu_n = 0$  for  $n \geq 3$  is a differential graded associative algebra. The  $\mu_n$  for  $n \geq 3$  record the higher homotopies that obstruct strict associativity after homological projection (Kadeishvili’s theorem: any dg algebra transfers to a minimal  $A_\infty$ -algebra on its cohomology).

The  $A_\infty$ -formalism originates with Stasheff’s associahedra [34]. Its modern use, as the natural habitat for derived categories, mirror symmetry, and deformation quantization, is due to Kontsevich and Soibelman [23] and Keller [18]. For a CY category  $C$ , the derived category  $D^b(C)$  lifts canonically to an  $A_\infty$ -enhancement; this enhancement is unique up to quasi-isomorphism (Lunts–Orlov, Canonaco–Stellari).

The non-minimal cochain-level model is the preferred input for the Vol III functor because cyclicity is visible chain-level, not only on cohomology.

*Remark 3.2* ( $A_\infty$ -category vs  $A_\infty$ -algebra). An  $A_\infty$ -category is the many-object generalization: objects  $\text{Ob}(C)$  and Hom-complexes  $\text{Hom}_C(X, Y)$  with composition maps  $\mu_n: \text{Hom}(X_0, X_1) \otimes \cdots \otimes \text{Hom}(X_{n-1}, X_n) \rightarrow \text{Hom}(X_0, X_n)[2 - n]$ . An  $A_\infty$ -algebra is the one-object case. In this chapter we state definitions for algebras; the category version is verbatim by tagging arguments with source and target objects.

## 3.2 Cyclic structures and CY dimension

*Definition 3.3* (Cyclic  $A_\infty$ -algebra). A cyclic  $A_\infty$ -algebra of dimension  $d$  is a pair  $((A, \{\mu_n\}_{n \geq 1}), \langle -, - \rangle)$ , where  $(A, \{\mu_n\})$  is an  $A_\infty$ -algebra and

$$\langle -, - \rangle: A \otimes A \longrightarrow k[-d]$$

is a non-degenerate graded symmetric pairing of degree  $-d$  (equivalently,  $\langle a, b \rangle$  is nonzero only when  $|a| + |b| = d$ ), satisfying the *cyclic invariance* identity

$$\langle \mu_n(a_1, a_2, \dots, a_n), a_{n+1} \rangle = (-1)^{\epsilon_n} \langle a_1, \mu_n(a_2, \dots, a_{n+1}) \rangle \quad \text{for all } n \geq 1,$$

with sign  $\epsilon_n = n + |a_1|(|a_2| + \cdots + |a_{n+1}|)$  determined by the Koszul rule.

The integer  $d$  is the *CY dimension* of the cyclic  $A_\infty$ -algebra. In the language of Costello [7, 8], a cyclic  $A_\infty$ -algebra of dimension  $d$  is precisely an algebra over the operad of framed chains on the moduli of disks with marked points, with a  $d$ -shifted symplectic structure. This operadic viewpoint is what produces, after homotopy transfer, the  $\mathbb{S}^d$ -framing of Hochschild homology used in Chapter 8.

**PROPOSITION 3.4** (*Cyclic invariance on the Hochschild complex*). <sup>[PE]</sup> Let  $(A, \langle -, - \rangle)$  be a cyclic  $A_\infty$ -algebra of dimension  $d$ . The pairing induces a non-degenerate pairing on the Hochschild complex

$$\langle -, - \rangle_{\text{HH}}: \text{HH}_\bullet(A) \otimes \text{HH}_{d-\bullet}(A) \longrightarrow k$$

implementing Poincaré duality of dimension  $d$ . Equivalently, the cyclic structure lifts Hochschild homology to negative cyclic homology with a class in  $\text{HC}_d^-(A)$  that is closed under the Connes differential.

*Attribution.* Costello [8] Thm. A; Kontsevich–Soibelman [23] Sec. 10. The AP-CY2 distinction is essential: the CY trace lives in  $\text{HC}_d^-(A)$ , not in  $\text{HH}_d \rightarrow k$ , because the  $S^1$ -invariance of the cyclic pairing is needed to produce the  $\mathbb{S}^d$ -framing.  $\square$

*Remark 3.5* (Geometric origin of  $d$ ). For  $X$  a smooth projective CY  $n$ -fold, the derived category  $D^b(\text{Coh}(X))$  carries a cyclic  $A_\infty$ -enhancement of dimension  $d = n$ . The pairing is the Serre-duality pairing

$$\langle -, - \rangle_{\text{Serre}}: \text{Ext}^i(\mathcal{F}, \cdot) \otimes \text{Ext}^{n-i}(\cdot, \mathcal{F}) \longrightarrow \text{Ext}^n(\mathcal{F}, \mathcal{F}) \xrightarrow{\text{tr}} H^n(X, \omega_X) \cong k,$$

using the trivialization  $\omega_X \cong \mathcal{O}_X$  that defines the CY structure. AP-CY1:  $d$  is the complex dimension, not the real dimension  $2n$ . The Serre functor  $\mathbb{S} = (-) \otimes \omega_X[n]$  becomes  $[n]$ -shift under the CY trivialization, which is what produces the  $d$ -shifted pairing.

*Remark 3.6* (Cyclic bar complex). The cyclic pairing enters the bar complex  $B(A) = T^c(s^{-1}\bar{A})$  through the cyclic quotient  $\text{CC}_\bullet(A) = B(A)/(1-t)$  where  $t$  is the signed cyclic rotation. The factor  $s^{-1}$  desuspends:  $|s^{-1}v| = |v| - 1$ . The augmentation ideal  $\bar{A} = \ker(\varepsilon)$  is used rather than  $A$  itself. The cyclic bar complex is the primary invariant of  $(A, \mu_n, \langle -, - \rangle)$  and is what II promotes to a factorization coalgebra on curves.



*Remark 3.7* (Cyclic symmetry at  $n = 2$ ). Specializing the cyclic invariance identity to  $n = 2$  gives

$$\langle \mu_2(a_1, a_2), a_3 \rangle = (-1)^{\epsilon_2} \langle a_1, \mu_2(a_2, a_3) \rangle,$$

which for a strictly associative cyclic algebra ( $\mu_n = 0$  for  $n \geq 3$ ) reduces to the standard invariance condition for a Frobenius algebra of degree  $d$ . The higher- $n$  cases are the  $A_\infty$ -correction:  $\mu_3, \mu_4, \dots$  must individually respect the pairing. In the  $n = 3$  case, the identity

$$\langle \mu_3(a_1, a_2, a_3), a_4 \rangle = (-1)^{\epsilon_3} \langle a_1, \mu_3(a_2, a_3, a_4) \rangle$$

is the first genuinely homotopical constraint and is what fails in a naive restriction of the Fukaya category to a sub-Lagrangian: cyclic invariance distinguishes the true CY input from merely  $A_\infty$ -enhanceable ones.

### 3.3 Examples

The four examples below cover the four CY dimensions  $d \in \{1, 1, 2, 3\}$  relevant to Vol III. Each fixes one numerical invariant visible to the functor  $\Phi$ .

*Example 3.8* ( $d = 1$  from a CY1 Lie algebra). Let  $\mathfrak{g}$  be a finite-dimensional simple Lie algebra with non-degenerate invariant form  $\langle -, - \rangle_{\mathfrak{g}}$ . The Chevalley–Eilenberg cochain complex  $C^\bullet(\mathfrak{g})$  is a dg commutative algebra of dimension  $\dim \mathfrak{g}$ ; applying Koszul duality twice produces a cyclic  $A_\infty$ -algebra of CY dimension 1 whose underlying vector space is  $\mathfrak{g}[1]$  with pairing  $\langle -, - \rangle_{\mathfrak{g}}$  (shifted to degree  $-1$ ). For  $\mathfrak{g} = \mathfrak{sl}_2$ , the resulting cyclic  $A_\infty$ -algebra has 3 generators and  $\mu_2$  given by the bracket; all higher  $\mu_n$  vanish. This is the simplest nontrivial cyclic  $A_\infty$ -algebra relevant to Vol III, and it recovers  $\widehat{\mathfrak{sl}}_2$  as its chiral image under  $\Phi$ .

*Example 3.9* (Explicit  $\mu_2$  for  $\mathfrak{sl}_2$ ). Continuing Example 3.8, the underlying graded space of the cyclic  $A_\infty$ -algebra for  $\mathfrak{sl}_2$  is the suspension  $\mathfrak{sl}_2[1]$ , with basis  $\{e[1], f[1], h[1]\}$  in degree  $-1$ . The binary operation  $\mu_2$  is the desuspended Lie bracket:

$$\mu_2(e[1], f[1]) = h[1], \quad \mu_2(h[1], e[1]) = 2e[1], \quad \mu_2(h[1], f[1]) = -2f[1],$$

with the other values determined by graded antisymmetry. The cyclic pairing has nonzero values

$$\langle e[1], f[1] \rangle = 1, \quad \langle h[1], h[1] \rangle = 2,$$

which is the Killing form shifted to degree  $-1$ , producing a CY structure of dimension  $d = 1$ . The  $A_\infty$ -relations reduce to the Jacobi identity of  $\mathfrak{sl}_2$ , and all  $\mu_n$  for  $n \geq 3$  vanish. This is the simplest nontrivial cyclic  $A_\infty$ -algebra in the volume; under  $\Phi$  it produces the affine Kac–Moody algebra  $\widehat{\mathfrak{sl}}_2$  at the critical level, recovering a classical landscape entry from the CY engine.

*Example 3.10* ( $d = 1$  from an elliptic curve). Let  $E$  be an elliptic curve over  $\mathbb{C}$ . The derived category  $D^b(\text{Coh}(E))$  has a minimal cyclic  $A_\infty$ -enhancement of dimension  $d = 1$ , computed explicitly by Polishchuk [30]. The generators are  $\mathcal{O}_E$  and a single degree-1 class in  $\text{Ext}^1(\mathcal{O}_E, \mathcal{O}_E) = H^1(E, \mathcal{O}_E) \cong \mathbb{C}$ . The Serre pairing  $\text{Ext}^1(\mathcal{O}_E, \mathcal{O}_E) \otimes \text{Hom}(\mathcal{O}_E, \mathcal{O}_E) \rightarrow \mathbb{C}$  realizes the  $d = 1$  cyclic structure. For higher-degree line bundles, Polishchuk’s theta-function computation gives explicit formulas for  $\mu_n, n \geq 3$ , in terms of modular forms on  $\Gamma_1(N)$ . This example sits at the boundary between  $d = 1$  (where the  $\mathbb{S}^1$ -framing is classical) and the modular-form territory that Vol I controls.

*Example 3.11* ( $d = 2$  from  $K_3$ ). Let  $X$  be a smooth projective  $K_3$  surface. The derived category  $D^b(\text{Coh}(X))$  is a cyclic  $A_\infty$ -category of dimension  $d = 2$  (Caldararu [3]). The Hochschild cohomology is

$$\text{HH}^\bullet(D^b(\text{Coh}(X))) \cong H^\bullet(X, \wedge^\bullet T_X) \cong H^\bullet(X, \mathbb{C}),$$

using the CY trivialization  $\wedge^2 T_X \cong \mathcal{O}_X$ . The HKR decomposition gives  $\text{HH}^0 \cong k^2$ ,  $\text{HH}^1 \cong k^{20}$ ,  $\text{HH}^2 \cong k^2$  (as computed in Example 2.12 of Chapter 2), so

$$\dim \text{HH}^\bullet(K_3) = 2 + 20 + 2 = 24,$$

recovering the Euler characteristic  $\chi(X) = 24$  familiar from lattice  $K_3$  geometry. The holomorphic Euler characteristic, which is the relevant invariant for the CY kappa-spectrum, is

$$\chi(\mathcal{O}_X) = 2.$$

This is the value  $\kappa_{\text{cat}}(K_3) = 2$  entering the Vol III kappa-table.

*Example 3.12* ( $d = 3$  from the quintic). Let  $Q \subset \mathbb{P}^4$  be a smooth quintic threefold.  $D^b(\text{Coh}(Q))$  is a cyclic  $A_\infty$ -category of dimension  $d = 3$ . By homological mirror symmetry (Kontsevich's conjecture, established for the quintic at genus 0 by Sheridan [33]), there is an  $A_\infty$ -equivalence

$$D^b(\text{Coh}(Q)) \simeq D^\pi \text{Fuk}(Q^\vee),$$

where  $Q^\vee$  is the mirror quintic and  $\text{Fuk}$  is the Fukaya category (wrapped, compact, depending on geometric input). Both sides are cyclic  $A_\infty$  of dimension  $d = 3$ , and HMS is an equivalence of cyclic  $A_\infty$ -categories: the Serre pairing on the B-side matches the Poincaré pairing on Lagrangian intersections on the A-side. The Hochschild cohomology

$$\text{HH}^\bullet(Q) \cong H^\bullet(Q, \wedge^\bullet T_Q)$$

has  $\dim \text{HH}^1 = 101$  (the Kodaira–Spencer space of the quintic) and  $\dim \text{HH}^2 = 1$ . The holomorphic Euler characteristic is  $\chi(\mathcal{O}_Q) = 0$ , so the naive  $\kappa_{\text{cat}}$  vanishes; the nontrivial chiral data enters through  $\text{HH}^1$ , not through the top pairing. This is the  $\text{CY}_3$  regime where the chiral algebra  $A_Q$  is not yet constructed: any result that passes through  $A_Q$  is conditional on  $\text{CY-}A_3$ .

### 3.4 The bridge to the Vol III functor

The cyclic  $A_\infty$ -structure is the complete input to the CY-to-chiral functor. The signature is

$$\Phi: \text{CY}_d\text{-Cat} \longrightarrow E_2\text{-ChirAlg}, \quad (C, \mu_n, \langle -, - \rangle) \longmapsto A_C,$$

and the data of  $(C, \mu_n, \langle -, - \rangle)$  factors through  $\Phi$  by the four-step passage in Chapter 8: Hochschild cohomology with Gerstenhaber bracket, Lie conformal algebra via the cyclic pairing, factorization envelope, and  $E_2$ -enhancement via the  $\mathbb{S}^d$ -framing. The cyclic pairing  $\langle -, - \rangle$  is what becomes the invariant form on  $A_C$ ; without it, the factorization envelope is not a chiral algebra with a trace.

**THEOREM 3.13** (*Cyclic  $A_\infty$  input determines  $\kappa_{\text{cat}}$  at  $d = 2$* ). <sup>[PH]</sup> Let  $C$  be a smooth proper cyclic  $A_\infty$ -category of dimension  $d = 2$ , and let  $A_C = \Phi(C)$  be its image under the  $d = 2$  CY-to-chiral functor (Theorem 8.1). Then the categorical modular characteristic satisfies

$$\kappa_{\text{cat}}(C) = \chi^{\text{CY}}(C) = \sum_i (-1)^i \dim \text{HH}^i(C) \Big|_{\text{top pairing}},$$

and this equals the chiral modular characteristic  $\kappa_{\text{ch}}(A_C)$  of the image chiral algebra at  $d = 2$  (Proposition 8.2; the Serre duality  $\mathbb{S}_C \simeq [2]$  kills the one-loop correction from quantization Step 4). For  $C = D^b(\text{Coh}(K3))$ , both invariants evaluate to  $\kappa_{\text{cat}}(K3) = 2$ . At  $d \geq 3$  the identification remains conjectural (Conjecture 8.3).

*Proof sketch.* The cyclic pairing of dimension  $d = 2$  produces a degree- $(-2)$  symmetric form on the Hochschild complex; its top-degree component is a linear map  $\text{HH}^2(C) \rightarrow k$  whose dimension (as a quotient of  $\text{HH}^\bullet$ ) is  $\chi(\mathcal{O}_X)$  in the geometric case. The CY-to-chiral functor sends this trace to the invariant form on  $A_C$ , and the resulting  $\kappa_{\text{cat}}$  coincides (up to  $\mathbb{S}^2$ -framing) with the coefficient of the  $E_2$ -algebra central extension. For  $K3$  the computation reduces to  $\chi(\mathcal{O}_{K3}) = 2$ ; see Chapter 8 Section 8.2 and the compute module `cy_to_chiral_functor.py` for the full verification.  $\square$

CONJECTURE 3.14 (*Cyclic  $A_\infty$  input determines  $\kappa_{\text{cat}}$  at  $d = 3$* ). <sup>[CJ]</sup> Let  $C$  be a smooth proper cyclic  $A_\infty$ -category of dimension  $d = 3$ . Assume Conjecture CY-A<sub>3</sub> (the  $d = 3$  CY-to-chiral functor exists with chain-level  $\mathbb{S}^3$ -framing). Then

$$\kappa_{\text{cat}}(C) = \chi^{\text{CY}}(C),$$

and for  $C = D^b(\text{Coh}(Q))$  with  $Q$  a smooth quintic threefold, this evaluates to a specific integer determined by the  $\mathbb{S}^3$ -framing obstruction. The conjecture is conditional on CY-A<sub>3</sub>; any downstream result passing through  $A_Q$  inherits this conditionality.

*Remark 3.15* (Scope: what cyclic  $A_\infty$  does not determine). The cyclic  $A_\infty$ -structure determines  $\kappa_{\text{cat}}$  and the chiral kappa  $\kappa_{\text{ch}}$  via the functor  $\Phi$ , but it does *not* determine the BKM kappa  $\kappa_{\text{BKM}}$  or the fiber kappa  $\kappa_{\text{fiber}}$  of the kappa-spectrum. Those kappas arise from additional data (lattice embedding for  $\kappa_{\text{fiber}}$ , Borchers product structure for  $\kappa_{\text{BKM}}$ ) that is not visible to the cyclic  $A_\infty$ -category alone. The four-way kappa-spectrum for  $K3 \times E$  is  $\{\kappa_{\text{cat}}, \kappa_{\text{ch}}, \kappa_{\text{BKM}}, \kappa_{\text{fiber}}\} = \{2, 3, 5, 24\}$ ; only the first two are extracted from the cyclic  $A_\infty$ -input.

*Remark 3.16* (Comparison with prior work). Four source threads feed the construction used here. Stasheff [34] introduced the associahedra and the higher homotopies  $\mu_n$ . Kontsevich [22] identified cyclic  $A_\infty$ -algebras with algebras over the operad of ribbon graphs, providing the link to moduli of curves with boundary. Costello [7, 8] proved that cyclic  $A_\infty$ -categories are equivalent to open topological conformal field theories and supplied the first rigorous construction of the associated chain-level trace. Kontsevich–Soibelman [23] axiomatized the CY structure in terms of the negative cyclic class and gave the formalism used in Part I. Keller [18] surveys the homological-algebra side. For explicit computations on projective varieties, Polishchuk [30] computed the cyclic  $A_\infty$ -structure on elliptic curves and on their products, and Caldararu [3] set up the Hochschild calculus for smooth proper CY categories. The Vol III role is the specific mapping of this input through the functor  $\Phi$ , producing chiral algebras whose modular characteristic can be computed and compared across the four kappas of the spectrum.

*Remark 3.17* (Costello’s operadic description). Costello [7] proves that cyclic  $A_\infty$ -algebras of dimension  $d$  are equivalent to algebras over the open topological conformal field theory operad with  $d$ -shifted framing. In this formulation, the  $\mathbb{S}^d$ -framing used in Chapter 8 is visible as the action of the framing circle on the moduli of disks with boundary marked points. For  $d = 2$  the framing is abelian ( $\pi_1(SO(2)) = \mathbb{Z}$ ) and the envelope step produces an honest  $E_2$ -algebra. For  $d = 3$  the framing is nonabelian ( $\pi_1(SO(3)) = \mathbb{Z}/2$ ) and the envelope step is the source of the  $\mathbb{S}^3$ -framing obstruction mentioned in AP-CY6.



## Chapter 4

# Hochschild Calculus for CY Categories

*Remark 4.1* (Convention: categorical Hochschild). In this chapter, “Hochschild” refers to the categorical Hochschild invariants of a dg category (the cyclic bar complex  $CC_*(C)$ ), which are topological in nature. The chiral Hochschild cohomology  $\text{ChirHoch}^*$  of Volume I (Theorem H) is the chiral upgrade incorporating OPE data and curve geometry. The CY-to-chiral functor  $\Phi$  of Chapter 8 maps the categorical Gerstenhaber bracket to the chiral convolution bracket.

A Calabi–Yau  $d$ -category has a non-degenerate Serre pairing. What does this pairing force on the Hochschild invariants? For a general dg category, Hochschild homology  $HH_\bullet(C)$  carries only a Connes  $B$ -operator and Hochschild cohomology  $HH^\bullet(C)$  carries only a Gerstenhaber bracket; the two are linked by a cap product, but no pairing relates them. The CY structure removes this deficiency.

The Serre pairing provides a quasi-isomorphism  $HH^\bullet(C) \simeq HH_{d-\bullet}(C)$  of degree  $-d$ , so that Hochschild cohomology inherits the  $B$ -operator and Hochschild homology inherits the Gerstenhaber bracket. The combination is a  $(2-d)$ -shifted Poisson (equivalently,  $BV_{2-d}$ ) structure on  $HH^\bullet(C)$ : bracket of degree  $1-d$ , operator of degree  $-1$ , compatibility equation  $[B, -] = \{-, -\}$ . At  $d = 2$  this is a 0-shifted Poisson structure (ordinary Poisson); at  $d = 3$ , a  $(-1)$ -shifted Poisson structure (the CY<sub>3</sub> case of Pantev–Toën–Vaquié–Vezzosi); at  $d = 1$ , a 1-shifted Poisson structure of the kind that drives factorization homology.

This chapter develops the Hochschild calculus underlying the CY-to-chiral functor of Chapter 8. The shifted Poisson structure on  $HH^\bullet(C)$  is the categorical precursor of the convolution Lie bracket on the modular deformation complex of Volume I: the Gerstenhaber bracket becomes the bar-cobar convolution bracket, the Connes  $B$ -operator becomes the modular differential, and the Serre pairing becomes the cyclic duality that controls the  $S^d$ -framing. The cyclic bar complex of Chapter 3 is the single-object specialization of the Hochschild complex constructed here.

### 4.1 Hochschild homology and cohomology

*Definition 4.2* (Hochschild complex). Let  $C$  be a dg category over  $k$ . The *Hochschild chain complex* of  $C$  is

$$CC_n(C) = \bigoplus_{X_0, \dots, X_n \in \text{Ob}(C)} \text{Hom}_C(X_n, X_0) \otimes \text{Hom}_C(X_{n-1}, X_n) \otimes \cdots \otimes \text{Hom}_C(X_0, X_1)$$

with the Hochschild differential

$$b(f_0 \otimes \cdots \otimes f_n) = \sum_{i=0}^{n-1} (-1)^{\epsilon_i} f_0 \otimes \cdots \otimes f_i f_{i+1} \otimes \cdots \otimes f_n \\ + (-1)^{\epsilon_n} f_n f_0 \otimes f_1 \otimes \cdots \otimes f_{n-1},$$

where  $\epsilon_i = |f_0| + \cdots + |f_i|$  (Koszul signs). The *Hochschild homology* is  $\mathrm{HH}_\bullet(C) = H_\bullet(\mathrm{CC}_\bullet(C), b)$ .

**Definition 4.3** (*Hochschild cohomology*). The *Hochschild cochain complex* of a dg category  $C$  is

$$\mathrm{CC}^n(C) = \prod_{X_0, \dots, X_n \in \mathrm{Ob}(C)} \mathrm{Hom}_k(\mathrm{Hom}_C(X_{n-1}, X_n) \otimes \cdots \otimes \mathrm{Hom}_C(X_0, X_1), \mathrm{Hom}_C(X_0, X_n))$$

with the Hochschild coboundary  $\delta$  dual to  $b$ . The *Hochschild cohomology* is  $\mathrm{HH}^\bullet(C) = H^\bullet(\mathrm{CC}^\bullet(C), \delta)$ .

**Remark 4.4** (Morita invariance). Both  $\mathrm{HH}_\bullet$  and  $\mathrm{HH}^\bullet$  are Morita invariant: if  $C$  and  $\mathcal{D}$  are derived Morita equivalent (i.e.,  $\mathrm{Perf}(C) \simeq \mathrm{Perf}(\mathcal{D})$  as dg categories), then  $\mathrm{HH}_\bullet(C) \simeq \mathrm{HH}_\bullet(\mathcal{D})$  and  $\mathrm{HH}^\bullet(C) \simeq \mathrm{HH}^\bullet(\mathcal{D})$ . This is essential for the CY programme, where different geometric models (Fukaya, derived coherent sheaves, matrix factorizations) of the same CY category must produce the same Hochschild invariants.

## 4.2 The Gerstenhaber bracket

**THEOREM 4.5** (*Gerstenhaber structure on  $\mathrm{HH}^\bullet$* ). <sup>[PE]</sup> The Hochschild cohomology  $\mathrm{HH}^\bullet(C)$  carries two compatible structures:

- (i) A graded commutative *cup product*  $\smile : \mathrm{HH}^p(C) \otimes \mathrm{HH}^q(C) \rightarrow \mathrm{HH}^{p+q}(C)$ , the composition of multilinear maps;
- (ii) A degree  $(-1)$  Lie bracket  $[\cdot, \cdot] : \mathrm{HH}^p(C) \otimes \mathrm{HH}^q(C) \rightarrow \mathrm{HH}^{p+q-1}(C)$  (the *Gerstenhaber bracket*), defined at the cochain level by

$$[f, g] = f \circ g - (-1)^{(|f|-1)(|g|-1)} g \circ f$$

where  $\circ$  denotes the operadic pre-Lie composition:

$$(f \circ g)(a_1, \dots, a_{p+q-1}) = \sum_{i=1}^p (-1)^\dagger f(a_1, \dots, a_{i-1}, g(a_i, \dots, a_{i+q-1}), a_{i+q}, \dots, a_{p+q-1}).$$

Together, the cup product and Gerstenhaber bracket make  $\mathrm{HH}^\bullet(C)$  a *Gerstenhaber algebra*: the bracket is a derivation of the cup product in each variable.

**Remark 4.6** (Gerstenhaber =  $E_2$  cohomology). By Kontsevich's formality theorem and the Deligne conjecture (proved by Kontsevich–Soibelman, McClure–Smith, and others),  $\mathrm{CC}^\bullet(C)$  carries a natural  $E_2$ -algebra structure whose cohomology-level shadow is the Gerstenhaber algebra  $(\mathrm{HH}^\bullet(C), \smile, [\cdot, \cdot])$ . The  $E_2$ -structure is the chain-level origin of the braided monoidal category structure on  $\mathrm{Rep}^{E_2}(Z_{\mathrm{ch}}^{\mathrm{der}}(A))$  (Chapter 13).

**PROPOSITION 4.7** (*Gerstenhaber bracket controls deformations*). <sup>[PE]</sup> The Gerstenhaber bracket controls the deformation theory of  $C$ :

- (i)  $\mathrm{HH}^2(C)$  classifies first-order deformations of  $C$  as a dg category;

- (ii) The obstruction to extending a first-order deformation  $\mu_1 \in \mathrm{HH}^2(C)$  to second order lies in  $\mathrm{HH}^3(C)$  and is given by the Gerstenhaber square  $\frac{1}{2}[\mu_1, \mu_1]$ ;
- (iii) A formal deformation is an element  $\mu = \sum_{n \geq 1} \hbar^n \mu_n \in \mathrm{HH}^2(C)[[\hbar]]$  satisfying the Maurer–Cartan equation  $\delta\mu + \frac{1}{2}[\mu, \mu] = 0$ .

The MC equation for deformations of  $C$  maps, under  $\Phi$ , to the MC equation  $D\Theta_A + \frac{1}{2}[\Theta_A, \Theta_A] = 0$  of the modular convolution algebra  $\mathfrak{g}_A^{\mathrm{mod}}$  (Volume I, thm:mc2-bar-intrinsic).

### 4.3 The Connes $B$ -operator and cyclic homology

*Definition 4.8 (Connes  $B$ -operator).* The Connes operator  $B: \mathrm{CC}_n(C) \rightarrow \mathrm{CC}_{n+1}(C)$  is defined by

$$B(f_0 \otimes \cdots \otimes f_n) = \sum_{i=0}^n (-1)^{ni} 1 \otimes f_i \otimes f_{i+1} \otimes \cdots \otimes f_{i-1}$$

(indices mod  $n+1$ ), composed with the insertion of the identity. It satisfies  $B^2 = 0$ ,  $bB + Bb = 0$ .

*Definition 4.9 (Cyclic, negative cyclic, and periodic cyclic homology).* From the mixed complex  $(\mathrm{CC}_\bullet(C), b, B)$  with  $|b| = -1$ ,  $|B| = +1$ ,  $b^2 = B^2 = bB + Bb = 0$ :

- (i) *Cyclic homology:*  $\mathrm{HC}_\bullet(C) = H_\bullet(\mathrm{CC}_\bullet(C) \otimes_{k[B]} k)$ ;
- (ii) *Negative cyclic homology:*  $\mathrm{HC}_\bullet^-(C) = H_\bullet(\mathrm{CC}_\bullet(C)[[u]], b + uB)$ , where  $|u| = 2$ ;
- (iii) *Periodic cyclic homology:*  $\mathrm{HP}_\bullet(C) = H_\bullet(\mathrm{CC}_\bullet(C)((u)), b + uB)$ .

The SBI sequence  $\cdots \rightarrow \mathrm{HH}_n \xrightarrow{B} \mathrm{HC}_{n+1} \xrightarrow{S} \mathrm{HC}_{n-1} \xrightarrow{I} \mathrm{HH}_{n-1} \rightarrow \cdots$  relates these.

*Example 4.10 (Smooth variety).* For  $C = \mathrm{Perf}(X)$  with  $X$  a smooth algebraic variety of dimension  $n$ , the Hochschild–Kostant–Rosenberg theorem gives

$$\mathrm{HH}_k(\mathrm{Perf}(X)) \simeq \bigoplus_{p-q=k} H^q(X, \Omega_X^p), \quad \mathrm{HH}^k(\mathrm{Perf}(X)) \simeq \bigoplus_{p+q=k} H^q(X, \bigwedge^p T_X).$$

Under HKR, the Connes  $B$ -operator is the de Rham differential  $d_{\mathrm{dR}}: \Omega^p \rightarrow \Omega^{p+1}$ , and the Gerstenhaber bracket is the Schouten–Nijenhuis bracket on polyvector fields.

### 4.4 The Hochschild-to-cyclic spectral sequence

**PROPOSITION 4.11** (*Hodge-to-de Rham degeneration for dg categories*). <sup>[PE]</sup> For a smooth proper dg category  $C$  over a field  $k$  of characteristic zero, the Hochschild-to-cyclic spectral sequence

$$E_1^{p,q} = \mathrm{HH}_{q-p}(C) \implies \mathrm{HC}_{q-p}(C)$$

degenerates at  $E_2$ .

*Remark 4.12* (Kaledin’s theorem). The degeneration was proved by Kaledin (2008) using Deligne–Illusie lifting methods in the dg setting. Kontsevich–Soibelman (2009) gave an independent proof via the theory of formal moduli problems. The degeneration is the categorical analogue of Hodge-to-de Rham degeneration for smooth projective varieties. In the CY programme, it is the categorical counterpart of the PBW spectral sequence collapse (Volume I, Koszulness characterization K2): both express the same underlying formality, one at the categorical level and one at the bar-complex level.

## 4.5 CY Hochschild duality

**THEOREM 4.13** (*CY Hochschild duality*). <sup>[PE]</sup> For a smooth CY category  $\mathcal{C}$  of dimension  $d$ , there is a canonical quasi-isomorphism

$$\mathrm{CC}^\bullet(\mathcal{C}) \simeq \mathrm{CC}_{\bullet+d}(\mathcal{C})$$

identifying the Hochschild cochain complex with the  $d$ -shifted Hochschild chain complex. On cohomology:

$$\mathrm{HH}^n(\mathcal{C}) \simeq \mathrm{HH}_{n+d}(\mathcal{C}).$$

Under this identification:

- (i) The Gerstenhaber bracket on  $\mathrm{HH}^\bullet$  corresponds to the *BV operator*  $\Delta_{\mathrm{BV}}: \mathrm{HH}_{n+d} \rightarrow \mathrm{HH}_{n+d+1}$  (the divergence operator associated to the CY volume form);
- (ii) The cup product on  $\mathrm{HH}^\bullet$  corresponds to the intersection product on  $\mathrm{HH}_{\bullet+d}$ ;
- (iii) The Connes  $B$ -operator on  $\mathrm{HH}_\bullet$  corresponds to the contraction with the Euler vector field under HKR.

*Proof sketch.* The CY condition furnishes an equivalence  $\mathcal{C} \simeq \mathcal{C}^\vee[d]$  between the category and its  $d$ -shifted dual. Smoothness (the diagonal bimodule  $C_\Delta$  is compact as a  $\mathcal{C} \otimes \mathcal{C}^{\mathrm{op}}$ -module) gives

$$\mathrm{CC}^\bullet(\mathcal{C}) = \mathrm{RHom}_{\mathcal{C} \otimes \mathcal{C}^{\mathrm{op}}}(C_\Delta, C_\Delta) \simeq C_\Delta^\vee \otimes_{\mathcal{C} \otimes \mathcal{C}^{\mathrm{op}}} C_\Delta \simeq \mathrm{CC}_\bullet(\mathcal{C}^\vee) \simeq \mathrm{CC}_{\bullet+d}(\mathcal{C})$$

using  $\mathcal{C}^\vee \simeq \mathcal{C}[-d]$  from the CY structure. The identification of the algebraic structures follows from the compatibility of the Serre functor with composition; see Kontsevich (ICM 2006), Keller (2006), Ginzburg (arXiv:0612139).  $\square$

## 4.6 The $(2 - d)$ -shifted Poisson structure

The CY Hochschild duality has a structural consequence:  $\mathrm{HH}^\bullet(\mathcal{C})$  carries a  $(2 - d)$ -shifted Poisson structure.

**THEOREM 4.14** ( *$(2 - d)$ -shifted Poisson structure*). <sup>[PE]</sup> For a smooth  $\mathrm{CY}_d$  category  $\mathcal{C}$ , the Hochschild cohomology  $\mathrm{HH}^\bullet(\mathcal{C})$  carries a  $(2 - d)$ -shifted Poisson structure: a Lie bracket of cohomological degree  $(2 - d)$

$$\{\cdot, \cdot\}_{2-d}: \mathrm{HH}^p(\mathcal{C}) \otimes \mathrm{HH}^q(\mathcal{C}) \rightarrow \mathrm{HH}^{p+q+2-d}(\mathcal{C})$$

satisfying the Jacobi identity and the Leibniz rule with respect to the cup product. This bracket is the CY-shifted Gerstenhaber bracket:

$$\{f, g\}_{2-d} = [f, g]_{\mathrm{Gerst}} \circ \sigma_d$$

where  $\sigma_d$  is the CY structure class in  $\mathrm{HC}_d^-(\mathcal{C}) \simeq \mathrm{HH}_0(\mathcal{C})$ .

*Remark 4.15* (The three dimensions). The shifted Poisson structure specializes by dimension:

- (a)  $d = 1$  (curves):  $(2 - d) = 1$ -shifted Poisson. The Gerstenhaber bracket has degree  $-1$ ; the shifted bracket has degree  $0$ . This is an ordinary Lie bracket on  $\mathrm{HH}^\bullet$ , recovering the classical Poisson bracket on deformations of the curve.



- (b)  $d = 2$  (surfaces):  $(2 - d) = 0$ -shifted Poisson. The shifted bracket has degree 0 and the cup product makes  $\mathrm{HH}^\bullet(C)$  a Poisson algebra in the ordinary sense. This is the Poisson structure that the CY-to-chiral functor  $\Phi$  promotes to the chiral Poisson (coisson) structure of the associated vertex algebra. The convolution Lie bracket on  $\mathfrak{g}_A^{\mathrm{mod}}$  (Volume I) is the chiral incarnation of this 0-shifted Poisson bracket.
- (c)  $d = 3$  (threefolds):  $(2 - d) = (-1)$ -shifted Poisson. The bracket has degree  $-1$ , making  $\mathrm{HH}^\bullet(C)$  a  $(-1)$ -shifted Poisson algebra = BV algebra (at the cochain level, a  $\mathrm{BV}_\infty$ -algebra). This is the BV structure underlying the BRST complex of the associated 3d holomorphic-topological theory. The chiral algebra produced by  $\Phi$  is  $E_1$ , not  $E_2$ : the  $(-1)$ -shifted Poisson bracket produces a Lie bracket on the loop space (= Hochschild chains), not a Poisson bracket on functions.

## 4.7 Categorical Hodge theory

*Definition 4.16 (Hodge filtration on periodic cyclic homology).* For a smooth proper dg category  $C$  over  $k = \mathbb{C}$ , the *categorical Hodge filtration* on the periodic cyclic homology  $\mathrm{HP}_\bullet(C)$  is the filtration

$$F^p \mathrm{HP}_n(C) = \mathrm{im}(\mathrm{HC}_{n+2p}^-(C) \rightarrow \mathrm{HP}_n(C))$$

induced by the natural map from negative to periodic cyclic homology. The associated graded is

$$\mathrm{gr}_F^p \mathrm{HP}_n(C) \simeq \mathrm{HH}_{n+2p}(C).$$

**PROPOSITION 4.17** (*Non-commutative Hodge-to-de Rham*). <sup>[PE]</sup> For a smooth proper  $\mathrm{CY}_d$  category  $C$ :

- (i) The Hodge filtration degenerates (Kaledin), giving  $\mathrm{HP}_n(C) \simeq \prod_p \mathrm{HH}_{n+2p}(C)$  as a filtered vector space;
- (ii) The CY structure class  $[\sigma] \in \mathrm{HC}_d^-(C)$  determines a preferred splitting of the Hodge filtration;
- (iii) In the commutative case  $C = \mathrm{Perf}(X)$ , the categorical Hodge filtration recovers the classical Hodge filtration on  $H_{\mathrm{dR}}^n(X)$  via HKR.

*Remark 4.18* (Bridge to the shadow obstruction tower). Under the CY-to-chiral functor  $\Phi$  (Part II), the categorical Hodge filtration maps to the weight filtration on the modular convolution algebra  $\mathfrak{g}_A^{\mathrm{mod}}$ . The associated graded pieces correspond to the degree-stratified components of the shadow obstruction tower:

| Categorical side                    | Chiral side                | Degree  |
|-------------------------------------|----------------------------|---------|
| $\mathrm{HH}_0(C)$ (CY class)       | $\kappa_{\mathrm{cat}}(C)$ | $r = 2$ |
| $\mathrm{HH}_1(C)$ (1st Hochschild) | Cubic shadow $C$           | $r = 3$ |
| $\mathrm{HH}_2(C)$ (2nd Hochschild) | Quartic resonance $Q$      | $r = 4$ |

The CY Hodge filtration degeneration (Kaledin) corresponds to the PBW spectral sequence collapse (Koszulness K2): both are manifestations of bar-cohomological formality (Koszulness, not SC formality) at different categorical levels.



# Chapter 5

## $E_1$ -Chiral Algebras

Braided output is too coarse for the first questions of Vol III. The quantum group, the Yangian, and the collision residue all live on an ordered  $E_1$  layer that remembers the direction of collisions. The CY-to-chiral functor  $\Phi$  reaches its braided  $E_2$  image only through that primitive step, so this chapter fixes the ordered conventions used in the rest of the volume.

*Remark 5.1* ( $E_1$  primacy for CY quantum groups). The  $E_1$ -chiral algebra (boundary) is the primitive object in this volume. The  $E_2$ -chiral algebra (bulk) is obtained from it by the Drinfeld center construction  $\mathcal{Z}(\text{Rep}^{E_1}(A)) \simeq \text{Rep}^{E_2}(\text{Drin}(A))$ . Quantum groups, Yangians, and braided tensor categories are natively  $E_1$  objects: the CoHA multiplication is ordered (short exact sequences have a preferred direction), and the  $R$ -matrix arises only in the Drinfeld double. The passage  $E_1 \rightarrow E_2$  is the higher-categorical analogue of the averaging map  $\text{av}: \mathfrak{g}^{E_1} \rightarrow \mathfrak{g}^{\text{mod}}$  from Vol I. This averaging is *lossy*:  $\text{av}(r(z)) = \kappa_{\text{ch}}$  forgets the  $R$ -matrix data. The  $E_1$ -bar  $B^{\text{ord}}(A)$  retains strictly more information than the  $E_\infty$ -bar  $B^\Sigma(A)$ .

### 5.1 Factorization algebras and the $E_n$ hierarchy

The first tension is immediate: a factorization algebra on a complex curve is topologically  $E_2$ , while the ordered bar and the Yangian are  $E_1$ . Holomorphy resolves the tension. Costello and Gwilliam [9] showed that the value on a small disk carries an  $E_n$ -algebra structure, and Lurie’s universal statement is  $\text{Fact}(\mathbb{R}^n) \simeq_{E_n}$ . For a complex curve  $C$  of complex dimension one,  $C$  has real dimension two, so factorization starts as  $E_2$  topologically before the holomorphic ordering picks an  $E_1$  slice.

**PROPOSITION 5.2** (*Holomorphic refinement, Costello-Gwilliam*). <sup>[PE]</sup> Let  $\mathcal{A}$  be a factorization algebra on a smooth complex curve  $C$  whose structure maps are holomorphic in the two-point configuration space. The associated  $E_2$ -algebra specializes to an  $E_1$ -algebra on the configuration space of ordered points along any real ray, with the ordering provided by the holomorphic argument.

The proof is operadic: the little two-disks operad contracts onto the little intervals operad along the real subspace, and holomorphy pins the contraction to a single slice. Beilinson and Drinfeld [1] studied the symmetric version of this specialization; they called the outcome a *chiral algebra*. Volume II elevates the ordered version.

*Definition 5.3* (*Swiss-cheese operad*). The Swiss-cheese operad  $\text{SC}^{\text{ch}, \text{top}}$  of Volume II is a *two-coloured* operad with colours  $\{\text{ch}, \text{top}\}$  (closed/holomorphic and open/topological). Its closed-to-closed operations are parametrised by  $\text{FM}_k(\mathbb{C})$  (the  $E_2$  operad); its open-to-open operations by  $E_1(\mathbb{R})$  (ordered configurations);

and its mixed operations (closed acting on open) by  $\mathrm{FM}_k(\mathbb{C}) \times E_1(m)$ . Open-to-closed operations are *empty* (directionality constraint). An  $\mathrm{SC}^{\mathrm{ch}, \mathrm{top}}$ -algebra is a pair  $(A, M)$  where  $A$  is an  $E_2$ -algebra (the closed colour),  $M$  is an  $E_1$ -algebra (the open colour), and  $A$  acts on  $M$  via the mixed operations.

$\mathrm{SC}^{\mathrm{ch}, \mathrm{top}}$  is *not* the tensor product  $E_1 \otimes E_2$ : the directionality constraint (no open-to-closed) and the mixed operations make it a genuinely two-coloured operad. Dunn additivity does not apply. The  $E_3$  upgrade requires a 3d holomorphic-topological theory (proved for Kac–Moody via holomorphic Chern–Simons; conjectural in general); the resulting  $E_3$  structure is *topological*, not chiral (the conformal vector at non-critical level kills the chiral direction via Sugawara; Volume I, the topologization theorem).

The closed colour carries  $E_2$  structure from  $\mathrm{FM}_k(\mathbb{C})$ : holomorphic, braided, and the home of the chiral algebra. The open colour carries  $E_1$  structure from ordered real configurations: topological, non-commutative, and the home of the ordered bar complex. Standard Beilinson–Drinfeld chiral algebras are  $E_\infty$ -chiral (symmetric OPE); the  $E_1$  open colour captures the additional ordering data. The  $\kappa_\bullet$ -spectrum of this volume distributes across the two colours:  $\kappa_{\mathrm{ch}}$  is read from the open colour (the ordered bar),  $\kappa_{\mathrm{cat}}$  from the closed colour (the Euler characteristic), and the Drinfeld center exchanges them.

*Remark 5.4* (Level-prefixed  $r$ -matrix). The classical  $r$ -matrix attached to an affine Kac–Moody  $E_1$ -chiral algebra at level  $k$  is

$$r(z) = \frac{k \Omega}{z},$$

not the level-stripped  $\frac{\Omega}{z}$ . The level survives the  $d \log$  absorption because the ordered bar complex builds in one factor of the level per collision. At  $k = 0$  the  $r$ -matrix vanishes identically. The collision residue of the Heisenberg  $r$ -matrix is  $k$ , not  $k/2$ , and the monodromy of the  $E_1$  representation category around a puncture is  $\exp(-2\pi i k)$ .

The level-prefix convention is the standing convention throughout this volume. Every time an  $r$ -matrix is written in the CY-geometric setting, the level constant prefactor is the image of the trace under the CY-to-chiral functor and must be tracked explicitly.

## 5.2 The ordered bar as the $E_1$ primitive

Braiding forgets collision order. The ordered bar keeps it. Let  $A$  be a chiral algebra on  $C$  with augmentation  $\varepsilon: A \rightarrow \Omega_C$  and augmentation ideal  $\bar{A} = \ker(\varepsilon)$ .

*Definition 5.5* (Ordered bar complex). The *ordered bar complex* of  $A$  is the cofree conilpotent tensor coalgebra

$$B^{\mathrm{ord}}(A) := T^c(s^{-1}\bar{A}) = \bigoplus_{n \geq 0} (s^{-1}\bar{A})^{\otimes n}$$

equipped with the *deconcatenation* coproduct

$$\Delta_{\mathrm{dec}}(a_1 \otimes \cdots \otimes a_n) = \sum_{k=0}^n (a_1 \otimes \cdots \otimes a_k) \otimes (a_{k+1} \otimes \cdots \otimes a_n)$$

and the bar differential built from the chiral product of  $A$ .

Deconcatenation is NOT the coshuffle coproduct on  $\mathrm{Sym}^c$ . Deconcatenation produces  $n + 1$  terms and respects the ordering; the coshuffle produces  $2^n$  terms and destroys it. The symmetric bar complex  $B^\Sigma(A)$  is the  $S_n$ -coinvariant quotient of  $B^{\mathrm{ord}}(A)$  with the coshuffle coproduct induced on invariants.

PROPOSITION 5.6 (*Averaging map*). <sup>[PE]</sup>The averaging map

$$\text{av}: B^{\text{ord}}(A) \longrightarrow B^{\Sigma}(A), \quad a_1 \otimes \cdots \otimes a_n \longmapsto \frac{1}{n!} \sum_{\sigma \in S_n} a_{\sigma(1)} \otimes \cdots \otimes a_{\sigma(n)}$$

is a morphism of cochain complexes and sends the  $E_1$  structure to the  $E_{\infty}$  structure. It is lossy: the kernel contains the  $R$ -matrix data of the holomorphic factor, and on degree two  $\text{av}(r(z)) = \kappa_{\text{ch}}$ .

Volume I establishes the map; Volume II identifies it as the  $E_1 \rightarrow E_{\infty}$  symmetrization. For Volume III purposes, the two consequences that matter are: (a) Yangians and quantum groups live on the  $E_1$  side and are quotiented by averaging; (b) the symmetric bar  $B^{\Sigma}$  is sufficient for computing the modular characteristic but insufficient for reconstructing the  $R$ -matrix.

Averaging the degree-two generator  $r(z)$  returns the scalar  $\kappa_{\text{ch}}$ , the unique  $S_2$ -coinvariant of the collision residue. When the same  $r(z)$  comes from the CY-to-chiral functor applied to  $D^b(\text{Coh}(K3 \times E))$ , the scalar is  $\kappa_{\text{ch}} = 3$  by additivity:  $\kappa_{\text{ch}}(K3) + \kappa_{\text{ch}}(E) = 2 + 1$ . This differs from the categorical Euler characteristic  $\kappa_{\text{cat}} = 2$ , the lattice-rank invariant  $\kappa_{\text{fiber}} = 24$ , and the BKM weight  $\kappa_{\text{BKM}} = 5$ . An unsubscripted symbol would conflate distinct invariants.

*Remark 5.7* (Three bars, one functor). The CY-to-chiral functor  $\Phi$  of Chapter 8 has three natural composites.  $\Phi_{E_1}$  produces  $B^{\text{ord}}(A_C)$ ; this is the ordered bar sensed by the Yangian.  $\text{av} \circ \Phi_{E_1}$  produces  $B^{\Sigma}(A_C)$ ; this is the symmetric bar sensed by the modular characteristic  $\kappa_{\text{ch}}$ . The full  $E_2$  enhancement  $\Phi_{E_2}$  produces the Drinfeld-centered bar  $B^{E_2}(A_C) = \mathcal{Z}(B^{\text{ord}}(A_C))$ ; this is the braided bar sensed by the quantum group. All three factor through  $\Phi_{E_1}$ . In this volume, the ordered bar is the primitive object.

### 5.3 $E_1$ -chiral algebras from CY categories

The CY input has to produce an ordered chiral object before any braided output can exist. For a  $\text{CY}_d$  category  $C$  in the sense of Kontsevich-Soibelman, the trace lives in  $\text{HC}_d^-(C)$ , not merely in Hochschild homology, and that trace feeds the ordered  $E_1$  structure first.

PROPOSITION 5.8 ( $E_1$  sector at  $d = 2$ ). <sup>[PH]</sup>Let  $C$  be a  $\text{CY}_2$  category with cyclic  $A_{\infty}$  structure and negative-cyclic trace. The ordered bar complex of the cyclic  $A_{\infty}$  algebra  $A_C$  carries a natural  $E_1$ -chiral structure whose holomorphic direction matches the Hochschild differential and whose ordered direction matches the cyclic action on the bar coalgebra.

*Proof.* The cyclic  $A_{\infty}$ -structure on  $C$  provides an associative product on the bar coalgebra  $B^{\text{ord}}(A_C) = T^c(s^{-1}\bar{A}_C)$  via OPE residues (Step 1 of the cyclic-to-chiral passage, Section 8.1 of Chapter 8). The holomorphic direction is the Hochschild differential  $b: \text{CC}_{\bullet}(C) \rightarrow \text{CC}_{\bullet-1}(C)$ , which descends to the bar differential  $d_{\text{bar}}$  on  $B^{\text{ord}}(A_C)$  via the standard identification of the bar construction with the Hochschild chain complex. The ordered direction is the  $S^1$ -action on the cyclic bar complex: the Connes  $B$ -operator cyclically permutes the bar entries, and its restriction to the ordered bar preserves the deconcatenation coproduct. The  $E_1$ -chiral structure is the factorization algebra on  $C$  obtained by the factorization envelope of the Lie conformal algebra  $\mathfrak{L}_C$  (Step 2), with the negative-cyclic trace in  $\text{HC}_2^-(C)$  providing the quantization datum (Step 4). At  $d = 2$ , no framing obstruction arises: the  $\mathbb{S}^2$ -framing is automatic (Kontsevich–Vlassopoulos). The full construction is Theorem 8.1 of Chapter 8.  $\square$

For  $C = D^b(\text{Coh}(K3))$ , the resulting  $K3$  chiral algebra has  $\kappa_{\text{ch}} = 2$ . The Borchers-side weight  $\kappa_{\text{BKM}} = 5$  belongs to the  $K3 \times E$  lift developed in Part III and must not be conflated with the  $K3$  value.

**CONJECTURE 5.9** ( $E_1$  sector at  $d = 3$ ). Let  $C$  be a  $CY_3$  category with a chain-level  $S^3$ -framing on  $HC_3^-(C)$  (the condition of Conjecture  $CY-A_3$ ). Then the ordered bar complex of the cyclic  $A_\infty$  algebra  $A_C$  carries an  $E_1$ -chiral structure whose representation category is a braided monoidal refinement of the BPS Yangian of  $C$ .

The conjecture is conditional on the chain-level framing; the unconditional statement requires the full  $CY-A_3$  programme. Every downstream result in III and IV that uses the BPS Yangian inherits this conditionality. The connection to Maulik-Okounkov and Costello-Witten cohomological Hall algebras is traced in Chapter 15.

**PROPOSITION 5.10** ( $E_2$  enhancement via Drinfeld center). <sup>[PE]</sup> If  $A$  is an  $E_1$ -chiral algebra with dualizable monoidal representation category  $\text{Rep}^{E_1}(A)$ , the Drinfeld center  $\mathcal{Z}(\text{Rep}^{E_1}(A))$  is a braided monoidal category equipped with an equivalence

$$\mathcal{Z}(\text{Rep}^{E_1}(A)) \simeq \text{Rep}^{E_2}(\text{Drin}(A))$$

where  $\text{Drin}(A)$  is the  $E_2$ -chiral algebra obtained by the categorified averaging map  $E_1\text{-Cat} \rightarrow E_2\text{-Cat}$ . This is a consequence of Lurie's identification of  $E_2$ -algebras with braided monoidal objects [25] and the Drinfeld center construction of Etingof–Nikshych–Ostrik.

This proposition provides the pathway  $\Phi_{E_1} \rightarrow \Phi_{E_2}$ : one computes the  $E_1$  chiral algebra first and then applies the Drinfeld center at the categorical level. At  $d = 2$  this produces the braided categories attached to the  $CY_2$  output of  $\Phi$ ; at  $d = 3$  the construction is conditional on Conjecture 5.9.

*Remark 5.11* (Convolution  $L_\infty$ -algebra on the  $E_1$  side). For an  $E_1$ -chiral algebra  $A$  and a coalgebra  $C$ , the convolution  $L_\infty$ -algebra  $\text{hom}_\alpha(C, A)$  of Volume I is the natural habitat of Maurer-Cartan elements controlling twisted morphisms. On the  $E_1$  side,  $\text{hom}_\alpha$  is a strict dg Lie algebra when the ordering of  $C$  is preserved (the strict model  $\text{str}$ ), and a genuine  $L_\infty$ -algebra when higher brackets are included ( $\infty$ ). The MC moduli of the two models coincide, but the full  $L_\infty$ -structure is required for homotopy transfer, formality statements, and gauge equivalence. For Vol III, the  $CY$ -to-chiral functor  $\Phi$  lands in the strict model  $\text{str}$  because the cyclic  $A_\infty$  structure on  $A_C$  is strict at degree two; higher-degree corrections require the  $L_\infty$  refinement.

*Remark 5.12* (Comparison with Beilinson-Drinfeld chiral algebras). Beilinson and Drinfeld [1] developed chiral algebras as a symmetric factorization formalism on the Ran space. Their chiral algebras lie on the  $E_\infty$  side of the locality hierarchy, even when they carry OPE poles. The  $E_1$ -chiral algebras of this chapter are a strict refinement: they remember collision order. The averaging map  $B^{\text{ord}} \rightarrow B^\Sigma$  forgets that extra data and returns to the Beilinson-Drinfeld world. Vol III's geometric output is ordered; its modular characteristic is symmetric.

*Remark 5.13* ( $E_1$ -chiral bialgebras versus  $E_\infty$ -vertex bialgebras). The  $E_n$ -hierarchy organises three bar complexes with three coalgebra structures, three Hopf structures, and three levels of quantum-group data. The comparison table is (cf. Construction 6.21 in Chapter 6 and Theorem 6.22 for the  $\mathbb{C}^3$  instance):

|                     | $E_1$ (labeled-ordered)               | $E_2$ (braided)                        | $E_\infty$ (symmetric)                          |
|---------------------|---------------------------------------|----------------------------------------|-------------------------------------------------|
| Bar complex         | $B^{\text{ord}} = T^c(s^{-1}\bar{A})$ | $B^{E_2}$ ( $B_n$ -action via $R(z)$ ) | $B^\Sigma = \text{Sym}^c(s^{-1}\bar{A})$        |
| Coproduct           | deconcatenation                       | braided deconcatenation                | coshuffle                                       |
| Terms at degree $n$ | $n + 1$                               | $B_n$ -orbits                          | $2^n$                                           |
| $R$ -matrix data    | full $r(z)$                           | braiding $\sigma$                      | none ( $\text{av}(r(z)) = \kappa_{\text{ch}}$ ) |
| Koszul dual         | $A^!$ (defect algebra)                | $A^!$ with $\sigma^{\text{rev}}$       | classical Koszul                                |
| Hopf structure      | $E_1$ -chiral bialgebra               | braided Hopf (Majid)                   | Li vertex bialgebra                             |
| Coproduct type      | Drinfeld $\Delta_z$                   | braided $\Delta_\sigma$                | symmetric $\Delta$                              |

*What is lost at each level.*

- (i)  $E_1 \rightarrow E_2$ : the labeled ordering is replaced by a braid-group action; half-braiding data (the choice of over- versus under-crossing) is forgotten.
- (ii)  $E_2 \rightarrow E_\infty$ : the braiding  $\sigma$  becomes symmetric; the  $R$ -matrix collapses to the scalar  $\kappa_{\text{ch}}$ . Every non-abelian quantum group datum is lost.
- (iii)  $E_1 \rightarrow E_\infty$  (direct, via  $\text{av}$ ): the full  $R$ -matrix, the Hopf structure, and the Drinfeld coproduct all vanish. What survives is the modular characteristic  $\kappa_{\text{ch}}$  and the shadow tower.

*Why Vol III works at  $E_1$ .* The entire quantum group programme (Yangians  $Y(\mathfrak{g})$ , quantum toroidal algebras  $U_{q,t}(\mathfrak{gl}_1)$ , Kazhdan–Lusztig categories  $\text{Rep}_q(\mathfrak{g})$  at roots of unity) lives natively at  $E_1$ . The passage to  $E_2$  is via the Drinfeld center  $\mathcal{Z}(\text{Rep}^{E_1}(A)) \simeq \text{Rep}^{E_2}(\text{Drin}(A))$  (Proposition 5.10); the passage to  $E_\infty$  forgets the quantum group entirely and returns to the classical vertex-algebraic world of Beilinson–Drinfeld. Li’s vertex-bialgebra framework is the  $E_\infty$  shadow of the  $E_1$ -chiral bialgebra: it retains the vertex-algebraic OPE structure but symmetrizes the coproduct, losing the  $R$ -matrix that distinguishes  $U_q(\mathfrak{g})$  from  $U(\mathfrak{g})$ . The chiral quantum group is the  $E_1$  substance; the vertex bialgebra is the  $E_\infty$  silhouette.

**PROPOSITION 5.14** (*Lossiness of the  $E_1 \rightarrow E_\infty$  forgetful functor on bialgebras*). <sup>[PH]</sup> Let  $(A, \mu, \Delta_z, R(z))$  be an  $E_1$ -chiral bialgebra with  $R$ -matrix  $R(z)$  and Drinfeld coproduct  $\Delta_z$ . The forgetful functor

$$\text{Forget}_{E_1 \rightarrow E_\infty} : E_1\text{-ChirBialg} \longrightarrow E_\infty\text{-VtxBialg}$$

sends  $(A, \mu, \Delta_z, R(z))$  to the Li vertex bialgebra  $(A, \mu, \text{av}(\Delta_z), 1)$  with *trivial* braiding. In particular:

- (a) The image has symmetric coproduct:  $\text{av}(\Delta_z) = \tau \circ \text{av}(\Delta_z)$ .
- (b) The  $R$ -matrix reduces to the scalar  $\text{av}(R(z))|_{z=0} = \kappa_{\text{ch}} \cdot \text{id}$ .
- (c) The braiding is *not* recoverable from the  $E_\infty$  image. It is recovered precisely by the Drinfeld center:  $\mathcal{Z}(\text{Rep}^{E_1}(A)) \simeq \text{Rep}^{E_2}(\text{Drin}(A))$ .

*Proof.* Claim (a): the averaging map  $\text{av} : B^{\text{ord}} \rightarrow B^\Sigma$  (Proposition 5.6) is  $S_n$ -coinvariant projection, so the image is  $S_n$ -invariant by construction; in particular the coproduct is symmetric. Claim (b): the degree-two component of  $\text{av}$  sends  $r(z) = R_{12}(z)$  to its  $S_2$ -coinvariant, which is the scalar  $\kappa_{\text{ch}}$  (loc. cit.). Claim (c): the fiber of  $\text{av}$  in degree two is  $\ker(\text{av}|_2) = \{f(z) : f(z) + f(-z) = 0\}$ , the space of antisymmetric collision data. The  $R$ -matrix lives in that kernel (it satisfies  $R_{12}(z) = -R_{21}(-z)$  by the Yang–Baxter skew symmetry). Hence  $R(z)$  is annihilated and cannot be recovered from the  $E_\infty$  image. By Proposition 5.10, the Drinfeld center reconstructs the braiding from the monoidal structure of  $\text{Rep}^{E_1}(A)$ ; this is the unique functorial recovery.  $\square$

## 5.4 Connection to Volume II

Volume II resolves the ordered side of the programme. Its seven parts develop the Swiss-cheese operad, the ordered bar complex, the classical  $r(z)$ -matrix at level  $k$ , the seven faces of  $r(z)$ , and the derived center of an  $E_1$ -chiral algebra. Vol III uses that machinery through the CY interface fixed here.

**PROPOSITION 5.15** (*Vol III composes with Vol II*). <sup>[PH]</sup> The CY-to-chiral functor  $\Phi$  of I factors as  $\Phi = \Phi_{E_1}^{\text{Vol II}} \circ \Phi_{\text{cyc}}^{\text{Vol III}}$ , where  $\Phi_{\text{cyc}}^{\text{Vol III}}$  takes a  $\text{CY}_d$  category to its cyclic  $A_\infty$  algebra and  $\Phi_{E_1}^{\text{Vol II}}$  takes a cyclic  $A_\infty$  algebra to an  $E_1$ -chiral algebra via the Swiss-cheese promotion.

*Proof.* The factorization is the decomposition of the four-step cyclic-to-chiral passage of Section 8.1 (Chapter 8) into two stages. The first stage  $\Phi_{\text{cyc}}^{\text{Vol III}}$  performs Steps 1–2: it takes a  $\text{CY}_d$  category  $\mathcal{C}$  with its cyclic  $A_\infty$ -structure and CY trace, extracts the Gerstenhaber bracket on  $\text{HH}^\bullet(\mathcal{C})$  to form the Lie conformal algebra  $\mathfrak{L}_{\mathcal{C}}$  (Step 1), and applies the factorization envelope to produce the factorization algebra  $\text{Fact}_X(\mathfrak{L}_{\mathcal{C}})$  on a curve  $X$  (Step 2). The output is a cyclic  $A_\infty$ -algebra with the data needed for the Vol II machine. The second stage  $\Phi_{E_1}^{\text{Vol II}}$  performs Steps 3–4: it takes the cyclic  $A_\infty$ -algebra, applies the  $\mathbb{S}^d$ -framing to obtain the  $E_n$ -enhancement (Step 3), and quantizes using the CY trace (Step 4), producing the  $E_1$ -chiral algebra  $A_{\mathcal{C}}$ . The Swiss-cheese promotion of Vol II equips this algebra with the two-coloured  $\text{SC}^{\text{ch}, \text{top}}$ -structure (Definition 5.3). The composition  $\Phi_{E_1}^{\text{Vol II}} \circ \Phi_{\text{cyc}}^{\text{Vol III}}$  recovers  $\Phi$  by construction.  $\square$

The detailed operadic content of  $\Phi_{E_1}^{\text{Vol II}}$  involves the three coalgebra structures, the difference between coshuffle and deconcatenation, the promotion from one-colour to two-colour, the mixed-sector dimension formula, the curved factor of two at positive genus, the averaging map lossiness, the bound on  $\text{ChirHoch}^*(\mathcal{C})$ , and the distinction between generating depth and algebraic depth.

*Remark 5.16* (Why the  $E_1$  layer cannot be skipped). The CY geometric data can be packaged directly into a braided factorization algebra (the  $E_2$  form) via Kontsevich-Soibelman COHA technology. The temptation is to skip the  $E_1$  intermediate and go from  $\mathcal{C}$  straight to the braided output. Three obstructions block that route. First, the convolution  $L_\infty$ -algebra  $\text{hom}_\alpha(\mathcal{C}, A)$  of Volume I fails to be a bifunctor in both slots simultaneously (RNW19);  $\text{MC}_3$  must be proved one slot at a time, and the slot that admits a proof is the  $E_1$  slot. Second, the  $r$ -matrix  $r_{\text{CY}}(z)$  cannot be extracted without seeing the ordered collisions, since the braiding of the  $E_2$  picture only remembers  $r(z)$  up to  $S_2$ -coinvariance. Third, the CoHA itself is an  $E_1$ -associative multiplication; writing it as an  $E_2$  product forgets the preferred direction on short exact sequences that is central to wall-crossing. These three obstructions are the same obstruction seen from three angles: the ordered bar is the only model that remembers the direction.

*Remark 5.17* (lambda-bracket convention in Vol III). Vol III writes classical shadow operations in lambda-bracket notation with divided powers:  $\{T_\lambda T\} = (c/12) \lambda^3$ . The divided-power prefactor  $1/3! = 1/6$  absorbs the OPE mode coefficient into the lambda-bracket rewrite: starting from the OPE mode  $T_{(3)}T$  and dividing by  $3!$  yields the stated  $c/12$  at order  $\lambda^3$ . Every formula imported from Vol I (which uses OPE mode notation) must be converted before appearing in Vol III. The CY-to-chiral functor  $\Phi$  is agnostic to the choice of convention, but its computed values of  $\kappa_{\text{ch}}$  are convention-dependent at the level of integral prefactors, and a Vol I formula transplanted without conversion will produce a wrong  $\kappa_{\text{ch}}$  by exactly a factor of 6.

*Remark 5.18* (Three-volume thesis). The three volumes are three faces of a single  $E_1$ - $E_1$  operadic Koszul duality. Volume I is the symmetric modular face: it develops  $B^\Sigma$ , the five theorems A-D+H, and the modular characteristic  $\kappa_{\text{ch}}$  in the uniform-weight setting. Volume II is the  $E_1$  open-colour face: it develops  $B^{\text{ord}}$ , the Swiss-cheese operad, the  $r(z)$ -matrix with its seven faces, and the three-dimensional holomorphic-topological bridge to quantum gravity. Volume III is the CY-geometric face: it develops the functor  $\Phi$  that produces the input algebra from a Calabi-Yau category, identifies  $\kappa_{\text{ch}}$  within the kappa-spectrum, and traces the quantum group back to its geometric origin in BPS state counts.

*Remark 5.19* (How this chapter is used downstream). Chapters 12, 13, and 19 use this chapter as the source of the ordered bar coalgebra and the averaging map. Chapter 8 uses Proposition 5.8 as the  $d = 2$  existence statement for  $A_{\mathcal{C}}$ . Every braided monoidal category that appears later is the Drinfeld center of an underlying ordered one.

## 5.5 Operadic dictionary for the CY reader

The operadic vocabulary of Volume II can be compressed into the following dictionary, which the CY reader can keep on a single page.



- $E_n$ : little  $n$ -disks operad.  $E_1$  is the operad of intervals on the line;  $E_2$  is the operad of disks in the plane;  $E_\infty$  is the colimit.  $E_1$  algebras are ordered and associative.  $E_2$  algebras are partially commutative and braided.  $E_\infty$  algebras are symmetric up to all higher homotopies; ordinary chiral algebras live here and still admit OPE poles. Dunn additivity:  $E_m \otimes E_n \simeq E_{m+n}$ .
- **Ordered bar**  $B^{\text{ord}}$ : the cofree conilpotent tensor coalgebra  $T^c(s^{-1}\bar{A})$  with deconcatenation coproduct. Retains degree ordering. Natural  $E_1$ -object. Source of Yangians and quantum groups.
- **Symmetric bar**  $B^\Sigma$ : the  $S_n$ -coinvariant quotient of  $B^{\text{ord}}$  with the coshuffle coproduct descended from the shuffle product on  $T^c$ . Natural  $E_\infty$ -object. Source of the modular characteristic  $\kappa_{\text{ch}}$ .
- **Harrison bar**  $B^{\text{Lie}}$ : the Harrison complex; Lie-coalgebra structure via the iterated commutator. Dual to the Chevalley-Eilenberg complex of the associated Lie algebra. Smallest of the three natural bar complexes on any commutative input.
- **Deconcatenation**: coproduct on  $T^c$  that splits a word into a prefix and a suffix. Produces  $n + 1$  terms in degree  $n$ . Respects ordering. The  $E_1$  coproduct.
- **Coshuffle**: coproduct on  $\text{Sym}^c$  dual to the shuffle product. Produces up to  $2^n$  terms. Destroys ordering. The  $E_\infty$  coproduct.
- **Swiss-cheese**  $\text{SC}^{\text{ch}, \text{top}}$ : two-coloured operad combining a chiral (holomorphic, closed) sector with an ordered (topological, open) sector. Product decomposition  $E_1(C) \times E_2(\text{top})$  isolates the  $E_1$  chiral factor.
- **Drinfeld center**  $\mathcal{Z}$ : endofunctor on monoidal categories that sends a monoidal category  $\mathcal{M}$  to the category of objects equipped with a half-braiding against every object of  $\mathcal{M}$ . The result is braided monoidal. Categorified averaging:  $\mathcal{Z}$  takes  $E_1$ -categories to  $E_2$ -categories.
- **$r$ -matrix**: the degree-two generator of the  $E_1$  Koszul duality, carrying a pole along the diagonal. At level  $k$ , the classical Kac-Moody  $r$ -matrix is  $r(z) = \frac{k\Omega}{z}$  and vanishes at  $k = 0$ .

The dictionary is not symmetric in  $E_1$  and  $E_\infty$ : the ordered bar is the primitive, and everything on the  $E_\infty$  side is a quotient. The CY-geometric functor  $\Phi$  lands primitively in the  $E_1$  layer; the  $E_\infty$  and  $E_2$  images are obtained by composing with averaging and Drinfeld center respectively.

*Remark 5.20* (Notation consistency across the three volumes). The three volumes use different coalgebra conventions in their displayed formulas. Vol I displays formulas in  $B^\Sigma$  with the symmetric coproduct; Vol II displays formulas in  $B^{\text{ord}}$  with deconcatenation; Vol III displays formulas in whichever form the CY functor produces, which is always  $B^{\text{ord}}$  at the source but frequently  $B^\Sigma$  after averaging to extract  $\kappa_{\text{ch}}$ . The reader who cross-references a formula between volumes must convert between the three coalgebra structures:  $B^{\text{ord}} \rightarrow B^\Sigma$  by averaging (dividing by  $n!$  and symmetrizing), and  $B^\Sigma \rightarrow B^{\text{Lie}}$  by taking the iterated commutator of the cofree tensor coalgebra. The three bars are NOT isomorphic even as complexes; they differ by  $S_n$ -coinvariant quotients, and the Euler characters diverge accordingly. See Vol II for the three-bar sequence.

## 5.6 $E_1$ -chiral Koszul duality: three standard families

Volume II proves  $E_1$ -chiral Koszul duality: the bar-cobar adjunction  $B^{\text{ord}} \dashv \Omega^{\text{ord}}$  is a Quillen equivalence on the Koszul locus, and the Koszul dual algebra  $A^\dagger$  satisfies  $\text{Rep}^{E_1}(A) \simeq \text{Rep}^{E_1}(A^\dagger)^{\text{rev}}$  as monoidal

categories. The Koszul conductor  $\rho_K(A)$  is the real number such that

$$\kappa_{\text{ch}}(A) + \kappa_{\text{ch}}(A^\dagger) = \rho_K(A). \quad (5.1)$$

The conductor  $\rho_K$  is a family invariant: it depends on the isomorphism class of the shadow tower, not on the particular algebra within the family. This section computes  $A^\dagger$ ,  $B^{\text{ord}}$ , and  $\rho_K$  explicitly for the three standard families that span the G/L/M classification: Heisenberg (class G,  $\rho_K = 0$ ), affine Kac–Moody  $V_k(\mathfrak{sl}_2)$  (class L,  $\rho_K = 0$ ), and Virasoro  $c$  (class M,  $\rho_K = 13$ ).

### Family I: Heisenberg $H_k$ (class G)

The rank-one Heisenberg vertex algebra  $H_k$  at level  $k \neq 0$  has a single strong generator  $J$  of conformal weight 1, with OPE  $J(z)J(w) \sim k/(z-w)^2$  and no nonlinear terms. The augmentation ideal  $\bar{H}_k = \langle J, \partial J, \partial^2 J, \dots \rangle$  is spanned by all derivatives, and  $\kappa_{\text{ch}}(H_k) = k$ .

**PROPOSITION 5.21** ( *$E_1$ -Koszul dual of the Heisenberg*). <sup>[PH]</sup> The  $E_1$ -chiral Koszul dual of  $H_k$  is  $H_k^\dagger = (\text{Sym}^{\text{ch}}(V^*), m_0 = -k\omega)$ , the curved commutative chiral algebra on the dual generator space (Vol I, §5.6; *not*  $H_{-k}$ , which has the same  $\kappa_{\text{ch}} = -k$  but is a Heisenberg Lie-type algebra, not commutative). The ordered bar complex  $B^{\text{ord}}(H_k) = T^c(s^{-1}\bar{H}_k)$  has trivial differential at all degrees, and the Koszul conductor is  $\rho_K = 0$ .

*Proof.* The bar differential  $d_{\text{bar}}$  on  $B_n^{\text{ord}}(H_k) = (s^{-1}\bar{H}_k)^{\otimes n}$  acts by summing adjacent OPE multiplications:

$$d_{\text{bar}}[a_1 | \dots | a_n] = \sum_{i=1}^{n-1} \pm [a_1 | \dots | \mu(a_i, a_{i+1}) | \dots | a_n].$$

For the Heisenberg,  $\mu(J, J) = 0$  (the singular OPE  $J(z)J(w) \sim k/(z-w)^2$  contributes to the metric/pairing, not to the multiplication  $\mu$ : the OPE has no first-order pole, so the normally-ordered product  $:JJ:$  has no correction from the singular part). Consequently  $d_{\text{bar}} = 0$  at all degrees. The bar complex is exact in the trivial sense:  $H^*(B^{\text{ord}}(H_k)) = B^{\text{ord}}(H_k)$  as a graded vector space.

The Koszul dual is then read from the linear dual of  $B^{\text{ord}}$ : the Verdier dual  $D_{\text{Ran}}(B^{\text{ord}}(H_k))$  is a co-commutative coalgebra on  $(s^{-1}\bar{H}_k)^*$  (cocommutative because the double-pole OPE produces symmetric coproducts). By Milnor–Moore, the cobar is the chiral symmetric algebra  $H_k^\dagger = \text{Sym}^{\text{ch}}(V^*)$ , equipped with the curvature  $m_0 = -k\omega$  encoding the level. The dual generator  $J^*$  has  $\kappa_{\text{ch}}(H_k^\dagger) = -k$  (the same numerical value as  $\kappa_{\text{ch}}(H_{-k})$ , but  $H_k^\dagger$  is commutative while  $H_{-k}$  is Lie-type; see Vol I, §5.6).

The conductor computation (for the Heisenberg/free-field family,  $\rho_K = 0$ ):

$$\kappa_{\text{ch}}(H_k) + \kappa_{\text{ch}}(H_k^\dagger) = k + (-k) = 0 = \rho_K.$$

□

The explicit bar complex at low degrees is:

$$B_1^{\text{ord}}(H_k) = s^{-1}\bar{H}_k = \langle [J], [\partial J], [\partial^2 J], \dots \rangle, \quad (5.2)$$

$$B_2^{\text{ord}}(H_k) = (s^{-1}\bar{H}_k)^{\otimes 2} = \langle [J|J], [J|\partial J], [\partial J|J], \dots \rangle, \quad d_{\text{bar}} = 0, \quad (5.3)$$

$$B_3^{\text{ord}}(H_k) = (s^{-1}\bar{H}_k)^{\otimes 3} = \langle [J|J|J], [J|J|\partial J], \dots \rangle, \quad d_{\text{bar}} = 0. \quad (5.4)$$

The dimension at degree  $n$  (truncated to conformal weight  $\leq N$ ) is  $\dim B_n^{\text{ord}} = p(N)^n$  where  $p(N)$  is the partition function (one basis vector per derivative  $\partial^m J$ ,  $m \geq 0$ , organized by conformal weight  $m+1$ ). The

generating function  $\sum_n \dim B_n^{\text{ord}} \cdot q^n$  at fixed weight truncation is a geometric series in  $p(N)$ , reflecting the triviality of the bar differential (class G: the shadow tower terminates at depth 2).

*Remark 5.22* (Self-duality at  $k = 1$ ). At  $k = 1$  the Koszul dual is  $H_{-1}$ , which is isomorphic to  $H_1$  after reversing the ordering of the bar (the level sign is absorbed by the rev functor on  $\text{Rep}^{E_1}$ ). At general  $k$ , the duality  $H_k \leftrightarrow H_{-k}$  is the chiral reflection of the parameter inversion  $(h_1, h_2, h_3) \rightarrow (-h_1, -h_2, -h_3)$  of the  $\Omega$ -background.

## Family II: Kac–Moody $V_k(\mathfrak{sl}_2)$ (class L)

The affine Kac–Moody vertex algebra  $V_k(\mathfrak{sl}_2)$  at non-critical level  $k \neq -2$  has three strong generators  $\{e, f, h\}$  of conformal weight 1 with OPE:

$$h(z)h(w) \sim \frac{2k}{(z-w)^2}, \quad h(z)e(w) \sim \frac{2e(w)}{z-w}, \quad e(z)f(w) \sim \frac{k}{(z-w)^2} + \frac{h(w)}{z-w}. \quad (5.5)$$

The critical fact is the *first-order pole* in  $e(z)f(w)$ : the OPE multiplication  $\mu(e, f) = h$  is nontrivial, so the bar differential is nonzero. The modular characteristic is  $\kappa_{\text{ch}}(V_k(\mathfrak{sl}_2)) = 3(k+2)/4$  (from the general formula  $\kappa_{\text{ch}}(V_k(\mathfrak{g})) = \dim(\mathfrak{g})(k+h^\vee)/(2h^\vee)$  with  $\dim(\mathfrak{sl}_2) = 3, h^\vee = 2$ ).

**PROPOSITION 5.23** ( $E_1$ -Koszul dual of  $V_k(\mathfrak{sl}_2)$ ). <sup>[PH]</sup> The  $E_1$ -chiral Koszul dual of  $V_k(\mathfrak{sl}_2)$  at non-critical level  $k \neq -2$  is  $V_{k'}(\mathfrak{sl}_2)$  at the Feigin–Frenkel dual level  $k' = -k - 2h^\vee = -k - 4$ . The Koszul conductor is  $\rho_K = 0$ .

*Proof.* The argument has three steps.

*Step 1: Bar differential.* The ordered bar complex has  $B_n^{\text{ord}}(V_k(\mathfrak{sl}_2)) = (s^{-1}\bar{V}_k)^{\otimes n}$ , where  $\bar{V}_k$  is spanned by  $\{e, f, h, \partial e, \partial f, \partial h, \dots\}$ . The bar differential is

$$d_{\text{bar}}[a_1 | \dots | a_n] = \sum_{i=1}^{n-1} \pm [a_1 | \dots | \mu(a_i, a_{i+1}) | \dots | a_n]$$

where  $\mu$  extracts the first-order pole of the OPE. At degree 2:

$$\begin{aligned} d_{\text{bar}}[e|f] &= +[h], & d_{\text{bar}}[f|e] &= -[h], \\ d_{\text{bar}}[h|e] &= +[2e], & d_{\text{bar}}[e|h] &= -[2e], \\ d_{\text{bar}}[h|f] &= -[2f], & d_{\text{bar}}[f|h] &= +[2f], \\ d_{\text{bar}}[e|e] &= 0, & d_{\text{bar}}[f|f] &= 0, & d_{\text{bar}}[h|h] &= 0. \end{aligned}$$

Generators  $[e|e], [f|f], [h|h]$  are bar cycles (diagonal terms have no first-order pole), while off-diagonal terms produce nontrivial boundaries. The bar cohomology  $H^1(B^{\text{ord}}(V_k)) = \bar{V}_k$  (the generators) and the image of  $d_{\text{bar}}: B_2^{\text{ord}} \rightarrow B_1^{\text{ord}}$  is 3-dimensional (spanned by  $h, 2e, 2f$ ).

*Step 2: Koszul dual OPE.* The Verdier dual of  $B^{\text{ord}}(V_k(\mathfrak{sl}_2))$  produces dual generators  $\{e^*, f^*, h^*\}$  with the *same*  $\mathfrak{sl}_2$  structure but inverted level. The key identity is the Feigin–Frenkel duality: the functor  $V_k(\mathfrak{g}) \mapsto V_{-k-2h^\vee}(\mathfrak{g})$  is the chiral level reflection, and its operadic implementation is exactly  $E_1$ -Koszul duality applied to the bar complex. For  $\mathfrak{sl}_2$ :

$$k' = -k - 2h^\vee = -k - 4.$$

The dual generators satisfy  $e^*(z)f^*(w) \sim k'/(z-w)^2 + h^*(w)/(z-w)$  with  $k' = -k - 4$ , reproducing the  $V_{k'}(\mathfrak{sl}_2)$  OPE.

*Step 3: Conductor.*

$$\begin{aligned}\kappa_{\text{ch}}(V_k) + \kappa_{\text{ch}}(V_{k'}) &= \frac{3(k+2)}{4} + \frac{3(k'+2)}{4} = \frac{3(k+2)}{4} + \frac{3(-k-4+2)}{4} \\ &= \frac{3(k+2)}{4} + \frac{3(-k-2)}{4} = \frac{3(k+2) - 3(k+2)}{4} = 0 = \rho_K.\end{aligned}$$

□

The low-degree bar complex is:

$$\begin{aligned}B_1^{\text{ord}} &= s^{-1}\bar{V}_k = \langle [e], [f], [h], [\partial e], [\partial f], [\partial h], \dots \rangle, \\ B_2^{\text{ord}} &= (s^{-1}\bar{V}_k)^{\otimes 2}, \quad \dim = 9 \text{ at weight } 1,\end{aligned}\tag{5.6}$$

$$d_{\text{bar}}([e|f]) = [h], \quad d_{\text{bar}}([h|e]) = [2e], \quad d_{\text{bar}}([h|f]) = [-2f],$$

$$B_3^{\text{ord}} = (s^{-1}\bar{V}_k)^{\otimes 3}, \quad \dim = 27 \text{ at weight } 1.\tag{5.7}$$

At degree 3 the bar differential has two summands (positions  $i = 1, 2$ ), and the bar cohomology carries the first nontrivial  $A_\infty$  operation  $m_3$  of  $V_k(\mathfrak{sl}_2)$ . The nonvanishing of  $m_3$  is equivalent to the non-formality of  $\mathfrak{sl}_2$  as a dg Lie algebra and places  $V_k(\mathfrak{sl}_2)$  in shadow class L (depth 2 tower with nontrivial  $m_3$ ).

*Remark 5.24* (Critical level  $k = -2$ ). At  $k = -2$  the modular characteristic  $\kappa_{\text{ch}}(V_{-2}(\mathfrak{sl}_2)) = 0$  and the Koszul dual level is  $k' = -(-2) - 4 = -2$ : the critical level is a fixed point of the Feigin–Frenkel involution. The critical-level vertex algebra  $V_{-2}(\mathfrak{sl}_2)$  is the home of geometric Langlands (see Chapter 23), and the self-duality under  $E_1$  Koszul reflects the Langlands self-duality of  $\mathfrak{sl}_2$ .

*Remark 5.25* (General  $\mathfrak{g}$ ). For a simple Lie algebra  $\mathfrak{g}$  with dual Coxeter number  $h^\vee$ , the same argument gives  $V_k(\mathfrak{g})^\dagger = V_{-k-2h^\vee}(\mathfrak{g})$  with conductor

$$\kappa_{\text{ch}}(V_k) + \kappa_{\text{ch}}(V_{k'}) = \frac{\dim \mathfrak{g}}{2h^\vee} [(k + h^\vee) + (-k - 2h^\vee + h^\vee)] = 0.$$

The vanishing  $\rho_K = 0$  is universal for the Kac–Moody family.

### Family III: Virasoro $_c$ (class M)

The Virasoro vertex algebra  $_c$  at central charge  $c$  has a single strong generator  $T$  (the stress-energy tensor) of conformal weight 2 with OPE

$$T(z)T(w) \sim \frac{c/2}{(z-w)^4} + \frac{2T(w)}{(z-w)^2} + \frac{\partial T(w)}{z-w}.\tag{5.8}$$

The first-order pole gives  $\mu(T, T) = \partial T$  (a nontrivial multiplication), the second-order pole gives the conformal weight assignment (Sugawara), and the fourth-order pole gives the central extension. The modular characteristic is  $\kappa_{\text{ch}}(c) = c/2$  (from  $\dim B_n^{\text{ord}} \sim p(n)$  and the Volume I shadow formula).

The Koszul duality of the Virasoro differs from the Heisenberg and Kac–Moody families in a fundamental way: the conductor is *nonzero*.

**CONJECTURE 5.26** ( $E_1$ -Koszul dual of the Virasoro). <sup>[CJ]</sup> The  $E_1$ -chiral Koszul dual of  $_c$  is  $_{c'}$  at central charge  $c' = 26 - c$ . The Koszul conductor is  $\rho_K = 13$ :

$$\kappa_{\text{ch}}(c) + \kappa_{\text{ch}}(c') = \frac{c}{2} + \frac{26 - c}{2} = 13 = \rho_K.$$

At the self-dual point  $c = 13$  (the unique fixed point of  $c \mapsto 26 - c$ ),  $_{13}$  is Koszul self-dual under  $c \mapsto 26 - c$ . At  $c = 26$  the dual is  $_0$ , and at  $c = 0$  the dual is  $_{26}$  (the bosonic string).

The conjecture status is necessary because the full  $E_1$ -chiral Koszul duality for the Virasoro requires controlling the bar differential to all arities, and the infinite shadow depth (class M) means the tower never terminates. The evidence is:

*Evidence.* (i) *Bar differential at degree 2.* Using the first-order pole  $\mu(T, T) = \partial T$ :

$$d_{\text{bar}}[T|T] = [\partial T].$$

The image  $\text{im}(d_{\text{bar}} : B_2^{\text{ord}} \rightarrow B_1^{\text{ord}})$  is 1-dimensional (spanned by  $[\partial T]$ ). The bar cycle  $[T|T] - [T|T]$  is trivial, so  $\ker(d_{\text{bar}}|_{B_2^{\text{ord}} \rightarrow B_1^{\text{ord}}}) = \langle [T|T] + [T|T] \rangle_{/\text{im}}$  is computed by the commutator  $[T|T] - [T|T] = 0$  (the ordered bar is noncommutative; only the symmetric  $[T|T] + [T|T]$  survives modulo  $d$ ).

(ii) *Bar differential at degree 3.*  $d_{\text{bar}}[T|T|T] = [\partial T|T] \pm [T|\partial T]$  (two summands from positions  $i = 1, 2$ ). The sign is determined by the Koszul convention (desuspension shifts degree by  $-1$ ):

$$d_{\text{bar}}[T|T|T] = [\partial T|T] - [T|\partial T].$$

This is a nonzero cycle in  $B_2^{\text{ord}}$  (its image under  $d_{\text{bar}} : B_2^{\text{ord}} \rightarrow B_1^{\text{ord}}$  is  $[\partial^2 T] - [\partial^2 T] = 0$ , so it is indeed a cycle), and it represents the  $A_\infty$  operation  $m_3(T, T, T)$ . The nonvanishing of this class at all orders is the hallmark of class M (infinite shadow depth).

(iii) *Conductor from the Sugawara construction.* The Virasoro at central charge  $c$  embeds into the Heisenberg via Sugawara:  $T = JJ :/(2k)$  at  $c = 1$ . Under Koszul duality, the level inverts ( $k \rightarrow -k$ ), so  $c \rightarrow c'$  with  $c + c' = 26$  determined by the ghost central charge of the bosonic string ( $bc$ -ghost system contributes  $c = -26$ , so the BRST-balanced point is  $c + c_{\text{ghost}} = 26$ ; equivalently,  $c' = 26 - c$  is the unique value such that  $c \otimes_{c'}$  has total  $\kappa_{\text{ch}} = 13$ , the value at which the Virasoro shadow tower of Vol I attains its self-dual fixed point).

(iv) *Kappa verification.*  $\kappa_{\text{ch}}(c) = c/2$  and  $\kappa_{\text{ch}}(26-c) = (26-c)/2$ , so

$$\kappa_{\text{ch}}(c) + \kappa_{\text{ch}}(26-c) = \frac{c}{2} + \frac{26-c}{2} = 13 = \rho_K.$$

The conductor  $\rho_K = 13$  is nonzero, reflecting the nontrivial conformal anomaly of the Virasoro. Compare with the Heisenberg ( $\rho_K = 0$ ) and Kac–Moody ( $\rho_K = 0$ ) families.  $\square$

The low-degree bar complex:

$$B_1^{\text{ord}}(c) = s^{-1}_c = \langle [T], [\partial T], [\partial^2 T], \dots \rangle, \quad (5.9)$$

$$B_2^{\text{ord}}(c) = (s^{-1}_c)^{\otimes 2}, \quad d_{\text{bar}}[T|T] = [\partial T], \quad (5.10)$$

$$B_3^{\text{ord}}(c) = (s^{-1}_c)^{\otimes 3}, \quad d_{\text{bar}}[T|T|T] = [\partial T|T] - [T|\partial T]. \quad (5.11)$$

*Remark 5.27* (The conductor and the G/L/M classification). The three families exhibit a clean correlation between shadow class and Koszul conductor:

| Family                 | Class                      | $\kappa_{\text{ch}}(A)$ | $\kappa_{\text{ch}}(A^\dagger)$ | $\rho_K$ |
|------------------------|----------------------------|-------------------------|---------------------------------|----------|
| $H_k$                  | G (depth 2)                | $k$                     | $-k$                            | 0        |
| $V_k(\mathfrak{sl}_2)$ | L (depth 2, $m_3 \neq 0$ ) | $\frac{3(k+2)}{4}$      | $-\frac{3(k+2)}{4}$             | 0        |
| $c$                    | M (infinite depth)         | $\frac{c}{2}$           | $\frac{26-c}{2}$                | 13       |

The pattern:  $\rho_K = 0$  for classes G and L (finite shadow depth, or finite-depth  $A_\infty$  operations), and  $\rho_K > 0$  for class M (infinite shadow depth). The nonzero conductor is the signature of the conformal anomaly:  $\rho_K = 13$  is one-half of the critical dimension  $c = 26$ , and the shadow tower never terminates because each order contributes a new independent correction to the Koszul sum.

*Remark 5.28* (Compute engine verification). The kappa complementarity  $\kappa_{\text{ch}}(A) + \kappa_{\text{ch}}(A^\dagger) = \rho_K$  for all three families is verified by the compute engines: `holomorphic_cs_chiral_engine.py` (Heisenberg,  $\rho_K = 0$ ), `e2_bar_complex.py` (Kac–Moody,  $k' = -k - 4$ ), and `entropy_koszul_complement_cy3.py` (Virasoro,  $\rho_K = 13$ ). The bar differentials at degrees 1–3 for  $V_k(\mathfrak{sl}_2)$  are independently verified in `e1_bar_cobar_cy3.py` (`compute_e1_bar_sl2`, 59 tests). The full Koszul duality computation for the three families is verified in `e1_koszul_three_families.py` (44 tests covering all three families, anti-pattern regression, and cross-family conductor verification).

## 5.7 $E_1$ -chiral bialgebras

The previous sections established the  $E_1$ -chiral algebra (product), the ordered bar coalgebra (coproduct), and the averaging map (symmetrization). This section assembles them into a single object: an  $E_1$ -chiral bialgebra. The definition does not exist in the literature. Classical Hopf algebras live in symmetric monoidal categories; Drinfeld’s quasi-Hopf algebras relax the coassociativity but remain in the same ambient category; Li’s vertex bialgebras use the  $E_\infty$ -chiral tensor product and the coshuffle coproduct, which destroys the quantum group  $R$ -matrix (Proposition 5.6). The correct categorical home for the chiral quantum group is an  $E_1$ -chiral bialgebra: a bialgebra object in the  $E_1$  sector of the Swiss-cheese operad  $\text{SC}^{\text{ch}, \text{top}}$ .

### The $E_1$ -chiral monoidal category

*Definition 5.29* ( $E_1$ -chiral monoidal category). <sup>[PH]</sup> Fix a smooth complex curve  $C$ . The  $E_1$ -chiral monoidal category  $(\mathcal{M}_C^{E_1}, \otimes_{E_1}, \mathbf{1})$  has:

- (i) *Objects*:  $E_1$ -chiral algebras on  $C$  in the sense of the open colour of  $\text{SC}^{\text{ch}, \text{top}}$  (Definition 5.3). These are factorization algebras on the configuration space  $C_n(\mathbb{R}) \subset C_n(C)$  of ordered real points along a ray in  $C$ .
- (ii) *Tensor product*: for  $E_1$ -chiral algebras  $A$  and  $B$ , the *ordered fusion product*

$$A \otimes_{E_1} B := \text{colim}_{z_1 < z_2} A(z_1) \otimes B(z_2),$$

taken over the configuration of two ordered real points in  $C$ . This is NOT symmetric:  $A \otimes_{E_1} B \neq B \otimes_{E_1} A$  in general. The ordering is fixed by the holomorphic argument (Proposition 5.2).

- (iii) *Unit*:  $\mathbf{1} = \Omega_C$ , the augmentation target. The augmentation  $\varepsilon: A \rightarrow \Omega_C$  is the counit of the  $E_1$ -chiral monoidal structure.
- (iv) *Associator*: the ordered fusion is strictly associative:  $(A \otimes_{E_1} B) \otimes_{E_1} C \simeq A \otimes_{E_1} (B \otimes_{E_1} C)$  via the identification  $C_3^{\text{ord}}(\mathbb{R}) \simeq \{z_1 < z_2 < z_3\}$ , which is contractible. This is the  $E_1$  associahedron collapsing to the linear order.

The category  $\mathcal{M}_C^{E_1}$  is monoidal but NOT braided: the  $E_1$ -operad has no braiding operations. This is the essential distinction from both the Beilinson–Drinfeld symmetric ( $E_\infty$ ) setting and the  $E_2$ -braided setting.

### Algebra and coalgebra objects

An  $E_1$ -chiral algebra  $A$  is an algebra object in  $\mathcal{M}_C^{E_1}$ : a product  $\mu: A \otimes_{E_1} A \rightarrow A$  (the OPE with ordered collision: the field at  $z_1 < z_2$  collides onto the field at  $z_2$  from the left, producing residues that differ from the  $z_2 < z_1$  collision by the  $R$ -matrix). The coalgebra side is the ordered bar complex.

*Definition 5.30 ( $E_1$ -chiral coalgebra).* <sup>[PH]</sup> An  $E_1$ -chiral coalgebra in  $\mathcal{M}_C^{E_1}$  is a coaugmented coassociative coalgebra  $(B, \Delta_{\text{dec}}, \eta)$  where:

- (i)  $B = T^c(V)$  is a cofree conilpotent tensor coalgebra on a graded vector space  $V$ ;
- (ii) The coproduct is the *deconcatenation* coproduct (Definition 5.5):

$$\Delta_{\text{dec}}(v_1 \otimes \cdots \otimes v_n) = \sum_{k=0}^n (v_1 \otimes \cdots \otimes v_k) \otimes (v_{k+1} \otimes \cdots \otimes v_n);$$

- (iii) The coaugmentation  $\eta: k \rightarrow B$  includes the degree-zero summand.

The canonical example is  $B^{\text{ord}}(A) = T^c(s^{-1}\bar{A})$  for an augmented  $E_1$ -chiral algebra  $A$ .

### The $E_1$ -chiral bialgebra axiom

*Definition 5.31 ( $E_1$ -chiral bialgebra).* <sup>[PH]</sup> An  $E_1$ -chiral bialgebra on a smooth curve  $C$  is a tuple  $(A, \mu, \Delta_z, \varepsilon, \eta)$  where:

- (i)  $(A, \mu, \varepsilon)$  is an augmented  $E_1$ -chiral algebra in  $\mathcal{M}_C^{E_1}$  (the product is the ordered OPE);
- (ii) For each  $z \in \text{Ran}(C)$ ,  $(A, \Delta_z, \eta)$  is an  $E_1$ -chiral coalgebra in the sense of Definition 5.30, where

$$\Delta_z: A \longrightarrow A \otimes_{E_1, z} A$$

is a family of coproducts parametrized by  $z$ , with  $\otimes_{E_1, z}$  denoting the ordered tensor product at collision point  $z$ . At  $z = 0$ , the coproduct  $\Delta_0$  coincides with the deconcatenation coproduct on  $B^{\text{ord}}(A)$ ; the  $z$ -deformation encodes the spectral data;

- (iii) *Bialgebra compatibility:*  $\Delta_z$  is a morphism of  $E_1$ -chiral algebras, i.e. the diagram

$$\begin{array}{ccc} A \otimes_{E_1} A & \xrightarrow{\mu} & A \\ \Delta_z \otimes \Delta_z \downarrow & & \downarrow \Delta_z \\ (A \otimes_{E_1, z} A) \otimes_{E_1} (A \otimes_{E_1, z} A) & \xrightarrow{\mu \otimes \mu} & A \otimes_{E_1, z} A \end{array}$$

commutes, where the bottom-left corner uses the interchange law for the  $E_1$ -monoidal structure;

- (iv) *Spectral coassociativity:* for  $z, w \in \text{Ran}(C)$  in general position, the coproduct satisfies

$$(\Delta_w \otimes \text{id}) \circ \Delta_{z+w} = (\text{id} \otimes \Delta_z) \circ \Delta_w. \quad (5.12)$$

At  $z = w = 0$  this reduces to ordinary coassociativity  $(\Delta_0 \otimes \text{id}) \circ \Delta_0 = (\text{id} \otimes \Delta_0) \circ \Delta_0$ . For generic  $z, w$  the identity encodes the Yang–Baxter equation: the  $R$ -matrix extracted from  $\Delta_z$  satisfies YBE if and only if spectral coassociativity holds (cf. Proposition 5.41).

The compatibility axiom (iii) is the  $E_1$ -chiral analogue of the classical bialgebra axiom “ $\Delta$  is an algebra map.” The spectral coassociativity axiom (iv) is the chiral analogue of the identity  $(\Delta \otimes 1) \circ \Delta = (1 \otimes \Delta) \circ \Delta$ , promoted to a family over the spectral parameter. It replaces Drinfeld’s quasi-coassociativity (which uses an associator  $\Phi$  to correct the failure of coassociativity in the symmetric monoidal setting): in the  $E_1$ -chiral setting, coassociativity holds on the nose, but with shifted spectral parameters. The parametrization by  $z \in \text{Ran}(C)$  is essential: the coproduct depends on the collision point, and the  $z$ -dependence carries the  $R$ -matrix data.

**PROPOSITION 5.32** (*Spectral coassociativity from factorization*). <sup>[PH]</sup>Let  $(A, \mu, \Delta_z, \varepsilon, \eta)$  be an  $E_1$ -chiral bialgebra on a smooth curve  $C$  in the sense of Definition 5.31. Then the spectral coassociativity axiom (iv) (5.12) is a consequence of the factorization property of  $A$  viewed as a factorization algebra on  $\text{Ran}(C)$ .

*Proof.* The argument is purely operadic; no mode computation is needed.

*Step 1: coproduct as factorization isomorphism.* The  $E_1$ -chiral algebra  $A$  is, by Proposition 5.2, a factorization algebra on the ordered configuration space of  $C$ . For disjoint finite subsets  $S_1, S_2 \subset C$  with  $\Re(S_1) < \Re(S_2)$ , the factorization structure provides a canonical isomorphism

$$\phi_{S_1, S_2} : A(S_1 \sqcup S_2) \xrightarrow{\sim} A(S_1) \otimes_{E_1} A(S_2).$$

The spectral coproduct  $\Delta_z$  is the specialization of  $\phi_{S_1, S_2}$  to the two-point configuration  $S_1 = \{p\}$ ,  $S_2 = \{p + z\}$ , where  $z$  records the relative position along the ordered ray.

*Step 2: triple factorization and the two bracketings.* For three disjoint subsets  $S_1, S_2, S_3$  with  $\Re(S_1) < \Re(S_2) < \Re(S_3)$ , the factorization property gives a canonical isomorphism

$$\phi_{S_1, S_2, S_3} : A(S_1 \sqcup S_2 \sqcup S_3) \xrightarrow{\sim} A(S_1) \otimes_{E_1} A(S_2) \otimes_{E_1} A(S_3).$$

This triple isomorphism factors through two binary compositions:

$$(L): \quad A(S_1 \sqcup S_2 \sqcup S_3) \xrightarrow{\phi_{S_1, S_2 \sqcup S_3}} A(S_1) \otimes_{E_1} A(S_2 \sqcup S_3) \xrightarrow{\text{id} \otimes \phi_{S_2, S_3}} A(S_1) \otimes_{E_1} A(S_2) \otimes_{E_1} A(S_3), \quad (5.13)$$

$$(R): \quad A(S_1 \sqcup S_2 \sqcup S_3) \xrightarrow{\phi_{S_1 \sqcup S_2, S_3}} A(S_1 \sqcup S_2) \otimes_{E_1} A(S_3) \xrightarrow{\phi_{S_1, S_2} \otimes \text{id}} A(S_1) \otimes_{E_1} A(S_2) \otimes_{E_1} A(S_3). \quad (5.14)$$

*Step 3: strict associativity of the  $E_1$ -monoidal structure.* The configuration space  $C_3^{\text{ord}}(\mathbb{R}) = \{z_1 < z_2 < z_3\}$  is contractible (it is an open cone). By Definition 5.29(iv), the  $E_1$ -monoidal category  $\mathcal{M}_C^{E_1}$  is therefore strictly associative: the two bracketings (5.13) and (5.14) coincide as maps from  $A(S_1 \sqcup S_2 \sqcup S_3)$  to the triple tensor product. This is the  $E_1$  associahedron collapsing to the point: the Stasheff polytope  $K_3$  is a single vertex for linear orders, so there is no associator to insert.

*Step 4: translation to spectral parameters.* Specialize to  $S_1 = \{p\}$ ,  $S_2 = \{p + w\}$ ,  $S_3 = \{p + w + z\}$ , so that  $w$  is the separation between  $S_1$  and  $S_2$  and  $z$  is the separation between  $S_2$  and  $S_3$ . The total separation between  $S_1$  and  $S_3$  is  $z + w$ . Then:

- Path (L) computes  $(\text{id} \otimes \Delta_z) \circ \Delta_w$ : first split at parameter  $w$  to separate  $S_1$  from  $S_2 \sqcup S_3$ , then split  $S_2 \sqcup S_3$  at parameter  $z$ .



- Path (R) computes  $(\Delta_w \otimes \text{id}) \circ \Delta_{z+w}$ : first split at parameter  $z + w$  to separate  $S_1 \sqcup S_2$  from  $S_3$ , then split  $S_1 \sqcup S_2$  at parameter  $w$ .

The equality (L) = (R), which is Step 3, is precisely the spectral coassociativity identity

$$(\Delta_w \otimes \text{id}) \circ \Delta_{z+w} = (\text{id} \otimes \Delta_z) \circ \Delta_w.$$

□

*Remark 5.33* (Why this is stronger than formal coassociativity). In the symmetric ( $E_\infty$ ) setting of Beilinson–Drinfeld, the factorization isomorphism is also associative, but the coproduct carries no spectral parameter and the resulting coassociativity is the classical identity  $(\Delta \otimes 1)\Delta = (1 \otimes \Delta)\Delta$ . In the quasi-Hopf ( $E_0$ ) setting of Drinfeld, the configuration space  $C_3$  is NOT contractible after quotienting by the symmetric group, so the two paths (L) and (R) differ by an associator  $\Phi$ . The  $E_1$  setting is intermediate: the ordered configuration space IS contractible (so no associator), but the collision parameters are retained (so the identity is spectral). This is the operadic reason that the  $E_1$ -chiral bialgebra achieves strict coassociativity with spectral parameters: the  $E_1$ -operad supplies exactly the right amount of homotopy coherence.

*Definition 5.34* ( $E_1$ -chiral Hopf algebra (antipode)). <sup>[PH]</sup> An  $E_1$ -chiral bialgebra  $(A, \mu, \Delta_z, \varepsilon, \eta)$  is an  $E_1$ -chiral Hopf algebra if there exists a morphism of  $E_1$ -chiral modules  $S: A \rightarrow A$  (the *antipode*) satisfying the Hopf axiom at  $z = 0$ :

$$\mu_{E_1} \circ (S \otimes \text{id}) \circ \Delta_0 = \eta \circ \varepsilon = \mu_{E_1} \circ (\text{id} \otimes S) \circ \Delta_0, \quad (5.15)$$

where  $\mu_{E_1}$  is the labeled-ordered product and  $\Delta_0 = \Delta_z|_{z=0}$  is the coproduct at the basepoint. The two equalities are independent conditions because  $\mu_{E_1}$  is not commutative. For  $z \neq 0$ , the antipode axiom generalizes to

$$\mu_{E_1} \circ (S \otimes \text{id}) \circ \Delta_z = \eta \circ \varepsilon \circ \pi_z, \quad (5.16)$$

where  $\pi_z: A \rightarrow A$  is the *spectral projection*: the composition  $A \xrightarrow{\Delta_z} A \otimes_{E_1, z} A \xrightarrow{\varepsilon \otimes \text{id}} A$  evaluated at the  $z$ -independent term, i.e.  $\pi_z = (\varepsilon \otimes \text{id}) \circ \Delta_z|_{z=0}$ , extracting the constant coefficient of the Laurent expansion in  $z$ . When  $A$  is connected ( $\varepsilon \circ \eta = \text{id}_k$ ), the basepoint axiom (5.15) determines  $S$  uniquely by induction on the coalgebra filtration, exactly as in the classical Hopf algebra setting.

*Remark 5.35* (Summary of the  $E_1$ -chiral Hopf axiom system). Definitions 5.31 and 5.34 assemble into a five-part axiom system for an  $E_1$ -chiral Hopf algebra  $(A, \mu, \Delta_z, \varepsilon, \eta, S)$  on a curve  $C$ :

- (H1)  *$E_1$ -chiral algebra*:  $(A, \mu, \varepsilon)$  is an augmented algebra in  $\mathcal{M}_C^{E_1}$ ; the product  $\mu$  is the labeled-ordered OPE (Definition 5.29).
- (H2)  *$E_1$ -chiral coalgebra*:  $\Delta_z: A \rightarrow A \otimes_{E_1, z} A$  is a  $z$ -parametrized coproduct with counit  $\varepsilon$  and coaugmentation  $\eta$  (Definition 5.30).
- (H3) *Bialgebra compatibility*:  $\Delta_z$  is a morphism of  $E_1$ -chiral algebras: the diagram of Definition 5.31(iii) commutes.
- (H4) *Spectral coassociativity*:  $(\Delta_w \otimes \text{id}) \circ \Delta_{z+w} = (\text{id} \otimes \Delta_z) \circ \Delta_w$  (equation (5.12)).
- (H5) *Hopf axiom*:  $\mu_{E_1} \circ (S \otimes \text{id}) \circ \Delta_0 = \eta \circ \varepsilon = \mu_{E_1} \circ (\text{id} \otimes S) \circ \Delta_0$  (equation (5.15)).

The axioms are strictly stronger than Li’s vertex-bialgebra framework ( $E_\infty$ -chiral, no spectral parameter, no antipode) and strictly weaker than Drinfeld’s quasi-Hopf formulation (which works in  $(\text{Vect}, \otimes_k)$  and requires an associator  $\Phi$ ). The  $E_1$ -chiral setting eliminates the associator: axiom (H4) holds on the nose because the  $E_1$ -monoidal category is strictly associative (the associahedron is contractible for linear orders).

### Why $E_1$ and not $E_\infty$

PROPOSITION 5.36 (*Averaging forgetful theorem*). <sup>[PH]</sup> The averaging map of Proposition 5.6 defines a functor

$$\text{av}: E_1\text{-ChirHopf}(C) \longrightarrow E_\infty\text{-ChirCoalg}(C)$$

that sends an  $E_1$ -chiral Hopf algebra  $(A, \mu, \Delta_z, \varepsilon, \eta, S)$  satisfying axioms **(H1)**–**(H5)** of Remark 5.35 to the  $E_\infty$ -chiral coalgebra  $(B^\Sigma(A), \Delta_{\text{cosh}})$  with the coshuffle coproduct  $\Delta_{\text{cosh}}$ . The functor forgets all Hopf data:

- (i) The  $R$ -matrix: on degree-two elements,  $\text{av}(r(z)) = \kappa_{\text{ch}}$  (a scalar), so the full  $z$ -dependent  $R$ -matrix  $r(z) = k \Omega/z$  is replaced by its  $S_2$ -coinvariant, which is the collision residue  $\kappa_{\text{ch}}$ .
- (ii) The antipode:  $S$  acts on ordered tensor factors; after symmetrization,  $S$  becomes the identity (on  $S_n$ -invariants, the reversal acts trivially).
- (iii) The coproduct  $z$ -dependence: the coshuffle coproduct is independent of the collision point  $z$ , while the deconcatenation coproduct  $\Delta_z$  depends on  $z$  through the ordered OPE residues.

In particular, the  $E_\infty$ -chiral coalgebra  $B^\Sigma(A)$  is NOT a bialgebra: it retains no Hopf structure. The coshuffle coproduct on  $B^\Sigma$  is compatible only with the commutative product, which  $A$  does not possess.

*Proof.* Statement (i) is Proposition 5.6. Statement (ii): the antipode  $S$  is an anti-homomorphism of the ordered product ( $S(\mu(a, b)) = \mu(S(b), S(a))$ ), so on  $S_n$ -coinvariants  $\text{av} \circ S = \text{av}$  (symmetrization absorbs the reversal of tensor factors). Statement (iii): the coshuffle coproduct is  $\Delta_{\text{cosh}} = \text{av}^{\otimes 2} \circ \Delta_{\text{dec}}$ , and  $\text{av}^{\otimes 2}$  kills the  $z$ -dependence because  $\sum_\sigma r(\sigma \cdot z) = n! \cdot \kappa_{\text{ch}}$ . The final claim (no bialgebra structure on  $B^\Sigma$ ) follows because the bialgebra compatibility diagram of Definition 5.31(iii) requires the interchange law for the  $E_1$  tensor product, which does not hold in the symmetric quotient.  $\square$

This is the central structural result: Li's vertex-bialgebra framework, which works with  $B^\Sigma$  and the coshuffle coproduct, cannot carry a Hopf structure. The quantum group  $R$ -matrix, the antipode, and the spectral-parameter dependence all live on the  $E_1$  side and are destroyed by symmetrization. The  $E_1$ -chiral bialgebra is the minimal structure that retains all three.

Remark 5.37 ( $A_\infty$  coproduct and the shadow tower). For class  $M$  algebras (infinite shadow depth), the formal Yangian coproduct  $\Delta_z^{\text{Yangian}}$  is the  $m_2$ -truncation of the full  $A_\infty$  coproduct:

$$\Delta_z^{A_\infty}(T) = \Delta_z^{\text{Yangian}}(T) + \hbar^2 \cdot \delta^{(3)}(T) + \hbar^3 \cdot \delta^{(4)}(T) + \dots$$

where  $\delta^{(k)}$  is the correction from the  $A_\infty$  operation  $m_k$ , with coefficient equal to the shadow invariant  $S_k$ . Concretely (`a_infinity_bar_wlinf.py`):  $m_3(T, T, T) = -2T$  gives  $\delta^{(3)}$  with coefficient  $\alpha = 2$ ;  $m_4(T, T, T, T) = \frac{40}{27}T$  gives  $\delta^{(4)}$  with coefficient  $S_4 = \frac{10}{27}$ . For class  $G$  (Heisenberg): all  $m_k = 0$  for  $k \geq 3$ , so the truncation is exact. For class  $L$  (Yangian):  $S_4 = 0$ , so the truncation terminates at finite depth. For class  $M$  (Virasoro): all  $S_k \neq 0$ , and the coproduct has genuinely infinite complexity. The shadow tower is therefore not merely a classification invariant: it encodes the  $A_\infty$  corrections to the chiral coproduct.

PROPOSITION 5.38 (*Universal coproduct formula from the Miura factorization*). <sup>[PH]</sup> Let  $A = W_{1+\infty}$  at level  $\Psi$  with transfer matrix  $T(u) = \sum_{s \geq 0} (-1)^s \psi_s u^{-s}$ , where  $\psi_s = e_s(\phi_1, \dots, \phi_N)$  is the  $s$ -th elementary symmetric polynomial in the free bosons. The Miura coproduct  $\Delta_z(T(u)) = T_L(u) \cdot T_R(u - z)$  gives:

$$\Delta_z(\psi_s) = \sum_{\substack{a+b+k=s \\ a, b, k \geq 0}} (-1)^k \binom{N_R - b}{k} z^k \psi_a^L \cdot \psi_b^R.$$

The coproduct at spin  $s$  is a polynomial in  $z$  of degree exactly  $s$ , with  $\frac{s(s+1)}{2}$  operator-product terms. The number of terms at  $z$ -power  $p$  is  $N(s, p) = s - p$ , with generating function

$$F(x, y) = \sum_{s \geq 1} \sum_{p=0}^{s-1} N(s, p) x^s y^p = \frac{x}{(1-x)^2(1-xy)}.$$

The leading  $z^{s-1}$  coefficient is  $\psi_1^R = J^R$  (single term). The subleading  $z^{s-2}$  coefficient is  $(s-1)\psi_2^R + \sum_k J_k^L J_{n-k}^R$  (universal at all spins).

*Proof.* Expand  $T_R(u - z) = \sum_b (-1)^b \psi_b^R(u - z)^{-b}$  using the binomial series  $(u - z)^{-b} = \sum_{k \geq 0} \binom{b-1+k}{k} z^k u^{-b-k}$ . Collecting the  $u^{-(a+b+k)}$  coefficient from the product  $T_L(u) \cdot T_R(u - z)$  gives the stated formula. The term count follows: at  $z$ -power  $p$ , the constraint  $a + b = s - p$  with  $a \geq 0$  gives  $s - p$  choices of  $(a, b)$ . The generating function  $F$  is obtained by geometric-series summation. See `compute/lib/chiral_coproduct_allspin_engine.py`.  $\square$

*Remark 5.39* (Spectral vs. worldsheet interpretation of  $z$ ). The spectral parameter  $z$  in  $\Delta_z$  is a shift of the transfer-matrix argument  $u \mapsto u - z$  in the Yangian mode algebra  $Y(\widehat{\mathfrak{gl}}_1)$ . It is *not* the worldsheet coordinate of a vertex operator insertion. The OPE singularity  $T(z)T(w) \sim c/2 \cdot (z-w)^{-4}$  involves the worldsheet variable; the coproduct  $\Delta_z$  involves the spectral variable. Setting  $z = 0$  in  $\Delta_z$  removes the spectral shift and gives a well-defined, cocommutative coproduct with no OPE singularity (verified in `test_hopf_axioms.py`, 50 tests). The two  $z$ 's live in different mathematical objects and must not be conflated.

**PROPOSITION 5.40** (*Coassociativity from the Ran space  $E_1$ -monoidal structure*). <sup>[PH]</sup> Let  $A$  be an  $E_1$ -factorization algebra on a smooth curve  $C$ . The factorization isomorphism  $\phi_{S_1, S_2}: A(S_1 \sqcup S_2) \xrightarrow{\sim} A(S_1) \otimes A(S_2)$  for disjoint  $S_1, S_2 \in \text{Ran}(C)$  is strictly coassociative: the two iterated coproducts

$$(\phi_{S_1, S_2} \otimes \text{id}) \circ \phi_{S_1 \sqcup S_2, S_3} = (\text{id} \otimes \phi_{S_2, S_3}) \circ \phi_{S_1, S_2 \sqcup S_3}$$

coincide as maps  $A(S_1 \sqcup S_2 \sqcup S_3) \rightarrow A(S_1) \otimes A(S_2) \otimes A(S_3)$ .

*Proof.* The ordered configuration space  $C_n^{\text{ord}}(\mathbb{R}) = \{x_1 < \dots < x_n\}$  is contractible for every  $n$  (it is convex). By the Ayala–Francis theorem (arXiv:1504.04007, Theorem 3.24), an  $E_1$ -algebra is equivalently a locally constant factorization algebra on  $\mathbb{R}$ , and the  $E_1$ -monoidal structure on  $\text{Ran}(C)$  is strictly associative: the space of parenthesizations of an  $n$ -fold labeled-ordered product is connected (contractible associahedron  $K_n \simeq *$ ). Applied to  $n = 3$ , the two binary decompositions  $(S_1 \sqcup S_2) \sqcup S_3$  and  $S_1 \sqcup (S_2 \sqcup S_3)$  are connected by the unique path in  $K_3 = *$ . Hence the two iterated factorization isomorphisms agree, which is coassociativity.  $\square$

## The Drinfeld center recovers $E_2$

The  $E_1$ -chiral bialgebra  $A$  is ordered and non-braided. The braided ( $E_2$ ) structure is recovered by the Drinfeld center.

**PROPOSITION 5.41** ( *$R$ -matrix from the Drinfeld center*). <sup>[PH]</sup> Let  $(A, \mu, \Delta_z, \varepsilon, \eta, S)$  be an  $E_1$ -chiral Hopf algebra. The Drinfeld center  $\mathcal{Z}(\text{Rep}^{E_1}(A))$  is a braided monoidal category (Proposition 5.10), and the braiding on  $\mathcal{Z}$  defines an  $R$ -matrix  $R(z) \in (A \otimes A)((z))$  characterized by the property: for all  $A$ -modules  $V, W$ , the half-braiding  $\sigma_{V, W}(z): V \otimes_{E_1} W \xrightarrow{\sim} W \otimes_{E_1} V$  acts by

$$\sigma_{V, W}(z)(v \otimes w) = \tau(R(z) \cdot (v \otimes w)), \quad (5.17)$$

where  $\tau$  is the transposition of tensor factors. The  $R$ -matrix lives in  $\mathcal{Z}(\text{Rep}^{E_1}(A))$ , not in  $A \otimes A$  directly: it is the half-braiding data encoding the obstruction to commutativity of the ordered tensor product. This resolves the apparent paradox that the  $R$ -matrix “lives in  $A \otimes A$ ” (which is ordered and has no braiding): the  $R$ -matrix is a morphism in the Drinfeld center, which IS braided.

*Proof.* The Drinfeld center  $\mathcal{Z}(\text{Rep}^{E_1}(A))$  consists of pairs  $(M, \sigma_M)$  where  $M$  is an  $A$ -module and  $\sigma_M: M \otimes_{E_1} N \xrightarrow{\sim} N \otimes_{E_1} M$  is a half-braiding natural in  $N$ . For  $M = A$  (the regular module), the half-braiding is constructed from the spectral coproduct: define  $\sigma_A(z)(a \otimes n) = \sum \Delta_z(a)_{(2)} \cdot n \otimes \Delta_z(a)_{(1)}$  (using Sweedler notation for  $\Delta_z$ ). The naturality of  $\sigma_A$  in the second slot follows from the bialgebra compatibility (axiom (iii) of Definition 5.31):  $\Delta_z$  is an algebra map, so  $\sigma_A$  intertwines the  $A$ -action. The invertibility of  $\sigma_A$  uses the antipode: the inverse half-braiding is  $\sigma_A^{-1}(z)(n \otimes a) = \sum S(\Delta_z(a)_{(1)}) \cdot n \otimes \Delta_z(a)_{(2)}$ , which is well-defined by the Hopf axiom (Definition 5.34). The  $R$ -matrix  $R(z) \in (A \otimes A)((z))$  is then the element implementing (5.17), extracted as the image of  $\sigma_A$  under the canonical map  $\text{End}(A \otimes_{E_1} A) \rightarrow (A \otimes A)((z))$ . The braiding satisfies the Yang–Baxter equation because the Drinfeld center is a braided monoidal category (a formal consequence of the definition). See Theorem 9.4 for the explicit verification at  $A = Y^+(\widehat{\mathfrak{gl}}_1)$ .  $\square$

## Examples

*Example 5.42 (Heisenberg  $E_1$ -chiral bialgebra).* The rank-one Heisenberg  $H_k$  ( $k \neq 0$ ) is an  $E_1$ -chiral Hopf algebra with:

- Product:  $\mu(J(z_1), J(z_2)) = :J(z_1)J(z_2): + k/(z_1 - z_2)^2$  (the time-ordered OPE, not the normally-ordered product).
- Coproduct:  $\Delta_z(J) = J \otimes 1 + 1 \otimes J$  (primitive). The Heisenberg coproduct is  $z$ -independent because the  $r$ -matrix  $r(z) = k/z$  has no nontrivial  $z$ -dependent correction to  $\Delta$ .
- Antipode:  $S(J) = -J$ , so  $\mu_{E_1}(S(J) \otimes 1 + 1 \otimes J) = -J + J = 0 = \eta(\varepsilon(J))$ .
- $R$ -matrix from Drinfeld center:  $R(z) = e^{k \log z \cdot h \otimes h}$  (the Heisenberg vertex  $R$ -matrix; here  $h$  is the Cartan generator identified with  $J$ ). After averaging,  $\text{av}(R(z)) = k$  (the level), consistent with  $\kappa_{\text{ch}}(H_k) = k$ .

The Heisenberg is class G:  $\Delta$  is primitive,  $S$  is the sign involution, and the Hopf structure is the simplest possible.

**CONJECTURE 5.43** ( $W_{1+\infty}$   $E_1$ -chiral bialgebra). <sup>[CJ]</sup> The  $W_{1+\infty}$  algebra at  $c = 1$  (equivalently, the affine Yangian  $Y(\widehat{\mathfrak{gl}}_1)$  via Procházka–Rapčák) carries an  $E_1$ -chiral Hopf algebra structure with:

- (i) *Coproduct*: for the generating field  $T(u)$  (the Yangian transfer matrix),  $\Delta_z(T(u)) = T(u) \otimes T(u)$  (group-like, NOT primitive). The deviation from primitivity is controlled by the Dimofte  $K$ -matrix (Remark 9.7):  $\Delta_z(W_n) = W_n \otimes 1 + K_{\mathcal{A}}(z) \otimes W_n + \dots$  for the higher spin fields  $W_n$ .
- (ii) *Antipode*:  $S(T(u)) = T(u)^{-1}$  (formal inverse of the transfer matrix). On modes:  $S(W_2) = -W_2$ ,  $S(W_3) = -W_3 + W_2^2$  (the Hopf antipode recursion).
- (iii)  *$R$ -matrix*: via the Drinfeld center, the universal  $R$ -matrix is the Yangian  $R$ -matrix of Theorem 9.5 with structure function  $g(u)$ .

- (iv) *Averaging*:  $\text{av}(\Delta_z) = \Delta_{\text{cosh}}$  (the coshuffle coproduct on  $B^\Sigma$ ), and  $\text{av}(R(u)) = g(0) = -1$  (the structure function at  $u = 0$ , a scalar). The quantum group data is destroyed.

This conjecture depends on the existence of the  $E_1$ -chiral bialgebra structure on  $W_{1+\infty}$  in the sense of Definition 5.31; the individual ingredients (Yangian coproduct,  $R$ -matrix, antipode) are established (Schiffmann–Vasserot, Procházka–Rapčák), but their assembly into an  $E_1$ -chiral bialgebra in the Swiss-cheese framework requires verifying the compatibility diagram (iii) of Definition 5.31.

*Remark 5.44* (The three-way resolution). The  $E_1$ -chiral bialgebra resolves the apparent conflict between three classical frameworks for quantum group structure on chiral algebras:

- (a) *Drinfeld quasi-Hopf*: lives in the symmetric monoidal category of vector spaces, with a non-coassociative coproduct corrected by the Drinfeld associator  $\Phi$ . This is the WRONG ambient category: chiral algebras are not vector spaces, and the associator  $\Phi$  is an artifact of forcing a braided structure into a symmetric setting.
- (b) *Li vertex bialgebra*: lives in the  $E_\infty$ -chiral monoidal category, with the coshuffle coproduct on  $B^\Sigma$ . This uses the CORRECT ambient category (chiral) but the WRONG structure level ( $E_\infty$ ), destroying the  $R$ -matrix.
- (c)  *$E_1$ -chiral bialgebra*: lives in the  $E_1$ -chiral monoidal category  $\mathcal{M}_C^{E_1}$ , with the deconcatenation coproduct on  $B^{\text{ord}}$ . This uses the correct ambient category AND the correct structure level. The  $R$ -matrix is recovered by passing to the Drinfeld center  $\mathcal{Z}(\text{Rep}^{E_1}(A))$ , which is the  $E_2$  enhancement.

The slogan: the  $E_1$ -chiral bialgebra is the Hopf algebra object in the category that chiral algebras actually inhabit when collision order is remembered.

*Remark 5.45* (Comparison table:  $E_1$  vs  $E_2$  vs  $E_\infty$  chiral bialgebras). Table 5.1 records the structural data at each level of the operad filtration  $E_1 \rightarrow E_2 \rightarrow E_\infty$ . The passage  $E_1 \rightarrow E_2$  is the Drinfeld center (Proposition 5.41); the passage  $E_2 \rightarrow E_\infty$  is the averaging map  $\text{av}: B^{\text{ord}} \rightarrow B^\Sigma$  (Proposition 5.6). Each arrow *loses* structure irreversibly: the first quotients out the ordering to gain a braiding; the second quotients out the braiding to gain full symmetry. The slogan:  $E_\infty$ -averaging kills the Hopf structure because  $\text{av}(r(z)) = \kappa_{\text{ch}}$ : the full  $z$ -dependent  $R$ -matrix collapses to a single scalar.

The table makes precise the content of Remark 5.44: Li’s vertex bialgebras occupy the rightmost column, where the coshuffle coproduct is  $z$ -independent and carries no  $R$ -matrix data. Drinfeld’s quasi-Hopf algebras do not appear in the table because they live in  $(\text{Vect}, \otimes_k)$ , not in a chiral monoidal category; the associator  $\Phi$  is the artifact of forcing a braided object into a symmetric ambient category. The  $E_1$ -chiral bialgebra (leftmost column) is the minimal framework retaining all quantum group data.

## The factorization-homology coproduct

The preceding subsections defined the spectral coproduct  $\Delta_z$  and proved coassociativity from the factorization property (Proposition 5.32). That proof is *operadic*: it uses the factorization isomorphism of the  $E_1$ -chiral algebra  $A$  on a curve. A deeper question is whether the coproduct itself can be *derived* from factorization homology, bypassing mode-level constructions such as the Miura factorization (which requires toric data and may not exist for general CY categories). The answer is yes: excision for factorization homology (Proposition 12.7) provides a canonical coproduct on any chiral algebra obtained by integrating a factorization algebra over a compact factor of a product CY manifold.

**(i) The  $E_3$  factorization algebra on  $\mathbb{C}^3$  and the Ran space diagonal.** Let  $\mathcal{F}$  be the  $E_3$  factorization algebra of Conjecture 7.9 on  $\mathbb{C}^3$ . For a disjoint union of open disks  $U_1 \sqcup U_2 \hookrightarrow \mathbb{C}^3$  in the Ran space  $\text{Ran}(\mathbb{C}^3)$ , the factorization structure of  $\mathcal{F}$  provides a canonical isomorphism

$$\mathcal{F}(U_1 \sqcup U_2) \xrightarrow{\sim} \mathcal{F}(U_1) \otimes \mathcal{F}(U_2).$$

| Feature                   | $E_1$ (ordered)                       | $E_2$ (braided)                      | $E_\infty$ (symmetric)                          |
|---------------------------|---------------------------------------|--------------------------------------|-------------------------------------------------|
| Operad                    | Little intervals                      | Little 2-disks                       | Comm. operad                                    |
| Symmetry at degree $n$    | Trivial (1)                           | Braid group $B_n$                    | $S_n$                                           |
| Bar complex               | $B^{\text{ord}} = T^c(s^{-1}\bar{A})$ | $B_{E_2} = T^c$ with $B_n$ -action   | $B^\Sigma = \text{Sym}^c(s^{-1}\bar{A})$        |
| Coproduct                 | Deconcatenation $\Delta_{\text{dec}}$ | $\Delta_X, \Delta_Y$ (two commuting) | Coshuffle $\Delta_{\text{cosh}}$                |
| $z$ -dependence           | $\Delta_z$ spectral family            | $R^{E_2}(z)$ braiding                | None ( $z$ -independent)                        |
| $R$ -matrix               | $r(z) = k \Omega/z + \dots$           | $R^{E_2}(z)$ (half-braiding)         | $\text{av}(r(z)) = \kappa_{\text{ch}}$ (scalar) |
| Koszul dual               | $A^!$ with reversed order             | $A^!$ with reversed braiding         | $A^!$ (no braiding data)                        |
| Hopf structure            | Full: $(\mu, \Delta_z, S)$            | Braided Hopf                         | None (Prop. 5.36)                               |
| Antipode $S$              | $S: A \rightarrow A$                  | Braided antipode                     | $\text{av}(S) = \text{id}$                      |
| Yang–Baxter               | Spectral coassoc. (H4)                | YBE from center                      | Trivial                                         |
| Rep. category             | Monoidal (not braided)                | Braided monoidal                     | Symmetric monoidal                              |
| <b>Lost at transition</b> | —                                     | Ordering; strict assoc.              | Braiding; $R$ -matrix; $S$ ; $z$ -dep.          |

Table 5.1: Twelve-row comparison of  $E_n$ -chiral bialgebra data. The  $E_1 \rightarrow E_2$  transition (Drinfeld center) trades the strict associativity of linear order for a braiding; the  $E_2 \rightarrow E_\infty$  transition (averaging) collapses  $r(z)$  to  $\kappa_{\text{ch}}$  and destroys the Hopf structure entirely. See Construction 6.21 for the bar-complex triple.

The *Ran diagonal*  $\delta: \text{Ran}(\mathbb{C}^3) \rightarrow \text{Ran}(\mathbb{C}^3) \times \text{Ran}(\mathbb{C}^3)$  (the map sending a finite set  $S$  to  $(S_1, S_2)$  for each decomposition  $S = S_1 \sqcup S_2$ ) produces a cosheaf-level coproduct: the global sections  $\mathcal{F}(\mathbb{C}^3)$  acquire a coalgebra structure via

$$\Delta^{\text{Ran}}: \mathcal{F}(U) \xrightarrow{\phi_{U_1, U_2}} \mathcal{F}(U_1) \otimes \mathcal{F}(U_2), \quad (5.18)$$

summed over all ways of decomposing  $U = U_1 \sqcup U_2$  into disjoint subsets. This is the factorization coproduct at the level of  $\mathbb{C}^3$ . The excision property (Proposition 12.7) guarantees that  $\Delta^{\text{Ran}}$  is well-defined: for a collar-gluing  $U = U_1 \cup_{N \times \mathbb{R}} U_2$  of a domain along a codimension-1 collar,

$$\int_U \mathcal{F} \simeq \int_{U_1} \mathcal{F} \otimes_{\int_{N \times \mathbb{R}} \mathcal{F}} \int_{U_2} \mathcal{F}.$$

No mode expansion or Miura-type factorization is needed: the excision gluing data is carried by the cosheaf structure of  $\mathcal{F}$  on  $\text{Ran}(\mathbb{C}^3)$ .

**(2)  $K3 \times E$ : the  $E$ -direction coproduct from cutting the elliptic curve.** For the product  $\text{CY}_3$  manifold  $X = K3 \times E$ , the factorization homology integral  $\int_{K3} \mathcal{F}|_{K3}$  of Section 14.10 produces a chiral algebra  $A_E$  on the residual elliptic curve  $E$ . The coproduct on  $A_E$  arises from cutting  $E$  into two intervals via excision. Concretely, choose a collar-gluing decomposition  $E = I_1 \cup_{S^0 \times \mathbb{R}} I_2$  (cutting  $E$  at two points into two intervals  $I_1, I_2$  with collar neighbourhoods). The excision equivalence gives

$$\int_E A_E \simeq \int_{I_1} A_E \otimes_{\int_{S^0 \times \mathbb{R}} A_E} \int_{I_2} A_E. \quad (5.19)$$

The map  $A_E \rightarrow \int_{I_1} A_E \otimes \int_{I_2} A_E$  is the *factorization coproduct*: it decomposes the algebra on the full curve into tensor factors localized on the two intervals. Since  $A_E$  is  $E_1$ -chiral on  $E$  (holomorphy provides the

ordering), the decomposition (5.19) respects the  $E_1$  ordering of the intervals, and the resulting coproduct is an  $E_1$ -chiral coalgebra map.

The essential point: this construction requires *only* that  $\int_{K_3} \mathcal{F}|_{K_3}$  exists as a factorization algebra on  $E$ . No Miura transform, no toric structure on the  $K_3$  factor, and no mode decomposition of the OPE are needed. The coproduct is a consequence of the factorization structure of  $A_E$  on the Ran space of  $E$ .

**(3) Coassociativity from the Ran monoidal structure.** The coassociativity of  $\Delta^{\text{Ran}}$  follows from the associativity of the Ran space monoidal structure, by the same operadic argument as Proposition 5.32. For three disjoint subsets  $S_1, S_2, S_3$ , the two bracketings

$$\begin{aligned} \mathcal{F}(S_1 \sqcup S_2 \sqcup S_3) &\xrightarrow{\phi_{S_1, S_2 \sqcup S_3}} \mathcal{F}(S_1) \otimes \mathcal{F}(S_2 \sqcup S_3) \xrightarrow{\text{id} \otimes \phi_{S_2, S_3}} \mathcal{F}(S_1) \otimes \mathcal{F}(S_2) \otimes \mathcal{F}(S_3), \\ \mathcal{F}(S_1 \sqcup S_2 \sqcup S_3) &\xrightarrow{\phi_{S_1 \sqcup S_2, S_3}} \mathcal{F}(S_1 \sqcup S_2) \otimes \mathcal{F}(S_3) \xrightarrow{\phi_{S_1, S_2} \otimes \text{id}} \mathcal{F}(S_1) \otimes \mathcal{F}(S_2) \otimes \mathcal{F}(S_3) \end{aligned}$$

coincide because Ran is a symmetric monoidal  $\infty$ -category under disjoint union, and the factorization algebra is a symmetric monoidal functor from  $\text{Disk}_{n/M}$  to chain complexes. No modes, no spectral parameters, no computation: the proof is a formal consequence of the symmetric monoidal structure of the Ran space.

When the  $E_1$ -ordering is imposed (restricting to ordered configurations along a real ray in  $E$ ), the coassociativity specializes to the spectral coassociativity identity  $(\Delta_w \otimes \text{id}) \circ \Delta_{z+w} = (\text{id} \otimes \Delta_z) \circ \Delta_w$  of equation (5.12), recovering the  $E_1$ -chiral bialgebra axiom (H4).

**CONJECTURE 5.46** (*Factorization-homology coproduct agrees with Miura for toric  $CY_3$* ). <sup>[CJ]</sup> Let  $X$  be a toric  $CY_3$  and let  $\mathcal{F}$  be the  $E_3$  factorization algebra of Conjecture 7.9 restricted to  $X$ . Let  $A_C = \int_{X/C} \mathcal{F}$  be the chiral algebra on a residual curve  $C$  obtained by factorization homology over the transverse directions.

- (i) The factorization-homology coproduct  $\Delta^{\text{Ran}}$  of (5.18), restricted to the  $E_1$ -sector, coincides with the Drinfeld coproduct  $\Delta_z$  of Definition 5.31(ii).
- (ii) For  $X = \mathbb{C}^3$  with  $\Omega$ -background  $(h_1, h_2, h_3)$  and gauge algebra  $\mathfrak{gl}_1$ , the Drinfeld coproduct on the affine Yangian  $Y(\widehat{\mathfrak{gl}}_1)$  is recovered from the Ran diagonal of the  $E_3$ -factorization algebra on  $\mathbb{C}^3$ : the Procházka–Rapčák Miura factorization of  $\mathcal{W}_{1+\infty}$  is the toric specialization of the general excision coproduct.

The conjecture is conditional on Conjecture 7.9 (the  $E_3$  factorization algebra on  $\mathbb{C}^3$ ). For the affine Yangian specifically, the Drinfeld coproduct is constructed independently (Guay–Nakajima–Wendlandt); the content of (ii) is that the two constructions agree under the identification  $\int_{\mathbb{C}^2} \mathcal{F}|_{\mathbb{C}^2} \simeq Y(\widehat{\mathfrak{gl}}_1)$ .

**Remark 5.47** (Well-definedness for general  $CY_3$ ). The factorization-homology coproduct  $\Delta^{\text{Ran}}$  is well-defined for *any*  $CY_3$  category  $\mathcal{C}$  for which the CY-to-chiral functor  $\Phi$  produces a factorization algebra, i.e., whenever the relevant CY-to-chiral statement is available: unconditionally at  $d = 2$  by Theorem CY-A<sub>2</sub>, and conditionally at  $d = 3$  via CY-A<sub>3</sub>.

- (a) *At  $d = 2$  (unconditional):* For a  $CY_2$  category (e.g.  $D^b(\text{Coh}(K3))$ ), the functor  $\Phi$  produces an  $E_2$ -chiral algebra (Theorem CY-A<sub>2</sub>), and the factorization-homology coproduct is the excision map for any collar-gluing of the underlying surface. This recovers the lattice vertex algebra coproduct (primitive on generators) without any toric structure.
- (b) *At  $d = 3$  (conditional on CY-A<sub>3</sub>):* The functor  $\Phi$  is conjectural, but if it produces a factorization algebra  $\mathcal{F}$  on a curve  $C$ , then the excision coproduct is automatically well-defined. The Miura factorization is the toric case

where  $\mathcal{F}$  admits a decomposition into free-field factors (one per toric leg of the web diagram); for non-toric  $\text{CY}_3$  (e.g. the quintic, a general complete intersection, or a Pfaffian CY), no such decomposition exists, and the factorization-homology coproduct is the *only* available construction.

- (c) *Obstructions:* The factorization-homology coproduct requires the chiral algebra  $A = \Phi(C)$  to be a factorization algebra (not merely a vertex algebra in the algebraic sense). This is automatic when  $A$  is obtained from the Costello–Gwilliam programme, but may fail for algebraically constructed vertex algebras that lack a factorization-algebra enhancement. In the CY setting, the Costello–Gwilliam construction is the native one, so no additional assumption is needed beyond CY-A.

The factorization-homology coproduct therefore provides a *universal* source of the  $E_1$ -chiral coalgebra structure: whenever the CY-to-chiral functor exists, the coproduct exists. The Miura factorization is the computational specialization in the toric case.

## 5.8 Closing: the ordered bar as the primitive of Vol III

Four facts control the later parts of Vol III. First, factorization algebras on a complex curve are  $E_2$  topologically but specialize to  $E_1$  when holomorphy fixes an ordering, and the Swiss-cheese splitting  $E_1(C) \times E_2(\text{top})$  makes that specialization precise. Second, the ordered bar complex  $B^{\text{ord}}(A) = T^c(s^{-1}\bar{A})$  with deconcatenation coproduct is the natural  $E_1$  primitive, and averaging to the symmetric bar  $B^{\Sigma}(A)$  is lossy because it kills the ordered  $R$ -matrix data. Third, the CY-to-chiral functor  $\Phi$  at  $d = 2$  produces an ordered bar whose Euler character is the Borcherds denominator and whose first shadow invariant is  $\kappa_{\text{ch}}$ , distinct from  $\kappa_{\text{BKM}}$ ,  $\kappa_{\text{cat}}$ , and  $\kappa_{\text{fiber}}$ . Fourth, the  $E_2$  enhancement requires the Drinfeld center construction, the categorified averaging map  $E_1\text{-Cat} \rightarrow E_2\text{-Cat}$ ; at  $d = 3$  the enhancement is conditional on the CY- $A_3$  programme.

This is the version of the  $E_1$ - $E_1$  operadic Koszul duality adapted to the CY setting: one form  $\eta = d \log(z_1 - z_2)$ , one relation (Arnold), one object  $\Theta_A$ , one equation  $D_*\Theta + \frac{1}{2}[\Theta, \Theta] = 0$ , all on the ordered bar of the cyclic  $A_\infty$  algebra  $A_C$  attached to the Calabi-Yau category  $\mathcal{C}$ . The symmetric bar is the modular shadow.

The remaining parts use these facts directly. II computes the modular trace  $\kappa_{\text{ch}}$  for each CY family, using the ordered bar as the computational model. III catalogues the families  $K3$ ,  $K3 \times E$ , toric  $\text{CY}_3$ , Fukaya, and quantum-group, and identifies the corresponding point in the kappa-spectrum. IV reads the seven faces of  $r_{\text{CY}}(z)$  from the  $E_1$  structure of Proposition 5.8. V collects the open problems, most of them variants of Conjecture 5.9 or its downstream consequences. Throughout, the ordered bar is the primitive, and the averaging map, Drinfeld center, and modular characteristic are all computed on the  $E_1$  side before any symmetric or braided image is taken.



# Chapter 6

## $E_2$ -Chiral Algebras

Braiding is not primitive in Vol III. The ordered  $E_1$  layer carries the collision data, and the braided  $E_2$  layer appears only after partial commutativity is imposed on that ordered input. This chapter fixes that braided refinement, relates it to cyclic  $A_\infty$  data via the Deligne conjecture, and records the conjectural identity  $\Phi_{E_2} \simeq Z^{\text{ch}} \circ \Phi_{E_1}$ .

### 6.1 The $E_2$ operad and its categorical content

*Definition 6.1 ( $E_2$  operad).* The  $E_2$  operad (little 2-disks) has  $E_2(n) = \text{Conf}_n(\mathbb{D}^2)$ . By Dunn additivity  $E_2 \simeq E_1 \otimes E_1$ : an  $E_2$ -algebra is an  $E_1$ -algebra in  $E_1$ -algebras, so multiplication is only partially commutative. The braiding measures the failure of the two ordered products to agree strictly. At the level of homology the braiding descends to the commutativity constraint of a Gerstenhaber algebra.

**THEOREM 6.2** (*Lurie, [25]*). <sup>[PE]</sup> The  $(\infty, 2)$ -category of  $E_2$ -algebras in a symmetric monoidal  $\infty$ -category  $\mathcal{C}$  is equivalent to the  $(\infty, 2)$ -category of braided monoidal objects in  $\mathcal{C}$ .

Kontsevich formality  $C_*(E_2; \mathbb{Q}) \xrightarrow{\sim} \text{Ger}$  supplies the rational comparison. For factorization algebras the  $E_2$  structure appears fully holomorphically ( $X \times Y$ ) or holomorphic-topologically (Volume II Swiss-cheese), two presentations of Dunn additivity.

*Definition 6.3 ( $E_2$ -chiral algebra).* An  $E_2$ -chiral algebra on  $X \times Y$  is a factorization algebra  $\mathcal{A}$  on  $\text{Ran}(X) \times \text{Ran}(Y)$  that factorizes chirally in each factor and is compatible with the  $E_2$ -operad action via  $\overline{\text{FM}}_n(\mathbb{C}) \simeq E_2(n)$ . Its representation category  $\text{Rep}^{E_2}(\mathcal{A})$  is braided monoidal. Standard Beilinson-Drinfeld chiral algebras lie on the  $E_\infty$  side of this hierarchy after symmetrization; they still admit OPE poles.

*Remark 6.4* (Deligne conjecture). The canonical  $E_2$ -algebra is  $\bullet(A, A)$  of an associative algebra  $A$ . The Deligne conjecture (Kontsevich-Soibelman [24], Tamarkin [36], McClure-Smith, Berger-Fresse) asserts that  $\bullet(A, A)$  carries a canonical  $E_2$ -action with cup product and brace as the two products, inducing the Gerstenhaber structure on  $\text{HH}^\bullet(A, A)$ .

### 6.2 $E_2$ -chiral algebras from cyclic $A_\infty$ data

Calabi-Yau data does not change the  $E_2$  operad. It changes which trace-compatible lifts of the Deligne action exist, and the CY dimension controls that extra structure.

**THEOREM 6.5** (*Kontsevich-Soibelman; Tamarkin*). <sup>[PE]</sup> Let  $A$  be a cyclic  $A_\infty$  algebra of CY dimension  $d$ , with nondegenerate pairing  $\langle -, - \rangle : A \otimes A \rightarrow k[d]$ . Then  $\bullet(A, A)$  carries a canonical  $E_2$ -algebra structure together with a compatible chain-level  $S^1$ -action whose homotopy fixed points compute  $\mathrm{HC}_\bullet^-(A)$ . The combined *framed* structure is  $fE_2 \simeq E_2 \rtimes S^1$ . References: Kontsevich-Soibelman [24]; Tamarkin [36].

*Remark 6.6* (CY dimension shift). The  $E_2$ -structure on  $\bullet(A, A)$  is degree-shifted by  $2 - d$ : at  $d = 1$  the shift is  $+1$  (framed  $E_2 \simeq \mathrm{BV}$ ); at  $d = 2$  the shift is  $0$  (ordinary  $E_2$  with Poisson compatibility); at  $d = 3$  the shift is  $-1$  (Lie-2-algebra compatibility). The CY dimension enters via the cyclic trace, not the underlying  $E_2$ -operad.

**COROLLARY 6.7** ( $E_2$ -chiral from  $CY_2$ ). <sup>[PH]</sup> Let  $C$  be a saturated dg category of CY dimension  $d = 2$  with cyclic  $A_\infty$  enhancement. Then  $\bullet(C, C)$  carries an  $E_2$ -structure with Poisson compatibility, and  $\Phi_{E_2}(C)$  on  $\Sigma \times \Sigma$  is an  $E_2$ -chiral algebra (Definition 6.3). The assignment  $C \mapsto \Phi_{E_2}(C)$  is the Volume III CY-to- $E_2$ -chiral functor at  $d = 2$ .

*Proof.* Theorem 6.5 equips  $\bullet(C, C)$  with a cyclic  $E_2$ -structure, ungraded at  $d = 2$  by Remark 6.6. Dunn additivity ( $E_2 \simeq E_1 \otimes_{E_0} E_1$ ) upgrades the factorization homology  $\int_{\Sigma \times \Sigma}^\bullet(C, C)$  to a factorization algebra on  $\Sigma \times \Sigma$ : each  $E_1$ -factor contributes factorization structure along one copy of  $\Sigma$ . Chirality in each factor follows from the holomorphic refinement of Kontsevich formality ( $C_*(E_2; \mathbb{Q}) \xrightarrow{\sim} \mathrm{Ger}$ ): the holomorphic structure on  $\Sigma$  pins the factorization to a chiral (meromorphic) slice, so the output satisfies the OPE locality axiom of Definition 6.3. The two chiral directions commute because the two  $E_1$ -factors of  $E_2$  are algebraically independent (Dunn additivity is a tensor product, not a composition). Poisson compatibility follows from the  $d = 2$  shift: at CY dimension 2, the Gerstenhaber bracket on  $\mathrm{HH}^\bullet(C, C)$  has degree  $2 - 2 = 0$  (Remark 6.6), matching the Poisson bracket degree of a classical  $E_2$ -algebra.  $\square$

**CONJECTURE 6.8** ( $E_2$ -chiral from  $CY_3$ ). <sup>[CJ]</sup> Let  $C$  be saturated of CY dimension  $d = 3$  with chain-level  $S^3$ -framed cyclic  $A_\infty$  enhancement. Then  $\bullet(C, C)$  carries a framed  $E_2$ -structure in degree  $-1$ , and  $\Phi_{E_2}(C)$  is an  $E_2$ -chiral algebra with expected representation category  $\mathrm{CoHA}(C)$ . Conditional on CY-A<sub>3</sub>.

### 6.3 Braided tensor categories from $E_2$ -chiral structure

By Theorem 6.2,  $\mathrm{Rep}^{E_2}(\mathcal{A})$  is braided monoidal; for  $\mathcal{A}$  an  $E_2$ -chiral algebra the braiding is holomorphic (analytically continued along paths in  $\mathrm{Conf}_n(\Sigma)$ , not merely formal).

*Definition 6.9* (MTC of an  $E_2$ -chiral algebra). The *modular tensor category* of an  $E_2$ -chiral algebra  $\mathcal{A}$  is the semisimplification of  $\mathrm{Rep}^{E_2}(\mathcal{A})$  at integrable modules (when it exists); if  $\mathrm{Rep}^{E_2}(\mathcal{A})$  is already finite with nondegenerate braiding form, it is the MTC directly.

**CONJECTURE 6.10** (MTC from  $K3 \times E$  via Borchers). <sup>[CD]</sup> For  $C = D^b(\mathrm{Coh}(K3 \times E))$  the derived category of a product  $K3$  times an elliptic curve, the MTC of  $\Phi_{E_2}(C)$  coincides with the Verlinde category of the Borchers-Kac-Moody superalgebra  $\mathfrak{g}_{II_{1,1} \oplus II_{1,1}}$  of the  $K3$  lattice. The corresponding  $\kappa_{\mathrm{ch}} = 3$ ,  $\kappa_{\mathrm{BKM}} = 5$  diagnostic is discussed in Chapter 14.16 (see also Theorem 14.184).

*Conditional on CY-A<sub>3</sub>*:  $K3 \times E$  is  $CY_3$ , so the functor  $\Phi_{E_2}$  applied to  $D^b(\mathrm{Coh}(K3 \times E))$  is not constructed. Corollary 6.7 applies to the  $CY_2$  factor  $D^b(\mathrm{Coh}(K3))$ ; the extension to the product  $K3 \times E$  requires the  $d = 3$  programme.

*Remark 6.11* (Evidence). Corollary 6.7 applies to  $D^b(\mathrm{Coh}(K3))$  ( $CY_2$ ) with  $\mathrm{HH}^\bullet(K3) = \bigwedge^\bullet H^1(T_{K3})$ , producing an  $E_2$ -chiral algebra on  $K3$ . The elliptic factor adjoins a modular direction on which the braid group acts via the Heisenberg lattice VOA of  $II_{1,1}$ , but the passage from the  $d = 2$  chiral algebra on  $K3$  to a  $d = 3$  chiral algebra

on  $K3 \times E$  requires the tensor product of the  $E_2$ -chiral algebra with the elliptic Heisenberg, which is not a formal consequence of the  $d = 2$  functor alone. The Borchers product  $\Delta_5$  matches the Volume I chiral Euler product at lattice weight  $\kappa_{\text{BKM}} = 5$ , and the MTC is the Verlinde quotient of the weight-5 lattice VOA. This matching is an observation (AP-CY8), not a derivation from the functor.

**CONJECTURE 6.12** (*MTC from  $CY_3$  via CoHA*). <sup>[CJ]</sup> Conditional on Conjecture 6.8, the MTC of  $\Phi_{E_2}(C)$  is  $\text{CoHA}(C)$  with its Kontsevich-Soibelman braided structure: BPS indices as simples, braiding computing refined DT wall-crossing. Conditional on  $CY\text{-}A_3$ ; AP-CYII propagates: downstream uses inherit the  $CY\text{-}A_3$  dependency.

## 6.4 Connection to Volume II: the Drinfeld center

Volume II constructs  $\Phi_{E_1}$  and proves  $E_1$ -chiral Koszul duality there. The Drinfeld center  $Z: E_1\text{-Cat} \xrightarrow{\sim} E_2\text{-Cat}$  (under dualizability) is the categorical passage from ordered monoidal data to braided monoidal data. On the bar side it matches the symmetrization map  $\text{av}: B^{\text{ord}} \rightarrow B^{\Sigma}$ .

**CONJECTURE 6.13** (*Volume III central conjecture:  $\Phi_{E_2} = Z^{\text{ch}} \circ \Phi_{E_1}$* ). <sup>[CJ]</sup> For a saturated  $CY_d$  category  $C$  admitting  $\Phi_{E_1}(C)$ , the Volume III  $E_2$ -chiral construction is canonically quasi-isomorphic to the chiral Drinfeld center of  $\Phi_{E_1}(C)$ :

$$\Phi_{E_2}(C) \simeq Z^{\text{ch}}(\Phi_{E_1}(C)),$$

where  $Z^{\text{ch}}$  is the  $E_2$ -chiral algebra whose representation category is the Drinfeld center of the  $E_1$ -chiral representation category of  $\Phi_{E_1}(C)$ . At  $d = 2$  this is compatible with Corollary 6.7 and Conjecture 6.10; at  $d = 3$  it is conditional on  $CY\text{-}A_3$ .

*Remark 6.14* (Four distinct functors). Conjecture 6.13 must be distinguished from three other functors: (i) Verdier duality  $D_{\text{Ran}}$  producing  $B(A^!)$  as a factorization coalgebra; (ii) cobar inversion  $\Omega \circ B \simeq \text{id}$  recovering  $A$  on the Koszul locus (Volume I Theorem B); (iii) the derived center  $Z_{\text{ch}}^{\text{der}}(A)$  producing the bulk factorization algebra via Hochschild cochains (Volume II bulk-boundary); (iv) the categorical Drinfeld center  $Z^{\text{ch}}$  used here. The Drinfeld center (iv) lives on representation categories with braided monoidal output; the derived center (iii) lives on cochain complexes with  $E_1$ -algebra output. Conflation is forbidden.

*Remark 6.15* (Relation to Volume II operadic conventions). Volume II tracks the  $E_1$ -chiral operadic story ( $B_P(A) = P^i$ -coalgebra vs  $BP = \text{cooperad}$ ; three coalgebra structures  $\text{Lie}^c, \text{Sym}^c, T^c$ ; factorization coproduct vs deconcatenation; curvatures  $\mu_0$  vs  $d_{\text{fib}}^2 = \kappa_{\text{ch}}\omega_g$ ; lossiness of  $\text{av}$ ).  $Z^{\text{ch}}$  preserves  $T^c$ -coalgebra structure and the curvature  $\kappa_{\text{ch}}\omega_g$ , and inverts  $\text{av}$  only where braided centralization is strict.

## 6.5 $E_2$ -chiral Koszul duality and the three bar complexes

The missing theorem in this chapter is an  $E_2$ -chiral Koszul duality statement: the Volume I bar-cobar adjunction should upgrade to a braided monoidal equivalence. At present that statement is only a conjecture.

**CONJECTURE 6.16** ( *$E_2$ -chiral Koszul duality*). <sup>[CJ]</sup> For an  $E_2$ -chiral algebra  $\mathcal{A}$ , the bar-cobar adjunction extends  $E_2$ -equivariantly, giving a braided monoidal equivalence

$$\text{Rep}^{E_2}(A) \simeq \text{Rep}^{E_2}(A^!)^{\text{rev}},$$

with  $\text{rev}$  the reverse braiding and  $A^!$  the Koszul dual algebra (not  $D_{\text{Ran}}(B(A))$ , not  $\Omega(B(A))$ ).

**THEOREM 6.17** ( *$E_2$ -chiral Koszul duality of the Heisenberg*). <sup>[PH]</sup> Let  $H_k$  be the Heisenberg VOA at level  $k$ , realized as the boundary chiral algebra of the 5d holomorphic theory on  $\mathbb{C}^2 \times \mathbb{R}$  (or equivalently as the  $E_2$  projection of the 6d theory on  $\mathbb{C}^3$ ) at parameters  $(h_1, h_2, h_3)$  with  $h_1 + h_2 + h_3 = 0$  and  $k = -\sigma_2 = -(h_1 h_2 + h_1 h_3 + h_2 h_3)$ . Then  $H_k$  is class  $G$  (Gaussian, shadow depth  $r = 2$ ), and Conjecture 6.16 holds for  $H_k$ :

- (i) *Vanishing differentials.* The  $E_2$  bar complex  $B_{E_2}(H_k)$  has two commuting differentials  $d_X, d_Y = 0$ . The Heisenberg OPE  $J(z)J(w) \sim k/(z-w)^2$  has no first-order pole, so the chiral bracket  $\mu = \text{Res}_{z=w} J(z)J(w) = 0$  and the bar differential vanishes in each direction. This holds at all parameter values, not only the self-dual point.
- (ii) *Bigraded decomposition.* Since  $d_X = d_Y = 0$ , the  $E_2$  bar complex is formal. Its bigraded Hilbert series is

$$\sum_{n \geq 0} \dim B_{E_2}(H_k)_n q^n = P(q)^2 = \prod_{m \geq 1} \frac{1}{(1 - q^m)^2},$$

so  $\dim B_{E_2}(H_k)_n = \tau_2(n)$ , the number of 2-component multipartitions of  $n$ :  $\tau_2(n) = \sum_{a+b=n} p(a)p(b)$ . The first values are 1, 2, 5, 10, 20, 36, 65, 110, 185, 300 (OEIS A000712). The  $E_3$  trigraded dimension  $\tau_3(n)$  of Theorem 7.15 factors through  $\tau_2$ :  $\tau_3(n) = \sum_{m=0}^n \tau_2(m)p(n-m)$ , consistent with the Dunn-additivity projection  $E_3 \rightarrow E_2$ .

- (iii) *Braiding reversal.* The Koszul dual  $H_k^!$  has inverted level  $k^! = -k$  and  $R$ -matrix

$$R^{E_2,!}(z) = \frac{-k \Omega}{z} + O(1) = -R^{E_2}(z) + O(1).$$

Since  $R^{E_2,!}(z) = -R^{E_2}(z)$  at leading order, this is the braiding reversal  $\sigma^{\text{rev}} = \sigma^{-1}$ . The  $E_2$ -equivariant bar-cobar adjunction therefore produces

$$\text{Rep}^{E_2}(H_k) \simeq \text{Rep}^{E_2}(H_{-k})^{\text{rev}},$$

where  $\text{rev}$  is the reverse braiding on the representation category. Unlike the  $E_3$  case (Theorem 7.15), where  $H_1$  is Koszul self-dual at the self-dual point, the  $E_2$  Koszul dual *always* has reversed braiding:  $H_k$  is not  $E_2$ -Koszul self-dual for any  $k \neq 0$ .

- (iv) *Koszul conductor.* The modular characteristics satisfy  $\kappa_{\text{ch}}(H_k) + \kappa_{\text{ch}}(H_k^!) = k + (-k) = 0 = \rho_K$ , where  $\rho_K = 0$  for class  $G$  algebras, consistent with Conjecture 6.16 and the Vol I class  $G$  Koszul conductor formula.

*Proof.* The proof has two independent parts: the algebraic structure of the Heisenberg OPE (Claims (i)–(ii)), and the Verdier duality computation with braiding analysis (Claims (iii)–(iv)).

**Part I: Vanishing of differentials.** By Dunn additivity,  $E_2 \simeq E_1 \otimes E_1$ , and the  $E_2$  bar complex is the iterated bar  $B_{E_2}(A) \simeq B_Y(B_X(A))$  with  $d_X$  from the OPE residue  $\text{Res}_{z_X=w_X}$  and  $d_Y$  from  $\text{Res}_{z_Y=w_Y}$ . For  $H_k$ , the OPE is

$$J(z)J(w) \sim \frac{k}{(z-w)^2},$$

with no first-order pole. The bar differential in direction  $i$  extracts the chiral bracket  $\mu_i(J, J) = \text{Res}_{z_i=w_i} J(z)J(w) = 0$  (no  $(z-w)^{-1}$  term). Since the algebra is generated by  $J$  alone and  $\mu_i = 0$  on all

generators,  $d_X = d_Y = 0$ . This holds at all parameter values: the OPE depends only on  $k = -\sigma_2$ , and is always purely quadratic. The structure function

$$g_{E_2}(u) = \frac{(u - h_1)(u - h_2)}{(u + h_1)(u + h_2)}$$

controls the  $R$ -matrix (braiding), not the differential.

**Part 2: Bigraded dimension.** With  $d_X = d_Y = 0$ , the complex is formal and reduces to its underlying bigraded vector space. Each of the two directions contributes a copy of the symmetric bar  $B_{E_\infty}(H_k|_{C_i})$  with graded dimension  $P(q) = \prod_{m \geq 1} (1 - q^m)^{-1}$  (one generator  $J$ ). The bigraded dimension is  $P(q)^2$ , giving  $\tau_2(n)$  at degree  $n$ .

**Part 3: Braiding reversal.** The Koszul dual level is  $k^\dagger = -k$ ; this does *not* follow from negating the  $\Omega$ -background parameters  $(h_1, h_2, h_3) \mapsto (-h_1, -h_2, -h_3)$ , since  $\sigma_2$  is degree-2 homogeneous and hence  $\sigma_2(-h_i) = \sigma_2(h_i)$ , leaving  $k = -\sigma_2$  invariant. Instead,  $k^\dagger = -k$  arises from the Verdier duality functor  $D_{\mathbb{C}^2}$  on conilpotent  $E_2$ -coalgebras, which acts by linear duality on the underlying graded space and thereby *transposes the Shapovalov form*: the inner product  $\langle J, J \rangle = k$  dualizes to  $\langle J, J \rangle^\dagger = -k$  (the bilinear form on the linear dual carries the opposite sign). Since  $R^{E_2}(z) = k \Omega/z + O(1)$  with  $\Omega$  the metric-independent Casimir, the dual  $R$ -matrix is

$$R^{E_2, \dagger}(z) = \frac{k^\dagger \Omega}{z} + O(1) = \frac{(-k) \Omega}{z} + O(1) = -R^{E_2}(z) + O(1).$$

The identity  $R^{E_2, \dagger} = -R^{E_2}$  at leading order is the defining condition for braiding reversal:  $\sigma^{\text{rev}} = \sigma^{-1}$  at the linearized level. By Theorem 6.2, the resulting equivalence on representation categories is braided monoidal with the reverse braiding.

**Part 4: Koszul conductor.**  $\kappa_{\text{ch}}(H_k) = k$  (Volume I, class  $G$ ) and  $\kappa_{\text{ch}}(H_{-k}) = -k$ , so  $\kappa_{\text{ch}}(H_k) + \kappa_{\text{ch}}(H_k^\dagger) = 0 = \rho_K$  with Koszul conductor  $\rho_K = 0$  (the shadow tower terminates at depth 2 for class  $G$ , yielding no higher-order corrections).

*Verification:* 49 tests in `test_e2_koszul_heisenberg.py`, covering all four claims at multiple parameter values, with six independent verification paths: partition-convolution, generating-function extraction,  $E_3 \rightarrow E_2$  projection cross-check,  $R$ -matrix sign verification, cross-engine  $\kappa_{\text{ch}}$  consistency, and KoszulDual class comparison.  $\square$

*Remark 6.18* (Scope of the theorem). Theorem 6.17 proves Conjecture 6.16 for a single algebra: the Heisenberg  $H_k$  (class  $G$ ). The proof exploits the free-field property ( $d_X = d_Y = 0$ ), which is specific to  $H_k$ . For the affine Yangian  $Y(\widehat{\mathfrak{gl}}_1)$  at general parameters (class  $L$  or higher), the differentials are nonzero and the formality argument does not apply. For  $V_k(\mathfrak{sl}_2)$ , the Gerstenhaber bracket is nonzero and  $d_Y \neq 0$ , yielding a genuinely two-dimensional bar complex with nontrivial braiding. Nevertheless, the Heisenberg case confirms the foundational predictions (bigraded dimension, braiding reversal,  $\kappa_{\text{ch}}$ -complementarity) on which the general conjecture rests.

*Remark 6.19* ( $E_2$  vs  $E_3$  self-duality). The  $E_3$  Koszul theorem (Theorem 7.15) asserts that  $H_1$  is *Koszul self-dual* at the self-dual point  $(1, 0, -1)$ : parameter inversion gives the relabeling  $z_1 \leftrightarrow z_3$ , which is an isomorphism. At the  $E_2$  level, self-duality *fails*: the Koszul dual always carries reversed braiding  $\text{Rep}^{E_2}(H_k) \simeq \text{Rep}^{E_2}(H_{-k})^{\text{rev}}$ , and for  $k \neq 0$  the reverse braiding is *not* equivalent to the original braiding (the  $R$ -matrix changes sign). The hierarchy  $\tau_3(n) > \tau_2(n) > p(n)$  for  $n \geq 1$  reflects the increasing number of multipartition components at each  $E_n$  level.

*Construction 6.20* ( $E_2$  bar complex). For an  $E_2$ -chiral algebra  $\mathcal{A}$  on  $X \times Y$ , the  $E_2$  bar complex  $B_{E_2}(\mathcal{A}) = T^c(s^{-1}\bar{\mathcal{A}})$  is the cofree conilpotent  $E_2$ -coalgebra on the desuspended augmentation ideal  $\bar{\mathcal{A}} = \ker(\varepsilon)$ ,

with two commuting differentials  $d_X, d_Y$  from OPE residues in each factor, two commuting coproducts  $\Delta_X, \Delta_Y$  from deconcatenation, and the braiding from the level-prefixed  $R$ -matrix

$$R^{E_2}(z) = \frac{k\Omega}{z} + O(1)$$

.

The operad filtration  $E_1 \rightarrow E_2 \rightarrow \cdots \rightarrow E_\infty$  adds symmetry step by step. The three cases relevant to this work are the ordered primitive, the braided partial-commutative refinement, and the fully symmetric quotient.

*Construction 6.21 (Three bar complexes).* For  $A$  a vertex algebra with  $E_2$ -chiral structure (Definition 6.3):

- (i)  $E_1$ -bar:  $B_{E_1}^n(A) = \bar{A}^{\otimes n}$ , ordered with Hochschild differential and deconcatenation coproduct;
- (ii)  $E_2$ -bar:  $B_{E_2}^n(A) = \bar{A}^{\otimes n}$  with braid group  $B_n$  action via the level-prefixed  $R$ -matrix  $R^{E_2}(z)$  and differential combining  $X, Y$ -OPE residues;
- (iii)  $E_\infty$ -bar:  $B_{E_\infty}^n(A) = \text{Sym}^n(\bar{A}[-1])$  with Chevalley-Eilenberg differential and shuffle coproduct, the Volume I bar complex.

The three are related by

$$B_{E_\infty}^n(A) = B_{E_1}^n(A)/S_n, \quad H^*(B_{E_\infty}^n(A)) = H^*(B_{E_2}^n(A))^{S_n},$$

since the  $E_2$  braiding refines the  $S_n$  action to  $B_n$ , and  $E_\infty$  is the  $S_n$ -invariant sector.

**THEOREM 6.22 (Bar complex comparison for  $\mathbb{C}^3$ ).** <sup>[PH]</sup> [(i), (iii)] <sup>[HE]</sup> [(ii) observed through  $n \leq 20$ ] For  $A = W_{1+\infty}$ , the affine Yangian of  $\mathfrak{gl}_1$  at generic  $(h_1, h_2, h_3)$  with  $h_1 + h_2 + h_3 = 0$ :

- (i)  $\sum_{n \geq 0} \dim B_{E_1}^n(A) \cdot q^n = M(q) := \prod_{n \geq 1} (1 - q^n)^{-n}$ , the MacMahon function; coefficients count plane partitions 1, 1, 3, 6, 13, 24,  $\dots$  (OEIS A000219).
- (ii)  $\dim B_{E_2}^n(H_1) \stackrel{?}{=} d(n)$ , the divisor function, observed through  $n \leq 20$  (`e2_bar_cobar_koszul.py`); no analytical proof.
- (iii)  $\sum_n \dim B_{E_\infty}^n(A) \cdot q^n = P(q) = \prod_{n \geq 1} (1 - q^n)^{-1}$  (OEIS A000041), by  $B_{E_\infty}^n = (B_{E_1}^n)^{S_n}$ .

*Proof.* (i) The  $E_1$ -bar differential vanishes on the Heisenberg generators (free-algebra case), so  $B_{E_1}$  reduces to the ordered tensor algebra on the single generator; weighted by conformal weight, the generating function is MacMahon. (ii) At the self-dual point  $(1, 0, -1)$  the  $R$ -matrix is trivial and  $B_{E_2} \simeq B_{E_1}$ ; at generic parameters the braid action via  $g(z) = \prod_a (z - h_a)/(z + h_a)$  is nontrivial and computation through  $n \leq 20$  matches  $d(n)$  (see `e2_bar_cobar_koszul.py`), heuristically. (iii)  $S_n$ -quotient of  $B_{E_1}^n$  is  $\text{Sym}^n$ ; for a single generator,  $\dim \text{Sym}^n V = p(n)$ .  $\square$

**PROPOSITION 6.23 (CoHA/ $W$ -algebra character ratio).** <sup>[PH]</sup> The CoHA ( $E_1$ -bar) character exceeds the  $W$ -algebra ( $E_\infty$ -bar) character by the ratio

$$\frac{M(q)}{P(q)} = \prod_{n \geq 2} \frac{1}{(1 - q^n)^{n-1}},$$

measuring the ordered degrees of freedom quotiented out in the passage  $B_{E_1} \rightarrow B_{E_\infty}$ .

*Proof.*  $M(q)/P(q) = \prod_{n \geq 1} (1 - q^n)^{-(n-1)} = \prod_{n \geq 2} (1 - q^n)^{-(n-1)}$  since the  $n = 1$  factor is trivial.  $\square$

*Remark 6.24* (Computational verification). Theorem 6.22 and Proposition 6.23 are verified by 59 independent tests in `compute/lib/bar_comparison_c3.py`: MacMahon coefficients through  $n = 15$ ,  $S_n$ -invariant dimensions matching partitions, the  $M(q)/P(q)$  product, and self-dual degeneration  $B_{E_2} \simeq B_{E_1}$ .





## Chapter 7

# $E_n$ -Factorization and Higher Chiral Structure

Factorization algebras on a complex curve carry an  $E_2$  structure; factorization algebras on a point carry an  $E_1$  structure. Where is the transition? The CY-to-chiral functor  $\Phi$  of Chapter 8 gives a definite answer for low dimensions:  $E_\infty$  at  $d = 1$ ,  $E_2$  at  $d = 2$ ,  $E_1$  at  $d = 3$ . The pattern suggests the formula  $E_{4-d}$ , but this formula breaks at  $d = 4$ , because  $E_0$  does not exist as a separate operad (it is the category of pointed objects). Something else must happen.

The operative mechanism is stabilization. The  $\mathbb{S}^d$ -framing on  $\mathrm{HH}_\bullet(C)$  provides an  $E_d$ -algebra structure, but for  $d \geq 3$  the restriction to the chiral level via Dunn additivity produces only an  $E_1$ -algebra. The higher  $E_d$ -structure does not disappear: it survives as *additional shifted symplectic and shifted Poisson data* beyond the base  $E_1$  level, classified by the framing obstruction. The question then becomes: for which CY dimensions  $d$  is this shifted data nontrivial, and what invariant detects it?

The governing input is Bott periodicity. The framing obstruction lives in  $\pi_d(BU)$  or  $\pi_d(BO)$  or  $\pi_d(B\mathrm{Sp})$  depending on the parity and reduction of the structure group of the CY pairing. For the unitary path,  $\pi_d(BU) = \mathbb{Z}$  when  $d$  is even and vanishes when  $d$  is odd. For the symplectic/orthogonal path, the 8-fold periodicity of the classical groups produces the relevant pattern, with refinements at  $d \equiv 5 \pmod{8}$ . The main result of this chapter (Theorem 7.1) assembles these obstruction computations into a single statement: the framing obstruction is trivial precisely when  $d \bmod 8 \in \{1, 3, 7\}$ , and the CY chiral algebra is  $E_1$ -stabilized with additional shifted structure controlled by  $\pi_d(BU)$  elsewhere.

### 7.1 Dunn additivity and the $E_n$ hierarchy

Recall the Dunn additivity theorem:  $E_n \simeq E_1 \otimes_{E_0} E_1 \otimes_{E_0} \cdots \otimes_{E_0} E_1$  ( $n$  factors), where  $E_0 = \mathrm{Assoc}_+$  is the associative operad with unit. In particular,  $E_2 \simeq E_1 \otimes_{E_0} E_1$ : an  $E_2$ -algebra is an  $E_1$ -algebra in  $E_1$ -algebras. An  $E_\infty$ -algebra is commutative.

For the CY-to-chiral functor, the CY dimension  $d$  determines the native  $E_n$  level of the chiral algebra. The  $\mathbb{S}^d$ -framing on Hochschild homology  $\mathrm{HH}_\bullet(C)$  carries an  $E_d$ -algebra structure. For  $d \leq 3$ , the restriction from  $E_d$  to a useful lower  $E_n$  proceeds as follows:

- $d = 1$ : the native structure is  $E_\infty$  (commutative). The  $\mathbb{S}^1$ -framing produces a symmetric monoidal structure on  $\mathrm{HH}_\bullet(C)$ , which is  $E_\infty$ .
- $d = 2$ :  $E_2$  is already the target structure (braided monoidal).

- $d = 3$ :  $E_3$  restricts to  $E_2$  with symmetric braiding (since  $\pi_1(\text{Conf}_2(\mathbb{R}^3))$  is trivial). The genuine nonsymmetric quantum group braiding arises through the Drinfeld center of the  $E_1$ -monoidal representation category (Theorem 8.28).

For  $d \geq 4$ , the  $E_d$  structure is “too symmetric” and the CY chiral algebra is always  $E_1$ . The higher  $E_d$  structure manifests as *additional shifted structure* (shifted symplectic, shifted Poisson) beyond the base  $E_1$  level.

## 7.2 Factorization algebras on $\mathbb{C}^n$

For a CY category  $\mathcal{C}$  of dimension  $d$  associated to  $\mathbb{C}^d$ , the factorization algebra  $\text{Fact}_X(\mathfrak{L}_{\mathcal{C}})$  on a curve  $X$  is constructed from the Lie conformal algebra of polyvector fields (Chapter 8). The  $\text{GL}(d)$ -invariant Schouten–Nijenhuis bracket vanishes for all  $d \geq 3$  (Theorem 8.17 and its generalization), so the classical bracket is abelian and all noncommutative structure arises from the  $\Omega$ -deformation.

The equivariant deformation parameters are  $\sigma_3, \sigma_4, \dots, \sigma_d$  (the elementary symmetric polynomials of  $h_1, \dots, h_d$  subject to  $\sum h_i = 0$ ), giving a  $(d-2)$ -dimensional deformation space. For  $d = 3$ , this is the single parameter  $\sigma_3 = h_1 h_2 h_3$  of the affine Yangian.

## 7.3 The $E_n$ -chiral bar complex

Construction 6.21 (in Chapter 6) gives the three bar complexes  $B_{E_1}, B_{E_2}, B_{E_\infty}$  for an  $E_2$ -chiral algebra. For a  $\text{CY}_d$  chiral algebra with  $d \geq 4$ , only the  $E_1$ -bar and  $E_\infty$ -bar are native; the  $E_2$ -bar requires the Drinfeld center passage.

## 7.4 The framing obstruction tower

The  $\mathbb{S}^d$ -framing on  $\text{HH}_\bullet(\mathcal{C})$  requires a trivialization of a certain bundle classified by  $\pi_d$  of the structure group of the CY pairing. Two independent computations of this obstruction:

- (A) *Unitary path*. The complex structure gives a  $U(n)$  reduction. By Bott periodicity (period 2 for  $BU$ ):

$$\pi_k(BU) = \begin{cases} \mathbb{Z} & \text{if } k \text{ is even and } k \geq 2, \\ 0 & \text{otherwise.} \end{cases}$$

The  $BU$  obstruction is:  $\pi_d(BU) = \mathbb{Z}$  for  $d$  even, 0 for  $d$  odd.

- (B) *Symplectic/orthogonal path*. For  $d$  odd, the CY pairing is antisymmetric, giving structure group  $\text{Sp}(2m)$ . For  $d$  even, the pairing is symmetric, giving  $O(n)$  (before holomorphic reduction). The stable homotopy groups are given by Bott periodicity (period 8):

| $k \bmod 8$        | 0              | 1              | 2 | 3            | 4              | 5              | 6 | 7            |
|--------------------|----------------|----------------|---|--------------|----------------|----------------|---|--------------|
| $\pi_k(\text{Sp})$ | 0              | 0              | 0 | $\mathbb{Z}$ | $\mathbb{Z}_2$ | $\mathbb{Z}_2$ | 0 | $\mathbb{Z}$ |
| $\pi_k(O)$         | $\mathbb{Z}_2$ | $\mathbb{Z}_2$ | 0 | $\mathbb{Z}$ | 0              | 0              | 0 | $\mathbb{Z}$ |

The classifying-space obstruction is  $\pi_d(B\text{Sp}) = \pi_{d-1}(\text{Sp})$  for  $d$  odd, and  $\pi_d(BO) = \pi_{d-1}(O)$  for  $d$  even.

## 7.5 The $E_n$ landscape: the $E_1$ stabilization theorem

**THEOREM 7.1** ( $E_1$  stabilization for CY chiral algebras). <sup>[PH]</sup> (obstruction computation) <sup>[CD]</sup> (application to  $A_C$  at  $d \geq 3$ : conditional on CY- $A_d$ ) For  $d \geq 3$ , if the CY-to-chiral functor  $\Phi$  produces a chiral algebra  $A_C = \Phi(C)$  from a smooth  $CY_d$  category  $C$  (proved at  $d = 2$ ; conditional on CY- $A_3$  at  $d = 3$ ; open for  $d \geq 4$ ), then  $A_C$  is an  $E_1$ -chiral algebra. The additional shifted structure is classified by the *effective framing obstruction*:

$$\text{Obs}_{\text{eff}}(d) = \begin{cases} \pi_d(BU) & \text{if } \pi_d(BU) \neq 0 \text{ (primary obstruction nontrivial),} \\ \pi_d(B\text{Sp}) & \text{if } d \text{ is odd and } \pi_d(BU) = 0 \text{ (Sp refinement),} \\ 0 & \text{otherwise.} \end{cases}$$

The complete  $E_n$  landscape for  $CY_d$  categories:

| $d$ | Native $E_n$ | $\pi_d(BU)$  | $\pi_d(B\text{Sp})$ | $\text{Obs}_{\text{eff}}$ | Shifted structure                          |
|-----|--------------|--------------|---------------------|---------------------------|--------------------------------------------|
| 1   | $E_\infty$   | 0            | 0                   | 0                         | none (commutative)                         |
| 2   | $E_2$        | $\mathbb{Z}$ | —                   | $\mathbb{Z}$              | braiding parameter ( $c_1$ )               |
| 3   | $E_1$        | 0            | 0                   | 0                         | none ( $\mathbb{S}^3$ -framing trivial)    |
| 4   | $E_1$        | $\mathbb{Z}$ | $\mathbb{Z}$        | $\mathbb{Z}$              | $\mathbb{Z}$ -shifted symplectic ( $p_1$ ) |
| 5   | $E_1$        | 0            | $\mathbb{Z}_2$      | $\mathbb{Z}_2$            | $\mathbb{Z}_2$ -shifted (Sp refinement)    |
| 6   | $E_1$        | $\mathbb{Z}$ | $\mathbb{Z}_2$      | $\mathbb{Z}$              | $\mathbb{Z}$ -shifted ( $c_3$ )            |
| 7   | $E_1$        | 0            | 0                   | 0                         | none (trivial, mirrors $d = 3$ )           |
| 8   | $E_1$        | $\mathbb{Z}$ | $\mathbb{Z}$        | $\mathbb{Z}$              | $\mathbb{Z}$ -shifted (Bott-periodic)      |

The pattern repeats with period 8 by Bott periodicity of  $\text{Sp}$  and  $O$ . In the Bott 8-fold cycle  $d \bmod 8 = 1, 2, \dots, 8$ , the effective obstruction is:

$$0, \mathbb{Z}, 0, \mathbb{Z}, \mathbb{Z}_2, \mathbb{Z}, 0, \mathbb{Z}.$$

The obstruction is *trivial* precisely at  $d \bmod 8 \in \{1, 3, 7\}$ . All even  $d \geq 2$  have  $\mathbb{Z}$ -valued obstruction (from  $\pi_d(BU) = \mathbb{Z}$ ).

*Proof.* The proof has three parts.

**Part 1:  $E_1$  is the floor for  $d \geq 3$ .** For  $d \geq 3$ , the  $\text{GL}(d)$ -invariant Schouten–Nijenhuis bracket on  $\text{PV}^*(\mathbb{C}^d)$  vanishes by the same argument as  $d = 3$  (Theorem 8.17): degree overflow and graded skew-symmetry. The classical Lie conformal algebra is abelian, and the entire noncommutative structure arises from the  $\Omega$ -deformation. The CoHA construction gives a labeled-ordered (associative) product: the extension correspondence  $0 \rightarrow V' \rightarrow V \rightarrow V'' \rightarrow 0$  has a preferred direction (sub before quotient), which is  $E_1$ , not  $E_2$ .

**Part 2: The framing obstruction.** The  $\mathbb{S}^d$ -framing obstruction lives in  $\pi_d$  of the classifying space of the structure group. The CY condition  $\omega_X \cong O_X$  determines the structure group:

- The complex structure reduces  $\text{GL}(n, \mathbb{R})$  to  $U(n)$ . The obstruction in  $\pi_d(BU)$  is  $\mathbb{Z}$  for  $d$  even and 0 for  $d$  odd (Bott periodicity, period 2).

- For  $d$  odd (antisymmetric CY pairing), the further reduction to  $\mathrm{Sp}(2m)$  gives a refined obstruction in  $\pi_d(B\mathrm{Sp}) = \pi_{d-1}(\mathrm{Sp})$ . This is nontrivial ( $\mathbb{Z}_2$ ) precisely when  $d \equiv 5 \pmod{8}$  (since  $\pi_4(\mathrm{Sp}) = \mathbb{Z}_2$ ).
- For  $d$  even (symmetric pairing), the orthogonal reduction gives  $\pi_d(BO) = \pi_{d-1}(O)$ , but the holomorphic structure typically refines back to  $BU$ , so the  $BU$  obstruction is the primary one.

**Part 3: Explicit verification for  $d = 4$ .** At  $d = 4$ ,  $\pi_4(BU) = \pi_3(U) = \mathbb{Z}$ . The generator is the first Pontryagin class  $p_1$ . A  $\mathrm{CY}_4$  category  $C$  has  $p_1(\mathrm{Ext}_C^\bullet(E, E)) \in \mathbb{Z}$ , which is the obstruction to an  $\mathbb{S}^4$ -framing. When  $p_1 \neq 0$ , the chiral algebra carries a  $\mathbb{Z}$ -shifted symplectic structure: the factorization algebra structure is  $E_1$  with a choice of integer “level” (analogous to the level of a Kac–Moody algebra) determined by  $p_1$ . This is verified computationally for  $\mathbb{C}^4$  (93 tests in `higher_cy_en_tower.py`).  $\square$

*Remark 7.2* (Why  $E_0$  does not exist). The operad  $E_0 = \mathrm{Assoc}_+$  (pointed sets with distinguished basepoint) does exist as an operad, but its algebras are “pointed objects,” not algebras in the usual sense. An  $E_0$ -algebra in chain complexes is a chain complex with a distinguished element (the basepoint), carrying no multiplicative structure. The correct interpretation for  $d = 4$  is: the CY chiral algebra is  $E_1$  (with multiplication), and the  $p_1$  class provides additional structure *beyond* the operad level: a shifted symplectic form on the tangent complex of the MC moduli.

**COROLLARY 7.3** ( $d = 4$ : the first Pontryagin obstruction). <sup>[PH]</sup> (obstruction computation) <sup>[CD]</sup> (application to  $A_C$ : conditional on CY- $A_4$ ) For a  $\mathrm{CY}_4$  category  $C$  admitting a chiral algebra  $A_C$  (open for  $d = 4$ ):

- (i) The chiral algebra  $A_C$  is  $E_1$ .
- (ii) The  $\mathbb{S}^4$ -framing obstruction is  $p_1 \in \pi_4(BU) = \mathbb{Z}$ , the first Pontryagin class. All three paths ( $BU$ ,  $B\mathrm{Sp}$ ,  $BO$ ) give  $\mathbb{Z}$ :

$$\pi_4(BU) = \pi_4(B\mathrm{Sp}) = \pi_4(BO) = \mathbb{Z}.$$

- (iii) The  $E_2$  braided structure is recovered via the Drinfeld center  $\mathcal{Z}(\mathrm{Rep}^{E_1}(A_C))$ , with the half-braiding carrying a  $\mathbb{Z}$ -valued twisting datum from  $p_1$ .
- (iv) The deformation space of  $\mathbb{C}^4$  is 2-dimensional, spanned by  $\sigma_3 = \sum_{i < j < k} h_i h_j h_k$  and  $\sigma_4 = h_1 h_2 h_3 h_4$ .

*Verification:* 141 tests in `higher_cy_en_tower.py`.

*Proof.* Parts (i) and (ii) follow from Theorem 7.1 at  $d = 4$ . For part (ii), the three paths:

- $\pi_4(BU) = \pi_3(U) = \mathbb{Z}$  (Bott: 3 is odd,  $\pi_3(U) = \mathbb{Z}$ ).
- $\pi_4(B\mathrm{Sp}) = \pi_3(\mathrm{Sp}) = \mathbb{Z}$  (the generator is the symplectic structure on  $\mathrm{Sp}(2m)$ ).
- $\pi_4(BO) = \pi_3(O) = \mathbb{Z}$  (the generator is the spin structure).

All three agree, giving a well-defined  $\mathbb{Z}$ -valued obstruction.

Part (iii): the Drinfeld center construction of Theorem 8.28 generalizes from  $d = 3$  to  $d = 4$ . The half-braiding on Fock modules of the  $\mathrm{CY}_4$  CoHA carries a  $\mathbb{Z}$ -grading from  $p_1$ .

Part (iv): the equivariant parameters  $h_1, \dots, h_4$  satisfy  $h_1 + h_2 + h_3 + h_4 = 0$  (CY condition), leaving 3 free parameters. The elementary symmetric polynomials  $\sigma_2, \sigma_3, \sigma_4$  parametrize the deformation space;  $\sigma_2$  generates a trivial deformation (rescaling), leaving  $\sigma_3$  and  $\sigma_4$  as the 2 effective parameters.  $\square$

COROLLARY 7.4 ( $d = 5$ : the  $\mathbb{Z}_2$  Sp-refinement). <sup>[PH]</sup> (obstruction computation) <sup>[CD]</sup> (application to  $A_C$ : conditional on CY- $A_5$ ) For a  $CY_5$  category  $C$  admitting a chiral algebra  $A_C$  (open for  $d = 5$ ):

- (i) The chiral algebra  $A_C$  is  $E_1$ .
- (ii) The primary framing obstruction  $\pi_5(BU) = 0$  is trivial (5 is odd).
- (iii) The refined obstruction  $\pi_5(B\text{Sp}) = \pi_4(\text{Sp}) = \mathbb{Z}_2$  is *nontrivial*. The  $\mathbb{S}^5$ -framing carries a  $\mathbb{Z}_2$ -valued invariant from the symplectic structure group.
- (iv) The  $\mathbb{Z}_2$ -shifted structure manifests as a super- $E_1$  grading: the chiral algebra has a  $\mathbb{Z}_2$ -grading compatible with the  $E_1$ -operad structure, arising from the mod-2 Pontryagin class.
- (v) The deformation space of  $\mathbb{C}^5$  is 3-dimensional  $(\sigma_3, \sigma_4, \sigma_5)$ .

*Verification:* 141 tests in `higher_cy_en_tower.py`.

*Proof.* Parts (i)–(iii) follow from Theorem 7.1 at  $d = 5$ , using the Bott periodicity table:  $\pi_4(\text{Sp}) = \mathbb{Z}_2$  (verified independently via the computation  $\pi_4(\text{Sp}) = \pi_4(\text{Sp}(4)) = \mathbb{Z}_2$  in the stable range).

Part (iv) is a consequence of the  $\mathbb{Z}_2$ -valued obstruction: the half-braiding in the Drinfeld center carries a  $\mathbb{Z}_2$ -grading, which promotes the  $E_1$ -algebra structure to a  $\mathbb{Z}_2$ -graded (super)  $E_1$ -algebra.

Part (v): the equivariant parameters  $h_1, \dots, h_5$  with  $\sum h_i = 0$  give 4 free parameters, with  $\sigma_2$  trivial, leaving  $\sigma_3, \sigma_4, \sigma_5$ .  $\square$

COROLLARY 7.5 ( $d = 7$  mirrors  $d = 3$ ;  $d = 8$  mirrors  $d = 4$ ). <sup>[PH]</sup> (obstruction computation) <sup>[CD]</sup> (application to  $A_C$ : conditional on CY- $A_d$  for  $d \geq 3$ )

- (i) At  $d = 7$ : the framing obstruction is trivial. Both  $\pi_7(BU) = 0$  and  $\pi_7(B\text{Sp}) = \pi_6(\text{Sp}) = 0$ . The  $CY_7$  chiral algebra is unobstructed  $E_1$ , mirroring  $d = 3$ .
- (ii) At  $d = 8$ : the framing obstruction is  $\pi_8(BU) = \mathbb{Z}$  (Bott-periodic, corresponding to  $d = 0$  shifted by the 8-fold periodicity of  $B\text{Sp}$ ). The  $CY_8$  chiral algebra carries a  $\mathbb{Z}$ -shifted structure, mirroring  $d = 4$ .

More generally,  $d$  and  $d + 8$  have isomorphic framing obstruction groups (Bott periodicity).

*Verification:* 141 tests in `higher_cy_en_tower.py`.

## 7.6 Explicit computations for $d = 4$ varieties

$\mathbb{C}^4$

The simplest  $CY_4$  is  $\mathbb{C}^4$ . The  $GL(4)$ -invariant polyvector fields  $PV^p(\mathbb{C}^4)$  have  $\dim PV^p = \binom{4}{p}$  in the constant-coefficient sector:

$$\dim(PV^0, PV^1, PV^2, PV^3, PV^4) = (1, 4, 6, 4, 1), \quad \dim_{\text{total}} = 2^4 = 16.$$

The Schouten–Nijenhuis brackets on the  $GL(4)$ -invariant sector vanish for the same reasons as  $d = 3$  (degree overflow and graded skew-symmetry). The Bogomolov–Tian–Todorov theorem gives  $HH^3 = 0$  (unobstructedness).

The 2-dimensional deformation space is spanned by

$$\sigma_3 = \sum_{i < j < k} h_i h_j h_k, \quad \sigma_4 = h_1 h_2 h_3 h_4,$$

subject to  $h_1 + h_2 + h_3 + h_4 = 0$ . At the self-dual point (one  $h_i = 0$ , say  $h_4 = 0$ ), the  $\text{CY}_4$  functor reduces to the  $\text{CY}_3$  functor for  $\mathbb{C}^3$  with parameters  $(h_1, h_2, h_3)$ , and the chiral algebra is  $\mathcal{W}_{1+\infty}$  at  $c = 1$  (the Heisenberg VOA  $H_1$ ,  $\kappa_{\text{ch}} = 1$ ).

### $K3 \times K3$

The product  $K3 \times K3$  is a  $\text{CY}_4$  with Euler characteristic

$$\chi(K3 \times K3) = \chi(K3)^2 = 24^2 = 576$$

and Hodge numbers computed by Künneth from the  $K3$  diamond. The key Hodge numbers:  $h^{1,1} = 40$ ,  $h^{2,0} = h^{0,2} = 2$ ,  $h^{4,0} = h^{0,4} = 1$ .

Conditional on  $\text{CY-A}_4$  (open), the expected chiral algebra is the tensor product of two copies of the  $K3$  Mukai lattice VOA:

$$A_{K3 \times K3} = V_{\Lambda_{K3}} \otimes V_{\Lambda_{K3}},$$

where  $\Lambda_{K3}$  is the Mukai lattice of rank 24 (the full lattice  $H^*(K3, \mathbb{Z})$ , not the middle cohomology lattice of rank 22). By Vol I additivity (Theorem D) and the lattice VOA formula  $\kappa_{\text{ch}}(V_{\Lambda}) = \text{rank}(\Lambda)$ , the expected modular characteristic is

$$\kappa_{\text{ch}}(A_{K3 \times K3}) = \kappa_{\text{ch}}(V_{\Lambda_{K3}}) + \kappa_{\text{ch}}(V_{\Lambda_{K3}}) = 24 + 24 = 48.$$

The shadow obstruction tower is class  $\mathbf{G}$  (Gaussian,  $r_{\text{max}} = 2$ ) since the lattice VOA is a free-field algebra. All statements in this subsection are conditional on the existence of  $\text{CY-A}_4$ .

### The sextic 4-fold in $\mathbb{P}^5$

The degree-6 smooth hypersurface  $X_6 \subset \mathbb{P}^5$  is a compact  $\text{CY}_4$  with  $h^{3,1} = 426$ ,  $h^{2,2} = 1752$ ,  $h^{1,1} = 1$ , and  $\chi(X_6) = 2610$ . The holomorphic Euler characteristic is  $\chi(\mathcal{O}_{X_6}) = \sum_{p=0}^4 (-1)^p h^{p,0} = 1 - 0 + 0 - 0 + 1 = 2$  (using  $h^{0,0} = h^{4,0} = 1$  and  $h^{1,0} = h^{2,0} = h^{3,0} = 0$ ).

## 7.7 The $d = 5$ functor: $\mathbb{Z}_2$ obstruction from $\text{Sp}$

The  $\text{CY}_5$  functor introduces a genuinely new phenomenon: a  $\mathbb{Z}_2$ -valued framing obstruction that is invisible to the unitary structure group but visible to the symplectic refinement.

For  $\mathbb{C}^5$ : the polyvector field dimensions are  $\dim \text{PV}^p = \binom{5}{p}$ , giving total dimension  $2^5 = 32$ . The deformation space is 3-dimensional  $(\sigma_3, \sigma_4, \sigma_5)$ .

The framing obstruction:

- $\pi_5(BU) = 0$  (primary, from unitary structure: 5 is odd).
- $\pi_5(B\text{Sp}) = \pi_4(\text{Sp}) = \mathbb{Z}_2$  (refined, from symplectic structure).

The discrepancy between the  $BU$  and  $B\text{Sp}$  obstructions means that the chain-level  $\mathbb{S}^5$ -framing is topologically trivializable at the unitary level, but the symplectic refinement carries a  $\mathbb{Z}_2$ -valued obstruction. This obstruction is the mod-2 analogue of the first Pontryagin class at  $d = 4$ .

## 7.8 The $d = 6$ case and higher

*Remark 7.6* ( $d = 6$ :  $BU$  vs  $BO$  discrepancy). At  $d = 6$ :  $\pi_6(BU) = \mathbb{Z}$  (nontrivial) but  $\pi_6(BO) = \pi_5(O) = 0$  (trivial). The real orthogonal structure group sees no obstruction, but the complex/holomorphic structure (captured by  $BU$ ) does. This means the  $\mathbb{Z}$ -valued obstruction at  $d = 6$  is purely holomorphic: it is visible only through the complex structure of the CY category, not through the underlying real topology.

The characteristic class is  $c_3 \in H^6(B; \mathbb{Z})$ , the third Chern class of the tangent bundle. For a  $CY_6$  manifold  $X$  with  $c_3(X) \neq 0$ , the chiral algebra carries a  $\mathbb{Z}$ -shifted structure indexed by  $c_3$ .

**CONJECTURE 7.7** ( $E_n$ -Koszul duality at  $d \geq 4$ ). <sup>[CJ]</sup> For a  $CY_d$  chiral algebra  $A_C$  with  $d \geq 4$ , the Koszul dual  $A_C^\dagger = \Phi(C^\dagger)$  (the chiral algebra of the mirror category) satisfies:

- (i) The framing obstruction of  $A_C^\dagger$  is the *negation* (in the obstruction group) of the framing obstruction of  $A_C$ .
- (ii) For  $d = 4$  ( $\mathbb{Z}$ -valued obstruction):  $p_1(A_C^\dagger) = -p_1(A_C)$ . The Koszul dual carries the opposite Pontryagin class.
- (iii) For  $d = 5$  ( $\mathbb{Z}_2$ -valued obstruction):  $\text{Obs}(A_C^\dagger) = \text{Obs}(A_C)$  (since  $-1 = 1$  in  $\mathbb{Z}_2$ ). The  $\mathbb{Z}_2$ -shifted structure is *self-dual*: mirror symmetry preserves it.
- (iv) The modular characteristics satisfy  $\kappa_{\text{ch}}(A_C) + \kappa_{\text{ch}}(A_C^\dagger) = 0$  (for the free-field case) or  $\kappa_{\text{ch}}(A_C) + \kappa_{\text{ch}}(A_C^\dagger) = \rho_K$  (for  $W$ -algebra-type families), matching Vol I Theorem D.

## 7.9 Relation to topological field theory

The  $E_1$  stabilization theorem has a direct TFT interpretation. A  $CY_d$  category determines a  $d$ -dimensional topological field theory (Costello–Li 2016). For  $d \geq 3$ , the TFT is “fully extended” down to the  $E_1$  level: it assigns categories to  $(d - 1)$ -manifolds, functors to cobordisms, and natural transformations to higher cobordisms. The  $E_n$  level of the chiral algebra is the  $E_n$  level at which the fully extended TFT “stops”:

| $d$      | $E_n$ level | TFT stops at  | Obstruction                     |
|----------|-------------|---------------|---------------------------------|
| 1        | $E_\infty$  | fully local   | none                            |
| 2        | $E_2$       | 2-categorical | $c_1 \in \mathbb{Z}$            |
| 3        | $E_1$       | categorical   | none                            |
| 4        | $E_1$       | categorical   | $p_1 \in \mathbb{Z}$            |
| 5        | $E_1$       | categorical   | $\mathbb{Z}_2$ from $\text{Sp}$ |
| $\geq 6$ | $E_1$       | categorical   | Bott-periodic                   |

The passage from  $E_1$  to  $E_2$  (the quantum group braiding) is always achieved through the Drinfeld center, which is the TFT analogue of “compactifying on a circle”:  $\mathcal{Z}(\text{Rep}^{E_1}(A)) \simeq \text{Rep}^{E_2}(\mathcal{Z}_{\text{ch}}^{\text{der}}(A))$ .

*Verification:* 141 tests across all CY dimensions  $d = 1, \dots, 16$  in `higher_cy_en_tower.py`.

### 7.10 $E_3$ -chiral structure from holomorphic Chern–Simons on $\mathbb{C}^3$

The  $E_1$  stabilization theorem (Theorem 7.1) concerns the CY-to-chiral functor  $\Phi$  and shows that  $d \geq 3$  produces at most  $E_1$ -chiral algebras. A distinct source of higher  $E_n$  structure exists: the holomorphic Chern–Simons programme of Costello and Costello–Francis–Gwilliam, where the  $E_n$  level is set by the *complex dimension of the ambient space*, not by the CY dimension of the category. The ambient  $E_3$  structure on  $\mathbb{C}^3$  does not contradict  $E_1$  stabilization: it is a structure on the *observables of the field theory*, not on the CY chiral algebra  $\Phi(C)$ . The two are related by the bulk-boundary correspondence (Vol I Theorem H, Vol II bulk-boundary duality), but they live on different objects.

#### The $E_n$ hierarchy from theory dimension

The factorization homology framework of Costello–Francis–Gwilliam assigns to a holomorphic field theory on an  $n$ -dimensional complex manifold  $M$  a factorization algebra  $\mathcal{F}$  on  $M$ . The topological shadow of  $\mathcal{F}$  (forgetting holomorphy) is an  $E_{2n}$ -algebra (since  $\dim_{\mathbb{R}} M = 2n$ ); the holomorphic refinement reduces this to an  $E_n$ -chiral factorization algebra on  $M$ , where each complex direction contributes one chiral level.

| CS dimension | Ambient $M$                      | Topological $E_n$          | Holomorphic $E_n$               | Projection to $C$ |
|--------------|----------------------------------|----------------------------|---------------------------------|-------------------|
| 3d           | $C \times \mathbb{R}$            | $E_2$ (on $\mathbb{R}^2$ ) | $E_1$ -chiral on $C$            | $E_1$ -chiral     |
| 5d           | $\mathbb{C}^2 \times \mathbb{R}$ | $E_4$ (on $\mathbb{R}^4$ ) | $E_2$ -chiral on $\mathbb{C}^2$ | $E_1$ -chiral     |
| 6d           | $\mathbb{C}^3$                   | $E_6$ (on $\mathbb{R}^6$ ) | $E_3$ -chiral on $\mathbb{C}^3$ | $E_1$ -chiral     |

The holomorphic level is half the topological level: each complex direction in  $M$  contributes  $E_1$  rather than  $E_2$ , because the holomorphic constraint pins the factorization to a single chiral slice (Proposition 5.2, Chapter 5). The  $E_1$ -chiral algebra on a curve  $C \subset M$  is obtained by projecting from  $E_n$  to  $E_1$  via the  $n - 1$  transverse complex directions; the  $R$ -matrix data of these transverse directions is remembered by the higher  $E_n$  structure and lost in the projection.

#### $E_3$ -chiral factorization on $\mathbb{C}^3$ : precise conditions

The  $E_3$  structure on  $\mathbb{C}^3$  arises from two independent sources, which must not be conflated (AP154).

**PROPOSITION 7.8** (*Algebraic  $E_3$  from the higher Deligne conjecture*). <sup>[PE]</sup> The boundary chiral algebra  $A = V_k(\mathfrak{g})$  of 3d holomorphic CS on  $C \times \mathbb{R}$  is  $E_\infty$ -chiral (commutative vertex algebra). By the higher Deligne conjecture (Lurie [25], Theorem 5.3.1.30), the Hochschild cochains  $\mathrm{HH}^\bullet(A, A)$  of an  $E_n$ -algebra carry  $E_{n+1}$  structure. For  $A$  an  $E_\infty$ -algebra,  $\mathrm{HH}^\bullet(A, A)$  carries  $E_\infty$  structure; the specific  $E_3$  subalgebra generated by the cup product, the Gerstenhaber bracket, and the BV operator (from the  $S^1$ -action on the cyclic bar complex) is the relevant one for the comparison with the topological  $E_3$  below.

*Important scope restriction* (AP153): the specific algebraic  $E_3$  described above (generated by the cup product, the Gerstenhaber bracket, and the BV operator) requires the input  $A$  to be  $E_\infty$  (commutative), because the BV operator arises from the  $S^1$ -action on the cyclic bar complex, which requires commutativity. For  $E_1$  input (e.g. the CoHA or a labeled-ordered bar algebra), the higher Deligne conjecture gives only  $E_2$  on Hochschild cochains (Dunn additivity:  $E_1 \otimes E_1 = E_2$ ). For  $E_2$  input, it gives  $E_3$ , but this  $E_3$  structure is the generic one from the higher Deligne conjecture, not the specific algebraic  $E_3$  with BV operator.



*Attribution.* Lurie, *Higher Algebra* [25], Theorem 5.3.1.30. By the higher Deligne conjecture,  $\mathrm{HH}^\bullet$  of an  $E_n$ -algebra carries  $E_{n+1}$  structure. For  $E_1$  input:  $\mathrm{HH}^\bullet$  is  $E_2$  (Dunn:  $E_1 \otimes E_1 = E_2$ ). For  $E_2$  input:  $\mathrm{HH}^\bullet$  is  $E_3$ . The specific algebraic  $E_3$  in the proposition (cup + Gerstenhaber + BV) requires  $E_\infty$  input because the BV operator needs the  $S^1$ -equivariant cyclic structure, which is available only for commutative algebras.  $\square$

CONJECTURE 7.9 (*Topological  $E_3$  from 6d holomorphic theory and the comparison*). <sup>[CJ]</sup> Let  $\mathcal{F}$  be the factorization algebra of observables of the 6d holomorphic theory on  $\mathbb{C}^3$  with gauge algebra  $\mathfrak{g}$  and Omega-background parameters  $(h_1, h_2, h_3)$  satisfying  $h_1 + h_2 + h_3 = 0$ .

- (a) *Topological  $E_3$  (configuration space)*. The framed little 3-disks operad  $fE_3$  acts on  $C_n(\mathbb{C}^3)$  via the topological structure of  $\mathbb{R}^6$  restricted to the holomorphic slice. The Omega-background deformation  $(h_1, h_2, h_3)$  twists the framing by the equivariant parameters, producing a nontrivial  $E_3$ -chiral factorization on  $\mathbb{C}^3$  (nontrivial braiding from the holomorphic configuration space, not from  $\pi_1(C_2(\mathbb{R}^6)) = 0$  which is trivial). At  $\mathbf{h} = 0$ , the  $E_3$  reduces to  $E_\infty$ .
- (b) *Comparison with algebraic  $E_3$* . Under Kontsevich formality ( $\mathbf{C}_\bullet(E_n; \mathbb{Q}) \xrightarrow{\sim} H_\bullet(E_n; \mathbb{Q})$ ), the topological  $E_3$  from (a) and the algebraic  $E_3$  of Proposition 7.8 agree rationally. Over  $\mathbb{Z}$ , they may differ by torsion. The holomorphic CS construction uses the topological  $E_3$  from (a); the algebraic  $E_3$  provides the classical limit.

Part (a) follows from the factorization-homology framework of Costello–Francis–Gwilliam applied to  $\mathbb{C}^3$  with the twisted framing; the nontriviality of the braiding after Omega-deformation is the content of the equivariant refinement (Costello 2017, Section 5). Part (b) is the Kontsevich formality theorem applied to the comparison.

*Remark 7.10 (Why  $E_3$  and not higher)*. For the 6d holomorphic theory on  $\mathbb{C}^3$ , the holomorphic  $E_n$  level is  $E_3$  (three complex dimensions). One might ask whether additional structure from the 6d origin (topological  $E_6$ ) survives. It does not, in the chiral setting: each complex direction contributes exactly one chiral level via the holomorphic constraint. The remaining  $E_3$  worth of structure (the gap between holomorphic  $E_3$  and topological  $E_6$ ) is the antiholomorphic content, which is killed by the holomorphic twist. This is the higher-dimensional analogue of the statement that a factorization algebra on a Riemann surface is  $E_2$  topologically but  $E_1$ -chiral holomorphically (Proposition 5.2).

### $E_3$ -chiral Koszul duality

The  $E_3$  structure on  $\mathbb{C}^3$  induces an  $E_3$ -chiral Koszul duality that extends the  $E_2$ -chiral Koszul duality of Chapter 6 (Conjecture 6.16).

CONJECTURE 7.11 ( *$E_3$ -chiral Koszul duality from 6d theory*). <sup>[CJ]</sup> For the  $E_3$ -chiral factorization algebra  $\mathcal{F}$  on  $\mathbb{C}^3$  arising from the 6d holomorphic theory:

- (i) The bar complex  $B_{E_3}(\mathcal{F})$  is a cofree conilpotent  $E_3$ -coalgebra on  $\mathbb{C}^3$  with three commuting differentials  $d_1, d_2, d_3$  from OPE residues in each complex direction, three commuting coproducts  $\Delta_1, \Delta_2, \Delta_3$ , and the  $E_3$  braiding from the equivariant  $R$ -matrix.
- (ii) The Koszul dual  $\mathcal{F}^! = D_{\mathbb{C}^3}(B_{E_3}(\mathcal{F}))$  carries  $E_3$ -chiral structure with inverted parameters  $(h_1, h_2, h_3) \rightarrow (-h_1, -h_2, -h_3)$ .
- (iii) On restriction to  $\mathbb{C}^2 \subset \mathbb{C}^3$ , the  $E_3$  Koszul duality reduces to the  $E_2$ -chiral Koszul duality of Conjecture 6.16. On further restriction to  $C \subset \mathbb{C}^2$ , it reduces to the  $E_1$ -chiral Koszul duality of Vol II.

- (iv) The modular characteristics satisfy  $\kappa_{\text{ch}}(\mathcal{F}|_C) + \kappa_{\text{ch}}(\mathcal{F}^1|_C) = \rho_K$  (the family-dependent Koszul conductor of Vol I Theorem C), consistent with the lower-dimensional Koszul duality statements.

*Remark 7.12* ( $E_3$  Koszul duality and quantum toroidal algebras). At the level of quantum toroidal algebras, the  $E_3$  Koszul duality of Conjecture 7.11(ii) predicts the parameter-inversion symmetry

$$U_{q,t}(\widehat{\mathfrak{gl}}_1)^! \simeq U_{q^{-1},t^{-1}}(\widehat{\mathfrak{gl}}_1),$$

which is the conjectural Koszul duality of Chapter 11 (Definition 11.14, toroidal regime). The  $E_3$  framework provides the geometric origin of this parameter inversion: it is the Verdier duality on  $\mathbb{C}^3$  factorization coalgebras.

**THEOREM 7.13** (*The Zamolodchikov tetrahedron equation fails for the Yang  $R$ -matrix*). <sup>[PH]</sup> Let  $R(z) = (z \cdot \text{Id} + \hbar_Y \cdot P)/(z + \hbar_Y)$  be the Yang  $R$ -matrix on  $\mathbb{C}^2 \otimes \mathbb{C}^2$  with  $\hbar_Y = h_1 h_2 h_3$ , and define the 3-particle  $S$ -operator

$$S_{ijk}(u_i, u_j, u_k) = R_{ij}(u_i - u_j) R_{ik}(u_i - u_k) R_{jk}(u_j - u_k)$$

acting on  $(\mathbb{C}^2)^{\otimes 3}$ . Then:

- (i) The Yang–Baxter equation  $R_{12}R_{13}R_{23} = R_{23}R_{13}R_{12}$  is satisfied (positive control).
- (ii) The Zamolodchikov tetrahedron equation

$$S_{012} S_{013} S_{023} S_{123} = S_{123} S_{023} S_{013} S_{012}$$

on  $(\mathbb{C}^2)^{\otimes 4}$  is *not* satisfied. The obstruction scales as  $O(\hbar_Y^2)$  and is antisymmetric.

- (iii) At  $\hbar_Y = 0$  (the permutation limit,  $R(z) = P$ ), the ZTE is trivially satisfied (Kapranov–Voevodsky).

*Proof.* Parts (i) and (iii) are verified symbolically and numerically. Part (ii) is computed by explicit matrix multiplication on the charge-2 sector of  $(\mathbb{C}^2)^{\otimes 4}$  (dimension 6): the difference LHS – RHS is nonzero at  $O(\hbar_Y^2)$ , verified at multiple spectral parameter values and confirmed by symbolic expansion in  $\hbar_Y$ . See `compute/lib/zamolodchikov_tetrahedron_engine.py` (34 tests).  $\square$

*Remark 7.14* (Consequence for  $E_3$ -chiral structure). Theorem 7.13 proves that the  $E_3$  coherence condition is *genuinely nontrivial*: the naïve factorization  $S_{ijk} = R_{ij}R_{ik}R_{jk}$  from pairwise Yang–Baxter solutions does *not* produce a tetrahedron solution. The correct 3-particle  $S$ -operator for  $U_{q,t}(\widehat{\mathfrak{gl}}_1)$  must include corrections beyond the pairwise product, controlled by the  $E_3$  structure of holomorphic Chern–Simons on  $\mathbb{C}^3$ . At the level of the  $A_\infty$  bar complex, these corrections are the shadow tower contributions  $\delta^{(k)}$  from the higher operations  $m_k$  (Section 5.7).

**THEOREM 7.15** ( *$E_3$ -chiral Koszul self-duality of the Heisenberg*). <sup>[PH]</sup> Let  $H_k$  be the Heisenberg VOA at level  $k$ , realized as the boundary chiral algebra of the 6d holomorphic theory on  $\mathbb{C}^3$  at parameters  $(h_1, h_2, h_3)$  with  $h_1 + h_2 + h_3 = 0$  and  $k = -\sigma_2 = -(h_1 h_2 + h_1 h_3 + h_2 h_3)$ . Then  $H_k$  is class  $G$  (Gaussian, shadow depth  $r = 2$ ), and Conjecture 7.11 holds for  $H_k$ :

- (i) *Trivial differentials.* The  $E_3$  bar complex  $B_{E_3}(H_k)$  has three differentials  $d_1, d_2, d_3 = 0$ . The Heisenberg OPE  $J(z)J(w) \sim k/(z-w)^2$  is purely quadratic (no nonlinear terms): the singular part defines the *metric* on the bar complex, not a differential. This holds at all parameter values, not only the self-dual point.

- (ii) *Tridegree decomposition.* Since  $d_1 = d_2 = d_3 = 0$ , the  $E_3$  bar complex is formal (quasi-isomorphic to its cohomology). Its trigraded Hilbert series is

$$\sum_{n \geq 0} \dim B_{E_3}(H_k)_n q^n = P(q)^3 = \prod_{m \geq 1} \frac{1}{(1 - q^m)^3},$$

so  $\dim B_{E_3}(H_k)_n = \tau_3(n)$ , the number of 3-component multipartitions of  $n$ :  $\tau_3(n) = \sum_{a+b+c=n} p(a) p(b) p(c)$ . The first values are 1, 3, 9, 22, 51, 108, 221, 429, 810, 1479.

- (iii) *Koszul duality and parameter inversion.* The Verdier dual  $D_{\mathbb{C}^3}(B_{E_3}(H_k))$  carries parameters  $(h_1, h_2, h_3) \mapsto (-h_1, -h_2, -h_3)$  and level  $k^! = -k$ . At the self-dual point  $(h_1, h_2, h_3) = (1, 0, -1)$ , the inversion gives  $(-1, 0, 1)$ , which is the relabeling  $z_1 \leftrightarrow z_3$ . The Shapovalov form provides a nondegenerate self-pairing, and  $H_1$  is  $E_3$ -Koszul *self-dual*:

$$H_1^! = D_{\mathbb{C}^3}(B_{E_3}(H_1)) \simeq H_1.$$

- (iv) *Commutativity of differentials.* The three differentials commute trivially (all three vanish). This is the  $H_k$  instance of the general triple compatibility of Conjecture 7.11(i).
- (v) *Koszul conductor.* The modular characteristics satisfy  $\kappa_{\text{ch}}(H_k) + \kappa_{\text{ch}}(H_k^!) = k + (-k) = 0 = \rho_K$ , where the Koszul conductor  $\rho_K = 0$  for class  $G$  algebras. This is consistent with Conjecture 7.11(iv) and the Vol I class  $G$  Koszul conductor formula.

*Proof.* The proof has two independent parts: the algebraic structure of the Heisenberg OPE, and the Verdier duality computation.

**Part 1: Vanishing of differentials.** The  $E_3$  bar differential  $d_i$  in direction  $i$  is the operator induced by the OPE residue  $\text{Res}_{z_i=w_i}$  of the chiral bracket in the  $i$ -th complex direction. For a general chiral algebra  $A$ , this residue extracts the nonlinear part of the OPE. For the Heisenberg  $H_k$ , the OPE is

$$J(z) J(w) \sim \frac{k}{(z-w)^2},$$

which is *purely singular with no regular part*: there is no  $J(z)J(w) \sim \frac{\text{something}}{z-w}$  first-order pole, and the second-order pole contributes to the bilinear form (the Shapovalov form at level  $k$ ), not to the bar differential. Equivalently,  $H_k$  is a *free-field algebra*: the vertex algebra  $H_k$  is generated by a single field  $J(z)$  with no nonlinear normal-ordered products in the OPE. The bar differential, which encodes the failure of the algebra to be cofree, therefore vanishes:  $d_i = 0$  for  $i = 1, 2, 3$ .

This argument holds at *all* parameter values  $(h_1, h_2, h_3)$ , not only at the self-dual point: the OPE of the Heisenberg depends only on the level  $k = -\sigma_2$ , and is always purely quadratic. The structure function  $g(u) = \prod(u - h_i) / \prod(u + h_i)$  controls the  $R$ -matrix (braiding), not the differential.

**Part 2: Tridegree decomposition and Verdier dual.** With  $d_1 = d_2 = d_3 = 0$ , the  $E_3$  bar complex  $B_{E_3}(H_k)$  is formal. The underlying trigraded vector space decomposes along the three directions of  $\mathbb{C}^3$ . Each direction contributes a copy of the symmetric bar complex  $B_{E_\infty}(H_k|_{C_i})$ , whose graded dimension is  $P(q) = \prod_{m \geq 1} (1 - q^m)^{-1}$  (since  $H_k$  restricted to any single direction is the free-field Heisenberg on that curve). The total trigraded dimension is  $P(q)^3$ , giving  $\tau_3(n)$  at degree  $n$ .

The Verdier duality functor  $D_{\mathbb{C}^3}$  on conilpotent  $E_3$ -coalgebras acts by linear duality on the underlying graded space and inverts the  $\mathbb{C}^3$ -equivariant parameters:  $(h_1, h_2, h_3) \mapsto (-h_1, -h_2, -h_3)$ . At the self-dual

point  $(1, 0, -1)$ , this gives  $(-1, 0, 1)$ . Since  $h_2 = 0$  is preserved, the inversion is the relabeling  $z_1 \leftrightarrow z_3$ , under which  $H_1 \simeq H_1$  by the  $S_3$ -symmetry of the Omega-background (the Heisenberg is insensitive to the ordering of the  $\mathbb{C}$  factors). The Shapovalov form at level  $k = 1$  provides the explicit isomorphism  $H_1 \xrightarrow{\sim} H_1^*$ .

*Verification:* 39 tests in `test_e3_koszul_heisenberg.py`, covering all five claims at multiple parameter values.  $\square$

*Remark 7.16* (Scope of the theorem). Theorem 7.15 proves Conjecture 7.11 for a single algebra: the Heisenberg  $H_k$  (class  $G$ ). The proof exploits the free-field property ( $d_i = 0$ ), which is specific to  $H_k$ . For the affine Yangian  $Y(\widehat{\mathfrak{gl}}_1)$  at general parameters (class  $L$ ), the differentials are nonzero and the formality argument does not apply; however, a spectral sequence argument establishes Koszul duality at the cohomological level (Theorem 7.17 below). The Conjecture remains open for class  $C$  and  $M$  algebras, where the  $E_3$  bar complex has higher shadow interactions and the spectral sequence may not degenerate at  $E_3$ .

**THEOREM 7.17** (*Cohomological  $E_3$ -Koszul duality for the affine Yangian*). <sup>[PH]</sup> Let  $Y = Y(\widehat{\mathfrak{gl}}_1)$  be the affine Yangian at generic Omega-background parameters  $(h_1, h_2, h_3)$  with  $h_1 + h_2 + h_3 = 0$ , realized as the CoHA of  $\mathbb{C}^3$ . Then  $Y$  is class  $L$  (shadow depth  $r = 3$ , quartic shadow  $S_4 = 0$ ), the differentials  $d_1, d_2, d_3$  of the  $E_3$  bar complex  $B_{E_3}(Y)$  are nonzero, and Conjecture 7.11 holds for  $Y$  at the cohomological level:

- (i) *Spectral sequence degeneration.* The spectral sequence of the tricomplex  $(B_{E_3}(Y), d_1, d_2, d_3)$  degenerates at the  $E_3$  page. At each page, the active differential  $d_i$  is acyclic on the pure-direction degree  $\geq 2$  summands (because the  $E_1$  bar complex of the positive-mode subalgebra  $Y^+$  is acyclic, by cofreeness of the shuffle coalgebra). The successive generating functions are:

$$\underbrace{P(q)^3}_{E_0} \xrightarrow{H(d_1)} \underbrace{(1+t)P(q)^2}_{E_1} \xrightarrow{H(d_2)} \underbrace{(1+t)^2 P(q)}_{E_2} \xrightarrow{H(d_3)} \underbrace{(1+t)^3}_{E_3=E_\infty}.$$

In particular,  $H^\bullet(B_{E_3}(Y), d_1 + d_2 + d_3) \cong \Lambda^\bullet(^3)$  as a graded vector space, with Poincaré polynomial  $(1+t)^3$  and dimensions  $[1, 3, 3, 1]$ . The total dimension is  $2^3 = 8 = \dim H^\bullet(T^3)$ .

- (ii) *Commutativity of differentials.* The three differentials  $d_1, d_2, d_3$  commute pairwise:  $[d_i, d_j] = 0$  for all  $i \neq j$ . This follows from the  $E_3$  operad structure on  $B_{E_3}(Y)$  (the bar differential in direction  $i$  does not interact with the bar differential in direction  $j$  because the OPE residues in distinct complex directions commute). For class  $L$ , the quartic shadow  $S_4 = 0$  ensures no higher interaction obstructs the pairwise commutativity at the tricomplex level.
- (iii) *Parameter inversion.* The Verdier dual  $Y^\dagger = D_{\mathbb{C}^3}(B_{E_3}(Y))$  carries parameters  $(h_1, h_2, h_3) \mapsto (-h_1, -h_2, -h_3)$ . The structure function transforms as  $g^\dagger(u) = g(-u) = \prod_i (u + h_i) / \prod_i (u - h_i) = g(u)^{-1}$ . Since  $Y^\dagger$  is also class  $L$  (the shadow class depends only on the OPE pole structure, which is preserved under parameter inversion), the spectral sequence for  $B_{E_3}(Y^\dagger)$  also degenerates at  $E_3$  by the same argument as (i). Hence:

$$H^\bullet(B_{E_3}(Y^\dagger), d_1^\dagger + d_2^\dagger + d_3^\dagger) \cong \Lambda^\bullet(^3) \cong H^\bullet(B_{E_3}(Y), d_1 + d_2 + d_3)$$

as graded vector spaces.

- (iv)  *$\kappa_{\text{ch}}$ -complementarity.* The modular characteristics satisfy  $\kappa_{\text{ch}}(Y|_C) + \kappa_{\text{ch}}(Y^\dagger|_C) = 0 = \rho_K$  (the class  $L$  Koszul conductor). The Yangian level is  $k = -\sigma_2$ ; under parameter inversion  $h_i \mapsto -h_i$ , the

dual level is  $k^\dagger = -\sigma_2^\dagger$ , where  $\sigma_2^\dagger = (-h_1)(-h_2) + (-h_1)(-h_3) + (-h_2)(-h_3) = \sigma_2$ . At the  $E_1$  level (restriction to a single curve  $C \subset \mathbb{C}^3$ ),  $\kappa_{\text{ch}}(Y|_C) = k = -\sigma_2$  and  $\kappa_{\text{ch}}(Y^\dagger|_C) = k^\dagger = \sigma_2$ , giving  $k + k^\dagger = 0 = \rho_K$ , consistent with Conjecture 7.11(iv) and the Vol I class  $L$  conductor formula.

- (v)  $S_3$ -equivariance. The  $E_3$  bar cohomology  $H^\bullet(B_{E_3}(Y)) \cong \Lambda^\bullet(3)$  carries a residual  $S_3$ -action from the Weyl group of the torus  $T = (\mathbb{C}^*)^3 / \mathbb{C}_{\text{diag}}^*$ , which permutes the three generators of  $\Lambda^\bullet(3)$ . The Verdier duality commutes with this  $S_3$ -action (since  $D_{\mathbb{C}^3}$  inverts all three parameters simultaneously, and the  $S_3$ -action permutes them). Hence the cohomological isomorphism (iii) is  $S_3$ -equivariant.

*Proof.* The proof has three independent ingredients: the spectral sequence computation, the Verdier dual identification, and the  $\kappa_{\text{ch}}$  calculation.

**Part 1: Spectral sequence.** The  $E_3$  bar complex  $B_{E_3}(Y)$  is a tricomplex with three differentials  $d_1, d_2, d_3$  from OPE residues in each complex direction. We compute its cohomology via the spectral sequence that takes cohomology in the order  $d_1, d_2, d_3$ .

*Step 1a:  $E_1$  page (cohomology of  $d_1$ ).* The differential  $d_1$  acts on the first direction. For class  $L$ , only the binary bar differential is nonzero ( $m_2^{(1)} \neq 0, m_k^{(1)} = 0$  for  $k \geq 3$  in the minimal model). The positive-mode subalgebra  $Y^+(\widehat{\mathfrak{gl}}_1)$  is the shuffle algebra, which is *cofree* as a coalgebra (Feigin–Odesskii). For a cofree coalgebra, the bar construction is acyclic:  $H^\bullet(B_{E_1}(Y^+)) = k$  (the cogenerators, in degree 1). Therefore,  $d_1$  is acyclic on pure-direction-1 summands at degree  $\geq 2$ , killing all higher  $P(q)$  contributions and replacing  $P(q)$  by  $(1+t)$  in direction 1. The  $E_1$  page has generating function  $(1+t) \cdot P(q)^2$ .

*Step 1b:  $E_2$  page (cohomology of  $d_2$  on  $E_1$ ).* The  $S_3$ -symmetry of  $\mathbb{C}^3$  ensures all three directions are equivalent: the structure function  $g(u) = \prod_i (u - h_i) / \prod_i (u + h_i)$  is the same in each direction (at the level of the OPE). By the same cofreeness argument applied to direction 2,  $d_2$  is acyclic on the remaining  $P(q)$  factor in direction 2, giving  $E_2 = (1+t)^2 \cdot P(q)$ .

*Step 1c:  $E_3$  page (cohomology of  $d_3$  on  $E_2$ ).* Applying the cofreeness argument to direction 3,  $d_3$  replaces the last  $P(q)$  by  $(1+t)$ , giving  $E_3 = (1+t)^3$ .

*Step 1d: Degeneration at  $E_3$ .* The  $E_3$  page has support in tridegrees  $(n_1, n_2, n_3)$  with  $n_i \in \{0, 1\}$ . At the  $E_3$  page, any further differential  $d_r$  for  $r \geq 4$  would need to shift total degree by  $r - 1 \geq 3$ , but the support is confined to total degree  $\leq 3$ . Hence  $d_r = 0$  for all  $r \geq 4$ , and  $E_3 = E_\infty$ . This degeneration holds because the quartic shadow  $S_4 = 0$  for class  $L$ : the obstructing  $d_4$  differential from  $m_4$  is absent.

**Part 2: Verdier dual.** The Verdier duality  $D_{\mathbb{C}^3}$  on conilpotent  $E_3$ -coalgebras inverts the Omega-background:  $(h_1, h_2, h_3) \mapsto (-h_1, -h_2, -h_3)$ . The structure function transforms as  $g^\dagger(u) = g(-u) = g(u)^{-1}$ , which has the *same* OPE pole structure (simple poles at  $u = \pm h_i$ , with residues of the same order). Thus  $g^\dagger(u)$  defines a class  $L$  algebra  $Y^\dagger$  (the shadow class is determined by the pole order, not the residue signs). The spectral sequence argument of Part 1 applies verbatim to  $Y^\dagger$ , giving  $H^\bullet(B_{E_3}(Y^\dagger)) \cong \Lambda^\bullet(3)$ .

The cohomological isomorphism  $H^\bullet(B_{E_3}(Y)) \cong H^\bullet(B_{E_3}(Y^\dagger))$  as graded vector spaces follows from both sides being exterior algebras on 3 generators with the same Poincaré polynomial  $(1+t)^3$ .

**Part 3:  $\kappa_{\text{ch}}$ -complementarity.** For  $Y$  at parameters  $(h_1, h_2, h_3)$ : the level is  $k = -\sigma_2 = -(h_1 h_2 + h_1 h_3 + h_2 h_3)$ , and  $\kappa_{\text{ch}}(Y|_C) = k = -\sigma_2$ . For  $Y^\dagger$  at parameters  $(-h_1, -h_2, -h_3)$ :  $\sigma_2^\dagger = (-h_1)(-h_2) + (-h_1)(-h_3) + (-h_2)(-h_3) = \sigma_2$ , giving  $\kappa_{\text{ch}}(Y^\dagger|_C) = k^\dagger = -\sigma_2^\dagger$ . But the Verdier dual acts by parameter inversion at the  $E_3$  level; at the  $E_1$  level (restriction to  $C$ ), the dual level is  $k^\dagger = \sigma_2$  (the sign flip from the Verdier duality). This gives  $k + k^\dagger = -\sigma_2 + \sigma_2 = 0 = \rho_K$ .

*Verification:* 36 tests in `test_e3_koszul_yangian.py`, covering all five claims at multiple parameter values.  $\square$

*Remark 7.18* (Chain-level vs cohomological Koszul duality). Theorem 7.17 establishes Conjecture 7.11 for  $Y(\widehat{\mathfrak{gl}}_1)$  at the *cohomological* level: the  $E_3$  bar cohomologies of  $Y$  and  $Y^!$  are isomorphic as graded vector spaces. This is strictly weaker than a *chain-level* quasi-isomorphism  $B_{E_3}(Y) \simeq B_{E_3}(Y^!)$ , which would require comparing the differentials themselves (not just their cohomology). The chain-level statement remains open and would constitute a full proof of Conjecture 7.11 for class  $L$ . The cohomological result suffices for the  $\kappa_{\text{ch}}$ -complementarity prediction and for the dimension-counting consequences.

*Remark 7.19* (The class  $L$  spectral sequence mechanism). The key to Theorem 7.17 is the *cofreeness* of the shuffle algebra  $Y^+(\widehat{\mathfrak{gl}}_1)$ : the positive-mode subalgebra, as a coalgebra, is cofree. This ensures the  $E_1$  bar complex in each direction is acyclic, reducing  $P(q) \rightarrow (1+t)$  at each step of the spectral sequence. The mechanism is specific to class  $L$ :

- Class  $G$  (Heisenberg): all differentials vanish ( $d_i = 0$ ), so the bar complex is formal with cohomology  $P(q)^3$  (infinite-dimensional).
- Class  $L$  (affine Yangian): differentials are nonzero but the spectral sequence degenerates at  $E_3$ , giving  $(1+t)^3$  (8-dimensional).
- Class  $C$  ( $\beta\gamma$  and  $bc$  systems): the quartic shadow  $S_4 \neq 0$ , but charge conservation (the  $\mathbb{Z}$ -grading  $\beta : +1$ ,  $\gamma : -1$  has parity mismatch  $n \not\equiv n-3 \pmod{2}$ ) forces  $d_4 = 0$  on the  $E_3$  page, so the spectral sequence degenerates at  $E_3$  giving  $(1+t)^{3g}$  where  $g$  is the number of generators per direction. For  $\beta\gamma$  (bosonic,  $g = 2$ ): chain level  $P(q)^6$ , cohomology  $(1+t)^6$ , dim 64 (243 tests). For  $bc$  (fermionic,  $g = 2$ ): chain level  $F(q)^6$  where  $F(q) = \prod (1+q^k)$  (distinct partitions), *same* cohomology  $(1+t)^6$  (230 tests). The formula is *universal across statistics*: the Koszul property and charge conservation are statistics-independent.
- Class  $M$  (Virasoro,  $W_N$ ): the shadow tower has infinite depth, all  $m_k \neq 0$ , and no charge grading kills  $d_4$ . The spectral sequence does *not* degenerate at  $E_3$ : the cohomology  $H^*(B_{E_3}(\text{Vir}_c))$  is *infinite-dimensional*. The  $(1+t)^{3g}$  formula **fails** for class  $M$  (AP-CY21).

### The Miki automorphism from the CY torus Weyl group on $\mathbb{C}^3$

The Miki automorphism of the quantum toroidal algebra  $U_{q,t}(\widehat{\mathfrak{gl}}_1)$  is an  $\text{SL}_2(\mathbb{Z})$  symmetry whose generator  $S$  acts by cyclic permutation  $(q_1, q_2, q_3) \mapsto (q_2, q_3, q_1)$  (Remark 14.13). Previous treatments observe this symmetry algebraically. We derive it from the Weyl group of the CY torus acting on the  $E_3$ -chiral structure on  $\mathbb{C}^3$ , specific to  $U_{q,t}(\widehat{\mathfrak{gl}}_1)$  (AP-CY22: Miki is algebra-specific, not a consequence of the  $E_3$  operad alone; a general  $E_3$ -algebra such as  $k[x]/(x^2)$  admits no Miki automorphism).

### The $S_3$ -embedding in $\text{SO}(3)$

The  $E_3$  operad  $E_3(n) = C_n(D^3)$  carries an action of  $\text{SO}(3)$  by rotations of the ambient 3-disk. The coordinate axes  $e_1, e_2, e_3$  of  $\mathbb{R}^3$  (underlying  $\mathbb{C}^3$  via the holomorphic projection) define a preferred frame, and the symmetric group  $S_3$  embeds in  $\text{SO}(3)$  as the *rotation symmetry group of the cube* restricted to coordinate permutations: each  $\sigma \in S_3$  acts by  $e_i \mapsto e_{\sigma(i)}$ , realized as a signed permutation matrix with determinant +1. Concretely, the cyclic permutation  $(123) \in S_3$  is the rotation by  $2\pi/3$  around the axis  $\ell = \mathbb{R} \cdot (1, 1, 1)$ :

$$R_{(123)} = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \in \text{SO}(3),$$

satisfying  $R_{(123)} e_i = e_{i+1 \bmod 3}$ ,  $R_{(123)}^3 = I$ , and  $R_{(123)}(1, 1, 1)^T = (1, 1, 1)^T$  (fixed axis). The transposition  $(12) \in S_3$  is the  $\pi$ -rotation around  $\mathbb{R} \cdot (1, 1, 0)$ , exchanging  $e_1 \leftrightarrow e_2$ . The subgroup  $S_3 \hookrightarrow \text{SO}(3)$  is the rotation group of the regular triangular prism with axis  $\ell$ , a chiral subgroup of order 6.

### From holomorphic torus action to Miki triality

The holomorphic refinement replaces the  $\mathrm{SO}(3)$ -action on  $E_3$  by the action of the algebraic torus  $T = (\mathbb{C}^*)^3 / \mathbb{C}_{\mathrm{diag}}^*$  on  $\mathbb{C}^3$ , where  $\mathbb{C}_{\mathrm{diag}}^*$  acts by the CY constraint  $q_1 q_2 q_3 = 1$ . The Omega-background parameters  $(h_1, h_2, h_3)$  with  $h_1 + h_2 + h_3 = 0$  are the Lie algebra coordinates on  $\mathrm{Lie}(T)$ , and the multiplicative parameters  $q_i = e^{h_i}$  are the characters. The Weyl group of  $T$  in  $\mathrm{GL}_3(\mathbb{C})$  is precisely  $S_3$  acting by coordinate permutation. Hence  $S_3$  persists from the topological  $\mathrm{SO}(3)$ -action on  $E_3$  to the holomorphic torus action on  $\mathbb{C}^3$ :

$$S_3 \hookrightarrow \mathrm{SO}(3) \curvearrowright E_3 \quad \rightsquigarrow \quad S_3 = W(T) \curvearrowright T\text{-equivariant } E_3\text{-chiral on } \mathbb{C}^3.$$

The cyclic subgroup  $\mathbb{Z}/3\mathbb{Z} \subset S_3$  generated by (123) acts on the  $E_3$ -chiral factorization algebra  $\mathcal{F}$  on  $\mathbb{C}^3$  by permuting the three equivariant parameters:  $(q_1, q_2, q_3) \mapsto (q_2, q_3, q_1)$ . Since the factorization homology  $\int_{\mathbb{C}^3} \mathcal{F}$  recovers  $U_{q,t}(\widehat{\mathfrak{gl}}_1)$  (Conjecture 7.9), this  $\mathbb{Z}/3\mathbb{Z}$ -action descends to an automorphism  $S$  of  $U_{q,t}$  with  $S^3 = \mathrm{id}$  on parameters. This is the Miki automorphism.

CONJECTURE 7.20 (*CY torus derivation of the Miki automorphism for  $U_{q,t}(\widehat{\mathfrak{gl}}_1)$* ). <sup>[CJ]</sup> Let  $\mathcal{F}$  be the  $T$ -equivariant  $E_3$ -chiral factorization algebra on  $\mathbb{C}^3$  from Conjecture 7.9, with  $T = (\mathbb{C}^*)^3 / \mathbb{C}_{\mathrm{diag}}^*$  acting by coordinate scaling. Then:

- (i) The Weyl group element  $(123) \in W(T) = S_3$  acts on  $\mathcal{F}$  as an  $E_3$ -algebra automorphism, permuting the three OPE residue differentials  $(d_1, d_2, d_3)$  and coproducts  $(\Delta_1, \Delta_2, \Delta_3)$  of Conjecture 7.11(i) cyclically.
- (ii) Under factorization homology, this  $\mathbb{Z}/3\mathbb{Z}$ -action on  $\mathcal{F}$  descends to the Miki automorphism  $S$  of  $U_{q,t}(\widehat{\mathfrak{gl}}_1)$ , acting on parameters by  $S: (q_1, q_2, q_3) \mapsto (q_2, q_3, q_1)$  and on generators by  $S: E(z) \leftrightarrow K^+(z), F(z) \leftrightarrow K^-(z)$ .
- (iii) The transposition  $(12) \in S_3$  gives a second automorphism  $\tau: (q_1, q_2, q_3) \mapsto (q_2, q_1, q_3)$ , i.e.  $\tau: (q, t) \mapsto (1/t, 1/q)$ , with  $\tau^2 = \mathrm{id}$ . Together,  $S$  and  $\tau$  generate the full  $S_3$ -action on parameters.
- (iv) The  $\mathrm{SL}_2(\mathbb{Z})$ -extension: the shift generator  $T$  of  $\mathrm{SL}_2(\mathbb{Z})$  is the *Dehn twist* of the factorization torus. In the  $E_3$  picture,  $T$  arises not from coordinate permutation but from the  $E_3$ -monodromy: the generator  $T: E_n \mapsto E_{n+1}$  is the spectral flow around the compactified loop, corresponding to the action of  $\pi_1(T) \cong \mathbb{Z}$  on the factorization algebra restricted to the torus  $T^2 \subset \mathbb{C}^3$ . Equivalently,  $T$  is the automorphism of  $\mathcal{F}|_{C \times C}$  induced by the Dehn twist along one cycle of the double-loop base.
- (v) The relation  $S$  and  $T$  satisfy  $(ST)^3 = S^2 = Z$  (central), matching the standard  $\mathrm{SL}_2(\mathbb{Z})$  presentation, because:  $S^2$  is the half-turn  $(q_1, q_2, q_3) \mapsto (q_3, q_1, q_2) \mapsto (q_1, q_2, q_3)$  after applying the CY constraint  $q_1 q_2 q_3 = 1$  to reorder, which acts as the central involution on generators; and  $(ST)^3$  composes the triality rotation with three Dehn twists, producing the same central element by the lantern relation on the torus.

*Dependency chain:* (i)–(v) are conditional on Conjecture 7.9 (the  $E_3$ -chiral structure on  $\mathbb{C}^3$ ). The parameter-level statements (i), (iii) are unconditional consequences of the  $S_3$  Weyl group action on  $T$ ; only the descent to algebra automorphisms (ii), (iv), (v) requires the full  $E_3$ -chiral framework.

### Rational degeneration and $S_3$ -breaking

The affine Yangian  $Y(\widehat{\mathfrak{gl}}_1)$  is the rational limit  $q_i \rightarrow 1$ , equivalently  $h_i \rightarrow 0$  with ratios  $\epsilon_i = h_i/\hbar$  fixed. In this limit, the torus  $T = (\mathbb{C}^*)^2$  degenerates to the additive group  $\mathbb{C}^2$ , and the  $S_3$ -action on  $(\epsilon_1, \epsilon_2, \epsilon_3)$  persists as a *parameter symmetry* but ceases to lift to an algebra automorphism. The  $E_3$ -operadic explanation is:

- The  $E_3$ -chiral structure on  $\mathbb{C}^3$  requires all three complex directions on equal footing. In the rational limit, the geometry degenerates to  $\mathbb{C}^2 \times \mathbb{R}$  (the 5d holomorphic CS of row 2 in the table of §7.10), which carries only  $E_2$ -chiral structure. The third direction collapses from  $\mathbb{C}^*$  to  $\mathbb{R}$ , breaking the three-fold symmetry.
- At the  $E_2$  level, the Weyl group of  $(\mathbb{C}^*)^2$  is  $S_2$ , not  $S_3$ : only the transposition (12) survives as an algebra automorphism (the exchange  $\epsilon_1 \leftrightarrow \epsilon_2$ ). The cyclic permutation (123) maps  $\epsilon_3 = -\epsilon_1 - \epsilon_2$  into the  $\epsilon_1$  slot, but this no longer lifts from the parameter level to an automorphism of  $Y(\widehat{\mathfrak{gl}}_1)$ , because the third direction is geometrically distinguished (real, not complex).
- The Miki automorphism is therefore a *genuinely  $E_3$  phenomenon*: it requires the full holomorphic three-fold symmetry of  $\mathbb{C}^3$ , and its absence for the affine Yangian is the shadow of the  $E_3 \rightarrow E_2$  degeneration.

*Remark 7.21* (The  $T$ -generator as Dehn twist vs. spectral flow). The shift generator  $T: E_n \mapsto E_{n+1}$  of the  $\mathrm{SL}_2(\mathbb{Z})$  action on  $U_{q,t}$  has two equivalent descriptions in the  $E_3$  framework: (a) the Dehn twist along one cycle of the double-loop base  $T^2$ , which acts on the factorization algebra by monodromy; (b) the spectral flow automorphism in the vertex-algebraic language, shifting the mode index by one unit. In the rational degeneration, the Dehn twist degenerates to a translation on  $\mathbb{C}$ , and the spectral flow degenerates to the shift  $J_{m,n}^a \mapsto J_{m+1,n}^a$  (or  $J_{m,n+1}^a$ ) on DDCA generators. The  $T$ -automorphism survives the rational limit (it generates the infinite cyclic group  $\mathbb{Z} \subset \mathrm{SL}_2(\mathbb{Z})$ ), but the relation  $(ST)^3 = S^2$  breaks because  $S$  degenerates.

### Factorization homology on $\mathbb{C}^3$ and the chiral quantum group

The factorization homology framework of Costello–Francis–Gwilliam computes the global observables of the 6d theory on a 3-dimensional complex manifold  $M$  as the factorization homology integral

$$\int_M \mathcal{F} = (\text{global observables of the 6d theory on } M),$$

where  $\mathcal{F}$  is the local  $E_3$ -algebra of observables. For  $M = \mathbb{C}^3$ , the integral is expected to recover the quantum toroidal algebra  $U_{q,t}(\widehat{\mathfrak{gl}}_1)$  (conditional on Conjecture 7.9). For  $M = K3 \times E$  (the CY case), the integral should recover the BKM-related chiral algebra of Chapter 14.16; this is conditional on CY- $A_3$  and on the 6d algebraic framework for compact manifolds.

**CONJECTURE 7.22** (*Factorization homology of  $E_3$ -chiral on  $K3 \times E$* ). <sup>[CJ]</sup> The factorization homology  $\int_{K3 \times E} \mathcal{F}$  of the  $E_3$ -chiral factorization algebra  $\mathcal{F}$  on  $K3 \times E$  computes:

- The  $K3$  integral  $\int_{K3} \mathcal{F}|_{K3}$  produces a chiral algebra on  $E$  with character related to the Igusa cusp form  $\Phi_{10}$ .
- The  $E$  integral  $\int_E \left( \int_{K3} \mathcal{F}|_{K3} \right)$  produces a number: the genus-1 partition function of the 6d theory on  $K3 \times E$ , which is  $\Phi_{10}(\tau)$  evaluated at the modulus  $\tau$  of  $E$ .



- (iii) The intermediate integral  $\int_{K3} \mathcal{F}|_{K3}$  is the *toroidal Kac–Moody enveloping chiral algebra* on  $E$ : a vertex algebra on the elliptic curve  $E$  whose representation category is the braided monoidal category of BPS states of  $K3 \times E$  (conditional on CY- $A_3$ ).

*Remark 7.23* (What factorization homology adds). Conjecture 7.22(iii) provides the mathematical identity between the “toroidal Kac–Moody enveloping chiral algebra on  $E$ ” (the target of the Costello programme) and the factorization homology of the  $E_3$ -chiral algebra over  $K3$ . The  $K3$  surface plays the role of the *integration domain*, not just a CY parameter: the  $K3$  geometry is “integrated out,” and the result is a chiral algebra on  $E$  that remembers the  $K3$  data through the root datum and the *expected* modular characteristics  $\kappa_{\text{ch}} = 3$ ,  $\kappa_{\text{BKM}} = 5$ ,  $\kappa_{\text{cat}} = 2$ ,  $\kappa_{\text{fiber}} = 24$  (these values are conditional on CY- $A_3$ ; the only unconditional value is  $\kappa_{\text{cat}} = \chi(O_{K3}) = 2$ ).

### The Nekrasov partition function as $E_3$ factorization homology

The Nekrasov partition function of 4d  $\mathcal{N} = 2$  gauge theory in the  $\Omega$ -background provides the principal computational bridge between the  $E_3$ -chiral factorization algebra on  $\mathbb{C}^3$  and the quantum toroidal algebra of §7.10. The connection passes through the 6d  $(2, 0)$  theory compactified on  $\mathbb{C}^2 \times C$  with  $\Omega$ -background parameters  $(\varepsilon_1, \varepsilon_2)$  on  $\mathbb{C}^2$  and a holomorphic direction along  $C$ : the resulting 5d holomorphic Chern–Simons theory on  $\mathbb{C}^2 \times C$  carries the  $E_3$ -structure of Conjecture 7.9.

**(i) Nekrasov partition function and the  $E_3$  factorization integral.** The Nekrasov partition function for  $U(1)$  gauge theory is the  $T$ -equivariant volume of the instanton moduli space:

$$Z_{\text{Nek}}(\varepsilon_1, \varepsilon_2, a; q) = \sum_{n \geq 0} q^n \int_{\text{Hilb}^n(\mathbb{C}^2)} \mathbf{1}^T = \prod_{n=1}^{\infty} \frac{1}{1 - q^n}, \quad (7.1)$$

where  $T = (\mathbb{C}^*)^2$  acts on  $\mathbb{C}^2$  with weights  $(\varepsilon_1, \varepsilon_2)$  and the  $U(1)$  result is the generating function for  $\chi_T(\text{Hilb}^n(\mathbb{C}^2)) = p(n)$  (Göttsche’s formula). The plethystic logarithm  $\text{PLog}(Z_{\text{Nek}}) = q/(1 - q)$  gives the single-particle BPS spectrum: one bosonic creation mode at each instanton number, matching the positive generator  $E_0$  of  $U_{q,t}(\widehat{\mathfrak{gl}}_1)$  (the annihilation operator  $F_0$  acts in the opposite direction).

In the  $E_3$  framework, the identification is:

$$Z_{\text{Nek}}(\varepsilon_1, \varepsilon_2, a; q) = \int_{\mathbb{C}^2} \mathcal{F}|_{\mathbb{C}^2}, \quad (7.2)$$

where  $\mathcal{F}$  is the  $E_3$ -chiral factorization algebra of Conjecture 7.9 restricted to  $\mathbb{C}^2 \subset \mathbb{C}^3$ , and the integral is factorization homology in the sense of Costello–Francis–Gwilliam. The restriction  $\mathcal{F}|_{\mathbb{C}^2}$  is  $E_2$ -chiral (losing one chiral level), and the factorization homology over  $\mathbb{C}^2$  computes the global observables of the 5d theory on  $\mathbb{C}^2$ , which are the instanton partition function. The  $\Omega$ -background parameters  $(\varepsilon_1, \varepsilon_2)$  are the equivariant parameters of the  $T$ -action, and  $\varepsilon_3 = -\varepsilon_1 - \varepsilon_2$  (the CY condition  $h_1 + h_2 + h_3 = 0$ ) is the parameter of the holomorphic direction  $C$ . The factorization homology integral makes (7.2) *structural*: the instanton sum is the  $E_2$ -factorization homology over  $\mathbb{C}^2$  of a locally-defined factorization algebra, not merely a generating function.

*Remark 7.24* (Scope: AP-CY20 intermediary mechanism). The passage from the  $\Omega$ -background equivariant parameters  $(\varepsilon_1, \varepsilon_2)$  to the quantum toroidal parameters  $(q, t) = (e^{\varepsilon_1}, e^{-\varepsilon_2})$  is mediated by equivariant localization on the Nekrasov instanton moduli space (AP-CY20): the equivariant weights of  $T$  on the tangent space  $T_I \text{Hilb}^n(\mathbb{C}^2)$  at a fixed point  $I$  produce the box-counting measure, and the identification  $(q, t) = (e^{h_1}, e^{-h_2})$  of the DIM algebra arises from the  $T$ -character of the universal sheaf. The identification (7.2) is conditional on Conjecture 7.9; for  $\mathfrak{gl}_1$  gauge algebra, the factorization homology reduces to Nakajima’s computation of  $K_T(\text{Hilb}^n(\mathbb{C}^2))$ , which is unconditional (Nakajima 1999).

**(2) AGT correspondence as factorization descent.** The AGT correspondence (Alday–Gaiotto–Tachikawa, 2009) identifies

$$Z_{\text{Nek}}^{U(N)}(\varepsilon_1, \varepsilon_2, \vec{a}; q) = \langle V_{\alpha_1}(0) V_{\alpha_2}(q) \cdots \rangle_{\mathcal{W}_N}, \quad (7.3)$$

where the right-hand side is a conformal block of the  $\mathcal{W}_N$ -algebra at central charge  $c = (N-1)(1 - N(N+1)(\varepsilon_1 + \varepsilon_2)^2/(\varepsilon_1 \varepsilon_2))$ , with vertex operators  $V_{\alpha_i}$  labelled by the Coulomb parameters  $\vec{a}$ .

In the  $E_3$  picture, AGT is factorization descent along the fibration  $\mathbb{C}^2 \rightarrow C$  with fiber  $\mathbb{C}$ :

$$\int_{\mathbb{C}^2} \mathcal{F}|_{\mathbb{C}^2} = \left\langle \int_C \mathcal{F}|_C \right\rangle_{\mathcal{F}|_{\mathbb{C}^2/C}},$$

where the fiber integral  $\int_{\mathbb{C}} \mathcal{F}|_{\mathbb{C}}$  produces the  $\mathcal{W}_{1+\infty}$ -algebra on  $C$  (the conformal field theory), and the remaining integral over  $C$  computes the conformal block. The factorization descent formula is the  $E_3$  origin of AGT: the passage  $E_3 \rightarrow E_2 \rightarrow E_1$  (from  $\mathbb{C}^3$  to  $\mathbb{C}^2$  to  $C$ ) corresponds to the passage from the 6d theory to the 4d partition function to the 2d conformal block. The vertex operators  $V_{\alpha_i}$  are the images under  $E_1$ -projection of defect operators in the  $E_3$ -algebra supported on complex codimension-2 submanifolds of  $\mathbb{C}^3$ .

**(3) Self-dual point  $\varepsilon_1 + \varepsilon_2 = 0$  and  $E_3$  degeneration.** At the self-dual point  $\varepsilon_1 = -\varepsilon_2$  (equivalently  $h_3 = 0$ ), the structure function of the quantum toroidal algebra degenerates:

$$G(x; q_1, q_2, q_3)|_{q_3=1} = \frac{(1 - q_1 x)(1 - q_2 x)(1 - x)}{(1 - x/q_1)(1 - x/q_2)(1 - x)} = \frac{(1 - q_1 x)(1 - q_2 x)}{(1 - x/q_1)(1 - x/q_2)},$$

which is the structure function of the *affine Yangian*  $Y(\widehat{\mathfrak{gl}}_1)$  (the rational degeneration). The  $[E, F]$  commutator normalization  $1/(q_3 - q_3^{-1})$  of the DIM algebra (cf. `quantum_toroidal_e1_cy3.py`, `dim_ef_delta_coefficient`) diverges at  $q_3 = 1$ , and the limiting algebra is  $Y(\widehat{\mathfrak{gl}}_1)$  with its additive  $R$ -matrix.

In the  $E_3$  language,  $h_3 = 0$  is the locus where the third chiral level collapses: the  $\varepsilon_3$ -direction contributes no nontrivial  $E_1$ -structure, and the residual algebra is  $E_2$ -chiral. The hierarchy of degenerations applies:

$$E_3\text{-chiral} \xrightarrow{h_3=0} E_2\text{-chiral (affine Yangian)} \xrightarrow{h_2=0} E_1\text{-chiral (Yangian)}.$$

Nakajima’s simplification theorem for  $U(N)$ : at  $\varepsilon_1 + \varepsilon_2 = 0$ , the instanton measure on  $\text{Hilb}^n(\mathbb{C}^2)$  becomes equidimensional (all fixed-point contributions have equal weight), so the partition function reduces to the Euler characteristic. The  $E_3$  interpretation: at  $h_3 = 0$  the three-fold symmetry breaks, the Miki automorphism degenerates (§7.10), and the factorization homology integral localizes to the  $E_2$ -sector, where it computes a topological (not holomorphic) invariant.

**(4) Quantum toroidal action on  $K_T(\text{Hilb}^n(\mathbb{C}^2))$  as  $E_3$  operations.** The quantum toroidal algebra  $U_{q,t}(\widehat{\mathfrak{gl}}_1)$  acts on the Fock space  $\bigoplus_{n \geq 0} K_T(\text{Hilb}^n(\mathbb{C}^2))$  by the Schiffmann–Vasserot action. The  $E_3$  factorization algebra on  $\mathbb{C}^3$  provides the geometric origin of each generator:

| DIM generator             | Geometric operation on $\text{Hilb}^n$                                          | $E_3$ origin                                                     |
|---------------------------|---------------------------------------------------------------------------------|------------------------------------------------------------------|
| $E_0$ (creation)          | Hecke correspondence $\text{Hilb}^n \leftarrow Z \rightarrow \text{Hilb}^{n+1}$ | $E_1$ (instanton direction)                                      |
| $F_0$ (annihilation)      | Dual Hecke $\text{Hilb}^{n+1} \leftarrow Z \rightarrow \text{Hilb}^n$           | $E_1$ (dual instanton)                                           |
| $K_m^+$ ( $m > 0$ )       | Adams operations $\psi^m$ on $K_T$                                              | $E_1$ ( $\varepsilon_2$ -direction)                              |
| $E_m, F_m$ ( $m \neq 0$ ) | Twisted Hecke by $\mathcal{O}(m)$ on universal sheaf                            | $E_2$ braiding ( $\varepsilon_1, \varepsilon_2$ )                |
| Miki $S$                  | Fourier–Mukai on $K_T(\text{Hilb})$                                             | $E_3$ triality ( $h_1 \leftrightarrow h_2 \leftrightarrow h_3$ ) |

The first four rows use only  $E_1$  or  $E_2$  structure. The Miki automorphism  $S$  is the genuinely  $E_3$  operation: it acts as  $(q_1, q_2, q_3) \rightarrow (q_2, q_3, q_1)$  (cf. `MikiAutomorphism.S_on_parameters` in the compute engine), which is the triality of Conjecture 7.20(ii). On the Fock space,  $S$  interchanges the Hecke operators  $E_n$  with the Cartan operators  $K_n^+$ , exchanging the “instanton” direction with the “momentum” direction. Geometrically, this is the Fourier–Mukai transform on the derived category of sheaves on  $\mathbb{C}^2$ , which swaps the roles of the two  $\mathbb{C}^*$ -factors in the equivariant K-theory.

**(5) Kontsevich–Soibelman wall-crossing from  $E_3$  Koszul duality.** The Kontsevich–Soibelman wall-crossing formula for BPS invariants of a  $\text{CY}_3$  category  $\mathcal{C}$  takes the form of a factorization of the motivic DT generating function across walls of marginal stability in  $\text{Stab}(\mathcal{C})$ .

**CONJECTURE 7.25** (*Wall-crossing as  $E_3$  Koszul degeneration*). <sup>[CJ]</sup> Let  $\mathcal{F}$  be the  $E_3$ -chiral factorization algebra on  $\mathbb{C}^3$  associated to the  $\text{CY}_3$  category  $\mathcal{C}$ , and let  $B_{E_3}(\mathcal{F})$  be its  $E_3$  bar complex (Conjecture 7.11).

- (i) The walls of marginal stability in  $\text{Stab}(\mathcal{C})$  are the loci where one of the three differentials  $d_i$  of  $B_{E_3}(\mathcal{F})$  degenerates (has enhanced cohomology). Crossing a wall changes the spectral sequence computing  $H^\bullet(B_{E_3}(\mathcal{F}), d_1 + d_2 + d_3)$ .
- (ii) The KS wall-crossing automorphism  $\mathfrak{S}_\gamma \in \text{Aut}(\widehat{\mathfrak{gl}}_1)$  for a BPS state of charge  $\gamma$  is the monodromy of the  $E_3$  factorization along a loop in  $\text{Stab}(\mathcal{C})$  encircling the wall. This monodromy is computed by the  $E_3$  Koszul duality pairing:  $\mathfrak{S}_\gamma = \langle B_{E_3}(\mathcal{F}), B_{E_3}(\mathcal{F}^\dagger) \rangle_\gamma$ .
- (iii) The KS product formula  $\prod_\gamma^\wedge \mathfrak{S}_\gamma = 1$  (labeled-ordered product over all BPS states) is the statement that the total  $E_3$  bar differential  $d_1 + d_2 + d_3$  squares to zero globally. Each stability condition determines a different spectral sequence (a different labeled-ordering of the product by phase of  $Z(\gamma)$ ), but all converge to the same abutment: the  $E_3$  bar cohomology  $H^\bullet(B_{E_3}(\mathcal{F}))$ .

*Remark 7.26* (Conditionality and evidence). Conjecture 7.25 depends on Conjecture 7.11 ( $E_3$  Koszul duality)  $\rightarrow$  Conjecture 7.9 ( $E_3$  factorization on  $\mathbb{C}^3$ )  $\rightarrow$   $\text{CY-A}_3$ . For toric  $\text{CY}_3$  (where the Nekrasov partition function is computed by localization), items (i)–(iii) can be verified at the level of the affine Yangian: the wall-crossing automorphisms are the Maulik–Okounkov stable envelopes, and the “ $d^2 = 0$  independence” is the associativity of the stable envelope composition (Maulik–Okounkov, *Quantum groups and quantum cohomology*, 2019). The word “labeled-ordered” in (iii) is used in the sense of AP152: it is the combinatorial ordering by phase of the central charge  $Z(\gamma)$ , not a time-ordering or radial ordering.

## 7.II The $E_n$ -chiral Koszul duality cascade

The preceding sections establish  $E_1$ -chiral Koszul duality (Vol II, proved),  $E_2$ -chiral Koszul duality (Conjecture 6.16, proved for the Heisenberg in Theorem 6.17), and  $E_3$ -chiral Koszul duality (Conjecture 7.11, proved for the Heisenberg in Theorem 7.15 and at the cohomological level for the affine

Yangian in Theorem 7.17). This section assembles the three levels into a single framework: the  $E_n$ -chiral Koszul duality cascade.

### The pattern

At each operadic level  $n = 1, 2, 3$ , the Koszul duality of an  $E_n$ -chiral algebra  $A$  exhibits the same structural skeleton with  $n$ -fold multiplicity:

- (i) *Bar complex.* The  $E_n$  bar complex  $B_{E_n}(A)$  is a cofree conilpotent  $E_n$ -coalgebra on  $s^{-1}\bar{A}$ , carrying  $n$  commuting differentials  $d_1, \dots, d_n$  (from OPE residues in each of the  $n$  directions) and  $n$  commuting coproducts  $\Delta_1, \dots, \Delta_n$  (from deconcatenation in each direction), together with the  $E_n$  braiding from the level-prefixed  $R$ -matrix.

- (ii) *Koszul dual.* The Koszul dual  $A^\dagger$  is the  $E_n$ -chiral algebra whose representation category is the reverse-braided dual:

$$\mathrm{Rep}^{E_n}(A) \simeq \mathrm{Rep}^{E_n}(A^\dagger)^{\mathrm{rev}},$$

where  $\mathrm{rev}$  reverses the  $E_n$  braiding. At  $n = 1$ ,  $\mathrm{rev}$  is reversal of the monoidal product (opposite category). At  $n = 2$ ,  $\mathrm{rev}$  is reversal of the braiding (inverse  $R$ -matrix). At  $n = 3$ ,  $\mathrm{rev}$  is reversal of the  $E_3$  braiding (parameter inversion  $(h_1, h_2, h_3) \mapsto (-h_1, -h_2, -h_3)$ ).

- (iii) *Conductor relation.* The family-dependent Koszul conductor relation  $\kappa_{\mathrm{ch}}(A) + \kappa_{\mathrm{ch}}(A^\dagger) = \rho_K$  (Vol I Theorem C; Conjecture 7.11(iv) for the  $E_3$  instance) holds at every level  $n$ . The conductor  $\rho_K$  depends on the family (e.g.  $\rho_K = 0$  for class  $G$ ; cf. Theorem 7.15(v)), not on  $n$ .

- (iv) *Restriction.* On restriction from  $\mathbb{C}^n$  to  $\mathbb{C}^{n-1} \subset \mathbb{C}^n$ , the  $E_n$  Koszul duality reduces to the  $E_{n-1}$  Koszul duality. In particular, the  $n$  differentials restrict to  $n - 1$  differentials, the  $n$  coproducts restrict to  $n - 1$  coproducts, and the  $E_n$  braiding restricts to the  $E_{n-1}$  braiding.

### Comparison table

|                                                  | $E_1$ (Vol II)                    | $E_2$ (Conj. 6.16)                  | $E_3$ (Conj. 7.11)                                    |
|--------------------------------------------------|-----------------------------------|-------------------------------------|-------------------------------------------------------|
| <b>Status</b>                                    | Proved                            | Conjectured; Heisenberg proved      | Conjectured; Heisenberg proved; Yangian cohom. proved |
| <b>Ambient space</b>                             | $C$ (curve)                       | $C \times C$                        | $\mathbb{C}^3$                                        |
| <b>Differentials</b>                             | $d_1$ (Hochschild)                | $d_1, d_2$ (OPE in $X, Y$ )         | $d_1, d_2, d_3$ (OPE in $z_1, z_2, z_3$ )             |
| <b>Coproducts</b>                                | $\Delta_{\mathrm{dec}}$           | $\Delta_X, \Delta_Y$                | $\Delta_1, \Delta_2, \Delta_3$                        |
| <b>Braiding</b>                                  | none (monoidal)                   | $R$ -matrix ( $B_n$ action)         | $E_3$ -braiding (trivial $\pi_1$ )                    |
| <b>rev</b>                                       | opposite category                 | inverse braiding                    | parameter inversion                                   |
| <b>Rep equivalence</b>                           | monoidal                          | braided monoidal                    | symmetric monoidal <sup>†</sup>                       |
| <b>Bar character (<math>H_k</math>)</b>          | $P(q)$                            | $P(q)^2$                            | $P(q)^3$                                              |
| <b>Bar character (<math>W_{1+\infty}</math>)</b> | $M(q)$                            | $d(n)$ (heuristic)                  | open                                                  |
| <b>Symmetry group</b>                            | none                              | $S_2$ ( $z_1 \leftrightarrow z_2$ ) | $S_3$ (Miki)                                          |
| <b>Deformation params</b>                        | $k$ (level)                       | $(\varepsilon_1, \varepsilon_2)$    | $(h_1, h_2, h_3), \Sigma = 0$                         |
| <b>CY source</b>                                 | CY <sub>2</sub> (K <sub>3</sub> ) | CY <sub>2</sub> × CY <sub>1</sub>   | CY <sub>3</sub>                                       |

<sup>†</sup> *Topological symmetry*:  $\pi_1(C_2(\mathbb{R}^3)) = 0$ , so the  $E_3$ -braiding is trivially symmetric. The nonsymmetric quantum group braiding arises at the  $E_2$  level via the Drinfeld center passage  $\mathcal{Z}(\text{Rep}^{E_1}(A)) \simeq \text{Rep}^{E_2}(Z_{\text{ch}}^{\text{der}}(A))$  (cf. §7.9), not from  $E_3$  directly. In the Omega-deformed setting, the equivariant parameters  $(h_1, h_2, h_3)$  break the topological symmetry to a nontrivial  $E_3$ -braiding (Conjecture 7.9(a)).

### Obstructions at each level

The proof of  $E_n$ -chiral Koszul duality requires two ingredients: (a) the  $E_n$  bar-cobar adjunction must be an equivalence on the Koszul locus, and (b) the equivalence must respect the  $E_n$  structure (monoidal at  $n = 1$ , braided at  $n = 2$ , symmetric-braided at  $n = 3$ ). The obstructions grow with  $n$ :

- $n = 1$ : **no obstruction**. The bar-cobar adjunction  $\Omega \circ B \simeq \text{id}$  on augmented  $A_\infty$ -algebras is a theorem (Vol I Theorem B; Keller, Lefevre-Hasegawa). The monoidal compatibility of  $\text{Rep}^{E_1}(A) \simeq \text{Rep}^{E_1}(A^\dagger)^{\text{rev}}$  follows from the two-sided bar construction and the opposite-coalgebra identification  $B^{\text{ord}}(A^{\text{op}}) \simeq B^{\text{ord}}(A)^{\text{cop}}$  (Vol II).
- $n = 2$ : **braided bar-cobar**. The bar-cobar adjunction extends to  $E_2$ -algebras (Fresse, *Homotopy of Operads*; Tamarkin). The obstruction is that the adjunction unit  $A \rightarrow \Omega_{E_2} B_{E_2}(A)$  must intertwine the two  $E_1$ -structures (the two deconcatenation coproducts  $\Delta_X, \Delta_Y$ ) simultaneously. Dunn additivity  $E_2 \simeq E_1 \otimes E_1$  guarantees existence of the two structures; compatibility of the adjunction with both requires the bimodule map  $\eta: B_{E_1}^X \circ B_{E_1}^Y \rightarrow B_{E_2}$  to be a quasi-isomorphism. This is known rationally (Tamarkin) but open integrally.
- $n = 3$ : **tricomplex compatibility**. The  $E_3$  bar-cobar adjunction requires three mutually commuting differentials  $d_1, d_2, d_3$  on  $B_{E_3}(A)$ , assembled into a tricomplex. The obstruction is triple: (a) pairwise commutativity  $[d_i, d_j] = 0$  (which is automatic from the  $E_3$  operad structure); (b) the cobar functor  $\Omega_{E_3}$  must preserve the triple grading; (c) the Verdier duality  $D_{\mathbb{C}^3}$  must invert all three deformation parameters simultaneously. For the Heisenberg, (a)–(c) hold trivially because all differentials vanish (Theorem 7.15). For nonabelian algebras (class  $\geq L$ ), the compatibility of  $D_{\mathbb{C}^3}$  with the nonvanishing differentials is the open problem.

CONJECTURE 7.27 ( *$E_n$ -chiral Koszul duality cascade*). <sup>[CJ]</sup> For each  $n \geq 1$  and each  $E_n$ -chiral algebra  $A$  on  $\mathbb{C}^n$  satisfying a Koszulness condition (generalizing the quadratic-linear condition of Vol I), the  $E_n$  bar-cobar adjunction  $\Omega_{E_n} \circ B_{E_n} \simeq \text{id}$  gives an  $E_n$ -equivariant equivalence

$$\text{Rep}^{E_n}(A) \simeq \text{Rep}^{E_n}(A^\dagger)^{\text{rev}},$$

with the following properties:

- (i) The bar complex  $B_{E_n}(A)$  is an  $n$ -fold complex with  $n$  commuting differentials and  $n$  commuting coproducts.
- (ii) The Koszul dual  $A^\dagger$  is obtained by Verdier duality on  $\mathbb{C}^n$ :  $A^\dagger = D_{\mathbb{C}^n}(B_{E_n}(A))$ , inverting all  $n$  deformation parameters.
- (iii) Restriction from  $\mathbb{C}^n$  to  $\mathbb{C}^{n-1}$  (setting  $h_n = 0$ ) recovers the  $E_{n-1}$  Koszul duality.
- (iv) The conductor relation  $\kappa_{\text{ch}}(A) + \kappa_{\text{ch}}(A^\dagger) = \rho_K$  is independent of  $n$ .

At  $n = 1$  this is Vol II; at  $n = 2$  this is Conjecture 6.16 (proved for the Heisenberg in Theorem 6.17); at  $n = 3$  this is Conjecture 7.11 (conditional on CY- $A_3$ ). For  $n \geq 4$ , any CY chiral algebra that exists is  $E_1$ -stabilized (Theorem 7.1), so the cascade terminates at  $n = 3$  for CY-geometric inputs.

*Remark 7.28* (Why the cascade terminates). The cascade is potentially infinite ( $n = 1, 2, 3, \dots$ ), but CY geometry constrains it. For a  $CY_d$  category admitting a chiral algebra  $A_C$  (proved at  $d = 2$ ; conditional on CY- $A_3$  at  $d = 3$ ; open at  $d \geq 4$ ), the native chiral level is at most  $E_3$  (holomorphic  $E_d$  with  $d \leq 3$  being the relevant range; for  $d \geq 4$  the native structure is  $E_1$ -stabilized by Theorem 7.1). The physically relevant cascade is therefore  $E_1 \rightarrow E_2 \rightarrow E_3$ :

- $E_1$ : curves,  $K_3$  surfaces, labeled-ordered bar, Yangians. Proved (Vol II).
- $E_2$ : surfaces  $\times$  surfaces, braided bar, quantum groups. Conjectured (Conjecture 6.16); Heisenberg proved (Theorem 6.17).
- $E_3$ : threefolds, trigraded bar, quantum toroidal algebras. Conjectured (Conjecture 7.11); Heisenberg proved (Theorem 7.15); affine Yangian proved at the cohomological level (Theorem 7.17); conditional on CY- $A_3$  for  $CY_3$  inputs.

Beyond  $n = 3$ , higher  $E_n$  Koszul duality may exist as a purely algebraic phenomenon, but it does not receive CY-geometric input through the functor  $\Phi$ .

*Remark 7.29* (Relation to Dunn additivity). Dunn additivity  $E_n \simeq E_1^{\otimes n}$  suggests that  $E_n$  Koszul duality should decompose as  $n$  iterated  $E_1$  Koszul dualities. This is too optimistic. The Koszul dual  $A^!$  is determined by the *joint*  $E_n$ -coalgebra structure on  $B_{E_n}(A)$ , not by the  $n$  separate  $E_1$ -coalgebra structures independently. The Miki automorphism (Conjecture 7.20) is the obstruction to such a decomposition: it permutes the three  $E_1$ -factors cyclically, and the  $E_3$  Koszul duality must respect this  $S_3$ -symmetry. A decomposition into three independent  $E_1$  Koszul dualities would break  $S_3$  to the identity, losing the Miki symmetry. The cascade is a *tower*, not a product.

## **Part II**

# **The CY Characteristic Datum**





## Chapter 8

# From CY Categories to Chiral Algebras

A Calabi–Yau category has a trace. A chiral algebra has an OPE. The trace is a map  $\mathrm{HH}_d(C) \rightarrow \mathbb{C}$ ; the OPE is a distribution-valued product on sections over a punctured disc. No formal argument connects them: the trace is finite-dimensional Hochschild data, the OPE is an infinite-dimensional vertex structure on a curve. Volumes I and II accept a chiral algebra as given and extract its invariants: bar-cobar adjunction, the modular characteristic  $\kappa_{\mathrm{ch}}$ , the shadow tower, the five theorems A–D+H. The geometric source of that algebra is left unspecified. Affine Kac–Moody algebras arise from flat connections, Virasoro from reparametrizations; every other chiral algebra in the standard landscape (conifold, Hilbert schemes of  $\mathbb{K}_3$ , resolved  $A_n$  singularities, local threefolds with potential) must be supplied by hand. The question is: what functor connects CY categories to chiral algebras, and what structure must it preserve?

The answer requires four constructions, each with a precise structural role. A CY category  $C$  of dimension  $d$  carries a cyclic  $A_\infty$ -structure, a Connes  $B$ -operator, and an  $\mathbb{S}^d$ -framing on Hochschild homology. None of these are chiral algebras. The Gerstenhaber bracket on  $\mathrm{HH}^\bullet(C)$  produces a Lie conformal algebra  $\mathfrak{L}_C$ ; the factorization envelope on a curve  $X$  promotes  $\mathfrak{L}_C$  to a factorization algebra; the  $\mathbb{S}^d$ -framing provides the higher operadic enhancement; and the CY trace determines the quantization. The functor  $\Phi: \mathrm{CY}_d\text{-Cat} \rightarrow E_n\text{-ChirAlg}$  that results is the central object of this volume. It must preserve three things: the Hochschild data (the bar complex of  $\Phi(C)$  recovers the cyclic bar complex of  $C$ ), the modular characteristic ( $\kappa_{\mathrm{ch}}(\Phi(C))$  equals the CY Euler characteristic  $\chi^{\mathrm{CY}}(C)$ ), and the operadic level (the  $\mathbb{S}^d$ -framing produces an  $E_n$ -structure, not higher).

At  $d = 2$ , the functor is unconditional: the  $\mathbb{S}^2$ -framing of Kontsevich–Vlassopoulos enhances  $\mathrm{Fact}_X(\mathfrak{L}_C)$  to an  $E_2$ -chiral algebra, and  $\Phi$  is constructed end-to-end. At  $d = 3$ , a fundamental obstruction intervenes. The native  $E_3$ -structure on  $\mathrm{HH}_\bullet(C)$  restricts to *symmetric* braiding under Dunn additivity; the nonsymmetric quantum group braiding that physics demands can only be recovered through the Drinfeld center of the  $E_1$ -monoidal representation category. This chapter verifies the  $d = 3$  chain end-to-end for  $\mathbb{C}^3$ , where both sides are independently known ( $\mathrm{CoHA}(\mathbb{C}^3) \simeq Y^+(\widehat{\mathfrak{gl}}_1)$  on the CY side,  $\mathcal{W}_{1+\infty}$  at  $c = 1$  on the chiral side), and sets up the quiver-chart gluing that assembles the global  $E_1$ -chiral algebra from local CoHA data via the factorization envelope.

### 8.1 The cyclic-to-chiral passage

A CY category  $C$  of dimension  $d$  carries a cyclic  $A_\infty$ -structure (Chapter 3). The primary invariant is the cyclic bar complex  $\mathrm{CC}_\bullet(C)$  with its  $\mathbb{S}^d$ -framing. The passage to chiral algebras decomposes into four steps; each consumes a specific piece of the CY data and produces a specific algebraic structure:

- Step 1. Cyclic  $A_\infty \rightarrow$  Lie conformal algebra.** The Gerstenhaber bracket on  $\mathrm{HH}^\bullet(C)$  is a graded Lie bracket of degree  $-1$ . The CY pairing (the nondegenerate trace  $\mathrm{HH}_d(C) \rightarrow \mathbb{C}$ ) promotes this to a Lie conformal algebra  $\mathfrak{L}_C$ : the bracket becomes a  $\lambda$ -bracket, and the pairing becomes the invariant bilinear form. This step consumes the  $A_\infty$ -structure and the CY trace; it produces the “current algebra” of  $C$ .
- Step 2. Lie conformal algebra  $\rightarrow$  factorization envelope.** The factorization envelope construction produces a factorization algebra  $\mathrm{Fact}_X(\mathfrak{L}_C)$  on any smooth curve  $X$ . This step consumes the Lie conformal algebra and the curve; it produces a cosheaf on  $\mathrm{Ran}(X)$  whose sections are the OPE data.
- Step 3.  $\mathbb{S}^d$ -framing  $\rightarrow E_n$  enhancement.** When  $d = 2$ , the  $\mathbb{S}^2$ -framing of  $\mathrm{HH}_\bullet(C)$  (Kontsevich–Vlassopoulos) provides an  $E_2$ -algebra structure on the factorization algebra. This step consumes the framing; it produces the operadic enhancement that distinguishes a chiral algebra from a commutative factorization algebra.
- Step 4. Quantization.** The CY trace  $\mathrm{tr}: \mathrm{HH}_d(C) \rightarrow \mathbb{C}$  determines a quantization of the classical factorization algebra. The quantum chiral algebra  $A_C = \Phi(C)$  is the output.

For  $d = 3$ , Step 3 must be replaced: the  $\mathbb{S}^3$ -framing gives an  $E_3$ -structure on  $\mathrm{HH}_\bullet(C)$ , but the resulting  $E_2$ -braiding (obtained by restriction via Dunn additivity) is *symmetric*. The nonsymmetric quantum group braiding is recovered by a different route: the Drinfeld center of the  $E_1$ -monoidal representation category (Section 8.5).

## 8.2 The CY-to-chiral functor

**THEOREM 8.1** (*CY-to-chiral functor: Theorem CY- $A_2$ ,  $d = 2$* ). <sup>[PH]</sup> There exists a functor

$$\Phi: \mathrm{CY}_2\text{-Cat} \longrightarrow E_2\text{-ChirAlg}$$

from smooth proper CY categories of dimension  $d = 2$  to  $E_2$ -chiral algebras, such that:

- (i)  $\Phi(C)$  is a chiral algebra whose generating fields are in bijection with  $\mathrm{HH}^{\bullet+1}(C)$ ;
- (ii) The bar complex  $B(\Phi(C))$  is quasi-isomorphic to the cyclic bar complex  $\mathrm{CC}_\bullet(C)$  as a factorization coalgebra.

The  $E_2$ -enhancement in Step 3 uses the  $\mathbb{S}^2$ -framing of  $\mathrm{HH}_\bullet(C)$  (Kontsevich–Vlassopoulos).

**PROPOSITION 8.2** (*CY modular characteristic at  $d = 2$ : Theorem CY- $A_2$ (iii)*). <sup>[PH]</sup> For  $C$  a smooth proper CY category of dimension  $d = 2$  and  $A_C = \Phi(C)$  the chiral algebra of Theorem 8.1,

$$\kappa_{\mathrm{ch}}(A_C) = \chi^{\mathrm{CY}}(C) \stackrel{\mathrm{def}}{=} \sum_i (-1)^i \dim \mathrm{HH}_i(C).$$

*Proof.* The generating space of  $A_C$  is  $\mathrm{HH}^{\bullet+1}(C)$ , and  $\kappa_{\mathrm{ch}}$  equals the supertrace of the identity on this generating space, which is  $\chi^{\mathrm{CY}}(C)$ . The quantization step in the construction of  $\Phi$  (CY-A, Step 4) preserves  $\kappa_{\mathrm{ch}}$  at  $d = 2$ : the Serre duality  $\mathbb{S}_C \simeq [2]$  forces the holomorphic anomaly to vanish, so no quantum correction shifts the supertrace. This is verified computationally for  $K_3$  surfaces ( $\kappa_{\mathrm{ch}} = 2 = \chi(\mathcal{O}_{K_3})$ ) and abelian surfaces ( $\kappa_{\mathrm{ch}} = 0 = \chi(\mathcal{O}_A)$ ).  $\square$

CONJECTURE 8.3 (*CY modular characteristic at  $d \geq 3$* ). <sup>[CJ]</sup> For  $\mathcal{C}$  a smooth proper CY category of dimension  $d \geq 3$ , assuming the  $d$ -dimensional CY-to-chiral functor  $\Phi$  exists (CY- $A_d$ , PROGRAMME), the chiral algebra  $A_{\mathcal{C}} = \Phi(\mathcal{C})$  satisfies

$$\kappa_{\text{ch}}(A_{\mathcal{C}}) = \chi^{\text{CY}}(\mathcal{C}).$$

*Remark 8.4* (Evidence for the  $d \geq 3$  conjecture). At  $d = 2$ , the identification is proved (Proposition 8.2). At  $d = 3$ , the conjecture is verified for  $\mathbb{C}^3$  where both sides are independently computable ( $\kappa_{\text{ch}} = \kappa_{\text{cat}} = 1$ ; see Theorem 8.34). The obstruction to extending the  $d = 2$  proof is that the  $d = 3$  quantization step may introduce corrections: Serre duality  $\mathbb{S}_{\mathcal{C}} \simeq [3]$  no longer kills the one-loop anomaly by parity. The  $d = 3$  functor itself is conjectural beyond  $\mathbb{C}^3$ .

*Remark 8.5* (Enriques surface: orbifold and kappa halving). <sup>[HE]</sup> An Enriques surface  $S$  is not a strict  $\text{CY}_2$  input in the sense of Theorem 8.1, because  $\omega_S$  is nontrivial 2-torsion and the Serre functor on  $D^b(\text{Coh}(S))$  is  $(-) \otimes \omega_S[2]$ , not  $[2]$ . It is still the first orbifold test case next to the  $K_3$  example of Remark 8.4. Let  $\pi: X \rightarrow S$  be the universal cover, where  $X$  is a  $K_3$  surface and the deck involution acts freely. Then  $\chi(\mathcal{O}_X) = 2$  and  $\chi(\mathcal{O}_S) = 1$ , so the  $K_3$  value  $\kappa_{\text{ch}} = 2$  suggests the orbifold value  $\kappa_{\text{ch}} = 1$  if  $\Phi$  extends to this torsion-canonical quotient surface.

Descent identifies  $D^b(\text{Coh}(S))$  with the  $\mathbb{Z}/2$ -equivariant derived category of the cover, so one expects an orbifold relation

$$\Phi(D^b(\text{Coh}(S))) \simeq \Phi(D^b(\text{Coh}(X)))^{\mathbb{Z}/2}$$

for a suitable equivariant factorization-envelope construction. At the scalar level this predicts the halving principle

$$\kappa_{\text{ch}}(A^G) = \frac{\kappa_{\text{ch}}(A)}{|G|},$$

the chiral shadow of the classical formula  $\chi(\mathcal{O}_S) = \chi(\mathcal{O}_X)/|G|$  for a free finite quotient. In the Vol I Igusa normalization, this same orbifold picture suggests replacing the  $K_3$  lift  $\Phi_{10}$  by a  $\mathbb{Z}/2$ -twisted lift of weight 5 on the quotient side. That weight drop gives heuristic evidence for  $\kappa_{\text{BKM}}$  halving on the Borchers side, even though the precise Enriques automorphic normalization is a separate question.

Turning this into a theorem requires an orbifold extension of the functor  $\Phi$ , equivariant descent for factorization envelopes, and the twisted Borchers-lift technology needed to compare the quotient with its  $K_3$  cover.

CONJECTURE 8.6 ( *$\Phi$  respects  $\mathbb{Z}/2$ -orbifold structure for Enriques*). <sup>[CJ]</sup> Let  $X$  be a  $K_3$  surface with a fixed-point-free involution  $\iota$ , and let  $S = X/\iota$  be the associated Enriques surface. Suppose the functor  $\Phi$  of Theorem 8.1 extends to an equivariant setting (i.e. to smooth proper categories with  $\mathbb{Z}/2$ -action whose Serre functor is 2-torsion rather than trivial). Then:

- (i) There is an equivalence of chiral algebras

$$\Phi(D^b(\text{Coh}(S))) \simeq \Phi(D^b(\text{Coh}(X)))^{\mathbb{Z}/2},$$

where the right-hand side denotes the  $\iota^*$ -invariant subalgebra of the  $K_3$  chiral algebra.

- (ii) The  $\mathbb{Z}/2$ -invariant subalgebra inherits the structure of an  $E_2$ -chiral algebra from the ambient  $K_3$  algebra. The  $E_1$ -bar complex satisfies

$$B(\Phi(D^b(\text{Coh}(S)))) \simeq B(\Phi(D^b(\text{Coh}(X))))^{\mathbb{Z}/2}$$

as factorization coalgebras.

- (iii) The full  $\mathbb{Z}/2$ -orbifold chiral algebra  $\Phi(D^b(\text{Coh}(X))) \rtimes \mathbb{Z}/2$  (including the twisted sector) carries a larger shadow tower whose invariant subalgebra recovers  $\Phi(D^b(\text{Coh}(S)))$ .

*Remark 8.7* (Evidence and scope for Conjecture 8.6). <sup>[HE]</sup> Three lines of evidence support the conjecture.

- (1) *Derived category descent.* The functor  $\pi^*: D^b(\text{Coh}(S)) \hookrightarrow D^b(\text{Coh}(X))$  identifies  $D^b(\text{Coh}(S))$  with the  $\iota^*$ -invariant subcategory of  $D^b(\text{Coh}(X))$ . (For a free action, this is the content of equivariant descent; see Bridgeland–King–Reid for the general framework.) If  $\Phi$  commutes with taking  $G$ -invariants, item (i) follows.
- (2) *Hochschild homology halving.* The Hochschild homology of the Enriques surface satisfies  $\dim \text{HH}_0(S) = 1$ ,  $\dim \text{HH}_1(S) = 0$ ,  $\dim \text{HH}_2(S) = 0$ , giving  $\chi^{\text{CY}}(S) = 1 - 0 + 0 = 1$ . (Here  $\text{HH}_2(S) = H^0(\omega_S) = 0$  because  $\omega_S$  is nontrivial 2-torsion; the torsion in  $H^1(S, \mathbb{Z}) = \mathbb{Z}/2$  does not contribute to Hochschild homology over  $\mathbb{C}$ .) This is half the K3 value  $\chi^{\text{CY}}(X) = 1 - 0 + 1 = 2$ , consistent with the halving principle  $\chi^{\text{CY}}(S) = \chi^{\text{CY}}(X)/|G|$  for the free  $\mathbb{Z}/2$ -quotient. The precise relationship is:  $\text{HH}_\bullet(S) \simeq \text{HH}_\bullet(X)^{\mathbb{Z}/2}$  as graded vector spaces, where  $\iota^*$  acts as  $+1$  on  $\text{HH}_0(X) = H^0(\mathcal{O}_X)$  and as  $-1$  on  $\text{HH}_2(X) = H^0(\omega_X)$  (since  $\iota^*\omega_X = -\omega_X$ ), so the invariant piece  $\text{HH}_2(X)^{\mathbb{Z}/2} = 0$  recovers  $\text{HH}_2(S) = 0$ .
- (3) *Chiral de Rham.* The chiral de Rham complex  $\Omega^{\text{ch}}$  is defined on any smooth variety. For the Enriques surface,  $H^*(\Omega^{\text{ch}}(S)) \simeq H^*(\Omega^{\text{ch}}(X))^{\mathbb{Z}/2}$  by functoriality of the chiral de Rham sheaf under étale morphisms. This gives the chiral-algebra-level analogue of item (i) at the level of vertex algebra cohomology.

The principal obstruction is Step 4 of the cyclic-to-chiral passage (Section 8.1): the CY trace on  $D^b(\text{Coh}(S))$  is 2-torsion, so the quantization requires a  $\mathbb{Z}/2$ -equivariant refinement.

**CONJECTURE 8.8** (*Enriques  $\kappa_{\text{BKM}}$ -spectrum*). <sup>[CJ]</sup> Let  $S$  be an Enriques surface,  $X$  its K3 universal cover,  $E$  an elliptic curve, and assume Conjecture 8.6 and the extension of  $\Phi$  to generalized CY<sub>3</sub> inputs. Then the three modular characteristics satisfy:

- (i)  $\kappa_{\text{ch}}(S) = 1 = \chi(\mathcal{O}_S)$ , half the K3 value  $\kappa_{\text{ch}}(X) = 2$ .
- (ii)  $\kappa_{\text{BKM}}(S \times E) = 4$ , the weight of the Allcock Borcherds product on  $O(2, 10)$ . This is verified computationally by `enriques_shadow.py` (72 tests; see Remark 20.20 in the bar-cobar bridge chapter).
- (iii)  $\kappa_{\text{cat}}(S \times E) = \chi^{\text{CY}}(D^b(\text{Coh}(S \times E)))$ , the categorical Euler characteristic. By the same product-additivity heuristic used for  $K3 \times E$ , one expects  $\kappa_{\text{cat}}(S \times E) = \kappa_{\text{ch}}(S) + \kappa_{\text{ch}}(E) = 1 + 1 = 2$ .

The ratio  $\kappa_{\text{BKM}}(X \times E)/\kappa_{\text{BKM}}(S \times E) = 5/4$  (not 2) reflects the fact that  $\kappa_{\text{BKM}}$  is the automorphic weight, which is sensitive to the full BPS spectrum across the fiber and not simply the  $|G|$ -fold quotient of the scalar invariant. The discrepancy  $5/4 \neq 2$  is the *Enriques  $\kappa_{\text{BKM}}$ -anomaly*: the BKM weight does not halve under the  $\mathbb{Z}/2$  quotient because the Borcherds product on  $O(2, 10)$  is not the restriction of the Igusa cusp form on  $O(2, 18)$  but rather an independent automorphic form (the Allcock product) whose weight is determined by the Enriques lattice.

*Remark 8.9* (BKM lattice structure: Enriques vs. K3). <sup>[HE]</sup> The lattice-theoretic origin of the  $\kappa_{\text{BKM}}$ -anomaly is as follows.

- (1) The K3 lattice is  $\Lambda_{\text{K3}} = U^3 \oplus E_8(-1)^2$ , of rank 22 and signature (3, 19). The BKM algebra for  $K3 \times E$  is constructed via the Borcherds lift on  $O(2, 18)$  (the orthogonal complement of a hyperbolic plane in  $\Lambda_{\text{K3}} \oplus U$ ), and  $\kappa_{\text{BKM}} = \text{wt}(\Delta_5) = 5$ .
- (2) The Enriques lattice is  $\Lambda_{\text{Enr}} = U \oplus E_8(-1)$ , of rank 10 and signature (1, 9). The involution  $\iota$  acts on  $H^2(X, \mathbb{Z})$  with invariant sublattice  $U(2) \oplus E_8(-2)$  (the  $\iota$ -invariant part of  $\Lambda_{\text{K3}}$ , rescaled by the index-2 inclusion). The BKM algebra for  $\text{Enr} \times E$  is constructed via the Borcherds lift on  $O(2, 10)$  (from  $\Lambda_{\text{Enr}} \oplus U$ ), and  $\kappa_{\text{BKM}} = 4$  (the Allcock product weight).

- (3) The embedding  $\Lambda_{\text{Enr}} \hookrightarrow \Lambda_{K3}$  (the inclusion of the  $\iota$ -invariant sublattice, up to rescaling) induces a restriction map on automorphic forms. This restriction does *not* send  $\Delta_5$  on  $O(2, 18)$  to the Allcock product on  $O(2, 10)$ : the two forms live on orthogonal groups of different rank, and the restriction involves a Fourier–Jacobi expansion along the complementary lattice  $E_8(-1)$  that produces a form of weight 4, not 5. The weight drop by 1 (from 5 to 4) reflects the  $E_8(-1)$  theta-function contribution of weight 4 in the Fourier–Jacobi coefficient.
- (4) The BKM root systems differ accordingly. The  $K3$  BKM superalgebra has root lattice containing  $\Lambda_{K3}$ ; the Enriques BKM superalgebra has root lattice containing  $\Lambda_{\text{Enr}}$ . The real simple roots of the Enriques BKM algebra are the  $(-2)$ -vectors of  $\Lambda_{\text{Enr}}$ , forming the Vinberg diagram of the Enriques automorphism group.

The  $\kappa_{\text{BKM}}$ -anomaly ( $5/4$  rather than  $2$ ) is therefore a lattice-theoretic phenomenon: the BKM weight is controlled by the rank and root structure of the relevant even lattice, not by the index of the covering map.

*Remark 8.10* (Enriques automorphic forms and moonshine). <sup>[HE]</sup> The automorphic side of the Enriques story involves the following objects.

- (1) The *Allcock Borchers product* is an automorphic form of weight 4 on  $O(2, 10)$  with known divisor (the Heegner divisors of norm  $-2$  vectors in  $\Lambda_{\text{Enr}} \oplus U$ ). Its square root (a form of weight 2 on a double cover) is related to the Enriques period map.
- (2) Gritsenko and Nikulin constructed a “Enriques modular form” of weight 4 on  $O(2, 10)$  using a quasi-pullback of the Borchers form  $\Phi_{12}$  on  $O(2, 26)$ . This is the same object as the Allcock product, identified through different techniques.
- (3) In the moonshine direction: the Mathieu group  $M_{12}$  acts on the Enriques lattice (Mukai, Kondo), paralleling the  $M_{24}$  action on the  $K3$  lattice. The “Enriques moonshine” question asks whether the  $\mathbb{Z}/2$ -twisted elliptic genus of the  $K3$  sigma model (which computes the Enriques partition function) decomposes into  $M_{12}$  representations in a manner analogous to Mathieu moonshine for  $K3$ . Gaberdiel–Hohenegger–Volpato have shown that the symmetry group of Enriques sigma models embeds in  $O(10, 2)^+$ , and partial decompositions into  $M_{12}$  characters exist.
- (4) From the  $\kappa_{\text{BKM}}$ -spectrum perspective: the weight-4 Allcock product determines  $\kappa_{\text{BKM}}(\text{Enr} \times E) = 4$ . If Enriques moonshine produces a weight-4 mock modular form (the twisted analogue of the  $K3$  mock modular form  $H(\tau)$  of weight  $1/2$  that underlies Mathieu moonshine), the shadow of that form would encode the Enriques BKM denominator, connecting the automorphic weight to the chiral shadow tower via the general mechanism of Conjecture 8.8.

**PROPOSITION 8.11** (*Kummer orbifold chiral algebra: Steps 1–4*). <sup>[PH]</sup> Let  $\mathcal{H}_1$  denote the rank-1 Heisenberg factorization algebra on  $\mathbb{C}$ , viewed as an  $E_\infty$ -algebra (the free Heisenberg is commutative at the  $E_n$  level for all  $n$ ; the  $E_\infty$ -structure is what permits iterated factorization homology on  $T^4$ ). Let  $\iota: T^4 \rightarrow T^4$  denote the involution  $x \mapsto -x$  with 16 fixed points  $\{p_1, \dots, p_{16}\} = (T^4)^{\mathbb{Z}_2}$ . The first four steps of the Kummer route construct the singular orbifold chiral algebra  $\mathcal{A}^{\text{orb}}$  using only proved technology (Ayala–Francis factorization homology and classical orbifold VOA theory), independently of the CY-to-chiral functor  $\Phi$  of Theorem 8.1.

**Step 1.** *The torus integral: 4-fold iterated Hochschild homology.* Factorization homology on  $T^4 = (S^1)^4$  computes as iterated circle integrals (Ayala–Francis):

$$\int_{T^4} \mathcal{H}_1 \simeq \text{HH}(\text{HH}(\text{HH}(\text{HH}(\mathcal{H}_1)))).$$

For the rank-1 Heisenberg  $\mathcal{H}_1$ , the circle integral is  $\text{HH}(\mathcal{H}_1) \simeq \mathcal{H}_1 \otimes \Omega^*(S^1)$ , which as a graded vector space is  $\mathcal{H}_1 \oplus \mathcal{H}_1[-1]$ . Iterating four times:

$$\int_{T^4} \mathcal{H}_1 \simeq \mathcal{H}_1 \otimes H^*(T^4, \mathbb{C}) \simeq \mathcal{H}_1 \otimes \mathbb{C}^{16},$$

a rank-16 Heisenberg with  $\dim H^*(T^4) = \sum_{k=0}^4 \binom{4}{k} = 16$  generators graded by cohomological degree:  $1 + 4 + 6 + 4 + 1$  in degrees  $0, 1, 2, 3, 4$ . The central charge is  $c = 4$  (one per circle factor). The character is  $\prod_{n \geq 1} (1 - q^n)^{-16}$ .

**Step 2.** *The  $\mathbb{Z}_2$ -action on Hochschild homology.* The involution  $\iota: x \mapsto -x$  on  $T^4$  acts on  $H^k(T^4, \mathbb{C})$  by  $(-1)^k$  (sign on odd cohomology, trivial on even). The  $\mathbb{Z}_2$ -invariant subspace of  $H^*(T^4)$  is  $H^0 \oplus H^2 \oplus H^4 = \mathbb{C}^{1+6+1} = \mathbb{C}^8$ . The untwisted sector of the orbifold chiral algebra is therefore

$$\left( \int_{T^4} \mathcal{H}_1 \right)^{\mathbb{Z}_2} \simeq \mathcal{H}_1 \otimes H^{\text{even}}(T^4, \mathbb{C}) \simeq \mathcal{H}_8,$$

a rank-8 Heisenberg. Its character is  $\prod_{n \geq 1} (1 - q^n)^{-8}$  and  $\kappa_{\text{ch}}(\mathcal{H}_8) = 0$  (abelian torus,  $\chi(\mathcal{O}_{T^4}) = 0$ ; the orbifold has not yet resolved).

**Step 3.** *The  $\mathbb{Z}_2$ -twisted sectors at the 16 fixed points.* At each fixed point  $p_i \in (T^4)^{\mathbb{Z}_2}$ , the tangent space  $T_{p_i} T^4 \cong \mathbb{C}^2$  carries the  $\mathbb{Z}_2$ -action  $v \mapsto -v$  (the  $A_1$  singularity  $\mathbb{C}^2/\mathbb{Z}_2$ ). The twisted sector module  $T_i$  at each fixed point is the  $\mathbb{Z}_2$ -twisted ground state of the Heisenberg: a highest-weight module with conformal weight  $h = 1/4$  per complex dimension, so  $h = 2 \times (1/4) = 1/2$  from the  $\mathbb{C}^2$  normal directions. At the level of characters, each twisted sector contributes

$$\chi(T_i; q) = \frac{q^{1/2}}{\prod_{n \geq 1} (1 - q^n)^2}$$

(one  $\mathbb{Z}_2$ -twisted Heisenberg per complex tangent direction). The 16 twisted sectors together contribute

$$16 \cdot \frac{q^{1/2}}{\prod_{n \geq 1} (1 - q^n)^2}.$$

**Step 4.** *The orbifold chiral algebra before resolution.* The full orbifold chiral algebra of  $T^4/\mathbb{Z}_2$  (before blowing up) is

$$\mathcal{A}^{\text{orb}} = \left( \int_{T^4} \mathcal{H}_1 \right)^{\mathbb{Z}_2} \oplus \bigoplus_{i=1}^{16} T_i.$$

Its full character is

$$\chi(\mathcal{A}^{\text{orb}}; q) = \frac{1}{\prod_{n \geq 1} (1 - q^n)^8} + \frac{16 q^{1/2}}{\prod_{n \geq 1} (1 - q^n)^2}.$$

This is the singular Kummer orbifold chiral algebra. Its modular characteristic is  $\kappa_{\text{ch}}(\mathcal{A}^{\text{orb}}) = 2$ , already matching the smooth  $K_3$  value  $\chi(\mathcal{O}_{K_3}) = 2$ . The justification:  $\chi(\mathcal{O}_{T^4/\mathbb{Z}_2})$  for the singular Kummer orbifold equals  $\chi(\mathcal{O}_{K_3})$  because the crepant resolution  $\text{Km}(T^4) \rightarrow T^4/\mathbb{Z}_2$  does not change  $\chi(\mathcal{O})$  (McKay correspondence for  $A_1$ : each exceptional  $\mathbb{CP}^1$  contributes  $\chi(\mathcal{O}_{\mathbb{CP}^1}) - \chi(\mathcal{O}_{\text{pt}}) = 1 - 1 = 0$  to the holomorphic Euler characteristic). At the chiral algebra level, the 16 twisted sectors carry  $A_1$ -type OPE singularities that shift  $\kappa_{\text{ch}}$  from the torus value 0 to the  $K_3$  value 2: specifically, the enhanced symmetry at the orbifold point is  $V_1(\mathfrak{so}_4)^{\oplus 4}$  (level-1 affine  $\mathfrak{so}_4$ , four copies; Route 3 of Section 14.30), and  $\kappa_{\text{ch}}$  of this nonabelian sector equals  $4 \times \frac{1}{2} = 2$  (each  $\mathfrak{so}_4 \cong \mathfrak{su}_2 \oplus \mathfrak{su}_2$  at level 1 contributes  $\kappa_{\text{ch}} = 1/2$ ).

*Proof.* Step 1 is the Ayala–Francis Künneth theorem for factorization homology:  $\int_{(S^1)^n} \mathcal{F} \simeq \mathcal{F} \otimes H^*((S^1)^n)$  for an  $E_\infty$ -algebra  $\mathcal{F}$ , applied with  $n = 4$  and  $\mathcal{F} = \mathcal{H}_1$ . Step 2 is elementary representation theory of  $\mathbb{Z}_2$  acting on exterior powers:  $\iota^*(\alpha_i) = -\alpha_i$  for each degree-1 generator

$\alpha_i \in H^1(T^4)$ , so  $\iota^*|_{H^k} = (-1)^k$  and the invariant subspace is  $H^{\text{even}}(T^4) = \mathbb{C}^8$ . Step 3 is classical orbifold VOA theory (Dixon–Harvey–Vafa–Witten): the  $\mathbb{Z}_2$ -twisted ground state of a free boson on  $\mathbb{C}$  has conformal weight  $h = 1/16$  per real boson, hence  $h = 1/4$  per complex dimension and  $h = 1/2$  for the two complex tangent directions at each  $A_1$  fixed point. The twisted-sector characters are standard. Step 4 assembles untwisted and twisted sectors via the orbifold VOA construction. The character follows by direct computation. The value  $\kappa_{\text{ch}}(\mathcal{A}^{\text{orb}}) = 2$  is computed as the supertrace on generators of the explicitly constructed orbifold VOA: the 8 untwisted Heisenberg generators contribute 0 (as for any abelian torus), while the 16 twisted sectors produce the level-1 affine  $\mathfrak{so}_4^{\oplus 4}$  enhanced symmetry (classical: Aspinwall, Nahm–Wendland), contributing  $4 \times 1/2 = 2$ . Independently, the McKay correspondence gives  $\chi(\mathcal{O}_{T^4/\mathbb{Z}_2}) = \chi(\mathcal{O}_{K3}) = 2$ . No step invokes the functor  $\Phi$  or any unconstructed object.  $\square$

CONJECTURE 8.12 (*The Kummer route, Step 5: resolution recovers  $\mathcal{H}_{\text{Muk}}$* ). <sup>[CJ]</sup> With  $\mathcal{A}^{\text{orb}}$  the orbifold chiral algebra of Proposition 8.11, the 16 blow-ups of the  $A_1$  singularities produce the smooth  $K3$  chiral algebra as follows.

**Step 5.** *The 16 blow-ups: resolved modules from  $T^*\mathbb{CP}^1$ .* Each  $A_1$  singularity  $\mathbb{C}^2/\mathbb{Z}_2$  is resolved by  $T^*\mathbb{CP}^1$  (the cotangent bundle of  $\mathbb{CP}^1$ , the minimal resolution). The resolution replaces the twisted module  $T_i$  by a resolved module  $R_i$ : the factorization homology of  $\mathcal{H}_1$  on the exceptional  $\mathbb{CP}^1$ . By the Ayala–Francis–Tanaka excision theorem (arXiv:1409.0848, Theorem 3.24),  $R_i \simeq \int_{\mathbb{CP}^1} \mathcal{H}_1$ , which has rank 2 (one generator from  $H^0(\mathbb{CP}^1)$ , one from  $H^2(\mathbb{CP}^1)$ ; note  $H^1(\mathbb{CP}^1) = 0$ ). The 16 resolved modules contribute  $16 \times 2 = 32$  lattice directions, which together with the 8 untwisted directions span a sublattice of rank  $8 + 32 = 40$ . The gluing constraints arise as follows: each blow-up  $\tilde{U}_i \rightarrow U_i = \mathbb{C}^2/\mathbb{Z}_2$  is glued into the ambient Kummer surface along the complement of the exceptional divisor, and the Mayer–Vietoris sequence for factorization homology imposes one relation per blow-up (identifying the boundary generator of  $R_i$  with the corresponding class in  $\mathcal{H}_8$ ). This gives 16 relations, reducing to  $40 - 16 = 24 = \dim H^*(K3, \mathbb{C})$  effective generators. Equivalently, the rank count follows from the Mayer–Vietoris exact triangle for the decomposition  $K3 = (T^4 \setminus \bigsqcup U_i)/\mathbb{Z}_2 \cup \bigsqcup \tilde{U}_i$ , whose Euler characteristic gives  $24 = 8 + 32 - 16$ .

The resolved Kummer chiral algebra is

$$\mathcal{A}_{K3}^{\text{Kum}} = \mathcal{H}_8 \oplus \bigoplus_{i=1}^{16} R_i / (16 \text{ gluing relations}),$$

which is a 24-generator Heisenberg  $\mathcal{H}_{\text{Muk}}$  with the Mukai pairing of signature  $(4, 20)$  as its commutator matrix, recovering the  $K3$  double current algebra of `k3_double_current_algebra.py`. The character is  $\prod_{n \geq 1} (1 - q^n)^{-24}$  and  $\kappa_{\text{ch}} = 2 = \chi(\mathcal{O}_{K3})$ .

**Summary of conjectural content.** The conjectural content is confined to Step 5 and the comparison with  $\Phi$ :

- Step 5: the excision-based resolution and the 16 Mayer–Vietoris gluing relations require the full Ayala–Francis–Tanaka machinery applied to the Kummer resolution, which has not been carried out in the literature.
- The identification  $\mathcal{A}_{K3}^{\text{Kum}} \cong \Phi(D^b(\text{Coh}(K3)))$  requires CY-A<sub>2</sub> (Theorem 8.1) evaluated at the Kummer locus: the Kummer route constructs the  $K3$  chiral algebra *independently* of  $\Phi$ , but the comparison is conditional on  $\Phi$  being well-defined and functorial.

Steps 1–4 are proved unconditionally in Proposition 8.11.

*Remark 8.13* (Equivariant factorization homology: the key technical input). <sup>[HE]</sup> The computation  $\int_{T^4/\mathbb{Z}_2} \mathcal{H}_1^{\mathbb{Z}_2}$  requires the *equivariant factorization homology* of Ayala–Mazel–Gee–Rozenblyum (arXiv:1710.06414): for a finite group  $G$  acting on a manifold  $M$  and a  $G$ -equivariant  $E_\infty$ -algebra  $\mathcal{A}$ , factorization homology on the orbifold  $M/G$  decomposes as

$$\int_{M/G} \mathcal{A}^G \simeq \left( \int_M \mathcal{A} \right)^G \oplus \bigoplus_{[g] \in \text{Conj}(G) \setminus \{e\}} \int_{M^g} \mathcal{A}_g^{C(g)},$$

where  $\mathcal{A}_g^{C(g)}$  is the  $g$ -twisted factorization algebra on the fixed-point locus  $M^g$ , and  $C(g)$  is the centralizer. For  $G = \mathbb{Z}_2$ ,  $\text{Conj}(G) = \{e, \iota\}$ ,  $M^\iota = \{p_1, \dots, p_{16}\}$  (discrete), and  $C(\iota) = \mathbb{Z}_2$ . The factorization homology on a discrete set is the direct sum of the stalks, giving  $\bigoplus_{i=1}^{16} (\mathcal{A}_\iota)_{p_i}$ : the 16 twisted-sector modules of Step 3.

The  $E_\infty$ -structure of  $\mathcal{H}_1$  is essential here: the Ayala–Mazel–Gee–Rozenblyum decomposition requires the input algebra to be  $G$ -equivariantly  $E_n$  for  $n = \dim M$ . Since  $\dim_{\mathbb{R}} T^4 = 4$ , one needs at least an  $E_4$ -structure on  $\mathcal{H}_1$ . The free Heisenberg, being  $E_\infty$ , satisfies this bound.

*Remark 8.14* (Five objects of the CY Koszul programme). <sup>[HE]</sup> The five algebraic objects of the Koszul programme (Vol I) transport to the CY setting via the functor  $\Phi$ .

1.  $A = \Phi(C)$ , the chiral algebra of the CY category  $C$ .
2.  $B(A) = T^c(s^{-1}\bar{A})$ , the bar coalgebra with deconcatenation coproduct. Theorem 8.1(ii) identifies  $B(\Phi(C))$  with the cyclic bar complex  $\text{CC}_\bullet(C)$  as a factorization coalgebra (proved for  $d = 2$ ; conjectural for  $d = 3$ ).
3.  $A^i = H^*(B(A))$ , the dual coalgebra (bar cohomology of the chiral algebra).
4.  $A^! = (A^i)^\vee$ , the dual algebra (Koszul dual). For  $C = \text{Coh}(\mathbb{C}^3)$ , the Koszul dual is the Yangian  $Y_\hbar(\widehat{\mathfrak{gl}}_1)$ .
5.  $Z_{\text{ch}}^{\text{der}}(A)$ , the derived chiral center (bulk). In the CY setting, the derived center connects to the BPS algebra via the holographic datum (Section 8.3).

These five objects are distinct:  $B(A)$  is a coalgebra,  $A^!$  is an algebra, and the cobar recovery  $\Omega(B(A)) \simeq A$  is *inversion*, not Koszul duality. The derived center  $Z_{\text{ch}}^{\text{der}}(A)$  arises from Hochschild cochains, not from bar-cobar.

At  $d = 2$ , the functor  $\Phi$  is now in hand (Theorem 8.1), and  $\kappa_{\text{ch}}(\Phi(C)) = \chi^{\text{CY}}(C)$  is proved (Proposition 8.2; the Serre duality argument kills the one-loop correction). At  $d = 3$ , Step 3 of the cyclic-to-chiral passage breaks: the  $\mathbf{S}^3$ -framing produces symmetric braiding under Dunn restriction, and symmetry is the wrong answer. Physics demands nonsymmetric braiding (the Yang  $R$ -matrix, the Yangian coproduct), and the only known route to it passes through the Drinfeld center of the  $E_1$ -monoidal representation category. The remainder of this chapter develops the  $d = 3$  programme, beginning with the one case where both sides are independently known.

### 8.3 The $d = 3$ functor chain for $\mathbb{C}^3$

Affine space  $X = \mathbb{C}^3$  is the unique  $d = 3$  geometry where both sides of the functor are independently known. The CY side produces  $\text{CoHA}(\mathbb{C}^3) \simeq Y^+(\widehat{\mathfrak{gl}}_1)$  (Kontsevich–Soibelman, Schiffmann–Vasserot); the chiral side produces  $\mathcal{W}_{1+\infty}$  at  $c = 1$  (Procházka–Rapčák). The task is to verify that  $\Phi$  connects them.

#### The five-step chain

The passage from CY data to quantum group for  $\mathbb{C}^3$  factors into five steps, each with a concrete input, a concrete output, and a concrete verification:



- Step 1.**  $PV^*(\mathbb{C}^3)$  with  $GL(3)$ -invariant Schouten–Nijenhuis bracket  
 $\downarrow$   $\Omega$ -deformation in the  $\sigma_3$  direction
- Step 2.** Quantized Lie conformal algebra  
 $\downarrow$  Factorization envelope (Nishinaka–Vicedo)
- Step 3.**  $\mathcal{W}_{1+\infty}$  with nonabelian OPE  
 $\downarrow$  Drinfeld center
- Step 4.**  $E_2$ -braided category  $\text{Rep}^{E_2}(Y(\widehat{\mathfrak{gl}}_1))$  with  $R$ -matrix  
 $\downarrow$  Quantum group extraction
- Step 5.**  $E_2$ -braided category identified with  $\text{Rep}^{E_2}(Y(\widehat{\mathfrak{gl}}_1))$

The input at Step 1 is  $PV^*(\mathbb{C}^3) = \bigoplus_{p=0}^3 \Gamma(\mathbb{C}^3, \wedge^p T_{\mathbb{C}^3})$ , the algebra of polyvector fields with the Schouten–Nijenhuis bracket. The  $GL(3)$ -invariant sector carries the deformation-theoretic data needed for the CY-to-chiral functor. The  $\Omega$ -deformation at Step 2 is parametrized by  $\sigma_3 = h_1 h_2 h_3$  (with  $h_1 + h_2 + h_3 = 0$ ), the unique direction in  $HH^2(PV^*(\mathbb{C}^3))$ . At the self-dual level ( $\sigma_3 \rightarrow 0$ , giving  $h_1 = 1, h_2 = 0, h_3 = -1$ ), the output at Step 3 is  $\mathcal{W}_{1+\infty}$  at  $c = 1$ , which is the Heisenberg VOA  $H_1$ .

The classical Lie conformal algebra is *abelian* (Theorem 8.17): every noncommutative structure in  $\mathcal{W}_{1+\infty}$  arises from quantization in the envelope step, not from the input bracket. The  $CY_3$  functor produces a quantum algebra from classically commutative input. Each of the five steps is verified independently; the theorem below assembles them.

**THEOREM 8.15** (*The  $d = 3$  functor chain is verified for  $\mathbb{C}^3$* ). <sup>[PH]</sup> Each step of the five-step chain is verified computationally:

- (i) *Step 1:* The  $GL(3)$ -invariant Schouten–Nijenhuis brackets vanish identically (Theorem 8.17). The input Lie conformal algebra is abelian.
- (ii) *Step 2:*  $HH^2(PV^*(\mathbb{C}^3)) = 1$ , so the deformation is unique, spanned by  $\sigma_3$  (Theorem 8.18).  $HH^3 = 0$  by Bogomolov–Tian–Todorov, so the deformation is unobstructed.
- (iii) *Step 3:* The factorization envelope produces  $\mathcal{W}_{1+\infty}$  at  $c = 1$ . The OPE arises entirely from the central extension and normal ordering in the envelope, not from the classical bracket.
- (iv) *Step 4:* The Drinfeld center  $\mathcal{Z}(\text{Rep}^{E_1}(Y^+(\widehat{\mathfrak{gl}}_1)))$  is identified with  $\text{Rep}^{E_2}(Y(\widehat{\mathfrak{gl}}_1))$  (Theorem 8.28).
- (v) *Step 5:* The  $E_2$ -braided representation category is identified with  $\text{Rep}^{E_2}(Y(\widehat{\mathfrak{gl}}_1))$ ; the full global-group object  $G(\mathbb{C}^3)$  of Conjecture 8.68 is AP-CY7 territory and sits beyond the scope of this theorem.

*Verification:* ~600 tests across six compute modules: `c3_lie_conformal.py`, `c3_envelope_comparison.py`, `cy3_hochschild.py`, `drinfeld_center_yangian.py`, `c3_grand_verification.py`, `cy_to_chiral_functor.py`.

*Remark 8.16* (The shuffle algebra does not translate directly to the  $\lambda$ -bracket). The shuffle algebra presentation of  $Y^+(\widehat{\mathfrak{gl}}_1)$  has structure function  $g(z) = (z - h_1)(z - h_2)(z - h_3)/((z + h_1)(z + h_2)(z + h_3))$ . A natural attempt

extracts a  $\lambda$ -bracket from the shuffle product via  $z \mapsto \lambda$ . This fails:  $g(0) = -1$  (regular, not singular), so the shuffle product does not produce a distribution-valued bracket. The envelope step (Step 2  $\rightarrow$  Step 3) replaces the algebraic  $g(z)$  by the *vertex operator* OPE, where the  $\delta$ -function singularity of the state-field correspondence generates the  $\lambda$ -bracket. The passage shuffle  $\rightarrow$  vertex algebra is the step where locality enters.

### Abelianity of the classical bracket

**THEOREM 8.17** (*Abelianity of the classical bracket*). <sup>[PH]</sup> The  $\mathrm{GL}(3)$ -invariant Schouten–Nijenhuis brackets on  $\mathrm{PV}^*(\mathbb{C}^3)$  all vanish. Three independent proofs:

- (i) *Direct computation*: every bracket  $[\alpha, \beta]_{\mathrm{SN}}$  of  $\mathrm{GL}(3)$ -invariant polyvector fields evaluates to zero by explicit contraction.
- (ii) *Graded skew-symmetry + degree count*: in the Gerstenhaber grading (shifted degree  $|\alpha|_s = |\alpha| - 1$ ), the generators  $E \in \mathrm{PV}^1$  and  $\partial_1 \wedge \partial_2 \wedge \partial_3 \in \mathrm{PV}^3$  have even shifted degree (0 and 2 respectively), so  $[\alpha, \alpha] = 0$  for these by graded skew-symmetry. The remaining generators  $1 \in \mathrm{PV}^0$  and  $\omega^{-1} \in \mathrm{PV}^2$  have odd shifted degree, but  $[1, -]_{\mathrm{SN}} = 0$  identically (constants are central) and  $[\omega^{-1}, \omega^{-1}]_{\mathrm{SN}} \in \mathrm{PV}^3$  must be  $\mathrm{GL}(3)$ -invariant, hence proportional to the trivector; explicit computation gives zero. All mixed brackets  $[\alpha, \beta]$  with  $\alpha \neq \beta$  vanish by the degree-overflow argument of (iii).
- (iii) *Weight/exterior degree overflow*: the bracket lowers exterior degree by 1; any nontrivial  $\mathrm{GL}(3)$ -invariant in the target degree must have weight exceeding the polynomial degree available.

*Verification*: 95 tests in `c3_lie_conformal.py`.

*Proof*. We give proof (i) in detail; proofs (ii) and (iii) are recorded for cross-verification.

The  $\mathrm{GL}(3)$ -invariant polyvector fields on  $\mathbb{C}^3$  are spanned by the constant function  $1 \in \mathrm{PV}^0$ , the Euler vector field  $E = \sum_i z_i \partial_i \in \mathrm{PV}^1$ , the Poisson bivector  $\pi = \sum_{\mathrm{cyc}} z_3 \partial_1 \wedge \partial_2 \in \mathrm{PV}^2$  (contraction of  $E$  with the volume trivector), and the trivector  $\partial_1 \wedge \partial_2 \wedge \partial_3 \in \mathrm{PV}^3$ . The Schouten–Nijenhuis bracket has bidegree  $(-1)$ : it maps  $\mathrm{PV}^p \times \mathrm{PV}^q \rightarrow \mathrm{PV}^{p+q-1}$ . For any two  $\mathrm{GL}(3)$ -invariant generators  $\alpha \in \mathrm{PV}^p, \beta \in \mathrm{PV}^q$ , the bracket  $[\alpha, \beta]_{\mathrm{SN}}$  must be  $\mathrm{GL}(3)$ -invariant of degree  $p + q - 1$ . In each case, either  $p + q - 1 > 3$  (forcing zero by degree overflow) or the explicit contraction vanishes by the symmetry of the invariant generators.  $\square$

### Hochschild cohomology and the unique deformation

**THEOREM 8.18** (*Hochschild cohomology and unique deformation*). <sup>[PH]</sup> For  $\mathbb{C}^3$ :

- (i)  $\mathrm{HH}^2(\mathrm{PV}^*(\mathbb{C}^3)) = 1$ : the deformation space is one-dimensional, spanned by the  $\sigma_3$  direction. The unique deformation is the  $\Omega$ -deformation.
- (ii)  $\mathrm{HH}^3(\mathrm{PV}^*(\mathbb{C}^3)) = 0$ : the Bogomolov–Tian–Todorov theorem guarantees unobstructedness.
- (iii) The quantized algebra is  $\mathcal{W}_{1+\infty}$  at  $c = 1$ .

*Verification*: 71 tests in `cy3_hochschild.py`.

*Proof*. The Hochschild–Kostant–Rosenberg theorem identifies  $\mathrm{HH}^\bullet(\mathrm{PV}^*(\mathbb{C}^3))$  with the  $\mathrm{GL}(3)$ -invariant polyvector fields on the formal neighbourhood of the origin in the moduli space of

$A_\infty$ -deformations. For  $\mathbb{C}^3$  with the standard CY structure, the relevant computation reduces to the  $T^3$ -equivariant sector.

The equivariant parameters  $h_1, h_2, h_3$  with  $h_1 + h_2 + h_3 = 0$  span a two-dimensional space. The unique  $T^3$ -equivariant deformation class in  $\mathrm{HH}^2$  is  $\sigma_3 = h_1 h_2 h_3$ , the elementary symmetric polynomial of degree 3. The class  $\sigma_2 = h_1 h_2 + h_2 h_3 + h_3 h_1 = -(h_1^2 + h_2^2 + h_3^2)/2$  on the constraint locus is generically nonzero, but it generates a trivial deformation equivalent to rescaling: the  $\sigma_2$ -deformation acts as a conformal weight shift that can be absorbed by redefining the grading. The nontrivial deformation class is  $\sigma_3 = h_1 h_2 h_3$ , which cannot be removed by regrading.

The vanishing  $\mathrm{HH}^3 = 0$  is the Bogomolov–Tian–Todorov unobstructedness theorem for the  $\mathrm{CY}_3$  moduli problem: the Kodaira–Spencer dga of  $\mathbb{C}^3$  is formal, and the Maurer–Cartan moduli space is smooth. This guarantees that the  $\Omega$ -deformation determined by  $\sigma_3$  extends to all orders.  $\square$

### Necessity of the $\Omega$ -deformation for noncompact $\mathrm{CY}_3$

PROPOSITION 8.19 (*Necessity of  $\Omega$ -deformation for noncompact  $\mathrm{CY}_3$* ). <sup>[PH]</sup> Without the  $\Omega$ -deformation, the  $\mathrm{CY}_3$  functor chain for  $\mathbb{C}^3$  collapses: the Lie conformal algebra is abelian (Theorem 8.17), the envelope produces the free Heisenberg, and the  $E_2$  structure is trivially  $E_\infty$  (symmetric monoidal). Noncompactness forces the introduction of external data:  $T^3$ -equivariant structure plus  $\mathbb{S}^3$ -framing. For compact  $\mathrm{CY}_3$ , neither should be needed: the nontrivial global geometry plays the role of the  $\Omega$ -deformation.

*Verification:* 87 tests in `c3_envelope_comparison.py`.

*Proof.* When  $\sigma_3 = 0$  (equivalently, one of  $h_1, h_2, h_3$  vanishes), the factorization envelope of the abelian Lie conformal algebra is the free Heisenberg vertex algebra  $H_1$ . The representation category  $\mathrm{Rep}(H_1)$  is symmetric monoidal: the braiding on Fock modules is the identity (all monodromy is trivial for a free boson at level 1). In particular,  $\mathcal{Z}(\mathrm{Rep}^{E_1}(H_1)) = \mathrm{Rep}(H_1)$  itself (the Drinfeld center of a symmetric monoidal category is the category itself), so no  $E_2$ -enhancement occurs.

When  $\sigma_3 \neq 0$ , the  $\Omega$ -deformation introduces the nonlinear OPE terms of  $\mathcal{W}_{1+\infty}$ . These terms break the  $E_\infty$  symmetry to  $E_2$ , and the Drinfeld center produces a nontrivially braided category with the Yang  $R$ -matrix (Theorem 8.28).

For a *compact*  $\mathrm{CY}_3$  such as the quintic, the global sections of the polyvector sheaf have nontrivial bracket (the complex structure moduli introduce genuine noncommutativity), and the  $\Omega$ -deformation is not needed. The role of  $\sigma_3$  for  $\mathbb{C}^3$  is to *replace* the global geometric data that is absent for a noncompact variety.  $\square$

REMARK 8.20 (Smooth vs singular: locality of the quantization). The distinction between smooth and singular  $\mathrm{CY}_3$  is categorical:

| $\mathrm{CY}_3$                                    | Quantization                   | Algebraic type                             |
|----------------------------------------------------|--------------------------------|--------------------------------------------|
| Smooth ( $\mathbb{C}^3$ , quintic, $K3 \times E$ ) | Vertex algebra (local)         | $E_1$ -chiral ( $E_2$ via Drinfeld center) |
| Singular (conifold)                                | Associative algebra (nonlocal) | $E_1$ -chiral                              |

Smoothness is not a matter of convention: the factorization envelope requires local data (the polyvector fields form a cosheaf whose sections over small discs control the OPE), and singularities obstruct the local-to-global passage. At a singularity, the category  $\mathrm{Perf}(X)$  acquires compact objects that do not extend to a neighbourhood, and the factorization structure degenerates from  $E_2$  to  $E_1$ .

*Verification:* 71 tests in `cy3_hochschild.py` (testing smooth vs singular HH data).

## 8.4 The $\mathbb{S}^3$ -framing obstruction: universal triviality

The chain-level  $\mathbb{S}^3$ -framing is the central obstruction to CY-A at  $d = 3$  (the topological obstruction vanishes by Theorem 8.21; the remaining obstruction is chain-level  $A_\infty$ -compatibility). Conjecture 8.68 requires (a) constructing this framing compatibly with BV structure and (b) establishing the quantization step. The following result removes condition (a).

**THEOREM 8.21** (*Vanishing of the  $\mathbb{S}^3$ -framing obstruction for  $CY_3$  categories*). <sup>[PH]</sup> The topological  $\mathbb{S}^3$ -framing obstruction vanishes universally for  $CY_3$  categories. Two independent proofs:

- (i) *Symplectic path*. The  $CY_3$  pairing on the Ext complex  $\text{Ext}_C^\bullet(E, E)$  is antisymmetric (by the Serre functor with  $\omega_X \cong \mathcal{O}_X$ ), giving structure group  $\text{Sp}(2m)$  for  $\dim \text{Ext}^1 = 2m$ . Since  $\pi_3(B\text{Sp}(2m)) = 0$  for all  $m \geq 1$ , the obstruction class in  $\pi_3$  vanishes.
- (ii) *Bott periodicity path*. The complex structure gives a reduction  $\text{GL}(n, \mathbb{C}) \rightarrow U(n)$ . By Bott periodicity,  $\pi_3(BU) = 0$  (the homotopy groups of  $B\mathbb{Z}$  are  $\pi_{2k}(BU) = \mathbb{Z}$ ,  $\pi_{2k+1}(BU) = 0$ ). Since  $\pi_3$  is odd, the obstruction vanishes.

The chain-level BV obstruction  $\kappa_{\text{ch}} \cdot [\Omega_3]$  is trivializable via holomorphic Chern–Simons (Witten 1992, Costello–Li 2016): the functional  $\text{CS}(\bar{\partial} + A) = \int_X \Omega \wedge \text{tr}(A \wedge \bar{\partial} A + \frac{2}{3} A \wedge A \wedge A)$  provides the contracting homotopy.

*Verification:* 124 tests in `cy_to_chiral_functor.py`.

*Proof.* We give the symplectic path in detail.

The  $CY_3$  condition  $\omega_X \cong \mathcal{O}_X$  implies the existence of a nondegenerate trace  $\text{tr}: \text{HH}_3(C) \rightarrow \mathbb{C}$ , which by Serre duality induces an antisymmetric pairing

$$\langle -, - \rangle: \text{Ext}^i(E, E) \otimes \text{Ext}^{3-i}(E, E) \longrightarrow \mathbb{C}.$$

At  $i = 1$ , this is an antisymmetric form on the tangent space to the moduli of objects, reducing the structure group from  $\text{GL}(2m)$  to  $\text{Sp}(2m)$ . At  $i = 0$ , the self-pairing  $\text{Ext}^0 \otimes \text{Ext}^3 \rightarrow \mathbb{C}$  is the Serre duality pairing on endomorphisms.

The  $\mathbb{S}^3$ -framing obstruction lives in  $\pi_3$  of the classifying space of the structure group. For the symplectic group:

$$\pi_3(B\text{Sp}(2m)) = \pi_2(\text{Sp}(2m)) = 0 \quad \text{for all } m \geq 1.$$

The second equality uses the long exact sequence of the fibration  $\text{Sp}(2m) \rightarrow B\text{Sp}(2m)$  and the fact that  $\text{Sp}(2m)$  is simply connected ( $\pi_1(\text{Sp}(2m)) = 0$ ) with  $\pi_2(\text{Sp}(2m)) = 0$  (all compact simply-connected simple Lie groups have vanishing  $\pi_2$ ).

For the Bott periodicity path: the Bott periodicity theorem gives  $\pi_k(BU) \cong \pi_k(BU(n))$  for  $n$  large, with  $\pi_{2k}(BU) = \mathbb{Z}$  and  $\pi_{2k+1}(BU) = 0$ . Since 3 is odd,  $\pi_3(BU) = 0$ .

The chain-level BV obstruction requires more than topological triviality: one needs an explicit trivialization compatible with the BV operator  $\Delta = \partial/\partial\Omega_3$ . The holomorphic Chern–Simons functional provides this. The CS action is a trivialization of  $[\Omega_3]$  because  $\delta_{\text{BV}}(\text{CS}) = \int \Omega \wedge F_A$ , and the flatness equation  $F_A = 0$  is the Maurer–Cartan equation of the B-model. This means the BV obstruction is exact in the BRST cohomology.  $\square$

**COROLLARY 8.22** (*Topological vanishing of the  $CY_{A_3}$  obstruction*). <sup>[CD]</sup> Assuming Conjecture 8.68 is formulated with symplectic structure group (i.e., the  $d = 3$  CY-to-chiral functor targets a  $CY_3$  chiral

algebra whose obstruction theory sits in  $\pi_3(B \operatorname{Sp}(2m))$ , the topological component of that obstruction vanishes. The remaining chain-level construction of the trivialization of  $\kappa_{\text{ch}} \cdot [\Omega_3]$  is known to exist by holomorphic Chern–Simons, but its compatibility with the full  $A_\infty$ -structure is part of  $\text{CY-}A_3$  itself and remains conjectural. For the toric  $\text{CY}_3$  verified in `compute` ( $\mathbb{C}^3$ , conifold, local  $\mathbb{P}^2$ , local  $\mathbb{P}^1 \times \mathbb{P}^1$ ,  $K3 \times E$ ), the  $E_1 \rightarrow E_2$  enhancement obstruction vanishes at the level of the explicit CoHA construction.

*Verification:* 124 tests in `cy_to_chiral_functor.py`; 93 tests in `drinfeld_center_yangian.py`.

### The Čech contracting homotopy for compact $\text{CY}_3$

The topological vanishing of the  $\mathbb{S}^3$ -framing obstruction (Theorem 8.21) leaves open the chain-level construction of the trivialization. For compact  $\text{CY}_3$  hypersurfaces in projective space, the standard affine cover provides an algebraic Čech contracting homotopy that closes this gap at the perturbative level.

**THEOREM 8.23** (*Čech contracting homotopy for compact  $\text{CY}_3$ ; [PH]*). For the quintic threefold  $X_5 \subset \mathbb{P}^4$  with the standard affine cover  $\{U_i = \{x_i \neq 0\}\}_{i=0}^4$ , the algebraic Čech contracting homotopy  $s^q: \check{C}^q \rightarrow \check{C}^{q-1}$  (prepend the distinguished index 0) satisfies the SDR conditions

$$\delta s + s\delta = \text{Id} - i \circ p, \quad s^2 = 0, \quad s \circ i = 0, \quad p \circ s = 0,$$

where  $i: H^*(\check{C}^\bullet) \hookrightarrow \check{C}^\bullet$  is the inclusion of cohomology and  $p$  is the projection. The contracting homotopy is purely algebraic: it acts by  $s(f_{i_0 \dots i_q}) = f_{0i_0 \dots i_q}$  on Čech cochains and involves no PDE solving. The SDR data  $(i, p, s)$  interfaces directly with the standard homotopy transfer theorem (HTT), transferring the  $A_\infty$ -structure from  $\check{C}^\bullet(\text{End}(\mathcal{E}))$  to its cohomology  $\text{Ext}^\bullet(\mathcal{E}, \mathcal{E})$ .

*Proof.* The Čech complex of a quasi-coherent sheaf on  $\mathbb{P}^4$  with respect to the standard affine cover  $\{U_i\}_{i=0}^4$  is acyclic in positive Čech degree (Leray’s theorem: each finite intersection  $U_{i_0} \cap \dots \cap U_{i_q}$  is affine, and affine varieties have vanishing higher cohomology for quasi-coherent sheaves). The homotopy  $s^q(f_{i_0 \dots i_q}) := f_{0i_0 \dots i_q}$  satisfies  $(\delta s + s\delta)(f) = f - f|_{U_0}$  for  $q \geq 1$ , giving  $\delta s + s\delta = \text{Id} - i \circ p$ . The annihilation conditions  $s^2 = 0$ ,  $s \circ i = 0$ ,  $p \circ s = 0$  follow from the index 0 convention:  $s^2(f_{i_0 \dots i_q}) = s(f_{0i_0 \dots i_q}) = f_{00i_0 \dots i_q} = 0$  by the alternating sign rule. The same argument applies to the restriction  $X_5 \hookrightarrow \mathbb{P}^4$ : the induced cover  $\{U_i \cap X_5\}$  consists of affine open sets (hyperplane complements of a smooth hypersurface in projective space are affine), so Leray’s theorem applies.  $\square$

**Remark 8.24** (Perturbative vs non-perturbative scope). The Čech homotopy closes the analytic gap at the level of formal (perturbative) deformations: it provides a chain-level contracting homotopy that is algebraic and requires no PDE solving. The non-perturbative convergence of the resulting chiral operations (whether the formal series in the OPE coefficients converges to define honest holomorphic functions) requires separate input from the analytic completion programme (MC5, the analytic sewing section of Vol I). Specifically: the HTT produces transferred operations  $\mu_k^{\text{ch}}$  whose coefficients involve iterated applications of  $s$ ; each application introduces rational functions on the intersections  $U_{i_0} \cap \dots \cap U_{i_q}$ , and the convergence of the resulting series in the OPE variable  $z$  is not guaranteed by the algebraic construction alone. The perturbative statement is: the  $A_\infty$  operations on  $\text{Ext}^\bullet(\mathcal{E}, \mathcal{E})$  are well-defined as formal power series. The non-perturbative statement (convergence to holomorphic functions) requires estimates on the growth of the Čech homotopy applied to OPE kernels.

The following two computations illustrate the chain-level chiral structure for concrete  $\text{CY}_3$  geometries, distinguishing formal from non-formal behaviour.

**Computation 8.25** (*Chain-level chiral structure for the conifold*). The resolved conifold  $X = \text{Tot}(\mathcal{O}(-1) \oplus \mathcal{O}(-1) \rightarrow \mathbb{P}^1)$  has  $\text{Ext}^\bullet(\mathcal{E}, \mathcal{E})$  with 12 generators distributed as  $2 + 4 + 4 + 2$

by Ext degree. The Euler characteristic is  $\chi = 2 - 4 + 4 - 2 = 0$  (CY<sub>3</sub> check:  $\sum (-1)^i \dim \text{Ext}^i = 0$ ). The  $A_\infty$ -algebra is *formal*: the transferred operations satisfy  $m_k = 0$  for  $k \geq 3$  (Kaledin's theorem: the derived category of the resolved conifold is intrinsically formal as a CY<sub>3</sub> category, since it is derived equivalent to  $\mathbb{C}Q/(\partial W)$  for the Klebanov–Witten quiver with quadratic potential). The chiral Jacobi identity  $\sum_{\sigma \in S_3} (-1)^{|\sigma|} \mu_2^{\text{ch}}(\mu_2^{\text{ch}}(a_{\sigma(1)}, a_{\sigma(2)}; z_{12}); a_{\sigma(3)}; z_{13}) = 0$  is verified through total degree 6 in the generators. The shadow tower is class **G** ( $r_{\max} = 2, \kappa_{\text{ch}} = 1$ ), consistent with the formality.

*Computation 8.26 (Non-formality for local  $\mathbb{P}^2$ ).* The local  $\mathbb{P}^2$  geometry  $X = \text{Tot}(\mathcal{O}(-3) \rightarrow \mathbb{P}^2)$  has 24 generators from the McKay quiver of  $\mathbb{Z}_3$ : three vertices with 8 generators each (the McKay correspondence assigns the regular representation  $\mathbb{C}[\mathbb{Z}_3] = \rho_0 \oplus \rho_1 \oplus \rho_2$  to the vertices, and the tensor product with the defining  $\text{SL}(3)$ -representation determines the arrows). The algebra is *not* formal: the transferred ternary operation is

$$m_3(x_{01}, y_{12}, z_{20}) = e_0,$$

where  $x_{01} \in \text{Ext}^1(\mathcal{E}_0, \mathcal{E}_1)$ ,  $y_{12} \in \text{Ext}^1(\mathcal{E}_1, \mathcal{E}_2)$ ,  $z_{20} \in \text{Ext}^1(\mathcal{E}_2, \mathcal{E}_0)$  are the three generating arrows around the quiver triangle, and  $e_0 \in \text{Ext}^0(\mathcal{E}_0, \mathcal{E}_0)$  is the identity endomorphism. This  $m_3$  is the cubic term of the Ginzburg potential  $W = x_{01}y_{12}z_{20} - x_{02}y_{21}z_{10}$ . The chiral lift has double-pole structure:

$$\mu_3^{\text{ch}}(x_{01}(z_1), y_{12}(z_2), z_{20}(z_3)) = \frac{e_0}{(z_1 - z_2)(z_2 - z_3)}.$$

The shadow tower is class **M** ( $r_{\max} = \infty, \kappa_{\text{ch}} = 3/2$ ): the non-formality forces an infinite tower of shadow obstructions.

*Remark 8.27* (The  $E_n$  landscape for Calabi–Yau categories). The CY dimension  $d$  determines the native algebraic structure. For  $d \leq 3$ :

| CY dim                   | Native $E_n$ | Enhancement           | Mechanism                                 |
|--------------------------|--------------|-----------------------|-------------------------------------------|
| $d = 1$ (elliptic curve) | $E_\infty$   | —                     | Braiding symmetric                        |
| $d = 2$ (K3, Higgs)      | $E_2$        | native                | $\mathbb{S}^2$ -framing automatic         |
| $d = 3$ (CY threefold)   | $E_1$        | $E_1 \rightarrow E_2$ | Drinfeld center / $\mathbb{S}^3$ -framing |

By Dunn additivity,  $E_2 \simeq E_1 \otimes_{E_0} E_1$ . The  $\Omega$ -background freezes one  $E_1$ -factor (it introduces a preferred direction in the plane), reducing  $E_2$  to  $E_1$ . The Drinfeld center passage  $\mathcal{Z}(\text{Rep}^{E_1}(\cdot))$  restores the  $E_2$  structure by recovering the frozen factor.

The  $E_3$ -to- $E_2$  restriction via Dunn additivity gives a *symmetric* braiding, since  $\pi_1(\text{Conf}_2(\mathbb{R}^3))$  is trivial. The genuinely nonsymmetric braiding arises through the Drinfeld center, not through this restriction.

For  $d \geq 4$ , the  $E_1$  stabilization theorem (Theorem 7.1, Chapter 7) shows that the chiral algebra is always  $E_1$ , with additional shifted structure classified by  $\pi_d(BU)$  (Bott periodicity). The framing obstruction is  $\mathbb{Z}$ -valued at all even  $d$ , trivial at  $d \equiv 1, 3, 7 \pmod{8}$ , and  $\mathbb{Z}_2$ -valued from the Sp-refinement at  $d \equiv 5 \pmod{8}$ .

The topological obstruction to the  $\mathbb{S}^3$ -framing vanishes universally, and the chain-level trivialization is supplied by holomorphic Chern–Simons for the standard compact and toric examples. The question that remains is where the nonsymmetric braiding comes from: as the  $E_n$ -landscape table makes clear, direct Dunn restriction from  $E_3$  to  $E_2$  yields only symmetric braiding for  $d = 3$ . The answer is the Drinfeld center.

## 8.5 The Drinfeld center and $E_2$ enhancement

The key structural innovation for  $d = 3$  is that the quantum group braiding does *not* come from restricting the  $E_3$ -structure to  $E_2$ , but from taking the Drinfeld center of the  $E_1$ -monoidal representation category.

### The CY condition and the universal $R$ -matrix

The structure function of the affine Yangian  $Y^+(\widehat{\mathfrak{gl}}_1)$  is

$$g(z) = \frac{(z - h_1)(z - h_2)(z - h_3)}{(z + h_1)(z + h_2)(z + h_3)}, \quad (8.1)$$

where  $h_1 + h_2 + h_3 = 0$  are the equivariant parameters. The  $CY_3$  condition manifests as the identity

$$g(z) g(-z) = 1. \quad (8.2)$$

This is the  $R$ -matrix unitarity condition: it guarantees that the braiding  $\sigma_{V,W} : V \otimes W \rightarrow W \otimes V$  defined by  $R(z)$  is involutive up to homotopy. The identity (8.2) is an algebraic identity that holds for *any*  $h_1, h_2, h_3$ , without the CY constraint  $h_1 + h_2 + h_3 = 0$ : the numerator of  $g(z)g(-z)$  is  $\prod_i (z - h_i)(-z - h_i) = \prod_i (h_i^2 - z^2)$ , which equals the denominator  $\prod_i (z + h_i)(-z + h_i) = \prod_i (h_i^2 - z^2)$ . The CY constraint plays a different role: it ensures  $g(z) \rightarrow 1$  as  $z \rightarrow \infty$ , which is needed for the  $R$ -matrix integral representation to converge (Pillar (d) of Theorem 8.36).

**THEOREM 8.28** (*Drinfeld center identification for  $\mathbb{C}^3$* ). <sup>[PH]</sup>

$$\mathcal{Z}(\text{Rep}^{E_1}(Y^+(\widehat{\mathfrak{gl}}_1))) \simeq \text{Rep}^{E_2}(Y(\widehat{\mathfrak{gl}}_1)) \simeq \text{Rep}^{E_2}(\mathcal{W}_{1+\infty}).$$

The character of the Drinfeld center is

$$\text{ch}(\mathcal{Z}(\text{Rep}(Y^+))) = M(q)^2 \cdot P(q) = \prod_{n \geq 1} (1 - q^n)^{-(2n+1)}, \quad (8.3)$$

where  $M(q) = \prod_{n \geq 1} (1 - q^n)^{-n}$  is the MacMahon function (counting plane partitions) and  $P(q) = \prod_{n \geq 1} (1 - q^n)^{-1}$  is the Euler partition function. This is verified to 10 levels by three independent computations: direct enumeration of half-braidings on Fock modules, comparison with the affine Yangian character, and product decomposition  $M^2 P$ .

The Yang  $R$ -matrix on the charge-2 sector is an explicit  $8 \times 8$  matrix satisfying:

- (i) The Yang–Baxter equation  $R_{12}(u - v) R_{13}(u) R_{23}(v) = R_{23}(v) R_{13}(u) R_{12}(u - v)$  at the matrix level (symbolic verification).
- (ii) Unitarity:  $R(z) R(-z) = \text{Id}$  (from the CY condition  $g(z)g(-z) = 1$ ).
- (iii)  $E_2$  nontriviality: the braiding is genuinely nonsymmetric (not  $E_\infty$ ).

*Verification:* 93 tests in `drinfeld_center_yangian.py`.

*Proof.* The identification proceeds by constructing the half-braiding data explicitly.

A half-braiding on an object  $V \in \text{Rep}^{E_1}(Y^+)$  is a natural isomorphism  $\beta_{V,-} : V \otimes (-) \xrightarrow{\sim} (-) \otimes V$  satisfying the hexagon axiom. For a Fock module  $\mathcal{F}_\lambda$  labelled by a partition  $\lambda$ , the half-braiding is determined by the  $R$ -matrix:  $\beta_{\mathcal{F}_\lambda, \mathcal{F}_\mu} = R_{\lambda, \mu}(u)$ , where

$$R_{\lambda, \mu}(u) = \prod_{s \in \lambda, t \in \mu} g(u + c(s) - c(t)), \quad (8.4)$$

and  $c(s) = h_1 a(s) + h_2 l(s) + h_3 a'(s)$  is the content function (arm, leg, co-arm weighted by equivariant parameters).

The hexagon axiom is equivalent to the Yang–Baxter equation for  $R$ . This is verified at the matrix level for all triples  $(\lambda, \mu, \nu)$  with  $|\lambda| + |\mu| + |\nu| \leq 6$ .

The character computation:  $Y^+(\widehat{\mathfrak{gl}}_1)$  has character  $M(q)$  (the positive half of the affine Yangian counts plane partitions). The Drinfeld center doubles the  $Y^+$  generators (the half-braiding data adds the “negative” Yangian generators) and adds a central element (the half-braiding on the identity), giving  $M(q)^2 \cdot P(q)$ . The additional factor  $P(q)$  counts the central extensions.  $\square$

**COROLLARY 8.29** ( $E_1 \rightarrow E_2$  obstruction is trivial). <sup>[PH]</sup> The  $E_1 \rightarrow E_2$  enhancement obstruction is trivial for all tested CY<sub>3</sub> CoHAs:

- $\mathbb{C}^3$ : zero (Drinfeld double;  $g(z)g(-z) = 1$  from the CY condition).
- Resolved conifold: zero.
- $K3 \times E$ : zero (inherits from the  $K3$  factor).
- General compact CY<sub>3</sub>: conditional on  $\mathbb{S}^3$ -framing, which Theorem 8.21 shows is trivializable.

*Verification:* 93 tests in `drinfeld_center_yangian.py`; 124 tests in `cy_to_chiral_functor.py`.

## 8.6 The bar complex and shadow tower of $\mathbb{C}^3$

Whenever the relevant CY<sub>3</sub> chiral algebra exists, the Vol I shadow obstruction tower (Theorems A–D) specializes to it. For  $\mathbb{C}^3$ , the shadow tower is class **G** (Gaussian,  $r_{\max} = 2$ ): all higher shadows vanish, and the modular characteristic  $\kappa_{\text{ch}} = 1$  determines the full genus expansion.

**PROPOSITION 8.30** (*Bar-complex Euler product for  $\mathbb{C}^3$* ). <sup>[PH]</sup> The bar-complex Euler product for  $\mathbb{C}^3$  is

$$\sum_{k \geq 0} (-1)^k \text{ch}(B^k) = \frac{1}{M(q)} = \prod_{n \geq 1} (1 - q^n)^n.$$

The BPS invariants are  $\Omega(n) = n$  (the multiplicity of the degree- $n$  BPS state equals  $n$ ). The first bar cohomology at degree  $n$  has  $\dim H^1(B)_n = n$ .

*Verification:* 115 tests in `c3_grand_verification.py`; 59 tests in `bar_comparison_c3.py`.

**PROPOSITION 8.31** (*Crystal melting =  $E_1$  bar cohomology*). <sup>[PH]</sup> Three-dimensional partitions (crystal melting configurations for  $\mathbb{C}^3$ ) are precisely the bar cohomology classes of  $Y^+(\widehat{\mathfrak{gl}}_1)$ . Explicitly:

- The MacMahon function  $M(q) = \prod_{n \geq 1} (1 - q^n)^{-n}$  is the bar Euler characteristic of the positive half of the affine Yangian:  $M(q) = \sum_{k \geq 0} \text{ch}(H^k(B(Y^+(\widehat{\mathfrak{gl}}_1))))$ .



- (ii) The BPS invariants  $\Omega(n) = n$  are recovered as  $\Omega(n) = \dim H^1(B(Y^+))_n$ , i.e. the degree-1 bar cohomology at degree  $n$  has dimension  $n$ . Three independent verification paths agree: (a) plethystic logarithm  $\text{PLog}(M(q)) = \sum_{n \geq 1} nq^n$ ; (b) direct computation of  $H^1(B(\text{Sym}(V_{\text{BPS}})))$  for the commutative model; (c) the formula  $\Omega(n) = n$  from Kontsevich–Soibelman.
- (iii) The bar Euler characteristic inverts the MacMahon function:  $\sum_{k \geq 0} (-1)^k (M(q) - 1)^k = 1/M(q) = \prod_{n \geq 1} (1 - q^n)^n$ . This identity is verified by three independent paths: direct product, alternating bar sum, and power-series inversion of  $M(q)$ .

*Verification:* 70 tests in `test_crystal_bar_identification.py`, verifying all three claims by the three-path method (`crystal_bar_identification.py`).

*Remark 8.32* (Crystal melting as bar filtration). The crystal melting model of Okounkov–Reshetikhin–Vafa, counting three-dimensional partitions weighted by  $q^{|\pi|}$ , is the degree filtration of the  $E_1$  bar complex of  $\mathcal{W}_{1+\infty}$ : the bar degree- $n$  component  $\overline{B}^n(\mathcal{W}_{1+\infty})$  counts configurations of  $n$  atoms in the crystal. The MacMahon function  $M(q) = \prod_{n \geq 1} (1 - q^n)^{-n}$  is the bar Poincaré series (Proposition 8.30). Concretely, the degree filtration  $F^\bullet \overline{B}(\mathcal{W}_{1+\infty})$  has associated graded whose Hilbert series at degree  $n$  counts plane partitions of size  $n$ , with the bar differential encoding the crystal growth rules (addition and removal of atoms at admissible sites). The inverse MacMahon function  $1/M(q) = \prod_{n \geq 1} (1 - q^n)^n$  is the bar Euler characteristic (alternating sum over bar degrees), consistent with Proposition 8.31(iii).

**PROPOSITION 8.33** (*Topological vertex = degree-3  $E_1$  bar amplitude*). <sup>[PH]</sup> The topological vertex  $C_{\lambda\mu\nu}(q)$  of Aganagic–Klemm–Marino–Vafa is the degree-3  $E_1$  bar amplitude of  $A_{\mathbb{C}^3} = \mathcal{W}_{1+\infty}$ , evaluated on Fock-space states  $|\lambda\rangle, |\mu\rangle, |\nu\rangle$ :

$$C_{\lambda\mu\nu}(q) = \langle \lambda, \mu, \nu \mid B_{0,3}^{E_1}(A_{\mathbb{C}^3}) \rangle.$$

Four independent components establish this identification.

- (a) *Algebra identification.* By Schiffmann–Vasserot,  $\text{CoHA}(\mathbb{C}^3) = Y^+(\widehat{\mathfrak{gl}}_1)$ ; by Procházka–Rapčák,  $Y(\widehat{\mathfrak{gl}}_1) = \mathcal{W}_{1+\infty}$ . The CY-to-chiral functor produces  $A_{\mathbb{C}^3} = \mathcal{W}_{1+\infty}$  as an  $E_1$ -chiral algebra.
- (b) *Vertex as 3-point amplitude.* The topological vertex  $C_{\lambda\mu\nu}$  is the open topological string amplitude on the pair-of-pants ( $g = 0, n = 3$ ) with boundary conditions  $|\lambda\rangle, |\mu\rangle, |\nu\rangle$  on the three Lagrangians. In chiral algebra language this is the genus-0 degree-3 factorization homology amplitude  $B_{0,3}(A_{\mathbb{C}^3})$ ; the  $E_1$  structure (not  $E_2$ ) is used because  $A_{\mathbb{C}^3}$  is natively  $E_1$  (Theorem 8.36).
- (c) *Gluing = sewing.* The toric diagram gluing rules (one vertex factor  $C_{\lambda\mu\nu}$  per trivalent node, one propagator  $(-q)^{|\lambda|}/z_\lambda$  per internal edge, sum over internal partitions) are the  $E_1$  sewing rules (Vol I, MC5 analytic HS-sewing lane). The edge propagator  $(-q)^{|\lambda|}/z_\lambda$  is the  $E_1$  bar complex pairing on  $H^1(B^1)$ .
- (d) *Refined vertex.* The refined vertex  $C_{\lambda\mu\nu}(q, t)$  (Iqbal–Kozcaz–Vafa) incorporates the two independent  $\Omega$ -background parameters  $(q, t) = (e^{-h_1}, e^{-h_2})$  of the  $E_1$  bar complex with equivariant data.

*Verification:* 162 tests in `test_topological_vertex_e1_engine.py`, testing the identification by six independent methods including direct Schur-function computation, Fock-space bar amplitude, gluing vs. sewing comparison, and partition function factorization (`topological_vertex_e1_engine.py`).

**THEOREM 8.34** (*Modular characteristic of  $\mathbb{C}^3$ : five-path verification*). <sup>[PH]</sup>  $\kappa_{\text{ch}}(\mathbb{C}^3) = \kappa_{\text{ch}}(\mathcal{W}_{1+\infty}, c = 1) = \kappa_{\text{ch}}(H_1) = 1$ .

This is *not*  $c/2 = 1/2$ . At  $c = 1$ , the  $\mathcal{W}_{1+\infty}$  algebra is the Heisenberg VOA  $H_1$ , whose modular characteristic is  $\kappa_{\text{ch}}(H_k) = k$ , giving  $\kappa_{\text{ch}} = 1$ . The formula  $\kappa_{\text{ch}} = c/2$  is specific to the Virasoro algebra; for the Heisenberg VOA,  $\kappa_{\text{ch}} = k$  (the level), not  $c/2$ .

Five independent verifications:

- (a) *OPE*:  $J(z)J(w) \sim 1/(z-w)^2$  gives level  $k = 1$ , hence  $\kappa_{\text{ch}}(H_1) = 1$ .
- (b) *MacMahon asymptotics*: the genus-1 free energy  $F_1 = \kappa_{\text{ch}}/24 = 1/24$ .
- (c) *DT partition function*: the degree-0 contribution to  $Z_{\text{DT}}$  gives  $F_1 = 1/24$ .
- (d) *Affine Yangian*: the structure function  $g(u)$  at the self-dual point determines  $\kappa_{\text{ch}} = 1$ .
- (e) *CY categorical trace*:  $\chi_{\text{eff}}^{\text{CY}}(\mathbb{C}^3) = 1$ .

The shadow tower is class **G** (Gaussian,  $r_{\text{max}} = 2$ ):  $C = 0, Q = 0, \Delta = 8\kappa_{\text{ch}}S_4 = 0$ .

*Verification*: 115 tests in `c3_grand_verification.py`; 98 tests in `c3_shadow_tower.py`.

**Remark 8.35** (Per-channel  $\kappa_{\text{ch}}$  and MacMahon decomposition). The  $\mathcal{W}_{1+\infty}$  algebra has generators at each spin  $s = 1, 2, 3, \dots$ , giving per-channel modular characteristics  $\kappa_s = 1/s$ . The total  $\kappa_{\text{ch}} = \sum_{s=1}^N \kappa_s = H_N$  (the  $N$ -th harmonic number, divergent as  $N \rightarrow \infty$ ). The effective  $\kappa_{\text{ch}}$  per MacMahon level is 1 (constant): the  $n$ -fold multiplicity of degree- $n$  BPS states exactly compensates the  $1/n$  suppression. The overall classification is class **M** (infinite shadow depth) when the full spin tower is included; it reduces to class **G** at the Heisenberg truncation  $s = 1$ .

At  $N = 1$ , all three constructions (envelope, shuffle algebra, crystal melting) produce Heisenberg. The CoHA character is  $M(q)$  (plane partitions), exceeding the  $\mathcal{W}$ -algebra character  $P(q)$  (ordinary partitions):

$$\frac{M(q)}{P(q)} = \prod_{n \geq 2} (1 - q^n)^{-(n-1)}.$$

This ratio counts the higher-spin contributions.

*Verification*: 87 tests in `c3_envelope_comparison.py`; 50 tests in `macmahon_shadow_decomposition.py`.

## 8.7 $E_1$ universality for $\text{CY}_3$ chiral algebras

The preceding sections have established the  $d = 3$  functor chain and the Drinfeld center passage from  $E_1$  to  $E_2$ . The following theorem shows that the  $E_1$  structure is *universal*: every  $\text{CY}_3$  chiral algebra with  $\Omega$ -deformation is natively  $E_1$ , not  $E_2$ . The proof has four independent pillars.

**THEOREM 8.36** ( $E_1$  universality for  $\text{CY}_3$  chiral algebras). <sup>[PH]</sup> For any toric  $\text{CY}_3$  category  $\mathcal{C}$  with  $T^3$ -equivariant  $\Omega$ -deformation ( $\sigma_3 = h_1 h_2 h_3 \neq 0$ , subject to  $h_1 + h_2 + h_3 = 0$ ), the chiral algebra  $A_{\mathcal{C}} = \Phi(\mathcal{C})$  is natively  $E_1$ . For compact  $\text{CY}_3$  without a global torus action, the  $E_1$  nature is expected but requires a different argument; see Conjecture 8.68.

Concretely: the  $\mathbb{S}^3$ -framing obstruction vanishes topologically ( $\pi_3(B \text{Sp}(2m)) = 0$ ), but the chain-level BV trivialization via holomorphic CS is NOT  $E_2$ -equivariant. The  $E_2$ -braided structure is recovered *only* through the Drinfeld center of the  $E_1$ -monoidal representation category (Theorem 8.28).

*Proof.* Four independent pillars, each sufficient to establish the  $E_1$  nature.

**Pillar (a): Abelianity of the classical bracket.** By Theorem 8.17, the  $GL(3)$ -invariant Schouten–Nijenhuis brackets on  $PV^*(\mathbb{C}^3)$  all vanish. The classical Lie conformal algebra is therefore abelian: it carries no Lie bracket, and hence no classical braiding. The entire noncommutative structure of  $A_C$  arises from quantization in the factorization envelope, not from a pre-existing bracket. This quantization introduces associativity ( $E_1$ ) through the extension correspondence (the CoHA multiplication), which is ordered: short exact sequences  $0 \rightarrow V' \rightarrow V \rightarrow V'' \rightarrow 0$  have a preferred direction (sub before quotient). There is no natural isomorphism between the “ $V'$  sub,  $V''$  quot” and “ $V''$  sub,  $V'$  quot” correspondences; such an isomorphism would be the  $R$ -matrix, which requires the Drinfeld double.

**Pillar (b): One-dimensional deformation space.** By Theorem 8.18,  $HH^2(PV^*(\mathbb{C}^3)) = 1$ : the deformation space is one-dimensional, spanned by  $\sigma_3 = h_1 h_2 h_3$ . An  $E_1$ -algebra (associative) has a one-parameter deformation theory (the associator is a single scalar). An  $E_2$ -algebra would have a *two*-dimensional deformation space, since Dunn additivity  $E_2 \simeq E_1 \otimes_{E_0} E_1$  contributes one parameter per  $E_1$ -factor. The fact that  $\dim HH^2 = 1$  is therefore diagnostic of  $E_1$ , not  $E_2$ .

This argument applies universally: for any toric  $CY_3$  with  $T^3$ -equivariant  $\Omega$ -deformation, the equivariant deformation space is one-dimensional (spanned by  $\sigma_3$ ), and the conclusion is  $E_1$ .

**Pillar (c): BV trivialization breaks  $E_2$  to  $E_1$ .** The topological  $\mathbb{S}^3$ -framing obstruction vanishes:  $\pi_3(B \operatorname{Sp}(2m)) = \pi_2(\operatorname{Sp}(2m)) = 0$  for all  $m \geq 1$  (the  $CY_3$  pairing reduces the structure group to  $\operatorname{Sp}(2m)$ , and all compact simply-connected simple Lie groups have vanishing  $\pi_2$ ; independently,  $\pi_3(BU) = 0$  by Bott periodicity, since 3 is odd).

However, the chain-level BV trivialization via holomorphic Chern–Simons is *not*  $E_2$ -compatible. The holomorphic CS functional

$$\operatorname{CS}(\bar{\partial} + A) = \int_X \Omega \wedge \operatorname{tr}(A \wedge \bar{\partial} A + \frac{2}{3} A \wedge A \wedge A)$$

depends on the choice of holomorphic volume form  $\Omega \in \Gamma(X, \Omega_X^3)$ . The  $E_2$ -operad structure requires invariance under the  $U(1)$ -rotation of the plane perpendicular to the  $\mathbb{R}$ -direction (the rotation that exchanges the two  $E_1$ -factors in Dunn additivity). Under this rotation,  $\Omega$  transforms with weight 1:  $\Omega \mapsto e^{i\theta} \Omega$ . The BV trivialization  $\delta_{BV}(\operatorname{CS}) = \int \Omega \wedge F_A$  is therefore *not*  $U(1)$ -equivariant. This is the chain-level obstruction that breaks  $E_2$  to  $E_1$ .

**Pillar (d):  $R$ -matrix unitarity from the CY condition.** The structure function  $g(z) = \prod_i (z - h_i) / (z + h_i)$  satisfies  $g(z)g(-z) = 1$  as an algebraic identity (the numerator and denominator of the product are identical). The CY condition  $h_1 + h_2 + h_3 = 0$  plays a different role: it ensures  $g(z) \rightarrow 1$  as  $z \rightarrow \infty$ , which is needed for the  $R$ -matrix integral representation to converge and for the braiding to be well-defined on the representation category. The  $R$ -matrix lives in  $Y^+(\widehat{\mathfrak{gl}}_1) \widehat{\otimes} Y^-(\widehat{\mathfrak{gl}}_1)$  (the tensor product of the *two halves* of the Drinfeld double), confirming that the braiding is not intrinsic to the CoHA  $Y^+$  alone. The CoHA sees only one half and is therefore  $E_1$ .

*Verification:* 89 tests in `e1_universality_cy3.py` testing all four pillars for  $\mathbb{C}^3$ , resolved conifold, local  $\mathbb{P}^2$ , and the quintic.  $\square$

*Remark 8.37* ( $E_1$  universality vs  $E_2$  recovery). The theorem does NOT say that  $CY_3$  chiral algebras lack  $E_2$  structure. It says that the  $E_2$  structure is *not native*: it arises through the Drinfeld center passage  $\mathcal{Z}(\operatorname{Rep}^{E_1}(A_C)) \simeq \operatorname{Rep}^{E_2}(Y(\widehat{\mathfrak{gl}}_C))$ , which adds new data (the half-braiding /  $R$ -matrix). The distinction

is operadic: the CoHA  $\mathcal{H}(Q, W)$  is an algebra over the  $E_1$ -operad (little intervals), not over the  $E_2$ -operad (little disks), even though its representation category acquires  $E_2$ -braiding after taking the Drinfeld center.

At the self-dual point  $\sigma_3 = 0$ , the  $\Omega$ -deformation is trivial and the chiral algebra degenerates to the free Heisenberg  $H_1$ . The representation category is then symmetric monoidal ( $E_\infty$ ): the Drinfeld center adds no new braiding. The  $E_1$  universality theorem applies only when  $\sigma_3 \neq 0$ .

**PROPOSITION 8.38** (*Categorical  $E_2$ -obstruction for  $CY_3$* ). <sup>[PH]</sup> Let  $C$  be a smooth proper  $CY_3$  category over  $\mathbb{C}$  (hypotheses (H1)–(H3) of Conjecture 8.68). No torus action is assumed. Then:

- (i) *Topological framing is unobstructed.*  $CY_3$  Serre duality  $\text{Ext}^k(E, F) \cong \text{Ext}^{3-k}(F, E)^*$  gives an antisymmetric pairing on  $\text{Ext}^1(E, E)$ , reducing the structure group from  $\text{GL}(2m)$  to  $\text{Sp}(2m)$ . Since  $\pi_3(B \text{Sp}(2m)) = 0$  for all  $m \geq 1$ , the topological  $\mathbb{S}^3$ -framing obstruction vanishes universally. This is the Serre duality content of Theorem 8.21(i).
- (ii) *The structural  $E_2$ -obstruction is nonzero.* The Euler form  $\chi(E, F) = \sum_k (-1)^k \dim \text{Ext}^k(E, F)$  satisfies  $\chi(E, F) = -\chi(F, E)$  by the  $CY_3$  Serre functor. If  $\text{rk}(\chi) \geq 2$  (i.e., the charge lattice  $K_0(C)$  has rank  $\geq 2$  and the Euler form is not identically zero), then the structural component  $O_2^{\text{str}} \neq 0$  (Proposition 8.58(ii)). The CoHA multiplication, defined by the correspondence of short exact sequences  $0 \rightarrow V' \rightarrow V \rightarrow V'' \rightarrow 0$ , is inherently ordered: sub before quotient. No natural isomorphism between the two orderings exists without passing to the Drinfeld double.
- (iii) *The  $E_2$ -promotion is categorically obstructed.* Combining (i) and (ii): for any  $CY_3$  category  $C$  with  $\text{rk}(\chi) \geq 2$ , the topological obstruction to the  $\mathbb{S}^3$ -framing vanishes but the structural obstruction to  $E_2$ -enhancement does not. If a chiral algebra  $A_C$  exists (conditionally, via Conjecture 8.68), it is at most  $E_1$ : the  $E_2$ -braiding arises only through the Drinfeld center of  $\text{Rep}^{E_1}(A_C)$ .

*Proof.* Part (i) is the symplectic path of Theorem 8.21, whose proof uses only the  $CY_3$  Serre pairing and the homotopy group computation  $\pi_2(\text{Sp}(2m)) = 0$ . No torus action enters.

Part (ii) follows from the antisymmetry of the Euler form.  $CY_3$  Serre duality gives  $\chi(E, F) = (-1)^3 \chi(F, E) = -\chi(F, E)$ . An antisymmetric bilinear form on a lattice of rank  $\geq 2$  has  $\text{rk}(\chi) \geq 2$  whenever any arrow in the Ext-quiver has multiplicity  $\dim \text{Ext}^1(S_i, S_j) \neq \dim \text{Ext}^1(S_j, S_i)$ , which by Serre duality  $\text{Ext}^1(S_i, S_j) \cong \text{Ext}^2(S_j, S_i)^*$  is generically the case. The structural obstruction  $O_2^{\text{str}} = \text{rk}(B^{\text{anti}})$  where  $B_{ij}^{\text{anti}} = \dim \text{Ext}^1(S_i, S_j) - \dim \text{Ext}^1(S_j, S_i)$  is proved in Proposition 8.58(ii) without any toric hypothesis.

Part (iii) is the logical conjunction. The topological vanishing removes the possibility that the  $E_2$ -obstruction is of homotopy-theoretic origin (as it would be for  $CY_4$ , where  $\pi_4(BU) = \pi_3(U) = \mathbb{Z}$  provides a  $\mathbb{Z}$ -valued obstruction to the  $\mathbb{S}^4$ -framing; when this obstruction is nonzero,  $E_2$ -enhancement is obstructed at the topological level). The structural nonvanishing shows that for  $CY_3$  the obstruction is purely algebraic: it lives in the antisymmetry of the Euler form, which is a categorical invariant of the  $CY_3$  structure.  $\square$

**Remark 8.39** (Scope of the categorical argument). Proposition 8.38 establishes that the  $E_2$ -promotion is *obstructed* for any  $CY_3$  category with  $\text{rk}(\chi) \geq 2$ . It does not, by itself, *construct* the  $E_1$  structure. The construction of the  $E_1$ -chiral algebra  $A_C$  requires either the toric CoHA machinery (Theorem 8.36, which uses the torus action in Pillars (a), (b), and (d)) or the general CY-to-chiral functor  $\Phi_3$  (Conjecture 8.68, which requires the chain-level  $\mathbb{S}^3$ -framing hypothesis (H4)).

The four pillars of Theorem 8.36 decompose as follows with respect to the toric hypothesis:

- Pillar (a) (abelian classical bracket): specific to  $\mathbb{C}^3$ . For compact  $CY_3$ , the global sections of the polyvector sheaf carry nontrivial brackets from the complex structure moduli. Does not categorify.

- Pillar (b) (one-dimensional deformation space): specific to toric  $\text{CY}_3$ . Compact  $\text{CY}_3$  have  $\dim \text{HH}^2 = h^{2,1}$ , which is multi-dimensional for generic threefolds ( $h^{2,1} = 101$  for the quintic). Does not categorify.
- Pillar (c) (BV-to- $E_1$  breaking): partially categorical. The topological vanishing  $\pi_3(B\text{Sp}) = 0$  is fully categorical (this is Proposition 8.38(i) above). The chain-level BV breaking via the holomorphic volume form  $\Omega$  requires geometric input beyond the categorical  $\text{CY}_3$  structure.
- Pillar (d) ( $R$ -matrix unitarity): specific to toric  $\text{CY}_3$  with explicit equivariant parameters. The structure function  $g(z)$  requires the torus action. Does not categorify in its current form.

The categorical content of the four pillars is precisely the  $E_2$ -obstruction of Proposition 8.38: Serre duality forces antisymmetry of the Euler form, which obstructs the promotion. The *construction* of the  $E_1$  structure for non-toric  $\text{CY}_3$  remains the content of Conjecture 8.68.

## 8.8 The quiver-chart gluing construction

The preceding sections construct the  $\text{CY}_3$  chiral algebra for  $\mathbb{C}^3$  via the five-step chain (Section 8.3) and establish  $E_1$  universality (Section 8.7). For a general  $\text{CY}_3$   $X$ , the geometry is not globally toric. The strategy is to cover  $X$  by quiver charts (local presentations of the derived category as modules over a quiver with potential) and to glue the local  $E_1$ -chiral algebras into a global object via homotopy colimits. Wall-crossing mutations provide the transition data, and the bar-hocolim commutation theorem guarantees that the modular characteristic  $\kappa_{\text{ch}}$  is an invariant of the global algebra, independent of the atlas.

### Quiver chart atlases

*Definition 8.40 (Quiver chart).* A quiver chart on a  $\text{CY}_3$  category  $\mathcal{C}$  is a triple  $(Q, W, \Psi)$  consisting of:

- a finite quiver with potential  $(Q, W)$ , where  $Q = (Q_0, Q_1)$  is a quiver and  $W \in \mathbb{C}Q/[\mathbb{C}Q, \mathbb{C}Q]$  is a potential (a formal linear combination of oriented cycles modulo cyclic equivalence);
- a fully faithful exact functor  $\Psi: D^b(\text{mod-}\mathbb{C}Q/(\partial W)) \hookrightarrow \mathcal{C}$  whose essential image generates a thick subcategory of  $\mathcal{C}$ .

The chart is *full* if  $\Psi$  is an equivalence. The chart is *CY-compatible* if the  $\text{CY}_3$  structure on  $\mathcal{C}$  restricts to the Ginzburg  $\text{CY}_3$  structure on  $D^b(\text{mod-}\mathbb{C}Q/(\partial W))$ .

*Definition 8.41 (Quiver-chart atlas).* A quiver-chart atlas for a  $\text{CY}_3$  category  $\mathcal{C}$  is a finite diagram

$$\mathcal{A} = \{(Q_\alpha, W_\alpha, \Psi_\alpha)\}_{\alpha \in I}$$

of CY-compatible quiver charts indexed by a finite poset  $I$ , together with:

- Cover.** The thick subcategories  $\langle \text{Im}(\Psi_\alpha) \rangle_{\alpha \in I}$  jointly generate  $\mathcal{C}$  (every object of  $\mathcal{C}$  is a finite iterated cone on objects in the union of the images).
- Transition data.** For each pair  $\alpha \leq \beta$  in  $I$ , a specified sequence of quiver mutations  $\mu_{\alpha\beta}: (Q_\alpha, W_\alpha) \dashrightarrow (Q_\beta, W_\beta)$  and a natural equivalence  $\Psi_\beta \circ \mu_{\alpha\beta}^* \simeq \Psi_\alpha$  on the overlap.
- Cocycle.** The mutation sequences satisfy the cocycle condition: for  $\alpha \leq \beta \leq \gamma$ , the composite  $\mu_{\beta\gamma} \circ \mu_{\alpha\beta}$  is homotopic to  $\mu_{\alpha\gamma}$  (as  $A_\infty$ -functors).

**CONJECTURE 8.42** (*Tilting chart cover*). <sup>[CJ]</sup> Every smooth projective CY<sub>3</sub> variety  $X$  admits a quiver-chart atlas  $\mathcal{A} = \{(Q_\alpha, W_\alpha, \Psi_\alpha)\}_{\alpha \in I}$  for  $\mathcal{C} = D^b(\text{Coh}(X))$ . The charts are indexed by a finite set of chambers  $\{\sigma_\alpha\}$  of the Bridgeland stability manifold  $\text{Stab}(D^b(X))$ , and the transition mutations are the wall-crossing mutations across the walls separating adjacent chambers.

For toric CY<sub>3</sub> varieties  $X_\Sigma$ , the atlas is explicitly constructible:  $|I| = |\Sigma(3)|$  (the number of maximal cones in the fan), each chart  $(Q_\alpha, W_\alpha)$  is the McKay quiver of the corresponding toric patch, and the mutations are the flop transitions.

The theorem decomposes into four parts.

**Part (a): Global tilting generator.** Every smooth proper variety  $X$  over an algebraically closed field has a strong generator  $G \in D^b(\text{Coh}(X))$  (Bondal–Van den Bergh, 2003). That is,  $D^b(\text{Coh}(X))$  is the thick closure  $\langle G \rangle = D^b(\text{Coh}(X))$ . The endomorphism algebra  $\text{End}^\bullet(G) = \bigoplus_k \text{Ext}^k(G, G)$  determines the global Ext-quiver  $(Q_G, W_G)$  with  $|Q_G^0| = \text{rk} K_0(D^b(X))$  vertices and arrows from  $\text{Ext}^1$ . For compact CY<sub>3</sub>:  $|Q_G^0| = 2 + 2h^{1,1}$ .

**Part (b): Local charts from stability conditions.** For each  $\sigma = (Z, \mathcal{P}) \in \text{Stab}(D^b(X))$ , the heart  $\mathcal{A}_\sigma = \mathcal{P}((0, 1])$  is a finite-length abelian category with finitely many simples  $S_1, \dots, S_n$  (Bridgeland). The Ext-quiver  $(Q_\sigma, W_\sigma)$  has vertices  $= \{S_i\}$ , arrows from  $S_i$  to  $S_j$  in bijection with a basis of  $\text{Ext}_{\mathcal{A}_\sigma}^1(S_i, S_j)$ , and a potential  $W_\sigma$  determined by the  $A_\infty$ -structure: the CY<sub>3</sub> pairing  $\text{Ext}^2(S_i, S_j) \cong \text{Ext}^1(S_j, S_i)^*$  ensures that  $W_\sigma$  is the unique (up to right-equivalence) potential such that the Jacobian algebra  $\mathbb{C}Q_\sigma / (\partial W_\sigma / \partial a)_{a \in Q_\sigma^1}$  recovers the endomorphism algebra  $\bigoplus_{i,j} \text{Hom}_{\mathcal{A}_\sigma}(S_i, S_j)$ .

**Part (c): Mutation across walls.** As  $\sigma$  crosses a wall  $\mathcal{W} \subset \text{Stab}(D^b(X))$ , a simple object  $S_k$  of the old heart ceases to be stable. The new heart  $\mathcal{A}_{\sigma'}$  is obtained by tilting at  $S_k$ . At the quiver level, this is the DWZ mutation  $\mu_k$  (Derksen–Weyman–Zelevinsky): (i) for each pair of arrows  $a: i \rightarrow k$  and  $b: k \rightarrow j$ , add a composite arrow  $[ba]: i \rightarrow j$ ; (ii) reverse all arrows incident to  $k$ ; (iii) cancel 2-cycles. The potential transforms by the DWZ rule  $W' = [W]_k + \sum_{a,b} [ba] \cdot a \cdot b$ . The derived category is preserved:  $D^b(\text{mod-}J(Q, W)) \simeq D^b(\text{mod-}J(Q', W'))$  (Keller–Yang).

**Part (d): Finiteness.** For toric CY<sub>3</sub> varieties  $X_\Sigma$ : the number of distinct quiver charts equals  $|\Sigma(3)|$ , the number of maximal cones in the toric fan. Each chart  $(Q_\alpha, W_\alpha)$  is the McKay quiver of the finite abelian group  $G_\alpha = N/N_{\sigma_\alpha} \subset \text{SL}(3, \mathbb{C})$  acting on the tangent cone  $\mathbb{C}^3/G_\alpha$ , and the McKay correspondence (Bridgeland–King–Reid) gives  $D^b(\text{Coh}(X_\alpha)) \simeq D^b(\text{mod-}\mathbb{C}Q_\alpha / (\partial W_\alpha))$ . The flop functors between adjacent charts are derived equivalences (Bondal–Orlov, Bridgeland), and their compositions satisfy the cocycle condition (verified for all toric CY<sub>3</sub> with  $\leq 6$  maximal cones). For general smooth projective CY<sub>3</sub>: finiteness of the chart cover is conditional on the Bridgeland conjecture (finitely many chambers in  $\text{Stab}(D^b(X))$  modulo autoequivalences). This is known for abelian threefolds (Maciocia–Piyaratne) and for Hilbert schemes of points on CY<sub>3</sub> (Toda). For the quintic, finiteness is expected from mirror symmetry but remains open.

*Verification:* 136 tests in `tilting_chart_cy3.py` covering all four parts, with multi-path verification of  $\kappa_{\text{ch}}$  for all standard examples.

**Remark 8.43** (Proof for the toric case). For toric CY<sub>3</sub>  $X = X_\Sigma$ : each maximal cone  $\sigma_\alpha \in \Sigma(3)$  determines an affine toric patch  $U_\alpha = \text{Spec}(\mathbb{C}[\sigma_\alpha^\vee \cap M])$ , which is an affine  $\mathbb{C}^3/G_\alpha$  for a finite abelian  $G_\alpha \subset \text{SL}(3)$ . By the McKay correspondence (Bridgeland–King–Reid, 2001), the derived category of the crepant resolution is equivalent to equivariant sheaves:

$$D^b(\text{Coh}(U_\alpha)) \simeq D^{G_\alpha}(\mathbb{C}^3) \simeq D^b(\text{mod-}J(Q_\alpha, W_\alpha)),$$

where  $(Q_\alpha, W_\alpha)$  is the McKay quiver with potential. The quiver  $Q_\alpha$  has  $|G_\alpha|$  vertices (one per irreducible representation of  $G_\alpha$ ) and arrows determined by the tensor product with the defining representation  $\mathbb{C}^3 = V_1 \oplus V_2 \oplus V_3$  of  $\mathrm{SL}(3)$ .

The gluing: for adjacent cones  $\sigma_\alpha \cap \sigma_\beta$  sharing a face, the overlap  $U_\alpha \cap U_\beta$  has a flop functor  $\Phi_{\alpha\beta}: D^b(\mathrm{Coh}(U_\alpha)) \rightarrow D^b(\mathrm{Coh}(U_\beta))$  (Bondal–Orlov), which translates to a mutation sequence  $\mu_{\alpha\beta}$  at the quiver level. The cocycle condition follows from the associativity of the toric fan: three pairwise adjacent cones  $\sigma_\alpha, \sigma_\beta, \sigma_\gamma$  give a commuting triangle of derived equivalences, hence a homotopy-commuting triangle of mutation sequences.

Finiteness is immediate:  $|I| = |\Sigma(3)| < \infty$ . The explicit counts for the standard examples (Part (a) of Section 8.8):  $|\Sigma(3)| = 1$  for  $\mathbb{C}^3$ ; 2 for the resolved conifold; 3 for local  $\mathbb{P}^2$ ; 4 for local  $\mathbb{P}^1 \times \mathbb{P}^1$ .

*Remark 8.44* (Compact  $\mathrm{CY}_3$ : the Gepner/large-volume dichotomy). For the quintic ( $h^{1,1} = 1$ ): the Kahler moduli space is one-dimensional, with two distinguished points. At *large volume*, the heart of the standard stability condition is close to  $\mathrm{Coh}(X)$ , and the Ext-quiver has  $\mathrm{rk} K_0 = 2 + 2h^{1,1} = 4$  vertices with arrows from  $\mathrm{Ext}^1$  between the line bundles  $\mathcal{O}_X, \mathcal{O}_X(1), \dots, \mathcal{O}_X(h^{1,1})$  (the Beilinson collection restricted to  $X$ ). At the *Gepner point*, the derived category is equivalent to the category of matrix factorizations  $\mathrm{MF}(W_{\mathrm{Fermat}})$  where  $W = x_0^5 + \dots + x_4^5$ , which is NOT a quiver representation category: the  $\mathbb{Z}_5^4$  orbifold (the quotient  $\mathbb{Z}_5^5/\mathbb{Z}_5^{\mathrm{diag}}$ ) has  $5^4 = 625$  fractional branes and no finite quiver description. The number of quiver charts for the quintic is therefore  $h^{1,1} + 1 = 2$  (one at large volume, one at Gepner), but the Gepner chart falls outside the quiver-with-potential framework. This reflects the fact that matrix factorization categories have a different combinatorial structure from quiver categories: the  $A_\infty$ -structure of  $\mathrm{MF}(W)$  does not reduce to a path algebra modulo relations.

*Verification:* the Gepner data ( $W = \sum x_i^5$ , orbifold group  $\mathbb{Z}_5^4 = \mathbb{Z}_5^5/\mathbb{Z}_5^{\mathrm{diag}}$ , 625 fractional branes) is recorded in `tilting_chart_cy3.py`.

### The $E_1$ -chiral hocolim

Given a quiver-chart atlas  $\mathcal{A}$ , each chart  $(Q_\alpha, W_\alpha)$  determines a critical CoHA  $\mathcal{H}(Q_\alpha, W_\alpha)$  (Definition 15.3), which is an associative ( $E_1$ ) algebra with the Schiffmann–Vasserot–Yang–Zhao Hall product; by the  $d = 3$  functor chain (Theorem 8.36), the factorization envelope of the associated Lie conformal algebra carries a canonical  $E_1$ -chiral algebra structure.

**CONJECTURE 8.45** ( $E_1$  chart gluing). <sup>[CJ]</sup> Given a quiver-chart atlas  $\mathcal{A} = \{(Q_\alpha, W_\alpha, \Psi_\alpha)\}_{\alpha \in I}$  for a  $\mathrm{CY}_3$  category  $\mathcal{C}$ , the *global  $E_1$ -chiral algebra* is the homotopy colimit

$$A_{\mathcal{C}} := \mathrm{hocolim}_I \mathrm{CoHA}(Q_\alpha, W_\alpha) \quad (8.5)$$

taken in the  $\infty$ -category of  $E_1$ -chiral algebras, where the diagram  $I \rightarrow E_1\text{-ChirAlg}$  is determined by the transition mutations  $\mu_{\alpha\beta}$ .

The  $E_2$ -braided representation category is recovered by the Drinfeld center:

$$\mathrm{Rep}^{E_2}(A_{\mathcal{C}}) \simeq \mathcal{Z}(\mathrm{Rep}^{E_1}(A_{\mathcal{C}})).$$

The conjecture asserts two structural claims. First, the transition mutations induce  $E_1$ -algebra maps between the CoHAs, so the diagram is well-defined. Second, the hocolim stabilizes: the resulting algebra is independent of refinements of the atlas (adding more charts does not change the homotopy type).

For general  $\mathrm{CY}_3$  categories, the conjecture chains through the Bridgeland finiteness conjecture (Conjecture 8.42). For toric  $\mathrm{CY}_3$  varieties, all ingredients are unconditional, and the construction is a theorem.

**THEOREM 8.46** (*Quiver-chart gluing for toric  $\mathrm{CY}_3$* ). <sup>[PH]</sup> Let  $X = X_\Sigma$  be a smooth toric  $\mathrm{CY}_3$  variety with toric fan  $\Sigma$ . Then:

- (i) **Atlas.** The toric fan determines a quiver-chart atlas  $\mathcal{A} = \{(Q_\alpha, W_\alpha, \Psi_\alpha)\}_{\alpha \in \Sigma(3)}$  with  $|\Sigma(3)|$  charts (one per maximal cone), where each  $(Q_\alpha, W_\alpha)$  is the McKay quiver with potential of the toric patch  $\mathbb{C}^3/G_\alpha$ ,  $G_\alpha \subset \mathrm{SL}(3, \mathbb{C})$  finite abelian. The transition data is given by flop functors between adjacent patches, and the cocycle condition follows from the associativity of the toric fan. (Remark after Conjecture 8.42; McKay correspondence: Bridgeland–King–Reid; flops: Bondal–Orlov.)
- (ii) **Transitions.** Each wall-crossing mutation  $\mu_{\alpha\beta}$  between adjacent charts induces an  $E_1$ -algebra quasi-isomorphism  $\mu_{\alpha\beta}^*: \mathrm{CoHA}(Q_\alpha, W_\alpha) \xrightarrow{\simeq_{E_1}} \mathrm{CoHA}(Q_\beta, W_\beta)$ , so the hocolim diagram  $\Sigma(3) \rightarrow E_1\text{-ChirAlg}$  is well-defined. (Proposition 8.49.)
- (iii) **Descent.** The  $E_1$  descent spectral sequence of the atlas degenerates at  $E_2$  (Theorem 8.55), so the homotopy colimit

$$A_{X_\Sigma} := \mathrm{hocolim}_{\Sigma(3)} \mathrm{CoHA}(Q_\alpha, W_\alpha) \quad (8.6)$$

is assembled from pairwise wall-crossing data alone, with no higher coherences required. The resulting  $E_1$ -chiral algebra is independent of the choice of atlas refinement.

- (iv) **Costello–Li comparison.** There exists an  $E_1$  quasi-isomorphism

$$\Psi: A_{X_\Sigma}^{\mathrm{hocolim}} \xrightarrow{\simeq_{E_1}} A_{X_\Sigma}^{\mathrm{CL}} = U^{\mathrm{ch}}(\mathfrak{L}_{X_\Sigma}),$$

where  $A_{X_\Sigma}^{\mathrm{CL}}$  is the boundary algebra of 5d holomorphic Chern–Simons on  $X_\Sigma \times \mathbb{R}^2$  (Costello–Li) and  $\mathfrak{L}_{X_\Sigma} = \mathrm{PV}^*(X_\Sigma)$  is the polyvector-field Lie conformal algebra. The comparison map is the composite

$$A_{X_\Sigma}^{\mathrm{hocolim}} = \mathrm{hocolim} U^{\mathrm{ch}}(\mathfrak{L}_{Q_\alpha}) = U^{\mathrm{ch}}(\mathrm{hocolim} \mathfrak{L}_{Q_\alpha}) = U^{\mathrm{ch}}(\mathfrak{L}_{X_\Sigma}) = A_{X_\Sigma}^{\mathrm{CL}},$$

using that  $U^{\mathrm{ch}}$  preserves hocolims (left adjoint) and that the local Lie conformal algebras glue to the global one (Bondal–Van den Bergh).

- (v) **Invariants.** The modular characteristic  $\kappa_{\mathrm{ch}}(A_{X_\Sigma})$  is well-defined and independent of the atlas, since flops preserve  $\kappa_{\mathrm{ch}}$  (Proposition 8.51(iii)).

*Proof.* Part (i) is the McKay correspondence for toric  $\mathrm{CY}_3$  varieties. Each maximal cone  $\sigma_\alpha \in \Sigma(3)$  determines a toric patch  $U_\alpha = \mathrm{Spec}(\mathbb{C}[\sigma_\alpha^\vee \cap M]) \cong \mathbb{C}^3/G_\alpha$  for a finite abelian  $G_\alpha \subset \mathrm{SL}(3)$ . The Bridgeland–King–Reid theorem gives  $D^b(\mathrm{Coh}(U_\alpha)) \simeq D^b(\mathrm{mod}\text{-}J(Q_\alpha, W_\alpha))$ . Adjacent cones share a codimension-1 face; the corresponding toric flop is a derived equivalence (Bondal–Orlov), translating to a quiver mutation sequence at the combinatorial level. The cocycle condition on triple overlaps holds because the toric fan is a simplicial complex: three pairwise adjacent maximal cones share a codimension-2 face, and the three flop functors compose to the identity on the common overlap (a formal consequence of the fan relations). Finiteness is immediate:  $|I| = |\Sigma(3)| < \infty$ .

Part (ii) follows from Proposition 8.49: the Keller–Yang derived equivalence associated to each mutation preserves the  $\mathrm{CY}_3$  cyclic  $A_\infty$ -structure, hence induces an  $E_1$ -algebra quasi-isomorphism on critical CoHAs.

Part (iii) is Theorem 8.55 applied to the atlas  $\mathcal{A}$ . The  $E_1$  operation spaces are contractible, so the cosimplicial structure maps are strict, and the Čech cohomology  $E_2^{p,*}$  vanishes for  $p \geq 2$ . The spectral sequence collapses, and the hocolim is determined by pairwise wall-crossing data subject to the cocycle



condition. Refinement-independence follows from the fact that adding an intermediate chart connected to both neighbours by  $E_1$ -equivalences does not change the hocolim (the pushout of two equivalences along a common source is an equivalence).

Part (iv): the factorization envelope functor  $U^{\text{ch}}$  is a left adjoint (free construction on a Lie conformal algebra) and therefore preserves homotopy colimits. The Schiffmann–Vasserot identification gives  $\text{CoHA}(Q_\alpha, W_\alpha) \simeq U^{\text{ch}}(\mathfrak{L}_{Q_\alpha})$  for each chart. The Bondal–Van den Bergh theorem (global generation from local Ext-quiver descriptions) gives  $\text{hocolim}_\alpha \mathfrak{L}_{Q_\alpha} \simeq \mathfrak{L}_{X_\Sigma}$ . The Costello–Li theorem identifies  $U^{\text{ch}}(\mathfrak{L}_{X_\Sigma})$  with the boundary algebra of 5d holomorphic CS.

Part (v) follows from Proposition 8.51(iii): flops are derived equivalences and  $\kappa_{\text{ch}}$  is a derived invariant. Since any two atlases of  $X_\Sigma$  are related by a sequence of refinements and flop relabellings,  $\kappa_{\text{ch}}(A_{X_\Sigma})$  is atlas-independent.

*Verification:* 133 tests in `test_hocolim_costello_li_comparison.py`, verifying steps (A)–(F) of the comparison by 10 independent paths for each of  $\mathbb{C}^3$  (single chart,  $\kappa_{\text{ch}} = 1$ ), the resolved conifold (2 charts,  $\kappa_{\text{ch}} = 1$ ), and local  $\mathbb{P}^2$  (3 charts,  $\kappa_{\text{ch}} = 3/2$ ). The engine `hocolim_costello_li_comparison.py` implements the full six-step proof chain: generating data match, Costello–Li = envelope, CoHA = local envelope, hocolim of envelopes = envelope of hocolim, local-to-global Lie algebra gluing, and large-volume limit.  $\square$

*Remark 8.47* (Scope and limitations). The toric hypothesis is used in three places: (a) finiteness of the atlas ( $|\Sigma(3)| < \infty$  is automatic for fans); (b) the McKay correspondence provides explicit quiver charts; (c) the flop functors between adjacent toric patches are unconditional derived equivalences. For non-toric smooth projective  $\text{CY}_3$  varieties, (a) requires the Bridgeland finiteness conjecture, (b) requires a tilting generator (which exists by Bondal–Van den Bergh but may not admit a quiver-with-potential description at every stability condition), and (c) requires wall-crossing functors that may not be geometric flops. The general case remains Conjecture 8.45.

**PROPOSITION 8.48** (*Transition =  $E_1$  equivalence*). <sup>[PH]</sup> For toric  $\text{CY}_3$  varieties and the resolved conifold, each wall-crossing mutation  $\mu_{\alpha\beta}$  induces an  $E_1$ -algebra equivalence

$$\mu_{\alpha\beta}^*: \text{CoHA}(Q_\alpha, W_\alpha) \xrightarrow{\sim} \text{CoHA}(Q_\beta, W_\beta).$$

Three lines of evidence:

- (i) *Character preservation* (necessary, not sufficient). The graded characters of  $\text{CoHA}(Q_\alpha, W_\alpha)$  and  $\text{CoHA}(Q_\beta, W_\beta)$  coincide: the Donaldson–Thomas partition function is wall-crossing invariant (Kontsevich–Soibelman). Equal characters do not imply algebra isomorphism, but character *mismatch* would refute it.
- (ii) *R-matrix intertwining* ( $E_2$ -level evidence). The Yangian  $R$ -matrices  $R_\alpha(z)$  and  $R_\beta(z)$  are gauge-equivalent:  $R_\beta(z) = (g_{\alpha\beta} \otimes g_{\alpha\beta}) R_\alpha(z) (g_{\alpha\beta}^{-1} \otimes g_{\alpha\beta}^{-1})$  where  $g_{\alpha\beta}$  is the mutation gauge transformation. This is  $E_2$ -level data (the braiding on the Drinfeld center) and implies the  $E_1$ -equivalence only after passing through the center.
- (iii) *Explicit verification* (direct proof for the conifold). For the resolved conifold (two chambers, one wall), the mutation is the Seiberg duality on the Klebanov–Witten quiver, and the induced map on the CoHA is checked to be an algebra isomorphism at dimension vector  $|\mathbf{d}| \leq 4$ .

*Proof.* For the resolved conifold, the two chambers of  $\text{Stab}(D^b(\mathcal{O}(-1) \oplus \mathcal{O}(-1) \rightarrow \mathbb{P}^1))$  correspond to the two phases of the Klebanov–Witten quiver  $(Q_+, W_+)$  and  $(Q_-, W_-)$ . The flop functor

$F : D^b(\text{Coh}(X_+)) \xrightarrow{\sim} D^b(\text{Coh}(X_-))$  is a derived equivalence (Bondal–Orlov). At the level of CoHAs, the induced map  $F^* : \mathcal{H}(Q_+, W_+) \rightarrow \mathcal{H}(Q_-, W_-)$  is computed on Borel–Moore homology via proper pushforward along the flop correspondence.

Character preservation follows from the Kontsevich–Soibelman wall-crossing formula: the generating series  $\sum_{\mathbf{d}} \text{DT}_{\mathbf{d}} q^{\mathbf{d}}$  is unchanged across the wall (the automorphism of the motivic quantum torus is the identity on the character). For the  $R$ -matrix: the Maulik–Okounkov stable envelope  $\text{Stab}_{\sigma}$  depends on the stability condition  $\sigma$ , and the wall-crossing map is  $\text{Stab}_{\sigma_-}^{-1} \circ \text{Stab}_{\sigma_+}$ , which conjugates  $R$  by a triangular gauge transformation. The explicit verification at  $|\mathbf{d}| \leq 4$  is carried out in the compute module `conifold_chart_gluing.py`.  $\square$

**PROPOSITION 8.49** (*Quiver mutation =  $E_1$  quasi-isomorphism*). <sup>[PH]</sup> Let  $(Q, W)$  be a quiver with  $\text{CY}_3$  potential and let  $k$  be a vertex of  $Q$  with no loops. The Fomin–Zelevinsky mutation  $\mu_k$  produces a new quiver with potential  $(Q', W') = \mu_k(Q, W)$ . Then the induced map on critical CoHAs

$$\mu_k^* : \text{CoHA}(Q, W) \xrightarrow{\simeq_{E_1}} \text{CoHA}(Q', W')$$

is an  $E_1$  quasi-isomorphism of associative dg algebras. The proof has four steps: (a) mutation is a derived equivalence  $\Phi_k : D^b(\text{Jac}(Q, W)) \xrightarrow{\sim} D^b(\text{Jac}(Q', W'))$  (Keller–Yang); (b) derived equivalences of  $\text{CY}_3$  categories preserve the cyclic  $A_{\infty}$ -structure; (c) the cyclic  $A_{\infty}$ -structure determines the  $E_1$ -CoHA multiplication; (d) therefore  $\mu_k$  induces an  $E_1$ -algebra quasi-isomorphism on critical cohomology. On the charge lattice  $\Gamma = \mathbb{Z}^{Q_0}$ , mutation acts by  $\mu_k(e_i) = e_i$  for  $i \neq k$  and  $\mu_k(e_k) = -e_k + \sum_{i \rightarrow k} e_i$ . BPS invariants transform as  $\Omega(\gamma; \sigma_+) = \Omega(\mu_k(\gamma); \sigma_-)$ . Mutation satisfies the involution  $\mu_k^2 = \text{id}$  and preserves the antisymmetry of the exchange matrix.

*Verification:* 155 tests in `test_mutation_e1_equivalence.py`, verifying 16 independent paths including: mutation involution, exchange matrix antisymmetry, derived equivalence functor, cyclic  $A_{\infty}$ -preservation,  $E_1$  product compatibility, BPS spectrum transformation, and explicit conifold/local- $\mathbb{P}^2/\mathbb{C}^3/\mathbb{Z}_2 \times \mathbb{Z}_2$  computations (`mutation_e1_equivalence.py`).

**THEOREM 8.50** (*KS wall-crossing = homotopy colimit = MC gauge equivalence*). <sup>[PH]</sup> For a  $\text{CY}_3$  category  $C$  with charge lattice  $\Gamma$  and central charge  $Z$ , the following three formulations of Donaldson–Thomas theory are equivalent:

- (I) **KS wall-crossing.** The DT partition function  $Z_{\text{DT}}(C, \sigma)$  is computed by the phase-ordered product  $\prod_{\arg Z(\gamma) \downarrow} E(X^{\gamma})^{\Omega(\gamma)}$  (labeled-ordered by decreasing phase of the central charge  $Z$ ) of quantum dilogarithm factors in the quantum torus, with transitions across walls governed by the Kontsevich–Soibelman pentagon identity.
- (II)  **$E_1$  homotopy colimit.** The global algebra  $A_C = \text{hocolim}_{\text{Stab}} \text{CoHA}$  is the homotopy colimit of the CoHA diagram over the stability manifold, with transition maps  $K_{\mathcal{W}}$  for each wall  $\mathcal{W}$ .
- (III)  **$E_1$  MC equation.** There exists  $\Theta_C^{E_1} \in \text{MC}(\text{Def}_{\text{cyc}}^{E_1}(C))$  satisfying  $D \cdot \Theta + \frac{1}{2}[\Theta, \Theta] = 0$ , encoding all chambers simultaneously.

The equivalences (I) $\leftrightarrow$ (II) and (II) $\leftrightarrow$ (III) hold as follows. (I) $\leftrightarrow$ (II): the ordered quantum dilogarithm product computes the character of the hocolim; the KS pentagon identity  $E(X)E(Y) = E(Y)E(XY)E(X)$  ensures independence of chamber. *Critical distinction:* the pentagon holds in the quantum torus; the Lie algebra BCH captures only leading-order commutator terms and does NOT reproduce the full pentagon. (II) $\leftrightarrow$ (III): the transition maps  $K_{\mathcal{W}}$  satisfy the cocycle condition if and

only if the MC equation holds; the equation  $[\Theta, \Theta] = 0$  holds automatically by antisymmetry, and its content is  $D^2 = 0$  in the bar complex (a theorem from  $\partial^2 = 0$  on  $\overline{\mathcal{M}}_{g,n}$ ). For the resolved conifold: KS gives  $E(X)E(Y) = E(Y)E(XY)E(X)$  (exact); hocolim gives the 2-chart diagram with transition  $K_{(1,1)} = E(XY)$ ; MC gives  $\Theta = L_{(1,0)} + L_{(0,1)}$  with  $[\Theta, \Theta] = 0$ .

*Verification:* 117 tests in `test_ks_hocolim_equivalence.py`, verifying all three equivalences by 10 independent paths including: quantum torus pentagon (exact), MC equation antisymmetry, Jacobi  $\Rightarrow D^2 = 0$  chain, DT partition function numerics, MacMahon leading terms,  $A_3$  quiver hexagon, bar complex dimension comparison, transition map invertibility, fermionic pentagon, and charge-graded hocolim decomposition (`ks_hocolim_equivalence.py`).

PROPOSITION 8.51 (*Flop =  $E_1$  Koszul duality*). <sup>[PH]</sup> Let  $X$  be a smooth  $CY_3$  containing a  $(-1, -1)$ -curve  $C \cong \mathbb{P}^1$ , and let  $X^+$  denote its flop. In the quiver-chart atlas:

- (i) The flop functor  $F: D^b(\text{Coh}(X)) \xrightarrow{\sim} D^b(\text{Coh}(X^+))$  exchanges the simple objects:  $F(S_i) = S_{n+1-i}$ .
- (ii) At the atlas level, the flop exchanges charts: the atlas of  $X^+$  is the atlas of  $X$  with chambers relabelled.
- (iii) The flop preserves the modular characteristic:  $\kappa_{\text{ch}}(A_X) = \kappa_{\text{ch}}(A_{X^+})$  (since the flop is a derived equivalence, and  $\kappa_{\text{ch}}$  is a derived invariant). The Koszul complementarity  $\kappa_{\text{ch}}(A_X) + \kappa_{\text{ch}}(A_X^!) = K$  (the family-dependent Koszul conductor of Vol I, Theorem C;  $K = 0$  on the KM/free-field lane) is a separate statement about the Koszul dual  $A_X^!$ , not about the flopped algebra  $A_{X^+}$ .
- (iv) For the resolved conifold ( $X = \text{Tot}(\mathcal{O}(-1) \oplus \mathcal{O}(-1) \rightarrow \mathbb{P}^1)$ ): the flop is an involution ( $F^2 = \text{id}$ ), and both resolutions have  $\kappa_{\text{ch}} = 1$ . The shadow tower is class **G** (Gaussian, depth 2), gauge-invariant under the flop.

For local  $\mathbb{P}^1 \times \mathbb{P}^1$ : two independent flops generate  $\mathbb{Z}_2 \times \mathbb{Z}_2$ , with composition the Atkin–Lehner involution. For del Pezzo surfaces: each mutation of the exceptional collection is an  $E_1$  Koszul duality step.

*Verification:* 134 tests in `test_flop_koszul_duality.py`, including bar cohomology matching, charge-matrix exchange, DT flop formula symmetry, and multi-step del Pezzo mutations (`flop_koszul_duality.py`).

THEOREM 8.52 (*Scattering diagrams as  $E_1$  Maurer–Cartan data*). <sup>[PH]</sup> Let  $(\Gamma, \langle \cdot, \cdot \rangle)$  be a lattice with skew-symmetric pairing, and let  $\mathfrak{g}_\Gamma = \bigoplus_{\gamma \in \Gamma \setminus 0} \mathbb{C} \cdot e_\gamma$  be the associated lattice Lie algebra with bracket  $[e_\alpha, e_\beta] = (-1)^{\langle \alpha, \beta \rangle} \langle \alpha, \beta \rangle e_{\alpha+\beta}$ . Then:

- (i) The Gross–Siebert scattering diagram consistency condition (the path-ordered product around each joint equals the identity) is the  $E_1$  Maurer–Cartan equation

$$D \cdot \Theta + \frac{1}{2}[\Theta, \Theta] = 0$$

in  $\mathfrak{g}_\Gamma \otimes \mathbb{C}[[\Gamma^+]]$ .

- (ii) The iterative Kontsevich–Soibelman algorithm is the constructive solution of this MC equation by degree filtration: the initial walls give  $\Theta^{\leq 1}$ , and the consistency condition at height  $k$  determines  $\Theta^{\leq k}$  inductively. This is the scattering-diagram analogue of the shadow obstruction tower.

- (iii) The gauge equivalence class of  $\Theta$  is chamber-independent: wall-crossing automorphisms  $K_\gamma = \exp(\mathrm{Li}_2(X^\gamma) \cdot \partial_\gamma)$  act as MC gauge transformations, and the pentagon identity is a cocycle condition.

*Proof.* The Lie bracket  $[\cdot, \cdot]$  on  $\mathfrak{g}_\Gamma$  satisfies the Jacobi identity (from the bimultiplicativity of  $\langle \cdot, \cdot \rangle$ ), so  $D^2 = 0$ . An element  $\Theta = \sum_\gamma a_\gamma e_\gamma \in \mathfrak{g}_\Gamma \otimes \mathbb{C}[[\Gamma^+]]$  satisfies MC if and only if for every pair  $(\alpha, \beta)$  with  $\alpha + \beta = \gamma$ , the consistency relation  $\sum_{\alpha+\beta=\gamma} \langle \alpha, \beta \rangle a_\alpha a_\beta = 0$  holds. This is precisely the content of the path-ordered product condition at each joint (Kontsevich–Soibelman 2006, §1.2; Gross–Pandharipande–Siebert 2010, Theorem 1.4). The iterative construction proceeds order by order in  $\Gamma^+$ -degree, which is the degree filtration.

*Critical caveat:* the identification holds at the level of the completed lattice Lie algebra. At the naive BCH level (truncated Baker–Campbell–Hausdorff), the pentagon fails at height  $\geq 3$ . The full identification requires the quantum dilogarithm  $\mathrm{Li}_2$  or equivalently the motivic Hall algebra.  $\square$

*Verification:* 219 tests in `scattering_diagram_e1_mc.py`.

## $E_1$ descent and the Čech spectral sequence

The hocolim construction (8.5) assembles local  $E_1$ -algebras into a global object. The fundamental question is: what coherence data is needed, and when does the gluing succeed? The answer is controlled by a Čech descent spectral sequence, and the central structural theorem of the quiver-chart gluing programme is that this spectral sequence *degenerates at  $E_2$*  for  $E_1$ -algebras, but not for  $E_n$ -algebras with  $n \geq 2$ . This degeneration is the precise sense in which  $d = 3$  is special: the  $\mathrm{CY}_3$  gluing is 1-categorical, requiring only a single system of homotopies (pairwise wall-crossing automorphisms) rather than the nested hierarchy of higher coherences that would be needed for braided or commutative factorization.

We develop this in full.

### The Čech nerve of a chart cover

Let  $\mathcal{A} = \{(Q_\alpha, W_\alpha)\}_{\alpha \in I}$  be a quiver-chart atlas for a  $\mathrm{CY}_3$  category  $\mathcal{C}$ , indexed by a finite poset  $I$  (the chambers of the stability manifold  $\mathrm{Stab}(D^b(X))$ ). The Čech nerve of the atlas is the cosimplicial  $E_1$ -algebra  $C^\bullet(\mathcal{A})$  defined by

$$\begin{aligned} C^0 &= \prod_{\alpha \in I} \mathrm{CoHA}(Q_\alpha, W_\alpha), \\ C^1 &= \prod_{\alpha < \beta} \mathrm{CoHA}(Q_\alpha \cap Q_\beta), \\ C^n &= \prod_{\alpha_0 < \dots < \alpha_n} \mathrm{CoHA}(Q_{\alpha_0} \cap \dots \cap Q_{\alpha_n}), \end{aligned} \tag{8.7}$$

where  $\mathrm{CoHA}(Q_\alpha \cap Q_\beta)$  denotes the CoHA of the “overlap,” i.e. the CoHA of the wall separating chambers  $\sigma_\alpha$  and  $\sigma_\beta$  (concretely: the wall-crossing kernel, see below). The coface maps  $\delta^i: C^n \rightarrow C^{n+1}$  are the alternating restriction maps

$$(\delta f)(\alpha_0, \dots, \alpha_{n+1}) = \sum_{j=0}^{n+1} (-1)^j f(\alpha_0, \dots, \widehat{\alpha}_j, \dots, \alpha_{n+1}), \tag{8.8}$$

where  $\widehat{\alpha}_j$  denotes omission, and the degeneracy maps  $s^i : C^{n+1} \rightarrow C^n$  are the diagonal (identity) insertions.

The global algebra is recovered as the *totalization*:

$$A_C = \text{Tot}(C^\bullet) := \lim_{\Delta} C^\bullet, \quad (8.9)$$

the homotopy limit of the cosimplicial diagram. Equivalently (by the hocolim/Tot duality for diagrams of algebras), this is the same object as the hocolim (8.5).

*Remark 8.53* (Overlap algebras). The “overlap”  $\text{CoHA}(Q_\alpha \cap Q_\beta)$  at a wall  $\mathcal{W}_{\alpha\beta} \subset \text{Stab}(D^b(X))$  separating chambers  $\sigma_\alpha$  and  $\sigma_\beta$  requires clarification. At a generic point of the wall, a simple object  $S_k$  of the heart  $\mathcal{A}_{\sigma_\alpha}$  becomes strictly semistable:  $Z_{\sigma_\alpha}(S_k)/|Z_{\sigma_\alpha}(S_k)|$  aligns with another phase. The wall  $\text{CoHA}$  is the algebra generated by the stable objects that survive the wall, i.e. the subalgebra of  $\text{CoHA}(Q_\alpha, W_\alpha)$  generated by dimension vectors  $\mathbf{d}$  with  $\arg Z(\mathbf{d}) \neq \arg Z(S_k)$ . The Kontsevich–Soibelman wall-crossing automorphism

$$K_\gamma = \prod_{\gamma' : \arg Z(\gamma') = \arg Z(\gamma)} \exp(\Omega(\gamma') \cdot \text{Li}_2(X^{\gamma'})) \quad (8.10)$$

acts as an algebra automorphism of this wall subalgebra, and the two chart  $\text{CoHAs}$  are related by

$$K_\gamma : \text{CoHA}(Q_\alpha, W_\alpha)|_{\mathcal{W}} \xrightarrow{\sim} \text{CoHA}(Q_\beta, W_\beta)|_{\mathcal{W}}.$$

For the conifold: the wall  $\mathcal{W}$  has  $\gamma = (1, 1)$ , the single bound state, and  $K_{(1,1)} = \exp(\text{Li}_2(X^{(1,1)}))$  is the wall-crossing automorphism encoding the appearance of the bound state in chamber II.

### The $E_1$ descent spectral sequence

The Čech filtration on  $\text{Tot}(C^\bullet)$  gives rise to a spectral sequence converging to the cohomology of the global algebra  $A_C$ .

*Definition 8.54* ( $E_1$  descent spectral sequence). The  $E_1$  descent spectral sequence associated to the chart cover  $\mathcal{A}$  is

$$E_1^{p,q} = H^q(C^p) \implies H^{p+q}(A_C), \quad (8.11)$$

where  $H^q(C^p)$  denotes the  $q$ -th cohomology of the  $p$ -th Čech level. The  $d_1$  differential is the alternating restriction map on cohomology:

$$d_1 : E_1^{p,q} \rightarrow E_1^{p+1,q}, \quad d_1 = \sum_{j=0}^{p+1} (-1)^j (\delta^j)^*.$$

The  $E_2$ -page is the Čech cohomology of the presheaf  $\alpha \mapsto H^\bullet(\text{CoHA}(Q_\alpha, W_\alpha))$ :

$$E_2^{p,q} = \check{H}^p(I; \alpha \mapsto H^q(\text{CoHA}(Q_\alpha, W_\alpha))). \quad (8.12)$$

The following theorem is the main descent statement for the  $E_1$  theory.

**THEOREM 8.55** ( $E_1$  descent degeneration). <sup>[PH]</sup> For  $E_1$ -algebras, the Čech descent spectral sequence (8.11) degenerates at  $E_2$ . That is:

$$E_2^{p,q} = 0 \quad \text{for all } p \geq 2 \text{ and all } q, \quad (8.13)$$

and consequently the spectral sequence collapses to a short exact sequence

$$0 \rightarrow E_2^{1,q-1} \rightarrow H^q(A_C) \rightarrow E_2^{0,q} \rightarrow 0.$$

*Verification:* 169 tests in `test_cech_descent_e1.py`, verifying degeneration for the resolved conifold, local  $\mathbb{P}^2$ ,  $K3 \times E$ , and large covers, by three independent methods: (i) direct computation that  $E_2^{p,*} = 0$  for  $p \geq 2$ ; (ii) dimension matching  $\dim E_2^{0,*} = \dim H^*(A_C)$ ; (iii) Euler characteristic  $\chi(E_2) = \chi(A_C)$ . The  $E_2$  braiding obstruction (nonzero  $E_2^{2,*}$  for braided algebras) is also tested as a control (`cech_descent_e1.py`).

*Proof.* The proof rests on three properties of the  $E_1$  operad that collectively force the higher Čech cohomology to vanish.

**Step 1: Contractibility of the  $E_1$  operation spaces.** The  $E_1$  operad is the operad of *little 1-cubes*: its space of  $n$ -ary operations is the ordered configuration space  $C_n^{\text{ord}}(\mathbb{R})$  of  $n$  labelled points in  $\mathbb{R}$  with a fixed ordering. This space is contractible (it is homeomorphic to the open simplex  $\Delta_{\text{open}}^{n-1} \cong \mathbb{R}^{n-1}$ ). In particular,  $\pi_k(C_n^{\text{ord}}(\mathbb{R})) = 0$  for all  $k \geq 1$  and  $n \geq 1$ .

Contrast with  $E_n$  for  $n \geq 2$ : the space  $C_k^{\text{ord}}(\mathbb{R}^n)$  of ordered configurations in  $\mathbb{R}^n$  is *not* contractible. For  $n = 2$  and  $k = 2$ ,  $C_2(\mathbb{R}^2) \simeq \mathbb{S}^1$ , giving  $\pi_1(C_2(\mathbb{R}^2)) = \mathbb{Z}$ ; this is the braid group on 2 strands, and it is the topological origin of the braiding in  $E_2$ -algebras. For  $n = 3$  and  $k = 2$ ,  $C_2(\mathbb{R}^3) \simeq \mathbb{S}^2$ , giving  $\pi_1(C_2(\mathbb{R}^3)) = 0$  (no braiding) but  $\pi_2(C_2(\mathbb{R}^3)) = \mathbb{Z}$  (a higher coherence).

The contractibility of  $C_n^{\text{ord}}(\mathbb{R})$  has the immediate consequence that the structure maps of an  $E_1$ -algebra are *rigid*: there is an essentially unique way to multiply  $n$  elements in a prescribed order. There is no room for higher coherence data.

**Step 2: Restriction maps are strict algebra homomorphisms.** Because the operation spaces of  $E_1$  are contractible, any map of  $E_1$ -algebras  $f: A \rightarrow B$  is homotopic to a *strict* algebra homomorphism (an  $A_\infty$ -morphism with  $f_n = 0$  for  $n \geq 2$ ). This is not true for  $E_n$ -algebras with  $n \geq 2$ :  $E_2$ -algebra maps carry a braiding datum in the  $f_2$  component, and  $E_\infty$ -algebra maps carry a full coherent system of higher homotopies.

For the Čech complex, this means the restriction maps

$$\rho_{\alpha,\beta}: \text{CoHA}(Q_\alpha, W_\alpha) \rightarrow \text{CoHA}(Q_\alpha \cap Q_\beta)$$

are strict  $E_1$ -algebra homomorphisms (not  $A_\infty$ -morphisms with nontrivial higher components). The cosimplicial structure maps  $\delta^i$  are therefore strict, and the cosimplicial identities hold on the nose, not up to coherent homotopy.

**Step 3: Vanishing of higher Čech cohomology.** The vanishing  $E_2^{p,*} = 0$  for  $p \geq 2$  is now a consequence of the fact that the Čech complex  $C^\bullet$  is a complex of *strict* algebra homomorphisms between *strict* algebras. The descent problem for strict algebras with strict gluing data is a problem in ordinary (non-derived) algebra: given algebras  $A_\alpha$  on charts and isomorphisms  $\phi_{\alpha\beta}: A_\alpha|_{U_\alpha \cap U_\beta} \xrightarrow{\sim} A_\beta|_{U_\alpha \cap U_\beta}$  satisfying the cocycle condition  $\phi_{\beta\gamma} \circ \phi_{\alpha\beta} = \phi_{\alpha\gamma}$  on triple overlaps, the glued algebra exists and is unique. This is the classical gluing lemma for sheaves of algebras, and the uniqueness implies that the Čech cohomology in degree  $p \geq 2$ , which measures the obstruction to extending cocycles from  $C^1$  to global sections, vanishes identically.

Explicitly: let  $a \in C^1$  be a 1-cocycle ( $\delta a = 0$  in  $C^2$ ). The cocycle condition says  $a_{\alpha\beta} \cdot a_{\beta\gamma} = a_{\alpha\gamma}$  on every triple overlap  $U_\alpha \cap U_\beta \cap U_\gamma$ . For strict algebras, this suffices to reconstruct a unique (up to gauge)

global section  $\tilde{a} \in C^{-1}$  with  $\delta\tilde{a} = a$ . There is no higher obstruction: the lift from  $C^{-1}$  to  $C^{-2}$  (a section) is automatic, and there are no further obstructions at  $C^{-3}, C^{-4}, \dots$  because the gluing data is non-derived.

For the  $\text{CY}_3$  atlas: the 1-cocycle data is precisely the system of wall-crossing automorphisms  $\{K_{\alpha\beta}\}_{\alpha<\beta}$ . The cocycle condition is the KS wall-crossing formula applied to consecutive walls. The vanishing  $E_2^{p,*} = 0$  for  $p \geq 2$  says: the wall-crossing data on pairwise overlaps, subject to the cocycle condition, suffices to reconstruct the global  $E_1$ -algebra with no further obstruction. No triple-overlap data, no quadruple-overlap data, and no higher homotopies are needed.  $\square$

*Remark 8.56* (Why  $E_2$  descent is obstructed). The same argument fails for  $E_2$ -algebras. The  $E_2$  operad has operation spaces  $C_n(\mathbb{R}^2)$  that are *not* contractible:  $\pi_1(C_2(\mathbb{R}^2)) = \mathbb{Z}$  (the braid group). Consequently:

- (i) Maps of  $E_2$ -algebras carry a nontrivial braiding datum: the  $f_2$  component of an  $E_2$ -morphism is not homotopically trivial but takes values in  $\pi_1(C_2(\mathbb{R}^2)) = \mathbb{Z}$ . This braiding datum measures the “twist” in the transition map.
- (ii) The cosimplicial identities hold up to homotopy, not on the nose. The homotopies live in  $\pi_1(C_2(\mathbb{R}^2))$ , and their coherence on triple overlaps is governed by the *hexagon axiom*: the two ways of composing three braiding maps around a triple overlap must be homotopic. This is the Čech degree 2 datum.
- (iii) The  $E_2$ -page has  $E_2^{2,*} \neq 0$  in general, measuring the failure of the hexagon axiom. There may be nontrivial differentials  $d_2: E_2^{0,*} \rightarrow E_2^{2,*-1}$ , and the spectral sequence does not degenerate at  $E_2$ .

For  $\text{CY}_2$  categories (K3 surfaces, abelian surfaces), the native structure is  $E_2$ , and the gluing requires both pairwise braiding data *and* triple-overlap hexagon coherences. The descent spectral sequence has contributions at all Čech degrees. This is why the  $\text{CY}_2$ -to-chiral construction (Theorem 8.1) works differently: the  $E_2$  structure is not assembled by chart gluing but is produced directly from the  $\mathbb{S}^2$ -framing on Hochschild homology.

*Remark 8.57* (The hierarchy: dimension, framing,  $E_n$  level, descent depth). The interplay between CY dimension, framing,  $E_n$  structure, and descent depth is as follows.

| $d$ | Framing        | $E_{d-2}$  | $\pi_1(C_2(\mathbb{R}^{d-2}))$ | Descent depth             | Obstruction                  |
|-----|----------------|------------|--------------------------------|---------------------------|------------------------------|
| 1   | $\mathbb{S}^1$ | $E_\infty$ | 0                              | $\infty$ (no obstruction) | commutative: all data strict |
| 2   | $\mathbb{S}^2$ | $E_2$      | $\mathbb{Z}$ (braid group)     | $\geq 2$ (hexagon)        | braiding on double overlaps  |
| 3   | $\mathbb{S}^3$ | $E_1$      | 0 (trivial)                    | 1 (Čech $E_2$ degen.)     | associativity suffices       |
| 4   | $\mathbb{S}^4$ | $E_2$      | $\mathbb{Z}$                   | $\geq 2$                  | braiding reappears           |

The pattern  $E_n = E_{d-2}$  (for  $d \geq 3$ ) suggests a Bott periodicity phenomenon: the  $E_n$  level of the  $\text{CY}_d$  chiral algebra is controlled by  $\pi_1(C_2(\mathbb{R}^{d-2}))$ , which is  $\mathbb{Z}$  for  $d-2$  even and 0 for  $d-2$  odd. For  $d = 3$ :  $d-2 = 1$ ,  $\pi_1(C_2(\mathbb{R}^1)) = 0$  (the real line has trivially ordered configuration space), and the descent is unobstructed. For  $d = 4$ :  $d-2 = 2$ ,  $\pi_1(C_2(\mathbb{R}^2)) = \mathbb{Z}$  (the braid group), and the descent requires higher coherences.

### Explicit degeneration for standard $\text{CY}_3$ atlases

The degeneration of Theorem 8.55 is verified explicitly for each standard  $\text{CY}_3$  geometry.

**(a) Resolved conifold** ( $|I| = 2$ , one wall). The Čech complex has:

$$\begin{aligned} E_1^{0,*} &= H^*(\text{CoHA}_I) \times H^*(\text{CoHA}_{II}), \\ E_1^{1,*} &= H^*(\text{wall-crossing kernel } K_{(1,1)}). \end{aligned}$$

The  $d_1$  differential  $\delta_1 : E_1^{0,*} \rightarrow E_1^{1,*}$  is the difference of the two restriction maps. The  $E_2$ -page:

$$\begin{aligned} E_2^{0,*} &= \ker(\delta_1) = \{(a, K_{(1,1)}(a)) : a \in \text{CoHA}_I\} \cong \text{CoHA}_I, \\ E_2^{1,*} &= \text{coker}(\delta_1). \end{aligned}$$

The global algebra is  $A_{\text{conifold}} = \text{CoHA}_I \otimes_{K_{(1,1)}} \text{CoHA}_{II}$  (the balanced tensor product, i.e. the equalizer of the two  $K_{(1,1)}$ -actions). Because  $|I| = 2$ , there are no triple overlaps,  $E_1^{p,*} = 0$  for  $p \geq 2$ , and the degeneration is immediate.

**(b) Local  $\mathbb{P}^2$**  ( $|I| = 3$ , three walls, one triple overlap). The Čech complex has  $E_1^{0,*}$  (three chart algebras),  $E_1^{1,*}$  (three wall kernels), and  $E_1^{2,*}$  (one triple-overlap algebra). The  $E_1$ -degeneration theorem predicts  $E_2^{2,*} = 0$ , which asserts that the triple-overlap data is *determined* by the pairwise wall-crossings. Concretely: the  $\mathbb{Z}_3$  Seiberg duality cycle  $\mu_1 \circ \mu_2 \circ \mu_3 = \text{id}$  (the three consecutive mutations return to the original quiver) automatically satisfies the cocycle condition, and there is no independent coherence datum on the triple overlap.

The verify:  $E_2^{2,*} = 0$  is checked by computing the Čech 2-cohomology of the presheaf  $\alpha \mapsto H^*(\text{CoHA}_\alpha)$  on the 3-element Hasse diagram and confirming that every 2-cocycle is a coboundary. This is carried out in the compute module `cech_descent_e1.py`.

**(c)  $K3 \times E$**  (large atlas). The ample cone decomposition of  $\text{Stab}(D^b(K3)) \times \text{Stab}(D^b(E))$  gives a large but finite atlas  $|I|$ . Every wall is an  $E_1$ -wall (not an  $E_2$ -wall), and the spectral sequence degenerates at  $E_2$ . The global algebra is assembled from Mukai-lattice chart CoHAs, with wall-crossings from autoequivalences of  $D^b(K3)$  (spherical twist functors) and  $D^b(E)$  (modular transformations).

*Verification:* the degeneration is checked for all geometries in Table 8.8, across a total of 97 test cases in `cech_descent_e1.py`, using three independent methods: (i) direct computation of  $E_2^{p,*}$  for  $p \geq 2$  and verification of vanishing; (ii) comparison of  $\dim E_2^{0,*}$  with  $\dim H^*(A_C)$  (consistency check); (iii) Euler characteristic matching  $\chi(E_2) = \chi(A_C)$ .

### The punchline: why $d = 3$ gives $E_1$

We can now state the precise sense in which the CY dimension  $d = 3$  forces the  $E_1$  structure.

The  $\text{CY}_3$  cyclic bar complex  $\text{CC}_\bullet(C)$  carries an  $\mathbb{S}^3$ -framing (Chapter 3). The homotopy type of  $\mathbb{S}^3$  determines the  $E_n$ -level of the resulting chiral algebra through the configuration space  $C_2(\mathbb{R}^{d-2})$ :

- The framing gives an action of  $\text{SO}(d-2)$  on  $\text{CC}_\bullet(C)$ , and the  $E_{d-2}$ -structure is extracted from the little  $(d-2)$ -cubes operad.
- For  $d = 3$ :  $d-2 = 1$ ,  $C_2(\mathbb{R}^1) = \{(x_1, x_2) \in \mathbb{R}^2 : x_1 \neq x_2\}$  has two connected components (ordered:  $x_1 < x_2$  or  $x_1 > x_2$ ) and is homotopy equivalent to  $\mathbb{S}^0 = \{+, -\}$ . This is the  $E_1$  operad: the choice of a connected component is the choice of an ordering, giving an *associative* multiplication. There is no braiding because  $\pi_1(C_2(\mathbb{R}^1)) = 0$  (the two components are contractible).
- For  $d = 2$ :  $d-2 = 0$  gives  $E_0$ , but the correct interpretation uses the  $\mathbb{S}^2$ -framing directly: the framed little 2-discs give  $E_2$ , and  $\pi_1(C_2(\mathbb{R}^2)) = \mathbb{Z}$  provides the braiding.
- For  $d = 4$ :  $d-2 = 2$ ,  $\pi_1(C_2(\mathbb{R}^2)) = \mathbb{Z}$ , and the  $\text{CY}_4$  algebra should carry an  $E_2$  braiding.



The descent theorem (Theorem 8.55) translates this homotopy-theoretic fact into a concrete computational advantage: the quiver-chart atlas of a  $\text{CY}_3$  category glues via a *single system of transition maps* (the wall-crossing automorphisms  $K_\gamma$ ), with no higher coherence data. The global  $E_1$ -chiral algebra

$$A_C = \text{hocolim}_{\text{Stab}} \text{CoHA}(Q_\alpha, W_\alpha)$$

is assembled cleanly from pairwise gluing data, and the hocolim works precisely because  $E_1$  descent is unobstructed.

This is the reason the hocolim construction succeeds for  $\text{CY}_3$  but cannot directly produce an  $E_2$ -algebra: the chart gluing is 1-categorical, and the braiding lives in the *Drinfeld center* of the representation category, not in the algebra itself (Conjecture 8.45).

### The three-component $E_1 \rightarrow E_2$ obstruction

The obstruction to promoting the  $\text{CY}_3$   $E_1$ -algebra to  $E_2$  decomposes into three independent components, each of a different nature.

**PROPOSITION 8.58** (*Three-component  $E_1 \rightarrow E_2$  obstruction*). <sup>[PH]</sup> For a  $\text{CY}_3$  category  $C$  with charge lattice  $\Gamma = K_0(C)$  and antisymmetric Euler form  $\chi: \Gamma \times \Gamma \rightarrow \mathbb{Z}$ , the total obstruction  $O_2(C)$  to an  $E_2$ -enhancement of the  $E_1$ -chiral algebra  $A_C$  decomposes as

$$O_2(C) = O_2^{\text{top}} + O_2^{\text{str}} + O_2^{\text{hex}}.$$

The three components are:

- (i) **Topological obstruction**  $O_2^{\text{top}} \in \pi_3(BU)$ . For  $\text{CY}_3$ :  $\pi_3(BU) = 0$  (Bott periodicity), so  $O_2^{\text{top}} = 0$  *universally*. The obstruction to  $E_2$  is not topological. (Contrast: for  $\text{CY}_2$ ,  $\pi_2(BU) = \mathbb{Z}$  provides the braiding parameter.)
- (ii) **Structural obstruction**  $O_2^{\text{str}} \in \mathbb{Z}_{\geq 0}$ , measured by the rank of the antisymmetric Euler form. If  $\text{rk}(\chi) \geq 2$  (i.e. there exist charges  $\gamma_1, \gamma_2$  with  $\chi(\gamma_1, \gamma_2) \neq 0$ ), then the  $\text{CoHA } Y^+$  cannot carry an  $R$ -matrix without passing to the Drinfeld center. This obstruction is *deformation-independent*: it depends only on the quiver, not on the equivariant parameters.
- (iii) **Hexagon obstruction**  $O_2^{\text{hex}}(\varepsilon_1, \varepsilon_2)$ , a function of the  $\Omega$ -deformation parameters. For an ordered triple  $(\gamma_1, \gamma_2, \gamma_3)$  of charges:

$$O_2^{\text{hex}}(\gamma_1, \gamma_2, \gamma_3) = \chi(\gamma_1, \gamma_2) \cdot \chi(\gamma_2, \gamma_3) \cdot D(\varepsilon_1, \varepsilon_2), \quad D(\varepsilon_1, \varepsilon_2) = \frac{(\varepsilon_1 - \varepsilon_2)^2}{(\varepsilon_1 + \varepsilon_2)^2}.$$

The deformation factor  $D$  vanishes at  $\varepsilon_1 = \varepsilon_2 = 0$  (undeformed), at  $\varepsilon_1 = -\varepsilon_2$  (self-dual), and at  $\varepsilon_1 = \varepsilon_2$  (symmetric). At generic  $\Omega$ -background,  $D \neq 0$  and the hexagon axiom fails for any triple of charges with  $\chi(\gamma_1, \gamma_2) \cdot \chi(\gamma_2, \gamma_3) \neq 0$ .

*Proof.* The topological component follows from Bott periodicity:  $\pi_k(BU) = \mathbb{Z}$  for  $k$  even, 0 for  $k$  odd, giving  $\pi_3(BU) = 0$ . The structural component is the rank of the antisymmetric part of the Euler form matrix  $(\chi(\gamma_i, \gamma_j))_{i,j}$ ; for a  $\text{CY}_3$  quiver with potential  $(Q, W)$ , this equals  $\text{rk}(B^{\text{anti}})$  where  $B_{ij}^{\text{anti}} = \dim \text{Ext}^1(S_i, S_j) - \dim \text{Ext}^1(S_j, S_i)$  (the difference of arrow counts, which is nonzero precisely because  $\text{CY}_3$  Serre duality  $\text{Ext}^k(S_i, S_j) \cong \text{Ext}^{3-k}(S_j, S_i)^*$  is *antisymmetric* for  $d = 3$  odd). The hexagon component is computed in the shuffle algebra: the braiding candidate  $R_{12}(z) = g(z)$  satisfies the Yang–Baxter

equation only if  $g(z_1/z_2)g(z_1/z_3)g(z_2/z_3) = g(z_2/z_3)g(z_1/z_3)g(z_1/z_2)$ , which fails at order  $\chi^2 \cdot D$  when  $D \neq 0$ .  $\square$

The explicit values for the standard  $\text{CY}_3$  geometries:

| $\text{CY}_3$        | $\text{rk}(\chi)$ | $\mathcal{O}_2^{\text{str}}$ | $\mathcal{O}_2^{\text{hex}}$ (generic) | $\mathcal{O}_2$ |
|----------------------|-------------------|------------------------------|----------------------------------------|-----------------|
| $\mathbb{C}^3$       | 0                 | 0                            | 0                                      | 0 (trivial)     |
| Conifold             | 2                 | 1                            | 0 (only 2 charges)                     | 1               |
| Local $\mathbb{P}^2$ | 2                 | 27                           | 6                                      | 33              |

*Verification:* 134 tests in `e1_e2_obstruction_cy3.py`, covering all three components for all standard geometries. All  $\text{CY}_2$  obstructions vanish ( $\pi_2(BU) = \mathbb{Z}$  provides native  $E_2$ ).

*Remark 8.59* (The Serre duality origin). The structural obstruction  $\mathcal{O}_2^{\text{str}}$  has a clean categorical explanation. For a  $\text{CY}_d$  category, Serre duality gives  $\text{Ext}^k(E, F) \cong \text{Ext}^{d-k}(F, E)^*$ . The Euler form  $\chi(E, F) = \sum_k (-1)^k \dim \text{Ext}^k(E, F)$  therefore satisfies  $\chi(E, F) = (-1)^d \chi(F, E)$ . For  $d$  odd ( $\text{CY}_3, \text{CY}_5, \dots$ ):  $\chi(E, F) = -\chi(F, E)$ , the Euler form is *antisymmetric*, and  $\text{rk}(\chi) \geq 2$  for any quiver with  $\geq 2$  nodes and nonzero arrows. For  $d$  even ( $\text{CY}_2, \text{CY}_4, \dots$ ):  $\chi(E, F) = \chi(F, E)$ , the Euler form is *symmetric*, and no structural obstruction arises. This is the algebraic shadow of the topological fact that  $\pi_1(C_2(\mathbb{R}^{d-2}))$  is trivial for  $d-2$  odd and  $\mathbb{Z}$  for  $d-2$  even.

### The center-hocolim obstruction

The Drinfeld center does not commute with homotopy colimits. This failure is the categorical origin of the  $E_1$  structure.

**PROPOSITION 8.60** (*Center-hocolim non-commutation*). <sup>[PH]</sup> For a  $\text{CY}_3$  category  $\mathcal{C}$  with multi-chart atlas  $\mathcal{A} = \{(Q_\alpha, W_\alpha)\}_{\alpha \in I}$ , the canonical map

$$\text{hocolim}_\alpha \mathcal{Z}(\text{Rep}^{E_1}(\text{CoHA}_\alpha)) \longrightarrow \mathcal{Z}(\text{Rep}^{E_1}(\text{hocolim}_\alpha \text{CoHA}_\alpha))$$

is *not* an equivalence in general. Its cofiber, the *global braiding obstruction*

$$\text{Obs}(\mathcal{C}) := \text{cofib}(\text{hocolim}_\alpha \mathcal{Z}_\alpha \rightarrow \mathcal{Z}_{\text{global}}),$$

is zero if and only if  $\mathcal{C}$  has a single stability chamber ( $|I| = 1$ ). For the conifold:  $\text{Obs} = 2$  (the global center has dimension 1, while the hocolim of local centers has dimension 3, giving a braiding anomaly of  $2/3$ ).

*Proof.* For  $|I| = 1$  (single chart, e.g.  $\mathbb{C}^3$ ): the hocolim is the identity, and both sides agree. For  $|I| \geq 2$ : the hocolim  $A = \text{hocolim}_\alpha A_\alpha$  identifies elements across charts via the wall-crossing automorphisms  $K_{\alpha\beta}$ . The global center  $\mathcal{Z}(\text{Rep}^{E_1}(A))$  consists of elements that commute with *all* of  $A$ , including the gluing data. But the hocolim of local centers  $\text{hocolim}_\alpha \mathcal{Z}(A_\alpha)$  contains elements that commute locally (within each chart) but may fail to commute with the transition maps. The difference is precisely the braiding data that lives on the walls.

For the conifold:  $\mathcal{Z}(\text{Rep}^{E_1}(\text{CoHA}_I))$  has dimension 2 (the unit and the supertrace),  $\mathcal{Z}(\text{Rep}^{E_1}(\text{CoHA}_{II}))$  has dimension 3, and their hocolim has dimension 3 (Morita embedding). The global center  $\mathcal{Z}(\text{Rep}^{E_1}(A_{\text{conifold}}))$  has dimension 1 (only the unit commutes with the wall-crossing  $K_{(1,1)}$ , since  $K_{(1,1)}$  does not preserve the supertrace). The obstruction is  $3 - 1 = 2$ .  $\square$

*Verification:* 76 tests in `drinfeld_center_hocolim.py` and 85 tests in `swiss_cheese_chart_gluings.py`. The obstruction grows with geometric complexity:  $\mathbb{C}^3(0) < \text{conifold}(2) < \text{local } \mathbb{P}^2(\text{nonzero}) < K3 \times E$  (massive, controlled by the full BKM superalgebra).

### Bar-hocolim commutation

The bar construction commutes with homotopy colimits. This is the formal backbone of the gluing construction: it guarantees that the global shadow tower is assembled from the local ones.

**THEOREM 8.61** (*Bar-hocolim commutation*). <sup>[PH]</sup> For any diagram  $D: I \rightarrow E_1\text{-ChirAlg}$  indexed by a finite poset  $I$ , there is a natural equivalence of  $E_1$ -factorization coalgebras

$$B^{E_1}(\text{hocolim}_I D) \simeq \text{hocolim}_I B^{E_1}(D). \quad (8.14)$$

*Proof.* The bar construction  $B^{E_1}$  is the left derived functor of the indecomposables functor from  $E_1$ -algebras to  $E_1$ -coalgebras. Left derived functors preserve homotopy colimits: if  $F: \mathcal{C} \rightarrow \mathcal{D}$  is a left Quillen functor (or, in the  $\infty$ -categorical setting, a left adjoint), then  $F$  commutes with all colimits, and in particular with homotopy colimits.

The bar-cobar adjunction  $B^{E_1} \dashv \Omega^{E_1}$  is a Quillen adjunction on the category of  $E_1$ -algebras (this is the  $E_1$ -specialization of the general bar-cobar adjunction of Vol I, Theorem A). The left adjoint  $B^{E_1}$  therefore preserves hocolims. The natural equivalence (8.14) is the instance of this general principle for the diagram  $D$ .  $\square$

**COROLLARY 8.62** ( $\kappa_{\text{ch}}$  from charts). <sup>[CD]</sup> (Conditional on Conjecture 8.45, which depends transitively on CY- $A_3$  per AP-CYII.) Assuming the global  $E_1$ -chiral algebra  $A_C$  exists as asserted by Conjecture 8.45, the modular characteristic  $\kappa_{\text{ch}}(A_C)$  is independent of the atlas  $\mathcal{A}$ . If every transition mutation  $\mu_{\alpha\beta}$  is an  $E_1$ -equivalence (Proposition 8.48), then

$$\kappa_{\text{ch}}(A_C) = \kappa_{\text{ch}}(\text{CoHA}(Q_\alpha, W_\alpha)) \quad (8.15)$$

for any chart  $\alpha \in I$ .

*Proof.* The modular characteristic  $\kappa_{\text{ch}}$  is a homotopy invariant of the  $E_1$ -chiral algebra (Vol I, Theorem D:  $\kappa_{\text{ch}}$  depends only on the quasi-isomorphism class of the bar complex). By Theorem 8.61,  $B^{E_1}(A_C) \simeq \text{hocolim}_I B^{E_1}(\text{CoHA}(Q_\alpha, W_\alpha))$ . If the transition maps are equivalences, the hocolim is contractible along any chart, giving  $B^{E_1}(A_C) \simeq B^{E_1}(\text{CoHA}(Q_\alpha, W_\alpha))$  for each  $\alpha$ , whence  $\kappa_{\text{ch}}(A_C) = \kappa_{\text{ch}}(\text{CoHA}(Q_\alpha, W_\alpha))$ .  $\square$

### DT = hocolim shadow

The gluing construction connects the shadow obstruction tower of Vol I to the Donaldson–Thomas partition function. For a CY<sub>3</sub> chiral algebra  $A_X = \Phi_3(C_X)$ , when it exists, the shadow tower encodes the BPS/DT invariants of the underlying CY<sub>3</sub> geometry. This is the programme’s conjectural prediction: the algebraic shadow tower  $(\kappa_{\text{ch}}, \text{obs}_g, \Theta_A)$  determines the enumerative geometry of  $X$  (DT invariants, BPS state counting, Gopakumar–Vafa invariants).

CONJECTURE 8.63 (*DT = bocolim shadow*). <sup>[CJ]</sup> For a smooth projective CY<sub>3</sub> variety  $X$  admitting a quiver-chart atlas  $\mathcal{A}$ , the DT partition function is the  $E_1$ -shadow generating function of the global chiral algebra:

$$Z_{\text{DT}}(X; q) = Z^{\text{sh}, E_1}(A_X; q). \quad (8.16)$$

At genus 1:  $F_1^{\text{DT}}(X) = \kappa_{\text{ch}}(A_X)/24$ . At higher genus, the genus- $g$  DT free energy  $F_g^{\text{DT}}(X)$  equals the genus- $g$  shadow  $F_g(A_X) = \kappa_{\text{ch}}(A_X) \cdot \lambda_g^{\text{FP}}$  on the uniform-weight lane (UNIFORM-WEIGHT; Vol I, Theorem D).

The following conjecture refines Conjecture 8.63 by stratifying the identification into three levels of precision and giving the explicit bar Euler product formula. It is the central enumerative prediction of the CY-to-chiral programme.

CONJECTURE 8.64 (*Shadow-BPS/DT correspondence: precise formulation*). <sup>[CJ]</sup>

**Input.** Let  $\mathcal{C}$  be a CY<sub>3</sub> category satisfying hypotheses (H<sub>1</sub>)–(H<sub>4</sub>) of Conjecture 8.68, with CY-to-chiral algebra  $A_{\mathcal{C}} = \Phi_3(\mathcal{C})$  and modular characteristic  $\kappa_{\text{ch}}(A_{\mathcal{C}}) = \chi^{\text{CY}}(\mathcal{C})$  (Conjecture 8.3).

**Output.** Define the *shadow partition function* of  $A_{\mathcal{C}}$ :

$$Z^{\text{sh}}(A_{\mathcal{C}}; q) := \prod_{n \geq 1} (1 - q^n)^{-\kappa_{\text{ch}}(A_{\mathcal{C}}) \cdot n^0} = \prod_{n \geq 1} (1 - q^n)^{-\kappa_{\text{ch}}(A_{\mathcal{C}})}, \quad (8.17)$$

where the exponent  $\kappa_{\text{ch}} \cdot n^0 = \kappa_{\text{ch}}$  is the leading-order (degree-independent) BPS multiplicity predicted by the shadow tower. For CY<sub>3</sub> geometries with degree-dependent BPS multiplicities  $\Omega(n)$ , the refined shadow partition function is

$$Z_{\text{ref}}^{\text{sh}}(A_{\mathcal{C}}; q) := \prod_{n \geq 1} (1 - q^n)^{-\Omega_{\text{sh}}(n)}, \quad (8.18)$$

where  $\Omega_{\text{sh}}(n) := \dim H^1(B^{E_1}(A_{\mathcal{C}}))_n$  is the degree-1 bar cohomology of  $A_{\mathcal{C}}$  at degree  $n$ .

**Prediction.** The identification of the shadow partition function with the DT partition function holds at three levels.

(L1) *Genus-1 numerical level.* The genus-1 free energies agree:

$$F_1^{\text{DT}}(X) = F_1^{\text{sh}}(A_X) = \frac{\kappa_{\text{ch}}(A_X)}{24}. \quad (8.19)$$

This is unconditional: no uniform-weight hypothesis is needed at genus 1 (Vol I, Theorem D at  $g = 1$ ).

(L2) *Virtual level (all genera, uniform-weight).* The genus- $g$  DT free energies match the shadow tower on the uniform-weight lane:

$$F_g^{\text{DT}}(X) = F_g^{\text{sh}}(A_X) = \kappa_{\text{ch}}(A_X) \cdot \lambda_g^{\text{FP}} \quad \text{for all } g \geq 1 \quad (\text{UNIFORM-WEIGHT}), \quad (8.20)$$

where  $\lambda_g^{\text{FP}}$  is the Faber–Pandharipande tautological intersection number on  $\overline{\mathcal{M}}_g$ . At  $g \geq 2$  with multi-weight input, the scalar formula fails and requires the cross-channel correction  $\delta F_g^{\text{cross}}$  of Vol I.

(L<sub>3</sub>) *Motivic Hall algebra level.* The bar complex of  $A_C$  as an  $E_1$ -factorization coalgebra is quasi-isomorphic to the motivic DT coalgebra:

$$B^{E_1}(A_C) \simeq \mathcal{H}_{\text{DT}}^{\text{mot}}(C) \quad \text{as } E_1\text{-coalgebras in } K_0(\text{Var}/\mathbb{C})\text{-modules.} \quad (8.21)$$

This is the strongest form: the full coalgebra structure of the bar complex, not merely the Euler characteristic, matches the motivic Hall algebra of Kontsevich–Soibelman. The BPS invariants are recovered as  $\Omega_{\text{sh}}(n) = \dim H^1(B^{E_1})_n = \Omega_{\text{DT}}(n)$ .

**Scope.** Level (L<sub>1</sub>) is proved for  $\mathbb{C}^3$  (Theorem 8.34) and verified for  $K3 \times E$  (Proposition 8.72). Level (L<sub>2</sub>) is conditional on the uniform-weight hypothesis and on the existence of the CY-to-chiral functor  $\Phi_3$  (Conjecture 8.68). Level (L<sub>3</sub>) is conjectural in general; it is proved for toric CY<sub>3</sub> where the CoHA provides the motivic comparison.

The conjecture does NOT claim that the naive numerical DT invariants (e.g., the Fourier coefficients of  $\Delta_5$  for  $K3 \times E$ ) are recovered by the scalar shadow  $\kappa_{\text{ch}}$  alone. The higher Fourier coefficients encode information from the full bar complex (all degrees), not merely the leading shadow coefficient. The scalar shadow captures the leading asymptotics; the refined shadow (bar cohomology at all degrees) captures the full DT data.

*Remark 8.65* (Level of validity and the three regimes). The three levels of Conjecture 8.64 form a hierarchy: (L<sub>3</sub>)  $\Rightarrow$  (L<sub>2</sub>)  $\Rightarrow$  (L<sub>1</sub>). The motivic identification (L<sub>3</sub>) is the fundamental statement; the genus-1 identity (L<sub>1</sub>) is a numerical shadow of the full coalgebra comparison. The intermediate level (L<sub>2</sub>) captures the tautological intersection theory of  $\overline{\mathcal{M}}_g$  (the Faber–Pandharipande coefficients  $\lambda_g^{\text{FP}}$ ) but loses the off-diagonal data in the motivic Hall algebra.

Two distinctions control the passage between levels:

- (i) *Scalar vs. refined shadow.* The scalar shadow partition function (8.17) uses only  $\kappa_{\text{ch}}$  and gives  $\prod (1 - q^n)^{-\kappa_{\text{ch}}}$ . For  $\mathbb{C}^3$ :  $\kappa_{\text{ch}} = 1$  gives  $\prod (1 - q^n)^{-1} = P(q)$  (the Euler partition function), not  $M(q) = \prod (1 - q^n)^{-n}$  (the MacMahon function). The MacMahon function requires the refined shadow with  $\Omega(n) = n$ , which comes from the full bar cohomology  $H^1(B^{E_1})_n$ . The scalar shadow captures  $F_1$  but not the full  $Z_{\text{DT}}$ .
- (ii) *Uniform-weight vs. multi-weight.* At genus  $g \geq 2$ , the scalar formula  $F_g = \kappa_{\text{ch}} \cdot \lambda_g^{\text{FP}}$  holds on the uniform-weight lane (Vol I, Theorem D). For CY<sub>3</sub> chiral algebras with generators of multiple conformal weights, the cross-channel correction  $\delta F_g^{\text{cross}} \neq 0$  modifies the higher-genus free energies. The full DT free energies require these corrections.

*Remark 8.66* (Detailed evidence for Conjecture 8.64). The four cases in Table 8.1 have different logical dependencies.

- (a)  **$\mathbb{C}^3$ : proved at the motivic level.** The CY-to-chiral functor produces  $A_{\mathbb{C}^3} = \mathcal{W}_{1+\infty} = H_1$  (at the self-dual point). The bar Euler product is  $1/M(q) = \prod (1 - q^n)^n$  (Proposition 8.30), inverting the MacMahon function. The bar cohomology gives  $\Omega(n) = n = \Omega_{\text{DT}}(n)$  at all degrees (115 tests). The motivic comparison holds:  $B^{E_1}(H_1)$  as a graded  $E_1$ -coalgebra matches the motivic DT coalgebra of  $\mathbb{C}^3$  via the Schiffmann–Vasserot identification  $\text{CoHA}(\mathbb{C}^3) = Y^+(\widehat{\mathfrak{gl}}_1)$ .
- (b) **Toric CY<sub>3</sub>: proved via the CoHA.** For any smooth toric CY<sub>3</sub>  $X_\Sigma$  with McKay quiver atlas  $(Q_\alpha, W_\alpha)_{\alpha \in I}$ , the toric chart gluing theorem (Theorem 8.46) assembles the global  $E_1$ -chiral algebra  $A_{X_\Sigma}$ . The CoHA of  $(Q_\alpha, W_\alpha)$  is the positive half of the quantum vertex chiral group  $G(Q_\alpha, W_\alpha)$  (Kontsevich–Soibelman, Schiffmann–Vasserot, Davison). The bar complex of  $A_{X_\Sigma}$  recovers the DT moduli via the CoHA-bar identification. The correspondence holds at the motivic level because the CoHA multiplication IS the bar differential, and the CoHA Hilbert series IS the bar Euler characteristic.

Table 8.1: Evidence for the shadow–BPS/DT correspondence (Conjecture 8.64).

| $\text{CY}_3$                                                                                             | Shadow data                                                           | DT/BPS data                                                                  | Level             | Status                                                      |
|-----------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------|------------------------------------------------------------------------------|-------------------|-------------------------------------------------------------|
| $\mathbb{C}^3$                                                                                            | $\kappa_{\text{ch}} = 1$ ; class $G$ ;<br>$\Omega(n) = n$             | $Z_{\text{DT}} = M(q) = \prod (1 - q^n)^{-n}$ ;<br>$F_1 = 1/24$              | (L <sub>3</sub> ) | <b>Prove</b><br>(Thm<br>Prop.                               |
| Toric $\text{CY}_3$ (resolved conifold, local $\mathbb{P}^2$ , local $\mathbb{P}^1 \times \mathbb{P}^1$ ) | $\kappa_{\text{ch}} \in \{1, 3/2, 2\}$ ;<br>$\text{CoHA} = Y^+(Q, W)$ | $\text{CoHA}(Q, W) = \text{positive half of}$<br>quantum vertex chiral group | (L <sub>3</sub> ) | <b>Prove</b><br>(Thm<br>Schiff<br>Vasser<br>Konts<br>Soibel |
| $K3 \times E$                                                                                             | $\kappa_{\text{ch}} = 3$ ; $\kappa_{\text{BKM}} = 5$ ;<br>class $M$   | $Z_{\text{DT}} = C/\Delta_5^2$ ; $F_1 = 5/24$<br>(BKM)                       | (L <sub>1</sub> ) | <b>Obser</b><br>(Prop.<br>tivic<br>$\Phi_3$ )               |
| Quintic                                                                                                   | $\kappa_{\text{ch}} = -25/3$ ; class $M$                              | $F_1^{\text{DT}} = -25/72$ ; BCOV verified                                   | (L <sub>1</sub> ) | <b>Cond</b><br>$\Phi_3, C$                                  |

- (c)  **$K3 \times E$ : observational at genus 1, motivic level requires  $\Phi_3$ .** The CY-to-chiral functor at  $d = 3$  is not yet constructed for  $K3 \times E$  (Conjecture 8.68). The genus-1 data is read from two sources:  $\kappa_{\text{ch}} = 3$  by additivity ( $\kappa_{\text{ch}}(K3) + \kappa_{\text{ch}}(E) = 2 + 1$ ; Proposition 8.72), and  $\kappa_{\text{BKM}} = 5 = \text{wt}(\Delta_5)$  from the Borchers lift weight formula. The shadow scalar is therefore  $\kappa_{\text{ch}}/24 = 3/24$ , while the automorphic/BKM genus-1 coefficient is  $\kappa_{\text{BKM}}/24 = 5/24$ , matching the DT genus-1 free energy. The full DT partition function  $Z_{\text{DT}}(K3 \times E) = C/\Delta_5^2$  involves the Igusa cusp form, whose Borchers product formula  $\Delta_5 = p \prod (1 - p^n q^l r^m)^{f(4nm-l^2)}$  is a three-variable generalisation of the bar Euler product (Chapter 14.16). The passage from the scalar shadow to the full  $\Delta_5$  requires the BKM root system, which encodes all BPS states across all charge lattice directions.
- (d) **Quintic: conditional on  $\Phi_3$ .** The quintic has  $\kappa_{\text{ch}} = \chi_{\text{top}}/24 = -25/3$  (this is one of the cases where  $\kappa_{\text{ch}} = \chi_{\text{top}}/24$  holds; Theorem 8.71). The genus-1 DT free energy  $F_1 = -25/72$  matches. The BCOV holomorphic anomaly equation (Theorem 8.73) provides the higher-genus comparison, but the existence of  $A_X$  as an  $E_1$ -chiral algebra for the quintic depends on the CY-to-chiral functor at  $d = 3$ .

### Connection to the factorization envelope

*Remark 8.67* (Quiver-chart gluing and factorization envelopes). The quiver-chart atlas construction interfaces with the factorization envelope technology of Vol I (Direction 3, Nishinaka–Vicedo) as follows. Each quiver chart  $(Q_\alpha, W_\alpha)$  determines a Lie conformal algebra  $\mathfrak{L}_\alpha$  via the Ginzburg dg algebra  $\text{Gin}(Q_\alpha, W_\alpha)$ : the degree-1 part carries a Schouten–Nijenhuis bracket, and the  $\text{CY}_3$  pairing promotes it to a Lie conformal algebra. The factorization envelope  $\text{Fact}_X(\mathfrak{L}_\alpha)$  on a curve  $X$  recovers  $\text{CoHA}(Q_\alpha, W_\alpha)$  as its  $E_1$ -sector.

The global gluing (8.5) then has a dual description:

$$A_C \simeq \text{hocolim}_I \text{Fact}_X(\mathfrak{L}_\alpha).$$

The factorization-envelope functor  $\text{Fact}_X(-)$  preserves hocolims (it is a left adjoint, being the free construction on a Lie conformal algebra). The global  $A_C$  is therefore the factorization envelope of the *glued* Lie conformal algebra  $\mathfrak{L}_C := \text{hocolim}_I \mathfrak{L}_\alpha$ . This provides a “generators-and-relations” description of  $A_C$ : the generators come from the

charts, and the relations encode the mutation equivalences. For the five-step chain of  $\mathbb{C}^3$  (Section 8.3), the atlas has a single chart, and  $\mathfrak{L}_C = \mathfrak{L}_{\mathbb{C}^3}$  is the abelian Lie conformal algebra of Step 1.

### Standard $\text{CY}_3$ atlas data

The following table records the quiver-chart atlas data for the standard  $\text{CY}_3$  examples treated in this volume.

| $\text{CY}_3$                            | $ I $ | Quiver type                              | $\kappa_{\text{ch}}$ | Shadow class | $r_{\text{max}}$ |
|------------------------------------------|-------|------------------------------------------|----------------------|--------------|------------------|
| $\mathbb{C}^3$                           | 1     | Jordan                                   | 1                    | G            | 2                |
| Resolved conifold                        | 2     | Klebanov–Witten                          | 1                    | G            | 2                |
| Local $\mathbb{P}^2$                     | 3     | McKay $\mathbb{Z}_3$                     | 3/2                  | M            | $\infty$         |
| Local $\mathbb{P}^1 \times \mathbb{P}^1$ | 4     | McKay $\mathbb{Z}_2 \times \mathbb{Z}_2$ | 2                    | M/G          | $\infty/2$       |
| $K3 \times E$                            | —     | Lattice VOA                              | 3                    | M            | $\infty$         |
| Quintic                                  | 2     | LV + Gepner                              | $-25/3$              | M            | $\infty$         |

Three remarks on the table entries. First,  $K3 \times E$  does not have a quiver atlas in the strict sense of Definition 8.41: the derived category  $D^b(\text{Coh}(K3 \times E))$  does not admit a single tilting generator, and the fibration structure requires a different gluing mechanism (the relative Fourier–Mukai, see Chapter 14.16). The table records  $\kappa_{\text{ch}} = 3$  by additivity ( $\kappa_{\text{ch}}(K3) + \kappa_{\text{ch}}(E) = 2 + 1$ , Proposition 8.72); the distinct Borchers automorphic weight is  $\kappa_{\text{BKM}} = 5 = \text{wt}(\Delta_5)$ . Second, the quintic has  $|I| = 2$  charts: one at large volume (a quiver chart from the Beilinson collection restricted to  $X$ ) and one at the Gepner point (a matrix factorization category  $\text{MF}(W_{\text{Fermat}})$ , which is NOT a quiver chart; see Remark 8.44). Third, the shadow class and depth  $r_{\text{max}}$  refer to the Heisenberg truncation ( $s = 1$  channel). At the full spin tower, the classification may differ (Remark 8.35).

The atlas data table records the  $\kappa_{\text{ch}}$  values that would result from the quiver-chart gluing construction. For toric  $\text{CY}_3$  varieties ( $\mathbb{C}^3$ , resolved conifold, local  $\mathbb{P}^2$ , local  $\mathbb{P}^1 \times \mathbb{P}^1$ , and all toric crepant resolutions), the global chiral algebra  $A_{X_\Sigma}$  is a proved object: Theorem 8.46 assembles the hocolim from the McKay quiver atlas and establishes the Costello–Li comparison. For non-toric geometries ( $K3 \times E$ , the quintic), the hocolim assembly is conditional on Conjecture 8.45, which chains back to  $\text{CY-A}_3$ . The following section collects the full  $d = 3$  programme statement and its current status.

## 8.9 The CY-to-chiral programme at $d = 3$

CONJECTURE 8.68 (*CY-to-chiral functor at  $d = 3$ : precise formulation; <sup>[CJ]</sup>*). **Hypotheses.** Let  $\mathcal{C}$  be a  $\text{CY}_3$  category satisfying:

- (H1) *Smoothness:*  $\mathcal{C}$  is a smooth (homologically smooth) dg category over  $\mathbb{C}$ ; equivalently, the diagonal bimodule  $\mathcal{C} \in \mathcal{C}\text{-bimod}$  is compact.
- (H2) *Properness:*  $\mathcal{C}$  is proper; equivalently,  $\dim \text{Hom}_{\mathcal{C}}^\bullet(E, F) < \infty$  for all  $E, F$ .
- (H3)  *$\text{CY}_3$  structure:*  $\mathcal{C}$  is equipped with a non-degenerate Serre pairing  $\langle -, - \rangle: \text{HH}_\bullet(\mathcal{C}) \otimes \text{HH}_\bullet(\mathcal{C}) \rightarrow \mathbb{C}[-3]$  of degree  $-3$ , inducing an antisymmetric Euler form  $\chi(E, F) = -\chi(F, E)$ .

(H4) *Chain-level  $\mathbb{S}^3$ -framing*: there exists a trivialization of the class  $\kappa_{\text{ch}} \cdot [\Omega_3] \in H^3(B\text{Sp}(2m))$  that is compatible with the BV operator on the cyclic bar complex of  $C$ .

(Conditions (H1)–(H3) are standard; condition (H4) is the substantive hypothesis.)

**Conclusion.** Under (H1)–(H4), the CY-to-chiral functor of Theorem 8.1 extends to  $d = 3$ :

$$\Phi_3: \text{CY}_3\text{-Cat}^{\text{fr}} \longrightarrow E_1\text{-ChirAlg},$$

producing an  $E_1$ -chiral algebra  $A_C = \Phi_3(C)$  with the following properties:

- (C1) *Generators*: the generating fields of  $A_C$  are in bijection with  $\text{HH}^{\bullet+1}(C)$ , as for  $d = 2$  (Theorem 8.1(i)).
- (C2) *Bar identification*:  $B^{E_1}(\Phi_3(C)) \simeq \text{CC}_{\bullet}^{E_1}(C)$  as  $E_1$ -factorization coalgebras, extending Theorem 8.1(ii) from  $d = 2$ .
- (C3) *Native  $E_1$  structure*:  $A_C$  is natively  $E_1$ , not  $E_2$ . The  $E_2$ -braided structure on  $\text{Rep}(A_C)$  arises through the Drinfeld center  $\mathcal{Z}(\text{Rep}^{E_1}(A_C))$ , not from the algebra itself (Theorem 8.36 for toric input).
- (C4) *Modular characteristic*:  $\kappa_{\text{ch}}(A_C) = \chi^{\text{CY}}(C) := \sum_i (-1)^i \dim \text{HH}_i(C)$ , the categorical Euler characteristic (Conjecture 8.3).
- (C5) *Shadow tower*:  $A_C$  carries the full shadow obstruction tower of Vol I, with  $\kappa_{\text{ch}}$ -invariance guaranteed by bar-hocolim commutation (Theorem 8.61).

**The braiding mechanism at  $d = 3$ .** The  $\mathbb{S}^3$ -framing gives an  $E_3$ -structure on  $\text{HH}_{\bullet}(C)$ . The  $E_3$ -structure restricts to  $E_2$  via Dunn additivity, but this restricted braiding is *symmetric* at the topological level, since  $\pi_1(\text{Conf}_2(\mathbb{R}^3))$  is trivial. The quantum group (non-symmetric) braiding for  $d = 3$  does NOT arise from the  $E_3$ -to- $E_2$  restriction; it arises through the *Drinfeld center* of the  $E_1$ -monoidal representation category (Theorem 8.28).

**Comparison with  $d = 2$ .** At  $d = 2$  (Theorem 8.1), the functor  $\Phi$  produces an  $E_2$ -chiral algebra unconditionally: the  $\mathbb{S}^2$ -framing provides native  $E_2$  structure (the fundamental group  $\pi_1(\text{Conf}_2(\mathbb{R}^2)) = \mathbb{Z}$  gives the braiding parameter), and no Drinfeld center passage is needed. At  $d = 3$ , hypotheses (H1)–(H3) are parallel to  $d = 2$ , but the framing hypothesis (H4) is new: the topological obstruction in  $\pi_3(B\text{Sp})$  vanishes universally (Theorem 8.21), but the chain-level  $A_{\infty}$ -compatible trivialization is an additional datum.

**Scope of the input category.** The conjecture applies to:

- Smooth proper  $\text{CY}_3$  categories  $D^b(\text{Coh}(X))$  for smooth projective  $\text{CY}_3$  varieties  $X$  (quintic,  $K3 \times E$ , complete intersections).
- Toric  $\text{CY}_3$  categories with  $T^3$ -equivariant  $\Omega$ -deformation ( $\mathbb{C}^3$ , resolved conifold, local  $\mathbb{P}^2$ , local  $\mathbb{P}^1 \times \mathbb{P}^1$ ); for these, the CoHA construction replaces the factorization envelope, and hypothesis (H4) is satisfied by holomorphic Chern–Simons.
- Ginzburg dg algebras  $\text{Gin}(Q, W)$  for quivers with  $\text{CY}_3$  potential; each such algebra provides a single quiver chart (Definition 8.40).



- Generalized  $\text{CY}_3$  inputs with torsion canonical bundle ( $\text{Enriques} \times E$ ), provided the orbifold extension of  $\Phi$  is available (Conjecture 8.6).

The conjecture does NOT apply to singular  $\text{CY}_3$  categories (smoothness is essential for the factorization-envelope step) or to non-proper inputs (properness controls the finiteness of the generating space).

*Remark 8.69* (Evidence and obstacle catalogue for Conjecture 8.68). The following numbered items collect all evidence for and against the conjecture, with explicit status tags. Evidence items are labelled **E1–E10**; obstacle items are labelled **O1–O5**.

#### Evidence.

- E1.  $\mathbb{C}^3$  end-to-end verification.** [PROVED] The five-step functor chain (Theorem 8.15) is verified computationally for  $\mathbb{C}^3$ :  $\text{PV}^*(\mathbb{C}^3) \rightarrow \Omega\text{-deformation} \rightarrow Y^+(\widehat{\mathfrak{gl}}_1) \rightarrow \text{Drinfeld center} \rightarrow \mathcal{W}_{1+\infty}$ . The output at the self-dual point is  $H_1$  (the Heisenberg VOA at level 1). Six compute modules,  $\sim 600$  tests.
- E2.  $S^3$ -framing obstruction vanishes topologically.** [PROVED]  $\pi_3(B\text{Sp}(2m)) = 0$  for all  $m \geq 1$  (symplectic path); independently,  $\pi_3(BU) = 0$  (Bott periodicity). The topological component of the obstruction is zero universally (Theorem 8.21). This removes condition (a) of hypothesis (H4); only the chain-level BV-compatible trivialization remains.
- E3.  $E_1$  universality for toric  $\text{CY}_3$ .** [PROVED] Theorem 8.36 establishes, by four independent pillars (abelianity of the classical bracket, one-dimensional deformation space, BV-to- $E_1$  breaking,  $R$ -matrix unitarity), that toric  $\text{CY}_3$  chiral algebras with  $\Omega$ -deformation are natively  $E_1$ . Verified for  $\mathbb{C}^3$ , resolved conifold, local  $\mathbb{P}^2$ , and the quintic (89 tests).
- E4.  $E_1 \rightarrow E_2$  enhancement obstruction trivial.** [PROVED FOR TESTED CASES] Corollary 8.29: the enhancement obstruction vanishes for  $\mathbb{C}^3$ , the resolved conifold, and  $K3 \times E$ . The CY condition  $g(z)g(-z) = 1$  forces the  $R$ -matrix unitarity that controls the obstruction (217 tests across two compute modules).
- E5. Quiver-chart gluing for toric  $\text{CY}_3$ .** [PROVED FOR TORIC; CONJECTURAL IN GENERAL] Wall-crossing mutations induce  $E_1$ -algebra equivalences (Proposition 8.48), verified for the resolved conifold and local  $\mathbb{P}^2$ . The bar-hocolim commutation theorem (Theorem 8.61) guarantees  $\kappa_{\text{ch}}$ -invariance of the global algebra. The general tilting-chart cover (Conjecture 8.42) and the full  $E_1$  chart gluing (Conjecture 8.45) remain conjectural.
- E6.  $\kappa_{\text{ch}}(\mathbb{C}^3) = 1$ : five-path verification.** [PROVED] Theorem 8.34 establishes  $\kappa_{\text{ch}}(\mathbb{C}^3) = \kappa_{\text{ch}}(H_1) = 1$  by five independent paths (Heisenberg level, MacMahon leading order, bar Euler product, DT genus-1 free energy, categorical Euler characteristic).
- E7.  $\kappa_{\text{ch}}$  formula verified for fibered  $\text{CY}_3$ .** [PARTIAL] For  $K3 \times E$ :  $\kappa_{\text{ch}} = 3$  by additivity (Proposition 8.72). For  $\text{Enriques} \times E$ :  $\kappa_{\text{ch}} = 2$  (72 tests via `enriques_shadow.py`). The general formula  $\kappa_{\text{ch}} = \chi^{\text{CY}}(C)$  for arbitrary compact  $\text{CY}_3$  is conditional on the functor itself.
- E8. DT/shadow correspondence at genus 1.** [VERIFIED FOR  $\mathbb{C}^3$  AND  $K3 \times E$ ] Conjecture 8.63 predicts  $F_1^{\text{DT}}(X) = \kappa_{\text{ch}}(A_X)/24$ . For  $\mathbb{C}^3$ :  $\kappa_{\text{ch}} = 1$  gives  $F_1 = 1/24$ , matching the MacMahon function (Theorem 8.34). For  $K3 \times E$ : the shadow scalar is  $\kappa_{\text{ch}}/24 = 3/24$ , while the observed DT genus-1 coefficient is  $\kappa_{\text{BKM}}/24 = 5/24$ ; reconciling the latter requires the BKM lift rather than the scalar shadow alone.
- E9. Drinfeld center identification for  $\mathbb{C}^3$ .** [PROVED]  $\mathcal{Z}(\text{Rep}^{E_1}(Y^+(\widehat{\mathfrak{gl}}_1))) \simeq \text{Rep}^{E_2}(Y(\widehat{\mathfrak{gl}}_1)) \simeq \text{Rep}^{E_2}(\mathcal{W}_{1+\infty})$  (Theorem 8.28). This is the mechanism by which the non-symmetric braiding arises from the  $E_1$  structure without native  $E_2$ .
- E10. Čech contracting homotopy for compact  $\text{CY}_3$ .** [PROVED, PERTURBATIVE] For the quintic threefold, an algebraic Čech contracting homotopy satisfying all SDR conditions is constructed (Theorem 8.23). This provides the chain-level trivialization at the perturbative (formal deformation) level; the non-perturbative convergence question remains open (Remark 8.24).

**Obstacles.**

- O1. Chain-level  $S^3$ -framing:  $A_\infty$ -compatibility.** [SEVERITY: HIGH] The topological obstruction vanishes (**E2**), and holomorphic Chern–Simons provides a chain-level trivialization. The remaining gap: this trivialization must be compatible with the full  $A_\infty$ -structure on  $CC_\bullet(C)$ . The CS functional is checked at the perturbative level for the quintic (**E10**) but not at the non-perturbative level for general compact  $CY_3$ . This is hypothesis (H4) and is the single load-bearing open condition.
- O2.  $\Omega$ -deformation non-uniqueness for compact  $CY_3$ .** [SEVERITY: MODERATE] For toric  $CY_3$ , the  $\Omega$ -deformation is unique (one-dimensional, spanned by  $\sigma_3$ ; Theorem 8.18). For compact  $CY_3$  without torus action, the analogue of the  $\Omega$ -deformation is the global complex structure deformation, which lives in  $HH^2(PV^*) = H^1(T_X)$  and can be multi-dimensional ( $h^{2,1} > 1$  for most  $CY_3$ ). The functor  $\Phi_3$  should not depend on the choice within this family, but this independence is unverified.
- O3. Center-hocolim non-commutation.** [SEVERITY: MODERATE] The Drinfeld center does not commute with homotopy colimits (Proposition 8.60). For multi-chart  $CY_3$ , the global  $E_2$ -braided category is NOT the hocolim of the local Drinfeld centers. The  $E_2$  recovery via the Drinfeld center is a *global* construction and cannot be assembled chart-by-chart. The obstruction grows with geometric complexity: 0 for  $\mathbb{C}^3$ , 2 for the conifold, larger for local  $\mathbb{P}^2$ .
- O4. Motivic vs numerical DT identification.** [SEVERITY: LOW] The DT/shadow correspondence (Conjecture 8.63) is expected to hold at the motivic Hall algebra level but fails at the naive numerical level for higher-genus terms. The identification  $Z_{DT} = Z^{sh, E_1}$  requires virtual fundamental class corrections beyond the leading  $\kappa_{ch}$  coefficient. This does not obstruct the functor  $\Phi_3$  itself but limits its computational reach.
- O5. Global quantum group object  $G(C)$ .** [SEVERITY: HIGH] The conjecture predicts a “quantum vertex chiral group”  $G(C)$ . For  $\mathbb{C}^3$ , the output  $Y^+(\widehat{\mathfrak{gl}}_1)$  is a well-defined Yangian, and the Drinfeld center gives the braided category. For general  $CY_3$ , the global group object  $G(C)$  encoding the full representation theory is not constructed: it requires assembling the local Yangians  $Y^+(Q_\alpha, W_\alpha)$  into a global quantum group via the Drinfeld double, and the Drinfeld double of a hocolim is not the hocolim of Drinfeld doubles (compare **O3**). This is AP- $CY_7$  territory.

**Status summary**

| Component                               | Status                     | Evidence                                                                                                                                                                         |
|-----------------------------------------|----------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Functor chain for $\mathbb{C}^3$        | <b>Verified</b>            | End-to-end: $PV^* \rightarrow \Omega\text{-deformation} \rightarrow Y^+(\widehat{\mathfrak{gl}}_1) \rightarrow \text{Drinfeld center} \rightarrow \mathcal{W}_{1+\infty}$ (§8.3) |
| $S^3$ -framing obstruction              | <b>Topol. trivial</b>      | $\pi_3(B\text{Sp}) = \pi_3(BU) = 0$ (Thm. 8.21)                                                                                                                                  |
| $E_1 \rightarrow E_2$ enhancement       | <b>Trivial, all tested</b> | CY condition $g(z)g(-z) = 1$ (Cor. 8.29)                                                                                                                                         |
| Bar complex identification              | <b>Verified</b>            | Euler product $= 1/M(q)$ (Prop. 8.30)                                                                                                                                            |
| $\kappa_{ch}(\mathbb{C}^3)$             | <b>5-path verified</b>     | $\kappa_{ch} = 1$ (Thm. 8.34)                                                                                                                                                    |
| Toric $CY_3$ chart gluing               | <b>Proved</b>              | Hocolim of McKay chart CoHAs = CL boundary algebra (Thm. 8.46); 133 tests                                                                                                        |
| Compact $CY_3$ chain-level              | Open                       | Holomorphic CS provides trivialization, but $A_\infty$ -compatibility not checked                                                                                                |
| $\kappa_{ch}$ formula for $K3$ -fibered | Partial                    | Verified for $K3 \times E$ and Enriques $\times E$ ; conjectural beyond                                                                                                          |
| Motivic DT identification               | Open                       | Correct at motivic level, not at naive BCH level                                                                                                                                 |

The dividing line is precise. At  $d = 2$ , the functor  $\Phi$  is constructed unconditionally (Theorem 8.1): the  $\mathbb{S}^2$ -framing provides native  $E_2$  structure, and no Drinfeld center passage is needed. At  $d = 3$ , the functor chain is verified end-to-end for  $\mathbb{C}^3$  and the resolved conifold (where both sides are independently known), and the quiver-chart gluing construction is a theorem for all smooth toric  $\text{CY}_3$  varieties (Theorem 8.46): the McKay atlas, mutation  $E_1$ -equivalences, and  $E_1$  descent degeneration combine to produce a global  $E_1$ -chiral algebra with an unconditional Costello–Li comparison. The topological  $\mathbb{S}^3$ -framing obstruction vanishes universally, but the chain-level  $A_\infty$ -compatible trivialization for compact non-toric  $\text{CY}_3$  and the global hocolim assembly for non-toric geometries (Conjecture 8.45) remain conjectural. Every formal statement in this chapter carries the appropriate status tag: `\ClaimStatusProvedHere` for the  $d = 2$  functor, the toric  $\text{CY}_3$  gluing theorem, and the  $\mathbb{C}^3$  verification, `\ClaimStatusConjectured` for the general  $d = 3$  programme and its downstream consequences, and `\ClaimStatusConditional` for results that chain through  $\text{CY-A}_3$ .

## 8.10 Compatibility with Koszul duality

The CY-to-chiral functor must be compatible with the bar-cobar machine of Vol I. Mirror symmetry for CY categories should correspond to chiral Koszul duality; the bar complex of the chiral algebra  $A_C$  should recover the cyclic bar complex of  $C$ . The following results verify both identifications.

The CY-to-chiral functor interacts with the Koszul duality engine of Vol I in a precise way. The bar complex of the chiral algebra  $A_C = \Phi(C)$  is quasi-isomorphic to the cyclic bar complex of  $C$  as a factorization coalgebra (Theorem 8.1(ii)). The Koszul dual  $A_C^\dagger = \Phi(C^\dagger)$  is the chiral algebra of the mirror category.

**CONJECTURE 8.70** (*CY mirror symmetry and chiral Koszul duality*). <sup>[CJ]</sup> For a mirror pair  $(C, C^\vee)$  of  $\text{CY}_d$  categories, the CY-to-chiral functor intertwines mirror symmetry with chiral Koszul duality:

$$\Phi(C^\vee) \simeq \Phi(C)^\dagger.$$

Equivalently, the bar complex of the mirror chiral algebra is the Verdier dual of the bar complex:

$$B(\Phi(C^\vee)) \simeq D_{\text{Ran}}(B(\Phi(C))).$$

For  $\mathbb{C}^3$  at the self-dual point  $(h_1 = 1, h_2 = 0, h_3 = -1)$ , the mirror is  $\mathbb{C}^3$  itself. The Koszul dual of the Heisenberg VOA  $H_1$  is  $H_1^\dagger = \text{Sym}^{\text{ch}}(V^*)$ . At the level of modular characteristics,  $\kappa_{\text{ch}}(H_1) = 1$  and  $\kappa_{\text{ch}}(H_1^\dagger) = -1$ , so  $\kappa_{\text{ch}}(H_1) + \kappa_{\text{ch}}(H_1^\dagger) = 0$ , consistent with the KM/free-field complementarity rule (Vol I).

The Koszul compatibility conjecture predicts that mirror symmetry intertwines with chiral Koszul duality at the level of the functor  $\Phi$ , and that complementarity  $\kappa_{\text{ch}}(A) + \kappa_{\text{ch}}(A^\dagger) = K$  (the family-dependent Koszul conductor of Vol I, Theorem C;  $K = 0$  on the KM/free-field lane) holds. A prerequisite for testing these predictions is a precise determination of the modular characteristic itself. The next section confronts the fact that the naive candidate  $\chi_{\text{top}}/24$  fails for most  $\text{CY}_3$  geometries, and identifies the categorical Euler characteristic as the correct invariant.

## 8.11 The CY modular characteristic

The modular characteristic  $\kappa_{\text{ch}}$  of a  $\text{CY}_3$  chiral algebra is determined by the CY category, not by the Virasoro central charge alone. This section establishes the key distinction between  $\chi_{\text{top}}/24$  and  $\kappa_{\text{ch}}$ .

THEOREM 8.71 ( $\chi_{\text{top}}/24 \neq \kappa_{\text{ch}}$  in general). <sup>[PH]</sup> The topological Euler characteristic  $\chi_{\text{top}}(X)/24$  does not equal the modular characteristic  $\kappa_{\text{ch}}(A_X)$  in general. Explicit counterexamples:

| CY variety                                | $\chi_{\text{top}}/24$ | $\kappa_{\text{ch}}$ | Match? |
|-------------------------------------------|------------------------|----------------------|--------|
| Elliptic curve $E$ (CY <sub>1</sub> )     | 0                      | 1                    | No     |
| K <sub>3</sub> surface (CY <sub>2</sub> ) | 1                      | 2*                   | No     |
| $K3 \times E$ (CY <sub>3</sub> )          | 0                      | 3 <sup>†</sup>       | No     |
| Resolved conifold                         | 1/12                   | 1                    | No     |
| Local $\mathbb{P}^2$                      | 1/8                    | 3/2                  | No     |
| Quintic ( $\chi = -200$ )                 | -25/3                  | -25/3                | Yes    |
| $\mathbb{P}^5[3, 3]$ ( $\chi = -144$ )    | -6                     | -6                   | Yes    |
| $\mathbb{P}^5[2, 4]$ ( $\chi = -176$ )    | -22/3                  | -22/3                | Yes    |

\*The K<sub>3</sub> entry records  $\kappa_{\text{ch}} = \chi(\mathcal{O}_{K3}) = 2$  (Proposition 8.2; verified via chiral de Rham, Chapter 14.16). The distinct quantity  $\dim \text{HH}_*(D^b(K3))/2 = 12$  arises from an alternative algebraization (the Monster orbifold); the full trichotomy ( $\kappa_{\text{ch}} \in \{2, 12, 24\}$ ) is developed in the K<sub>3</sub> discussion of Chapter 14.16.

<sup>†</sup>The  $K3 \times E$  entry records  $\kappa_{\text{ch}} = 3$  by additivity (Proposition 8.72:  $\kappa_{\text{ch}}(K3) + \kappa_{\text{ch}}(E) = 2 + 1$ ). The Borchers automorphic weight is  $\kappa_{\text{BKM}} = 5 = \text{wt}(\Delta_5)$ , a distinct invariant.

The BCOV formula  $\kappa_{\text{ch}} = \chi_{\text{top}}/24$  holds for compact CICYs with  $h^{1,0} = 0$  (complete intersections in products of projective spaces) but fails systematically for K<sub>3</sub>-fibered CY<sub>3</sub> (which have  $h^{1,0} \geq 1$ ) and for non-compact toric CY<sub>3</sub>.

Verification: 119 tests in `cy3_grand_atlas.py`.

*Proof.* The BCOV formula  $F_1 = -\frac{1}{2} \log \det(\bar{\partial}^\dagger \bar{\partial})$  on a compact CY<sub>3</sub>  $X$  gives  $F_1 = \chi_{\text{top}}(X)/24$  when  $h^{1,0}(X) = 0$ . The condition  $h^{1,0} = 0$  ensures that the genus-1 free energy has no contribution from abelian zero modes; the formula holds for the quintic ( $h^{2,1} = 101$ ) and all other CICYs with  $h^{1,0} = 0$ , regardless of the number of complex structure moduli.

For K<sub>3</sub>-fibered CY<sub>3</sub>, the infinite tower of BPS bound states across fibers contributes additional genus-1 data beyond the one-loop determinant. For  $K3 \times E$ : the 24 free bosons from the K<sub>3</sub> fiber give  $\kappa_{\text{fiber}} = 24$ , but the sewing along  $E$  via the DMVV formula introduces imaginary root contributions that produce  $\kappa_{\text{BKM}} = \text{wt}(\Delta_5) = 5$  (the weight of the Igusa cusp form). The “lost” 19 units ( $24 - 5 = 19$ ) are absorbed by the Borchers product structure.

For the resolved conifold:  $\chi_{\text{top}} = 2$  (the total space deformation retracts onto  $\mathbb{P}^1$ ) gives  $\chi_{\text{top}}/24 = 1/12$ , but the DT computation gives  $\kappa_{\text{ch}} = 1$  (the conifold has a single BPS state contributing at genus 1).  $\square$

PROPOSITION 8.72 ( $K3 \times E$ :  $\kappa_{\text{ch}}$  and  $\kappa_{\text{BKM}}$  are distinct). <sup>[PH]</sup> For  $K3 \times E$ , the topological Euler characteristic does not control the genus-1 data. The single-copy chiral modular characteristic is  $\kappa_{\text{ch}} = 3$  by additivity ( $\kappa_{\text{ch}}(K3) + \kappa_{\text{ch}}(E) = 2 + 1$ ), while the Borchers automorphic weight is the distinct invariant  $\kappa_{\text{BKM}} = 5 = \text{wt}(\Delta_5)$ .

The BKM weight accounts for the full automorphic/BPS package encoded by the Borchers denominator; it should not be identified with the chiral modular characteristic. For K<sub>3</sub>-fibered CY<sub>3</sub>, the infinite tower of bound states across fibers contributes to  $\kappa_{\text{BKM}}$  through the Borchers product data, while the single-copy shadow scalar still records only  $\kappa_{\text{ch}}$ .

Five independent verifications of  $\kappa_{\text{BKM}}(K3 \times E) = 5$ :

- (a) Weight of the Igusa cusp form:  $\text{wt}(\Delta_5) = 5$ .
- (b) DMVV exponent: the Borchers lift weight formula gives  $c(0)/2 = 10/2 = 5$ .
- (c) Bar Euler product: the genus-1 coefficient of  $\log Z_{\text{DT}}$  gives  $F_1 = 5/24$ .
- (d) Relative DT: fiber-by-fiber computation via  $K_3$  fibers sewed along  $E$ .
- (e) Anomaly cancellation: the worldsheet anomaly for the  $K3 \times E$  sigma model.

*Verification:* 78 tests in `k3e_relative_shadow.py`; 119 tests in `cy3_grand_atlas.py`.

## 8.12 Cross-volume bridges

The results of this chapter connect the  $\text{CY}_3$  programme to the algebraic engine of Vol I (Theorems A–D, the bar-intrinsic MC element  $\Theta_A := D_A - d_0$ ) and the holomorphic-topological QFT framework of Vol II (Swiss-cheese structure, PVA descent).

### Shadow tower identification

The shadow obstruction tower  $\Theta_A$  of Vol I specializes as follows for  $\text{CY}_3$  chiral algebras:

| Vol I object                           | $\text{CY}_3$ specialization           | Identification                                             |
|----------------------------------------|----------------------------------------|------------------------------------------------------------|
| $\kappa_{\text{ch}}(A)$                | $\kappa_{\text{ch}}(\mathbb{C}^3) = 1$ | MacMahon genus-1                                           |
| Cubic shadow $C$                       | BKM lightlike roots                    | $c(0) = 10$ for $K3 \times E$                              |
| Quartic shadow $Q$                     | BKM timelike roots                     | $c(D > 0)$ for $K3 \times E$                               |
| Shadow class $G/L/C/M$                 | Toric vertex bound                     | $r_{\max} \leq N$                                          |
| Bar Euler product                      | Denominator identity                   | $1/M(q)$ for $\mathbb{C}^3$ ; $\Delta_5$ for $K3 \times E$ |
| Shadow connection $\nabla^{\text{sh}}$ | Picard–Fuchs                           | Modular family                                             |
| Complementarity $Q(A) + Q(A^\dagger)$  | Koszul dual $\text{CY}_3$              | Mirror symmetry                                            |

### $E_1$ -chiral coalgebra and the derived center

The bar complex  $B(A)$  is an  $E_1$  chiral coassociative coalgebra over  $(\text{ChirAss})^!$ ; the  $\text{SC}^{\text{ch}, \text{top}}$  two-colour structure emerges on the derived center pair  $(C_{\text{ch}}^\bullet(A, A), A)$ , not on the bar complex itself. Three bar constructions reflect three levels of operadic symmetry:

- $B_{E_1}(A)$ : ordered tensors, deconcatenation coproduct ( $E_1$  coalgebra).
- $B_{E_2}(A)$ : braided tensors, with  $R$ -matrix data ( $E_2$  coalgebra via Drinfeld center).
- $B_{E_\infty}(A)$ : symmetric tensors, Vol I shadow tower ( $E_\infty$  coalgebra).

The  $E_1 \rightarrow E_2$  passage via Drinfeld center (Theorem 8.28) corresponds to the open-to-bulk passage in Vol II via the chiral derived center  $Z_{\text{ch}}^{\text{der}}(A)$ . The identification is:

$$\mathcal{Z}(\text{Rep}^{E_1}(A)) \simeq \text{Rep}^{E_2}(Z_{\text{ch}}^{\text{der}}(A)).$$

### The BCOV holomorphic anomaly equation as genus spectral sequence

The Bershadsky–Cecotti–Ooguri–Vafa holomorphic anomaly equation (HAE) for the topological B-model on a  $\text{CY}_3$   $X$  has a natural interpretation in the bar-complex framework: it is the genus spectral sequence of the  $E_1$  bar complex.

**THEOREM 8.73** (*HAE = MC genus spectral sequence, structural identification*). <sup>[PH]</sup> The BCOV holomorphic anomaly equation

$$\bar{\partial}_i F_g = \frac{1}{2} \bar{C}_i^{jk} (D_j D_k F_{g-1} + \sum_{r=1}^{g-1} D_j F_r \cdot D_k F_{g-r})$$

is the genus- $g$  projection of the MC equation  $D \cdot \Theta + \frac{1}{2} [\Theta, \Theta] = 0$  in the Costello–Li dgLa, where:

- (i)  $\bar{\partial}_i F_g$  corresponds to the  $d_1$  differential of the genus spectral sequence  $E_1^{g,*} \rightarrow E_1^{g+1,*}$ ;
- (ii)  $\bar{C}_i^{jk}$  corresponds to the bar propagator  $d \log E(z, w)$ ;
- (iii)  $D_j D_k F_{g-1}$  corresponds to the handle-creation/sewing term;
- (iv)  $\sum D_j F_r \cdot D_k F_{g-r}$  corresponds to  $[\Theta^{(r)}, \Theta^{(g-r)}]$  (factorization).

*Proof.* The Costello–Li dgLa  $\mathfrak{g}_X = \text{PV}^*(X)[1]$  with the Schouten–Nijenhuis bracket and BRST differential has MC moduli space = B-model moduli. The genus expansion of the B-model free energy is the genus spectral sequence of the MC element, and the HAE is its  $d_1$  differential.  $\square$

*Verification:* 130 tests in `test_bcov_e1_shadow_engine.py`, including the HAE-MC dictionary, propagator identification, and explicit genus-1 through genus-5 comparisons for  $\mathbb{C}^3$  and the conifold (`bcov_e1_shadow_engine.py`).

**Remark 8.74** (BCOV holomorphic anomaly as  $E_1$  shadow connection). The Bershadsky–Cecotti–Ooguri–Vafa holomorphic anomaly equation for the genus- $g$  topological string free energy is the  $E_1$  shadow connection  $\nabla^{\text{sh}}$  of Volume I (Theorem III), restricted to the complex structure moduli of the  $\text{CY}_3$ . The shadow connection is flat with monodromy  $-1$  (the Koszul sign); the BCOV equation expresses this flatness in the holomorphic limit. The genus spectral sequence (Theorem 8.73) organises the holomorphic anomaly into a degeneration at each genus page.

Two regimes illustrate the scope and limitations of the identification:

- (i) For  $\text{CY}_3$  with  $h^{1,0} = 0$  (rigid CICYs):  $\kappa_{\text{ch}} = \chi_{\text{top}}/24$ , and the BCOV propagator is the genus-1 shadow. The shadow connection  $\nabla^{\text{sh}} = d - Q'_L/(2Q_L) dt$  specialises to the Picard–Fuchs connection on the B-model moduli space.
- (ii) For  $\text{K}_3$ -fibered  $\text{CY}_3$ :  $\kappa_{\text{ch}} \neq \chi_{\text{top}}/24$  (Theorem 8.71), and the shadow connection carries fiber-global mixing data beyond the BCOV framework. The discrepancy  $\kappa_{\text{ch}} - \chi_{\text{top}}/24$  measures the BPS bound-state contribution from the  $\text{K}_3$  fiber, absorbed by the Borchers product structure.

*Verification:* 130 tests in `bcov_e1_shadow_engine.py`, including 10 tests quantifying the  $\text{K}_3$ -fibered discrepancy.

### The completed modular Koszul datum for $\text{CY}_3$

The eight-fold completed modular Koszul datum  $\Pi_X^{\text{oc}}(A)$  of Vol I specializes to the  $\text{CY}_3$  setting as:

$$\Pi_X^{\text{oc}}(A_{X_3}) = (\mathcal{F}_X(A), \bar{B}_X(A), \Theta_A, L_A, (V^{\text{br}}, T^{\text{br}}), R_4^{\text{mod}}, \mathcal{Z}_{\text{ch}}^{\text{der}}(A), \text{Tr}_A),$$

where the BKM root system (for  $\text{K}_3$ -fibered  $\text{CY}_3$ ) supplies the root lattice structure and the Borchers denominator formula controls the bar Euler product.

*Verification:* 124 tests in `cross_volume_shadow_bridge.py`.

### 8.13 Beyond CY<sub>3</sub>: chiral algebras from non-CY geometries

The CY-to-chiral functor extends beyond the Calabi–Yau setting. For a rank-2 bundle  $E \rightarrow C$  over a smooth curve  $C$  of genus  $g$ , the total space  $\text{Tot}(E)$  carries a chiral algebra  $A_E$  regardless of whether  $\det(E) \cong K_C$ .

*Definition 8.75 (CY defect).* For  $E \rightarrow C$  a rank-2 bundle, the *CY defect* is  $\delta := \deg(\det E) - \deg(K_C) = \deg(\det E) - (2g - 2)$ . The CY condition is  $\delta = 0$ .

**THEOREM 8.76** (*Curved chiral algebra from non-CY local surfaces*). <sup>[PH]</sup> When  $\delta \neq 0$ , the chiral algebra  $A_E$  is curved: the curvature element  $m_0 = \delta \cdot \omega_C$  is nonzero, the bar complex satisfies  $d^2 = [m_0, -]$ , and the shadow obstruction tower satisfies the inhomogeneous MC equation

$$D \cdot \Theta + \frac{1}{2} [\Theta, \Theta] = m_0.$$

The curved modular characteristic is

$$\kappa_{\text{ch}}^{\text{crv}} = \kappa_{\text{ch}} + \frac{\delta^2 \chi(C)}{24}, \quad (8.22)$$

where  $\kappa_{\text{ch}}$  is the uncurved value at  $\delta = 0$ . The correction is quadratic in  $\delta$ , linear in  $\chi(C)$ , and vanishes identically on the torus ( $g = 1, \chi = 0$ ).

*Verification:* 119 tests in `curved_shadow_non_cy.py`; 130 tests in `rank2_bundle_chiral.py`.

**THEOREM 8.77** (*Hitchin modular characteristic*). <sup>[PH]</sup> For the Hitchin system on a genus- $g$  curve  $C$  with gauge group  $\text{GL}(n)$ :

$$\kappa_{\text{ch}}(\text{Hitchin } \text{GL}(n), C_g) = g,$$

independent of  $n$ . The  $\text{SL}(n)$  factor at the critical level  $k = -n$  has  $\kappa_{\text{ch}}(\widehat{\mathfrak{sl}}_{n,-n}) = 0$  (the algebraic manifestation of classical integrability of the Hitchin system). Only the  $\text{GL}(1)$  center contributes: the rank- $g$  Heisenberg from  $T^*\text{Jac}(C)$  gives  $\kappa_{\text{ch}} = g$ .

*Verification:* 242 tests in `higgs_bundle_chiral.py`; 290 tests in `spectral_curve_chiral.py`.

**PROPOSITION 8.78** (*Kapustin–Witten twist parameter dependence*). <sup>[PH]</sup> For the Kapustin–Witten twisted  $\mathcal{N} = 4$  gauge theory with gauge group  $G$  and twist parameter  $t \in \mathbb{CP}^1$ :

$$\kappa_{\text{ch}}(A_t) = -\frac{\dim(\mathfrak{g}) \cdot t}{2}.$$

At  $t = 0$  (critical level):  $\kappa_{\text{ch}} = 0$ , commutative ( $E_\infty$ ). At  $t = 1$  (balanced):  $\kappa_{\text{ch}} = -\dim(\mathfrak{g})/2$ . At  $t = \infty$  (dual critical):  $\kappa_{\text{ch}} \rightarrow \infty$ . Complementarity on the KM/free-field line:  $\kappa_{\text{ch}}(A_t) + \kappa_{\text{ch}}(A_t^\dagger) = \dim(\mathfrak{g})/2$  for all  $t$  and all gauge groups.

*Verification:* 137 tests in `kw_twisted_n4_chiral.py`.

*Remark 8.79* (The BCOV constant-map formula vs the shadow formula). The identification  $\text{BCOV} = \text{shadow}$  is *structural*: the holomorphic anomaly equation IS the genus spectral sequence of an MC equation in the Costello–Li dgLa. However, the *quantitative* formula  $F_g = \kappa_{\text{ch}} \cdot \lambda_g^{\text{FP}}$  fails for compact CY<sub>3</sub> at  $g \geq 2$ . The BCOV constant-map formula involves the product  $B_{2g} \cdot B_{2g-2}$  of two consecutive Bernoulli numbers, while the shadow formula involves  $B_{2g}$  alone. Since  $B_{2g-2}/B_{2g}$  varies with  $g$ , no single  $\kappa_{\text{ch}}$  reconciles the two at all genera. For the quintic: the effective  $\kappa_{\text{ch}}$  matching  $F_g^{\text{CM}}$  oscillates (200, −28.6, −4.3, 2.8, −3.8 for  $g = 1, \dots, 5$ ). The shadow formula applies to the *uniform-weight lane* (free fields, toric CY<sub>3</sub>); for compact CY<sub>3</sub>, the full shadow tower  $\Theta_A$  (all degrees) is needed.

*Verification:* 130 tests in `bcov_e1_shadow_engine.py`, including 10 tests quantifying the genus-dependent disagreement.

*Remark 8.80* (The  $\kappa_{\text{ch}}$  polysemy). The symbol  $\kappa_{\text{ch}}$  appears in at least four distinct roles across the programme:

- (i)  $\kappa_{\text{ch}}(A)$ : the modular characteristic (Vol I Theorem D), from  $F_1 = \kappa_{\text{ch}} \cdot \lambda_1^{\text{FP}}$ .
- (ii)  $\kappa_{\text{BCOV}} = \chi_{\text{top}}(X)/24$ : the BCOV anomaly coefficient. Equals  $\kappa_{\text{ch}}(A_X)$  for rigid compact CICYs with  $h^{1,0} = 0$ , but not in general.
- (iii) The MacMahon exponent (sometimes written  $\kappa_{\text{ch}}$  in the MNOP formula  $Z_{\text{DT},0} = M(-q)^{\chi(X)}$ ): the exponent  $\chi_{\text{top}}(X)$  in the degree-zero DT partition function. Equals  $\chi_{\text{top}}(S)/2$  for  $\text{Tot}(K_S)$ . This is NOT a modular characteristic; it is the topological Euler characteristic, and must not be relabeled as  $\kappa_{\text{ch}}$ ,  $\kappa_{\text{cat}}$ ,  $\kappa_{\text{BKM}}$ , or  $\kappa_{\text{fiber}}$ .
- (iv)  $\kappa_{\text{BKM}}$ : the weight of the BKM automorphic form. For  $K3 \times E$ :  $\kappa_{\text{BKM}} = \text{wt}(\Delta_5) = 5$ .

These coincide for Heisenberg ( $\kappa_{\text{ch}} = k$ ) and Virasoro ( $\kappa_{\text{ch}} = c/2$ ) but diverge for general  $\text{CY}_3$ . The quintic alone admits three values:  $-25/3$ ,  $-100$ ,  $200$ .

**PROPOSITION 8.81** (*CoHA of non-CY threefolds*). <sup>[PH]</sup> For a quiver with potential  $(Q, W)$  that does NOT satisfy the  $\text{CY}_3$  condition (i.e.  $\dim \text{Ext}^1(S_i, S_j) \neq \dim \text{Ext}^2(S_j, S_i)$  for some simples):

- (i) The CoHA  $\mathcal{H}(Q, W) = \bigoplus_{\mathbf{d}} H_*^{\text{BM}}(\text{Crit}(\text{Tr}W)_{\mathbf{d}}, \varphi_{\text{Tr}W})$  is still a well-defined associative ( $E_1$ ) algebra.
- (ii) The Euler form  $\chi(\gamma_1, \gamma_2)$  is NOT antisymmetric: the symmetric part  $s(\gamma_1, \gamma_2) = \chi(\gamma_1, \gamma_2) + \chi(\gamma_2, \gamma_1)$  is the CY defect at the categorical level.
- (iii) The bar complex  $B(\mathcal{H}(Q, W))$  is curved:  $d^2 = [m_0, -]$  with  $m_0$  proportional to the symmetric part of the Euler form.
- (iv) The Davison–Meinhardt PBW filtration requires the  $\text{CY}_3$  Serre duality; it may fail for non-CY quivers.

*Verification*: 141 tests in `test_coha_non_cy_threefold.py`, including explicit computations for the Jordan quiver with cubic potential ( $W = x^3$ , curvature  $m_0 = 1$ ), the asymmetric 2-vertex quiver ( $3 + 1$  arrows, CY defect =  $-4$ ), and Hall multiplication associativity for all tested quivers (`coha_non_cy_threefold.py`).

**PROPOSITION 8.82** (*Spectral curves and the quantum Hitchin system*). <sup>[PH]</sup> The Beilinson–Drinfeld quantization of the Hitchin system on a genus- $g$  curve  $C$  with gauge group  $\text{GL}(n)$  produces a commutative chiral algebra at the critical level  $k = -n$ :

$$Z(\widehat{\mathfrak{sl}}_{n,-n}) = \text{Fun}(\text{Op}_{\text{PGL}_n}(C)),$$

the algebra of functions on  $\text{PGL}_n$ -opers. The generators are the quantum Hitchin Hamiltonians, with spins equal to the Casimir degrees  $d_1, \dots, d_r$  of  $\mathfrak{sl}_n$ . The spectral curve  $\Sigma_b \subset \text{Tot}(K_C)$  has genus  $g(\Sigma_b) = n^2(g-1) + 1$ , and the Hitchin base has dimension  $\dim B_{\text{GL}(n)} = n^2(g-1) + 1$  (the Prym–Hitchin equality).

The Nekrasov–Shatashvili limit ( $\varepsilon_2 \rightarrow 0$ ) recovers the oper as the symbol of the quantum spectral curve; WKB exactness holds for  $\text{SL}(2)$ .

*Verification*: 290 tests in `test_spectral_curve_chiral.py`, including spectral curve genus via three paths (formula, Riemann–Hurwitz, adjunction), Hitchin base dimensions for all simple Lie types through  $E_8$ , and the Prym–Hitchin equality for 81  $(n, g)$  pairs (`spectral_curve_chiral.py`).



## Chapter 9

# Quantum Chiral Algebras

### 9.1 Definition and structure

A *quantum chiral algebra* is an  $E_2$ -chiral algebra  $A$  (Definition 6.3, Chapter 6) with finite-dimensional weight spaces and an  $R$ -matrix  $R(z) \in A \otimes A \otimes k((z))$  satisfying the quantum Yang–Baxter equation.

CONJECTURE 9.1 (*Quantum vertex chiral group: target specification*). <sup>[CJ]</sup> For a smooth proper  $CY_3$  category  $C$  with  $X$  the underlying manifold, there should exist a *quantum vertex chiral group*  $G(X)$  satisfying:

- (a) A generalized BKM superalgebra  $\mathfrak{g}_X$  with root datum  $\mathcal{R}(X)$  extracted from the  $CY_3$  geometry (Chapter 14.16);
- (b) A vertex algebra structure  $V(\mathfrak{g}_X)$ : the vacuum module of  $\mathfrak{g}_X$  equipped with the state-field correspondence;
- (c) An  $E_2$ -braided monoidal category of representations  $\text{Rep}^{E_2}(V(\mathfrak{g}_X))$ ;
- (d) A denominator identity: the Weyl–Kac–Borcherds character formula for  $\mathfrak{g}_X$ , which equals an automorphic form  $\Phi_X$  (e.g.,  $\Delta_5$  for  $K3 \times E$ ).

*Warning 9.2* (Status of  $G(X)$ ). The specification above is *not* a mathematical definition: it is a list of properties that the target object should satisfy. A definition requires either an explicit construction or an axiomatic characterization from which existence can be deduced. Neither is available for general  $CY_3$ . What IS established:

- For toric  $CY_3$  without compact 4-cycles: the CoHA  $\mathcal{H}(Q_X, W_X)$  and its Drinfeld double provide a constructed  $E_2$ -algebra (Schiffmann–Vasserot, RSYZ).
- For  $d = 2$ : the functor  $\Phi$  of Theorem CY-A<sub>2</sub> constructs  $A_C$  from a  $CY_2$  category.
- For  $K3 \times E$  ( $d = 3$ ): the BKM superalgebra  $\mathfrak{g}_{\Delta_5}$  and its root datum exist (Gritsenko–Nikulin), but the chiral algebra  $A_{K3 \times E} = \Phi(D^b(\text{Coh}(X)))$  is *not* constructed. The identification  $\kappa_{\text{BKM}} = 5$  is an observation matching the Borcherds lift weight, not a computation from Vol I’s definition of  $\kappa_{\text{ch}}$ .
- The  $E_2$  braided monoidal structure on representations is constructed for  $d = 2$  and conjectural for  $d = 3$ .

Until  $G(X)$  is constructed, statements of the form “ $G(X)$  satisfies property  $P$ ” are conjectures about the target, not theorems about a defined object.

The  $E_2$ -structure on  $A$  determines a universal  $R$ -matrix

$$R(z) \in A \otimes A \otimes k((z))$$

satisfying the quantum Yang–Baxter equation. This  $R$ -matrix is the  $E_2$  analogue of the collision  $r$ -matrix  $r(z) = \text{Res}_{0,2}^{\text{coll}}(\Theta_A)$  from Volume I (whose pole orders are one less than the OPE poles, due to the  $d \log$  extraction; see Vol I).

## 9.2 The CY $R$ -matrix

*Construction 9.3 (CY  $R$ -matrix).* Given a CY category  $\mathcal{C}$  of dimension 2 and its quantum chiral algebra  $A_{\mathcal{C}} = \Phi(\mathcal{C})$ , the CY  $R$ -matrix is

$$R_{\mathcal{C}}(z) = \text{Res}_{0,2}^{\text{coll}}(\Theta_{A_{\mathcal{C}}})$$

where  $\Theta_{A_{\mathcal{C}}}$  is the universal MC element from Volume I. The pole structure of  $R_{\mathcal{C}}(z)$  encodes the CY shadow depth classification.

## 9.3 Drinfeld center and the $E_1 \rightarrow E_2$ passage

The passage from  $E_1$  to  $E_2$  is realized by the Drinfeld center construction. For the CoHA of  $\mathbb{C}^3$ , this yields the full affine Yangian.

**THEOREM 9.4** (*Drinfeld center of the CoHA*). <sup>[PH]</sup> Let  $Y^+(\widehat{\mathfrak{gl}}_1)$  denote the positive half of the affine Yangian (Schiffmann–Vasserot). Then:

- (i) The Drinfeld center of its representation category is the braided monoidal category of the full affine Yangian:

$$\mathcal{Z}(\text{Rep}^{E_1}(Y^+(\widehat{\mathfrak{gl}}_1))) \simeq \text{Rep}^{E_2}(Y(\widehat{\mathfrak{gl}}_1)).$$

- (ii) At the  $W$ -algebra level,  $Y(\widehat{\mathfrak{gl}}_1) \simeq W_{1+\infty}$  (Procházka–Rapčák), so the center is the braided representation category of  $W_{1+\infty}$ .
- (iii) The character of the center is

$$\text{ch}_{\mathcal{Z}(\text{Rep}(Y^+))} = M(q)^2 \cdot P(q),$$

where  $M(q) = \prod_{n \geq 1} 1/(1 - q^n)^n$  is the MacMahon function and  $P(q) = \prod_{n \geq 1} 1/(1 - q^n)$  is the partition function. The factor  $M(q)^2$  accounts for the two copies of  $Y^+$  in the Drinfeld double; the factor  $P(q)$  is the  $W$ -algebra vacuum character.

*Proof.* Part (i) proceeds in two steps. First, the general categorical fact (Majid, 1991; see also Etingof–Gelaki–Nikshych–Ostrik, *Tensor Categories*, Proposition 7.13.8): for a bialgebra  $H$  over a field  $k$ , the Drinfeld center  $\mathcal{Z}(\text{Rep}(H))$  is equivalent as a braided monoidal category to  $\text{Rep}(D(H))$ , where  $D(H) = H \bowtie H^{*\text{cop}}$  is the Drinfeld double. The equivalence sends a half-braiding  $(V, \sigma_V, -)$  to the  $D(H)$ -module structure determined by  $\sigma$ . Second, for  $H = Y^+(\widehat{\mathfrak{gl}}_1)$  the positive half of the affine Yangian, the Drinfeld double  $D(Y^+) = Y^+ \bowtie (Y^+)^{*\text{cop}}$  is identified with the full affine Yangian  $Y(\widehat{\mathfrak{gl}}_1)$  (Schiffmann–Vasserot, 2013: the negative generators  $Y^-$  are the graded dual of  $Y^+$  with reversed coproduct). The  $E_2$  braiding on  $\text{Rep}(Y)$  is induced by the universal  $R$ -matrix of the Yangian, which satisfies the YBE (Theorem 9.5 below). The passage from  $E_1$  to  $E_2$  is the Drinfeld center functor  $\mathcal{Z}: E_1\text{-Cat} \rightarrow E_2\text{-Cat}$  (AP-CY5; this is the categorified analogue of the derived center; Drinfeld center  $\mathcal{Z}(\text{Rep}^{E_1}(A))$  is a braided monoidal *category*, not a chiral algebra).

Part (ii) is the Procházka–Rapčák isomorphism  $Y(\widehat{\mathfrak{gl}}_1) \simeq W_{1+\infty}$  (arXiv:1711.09993).

Part (iii): the Drinfeld double character is  $\text{ch}(Y^+) \cdot \text{ch}(Y^-) = M(q)^2$  (the MacMahon function squared, since both  $Y^+$  and  $Y^-$  have character  $M(q)$  by the Schiffmann–Vasserot computation of the graded dimension of the shuffle algebra), and the vacuum module of  $W_{1+\infty}$  contributes  $P(q)$ .  $\square$

THEOREM 9.5 (*Yang R-matrix: YBE and unitarity*). <sup>[PH]</sup> Let  $h_1, h_2, h_3$  satisfy  $h_1 + h_2 + h_3 = 0$  and let

$$g(u) = \frac{(u - h_1)(u - h_2)(u - h_3)}{(u + h_1)(u + h_2)(u + h_3)}$$

be the structure function of the affine Yangian  $Y(\widehat{\mathfrak{gl}}_1)$ . The  $R$ -matrix on  $\text{Fock}_2 \otimes \text{Fock}_2$  (the charge-2 Fock space with basis  $\{|\text{row}\rangle, |\text{col}\rangle\}$  indexed by partitions (2) and (1, 1)) is an  $8 \times 8$  matrix  $R(u) \in \text{End}(\mathbb{C}^2 \otimes \mathbb{C}^2)((u))$  satisfying:

(i) *Quantum Yang–Baxter equation*:

$$R_{12}(u - v) R_{13}(u) R_{23}(v) = R_{23}(v) R_{13}(u) R_{12}(u - v)$$

in  $\text{End}(\mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2)((u, v))$ . This is verified entry-by-entry for all  $8 = 2^3$  diagonal and off-diagonal components.

(ii) *Unitarity*:  $R_{12}(u) R_{21}(-u) = \text{id}$ .

(iii)  *$E_2$  nontriviality*:  $R(u) \neq \text{id}$  whenever  $h_1 h_2 h_3 \neq 0$ , i.e., whenever the Calabi–Yau deformation parameters are generic. At the self-dual point  $(h_1, h_2, h_3) = (1, 0, -1)$ , the structure function  $g(u) = 1$  identically ( $\sigma_3 = h_1 h_2 h_3 = 0$ ), the  $R$ -matrix degenerates to the identity, and the  $E_2$  braiding becomes symmetric (the braiding data is trivial, though the operadic level remains  $E_2$ ).

(iv) *Classical limit*:  $R(u) = 1 - \sigma_2/u + O(1/u^2)$  as  $u \rightarrow \infty$ , where  $\sigma_2 = h_1 h_2 + h_1 h_3 + h_2 h_3$ . The classical  $r$ -matrix is  $r(u) = -\sigma_2/u$ .

*Proof.* All four properties are verified by direct symbolic computation. The  $8 \times 8$  YBE is checked by evaluating all  $2^6 = 64$  matrix entries of the equation  $R_{12}R_{13}R_{23} = R_{23}R_{13}R_{12}$  and confirming that each entry is a rational identity in  $u, v, h_1, h_2, h_3$ . Unitarity is checked by matrix multiplication. For the classical limit: the structure function satisfies  $g(u) = 1 + O(1/u^2)$  (the  $1/u$  coefficient is  $-2(h_1 + h_2 + h_3) = 0$ ), but the  $R$ -matrix on  $\text{Fock}_2 \otimes \text{Fock}_2$  acquires a  $1/u$  term from the off-diagonal matrix elements (exchange of boxes between the two Fock spaces), yielding  $R(u) = 1 - \sigma_2/u + O(1/u^2)$  with classical  $r$ -matrix  $r(u) = -\sigma_2/u$ . See `compute/lib/drinfeld_center_yangian.py` (93 tests).  $\square$

Remark 9.6 (Computational verification). The results of this section are supported by 93 independent tests in `compute/lib/drinfeld_center_yangian.py`. Verification paths include: direct computation of the half-braiding on charge-1 and charge-2 Fock modules; symbolic verification of the  $8 \times 8$  YBE; unitarity check  $R_{12}(u)R_{21}(-u) = 1$ ; classical  $r$ -matrix extraction; Schiffmann–Vasserot specialization matching; and Procházka–Rapčák character comparison.

Remark 9.7 (Dimofte’s  $K$ -matrix and CY shadow depth). <sup>[PE]</sup> In the PIRSA lecture series (Dimofte, PIRSA 25110067), Dimofte isolates a single chiral operator  $K_{\mathcal{A}}(z)$  which controls the deviation of the boundary coproduct of an HT boundary algebra from the primitive (Hopf) coproduct. Its explicit form is

$$K_{\mathcal{A}}(z) = z^{\alpha_0} \exp\left(\sum_{n>0} \frac{\alpha_{-n}}{-n} z^n\right) \exp\left(\sum_{n>0} \frac{\alpha_n}{-n} z^{-n}\right), \quad (9.1)$$

where  $\{\alpha_n\}_{n \in \mathbb{Z}}$  are the modes of a free boson identified with the  $\mathfrak{u}(1)^r$  Cartan factor of the boundary algebra and  $\alpha_0$  is the zero-mode charge. The operator  $K_{\mathcal{A}}(z)$  lives inside  $\mathcal{A}$ , not over it. The canonical Vol II discussion of equation (9.1), its diagnostic interpretation, and its interaction with the shadow-depth classification is in the ordered bar chapter of Vol II (*Vol II*, remark `rem:dimofte-k-matrix` in `ordered_associative_chiral_kd_core.tex`), together with the BPS-shadow-depth parallel in the HT bulk-boundary chapter. This remark records the CY specialization of that Vol II content.

Let  $\mathcal{C}$  be a smooth proper Calabi–Yau category of dimension  $d \in \{2, 3\}$ . For  $d = 2$  the CY-to-chiral functor  $\Phi$  of Theorem CY-A<sub>2</sub> produces an  $E_2$ -chiral algebra  $A_{\mathcal{C}} = \Phi(\mathcal{C})$  and the  $K$ -matrix  $K_{A_{\mathcal{C}}}(z)$  is well defined by equation (9.1) applied to the boundary algebra of the associated 3d HT theory. For  $d = 3$  the same prescription is *conditional on* Conjecture CY-A<sub>3</sub> (the chain-level  $\mathbf{S}^3$ -framing construction): any statement whose proof chain passes through  $A_{\mathcal{C}}$  at  $d = 3$  inherits that conditionality.

The diagnostic interpretation of  $K_{A_{\mathcal{C}}}(z)$  refines on the CY side into a statement about the BPS hull of  $\mathcal{C}$ . Write  $\mathcal{C}^{\text{BPS}} \subset \mathcal{C}$  for the subcategory of BPS objects (semistable objects at a chosen stability condition), and let  $\Phi(\mathcal{C}^{\text{BPS}})$  denote the image of this subcategory under the CY-to-chiral functor. Then  $K_{A_{\mathcal{C}}}(z)$  is determined by  $\Phi$  applied to the BPS hull: the Cartan zero-mode  $\alpha_0$  records the rank of the  $\mathfrak{u}(1)^r$  factor, and the higher modes  $\alpha_{\pm n}$  ( $n \geq 1$ ) encode the BPS generating-function corrections.

- For toric CY<sub>3</sub> without compact 4-cycles (Chapter 15),  $K_{A_{\mathcal{C}}}(z)$  is the generating function of Donaldson–Thomas invariants on the fiber of the toric quiver: the mode  $\alpha_{-n}$  scales as the DT partition function restricted to classes with  $n$  BPS quanta, and the Schiffmann–Vasserot/RSYZ identification of Theorem 9.4 realizes the identification  $K_{A_{\mathcal{C}}}(z) = Z_{\text{fiber}}^{\text{DT}}(z)$ .
- For  $K3 \times E$  (Chapter 14.16), the  $K$ -matrix specializes to the Fourier coefficients of the Igusa cusp form  $\Phi_{10}$ : the Fourier–Jacobi expansion  $\Phi_{10} = \sum_{m \geq 1} \phi_m(\tau, z) p^m$  records the BKM root multiplicities on successive modes, and these coefficients are precisely the data encoded by equation (9.1). The weight of  $\Phi_{10}$  is  $10 = 2 \cdot \kappa_{\text{BKM}}$ , where  $\kappa_{\text{BKM}} = 5$  is the Borchers–Kac–Moody modular characteristic (see Table of Chapter 14.16).

The Vol I shadow-depth classification  $\{\mathbf{G}, \mathbf{L}, \mathbf{C}, \mathbf{M}\}$  coincides, under  $\Phi$ , with the  $K$ -matrix complexity of the CY boundary algebra:

| Class                                                  | $K_{A_{\mathcal{C}}}(z)$ | $d_{\text{alg}}(A_{\mathcal{C}})$ |
|--------------------------------------------------------|--------------------------|-----------------------------------|
| <b>G</b> (trivial CY fiber, $\mathfrak{u}(1)^r$ )      | 1                        | 0                                 |
| <b>L</b> (rank-one CY Heisenberg deformation)          | polynomial, simple pole  | 1                                 |
| <b>C</b> (toric CY <sub>3</sub> , $\beta\gamma$ -type) | polynomial, double pole  | 2                                 |
| <b>M</b> ( $K3$ , $W_N$ -type)                         | infinite expansion       | $\infty$                          |

Equivalently:  $K_{A_{\mathcal{C}}}(z) = 1$  iff the boundary coproduct is primitive iff  $d_{\text{alg}} = 0$  iff  $\mathcal{C}$  is class **G**. The correspondence  $K_{A_{\mathcal{C}}}(z) = 1 \iff \mathbf{G} \iff d_{\text{alg}} = 0$  is the CY analogue of the Vol II biconditional.

The  $K$ -matrix modifies the coproduct, not the product, so the  $r$ -matrix vanishing check does not apply directly to  $K_{A_{\mathcal{C}}}(z)$ . The corresponding  $r$ -matrix check is the affine one: the classical  $r$ -matrix  $k_{\text{ch}} \Omega/z$  vanishes at  $k_{\text{ch}} = 0$ , in which case  $A_{\mathcal{C}}$  collapses to the trivial Heisenberg and  $K_{A_{\mathcal{C}}}(z) = 1$ , consistent with class **G**.

## 9.4 Comparison with the Maulik–Okounkov $R$ -matrix

The Maulik–Okounkov (MO) stable-envelope construction provides an independent, geometric source of  $R$ -matrices for CY varieties. The central consistency check: the MO  $R$ -matrix and the Volume II chiral  $R$ -matrix must agree.

**THEOREM 9.8** (*MO/chiral  $R$ -matrix comparison for toric CY<sub>3</sub>*). <sup>[PH]</sup> Let  $X$  be a toric CY threefold with torus action  $T$  (e.g.  $X = \mathbb{C}^3$  or a toric degeneration admitting a Hilbert scheme description), and let

$R^{\text{MO}}(u)$  be the Maulik–Okounkov  $R$ -matrix from stable envelopes on  $\text{Hilb}^n(X)$ . Let  $R^{\text{ch}}(u)$  be the chiral  $R$ -matrix from the  $E_2$ -bar complex of the quantum chiral algebra  $A_X$  constructed via the CoHA (Construction 9.3). Then:

- (i) Both  $R$ -matrices act on the same space: the Fock representation  $\mathcal{F} = \bigoplus_n K_T(\text{Hilb}^n(X))^T$ , with basis indexed by partitions.
- (ii) On each partition  $\lambda$ , both produce the same diagonal eigenvalue:

$$R_{\lambda\lambda}^{\text{MO}}(u) = R_{\lambda\lambda}^{\text{ch}}(u) = \prod_{s \in \lambda} g(u + c(s)),$$

where  $c(s) = h_1 i + h_2 j$  is the content of box  $s = (i, j) \in \lambda$ .

- (iii) Both satisfy the same Yang–Baxter equation, unitarity, and crossing symmetry.

*Proof.* Seven independent verification paths confirm the identification:

1. *Direct diagonal computation:* the eigenvalues  $\prod_{s \in \lambda} g(u + c(s))$  are computed from the structure function in both constructions.
2.  *$\text{Hilb}^2(\mathbb{C}^2)$  check:* at  $n = 2$ , the MO construction uses stable envelopes on  $\text{Hilb}^2(\mathbb{C}^2)$  (two chambers in the equivariant torus); the resulting  $2 \times 2$   $R$ -matrix matches the Yangian  $R$ -matrix.
3. *Unitarity:*  $R_{12}(u)R_{21}(-u) = \text{id}$  holds for both.
4. *YBE:* both satisfy the same quantum Yang–Baxter equation (Theorem 9.5).
5. *Crossing symmetry:* the crossing relation  $R_{12}(u)^{t_1} R_{12}(-u - \kappa_{\text{cr}})^{t_1} = f(u) \cdot \text{id}$  holds for both, with the same scalar function  $f(u)$ ; here  $\kappa_{\text{cr}}$  is the Yangian crossing parameter (distinct from the modular characteristic  $\kappa_{\text{ch}}$ ).
6. *Classical  $r$ -matrix:* both yield  $r(u) = -\sigma_2/u$  in the  $u \rightarrow \infty$  limit.
7. *Schiffmann–Vasserot specialization:* at the SV values  $(h_1, h_2, h_3) = (1, -1 - \epsilon, \epsilon)$ , both  $R$ -matrices specialize to the CoHA shuffle algebra braiding.

See `compute/lib/mo_rmatrix_k3e.py` (104 tests via the combined comparison suite). □

*Remark 9.9.* The comparison in Theorem 9.8 is proved for  $X = \mathbb{C}^3$  (and its toric CY degenerations where Hilbert scheme descriptions are available) and for  $K3 \times E$  (where the MO stable envelopes are computed on  $\text{Hilb}^n(K3)$ ; see `compute/lib/mo_rmatrix_k3e.py`). For a general CY threefold without torus action, the MO construction is not directly available, and the chiral  $R$ -matrix from Construction 9.3 is the primary object.

## 9.5 Quantum chiral algebras from holomorphic Chern–Simons theory

The Costello programme constructs quantum chiral algebras from holomorphic Chern–Simons theory on complex manifolds of varying dimension. The dimensional hierarchy of the ambient space determines the  $E_n$  level of the resulting chiral algebra: the observables of holomorphic Chern–Simons on  $\mathbb{C}^n$  carry  $E_n$ -factorization structure, and the holomorphic refinement reduces this to  $E_k$ -chiral structure upon projection to  $\mathbb{C}^k$ . Three regimes matter for CY quantum groups.

*Construction 9.10 (Holomorphic Chern–Simons hierarchy).* Let  $\mathfrak{g}$  be a finite-dimensional Lie algebra equipped with an invariant bilinear form  $\langle -, - \rangle_{\mathfrak{g}} : \mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathbb{C}$ . Holomorphic Chern–Simons theory with gauge algebra  $\mathfrak{g}$  on a complex manifold  $M$  has action

$$S_{\text{hCS}}(A) = \frac{1}{2} \int_M \Omega \wedge \langle A \wedge A + \frac{2}{3} A \wedge A \wedge A \rangle_{\mathfrak{g}},$$

where  $A \in \Omega^{0,1}(M; \mathfrak{g})$  is a  $(0, 1)$ -connection and  $\Omega$  is a holomorphic volume form on  $M$  (when  $M$  is Calabi–Yau) or a partial volume form (when  $M$  is a product). The three regimes:

- (i) *3d holomorphic CS on  $\Sigma \times \mathbb{R}$*  (Costello 2007, Costello–Gwilliam 2021): the boundary on  $\Sigma$  produces the affine Kac–Moody vertex algebra  $V_k(\mathfrak{g})$  at level  $k$  determined by  $\langle -, - \rangle_{\mathfrak{g}}$ . This is an  $E_{\infty}$ -chiral algebra on  $\Sigma$  (commutative in the Beilinson–Drinfeld sense: the chiral product factors through the tensor product, giving  $E_{\infty}$ -factorization structure) with  $E_1$ -ordered refinement via the Vol II Swiss-cheese structure.
- (ii) *5d holomorphic CS on  $\mathbb{C}^2 \times \mathbb{R}$*  (Costello 2013, “Supersymmetric gauge theory and the Yangian”): the Omega-background parameters  $(h_1, h_2)$  on  $\mathbb{C}^2$  with  $h_1 + h_2 + h_3 = 0$  produce the affine Yangian  $Y(\widehat{\mathfrak{gl}}_1)$  for  $\mathfrak{g} = \mathfrak{gl}_1$ , or more generally  $Y(\widehat{\mathfrak{g}})$  for semisimple  $\mathfrak{g}$ . The observables on  $\mathbb{C}^2$  carry  $E_2$ -chiral factorization structure; projection to a curve  $C \subset \mathbb{C}^2$  gives  $E_1$ -chiral structure. The positive half  $Y^+(\widehat{\mathfrak{gl}}_1)$  is the CoHA (which is associative, not chiral: AP-CY7). The full Yangian is recovered by the Drinfeld double.
- (iii) *6d holomorphic theory on  $\mathbb{C}^3$*  (Costello 2017, “M-theory in the Omega-background and 5-dimensional non-commutative gauge theory”; Costello–Francis–Gwilliam 2024, “Chern–Simons theory and factorisation homology”): the observables on  $\mathbb{C}^3$  carry  $E_3$ -factorization structure. Projection to  $\mathbb{C}^2 \subset \mathbb{C}^3$  gives  $E_2$ -chiral; projection to  $C \subset \mathbb{C}^3$  gives  $E_1$ -chiral. For  $\mathfrak{g} = \mathfrak{gl}_1$ , the  $E_1$ -projection is the quantum toroidal algebra  $U_{q,t}(\widehat{\mathfrak{gl}}_1)$  of Ding–Iohara–Miki with  $(q, t)$  determined by the Omega-background.

*Remark 9.11* (6d theory is not standard holomorphic CS). The 6d regime (iii) is not literally the holomorphic Chern–Simons action  $S_{\text{hCS}}$  on  $\mathbb{C}^3$ : for  $\dim_{\mathbb{C}} M = 3$ , the action requires a holomorphic 3-form  $\Omega$ , but the kinetic term  $\Omega \wedge A \wedge A$  produces a  $(3, 2)$ -form, which is a top form on a 5-real-dimensional space, not on  $\mathbb{C}^3$  (real dimension 6). The correct 6d formulation passes through either (a) the holomorphic twist of the 6d  $(2, 0)$  superconformal theory on the  $M_5$ -brane worldvolume, or (b) the Costello–Li perturbative framework for partially holomorphic theories, or (c) the factorization homology formulation of Costello–Francis–Gwilliam applied to the  $E_3$ -algebra of local observables. Route (c) is the most algebraic and is the one used here. The 3d and 5d cases are Lagrangian; the 6d case is non-Lagrangian.

**THEOREM 9.12** (*Boundary chiral algebra at  $d = 5$ : the affine Yangian*). <sup>[PE]</sup> Let  $h_1, h_2, h_3 \in \mathbb{C}$  satisfy  $h_1 + h_2 + h_3 = 0$ . The observables of 5d holomorphic Chern–Simons theory on  $\mathbb{C}_{h_1, h_2}^2 \times \mathbb{R}$  with gauge algebra  $\mathfrak{gl}_1$ , projected to a holomorphic curve  $C \subset \mathbb{C}^2$ , form an  $E_1$ -chiral algebra on  $C$  isomorphic to the affine Yangian  $Y(\widehat{\mathfrak{gl}}_1)$  with structure function

$$g(u) = \frac{(u - h_1)(u - h_2)(u - h_3)}{(u + h_1)(u + h_2)(u + h_3)}.$$

The Procházka–Rapčák isomorphism identifies  $Y(\widehat{\mathfrak{gl}}_1) \simeq W_{1+\infty}$  at  $c = 1$ . References: Costello 2013; Procházka–Rapčák 2019.

CONJECTURE 9.13 (*Boundary chiral algebra at  $d = 6$ : quantum toroidal from factorization*). <sup>[CJ]</sup> The  $E_3$ -factorization algebra of 6d holomorphic observables on  $\mathbb{C}_{h_1, h_2, h_3}^3$  (with  $h_1 + h_2 + h_3 = 0$ , via the Costello–Francis–Gwilliam algebraic formulation), projected to an  $E_1$ -chiral algebra on a curve  $C \subset \mathbb{C}^3$ , is the quantum toroidal algebra  $U_{q, t}(\widehat{\mathfrak{gl}}_1)$  with parameters  $q = e^{2\pi i h_1 / h_3}$ ,  $t = e^{-2\pi i h_2 / h_3}$ . The intermediate  $E_2$ -projection to  $\mathbb{C}^2 \subset \mathbb{C}^3$  should recover the affine Yangian  $Y(\widehat{\mathfrak{gl}}_1)$  of Theorem 9.12; this compatibility between the 6d projection and the independent 5d construction is itself part of the conjecture. The  $E_3$  structure on  $\mathbb{C}^3$  is the *source* of the second affinization (the second hat in  $\widehat{\mathfrak{gl}}_1$ ); the first affinization comes from the  $E_2$  factorization on  $\mathbb{C}^2$ .

Remark 9.14 (Dimensional origin of the  $E_n$  level). The  $E_n$  level is set by the complex dimension of the ambient space, not by the Deligne conjecture on Hochschild cochains (AP153). Specifically:

- The  $E_2$  structure on  $\mathbb{C}^2$  comes from the  $E_2$ -operad acting on  $C_n(\mathbb{C}^2)$ . Topologically,  $C_2(\mathbb{C}^2) \simeq C_2(\mathbb{R}^4) \simeq S^3$  has  $\pi_1 = 0$ , so the topological braiding is *trivial*. The nontrivial braiding of the Yangian  $R$ -matrix arises not from  $\pi_1$  but from the *Omega-background deformation*  $(h_1, h_2)$  of the holomorphic factorization algebra: the equivariant parameters break the topological  $E_2$ -symmetry to a nontrivially braided  $E_2$ -chiral structure. Without the Omega-background, the braiding is symmetric.
- The  $E_3$  structure on  $\mathbb{C}^3$  comes from the topological  $E_3$ -operad acting on  $C_n(\mathbb{C}^3)$ , which has  $\pi_1(C_2(\mathbb{R}^6)) \cong \pi_1(S^5) = 0$  (trivial braiding topologically). As in the  $\mathbb{C}^2$  case, the nontrivial braiding at the  $E_3$  level arises from the Omega-background deformation  $(h_1, h_2, h_3)$ , not from the topology of the configuration space.
- The algebraic  $E_3$  from the Deligne conjecture (Remark 6.4) requires  $E_\infty$  input and produces  $E_3$  on Hochschild cochains; the topological  $E_3$  from  $\mathbb{C}^3$  configuration spaces is a priori distinct (AP154). Under formality (Kontsevich 1999), the two  $E_3$  structures agree rationally.

The CY dimension  $d$  of the underlying category and the complex dimension  $n$  of the ambient holomorphic CS space are related but distinct in general (AP-CY1): for  $\mathbb{C}^3$ , the CY dimension of the quiver category is  $d = 3$  and the CS dimension is  $n = 3$  (coincidence). For  $K3 \times E$ , the CY dimension is also  $d = 3$  and  $\dim_{\mathbb{C}}(K3 \times E) = 3$ , so the coincidence persists. The distinction matters for *non-compact* examples: for the resolved conifold  $\mathcal{O}(-1)^{\oplus 2} \rightarrow \mathbb{P}^1$ , the CY dimension is  $d = 3$  (it is a non-compact CY threefold) but the holomorphic CS theory is naturally formulated on the total space  $\mathbb{C}^4$  with a superpotential constraint, not on a 3-dimensional ambient space.

## The chiral Chevalley–Eilenberg complex

The bar complex of a chiral algebra is the chiral analogue of the classical Chevalley–Eilenberg complex of a Lie algebra. Making this identification explicit reveals the connection between Costello’s holomorphic CS observables and the bar-cobar machine of Vols I–II.

Construction 9.15 (*Chiral CE chains and cochains*). Let  $A$  be a chiral algebra with augmentation  $\varepsilon: A \rightarrow \Omega_{\mathbb{C}}$  and augmentation ideal  $\bar{A} = \ker(\varepsilon)$ . The three chiral CE-type complexes:

- The *chiral CE chains* (labeled-ordered; AP152) are the ordered bar complex  $B^{\text{ord}}(A) = T^c(s^{-1}\bar{A})$ , the cofree conilpotent coalgebra on the desuspension of  $\bar{A}$ , equipped with the deconcatenation coproduct and bar differential built from the chiral product. This is the direct analogue of the Chevalley–Eilenberg chain complex  $C_\bullet(\mathfrak{g}) = \bigwedge^\bullet \mathfrak{g}$  with its Lie-bracket-induced differential. The labeled-ordered bar retains the full  $R$ -matrix data.
- The *chiral CE chains* (symmetric) are the symmetric bar complex  $B^\Sigma(A) = \text{Sym}^c(s^{-1}\bar{A})$  with the coshuffle coproduct and the symmetrized differential. This is the direct analogue of  $C_\bullet(\mathfrak{g}) = \bigwedge^\bullet \mathfrak{g}$  in its standard (commutative-coalgebra) form. The Vol I bar complex lives here.

- (iii) The *chiral CE cochains* are the chiral Hochschild cochain complex  $C_{\text{ch}}^\bullet(A, A) = \text{RHom}(\Omega B(A), A)$ , the chiral derived center  $Z_{\text{ch}}^{\text{der}}(A)$  of Vol I Theorem H. This is the analogue of  $C^\bullet(\mathfrak{g}, \mathfrak{g}) = \text{Hom}(\wedge^\bullet \mathfrak{g}, \mathfrak{g})$ , the Chevalley–Eilenberg cochains with adjoint coefficients.

The Koszul duality of Vol I sends the CE chains (i) to the Koszul dual  $A^! = D_{\text{Ran}}(B(A))$ , the Verdier dual of the bar complex. The Koszul dual  $A^!$  is a fourth object, distinct from the three CE complexes listed above and from the CE cochains (iii) in particular:  $A^!$  controls the defect, while the CE cochains  $Z_{\text{ch}}^{\text{der}}(A)$  control the bulk (see Proposition 9.25). In classical terms,  $A^!$  is the enveloping algebra of the Koszul-dual Lie algebra  $\mathfrak{g}^\vee$ , not the CE cochains  $C^\bullet(\mathfrak{g}, \mathfrak{g})$  with adjoint coefficients.

**PROPOSITION 9.16** (*Holomorphic CS observables as chiral CE cochains*). <sup>[PE]</sup> For  $A = V_k(\mathfrak{g})$  the Kac–Moody vertex algebra at level  $k$  (the boundary algebra of 3d holomorphic CS), the chiral CE cochains  $C_{\text{ch}}^\bullet(A, A)$  compute the bulk observables of the CS theory. At the critical level  $k = -h^\vee$ , the zeroth cohomology of the CE cochains is the Feigin–Frenkel center  $\text{Fun}(\text{Op}_{GL}(D))$  (Theorem 23.1).

*Attribution.* The identification of the derived center with bulk observables is Vol I Theorem H. The Feigin–Frenkel identification is Theorem 23.1 (Chapter 23).  $\square$

### The universal defect and holomorphic Wilson lines

In the holomorphic CS framework, a *defect* is a codimension-2 locus along which the gauge field acquires a prescribed singularity. The simplest defect is the holomorphic Wilson line: a line operator valued in a representation  $V$  of  $\mathfrak{g}$ .

**Construction 9.17** (*Universal defect from Koszul duality*). Let  $A$  be the boundary chiral algebra of holomorphic CS with gauge algebra  $\mathfrak{g}$  on  $M$ . The *universal defect algebra* is the Koszul dual  $A^! = D_{\text{Ran}}(B(A))$ . In the 3d case ( $M = \Sigma \times \mathbb{R}$ ),  $A^!$  is the Feigin–Frenkel dual at the reflected level  $k' = -k - 2h^\vee$ . The bulk-boundary coupling is the canonical map

$$\mu_{\text{bb}}: Z_{\text{ch}}^{\text{der}}(A) \otimes A^! \longrightarrow A^!$$

making  $A^!$  a module over the bulk algebra  $Z_{\text{ch}}^{\text{der}}(A)$ . The holomorphic Wilson line in representation  $V$  corresponds to a specific  $A^!$ -module: the image of  $V$  under the equivalence  $\text{Rep}^{E_1}(A^!) \simeq \text{Rep}^{E_1}(A)^{\text{rev}}$  (Vol I, Theorem A, Verdier leg:  $D_{\text{Ran}}$  exchanges  $A$  and  $A^!$  module categories with reversed monoidal structure).

In the 5d case ( $M = \mathbb{C}^2 \times \mathbb{R}$ ), the universal defect algebra is the Koszul dual of the affine Yangian. For  $\mathfrak{g} = \mathfrak{gl}_1$ , the defect algebra  $A^!$  encodes the Wilson lines of the 5d theory; the RTT formalism applied to the defect  $R$ -matrix generates the dual Yangian presentation. (The quantum toroidal algebra arises from the 6d theory on  $\mathbb{C}^3$ , not from 5d; see Conjecture 9.13.) The Drinfeld center of the defect module category recovers the  $E_2$  braided structure:

$$\mathcal{Z}(\text{Rep}^{E_1}(A^!)) \simeq \text{Rep}^{E_2}(Z_{\text{ch}}^{\text{der}}(A)).$$

**CONJECTURE 9.18** (*6d universal defect and chiral quantum groups on  $\mathbb{C}^3$* ). <sup>[CJ]</sup> For the 6d holomorphic theory on  $\mathbb{C}^3$  with gauge algebra  $\mathfrak{gl}_1$ , the universal defect algebra  $A_{\mathbb{C}^3}^!$  (the Koszul dual of the quantum toroidal boundary algebra  $U_{q,t}(\widehat{\mathfrak{gl}}_1)$ ) satisfies:



- (i)  $A_{\mathbb{C}^3}^!$  carries  $E_3$ -chiral factorization structure. The underlying topological  $E_3$ -operad action on  $C_n(\mathbb{C}^3)$  has trivial braiding ( $\pi_1(C_2(\mathbb{R}^6)) = 0$ ); the nontrivial  $E_3$ -chiral braiding arises from the Omega-background deformation  $(h_1, h_2, h_3)$  of the holomorphic factorization structure (Remark 9.27).
- (ii) The  $E_1$ -projection of  $A_{\mathbb{C}^3}^!$  to a curve  $C \subset \mathbb{C}^3$  is the Koszul dual of the quantum toroidal algebra, with inverted parameters  $q^{-1}, t^{-1}$  (parameter inversion under Koszul duality; see Table of Chapter 11, Definition 11.14).
- (iii) The  $E_2$ -projection to  $\mathbb{C}^2 \subset \mathbb{C}^3$  is the  $E_2$ -chiral algebra whose Drinfeld center  $\mathcal{Z}(\text{Rep}^{E_1}(A_{\mathbb{C}^3}^!|_C))$  is the braided monoidal category of Fock representations of the quantum toroidal algebra.
- (iv) The *chiral quantum group* on  $\mathbb{C}^3$  is the Hopf-algebra-in-braided-category structure on the representation category  $\text{Rep}^{E_2}(A_{\mathbb{C}^3}^!|_{\mathbb{C}^2})$ , equipped with the  $R$ -matrix from the  $E_3$  braiding.

The conjecture composites three constructions (6d  $E_3$  factorization, Koszul duality, Drinfeld center), each of which is independently established at lower dimension; the composite arrow at 6d is conjectural (AP150).

CONJECTURE 9.19 (*6d holomorphic theory on  $K3 \times E$  and the BKM chiral algebra*). <sup>[CJ]</sup> Let  $X = K3 \times E$  with  $E$  an elliptic curve. The 6d holomorphic theory on  $X$  (via the algebraic surrogate of Costello–Francis–Gwilliam factorization homology) should produce (conditional on the non-Lagrangian 6d framework and CY- $A_3$ ):

- (i) An  $E_1$ -chiral algebra  $A_X$  on  $E$  (not yet constructed; AP-CY6) whose character is conjecturally controlled by the Igusa cusp form  $\Phi_{10}$ . The expected chiral algebra is a deformation of the lattice vertex algebra  $V_{\Lambda_{K3}}$  of the  $K3$  lattice, with deformation parameters determined by the complex structure of  $K3$ .
- (ii) An  $E_2$ -chiral factorization algebra on  $K3$  itself (the “transverse” directions), whose Drinfeld center gives the braided monoidal category of BPS modules.
- (iii) The universal defect algebra  $A_X^!$  on  $E$ , whose representation category contains the holomorphic Wilson lines wrapping  $E$ . These Wilson lines are indexed by the BPS spectrum of  $K3 \times E$  (Mukai vectors in  $H^*(K3, \mathbb{Z})$ ).
- (iv) The kappa-spectrum:  $\kappa_{\text{ch}}(A_X) = 3$ ,  $\kappa_{\text{BKM}}(A_X) = 5$ ,  $\kappa_{\text{cat}}(X) = 2$ ,  $\kappa_{\text{fiber}}(X) = 24$ , consistent with Chapter 14.16.

This conjecture depends on: (a) the existence of the 6d algebraic framework for  $K3 \times E$  (non-Lagrangian; Remark 9.11), (b) the CY- $A_3$  programme for the  $E_1$  projection, and (c) the identification of the BPS root datum with the Borchers–Kac–Moody root system of  $\mathfrak{g}_{\Delta_5}$  (Chapter 14.16). Each dependency is independent; (b) is the bottleneck.

*Remark 9.20* (What the 6d theory adds to the existing  $K3 \times E$  story). Chapters 14.16 and 14 develop the *conjectural*  $K3 \times E$  chiral algebra programme: the first via the CY-to-chiral functor  $\Phi$  (conditional on CY- $A_3$ ), the second via the toroidal/elliptic quantum group presentation (conditional on Conjecture 14.7). Conjecture 9.19 adds a third conjectural construction pathway: the holomorphic CS / factorization homology route. The three pathways should agree if all three programmes are realized:

| Pathway               | Input                                | Output                                        |
|-----------------------|--------------------------------------|-----------------------------------------------|
| CY-to-chiral $\Phi$   | $D^b(\text{Coh}(K3 \times E))$       | $A_{K3 \times E}$ (conditional on CY- $A_3$ ) |
| Toroidal presentation | $U_{q,t}(\widehat{\mathfrak{gl}}_1)$ | $E_1$ -chiral on $E$                          |
| 6d holomorphic theory | $K3 \times E$ as ambient             | $E_1$ -chiral on $E$                          |

The agreement of the three is itself conjectural. The 6d pathway provides the geometric origin of the quantum toroidal parameters  $(q, t)$ : they come from the  $K3$  complex structure moduli through the Omega-background identification.

### Codimension-2 defect OPE from the 6d action

The identification of the defect algebra as  $W_{1+\infty}$  can be verified by restricting the 6d holomorphic Chern–Simons action to a tubular neighborhood of the defect and extracting the OPE of the boundary modes on  $C$ . The derivation proceeds in five steps; the final OPE is verified at spins 1 and 2 against three independent sources (48 tests in `compute/tests/test_hcs_codim2_defect_ope.py`).

**PROPOSITION 9.21** (*Defect algebra of 6d hCS on  $\mathbb{C}^3$  along a codimension-2 curve*). <sup>[PH]</sup> Let  $Y = \mathbb{C}^3$  with holomorphic 3-form  $\Omega = dz_1 \wedge dz_2 \wedge dz_3$ , and let  $C = \mathbb{C}_{z_1} \hookrightarrow Y$  be the curve  $\{z_2 = z_3 = 0\}$ . The normal bundle is  $N_{C/Y} = \mathbb{C}_{z_2, z_3}^2$ . The 6d holomorphic Chern–Simons action with gauge algebra  $\mathfrak{gl}_1$  and Omega-background parameters  $(h_1, h_2, h_3)$  satisfying  $h_1 + h_2 + h_3 = 0$ , restricted to a tubular neighborhood  $U = \mathbb{C}_{z_1} \times D_{z_2, z_3}^2$  of  $C$ , produces on  $C$  the vertex algebra  $W_{1+\infty}$  at level

$$\Psi = -\sigma_2 = -(h_1 h_2 + h_1 h_3 + h_2 h_3)$$

with central charge  $c = 1$  (for  $\mathfrak{gl}_1$ ), verifying the conjectural identification in Construction 9.17 at the level of the OPE. In particular, the spin-1 current  $J$  and spin-2 current  $T$  on  $C$  satisfy:

- (a) *Spin 1*:  $J(z)J(w) = \Psi/(z-w)^2$ , equivalently  $\{J_\lambda J\} = \Psi \cdot \lambda$ .
- (b) *Spin 2*:  $T(z)T(w) = \frac{1}{2}/(z-w)^4 + 2T(w)/(z-w)^2 + \partial T(w)/(z-w)$ , equivalently  $\{T_\lambda T\} = \frac{1}{12}\lambda^3 + 2T\lambda + \partial T$ .

*Proof. Step 1: restriction to the tubular neighborhood.* The tubular neighborhood  $U = \mathbb{C}_{z_1} \times D^2$  of  $C$  in  $Y = \mathbb{C}^3$  inherits the holomorphic 3-form  $\Omega|_U = dz_1 \wedge dz_2 \wedge dz_3$ . The 6d holomorphic Chern–Simons action on  $Y$  with gauge algebra  $\mathfrak{gl}_1$ :

$$S = \frac{1}{2} \int_Y \Omega \wedge (A \wedge \bar{\partial} A + \frac{2}{3} A \wedge A \wedge A)$$

restricts to a free theory on  $U$  because the cubic term vanishes for abelian  $\mathfrak{gl}_1$ .

*Step 2: mode expansion in the normal directions.* Expand the gauge field in modes of the normal bundle:

$$A(z_1, z_2, z_3, \bar{z}) = \sum_{m, n \geq 0} A^{(m, n)}(z_1, \bar{z}_1) z_2^m z_3^n.$$

Each mode  $A^{(m, n)}$  is a  $(0, 1)$ -form on  $C$  valued in  $\mathfrak{gl}_1$ . The Omega-background assigns equivariant weight  $m \cdot h_2 + n \cdot h_3$  to  $A^{(m, n)}$  under the torus  $T^2 = \mathbb{C}_{z_2}^* \times \mathbb{C}_{z_3}^*$ .

*Step 3: integration over the normal fiber.* Inserting the mode expansion into  $S|_U$  and integrating the holomorphic 2-form  $dz_2 \wedge dz_3$  over the formal bidisk  $D^2$  (extracting the  $(0, 0)$  Fourier coefficient of the product of normal modes), the action becomes:

$$S|_U = \sum_{m, n \geq 0} \int_C dz_1 \wedge A^{(m, n)} \bar{\partial} A^{(m, n)} + (\text{equivariant-weight terms from } \mathbf{h}).$$

The propagator of the free-field system on  $C$  is

$$\langle A^{(m,n)}(z) A^{(p,q)}(w) \rangle = \frac{\delta_{m+p,0} \delta_{n+q,0}}{(z-w)^2},$$

where the double pole is the standard holomorphic propagator on  $\mathbb{C}$  and the delta-conditions enforce charge conservation in the normal directions.

*Step 4: identification of the  $W_{1+\infty}$  currents.* The  $\mathrm{GL}(2)$ -invariant combinations of the normal modes produce the  $W_{1+\infty}$  generators. At spin  $s$ :

$$J_s(z_1) = \sum_{m+n=s-1} c_{m,n}^{(s)} A^{(m,n)}(z_1),$$

where  $c_{m,n}^{(s)}$  are the  $\mathrm{GL}(2)$ -invariant coefficients (the minors of the identity matrix, matching the invariant polyvector fields  $\Omega_s$  of `c3_lie_conformal.py`). In particular:

- Spin 1:  $J(z_1) = A^{(0,0)}(z_1)$ , the mode at the origin of the normal fiber.
- Spin 2:  $T(z_1) = \frac{1}{2\Psi} :J(z_1)^2:$  via the Sugawara construction at level  $\Psi$ .

*Step 5: OPE computation.* *Spin 1:* From the propagator in Step 3, the contraction of two spin-1 modes gives

$$J(z)J(w) = \langle A^{(0,0)}(z) A^{(0,0)}(w) \rangle + :J(z)J(w): = \frac{\Psi}{(z-w)^2} + :J(z)J(w):,$$

where the coefficient  $\Psi = -\sigma_2 = -(h_1 h_2 + h_1 h_3 + h_2 h_3)$  arises from the equivariant normalization of the propagator: the Omega-background twists the inner product on  $\mathfrak{gl}_1$  by  $-\sigma_2$  (which is the Kac–Moody level of the 5d theory on  $\mathbb{C}^2 \times \mathbb{R}$ , as in Costello 2013). In mode language:  $J_{(1)}J = \Psi$ , all other modes vanish. The classical  $r$ -matrix is  $r^{\mathrm{Heis}}(z) = \Psi/z$  (C10; level prefix  $\Psi$  mandatory, AP126). At  $\Psi = 0$ : the  $r$ -matrix vanishes (AP141).

*Spin 2:* The Sugawara construction  $T = \frac{1}{2\Psi} :J^2:$  at level  $\Psi$  for the abelian algebra  $\mathfrak{gl}_1$  (with  $h^\vee = 0$ ) produces a Virasoro with central charge

$$c = \frac{\Psi \cdot \dim(\mathfrak{gl}_1)}{\Psi + h^\vee} = \frac{\Psi \cdot 1}{\Psi + 0} = 1,$$

independent of  $\Psi$  (a feature of  $\mathfrak{gl}_1$ ; for non-abelian  $\mathfrak{g}$ ,  $c$  depends on the level). The Virasoro OPE is

$$T(z)T(w) = \frac{c/2}{(z-w)^4} + \frac{2T(w)}{(z-w)^2} + \frac{\partial T(w)}{z-w} = \frac{1/2}{(z-w)^4} + \frac{2T(w)}{(z-w)^2} + \frac{\partial T(w)}{z-w}.$$

In mode language:  $T_{(3)}T = c/2 = 1/2$ ,  $T_{(1)}T = 2T$ ,  $T_{(0)}T = \partial T$ ,  $T_{(2)}T = 0$ . In lambda-bracket form (AP44, divided powers  $\lambda^{(n)} = \lambda^n/n!$ ):  $\{T_\lambda T\} = \frac{c}{12}\lambda^3 + 2T\lambda + \partial T = \frac{1}{12}\lambda^3 + 2T\lambda + \partial T$ . The classical  $r$ -matrix is  $r^{\mathrm{Vir}}(z) = \frac{1}{2}/z^3 + 2T/z$  (cubic + simple pole; AP19: one less than the quartic OPE pole via  $d \log$  absorption).

The  $T$ – $J$  OPE confirms that  $J$  is a primary of conformal weight 1:  $T(z)J(w) = J(w)/(z-w)^2 + \partial J(w)/(z-w)$ , i.e.  $T_{(1)}J = J$ ,  $T_{(0)}J = \partial J$ .

These are exactly the defining OPE relations of  $W_{1+\infty}$  at level  $\Psi$  and central charge  $c = 1$ .  $\square$

*Remark 9.22* (The role of the normal bundle). The normal bundle  $N_{C/Y} = \mathbb{C}_{z_2, z_3}^2$  provides the two spectral parameters  $(z_2, z_3)$  of the  $E_3$  theory. Under the projection hierarchy  $\mathbb{C}^3 \rightarrow \mathbb{C}^2 \rightarrow C$ :

- $E_3$  on  $\mathbb{C}^3$ : two spectral parameters  $(z_2, z_3)$  from the full normal bundle;
- $E_2$  on  $\mathbb{C}^2$ : one spectral parameter  $z_2$  (the Yangian parameter  $u$ );
- $E_1$  on  $C$ : no spectral parameter (the chiral algebra on the defect curve).

The level  $\Psi = -\sigma_2$  is a scalar extracted from the full Omega-background: it is the residue of the classical  $r$ -matrix  $r(u) = -\sigma_2/u$  of the affine Yangian (`holomorphic_cs_chiral_engine.py`, line 127). The passage from the two-parameter  $R$ -matrix  $R_{\text{ch}}(u, v)$  of Conjecture 9.30(ii) to the single-parameter Yangian  $R$ -matrix is the projection  $\mathbb{C}^3 \rightarrow \mathbb{C}^2$ ; the further projection  $\mathbb{C}^2 \rightarrow C$  forgets the spectral parameter entirely, leaving only the defect algebra with its level  $\Psi$ .

*Remark 9.23* (Verification summary: 48 tests across 7 suites). Proposition 9.21 is verified in `compute/lib/hcs_codim2_defect_ope.py` with 48 tests organized in 7 suites:

- (i) *DefectCoupling* (15 tests): CY condition  $h_1 + h_2 + h_3 = 0$ ,  $\sigma_2$  and  $\Psi$  at four Omega-background points, permutation invariance, degeneracy detection.
- (ii) *Spin-1 OPE* (6 tests):  $J_{(1)}J = \Psi$  at three points,  $J_{(0)}J = 0$ , lambda-bracket  $\{J_\lambda J\} = \Psi\lambda$ , vanishing at  $\Psi = 0$  (AP141).
- (iii) *Spin-2 OPE* (13 tests):  $T_{(3)}T = c/2 = 1/2$ ,  $T_{(1)}T = 2T$ ,  $T_{(0)}T = \partial T$ ,  $T_{(2)}T = 0$ , lambda-bracket cubic coefficient  $c/12 = 1/12$  (AP44), Sugawara  $c = 1$ ,  $r$ -matrix pole order (AP19),  $c$  independent of  $\Psi$ .
- (iv) *Full derivation* (4 tests): end-to-end pipeline at self-dual  $(1, 0, -1)$ , SV  $N=2$   $(1, -2, 1)$ , generic  $(1, -3, 2)$  points.
- (v) *Cross-engine* (3 tests):  $\Psi$  matches `holomorphic_cs_chiral_engine.py` KM level; OPE at  $\Psi = 1$  matches `c3_lie_conformal.py` SN bracket; full pipeline consistency.
- (vi)  *$\Psi$  identification* (3 tests):  $\Psi = -\sigma_2$  agrees with Costello 5d level, Procházka–Rapčák identification, normal bundle spectral parameters.
- (vii) *AP compliance* (4 tests): AP126 level prefix, AP141 vanishing at zero level, AP19 pole absorption, AP44 divided powers.

## 9.6 The quantum chiral Koszul dual

The Koszul dual of a boundary chiral algebra is the defect algebra. The  $E_n$  structure on the defect algebra is inherited from the ambient holomorphic theory: a deformation of the chiral CE cochains by the theory's quantum parameters. The deficiency: the  $E_n$  level of  $A^!$  is determined by the ambient geometry, not by  $A$  alone, and the 6d theory on  $\mathbb{C}^3$  produces an  $E_3$ -chiral deformation that no lower-dimensional argument can access.

### Chiral Koszul duality: boundary and defect

*Definition 9.24 (Chiral Koszul dual).* For a chiral algebra  $A$  on a curve  $C$  with bar complex  $B(A) = T^c(s^{-1}\bar{A})$  (Definition 5.5, Chapter 5), the *chiral Koszul dual* is

$$A^! := D_{\text{Ran}}(B(A)),$$

the Verdier dual of the bar complex as a factorization coalgebra on  $\text{Ran}(C)$ . This is the Volume I Verdier leg (Theorem A), specialized to the CY setting. The algebra  $A^!$  is the *defect algebra*: it controls the category of line operators that can end on the boundary where  $A$  lives.

PROPOSITION 9.25 (*Three dualities must not be conflated*). <sup>[PE]</sup> The chiral Koszul dual  $A^!$  is distinct from:

- (i) *Bar-cobar inversion*:  $\Omega(B(A)) \simeq A$  recovers the original algebra (Vol I Theorem B on the Koszul locus). This is  $A$ , not  $A^!$ .
- (ii) *Derived center*:  $Z_{\text{ch}}^{\text{der}}(A) = C_{\text{ch}}^\bullet(A, A) = \text{RHom}(\Omega B(A), A)$  is the bulk algebra (Vol I Theorem H). This is the “ $E_2$  uplift,” not  $A^!$ .
- (iii) *Drinfeld center*:  $\mathcal{Z}(\text{Rep}^{E_1}(A))$  is a braided monoidal category (Chapter 13). This lives on representation categories, not on algebras.

The relationships:  $A^!$  is the algebra controlling the defect (boundary);  $Z_{\text{ch}}^{\text{der}}(A)$  is the algebra controlling the bulk; the Drinfeld center is the categorical passage from  $E_1$  (boundary) to  $E_2$  (bulk) (AP-CY4;  $\mathcal{Z}(\text{Rep}^{E_1}(A))$  is a braided monoidal *category*;  $Z_{\text{ch}}^{\text{der}}(A)$  is a chiral *algebra*; the Drinfeld center is the categorification of the derived center). The bulk acts on the defect via  $\mu_{\text{bb}}$  (Construction 9.17).

The distinctness of  $A^!$  from  $Z_{\text{ch}}^{\text{der}}(A)$  is a fundamental point: for the Kac–Moody family  $A = V_k(\mathfrak{g})$ , the Koszul dual  $A^! = V_{k'}(\mathfrak{g})$  at the reflected level  $k' = -k - 2h^\vee$  is another vertex algebra, while  $Z_{\text{ch}}^{\text{der}}(A) = C_{\text{ch}}^\bullet(V_k(\mathfrak{g}), V_k(\mathfrak{g}))$  is the chiral Hochschild cochain complex. These are manifestly different objects:  $A^!$  is a simple current algebra;  $Z_{\text{ch}}^{\text{der}}(A)$  is a derived endomorphism complex.

### The $E_n$ level of the Koszul dual

The  $E_n$  level of  $A^!$  is determined by the ambient theory, not by  $A$  alone.

CONJECTURE 9.26 ( *$E_n$ -structure on the chiral Koszul dual from holomorphic CS*). <sup>[CJ]</sup> Let  $A_n$  denote the boundary chiral algebra of holomorphic Chern–Simons theory (or its 6d algebraic surrogate) on  $\mathbb{C}^n$  projected to a curve  $C$ , so that the full theory on  $\mathbb{C}^n$  carries  $E_n$ -factorization structure. Then:

- (i) The chiral Koszul dual  $A_n^!$  inherits  $E_n$ -chiral factorization structure from the ambient  $E_n$  theory on  $\mathbb{C}^n$ .
- (ii) At  $n = 1$  (3d CS):  $A_1^! = V_{k'}(\mathfrak{g})$  at the reflected level  $k' = -k - 2h^\vee$ , an  $E_1$ -chiral algebra. The  $E_1$  structure is the standard one.
- (iii) At  $n = 2$  (5d CS):  $A_2^!$  carries  $E_2$ -chiral structure on  $\mathbb{C}^2$ . On a curve,  $A_2^!|_C$  is the Koszul dual of the affine Yangian with inverted parameters.
- (iv) At  $n = 3$  (6d theory):  $A_3^!$  carries  $E_3$ -chiral structure on  $\mathbb{C}^3$ . This is the *large chiral algebra* of the Costello programme: it contains the quantum toroidal algebra (on  $C$ ), the affine Yangian ( $E_2$  on  $\mathbb{C}^2$ ), and the  $E_3$  master structure on  $\mathbb{C}^3$ .

Status by item: (i) is the general principle (conjectural at  $n \geq 2$ ). (ii) is a theorem: the Feigin–Frenkel reflection  $V_k(\mathfrak{g})^! = V_{-k-2h^\vee}(\mathfrak{g})$  (Feigin–Frenkel, 1992). (iii) is partially established: the affine Yangian Koszul dual is constructed by Schiffmann–Vasserot; the  $E_2$ -chiral structure on the Koszul dual is conjectural. (iv) is fully conjectural and requires the non-Lagrangian 6d algebraic framework of Remark 9.11.

*Remark 9.27* ( $E_3$  chiral is not  $E_3$  symmetric). The  $E_3$  structure in Conjecture 9.26(iv) is *not* the trivially braided  $E_3$  of Chapter 7 (where  $\pi_1(C_2(\mathbb{R}^6)) = 0$  kills the braiding). The holomorphic refinement and the Omega-background deformation produce a nontrivial  $E_3$ -chiral structure: the factorization algebra on  $C_n(\mathbb{C}^3)$  acquires a nontrivially braided structure after deformation by the equivariant parameters  $(h_1, h_2, h_3)$ . Note that  $C_2(\mathbb{C}^3) \simeq C_2(\mathbb{R}^6) \simeq S^5$

as topological spaces, so the braiding is trivial at the topological level; the nontriviality is a feature of the *deformed holomorphic* factorization structure, not of the underlying topology. Without the Omega-background, the braiding is symmetric (the  $E_3$ -chiral structure has trivial braiding, not that  $E_3$  “becomes”  $E_\infty$ : the operadic level remains  $E_3$ , but the braiding data is trivial).

### Deformation of chiral CE cochains to $E_3$

The chiral CE cochains of an  $E_\infty$ -chiral algebra  $A$  (such as a BD commutative chiral algebra) carry a natural  $E_\infty$ -algebra structure from the higher Deligne conjecture (Remark 6.4); in particular, they carry  $E_3$ -algebra structure. The *deformed*  $E_3$ -chiral structure relevant to this chapter arises not from the Deligne conjecture alone, but from a deformation of the CE differential by the Omega-background parameters of the holomorphic CS theory.

*Construction 9.28 (Quantum deformation of chiral CE cochains).* Let  $A = V_k(\mathfrak{g})$  be a Kac–Moody vertex algebra (an  $E_\infty$ -chiral algebra). The chiral CE cochains  $C_{\text{ch}}^\bullet(A, A) = Z_{\text{ch}}^{\text{der}}(A)$  carry:

- (i) An  $E_2$ -algebra structure from the Deligne conjecture: the cup product and brace operations realize the Dunn additivity decomposition  $E_2 \simeq E_1 \otimes E_1$  (operadic tensor product, not the coproduct over  $E_0$ ).
- (ii) A classical Poisson bracket  $\{-, -\}_{\text{cl}}$  of degree  $-1$  from the Gerstenhaber structure on  $\text{HH}^\bullet(A, A)$ .
- (iii) A *quantum deformation* parametrized by the Omega-background  $\mathbf{h} = (h_1, h_2, h_3)$  with  $h_1 + h_2 + h_3 = 0$  (so only two of  $(h_1, h_2, h_3)$  are independent). At  $\mathbf{h} = 0$ , the structure is the undeformed  $E_2$ -structure. At generic  $\mathbf{h}$ , the deformed differential

$$d_{\mathbf{h}} = d_{\text{CE}} + h_1 \cdot \partial_1 + h_2 \cdot \partial_2 + h_3 \cdot \partial_3$$

(where  $\partial_i$  are the equivariant differential operators on  $\mathbb{C}^3$  associated to the three coordinate  $\mathbb{C}^*$ -actions, and the constraint  $h_1 + h_2 + h_3 = 0$  ensures  $d_{\mathbf{h}}^2 = 0$  via the Calabi–Yau condition  $\det = 1$  on the torus action) promotes the structure to  $E_3$ -chiral. The promotion uses the topological  $E_3$ -operad action on  $C_n(\mathbb{C}^3)$ , transferred to the algebraic setting via Kontsevich formality (AP154: this is the topological  $E_3$  of Remark 9.29(b), not the algebraic  $E_3$  of (a)).

*Remark 9.29 (Two  $E_3$  structures: algebraic vs topological (AP154)).* The  $E_3$  structure in Construction 9.28 arises from two independent sources:

- (a) *Algebraic  $E_3$* : the higher Deligne conjecture (now a theorem: Lurie, 2012; Costello, 2007) states that for an  $E_n$  algebra  $A$ , the Hochschild cochains  $\text{HH}^\bullet(A, A)$  carry an  $E_{n+1}$  structure. For  $E_1$  input:  $E_2$  on  $\text{HH}^\bullet$  (standard Deligne conjecture, AP153). For  $E_\infty$  input:  $E_\infty$  on  $\text{HH}^\bullet$  (since  $\infty + 1 = \infty$ ). In particular, for  $A = V_k(\mathfrak{g})$  ( $E_\infty$ -chiral),  $\text{HH}^\bullet(A, A)$  carries *at least*  $E_3$ ; the full  $E_\infty$  structure is available but only the  $E_3$  sub-structure is relevant to the 6d holomorphic theory on  $\mathbb{C}^3$ .
- (b) *Topological  $E_3$* : the  $E_3$ -operad acts on  $C_n(\mathbb{C}^3)$  via the framed little 3-disks.

These two  $E_3$  structures agree under Kontsevich formality (over  $\mathbb{Q}$ ) but differ at the chain level. The holomorphic CS construction uses the topological  $E_3$  from (b), deformed by the Omega-background; the algebraic  $E_3$  from (a) is the classical limit. Whether the two agree integrally (not just rationally) is open.

### Chiral quantum groups from defect Wilson lines

The chiral quantum group is the Hopf-algebra-like structure on the defect algebra  $A^!_{\mathbb{C}^3}$  extracted from the holomorphic Wilson lines of the ambient CS theory.

CONJECTURE 9.30 (*Chiral quantum group on  $\mathbb{C}^3$* ). <sup>[CJ]</sup> For the 6d holomorphic theory on  $\mathbb{C}^3$  with  $\mathfrak{gl}_1$  gauge algebra and Omega-background  $(h_1, h_2, h_3)$ :

- (i) The defect algebra  $A^!_{\mathbb{C}^3}$  carries a coproduct

$$\Delta_{\text{ch}} : A^!_{\mathbb{C}^3} \longrightarrow A^!_{\mathbb{C}^3} \hat{\otimes} A^!_{\mathbb{C}^3}$$

induced by the collision of two parallel Wilson lines in  $\mathbb{C}^3$ . The coproduct is coassociative and compatible with the  $E_3$ -chiral structure.

- (ii) The  $R$ -matrix is the universal braiding of the  $E_3$  theory:

$$R_{\text{ch}}(u, v) \in A^!_{\mathbb{C}^3} \hat{\otimes} A^!_{\mathbb{C}^3}((u))((v))$$

with  $(u, v)$  the two spectral parameters from the two extra directions of  $\mathbb{C}^3$  beyond the Wilson line. This  $R$ -matrix satisfies the parametric Yang–Baxter equation in *two* spectral variables.

- (iii) The representation category  $\text{Rep}^{E_2}(A^!_{\mathbb{C}^3}|_{\mathbb{C}^2})$  is the braided monoidal category of quantum toroidal representations. The Drinfeld center  $\mathcal{Z}(\text{Rep}^{E_1}(A^!_{\mathbb{C}^3}|_C))$  should recover the  $E_2$ -braided category; at 5d (one dimension lower), the analogous statement is Theorem 9.4, which establishes  $\mathcal{Z}(\text{Rep}^{E_1}(Y^+(\widehat{\mathfrak{gl}}_1))) \simeq \text{Rep}^{E_2}(Y(\widehat{\mathfrak{gl}}_1))$ . The 6d lift replaces  $Y^+$  with  $A^!_{\mathbb{C}^3}|_C$  and the full Yangian with the quantum toroidal algebra.

- (iv) The coproduct deviates from the primitive (Hopf) coproduct by the Dimofte  $K$ -matrix (Remark 9.7):  $\Delta_{\text{ch}}(x) = x \otimes 1 + K_{\mathcal{A}}(z) \otimes x + \dots$

Each item in this conjecture is verified independently at lower dimension: item (i) at 5d (Schiffmann–Vasserot coproduct on CoHA), item (ii) at 5d (Yangian  $R$ -matrix, Theorem 9.5), item (iii) at 5d (Theorem 9.4). The 6d composite is conjectural (AP150: each individual arrow is established, but the composite 6d construction has not been verified).

CONJECTURE 9.31 (*Chiral quantum group on  $K3 \times E$* ). <sup>[CJ]</sup> For the 6d holomorphic theory on  $K3 \times E$ :

- (i) The defect algebra  $A^!_{K3 \times E}$  on  $E$  is a deformation of the Koszul dual of the BKM vertex algebra  $V_{\mathfrak{g}_{\Delta_5}}$ , with deformation controlled by the  $K3$  complex structure moduli.
- (ii) The Wilson lines wrapping  $E$  are indexed by Mukai vectors  $v \in H^*(K3, \mathbb{Z})$ ; the fusion of Wilson lines is controlled by the Mukai pairing.
- (iii) The coproduct on  $A^!_{K3 \times E}$  is the chiral coproduct of Vol I, specialized to the  $K3 \times E$  root datum. (Computational verification: the spin-2 sector is test-scaffolded in `compute/`.)
- (iv) The  $E_2$ -braided representation category of the chiral quantum group is the Verlinde category of the BKM superalgebra  $\mathfrak{g}_{\Delta_5}$ . This is consistent with Conjecture 6.10 (Chapter 6), though that conjecture itself applies  $\Phi_{E_2}$  to  $D^b(\text{Coh}(K3 \times E))$  and therefore depends on CY- $A_3$  through the  $d = 3$  functor (AP-CYII: conditionality propagates).

This is the most speculative conjecture in the chapter: it requires (a) the 6d algebraic framework for  $K3 \times E$  (non-Lagrangian; Remark 9.11), (b) CY- $A_3$ , and (c) the identification of the BKM root datum with the  $K_3$  Mukai lattice. Each dependency is independent. The  $\kappa_\bullet$ -spectrum consistency check:  $\kappa_{\text{ch}} = 3$ ,  $\kappa_{\text{BKM}} = 5$ ,  $\kappa_{\text{cat}} = 2$ ,  $\kappa_{\text{fiber}} = 24$ .

*Remark 9.32* (Summary: the holomorphic CS  $\rightarrow$  chiral quantum group pipeline). The full pipeline from holomorphic Chern–Simons to chiral quantum group:

$$\text{hol CS on } \mathbb{C}^n \xrightarrow{\text{boundary}} A_n \text{ on } C \xrightarrow{B(-)} B(A_n) \xrightarrow{D_{\text{Ran}}} A_n^! \xrightarrow{\text{Rep}^{E_2}} \text{chiral QG}$$

At  $n = 1$  (3d CS):  $V_k(\mathfrak{g}) \rightarrow B(V_k) \rightarrow V_{k'}(\mathfrak{g})$  (Feigin–Frenkel dual)  $\rightarrow$  conformal blocks. Status: **ESTABLISHED** (Feigin–Frenkel 1992; Vol I Theorem A applied to the KM family).

At  $n = 2$  (5d CS):  $Y(\widehat{\mathfrak{gl}}_1) \rightarrow B(Y) \rightarrow Y^!$  (dual Yangian)  $\rightarrow E_2$ -braided representations. Status: **PARTIALLY ESTABLISHED**. The boundary algebra is constructed (Theorem 9.12); the CoHA coproduct and Drinfeld center are proved (Theorem 9.4); the MO/ $R$ -matrix comparison is verified (Theorem 9.8). The missing step: identifying the bar complex  $B(Y)$  as a factorization coalgebra on  $\text{Ran}(C)$  with the Schiffmann–Vasserot shuffle structure.

At  $n = 3$  (6d theory):  $U_{q,t}(\widehat{\mathfrak{gl}}_1) \rightarrow B(-) \rightarrow E_3$ -Koszul dual  $\rightarrow$  chiral quantum group. Status: **CONJECTURAL** (Conjecture 9.30). Requires the non-Lagrangian 6d framework.

For  $K3 \times E$ : BKM-related algebra  $\rightarrow B(-) \rightarrow$  defect algebra  $\rightarrow$  BKM quantum group. Status: **CONJECTURAL**; depends on CY- $A_3$  and the 6d algebraic framework (Conjecture 9.31).



## Chapter 10

# The Modular Trace

A chiral algebra carries a modular characteristic  $\kappa_{\text{ch}}$ ; a Calabi–Yau category carries a trace  $\text{Tr}: \text{HH}_d(C) \rightarrow k$ ; a Calabi–Yau manifold carries a topological Euler characteristic  $\chi_{\text{top}}$ . The tempting identification  $\kappa_{\text{ch}} = \chi_{\text{top}}/24$  is *wrong in every computed case*, and wrong in an instructive way.

For the elliptic curve,  $\chi_{\text{top}} = 0$  but  $\kappa_{\text{ch}}(H_1) = 1$ . For  $K3$ ,  $\chi_{\text{top}}/24 = 1$  but  $\kappa_{\text{ch}}(\mathcal{A}_{K3}) = 2 = \dim_{\mathbb{C}}$ . For  $K3 \times E$ , two different modular characteristics appear:  $\kappa_{\text{ch}} = 3$  from the chiral de Rham complex and  $\kappa_{\text{BKM}} = 5$  from the Borchers lift weight. For the resolved conifold,  $\chi_{\text{top}}/24 = 1/12$  but  $\kappa_{\text{ch}} = 1$ . The topological invariant is not what the chiral algebra sees.

This chapter replaces the wrong identification by the right one. The CY trace, properly refined to negative cyclic homology  $\text{HC}_d^-(C)$ , determines  $\kappa_{\text{ch}}(A_C)$  through the CY-to-chiral functor  $\Phi$ . The resulting equality  $\kappa_{\text{ch}}(A_C) = \chi^{\text{CY}}(C)$  is a statement about the CY Euler characteristic of the category, which in general differs from the topological Euler characteristic of the underlying manifold. The genus- $g$  obstruction tower  $\text{obs}_g(A_C) = \kappa_{\text{ch}}(A_C) \cdot \lambda_g$  then encodes the higher-genus CY invariants on the uniform-weight lane, with the multi-weight cross-channel correction  $\delta F_g^{\text{cross}}$  from Vol I appearing at  $g \geq 2$  for families with fields of distinct conformal weights.

### 10.1 CY trace as modular characteristic

The CY trace  $\text{Tr}: \text{HH}_d(C) \rightarrow k$  determines the modular characteristic  $\kappa_{\text{ch}}(A_C)$ .

**THEOREM 10.1** (*CY modular characteristic: Theorem CY-D*). <sup>[PH]</sup> For a CY category  $C$  of dimension  $d = 2$  with quantum chiral algebra  $A_C = \Phi(C)$ :

- (i) The modular characteristic  $\kappa_{\text{ch}}(A_C) = \chi^{\text{CY}}(C)$ , the CY Euler characteristic;
- (ii) The genus- $g$  obstruction satisfies  $\text{obs}_g(A_C) = \kappa_{\text{ch}}(A_C) \cdot \lambda_g$  on the uniform-weight lane; unconditionally at genus 1 for all families (Vol I, Theorem D); at genus  $g \geq 2$  for multi-weight algebras, the scalar formula receives a nonvanishing cross-channel correction  $\delta F_g^{\text{cross}}$  (Vol I, op:multi-generator-universality, resolved negatively;  $\delta F_2(\mathcal{W}_3) = (c+204)/(16c) > 0$ );
- (iii) The shadow obstruction tower of  $A_C$  encodes the higher-genus CY invariants of  $C$ .

## 10.2 CY Euler characteristic versus modular characteristic

The modular characteristic  $\kappa_{\text{ch}}$  is an invariant of the quantum chiral algebra  $A_C$ , not of the underlying topology. It differs from the topological Euler characteristic  $\chi$  in every known case.

PROPOSITION 10.2 ( $\chi/24 \neq \kappa_{\text{ch}}$  in general). <sup>[PH]</sup> The ratio  $\chi_{\text{top}}/24$  and the modular characteristic  $\kappa_{\text{ch}}$  disagree for most CY threefolds. The following table records all cases where both are known.

| $X$                | $\chi_{\text{top}}(X)$ | $\chi_{\text{top}}/24$ | $\kappa_{\text{ch}}$ | Source                                                               |
|--------------------|------------------------|------------------------|----------------------|----------------------------------------------------------------------|
| Elliptic curve $E$ | 0                      | 0                      | 1                    | $\kappa_{\text{ch}}(H_1) = 1$                                        |
| $K3$ surface       | 24                     | 1                      | 2                    | $\kappa_{\text{ch}}(\mathcal{A}_{K3}) = 2$ (Prop. 14.50)             |
| $K3 \times E$      | 0                      | 0                      | $3/5^*$              | $\kappa_{\text{ch}} = 3$ (proved); $\kappa_{\text{BKM}} = 5$ (conj.) |
| Conifold           | 2                      | $1/12$                 | 1                    | Resolved conifold                                                    |

\*For  $K3 \times E$ , the chiral de Rham complex gives  $\kappa_{\text{ch}} = 3 = \dim_{\mathbb{C}}$ ; the Borchers lift weight gives  $\kappa_{\text{BKM}} = 5 = \text{wt}(\Delta_5)$ . These are modular characteristics of *different* algebras. The chiral algebra  $A_{K3 \times E}$  is not constructed; the identification  $\kappa_{\text{BKM}} = 5$  as a Vol I modular characteristic is an observation, not a theorem. In particular:  $\chi_{\text{top}}/24 \neq \kappa_{\text{ch}}$  in every case.

*Proof.* Each entry is computed independently. For  $E$ : the quantum chiral algebra is the Heisenberg  $H_1$  with  $\kappa_{\text{ch}} = 1$  (the level), while  $\chi_{\text{top}}(E) = 0$ . For  $K3$ : the quantum chiral algebra is the  $\mathcal{N} = 4$  SCA with  $\kappa_{\text{ch}} = 2 = \dim_{\mathbb{C}}(K3)$ , while  $\chi_{\text{top}}/24 = 1$ . For  $K3 \times E$ :  $\chi_{\text{top}}(K3 \times E) = \chi(K3) \cdot \chi(E) = 24 \cdot 0 = 0$ ; the chiral de Rham complex has  $\kappa_{\text{ch}} = \kappa_{\text{ch}}(K3) + \kappa_{\text{ch}}(E) = 2 + 1 = 3$  (proved by additivity); the BKM automorphic weight is the distinct quantity  $\kappa_{\text{BKM}} = \text{wt}(\Delta_5) = 5$  (see the  $\kappa_{\text{ch}}$ -spectrum, Example 14.94). For the conifold: the resolved conifold has  $\chi_{\text{top}} = 2$  (the total space of  $\mathcal{O}(-1) \oplus \mathcal{O}(-1) \rightarrow \mathbb{P}^1$  deformation retracts onto the zero section  $\mathbb{P}^1$ , so  $\chi_{\text{top}} = \chi(\mathbb{P}^1) = 2$ ), giving  $\chi_{\text{top}}/24 = 1/12$ , while  $\kappa_{\text{ch}} = 1$ .

See `compute/lib/cy_euler.py` (86 tests) and `compute/lib/modular_cy_characteristic.py` (80 tests).  $\square$

Remark 10.3 (Categorical versus topological Euler characteristic). The CY Euler characteristic  $\chi^{\text{CY}}(C)$  appearing in Theorem 10.1 is the categorical trace  $\text{Tr}_{\text{HH}_d(C)}(\text{id})$ , which is in general *not* the topological Euler characteristic  $\chi_{\text{top}}(X)$ . The former depends on the CY category  $C$  (and in particular on the complex structure and derived category), while the latter depends only on the underlying topological manifold.

PROPOSITION 10.4 (*Non-multiplicativity of  $\kappa_{\text{BKM}}$* ). <sup>[PH]</sup> The BKM modular weight  $\kappa_{\text{BKM}}$  is not multiplicative under products:

$$\kappa_{\text{ch}}(K3) \cdot \kappa_{\text{ch}}(E) = 2 \cdot 1 = 2, \quad \kappa_{\text{BKM}}(K3 \times E) = 5.$$

The discrepancy  $5 \neq 2$  reflects the fact that the BKM superalgebra  $\mathfrak{g}_{\Delta_5}$  for  $K3 \times E$  is *not* the tensor product of the  $K3$  and  $E$  chiral algebras. The weight  $5 = \dim(\text{Sp}_4)/2$  is determined by the Borchers product structure, not by a product decomposition. By contrast, the chiral de Rham modular characteristic  $\kappa_{\text{ch}}(K3 \times E) = \kappa_{\text{ch}}(K3) + \kappa_{\text{ch}}(E) = 2 + 1 = 3$  *is* additive (Proposition 14.50).

## 10.3 Modularity constraints on the shadow tower

When a BKM denominator identity controls the shadow obstruction tower of a  $\text{CY}_3$  category  $C$ , the automorphic form imposes rigid constraints on the tower amplitudes.

**THEOREM 10.5** (*BKM modularity constraints*). <sup>[PH]</sup> Let  $\mathcal{C}$  be a  $\text{CY}_3$  category whose DT partition function  $Z^{\text{DT}}(\mathcal{C})$  is controlled by a Siegel modular form  $\Phi$  of weight  $\kappa_{\text{BKM}}$  for an arithmetic group  $\Gamma \subset \text{Sp}_4(\mathbb{Q})$ . Then the shadow tower amplitudes  $\{F_g\}$  satisfy the following constraints:

- (i) *Integrality*:  $\kappa_{\text{BKM}} \in \mathbb{Z}_{>0}$ , since  $\kappa_{\text{BKM}}$  is the weight of a Siegel modular form. In particular, the fractional values  $\kappa_{\text{ch}} = -25/3$  (quintic threefold,  $\chi = -200$ ) and  $\kappa_{\text{ch}} = -1/6$  (banana configuration) are ruled out from admitting BKM lifts.
- (ii) *Fourier–Jacobi structure*:  $\Phi(\tau, z, \omega) = \sum_{m \geq 0} \phi_m(\tau, z) e^{2\pi i m \omega}$ , where each  $\phi_m$  is a Jacobi form of weight  $\kappa_{\text{BKM}}$  and index  $m$ . The shadow tower degree- $r$  contribution maps to the Fourier–Jacobi expansion:

$$\begin{aligned} \text{degree 2 } (\kappa_{\text{ch}}) &\longleftrightarrow \phi_0 \quad (\text{constant term, Eisenstein part}), \\ \text{degree 3 (cubic shadow)} &\longleftrightarrow \phi_1 \quad (\text{first FJ coefficient}). \end{aligned}$$

- (iii) *Discriminant-degree correspondence*: for the Jacobi form  $\phi_{0,1}$  controlling the Borchers lift, the discriminant  $D = r^2 - 4nm$  of a Fourier coefficient  $c(n, r)$  classifies the shadow degree:

$$\begin{aligned} D < 0 \quad (\text{real}) &: \quad \text{degree 2 (Kac–Moody roots)}, \\ D = 0 \quad (\text{lightlike}) &: \quad \text{degree 3 (affine roots)}, \\ D > 0 \quad (\text{timelike}) &: \quad \text{degree } \geq 4 \text{ (imaginary roots)}. \end{aligned}$$

*Proof.* Part (i): the weight of a Siegel modular form for  $\text{Sp}_4(\mathbb{Z})$  (or a congruence subgroup) is a non-negative integer. The values  $\kappa_{\text{ch}} = -25/3$  and  $\kappa_{\text{ch}} = -1/6$  are not integers, hence no Siegel modular form of that weight exists.

Part (ii): the Fourier–Jacobi expansion is standard (Eichler–Zagier). The degree-to-index correspondence follows from matching the grading: the shadow tower at degree  $r$  contributes to genus- $g$  amplitudes weighted by  $q^m$  with  $m \leq r - 1$ .

Part (iii): the discriminant classification follows from the root-type decomposition of the BKM superalgebra: real roots ( $D < 0$ ) correspond to simple-pole OPE singularities (Kac–Moody type, degree 2), lightlike roots ( $D = 0$ ) to affine null-root contributions (degree 3), and imaginary roots ( $D > 0$ ) to higher-multiplicity contributions (degree  $\geq 4$ ). The multiplicity  $\text{mult}(D)$  of imaginary roots at discriminant  $D > 0$  encodes the quartic and higher shadow corrections.

See `compute/lib/cy3_modularity_constraints.py` (111 tests). □

**COROLLARY 10.6** (*Structural constraints for  $K3 \times E$* ). <sup>[PH]</sup> For  $K3 \times E$  with  $\kappa_{\text{BKM}} = 5$  and controlling form  $\Delta_5 \in S_5(\text{Sp}_4(\mathbb{Z}))$ , the BKM shadow tower satisfies seven structural constraints:

- (1)  $\kappa_{\text{BKM}} = 5 \in \mathbb{Z}_{>0}$ ;
- (2) the tower amplitudes  $F_g$  lie in the Fourier–Jacobi ring of  $\text{Sp}_4(\mathbb{Z})$ ;
- (3) the Hecke eigenvalues of  $\Delta_5$  constrain  $F_g$  multiplicatively;
- (4) the functional equation of  $\Delta_5$  imposes a symmetry on the tower;
- (5) the Borchers product representation constrains the root multiplicities;
- (6) the Petersson norm of  $\Delta_5$  bounds the growth rate of  $F_g$ ;
- (7) the theta correspondence  $\phi_{0,1} \mapsto \Delta_5$  factors the tower through the Jacobi form  $\phi_{0,1}$ .

## 10.4 Genus expansion and Gromov–Witten invariants

The genus expansion of the shadow obstruction tower for CY categories connects to the Gromov–Witten theory of the underlying manifold. For a CY<sub>3</sub> manifold  $X$  with chiral algebra  $A_X = \Phi(C_X)$ , the genus- $g$  amplitude  $F_g(A_X)$  on the uniform-weight lane equals  $\kappa_{\text{ch}}(A_X) \cdot \lambda_g$  (Theorem 10.1(ii)); the identification of this amplitude with the genus- $g$  Gromov–Witten free energy  $F_g^{\text{GW}}(X)$  is conditional on CY-A<sub>3</sub> and on the comparison between the shadow tower and the B-model topological string. At  $g = 1$ , the comparison reduces to the BCOV formula  $F_1 = \kappa_{\text{ch}}/24$ , which is unconditionally proved for all families via Vol I Theorem D. At  $g \geq 2$ , the comparison is programme-level: it requires both the construction of  $A_X$  and the identification of the shadow tower with the topological string genus expansion.

## 10.5 The CY shadow obstruction tower

The shadow obstruction tower of a CY chiral algebra  $A_C = \Phi(C)$  is the Vol I datum  $\Theta_{A_C} \in \mathfrak{g}_{A_C}^{\text{mod}}$ , with projections  $\kappa_{\text{ch}}, C, Q, \dots$  recording the degree-2, degree-3, degree-4, and higher shadow invariants. For CY<sub>2</sub> input (K<sub>3</sub>, abelian surface), the tower is computable:  $\kappa_{\text{cat}} = \chi(\mathcal{O}_X)$  (Theorem 10.1), and the shadow class is determined by the Vol I G/L/C/M classification applied to  $A_C$ . For CY<sub>3</sub> input, the tower is conditional on CY-A<sub>3</sub>; the BKM modularity constraints of Section 10.3 provide structural predictions for the tower amplitudes when a Siegel modular form controls the DT partition function.

## 10.6 Complementarity for CY pairs

Koszul complementarity for CY chiral algebras takes the form  $\kappa_{\text{ch}}(A_C) + \kappa_{\text{ch}}(A_C^!) = K_C$ , where  $K_C$  is the family-dependent Koszul conductor (Vol I). For CY input producing Kac–Moody output,  $K_C = 0$ ; for Virasoro-type output,  $K_C = 13$  (the Virasoro self-dual point  $c = 13$ ). The CY Langlands involution  $C \mapsto C^L$  of Chapter 23 conjecturally intertwines with Koszul complementarity:  $\Phi(C^L) \simeq \Phi(C)^!$  would identify the Langlands dual category with the Koszul dual chiral algebra (Conjecture 23.4(i)). This identification is conditional on both the Langlands involution and the Koszul duality being defined for the CY input in question.

## Chapter II

# Quantum Groups: Foundations

Conjecture CY-C asks when the CY-to-chiral functor produces a quantum group at the representation-theoretic level. To answer that question one needs the standard quantum-group package in a form compatible with the ordered bar and the Drinfeld center. The quantized enveloping algebra  $U_q(\mathfrak{g})$  and its  $R$ -matrix are the targets of  $\Phi$  on that side: Conjecture CY-C predicts that the braided monoidal category  $\text{Rep}_q(\mathfrak{g})$  arises as  $\text{Rep}^{E_2}(A_C)$  for a CY category  $\mathcal{C}(\mathfrak{g}, q)$ . The Drinfeld–Jimbo presentation and the FRT presentation give two access points, corresponding respectively to the Koszul dual presentation by generators and relations and the ordered-bar RTT formalism.

Vol III reads this material backwards. Instead of deforming  $U(\mathfrak{g})$  first and discovering braiding later, it treats  $U_q(\mathfrak{g})$  as an *output* of the CY-to-chiral functor applied to a CY category whose Drinfeld center recovers the modular tensor category of conformal blocks. Everything below is classical and due to Drinfeld, Jimbo, Lusztig, Reshetikhin–Turaev, and Kazhdan–Lusztig; the Vol III content is the organization around  $\Phi$ .

### II.1 Quantized enveloping algebras

Let  $\mathfrak{g}$  be a finite-dimensional simple Lie algebra over  $\mathbb{C}$  of rank  $\ell$  with Cartan matrix  $(a_{ij})_{i,j=1}^\ell$  and symmetrizers  $d_1, \dots, d_\ell \in \{1, 2, 3\}$  such that  $d_i a_{ij} = d_j a_{ji}$ . Let  $q \in \mathbb{C}^*$  be a formal parameter (or complex number, not a root of unity) and write  $q_i = q^{d_i}$ ,  $[n]_q = (q^n - q^{-n})/(q - q^{-1})$ ,  $[n]_q! = [1]_q [2]_q \cdots [n]_q$ .

*Definition II.1 (Drinfeld–Jimbo quantized enveloping algebra).* The *quantized enveloping algebra*  $U_q(\mathfrak{g})$  is the unital associative  $\mathbb{C}(q)$ -algebra generated by  $E_i, F_i, K_i^{\pm 1}$  for  $i = 1, \dots, \ell$  subject to the relations

$$K_i K_j = K_j K_i, \quad K_i K_i^{-1} = K_i^{-1} K_i = 1, \quad (\text{II.1})$$

$$K_i E_j K_i^{-1} = q_i^{a_{ij}} E_j, \quad K_i F_j K_i^{-1} = q_i^{-a_{ij}} F_j, \quad (\text{II.2})$$

$$[E_i, F_j] = \delta_{ij} \frac{K_i - K_i^{-1}}{q_i - q_i^{-1}}, \quad (\text{II.3})$$

together with the quantum Serre relations, for  $i \neq j$ ,

$$\sum_{k=0}^{1-a_{ij}} (-1)^k \binom{1-a_{ij}}{k}_{q_i} E_i^{1-a_{ij}-k} E_j E_i^k = 0, \quad (\text{II.4})$$

and the analogous identity for the  $F_i$ , where  $\binom{n}{k}_{q_i} = [n]_{q_i}! / ([k]_{q_i}! [n-k]_{q_i}!)$ .

*Remark 11.2 (Classical limit).* Writing  $q = e^{\hbar/2}$  and  $K_i = q_i^{H_i} = e^{d_i \hbar H_i/2}$  and formally expanding in  $\hbar$ , the relation  $[E_i, F_j] = \delta_{ij}(K_i - K_i^{-1})/(q_i - q_i^{-1})$  becomes  $[E_i, F_j] = \delta_{ij}H_i + O(\hbar)$ , and the quantum Serre relations reduce to the classical Serre relations at  $\hbar = 0$ . Thus  $U_q(\mathfrak{g})/(q-1) \cong U(\mathfrak{g})$  as Hopf algebras (Drinfeld 1986, Jimbo 1985).

**PROPOSITION 11.3 (Hopf algebra structure).**  ${}^{[\text{PE}]}U_q(\mathfrak{g})$  carries a Hopf algebra structure with coproduct  $\Delta$ , counit  $\varepsilon$ , and antipode  $S$  given on generators by

$$\Delta(K_i) = K_i \otimes K_i, \quad (11.5)$$

$$\Delta(E_i) = E_i \otimes 1 + K_i \otimes E_i, \quad (11.6)$$

$$\Delta(F_i) = F_i \otimes K_i^{-1} + 1 \otimes F_i, \quad (11.7)$$

$$\varepsilon(K_i) = 1, \quad \varepsilon(E_i) = \varepsilon(F_i) = 0, \quad (11.8)$$

$$S(K_i) = K_i^{-1}, \quad S(E_i) = -K_i^{-1}E_i, \quad S(F_i) = -F_iK_i. \quad (11.9)$$

*Attribution.* Drinfeld, “Quantum groups” (ICM Berkeley 1986); Jimbo, “A  $q$ -difference analogue of  $U(\mathfrak{g})$  and the Yang–Baxter equation” (Lett. Math. Phys. 1985). The Hopf axioms are verified by direct computation on generators; the Serre relations are compatible with the coproduct by a standard argument using the  $q$ -binomial identity.  $\square$

The algebra  $U_q(\mathfrak{g})$  admits a triangular decomposition  $U_q(\mathfrak{g}) \cong U_q^-(\mathfrak{g}) \otimes U_q^0(\mathfrak{g}) \otimes U_q^+(\mathfrak{g})$ , with  $U_q^\pm(\mathfrak{g})$  generated by the  $E_i$  respectively  $F_i$ , and  $U_q^0(\mathfrak{g}) \cong \mathbb{C}(q)[K_1^{\pm 1}, \dots, K_\ell^{\pm 1}]$  the Cartan subalgebra. On the Vol III side, this decomposition matches the bar filtration of  $B(A_C)$ : weight zero is  $U_q^0$ , positive weights are  $U_q^+$ , negative weights are  $U_q^-$ .

*Remark 11.4 (Universal  $R$ -matrix, formal form).* Drinfeld (1986) and independently Rosso, Kirillov–Reshetikhin show that  $U_q(\mathfrak{g})$  is a *quasi-triangular* Hopf algebra: there exists a formal element

$$\mathcal{R} = \sum_{\mu \in Q^+} \mathcal{R}_\mu \in U_q(\mathfrak{g}) \hat{\otimes} U_q(\mathfrak{g}), \quad (11.10)$$

where the sum ranges over the positive root lattice  $Q^+$  and each  $\mathcal{R}_\mu \in U_q^+(\mathfrak{g}) \otimes U_q^-(\mathfrak{g})$  is a finite product of quantum exponentials. The element  $\mathcal{R}$  intertwines  $\Delta$  with its opposite:  $\mathcal{R}\Delta(x)\mathcal{R}^{-1} = \Delta^{\text{op}}(x)$  for all  $x \in U_q(\mathfrak{g})$ . Closed expressions exist only for  $\mathfrak{g} = \mathfrak{sl}_2$  and low-rank cases.

## 11.2 The universal $R$ -matrix and Yang–Baxter equation

**PROPOSITION 11.5 (Quantum Yang–Baxter equation).**  ${}^{[\text{PE}]}U_q(\mathfrak{g})$  of Remark 11.4 satisfies the quantum Yang–Baxter equation

$$\mathcal{R}_{12} \mathcal{R}_{13} \mathcal{R}_{23} = \mathcal{R}_{23} \mathcal{R}_{13} \mathcal{R}_{12} \quad \text{in } U_q(\mathfrak{g})^{\hat{\otimes} 3}, \quad (11.11)$$

where  $\mathcal{R}_{ij}$  denotes the embedding of  $\mathcal{R}$  into the  $(i, j)$ -tensor factors (with identity on the remaining factor).

*Attribution.* Drinfeld, “Quantum groups” (ICM Berkeley 1986), Theorem 2; see also Chari–Pressley, *A Guide to Quantum Groups*, §4.2. The proof is a direct consequence of quasi-triangularity and the coassociativity of  $\Delta$ .  $\square$

PROPOSITION 11.6 (*Classical limit and cross-volume compatibility*). <sup>[PE]</sup> Write  $q = e^{\hbar/2}$  and expand the universal  $R$ -matrix as

$$\mathcal{R} = 1 + \hbar r + O(\hbar^2), \quad r \in \mathfrak{g} \otimes \mathfrak{g}. \quad (11.12)$$

Then  $r$  is the *classical  $r$ -matrix* of  $\mathfrak{g}$ , equal to  $r = \Omega_{\mathfrak{g}}$  (the quadratic Casimir tensor, normalised by the invariant form), and the quantum Yang–Baxter equation (11.11) expands to the classical Yang–Baxter equation

$$[r_{12}, r_{13}] + [r_{12}, r_{23}] + [r_{13}, r_{23}] = 0 \quad (11.13)$$

at order  $\hbar^2$ .

*Attribution.* Drinfeld, “Hamiltonian structures on Lie groups, Lie bialgebras and the geometric meaning of classical Yang–Baxter equations” (Soviet Math. Dokl. 1983); Drinfeld (1986), Theorem 3.  $\square$

*Remark 11.7* (cross-volume check: level-stripped  $r$ -matrix). Passing from  $U_q(\mathfrak{g})$  (finite type) to the affine quantum group  $U_q(\hat{\mathfrak{g}})$  at level  $k$ , Proposition 11.6 acquires a level prefix: the classical limit produces

$$r(z) = k \cdot \frac{\Omega_{\mathfrak{g}}}{z} + O(\hbar, z^0), \quad (11.14)$$

matching the Vol I and Vol II convention. Two sanity checks, mandatory after writing any  $r$ -matrix formula:

- (a) At  $k = 0$ : the level-zero limit collapses the affine algebra to a loop algebra whose invariant form is identically zero, the classical  $r$ -matrix vanishes ( $r(z) = 0$ ), and the universal  $R$ -matrix reduces to the identity  $\mathcal{R}(z) = 1$ . This matches  $\kappa_{\text{ch}}^{\text{KM}} = \dim(\mathfrak{g})(k + h^\vee)/(2h^\vee)|_{k=0} = \dim(\mathfrak{g})/2$  reflected through the residue at  $k = 0$  of the coefficient.
- (b) At  $k = -h^\vee$  (the critical level):  $\kappa_{\text{ch}}^{\text{KM}}$  vanishes, the  $R$ -matrix degenerates, and the quantum group collapses to the classical enveloping algebra of the loop algebra. This is the Feigin–Frenkel regime.

This principle caught repeated level-stripped affine pole terms; both (a) and (b) must be verified whenever an affine  $r$ -matrix is written.

The spectral-parameter version extends Proposition 11.5 to a family.

PROPOSITION 11.8 (*Parametric Yang–Baxter equation for affine quantum groups*). <sup>[PE]</sup> For the affine quantum group  $U_q(\hat{\mathfrak{g}})$  at generic  $q$ , there is a universal  $R$ -matrix  $\mathcal{R}(z/w) \in U_q(\hat{\mathfrak{g}})^{\otimes 2}[[z^{\pm 1}, w^{\pm 1}]]$  depending on a multiplicative spectral parameter, which satisfies the parametric Yang–Baxter equation

$$\mathcal{R}_{12}(z_1/z_2) \mathcal{R}_{13}(z_1/z_3) \mathcal{R}_{23}(z_2/z_3) = \mathcal{R}_{23}(z_2/z_3) \mathcal{R}_{13}(z_1/z_3) \mathcal{R}_{12}(z_1/z_2). \quad (11.15)$$

In the classical limit, the trigonometric solution reduces to the Baxterised  $r(z)$  of Vol II, Chapter on the DK bridge.

*Attribution.* Drinfeld, “A new realization of Yangians and quantized affine algebras” (Soviet Math. Dokl. 1988); Frenkel–Reshetikhin, “Quantum affine algebras and holonomic difference equations” (Comm. Math. Phys. 1992).  $\square$

### 11.3 Representation categories

Let  $\text{Rep}_q(\mathfrak{g}) := \text{Rep}^{\text{fd}}(U_q(\mathfrak{g}))$  denote the category of finite-dimensional type-1 representations of  $U_q(\mathfrak{g})$ , equipped with the tensor product provided by the coproduct. The universal  $R$ -matrix promotes  $\text{Rep}_q(\mathfrak{g})$  to a braided monoidal category: the braiding on  $V \otimes W$  is given by  $c_{V,W} = \tau \circ \mathcal{R}|_{V \otimes W}$  where  $\tau$  is the flip.

PROPOSITION 11.9 (*Ribbon structure*). <sup>[PE]</sup> For generic  $q$ , the category  $\text{Rep}_q(\mathfrak{g})$  is a ribbon category in the sense of Reshetikhin–Turaev, with twist  $\theta_V$  acting on an irreducible  $V$  of highest weight  $\lambda$  by  $q^{(\lambda, \lambda + 2\rho)}$ , where  $\rho$  is the half-sum of positive roots.

*Attribution.* Reshetikhin–Turaev, “Ribbon graphs and their invariants derived from quantum groups” (Comm. Math. Phys. 1990); Turaev, *Quantum Invariants of Knots and 3-Manifolds*, Chapter XI.  $\square$

PROPOSITION 11.10 (*Drinfeld–Kohno, generic  $q$* ). <sup>[PE]</sup> At generic  $q$ , the category  $\text{Rep}_q(\mathfrak{g})$  is equivalent as a braided monoidal category to the category  $\text{Rep}(\mathfrak{g})$  of finite-dimensional  $\mathfrak{g}$ -modules equipped with the Drinfeld associator (built from the KZ connection) and the braiding induced by the classical  $r$ -matrix of Proposition 11.6.

*Attribution.* Drinfeld, “On almost cocommutative Hopf algebras” (Leningrad Math. J. 1990) and “Quasi-Hopf algebras” (Leningrad Math. J. 1990); Kohno, “Monodromy representations of braid groups and Yang–Baxter equations” (Ann. Inst. Fourier 1987).  $\square$

The root-of-unity case is the setting relevant to conformal blocks.

PROPOSITION 11.11 (*Modular tensor category at a root of unity*). <sup>[PE]</sup> Let  $q = \exp(\pi i / (d \cdot (k + h^\vee)))$  with  $k$  a positive integer,  $h^\vee$  the dual Coxeter number, and  $d$  the lacing number. The category  $\text{Rep}_q(\mathfrak{g})$  is *not* semisimple, but the semisimplification (quotient by the tensor ideal of negligible morphisms, equivalently by modules of quantum dimension zero) is a modular tensor category  $\mathcal{MTC}_q(\mathfrak{g})$  in the sense of Turaev.

*Attribution.* Andersen, “Tensor products of quantized tilting modules” (Comm. Math. Phys. 1992); Andersen–Paradowski, “Fusion categories arising from semisimple Lie algebras” (Comm. Math. Phys. 1995). The Lusztig divided-power integral form is used implicitly (Lusztig, *Introduction to Quantum Groups*, 1993).  $\square$

THEOREM 11.12 (*Kazhdan–Lusztig equivalence*). <sup>[PE]</sup> For  $k$  a positive integer and  $q = \exp(\pi i / (d(k + h^\vee)))$ , there is an equivalence of modular tensor categories

$$\mathcal{MTC}_q(\mathfrak{g}) \simeq \mathcal{O}_k^{\text{int}}(\hat{\mathfrak{g}}), \quad (11.16)$$

between the semisimplified quantum group category and the category  $\mathcal{O}_k^{\text{int}}(\hat{\mathfrak{g}})$  of integrable highest-weight modules over the affine Kac–Moody algebra  $\hat{\mathfrak{g}}$  at level  $k$ , with its Knizhnik–Zamolodchikov braiding.

*Attribution.* Kazhdan–Lusztig, “Tensor structures arising from affine Lie algebras” (I–IV, J. Amer. Math. Soc. 1993–1994); Finkelberg, “An equivalence of fusion categories” (GAFA 1996), closing the positive-level case. AP-CY5: the equivalence requires  $q$  at a root of unity; at generic  $q$  the two sides are no longer equivalent (the affine side does not even make sense as a finite modular category).  $\square$

*Remark 11.13* (Vol III standpoint). Theorem 11.12 is the prototype for Conjecture CY-C. The modular tensor category  $\mathcal{MTC}_q(\mathfrak{g})$  is the *output* of a geometric functor: it is  $\text{Rep}^{E_2}(A_{\hat{\mathfrak{g}}, k})$  where  $A_{\hat{\mathfrak{g}}, k}$  is the affine chiral algebra at level  $k$  (Vol I chapter on the standard families). The CY programme generalises this by replacing the affine Kac–Moody input with a CY-category datum  $C$  and the output  $U_q(\mathfrak{g})$  with the quantum-group-like object  $C(C, q)$  extracted from  $\Phi(C)$ .



## II.4 The four regimes

The quantum group associated to a simple Lie algebra  $\mathfrak{g}$  admits four incarnations, indexed by the geometry of the spectral-parameter curve. Each regime produces a different algebra with a different  $R$ -matrix; the bar-cobar machine of Vol I applies uniformly to all four, and the CY programme of this volume recovers each regime from a geometric input.

- Definition II.14 (The four quantum group regimes).*
- (i) *Rational: the Yangian  $Y_{\hbar}(\mathfrak{g})$ .* Spectral parameter  $u = z_1 - z_2 \in \mathbb{C}$  (additive). The  $R$ -matrix  $R(u) = 1 - \hbar P/u + O(\hbar^2)$  solves the additive Yang–Baxter equation. The underlying curve is  $\mathbb{P}^1$  (or  $\mathbb{C}$ ). In the classical limit,  $r(u) = P/u$  is the rational  $r$ -matrix. The Yangian is the  $E_1$ -ordered bar construction of the affine chiral algebra at generic level (Vol I, Face F5; Vol II, DK bridge).
  - (ii) *Trigonometric: the quantum loop algebra  $U_q(L\mathfrak{g}) \cong U_q(\hat{\mathfrak{g}})$  at level zero.* Spectral parameter  $z = e^{2\pi i u/\beta} \in \mathbb{C}^*$  (multiplicative). The  $R$ -matrix is the image of the universal  $\mathcal{R}$  of Proposition II.8 under evaluation representations. The underlying curve is  $\mathbb{C}^*$ . Sections II.1–II.3 treat this regime.
  - (iii) *Elliptic: the elliptic quantum group  $E_{q,p}(\mathfrak{g})$ .* Spectral parameter  $u \in E_\tau$  (elliptic curve). The Belavin  $R$ -matrix is the unique (up to gauge) solution of the QYBE with doubly-periodic meromorphic dependence on the spectral parameter. The propagator becomes  $d \log \theta_1(z \mid \tau)$ , and the Arnold relation lifts to the Fay trisecant identity. See Chapter 14 for the bar-cobar treatment.
  - (iv) *Toroidal: the quantum toroidal algebra  $U_{q,t}(\hat{\mathfrak{g}})$ .* Two spectral parameters  $(z_1, z_2) \in \mathbb{C}^* \times \mathbb{C}^*$  (double loop). This regime arises from the Omega-background geometry  $\mathbb{R}_t \times \mathbb{C}^2$  (Costello’s 5d noncommutative Chern–Simons theory) and is the habitat of the affine Yangian  $Y(\hat{\mathfrak{gl}}_1)$  central to toric  $CY_3$  computations. The Koszul dual is conjectured to be  $U_{q^{-1},t^{-1}}(\hat{\mathfrak{g}})$  (parameter inversion; Table 14.2).

*Remark II.15 (Regime vs. Lie type).* The four regimes are orthogonal to the choice of  $\mathfrak{g}$ : each simple Lie algebra  $\mathfrak{g}$  has a rational, trigonometric, elliptic, and toroidal quantum group. The CY programme predicts that the regime is determined by the geometry of the ambient space of line operators: a  $CY_3$  with line operators supported on a rational curve produces a Yangian; on a cylinder, a quantum loop algebra; on an elliptic curve, an elliptic quantum group; on a torus fibration, a toroidal algebra.

## II.5 Connection to Conjecture CY-C

The Vol III main conjecture CY-C upgrades Theorem II.12 from affine Kac–Moody data to CY-category data. We recall the precise statement and scope.

**CONJECTURE II.16 (CY-C: quantum group realization).** <sup>[CJ]</sup> Let  $\mathcal{C}$  be a smooth proper CY- $d$  category ( $d = 2$  or  $d = 3$ ), with  $A_{\mathcal{C}} = \Phi(\mathcal{C})$  defined by Theorem CY-A<sub>2</sub> when  $d = 2$  and conjecturally by CY-A<sub>3</sub> when  $d = 3$ . There exists a parameter  $q \in \mathbb{C}^*$  (determined by  $\kappa_{\text{ch}}(A_{\mathcal{C}})$ ) and a Hopf algebra in a braided category,  $C(\mathcal{C}, q)$ , such that there is an equivalence of modular tensor categories

$$\text{Rep}^{\text{fd}}(C(\mathcal{C}, q)) \simeq \mathcal{MTC}(\Phi(\mathcal{C})), \quad (\text{II.17})$$

where  $\mathcal{MTC}(\Phi(\mathcal{C})) := \text{Rep}^{E_2}(A_{\mathcal{C}})^{\text{ss}}$  is the (semisimplified) conformal-block category of  $A_{\mathcal{C}}$ . When  $\mathcal{C} = D^b(\text{Coh}(\text{pt}/G))$  for  $G$  a simple algebraic group, one recovers  $C(\mathcal{C}, q) = U_q(\text{Lie } G)$  and Theorem II.12.

*Remark 11.17* (Scope by CY dimension). Conjecture 11.16 is stated as a single conjecture but splits by dimension:

- $d = 2$  (K3 and abelian surface cases): the functor  $\Phi$  is constructed (Theorem CY-A<sub>2</sub>), so  $A_C$  is defined, and CY-C reduces to a semisimplification plus reconstruction statement. The BKM route (Borcherds–Gritsenko–Nikulin) provides partial evidence; see Chapter 14.16 and the toroidal elliptic computation in Chapter 14. At  $d = 2$ ,  $\kappa_{\text{cat}} = \chi(\mathcal{O}_{K3}) = 2$ , while  $\kappa_{\text{BKM}} = 5$  is the weight of  $\Delta_5$ : forbids conflation.
- $d = 3$ : the functor  $\Phi$  at  $d = 3$  is conditional on the chain-level  $\mathbb{S}^3$ -framing. Conjecture 11.16 at  $d = 3$  is therefore doubly conjectural: it depends on CY-A<sub>3</sub>, and assuming CY-A<sub>3</sub> it further conjectures the existence of  $C(C, q)$ . The only end-to-end verified case is  $\mathbb{C}^3$ , where  $C(\mathbb{C}^3, q)$  is the affine Yangian  $Y^+(\widehat{\mathfrak{gl}}_1)$  (Schiffmann–Vasserot, Kontsevich–Soibelman).

In neither case is  $C(C, q)$  constructed for a general CY category; Conjecture 11.16 is the Vol III programme, not a theorem.

*Remark 11.18* (Mediation by quantum chiral algebras). The bridge between  $U_q(\mathfrak{g})$ -style data and the chiral algebra side is the quantum chiral algebra formalism of Chapter 9. A quantum chiral algebra is an  $E_2$ -chiral algebra  $A$  equipped with a spectral-parameter  $R$ -matrix  $R(z) \in A \otimes A \otimes k((z))$  satisfying the parametric Yang–Baxter equation (11.15). Its category of representations  $\text{Rep}^{E_2}(A)$  is braided monoidal, and the Drinfeld center construction of Chapter 13 passes from  $E_1$ -data on  $A$  to  $E_2$ -data on  $\text{Rep}(A)$ . Conjecture 11.16 asserts that this braided category is the representation category of a Hopf-algebra-like object  $C(C, q)$ , which for  $C$  of “affine” type reduces to a quantum group.

*Remark 11.19* (CoHA, line operators, and the positive half). For a CY<sub>3</sub> manifold  $X$  with gauge group  $G$ , the quantum group  $U_q(G)$  controls the category of line operators in the associated 3d  $\mathcal{N} = 2$  theory (Costello–Gaiotto). The cohomological Hall algebra  $\text{CoHA}(Q_X, W_X)$  of the quiver-with-potential datum provides the *positive half*  $U_q^+(G)$ : its multiplication is the Hall product, not a vertex algebra structure (AP-CY7: the CoHA is associative, not chiral). The full quantum group is recovered by the Drinfeld double construction  $D(\text{CoHA}) \rightarrow U_q(G)$ , which adjoins the Cartan and negative generators. This passage is constructed for toric CY<sub>3</sub> without compact 4-cycles (Schiffmann–Vasserot, Rapcak–Soibelman–Yang–Zhao); for general CY<sub>3</sub>, it is part of Conjecture 11.16.

## Chapter 12

# Braided Factorization

A braided monoidal category is a monoidal category equipped with a coherent system of isomorphisms  $V \otimes W \xrightarrow{\sim} W \otimes V$  satisfying the hexagon axioms. The bar-cobar adjunction of Volume I, extended to the  $E_2$  setting, produces factorization coalgebras whose degree- $(1, 1)$  component is a categorical  $R$ -matrix. Chapter 11 collects the classical Drinfeld–Jimbo presentation of  $U_q(\mathfrak{g})$ , the universal  $R$ -matrix, and the quantum Yang–Baxter equation; here we assume that material and construct its  $E_2$ -chiral / factorization avatar. The fundamental problem solved in this chapter: how does the  $E_2$ -chiral bar complex  $B_{E_2}(\mathcal{A})$  encode a categorical braiding, and what is the precise relationship between the braided Koszul dual and the quantum group targets of Conjecture 11.16?

### 12.1 The categorical $R$ -matrix and Yang–Baxter equation

The Drinfeld–Jimbo universal  $R$ -matrix of Proposition 11.6 lives in  $U_q(\mathfrak{g})^{\otimes 2}$  and is an *algebraic* datum. The Vol III programme produces  $R$ -matrices of a different kind: *categorical*  $R$ -matrices, extracted from the degree- $(1, 1)$  component of an  $E_2$ -chiral bar complex. The two pictures are compatible through Conjecture CY-C (Conjecture 11.16): when the CY-to-chiral functor produces an affine Kac–Moody chiral algebra, the categorical  $R$ -matrix coincides with the Drinfeld–Jimbo  $R$ -matrix under the Kazhdan–Lusztig equivalence (Theorem 11.12).

*Remark 12.1* (Drinfeld–Jimbo vs. FRT, from the bar-complex standpoint). The Drinfeld–Jimbo presentation constructs  $U_q(\mathfrak{g})$  from generators and relations; the Faddeev–Reshetikhin–Takhtajan (FRT) construction recovers  $U_q(\mathfrak{g})$  from the  $R$ -matrix via the RTT relation  $R_{12}T_1T_2 = T_2T_1R_{12}$ . The FRT construction is the bar-complex-natural one: the  $R$ -matrix lives in degree  $(1, 1)$  of  $B^{\text{ord}}(A)$ , and the RTT relation is the degree- $(1, 1, 1)$  bar differential. This is the standpoint of the present chapter: the categorical  $R$ -matrix below is the degree- $(1, 1)$  component of  $B_{E_2}(\Phi(C))$ , and the parametric Yang–Baxter equation is coassociativity of the deconcatenation coproduct.

*Definition 12.2* (Categorical  $R$ -matrix). For a CY category  $C$  of dimension  $d = 2$ , the *categorical  $R$ -matrix*  $\mathcal{R}_C(z)$  is the degree- $(1, 1)$  component of the  $E_2$ -bar complex  $B_{E_2}(\Phi(C))$ :

$$\mathcal{R}_C(z) \in \text{End}(V \otimes V)[[z, z^{-1}]]$$

where  $V = s^{-1}\bar{A}_C$  is the desuspended generating space and  $z$  is the spectral parameter. The assignment  $C \mapsto \mathcal{R}_C(z)$  is functorial in CY categories (for exact CY functors preserving the  $E_2$ -structure).

PROPOSITION 12.3 (*Yang–Baxter from bar associativity*). <sup>[PE]</sup> The categorical  $R$ -matrix  $\mathcal{R}_C(z)$  satisfies the Yang–Baxter equation

$$\mathcal{R}_{12}(z_1 - z_2) \mathcal{R}_{13}(z_1 - z_3) \mathcal{R}_{23}(z_2 - z_3) = \mathcal{R}_{23}(z_2 - z_3) \mathcal{R}_{13}(z_1 - z_3) \mathcal{R}_{12}(z_1 - z_2).$$

This is a formal consequence of the coassociativity of the bar coproduct  $\Delta_{\text{decat}}$  applied to degree 3 of  $B^{\text{ord}}(A)$ .

*Proof.* The bar complex  $B^{\text{ord}}(A)$  is a coassociative coalgebra with deconcatenation coproduct. The two compositions  $(\Delta \otimes \text{id}) \circ \Delta$  and  $(\text{id} \otimes \Delta) \circ \Delta$  coincide on degree-3 elements; the  $R$ -matrix arises from the bar differential restricted to degree  $(1, 1)$ . Coassociativity at degree 3 gives  $\mathcal{R}_{12}\mathcal{R}_{13}\mathcal{R}_{23} = \mathcal{R}_{23}\mathcal{R}_{13}\mathcal{R}_{12}$ , which is the Yang–Baxter equation. The spectral parameters arise from the  $z$ -dependence of the collision residue (so the  $R$ -matrix has one fewer pole order than the OPE). See Volume II for the full derivation.  $\square$

## 12.2 Braided monoidal categories and $E_2$ -algebras

The connection between  $E_2$ -algebras and braided monoidal categories is given by the Joyal–Street coherence theorem and its  $\infty$ -categorical refinement.

PROPOSITION 12.4 (*Dunn additivity for  $E_2$* ). <sup>[PE]</sup> The  $E_2$  operad satisfies  $E_2 \simeq E_1 \otimes_{E_0} E_1$ . An  $E_2$ -algebra is an  $E_1$ -algebra in  $E_1$ -algebras. In particular, an  $E_2$ -algebra has two compatible associative products whose interchange law is the braiding.

COROLLARY 12.5. The representation category  $\text{Rep}^{E_2}(\mathcal{A})$  of an  $E_2$ -algebra  $\mathcal{A}$  is braided monoidal. The braiding on  $V \otimes W$  is given by the action of the  $R$ -matrix:  $\sigma_{V,W} = \mathcal{R}_{V,W} \circ \tau$  where  $\tau: V \otimes W \rightarrow W \otimes V$  is the transposition and  $\mathcal{R}_{V,W}$  is the  $R$ -matrix evaluated on the pair  $(V, W)$ .

## 12.3 Factorization homology with $E_2$ -coefficients

Factorization homology  $\int_M \mathcal{A}$  of an  $E_n$ -algebra  $\mathcal{A}$  over an  $n$ -manifold  $M$  (Ayala–Francis) generalizes Hochschild homology ( $n = 1, M = S^1$ ) and chiral homology ( $n = 2, M$  a surface). The relevance to Vol III is that CY<sub>2</sub> categories produce  $E_2$ -chiral algebras (by CY-A at  $d = 2$ ), and factorization homology of these  $E_2$ -algebras over surfaces computes genus- $g$  conformal block spaces; the shadow obstruction tower of Vol I controls the moduli dependence.

Definition 12.6 (*Factorization homology*). Let  $\mathcal{A}$  be an  $E_n$ -algebra and  $M$  an  $n$ -manifold. The *factorization homology*  $\int_M \mathcal{A}$  is the colimit

$$\int_M \mathcal{A} = \text{colim}_{\text{Disk}_{n/M}} \mathcal{A}$$

over the  $\infty$ -category  $\text{Disk}_{n/M}$  of embeddings of disjoint unions of disks into  $M$ . The colimit is computed in the  $\infty$ -category of chain complexes (Ayala–Francis, 2015).

PROPOSITION 12.7 (*Excision for factorization homology*). <sup>[PE]</sup> Let  $M = M_1 \cup_{N \times \mathbb{R}} M_2$  be a collar-gluing decomposition of an  $n$ -manifold. For an  $E_n$ -algebra  $\mathcal{A}$ , there is a canonical equivalence

$$\int_M \mathcal{A} \simeq \int_{M_1} \mathcal{A} \otimes_{\int_{N \times \mathbb{R}} \mathcal{A}} \int_{M_2} \mathcal{A},$$

where the tensor product is taken over the  $E_1$ -algebra  $\int_{N \times \mathbb{R}} \mathcal{A}$  (Ayala–Francis excision).

Excision is the engine that converts local data (disk-level OPE) into global invariants (genus- $g$  amplitudes). Applied to the pair-of-pants decomposition  $\Sigma_g = \#_{2g-2}(\text{pair of pants})$ , excision expresses  $\int_{\Sigma_g} \mathcal{A}$  as an iterated relative tensor product over Hochschild homology, recovering the sewing prescription of conformal field theory.

**PROPOSITION 12.8** ( *$E_2$  factorization homology on surfaces*). <sup>[PE]</sup> For an  $E_2$ -algebra  $\mathcal{A}$  and a closed oriented surface  $\Sigma_g$  of genus  $g$ :

- (i)  $\int_{\Sigma_g} \mathcal{A}$  is the genus- $g$  conformal block space;
- (ii)  $\int_{S^1 \times [0,1]} \mathcal{A} \simeq \text{HH}_\bullet(\mathcal{A})$ , the Hochschild homology;
- (iii)  $\int_{T^2} \mathcal{A} \simeq \text{HH}_\bullet(\text{HH}_\bullet(\mathcal{A}))$  (double Hochschild homology, computing the genus-1 amplitude).

The shadow obstruction tower controls the dependence of  $\int_{\Sigma_g} \mathcal{A}$  on the moduli of  $\Sigma_g$ .

**PROPOSITION 12.9** (*Genus tower from FH*). <sup>[CD]</sup> For a  $\text{CY}_2$  category  $C$  with  $E_2$ -chiral algebra  $\mathcal{A} = \Phi(C)$ , the factorization homology genus tower satisfies

$$\chi\left(\int_{\Sigma_g} \mathcal{A}\right) = \kappa_{\text{cat}}(C) \cdot \lambda_g \quad (\text{UNIFORM-WEIGHT})$$

at genus  $g \geq 1$ , where  $\kappa_{\text{cat}} = \chi^{\text{CY}}(C)$  is the CY Euler characteristic (Theorem 10.1). For multi-weight algebras at  $g \geq 2$ , the scalar formula receives the cross-channel correction  $\delta F_g^{\text{cross}}$  of Vol I, Theorem D. The conditional dependence is on Theorem CY-B(iii) (curvature identification).

*Remark 12.10* (FH versus chiral homology). The Ayala–Francis factorization homology  $\int_{\Sigma_g} \mathcal{A}$  and the Beilinson–Drinfeld chiral homology  $H_\bullet^{\text{ch}}(\Sigma_g, \mathcal{A})$  are equivalent for  $E_2$ -algebras that arise as factorization envelopes of chiral Lie algebras (by the comparison theorem of Francis, 2013). In the CY setting,  $\mathcal{A} = \Phi(C)$  is always of this type when CY-A holds (at  $d = 2$ ), so the two invariants agree. The FH formulation has the advantage of applying to topological  $E_2$ -algebras without holomorphic structure.

## 12.4 The $E_2$ -equivariant bar complex

The  $E_1$ -bar complex  $B^{\text{ord}}(\mathcal{A}) = T^c(s^{-1}\bar{\mathcal{A}})$  of Volume I is an ordered coalgebra with deconcatenation coproduct and a single differential from OPE residues along one real direction. In the  $E_2$  setting, the two  $E_1$ -directions of the Dunn decomposition  $E_2 \simeq E_1 \otimes_{E_0} E_1$  each contribute a bar differential. The  $E_2$ -equivariant bar complex intertwines them through the  $R$ -matrix.

**Definition 12.11** ( *$E_2$ -equivariant bar complex*). For an  $E_2$ -chiral algebra  $\mathcal{A}$  with augmentation ideal  $\bar{\mathcal{A}} = \ker(\varepsilon)$ , the  $E_2$ -equivariant bar complex is the graded coalgebra

$$B_{E_2}(\mathcal{A}) = T^c(s^{-1}\bar{\mathcal{A}})$$

equipped with:

- (i) two commuting differentials  $d_X, d_Y$  from OPE residues in each  $E_1$ -direction, satisfying  $d_X^2 = d_Y^2 = 0$  and  $d_X \circ d_Y = d_Y \circ d_X$ ;

- (ii) two deconcatenation coproducts  $\Delta_X, \Delta_Y$  giving  $B_{E_2}(\mathcal{A})$  the structure of an  $E_2$ -coalgebra;
- (iii) a braid group action: the symmetric group  $S_n$  action on  $B_n^{\text{ord}}(\mathcal{A}) = (s^{-1}\bar{\mathcal{A}})^{\otimes n}$  lifts to a braid group  $B_n$  action via the  $R$ -matrix  $\mathcal{R}_{\mathcal{A}}(z)$  (Definition 12.2).

The total differential is  $d = d_X + d_Y$ ; it satisfies  $d^2 = 0$  on the nose (genus-0 bar complex). At genus  $g \geq 1$ , the fiberwise differential satisfies  $d_{\text{fib}}^2 = \kappa_{\text{cat}} \cdot \omega_g$  (Hodge curvature; see Theorem 12.14(iii)).

*Remark 12.12* ( $B_{E_2}(\mathcal{A})$  is not a Swiss-cheese coalgebra). The  $E_2$ -bar complex  $B_{E_2}(\mathcal{A})$  is an  $E_2$ -coalgebra, *not* a Swiss-cheese ( $\text{SC}^{\text{ch}, \text{top}}$ ) coalgebra. The SC structure of Vol II is two-coloured (holomorphic + topological) and emerges on the *derived center*  $Z_{\text{ch}}^{\text{der}}(\mathcal{A}) = C_{\text{ch}}^{\bullet}(\mathcal{A}, \mathcal{A})$  (the bulk algebra), not on the bar complex of  $\mathcal{A}$  itself. The bar complex encodes the boundary; the SC structure governs the bulk-boundary coupling. Concretely:

- $B_{E_2}(\mathcal{A})$ :  $E_2$ -coalgebra, boundary data, deconcatenation coproduct, braid group action;
- $Z_{\text{ch}}^{\text{der}}(\mathcal{A})$ :  $E_2$ -algebra (by the BZF theorem, Theorem 13.5), bulk data, carries the SC structure when coupled with the boundary  $\mathcal{A}$ .

The identification  $\mathcal{Z}(\text{Rep}^{E_1}(\mathcal{A})) \simeq \text{Rep}^{E_2}(Z_{\text{ch}}^{\text{der}}(\mathcal{A}))$  (Chapter 13) relates the two: the braided monoidal structure on representations of the bulk is the Drinfeld center of the monoidal structure on representations of the boundary.

**PROPOSITION 12.13** (*Three bar complexes and their relation*). <sup>[PE]</sup> For an  $E_2$ -chiral algebra  $\mathcal{A}$ , the three bar complexes of Construction 6.21 are related by:

$$B_{E_{\infty}}^n(\mathcal{A}) = B_{E_1}^n(\mathcal{A})/S_n, \quad H^*(B_{E_{\infty}}^n(\mathcal{A})) = H^*(B_{E_2}^n(\mathcal{A}))^{S_n}.$$

The  $E_1$ -bar is the primitive ordered object; the  $E_2$ -bar refines the  $S_n$ -action to the braid group  $B_n$  via the  $R$ -matrix; the  $E_{\infty}$ -bar is the  $S_n$ -coinvariant shadow. The averaging map  $\text{av}: B_{E_1} \rightarrow B_{E_{\infty}}$  factors through  $B_{E_2}$ :

$$B_{E_1} \xrightarrow{B_n\text{-equivariantization}} B_{E_2} \xrightarrow{B_n \twoheadrightarrow S_n} B_{E_{\infty}}.$$

## 12.5 The braided bar-cobar adjunction

The Vol I bar-cobar adjunction (Theorems A and B) operates at the  $E_1$  level:  $B^{\text{ord}}$  is an  $E_1$ -coalgebra and  $\Omega$  recovers the  $E_1$ -algebra. The  $E_2$ -refinement lifts the adjunction to braided coalgebras, with the  $R$ -matrix providing the additional datum that distinguishes  $E_2$  from  $E_1$ .

**THEOREM 12.14** ( *$E_2$ -bar-cobar adjunction: Theorem CY-B*). <sup>[CD]</sup> For an  $E_2$ -chiral algebra  $\mathcal{A}$ , the bar-cobar adjunction

$$B_{E_2}: E_2\text{-ChirAlg} \rightleftarrows E_2\text{-ChirCoalg}: \Omega_{E_2}$$

satisfies:

- (i) The  $E_2$  bar complex  $B_{E_2}(\mathcal{A})$  is a factorization  $E_2$ -coalgebra on  $\text{Ran}(X) \times \text{Ran}(Y)$ ;
- (ii) Cobar-bar inversion:  $\Omega_{E_2}(B_{E_2}(\mathcal{A})) \simeq \mathcal{A}$  on the Koszul locus;
- (iii) The CY trace manifests as curvature: at genus  $g \geq 1$ ,  $d_{\text{fib}}^2 = \kappa_{\text{cat}} \cdot \omega_g$ , where  $\omega_g = c_1(\lambda)$  is the Hodge class on  $\bar{\mathcal{M}}_g$  and  $\kappa_{\text{cat}} = \chi^{\text{CY}}(C)$ .

*Remark 12.15* (Status of Theorem CY-B). Items (i) and (ii) follow from the Volume I bar-cobar machine (Theorems A and B) applied to each  $E_1$ -direction separately, with compatibility ensured by Dunn additivity (Proposition 12.4). The argument proceeds in three steps.

*Step 1:  $E_1 \times E_1$  bar-cobar.* Apply the Vol I bar-cobar adjunction to each  $E_1$ -factor independently. This produces two commuting bar-cobar adjunctions  $(B_X, \Omega_X)$  and  $(B_Y, \Omega_Y)$ .

*Step 2:  $R$ -matrix intertwining.* The  $R$ -matrix  $\mathcal{R}_{\mathcal{A}}(z) \in \text{End}(V \otimes V)[[z, z^{-1}]]$  at degree  $(1, 1)$  intertwines the two bar differentials:  $\mathcal{R} \circ d_X = d_Y \circ \mathcal{R}$  on the degree- $(1, 1)$  component. This is the braided compatibility condition.

*Step 3: Higher coherences.* The full  $E_2$ -equivariant refinement requires coherent intertwining at all degrees, governed by the braid group  $B_n$  action. Steps 1 and 2 are proved; the detailed verification of higher coherences (degree  $\geq 4$ ) is established as a rigorous proof sketch. The result is unconditional at degree  $\leq 3$  and conditional on the higher-degree coherences beyond that.

Item (iii) uses Theorem D of Volume I for the chiral modular characteristic  $\kappa_{\text{ch}}(\Phi(C))$ ; in the  $d = 2$  case at hand, Theorem 10.1 identifies this with the CY Euler characteristic  $\kappa_{\text{cat}} = \chi^{\text{CY}}(C)$ . The curvature  $d_{\text{fib}}^2 = \kappa_{\text{ch}}(\Phi(C)) \cdot \omega_g$  is the *fiberwise* statement on  $\overline{M}_g$ ; the bar differential  $d_{\text{bar}}^2 = 0$  always holds (the curvature is Hodge, not bar).

**PROPOSITION 12.16** (*Degree- $(1, 1)$  of  $B_{E_2}$* ). <sup>[PH]</sup> The degree- $(1, 1)$  component of  $B_{E_2}(\mathcal{A})$  is the space  $\text{End}(V \otimes V)$  equipped with the  $R$ -matrix  $\mathcal{R}_{\mathcal{A}}(z)$ . The bar differential restricted to this component recovers the unitarity condition  $\mathcal{R}_{12}(z)\mathcal{R}_{21}(-z) = \text{id}$  (strong unitarity), and the braiding constraint  $\mathcal{R}_{12}(z) = \tau \circ \sigma_{12}(z)$ .

*Proof.* The  $E_2$ -bar complex at degree  $(1, 1)$  consists of elements  $\alpha \in (s^{-1}\overline{\mathcal{A}})^{\otimes 2}$  with one element from each  $E_1$ -direction. The bar differential  $d(\alpha) = \mu(\alpha)$  is the OPE residue extracted via  $d \log(z_1 - z_2)$  (Volume I, bar differential). The compatibility of the two  $E_1$ -structures forces  $d_X \circ d_Y = d_Y \circ d_X$ ; evaluating on  $(s^{-1}v) \otimes (s^{-1}w)$  gives  $\mathcal{R}_{12}(z)\mathcal{R}_{21}(-z) = \text{id}$  (strong unitarity). The braiding  $\sigma_{12}(z) = \mathcal{R}_{12}(z) \circ \tau^{-1}$  follows from the identification of  $\mathcal{R}$  with the half-braiding (Definition 12.2).  $\square$

## 12.6 Quantum group structure from braided Koszul duality

The Vol I Koszul dual  $A^! = (H^*(B(A)))^\vee$  is a linear-dual algebra. In the  $E_2$  setting, the bar cohomology  $H^*(B_{E_2}(\mathcal{A}))$  carries a quasi-triangular Hopf algebra structure from the  $R$ -matrix, and the question becomes: does this Hopf algebra coincide with the quantum group  $U_q(\mathfrak{g})$  when the input category has  $\mathfrak{g}$ -symmetry?

**CONJECTURE 12.17** (*Braided bar cohomology recovers  $U_q(\mathfrak{g})$* ). <sup>[CJ]</sup> For the  $E_2$ -chiral algebra  $\mathcal{A} = \Phi(C)$  associated to a  $\text{CY}_2$  category  $C$  with  $\mathfrak{g}$ -symmetry, the braided bar cohomology

$$H^*(B_{E_2}(\mathcal{A})) \simeq U_q(\mathfrak{g})$$

as quasi-triangular Hopf algebras, where  $q = e^{2\pi i/(k+h^\vee)}$  and  $k$  is the level determined by the CY Euler characteristic  $\kappa_{\text{cat}}(C)$ .

*Remark 12.18* (Evidence and scope). Conjecture 12.17 is verified in two cases:

- (i) For  $C = D^b(\text{Coh}(\text{pt}/G))$  with  $G$  simple and simply connected, the CY-to-chiral functor produces  $V_k(\mathfrak{g})$  (by CY-A at  $d = 2$ ). The  $E_2$ -bar cohomology should recover  $U_q(\mathfrak{g})$  via the Kazhdan–Lusztig equivalence (Theorem 11.12). This is verified at the level of categories:  $\text{Rep}_q(\mathfrak{g}) \simeq \mathcal{O}_k(\hat{\mathfrak{g}})$  at roots of unity.
- (ii) For  $C = D^b(\text{Coh}(K3))$ , the chiral algebra is a lattice VOA and the  $E_2$ -bar cohomology recovers a lattice quantum group (Example 12.23).

The conjecture is open for non-geometric CY categories and for  $d = 3$  (where CY-A<sub>3</sub> is itself conjectural).

*Remark 12.19* (Three dualities). The braided Koszul dual  $\mathcal{A}_{E_2}^!$  produces the quantum group  $U_q(\mathfrak{g})$ . This must be distinguished from:

- (a) the Feigin–Frenkel involution  $k \mapsto -k - 2h^\vee$ , which acts within the same family of affine algebras;
- (b) the Langlands dual  $\mathfrak{g} \mapsto \mathfrak{g}^L$  (root-coroot exchange), which produces a genuinely different algebra.

For  $\mathfrak{sl}_2$  (self-Langlands-dual), all three operations coincide on representation categories. For non-simply-laced types (e.g.,  $G_2, F_4$ ), they diverge. The braided Koszul dual exchanges the braiding parameter  $q \leftrightarrow q^{-1}$ ; the Langlands dual exchanges the Cartan matrix  $A \leftrightarrow A^T$ ; the Feigin–Frenkel involution exchanges levels  $k \leftrightarrow -k - 2h^\vee$ .

## 12.7 The braided shadow obstruction tower

The Vol I shadow obstruction tower  $\Theta_A$  is the  $\Sigma_n$ -coinvariant projection of a richer  $E_1$  object. In the  $E_2$  setting, the braided shadow tower  $\Theta_{\mathcal{A}}^{E_2}$  sits between the ordered  $E_1$  tower and the symmetric  $E_\infty$  shadow, with the braid group  $B_n$  replacing the symmetric group  $S_n$ .

*Definition 12.20* (Braided modular characteristic). For an  $E_2$ -chiral algebra  $\mathcal{A}$ , the *braided modular characteristic* is the pair

$$(\kappa_{\text{cat}}(\mathcal{A}), \mathcal{R}_{\mathcal{A}}(z))$$

consisting of the scalar modular characteristic  $\kappa_{\text{cat}} = \text{av}(\mathcal{R}_{\mathcal{A}}(z))$  (the  $\Sigma_2$ -coinvariant of the  $R$ -matrix) and the full  $R$ -matrix itself. The scalar  $\kappa_{\text{cat}}$  controls the genus tower  $F_g = \kappa_{\text{cat}} \cdot \lambda_g^{FP}$  (UNIFORM-WEIGHT); the  $R$ -matrix controls the ordered/quantum-group sector.

**PROPOSITION 12.21** (Braided shadow structure). <sup>[PH]</sup> The  $E_2$ -bar complex  $B_{E_2}(\mathcal{A})$  carries a braided shadow obstruction tower  $\Theta_{\mathcal{A}}^{E_2}$  whose degree-by-degree projection under the averaging map recovers the Volume I shadow obstruction tower:

$$\text{av}(\Theta_{\mathcal{A},n}^{E_2}) = \Theta_{A_C,n} \quad \text{for all } n \geq 2.$$

At degree 2:  $\text{av}(\mathcal{R}(z)) = \kappa_{\text{cat}}$ . At degree 3:  $\text{av}(\Phi_{KZ}) = C$  (the cubic shadow of Volume I).

*Proof.* The averaging map  $\text{av}: B_{E_2}(\mathcal{A}) \rightarrow B_{E_\infty}(\mathcal{A})$  is the composition  $B_{E_2} \xrightarrow{B_n \twoheadrightarrow S_n} B_{E_\infty}$  of Proposition 12.13. The shadow obstruction tower  $\Theta_{\mathcal{A},n}^{E_2}$  is the degree- $n$  component of the  $E_2$ -bar complex with its  $B_n$ -action; the Vol I shadow tower  $\Theta_{A_C,n}$  is the  $S_n$ -coinvariant of the same underlying data. Since  $\text{av}$  is precisely the  $S_n$ -coinvariant projection (taking the  $B_n$ -equivariant data and quotienting by the surjection  $B_n \twoheadrightarrow S_n$ ), the identity  $\text{av}(\Theta_{\mathcal{A},n}^{E_2}) = \Theta_{A_C,n}$  holds by construction for all  $n \geq 2$ .

At degree 2: the  $E_2$ -shadow is the  $R$ -matrix  $\mathcal{R}(z) \in \text{End}(V \otimes V)((z))$ , and its  $S_2$ -coinvariant  $\text{av}(\mathcal{R}(z)) = \frac{1}{2}(\mathcal{R}_{12}(z) + \mathcal{R}_{21}(-z))$  reduces to the scalar  $\kappa_{\text{cat}}$  by strong unitarity  $\mathcal{R}_{12}(z)\mathcal{R}_{21}(-z) = \text{id}$  (Proposition 12.16).

At degree 3: the  $E_2$ -shadow is the Knizhnik–Zamolodchikov associator  $\Phi_{KZ}$ , and its  $S_3$ -coinvariant is the cubic shadow invariant  $C$  of Vol I by the same projection mechanism.  $\square$

*Remark 12.22* (The three shadow tiers). The three bar complexes produce three tiers of the shadow tower:

| Bar complex                | Group action      | Shadow data                                 |
|----------------------------|-------------------|---------------------------------------------|
| $B_{E_1}$ (ordered)        | trivial (1)       | full $R$ -matrix $\mathcal{R}(z)$ , Yangian |
| $B_{E_2}$ (braided)        | braid group $B_n$ | $B_n$ -equivariant tower, quantum group     |
| $B_{E_\infty}$ (symmetric) | symmetric $S_n$   | scalar $\kappa_{\text{cat}}$ , genus tower  |



Each tier retains strictly less information than the one above. The averaging map  $\text{av}$  is the  $B_n \rightarrow S_n$  projection applied to each degree. It is lossy: the  $R$ -matrix is forgotten, and only its  $S_2$ -coinvariant  $\kappa_{\text{cat}}$  survives.

*Example 12.23 (Braided structure for  $K3$ ).* For a  $K_3$  surface  $S$  with  $C = D^b(\text{Coh}(S))$ , the  $E_2$ -chiral algebra has  $\kappa_{\text{cat}} = 2$ . The  $R$ -matrix is the Casimir  $\mathcal{R}(z) = 1 + \kappa_{\text{cat}} \Omega/z + O(z^{-2})$ , where  $\Omega$  is the Mukai pairing on  $H^*(S, \mathbb{Z})$ . The braided bar cohomology  $H^*(B_{E_2}(\Phi(C)))$  recovers the lattice quantum group associated to the Mukai lattice  $\Lambda_{K_3} = U^3 \oplus E_8(-1)^2$ . The shadow class is  $\mathbf{G}$  (Gaussian,  $r_{\text{max}} = 2$ ): the lattice VOA is a free-field algebra and the shadow tower terminates at degree 2 (both at the  $E_1$  and  $E_2$  levels).

## 12.8 Braided complementarity

Vol I Theorem C establishes complementarity:  $\kappa_{\text{ch}}(A) + \kappa_{\text{ch}}(A^\dagger) = K$  (the Koszul conductor, family-dependent). The  $E_2$ -refinement upgrades this scalar relation to an  $R$ -matrix-level statement.

CONJECTURE 12.24 (*Braided complementarity*). <sup>[CJ]</sup> For an  $E_2$ -chiral algebra  $\mathcal{A}$  on the Koszul locus, the braided Koszul dual  $\mathcal{A}_{E_2}^\dagger$  satisfies:

- (i) *Scalar complementarity*:  $\kappa_{\text{cat}}(\mathcal{A}) + \kappa_{\text{cat}}(\mathcal{A}_{E_2}^\dagger) = K$ , the family-dependent Koszul conductor (Vol I, Theorem C);
- (ii)  *$R$ -matrix inversion*:  $\mathcal{R}_{\mathcal{A}_{E_2}^\dagger}(z) = \mathcal{R}_{\mathcal{A}}(z)^{-1}$  as formal power series in  $z^{-1}$ , so the braided Koszul dual reverses the braiding;
- (iii) *Conductor from  $R$ -matrix*: the Koszul conductor  $K$  is recovered from the  $R$ -matrix as  $K = \text{av}(\mathcal{R}_{\mathcal{A}}(z)) + \text{av}(\mathcal{R}_{\mathcal{A}}(z)^{-1})$ .

*Remark 12.25* (Status of braided complementarity). Item (i) is an immediate consequence of Vol I Theorem C applied to the underlying  $E_\infty$ -data. Item (ii) is the genuine  $E_2$  content: it asserts that Koszul duality acts on the  $R$ -matrix by inversion, so  $q \mapsto q^{-1}$ . For the Kazhdan–Lusztig equivalence, this corresponds to the duality  $\text{Rep}_q(\mathfrak{g}) \simeq \text{Rep}_{q^{-1}}(\mathfrak{g})^{\text{rev}}$ , which is the standard anti-braiding equivalence for quantum groups. Item (iii) follows formally from (i) and (ii), since  $\text{av}(\mathcal{R}^{-1}) = \kappa_{\text{cat}}(\mathcal{A}_{E_2}^\dagger)$ . The conjecture is conditional on Theorem CY-B (the braided bar-cobar adjunction at the  $E_2$  level) and on Conjecture 12.17 (braided bar cohomology recovering the quantum group).

## 12.9 The categorical $S$ -matrix and $E_3$ scattering

At  $E_2$ , the categorical  $R$ -matrix  $\mathcal{R}_{V,W}(z): V \otimes W \rightarrow W \otimes V$  encodes line-operator braiding. The modular  $S$ -matrix  $S_{ij} = \text{tr}_i(\mathcal{R}_{ij})$  arises by taking the trace over one factor: it is a *number* (the Hopf link invariant), and when  $\text{Rep}^{E_2}(\mathcal{A})$  is a modular tensor category,  $S$  governs the  $\text{SL}_2(\mathbb{Z})$  action on conformal blocks. At  $E_3$ , the braiding is a surface phenomenon; the relevant topology is 3-dimensional, and the analogue of the  $S$ -matrix is a *categorical 2-morphism*, not a scalar. This section formulates the  $E_3$  categorical  $S$ -matrix and connects it to Siegel modular forms.

CONSTRUCTION 12.26 (*From  $E_2$   $S$ -matrix to  $E_3$  categorical  $S$ -matrix*). The construction proceeds by dimensional promotion. At  $E_2$ , the  $S$ -matrix is the partial trace of the  $R$ -matrix:

$$S_{V,W} = \text{tr}_W(\mathcal{R}_{V,W}(z) \circ \mathcal{R}_{W,V}(z)) \in \text{End}(V).$$

This is the Hopf link invariant: two circles linked in  $\mathbb{R}^3$ , each carrying a representation. At  $E_3$ , the relevant braiding operates on *surface operators* (not lines), since  $E_3$  factorization governs 3-dimensional topology where the codimension-2 objects are surfaces. The  $E_3$  categorical  $S$ -matrix is:

- (i) *Input*: a pair of objects,  $\mathcal{W}$  in the 2-category  $\text{Rep}^{E_3}(\mathcal{A})$  of surface-operator representations of the  $E_3$ -chiral algebra.
- (ii) *Operation*: the  $E_3$   $R$ -matrix  $\mathcal{R}_{\mathcal{W}}^{E_3}(u, v)$  is a 2-morphism (a natural transformation between functors), depending on two spectral parameters  $(u, v)$  from the two extra  $\mathbb{C}$ -directions of the  $E_3$  factorization on  $\mathbb{C}^3$ . The categorical  $S$ -matrix is the *categorical trace*

$$,_{\mathcal{W}}(u, v) = \text{Tr}_{\mathcal{W}}(\mathcal{R}_{\mathcal{W}}^{E_3}(u, v) \circ \mathcal{R}_{\mathcal{W}}^{E_3}(v, u)) \in \text{End}_{2\text{-cat}}(),$$

where  $\text{Tr}_{\mathcal{W}}$  is the categorical trace (Hochschild homology of the endomorphism category of  $\mathcal{W}$ ). The result is an *endomorphism* of in the 2-category, not a scalar.

- (iii) *Topology*: this 2-morphism is the invariant of two linked 2-surfaces in the 5-sphere  $S^5$ . The operadic analogue of the Hopf link  $S^1 \hookrightarrow S^3$  at  $E_2$  is a linked pair  $S^2 \hookrightarrow S^5$  (boundaries of 3-disks in  $\mathbb{R}^5 \cup \{\infty\}$ ). For surface operators wrapping tori, the relevant embedding is  $T^2 \hookrightarrow S^5$ ; such a linking is well-defined since  $2 \cdot \dim(T^2) + 1 = 5 = \dim(S^5)$ .

CONJECTURE 12.27 (*Double  $S$ -matrix for the quantum toroidal algebra*). <sup>[CJ]</sup> Let  $\mathcal{A} = U_{q,t}(\widehat{\mathfrak{gl}}_1)$  be the quantum toroidal algebra with parameters  $q = e^{2\pi i \tau_1}$ ,  $t = e^{2\pi i \tau_2}$ , and let  $\mathcal{F}_1 = \text{Fock}_1$  be the charge-1 Fock module. The  $E_3$  categorical  $S$ -matrix on the pair  $(\mathcal{F}_1, \mathcal{F}_1)$ , evaluated as a scalar via the rank-1 decategorification  $\text{Tr}_{\mathcal{F}_1} \rightarrow \text{tr}_{\mathcal{F}_1}$ , is the *double  $S$ -matrix*

$$S(\tau_1, \tau_2) = \text{tr}_{\mathcal{F}_1}(q^{L_0} t^{L'_0} \mathcal{R}^{E_3}(u, v) \circ \mathcal{R}^{E_3}(v, u)) = \prod_{n \geq 1} \frac{1}{(1 - q^n)^2 (1 - t^n)^2 (1 - q^n t^n)^2},$$

where  $L_0, L'_0$  are the two Virasoro zero-modes of the quantum toroidal algebra corresponding to the two loop directions. The product is the square of the MacMahon-type generating function, reflecting the two spectral parameters of the  $E_3$   $R$ -matrix  $R_{\text{ch}}(u, v)$  of Conjecture 9.30(ii).

CONJECTURE 12.28 ( *$\text{Sp}_4(\mathbb{Z})$  modularity of the  $E_3$   $S$ -matrix*). <sup>[CJ]</sup> The double  $S$ -matrix  $S(\tau_1, \tau_2)$  extends to a function of the Siegel upper half-plane  $\mathcal{H}_2 = \{Z = \begin{pmatrix} \tau_1 & z \\ z & \tau_2 \end{pmatrix} : \Im(Z) > 0\}$  via the off-diagonal parameter  $z = u - v$  (the relative spectral parameter). The extended function  $\hat{S}(\tau_1, z, \tau_2)$  transforms as a Siegel modular form of weight  $-2\kappa_{\text{cat}}$  for a subgroup of  $\text{Sp}_4(\mathbb{Z})$  (the  $E_3$  analogue of the  $E_2$  relation  $\text{wt}(S_{ij}) = -\kappa_{\text{cat}}$ ). The modular group promotion is:

| $E_n$ level | Modular group                                             | $S$ -matrix type                       |
|-------------|-----------------------------------------------------------|----------------------------------------|
| $E_2$       | $\text{SL}_2(\mathbb{Z})$ acting on $\tau$                | scalar $S_{ij}$ , elliptic             |
| $E_3$       | $\text{Sp}_4(\mathbb{Z})$ acting on $(\tau_1, z, \tau_2)$ | categorical $,_{\mathcal{W}}$ , Siegel |

The mechanism:  $E_2$  factorization homology on  $T^2$  (the torus, moduli  $\tau$ ) gives the genus-1 amplitude with  $\text{SL}_2(\mathbb{Z})$  symmetry.  $E_3$  factorization homology on the 3-manifold  $\Sigma_2 \times S^1$  (where  $\Sigma_2$  is a genus-2 surface) produces the genus-2 Siegel period matrix  $Z \in \mathcal{H}_2$ , inheriting  $\text{Sp}_4(\mathbb{Z})$  symmetry from the mapping class group  $\text{MCG}(\Sigma_2) \twoheadrightarrow \text{Sp}_4(\mathbb{Z})$ . (Note:  $\Sigma_2 \times S^1$  is *not* the 3-torus  $T^3 = T^2 \times S^1$ ; the latter has  $\text{MCG}(T^2) = \text{SL}_2(\mathbb{Z})$ , not  $\text{Sp}_4(\mathbb{Z})$ .)

CONJECTURE 12.29 (*Connection to  $\Phi_{10}$ : the BKM  $S$ -matrix*). <sup>[CJ]</sup> For  $X = K3 \times E$ , the  $E_3$  categorical  $S$ -matrix of the conjectural chiral algebra  $A_X$  (Conjecture 9.19, conditional on CY-A<sub>3</sub>) is controlled by the Igusa cusp form  $\Phi_{10}$ . Specifically:

- (i) The Borcherds denominator identity for the BKM superalgebra  $\mathfrak{g}_{\Delta_5}$  is

$$\Phi_{10}(\tau_1, z, \tau_2) = e^{2\pi i(\tau_1 + z + \tau_2)} \prod_{(n, \ell, m) > 0} (1 - e^{2\pi i(n\tau_1 + \ell z + m\tau_2)})^{c(4nm - \ell^2)},$$

where  $c(D)$  are the Fourier coefficients of the weak Jacobi form  $\phi_{0,1}(\tau, z)$  of weight 0 and index 1 (the  $K_3$  elliptic genus). This is the Borcherds multiplicative lift of  $\phi_{0,1}$ ; the relation to the additive (Gritsenko) lift is  $\phi_{0,1} = \phi_{10,1}/\eta^{24}$ , where  $\phi_{10,1} = \eta^{18} \vartheta_1^2$  is the weight-10 Jacobi cusp form (the first Fourier–Jacobi coefficient of  $\Phi_{10}$ ). This product is a Siegel modular form of weight  $\kappa_{\text{BKM}} = 5$  for  $\text{Sp}_4(\mathbb{Z})$ .

- (ii) The  $E_3$  categorical  $S$ -matrix, decategorified to a scalar function on  $\mathcal{H}_2$ , satisfies

$$\hat{S}_{K3 \times E}(\tau_1, z, \tau_2) = \Phi_{10}(\tau_1, z, \tau_2)^{-1} \cdot (\text{Eisenstein factor}),$$

where  $\Phi_{10}^{-1}$  is the Borcherds product *inverted*: its poles on the Humbert surfaces  $\{4nm - \ell^2 = 0\}$  are the *resonances* of the categorical scattering matrix, corresponding to BPS bound states becoming massless. The Eisenstein factor is a (normalizing) Siegel Eisenstein series of weight  $2\kappa_{\text{BKM}} - 2\kappa_{\text{cat}} = 10 - 4 = 6$  which ensures the correct transformation law  $\text{wt}(\hat{S}) = -10 + 6 = -2\kappa_{\text{cat}}$ .

- (iii) The Fourier–Jacobi expansion  $\hat{S}^{-1} = \Phi_{10} \cdot E^{-1} = \sum_{m \geq 1} \phi_m(\tau_1, z) e^{2\pi i m \tau_2}$  encodes the shadow tower: the Jacobi form  $\phi_m$  of index  $m$  is the degree- $m$  braided shadow obstruction  $\Theta_{\mathcal{A}, m}^{E_2}$  of Proposition 12.21, now promoted from an elliptic modular object to a Siegel modular one by the  $E_3$  structure.
- (iv) The weight:  $\Phi_{10}$  has weight  $10 = 2\kappa_{\text{BKM}}$ , and  $\kappa_{\text{BKM}} = 5$ . The  $E_3$   $S$ -matrix has modular weight  $-10$  (as  $\Phi_{10}^{-1}$ ), shifted by the Eisenstein normalizer of weight  $\text{wt}(E) = 2\kappa_{\text{BKM}} - 2\kappa_{\text{cat}} = 10 - 4 = 6$ . The net weight of  $\hat{S}$  is  $-10 + 6 = -4 = -2\kappa_{\text{cat}}$ . The formula  $\text{wt}(\hat{S}) = -2\kappa_{\text{cat}}$  is the  $E_3$  analogue of the  $E_2$  relation  $\text{wt}(S_{ij}) = -\kappa_{\text{cat}}$  (where the modular  $S$ -matrix of an MTC has weight  $-c/2$ ,  $c = 2\kappa_{\text{cat}}$  in our convention): the  $E_3$  promotion doubles the weight, as it doubles the genus ( $g = 1 \rightarrow g = 2$ ).

This conjecture depends on CY- $A_3$  (for the existence of  $A_X$ ), on Conjecture 9.19 (for the 6d framework), and on the identification of the BKM root datum with the  $K3 \times E$  BPS spectrum. Each dependency is independent; CY- $A_3$  is the bottleneck.

*Remark 12.30* (The  $E_2 \rightarrow E_3$  modular promotion mechanism). The passage from  $\text{SL}_2(\mathbb{Z})$  to  $\text{Sp}_4(\mathbb{Z})$  is not an *ad hoc* enlargement: it is forced by the factorization homology of the  $E_3$ -algebra over the mapping class group of 3-manifolds. The concrete mechanism:

- At  $E_2$ : factorization homology  $\int_{T^2} \mathcal{A}$  computes the genus-1 amplitude. The moduli space of  $T^2$  is  $\text{SL}_2(\mathbb{Z}) \backslash \mathcal{H}_1$ . The  $S$ -transformation  $\tau \mapsto -1/\tau$  and  $T$ -transformation  $\tau \mapsto \tau + 1$  generate  $\text{SL}_2(\mathbb{Z})$ . The  $S$ -matrix is the  $S$ -transformation of conformal blocks.
- At  $E_3$ : factorization homology  $\int_{\Sigma_2 \times S^1} \mathcal{A}$  computes the genus-2 Siegel amplitude. The moduli space of genus-2 Riemann surfaces is  $\text{Sp}_4(\mathbb{Z}) \backslash \mathcal{H}_2$ . The  $E_3$   $S$ -matrix is the  $\text{Sp}_4(\mathbb{Z})$  transformation matrix of genus-2 conformal blocks. The two spectral parameters  $(\tau_1, \tau_2)$  are the diagonal entries of the period matrix; the off-diagonal entry  $z$  is the relative spectral parameter from the  $E_3$   $R$ -matrix  $R_{\text{ch}}(u, v)$ .
- The Fourier–Jacobi expansion  $\Phi_{10} = \sum_m \phi_m p^m$  is the  $E_2 \hookrightarrow E_3$  restriction: each  $\phi_m(\tau_1, z)$  is an  $\text{SL}_2(\mathbb{Z})$  Jacobi form (the  $E_2$  datum), and the assembly into a Siegel form is the  $E_3$  datum. The index  $m$  is the Fourier–Jacobi charge, which in the categorical picture is the degree of the braided shadow tower  $\Theta_{\mathcal{A}, m}^{E_2}$ .

The factorization homology perspective explains *why* the genus-2 modular group appears:  $E_3$  factorization on a 3-manifold  $M^3$  produces invariants that depend on the moduli of  $M^3$ ; for  $M^3 = \Sigma_g \times S^1$ , these moduli include the Siegel period matrix of  $\Sigma_g$ . At  $g = 2$ , the period matrix lives in  $\mathcal{H}_2$  with  $\mathrm{Sp}_4(\mathbb{Z})$  symmetry.

*Remark 12.31* (Scope and status). Construction 12.26 is a mathematical definition (of the categorical  $S$ -matrix as a categorical trace of the  $E_3$   $R$ -matrix). Conjectures 12.27, 12.28, and 12.29 are conjectural: the first requires the existence of the  $E_3$   $R$ -matrix on Fock modules (Conjecture 9.30); the second requires the  $\mathrm{Sp}_4(\mathbb{Z})$  modularity of the resulting function; the third requires the full  $K3 \times E$  programme (CY- $A_3$  + BKM identification). All three conjectures are conditional on CY- $A_3$  through the dependency chain  $A_X \rightarrow R^{E_3} \rightarrow \hat{S}$ . None of the conjectures above is a theorem. They are formulated here because the structural parallels ( $E_2/\mathrm{SL}_2(\mathbb{Z})/S_{ij}$  vs.  $E_3/\mathrm{Sp}_4(\mathbb{Z})/\hat{S}$ ) are sufficiently rigid to be falsifiable: compute the  $E_3$  factorization homology of the quantum toroidal Fock module over  $\Sigma_2 \times S^1$  and check whether  $\mathrm{Sp}_4(\mathbb{Z})$  modularity holds.

## Chapter 13

# The Drinfeld Center and Bulk Algebras

The passage from boundary to bulk requires the Drinfeld center. An  $E_1$ -chiral algebra  $A$  has a monoidal representation category  $\text{Rep}^{E_1}(A)$ ; this category is not braided, because the  $E_1$  operad sees only one real direction and the fundamental group  $\pi_1(\text{Conf}_2(\mathbb{R})) = \mathbb{Z}$  produces only a monoidal, not braided, structure. The Drinfeld center  $\mathcal{Z}(\text{Rep}^{E_1}(A))$  universally adjoins a braiding; the resulting braided monoidal category is equivalent to  $\text{Rep}^{E_2}(Z_{\text{ch}}^{\text{der}}(A))$ , where  $Z_{\text{ch}}^{\text{der}}(A) = C_{\text{ch}}^\bullet(A, A)$  is the chiral derived center of Volume I. This identification is the categorical bulk-boundary correspondence.

### 13.1 The Drinfeld center of a monoidal category

*Definition 13.1 (Drinfeld center).* Let  $(C, \otimes, \mathbf{1})$  be a monoidal category. The *Drinfeld center*  $\mathcal{Z}(C)$  is the category whose objects are pairs  $(X, \beta_{X,-})$  where:

- (i)  $X \in C$  is an object;
- (ii)  $\beta_{X,-} : X \otimes (-) \xrightarrow{\sim} (-) \otimes X$  is a natural isomorphism (the *half-braiding*);
- (iii) the half-braiding satisfies the hexagon axiom: for all  $Y, Z \in C$ ,

$$\beta_{X, Y \otimes Z} = (\text{id}_Y \otimes \beta_{X, Z}) \circ (\beta_{X, Y} \otimes \text{id}_Z).$$

A morphism  $(X, \beta_X) \rightarrow (Y, \beta_Y)$  is a morphism  $f : X \rightarrow Y$  in  $C$  that intertwines the half-braidings. The Drinfeld center is always braided monoidal, with braiding  $\sigma_{(X, \beta_X), (Y, \beta_Y)} = \beta_{X, Y}$ .

**PROPOSITION 13.2 (Universal property).** <sup>[PE]</sup> The Drinfeld center  $\mathcal{Z}(C)$  is the universal braided monoidal category receiving a central (i.e., braiding-compatible) monoidal functor from  $C$ . The forgetful functor  $U : \mathcal{Z}(C) \rightarrow C$  is faithful and monoidal.

*Example 13.3 (Finite group).* For  $C = \text{Rep}(G)$  with  $G$  a finite group, the Drinfeld center  $\mathcal{Z}(\text{Rep}(G)) \simeq \text{Rep}(D(G))$  is the representation category of the Drinfeld double  $D(G) = \mathbb{C}[G] \rtimes \mathbb{C}(G)$ . When  $G$  is abelian,  $D(G) \simeq \mathbb{C}[G \times \hat{G}]$  and  $\mathcal{Z}(\text{Rep}(G)) \simeq \text{Rep}(G \times \hat{G})$ , the modular tensor category governing the  $G$ -Dijkgraaf–Witten theory.

*Example 13.4 (Quantum group).* For  $C = \text{Rep}_q(\mathfrak{g})$  at generic  $q$ , the Drinfeld center  $\mathcal{Z}(\text{Rep}_q(\mathfrak{g})) \simeq \text{Rep}_q(\mathfrak{g}) \boxtimes \text{Rep}_{q^{-1}}(\mathfrak{g})$ , recovering the Drinfeld double  $D(U_q(\mathfrak{g})) \simeq U_q(\mathfrak{g}) \otimes U_q(\mathfrak{g}^{\text{op}})$ . At roots of unity,  $\mathcal{Z}(C_q) = C_q \boxtimes \overline{C_q}$  is the Drinfeld center of the modular tensor category, and the  $S$ -matrix of  $\mathcal{Z}(C_q)$  governs the Reshetikhin–Turaev invariant of the torus  $T^2$ .

### 13.2 Drinfeld center as chiral derived center

The categorical Drinfeld center and the algebraic chiral derived center are identified by the Ben-Zvi–Francis–Nadler theorem.

**THEOREM 13.5** (*Ben-Zvi–Francis–Nadler; Lurie*). <sup>[PE]</sup> For an  $E_1$ -algebra  $A$  in a symmetric monoidal stable  $\infty$ -category, there is an equivalence of braided monoidal  $\infty$ -categories

$$\mathcal{Z}(\mathrm{Rep}^{E_1}(A)) \simeq \mathrm{Rep}^{E_2}(Z_{\mathrm{ch}}^{\mathrm{der}}(A))$$

where  $Z_{\mathrm{ch}}^{\mathrm{der}}(A) = \mathrm{RHom}_{A\text{-bimod}}(A, A)$  is the derived center (Hochschild cochains) of  $A$ .

**COROLLARY 13.6** (*Chiral derived center = Drinfeld center*). <sup>[PH]</sup> For an  $E_1$ -chiral algebra  $A$ , the chiral derived center  $Z_{\mathrm{ch}}^{\mathrm{der}}(A) = C_{\mathrm{ch}}^{\bullet}(A, A)$  of Volume I (Theorem H) satisfies

$$\mathcal{Z}(\mathrm{Rep}^{E_1}(A)) \simeq \mathrm{Rep}^{E_2}(Z_{\mathrm{ch}}^{\mathrm{der}}(A)).$$

The Drinfeld center is the categorical incarnation of the universal bulk algebra.

*Proof.* The chiral derived center is  $Z_{\mathrm{ch}}^{\mathrm{der}}(A) = \mathrm{RHom}_{\Omega^{\mathrm{ch}}(B(A)) \otimes \Omega^{\mathrm{ch}}(B(A))^{\mathrm{op}}}(A, A)$ . By bar-cobar inversion (Volume I, Theorem B),  $\Omega^{\mathrm{ch}}(B(A)) \simeq A$  on the Koszul locus, so  $Z_{\mathrm{ch}}^{\mathrm{der}}(A) \simeq \mathrm{RHom}_{A \otimes A^{\mathrm{op}}}(A, A) = \mathrm{HH}^{\bullet}(A)$ . The Ben-Zvi–Francis–Nadler theorem (Theorem 13.5) then gives the stated equivalence.  $\square$

**Warning 13.7** (Three functors on  $B(A)$ ). The Drinfeld center / chiral derived center is a FOURTH operation on the bar complex, distinct from the three of Volume I:

- (a) *Bar-cobar inversion*:  $\Omega(B(A)) \simeq A$  (recovers the original algebra);
- (b) *Verdier duality*:  $D_{\mathrm{Ran}}(B(A)) \simeq B(A^{\dagger})$  (produces the bar of the Koszul dual);
- (c) *Koszul dual extraction*:  $H^*(B(A))^{\vee} = A^{\dagger}$  (the strict Koszul dual algebra);
- (d) *Derived center*:  $Z_{\mathrm{ch}}^{\mathrm{der}}(A) = C_{\mathrm{ch}}^{\bullet}(A, A) = \mathrm{RHom}(\Omega(B(A)), A)$  (the universal bulk algebra).

The Drinfeld center  $\mathcal{Z}(\mathrm{Rep}^{E_1}(A))$  is  $\mathrm{Rep}^{E_2}(Z_{\mathrm{ch}}^{\mathrm{der}}(A))$ : it is functor (d), not (a), (b), or (c). Bar-cobar inversion recovers the boundary; the Drinfeld center produces the bulk.

**Remark 13.8** (Three centers: categorical, algebraic, and symmetric). <sup>[PE]</sup> Three “center” constructions arise in the passage from  $E_n$ -algebra to  $E_{n+1}$ -algebra. They are distinct objects that agree only in the lowest case.

|                  | Drinfeld center $\mathcal{Z}(C)$ | Derived center $\mathrm{HH}^{\bullet}(B, B)$ | Müger center $\mathcal{Z}_2(C)$      |
|------------------|----------------------------------|----------------------------------------------|--------------------------------------|
| <i>Input</i>     | $E_1$ -monoidal $C$              | $E_n$ -algebra $B$                           | braided monoidal $C$                 |
| <i>Output</i>    | $E_2$ -braided category          | $E_{n+1}$ -algebra                           | full subcategory of $C$              |
| <i>Mechanism</i> | half-braidings                   | Hochschild cochains                          | objects with trivial double braiding |
| <i>Promotion</i> | $E_1 \rightarrow E_2$            | $E_n \rightarrow E_{n+1}$                    | <b>none</b> : detects obstructions   |

**Agreement at  $E_1 \rightarrow E_2$ .** For an  $E_1$ -algebra  $A$  in a stable  $\infty$ -category, the BZFN theorem (Theorem 13.5) identifies  $\mathcal{Z}(\mathrm{Rep}^{E_1}(A)) \simeq \mathrm{Rep}^{E_2}(\mathrm{HH}^{\bullet}(A, A))$ : the categorical and algebraic centers produce the same  $E_2$ -structure. This is the identification used throughout this chapter.

**Divergence at  $E_2 \rightarrow E_3$ .** For an  $E_2$ -algebra  $B$ , the Hochschild cochains  $\mathrm{HH}^{\bullet}(B, B)$  carry an  $E_3$ -structure by Dunn additivity (subject to the input conditions of AP153:  $E_3$  on cochains requires  $B$  at least  $E_{\infty}$ ; for  $E_1$ -input the cochains carry only  $E_2$ ). The iterated Drinfeld center  $\mathcal{Z}(\mathcal{Z}(C))$  does *not* produce  $E_3$ : for a braided category  $C$ , one has  $\mathcal{Z}(C) \simeq C \boxtimes C^{\mathrm{rev}}$  (the Deligne product with the reverse braiding), which is an  $E_2$ -doubling, not an  $E_3$ -promotion.

**The Müger center is an obstruction detector.** The Müger center  $\mathcal{Z}_2(C) = \{X \in C : \sigma_{Y,X} \circ \sigma_{X,Y} = \text{id}_{X \otimes Y} \forall Y\}$  is the subcategory of “transparent” objects. It detects degeneracy of the braiding, not a promotion:  $\mathcal{Z}_2(C) \simeq \text{Vect}$  if and only if  $C$  is non-degenerate (and hence a candidate modular tensor category). When  $\mathcal{Z}_2(C)$  is nontrivial, the braiding has a symmetric kernel that obstructs modularity.

*References.* Drinfeld center universality: Lurie, *Higher Algebra*, §5.3.1. BZFN identification: Ben-Zvi–Francis–Nadler, *Integral transforms and Drinfeld centers* (2010). Müger center: Müger, *Galois theory for braided tensor categories* (2003).

*Remark 13.9* (Three centers: agreement, divergence, and obstruction). Three operations called “center” play fundamentally different roles.

**(a) Drinfeld center  $\mathcal{Z}(C)$ : categorical,  $E_1 \rightarrow E_2$ .** Given an  $E_1$ -monoidal category  $C$ , the Drinfeld center  $\mathcal{Z}(C)$  universally adjoins half-braidings to produce a braided monoidal category. It is intrinsically a *one-step* promotion:  $E_1$ -monoidal input,  $E_2$ -braided output. The mechanism is purely categorical (no cochains, no resolution).

**(b) Derived center  $\text{HH}^*(B, B)$ : algebraic,  $E_n \rightarrow E_{n+1}$ .** For an  $E_n$ -algebra  $B$ , the Hochschild cochain complex  $\text{HH}^*(B, B) = \text{RHom}_{B \otimes B^{\text{op}}}(B, B)$  carries an  $E_{n+1}$ -structure by Dunn additivity (subject to AP153;  $E_3$  on cochains requires  $B$  at least  $E_\infty$ ; for  $E_1$ -input the cochains are  $E_2$ ). This promotion iterates: applied to an  $E_2$ -algebra  $B$  with sufficient commutativity,  $\text{HH}^*(B, B)$  is genuinely  $E_3$ .

**Agreement at  $E_1 \rightarrow E_2$  (BZFN).** At the lowest step, (a) and (b) coincide: the BZFN theorem (Theorem 13.5) gives  $\mathcal{Z}(\text{Rep}^{E_1}(A)) \simeq \text{Rep}^{E_2}(\text{HH}^*(A, A))$ . This is the identification used throughout this chapter and the CY programme.

**Divergence at  $E_2 \rightarrow E_3$ .** The two operations diverge at the next step. For a braided category  $\mathcal{B}$ , the iterated Drinfeld center satisfies  $\mathcal{Z}(\mathcal{B}) \simeq \mathcal{B} \boxtimes \mathcal{B}^{\text{rev}}$ , the Deligne product with the reverse braiding. This is an  $E_2$ -doubling, not an  $E_3$ -promotion:  $\mathcal{Z}(\mathcal{Z}(C))$  for braided  $C$  gives  $C \boxtimes C^{\text{rev}}$ , which has no  $E_3$ -structure. The algebraic derived center  $\text{HH}^*(B, B)$  does promote to  $E_3$  (when the input satisfies AP153). Thus Drinfeld centers do not iterate to higher  $E_n$ -structures; derived centers do.

**(c) Müger center  $\mathcal{Z}_2(C)$ : obstruction, NOT promotion.** For a braided  $C$ , the Müger center  $\mathcal{Z}_2(C) = \{X : \sigma_{Y,X} \circ \sigma_{X,Y} = \text{id} \forall Y\}$  is the subcategory of transparent objects. It detects degeneracy of the braiding:  $\mathcal{Z}_2(C) \simeq \text{Vect}$  iff  $C$  is non-degenerate (a modularity prerequisite). The Müger center produces no new algebraic structure; it measures the *failure* of the existing braiding to be non-degenerate.

### The $\mathbb{C}^3$ theorem: Yangian and $\mathcal{W}_{1+\infty}$

A concrete instantiation of Corollary 13.6 is for the CY<sub>3</sub> chiral algebra of  $\mathbb{C}^3$ .

**THEOREM 13.10** (*Drinfeld center identification for  $\mathbb{C}^3$ ; Schiffmann–Vasserot, Maulik–Okounkov*). <sup>[PE]</sup>

$$\mathcal{Z}(\text{Rep}^{E_1}(Y^+(\widehat{\mathfrak{gl}}_1))) \simeq \text{Rep}^{E_2}(Y(\widehat{\mathfrak{gl}}_1)) \simeq \text{Rep}^{E_2}(\mathcal{W}_{1+\infty}).$$

The positive half-Yangian  $Y^+(\widehat{\mathfrak{gl}}_1)$  is an  $E_1$ -chiral algebra (the CoHA of  $\mathbb{C}^3$  after Schiffmann–Vasserot); its representation category is monoidal but not braided. The full Yangian  $Y(\widehat{\mathfrak{gl}}_1)$  is the chiral derived center  $\mathcal{Z}_{\text{ch}}^{\text{der}}(Y^+(\widehat{\mathfrak{gl}}_1))$ , and the  $E_2$ -braided structure on its representation category arises from the Drinfeld center construction. The equivalence  $Y(\widehat{\mathfrak{gl}}_1) \simeq \mathcal{W}_{1+\infty}$  (at  $c = 1$ , the CY<sub>3</sub> central charge) identifies the bulk algebra with the  $\mathcal{W}_{1+\infty}$ -algebra.

*Remark 13.11* (Attribution). The  $E_1$ -nature of  $Y^+(\widehat{\mathfrak{gl}}_1)$  is due to Schiffmann–Vasserot (2013) and the identification of the full Yangian as the Drinfeld double is due to Maulik–Okounkov (2019). The vertex algebra isomorphism  $Y(\widehat{\mathfrak{gl}}_1) \simeq \mathcal{W}_{1+\infty}$  at  $c = 1$  follows from the work of Procházka–Rapčák (2019) and Litvinov (2020). The formulation as a Drinfeld center identification, connecting the categorical and algebraic perspectives, is the content of Theorem 8.28 in Chapter 8.

### The Ben-Zvi–Francis–Nadler loop space identification

The BZFN theorem admits a geometric reformulation in terms of loop spaces that makes the passage from  $E_1$  to  $E_2$  explicit.

**THEOREM 13.12** (*BZFN loop space identification; Ben-Zvi–Francis–Nadler*). <sup>[PE]</sup> For a smooth algebraic stack  $X$ , there is an equivalence of braided monoidal  $\infty$ -categories

$$\mathcal{Z}(\mathrm{QCoh}(X)) \simeq \mathrm{QCoh}(X)$$

where  $X = \mathrm{Map}(S^1, X)$  is the (derived) free loop space of  $X$ . The braided monoidal structure on  $\mathrm{QCoh}(X)$  arises from the  $S^1$ -rotation action on the loop space.

*Remark 13.13* (Connection to the derived center). When  $X = BG$  for a reductive group  $G$ , the loop space  $(BG) = G/G$  (the adjoint quotient), and  $\mathrm{QCoh}((BG)) \simeq \mathrm{QCoh}(G/G)$  recovers the character sheaves of Lusztig. For  $X = \mathrm{Spec}(A)$ , the loop space  $X = \mathrm{Spec}(\mathrm{HH}_\bullet(A))$  (derived Hochschild homology), and the BZFN equivalence reduces to Theorem 13.5: the Drinfeld center of  $A$ -modules is computed by the Hochschild cochains of  $A$  (which are Koszul dual to the Hochschild chains defining  $X$ ).

### 13.3 Bulk-boundary correspondence

**PROPOSITION 13.14** (*Information flow by genus*). <sup>[PH]</sup> The bulk-boundary correspondence  $\mathcal{Z}(\mathrm{Rep}^{E_1}(A)) \simeq \mathrm{Rep}^{E_2}(Z_{\mathrm{ch}}^{\mathrm{der}}(A))$  exhibits genus-dependent information flow:

- (i) *Genus 0*: Information flows one way, from bulk to boundary. The forgetful functor  $\mathcal{Z}(C) \rightarrow C$  has no natural section: the Swiss-cheese operad forbids open-to-closed maps at genus 0 (Volume II, thm:thqg-swiss-cheese);
- (ii) *Genus 1*: The annulus trace  $\Delta_{\mathrm{ns}}(\mathrm{Tr}_A) = \kappa_{\mathrm{cat}} \cdot \lambda_1$  (Volume I, thm:annulus-trace) provides the first open-to-closed map. The Dennis trace  $K(C) \rightarrow \mathrm{HH}_\bullet(C)$  maps K-theory of the open sector into the bulk via the SBI sequence;
- (iii) *Higher genus*: The full genus tower progressively restores bidirectionality. The clutching maps  $\mathrm{cl}_{g,n}: \mathrm{HH}_\bullet(A)^{\otimes n} \rightarrow Z_{\mathrm{ch}}^{\mathrm{der}}(A)$  on the modular operad provide genus- $g$  open-to-closed data.

*Proof.* Item (i): at the operad level,  $\mathrm{SC}^{\mathrm{ch}, \mathrm{top}}$  has no composition open  $\rightarrow$  closed at genus 0 (the Kontsevich no-open-to-closed axiom).

Item (ii): the Dennis trace  $K(C) \rightarrow \mathrm{HH}_\bullet(C)$  factors through negative cyclic homology  $\mathrm{HC}_\bullet^-(C)$ . The CY duality  $\mathrm{HH}^\bullet(C) \simeq \mathrm{HH}_{\bullet+d}(C)$  (Theorem 4.13) and the identification  $\mathrm{HH}^\bullet(C) \simeq Z_{\mathrm{ch}}^{\mathrm{der}}(A)$  (Corollary 13.6) give the map into the bulk. The annulus trace is the genus-1 specialization: the  $S^1$ -trace on  $\mathrm{HH}_0(A)$  gives  $\kappa_{\mathrm{cat}} \cdot \lambda_1 \in H^2(\overline{\mathcal{M}}_{1,1})$ .

Item (iii): the modular operad structure on  $\{H^*(\overline{\mathcal{M}}_{g,n})\}$  provides clutching maps  $\gamma_{g_1, n_1} \circ_{g_2, n_2}$  that compose open-sector data (living on boundaries of  $\overline{\mathcal{M}}_{g,n}$ ) into closed-sector invariants. Each genus  $g$  introduces new clutching compositions (from codimension-1 boundary strata of  $\overline{\mathcal{M}}_{g,n}$ ), progressively enlarging the image of the open-to-closed map.  $\square$



### 13.4 CY enhancement of the Drinfeld center

**THEOREM 13.15** (*CY Drinfeld center*). <sup>[PE]</sup> Let  $\mathcal{C}$  be a smooth proper  $\text{CY}_d$  category with quantum chiral algebra  $A_{\mathcal{C}} = \Phi(\mathcal{C})$ . The Drinfeld center  $\mathcal{Z}(\text{Rep}^{E_1}(A_{\mathcal{C}}))$  carries a CY structure of dimension  $d + 1$ . Under the identification  $\mathcal{Z}(\text{Rep}^{E_1}(A_{\mathcal{C}})) \simeq \text{Rep}^{E_2}(Z_{\text{ch}}^{\text{der}}(A_{\mathcal{C}}))$ , this  $\text{CY}_{d+1}$  structure arises from the  $(1 - d)$ -shifted symplectic structure on the derived center (Pantev–Toën–Vaquié–Vezzosi).

*Remark 13.16* (Dimensional hierarchy). The shift  $d \rightarrow d + 1$  is the categorical holographic principle:

| CY dim  | Boundary      | Bulk          | Physical                     |
|---------|---------------|---------------|------------------------------|
| $d = 1$ | $E_1$ -chiral | $\text{CY}_2$ | 2d CFT / 3d TFT              |
| $d = 2$ | $E_2$ -chiral | $\text{CY}_3$ | 3d HT / 4d $\mathcal{N} = 2$ |
| $d = 3$ | $E_1$ -chiral | $\text{CY}_4$ | 4d $\mathcal{N} = 1$ gauge   |

For  $d = 3$ , the boundary algebra is  $E_1$  (not  $E_2$ ): the  $\mathbb{S}^3$ -framing obstruction (Conjecture  $\text{CY-A}_3$ ) breaks  $E_2$  to  $E_1$ . The braiding is recovered via  $\mathcal{Z}(\text{Rep}^{E_1}(A))$ , not from the boundary  $A$  directly.

### 13.5 Categorical Theorem H

Volume I Theorem H establishes that chiral Hochschild cohomology  $\text{ChirHoch}^*(A)$  is polynomial (concentrated in degrees  $\{0, 1, 2\}$ ) for modular Koszul algebras, with total dimension  $\leq 4$ . The categorical analogue replaces  $\text{ChirHoch}^*$  with  $\text{HH}^\bullet(\mathcal{C})$ .

**THEOREM 13.17** (*Categorical Theorem H*). <sup>[CD]</sup> For a smooth proper  $\text{CY}_d$  category  $\mathcal{C}$  whose quantum chiral algebra  $A_{\mathcal{C}} = \Phi(\mathcal{C})$  is modular Koszul:

- (i)  $\text{HH}^k(\mathcal{C}) = 0$  for  $k > d$  (concentration, from smoothness and properness);
- (ii)  $\text{HH}^0(\mathcal{C}) \simeq k$  (the identity natural transformation);
- (iii)  $\text{HH}^1(\mathcal{C})$  classifies first-order deformations of  $\mathcal{C}$  as a dg category;
- (iv)  $\text{HH}^2(\mathcal{C})$  is the obstruction space for deformations.

The Gerstenhaber bracket  $[-, -]: \text{HH}^p \otimes \text{HH}^q \rightarrow \text{HH}^{p+q-1}$  on  $\text{HH}^\bullet(\mathcal{C})$  maps, under  $\Phi$ , to the convolution bracket on  $\mathfrak{g}_{A_{\mathcal{C}}}^{\text{mod}}$  of Volume I.

*Remark 13.18* (Status). Items (i)–(iv) are standard consequences of smoothness and properness (Keller, Kontsevich). The conditional content is the identification of the Gerstenhaber bracket with the Volume I convolution bracket under  $\Phi$ , established for  $d = 2$  and conjectural for  $d = 3$ .

### 13.6 Quantum group reconstruction from the center

**PROPOSITION 13.19** (*Tannakian reconstruction*). <sup>[PE]</sup> Let  $\mathcal{B}$  be a rigid braided monoidal category over  $\mathbb{C}$  equipped with a fiber functor  $\omega: \mathcal{B} \rightarrow \text{Vect}_{\mathbb{C}}$ . The Tannakian reconstruction gives a quasi-triangular Hopf algebra  $H = \text{End}(\omega)$  with  $\mathcal{B} \simeq \text{Rep}(H)$  as braided monoidal categories. When  $\mathcal{B} = \mathcal{Z}(\text{Rep}^{E_1}(A))$  for an  $E_1$ -chiral algebra  $A$  with  $\mathfrak{g}$ -symmetry,  $H \simeq U_q(\mathfrak{g})$  with  $q$  determined by  $\kappa_{\text{cat}}$ .

*Remark 13.20* (Non-semisimple reconstruction). At roots of unity ( $q^N = 1$ ), the category  $\text{Rep}_q(\mathfrak{g})$  is non-semisimple: indecomposable but non-simple modules appear. The standard fiber functor is not exact on the full category. Two remedies:

- (a) Restrict to the semisimplified quotient  $\overline{\text{Rep}}_q(\mathfrak{g})$  (modular tensor category of Reshetikhin–Turaev);
- (b) Use the modified trace of Geer–Patureau-Mirand–Turaev, which replaces the categorical trace for non-semisimple ribbon categories.

For the CY programme, approach (b) is preferred: the full non-semisimple structure encodes the complete BPS spectrum (Chapter 15), and the modified trace is the categorical analogue of the regularized partition function.

### 13.7 The Drinfeld double and the slab

*Definition 13.21* (*Drinfeld double*). For a finite-dimensional Hopf algebra  $H$ , the *Drinfeld double*  $D(H) = H \bowtie H^{*\text{cop}}$  is the quasi-triangular Hopf algebra built from  $H$  and the co-opposite of its dual. Its representation category is  $\mathcal{Z}(\text{Rep}(H))$ , the Drinfeld center of  $\text{Rep}(H)$ .

*Remark 13.22* (The slab is a bimodule). In the 3d holomorphic-topological setting (Volume II), the Drinfeld double arises from the *slab*: a strip  $X \times [0, 1]$  with two boundary components  $X \times \{0\}$  and  $X \times \{1\}$ . The algebra of operators on the slab is a bimodule for the two boundary algebras  $A_0$  and  $A_1$ . The Drinfeld double  $D(A) = A \bowtie A^{\dagger, \text{cop}}$  appears when  $A_0 = A$  and  $A_1 = A^\dagger$  (the Koszul dual). The slab is NOT a Swiss-cheese disk (which has one closed and one open boundary component); it is a cylinder with two open boundary components.

**CONJECTURE 13.23** (*Slab = Drinfeld double*). <sup>[CJ]</sup> For an  $E_1$ -chiral algebra  $A$  on  $X \times \mathbb{R}$ , the slab algebra  $\mathcal{A}_{\text{slab}} = \text{End}_{A-A^\dagger}(A \otimes_{A^\dagger} A)$  is isomorphic to the Drinfeld double  $D(A)$ . The slab  $R$ -matrix  $\mathcal{R}_{\text{slab}} = \mathcal{R}_A \otimes \mathcal{R}_{A^\dagger}^{-1}$  is the universal  $R$ -matrix of  $D(A)$ , and its representation category is the Drinfeld center  $\mathcal{Z}(\text{Rep}^{E_1}(A))$ .

**CONJECTURE 13.24** (*Quantum vertex chiral group*). <sup>[CJ]</sup> For a  $\text{CY}_2$  category  $\mathcal{C}$  with  $\mathfrak{g}$ -symmetry, the Drinfeld center  $\mathcal{Z}(\text{Rep}^{E_1}(\Phi(\mathcal{C})))$  determines a quantum vertex chiral group  $G(\mathcal{C})$  unifying four algebraic structures:

- (i)  $U_q(\mathfrak{g})$ : the quantum group (fiber at a point);
- (ii)  $Y(\mathfrak{g})$ : the Yangian ( $E_1$ -sector, Volume II, MC3);
- (iii)  $U_q(\widehat{\mathfrak{g}})$ : the affine quantum group (loop algebra sector);
- (iv) The Etingof–Kazhdan quantum vertex algebra (the genuinely nonlocal atom, Volume II).

The four incarnations arise from different geometric specializations of the  $E_2$ -factorization structure on  $\text{Ran}(X) \times \text{Ran}(Y)$ .

**CONJECTURE 13.25** (*Drinfeld center equals bulk*). <sup>[CJ]</sup> Let  $A$  be a modular Koszul  $E_1$ -chiral algebra and let  $U_A = A \bowtie A^\dagger$  denote the Drinfeld double of the Koszul pair  $(A, A^\dagger)$ . Then there is an equivalence of  $E_2$ -algebras

$$\mathcal{Z}(U_A) \simeq \mathcal{Z}_{\text{ch}}^{\text{der}}(A)$$

identifying the center of the Drinfeld double with the chiral derived center (the universal bulk algebra of Volume I).

*Remark 13.26* (Three obstructions). Three independent obstructions prevent a direct proof of Conjecture 13.25:

- (i) *Pointwise reduction failure for class M*. For algebras of class M (Virasoro,  $\mathcal{W}_N$ ), the shadow tower has infinite depth: all operations  $m_k$  are nonzero. The Drinfeld double  $U_A = A \bowtie A^\dagger$  is defined at the chain level, but the passage from the Drinfeld center  $\mathcal{Z}(\text{Rep}^{E_1}(U_A))$  to the algebraic center  $\mathcal{Z}(U_A)$  requires a pointwise reduction (evaluating the half-braiding on each object) that may not converge when the shadow tower is infinite.
- (ii) *Factorization property of  $A^\dagger$* . The Koszul dual  $A^\dagger$  is an algebra, not a factorization algebra on  $\text{Ran}(X)$  in general. The Drinfeld double construction requires both  $A$  and  $A^\dagger$  to be factorization algebras (so that  $U_A$  factors on  $\text{Ran}(X) \times \text{Ran}(Y)$ ). For class G and L algebras,  $A^\dagger$  inherits a factorization structure from the bar-cobar inversion; for class C and M, this is open.
- (iii) *Bar-cobar/Hochschild compatibility*. The chiral derived center  $Z_{\text{ch}}^{\text{der}}(A) = C_{\text{ch}}^\bullet(A, A)$  is defined via chiral Hochschild cochains. The Drinfeld double center  $\mathcal{Z}(U_A)$  is defined via the half-braiding construction on  $A \bowtie A^\dagger$ . The two constructions agree when the bar-cobar adjunction of Volume I (Theorem A) is compatible with the Hochschild complex in the precise sense that  $\text{RHom}_{U_A}(A, A) \simeq \text{RHom}_{A \otimes A^{\text{op}}}(A, A)$ . This compatibility is proved for class G (where  $A^\dagger$  is Koszul dual in the classical sense) and conjectural in general.

*Remark 13.27* (Verification for the Heisenberg algebra). For  $A = H_k$  (Heisenberg, class G), the conjecture is verified at the level of numerical invariants. The Drinfeld double is  $\text{Drin}(H_k) = H_k \otimes H_{-k}/(c = c')$ , dimension 3, with brackets  $[J, J'] = k \cdot c$ . The naive Lie centre  $Z(\text{Drin}(H_k))$  has dimension 1 at generic  $k$ . The chiral derived centre is  $Z_{\text{ch}}^{\text{der}}(H_k) \cong \mathbb{C} \oplus \mathbb{C}[-1] \oplus \mathbb{C}[-2]$ , dimension 3: the discrepancy (1 vs 3) reflects  $\text{Ext}^1(J, \text{the derivation})$  and  $\text{Ext}^2(J, J)$  (the level deformation) invisible to the commutant. Five consistency checks match across both sides:  $r$ -matrix coefficient  $= k$ ,  $\kappa_{\text{ch}} = k$ ,  $\chi = 1$ , shadow class  $G$  / depth 2, complementarity  $k + (-k) = 0$ . Verified at 6 level values ( $k = 0, 1, -1, 1/2, 7, -3/2$ ) with 72 compute tests. For class G the  $E_2$  structure on both sides is determined by the single parameter  $k$ , so the conjecture verification reduces to matching this one numerical invariant.

## 13.8 The $E_n$ -circle and the categorified averaging map

The Drinfeld center is not an isolated construction: it is the  $E_1 \rightarrow E_2$  arrow in the  $E_n$ -circle, the systematic passage between  $E_n$ -structures of different degrees.

*Definition 13.28* ( $E_n$ -center). For  $0 \leq m < n$ , the  $E_n$ -center of an  $E_m$ -monoidal category  $C$  is the  $E_n$ -monoidal category

$$\mathcal{Z}_{E_n/E_m}(C) := \text{End}_{C\text{-BiMod}_{E_n}\text{-}C}(C)$$

obtained by taking bimodule endomorphisms in the  $E_n$ -sense.

- $m = 0, n = 1$ : the ordinary center  $\mathcal{Z}(C) = Z(C)$  of a category (monoidal center).
- $m = 1, n = 2$ : the Drinfeld center  $\mathcal{Z}(C)$ , recovering  $E_2$ -braiding from  $E_1$ -monoidal structure. This is the case relevant to this chapter.
- $m = n - 1$ , general  $n$ : Dunn additivity gives  $\mathcal{Z}_{E_n/E_{n-1}}(C) \simeq C$  when  $C$  is already  $E_n$ -monoidal and non-degenerate. The center construction “does nothing” when the target degree is already present.

**PROPOSITION 13.29** (*Drinfeld center as categorified averaging*). <sup>[PH]</sup> The Drinfeld center  $\mathcal{Z}: E_1\text{-Cat} \rightarrow E_2\text{-Cat}$  is the categorification of the averaging map  $\text{av}: \mathfrak{g}^{E_1} \rightarrow \mathfrak{g}^{\text{mod}}$  of Volume I. The correspondence is:

| Volume I (algebraic)                                                  | This chapter (categorical)                               | Level     |
|-----------------------------------------------------------------------|----------------------------------------------------------|-----------|
| $\mathfrak{g}_A^{E_1}$ (convolution Lie)                              | $\text{Rep}^{E_1}(A)$ (monoidal)                         | $E_1$     |
| $\text{av}: \mathfrak{g}^{E_1} \rightarrow \mathfrak{g}^{\text{mod}}$ | $\mathcal{Z}: E_1\text{-Cat} \rightarrow E_2\text{-Cat}$ | averaging |
| $\kappa_{\text{ch}} = \text{av}(r(z))$ (scalar shadow)                | braiding on $\mathcal{Z}(C)$ (categorical shadow)        | output    |

The averaging map is lossy:  $\text{av}$  discards the  $\Sigma_n$ -equivariant structure of the ordered bar  $B^{\text{ord}}(A)$ , retaining only the  $\Sigma_n$ -coinvariant projection (the scalar  $\kappa_{\text{ch}}$ ). The Drinfeld center is the categorification of this projection: it extracts the  $E_2$ -braided content from the full  $E_1$ -monoidal structure.

*Proof.* The algebraic averaging map  $\text{av}: \mathfrak{g}_A^{E_1} \rightarrow \mathfrak{g}_A^{\text{mod}}$  is defined as the  $\Sigma_n$ -coinvariant projection on the ordered convolution algebra. At degree 2,  $\text{av}(r(z)) = \kappa_{\text{ch}}$  extracts the scalar shadow from the  $r$ -matrix (Volume I, §5).

The categorical analogue: the  $E_1$ -monoidal category  $C = \text{Rep}^{E_1}(A)$  carries a tensor product that is associative but not braided. The Drinfeld center  $\mathcal{Z}(C)$  extracts the braided monoidal structure by imposing half-braiding compatibility (the categorical analogue of  $\Sigma_n$ -symmetrization). The braiding on  $\mathcal{Z}(C)$ , when restricted to the forgetful image  $U(\mathcal{Z}(C)) \subset C$ , recovers the  $R$ -matrix data that categorifies  $r(z)$ . The scalar  $\kappa_{\text{ch}}$  is the Euler characteristic of the braiding: the categorical trace  $\text{tr}_{\mathcal{Z}}(\sigma) = \kappa_{\text{cat}}$  gives the categorical shadow of the  $R$ -matrix.  $\square$

### 13.9 The $E_3 \rightarrow E_2$ restriction and symmetric braiding

The Drinfeld center produces non-symmetric braiding from  $E_1$ -monoidal input. This is a fundamentally different mechanism from the  $E_3 \rightarrow E_2$  restriction, which produces only symmetric braiding. The distinction is the mechanism behind the  $E_1$  stabilization theorem (Chapter 8, Theorem 7.1).

**PROPOSITION 13.30** ( *$E_3$  restriction gives symmetric braiding*). <sup>[PE]</sup> Let  $C$  be an  $E_3$ -monoidal category. The restriction of the  $E_3$ -structure to  $E_2$  produces a braided monoidal category whose braiding is *symmetric*:  $\sigma_{X,Y} \circ \sigma_{Y,X} = \text{id}_{X \otimes Y}$  for all objects  $X, Y$ .

*Proof.* The braiding on an  $E_2$ -monoidal category is classified by  $\pi_1(\text{Conf}_2(\mathbb{R}^2)) \cong \mathbb{Z}$ : the generator is the counterclockwise winding of one point around another in the plane. When the  $E_2$ -structure is the restriction of an  $E_3$ -structure, the braiding is classified by  $\pi_1(\text{Conf}_2(\mathbb{R}^3))$ , which is trivial (two points in  $\mathbb{R}^3$  can be unlinked by moving in the third direction). The double braiding  $\sigma_{X,Y} \circ \sigma_{Y,X}$  corresponds to the generator of  $\pi_1(\text{Conf}_2(\mathbb{R}^2))$  traversed twice; in  $\mathbb{R}^3$  this loop is contractible, so the double braiding is the identity.  $\square$

**Remark 13.31** (Two sources of  $E_2$ -braiding for  $\text{CY}_3$ ). For a  $\text{CY}_3$  category  $C$ , two  $E_2$ -braided structures are available, and they are fundamentally different:

- (a)  *$E_3$ -restriction.* The  $\mathbb{S}^3$ -framing on  $\text{HH}_\bullet(C)$  gives an  $E_3$ -structure (conditional on chain-level trivialization of the framing obstruction). Restricting  $E_3$  to  $E_2$  via Dunn additivity gives a braided monoidal category with *symmetric* braiding ( $\pi_1(\text{Conf}_2(\mathbb{R}^3)) = 0$ ). This symmetric braiding is the “wrong” answer for quantum groups: it sees only the classical limit  $q = 1$ .
- (b) *Drinfeld center.* The  $E_1$ -chiral algebra  $A = \Phi(C)$  has a monoidal representation category  $\text{Rep}^{E_1}(A)$ , and the Drinfeld center  $\mathcal{Z}(\text{Rep}^{E_1}(A))$  produces an  $E_2$ -braided category with *non-symmetric* braiding ( $\pi_1(\text{Conf}_2(\mathbb{R}^2)) \cong \mathbb{Z}$ ). This non-symmetric braiding is the quantum group  $R$ -matrix at  $q \neq 1$ .

The passage (a)  $\rightarrow$  (b) is the  $E_1$  stabilization: the  $E_3$ -structure on the  $\text{CY}_3$  Hochschild complex is “too symmetric” to see quantum groups, and one must descend to  $E_1$  (by choosing a preferred direction, i.e., the  $\Omega$ -background) before applying the Drinfeld center to recover the non-trivial braiding. This is the categorical content of the  $E_1$  stabilization theorem: the genuine quantum group braiding arises not from restricting a higher  $E_n$ -structure, but from the Drinfeld center of a lower one.

*Remark 13.32* (The  $E_n$ -circle in the CY programme). The pattern generalizes across CY dimensions as follows:

| CY dim $d$ | Native $E_n$ on $\text{HH}_\bullet$ | Braiding type | Quantum group                       |
|------------|-------------------------------------|---------------|-------------------------------------|
| $d = 1$    | $E_1$ (trivially)                   | none          | Drinfeld center gives $E_2$         |
| $d = 2$    | $E_2$                               | non-symmetric | direct (the “sweet spot”)           |
| $d = 3$    | $E_3$                               | symmetric     | must descend to $E_1$ , then center |
| $d = 4$    | $E_4$                               | symmetric     | further descent needed              |

The dimension  $d = 2$  is distinguished: the native  $E_2$ -structure already carries non-symmetric braiding, so no descent or center construction is needed. For  $d \geq 3$ , the  $E_n$ -structure with  $n \geq 3$  gives only symmetric braiding, and the non-symmetric quantum group braiding requires the  $E_1$  stabilization followed by the Drinfeld center. For  $d = 1$ , there is no native braiding at all, and the Drinfeld center is the only source.

This explains why quantum groups are ubiquitous in the  $d = 2$  CY landscape ( $K_3$  surfaces, quiver varieties) and require additional structure ( $\Omega$ -deformation, stability conditions) in the  $d = 3$  CY landscape (Calabi–Yau threefolds).



## **Part III**

# **The CY Landscape**





## Chapter 14

# Toroidal and elliptic algebras

The chiral algebras of Vols I–II each depend on a single deformation parameter: the level  $k$ , the central charge  $c$ , or the spectral parameter  $u$ . This chapter enters the regime where one parameter is not enough. The toroidal algebra  $U_{q,t}(\hat{\mathfrak{g}})$  carries two independent parameters  $(q, t)$ , the elliptic quantum group  $E_{q,p}(\mathfrak{g})$  replaces the rational  $R$ -matrix by a doubly-periodic one, and the double affine Hecke algebra (DAHA) intertwines spectral and modular directions. In these algebras the modular parameter  $\tau$  and the spectral parameter  $u$  are coupled through the Fay trisecant identity and the quasi-periodicity of theta functions.

The bar-cobar framework meets this two-parameter world in two ways. Direction (a) keeps the base a single curve, now of genus 1: the propagator becomes elliptic, the Arnold relation generalizes to the Fay trisecant identity (Theorem 14.20), and the bar differential acquires Eisenstein-series corrections indexed by  $\tau$ . Direction (b) seeks genuine surface-factorization objects on  $\mathbb{C}^* \times \mathbb{C}^*$ , requiring a two-dimensional factorization framework beyond the curve-chiral hierarchy of Vols I–II. Only direction (a) is developed here.

The Fay identity replaces the Arnold relation at genus 1; it is both an identity of theta functions and a geometric consequence of the degeneration  $E_\tau \rightarrow \mathbb{P}^1$  as  $\tau \rightarrow i\infty$ .

*Remark 14.1* (Two programme tracks). Track (a) keeps the base an algebraic curve and constructs an elliptic  $E_1$ -chiral realization on  $E_\tau$ . Track (b) seeks a genuine surface-factorization object on  $\mathbb{C}^* \times \mathbb{C}^*$ . Only track (a) is developed here; track (b) is a completion target.

*Remark 14.2* (The generalization principle: Arnold to Fay). The Arnold relation (Vol I, Theorem III), which governs the genus-0 bar differential, admits two distinct generalizations. Direction (a): via clutching morphisms and the shadow obstruction tower  $\Theta_A^{\leq r}$  (Vol I, Definition III), one passes to the modular regime on stable curves, recovering the higher-genus bar complex developed in Vol I (Vol I, Remark III). Direction (b): via the Fay trisecant identity (Theorem 14.20), one replaces the additive (logarithmic) propagator  $\omega = dz/(z - w)$  by the elliptic (multiplicative) propagator built from the Weierstrass  $\sigma$ -function, entering the toroidal regime. The present chapter develops direction (b): the systematic passage from logarithmic to elliptic propagators and its consequences for bar-cobar duality.

*Remark 14.3* (Rational, trigonometric, elliptic: curve geometry). The three quantum group families arise from the geometry of the underlying curve:

| <i>R</i> -matrix type | Quantum group                       | Spectral parameter                | Curve          |
|-----------------------|-------------------------------------|-----------------------------------|----------------|
| Rational              | Yangian $Y(\mathfrak{g})$           | $u = z_1 - z_2$                   | $\mathbb{P}^1$ |
| Trigonometric         | Quantum loop $U_q(L\mathfrak{g})$   | $q = e^{2\pi i(z_1 - z_2)/\beta}$ | $\mathbb{C}^*$ |
| Elliptic              | Elliptic QG $E_{q,p}(\mathfrak{g})$ | $u \in E_\tau$                    | $E_\tau$       |

The passage rational  $\rightarrow$  trigonometric  $\rightarrow$  elliptic parallels additive  $\rightarrow$  multiplicative  $\rightarrow$  modular spectral parameter.

*Construction 14.4 (Elliptic shadow data).* Toroidal and elliptic quantum group algebras carry shadow data in the elliptic regime:

- (i) The elliptic propagator  $K^{\text{ell}}(z, w; \tau) = \vartheta_1(z - w; \tau) / \vartheta'_1(0; \tau)$  replaces the rational propagator  $1/(z - w)$ .
- (ii) The Fay trisecant identity for  $\vartheta_1$  is the genus-1 MC equation  $D^{(1)}\Theta + \frac{1}{2}[\Theta^{(0)}, \Theta^{(0)}] + \kappa_{\text{ch}} \omega_1 = 0$ .
- (iii) The Arnold relation  $K_{12}K_{23} + K_{23}K_{31} + K_{31}K_{12} = 0$  (Fay identity) specializes the CYBE to the elliptic regime.

## 14.1 Double affine setup

### Definition

*Definition 14.5 (Toroidal algebra).* The *toroidal algebra* (or double affine algebra)  $U_{q,t}(\hat{\mathfrak{g}})$  is generated by:

- Horizontal currents  $x_{i,n}^{\pm}, \psi_{i,n}^{\pm}$  (affine in one direction)
- Vertical currents  $e_i(z), f_i(z), \psi_i(z)$  (affine in the other direction)

subject to:

1. *Drinfeld-type relations* in each direction:  $[h_{i,m}, x_{j,n}^{\pm}] = \pm a_{ij} x_{j,m+n}^{\pm}, [x_{i,m}^+, x_{j,n}^-] = \delta_{ij} (\psi_{i,m+n}^+ - \psi_{i,m+n}^-) / (q_i - q_i^{-1})$
2. *Cross relations* coupling horizontal and vertical:  $\psi_i(z) e_j(w) = g_{ij}(z/w) e_j(w) \psi_i(z)$  where  $g_{ij}(u) = (u - q^{a_{ij}}) / (u - q^{-a_{ij}})$
3. *Serre relations* in both directions

with two independent parameters  $q$  and  $t = q^k$ . See Ginzburg–Kapranov–Vasserot for the complete presentation.

**PROPOSITION 14.6** (*Toroidal OPE [15];<sup>[PE]</sup>*). The OPE for toroidal currents involves both parameters:

$$e_i(z) f_j(w) = \frac{\delta_{ij}}{(1-q)(1-t^{-1})} \cdot \frac{\psi_i^+(w) - \psi_i^-(w)}{z-w} + \text{rational terms in } q, t$$

*Proof.* We derive the OPE from the defining relations of Definition 14.5.

*Step 1: Mode commutator.* The Drinfeld-type relation (1) gives:

$$[e_{i,m}, f_{j,n}] = \delta_{ij} \frac{\psi_{i,m+n}^+ - \psi_{i,m+n}^-}{q_i - q_i^{-1}}.$$

Form the generating functions  $e_i(z) = \sum_m e_{i,m} z^{-m-1}$ ,  $f_j(w) = \sum_n f_{j,n} w^{-n-1}$ , and  $\psi_i^{\pm}(w) = \sum_n \psi_{i,n}^{\pm} w^{-n}$ .

*Step 2: Extracting the OPE.* Contracting the commutator against  $z^{-m-1} w^{-n-1}$  and summing:

$$e_i(z) f_j(w) - :e_i(z) f_j(w): = \delta_{ij} \sum_m \frac{\psi_{i,m}^+(w) - \psi_{i,m}^-(w)}{q_i - q_i^{-1}} \cdot \frac{w^{-m}}{z-w}.$$

For the toroidal algebra with parameters  $q$  and  $t = q^k$ , we have  $q_i = q^{d_i}$  where  $d_i = (\alpha_i, \alpha_i)/2$ . The singular part of the OPE is the residue at  $z = w$ :

$$e_i(z)f_j(w) \sim \frac{\delta_{ij}}{q_i - q_i^{-1}} \cdot \frac{\psi_i^+(w) - \psi_i^-(w)}{z - w}.$$

*Step 3: Two-parameter structure.* The cross relation (2) introduces the second parameter. From  $\psi_i(z)e_j(w) = g_{ij}(z/w)e_j(w)\psi_i(z)$  with  $g_{ij}(u) = (u - q^{a_{ij}})/(u - q^{-a_{ij}})$ , the normal ordering of  $\psi$  against  $e, f$  currents produces corrections rational in both  $q$  and  $t$ . Specifically, expanding  $1/(q_i - q_i^{-1}) = 1/((q^{d_i} - q^{-d_i}))$  and using  $t = q^k$  to express the central element as  $c = (1 - q)(1 - t^{-1})/(q_i - q_i^{-1})$  (the standard Ding–Iohara normalization). See [10] for the complete change-of-basis computation.  $\square$

**CONJECTURE 14.7** (*Elliptic-curve toroidal realization;  $^{[CJ]}$* ). The toroidal algebra  $U_{q,t}(\hat{\mathfrak{g}})$  admits a curve-based  $E_1$ -chiral realization on the elliptic curve  $E_\tau$  (with modulus  $\tau = \frac{\log t}{\log q}$ ).

**CONJECTURE 14.8** (*Double-affine surface-factorization realization;  $^{[CJ]}$* ). The same toroidal data should extend to a doubly-graded surface-factorization or double-affine object on  $\mathbb{C}^* \times \mathbb{C}^*$ , with one direction encoding the spectral parameter and the other the elliptic coordinate. (This is the toroidal/elliptic extension flank of Vol I, Conjecture III, not the standard  $W_\infty$ /Yangian finite-packet lane.)

*Remark 14.9* (Scope). Conjecture 14.7 is conjectural ( $^{[CJ]}$ ): a complete proof requires verifying associativity of the  $E_1$ -chiral product directly from the OPE relations, since the BD locality approach applies only to commutative chiral algebras. The  $\mathbb{C}^* \times \mathbb{C}^*$  realization (Conjecture 14.8) is a separate surface-factorization problem within the MC4 elliptic/double-affine extension problem.

*Homotopy template* (Vol I, Convention III): Type I (existence).

*Remark 14.10* (Evidence). We describe the conjectural  $E_1$ -chiral structure.

*Step 1: R-matrix braiding.* The cross relation (2) of Definition 14.5 gives the structure function  $g_{ij}(z/w) = (z/w - q^{a_{ij}})/(z/w - q^{-a_{ij}})$ . Define the  $R$ -matrix-valued braiding on generating currents by

$$\sigma_{12} \circ \mu(e_i(z), e_j(w)) = g_{ij}(z/w) \cdot \mu(e_j(w), e_i(z)).$$

Since  $g_{ij}(u)$  is a rational function with  $g_{ij}(u) \neq 1$  for generic  $q$  and  $a_{ij} \neq 0$ , the braiding is non-trivial: the algebra is associative but not commutative, hence strictly  $E_1$ .

*Step 2: Associativity (gap).* The  $R$ -matrix  $R_{ij}(u) = g_{ij}(u) \cdot \text{Id}$  satisfies the Yang–Baxter equation trivially (being scalar-valued). The genuine content of associativity lies in verifying that the full matrix-valued OPE structure, including the Serre-type relations of Definition 14.5, defines an algebra over  $\text{Ass}^{\text{ch}}$  (Vol I, Definition III). This requires an explicit verification that triple OPEs are consistent, which we do not carry out here.

*Step 3: Realization on  $E_\tau$ .* Set  $\tau = \log t / \log q$  and let  $E_\tau = \mathbb{C}^*/q^{\mathbb{Z}}$ . The generating currents  $e_i(z), f_i(z), \psi_i(z)$  are meromorphic on  $\mathbb{C}^*$  with quasi-periodicity  $e_i(qz) = q^{d_i}e_i(z)$  (from the grading), so they define sections of line bundles on  $E_\tau$ . The OPE of Proposition 14.6 has poles at  $z/w \in q^{\mathbb{Z}}$ , which descend to a single point on  $E_\tau$ . This defines a candidate  $E_1$ -chiral algebra structure on  $E_\tau$ , pending verification of Step 2.

*Step 4: Surface-factorization horizon on  $\mathbb{C}^* \times \mathbb{C}^*$ .* The horizontal modes  $x_{i,n}^\pm$  depend on the spectral parameter  $u$  and the vertical modes  $e_i(z)$  depend on  $z$ . Together they give fields on  $\mathbb{C}_u^* \times \mathbb{C}_z^*$  with the two independent gradings (by horizontal and vertical mode number) providing the doubly-graded structure. This belongs to a separate conjectural surface-factorization or double-affine track, not automatically a curve-based  $E_1$ -chiral algebra. A correct formulation would require a bona fide two-parameter factorization framework.

Conditional on Step 2 (i.e., the verification that the toroidal OPE defines an associative  $\text{Ass}^{\text{ch}}$ -algebra structure, which requires checking consistency of triple OPEs with the Serre-type relations), the  $E_1$ -chiral Koszul duality framework (Vol I, Theorem III) applies to the elliptic-curve realization of Conjecture 14.7. The  $\mathbb{C}^* \times \mathbb{C}^*$  surface track belongs instead to Conjecture 14.8.

### Koszul dual of the toroidal algebra

CONJECTURE 14.II (*Toroidal Koszul dual*; <sup>[CJ]</sup>). The  $E_1$ -chiral Koszul dual of the toroidal algebra  $U_{q,t}(\hat{\mathfrak{g}})$  is isomorphic to  $U_{q^{-1},t^{-1}}(\hat{\mathfrak{g}})$ , the toroidal algebra with inverted parameters. (Contributing to the toroidal/elliptic extension flank of Vol I, Conjecture III.)

Remark 14.12 (Evidence). Conditional on Conjecture 14.7, the evidence is threefold:

- (i) *RTT presentation*. By  $E_1$ -chiral Koszul duality (Vol I, Theorem III), the dual has  $R$ -matrix  $R(u; q, t)^{-1} = R(u; q^{-1}, t^{-1})$  (Ding–Iohara inversion; cf. Vol I, Theorem III).
- (ii) *Affine degeneration*. In the limit  $t \rightarrow 0$ ,  $U_{q,t}(\hat{\mathfrak{g}}) \rightarrow U_q(\hat{\mathfrak{g}})$  and the dual reduces to  $U_{q^{-1}}(\hat{\mathfrak{g}})$ , consistent with Vol I, Theorem III.
- (iii) *Central charge complementarity*. The inversion  $q \rightarrow q^{-1}$  sends  $\tau \rightarrow -\tau$ , reversing the complex structure of  $E_\tau$ , the elliptic analogue of the Verdier intertwining (Vol I, Theorem A; Theorem III).

The Fay identity (Proposition 14.21) serves as the elliptic Arnold relation. The universal MC class predicts  $\Theta_{U_{q,t}} + \Theta_{U_{q^{-1},t^{-1}}} = 0$ .

*Homotopy template* (Vol I, Convention III): Type II (equivalence).

Remark 14.13 (Miki automorphism and Koszul duality). The quantum toroidal algebra  $U_{q,t}(\hat{\mathfrak{g}})$  carries an  $\mathrm{SL}_2(\mathbb{Z})$  automorphism group, discovered by Miki [26] and implicit in the Ding–Iohara construction [10]. Write  $(q_1, q_2, q_3)$  for the three parameters satisfying  $q_1 q_2 q_3 = 1$ , with  $q = q_1$  and  $t = q_2^{-1}$  in the  $(q, t)$  convention. The *Miki automorphism*  $S$  is the generator of  $\mathrm{SL}_2(\mathbb{Z})$  that acts by cyclic permutation:

$$S: (q_1, q_2, q_3) \mapsto (q_2, q_3, q_1),$$

so that  $S^3 = \mathrm{id}$  on parameters (triality). At the level of generators,  $S$  exchanges the horizontal subalgebra (generated by  $E(z), F(z)$ ) with the vertical subalgebra (generated by  $K^\pm(z)$ ):

$$S: E(z) \leftrightarrow K^+(z), \quad F(z) \leftrightarrow K^-(z) \quad (\text{up to normalization}).$$

This is the algebraic incarnation of the exchange of the two  $\mathbb{C}^*$  factors in the torus  $\mathbb{C}^* \times \mathbb{C}^*$  underlying the double loop.

Three properties connect  $S$  to the Koszul duality of Conjecture 14.II:

- (a) *Parameter inversion*. In  $(q, t)$  coordinates,  $S$  acts by  $q \mapsto 1/t$ ,  $t \mapsto q/t$ , and  $S^2$  acts by  $q \mapsto t/q$ ,  $t \mapsto 1/q$ . The composite  $S^3 = \mathrm{id}$  gives the triality. The Koszul involution  $(q, t) \mapsto (q^{-1}, t^{-1})$  is NOT one of the six  $\mathrm{SL}_2(\mathbb{Z})$  images; it is a separate involution that commutes with the  $\mathrm{SL}_2(\mathbb{Z})$  action. The Koszul involution sends  $(q_1, q_2, q_3) \mapsto (q_1^{-1}, q_2^{-1}, q_3^{-1})$  (total inversion, preserving  $q_1 q_2 q_3 = 1$ ).
- (b) *Affine Yangian obstruction*. The affine Yangian  $Y(\hat{\mathfrak{gl}}_1)$  is the rational degeneration  $q_1 \rightarrow 1$ ; in this limit, the three parameters collapse to two  $(\epsilon_1, \epsilon_2)$  with  $\epsilon_1 + \epsilon_2 + \epsilon_3 = 0$ , and the  $\mathrm{SL}_2(\mathbb{Z})$  degenerates to at most an  $S_3$  permutation on  $(\epsilon_1, \epsilon_2, \epsilon_3)$ , which does not lift to a full algebra automorphism. The Miki automorphism is a genuinely quantum-toroidal phenomenon; it requires both loops to be on equal footing (cf. the compute engine `quantum_toroidal_e1_cy3.py`, which verifies  $S^3 = \mathrm{id}$  numerically at multiple specialisations and confirms the absence of  $S$  for the affine Yangian).
- (c) *Spectral-parameter exchange for the DDCA*. In the rational limit, the double deformation current algebra (DDCA) carries two spectral parameters  $(u, v)$ . The rational shadow of  $S$  should exchange these two parameters, intertwining the horizontal and vertical RTT presentations. This exchange is compatible with the  $R$ -matrix inversion  $R(u; q, t)^{-1} = R(u; q^{-1}, t^{-1})$  appearing in item (i) of Remark 14.12: the Miki automorphism provides a *second* route to the Koszul dual, via the internal symmetry of  $U_{q,t}$  rather than the external bar-cobar construction. The precise rational degeneration identifying the DDCA as the  $\hbar_1, \hbar_2 \rightarrow 0$  limit of the toroidal algebra is Conjecture 14.197; the spectral-parameter exchange  $\sigma: (u, v) \mapsto (v, u)$  of the DDCA as the rational shadow of  $S$  is Remark 14.199.

The distinction between  $S$  (internal  $\mathrm{SL}_2(\mathbb{Z})$  automorphism) and Koszul duality (external bar-cobar inversion) is essential:  $S$  is an automorphism of  $U_{q,t}$ , while Koszul duality produces a *different* algebra  $U_{q^{-1},t^{-1}}$ . At the level of parameters, the Koszul involution  $(q_1, q_2, q_3) \mapsto (q_1^{-1}, q_2^{-1}, q_3^{-1})$  commutes with the cyclic permutation  $S$ , so the  $\mathrm{SL}_2(\mathbb{Z})$  symmetry of  $U_{q,t}$  descends to an  $\mathrm{SL}_2(\mathbb{Z})$  symmetry of the Koszul dual  $U_{q^{-1},t^{-1}}$  with identical parameter action.

### Connection to the three main theorems of Vol I

*Remark 14.14* (Toroidal three theorems). Conditional on Conjecture 14.7, the expected toroidal analogues of the three main theorems of Vol I are: *Theorem A* (Vol I, bar-cobar adjunction):  $\bar{B}^{\mathrm{ell}}(U_{q,t})$  computes the Koszul dual coalgebra (nilpotency via Proposition 14.21). *Theorem B* (Vol I, bar-cobar inversion):  $\Omega(\bar{B}(U_{q,t})) \xrightarrow{\sim} U_{q,t}$  by the rational spectral sequence with elliptic corrections (Theorem 14.24). *Theorem C* (Vol I, complementarity):  $\mathrm{obs}_g(U_{q,t}) + \mathrm{obs}_g(U_{q^{-1},t^{-1}})$  controlled by  $H^*$  of a framed moduli space (the  $E_1$  replacement for  $\overline{\mathcal{M}}_g$ ; cf. Vol I, Remark III). The universal MC class  $\Theta_{U_{q,t}}$  should govern the genus expansion, with  $(q, t) \mapsto (q^{-1}, t^{-1})$  reflected in Verdier duality.

*Remark 14.15* (Lattice evidence for toroidal genus theory). The curvature-braiding orthogonality theorem for quantum lattice VOAs (Vol I, Theorem III) provides structural evidence for the toroidal case. For the lattice, the genus-1 curvature  $\kappa_{\mathrm{ch}} = \mathrm{rank}(\Lambda)$  lives entirely in the Cartan (Heisenberg) sector and is independent of the deformation cocycle  $\varepsilon_{N,q}$ . The toroidal algebra  $U_{q,t}(\hat{\mathfrak{g}})$  has a similar double-loop structure: the inner loop ( $\hat{\mathfrak{g}}$ ) is the Cartan analog, while the outer loop (the second affinization) provides the braiding. The orthogonality principle predicts that the toroidal genus-1 curvature should be controlled by the inner-loop data alone, with the dynamical parameter  $\lambda \in H^1(E_\tau)$  entering the braiding axis rather than the modular axis. The dynamical YBE for  $R(u, \lambda)$  would then be the toroidal analog of the modified Arnold relation at genus 1, with the dynamical shift  $\lambda \rightarrow \lambda - h^{(i)}$  replacing the B-cycle monodromy correction.

## 14.2 Elliptic quantum groups

### Felder's elliptic quantum group

*Definition 14.16* (Elliptic quantum group). The *elliptic quantum group*  $E_{q,p}(\mathfrak{g})$  (Felder) is defined by:

- The dynamical R-matrix  $R(u, \lambda)$  depending on spectral parameter  $u$  and dynamical parameter  $\lambda$ .
- The RLL relations:

$$R(u - v, \lambda) L_1(u, \lambda) L_2(v, \lambda - h^{(1)}) = L_2(v, \lambda) L_1(u, \lambda - h^{(2)}) R(u - v, \lambda)$$

*Definition 14.17* (Elliptic R-matrix). The elliptic R-matrix for  $\mathfrak{sl}_2$  is:

$$R(u, \lambda) = \begin{pmatrix} a(u) & 0 & 0 & 0 \\ 0 & b(u, \lambda) & c(u, \lambda) & 0 \\ 0 & c(u, -\lambda) & b(u, -\lambda) & 0 \\ 0 & 0 & 0 & a(u) \end{pmatrix}$$

where  $a(u)$ ,  $b(u, \lambda)$ ,  $c(u, \lambda)$  are expressed via theta functions:

$$\begin{aligned} a(u) &= \frac{\theta(u + \eta)}{\theta(\eta)} \\ b(u, \lambda) &= \frac{\theta(u)\theta(\lambda + \eta)}{\theta(\eta)\theta(\lambda)} \\ c(u, \lambda) &= \frac{\theta(u + \lambda)\theta(\eta)}{\theta(\lambda)\theta(u + \eta)} \end{aligned}$$

*Remark 14.18* (Elliptic quantum groups and bar complex). We expect <sup>[CJ]</sup>, conditional on Conjecture 14.7): (i)  $\bar{B}(U_{q,t})$  on  $E_\tau$  computes the chiral coalgebra  $E_{q,p}(\mathfrak{g})$  (elliptic Kazhdan–Lusztig); (ii) the dynamical parameter  $\lambda$  (Definition 14.16) is the B-cycle monodromy of the elliptic propagator (elliptic analogue of  $k \rightarrow k + h^\vee$ ); (iii) the dynamical YBE for  $R(u, \lambda)$  is the elliptic deformation of  $d^2 = 0$  (Proposition 14.21), with the dynamical parameter from  $H^1(E_\tau) \neq 0$ .

### 14.3 Elliptic R-matrices and theta functions

#### Theta function identities

*Definition 14.19* (Jacobi theta function). The Jacobi theta function is:

$$\theta(u|\tau) = \sum_{n \in \mathbb{Z}} (-1)^n e^{\pi i \tau n(n-1) + 2\pi i n u} = (e^{2\pi i u}; p)_\infty (p e^{-2\pi i u}; p)_\infty (p; p)_\infty$$

where  $p = e^{2\pi i \tau}$  and  $(a; q)_\infty = \prod_{n \geq 0} (1 - a q^n)$ .

**THEOREM 14.20** (*Fay identity at genus 1* [14]; <sup>[PE]</sup>). For any four points  $u_1, u_2, u_3, u_4$  on an elliptic curve  $E_\tau$ , the theta function satisfies the Fay trisecant identity:

$$\theta(u_1 - u_2) \theta(u_3 - u_4) - \theta(u_1 - u_3) \theta(u_2 - u_4) + \theta(u_1 - u_4) \theta(u_2 - u_3) = 0$$

This is a *bilinear* identity (each term is a product of two theta functions), which at genus 1 is the degenerate case of the general Fay trisecant identity on higher-genus curves. The identity is equivalent to the Riemann addition formula and reduces to the Ptolemy relation  $\sin(\alpha - \beta) \sin(\gamma - \delta) - \sin(\alpha - \gamma) \sin(\beta - \delta) + \sin(\alpha - \delta) \sin(\beta - \gamma) = 0$  in the rational limit  $\tau \rightarrow i\infty$ .

This identity underlies the Yang–Baxter equation for elliptic R-matrices: the star-triangle relation is a consequence of the Fay identity applied to the structure functions of the elliptic algebra.

#### Fay identity and bar nilpotency

**PROPOSITION 14.21** (*Fay identity implies elliptic  $d^2 = 0$* ; <sup>[PH]</sup>). On  $\bar{C}_3(E_\tau)$ , the elliptic bar differential satisfies  $d^2 = 0$ . The key algebraic input is the Fay trisecant identity (Theorem 14.20), which plays the role of the Arnold relation in the rational case (Vol I, Theorem III).

*Proof.* The elliptic bar differential on degree-2 elements uses the propagator  $\eta_{ij}^{\text{ell}} = d \log \theta_1(z_i - z_j | \tau)$ . The computation of  $d^2$  on a degree-1 element  $[a] \otimes 1$  produces terms on  $\bar{C}_3(E_\tau)$  involving the 2-form  $\eta_{12}^{\text{ell}} \wedge \eta_{23}^{\text{ell}} + \eta_{23}^{\text{ell}} \wedge \eta_{31}^{\text{ell}} + \eta_{31}^{\text{ell}} \wedge \eta_{12}^{\text{ell}}$ , exactly as in the rational case (cf. Vol I, §III).

The rational Arnold relation  $\eta_{12} \wedge \eta_{23} + \eta_{23} \wedge \eta_{31} + \eta_{31} \wedge \eta_{12} = 0$  (where  $\eta_{ij} = d \log(z_i - z_j)$ ) is a consequence of the partial fraction identity  $(z_1 - z_2)^{-1} (z_2 - z_3)^{-1} + \text{cyclic} = 0$ .

The elliptic replacement for  $(z_i - z_j)^{-1}$  is  $\zeta_\tau(z_i - z_j) := \partial_z \log \theta_1(z_i - z_j | \tau)$ , the Weierstrass  $\zeta$ -function on  $E_\tau$ . The corresponding three-term identity is

$$d\zeta_\tau(z_1 - z_2) \wedge d\zeta_\tau(z_2 - z_3) + d\zeta_\tau(z_2 - z_3) \wedge d\zeta_\tau(z_3 - z_1) + d\zeta_\tau(z_3 - z_1) \wedge d\zeta_\tau(z_1 - z_2) = 0. \quad (14.1)$$

This is the differential-form shadow of the Fay identity (Theorem 14.20). Differentiate

$$\theta(u_1 - u_2) \theta(u_3 - u_4) - \theta(u_1 - u_3) \theta(u_2 - u_4) + \theta(u_1 - u_4) \theta(u_2 - u_3) = 0$$

with respect to  $u_1$  and  $u_4$ , then set  $u_4 = u_1$ . The resulting trisecant consequence is

$$\begin{aligned} \zeta_\tau(u_1 - u_2) \zeta_\tau(u_2 - u_3) + \zeta_\tau(u_2 - u_3) \zeta_\tau(u_3 - u_1) + \zeta_\tau(u_3 - u_1) \zeta_\tau(u_1 - u_2) \\ = \wp_\tau(u_1 - u_2) + \wp_\tau(u_2 - u_3) + \wp_\tau(u_3 - u_1) \end{aligned}$$

where  $\wp_\tau = -\zeta'_\tau$  is the Weierstrass  $\wp$ -function. Now take exterior derivatives in the difference variables. Each  $\wp_\tau(z_i - z_j)$  depends on a single difference, so the mixed wedges on the right vanish or cancel pairwise. The left-hand side is then exactly (14.1).

The remaining terms in  $d^2$  (involving OPE data of the chiral algebra  $\mathcal{A}$ ) cancel by the same mechanism as in the rational case: the bar differential decomposes as  $d = d_{\text{local}} + d_{\text{global}}$ , where  $d_{\text{local}}^2 = 0$  by the OPE associativity of  $\mathcal{A}$ , and the cross-terms  $d_{\text{local}} \circ d_{\text{global}} + d_{\text{global}} \circ d_{\text{local}} = 0$  by the compatibility of the OPE with the propagator (which holds for both the rational and elliptic propagators, since the residue at a simple pole is the same:  $\text{Res}_{z=0} \zeta_\tau(z) dz = 1$ ).  $\square$

*Remark 14.22 (Rational limit).* In the rational limit  $\tau \rightarrow i\infty$ , the theta function  $\theta_1(z|\tau) \rightarrow 2 \sin(\pi z) \rightarrow 2\pi z$  (for  $|z|$  small), so  $\zeta_\tau(z) \rightarrow 1/z$  and  $\wp_\tau(z) \rightarrow 1/z^2$ . The identity (14.1) reduces to the Arnold relation  $\eta_{12} \wedge \eta_{23} + \eta_{23} \wedge \eta_{31} + \eta_{31} \wedge \eta_{12} = 0$  (Vol I, Theorem III), and the bar nilpotency reduces to Vol I, Theorem III.

### Bar complex with theta functions

*Construction 14.23 (Elliptic bar complex).* The bar complex of an elliptic chiral algebra incorporates theta functions:

*Forms.* Replace  $d \log(z_1 - z_2)$  with:

$$\eta_{12} = d \log \theta(z_1 - z_2|\tau)$$

the logarithmic derivative of theta.

*Differential.* The residue is computed at the zeros of  $\theta$ :

$$d_{\text{res}}([a|b] \otimes \eta_{12}) = \sum_{\theta(z_0)=0} \text{Res}_{z_1=z_0+z_2} [\cdots]$$

The zeros of  $\theta(u|\tau)$  are at  $u = m + n\tau$  for  $m, n \in \mathbb{Z}$  (the lattice points).

**THEOREM 14.24 (Elliptic vs rational homology; <sup>[PH]</sup>).** Let  $\mathcal{A}$  be a chiral algebra of central charge  $c$  on an elliptic curve  $E_\tau$ . The homology of the elliptic bar complex admits a decomposition:

$$H_n(\bar{B}^{\text{ell}}(\mathcal{A})) \cong H_n(\bar{B}^{\text{rat}}(\mathcal{A})) \oplus \bigoplus_{k \geq 1} \mathcal{E}_n^{(k)}(\mathcal{A}, \tau)$$

where the elliptic corrections  $\mathcal{E}_n^{(k)}(\mathcal{A}, \tau)$  are  $\mathcal{A}$ -modules whose dependence on  $\tau$  is through (quasi-)modular forms. Specifically:

- (i) The leading correction  $\mathcal{E}_n^{(1)}$  is proportional to  $E_2(\tau)$ , the weight-2 quasi-modular Eisenstein series, arising from the regularization of the elliptic propagator  $\partial_z \log \theta(z|\tau)$  relative to the rational propagator  $1/z$  (cf. Construction 14.23).
- (ii) Higher corrections  $\mathcal{E}_n^{(k)}$  for  $k \geq 2$  lie in the space of modular forms  $M_{2k}(\text{SL}_2(\mathbb{Z}))$  of weight  $2k$ ; the first truly modular correction is an  $E_4(\tau)$ -coefficient.

- (iii) The conformal dimension  $h$  of the states entering the bar complex constrains the weights: a state of dimension  $h$  contributes to  $\mathcal{E}_n^{(k)}$  only for  $k \leq h + n$ .
- (iv) At the rational limit  $\tau \rightarrow i\infty$  (i.e.,  $q = e^{2\pi i\tau} \rightarrow 0$ ), the Eisenstein series  $E_{2k}(\tau) \rightarrow 1$  and all corrections degenerate to constants, recovering  $H_n(\bar{B}^{\text{rat}}(\mathcal{A}))$  up to scalar renormalization.

For the Heisenberg algebra ( $c = 1$ ), the corrections reduce to the Eisenstein series hierarchy computed in Vol I, §III.

*Proof.* We construct the decomposition via a perturbative spectral sequence comparing the elliptic and rational bar differentials.

*Step 1 (Propagator decomposition).* The elliptic bar complex (Construction 14.23) uses the propagator  $\eta_{ij}^{\text{ell}} = d \log \theta_1(z_i - z_j | \tau)$ , while the rational bar complex uses  $\eta_{ij}^{\text{rat}} = d \log(z_i - z_j)$ . Near  $z_i = z_j$ , both propagators have the same simple pole with residue 1, so their difference  $\delta_{ij}(\tau) := \eta_{ij}^{\text{ell}} - \eta_{ij}^{\text{rat}}$  is a regular 1-form. The classical expansion of the Weierstrass  $\zeta$ -function  $\zeta(z; \Lambda_\tau)$  in terms of the Eisenstein–Weierstrass series  $G_{2k}(\tau)$  gives the Laurent expansion:

$$\partial_z \log \theta_1(z | \tau) = \frac{1}{z} + \sum_{k=1}^{\infty} c_k(\tau) z^{2k-1},$$

where the correction coefficients  $c_k(\tau)$  are expressed in terms of Eisenstein series via  $G_{2k}(\tau) = 2\zeta_R(2k) E_{2k}(\tau)$ :

- $c_1(\tau) = -\frac{\pi^2}{3} E_2(\tau) \in \widetilde{M}_2(\text{SL}_2(\mathbb{Z}))$  (quasi-modular, weight 2), since  $c_1 = -2\eta_1$  where  $\eta_1 = \zeta(1/2; \Lambda_\tau) = \frac{\pi^2}{6} E_2(\tau)$ ;
- $c_k(\tau) = -G_{2k}(\tau) \in M_{2k}(\text{SL}_2(\mathbb{Z}))$  for  $k \geq 2$  (holomorphic modular forms).

The expansion converges absolutely for  $|z| < \min_{\omega \in \Lambda_\tau \setminus \{0\}} |\omega|$ .

*Step 2 (Correction-order filtration).* Define the *correction-order filtration*  $\mathcal{F}^p \bar{B}_n^{\text{ell}}(\mathcal{A})$  by declaring that an element has filtration degree  $\geq p$  if it arises from at least  $p$  insertions of the regular correction terms  $\{c_k(\tau) z^{2k-1}\}_{k \geq 1}$  in the propagator expansion. Equivalently, replace each propagator by  $1/z + t \cdot (c_1 z + c_2 z^3 + \dots)$  and expand in powers of the formal parameter  $t$ ; then  $\mathcal{F}^p$  consists of terms of order  $\geq t^p$ .

Since the bar differential  $d^{\text{ell}}$  at degree  $n$  involves at most  $n$  propagator factors, we have  $\mathcal{F}^{n+1} \bar{B}_n^{\text{ell}} = 0$ . The filtration is compatible with the differential:  $d^{\text{ell}}(\mathcal{F}^p) \subset \mathcal{F}^p$  (inserting the bar differential does not decrease the number of correction insertions).

*Step 3 (Spectral sequence:  $E_0$  and  $E_1$  pages).* The associated graded  $\text{gr}_{\mathcal{F}}^0 \bar{B}^{\text{ell}}(\mathcal{A})$  is the bar complex computed using only the singular part  $1/z$  of the propagator. Since the residue extraction in the bar differential (which encodes the OPE data of  $\mathcal{A}$ ) depends only on the polar part, the differential on  $\text{gr}^0$  coincides with the rational bar differential  $d^{\text{rat}}$ . Thus:

$$E_1^{0,n} = H_n(\bar{B}^{\text{rat}}(\mathcal{A})).$$

More generally,  $E_1^{p,n} = H_n(\bar{B}^{\text{rat}}(\mathcal{A})) \otimes \text{Sym}_{\geq 1}^p \{c_k(\tau)\}$ , where the symmetric product is taken over all propagator correction coefficients contributing total weight  $p$ .

*Step 4 (Modular structure via Zhu's theorem).* The differential  $d_r : E_r^{p,n} \rightarrow E_r^{p+r,n+1}$  on the  $r$ -th page arises from the interaction of  $r$  correction insertions with the bar differential. By Zhu's modular invariance theorem [39], the  $n$ -point genus-1 correlation functions  $\langle V_1(z_1) \cdots V_n(z_n) \rangle_{E_\tau}$  of a rational



vertex algebra  $\mathcal{A}$  transform as components of a vector-valued modular form for  $\mathrm{SL}_2(\mathbb{Z})$ . Since the bar complex at degree  $n$  is built from such correlation functions composed with the propagator form, and the correction coefficients  $c_k(\tau)$  are themselves (quasi-)modular, the differentials  $d_r$  respect the modular weight grading. Consequently, the  $E_\infty$  terms inherit a weight decomposition from the ring  $\tilde{M}_*(\mathrm{SL}_2(\mathbb{Z}))$  of quasi-modular forms.

*Step 5 (Convergence and splitting).* The spectral sequence converges because the filtration is bounded:  $\mathcal{F}^{n+1}\bar{B}_n^{\mathrm{ell}} = 0$  at each bar degree  $n$ . At convergence,  $H_n(\bar{B}^{\mathrm{ell}}(\mathcal{A}))$  admits the filtration

$$0 = \mathcal{F}^{n+1}H_n \subset \mathcal{F}^n H_n \subset \cdots \subset \mathcal{F}^1 H_n \subset \mathcal{F}^0 H_n = H_n(\bar{B}^{\mathrm{ell}}(\mathcal{A}))$$

with successive quotients  $\mathcal{E}_n^{(k)} := E_\infty^{k,n}$ . The filtration *splits*: the modular weight provides a grading on the coefficient ring (quasi-modular forms of different weights are linearly independent over  $\mathbb{C}$ ), and the  $E_\infty$  terms in different weights cannot be related by extensions. We therefore obtain the direct sum decomposition  $H_n(\bar{B}^{\mathrm{ell}}(\mathcal{A})) \cong H_n(\bar{B}^{\mathrm{rat}}(\mathcal{A})) \oplus \bigoplus_{k \geq 1} \mathcal{E}_n^{(k)}(\mathcal{A}, \tau)$  as claimed.

Properties (i)–(iv) now follow from the structure of the correction coefficients:

- (i) The leading correction involves  $c_1(\tau) = -(\pi^2/3) E_2(\tau)$  (quasi-modular, weight 2).
- (ii) Higher corrections involve products of  $c_k(\tau) \in M_{2k}(\mathrm{SL}_2(\mathbb{Z}))$  for  $k \geq 2$ .
- (iii) A bar element of degree  $n$  involves at most  $n$  propagator factors; a state of conformal dimension  $h$  constrains the Laurent order of each propagator insertion to  $\leq 2h - 1$ , giving the bound  $\mathcal{E}_n^{(k)} = 0$  for  $k > h + n$ .
- (iv) As  $\tau \rightarrow i\infty$ :  $q \rightarrow 0$  and  $E_{2k}(\tau) \rightarrow 1$ , so all corrections reduce to constants, recovering  $H_n(\bar{B}^{\mathrm{rat}}(\mathcal{A}))$  up to renormalization.

*Verification for the Heisenberg algebra.* For  $\mathcal{A} = \mathcal{H}(c = 1)$ , the genus-1 two-point function is  $\langle a(z_1)a(z_2) \rangle_{E_\tau} = \kappa_{\mathrm{ch}} G_\tau(z_1 - z_2)$  with  $G_\tau(z) = \zeta_\tau(z) - (\pi^2 E_2(\tau)/3)z$  (Vol I, Theorem III). The leading correction  $\mathcal{E}_n^{(1)} \propto E_2(\tau)$  arises from the  $c_1$  term. The complete Eisenstein hierarchy of Vol I, §III gives  $\mathcal{E}_n^{(k)} \propto E_{2k}(\tau)$ , confirming the general pattern.  $\square$

*Remark 14.25 (Scope).* The proof assembles three ingredients: (i) the Laurent expansion of the Weierstrass  $\zeta$ -function in terms of Eisenstein series ([PE], classical); (ii) the convergence of the correction-order spectral sequence (bounded filtration at each bar degree); and (iii) Zhu's modular invariance theorem for genus-1  $n$ -point functions of rational vertex algebras [39] ([PE]). The combination and the identification of the splitting mechanism (Step 5) are new to this volume. The Heisenberg verification uses the explicit computations of Vol I, §III.

## 14.4 Explicit elliptic bar complex for the Heisenberg algebra

### Bar elements and the elliptic propagator

We work on  $E_\tau = \mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$  with the elliptic propagator

$$\eta_{ij}^{\mathrm{ell}} = d \log \theta_1(z_i - z_j | \tau) = \zeta_\tau(z_i - z_j) (dz_i - dz_j), \quad (14.2)$$

where  $\zeta_\tau(z) = \partial_z \log \theta_1(z | \tau)$  is the Weierstrass zeta function on  $E_\tau$ .

*Computation 14.26 (Degree-1 bar elements).* In bar degree 1, the generators of  $\bar{B}_1^{\text{ell}}(\mathcal{H}_{\kappa_{\text{ch}}})$  are:

$$[a_m] \otimes 1 \in \bar{B}_1^{\text{ell}}(\mathcal{H}_{\kappa_{\text{ch}}}), \quad m \in \mathbb{Z},$$

where  $a_m$  denotes the mode of the Heisenberg current  $a(z) = \sum_m a_m z^{-m-1}$ . The bar differential on degree-1 elements is:

$$d^{\text{ell}}([a_m] \otimes 1) = 0,$$

since the Heisenberg algebra has trivial unary OPE ( $a_{(n)}a = 0$  for  $n \geq 1$  except  $a_{(1)}a = \kappa_{\text{ch}}$ , and the residue extraction on a single element yields zero). This is identical to the rational case.

*Computation 14.27 (Degree-2 bar elements and the elliptic differential).* In bar degree 2, the generators are:

$$[a_m|a_n] \otimes \eta_{12}^{\text{ell}} \in \bar{B}_2^{\text{ell}}(\mathcal{H}_{\kappa_{\text{ch}}}),$$

where  $\eta_{12}^{\text{ell}} = \zeta_{\tau}(z_1 - z_2)(dz_1 - dz_2)$ . The bar differential acts by residue extraction:

$$\begin{aligned} d^{\text{ell}}([a_m|a_n] \otimes \eta_{12}^{\text{ell}}) &= \text{Res}_{z_1=z_2} \left[ a_m(z_1) a_n(z_2) \eta_{12}^{\text{ell}} \right] \\ &= \text{Res}_{\epsilon=0} \left[ \frac{\kappa_{\text{ch}}}{\epsilon^2} \zeta_{\tau}(\epsilon) d\epsilon \right] \cdot z_2^{-m-n-2} dz_2, \end{aligned} \quad (14.3)$$

where  $\epsilon = z_1 - z_2$ . We use the Laurent expansion of  $\zeta_{\tau}$  (established in the proof of Theorem 14.24, Step 1):

$$\zeta_{\tau}(\epsilon) = \frac{1}{\epsilon} - \frac{\pi^2}{3} E_2(\tau) \epsilon - \frac{\pi^4}{45} E_4(\tau) \epsilon^3 - \frac{2\pi^6}{945} E_6(\tau) \epsilon^5 - \dots \quad (14.4)$$

Substituting into the residue:

$$\begin{aligned} \text{Res}_{\epsilon=0} \left[ \frac{\kappa_{\text{ch}}}{\epsilon^2} \zeta_{\tau}(\epsilon) d\epsilon \right] &= \text{Res}_{\epsilon=0} \left[ \frac{\kappa_{\text{ch}}}{\epsilon^2} \left( \frac{1}{\epsilon} - \frac{\pi^2}{3} E_2(\tau) \epsilon - \dots \right) d\epsilon \right] \\ &= \text{Res}_{\epsilon=0} \left[ \frac{\kappa_{\text{ch}}}{\epsilon^3} - \frac{\kappa_{\text{ch}} \pi^2}{3} E_2(\tau) \frac{1}{\epsilon} - \dots \right] d\epsilon \\ &= -\frac{\kappa_{\text{ch}} \pi^2}{3} E_2(\tau). \end{aligned} \quad (14.5)$$

The  $1/\epsilon^3$  term has zero residue; the first nonzero contribution is the  $E_2(\tau)$  term. In the rational case ( $\zeta_{\tau}(\epsilon) \rightarrow 1/\epsilon$ ), the residue of  $\kappa_{\text{ch}} \epsilon^{-3} d\epsilon$  vanishes, so the rational bar differential on this element is zero. The entire degree-2 bar differential is the elliptic correction:

$$d^{\text{ell}}([a_m|a_n] \otimes \eta_{12}^{\text{ell}}) = -\frac{\kappa_{\text{ch}} \pi^2}{3} E_2(\tau) \cdot z_2^{-m-n-2} dz_2. \quad (14.6)$$

*Computation 14.28 (Degree-2 curvature).* The elliptic bar complex of  $\mathcal{H}_{\kappa_{\text{ch}}}$  has curvature element  $m_0^{\text{ell}} \in \bar{B}_0^{\text{ell}}(\mathcal{H}_{\kappa_{\text{ch}}})$ . From the genus-1 propagator (Vol I, Theorem III), the two-point function on  $E_{\tau}$  is  $\langle a(z_1) a(z_2) \rangle_{E_{\tau}} = \kappa_{\text{ch}} G_{\tau}(z_1 - z_2)$ , where  $G_{\tau}(z) = \zeta_{\tau}(z) - (\pi^2 E_2(\tau)/3) z$ . The curvature arises from the failure of double periodicity (cf. the proof of Vol I, Theorem III):

$$m_0^{\text{ell}} = \kappa_{\text{ch}} \cdot \frac{\pi^2 E_2(\tau)}{3}. \quad (14.7)$$

Expanding in the nome  $q = e^{2\pi i\tau}$ :

$$m_0^{\text{ell}} = \frac{\kappa_{\text{ch}}\pi^2}{3} (1 - 24q - 72q^2 - 96q^3 - \dots). \quad (14.8)$$

At the rational limit  $\tau \rightarrow i\infty$  ( $q \rightarrow 0$ ),  $E_2(\tau) \rightarrow 1$  and  $m_0^{\text{ell}} \rightarrow \kappa_{\text{ch}}\pi^2/3$ , recovering the rational curvature (up to the normalization convention for the propagator on  $\mathbb{C}$  versus  $E_\tau$ ; on  $\mathbb{C}$  the propagator  $1/z$  has no  $B$ -cycle and hence  $m_0^{\text{rat}} = 0$ ).

### Comparison table: elliptic versus rational

Table 14.1: Elliptic versus rational bar differentials for  $\mathcal{H}_{\kappa_{\text{ch}}}$  at low degrees

| Degree                        | Rational $d^{\text{rat}}$ | Elliptic $d^{\text{ell}}$                                                                       | Correction                        |
|-------------------------------|---------------------------|-------------------------------------------------------------------------------------------------|-----------------------------------|
| 1                             | 0                         | 0                                                                                               | —                                 |
| 2                             | 0                         | $-\frac{\kappa_{\text{ch}}\pi^2}{3} E_2(\tau)$                                                  | $E_2(\tau)$ (quasi-modular, wt 2) |
| 3                             | 0                         | $-\frac{\kappa_{\text{ch}}\pi^4}{45} E_4(\tau)$                                                 | $E_4(\tau)$ (modular, wt 4)       |
| 4                             | 0                         | $-\frac{2\kappa_{\text{ch}}\pi^6}{945} E_6(\tau)$                                               | $E_6(\tau)$ (modular, wt 6)       |
| 5                             | 0                         | $-\frac{\kappa_{\text{ch}}\pi^8}{4725} E_8(\tau)$                                               | $E_8(\tau)$ (modular, wt 8)       |
| 6                             | 0                         | $-\frac{2\kappa_{\text{ch}}\pi^{10}}{93555} E_{10}(\tau)$                                       | $E_{10}(\tau)$ (modular, wt 10)   |
| 7                             | 0                         | $-\frac{1382\kappa_{\text{ch}}\pi^{12}}{638512875} E_{12}(\tau)$                                | $E_{12}(\tau)$ (modular, wt 12)   |
| 8                             | 0                         | $-\frac{4\kappa_{\text{ch}}\pi^{14}}{18243225} E_{14}(\tau)$                                    | $E_{14}(\tau)$ (modular, wt 14)   |
| <b>General <math>n</math></b> | 0                         | $-\kappa_{\text{ch}} \cdot \frac{2^{2n-2}  B_{2n-2} }{(2n-2)!} \cdot \pi^{2n-2} E_{2n-2}(\tau)$ | $E_{2n-2}(\tau)$ (wt $2n-2$ )     |

The vanishing of the rational bar differential at all degrees reflects that the Heisenberg algebra on  $\mathbb{C}$  (genus 0) has no curvature ( $m_0^{\text{rat}} = 0$ ): the propagator  $1/z$  is strictly periodic on  $\mathbb{C}$  (there is no  $B$ -cycle). On the elliptic curve  $E_\tau$ , the  $B$ -cycle monodromy of  $\zeta_\tau$  introduces the Eisenstein corrections. At degree  $n$ , the coefficient  $c_{n-1}(\tau)$  in the Laurent expansion (14.4) controls the bar differential; since  $c_k(\tau) \in M_{2k}(\text{SL}_2(\mathbb{Z}))$  for  $k \geq 2$  and  $c_1(\tau) \in \tilde{M}_2(\text{SL}_2(\mathbb{Z}))$  (quasi-modular), the corrections at degree  $n \geq 3$  are genuine modular forms.

PROPOSITION 14.29 (*Decomposition of the elliptic bar complex; <sup>[PH]</sup>*). The elliptic bar complex of  $\mathcal{H}_{\kappa_{\text{ch}}}$  at degree  $n$  admits the decomposition:

$$\bar{B}_n^{\text{ell}}(\mathcal{H}_{\kappa_{\text{ch}}}) = \bar{B}_n^{\text{rat}}(\mathcal{H}_{\kappa_{\text{ch}}}) \oplus E_2(\tau) \cdot \bar{B}_n^{(1)} \oplus E_4(\tau) \cdot \bar{B}_n^{(2)} \oplus \dots \oplus E_{2n-2}(\tau) \cdot \bar{B}_n^{(n-1)} \quad (14.9)$$

with only finitely many nonzero terms. Specifically,  $\bar{B}_n^{(k)} = 0$  for  $k \geq n$ .

*Proof. Step 1 (Finite propagator expansion).* The elliptic bar complex at degree  $n$  involves  $\binom{n}{2}$  propagator factors  $\eta_{ij}^{\text{ell}}$ , one for each pair  $(i, j)$  with  $1 \leq i < j \leq n$ . Each propagator is expanded via (14.4) as  $\zeta_\tau(\epsilon_{ij}) = \epsilon_{ij}^{-1} + \sum_{k \geq 1} c_k(\tau) \epsilon_{ij}^{2k-1}$ . The bar differential at degree  $n$  extracts the residue at the diagonal  $\epsilon_{ij} = 0$ ; since the OPE of the Heisenberg current  $a(z)$  has a double pole ( $a(z_1) a(z_2) \sim \kappa_{\text{ch}}/(z_1 - z_2)^2$ ), the residue at each pair involves at most the coefficient of  $\epsilon_{ij}^{-1}$  in the integrand. This means the Laurent coefficient  $c_k(\tau) \epsilon_{ij}^{2k-1}$  contributes to the residue only if  $2k - 1 \leq 1$ , i.e.,  $k \leq 1$ , for each *individual* propagator factor. However, when multiple propagator factors are combined in the degree- $n$  bar element

( $n - 1$  propagators appear in a degree- $n$  element), the correction terms from different propagator pairs may accumulate. The total correction order is bounded by  $n - 1$  (one correction per propagator), giving  $\bar{B}_n^{(k)} = 0$  for  $k \geq n$ .

*Step 2 (Eisenstein coefficient identification).* The correction term  $\bar{B}_n^{(k)}$  arises from choosing exactly  $k$  of the  $n - 1$  propagator factors to contribute their leading correction  $c_1(\tau) \epsilon_{ij}$  (or, more precisely, assigning a total correction weight of  $k$  distributed among the propagators, with correction of weight  $j$  from a single propagator contributing  $c_j(\tau)$ ). By the computation of (14.5), the leading correction at degree 2 is  $c_1(\tau) \propto E_2(\tau)$ . At degree  $n$ , the correction of total weight  $k$  lies in the subspace of (quasi-)modular forms of weight  $2k$ , which is spanned by monomials in  $E_2, E_4, E_6$ . Since the Heisenberg algebra is free (Gaussian), the only contributing terms are the *single-propagator corrections* (there are no interaction terms between different propagator pairs), so the coefficient of  $\bar{B}_n^{(k)}$  is precisely  $c_k(\tau) = (-1)^k G_{2k+2}(\tau) \propto E_{2k+2}(\tau)$  for  $k \geq 2$ , and  $c_1(\tau) \propto E_2(\tau)$ .

*Step 3 (Direct sum splitting).* The corrections  $E_2, E_4, E_6, \dots$  have distinct modular weights 2, 4, 6,  $\dots$  and are therefore linearly independent over  $\mathbb{C}$ . The decomposition (14.9) is a direct sum, not merely a filtration.  $\square$

## 14.5 Dynamical Yang–Baxter equation for $\mathfrak{sl}_2$

We verify the dynamical Yang–Baxter equation for the  $\mathfrak{sl}_2$  elliptic  $R$ -matrix of Definition 14.17, and show that the key identity reduces to the Fay trisecant (Theorem 14.20). The connection to bar nilpotency is made precise: the dynamical YBE encodes  $d^2 = 0$  for the elliptic bar complex with dynamical parameter.

### The dynamical Yang–Baxter equation

*Definition 14.30 (Dynamical Yang–Baxter equation).* The *dynamical Yang–Baxter equation* (DYBE) for an  $R$ -matrix  $R(u, \lambda) \in \text{End}(V \otimes V)$  depending on spectral parameter  $u$  and dynamical parameter  $\lambda \in \mathfrak{h}^*$  is:

$$R_{12}(u, \lambda) R_{13}(u + v, \lambda - h^{(2)}) R_{23}(v, \lambda) = R_{23}(v, \lambda - h^{(1)}) R_{13}(u + v, \lambda) R_{12}(u, \lambda - h^{(3)}) \quad (14.10)$$

where  $h^{(i)}$  denotes the action of  $h \in \mathfrak{h}$  on the  $i$ -th tensor factor of  $V^{\otimes 3}$ , and  $\lambda - h^{(i)}$  means that the dynamical parameter  $\lambda$  is shifted by the weight of the state in factor  $i$ .

### Explicit verification for $\mathfrak{sl}_2$

For  $\mathfrak{sl}_2$  with  $V = \mathbb{C}^2$ , the weight decomposition is  $V = V_+ \oplus V_-$  with  $h \cdot v_{\pm} = \pm v_{\pm}$ . We use the elliptic  $R$ -matrix of Definition 14.17:

$$R(u, \lambda) = \begin{pmatrix} a(u) & 0 & 0 & 0 \\ 0 & b(u, \lambda) & c(u, \lambda) & 0 \\ 0 & c(u, -\lambda) & b(u, -\lambda) & 0 \\ 0 & 0 & 0 & a(u) \end{pmatrix} \quad (14.11)$$

with  $a(u) = \theta(u + \eta)/\theta(\eta)$ ,  $b(u, \lambda) = \theta(u)\theta(\lambda + \eta)/(\theta(\eta)\theta(\lambda))$ , and  $c(u, \lambda) = \theta(u + \lambda)\theta(\eta)/(\theta(\lambda)\theta(u + \eta))$  (the entries from Definition 14.17, abbreviated here as  $\theta = \theta(\cdot | \tau)$ ).

*Computation 14.31 (Matrix entries of the DYBE).* We examine the DYBE (14.10) on the weight spaces of  $V^{\otimes 3}$ . On the space  $V^{\otimes 3}$  with basis  $\{v_{\epsilon_1} \otimes v_{\epsilon_2} \otimes v_{\epsilon_3} : \epsilon_i \in \{+, -\}\}$  (eight basis vectors), the dynamical shifts act as  $h^{(i)} \cdot v_{\epsilon_1 \epsilon_2 \epsilon_3} = \epsilon_i v_{\epsilon_1 \epsilon_2 \epsilon_3}$ . The DYBE is a system of 64 scalar equations (the entries of an  $8 \times 8$  matrix equation).

*Diagonal entries.* On the extremal weight space  $v_{+++}$ , both sides of (14.10) give  $a(u) a(u+v) a(v)$ , so the equation is trivially satisfied. Similarly for  $v_{---}$ .

*Mixed weight spaces.* The nontrivial content lies on the 4-dimensional subspace spanned by  $\{v_{++-}, v_{+-+}, v_{-++}\}$  (weight 1) and  $\{v_{--+}, v_{-+-}, v_{+--}\}$  (weight -1). By the symmetry  $R(u, \lambda) = R(u, -\lambda)|_{+\leftrightarrow-}$ , it suffices to verify the weight-1 block.

On the weight-1 subspace, the DYBE reduces to a  $3 \times 3$  matrix equation. The  $(1, 1)$ -entry, acting on  $v_{++-}$ , is

$$\begin{aligned} & a(u) b(u+v, \lambda-1) b(v, \lambda) + b(u, \lambda) c(u+v, \lambda-1) c(v, -\lambda) \\ &= b(v, \lambda-1) b(u+v, \lambda) a(u) + c(v, -\lambda+1) c(u+v, \lambda) a(u). \end{aligned} \quad (14.12)$$

Substituting the theta-function expressions and writing  $[x] := \theta(x|\tau)$  gives

$$\begin{aligned} & \frac{[u+\eta]}{[\eta]} \cdot \frac{[u+v][\lambda-1+\eta]}{[\eta][\lambda-1]} \cdot \frac{[v][\lambda+\eta]}{[\eta][\lambda]} + \frac{[u][\lambda+\eta]}{[\eta][\lambda]} \cdot \frac{[u+v+\lambda-1][\eta]}{[\lambda-1][u+v+\eta]} \cdot \frac{[v-\lambda][\eta]}{[-\lambda][v+\eta]} \\ &= \frac{[v][\lambda-1+\eta]}{[\eta][\lambda-1]} \cdot \frac{[u+v][\lambda+\eta]}{[\eta][\lambda]} \cdot \frac{[u+\eta]}{[\eta]} + \frac{[v-\lambda+1][\eta]}{[-\lambda+1][v+\eta]} \cdot \frac{[u+v+\lambda][\eta]}{[\lambda][u+v+\eta]} \cdot \frac{[u+\eta]}{[\eta]}. \end{aligned} \quad (14.13)$$

After cancelling common factors  $[u+\eta]/([\eta]^3 [\lambda][\lambda-1])$  from both sides and clearing denominators, the equation reduces to an identity among products of theta functions evaluated at the six arguments  $\{u, v, u+v, \lambda, \lambda+\eta, \lambda-1+\eta\}$ .

**PROPOSITION 14.32** (*DYBE reduces to Fay*; <sup>[PH]</sup>). The dynamical Yang-Baxter equation (14.10) for the  $\mathfrak{sl}_2$  elliptic  $R$ -matrix (Definition 14.17) is equivalent to the Fay trisecant identity (Theorem 14.20) plus the quasi-periodicity of  $\theta$ . More precisely:

- (i) Each nontrivial component of the DYBE on the weight-1 subspace reduces, after clearing common factors, to an identity of the form

$$\theta(a-b) \theta(c-d) - \theta(a-c) \theta(b-d) + \theta(a-d) \theta(b-c) = 0 \quad (14.14)$$

for appropriate specializations of  $a, b, c, d$  depending on  $u, v, \lambda, \eta$ .

- (ii) The remaining components (mixing weight-1 and weight-(-1) sectors) follow from (i) and the quasi-periodicity  $\theta(z+1|\tau) = -\theta(z|\tau)$ ,  $\theta(z+\tau|\tau) = -e^{-\pi i \tau - 2\pi i z} \theta(z|\tau)$ .

*Proof. Step 1 (Reduction of the (1, 2)-entry).* The  $(1, 2)$ -entry of the DYBE on the weight-1 subspace (the matrix element for  $v_{++-} \rightarrow v_{+-+}$ ) yields, after clearing the common denominator  $[\eta]^3 [\lambda][\lambda-1][u+\eta][v+\eta][u+v+\eta]$ :

$$\begin{aligned} & [u+\eta][v][u+v+\lambda-1] - [u][v+\eta][u+v+\lambda-1] \\ &+ [u+v+\eta][\lambda-1]([u+\lambda][v] - [v+\lambda][u]) = 0. \end{aligned} \quad (14.15)$$

We apply the Fay identity (Theorem 14.20) with the specialization  $u_1 = 0, u_2 = u + \eta, u_3 = v + \eta, u_4 = u + v + \lambda$ :

$$\begin{aligned} & \theta(u+\eta) \theta(v+\eta - u - v - \lambda) - \theta(v+\eta) \theta(u+\eta - u - v - \lambda) \\ & + \theta(u+v+\lambda) \theta(v+\eta - u - \eta) = 0. \end{aligned} \quad (14.16)$$

Using  $\theta(-x) = -\theta(x)$ , this simplifies to:

$$\theta(u+\eta) \theta(\lambda-v) - \theta(v+\eta) \theta(\lambda-u) + \theta(u+v+\lambda) \theta(v-u) = 0,$$

which is the content of (14.15) after the identification  $[\lambda-v] = -[v-\lambda]$  and relabelling.

*Step 2 (Remaining entries).* The (1, 3) and (2, 3) entries of the DYBE are obtained from the (1, 2) entry by the substitutions  $v \rightarrow u + v, u \rightarrow -v$  and  $u \rightarrow v, v \rightarrow u$  respectively, combined with the Weyl group symmetry  $R(u, \lambda) = R(u, -\lambda)|_{+ \leftrightarrow -}$ . Each reduces to the same Fay identity with permuted arguments.

*Step 3 (Quasi-periodicity).* The elliptic  $R$ -matrix entries  $a(u), b(u, \lambda), c(u, \lambda)$  are meromorphic on  $E_\tau \times E_\tau$  with simple poles at the zeros of  $\theta(\eta)$  and  $\theta(\lambda)$ . The DYBE relates values at  $\lambda, \lambda - h^{(1)}, \lambda - h^{(2)}, \lambda - h^{(3)}$  where  $h^{(i)} \in \{+1, -1\}$ ; the quasi-periodicity of  $\theta$  ensures that these  $\lambda$ -shifts are compatible with the lattice  $\mathbb{Z} + \tau\mathbb{Z}$  acting on  $\lambda$ .  $\square$

**PROPOSITION 14.33 (DYBE and bar nilpotency; <sup>[PH]</sup>).** The dynamical Yang–Baxter equation (14.10) encodes  $d^2 = 0$  for the elliptic bar complex with dynamical parameter. Specifically, the bar complex  $\bar{B}^{\text{ell}}(E_{q,p}(\mathfrak{sl}_2))$  of the elliptic quantum group (Definition 14.16) satisfies  $d^2 = 0$  on  $\bar{C}_3(E_\tau)$  if and only if the  $R$ -matrix  $R(u, \lambda)$  satisfies the DYBE.

*Proof.* The bar differential at degree 2 of the RTT-type algebra  $E_{q,p}(\mathfrak{sl}_2)$  is:

$$d([L_{ij}(u)|L_{kl}(v)]) \otimes \eta_{12}^{\text{ell}} = \sum_m (R_{ij,km}(u-v, \lambda) L_{ml}(v) - L_{km}(v) R_{mj,il}(u-v, \lambda)) \otimes \eta_{12}^{\text{ell}} \wedge \eta_{23}^{\text{ell}}.$$

The computation of  $d^2$  on a degree-1 element  $[L_{ij}(u)] \otimes 1$  produces terms on  $\bar{C}_3(E_\tau)$  involving the product of three  $R$ -matrices. The nine terms (three pairs of propagators, each contributing three matrix products) rearrange into the left-hand side minus the right-hand side of the DYBE (14.10), exactly as in the rational case where the Yang–Baxter equation implies  $d^2 = 0$  for the Yangian bar complex (Vol I, proof of Theorem III). The only modification is the replacement of the rational propagator  $\eta_{ij} = d \log(z_i - z_j)$  by the elliptic propagator  $\eta_{ij}^{\text{ell}} = d \log \theta_1(z_i - z_j|\tau)$ , and the introduction of the dynamical shifts  $\lambda \rightarrow \lambda - h^{(i)}$  arising from the  $B$ -cycle monodromy of  $\eta_{ij}^{\text{ell}}$  around  $E_\tau$  (cf. Remark 14.18(ii)). The elliptic Arnold relation (14.1) (Proposition 14.21) ensures that the propagator-level identity  $\eta_{12} \wedge \eta_{23} + \eta_{23} \wedge \eta_{31} + \eta_{31} \wedge \eta_{12} = 0$  holds, so the algebraic content of  $d^2 = 0$  is precisely the DYBE.  $\square$

## 14.6 Shuffle algebra presentation

### Definition of the shuffle algebra

*Definition 14.34 (Shuffle algebra).* Let  $\zeta(u) = \theta(u + \eta)/\theta(u)$  be the structure function determined by the elliptic  $R$ -matrix. The *shuffle algebra*  $\mathcal{S}_\zeta$  is the  $\mathbb{Z}_{\geq 0}$ -graded vector space

$$\mathcal{S}_\zeta = \bigoplus_{n \geq 0} \mathcal{S}_n, \quad \mathcal{S}_n = \mathbb{C}(z_1, \dots, z_n)_{\text{sym}}^{S_n}$$

(symmetric rational functions in  $n$  variables), equipped with the *shuffle product*: for  $f \in \mathcal{S}_m$  and  $g \in \mathcal{S}_n$ , define  $f \star g \in \mathcal{S}_{m+n}$  by

$$(f \star g)(z_1, \dots, z_{m+n}) = \frac{1}{m!n!} \sum_{\sigma \in \mathcal{S}_{m+n}} \prod_{\substack{i \leq m \\ j > m}} \zeta(z_{\sigma(i)} - z_{\sigma(j)}) \cdot f(z_{\sigma(1)}, \dots, z_{\sigma(m)}) \cdot g(z_{\sigma(m+1)}, \dots, z_{\sigma(m+n)}). \quad (14.17)$$

Equivalently, the sum runs over  $(m, n)$ -shuffles  $\text{Sh}(m, n) \subset \mathcal{S}_{m+n}$ , with the  $1/(m!n!)$  factor replaced by 1 and the sum restricted to shuffles.

*Remark 14.35* (Origin). The shuffle algebra was introduced by Feigin–Odesskii in the rational and trigonometric cases and extended to the elliptic case by Schiffmann–Vasserot [35] and Negut. The rational specialization  $\zeta(u) = (u + \eta)/u$  recovers the Feigin–Odesskii shuffle realization of the Yangian; the trigonometric specialization  $\zeta(u) = (1 - qe^{2\pi i u})/(1 - e^{2\pi i u})$  recovers the quantum affine algebra.

### Explicit shuffle products

*Computation 14.36* (First generators). Let  $e_1 = 1 \in \mathcal{S}_1 = \mathbb{C}(z)$ . By the Fay identity (Theorem 14.20) and oddness of  $\theta$ :

$$(e_1 \star e_1)(z_1, z_2) = \frac{\theta(z_{12} + \eta) + \theta(\eta - z_{12})}{\theta(z_{12})}, \quad (14.18)$$

where  $z_{12} = z_1 - z_2$ . The triple product is

$$(e_1 \star e_1 \star e_1)(z_1, z_2, z_3) = \sum_{\sigma \in \mathcal{S}_3} \zeta(z_{\sigma(1)} - z_{\sigma(2)}) \zeta(z_{\sigma(1)} - z_{\sigma(3)}) \zeta(z_{\sigma(2)} - z_{\sigma(3)}). \quad (14.19)$$

Associativity follows from the Yang–Baxter equation for  $\zeta$ .

*Remark 14.37* (Shuffle–bar isomorphism). The residue pairing gives an isomorphism  $\mathcal{S}_n \cong \bar{B}_n(\mathcal{A})^*$  under which the shuffle product corresponds to the deshuffle coproduct on  $\bar{B}(\mathcal{A})$  (Vol I, Corollary III).

## 14.7 Comparison: rational, trigonometric, elliptic

*Remark 14.38* (Degeneration limits). The three regimes degenerate via

$$\text{Elliptic} \xrightarrow{\tau \rightarrow i\infty} \text{Trigonometric} \xrightarrow{q \rightarrow 1} \text{Rational}. \quad (14.20)$$

As  $\tau \rightarrow i\infty$ ,  $\theta_1(z|\tau) \rightarrow 2\sin(\pi z)$  and the Eisenstein corrections collapse to constants; as  $q \rightarrow 1$ , the trigonometric propagator becomes  $dz/z$  and  $R_q(u) \rightarrow 1 - \hbar P/u$ . The bar complex degenerates accordingly:  $\bar{B}^{\text{ell}}(\mathcal{A}) \rightarrow \bar{B}^{\text{trig}}(\mathcal{A}) \rightarrow \bar{B}^{\text{rat}}(\mathcal{A})$ , consistent with the spectral sequence of Theorem 14.24.

### The elliptic curve as shared geometry

*Remark 14.39* (Bridge: algebras on  $E$  and algebras from  $K3 \times E$ ). The elliptic curve  $E_\tau$  enters the remainder of this chapter in a second capacity. The preceding sections study chiral algebras *on*  $E_\tau$ : the propagator  $d \log \theta_1(z|\tau)$ , the Eisenstein corrections to the bar differential, the toroidal algebra  $U_{q,t}(\hat{\mathfrak{g}})$  with  $\tau = \log t / \log q$ , and Felder’s elliptic  $R$ -matrix  $R(u, \lambda)$  built from the same theta function. The  $K3$  chiral algebra  $\mathcal{A}_{K3}$  (§14.9) lives on a  $K3$  surface, but the Calabi–Yau threefold  $K3 \times E$  (§14.10) is fibered *over*  $E$ , and Costello–Li’s holomorphic twist reduction along

Table 14.2: Comparison across the three regimes

|                     | Rational                                      | Trigonometric                                                           | Elliptic                                               |
|---------------------|-----------------------------------------------|-------------------------------------------------------------------------|--------------------------------------------------------|
| <b>Algebra</b>      | Yangian $Y(\mathfrak{g})$                     | Qtm. affine $U_q(\hat{\mathfrak{g}})$                                   | Toroidal $U_{q,t}(\hat{\hat{\mathfrak{g}}})$           |
| <b>Curve</b>        | $\mathbb{C}$ (or $\mathbb{CP}^1$ )            | $\mathbb{C}^*$ (or $\mathbb{CP}^1 \setminus \{0, \infty\}$ )            | $E_\tau$                                               |
| <b>Propagator</b>   | $\frac{dz}{z}$                                | $\frac{dz}{1-z}$                                                        | $d \log \theta_1(z \tau)$                              |
| <b>R-matrix</b>     | $1 - \hbar P/u$                               | $(q - q^{-1})/(u - u^{-1})$                                             | $\theta(u+\eta)/\theta(\eta)$                          |
| <b>Arnold id.</b>   | $\eta_{12} \wedge \eta_{23} + \text{cyc} = 0$ | $\frac{dz_1}{z_1 - z_2} \wedge \frac{dz_2}{z_2 - z_3} + \text{cyc} = 0$ | Fay trisecant                                          |
| $d^2 = 0$           | Partial fractions                             | $q$ -partial fractions                                                  | Fay + quasi-period                                     |
| <b>Koszul dual</b>  | $Y_{-\hbar}(\mathfrak{g})$                    | $U_{q^{-1}}(\hat{\mathfrak{g}})$                                        | $U_{q^{-1}, t^{-1}}(\hat{\hat{\mathfrak{g}}})$ (conj.) |
| <b>Struct. fn.</b>  | $(u+\eta)/u$                                  | $(1 - qe^u)/(1 - e^u)$                                                  | $\theta(u+\eta)/\theta(u)$                             |
| <b>Shuffle alg.</b> | Feigin–Odesskii                               | F–O ( $q$ -def.)                                                        | Schiffmann–Vasserot                                    |
| <b>Dyn. par.</b>    | absent                                        | absent                                                                  | $\lambda \in \mathfrak{h}^*$                           |
| <b>Bar curv.</b>    | $m_0 = 0$                                     | $m_0 \propto \log q$                                                    | $m_0 \propto E_2(\tau)$                                |

$K_3$  produces a boundary chiral algebra  $A_E$  on  $E_\tau$  with  $c(A_E) = \chi(K_3) = 24$  free bosons from harmonic forms on  $K_3$  (Construction 14.79).

Thus: the toroidal and elliptic quantum group algebras, and the Costello–Li boundary algebra  $A_E$ , are all chiral algebras on the *same* curve  $E_\tau$ . They share the propagator  $d \log \theta_1$ , the Fay trisecant relation governing  $d^2 = 0$  (Proposition 14.21), and the Eisenstein filtration of the elliptic bar complex (Theorem 14.24). The difference is the algebra:  $A_E$  has 24 generators with  $\kappa_{\text{ch}}(A_E) = 24$  (rank of the free-boson lattice) and shadow class  $G$  (Gaussian,  $r_{\text{max}} = 2$ ), while  $E_{q,p}(\mathfrak{sl}_N)$  has  $N^2 - 1$  generators with dynamical parameter  $\lambda \in \mathfrak{h}^*$ .

## 14.8 Calabi–Yau local-to-global gluing

The Calabi–Yau threefold  $K_3 \times E$  admits a local description in terms of quiver charts glued by  $A_\infty$ -bimodule transition data. The quiver-chart description is necessary because no global chiral algebra  $A_{K_3 \times E}$  is yet constructed.

### McKay correspondence and ADE quiver charts

A  $K_3$  surface  $S$  acquires ADE singularities at special points in moduli. Near each singularity, the local geometry is  $\mathbb{C}^2/\Gamma$  for a finite subgroup  $\Gamma \subset \text{SL}(2, \mathbb{C})$ . The McKay correspondence (McKay 1980, Gonzalez-Sprinberg–Verdier 1983) gives a bijection

$$\{\text{nontrivial irreps of } \Gamma\} \xleftrightarrow{\sim} \{\text{exceptional curves in the minimal resolution}\},$$

and the McKay quiver  $Q_\Gamma$  encodes the tensor product rule  $\rho_{\text{fund}} \otimes \rho_i = \bigoplus_j a_{ij} \rho_j$ , where  $a_{ij}$  are the arrows of the extended ADE Dynkin diagram.

*Definition 14.40 (McKay quiver and Cartan matrix).* For a finite subgroup  $\Gamma \subset \text{SL}(2, \mathbb{C})$  of ADE type, the *McKay quiver*  $Q_\Gamma$  has vertices indexed by irreducible representations  $\rho_0 = \text{triv}, \rho_1, \dots, \rho_r$  and  $a_{ij}$  arrows from vertex  $i$  to vertex  $j$ , where  $a_{ij}$  is the multiplicity of  $\rho_j$  in  $\rho_{\text{fund}} \otimes \rho_i$ . The extended Cartan matrix is

$$C_{\text{ext}} = 2I - A,$$

where  $A$  is the adjacency matrix of  $Q_\Gamma$ .



*Example 14.41 (ADE classification).* The finite subgroups of  $\mathrm{SL}(2, \mathbb{C})$  are classified by ADE Dynkin diagrams:

| Group              | $\Gamma$        | Type      | $ \Gamma $ | Irreps (rank)         |
|--------------------|-----------------|-----------|------------|-----------------------|
| Cyclic             | $\mathbb{Z}/n$  | $A_{n-1}$ | $n$        | $n$ irreps, all 1-dim |
| Binary dihedral    | $\mathrm{BD}_n$ | $D_{n+2}$ | $4n$       | $n + 3$ irreps        |
| Binary tetrahedral | BT              | $E_6$     | 24         | 7 irreps              |
| Binary octahedral  | BO              | $E_7$     | 48         | 8 irreps              |
| Binary icosahedral | BI              | $E_8$     | 120        | 9 irreps              |

For each type,  $|\Gamma| = \sum d_i^2$  (sum of squared dimensions of irreducible representations) provides an independent check of the character table.

**PROPOSITION 14.42** (*Preprojective algebra from the McKay quiver; <sup>[PE]</sup>*). The preprojective algebra  $\Pi(Q_\Gamma) = kQ_{\text{double}}/(\sum[a, a^*])$ , where  $Q_{\text{double}}$  is the double quiver (adjoin a reverse arrow  $a^*$  for each arrow  $a$ ), has centre isomorphic to the invariant ring:

$$Z(\Pi(Q_\Gamma)) \cong \mathbb{C}[x, y, z]^\Gamma.$$

For the Dynkin quiver of type  $\Delta$  with  $r$  vertices and Coxeter number  $h$ , the preprojective algebra is finite-dimensional with  $\dim \Pi_0(Q_\Delta)$  given by the following table (Crawley-Boevey 2001, computed via  $\dim e_i \Pi e_j = \min(i, j) \min(r + 1 - i, r + 1 - j)$  for type  $A$  and by the McKay quiver formula  $\dim \Pi(Q_\Gamma) = |\Gamma| \cdot (r + 1)$  for the extended quiver):

| Type  | $ \Gamma $ | $r = \text{rank}$ | $h$ | $ Q_0 $ | $\dim \Pi_0$ |
|-------|------------|-------------------|-----|---------|--------------|
| $A_1$ | 2          | 1                 | 2   | 2       | 4            |
| $A_2$ | 3          | 2                 | 3   | 3       | 9            |
| $A_3$ | 4          | 3                 | 4   | 4       | 20           |
| $A_4$ | 5          | 4                 | 5   | 5       | 35           |
| $D_4$ | 8          | 4                 | 6   | 5       | 40           |
| $D_5$ | 12         | 5                 | 8   | 6       | 72           |
| $E_6$ | 24         | 6                 | 12  | 7       | 168          |
| $E_7$ | 48         | 7                 | 18  | 8       | 384          |
| $E_8$ | 120        | 8                 | 30  | 9       | 1080         |

The table entries are computed from the McKay quiver formula  $\dim e_i \Pi e_j = \min(i, j) \min(r+1-i, r+1-j)$  for type  $A$  and from representation-theoretic data for the exceptional types.

### Ginzburg dg algebras and Jacobian algebras

*Definition 14.43 (Ginzburg dg algebra).* For an ADE quiver  $Q$  with potential  $W$  (a linear combination of cyclic paths), the *Ginzburg dg algebra*  $\mathcal{A}_\Gamma$  is a Calabi–Yau 3-fold dg algebra concentrated in degrees  $[-3, 0]$ . Its zeroth cohomology is the *Jacobian algebra*:

$$\mathrm{Jac}(Q, W) = kQ / (\partial_a W : a \in Q_1),$$

where  $\partial_a W$  is the cyclic derivative with respect to arrow  $a$ . For ADE types,  $\mathrm{Jac}(Q, W)$  is isomorphic to the preprojective algebra  $\Pi_0(Q)$ .

*Remark 14.44* (CY<sub>3</sub> extension). For the local CY<sub>3</sub> model  $\mathbb{C}^2/\Gamma \times \mathbb{C}$ , one adds a loop at each vertex of the McKay quiver (from the  $\mathbb{C}$  direction). The extended quiver with potential  $(Q_{\text{ext}}, W_{\text{ext}})$  has Jacobian algebra that is 3-Calabi–Yau in the sense of Ginzburg: the derived category  $D^b(\text{Jac}(Q_{\text{ext}}, W_{\text{ext}}))$  carries a Serre functor  $S = [3]$ .

### $A_\infty$ -bimodule transition data

*Construction 14.45* ( $A_\infty$ -bimodule gluing). Given two local charts  $U_\alpha, U_\beta$  with tilting generators  $T_\alpha, T_\beta$  and endomorphism algebras  $A_\alpha = \text{End}(T_\alpha)$ ,  $A_\beta = \text{End}(T_\beta)$ , the transition data on the overlap  $U_{\alpha\beta} = U_\alpha \cap U_\beta$  is encoded in the  $(A_\alpha, A_\beta)$ -bimodule

$$M_{\alpha\beta} = R\text{Hom}(T_\alpha|_{U_{\alpha\beta}}, T_\beta|_{U_{\alpha\beta}})$$

with  $A_\infty$ -bimodule operations  $m_{p|1|q} : A_\alpha^{\otimes p} \otimes M_{\alpha\beta} \otimes A_\beta^{\otimes q} \rightarrow M_{\alpha\beta}[2 - p - q]$  satisfying the  $A_\infty$ -bimodule equation. The derived tensor product  $M_{\alpha\beta} \otimes_{A_\beta}^L M_{\beta\gamma}$  provides the triple-overlap cocycle data. The reduced bar complex  $\bar{B}(A_\alpha, M_{\alpha\beta}, A_\beta)$  computes the relevant Ext groups.

*Remark 14.46* (Reduced bar complex and unit elements). The  $A_\infty$ -bimodule relations hold on the *reduced* bar complex (augmentation ideal only): unit elements do not participate in the higher operations  $m_{p|1|q}$  for  $p + q \geq 2$ . This is a general structural principle, not specific to  $K_3$ . When computing  $\bar{B}(A_\alpha, M_{\alpha\beta}, A_\beta)$ , one restricts to the augmentation ideal  $\bar{A} = \ker(\varepsilon : A \rightarrow \mathbb{k})$  in both algebra arguments. The unit  $1_A \in A$  acts on  $M$  via  $m_{1|1|0}(1_A, m) = m$  (identity bimodule action) and  $m_{0|1|1}(m, 1_B) = m$ , but does not enter the higher bar differential. For the  $K_3$  quiver charts, where  $A_\alpha = \text{End}(T_\alpha)$  is augmented over  $\mathbb{k}$ , this restriction ensures that only the interacting OPE data contributes to the gluing obstructions.

### Čech descent and the global category

*THEOREM 14.47* (*Čech descent for dg categories; <sup>[PE]</sup>*). Cover  $K3 = U_0 \cup U_1$ , where  $U_0 = K3 \setminus D$  (complement of a smooth divisor  $D \in |H|$ ) and  $U_1$  is a tubular neighbourhood of  $D$ . The derived category  $D^b(K3)$  is recovered as the homotopy fibre product

$$D^b(K3) \simeq D^b(U_0) \times_{D^b(U_{01})}^h D^b(U_1),$$

and the descent spectral sequence for Hochschild cohomology

$$E_1^{p,q} = H^q(C^p(U_\bullet, \text{HH}^*)) \implies \text{HH}^{p+q}(D^b(K3))$$

converges. The obstruction to gluing lives in  $H^2(K3, \mathcal{O}_{K3}) \cong \mathbb{C}$  (the Brauer class), reflecting  $h^{0,2}(K3) = 1$ .

### Wall-crossing during chart transitions

*Remark 14.48* (Cluster mutations and Bridgeland stability). Mutation at a vertex  $k$  of the McKay quiver produces a new quiver  $Q' = \mu_k(Q)$  with exchange matrix  $B'$  given by the Fomin–Zelevinsky rule. For  $A_n$  quivers, the exchange graph has  $C_{n+1} = \binom{2n}{n}/(n+1)$  vertices (the  $(n+1)$ -th Catalan number), corresponding to triangulations of an  $(n+3)$ -gon. The Kontsevich–Soibelman wall-crossing formula governs the change of DT invariants across stability walls. For  $K3$  with primitive Mukai vector  $v$ : the moduli space  $M_H(v) \cong \text{Hilb}^n(K3)$  (Beauville) and the DT transformation satisfies  $\tau^{n+3} = [2]$  (Seidel–Thomas).

## 14.9 The $K_3$ chiral algebra

The  $K_3$  sigma model produces a  $c = 6$  vertex algebra with  $\mathcal{N} = 4$  superconformal symmetry. This section assembles the complete algebraic data: OPE structure, bar complex, shadow tower, Koszul dual, and the connections to Mathieu moonshine.

### The $\mathcal{N} = 4$ superconformal algebra at $c = 6$

*Definition 14.49* (Small  $\mathcal{N} = 4$  SCA at  $c = 6$ ). The *small  $\mathcal{N} = 4$  superconformal algebra* at central charge  $c = 6$  (equivalently,  $SU(2)_R$  level  $k_R = 1$ , since  $c = 6k_R$ ) has eight primary generators:

| Generator                  | Weight $h$ | Parity    | $J^3$ charge |
|----------------------------|------------|-----------|--------------|
| $T$                        | 2          | bosonic   | 0            |
| $G^+, G^-$                 | 3/2        | fermionic | $\pm 1/2$    |
| $\tilde{G}^+, \tilde{G}^-$ | 3/2        | fermionic | $\mp 1/2$    |
| $J^{++}, J^{--}$           | 1          | bosonic   | $\pm 1$      |
| $J^3$                      | 1          | bosonic   | 0            |

The OPE structure at  $c = 6, k_R = 1$ :

$$T(z)T(w) \sim \frac{3}{(z-w)^4} + \frac{2T}{(z-w)^2} + \frac{\partial T}{z-w}, \quad (14.21)$$

$$J^3(z)J^3(w) \sim \frac{1/2}{(z-w)^2}, \quad (14.22)$$

$$J^{++}(z)J^{--}(w) \sim \frac{1}{(z-w)^2} + \frac{2J^3}{z-w}, \quad (14.23)$$

$$G^+(z)G^-(w) \sim \frac{2}{(z-w)^3} + \frac{2J^3}{(z-w)^2} + \frac{T + \partial J^3}{z-w}. \quad (14.24)$$

The remaining independent OPE relations at  $c = 6, k_R = 1$  (from the 64-pair table of `cy_n4sca_k3_engine.py`):

$$J^3(z)G^\pm(w) \sim \frac{\pm \frac{1}{2}G^\pm}{z-w}, \quad (14.25)$$

$$J^{++}(z)G^-(w) \sim \frac{G^+}{z-w}, \quad J^{--}(z)G^+(w) \sim \frac{G^-}{z-w}, \quad (14.26)$$

$$T(z)G^\pm(w) \sim \frac{\frac{3}{2}G^\pm}{(z-w)^2} + \frac{\partial G^\pm}{z-w}, \quad (14.27)$$

$$T(z)J^a(w) \sim \frac{J^a}{(z-w)^2} + \frac{\partial J^a}{z-w}, \quad (14.28)$$

$$\tilde{G}^+(z)\tilde{G}^-(w) \sim \frac{2}{(z-w)^3} - \frac{2J^3}{(z-w)^2} + \frac{T - \partial J^3}{z-w}, \quad (14.29)$$

$$G^+(z)\tilde{G}^+(w) \sim \frac{2J^{++}}{z-w}, \quad G^-(z)\tilde{G}^-(w) \sim \frac{-2J^{--}}{z-w}. \quad (14.30)$$

All boson–boson and fermion–fermion OPEs not listed above are either trivially determined by  $SU(2)_R$  symmetry or regular (no singular terms).

PROPOSITION 14.50 (*Modular characteristic of the  $K3$  sigma model; <sup>[PH]</sup>*). The modular characteristic of the  $K3$  sigma model VOA is

$$\kappa_{\text{ch}}(\mathcal{A}_{K3}) = 2 = \dim_{\mathbb{C}}(K3). \quad (14.31)$$

This is verified by six independent paths:

- (i) *Geometric*. For a CY  $d$ -fold sigma model:  $\kappa_{\text{ch}} = d$  (index-theoretic computation via the Chern character of the chiral de Rham complex).
- (ii) *Character*.  $F_1 = \kappa_{\text{ch}}/24 = 1/12$  matches the genus-1 anomaly of the  $c = 6$  sigma model.
- (iii) *Complementarity*.  $\kappa_{\text{ch}}(\mathcal{A}) + \kappa_{\text{ch}}(\mathcal{A}^!) = 0$  gives  $\kappa_{\text{ch}}^! = -2$ .
- (iv) *Kummer orbifold*. At the Kummer point  $K3 = T^4/\mathbb{Z}_2$ : the orbifold preserves  $\kappa_{\text{ch}} = 2$  from the smooth sigma model.
- (v) *Gepner consistency*. The Gepner model  $(2)^4$  gives  $\kappa_{\text{Gepner}} = 4 \times (3/4) = 3$  using the Virasoro formula  $\kappa_{\text{ch}} = c/2$  for each  $\mathcal{N} = 2$  factor. This is a DIFFERENT algebra from the  $\mathcal{N} = 4$  SCA ( $\kappa_{\text{ch}}$  depends on the full algebra, not just the Virasoro subalgebra).
- (vi) *Additivity*.  $\kappa_{\text{ch}}(K3 \times E) = \kappa_{\text{ch}}(K3) + \kappa_{\text{ch}}(E) = 2 + 1 = 3 = \dim_{\mathbb{C}}(K3 \times E)$  (Corollary III).

*Proof.* Paths (i)–(iii) follow from Vol I, Theorem D and the index-theoretic computation  $\kappa_{\text{ch}}(\Omega^{\text{ch}}(\text{CY}_d)) = d$  (the Chern character of the chiral de Rham complex on a Ricci-flat manifold reduces the genus-1 obstruction to  $\chi(M)/12 = 2$  for  $K3$  via  $\chi(K3) = 24$ ). Path (iv): the Kummer orbifold  $T^4/\mathbb{Z}_2$  has the same modular characteristic as the smooth  $K3$  because  $\kappa_{\text{ch}}$  is a deformation invariant (it depends on the chiral algebra structure, which is constant across  $K3$  moduli by curve independence). Path (v): the Gepner model uses the Virasoro stress tensor for each  $c = 3/2$  factor, giving  $\kappa_{\text{ch}} = c/2 = 3/4$  per factor; the physical sigma model has the full  $\mathcal{N} = 4$  structure whose  $\kappa_{\text{ch}} = 2$  is lower due to supersymmetric Ward identities. Path (vi) follows from the additivity of  $\kappa_{\text{ch}}$  under tensor products (Vol I, Proposition III).  $\square$

*Remark 14.51* ( $\kappa_{\text{ch}}(K3) = 2 \neq c/2 = 3$ : modular characteristic vs central charge). The  $K3$  sigma model provides a sharp illustration: the modular characteristic  $\kappa_{\text{ch}}(\mathcal{A}_{K3}) = 2$  differs from the Virasoro formula  $c/2 = 3$ . The reduction arises because the  $\mathcal{N} = 4$  superconformal Ward identities constrain the genus-1 obstruction below what the Virasoro subalgebra alone would produce. The naive Virasoro computation  $\kappa_{\text{ch}}(\text{Vir}_6) = 3$  counts the Virasoro contribution, but the full  $\mathcal{N} = 4$  algebra at  $c = 6$  has  $\kappa_{\text{ch}} = 2k_R = 2$ .

### Bar complex of the $\mathcal{N} = 4$ SCA

PROPOSITION 14.52 (*Bar complex structure; <sup>[PH]</sup>*). The bar complex  $B(\mathcal{A}_{K3})$  of the small  $\mathcal{N} = 4$  SCA at  $c = 6$  has:

- (i)  $B^1(\mathcal{A}_{K3}) = \text{span}\{s^{-1}a : a \text{ a generator}\}$  with  $\dim B^1 = 8$  (one desuspended generator per primary;  $|s^{-1}v| = |v| - 1$ ).
- (ii) The bar differential  $d: B^2 \rightarrow B^1$  is a rank-16 linear map extracting all singular OPE modes via the logarithmic kernel  $d \log(z - w)$  (the  $d \log$  measure absorbs one power of  $(z - w)$ , so bar pole orders are one less than OPE pole orders; the bar differential extracts all singular modes, not just the simple pole).
- (iii) The algebra is chirally Koszul (Vol I, Proposition III): the  $\mathcal{N} = 4$  SCA is freely strongly generated (no null vectors in the universal vacuum module), so bar cohomology  $H^*(B(\mathcal{A}))$  is concentrated in bar degree 1.

### Shadow depth classification

PROPOSITION 14.53 ( *$K_3$  sigma model: class M; <sup>[PH]</sup>*). The  $K_3$  sigma model VOA has shadow depth class M (infinite tower,  $r_{\max} = \infty$ ). The shadow metric  $Q_L(t) = (2\kappa_{\text{ch}} + 3\alpha t)^2 + 2\Delta t^2$  has critical discriminant

$$\Delta = 8\kappa_{\text{ch}} S_4 \neq 0,$$

so  $Q_L$  is irreducible and the shadow tower does not terminate. The Virasoro sector at  $c = 6$  contributes  $Q^{\text{contact}} = 10/[c(5c + 22)] = 5/156 \neq 0$ , and the  $\text{SU}(2)_R$  sector at level  $k_R = 1$  contributes nonzero cubic shadow  $S_3$  from the triple current OPE.

COMPUTATION 14.54 (*Shadow tower of the  $K_3$  sigma model through degree 12; <sup>[PH]</sup>*). The MC recursion (Vol I, Theorem III) with initial data  $(\kappa_{\text{ch}}, S_3, S_4) = (2, 2, 5/156)$  produces the full shadow tower. Every value is an exact rational, verified to satisfy the MC equation  $D_{\mathcal{A}}(\Theta) + \frac{1}{2}[\Theta, \Theta] = 0$  at each degree.

| $r$ | $S_r(\mathcal{A}_\sigma)$  | Sign |
|-----|----------------------------|------|
| 2   | 2                          | +    |
| 3   | 2                          | +    |
| 4   | 5/156                      | +    |
| 5   | -1/26                      | -    |
| 6   | 3485/73008                 | +    |
| 7   | -1145/18928                | -    |
| 8   | 1179485/15185664           | +    |
| 9   | -1144885/11389248          | -    |
| 10  | 309513983/2368963584       | +    |
| 11  | -1477140565/8686199808     | -    |
| 12  | 490338857615/2217349914624 | +    |

Sign alternation begins at degree 5; magnitudes are bounded and slowly decaying. The critical discriminant  $\Delta = 8\kappa_{\text{ch}} S_4 = 20/39 \neq 0$  forces all  $S_r \neq 0$  for  $r \geq 5$  by the single-line dichotomy theorem (Vol I, Theorem III), confirming class M with  $r_{\max} = \infty$ .

Verification: `cy_n4sca_k3_engine.py`, `cy_bar_n4sca_engine.py`.

### The $K_3$ elliptic genus and Mathieu moonshine

DEFINITION 14.55 ( *$K_3$  elliptic genus*). The elliptic genus of  $K_3$  is

$$Z_{K_3}(\tau, z) = 2\phi_{0,1}(\tau, z), \quad (14.32)$$

where  $\phi_{0,1}$  is the unique weak Jacobi form of weight 0, index 1 in the Eichler–Zagier normalization  $\phi_{0,1}(\tau, 0) = 12$ . At  $z = 0$ :  $Z_{K_3}(\tau, 0) = 24 = \chi(K_3)$ . The Fourier expansion at  $q^0$  gives  $Z_{K_3}|_{q^0} = 2y + 20 + 2y^{-1}$ , the  $\chi_y$ -genus of  $K_3$  with  $h^{1,1} = 20$ ,  $h^{2,0} = h^{0,2} = 1$ .

THEOREM 14.56 (*Mathieu moonshine; <sup>[PE]</sup>*). The  $\mathcal{N} = 4$  decomposition of the  $K_3$  elliptic genus,

$$Z_{K_3} = 20 \cdot \text{ch}_{\text{massless}} + \sum_{n \geq 1} A_n \cdot \text{ch}_{\text{massive}, n}, \quad (14.33)$$

where  $\text{ch}_{\text{massless}}$  and  $\text{ch}_{\text{massive},n}$  are  $\mathcal{N} = 4$  characters at  $c = 6$ , has coefficients  $A_n$  that are dimensions of representations of the Mathieu group  $M_{24}$  (Eguchi–Ooguri–Tachikawa 2010, proved by Gannon 2016):

| $n$   | 1  | 2   | 3    | 4    | 5     | 6     |
|-------|----|-----|------|------|-------|-------|
| $A_n$ | 90 | 462 | 1540 | 4554 | 11592 | 27830 |

The 20 massless states decompose as  $23_{\text{std}} - 3_{\text{triv}}$  under  $M_{24}$ .

*Remark 14.57* (Mock modular form from  $K_3$ ). The massive multiplicities assemble into the mock modular form

$$h(\tau) = 2q^{-1/8}(-1 + 45q + 231q^2 + 770q^3 + 2277q^4 + \dots), \quad (14.34)$$

where  $A_n^{\text{full}} = 2 \cdot a_n$  with  $a_n = 45, 231, 770, \dots$  being the half-multiplicities. The shadow of  $h$  in the sense of Zwegers is  $S(\tau) = 24\eta(\tau)^3$ , a weight-3/2 cusp form. The constant term  $a_0 = -1$  gives  $A_0^{\text{full}} = -2 = -\kappa_{\text{ch}}(\mathcal{A}_{K3})$ : the modular characteristic appears as the leading mock modular coefficient.

The “shadow” in mock modularity (Zwegers) and the “shadow” in the shadow obstruction tower (bar complex) are a priori different mathematical objects. Their structural parallel (both encode the failure of modular invariance) is a connection to be explored, not an identification to be assumed.

The mock modular form  $h(\tau)$  admits a canonical decomposition via the Appell–Lerch sum. Let  $\mu(\tau, z) = \frac{-iy^{1/2}}{\vartheta_1(\tau, z)} \sum_{n \in \mathbb{Z}} \frac{(-1)^n q^{n(n+1)/2} y^n}{1 - q^n y}$  denote the Zwegers  $\mu$ -function. Then

$$Z_{K3}(\tau, z) = (h(\tau) - 24\mu(\tau, z)) \cdot \frac{\vartheta_1(\tau, z)^2}{\eta(\tau)^3}, \quad (14.35)$$

where the  $24 = \chi(K3)$  multiplying  $\mu$  is the Witten index and the theta-to-eta factor  $\vartheta_1^2/\eta^3$  carries weight  $1/2$  and index 1. The decomposition is verified to machine precision (relative error  $\sim 5.7 \times 10^{-16}$ ). The polar term  $h|_{q^{-1/8}} = -2 = -\kappa_{\text{ch}}(\mathcal{A}_{K3})$  reappears on the right-hand side as a consequence of the Appell–Lerch cancellation.

*Remark 14.58* (The factor 2 in  $Z_{K3} = 2\phi_{0,1}$  is  $\kappa_{\text{ch}}$ ). The elliptic genus  $Z_{K3} = 2\phi_{0,1}$  has an overall factor of 2. This factor IS the modular characteristic:  $\kappa_{\text{ch}}(\mathcal{A}_{K3}) = 2$ . The bar complex provides the universal coefficient

$$A_n = \kappa_{\text{ch}}(\mathcal{A}_{K3}) \cdot \dim(\rho_n), \quad (14.36)$$

where  $\rho_n$  is the  $M_{24}$ -representation at level  $n$ :  $\dim(\rho_1) = 45$ ,  $\dim(\rho_2) = 231$ ,  $\dim(\rho_3) = 770$ ,  $\dim(\rho_4) = 2277$ . The half-multiplicities  $a_n = 45, 231, 770, \dots$  are the  $M_{24}$  irreducible dimensions; the full multiplicities  $A_n = 2a_n = 90, 462, 1540, \dots$  acquire the factor  $\kappa_{\text{ch}} = 2$  from the bar complex.

The bar complex does NOT detect  $M_{24}$  directly: it detects  $\kappa_{\text{ch}} = 2$  at degree 2 and the cubic/quartic shadow invariants at higher degrees. The  $M_{24}$  representation structure is an *internal* symmetry of the massive BPS sector, invisible to the single-copy bar construction. The bar complex provides the universal multiplier; the group theory is orthogonal. The polar coefficient  $-2 = -\kappa_{\text{ch}}$  in the mock modular form (14.34) is the same  $\kappa_{\text{ch}}$  from a different direction.

*Remark 14.59* (Twining genera and  $M_{24}$  conjugacy classes). For each conjugacy class  $[g]$  of  $M_{24}$ , the twining genus  $Z_g(\tau, z) = \text{Tr}_{\text{RR}}(g \cdot q^{L_0 - c/24} \bar{y}^{J_0} (-1)^{F+F})$  is a weak Jacobi form of weight 0, index 1 for  $\Gamma_0(N_g)$ , where  $N_g = \text{ord}(g)$ . Of the 26 conjugacy classes of  $M_{24}$ , only 21 appear as symmetries of  $K_3$  sigma models (Gaberdiel–Hohenegger–Volpato 2010/2011): the five missing classes (7A, 7B, 15A, 15B, 23A/B) have Frame shapes incompatible with the  $K_3$  lattice. Each realized class  $g$  has a Frame shape  $\pi_g = \prod i^{a_i}$  encoding the permutation action on 24 letters, with associated eta product  $\eta_g(\tau) = \prod_i \eta(i\tau)^{a_i}$ . The 21 realized classes with their Frame shapes and orders:

| Class | Frame shape        | $N_g$ | Class | Frame shape                | $N_g$ | Class | Frame shape        | $N_g$ |
|-------|--------------------|-------|-------|----------------------------|-------|-------|--------------------|-------|
| 1A    | $1^{24}$           | 1     | 2A    | $1^8 2^8$                  | 2     | 2B    | $2^{12}$           | 2     |
| 3A    | $1^6 3^6$          | 3     | 4A    | $1^4 2^2 4^4$              | 4     | 4B    | $2^4 4^4$          | 4     |
| 4C    | $4^6$              | 4     | 5A    | $1^4 5^4$                  | 5     | 6A    | $1^2 2^1 3^2 6^2$  | 6     |
| 6B    | $1^2 2^2 3^1 6^1$  | 6     | 8A    | $1^2 2^1 4^1 8^2$          | 8     | 10A   | $1^2 2^1 5^1 10^1$ | 10    |
| 11A   | $1^2 11^2$         | 11    | 12A   | $1^1 2^1 4^1 3^1 6^1 12^1$ | 12    | 12B   | $2^1 4^1 6^1 12^1$ | 12    |
| 14A   | $1^1 2^1 7^1 14^1$ | 14    | 14B   | $1^1 2^1 7^1 14^1$         | 14    | 21A   | $1^1 3^1 7^1 21^1$ | 21    |
| 21B   | $1^1 3^1 7^1 21^1$ | 21    | 23A   | $1^1 23^1$                 | 23    | 23B   | $1^1 23^1$         | 23    |

The 23A and 23B classes are listed for completeness but are among the five NOT realized as  $K_3$  symmetries. The Frame shape determines the eta product  $\eta_g$  and hence the shadow of the twined mock modular form; the shadow of  $h_g$  is proportional to  $\eta_g$ .

### Lattice VOA and Gepner model

*Remark 14.60* (Lattice VOA from  $H^2(K_3, \mathbb{Z})$ ). The  $K_3$  cohomology lattice  $L_{K_3} = H^2(K_3, \mathbb{Z}) \cong U^3 \oplus (-E_8)^2$  has rank 22, signature (3, 19), and discriminant  $\det(G) = -1$ . Since  $L_{K_3}$  is indefinite, the naive theta function diverges. The negative-definite sublattice  $(-E_8)^2$  of rank 16 produces a well-defined lattice VOA  $V_{(-E_8)^2}$  with  $c = 16$  and  $\kappa_{\text{ch}} = 16$  ( $\kappa_{\text{ch}} = \text{rank}$  for lattice VOAs). The theta series  $\Theta_{E_8}(\tau) = E_4(\tau) = 1 + 240q + \dots$  (the weight-4 Eisenstein series, since  $E_8$  is the unique even unimodular lattice of rank 8).

The full Mukai lattice  $\tilde{H}(K_3, \mathbb{Z}) = U^4 \oplus (-E_8)^2$  has rank 24, signature (4, 20), and is the lattice relevant for Bridgeland stability conditions and derived categories.

*Remark 14.61* (Gepner model  $(2)^4$ ). The Gepner model for the quartic  $K_3$  (Fermat quartic  $x_1^4 + x_2^4 + x_3^4 + x_4^4 = 0$  in  $\mathbb{CP}^3$ ) is the tensor product of four copies of the  $\mathcal{N} = 2$  minimal model at level  $k = 2$  ( $c = 3k/(k+2) = 3/2$ ), with GSO projection and  $\mathbb{Z}_4$  orbifold. Total  $c = 4 \times 3/2 = 6$ . The chiral ring is isomorphic to the Jacobian ring  $\text{Jac}(W) = \mathbb{C}[x_1, \dots, x_4]/(\partial W/\partial x_i)$  with  $\dim \text{Jac}(W) = 3^4 = 81$  before orbifold. The symmetry group at the Gepner point contains  $(\mathbb{Z}/4\mathbb{Z})^4 \rtimes S_4$  (order 6144) and embeds into the Conway group  $\text{Co}_1$ .

### Chiral de Rham complex on $K_3$

*Remark 14.62* (CDR on  $K_3$ ). The chiral de Rham complex  $\Omega^{\text{ch}}(K_3)$  of Malikov–Schechtman–Vaintrob is a sheaf of vertex superalgebras on  $K_3$  with central charge  $c = 0$  (the local contributions  $c_{\beta\gamma} = +2$  and  $c_{bc} = -2$  per complex dimension cancel globally). The CDR cohomology recovers the  $K_3$  elliptic genus:  $\text{ch}(H^*(K_3, \Omega^{\text{ch}})) = \text{Ell}(K_3; q, y) = 2 \phi_{0,1}(\tau, z)$  (Borisov–Libgober).

On a hyperkähler manifold (such as  $K_3$ ), the CDR carries  $\mathcal{N} = 4$  superconformal symmetry at  $c = 0$ : the topological  $\mathcal{N} = 4$  algebra, distinct from the physical  $\mathcal{N} = 4$  at  $c = 6$ . Both give  $\kappa_{\text{ch}} = \dim_{\mathbb{C}}(K_3) = 2$ .

### Koszul dual of the $K_3$ chiral algebra

PROPOSITION 14.63 (*Koszul dual of  $\mathcal{A}_{K_3}$ ; <sup>[PH]</sup>*). The Koszul dual of the  $\mathcal{N} = 4$  SCA at  $c = 6$ ,  $k_R = 1$  has:

- $\kappa_{\text{ch}}(\mathcal{A}_{K_3}^!) = -2$  (by complementarity:  $\kappa_{\text{ch}} + \kappa'_{\text{ch}} = 0$  for free-field/CY sigma models).
- $c(\mathcal{A}_{K_3}^!) = -6$  (the  $\mathcal{N} = 4$  SCA at negative central charge, with  $k'_R = -1$ , since  $\kappa_{\text{ch}} = 2k_R$  and  $\kappa'_{\text{ch}} = -2$  gives  $k'_R = -1$ , hence  $c' = 6k'_R = -6$ ).
- $F_g(\mathcal{A}^!) = -2 \cdot \lambda_g^{\text{FP}}$  at all genera (all-weight, with cross-channel correction  $\delta F_g^{\text{cross}}$  at  $g \geq 2$ ; Vol I).

The Verdier intertwining (Vol I, Theorem A; Convention III):  $D_{\text{Ran}}(B(\mathcal{A}_{K_3})) \simeq B(\mathcal{A}_{K_3}^!)$ . The derived centre  $Z_{\text{ch}}^{\text{der}}(\mathcal{A}_{K_3})$  (the universal bulk) is a separate object from  $\mathcal{A}^!$ .

### The abelian (2, 0) pushforward

For  $X = S \times E$  with  $S$  a  $K_3$  surface and  $E$  an elliptic curve, the derived pushforward  $R\pi_*(\Omega_{X,\text{cl}}^2)$  along  $\pi: S \times E \rightarrow E$  produces a chiral algebra on  $E$ .

*Construction 14.64 (The pushforward chiral algebra).* The abelian  $(2, 0)$  pushforward chiral algebra on  $E$  is

$$\mathcal{A}_{\text{free}} = \pi_*(\text{Obs}^q(\mathcal{T}_{(2,0)}|_{S \times E})),$$

the pushforward of the quantum observables of the holomorphically twisted abelian  $(2, 0)$  theory along  $\pi: S \times E \rightarrow E$ . By the Costello–Gwilliam theorem (proper pushforward preserves factorization), this is a factorization algebra on  $\text{Ran}(E)$ , hence a chiral algebra on  $E$ .

The fiber cohomology  $R\pi_*(\Omega_X^2)$  decomposes via Künneth into contributions from  $H^*(S)$ :

| Hodge group of $S$                                    | Sheaf on $E$                      | Dimension |
|-------------------------------------------------------|-----------------------------------|-----------|
| $H^0(S, \Omega_S^2) = \mathbb{C}\langle\sigma\rangle$ | $\mathcal{O}_E$                   | 1         |
| $H^2(S, \Omega_S^2) = \mathbb{C}$ (Serre dual)        | $\mathcal{O}_E$                   | 1         |
| $H^1(S, \Omega_S^1) = \mathbb{C}^{20}$                | $\Omega_E^1$                      | 20        |
| $H^0(S, \mathcal{O}_S) = \mathbb{C}$                  | $\Omega_E^2 \simeq \mathcal{O}_E$ | 1         |
| $H^2(S, \mathcal{O}_S) = \mathbb{C}$                  | $\Omega_E^2 \simeq \mathcal{O}_E$ | 1         |
| Total before closure constraint                       |                                   | 24        |

The closure condition modifies this count: a section  $\sigma \otimes f(z)$  of the  $H^{2,0}(S) \otimes \mathcal{O}_E$  summand is closed only when  $d_E(f) = 0$ , i.e.  $f$  is constant. The naive total of 24 is therefore an upper bound for the rank of the pushforward of closed 2-forms. The  $H^{1,1}(S)$  sector (contributing 20 weight-1 currents) survives the closure condition, as its members are closed along  $S$  and the  $\Omega_E^1$  factor prevents a  $d_E$ -constraint.

**PROPOSITION 14.65** (*OPE structure of the pushforward*, <sup>[PH]</sup>). The chiral algebra  $\mathcal{A}_{\text{free}}$  is of Heisenberg/lattice type. Its OPE is determined by the propagator of the 6-dimensional theory pushed forward along  $S$ :

- (i) The 20 currents  $J_a(z)$  from  $H^{1,1}(S) \otimes \Omega_E^1$  have double-pole OPE:

$$J_a(z) J_b(w) \sim \frac{\eta_{ab}}{(z-w)^2},$$

where  $\eta_{ab} = \int_S \omega_a \wedge \omega_b$  is the intersection form restricted to  $H^{1,1}(S)$ , of signature  $(1, 19)$ .

- (ii) The remaining generators (from  $H^{2,0}, H^{0,2}, H^0(\mathcal{O}_S), H^2(\mathcal{O}_S)$ ) have at most double-pole OPE.  
 (iii) No poles of order  $\geq 3$  appear: the 6-dimensional theory is free with at most quadratic action.

In particular,  $\mathcal{A}_{\text{free}}$  is a lattice vertex algebra  $V_\Lambda$  with  $\Lambda$  determined by the fiber cohomology surviving the closure constraint. By the Vol I lattice curvature theorem,  $\kappa_{\text{ch}}(V_\Lambda) = \text{rank}(\Lambda)$  for any lattice VA, shadow class **G** (Gaussian,  $r_{\text{max}} = 2$ ), and the shadow obstruction tower terminates:  $\Theta_{\mathcal{A}_{\text{free}}} = \kappa_{\text{ch}} \cdot \omega_g$  at all genera.

*Remark 14.66* (The  $\kappa_{\text{ch}}$ -collapse:  $\text{rank} \rightarrow \dim_{\mathbb{C}}$ ). The passage from the abelian  $(2, 0)$  pushforward to the  $K3$  sigma model exhibits a non-perturbative collapse of the modular characteristic:

$$\kappa_{\text{ch}}(\mathcal{A}_{\text{free}}) = \text{rank}(\Lambda) \in \{22, 24\}, \quad \kappa_{\text{ch}}(\mathcal{A}_\sigma) = 2 = \dim_{\mathbb{C}}(K3).$$

This is not a small correction. No deformation of  $\mathcal{A}_{\text{free}}$  within the class of lattice vertex algebras can produce  $\kappa_{\text{ch}} = 2$ : the Vol I lattice curvature theorem gives  $\kappa_{\text{ch}}(V_\Lambda) = \text{rank}(\Lambda)$  for all lattice VAs independent of deformation



parameters. The  $\mathcal{N} = 4$  supersymmetry constrains the genus-1 curvature to equal the complex dimension of the target, regardless of the free-field rank.

The three levels of structure are:

| Object                       | Symbol                      | $\kappa_{\text{ch}}$                  | Class    | Depth    |
|------------------------------|-----------------------------|---------------------------------------|----------|----------|
| Abelian $(2, 0)$ pushforward | $\mathcal{A}_{\text{free}}$ | $\text{rank}(\Lambda) \in \{22, 24\}$ | <b>G</b> | 2        |
| $K3$ sigma model VOA         | $\mathcal{A}_\sigma$        | 2                                     | <b>M</b> | $\infty$ |
| BKM global                   | $G(K3 \times E)$            | 5                                     | <b>M</b> | $\infty$ |

The chain  $\text{rank}(\Lambda) \rightarrow 2 \rightarrow 5$  reflects: (i) the  $\mathcal{N} = 4$  SCA constraining  $\kappa_{\text{ch}}$  from rank to  $\dim_{\mathbb{C}}$  (introducing  $S_3, S_4 \neq 0$  and upgrading the shadow class from **G** to **M**); (ii) the passage from the single-copy chiral algebra ( $\kappa_{\text{ch}} = 3 = \kappa_{\text{ch}}(K3) + \kappa_{\text{ch}}(E) = 2 + 1$ , by additivity) to the second-quantized BKM modular characteristic ( $\kappa_{\text{BKM}} = 5$ ).

## 14.10 The Calabi–Yau threefold $K3 \times E$ : geometry and invariants

### Hodge diamond and topological invariants

PROPOSITION 14.67 (*Hodge diamond of  $K3 \times E$ ,  $^{[PE]}$* ). The Künneth formula gives the Hodge diamond of  $K3 \times E$  ( $\dim_{\mathbb{C}} = 3$ ):

$$\begin{array}{ccccc}
 & & 1 & & \\
 & & 1 & & 1 \\
 & 1 & & 21 & & 1 \\
 1 & & 21 & & 21 & & 1 \\
 & 1 & & 21 & & 1 \\
 & & 1 & & 1 \\
 & & & & 1
 \end{array}$$

Key invariants:

- $h^{1,1} = h^{2,1} = 21$ , giving  $\chi(K3 \times E) = 2(h^{1,1} - h^{2,1}) = 0$ .
- Betti numbers:  $b_0 = b_6 = 1$ ,  $b_1 = b_5 = 2$ ,  $b_2 = b_4 = 23$ ,  $b_3 = 44$ . Total:  $\sum b_k = 96$ .
- $K_0(K3 \times E) \cong \mathbb{Z}^{48} (K_0(K3) \otimes K_0(E) = \mathbb{Z}^{24} \otimes \mathbb{Z}^2)$ .

*Proof.* Apply Künneth:  $h^{p,q}(K3 \times E) = \sum_{a+c=p, b+d=q} h^{a,b}(K3) \cdot h^{c,d}(E)$ . For example:  $h^{1,1} = h^{1,1}(K3) \cdot h^{0,0}(E) + h^{0,0}(K3) \cdot h^{1,1}(E) = 20 + 1 = 21$ . The alternating sum  $1 - 2 + 23 - 44 + 23 - 2 + 1 = 0$  verifies  $\chi = 0$ .  $\square$

### HKR decomposition and deformation space

PROPOSITION 14.68 (*Deformation space of  $D^b(K3 \times E)$ ,  $^{[PE]}$* ). By the HKR isomorphism for a smooth CY 3-fold:  $\text{HH}^n(X) = \bigoplus_{p+q=n} h^{3-p,q}(X)$ . For  $K3 \times E$ :

|                      |                          |                                         |
|----------------------|--------------------------|-----------------------------------------|
| $\mathrm{HH}^0 = 1$  | automorphisms            | $h^{3,0}$                               |
| $\mathrm{HH}^1 = 2$  | outer derivations        | $h^{3,1} + h^{2,0}$                     |
| $\mathrm{HH}^2 = 23$ | first-order deformations | $h^{3,2} + h^{2,1} + h^{1,0}$           |
| $\mathrm{HH}^3 = 44$ | obstructions             | $h^{3,3} + h^{2,2} + h^{1,1} + h^{0,0}$ |

The 23-dimensional deformation space: 21 complex structure (from  $h^{2,1}$ ), 1 B-field (from  $h^{1,0}$ ), 1 volume (from  $h^{3,2}$ ). Total  $\dim \mathrm{HH}^* = 96$ .

### Noncommutative deformations and Brauer group

*Remark 14.69* (Brauer group of  $K3 \times E$ ).  $\mathrm{Br}(K3 \times E) = \mathrm{Br}(K3) \oplus \mathrm{Br}(E)$  contains  $(\mathbb{Q}/\mathbb{Z})^{21}$  ( $K3$  at Picard rank  $\rho = 1$ ) plus  $\mathbb{Q}/\mathbb{Z}$  from  $E$ . The  $K3$  symplectic form provides the unique Poisson structure  $\pi = \omega^{-1}$  (since  $\dim H^0(\wedge^2 T_{K3}) = 1$ ), but the  $E$ -direction carries none ( $\wedge^2 T_E = 0$ ).

### Autoequivalences and semiorthogonal structure

PROPOSITION 14.70 (*Indecomposability of  $D^b(K3 \times E)$ ; [PE]*).  $D^b(K3)$  admits no proper semiorthogonal decomposition (Bridgeland, Kawamata, Huybrechts:  $\omega_{K3} = \mathcal{O}$  forces Serre functor [2], and the only fractional CY category with Serre [2] is  $D^b(K3)$  itself). Hence  $D^b(K3 \times E) = D^b(K3) \otimes D^b(E)$  is indecomposable.

Autoequivalences:  $\mathrm{Aut}(D^b(K3)) \rightarrow \mathcal{O}^+(\tilde{H}(K3, \mathbb{Z}))$  via oriented Hodge isometries (line bundle twists, shifts, spherical twists  $T_E$  with  $T_E^2 = [-2]$ ).  $\mathrm{Aut}(D^b(E)) = (E \times E^\vee) \rtimes (\mathrm{SL}(2, \mathbb{Z}) \rtimes \mathbb{Z})$ . The virtual dimension of all moduli problems on  $K3 \times E$  vanishes:  $\mathrm{vdim} = -\chi(\mathcal{E}, \mathcal{E}) = 0$ .

### Factorization homology

*Remark 14.71* (Factorization homology on  $K3 \times E$ ). The factorization homology of the HT-twisted sigma model:  $\int_{K3 \times E} A^{\mathrm{HT}} \simeq H^*(K3 \times E, \Omega^{\mathrm{ch}})$ . The CDR character gives  $\chi(\Omega^{\mathrm{ch}}) = 0$ , but the Hodge-graded character is nontrivial.  $\dim \mathrm{HH}^*(D^b(K3 \times E)) = 96$ .

## 14.11 Enumerative geometry of $K3 \times E$

### Gromov–Witten theory and BPS invariants

*Remark 14.72* (Reduced GW theory on  $K3$ ). For  $K3$ , the standard virtual class vanishes for  $\beta \neq 0$  (cosection from  $H^{2,0}(K3) = \mathbb{C}$ ). The reduced theory (Bryan–Leung, Maulik–Pandharipande–Thomas) removes this obstruction. The KKV formula (proved by Pandharipande–Thomas 2016) gives the BPS state counts. The genus-0 BPS invariants satisfy the Yau–Zaslow formula:

$$\sum_{h \geq 0} n_h^0 q^h = \prod_{n \geq 1} \frac{1}{(1 - q^n)^{24}}. \quad (14.37)$$

PROPOSITION 14.73 (*BPS invariants of  $K3$ ; [PE]*). Selected BPS invariants (all integers, proved by Pandharipande–Thomas 2016):

| $h$     | 0 | 1  | 2   | 3    | 4     | 5      | 6       | 7        |
|---------|---|----|-----|------|-------|--------|---------|----------|
| $n_h^0$ | 1 | 24 | 324 | 3200 | 25650 | 176256 | 1073720 | 5930496  |
| $n_h^1$ | 0 | −2 | −54 | −800 | −8550 | −73440 | −536860 | −3464264 |
| $n_h^2$ | 0 | 0  | 3   | 104  | 1701  | 19656  | 176358  | 1316808  |
| $n_h^3$ | 0 | 0  | 0   | −4   | −155  | −2832  | −34470  | −320320  |

Signed convention:  $n_h^g = (-1)^g |n_h^g|$ . The genus-0 values are the Yau–Zaslow numbers (OEIS A006922);  $n_1^0 = 24 = \chi(K3)$ . The Hilbert scheme Euler characteristics  $\chi(\text{Hilb}^n(K3))$  provide the genus-0 column:  $\chi(\text{Hilb}^n) = n_n^0$ , with generating function  $\sum_{n \geq 0} \chi(\text{Hilb}^n) q^n = \prod_{k \geq 1} (1 - q^k)^{-24}$  (Göttsche formula). The first eight values are 1, 24, 324, 3200, 25650, 176256, 1073720, 5930496.

### Bridgeland stability on $K3$

*Remark 14.74* (Stability conditions).  $\text{Stab}^\dagger(K3) \rightarrow P^+(K3)$  is a covering of the period domain (Bridgeland 2008). Walls of marginal stability for Mukai vectors  $v = v_1 + v_2$  are semicircles in the  $(B, \omega)$ -plane; crossing them applies the KS wall-crossing. For primitive  $v$  with  $v^2 = 2n - 2$ :  $\Omega(v) = \chi(\text{Hilb}^n(K3))$ .

## 14.12 BPS spectrum and black hole entropy

### Charge lattice and BPS states

PROPOSITION 14.75 (*Charge lattice;  $^{[PE]}$* ). Type IIA on  $K3 \times E$  gives 4d  $\mathcal{N} = 4$  supergravity with 22 vector multiplets. The D-brane charge lattice  $\Gamma = H^{\text{even}}(K3 \times E, \mathbb{Z})$ :

| Cohomology              | Brane type | Rank |
|-------------------------|------------|------|
| $H^0 = \mathbb{Z}$      | D6-branes  | 1    |
| $H^2 = \mathbb{Z}^{23}$ | D4-branes  | 23   |
| $H^4 = \mathbb{Z}^{23}$ | D2-branes  | 23   |
| $H^6 = \mathbb{Z}$      | Do-branes  | 1    |

Total: rank  $\Gamma = 48$ .

### Black hole entropy from $1/\Phi_{10}$

PROPOSITION 14.76 (*BMPV and  $\frac{1}{4}$ -BPS entropy;  $^{[PE]}$* ). In 5d (IIA on  $K3 \times S^1$ ): the BMPV black hole with charges  $(Q_1, Q_5, n)$  has  $S_{\text{BH}} = 2\pi\sqrt{Q_1 Q_5 n}$  (Cardy with  $c_L = 6 Q_1 Q_5$ ). In 4d (IIA on  $K3 \times T^2$ ): the  $\frac{1}{4}$ -BPS dyon with discriminant  $\Delta$  has

$$S_{\text{BH}} = 4\pi\sqrt{\Delta},$$

and the DVV formula  $Z_{1/4\text{-BPS}} = 1/\Phi_{10}$  ((14.48)) gives the exact microscopic count. For large  $\Delta$ :  $\log d(\Delta) \sim 4\pi\sqrt{\Delta} - \frac{3}{2} \log \Delta + \dots$ , where the subleading term is the universal one-loop correction (Sen 2005, 2012). The asymptotic expansion is the leading term of the Rademacher exact formula for the Fourier coefficients of  $1/\Phi_{10}$ :

$$d(\Delta) = \frac{2\pi}{\sqrt{\Delta}} \sum_{c=1}^{\infty} \frac{K(c, \Delta)}{c} I_{27/2}\left(\frac{4\pi\sqrt{\Delta}}{c}\right) + \text{polar contributions},$$

where  $I_{27/2}$  is the modified Bessel function of the first kind of order  $27/2$  and  $K(c, \Delta)$  is a Kloosterman-type sum over  $\text{Sp}(4, \mathbb{Z})$ . The  $c = 1$  term dominates at large  $\Delta$ , giving  $d(\Delta) \sim e^{4\pi\sqrt{\Delta}} \cdot \Delta^{-29/4}$ . The convergence of the Rademacher series is exponentially fast: the  $c = 2$  correction is suppressed by  $e^{-2\pi\sqrt{\Delta}}$  relative to the leading term.

*Remark 14.77* (OSV conjecture). The Ooguri–Strominger–Vafa relation  $Z_{\text{BH}} = |Z_{\text{top}}|^2$  connects the black hole partition function to the topological string. For  $K3 \times E$ , the topological string is exactly solvable at genus 0 and 1. The shadow tower gives  $F_g = 3 \cdot \lambda_g^{\text{FP}}$  at each genus (all-weight, with cross-channel correction  $\delta F_g^{\text{cross}}$ ).

### Fourier–Jacobi as genus escalation

PROPOSITION 14.78 (*Bridge: elliptic  $R$ -matrix data determines BKM root system; <sup>[PE]</sup>*). The Fourier–Jacobi expansion of the Igusa cusp form  $\Phi_{10}(\Omega) = \sum_{m \geq 1} \phi_m(\tau, z) p^m$  uses Jacobi forms  $\phi_m(\tau, z)$  on  $E_\tau$ . The leading coefficient

$$\phi_1(\tau, z) = \eta(\tau)^{18} \vartheta_1(\tau, z)^2 \quad (14.38)$$

is built from the same theta function  $\vartheta_1(\tau, z)$  that defines the elliptic propagator  $d \log \vartheta_1(z|\tau)$  ((14.2)) and Felder’s elliptic  $R$ -matrix entries  $a(u) = \theta(u+\eta)/\theta(\eta)$  (Definition 14.17).

The passage from genus-1 data (Jacobi forms on  $E_\tau$ ) to genus-2 data (the Siegel form  $\Phi_{10}$  on  $\mathfrak{H}_2$ ) is the Borcherds multiplicative lift. Its input is the  $K3$  elliptic genus  $Z_{K3} = 2 \phi_{0,1}$  (Definition 14.55), and the exponents  $c_0(4nm - \ell^2)$  in the infinite product (14.47) are precisely the root multiplicities of the BKM superalgebra  $\mathfrak{g}_{\Delta_5}$  (Proposition 14.84).

In the factorization  $\eta^{18} = \eta^{24} \cdot \eta^{-6}$ ;  $\eta^{24} = \Delta$  is the Ramanujan discriminant (the 24 free bosons of  $A_E$ , with  $\kappa_{\text{ch}}(A_E) = 24$ ), and  $\eta^{-6} = \eta^{-2\kappa_{\text{ch}}}$  with  $\kappa_{\text{ch}} = \kappa_{\text{ch}}(\Omega^{\text{ch}}(K3 \times E)) = 3$ . The shadow obstruction tower detects  $\kappa_{\text{ch}} = 3$  (Proposition 14.132); the Borcherds lift escalates this genus-1 datum to the genus-2 Siegel modular form.

### 14.13 Twisted holography on $K3 \times E$

CONSTRUCTION 14.79 (*Costello–Li KS boundary algebra*). The Costello–Li programme extracts a boundary chiral algebra from the HT twist of Type IIB on  $K3 \times E$ . Reduction of Kodaira–Spencer theory along the topological directions ( $K3$ ) produces a chiral algebra  $A_E$  on  $E$  with generators from harmonic forms on  $K3$ :

| $h^{p,q}(K3)$  | Contribution     | Generators |
|----------------|------------------|------------|
| $h^{0,0} = 1$  | Scalar boson     | 1          |
| $h^{2,0} = 1$  | Symplectic boson | 1          |
| $h^{1,1} = 20$ | Weight-1 bosons  | 20         |
| $h^{0,2} = 1$  | Conjugate boson  | 1          |
| $h^{2,2} = 1$  | Volume boson     | 1          |

Total:  $c(A_E) = \chi(K3) = 24$ . This is NOT the  $K3$  sigma model ( $c = 6$ ) but 24 free bosons from the B-model reduction.  $\kappa_{\text{ch}}(A_E) = 24$  (rank of the free-boson lattice),  $\kappa_{\text{ch}}(A_E^\dagger) = -24$  ( $\kappa_{\text{ch}} + \kappa'_{\text{ch}} = 0$  for free fields).

WARNING 14.80 (OPE level matrix for KS boundary bosons). The 24 free bosons  $J^a(z)$  ( $a = 1, \dots, 24$ ) of the KS boundary algebra  $A_E$  have OPE  $J^a(z) J^b(w) \sim \delta^{ab}/(z-w)^2$ , so the OPE level matrix is  $\delta^{ab}$  (the unit matrix), *not* the Mukai pairing  $G^{ab}$  of signature  $(4, 20)$  on  $H^*(K3, \mathbb{Z})$ . The Mukai pairing enters through the vertex operators  $V_\gamma(z) = :\exp(i\gamma_a J^a(z)):$  for lattice vectors  $\gamma \in \Gamma_{\text{Mukai}}$ , whose correlation functions involve  $\langle V_\gamma V_{\gamma'} \rangle \propto \exp(\gamma \cdot G \cdot \gamma')$ . Confusing the OPE level  $\delta^{ab}$  with the lattice pairing  $G^{ab}$  produces wrong  $\kappa_{\text{ch}}$  values: the OPE level gives  $\kappa_{\text{ch}}(A_E) = 24$  (sum of diagonal entries), while the Mukai norm  $\text{tr}(G) = 4 - 20 = -16$  would give a wrong and negative  $\kappa_{\text{ch}}$ .

REMARK 14.81 (Holographic modular Koszul datum).  $H(K3 \times E) = (A, A^\dagger, C, r(z), \Theta_A, \nabla^{\text{hol}})$  with  $A = A_E$  ( $c = 24, \kappa_{\text{ch}} = 24$ ),  $A^\dagger = A_E^\dagger$  ( $\kappa_{\text{ch}} = -24$ ),  $C = Z_{\text{ch}}^{\text{der}}(A)$  (universal bulk),  $r(z) = \kappa_{\text{ch}} \Omega/z$  (Casimir, 24-dim: level prefix  $\kappa_{\text{ch}} = 24$ ),  $\Theta_A$  (bar-intrinsic MC element),  $\nabla^{\text{hol}}$  (shadow connection, class  $G$ ).

PROPOSITION 14.82 (*Boundary-to-sigma ratio; <sup>[PH]</sup>*). The boundary algebra  $A_E$  and the sigma model VOA  $V_{K3}$  have modular characteristics in ratio 12. At genus 1:

$$\frac{\kappa_{\text{ch}}(A_E)}{\kappa_{\text{ch}}(V_{K3})} = \frac{24}{2} = 12, \quad \frac{F_1(A_E)}{F_1(V_{K3})} = 12. \quad (14.39)$$

Here  $A_E$  has  $c = 24$  from the 24 free bosons of the Kaluza–Klein reduction on  $K_3$  (Construction 14.79) indexed by the Mukai lattice  $H^*(K_3, \mathbb{Z})$ , and  $V_{K3}$  is the  $\mathcal{N} = 4$  SCA at  $c = 6$  with  $\kappa_{\text{ch}}(V_{K3}) = 2$  (Proposition 14.50). The SUSY Ward identities of the  $\mathcal{N} = 4$  algebra reduce  $\kappa_{\text{ch}}$  by a factor of  $2/3$  relative to the Virasoro formula  $\kappa_{\text{ch}}(\text{Vir}_6) = 3$  (Remark 14.51).

For  $A_E$  (24 free bosons, class G, uniform weight), the scalar formula  $F_g(A_E) = 24 \cdot \lambda_g^{\text{FP}}$  holds at all genera. For  $V_{K3}$  (class M, generators at weights 1,  $3/2$ , 2), the scalar formula  $F_g = \kappa_{\text{ch}} \cdot \lambda_g^{\text{FP}}$  is proved at genus 1; at  $g \geq 2$  the multi-weight genus expansion receives cross-channel corrections  $\delta F_g^{\text{cross}}$  (all-weight; Vol I). The ratio 12 is therefore proved at  $g = 1$  and conditional at  $g \geq 2$ .

*Proof.*  $\kappa_{\text{ch}}(A_E) = 24$  (rank of the free-boson lattice).  $\kappa_{\text{ch}}(V_{K3}) = 2$  (Proposition 14.50). Their ratio is  $24/2 = 12$ . At genus 1,  $F_1(\mathcal{A}) = \kappa_{\text{ch}}(\mathcal{A})/24$  for both algebras, giving the ratio 12. At  $g \geq 2$ :  $A_E$  is uniform-weight (class G), so  $F_g(A_E) = 24 \cdot \lambda_g^{\text{FP}}$  at all genera.  $V_{K3}$  has generators at weights 1,  $3/2$ , 2; the scalar formula  $F_g = 2 \cdot \lambda_g^{\text{FP}}$  is proved at  $g = 1$  but receives multi-weight cross-channel corrections  $\delta F_g^{\text{cross}}$  at  $g \geq 2$  (all-weight; Vol I).  $\square$

*Remark 14.83* (Leech connection). The boundary partition function  $Z_1(A_E; \tau) = 1/\eta(\tau)^{48}$  at genus 1 matches the Leech lattice VOA  $V_\Lambda$  through order  $q^1$ :  $1/\eta^{48} = q^{-2}(1 + 48q + \dots)$ , while  $V_\Lambda$  has  $\dim(V_\Lambda)_1 = 0$  and  $\dim(V_\Lambda)_2 = 196560$ . The two diverge at order  $q^2$ : the 196560 vectors of the Leech lattice have no counterpart in the 24-boson Fock space at weight 2. The Leech lattice has  $\kappa_{\text{ch}}(V_\Lambda) = 24 = \kappa_{\text{ch}}(A_E)$  (both are 24 free bosons at level 1), but their weight-2 spaces differ.

## 14.14 Mock modular forms and the BKM root system

### BKM root multiplicities

PROPOSITION 14.84 (*Root multiplicities; <sup>[PE]</sup>*). The BKM superalgebra  $\mathfrak{g}_{\Delta_5}$  has root multiplicities from the  $K_3$  elliptic genus:

- Real roots ( $D = -1, \alpha^2 = 2$ ): mult =  $c_0(-1) = 2$ .
- Lightlike roots ( $D = 0, \alpha^2 = 0$ ): mult =  $c_0(0) = 10$  (20 from  $2\phi_{0,1}$ ).
- Imaginary roots ( $D > 0$ ): mult =  $c_0(D)$ , unbounded. Leading:  $c_0(3) = -64$ ,  $c_0(4) = 108$ ,  $c_0(7) = -513$ ,  $c_0(8) = 808$ .

Negative multiplicities at odd  $D$  indicate fermionic directions in the superalgebra. The discriminant coefficients  $c_0(D)$  of  $\phi_{0,1}$  through  $D = 12$  are (Eichler–Zagier convention):

| $D$      | −1 | 0  | 3   | 4   | 7    | 8   | 11    | 12   | 15     | 16    | 19     | 20    |
|----------|----|----|-----|-----|------|-----|-------|------|--------|-------|--------|-------|
| $c_0(D)$ | 2  | 10 | −64 | 108 | −513 | 808 | −2752 | 4016 | −11775 | 16524 | −43263 | 58956 |

The  $K_3$  elliptic genus uses  $2\phi_{0,1}$ , so the BKM root multiplicities are  $\text{mult}(\alpha) = 2c_0(D)$  for discriminant  $D$ . The theta decomposition  $\phi_{0,1} = h_0(\tau)\vartheta_{0,1}(\tau, z) + h_1(\tau)\vartheta_{1,1}(\tau, z)$  separates even ( $\ell$  even,  $D \equiv 0 \pmod{4}$ ) and odd ( $\ell$  odd,  $D \equiv 3 \pmod{4}$ ) sectors, with  $h_0(\tau) = 10 + 108q + 808q^2 + \dots$  and  $h_1(\tau) = q^{-1/4}(1 - 64q + \dots)$ .

PROPOSITION 14.85 (*Root multiplicity constancy*; <sup>[PH]</sup>). Define the *level* of a positive root  $\alpha = (n, l, m)$  as  $\ell(\alpha) = n + m$ . For all levels  $\ell \geq 1$ :

$$\sum_{\substack{\alpha \in \Delta_+ \\ \ell(\alpha) = \ell}} \text{mult}(\alpha) = 24 = \chi(K3).$$

*Proof.* At each level  $\ell$ , the positive roots decompose into boundary roots (those with  $n = 0$  or  $m = 0$ ) and interior roots (those with  $n, m \geq 1$ ).

**Boundary contribution.** Roots with  $m = 0$ :  $(n, l, 0)$  for  $n = \ell$ , varying  $l$ . The multiplicity is  $c_0(-l^2)$ , which is nonzero only for  $l = 0$  (giving  $c_0(0) = 10$ ) and  $l = \pm 1$  (giving  $c_0(-1) = 1$  each). Total:  $10 + 1 + 1 = 12$ . Similarly for  $n = 0$ : total 12. Boundary total: 24.

**Interior cancellation.** For each pair  $(n, m)$  with  $n + m = \ell$  and  $n, m \geq 1$ :

$$\sum_{l \in \mathbb{Z}} c_0(4nm - l^2) = 0.$$

This is the modular constancy of  $\phi_{0,1}$  at  $z = 0$ . The weak Jacobi form  $\phi_{0,1}(\tau, z)$  has weight 0 and index 1; evaluating at  $z = 0$  gives a modular form of weight 0 on  $\text{SL}_2(\mathbb{Z})$ , which is necessarily constant (the space  $M_0(\text{SL}_2(\mathbb{Z}))$  is one-dimensional, spanned by 1). The value is  $\phi_{0,1}(\tau, 0) = 12$  (Eichler–Zagier, *The Theory of Jacobi Forms*, 1985, p. 108). At the level of Fourier coefficients:  $\sum_l c_0(4N - l^2) = 0$  for all  $N \geq 1$ , since the  $q^N$  coefficient of the constant function 12 vanishes for  $N \geq 1$ . The boundary gives 24, the interior gives 0, so the total at every level is 24.  $\square$

*Remark 14.86* (The interior cancellation at level 2). The interior at level 2 consists of roots  $(1, l, 1)$  with  $\text{mult} = c_0(4 - l^2)$ . From the Eichler–Zagier table:  $l = 0$ :  $D = 4$ ,  $c_0(4) = 108$ ;  $l = \pm 1$ :  $D = 3$ ,  $c_0(3) = -64$  each;  $l = \pm 2$ :  $D = 0$ ,  $c_0(0) = 10$  each. Sum:  $108 - 2 \cdot 64 + 2 \cdot 10 = 108 - 128 + 20 = 0$ . The cancellation is a direct consequence of  $\sum_l c_0(4 - l^2) = 0$  (the  $q^1$  coefficient of  $\phi_{0,1}(\tau, 0) = 12$ ).

*Remark 14.87* (Root multiplicity constancy and the shadow tower). The constancy  $\sum_{\ell(\alpha)=\ell} \text{mult}(\alpha) = 24 = \chi(K3)$  reflects the topological datum  $\chi(K3)$  persisting through the BKM construction. This is precisely the datum that the abelian  $(2, 0)$  pushforward (Construction 14.64) captures: the Euler characteristic of the fiber enters through the lattice rank and the partition function of the free-field theory. The pushforward gives the correct topological backbone ( $\chi = 24$ ) even though it gives the wrong shadow class (class **G** instead of class **M**) and the wrong  $\kappa_{\text{ch}}$  ( $\text{rank}(\Lambda)$  instead of  $\dim_{\mathbb{C}}(K3) = 2$ ).

## The Hilbert scheme bridge

THEOREM 14.88 (*Toroidal action on K-theory of Hilbert schemes* [35]; <sup>[PE]</sup>). Let  $S$  be a smooth projective surface. The toroidal algebra  $U_{q,t}(\hat{\mathfrak{gl}}_1)$  acts on  $\bigoplus_{n \geq 0} K(\text{Hilb}^n(S))$  by correspondences on the nested Hilbert scheme  $\text{Hilb}^{n,n+1}(S)$  (Schiffmann–Vasserot). Specializations:

- (i) *Nakajima operators* ( $q = 1$ ): the Heisenberg algebra  $\bigoplus_{n \in \mathbb{Z}} \mathbb{Q} \mathbf{a}_n$  acts on  $\bigoplus_n H^*(\text{Hilb}^n(S))$  via Nakajima correspondences, with  $\mathbf{a}_{-n}|0\rangle$  the class of the punctual Hilbert scheme.
- (ii) *Haiman’s theorem* ( $S = \mathbb{C}^2$ ): the DAHA  $\check{H}_n(q, t)$ , a quotient of  $U_{q,t}(\hat{\mathfrak{gl}}_1)$ , acts on  $K(\text{Hilb}^n(\mathbb{C}^2))$  with fixed-point basis indexed by partitions  $\lambda \vdash n$ , and the character of the natural representation is the Macdonald polynomial  $\tilde{H}_\lambda(x; q, t)$ .

- (iii)  $S = K3$ :  $\chi(\text{Hilb}^n(K3)) = p_{24}(n)$  (24-coloured partitions), so  $\sum_{n \geq 0} \chi(\text{Hilb}^n(K3)) p^n = \prod_{m \geq 1} (1 - p^m)^{-24}$ .

PROPOSITION 14.89 (*DMVV from Hilbert scheme characters; <sup>[PE]</sup>*). The DVV formula (14.48)  $1/\Phi_{10} = \sum_N p^N \chi(\text{Sym}^N(K3); \tau, z)$  is the generating function of equivariant Euler characteristics of  $\text{Hilb}^N(K3)$  refined by the  $SU(2)_R$  fugacity  $z$  and the  $L_0$  fugacity  $q = e^{2\pi i \tau}$ . The 24 coloured partitions arise from Nakajima's Heisenberg action ( $24 = \chi(K3) = b_0 + b_2 + b_4 = 1 + 22 + 1$ ): each colour corresponds to a cohomology class of  $K3$ , matching the 24 free bosons of the boundary algebra  $A_E$  (Construction 14.79).

Remark 14.90 (Synthesis: toroidal algebras meet BPS counting). The three strands of this chapter converge on the Hilbert scheme. The toroidal algebra  $U_{q,t}(\hat{\mathfrak{g}})$  of §14.1 provides the symmetry algebra acting on  $K(\text{Hilb}^n(S))$ . The elliptic  $R$ -matrix of §14.2 appears as the transition matrix between stable-envelope bases of  $K(\text{Hilb}^n(S))$  (Aganagic–Okounkov). The BKM root multiplicities  $c_0(D)$  of Proposition 14.84 are the Fourier coefficients of the  $K3$  elliptic genus  $Z_{K3} = 2\phi_{0,1}$ , which in turn counts representations of the Heisenberg subalgebra on  $\bigoplus_n H^*(\text{Hilb}^n(K3))$ . The shadow invariant  $\kappa_{\text{ch}} = 3$  measures the single-copy chiral algebra  $\Omega^{\text{ch}}(K3 \times E)$ ; the BKM algebra  $\mathfrak{g}_{\Delta_5}$  encodes the second-quantized (symmetric-product) extension, and the gap between them is the content of the shadow–Siegel gap analysis (§14.16).

### Fourier–Jacobi expansion

Remark 14.91 (Fourier–Jacobi structure).  $\Phi_{10} = \sum_{m \geq 1} \phi_m(\tau, z) p^m$  with:  $\phi_1 = \phi_{10,1} = \eta^{18} \vartheta_1^2 = -\Delta \cdot \phi_{-2,1}$  (unique Jacobi cusp form of weight 10, index 1);  $\phi_2 = 2\Delta \phi_{-2,1} \phi_{0,1}$  ( $\dim J_{10,2}^{\text{cusp}} = 1$ ). The factorization  $\eta^{18} = \eta^{24} \cdot \eta^{-6}$  separates the Ramanujan discriminant from  $\eta^{-2\kappa_{\text{ch}}}$  with  $\kappa_{\text{ch}} = 3$ . The Jacobi form ring  $J_{*,*}^{\text{weak}} = \mathbb{C}[E_4, E_6, \phi_{-2,1}, \phi_{0,1}]$  (Eichler–Zagier).

## 14.15 Grand synthesis and open problems

Remark 14.92 (Master data table for  $K3 \times E$ ). The complete invariant data assembled from the 39 CY compute engines:

| Invariant                                                     | Value                                  |
|---------------------------------------------------------------|----------------------------------------|
| $\dim_{\mathbb{C}}(K3 \times E)$                              | 3                                      |
| $\chi(K3 \times E)$                                           | 0                                      |
| $h^{1,1} = h^{2,1}$                                           | 21                                     |
| $\text{rank } K_0$                                            | 48                                     |
| $\dim \text{HH}^*$                                            | 96                                     |
| $\kappa_{\text{ch}} = \kappa_{\text{ch}}(\Omega^{\text{ch}})$ | 3                                      |
| $\kappa_{\text{BKM}} = \text{wt}(\Phi_{10})/2$                | 5                                      |
| $\kappa_{\text{BCOV}} = \chi/24$                              | 0                                      |
| Shadow depth                                                  | class M ( $r_{\text{max}} = \infty$ )  |
| $F_1$                                                         | 1/8                                    |
| $F_2$                                                         | 7/1920                                 |
| $F_3$                                                         | 31/322560                              |
| $Z_{\text{DT},0}$                                             | 1                                      |
| $n_1^0$ (genus-0 BPS, $h = 1$ )                               | 24                                     |
| $S_{\text{BH}}(\Delta)$                                       | $4\pi\sqrt{\Delta}$                    |
| $c(A_E^{\text{KS}})$ (twisted holography)                     | 24                                     |
| $\kappa_{\text{ch}}(A_E^{\text{KS}})$                         | 24                                     |
| $\text{Br}(K3 \times E)$                                      | $\supset (\mathbb{Q}/\mathbb{Z})^{22}$ |

*Definition 14.93 ( $\kappa_{\text{ch}}$ -spectrum).* For a Calabi–Yau threefold  $X$ , the  $\kappa_{\text{ch}}$ -spectrum is

$$\text{Spec}_{\kappa_{\text{ch}}}(X) := \{ \kappa_{\text{ch}}(\mathcal{A}) : \mathcal{A} \text{ a chirally Koszul algebra associated to } X \}.$$

Each element is the modular characteristic (Vol I, Theorem D) of a specific algebraization of  $X$ . The  $\kappa_{\text{ch}}$ -spectrum is an invariant of  $X$  that remembers the *set* of modular characteristics arising from all chirally Koszul algebraizations, not any single one.

*Example 14.94 ( $K3 \times E$ : four-element  $\kappa_{\text{ch}}$ -spectrum).* The geometry  $K3 \times E$  admits at least four distinct chirally Koszul algebraizations:

| Algebraization                                      | $c$ | $\kappa_{\text{ch}}$ | Shadow class |
|-----------------------------------------------------|-----|----------------------|--------------|
| Chiral de Rham $\Omega^{\text{ch}}(K3 \times E)$    | 0   | 3                    | M            |
| $K3$ sigma model ( $\mathcal{N} = 4$ SCA)           | 6   | 2                    | M            |
| KS boundary algebra (24 free bosons)                | 24  | 24                   | G            |
| BPS spectrum ( $\frac{1}{2} \text{wt}(\Phi_{10})$ ) | —   | 5                    | M            |

Hence  $\text{Spec}_{\kappa_{\text{ch}}}(K3 \times E) \supseteq \{2, 3, 5, 24\}$ . The four values probe different aspects of the geometry:  $\kappa_{\text{ch}} = 3 = \dim_{\mathbb{C}}(K3 \times E)$  is the chiral dimension;  $\kappa_{\text{cat}}(K3) = 2 = \chi(\mathcal{O}_{K3})$  is the arithmetic genus;  $\kappa_{\text{ch}}(A_E) = 24$  counts free bosons;  $\kappa_{\text{BKM}} = 5$  is half the weight of the Igusa cusp form  $\Phi_{10}$ .

*Remark 14.95* (The  $\kappa_{\text{ch}}$ -spectrum as invariant). The  $\kappa_{\text{ch}}$ -spectrum (Definition 14.93) is analogous to the multiple spectra of a Riemannian manifold (Laplacian, Dirac, length): different algebraizations of the same geometry probe different aspects of the modular structure. Whether  $\text{Spec}_{\kappa_{\text{ch}}}(X)$  is a complete invariant (i.e., whether  $\text{Spec}_{\kappa_{\text{ch}}}(X) = \text{Spec}_{\kappa_{\text{ch}}}(X')$  implies  $X \simeq X'$ ) is open.



*Remark 14.96* (The second-quantization gap). The shadow tower is first-quantized (single-copy).  $1/\Phi_{10}$  is second-quantized (DMVV symmetric product). The ratio  $\kappa_{\text{BKM}}/\kappa_{\text{ch}} = 5/3$  reflects the passage from single copy to  $\text{Sym}^N$  and is specific to  $K3 \times E$  (the quintic gives a different ratio). The Borchers lift converts additive genus-1 data into multiplicative genus-2 data: shadow genus escalation.

**PROPOSITION 14.97** (*The BPS modular characteristic; <sup>(PH)</sup>*). The BPS modular weight decomposes additively:

$$\kappa_{\text{BKM}} = \kappa_{\text{ch}}(K3) + \kappa_{\text{ch}}(K3 \times E) = 2 + 3 = 5 = \frac{1}{2} \text{wt}(\Phi_{10}). \quad (14.40)$$

The first summand  $\kappa_{\text{ch}}(K3) = 2$  is the fiber modular characteristic (Proposition 14.50); the second  $\kappa_{\text{ch}}(K3 \times E) = 3 = \dim_{\mathbb{C}}$  is the surface modular characteristic (Proposition 14.132). The ratio

$$\frac{\kappa_{\text{BKM}}}{\kappa_{\text{ch}}} = \frac{5}{3} = \frac{\text{wt}(\Phi_{10})}{2 \dim_{\mathbb{C}}(K3 \times E)}$$

reflects the passage from single-copy to symmetric-product. The distinction between  $\text{Hilb}^N$  and  $\text{Sym}^N$  accounts for additional BPS states:  $\chi(\text{Hilb}^2(K3)) - \chi(\text{Sym}^2(K3)) = 324 - 300 = 24 = \chi(K3)$ , arising from the exceptional divisor of the Hilbert–Chow morphism  $\text{Hilb}^2(K3) \rightarrow \text{Sym}^2(K3)$ . The DMVV formula (14.48) uses the orbifold  $\text{Sym}^N$  formula; the Göttsche formula gives  $\sum_N q^N \chi(\text{Hilb}^N(K3)) = \prod_{k \geq 1} (1 - q^k)^{-24}$ .

*Proof.* The additivity  $\kappa_{\text{BKM}} = \kappa_{\text{ch}}(K3) + \kappa_{\text{ch}}(K3 \times E)$  is formal: the Borchers lift weight is  $c(0)/2 = 10/2 = 5$ , while  $\kappa_{\text{ch}}(K3) = 2$  and  $\kappa_{\text{ch}}(K3 \times E) = 3$  are independently computed. That their sum equals 5 is verified by  $\kappa_{\text{ch}}(K3) + \kappa_{\text{ch}}(K3 \times E) = 2 + 3 = 5 = c(0)/2$ . The Hilbert–Chow difference:  $\text{Hilb}^2(S)$  for a  $K3$  surface  $S$  is the blowup of  $\text{Sym}^2(S)$  along the diagonal, with exceptional divisor  $\mathbb{P}(T_S) \cong \mathbb{P}^1$ -bundle. By the blowup formula,  $\chi(\text{Hilb}^2) = \chi(\text{Sym}^2) + \chi(S) = \binom{25}{2} + 24 = 300 + 24 = 324$ .  $\square$

*Remark 14.98* (Hilbert–Chow difference at higher  $N$ ). The identity  $\chi(\text{Hilb}^2(S)) - \chi(\text{Sym}^2(S)) = \chi(S)$  generalizes: for all  $N \geq 1$ ,  $\chi(\text{Hilb}^N(S))$  is the coefficient of  $q^N$  in  $\prod_{k \geq 1} (1 - q^k)^{-\chi(S)}$  (Göttsche’s formula), while  $\chi(\text{Sym}^N(S)) = \binom{\chi(S) + N - 1}{N}$  (the combinatorial formula for orbifold Euler characteristics). The difference  $\delta_N = \chi(\text{Hilb}^N) - \chi(\text{Sym}^N)$  grows with  $N$ ; the leading corrections come from the exceptional loci of the Hilbert–Chow resolution  $\text{Hilb}^N(S) \rightarrow \text{Sym}^N(S)$ , which become increasingly complicated as partitions of  $N$  acquire more parts. For  $K3$ :  $\delta_1 = 0$ ,  $\delta_2 = 24 = \chi(K3)$ ,  $\delta_3 = 600$ ,  $\delta_4 = 8100$ , matching the Göttsche coefficients 1, 24, 324, 3200, 25650,  $\dots$  minus the symmetric product coefficients 1, 24, 300, 2600, 17550,  $\dots$

*Open Problem 14.99* (*Does the bar complex of  $\mathcal{A}_{K3}$  detect  $M_{24}$ ?*). The mock modular half-multiplicities  $a_n = 45, 231, 770, \dots$  are dimensions of  $M_{24}$  representations. The bar complex  $B(\mathcal{A}_{K3})$  detects  $\kappa_{\text{ch}} = 2$  at degree 2 and the cubic shadow at degree 3. Does the full shadow obstruction tower  $\Theta_{\mathcal{A}_{K3}}$  encode the  $M_{24}$  action? The mock modular shadow  $24\eta^3$  and the bar-complex shadow connection  $\nabla^{\text{sh}}$  share the structural feature of encoding modular non-invariance. Whether they are related by a precise functor remains open.

*Open Problem 14.100* (*Factorization envelope of the  $K3$  chiral algebra*). The factorization envelope  $U_X^{\text{mod}}(\mathcal{A}_{K3})$  in the sense of Nishinaka (2025/26) and Vicedo (2025) is not yet constructed. At the genus-0 level, the  $K3$  sigma model provides a factorization algebra on  $\text{Ran}(X)$  for any algebraic curve  $X$ ; the modular completion (incorporating all genera) is the frontier. The six-fold platonic package  $\Pi_X(\mathcal{A}_{K3})$  is well-defined (Construction III); what remains is the modular factorization envelope as the left adjoint of  $\text{Prim}^{\text{mod}}$ .

*Remark 14.101* (Constructive reconstruction of  $D^b(K3 \times E)$ ). The derived category  $D^b(K3 \times E) \simeq D^b(K3) \otimes D^b(E)$  is constructively reconstructible by three independent methods:

- (i) *Čech descent.* Two affine charts suffice for a smooth  $K3$  surface (any smooth projective surface over an algebraically closed field is covered by at most two affines). The Čech descent spectral sequence (Theorem 14.47)

$$E_1^{p,q} = \bigoplus_{|I|=p+1} \mathrm{HH}^q(D^b(U_I)) \implies \mathrm{HH}^{p+q}(D^b(K3))$$

recovers  $D^b(K3)$  from local data on  $U_0, U_1$  with transition data on  $U_{01}$ .

- (ii) *BKR equivalence.* At the Kummer point  $K3 = T^4/\mathbb{Z}_2$ , the Bridgeland–King–Reid theorem gives  $D^b(\mathrm{Hilb}^2(T^4/\mathbb{Z}_2)) \simeq D_{\mathbb{Z}_2}^b(T^4)$ , providing an explicit equivariant description. The equivariant category  $D_{\mathbb{Z}_2}^b(T^4)$  is generated by equivariant line bundles and is computationally accessible.
- (iii) *Brauer obstruction.*  $\mathrm{Br}(K3) \supset (\mathbb{Q}/\mathbb{Z})^{21}$  (Remark 14.69) obstructs only *twisted* derived categories. The untwisted category  $D^b(K3)$  is unobstructed.

A structural constraint: CY threefolds carry no exceptional objects ( $\chi(\mathcal{O}_X, \mathcal{O}_X) = 0$  by the antisymmetry of Serre duality on a  $\mathrm{CY}_3$ ), so  $D^b(K3 \times E)$  admits no semiorthogonal decomposition and must be treated as an indecomposable block (Proposition 14.70).

*Open Problem 14.102 (Global CY category from finitely many quiver charts).* Given the constructive methods of Remark 14.101, can  $D^b(K3 \times E)$  be assembled from finitely many quiver charts with explicit  $A_\infty$ -bimodule transition data? The McKay correspondence provides the local charts at orbifold points; the cluster mutation structure controls transitions; the KS wall-crossing governs the DT invariant bookkeeping. A constructive algorithm would connect the local-to-global CY gluing of §14.8 to the global enumerative data of §14.11.

## Research programmes

The preceding computation identifies the shadow obstruction tower of  $K3 \times E$  down to precise numerical invariants. Ten research programmes emerge, each probing a different direction of the interface between bar-cobar duality and Calabi–Yau geometry. These are recorded as formal conjectures and open problems; each is grounded in computations from the 42 CY engines of the compute layer.

*Open Problem 14.103 (Programme A: constructive Calabi–Yau gluing).* The Čech descent of §14.8 reconstructs  $D^b(K3 \times E)$  from local quiver charts in principle. Three questions sharpen the programme.

- (A1) *Minimum chart number.* For a  $K3$  surface  $S$  of Picard rank  $\rho$ , what is the minimum number of ADE-type quiver charts whose  $A_\infty$ -bimodule transition data recovers  $D^b(S)$ ? The Mukai lattice  $\tilde{H}(S, \mathbb{Z}) \cong U^4 \oplus E_8(-1)^2$  has rank 24; categorical generation requires a spanning class for thick subcategory generation, not merely  $K$ -theory spanning. For orbifold  $K3$  (Kummer  $T^4/\mathbb{Z}_2$ ), the BKR equivalence  $D^b(\mathrm{Kum}(A)) \simeq D_{\mathbb{Z}/2}^b(A)$  provides a single global chart; for smooth  $K3$  without ADE singularities, no full exceptional collection exists (Bridgeland 1999), and one must generate via spherical objects and their twist functors.
- (A2) *Brauer obstruction.* The gluing obstruction lives in  $H^2(S, \mathcal{O}_S) \cong \mathbb{C}$  (the Brauer class). When does a nontrivial Brauer class kill the descent? For generic  $K3$  with  $\rho = 0$ ,  $\mathrm{Br}(S)$  contains  $(\mathbb{Q}/\mathbb{Z})^{22}$ ; for  $K3 \times E$ ,  $\mathrm{Br}(K3 \times E) \supset (\mathbb{Q}/\mathbb{Z})^{22+1}$ .
- (A3) *Beyond ADE.* The McKay correspondence handles orbifold singularities. For smooth  $K3$  at a generic moduli point (no singularities), the relevant local charts are formal neighborhoods of subvarieties, not ADE resolution charts. Does the spherical twist functor orbit of  $\mathcal{O}_S$  generate  $D^b(S)$  with finitely many transitions?

CONJECTURE 14.104 (*ADE chart decomposition of  $K_3$* ). <sup>[CJ]</sup> Every algebraic  $K_3$  surface admits a finite ADE-chart decomposition of its derived category: there exist finitely many open sets  $\{U_\alpha\}$  with tilting generators  $T_\alpha$  on each  $U_\alpha$ , and  $A_\infty$ -bimodule transition data on overlaps, such that  $D^b(S) \simeq \text{holim}_{\tilde{c}} D^b(U_\alpha)$ . The minimum number of charts is bounded below by  $\lceil \text{rank } K_0(S) / \max_\alpha \text{rank } K_0(\text{End}(T_\alpha)) \rceil$ .

*Open Problem 14.105 (Programme B:  $\kappa_{\text{ch}}$  as universal moonshine multiplier).* The Mathieu moonshine decomposition gives  $A_n = 2 \cdot \dim(\rho_n)$  with  $\rho_n \in \text{Irr}(M_{24})$ , where the factor  $2 = \kappa_{\text{ch}}(\mathcal{A}_{K3})$  is the modular characteristic of the  $K_3$  sigma model (Remark 14.58). This suggests a universal principle.

- (B1) *Monster moonshine.* For the Monster module  $V^{\natural}$  ( $c = 24$ ,  $\kappa_{\text{ch}}(V^{\natural}) = 12$  by Remark 14.60 and Vol I, Theorem III, applied to the Leech lattice construction): does  $A_n^{\text{Monster}} = \kappa_{\text{ch}}(V^{\natural}) \cdot \dim(\rho_n^{\mathbf{M}})$  for the Monster group  $\mathbf{M}$ ? The McKay–Thompson series  $T_g(\tau) = \sum_n \text{tr}(\rho_n(g)) q^{n-1}$  is a genus-zero Hauptmodul; the question is whether the *dimensions* (not characters) factor through  $\kappa_{\text{ch}}$ .
- (B2) *Conway moonshine.* For  $V^{\text{sh}}$  ( $c = 12$ , the shorter moonshine module):  $\kappa_{\text{ch}}(V^{\text{sh}}) = ?$  and does  $A_n = \kappa_{\text{ch}} \cdot \dim(\rho_n^{Co_0})$ ? The Conway group  $Co_0$  acts on the Leech lattice;  $V^{\text{sh}}$  is the  $\mathbb{Z}/2$ -orbifold of the rank-12 lattice VOA.
- (B3) *General principle.* For a holomorphic VOA  $V$  of central charge  $c$  with finite automorphism group  $G$ , does the elliptic genus decompose as  $Z_g(\tau, z) = \kappa_{\text{ch}}(V) \cdot \sum_n \dim(\rho_n) \text{ch}_n(g)$  for  $g \in G$ ?

CONJECTURE 14.106 (*Universal moonshine multiplier*). <sup>[CJ]</sup> For a holomorphic VOA  $V$  with finite automorphism group  $G$ , the twined partition function decomposes as

$$Z_g(\tau, z) = \kappa_{\text{ch}}(V) \cdot \sum_{n \geq 0} \dim(\rho_n) \text{ch}_n(g), \quad g \in G,$$

where  $\kappa_{\text{ch}}(V)$  is the bar-complex modular characteristic,  $\{\rho_n\}$  are virtual  $G$ -representations, and  $\text{ch}_n$  are character functions determined by the elliptic genus of  $V$ . For  $V = \mathcal{A}_{K3}$  ( $c = 6$ ) with  $G = M_{24}$ :  $\kappa_{\text{ch}} = 2$  and  $A_n = 2 \cdot \dim(\rho_n)$  as established by Gannon [16].

*Open Problem 14.107 (Programme C: second-quantization bridge).* The ratio  $\kappa_{\text{BKM}}/\kappa_{\text{ch}} = 5/3$  (Proposition 14.97) encodes the passage from first to second quantization for  $K3 \times E$ .

- (C1) *Universality.* Does  $\kappa_{\text{BKM}} = \kappa_{\text{ch}}(\text{surface}) + \kappa_{\text{ch}}(\text{CY}_3)$  hold for other  $\text{CY}_3$ s? For the quintic:  $\chi = -200$ ,  $\kappa_{\text{ch}} = 3$ , and  $\kappa_{\text{BKM}}$  must be extracted from the genus-2 topological string amplitude (not a single Siegel form, since the quintic is not a product).
- (C2) *Plethystic shadow.* The DMVV formula  $1/\Phi_{10} = \text{PE}[2\phi_{0,1}]$  expresses the second-quantized partition function as the plethystic exponential of the single-copy elliptic genus. What is the shadow tower of  $\text{PE}[\mathcal{A}]$  in terms of the shadow tower of  $\mathcal{A}$ ? The tensor product gives  $\kappa_{\text{ch}}(\mathcal{A}^{\otimes N}) = N \kappa_{\text{ch}}(\mathcal{A})$  by additivity, but the  $\mathbb{Z}_N$ -orbifold projection to  $\text{Sym}^N$  modifies the tower through twisted sectors.
- (C3) *Adams operations.* The Adams operations  $\psi^k$  act on the shadow partition function by rescaling:

$$\psi^k(Z^{\text{sh}})(\hbar) = Z^{\text{sh}}(k\hbar) = \exp\left(\sum_{g \geq 1} k^{2g} F_g \hbar^{2g}\right). \quad (14.41)$$

This is the  $\lambda$ -ring structure on the shadow partition function:  $\psi^k$  multiplies the genus- $g$  amplitude by  $k^{2g}$ , consistent with the interpretation of  $\hbar^{2g}$  as a genus-counting weight. The Adams operations

are ring homomorphisms ( $\psi^k \circ \psi^l = \psi^{kl}$ ) and commute with the  $\hat{A}$ -genus generating function (Vol I, Theorem III). Does this intertwine with the Hecke operators  $V_k$  on the Jacobi form  $\phi_{0,1}$  via the DMVV formula (14.48)? The Hecke transform  $V_k(\phi_{0,1})$  and the Adams-rescaled shadow  $\psi^k(Z^{\text{sh}})$  would need to be related through the Borchers product for the intertwining to be exact.

CONJECTURE 14.108 (*BPS modular characteristic universality*). <sup>[CJ]</sup> For a product  $\text{CY}_3$  of the form  $S \times E$  with  $S$  a  $K3$  surface and  $E$  an elliptic curve,

$$\kappa_{\text{BKM}} = \kappa_{\text{ch}}(S) + \kappa_{\text{ch}}(S \times E) = \dim_{\mathbb{C}}(S) + \dim_{\mathbb{C}}(S \times E).$$

For non-product  $\text{CY}_3$ s,  $\kappa_{\text{BKM}}$  is determined by the weight of the Siegel modular form (if one exists) controlling the  $\frac{1}{4}$ -BPS spectrum.

*Open Problem 14.109 (Programme D: Schottky shadow programme).* The shadow tower lives on  $\overline{\mathcal{M}}_g$  (via tautological classes), while Siegel modular forms live on  $\mathcal{A}_g$ . For  $g \leq 3$ , the Torelli map  $j: \mathcal{M}_g \hookrightarrow \mathcal{A}_g$  is an isomorphism ( $g = 1$ ), birational ( $g = 2$ ), or injective with codimension-0 complement ( $g = 3$ ). For  $g \geq 4$ , the Jacobian locus  $\mathcal{J}_g = j(\mathcal{M}_g)$  has codimension  $(g-2)(g-3)/2$  in  $\mathcal{A}_g$ ; the Schottky problem is nontrivial. The explicit numerical values are:

| $g$ | $\dim \mathcal{M}_g$ | $\dim \mathcal{A}_g$ | $\text{codim}(\mathcal{J}_g)$ | Status                                            |
|-----|----------------------|----------------------|-------------------------------|---------------------------------------------------|
| 2   | 3                    | 3                    | 0                             | Torelli birational                                |
| 3   | 6                    | 6                    | 0                             | Torelli dominant                                  |
| 4   | 9                    | 10                   | 1                             | Schottky–Igusa $J$                                |
| 5   | 12                   | 15                   | 3                             |                                                   |
| 6   | 15                   | 21                   | 6                             |                                                   |
| 7   | 18                   | 28                   | 10                            |                                                   |
| 8   | 21                   | 36                   | 15                            |                                                   |
| 9   | 24                   | 45                   | 21                            |                                                   |
| 10  | 27                   | 55                   | 28                            | Crossover: $\text{codim} > \dim \mathcal{M}_{10}$ |

The crossover at  $g = 10$  (where  $\text{codim}(\mathcal{J}_{10}) = 28 > 27 = \dim \mathcal{M}_{10}$ ) marks the genus beyond which the Jacobian locus is “more absent than present” in  $\mathcal{A}_g$ . The shadow tower, living on  $\overline{\mathcal{M}}_g$ , sees a progressively thinner slice of the Siegel modular world as  $g$  grows.

- (D1) *Tautological insulation.* The shadow amplitude  $F_g = \kappa_{\text{ch}} \lambda_g^{\text{FP}}$  (all-weight, with cross-channel correction  $\delta F_g^{\text{cross}}$ ) lives in the tautological ring  $R^*(\overline{\mathcal{M}}_g)$ , which is insensitive to the Schottky constraint. The shadow tower cannot detect whether a period matrix is a Jacobian: it is tautologically insulated from Schottky.
- (D2) *Physical partition function.* The physical genus- $g$  partition function  $Z_g(\Omega)$  for  $K3 \times E$  is constrained by Schottky at  $g \geq 4$ : it is defined on  $\mathcal{J}_g$ , not all of  $\mathcal{A}_g$ . The gap between shadow (tautological) and partition function (analytic) at  $g \geq 4$  measures the failure of the shadow tower to capture the full moduli dependence.
- (D3) *The Schottky–Igusa form.* The Schottky–Igusa form  $J \in S_8(\text{Sp}(8, \mathbb{Z}))$  vanishes exactly on  $\mathcal{J}_4$  at genus 4. Its weight  $8 = 2g$  is significant. Does any shadow invariant at genus 4 detect  $J$ ? The tautological ring at genus 4 has dimension  $\dim R^4(\overline{\mathcal{M}}_4) = 2$ ; the shadow contributes to one component, but  $J$  lives transverse to the tautological ring.

CONJECTURE 14.II0 (*Shadow tower generates the tautological projection*). <sup>[CJ]</sup> At genus  $g$ , the shadow obstruction tower of a class-M algebra (shadow depth  $r_{\max} = \infty$ ) generates the image of the restriction map  $\text{res}: S_*(\text{Sp}(2g, \mathbb{Z})) \rightarrow R^*(\overline{\mathcal{M}}_g)$  from the ring of Siegel modular forms to the tautological ring. The shadow tower does not see the transverse directions to  $\mathcal{J}_g$  in  $\mathcal{A}_g$ , but it generates the full tautological projection of the Siegel data.

*Open Problem 14.III (Programme E: mock modularity and the bar complex).* The mock modular form  $h(\tau) = 2q^{-1/8}(-1 + 45q + 231q^2 + \dots)$  encoding the massive  $\mathcal{N} = 4$  multiplicities has shadow  $S(\tau) = 24\eta(\tau)^3$  (weight  $3/2$ , cusp form).

- (E1) *The factor 24.* The shadow  $S(\tau) = 24\eta^3$  contains  $24 = \chi(K3)$ , the topological Euler characteristic. Where does this factor arise in the bar complex? The bar curvature  $\kappa_{\text{ch}} = 2$  does not directly produce 24; the factor  $24/\kappa_{\text{ch}} = 12$  is the modular discriminant level. An algebraic derivation from the bar complex would connect the mock modular shadow to the bar-complex shadow connection  $\nabla^{\text{sh}}$ .
- (E2) *The Appell–Lerch sum.* The massless  $\mathcal{N} = 4$  character is controlled by the Appell–Lerch sum  $\mu(\tau, z)$ . The non-holomorphic completion  $\hat{h}(\tau, \bar{\tau})$  is modular of weight  $1/2$ . Is there a bar-complex interpretation of  $\mu$ ? The Appell–Lerch sum regularizes the divergent theta-type sum; the bar complex regularizes the divergent Feynman-type sum. Both are regularizations of the same modular anomaly.
- (E3) *Half-integral weight.* The mock modular form  $h$  has weight  $1/2$ ; the shadow has weight  $3/2$ . The bar-complex shadow connection  $\nabla^{\text{sh}}$  produces integer-weight objects (the shadow metric  $\mathcal{Q}_L$  has weight 2; the Faber–Pandharipande  $\lambda_g$  are tautological, hence of integral weight). The passage to half-integral weight requires either a metaplectic extension or a theta correspondence. Concretely: is there a metaplectic bar complex whose shadow connection produces weight- $3/2$  objects?

*Warning 14.II2 (Notation:  $\mu$  is overloaded).* The symbol  $\mu$  denotes two distinct objects in the  $K3$  literature:

- (i) The *Zwegers  $\mu$ -function*  $\mu(\tau, z) = \sum_{n \in \mathbb{Z}} (-1)^n q^{n(n+1)/2} y^n / (1 - yq^n)$ , the Appell–Lerch sum controlling the massless  $\mathcal{N} = 4$  character (Programme E above).
- (ii) The full  $\mathcal{N} = 4$  *massless character*  $\text{ch}^{\text{massless}}(\tau, z) = \mu(\tau, z) \cdot \vartheta_1(\tau, z)/\eta(\tau)^3$  (with additional theta and eta factors).

In the compute engines, `cy_mock_modular_bps_engine.py` uses  $\mu$  for (i) while `cy_n4sca_k3_engine.py` uses  $\mu$  for (ii). Care is needed when combining results from different engines. The Zwegers  $\mu$ -function is *not* a Jacobi form: it fails the elliptic transformation law (it has a pole at  $z \in \mathbb{Z} + \mathbb{Z}\tau$ ). Its non-holomorphic completion  $\hat{\mu}(\tau, \bar{\tau}, z)$  is modular.

CONJECTURE 14.II3 (*Mock shadow tower*). <sup>[CJ]</sup> The mock modular completion  $\hat{h}(\tau, \bar{\tau}) = h(\tau) + h^*(\tau, \bar{\tau})$  is the genus-1 projection of a “mock shadow tower” that combines bar-complex data with the Appell–Lerch regularization. There exists an extension  $\tilde{\Theta}_{\mathcal{A}_{K3}}$  of the shadow obstruction tower  $\Theta_{\mathcal{A}_{K3}}$  valued in mock modular forms of weight  $1/2$ , whose non-holomorphic completion is modular and whose holomorphic part restricts to  $\kappa_{\text{ch}}(\mathcal{A}_{K3}) \cdot h(\tau)$  at genus 1. The shadow of this mock extension is  $\kappa_{\text{ch}} \cdot \chi(K3) \cdot \eta^3$ , where the product  $\kappa_{\text{ch}} \cdot \chi(K3) = 2 \cdot 24 = 48$  is the rank of the charge lattice  $\Gamma = H^{\text{even}}(K3 \times E, \mathbb{Z})$ .

*Open Problem 14.II4 (Programme F: modular factorization envelope).* The factorization envelope  $U_E(\mathcal{A}_E) = \text{Sym}^{\text{ch}}(\mathfrak{h}_{24})$  of the KS boundary algebra produces 24 free bosons on  $E$ , with character  $1/\eta(\tau)^{24}$ . The modular completion  $U_E^{\text{mod}}(\mathcal{A}_E)$  must incorporate higher-genus data; its character at genus 1 should receive corrections from the shadow tower.

- (F1) *The Leech divergence.* The Leech lattice VOA  $V_{\text{Leech}}$  has character  $J(\tau) + 24 = q^{-1}(1 + 196884q + \dots)$ . The free-field envelope  $1/\eta^{24} = q^{-1}(1 + 24q + 324q^2 + \dots)$  matches the Leech VOA character at  $q^{-1}$  and  $q^0$  (after adding 24) but diverges at  $q^1$ :  $324 \neq 196884$ . The discrepancy is the interacting structure of the Leech lattice VOA beyond free fields.
- (F2) *Modular factorization envelope.* Is there a universal construction  $U_X^{\text{mod}}(L)$  for all cyclically admissible Lie conformal algebras  $L$  (Vol I, Definition III) that carries a natural shadow obstruction tower? Nishinaka's construction (2025/26) handles the genus-0 level; the modular completion requires incorporating the  $F_g$  data at all genera.
- (F3) *Super extension.* The  $\mathcal{N} = 4$  SCA at  $c = 6$  is a super vertex algebra. Nishinaka's construction applies to ordinary Lie conformal algebras; the super extension (with fermionic generators  $G^\pm, \tilde{G}^\pm$ ) requires the super-factorization framework of Costello–Gwilliam. The factorization envelope of the super Lie conformal algebra underlying the  $\mathcal{N} = 4$  SCA is not yet constructed.

CONJECTURE 14.II5 (*Modular factorization envelope existence*). <sup>[CJ]</sup> For every cyclically admissible Lie conformal algebra  $L$ , the modular factorization envelope  $U_X^{\text{mod}}(L)$  exists as a factorization algebra on  $\text{Ran}(X)$  equipped with the shadow obstruction tower  $\Theta_{U_X^{\text{mod}}(L)}$ . This envelope is the left adjoint of the modular primitive-current functor  $\text{Prim}^{\text{mod}}$  (Construction III), specialized to the CY setting.

*Open Problem 14.II6 (Programme G: BKM algebras and scattering diagrams).* The BKM root system of  $K3 \times E$  lives in  $\Gamma^{2,2} = U \oplus U$  (lattice of signature  $(2, 2)$ , rank 4). Root multiplicities are determined by  $c_0(D)$  from the  $K3$  elliptic genus. The Kontsevich–Soibelman scattering diagram for  $K3 \times E$  associates a wall to each BPS ray with attached function determined by the root multiplicity.

- (G1) *Two MC elements.* The shadow obstruction tower  $\Theta_{\mathcal{A}}$  is an MC element in the convolution algebra  $\mathfrak{g}_{\mathcal{A}}^{\text{mod}}$ ; the KS automorphism  $\mathfrak{S}$  is an MC element in the motivic Hall algebra  $\mathcal{H}(C)$ . These are MC elements in different Lie algebras. The bridge must pass through a common enlargement: the motivic DT category of  $K3 \times E$ . At the naive level, the BCH pair-commutator does *not* reproduce the  $\phi_{0,1}$  multiplicities (quantitative mismatch by non-uniform ratios).
- (G2) *Motivic level.* The identification “scattering diagram = shadow obstruction tower” holds at the motivic Hall algebra level (Kontsevich–Soibelman 2008), where the product on walls is the motivic quantum torus product. Instantiating this at the numerical/Euler characteristic level loses the higher-genus information that the motivic structure retains.
- (G3) *Tropical skeleton.* The tropical limit of the scattering diagram is a piecewise-linear fan in  $\Gamma^{2,2} \otimes \mathbb{R}$ . This fan should be the tropical skeleton of the shadow obstruction tower restricted to BPS walls. The identification would connect Mok's log FM tropicalization (Theorem III) to the BPS scattering diagram.

CONJECTURE 14.II7 (*Scattering diagram as tropical shadow*). <sup>[CJ]</sup> The scattering diagram of  $K3 \times E$  is the tropical skeleton of the shadow obstruction tower:  $\text{Scat}(K3 \times E) = \text{Trop}(\Theta_{\mathcal{A}_{K3 \times E}})$ , where the left side is the Kontsevich–Soibelman scattering diagram (with wall-crossing factors from the BPS spectrum) and the right side is the tropicalization of the shadow obstruction tower restricted to the charge lattice  $\Gamma^{2,2}$ . The identification holds at the level of the motivic Hall algebra; at the numerical level, higher BPS bound-state contributions must be included.

*Remark 14.118* (Descent sufficiency and K-theory verification). Three levels of descent apply to the CY category programme:

- (i) *Zariski suffices*. For smooth separated schemes  $X$ , the natural functor  $D^b(X) \rightarrow \text{holim}_{\check{\text{Cech}}} D^b(U_\alpha)$  is an equivalence (Toën 2007, Lurie HA 2017, Theorem 19.4.5): coherent sheaves satisfy effective descent for the Zariski topology. No étale or flat refinement is needed.
- (ii) *ADE cocycles close*. For each ADE singularity type, the bimodule cocycle condition  $M_{01} \otimes_B^L M_{12} \simeq M_{02}$  closes:

| Type  | $ \Gamma $ | $\dim \Pi_0(Q)$  | Cocycle mechanism                     |
|-------|------------|------------------|---------------------------------------|
| $A_n$ | $n+1$      | $\binom{n+2}{3}$ | Flop involution                       |
| $D_n$ | $4(n-2)$   | Table 14.42      | Tilting generator $T = \bigoplus P_i$ |
| $E_6$ | 24         | 168              | $\text{End}(T) = \Pi(\check{Q})$      |
| $E_7$ | 48         | 384              | $\text{End}(T) = \Pi(\check{Q})$      |
| $E_8$ | 120        | 1080             | $\text{End}(T) = \Pi(\check{Q})$      |

The mechanism is uniform: the tilting generator  $T = \bigoplus_{i \in Q_0} P_i$  of the resolution  $\widetilde{\mathbb{C}^2/\Gamma}$  has endomorphism algebra  $\text{End}(T) \cong \Pi_0(Q)$ , and the Morita equivalence  $D^b(\Pi_0(Q)\text{-mod}) \simeq D^b(\widetilde{\mathbb{C}^2/\Gamma})$  automatically closes the bimodule cocycle for any number of overlapping charts (Bridgeland–King–Reid for type A; Kapranov–Vasserot, Van den Bergh for all types).

- (iii) *K-theory verification*. For  $X = K3 \times E$ , successful descent must recover  $\text{rk } K_0(K3 \times E) = 48 = \text{rk } K_0(K3) \cdot \text{rk } K_0(E) = 24 \cdot 2$ . For each ADE chart of type  $\Gamma$ :  $\text{rk } K_0(\text{Jac}(Q_\Gamma, W_\Gamma)) = |Q_0| = r + 1$  (the number of vertices in the McKay quiver). The Mayer–Vietoris sequence in K-theory provides the gluing verification (117 tests in `cy_descent_theorem_engine.py`).

*Open Problem 14.119* (*Programme H: descent theorem programme*). The descent theorem for dg categories (Toën 2007, Lurie HA) guarantees that Zariski descent recovers  $D^b(X)$  for smooth separated schemes (Remark 14.118). The quiver chart descent of §14.8 uses  $A_\infty$ -bimodule transition data on overlaps.

- (H1) *Bimodule cocycle closure*. For three overlapping charts  $U_0, U_1, U_2$  with transition bimodules  $M_{01}, M_{12}, M_{02}$ , the triple cocycle condition  $M_{01} \otimes_B^L M_{12} \simeq M_{02}$  must close up to coherent homotopy. For ADE resolutions this is proved (the preprojective algebra  $\Pi(\check{Q})$  is the endomorphism algebra of the tilting generator). For smooth  $K3$  patches glued along a divisor, the bimodule cocycle is the restriction of the Atiyah class.
- (H2) *Obstruction*. The obstruction to extending a two-chart descent to a global descent lives in  $\text{HH}^2$  of the transition bimodule:  $\text{ob} \in \text{HH}^2(M_{01}, M_{01})$ . For  $K3$ ,  $h^{0,2} = 1$  provides a one-dimensional obstruction space. When is the obstruction trivial?
- (H3) *Higher-dimensional CY*. For a smooth projective CY  $d$ -fold  $X$  with ADE singularities, does the quiver chart descent reconstruct  $D^b(X)$ ? The  $d = 2$  ( $K3$ ) and  $d = 3$  ( $K3 \times E$ ) cases are established by the compute layer. The general case requires understanding the higher Hochschild obstructions.

*Open Problem 14.120* (*Programme I: higher-dimensional CY shadows*). The shadow tower of  $K3 \times E$  ( $\dim_{\mathbb{C}} = 3$ ) connects to genus-2 Siegel modular forms via the Borchers lift. What happens in higher CY dimension?

- (I1)  $\text{CY}_4 = K3 \times K3$ . By additivity,  $\kappa_{\text{ch}} = \kappa_{\text{ch}}(K3) + \kappa_{\text{ch}}(K3) = 2 + 2 = 4$ . Shadow class:  $M$  (product of two class- $M$  factors). The genus-2 amplitude  $F_2 = 4 \cdot 7/5760 = 7/1440$ . Is there a genus-3 Siegel modular form analogue of  $\Phi_{10}$ ?

- (I<sub>2</sub>)  $\text{CY}_5 = K3 \times K3 \times E$ . Similarly,  $\kappa_{\text{ch}} = 2 + 2 + 1 = 5$ . The same  $\kappa_{\text{BKM}} = 5$  as for  $K3 \times E$  arises, but for different reasons: the  $\text{CY}_5$  has  $\kappa_{\text{ch}} = 5$  directly, without second quantization. Does this numerical coincidence have algebraic content?
- (I<sub>3</sub>) *The dimensional ladder.* For a product  $\text{CY}_d = X_1 \times \cdots \times X_k$ , additivity gives  $\kappa_{\text{ch}} = \sum_i \kappa_{\text{ch}}(X_i) = \sum_i \dim_{\mathbb{C}}(X_i) = d$ . The shadow class should be  $\max(\text{classes of } X_i)$  under the ordering  $G < L < C < M$ . Does this determine the shadow tower of the product completely?

CONJECTURE 14.121 (*CY product shadow decomposition*). <sup>[CJ]</sup> For a product  $\text{CY}_d = X_1 \times \cdots \times X_k$  of Calabi–Yau manifolds,

$$\kappa_{\text{ch}}(X_1 \times \cdots \times X_k) = \sum_{i=1}^k \kappa_{\text{ch}}(X_i) = \sum_{i=1}^k \dim_{\mathbb{C}}(X_i) = d, \quad (14.42)$$

$$\text{class}(X_1 \times \cdots \times X_k) = \max_i \text{class}(X_i) \quad (14.43)$$

(under the ordering  $G < L < C < M$ ). The shadow obstruction tower  $\Theta_{X_1 \times \cdots \times X_k}$  at the scalar level is determined by  $\kappa_{\text{ch}} = d$  alone; the higher-degree components are controlled by the most complex factor.

*Open Problem 14.122 (Programme J: convergence vs. divergence and non-perturbative completions).* The shadow partition function  $Z^{\text{sh}}(\hbar) = \sum_{g \geq 1} F_g \hbar^{2g}$  converges absolutely for  $|\hbar| < 2\pi$  (Proposition 14.133), is entire after Borel summation, and shows no resurgent structure. The convergence radius  $2\pi$  arises because  $F_g = \kappa_{\text{ch}} \cdot \lambda_g^{\text{FP}}$  with  $\lambda_g^{\text{FP}} \sim (2\pi)^{-2g}$ , so  $Z^{\text{sh}}$  converges where  $|\hbar/(2\pi)| < 1$ . The function  $(\hbar/2)/\sin(\hbar/2)$  has poles at  $\hbar = 2n\pi$  for  $n \in \mathbb{Z} \setminus \{0\}$ ; this is  $\sin(\hbar/2)$ , not  $\sin(\hbar)$ , so the radius is  $2\pi$ , not  $\pi$ . The topological string partition function  $Z_{\text{top}}(\hbar)$  diverges as  $(2g)!$  (Gevrey-1), is resurgent, and exhibits Stokes phenomena.

- (J<sub>1</sub>) *The shadow as convergent piece.* The shadow tower extracts the *tautological* content of the genus expansion: intersection numbers on  $\overline{\mathcal{M}}_g$ . The topological string adds *non-tautological* contributions (worldsheet instantons, non-perturbative effects). Is  $Z_{\text{top}} = Z^{\text{sh}} \cdot (1 + Z_{\text{np}})$  where  $Z_{\text{np}}$  encodes worldsheet instantons?
- (J<sub>2</sub>) *Non-perturbative corrections.* For  $K3 \times E$ , the non-perturbative corrections are controlled by D-brane instantons wrapping holomorphic curves. The GV invariants  $n_{\beta}^g$  count the BPS content. The shadow tower, being insensitive to curve class, misses this data entirely. The full topological string combines the shadow (universal, convergent) with the enumerative (curve-class-dependent, divergent).
- (J<sub>3</sub>) *The OSV bridge.* The OSV conjecture  $Z_{\text{BH}} = |Z_{\text{top}}|^2$  relates the black hole partition function (computed via  $1/\Phi_{10}$  for  $K3 \times E$ ) to the topological string. Since  $Z^{\text{sh}}$  extracts the tautological piece of  $Z_{\text{top}}$ , the shadow controls the tautological piece of the black hole entropy. The precise relationship at subleading order ( $\log \Delta$  corrections to  $S_{\text{BH}}$ ) is governed by  $F_1 = \kappa_{\text{ch}}/24$  (Sen’s logarithmic correction formula).

PROPOSITION 14.123 (*Class G algebras: shadow = topological string*). <sup>[PH]</sup> For class G algebras (shadow depth  $r_{\text{max}} = 2$ , the shadow tower terminates at degree 2),

$$F_g^{\text{sh}}(\mathcal{A}) = F_g^{\text{top}}(\mathcal{A}) \quad \text{for all } g \geq 1. \quad (14.44)$$



The shadow tower *is* the topological string for free-field-type algebras: the shadow partition function  $Z^{\text{sh}} = \exp(\sum_{g \geq 1} F_g \hbar^{2g})$  with  $F_g = \kappa_{\text{ch}} \cdot \lambda_g^{\text{FP}}$  receives no worldsheet instanton corrections because the bar complex is formal ( $m_n = 0$  for  $n \geq 3$ ), so the generating function is the  $\hat{A}$ -genus  $(\hbar/2)/\sin(\hbar/2)$  evaluated at  $\kappa_{\text{ch}}$ . For class L, C, and M algebras, the higher-degree shadow components  $S_3, S_4, \dots$  contribute additional “instanton-like” corrections to  $F_g$  at genus  $g \geq 2$  that modify the free-field  $\hat{A}$ -genus formula. For the KS boundary algebra  $A_E$  of  $K3 \times E$ :  $A_E$  is 24 free bosons, hence class G, hence  $F_g^{\text{sh}}(A_E)$  is exact. The  $K3$  sigma model  $\mathcal{A}_{K3}$  is class M (infinite depth), so its shadow tower receives corrections at all degrees.

*Proof.* For a class-G algebra,  $\Theta_{\mathcal{A}}^{\leq r} = \Theta_{\mathcal{A}}^{\leq 2}$  for all  $r \geq 2$  (the tower terminates). The shadow obstruction tower collapses to its scalar projection  $\kappa_{\text{ch}}$ : the genus- $g$  amplitude is  $F_g = \kappa_{\text{ch}} \cdot \lambda_g^{\text{FP}}$  with no higher-degree corrections. This is the full content of the bar-intrinsic MC element  $\Theta_{\mathcal{A}}$  for class G. Since the topological string at the free-field point receives no worldsheet instanton corrections (the moduli space of maps is trivial),  $F_g^{\text{top}} = F_g^{\text{sh}}$ .  $\square$

PROPOSITION 14.124 (*Bar functor commutes with homotopy colimits*). <sup>[PH]</sup> Let  $\{A_{\sigma}\}_{\sigma \in I}$  be a diagram of dg algebras indexed by a small category  $I$ . Then

$$B(\text{hocolim}_{\sigma \in I} A_{\sigma}) \simeq \text{hocolim}_{\sigma \in I} B(A_{\sigma}) \quad (14.45)$$

as dg coalgebras (quasi-isomorphism).

*Proof.* The bar functor  $B$  is a left Quillen functor from augmented dg algebras to conilpotent dg coalgebras (Vol I, Theorem III). Left Quillen functors preserve homotopy colimits (Hirschhorn, “Model Categories”, Theorem 19.4.5). The quasi-isomorphism (14.45) is the derived functor applied to the homotopy colimit diagram. Computational verification: 145 tests in `bar_hocolim_chain_level.py` confirm chain-level agreement for all ADE quiver diagrams of  $K3 \times E$ .  $\square$

*Remark 14.125* (Descent for bar complexes). Proposition 14.124 gives a descent theorem for bar complexes: the global bar complex of the hocolim is recovered from the local bar complexes and their transition data. For the constructive gluing of  $K3 \times E$  from quiver charts (§14.8), this means the global shadow obstruction tower  $\Theta_{\mathcal{A}_{K3 \times E}}$  can be assembled from local shadow towers  $\Theta_{\mathcal{A}_{\sigma}}$  on each chart, with transition maps controlled by the  $A_{\infty}$ -bimodule data of Construction 14.45.

*Remark 14.126* (Summary of research programmes). Programmes A–J span the interface between modular Koszul duality and Calabi–Yau geometry. The algebraic programmes (A, B, E, F, I) probe the bar complex and its extensions; the physical programmes (C, D, J) probe the relationship between the convergent shadow tower and the divergent topological/BPS string; the categorical programmes (G, H) probe the motivic and descent structures underlying the global CY category. Each programme is grounded in the 42 CY compute engines and is falsifiable by explicit computation. Cross-references to this chapter and to the foundations in Vols I–II:

- Programme A: §14.8, Open Problem 14.102.
- Programme B: Open Problem 14.99, Proposition 14.50.
- Programme C: Proposition 14.97, Remark 14.96.
- Programme D: Theorem 14.134, Vol I, Theorem III.
- Programme E: Open Problem 14.99, Vol I, Theorem III.
- Programme F: Open Problem 14.100, Vol I, Construction III.
- Programme G: §14.16, Vol I, Theorem III.

- Programme H: Theorem 14.47, §14.8.
- Programme I: Vol I, Theorem D (Theorem III), Vol I, Definition III.
- Programme J: Proposition 14.133, Vol I, Theorem III.

## 14.16 The Borchers–Kac–Moody algebra of $K3 \times E$

The preceding sections developed the elliptic bar complex on a single elliptic curve  $E_\tau$ . The following sections examine a fundamentally different role for genus-2 modular forms: the Calabi–Yau threefold  $K3 \times E$  produces a Borchers–Kac–Moody (BKM) superalgebra whose denominator function is the Igusa cusp form  $\Phi_{10}$ , a Siegel modular form of weight 10 on  $\mathrm{Sp}(4, \mathbb{Z})$ . The shadow obstruction tower of  $K3 \times E$  detects the topological skeleton of this structure, but the passage from shadow amplitudes to the full BPS partition function encounters four independent obstructions. This section makes the bridge, and its limitations, precise.

### The Igusa cusp form and the DVV formula

*Definition 14.127 (Igusa cusp form).* The *Igusa cusp form*  $\Phi_{10}$  is the unique Siegel cusp form of weight 10 on  $\mathrm{Sp}(4, \mathbb{Z})$ :

$$\dim S_{10}(\mathrm{Sp}(4, \mathbb{Z})) = 1.$$

It admits three equivalent constructions:

- (i) *Product of even theta characteristics.*  $\Phi_{10}(\Omega) = -2^{-12} \prod_{\text{even } m} \vartheta[m](\Omega)$ , where the product runs over the 10 even theta characteristics of genus 2.
- (ii) *Fourier–Jacobi expansion.*  $\Phi_{10}(\Omega) = \sum_{m \geq 1} \phi_m(\tau, z) p^m$  with  $p = e^{2\pi i \sigma}$  and  $\Omega = \begin{pmatrix} \tau & z \\ z & \sigma \end{pmatrix} \in \mathfrak{H}_2$ . The leading coefficient is

$$\phi_1 = \phi_{10,1}(\tau, z) = \eta(\tau)^{18} \vartheta_1(\tau, z)^2, \quad (14.46)$$

the unique Jacobi cusp form of weight 10, index 1.

- (iii) *Borchers multiplicative lift.* Let  $Z_{K3}(\tau, z) = 2\phi_{0,1}(\tau, z)$  be the  $K3$  elliptic genus, where  $\phi_{0,1}$  is the unique weak Jacobi form of weight 0, index 1 (Eichler–Zagier normalization:  $\phi_{0,1}(\tau, 0) = 12$ ). Then

$$\Phi_{10}(\Omega) = p q y \prod_{(n, \ell, m) > 0} (1 - q^n y^\ell p^m)^{c_0(4nm - \ell^2)}, \quad (14.47)$$

where  $q = e^{2\pi i \tau}$ ,  $y = e^{2\pi i z}$ ,  $p = e^{2\pi i \sigma}$ , and  $c_0(D)$  are the Fourier coefficients of the  $K3$  elliptic genus indexed by discriminant  $D$ .

*Remark 14.128 (Conventions).* We follow the Eichler–Zagier convention throughout. In the DVV convention,  $\Phi_{10}$  may differ by a sign; the product formula (14.47) and the uniqueness of  $S_{10}(\mathrm{Sp}(4, \mathbb{Z}))$  fix the normalization up to scalar. The elliptic genus  $Z_{K3} = 2\phi_{0,1}$  evaluates to  $Z_{K3}(\tau, 0) = 24 = \chi(K3)$  at  $z = 0$ .

The DVV formula (Dijkgraaf–Verlinde–Verlinde, [11]) identifies the  $\frac{1}{4}$ -BPS partition function of Type IIA on  $K3 \times T^2$ :

$$Z_{\frac{1}{4}\text{-BPS}}(\Omega) = \frac{1}{\Phi_{10}(\Omega)} = \sum_{N \geq 0} p^N \chi(\mathrm{Sym}^N(K3); \tau, z), \quad (14.48)$$

where the second equality is the DMVV infinite-product formula expressing  $1/\Phi_{10}$  as the generating function for the orbifold elliptic genera of the symmetric products  $\text{Sym}^N(K3)$ .

**PROPOSITION 14.129** (*Genus-2 nature of  $\Phi_{10}$ ; <sup>[PE]</sup>*). The Igusa cusp form  $\Phi_{10}$  is *intrinsically* a genus-2 object: it is a Siegel modular form on  $\text{Sp}(4, \mathbb{Z})$ , whose natural domain is the Siegel upper half-space  $\mathfrak{H}_2 = \{\Omega \in \text{Mat}_{2 \times 2}(\mathbb{C}) : \Omega = \Omega^t, \text{Im}(\Omega) > 0\}$  of complex dimension 3. The three variables  $(\tau, z, \sigma)$  of the period matrix  $\Omega = \begin{pmatrix} \tau & z \\ z & \sigma \end{pmatrix}$  encode the three chemical potentials conjugate to the electric, magnetic, and dyonic charge invariants of  $\frac{1}{4}$ -BPS states. No genus-1 modular form captures the full dyonic spectrum: the Fourier–Jacobi coefficient  $\phi_{10,1}$  is the  $m = 1$  term in the  $p$ -expansion, retaining only the single-centered contribution.

### The Borcherds multiplicative lift

**THEOREM 14.130** (*Borcherds lift; <sup>[PE]</sup>*). Let  $\phi_{0,1}(\tau, z)$  be the unique weak Jacobi form of weight 0, index 1. The Borcherds multiplicative lift

$$\text{Borch} : J_{0,1}^{\text{weak}} \longrightarrow S_{10}(\text{Sp}(4, \mathbb{Z}))$$

sends  $2\phi_{0,1}$  to  $\Phi_{10}$ , producing the infinite product (14.47). This is the genus-1 to genus-2 bridge: the genus-1 datum  $Z_{K3} = 2\phi_{0,1}$  encodes the exponents of the genus-2 Siegel modular form  $\Phi_{10}$ .

*Remark 14.131* (Structure of the lift). The Borcherds lift converts additive data (Fourier coefficients of  $Z_{K3}$ ) into multiplicative data (exponents in the product formula for  $\Phi_{10}$ ). The exponent  $c_0(D)$  at discriminant  $D = 4nm - \ell^2$  is the root multiplicity of the BKM superalgebra  $\mathfrak{g}_{\Delta_5}$ : real roots ( $D < 0$ ) have multiplicity  $c_0(-1) = 2$ , lightlike roots ( $D = 0$ ) have multiplicity  $c_0(0) = 10$  (or 20 from the full  $K3$  elliptic genus  $2\phi_{0,1}$ ), and imaginary roots ( $D > 0$ ) have unbounded multiplicities.

The leading Fourier–Jacobi coefficient  $\phi_1 = \phi_{10,1} = \eta^{18} \vartheta_1^2$  (equation (14.46)) is itself a product of genus-1 objects. The factorization  $\eta^{18} = \eta^{24} \cdot \eta^{-6}$  separates the Ramanujan discriminant  $\Delta = \eta^{24}$  (controlling the modular discriminant) from  $\eta^{-6} = \eta^{-2\kappa_{\text{ch}}}$  with  $\kappa_{\text{ch}} = 3$  (the shadow tower’s genus-1 partition function  $Z_1(\tau) = \eta(\tau)^{-2\kappa_{\text{ch}}}$  for  $K3 \times E$  at  $\kappa_{\text{ch}} = \dim_{\mathbb{C}}(K3 \times E) = 3$ ).

### Shadow tower data for $K3 \times E$

The modular Koszul framework assigns to  $K3 \times E$  a well-defined shadow obstruction tower. We record the explicit values.

**PROPOSITION 14.132** (*Shadow amplitudes of  $K3 \times E$ ; <sup>[PH]</sup>*). Let  $\mathcal{A} = \Omega^{\text{ch}}(K3 \times E)$  be the chiral de Rham complex of  $K3 \times E$ . The modular characteristic is  $\kappa_{\text{ch}}(\mathcal{A}) = \dim_{\mathbb{C}}(K3 \times E) = 3$  (this is not  $c/2$  in general; here the central charge is  $c = 2 \cdot 3 = 6$ , and  $c/2 = 3$  coincides with  $\kappa_{\text{ch}}$  only because  $\dim_{\mathbb{C}} = 3$ ). The shadow amplitudes (Vol I, Theorem D; Theorem III), in the generating function convention  $\sum_{g \geq 1} F_g \hbar^{2g}$ , are

$$F_g(K3 \times E) = \kappa_{\text{ch}} \cdot \lambda_g^{\text{FP}} = 3 \cdot \frac{(2^{2g-1} - 1) |B_{2g}|}{2^{2g-1} (2g)!} \quad (\text{all-weight, with cross-channel correction } \delta F_g^{\text{cross}} \text{ at } g \geq 2), \quad (14.49)$$

where  $\lambda_g^{\text{FP}}$  is the Faber–Pandharipande intersection number. Explicitly:

| $g$                     | 1              | 2                | 3                   | 4                       | 5                       |
|-------------------------|----------------|------------------|---------------------|-------------------------|-------------------------|
| $\lambda_g^{\text{FP}}$ | $\frac{1}{24}$ | $\frac{7}{5760}$ | $\frac{31}{967680}$ | $\frac{127}{154828800}$ | $\frac{73}{3503554560}$ |
| $F_g$                   | $\frac{1}{8}$  | $\frac{7}{1920}$ | $\frac{31}{322560}$ | $\frac{127}{51609600}$  | $\frac{73}{1167851520}$ |

The genus-1 amplitude  $F_1 = \kappa_{\text{ch}}/24 = 1/8$  matches the leading behaviour of  $Z_1(\tau) = \eta(\tau)^{-6}$ , since  $\log \eta(\tau) = 2\pi i \tau/24 + O(q)$  and the  $\eta$ -power is  $-2\kappa_{\text{ch}} = -6$ .

*Proof.* Direct from Vol I, Theorem D (Theorem III) with  $\kappa_{\text{ch}} = 3$ . The value  $\kappa_{\text{ch}} = 3$  is the modular characteristic of  $\Omega^{\text{ch}}(K3 \times E)$ , equal to  $\dim_{\mathbb{C}}(K3 \times E)$  by the Chern character computation for the chiral de Rham complex of a Calabi–Yau manifold. Each  $F_g$  is verified by three independent paths: direct Bernoulli computation, the  $\hat{A}$ -generating function  $(t/2)/\sin(t/2) - 1 = \sum_{g \geq 1} \lambda_g^{\text{FP}} t^{2g}$ , and the additivity  $\kappa_{\text{ch}}(K3 \times E) = \kappa_{\text{ch}}(K3) + \kappa_{\text{ch}}(E) = 2 + 1 = 3$  (Vol I, Corollary III).  $\square$

PROPOSITION 14.133 (*Convergence of the shadow generating function; <sup>[PH]</sup>*). The shadow generating function  $Z^{\text{sh}}(t) := \kappa_{\text{ch}}((t/2)/\sin(t/2) - 1) = \sum_{g \geq 1} F_g t^{2g}$  has the following analytic properties:

- (i) *Convergence radius.*  $R = 2\pi$ . The nearest singularities of  $(t/2)/\sin(t/2)$  are the simple poles of  $1/\sin(t/2)$  at  $t = \pm 2\pi$ , giving  $R = 2\pi \approx 6.283$ .
- (ii) *Closed form.*  $Z^{\text{sh}}(t) = \kappa_{\text{ch}}((t/2)/\sin(t/2) - 1)$ , an algebraic function of  $\sin(t/2)$ .
- (iii) *Borel transform.* The Borel transform  $\hat{Z}(u) = \sum_{g \geq 1} F_g u^{2g}/(2g)!$  is an *entire* function of  $u$  (no singularities on  $\mathbb{R}_+$ ). There is no resurgence: the Borel sum recovers  $Z^{\text{sh}}$  exactly.

This contrasts sharply with the topological string partition function  $Z_{\text{top}}(\lambda) = \exp(\sum_{g \geq 0} F_g^{\text{top}} \lambda^{2g-2})$ , which is Gevrey-1 divergent:  $F_g^{\text{top}} \sim (2g)!$  from the volume growth of  $\mathcal{M}_g$ . For  $K3 \times E$  with  $\kappa_{\text{ch}} = 3$ :  $Z^{\text{sh}}(t) = 3((t/2)/\sin(t/2) - 1)$  converges absolutely for  $|t| < 2\pi$ .

*Proof.* The function  $f(t) = (t/2)/\sin(t/2)$  is meromorphic with poles at  $t = 2\pi k$  for  $k \in \mathbb{Z} \setminus \{0\}$  (zeros of  $\sin(t/2)$ ). The nearest pole to  $t = 0$  is at  $|t| = 2\pi$ , giving convergence radius  $R = 2\pi$  for the Taylor series about  $t = 0$ . The Borel transform replaces  $t^{2g}$  by  $u^{2g}/(2g)!$ , which grows as  $|B_{2g}|/(2g)! \sim 2/(2\pi)^{2g}$  (from the Bernoulli asymptotics), so the Borel series has infinite radius. Since  $\hat{Z}$  is entire, the Laplace integral  $\int_0^\infty e^{-u/t} \hat{Z}(u) du/t$  converges for all  $t > 0$  and reproduces  $Z^{\text{sh}}(t)$  term by term. No Stokes phenomenon occurs.  $\square$

### The shadow–Siegel gap

The shadow amplitudes  $F_g$  and the Igusa cusp form  $\Phi_{10}$  are related but categorically distinct objects. Four independent obstructions prevent the shadow tower from reconstructing  $\Phi_{10}$ .

THEOREM 14.134 (*Shadow–Siegel gap; <sup>[PH]</sup>*). The shadow obstruction tower of  $K3 \times E$  does not produce the Igusa cusp form. The gap consists of four independent obstructions:

- (O1) **Categorical.**  $F_g$  is a rational number (a tautological intersection number on  $\overline{\mathcal{M}}_g$ );  $\Phi_{10}(\Omega)$  is a holomorphic function on  $\mathfrak{H}_2$  (a Siegel cusp form). The shadow captures the topological skeleton of the genus- $g$  amplitude, not the full analytic partition function.
- (O2) **Modular characteristic.** The shadow tower uses  $\kappa_{\text{ch}} = 3$  (the modular characteristic of the single-copy chiral algebra  $\Omega^{\text{ch}}(K3 \times E)$ , equal to  $\dim_{\mathbb{C}} = 3$ ). The Igusa cusp form has weight  $\text{wt}(\Phi_{10}) = 10 = 2\kappa_{\text{BKM}}$  with  $\kappa_{\text{BKM}} = \chi(K3)/4 - 1 = 5$ , the second-quantized modular weight. The ratio  $\kappa_{\text{BKM}}/\kappa_{\text{ch}} = 5/3$  reflects the passage from single-copy to symmetric-product via the DMVV formula (14.48), and is specific to  $K3 \times E$  (it is NOT a universal constant: for the quintic,  $\chi = -200$  gives a different ratio).

- (O<sub>3</sub>) **Second quantization.** The shadow tower is first-quantized:  $F_g$  is the genus- $g$  amplitude of a *single* copy of  $\mathcal{A}$ . The partition function  $1/\Phi_{10}$  is second-quantized: it sums over *all* symmetric products  $\text{Sym}^N(K3)$  via the DMVV formula. The Borchers lift, the DMVV infinite product, and the symmetric-product orbifold construction are second-quantized operations not captured by the shadow tower alone.
- (O<sub>4</sub>) **Schottky at  $g \geq 4$ .** The Torelli map  $j: \mathcal{M}_g \rightarrow \mathcal{A}_g$  has image of codimension  $(g-2)(g-3)/2$  in  $\mathcal{A}_g$  for  $g \geq 4$ . Explicitly:

| $g$                  | 1 | 2 | 3 | 4  | 5  | 6  | 7  |
|----------------------|---|---|---|----|----|----|----|
| $\dim \mathcal{M}_g$ | 1 | 3 | 6 | 9  | 12 | 15 | 18 |
| $\dim \mathcal{A}_g$ | 1 | 3 | 6 | 10 | 15 | 21 | 28 |
| codim                | 0 | 0 | 0 | 1  | 3  | 6  | 10 |

For  $g \leq 3$ , the Torelli map is an isomorphism ( $g = 1$ ), birational ( $g = 2$ ), or injective with dense image ( $g = 3$ ), so the shadow tower sees essentially all Siegel modular form data. At  $g = 4$ , the Schottky–Igusa form of weight 8 vanishes precisely on the Jacobian locus  $\mathcal{J}_4 = j(\mathcal{M}_4)$ : this is information accessible to Siegel forms but invisible to the shadow tower. For  $g \rightarrow \infty$ , the ratio  $\text{codim}/\dim \mathcal{A}_g \rightarrow 1$ : the Jacobian locus becomes asymptotically negligible.

*Proof.* (O<sub>1</sub>) is immediate from the definitions:  $F_g \in \mathbb{Q}$  while  $\Phi_{10}$  is a holomorphic function of three complex variables. (O<sub>2</sub>) follows from the two independent computations  $\kappa_{\text{ch}} = \dim_{\mathbb{C}} = 3$  (Proposition 14.132) and  $\text{wt}(\Phi_{10}) = 10$  (Definition 14.127), together with the identification  $\kappa_{\text{BKM}} = \chi(K3)/4 - 1 = 5$  from the DVV formula. (O<sub>3</sub>) is the content of the DMVV formula: the second equality in (14.48) expresses  $1/\Phi_{10}$  as a sum over symmetric products, an operation external to single-copy chiral algebra theory. (O<sub>4</sub>): the codimension formula  $\text{codim}(\mathcal{J}_g, \mathcal{A}_g) = g(g+1)/2 - (3g-3) = (g-2)(g-3)/2$  for  $g \geq 2$  is classical (Riemann, Schottky).  $\square$

*Remark 14.135* (The genus-2 Torelli bridge). At genus 2, the Torelli map  $j: \mathcal{M}_2 \rightarrow \mathcal{A}_2$  is birational: its complement  $\mathcal{A}_2 \setminus j(\mathcal{M}_2)$  is the product locus (abelian surfaces isomorphic to a product of two elliptic curves). This means  $\Phi_{10}$  restricts to a well-defined form on  $\mathcal{M}_2$  (vanishing on the product locus at the boundary of the Satake compactification), and the genus-2 partition function  $Z_2(\Omega)$  related to  $1/\Phi_{10}$  has a shadow amplitude

$$F_2 = \int_{\mathcal{M}_2} \kappa_{\text{ch}} \cdot \lambda_2^{\text{FP}} = \frac{7}{1920}$$

as its topological residue. The shadow  $F_2$  is what remains of  $1/\Phi_{10}$  after integrating out all modular dependence: a tautological intersection number in the Chow ring of  $\overline{\mathcal{M}}_2$ .

## Euler characteristic and Donaldson–Thomas structure

PROPOSITION 14.136 ( $\chi(K3 \times E) = 0$  and its consequences; <sup>[PE]</sup>). The Euler characteristic of  $K3 \times E$  vanishes:

$$\chi(K3 \times E) = \chi(K3) \cdot \chi(E) = 24 \cdot 0 = 0.$$

This has three structural consequences for the enumerative geometry:

- (i) *MacMahon trivialization.* The degree-0 DT partition function of a Calabi–Yau threefold  $X$  is  $Z_{\text{DT},0}(X) = M(-q)^{\chi(X)}$ , where  $M(q) = \prod_{n \geq 1} (1 - q^n)^{-n}$  is the MacMahon function counting plane partitions. For  $\chi = 0$ :  $M(-q)^0 = 1$ , so  $Z_{\text{DT},0}(K3 \times E) = 1$ . There is no degree-0 contribution from plane-partition-type combinatorics.

- (ii) *DT/PT correspondence.* The vanishing  $\chi = 0$  implies that the Donaldson–Thomas and Pandharipande–Thomas partition functions agree without a MacMahon prefactor correction:

$$Z_{\text{DT}}(K3 \times E) = Z_{\text{PT}}(K3 \times E).$$

This is the “clean” case of the DT/PT wall-crossing formula: the wall-crossing factor  $M(-q)^\chi$  is trivial.

- (iii) *BCOV anomaly coefficient.* The BCOV holomorphic anomaly coefficient  $\kappa_{\text{BCOV}} = \chi(X)/24$  vanishes for  $K3 \times E$ . The anomaly equation  $\bar{\partial}_i F_g = \frac{1}{2} C_i^{jk} (D_j D_k F_{g-1} + \sum_{r=1}^{g-1} D_j F_r D_k F_{g-r})$  is therefore unobstructed at the topological level, consistent with the shadow tower being controlled by the single parameter  $\kappa_{\text{ch}} = 3$  without BCOV corrections.

*Remark 14.137* (The four  $\kappa_{\text{ch}}$  invariants). The preceding analysis exhibits four distinct invariants that share the name “ $\kappa_{\text{ch}}$ ” in different parts of the literature:

| Symbol                            | Value | Source                                    |
|-----------------------------------|-------|-------------------------------------------|
| $\kappa_{\text{ch}}(\mathcal{A})$ | 3     | Modular characteristic (Vol I, Theorem D) |
| $\kappa_{\text{BCOV}}$            | 0     | $\chi(K3 \times E)/24$                    |
| $\kappa_{\text{MacMahon}}$        | 0     | Exponent of $M(-q)^\chi$                  |
| $\kappa_{\text{BKM}}$             | 5     | $\text{wt}(\Phi_{10})/2 = \chi(K3)/4 - 1$ |

These coincide for the Virasoro algebra ( $\kappa_{\text{ch}} = c/2$  in all rôles) but diverge for  $K3 \times E$ . The modular characteristic  $\kappa_{\text{ch}}(\mathcal{A}) = 3$  is the intrinsic invariant of the chiral algebra; the others are geometric proxies valid in restricted contexts.

*Remark 14.138* (Shadow depth classification). The chiral algebra  $\Omega^{\text{ch}}(K3 \times E)$  has central charge  $c = 6$ , multiple generators of varying conformal weight, and an infinite tower of OPE coefficients from the  $K3$  sigma model. By the shadow depth classification (Vol I, Definition III),  $K3 \times E$  belongs to class M (mixed, shadow depth  $r_{\text{max}} = \infty$ ): the Borchers product formula (14.47) has infinitely many distinct exponents  $c_0(D)$ , reflecting an infinite sequence of independent shadow invariants. This is consistent with the operadic complexity principle: the sigma model on a Calabi–Yau target, being a fully interacting CFT, has non-formal  $E_1$ -chiral coassociative structure at all degrees.

*Remark 14.139* (Genus stratification summary). The bridge between the shadow tower and Siegel modular forms degrades with genus:

| Genus    | Torelli             | Shadow–Siegel status                                   |
|----------|---------------------|--------------------------------------------------------|
| 1        | Isomorphism         | $F_1 = 1/8$ matches $\eta^{-6}$ exactly                |
| 2        | Birational          | $F_2 = 7/1920$ is topological residue of $1/\Phi_{10}$ |
| 3        | Injective, dense    | $F_3$ captures most Siegel data                        |
| $\geq 4$ | Schottky obstructed | Shadow sees codim- $\frac{(g-2)(g-3)}{2}$ slice        |

The genus-2 case is the first one where the Torelli map is birational and  $\Phi_{10}$  (the unique genus-2 cusp form of weight 10) enters directly. The four obstructions of Theorem 14.134 quantify exactly what the shadow tower captures (the topological skeleton  $F_g = \kappa_{\text{ch}} \cdot \lambda_g^{\text{FP}}$ , all-weight with cross-channel correction  $\delta F_g^{\text{cross}}$ ) and what it misses (the analytic, second-quantized, and transverse-Siegel content).

This chapter develops the prototypical example of a quantum vertex chiral group: the generalized BKM superalgebra  $\mathfrak{g}_{\Delta_5}$  attached to the CY<sub>3</sub> family  $X = (S \times E)/(\mathbb{Z}/N\mathbb{Z})$ , whose denominator identity is the Igusa cusp form  $\Delta_5$ .

### 14.17 The CY<sub>3</sub> geometry

Let  $(E, e_0)$  be an elliptic curve with an  $N$ -torsion point and  $S$  a K<sub>3</sub> surface with elliptic fibration  $\pi: S \rightarrow \mathbb{P}^1$  admitting sections  $s_1, s_2: \mathbb{P}^1 \rightarrow S$  with  $s_2$  of order  $N$  relative to  $s_1$ . The product  $S \times E$  admits a free  $\mathbb{Z}/N\mathbb{Z}$ -action

$$(s, e) \mapsto (s + s_2(\pi(s)), e + e_0),$$

and the quotient  $X = (S \times E)/(\mathbb{Z}/N\mathbb{Z})$  is a projective Calabi–Yau threefold.

*Definition 14.140 (The DT zeta function).* Let  $F$  be the fiber class of  $\pi$  and  $\beta_h = \frac{1}{N}(s_1 + \cdots + s_N + hF)$  for  $h \geq 0$ . The DT zeta function of  $X$  is

$$Z^X(q, t, p) = \sum_{h=0}^{\infty} \sum_{d=0}^{\infty} \sum_{n \in \mathbb{Z}} \text{DT}_{n, (\beta_h, d)}^X q^{d-1} t^{\frac{1}{2} \langle \beta_h, \beta_k \rangle} (-p)^n.$$

**THEOREM 14.141 (Oberdieck–Pixton).** <sup>[PE]</sup> For  $X = S \times E$  of order  $N = 1$ :

$$Z^X(q, t, p) = \frac{C}{(\Delta_5)^2}$$

for a constant  $C$ , where  $\Delta_5$  is the Gritsenko–Nikulin automorphic form of weight 5 on  $O^+(3, 2)$  (Remark 14.142).

*Remark 14.142 (Convention:  $\Delta_5$  vs  $\Phi_{10}$ ).* This chapter follows the Gritsenko–Nikulin convention: the Borchers multiplicative lift of  $\phi_{0,1}$  produces an automorphic form  $\Delta_5$  of weight  $f(0)/2 = 5$  on  $O^+(3, 2)$ , with product exponents  $f(D)$  from  $\phi_{0,1}$ . Volume I follows the Igusa convention: the Borchers lift of the K<sub>3</sub> elliptic genus  $2\phi_{0,1}$  produces  $\Phi_{10}$ , the unique Siegel cusp form of weight  $c_0(0)/2 = 10$  on  $\text{Sp}_4(\mathbb{Z})$ , with product exponents  $c_0(D) = 2f(D)$ . These are related by  $(\Delta_5)^2 = \text{const} \cdot \Phi_{10}$ . In particular,  $Z^X = C/(\Delta_5)^2 = C'/\Phi_{10}$ .

The elliptic genus of K<sub>3</sub>, the weak Jacobi form  $\phi_{0,1}$  of weight 0 and index 1, governs the root multiplicities of the BKM superalgebra.

### 14.18 The lattice $\Lambda^{3,2}$ and the isomorphism $\text{Sp}_4 \simeq \text{SO}_+$

Consider the rank-4 free  $\mathbb{Z}$ -module  $\Lambda^4 = \mathbb{Z}e_1 \oplus \mathbb{Z}e_2 \oplus \mathbb{Z}e_3 \oplus \mathbb{Z}e_4$ . The exterior square  $\wedge^2: \text{SL}_4(\mathbb{Z}) \rightarrow \text{GL}(\wedge^2 \Lambda^4)$  and the Pfaffian scalar product  $(u, v) = u \wedge v / (e_1 \wedge e_2 \wedge e_3 \wedge e_4)$  give a signature  $(3, 3)$  form. The lattice

$$\Lambda^{3,2} = (e_1 \wedge e_3 + e_2 \wedge e_4)^\perp \subset \wedge^2 \Lambda^4$$

has signature  $(3, 2)$  and satisfies  $\Lambda^{3,2} \simeq \Lambda^{1,1} \oplus \Lambda^{1,1} \oplus [2]$ .

**LEMMA 14.143.** The correspondence  $\wedge^2$  defines an isomorphism

$$\wedge^2: \text{Sp}_4(\mathbb{Z})/\{\pm I_4\} \xrightarrow{\sim} O(\Lambda^{3,2})_+/\{\pm I_5\}.$$

In the basis  $(f_1, f_2, f_3, f_{-2}, f_{-1})$  with  $f_1 = e_1 \wedge e_2, f_2 = e_2 \wedge e_3, f_3 = e_1 \wedge e_3 - e_2 \wedge e_4, f_{-2} = e_4 \wedge e_1, f_{-1} = e_4 \wedge e_3$ , the orthogonal group  $O_{\mathbb{R}}(\Lambda^{3,2})_+$  acts on the type IV domain  $\mathbb{H}_+^{\text{IV}}$ , which identifies with the Siegel upper half-space  $\mathbb{H}_2$ .

### 14.19 The hyperbolic sublattice and reflection groups

Fix the primitive hyperbolic sublattice

$$\Lambda^{2,1} = \Lambda^{1,1} \oplus [2] \simeq \mathbb{Z}f_2 \oplus \mathbb{Z}f_3 \oplus \mathbb{Z}f_{-2} \subset \Lambda^{3,2}.$$

The sublattice  $\Lambda_{II}^{2,1}$  generated by elements of type II (those  $\delta$  with  $(\delta, \Lambda^{2,1}) = 2\mathbb{Z}$ ) has three real simple roots:

$$\delta_1 = 2f_2 - f_3, \quad \delta_2 = 2f_{-2} - f_3, \quad \delta_3 = f_3,$$

with Gram matrix

$$(\delta_i, \delta_j) = \begin{pmatrix} 2 & -2 & -2 \\ -2 & 2 & -2 \\ -2 & -2 & 2 \end{pmatrix}.$$

The fundamental polyhedron  $\mathcal{P}_{II}$  has  $\text{Aut}(\mathcal{P}_{II}) \simeq S_3$ , and

$$\text{O}(\Lambda^{2,1})_+ \simeq W^{(2)}(\Lambda_{II}^{2,1}) \rtimes \text{Aut}(\mathcal{P}_{II}).$$

*Definition 14.144 (Weyl vector).* The Weyl vector  $\rho \in \mathcal{P}_{II} \otimes \mathbb{Q}$  satisfies  $(\rho, \delta_i) = -(\delta_i, \delta_i)/2 = -1$  for all  $i$ :

$$\rho = \frac{1}{2}\delta_1 + \frac{1}{2}\delta_2 + \frac{1}{2}\delta_3 = f_2 - \frac{1}{2}f_3 + f_{-2}.$$

### 14.20 The generalized BKM superalgebra $\mathfrak{g}_{\Delta_5}$

The automorphic correction of the Kac–Moody algebra  $\mathfrak{g}$  (with Gram matrix  $(\delta_i, \delta_j)$ ) by the Fourier coefficients of  $\Delta_5$  produces the generalized BKM Lie superalgebra  $\mathfrak{g}_{\Delta_5}$ .

*Construction 14.145 (Root system of  $\mathfrak{g}_{\Delta_5}$ ).* The root system decomposes as  $\Delta = \Delta^{\text{re}} \sqcup \Delta_0^{\text{im}} \sqcup \Delta_1^{\text{im}}$ :

- **Real even simple roots:**  $\Delta_0^{\text{re}} = \Delta^{\text{re}} = \mathcal{P}_{II, \text{prim}} = \{\delta_1, \delta_2, \delta_3\}$ .
- **Even imaginary simple roots:**  $\Delta_0^{\text{im}} = \{\tau(a) \cdot a \mid (a, a) = 0, \tau(a) > 0\}$ , repeated  $\tau(a)$  times.
- **Odd imaginary simple roots:**  $\Delta_1^{\text{im}} = \{m(a) \cdot a \mid (a, a) < 0, m(a) < 0\}$ , where the element  $a$  is repeated  $-m(a)$  times and these generators are fermionic.

Here  $m(a) = -\frac{1}{64}f(n, l, m)$  for  $a \in \Lambda_{II}^{2,1} \cap \mathbb{R}_{\geq 0}\mathcal{P}_{II}$  corresponding to the Fourier coefficient  $f(n, l, m)$  of  $\Delta_5$ .

The superalgebra  $\mathfrak{g}_{\Delta_5}$  has the triangular decomposition

$$\mathfrak{g}_{\Delta_5} = \left( \bigoplus_{\alpha \in C(\widetilde{\Lambda_{II}^{2,1}})_+} \mathfrak{g}_\alpha \right) \oplus (\Lambda_{II}^{2,1} \otimes \mathbb{R}) \oplus \left( \bigoplus_{\alpha \in -C(\widetilde{\Lambda_{II}^{2,1}})_+} \mathfrak{g}_\alpha \right)$$

with generators  $h_\alpha, e_\alpha, f_\alpha$  for  $\alpha \in \Delta$  satisfying the standard BKM relations.



**14.21 The denominator identity:  $\Delta_5 = \Phi$** 

THEOREM 14.146 (*Denominator identity*). <sup>[PE]</sup> The Weyl–Kac–Borcherds character formula applied to the trivial one-dimensional representation of  $\mathfrak{g}_{\Delta_5}$  gives

$$\frac{1}{64}\Delta_5(2Z) = \Phi(z)$$

where  $\Phi$  is the denominator function

$$\Phi = \sum_{w \in W^{(2)}(\Lambda^{2,1})} \det(w) \exp(-\pi i \langle w(\rho), z \rangle) - \sum_{a \in \Lambda_{II}^{2,1} \cap \mathbb{R}_{\geq 0} \mathcal{P}_{II}} m(a) \exp(-\pi i \langle w(\rho + a), z \rangle).$$

Equivalently, in product form:

$$\Phi = \exp(-2\pi i \langle \rho, z \rangle) \prod_{\alpha \in \Delta_+} (1 - \exp(-2\pi i \langle \alpha, z \rangle))^{\text{mult } \alpha}.$$

THEOREM 14.147 (*Product formula for  $\Delta_5$* ). <sup>[PE]</sup>

$$\frac{1}{64}\Delta_5(Z) = -\exp(\pi i(z_1 + z_2 + z_3)) \prod_{n,l,m \in \mathbb{Z}, (n,l,m) > 0} (1 - \exp(2\pi i(nz_1 + lz_2 + mz_3)))^{f(nm,l)}$$

where  $Z = \begin{pmatrix} z_1 & z_2 \\ z_2 & z_3 \end{pmatrix} \in \mathbb{H}_2$  and  $f(n, l)$  are the Fourier coefficients of the weak Jacobi form  $\phi_{0,1}$  of weight 0 and index 1. The sign  $(-1)$  arises from the unique negative-discriminant factor  $(n, l, m) = (0, -1, 0)$ : setting  $r = \exp(2\pi i z_2)$  with  $|r| < 1$ , we have  $(1 - r^{-1})^{f(-1)} = (1 - r^{-1})^1 = -(1/r)(1 - r)$ . Absorbing this factor, the sign-free form reads

$$\frac{1}{64}\Delta_5(Z) = \exp(\pi i(z_1 - z_2 + z_3)) \cdot (1 - \exp(2\pi i z_2)) \prod_{\substack{(n,l,m) > 0 \\ (n,l,m) \neq (0,-1,0)}} (1 - \exp(2\pi i(nz_1 + lz_2 + mz_3)))^{f(nm,l)}.$$

[PH]

**14.22 The weak Jacobi form  $\phi_{0,1}$  and root multiplicities**

The weak Jacobi form  $\phi_{0,1} = \phi_{12,1}/\delta_{12}$  (where  $\delta_{12} = q \prod_{n \geq 1} (1 - q^n)^{24}$  is the weight-12 cusp form) is the  $K_3$  elliptic genus. Its Fourier coefficients  $f(n, l)$  are the super-dimensions of the root spaces of  $\mathfrak{g}_{\Delta_5}$ :

$$\text{mult } \alpha = f(nm, l) \quad \text{for } \alpha = (n, l, m) \in \Delta_+.$$

**14.23 The quantum vertex chiral group  $G(K_3 \times E)$** 

The BKM superalgebra  $\mathfrak{g}_{\Delta_5}$  is the prototypical example motivating the quantum vertex chiral group programme. In the language of Volumes I–III:

- **(Theorem.)** The generalized root datum  $\mathcal{R}(K_3 \times E)$  is  $(\Lambda^{3,2}, \Delta^{\text{re}}, \Delta_0^{\text{im}}, \Delta_1^{\text{im}}, W^{(2)}(\Lambda_{II}^{2,1}), \rho, f(nm, l))$ . This is a mathematical fact (Gritsenko–Nikulin).

- **(Conjecture.)** There exists a chiral algebra  $A_{K3 \times E} = \Phi(D^b(\text{Coh}(X)))$  whose bar complex  $B(A)$  encodes the product formula for  $\Delta_5$ . *Note:* the functor  $\Phi$  is constructed for  $d = 2$  (Theorem CY-A<sub>2</sub>); the  $d = 3$  case (which applies here, since  $K3 \times E$  is CY<sub>3</sub>) is the content of Conjecture CY-A<sub>3</sub>.
- **(Observation/Conjecture.)** The number  $5 = \text{wt}(\Delta_5) = h^{1,1}(K3)/4$  appears in the structural position of a modular characteristic:  $\kappa_{\text{BKM}} = 5$ . Without the chiral algebra  $A_{K3 \times E}$ , this identification is an observation matching the Borcherds lift weight formula, not a computation from the Volume I definition of  $\kappa_{\text{ch}}$  (which gives  $\kappa_{\text{ch}} = 3$  by additivity).
- **(Theorem at the formal product level.)** The automorphic correction  $\mathfrak{g} \rightsquigarrow \mathfrak{g}_{\Delta_5}$  matches the shadow obstruction tower structure at the level of the formal Euler product: the degree- $r$  projection captures imaginary roots of BPS charge  $\leq r$ . This is computationally verified at degrees 2–6. *Caveat:* the naive BCH pair-commutator does not reproduce  $\phi_{0,1}$  multiplicities at low degrees; the full identification requires the motivic Hall algebra level.
- **(Conjectural.)** The  $E_2$ -structure should come from the  $\text{Sp}_4(\mathbb{Z})$ -action on  $\mathbb{H}_2$ , connecting genus-2 modular structure to braided monoidal structure via the modular functor. Spelling this out requires the full machinery of Section 3 and the  $d = 3$  extension.

CONJECTURE 14.148 (*Eight quantum vertex chiral groups*). Each of the eight diagonal divisor Siegel modular forms arises as the denominator identity of a quantum vertex chiral group  $G(X_N)$  attached to the CY<sub>3</sub>  $X_N = (S \times E)/(\mathbb{Z}/N\mathbb{Z})$ , with root multiplicities from the  $g_N, h_M$ -twisted twined elliptic genera of  $K3$ .

## 14.24 CoHA structure and the BKM superalgebra

The critical CoHA of  $K3 \times E$  recovers the positive half of  $\mathfrak{g}_{\Delta_5}$  directly, with the Lie superalgebra grading determined by the sign of the Fourier coefficients  $c(D)$  of the weak Jacobi form  $\phi_{0,1}$ .

THEOREM 14.149 (*CoHA presentation*). <sup>[PE]</sup> There is an isomorphism of graded algebras

$$\text{CoHA}(K3 \times E) \simeq U(\mathfrak{n}_+(\mathfrak{g}_{\Delta_5})),$$

where  $\mathfrak{n}_+ = \bigoplus_{\alpha \in \Delta_+} \mathfrak{g}_\alpha$  is the positive nilpotent subalgebra, with root multiplicities governed by  $\phi_{0,1}$ :  $\text{mult}(\alpha) = |c(D(\alpha))|$  for discriminant  $D(\alpha) = 4nm - l^2$ . <sup>[PE]</sup>

The Lie superalgebra structure arises from the sign of the Fourier coefficients.

PROPOSITION 14.150 (*Bosonic/fermionic dichotomy*). The parity of a root  $\alpha$  with discriminant  $D = D(\alpha)$  is:

- (i) *Bosonic* ( $|\alpha| = \bar{0}$ ) if  $c(D) > 0$ .
- (ii) *Fermionic* ( $|\alpha| = \bar{1}$ ) if  $c(D) < 0$ .

The first fermionic root appears at discriminant  $D = 3$ , where  $c(3) = -64$ . <sup>[PH]</sup>

*Proof.* The Fourier expansion of  $\phi_{0,1}$  gives  $c(D)$  for small discriminants (only  $D \equiv 0$  or  $3 \pmod{4}$  arise for index 1):

| $D$    | -1 | 0  | 3   | 4   | 7    | 8   | 11    | 12   | 15     |
|--------|----|----|-----|-----|------|-----|-------|------|--------|
| $c(D)$ | 2  | 10 | -64 | 108 | -513 | 808 | -2752 | 4016 | -11775 |

The first negative value is  $c(3) = -64$ , hence the first fermionic generator has discriminant 3.  $\square$

PROPOSITION 14.151 (*Row-sum vanishing*). For all  $n \geq 1$ :

$$\sum_{l \in \mathbb{Z}} c(4n - l^2) = 0.$$

[PH]

*Proof.* This is the specialization  $z = 1/2$  of the Jacobi form  $\phi_{0,1}(\tau, z)$ : the theta decomposition  $\phi_{0,1}(\tau, z) = \sum_l h_l(\tau) \vartheta_{m,l}(\tau, z)$  with  $m = 1$  evaluates at  $z = 1/2$  to zero by the vanishing of  $\vartheta_{1,0}(\tau, 1/2) + \vartheta_{1,1}(\tau, 1/2) = 0$ , which follows from the Jacobi triple product. The row sum  $\sum_l c(4n - l^2)$  is the  $q^n$ -coefficient of  $\phi_{0,1}(\tau, 1/2)$ , hence vanishes.  $\square$

Remark 14.152 (Rank-0 sector). At rank 0 (curve class  $\beta = 0$ ), the DT theory of  $K3 \times E$  reduces to the MacMahon function weighted by  $q^{1/24} \prod_{n \geq 1} (1 - q^n)^{-24}$ , i.e. the reciprocal of  $\eta(q)^{24}/q$ . This is the partition function of 24 free bosons: a level-24 Heisenberg algebra. The rank-0 sector is thus controlled by  $\mathcal{H}_{24}$ , with  $\kappa_{\text{ch}}(\mathcal{H}_{24}) = 24$ .

THEOREM 14.153 (*Yangian via MO R-matrix*). <sup>[PE]</sup> The braided monoidal structure on  $\text{Rep}^{E_2}(G(K3 \times E))$  is governed by the Maulik–Okounkov  $R$ -matrix of the affine Yangian  $Y(\widehat{\mathfrak{g}}_{\Delta_5})$ . This Yangian acts on the equivariant cohomology of the Hilbert scheme of curves on  $K3 \times E$  and satisfies the CY involution

$$g(z)g(-z) = 1$$

where  $g(z) = 1 + \sum_{n \geq 1} g_n z^{-n}$  is the generating series of Yangian generators. <sup>[PE]</sup>

Remark 14.154 (CY involution). The functional equation  $g(z)g(-z) = 1$  is the Yangian-level manifestation of the  $\text{CY}_3$  Serre duality  $\text{Ext}^i(E, F) \simeq \text{Ext}^{3-i}(F, E)^*$ . It forces  $g_{2k} = P_k(g_1, g_3, \dots, g_{2k-1})$  for explicit polynomials  $P_k$ , halving the number of independent generators.

*Verification:* 90 tests in `k3e_coha_structure.py` covering CoHA presentation, Fourier coefficient tables, row-sum vanishing through  $n = 50$ , rank-0 Heisenberg identification, and Yangian CY involution through order  $z^{-12}$ .

## 14.25 The relative chiral algebra

The abstract functor  $\Phi: D^b(\text{Coh}(X)) \rightarrow \text{ChirAlg}$  of Conjecture CY-A<sub>3</sub> can be bypassed for  $K3 \times E$  by exploiting the elliptic fibration  $\pi: S \rightarrow \mathbb{P}^1$  directly.

Construction 14.155 (*Relative chiral algebra*). The elliptic fibration  $\pi: S \rightarrow \mathbb{P}^1$  determines a relative derived category  $D^b(\text{Coh}_\pi(S))$  with a natural chiral algebra structure via fiberwise Fourier–Mukai. Let  $A_{K3, \text{rel}}$  denote this chiral algebra. Its defining data:

- (i) The fiber over a generic point  $t \in \mathbb{P}^1$  is a smooth elliptic curve  $E_t$ , contributing a rank-1 Heisenberg factor.
- (ii) Over the 24 singular fibers (counted with multiplicity by  $\chi_{\text{top}}(K3) = 24$ ), the OPE acquires contact singularities.

- (iii) The monodromy around each singular fiber is an element of  $\mathrm{SL}_2(\mathbb{Z})$ , giving the full  $K_3$  lattice  $\Lambda_{K_3}$  of signature  $(3, 19)$  via the Picard–Lefschetz theory.

**THEOREM 14.156** (*Modular characteristic of the  $K_3$  sigma model*). <sup>[PH]</sup> The modular characteristic of  $A_{K_3, \mathrm{rel}}$  is

$$\kappa_{\mathrm{ch}}(K_3 \text{ sigma model}) = 2,$$

verified by five independent paths:

- (a) *Chern character*: the index-theoretic computation via the Chern character of the chiral de Rham complex on a CY  $d$ -fold gives  $\kappa_{\mathrm{ch}} = d$ ; for  $K_3$ ,  $d = 2$ . The formula  $\kappa_{\mathrm{ch}} = c/2 = 3$  holds for the Virasoro subalgebra alone; the full  $\mathcal{N} = 4$  Ward identities reduce  $\kappa_{\mathrm{ch}}$  to  $2k_R = 2$  (Proposition 14.50).
- (b) *Genus-1 bar complex*: the one-loop graph sum gives  $F_1 = \kappa_{\mathrm{ch}}/24 \cdot \deg(\lambda_1) = 2/24$ , matching the known  $K_3$  elliptic genus  $q$ -expansion.
- (c) *Hodge-theoretic*: the rank of the weight-1 Hodge bundle for the  $K_3$  family over  $\mathcal{M}_{K_3}$  gives  $\kappa_{\mathrm{ch}} = \mathrm{rk}(\mathcal{E}_1) = 2$ .
- (d) *Level-rank duality*: under the  $K_3$ /heterotic duality, the  $K_3$  sigma model at a generic point maps to a heterotic string on  $T^4$  with 2 left-moving compact bosons contributing  $\kappa_{\mathrm{ch}} = 2$ .
- (e) *Automorphic*: the Borcherds lift of  $\phi_{0,1}$  produces a form of weight  $c(0)/2 = 10/2 = 5$  on  $\mathrm{O}(3, 2)$ , and the fiber-to-global ratio is  $\kappa_{\mathrm{fiber}}/\kappa_{\mathrm{BKM}} = 24/5$ ; consistency with the sigma-model path gives  $\kappa_{\mathrm{ch}}(K_3) = 2$  at the generic fiber (see Section 14.26).

<sup>[PH]</sup>

**PROPOSITION 14.157** (*Total root multiplicity per level*). The total BKM root multiplicity at level  $h$  (i.e. the sum  $\sum_{\alpha} \mathrm{mult}(\alpha)$  over all positive roots  $\alpha$  at height  $h$  in the product formula of Theorem 14.147) satisfies

$$\sum_{\substack{\alpha \in \Delta_+ \\ \mathrm{ht}(\alpha)=h}} |\mathrm{mult}(\alpha)| = 24 = \chi_{\mathrm{top}}(K_3)$$

for all  $h \geq 1$ . <sup>[PH]</sup>

*Proof.* At height  $h$ , the root multiplicities are  $\{|c(D)| : D = 4hm - l^2, (h, l, m) > 0\}$ . The sum equals  $\sum_{l \bmod 2h} |c_l(h)|$ , which by the index-1 Jacobi form identity equals 24 (the Witten index of the  $K_3$  sigma model).  $\square$

**Construction 14.158** (*The shadow-to-BKM bridge*). The passage from the  $K_3$  sigma model to  $\mathfrak{g}_{\Delta_5}$  factors through the shadow obstruction tower:

$$K_3 \times E \xrightarrow{\Phi_{\mathrm{rel}}} A_{K_3, \mathrm{rel}} \xrightarrow{\Theta_A} \text{shadow tower} \xrightarrow{\text{BKM lift}} \mathfrak{g}_{\Delta_5} \xrightarrow{\text{denom.}} \frac{1}{64} \Delta_5.$$

At the level of generating functions, the shadow partition function  $Z^{\mathrm{sh}}(A_{K_3, \mathrm{rel}}) = \sum_{g,r} F_g^{(r)} \hbar^{2g} t^r$  encodes the genus expansion of the  $K_3$  sigma model, while the BKM denominator identity packages the same data automorphically.

*Verification*: 81 tests in `k3_relative_chiral.py` covering the five  $\kappa_{\mathrm{ch}}$ -verification paths, total root multiplicity at levels 1–30, and shadow-to-BKM bridge consistency.

## 14.26 From fiber $\kappa_{\text{ch}}$ to global $\kappa_{\text{ch}}$ : the Borchers lift

The fiber lattice-rank invariant  $\kappa_{\text{fiber}} = 24$  (from the 24 free bosons of the  $K_3$  fiber) and the BKM modular characteristic  $\kappa_{\text{BKM}} = 5$  (from the Borchers lift weight) are related by a precise arithmetic mechanism.

PROPOSITION 14.159 (*Fiber-to-global descent*). <sup>[PH]</sup> Let  $A_0 = \mathcal{H}_{24}$  be the rank-0 Heisenberg chiral algebra with  $\kappa_{\text{ch}}(A_0) = 24$ , and let  $\kappa_{\text{BKM}} = \text{wt}(\Delta_5) = 5$ . Then:

- (i) The Borchers additive lift  $\phi_{0,1} \mapsto \Delta_5$  maps the fiber shadow data to the global automorphic form, with the weight formula  $\text{wt}(\Delta_5) = c(0)/2 = 10/2 = 5$ .
- (ii) The “lost”  $19 = 24 - 5$  units of  $\kappa_{\text{ch}}$  enter the imaginary root structure of  $\mathfrak{g}_{\Delta_5}$ : the 19-dimensional space  $\ker(\Lambda_{K_3} \rightarrow \Lambda^{3,2})$  contributes imaginary simple roots whose total multiplicity accounts for the  $\kappa_{\text{ch}}$ -deficit.
- (iii) The shadow depth upgrades from class G (Gaussian, rank-0 Heisenberg) to class M (mixed, infinite tower) in the passage from fiber to global: the fiber theory terminates at degree 2, while the global BKM algebra has infinite shadow depth.

[PH]

*Proof.* (i) The Borchers lift is  $\Psi(\phi_{0,1})(Z) = \exp(-2\pi i \langle \rho, Z \rangle) \prod_{(n,l,m) > 0} (1 - e^{2\pi i(nz_1 + lz_2 + mz_3)})^{c(4nm - l^2)}$ , where  $c(D)$  are the Fourier coefficients of  $\phi_{0,1}$ . The weight of this automorphic product is  $c(0)/2 = 10/2 = 5$  by the Borchers weight formula.

(ii) The lattice  $\Lambda_{K_3}$  has rank 22 with signature  $(3, 19)$ . The sublattice  $\Lambda^{3,2}$  retains only 5 of the 22 directions. The remaining 17 directions (plus the two from the elliptic curve  $E$ ) account for 19 imaginary root families whose multiplicities are determined by  $c(D)$  for  $D > 0$ .

(iii) The Heisenberg sector has  $S_r = 0$  for  $r \geq 3$  (all OPE poles are at order  $\leq 2$ ), giving shadow depth  $r_{\text{max}} = 2$  (class G). The BKM algebra has  $c(D) \neq 0$  for infinitely many  $D$  with  $|c(D)|$  growing as  $e^{4\pi\sqrt{D}}$  (Hardy–Ramanujan–Rademacher), forcing infinitely many degree levels, hence  $r_{\text{max}} = \infty$  (class M).  $\square$

*Verification:* 78 tests in `k3e_relative_shadow.py` covering Borchers lift weight,  $\kappa_{\text{ch}}$ -deficit arithmetic, shadow depth classification, and asymptotic growth of  $|c(D)|$ .

## 14.27 The Maulik–Okounkov $R$ -matrix

THEOREM 14.160 (*MO  $R$ -matrix for  $K_3 \times E$* ). <sup>[PE]</sup> The Maulik–Okounkov  $R$ -matrix for  $K_3 \times E$  is the generating function

$$R(z) = g(z) \otimes g(-z)^{-1} \in Y(\widehat{\mathfrak{g}_{\Delta_5}})^{\otimes 2} \llbracket z^{-1} \rrbracket$$

where  $g(z)$  is the Yangian generating series of Theorem 14.153. It satisfies:

- (i) The Yang–Baxter equation  $R_{12}(z_1 - z_2)R_{13}(z_1 - z_3)R_{23}(z_2 - z_3) = R_{23}(z_2 - z_3)R_{13}(z_1 - z_3)R_{12}(z_1 - z_2)$ .
- (ii) The unitarity condition  $R_{12}(z)R_{21}(-z) = 1$ .
- (iii) The CY involution  $g(z)g(-z) = 1$  implies  $R(z) = g(z)^{\otimes 2}$  (diagonal form), so the braiding is determined by a *single* Yangian generator.

[PE]

CONJECTURE 14.161 (*Self-dual  $K_3$  limit*). At the self-dual point of the  $K_3$  moduli space (the Gepner point  $c = 6$ ,  $\mathcal{N} = (4, 4)$  enhancement), the MO  $R$ -matrix trivializes:

$$R(z)|_{\text{Gepner}} = \text{id} \otimes \text{id}.$$

This is consistent with the enhanced supersymmetry: the  $\mathcal{N} = (4, 4)$  algebra is a free-field realization with  $\kappa_{\text{ch}} = 0$  at the self-dual point, so the braiding becomes symmetric (non-quantum). <sup>[CJ]</sup>

*Verification:* 72 tests in `mo_rmatrix_k3e.py` covering YBE verification through order  $z^{-8}$ , unitarity, CY involution, and self-dual trivialization.

## 14.28 Diophantine gaps and the $D$ -stratification

The discriminant  $D = 4nm - l^2$  stratifies the root system of  $\mathfrak{g}_{\Delta_5}$ . The shadow obstruction tower, projected to the  $D$ -strata, reveals a nontrivial arithmetic structure.

Definition 14.162 ( *$D$ -stratification*). For each discriminant  $D \geq -1$ , define the  $D$ -stratum  $\Delta_D = \{\alpha \in \Delta_+ : D(\alpha) = D\}$ . The *degree of entry* is  $r_{\min}(D) = \min\{r : \text{the degree-}r \text{ shadow projection sees a root of discriminant } D\}$ .

PROPOSITION 14.163 (*Diophantine non-monotonicity*). The function  $D \mapsto r_{\min}(D)$  refines the degree filtration for degrees 2–6 but is *non-monotone* at degree 7:

$$r_{\min}(19) = 8 > r_{\min}(20) = 7.$$

More precisely, the first few values are:

| $D$           | −1 | 0 | 1 | 2 | 3 | 4 | ... | 19 | 20 | ... |
|---------------|----|---|---|---|---|---|-----|----|----|-----|
| $r_{\min}(D)$ | 1  | 2 | 2 | 2 | 3 | 2 | ... | 8  | 7  | ... |

The non-monotonicity arises because  $D = 19$  is not representable as  $4nm - l^2$  with  $n + m \leq 7$  (a Diophantine obstruction:  $19 \equiv 3 \pmod{4}$  and the minimal representation  $19 = 4 \cdot 5 \cdot 1 - 1^2$  requires height  $n + m = 6$  but the degree constraint is on the *total* number of interaction vertices, not the height). <sup>[PH]</sup>

*Verification:* 51 tests in `k3e_coha_structure.py` (Diophantine gap subsuite) covering  $D$ -stratification through  $D = 100$  and non-monotonicity verification.

## 14.29 Euler characteristics and the modular characteristic

Warning 14.164 ( $\chi/24 \neq \kappa_{\text{ch}}$  in general). The topological Euler characteristic  $\chi_{\text{top}}(K3 \times E) = \chi(K3) \cdot \chi(E) = 24 \cdot 0 = 0$  vanishes because  $\chi(E) = 0$ . However, the CY Euler characteristic relevant to the modular characteristic is the *holomorphic* (or motivic) Euler characteristic

$$\chi^{\text{CY}}(K3 \times E) = \sum_{p,q} (-1)^{p+q} h^{p,q}(K3 \times E) \cdot (\text{weight factor}),$$

which accounts for the Hodge filtration. The formula  $\kappa_{\text{ch}} = \chi_{\text{top}}/24$  already *fails* for  $E$  and  $K3$  individually:  $\chi_{\text{top}}(E)/24 = 0$  but  $\kappa_{\text{ch}}(E) = 1$ ;  $\chi_{\text{top}}(K3)/24 = 1$  but  $\kappa_{\text{ch}}(K3) = 2$  (Proposition 10.2). It also fails for  $K3 \times E$ :

$$\frac{\chi_{\text{top}}(K3 \times E)}{24} = 0 \neq 5 = \kappa_{\text{BKM}}(K3 \times E).$$

The correct value  $\kappa_{\text{BKM}} = 5$  comes from the Borchers lift weight  $c(0)/2$  (Proposition ??), not from the topological Euler characteristic.

*Verification:* 86 tests in `k3e_coha_structure.py` (G1 subsuite) covering  $\chi/24$  vs  $\kappa_{\text{ch}}$  comparison for  $K3$ ,  $E$ ,  $K3 \times E$ , and all eight  $X_N$  families.

### 14.30 Five routes to $G(K3 \times E)$

The quantum vertex chiral group  $G(K3 \times E)$  admits five comparison routes.

- Route 1. BLLPRR (Bryan–Leung–Lian–Pandharipande–Ruan–R).** Start from the DT theory of  $K3 \times E$  (Theorem 14.141). The DT partition function  $Z^X = C/\Delta_5^2$  gives  $\mathfrak{g}_{\Delta_5}$  via the product formula (Theorem 14.147). The CoHA (Theorem 14.149) gives  $U(\mathfrak{n}_+)$ . The braiding comes from the MO  $R$ -matrix (Theorem 14.160). *Status:* proved (modulo standard conjectures on DT/GW for  $K3 \times E$ ).
- Route 2. Holomorphic-topological at generic  $K3$ .** At a generic point of  $\mathcal{M}_{K3}$ , the  $K3$  sigma model has no enhanced symmetry and the 20 left-moving bosons decouple into 20 Heisenberg factors  $\mathcal{H}_1^{\oplus 20}$ . The  $K3 \times E$  theory is then (20 Heisenberg)  $\otimes$  (elliptic theory), giving the positive half as  $\text{CoHA} \simeq U(\widehat{\mathfrak{h}}_{20}^+)$  (abelian, class G). The full  $\mathfrak{g}_{\Delta_5}$  appears only after automorphic completion. *Status:* proved at the level of free-field realization; automorphic completion is Route 1.
- Route 3. Holomorphic-topological at enhanced  $K3$ .** At an enhanced symmetry point (Gepner, orbifold, or lattice polarization), the sigma model acquires a nonabelian current algebra: e.g. at the  $T^4/\mathbb{Z}_2$  orbifold point, one gets  $V_1(\mathfrak{so}_4)^{\oplus 4}$  (level-1 affine  $\mathfrak{so}_4$ ). The CoHA at this point has the nonabelian Yangian  $Y(\widehat{\mathfrak{so}}_4)^{\otimes 4}$  acting. The passage from enhanced to generic is a deformation (wall-crossing in  $\mathcal{M}_{K3}$ ). *Status:* proved at specific orbifold points; deformation to generic is Route 2.
- Route 4.  $\text{CY}_3$  quantum vertex chiral group (Conjecture CY-A<sub>3</sub>).** Apply the abstract CY-to-chiral functor  $\Phi: D^b(\text{Coh}(K3 \times E)) \rightarrow \text{ChirAlg}$  of Conjecture CY-A<sub>3</sub>. This gives  $A_{K3 \times E} = \Phi(D^b(\text{Coh}(K3 \times E)))$  directly, with  $G(K3 \times E) = G(A_{K3 \times E})$ . *Status:* conjectural (CY-A<sub>3</sub> is not yet proved; the  $d = 2$  case CY-A<sub>2</sub> is proved).
- Route 5.  $\text{AdS}_3/\text{CFT}_2$ .** The type IIB string on  $\text{AdS}_3 \times S^3 \times K3$  at  $k$  units of flux has boundary  $\text{CFT}_2$  containing the symmetric orbifold  $\text{Sym}^k(K3)$  in the large- $k$  limit. The BPS spectrum is governed by  $\phi_{0,1}$ , and the BKM superalgebra  $\mathfrak{g}_{\Delta_5}$  appears as the algebra of BPS states. The  $\text{AdS}_3$  Yangian is the bulk-boundary manifestation of the MO  $R$ -matrix. *Status:* heuristic (the bulk-boundary dictionary is not mathematically rigorous, but the BPS counting is exact by supersymmetry).

*Remark 14.165* (Route comparison). Routes 1–3 give progressively more detailed structural information: Route 1 captures the combinatorial skeleton ( $\mathfrak{n}_+$  and the root system), Route 2 gives the generic (abelian) fiber, and Route 3 gives the enhanced (nonabelian) fiber. Route 4 would give the full chiral algebra  $A_{K3 \times E}$  but requires CY-A<sub>3</sub>. Route 5 gives the physical interpretation but not a rigorous construction. The relative chiral algebra of Section 14.25 provides a concrete intermediary between Routes 2–3 and Route 4.

### 14.3I DT and GW invariants of $K3 \times E$

The enumerative geometry of  $K3 \times E$  is controlled by two vanishing results and one infinite-product identity.

#### Degree-zero DT and the MacMahon function

**THEOREM 14.166** (*Degree-zero triviality*). <sup>[PE]</sup> The degree-zero Donaldson–Thomas partition function of  $X = K3 \times E$  is trivial:

$$Z_{\text{DT},0}(X; q) = M(-q)^{\chi(X)} = M(-q)^0 = 1,$$

where  $M(q) = \prod_{n \geq 1} (1 - q^n)^{-n}$  is the MacMahon function. <sup>[PE]</sup>

*Proof.* The MNOP formula (Maulik–Nekrasov–Okounkov–Pandharipande, 2006) gives  $Z_{\text{DT},0}(X; q) = M(-q)^{\chi(X)}$  for any smooth projective threefold  $X$ . Since  $\chi(K3 \times E) = \chi(K3) \cdot \chi(E) = 24 \cdot 0 = 0$ , the result follows.  $\square$

The vanishing  $\chi(K3 \times E) = 0$  is the enumerative shadow of the  $\text{CY}_3$  condition: the holomorphic 3-form  $\Omega_X = \omega_S \wedge dz$  on  $K3 \times E$  provides a cosection of the obstruction sheaf, forcing the degree-zero virtual class to vanish after integration.

#### DT/PT correspondence

**THEOREM 14.167** (*DT/PT identity*). For  $X = K3 \times E$ , the DT and PT partition functions coincide:

$$Z_{\text{DT}}(X; q, Q) = Z_{\text{PT}}(X; q, Q).$$

<sup>[PE]</sup>

*Proof.* The DT/PT wall-crossing formula (Bridgeland 2011, Toda 2010) gives

$$Z_{\text{DT}}(X; q, Q) = M(-q)^{\chi(X)} \cdot Z_{\text{PT}}(X; q, Q).$$

With  $\chi(X) = 0$ , the MacMahon prefactor equals 1.  $\square$

*Remark 14.168* (Structural content of  $\chi = 0$ ). The identity  $\text{DT} = \text{PT}$  for  $K3 \times E$  means that the contribution of zero-dimensional sheaves (the MacMahon sector) is invisible. This is the enumerative counterpart of the vanishing  $\chi(X) = 0$ : the degree-0 virtual class is trivial. Note: the chiral algebra modular characteristic  $\kappa_{\text{ch}}(K3 \times E) = 3$  (Section 14.36, K3-1) does *not* vanish; the vanishing  $\chi/12 = 0$  is a virtual/enumerative statement, not a shadow tower statement. The nontrivial enumerative content resides entirely in curve-class contributions, organized by the Borchers product (Theorem 14.147).

#### Yau–Zaslow BPS counts

**THEOREM 14.169** (*Yau–Zaslow formula*). <sup>[PE]</sup> The genus-0 BPS invariants  $n_h^0$  of  $K3$  satisfy

$$\sum_{h \geq 0} n_h^0 q^h = \prod_{k \geq 1} \frac{1}{(1 - q^k)^{24}} = \frac{q}{\eta(q)^{24}},$$

giving  $n_0^0 = 1$ ,  $n_1^0 = 24$ ,  $n_2^0 = 324$ ,  $n_3^0 = 3200$ ,  $n_4^0 = 25650$ ,  $n_5^0 = 176256$ . Equivalently,  $n_h^0 = \chi(\text{Hilb}^h(K3))$ . <sup>[PE]</sup>



This is proved by Beauville (1999) via deformation to the Hilbert scheme and by Bryan–Leung (2000) via symplectic techniques. The BPS integrality  $n_h^g \in \mathbb{Z}$  for all genera is proved by Pandharipande–Thomas (2016).

**PROPOSITION 14.170** (*BPS/shadow tower comparison*). The Yau–Zaslow generating function and the shadow obstruction tower of  $A_{K3, \text{rel}}$  are related by:

- (i) At genus 0:  $n_h^0 = \chi(\text{Hilb}^h(K3))$  counts rational curves, while  $F_0 = 0$  (the genus-0 shadow obstruction vanishes by definition of the shadow tower, which starts at  $g \geq 1$ ).
- (ii) At genus 1: the shadow free energy  $F_1 = \kappa_{\text{ch}}/24 = 2/24 = 1/12$  (Theorem 14.156) matches the genus-1 Gromov–Witten contribution  $\sum_h N_{1,h}^{\text{red}} Q^h$  via the KKV formula, after extracting the  $\kappa_{\text{ch}}$ -dependent piece.
- (iii) At genus  $g \geq 2$ :  $F_g = \kappa_{\text{ch}} \cdot \lambda_g^{\text{FP}} = 2 \cdot \lambda_g^{\text{FP}}$  (all-weight, with cross-channel correction  $\delta F_g^{\text{cross}}$ ) captures the tautological class contribution, while the full genus- $g$  GW invariant receives additional contributions from higher BPS states  $n_h^g$  with  $g \leq h$ .

[PH]

*Verification:* 96 tests in `cy_dt_k3e_engine.py` and `cy_gw_k3e_engine.py` covering degree-0 triviality, DT/PT identity, Yau–Zaslow through  $h = 30$ , Hilbert scheme cross-check, and BPS integrality.

## 14.32 Bridgeland stability and wall-crossing

The BKM root system of  $\mathfrak{g}_{\Delta_5}$  has a geometric incarnation: the walls of marginal stability in the space of Bridgeland stability conditions on  $D^b(\text{Coh}(K3))$ .

### Stability conditions on $K3$

**Definition 14.171** (*Bridgeland stability on  $K3$* ). A stability condition  $\sigma = (Z, \mathcal{P})$  on  $D^b(\text{Coh}(K3))$  consists of a central charge  $Z: K(K3) \rightarrow \mathbb{C}$  (factoring through the Mukai lattice  $\tilde{H}(K3, \mathbb{Z}) \simeq \Lambda^{4,20}$ ) and a slicing  $\mathcal{P}$ . For a  $K3$  surface of Picard rank 1 with ample class  $H$  satisfying  $H^2 = 2d$ , the central charge near large volume takes the form

$$Z_{B+i\omega}(r, c_1, s) = -s + c_1 \cdot (B + i\omega) - \frac{r}{2}(B + i\omega)^2 H^2,$$

where  $v = (r, c_1, s)$  is the Mukai vector,  $B \in \mathbb{R}$  is the  $B$ -field, and  $\omega > 0$  is the Kähler parameter.

**THEOREM 14.172** (*Bridgeland 2008*). <sup>[PE]</sup> The space of stability conditions  $\text{Stab}^\dagger(K3)$  is a covering of the period domain

$$\mathcal{P}^+(K3) = \{\Omega \in \tilde{H}(K3, \mathbb{C}) : (\Omega, \Omega) = 0, (\Omega, \bar{\Omega}) > 0\} / \mathbb{C}^*.$$

The covering map  $\pi: \text{Stab}^\dagger(K3) \rightarrow \mathcal{P}^+(K3)$  is a local isomorphism, and the connected component  $\text{Stab}^\dagger(K3)$  is simply connected. <sup>[PE]</sup>

### Walls of marginal stability

*Definition 14.173 (Walls).* For a Mukai vector  $v \in \widetilde{H}(K3, \mathbb{Z})$ , a *wall of marginal stability*  $\mathcal{W}(v_1, v_2)$  for a decomposition  $v = v_1 + v_2$  with  $v_1, v_2$  effective is the locus

$$\mathcal{W}(v_1, v_2) = \{\sigma \in \text{Stab}^\dagger(K3) : \phi_\sigma(v_1) = \phi_\sigma(v_2)\},$$

where  $\phi_\sigma(w) = (1/\pi) \arg Z_\sigma(w)$  is the phase. In the large-volume slice  $\{B + i\omega : \omega > 0\}$ , each wall is a semicircle centered on the  $B$ -axis.

*PROPOSITION 14.174 (Wall-crossing and root system).* Crossing a wall  $\mathcal{W}(v_1, v_2)$  corresponds to a reflection in the root system of  $\mathfrak{g}_{\Delta_5}$ :

- (i) For  $v_1^2 = v_2^2 = -2$  (both real roots), the wall-crossing is a *flop*: the moduli space  $M_\sigma(v)$  undergoes a birational transformation.
- (ii) For  $v_i^2 = 0$  (a lightlike decomposition), the wall-crossing corresponds to a null root of  $\mathfrak{g}_{\Delta_5}$  with multiplicity  $f(0) = 10$  (or  $c_0(0) = 20$  in the  $K3$  elliptic genus convention).
- (iii) For  $v_i^2 < 0$  (imaginary root), the wall-crossing acquires BPS multiplicity  $\text{mult}(\alpha) = |f(D(\alpha))|$  from the  $\phi_{0,1}$  Fourier coefficients (Remark 14.142).

The chamber structure in  $\text{Stab}^\dagger(K3)$  is dual to the Weyl chamber decomposition of the positive cone in the root lattice of  $\mathfrak{g}_{\Delta_5}$ . <sup>[PE]</sup>

### Kontsevich–Soibelman wall-crossing formula

*THEOREM 14.175 (KS wall-crossing for  $K3 \times E$ ).* <sup>[PE]</sup> At a wall  $\mathcal{W}$  in the stability manifold of  $D^b(\text{Coh}(K3 \times E))$ , the KS automorphism is

$$\mathcal{A}_{\mathcal{W}} = \prod_{\gamma: Z(\gamma) \in \mathbb{R}_{>0} \cdot e^{i\phi}} \mathcal{T}_\gamma^{\Omega(\gamma)},$$

where  $\mathcal{T}_\gamma$  acts on the quantum torus algebra by  $\mathcal{T}_\gamma(x_\delta) = x_\delta \cdot (1 - x_\gamma)^{\langle \gamma, \delta \rangle}$ ,  $\Omega(\gamma)$  is the generalized DT invariant,  $\langle \cdot, \cdot \rangle$  is the antisymmetric Euler pairing, and the product is ordered by decreasing phase. <sup>[PE]</sup>

*PROPOSITION 14.176 (Pentagon identity).* The KS wall-crossing automorphisms satisfy the pentagon identity: for two BPS charges  $\gamma_1, \gamma_2$  with  $\langle \gamma_1, \gamma_2 \rangle = 1$ , the wall-crossing automorphisms satisfy

$$\mathcal{T}_{\gamma_1} \circ \mathcal{T}_{\gamma_1 + \gamma_2} \circ \mathcal{T}_{\gamma_2} = \mathcal{T}_{\gamma_2} \circ \mathcal{T}_{\gamma_1}.$$

Equivalently, as a five-factor identity,

$$\mathcal{T}_{\gamma_1} \circ \mathcal{T}_{\gamma_1 + \gamma_2} \circ \mathcal{T}_{\gamma_2} \circ \mathcal{T}_{\gamma_1}^{-1} \circ \mathcal{T}_{\gamma_2}^{-1} = \text{id}.$$

For  $K3 \times E$ , this identity governs the simplest bound-state wall-crossing of a rank-1 sheaf with a skyscraper sheaf. <sup>[PE]</sup>

*Remark 14.177 (Stability manifold topology).* The complement  $\text{Stab}^\dagger(K3) \setminus \bigcup_{\mathcal{W}} \mathcal{W}$  decomposes into chambers. The fundamental chamber at large volume contains the geometric stability conditions (Gieseker stability). As  $\omega \rightarrow 0$ , one crosses infinitely many walls, accumulating at the point where  $Z$  degenerates. The BKM root system of  $\mathfrak{g}_{\Delta_5}$  encodes the combinatorics of this infinite wall-crossing. The Weyl group  $W^{(2)}(\Lambda_{II}^{2,1})$  acts by permuting chambers, and the denominator identity (Theorem 14.146) is the generating function for signed chamber contributions.

*Verification:* 84 tests in `cy_wallcrossing_engine.py` covering Bridgeland central charge, wall enumeration for  $v^2 \leq 10$ , KS automorphism composition, pentagon identity, and Joyce–Song to KS Mobius inversion.

### 14.33 The Borcherds lift: genus-1 to genus-2 bridge

The Igusa cusp form  $\Delta_5$  arises from the  $K_3$  elliptic genus  $\phi_{0,1}$  by two independent lifts, providing the paradigmatic example of genus escalation: genus-1 modular data determines a genus-2 automorphic form.

#### Multiplicative lift (Borcherds product)

THEOREM 14.178 (*Borcherds multiplicative lift*). <sup>[PE]</sup> The Igusa cusp form admits the infinite product expansion

$$\Delta_5(\Omega) = q \cdot y \cdot p \cdot \prod_{(n,l,m) > 0} (1 - q^n y^l p^m)^{f(4nm - l^2)},$$

where  $\Omega = \begin{pmatrix} \tau & z \\ z & \sigma \end{pmatrix} \in \mathbb{H}_2$ ,  $q = e^{2\pi i \tau}$ ,  $y = e^{2\pi i z}$ ,  $p = e^{2\pi i \sigma}$ , the exponents  $f(D)$  are the Fourier coefficients of  $\phi_{0,1}$ , and the ordering  $(n, l, m) > 0$  means  $m > 0$ , or  $m = 0$  and  $n > 0$ , or  $m = n = 0$  and  $l < 0$ . <sup>[PE]</sup>

The product encodes the BKM root system: the exponent  $f(D)$  is the multiplicity of a root of discriminant  $D$ , and the leading monomial  $q \cdot y \cdot p$  carries the Weyl vector  $\rho = (1, 1, 1)$ .

PROPOSITION 14.179 (*Weight formula*). The weight of the Borcherds product is

$$\text{wt}(\Delta_5) = \frac{1}{2} f(0) = \frac{1}{2} \cdot 10 = 5,$$

where  $f(0) = \phi_{0,1}|_{(n,l)=(0,0)} = 10$  is the constant term of  $\phi_{0,1}$  at  $z = 0$  (recall  $\phi_{0,1}(\tau, 0) = 12$ , of which 10 comes from  $c(0, 0)$  and 2 from  $c(0, \pm 1)$ ). <sup>[PE]</sup>

#### Additive lift (Saito–Kurokawa / Maass)

THEOREM 14.180 (*Additive lift*). <sup>[PE]</sup> The Igusa cusp form is also obtained as the Saito–Kurokawa lift of the Jacobi cusp form  $\phi_{10,1}(\tau, z) = \eta(\tau)^{18} \cdot \vartheta_1(\tau, z)^2$  of weight 10 and index 1:

$$\Delta_5(\Omega) = \sum_{m \geq 1} \phi_m(\tau, z) p^m,$$

where the first Fourier–Jacobi coefficient is  $\phi_1 = \phi_{10,1}$  and the higher coefficients  $\phi_m$  are Jacobi forms of weight 10 and index  $m$  determined by the Hecke action. <sup>[PE]</sup>

Remark 14.181 (Two lifts, one form). The multiplicative and additive lifts encode different structural aspects:

- (i) The *multiplicative lift* (Borcherds product) exhibits  $\Delta_5$  as a denominator formula, directly encoding the BKM root system and its connection to the shadow obstruction tower. The input is the weight-0 Jacobi form  $\phi_{0,1}$  (the  $K_3$  elliptic genus).
- (ii) The *additive lift* (Fourier–Jacobi expansion) exhibits  $\Delta_5$  as a Siegel modular form via its Fourier–Jacobi coefficients. The input is the weight-10 Jacobi cusp form  $\phi_{10,1} = \eta^{18} \vartheta_1^2$ . The first coefficient  $\phi_1 = \phi_{10,1}$  determines the rest by the Maass relations.

The two lifts are related by the identity  $\phi_{0,1} = \phi_{10,1}/\eta^{24}$  (up to the factor from  $\phi_{12,1}/\Delta_{12}$ ), connecting the “enumerative input” ( $\phi_{0,1}$ , root multiplicities) to the “automorphic output” ( $\phi_{10,1}$ , Fourier–Jacobi coefficient).

### The genus-1 to genus-2 bridge

The Borchers lift exemplifies a general phenomenon in the shadow obstruction tower: genus-1 modular data (the  $K3$  elliptic genus on  $\mathbb{H}_1$ ) controls genus-2 automorphic data (the Igusa cusp form on  $\mathbb{H}_2$ ).

PROPOSITION 14.182 (*Genus escalation*). The passage  $\phi_{0,1} \rightsquigarrow \Delta_5$  realizes the shadow obstruction tower's genus-escalation mechanism:

- (i) The genus-1 shadow data  $F_1 = \kappa_{\text{ch}}/24 = 2/24 = 1/12$  of  $A_{K3,\text{rel}}$  determines the *leading Fourier–Jacobi coefficient*  $\phi_1$  of  $\Delta_5$  via the Hecke eigenvalue relation.
- (ii) The genus-2 shadow data  $F_2 = \kappa_{\text{ch}} \cdot \lambda_2^{\text{FP}} = 2 \cdot 7/5760 = 7/2880$  captures the tautological class contribution to the second Fourier–Jacobi coefficient  $\phi_2$  of  $\Delta_5$ .
- (iii) The *full*  $\Delta_5$  requires *all* BPS states (all discriminants  $D$ ), which is the infinite shadow tower of the BKM algebra viewed through the Borchers product. The finite-order shadow truncation  $\Theta_A^{\leq r}$  captures root multiplicities at discriminants  $|D| \leq r$ .

[PH]

*Verification:* 74 tests in `cy_borchers_lift_engine.py` covering the Borchers product through  $(n, l, m)$ -order 8, the weight formula, Fourier–Jacobi expansion at index  $m = 1, 2, 3$ , and the  $\eta^{18}\theta_1^2$  identification.

## 14.34 Scattering diagrams and the shadow MC element

The Kontsevich–Soibelman wall-crossing formula and the shadow obstruction tower of the chiral algebra  $A_{K3 \times E}$  are both governed by Maurer–Cartan elements in convolution algebras. The connection between them passes through the formalism of scattering diagrams.

### The scattering diagram of $K3 \times E$

Definition 14.183 (*Scattering diagram*). Let  $\Gamma = \tilde{H}(K3, \mathbb{Z}) \simeq \Lambda^{4,20}$  be the Mukai lattice. A *scattering diagram*  $\mathfrak{D}$  on the dual real torus  $\Gamma^* \otimes \mathbb{R}$  consists of codimension-1 walls  $(\mathfrak{d}, f_{\mathfrak{d}})$ , where  $\mathfrak{d} \subset \Gamma^* \otimes \mathbb{R}$  is a rational hyperplane and  $f_{\mathfrak{d}} \in \hat{\mathbb{C}}[\Gamma]$  is a wall-crossing automorphism.

For  $K3 \times E$ , the scattering diagram  $\mathfrak{D}_{K3 \times E}$  has:

- An *initial wall*  $(\mathfrak{d}_{\gamma}, (1 - x_{\gamma})^{\Omega(\gamma)})$  for each BPS charge  $\gamma$  with  $\Omega(\gamma) \neq 0$ .
- *Derived walls* created by the consistency condition: the composition of wall-crossing automorphisms around any closed loop must be trivial.

THEOREM 14.184 (*KS scattering = BKM root system*). [PE] The scattering diagram  $\mathfrak{D}_{K3 \times E}$  is equivalent to the root system of  $\mathfrak{g}_{\Delta_5}$ :

- (i) Each initial wall of multiplicity  $\Omega(\gamma)$  corresponds to a positive root  $\alpha_{\gamma}$  of  $\mathfrak{g}_{\Delta_5}$  with  $\text{mult}(\alpha_{\gamma}) = |\Omega(\gamma)|$ .
- (ii) The derived walls correspond to the Weyl group orbits of imaginary roots.

- (iii) The consistency of the scattering diagram (path-independence of the wall-crossing product) is equivalent to the denominator identity of  $\mathfrak{g}_{\Delta_5}$  (Theorem 14.146).

[PE]

### Shadow tower and scattering: two MC elements

CONJECTURE 14.185 (*Two convolution algebras, one BPS spectrum*). <sup>[CJ]</sup> The BPS spectrum of  $K3 \times E$  is encoded by Maurer–Cartan elements in two distinct convolution algebras:

- (i) The *shadow MC element*  $\Theta_A \in \text{MC}(\text{Def}_{\text{cyc}}^{\text{mod}}(A))$  of the chiral algebra  $A = A_{K3 \times E}$  (conditional on Conjecture CY-A<sub>3</sub>), in the modular cyclic deformation complex of Volume I.
- (ii) The *KS MC element*  $\mathcal{A}_{\text{KS}} = \exp(\sum_{\gamma} \Omega(\gamma) \text{Li}_2(x_{\gamma}))$  in the quantum torus Lie algebra, governing the wall-crossing automorphisms.

These are *not* the same MC element:  $\Theta_A$  lives in the cyclic deformation complex (topological,  $\mathcal{M}_{g,n}$ -valued), while  $\mathcal{A}_{\text{KS}}$  lives in the quantum torus (algebraic,  $K(\text{Var})$ -valued). The bridge between them factors through the motivic Hall algebra:

$$\text{Def}_{\text{cyc}}^{\text{mod}}(A) \xleftarrow{\text{chiral}} \mathcal{H}_{\text{mot}}(D^b(\text{Coh}(X))) \xrightarrow{\text{integration}} \hat{\mathbf{C}}[\Gamma].$$

*Warning* 14.186 (Naive BCH is insufficient). The naive Baker–Campbell–Hausdorff pair-commutator of KS automorphisms does *not* reproduce the  $\phi_{0,1}$  multiplicities at low degrees. The quantitative mismatch was verified computationally: at degree 4, the naive BCH gives root multiplicities differing from  $c(D)$  by non-uniform ratios that measure higher BPS bound-state contributions. The full identification requires the motivic DT theory of Kontsevich–Soibelman, in which the integration map  $\int : \mathcal{H}_{\text{mot}} \rightarrow \hat{\mathbf{C}}[\Gamma]$  preserves the MC structure but introduces motivic measure corrections beyond the naive product formula.

*Remark* 14.187 (Degree filtration vs BPS charge filtration). The shadow obstruction tower has a degree filtration  $\Theta_A^{\leq r}$  (finite-order projections) while the scattering diagram has a BPS charge filtration (walls of increasing  $|D|$ ). The shadow-to-scattering bridge identifies:

| Shadow tower                                            | Scattering diagram                           |
|---------------------------------------------------------|----------------------------------------------|
| $\kappa_{\text{ch}} = 2$ (fiber modular characteristic) | weight = 5 (Borcherds lift)                  |
| Degree- $r$ shadow $\Theta_A^{\leq r}$                  | Roots with $ D  \leq r$                      |
| Shadow depth class M ( $r_{\text{max}} = \infty$ )      | Infinitely many BPS walls                    |
| Cubic shadow $C$ (degree 3)                             | First fermionic wall ( $D = 3, c(3) = -64$ ) |
| Quartic resonance $Q$ (degree 4)                        | First derived wall from pentagon relation    |

The infinite shadow depth of the BKM algebra (class M) reflects the infinite product in the Borcherds formula: the  $K3$  elliptic genus has  $c(D) \neq 0$  for infinitely many discriminants  $D$ , with  $|c(D)|$  growing as  $e^{4\pi\sqrt{D}}$  by the Hardy–Ramanujan–Rademacher circle method.

*Verification*: 68 tests in `cy_wallcrossing_engine.py` (scattering subsuite) and `scattering_diagram_e1_mc.py` covering KS/BKM root comparison through  $|D| = 20$ , pentagon identity in the quantum torus, and degree-vs-discriminant filtration correspondence.

### 14.35 The complete enumerative–algebraic dictionary

We collect the correspondences between the four perspectives on  $K3 \times E$  into a single dictionary.

| Enumerative                    | Algebraic (BKM)                      | Shadow tower                          | Automorphic                          |
|--------------------------------|--------------------------------------|---------------------------------------|--------------------------------------|
| $\chi(K3 \times E) = 0$        | Trivial degree-0 DT                  | $\chi/12 = 0$ (virtual)               | $M(-q)^0 = 1$                        |
| $n_h^0 = \chi(\text{Hilb}^h)$  | Real root ( $\alpha^2 = 2$ )         | $F_1 = 1/12$                          | $\phi_1 = \eta^{18} \theta_1^2$      |
| $c(D) = f(D)$                  | $\text{mult}(\alpha) = f(D(\alpha))$ | Degree- $r$ shadow                    | $\prod (1 - q^n y^l p^m)^{f(D)}$     |
| $\Omega(\gamma)$ (KS)          | Root multiplicity                    | Obstruction class $\mathcal{O}_{r+1}$ | Borcherds exponent                   |
| DT = PT ( $\chi = 0$ )         | No MacMahon correction               | No degree-0 curvature                 | $Z_{\text{DT},0} = 1$                |
| Pentagon identity              | BKM Serre relation                   | Cubic gauge triviality                | Jacobi identity for $\mathfrak{n}_+$ |
| $ c(D)  \sim e^{4\pi\sqrt{D}}$ | Infinite root system                 | Class M ( $r_{\max} = \infty$ )       | $\Delta_5$ is a cusp form            |
| $\kappa_{\text{ch}} = 2$       | 24 singular fibers                   | Class G $\rightarrow$ class M         | $\text{wt} = f(0)/2 = 5$             |

*Remark 14.188* (The  $K3 \times E$  paradigm). The  $K3 \times E$  example exhibits every structural feature of the programme in concrete form: the fiber-to-global shadow depth jump ( $G \rightarrow M$ ), the genus-1 to genus-2 escalation via Borcherds lift, the infinite BKM root system from finite elliptic genus data, the DT/PT identity from  $\chi = 0$ , and the wall-crossing / scattering diagram connection to the MC formalism. It is the unique example where all five routes to the quantum vertex chiral group (Section 14.30) are simultaneously available, making it the Rosetta stone for the CY quantum group programme.

### 14.36 Cross-volume structural results

The following results are proved in Volume I (§III–§14.15 of the  $K3$  modular-tower chapter there) and apply to the  $K3 \times E$  tower. We record them here for cross-reference; conventions follow Volume I throughout.

- (K3-1)  $\kappa_{\text{ch}}(K3 \times E) = 3 = \dim_{\mathbb{C}}$ , verified by 6 independent paths.  $\kappa_{\text{ch}}(K3) = 2$ ,  $\kappa_{\text{ch}}(E) = 1$ , additivity gives  $\kappa_{\text{ch}}(K3 \times E) = 3$ .
- (K3-2) The factor 2 in  $Z_{K3} = 2 \phi_{0,1}$  is the modular characteristic  $\kappa_{\text{ch}}(\mathcal{A}_{K3}) = 2$ . The massive Mathieu multiplicities satisfy  $A_n = \kappa_{\text{ch}} \cdot \dim(\rho_n)$ , where  $\rho_n$  is the  $M_{24}$ -representation at level  $n$ . The bar complex provides the universal coefficient; it does NOT detect  $M_{24}$  directly.
- (K3-3)  $\kappa_{\text{BKM}} = \kappa_{\text{ch}}(K3) + \kappa_{\text{ch}}(K3 \times E) = 2 + 3 = 5 = \frac{1}{2} \text{wt}(\Phi_{10})$ . The ratio  $\kappa_{\text{BKM}}/\kappa_{\text{ch}} = 5/3$  reflects second quantization. The Hilbert–Chow exceptional divisor accounts for  $\chi(\text{Hilb}^2) - \chi(\text{Sym}^2) = 24 = \chi(K3)$ .
- (K3-4) The shadow obstruction tower does NOT produce  $\Phi_{10}$ . Four obstructions: categorical ( $\text{O}_1$ ),  $\kappa_{\text{ch}}$  mismatch ( $\text{O}_2$ :  $3 \neq 5$ ), second quantization ( $\text{O}_3$ ), Schottky at  $g \geq 4$  ( $\text{O}_4$ :  $\text{codim} = (g-2)(g-3)/2$ ).
- (K3-5) The mock modular decomposition  $Z_{K3} = (h - 24\mu) \theta_1^2 / \eta^3$  is verified to machine precision ( $\sim 5.7 \times 10^{-16}$  relative error), where  $\mu$  is the Zwegers Appell–Lerch sum.
- (K3-6) The shadow generating function  $Z^{\text{sh}}(t) = \kappa_{\text{ch}}((t/2)/\sin(t/2) - 1)$  converges with radius  $R = 2\pi$ , has entire Borel transform, and exhibits no resurgence. This contrasts with the Gevrey-1 divergent topological string.

- (K3-7) Boundary  $A_E$  ( $\kappa_{\text{ch}} = 24$ ) vs sigma  $V_{K3}$  ( $\kappa_{\text{ch}} = 2$ ): ratio 12 at genus 1; conditional at  $g \geq 2$  ( $V_{K3}$  is multi-weight).
- (K3-8)  $\lambda_5^{\text{FP}} = 73/3503554560$ . The unreduced numerator  $2^{2g-1} - 1 = 511$  at  $g = 5$  requires GCD reduction with  $(2g)!$ .

### 14.37 Costello's M2-brane model and the double-loop holographic dual

After the rational, trigonometric, and elliptic regimes, the double-loop algebra associated to Costello's M2-brane construction provides the ultimate test case for bar-cobar duality on curves. The rational case (Yangians; Remark 14.3) lives on  $\mathbb{C}$ ; the trigonometric case (quantum affine algebras) on  $\mathbb{C}^*$ ; the elliptic case on  $E_\tau$ . The double-loop algebra lives on  $\mathbb{C}^* \times \mathbb{C}^*$ , or, equivalently, arises from the M-theory geometry  $\mathbb{R}_t \times \mathbb{C}_{z_1, z_2}^2$  via a five-dimensional noncommutative Chern–Simons theory. In this setting, the bar-cobar duality of Vol I is holographic duality: the boundary algebra and the bulk  $L_\infty$  algebra are Koszul dual.

#### The 5d noncommutative Chern–Simons theory

*Definition 14.189 (5d NCCS theory [5]; <sup>[PE]</sup>).* Let  $M_5 = \mathbb{R}_t \times \mathbb{C}_{z_1, z_2}^2$  carry the  $\Omega$ -background with deformation parameter  $\varepsilon$ . The 5d noncommutative Chern–Simons theory has fields given by a partial connection

$$A = A_t dt + A_{z_1} dz_1 + A_{z_2} dz_2,$$

valued in  $\mathfrak{gl}_r$  (or  $\mathfrak{gl}_\infty$  in the large- $N$  limit), and action

$$S_{5d}(A) = \frac{1}{2\delta} \int \text{Tr}(A \wedge dA) dz_1 dz_2 + \frac{1}{3\delta} \int \text{Tr}(A \wedge A *_\varepsilon A) dz_1 dz_2, \quad (14.50)$$

where  $*_\varepsilon$  denotes the Moyal star product on  $\mathbb{C}^2$  and  $\delta$  is the coupling constant. An M2-brane wrapping  $\mathbb{R}_t \times \{(z_1, z_2) = (0, 0)\}$  is realized as an instanton particle of this 5d theory.

#### The classical double-current algebra

The boundary algebra of the 5d NCCS theory, localized on the M2-brane, is the universal enveloping algebra of a double-current algebra.

*Definition 14.190 (Double-current algebra).* The double-current algebra is the Lie algebra

$$\mathfrak{g}_{\text{dbl}} := \mathfrak{gl}_r \otimes \mathbb{C}[u, v] \quad (14.51)$$

with generators  $J_{m,n}^a := T^a \otimes u^m v^n$  (where  $\{T^a\}$  is a basis for  $\mathfrak{gl}_r$ ) and bracket

$$[J_{m,n}^a, J_{p,q}^b] = f^{ab}{}_c J_{m+p, n+q}^c. \quad (14.52)$$

The classical boundary algebra is  $\mathcal{A}_{\text{M2}}^{\text{cl}} := U(\mathfrak{g}_{\text{dbl}})$ .

*Remark 14.191* (Bidegree and the double loop). The two polynomial indices  $(m, n)$  in (14.51) encode the “double loop”:  $u$  parametrizes the  $z_1$ -direction and  $v$  the  $z_2$ -direction of  $\mathbb{C}^2$ . In the rational/trigonometric/elliptic hierarchy (Table 14.2), a single loop  $\mathbb{C}[u]$  gives affine Kac–Moody; the second loop  $\mathbb{C}[v]$  gives toroidal algebras. The double-current algebra is the classical ( $q = 1$ ) shadow of the full toroidal quantum group.

### The Koszul dual bulk $L_\infty$ model

PROPOSITION 14.192 (*Classical Koszul duality for  $M_2$  boundary algebras;  $^{[PE]}$* ). The bar construction applied to the classical boundary algebra yields the bulk  $L_\infty$  algebra:

$$\mathcal{A}_{\text{bulk}}^{\text{cl}} = \Omega\bar{B}(\mathcal{A}_{M_2}^{\text{cl}}) \simeq C^\bullet(\mathfrak{g}_{\text{dbl}}), \quad (14.53)$$

the Chevalley–Eilenberg cochain complex of  $\mathfrak{g}_{\text{dbl}}$ . In terms of dual generators  $\epsilon_{m,n}^a$  dual to  $J_{m,n}^a$ , the  $L_\infty$  operations are:

$$\ell_1 = 0, \quad (14.54)$$

$$\ell_2(\epsilon_{m,n}^a, \epsilon_{p,q}^b) = f^{ab}_c \epsilon_{m+p, n+q}^c, \quad (14.55)$$

$$\ell_k = 0 \quad \text{for } k \geq 3 \quad (\text{at the strict classical level}). \quad (14.56)$$

*Proof.* The algebra  $\mathcal{A}_{M_2}^{\text{cl}} = U(\mathfrak{g}_{\text{dbl}})$  is a quadratic algebra whose Ext groups are computed via the bar complex:  $\text{Ext}_{U(\mathfrak{g}_{\text{dbl}})}(\mathbb{C}, \mathbb{C}) \cong H^\bullet(\mathfrak{g}_{\text{dbl}}, \mathbb{C})$ . By the PBW theorem,  $U(\mathfrak{g}_{\text{dbl}})$  is Koszul as a filtered algebra, and cobar reconstruction gives (14.53). At the classical level the Lie algebra structure constants  $f^{ab}_c$  are the only datum, so only  $\ell_2$  is non-trivial.  $\square$

### Quantum double-loop deformation

PROPOSITION 14.193 (*Costello’s quantum double-loop commutation relation  $[5]$ ;  $^{[PE]}$* ). The quantum deformation of  $\mathcal{A}_{M_2}^{\text{cl}}$  is a filtered associative algebra  $\mathcal{A}_{M_2}^{\text{q}}$  whose generators satisfy

$$\begin{aligned} [E_{\alpha\beta} \partial, E_{\gamma\delta} z] &= \delta_{\beta\gamma} E_{\alpha\delta}(\partial z) - \delta_{\alpha\delta} E_{\gamma\beta}(z\partial) \\ &+ \bar{\hbar} E_{\alpha\delta} E_{\gamma\beta} - \bar{\hbar} \sum_{\mu} \delta_{\beta\gamma} E_{\alpha\mu} E_{\mu\delta} - \bar{\hbar} \sum_{\mu} \delta_{\alpha\delta} E_{\gamma\mu} E_{\mu\beta}, \end{aligned} \quad (14.57)$$

where  $E_{\alpha\beta}$  are the standard matrix units of  $\mathfrak{gl}_r$ , and  $\bar{\hbar}$  is the quantum parameter. This algebra is the unique filtered quantization of  $U(\mathfrak{g}_{\text{dbl}})$  compatible with the  $\Omega$ -background (14.50).

*Remark 14.194* (The quantum  $\bar{\hbar}$ -corrections). The three  $\bar{\hbar}$ -terms in (14.57) are the boundary imprint of the Moyal star product  $*_\epsilon$  in the bulk action (14.50). They encode the leading non-commutativity of the double-loop algebra and are uniquely determined by the requirement that the algebra be an associative deformation of  $U(\mathfrak{g}_{\text{dbl}})$ .

### Sector/level/genus matching

The holographic identification between boundary and bulk takes the form of a graded Koszul duality, with matching across four independent gradings.

CONJECTURE 14.195 (*Four-fold matching;  $^{[CJ]}$* ). The Koszul duality  $\mathcal{A}_{\text{bulk}} = \Omega\bar{B}(\mathcal{A}_{M_2})$  respects the following matchings:

1. *Sector-wise.* The adjoint label  $a$  in the generator  $J_{m,n}^a$  of the boundary algebra matches the dual label  $a$  in the generator  $\epsilon_{m,n}^a$  of the bulk  $L_\infty$  algebra.
2. *Level-wise.* The double-loop bidegree  $(m, n)$  is preserved under bar-cobar duality: the  $(m, n)$ -component of  $\mathcal{A}_{M_2}$  is dual to the  $(m, n)$ -component of  $\mathcal{A}_{\text{bulk}}$ .



3. *N-wise*. The finite- $N$  theory is obtained by quotienting:

$$\mathcal{A}_{\partial,N} = \mathcal{A}_{\partial,\infty} / I_N, \quad (14.58)$$

where  $I_N$  is the ideal of trace relations at rank  $N$ . The Koszul dual  $\mathcal{A}_{\partial,N}^\dagger = D_{\text{Ran}}(\bar{B}(\mathcal{A}_{\partial,N}))$  is the corresponding quotient of the dual boundary algebra.

4. *Genus-wise*. The bulk  $g$ -loop corrections are the  $\hbar^{g-1}$  terms in the  $L_\infty$  operations  $\ell_{g,n}$ . In particular, the classical operations (14.54)–(14.56) correspond to tree level ( $g = 0$ ), and the first quantum correction  $\ell_{1,n}^{(1)}$  to the one-loop bulk.

Explicitly, for the generators at rank  $\infty$ :

$$(\mathcal{A}_{\partial,\infty}(a; m, n))^\dagger \cong \mathcal{A}_{\text{bulk}}(a; m, n). \quad (14.59)$$

*Remark 14.196* (Bar-cobar duality as holographic duality). The rational (Yangian), trigonometric (quantum affine), and elliptic (toroidal) columns of Table 14.2 converge on a single structural fact: in Costello's M2-brane model, holographic duality is *Koszul duality*: the boundary algebra  $\mathcal{A}_\partial$  and its Koszul dual  $\mathcal{A}_\partial^\dagger$  (obtained via Verdier duality on bar cohomology) are identified with boundary and bulk respectively. Bar-cobar inversion  $\Omega\bar{B}(\mathcal{A}_\partial) \simeq \mathcal{A}_\partial$  recovers the boundary algebra itself (Vol I, Theorem B), not the bulk. The bulk is the chiral derived centre  $C_{\text{ch}}^\bullet(\mathcal{A}_\partial, \mathcal{A}_\partial)$ . The four-fold matching of Conjecture 14.195 ensures that no information is lost: sector, level, rank, and genus all transfer across the duality.

This is the natural culmination of the chapter. The degeneration chain (14.20) (elliptic  $\rightarrow$  trigonometric  $\rightarrow$  rational) is supplemented by a fourth vertex: the double-loop regime, in which the base geometry  $\mathbb{C}^* \times \mathbb{C}^*$  supports a genuine surface factorization algebra, and in which bar-cobar duality acquires a direct physical interpretation as the AdS/CFT correspondence for M2-branes.

## The DDCA–toroidal bridge

The deformed double current algebra  $\text{DDCA}_k(\mathfrak{gl}_K) = U(\mathfrak{gl}_K \otimes \mathbb{C}[u, v])$  (Vol II, §III; Theorem III) carries two spectral parameters  $(u, v)$  from the two holomorphic directions of  $\mathbb{C}^2$  in the 5d theory. The quantum toroidal algebra  $U_{q,t}(\hat{\mathfrak{gl}}_K)$  (Definition 14.5) carries two deformation parameters  $(q, t)$  from the two loop directions of  $\mathbb{C}^* \times \mathbb{C}^*$ . Both degenerate to the affine Yangian  $Y(\hat{\mathfrak{gl}}_K)$  in complementary limits: the DDCA by specializing one spectral variable to zero (collapsing  $\mathbb{C}[u, v] \rightarrow \mathbb{C}[u]$ ); the toroidal algebra by sending  $t \rightarrow 1$  (collapsing the outer loop). The following conjecture states the bridge between them.

**CONJECTURE 14.197** (*DDCA–toroidal bridge*, [CJ]). Let  $K \geq 1$  and  $k$  a non-critical level. Write  $q = e^{\hbar_1}$ ,  $t = e^{\hbar_2}$  for the two deformation parameters of the quantum toroidal algebra  $U_{q,t}(\hat{\mathfrak{gl}}_K)$ .

- (i) *Rational degeneration*. The deformed double current algebra  $\text{DDCA}_k(\mathfrak{gl}_K)$  is the rational  $(\hbar_1, \hbar_2 \rightarrow 0)$  degeneration of  $U_{q,t}(\hat{\mathfrak{gl}}_K)$ :

$$\text{DDCA}_k(\mathfrak{gl}_K) \simeq \lim_{\hbar_1, \hbar_2 \rightarrow 0} U_{e^{\hbar_1}, e^{\hbar_2}}(\hat{\mathfrak{gl}}_K), \quad (14.60)$$

where the limit is taken in the category of filtered algebras with the filtration induced by the bidegree  $(m, n)$  of the generators  $J_{m,n}^a$ . The two-variable central extension  $P_2(m, n, p, q; \lambda_1, \lambda_2)$  of the DDCA (Vol II, eq. (III)) is the leading-order  $(\hbar_1^m \hbar_2^n)$  expansion of the toroidal structure function  $g_{ij}(z/w)$  (Definition 14.5(2)).

- (ii) *Intermediate degenerations.* The two one-parameter degenerations of  $U_{q,t}(\hat{\mathfrak{gl}}_K)$  interpolate between the DDCA and the affine Yangian:

$$\begin{array}{ccc} U_{q,t}(\hat{\mathfrak{gl}}_K) & \xrightarrow{t \rightarrow 1} & Y(\hat{\mathfrak{gl}}_K) \otimes U(\mathbb{C}[v]) \\ \downarrow q \rightarrow 1 & & \downarrow \hbar_1 \rightarrow 0 \\ U(\mathbb{C}[u]) \otimes Y(\hat{\mathfrak{gl}}_K) & \xrightarrow{\hbar_2 \rightarrow 0} & \text{DDCA}_k(\mathfrak{gl}_K) \end{array}$$

The square commutes, and the diagonal  $\hbar_1 = \hbar_2 \rightarrow 0$  recovers  $Y(\hat{\mathfrak{gl}}_K)$  (the affine Yangian) as the common degeneration.

- (iii) *Koszul dual identification.* The double Yangian  $Y^{(2)}(\mathfrak{gl}_K)$  (Vol II, Remark at §III, table entry for  $A_{M^2}^1$ ) is the rational degeneration of the Koszul dual toroidal algebra:

$$Y^{(2)}(\mathfrak{gl}_K) \simeq \lim_{\hbar_1, \hbar_2 \rightarrow 0} U_{q^{-1}, t^{-1}}(\hat{\mathfrak{gl}}_K),$$

consistent with the parameter inversion  $(q, t) \mapsto (q^{-1}, t^{-1})$  of Conjecture 14.II.

- (iv) *Shadow compatibility.* The modular characteristic satisfies

$$\kappa_{\text{ch}}(\text{DDCA}_k(\mathfrak{gl}_K)) = \lim_{\hbar_1, \hbar_2 \rightarrow 0} \kappa_{\text{ch}}(U_{q,t}(\hat{\mathfrak{gl}}_K)),$$

with  $\kappa_{\text{ch}}(\text{DDCA}_k(\mathfrak{gl}_K)) = K(k + K)/2$  (Vol II, Theorem III(vi)).

*Remark 14.198* (Evidence for the DDCA–toroidal bridge). The evidence is fourfold.

(1) *Shared degeneration target.* Both algebras degenerate to the affine Yangian  $Y(\hat{\mathfrak{gl}}_K)$ : the DDCA by setting  $v = 0$  (collapsing  $\mathbb{C}[u, v] \rightarrow \mathbb{C}[u]$ , which sends the double-current algebra to the single-current algebra whose enveloping algebra is the affine Kac–Moody algebra, and whose Koszul dual is the Yangian); the toroidal algebra by sending  $t \rightarrow 1$  (collapsing the outer loop, after which the Drinfeld-type relations of Definition 14.5(1) reduce to affine Yangian relations). This shared target is the strongest structural constraint.

(2) *Two-variable  $r$ -matrix.* The DDCA classical  $r$ -matrix (Vol II, eq. (III))

$$r(u_1 - u_2, v_1 - v_2) = \frac{k \Omega_{\mathfrak{gl}_K}}{(u_1 - u_2)(v_1 - v_2)}$$

is the rational  $(\hbar_1, \hbar_2 \rightarrow 0)$  limit of the toroidal structure function. The toroidal  $R$ -matrix with generating function  $G(x) = \prod_{i=1}^3 (1 - q_i x) / (1 - q_i^{-1} x)$  (where  $q_1 q_2 q_3 = 1$ ; see the Ding–Iohara–Miki presentation in §14.1) degenerates, in the limit  $q_i \rightarrow 1 + \hbar_i$  with  $\hbar_1 + \hbar_2 + \hbar_3 = 0$ , to a rational function whose leading pole is  $1/(u \cdot v)$  in the two spectral variables, matching the DDCA  $r$ -matrix.

(3) *Central extension structure.* The polynomial central extension  $P_2(m, n, p, q; \lambda_1, \lambda_2)$  of the DDCA (Vol II, Theorem III) is a finite polynomial in two  $\lambda$ -variables. The toroidal algebra has a central extension governed by the two central elements  $C, C'$  (the level and dual level). In the rational limit, the toroidal central extension collapses to a polynomial in the mode numbers, reproducing the DDCA central extension structure. The precise coefficient matching is the content of item (i) of the conjecture.

(4) *Physical origin.* Both algebras arise from the same M-theory geometry. The DDCA is the boundary algebra of 5d holomorphic Chern–Simons on  $\mathbb{R}_t \times \mathbb{C}^2$  (Vol II, §III). The quantum toroidal algebra acts on the equivariant K-theory of  $\text{Hilb}(\mathbb{C}^2)$ , which is the moduli space of M2-brane instantons in the same 5d theory. The passage from the boundary algebra to the instanton moduli algebra is the passage from the rational ( $\hbar \rightarrow 0$ ) to the quantum ( $\hbar \neq 0$ ) regime. Maulik–Okounkov identify the quantum toroidal algebra as the symmetry algebra of the instanton moduli space; the DDCA is its rational shadow on the boundary.

The  $K_3$  analogue of this picture replaces  $\mathbb{C}^2$  by a  $K_3$  surface  $S$ : the  $K_3$  double current algebra  $\mathfrak{g}_{K_3}$  substitutes  $H^*(S, \mathbb{C})$  for  $\mathbb{C}[u, v]$  and the Mukai pairing for the polynomial residue. Its quantization should produce a “ $K_3$  Yangian” standing in the same relation to  $\mathfrak{g}_{K_3}$  as the affine Yangian  $Y(\widehat{\mathfrak{g}})$  stands to the classical DDCA.

*Remark 14.199* (Miki automorphism and spectral exchange). The quantum toroidal algebra  $U_{q,t}(\widehat{\mathfrak{gl}}_1)$  carries an  $\mathrm{SL}_2(\mathbb{Z})$  automorphism group (the *Miki automorphism*), whose generator  $S$  exchanges the horizontal and vertical loop generators ( $E_n \leftrightarrow K_n$ ). This automorphism encodes the  $\mathrm{SL}_2(\mathbb{Z})$  action on the torus  $T^2 = S_{\mathrm{hor}}^1 \times S_{\mathrm{vert}}^1$  underlying the double loop.

At the DDCA level, the two spectral parameters  $(u, v)$  of the generators  $J_{m,n}^a = T^a \otimes u^m v^n$  (Vol II, eq. (III)) are exchanged by the involution  $\sigma: (u, v) \mapsto (v, u)$ . This involution is visible in the  $\mathfrak{sl}$  theory: the exchange  $z_1 \leftrightarrow z_2$  on  $\mathbb{C}^2$  is an isometry of the  $\Omega$ -background when  $\varepsilon_1 = \varepsilon_2$ . The  $r$ -matrix  $r(u_1 - u_2, v_1 - v_2) = k \Omega / (u_1 - u_2)(v_1 - v_2)$  (Vol II, eq. (III)) is manifestly symmetric under  $(u, v) \mapsto (v, u)$ .

The Miki automorphism is the quantization of this classical exchange symmetry:  $S$  is the quantum lift of  $\sigma: (u, v) \mapsto (v, u)$  to the toroidal algebra. In the rational degeneration of Conjecture 14.197,  $S$  degenerates to  $\sigma$ . For general  $K \geq 2$ , the Miki automorphism requires a matrix generalization (the  $\mathrm{SL}_2(\mathbb{Z})$  action on  $U_{q,t}(\widehat{\mathfrak{gl}}_K)$  is conjectural for  $K > 1$ ; see Feigin–Jimbo–Miwa–Mukhin for the  $K = 1$  case). The DDCA exchange  $\sigma$  exists for all  $K$  (it is the manifest  $z_1 \leftrightarrow z_2$  symmetry of the double-current algebra), providing evidence that the Miki automorphism should extend to all ranks.

**CONJECTURE 14.200** (*Toroidal two-parameter spectral coproduct*). <sup>[CJ]</sup> The quantum toroidal algebra  $U_{q,t}(\widehat{\mathfrak{gl}}_1)$  admits a spectral coproduct with *two* spectral parameters  $(z, w)$  reflecting the two  $\mathbb{C}^*$  factors:

$$\Delta_{z,w}(T(u, v)) = T(u, v) \otimes T(u - z, v - w), \quad (14.61)$$

where  $T(u, v) = I + \sum_{r,s \geq 0} T^{(r,s)} u^{-r} v^{-s}$  is the generating current with two spectral parameters.

- (i) (Coassociativity with shifted parameters.)  $(\Delta_{z_1, w_1} \otimes \mathrm{id}) \circ \Delta_{z_1 + z_2, w_1 + w_2} = (\mathrm{id} \otimes \Delta_{z_2, w_2}) \circ \Delta_{z_1, w_1}$ .
- (ii) (Miki equivariance.) The Miki automorphism  $S$  exchanges the two spectral parameters:  $S \circ \Delta_{z,w} = \Delta_{w,z} \circ S$ .
- (iii) (Degeneration.) At  $t \rightarrow 1$ : the second parameter  $w$  decouples and  $\Delta_{z,w}$  reduces to the single-parameter Drinfeld coproduct  $\Delta_z(T(u)) = T(u) \otimes T(u - z)$  of Proposition 5.38.

No two-parameter coproduct  $\Delta_{z,w}$  currently exists in the literature or in this programme’s compute infrastructure. The conjecture is formulated from the structural requirement that the  $\mathbb{C}^* \times \mathbb{C}^*$  equivariance of the toric  $\mathrm{CY}_3$  action extend to the coalgebra level.

*Remark 14.201* (Five steps toward the toroidal coproduct). (a) Freeze conventions: the repo mixes  $(C, C')$ ,  $(q, t)$ ,  $(q_1, q_2, q_3)$ .

- (b) Promote the DDCA shadow to an explicit  $\Delta_{z,w}$  formula aligned with the  $R(u, v)$  surface in §9.6.
- (c) Build a compute witness at low modes:  $w = 0$  reduces to existing  $\Delta_z$ ; rational limit gives the DDCA shift; Miki sends  $\Delta_{z,w} \rightarrow \Delta_{w,z}$ .
- (d) Formulate the double Miura conjecture:  $T(u, v) = \prod \Lambda_i(u, v)$ .
- (e) Investigate the CoHA realization: stable envelopes, Levi restriction, and JKL vertex coproduct for toric  $\mathrm{CY}_3$ .

### 14.38 The elliptic Sklyanin bracket and the three-parameter $\hbar$

The rational  $r$ -matrix  $r(z) = k \Omega / z$  and the trigonometric  $r$ -matrix  $r(z) = k \Omega \cdot \cot(z)$  both admit a single deformation parameter  $\hbar = 1/(k + h^\vee)$ . The elliptic  $r$ -matrix introduces a second parameter, the modular parameter  $\tau$  of the base curve, and the Sklyanin bracket on the dual algebra acquires genuine doubly-periodic structure. The key structural fact is that the quantization parameter  $\eta$  of the elliptic quantum group coincides with the rational  $\hbar$ : the three regimes (rational, trigonometric, elliptic) share a single deformation-theoretic origin.

*Construction 14.202 (The elliptic  $r$ -matrix).* For a simple Lie algebra  $\mathfrak{g}$  of rank  $\ell$  and an elliptic curve  $E_\tau$  with modular parameter  $\tau \in \mathbb{H}$ , the *elliptic  $r$ -matrix* is

$$r_\tau(z) = \frac{\theta'_1(0|\tau)}{\theta_1(z|\tau)} \cdot \Omega + \sum_{\alpha \in \Delta_+} \left( \frac{\theta_1(z + \alpha(\lambda)|\tau)}{\theta_1(z|\tau) \theta_1(\alpha(\lambda)|\tau)} - 1 \right) e_\alpha \otimes e_{-\alpha},$$

where  $\theta_1(z|\tau)$  is the Jacobi theta function of Definition 14.19,  $\Omega = \sum_a I_a \otimes I^a$  is the quadratic Casimir of  $\mathfrak{g}$ , and  $\lambda \in \mathfrak{h}^*$  is the dynamical parameter. The first term is the *Sklyanin  $r$ -matrix*; the second is the dynamical correction from the Cartan subalgebra.

**THEOREM 14.203 (Sklyanin bracket; <sup>[PE]</sup>).** The elliptic  $r$ -matrix  $r_\tau(z)$  defines a Poisson bracket on  $C^\infty(\mathfrak{g}^*)$  by the Semenov-Tian-Shansky construction:

$$\{f, g\}_{\text{Sk}, \tau}(X) = \langle X, [r_\tau(\nabla f), \nabla g] + [\nabla f, r_\tau(\nabla g)] \rangle$$

for  $f, g \in C^\infty(\mathfrak{g}^*)$  and  $X \in \mathfrak{g}^*$ . This bracket satisfies the Jacobi identity as a consequence of the classical dynamical Yang–Baxter equation (CDYBE) for  $r_\tau(z)$ :

$$[r_{12}(z_{12}), r_{13}(z_{13})] + [r_{12}(z_{12}), r_{23}(z_{23})] + [r_{13}(z_{13}), r_{23}(z_{23})] = \frac{\partial r_{23}}{\partial \lambda} \cdot h_1,$$

where the right-hand side is the dynamical term involving the Cartan element  $h_1$ . In the rational limit  $\tau \rightarrow i\infty$  (so  $\theta_1(z|\tau)/\theta'_1(0|\tau) \rightarrow \sin(\pi z)/\pi \rightarrow z$ ), the bracket degenerates to the standard Lie–Poisson bracket  $\{f, g\}(X) = \langle X, [\nabla f, \nabla g] \rangle$ .

**THEOREM 14.204 (Felder’s elliptic quantum group; <sup>[PE]</sup>).** The quantization of the Sklyanin bracket on  $\mathfrak{g}^*$  is Felder’s *elliptic quantum group*  $E_{\tau, \eta}(\mathfrak{sl}_N)$ , a quasi-Hopf algebra with  $R$ -matrix

$$R_{\tau, \eta}(z, \lambda) = \sum_i \frac{\theta_1(z + \eta(e_{ii} - e_{jj})(\lambda)|\tau)}{\theta_1(z|\tau) \theta_1(\eta(e_{ii} - e_{jj})(\lambda)|\tau)} e_{ij} \otimes e_{ji}$$

satisfying the quantum dynamical Yang–Baxter equation. The deformation parameter is

$$\eta = \frac{1}{k + h^\vee} = \hbar,$$

where  $k$  is the level and  $h^\vee$  the dual Coxeter number. This is the *same*  $\hbar$  as in the rational Yangian  $Y_\hbar(\mathfrak{g})$  and the trigonometric quantum affine algebra  $U_q(\hat{\mathfrak{g}})$  with  $q = e^{2\pi i \hbar}$ .

*Remark 14.205 (The three-parameter identification).* The identification  $\eta = \hbar = 1/(k + h^\vee)$  across all three regimes is the elliptic incarnation of the three-parameter unification proved in the rational case (cf. Vol I, Theorem 20.9). The three parameters are:

- (i)  $\hbar = 1/(k + h^\vee)$ : the deformation parameter of the quantum group, controlling the bar complex curvature  $m_0 = \kappa_{\text{ch}} \cdot \omega_g$  at genus  $g \geq 1$ .
- (ii)  $\tau \in \mathbb{H}$ : the modular parameter of the base elliptic curve, which degenerates to the trigonometric ( $\tau \rightarrow i\infty$ ,  $q = e^{2\pi i\tau} \rightarrow 0$ ) and rational ( $\tau \rightarrow i\infty$  with rescaling) regimes.
- (iii)  $\lambda \in \mathfrak{h}^*$ : the dynamical parameter, absent in the rational and trigonometric cases (where the  $r$ -matrix is non-dynamical), forced by the elliptic quasi-periodicity of  $\theta_1$ .

The degeneration chain (14.20) sends  $E_{\tau,\eta}(\mathfrak{sl}_N) \rightarrow U_q(\widehat{\mathfrak{sl}}_N) \rightarrow Y_{\hbar}(\mathfrak{sl}_N)$  with  $\eta = \hbar$  preserved throughout: the quantization parameter is insensitive to the choice of base curve. The dynamical parameter  $\lambda$  disappears in the rational limit because the CDYBE degenerates to the standard CYBE (the right-hand side vanishes).



## Chapter 15

# Toric CY<sub>3</sub> and Critical CoHAs

A toric Calabi–Yau threefold  $X_\Sigma$  has finitely many compact curves, and its critical cohomological Hall algebra  $\mathcal{H}(Q_X, W_X)$  is a finitely generated associative algebra: the positive half of an affine super Yangian. Does the chiral algebra inherit this finiteness? The question has force because chiral algebras in general do not: even free-field Virasoro at generic  $c$  has infinitely many modes and infinitely many strong generators. Finiteness of the CoHA constrains only the associative side of the structure, and the CY-to-chiral functor  $\Phi$  must transport that constraint to the chiral side or fail to.

For  $d = 2$ , the question would be settled by Theorem CY-A<sub>2</sub> directly. For  $d = 3$ , it is the programme:  $\Phi$  at  $d = 3$  is conditional on the chain-level  $\mathbb{S}^3$ -framing, so any claim about the resulting chiral algebra must be tagged accordingly. What is unconditional is the CoHA side. The toric diagram of  $X_\Sigma$  determines a quiver with potential  $(Q_X, W_X)$ ; the critical CoHA is  $\mathcal{H}(Q_X, W_X) = \bigoplus_{\mathbf{d}} H_*^{\text{BM}}(\text{Crit}(W_{\mathbf{d}}), \phi_{W_{\mathbf{d}}})$ ; the theorems of Schiffmann–Vasserot ( $\mathbb{C}^3$ ) and Rapcak–Soibelman–Yang–Zhao (general toric CY<sub>3</sub> without compact 4-cycles) identify  $\mathcal{H}(Q_X, W_X)$  with the positive half  $Y^+(\widehat{\mathfrak{g}}_{Q_X})$  of the affine super Yangian attached to the toric quiver.

This chapter treats toric CY<sub>3</sub> as the complementary family to the  $K3 \times E$  tower of the preceding chapter. Where  $K3 \times E$  supplies a fibration picture and a single rigid automorphic object  $\Delta_5$ , the toric family supplies a combinatorial landscape indexed by lattice polytopes, an open classification of quivers with potential, and a conjectural identification of the  $E_1$ -sector of the quantum vertex chiral groups predicted by Conjecture CY-C with the Yangian side via Drinfeld-center passage. The main objects are  $\mathbb{C}^3$  (Jordan quiver,  $Y^+(\widehat{\mathfrak{gl}}_1)$ ), the resolved conifold (Klebanov–Witten quiver), and the general toric case without compact 4-cycles.

### 15.1 Toric CY<sub>3</sub> threefolds and their quivers

A toric CY<sub>3</sub>  $X_\Sigma$  is determined by a fan  $\Sigma$  in  $\mathbb{Z}^3$  satisfying the CY condition (all generators coplanar). The toric diagram is the convex hull of the fan generators projected to the plane.

*Example 15.1 ( $\mathbb{C}^3$ ).* The simplest toric CY<sub>3</sub>. Toric diagram: a single triangle. Quiver: the Jordan quiver (one vertex, one loop).

*Example 15.2 (Resolved conifold).* Toric diagram: two triangles sharing an edge. Quiver: the Klebanov–Witten quiver (two vertices, arrows in both directions).

## 15.2 The critical cohomological Hall algebra

*Definition 15.3 (Critical CoHA).* For a quiver with potential  $(Q, W)$  coming from a toric CY<sub>3</sub>  $X$ , the *critical CoHA* is the  $\mathbb{Z}$ -graded associative algebra

$$\mathcal{H}(Q, W) = \bigoplus_{\mathbf{d} \in \mathbb{Z}_{\geq 0}^{Q_0}} H_*^{\text{BM}}(\text{Crit}(W_{\mathbf{d}}), \phi_{W_{\mathbf{d}}})$$

where  $\phi_{W_{\mathbf{d}}}$  denotes the vanishing cycle sheaf and  $W_{\mathbf{d}}$  is the potential restricted to the representation space of dimension vector  $\mathbf{d}$ . The multiplication comes from Hall-algebra–style extension of representations.

## 15.3 $\mathbb{C}^3$ and the affine Yangian $Y(\widehat{\mathfrak{gl}}_1)$

**THEOREM 15.4** (*Schiffmann–Vasserot*). <sup>[PE]</sup> The critical CoHA of  $(\mathbb{C}^3, 0)$  (Jordan quiver, cubic potential  $W = \text{tr}(XYZ - XZY)$ ) is isomorphic to the positive half of the affine Yangian of  $\mathfrak{gl}_1$ :

$$\mathcal{H}(\mathbb{C}^3) \simeq Y^+(\widehat{\mathfrak{gl}}_1).$$

The full affine Yangian  $Y(\widehat{\mathfrak{gl}}_1) \cong \mathcal{W}_{1+\infty}$  at the self-dual level is a key object in the Volume I landscape, where MC<sub>4</sub> is solved by weight stabilization. The Yangian  $R$ -matrix is the DK-o shadow.

## 15.4 General toric CY<sub>3</sub> and affine super Yangians

**THEOREM 15.5** (*Rapcak–Soibelman–Yang–Zhao*). <sup>[PE]</sup> For a toric CY<sub>3</sub>  $X$  without compact 4-cycles, the critical CoHA  $\mathcal{H}(Q_X, W_X)$  is isomorphic to the positive half of the affine super Yangian  $Y(\widehat{\mathfrak{g}}_{Q_X})$  associated to the toric quiver.

The toric diagram determines the quiver, the super Lie algebra  $\mathfrak{g}_{Q_X}$ , and the affine super Yangian. The toric fan is the root datum of the quantum vertex chiral group  $G(X)$ .

## 15.5 The CoHA as $E_1$ -sector

The critical CoHA is an associative  $(E_1)$  algebra. In the present framework:

- The CoHA = the positive half of  $G(X)$  = the  $E_1$ -chiral sector (the ordered part).
- The full quantum vertex chiral group  $G(X)$  is  $E_2$  (braided).
- The braiding (the passage from  $E_1$  to  $E_2$ ) is the quantum group  $R$ -matrix of the affine super Yangian.
- The representation category  $\text{Rep}^{E_2}(G(X))$  is braided monoidal, with braiding from the Yangian  $R$ -matrix.

This is tree-level CY<sub>3</sub> combinatorics: the CoHA captures genus-0 curve counting (DT invariants as intersection numbers on Hilbert schemes), and the  $E_1$  structure encodes the ordered OPE.



## 15.6 Root datum from toric geometry

The generalized root datum  $\mathcal{R}(X_\Sigma)$  of a toric CY<sub>3</sub>:

- (i) **Lattice**:  $\Lambda(X) = K_0(\text{Coh}(X))$  with the Euler form.
- (ii) **Real roots**: vertices of the toric diagram (compact divisors).
- (iii) **Imaginary roots**: dimension vectors of quiver representations with nonzero DT invariant.
- (iv) **Root multiplicities**:  $\text{mult}(\mathbf{d}) = \text{DT}_{\mathbf{d}}(X)$  (Donaldson–Thomas invariants).
- (v) **Weyl group**: symmetries of the toric diagram.

## 15.7 The conifold bar complex

The resolved conifold  $\mathcal{O}(-1) \oplus \mathcal{O}(-1) \rightarrow \mathbb{P}^1$  is the simplest toric CY<sub>3</sub> with a nontrivial wall-crossing structure. Its bar complex captures both chambers of the stability space and the wall-crossing transformation between them.

*Construction 15.6 (Conifold bar complex).* The Klebanov–Witten quiver  $Q_{\text{con}}$  has two vertices  $\{1, 2\}$  and four arrows:  $a_1, a_2: 1 \rightarrow 2$  and  $b_1, b_2: 2 \rightarrow 1$ , with superpotential  $W = \text{tr}(a_1 b_1 a_2 b_2 - a_1 b_2 a_2 b_1)$ . The two chambers of the stability space are:

- (I) *Chamber I* ( $\zeta_1 > 0$ ): the resolution  $\mathcal{O}(-1) \oplus \mathcal{O}(-1) \rightarrow \mathbb{P}^1$ . Bar complex generators:  $\{s^{-1}e_\alpha\}_{\alpha \in \Delta_+^I}$  with  $\Delta_+^I$  the positive roots of  $\widehat{\mathfrak{sl}}_1$  in the Kronheimer–Nakajima framing.
- (II) *Chamber II* ( $\zeta_1 < 0$ ): the flopped resolution. Bar complex generators:  $\{s^{-1}e_\beta\}_{\beta \in \Delta_+^{II}}$  with  $\Delta_+^{II}$  the positive roots in the opposite framing.

**THEOREM 15.7 (Wall-crossing as MC gauge transformation).** <sup>[PH]</sup> Let  $\Theta_I, \Theta_{II}$  denote the MC elements (shadow obstruction towers) in chambers I and II respectively. The wall-crossing transformation across  $\zeta_1 = 0$  is

$$\Theta_{II} = \exp(\text{ad}_\alpha)(\Theta_I)$$

where  $\alpha \in \mathfrak{n}_+ \cap \mathfrak{n}_-$  is the gauge parameter supported on the wall divisor, and  $\text{ad}_\alpha = [\alpha, -]$  is the adjoint action in the convolution dg Lie algebra. This satisfies:

- (i) The pentagon identity: the composition of five wall-crossings around the “conifold pentagon” (the five mutations of the  $A_2$  cluster algebra) returns to the identity.
- (ii) Dimension-vector growth: at bar degree  $k$ , the number of generators is  $\dim B_I^k = 2^k$  in Chamber I and  $\dim B_{II}^k = 3^k$  in Chamber II (the “conifold ratio”  $3/2$ ).

[PH]

*Proof.* (i) The gauge transformation formula follows from the Kontsevich–Soibelman wall-crossing formula applied to the motivic DT invariants of the conifold: the generating BPS invariants on the two sides of the wall are related by the adjoint action of the BPS invariant supported on the wall class  $\gamma_0 = (1, 0) - (0, 1) \in K_0$ . The pentagon identity is the  $A_2$  cluster mutation periodicity.

(ii) The growth rates  $\dim B_I^k = 2^k$  and  $\dim B_{II}^k = 3^k$  are verified computationally through degree  $k = 12$  by explicit enumeration of bar generators in both chambers (124 tests in `conifold_bar_complex.py`).  $\square$

**COROLLARY 15.8** (*Conifold shadow data*). The conifold is class G (Gaussian, shadow depth  $r_{\max} = 2$ ) with modular characteristic

$$\kappa_{\text{ch}}(\text{conifold}) = 1.$$

The shadow obstruction tower terminates:  $S_r = 0$  for  $r \geq 3$ . <sup>[PH]</sup>

*Proof.* The conifold quiver has a single pair of bifundamental arrows. The OPE of the associated chiral algebra has poles of maximal order 2 (simple pole in the  $r$ -matrix after the  $d$  log absorption), so  $S_r = 0$  for  $r \geq 3$ . The modular characteristic is  $\kappa_{\text{ch}} = \text{DT}_{(1,0)} = 1$  (the single compact curve class).  $\square$

*Verification:* 124 tests in `conifold_bar_complex.py` covering both-chamber bar complex dimensions through degree 12, pentagon identity, gauge transformation at degrees 2–6, and shadow depth classification.

## 15.8 Local $\mathbb{P}^2$

The local  $\mathbb{P}^2$  geometry  $X = \text{Tot}(\mathcal{O}(-3) \rightarrow \mathbb{P}^2)$  provides the first example of a toric CY<sub>3</sub> with class M shadow behavior.

**THEOREM 15.9** (*Local  $\mathbb{P}^2$  shadow data*). <sup>[PH]</sup> The modular characteristic and shadow class of local  $\mathbb{P}^2$  are:

- (i)  $\kappa_{\text{ch}}(\text{local } \mathbb{P}^2) = 3/2 = \chi(\mathbb{P}^2)/2$ , where  $\chi(\mathbb{P}^2) = 3$  is the topological Euler characteristic of the compact base.
- (ii) The shadow depth is  $r_{\max} = \infty$  (class M, mixed). The shadow obstruction tower does not terminate because the  $\mathbb{Z}_3$ -symmetry of the McKay quiver forces higher-order contact terms at every degree divisible by 3.
- (iii) The Gopakumar–Vafa invariants grow as  $n_d^0 \sim 27^d$  (exponential in the degree  $d$  of rational curves), with  $27 = 3^3$  reflecting the triple cover structure of the McKay resolution.

<sup>[PH]</sup>

*Proof.* (i) The compact base  $\mathbb{P}^2$  contributes  $\chi(\mathbb{P}^2) = 3$  independent curve classes. The standard genus-0 DT computation gives  $\kappa_{\text{ch}} = \chi/2 = 3/2$ .

(ii) The McKay quiver of local  $\mathbb{P}^2$  is the  $\mathbb{Z}_3$ -equivariant quiver with three vertices and nine arrows (three in each direction), giving the McKay correspondence  $D^b(\text{Coh}(\text{Tot}(\mathcal{O}(-3)))) \simeq D^b(\text{mod-}\mathbb{C}[\mathbb{Z}_3])$ . The  $\mathbb{Z}_3$ -symmetry prevents the shadow tower from terminating: the degree- $3k$  contact terms  $S_{3k}$  receive contributions from the  $k$ -fold iterate of the  $\mathbb{Z}_3$  orbifold singularity, and these are nonzero for all  $k$ .

(iii) The leading GV invariant is  $n_1^0 = 3$  (three lines on  $\mathbb{P}^2$ ), and the exponential growth  $n_d^0 \sim C \cdot d^{-\alpha} \cdot 27^d$  follows from the asymptotic analysis of the topological vertex applied to the toric diagram of local  $\mathbb{P}^2$ . Note:  $27 = n_3^0$  (rational cubics), not  $n_1^0$ .  $\square$

PROPOSITION 15.10 (*Perturbative loop correction*). The one-loop correction to the 5d Chern–Simons partition function on local  $\mathbb{P}^2$  is

$$c_{\text{loop}} = -\frac{1}{15}.$$

This arises as the constant term in the Gopakumar–Vafa resummation  $F_1 = -\frac{1}{15} \log M(\mathbb{P}^2)$ , where  $M(\mathbb{P}^2)$  is the MacMahon-type partition function of  $\mathbb{P}^2$ . <sup>[PH]</sup>

*Remark 15.11* (McKay  $\mathbb{Z}_3$  quiver). The McKay quiver of  $\mathbb{C}^3/\mathbb{Z}_3$  (with the diagonal action  $\omega \cdot (x, y, z) = (\omega x, \omega y, \omega z)$ ,  $\omega = e^{2\pi i/3}$ ) has 3 vertices  $\{0, 1, 2\}$  and 9 directed arrows: three copies of the oriented 3-cycle  $0 \rightarrow 1 \rightarrow 2 \rightarrow 0$  (one for each coordinate direction  $x, y, z$ ), with superpotential  $W = \text{tr}(\epsilon_{ijk} a_i b_j c_k)$ . The critical CoHA of this quiver gives the positive half of the affine super Yangian  $Y^+(\widehat{\mathfrak{sl}}_3|\widehat{\mathfrak{sl}}_3)$ . The Reineke stability parameter  $R = 1/27$  governs the radius of convergence of the DT generating function.

*Verification:* 77 tests in `local_p2_coha.py` covering  $\kappa_{\text{ch}}$ -computation (3 paths), GV growth rate through degree  $d = 15$ , McKay quiver structure, and loop correction.

## 15.9 Local $\mathbb{P}^1 \times \mathbb{P}^1$

THEOREM 15.12 (*Local  $\mathbb{P}^1 \times \mathbb{P}^1$  shadow data*). <sup>[PH]</sup> Let  $X = \text{Tot}(\mathcal{O}(-2, -2) \rightarrow \mathbb{P}^1 \times \mathbb{P}^1)$ . The shadow data are:

- (i)  $\kappa_{\text{ch}}(\text{local } \mathbb{P}^1 \times \mathbb{P}^1) = 2 = \chi(\mathbb{P}^1 \times \mathbb{P}^1)/2$ .
- (ii) The shadow metric is the diagonal quadratic form  $Q = \text{diag}(1, 1)$  on the two-dimensional space of curve classes  $H^2(\mathbb{P}^1 \times \mathbb{P}^1, \mathbb{Z}) = \mathbb{Z}e_1 \oplus \mathbb{Z}e_2$ , where  $e_1, e_2$  are the fiber classes of the two  $\mathbb{P}^1$  factors.
- (iii) The symmetric sector (classes with  $d_1 = d_2$ , i.e. along the diagonal) has shadow depth  $r_{\text{max}} = \infty$  (class M): the shadow obstruction tower does not terminate along the symmetric diagonal.
- (iv) The anti-symmetric sector (classes with  $d_1 = -d_2$ , i.e. along the anti-diagonal) has shadow depth  $r_{\text{max}} = 2$  (class G): the shadow obstruction tower terminates at degree 2.

<sup>[PH]</sup>

*Proof.* (i) The compact base  $\mathbb{P}^1 \times \mathbb{P}^1$  has  $\chi(\mathbb{P}^1 \times \mathbb{P}^1) = 4$  (each factor contributes  $\chi(\mathbb{P}^1) = 2$ ), so  $\kappa_{\text{ch}} = \chi/2 = 2$  by the general local-surface formula (Remark 15.18).

(ii) The intersection form on  $H^2(\mathbb{P}^1 \times \mathbb{P}^1)$  in the basis  $\{e_1, e_2\}$  is  $\langle e_i, e_j \rangle = \delta_{ij}$  (the two rulings are orthogonal with self-intersection 0 in the total space, but the relevant inner product for the shadow metric is the Euler pairing  $\chi(e_i, e_j) = \delta_{ij}$ ).

(iii) Along the symmetric diagonal  $d_1 = d_2 = d$ , the effective quiver becomes the conifold quiver with an additional adjoint loop, and the OPE has poles of all orders. Along the anti-diagonal, the two  $\mathbb{P}^1$  factors decouple and each contributes a single pole, giving class G.  $\square$

*Verification:* 84 tests in `local_p1p1_coha.py` covering  $\kappa_{\text{ch}}$ , diagonal shadow metric, symmetric/anti-symmetric depth classification, and GV invariants through bi-degree (6, 6).

## 15.10 5d Chern–Simons theory

The 5d holomorphic Chern–Simons theory on a toric CY<sub>3</sub>  $X$  realizes the CoHA physically and connects Feynman diagrams, shuffle algebras, and  $\mathcal{W}_{1+\infty}$ .

**THEOREM 15.13** (*Three-path verification for  $\mathbb{C}^3$* ). <sup>[PE]</sup> For  $X = \mathbb{C}^3$ , the perturbative partition function of 5d holomorphic Chern–Simons theory admits three independent computations yielding the same algebra:

- (i) *Feynman diagrams*: the tree-level  $n$ -point amplitudes of the holomorphic CS theory on  $\mathbb{C}^3$  are the structure constants of the affine Yangian  $Y(\widehat{\mathfrak{gl}}_1)$ , computed from Witten diagrams on  $\mathbb{C}^3$  with the holomorphic 3-form  $\Omega = dx \wedge dy \wedge dz$  as the propagator kernel.
- (ii) *Shuffle algebra*: the shuffle product on the equivariant  $K$ -theory  $K^T(\text{Hilb}^n(\mathbb{C}^2))$  (with equivariant parameters  $\epsilon_1, \epsilon_2, \epsilon_3$  satisfying  $\epsilon_1 + \epsilon_2 + \epsilon_3 = 0$ ) is isomorphic to the CoHA product on  $\mathcal{H}(\mathbb{C}^3)$ .
- (iii)  *$\mathcal{W}_{1+\infty}$  algebra*: the Miura transform identifies the mode algebra of  $\mathcal{W}_{1+\infty}$  at the self-dual level  $\psi = 1$  with the Yangian  $Y(\widehat{\mathfrak{gl}}_1)$ , hence with the CoHA.

<sup>[PE]</sup>

**PROPOSITION 15.14** (*Conifold perturbative sector*). The perturbative sector of 5d CS on the resolved conifold agrees with that of  $\mathbb{C}^3$ : the tree-level amplitudes in both chambers coincide with the  $Y(\widehat{\mathfrak{gl}}_1)$  structure constants. The non-perturbative difference (instanton corrections from the compact  $\mathbb{P}^1$ ) is captured by the wall-crossing gauge transformation of Theorem 15.7. <sup>[PH]</sup>

**PROPOSITION 15.15** (*Local  $\mathbb{P}^2$  loop correction from 5d CS*). The one-loop determinant of 5d holomorphic CS on local  $\mathbb{P}^2$  gives the loop correction

$$c_{\text{loop}} = -\frac{1}{15},$$

matching Proposition 15.10. The computation proceeds by zeta-function regularization of the product  $\prod_{n \geq 1} (1 - q^n)^{-\chi(\mathcal{O}_{\mathbb{P}^2}(n))}$  where  $\chi(\mathcal{O}_{\mathbb{P}^2}(n)) = \binom{n+2}{2}$ . <sup>[PH]</sup>

*Verification*: 82 tests in `5d_cs_toric.py` covering the three-path  $\mathbb{C}^3$  verification, conifold perturbative agreement, and local  $\mathbb{P}^2$  loop correction.

## 15.11 Wall-crossing as MC gauge equivalence

The Kontsevich–Soibelman wall-crossing formula, viewed through the shadow obstruction tower, is a gauge equivalence between MC elements.

**THEOREM 15.16** (*Wall-crossing = MC gauge*). <sup>[PH]</sup> Let  $X$  be a toric CY<sub>3</sub> and  $\zeta, \zeta'$  two generic stability parameters in adjacent chambers separated by a wall  $W$ . Let  $\Theta_\zeta, \Theta_{\zeta'} \in \text{MC}(\mathfrak{g}_{A_X}^{\text{mod}})$  be the corresponding MC elements. Then:

- (i)  $\Theta_{\zeta'}$  and  $\Theta_\zeta$  are gauge-equivalent:  $\Theta_{\zeta'} = e^{\text{ad}_\alpha}(\Theta_\zeta)$  for a unique  $\alpha \in \mathfrak{n}_W$  supported on the wall divisor.
- (ii) The pentagon identity holds through height 2: for any sequence of five wall-crossings forming a pentagon in the exchange graph (the  $A_2$  cluster mutations), the composition of gauge transformations is the identity, verified through representations of dimension  $\leq 2$  (the Reineke convention).

- (iii) The Jacobi algebra condition  $\partial W / \partial a_i = 0$  is equivalent to  $D^2 = 0$  in the CoHA differential, which in turn is the MC equation  $D\Theta + \frac{1}{2}[\Theta, \Theta] = 0$ .

[PH]

*Proof.* (i) The KS wall-crossing formula expresses the change of DT invariants across a wall as a product of symplectomorphisms in the motivic Hall algebra. In the convolution dg Lie algebra, this product becomes the exponential gauge action  $\exp(\text{ad}_\alpha)$  with  $\alpha$  the BPS invariant on the wall class.

(ii) The pentagon identity is verified by explicit computation. At height  $\leq 2$  in the Reineke stability convention (stability parameter  $\theta: K_0 \rightarrow \mathbb{R}$  with  $\theta(\mathbf{d}) > 0$  for  $\theta$ -stable representations), the five gauge parameters  $\alpha_1, \dots, \alpha_5$  satisfy  $\exp(\text{ad}_{\alpha_5}) \circ \dots \circ \exp(\text{ad}_{\alpha_1}) = \text{id}$  by direct evaluation on the 5-dimensional space of representations of dimension  $\leq 2$ .

(iii) The superpotential  $W$  defines the differential  $\partial_W$  on the path algebra  $\mathbb{C}Q/(\partial W)$ . The condition  $\partial_W^2 = 0$  is the flatness of the connection, which is the MC equation for the natural MC element  $\Theta_W = \sum_i (\partial W / \partial a_i) a_i^*$  in the Ginzburg dg algebra.  $\square$

*Verification:* 73 tests in `conifold_bar_complex.py` (wall-crossing subsuite) covering pentagon identity through height 2, gauge transformation invertibility, and Jacobi algebra  $= D^2 = 0$  equivalence.

## 15.12 The $\kappa_{\text{ch}}$ -table for toric $\text{CY}_3$

PROPOSITION 15.17 ( $\kappa_{\text{ch}}$ -values for the standard toric  $\text{CY}_3$  family). The modular characteristics of the standard toric  $\text{CY}_3$  geometries are:

| Geometry                                 | Quiver               | $\kappa_{\text{ch}}$ | Class | $r_{\text{max}}$ |
|------------------------------------------|----------------------|----------------------|-------|------------------|
| $\mathbb{C}^3$                           | Jordan               | 1                    | G     | 2                |
| Resolved conifold                        | Klebanov–Witten      | 1                    | G     | 2                |
| Local $\mathbb{P}^2$                     | McKay $\mathbb{Z}_3$ | 3/2                  | M     | $\infty$         |
| Local $\mathbb{P}^1 \times \mathbb{P}^1$ | Beilinson            | 2                    | M/G   | $\infty/2$       |

For local  $\mathbb{P}^1 \times \mathbb{P}^1$ , the class depends on the sector: M along the symmetric diagonal, G along the anti-symmetric diagonal (Theorem 15.12). [PH]

*Remark 15.18* (Patterns in the  $\kappa_{\text{ch}}$ -table). Three structural patterns emerge:

- (i)  *$\kappa_{\text{ch}}$  from the compact base:* for local  $\text{CY}_3$  geometries of the form  $X = \text{Tot}(K_S \rightarrow S)$  over a smooth projective surface  $S$ , the modular characteristic is  $\kappa_{\text{ch}} = \chi(S)/2$ , giving  $\kappa_{\text{ch}}(\text{local } \mathbb{P}^2) = 3/2$  ( $\chi(\mathbb{P}^2) = 3$ ) and  $\kappa_{\text{ch}}(\text{local } \mathbb{P}^1 \times \mathbb{P}^1) = 2$  ( $\chi(\mathbb{P}^1 \times \mathbb{P}^1) = 4$ ). For geometries not of the form  $\text{Tot}(K_S)$ , the value is computed from DT invariants directly:  $\kappa_{\text{ch}}(\mathbb{C}^3) = 1$  (from the MacMahon plethystic logarithm) and  $\kappa_{\text{ch}}(\text{conifold}) = 1$  (from the single compact curve class). Note: the conifold is  $\text{Tot}(\mathcal{O}(-1)^{\oplus 2} \rightarrow \mathbb{P}^1)$ , which is *not*  $\text{Tot}(K_{\mathbb{P}^1}) = \text{Tot}(\mathcal{O}(-2) \rightarrow \mathbb{P}^1)$ , so the  $\chi(S)/2$  formula does not apply to it directly.
- (ii) *Shadow class from the quiver:* class G occurs when the quiver has a single nontrivial vertex (Jordan, KW) and class M occurs when the quiver has  $\mathbb{Z}_N$ -symmetry for  $N \geq 3$  (McKay). The conifold, despite having two vertices, is class G because the Klebanov–Witten quiver has a  $\mathbb{Z}_2$ -symmetry that terminates the tower at degree 2.
- (iii) *Wall-crossing preserves  $\kappa_{\text{ch}}$ :* the modular characteristic is chamber-independent (it depends only on the topology of the compact base, not on the stability parameter). This is a manifestation of the gauge invariance of Theorem 15.16.

### 15.13 Gaudin Hamiltonians from toric CY<sub>3</sub> collision residues

The collision residue  $r(z) = \text{Res}_{0,2}^{\text{coll}}(\Theta_A)$  of the MC element extracts, at genus 0 and degree 2, the classical  $r$ -matrix of the quantum group associated to a CY<sub>3</sub> geometry. For toric CY<sub>3</sub> threefolds this  $r$ -matrix generates commuting Gaudin Hamiltonians on configuration spaces, providing the integrable-systems face of the holographic datum.

*Construction 15.19 (Toric CY<sub>3</sub> Gaudin Hamiltonians).* Let  $X$  be a toric CY<sub>3</sub> with associated affine super Yangian  $Y(\widehat{\mathfrak{g}}_{Q_X})$  (Theorem 15.5). For  $n$  distinct points  $z_1, \dots, z_n \in \mathbb{C}$ , the *Gaudin Hamiltonians* are the pairwise sums

$$H_i = \sum_{j \neq i} \frac{\Omega_{ij}^{Y(\widehat{\mathfrak{g}}_{Q_X})}}{z_i - z_j}, \quad i = 1, \dots, n,$$

where  $\Omega_{ij}^{Y(\widehat{\mathfrak{g}}_{Q_X})} \in \text{End}(V_1 \otimes \dots \otimes V_n)$  is the image of the quadratic Casimir  $\Omega \in Y(\widehat{\mathfrak{g}}_{Q_X}) \otimes Y(\widehat{\mathfrak{g}}_{Q_X})$  acting in the  $i$ -th and  $j$ -th tensor factors.

**PROPOSITION 15.20 (Commutativity of Gaudin Hamiltonians).** The operators  $H_1, \dots, H_n$  pairwise commute:  $[H_i, H_j] = 0$  for all  $i, j$ . The proof reduces to the classical Yang–Baxter equation for  $r(z)$ , which follows from  $D^2 = 0$  in the bar complex projected to genus 0, degree 3. <sup>[PH]</sup>

*Proof.* The commutator  $[H_i, H_j]$  expands into triple sums involving  $[\Omega_{ik}, \Omega_{jk}]/(z_i - z_k)(z_j - z_k)$  and cyclic permutations. The classical Yang–Baxter equation  $[r_{12}(z_{12}), r_{13}(z_{13})] + [r_{12}(z_{12}), r_{23}(z_{23})] + [r_{13}(z_{13}), r_{23}(z_{23})] = 0$  (where  $z_{ij} = z_i - z_j$ ) forces exact cancellation. The CYBE itself is the degree-3 projection of the MC equation  $D\Theta + \frac{1}{2}[\Theta, \Theta] = 0$  at genus 0 (cf. Vol I, Theorem 20.9).  $\square$

For  $X = \mathbb{C}^3$  (the Jordan quiver), the Yangian is  $Y(\widehat{\mathfrak{gl}}_1)$  and the Casimir is the standard  $\Omega = \sum_a J_a \otimes J^a$  with  $J_a$  the generators of  $\widehat{\mathfrak{gl}}_1$ . The Gaudin system on  $n$  points then coincides with the  $\widehat{\mathfrak{gl}}_1$  Gaudin model studied by Rapcak–Soibelman–Yang–Zhao in the context of  $\mathcal{W}_{1+\infty}$  at the self-dual level. The spectral curve of the Gaudin system is the Seiberg–Witten curve of the 4d  $\mathcal{N} = 2$  theory obtained by compactification of the 5d holomorphic Chern–Simons theory on  $X$ .

**PROPOSITION 15.21 (Wall-crossing as Gaudin gauge equivalence).** Let  $\zeta, \zeta'$  be generic stability parameters in adjacent chambers. The wall-crossing gauge transformation  $\Theta_{\zeta'} = e^{\text{ad}_\alpha}(\Theta_\zeta)$  (Theorem 15.16) induces a gauge equivalence of the associated Gaudin systems: the Hamiltonians  $H_i^\zeta$  and  $H_i^{\zeta'}$  are conjugate by the gauge operator  $e^\alpha \in \exp(\mathfrak{n}_W)$ , hence have identical spectra. <sup>[PH]</sup>

*Proof.* The Gaudin Hamiltonians are extracted from the degree-2 projection of  $\Theta$ ; the gauge transformation  $e^{\text{ad}_\alpha}$  acts on this projection by conjugation, preserving commutativity and spectra.  $\square$

*Remark.* The Gaudin construction extends to the resolved conifold (Klebanov–Witten Yangian) and to local  $\mathbb{P}^2$  (McKay super Yangian), giving integrable spin chains whose Bethe ansatz equations encode DT invariants. The integrability of the Gaudin system is the dynamical manifestation of  $D^2 = 0$  in the bar complex.

## 15.14 The chiral quantum group of a toric CY<sub>3</sub>

The preceding sections establish each component of the quantum group structure independently: the CoHA bialgebra (Theorems 15.4 and 15.5), the Maulik–Okounkov  $R$ -matrix, the Drinfeld center passage from  $E_1$  to  $E_2$  (Theorem 8.28 for  $\mathbb{C}^3$ ), and the quiver-chart gluing for toric CY<sub>3</sub> (Theorem 8.46). The theorem below assembles these components into a single chiral quantum group structure for all toric CY<sub>3</sub> without compact 4-cycles. For  $\mathbb{C}^3$  the five-step functor chain (Theorem 8.15) verifies every step end-to-end; the general toric case inherits the algebraic structure from the RSYZ identification and the chart-gluing theorem, with the Drinfeld center identification conditional on Conjecture CY-C outside  $\mathbb{C}^3$ .

**THEOREM 15.22** (*Chiral quantum group structure for toric CY<sub>3</sub>*). <sup>[PH]</sup> Let  $X = X_\Sigma$  be a smooth toric CY<sub>3</sub> without compact 4-cycles, with toric quiver with potential  $(Q_X, W_X)$ . The following five-component chiral quantum group datum  $\mathcal{G}(X)$  exists and is uniquely determined by the toric fan  $\Sigma$ :

(I)  **$E_1$ -bialgebra (CoHA).** The critical CoHA

$$\mathcal{H}(Q_X, W_X) \simeq Y^+(\widehat{\mathfrak{g}}_{Q_X})$$

is the positive half of the affine super Yangian  $Y^+(\widehat{\mathfrak{g}}_{Q_X})$  (Schiffmann–Vasserot for  $\mathbb{C}^3$ ; Rapcak–Soibelman–Yang–Zhao for general toric). The CoHA carries an associative  $E_1$  multiplication (the Hall product) and a coassociative coproduct (the Yang–Zhao localization coproduct), making  $\mathcal{H}(Q_X, W_X)$  a topological bialgebra over the charge lattice  $\Gamma(X) = K_0(\text{Coh}(X))$ .

(II)  **$R$ -matrix.** The Maulik–Okounkov stable-envelope construction produces a Yang  $R$ -matrix  $R^{\text{MO}}(z) \in \text{End}(V \otimes V)((z))$  for each pair of representations  $V$  of  $Y^+(\widehat{\mathfrak{g}}_{Q_X})$ , satisfying:

- (a) the Yang–Baxter equation  $R_{12}(u-v) R_{13}(u) R_{23}(v) = R_{23}(v) R_{13}(u) R_{12}(u-v)$ ;
- (b) unitarity  $R(z) R(-z) = \text{Id}$  (from the CY condition);
- (c) wall-crossing covariance: for adjacent stability chambers  $\sigma, \sigma'$ , the  $R$ -matrices are gauge-equivalent  $R^{\sigma'}(z) = (g \otimes g) R^\sigma(z) (g^{-1} \otimes g^{-1})$  with  $g$  the MO wall-crossing operator (Proposition 8.48(ii)).

(III) **Drinfeld double.** The Drinfeld double

$$D(\mathcal{H}(Q_X, W_X)) = Y(\widehat{\mathfrak{g}}_{Q_X}) = Y^+(\widehat{\mathfrak{g}}_{Q_X}) \otimes Y^0(\widehat{\mathfrak{g}}_{Q_X}) \otimes Y^-(\widehat{\mathfrak{g}}_{Q_X}) \quad (15.1)$$

is the full affine super Yangian, where  $Y^-$  is the CoHA of the opposite quiver  $(Q_X^{\text{op}}, W_X^{\text{op}})$  and  $Y^0$  is the Cartan subalgebra generated by the charge lattice  $\Gamma(X)$ . The cross-relations between  $Y^+$  and  $Y^-$  are determined by the  $R$ -matrix of (II). For  $\mathbb{C}^3$ :  $Y(\widehat{\mathfrak{gl}}_1) \simeq \mathcal{W}_{1+\infty}$  (Procházka–Rapčák).

(IV)  **$E_1$ -chiral algebra via chart gluing.** The global  $E_1$ -chiral algebra

$$A_{X_\Sigma} = \text{hocolim}_{\Sigma(3)} \text{CoHA}(Q_\alpha, W_\alpha) \quad (15.2)$$

is assembled from the quiver-chart atlas  $\{(Q_\alpha, W_\alpha)\}_{\alpha \in \Sigma(3)}$  by the chart-gluing theorem (Theorem 8.46). Each transition is an  $E_1$  quasi-isomorphism (Proposition 8.49). The descent spectral sequence degenerates at  $E_2$  (Theorem 8.55), so the hocolim is determined by pairwise wall-crossing data alone.

- (V) **Modular characteristic and shadow tower.** The modular characteristic  $\kappa_{\text{ch}}(A_{X_\Sigma}) = \chi(S)/2$  for  $X_\Sigma = \text{Tot}(K_S \rightarrow S)$  (Proposition 15.17). It is independent of the stability chamber (Proposition 8.51(iii)). The full shadow obstruction tower  $\Theta_{A_X}$  satisfies the MC equation  $D\Theta + \frac{1}{2}[\Theta, \Theta] = 0$ , with the shadow class (G, L, C, or M) determined by the quiver combinatorics (Proposition 15.17).

**Compatibility.** The five components are mutually consistent:

- (a) The  $R$ -matrix of (II) is the collision residue of the MC element:  $r(z) = \text{Res}_{0,2}^{\text{coll}}(\Theta_{A_X})$ , and the classical Yang–Baxter equation for  $r(z)$  follows from  $D^2 = 0$  at genus 0, degree 3 (Proposition 15.20).
- (b) The Kontsevich–Soibelman wall-crossing formula is equivalent to the MC gauge equivalence  $\Theta_{\zeta'} = e^{\text{ad}_\alpha}(\Theta_\zeta)$  between adjacent chambers (Theorem 8.50).
- (c) The Costello–Li comparison  $A_{X_\Sigma}^{\text{hocolim}} \simeq_{E_1} U^{\text{ch}}(\mathfrak{g}_{X_\Sigma})$  identifies the CoHA chart-gluing with the boundary algebra of 5d holomorphic Chern–Simons on  $X_\Sigma \times \mathbb{R}^2$  (Theorem 8.46(iv)).

*Proof.* Component (I) is the combination of Theorem 15.4 ( $\mathbb{C}^3$ ) and Theorem 15.5 (general toric), which identify the critical CoHA with  $Y^+(\widehat{\mathfrak{g}}_{Q_X})$ . The coproduct is the localization coproduct of Yang–Zhao, defined by restriction to the attracting set of a one-parameter subgroup in the equivariant cohomology of the representation variety.

Component (II): the Maulik–Okounkov stable-envelope  $R$ -matrix is constructed for all toric CY<sub>3</sub> in the equivariant  $K$ -theory of Nakajima quiver varieties. The Yang–Baxter equation is proved in MO by the geometric argument (Steinberg correspondence). Unitarity follows from the CY condition: the structure function  $g(z) = \prod_i (z - h_i)/(z + h_i)$  satisfies  $g(z)g(-z) = 1$  (see equation (8.2)). Wall-crossing covariance follows from the stable-envelope wall-crossing formula of MO.

Component (III): the Drinfeld double construction is standard (Drinfeld). The identification  $D(\mathcal{H}) = Y(\widehat{\mathfrak{g}}_{Q_X})$  for toric CY<sub>3</sub> follows from the triangular decomposition of the affine super Yangian (RSYZ, building on Schiffmann–Vasserot for  $\mathbb{C}^3$ ). The opposite CoHA  $\mathcal{H}^{\text{op}}$  is the CoHA of  $(Q_X^{\text{op}}, W_X^{\text{op}})$ , which provides the negative Borel. The Cartan  $Y^0$  is the group algebra of  $\Gamma(X) = K_0(\text{Coh}(X))$ .

Component (IV) is Theorem 8.46, proved above.

Component (V): the  $\kappa_{\text{ch}}$ -values are established in Proposition 15.17. Chamber-independence follows from Proposition 8.51(iii). The MC equation for  $\Theta_{A_X}$  is proved by the same argument as for the  $\mathbb{C}^3$  case, since the MC equation is a formal consequence of  $D^2 = 0$  on  $\overline{\mathcal{M}}_{g,n}$  and does not depend on the specific toric geometry.

Compatibility (a): the collision residue extraction  $r(z) = \text{Res}_{0,2}^{\text{coll}}(\Theta_A)$  is a general construction from Vol I (Chapter 8, seven-faces formalism). The CYBE is the degree-3 projection of the MC equation at genus 0.

Compatibility (b) is Theorem 8.50.

Compatibility (c) is Theorem 8.46(iv). □

*Remark 15.23* (The  $E_2$  structure and Conjecture CY-C). Theorem 15.22 constructs the  $E_1$ -chiral algebra  $A_{X_\Sigma}$  unconditionally for all toric CY<sub>3</sub>. The  $E_2$ -braided representation category  $\text{Rep}^{E_2}(\mathcal{G}(X))$  is recovered by the Drinfeld center:  $\mathcal{Z}(\text{Rep}^{E_1}(A_{X_\Sigma})) \simeq \text{Rep}^{E_2}(Y(\widehat{\mathfrak{g}}_{Q_X}))$ . For  $\mathbb{C}^3$  this is proved (Theorem 8.28). For general toric CY<sub>3</sub>, the Drinfeld center of the CoHA representation category is identified with the braided representation category of the full affine super Yangian: this is the content of Conjecture CY-C restricted to the toric case (Conjecture 19.10). The difficulty is that the Drinfeld center does not commute with homotopy colimits (Remark 8.69, obstacle **O3**): the global  $E_2$ -braided category for a multi-chart toric CY<sub>3</sub> cannot be assembled chart-by-chart from the local Drinfeld centers.



*Remark 15.24* (Comparison:  $\mathbb{C}^3$  versus general toric CY<sub>3</sub>). For  $\mathbb{C}^3$ , the five-step functor chain (Theorem 8.15) provides stronger results:

- (i) The classical bracket is abelian (Theorem 8.17), so all noncommutative structure arises from the  $\Omega$ -deformation. For general toric CY<sub>3</sub> with orbifold singularities, the McKay quiver can have a nonabelian classical bracket.
- (ii) The  $\Omega$ -deformation is unique ( $\mathrm{HH}^2 = 1$ , Theorem 8.18). For general toric CY<sub>3</sub>, the deformation space may be multi-dimensional (spanned by the independent equivariant parameters subject to the CY constraint).
- (iii) The Drinfeld center identification is proved end-to-end for  $\mathbb{C}^3$ :  $\mathcal{Z}(\mathrm{Rep}^{E_1}(Y^+(\widehat{\mathfrak{gl}}_1))) \simeq \mathrm{Rep}^{E_2}(\mathcal{W}_{1+\infty})$ . For general toric CY<sub>3</sub>, this identification is conditional on Conjecture CY-C.
- (iv) The output algebra  $\mathcal{W}_{1+\infty}$  has a vertex algebra presentation (Procházka–Rapčák). For general toric CY<sub>3</sub>, the full Yangian  $Y(\widehat{\mathfrak{g}}_{Q_X})$  is defined as a shuffle algebra or RTT algebra, but an explicit vertex algebra presentation (strong generators, OPE) exists only for  $\mathbb{C}^3$  and the conifold.

## 15.15 The chiral quantum group equivalence for toric CY<sub>3</sub>

Theorem 15.22 assembles five components of the chiral quantum group datum for toric CY<sub>3</sub> without compact 4-cycles. The components are proved independently: the CoHA bialgebra by Schiffmann–Vasserot and RSYZ, the  $R$ -matrix by Maulik–Okounkov, the Drinfeld double by triangular decomposition, the chart gluing by descent, the shadow tower by the MC equation. These components are *a priori* separate structures on the same algebra. The Vol I chiral quantum group equivalence asserts that three of them (the vertex  $R$ -matrix, the chiral  $A_\infty$ -structure, and the chiral coproduct) determine each other on the Koszul locus. The theorem below specializes the abstract equivalence to the toric CY<sub>3</sub> setting, where both sides are independently known, yielding the strongest unconditional statement: the RSYZ bialgebra coproduct, the MO  $R$ -matrix, and the bar-differential  $A_\infty$ -structure are three faces of a single object.

**THEOREM 15.25** (*Chiral quantum group equivalence for toric CY<sub>3</sub>*). <sup>[PH]</sup> Let  $X = X_\Sigma$  be a smooth toric CY<sub>3</sub> without compact 4-cycles, with toric quiver  $(Q_X, W_X)$  and critical CoHA  $\mathcal{H}(Q_X, W_X) \simeq Y^+(\widehat{\mathfrak{g}}_{Q_X})$  (Theorem 15.5). Let  $A_X = \Phi(\mathrm{Coh}(X))$  denote the  $E_1$ -chiral algebra of Theorem 15.22(IV), with ordered bar complex  $\overline{B}^{\mathrm{ord}}(A_X) = T^c(s^{-1}\overline{A}_X)$  (deconcatenation coproduct,  $|s^{-1}v| = |v| - 1$ ). On the Koszul locus, the following three structures on  $A_X$  determine each other:

- (I) **Vertex  $R$ -matrix.** The Maulik–Okounkov stable-envelope  $R$ -matrix  $R^{\mathrm{MO}}(z) \in \mathrm{End}(V \otimes V)((z))$  of Theorem 15.22(II), satisfying the Yang–Baxter equation, unitarity  $R(z)R(-z) = \mathrm{Id}$ , and the shift condition.
- (II) **Chiral  $A_\infty$ -structure.** The bar differential on  $\overline{B}^{\mathrm{ord}}(A_X)$  determines chiral  $A_\infty$ -maps  $m_k^{\mathrm{ch}}: A_X^{\otimes k} \rightarrow A_X((\lambda_1)) \cdots ((\lambda_{k-1}))$  satisfying the Stasheff identities in the chiral endomorphism operad  $\mathrm{End}_{A_X}^{\mathrm{ch}}$ .
- (III) **Chiral coproduct.** The deconcatenation coproduct on  $\overline{B}^{\mathrm{ord}}(A_X)$  yields, via cobar recovery  $\Omega(\overline{B}^{\mathrm{ord}}(A_X)) \xrightarrow{\sim} A_X$  (on the Koszul locus), a chiral coproduct  $\Delta^{\mathrm{ch}}: A_X \rightarrow (A_X \hat{\otimes} A_X)((z))$ , coassociative up to conjugation by the Drinfeld associator  $\Phi \in \exp(\hat{\mathfrak{t}}_3)$ .

The equivalences are:

- (a) (I)  $\rightarrow$  (II): restriction of the vertex OPE to the real Stasheff associahedra  $K_n \hookrightarrow \overline{\mathrm{FM}}_n^{\mathrm{ord}}(\mathbb{C})$ ;

- (b) (II)  $\rightarrow$  (III): cobar applied to the deconcatenation coproduct on  $\overline{B}^{\text{ord}}(A_X)$ ;
- (c) (III)  $\rightarrow$  (I): the chiral Drinfeld formula  $R(z) = \sigma \circ \Delta^{\text{ch}} \circ (\Delta^{\text{ch,op}})^{-1}$ .

**Concrete identifications.** Under the RSYZ isomorphism  $\mathcal{H}(Q_X, W_X) \simeq Y^+(\widehat{\mathfrak{g}}_{Q_X})$ :

- (i) The vertex  $R$ -matrix (I) is identified with the MO stable-envelope  $R$ -matrix of the Nakajima quiver variety  $\mathfrak{M}(Q_X, \mathbf{d})$ ;
- (ii) The chiral  $A_\infty$ -maps (II) are identified with the CoHA multiplication and higher compositions, read off from the bar differential  $d_{\overline{B}^{\text{ord}}} = \sum_{k \geq 1} b_k$  where  $b_k$  is the codegree- $k$  component encoding the  $k$ -fold Hall product;
- (iii) The chiral coproduct (III) is identified with the Yang–Zhao localization coproduct on the CoHA, defined by restriction to attracting sets of one-parameter subgroups in equivariant cohomology;
- (iv) The classical  $r$ -matrix  $r(z) = \text{Res}_{0,2}^{\text{coll}}(\Theta_{A_X})$  (collision residue of the MC element) is the classical limit of  $R^{\text{MO}}(z)$ .

*Proof.* The abstract equivalence (I)  $\leftrightarrow$  (II)  $\leftrightarrow$  (III) is the Vol I chiral quantum group equivalence, proved for any  $E_1$ -chiral algebra on the Koszul locus. The content of the present theorem is the identification of each abstract vertex with the independently constructed toric CY<sub>3</sub> structure.

*Component (I):  $R$ -matrix.* The  $E_1$ -chiral algebra  $A_X$  carries a vertex  $R$ -matrix by the general theory (item (I) of that equivalence). For toric CY<sub>3</sub>, the Maulik–Okounkov stable-envelope construction provides an independent  $R$ -matrix on  $Y^+(\widehat{\mathfrak{g}}_{Q_X})$ -modules. The two coincide because both solve the same Yang–Baxter equation with the same classical limit  $r(z) = \text{Res}_{0,2}^{\text{coll}}(\Theta_{A_X})$  and the same unitarity constraint  $R(z)R(-z) = \text{Id}$  (from the CY condition, equation (8.2)). Uniqueness of the quantum  $R$ -matrix from the classical  $r$ -matrix on the Koszul locus is standard (Etingof–Kazhdan).

*Component (II):  $A_\infty$ .* The bar differential on  $\overline{B}^{\text{ord}}(A_X) = T^c(s^{-1}\bar{A}_X)$  encodes the chiral  $A_\infty$ -maps  $m_k^{\text{ch}}$  by definition. On the CoHA side, the Hall product is  $m_2^{\text{ch}}$ , and the higher compositions  $m_k^{\text{ch}}$  for  $k \geq 3$  are the higher-order contact terms arising from the bar-complex boundary strata (associahedron facets). For  $\mathbb{C}^3$ , all  $m_k^{\text{ch}}$  with  $k \geq 3$  vanish: the CoHA is strictly associative (class G, shadow depth  $r_{\max} = 2$ ). For local  $\mathbb{P}^2$  (class M), the  $m_k^{\text{ch}}$  are nonzero for all  $k$  and encode the full infinite shadow tower.

*Component (III): coproduct.* The Yang–Zhao localization coproduct on  $\mathcal{H}(Q_X, W_X)$  is coassociative and compatible with the Hall multiplication. The deconcatenation coproduct on  $\overline{B}^{\text{ord}}(A_X)$  produces, via the cobar functor  $\Omega$ , a chiral coproduct on  $A_X$  that is coassociative up to the Drinfeld associator. On the Koszul locus,  $\Omega(\overline{B}^{\text{ord}}(A_X)) \xrightarrow{\sim} A_X$  is a quasi-isomorphism (Theorem B of Vol I), and the transported coproduct is identified with the Yang–Zhao coproduct by the universal property of the localization construction (the attracting-set restriction computes the same homotopy-coherent coproduct as the bar-cobar transport).

*Concrete identification (iv).* The classical  $r$ -matrix is the collision residue of the MC element  $\Theta_{A_X}$  (Vol I seven-faces formalism). The classical limit  $R(z) = \text{Id} + \hbar r(z) + O(\hbar^2)$  holds by the MO construction (the stable envelope  $R$ -matrix is a quantization of the geometric  $r$ -matrix with classical limit given by the torus-fixed point localization).  $\square$

**COROLLARY 15.26** ( $\mathbb{C}^3$ : the complete chiral quantum group). <sup>[PH]</sup> For  $X = \mathbb{C}^3$  (Jordan quiver,  $\mathcal{H} \simeq Y^+(\widehat{\mathfrak{gl}}_1)$ ), the chiral quantum group equivalence of Theorem 15.25 specializes to:

- (i) The vertex  $R$ -matrix is the MO stable-envelope  $R$ -matrix of  $\text{Hilb}^n(\mathbb{C}^2)$ .
- (ii) The chiral  $A_\infty$ -structure is strictly associative:  $m_k^{\text{ch}} = 0$  for  $k \geq 3$  (class G,  $r_{\max} = 2$ ).
- (iii) The chiral coproduct coincides with the shuffle coproduct on  $Y^+(\widehat{\mathfrak{gl}}_1)$ .
- (iv) The Drinfeld double  $D(Y^+(\widehat{\mathfrak{gl}}_1)) = Y(\widehat{\mathfrak{gl}}_1) \simeq \mathcal{W}_{1+\infty}$  (Procházka–Rapčák).
- (v) The Drinfeld center  $\mathcal{Z}(\text{Rep}^{E_1}(Y^+(\widehat{\mathfrak{gl}}_1))) \simeq \text{Rep}^{E_2}(Y(\widehat{\mathfrak{gl}}_1))$  (Theorem 8.28).

All five components are unconditional. The  $\mathbb{C}^3$  case is the unique toric CY<sub>3</sub> where the Drinfeld center identification is proved; for general toric, component (v) is conditional on Conjecture 19.10.

*Proof.* Items (i)–(iii) follow from Theorem 15.25 specialized to the Jordan quiver. Item (ii) uses the class G classification of  $\mathbb{C}^3$  (Corollary 15.8 for the conifold; for  $\mathbb{C}^3$  itself,  $S_r = 0$  for  $r \geq 3$  because the single-field Heisenberg OPE has  $r_{\max} = 2$ ). Item (iv) is the triangular decomposition  $Y = Y^+ \otimes Y^0 \otimes Y^-$  combined with the Procházka–Rapčák identification. Item (v) is Theorem 8.28.  $\square$

**COROLLARY 15.27** (*Conifold: wall-crossing preserves the triangle*). <sup>[PH]</sup> For the resolved conifold  $X = \mathcal{O}(-1) \oplus \mathcal{O}(-1) \rightarrow \mathbb{P}^1$  (Klebanov–Witten quiver), the chiral quantum group equivalence of Theorem 15.25 holds in both stability chambers  $\zeta_1 > 0$  and  $\zeta_1 < 0$ . The wall-crossing gauge transformation  $\Theta_{\text{II}} = e^{\text{ad}_\alpha}(\Theta_{\text{I}})$  of Theorem 15.7 preserves each vertex of the triangle:

- (i) The  $R$ -matrices in the two chambers are gauge-equivalent:  $R^{\text{II}}(z) = (g \otimes g) R^{\text{I}}(z) (g^{-1} \otimes g^{-1})$ .
- (ii) The  $A_\infty$ -structures are  $A_\infty$ -quasi-isomorphic via the transfer induced by the gauge equivalence.
- (iii) The chiral coproducts are intertwined by the same gauge operator  $g = e^\alpha$ .

In particular,  $\kappa_{\text{ch}} = 1$  is chamber-independent (Proposition 15.17).

*Proof.* The MC gauge equivalence  $\Theta_{\text{II}} = e^{\text{ad}_\alpha}(\Theta_{\text{I}})$  preserves the MC equation and hence each derived structure. The  $R$ -matrix transforms by conjugation (compatibility (a) of Theorem 15.22). The  $A_\infty$  transfer is the homotopy transfer theorem applied to the quasi-isomorphism between the bar complexes in the two chambers (the gauge transformation is a filtered quasi-isomorphism on  $\overline{B}^{\text{ord}}$ ). The coproduct intertwining follows from the coassociativity of deconcatenation and the naturality of the cobar functor under quasi-isomorphisms of dg coalgebras.  $\square$

**Remark 15.28** (Extension to compact CY<sub>3</sub>). The toric hypothesis enters Theorem 15.22 in four places:

- (a) *Existence of CoHA*: the critical CoHA is constructed for quivers with potential, which are available for toric CY<sub>3</sub> via the McKay correspondence. For compact CY<sub>3</sub> without torus action (quintic, generic  $K3 \times E$ ), no CoHA construction is available; the entry point is instead the factorization envelope of the polyvector-field Lie conformal algebra (Costello–Li).
- (b) *RSYZ identification*:  $\mathcal{H}(Q_X, W_X) \simeq Y^+(\widehat{\mathfrak{gl}}_{Q_X})$  is specific to toric geometry (the shuffle-algebra presentation uses equivariant localization).
- (c) *MO  $R$ -matrix*: the stable-envelope construction requires a torus action on the Nakajima quiver variety. For compact CY<sub>3</sub>, the  $R$ -matrix must come from a different source (the chain-level  $\mathbb{S}^3$ -framing of Conjecture 8.68).
- (d) *Chart gluing*: the toric fan provides a finite atlas; for general CY<sub>3</sub>, the tilting-chart cover is conjectural (Conjecture 8.42).

The extension to compact  $CY_3$  (including  $K3 \times E$ , the quintic, and Enriques  $\times E$ ) is the content of Conjecture 8.68, with the chain-level  $\mathbb{S}^3$ -framing as the single remaining obstruction.

## Chapter 16

# Fukaya Categories

Symplectic geometry gives Calabi–Yau categories in every dimension relevant to Vol III, but the CY-to-chiral functor sees a sharp boundary at  $d = 3$ . For  $d = 2$  the Fukaya category feeds  $\Phi$  unconditionally and produces an  $E_2$ -chiral algebra. For  $d = 3$  the functor stops at the ordered  $E_1$  layer unless the chain-level  $\mathbb{S}^3$ -framing exists. The Fukaya category  $\mathrm{Fuk}(X)$  is the test case for that boundary: its objects are Lagrangians, its morphisms are Floer complexes, its higher compositions count pseudoholomorphic polygons, and for Calabi–Yau  $X$  it carries the cyclic structure required by Chapter 3. The modular characteristic  $\kappa_{\mathrm{cat}}$  extracted from  $\Phi(\mathrm{Fuk}(X))$  records symplectic information carried by  $X$ .

### 16.1 The Fukaya category: definition and $A_\infty$ -structure

*Definition 16.1 (Fukaya category).* Let  $(X^{2n}, \omega)$  be a compact symplectic manifold. The *Fukaya category*  $\mathrm{Fuk}(X)$  is the  $A_\infty$ -category whose:

- (i) *Objects* are compact, oriented, graded Lagrangian submanifolds  $L \subset X$  equipped with a flat  $U(1)$ -connection (a local system);
- (ii) *Morphism spaces* are Floer cochain complexes: for transverse  $L_0, L_1$ ,

$$\mathrm{Hom}_{\mathrm{Fuk}(X)}(L_0, L_1) = \mathrm{CF}^\bullet(L_0, L_1) = \bigoplus_{p \in L_0 \cap L_1} \Lambda_{\mathrm{nov}} \cdot p,$$

where  $\Lambda_{\mathrm{nov}} = \mathbb{C}[[T]][T^{-1}]$  is the Novikov field and the grading on  $p$  is the Maslov index;

- (iii) *Higher compositions* are  $\mu_k : \mathrm{CF}^\bullet(L_0, L_1) \otimes \cdots \otimes \mathrm{CF}^\bullet(L_{k-1}, L_k) \rightarrow \mathrm{CF}^\bullet(L_0, L_k)[2-k]$ , defined by counting pseudoholomorphic  $(k+1)$ -gons:

$$\mu_k(p_1, \dots, p_k) = \sum_{\substack{u: D^2 \rightarrow X \\ \bar{\partial}_J u = 0}} T^{\int u^* \omega} \cdot q(u)$$

where  $q(u)$  is the output intersection point and the sum is over  $J$ -holomorphic disks  $u$  with boundary on  $L_0 \cup \cdots \cup L_k$  and corners at  $p_1, \dots, p_k$ .

The  $A_\infty$ -relations  $\sum_{i,j} \pm \mu_{k-j+1}(\dots, \mu_j(\dots), \dots) = 0$  follow from the compactification of moduli spaces of pseudoholomorphic polygons (Fukaya–Oh–Ohta–Ono).

*Remark 16.2* (Analytic foundations). The rigorous construction of  $\mathrm{Fuk}(X)$  requires virtual perturbation techniques (Kuranishi structures, polyfolds, or algebraic approaches) to handle transversality for the moduli spaces of holomorphic polygons. The foundational references are Fukaya–Oh–Ohta–Ono for Kuranishi structures, Hofer–Wysocki–Zehnder for polyfolds, and Pardon for the algebraic virtual fundamental class. For this volume, the  $A_\infty$ -structure on  $\mathrm{Fuk}(X)$  is assumed constructed; the CY-to-chiral functor  $\Phi$  takes this structure as input.

**PROPOSITION 16.3** (*CY structure on the Fukaya category*). <sup>[PE]</sup> For a compact CY manifold  $(X^{2n}, \omega, \Omega)$  of complex dimension  $n$ , the Fukaya category  $\mathrm{Fuk}(X)$  is CY of dimension  $n$ :

- (i) The CY pairing  $\langle \cdot, \cdot \rangle : \mathrm{CF}^\bullet(L_0, L_1) \otimes \mathrm{CF}^\bullet(L_1, L_0) \rightarrow \Lambda_{\mathrm{nov}}[-n]$  is the Poincaré duality pairing on Lagrangian intersections;
- (ii) The cyclic invariance of  $\mu_k$  with respect to this pairing follows from the boundary of a pseudoholomorphic polygon being a cycle on  $L_0 \cup \cdots \cup L_k$ ;
- (iii) Non-degeneracy follows from Poincaré duality on compact oriented manifolds.

## 16.2 CY 1-folds: elliptic curves

*Example 16.4* (*Elliptic curve*). For an elliptic curve  $E_\tau = \mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$  with the standard Kähler form  $\omega = \frac{i}{2}dz \wedge d\bar{z}$ :

- (i)  $\mathrm{Fuk}(E_\tau)$  is generated by the Lagrangian circles  $L_\alpha = \{z \in E_\tau : \Im(e^{-i\alpha}z) = c\}$ ;
- (ii)  $\mathrm{Fuk}(E_\tau)$  is CY of dimension 1;
- (iii) The Hochschild homology  $\mathrm{HH}_\bullet(\mathrm{Fuk}(E_\tau)) \simeq H_{\mathrm{dR}}^*(E_\tau)$  recovers the de Rham cohomology (Abouzaid);
- (iv) The CY-to-chiral functor produces the Heisenberg vertex algebra  $H_k$  at level  $k = \mathrm{vol}(E_\tau)$ ;
- (v) The modular characteristic is  $\kappa_{\mathrm{cat}}(\Phi(\mathrm{Fuk}(E_\tau))) = k$  (the level, by the Heisenberg formula).

*Remark 16.5* (Homological mirror symmetry for  $E_\tau$ ). Polishchuk–Zaslow proved  $\mathrm{Fuk}(E_\tau) \simeq D^b(\mathrm{Coh}(E_{\tau'}))$  where  $\tau' = -1/\tau$  (mirror elliptic curve). Under this equivalence, the Heisenberg algebra  $H_k$  is the chiral algebra of the free boson compactified on the dual circle. The modular characteristic is preserved:  $\kappa_{\mathrm{cat}}(\Phi(\mathrm{Fuk}(E_\tau))) = \kappa_{\mathrm{cat}}(\Phi(D^b(\mathrm{Coh}(E_{\tau'})))) = k$ .

## 16.3 CY 2-folds: K3 and abelian surfaces

**PROPOSITION 16.6** (*K3 surface*). <sup>[PH]</sup> For a K3 surface  $S$  with the Ricci-flat Kähler metric:

- (i)  $\mathrm{Fuk}(S)$  is CY of dimension 2;
- (ii) The functor  $\Phi$  (Theorem CY-A<sub>2</sub>) produces an  $E_2$ -chiral algebra  $A_{\mathrm{Fuk}(S)}$ ;
- (iii) The modular characteristic is  $\kappa_{\mathrm{cat}}(\Phi(\mathrm{Fuk}(S))) = \chi(\mathcal{O}_S) = 2$ ;
- (iv) The Hochschild homology  $\mathrm{HH}_\bullet(\mathrm{Fuk}(S)) \simeq H^{*+2}(S, \mathbb{C}) \simeq \mathbb{C}^{24}$  (by HKR and CY duality);
- (v) The  $R$ -matrix  $\mathcal{R}_{\mathrm{Fuk}(S)}(z) = 1 + \frac{\kappa_{\mathrm{cat}}\Omega}{z} + O(1)$  with  $\kappa_{\mathrm{cat}} = \chi(\mathcal{O}_S) = 2$  encodes the Mukai pairing  $\Omega$  on  $H^*(S, \mathbb{Z})$ .

*Proof.* Item (i):  $S$  is a compact Kähler manifold with  $c_1(S) = 0$  and  $\dim_{\mathbb{C}} S = 2$ , so  $\text{Fuk}(S)$  is  $\text{CY}_2$  (Proposition 16.3).

Item (ii): Theorem CY-A<sub>2</sub> applies since  $d = 2$  and the  $\mathbb{S}^2$ -framing is unconditional.

Item (iii): the CY Euler characteristic is  $\chi^{\text{CY}}(\text{Fuk}(S)) = \chi(\mathcal{O}_S) = 1 - 0 + 1 = 2$  (using  $h^{0,0} = 1$ ,  $h^{1,0} = 0$ ,  $h^{2,0} = 1$  for  $K_3$ ). By Theorem CY-D (Theorem 10.1),  $\kappa_{\text{cat}} = \chi^{\text{CY}} = 2$ .

Item (iv): HKR gives  $\text{HH}_k(\text{Perf}(S)) \simeq \bigoplus_{p-q=k} H^q(S, \Omega^p)$ . The Hodge diamond of  $K_3$  has  $h^{p,q}(S) = 1, 0, 1, 0, 20, 0, 1, 0, 1$  (reading  $p + q = 0, 1, \dots, 4$ ), giving total  $\dim \text{HH}_{\bullet} = 24$ . CY duality  $\text{HH}^{\bullet} \simeq \text{HH}_{\bullet+2}$  shifts by  $d = 2$ .

Item (v): the  $R$ -matrix at degree  $(1, 1)$  of  $B^{\text{ord}}(A)$  has leading term  $\frac{\kappa_{\text{cat}} \Omega}{z}$  where  $\Omega \in \text{End}(V \otimes V)$  is the Casimir of the Mukai lattice  $\Lambda_{K3} = U^3 \oplus E_8(-1)^2$ . The subleading terms encode higher-order OPE data.  $\square$

*Example 16.7 (Abelian surface).* For an abelian surface  $A = E_1 \times E_2$  (product of two elliptic curves):

- (i)  $\text{Fuk}(A)$  is CY of dimension 2;
- (ii)  $\kappa_{\text{cat}}(\Phi(\text{Fuk}(A))) = \chi(\mathcal{O}_A) = 0$  (since  $h^{1,0}(A) = 2$ ,  $h^{2,0}(A) = 1$ , giving  $\chi = 1 - 2 + 1 = 0$ );
- (iii) The modular characteristic vanishes:  $\kappa_{\text{cat}} = 0$ , so the bar complex is uncurved at all genera ( $d^2 = 0$  strictly).

The abelian surface is the  $\text{CY}_2$  analogue of the critical point  $c = 26$  for Virasoro: the vanishing of  $\kappa_{\text{cat}}$  makes the genus tower trivial at the scalar level. The full shadow obstruction tower may still be nontrivial at higher degrees.

## 16.4 CY 3-folds

For  $d = 3$ , the CY-to-chiral functor  $\Phi$  is conditional on the chain-level  $\mathbb{S}^3$ -framing (Conjecture CY-A<sub>3</sub>). When  $\Phi$  exists, the chiral algebra is  $E_1$  (not  $E_2$ ): the structural  $E_2$ -obstruction from the antisymmetric Euler form (Proposition 8.38) is generically nontrivial, while the topological  $\mathbb{S}^3$ -framing obstruction vanishes ( $\pi_3(BU) = 0$  by Bott periodicity).

*Remark 16.8* (Bott periodicity verification). The stable unitary classifying space satisfies  $\Omega BU \simeq U$ , so

$$\pi_k(BU) \cong \pi_{k-1}(U).$$

Bott periodicity for  $U$  gives

$$\pi_{2m-1}(U) = \mathbb{Z}, \quad \pi_{2m}(U) = 0 \quad (m \geq 1),$$

hence

$$\pi_{2m}(BU) = \mathbb{Z}, \quad \pi_{2m+1}(BU) = 0.$$

In the low dimensions relevant here:

$$\pi_2(BU) = \mathbb{Z}, \quad \pi_3(BU) = 0, \quad \pi_4(BU) = \mathbb{Z}.$$

These groups have different meanings in adjacent CY dimensions. At  $d = 2$ , the nonzero class  $\pi_2(BU) = \mathbb{Z}$  records the native braiding parameter of the surface-level  $E_2$  theory, so Theorem CY-A<sub>2</sub> is not a vanishing statement. At  $d = 3$ , the vanishing of  $\pi_3(BU)$  removes only the topological framing obstruction; the native  $E_2$  obstruction is the antisymmetric Euler form of Proposition 8.38. At  $d = 4$ , the group  $\pi_4(BU) = \mathbb{Z}$  is an obstruction group for the  $\mathbb{S}^4$ -framing, not a guarantee of an  $E_2$  enhancement: Chapter 7 still yields only the  $E_1$ -stabilized conclusion with extra shifted data.

CONJECTURE 16.9 (*CY<sub>3</sub> Fukaya chiral algebra*). <sup>[CJ]</sup> For a compact CY threefold  $X$  with Fukaya category  $\mathrm{Fuk}(X)$ :

- (i) The CY-to-chiral functor  $\Phi(\mathrm{Fuk}(X))$  produces an  $E_1$ -chiral algebra  $A_{\mathrm{Fuk}(X)}$ , conditional only on the chain-level  $\mathbb{S}^3$ -framing (the topological obstruction already vanishes);
- (ii) The modular characteristic satisfies  $\kappa_{\mathrm{cat}}(\Phi(\mathrm{Fuk}(X))) = \chi(\mathcal{O}_X)$ ;
- (iii) The braided monoidal category  $\mathcal{Z}(\mathrm{Rep}^{E_1}(A_{\mathrm{Fuk}(X)}))$  (recovered via the Drinfeld center, Chapter 13) encodes the DT/GW invariants of  $X$ ;
- (iv) The open-string sector ( $\mathrm{Fuk}(X)$  with Lagrangian boundary conditions) connects to the Swiss-cheese structure of Volume II.

*Remark 16.10* (What remains to remove the CY<sub>3</sub> condition). The topological part of Conjecture 16.9 is already gone: both Bott periodicity and the symplectic reduction path give vanishing of the  $\mathbb{S}^3$ -framing obstruction. The remaining gap is chain-level. One must construct an explicit trivialization of the negative-cyclic CY class on the cyclic bar complex of  $\mathrm{Fuk}(X)$  and prove that this trivialization intertwines the full family of polygon operations  $\mu_k$ . Equivalently, the missing datum is an  $\mathbb{S}^3$ -framing on  $\mathrm{HH}_\bullet(\mathrm{Fuk}(X))$  compatible with the Connes operator and the cyclic  $A_\infty$  structure. Two plausible routes are known: transfer of a chain-level trivialization across a proved HMS equivalence, or a direct continuation-map construction on the Fukaya cyclic bar complex. Neither route is completed in this manuscript, so the condition is reformulated sharply but not removed.

*Example 16.11* (*Quintic threefold*). For the quintic threefold  $X_5 \subset \mathbb{P}^4$  defined by a generic degree-5 polynomial:

- (i)  $\chi(X_5) = -200$  (topological Euler characteristic);
- (ii)  $\chi(\mathcal{O}_{X_5}) = 0$  (since  $h^{0,0} = 1, h^{1,0} = 0, h^{2,0} = 0, h^{3,0} = 1$ , giving  $\chi = 1 - 0 + 0 - 1 = 0$ );
- (iii) The predicted modular characteristic is  $\kappa_{\mathrm{cat}} = \chi(\mathcal{O}_{X_5}) = 0$ .

The vanishing  $\kappa_{\mathrm{cat}} = 0$  for the quintic is the CY<sub>3</sub> analogue of the abelian surface ( $\kappa_{\mathrm{cat}} = 0$  for CY<sub>2</sub>). The genus tower is trivial at the scalar level; the full shadow obstruction tower may be nontrivial if higher-degree shadows are nonzero.

*Example 16.12* (*Resolved conifold*). The resolved conifold  $\mathcal{O}(-1) \oplus \mathcal{O}(-1) \rightarrow \mathbb{P}^1$  is non-compact with  $\chi_{\mathrm{top}} = 2$ . The wrapped Fukaya category  $\mathcal{W}(\mathrm{conifold})$  is generated by a single Lagrangian  $S^3$ -cycle. The chiral algebra has  $\kappa_{\mathrm{cat}} = 1$ , and the CoHA of the conifold (Chapter 15) recovers the affine Yangian  $Y(\widehat{\mathfrak{gl}}_1)$  at level 1.

## 16.5 Wrapped Fukaya categories

*Definition 16.13* (*Wrapped Fukaya category*). For a Liouville manifold  $(X, \lambda)$  (an exact symplectic manifold with a convex end), the *wrapped Fukaya category*  $\mathcal{W}(X)$  modifies  $\mathrm{Fuk}(X)$  by allowing wrapping at infinity: the Floer differential and higher compositions include Hamiltonian orbits of arbitrarily large period along the contact boundary  $\partial X$ . The morphism space is the direct limit

$$\mathrm{Hom}_{\mathcal{W}(X)}(L_0, L_1) = \varinjlim_{H \rightarrow \infty} \mathrm{CF}^\bullet(L_0, \phi_H(L_1))$$

where  $\phi_H$  is the time-1 flow of a Hamiltonian  $H$  that is linear with increasing slope at infinity.



**THEOREM 16.14** (*Abouzaid: cotangent bundles*). <sup>[PE]</sup> For a closed oriented manifold  $M$ , the wrapped Fukaya category of the cotangent bundle satisfies

$$\mathcal{W}(T^*M) \simeq \text{Mod}_{C_*(\Omega M)},$$

where  $C_*(\Omega M)$  is the chain algebra of the based loop space. Under this equivalence:

- (i) The zero section  $M \hookrightarrow T^*M$  corresponds to the trivial module  $\mathbb{C}$  over  $C_*(\Omega M)$ ;
- (ii) The wrapping corresponds to the bar differential on  $B(C_*(\Omega M))$ ;
- (iii) The CY pairing (when  $M$  is orientable of dimension  $n$ ) comes from Poincaré duality on  $M$ .

*Remark 16.15* (Wrapped categories and the CY-to-chiral functor). The wrapped Fukaya category is smooth but typically not proper (morphism spaces are infinite-dimensional). The CY-to-chiral functor  $\Phi$  requires a cyclic  $A_\infty$ -structure with a non-degenerate pairing; for non-compact targets, this requires either:

- (a) A compactly-supported variant (using the compact Fukaya category  $\text{Fuk}_{\text{cpt}}(X)$  as a subquotient of  $\mathcal{W}(X)$ );
- (b) An analytic completion (the sewing envelope of Volume I, MC<sub>5</sub> analytic HS-sewing lane) to handle the non-compact directions.

For cotangent bundles  $T^*M$  with  $M$  compact, approach (a) suffices:  $\text{Fuk}_{\text{cpt}}(T^*M)$  is proper, and  $\Phi$  produces the chiral algebra of the  $\sigma$ -model on  $M$ .

## 16.6 Homological mirror symmetry and the CY-to-chiral functor

**PROPOSITION 16.16** (*HMS compatibility with  $\Phi$* ). <sup>[CD]</sup> Let  $X$  and  $X^\vee$  be mirror CY manifolds related by homological mirror symmetry  $\text{Fuk}(X) \simeq D^b(\text{Coh}(X^\vee))$ . Then the CY-to-chiral functor is compatible with HMS:

$$\Phi(\text{Fuk}(X)) \simeq \Phi(D^b(\text{Coh}(X^\vee)))$$

as  $E_2$ -chiral algebras (for  $d = 2$ ) or  $E_1$ -chiral algebras (for  $d = 3$ , conditional on  $\mathbb{S}^3$ -framing for both sides). In particular, the modular characteristics agree:  $\kappa_{\text{cat}}(\Phi(\text{Fuk}(X))) = \kappa_{\text{cat}}(\Phi(D^b(\text{Coh}(X^\vee))))$ .

*Remark 16.17* (Status of HMS compatibility). The compatibility follows formally from the fact that  $\Phi$  is Morita invariant: equivalent CY categories produce equivalent chiral algebras. The conditional content is HMS itself, which is proved for:

- (a) Elliptic curves (Polishchuk–Zaslow);
- (b) Abelian varieties (Fukaya, Kontsevich–Soibelman);
- (c) K3 surfaces (Seidel, Smith; partial results);
- (d) Quintic threefold (Sheridan, for a suitable formulation);
- (e) Toric varieties (Abouzaid, Fang–Liu–Treumann–Zaslow).

For cases where HMS is proved, the compatibility of  $\Phi$  with HMS is an unconditional theorem.

## 16.7 The quantum chiral algebra of a Fukaya category

*Construction 16.18 (Assembling  $\Phi(\mathrm{Fuk}(X))$ ).* For a CY manifold  $X$  of dimension  $d$ , the quantum chiral algebra  $A_{\mathrm{Fuk}(X)} = \Phi(\mathrm{Fuk}(X))$  is assembled in three steps (matching the abstract construction of Theorem CY-A<sub>2</sub> in the proved  $d = 2$  case, and of the CY-A<sub>3</sub> programme at  $d = 3$ ):

1. *Cyclic  $A_\infty$ -extraction:* The Fukaya  $A_\infty$ -structure  $\{\mu_k\}_{k \geq 1}$  and Poincaré pairing give the input cyclic  $A_\infty$ -algebra (Chapter 3);
2. *Hochschild-to-chiral bridge:* The categorical Hochschild cochains  $\mathrm{CC}^\bullet(\mathrm{Fuk}(X), \mathrm{Fuk}(X))$  carry the Gerstenhaber bracket and Connes operator; under the bar dictionary these map to the chiral Hochschild cochains of the output chiral algebra, and on cohomology give the Lie conformal algebra  $L_{\mathrm{Fuk}(X)}$  (Chapter 4 and Proposition 16.19);
3. *Factorization envelope:* The factorization envelope of  $L_{\mathrm{Fuk}(X)}$  on a curve  $C$  gives the chiral algebra  $A_{\mathrm{Fuk}(X)}$  on  $\mathrm{Ran}(C)$ .

The bar complex  $B(A_{\mathrm{Fuk}(X)})$ , the shadow obstruction tower  $\Theta_{A_{\mathrm{Fuk}(X)}}$ , and the modular characteristic  $\kappa_{\mathrm{cat}}$  are then computed from  $A_{\mathrm{Fuk}(X)}$  via the Volume I machinery.

**PROPOSITION 16.19 (Fukaya Hochschild bridge).** <sup>[PH]</sup> Let  $X$  be a compact CY surface and set  $A_{\mathrm{Fuk}(X)} = \Phi(\mathrm{Fuk}(X))$ . Then:

- (i) The categorical Hochschild chains and the ordered chiral bar complex agree:

$$\mathrm{CC}_\bullet(\mathrm{Fuk}(X)) \xrightarrow{\sim} B(A_{\mathrm{Fuk}(X)})$$

as factorization coalgebras on  $\mathrm{Ran}(C)$ ;

- (ii) The induced map on cohomology

$$\mathrm{HH}^\bullet(\mathrm{Fuk}(X)) \longrightarrow \mathrm{ChirHoch}^*(A_{\mathrm{Fuk}(X)})$$

sends the Gerstenhaber bracket to the chiral convolution bracket and, under the CY<sub>2</sub> identification  $\mathrm{HH}^\bullet(\mathrm{Fuk}(X)) \simeq \mathrm{HH}_{2-\bullet}(\mathrm{Fuk}(X))$ , sends the Connes operator to the modular differential;

- (iii) The categorical Hochschild cochains map to the chiral derived center:

$$\mathrm{CC}^\bullet(\mathrm{Fuk}(X), \mathrm{Fuk}(X)) \longrightarrow \mathrm{RHom}(\Omega B(A_{\mathrm{Fuk}(X)}), A_{\mathrm{Fuk}(X)}) = Z_{\mathrm{ch}}^{\mathrm{der}}(A_{\mathrm{Fuk}(X)}).$$

Thus the bulk operators seen categorically by  $\mathrm{Fuk}(X)$  survive as bulk operators of the chiral algebra.

*Proof.* This is Theorem 22.22 specialized to  $C = \mathrm{Fuk}(X)$ . Part (i) is the bar dictionary. For part (ii), the CY<sub>2</sub> pairing identifies Hochschild cochains with shifted chains, so the Gerstenhaber bracket is transported to the convolution bracket on  $\mathrm{Hom}_{\mathrm{Ran}}(B(A_{\mathrm{Fuk}(X)}), A_{\mathrm{Fuk}(X)})$ , while the Connes operator corresponds to the degree-preserving component of the bar differential, namely the modular differential. Part (iii) is the cochain-level map of Theorem 22.22(iii), whose target is the chiral derived center by definition.  $\square$

*Remark 16.20* (Secondary proof sketch for the Fukaya Hochschild bridge). Fix a split-generator of  $\mathrm{Fuk}(X)$  and a cyclic  $A_\infty$  model  $A_L$ . Then one can rederive Proposition 16.19 directly on that model:  $\mathrm{CC}_\bullet(\mathrm{Fuk}(X))$  is the cyclic bar coalgebra of  $A_L$ , the ordered deconcatenation coproduct on that bar coalgebra produces the same convolution complex that defines the chiral Hochschild cochains, and the cyclic rotation operator defining Connes  $B$  becomes the genus-preserving part of the chiral bar differential after the factorization envelope. This is a genuine second route because it uses only a concrete cyclic  $A_\infty$  model and the bar-cobar adjunction, not Theorem 22.22. The remaining bookkeeping is the degree-by-degree sign comparison between the cyclic bar rotation and the modular differential.

**PROPOSITION 16.21** (*CY-to-chiral:  $d = 2$  proved,  $d = 3$  conditional*). <sup>[CD]</sup> The CY-to-chiral functor  $\Phi$  applied to Fukaya categories:

- (i) For  $d = 2$  ( $K3$ , abelian surfaces):  $\Phi(\mathrm{Fuk}(X))$  is an  $E_2$ -chiral algebra. All three steps of Construction 16.18 are unconditional. The relevant Bott group is  $\pi_2(BU) = \mathbb{Z}$ , which records the native braiding parameter of the surface theory rather than an obstruction to existence;
- (ii) For  $d = 3$  (CY threefolds): conditional on the chain-level  $\mathbb{S}^3$ -framing,  $\Phi(\mathrm{Fuk}(X))$  produces an  $E_1$ -chiral algebra. Steps 1–2 are unconditional and already determine the ordered  $E_1$  sector. The topological  $\mathbb{S}^3$ -framing obstruction vanishes ( $\pi_3(BU) = 0$  by Bott periodicity), but the structural  $E_2$ -obstruction from the antisymmetric Euler form is generically nonzero (Proposition 8.38). The braiding is recovered via the Drinfeld center  $\mathcal{Z}(\mathrm{Rep}^{E_1}(A))$  (Chapter 13).

*Proof.* For  $d = 2$ , the operative input is Theorem CY-A<sub>2</sub>: it proves directly that a  $\mathrm{CY}_2$  category produces an  $E_2$ -chiral algebra, and Construction 16.18 applies without any extra  $d = 3$ -type hypothesis. The Bott computation of Remark 16.8 shows that  $\pi_2(BU) = \mathbb{Z}$ , not 0; in dimension 2 that group measures the native braiding parameter already built into the  $E_2$  output.

For  $d = 3$ , there are two independent topological vanishing arguments. The first is the Bott path:  $\pi_3(BU) = \pi_2(U) = 0$ . The second is the symplectic path:  $\mathrm{CY}_3$  Serre duality gives an antisymmetric pairing on  $\mathrm{Ext}^1(E, E)$ , reducing the structure group to  $\mathrm{Sp}(2m)$ , and  $\pi_3(B\mathrm{Sp}(2m)) = \pi_2(\mathrm{Sp}(2m)) = 0$ . Thus the topological  $\mathbb{S}^3$ -framing obstruction vanishes universally. The reason  $\mathrm{CY}_3$  Fukaya algebras are  $E_1$  rather than native  $E_2$  is therefore structural, not topological: Serre duality forces  $\chi(E, F) = -\chi(F, E)$ , and Proposition 8.38 shows that this antisymmetric Euler form obstructs the  $E_2$  promotion. Steps 1–2 of Construction 16.18 therefore survive unconditionally and determine the ordered  $E_1$  sector. The passage from that ordered sector to the full chiral algebra  $A_{\mathrm{Fuk}(X)} = \Phi(\mathrm{Fuk}(X))$  remains conditional on the chain-level  $\mathbb{S}^3$ -framing, and the braided structure is recovered only afterwards through the Drinfeld center.  $\square$

*Remark 16.22* (Secondary proof paths for Proposition 16.21). Item (i) admits an independent B-model verification whenever HMS is proved: transport the CY-to-chiral result from  $D^b(\mathrm{Coh}(X^\vee))$  across the equivalence of Propositions 16.16 and 16.23. Item (ii) already contains two independent topological proofs inside the proposition itself: the Bott periodicity path through  $B\mathrm{U}$  and the Serre-duality reduction to  $\mathrm{Sp}$ . What neither route supplies is the missing chain-level trivialization compatible with the Fukaya  $A_\infty$  operations.

## 16.8 Homological mirror symmetry and chiral algebra equivalence

Kontsevich’s homological mirror symmetry conjecture identifies the symplectic and algebraic categorifications of a CY manifold:  $\mathrm{Fuk}(X) \simeq D^b(\mathrm{Coh}(X^\vee))$ . Under the CY-to-chiral functor  $\Phi$ , this becomes an equivalence of quantum chiral algebras.

**PROPOSITION 16.23** (*HMS implies chiral algebra equivalence*). <sup>[CD]</sup> Let  $X$  and  $X^\vee$  be mirror CY manifolds of dimension  $d$ . If the HMS equivalence  $\mathrm{Fuk}(X) \simeq D^b(\mathrm{Coh}(X^\vee))$  holds as an equivalence of  $\mathrm{CY}_d$

$A_\infty$ -categories (preserving the cyclic structure), then the CY-to-chiral functor produces an equivalence of quantum chiral algebras:

$$A_{\text{Fuk}(X)} \simeq A_{D^b(\text{Coh}(X^\vee))}.$$

In particular, the modular characteristics agree:  $\kappa_{\text{cat}}(\text{Fuk}(X)) = \kappa_{\text{cat}}(D^b(\text{Coh}(X^\vee)))$ , the bar complexes are quasi-isomorphic as factorization coalgebras, and the shadow obstruction towers coincide at all degrees.

*Proof sketch.* The CY-to-chiral functor  $\Phi$  factors through the cyclic  $A_\infty$ -structure (Chapter 3). An equivalence of  $\text{CY}_d$  categories that preserves the cyclic pairing induces an isomorphism of cyclic bar complexes, hence of factorization coalgebras after applying the factorization envelope. The chiral algebra is determined by its factorization coalgebra (Volume I, Theorem A), so the chiral algebras are equivalent. The modular characteristic  $\kappa_{\text{cat}}$  is a derived invariant of the CY category (Theorem 10.1), hence preserved by any CY equivalence.  $\square$

*Remark 16.24* (Numerical HMS tests via shadow invariants). The shadow obstruction tower provides computable invariants that can be independently evaluated on the symplectic (A-model) and algebraic (B-model) sides. For CY surfaces, the modular characteristic  $\kappa_{\text{cat}}$  and the shadow depth  $r_{\text{max}}$  are accessible on both sides:

- (i) *A-model:*  $\kappa_{\text{cat}}$  is computed from the Fukaya category's cyclic bar complex; the shadow depth is determined by the  $A_\infty$ -operations  $\mu^k$  (shadow depth  $r_{\text{max}} = k_0$  when  $\mu^k = 0$  for all  $k > k_0$  up to homotopy);
- (ii) *B-model:*  $\kappa_{\text{cat}} = \chi(\mathcal{O}_{X^\vee})$  from algebraic geometry; the shadow depth is determined by the Ext algebra structure on  $D^b(\text{Coh}(X^\vee))$ .

Agreement of these invariants provides a quantitative test of HMS beyond Hodge-number matching.

*Remark 16.25* (SYZ fibrations and the Fukaya–chiral correspondence). For CY threefolds admitting a special Lagrangian  $T^3$ -fibration  $\pi: X \rightarrow B$  (the SYZ picture), the Fukaya category decomposes along the base: the fiber Fukaya categories  $\text{Fuk}(\pi^{-1}(b))$  for smooth fibers  $b \in B_{\text{sm}}$  are  $\text{CY}_0$  (zero-dimensional CY categories, i.e., semisimple), while the singular fibers contribute non-trivial  $A_\infty$ -operations. The CY-to-chiral functor applied fiberwise produces a family of chiral algebras over  $B$ , with the singular-fiber contributions encoding instanton corrections to the mirror map. The total chiral algebra  $A_{\text{Fuk}(X)}$  is assembled as a homotopy colimit over the base, paralleling the stability-condition assembly of Chapter 17.

## 16.9 Open-string sector and the Swiss-cheese structure

The Fukaya category with a choice of Lagrangian  $L$  provides the open-string data for the Volume II Swiss-cheese structure: the endomorphism algebra  $\text{End}_{\text{Fuk}}(L)$  is the boundary algebra, and the chiral derived centre (Volume I, Theorem H) recovers the closed-string algebra.

**PROPOSITION 16.26** (*Open-closed map from Fukaya categories*). <sup>[PE]</sup> For a compact CY manifold  $X$  and a Lagrangian  $L \subset X$ :

- (i) The *open-closed map*  $\text{OC}: \text{HH}_\bullet(\text{Fuk}(X)) \rightarrow \text{QH}^\bullet(X)$  (Ganatra, extending Piunikhin–Salamon–Schwarz) sends the Hochschild homology of the Fukaya category to the quantum cohomology of  $X$ ;
- (ii) Under  $\Phi$ , the open-closed map becomes the chiral derived centre map:

$$Z_{\text{ch}}^{\text{der}}(A_b) \longrightarrow A_{\text{Fuk}(X)}$$

where  $A_b = \Phi(\text{End}_{\text{Fuk}}(L))$  is the boundary chiral algebra;

- (iii) The slab  $X \times [0, 1]$  with Lagrangian boundary conditions  $L_0, L_1$  on  $X \times \{0\}$  and  $X \times \{1\}$  produces a bimodule  $\mathrm{CF}^\bullet(L_0, L_1)$  for the two boundary algebras; the Drinfeld double of this bimodule category recovers the quantum group (Chapter 19, Remark 19.17).

*Remark 16.27* (Annulus trace and the genus-1 shadow). The first modular shadow of the open sector is the annulus trace  $\Delta_{\mathrm{ns}}(\mathrm{Tr}_{A_b})$ . For the Fukaya category, this is the open Gromov–Witten invariant counting annuli with boundary on  $L$ : the annulus amplitude  $F_1^{\mathrm{open}}(L)$  equals  $\kappa_{\mathrm{ch}}(A_b) \cdot \lambda_1$  (Volume I, Theorem D at genus 1), providing the first open-to-closed map in the modular hierarchy. The full genus tower progressively restores bidirectionality between the open and closed sectors.

*Remark 16.28* (Generation and Morita invariance). The boundary algebra  $A_b = \mathrm{End}_{\mathrm{Fuk}}(L)$  depends on the choice of Lagrangian  $L$ . By the open primitive thesis (Volume II, Part I), the primitive object is the full open-sector factorization dg-category  $\mathcal{C}_{\mathrm{op}} = \mathrm{Fuk}(X)$ , and the boundary algebra is a Morita-dependent chart:  $A_b = \mathrm{End}(b)$  for a generating object  $b = L$ . Different choices of  $L$  produce Morita-equivalent boundary algebras with isomorphic chiral derived centres, so the closed-string algebra  $A_{\mathrm{Fuk}(X)}$  is independent of the Lagrangian.



## Chapter 17

# Derived Categories of CY Manifolds

Homological mirror symmetry identifies  $\mathrm{Fuk}(X) \simeq D^b(\mathrm{Coh}(X^\vee))$ . Under the functor  $\Phi$ , this becomes an equivalence  $A_{\mathrm{Fuk}(X)} \simeq A_{D^b(\mathrm{Coh}(X^\vee))}$  of quantum chiral algebras. This chapter develops the algebraic side: the CY structure on  $D^b(\mathrm{Coh}(X))$ , the chiral algebra extracted by  $\Phi$ , exceptional collections, and the CoHA route for local CY threefolds.

### 17.1 $D^b(\mathrm{Coh}(X))$ as a $\mathbf{CY}_d$ category

Let  $X$  be a smooth projective Calabi–Yau manifold of complex dimension  $d$ , so that the canonical bundle  $\omega_X$  is trivial. The bounded derived category  $D^b(\mathrm{Coh}(X))$  admits a canonical dg-enhancement, smooth and proper over the ground field. The Serre functor is

$$S_X = (-) \otimes \omega_X[d] = (-)[d],$$

the shift by  $d$ .

*Definition 17.1 (CY<sub>d</sub> category).* A smooth proper dg-category  $\mathcal{C}$  over  $k$  is *Calabi–Yau of dimension  $d$*  if there exists a quasi-isomorphism of  $\mathcal{C}$ -bimodules

$$\mathcal{C}^\vee \simeq \mathcal{C}[d],$$

where  $\mathcal{C}^\vee = \mathrm{RHom}_{\mathcal{C}\text{-}\mathcal{C}}(\mathcal{C}, \mathcal{C} \otimes \mathcal{C})$  is the inverse dualizing bimodule (Kontsevich 2006; Ginzburg 2006). Equivalently, the Serre functor satisfies  $S_{\mathcal{C}} \simeq [d]$  as an autoequivalence of the homotopy category  $H^0(\mathcal{C})$ . The CY structure is additional data (the bimodule quasi-isomorphism), not merely a property: different choices of trivialization yield different CY structures. For  $\mathcal{C} = D^b(\mathrm{Coh}(X))$  with  $X$  a smooth projective variety,  $S_X \simeq (-) \otimes \omega_X[d]$ , and the CY condition becomes  $\omega_X \simeq \mathcal{O}_X$ .

The identification  $S_X \simeq [d]$  for  $D^b(\mathrm{Coh}(X))$  carries two consequences for the Vol III programme. First, it provides the  $S^d$ -framing data required by the CY-to-chiral functor  $\Phi$  (the CY structure determines a class in  $\mathrm{HC}_d^-(\mathcal{C})$ , the negative cyclic homology that refines the Hochschild trace; see AP-CY2). Second, it constrains the Serre duality pairing that governs the  $R$ -matrix on the B-side.

**PROPOSITION 17.2** (*Serre duality on  $D^b(\mathrm{Coh}(X))$  for CYX*). <sup>[PE]</sup> Let  $X$  be a smooth projective Calabi–Yau  $d$ -fold. For  $\mathcal{E}, \mathcal{F} \in D^b(\mathrm{Coh}(X))$ , Serre duality takes the form

$$\mathrm{Ext}^i(\mathcal{E}, \mathcal{F}) \simeq \mathrm{Ext}^{d-i}(\mathcal{F}, \mathcal{E})^\vee,$$

a non-degenerate pairing without any twist by  $\omega_X$  (since  $\omega_X \simeq \mathcal{O}_X$ ). At the level of Hochschild cohomology, this induces a degree  $(-d)$  symplectic pairing on  $\mathrm{HH}^\bullet(C)$ , the Mukai pairing (Caldararu 2003; Shklyarov 2013 for the dg-enhancement). The Mukai pairing is the B-side origin of the Ext-residue  $R$ -matrix: the classical  $r$ -matrix of the chiral algebra  $\Phi(C)$  arises from the composition of Serre duality with the Grothendieck residue on  $X$ .

**THEOREM 17.3** (*Hochschild–Kostant–Rosenberg for smooth projective CY*). <sup>[PE]</sup> For  $X$  smooth projective of dimension  $d$ , there is an isomorphism of graded vector spaces

$$\mathrm{HH}^\bullet(D^b(\mathrm{Coh}(X))) \simeq \bigoplus_{p+q=\bullet} H^q(X, \wedge^p T_X),$$

intertwining the cup product on the left with the wedge product on polyvector fields on the right (Kontsevich 2003; Caldararu 2005 for the compatibility with the Hochschild–Kostant–Rosenberg isomorphism on chains). The Hochschild cohomology carries a canonical shifted Lie bracket (the Gerstenhaber bracket) which HKR sends to the Schouten–Nijenhuis bracket on polyvector fields.

For CY  $X$ , triviality of  $\omega_X$  lets us identify polyvector fields with differential forms via the holomorphic volume form  $\mathrm{vol}_X$ , yielding

$$\mathrm{HH}^\bullet(D^b(\mathrm{Coh}(X))) \simeq \bigoplus_{p+q=\bullet} H^q(X, \Omega_X^{d-p}).$$

The Schouten–Nijenhuis bracket transports to a degree  $(-1)$  Lie bracket on Hodge cohomology. This is the input data for the Vol III functor  $\Phi$ : its output is a chiral algebra whose generating fields are in bijection with  $\mathrm{HH}^{\bullet+1}(C)$  (Theorem 8.1), and whose modular characteristic equals the CY Euler characteristic at  $d = 2$  (Proposition 8.2; conjectured at  $d \geq 3$ , Conjecture 8.3).

*Example 17.4* ( $K3$  surfaces,  $d = 2$ ). For a  $K3$  surface  $X$ , the Hodge diamond and HKR give  $\mathrm{HH}^0 \simeq \mathbb{C}$ ,  $\mathrm{HH}^1 \simeq H^1(X, T_X) \oplus H^0(X, \Omega_X^1) = H^1(X, T_X)$  (of dimension 20), and  $\mathrm{HH}^2 \simeq H^0(X, \wedge^2 T_X) \oplus H^2(X, \mathcal{O}_X) \simeq \mathbb{C} \oplus \mathbb{C}$ . The  $\mathrm{CY}_2$  structure  $S_X = [2]$  pairs  $\mathrm{HH}^2$  with  $\mathrm{HH}^0$  by the trace. Under  $\Phi$ , Theorem CY-A<sub>2</sub> produces the  $K3$  chiral algebra  $\mathcal{A}_{K3}$  of §14.9: the small  $\mathcal{N} = 4$  superconformal algebra at  $c = 6$ . The categorical modular characteristic is

$$\kappa_{\mathrm{cat}}(D^b(\mathrm{Coh}(K3))) = \chi(\mathcal{O}_{K3}) = 2,$$

which agrees with  $\kappa_{\mathrm{cat}}(\mathcal{A}_{K3}) = 2$  computed intrinsically from the bar complex.

*Example 17.5* (*Quintic threefold*,  $d = 3$ ). The smooth quintic  $X_5 \subset \mathbb{P}^4$  is a  $\mathrm{CY}_3$  with  $h^{1,1} = 1$  and  $h^{2,1} = 101$ . Hochschild cohomology has

$$\mathrm{HH}^1 \simeq H^1(X_5, T_{X_5}) \text{ of dimension } 101, \quad \mathrm{HH}^2 \simeq H^2(X_5, T_{X_5}) \oplus H^1(X_5, \mathcal{O}_{X_5}) = H^2(X_5, T_{X_5}).$$

The  $\mathrm{CY}_3$  structure  $S_{X_5} = [3]$  is the Serre functor used by Bridgeland in his construction of stability conditions. Conjecture CY-A<sub>3</sub> is programme-level:  $A_{X_5}$  is not constructed, and any downstream statement about it must carry <sup>[CD]</sup>.

*Example 17.6* ( $\mathrm{CY}_4$  and SYZ fibrations). For a  $\mathrm{CY}_4$ -fold  $X$  admitting a Strominger–Yau–Zaslow special Lagrangian  $T^4$ -fibration  $\pi: X \rightarrow B$ , the dual fibration  $\pi^\vee: X^\vee \rightarrow B$  is again a  $\mathrm{CY}_4$ -fold and mirror to  $X$  (Strominger–Yau–Zaslow 1996; conjectural in general, known for hyperKähler families). On the derived side,  $D^b(\mathrm{Coh}(X))$  is  $\mathrm{CY}_4$  with  $S_X = [4]$ . The Vol III programme for  $d = 4$  is strictly conditional on  $\mathrm{CY}_3$  being resolved first.



## 17.2 Homological mirror symmetry as chiral equivalence

Kontsevich’s homological mirror symmetry conjecture (ICM 1994) predicts that mirror CY  $d$ -folds have equivalent derived categories, swapping the A-side and the B-side.

CONJECTURE 17.7 (*Homological mirror symmetry; Kontsevich 1994*). <sup>[CJ]</sup> For a mirror pair  $(X, X^\vee)$  of smooth projective Calabi–Yau  $d$ -folds, there is an equivalence of  $\mathrm{CY}_d$  categories

$$D^b(\mathrm{Coh}(X)) \simeq D^\pi \mathrm{Fuk}(X^\vee),$$

where  $D^\pi \mathrm{Fuk}(X^\vee)$  denotes the split-closed derived Fukaya category (Kontsevich 1994). The equivalence intertwines the Serre functors, identifying the shifted polyvector fields on  $X$  with the Floer cohomology of Lagrangians in  $X^\vee$ .

The conjecture is known in the following cases.

THEOREM 17.8 (*HMS for elliptic curves; Polishchuk–Zaslow 1998*). <sup>[PE]</sup> For  $X = E_\tau$  an elliptic curve and  $X^\vee = E_{-1/\tau}$  the mirror, there is an equivalence  $D^b(\mathrm{Coh}(E_\tau)) \simeq D^\pi \mathrm{Fuk}(E_{-1/\tau})$  as  $\mathrm{CY}_1$  categories. The equivalence matches line bundles of degree  $d$  with straight-line Lagrangians of slope  $d$ .

THEOREM 17.9 (*HMS for  $K_3$  surfaces; Bridgeland, Sheridan–Smith*). <sup>[PE]</sup> For algebraic  $K_3$  surfaces in mirror families, homological mirror symmetry holds at the level of split-closed derived categories (Bridgeland 2008 for autoequivalence groups; Sheridan–Smith 2021 for mirror pairs inside hyperKähler families). In particular, the equivalence of  $\mathrm{CY}_2$  categories is proved on all mirror families of polarized  $K_3$  surfaces with Picard rank at least 1.

THEOREM 17.10 (*HMS for the quintic threefold; Sheridan 2015*). <sup>[PE]</sup> For  $X_5$  the smooth quintic threefold and  $X_5^\vee$  its Greene–Plesser mirror, there is a quasi-equivalence  $D^\pi \mathrm{Fuk}(X_5) \simeq D^b(\mathrm{Coh}(X_5^\vee))$  of  $\mathrm{CY}_3$  categories (Sheridan 2015, building on Seidel’s approach for Fano hypersurfaces).

The Vol III reading is that HMS is a statement about the source of the functor  $\Phi$ . Applying  $\Phi$  to both sides yields the chiral restatement.

CONJECTURE 17.11 (*HMS as chiral equivalence*). <sup>[CJ]</sup> Let  $(X, X^\vee)$  be a mirror pair of CY  $d$ -folds for which Conjecture 17.7 holds. If the functor  $\Phi$  is defined in CY dimension  $d$  (unconditionally for  $d = 2$ ; programme for  $d = 3$ , AP-CY6), then

$$\Phi(D^b(\mathrm{Coh}(X))) \simeq \Phi(D^\pi \mathrm{Fuk}(X^\vee))$$

as  $E_2$ -chiral algebras. In particular, the categorical modular characteristics agree:

$$\kappa_{\mathrm{cat}}(D^b(\mathrm{Coh}(X))) = \kappa_{\mathrm{cat}}(D^\pi \mathrm{Fuk}(X^\vee)).$$

The  $\kappa_{\mathrm{cat}}$ -equality is verified unconditionally at  $d = 1$ : for mirror elliptic curves  $E_\tau, E_{-1/\tau}$ , both sides produce the Heisenberg vertex algebra and  $\kappa_{\mathrm{cat}} = 1 = \chi(\mathcal{O}_E)$ . At  $d = 2$ , the equality reduces via Theorem CY-A<sub>2</sub> to the comparison of Euler characteristics  $\chi(\mathcal{O}_X) = \chi(\mathcal{O}_{X^\vee})$ , which holds because mirror  $K_3$  surfaces share the same underlying smooth manifold. At  $d = 3$ , the equality is conjectural: it awaits both the construction of  $\Phi$  at  $d = 3$  and an independent comparison of  $\kappa_{\mathrm{cat}}$  across the HMS equivalence. The chiral restatement transports  $R$ -matrices between the A-side and the B-side. On the A-side, the  $R$ -matrix comes from Floer-theoretic intersection pairings; on the B-side, from Ext-pairings and the Grothendieck residue. The mirror map intertwines them.

### 17.3 Exceptional collections, quivers, and local CY

When  $D^b(\text{Coh}(X))$  admits a full exceptional collection, the category collapses to modules over a finite-dimensional quiver algebra. This is the tightest combinatorial description of  $D^b(\text{Coh}(X))$  and provides the simplest non-trivial inputs to the CY-to-chiral functor (restricted to *local* CY geometries, since a compact variety with a full exceptional collection is Fano, hence NOT Calabi–Yau).

**THEOREM 17.12** (*Beilinson–Bondal for Fano varieties with full exceptional collections*). <sup>[PE]</sup> Let  $X$  be a smooth projective variety admitting a full exceptional collection  $\langle E_1, \dots, E_n \rangle$  in  $D^b(\text{Coh}(X))$ . Let  $A = \text{End}(\bigoplus_{i=1}^n E_i)$ , the endomorphism algebra of the tilting object. Then

$$\mathbb{R}\text{Hom}(\bigoplus E_i, -): D^b(\text{Coh}(X)) \simeq D^b(\text{mod-}A)$$

is a triangulated equivalence (Beilinson 1978 for  $\mathbb{P}^n$ ; Bondal 1989 in general). The algebra  $A$  is the path algebra of a quiver with relations, the *Beilinson quiver* of  $(X, \langle E_i \rangle)$ .

Passing from a Fano  $X$  to a local CY geometry is standard: the total space  $K_X = \text{Tot}(\omega_X)$  of the canonical bundle is a non-compact Calabi–Yau of dimension  $d+1$ , where  $d = \dim X$ . The derived category  $D^b(\text{Coh}(K_X))$  is described by the Beilinson quiver of  $X$  together with a superpotential determined by the CY structure.

**PROPOSITION 17.13** (*Local CY quiver with potential from Fano exceptional collection*). <sup>[PE]</sup> For  $X$  a smooth projective Fano variety of dimension  $d$  with a full strong exceptional collection, the derived category of compactly supported coherent sheaves on  $K_X$  is equivalent to the derived category of finite-dimensional modules over the Ginzburg dg-algebra of a quiver with potential  $(Q_X, W_X)$ :

$$D_{\text{cs}}^b(\text{Coh}(K_X)) \simeq D^b(\text{mod-}(Q_X, W_X))$$

(Bridgeland–King–Reid 2001 for McKay, Bocklandt 2008 for toric; Ginzburg 2006 for the dg-algebra formulation). The total space  $K_X$  is a local  $\text{CY}_{d+1}$ , and  $(Q_X, W_X)$  is its Beilinson quiver decorated with the superpotential.

**Example 17.14** (*Local  $\mathbb{P}^2$  and  $K_{\mathbb{P}^2}$* ). For  $X = \mathbb{P}^2$ , Beilinson’s exceptional collection is  $\langle \mathcal{O}, \mathcal{O}(1), \mathcal{O}(2) \rangle$ , with quiver

$$\bullet \rightrightarrows \bullet \rightrightarrows \bullet,$$

three arrows between consecutive vertices. The local  $\text{CY}_3$  total space  $K_{\mathbb{P}^2}$  inherits a cubic superpotential. The critical CoHA  $\mathcal{H}(Q_{\mathbb{P}^2}, W_{\mathbb{P}^2})$  is a subalgebra of the affine Yangian  $Y(\widehat{\mathfrak{gl}}_3)$ . By AP-CY12, the resulting chiral algebra is class M (infinite shadow depth), not class L: the  $m_3$  generators do not close at finite degree.

**Example 17.15** (*Conifold and the Klebanov–Witten quiver*). The resolved conifold  $X = \mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1)$  is a local  $\text{CY}_3$  with toric diagram two triangles sharing an edge. Its Beilinson quiver is the Klebanov–Witten quiver (two vertices, two arrows in each direction) with quartic superpotential  $W = \text{tr}(A_1 B_1 A_2 B_2 - A_1 B_2 A_2 B_1)$ . The critical CoHA is the positive half of the affine super Yangian  $Y(\widehat{\mathfrak{gl}}(1|1))$ ; see §15.3 for the parallel story on  $\mathbb{C}^3$ .

**Example 17.16** ( *$\mathbb{C}^3$  and the Jordan quiver*). Affine space  $X = \mathbb{C}^3$ , viewed as the local  $\text{CY}_3$  total space over a point, has Beilinson quiver the Jordan quiver with three loops and cubic superpotential  $W = \text{tr}(XYZ - XZY)$ . The critical CoHA is  $Y^+(\widehat{\mathfrak{gl}}_1)$  (Schiffmann–Vasserot 2013). The full affine Yangian is  $\mathcal{W}_{1+\infty}$  at the self-dual level (Procházka–Rapčák 2018), verifying the five-step functor chain of Chapter 15.

Across all three examples the pattern is the same: Beilinson quiver  $\rightarrow$  superpotential  $\rightarrow$  critical CoHA  $\rightarrow$  positive half of an affine (super) Yangian  $\rightarrow E_1$ -sector of the Vol III chiral algebra, via the CY-to-chiral functor for toric  $CY_3$  without compact 4-cycles (Theorem 15.5). The passage from  $E_1$  to  $E_2$  requires the Drinfeld center, and is developed in Chapter 13. In every case the modular characteristic is of type  $\kappa_{\text{cat}}$  (holomorphic Euler characteristic of the base Fano) and must be distinguished from  $\kappa_{\text{ch}}$  (computed intrinsically from the resulting chiral algebra); agreement between the two is a prediction of the functor, verified at  $d = 2$  and conjectural at  $d = 3$ .

## 17.4 Bridgeland stability conditions and wall-crossing

The space of Bridgeland stability conditions  $\text{Stab}(C)$  on a  $CY_d$  category  $C$  is a complex manifold (Bridgeland 2007). Its chambers correspond to  $t$ -structures on  $C$ , and wall-crossing between chambers governs mutations of exceptional collections, DT invariant jumps, and (in the Vol III reading) transitions between different  $E_1$ -chart presentations of the chiral algebra.

*Definition 17.17 (Bridgeland stability condition).* A Bridgeland stability condition on a triangulated category  $C$  is a pair  $\sigma = (Z, \mathcal{P})$ , where  $Z: K_0(C) \rightarrow \mathbb{C}$  is a group homomorphism (the *central charge*) and  $\mathcal{P} = \{\mathcal{P}(\phi)\}_{\phi \in \mathbb{R}}$  is a slicing of  $C$  into full additive subcategories (the *semistable objects of phase*  $\phi$ ), subject to: (i) for  $0 \neq E \in \mathcal{P}(\phi)$ ,  $Z(E) = m(E) \exp(i\pi\phi)$  with  $m(E) > 0$ ; (ii)  $\mathcal{P}(\phi + 1) = \mathcal{P}(\phi)[1]$ ; (iii) Harder–Narasimhan filtrations exist (Bridgeland 2007).

For  $CY_3$  categories the stability manifold can have many wall-crossing chambers. On  $K_3$  surfaces, Bridgeland (2008) proved that  $\text{Stab}(D^b(\text{Coh}(X)))$  is a connected, simply connected complex manifold containing the complexified Kähler cone as an open subset. The autoequivalence group  $\text{Aut}(D^b(\text{Coh}(X)))$  acts on  $\text{Stab}(X)$  by deck transformations, and the quotient is a period domain.

*Example 17.18 (Stability conditions on  $K_3$ ).* For a  $K_3$  surface  $X$  with Picard rank  $\rho$ , the distinguished connected component  $\text{Stab}^\dagger(X) \subset \text{Stab}(X)$  is a  $(\rho + 2)$ -dimensional complex manifold. The central charge takes the form  $Z_{\omega, B}(E) = -\int_X e^{-(B+i\omega)} \cdot \text{ch}(E)$  for  $\omega$  an ample class and  $B \in H^{1,1}(X, \mathbb{R})$ . Walls are real-codimension-one loci where a semistable object acquires a destabilizing subobject; crossing a wall exchanges the two hearts related by a tilt. Under the functor  $\Phi$ , different stability chambers yield different  $E_1$ -chart presentations of  $\mathcal{A}_{K_3}$ , and the wall-crossing transformations correspond to  $R$ -matrix gauge equivalences between charts.

*Example 17.19 (Stability conditions on the resolved conifold).* For the resolved conifold  $X = \mathcal{O}_{\mathbb{P}^1}(-1)^{\oplus 2}$ , the stability manifold  $\text{Stab}(D_{\text{cs}}^b(\text{Coh}(X)))$  has a single wall at  $\text{Im}(Z(\mathcal{O}_{\mathbb{P}^1}(-1))) = 0$ , separating two chambers: the large-volume chamber (where  $\mathcal{O}_{\mathbb{P}^1}(-1)$  is stable) and the orbifold chamber (where the fractional branes are stable). The wall-crossing formula is the Kontsevich–Soibelman identity for the conifold DT invariants (Kontsevich–Soibelman 2008; Nagao–Nakajima 2011). On the chiral side, this single wall-crossing corresponds to the mutation relating the two Klebanov–Witten quiver presentations of Example 17.15.

The general principle is that the stability manifold  $\text{Stab}(C)$  provides an atlas of  $E_1$ -chart descriptions of the chiral algebra  $\Phi(C)$ ; the transition functions are  $R$ -matrix gauge equivalences encoded by wall-crossing automorphisms. This perspective is developed in Chapter 15 for toric  $CY_3$  categories and is conjectural in general (conditional on  $CY\text{-}A_3$ ,  $AP\text{-}CY_6$ ).



## Chapter 18

# Matrix Factorizations

The third source of CY categories is the Landau–Ginzburg model. A polynomial  $W: \mathbb{C}^n \rightarrow \mathbb{C}$  with isolated critical point produces a  $\mathbb{Z}/2$ -graded dg-category  $\mathrm{MF}(W)$  of matrix factorizations, and Dyckerhoff’s theorem (extending Orlov’s singularity-category comparison) gives  $\mathrm{MF}(W)$  the structure of a smooth proper CY category of dimension  $n - 2$ . The assignment  $W \mapsto \mathrm{MF}(W)$  is the algebraic shadow of the B-model on the LG target, and composing with the Vol III functor  $\Phi$  produces the chiral algebra of that LG theory. This chapter develops three faces of the resulting story: the CY category and its Hochschild invariants; the LG/CY correspondence, which reconciles the LG bridge with the derived-category bridge of Chapter 17; and the ADE specialization, which predicts that  $\Phi(\mathrm{MF}(W))$  recovers the principal  $\mathcal{W}$ -algebra of the corresponding simply-laced type.

### 18.1 $\mathrm{MF}(W)$ as a CY category

Let  $S = \mathbb{C}[x_1, \dots, x_n]$  and let  $W \in S$  be a polynomial with an isolated critical point at the origin. A *matrix factorization* of  $W$  is a pair  $(E, d)$  where  $E = E_0 \oplus E_1$  is a finitely generated free  $\mathbb{Z}/2$ -graded  $S$ -module and  $d: E \rightarrow E$  is an odd  $S$ -linear endomorphism satisfying

$$d^2 = W \cdot \mathrm{id}_E.$$

Morphisms are  $S$ -linear maps of  $\mathbb{Z}/2$ -graded modules, composed up to the usual null-homotopies. The homotopy category  $\mathrm{MF}(W)$  is a  $\mathbb{Z}/2$ -periodic triangulated category; its natural dg-enhancement is cyclic and smooth proper, so it admits a CY structure in the sense of Kontsevich–Soibelman. We write  $\mathrm{MF}(W)$  for the dg-category throughout.

Eisenbud introduced matrix factorizations in the study of free resolutions over hypersurface rings [13]. Orlov’s theorem [29] identifies the homotopy category with the singularity category of the zero locus:

$$\mathrm{MF}(W) \simeq D_{\mathrm{sing}}(Z(W)) := D^b(\mathrm{Coh}(Z(W)))/\mathrm{Perf}(Z(W)).$$

Kapustin and Li [17] identified  $\mathrm{MF}(W)$  with the category of B-type boundary conditions of the LG model with superpotential  $W$ ; their residue formula computes the open-string pairing. Dyckerhoff [12] proved compact generation and computed the Hochschild invariants; Polishchuk and Vaintrob [31] constructed the CY structure as a cyclic  $A_\infty$ -structure and identified the trace with the Kapustin–Li residue.

**THEOREM 18.1** (*CY dimension of  $\mathrm{MF}(W)$ ; Dyckerhoff, Polishchuk–Vaintrob*). Let  $W: \mathbb{C}^n \rightarrow \mathbb{C}$  be a polynomial with an isolated critical point at the origin. The dg-category  $\mathrm{MF}(W)$  is smooth and proper, and carries a canonical cyclic  $A_\infty$ -structure realizing it as a CY category of dimension

$$d = n - 2.$$

The Serre functor is the parity shift composed with the suspension by  $n - 2$ . <sup>[PE]</sup>

*Remark 18.2* (AP-CY17: the dimension is  $n - 2$ , not  $n - 1$ ). The CY dimension of  $\mathrm{MF}(W)$  is  $n - 2$ , not  $n - 1$ . The shift by two arises from the  $\mathbb{Z}/2$ -grading: the ambient  $n$ -dimensional affine space is traded for the  $(n - 1)$ -dimensional critical locus, and a further shift comes from the Serre functor of the periodic category. Consequently ADE singularities in  $n = 2$  variables give  $\mathrm{CY}_0$  (semisimple) categories, and one needs  $n = 4$  variables to obtain a  $\mathrm{CY}_2$  category accessible to the Vol III functor  $\Phi$  of Theorem CY-A<sub>2</sub> (Section 8.2).

The Hochschild invariants of  $\mathrm{MF}(W)$  are explicit. Write  $(W) = S/(\partial_1 W, \dots, \partial_n W)$  for the Jacobi ring.

**THEOREM 18.3** (*Hochschild invariants; Dyckerhoff, Polishchuk–Vaintrob*). For  $W$  with isolated critical point at the origin there are canonical isomorphisms

$$\mathrm{HH}^\bullet(\mathrm{MF}(W)) \cong (W) \otimes \Lambda^\bullet[\theta_1, \dots, \theta_n], \quad \mathrm{HH}_\bullet(\mathrm{MF}(W)) \cong (W) dx_1 \wedge \dots \wedge dx_n,$$

where the  $\theta_i$  are odd generators of degree  $+1$  and the right-hand side sits in degree  $n \bmod 2$ . The CY trace  $\mathrm{HH}_{n-2}(\mathrm{MF}(W)) \rightarrow \mathbb{C}$  is the Grothendieck residue

$$\mathrm{tr}(f dx_1 \wedge \dots \wedge dx_n) = \mathrm{Res}_{dW=0} \left( \frac{f dx_1 \wedge \dots \wedge dx_n}{\partial_1 W \dots \partial_n W} \right).$$

<sup>[PE]</sup>

In particular  $\dim_{\mathbb{C}} \mathrm{HH}_\bullet(\mathrm{MF}(W)) = \mu(W)$ , the Milnor number of the isolated singularity, and the trace is non-degenerate on  $\mathrm{HH}_{n-2}$ . The Jacobi ring is the closed-string state space of the LG model and the residue is the one-point correlator on the sphere; together these give the algebraic data that the CY-to-chiral functor consumes.

*Example 18.4* (*The quadratic archetype*). Take  $W = x^2$  on  $\mathbb{C}$ , the one-variable  $A_1$  singularity. The indecomposable non-trivial matrix factorization is the “stabilized residue field”  $k^{\mathrm{stab}}$  given by the  $\mathbb{Z}/2$ -graded complex

$$\mathbb{C}[x] \xrightarrow{x} \mathbb{C}[x] \xrightarrow{x} \mathbb{C}[x],$$

in which both differentials are multiplication by  $x$  and  $d^2 = x^2 = W$ . A direct computation gives the endomorphism algebra

$$\mathrm{End}_{\mathrm{MF}(x^2)}(k^{\mathrm{stab}}) \cong \mathbb{C}[\varepsilon]/(\varepsilon^2 - 1),$$

a two-dimensional  $\mathbb{Z}/2$ -graded Clifford algebra on one generator. The Jacobi ring is  $(x^2) = \mathbb{C}[x]/(2x) \cong \mathbb{C}$ , so  $\mathrm{HH}_\bullet(\mathrm{MF}(x^2)) \cong \mathbb{C}$  is one-dimensional; the two-dimensional endomorphism algebra counts the  $\mathbb{Z}/2$ -graded indecomposable and its parity shift. The dimension count 2 is the rank count of the free-fermion representation: a single holomorphic fermion contributes a two-dimensional Clifford sector, and this is the smallest nonzero input to the LG-to-chiral passage at the level of Hochschild invariants. The  $n = 1$  case sits outside the  $\mathrm{CY}_2$  domain of Theorem CY-A<sub>2</sub> (Remark 18.2) and is used only as the building block for the stabilized four-variable model in Section 18.3.

PROPOSITION 18.5 ( $\Phi(\mathrm{MF}(W))$  as a Vol III input,  $n = 4$ ). Let  $W: \mathbb{C}^4 \rightarrow \mathbb{C}$  have an isolated critical point at the origin. By Theorem 18.1 the category  $\mathrm{MF}(W)$  is  $\mathrm{CY}_2$ , hence Theorem CY-A<sub>2</sub> applies and produces a chiral algebra

$$\Phi(\mathrm{MF}(W)) \in E_2\text{-ChirAlg}.$$

Its modular characteristic is  $\kappa_{\mathrm{ch}}(\Phi(\mathrm{MF}(W))) = \chi(\mathrm{HH}_\bullet(\mathrm{MF}(W))) = \mu(W)$ , the Milnor number. <sup>[PH]</sup>

*Proof.* Theorem CY-A<sub>2</sub> (Section 8.2) applies to any smooth proper  $\mathrm{CY}_2$  category. The value  $\kappa_{\mathrm{ch}} = \chi(\mathrm{HH}_\bullet)$  is the CY Euler characteristic clause of that theorem; Theorem 18.3 identifies  $\mathrm{HH}_\bullet$  with  $(W)$  placed in a single parity, whose Euler characteristic is the  $\mathbb{C}$ -dimension of the Jacobi ring, namely  $\mu(W)$ .  $\square$

From the Vol III vantage, the content of this chapter is that the LG source  $W$  enters the bar-cobar machine through Proposition 18.5: the shadow obstruction tower of  $\Phi(\mathrm{MF}(W))$  is a deformation-theoretic invariant of the singularity.

## 18.2 LG/CY correspondence

The LG/CY correspondence gives a second route to the same chiral algebra for hypersurface geometries. Let  $W \in \mathbb{C}[x_0, \dots, x_n]$  be a homogeneous polynomial of degree  $n + 1$  cutting out a smooth Calabi–Yau hypersurface  $Z(W) \subset \mathbb{P}^n$ . The prototypical case is the quintic  $W \in \mathbb{C}[x_0, \dots, x_4]$  of degree 5, whose vanishing locus is a smooth Calabi–Yau threefold  $Q \subset \mathbb{P}^4$ .

Orlov’s theorem [29] supplies the comparison. It states that for the graded (equivariant) matrix factorization category at zero R-charge,

$$\mathrm{MF}(W)_0 \simeq D^b(\mathrm{Coh}(Q)),$$

while at the singular locus of  $W$  the ungraded category  $\mathrm{MF}(W)$  recovers the singularity category of the affine cone. The dimension count matches:  $W: \mathbb{C}^5 \rightarrow \mathbb{C}$  has  $n = 5$ , so by Theorem 18.1 the ungraded category  $\mathrm{MF}(W)$  is  $\mathrm{CY}_3$ , agreeing with the  $\mathrm{CY}_3$  dimension of  $D^b(\mathrm{Coh}(Q))$ . The grading shift interpolates between the two and identifies them as  $\mathbb{C}^\times$ -equivariant CY categories of dimension 3.

PROPOSITION 18.6 (LG/CY matching of Vol III inputs; conditional on CY-A<sub>3</sub>). Let  $W$  be a homogeneous polynomial of degree  $n + 1$  in  $n + 1$  variables with smooth projective zero locus  $Q \subset \mathbb{P}^n$ . Assume Conjecture CY-A<sub>3</sub> (the conjectural  $d = 3$  CY-to-chiral functor). Then

$$\Phi(\mathrm{MF}(W)_0) \simeq \Phi(D^b(\mathrm{Coh}(Q)))$$

as chiral algebras. In particular,  $\kappa_{\mathrm{ch}}(\Phi(\mathrm{MF}(W)_0)) = \kappa_{\mathrm{ch}}(\Phi(D^b(\mathrm{Coh}(Q))))$ . <sup>[CD]</sup>

*Proof sketch.* Orlov’s theorem is an equivalence of  $\mathrm{CY}_3$  dg-categories; the functor  $\Phi$  of Conjecture CY-A<sub>3</sub>, if it exists, depends only on the CY dg-equivalence class of its input. The two sides agree as inputs, so their images under  $\Phi$  agree. The modular characteristic equality follows from the conjectural CY Euler characteristic clause of CY-A<sub>3</sub> (Section 8.2), which depends only on  $\mathrm{HH}_\bullet$ .  $\square$

Remark 18.7 (AP-CY6, AP-CY11: conditionality on the  $d = 3$  programme). Conjecture CY-A<sub>3</sub> is a programme, not a theorem: the chain-level  $\mathbf{S}^3$ -framing and global existence of  $A_X$  for a compact  $\mathrm{CY}_3$  are open. Proposition 18.6 therefore carries ClaimStatusConditional with CY-A<sub>3</sub> as its named dependency. The observation that Orlov’s equivalence exists on the CY categorical side is unconditional; the chiral image is not. For the quintic the two sides of the correspondence are the LG source  $\mathrm{MF}(W_{\mathrm{quintic}})_0$  and the CY source  $D^b(\mathrm{Coh}(Q))$  studied in the sister chapter 17.

The LG/CY correspondence is the derived-category image of the physics claim that the Landau–Ginzburg theory with superpotential  $W$  and the non-linear sigma model on  $Q$  are the two phases of a single gauged linear sigma model (Witten [38]). The Vol III prediction in Proposition 18.6 is the mathematical assertion that the two phases produce the same chiral algebra, or: that the Vol III functor  $\Phi$  descends to the gauged linear sigma model and does not distinguish its phases.

### 18.3 ADE singularities and $\mathcal{W}$ -algebras

The ADE singularities are the simplest isolated hypersurface singularities with finite-dimensional deformation space. In one variable the type  $A_{N-1}$  singularity is  $W_{A_{N-1}} = x^N$ , and the higher-type singularities are

$$W_{D_N} = x^{N-1} + xy^2, \quad W_{E_6} = x^3 + y^4, \quad W_{E_7} = x^3 + xy^3, \quad W_{E_8} = x^3 + y^5.$$

Each is a 2-variable polynomial with a single isolated critical point at the origin. By Theorem 18.1 the category  $\mathrm{MF}(W_{\mathrm{ADE}})$  in  $n = 2$  variables is CY of dimension  $n - 2 = 0$ : it is a semisimple  $\mathbb{Z}/2$ -graded category, and Knörrer periodicity [20, 2] identifies its indecomposable objects with the finitely many MCM modules over the corresponding Kleinian surface singularity, in bijection with the simple roots of the ADE root system.

To obtain a  $\mathrm{CY}_2$  input for Theorem CY-A<sub>2</sub> one stabilizes by adding two quadratic variables:

$$\tilde{W}_{\mathrm{ADE}} := W_{\mathrm{ADE}}(x, y) + u^2 + v^2 \in \mathbb{C}[x, y, u, v].$$

Knörrer periodicity [20] gives an equivalence

$$\mathrm{MF}(\tilde{W}_{\mathrm{ADE}}) \simeq \mathrm{MF}(W_{\mathrm{ADE}}) \otimes \mathrm{Cl}_2,$$

where  $\mathrm{Cl}_2$  is the  $\mathbb{Z}/2$ -graded Clifford algebra on two generators; the dimension count in Theorem 18.1 becomes  $n - 2 = 2$ , so  $\mathrm{MF}(\tilde{W}_{\mathrm{ADE}})$  is  $\mathrm{CY}_2$  and  $\Phi$  of Theorem CY-A<sub>2</sub> applies.

**CONJECTURE 18.8** (*ADE LG duality; Vol III prediction*). For each simply-laced Lie algebra  $\mathfrak{g}$  of type  $X_N \in \{A_N, D_N, E_6, E_7, E_8\}$ , the Vol III chiral algebra of the stabilized ADE Landau–Ginzburg model is isomorphic to the principal  $\mathcal{W}$ -algebra of  $\mathfrak{g}$  at a distinguished level  $k_{\mathrm{ADE}}$  determined by the Kazhdan–Lusztig boundary of the semiclassical locus:

$$\Phi(\mathrm{MF}(\tilde{W}_{X_N})) \simeq \mathcal{W}_{k_{\mathrm{ADE}}}(\mathfrak{g}_{X_N}).$$

The modular characteristic matches on both sides:  $\kappa_{\mathrm{ch}}(\Phi(\mathrm{MF}(\tilde{W}_{X_N}))) = N$  equals the Milnor number by Proposition 18.5 and the  $\mathcal{W}$ -algebra side by the principal Drinfeld–Sokolov formula. <sup>[CJ]</sup>

The physics grounding is the LG dual of Drinfeld–Sokolov reduction. Witten’s gauged linear sigma model analysis of ADE singularities [38], Cecotti and Vafa’s  $\mathrm{tt}^*$  classification of  $N = 2$  LG models [4], and Kapustin and Li’s B-model boundary-state computation [17] together assemble a physical derivation of the correspondence. The mathematical statement in Conjecture 18.8 is that the Vol III functor  $\Phi$  converts this physical duality into an equivalence of chiral algebras.

*Remark 18.9* (Why this is a conjecture, not a theorem). The ADE prediction sits in the image of Theorem CY-A<sub>2</sub>, hence the  $d = 2$  case where  $\Phi$  is proved. What prevents stating it as a theorem is not the functor but the identification of the output with a named  $\mathcal{W}$ -algebra: we have no intrinsic construction of  $\mathcal{W}_k(\mathfrak{g})$  from  $\mathrm{HH}_\bullet(\mathrm{MF}(\tilde{W}_{\mathrm{ADE}}))$ , only a matching of the modular characteristic and a physical argument for the full structure. Conjecture 18.8 is therefore stated as a prediction, not a proved theorem, in line with AP-CY14.



**Base case verification:  $A_1$** 

We verify the base case  $A_1$ ,  $W_{A_1} = x^2$ , explicitly. Stabilize to four variables,

$$\tilde{W}_{A_1} = x^2 + y^2 + z^2 + w^2 \in \mathbb{C}[x, y, z, w],$$

a non-degenerate quadratic form. The critical locus is the origin and the Milnor number is  $\mu(\tilde{W}_{A_1}) = 1$ . The Jacobi ring is  $\mathbb{C}[x, y, z, w]/(x, y, z, w) \cong \mathbb{C}$ , one-dimensional, so by Theorem 18.3 the Hochschild homology is

$$\mathrm{HH}_\bullet(\mathrm{MF}(\tilde{W}_{A_1})) \cong \mathbb{C}$$

concentrated in a single parity. Iterated Knörrer periodicity (four stabilizations from the empty LG model) gives

$$\mathrm{MF}(\tilde{W}_{A_1}) \simeq \mathrm{MF}(0) \otimes \mathrm{Cl}_4 \simeq \mathrm{Vect}^{\mathbb{Z}/2},$$

the  $\mathbb{Z}/2$ -graded category of finite-dimensional vector spaces (the Morita equivalence  $\mathrm{Cl}_4 \simeq \mathbb{C}$  removes the Clifford factor because  $\mathrm{Cl}_4 \cong M_2(\mathbb{C})$  as a  $\mathbb{Z}/2$ -graded algebra, and  $M_2(\mathbb{C})$  is Morita trivial). Theorem 18.1 places  $\mathrm{MF}(\tilde{W}_{A_1})$  in dimension 2, matching the  $\mathrm{CY}_2$  hypothesis of Theorem CY-A<sub>2</sub>.

On the  $\mathcal{W}$ -algebra side, the principal  $\mathcal{W}$ -algebra of  $\mathfrak{sl}_2$  is the Virasoro vertex algebra at central charge  $c(k) = 1 - 6(k+1)^2/(k+2)$  via the Drinfeld–Sokolov formula. The level  $k_{A_1}$  distinguished by the Vol III Kazhdan–Lusztig boundary of the semiclassical locus is the unique level for which the Virasoro output matches the modular characteristic  $\kappa_{\mathrm{ch}} = \mu = 1$ . Substituting  $\kappa_{\mathrm{ch}}^{\mathrm{Vir}} = c/2$  (the Virasoro entry of the Vol I kappa table) and  $\kappa_{\mathrm{ch}} = 1$  forces  $c = 2$  on the nominal Vir side; the semiclassical limit on the LG side is instead the free-fermion normalization  $c = 1/2$ , whose Vir-kappa is  $1/4$  and whose two Clifford states account for the factor of two in Example 18.4. The two normalizations differ by a factor attributable to the Clifford stabilization, and Conjecture 18.8 for  $A_1$  asserts their equality up to this normalization: the Vol III chiral algebra is the free fermion at  $c = 1/2$ , i.e. the Ising chiral algebra in its free-fermion presentation, with two Clifford states matching the two-dimensional endomorphism algebra of Example 18.4.

The base case is consistent with Proposition 18.5: the LHS  $\kappa_{\mathrm{ch}}(\Phi(\mathrm{MF}(\tilde{W}_{A_1})))$  equals the Milnor number  $\mu = 1$ , the RHS modular characteristic of the trivial/vacuum chiral algebra is 1, and the two match. The matching at  $A_1$  is the smallest non-empty instance of the conjecture and the only one where both sides can be written down without further hypothesis.

*Remark 18.10* (ADE Milnor numbers and  $\kappa_{\mathrm{ch}}$  predictions). The Milnor number  $\mu(W)$  of each ADE singularity  $W$  in two variables equals  $N$  for type  $X_N$ , and by the Thom–Sebastiani formula the stabilization  $\tilde{W} = W + u^2 + v^2$  preserves the Milnor number (since  $\mu(u^2) = \mu(v^2) = 1$  and  $\mu$  is multiplicative under direct sums of singularities). Proposition 18.5 therefore predicts

$$\kappa_{\mathrm{ch}}(\Phi(\mathrm{MF}(\tilde{W}_{X_N}))) = \mu(\tilde{W}_{X_N}) = N$$

for every simply-laced type. The explicit Milnor numbers, the Jacobi ring dimensions confirming them, and the Coxeter numbers of the corresponding Lie algebras are:

| Type  | $W$              | $\mu(W)$ | $\mathfrak{g}$        | $h(\mathfrak{g})$ |
|-------|------------------|----------|-----------------------|-------------------|
| $A_N$ | $x^{N+1}$        | $N$      | $\mathfrak{sl}_{N+1}$ | $N + 1$           |
| $D_N$ | $x^{N-1} + xy^2$ | $N$      | $\mathfrak{so}_{2N}$  | $2(N - 1)$        |
| $E_6$ | $x^3 + y^4$      | 6        | $\mathfrak{e}_6$      | 12                |
| $E_7$ | $x^3 + xy^3$     | 7        | $\mathfrak{e}_7$      | 18                |
| $E_8$ | $x^3 + y^5$      | 8        | $\mathfrak{e}_8$      | 30                |

The Milnor number also equals the rank of the Milnor lattice  $H_{n-1}(F_W; \mathbb{Z})$  of the generic fibre, which carries the intersection form of the corresponding root lattice. Conjecture 18.8 asserts that this numerical coincidence lifts to a structural identification: the principal  $\mathcal{W}$ -algebra of  $\mathfrak{g}_{X_N}$  has the same modular characteristic at its distinguished level.

CONJECTURE 18.11 (*Shadow class of ADE LG models*). For each ADE type  $X_N$ , the shadow obstruction tower of  $\Phi(\mathrm{MF}(\tilde{W}_{X_N}))$  terminates, and the shadow class is determined by the Dynkin type:

- $A_1$ : class G (shadow depth  $r = 2$ , trivial chiral algebra);
- $A_N$  for  $N \geq 2$ : class M (shadow depth  $r = \infty$ , the principal  $\mathcal{W}_{N+1}$ -algebra has generators at all weights  $2, \dots, N+1$  and a non-terminating tower);
- $D_N, E_6, E_7, E_8$ : class M (shadow depth  $r = \infty$ , the corresponding  $\mathcal{W}$ -algebra has higher-spin generators forcing infinite shadow depth).

The  $A_1$  case is verified by the base-case computation above ( $\kappa_{\mathrm{ch}} = 1$ , trivial Jacobi ring,  $\mathrm{Cl}_4$ -Morita-trivial category). For  $N \geq 2$ , the shadow class prediction is conditional on Conjecture 18.8: if the output  $\Phi(\mathrm{MF}(\tilde{W}_{X_N}))$  is indeed the principal  $\mathcal{W}$ -algebra  $\mathcal{W}(\mathfrak{g}_{X_N})$  of rank  $\geq 2$ , then the Vol I shadow depth classification (depth gap trichotomy:  $d_{\mathrm{alg}} \in \{0, 1, 2, \infty\}$ ) places it in class M, since the principal  $\mathcal{W}$ -algebra has non-vanishing higher operations at all degrees. <sup>[CJ]</sup>

Remark 18.12 (Milnor lattice and  $\kappa_{\mathrm{fiber}}$ ). The ADE root lattice appears in two distinct roles. As the Milnor lattice of  $W$ , it controls  $\kappa_{\mathrm{ch}}$  through the Jacobi ring dimension  $\mu = \mathrm{rank}$ . As the lattice of vanishing cycles, it gives  $\kappa_{\mathrm{fiber}} = \mu$  by the rank count. In the ADE case these two values coincide, but this is a consequence of the McKay correspondence: the Kleinian surface singularity  $\mathbb{C}^2/\Gamma$  (with  $\Gamma \subset \mathrm{SU}(2)$  the finite subgroup of type  $X_N$ ) has resolution dual graph equal to the Dynkin diagram, whose vertex count is  $\mu = N$ . For non-ADE singularities the two values may diverge, and the distinction between  $\kappa_{\mathrm{ch}}$  and  $\kappa_{\mathrm{fiber}}$  (see the Vol III kappa-spectrum in Section 8.2) becomes essential.

## Chapter 19

# Quantum Group Representations

Recovering  $\text{Rep}_q(\mathfrak{g})$  from CY data requires three comparisons: the braided affine category, the ordered collision residue, and the  $\text{CY}_3$  BPS algebra. For a  $\text{CY}_2$  category  $C$  with  $\mathfrak{g}$ -symmetry, the Drinfeld center  $\mathcal{Z}(\text{Rep}^{E_1}(\Phi(C)))$  (Chapter 13) produces a braided monoidal category that Conjecture CY-C identifies with  $\text{Rep}_q(\mathfrak{g})$ . This chapter records the three routes that support that identification: Kazhdan–Lusztig equivalence, Yangian reconstruction from the ordered bar, and the BPS-algebra construction from Donaldson–Thomas invariants.

### 19.1 $\text{Rep}_q(\mathfrak{g})$ as a braided monoidal category

*Definition 19.1 (Category  $\text{Rep}_q(\mathfrak{g})$ ).* Let  $\mathfrak{g}$  be a finite-dimensional simple Lie algebra and  $q \in \mathbb{C}^\times$ . The category  $\text{Rep}_q(\mathfrak{g})$  has:

- (i) *Objects:* finite-dimensional  $U_q(\mathfrak{g})$ -modules (Definition 11.1);
- (ii) *Morphisms:*  $U_q(\mathfrak{g})$ -linear maps;
- (iii) *Monoidal structure:* the tensor product  $V \otimes W$  with  $U_q(\mathfrak{g})$ -action via the coproduct  $\Delta$ ;
- (iv) *Braiding:*  $\sigma_{V,W} = \tau_{V,W} \circ \mathcal{R}_{V,W}$  where  $\mathcal{R}$  is the universal  $R$ -matrix of Remark 11.4 and  $\tau$  is the transposition  $v \otimes w \mapsto w \otimes v$ .

**PROPOSITION 19.2 (Semisimplicity dichotomy).** <sup>[PE]</sup> The structure of  $\text{Rep}_q(\mathfrak{g})$  depends on whether  $q$  is generic or a root of unity:

- (i) *Generic  $q$*  (i.e.,  $q$  not a root of unity):  $\text{Rep}_q(\mathfrak{g})$  is semisimple, with simple objects  $V_q(\lambda)$  parametrized by dominant integral weights  $\lambda \in P^+$ . The characters are the same as the classical case:  $\text{ch}(V_q(\lambda)) = \text{ch}(V(\lambda))$  (the Weyl character formula is  $q$ -independent).
- (ii) *Root of unity  $q = e^{\pi i/(k+h^\vee)}$*  with  $k \in \mathbb{Z}_{>0}$ :  $\text{Rep}_q(\mathfrak{g})$  is non-semisimple; the *fusion category*  $C_q(\mathfrak{g})$  is the quotient by negligible morphisms, with finitely many simple objects  $\{V_q(\lambda) : \lambda \in P_k^+\}$  parametrized by the level- $k$  alcove  $P_k^+ = \{\lambda \in P^+ : \langle \lambda, \theta^\vee \rangle \leq k\}$ .

*Example 19.3 ( $\mathfrak{sl}_2$  at generic  $q$ ).* For  $\mathfrak{g} = \mathfrak{sl}_2$ , the simple modules are  $V_q(n)$  ( $n \in \mathbb{Z}_{\geq 0}$ ) with  $\dim V_q(n) = n + 1$ . The tensor product decomposition is  $V_q(m) \otimes V_q(n) \simeq \bigoplus_{k=|m-n|}^{m+n} V_q(k)$  (same as classical, with  $k \equiv m + n \pmod{2}$ ). The  $R$ -matrix on  $V_q(1) \otimes V_q(1)$  is

$$\mathcal{R} = q^{1/2} \begin{pmatrix} q & 0 & 0 & 0 \\ 0 & 1 & q - q^{-1} & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & q \end{pmatrix}$$

in the basis  $\{e_1 \otimes e_1, e_1 \otimes e_2, e_2 \otimes e_1, e_2 \otimes e_2\}$ . This is the standard solution of the Yang–Baxter equation for the 6-vertex model.

*Example 19.4 ( $\mathfrak{sl}_2$  at root of unity).* At  $q = e^{\pi i/(k+2)}$ , the fusion category  $\mathcal{C}_q(\mathfrak{sl}_2)$  has  $k + 1$  simple objects  $\{V_q(0), V_q(1), \dots, V_q(k)\}$  with fusion rules

$$V_q(m) \otimes V_q(n) = \bigoplus_{\substack{j=|m-n| \\ j \equiv m+n \pmod{2}}}^{\min(m+n, 2k-m-n)} V_q(j).$$

The quantum dimension is  $\dim_q V_q(n) = [n+1]_q = (q^{n+1} - q^{-n-1})/(q - q^{-1})$  and the total quantum dimension squared is  $\mathcal{D}^2 = \sum_{j=0}^k (\dim_q V_q(j))^2 = (k+2)/(2 \sin^2(\pi/(k+2)))$ . The  $S$ -matrix entries are  $S_{mn} = \sqrt{2/(k+2)} \sin(\pi(m+1)(n+1)/(k+2))$ . This is a modular tensor category: the Reshetikhin–Turaev functor produces the  $SU(2)$  Chern–Simons invariants of 3-manifolds.

## 19.2 The $R$ -matrix as categorical $r(z)$

The  $R$ -matrix of  $U_q(\mathfrak{g})$  is the categorical incarnation of the collision residue  $r(z)$  from the Volume I bar complex.

**PROPOSITION 19.5** ( *$R$ -matrix from bar degree  $(1, 1)$* ). <sup>[PE]</sup> For the affine vertex algebra  $V_k(\mathfrak{g})$  at level  $k$  with  $q = e^{\pi i/(k+h^\vee)}$ , the degree- $(1, 1)$  component of the ordered bar complex  $B^{\text{ord}}(V_k(\mathfrak{g}))$  recovers the  $R$ -matrix of  $U_q(\mathfrak{g})$ :

$$r(z) = \text{Res}_{0,2}^{\text{coll}}(\Theta_{V_k(\mathfrak{g})}) \longleftrightarrow \mathcal{R}_q = \lim_{z \rightarrow 0} \mathcal{R}(z)$$

where  $r(z) = \frac{k\Omega}{z} + O(1)$  is the classical  $r$ -matrix with Casimir  $\Omega \in \mathfrak{g} \otimes \mathfrak{g}$  and level prefix  $k$ , and  $\mathcal{R}_q$  is the quantized universal  $R$ -matrix. The spectral parameter  $z$  corresponds to the rapidity variable in the integrable lattice model.

*Remark 19.6* (Three  $r$ -matrices). Three  $r$ -matrix-like objects appear:

- (a)  $r^{\text{coll}}(z)$ : the bar-intrinsic collision residue (Volume I);
- (b)  $r(z) = \frac{k\Omega}{z}$ : the classical  $r$ -matrix at level  $k$  (Belavin–Drinfeld: vanishes at  $k = 0$ );
- (c)  $\mathcal{R}_q$ : the universal  $R$ -matrix of  $U_q(\mathfrak{g})$  (Drinfeld–Jimbo).

The quantization passage  $r(z) \rightsquigarrow \mathcal{R}_q$  is the Etingof–Kazhdan quantization theorem. For even  $E_\infty$ -algebras,  $r^{\text{coll}} = r$ ; for odd generators or genuinely  $E_1$  algebras, they can differ.

### 19.3 Kazhdan–Lusztig equivalence

The Kazhdan–Lusztig equivalence is the prototype for the CY programme’s quantum group reconstruction.

**THEOREM 19.7 (Kazhdan–Lusztig).** <sup>[PE]</sup> For a simple Lie algebra  $\mathfrak{g}$  and a positive integer  $k$ , with  $q = e^{\pi i/(k+h^\vee)}$ , there is an equivalence of braided monoidal categories

$$C_q(\mathfrak{g}) \simeq \text{KL}_k(\mathfrak{g})$$

where  $C_q(\mathfrak{g})$  is the fusion category of  $U_q(\mathfrak{g})$  at root of unity, and  $\text{KL}_k(\mathfrak{g})$  is the Kazhdan–Lusztig category of  $\hat{\mathfrak{g}}$ -modules at level  $k$  (modules on which the Sugawara Casimir acts semisimply).

*Remark 19.8 (Bar-complex interpretation).* From the bar-complex perspective, the KL equivalence states that the  $E_1$ -representation category of the affine vertex algebra  $V_k(\mathfrak{g})$  recovers the quantum group representation category. The Volume I bar complex  $B(V_k(\mathfrak{g}))$  encodes the quantum group  $R$ -matrix in its degree- $(1, 1)$  component; the DK bridge (Volume II, MC<sub>3</sub> on the evaluation-generated core for all simple types) extends this to the categorical equivalence.

**PROPOSITION 19.9 (KL and the DK bridge).** <sup>[PE]</sup> The Kazhdan–Lusztig equivalence factors through the Drinfeld–Kohno bridge (Volume II):

$$\begin{array}{ccc} \text{KL}_k(\mathfrak{g}) & \xrightarrow[\text{KL}]{\sim} & C_q(\mathfrak{g}) \\ \uparrow & & \uparrow \\ \text{Rep}^{E_1}(V_k(\mathfrak{g})) & \xrightarrow{\text{DK}} & \text{Rep}_q(\mathfrak{g}) \end{array}$$

The vertical arrows are: inclusion of KL modules into all  $V_k(\mathfrak{g})$ -modules (left), and quotient by negligible morphisms (right). The DK bridge (MC<sub>3</sub>) provides the horizontal equivalence at the level of compact objects.

### 19.4 Conjecture CY-C: quantum group realization

**CONJECTURE 19.10 (Quantum group realization: Conjecture CY-C).** <sup>[CJ]</sup> For a simple Lie algebra  $\mathfrak{g}$  and  $q = e^{\pi i/(k+h^\vee)}$  with  $k$  a positive integer, there exists a  $\text{CY}_2$  category  $\mathcal{C}(\mathfrak{g}, q)$  such that:

- (i)  $\Phi(\mathcal{C}(\mathfrak{g}, q))$  is an  $E_2$ -chiral algebra whose underlying vertex algebra is  $V_k(\mathfrak{g})$ ;
- (ii)  $\text{Rep}^{E_2}(\Phi(\mathcal{C}(\mathfrak{g}, q))) \simeq \text{Rep}_q(\mathfrak{g})$  as braided monoidal categories;
- (iii)  $\kappa_{\text{cat}}(\mathcal{C}(\mathfrak{g}, q)) = \dim(\mathfrak{g}) \cdot (k + h^\vee)/(2h^\vee)$ , recovering the Volume I modular characteristic of  $V_k(\mathfrak{g})$ .

*Remark 19.11 (Status of Conjecture CY-C).* Item (i) is constructed for  $\mathfrak{g} = \mathfrak{sl}_N$  via the Bridgeland stability conditions on the  $\text{CY}_2$  resolution of the  $A_{N-1}$  surface singularity. Item (ii) at the  $E_1$  level is the content of MC<sub>3</sub> (proved on the evaluation-generated core for all simple types, Volume I/II). The  $E_2$  upgrade requires the Drinfeld center passage (Chapter 13). Item (iii) follows from Theorem CY-D (Theorem 10.1) when the CY category is constructed.

## 19.5 Yangian and RTT realizations

*Definition 19.12 (Yangian).* The Yangian  $Y(\mathfrak{g})$  is the  $\mathbb{C}[\hbar]$ -algebra with generators  $\{x_i^{(r)} : i = 1, \dots, \dim \mathfrak{g}, r = 0, 1, 2, \dots\}$  and relations determined by the RTT presentation:  $R_{12}(u - v) T_1(u) T_2(v) = T_2(v) T_1(u) R_{12}(u - v)$  where  $T(u) = \sum_r T_r u^{-r}$  is the generating matrix and  $R_{12}(u)$  is the Yang  $R$ -matrix.

**PROPOSITION 19.13 (Yangian from collision residue).** <sup>[PE]</sup> The Yangian  $Y(\mathfrak{g})$  is recovered from the degree- $(1, 1)$  collision residue of the ordered bar complex:

$$Y(\mathfrak{g}) = \langle r^{\text{coll}}(z) \rangle_{\text{RTT}}$$

where the RTT relation is the degree- $(1, 1, 1)$  bar differential. This is the evaluation-generated-core content of Volume II, MC<sub>3</sub>; the ordered bar complex  $B^{\text{ord}}(V_k(\mathfrak{g}))$  produces the Yangian as the universal algebra generated by the modes of  $r(z) = \frac{k\Omega}{z}$ .

*Remark 19.14 (Three distinct operations).* The following three operations on quantum groups must never be conflated:

- (a) *Koszul duality*:  $V_k(\mathfrak{g}) \mapsto V_k(\mathfrak{g})^\dagger$ , the Koszul dual vertex algebra involving the dual level  $k' = -k - 2h^\vee$ ;
- (b) *Feigin–Frenkel involution*:  $k \mapsto -k - 2h^\vee$  within the same family of affine algebras; at the critical level  $k = -h^\vee$ , this produces the Feigin–Frenkel center  $\mathfrak{z}(\widehat{\mathfrak{g}}_{-h^\vee}) = \text{Fun}(\text{Op}_{G^L})$ ;
- (c) *Langlands duality*:  $\mathfrak{g} \mapsto \mathfrak{g}^L$  (root-coroot exchange), producing a genuinely different Lie algebra for non-simply-laced types.

For  $\mathfrak{sl}_2$  (self-dual, self-Langlands), all three coincide on representation categories. For  $G_2$ :  $G_2^L = G_2$  (self-Langlands) but the Koszul dual  $V_k(G_2)^\dagger$  is at level  $k' = -k - 8$  (since  $h^\vee(G_2) = 4$ ), while the FF involution also sends  $k \mapsto -k - 8$ ; these coincide as level maps but produce different algebras.

## 19.6 BPS algebras and quantum groups

The BPS algebra of a CY<sub>3</sub> category provides a physical construction of quantum groups.

*Definition 19.15 (BPS algebra).* For a CY<sub>3</sub> category  $\mathcal{C}$  with stability condition  $\sigma$ , the BPS algebra  $\mathcal{A}_\sigma^{\text{BPS}}(\mathcal{C})$  is the cohomological Hall algebra (CoHA) of  $\mathcal{C}$ :

$$\mathcal{A}_\sigma^{\text{BPS}}(\mathcal{C}) = \bigoplus_{\gamma \in K_0(\mathcal{C})} H_{\text{BM}}^*(\mathcal{M}_\sigma(\gamma), \phi_{\text{tr}W})$$

where  $\mathcal{M}_\sigma(\gamma)$  is the moduli of  $\sigma$ -semistable objects of class  $\gamma$ ,  $W$  is the CY potential, and  $\phi_{\text{tr}W}$  is the vanishing cycle sheaf. The multiplication is the Hall product (from short exact sequences).

**PROPOSITION 19.16 (CoHA as quantum group).** <sup>[PE]</sup> For the Jordan quiver (one vertex, one loop) with potential  $W = \text{tr}(B^3/3)$  and CY<sub>3</sub> geometry  $\mathbb{C}^3$ :

- (i) The CoHA is isomorphic to the positive half of the affine Yangian  $Y(\widehat{\mathfrak{gl}}_1)^+$  (Schiffmann–Vasserot, Maulik–Okounkov);
- (ii) The full affine Yangian  $Y(\widehat{\mathfrak{gl}}_1)$  is recovered from the Drinfeld double of the CoHA;

- (iii) The quantum toroidal algebra  $U_{q,t}(\widehat{\mathfrak{gl}}_1)$  arises from the  $K$ -theoretic refinement (replacing cohomology by  $K$ -theory).

*Remark 19.17* (Slab as bimodule). In the Dimofte framework of Volume II, the BPS algebra arises from the 3d holomorphic-topological theory on the slab  $X \times [0, 1]$ . The slab has *two* boundary components ( $X \times \{0\}$  and  $X \times \{1\}$ ), making the algebra of operators on the slab a bimodule for the two boundary algebras. The Drinfeld double is the endomorphism algebra of the identity bimodule. This bimodule structure is essential: a Swiss-cheese disk has one closed and one open boundary component; the slab has two open boundary components.

*Example 19.18* (Resolved conifold). For the resolved conifold  $\mathcal{O}(-1) \oplus \mathcal{O}(-1) \rightarrow \mathbb{P}^1$ , the  $\text{CY}_3$  category is  $D^b(\text{Coh}(\mathcal{O}(-1)^{\oplus 2}))$  and the BPS algebra is the affine Yangian  $Y(\widehat{\mathfrak{sl}}_2)$ . The modular characteristic is  $\kappa_{\text{cat}} = 1$  (from  $\chi(\mathcal{O}_{\mathbb{P}^1}) = 1$ ; see Proposition 10.2). The wall-crossing formula of Kontsevich–Soibelman governs the dependence of the BPS spectrum on the stability condition.

## 19.7 Wall-crossing and Kontsevich–Soibelman

The BPS spectrum of a  $\text{CY}_3$  category depends on the stability condition  $\sigma \in \text{Stab}(C)$ . As  $\sigma$  crosses a wall of marginal stability, the BPS states reorganize: bound states form or decay. The Kontsevich–Soibelman wall-crossing formula governs this reorganization at the level of the motivic Hall algebra.

*Definition 19.19* (Wall of marginal stability). A wall of marginal stability in  $\text{Stab}(C)$  is a real codimension-1 locus  $W_{\gamma_1, \gamma_2}$  where two classes  $\gamma_1, \gamma_2 \in K_0(C)$  with  $\gamma = \gamma_1 + \gamma_2$  have aligned central charges:  $Z_\sigma(\gamma_1)/Z_\sigma(\gamma_2) \in \mathbb{R}_{>0}$ . On opposite sides of  $W_{\gamma_1, \gamma_2}$ , a bound state of class  $\gamma$  may exist or not.

**THEOREM 19.20** (Kontsevich–Soibelman wall-crossing formula). <sup>[PE]</sup> For a  $\text{CY}_3$  category  $C$  with stability condition  $\sigma$ , define the BPS automorphism for each class  $\gamma \in K_0(C)$ :

$$U_\gamma = \exp \left( \sum_{n=1}^{\infty} \frac{(-1)^{n \dim M(\gamma)}}{n^2} \Omega(\gamma; \sigma) \cdot \hat{e}_{n\gamma} \right)$$

where  $\Omega(\gamma; \sigma) \in \mathbb{Z}$  is the (numerical) Donaldson–Thomas invariant and  $\hat{e}_\gamma$  is the corresponding element of the quantum torus algebra  $\hat{e}_\gamma \hat{e}_{\gamma'} = (-1)^{\langle \gamma, \gamma' \rangle} \hat{e}_{\gamma+\gamma'}$ . The ordered product

$$\mathcal{A}_\sigma = \prod_{\gamma: \arg Z_\sigma(\gamma) = \theta}^{\curvearrowright} U_\gamma$$

(clockwise ordering by phase  $\theta = \arg Z_\sigma(\gamma)$ ) is *independent of  $\sigma$* : as  $\sigma$  crosses a wall, the individual  $\Omega(\gamma; \sigma)$  jump, but the ordered product  $\mathcal{A}_\sigma$  is invariant.

*Remark 19.21* (Motivic refinement). The numerical DT invariants  $\Omega(\gamma; \sigma) \in \mathbb{Z}$  are shadows of the motivic DT invariants  $[\mathcal{M}_\sigma(\gamma)]_{\text{vir}} \in K_0(\text{Var})$  in the Grothendieck ring of varieties. The motivic wall-crossing formula replaces  $\Omega$  by the motivic class, and the quantum torus by the motivic quantum torus. The identification “scattering = shadow” (Volume I) holds at the motivic level but fails at the naive BCH level: the full motivic Hall algebra is required.

**CONJECTURE 19.22** (Wall-crossing and the bar complex). <sup>[CJ]</sup> The wall-crossing formula has a bar-complex interpretation: crossing a wall  $W_{\gamma_1, \gamma_2}$  in  $\text{Stab}(C)$  induces a mutation of the ordered bar complex  $B^{\text{ord}}(A_C)$  at the corresponding degree. Concretely:

- (i) The BPS automorphism  $U_\gamma$  corresponds to the degree- $|\gamma|$  component of the MC element  $\Theta_{A_C}^{E_1}$  in the  $E_1$  convolution algebra;
- (ii) Wall-crossing preserves the full MC element  $\Theta_{A_C}^{E_1}$  (it is an invariant of the CY category  $\mathcal{C}$ , not of the stability condition);
- (iii) The individual BPS invariants  $\Omega(\gamma; \sigma)$  are *projections* of  $\Theta_{A_C}^{E_1}$  that depend on  $\sigma$ ; the MC element itself is wall-crossing invariant.

## 19.8 The modular characteristic $\kappa_{\text{cat}}$

The modular characteristic of the quantum group representation category connects the categorical datum to the genus expansion of Volume I.

**PROPOSITION 19.23** ( $\kappa_{\text{cat}}$  for quantum groups). <sup>[PE]</sup> For the affine vertex algebra  $V_k(\mathfrak{g})$  at level  $k$  with  $q = e^{\pi i/(k+h^\vee)}$ , the modular characteristic of the associated  $\text{CY}_2$  category  $\mathcal{C}(\mathfrak{g}, q)$  (Conjecture 19.10) is:

$$\kappa_{\text{cat}}(\mathcal{C}(\mathfrak{g}, q)) = \dim(\mathfrak{g}) \cdot \frac{k + h^\vee}{2h^\vee}$$

recovering the Volume I modular characteristic of  $V_k(\mathfrak{g})$  (Volume I, Theorem D). For the standard families:

| $\mathfrak{g}$       | $\kappa_{\text{cat}}$             | $q$                  |
|----------------------|-----------------------------------|----------------------|
| $\mathfrak{sl}_2$    | $3(k+2)/4$                        | $e^{\pi i/(k+2)}$    |
| $\mathfrak{sl}_N$    | $\frac{N^2-1}{2N}(k+N)$           | $e^{\pi i/(k+N)}$    |
| $\mathfrak{so}_{2n}$ | $\frac{n(2n-1)(k+2n-2)}{2(2n-2)}$ | $e^{\pi i/(k+2n-2)}$ |
| $E_8$                | $\frac{248(k+30)}{60}$            | $e^{\pi i/(k+30)}$   |

**Remark 19.24** ( $\kappa_{\text{cat}}$  versus other invariants). Three distinct invariants must be distinguished:

- (i)  $\kappa_{\text{cat}}$ : the modular characteristic of the CY category, equal to the genus-1 Hodge class coefficient of the associated chiral algebra (Volume I, Theorem D);
- (ii)  $\kappa_{\text{ch}}$ : the modular characteristic computed from the chiral algebra directly (may differ from  $\kappa_{\text{cat}}$  when the CY-to-chiral functor involves additional data);
- (iii)  $\kappa_{\text{BKM}}$ : the BKM modular weight, a number-theoretic invariant arising from the BKM denominator formula (Chapter 14; distinct from both  $\kappa_{\text{cat}}$  and  $\kappa_{\text{ch}}$ ).

For  $\text{CY}_2$  categories with  $\mathfrak{g}$ -symmetry, all three can be independently computed. For the  $K3 \times E$  example:  $\kappa_{\text{ch}} = 3$ ,  $\kappa_{\text{BKM}} = 5$ ,  $\kappa_{\text{cat}} = 3$  (Proposition 10.4).

**PROPOSITION 19.25** (*Complementarity for  $\kappa_{\text{cat}}$* ). <sup>[PE]</sup> Under Koszul duality  $V_k(\mathfrak{g}) \mapsto V_{k'}(\mathfrak{g})$  with  $k' = -k - 2h^\vee$  (Feigin–Frenkel involution):

$$\kappa_{\text{cat}}(\mathcal{C}(\mathfrak{g}, q)) + \kappa_{\text{cat}}(\mathcal{C}(\mathfrak{g}, q')) = 0$$

where  $q' = e^{\pi i/(k'+h^\vee)} = e^{-\pi i/(k+h^\vee)}$ . This is the CY incarnation of the Volume I complementarity  $\kappa_{\text{ch}} + \kappa'_{\text{ch}} = 0$  for KM/free fields (Volume I, Theorem C).

*Proof.* Under  $k \mapsto k' = -k - 2h^\vee$ :  $\kappa'_{\text{cat}} = \dim(\mathfrak{g})(k' + h^\vee)/(2h^\vee) = \dim(\mathfrak{g})(-k - h^\vee)/(2h^\vee) = -\dim(\mathfrak{g})(k + h^\vee)/(2h^\vee) = -\kappa_{\text{cat}}$ .  $\square$



## 19.9 Shadow obstruction tower for quantum group categories

The shadow obstruction tower  $\Theta_{AC}$  (Volume I) acquires categorical meaning through the quantum group lens.

PROPOSITION 19.26 (*Shadow depth from  $R$ -matrix pole structure*). <sup>[PE]</sup> For  $V_k(\mathfrak{g})$  with  $R$ -matrix  $R(z) = 1 + r(z) + O(r^2)$  where  $r(z) = \frac{k\Omega}{z}$ :

- (i) The shadow depth  $r_{\max} = 3$  (class L): the collision residue  $r(z)$  has a single pole, the cubic shadow  $C$  is nonzero (from the Lie bracket), and the quartic resonance class  $Q$  vanishes by semisimplicity of  $\mathfrak{g}$ ;
- (ii) The averaging map (Volume I) sends  $r(z) = \frac{k\Omega}{z} \mapsto \kappa_{\text{cat}}$ : the full  $z$ -dependent profile is killed by  $\Sigma_2$ -coinvariance, leaving only the scalar modular characteristic;
- (iii) The KZ associator  $\Phi_{KZ}$  (the degree-3 component of  $\Theta_{AC}^{E_1}$ ) maps under averaging to the cubic shadow  $C$ .

*Remark 19.27* (From quantum groups to the shadow tower). The passage from quantum group data to the shadow obstruction tower inverts the construction of this chapter: the quantum group  $U_q(\mathfrak{g})$  determines the  $R$ -matrix, which is the degree- $(1, 1)$  component of the  $E_1$  MC element  $\Theta^{E_1}$ , whose  $\Sigma_n$ -coinvariant projections are the shadow tower. The reconstruction problem (from quantum group to chiral algebra) is solved on the evaluation-generated core by the DK bridge (MC<sub>3</sub> for all simple types on that core): the categorical data of  $\text{Rep}_q(\mathfrak{g})$  uniquely determines the chiral algebra  $V_k(\mathfrak{g})$  up to the completion issues of MC<sub>4</sub>.



## **Part IV**

# **The Seven Faces of $r_{\text{CY}}(z)$**



## Chapter 20

# The Bar-Cobar Bridge to Volume I

The shadow obstruction tower  $\Theta_A$  of Volume I applies to chiral algebras arising from the cyclic bar complex of a Calabi–Yau category. The missing link: a concrete dictionary between the CY cyclic bar complex  $CC_\bullet(C)$  and the chiral bar complex  $B(A)$ , together with the shadow tower of  $\mathbb{C}^3$  where both sides are explicit, the passage from finite to infinite shadow depth under the factorization envelope, and the open string field theory realization of Koszul duality.

### 20.1 CY bar complex vs chiral bar complex

Let  $C$  be a smooth proper  $CY_3$  category with Serre functor  $S_C \simeq [3]$ , and let  $A = A_C$  denote the chiral algebra produced by the CY-to-chiral functor  $\Phi$  (Theorem CY-A<sub>2</sub> for  $d = 2$ ; conjectural for  $d = 3$ ). The cyclic bar complex  $CC_\bullet(C)$  and the chiral bar complex  $B(A)$  are related by a canonical quasi-isomorphism

$$CC_\bullet(C) \simeq B(A_C) \quad (\text{CY-A(ii)})$$

as factorization coalgebras on  $\text{Ran}(X)$ .

The precise dictionary is as follows.

**PROPOSITION 20.1** (*Bar complex dictionary*). Under the identification CY-A(ii):

- (i) The cyclic differential  $b + B$  on  $CC_\bullet(C)$  corresponds to the bar differential  $d_B$  on  $B(A)$ .
- (ii) The Connes operator  $B: HH_n(C) \rightarrow HH_{n+1}(C)$  corresponds to the degree-preserving component of  $d_B$  (the internal differential), while the Hochschild differential  $b$  corresponds to the degree-lowering component (the bar contraction).
- (iii) The  $S^1$ -action on  $CC_\bullet(C)$  (the cyclic rotation) corresponds to the factorization coalgebra structure on  $B(A)$ , with the cocomposition maps  $\Delta_\Gamma$  indexed by stable graphs  $\Gamma$ .
- (iv) The Hodge filtration on  $HH_*(C)$  corresponds to the degree filtration on  $B(A)$ : the degree- $r$  piece  $B^{(r)}(A)$  captures Hochschild chains of tensor length  $\leq r$ .

[PH]

**Remark 20.2** (Three functors, three outputs). The bar complex  $B(A)$  is a factorization *coalgebra*. Three distinct functors produce three distinct objects from  $B(A)$ :

- (1)  $\Omega(B(A)) \simeq A$  recovers the original algebra (bar-cobar inversion, Theorem B of Volume I).
- (2)  $D_{\text{Ran}}(B(A)) \simeq B(A^\dagger)$  is the Verdier dual, a factorization *algebra* identified with the bar of the Koszul dual  $A^\dagger$  (Theorem A, Convention conv:bar-coalgebra-identity).
- (3)  $\mathcal{Z}_{\text{ch}}^{\text{der}}(A) = \text{RHom}(\Omega(B(A)), A)$  is the chiral derived center, computing the universal bulk (Theorem H).

These are not the same operation, and the CY cyclic bar complex  $\text{CC}_\bullet(C)$  corresponds to  $B(A)$  itself, not to the derived center and not to the cobar.

## 20.2 The shadow tower of $\mathbb{C}^3$ : MacMahon meets Koszul

The affine threefold  $\mathbb{C}^3$  is the simplest case where both sides of the CY-to-chiral correspondence are explicit.

**THEOREM 20.3** (*Shadow tower of  $\mathbb{C}^3$* ). <sup>[PH]</sup> The shadow obstruction tower of  $W_{1+\infty}$  at  $c = 1$  satisfies:

- (i) The bar Euler product equals the inverse MacMahon function:

$$\sum_{n \geq 0} \dim B_{E_\infty}^{(n)}(A) \cdot q^n = \frac{1}{M(q)}, \quad M(q) = \prod_{n \geq 1} \frac{1}{(1 - q^n)^n}.$$

- (ii) The modular characteristic is  $\kappa_{\text{ch}} = 1$ , verified by five independent paths:

- (a) genus-1 free energy:  $F_1 = \kappa_{\text{ch}}/24 = 1/24$ ;
- (b) degree-0 MacMahon exponent:  $\log M(q) = \sum \sigma_2(k) q^k / k$  matches  $\kappa_{\text{ch}} = 1$ ;
- (c) channel decomposition:  $\sum_{s \geq 1} \kappa_s^{\text{eff}} = \sum_{s \geq 1} s \cdot (1/s) = \sum 1 = +\infty$  (at finite  $N$ ,  $\kappa_{\text{ch}}(W_N) = c(H_N - 1)$ ; the total  $\kappa_{\text{ch}}(W_{1+\infty}, c = 1) = \kappa_{\text{ch}}(H_1) = 1$ );
- (d) Hodge-theoretic: via the categorical trace  $\chi^{\text{CY}}(\text{Perf}(\mathbb{C}^3))$ ;
- (e) complementarity:  $\kappa_{\text{ch}}(\mathbb{C}^3) + \kappa_{\text{ch}}(\text{dual}) = 0$  (free-field anti-symmetry).

- (iii) Per-channel modular characteristics: for the spin- $s$  channel of  $W_{1+\infty}$ ,

$$\kappa_s = \frac{c}{s} = \frac{1}{s}, \quad \kappa_s^{\text{eff}} = s \cdot \kappa_s = 1 \quad (\text{constant per MacMahon level}).$$

This constancy is the shadow-DT Rosetta stone: each energy level contributes equally to the modular characteristic, and the MacMahon function  $M(q) = \prod (1 - q^n)^{-n}$  is the partition function of a system with effective  $\kappa_{\text{ch}} = 1$  per energy level.

Verification: `compute/lib/c3_shadow_tower.py` (73 tests), `compute/lib/macmahon_shadow_decomposition.`

**Remark 20.4** (Envelope creates infinite depth from abelian input). The Heisenberg algebra  $H_1$  (a single free boson) has shadow depth class G with  $r_{\text{max}} = 2$ : the tower terminates at degree 2, and  $\kappa_{\text{ch}} = 1$  is the only nonvanishing shadow invariant.

The vertex algebra  $W_{1+\infty}$  at  $c = 1$ , the factorization envelope of the Lie conformal algebra of polyvector fields on  $\mathbb{C}^3$ , has shadow depth class M with  $r_{\text{max}} = \infty$ . The spin-2 channel (Virasoro at  $c = 1$ ) already has infinite shadow depth, with  $\kappa_2 = 1/2$ ,  $\alpha_2 = 2$ ,  $S_4 = 10/27$ .

This discrepancy between input (class G, depth 2) and output (class M, depth  $\infty$ ) reflects the non-conservativity of the factorization envelope on shadow invariants. Normal ordering, Wick contractions, and OPE singular terms generate higher-degree shadows invisible at the  $E_1$ -level. Each stage of  $E_1 \rightarrow E_2 \rightarrow E_\infty$ -chiral introduces genuinely new invariants.

### 20.3 Open string field theory: Koszul duality at chain level

For  $\mathbb{C}^3$ , the cyclic  $A_\infty$ -algebra is the exterior algebra  $A = \bigwedge^*(V)$  with  $V = \mathbb{C}^3$ , equipped with the  $CY_3$  trace  $\langle a, b \rangle = \text{Tr}(a \wedge b)$  (projection to top degree).

**THEOREM 20.5** (*Chain-level Koszul duality for  $\mathbb{C}^3$* ). [PH] The Koszul duality pair for  $A = \bigwedge^*(\mathbb{C}^3)$  satisfies:

- (i)  $A^! = \text{Sym}^*(\mathbb{C}^{3,*})$ , the symmetric algebra on the dual vector space. This is the  $\text{Com}^! = \text{Lie}$  incarnation at the chain level: the exterior algebra (free  $\text{Com}$ -algebra on  $V$ ) is Koszul dual to the symmetric algebra (free  $\text{Lie}$ -algebra envelope on  $V^*$ ).
- (ii) The bar complex  $B(\bigwedge^* V)$  has cohomology concentrated in bar degree 1:

$$H^*(B(\bigwedge^* V)) \simeq \text{Sym}^*(V^*[-1]),$$

confirming that  $A$  is Koszul (chirally and classically).

- (iii) The bar-cobar round-trip recovers  $A$  itself:

$$\Omega(B(\bigwedge^* V)) \xrightarrow{\sim} \bigwedge^* V.$$

Verification: `compute/lib/bar_comparison_c3.py`, `compute/lib/e2_bar_cobar_koszul.py` (73 tests).

### 20.4 From shuffle product to $\lambda$ -bracket: the seven-step chain

The Schiffmann–Vasserot shuffle presentation of  $Y^+(\widehat{\mathfrak{gl}}_1)$  does *not* directly produce the  $\lambda$ -bracket of  $R_{\mathbb{C}^3}$ . Seven comparison steps are required.

**Construction 20.6** (*The seven-step chain*). The chain connecting the shuffle algebra to the chiral algebra:

1. **Shuffle algebra**  $\text{Sh}(h_1, h_2, h_3)$ : the associative algebra with generators  $e(z)$  at each spectral parameter  $z$  and multiplication by the rational shuffle kernel  $\zeta(z_i - z_j) = \prod_a (z_i - z_j + h_a) / (z_i - z_j - h_a)$ .
2. **Affine Yangian**  $Y(\widehat{\mathfrak{gl}}_1)$ : the RTT-presentation with generators  $e_j, f_j, \psi_j$  satisfying the affine Yangian relations. Schiffmann–Vasserot proved  $\text{Sh}^+ \simeq Y^+(\widehat{\mathfrak{gl}}_1)$ .
3.  $W_{1+\infty}$  **modes**: the Feigin–Frenkel realization of  $W_{1+\infty}$  as the limit  $\lim_{N \rightarrow \infty} W_N$ , with explicit mode algebra generators  $W_n^s$  for spin  $s$  and mode  $n$ .
4. **OPE modes**: the vertex algebra  $\text{OPE } W^s(z) W^{s'}(w) \sim \sum_n C_n^{ss'}(W) (z - w)^{-n-1}$ , extracting the singular coefficients  $C_n^{ss'}$ .
5.  **$\lambda$ -bracket**:  $\{W_\lambda^s W^{s'}\} = \sum_n \lambda^{(n)} C_n^{ss'}(W)$ , where  $\lambda^{(n)} = \lambda^n / n!$  is the divided power.
6. **Lie conformal algebra**  $R_{\mathbb{C}^3}$ : the  $\text{GL}(3)$ -invariant subalgebra of the Schouten–Nijenhuis Lie conformal algebra on polyvector fields  $\text{PV}^*(\mathbb{C}^3)$ .

7. **Factorization envelope**  $U^{\text{ch}}(R_{\mathbb{C}^3})$ : the enveloping vertex algebra, which by the Nishinaka–Vicedo construction recovers  $W_{1+\infty}$ .

Steps  $1 \rightarrow 2$  is Schiffmann–Vasserot. Steps  $2 \rightarrow 3$  is Prochazka–Rapcak. Steps  $3 \rightarrow 4$  is the standard mode-to-OPE dictionary. Step  $4 \rightarrow 5$  requires the divided-power convention. Steps  $5 \rightarrow 6$  is the Kontsevich–Soibelman identification of  $R_{\mathbb{C}^3}$  as the  $\text{GL}(3)$ -invariant part of  $\text{PV}^*(\mathbb{C}^3)$  with the Schouten–Nijenhuis bracket. Step  $6 \rightarrow 7$  is the factorization envelope.

The composition  $1 \rightarrow 7$  is the full CY-to-chiral functor for  $\mathbb{C}^3$ .

Verification: `compute/lib/c3_lie_conformal.py, compute/lib/c3_envelope_comparison.py` (52 tests).

*Warning 20.7* (Shuffle product  $\neq \lambda$ -bracket). The shuffle product on  $\text{Sh}(h_1, h_2, h_3)$  is an associative product on symmetric functions of the spectral parameters, while the  $\lambda$ -bracket is a Lie conformal bracket on fields. The two are related by the seven-step chain above, not by a direct identification. In particular, the shuffle kernel  $\zeta(z)$  is *not* the OPE kernel: the OPE has poles at  $z = 0$  (collision singularities), while the shuffle kernel has poles at  $z = \pm h_a$  (Yangian structure function poles). These coincide only after the identification of the parameters  $h_a$  with the deformation parameters of  $W_{1+\infty}$  and the passage from additive to multiplicative spectral parameters.

## 20.5 Koszul duality: CY vs chiral

The chiral Koszul duality of Volume I (Theorems A and B) descends to CY Koszul duality.

CONJECTURE 20.8 (*CY Koszul duality dictionary*). <sup>[CJ]</sup> For a  $\text{CY}_3$  category  $C$  with chiral algebra  $A = A_C$ :

- (i) The chiral Koszul dual  $A^!$  corresponds to the *mirror* CY category:

$$A_C^! \simeq A_{C^!}$$

where  $C^!$  is the Koszul dual CY category (for  $C = D^b(\text{Coh}(X))$  of a CY manifold  $X$ , this is  $C^! \simeq \text{Fuk}(X^\vee)$  under homological mirror symmetry).

- (ii) The complementarity theorem (Theorem C) gives:

$$Q_g(A_C) + Q_g(A_{C^!}) = H^*(\overline{\mathcal{M}}_g, \mathcal{Z}(A_C))$$

where  $\mathcal{Z}(A_C) = \mathcal{Z}_{\text{ch}}^{\text{der}}(A_C)$  is the chiral derived center.

- (iii) The modular characteristics satisfy:

$$\kappa_{\text{ch}}(A_C) + \kappa_{\text{ch}}(A_{C^!}) = 0$$

when  $C$  arises from a free-field or lattice construction (anti-symmetry), and more generally  $\kappa_{\text{ch}} + \kappa'_{\text{ch}} = \rho \cdot K$  for algebras with nontrivial contact terms (Theorem D).



## 20.6 The five theorems in the CY setting

We summarize the status of the five main theorems of Volume I when specialized to chiral algebras arising from CY categories.

**THEOREM** (*The five theorems for CY chiral algebras*). Let  $A = A_C$  be the chiral algebra of a  $\text{CY}_3$  category  $C$ . Then:

**Theorem A (adjunction).** The bar-cobar adjunction  $B \dashv \Omega$  restricts to CY chiral algebras:  $B(A_C)$  is a factorization coalgebra on  $\text{Ran}(X)$ , and  $D_{\text{Ran}}(B(A_C)) \simeq B(A_C^\dagger)$ . The CY identification  $\text{CY-A(ii)}$  gives  $\text{CC}_\bullet(C) \simeq B(A_C)$ .

**Theorem B (inversion).** Bar-cobar inversion  $\Omega(B(A_C)) \xrightarrow{\sim} A_C$  holds on the Koszul locus. For CY categories, chirally Koszul is equivalent to the formality of  $\text{CC}_\bullet(C)$  as a dg coalgebra.

**Theorem C (complementarity).** The CY Euler characteristic  $\chi(C)$  splits into complementary halves:  $Q_g(C) + Q_g(C^\dagger)$  recovers the full Hochschild cohomology.

**Theorem D (modular characteristic).** The genus- $g$  obstruction is  $\text{obs}_g(A_C) = \kappa_{\text{ch}}(A_C) \cdot \lambda_g^{\text{FP}}$  on the uniform-weight lane, where  $\kappa_{\text{ch}}(A_C) = \chi^{\text{CY}}(C)$  is the CY modular characteristic. For rigid compact CICYs with  $h^{1,0} = 0$ , the BCOV prediction gives  $\kappa_{\text{ch}} = \chi_{\text{top}}/24$ ; for  $K_3$ -fibered  $\text{CY}_3$ , this fails (Theorem 8.71).

**Theorem H (Hochschild).**  $\text{ChirHoch}^*(A_C)$  is polynomial in degrees  $\{0, 1, 2\}$ , and the CY Hochschild calculus of  $C$  (HKR decomposition, Connes operator, categorical Hodge structure) is faithfully reflected in  $\text{ChirHoch}^*(A_C)$ .

## 20.7 Compact $\text{CY}_3$ examples

The following compact  $\text{CY}_3$  families illustrate the range of shadow tower behaviour. In each case, the predicted modular characteristic is  $\kappa_{\text{ch}} = \chi_{\text{top}}/24$  (BCOV prediction); the shadow depth class, Hochschild data, and BKM structure vary.

### The banana manifold: $\kappa_{\text{ch}} = 0$ with nontrivial tower

The banana manifold  $X_{\text{ban}}$  (Schoen's  $\text{CY}_3$ , the fiber product of two rational elliptic surfaces over  $\mathbb{P}^1$ ) has Hodge numbers  $h^{1,1} = h^{2,1} = 19$  and Euler characteristic  $\chi = 0$ .

**PROPOSITION 20.9** (*Shadow tower of the banana manifold*). The shadow obstruction tower of  $X_{\text{ban}}$  has:

- (i)  $\kappa_{\text{ch}} = \chi/24 = 0$ . The degree-2 scalar shadow *vanishes*.
- (ii)  $S_4 = -44$  (quartic shadow from the genus-0 GV invariants  $n_{d_1, d_2}^0 = -2$  of the banana curve classes).
- (iii) The shadow tower starts at degree 4, not degree 2: the cubic shadow is invisible when  $\kappa_{\text{ch}} = 0$  (it enters as  $\kappa_{\text{ch}} \cdot \alpha$ , which vanishes).

- (iv) The critical discriminant  $\Delta = 8\kappa_{\text{ch}}S_4 = 0$  (since  $\kappa_{\text{ch}} = 0$ ). The standard single-line classification (G/L/C/M) does not directly apply.
- (v) Shadow depth class: M (infinite tower) shifted to degree 4. The BKY partition function involves infinitely many banana-curve BPS states, generating an infinite shadow tower whose leading term is quartic.

[PH]

*Remark 20.10* ( $\kappa_{\text{ch}} = 0$  does not imply  $\Theta_A = 0$ ). The banana manifold is the prototypical example:  $\kappa_{\text{ch}} = 0$  does not imply  $\Theta_A = 0$ . The vanishing of  $\kappa_{\text{ch}}$  means the bar complex is uncurved ( $d^2 = 0$  at the leading scalar level), but the higher-degree components of  $\Theta_A$  (the quartic shadow  $Q$  and beyond) are nonvanishing, sourced by the instanton contributions (genus-0 GV invariants of the banana curves).

Three  $\text{CY}_3$  examples with  $\kappa_{\text{ch}} = 0$  show qualitatively different behaviour:

- Heisenberg at  $k = 0$ :  $\kappa_{\text{ch}} = 0$ , trivially uncurved, class G.
- Virasoro at  $c = 0$ :  $\kappa_{\text{ch}} = 0$ , but higher shadows from contact terms (class M).
- Banana manifold:  $\kappa_{\text{ch}} = 0$ , but higher shadows from *instantons* (class M, shifted to degree 4).

Verification: `compute/lib/banana_shadow.py` (63 tests).

### Enriques $\times E$ : the $\mathbb{Z}_2$ quotient of $\text{K}_3 \times E$

Let  $S_{\text{Enr}} = \text{K}_3/\iota$  be an Enriques surface ( $\iota$  a fixed-point-free involution), and  $E$  an elliptic curve. The product  $S_{\text{Enr}} \times E$  is a generalized  $\text{CY}_3$  (the canonical bundle  $K_{S_{\text{Enr}} \times E}$  is 2-torsion, so  $h^{3,0} = 0$ ). Nevertheless, the shadow tower machinery applies to the associated vertex algebra, and the Borchers lift on  $O(2, 10)$  provides the automorphic structure.

**THEOREM 20.11** (*Shadow tower of Enriques  $\times E$* ). [PH] The shadow data of Enriques  $\times E$  is:

- (i)  $\kappa_{\text{BKM}}(\text{Enr} \times E) = 4$ , the weight of the Enriques Borchers product  $\Psi$  on  $O(2, 10)$  (Allcock 2000).
- (ii) Root multiplicities come from the Enriques elliptic genus  $\phi_{\text{Enr}} = \frac{1}{2}\phi_{0,1}$ : the Fourier coefficients are  $f_{\text{Enr}}(n, l) = f_{\text{K}_3}(n, l)/2$  (all integers since  $f_{\text{K}_3}$  has even coefficients).
- (iii) The ratio  $\kappa_{\text{BKM}}(\text{K}_3 \times E)/\kappa_{\text{BKM}}(\text{Enr} \times E) = 5/4$ .
- (iv) Shadow depth class: M (infinite tower), driven by the infinite BKM root spectrum.

*Remark 20.12* (The  $\mathbb{Z}_2$  quotient on  $\kappa_{\text{BKM}}$ ). The passage from  $\text{K}_3 \times E$  ( $\kappa_{\text{BKM}} = 5$ ) to Enriques  $\times E$  ( $\kappa_{\text{BKM}} = 4$ ) under the  $\mathbb{Z}_2$  quotient does *not* halve  $\kappa_{\text{BKM}}$ . Rather, the weight of the Borchers product drops by 1: the  $O(2, 10)$  lattice for the Enriques surface has a different constant term in the Borchers input form than the  $O(3, 19)$  lattice for  $\text{K}_3$ , and the weight formula gives  $8/2 = 4$  instead of  $10/2 = 5$ . The ratio  $5/4$  is not a simple topological quotient; it reflects the lattice-theoretic structure of the Borchers lift.

Verification: `compute/lib/enriques_shadow.py` (72 tests).

### The quintic threefold: non-integer $\kappa_{\text{ch}}$ and BKM obstruction

The quintic hypersurface  $Q \subset \mathbb{P}^4$  has  $h^{1,1} = 1$ ,  $h^{2,1} = 101$ ,  $\chi = -200$ .

PROPOSITION 20.13 (*Quintic shadow tower*). The shadow data of the quintic is:

- (i)  $\kappa_{\text{ch}} = \chi/24 = -25/3$  (non-integer).
- (ii) The non-integrality of  $\kappa_{\text{ch}}$  obstructs a naive BKM superalgebra: BKM root multiplicities must be integers, and the standard denominator-identity machinery requires integral Borcherds lift weight. The quintic has no natural Borcherds–Kac–Moody structure.
- (iii) The shadow metric  $Q_L(t) = 4\kappa_{\text{ch}}^2 + 12\kappa_{\text{ch}}\alpha t + (9\alpha^2 + 16\kappa_{\text{ch}}S_4)t^2$  is well-defined for fractional  $\kappa_{\text{ch}}$ . With  $\kappa_{\text{ch}} = -25/3$ :  $Q_L(0) = 2500/9 > 0$ .
- (iv) The GW invariants are known to high degree:

$$n_1^0 = 2875, \quad n_2^0 = 609250, \quad n_3^0 = 317206375, \quad n_4^0 = 242467530000.$$

These are BPS state counts via the Gopakumar–Vafa formula.

- (v) Shadow depth class: M (infinite tower). The quintic has an infinite BPS spectrum with curves of all degrees and genera; the shadow tower is controlled by the BCOV holomorphic anomaly equation.

[PH]

Remark 20.14 ( $\chi/24$  integrality survey). The quantity  $\chi/24$  is an integer if and only if  $12 \mid (h^{1,1} - h^{2,1})$ . Among standard complete intersections:

- $\mathbb{P}^5[3, 3]$ :  $\chi = -144$ ,  $\kappa_{\text{ch}} = -6$  (integer).
- $\mathbb{P}^5[2, 4]$ :  $\chi = -176$ ,  $\kappa_{\text{ch}} = -22/3$  (non-integer).
- $\mathbb{P}^7[2, 2, 2, 2]$ :  $\chi = -128$ ,  $\kappa_{\text{ch}} = -16/3$  (non-integer).
- Quintic:  $\chi = -200$ ,  $\kappa_{\text{ch}} = -25/3$  (non-integer).

Non-integrality of  $\kappa_{\text{ch}}$  is the generic situation for compact CY<sub>3</sub> manifolds. A BKM structure requires additional input beyond the BCOV prediction.

Verification: `compute/lib/quintic_shadow_obstruction.py` (86 tests).

### Hochschild homology of compact CY<sub>3</sub> examples

The Hochschild homology  $\text{HH}_*(D^b(\text{Coh}(X)))$  is computed from the HKR decomposition  $\text{HH}_p \simeq \bigoplus_q H^q(X, \Omega_X^{3-p})$  using the CY<sub>3</sub> isomorphism  $\bigwedge^p T_X \simeq \Omega_X^{3-p}$ .

PROPOSITION 20.15 (*Hochschild data of compact CY<sub>3</sub> examples*). (i)  $X = \text{K3} \times E$ :  $\dim \text{HH}_* = 96$ , with Hodge decomposition

$$(\dim \text{HH}_0, \dim \text{HH}_1, \dim \text{HH}_2, \dim \text{HH}_3) = (4, 44, 44, 4).$$

The palindromic symmetry  $\dim \text{HH}_p = \dim \text{HH}_{3-p}$  reflects Serre duality.

(ii)  $X = \mathcal{Q}$  (quintic):  $\dim \mathrm{HH}_* = 208$ , with

$$(\dim \mathrm{HH}_0, \dim \mathrm{HH}_1, \dim \mathrm{HH}_2, \dim \mathrm{HH}_3) = (2, 102, 102, 2).$$

The concentration in degrees 1 and 2 reflects  $h^{1,1} = 1$  and  $h^{2,1} = 101$ : moduli dominate, and the CY structure contributes only the top and bottom forms.

[PH]

Verification: `compute/lib/cy3_hochschild.py` (71 tests).

## 20.8 Integrable systems and lattice models

The shadow obstruction tower admits a dictionary with classical integrable systems, where the conserved quantities  $I_r$  correspond to shadow invariants at each degree.

### The Calogero–Moser/Bethe dictionary

The Calogero–Moser system with  $N$  particles on a line and inverse-square interaction  $V(x) = \sum_{i < j} g^2 / (x_i - x_j)^2$  has  $N$  independent conserved quantities  $I_1, I_2, \dots, I_N$  (the Sekiguchi operators), with  $I_2 = H$  (the Hamiltonian) and  $I_k$  of degree  $k$  in the momenta.

CONJECTURE 20.16 (*Shadow-integrable dictionary*). [CJ] There is a dictionary between the shadow obstruction tower invariants and the Calogero–Moser conserved quantities:

$$\begin{aligned} I_2 \text{ (quadratic Hamiltonian)} &\longleftrightarrow \kappa_{\mathrm{ch}} \text{ (modular characteristic),} \\ I_3 \text{ (cubic integral)} &\longleftrightarrow C \text{ (cubic shadow),} \\ I_4 \text{ (quartic integral)} &\longleftrightarrow Q \text{ (quartic shadow / contact invariant).} \end{aligned}$$

The Bethe ansatz equations for the Calogero–Moser system correspond to the Maurer–Cartan equation  $D\Theta + \frac{1}{2}[\Theta, \Theta] = 0$  of the modular convolution algebra, projected to each degree: the  $r$ -th conserved quantity  $I_r$  is the genus-0 degree- $r$  projection of the MC element  $\Theta_A$ .

*Remark 20.17.* The dictionary is *structural*, not a theorem. It captures the parallel between two filtration structures: the degree filtration on  $\Theta_A$  (controlling the shadow depth classification G/L/C/M) and the degree filtration on the conserved quantities  $I_r$  (controlling the integrability class of the Calogero–Moser system). The coupling constant  $g^2$  maps to  $\kappa_{\mathrm{ch}}$ , the number of particles  $N$  maps to the number of strong generators, and the Bethe roots map to the shadow metric zeros.

Verification: `compute/lib/cross_volume_shadow_bridge.py` (60 tests).

### Lattice finite-size corrections

For a critical lattice model (e.g., the XXX Heisenberg spin chain of length  $L$ ), the finite-size corrections to the ground-state energy are controlled by the central charge  $c$  and conformal dimensions of the associated CFT:

$$E_0(L) = L\varepsilon_\infty - \frac{\pi c}{6L} + O(L^{-3}).$$

More generally, the full finite-size spectrum organizes into an asymptotic expansion

$$E_n(L) - L\varepsilon_\infty = \frac{2\pi}{L} \left( -\frac{c}{12} + h_n + \bar{h}_n \right) + \sum_{k \geq 2} \frac{a_k(n)}{L^{2k-1}}.$$

**CONJECTURE 20.18** (*Finite-size corrections as shadow tower*). The finite-size correction coefficients  $a_k(n)$  correspond to shadow tower data: the leading coefficient  $a_1 = -c/12 = -\kappa_{\text{ch}}/6$  recovers  $\kappa_{\text{ch}}$ , and the higher coefficients  $a_k$  for  $k \geq 2$  encode the degree- $(2k)$  shadow projections. The decay rate of the corrections is  $q = e^{-2\pi/L}$ , matching the shadow tower variable. <sup>[HE]</sup>

Verification: `compute/lib/cross_volume_shadow_bridge.py` (63 tests).

### Crystal melting: combinatorial vs bar-algebraic degree

The MacMahon function  $M(q)$  counts plane partitions (3D Young diagrams), and the crystal melting model of Okounkov–Reshetikhin–Vafa interprets the DT partition function of  $\mathbb{C}^3$  as a statistical-mechanical partition function on the crystal lattice  $\mathbb{Z}_{\geq 0}^3$ .

*Warning 20.19* (Combinatorial degree  $\neq$  bar degree). The crystal melting “degree” (the number of boxes removed from the corner of the crystal) does *not* match the bar complex degree (the tensor length in  $B^{(n)}(A)$ ).

The crystal partition function organizes by the number of boxes:

$$M(q) = \sum_{\pi \in \mathcal{PP}} q^{|\pi|}, \quad |\pi| = \text{number of boxes in plane partition } \pi.$$

The bar complex organizes by tensor length:

$$B(A) = \bigoplus_{n \geq 1} \bar{A}^{\otimes n}[n], \quad \bar{A} = \ker(\varepsilon), \quad \text{degree } n = \text{number of tensor factors}.$$

These two gradings are *not* compatible: a plane partition with  $k$  boxes does not correspond to a bar element of degree  $k$ . Rather, the crystal melting expansion is an exponential reorganization of the bar complex:  $\log M(q) = \sum_k \sigma_2(k) q^k / k$  is the free energy, while the bar complex detects the individual OPE modes.

The fundamental mismatch: crystal melting is a *tensor-algebraic* construction (counting 3D diagrams in  $\mathbb{Z}^3$ ), while the bar complex is a *factorization-algebraic* construction (organized by collision patterns on  $\text{Ran}(X)$ ). The two agree in the graded *character* (both reproduce  $M(q)$  or its inverse) but not in the *degree structure*.

Verification: `compute/lib/crystal_bar_identification.py` (70 tests).

## 20.9 Grand atlas of CY<sub>3</sub> shadow invariants

Table 20.1 collects the shadow tower data for 18 CY<sub>3</sub> families spanning the full range of shadow behaviour.

Key patterns visible in the atlas:

1. **Toric depth bound:** for toric CY<sub>3</sub> with  $N$  vertices in the web diagram, the shadow depth satisfies  $r_{\text{max}} \leq N$ . The cases  $N = 1$  (trivial), 2 (Gaussian), 3 (Lie), 4 (contact) are realized;  $N \geq 5$  gives class M.
2. **Compact CY<sub>3</sub> are generically class M:** all compact CY<sub>3</sub> with  $\chi \neq 0$  have infinite shadow depth, driven by the infinite BPS spectrum.
3.  **$\kappa_{\text{ch}}$  integrality:** among the atlas families,  $\mathbb{P}^5[3, 3]$  ( $\kappa_{\text{ch}} = -6$ ),  $K_3 \times E$  ( $\kappa_{\text{BKM}} = 5$ ), Enriques  $\times E$  ( $\kappa_{\text{BKM}} = 4$ ), and  $\text{BV}(20, 2, 0)$  ( $\kappa_{\text{ch}} = 5$ ) have integral  $\kappa_{\text{ch}}$ . For compact CICYs,  $\kappa_{\text{ch}} = \chi/24$  is integral only when  $24 \mid \chi$ ; for K<sub>3</sub>-fibered geometries,  $\kappa_{\text{BKM}}$  (the automorphic weight) is always integral.

Table 20.1: Grand atlas of  $\text{CY}_3$  shadow invariants. Columns:  $\text{CY}_3$  family, topological Euler characteristic  $\chi$ , Hodge numbers  $h^{1,1}$  and  $h^{2,1}$ , modular characteristic  $\kappa_{\text{ch}}$ , shadow depth class, and data source. For non-compact toric  $\text{CY}_3$ :  $\kappa_{\text{ch}} = \chi(S)/2$  where  $S$  is the compact base (Proposition 15.17). For compact CICYs:  $\kappa_{\text{ch}} = \chi_{\text{top}}/24$ . For  $\text{K}_3$ -fibered geometries: the column records  $\kappa_{\text{BKM}}$  (the automorphic weight), not  $\kappa_{\text{ch}}$ ; see Remark 20.12. Families above the first rule are non-compact toric; between the rules are  $\text{K}_3$ -fibered; below are compact complete intersections and special families. BV = Borcea–Voisin.

| $\text{CY}_3$                              | $\chi$ | $h^{1,1}$ | $h^{2,1}$ | $\kappa_{\text{ch}}$ | Class        | Source                    |
|--------------------------------------------|--------|-----------|-----------|----------------------|--------------|---------------------------|
| $\mathbb{C}^3$                             | —      | —         | —         | 1                    | G            | MacMahon                  |
| Resolved conifold                          | —      | 1         | 0         | 1                    | G            | KS wall-crossing          |
| Local $\mathbb{P}^2$                       | —      | 1         | 0         | 3/2                  | M            | McKay( $\mathbb{Z}_3$ )   |
| Local $\mathbb{P}^1 \times \mathbb{P}^1$   | —      | 2         | 0         | 2                    | M            | McKay( $\mathbb{Z}_2^2$ ) |
| Local $F_1$                                | —      | 2         | 0         | −2                   | M            | Hirzebruch                |
| $\text{K}_3 \times E$                      | 0      | 21        | 21        | 5 <sup>†</sup>       | M            | $\Delta_5$ / BKM          |
| Enriques $\times E$                        | 0      | 11        | 11        | 4 <sup>†</sup>       | M            | Allcock / $O(2, 10)$      |
| Quintic                                    | −200   | 1         | 101       | −25/3                | M            | BCOV                      |
| $\mathbb{P}^5[3, 3]$                       | −144   | 1         | 73        | −6                   | M            | BCOV                      |
| $\mathbb{P}^5[2, 4]$                       | −176   | 1         | 89        | −22/3                | M            | BCOV                      |
| $\mathbb{P}^6[2, 2, 3]$                    | −144   | 1         | 73        | −6                   | M            | BCOV                      |
| $\mathbb{P}^7[2, 2, 2, 2]$                 | −128   | 1         | 65        | −16/3                | M            | BCOV                      |
| Bicubic $\mathbb{P}^2 \times \mathbb{P}^2$ | −162   | 2         | 83        | −27/4                | M            | BCOV                      |
| Banana (Schoen)                            | 0      | 19        | 19        | 0                    | $M^{\geq 4}$ | BKY                       |
| BV(1, 1, 1)                                | −108   | 0         | 54        | −9/2                 | M            | Voisin–Borcea             |
| BV(10, 0, 0)                               | 0      | 35        | 35        | 0                    | ?            | Voisin–Borcea             |
| BV(10, 8, 0)                               | 0      | 19        | 19        | 0                    | ?            | Voisin–Borcea             |
| BV(20, 2, 0)                               | 120    | 61        | 1         | 5                    | M            | Voisin–Borcea             |

<sup>†</sup>These entries record  $\kappa_{\text{BKM}}$  (Borcherds automorphic weight), not  $\kappa_{\text{ch}}$ ; for  $\text{K}_3 \times E$ ,  $\kappa_{\text{ch}} = 3$ .

- BKM structure requires integral automorphic weight:**  $\text{K}_3$ -fibered  $\text{CY}_3$ s with  $\chi = 0$  carry BKM superalgebras via Borcherds lifts; compact CICYs with  $\kappa_{\text{ch}} \notin \mathbb{Z}$  do not.
- $\kappa_{\text{ch}} = 0$  **does not imply trivial tower:** the banana manifold and the balanced Borcea–Voisin families ( $r = 10$ ) have  $\kappa_{\text{ch}} = 0$  but potentially nontrivial higher-degree shadows sourced by instantons.
- Borcea–Voisin as  $\kappa_{\text{ch}}$ -interpolation:** the BV family parametrized by  $(r, a, \delta)$  gives  $\kappa_{\text{ch}} = (r - 10)/2$ , interpolating between  $\kappa_{\text{ch}} = -9/2$  (minimal,  $r = 1$ ) and  $\kappa_{\text{ch}} = 5$  (maximal,  $r = 20$ , recovering  $\text{K}_3 \times E$ ).

Verification: `compute/lib/cy3_grand_atlas.py, compute/tests/test_cy3_grand_atlas.py` (119 tests).

*Remark 20.20* (Enriques  $\times E$  kappa anomaly). The grand atlas uses  $\kappa_{\text{BKM}}(\text{Enr} \times E) = 4$ , the weight of the Allcock Borcherds product on  $O(2, 10)$ , verified by `enriques_shadow.py` (72 tests). This is the authoritative value. The naive BCOV formula  $\kappa_{\text{ch}} = \chi/24$  does not apply because Enriques  $\times E$  is a *generalized*  $\text{CY}_3$  ( $h^{3,0} = 0$ , torsion canonical). The ratio  $\kappa_{\text{BKM}}(\text{K}_3 \times E)/\kappa_{\text{BKM}}(\text{Enr} \times E) = 5/4$ .

## Chapter 2I

# The seven faces of $r_{CY}(z)$ for Calabi–Yau chiral algebras

*For a Calabi–Yau chiral algebra arising from the CY-to-chiral functor, the binary collision residue  $r_{CY}(z) = \text{Res}_{0,2}^{\text{coll}}(\Theta_{CY})$  has seven equivalent realizations specific to the CY setting: as the CY bar-cobar twisting morphism, as a CoHA / perverse-coherent-sheaves shadow, as the classical CY Poisson coisson, as the Maulik–Okounkov R-matrix (for  $K3 \times E$ ), as the affine super Yangian  $Y(\widehat{\mathfrak{gl}}_1)$  r-matrix (for toric  $CY_3$ ), as the elliptic Sklyanin bracket (for toroidal CY), and as a Gaudin model arising from  $CY_3$  collision residues. The seven-face identification in the CY setting specializes Vol I’s general theorems to the  $CY_3$  cases.*

### 2I.1 The $CY_3$ collision residue

The collision residue is a single algebraic operation extracted from the universal Maurer–Cartan element  $\Theta_A$  of a chiral algebra. In the CY setting, this operation acquires geometric meaning: the residue is computed against the holomorphic CY volume form, and the resulting binary operation is the classical limit of the quantum vertex chiral group braiding.

*Definition 2I.1 (Binary collision residue, CY version).* Let  $\mathcal{C}$  be a smooth proper Calabi–Yau category of dimension  $d$  with holomorphic volume form  $\Omega_{\mathcal{C}}$ . Let  $A_{\mathcal{C}}$  denote the chiral algebra produced by the CY-to-chiral functor (Theorem CY-A<sub>2</sub> at  $d = 2$ ; Conjecture CY-A<sub>3</sub> at  $d = 3$ ). Let  $\Theta_{A_{\mathcal{C}}} \in \text{MC}(\mathfrak{g}_{A_{\mathcal{C}}}^{\text{mod}})$  denote its universal Maurer–Cartan element (cf. Vol I, Theorem III). The *binary CY collision residue* is

$$r_{CY}(z) := \text{Res}_{0,2}^{\text{coll}}(\Theta_{A_{\mathcal{C}}}) \in A_{\mathcal{C}} \otimes A_{\mathcal{C}}[[z^{-1}]],$$

the genus-0, degree-2 residue along the collision divisor of  $\overline{C}_2(\text{Spec } \mathbb{C})$ , paired against  $\Omega_{\mathcal{C}}$  in the CY direction.

*Remark 2I.2* (The CY direction is a separate slot). The collision residue extracts the  $z^{-1}$  part of  $\Theta$  along the chiral curve direction. In the CY setting,  $\Theta$  has *two* gradings: degree in the chiral direction (controlled by the curve  $X$ ) and Hodge weight in the CY direction (controlled by  $\mathcal{O}_{\mathcal{C}}$ ). The CY volume form  $\Omega_{\mathcal{C}}$  kills the Hodge weight by integration; what remains is a chiral binary operation valued in  $A_{\mathcal{C}} \otimes A_{\mathcal{C}}$ . This is the precise sense in which  $r_{CY}$  couples the two directions.

*Remark 2I.3* (Three invariants must not be conflated). For any chiral algebra  $A$ , the binary residue carries three *distinct* numerical invariants which the CY setting will illuminate:

- (i)  $p_{\max}(A)$ , the maximal pole order of generator-on-generator OPEs (cf. Vol I, Definition 20.9);
- (ii)  $k_{\max}(A) = p_{\max}(A) - 1$ , the collision depth of  $r_{CY}$  itself, after  $d$  log-absorption (cf. Vol I, Definition 20.9);
- (iii)  $r_{\max}(A)$ , the shadow depth, the degree at which the obstruction tower  $\Theta_A^{\leq r}$  terminates (Vol I, Definition 20.9).

The  $\beta\gamma$  system is the archetypal witness of their independence:  $p_{\max}(\beta\gamma) = 1$ ,  $k_{\max}(\beta\gamma) = 0$ ,  $r_{\max}(\beta\gamma) = 4$  (class C). Conflating these produces incorrect classifications, and the seven-face programme will distinguish them at every face.

## 21.2 Face I: the CY bar-cobar twisting morphism

The first face is the home framework of this volume. The CY-to-chiral functor sends a Calabi–Yau category  $\mathcal{C}$  to a chiral algebra  $A_{\mathcal{C}}$ , and the bar construction sends  $A_{\mathcal{C}}$  to a factorization coalgebra  $\overline{B}(A_{\mathcal{C}})$  on  $\text{Ran}(X)$ . The collision residue is the binary component of the universal twisting morphism between  $\overline{B}(A_{\mathcal{C}})$  and  $A_{\mathcal{C}}$ .

**THEOREM 21.4** (*Face I: CY bar-cobar realization,  $d = 2$* ). <sup>[PH]</sup> Let  $A_{\mathcal{C}}$  be the chiral algebra of a CY<sub>2</sub> category  $\mathcal{C}$  produced by the CY-to-chiral functor  $\Phi$  (Theorem CY-A<sub>2</sub>). Then:

- (i) The convolution dg Lie algebra  $\mathfrak{g}_{A_{\mathcal{C}}}^{\text{mod}}$  identifies with the homotopy Lie algebra of twisting morphisms  $\text{hom}_{\alpha}(\overline{B}(A_{\mathcal{C}}), A_{\mathcal{C}})$  between bar coalgebra and chiral algebra (cf. Vol I, Theorem III);
- (ii) The universal MC element  $\Theta_{A_{\mathcal{C}}}$  is the universal twisting morphism;
- (iii) The binary collision residue  $r_{CY}(z) = \text{Res}_{0,2}^{\text{coll}}(\Theta_{A_{\mathcal{C}}})$  is the leading degree-2 component of this twisting morphism, evaluated in the CY direction against  $\Omega_{\mathcal{C}}$ .

*Proof sketch.* For (i), the cyclic bar complex  $\text{CC}_{\bullet}(\mathcal{C})$  identifies with  $\overline{B}(A_{\mathcal{C}})$  as a factorization coalgebra (Theorem 20.1). The convolution identification is the application of Vol I’s master twisting-morphism theorem to this coalgebra-algebra pair. For (ii),  $\Theta_{A_{\mathcal{C}}}$  satisfies the MC equation  $D\Theta + \frac{1}{2}[\Theta, \Theta] = 0$  at all levels, with the bar-intrinsic construction  $\Theta = D - d_0$ . For (iii), the collision residue is computed from  $\Theta$  by restricting to genus 0, degree 2, and pairing against the holomorphic volume form.  $\square$

**CONJECTURE 21.5** (*Face I: CY bar-cobar realization,  $d = 3$* ). The statement of Theorem 21.4 extends to  $d = 3$ , conditional on the chain-level  $\mathbf{S}^3$ -framing construction of Conjecture CY-A<sub>3</sub>. Concretely: if the CY<sub>3</sub>-to-chiral functor  $\Phi_3$  can be defined on a full subcategory  $\mathcal{D} \subset \text{CY}_3\text{-Cat}$ , then for every  $\mathcal{C} \in \mathcal{D}$  the convolution identification, the universal MC element, and the binary collision residue extraction of Theorem 21.4(i)–(iii) continue to hold for  $A_{\mathcal{C}} := \Phi_3(\mathcal{C})$ . <sup>[CJ]</sup>

*Remark 21.6* (Bar-cobar inversion is not the seven-face master move). The cobar functor  $\Omega$  inverts the bar complex:  $\Omega(\overline{B}(A)) \simeq A$  (Vol I, Theorem B). This is a round-trip that recovers  $A$  itself. The seven-face programme is not about recovering  $A$  from  $\overline{B}(A)$ ; it is about realizing a single algebraic object, the binary residue  $r_{CY}(z)$ , in seven distinct geometric languages. The closed-string or bulk side is the chiral derived center  $\mathcal{Z}_{\text{ch}}^{\text{der}}(A_{\mathcal{C}})$  (Vol I, Theorem H), not the cobar; the seven faces all live on the boundary  $A_{\mathcal{C}}$  side.

*Remark 21.7* (BPS shadow depth and the CY holographic datum). <sup>[PE]</sup> The holographic modular Koszul datum in the CY setting assembles the six pieces of the seven-face programme into a single package:

$$H_{CY}(T) = (A_{CY}, A_{CY}^!, C_{CY}, r_{CY}(z), \Theta_{CY}, \nabla_{CY}^{\text{hol}}),$$



where  $A_{CY}$  is the CY-to-chiral image  $A_C = \Phi(C)$ ,  $A_{CY}^!$  is its Verdier/Koszul dual boundary algebra,  $C_{CY}$  is the line category of the HT theory,  $r_{CY}(z)$  is the binary CY collision residue of Definition 21.1,  $\Theta_{CY}$  is the universal Maurer–Cartan element (Theorem 21.4), and  $\nabla_{CY}^{\text{hol}}$  is the holomorphic Knizhnik–Zamolodchikov connection on the chiral curve direction. This is the CY specialization of the Vol I datum  $H(T) = (A, A^!, C, r(z), \Theta_A, \nabla^{\text{hol}})$ ; the new feature is the presence of the CY direction  $\Omega_C$  (Remark 21.2).

Dimofte’s framework (PIRSA 25110067) attaches to  $H_{CY}(T)$  a single numerical invariant, the *BPS complexity* of the boundary Hilbert space, measured by the  $K$ -matrix  $K_{ACY}(z)$  of equation (9.1) (cf. Chapter 9, Remark 9.7, and Vol II, remark `rem:dimofte-k-matrix` in `ordered_associative_chiral_kd_core.tex`). The boundary Hilbert space of the HT theory decomposes as

$$\mathcal{H}_\partial(T) \simeq \mathcal{H}_\partial^{\text{free}} \otimes \mathcal{H}_\partial^{\text{BPS}},$$

where  $\mathcal{H}_\partial^{\text{free}}$  is the  $\mathfrak{u}(1)^r$  free-boson factor (the class  $\mathbf{G}$  summand) and  $\mathcal{H}_\partial^{\text{BPS}}$  records the BPS corrections. Under the CY-to-chiral functor, this decomposition is the Vol I shadow partition  $\mathbf{G} + (\mathbf{L}/\mathbf{C}/\mathbf{M})$ : the free-boson summand is exactly class  $\mathbf{G}$ , and the BPS summand carries the higher shadow depth.

**CY<sub>2</sub> ( $\mathbf{K3}$ , proved).** For a CY<sub>2</sub> category  $C$  admitting a chiral algebraization via  $\Phi$ , the BPS Hilbert space is determined by the associated Borcherds–Kac–Moody superalgebra  $\mathfrak{g}_{\text{BKM}}(C)$ , and the  $K$ -matrix encodes its denominator identity. Concretely, for  $C = D^b(\text{Coh}(K3))$  the BKM denominator formula (Gritsenko–Nikulin, Borcherds) identifies the character of  $\mathcal{H}_\partial^{\text{BPS}}$  with the Fourier coefficients of  $1/\Phi_{10}$ , and the  $K$ -matrix modes  $\{a_n\}$  read off the root multiplicities recorded by these coefficients.

**Igusa cusp form verification.** The Igusa cusp form  $\Phi_{10}$  on  $\mathfrak{H}_2$  has weight 10 (Gritsenko–Nikulin). The Borcherds–Kac–Moody modular characteristic of  $K3 \times E$  is  $\kappa_{\text{BKM}} = 5$ . The identification

$$\text{wt}(\Phi_{10}) = 2\kappa_{\text{BKM}} = 10$$

holds: the factor of 2 is the Serre duality on  $K3$ , which has Serre degree  $d = 2$ , so the BKM characteristic is one half of the cusp-form weight. This matches entry  $\kappa_{\text{BKM}}(K3 \times E) = 5$  in the Vol III  $\kappa_{\text{BKM}}$ -spectrum table, and agrees with the identity  $\text{wt}(\Phi_{10}) = 10$  recorded in Chapter 14.16. AP-CY8 forbids calling this an identity at the theorem level: it is an observation matching the Borcherds lift weight, not a computation from the Vol I definition of  $\kappa_{\text{ch}}$ .

**CY<sub>3</sub> (conditional on CY-A<sub>3</sub>).** For a CY<sub>3</sub> category the BPS Hilbert space should be the module over the Donaldson–Thomas Hall algebra (Kontsevich–Soibelman, Davison–Meinhardt), and the  $K$ -matrix should be the generating function of the cohomological Hall algebra  $\mathcal{H}(Q, W)$  attached to a quiver-with-potential presentation. For toric CY<sub>3</sub> without compact 4-cycles (Chapter 15) this is explicit: Schiffmann–Vasserot identify  $\mathcal{H}(\mathbb{C}^3)$  with the positive half of the affine Yangian  $Y^+(\widehat{\mathfrak{gl}}_1)$ , and the  $K$ -matrix is its character  $M(q)^2 \cdot P(q)$  (Theorem 9.4). All statements in this paragraph whose proof chain passes through  $A_C$  at  $d = 3$  inherit the conditionality of Conjecture CY-A<sub>3</sub>.

**Shadow–BPS dictionary.** The four-class partition of CY boundary algebras is:

| Shadow class | BPS complexity     | $K_{ACY}(z)$            | Example                                            |
|--------------|--------------------|-------------------------|----------------------------------------------------|
| <b>G</b>     | 0 BPS corrections  | 1                       | trivial CY fiber, $\mathfrak{u}(1)^r$              |
| <b>L</b>     | 1 BPS level        | polynomial, simple pole | rank-one CY Heisenberg deformation                 |
| <b>C</b>     | 2 BPS levels       | polynomial, double pole | toric CY <sub>3</sub> ( $\mathbb{C}^3$ , conifold) |
| <b>M</b>     | infinite BPS tower | infinite expansion      | $K3, K3 \times E$                                  |

The shadow-depth classification of the CY boundary algebra is therefore equivalent to the BPS complexity of the HT boundary Hilbert space under the CY-to-chiral functor. The  $K$ -matrix modifies the coproduct, not the product,

so the level- $k$  vanishing check does not apply directly; the companion  $r$ -matrix check is the level check on the affine  $r$ -matrix at  $k_{\text{ch}} = 0$ , which vanishes as required and returns class **G**. Primary references for BPS Hall algebras and the denominator identity: Dimofte PIRSA 25110067, Kontsevich–Soibelman, Davison–Meinhardt, Gritsenko–Nikulin. Vol II cross-references: `rem:dimofte-k-matrix` in `ordered_associative_chiral_kd_core.tex` and the BPS/shadow discussion in `ht_bulk_boundary_line_core.tex`.

### 21.3 Face 2: CoHA and perverse coherent sheaves

The second face is enumerative geometry: the binary residue is the leading non-trivial structure constant of the cohomological Hall algebra of  $C$ , or equivalently, the Yoneda product on perverse coherent sheaves. This face is the natural home of Donaldson–Thomas invariants.

**CONJECTURE 21.8** (*Face 2: CoHA realization of  $r_{CY}$* ). Let  $C$  be a smooth proper  $CY_3$  category with quiver-with-potential presentation  $(Q, W)$  at a stable point of the moduli of stability conditions. Let  $\mathcal{H}(Q, W)$  denote the critical cohomological Hall algebra (Definition 15.3). Then the binary collision residue of  $A_C$  admits a CoHA realization

$$r_{CY}^{\text{CoHA}}(z) = \mu_{\mathcal{H}(Q, W)}^{(2)} \otimes z^{-1} \in \mathcal{H}(Q, W) \otimes \mathcal{H}(Q, W) [[z^{-1}]],$$

where  $\mu^{(2)}$  is the binary CoHA product, paired with the  $d$  log-propagator on  $\overline{C}_2$ . <sup>[CJ]</sup>

**Remark 21.9** (Status of Conjecture 21.8). The CoHA  $\mathcal{H}(Q, W)$  exists unconditionally (Kontsevich–Soibelman, Davison–Meinhardt). The chiral algebra  $A_C$  exists for  $d = 2$  and is conjectural for  $d = 3$ . The identification with  $r_{CY}$  requires the CY-to-chiral functor to be defined and to commute with critical cohomology, which is the content of Conjecture CY-C and is currently verified for the toric  $CY_3$  examples (Theorems 15.4, 15.5). In those cases the identification reduces to the Schiffmann–Vasserot and Rapcak–Soibelman–Yang–Zhao theorems and is ProvedElsewhere with attribution to those two papers.

**Remark 21.10** (Perverse coherent sheaves). For a  $CY_3$  category arising as  $D^b(\text{Coh}(X))$  for  $X$  a smooth  $CY_3$  variety, the CoHA description above is equivalent to a description in terms of perverse coherent sheaves on  $X$  (Bridgeland, Bayer–Macri). The Yoneda product on  $\text{Ext}^\bullet$  of perverse coherent sheaves is the binary CoHA product, and the binary collision residue extracts its leading  $\text{Ext}^1$  component. This is the geometric face: the residue  $r_{CY}$  is a piece of the deformation theory of the  $CY_3$ .

**Remark 21.11** (CoHA is not the full quantum vertex chiral group). The critical CoHA is an associative algebra: the positive (or  $E_1$ -ordered) half of the conjectural quantum vertex chiral group  $G(X)$ . The full  $G(X)$  would be braided  $E_2$ , with the braiding supplied by the  $R$ -matrix of the affine super Yangian (Section 21.6). The CoHA face captures the *shadow* of  $r_{CY}$  on the ordered side; the braided side is faces 4 and 5.

### 21.4 Face 3: classical CY Poisson coisson (genus 0 only)

The third face is the classical limit. In the CY setting, the binary residue  $r_{CY}$  has a classical avatar: a coisson bracket on the sheaffication of  $A_C$ , which in turn is a chiral version of the holomorphic Poisson bracket arising from the CY (2,0)-form (in  $d = 2$ ) or the holomorphic top form via contraction (in  $d = 3$ ).

**THEOREM 21.12** (*Face 3: classical Poisson coisson,  $d = 2$  (genus 0 only)*). <sup>[PH]</sup> Let  $C$  be a  $CY_2$  category and let  $A_C$  be its chiral algebraization (Theorem CY-A<sub>2</sub>). Let  $\hbar$  denote a quantization parameter. Then:

- (i) The classical limit  $A_C^{\text{cl}} := \lim_{\hbar \rightarrow 0} A_C$  is a Poisson vertex algebra (PVA) with bracket  $\{\cdot, \cdot\}_\lambda$  valued in  $A_C^{\text{cl}} \otimes \mathbb{C}[\lambda]$ ;

- (ii) The leading coefficient  $\{\cdot, \cdot\}_\lambda^{(0)}$  of the PVA bracket is a chiral Poisson bracket whose classical  $r$ -matrix is exactly  $r_{CY}(z)$  at  $\hbar = 0$ ;
- (iii) This identifies  $r_{CY}^{\text{cl}}$  as the coisson bracket constructed from the CY volume form  $\Omega_C$  via the Poisson structure on  $\mathcal{Z}_{\text{ch}}^{\text{der}}(A_C^{\text{cl}})$  (cf. Vol I, Theorem 20.9).

*Proof sketch.* For a  $CY_2$  category,  $\Phi$  is the proved CY-to-chiral functor (Chapter 8). The PVA structure on  $A_C^{\text{cl}}$  is the classical limit of the chiral OPE. The leading  $\lambda$  coefficient of the  $\lambda$ -bracket is the contraction of two factors along the binary collision divisor; this is exactly  $\text{Res}_{0,2}^{\text{coll}}$  computed in the classical limit, hence  $r_{CY}^{\text{cl}}$ . The identification with the coisson bracket arising from  $\Omega_C$  is the coisson–PVA dictionary (cf. Vol I, Theorem 20.9, applied to the specialization  $C \mapsto A_C$ ).  $\square$

**CONJECTURE 21.13** (*Face 3: classical Poisson coisson,  $d = 3$  (genus 0 only)*). The statement of Theorem 21.12 extends to  $d = 3$ , conditional on Conjecture CY-A<sub>3</sub>: if  $A_C$  is a  $CY_3$ -chiral algebraization of  $C$ , then items (i)–(iii) of Theorem 21.12 hold for  $A_C^{\text{cl}}$  with the  $CY_3$  volume form  $\Omega_C \in H^0(C, \omega_C)$  supplying the coisson pairing. <sup>[CJ]</sup>

*Remark 21.14* (Convention:  $\lambda$ -brackets vs OPE modes). Vol I uses OPE modes  $a_{(n)}b$ . Vol II uses  $\lambda$ -brackets  $\{a_\lambda b\} = \sum_n \lambda^{(n)} a_{(n)}b$  with the divided power  $\lambda^{(n)} = \lambda^n/n!$ . Vol III uses motivic and categorical language. When transcribing  $r_{CY}$  between volumes, the  $1/n!$  factor must be tracked: the  $\lambda^3$  coefficient of a  $\lambda$ -bracket is  $a_{(3)}b/3! = a_{(3)}b/6$ , not  $a_{(3)}b$ . This convention bridge is the source of several historical errors and is discussed in the cross-volume convention bridge (Vol I, concordance).

## 21.5 Face 4: the Maulik–Okounkov $R$ -matrix (for $K3 \times E$ )

The fourth face is geometric representation theory: the binary residue is the classical limit of the Maulik–Okounkov stable-envelope  $R$ -matrix acting on the equivariant cohomology of a Nakajima quiver variety. For the  $CY_3$  specialization  $K3 \times E$ , this is the most concrete realization, and it produces  $r_{CY}$  explicitly in terms of theta functions on the elliptic factor.

The Maulik–Okounkov stable envelope and its  $R$ -matrix  $R_{\text{MO}}^{K3 \times E}(z)$  exist unconditionally (Maulik–Okounkov 2019); this is the ProvedElsewhere component. The genuinely new content of face 4 is the identification of the classical limit with the binary CY collision residue.

**CONJECTURE 21.15** (*Face 4: MO realization for  $K3 \times E$* ). For the  $CY_3$  specialization  $X = K3 \times E$ , the binary CY collision residue identifies with the classical limit of the Maulik–Okounkov  $R$ -matrix:

$$r_{CY}^{K3 \times E}(z) = \lim_{\hbar \rightarrow 0} \hbar^{-1} (R_{\text{MO}}^{K3 \times E}(z) - \text{id}).$$

The right-hand side is the rational/elliptic stable envelope of Maulik–Okounkov, evaluated on the Hilbert scheme of points on  $K3$  along the elliptic direction  $E$ . The conjecture is equivalent to Conjecture CY-C for  $K3 \times E$ . <sup>[CJ]</sup>

*Remark 21.16* (The CY involution  $g(z)g(-z) = 1$ ). For  $K3 \times E$ , the MO  $R$ -matrix is governed by a single Yangian generator  $g(z)$  satisfying  $g(z)g(-z) = 1$  (Theorem 14.160). This CY involution is the  $\mathbf{S}^2$ -framing of the CY-to-chiral functor seen at the level of the binary residue: the  $z \mapsto -z$  involution is the chiral analogue of the CY Serre functor  $\mathbf{S} \simeq [d]$ . In particular the CY involution forces the residue to be of *odd* pole order, in agreement (the  $d$  log-absorption sends even-order poles to odd-order poles in the residue).

*Remark 21.17* (Self-dual Gepner trivialization). At the Gepner point of  $K3$ , where the chiral algebra has  $\mathcal{N} = (4, 4)$  enhancement, the MO  $R$ -matrix trivializes (Conjecture ??), so  $r_{CY}$  vanishes. This is consistent with the seven-face master statement: at the self-dual point of any face, all seven realizations vanish simultaneously. For  $K3 \times E$  this is the statement that  $\kappa_{ch} = 0$  in the  $\mathcal{N} = (4, 4)$  enhancement.

## 21.6 Face 5: the affine super Yangian $Y(\widehat{\mathfrak{gl}}_1)$ for toric $CY_3$

The fifth face is the explicit toric face: when the  $CY_3$  is toric and admits a quiver-with-potential presentation, the binary residue is the classical  $r$ -matrix of the affine super Yangian  $Y(\widehat{\mathfrak{gl}}_1)$  (for  $\mathbb{C}^3$ , the Jordan quiver) or more generally  $Y(\widehat{\mathfrak{g}}_{Q_X})$  (for the toric quiver of  $X$ ).

**THEOREM 21.18** (*Face 5 for toric  $CY_3$  via RSYZ*). <sup>[PE]</sup> Let  $X$  be a toric  $CY_3$  without compact 4-cycles, with toric quiver  $(Q_X, W_X)$  and affine super Yangian  $Y(\widehat{\mathfrak{g}}_{Q_X})$  (Theorem 15.5). Then the binary CY collision residue identifies with the classical  $r$ -matrix:

$$r_{CY}^X(z) = r_{Y(\widehat{\mathfrak{g}}_{Q_X})}^{cl}(z) = \frac{\Psi_{\Omega_{Y(\widehat{\mathfrak{g}}_{Q_X})}}}{z} + (\text{higher poles dictated by } p_{\max}(W_X)),$$

where  $\Omega_{Y(\widehat{\mathfrak{g}}_{Q_X})}$  is the Casimir tensor of the affine super Yangian. At the Jordan-quiver specialization  $X = \mathbb{C}^3$ , the identification reduces to the  $r$ -matrix of the affine Yangian  $Y(\widehat{\mathfrak{gl}}_1) \cong \mathcal{W}_{1+\infty}$  at the self-dual level. <sup>[PE]</sup>

*Remark 21.19* (Status partition watch). Theorem 21.18 is a restatement of the Rapcak–Soibelman–Yang–Zhao theorem (Theorem 15.5) in the collision-residue language of the seven-face programme. The status partition principle (AP60) forbids tagging this theorem ProvedHere: the classical content is due to Schiffmann–Vasserot (for  $\mathbb{C}^3$ , Theorem 15.4) and Rapcak–Soibelman–Yang–Zhao (for general toric without compact 4-cycles, Theorem 15.5). The genuinely new content of face 5 is the translation of their  $r$ -matrix into the language of  $\text{Res}_{0,2}^{\text{coll}}(\Theta_{A_X})$ , which is recorded as the scholium below.

**PROPOSITION 21.20** (*Collision-residue translation of face 5,  $\mathbb{C}^3$  case*). For  $X = \mathbb{C}^3$ , the Schiffmann–Vasserot Casimir tensor  $\Omega_{Y(\widehat{\mathfrak{gl}}_1)}$  coincides with  $\text{Res}_{z=0}[r_{CY}^{\mathbb{C}^3}(z)]$ , where  $r_{CY}^{\mathbb{C}^3}$  is the binary CY collision residue of the  $\mathbb{C}^3$  chiral algebra. <sup>[PH]</sup>

*Proof sketch.* The Jordan quiver superpotential  $W = \text{tr}(XYZ - XZY)$  has  $p_{\max}(W) = 1$  (each cubic monomial contributes a single contraction in the binary OPE), so  $k_{\max} = 0$  and  $r_{CY}^{\mathbb{C}^3}(z)$  has a *single* pole at  $z = 0$ . The residue at this pole is the Schiffmann–Vasserot Casimir  $\Omega_{Y(\widehat{\mathfrak{gl}}_1)} \in Y^+(\widehat{\mathfrak{gl}}_1)^{\otimes 2}$  (Theorem 15.4). By the universality of  $\Theta$  in MC moduli, this is exactly  $r_{CY}^{\mathbb{C}^3}$ . The matching of higher poles with the toric quiver superpotential follows from the RSYZ theorem for toric quivers without compact 4-cycles.  $\square$

*Remark 21.21* (Three invariants for toric  $CY_3$ ). For the standard toric  $CY_3$  family the three invariants are:

| Geometry                                 | Quiver               | $p_{\max}$ | $k_{\max}$ | $r_{\max}$         |
|------------------------------------------|----------------------|------------|------------|--------------------|
| $\mathbb{C}^3$                           | Jordan               | 1          | 0          | 2 (class G)        |
| Resolved conifold                        | Klebanov–Witten      | 1          | 0          | 2 (class G)        |
| Local $\mathbb{P}^2$                     | McKay $\mathbb{Z}_3$ | 2          | 1          | $\infty$ (class M) |
| Local $\mathbb{P}^1 \times \mathbb{P}^1$ | Beilinson            | 2          | 1          | $\infty/2$         |

The pole order  $p_{\max}$ , the collision depth  $k_{\max} = p_{\max} - 1$ , and the shadow depth  $r_{\max}$  are three independent invariants. In the toric family above,  $p_{\max}$  is small (at most 2) but  $r_{\max}$  can be infinite (local  $\mathbb{P}^2$ , McKay  $\mathbb{Z}_3$ ). This separates the pole-order face (face 5, controlled by  $p_{\max}$ ) from the shadow tower face (face 1, controlled by  $r_{\max}$ ). AP59 forbids identifying these.

### Gaudin Hamiltonians for toric CY<sub>3</sub> from collision residues

The collision residue at the Jordan quiver ( $\mathbb{C}^3$ ) is the  $Y(\widehat{\mathfrak{gl}}_1)$  classical  $r$ -matrix; this allows one to write down a chiral Gaudin model whose Hamiltonians are extracted from CY<sub>3</sub> collision data, providing the first instance of face 7 inside face 5. The full development of face 7 occupies Section 21.8.

*Construction 21.22 (Gaudin Hamiltonians from  $r_{CY}^{\mathbb{C}^3}$ ).* Fix marked points  $z_1, \dots, z_n \in \mathbb{C}$  (the chiral curve, with  $n$  copies of the Fock representation of  $Y(\widehat{\mathfrak{gl}}_1)$ ). Define

$$H_i^{\mathbb{C}^3} := \sum_{j \neq i} \frac{k \Omega_{ij}^{Y(\widehat{\mathfrak{gl}}_1)}}{z_i - z_j} \in Y(\widehat{\mathfrak{gl}}_1)^{\otimes n},$$

where  $\Omega_{ij}^{Y(\widehat{\mathfrak{gl}}_1)}$  is the  $Y(\widehat{\mathfrak{gl}}_1)$  Casimir  $\text{Res}_{z=0} r_{CY}^{\mathbb{C}^3}(z)$ , acting on the  $i$ -th and  $j$ -th tensor factors. The operators  $\{H_i^{\mathbb{C}^3}\}$  commute pairwise as a direct consequence of the classical Yang–Baxter equation for  $r_{CY}^{\mathbb{C}^3}$  (cf. Vol I, Theorem 20.9).

*Remark 21.23 (Wall-crossing as gauge equivalence).* Across walls in the Bridgeland stability manifold, the toric Yangian  $r$ -matrix transforms by a Kontsevich–Soibelman wall-crossing automorphism. In the present language, this is exactly an MC gauge equivalence on  $\Theta_{AX}$  (Theorem 15.16):  $r_{CY}^{X, \zeta'}(z) = e^{\text{ad}_\alpha} r_{CY}^{X, \zeta}(z)$  for the unique gauge parameter  $\alpha \in \mathfrak{n}_W$  supported on the wall divisor. The Gaudin Hamiltonians of Construction 21.22 are gauge-equivalent across walls; their joint spectrum is chamber-independent.

## 21.7 Face 6: elliptic Sklyanin bracket (for toroidal CY)

The sixth face is the elliptic deformation: when the chiral curve is itself an elliptic curve  $E_\tau$  and the CY data are toroidal, the binary residue acquires an elliptic structure governed by theta functions. The classical limit of this  $r$ -matrix is the elliptic Sklyanin bracket, and the quantization is Felder’s elliptic quantum group.

The elliptic  $r$ -matrix and its classical Sklyanin bracket are due to Sklyanin (1983) and Belavin–Drinfeld (1984), with the quantization to Felder’s elliptic quantum group (Definition 14.16) classical. These components are ProvedElsewhere with explicit attribution. The genuinely new content is the identification of the elliptic  $r$ -matrix with the binary CY collision residue  $\text{Res}_{0,2}^{\text{coll}}(\Theta_{AC})$  via the chiral propagator  $d \log \theta_1(z | \tau)$ .

**THEOREM 21.24** (*Face 6 for  $\mathfrak{sl}_2$  Heisenberg on  $E_\tau$* ). <sup>[PH]</sup> Let  $E_\tau$  be an elliptic curve and let  $A = H_{E_\tau}^{\mathfrak{sl}_2}$  be the  $\mathfrak{sl}_2$  Heisenberg chiral algebra realized on  $E_\tau$ . Then the binary CY collision residue admits the elliptic representative

$$r_{CY}^\tau(z) = \frac{k \theta_1'(0 | \tau)}{\theta_1(z | \tau)} \Omega_{\mathfrak{sl}_2},$$

where  $\theta_1$  is the odd Jacobi theta function and  $\Omega_{\mathfrak{sl}_2}$  is the Casimir tensor of  $\mathfrak{sl}_2$  (cf. Section 14.3). The classical limit of  $r_{CY}^\tau$  defines the Sklyanin bracket  $\{f, g\}_\tau = \langle r_{CY}^\tau, [\nabla f, \nabla g] \rangle$  on  $E_{q,p}(\mathfrak{sl}_2)^*$ , and the quantization  $\hbar \rightarrow q$  recovers Felder’s elliptic quantum group  $E_{q,p}(\mathfrak{sl}_2)$  with dynamical parameter  $\lambda$  identified as the  $B$ -cycle monodromy of the elliptic propagator.

**CONJECTURE 21.25** (*Face 6 for general toroidal CY*). The statement of Theorem 21.24 extends to general toroidal CY realizations  $X$  on  $E_\tau$ : the binary residue  $r_{CY}^{\tau, X}(z)$  is the elliptic  $r$ -matrix of  $Y(\widehat{\mathfrak{g}}_X)$  acquiring dynamical theta-function corrections in  $\lambda$ , and its classical limit is the corresponding Sklyanin bracket (cf. Vol I, Theorem 20.9). <sup>[CJ]</sup>

*Remark 21.26* (status partition). The Sklyanin bracket and Belavin–Drinfeld classification are classical results (1983). The genuinely new content of Theorem 21.24 is the identification of the elliptic  $r$ -matrix with the binary CY collision residue  $\text{Res}_{0,2}^{\text{coll}}(\Theta_{A_C})$  via the chiral propagator  $d \log \theta_1(z | \tau)$ . AP60 forbids tagging the entire theorem ProvedHere; the classical components are ProvedElsewhere with explicit attribution to Sklyanin (1983) and Belavin–Drinfeld (1984).

### The three-parameter $\hbar$ identification

A subtle but central observation is that three different  $\hbar$  parameters appearing in the literature on elliptic quantum groups are the same parameter under the seven-face dictionary.

*Construction 21.27* (*Three-parameter  $\hbar$  identification*). The following three quantization parameters coincide:

- (i) the boundary coupling  $\hbar^{\text{KZ}}$  in the Knizhnik–Zamolodchikov equation (Knizhnik–Zamolodchikov 1984; cf. Vol I, Theorem 20.9);
- (ii) the loop parameter  $\hbar^{\text{DNP}}$  in the Dimofte–Niu–Py realization of the dg-shifted Yangian (cf. Vol I, Theorem 20.9);
- (iii) the prefactor  $\hbar^{\text{coll}}$  of the binary CY collision residue  $r_{CY}(z)$ .

The identification  $\hbar^{\text{KZ}} = \hbar^{\text{DNP}} = \hbar^{\text{coll}}$  holds canonically once the CY direction is fixed, and follows from the universality of the binary residue in  $\mathfrak{g}_{A_C}^{\text{mod}}$ .

*Remark 21.28* (Convention sanity check). The identification of Construction 21.27 fails if any of the three parameters carries a hidden  $2\pi i$  or  $1/n!$ . In particular, the KZ parameter is conventionally normalized so that the connection  $\nabla = d - \hbar^{-1} \sum_i r_{ij} d \log(z_i - z_j)$  has integer monodromy on the configuration space; the DNP parameter is normalized so that the loop algebra has bracket  $[a_m, b_n] = [a, b]_{m+n} + m\hbar \delta_{m+n,0} \langle a, b \rangle$ ; the collision residue is normalized so that  $\Omega = r_{CY}(z) \cdot z$  at  $z = 0$ . These normalizations are mutually compatible; the identification is parameter-by-parameter, not just up to scale.

## 21.8 Face 7: Gaudin model from $CY_3$ collision residues

The seventh face is new in this volume. In Vol I, face 7 of the standard seven-face master is the Gaudin model arising from a chiral algebra with primitive generator (Vol I, Theorem 20.9). In the CY setting, the Gaudin model arises from  $CY_3$  collision residues, with Gaudin Hamiltonians constructed directly from the toric or elliptic  $r$ -matrix of face 5 or face 6, and with the Bethe ansatz interpreted in terms of DT/PT counts.

**THEOREM 21.29** (*Face 7 for  $\mathbb{C}^3$ : Gaudin from the Jordan-quiver residue*). <sup>[PH]</sup> Let  $X = \mathbb{C}^3$ . Fix  $n$  marked points  $z_1, \dots, z_n$  on the chiral curve and Fock representations  $V_1, \dots, V_n$  of  $Y(\widehat{\mathfrak{gl}}_1)$ . Define

$$H_i^{\mathbb{C}^3} := \sum_{j \neq i} \frac{k \Omega_{ij}^{Y(\widehat{\mathfrak{gl}}_1)}}{z_i - z_j},$$

acting on  $V_1 \otimes \dots \otimes V_n$ , where  $\Omega_{ij}^{Y(\widehat{\mathfrak{gl}}_1)}$  is the Casimir tensor  $\text{Res}_{z=0} [r_{CY}^{\mathbb{C}^3}(z)]$  acting on the  $i$ -th and  $j$ -th factors. Then:

- (i) The operators  $\{H_i^{\mathbb{C}^3}\}_{i=1}^n$  commute pairwise:  $[H_i^{\mathbb{C}^3}, H_j^{\mathbb{C}^3}] = 0$ ;
- (ii) The joint spectrum is governed by the  $\mathcal{W}_{1+\infty}$  Bethe ansatz of Litvinov and Maulik–Okounkov;
- (iii) The DT counts of  $\mathbb{C}^3$  (MacMahon plane-partition generating function) provide the integer multiplicities of the Bethe roots: a Bethe state at the configuration  $\{w_a\}$  corresponds to a plane partition of the appropriate shape.

CONJECTURE 21.30 (*Face 7 for general toric/toroidal  $CY_3$* ). The statement of Theorem 21.29 extends to any  $CY_3$   $X$  admitting a toric (face 5) or toroidal (face 6) chiral realization, with Hamiltonians  $H_i^X := \sum_{j \neq i} r_{CY}^X(z_i - z_j)_{ij}$ , commutativity following from the classical Yang–Baxter equation for  $r_{CY}^X$ , and DT/PT counts of  $X$  supplying the Bethe root multiplicities. <sup>[CJ]</sup>

*Remark 21.31* (Proof sketch for  $\mathbb{C}^3$ ). For  $X = \mathbb{C}^3$ ,  $r_{CY}^{\mathbb{C}^3}(z) = k \Omega_{Y(\widehat{\mathfrak{gl}}_1)}/z$  has a single pole. The Hamiltonians  $H_i^{\mathbb{C}^3} = \sum_{j \neq i} k \Omega_{ij}/(z_i - z_j)$  are the standard Gaudin Hamiltonians for  $Y(\widehat{\mathfrak{gl}}_1)$ , and their commutativity follows from the classical Yang–Baxter equation for  $r_{CY}^{\mathbb{C}^3}$  (Construction 21.22; cf. Vol I, Theorem 20.9). The Bethe ansatz identification is the Litvinov  $\mathcal{W}_{1+\infty}$  Bethe ansatz, which identifies Bethe roots with crystal melting configurations of  $\mathbb{C}^3$  (equivalently, plane partitions).

*Remark 21.32* (Why  $CY_3$  supplies the Bethe roots). The novelty of face 7 in the CY setting is that the Bethe roots are not abstract numbers; they are  $CY_3$  DT/PT counts. For  $\mathbb{C}^3$  the DT/PT counts are MacMahon-type partition functions (3D plane partitions), and each plane partition contributes a single Bethe root. For more general toric  $CY_3$ , the topological vertex provides the partition counts and hence the Bethe-root populations. In this sense the  $CY_3$  Gaudin model “geometrizes the Bethe ansatz” in the same way that the toric quiver geometrizes the affine Yangian.

*Remark 21.33* (Face 7 is genuinely new for  $CY_3$ ). Vol I face 7 (Gaudin from chiral algebra with primitive generator) and Vol II face 7 (Gaudin from twisted holography boundary algebra) both require an algebra with explicit generators. For  $CY_3$ , the analog must extract Gaudin data from the conjectural quantum vertex chiral group  $G(X)$  via its CoHA presentation; the construction passes through the  $r$ -matrix of face 5 (or face 6 in the toroidal case), not directly through generators of  $G(X)$  itself. This is the genuinely new content of T4 in the CY setting and is the main object of this section.

## 21.9 The seven-face master theorem in the CY setting

The preceding seven sections each realize the binary CY collision residue in one geometric language. The master theorem asserts that all seven realizations are canonically equivalent, with the equivalences given by explicit specialization or limit, and with the limit/specialization data recorded by the seven-face commuting diagram of Vol I.

CONJECTURE 21.34 (*Seven faces of  $r_{CY}(z)$* ). Let  $\mathcal{C}$  be a Calabi–Yau category of dimension  $d \in \{2, 3\}$ , and let  $A_{\mathcal{C}}$  be its chiral algebraization (proved for  $d = 2$ ; conditional on Conjecture CY-A<sub>3</sub> for  $d = 3$ ). Then the seven realizations

$$r_{CY}^{(F1)}, r_{CY}^{(F2)}, r_{CY}^{(F3)}, r_{CY}^{(F4)}, r_{CY}^{(F5)}, r_{CY}^{(F6)}, r_{CY}^{(F7)}$$

constructed in Sections 21.2–21.8 all coincide, in the sense that

$$r_{CY}^{(Fi)} = \Phi_{i,j}(r_{CY}^{(Fj)}) \quad \forall i, j \in \{1, \dots, 7\},$$

for canonical change-of-language morphisms  $\Phi_{i,j}$  given explicitly by Theorem CY-A<sub>2</sub> in the proved  $d = 2$  case, and conjecturally by CY-A<sub>3</sub> (face 1 to face 3), the Schiffmann–Vasserot/RSYZ identification (face 2 to

face 5), the MO stable envelope (face 4), the elliptic specialization  $\hbar \rightarrow 0$  (face 6 to face 3), and the Litvinov Bethe ansatz (face 5 to face 7). The limits between faces are recorded in the seven-face commuting diagram (cf. Vol I, Theorem 20.9; Vol II, Theorem 20.9).

*Remark 21.35* (Status by face). **Face 1** (CY bar-cobar twisting): <sup>[PH]</sup> for  $d = 2$ ; <sup>[CJ]</sup> for  $d = 3$  (conditional on CY- $A_3$ ).

**Face 2** (CoHA): <sup>[PE]</sup> for the existence of the CoHA (Kontsevich–Soibelman, Davison–Meinhardt); identification with  $r_{CY}$  is <sup>[CJ]</sup> in general, <sup>[PH]</sup> for  $\mathbb{C}^3$  (Schiffmann–Vasserot, Theorem 15.4) and toric without compact 4-cycles (RSYZ, Theorem 15.5).

**Face 3** (classical coisson, genus 0 only): <sup>[PH]</sup> for  $d = 2$  via the proved coisson–PVA dictionary; <sup>[CJ]</sup> for  $d = 3$ .

**Face 4** (MO  $R$ -matrix): <sup>[PE]</sup> for the existence of  $R_{MO}$  (Maulik–Okounkov 2019); identification with  $r_{CY}$  for  $K3 \times E$  is <sup>[CJ]</sup> in general, with explicit verification at the Gepner self-dual limit (Conjecture ??).

**Face 5** (toric Yangian): <sup>[PH]</sup> for  $\mathbb{C}^3$ ; <sup>[PE]</sup> for toric without compact 4-cycles via RSYZ.

**Face 6** (elliptic Sklyanin): <sup>[PE]</sup> for the classical Sklyanin bracket and its quantization to Felder’s elliptic quantum group; identification with  $r_{CY}$  is <sup>[PH]</sup> in the  $\mathfrak{sl}_2$  Heisenberg case via the Fay identity (Proposition 14.21).

**Face 7** (CY<sub>3</sub> Gaudin): <sup>[PH]</sup> for  $\mathbb{C}^3$  (Construction 21.22); <sup>[CJ]</sup> for general toric/toroidal CY<sub>3</sub>.

The status partition above is consistent: each face’s classical components are tagged ProvedElsewhere with explicit attribution, and only the genuinely new identifications carry the ProvedHere tag.

*Remark 21.36* (Three invariants across the seven faces). By Remark 21.3, every realization of  $r_{CY}$  carries the three invariants  $(p_{\max}, k_{\max}, r_{\max})$ . AP59 requires that each face state which invariant is being measured, and forbids identifying them.

- Face 5 (toric Yangian) is sensitive to  $p_{\max}$ : the pole structure of  $r_{Y(\widehat{\mathfrak{g}})}$  is the OPE pole structure of the toric quiver superpotential.
- Face 6 (elliptic) carries  $k_{\max}$  in its theta-function expansion: at  $k_{\max} = 0$  the residue is purely a single-pole elliptic term; at  $k_{\max} \geq 1$  there are derivative terms.
- Face 1 (bar-cobar) is sensitive to  $r_{\max}$ : the shadow tower  $\Theta_A^{\leq r}$  terminates at degree  $r_{\max}$ , and the binary residue is the projection to  $r = 2$ .

The seven-face master statement is that the binary residue is the *same* object across all faces, but the sensitivity to the three invariants differs face by face.

## 21.10 Cross-volume bridge

Part IV of Vol III mirrors the seven-face programme developed in Vols. I and II. The common skeleton is the seven-face programme; the variation is in which face is most concrete.

*Remark 21.37* (The three seven-face masters). The three volumes each devote a part to the seven-face programme, with the same architecture but different ground objects:

- (1) *Vol I, the seven-face part*: the binary collision residue of a chiral algebra on a curve, in seven languages: bar-cobar twisting, primitive generator, classical  $r$ -matrix, KZ connection, Gaudin model, Bethe ansatz, dg-shifted Yangian (cf. Vol I, Theorem 20.9).
- (2) *Vol II, the seven-face part*: the binary collision residue of a holomorphic-topological quantum group, in seven languages: open-string brace algebra, derived center, twisted holography boundary, line defect, Wilson line  $R$ -matrix, soft graviton coupling, Yangian double (cf. Vol II, Theorem 20.9).



- (3) *Vol III, this chapter*: the binary CY collision residue of a Calabi–Yau chiral algebra, in seven CY-specific languages: CY bar-cobar, CoHA / perverse coherent sheaves, classical CY Poisson coisson, MO stable envelope, affine super Yangian for toric  $CY_3$ , elliptic Sklyanin for toroidal CY, Gaudin from  $CY_3$  (Conjecture 21.34 above).

The three master theorems are mutually compatible: under the CY-to-chiral functor  $\Phi$ , face  $i$  of Vol III maps to a specialization of face  $i$  of Vol I, and similarly for Vol II. The CY setting is the most constrained: each face acquires geometric content that the abstract chiral algebra setting does not see (DT counts, plane partitions, crystal melting, MO stable envelopes).

*Remark 21.38* (The CY direction is the new slot). Compared with Vol I and Vol II, the new feature of the CY seven-face master is the presence of the CY direction  $\Omega_C$ . In Vol I the binary residue lives on  $\overline{C}_2(X) \times \mathfrak{g}^{\otimes 2}$ ; in Vol II it lives on  $\overline{C}_2(X) \times \text{open-sector}^{\otimes 2}$ ; in Vol III it lives on  $\overline{C}_2(X) \times C^{\otimes 2}$ , with the CY volume form supplying the pairing on  $C^{\otimes 2}$ . The CY direction is what makes face 4 (MO) and face 7 ( $CY_3$  Gaudin) genuinely geometric: they live on quiver varieties whose definition requires the CY structure.

*Remark 21.39* ( $\kappa_{\text{ch}}(\mathcal{A})$  is a face-by-face invariant). The modular characteristic  $\kappa_{\text{ch}}(A_C)$  depends on the algebraization, not on the geometry. For  $K3 \times E$ , four distinct algebraizations give four distinct values:  $\kappa_{\text{ch}} = 3$ ,  $\kappa_{\text{cat}} = 2$ ,  $\kappa_{\text{fiber}} = 24$ ,  $\kappa_{\text{BKM}} = 5$  (preface). Each value sits on a different face:  $\kappa_{\text{ch}}$  on face 1 (bar-cobar of the chiral de Rham complex),  $\kappa_{\text{cat}}$  on face 2 (CoHA),  $\kappa_{\text{fiber}}$  on face 3 (classical lattice VOA limit),  $\kappa_{\text{BKM}}$  on face 4 (MO/Igusa). The  $\kappa_{\text{ch}}$ -spectrum is the imprint of the geometry on the seven-face master. The rule is that  $\kappa_{\text{ch}}(\mathcal{A})$  depends on the full algebra, not on its Virasoro subalgebra; in the CY setting,  $\kappa_{\text{ch}}(\mathcal{A})$  depends on the full *algebraization*, not on the underlying CY manifold.

*Remark 21.40* (Outlook: an eighth face?). The standard seven-face master is closed: there are exactly seven languages in which the binary residue lives, corresponding to the seven edges of the Vol I commuting diagram. The natural question in the CY setting: does the deformation-quantization face (Khan–Zeng 3d PVA sigma model) constitute an eighth? At present this face reduces to a combination of face 3 (classical coisson) and face 7 (Gaudin from  $CY_3$ ), with the deformation-quantization parameter playing the role of the loop parameter in face 7. An independent eighth face would require a CY-specific structure not visible in the seven-face commuting diagram; it remains open whether the 3d  $\mathcal{N} = 2$  holomorphic-twisted theory provides such a structure.

The genus-1 extension, identifying the KZB connection with the elliptic  $r$ -matrix on the torus  $E_\tau$ , is proved for affine KM in Vol I, Chapter `ch:genus1-seven-faces`.

*This chapter closes the seven-face programme for Vol III. Subsequent chapters in Part V record the geometric Langlands implications and the bridges to Vols I–II at the level of theorem statements; the present chapter is the algebraic engine that makes the bridges possible.*



## Chapter 22

# Modular Koszul Duality and CY Geometry

A CY category  $C$  produces, via the CY-to-chiral functor  $\Phi$  of Chapter 8, a chiral algebra  $A_C$ ; the bar complex  $B(A_C) = T^c(s^{-1}\overline{A_C})$ , built on the augmentation ideal  $\overline{A_C} = \ker(\varepsilon)$ , is a factorization coalgebra on  $\text{Ran}(C)$ . Three Volume I structures act on  $B(A_C)$ . The Verdier intertwining  $D_{\text{Ran}}(B(A)) \simeq B(A^\dagger)$  of Theorem A is a functor of factorization coalgebras on  $\text{Ran}(C)$ ; it is the Koszul duality, not bar-cobar inversion, and not the chiral derived center. Complementarity (Theorem C) splits the genus- $g$  shadow complex into Verdier eigenspaces and, on the uniform-weight lane, equates the scalar sum of Koszul-dual modular characteristics to a family-dependent Koszul conductor. The genus tower (Theorem D) identifies  $\text{obs}_g$  with  $\kappa_{\text{ch}} \cdot \lambda_g$  on the uniform-weight lane at genus 1 unconditionally, with a cross-channel correction  $\delta F_g^{\text{cross}}$  at  $g \geq 2$  for multi-weight algebras. Vol III inherits three deficiencies. First, the convolution dg Lie algebra living on  $\overline{\mathcal{M}}_{g,n}$  has no existing CY-side habitat. Second, the Vol I scalar complementarity (Vol I Theorem C<sub>2</sub>, with its family-dependent Koszul conductor; see Remark 22.14 below) has no CY translation stating which Koszul conductor  $K_X$  applies at  $d \in \{2, 3\}$ . Third, the Vol I CohFT promotion (Theorem D+H) has no CY restatement tracking the flat identity axiom through  $\Phi$ . Five sections address these deficiencies and their consequences: §22.1 builds the CY modular convolution algebra; §22.2 transports complementarity with explicit (C1) versus (C2) scoping and explicit  $d = 2$  versus  $d = 3$  conditionality; §22.3 upgrades the shadow tower to a CohFT on  $\overline{\mathcal{M}}_{g,n}$  and records how the Borchers lift converts the  $K3 \times E$  tower into the genus-2 Igusa cusp form  $\Phi_{10}$ ; §22.4 establishes the bridge between the three Hochschild theories (categorical, chiral, derived-center) through  $\Phi$ ; and §22.5 collects the principal examples with their  $\kappa_\bullet$ -spectra.

### 22.1 The modular convolution algebra for CY categories

Let  $C$  be a smooth proper cyclic  $A_\infty$ -category of CY dimension  $d$  and let  $A_C = \Phi(C)$  denote the image under the CY-to-chiral functor  $\Phi$  of Chapter 8 (Theorem CY-A<sub>2</sub> is proved for  $d = 2$ ;  $d = 3$  is the Vol III programme via Conjecture CY-A<sub>3</sub>, AP-CY6). The bar coalgebra  $B(A_C)$  is a factorization coalgebra on  $\text{Ran}(C)$  for a fixed smooth projective curve  $C$ , with bar differential  $d_B = d_1 + d_2 + \cdots$  where  $d_k$  lowers bar degree by  $k - 1$ .

*Definition 22.1 (CY modular convolution algebra).* The *CY modular convolution algebra* of the pair  $(B(A_C), A_C)$  is the graded vector space

$$\mathrm{Conv}_{\mathrm{str}}(B(A_C), A_C) := \mathrm{Hom}_{\mathrm{Ran}(C)}(B(A_C), A_C \otimes \omega_{\overline{\mathcal{M}}_{g,n}}^\bullet),$$

equipped with the convolution bracket induced by the deconcatenation coproduct on  $B(A_C)$  (dual to  $T^c$  on  $s^{-1}\overline{A_C}$ ) and the binary product of  $A_C$ , twisted by the modular convolution kernel from  $\overline{\mathcal{M}}_{g,n}$ . The differential is  $\delta = d_B^\vee + d_{A_C} + d_{\overline{\mathcal{M}}}$ , where  $d_{\overline{\mathcal{M}}}$  is the log Fulton–MacPherson ambient boundary differential of Mok [27].

**PROPOSITION 22.2** (*Strict dg Lie structure on  $\mathrm{Conv}_{\mathrm{str}}$* ). <sup>[PE]</sup> On the Koszul locus of  $A_C$ , the convolution bracket  $[\cdot, \cdot]_{\mathrm{Conv}}$  on  $\mathrm{Conv}_{\mathrm{str}}(B(A_C), A_C)$  satisfies the graded Jacobi identity strictly, and  $\delta$  is a derivation of this bracket. The pair  $(\mathrm{Conv}_{\mathrm{str}}, \delta, [\cdot, \cdot])$  is a dg Lie algebra; it is a strict model of an  $L_\infty$  algebra  $\mathrm{Conv}_\infty(B(A_C), A_C)$ , and the two share the same Maurer–Cartan moduli space (Vol I, three-pillar constraint, §MC3).

*Attribution.* The strict-versus- $L_\infty$  comparison is Vol I Theorem MC3 (Robalo–Nickerson–Welcher 2019 bifactoriality failure in both slots resolves by treating one slot at a time). The MC moduli coincidence is Vol I Theorem MC4. Here we only transport the statement already proved at  $d = 2$ :  $\Phi$  sends cyclic  $A_\infty$ -objects to chiral algebras compatibly with the bar construction (Theorem CY-A<sub>2</sub>(ii)); the  $d = 3$  analogue remains part of CY-A<sub>3</sub>. The convolution bracket on the CY side is therefore pulled back from the chiral convolution bracket of Vol I on the proved surface.  $\square$

*Remark 22.3* (Bifactoriality warning). The convolution  $\mathrm{Hom}_\alpha(C, A)$  is *not* a bifunctor in both slots simultaneously (Vol I, three-pillar constraint (2); Robalo–Nickerson–Welcher 2019). MC3 holds one slot at a time. In the CY setting this means: varying  $C$  along a path of CY categories while holding  $\Phi(C)$  fixed, or varying  $A_C$  along a chain homotopy while holding  $C$  fixed, each gives a homotopy equivalence of MC moduli; the simultaneous variation does not (*a priori*) define a coherent bifunctor.

**THEOREM 22.4** (*CY shadow obstruction tower as MC element*). <sup>[PE]</sup> The Vol I shadow obstruction tower

$$\Theta_{A_C} = D_{A_C} - d_0 \in (\mathrm{Conv}_{\mathrm{str}}(B(A_C), A_C))$$

is a Maurer–Cartan element of the CY modular convolution algebra, solving

$$\delta \Theta_{A_C} + \frac{1}{2} [\Theta_{A_C}, \Theta_{A_C}]_{\mathrm{Conv}} = 0$$

at genus  $g \geq 0$  and degree  $n \geq 2$  with  $2g - 2 + n > 0$ . The degree filtration yields finite-order projections  $\Theta_{A_C}^{\leq r}$  converging to  $\Theta_{A_C}$  as  $r \rightarrow \infty$ .

*Attribution.* Vol I Theorem thm:recursive-existence establishes the all-degree inverse limit. The translation to  $\Phi$ -images follows from Proposition 22.2 and functoriality of the bar construction under the CY-to-chiral map.  $\square$

*Remark 22.5* (What lives where). The convolution dg Lie algebra  $\mathrm{Conv}_{\mathrm{str}}(B(A_C), A_C)$  is the ambient home of  $\Theta_{A_C}$ ; it is distinct from the chiral derived center  $\mathcal{Z}_{\mathrm{ch}}^{\mathrm{der}}(A_C) = \mathrm{RHom}(\Omega B(A_C), A_C)$ , which computes the bulk observables (Theorem H). The three functors  $\Omega, D_{\mathrm{Ran}}, C_{\mathrm{ch}}^\bullet(A, A)$  produce three distinct outputs from  $B(A_C)$ , and the convolution algebra is none of them: it is the *working surface* on which  $\Theta_{A_C}$  solves the master equation.

*Definition 22.6 (Categorical modular characteristic).* For a smooth proper  $\text{CY}_d$  category  $C$ , the *categorical modular characteristic* is

$$\kappa_{\text{cat}}(C) := \chi^{\text{CY}}(C) \stackrel{\text{def}}{=} \sum_{i=0}^d (-1)^i \dim \text{HH}_i(C).$$

This is the holomorphic Euler characteristic of  $C$  as a CY category (equivalently, for  $C = D^b(\text{Coh}(X))$  with  $X$  a smooth projective  $\text{CY}_d$  manifold,  $\kappa_{\text{cat}}(C) = \chi(\mathcal{O}_X) = \sum_i (-1)^i h^{0,i}(X)$ ). It is distinct *a priori* from the chiral modular characteristic  $\kappa_{\text{ch}}(A_C)$  of the chiral algebra  $A_C = \Phi(C)$ ; the two coincide at  $d = 2$  (Proposition 22.7) and are conjectured to coincide at  $d \geq 3$  (Conjecture 8.3; see Remark 22.8 below).

**PROPOSITION 22.7** ( $\kappa_{\text{cat}} = \kappa_{\text{ch}}$  under  $\Phi$  at  $d = 2$ ). <sup>[PH]</sup> For  $C$  a smooth proper  $\text{CY}_2$  category with chiral algebra  $A_C = \Phi(C)$  (CY-A at  $d = 2$ , PROVED),

$$\kappa_{\text{ch}}(A_C) = \kappa_{\text{cat}}(C) = \chi^{\text{CY}}(C).$$

Consequently, for every genus  $g \geq 1$  and on the uniform-weight lane,

$$\text{obs}_g(A_C) = \kappa_{\text{cat}}(C) \cdot \lambda_g \quad (g \geq 1, \text{ UNIFORM-WEIGHT});$$

at genus  $g \geq 2$  for multi-weight algebras the scalar formula receives a nonvanishing cross-channel correction  $\delta F_g^{\text{cross}}$  (ALL-WEIGHT, with cross-channel correction  $\delta F_g^{\text{cross}}$ ).

*Proof.* The free-field argument: the generating space of  $A_C$  is  $\text{HH}^{\bullet+1}(C)$ , and  $\kappa_{\text{ch}}$  equals the supertrace of the identity on this generating space, which is  $\chi^{\text{CY}}(C) = \kappa_{\text{cat}}(C)$ . The quantization step in the construction of  $\Phi$  (CY-A, Step 4) preserves  $\kappa_{\text{ch}}$  at  $d = 2$ : the holomorphic anomaly cancellation at  $d = 2$  (Serre duality  $\mathbb{S}_C \simeq [2]$ ) guarantees that no quantum correction shifts the supertrace. The genus- $g$  obstruction formula is Vol I Theorem D applied to  $A_C$ ; the substitution  $\kappa_{\text{ch}} = \kappa_{\text{cat}}$  follows from the first part.  $\square$

*Remark 22.8* ( $\kappa_{\text{cat}}$  versus  $\kappa_{\text{ch}}$ ). The categorical modular characteristic  $\kappa_{\text{cat}}(C)$  is a topological invariant of the CY category  $C$  (it depends only on the Hochschild homology dimensions). The chiral modular characteristic  $\kappa_{\text{ch}}(A_C)$  is an analytic invariant of the chiral algebra  $A_C$  (it depends on the OPE structure and the generating field content). Proposition 22.7 identifies them at  $d = 2$ ; at  $d = 3$  the identification is Conjecture 8.3 and is verified for  $\mathbb{C}^3$  (§22.5). Both are distinct from  $\kappa_{\text{BKM}}$  (the BKM algebra weight) and  $\kappa_{\text{fiber}}$  (the lattice rank); the four values constitute the  $\kappa_{\bullet}$ -spectrum (Remark 8.8o).

## 22.2 CY complementarity

Volume I Theorem C has two components. The eigenspace statement (C1) asserts that the genus- $g$  shadow complex of a Koszul pair  $(A, A^\dagger)$  splits into complementary eigenspaces for the Verdier involution; this holds unconditionally. The scalar statement (C2) asserts the sum

$$\kappa_{\text{ch}}(A) + \kappa_{\text{ch}}(A^\dagger) = \rho \cdot K,$$

where  $K$  is the Koszul conductor and  $\rho$  the anomaly ratio; this holds only on the *uniform-weight lane* (all generators of  $A$  of equal conformal weight), and at  $g \geq 2$  multi-weight algebras incur a nonvanishing cross-channel correction  $\delta F_g^{\text{cross}}$ . This section transports both statements to CY categories via the functor  $\Phi$ .

### The $d = 2$ theorem

**THEOREM 22.9** (*CY complementarity at  $d = 2$* ). <sup>[PH]</sup> Let  $\mathcal{C}$  be a smooth proper cyclic  $A_\infty$ -category of CY dimension  $d = 2$  with Serre duality  $\mathbb{S}_{\mathcal{C}} \simeq [2]$ , and let  $A_{\mathcal{C}} = \Phi(\mathcal{C})$  be its quantum chiral algebra (CY-A at  $d = 2$ , PROVED). Let  $\mathcal{C}^\dagger$  denote the Koszul dual  $\text{CY}_2$  category (for  $\mathcal{C} = D^b(\text{Coh}(X))$  with  $X$  a  $K_3$  surface,  $\mathcal{C}^\dagger \simeq \text{Fuk}(X)$  under homological mirror self-duality). Then:

(C1<sup>CY</sup>) *Eigenspace complementarity*. For every genus  $g \geq 1$  and every degree  $n \geq 1$  with  $2g - 2 + n > 0$ , the genus- $g$  shadow complex satisfies

$$Q_g^n(A_{\mathcal{C}}) \oplus Q_g^n(A_{\mathcal{C}^\dagger}) \simeq H^\bullet(\overline{\mathcal{M}}_{g,n}, \mathcal{Z}_{\text{ch}}^{\text{der}}(A_{\mathcal{C}})),$$

as a direct sum of  $\pm 1$  eigenspaces for the Verdier involution induced by Serre duality on  $D^b(\text{Coh}(\mathcal{C}))$ . This is unconditional in the  $\text{CY}_2$  case.

(C2<sup>CY</sup>) *Scalar complementarity (restricted scope)*. Under the CY uniform-weight hypothesis (all generating fields of  $A_{\mathcal{C}}$  at equal conformal weight) and in the free-field/lattice Koszul class (the KM/Heisenberg branch, where the Koszul conductor  $K = c(A_{\mathcal{C}}) + c(A_{\mathcal{C}^\dagger})$  vanishes),

$$\kappa_{\text{ch}}(A_{\mathcal{C}}) + \kappa_{\text{ch}}(A_{\mathcal{C}^\dagger}) = 0 \quad (\text{free-field/KM class}).$$

Away from the free-field class, the sum equals the family-dependent Koszul conductor and is nonzero in general: for the Virasoro class the analogous sum is  $\kappa_{\text{ch}} + \kappa'_{\text{ch}} = 13$  (not 0), and for  $\mathcal{W}_N$  it equals  $c \cdot (H_N - 1)$  where  $H_N = \sum_{j=1}^N 1/j$ . For  $\mathcal{C} = D^b(\text{Coh}(K_3))$  specifically,  $\kappa_{\text{ch}}(\mathcal{A}_{K_3}) = \chi^{\text{CY}}(K_3) = 2$  (Theorem CY-D, §10.1); the relevant chiral algebra is the  $\widehat{\mathfrak{sl}}_2$  subalgebra at level  $k = 1$  of the  $\mathcal{N} = 4$  superconformal algebra, which lies in the free-field/KM class with  $K = 0$ , so the Verdier involution induced by the Mukai pairing gives  $\kappa'_{\text{ch}} = -2$  and the scalar sum vanishes:  $\kappa_{\text{ch}} + \kappa'_{\text{ch}} = 0$  ( $K_3$  value, free-field/KM branch; NOT universal across all  $\text{CY}_2$  categories).

*Sketch.* (C1<sup>CY</sup>): the eigenspace decomposition is the  $\Phi$ -image of Vol I Theorem C1. The functor  $\Phi$  is compatible with the Verdier involution (Chapter 8, Proposition on Serre-functor intertwining), so the direct sum decomposition of Vol I pulls back to a decomposition of  $Q_g^n(A_{\mathcal{C}}) \oplus Q_g^n(A_{\mathcal{C}^\dagger})$  indexed by Serre eigenvalues.

(C2<sup>CY</sup>): under uniform weight, Vol I Theorem C2 gives  $\kappa_{\text{ch}} + \kappa'_{\text{ch}} = 0$  for the Heisenberg/free-field classes. For  $K_3$ ,  $\mathcal{A}_{K_3}$  is the  $\mathcal{N} = 4$  superconformal algebra with  $c = 6$ , and all generators sit at conformal weights  $\{1/2, 1, 3/2, 2\}$ ; the uniform-weight lane of interest is the  $\widehat{\mathfrak{sl}}_2$  subalgebra at level  $k = 1$ , where  $\kappa_{\text{ch}} = 2$  is attained. The Verdier involution under the  $K_3$  Mukai lattice self-duality  $\Lambda_{K_3} \simeq \Lambda_{K_3}^\vee$  acts by a sign on odd-degree Hochschild classes, giving  $\kappa'_{\text{ch}} = -2$ . The sum vanishes.  $\square$

*Remark 22.10* (Verification against CY-D). Substituting  $d = 2$  ( $K_3$ ) into  $\kappa_{\text{ch}}(A_{\mathcal{C}}) = \chi^{\text{CY}}(\mathcal{C})$  (Theorem 10.1) gives  $\kappa_{\text{ch}}(\mathcal{A}_{K_3}) = \chi^{\text{CY}}(K_3) = 2$ , which agrees with the independently verified chiral de Rham computation (§10.1, Proposition 14.50). Theorem 22.9 is therefore consistent with the five-path verification of  $\kappa_{\text{ch}}(K_3) = 2$  (compute/lib/modular\_cy\_characteristic.py, 80 tests).

*Remark 22.11* (Categorical complementarity). Proposition 22.7 allows the scalar complementarity (C2<sup>CY</sup>) to be restated on the categorical side. The *CY Koszul conductor* is

$$K_{\text{cat}} := \kappa_{\text{cat}}(\mathcal{C}) + \kappa_{\text{cat}}(\mathcal{C}^\dagger) = \chi^{\text{CY}}(\mathcal{C}) + \chi^{\text{CY}}(\mathcal{C}^\dagger).$$

On the free-field/KM branch (which includes  $K_3$  under the Mukai self-duality),  $K_{\text{cat}} = 0$ : the Koszul dual categorical characteristic is  $\kappa_{\text{cat}}(\mathcal{C}^\dagger) = -\kappa_{\text{cat}}(\mathcal{C})$ . For  $K_3$  specifically,  $\kappa_{\text{cat}}(D^b(\text{Coh}(K_3))) = 2$  and  $\kappa_{\text{cat}}(\text{Fuk}(K_3)) = -2$  under the Mukai-HMS involution, yielding  $K_{\text{cat}} = 0$ . Away from the free-field class,  $K_{\text{cat}}$  is nonzero and family-dependent (see Vol I, Theorem C2).

### The $d = 3$ conjecture

CONJECTURE 22.12 (*CY complementarity at  $d = 3$* ). <sup>[CJ]</sup> Let  $X$  be a compact Calabi–Yau threefold and  $C = D^b(\text{Coh}(X))$  its bounded derived category, with cyclic  $A_\infty$  structure from the Serre trace. Assume CY- $A_3$ , so that  $A_X = \Phi(C)$  exists as an  $E_2$ -chiral algebra. Let  $C^\dagger$  denote the Koszul dual  $\text{CY}_3$  category, conjectured (under homological mirror symmetry) to be  $\text{Fuk}(X^\vee)$  for the mirror threefold  $X^\vee$ . Then:

(C1<sup>CY<sub>3</sub></sup>) The genus- $g$  shadow complexes satisfy the eigenspace complementarity

$$Q_g^n(A_X) \oplus Q_g^n(A_{X^\vee}) \simeq H^\bullet(\overline{\mathcal{M}}_{g,n}, \mathcal{Z}_{\text{ch}}^{\text{der}}(A_X)),$$

conditionally on CY- $A_3$  (the conditionality propagates, AP-CY $\Pi$ ).

(C2<sup>CY<sub>3</sub></sup>) Under the CY uniform-weight hypothesis (which is *not* generically satisfied for compact  $\text{CY}_3$ : the chiral de Rham complex has fields in weights  $\{1/2, 1, 3/2, 2, \dots\}$ ), and scope (the Koszul conductor is family-dependent: free-field/KM gives  $K = 0$ , Virasoro gives  $K = 26$  with scalar sum 13,  $\mathcal{W}_N$  gives  $K_N = 4N^3 - 2N - 2$  with scalar sum  $c(H_N - 1)$ ),

$$\kappa_{\text{ch}}(A_X) + \kappa_{\text{ch}}(A_{X^\vee}) = \rho \cdot K_X \quad (\text{CY}_3, \text{family-dependent, nonzero in general}),$$

where  $K_X = c(A_X) + c(A_{X^\vee})$  is the CY Koszul conductor and  $\rho$  is the CY anomaly ratio. For  $X = X_{\text{quintic}}$  with  $\chi_{\text{top}} = -200$ , the BCOV prediction  $\kappa_{\text{ch}} = \chi_{\text{top}}/24 = -25/3$  would give a scalar sum of  $-50/3$  on the self-mirror diagonal; the conjecture predicts this equals  $\rho \cdot K_{\text{quintic}}$ .

*Remark 22.13* (Why  $d = 3$  is a conjecture, not a theorem). Three independent obstructions block upgrading Conjecture 22.12 to a theorem: (a) AP-CY6, the chiral algebra  $A_X$  is not constructed at  $d = 3$ ; (b) the uniform-weight hypothesis fails for compact  $\text{CY}_3$  (chiral de Rham is multi-weight, so gives  $\delta F_g^{\text{cross}} \neq 0$  at  $g \geq 2$ ); (c) the BKM automorphic correction at  $d = 3$  generates infinitely many imaginary root generators (§22.3 below), so even stating the Koszul conductor  $K_X$  requires resolving the degree- $r$  shadow identification of theory\_automorphic\_shadow.

*Remark 22.14* (Degeneracy at vanishing modular characteristic warning). When  $\kappa_{\text{ch}}(A_C)$  vanishes (banana manifold, Heisenberg at level  $k = 0$ , Virasoro at  $c = 0$ ; see §20.7), the free-field/KM branch of the scalar complementarity forces the Koszul-dual characteristic  $\kappa'_{\text{ch}}$  to vanish likewise, but this does *not* imply  $\Theta_{A_C}$  vanishes. Higher-degree shadow components can remain nonzero, sourced in the banana case by genus-0 GV invariants. The leading scalar complementarity degenerates; the full tower complementarity continues to encode nontrivial Koszul duality data. Note: the Virasoro-at- $c = 0$  example sits on the free-field boundary of the  $c = 13$  self-dual Virasoro family, where the generic scalar sum is  $\kappa_{\text{ch}} + \kappa'_{\text{ch}} = 13$ ;  $c = 0$  is the exceptional point of that family.

## 22.3 The CY shadow CohFT

Volume I Theorem D promotes the shadow obstruction tower to a cohomological field theory (CohFT) on  $\overline{\mathcal{M}}_{g,n}$  under a flat identity axiom. requires the flat identity to live in the generating space (vacuum in  $V$ ); we list this conditionality at every cross-reference.

### CohFT structure

THEOREM 22.15 (*CY shadow CohFT*). <sup>[PH]</sup> Let  $C$  be a smooth proper cyclic  $A_\infty$ -category of CY dimension  $d$ , and let  $A_C = \Phi(C)$ . **Assume:** the vacuum  $\mathbf{1}_{A_C}$  lies in the generating space of  $A_C$  (equivalently,  $\mathbf{1}_{A_C}$  is flat for the Dubrovin connection of the shadow tower). Then the degree- $n$  shadow classes

$$\Omega_{g,n}(A_C) := \pi_{g,n,*}(\Theta_{A_C}^{(g,n)}) \in H^\bullet(\overline{\mathcal{M}}_{g,n})^{\otimes n}$$

assemble into a CohFT: the pullbacks along the boundary maps  $\overline{\mathcal{M}}_{g_1, n_1+1} \times \overline{\mathcal{M}}_{g_2, n_2+1} \rightarrow \overline{\mathcal{M}}_{g_1+g_2, n_1+n_2}$  and the loop  $\overline{\mathcal{M}}_{g-1, n+2} \rightarrow \overline{\mathcal{M}}_{g, n}$  factor through the Koszul pair  $(A_C, A_C^\dagger)$ , and the unit axiom holds by the flat identity assumption.

*Sketch.* Vol I Theorem D+H constructs the CohFT structure on the shadow tower of a chiral algebra with flat identity. The functor  $\Phi$  preserves the flat identity (it sends the cyclic unit of  $C$  to the vacuum of  $A_C$ ), so under the hypothesis, the pulled-back shadow classes on  $\overline{\mathcal{M}}_{g, n}$  satisfy the CohFT axioms. The level at which the statement holds (ambient versus convolution, §20.1): this is the ambient level, built via Mok's log Fulton–MacPherson boundary differential.  $\square$

*Remark 22.16* (conditionality). The flat identity hypothesis is conditional and must be stated at every cross-reference. Three scenarios where it fails: (i) CY categories without a categorical unit (rare but possible for nonunital  $A_\infty$  models); (ii) vertex algebras where the vacuum does not lie in the generating space (e.g. coset constructions); (iii)  $W$ -algebras with nontrivial BRST cohomology at degree zero. Every theorem that invokes Theorem 22.15 downstream (e.g. the Igusa cusp form recovery below) inherits this hypothesis.

### The $K3 \times E$ CohFT and the Igusa cusp form

**THEOREM 22.17** ( *$K3 \times E$  shadow CohFT and  $\Phi_{10}$* ). <sup>[PE]</sup> Let  $X = K3 \times E$ , with chiral algebra  $\mathcal{A}_{K3} \otimes H_1$  (chiral de Rham complex of  $K3$  tensored with the Heisenberg algebra of  $E$ ),  $\kappa_{\text{ch}}(K3 \times E) = 3$  by additivity (Proposition 10.4;  $K3$ -I of §14.36). Assume the flat identity hypothesis. Then:

- (i) The shadow CohFT  $\Omega_{g, n}(\mathcal{A}_{K3} \otimes H_1)$  exists at all  $g \geq 1$  and  $n \geq 1$  with  $2g - 2 + n > 0$  (Theorem 22.15).
- (ii) The Borcherds multiplicative lift transports the genus-1 datum (the elliptic genus of  $K3$ ) to the genus-2 Siegel modular form

$$\Phi_{10}(\Omega) = \text{Borch}(\phi_{-2,1})(\Omega), \quad \text{wt}(\Phi_{10}) = 10,$$

where  $\phi_{-2,1}$  is the weak Jacobi form of weight  $-2$  and index  $1$ ; this is the Igusa cusp form, §14.31 and chapters/examples/toroidal\_elliptic.tex equation (5.1).

- (iii) The BPS modular characteristic  $\kappa_{\text{BKM}} = 5 = \text{wt}(\Phi_{10})/2$  is distinct from the chiral characteristic  $\kappa_{\text{ch}} = 3$  (the  $\kappa_\bullet$ -spectrum polysemy, Remark 8.80; neither value is universal).
- (iv) The shadow obstruction tower of  $\mathcal{A}_{K3} \otimes H_1$  does *not* by itself reproduce  $\Phi_{10}$ : four obstructions ( $K3$ -4 of §14.36) separate the shadow tower output from  $\Phi_{10}$ . Namely, (O<sub>1</sub>) a categorical obstruction, (O<sub>2</sub>) the  $\kappa_{\text{ch}}/\kappa_{\text{BKM}}$  mismatch  $3 \neq 5$ , (O<sub>3</sub>) second quantization (the Hilbert–Chow exceptional divisor), and (O<sub>4</sub>) the Schottky obstruction at  $g \geq 4$  of codimension  $(g-2)(g-3)/2$ . The Borcherds lift supplies precisely the combinatorial data needed to bridge these four obstructions.

*Attribution.* Parts (i), (iii), (iv) are Vol I/II results (§14.36, items  $K3$ -1 through  $K3$ -13, proved in the  $K3$  modular-tower chapter of Vol I and in the holography part of Vol II). Part (ii) is the Borcherds lift of Borcherds (1998); its use as a modular characteristic is the observation of AP-CY8, not a theorem of Vol III. The conditionality propagates from Theorem 22.15.  $\square$

*Remark 22.18* (Kappa-spectrum verification for  $K3 \times E$ ). Substitute  $d = 2$  for  $K3$  into the CY-D formula  $\kappa_{\text{ch}}(A_C) = \chi^{\text{CY}}(C)$ : this gives  $\kappa_{\text{ch}}(\mathcal{A}_{K3}) = 2$ . Tensor-product additivity (Proposition 10.4) for the chiral de Rham complex then determines  $\kappa_{\text{ch}}(K3 \times E) = 3$  (namely, the value 2 for  $K3$  combined additively with the value 1 for the elliptic



curve  $E$  yields 3). Note: this is additivity of the modular characteristic under product factorization, not the Koszul-dual scalar sum of Theorem 22.9 ( $C_2^{CY}$ ); the two operations are distinct and should not be confused. The value  $\kappa_{\text{BKM}} = 5$  is the Borchers weight, independently equal to half the weight of  $\Phi_{10}$ ; this corresponds to the BKM superalgebra  $\mathfrak{g}_{\Delta_5}$ , not to the chiral algebra of  $K3 \times E$ . The four members of the spectrum  $\{\kappa_{\text{cat}}, \kappa_{\text{ch}}, \kappa_{\text{BKM}}, \kappa_{\text{fiber}}\} = \{2, 3, 5, 24\}$  each arise from a distinct chiral algebraization: bare kappa is forbidden in Vol III.

### Toric $CY_3$ and the DT CohFT

**CONJECTURE 22.19** (*Toric  $CY_3$  shadow CohFT*). <sup>[CJ]</sup> Let  $X_\Sigma$  be a smooth toric  $CY_3$  with fan  $\Sigma$ . Conditional on  $CY\text{-}A_3$ , the shadow CohFT  $\Omega_{g,n}(A_{X_\Sigma})$  exists and satisfies:

- (i) The genus-0 generating function of  $\Omega_{0,n}$  equals the topological vertex of Aganagic–Klemm–Mariño–Vafa;
- (ii) The full genus expansion equals the DT partition function  $Z^{\text{DT}}(X_\Sigma)$ , which is the MacMahon prefactor for  $X_\Sigma = \mathbb{C}^3$  and the refined MacMahon for the resolved conifold (§20.2).

*Remark 22.20* (Scope). Conjecture 22.19 is double-conditional: on  $CY\text{-}A_3$  and on the flat identity hypothesis (Remark 22.16). The resolved conifold case is the least obstructed:  $\kappa_{\text{ch}} = 1$  is known (§10.1), and the  $W_{1+\infty}$  vertex algebra has a flat vacuum. For  $X_\Sigma = \mathbb{C}^3$ , the verification is partial: the MacMahon identification of the bar Euler product is Theorem 20.3, but the upgrade to a full CohFT on  $\overline{\mathcal{M}}_{g,n}$  awaits resolution of the  $d = 3$  degree-to-depth correspondence (theory\_automorphic\_shadow, §22.3).

## 22.4 The Hochschild bridge

Three Hochschild theories act on a  $CY$  category  $\mathcal{C}$  with chiral algebra  $A_{\mathcal{C}} = \Phi(\mathcal{C})$ , and distinguishing them is essential for the bridge to Volume I.

**Definition 22.21** (*Three Hochschild theories*). Let  $\mathcal{C}$  be a smooth proper  $CY_d$  category with chiral algebra  $A_{\mathcal{C}} = \Phi(\mathcal{C})$ .

- (i) *Categorical Hochschild*. The cyclic bar complex  $CC_\bullet(\mathcal{C})$  with Hochschild differential  $b$  and Connes operator  $B$ . This is the topological invariant of  $\mathcal{C}$  as a dg category: it carries the Gerstenhaber bracket and the  $(2 - d)$ -shifted Poisson structure from the Serre pairing (Chapter 4).
- (ii) *Chiral Hochschild*. The chiral Hochschild cohomology  $\text{ChirHoch}^*(A_{\mathcal{C}})$  of Volume I (Theorem H): concentrated in cohomological degrees  $\{0, 1, 2\}$  with polynomial Hilbert series. This is the obstruction-theoretic invariant governing the shadow tower of  $A_{\mathcal{C}}$ .
- (iii) *Derived center*. The chiral derived center  $\mathcal{Z}_{\text{ch}}^{\text{der}}(A_{\mathcal{C}}) = \text{RHom}(\Omega \mathcal{B}(A_{\mathcal{C}}), A_{\mathcal{C}})$ : the universal bulk algebra, computed by chiral Hochschild *cochains* (not cohomology). This is the target of the holographic datum (Vol II).

These are distinct invariants. The categorical Hochschild lives on  $\mathcal{C}$  (no curve geometry); the chiral Hochschild lives on  $A_{\mathcal{C}}$  (curve geometry through the OPE); the derived center is the endomorphism algebra of the identity functor in the factorization category (curve geometry through  $\text{Ran}(X)$ ).

The bridge theorem identifies how  $\Phi$  intertwines the categorical and chiral sides.

**THEOREM 22.22** (*Hochschild bridge*). <sup>[PH]</sup> Let  $\mathcal{C}$  be a smooth proper CY<sub>2</sub> category and  $A_{\mathcal{C}} = \Phi(\mathcal{C})$  (CY-A at  $d = 2$ , PROVED). Then  $\Phi$  induces:

- (i) A quasi-isomorphism of factorization coalgebras  $\mathrm{CC}_{\bullet}(\mathcal{C}) \xrightarrow{\sim} B(A_{\mathcal{C}})$  on  $\mathrm{Ran}(X)$  (this is CY-A(ii), Proposition 20.1).
- (ii) A map on cohomology  $\mathrm{HH}^{\bullet}(\mathcal{C}) \rightarrow \mathrm{ChirHoch}^*(A_{\mathcal{C}})$  that sends the Gerstenhaber bracket on  $\mathrm{HH}^{\bullet}(\mathcal{C})$  to the convolution bracket on  $\mathrm{ChirHoch}^*(A_{\mathcal{C}})$ , and sends the Connes  $B$ -operator to the modular differential.
- (iii) A map  $\mathrm{HH}^{\bullet}(\mathcal{C}, \mathcal{C}) \rightarrow \mathcal{Z}_{\mathrm{ch}}^{\mathrm{der}}(A_{\mathcal{C}})$  from the categorical Hochschild cochains (the endomorphism algebra of the identity bimodule) to the chiral derived center, compatible with the Gerstenhaber product on the source and the chiral bracket on the target.

*Sketch.* Part (i) is Proposition 20.1. For (ii): the Gerstenhaber bracket on  $\mathrm{HH}^{\bullet}(\mathcal{C})$  is a Lie bracket of degree  $-1$ . Under  $\Phi$ , this bracket becomes the convolution bracket of Vol I by functoriality of the bar construction: the deconcatenation coproduct on  $B(A_{\mathcal{C}})$  and the binary product of  $A_{\mathcal{C}}$  assemble into a bracket on  $\mathrm{Hom}_{\mathrm{Ran}}(B(A_{\mathcal{C}}), A_{\mathcal{C}})$ , which restricts to  $\mathrm{ChirHoch}^*(A_{\mathcal{C}})$  on the cohomology. The Connes operator  $B$  on  $\mathrm{CC}_{\bullet}(\mathcal{C})$  corresponds, under the bar dictionary, to the degree-preserving component of the bar differential (Proposition 20.1(ii)), which is the modular differential on the chiral side. For (iii): the categorical Hochschild cochains  $\mathrm{CC}^{\bullet}(\mathcal{C}, \mathcal{C})$  map to  $\mathrm{RHom}(\Omega B(A_{\mathcal{C}}), A_{\mathcal{C}})$  by applying  $\Phi$  to each hom-space and using the bar-cobar quasi-isomorphism  $\Omega B(A_{\mathcal{C}}) \simeq A_{\mathcal{C}}$  (Vol I Theorem B on the Koszul locus).  $\square$

*Remark 22.23* (Which Hochschild controls which invariant). The categorical Hochschild  $\mathrm{HH}_{\bullet}(\mathcal{C})$  controls the deformation theory of  $\mathcal{C}$  as a CY category:  $\mathrm{HH}^2(\mathcal{C})$  parametrizes first-order deformations,  $\mathrm{HH}^1(\mathcal{C})$  the infinitesimal automorphisms. The chiral Hochschild  $\mathrm{ChirHoch}^*(A_{\mathcal{C}})$  controls the shadow tower of  $A_{\mathcal{C}}$ : the obstruction classes  $\mathrm{obs}_g$  live in  $\mathrm{ChirHoch}^2(A_{\mathcal{C}}) \otimes H^{\bullet}(\overline{\mathcal{M}}_{g,n})$ . The derived center  $\mathcal{Z}_{\mathrm{ch}}^{\mathrm{der}}(A_{\mathcal{C}})$  is the bulk algebra of the holographic datum: its elements are the operators that commute with all boundary insertions. Theorem 22.22 guarantees that the categorical deformation data of  $\mathcal{C}$  is faithfully transmitted, through  $\Phi$ , to the chiral obstruction data of  $A_{\mathcal{C}}$ .

**CONJECTURE 22.24** (*Hochschild bridge at  $d = 3$* ). <sup>[CJ]</sup> For  $\mathcal{C}$  a smooth proper CY<sub>3</sub> category, conditional on CY-A<sub>3</sub> (the conditionality propagates, AP-CYII), the maps (i)–(iii) of Theorem 22.22 extend to  $d = 3$ . The  $(-1)$ -shifted Poisson structure on  $\mathrm{HH}^{\bullet}(\mathcal{C})$  (Pantev–Toën–Vaquié–Vezzosi) maps to the genus-0 contribution of the convolution bracket on  $\mathrm{ChirHoch}^*(A_{\mathcal{C}})$ ; the genus- $g \geq 1$  components of the convolution bracket have no direct categorical-Hochschild source and arise from the curve geometry of  $\mathrm{Ran}(X)$  through  $\Phi$ .

## 22.5 Principal examples

The modular Koszul bridge produces concrete numerical data for each CY category: the  $\kappa_{\bullet}$ -spectrum, the shadow class, and the genus- $g$  obstruction tower. This section collects the principal examples.

### $K3 \times E$

The product  $K3 \times E$  ( $K3$  surface times an elliptic curve) is CY<sub>3</sub> with independently computable chiral algebra  $\mathcal{A}_{K3} \otimes H_1$ . The  $\kappa_{\bullet}$ -spectrum (the four distinct modular characteristics arising from four distinct chiral algebraizations) is:

| Subscript               | Source                                             | Value | Derivation                                                           |
|-------------------------|----------------------------------------------------|-------|----------------------------------------------------------------------|
| $\kappa_{\text{cat}}$   | $\chi^{\text{CY}}(K3) = \chi(O_{K3})$              | 2     | Hodge diamond: $h^{0,0} - h^{1,0} + h^{2,0} = 1 - 0 + 1$             |
| $\kappa_{\text{ch}}$    | $\kappa_{\text{ch}}(\mathcal{A}_{K3} \otimes H_1)$ | 3     | Additivity: $\kappa_{\text{ch}}(K3) + \kappa_{\text{ch}}(E) = 2 + 1$ |
| $\kappa_{\text{BKM}}$   | $\text{wt}(\Phi_{10})/2$                           | 5     | BKM superalgebra weight                                              |
| $\kappa_{\text{fiber}}$ | $\text{rank}(\Lambda_{K3})$                        | 24    | $K3$ lattice rank                                                    |

The obstruction class at genus  $g$  (UNIFORM-WEIGHT, for the  $\widehat{\mathfrak{sl}}_2$  subalgebra at level  $k = 1$ , which is the uniform-weight sector of the  $\mathcal{N} = 4$  SCA):

$$\text{obs}_g = \kappa_{\text{ch}} \cdot \lambda_g = 3\lambda_g \quad (g \geq 1, \text{ UNIFORM-WEIGHT}).$$

For the full  $\mathcal{N} = 4$  algebra (multi-weight, generators at conformal weights  $\{1/2, 1, 3/2, 2\}$ ), the scalar formula acquires a correction  $\delta F_g^{\text{cross}} \neq 0$  at  $g \geq 2$  (ALL-WEIGHT, with cross-channel correction).

The complementarity data:  $\kappa_{\text{ch}}(\mathcal{A}_{K3}) + \kappa_{\text{ch}}(\mathcal{A}_{K3}^1) = 0$  on the free-field/KM branch (Theorem 22.9,  $(C_2^{\text{CY}})$ ).

### $\mathbb{C}^3$ and $\mathcal{W}_{1+\infty}$

The noncompact  $\text{CY}_3$  category  $\mathcal{C} = D^b(\text{Coh}(\mathbb{C}^3))$  produces, on the CY side, the  $\text{CoHA}(\mathbb{C}^3) \simeq Y^+(\widehat{\mathfrak{gl}}_1)$  (the positive half of the affine Yangian of  $\widehat{\mathfrak{gl}}_1$ ). The chiral algebra is  $\mathcal{W}_{1+\infty}$  at  $c = 1$  (Schiffmann–Vasserot, Maulik–Okounkov). The  $\kappa_{\bullet}$ -values:

| Subscript             | Value | Source                                                         |
|-----------------------|-------|----------------------------------------------------------------|
| $\kappa_{\text{cat}}$ | 1     | $\chi^{\text{CY}}(\mathbb{C}^3) = 1$ (noncompact, regularized) |
| $\kappa_{\text{ch}}$  | 1     | $\kappa_{\text{ch}}(\mathcal{W}_{1+\infty} _{c=1})$            |

Here  $\kappa_{\text{cat}} = \kappa_{\text{ch}}$ : the categorical and chiral modular characteristics coincide. This is the content of Conjecture 8.3 (CY-A(iii)) specialized to  $\mathbb{C}^3$ , where both sides are independently computable. The shadow tower of  $\mathcal{W}_{1+\infty}$  at  $c = 1$  has class M (infinite shadow depth), with the bar Euler product recovering the MacMahon function  $M(q) = \prod_{n \geq 1} (1 - q^n)^{-n}$  (Theorem 20.3). The shadow CohFT is conditional on CY-A<sub>3</sub> and the flat identity hypothesis (Conjecture 22.19).

The obstruction class (UNIFORM-WEIGHT, at  $c = 1$  where  $\mathcal{W}_{1+\infty}$  is freely generated by a single field of weight 1):

$$\text{obs}_g = \kappa_{\text{ch}} \cdot \lambda_g = \lambda_g \quad (g \geq 1, \text{ UNIFORM-WEIGHT}).$$

*Remark 22.25* (Shadow depth versus Koszulness).  $\mathcal{W}_{1+\infty}$  at  $c = 1$  is Koszul (PBW degree concentration in the bar complex) but has class M (infinite shadow depth: all  $m_k^{\text{shadow}} \neq 0$ ). Koszulness is a cohomological property of the bar complex; shadow depth measures the vanishing pattern of the obstruction tower. The two invariants are independent (Vol I, AP14).

The five sections transport the Vol I modular Koszul machine into the CY geometric realm: the convolution algebra of §22.1 is the working surface, the complementarity of §22.2 is the duality statement, the CohFT of §22.3 is the genus tower, the Hochschild bridge of §22.4 identifies which Hochschild theory controls which invariant, and the examples of §22.5 verify the  $\kappa_{\bullet}$ -spectrum against independent computations. The  $d = 2$  case is unconditional (CY-A proved, Theorem 22.9); the  $d = 3$  case is the Vol III programme (Conjecture 22.12, Conjecture 22.19, Conjecture 22.24). Verification of every  $\kappa_{\bullet}$ -value uses the independent paths of `compute/lib/modular_cy_characteristic.py` and `compute/lib/cy_euler.py`, cross-checked against the  $\kappa_{\bullet}$ -spectrum  $\{\kappa_{\text{cat}}, \kappa_{\text{ch}}, \kappa_{\text{BKM}}, \kappa_{\text{fiber}}\} = \{2, 3, 5, 24\}$  for  $K3 \times E$ .



## **Part V**

# **The CY Frontier**



## Chapter 23

# Geometric Langlands and CY Quantum Groups

The functor  $\Phi$  of II sends a Calabi–Yau category to an  $E_2$ -chiral algebra; the bar complex of the output (Volume I, Theorem A) is the factorization invariant on which geometric Langlands is ultimately a statement. This chapter traces the thread. At the critical level the Feigin–Frenkel theorem identifies the chiral center with the algebra of  $G^L$ -opers; the Verdier intertwining of Volume I Theorem A then relates local geometric Langlands to the four-functor picture (bar, cobar, Verdier, derived center). For Calabi–Yau input, the analogue is conjectural: a Langlands dual of a CY  $d$ -category should realize the mirror of its  $\Phi$ -image. The chapter is entirely FRONTIER material. Every formal statement uses `\begin{conjecture}` unless it is a literal citation of Feigin–Frenkel (1992) or Frenkel–Gaitsgory (2006), in which case it is tagged <sup>[PE]</sup>.

### 23.1 The Feigin–Frenkel center at the critical level

Let  $\mathfrak{g}$  be a simple finite-dimensional complex Lie algebra and  $\hat{\mathfrak{g}}_k$  its affine Kac–Moody algebra at level  $k$ . The vacuum vertex algebra  $V_k(\mathfrak{g})$  is the universal chiral algebra generated by the currents  $J^a(z)$  with the Kac–Moody OPE. The *critical level* is  $k_c = -h^\vee$ , where  $h^\vee$  is the dual Coxeter number.

**THEOREM 23.1** (*Feigin–Frenkel, 1992*). <sup>[PE]</sup> At the critical level  $k = -h^\vee$ , the center of the vacuum vertex algebra is canonically isomorphic to the algebra of functions on the space of  $G^L$ -opers on the formal disk:

$$\mathfrak{z}(\hat{\mathfrak{g}}) := Z(V_{-h^\vee}(\mathfrak{g})) \xrightarrow{\sim} \mathrm{Fun}(\mathrm{Op}_{G^L}(D)).$$

Here  $G^L$  is the Langlands dual group and  $\mathrm{Op}_{G^L}(D)$  is the scheme of  $G^L$ -opers on the formal disk  $D = \mathrm{Spec} \mathbb{C}[[t]]$ .

The isomorphism is the content of Feigin–Frenkel (1992), with the full exposition in Frenkel’s book *Langlands Correspondence for Loop Groups* (2007). Off the critical level the center is trivial; at  $k = -h^\vee$  it becomes a commutative vertex subalgebra of maximal size, and the resulting ind-scheme is  $\mathrm{Op}_{G^L}(D)$ .

**Free-field resolution and the bar complex.** The proof of the Feigin–Frenkel isomorphism proceeds via the Wakimoto realization: a free-field embedding  $V_{-h^\vee}(\mathfrak{g}) \hookrightarrow \Pi_{-h^\vee}$  into a  $\beta\gamma$ -system tensored with a Heisenberg algebra, followed by a BRST reduction. The bar complex  $B(V_{-h^\vee}(\mathfrak{g})) = T^c(s^{-1}\overline{V_{-h^\vee}(\mathfrak{g})})$  of the critical-level vacuum algebra carries the deconcatenation coproduct of Volume I. The Wakimoto

free-field embedding induces a map  $B(V_{-h^\vee}(\mathfrak{g})) \rightarrow B(\Pi_{-h^\vee})$  of factorization coalgebras. Since  $\Pi_{-h^\vee}$  is a tensor product of free-field algebras, its bar complex is computed by the abelian (class G) shadow tower, where all operations above degree two vanish. The nontrivial content of the Feigin–Frenkel isomorphism, from the bar-complex perspective, is that the BRST cohomology of the Wakimoto complex computes  $\text{Fun}(\text{Op}_{GL}(D))$  as the Verdier-dual of a specific summand of  $B(V_{-h^\vee}(\mathfrak{g}))$ .

**Locating Feigin–Frenkel among the four functors.** Volume I Theorem A organizes four distinct operations on chiral algebras over  $\text{Ran}(X)$ . They must never be conflated (Volume I).

- (a) The bar functor  $B$ , producing a factorization coalgebra.
- (b) Verdier duality  $D_{\text{Ran}}$  applied to the bar, producing the linear-dual algebra denoted  $A^\dagger$ .
- (c) Inversion, returning the original algebra up to quasi-isomorphism on the Koszul locus.
- (d) The derived chiral center  $Z_{\text{ch}}^{\text{der}}$ , computed as chiral Hochschild cochains; this is the bulk.

Feigin–Frenkel is a statement about the chiral center  $\mathfrak{z}(\hat{\mathfrak{g}}) \subset V_{-h^\vee}(\mathfrak{g})$ . The relevant legs of the four-functor picture are the derived center of item (d), and the Verdier leg of item (b). It is not an instance of inversion (item (c)). In particular one should not describe Feigin–Frenkel as “bar followed by cobar produces the spectral side.” For  $A = V_{-h^\vee}(\mathfrak{g})$ , the bar complex  $B(A)$  carries the deconcatenation coproduct of  $T^c(s^{-1}\bar{A})$  (Volume I); the Verdier-dual complex is the habitat in which  $\text{Fun}(\text{Op}_{GL})$  should be located.

At the critical level  $\kappa_{\text{ch}}(V_{-h^\vee}(\mathfrak{g})) = \dim(\mathfrak{g}) \cdot (k + h^\vee) / (2h^\vee) = 0$ . In the AP126 trace-form convention the classical  $r$ -matrix is still  $r(z) = k \Omega / z$ , so at  $k = -h^\vee$  it is  $-h^\vee \Omega / z$ , not 0. The critical phenomenon is the vanishing of  $\kappa_{\text{ch}}$  and the resulting shadow/center degeneration encoded by  $\Theta_A$ , not a level-stripped disappearance of the affine collision residue.

**CONJECTURE 23.2** (*Critical-level Verdier-intertwining*; <sup>[CJ]</sup>). Let  $A = V_{-h^\vee}(\mathfrak{g})$  and write  $A^\dagger$  for the Verdier-dual chiral algebra  $D_{\text{Ran}}(B(A))$  of Volume I Theorem A. At the critical level the chiral-algebra inclusion  $\mathfrak{z}(\hat{\mathfrak{g}}) \hookrightarrow A$ , combined with the Feigin–Frenkel isomorphism of Theorem 23.1, implies (does not iff) the existence of a factorization-coalgebra map  $\text{Fun}(\text{Op}_{GL}(D)) \rightarrow B(A^\dagger)$  on  $\text{Ran}(X)$ . The conjecture is that this map is a quasi-isomorphism on the Volume I Koszul locus. This is a statement about the Verdier leg of the four-functor picture, not about the inversion leg  $\Omega \circ B$ .

The forward implication (Feigin–Frenkel implies a bar-coalgebra map) follows from Theorem 23.1 and the factorization construction of  $\text{Op}_{GL}$ . The converse (that the map is a quasi-isomorphism) is open.

**Remark on the Koszul locus.** Volume I defines the Koszul locus as the set of chiral algebras  $A$  for which the inversion map  $\Omega(B(A)) \rightarrow A$  is a quasi-isomorphism. For  $V_k(\mathfrak{g})$  with  $\mathfrak{g}$  simple and  $k$  generic, membership in the Koszul locus is a theorem of Volume I (the Kac–Moody family is class L, shadow depth  $r = 3$ ). At the critical level  $k = -h^\vee$  the chiral modular characteristic  $\kappa_{\text{ch}}$  vanishes, and membership requires a separate argument because the shadow metric degenerates there even though the AP126 trace-form  $r$ -matrix remains nonzero. The status of  $V_{-h^\vee}(\mathfrak{g})$  on the Koszul locus is therefore itself conjectural and feeds into Conjecture 23.2.



**Remark on the opposite level.** Vol I records that the Koszul conductor for the Kac–Moody family is  $\kappa_{\text{ch}} + \kappa'_{\text{ch}} = 0$  where  $\kappa'_{\text{ch}} = \kappa_{\text{ch}}(V_{k'}(\mathfrak{g}))$  with  $k' = -k - 2h^\vee$  (the Feigin–Frenkel reflection). The two critical-level points  $k = -h^\vee$  and  $k' = -h^\vee$  coincide: the critical level is the fixed point of the Feigin–Frenkel reflection. This is the algebraic reason the critical level is distinguished, and the source of the self-duality language that must not be transposed to universal form.

**D-modules on  $\text{Bun}_G$  and chiral localization.** The Frenkel–Gaitsgory programme (2006–) globalizes the Feigin–Frenkel center from the formal disk to an algebraic curve  $X$ . For each smooth projective curve  $X$  and reductive group  $G$ , the category  $\hat{\mathfrak{g}}_{\text{crit}}\text{-mod}^{\text{Ran}(X)}$  of chiral modules for the critical-level vacuum algebra over  $\text{Ran}(X)$  admits a localization functor

$$\Delta_X: \hat{\mathfrak{g}}_{\text{crit}}\text{-mod}^{\text{Ran}(X)} \longrightarrow D\text{-mod}(\text{Bun}_G(X))$$

whose essential image at the abelian level is the category of Hecke eigensheaves. The localization  $\Delta_X$  is a chiral-algebraic shadow of the Beilinson–Drinfeld construction: the factorization structure of the vacuum module on  $\text{Ran}(X)$  produces, upon localization, the D-module structure on  $\text{Bun}_G$ . The bar complex  $B(V_{-h^\vee}(\mathfrak{g}))$  on  $\text{Ran}(X)$  maps, under the Verdier leg of Theorem A, to the factorization algebra that controls the spectral side. The passage from factorization coalgebra to D-module on  $\text{Bun}_G$  is the content of  $\Delta_X$ ; its compatibility with the four-functor picture of Volume I is the subject of Conjecture 23.3.

**CONJECTURE 23.3** (*Factorization-to-D-module compatibility*; <sup>[CJ]</sup>). The localization functor  $\Delta_X$  intertwines the Verdier leg  $D_{\text{Ran}} \circ B$  of Volume I Theorem A with the de Rham functor on  $\text{Bun}_G(X)$ . Concretely: for  $A = V_{-h^\vee}(\mathfrak{g})$ , the D-module  $\Delta_X(A^\dagger)$  on  $\text{Bun}_G(X)$  is the Hecke eigensheaf associated to the trivial  $G^L$ -local system, and the factorization-coalgebra structure of  $B(A)$  on  $\text{Ran}(X)$  determines the Hecke property via the chiral Verdier intertwining. This is an “implies” statement: the factorization structure determines the Hecke property, but the converse (recovering the factorization structure from the Hecke property alone) is open.

## 23.2 CY Langlands for $d = 1, 2, 3$

The statement of Feigin–Frenkel is that the *center* of a chiral algebra at a distinguished level encodes opers for the Langlands dual group. For a CY category  $\mathcal{C}$ , the functor  $\Phi$  produces a chiral algebra  $A_{\mathcal{C}}$  whose bar complex is the cyclic bar complex of  $\mathcal{C}$  (Theorem 8.1); a CY analogue of Langlands duality should therefore manifest as a pairing  $\mathcal{C} \leftrightarrow \mathcal{C}^L$  on the source of  $\Phi$ .

**CONJECTURE 23.4** (*CY Langlands*; <sup>[CJ]</sup>). For each dimension  $d \in \{1, 2, 3\}$  there exists an involution  $\mathcal{C} \mapsto \mathcal{C}^L$  on smooth proper CY  $d$ -categories, the *Langlands dual*, such that:

- (i) The Verdier-dual chiral algebra  $\Phi(\mathcal{C})^\dagger := D_{\text{Ran}}(B(\Phi(\mathcal{C})))$  is quasi-isomorphic to  $\Phi(\mathcal{C}^L)$  (compatibility of the Langlands involution with the Verdier leg of Vol I Theorem A); equivalently, the modular tensor categories  $\Phi(\mathcal{C})$  and  $\Phi(\mathcal{C}^L)$  are related by the geometric Langlands equivalence of Frenkel–Gaitsgory (2006).
- (ii) The chiral modular characteristics  $\kappa_{\text{ch}}(\Phi(\mathcal{C}))$  and  $\kappa_{\text{ch}}(\Phi(\mathcal{C}^L))$  satisfy a family-dependent Koszul conductor relation. For input giving rise to Kac–Moody output, the conductor is  $\kappa_{\text{ch}} + \kappa'_{\text{ch}} = 0$  (the KM/free-field case of Volume I); for Virasoro-type output it is 13 (the self-dual point  $c = 13$ ). The general CY Koszul conductor  $\rho_K(\mathcal{C})$  is family-dependent: bare universality is forbidden.

**The case  $d = 1$ .** Smooth proper CY 1-categories are, up to Morita equivalence, the derived category of coherent sheaves on a smooth projective curve of genus  $g = 1$ . The functor  $\Phi$  on  $d = 1$  input is not covered by Theorem 8.1 (which is stated at  $d = 2$ ); the expected output is the Heisenberg-like current algebra of the elliptic curve. The involution  $C \mapsto C^L$  at  $d = 1$  is the identity. This is consistent with geometric Langlands on an elliptic curve being the Fourier–Mukai self-duality (Polishchuk–Rothstein, 2001).

**The case  $d = 2$ .** Smooth proper CY 2-categories include  $D^b(\text{Coh}(X))$  for  $X$  a K3 or abelian surface, and the Fukaya category of the same. The conjectured  $d = 2$  Langlands is the  $K3$  mirror map: Mukai’s Hodge-theoretic mirror  $X \leftrightarrow X^\vee$  for K3 surfaces should induce an involution on CY 2-categories which, via  $\Phi$ , corresponds to the  $d = 2$  Langlands equivalence. This is not proved. A first milestone is to verify Conjecture 23.4(i) on the sublocus of lattice-polarized K3 categories, where the mirror map is the Dolgachev–Nikulin involution on the Néron–Severi lattice.

**The case  $d = 3$ .** At  $d = 3$  the functor  $\Phi$  is itself programmatic; CY- $A_3$  is conditional on chain-level  $\mathbf{S}^3$ -framing. The Langlands conjecture is correspondingly the most speculative. Because BPS invariants are the primary output of the  $d = 3$  functor (via CoHA), the expected statement is that Langlands duality exchanges mirror BPS algebras. Concretely: for a CY 3-fold  $X$  with mirror  $X^\vee$ , the CoHA  $G(X)$  and  $G(X^\vee)$  should become Langlands dual as  $E_1$ -chiral algebras under  $\Phi$ . Every statement of this form passes through an unconstructed object and must remain a conjecture.

**The Hitchin realization at  $d = 2$ .** Let  $C$  be a smooth projective curve and  $\mathbf{G}$  a reductive group. The Hitchin moduli space  $\mathcal{M}_H(C, \mathbf{G})$  is a smooth hyperkähler manifold whose derived category of coherent sheaves  $D^b(\text{Coh}(\mathcal{M}_H(C, \mathbf{G})))$  is a CY 2-category (in the sense of the Mukai pairing on K3-type derived categories; the precise CY dimension depends on  $\dim \mathcal{M}_H$ ). The Hausel–Thaddeus and Donagi–Pantev mirror symmetry for Hitchin systems identifies

$$\mathcal{M}_H(C, \mathbf{G})^\vee \simeq \mathcal{M}_H(C, \mathbf{G}^L)$$

at the level of generic Hitchin fibers (dual abelian varieties) and, conjecturally, at the derived-categorical level via homological mirror symmetry:  $D^b(\text{Coh}(\mathcal{M}_H(\mathbf{G}))) \simeq \text{Fuk}(\mathcal{M}_H(\mathbf{G}^L))$ . Applying  $\Phi$  to both sides produces a pair of  $E_2$ -chiral algebras.

**CONJECTURE 23.5** (*Hitchin-system CY Langlands; [CJ]*). The CY Langlands involution of Conjecture 23.4, restricted to Hitchin-type CY 2-categories, is the Donagi–Pantev mirror map:  $\Phi(D^b(\text{Coh}(\mathcal{M}_H(\mathbf{G}))))^! \simeq \Phi(\text{Fuk}(\mathcal{M}_H(\mathbf{G}^L)))$  as  $E_2$ -chiral algebras on  $\text{Ran}(X)$ . The chiral modular characteristics satisfy  $\kappa_{\text{ch}}(\Phi(\mathcal{M}_H(\mathbf{G}))) + \kappa_{\text{ch}}(\Phi(\mathcal{M}_H(\mathbf{G}^L))) = 0$  (the KM-type Koszul conductor, not the Virasoro conductor 13). This is the  $d = 2$  Hitchin specialization; it implies (does not iff) the general  $d = 2$  CY Langlands.

**The Kapustin–Witten picture.** A physical interpretation of Conjecture 23.4 comes from the Kapustin–Witten realization of geometric Langlands as S-duality of  $\mathcal{N} = 4$  super-Yang–Mills on a product  $\Sigma \times C$  (Kapustin–Witten, 2007). In that picture, the Langlands dual group arises as the S-dual gauge group and the category of D-modules on  $\text{Bun}_G$  arises from the category of branes of an A-model or B-model twist. The CY category entering the construction is the Fukaya category of  $\text{Hit}(C, G)$  (the Hitchin moduli space) on one side and  $D^b(\text{Coh}(\text{Hit}(C, G^L)))$  on the other. The mirror symmetry of Hitchin moduli

spaces under hyperkähler rotation (Hausel–Thaddeus, 2003; Donagi–Pantev, 2012) provides a concrete CY 2-category instance where the  $d = 2$  Langlands involution of Conjecture 23.4 is testable. The  $\Phi$ -image on each side should recover, via Volume I, the spectral and automorphic sides of the Beilinson–Drinfeld construction.

### 23.3 Quantum geometric Langlands and the shadow parameter

Quantum geometric Langlands (QGL), introduced by Feigin–Frenkel and developed by Frenkel–Gaitsgory (2006), is a one-parameter deformation of classical geometric Langlands. The parameter  $\kappa_{\text{QGL}}$  is a complex number. The two limits recover known correspondences:

- $\kappa_{\text{QGL}} \rightarrow \infty$  (the critical level  $k = -h^\vee$ ): the classical geometric Langlands correspondence of Beilinson–Drinfeld, with Feigin–Frenkel centers as the spectral side.
- $\kappa_{\text{QGL}} \rightarrow 0$ : the classical de Rham geometric Langlands in Gaitsgory’s formulation.

At finite  $\kappa_{\text{QGL}}$  the spectral side is a category of *twisted  $D$ -modules* on  $\text{Bun}_G$  at level  $\kappa_{\text{QGL}}$ , and the automorphic side is the corresponding category at level  $-1/\kappa_{\text{QGL}}$  for the Langlands dual group. The quantum correspondence is stated as an equivalence of DG categories (Frenkel, *Lectures on Quantum Geometric Langlands*, 2005; Gaitsgory’s ongoing programme).

**The identification conjecture.** The parameter  $\kappa_{\text{QGL}}$  acts on the  $E_2$ -chiral algebra  $V_k(\mathfrak{g})$  via  $\kappa_{\text{QGL}} = k + h^\vee$ . Volume I defines a distinct parameter  $\kappa_{\text{ch}}$ , the chiral modular characteristic (Volume I, Theorem D), which for  $V_k(\mathfrak{g})$  evaluates to  $\kappa_{\text{ch}} = \dim(\mathfrak{g})(k + h^\vee)/(2h^\vee)$ . The two are proportional:

$$\kappa_{\text{ch}}(V_k(\mathfrak{g})) = \frac{\dim(\mathfrak{g})}{2h^\vee} \cdot \kappa_{\text{QGL}}.$$

At the critical level both vanish. This proportionality is already unconditional for  $V_k(\mathfrak{g})$ ; the conjectural extension is to general CY input.

**CONJECTURE 23.6** (*QGL parameter equals shadow obstruction; <sup>[CJ]</sup>*). For every CY category  $\mathcal{C}$  of dimension  $d = 2$ , the quantum geometric Langlands parameter of the Frenkel–Gaitsgory equivalence applied to  $\Phi(\mathcal{C})$  coincides, up to the dimensional normalization  $\dim(\mathfrak{g})/(2h^\vee)$  in the Kac–Moody case, with the Vol I chiral shadow parameter  $\kappa_{\text{ch}}(\Phi(\mathcal{C}))$  of the CY 2-category. Equivalently, the Frenkel–Gaitsgory QGL deformation is the shadow obstruction tower of Vol I (Volume I, Theorem D) evaluated along the critical-level family.

Consequence of Conjecture 23.6: the  $G/L/C/M$  shadow class of  $\Phi(\mathcal{C})$  controls the convergence of the QGL deformation series in  $\kappa_{\text{QGL}}$ . Class  $G$  (finite tower,  $r = 2$ ) would correspond to polynomial deformations; class  $L$  ( $r = 3$ , the Kac–Moody generic class) to rational-in- $\kappa_{\text{QGL}}$  deformations; class  $C$  ( $r = 4$ ,  $\beta\gamma$ -type) to deformations with finite transcendence degree; and class  $M$  (infinite tower, Virasoro and  $W_N$ ) to formal power series with genuine transcendental content. The critical-level limit  $\kappa_{\text{ch}} \rightarrow 0$  is then the degeneration point of Volume I at  $k = -h^\vee$ , consistent with Conjecture 23.2.

The conjecture implies (does not iff): the QGL duality exchanges  $(\kappa_{\text{QGL}}, G) \leftrightarrow (-1/\kappa_{\text{QGL}}, G^L)$ , which under Conjecture 23.6 should correspond to an involution on the space of CY 2-categories with a distinguished  $\mathfrak{g}$ -action. This is the  $d = 2$  specialization of Conjecture 23.4.

**CONJECTURE 23.7** (*Shadow convergence class determines QGL analytic type; <sup>[CJ]</sup>*). Let  $C$  be a smooth proper CY 2-category with  $\Phi(C)$  in shadow class  $X \in \{G, L, C, M\}$  (Volume I classification). The QGL deformation series of the Frenkel–Gaiitsgory equivalence applied to  $\Phi(C)$ , viewed as a formal power series in  $\kappa_{\text{QGL}}$ , has the following analytic type:

- (a) class G ( $r = 2$ ): the series is polynomial in  $\kappa_{\text{QGL}}$ , terminating at degree 1;
- (b) class L ( $r = 3$ ): the series is a rational function of  $\kappa_{\text{QGL}}$  with poles at roots of the Kac–Moody Kac determinant;
- (c) class C ( $r = 4$ ): the series converges in a disk  $|\kappa_{\text{QGL}}| < R_C$  with  $R_C$  determined by the  $\beta\gamma$ -type spectral radius;
- (d) class M ( $r = \infty$ ): the series is Gevrey-1 divergent and requires Borel resummation, with Stokes data encoding the shadow tower.

Conditional on Conjecture 23.6. At the critical level  $\kappa_{\text{QGL}} = 0$  all four types agree: the deformation is trivial and classical geometric Langlands holds.

**The Costello–Gaiotto perspective.** Costello and Gaiotto derive quantum geometric Langlands from a twisted compactification of 4d  $\mathcal{N} = 4$  super-Yang–Mills on a Riemann surface  $C$  (Costello, 2013; Costello–Gaiotto, 2020). In their framework the QGL parameter is the complexified gauge coupling  $\Psi = k + h^\vee$ , and the duality  $\Psi \leftrightarrow -1/\Psi$  is S-duality of the 4d theory. The chiral algebra on  $C$  is produced by the holomorphic-topological twist of Volume II; the bar complex of this chiral algebra (Volume I, Theorem A) encodes the factorization structure that Costello–Gaiotto identify with the Hecke action. For CY 2-category input, the expected statement is that the CY functor  $\Phi$  applied to the B-model category of the Hitchin system reproduces the Costello–Gaiotto chiral algebra at the corresponding value of  $\Psi$ . This identification is conditional on Conjecture 23.5 and the CY- $A_2$  functor (proved).

## 23.4 Open problems

The following five problems are the concrete frontier of CY Langlands. Each is stated as an open problem, not a conjecture, because the evidence is not yet sufficient to commit to a definite equivalence.

- (i) **K3 mirror map as  $d = 2$  Langlands.** Is the Mukai–Dolgachev–Nikulin mirror map on lattice-polarized K3 categories the  $d = 2$  Langlands involution of Conjecture 23.4? A first test: verify that  $\Phi$  sends mirror K3 pairs to Koszul-dual  $E_2$ -chiral algebras in the Volume I sense.
- (ii) **BPS Langlands for  $d = 3$ .** For a CY 3-fold  $X$  with mirror  $X^\vee$ , are the CoHAs  $G(X)$  and  $G(X^\vee)$  Langlands dual as  $E_1$ -chiral algebras? This passes through CY- $A_3$  and is therefore conditional on the  $\mathbb{S}^3$ -framing programme.
- (iii) **QGL parameter equals shadow  $\kappa_{\text{ch}}$ .** Prove Conjecture 23.6 for the Kac–Moody family as a theorem (it is currently only a proportionality of definitions), then extend to  $\Phi(C)$  for CY 2-categories  $C$ . The key step is to identify the shadow tower of  $\Phi(C)$  with the Gaiitsgory deformation at finite  $\kappa_{\text{QGL}}$ .
- (iv) **Categorified Feigin–Frenkel.** Does there exist a CY 2-category  $C_{\text{FF}}$  whose image  $\Phi(C_{\text{FF}})$  is, at the critical level, equivalent to the Feigin–Frenkel center  $\text{Fun}(\text{Op}_{GL}(D))$  viewed as a constant  $E_2$ -chiral algebra? If so,  $C_{\text{FF}}$  would be a geometric realization of opers at the categorical level.

- (v) **Geometric Satake from CY.** The geometric Satake equivalence (Mirković–Vilonen, 2007) identifies the category of perverse sheaves on the affine Grassmannian with  $\text{Rep}(G^L)$ . Can the equivalence be derived from a CY 3-category  $C_{\text{Sat}}$  via  $\Phi$ , so that  $\text{Rep}^{E_2}(\Phi(C_{\text{Sat}})) \simeq \text{Perv}(\text{Gr}_G)$ ? Supporting evidence: both sides carry braided monoidal structures; both admit fusion constructions; both are controlled by an  $\mathfrak{sl}_2$ -triple at the critical level.

Problems (i)–(iii) are the  $d = 2$  core and can in principle be attacked with the technology of Volume I and the  $d = 2$  functor of Theorem 8.1. Problems (iv)–(v) lie beyond the current reach: (iv) asks to reverse-engineer a CY 2-category from a target chiral algebra, and (v) asks to place the geometric Satake equivalence, which is the foundational theorem of the geometric Langlands programme, inside the CY–chiral dictionary. The payoff would be a derivation of local geometric Langlands from Calabi–Yau geometry alone.

**Status summary.** The material of this chapter is organized in three status tiers. The Feigin–Frenkel theorem (Theorem 23.1) is <sup>[PE]</sup>; it is cited from Feigin–Frenkel (1992) and Frenkel (2007). Conjecture 23.2 (critical-level Verdier-intertwining), Conjecture 23.4 (CY Langlands), Conjecture 23.5 (Hitchin-system specialization), Conjecture 23.6 (QGL parameter equals shadow), and Conjecture 23.7 (shadow convergence class determines QGL analytic type) are <sup>[CJ]</sup>; each is stated with “implies” rather than “iff” where the converse is not available. The five open problems of Section 23.4 are not conjectures: they are research questions whose expected answers the author cannot yet commit to. Downstream work should cite Theorem 23.1 freely, cite the conjectures only with the qualifier “conjecturally,” and refer to the open problems as motivation, not as established background.

**Dependencies on Volumes I and II.** The chapter uses four inputs from the earlier volumes. From Volume I: the bar functor  $B$  and the Verdier leg  $D_{\text{Ran}} \circ B$  of Theorem A (four-functor picture), the shadow tower  $\Theta_A$  and its G/L/C/M classification (Theorem D), the Koszul locus definition, and the Kac–Moody chiral modular characteristic  $\kappa_{\text{ch}}(V_k(\mathfrak{g})) = \dim(\mathfrak{g})(k + h^\vee)/(2h^\vee)$ . From Volume II: the  $E_1$ -chiral realization of CoHAs at  $d = 3$ , used in the BPS Langlands conjecture. Nothing in this chapter depends on the  $d = 3$  functor of Volume III being unconditional; every  $d = 3$  statement is explicitly tagged conditional on CY-A<sub>3</sub>.



# Appendix A

## Conventions

### A.1 Grading conventions

All grading is **cohomological**: differentials have degree  $+1$ . The bar construction uses desuspension (matching Volumes I–II).

### A.2 Sign conventions

Signs follow the Koszul rule throughout. For the curved  $A_\infty$ -structure:  $\mu_1^2(a) = [\mu_0, a]$  (commutator, minus sign).

### A.3 Categorical conventions

- $\mathbf{dgCat}$  denotes the  $\infty$ -category of small dg categories over  $k$ .
- $\mathbf{CY}_d\text{-Cat}$  denotes the  $\infty$ -category of smooth  $d$ -dimensional CY categories.
- Tensor products of dg categories are derived unless stated otherwise.
- “Equivalence” means quasi-equivalence of dg categories (equivalence of associated  $\infty$ -categories).

### A.4 Operadic conventions

- $E_n$  denotes the little  $n$ -disks operad (Boardman–Vogt).
- $\overline{\mathbf{FM}}_k(\mathbb{C})$  denotes the Fulton–MacPherson compactification of  $\mathbf{Conf}_k(\mathbb{C})$ . Note:  $\overline{\mathbf{FM}}$  is a blowup along diagonals, NOT the complement of the diagonal (AP, Vol I).
- $\mathbf{SC}^{\text{ch, top}}$  denotes the Swiss-cheese operad from Volume II.

### A.5 Bar complex and Koszul duality conventions

- $B(A)$  is the bar complex, a factorization *coalgebra* on  $\mathbf{Ran}(X)$ .

- $\Omega(B(A)) \simeq A$  is bar-cobar *inversion*: it recovers the original algebra (Vol I, Theorem B). This is NOT Koszul duality.
- $A^\dagger = (H^*(B(A)))^\vee$  is the Koszul dual *algebra* (linear dual of bar cohomology).
- $D_{\text{Ran}}(B(A)) \simeq B(A^\dagger)$  is the Verdier dual (Vol I, Theorem A, Convention conv:bar-coalgebra-identity).
- The collision  $r$ -matrix  $r(z) = \text{Res}_{0,2}^{\text{coll}}(\Theta_A)$  has pole orders *one less* than the OPE, because the bar construction extracts residues along  $d \log(z_i - z_j)$ , which absorbs one power of  $(z - w)$  (Vol I).



# Bibliography

- [1] A. Beilinson and V. Drinfeld, *Chiral Algebras*, American Mathematical Society Colloquium Publications, vol. 51, American Mathematical Society, Providence, RI, 2004.
- [2] R.-O. Buchweitz, G.-M. Greuel, and F.-O. Schreyer, *Cohen–Macaulay modules on hypersurface singularities II*, *Invent. Math.* **88** (1987), no. 1, 165–182.
- [3] A. Căldăraru, *The Mukai pairing, II: the Hochschild–Kostant–Rosenberg isomorphism*, *Adv. Math.* **194** (2005), no. 1, 34–66, arXiv:math/0308080.
- [4] S. Cecotti and C. Vafa, *On classification of  $N = 2$  supersymmetric theories*, *Comm. Math. Phys.* **158** (1993), no. 3, 569–644, arXiv:hep-th/9211097.
- [5] K. Costello, *Holography and Koszul duality: the example of the M2 brane*, preprint, 2017, arXiv:1705.02500.
- [6] K. Costello, *Topological conformal field theories and Calabi–Yau categories*, *Adv. Math.* **210** (2007), no. 1, 165–214, arXiv:math/0412149.
- [7] K. Costello, *Topological conformal field theories and Calabi–Yau categories*, *Adv. Math.* **210** (2007), no. 1, 165–214, arXiv:math/0412149.
- [8] K. Costello, *Topological conformal field theories and Calabi–Yau categories*, *Adv. Math.* **210** (2007), no. 1, 165–214, arXiv:math/0412149.
- [9] K. Costello and O. Gwilliam, *Factorization Algebras in Quantum Field Theory, Volume 1*, New Mathematical Monographs, vol. 31, Cambridge University Press, Cambridge, 2017; *Factorization Algebras in Quantum Field Theory, Volume 2*, New Mathematical Monographs, vol. 41, Cambridge University Press, Cambridge, 2021.
- [10] J. Ding and K. Iohara, *Generalization of Drinfeld quantum affine algebras*, *Lett. Math. Phys.* **41** (1997), no. 2, 181–193.
- [11] R. Dijkgraaf, G. Moore, E. Verlinde, and H. Verlinde, *Elliptic genera of symmetric products and second quantized strings*, *Comm. Math. Phys.* **185** (1997), 197–209, arXiv:hep-th/9608096.
- [12] T. Dyckerhoff, *Compact generators in categories of matrix factorizations*, *Duke Math. J.* **159** (2011), no. 2, 223–274, arXiv:0904.4713.
- [13] D. Eisenbud, *Homological algebra on a complete intersection, with an application to group representations*, *Trans. Amer. Math. Soc.* **260** (1980), no. 1, 35–64.

- [14] J. D. Fay, *Theta Functions on Riemann Surfaces*, Lecture Notes in Mathematics, vol. 352, Springer-Verlag, Berlin-New York, 1973.
- [15] E. Frenkel and D. Ben-Zvi, *Vertex Algebras and Algebraic Curves*, Mathematical Surveys and Monographs, vol. 88, American Mathematical Society, Providence, RI, 2004.
- [16] T. Gannon, *Much ado about Mathieu*, Adv. Math. **301** (2016), 322–358, arXiv:1211.5531.
- [17] A. Kapustin and Y. Li, *D-branes in Landau–Ginzburg models and algebraic geometry*, J. High Energy Phys. **2003** (2003), no. 12, 005, arXiv:hep-th/0210296.
- [18] B. Keller, *Introduction to A-infinity algebras and modules*, Homology Homotopy Appl. **3** (2001), no. 1, 1–35.
- [19] B. Keller, *On differential graded categories*, in *International Congress of Mathematicians, Vol. II*, 151–190, European Mathematical Society, Zürich, 2006, arXiv:math/0601185.
- [20] H. Knörrer, *Cohen–Macaulay modules on hypersurface singularities I*, Invent. Math. **88** (1987), no. 1, 153–164.
- [21] M. Kontsevich, *Homological algebra of mirror symmetry*, in *Proceedings of the International Congress of Mathematicians, Vol. I, II (Zürich, 1994)*, 120–139, Birkhäuser, Basel, 1995, arXiv:alg-geom/9411018.
- [22] M. Kontsevich, *Feynman diagrams and low-dimensional topology*, in *First European Congress of Mathematics, Vol. II (Paris, 1992)*, Progr. Math., vol. 120, 97–121, Birkhäuser, Basel, 1994.
- [23] M. Kontsevich and Y. Soibelman, *Notes on A-infinity algebras, A-infinity categories and non-commutative geometry. I*, in *Homological Mirror Symmetry*, Lecture Notes in Phys., vol. 757, 153–219, Springer, Berlin, 2009, arXiv:math/0606241.
- [24] M. Kontsevich and Y. Soibelman, *Deformations of algebras over operads and the Deligne conjecture*, in *Conférence Moshé Flato 1999, Vol. I (Dijon)*, Math. Phys. Stud., vol. 21, 255–307, Kluwer Acad. Publ., Dordrecht, 2000, arXiv:math/0001151.
- [25] J. Lurie, *Higher Algebra*, manuscript, 2017, available at <https://www.math.ias.edu/~lurie/papers/HA.pdf>.
- [26] K. Miki, *A  $(q, \gamma)$  analog of the  $W_{1+\infty}$  algebra*, J. Math. Phys. **48** (2007), no. 12, 123520.
- [27] C.-P. Mok, *Logarithmic Fulton–MacPherson configuration spaces*, arXiv:2503.17563, 2025.
- [28] S. Mukai, *On the moduli space of bundles on K3 surfaces. I*, in *Vector Bundles on Algebraic Varieties (Bombay, 1984)*, Tata Inst. Fund. Res. Stud. Math., vol. 11, 341–413, Tata Inst. Fund. Res., Bombay, 1987.
- [29] D. Orlov, *Triangulated categories of singularities and equivalences between Landau–Ginzburg models*, Sb. Math. **197** (2006), no. 12, 1827–1840, arXiv:math/0503630.
- [30] A. Polishchuk,  *$A_\infty$ -algebra of an elliptic curve and Eisenstein series*, Comm. Math. Phys. **301** (2011), no. 3, 709–722, arXiv:0907.5484.

- [31] A. Polishchuk and A. Vaintrob, *Chern characters and Hirzebruch–Riemann–Roch formula for matrix factorizations*, Duke Math. J. **161** (2012), no. 10, 1863–1926, arXiv:1002.2116.
- [32] P. Seidel, *Fukaya Categories and Picard–Lefschetz Theory*, Zurich Lectures in Advanced Mathematics, European Mathematical Society, Zürich, 2008.
- [33] N. Sheridan, *Homological mirror symmetry for Calabi–Yau hypersurfaces in projective space*, Invent. Math. **199** (2015), no. 1, 1–186, arXiv:1111.0632.
- [34] J. D. Stasheff, *Homotopy associativity of H-spaces. I, II*, Trans. Amer. Math. Soc. **108** (1963), 275–292; 293–312.
- [35] O. Schiffmann and E. Vasserot, *Cherednik algebras, W-algebras and the equivariant cohomology of the moduli space of instantons on  $\mathbb{A}^2$* , Publ. Math. IHÉS **118** (2013), 213–342, arXiv:1202.2756.
- [36] D. Tamarkin, *What do dg-categories form?*, Compos. Math. **143** (2007), no. 5, 1335–1358, arXiv:math/0606553.
- [37] R. Lorgat,  *$A_\infty$  Chiral Algebras and 3D Holomorphic-Topological QFT*, Modular Koszul Duality Programme, Vol. II, in preparation, 2026.
- [38] E. Witten, *Phases of  $N = 2$  theories in two dimensions*, Nuclear Phys. B **403** (1993), no. 1–2, 159–222, arXiv:hep-th/9301042.
- [39] Y. Zhu, *Modular invariance of characters of vertex operator algebras*, J. Amer. Math. Soc. **9** (1996), no. 1, 237–302.