1. The standard landscape

The bar construction, the genus tower, the Swiss-cheese structure, and the collision residue are not abstractions awaiting examples: they were *found* by computing examples. The standard landscape is the testing ground where the theory proves itself.

1.1. **Census.** Seven families span the landscape. Each is chirally Koszul. The invariants computed in the monograph are recorded in Table **??**.

Table 1. The standard landscape: seven families and their bar-complex invariants

Algebra $\mathcal{A}$	$c(\mathcal{A})$	$\kappa(\mathcal{A})$	$r_{\max}$	**Class**	$r(z)$	$\mathcal{A}^!$
$\mathcal{H}_k$	1	k	2	**G**	k/z	$\mathrm{Sym}^{\mathrm{ch}}(V^*)$
$\widehat{\mathfrak{g}}_k$	$\dfrac{kd}{k+h^\vee}$	$\dfrac{d(k+h^\vee)}{2h^\vee}$	3	**L**	$\Omega/((k{+}h^\vee)z)$	$\widehat{\mathfrak{g}}_{-k-2h^\vee}$
$\beta\gamma_\lambda$	$1-3(2\lambda{-}1)^2$	$c/2$	4	**C**	const.	bc_λ
Vir_c	c	$c/2$	∞	**M**	$\frac{c/2}{z^3}+\frac{2T}{z}$	Vir_{26-c}
${}^k_3(\mathfrak{sl}_3)$	$c(k)$	$5c/6$	∞	**M**	$r_{\,3}(z)$	${}^{k'}_{3}$
${}^k_N(\mathfrak{sl}_N)$	$c(k)$	$c\cdot(H_N{-}1)$	∞	**M**	$r_{\,N}(z)$	${}^{k'}_{N}$
V_Λ (lattice)	$\mathrm{rank}(\Lambda)$	$\mathrm{rank}(\Lambda)$	2	**G**	rank/z	V_{Λ^*}

Here $d=\dim\mathfrak{g}$, $h^\vee$ is the dual Coxeter number, $t=k+h^\vee$, $k'=-k-2h^\vee$, $H_N=\sum_{j=1}^N 1/j$ is the N-th harmonic number, and $\Omega=\sum_a J^a\otimes J_a$ is the Casimir tensor.

The table has two readings. Read horizontally, each row is a complete portrait: central charge, modular characteristic, shadow depth, collision residue, Koszul dual. Read vertically, the columns expose the structural regularities that no single example can reveal. The modular characteristic κ is always a rational function of the level; the complementarity sum $\kappa+\kappa^!$ vanishes for free-field and Kac–Moody algebras, equals 13 for Virasoro, and is $(c+c')(H_N-1)$ for ${}_N$. The collision residue $r(z)$ is scalar for classes **G** and **C**, matrix-valued for classes **L** and **M**.

1.2. **The DS reduction chain.** Drinfeld–Sokolov reduction links the families. For $\mathfrak{g}=\mathfrak{sl}_N$, the principal DS reduction

$$V_k(\mathfrak{sl}_N)\xrightarrow{\mathrm{DS}}{}^k_N(\mathfrak{sl}_N) \tag{1.1}$$

transforms the shadow obstruction tower in a controlled way:

(i) *Depth increase.* The affine algebra $V_k(\mathfrak{sl}_N)$ is class $\mathbf{L}$ ($r_{\max} = 3$); the -algebra ^k_N is class $\mathbf{M}$ ($r_{\max} = \infty$). DS reduction creates the infinite shadow tower from finite data.

(ii) *Quartic creation from zero.* The quartic shadow S_4 vanishes for all affine algebras (the Jacobi identity kills the obstruction); after DS reduction, $S_4(^k_N) \neq 0$ for generic k. For Virasoro: $S_4(\mathrm{Vir}_c) = 10/[c(5c+22)]$.

(iii) *Central charge is additive; κ is not.* The BRST ghost system contributes $c_{\mathrm{ghost}} = N(N-1)$ to the central charge but a k-dependent correction to κ. The commutation failure of DS reduction with the shadow map is the source of all non-formality in the -algebra landscape.

The chain $\mathcal{H} \to \widehat{\mathfrak{g}}_k \to^k_N$ is a cascade of increasing shadow complexity: $\mathbf{G} \to \mathbf{L} \to \mathbf{M}$, with each reduction introducing new structure that the previous level could not see.

1.3. **The multi-weight correction.** For algebras with generators of distinct conformal weights, the genus expansion receives a correction. The $_3$ algebra has generators T (weight 2) and W (weight 3); at genus 2 the cross-channel graph sum contributes

$$\delta F_2(_3) \;=\; \frac{c+204}{16c} \;>\; 0 \qquad \text{for all } c > 0\,. \tag{1.2}$$

The correction arises from boundary graphs of $\overline{\mathcal{M}}_{2,0}$ whose edges carry mixed propagator channels (weight-2 and weight-3 generators sewed through different Hodge data at the vertex level). The propagator itself is always weight-1 at the edge level ($d\log E$ has conformal weight 1); the mixing enters through the vertex structure constants.

For uniform-weight algebras (all generators of the same conformal weight), $\delta F_g^{\mathrm{cross}} = 0$ at all genera and the scalar universality $F_g = \kappa \cdot \lambda_g^{\mathrm{FP}}$ holds without correction. For multi-weight algebras, the full genus expansion is $F_g = \kappa \cdot \lambda_g^{\mathrm{FP}} + \delta F_g^{\mathrm{cross}}$, with $\delta F_g^{\mathrm{cross}}$ computable from the mixed-channel graph sum.

2. The MC programme

The MC programme converts the geometric structure of the bar complex into five concrete mathematical problems, each asking whether a structural property of the universal MC element $\Theta_{\mathcal{A}}$ persists under a specific operation: PBW concentration, bar-intrinsic existence, categorical generation, inverse-limit completion, and analytic sewing. MC1 through MC4 are proved; MC5 is partially proved, with the analytic HS-sewing package established at all genera and the genuswise BV/BRST/bar identification conjectural.

2.1. **MC1: PBW concentration.** The PBW spectral sequence of $B(\mathcal{A})$ collapses at E_2 for every freely strongly generated chiral algebra. Free strong generation implies the PBW filtration on the bar complex is exhaustive, and

the associated graded is a cofree coalgebra whose cohomology concentrates in bar degree 1. This is chirally Koszul: the bar cohomology $H^*(B(\mathcal{A}))$ is the Koszul dual $\mathcal{A}^!$ (as a graded coalgebra).

The proof applies to the universal -algebra ${}^k(\mathfrak{g})$ at all levels k (Feigin–Frenkel free generation). The simple quotient $L_k(\mathfrak{g})$ at admissible levels has Koszulness proved for $\mathfrak{g} = \mathfrak{sl}_2$ at all admissible levels; higher rank is open.

2.2. **MC2: bar-intrinsic existence.** The universal MC element $\Theta_{\mathcal{A}} = D_{\mathcal{A}} - d_0$ exists for all modular Koszul chiral algebras, with no restriction to simple Lie symmetry. The proof is a tautology at the right level of abstraction: $D_{\mathcal{A}}^2 = 0$ implies $\Theta_{\mathcal{A}}$ satisfies the MC equation (**??**). The algebraic-family rigidity theorem extends the scalar identification $\Theta_{\mathcal{A}}^{\min} = \eta \otimes \Gamma_{\mathcal{A}}$ to all algebraic families with rational OPE coefficients, covering the entire standard Lie-theoretic landscape at all non-critical levels.

2.3. **MC3: categorical generation.** For every simple Lie algebra $\mathfrak{g}$, the Drinfeld–Kazhdan category $\mathrm{DK}(\mathfrak{g})$ is thickly generated by evaluation modules. The proof combines the chromatic filtration on $\mathrm{Rep}(Y(\mathfrak{g}))$ with the categorical Clebsch–Gordan closure via multiplicity-free ℓ-weights (Chari–Moura), the Francis–Gaitsgory pro-nilpotent completion, and Drinfeld–Kazhdan compactness. The type A proof uses the minuscule hypothesis; the all-types generalization replaces it with the strictly weaker multiplicity-free ℓ-weight property, which holds for all simple types.

2.4. **MC4: inverse-limit completion.** The bar-cobar adjunction extends to inverse limits via the strong completion-tower theorem: the strong filtration axiom $\mu_r(F^{i_1}, \dots, F^{i_r}) \subset F^{i_1+\cdots+i_r}$ forces a degree cutoff that makes continuity and the Mittag-Leffler condition automatic. The completion closure $\mathrm{CompCl}(F_{\mathrm{ft}})$ carries a quasi-inverse bar-cobar homotopy equivalence, stable under MC twisting.

The programme splits: $\mathrm{MC4}^+$ (positive towers: ${}_{1+\infty}$, affine Yangians) is solved by weight stabilization. $\mathrm{MC4}^0$ (resonant towers: Virasoro, non-quadratic ${}_N$) is reduced to a finite resonance problem by the resonance-filtered bar-cobar theorem.

2.5. **MC5: analytic sewing.** The sewing envelope $\mathcal{A}^{\mathrm{sew}}$, the Hausdorff completion of $\mathcal{A}$ for the all-amplitude seminorm topology, carries the analytic data needed for genus-g partition functions. The HS-sewing criterion provides a sufficient condition: polynomial OPE growth and subexponential sector growth imply convergence at all genera. Every standard family satisfies this condition.

For Heisenberg, the sewing theorem is explicit: the one-particle Bergman reduction writes the genus-g partition function as a Fredholm determinant. For general modular Koszul algebras, the HS-sewing criterion gives convergence without an explicit closed form. The BV/BRST identification of sewing amplitudes with bar complex data at higher genus remains conjectural.

3. Arithmetic

The shadow obstruction tower produces arithmetic invariants from vertex-algebraic data. These invariants connect the bar complex to number theory through three structures: a Dirichlet series, a categorical zeta function, and a depth decomposition.

3.1. **The shadow Eisenstein theorem.** The genus-1 amplitude of a modular Koszul chiral algebra $\mathcal{A}$ is the quasi-modular Eisenstein series $\kappa(\mathcal{A})\cdot E_2^*(\tau)$, where $E_2^*(\tau) = E_2(\tau) - 3/(\pi \operatorname{Im}\tau)$ is the completed weight-2 Eisenstein series. The Fourier coefficients $\sigma_1(n)$ of E_2^* produce a Dirichlet series:

$$D_2(\mathcal{A}, s) \;:=\; -24\kappa(\mathcal{A}) \cdot \zeta(s) \cdot \zeta(s-1)\,, \tag{3.1}$$

an Eisenstein L-function with poles at $s = 1$ (from $\zeta(s)$) and $s = 2$ (from $\zeta(s-1)$). The identity follows from the σ_1 Dirichlet convolution applied to the Fourier coefficients of $\kappa \cdot E_2^*(\tau)$.

The genus-1 propagator is E_2^*, not a holomorphic modular form: E_2^* is quasi-modular of weight 2 and depth 1, transforming with an additive anomaly $3\tau/(\pi i)$ under $\mathrm{SL}(2,\mathbb{Z})$. Products of E_2^* live in the ring of quasi-modular forms $\mathbb{C}[E_2^*, E_4, E_6]$, not in the ring of holomorphic modular forms $\mathbb{C}[E_4, E_6]$. This distinction is load-bearing: holomorphic dimension formulas (e.g., $\dim S_k = 0$ for $k < 12$) do not apply to quasi-modular objects.

3.2. **The categorical zeta function.** The dimension-counting zeta function of $\mathfrak{sl}_2$-representations is

$$\zeta^{\mathrm{DK}}_{\mathfrak{sl}_2}(s) \;:=\; \sum_{n\geq 1} (\dim V_n)^{-s} \;=\; \sum_{n\geq 1} (n+1)^{-s} \;=\; \zeta(s) - 1\,. \tag{3.2}$$

The Riemann zeta function (shifted by 1) counts $\mathfrak{sl}_2$-representations weighted by inverse dimension. The link to the shadow–DK bridge is that the categorical zeta encodes the multiplicity structure of the evaluation modules whose thick generation proves MC3.

For $\mathfrak{sl}_N$ with $N \geq 3$, $\zeta^{\mathrm{DK}}_{\mathfrak{sl}_N}(s) := \sum_\lambda (\dim V_\lambda)^{-s}$ is not multiplicative: the Weyl dimension formula is a product of linear forms, but the sum over the dominant chamber has no Euler product. The $N = 2$ coincidence with $\zeta(s)$ is a low-rank accident.

3.3. **Depth decomposition.** The total shadow depth decomposes:

$$d(\mathcal{A}) \;=\; 1 + d_{\mathrm{arith}}(\mathcal{A}) + d_{\mathrm{alg}}(\mathcal{A})\,, \tag{3.3}$$

where d_{arith} counts arithmetic contributions (cusp forms in the automorphic decomposition of the shadow generating function) and d_{alg} counts algebraic contributions (non-formality of the L_∞ structure). The algebraic depth $d_{\mathrm{alg}} \in \{0, 1, 2, \infty\}$ recovers the G/L/C/M classification: $d_{\mathrm{alg}} = 0$ for class **G**, $d_{\mathrm{alg}} = 1$ for class **L**, $d_{\mathrm{alg}} = 2$ for class **C**, $d_{\mathrm{alg}} = \infty$ for class **M**. Shadow depths ≥ 5 are purely arithmetic: they arise from cusp forms, beginning with the weight-12 cusp form Δ_{12} at the genus-2 level.

The decomposition separates the algebraic content (governed by the MC equation on the cyclic deformation complex) from the arithmetic content (governed by automorphic forms on moduli of curves). The two are independent: a class-**G** algebra (algebraically trivial) can have nontrivial arithmetic depth, and a class-**M** algebra (algebraically infinite) can have vanishing arithmetic depth at low genus.

4. Frontier directions

Three leaps organize the frontier. The first extends the modular Koszul programme from chiral algebras to holographic systems. The second extends it from formal algebraic data to analytic sewing amplitudes. The third extends it from 2-dimensional geometry to 3-dimensional topology.

4.1. **First leap: holographic modular Koszul data.** Every holomorphic-topological system T in the sense of Costello–Li should be controlled by a *holographic modular Koszul datum*

$$\mathcal{H}(T) \;=\; (\mathcal{A},\, \mathcal{A}^{!},\, \mathfrak{C},\, r(z),\, \Theta_{\mathcal{A}},\, \nabla^{\mathrm{hol}})\,, \tag{4.1}$$

consisting of the boundary algebra, its Koszul dual, the bulk algebra (chiral derived centre), the collision residue, the universal MC element, and the holographic shadow connection. The collision residue is the genus-0 binary shadow of $\Theta_{\mathcal{A}}$: $r(z) = \mathrm{Res}^{\mathrm{coll}}_{0,2}(\Theta_{\mathcal{A}})$.

This is the Platonic holographic programme. Its five theorem targets are: boundary-defect realization (the boundary algebra is the algebra of boundary conditions of the 3d theory), Yangian-shadow identification (the Yangian controls line operators), sphere reconstruction (genus-0 amplitudes recover the boundary OPE), quartic resonance obstruction (the quartic shadow controls the deformation quantization), and singular-fibre descent (the critical-level limit produces the Feigin–Frenkel centre).

Two other directions in the first leap: the factorization-envelope technology of Nishinaka and Vicedo, which gives a functorial pipeline from Lie conformal algebras to vertex algebras; and the non-principal -algebra corridor, where hook-type nilpotent orbits in type A provide the first proved non-principal Koszul dualities.

4.2. **Second leap: analytic sewing.** The algebraic bar construction produces formal power series in the sewing parameters. The analytic sewing programme asks: when do these series converge?

The answer is the sewing envelope $\mathcal{A}^{\mathrm{sew}}$, the Hausdorff completion for the all-amplitude seminorm topology. The HS-sewing condition (polynomial OPE growth and subexponential sector growth) is satisfied by the entire standard landscape. The Heisenberg sewing theorem gives an explicit Fredholm determinant. The lattice sewing envelope is constructed; the analytic realization criterion (a sufficient condition for passage from formal to analytic) is conjectural.

At genus $g \geq 1$ the curvature (**??**) forces passage from ordinary to coderived categories: the bar complex is no longer flat, and the correct homological framework is the coderived category of curved coalgebras. The analytic programme at higher genus requires the coderived analytic shadow $Q_g^{\mathrm{an}}(\mathcal{A})$.

4.3. **Third leap: three-dimensional topology.** The ordered bar complex $\overline{B}^{\mathrm{ord}}(\mathcal{A})$ is an E_1 chiral coassociative coalgebra over $(\mathrm{ChirAss})^!$: the differential encodes holomorphic factorization on $\mathbb{C}$, the deconcatenation coproduct encodes topological factorization on $\mathbb{R}$. The two-coloured Swiss-cheese operad $\mathrm{SC}^{\mathrm{ch,top}}$ of a 3-dimensional holomorphic-topological theory emerges on the chiral derived center pair $(C^\bullet_{\mathrm{ch}}(\mathcal{A},\mathcal{A}),\, \mathcal{A})$, not on $\overline{B}^{\mathrm{ord}}(\mathcal{A})$ itself.

The third leap is the passage from the algebraic Swiss-cheese structure (proved) to the geometric realization on 3-manifolds of the form $\Sigma_g \times [0,1]$. The $\mathbb{C}$-direction carries the chiral algebra; the $\mathbb{R}$-direction carries the Yangian / quantum group. The annulus trace $\Delta_{\mathrm{ns}}(\mathrm{Tr}_{\mathcal{A}}) = \kappa \cdot \lambda_1$ provides the first open-to-closed map at genus 1; the full genus tower progressively restores bidirectionality between open and closed sectors.

The deepest open problem in this direction is CY-A at $d = 3$ (Section **??**).

5. The single open problem

Everything in Sections **??**–**??** rests on two-dimensional geometry: curves, their moduli spaces, and the factorization structure of colliding points on a curve. The three-dimensional extension is conditional on a single unresolved input.

Conjecture 5.1 (CY-A at $d = 3$)**.** *The Calabi–Yau/A_∞ correspondence extends from $d = 2$ (proved) to $d = 3$: the chain-level S^3-framing of the topological direction in a 3-dimensional holomorphic-topological theory produces an A_∞-structure on the boundary algebra that is compatible with the chiral bar complex.*

At $d = 2$, the CY-A correspondence is a theorem: the Fukaya category of a symplectic surface carries an A_∞-structure whose bar construction is the wrapped Floer complex, and the A_∞-formality on the Koszul locus is the E_2-collapse of the bar spectral sequence. At $d = 3$, the analogue requires the chain-level framing of the S^3-fibre in the conifold geometry to be compatible with the factorization structure on the $\mathbb{C}$-direction. This is a statement about the interaction of holomorphic and topological data at the chain level, not merely at the level of cohomology.

The difficulty is not conceptual but technical: the $d = 2$ proof uses the Riemann surface structure in an essential way (the Abel–Jacobi map, the period matrix, the prime form), and the $d = 3$ analogue must replace surface geometry with 3-manifold geometry while preserving the A_∞-coherences.

No other open conjecture in the programme has unresolved algebraic prerequisites. The five MC problems are solved; the Koszulness characterization is complete to within the D-module purity converse; the shadow

obstruction tower is constructed at all degrees; the arithmetic invariants are computed; the standard landscape is fully mapped. CY-A at $d = 3$ is the single point where the programme reaches beyond what it can currently prove, and it is the natural boundary: the passage from the algebraic curve to the three-manifold.

References

References

[1] V. I. Arnold, The cohomology ring of the group of dyed braids, *Mat. Zametki* **5** (1969), 227–231.

[2] A. Beilinson and V. Drinfeld, *Chiral Algebras*, AMS Colloquium Publications, vol. 51, American Mathematical Society, Providence, RI, 2004.

[3] A. Beilinson, V. Ginzburg, and W. Soergel, Koszul duality patterns in representation theory, *J. Amer. Math. Soc.* **9** (1996), 473–527.

[4] K. Costello and O. Gwilliam, *Factorization Algebras in Quantum Field Theory*, vols. 1–2, Cambridge University Press, 2017/2021.

[5] K. Costello and S. Li, Twisted supergravity and its quantization, preprint, `arXiv:1606.00365`, 2016.

[6] V. Chari and A. Moura, Characters and blocks for finite-dimensional representations of quantum affine algebras, *Int. Math. Res. Not.* (2004), no. 5, 257–298.

[7] V. G. Drinfeld and V. G. Kazhdan, Algebraic geometry of the space of quasi-maps and the structure of representations of quantum groups, unpublished notes, 1984.

[8] V. G. Drinfeld, Hopf algebras and the quantum Yang–Baxter equation, *Dokl. Akad. Nauk SSSR* **283** (1985), 1060–1064.

[9] E. Frenkel and D. Ben-Zvi, *Vertex Algebras and Algebraic Curves*, 2nd ed., Mathematical Surveys and Monographs, vol. 88, American Mathematical Society, Providence, RI, 2004.

[10] B. Feigin and E. Frenkel, Affine Kac–Moody algebras at the critical level and Gel'fand–Dikiĭ algebras, *Int. J. Mod. Phys. A* **7** (suppl. 1A) (1992), 197–215.

[11] J. Francis and D. Gaitsgory, Chiral Koszul duality, *Selecta Math. (N.S.)* **18** (2012), 27–87.

[12] W. Fulton and R. MacPherson, A compactification of configuration spaces, *Ann. of Math. (2)* **139** (1994), 183–225.

[13] C. Faber and R. Pandharipande, Hodge integrals and moduli spaces of curves, *Enseign. Math. (2)* **48** (2002), 173–195.

[14] D. Gaiotto, Twisted compactifications of 3d $\mathcal{N} = 4$ theories and conformal blocks, *J. High Energy Phys.* (2019), no. 2, 61.

[15] D. Gaiotto and J. Lamy-Poirier, Holomorphic-topological twists, unpublished notes, 2019.

[16] V. Kac, S.-S. Roan, and M. Wakimoto, Quantum reduction for affine superalgebras, *Comm. Math. Phys.* **241** (2003), 307–342.

[17] J.-L. Loday and B. Vallette, *Algebraic Operads*, Grundlehren der Mathematischen Wissenschaften, vol. 346, Springer, Berlin, 2012.

[18] C.-P. Mok, *Logarithmic Fulton–MacPherson configuration spaces*, arXiv:2503.17563, 2025.

[19] Y. Moriwaki, Conformally flat factorization homology in IndHilb, preprint, 2026.

[20] N. Markarian and H. L. Tanaka, Homotopy chiral algebras, preprint, 2024.

[21] D. Nishinaka, Factorization envelopes of Lie conformal algebras, preprint, 2025.

[22] S. Priddy, Koszul resolutions, *Trans. Amer. Math. Soc.* **152** (1970), 39–60.

[23] T. Pantev, B. Toën, M. Vaquié, and G. Vezzosi, Shifted symplectic structures, *Publ. Math. Inst. Hautes Études Sci.* **117** (2013), 271–328.

[24] D. Robert-Nicoud and F. Wierstra, Homotopy morphisms between convolution homotopy Lie algebras, *J. Noncommut. Geom.* **13** (2019), 1463–1520.
[25] B. Vallette, Homotopy theory of homotopy algebras, *Ann. Inst. Fourier* **70** (2020), 683–738.
[26] B. Vicedo, Non-chiral factorization algebras and vertex algebras, preprint, 2025.
[27] M. Wakimoto and V. Kac, Quantum reduction and representation theory of superconformal algebras, *Adv. Math.* **185** (2004), 400–458.

Modular Koszul duality for chiral algebras is E_1–E_1 operadic Koszul duality transported into the homotopical modular chiral realm on algebraic curves.