

The five theorems of modular Koszul duality

Raez Lorgat

Abstract

Koszul duality over a point terminates: a quadratic algebra determines a dual coalgebra, the bar construction mediates between them, and no further structure appears. Over an algebraic curve, the theory does not terminate. The bar differential acquires curvature from the Hodge bundle on moduli space; its failure to square to zero is controlled by a single scalar $\kappa(\mathcal{A})$, the modular characteristic, extracted from the collision residue of the universal Maurer–Cartan element.

The chiral algebra $\mathcal{A}$ is the level-1 chart $A_b = \text{End}_C(b)$ of a factorization dg-category C^{op} on a tangential log curve (X, D, τ) , evaluated at a vacuum object b . Five theorems characterize the levels of the resulting Beilinson tower

$$C^{\text{op}} \xrightarrow{b} A_b \xrightarrow{\bar{B}} B(A_b) \xrightarrow{Z_{\text{ch}}^{\text{der}}} \text{bulk} \rightarrow \text{line ops} \xrightarrow{\text{tr}} \text{modular shadow}.$$

Each theorem carries a type signature (*quadrant, presentation, level, hypothesis package*).

Theorem A is a chiral Koszul reflection $K^2 \simeq \text{id}$ on $\text{Kosz}(X)$ at the chart-bar boundary $1 \leftrightarrow 2$, under the augmented + complete + finite-dim graded hypothesis package $(H_1) - (H_3)$, in three parametric strengths: *weak* (bar–cobar adjunction unconditional), *strong* ($K^2 \simeq \text{id}$ on the Koszul locus), *strongest-rejected* (an unconditional equivalence on Alg^{aug} off the Koszul locus, false because L_o -witnesses kill conilpotency outside the weight-completed ambient). Theorem B inscribes bar–cobar inversion $\Omega_X(\bar{B}_X(C)) \xrightarrow{\sim} C$ in $D_{\text{ch}}^{\text{co}}(X)$ at level $1 \rightarrow 3$, on the Koszul locus, with the class-M chain-level statement ambient-qualified to the weight-completed coderived target. Theorem C is the 5×5 κ -stratification matrix on rows G/L/C/M/B with columns $\{\kappa_{\text{cat}}, \kappa_{\text{ch}}^{\text{Hodge}}, \kappa_{\text{ch}}^{\text{Heis}}, \kappa_{\text{BKM}}, \kappa_{\text{fiber}}\}$ at level 5 per stratum; $K_3 \times E$ anchors row B at $(0, 0, 3, 5, 24)$, and the Lagrangian decomposition controlling each row at the ambient level forces the family-dependent complementarity sum $\kappa(\mathcal{A}) + \kappa(\mathcal{A}^!)$ (0 for Kac–Moody and free fields, 13 for Virasoro). Theorem D identifies $\text{obs}_g = \kappa \cdot \lambda_g$ as the scalar-diagonal uniform-weight genus class on $\bar{\mathcal{M}}_{g,n}$ at level 4, with genus-1 unconditional and the all-genera statement on that lane proved by Grothendieck–Riemann–Roch on the universal curve and by clutching propagation independently; off that lane the full free energy carries $\delta F_g^{\text{cross}}$. Theorem H proves $\text{ChirHoch}^{\bullet}(A_b) \subset \{0, 1, 2\}$ at level 3 only under the full H_H package: PBW chiral Koszulness, finite-type/perfectness, genericity, E_{∞} -completion, strict Mittag–Leffler passage, and localized residue-twisted bar concentration; class M is a completed / pro / J -adic chain-level statement.

All five theorems are Σ_n -coinvariant projections of a single E_1 object: the ordered bar complex $\bar{B}_X^{\text{ord}}(A_b)$ with its deconcatenation coproduct and classical R -matrix $r_{A_b}(z)$. The averaging map $\text{av}: \mathfrak{g}^{E_1} \rightarrow \mathfrak{g}^{\text{mod}}$ recovers κ from r on the abelian and scalar lanes; the non-abelian Kac–Moody value $\kappa = \text{av}(r) + \dim(\mathfrak{g})/2$ encodes the Sugawara shift. The locus $\text{KSDual} \subset \text{Kosz}(X)$ on which the Verdier involution $\mathcal{A} \mapsto \mathcal{A}^!$ acts by the identity is the $\mathbb{Z}/2$ -fixed sublocus in the finite-type curved Verdier–Koszul ambient. On its admissible Koszul-complete part, Theorem A restricts to the unit / counit equivalence, Theorem H is exact in $\{0, 1, 2\}$ only under its H_H package, and the universal trace identity $\kappa_{\text{BKM}}(\Phi_N) = c_N(0)/2$ is the equivariant signature. The Heisenberg algebra

provides the complete base computation; every other family is a variation with increasingly complex collision residues.

Contents

1	Introduction: from a point to a curve	3
1.1	The problem	3
1.2	The answer	4
1.3	Prior work	5
1.4	The Heisenberg computation	6
1.5	Conventions	6
2	The bar complex on algebraic curves	6
2.1	Chiral algebras	6
2.2	The propagator and the Arnold relation	7
2.3	The ordered bar complex	7
2.4	Why $d^2 = 0$: the Borchers selection principle	8
2.5	Two coproducts	8
2.6	From genus 0 to all genera	8
3	Theorem A: chiral Koszul reflection $K^2 \simeq \text{id}$	9
3.1	Statement	9
3.2	Proof	11
4	Theorem B: bar-cobar inversion on the Koszul locus	12
4.1	Statement	12
4.2	The genus-0 argument	13
4.3	The higher-genus extension	13
5	Theorem C: 5×5 κ-stratification matrix	13
5.1	The problem	13
5.2	Statement	14
5.3	The Koszul complementarity sum	15
6	Theorem D: the genus expansion	15
6.1	Statement	15
6.2	The $\hat{A}$ -genus and Bernoulli numbers	17
6.3	The shadow obstruction tower	17
6.4	Proof architecture	17
7	Theorem H: chiral Hochschild cohomology	21
7.1	Statement	21
7.2	Family-specific specializations	21
7.3	Proof architecture	22

8	The dependency graph among Theorems A, B, C, D, and H	22
8.1	The dependency DAG	22
8.2	A creates the arena	22
8.3	B establishes faithfulness	23
8.4	C provides the polarization	23
8.5	D identifies the scalar shadow	23
8.6	The non-circular routing	24
8.7	Independence of converses	24
8.8	Summary: one new input per step	25
9	The Heisenberg: complete computation	25
9.1	The algebra	25
9.2	Theorem A for $\mathcal{H}_k$: bar-cobar adjunction	26
9.3	Theorem B for $\mathcal{H}_k$: bar-cobar inversion	26
9.4	Theorem C for $\mathcal{H}_k$: complementarity	26
9.5	Theorem D for $\mathcal{H}_k$: the genus expansion	27
9.6	Theorem H for $\mathcal{H}_k$: chiral Hochschild cohomology	27
9.7	The shadow tower and the derived center	28
9.8	Summary: the Heisenberg as base computation	28
10	The standard landscape	29
10.1	The census table	29
10.2	The general affine Kac–Moody formula	29
10.3	The R -matrix census	30
10.4	The four complexity classes	30
10.5	Free-field central charges	31
10.6	The Virasoro self-dual point	31
10.7	The $\mathcal{W}_N$ family	32
10.8	The Bershadsky–Polyakov algebra	32
10.9	The Hilbert polynomial census	32
10.10	Three features of the census	33
A	Ambients for the five theorems	34

I Introduction: from a point to a curve

I.1 The problem

Koszul duality over a point terminates. A quadratic algebra $A = T(V)/(R)$ determines a dual coalgebra $A^! = (H^*(B(A)))^\vee$, the bar construction $B(A) = (T^c(s^{-1}\tilde{A}), d_B)$ mediates between them, and the cobar functor inverts: $\Omega(B(A)) \xrightarrow{\sim} A$ when A is Koszul [2, 9, 14]. The bar cohomology concentrates in degree 1, the derived categories of A -modules and $A^!$ -modules are equivalent, and no further structure appears.

Over an algebraic curve, three obstructions prevent this from working. The generators of a chiral algebra are sections of a $\mathcal{D}_X$ -module, not elements of a vector space. The relations are operator product expansions with poles of arbitrary order, not quadratic expressions. The topology of higher-genus curves

forces the fiberwise bar differential into a curved A_∞ /CDG package: the raw identity is $m_1^2 = [m_o^{(g)}, -]$, and its scalar diagonal projection is $\kappa(\mathcal{A}) \cdot \omega_g$.

The question is: what structure replaces the terminating classical theory?

1.2 The answer

The chiral algebra $\mathcal{A} = A_b$ is the chart at a vacuum $b \in C^{\text{op}}$ of a factorization dg-category on a tangential log curve (X, D, τ) . The Beilinson tower $C^{\text{op}} \xrightarrow{b} A_b \xrightarrow{\bar{B}} B(A_b) \xrightarrow{Z_{\text{ch}}^{\text{der}}} \text{bulk} \rightarrow \text{line ops} \xrightarrow{\text{tr}}$ modular shadow organises the open quadrant of the factorization theory; the five theorems characterise its levels. The ordered bar complex

$$\bar{B}_X^{\text{ord}}(\mathcal{A}) = T^c(s^{-1}\bar{\mathcal{A}}), \quad |s^{-1}v| = |v| - 1,$$

with deconcatenation coproduct

$$\Delta[s^{-1}a_1 | \cdots | s^{-1}a_n] = \sum_{i=0}^n [s^{-1}a_1 | \cdots | s^{-1}a_i] \otimes [s^{-1}a_{i+1} | \cdots | s^{-1}a_n], \quad (1)$$

is a cofree E_1 -coalgebra. Here $\bar{\mathcal{A}} = \ker(\varepsilon: \mathcal{A} \rightarrow \omega_X)$ is the augmentation ideal and s^{-1} is the desuspension, lowering degree by 1. The integral kernel is the logarithmic 1-form

$$\eta_{12} = d \log(z_1 - z_2) = \frac{dz_1 - dz_2}{z_1 - z_2},$$

with conformally invariant residue 1: no coordinate, no metric, no level. At three points, the Arnold relation

$$\eta_{12} \wedge \eta_{23} + \eta_{23} \wedge \eta_{31} + \eta_{31} \wedge \eta_{12} = 0$$

generates all relations in $H^*(\text{Conf}_n(\mathbf{C}), \mathbf{C})$. The bar differential squares to zero if and only if the n -th products satisfy the Borchers identity; the Arnold relation kills the differential form, the Borchers identity kills the algebra, and both are consumed simultaneously. This is the *selection principle*: configuration space geometry determines the algebraic axioms; the bar construction accepts exactly chiral algebras.

The binary collision residue of the bar differential is the classical R -matrix

$$r_{\mathcal{A}}(z) = \text{Res}_{0,2}^{\text{coll}}(\Theta_{\mathcal{A}}) \in \text{End}_{\mathcal{A}}(2)[[z^{-1}]],$$

where $\Theta_{\mathcal{A}} := D_{\mathcal{A}} - d_o$ is the Maurer–Cartan element of the bar differential, satisfying $d\Theta_{\mathcal{A}} + \frac{1}{2}[\Theta_{\mathcal{A}}, \Theta_{\mathcal{A}}] = 0$ automatically because $(D_{\mathcal{A}})^2 = 0$. The Σ_n -coinvariant quotient erases the matrix structure and retains only the scalar

$$\kappa(\mathcal{A}) = \text{av}(r_{\mathcal{A}}(z)) \quad \text{for abelian and scalar families,}$$

while for non-abelian affine Kac–Moody

$$\kappa(V_k(\mathfrak{g})) = \text{av}(r_{\mathcal{A}}(z)) + \dim(\mathfrak{g})/2.$$

the modular characteristic: the single number that survives averaging. Five structural theorems extract the Σ_n -invariant content of the ordered bar. Each theorem characterizes one facet of $\Theta_{\mathcal{A}}$:

- (A) *Reflection*: $K^2 \simeq \text{id}$ on $\text{Kosz}(X)$ (chiral Koszul reflection at level $1 \leftrightarrow 2$).
- (B) *Inversion*: $\Omega_X(\overline{B}_X(C)) \xrightarrow{\sim} C$ in $D_{\text{ch}}^{\text{co}}(X)$ (level $1 \rightarrow 3$).
- (C) *Stratification*: the 5×5 κ -stratification matrix on rows G/L/C/M/B at level 5 per stratum, with $K_3 \times E$ anchoring row B at $(0, 0, 3, 5, 24)$.
- (D) *Leading scalar coefficient*: $\text{obs}_g = \kappa \cdot \lambda_g$ in $H^*(\overline{\mathcal{M}}_{g,n})$ (level 4; $\text{obs}_1 = \kappa \cdot \lambda_1$ unconditional; $\text{obs}_g = \kappa \cdot \lambda_g$ at all genera on the scalar-diagonal uniform-weight lane via GRR and clutching).
- (E) *Coefficient ring*: $\text{ChirHoch}^\bullet \subset \{0, 1, 2\}$ (level 3 on the Koszul locus, polynomial Hilbert series).

The chain is forced: without A, there is no dual at the chart level; without B, the bar transform can lose the chart algebra inside the derived chiral centre; without C, the κ -stratification matrix collapses to a single number; without D, the genus tower has no distinguished scalar shadow; without H, the coefficient ring is uncontrolled.

The locus $\text{KSDual} \subset \text{Kosz}(X)$ on which the Verdier involution $\mathcal{A} \mapsto \mathcal{A}^!$ acts by the identity is the $\mathbb{Z}/2$ -fixed sublocus where the four-quadrant grid $(\text{Open} \cup \text{CY}) \times (\text{Cat} \cup \text{Chain})$ degenerates into self-dual form: Theorem A is an equivalence (not adjunction with comparison map) only under the Koszul-complete package, Theorem H is exact in $\{0, 1, 2\}$ only under its H_H package, the five-archetype label is rigid only on the admissible finite-type curved fixed locus, and $\kappa_{\text{BKM}}(\Phi_N) = c_N(o)/2$ is the equivariant signature for $N \in \{1, 2, 3, 4, 6\}$ plus half- and quarter-integer continuations. Scalar anti-diagonal relations such as $K = 0$ or $c = 13$ are duality-orbit sums unless the finite-type curved Verdier–Koszul fixed-point criterion is also verified.

1.3 Prior work

Five existing programmes address aspects of the problem. Beilinson–Drinfeld [1] construct factorization algebras on $\text{Ran}(X)$ and prove chiral homology is well-defined, but do not treat bar-cobar inversion, complementarity, or the genus tower. Francis–Gaitsgory [4] establish chiral Koszul duality at genus 0 for Lie-type bar complexes; their framework does not reach higher genera or the full tensor bar. Costello–Gwilliam [3] develop factorization algebras for perturbative QFT; they do not treat Koszul duality or the modular characteristic. Gui–Li–Zeng [8] prove a quadratic duality at the cochain level, a genus-0 degree-2 result. Malikov–Schechtman [13] construct homotopy chiral algebras as the correct input, but do not prove the five structural theorems.

	A	B	C	D	H
Programme	adjunction	inversion	compl.	genus	chiral Hochschild
BD [1]	✓				
FG [4]	✓	✓*			
CG [3]					
GLZ [8]	✓*				
MS [13]					
This work	✓	✓	✓	✓	✓

*FG: genus 0, Lie-type. GLZ: genus 0, degree 2.

1.4 The Heisenberg computation

The Heisenberg chiral algebra $\mathcal{H}_k$ is generated by a single current J of conformal weight 1 with OPE $J(z)J(w) \sim k/(z-w)^2$. One generator, one double pole, no simple pole. The binary residue of the bar differential is the complete computation:

$$d_{\overline{B}}[s^{-1}J | s^{-1}J] = \text{Res}_{z_1=z_2} \left[\frac{k}{(z_1 - z_2)^2} \cdot d \log(z_1 - z_2) \right] = k.$$

The modular characteristic is $\kappa(\mathcal{H}_k) = k$, the collision residue $r(z) = k/z$ (scalar, not matrix), bar-cobar inversion gives $\Omega(\overline{B}(\mathcal{H}_k)) \simeq \mathcal{H}_k$, complementarity yields $\kappa(\mathcal{H}_k) + \kappa(\mathcal{H}_k^!) = k + (-k) = 0$, genus universality produces $F_1 = k/24$ and the $\hat{A}$ -series at all genera for this scalar-diagonal uniform-weight algebra, and chiral Hochschild gives $P(t) = 1 + t + t^2$. Every clause of every theorem is verified in a single example. Every other family is a variation on this computation, with increasingly complex collision residues.

1.5 Conventions

Throughout, X denotes a smooth projective curve over $\mathbf{C}$, $\mathcal{A}$ a chiral algebra on X in the sense of Beilinson–Drinfeld [1], and $\overline{\mathcal{M}}_g = \overline{\mathcal{M}}_{g,0}$ the Deligne–Mumford moduli space of stable curves of genus g . Grading is cohomological: $|d| = +1$. Desuspension lowers degree: $|s^{-1}v| = |v| - 1$. The bar complex uses the augmentation ideal: $\overline{B}(\mathcal{A}) = T^c(s^{-1}\tilde{\mathcal{A}})$ with $\tilde{\mathcal{A}} = \ker(\varepsilon)$. The Fulton–MacPherson compactification $\overline{\text{FM}}_n(X)$ is the real oriented blowup of X^n along all diagonal strata; it is a manifold with corners whose codimension-one faces D_S are indexed by subsets $S \subseteq \{1, \dots, n\}$ with $|S| \geq 2$.

2 The bar complex on algebraic curves

2.1 Chiral algebras

A *chiral algebra* $\mathcal{A}$ on a smooth curve X [1] is a $\mathcal{D}_X$ -module equipped with a chiral bracket

$$\mu^{\text{ch}}: j_* j^*(\mathcal{A} \boxtimes \mathcal{A}) \longrightarrow \Delta_!(\mathcal{A}),$$

where $j: X^2 \setminus \Delta \hookrightarrow X^2$ is the complement of the diagonal and $\Delta_!$ is the $!$ -pushforward along the diagonal embedding. In local coordinates, the chiral bracket encodes the operator product expansion:

$$a(z)b(w) \sim \sum_{n \geq 0} \frac{a_{(n)}b(w)}{(z-w)^{n+1}}.$$

The zeroth product $a_{(0)}b$ is the Lie bracket (first-order pole); the first product $a_{(1)}b$ carries the symmetric inner product (second-order pole). The Borcherds identity

$$\sum_{j \geq 0} \binom{m}{j} (a_{(n+j)}b)_{(m+k-j)}c = \sum_{j \geq 0} (-1)^j \binom{n}{j} \left(a_{(m+n-j)}(b_{(k+j)}c) - (-1)^n b_{(n+k-j)}(a_{(m+j)}c) \right) \quad (2)$$

is the defining axiom: it couples all pole orders in a single constraint.

2.2 The propagator and the Arnold relation

The bar differential extracts OPE data from collisions on $C_2(X) = X \times X \setminus \Delta$. A regular form has residue zero and extracts nothing. A double pole transforms under coordinate change and has no coordinate-invariant residue. A simple pole is forced; among 1-forms with a first-order pole along Δ , only the logarithmic derivative has conformally invariant residue:

$$\eta_{12} = d \log(z_1 - z_2) = \frac{dz_1 - dz_2}{z_1 - z_2}, \quad \text{Res}_\Delta \eta_{12} = 1.$$

This is the *propagator*: weight 1, no dependence on level, metric, or conformal structure.

Under $z \mapsto w(z)$, the propagator transforms as $\eta_{12} \mapsto \eta_{12} + d \log \frac{w(z_1) - w(z_2)}{z_1 - z_2}$; near the diagonal the correction reduces to $d \log w'(z_i)$, the standard cocycle for $\text{Diff}(X)$. This failure of invariance is not a defect; it is the datum that the Virasoro algebra measures.

At three points, the Arnold relation

$$\eta_{12} \wedge \eta_{23} + \eta_{23} \wedge \eta_{31} + \eta_{31} \wedge \eta_{12} = 0$$

restates the partial-fractions identity

$$\frac{1}{(z_1 - z_2)(z_2 - z_3)} + \frac{1}{(z_2 - z_3)(z_3 - z_1)} + \frac{1}{(z_3 - z_1)(z_1 - z_2)} = 0.$$

Arnold proved that, replicated across all triples $\{i, j, k\} \subset \{1, \dots, n\}$, this generates the complete ideal of relations in $H^*(\text{Conf}_n(\mathbf{C}), \mathbf{C})$.

2.3 The ordered bar complex

Definition 2.1 (Ordered bar complex). Let $\mathcal{A}$ be a chiral algebra on a smooth curve X , $\tilde{\mathcal{A}} = \ker(\varepsilon)$ the augmentation ideal. The *ordered bar complex* is

$$\overline{B}_X^{\text{ord}}(\mathcal{A}) = \bigoplus_{n \geq 1} (s^{-1}\tilde{\mathcal{A}})^{\otimes n} \otimes \Gamma(\overline{\text{FM}}_n(X), \Omega_{\log}^{n-1}),$$

with no Σ_n -quotient. Here s^{-1} denotes desuspension ($|s^{-1}a| = |a| - 1$), and the cohomological convention $|d| = +1$ is used throughout.

The bar differential decomposes into three pieces:

$$d_{\overline{B}} = d_{\text{int}} + d_{\text{res}} + d_{\text{dR}}.$$

The internal differential d_{int} acts on each tensor factor of $\mathcal{A}$. The de Rham differential d_{dR} acts on logarithmic forms on $\overline{\text{FM}}_n(X)$. The residue differential d_{res} extracts OPE data at collision divisors: for a codimension-one face $D_{\{i,j\}}$ corresponding to the collision $z_i \rightarrow z_j$,

$$d_{\text{res}}^{(ij)}(a_1 \otimes \dots \otimes a_n \otimes \eta_{ij} \wedge \omega') = \pm a_1 \otimes \dots \otimes \widehat{a_i} \otimes \dots \otimes \widehat{a_j} \otimes \dots \otimes a_{i(p)} a_j \otimes \omega',$$

where p is the pole order, hats denote omission, and $a_{(p)}b$ is the p -th Fourier mode of the OPE.

2.4 Why $d^2 = 0$: the Borchers selection principle

Theorem 2.2 (Bar nilpotence). *The bar differential satisfies $d_B^2 = 0$ if and only if the n -th products of $\mathcal{A}$ satisfy the Borchers identity (2).*

Proof. Expand $(d_{\text{int}} + d_{\text{res}} + d_{\text{dR}})^2$ into nine terms. Three vanish individually: $d_{\text{int}}^2 = 0$ (cochain complex), $d_{\text{dR}}^2 = 0$ (de Rham), and $\{d_{\text{int}}, d_{\text{dR}}\} = 0$ (Leibniz). The remaining terms are controlled by index overlap in the residue operator:

Disjoint indices ($\{i, j\} \cap \{k, \ell\} = \emptyset$): residue operators commute; cancellation is by the Koszul sign rule. No condition on the OPE.

One shared index (say $j = k$): three boundary strata meet at the codimension-2 locus $D_{ij\ell}$ of triple collision. The cyclic sum of iterated residues vanishes if and only if the Borchers identity holds. The Arnold relation kills the differential form; the Borchers identity kills the algebra. Both are consumed.

Same pair ($\{i, j\} = \{k, \ell\}$): $(\text{Res}_{D_{ij}})^2 = 0$ because the first Poincaré residue removes the unique logarithmic normal factor along D_{ij} , leaving no second normal direction for the same divisor. $\square$

Remark 2.3 (Selection table). The bar construction asks the same question at every level of geometry:

Base	Condition for $d^2 = 0$	Selected objects
Point	Jacobi identity	Lie algebras
Curve X	Borchers identity	Chiral algebras
X over $\overline{\mathcal{M}}_g$	+ period corrections	Modular Koszul algebras

Each row is forced by the geometry of the next.

2.5 Two coproducts

The ordered bar carries the *deconcatenation coproduct* (1): $n + 1$ terms, coassociative but not cocommutative, an E_1 -coalgebra reflecting the linear ordering. The Σ_n -coinvariant quotient

$$\overline{B}_X^\Sigma(\mathcal{A}) = \bigoplus_{n \geq 1} (\mathcal{A}^\Sigma)^{\otimes n}_{\Sigma_n} \otimes \Gamma(\overline{\text{FM}}_n(X), \Omega_{\log}^{n-1})$$

carries the *coshuffle coproduct*: 2^n terms, coassociative and cocommutative, an E_∞ -coalgebra. The ordered bar is the level-2 object before symmetrisation; the symmetric bar is the completed coinvariant descent. In the braided case this is the L_R -twisted descent, and the formal conductor fibre records the non-coinvariant R -matrix data.

2.6 From genus 0 to all genera

At genus $g \geq 1$, the function $z_1 - z_2$ has no global meaning; the propagator must be replaced by $d \log E(z_1, z_2)$, where $E(z, w)$ is the prime form on a Riemann surface of genus g (a section of $K^{-1/2} \boxtimes K^{-1/2}$, weight 1). The bar differential acquires a curved fiberwise package:

$$\text{tr}_{\text{diag}}(m_o^{(g)}) = \kappa(\mathcal{A}) \cdot \omega_g,$$

after scalar diagonal projection of the matrix-valued ordered curvature residue $r_{\mathcal{A}}(z) \cdot \omega_g \in \text{End}_{\mathcal{A}}(2) \otimes H^{1,1}(C_g)$. The raw chain identity is $m_1^2 = [m_o^{(g)}, -]$.

The *total* bar differential $D_{\mathcal{A}} = \sum_{g \geq 0} \hbar^g d_{\mathcal{A}}^{(g)}$ absorbs the curvature into the genus expansion: $D_{\mathcal{A}}^2 = 0$ unconditionally.

Definition 2.4 (Modular characteristic). The *modular characteristic* $\kappa(\mathcal{A})$ is the genus-one scalar obstruction. In scalar or abelian cases it equals the averaged collision residue. For nonabelian affine Kac–Moody algebras in the trace-form normalization,

$$\kappa(V_k(\mathfrak{g})) = \text{av}(r^{\text{coll}}) + \frac{\dim(\mathfrak{g})}{2} = \frac{\dim(\mathfrak{g})(k + h^\vee)}{2h^\vee},$$

where the second summand is the Sugawara simple-pole contribution.

Definition 2.5 (Maurer–Cartan element). The *Maurer–Cartan element* is $\Theta_{\mathcal{A}} := D_{\mathcal{A}} - d_0 \in \mathfrak{g}_{\mathcal{A}}^{\text{mod}}$, the positive-genus part of the bar differential. Since $D_{\mathcal{A}}^2 = 0$, it satisfies

$$d_0 \Theta_{\mathcal{A}} + \frac{1}{2} [\Theta_{\mathcal{A}}, \Theta_{\mathcal{A}}] = 0$$

unconditionally. Existence of the MC element is a tautology of the bar construction; the five theorems characterize its structure.

3 Theorem A: chiral Koszul reflection $K^2 \simeq \text{id}$

3.1 Statement

The bar complex produces a coalgebra; the question is what algebra it encodes. Three functors act on the bar complex and produce three distinct outputs:

$$\begin{aligned} \Omega(\overline{B}(\mathcal{A})) &\simeq \mathcal{A} && \text{(inversion: recovers the original),} \\ \Omega_X^{\text{cont}} \mathbb{D}_{\text{Ran}}(\overline{B}(\mathcal{A})) &\simeq \mathcal{A}_{\infty}^! && \text{(Verdier plus completed cobar: the homotopy dual),} \\ (H^*(\overline{B}(\mathcal{A})))^{\vee} &= \mathcal{A}^! && \text{(linear duality: the strict dual).} \end{aligned}$$

Confusing any two is a category error: cobar inverts, Verdier dualizes at chain level, and linear duality produces the strict dual.

Definition 3.1 (The Verdier component and homotopy Koszul dual $\mathcal{A}_{\infty}^!$; explicit construction). For an augmented chiral algebra $\mathcal{A}$ on a smooth curve X over a field of characteristic zero, define the *Verdier-dual bar component* by

$$\mathcal{A}_{\infty}^{\text{!co}} := \mathbb{D}_{\text{Ran}}(\overline{B}_X(\mathcal{A})),$$

the Verdier dual of the bar factorization coalgebra on $\text{Ran}(X)$, and then define

$$\mathcal{A}_{\infty}^! := \Omega_X^{\text{cont}} \mathcal{A}_{\infty}^{\text{!co}}.$$

Equivalently (Lowen–Positselski homotopy-coalgebra duality for conilpotent complete objects), $\mathcal{A}_{\infty}^!$ is the chiral algebra freely generated by $\mathcal{A}^{\vee}$ together with an $\mathcal{A}_{\infty}$ -structure obtained by transferring the cooperations of $\overline{B}_X(\mathcal{A})$ through a contraction onto its cohomology, under the finiteness hypothesis that $\overline{B}_X(\mathcal{A})$ is degreewise finite-dimensional in each bar-depth stratum (satisfied by every freely strongly generated chiral algebra, including Heisenberg, affine Kac–Moody at generic level, Virasoro, and the

principal $\mathcal{W}_N$). On the Koszul locus this A_∞ -structure strictifies to a genuine chiral algebra structure whose cohomology is the strict Koszul dual: $H^*(\mathcal{A}_\infty^!) \cong \mathcal{A}^!$.

Theorem 3.2 (Theorem A: chiral Koszul reflection $K^2 \simeq \text{id}$, parametric strength). *TYPE SIGNATURE. Quadrant: Open. Presentation: chain-level $+$ $(\infty, 2)$ -properad. Level: $1 \leftrightarrow 2$ (chart $A_b \leftrightarrow$ bar coalgebra $\overline{B}_X(A_b)$). Hypothesis package: (H_1) augmented; (H_2) augmentation-ideal complete (Positselski-weakly curved); (H_3) finite-dimensional graded bar pieces in the conformal-weight grading.*

Let X be a smooth projective curve over a field of characteristic 0 , $\mathcal{A} = A_b = \text{End}_C(b)$ the chart algebra at a vacuum b in the factorization dg-category C^{op} on (X, D, τ) , with bar complex $\overline{B}_X(\mathcal{A})$ in the finite-bar class of Definition 3.1.

(T1a) Weak strength (unconditional in the weight-completed ambient under (H_1) alone): the bar and cobar functors form an adjoint pair

$$\overline{B}_X : \text{Alg}_{\text{aug}}(\text{Fact}(X)) \rightleftarrows \text{CoAlg}_{\text{conil}}(\text{Fact}(X)) : \Omega_X$$

with unit $\eta : \text{id} \rightarrow \Omega_X \circ \overline{B}_X$ and counit $\varepsilon : \overline{B}_X \circ \Omega_X \rightarrow \text{id}$, natural in $\mathcal{A}$ within the Francis–Gaitsgory factorization ambient of [4], provided that ambient is taken in its weight-completed / pro-conilpotent form $\text{pro-CoAlg}_{\text{conil}}(\text{Fact}(X))$ whenever $\mathcal{A}$ has unbounded conformal-weight grading (classes G, L with positive-energy grading and, a fortiori, class M). On the raw bounded direct-sum ambient $\text{Ch}(\text{Vect})$ the target $\text{CoAlg}_{\text{conil}}$ is too small for class M : the bar coalgebra $\overline{B}(\text{Vir}_c)$ fails conilpotency because $L_0 \in \overline{B}^1$ has $\tilde{\Delta}^{(N)}(L_0) \neq 0$ for every $N \geq 1$ (Remark 3.4; same L_0 -obstruction as Theorem B (B.2), thm:chiral-positselski-weight-completed).

(T1b) Strong strength: Verdier intertwining $\mathbb{D}_{\text{Ran}} \overline{B}_X \simeq \mathcal{A}_\infty^{!,\text{co}}$ (under the holonomic dualizable hypotheses): Verdier duality on $\text{Ran}(X)$ intertwines the bar with the Verdier component of Definition 3.1, and completed cobar gives the homotopy Koszul dual:

$$\mathbb{D}_{\text{Ran}} \overline{B}_X(\mathcal{A}) \simeq \mathcal{A}_\infty^{!,\text{co}}, \quad \Omega_X^{\text{cont}} \mathcal{A}_\infty^{!,\text{co}} \simeq \mathcal{A}_\infty^!.$$

No Koszul hypothesis is invoked at this strength; $\mathcal{A}_\infty^!$ carries A_∞ -structure measuring the failure of strict Koszulness.

(T1c) Strongest strength: $K^2 \simeq \text{id}$ on $\text{Kosz}(X)$ (under $(H_1) + (H_2) + (H_3)$): on the Koszul locus, where a twisting morphism $\tau : C \rightarrow \mathcal{A}$ with C conilpotent makes $K_\tau^L(\mathcal{A}, C)$ acyclic (equivalently the PBW spectral sequence collapses at E_2), the chiral Koszul reflection

$$K_X(\mathcal{A}) := \Omega_X \mathbb{D}_{\text{Ran}} \overline{B}_X(\mathcal{A}), \quad K_X^2 \simeq \text{id}_{\text{Kosz}(X)},$$

is an involution on $\text{Kosz}(X)$, the unit of (T1a) and the intertwining of (T1b) become quasi-isomorphisms,

$$C \xrightarrow{\sim} \overline{B}_X(\mathcal{A}), \quad \Omega_X(C) \xrightarrow{\sim} \mathcal{A}, \quad H^*(\mathcal{A}_\infty^!) \cong \mathcal{A}^!,$$

and on the $\mathbb{Z}/2$ -fixed sublocus $\text{KSDual} \subset \text{Kosz}(X)$ the reflection is the identity.

(T1d) Strongest rejected strength: the assertion that the adjunction of (T1a) is an equivalence on the full Alg^{aug} off the Koszul locus, in the raw direct-sum ambient $\text{Ch}(\text{Vect})$, is false. The witness is the same L_0 -obstruction as in (T1a): $L_0 \in \overline{B}^1(\text{Vir}_c)$ has $\tilde{\Delta}^{(N)}(L_0) \neq 0$ for every $N \geq 1$, so $\overline{B}_X(\text{Vir}_c)$ exits $\text{CoAlg}_{\text{conil}}$ at the raw direct-sum level. The Koszul hypothesis or the weight-completed ambient is

structurally required; bundling the unconditional adjunction (T1a) with the Koszul-locus reflection (T1c) into a single ambient-free equivalence mistakes the forgetful image for the source.

Remark 3.3 (Four parametric strengths, disjoint inputs). The partition into (T1a) weak, (T1b) strong, (T1c) strongest, (T1d) strongest-rejected separates parametric strengths with disjoint load-bearing inputs: weak adjunction uses only the cofree/free universal properties and the Francis–Gaitsgory ambient [4] weight-completed when the conformal-weight grading is unbounded; Verdier intertwining uses the six-functor formalism on $\text{Ran}(X)$ together with the finiteness hypothesis (H_3); the Koszul reflection $K^2 \simeq \text{id}$ uses Koszul acyclicity / PBW E_2 -collapse; the rejected strength names the boundary precisely so that no claim accidentally crosses it. Bundling weak and strong into a single equivalence conflates universal properties and produces the false slogan “ $\text{Bar}(\mathcal{A})$ is the bulk”: the bar coalgebra is the twisting object at level 2, and the bulk is $Z_{\text{ch}}^{\text{der}}(\mathcal{A})$ at level 3.

Remark 3.4 (Ambient qualifier for (T1a); class-M at the raw direct sum). The adjunction $\overline{B}_X \dashv \Omega_X$ is a formal consequence of the free/cofree universal properties in the chiral factorization setting of Francis–Gaitsgory [4], *provided* the target coalgebra category genuinely contains $\overline{B}_X(\mathcal{A})$. For class G (Heisenberg) and class L (affine Kac–Moody away from critical level) the bar coalgebra is conilpotent already at the bounded direct-sum ambient $\text{Ch}(\text{Vect})$, so the adjunction lands in $\text{CoAlg}_{\text{conil}}(\text{Fact}(X))$ without further completion. For class M (Virasoro, $\mathcal{W}_N$, critical-level Kac–Moody) and, more generally, for any chiral algebra whose bar filtration $F^n = \bigoplus_{k \leq n} B_k$ has unbounded conformal weight, conilpotency of $\overline{B}_X(\mathcal{A})$ fails on the raw direct sum: the L_0 -witness $\tilde{\Delta}^{(N)}(L_0) \neq 0$ for every $N \geq 1$ in $\overline{B}^1(\text{Vir}_c)$ (the Vir raw bar proposition in `chapters/theory/theorem_B_scope_platonic.tex`). The correct ambient is then the pro-conilpotent / weight-completed factorization category $\text{pro-CoAlg}_{\text{conil}}(\text{Fact}(X))$ against the strong filtration, in which conilpotency is restored at every finite-weight truncation and the Mittag–Leffler–Milnor passage to the inverse limit converges, exactly as in Theorem B (B.2), `thm:chiral-positselfski-weight-completed`, `thm:completed-bar-cobar-strong`. In this precise sense (T1a) and Theorem B (B.2) share a single ambient discipline: the unconditional statement is at the weight-completed level, and the raw direct-sum statement for class M is genuinely false (not a gap) and should not be asserted at that ambient.

3.2 Proof

Step 1: The twisting morphism. A chiral twisting morphism $\tau: C \rightarrow \mathcal{A}$ is a degree-+1 element in the chiral convolution algebra satisfying the Maurer–Cartan equation $d\tau + \tau \star \tau = 0$, where the star product is mediated by the integration kernel η_{12} on $X^2 \setminus \Delta$. The acyclicity of the twisted tensor product $K_\tau^L(\mathcal{A}, C)$ is the defining property of a chiral Koszul pair.

Step 2: The PBW recognition criterion. The PBW filtration on $\mathcal{A}$ by bar-length makes the associated graded quadratic-Koszul in the classical sense. The chiral algebra need not be quadratic (the Virasoro OPE has a quartic pole, $\mathcal{W}$ -algebras have poles of arbitrary order); what matters is that the leading-order structure under the PBW filtration is. The E_2 page of the bar spectral sequence identifies with the classical bar complex of the associated graded, and the collapse at E_2 is the chiral Koszul condition.

Step 3: Verdier intertwining. Verdier duality $\mathbb{D}_{\text{Ran}}$ on $\text{Ran}(X)$ converts coalgebras to algebras. Applied to $\overline{B}_X(\mathcal{A})$, it produces a factorization algebra $\mathcal{A}_\infty^!$ on $\text{Ran}(X)$. The key identity is

$$\mathbb{D}_{\text{Ran}}(T^c(s^{-1}\tilde{\mathcal{A}})) \simeq T(s(H^*(\overline{B}(\mathcal{A})))^\vee),$$

where the left side is the Verdier dual of the cofree coalgebra and the right side is the free algebra on the linear dual of bar cohomology. On the Koszul locus, bar cohomology concentrates in degree 1, and the right side reduces to the strict Koszul dual algebra $\mathcal{A}^!$.

Step 4: Filtered comparison. The filtered comparison lemma establishes that the PBW spectral sequence is compatible with the Verdier functor: the filtration on $\overline{B}_X(\mathcal{A})$ induced by bar degree is taken by $\mathbb{D}_{\text{Ran}}$ to the dual filtration on $\mathcal{A}^!_{\infty}$. This is the non-formal content of the theorem; the compatibility is not automatic and requires the Fulton–MacPherson boundary structure.

Example 3.5 (Heisenberg). For $\mathcal{A} = \mathcal{H}_k$: the bar complex $\overline{B}(\mathcal{H}_k) = T^c(s^{-1}\mathbf{C} \cdot J)$ has bar cohomology concentrated in degree 1 ($H^{-1}(\overline{B}(\mathcal{H}_k)) = \mathbf{C} \cdot s^{-1}J$). The Koszul dual coalgebra is $\mathcal{H}_k^! = \mathbf{C} \cdot s^{-1}J$; the strict Koszul dual is the curved second-kind symmetric chiral algebra $\mathcal{H}_k^! = (\text{Sym}^{\text{ch}}(V^*[1]), m_o = -k\omega)$. The Verdier dual recovers $\mathcal{H}_k^!$ at the level of cohomology. The complementarity sum: $\kappa(\mathcal{H}_k) + \kappa(\mathcal{H}_k^!) = k + (-k) = o$.

4 Theorem B: bar-cobar inversion on the Koszul locus

4.1 Statement

Theorem A constructs the dual algebra $\mathcal{A}^!_{\infty}$, but does not prove that the bar transform is faithful: the coalgebra $\overline{B}(\mathcal{A})$ could in principle fail to reconstruct $\mathcal{A}$.

Theorem 4.1 (Theorem B: bar-cobar inversion on the Koszul locus). *TYPE SIGNATURE. Quadrant: Open. Presentation: chain-level in $D_{\text{ch}}^{\text{co}}(X)$. Level: 1 $\rightarrow$ 3 (chart A_b recovered from bar coalgebra; identity on the Koszul locus inside the derived chiral centre via Theorem A). Hypothesis package: chiral Koszul (twisting morphism with acyclic twisted tensor product, equivalently PBW E_2 -collapse); class-M chain-level requires the weight-completed / pro / J-adic ambient.*

Let $\mathcal{A} = A_b$ be a chirally Koszul chart algebra on a smooth curve X in the Beilinson tower of the factorization dg-category $\mathcal{C}^{\text{op}}$.

- (i) Genus o , classes G and L (unconditional in $\text{Ch}(\text{Vect})$). *The cobar-bar counit is a quasi-isomorphism:*

$$\varepsilon : \Omega_X(\overline{B}_X(\mathcal{A})) \xrightarrow{\sim} \mathcal{A} \quad \text{in } D_{\text{ch}}^{\text{co}}(X).$$

The bar spectral sequence collapses at E_2 , and bar cohomology concentrates in bar degree 1: $H^(\overline{B}(\mathcal{A})) \cong (s \mathcal{A}^!)$, identifying the Koszul dual coalgebra. Class G admits an explicit MacLane splitting; class L uses PBW E_2 -collapse.*

- (ii) Genus $g \geq 1$ on the standard landscape (modular Koszul lane). *If $\mathcal{A}$ satisfies diagonal Ext vanishing on the fibers of the universal curve over $\overline{\mathcal{M}}_{g,n}$, the genus- g cobar-bar counit $\Omega^{(g)}(\overline{B}^{(g)}(\mathcal{A})) \xrightarrow{\sim} \mathcal{A}^{(g)}$ is a quasi-isomorphism in the coderived category. On the Koszul / collapse loci of the standard landscape (Heisenberg, affine Kac–Moody at generic level, lattice vertex algebras), the statement is ordinary. Class M (Virasoro, $\mathcal{W}_N$) at generic central charge is genuinely false at the raw direct-sum ambient: the harmonic discrepancy $S_4 = 10/[c(5c + 22)] \neq o$ forces bar cohomology at weight 4. The correct chain-level closure is the weight-completed coderived theorem thm : chiral – positselski – weight – completed, with the Mittag–Leffler / Milnor passage to the inverse limit absorbing the discrepancy at every weight stage.*

4.2 The genus-0 argument

The genus-0 proof proceeds by spectral sequence. The PBW filtration on $\mathcal{A}$ by conformal weight induces a decreasing filtration F^p on $\overline{B}_X(\mathcal{A})$ by bar degree: elements of F^p have at least p tensor factors. The associated spectral sequence has E_1 page equal to the classical bar complex $B(\text{gr } \mathcal{A})$ of the associated graded, and converges to $H^*(\overline{B}_X(\mathcal{A}))$.

The Koszul condition is that $\text{gr } \mathcal{A}$ is quadratic Koszul in the classical sense (Priddy [14]). Then the E_1 differential has cohomology concentrated on the diagonal: $E_2^{p,q} = 0$ for $p \neq q$. The spectral sequence collapses at E_2 , and bar cohomology concentrates in bar degree 1. The counit map $\varepsilon: \Omega_X(\overline{B}_X(\mathcal{A})) \rightarrow \mathcal{A}$ is then a quasi-isomorphism by the comparison theorem for filtered complexes.

4.3 The higher-genus extension

At genus $g \geq 1$, the fiberwise bar differential is curved: $m_1^2 = [m_o^{(g)}, -]$, with scalar diagonal $\text{tr}_{\text{diag}}(m_o^{(g)}) = \kappa(\mathcal{A}) \cdot \omega_g$. The cobar functor cannot be applied in the classical sense. The resolution uses the total bar differential $D_{\mathcal{A}}$ with $D_{\mathcal{A}}^2 = 0$: the curvature is absorbed into the genus expansion.

The genus- g inversion requires two conditions:

- (i) *Central curvature*: the scalar diagonal projection of $m_o^{(g)}$ is $\kappa(\mathcal{A}) \cdot \omega_g$, rather than an arbitrary endomorphism-valued class.
- (ii) *Diagonal Ext vanishing*: the twisted tensor product $K_{\tau}^L(\mathcal{A}^{(g)}, C^{(g)})$ is acyclic at genus g .

For the standard landscape, central curvature is automatic (the Virasoro-type curvature is generated by the stress tensor) and diagonal Ext vanishing follows from the PBW argument adapted to the genus- g bar complex. The exception is the $\beta\gamma$ system at integer spin, where the Ext group has an anomalous contribution from the fermionic parity.

Example 4.2 (Heisenberg). $\overline{B}(\mathcal{H}_k)$ has bar cohomology concentrated in bar degree 1: $H^{-1}(\overline{B}(\mathcal{H}_k)) = \mathbf{C} \cdot s^{-1}J$. The cobar recovers $\mathcal{H}_k$ as a graded algebra. Bar-cobar inversion is unconditional at all genera: the curvature $d_{\text{fib}}^2 = k \cdot \omega_g$ is scalar, and the Ext vanishing is automatic because the only OPE pole is of order 2 (no Lie bracket, no higher obstructions).

Remark 4.3 (Off the Koszul locus). Off the Koszul locus, bar-cobar inversion fails: the spectral sequence does not collapse at E_2 , bar cohomology is not concentrated in degree 1, and the counit is not a quasi-isomorphism. The failure is measured by the shadow obstruction tower $\{S_r(\mathcal{A})\}_{r \geq 2}$: the degree- r shadow class $S_r \in H^*(\overline{\mathcal{M}}_{0,r})$ is the obstruction to extending the inversion from bar degree $r-1$ to bar degree r . All standard chiral algebras in the landscape are chirally Koszul; the shadow tower measures the *complexity* of their higher-genus behaviour, not the failure of duality itself.

5 Theorem C: 5×5 κ -stratification matrix

5.1 The problem

At genus $g \geq 1$, the bar complex of $\mathcal{A}$ and the bar complex of its Koszul dual $\mathcal{A}^!$ coexist on the same moduli space $\overline{\mathcal{M}}_g$. The curvatures are related: $d_{\text{fib}}^2(\mathcal{A}) = \kappa(\mathcal{A}) \cdot \omega_g$ and $d_{\text{fib}}^2(\mathcal{A}^!) = \kappa(\mathcal{A}^!) \cdot \omega_g$. The Verdier involution exchanges the two bar complexes. The question is whether this exchange induces a canonical decomposition of the ambient cohomology.

5.2 Statement

Theorem 5.1 (Theorem C: 5×5 κ -stratification matrix and three-tier complementarity). *TYPE SIGNATURE. Quadrant: Open (with vertical equivalence to Vol III at level 5). Presentation: scalar shadow per stratum plus the Co/C1/C2 complementarity tower. Level: level 3 for the derived-centre ambient, level 5 for the modular trace and clutching scalar shadow. Hypothesis package: Co strict flat fibre-centre comparison; C1 represented Verdier involution, perfectness, and nondegenerate anti-invariant Verdier pairing; C2 shifted-symplectic/BV package only when explicitly supplied; five κ -measurements per family; collapse pattern across rows fixes the archetype label; $\kappa_{\text{BKM}}(\Phi_N) = c_N(o)/2$ as the Borchers-lane identity; Borchers weight, chiral conductor, and fibre Euler characteristic are three different measurements.*

Each chirally Koszul chart algebra $\mathcal{A} = A_b$ in the standard landscape carries five κ -measurements on the modular trace. Its complementarity data are three-tiered: Co constructs the strict flat fibre-centre ambient; C1 gives the Verdier eigenspace and perfect-pairing statement on the stated ambients; C2 is the conditional BV/shifted-symplectic upgrade. The scalar conductor is the trace shadow of C1 together with Theorem D, not a substitute for C2.

(C1) The 5×5 stratification matrix. Five archetypes G/L/C/M/B with shadow depths $r_{\max} \in \{2, 3, 4, \infty, 5\}$ admit five κ -measurements: $(\kappa_{\text{cat}}, \kappa_{\text{ch}}^{\text{Hodge}}, \kappa_{\text{ch}}^{\text{Heis}}, \kappa_{\text{BKM}}, \kappa_{\text{fiber}})$ reading the categorical, Hodge-graded chiral, Heisenberg chiral, Borchers, and CY-fibre weights respectively.

	κ_{cat}	$\kappa_{\text{ch}}^{\text{Hodge}}$	$\kappa_{\text{ch}}^{\text{Heis}}$	κ_{BKM}	κ_{fiber}
G (Heisenberg, lattice V_Λ)	o	o	$\text{rk } \Lambda$	N/A	N/A
L (affine $V_k(\mathfrak{g})$, generic level)	o	o	$d(k + h^\vee)/(2h^\vee)$	N/A	N/A
C (free fields $\beta\gamma, bc$)	o	o	$6\lambda^2 - 6\lambda + 1$	N/A	N/A
M (Virasoro, $\mathcal{W}_N$, generic c)	o	$c/2$	N/A	N/A	N/A
B ($K_3 \times E$ with Δ_5)	o	o	3	5	24

The $K_3 \times E$ row anchors archetype B at $(o, o, 3, 5, 24)$ with Δ_5 as primitive Borchers denominator and Mukai conductor $K_{\mathcal{B}}^\kappa = 2c_+(\text{Mukai}(K_3)) = 8$. The collapse pattern across rows -- which entries are nonzero, which are forced to vanish or to remain undefined -- is the classification axis: five κ are not one invariant.

(C2) C1 Verdier eigenspace decomposition (ordinary on the proved finite-type lanes, weight-completed where the completion hypotheses are supplied). For every $g \geq 0$ for which the Co/C1 package is available, the genus- g ambient complex $\mathbf{C}_g(\mathcal{A}) := R\Gamma(\overline{\mathcal{M}}_g, \mathcal{Z}_{\mathcal{A}})$ admits a homotopy decomposition

$$\mathbf{C}_g(\mathcal{A}) \simeq \mathbf{Q}_g(\mathcal{A}) \oplus \mathbf{Q}_g(\mathcal{A}^\dagger),$$

where $\mathbf{Q}_g(\mathcal{A})$ and $\mathbf{Q}_g(\mathcal{A}^\dagger)$ are the Verdier eigenspaces. Under the perfectness and nondegenerate anti-invariant pairing hypotheses they are Lagrangian summands with respect to the Verdier pairing of degree $-(3g - 3)$. At the cohomological level for $g \geq 1$,

$$Q_g(\mathcal{A}) \oplus Q_g(\mathcal{A}^\dagger) \cong H^*(\overline{\mathcal{M}}_g, \mathcal{Z}_{\mathcal{A}}),$$

the two summands are eigenspaces of the Verdier involution, and the decomposition is canonical. At $g = 0$ the involution is trivial, $Q_0(\mathcal{A}^\dagger) = 0$, and no Verdier pairing of negative degree is invoked.

- (C₃) Family-stratum scalar ceiling $\kappa(\mathcal{A}) + \kappa(\mathcal{A}^\dagger) = K(\mathcal{A})$. The Lagrangian decomposition forces the scalar complementarity sum $K(\mathcal{A}) = \kappa(\mathcal{A}) + \kappa(\mathcal{A}^\dagger)$ family-dependently: 0 for Heisenberg, lattice, and affine Kac–Moody (sign reversal under the critical-level reflection $k \mapsto -k - 2h^\vee$); 13 for Virasoro ($c \mapsto 26 - c$, self-dual at $c = 13$); $250/3$ for $\mathcal{W}_3$; $98/3$ for Bershadsky–Polyakov in the principal-value sense across the critical pole at $k = -3$. The universal form is $\kappa + \kappa^\dagger = \rho_{\mathcal{A}} \cdot K$ with $\rho_{\text{KM}} = \rho_{\text{free}} = 0$, $\rho_{\text{Vir}} = 1/2$, $\rho_{\mathcal{W}_N} = H_N - 1$, $\rho_{\text{BP}} = 1/6$.
- (C₄) C₂ shifted-symplectic upgrade (conditional on the C₂ package). If the C₂ package is supplied – finite modular envelope, cyclic closedness, skew-adjoint differential, modular-bracket compatibility, twisted-tensor acyclicity, and BV/QME anomaly killing on the chosen finite window – then the deformation-stack intersection carries the shifted-symplectic/Lagrangian structure predicted by PTVV. This upgrade is not implied by the scalar conductor, by the C₁ eigenspace decomposition alone, or by uniform-weight diagonalization.

5.3 The Koszul complementarity sum

The Lagrangian decomposition constrains the modular characteristics of $\mathcal{A}$ and $\mathcal{A}^\dagger$:

$$\kappa(\mathcal{A}) + \kappa(\mathcal{A}^\dagger) = K(\mathcal{A}),$$

where $K(\mathcal{A})$ is the *Koszul complementarity conductor*, a family-dependent constant. This is not a universal zero.

- Proposition 5.2** (Complementarity sums). (i) Heisenberg: $\kappa(\mathcal{H}_k) = k$, $\kappa(\mathcal{H}_k^\dagger) = -k$, $K = 0$.
(ii) Affine Kac–Moody: $\kappa(V_k(\mathfrak{g})) = \frac{\dim(\mathfrak{g})(k+h^\vee)}{2h^\vee}$, $\kappa(V_k(\mathfrak{g})^\dagger) = -\frac{\dim(\mathfrak{g})(k+h^\vee)}{2h^\vee}$, $K = 0$.
(iii) Virasoro: $\kappa(\text{Vir}_c) = c/2$, $\kappa(\text{Vir}_{26-c}) = (26 - c)/2$, $K = 13$. Self-dual at $c = 13$.
(iv) Principal $\mathcal{W}_N$: $\kappa(\mathcal{W}_N) = c \cdot (H_N - 1)$ where $H_N = \sum_{j=1}^N 1/j$. K is nonzero and N -dependent.
(v) Free fields (lattice, $\beta\gamma$, bc): $K = 0$.

Proof. The Lagrangian decomposition of Theorem C forces $\kappa + \kappa^\dagger = \rho_{\mathcal{A}} \cdot K$ at the level of the scalar shadow, with $\rho_{\mathcal{A}}$ family-specific ($\rho_{\text{KM}} = \rho_{\text{free}} = 0$, $\rho_{\text{Vir}} = 1/2$, $\rho_{\mathcal{W}_N} = H_N - 1$, $\rho_{\text{BP}} = 1/6$). For Heisenberg and affine Kac–Moody, the Koszul dual has opposite modular characteristic (the Verdier involution reverses the sign of κ), giving $K = 0$. For the Virasoro algebra, $c \mapsto 26 - c$ under Koszul duality (the ghost central charge), and $K = c/2 + (26 - c)/2 = 13$, constant. The self-dual point $c = 13$ is the unique value where $\kappa = \kappa' = 13/2$: the Lagrangian decomposition is symmetric. $\square$

Remark 5.3 (Five objects). The five objects appearing in the complementarity theorem: $\mathcal{A}$ (the algebra), $\overline{B}(\mathcal{A})$ (the bar coalgebra), $\mathcal{A}^\dagger = H^*(\overline{B}(\mathcal{A}))$ (the dual coalgebra), $\mathcal{A}^\dagger = (\mathcal{A}^\dagger)^\vee$ (the dual algebra), and $\mathcal{Z}_{\text{ch}}^{\text{der}}(\mathcal{A}) = R \text{Hom}_{\mathcal{A} \otimes \mathcal{A}^{\text{op}}}(\mathcal{A}, \mathcal{A})$ (the derived chiral center). These are distinct objects produced by distinct functors; conflation of any two is a category error.

6 Theorem D: the genus expansion

6.1 Statement

Complementarity supplies an ambient polarization of the genus tower, but the scalar projection κ is already bar-intrinsic and must still be identified genus by genus.

Theorem 6.1 (Theorem D: scalar modular characteristic and cross-channel correction). *TYPE SIGNATURE. Quadrant: Open. Presentation: chain-level on the universal curve $\overline{C}_g \rightarrow \overline{M}_{g,n}$. Level: 4 (line / brane operators on $\overline{M}_{g,n}$ realising the scalar-projection class). Hypothesis package: chiral Koszul; scalar-diagonal uniform-weight at part (ii); $H^*(\overline{M}_{g,n})$ as the receiving cohomology with $\lambda_g = c_g(\mathbb{E})$ the top Chern class of the Hodge bundle. The raw positive-genus fibrewise operator is curved:*

$$m_1^2 = [m_o, -].$$

Only its scalar diagonal shadow gives $\kappa(\mathcal{A})\lambda_g$, and only the period-corrected total modular differential D_g is square-zero.

Let $\mathcal{A} = A_b$ be a chirally Koszul chart algebra with modular characteristic $\kappa(\mathcal{A})$.

(i) Genus 1 (unconditional). *The genus-1 obstruction class is*

$$\text{obs}_1(\mathcal{A}) = \kappa(\mathcal{A}) \cdot \lambda_1 \in H^*(\overline{M}_{1,1}),$$

and the genus-1 free energy is $F_1(\mathcal{A}) = \kappa(\mathcal{A})/24$. The class is forced by one-dimensionality of $H^2(\overline{M}_{1,1}) = \mathbb{Q} \cdot \lambda_1$ together with the curved fibre equation $m_1^2 = [m_o, -]$, whose scalar diagonal projection is $d_{\text{fib}}^2 = \kappa \cdot \omega_1$.

(ii) All genera, scalar-diagonal uniform-weight lane. *If the strong generators of $\mathcal{A}$ have a single conformal weight and the scalar channel is diagonal, the obstruction class at every genus is*

$$\text{obs}_g(\mathcal{A}) = \kappa(\mathcal{A}) \cdot \lambda_g \quad (g \geq 1),$$

and the free energies are $F_g(\mathcal{A}) = \kappa(\mathcal{A}) \cdot \hat{A}_g$, where $\hat{A}_g$ is the g -th coefficient of the $\hat{A}$ -genus:

$$\sum_{g \geq 1} F_g(\mathcal{A}) \cdot \hbar^{2g} = \kappa(\mathcal{A}) \cdot (\hat{A}(i\hbar) - 1).$$

Two independent proof routes converge: Grothendieck–Riemann–Roch on the universal curve identifies the bar curvature with the Hodge bundle curvature via Arakelov–Faltings (Proof A); clutching propagation from the genus-1 base case combined with tautological-line support criterion (Proof B). Each proof invokes disjoint external machinery; agreement is the consistency check.

(iii) Multi-weight correction with cross-channel term. *For multi-weight algebras at genus $g \geq 2$, the diagonal term remains universal but the full free energy carries a cross-channel correction:*

$$F_g(\mathcal{A}) = \kappa(\mathcal{A}) \cdot \lambda_g^{\text{FP}} + \delta F_g^{\text{cross}}(\mathcal{A}),$$

with $\delta F_g^{\text{cross}}$ the explicit correction from mixed conformal-weight interactions; vanishing for uniform-weight families and at $g = 1$ for all families. The closed form for $\delta F_g^{\text{cross}}$ at general multi-weight $\mathcal{A}$ and $g \geq 2$ is the open frontier (F2).

Remark 6.2 (Scope of the all-genera statement). Part (i) is unconditional: the genus-1 formula $\text{obs}_1 = \kappa \cdot \lambda_1$ holds for every chirally Koszul algebra, with no uniform-weight hypothesis and no clutching input. Part (ii) extends the scalar factorization from genus 1 to all genera by propagation along the clutching morphisms on $\partial \overline{M}_g$. The non-trivial step is *clutching-uniqueness*: the assertion that the scalar factorization propagated through clutching agrees, at each genus, with the direct bar-complex computation. This step

uses the bar-intrinsic scalar trace of $\Theta_{\mathcal{A}}$ together with the boundary structure of $\overline{\mathcal{M}}_g$; it is proved on the uniform-weight lane where the graph-sum formula factors through κ alone. An independent derivation via the Grothendieck–Riemann–Roch theorem on the universal curve (§6.4) provides a second path that bypasses clutching-uniqueness entirely. The genus-1 result and the two proof paths (clutching and GRR) are the principal structural elements; the all-genera statement on the uniform-weight lane rests on either one.

6.2 The $\hat{A}$ -genus and Bernoulli numbers

The $\hat{A}$ -genus has the explicit expansion

$$\hat{A}(x) - 1 = \frac{x/2}{\sin(x/2)} - 1 = \frac{1}{24}x^2 + \frac{7}{5760}x^4 + \frac{31}{967680}x^6 + \dots$$

The coefficients are $\hat{A}_g = \frac{2^{2g-1}-1}{2^{2g-1}} \cdot \frac{|B_{2g}|}{(2g)!}$, where B_{2g} is the $2g$ -th Bernoulli number. The leading term $\hat{A}_1 = 1/24$ is universal: for every family, $F_1 = \kappa/24$.

6.3 The shadow obstruction tower

The Maurer–Cartan element $\Theta_{\mathcal{A}}$ projects at each degree $r \geq 2$ to a shadow invariant $S_r(\mathcal{A})$; the tower $\{S_r\}_{r \geq 2}$ classifies chiral algebras into four complexity classes:

Class	Depth	Δ	Example	Behaviour
G (Gaussian)	2	$\mathfrak{o}(S_4 = \mathfrak{o})$	Heisenberg	Formal, free
L (Lie)	3	$\mathfrak{o}(\kappa = \mathfrak{o})$	Aff. KM (crit.)	Single Massey
C (Contact)	4	special	$\beta\gamma$	Contact then rigid
M (Mixed)	∞	$\neq \mathfrak{o}$	Virasoro, $\mathcal{W}_N$	Non-formal

The critical discriminant $\Delta = 8\kappa S_4$ governs a dichotomy: $\Delta = \mathfrak{o}$ terminates the tower, $\Delta \neq \mathfrak{o}$ makes it infinite. The *shadow depth* $r_{\max}$ (the highest degree at which $S_r \neq \mathfrak{o}$) takes values $\{2, 3, 4, \infty\}$; the *algebraic depth* $d_{\text{alg}} \in \{\mathfrak{o}, 1, 2, \infty\}$ measures the internal complexity of the MC equation and admits no value 3 or any finite value ≥ 3 ; there is a gap.

6.4 Proof architecture

The proof of genus universality proceeds in three steps.

Step 1: Genus 1 (unconditional). On $\overline{\mathcal{M}}_{1,1}$, the self-loop graph Γ_{irr} contributes $\text{tr}_{\text{diag}}(m_{\mathfrak{o}}^{(1)}) = \kappa(\mathcal{A}) \cdot \omega_1$, and integration over $\overline{\mathcal{M}}_{1,1}$ gives $F_1 = \kappa/24$. The only invariant that contributes at genus 1 is κ itself (degree 2 of the MC element); all higher shadow coefficients S_r with $r \geq 3$ are invisible at genus 1.

Uniform-weight factorization. For uniform-weight algebras, the scalar obstruction class at every genus equals $\kappa(\mathcal{A}) \cdot \lambda_g$ (Proposition 6.4). Two verification routes are given there: (A) via Grothendieck–Riemann–Roch on the universal curve, identifying the bar curvature with the Hodge bundle curvature through the Arakelov–Faltings theorem; (B) via clutching propagation from genus 1, using the Whitney sum formula and the injectivity of the trace on the one-dimensional tautological line.

Multi-weight correction. When generators have distinct conformal weights, cross-channel propagators produce mixed terms that do not factor through κ alone. The correction $\delta F_g^{\text{cross}}$ is explicit and computable from the full R -matrix data of the ordered bar.

Remark 6.3 (Genus-1 anchor). The genus-1 formula $\text{obs}_1 = \kappa \cdot \lambda_1$ holds for every chirally Koszul algebra, with no uniform-weight hypothesis and no clutching input. The genus-1 result of Theorem D(i) is independent of the all-genera mechanism.

The passage from genus 1 to all genera is the content of the following proposition, which closes the all-genera extension by two independent routes.

Proposition 6.4 (Scalar obstruction equals Hodge Euler class). *Let $\mathcal{A}$ be a chirally Koszul chiral algebra on the Koszul locus whose strong generators all have the same conformal weight and whose scalar channel is diagonal. Let $\pi : \overline{C}_g \rightarrow \overline{M}_g$ be the universal curve, $\mathbb{E} = R^\circ \pi_* \omega_\pi$ the Hodge bundle of rank g , and $\lambda_g = c_g(\mathbb{E})$ its top Chern class. Then for all $g \geq 1$:*

$$\text{obs}_g(\mathcal{A}) = \kappa(\mathcal{A}) \cdot \lambda_g \quad \text{for scalar-diagonal uniform-weight algebras.}$$

Proof. Two independent proofs are given. Each uses the genus-1 identification $\text{obs}_1 = \kappa \cdot \lambda_1$ as base case and extends to all genera by a different mechanism.

Lemma 6.5 (Uniform-weight scalar factorization of the bar complex; shared input to Proofs A and B). *On the uniform-weight lane, the bar complex of a modular Koszul chiral algebra $\mathcal{A}$ over a fixed smooth projective curve Σ_g factorizes as an external tensor product*

$$\overline{B}(\mathcal{A})|_{\Sigma_g} \simeq \overline{B}^{\text{scalar}}(\mathcal{A}) \otimes_{\mathbb{Q}} \Omega_{\Sigma_g}^\bullet(\log D)|_{\text{unif-wt}},$$

with $\overline{B}^{\text{scalar}}(\mathcal{A})$ a one-dimensional algebraic factor carrying the curvature scalar $\kappa(\mathcal{A})$ (the modular characteristic), and $\Omega_{\Sigma_g}^\bullet(\log D)$ the logarithmic de Rham complex with log-poles on the collision divisor $D \subset \Sigma_g^{\times n}$. As the curve varies over $\overline{M}_g$, the zero-mode sector of the geometric factor is the direct image $R^\circ \pi_ \omega_\pi = \mathbb{E}$ (the Hodge bundle), and the scalar virtual class in $K^\circ(\overline{M}_g) \otimes \mathbb{Q}$ is*

$$[\overline{B}_{\text{scalar}}^{(g)}(\mathcal{A})]^{\text{vir}} = \kappa(\mathcal{A}) \cdot \lambda_{-1}(\mathbb{E}).$$

The obstruction class is the degree- g component of its Chern character:

$$\text{obs}_g(\mathcal{A}) = \kappa(\mathcal{A}) c_g(\mathbb{E}) = \kappa(\mathcal{A}) \lambda_g.$$

Proof. The uniform-weight hypothesis forces every generator of $\mathcal{A}$ to have the same conformal weight, so every OPE propagator reduces to a scalar multiple of the single weight-1 form $d \log E(z, w)$ (the prime form differential). The bar differential is therefore a tensor of an algebraic piece (scalar channel, recording the Casimir coefficient) with a purely geometric piece (log-de-Rham). The zero-mode reduction from the full $\Omega_{\Sigma_g}^\bullet$ to $R^\circ \pi_* \omega_\pi = \mathbb{E}$ is the standard Hodge-theoretic restriction at the constant term. The virtual class equation then follows because the scalar channel contributes the constant $\kappa(\mathcal{A})$ while the geometric channel contributes $\lambda_{-1}(\mathbb{E})$. By the splitting principle, $\text{ch}_g(\lambda_{-1}(\mathbb{E})) = c_g(\mathbb{E})$ after the standard obstruction-orientation convention. $\square$

Remark 6.6 (Two proofs, one input). Lemma 6.5 is the common load-bearing input to the two proofs of Theorem T4 that follow. Proof A derives $\text{obs}_g(\mathcal{A}) = \kappa(\mathcal{A}) \cdot \lambda_g$ from this input by Grothendieck–Riemann–Roch plus the Arakelov–Faltings identification of the Hodge curvature. Proof B derives

it from the same input by clutching propagation from the unconditional genus-1 base case. The two proofs are independent post-input: they invoke disjoint external machinery (GRR/Arakelov vs. Getzler–Kapranov modular operad), and any error in one is detected by the other. The two proofs therefore provide *redundant verification* of the same numerical identity, not *independent derivations* of its structural origin -- the latter is furnished uniquely by Lemma 6.5.

Proof A: Grothendieck–Riemann–Roch on the universal curve. This proof identifies the bar-complex obstruction with a characteristic class of the Hodge bundle directly, without induction on the genus. It takes Lemma 6.5 as input.

Virtual class on moduli. On the uniform-weight lane, the bar complex over a fixed curve Σ_g has a one-dimensional algebraic factor (the scalar channel) tensored with a geometric factor given by the de Rham complex of Σ_g with coefficients in ω_{Σ_g} , because every propagator is the standard weight-1 form $d \log E(z, w)$. As the curve varies over $\overline{\mathcal{M}}_g$, the zero-mode sector of this geometric factor is the direct image $R^\circ \pi_* \omega_\pi = \mathbb{E}$. The scalar virtual class is therefore

$$[\overline{B}_{\text{scalar}}^{(g)}(\mathcal{A})]^{\text{vir}} = \kappa(\mathcal{A}) \cdot \lambda_{-1}(\mathbb{E}) \quad \text{in } K^\circ(\overline{\mathcal{M}}_g) \otimes \mathbb{Q}.$$

Mumford’s GRR computation. Grothendieck–Riemann–Roch on the universal curve gives

$$\text{ch}(R\pi_* \omega_\pi) = \pi_*(\text{ch}(\omega_\pi) \cdot \text{Td}(T_\pi)).$$

The degree- g component of the Chern character of the rank- g bundle $\mathbb{E}$, combined with the Newton identities relating Chern characters to Chern classes, gives

$$c_g(\mathbb{E}) = \lambda_g \in \text{CH}^g(\overline{\mathcal{M}}_g).$$

This is Mumford’s computation [11].

Chern–Weil identification. The Hodge bundle carries the Arakelov metric, whose restriction to each fiber $H^\circ(\Sigma_g, \Omega^1)$ is the L^2 inner product. The Arakelov–Faltings theorem [12] identifies the curvature of $(\mathbb{E}, h_{\text{Ar}})$ with the Arakelov $(1, 1)$ -form:

$$\Theta_{\mathbb{E}}|_{\text{fiber}} = \omega_g^{\text{Ar}} = \frac{i}{2} \sum_{\alpha, \beta=1}^g (\text{Im } \Omega)_{\alpha\beta}^{-1} \omega_\alpha \wedge \overline{\omega}_\beta.$$

The quantum Arnold relations give a curved fiberwise bar package $m_1^2 = [m_o^{(g)}, -]$ whose scalar diagonal projection is $\kappa(\mathcal{A}) \cdot \omega_g^{\text{Ar}}$. The Arakelov–Faltings theorem identifies ω_g^{Ar} with the curvature of the Hodge bundle. Chern–Weil then yields

$$\text{obs}_g(\mathcal{A}) = \text{ch}_g(\kappa(\mathcal{A})\lambda_{-1}(\mathbb{E})) = \kappa(\mathcal{A}) c_g(\mathbb{E}) = \kappa(\mathcal{A}) \cdot \lambda_g.$$

No clutching argument is used. The only external inputs are Mumford’s GRR computation of the Chern classes of $\mathbb{E}$ and the Arakelov–Faltings identification of the Hodge curvature.

Proof B: clutching propagation from genus 1. This proof uses the boundary stratification of $\overline{\mathcal{M}}_g$ and proceeds by induction on g .

Base case. $\text{obs}_1 = \kappa \cdot \lambda_1$ is proved unconditionally (the genus-1 tautological group $H^2(\overline{\mathcal{M}}_{1,1}) \cong \mathbb{Q} \cdot \lambda_1$ is one-dimensional).

Inductive step. Assume $\text{obs}_h = \kappa \cdot \lambda_h$ for all $h < g$. The Maurer–Cartan element $\Theta_{\mathcal{A}} \in \text{MC}(\text{Def}_{\text{cyc}}(\mathcal{A}) \widehat{\otimes} \mathfrak{G}^{\text{mod}})$ automatically satisfies the clutching factorization by the Getzler–Kapranov correspondence [7, Theorem 5.4]: the MC equation on the Feynman transform encodes exactly the clutching compatibility conditions.

The separating clutching $\xi_{\text{sep}}: \overline{\mathcal{M}}_{g_1,1} \times \overline{\mathcal{M}}_{g_2,1} \rightarrow \overline{\mathcal{M}}_g$ (with $g_1 + g_2 = g$) gives

$$\xi_{\text{sep}}^*(\text{obs}_g) = \kappa \cdot \xi_{\text{sep}}^*(\lambda_g),$$

because the scalar channel is one-dimensional and the clutching propagator is normalized before multiplication by the scalar curvature κ . The equality uses the Whitney sum formula $\xi_{\text{sep}}^*(\lambda_g) = \sum_{i+j=g} \lambda_i \otimes \lambda_j$ for the Hodge bundle.

The non-separating clutching $\xi_{\text{nsep}}: \overline{\mathcal{M}}_{g-1,2} \rightarrow \overline{\mathcal{M}}_g$ gives

$$\xi_{\text{nsep}}^*(\text{obs}_g) = \kappa \cdot \xi_{\text{nsep}}^*(\lambda_g),$$

by the inductive hypothesis applied to the genus- $(g-1)$ data at the self-node vertex.

The trace functional (integration over $\overline{\mathcal{M}}_g$) satisfies $\int_{\overline{\mathcal{M}}_g} \text{obs}_g = F_g = \kappa \cdot \hat{A}_g$ (from the bar-complex graph sum on the uniform-weight lane), and $\int_{\overline{\mathcal{M}}_g} \lambda_g = \hat{A}_g$ (the Faber–Pandharipande evaluation). Hence

$$\ell_g(\text{obs}_g) = \kappa \cdot \ell_g(\lambda_g).$$

It remains to show that the joint map $J_g = (\ell_g, \text{clut}_g): \mathcal{W}_g \rightarrow \mathbb{C} \oplus B_g$ is injective on the relevant subspace $U_g \ni \text{obs}_g, \lambda_g$. On the uniform-weight lane, U_g is the one-dimensional tautological line $\mathbb{Q} \cdot \lambda_g$ (the scalar channel produces only classes built from the Hodge bundle; no higher Hodge bundles $R^o \pi_* \omega_\pi^{\otimes h}$ with $h \neq 1$ enter). On this one-dimensional space, the trace ℓ_g is already injective (since $\ell_g(\lambda_g) = \hat{A}_g \neq 0$), so J_g is injective a fortiori.

By the tautological-line support criterion, the injectivity of J_g together with the clutching and trace compatibilities gives $\text{obs}_g = \kappa \cdot \lambda_g$.

Agreement of the two proofs. Proof A produces the identification directly from the Hodge bundle structure on the universal curve. Proof B produces it inductively from the boundary stratification. The two proofs have no logical dependency on each other: Proof A uses GRR and Arakelov–Faltings; Proof B uses Getzler–Kapranov and the tautological-line support criterion. Their agreement is a consistency check on the framework. $\square$

Remark 6.7 (Resolution of the clutching-uniqueness gap). The all-genera extension of Theorem D depends on clutching compatibility together with support on the scalar tautological line. Proof B above establishes this support by showing that the relevant subspace U_g is one-dimensional on the uniform-weight lane (the scalar channel sees only the Hodge bundle), so injectivity of the joint map is automatic. Proof A bypasses the question entirely. The all-genera statement of Theorem D(ii) is proved on the scalar-diagonal uniform-weight lane by either route.

Example 6.8 (Heisenberg at all genera). $\kappa(\mathcal{H}_k) = k$ for the scalar-diagonal single generator of conformal weight 1). The generating function:

$$\sum_{g \geq 1} F_g(\mathcal{H}_k) \cdot \hbar^{2g} = k \cdot \left(\frac{\hbar/2}{\sin(\hbar/2)} - 1 \right) = \frac{k}{24} \hbar^2 + \frac{7k}{5760} \hbar^4 + \frac{31k}{967680} \hbar^6 + \dots$$

The chiral partition function $Z_{\text{ch}}(\mathcal{H}_k) = 1/\eta(\tau)$ (where $\eta(\tau) = q^{1/24} \prod_{n \geq 1} (1 - q^n)$) gives $F_1 = k/24$, consistent with the universal formula.

7 Theorem H: chiral Hochschild cohomology

7.1 Statement

Theorem D identifies a universal genus expansion governed by κ ; the question is what constrains the coefficient ring in which κ lives.

Theorem 7.1 (Theorem H: H_H -conditional chiral Hochschild concentration). *TYPE SIGNATURE. Quadrant: Open. Presentation: chain-level $\text{ChirHoch}^\bullet(A_b) = H^\bullet(\text{CoDer}(\overline{B}(A_b)))$. Level: 3 (derived chiral centre $Z_{\text{ch}}^{\text{der}}(A_b)$, the algebraic closed-sector object in the Beilinson tower). Hypothesis package H_H : PBW chiral Koszulness, finite-type/perfect diagonal Hochschild complexes, genericity, E_∞ -chiral PBW completion, strict Mittag-Leffler passage, and localized residue-twisted bar concentration.*

Let $\mathcal{A} = A_b$ lie in the standard PBW chiral Koszul locus with the full package H_H , away from critical and non-generic loci. The chiral Hochschild cohomology is concentrated in degrees $\{0, 1, 2\}$:

$$\text{ChirHoch}^n(\mathcal{A}) = 0 \quad \text{for } n \notin \{0, 1, 2\},$$

and the Hilbert polynomial is

$$P_{\mathcal{A}}(t) = \dim Z(\mathcal{A}) + \dim \text{ChirHoch}^1(\mathcal{A}) \cdot t + \dim Z(\mathcal{A}^1) \cdot t^2. \quad (3)$$

The Verdier intertwining of Theorem A pairs degree 0 with degree 2, identifying $Z(\mathcal{A}^1) = \text{Hom}(Z(\mathcal{A}), k)$ on the H_H surface. The Feigin–Frenkel companion at $k = -h^\vee$ (critical level) is outside the regime: the centre is infinite-dimensional, and the $\{0, 1, 2\}$ concentration fails in any ambient.

Remark 7.2 (Three Hochschild theories: chiral, topological, categorical). Three inequivalent Hochschild cohomology theories coexist: chiral, topological, and categorical. *Chiral Hochschild* $\text{ChirHoch}^*(\mathcal{A}) = H^*(\text{CoDer}(\overline{B}(\mathcal{A})))$ is the cohomology of coderivations of the bar coalgebra; it lives on the curve X and has amplitude $[0, 2]$ under the H_H package of Theorem H. *Topological Hochschild* HH_{top}^* is factorization homology on S^1 ; it lives in one dimension higher. *Categorical Hochschild* $\text{HH}^*(\text{Rep}(\mathcal{A}))$ governs the Drinfeld center and the E_2 structure. Conflating any two produces errors.

7.2 Family-specific specializations

Proposition 7.3 (Hilbert series for standard families). (i) Heisenberg $\mathcal{H}_k$: $P(t) = 1 + t + t^2$. *Centre = $\mathbb{C}$, outer derivation $D(J) = 1$, level deformation in degree 2.*

(ii) Affine Kac–Moody $V_k(\mathfrak{g})$ (generic level): *Under the affine H_H package, after the zero-mode inner quotient, $\text{ChirHoch}^1(V_k(\mathfrak{g})) = 0$ and $P_{V_k(\mathfrak{g})}(t) = 1 + t^2$. The adjoint $\mathfrak{g}$ is retained as zero-mode derivation prequotient and bar-dual generator count.*

(iii) Virasoro Vir_c (generic): $P(t) = 1 + t^2$. $\text{ChirHoch}^1 = 0$ (no outer chiral derivations).

(iv) Principal $\mathcal{W}_N$ (generic): $P(t) = 1 + t^2$.

(v) Critical level $k = -h^\vee$: *Outside the regime of Theorem H. The Feigin–Frenkel center makes the cohomology unbounded.*

7.3 Proof architecture

The concentration proof factors through the PBW Koszulity criterion and the completed chiral Hochschild model. Bar concentration and Fulton–MacPherson tower collapse are parallel consequences of PBW collapse. The identification of the Hilbert polynomial with the dimensions of the centre and its dual follows from the Verdier intertwining pairing degree 0 with degree 2.

The configuration-space collapse uses Fulton–MacPherson formality: the spectral sequence from the FM filtration on the coderivation complex collapses, and the E_2 page identifies with the cohomology of the classical coderivation complex of the associated graded.

8 The dependency graph among Theorems A, B, C, D, and H

The five theorems are not five unrelated assertions about the bar complex. They form a dependency graph: each theorem supplies one load-bearing object or comparison used by at least one later theorem, but the graph is not a total order.

8.1 The dependency DAG

The logical dependencies are:

$$\boxed{A} \xrightarrow{\text{dual object}} \boxed{B} \quad \boxed{B} \longrightarrow \boxed{H}, \quad \boxed{B} \longrightarrow \boxed{C}, \quad \boxed{B} \longrightarrow \boxed{D}.$$

Theorem C and Theorem D use overlapping bar, duality, and scalar inputs, but Theorem D is not a theorem-level consequence of Theorem C. Theorem H uses PBW/Orlik–Solomon input together with the completed Hochschild model.

8.2 A creates the arena

Proposition 8.1 (*A is necessary for B–H*). *Without Theorem A, the objects of Theorems B–H do not exist.*

Proof. Theorem B asserts that the bar-cobar counit $\varepsilon: \Omega(\overline{B}(\mathcal{A})) \rightarrow \mathcal{A}$ is a quasi-isomorphism. The cobar functor Ω is defined on conilpotent coalgebras in the factorization category $\text{Fact}(X)$; its existence and the adjunction with $\overline{B}$ are the content of Theorem A(i). Without the adjunction, the counit ε is undefined.

Theorem C decomposes $H^*(\overline{M}_g, \mathcal{Z}_{\mathcal{A}}) \cong Q_g(\mathcal{A}) \oplus Q_g(\mathcal{A}^!)$. The summand $Q_g(\mathcal{A}^!)$ involves the Koszul dual $\mathcal{A}_{\infty}^!$, whose existence is Theorem A(ii): the Verdier intertwining $\mathbb{D}_{\text{Ran}} \overline{B}(\mathcal{A}) \simeq \mathcal{A}_{\infty}^{!,\text{co}}$ and $\Omega_X^{\text{cont}} \mathcal{A}_{\infty}^{!,\text{co}} \simeq \mathcal{A}_{\infty}^!$. Without this intertwining, the dual object is unavailable.

Theorems D and H depend on C and B respectively; their dependence on A is inherited. $\square$

The geometric input that Theorem A requires, beyond the bar complex itself, is the Fulton–MacPherson compactification $\overline{\text{FM}}_n(X)$ and its boundary divisor structure: the twisting morphism τ is mediated by integration kernels on $\overline{\text{FM}}_n(X)$, and the Verdier intertwining uses the Poincaré–Verdier duality on $\text{Ran}(X)$.

8.3 B establishes faithfulness

Proposition 8.2 (B is necessary for C and strict-dual identification). *Without Theorem B, the Lagrangian decomposition of Theorem C is vacuous and the relation between the homotopy dual and the strict Koszul dual is ungrounded.*

Proof. Theorem A constructs the homotopy Koszul dual $\mathcal{A}'_\infty$, but the construction produces a factorization algebra that a priori differs from $\mathcal{A}'$. Theorem B proves that bar-cobar inversion is faithful: $\overline{B}$ is injective on quasi-isomorphism classes. This identifies $\mathcal{A}$ and $\mathcal{A}'$ as the two canonical algebras recoverable from the bar coalgebra, and establishes that Verdier duality on $\overline{B}(\mathcal{A})$ produces the *correct* dual.

Without this identification, the decomposition $Q_g(\mathcal{A}) \oplus Q_g(\mathcal{A}')$ in Theorem C involves an object $\mathcal{A}'_\infty$ whose relation to the strict dual $\mathcal{A}'$ is uncontrolled: the summands may not be complementary. Theorem D is bar-intrinsic and does not use this identification to produce the scalar genus tower, but the strict-dual interpretation of complementarity and the downstream duality package do require it. $\square$

The geometric input is the *PBW filtration*: the spectral sequence from conformal-weight filtration collapses at E_2 , reducing the problem to classical Koszul duality of the associated graded. This is the only step that involves the conformal weight grading; Theorem A uses only the chiral bracket and $\mathcal{D}_X$ -module structure.

8.4 C provides the polarization

Proposition 8.3 (C is necessary for the ambient polarization package). *Without Theorem C, the genus tower has no canonical factorization into $\mathcal{A}$ and $\mathcal{A}'$ contributions, and the complementarity interpretation of the scalar genus tower is unavailable.*

Proof. At genus $g \geq 1$, the bar differential is curved with scalar diagonal curvature $\kappa \cdot \omega_g$, and the Maurer–Cartan element $\Theta_{\mathcal{A}}$ encodes all genus corrections. But the ambient complex $\mathbf{C}_g(\mathcal{A}) = R\Gamma(\overline{\mathcal{M}}_g, \mathcal{Z}_{\mathcal{A}})$ receives contributions from both $\mathcal{A}$ and its dual. Without the Lagrangian decomposition, these contributions are entangled: one loses the canonical ambient polarization into $\mathcal{A}$ and $\mathcal{A}'$ sectors.

Theorem C provides the polarization: the Verdier involution decomposes $\mathbf{C}_g$ into eigenspaces, and the scalar projection of this decomposition yields the complementarity constraint $\kappa + \kappa' = \rho_{\mathcal{A}} \cdot K$ (family-specific $\rho_{\mathcal{A}}$: 0 for Kac–Moody and free fields, $1/2$ for Virasoro, $H_N - 1$ for principal $\mathcal{W}_N$, $1/6$ for Bershadsky–Polyakov). Theorem D still extracts the universal scalar shadow bar-intrinsically from the genus-1 curvature together with clutching/GRR propagation; Theorem C explains how that scalar sits inside the ambient complex and how duality constrains it. $\square$

The geometric input is the *Verdier involution*: the existence of a functorial duality on sheaves on $\overline{\mathcal{M}}_g$ that exchanges the bar complex of $\mathcal{A}$ with that of $\mathcal{A}'$. This requires the moduli space $\overline{\mathcal{M}}_g$ as a smooth Deligne–Mumford stack with Poincaré duality.

8.5 D identifies the scalar shadow

Proposition 8.4 (D is necessary for H). *Without Theorem D, the coefficient ring of the genus tower is unidentified, and Theorem H cannot distinguish polynomial behaviour from more complex coefficient growth.*

Proof. The genus expansion $\sum_g F_g(\mathcal{A}) \cdot \hbar^{2g} = \kappa \cdot (\hat{A}(i\hbar) - 1)$ shows that all genus invariants are determined by a single number κ . The Maurer–Cartan element $\Theta_{\mathcal{A}}$ lives in the convolution algebra $\mathfrak{g}_{\mathcal{A}}^{\text{mod}}$, whose coefficient ring is the chiral Hochschild cohomology $\text{ChirHoch}^*(\mathcal{A})$. Theorem D demonstrates that the genus tower is not an independent sequence of invariants but a single series governed by κ ; this forces the question: what constrains κ itself?

The answer is Theorem H. The modular characteristic κ lives in degree 2 of $\text{ChirHoch}^*(\mathcal{A})$. The deformations of κ are parametrized by $\text{ChirHoch}^2(\mathcal{A})$; the obstructions are in $\text{ChirHoch}^3(\mathcal{A})$. Under the full H_H package, Theorem H gives $\text{ChirHoch}^3 = 0$; on that surface the deformation theory is unobstructed and the coefficient ring is polynomial. Without Theorem D’s identification of κ as the universal scalar, one would not know that the coefficient ring has a single generator in degree 2; one would only know that genus invariants exist. $\square$

The geometric input is *Faber–Pandharipande integration*: the evaluation of λ_g on $\overline{\mathcal{M}}_g$ produces the numerical coefficients $\hat{A}_g$, and the factorization through stable graphs uses the clutching morphisms on the boundary of $\overline{\mathcal{M}}_g$.

8.6 The non-circular routing

The chain $A \Rightarrow B \Rightarrow C \Rightarrow D \Rightarrow H$ might appear circular: Theorem A uses the bar complex, whose higher-genus structure involves κ (which is computed in Theorem D). The resolution is that the chain has a non-circular anchor.

Remark 8.5 (Primitive non-circular anchor). The bar complex $\overline{B}(\mathcal{A}) = T^c(\mathcal{A})$ and the Maurer–Cartan element $\Theta_{\mathcal{A}} = D_{\mathcal{A}} - d_0$ are defined intrinsically, without reference to any theorem. The MC equation $d_0 \Theta + \frac{1}{2}[\Theta, \Theta] = 0$ is a tautology of the bar differential $D_{\mathcal{A}}^2 = 0$.

Theorem A uses only genus-0 data: the twisting morphism and the Verdier functor on $\text{Ran}(X)$. Theorem B uses the PBW filtration at genus 0. Theorem C uses the Verdier involution on $\overline{\mathcal{M}}_g$ applied to the objects constructed in A and identified in B. Theorem D uses the bar-intrinsic scalar trace of $\Theta_{\mathcal{A}}$, the genus-1 curvature normalization, and the clutching/GRR propagation to identify the scalar shadow κ , which was already present in the definition of $\Theta_{\mathcal{A}}$ but not yet identified as universal. Theorem C gives a complementary ambient interpretation of the same scalar, but is not the load-bearing input to Theorem D. Theorem H uses the concentration result from Koszul duality (Theorem B) together with the scalar package of Theorem D to bound the coderivation complex.

The routing is: bar-intrinsic MC element (definition) $\rightarrow$ adjunction/intertwining (A), then two downstream branches. One branch is inversion (B) $\rightarrow$ Verdier polarization (C). The other is scalar trace + genus-1 curvature + clutching/GRR $\rightarrow$ scalar extraction (D). Theorem H reads the coefficient ring against these outputs. No step requires a result proved later.

8.7 Independence of converses

The converses of the five implications fail. This is not a deficiency; it reflects the fact that each theorem addresses a strictly different aspect of the bar complex.

- (i) *H does not imply D.* A polynomial Hilbert series $P(t) = 1 + t^2$ (Virasoro) does not determine $\kappa = c/2$; it constrains $\dim \text{ChirHoch}^2 = 1$ (the space where κ lives has dimension 1) but not the value.

- (ii) *D does not imply C.* The formula $\text{obs}_g = \kappa \cdot \lambda_g$ could hold without a Lagrangian decomposition: one could imagine a non-decomposable ambient complex whose scalar projection still factors through κ . Theorem C is stronger; it provides a complementary ambient interpretation of the factorization, but not the load-bearing proof of Theorem D.
- (iii) *C does not imply B.* The Lagrangian decomposition $Q_g(\mathcal{A}) \oplus Q_g(\mathcal{A}^!)$ is meaningful even if bar-cobar inversion fails; one could have a Lagrangian complement indexed by a coalgebra that does not reconstruct $\mathcal{A}$. Inversion is a separate condition.
- (iv) *B does not imply A.* One could define bar-cobar inversion without the adjunction formalism: the counit ε could be a quasi-isomorphism without being part of an adjoint pair. The adjunction gives functoriality; inversion alone does not.

8.8 Summary: one new input per step

Theorem	Output	New input	Problem solved
A	dual object $\mathcal{A}_\infty^!$	FM boundary	no dual without adjunction
B	inversion $\Omega(\overline{B}(\mathcal{A})) \simeq \mathcal{A}$	PBW filtration	bar could lose boundary
C	$Q_g(\mathcal{A}) \oplus Q_g(\mathcal{A}^!)$	Verdier involution on $\overline{\mathcal{M}}_g$	no canonical polarization
D	$\text{obs}_g = \kappa \cdot \lambda_g$	Faber–Pandharipande	no distinguished scalar
H	H_H -conditional $P(t) = 1 + t^2$ after the zero-mode inner quotient	coderivation complex	coefficient ring uncontractible

Each row adds exactly one geometric tool to the preceding rows. The five-step chain is the minimum path from the bare bar complex to the complete structural description of the modular Koszul dual.

9 The Heisenberg: complete computation

The Heisenberg algebra $\mathcal{H}_k$ is the simplest chiral algebra that is not a Lie algebra. It has one generator, one double pole, and no simple pole. Every clause of every theorem is computable in closed form. This section carries out the computation: the five theorems verified in a single place, with no step deferred.

9.1 The algebra

The Heisenberg chiral algebra $\mathcal{H}_k$ at level $k \in \mathbf{C}$ is generated by a single current J of conformal weight 1, with the operator product expansion

$$J(z)J(w) \sim \frac{k}{(z-w)^2}.$$

One generator, one OPE pole of order 2, no simple pole. The zeroth product $J_{(0)}J = 0$ (no Lie bracket: the algebra is abelian). The first product $J_{(1)}J = k$ (the level is the inner product). The augmentation ideal is $\overline{\mathcal{H}}_k = \mathbf{C} \cdot J$, one-dimensional.

The classical R -matrix is scalar:

$$r^{\mathcal{H}}(z) = \frac{k}{z},$$

obtained from the OPE double pole by d log absorption: $\text{pole}_r = \text{pole}_{\text{OPE}} - 1 = 2 - 1 = 1$. The modular characteristic is $\kappa(\mathcal{H}_k) = \text{av}(r^{\mathcal{H}}(z)) = k$. Since $r(z)$ is already scalar, the averaging map is the identity:

no matrix structure to erase, no Casimir tensor to contract. The Heisenberg is the unique family where the ordered and symmetric bars carry exactly the same information.

9.2 Theorem A for $\mathcal{H}_k$: bar-cobar adjunction

The bar complex is the cofree coalgebra on the desuspended augmentation ideal:

$$\overline{B}(\mathcal{H}_k) = T^c(s^{-1}\mathbf{C} \cdot J) = \bigoplus_{n \geq 1} (s^{-1}J)^{\otimes n}, \quad |s^{-1}J| = 1 - 1 = 0.$$

The bar differential at degree 2 is the complete computation:

$$d_{\overline{B}}[s^{-1}J | s^{-1}J] = \text{Res}_{z_1=z_2} \left[\frac{k}{(z_1 - z_2)^2} \cdot d \log(z_1 - z_2) \right] = k.$$

At degree $n \geq 3$, the bar differential involves iterated residues that reduce to degree $n - 1$ by the Arnold relation. The bar cohomology concentrates in bar degree 1:

$$H^*(\overline{B}(\mathcal{H}_k)) = \mathbf{C} \cdot s^{-1}J \quad (\text{degree } -1).$$

The Koszul dual coalgebra is $\mathcal{H}_k^! = \mathbf{C} \cdot s^{-1}J$ (one-dimensional, in degree -1). The strict Koszul dual is the curved symmetric chiral algebra

$$\mathcal{H}_k^! = \text{Sym}^{\text{ch}}(V^*), \quad m_o = -k\omega,$$

with curvature $m_o = -k\omega$ encoding the dual level. The Verdier intertwining of Theorem A(ii) recovers $\mathcal{H}_k^!$ at the cohomological level: $\mathbb{D}_{\text{Ran}} \overline{B}(\mathcal{H}_k) \simeq (\mathcal{H}_k^!)_{\infty}$.

9.3 Theorem B for $\mathcal{H}_k$: bar-cobar inversion

The cobar of the bar recovers the original algebra:

$$\Omega(\overline{B}(\mathcal{H}_k)) \xrightarrow{\sim} \mathcal{H}_k.$$

The PBW spectral sequence collapses at E_2 because the bar cohomology is already concentrated in degree 1: the associated graded is quadratic Koszul in the classical sense (the relation space is one-dimensional, generated by $J \otimes J - k$).

Bar-cobar inversion for $\mathcal{H}_k$ is *unconditional at all genera*. At genus $g \geq 1$, the fiberwise bar differential satisfies $d_{\text{fib}}^2 = k \cdot \omega_g$ with scalar curvature. The Ext vanishing condition of Theorem 4.1(ii) is automatic: the OPE has a single double pole and no simple pole, so there are no Lie brackets to obstruct and no higher-order collisions to produce non-trivial Ext groups. The absence of a simple pole ($J_{(o)}J = 0$) is the algebraic reason that $\mathcal{H}_k$ is the simplest non-trivial test case for every theorem in the programme.

9.4 Theorem C for $\mathcal{H}_k$: complementarity

The Koszul dual has modular characteristic $\kappa(\mathcal{H}_k^!) = -k$: the curvature $m_o = -k\omega$ encodes the opposite level. The complementarity sum is

$$\kappa(\mathcal{H}_k) + \kappa(\mathcal{H}_k^!) = k + (-k) = 0 = K(\mathcal{H}_k).$$

The Koszul complementarity conductor $K = 0$: the Heisenberg is in the class of algebras where duality reverses the sign of κ .

The Lagrangian decomposition of Theorem 5.1 is visible by inspection. The center local system $\mathcal{Z}_{\mathcal{H}_k}$ on $\overline{\mathcal{M}}_g$ is the constant sheaf (the center of $\mathcal{H}_k$ is $\mathbf{C}$, with no monodromy). The ambient complex $\mathbf{C}_g(\mathcal{H}_k) = R\Gamma(\overline{\mathcal{M}}_g, \mathcal{Z}_{\mathcal{H}_k}) = H^*(\overline{\mathcal{M}}_g, \mathbf{C})$ decomposes into the Hodge eigenspaces of the Verdier involution, and the two Lagrangian summands $\mathcal{Q}_g(\mathcal{H}_k)$ and $\mathcal{Q}_g(\mathcal{H}_k^\dagger)$ are exchanged by $\kappa \mapsto -\kappa$.

9.5 Theorem D for $\mathcal{H}_k$: the genus expansion

The Heisenberg is a uniform-weight algebra (single generator of conformal weight 1), so the genus expansion takes the unconditional form

$$\text{obs}_g(\mathcal{H}_k) = k \cdot \lambda_g \quad (g \geq 1) \quad \text{for this uniform-weight algebra.}$$

No cross-channel correction: $\partial F_g^{\text{cross}} = 0$ because there is only one conformal weight.

The genus-1 free energy:

$$F_1(\mathcal{H}_k) = \frac{k}{24}.$$

The full generating function is

$$\sum_{g \geq 1} F_g(\mathcal{H}_k) \cdot \hbar^{2g} = k \cdot \left(\frac{\hbar/2}{\sin(\hbar/2)} - 1 \right) = \frac{k}{24} \hbar^2 + \frac{7k}{5760} \hbar^4 + \frac{31k}{967680} \hbar^6 + \dots$$

with coefficients $F_g(\mathcal{H}_k) = k \cdot \frac{2^{2g-1}-1}{2^{2g-1}} \cdot \frac{|B_{2g}|}{(2g)!}$, where B_{2g} is the $2g$ -th Bernoulli number.

Remark 9.1 (The chiral partition function). The genus-1 chiral partition function of $\mathcal{H}_k$ is

$$Z_{\text{ch}}(\mathcal{H}_k) = \frac{1}{\eta(\tau)}, \quad \eta(\tau) = q^{1/24} \prod_{n \geq 1} (1 - q^n),$$

where $q = e^{2\pi i \tau}$. The leading exponent $q^{-1/24}$ encodes $F_1 = k/24$ (at $k = 1$). The product form is the Euler function; its reciprocal counts states at each energy level.

Remark 9.2 (R -matrix and averaging). The R -matrix $r^{\mathcal{H}}(z) = k/z$ is already scalar, so $\text{av}(r^{\mathcal{H}}(z)) = k = \kappa(\mathcal{H}_k)$ with no information lost. For every other family, the averaging map av compresses a matrix-valued function of z into a single number. The Heisenberg is the unique base case where the compression is lossless. For non-abelian affine Kac–Moody, the averaging discards the Casimir tensor Ω and the z -dependence; for the Virasoro algebra, it discards the stress tensor T and the cubic pole structure. The ordered bar sees the full R -matrix; the symmetric bar sees only κ .

9.6 Theorem H for $\mathcal{H}_k$: chiral Hochschild cohomology

The chiral Hochschild cohomology of $\mathcal{H}_k$ is three-dimensional:

$$\text{ChirHoch}^n(\mathcal{H}_k) = \begin{cases} \mathbf{C} & n = 0, \\ \mathbf{C} & n = 1, \\ \mathbf{C} & n = 2, \\ 0 & n \geq 3. \end{cases}$$

The Hilbert polynomial is $P_{\mathcal{H}_k}(t) = 1 + t + t^2$.

The three generators have transparent algebraic meaning:

- (i) *Degree 0: the center.* $\text{ChirHoch}^0(\mathcal{H}_k) = Z(\mathcal{H}_k) = \mathbf{C}$. The center consists of scalars; J has no zeroth product ($J_{(0)} = 0$), so J itself is central, but J is not a scalar; the degree-0 chiral Hochschild class is the identity.
- (ii) *Degree 1: the outer derivation.* $\text{ChirHoch}^1(\mathcal{H}_k) = \mathbf{C}$, generated by the derivation $D: J \mapsto 1$ that maps the current to the vacuum. This derivation is outer: it is not of the form $a \mapsto a_{(0)}b$ for any $b \in \mathcal{H}_k$, because $J_{(0)} = 0$ (no simple pole in the OPE). The existence of an outer derivation is the Heisenberg's distinguishing feature relative to the Virasoro algebra, where $\text{ChirHoch}^1 = 0$.
- (iii) *Degree 2: the level deformation.* $\text{ChirHoch}^2(\mathcal{H}_k) = \mathbf{C}$, generated by the infinitesimal deformation of the OPE coefficient $k \mapsto k + \epsilon$. This is the dual of the center: the Verdier pairing identifies $\text{ChirHoch}^2(\mathcal{H}_k) \cong Z(\mathcal{H}_k^!)^\vee = \mathbf{C}$, consistent with the general formula (3).

9.7 The shadow tower and the derived center

The shadow tower $\{S_r(\mathcal{H}_k)\}_{r \geq 2}$ terminates at $r = 2$:

$$S_2 = \kappa = k, \quad S_r = 0 \text{ for all } r \geq 3.$$

The critical discriminant $\Delta = 8\kappa S_4 = 0$: the Heisenberg is class G (Gaussian), the simplest archetype. The shadow tower carries no information beyond κ itself.

The derived chiral center is $\mathcal{Z}_{\text{ch}}^{\text{der}}(\mathcal{H}_k) = \text{ChirHoch}^*(\mathcal{H}_k) = \mathbf{C} \oplus \mathbf{C}[1] \oplus \mathbf{C}[2]$, three-dimensional ($k \neq 0$). This is strictly larger than the naive center $Z(\mathcal{H}_k) = \mathbf{C}$: the Ext^1 and Ext^2 contributions are invisible to the commutant.

Remark 9.3 (Abelian Chern–Simons). The Heisenberg algebra $\mathcal{H}_k$ is the boundary algebra of abelian Chern–Simons theory at level k : the 3-dimensional bulk is $U(1)_k$ Chern–Simons on a 3-manifold M^3 with $\partial M^3 = \Sigma$. The modular characteristic $\kappa = k$ is the Chern–Simons level; the genus expansion $F_g = k \cdot \hat{A}_g$ is the perturbative expansion of the Chern–Simons partition function. The complementarity sum $K = 0$ reflects the topological nature of the theory: bulk and boundary carry opposite orientations.

9.8 Summary: the Heisenberg as base computation

Theorem	Statement for $\mathcal{H}_k$	Key simplification
A	$\overline{B}(\mathcal{H}_k) = T^c(s^{-1}\mathbf{C} \cdot J)$, $\mathcal{H}_k^! = (\text{Sym}^{\text{ch}}(V^*[1]), m_0 = -k\omega)$	one generator, degree-1 concentration
B	$\Omega(\overline{B}(\mathcal{H}_k)) \simeq \mathcal{H}_k$, all genera	no Lie bracket, Ext automatic
C	$\kappa + \kappa' = 0$	Verdier reverses sign
D	$F_g = k \cdot \hat{A}_g$ for this uniform-weight algebra	single weight, no cross-channel
H	$P(t) = 1 + t + t^2$	outer derivation from abelian OPE

Every other family is a deformation of this computation. Affine Kac–Moody adds a simple pole (Lie bracket); the Virasoro algebra adds a quartic pole (stress tensor) and the cubic R -matrix structure; the $\mathcal{W}$ -algebras add higher-spin generators and multi-channel interactions. The five theorems persist, but their proofs become harder: bar cohomology is still concentrated in degree 1 (Koszul), but the spectral sequence argument requires PBW-type tools; the complementarity sum is no longer zero; the genus

expansion acquires cross-channel corrections for multi-weight algebras; the chiral Hochschild loses its degree-1 component. The Heisenberg computation is the base case that makes the pattern visible.

10 The standard landscape

The five theorems hold for every chirally Koszul chiral algebra. The *standard landscape* is the collection of families for which every invariant has been computed: the modular characteristic κ , the Koszul complementarity conductor K , the critical discriminant Δ , the complexity class, the shadow depth $r_{\max}$, and the classical R -matrix. This section records the census.

10.1 The census table

In the table: $H_N = \sum_{j=1}^N 1/j$ is the N -th harmonic number; $K = \kappa(\mathcal{A}) + \kappa(\mathcal{A}^\dagger)$ is the Koszul complementarity conductor; $\Delta = 8\kappa S_4$ is the critical discriminant; $r_{\max}$ is the shadow depth (the highest degree at which $S_r \neq 0$, or ∞ if the tower does not terminate).

10.2 The general affine Kac–Moody formula

For a simple Lie algebra $\mathfrak{g}$ of dimension $d = \dim \mathfrak{g}$ and dual Coxeter number $h^\vee$, the affine Kac–Moody vertex algebra $V_k(\mathfrak{g})$ at level k has

$$\kappa(V_k(\mathfrak{g})) = \frac{d \cdot (k + h^\vee)}{2h^\vee}.$$

The formula is derived from the trace-form classical R -matrix $r(z) = k \Omega_{\text{tr}}/z$:

$$\text{av}(r(z)) = \frac{k \cdot d}{2h^\vee} = \kappa - \frac{d}{2},$$

where the shift $d/2$ is the Sugawara contribution from the simple-pole self-contraction through the adjoint Casimir eigenvalue $2h^\vee$. The full modular characteristic is $\kappa = \text{av}(r(z)) + d/2 = d(k + h^\vee)/(2h^\vee)$.

Two boundary checks:

- (i) At $k = 0$ (abelian limit): $\kappa(V_0(\mathfrak{g})) = d/2 \neq 0$. The R -matrix $r(z) = 0$ vanishes, but the Sugawara shift persists.
- (ii) At $k = -h^\vee$ (critical level): $\kappa(V_{-h^\vee}(\mathfrak{g})) = 0$. All monodromy becomes trivial; the Feigin–Frenkel center appears; Koszulness fails.

The affine scalar companion is obtained by the critical-level reflection $k \mapsto -k - 2h^\vee$; the chiral Koszul dual still passes through the Verdier-dual bar and chiral CE lane. Consequently $\kappa(V_k(\mathfrak{g})^\dagger) = -d(k + h^\vee)/(2h^\vee) = -\kappa$, so $K = 0$ for all affine Kac–Moody algebras. The complementarity sum $c + c'$ is $2d$, independent of k .

	$\mathfrak{g}$	d	$h^\vee$	κ	$c + c'$	$\kappa _{k=0}$
Example 10.1 (Specific Lie algebras).	$\mathfrak{sl}_2$	3	2	$3(k+2)/4$	6	$3/2$
	$\mathfrak{sl}_3$	8	3	$4(k+3)/3$	16	4
	$\mathfrak{so}_8$	28	6	$7(k+6)/3$	56	14
	G_2	14	4	$7(k+4)/4$	28	7
	F_4	52	9	$26(k+9)/9$	104	26
	E_6	78	12	$13(k+12)/4$	156	39
	E_7	133	18	$133(k+18)/36$	266	$133/2$
	E_8	248	30	$62(k+30)/15$	496	124

10.3 The R -matrix census

The classical R -matrix $r_{\mathcal{A}}(z) = \text{Res}_{0,2}^{\text{coll}}(\Theta_{\mathcal{A}})$ is the binary collision residue. Its pole structure is determined by the OPE pole order via d log absorption:

$$\text{pole}_r = \text{pole}_{\text{OPE}} - 1.$$

Family	OPE pole	r -matrix pole	$r(z)$
$\mathcal{H}_k$	2	1	k/z (scalar)
$V_k(\mathfrak{g})$	2	1	$k \Omega_{\text{tr}}/z$ (matrix, trace-form)
Vir_c	4	3	$(c/2)/z^3 + 2T/z$ (cubic + simple)
$\mathcal{W}_N$	$2N$	$2N - 1$	matrix-valued, $N-1$ channels
$\beta\gamma$	1	0	constant
bc	1	0	Clifford-valued

The Virasoro R -matrix has a cubic pole and a simple pole. The cubic pole encodes the central charge; the simple pole encodes the stress tensor. It is *not* quartic: the quartic pole of the $T(z)T(w)$ OPE is absorbed by d log.

10.4 The four complexity classes

The shadow obstruction tower $\{S_r\}_{r \geq 2}$ partitions the landscape into four classes. The partition is determined by two invariants: S_4 (the quartic shadow coefficient) and κ .

Definition 10.2 (Complexity class). Let $\Delta = 8\kappa S_4$ be the critical discriminant. The four classes are:

- (a) **Class G** (Gaussian): $S_r = 0$ for all $r \geq 3$. Shadow depth $r_{\max} = 2$. The shadow tower carries only κ ; no nonlinear corrections. $\Delta = 0$ because $S_4 = 0$.
- (b) **Class L** (Lie): $S_3 \neq 0, S_4 = 0$. Shadow depth $r_{\max} = 3$. A single Massey product appears at degree 3 (the Lie bracket), then the tower terminates. $\Delta = 0$ because $S_4 = 0$.
- (c) **Class C** (Contact): $S_3 = 0, S_4 \neq 0$, but $\kappa = 0$ or special cancellation gives $\Delta = 0$. Shadow depth $r_{\max} = 4$. The $\beta\gamma$ system is the witness: $r_{\max} = 4$ but $S_r = 0$ for $r \geq 5$.
- (d) **Class M** (Mixed): $\Delta \neq 0$. Shadow depth $r_{\max} = \infty$. The tower does not terminate; every S_r is nonzero. The Virasoro algebra and all principal $\mathcal{W}_N$ -algebras are class M.

The critical discriminant Δ controls the downstream depth on a fixed primary line. It is not a global classifier by itself. The G/L/C/M taxonomy is fixed by the shadow tower together with the family-specific collision data.

Class	S_3	S_4	Δ	Families
G	o	o	o	Heisenberg, lattice, free fermion
L	$\neq$ o	o	o	Affine Kac–Moody (all $\mathfrak{g}$, generic k)
C	o	$\neq$ o	o	$\beta\gamma$
M	$\neq$ o	$\neq$ o	$\neq$ o	Virasoro, $\mathcal{W}_N$, Bershadsky–Polyakov

10.5 Free-field central charges

The free-field families provide essential boundary data.

Fermionic bc ghosts (conformal weights λ and $1 - \lambda$):

$$c_{bc}(\lambda) = 1 - 3(2\lambda - 1)^2.$$

At $\lambda = 1/2$: $c_{bc} = 1$ (a single Dirac fermion). At $\lambda = 2$: $c_{bc} = -26$ (the reparametrization ghosts of the bosonic string).

Bosonic $\beta\gamma$ system (conformal weights λ and $1 - \lambda$):

$$c_{\beta\gamma}(\lambda) = 2(6\lambda^2 - 6\lambda + 1).$$

At $\lambda = 1/2$: $c_{\beta\gamma} = -1$ (the symplectic boson). At $\lambda = 2$: $c_{\beta\gamma} = 26$.

The complementarity relation

$$c_{\beta\gamma}(\lambda) + c_{bc}(\lambda) = 0 \quad \text{for all } \lambda$$

is the ghost cancellation: bosonic and fermionic partners contribute opposite central charges. At $\lambda = 2$, this is the string-theoretic statement $26 + (-26) = 0$.

10.6 The Virasoro self-dual point

Under Koszul duality, the Virasoro algebra satisfies $c \mapsto 26 - c$. The modular characteristic:

$$\kappa(\text{Vir}_c) = c/2, \quad \kappa(\text{Vir}_{26-c}) = (26 - c)/2.$$

The complementarity sum is $K = c/2 + (26 - c)/2 = 13$, constant and independent of c . The self-dual point is

$$c = 13: \quad \kappa = \kappa' = 13/2.$$

This is the unique point where the Lagrangian decomposition of Theorem 5.1 is symmetric: the two summands $Q_g(\text{Vir}_{13})$ and $Q_g(\text{Vir}_{13}^!)$ are isomorphic.

Remark 10.3 (The self-dual point is $c = 13$, not $c = 26$). The complementarity sum $c + c' = 26$ involves both central charges; the self-dual *point* is $c = c' = 13$, the fixed point of $c \mapsto 26 - c$. Confusing the sum with the fixed point is a persistent error.

10.7 The $\mathcal{W}_N$ family

The principal $\mathcal{W}$ -algebra $\mathcal{W}_N$ is the Drinfeld–Sokolov reduction of $\widehat{\mathfrak{sl}}_N$. It has $N - 1$ strong generators of conformal weights $2, 3, \dots, N$. The modular characteristic:

$$\kappa(\mathcal{W}_N) = c \cdot (H_N - 1), \quad H_N = \sum_{j=1}^N \frac{1}{j}.$$

At $N = 2$: $H_2 - 1 = 3/2 - 1 = 1/2$, so $\kappa(\mathcal{W}_2) = c/2 = \kappa(\text{Vir}_c)$, consistent with $\mathcal{W}_2 = \text{Vir}$. At $N = 3$: $H_3 - 1 = 1 + 1/2 + 1/3 - 1 = 5/6$, so $\kappa(\mathcal{W}_3) = 5c/6$.

All $\mathcal{W}_N$ -algebras with $N \geq 2$ are class M: the shadow tower is infinite and the critical discriminant $\Delta \neq 0$. As multi-weight algebras (generators at weights $2, \dots, N$), at genus $g \geq 2$ the full free energy acquires a cross-channel correction $\delta F_g^{\text{cross}}$ from mixed conformal-weight interactions. The scalar formula $\text{obs}_g = \kappa \cdot \lambda_g$ is unconditional at genus 1; at higher genera the diagonal term persists but $\delta F_g^{\text{cross}} \neq 0$ in the multi-weight channel.

10.8 The Bershadsky–Polyakov algebra

The Bershadsky–Polyakov algebra $\text{BP}_k = \mathcal{W}_3^{(2)}(k)$ is the subregular DS reduction of $\widehat{\mathfrak{sl}}_3$. It has four generators: J (weight 1), $G^\pm$ (weight $3/2$), and T (weight 2). The central charge:

$$c(\text{BP}_k) = 2 - \frac{24(k+1)^2}{k+3}.$$

The modular characteristic $\kappa(\text{BP}_k) = c/6$, and the Koszul complementarity conductor:

$$K(\text{BP}) = \kappa(\text{BP}_k) + \kappa(\text{BP}_{-k-6}) = \frac{98}{3}.$$

The self-dual level $k = -3$ coincides with the critical level $-h^\vee(\mathfrak{sl}_3)$, where $c(\text{BP}_k)$ has a simple pole; the value $\kappa(\text{BP}_{-3}) = 49/3$ quoted here is the *principal-value symmetric limit* $\lim_{\varepsilon \rightarrow 0} \frac{1}{2} [\kappa(\text{BP}_{-3+\varepsilon}) + \kappa(\text{BP}_{-3-\varepsilon})]$ across the pole, not a regular function value. The full Koszul conductor is a polynomial identity in $\mathbb{Q}(k)$: $K_{\text{BP}}(k) := c(k) + c(-k-6) \equiv 196$ as an element of $\mathbb{Q}(k)$; equivalently $c(k) - 98$ is an odd function of $(k+3)$. The full statement is the Bershadsky–Polyakov conductor theorem.

10.9 The Hilbert polynomial census

Proposition 10.4 (Hilbert polynomials across the landscape). *The chiral Hochschild Hilbert polynomial for the standard families:*

<i>Family</i>	$P(t)$	$\dim \text{ChirHoch}^1$
$\mathcal{H}_k$ (Heisenberg)	$1 + t + t^2$	1
Lattice V_Λ (rank = r)	$1 + r t + t^2$	r
$V_k(\mathfrak{g})$ (affine Kac–Moody)	$1 + t^2$ after zero-mode inner quotient	0; adjoint prequotient $\dim(\mathfrak{g})$
Vir_c (generic)	$1 + t^2$	0
$\mathcal{W}_N$ (generic)	$1 + t^2$	0
BP_k (generic)	$1 + t + t^2$	1
$\beta\gamma$	$1 + t + t^2$	1
bc	$1 + t^2$	0

The degree-1 component $\text{ChirHoch}^1(\mathcal{A})$ is the space of chiral derivations modulo inner derivations. Its dimension distinguishes families; the computation rests on the following Garland–Lepowsky-type identification.

Lemma 10.5 (Chiral H^1 for freely-generated VOAs at generic level). *Let $\mathcal{A}$ be a freely strongly-generated chiral algebra with generating space $V = \bigoplus_i V_{h_i}$ at conformal weights $h_i \geq 1$, at a generic (non-critical, non-resonant) level. Then*

$$\text{ChirHoch}^1(\mathcal{A}) \cong V_1$$

is the space of weight-1 strong generators. Equivalently, $\dim \text{ChirHoch}^1(\mathcal{A})$ equals the number of current-type (spin-1) strong generators.

Proof. Chiral derivations $D: \mathcal{A} \rightarrow \mathcal{A}$ of weight 0 are determined by their values on strong generators. At generic level, the resolution $\bar{B}(\mathcal{A})$ is concentrated in bar degrees $\{0, 1, 2\}$ (Theorem H), so $\text{ChirHoch}^1 = \text{Der}(\mathcal{A})/\text{Inn}(\mathcal{A})$ with no higher corrections. Inner derivations are modes of weight-1 fields acting as $\text{ad}(J_0^a)$; by the Garland–Lepowsky argument ([6], adapted to chiral algebras via [5]), every weight-0 derivation annihilating the vacuum and commuting with L_{-1} is inner on the subalgebra generated by weight ≥ 2 fields. Hence $\text{ChirHoch}^1(\mathcal{A}) = V_1$ modulo the span of J_0^a -actions that are already inner; for currents at generic level, the Kac–Moody relations make every J^a an independent outer derivation of the non-current subalgebra. $\square$

Specialising: $V_k(\mathfrak{g})$ has $V_1 = \mathfrak{g}$ (currents J^a), giving $\dim \text{ChirHoch}^1 = \dim(\mathfrak{g})$; Vir_c has $V_1 = 0$ (only $L_{-2}\mathbf{1}$ at weight 2), giving $\dim \text{ChirHoch}^1 = 0$; the principal $\mathcal{W}_N$ has primary generators at weights $2, 3, \dots, N$ with $V_1 = 0$ (the Cartan is frozen by the Drinfeld–Sokolov reduction), giving $\dim \text{ChirHoch}^1(\mathcal{W}_N) = 0$. The degree-0 and degree-2 terms are always present (center and dual center); the degree-1 term is the spin-1 generator count.

10.10 Three features of the census

Three structural features of the census table (Table 1) are visible only across the full landscape.

Shadow depth is not monotone in OPE pole order. The $\beta\gamma$ system has OPE pole order $p_{\max} = 1$ (a single simple pole) yet shadow depth $r_{\max} = 4$, while affine Kac–Moody has $p_{\max} = 2$ yet $r_{\max} = 3$. Shadow depth is an invariant of the obstruction tower, not of the OPE; the $\beta\gamma$ system separates the two invariants.

The complementarity conductor K is family-specific, not universal. It is 0 for Kac–Moody and free fields, 13 for Virasoro, 250/3 for $\mathcal{W}_3$, 98/3 for Bershadsky–Polyakov. The relation is a family-specific identity with universal form $\kappa + \kappa^! = \rho_{\mathcal{A}} \cdot K$, where $\rho_{\mathcal{A}}$ is the per-family leading κ -coefficient ($\rho_{\text{KM}} = \rho_{\text{free}} = 0$,

$\varrho_{\text{Vir}} = 1/2$, $\varrho_{\mathcal{W}_N} = H_N - 1$, $\varrho_{\text{BP}} = 1/6$); the bare equation $\kappa + \kappa^! = K$ (without ϱ) holds for no standard family.

The R -matrix column compresses to the κ column under averaging. Every entry demonstrates the same phenomenon: $\text{av}(r_{\mathcal{A}}(z)) = \kappa(\mathcal{A})$ (for abelian algebras, directly; for non-abelian Kac–Moody, after accounting for the Sugawara shift). The matrix-valued, z -dependent classical R -matrix of the ordered bar collapses to a single scalar under the Σ_2 -coinvariant projection. Every family with $r_{\max} \geq 3$ has an ordered bar that carries strictly more information than the symmetric bar; the five theorems extract the Σ_n -invariant content of this richer structure.

References

- [1] A. Beilinson and V. Drinfeld, *Chiral Algebras*, AMS Colloquium Publications, vol. 51, 2004.
- [2] A. Beilinson, V. Ginzburg, and W. Soergel, Koszul duality patterns in representation theory, *J. Amer. Math. Soc.* **9** (1996), no. 2, 473–527.
- [3] K. Costello and O. Gwilliam, *Factorization Algebras in Quantum Field Theory*, vols. 1–2, Cambridge University Press, 2017–2021.
- [4] J. Francis and D. Gaitsgory, Chiral Koszul duality, *Selecta Math. (N.S.)* **18** (2012), no. 1, 27–87.
- [5] E. Frenkel and D. Ben-Zvi, *Vertex Algebras and Algebraic Curves*, second edition, Mathematical Surveys and Monographs, vol. 88, American Mathematical Society, 2004.
- [6] H. Garland and J. Lepowsky, Lie algebra homology and the Macdonald–Kac formulas, *Invent. Math.* **34** (1976), 37–76.
- [7] E. Getzler and M. M. Kapranov, Modular operads, *Compositio Math.* **110** (1998), 65–126.
- [8] J. Gui, H. Li, and J. Zeng, Quadratic duality for chiral algebras, *Adv. Math.* **451** (2024), Paper no. 109791.
- [9] J.-L. Loday and B. Vallette, *Algebraic Operads*, Grundlehren der mathematischen Wissenschaften, vol. 346, Springer, 2012.
- [10] S. C. Mok, Logarithmic Fulton–MacPherson configuration spaces, arXiv:2503.17563, 2025.
- [11] D. Mumford, Towards an enumerative geometry of the moduli space of curves, in *Arithmetic and Geometry*, vol. II, Birkhäuser, 1983, 271–328.
- [12] G. Faltings, Calculus on arithmetic surfaces, *Ann. of Math. (2)* **119** (1984), 387–424.
- [13] F. Malikov and V. Schechtman, Homotopy chiral algebras, arXiv:2408.16787, 2024.
- [14] S. Priddy, Koszul resolutions, *Trans. Amer. Math. Soc.* **152** (1970), 39–60.
- [15] L. Positselski, *Two kinds of derived categories, Koszul duality, and comodule-contramodule correspondence*, Mem. Amer. Math. Soc. **212** (2011), no. 996.

A Ambients for the five theorems

The following ambient conditions sharpen the five theorem statements: ordinary direct-sum claims are separated from weight-completed coderived claims, shifted-symplectic upgrades are kept under their perfectness hypotheses, and uniform-weight genus identities are distinguished from all-weight corrections.

(A) Theorem A strengthenings

(A.1) Theorem $A^{\infty,2}$ (properad equivalence). The bar-cobar adjunction of Theorem A lifts to an $(\infty, 2)$ -adjoint equivalence at the *properad* level in the Francis–Gaitsgory factorization ambient, inscribed as `thm:A-infinity-2` in Vol I Chapter chapters/theory/theorem_A_infinity_2.tex. The pole-free restriction recovers Loday–Vallette’s $(\text{Ass}, \text{Ass}^!)$ Koszul pair via $(D_X\text{-mod}, \otimes^!) \hookrightarrow \text{Fact}(X)$;

the R -twisted Σ_n -descent lemma `lem:R-twisted-descent` (scoped to $R(z)$ satisfying the classical Yang–Baxter equation and unitarity $R(z) R^{op}(-z) = \text{id}$; automatic for rational Yangians and generic- q trigonometric $U_q(\widehat{\mathfrak{g}})$, vacuous when $R(z) = \tau$) relates $B^{\text{ord}}(\mathcal{A})$ to $B^\Sigma(\mathcal{A})$ at properad level and upgrades Theorem A ^{E_i} of the E_i primacy standalone to a corollary of A ^{$\infty,2$} . Fourteen downstream corollaries (classical Theorem A, bar-cobar adjunction, Vol II bridge I.3.2, Vol III $\Phi, \dots$) follow.

(A.2) Modular-family Theorem A on $\overline{\mathcal{M}}_{g,n}$. The modular-family extension of Theorem A to the full Deligne–Mumford moduli $\overline{\mathcal{M}}_{g,n}$ (including boundary) is now unconditional on the standard landscape. Three independent ingredients converge: (i) hypothesis (c) base-change proved via the Beilinson–Drinfeld holonomic formalism combined with the Vol II six-functor formalism on the relative Ran prestack; (ii) a nodal sewing theorem via Mok’s logarithmic Fulton–MacPherson compactification and Francis–Gaitsgory factorization-gluing, with Huang–Segal sewing convergence carrying chain-level matching at the nodes; (iii) globalization to the full $\overline{\mathcal{M}}_{g,n}$ via the K -theoretic filtration by collision depth, with associated graded equal to Arnold–Orlik–Solomon and the Euler characteristic computed via $\lambda_{-1}(\text{bundle})$ (same template as Theorem D).

(A.3) PTVV/factorization-homology alternative (Hoi). The factorization-homology alternative proof path for Theorem A, using Costello–Gwilliam factorization homology combined with Pantev–Toën–Vaquié–Vezzosi 1-shifted symplectic Lagrangian correspondence, upgrades from a sketch (Hoi) to a proposition in the rectified manuscript. The alternative is *genuinely independent*: it invokes neither Francis–Gaitsgory nor the Verdier-pair hypothesis.

(A.4) Six-fold TFAE on the Koszul locus, with a class-G seventh equivalence, at genus 0. The genus-0 clause of Theorem A is sharpened through the hub-and-spoke theorem `thm:ftm-seven-fold-tfae-via-hub-spoke` (Vol I chapters/theory/ftm_seven_fold_tfae_platonic.tex): five spokes (Koszul morphism, counit quasi-isomorphism, unit weak equivalence, twisted tensor acyclic, bar concentrated in weight 1) are each bidirected to the PBW E_2 -collapse hub on the full Koszul locus, and a sixth spoke (Swiss-cheese formality) is bidirected to the hub on the class-G stratum. The ten transitive bidirections of the six-fold TFAE thus follow from five proved spokes, with a sixth class-G-scoped spoke adjoining the seventh equivalence. Spoke 4 (twisted tensor $\Leftrightarrow$ PBW) is the load-bearing non-tautology, witnessed by $V_k(\mathfrak{sl}_2)$ at generic level. The SC-formality spoke is genuinely class-G scoped: class L/M members are Koszul but not SC-formal. Spoke 5 (bar weight-1 concentration) extends to $g \geq 1$ only for class G; outside class G the fiberwise obstruction $d_{\text{fib}}^2 = \kappa \cdot \omega_g$ is nonzero.

(B) Theorem B scope refinements

(B.1) Weight-completed chiral Positselski inscribed. The coderived bar-cobar adjunction is inscribed as `thm:chiral-positselski-weight-completed` (Vol I chapters/theory/theorem_B_scope_platonic.tex), which carries the legitimate phantomsection aliases `thm:chiral-positselski-7-2` and `thm:chiral-positselski-5-3` at the theorem’s home. The statement is unconditional on the *weight-completed* bar coalgebra for every chiral algebra with positive-energy conformal-weight grading and finite-dimensional conformal weight spaces. The five-step proof uses the chiral Φ/Ψ adjoints on the weight truncations $C/F^{\leq w}$, the chiral co-contra correspondence `thm:chiral-co-contra-correspondence` at each finite stage, and a Mittag–Leffler Milnor passage to the inverse limit. The chiral adaptations of Positselski [15, Theorem 4.6, Theorem 5.3] are invoked at citation level for the inverse-limit compatibility of the coderived and contraderived constructions; the chiral tensor $\otimes_{\text{ch}}$ preserves filtered colimits in each variable, making the adaptations immediate.

(B.2) Direct-sum chain-level class M and the weight-completed theorem. The chain-level class-M statement does not hold at the direct-sum level: at generic central charge, $S_4(\text{Vir}_c) = 10/[c(5c + 22)] \neq 0$ forces bar cohomology at weight 4, so strict chain-level quasi-isomorphism cannot hold. This is the correct scope, not a gap. The correct chain-level closure is in the *weight-completed* category via the strong filtration $F^n = \bigoplus_{k \leq n} B_k$, established as `thm:completed-bar-cobar-strong` in Vol I chapters/theory/theorem_B_scope_platonic.tex. The harmonic discrepancy at each weight stage N is a finite sum absorbed by an explicit chain homotopy $h \cdot m_o^{j-1}$; the Milnor/Mittag-Leffler passage to inverse limit converges.

(B.3) Class G/class L chain-level explicit. Class G Heisenberg chain-level inversion is unconditional via explicit MacLane splitting σ_{Heis} on $\text{Sym}(V_{[1]})$. Class L affine Kac–Moody at non-critical level chain-level inversion is unconditional via the PBW filtration and E_2 -collapse lifting.

(C) Theorem C: ambients

Nine refinements to Theorem C are recorded in the Theorem C refinement chapter. The list is as follows.

(C.1) The brace-dg-algebra-level identification $Z_{\text{ch}}^{\text{der}}(\mathcal{A}) \cong Z_{\text{ch}}^{\text{der}}(\mathcal{A}^\dagger)$ as E_2 -algebras after Deligne–Tamarkin is recorded as the derived-centre Koszul-equivalence conjecture. The H^0 -level naive centre recovery with the Koszul pairing is proved unconditionally by the naive-centre Koszul-identification lemma.

(C.2) Perfectness on the standard landscape is a conditional proposition in the fiber-finiteness hypothesis. It is unconditional for the Heisenberg family and for affine Kac–Moody at positive integer or boundary admissible level, via Tsuchiya–Ueno–Yamada propagation of finite conformal blocks in the first case and Arakawa C_2 -cofiniteness in the second. For affine Kac–Moody at generic non-critical k outside these loci, the boundary-stratum finiteness is recorded as the generic Kac–Moody boundary-perfectness conjecture. For class M the boundary-stratum fiber-finiteness is recorded as the class M boundary-perfectness conjecture.

(C.3) C_1 reflexivity on the Koszul locus is unconditional for the class G and class L loci, and is conditional on the class M or generic Kac–Moody boundary-perfectness conjectures for the rest of the standard landscape.

(C.4) There is no $+3$ shift contradiction at $g = 0$: the Verdier pairing of degree $-(3g-3)$ belongs to the $g \geq 1$ clause of quantum complementarity; at $g = 0$ the involution is trivial, $Q_o(\mathcal{A}^\dagger) = 0$, and no Verdier pairing is invoked.

(C.5) The C_2 hypotheses are pinned: BV package + Verdier + bar-chart lift only. Uniform-weight is *not* a C_2 hypothesis but belongs to the scalar-complementarity corollary.

(C.6) C_o is unconditional for the class G and class L families; the extension to the class M standard landscape is conditional on class M boundary-perfectness.

(C.7) The MC5 strengthening closes class-M C_2 (iii) in the weight-completed category.

(C.8) The explicit cross-channel correction at $(W_3, g=2)$ is inscribed: $\partial F_2^{\text{cross}}(W_3) = (c + 204)/(16c)$ in the explicit W_3 , genus-2 correction proposition.

(C.9) The mapping-stack $R\text{Map}(\overline{M}_g, BG_{\mathcal{A}})$ Pantev–Toën–Vaquié–Vezzosi presentation has two separate clauses. The shifted-symplectic form on the mapping stack is constructed independently of the eigenspace decomposition; the Lagrangian complementarity clause forwards to the quantum-complementarity theorem and remains conjectural until an intrinsic derived-stack construction of the Verdier anti-symplectic involution is given.

(D) Theorem D: globalization mechanisms

Part (i), the genus-1 formula $\text{obs}_1 = \kappa \cdot \lambda_1$, is unconditional.

(D.1) Virtual-class globalization. Via K -theoretic filtration, $\lambda_{-1}(E)$, and the top-degree Chern-character identity $\text{ch}_g(\lambda_{-1}(E)) = c_g(E)$ (splitting principle).

(D.2) Faltings citation retargeted. The Arakelov–Faltings identification used in Proof A is retargeted from the unpublished Faltings source to Arakelov 1974 plus Faltings 1984 §2 plus Soulé 1992 Ch. III (fiberwise Chern-curvature formula for the Hodge bundle in the Arakelov metric).

(D.3) BGS $c_1(\det E) \rightarrow c_g(E)$ bridge. Via the splitting principle combined with scalar-channel linearity: Chern–Weil for a scalar-endomorphism-valued curvature reads off the top Chern class with a *linear* scalar coefficient, not a multiplicative one.

(D.4) Clutching compatibility. The clutching proposition uses the genus-1 base, separating-clutching additivity, nonseparating-clutching compatibility, and tautological-ring purity to identify the same scalar obstruction class in the tautological socle. No division by κ is used; when $\kappa = 0$ the scalar obstruction is the zero class and satisfies the same boundary restrictions.

(D.5) Derived-stack route. The factorization-homology plus PTVV shifted-symplectic route is a derived-categorical compatibility check. It does not remove the scalar-diagonal uniform-weight hypothesis from the proved statement of Theorem D(ii).

Consequently Theorem D(ii) is proved on the scalar-diagonal uniform-weight lane by the GRR and clutching arguments of §6.4.

(H) Theorem H: Step 3 circularity rerouted

(H.1) PBW Koszulity criterion. The circularity risk at Step 3 of the $\{0, 1, 2\}$ concentration proof is rerouted through the H_H finite-window lane: PBW chiral Koszulness, localized residue-twisted bar concentration, strict Mittag–Leffler passage, and ordered-to-symmetric averaging. Shelton–Yuzvinsky/Priddy supplies the Koszul-complex acyclicity of the braid-arrangement Orlik–Solomon fibre. It does not contract the Arnold cohomology algebra itself and does not by itself prove Hochschild concentration. Bar-concentration and FM-tower collapse are linked by the localized residue-twisted spectral sequence, not by a bare Orlik–Solomon formality slogan.

(H.2) Independent verification paths. The disjoint-source checks are family certificates for the H_H mechanism: Heisenberg through the completed curved-dual finite-window certificate, Virasoro through Wang/Feigin–Fuchs generic cohomology on the stated lane, and affine $\mathfrak{sl}_2$ through Whitehead plus the zero-mode inner quotient. They are evidence for the stated package, not replacements for the collision-depth proof.

(H.3) E_1 -variant provable. The E_1 variant `thm:hochschild-concentration-E1` (for Yangian input) is provable on the ordered $H_H^{E_1}$ analogue: FM^{ord} , pure-braid Orlik–Solomon Koszul-complex input, the ordered Koszul dual, and the residue-twisted acyclicity hypothesis; the symmetric-bar obstruction $R(z) \neq \tau$ is the quantum-group content. The alternative `Ho5` is retrievable as the sharp-Hilbert-series statement from the `fm-tower-collapse  $E_2$` page.

Structural adjacents

(Er) Iterated-Sugawara tower and topologization (Vol II). `thm:iterated-sugawara-construction` and `thm:e-infinity-topologization-ladder` (Vol II chapters/connections/`e_infinity_topologization.tex`) promote the Sugawara construction of a single stress tensor to an infinite tower of inner higher-spin

Casimir tensors, each of which raises the topologization level by one: k inner stress tensors yield E_{k+2}^{top} . Virasoro at non-critical level gives E_3^{top} ; finite $\mathcal{W}_N$ gives E_{N+1}^{top} . The $\mathcal{W}_\infty$ endpoint requires the separate inverse-limit and completion hypotheses stated in Vol II.

(E2) Chiral higher Deligne (Vol II). `thm:chiral-higher-deligne` (Vol II chapters/theory/chiral_higher_deligne.tex) identifies $Z_{\text{ch}}^{\text{der}}(\mathcal{A})$ as the universal E_3 -chiral algebra acted on by $\text{SC}^{\text{ch},\text{top}}$ via heptagon edges $(3) \leftrightarrow (4) \leftrightarrow (5)$. This supplies a conditional second route to Theorem H once the chiral E_3 -PBW theorem is established. The theorem proved in this volume remains the ordered-bar/Orlik–Solomon concentration statement.

(E3) Periodic CDG for admissible KL (Vol I). `thm:periodic-cdg-is-koszul-compatible` (Vol I chapters/theory/periodic_cdg_admissible.tex) closes the admissible-level Kazhdan–Lusztig case: at $k + h^\vee = p/q$, the periodic-CDG filtration $F^n = \ker(Q_{\text{adm}}^n)$ on KL_k^{adm} is compatible with chiral Koszul duality; the chiral adjunction $\Omega^{\text{ch}} \dashv B^{\text{ch}}$ descends unconditionally.

Residual open frontiers

The five theorems require the following scope conditions and open obligations.

(F.1) $\mathcal{W}(p)$ triplet tempering. The Vol II statement `thm:tempered-stratum-contains-wp` is conjectural. The Zhu-bounded-Massey proof chain fails: Gurarie 1993 and Flohr 1996 construct logarithmic-CFT amplitudes with unbounded Massey products despite finite-dimensional Zhu algebra. The corrected tempered scope is: principal, non-logarithmic, non-minimal standard landscape; $\mathcal{W}(p)$ is open pending an Adamović–Milas character-amplitude bound.

(F.2) Non-tempered stratum. Emptiness of the non-tempered stratum in the C_2 -cofinite standard landscape holds on the non-logarithmic subset (Virasoro, $\mathcal{W}_N$, Schellekens, Monster, irrational cosets); logarithmic $\mathcal{W}(p)$ remains open.

(F.3) Riccati-arithmetic primes. Primes 1423, 3067, 23, 43, 419 have no primary-source Bernoulli witness for Kummer irregularity. They still appear in S_p numerators as Riccati-arithmetic characteristic primes rather than Kummer-arithmetic primes. The Riccati emergence set is $\{37, 691, 811\}$.

(F.4) Super-Yangian complementarity. The super-Yangian complementarity sum splits by normalisation. The super-trace form $\kappa^{\text{str}} + \kappa^{\text{str},!} = 0$ is proved for $Y_h(\mathfrak{sl}(m|n))$ on the sub-Sugawara line $m \neq n$ via super-Sugawara rational-identity cancellation (the Nazarov centrality and super-trace complementarity proposition). The Berezinian form $\kappa^{\text{sBer}} + \kappa^{\text{sBer},!} = \max(m, n)$ at the Sugawara-shifted dual level is conjectural, conditional on a closed-form derivation of the $\frac{1}{2} \max(m, n)$ shift magnitude induced by the central automorphism σ_{sBer} . Nazarov 1991 and Gow 2006 supply centrality of $\text{sBer}(T(u))$ and non-degeneracy of the Berezinian pairing, not the shift magnitude in closed form; the identity is verified symbolically at small rank as evidence, not derivation.

(F.5) CY-C pentagon stratifies generator rank, not κ_{ch} (Vol III). The Vol III pentagon of five intertwiners stratifies the generator-lattice rank $\rho^{R_i} \in \{3, 12, 24\}$, not the chiral modular characteristic. κ_{ch} is route-independent and equal to zero for $K3 \times E$ by the Hodge supertrace. This is a CY-side correction with no effect on the five theorems of Vol I at their stated scopes.

Table 1: Invariants of the standard landscape. All R -matrices in the trace-form convention; Ω is the inverse Killing form Casimir; $d = \dim \mathfrak{g}$, $t = k + h^\vee$.

Family	c	κ	K	Δ	Class	$r_{\max}$	R -matrix
<i>Abelian (class G, Gaussian)</i>							
$\mathcal{H}_k$ (Heisenberg)	1	k	o	o	G	2	k/z
Free fermion ψ	1/2	1/4	o	o	G	2	$\frac{1}{2z}$
Lattice V_Λ	$\text{rank}(\Lambda)$	$\text{rank}(\Lambda)$	o	o	G	2	Heisenberg
<i>Affine Kac–Moody (class L, Lie)</i>							
$\widehat{\mathfrak{sl}}_2$ at level k	$\frac{3k}{k+2}$	$\frac{3(k+2)}{4}$	o	o	L	3	$k \Omega_{\text{tr}}/z$
$\widehat{\mathfrak{sl}}_3$ at level k	$\frac{8k}{k+3}$	$\frac{4(k+3)}{3}$	o	o	L	3	$k \Omega_{\text{tr}}/z$
$\widehat{\mathfrak{so}}_8(D_4)$	$\frac{28k}{k+6}$	$\frac{7(k+6)}{3}$	o	o	L	3	$k \Omega_{\text{tr}}/z$
$\widehat{G}_2$	$\frac{14k}{k+4}$	$\frac{7(k+4)}{4}$	o	o	L	3	$k \Omega_{\text{tr}}/z$
$\widehat{F}_4$	$\frac{52k}{k+9}$	$\frac{26(k+9)}{9}$	o	o	L	3	$k \Omega_{\text{tr}}/z$
$\widehat{E}_6$	$\frac{78k}{k+12}$	$\frac{13(k+12)}{4}$	o	o	L	3	$k \Omega_{\text{tr}}/z$
$\widehat{E}_7$	$\frac{133k}{k+18}$	$\frac{133(k+18)}{36}$	o	o	L	3	$k \Omega_{\text{tr}}/z$
$\widehat{E}_8$	$\frac{248k}{k+30}$	$\frac{62(k+30)}{15}$	o	o	L	3	$k \Omega_{\text{tr}}/z$
<i>Free fields (class C/G)</i>							
$\beta\gamma$ (weight λ)	$2(6\lambda^2 - 6\lambda + 1)$	$6\lambda^2 - 6\lambda + 1$	o	special	C	4	constant
bc ghosts (weight λ)	$1 - 3(2\lambda - 1)^2$	$c_{bc}/2$	o	special	C	4	Clifford
<i>$\mathcal{W}$-algebras and Virasoro (class M, Mixed)</i>							
Vir_c	c	$c/2$	13	$\neq 0$	M	∞	$\frac{c/2}{z^3} + \frac{2T}{z}$
$\mathcal{W}_3(\mathfrak{sl}_3 \text{ DS})$	c	$5c/6$	$250/3$	$\neq 0$	M	∞	matrix-valued
$\mathcal{W}_N(\mathfrak{sl}_N \text{ DS})$	c	$c(H_N - 1)$	N -dep.	$\neq 0$	M	∞	matrix-valued
<i>Non-principal $\mathcal{W}$-algebras</i>							
BP_k (Bersh.–Pol.)	$2 - \frac{24(k+1)^2}{k+3}$	$c/6$	$98/3$	$\neq 0$	M	∞	multi-channel