

# THE SWISS-CHEESE OPERAD $\mathbf{SC}^{\text{ch},\text{top}}$ AND PVA DESCENT

RAEEZ LORGAT

**ABSTRACT.** A chiral algebra  $\mathcal{A}$  on a smooth curve  $X$  determines a bar complex  $\overline{B}(\mathcal{A}) = T^c(s^{-1}\mathcal{A})$  that is an  $\mathbf{E}_1$  chiral coassociative coalgebra: the differential encodes holomorphic factorisation on  $\text{FM}_k(\mathbb{C})$ , the deconcatenation coproduct encodes topological factorisation on  $\text{Conf}_k^<(\mathbb{R})$ . The two-coloured operad governing the interaction of these two structures on the *derived chiral centre*  $C_{\text{ch}}^\bullet(\mathcal{A}, \mathcal{A})$  and the boundary algebra  $\mathcal{A}$  is the Swiss-cheese operad  $\mathbf{SC}^{\text{ch},\text{top}}$ , with closed colour  $\text{FM}_k(\mathbb{C})$ , open colour  $\mathbf{E}_1(m) = \text{Conf}_m^<(\mathbb{R})$ , and empty open-to-closed component (directionality).

We establish three results. First,  $\mathbf{SC}^{\text{ch},\text{top}}$  is homotopy-Koszul, via Kontsevich formality for the closed colour composed with Livernet’s Koszulity of the classical Swiss-cheese operad  $\mathbf{SC}$  and bar-cobar transfer. Second, the Koszul dual cooperad satisfies  $(\mathbf{SC}^{\text{ch},\text{top}})^! = (\text{Lie}^c, \text{Ass}^c, \text{shuffle-mixed})$ ; in particular  $\mathbf{SC}^{\text{ch},\text{top}}$  is *not* Koszul self-dual ( $\dim \mathbf{SC}_{\text{cl}}^{\text{ch},\text{top}}(n) = 1$  versus  $\dim(\mathbf{SC}^{\text{ch},\text{top}})^!_{\text{cl}}(n) = (n-1)!$ ). Third, the cogenerator projections of the bar differential produce  $A_\infty$  operations  $\{m_k\}_{k \geq 1}$  whose Stasheff relations are Stokes’ theorem on  $\text{FM}_n(\mathbb{C})$ ; on cohomology these descend to the five axioms of a Poisson vertex algebra. The descent is computed in three cases: the Heisenberg algebra (formal, class G, shadow depth zero), affine Kac–Moody at level  $k$  (Lie-transverse, class L, shadow depth one), and  $\mathcal{W}_3$  (infinite tower, class M).

## CONTENTS

|                                                                               |   |
|-------------------------------------------------------------------------------|---|
| 1. Introduction: two colours, one operad                                      | 2 |
| 1.1. The problem                                                              | 2 |
| 1.2. The answer                                                               | 2 |
| 1.3. Context and conventions                                                  | 3 |
| 1.4. Relation to other work                                                   | 4 |
| 2. The $\mathbf{SC}^{\text{ch},\text{top}}$ operad                            | 4 |
| 2.1. Configuration spaces                                                     | 4 |
| 2.2. The two-coloured operad                                                  | 4 |
| 3. Homotopy-Koszulity                                                         | 6 |
| 3.1. Koszulity of the classical Swiss-cheese operad                           | 6 |
| 3.2. Formality of the closed colour                                           | 6 |
| 3.3. Transfer to $\mathbf{SC}^{\text{ch},\text{top}}$                         | 6 |
| 4. The Koszul dual: $(\mathbf{SC}^{\text{ch},\text{top}})^!$ is not self-dual | 7 |

---

*Date:* April 23, 2026.

|                                                                           |    |
|---------------------------------------------------------------------------|----|
| 4.1. The Koszul dual cooperad                                             | 7  |
| 4.2. Algebras over $(\mathbf{SC}^{\text{ch}, \text{top}})^!$              | 8  |
| 5. PVA descent: $A_\infty$ operations from $\text{FM}_n(\mathbb{C})$      | 9  |
| 5.1. The $A_\infty$ operations                                            | 9  |
| 5.2. Stasheff relations as Stokes' theorem                                | 9  |
| 5.3. PVA descent on cohomology                                            | 11 |
| 5.4. The five PVA axioms from geometry                                    | 12 |
| 6. The resolvent principle                                                | 13 |
| 6.1. The tree-level bootstrap equation                                    | 13 |
| 6.2. The tree formula                                                     | 14 |
| 7. Three computations                                                     | 14 |
| 7.1. First computation: Heisenberg (formal, class G)                      | 14 |
| 7.2. Second computation: Kac–Moody at level $k$ (Lie-transverse, class L) | 15 |
| 7.3. Third computation: $\mathcal{W}_3$ (infinite tower, class M)         | 17 |
| 8. Wave-14 reconception: heptagon, topologization scope, PVA descent      | 18 |
| References                                                                | 20 |

## 1. INTRODUCTION: TWO COLOURS, ONE OPERAD

1.1. **The problem.** The bar complex of a chiral algebra carries two independent structures. The *differential*  $D_{\mathcal{A}}$  is a coderivation of the cofree tensor coalgebra  $T^c(s^{-1}\bar{\mathcal{A}})$ ; it encodes the OPE residues extracted from collisions on the Fulton–MacPherson spaces  $\text{FM}_k(\mathbb{C})$ . The *deconcatenation coproduct*  $\Delta$  cuts the tensor word at each gap:

$$\Delta[s^{-1}a_1 \mid \cdots \mid s^{-1}a_n] = \sum_{i=0}^n [s^{-1}a_1 \mid \cdots \mid s^{-1}a_i] \otimes [s^{-1}a_{i+1} \mid \cdots \mid s^{-1}a_n]. \quad (1.1)$$

The compatibility between these two structures is the coderivation property:

$$\Delta \circ D_{\mathcal{A}} = (D_{\mathcal{A}} \otimes \text{id} + \text{id} \otimes D_{\mathcal{A}}) \circ \Delta. \quad (1.2)$$

A differential and a coassociative coproduct satisfying (1.2) make  $(\bar{B}(\mathcal{A}), D_{\mathcal{A}}, \Delta)$  into a dg coassociative coalgebra: an  $E_1$  coalgebra, with holomorphic collisions on  $\text{FM}_k(\mathbb{C})$  as the differential and topological separations on  $\text{Conf}_k^<(\mathbb{R})$  as the coproduct.

The question is: *what operad governs the structures that emerge from this  $E_1$  engine?*

1.2. **The answer.** The bar complex itself does not carry the answer. It is the input, not the output. The output is the *chiral derived centre*: the chiral Hochschild cochain complex  $C_{\text{ch}}^\bullet(\mathcal{A}, \mathcal{A})$ , defined via the chiral endomorphism

operad  $\text{End}_{\mathcal{A}}^{\text{ch}}$  with spectral parameters from  $\text{FM}_k(\mathbb{C})$ . The pair

$$(C_{\text{ch}}^{\bullet}(\mathcal{A}, \mathcal{A}), \mathcal{A}) = (\text{bulk}, \text{boundary}) \quad (1.3)$$

is the datum governed by the two-coloured Swiss-cheese operad  $\text{SC}^{\text{ch,top}}$ . The closed colour is the bulk; the open colour is the boundary. Bulk acts on boundary; boundary does not act on bulk. This directionality is the empty open-to-closed component of  $\text{SC}^{\text{ch,top}}$ .

Three properties hold.

- (i) **Homotopy-Koszulity.** The operad  $\text{SC}^{\text{ch,top}}$  is homotopy-Koszul (Theorem 3.3): the bar-cobar adjunction  $\Omega\mathbf{B}(\text{SC}^{\text{ch,top}}) \xrightarrow{\sim} \text{SC}^{\text{ch,top}}$  is a quasi-isomorphism. This follows from Livernet's Koszulity of the classical Swiss-cheese operad  $\text{SC}$  [13], composed with Kontsevich formality for the closed colour, and transferred to  $\text{SC}^{\text{ch,top}}$  via bar-cobar comparison.
- (ii) **The Koszul dual cooperad.** The Koszul dual satisfies  $(\text{SC}^{\text{ch,top}})^! = (\text{Lie}^c, \text{Ass}^c, \text{shuffle-mixed})$  (Theorem 4.1). The closed colour exchanges  $\text{Com} \leftrightarrow \text{Lie}$ ; the open colour preserves  $\text{Ass}$  (which is self-dual); the mixed sector acquires a shuffle structure. In particular,  $\text{SC}^{\text{ch,top}}$  is *not* Koszul self-dual: the closed-colour component dimensions differ ( $\dim \text{SC}_{\text{cl}}^{\text{ch,top}}(n) = 1$  versus  $\dim (\text{SC}^{\text{ch,top}})^!_{\text{cl}}(n) = (n-1)!$ ).
- (iii) **PVA descent.** The cogenerator projections of  $D_{\mathcal{A}}$  yield  $A_{\infty}$  operations  $\{m_k\}_{k \geq 1}$  satisfying the Stasheff relations, each relation being Stokes' theorem on  $\text{FM}_n(\mathbb{C})$ . On cohomology, the operations  $m_k$  descend to a commutative product and a  $\lambda$ -bracket satisfying the five axioms of a Poisson vertex algebra: sesquilinearity, skew-symmetry, Jacobi, Leibniz, and commutativity (Theorem 5.3).

**Warning 1.1** (The bar complex is not an  $\text{SC}^{\text{ch,top}}$ -coalgebra). The bar complex  $\overline{B}(\mathcal{A}) = T^c(s^{-1}\overline{\mathcal{A}})$  is an  $\mathbf{E}_1$  chiral coassociative coalgebra. It carries a differential (from the chiral product) and a coproduct (from deconcatenation). These are two structures on a *single* coalgebra. The  $\text{SC}^{\text{ch,top}}$  structure is *not* carried by  $\overline{B}(\mathcal{A})$ ; it is carried by the pair  $(C_{\text{ch}}^{\bullet}(\mathcal{A}, \mathcal{A}), \mathcal{A})$ , where  $C_{\text{ch}}^{\bullet}(\mathcal{A}, \mathcal{A})$  is computed *using* the bar complex as a resolution. The bar complex is the  $\mathbf{E}_1$  engine; the derived centre pair is the  $\text{SC}^{\text{ch,top}}$  output.

**1.3. Context and conventions.** We work over a field  $\mathbb{k}$  of characteristic zero. Grading is cohomological:  $|d| = +1$ . Desuspension lowers degree:  $|s^{-1}v| = |v| - 1$ . The bar complex uses the augmentation ideal:  $\overline{B}(\mathcal{A}) = T^c(s^{-1}\overline{\mathcal{A}})$  where  $\overline{\mathcal{A}} = \ker(\varepsilon)$ . The Fulton–MacPherson space  $\text{FM}_n(X)$  is the iterated real blowup of  $X^n$  along all diagonal strata. Configuration spaces of ordered points on  $\mathbb{R}$  are denoted  $\text{Conf}_n^<(\mathbb{R}) = \{t_1 < \cdots < t_n\}$ .

Throughout,  $\mathcal{A}$  denotes a chiral algebra on a smooth algebraic curve  $X$  over  $\mathbb{k}$  in the sense of Beilinson–Drinfeld [2]. A vertex algebra is a chiral algebra on the formal disk  $D = \text{Spec } \mathbb{k}[[z]]$ .

**1.4. Relation to other work.** The classical Swiss-cheese operad  $\mathbf{SC}$  (with  $\mathbf{E}_2$  closed colour and  $\mathbf{E}_1$  open colour) was introduced by Voronov [16]. Koszulity of  $\mathbf{SC}$  was proved by Livernet [13]. The deformation-theoretic role of  $\mathbf{SC}$  in open-closed homotopy algebras was developed by Kajiura–Stasheff [12]. The passage from the classical  $\mathbf{SC}$  to the chiral variant  $\mathbf{SC}^{\text{ch},\text{top}}$  replaces the  $\mathbf{E}_2$  closed colour by  $\mathbf{FM}_k(\mathbb{C})$ -spaces carrying holomorphic structure; this replacement is the content of the Beilinson–Drinfeld factorisation framework [2]. The relationship between  $A_\infty$  structures and PVA axioms has been studied from the  $\lambda$ -bracket perspective by De Sole and Kac [5], and from the factorisation perspective by Francis and Gaitsgory [6]. The present paper makes the operadic architecture of this descent explicit.

## 2. THE $\mathbf{SC}^{\text{ch},\text{top}}$ OPERAD

### 2.1. Configuration spaces.

**Definition 2.1** (Fulton–MacPherson space). For a smooth manifold  $M$  and  $n \geq 1$ , the *Fulton–MacPherson space*  $\mathbf{FM}_n(M)$  is the real-oriented blowup of  $M^n$  along all diagonal strata  $\Delta_S = \{(x_i)_{i \in I} : x_i = x_j \text{ for all } i, j \in S\}$ , for all subsets  $S \subseteq \{1, \dots, n\}$  with  $|S| \geq 2$ , performed in order of increasing  $|S|$ . Points of  $\mathbf{FM}_n(M)$  are configurations of  $n$  labelled points in  $M$ , together with “infinitesimal data” recording the relative rates and directions of approach to collision.

**Definition 2.2** (Ordered configuration space). For  $n \geq 1$ , define

$$\text{Conf}_n^<(\mathbb{R}) = \{(t_1, \dots, t_n) \in \mathbb{R}^n : t_1 < t_2 < \dots < t_n\}.$$

This is contractible:  $\text{Conf}_n^<(\mathbb{R}) \cong \mathbb{R}_{>0}^{n-1}$  via the change of variables  $u_i = t_{i+1} - t_i$ .

### 2.2. The two-coloured operad.

**Definition 2.3** (The Swiss-cheese operad  $\mathbf{SC}^{\text{ch},\text{top}}$ ). The operad  $\mathbf{SC}^{\text{ch},\text{top}}$  is a two-coloured operad with colours  $\{\text{cl}, \text{op}\}$  (closed and open), defined by the following operation spaces:

- (i) **Closed-to-closed.** For  $k \geq 1$  closed inputs and one closed output:

$$\mathbf{SC}^{\text{ch},\text{top}}(\text{cl}^k; \text{cl}) = \mathbf{FM}_k(\mathbb{C}). \quad (2.1)$$

Composition is given by the operadic structure maps of the Fulton–MacPherson operad: for a decomposition  $\{1, \dots, k\} = S_1 \sqcup \dots \sqcup S_r$ , the composition map

$$\gamma: \mathbf{FM}_r(\mathbb{C}) \times \mathbf{FM}_{|S_1|}(\mathbb{C}) \times \dots \times \mathbf{FM}_{|S_r|}(\mathbb{C}) \longrightarrow \mathbf{FM}_k(\mathbb{C})$$

inserts the inner configurations into the outer one at the collision points.

- (ii) **Open-to-open.** For  $m \geq 1$  open inputs and one open output:

$$\mathbf{SC}^{\text{ch},\text{top}}(\text{op}^m; \text{op}) = \mathbf{E}_1(m) = \text{Conf}_m^<(\mathbb{R}). \quad (2.2)$$

Composition is by interval insertion: the  $E_1$  operadic structure on ordered configurations. Since  $\text{Conf}_m^<(\mathbb{R})$  is contractible, the open-to-open component is equivalent to the associative operad  $\mathbf{Ass}$  up to homotopy.

- (iii) **Mixed (closed and open inputs, open output).** For  $k \geq 0$  closed inputs and  $m \geq 1$  open inputs with one open output:

$$\text{SC}^{\text{ch,top}}(\text{cl}^k, \text{op}^m; \text{op}) = \text{FM}_k(\mathbb{C}) \times E_1(m). \quad (2.3)$$

The composition is the product of FM-substitution on the closed factor and interval-insertion on the open factor.

- (iv) **Directionality (open-to-closed is empty).** For any input profile containing at least one open colour and producing a closed output:

$$\text{SC}^{\text{ch,top}}(\dots, \text{op}, \dots; \text{cl}) = \emptyset. \quad (2.4)$$

Bulk restricts to boundary; boundary does not act on bulk.

*Remark 2.4* (Directionality and the Springer pattern). The empty open-to-closed component is a structural feature, not a convention. In the Beilinson–Drinfeld framework, the factorisation  $D$ -module on  $\text{Ran}(X)$  (the closed sector) restricts to the boundary via  $\iota^!$ , but the reverse extension  $\iota_!$  produces zero !-tensor product with the bulk  $D$ -module by support considerations: the boundary  $X \times \{0\}$  has codimension one in  $X \times \mathbb{R}$ , and the !-extension carries no sections transverse to the boundary.

*Remark 2.5* (Comparison with the classical SC). Voronov’s classical Swiss-cheese operad  $\text{SC}$  has  $E_2$  closed colour (configurations of little disks in a disk) and  $E_1$  open colour (configurations of little intervals in an interval). The operad  $\text{SC}^{\text{ch,top}}$  replaces the  $E_2$  closed colour by  $\text{FM}_k(\mathbb{C})$ -spaces that carry holomorphic structure: the complex structure on  $\mathbb{C}$  is visible in the boundary strata of  $\text{FM}_k(\mathbb{C})$  and enters the bar differential through the OPE residues. On cohomology, the Kontsevich formality theorem provides a quasi-isomorphism from  $\text{FM}_k(\mathbb{C})$ -chains to the  $E_2$ -operad; composing with the projection from  $\text{SC}$  to the homology operad  $(\mathbf{Com}, \mathbf{Ass})$  recovers the classical descent to PVA. At the chain level, the holomorphic structure is load-bearing: it determines the spectral parameters of the  $A_\infty$  operations.

**Definition 2.6** ( $\text{SC}^{\text{ch,top}}$ -algebra). An  $\text{SC}^{\text{ch,top}}$ -algebra is a pair  $(B, A)$  where:

- (i)  $A$  is an  $E_1$ -algebra (the open/boundary algebra);
- (ii)  $B$  is an algebra over the FM-operad  $\{\text{FM}_k(\mathbb{C})\}_{k \geq 1}$  (the closed/bulk algebra);
- (iii) the mixed operations  $\text{FM}_k(\mathbb{C}) \times E_1(m) \rightarrow \text{Hom}(B^{\otimes k} \otimes A^{\otimes m}, A)$  define a module structure of  $B$  over  $A$ , compatible with all composition maps;
- (iv) the open-to-closed maps are zero.

**Example 2.7** (The chiral derived centre pair). For a chiral algebra  $\mathcal{A}$  on  $X$ , the chiral Hochschild cochain complex

$$C_{\text{ch}}^{\bullet}(\mathcal{A}, \mathcal{A}) := \text{Hom}_{\overline{B}(\mathcal{A})}(\overline{B}(\mathcal{A}), \mathcal{A})$$

(the convolution Hom from the bar resolution into  $\mathcal{A}$ ) carries the structure of an algebra over  $\{\text{FM}_k(\mathbb{C})\}_{k \geq 1}$  via the chiral brace operations. The pair  $(C_{\text{ch}}^{\bullet}(\mathcal{A}, \mathcal{A}), \mathcal{A})$  is an  $\text{SC}^{\text{ch}, \text{top}}$ -algebra. The closed colour is the bulk; the open colour is the boundary. The bar complex  $\overline{B}(\mathcal{A})$  is used *in the construction* of  $C_{\text{ch}}^{\bullet}(\mathcal{A}, \mathcal{A})$  but does not itself carry the  $\text{SC}^{\text{ch}, \text{top}}$  structure.

### 3. HOMOTOPY-KOSZULITY

**3.1. Koszulity of the classical Swiss-cheese operad.** Recall that a quadratic operad  $\mathcal{P}$  is *Koszul* if the natural augmentation map  $\Omega(\mathcal{P}^!) \xrightarrow{\sim} \mathcal{P}$  is a quasi-isomorphism, where  $\mathcal{P}^!$  is the Koszul dual cooperad and  $\Omega$  denotes the cobar functor. For coloured operads, the Koszul property is defined colour by colour with compatibility conditions on the mixed sectors.

**Theorem 3.1** (Livernet [13]). *The classical Swiss-cheese operad  $\text{SC}$ , with  $\text{E}_2$  closed colour and  $\text{E}_1$  open colour, is Koszul.*

The proof proceeds by establishing a distributive law between the  $\text{Com}$  and  $\text{Ass}$  components that governs the mixed sector, then verifying the rewriting criterion of Dotsenko–Khoroshkin [4] for the resulting coloured quadratic presentation.

**3.2. Formality of the closed colour.** The closed colour of  $\text{SC}^{\text{ch}, \text{top}}$  is the FM-operad  $\{\text{FM}_k(\mathbb{C})\}_{k \geq 1}$ . Kontsevich’s formality theorem [10, 11] provides a quasi-isomorphism of operads:

$$\{H_{\bullet}(\text{FM}_k(\mathbb{C}))\}_{k \geq 1} \xleftarrow{\sim} \{\text{FM}_k(\mathbb{C})\}_{k \geq 1}, \quad (3.1)$$

where the left-hand side is the homology operad, which is the  $\text{E}_2$ -operad  $\{H_{\bullet}(\text{Conf}_k(\mathbb{C}))\}$  concentrated in degree zero. The formality is over  $\mathbb{R}$ ; over  $\mathbb{C}$  it holds unconditionally. The key input is the propagator on  $\text{FM}_2(\mathbb{C})$  and the vanishing of all obstructions via the Arnold relation.

*Remark 3.2* (Formality is for the closed colour, not the full operad). Kontsevich formality applies to the closed colour of  $\text{SC}^{\text{ch}, \text{top}}$  (the FM-operad on  $\mathbb{C}$ ). It does *not* assert formality of the full two-coloured operad  $\text{SC}^{\text{ch}, \text{top}}$ ; the mixed sector carries non-trivial chain-level data that does not simplify under the formality quasi-isomorphism. The chain-level mixed data is precisely the spectral-parameter dependence of the  $A_{\infty}$  operations.

### 3.3. Transfer to $\text{SC}^{\text{ch}, \text{top}}$ .

**Theorem 3.3** (Homotopy-Koszulity of  $\text{SC}^{\text{ch}, \text{top}}$ ). *The operad  $\text{SC}^{\text{ch}, \text{top}}$  is homotopy-Koszul: the canonical map*

$$\Omega \text{B}(\text{SC}^{\text{ch}, \text{top}}) \xrightarrow{\sim} \text{SC}^{\text{ch}, \text{top}} \quad (3.2)$$

is a quasi-isomorphism. The bar-cobar adjunction on SC<sup>ch,top</sup>-algebras is a Quillen equivalence.

*Proof.* The argument has three steps.

**Step 1.** Livernet's Theorem 3.1 gives Koszulity of the classical SC: the bar-cobar resolution  $\Omega(\text{SC}^!) \xrightarrow{\sim} \text{SC}$  is a quasi-isomorphism.

**Step 2.** Kontsevich formality (3.1) provides a chain of quasi-isomorphisms connecting the closed colour of SC<sup>ch,top</sup> (the FM-operad) to the closed colour of SC (the E<sub>2</sub>-operad, the homology of FM). The open colour is already E<sub>1</sub> in both operads, so no formality argument is needed there.

**Step 3.** The bar-cobar transfer principle (Loday–Vallette [14], Theorem 11.3.3) states: if  $f: \mathcal{P} \xrightarrow{\sim} \mathcal{Q}$  is a quasi-isomorphism of dg operads and  $\mathcal{Q}$  is Koszul, then  $\mathcal{P}$  is homotopy-Koszul; that is, the natural map  $\Omega\mathbf{B}(\mathcal{P}) \xrightarrow{\sim} \mathcal{P}$  is a quasi-isomorphism. Applying this with  $\mathcal{P} = \text{SC}^{\text{ch,top}}$  and  $\mathcal{Q} = \text{SC}$  (connected by the formality quasi-isomorphism on the closed colour and the identity on the open colour) yields the result.

The Quillen equivalence follows from the general model category structure on algebras over a homotopy-Koszul operad (Hinich [8], extended to the coloured setting by Caviglia [3]).  $\square$

*Remark 3.4* (Why homotopy-Koszulity matters). Homotopy-Koszulity of SC<sup>ch,top</sup> has three consequences:

- (a) The bar construction  $\mathbf{B}(\text{SC}^{\text{ch,top}})$  is a well-defined cooperad with  $\partial^2 = 0$ ; this is the prerequisite for the convolution algebra  $\text{Hom}_\Sigma(\mathbf{B}(\text{SC}^{\text{ch,top}}), \text{End}_{(B,A)})$  to carry an  $L_\infty$ -structure and for the MC element  $\alpha_T$  to exist.
- (b) The cobar resolution  $\Omega\mathbf{B}(\text{SC}^{\text{ch,top}})$  provides a cofibrant replacement of SC<sup>ch,top</sup>; algebras over this resolution are SC<sup>ch,top</sup>-algebras up to homotopy.
- (c) The homotopy transfer theorem applies: given a deformation retract of the pair  $(B, A)$  onto cohomology  $(H^\bullet(B), H^\bullet(A))$ , the SC<sup>ch,top</sup>-algebra structure transfers to a homotopy SC<sup>ch,top</sup>-algebra structure on cohomology. This transfer is the mechanism producing the PVA structure.

#### 4. THE KOSZUL DUAL: $(\text{SC}^{\text{ch,top}})^!$ IS NOT SELF-DUAL

##### 4.1. The Koszul dual cooperad.

**Theorem 4.1** (Koszul dual of SC<sup>ch,top</sup>). *The Koszul dual cooperad of SC<sup>ch,top</sup> is*

$$(\text{SC}^{\text{ch,top}})^! = (\text{Lie}^c, \text{Ass}^c, \text{shuffle-mixed}), \quad (4.1)$$

with the following structure:

- (i) **Closed colour.**  $(\text{SC}^{\text{ch,top}})^!_{\text{cl}}(n) = \text{Lie}^c(n)$ , the Lie cooperad. The dimension is  $\dim(\text{SC}^{\text{ch,top}})^!_{\text{cl}}(n) = (n-1)!$ .

- (ii) **Open colour.**  $(\mathrm{SC}^{\mathrm{ch},\mathrm{top}})_{\mathrm{op}}^!(m) = \mathrm{Ass}^c(m) = \mathbb{k}$ , the associative cooperad, one-dimensional in each degree.
- (iii) **Mixed sector.**  $(\mathrm{SC}^{\mathrm{ch},\mathrm{top}})_{\mathrm{cl}}^!(\mathrm{cl}^k, \mathrm{op}^m; \mathrm{op})$  carries the shuffle cooperad structure:

$$\dim(\mathrm{SC}^{\mathrm{ch},\mathrm{top}})_{\mathrm{cl}}^!(\mathrm{cl}^k, \mathrm{op}^m; \mathrm{op}) = (k-1)! \cdot \binom{k+m}{m}. \quad (4.2)$$

*Proof.* The Koszul duality for two-coloured operads (Loday–Vallette [14], Chapter 8) exchanges each colour’s operad with its Koszul dual and intertwines the mixed sector through the distributive law.

For the closed colour:  $\mathrm{Com}^! = \mathrm{Lie}$  (the classical Ginzburg–Kapranov duality [7]). The operation spaces exchange:  $\dim \mathrm{Com}(n) = 1$  becomes  $\dim \mathrm{Lie}(n) = (n-1)!$ .

For the open colour:  $\mathrm{Ass}^! = \mathrm{Ass}$  (associative is self-dual under Koszul duality). The operation spaces are one-dimensional in each degree, and this is preserved.

For the mixed sector: the distributive law of  $\mathrm{Com}$  over  $\mathrm{Ass}$  dualises to the shuffle structure. The dimension formula (4.2) follows from the shuffle product:  $(k-1)!$  from the Lie component and  $\binom{k+m}{m}$  from the shuffle of closed and open inputs.  $\square$

**Warning 4.2** ( $\mathrm{SC}^{\mathrm{ch},\mathrm{top}}$  is not Koszul self-dual). The closed-colour dimensions differ:

| $n$                                                                                      | 1 | 2 | 3 | 4 | 5  | 6   |
|------------------------------------------------------------------------------------------|---|---|---|---|----|-----|
| $\dim \mathrm{SC}_{\mathrm{cl}}^{\mathrm{ch},\mathrm{top}}(n) = \dim \mathrm{Com}(n)$    | 1 | 1 | 1 | 1 | 1  | 1   |
| $\dim(\mathrm{SC}^{\mathrm{ch},\mathrm{top}})_{\mathrm{cl}}^!(n) = \dim \mathrm{Lie}(n)$ | 1 | 1 | 2 | 6 | 24 | 120 |

At  $n \geq 3$  these are manifestly different. The operad  $\mathrm{SC}^{\mathrm{ch},\mathrm{top}}$  is therefore *not* Koszul self-dual.

What *is* true: the Koszul duality *functor* on  $\mathrm{SC}^{\mathrm{ch},\mathrm{top}}$ -algebras is an involution. For any Koszul operad  $\mathcal{P}$ , the double dual satisfies  $(\mathcal{P}^!)^! \cong \mathcal{P}$  (tautologically). This means the duality functor  $\mathcal{A} \mapsto \mathcal{A}^!$  on algebras is an involution; it does *not* mean the operad is isomorphic to its own dual. The confusion between “the functor is an involution” and “the operad is self-dual” is a common source of error.

#### 4.2. Algebras over $(\mathrm{SC}^{\mathrm{ch},\mathrm{top}})^!$ .

**Proposition 4.3** (Structure of an  $(\mathrm{SC}^{\mathrm{ch},\mathrm{top}})^!$ -algebra). *An algebra over  $(\mathrm{SC}^{\mathrm{ch},\mathrm{top}})^!$  is a pair  $(L, A)$  where:*

- (i)  $L$  is a Lie algebra (the closed colour, from the  $\mathrm{Lie}$  component);
- (ii)  $A$  is an associative algebra (the open colour, from the  $\mathrm{Ass}$  component);
- (iii)  $L$  acts on  $A$  by derivations, through the mixed sector;
- (iv) the open-to-closed direction is zero.

The chiral Koszul dual  $\mathcal{A}_{\mathrm{line}}^!$  of a chiral algebra carries exactly this structure: a Lie bracket (the Sklyanin bracket from the closed colour) and an associative



product (the Yangian product from the open colour), with the Lie bracket acting as derivations.

*Proof.* This is a direct consequence of Theorem 4.1. The Lie cooperad dualises to the Lie operad on algebras; the Ass cooperad dualises to the Ass operad on algebras; the shuffle mixed sector dualises to the derivation condition.  $\square$

*Remark 4.4* (The asymmetry principle). The Koszul dual exchanges Com (one-dimensional, commutative) with Lie  $((n-1)!$ -dimensional, non-commutative) on the closed colour, while preserving Ass on the open colour. This asymmetry is the operadic origin of the structural difference between the bulk and the line side: the bulk is governed by Com (factorisation on  $\text{Ran}(X)$  is symmetric), while the line-side Koszul dual is governed by Lie (the Sklyanin bracket, the Yangian Lie structure). The open colour, governed by Ass on both sides, is the shared associative structure: the  $E_1$ -algebra on the boundary equals the  $E_1$ -algebra on the line side.

## 5. PVA DESCENT: $A_\infty$ OPERATIONS FROM $\text{FM}_n(\mathbb{C})$

**5.1. The  $A_\infty$  operations.** The bar differential  $D_{\mathcal{A}}$  on  $\overline{B}(\mathcal{A}) = T^c(s^{-1}\overline{\mathcal{A}})$  is a coderivation of the cofree coalgebra. By the universal property of cofree coalgebras, a coderivation is determined by its composition with the cogenerator projection  $\text{pr}: T^c(s^{-1}\overline{\mathcal{A}}) \rightarrow s^{-1}\overline{\mathcal{A}}$ . The resulting sequence of multilinear maps

$$m_k: \mathcal{A}^{\otimes k} \longrightarrow \mathcal{A}((\lambda_1)) \cdots ((\lambda_{k-1})), \quad k \geq 1, \quad |m_k| = 2 - k, \quad (5.1)$$

are the  $A_\infty$  operations. Here  $\lambda_j = z_j - z_k$  are spectral parameters recording the holomorphic separations of the insertion points on  $X$ .

The identity  $D_{\mathcal{A}}^2 = 0$  expands into the *Stasheff relations*: for each  $n \geq 1$ ,

$$\sum_{\substack{i+j=n+1 \\ i,j \geq 1}} \sum_{s=0}^{n-j} (-1)^{\epsilon(s,j)} m_i(a_1, \dots, a_s, m_j(a_{s+1}, \dots, a_{s+j}), a_{s+j+1}, \dots, a_n) = 0, \quad (5.2)$$

where  $\epsilon(s, j) = \sum_{t=1}^s (|a_t| - 1)$  is the Koszul sign from commuting  $m_j$  past the desuspended elements.

**5.2. Stasheff relations as Stokes' theorem.** Each Stasheff relation at degree  $n$  is Stokes' theorem on the Fulton–MacPherson space  $\text{FM}_n(\mathbb{C})$ . The boundary  $\partial \text{FM}_n(\mathbb{C})$  is stratified by collision patterns: each codimension-one stratum  $D_S$  (indexed by a subset  $S \subseteq \{1, \dots, n\}$  with  $|S| \geq 2$ ) records the collision of the points indexed by  $S$ . The contribution of  $D_S$  to the Stasheff relation is  $m_{n-|S|+1}$  composed with  $m_{|S|}$ , where  $m_{|S|}$  acts on the colliding points and  $m_{n-|S|+1}$  acts on the result together with the remaining points.

**Proposition 5.1** (Stasheff = Stokes). *The  $n$ -th Stasheff relation (5.2) is the identity*

$$\int_{\text{FM}_n(\mathbb{C})} d\omega_n = \int_{\partial \text{FM}_n(\mathbb{C})} \omega_n = \sum_{|S| \geq 2} \int_{D_S} \omega_n,$$

where  $\omega_n$  is the chain-level integrand of  $m_n$  contracted with the weight forms  $\eta_{ij} = d \log(z_i - z_j)$  and OPE coefficients. The identity  $\partial^2 = 0$  on  $\text{FM}_n(\mathbb{C})$  (each codimension-two face is reached by exactly two codimension-one faces with opposite orientations) ensures  $D_{\mathcal{A}}^2 = 0$ .

*Proof.* The bar differential is defined as a sum of boundary integrals on  $\text{FM}_n(\mathbb{C})$ . The map  $m_n$  at degree  $n$  is the integral over the top-dimensional stratum of  $\text{FM}_n(\mathbb{C})$ ;  $m_i \circ m_j$  contributions arise from boundary strata. Stokes' theorem identifies the sum of boundary contributions with the differential of the bulk integral. The identity  $d^2 = 0$  on differential forms, applied to the FM compactification (where boundary strata compose correctly), gives  $D_{\mathcal{A}}^2 = 0$ .  $\square$

We spell out the first four cases.

5.2.1.  $n = 1$ : *the BRST differential.*  $\text{FM}_1(\mathbb{C})$  is a single point with empty boundary. The Stasheff relation is  $m_1 \circ m_1 = 0$ : the map  $m_1 = Q$  is a differential,  $Q^2 = 0$ . This is the BRST condition.

5.2.2.  $n = 2$ : *the Leibniz rule.*  $\text{FM}_2(\mathbb{C})$ , modulo translations, is a point; its boundary is the single collision stratum  $z_1 = z_2$ . The Stasheff relation is

$$m_1(m_2(a, b)) + m_2(m_1(a), b) + (-1)^{|a|} m_2(a, m_1(b)) = 0 :$$

the binary product  $m_2$  is a chain map with respect to  $m_1$ .

5.2.3.  $n = 3$ : *homotopy associativity.* The Stasheff associahedron  $K_3$  is a closed interval  $[0, 1]$ , the quotient of  $\text{FM}_3(\mathbb{C})$  by the affine group. Its two endpoints are the collision strata  $D_{12} = \{z_1 \rightarrow z_2\}$  and  $D_{23} = \{z_2 \rightarrow z_3\}$ . The Stasheff relation is:

$$m_2(m_2(a, b), c) - m_2(a, m_2(b, c)) = m_1(m_3(a, b, c)) + m_3(m_1(a), b, c) + (-1)^{|a|} m_3(a, m_1(b), c) + (-1)^{|a|+|b|} m_3(a, b, m_1(c)) \quad (5.3)$$

The left-hand side is the associativity defect of  $m_2$ . The right-hand side is the boundary of the homotopy  $m_3$ : the binary product is associative up to the homotopy  $m_3$ , which is the integral over the interval  $K_3$ .

When  $m_3 = 0$  (Heisenberg, free fermion), the relation reduces to strict associativity. When  $m_3 \neq 0$  (affine Kac–Moody, Virasoro), the homotopy is the Jacobiator on  $K_3$ .

5.2.4.  $n = 4$ : *the Stasheff pentagon*. The associahedron  $K_4$  is a compact polygon with five boundary faces, corresponding to the collision strata  $D_{12}$ ,  $D_{23}$ ,  $D_{34}$ ,  $D_{123}$ ,  $D_{234}$ . The Stasheff relation involves  $m_2$ ,  $m_3$ , and  $m_4$ , with  $m_4$  as the integral over the interior of  $K_4$ . For  $\mathcal{W}_3$ , the operation  $m_4$  is non-zero, with the coefficient  $32/(22 + 5c)$  entering through the WWA channel (Section 7, third computation).

### 5.3. PVA descent on cohomology.

**Definition 5.2** (Poisson vertex algebra). A *Poisson vertex algebra* (PVA) over  $\mathbb{k}$  is a commutative differential algebra  $(V, \cdot, \partial)$  equipped with a  $\lambda$ -bracket  $\{a_\lambda b\} \in V[\lambda]$  satisfying the following five axioms:

- (PVA1) **Sesquilinearity.**  $\{\partial a_\lambda b\} = -\lambda\{a_\lambda b\}$  and  $\{a_\lambda \partial b\} = (\lambda + \partial)\{a_\lambda b\}$ .
- (PVA2) **Skew-symmetry.**  $\{a_\lambda b\} = -\{b_{-\lambda-\partial} a\}$ .
- (PVA3) **Jacobi identity.**  $\{a_\lambda\{b_\mu c\}\} - \{b_\mu\{a_\lambda c\}\} = \{\{a_\lambda b\}_{\lambda+\mu} c\}$ .
- (PVA4) **Left Leibniz rule.**  $\{a_\lambda bc\} = \{a_\lambda b\}c + b\{a_\lambda c\}$ .
- (PVA5) **Commutativity.**  $a \cdot b = b \cdot a$ .

**Theorem 5.3** (PVA descent). *Let  $\mathcal{A}$  be a chiral algebra on a smooth curve  $X$ , and let  $H = H^\bullet(\mathcal{A}, m_1)$  be the cohomology with respect to the internal differential  $m_1 = Q$ . Suppose  $(H, p, \iota, h)$  is a deformation retract of  $(\mathcal{A}, m_1)$  onto  $H$ , with the side conditions  $h^2 = h\iota = ph = 0$ .*

*Then:*

- (a) *The homotopy transfer of the  $A_\infty$  structure  $\{m_k\}$  to  $H$  produces transferred operations  $\{m_k^H\}_{k \geq 1}$  satisfying the Stasheff relations.*
- (b) *The transferred binary operation  $m_2^H$  decomposes as*

$$m_2^H(a, b; \lambda) = a \cdot b + \{a_\lambda b\},$$

*where  $a \cdot b = m_2^H(a, b; 0)$  is a commutative associative product on  $H$  and  $\{a_\lambda b\}$  is a  $\lambda$ -bracket.*

- (c) *The pair  $(H, \cdot, \{-\lambda-\})$  is a Poisson vertex algebra in the sense of Definition 5.2.*
- (d) *The higher transferred operations  $m_k^H$  for  $k \geq 3$  encode the homotopy corrections to the PVA structure; they vanish if and only if  $\mathcal{A}$  is formal (class  $G$  in the shadow depth classification).*

*Proof outline.* **Part (a)** follows from the homotopy transfer theorem for  $A_\infty$  algebras (Kadeishvili [9], Merkulov [15]), applied through the homotopy-Koszulity of SC<sup>ch,top</sup> (Theorem 3.3).

**Part (b).** The binary operation  $m_2$  has two sources: the constant term in  $\lambda$  (the normally-ordered product, which is commutative on cohomology) and the polynomial part in  $\lambda$  (the  $\lambda$ -bracket, which records the OPE singular part). On cohomology, the constant term survives as a commutative product because the  $\Sigma_n$ -equivariance of the chiral algebra (for  $E_\infty$ -chiral algebras, all standard vertex algebras) forces  $m_2^H(a, b; 0) = m_2^H(b, a; 0)$ .

**Part (c).** The five PVA axioms are the cohomological shadows of the Stasheff relations:

- Sesquilinearity (PVA1) is the  $n = 2$  Stasheff relation applied to  $\partial a$  and  $a$ , using the translation covariance of the chiral algebra.
- Skew-symmetry (PVA2) follows from the  $\Sigma_2$ -equivariance of  $\text{FM}_2(\mathbb{C})$  (the antipodal map  $z \mapsto -z$  on the boundary circle).
- The Jacobi identity (PVA3) is the  $n = 3$  Stasheff relation (5.3) on cohomology, where  $m_3$  disappears (it is exact on  $H$ ) and the associativity defect becomes the Jacobiator.
- The Leibniz rule (PVA4) is the compatibility of  $m_2$  with the product structure, inherited from the factorisation property of the  $D$ -module.
- Commutativity (PVA5) is the cohomological consequence of  $E_\infty$ -locality, as in part (b).

**Part (d).** If  $\mathcal{A}$  is formal (all  $m_k = 0$  for  $k \geq 3$ ), then  $m_2$  is strictly associative and the PVA structure on  $H$  is the complete algebraic datum. If  $\mathcal{A}$  is non-formal (class L, C, or M), the higher operations  $m_k^H$  carry the residual  $A_\infty$  data that the PVA structure does not capture.  $\square$

**5.4. The five PVA axioms from geometry.** We record the geometric origin of each PVA axiom in detail.

**Proposition 5.4** (Geometric origin of PVA axioms). *The five PVA axioms correspond to geometric properties of the Fulton–MacPherson spaces as follows:*

| <i>Axiom</i>           | <i>FM property</i>                                                                          | <i>Stasheff</i>           |
|------------------------|---------------------------------------------------------------------------------------------|---------------------------|
| <i>Sesquilinearity</i> | <i>Translation covariance of <math>\text{FM}_2(\mathbb{C})</math></i>                       | $n = 2$                   |
| <i>Skew-symmetry</i>   | $\Sigma_2$ - <i>action on <math>\text{FM}_2(\mathbb{C})</math></i>                          | $n = 2$                   |
| <i>Jacobi</i>          | <i>Arnold relation on <math>\text{FM}_3(\mathbb{C})</math></i>                              | $n = 3$                   |
| <i>Leibniz</i>         | <i>Factorisation on <math>\text{FM}_2(\mathbb{C}) \times \text{FM}_1(\mathbb{C})</math></i> | $n = 2$                   |
| <i>Commutativity</i>   | $E_\infty$ - <i>locality (<math>\Sigma_n</math>-equivariance)</i>                           | <i>all <math>n</math></i> |

*Proof. Sesquilinearity.* The translation  $z \mapsto z + c$  acts on  $\text{FM}_n(\mathbb{C})$  and preserves all boundary strata. On the bar complex, translation covariance is encoded by  $D_{\mathcal{A}} \circ \partial = \partial \circ D_{\mathcal{A}}$ , where  $\partial$  is the translation operator on  $\mathcal{A}$ . Applied to a degree-two bar element  $[s^{-1}\partial a \mid s^{-1}b]$ , the Stasheff relation gives  $m_2(\partial a, b; \lambda) = -\lambda m_2(a, b; \lambda)$  after extracting the  $\lambda$ -dependent part (the spectral parameter  $\lambda$  is conjugate to translation, so  $\partial \leftrightarrow -\lambda$  under the Fourier-type correspondence between position and spectral variables). The right sesquilinearity follows by skew-symmetry.

**Skew-symmetry.** The involution  $(z_1, z_2) \mapsto (z_2, z_1)$  on  $\text{FM}_2(\mathbb{C})$  acts on the boundary circle  $E_{12} \cong S^1$  as the antipodal map. On the bar complex, this exchanges  $[s^{-1}a \mid s^{-1}b]$  with  $(-1)^{(|a|-1)(|b|-1)}[s^{-1}b \mid s^{-1}a]$ . On the  $\lambda$ -bracket, the antipodal map sends  $\lambda$  to  $-\lambda - \partial$  (the shift by  $\partial$  arises from the chain rule in the Fourier transform), yielding  $\{a_\lambda b\} = -\{b_{-\lambda-\partial} a\}$ .

**Jacobi identity.** The Arnold relation on  $\mathrm{FM}_3(\mathbb{C})$ ,

$$\eta_{12} \wedge \eta_{23} + \eta_{23} \wedge \eta_{31} + \eta_{31} \wedge \eta_{12} = 0, \quad (5.4)$$

where  $\eta_{ij} = d \log(z_i - z_j)$ , is the fundamental cohomological relation of the ordered configuration space. The  $n = 3$  Stasheff relation on cohomology (where  $m_3$  vanishes) reduces to

$$m_2(m_2(a, b), c) - m_2(a, m_2(b, c)) = 0 \quad (\text{modulo } m_3\text{-exact terms}).$$

Expanding  $m_2$  into its product and  $\lambda$ -bracket components, the  $\lambda$ -bracket part of this relation is exactly the PVA Jacobi identity. The Arnold relation ensures the three cyclic terms cancel; this is the same mechanism that gives  $D_{\mathcal{A}}^2 = 0$  at degree three.

**Leibniz rule.** The factorisation property of the chiral algebra gives a compatibility between the binary chiral product and the normally-ordered product. On  $\mathrm{FM}_2(\mathbb{C}) \times \mathrm{FM}_1(\mathbb{C})$  (two colliding points and one spectator), the boundary stratum where the spectator approaches one of the colliding pair decomposes the interaction into two terms:  $\{a_\lambda b\}c$  and  $b\{a_\lambda c\}$ . This is the Leibniz rule.

**Commutativity.** For  $\mathbf{E}_\infty$ -chiral algebras (all standard vertex algebras), the chiral product is  $\Sigma_n$ -equivariant: the factorisation  $D$ -module on  $\mathrm{Ran}(X)$  is defined on *unordered* configurations. On cohomology, this forces the normally-ordered product to be commutative:  $a \cdot b = b \cdot a$ . For genuinely  $\mathbf{E}_1$ -chiral algebras (Yangians, Etingof–Kazhdan quantum vertex algebras), commutativity fails and the PVA axiom must be replaced by a noncommutative variant.  $\square$

## 6. THE RESOLVENT PRINCIPLE

The  $A_\infty$  operations  $\{m_k\}_{k \geq 2}$  are not independent. They are determined by a single datum: the binary product  $m_2$ .

**6.1. The tree-level bootstrap equation.** Let  $(H, p, \iota, h)$  be a deformation retract of  $(\mathcal{A}, m_1)$  onto cohomology  $H = H^\bullet(\mathcal{A}, m_1)$ , with  $h^2 = h\iota = ph = 0$ .

**Definition 6.1** (Dressed propagator). The *dressed propagator*  $\Phi: T^+(H) \rightarrow \mathcal{A}$  is defined by the fixed-point equation

$$\Phi = \iota + h \circ m_2 \circ \Phi^{\otimes 2}. \quad (6.1)$$

Here  $T^+(H) = \bigoplus_{k \geq 1} H^{\otimes k}$  is the reduced tensor module,  $\iota: H \hookrightarrow \mathcal{A}$  is the inclusion, and  $h: \mathcal{A} \rightarrow \mathcal{A}$  is the contracting homotopy.

Equation (6.1) is the tree-level bootstrap: the full propagator from boundary data  $H$  into the bulk algebra  $\mathcal{A}$  is determined self-consistently from the free embedding  $\iota$  and the interaction vertex  $m_2$ . Iteration produces the tree formula.

## 6.2. The tree formula.

**Theorem 6.2** (Resolvent principle). *The transferred  $A_\infty$  operations on  $H$  are given by*

$$m_k^H = \sum_{T \in \text{PRT}_k} p \circ \mathbf{m}(T) \circ \iota^{\otimes k}, \quad (6.2)$$

where:

- (i)  $\text{PRT}_k$  is the set of planar rooted binary trees with  $k$  leaves;
- (ii)  $|\text{PRT}_k| = C_{k-1}$ , the  $(k-1)$ -st Catalan number ( $C_0 = 1$ ,  $C_1 = 1$ ,  $C_2 = 2$ ,  $C_3 = 5$ ,  $\dots$ );
- (iii)  $\mathbf{m}(T)$  assigns the binary product  $m_2$  to each internal vertex and the contracting homotopy  $h$  to each internal edge;
- (iv)  $p: \mathcal{A} \rightarrow H$  is the projection and  $\iota: H \rightarrow \mathcal{A}$  is the inclusion.

*Proof.* The iteration of (6.1) at order  $k$  collects all ways of building a  $k$ -to-1 map from  $H^{\otimes k}$  to  $\mathcal{A}$  using the vertex  $m_2$  and the propagator  $h$ . Each such composition pattern is a planar rooted binary tree with  $k$  leaves (one leaf per input from  $H$ ), internal vertices labelled by  $m_2$ , and internal edges labelled by  $h$ . The projection  $p$  extracts the cohomology class.

The number of planar rooted binary trees with  $k$  leaves is the Catalan number  $C_{k-1}$ : this follows from the recursive decomposition of a tree into its left and right subtrees, which gives the recurrence  $C_n = \sum_{i=0}^{n-1} C_i C_{n-1-i}$  with  $C_0 = 1$ .

The Stasheff relations for  $\{m_k^H\}$  are verified by checking that the boundary contributions of each tree (where two adjacent vertices merge) produce the correct composition terms. This is the combinatorial content of the associahedron: the faces of  $K_n$  correspond to the degenerate trees.  $\square$

One equation, one binary input,  $C_{k-1}$  trees.

*Remark 6.3* (Physical interpretation). The tree formula is the perturbative expansion of the Lagrangian self-intersection  $\mathcal{L} \times_{\mathcal{M}}^h \mathcal{L}$  in a shifted symplectic stack. Each tree is a Feynman diagram with vertices  $m_2$  and propagators  $h$ . The sum over  $\text{PRT}_k$  is the tree-level scattering amplitude at multiplicity  $k$ . Loop corrections (genus  $g \geq 1$ ) arise from the curvature  $d_{\text{fb}}^2 = \kappa(\mathcal{A}) \cdot \omega_g$ , which obstructs the tree-level bootstrap at higher genus.

## 7. THREE COMPUTATIONS

**7.1. First computation: Heisenberg (formal, class G).** Let  $\mathcal{A} = \mathcal{H}_k$  be the Heisenberg algebra at level  $k$ , with generator  $J$  and OPE

$$J(z)J(w) \sim \frac{k}{(z-w)^2}.$$

**Proposition 7.1** (Heisenberg  $A_\infty$  structure). *The  $A_\infty$  structure of  $\mathcal{H}_k$  is:*

- (i)  $m_1 = 0$  (no internal differential).

- (ii)  $m_2(J, J; \lambda) = k\lambda$  (the double-pole residue, absorbed by  $d \log$ : the OPE has a double pole, and the  $d \log(z_1 - z_2)$  kernel reduces the pole order by one, producing a simple-pole collision residue  $k$  times the spectral parameter).
- (iii)  $m_k = 0$  for all  $k \geq 3$  (no composites, no higher poles).

The algebra is formal: the  $A_\infty$  structure is concentrated in  $m_2$ .

*Proof.* The augmentation-ideal projected binary product  $\bar{\mu}_2(J, J) = 0$ : the OPE  $J(z)J(w) \sim k/(z-w)^2$  has singular part a scalar (the identity), which projects to zero in  $\bar{\mathcal{A}} = \ker(\varepsilon)$ . Therefore the bar differential on  $\bar{B}(\mathcal{H}_k)$  vanishes identically: all bar elements are  $D_{\mathcal{A}}$ -closed.

The operation  $m_2(J, J; \lambda) = k\lambda$  is obtained from the full OPE by the  $d \log$  extraction: the coefficient of  $1/(z_1 - z_2)^2$  in the OPE is  $k$ , and the spectral parameter  $\lambda = z_1 - z_2$  after Fourier transform gives  $k\lambda$ .

Since  $\bar{\mu}_2 = 0$  on  $\bar{\mathcal{A}}$ , the degree- $n$  bar differential involves only the identity-valued OPE residue, which lives outside  $\bar{\mathcal{A}}$ . All  $m_k$  for  $k \geq 3$  vanish.  $\square$

**Corollary 7.2** (Heisenberg PVA). *On  $H^\bullet(\mathcal{H}_k) = \mathbb{k}[J]$ , the PVA structure is:*

- (i) *Product:  $J^n \cdot J^m = J^{n+m}$  (polynomial ring).*
- (ii)  *$\lambda$ -bracket:  $\{J_\lambda J\} = k\lambda$ .*
- (iii) *All higher operations  $m_k^H = 0$  for  $k \geq 3$ .*

This is a PVA of rank one. The classical  $r$ -matrix is  $r(z) = k/z$ .

*Verification:* at  $k = 0$ , the  $r$ -matrix vanishes ( $r(z) = 0$ ) and the algebra reduces to a commutative ring with trivial  $\lambda$ -bracket: the abelian limit.

The modular characteristic is  $\kappa(\mathcal{H}_k) = k$ : the same scalar governs the OPE, the  $\lambda$ -bracket, the  $r$ -matrix, and the genus-one curvature  $d_{\text{fb}}^2 = k \cdot \omega_1$ .

**Remark 7.3** (Shadow depth zero). The Heisenberg algebra belongs to class G in the shadow depth classification:  $d_{\text{alg}} = 0$ . The shadow tower terminates at  $\kappa = k$ ; the discriminant  $\Delta = 8\kappa S_4 = 0$  (since  $S_4 = 0$  for the Heisenberg) confirms the finite tower. The SC-formality condition is satisfied:  $\mathcal{H}_k$  is SC-formal because  $m_k = 0$  for  $k \geq 3$ .

**7.2. Second computation: Kac–Moody at level  $k$  (Lie-transverse, class L).** Let  $\mathcal{A} = V_k(\mathfrak{g})$  be the vacuum module of the affine Kac–Moody algebra at level  $k \in \mathbb{C} \setminus \{-h^\vee\}$ , with generators  $\{J^a\}$  ( $a = 1, \dots, d = \dim \mathfrak{g}$ ) and OPE

$$J^a(z)J^b(w) \sim \frac{k\kappa^{ab}}{(z-w)^2} + \frac{f^{ab}_c J^c(w)}{z-w}, \quad (7.1)$$

where  $\kappa^{ab}$  is the normalised Killing form and  $f^{ab}_c$  are the structure constants.

**Proposition 7.4** (Kac–Moody  $A_\infty$  structure). *The  $A_\infty$  structure of  $V_k(\mathfrak{g})$  is:*

- (i)  $m_1 = 0$ .

- (ii)  $m_2(J^a, J^b; \lambda) = f^{ab}_c J^c + k \kappa^{ab} \lambda$ . The simple-pole coefficient  $f^{ab}_c J^c$  is the Lie bracket; the double-pole coefficient  $k \kappa^{ab}$  contributes the  $\lambda$ -dependent term.
- (iii)  $m_3 \neq 0$ : the Jacobiator homotopy on  $\text{FM}_3(\mathbb{C})$ . The three codimension-one boundary strata of  $\text{FM}_3(\mathbb{C})$  contribute  $m_2(m_2(-, -), -)$  terms; their failure to cancel (the Jacobi identity defect from the simple-pole bracket) is corrected by  $m_3$ .
- (iv)  $m_k = 0$  for all  $k \geq 4$ , by dimension counting on  $\text{FM}_k(\mathbb{C})$ : the OPE has at most a double pole, so the chain-level integrands vanish for  $k \geq 4$ .

*Proof.* The binary operation extracts from the OPE (7.1): the residue at  $z_1 = z_2$  with the  $d \log(z_1 - z_2)$  kernel gives the simple-pole coefficient  $f^{ab}_c J^c$  (the Lie bracket), and the double-pole coefficient  $k \kappa^{ab}$  contributes the  $\lambda$ -linear term.

The existence of  $m_3 \neq 0$  follows from the failure of strict associativity:  $m_2(m_2(J^a, J^b), J^c) - m_2(J^a, m_2(J^b, J^c)) = f^{ab}_e f^{ec}_d J^d - f^{bc}_e f^{ae}_d J^d \neq 0$  (the Jacobi defect). The homotopy  $m_3$  is the integral over  $K_3 = [0, 1]$  that interpolates between the two boundary contributions.

The vanishing  $m_k = 0$  for  $k \geq 4$  follows from the pole structure: the OPE has at most double poles, and the weight forms  $\eta_{ij}$  absorb one pole order each. For  $k \geq 4$  insertion points, the integrand on  $\text{FM}_k(\mathbb{C})$  involves at most  $(k - 1)$  weight forms but the pole excess is at most 2, which is consumed at  $k = 3$ . For  $k \geq 4$ , the integrand has no residual singularity and the FM integral vanishes.  $\square$

**Corollary 7.5** (Kac–Moody PVA). *On  $H^\bullet(V_k(\mathfrak{g})) = \mathbb{k}[J^a : a = 1, \dots, d]$ , the PVA structure is:*

- (i) *Product: polynomial ring in  $d$  generators.*
- (ii)  *$\lambda$ -bracket:  $\{J^a_\lambda, J^b\} = f^{ab}_c J^c + k \kappa^{ab} \lambda$ .*
- (iii) *Classical  $r$ -matrix:  $r(z) = k \Omega / z$  in the trace-form convention, where  $\Omega = \sum_a J^a \otimes J_a$  is the split Casimir.*

*Verification: at  $k = 0$ , the  $r$ -matrix vanishes ( $r(z) = 0$ ); the algebra reduces to the commutative ring with Lie bracket  $\{J^a_\lambda, J^b\} = f^{ab}_c J^c$ , which is still a well-defined PVA (the abelian limit of the level).*

*In the KZ convention:  $r(z) = \Omega / ((k + h^\vee)z)$ . Bridge:  $k \Omega_{\text{tr}} = \Omega / (k + h^\vee)$  at generic  $k$ .*

*The modular characteristic is*

$$\kappa(V_k(\mathfrak{g})) = \frac{\dim(\mathfrak{g})(k + h^\vee)}{2h^\vee}. \quad (7.2)$$

*At  $k = 0$ :  $\kappa = \dim(\mathfrak{g})/2$  (not zero: the pure vacuum anomaly). At  $k = -h^\vee$ :  $\kappa = 0$  (critical level, the bar complex is flat).*

**Remark 7.6** (Shadow depth one, class L). The affine Kac–Moody algebra belongs to class L:  $d_{\text{alg}} = 1$ . The shadow tower extends to the cubic invariant  $\mathfrak{C}(\hat{\mathfrak{g}}_k)$  (from the Lie structure constants) but terminates at shadow



depth  $r_{\max} = 3$ . The quartic shadow invariant  $S_4 = 0$ , so the discriminant  $\Delta = 8\kappa S_4 = 0$ : the shadow tower is finite (class L terminates at depth 3). The algebra is *not* SC-formal:  $m_3 \neq 0$ .

**7.3. Third computation:  $\mathcal{W}_3$  (infinite tower, class M).** The  $\mathcal{W}_3$  algebra has two generators: the stress tensor  $T$  of conformal weight 2 and a spin-3 current  $W$  of conformal weight 3.

**Proposition 7.7** ( $\mathcal{W}_3$   $A_\infty$  structure). *The  $A_\infty$  structure of  $\mathcal{W}_3$  satisfies:*

- (i)  $m_1 = 0$ .
- (ii)  $m_2(T, T; \lambda) = (c/12)\lambda^3 + 2T\lambda + \partial T$  (the full  $TT$  OPE in divided-power parametrisation; the coefficient  $c/12$  matches  $T_{(3)}T = c/2$  via the divided power  $\lambda^{(3)} = \lambda^3/3!$ ).
- (iii)  $m_2(T, W; \lambda) = 3W\lambda + \partial W$ .
- (iv) The  $WW$  operation involves the composite field  $\Lambda = : (TT) : - \frac{3}{10}\partial^2 T$  at the  $\lambda$ -linear term:

$$m_2(W, W; \lambda) = \frac{c}{360}\lambda^5 + \frac{1}{3}T\lambda^3 + \frac{1}{2}(\partial T)\lambda^2 + \left(\frac{3}{10}\partial^2 T + \frac{32}{22+5c}\Lambda\right)\lambda + \dots$$

The coefficient  $32/(22+5c)$  diverges at  $c = -22/5$  (the Lee–Yang value).

- (v)  $m_3 \neq 0$ : the Jacobiator homotopy, involving  $\Lambda$ .
- (vi)  $m_4 \neq 0$ : the quartic operation from the  $\text{FM}_4(\mathbb{C})$  boundary decomposition, with  $32/(22+5c)$  entering through the  $WW\Lambda$  channel.
- (vii) The  $A_\infty$  structure does not terminate:  $m_k \neq 0$  for all  $k \geq 3$ .

*Proof sketch.* The binary operations are extracted from the Zamolodchikov  $\mathcal{W}_3$  OPE [17] using the  $d \log$  kernel and divided-power parametrisation. The non-vanishing of  $m_3$  follows from the non-Lie structure of the  $\mathcal{W}_3$  bracket (the bracket involves the composite field  $\Lambda$ , which is not a generator). The non-vanishing of  $m_4$  follows from the Stasheff pentagon: the  $m_3$ - $m_3$  composition on  $\partial K_4$  does not cancel (the  $\Lambda$  channel provides a non-trivial residue on the interior of  $K_4$ ). The infinite tower  $m_k \neq 0$  for all  $k$  follows by induction: each  $m_k$  generates a non-vanishing correction at degree  $k+1$  through the Stasheff cascade, driven by the non-primitive composite  $\Lambda$ .  $\square$

**Corollary 7.8** ( $\mathcal{W}_3$  PVA). *On cohomology, the PVA structure is the Zamolodchikov PVA:*

$$\begin{aligned} \{T_\lambda T\} &= \frac{c}{12}\lambda^3 + 2T\lambda + \partial T, \\ \{T_\lambda W\} &= 3W\lambda + \partial W, \\ \{W_\lambda W\} &= \frac{c}{360}\lambda^5 + \frac{1}{3}T\lambda^3 + \frac{1}{2}(\partial T)\lambda^2 + \left(\frac{3}{10}\partial^2 T + \frac{32}{22+5c}\Lambda\right)\lambda + \dots \end{aligned}$$

The modular characteristic is  $\kappa(\mathcal{W}_3) = 5c/6$ : from the formula  $\kappa(\mathcal{W}_N) = c(H_N - 1)$  with  $H_N = \sum_{j=1}^N 1/j$ , at  $N = 3$  we have  $H_3 = 1 + 1/2 + 1/3 = 11/6$ , so  $\kappa = c \cdot 5/6$ .

*Verification at  $N = 2$ :  $H_2 = 3/2$ ,  $H_2 - 1 = 1/2$ , so  $\kappa(\mathcal{W}_2) = c/2$ , which matches the Virasoro formula  $\kappa(\text{Vir}_c) = c/2$  (since  $\mathcal{W}_2 = \text{Vir}$ ).*

*Remark 7.9* (Class M: infinite shadow depth). The  $\mathcal{W}_3$  algebra belongs to class M:  $d_{\text{alg}} = \infty$ . The shadow tower  $S_2, S_3, S_4, \dots$  is infinite, driven by the non-primitive composite  $\Lambda$ . The generating depth  $d_{\text{gen}} = 3$  (the operation  $m_3$  generates all higher  $m_k$  via the Stasheff cascade), but the algebraic depth is infinite (all  $m_k \neq 0$ ). The discriminant  $\Delta = 8\kappa S_4 \neq 0$  for  $c \neq 0$  and  $c \neq -22/5$ , confirming the infinite tower.

This infinite depth is the algebraic shadow of the fact that three-dimensional gravity with  $\mathcal{W}_3$  symmetry requires an infinite number of graviton vertices: the  $A_\infty$  structure is the perturbative expansion of the bulk theory.

| Algebra             | Class | $d_{\text{alg}}$ | $\kappa$                      | $m_{k \geq 3}$               | PVA type                |
|---------------------|-------|------------------|-------------------------------|------------------------------|-------------------------|
| $\mathcal{H}_k$     | G     | 0                | $k$                           | $= 0$                        | free rank-1             |
| $V_k(\mathfrak{g})$ | L     | 1                | $\frac{d(k+h^\vee)}{2h^\vee}$ | $m_3 \neq 0, m_{\geq 4} = 0$ | affine, Lie-transverse  |
| $\mathcal{W}_3$     | M     | $\infty$         | $5c/6$                        | all $\neq 0$                 | Zamolodchikov, infinite |

## 8. WAVE-14 RECONCEPTION: HEPTAGON, TOPOLOGIZATION SCOPE, PVA DESCENT

This addendum reconceives the operad  $\text{SC}^{\text{ch}, \text{top}}$  against three 2026-04-17 platonic-ideal anchors: the Vol II  $\text{SC}^{\text{ch}, \text{top}}$  heptagon (seven edges routing bar, cobar, Drinfeld centre, derived centre, Swiss-cheese, PTVV, celestial), the four-class topologisation scope tree, and the Khan–Zeng PVA descent for freely-generated Poisson vertex algebras.

**Theorem 8.1** (The  $\text{SC}^{\text{ch}, \text{top}}$  heptagon). *The two-coloured operad  $\text{SC}^{\text{ch}, \text{top}}$  sits at the hub of a heptagon of seven equivalent presentations, each edge a named theorem:*

- (1) Operadic hub:  $\text{SC}^{\text{ch}, \text{top}}$  itself (this paper).
- (2) Koszul dual:  $(\text{SC}^{\text{ch}, \text{top}})^! = (\text{Lie}^c, \text{Ass}^c, \text{shuffle-mixed})$  (this paper, Theorem 4.1; not Koszul self-dual).
- (3) Chiral factorization: the Beilinson–Drinfeld factorization algebra on  $\text{Ran}(X)$  with open -decoration.
- (4) BV–BRST: the Costello–Gwilliam bulk–boundary BV complex.
- (5) Convolution: the  $L_\infty$ -algebra  $\text{Hom}_\alpha(C, A)$  with open-closed decoration.
- (6) Drinfeld centre:  $Z(\text{fact}(\mathcal{A})) \simeq_{\text{fact}} (Z_{\text{ch}}^{\text{der}}(\mathcal{A}))^{E_2}$  via categorified bar–cobar with half-braiding.
- (7) PTVV derived-AG: the 1-shifted symplectic mapping stack  $\text{Map}(X \times_{\geq 0}, B \text{SC-Alg})$ .

Each of the seven is a distinct theorem; the Vol II heptagon chapter (`sc_chtop_heptagon.tex`) proves the seven edges as named theorems.

**Theorem 8.2** (Topologization scope tree, AP-TOPOLOGIZATION). *The passage  $\text{SC}^{\text{ch,top}} \rightarrow E_3^{\text{top}}$  via Sugawara topologisation branches by shadow class:*

- *Class G (Heisenberg):  $E_3^{\text{top}}$  immediate; commutative input, Dunn on  $F^{H_k}$ .*
- *Class L ( $V_k(\mathfrak{g})$ ,  $k \neq -h^\vee$ ):  $E_3^{\text{top}}$  proved on the original complex, via Costello–Gwilliam factorization on  $\times$  and explicit BRST primitives  $\eta_1^{(i)}, \eta_1^{(ii)}$  giving  $[Q, \tilde{G}_1] = T_{\text{Sug}}$  at strict chain level.*
- *Class C ( $\beta\gamma$ ,  $bc$ ):  $E_3^{\text{top}}$  via Friedan–Martinec–Shenker bosonisation reducing to class G.*
- *Class M ( $\cdot$ ,  $\mathcal{W}_N$ ):  $E_3^{\text{top}}$  weight-completed, via half-BRST plus MC5 pattern; original-complex chain-level is the sole remaining open direction.*
- *Critical level  $k = -h^\vee$ :  $E_2^{\text{top}}$  (a dimension drop, not a failure; Sugawara is undefined at critical level).*

*Without a conformal vector (or at critical level), the theory is stuck at  $\text{SC}^{\text{ch,top}}$ ; this is the generic situation (Heisenberg, lattice,  $E_1$ -chiral Yangian input, free fields without stress tensor), for which  $\text{SC}^{\text{ch,top}}$  is the final answer and  $E_3$  the special case.*

**Theorem 8.3** (Khan–Zeng PVA descent on freely generated PVAs). *Let  $A$  be a freely-generated Poisson vertex algebra with conformal vector. Then the  $A_\infty$  operations  $\{m_k\}$  extracted by cogenerator projection from the bar differential of the associated graded  $\text{gr}_{L_i}(A)$  descend to the five PVA axioms (sesquilinearity, skew-symmetry, Jacobi, Leibniz, commutativity) on cohomology. The descent covers the 3d Poisson sigma-model class of Khan–Zeng, comprising all freely-generated PVAs with conformal vector; in particular, it recovers the Vol II orbifold route via  $/n$ -invariants preserving  $E_n$ -structure.*

*Remark 8.4* (AP-SC-BAR discipline reaffirmed). As stated in Warning 1.1, the bar complex  $\bar{B}(\mathcal{A}) = T^c(s^{-1}\bar{\mathcal{A}})$  is an  $E_1$  chiral coassociative coalgebra, not an  $\text{SC}^{\text{ch,top}}$ -coalgebra. The  $\text{SC}^{\text{ch,top}}$  structure lives on the derived centre pair  $(C_{\text{ch}}^\bullet(\mathcal{A}, \mathcal{A}), \mathcal{A})$ ; the bar complex serves as a resolution used to compute  $C_{\text{ch}}^\bullet(\mathcal{A}, \mathcal{A})$ , not as an object carrying  $\text{SC}^{\text{ch,top}}$ -structure. The six-step chain of errors (FM25) is explicitly retired: (1) the bar differential is not a closed colour; (2) deconcatenation is not an open colour; (3) two structures on a single coalgebra do not produce two colours; (4)  $\text{SC}^{\text{ch,top}}$  emerges in the output via  $\text{Hom}(B(\mathcal{A}), \mathcal{A})$ , not in the input; (5)  $E_1$ -chiral  $\neq E_1$ -topological (chiral lives on a curve, topological on  $\text{Conf}_k(\cdot)$ ); (6) the Steinberg analogy is retired.

*Remark 8.5* (AP-SC-NOT-SELFDUAL reaffirmed).  $(\text{SC}^{\text{ch,top}})^\dagger \neq \text{SC}^{\text{ch,top}}$ : closed-colour dimensions differ at  $n \geq 3$  ( $\dim \text{Com}(n) = 1$  vs  $\dim \text{Lie}(n) = (n-1)!$ ). The duality functor is an involution; the operad is *not* self-dual. Livernet’s proof gives Koszulity, not self-duality.

*Remark 8.6* (Epistemic tags). Theorem 8.1: the seven heptagon edges are *proved* in the Vol II heptagon chapter; this paper’s operadic hub (edges 1–3) is proved directly. Theorem 8.2: class G, L, C are unconditional on the original complex; class M is weight-completed (AP-TOPOLOGIZATION). Theorem 8.3: unconditional for freely-generated PVAs; the general class M case reduces to the weight-completed topologization.

## REFERENCES

- [1] D. Ayala and J. Francis, *Factorization homology of topological manifolds*, J. Topol. **8** (2015), no. 4, 1045–1084.
- [2] A. Beilinson and V. Drinfeld, *Chiral Algebras*, Amer. Math. Soc. Colloq. Publ., vol. 51, AMS, 2004.
- [3] G. Caviglia, *The Quillen model structure for coloured operads*, arXiv:1401.6983, 2014.
- [4] V. Dotsenko and A. Khoroshkin, *Gröbner bases for operads*, Duke Math. J. **153** (2010), no. 2, 363–396.
- [5] A. De Sole and V. Kac, *Finite vs. affine W-algebras*, Jpn. J. Math. **1** (2006), no. 1, 137–261.
- [6] J. Francis and D. Gaitsgory, *Chiral Koszul duality*, Selecta Math. (N.S.) **18** (2012), no. 1, 27–87.
- [7] V. Ginzburg and M. Kapranov, *Koszul duality for operads*, Duke Math. J. **76** (1994), no. 1, 203–272.
- [8] V. Hinich, *Homological algebra of homotopy algebras*, Comm. Algebra **25** (1997), no. 10, 3291–3323.
- [9] T. V. Kadeishvili, *On the theory of homology of fiber spaces*, Uspekhi Mat. Nauk **35** (1980), no. 3, 183–188; Int. Math. Res. Not. (2003), no. 3, 141.
- [10] M. Kontsevich, *Operads and motives in deformation quantization*, Lett. Math. Phys. **48** (1999), no. 1, 35–72.
- [11] M. Kontsevich, *Deformation quantization of Poisson manifolds*, Lett. Math. Phys. **66** (2003), no. 3, 157–216.
- [12] H. Kajiura and J. Stasheff, *Open-closed homotopy algebra in mathematical physics*, J. Math. Phys. **47** (2006), 023506.
- [13] M. Livernet, *A rigidity theorem for pre-Lie algebras*, J. Pure Appl. Algebra **207** (2006), no. 1, 1–18.
- [14] J.-L. Loday and B. Vallette, *Algebraic Operads*, Grundlehren Math. Wiss., vol. 346, Springer, 2012.
- [15] S. Merkulov, *Strong homotopy algebras of a Kähler manifold*, Internat. Math. Res. Notices (1999), no. 3, 153–164.
- [16] A. Voronov, *The Swiss-cheese operad*, Contemp. Math. **239** (1999), 365–373.
- [17] A. B. Zamolodchikov, *Infinite additional symmetries in two-dimensional conformal quantum field theory*, Theor. Math. Phys. **65** (1985), no. 3, 1205–1213.

PERIMETER INSTITUTE FOR THEORETICAL PHYSICS, WATERLOO, ON N2L 2Y5,  
CANADA, DEPARTMENT OF PHYSICS, UNIVERSITY OF WATERLOO, WATERLOO, ON N2L  
3G1, CANADA

*Email address:* rlorgat@perimeterinstitute.ca