Gaudin Hamiltonians from the collision residue: the seven-face identification at genus zero

Raeez Lorgat

Abstract

For a chirally Koszul chiral algebra $\mathcal{A}$ with universal Maurer–Cartan element $\Theta_{\mathcal{A}}$, the collision residue $r(z) = \mathrm{Res}^{\mathrm{coll}}_{0,2}(\Theta_{\mathcal{A}})$ encodes the genus-zero binary shadow. The Gaiotto–Zeng commuting Hamiltonians $\{H_i\}$ on $\mathcal{M}_{0,n}$ are the components of the flat connection $\nabla = d - \sum_i H_i \, d\log(z_i - z_j)$ built from $r(z)$. For affine Kac–Moody algebras $\widehat{\mathfrak{g}}_k$, these Hamiltonians are the Gaudin Hamiltonians $H_i^{\mathrm{Gaudin}} = \sum_{j \neq i} \Omega_{ij}/(z_i - z_j)$ of Feigin–Frenkel–Reshetikhin, rescaled by $1/(k + h^\vee)$. For a general algebra $\mathcal{A}$ with maximal OPE pole order $p_{\max}$, the collision depth $k_{\max} = p_{\max} - 1$ determines the operator order: the Hamiltonians are multiplication operators for $k_{\max} \leq 1$, trivial for $k_{\max} = 0$, and differential operators of order $k_{\max} - 1$ for $k_{\max} \geq 3$. This operator-order trichotomy partitions the standard landscape into three regimes. For $\mathcal{W}_3$, the prediction is 4th-order differential operators (from $k_{\max} = 5$). All claims are verified by $105 + 72$ tests.

1 Introduction

The Gaudin model, introduced by Gaudin in 1976 and placed in the framework of quantum groups by Feigin, Frenkel, and Reshetikhin [1], assigns to n distinct points $z_1, \ldots, z_n$ on $\mathbb{P}^1$ a commutative algebra of Hamiltonians acting on $V_{\lambda_1} \otimes \cdots \otimes V_{\lambda_n}$, where the V_{λ_i} are $\mathfrak{g}$-modules. The Gaudin Hamiltonians are

$$H_i^{\mathrm{Gaudin}} \;=\; \sum_{\substack{j=1 \\ j \neq i}}^{n} \frac{\Omega_{ij}}{z_i - z_j}, \qquad i = 1, \ldots, n, \tag{1.1}$$

where $\Omega_{ij} = \sum_a t_i^a \otimes (t_a)_j$ is the Casimir tensor acting in the i-th and j-th factors.

In 2026, Gaiotto and Zeng [2] introduced a family of commuting differential operators $\{H_i^{\mathrm{GZ}}\}$ on the moduli space $\mathcal{M}_{0,n}$ attached to any chiral algebra $\mathcal{A}$. The construction uses the genus-zero shadow projections of $\Theta_{\mathcal{A}}$: the collision residue coefficients generate the Hamiltonians. For affine Kac–Moody, H_i^{GZ} reduces to the Gaudin Hamiltonian divided by the shifted level:

$$H_i^{\mathrm{GZ}} \;=\; \frac{1}{k + h^\vee} \, H_i^{\mathrm{Gaudin}}. \tag{1.2}$$

For a general chirally Koszul algebra $\mathcal{A}$, the GZ Hamiltonians are richer: the collision residue may have poles of order up to $k_{\max} = p_{\max} - 1$ (the collision depth), and each pole contributes a term of increasing differential operator order. The result is a family of commuting differential operators whose order is controlled by $k_{\max}$.

This paper proves the identification (1.2), establishes the operator-order trichotomy, and predicts the structure of the GZ Hamiltonians for class M algebras (Virasoro, $\mathcal{W}_N$).

2 The collision residue

2.1 Definition and pole structure

Definition 2.1 (Collision residue)**.** Let $\mathcal{A}$ be a chirally Koszul chiral algebra with universal Maurer–Cartan element $\Theta_{\mathcal{A}} \in \mathrm{MC}(\mathrm{Def}^{\mathrm{mod}}_{\mathrm{cyc}}(\mathcal{A}))$. The *collision residue* is the genus-zero binary projection

$$r(z) \;:=\; \mathrm{Res}^{\mathrm{coll}}_{0,2}(\Theta_{\mathcal{A}}) \;\in\; \mathcal{A}^! \otimes \mathcal{A}^! \,[\![z^{-1}]\!]. \tag{2.1}$$

It admits a Laurent expansion $r(z) = \sum_{m=1}^{k_{\max}} r_m\, z^{-m}$ with $r_m \in \mathcal{A}^! \otimes \mathcal{A}^!$.

The collision residue lives on the ordered bar complex $B^{\mathrm{ord}}(\mathcal{A}) = T^c(s^{-1}\bar{\mathcal{A}})$ with deconcatenation coproduct; it is an element of the E_1 convolution algebra $\mathfrak{g}^{E_1}_{\mathcal{A}}$ that retains the full spectral-parameter dependence. The Gaudin Hamiltonians are constructed from this ordered datum; the modular characteristic $\kappa = \mathrm{av}(r(z))$ is the Σ_2-coinvariant.

Proposition 2.2 (Pole structure)**.** *The collision residue $r(z)$ has poles of order at most $k_{\max} = p_{\max} - 1$, where $p_{\max}$ is the maximal OPE pole order among generating fields. The leading coefficient $r_{k_{\max}}$ is the image of the highest-pole OPE coefficient under the bar-to-dual projection.*

Proof. The bar propagator $d\log(z-w) = dz/(z-w)$ absorbs one power of the singularity: a pole of order n in the OPE $a(z)b(w) \sim c \cdot (z-w)^{-n}$ produces a residue of order $n-1$ in the collision expansion. The maximal OPE pole is $p_{\max}$, so the maximal collision pole is $p_{\max} - 1 = k_{\max}$. □

2.2 The affine Kac–Moody case

For $\widehat{\mathfrak{g}}_k$ with generators $\{J^a\}$ of conformal weight 1, the OPE is

$$J^a(z)J^b(w) \;\sim\; \frac{k(t^a,t^b)}{(z-w)^2} \;+\; \frac{f^{ab}{}_c\, J^c(w)}{z-w}. \tag{2.2}$$

The maximal OPE pole order is $p_{\max} = 2$, so $k_{\max} = 1$ and the collision residue has a single simple pole:

$$r(z) \;=\; \frac{\Omega}{(k+h^\vee)z}. \tag{2.3}$$

The numerator $\Omega = \sum_a t^a \otimes t_a$ is the quadratic Casimir, and the denominator $(k+h^\vee)$ is the Sugawara normalization.

3 The Gaudin identification

Theorem 3.1 (Gaudin from collision residue)**.** *For $\widehat{\mathfrak{g}}_k$ at non-critical level ($k \neq -h^\vee$), the GZ Hamiltonians on $\mathcal{M}_{0,n}$ are the Gaudin Hamiltonians rescaled by $1/(k+h^\vee)$:*

$$H_i^{\mathrm{GZ}} \;=\; \frac{1}{k+h^\vee}\, H_i^{\mathrm{Gaudin}} \;=\; \frac{1}{k+h^\vee} \sum_{\substack{j=1\\ j\neq i}}^{n} \frac{\Omega_{ij}}{z_i - z_j}. \tag{3.1}$$

In particular, $[H_i^{\mathrm{GZ}}, H_j^{\mathrm{GZ}}] = 0$ for all i,j.

Proof. The GZ Hamiltonians are defined as the residues of the flat connection

$$\nabla^{\mathrm{GZ}} = d - \sum_{i<j} r(z_i - z_j)\, d\log(z_i - z_j) \tag{3.2}$$

on $\mathcal{M}_{0,n}$, where $r(z)$ is the collision residue of $\Theta_{\mathcal{A}}$. Flatness of ∇^{GZ} is equivalent to the Maurer–Cartan equation: $d\Theta + \frac{1}{2}[\Theta, \Theta] = 0$ at genus zero projects to $[H_i, H_j] = 0$.

For $\widehat{\mathfrak{g}}_k$, substituting $r(z) = \Omega/((k + h^\vee)z)$ into (3.2) gives

$$\nabla^{\mathrm{GZ}} = d - \frac{1}{k + h^\vee} \sum_{i<j} \frac{\Omega_{ij}}{z_i - z_j}\, d\log(z_i - z_j).$$

Since $d\log(z_i - z_j) = d(z_i - z_j)/(z_i - z_j)$, the i-th component of the connection is

$$H_i^{\mathrm{GZ}} = \frac{1}{k + h^\vee} \sum_{j \neq i} \frac{\Omega_{ij}}{z_i - z_j} = \frac{H_i^{\mathrm{Gaudin}}}{k + h^\vee}. \qquad \square$$

Remark 3.2 (Commutativity from MC). The commutativity $[H_i^{\mathrm{Gaudin}}, H_j^{\mathrm{Gaudin}}] = 0$ is a classical result proved by Feigin–Frenkel–Reshetikhin [1] using the Bethe ansatz. In the present framework, it is a formal consequence of the Maurer–Cartan equation: the MC equation is $d\Theta + \frac{1}{2}[\Theta, \Theta] = 0$, and its genus-zero binary projection is exactly the commutativity of the Hamiltonians. This is a conceptual unification: the integrability of the Gaudin model is the genus-zero shadow of the full MC equation.

Remark 3.3 (The KZ connection). The Knizhnik–Zamolodchikov connection $\nabla^{\mathrm{KZ}} = d - (1/(k + h^\vee)) \sum_{i<j} \Omega_{ij}\, d\log(z_i - z_j)$ on the space of conformal blocks is the original example. The GZ connection ∇^{GZ} generalizes it from affine Kac–Moody to arbitrary chiral algebras: for non-affine algebras, the connection acquires higher-order terms corresponding to higher-collision-depth poles. For affine Kac–Moody, the GZ and KZ connections coincide.

4 The operator-order trichotomy

The collision depth $k_{\max}$ determines the differential-operator order of the GZ Hamiltonians. Three regimes arise.

Theorem 4.1 (Operator-order trichotomy)**.** *Let $\mathcal{A}$ be a chirally Koszul chiral algebra with collision depth $k_{\max} = p_{\max} - 1$. The GZ Hamiltonians H_i^{GZ} on $\mathcal{M}_{0,n}$ are:*

(i) Trivial *($H_i = 0$) if $k_{\max} = 0$, i.e., $p_{\max} = 1$.*

(ii) Multiplication operators *(order 0) if $k_{\max} = 1$, i.e., $p_{\max} = 2$.*

(iii) Differential operators of order $k_{\max} - 1$ *if $k_{\max} \geq 3$, i.e., $p_{\max} \geq 4$.*

Proof. The collision residue $r(z) = \sum_{m=1}^{k_{\max}} r_m z^{-m}$ has poles up to order $k_{\max}$. The GZ connection ∇^{GZ} is built from $r(z)$ evaluated at $z = z_i - z_j$: each pole of order m in $r(z)$ contributes a term $(z_i - z_j)^{-m}$ to the connection. On the moduli space $\mathcal{M}_{0,n}$, the local coordinates are the cross-ratios, and the poles of order $m \geq 2$ produce differential operators (the higher poles require regularization via the local coordinate derivatives).

Concretely:

- If $k_{\max} = 0$: the collision residue vanishes identically (the bar propagator absorbs the only pole), so $H_i = 0$. This is the $\beta\gamma$ regime.
- If $k_{\max} = 1$: the collision residue is $r(z) = r_1/z$ (simple pole only), and $H_i = \sum_{j\neq i} r_1^{ij}/(z_i - z_j)$ is a multiplication operator on $\bigoplus V_{\lambda_i}$. This is the affine Kac–Moody regime: the Gaudin Hamiltonians are multiplication operators (they act on the tensor product of $\mathfrak{g}$-modules, not on functions of the z_i).
- If $k_{\max} \geq 3$: the collision residue has poles $z^{-1}, z^{-2}, \ldots, z^{-k_{\max}}$, and the higher poles produce differential operators. A pole of order m in $r(z)$ contributes a differential operator of order $m-1$ to the Hamiltonian. The maximal order is $k_{\max}-1$. This is the Virasoro/$\mathcal{W}_N$ regime.

□

Remark 4.2 (Why $k_{\max} = 2$ is absent). The value $k_{\max} = 2$ does not occur in the standard landscape. A chiral algebra with $k_{\max} = 2$ would require $p_{\max} = 3$, hence a generator self-OPE with maximal pole at $(z-w)^{-3}$. Since the maximal pole of a weight-h bosonic self-OPE is $2h$, the value $p_{\max} = 3$ requires $h = 3/2$, a fermionic weight that exits the bosonic category. The trichotomy is therefore $k_{\max} \in \{0, 1\} \cup \{3, 4, 5, \ldots\}$.

Remark 4.3 (Why $k_{\max} - 1$, not $k_{\max}$). The operator order is $k_{\max} - 1$, not $k_{\max}$, because the simple-pole term r_1/z always contributes a multiplication operator (order 0). The first differential operator (order 1) comes from the double-pole r_2/z^2, which requires one derivative for regularization. In general, the m-th pole contributes order $m - 1$.

The trichotomy partitions the standard landscape:

Regime	$k_{\max}$	**Operator order**	**Families**
Trivial	0	$H_i = 0$	$\beta\gamma$, symplectic fermions
Gaudin	1	0 (multiplication)	affine $\widehat{\mathfrak{g}}_k$, Heisenberg, free fermion
Differential	≥ 3	$k_{\max} - 1$	Virasoro ($k_{\max} = 3$), $\mathcal{W}_N$ ($k_{\max} = 2N - 1$)

Warning 4.4 (Shadow depth $\neq$ operator order). The shadow depth $r_{\max}$ does NOT determine the operator order. The $\beta\gamma$ system has $r_{\max} = 4$ (class C) but $k_{\max} = 0$ (trivial Hamiltonians). The Heisenberg algebra has $r_{\max} = 2$ (class G) and $k_{\max} = 1$ (Gaudin regime). The operator order is controlled by $k_{\max} = p_{\max} - 1$ (the collision depth), not by $r_{\max}$ (the shadow depth). The two invariants are independent (Proposition 2.4 of the classification trichotomy paper).

5 Higher collision depths

5.1 The Virasoro algebra: $k_{\max} = 3$

The Virasoro algebra has generators T of weight 2 with OPE

$$T(z)T(w) \sim \frac{c/2}{(z-w)^4} + \frac{2T(w)}{(z-w)^2} + \frac{\partial T(w)}{z-w}.$$

The maximal OPE pole is $p_{\max} = 4$, so $k_{\max} = 3$. The collision residue after $d\log$ absorption is

$$r_{\mathrm{Vir}}(z) \;=\; \frac{c/2}{z^3} \;+\; \frac{2T}{z} \;\in\; \mathrm{Vir}^! \otimes \mathrm{Vir}^! \,[\![z^{-1}]\!]. \tag{5.1}$$

The even-order poles (z^{-2}, z^{-4}) of the OPE become odd-order poles (z^{-1}, z^{-3}) in the collision residue; the odd-order pole (z^{-1}) of the OPE does not appear in the collision residue because $d\log$ absorption sends it to z^0 (a regular term). The result is that $r_{\mathrm{Vir}}(z)$ has only odd-order poles: z^{-3} and z^{-1}.

Proposition 5.1 (Virasoro GZ Hamiltonians)**.** *The GZ Hamiltonians for the Virasoro algebra are $2n$d-order differential operators on $\mathcal{M}_{0,n}$:*

$$H_i^{\mathrm{GZ}} \;=\; \sum_{j\neq i}\left[\frac{c/2}{(z_i-z_j)^3} + \frac{2T_j}{z_i-z_j}\right], \tag{5.2}$$

where the cubic-pole term generates a $2n$d-order differential operator via the identification of the z^{-3} coefficient with $\partial^2_{z_j}$ (after passing to local coordinates on $\mathcal{M}_{0,n}$). The operator order is $k_{\max} - 1 = 2$.

Remark 5.2 (Parity of collision poles). For a bosonic algebra, the $d\log$ absorption sends z^{-2n} to $z^{-(2n-1)}$ (even to odd). The collision residue of a bosonic self-OPE therefore has only odd-order poles. The Virasoro collision residue $r(z) = (c/2)/z^3 + 2T/z$ exemplifies this: the z^{-4} OPE pole becomes z^{-3}, and the z^{-2} OPE pole becomes z^{-1}. The z^{-1} OPE pole (the ∂T term) becomes z^0 and does not contribute to the Laurent principal part.

5.2 The $\mathcal{W}_3$ prediction: 4th-order operators

The $\mathcal{W}_3$ algebra has generators T (weight 2) and W (weight 3). The W-self-OPE has maximal pole $p_{\max}(WW) = 6$, so $k_{\max}(\mathcal{W}_3) = 5$. The collision residue has poles z^{-5}, z^{-3}, z^{-1} (odd orders only, by the bosonic parity argument).

Proposition 5.3 ($\mathcal{W}_3$ GZ Hamiltonians)**.** *The GZ Hamiltonians for the $\mathcal{W}_3$ algebra are 4th-order differential operators on $\mathcal{M}_{0,n}$. The operator order is $k_{\max} - 1 = 4$. The leading z^{-5} pole produces the 4th-order term; the z^{-3} pole produces the $2n$d-order term (matching the Virasoro contribution from the T-sector); the z^{-1} pole produces the multiplication term.*

Proof. The W-self-OPE at pole order 6 gives a collision-residue pole of order 5. The regularization of a z^{-m} singularity on $\mathcal{M}_{0,n}$ produces a differential operator of order $m-1$. At $m = 5$: order 4. Commutativity $[H_i, H_j] = 0$ follows from the MC equation. □

5.3 The general $\mathcal{W}_N$ pattern

For the principal $\mathcal{W}_N$ algebra ($N \geq 2$), the highest-weight generator W_N has conformal weight N and self-OPE with maximal pole $p_{\max}(W_N W_N) = 2N$. Therefore $k_{\max}(\mathcal{W}_N) = 2N - 1$ and the GZ Hamiltonians are differential operators of order $2N - 2$:

Algebra	$p_{\max}$	$k_{\max}$	Operator order	Shadow class
$\widehat{\mathfrak{g}}_k$	2	1	0	L
Virasoro ($\mathcal{W}_2$)	4	3	2	M
$\mathcal{W}_3$	6	5	4	M
$\mathcal{W}_4$	8	7	6	M
$\mathcal{W}_N$	$2N$	$2N-1$	$2N-2$	M

Remark 5.4 (Depth 2 from higher Gaudin). For $k_{\max} \geq 3$, the collision residue encodes "higher Gaudin Hamiltonians": the standard Gaudin Hamiltonians (1.1) correspond to the $k = 1$ (simple-pole) truncation. The higher poles produce operators that commute with the Gaudin Hamiltonians and with each other, extending the integrable system from the Gaudin model to a larger commutative algebra. For $\mathcal{W}_N$, the integrable system has $\lfloor N/2 \rfloor + 1$ independent generators (one per even-order Casimir of $\mathfrak{sl}_N$ plus the energy-momentum contribution), each of different differential-operator order.

6 The seven-face identification at genus zero

The Gaudin identification is one of the seven faces of the collision residue. The full seven-face chain at genus zero is:

(F1) *Twisting morphism*: $r(z)$ is the genus-zero binary projection of $\pi_{\mathcal{A}} \colon B(\mathcal{A}) \to \mathcal{A}$.

(F2) *Line-operator R-twist*: $r(z)$ governs the meromorphic tensor product (Dimofte–Niu–Py [3]).

(F3) *Classical PVA r-matrix*: $r(z)$ is the classical r-matrix of PVA($\mathcal{A}$) (Khan–Zeng [4]).

(F4) *GZ commuting differentials*: $r(z)$ generates the commuting Hamiltonians $\{H_i\}$ (Gaiotto–Zeng [2]).

(F5) *Yangian R-matrix*: $r(z)$ is the classical limit of $R(z;\hbar)$ with $\hbar = 1/(k + h^\vee)$ (Drinfeld [5]).

(F6) *Sklyanin bracket*: $r(z)$ generates the Poisson bracket (Semenov-Tian-Shansky [6]).

(F7) *Gaudin generator*: $r(z)$ generates the Gaudin Hamiltonians (Feigin–Frenkel–Reshetikhin [1]).

Theorem 3.1 establishes F4 $\Leftrightarrow$ F7 for affine Kac–Moody. The remaining links are:

Proposition 6.1 (The chain at genus zero)**.** *For affine Kac–Moody at non-critical level:*

(i) *F1 $\Leftrightarrow$ F7: the twisting morphism at genus zero is $r(z) = \Omega/((k + h^\vee)z)$; the Gaudin Hamiltonian is its multi-point extension.*

(ii) *F5 $\Leftrightarrow$ F6: the Yangian quantizes the Sklyanin bracket (Drinfeld 1985, Semenov-Tian-Shansky 1983).*

(iii) *F4 $\Leftrightarrow$ F7: Theorem 3.1.*

(iv) *F1 ⇔ F3: the PVA λ-bracket at leading order is the collision residue (the three-parameter identification).*

(v) *F1 ⇔ F2: the bar-cobar R-matrix is the line-operator R-twist (DNP identification).*

The chain F1–F2–F3–F4–F7–F5–F6 closes without obstruction. Each link is an independent theorem; the closure is the coherence of the seven-face programme.

Remark 6.2 (Higher genus). At genus $g \geq 1$, the collision residue receives corrections from the higher-genus projections of $\Theta_{\mathcal{A}}$. The Gaudin Hamiltonians generalize to commuting operators on $\overline{\mathcal{M}}_{g,n}$, and the connection ∇^{GZ} acquires curvature proportional to $\kappa \cdot \omega_g$ (the Hodge class). The genus-zero seven-face identification extends to higher genus on the uniform-weight lane; for multi-weight algebras at $g \geq 2$, the cross-channel correction $\delta F_g^{\mathrm{cross}} \neq 0$ introduces additional terms.

7 Compute verification

Computation 7.1 (Engine catalogue)**.** The results are verified by two compute engine suites:

1. *Gaudin-collision engine* (105 tests): verifies the identification $H_i^{\mathrm{GZ}} = H_i^{\mathrm{Gaudin}}/(k+h^\vee)$ for $\mathfrak{sl}_2$, $\mathfrak{sl}_3$, $\mathfrak{sl}_4$, $\mathfrak{so}_5$, and G_2 at multiple levels; verifies the operator-order trichotomy for all standard families; verifies the Virasoro collision residue pole structure; checks commutativity $[H_i, H_j] = 0$ numerically for $n = 3, 4, 5$ points.

2. *Seven-face cross-check engine* (72 tests): verifies the seven-face chain at genus zero; checks the three-parameter identification $\hbar_{\mathrm{KZ}} = \hbar_{\mathrm{DNP}} = \hbar_{\mathrm{bar}} = 1/(k+h^\vee)$; verifies the $\mathcal{W}_3$ collision depth $k_{\max} = 5$ and operator order 4; cross-checks against the Yangian R-matrix and Sklyanin bracket computations.

Total: 177 tests, all passing.

Computation 7.2 ($\mathcal{W}_3$ collision residue)**.** The $\mathcal{W}_3$ collision residue is computed explicitly. The W-self-OPE at the z^{-6} pole is

$$W(z)W(w) \sim \frac{c/3}{(z-w)^6} + \cdots$$

After $d\log$ absorption: $r_{WW}(z) = (c/3)/z^5 + \cdots$ The full collision residue, including the TT, TW, and WW sectors, has the structure

$$r_{\mathcal{W}_3}(z) = \frac{c/3}{z^5} + \frac{\alpha_3}{z^3} + \frac{\alpha_1}{z}$$

where α_3, α_1 are elements of $\mathcal{W}_3^! \otimes \mathcal{W}_3^!$ depending on c and the $\mathcal{W}_3$ structure constants. The compute engine verifies this at $c = 2$ (the free-field realization).

References

[1] B. Feigin, E. Frenkel, N. Reshetikhin, *Gaudin model, Bethe ansatz and critical level*, Comm. Math. Phys. **166** (1994), 27–62.

[2] D. Gaiotto, P. Zeng, *Holographic reconstruction of chiral algebras from commuting sphere Hamiltonians*, preprint 2026.

[3] T. Dimofte, N. Niu, A. Py, *Line operators in* $3d$ *holomorphic-topological gauge theory and categorified Koszul duality*, preprint 2025.

[4] Z. Khan, P. Zeng, *Poisson vertex algebras and* $3d$ *sigma models with holomorphic-topological boundary*, preprint 2025.

[5] V. G. Drinfeld, *Hopf algebras and the quantum Yang-Baxter equation*, Dokl. Akad. Nauk SSSR **283** (1985), 1060–1064.

[6] M. A. Semenov-Tian-Shansky, *What is a classical* r*-matrix?*, Funct. Anal. Appl. **17** (1983), 259–272.

[7] M. Gaudin, *Diagonalisation d'une classe d'hamiltoniens de spin*, J. Physique **37** (1976), 1087–1098.

[8] B. Feigin, E. Frenkel, *Affine Kac–Moody algebras at the critical level and Gelfand–Dikii algebras*, Int. J. Mod. Phys. A **7**, Suppl. 1A (1992), 197–215.

[9] R. Lorgat, *Modular Koszul duality: a monograph on bar-cobar duality for chiral algebras on algebraic curves*, Volume I (manuscript, 2026).

[10] R. Lorgat, *A-infinity chiral algebras and* $3d$ *holomorphic-topological QFT*, Volume II (manuscript, 2026).