

Algebraic \mathcal{D} -modules

Pavel Etingof

E-mail address: etingof@math.mit.edu

ABSTRACT. Typeset notes from a course at MIT.

Contents

Part 1. Lecture 1: Algebraic \mathcal{D}-modules	7
1. Lecture 1	8
Part 2. Lecture 2: Hilbert Polynomials of filtered algebras	9
2. Lecture 2	10
Part 3. Lecture 3: Further properties of the algebra \mathcal{D}	11
3. section Plan	12
Part 4. Lecture 4: \mathcal{O}-coherent \mathcal{D}-modules, 1	13
4. section Plan	14
Part 5. Lecture 5: \mathcal{O}-coherent \mathcal{D}-modules, 2	15
5. section Plan	16
Part 6. Lecture 6: Functional dimension and homological algebra	17
6. section Plan	18
Part 7. Lecture 7: \mathcal{D}-modules on general affine varieties	19
7. section Plan	20
Part 8. Lecture 8: Functorial Yoga, 1	21
8. section Plan	22
Part 9. Lecture 9: Proof of Kashiwara's theorem	23
9. section Plan	24
Part 10. Lecture 10: Theorem on Preservation of Holonomicity	25
10. section Plan	26
Part 11. Lecture 11: \mathcal{D}-modules on general varieties	27
11. section Plan	28
Part 12. Lecture 12: Equivariant \mathcal{D}-modules	29

12. section Plan	30
Part 13. Lecture 13: Derived Categories, 1	31
13. section Plan	32
Part 14. Lecture 14: Derived Categories, 2	33
14. section Plan	34
Part 15. Lecture 15: Derived Functors	35
15. section Plan	36
Part 16. Lecture 16: Functorial Yoga, 2	37
16. section Plan	38
Part 17. Lecture 17: The derived category of holonomic \mathcal{D}-modules	39
17. section Plan	40
Part 18. Lecture 18: Functorial Yoga, 3	41
18. section Plan	42
Part 19. Lecture 19: \mathcal{D}-modules with regular singularities, 1	43
19. section Plan	44
Part 20. Lecture 20: \mathcal{D}-modules with regular singularities, 2	45
20. More on the Riemann-Hilbert Map	46
21. Example 1: case of a complete curve	46
22. Example 2: the projective line minus four points	47
23. Holonomic \mathcal{D} -modules with RS in higher dimensions	48

Part 1

Lecture 1: Algebraic \mathcal{D} -modules

lec1

1. Lecture 1

Part 2

Lecture 2: Hilbert Polynomials of filtered algebras

lec2

2. Lecture 2

Part 3

Lecture 3: Further properties of the algebra \mathcal{D}

lec3

3. section Plan

Part 4

Lecture 4: \mathcal{O} -coherent \mathcal{D} -modules, 1

lec4

4. section Plan

Part 5

Lecture 5: \mathcal{O} -coherent \mathcal{D} -modules, 2

lec5

5. section Plan

Part 6

Lecture 6: Functional dimension and homological algebra

lec6

6. section Plan

Part 7

Lecture 7: \mathcal{D} -modules on general affine varieties

lec7

7. section Plan

Part 8

Lecture 8: Functorial Yoga, 1

lec8

8. section Plan

Part 9

Lecture 9: Proof of Kashiwara's theorem

lec9

9. section Plan

Part 10

Lecture 10: Theorem on Preservation of Holonomicity

lec10

10. section Plan

Part 11

Lecture 11: \mathcal{D} -modules on general varieties

lec11

11. section Plan

Part 12

Lecture 12: Equivariant \mathcal{D} -modules

lec12

12. section Plan

Part 13

Lecture 13: Derived Categories, 1

lec13

13. section Plan

Part 14

Lecture 14: Derived Categories, 2

lec14

14. section Plan

Part 15

Lecture 15: Derived Functors

lec15

15. section Plan

Part 16

Lecture 16: Functorial Yoga, 2

lec16

16. section Plan

Part 17

Lecture 17: The derived category of holonomic \mathcal{D} -modules

lec17

17. section Plan

Part 18

Lecture 18: Functorial Yoga, 3

lec18

18. section Plan

Part 19

Lecture 19: \mathcal{D} -modules with regular singularities, 1

lec19

19. section Plan

Part 20

Lecture 20: \mathcal{D} -modules with regular singularities, 2

20. More on the Riemann-Hilbert Map

Last lecture we considered the Riemann Hilbert map

$$\mathbf{RH} : \begin{array}{l} \text{Algebraic vector bundles on } X \\ \text{with an RS flat connection} \end{array} \longrightarrow \mathbf{Rep}\pi_1(X)$$

which assigns to each bundle (E, ∇) the monodromy representation of ∇ . Note that both categories for fixed rank r have a moduli space of objects which generically is an algebraic variety, and so in particular, a complex manifold. However, the map **RH** is not algebraic; it is only holomorphic.

Let's consider two examples of what this map does.

21. Example 1: case of a complete curve

Let X be a projective curve of genus g , then the moduli space \mathcal{M}_{dR} of line bundles with connection looks as follows: we have a map $\mathcal{M}_{dR} \longrightarrow \mathbf{Pic}_0(X) = \mathbf{Jac}(X)$ whose fiber is \mathbf{A}^g , an affine space bundle. Here \mathbf{A}^g is a torsor over $H^0(X, \Omega)$, and in particular is an algebraic variety of dimension $2g$.

On the other hand, the betti moduli space \mathcal{M}_b is the moduli space of representations of $\pi_1(X)$. Once we fix generators for $\pi_1(X)$:

$$a_1, \dots, a_g, b_1, \dots, b_g, \prod [a_i, b_i] = 1,$$

then we can identify

$$\mathcal{M}_b \simeq (\mathbf{C}^*)^{2g}.$$

Here the **RH** map is a holomorphic isomorphism

$$\mathcal{M}_{dR} \simeq \mathcal{M}_b.$$

Clearly it cannot be algebraic, since we have

LEMMA 21.1. *Any regular map $\mathbf{C}^* \rightarrow \mathbf{Jac}(X)$ is constant.*

PROOF. The map extends to

$$\mathbf{CP}^1 \rightarrow \mathbf{Jac}(X),$$

passage to the universal cover then yields

$$\mathbf{CP}^1 \rightarrow \mathbf{C}^n.$$

Now Liouville's theorem shows this map is constant. □

In more detail, **RH** is inverse to a map

$$f : \mathcal{M}_b \rightarrow \mathcal{M}_{dR}.$$

To construct f , let's construct

$$\pi : (\mathbf{C}^*)^{2g} \rightarrow \mathbf{Jac}(X) = \mathbf{Pic}_0(X);$$

which is an affine space bundle. To define f , consider the $4g$ -gon:




fig-4g-gon.png

Given

$$(\alpha_1, \beta_1, \dots, \alpha_g, \beta_g) \in (\mathbf{C}^*)^{2g}$$

we consider the trivial line bundle on the polygon and glue together a line bundle on X by using α_1 along a_1 , β_1 along b_1 etc.

22. Example 2: the projective line minus four points

Let $X = \mathbf{P}^1 - \{0, 1, \lambda, \infty\}$ with $\lambda \neq 0, 1, \infty$ and let's restrict to connections with trivial determinant. Now \mathcal{M}_{dR} has an open set \mathcal{M}_{dR}° of connections which have first order poles on the trivial bundle. Let's also restrict further to fixed monodromy; i.e. \mathcal{M}_{dR}° is the set of connections

$$\nabla = \partial - \frac{a_0}{z} - \frac{a_1}{z-1} - \frac{a_\lambda}{z-\lambda}, \mathbf{Tr}(a_j) = 0$$

and let $a_\infty = -a_0 - a_1 - a_\lambda$. Denote by

$$\mathcal{M}_{dR}^\circ(\alpha_0, \alpha_1, \alpha_\lambda, \alpha_\infty) = \left\{ \nabla, a_j \sim \begin{pmatrix} \alpha_j & 0 \\ 0 & -\alpha_j \end{pmatrix} \right\} \quad \text{then letting}$$

$$A_j \sim \begin{pmatrix} e^{2\pi i \alpha_j} & 0 \\ 0 & e^{-2\pi i \alpha_j} \end{pmatrix} \quad \text{we see the map}$$

$$\mathbf{RH} : \mathcal{M}_b(\alpha_0, \alpha_1, \alpha_\lambda, \alpha_\infty) \longrightarrow \{A_0, A_1, A_\lambda, A_\infty, A_0 A_1, A_\lambda A_\infty = 1\}$$

This map is highly transcendental.

Namely, let $\mathcal{P} \in \mathcal{M}_b(\alpha_0, \alpha_1, \alpha_\lambda, \alpha_\infty)$ and consider the point $Q_\lambda = \mathbf{RH}_\lambda^{-1}(\mathcal{P}) \in \mathcal{M}_{dR}^\circ(\alpha_0, \alpha_1, \alpha_\lambda, \alpha_\infty)$. This defines a flow on $\mathcal{M}_{dR}^\circ(\alpha_0, \alpha_1, \alpha_\lambda, \alpha_\infty)$ known as the Painlevé-6 flow.

23. Holonomic \mathcal{D} -modules with RS in higher dimensions

23.1. Constructible sheaves and complexes. Let X be a \mathbf{C} -algebraic variety. Denote by X^{an} the corresponding analytic variety considered with the classical topology. Let \mathbf{C}_X be the constant sheaf on X^{an} and $\mathbf{Sh}(X^{an})$ the category of \mathbf{C}_X -modules i.e. sheaves of \mathbf{C} -vector spaces. The derived category of bounded \mathbf{C}_X -complexes will be denoted $\mathbf{D}(X^{an})$.

DEFINITION 23.1. A \mathbf{C}_X -module \mathcal{F} is constructible if there exists a stratification

$$X = \cup_i X_i$$

of X by locally closed algebraic subvarieties such that $\mathcal{F}|_{X_i^{an}}$ is a locally constant complex of finite dimensional vector spaces.

REMARK 23.2. Note that a \mathbf{C}_X -complex is constructible if all of its cohomology sheaves are constructible as \mathbf{C}_X -vector spaces.

The full subcategory of $\mathbf{D}(X^{an})$ consisting of constructible complexes will be denoted by $\mathbf{D}_{con}(X^{an})$.

Any morphism $\pi : X \rightarrow Y$ of algebraic varieties induces a continuous map $\pi^{an} : X^{an} \rightarrow Y^{an}$, and we can consider the functors

$$\pi_!, \pi_* : \mathbf{D}(X^{an}) \longrightarrow \mathbf{D}(Y^{an})$$

$$\pi^*, \pi^! : \mathbf{D}(Y^{an}) \longrightarrow \mathbf{D}(X^{an})$$

We also have

$$\mathbb{D} : \mathbf{D}(X^{an}) \longrightarrow \mathbf{D}(X^{an})$$

We have

THEOREM 23.3. *These functors preserve the subcategory of derived constructible sheaves \mathbf{D}_{con} , and on them we have*

$$\mathbb{D}^2 = \mathbf{Id}$$

$$\mathbb{D}\pi^*\mathbb{D} = \pi^!$$

$$\mathbb{D}\pi_*\mathbb{D} = \pi_!$$

and $\mathbb{D}M = \mathbf{RHom}_{an}(M, \mathbf{C}_X)$.

23.2. De Rham Functor. Let \mathcal{O}_X^{an} be the structure sheaf of X^{an} . We will assign to each \mathcal{O}_X -module M the corresponding *analytic sheaf* of \mathcal{O}_X^{an} -modules M^{an} , which is locally given by

$$M^{an} = \mathcal{O}_X^{an} \otimes_{\mathcal{O}_X} M$$

.

This defines an exact functor

$$an : M(\mathcal{O}_X) \rightarrow M(\mathcal{O}_{X^{an}})$$

and in particular an exact functor

$$an : M(\mathcal{D}_X) \rightarrow M(\mathbf{D}_X^{an})$$

, where \mathcal{D}_X^{an} is the sheaf of analytic differential operators.

DEFINITION 23.4. *The De Rham Functor*

$$\mathbf{DR} : \mathbf{D}^b(\mathcal{D}_X) \rightarrow \mathbf{D}^b(X^{an}) = \mathbf{D}^b(\mathbf{Sh}(X^{an}))$$

is

$$\mathbf{DR}(M^\circ) = \Omega_X^{an} \otimes_{\mathcal{D}_X^{an}} (M^\circ)^{an}$$

REMARK 23.5. Since $\mathbf{dR}(\mathcal{D}_X)$ is a locally projective resolution of Ω_X we have

$$\mathbf{DR}(M^\circ) = \mathbf{dR}(\mathcal{D}_X^{an}) \otimes_{\mathcal{D}_X^{an}} (M^\circ)^{an}[\dim X]$$

In particular, if M is an \mathcal{O} -coherent \mathcal{D}_X -module corresponding to a vector bundle with a flat connection and $\mathcal{L} = M^{flat}$ is the local system of flat sections, then

$$\mathbf{DR}(M) = \mathcal{L}[\dim X]$$

by Poincaré's lemma.

Here is the main theorem about the connection between \mathcal{D} -modules and constructible sheaves:

- THEOREM 23.6.
 - $\mathbf{DR}(\mathbf{D}_{hol}(\mathcal{D}_X)) \subset \mathbf{D}_{con}(X^{an})$, and on \mathbf{D}_{hol} \mathbf{DR} commutes with both tensor product and \mathbf{D} .
 - On the subcategory \mathbf{D}_{rs} the functor \mathbf{DR} commutes with all of the above functors
 - $\mathbf{DR} : \mathbf{D}_{rs}(\mathcal{D}_X) \rightarrow \mathbf{D}_{con}(X^{an})$ is an equivalence.

23.3. Definitions concerning vector bundles with flat connection. Let M be a vector bundle on X with a flat connection.

DEFINITION 23.7. M lies in \mathbf{D}_{rs} if the restriction of M to any curve has regular singularities.

DEFINITION 23.8. An irreducible M in \mathbf{D}_{hol} has *regular singularities* if $M = j_{!*}L$ for L a vector bundle with flat connection and regular singularities.

REMARK 23.9. An object $M \in \mathbf{D}_{hol}$ has regular singularities if and only if all composition factors have regular singularities.