

The Igusa Square Root

The Borcherds determinant of $\phi_{0,1}$, its denominator algebra, and the $K_3 \times E$ realization problem

Raez Lorgat

April 2026

FRONTISPIECE

The Igusa cusp form Δ_5 is one object. The genus-two automorphic section, the denominator of a generalized Borcherds–Kac–Moody superalgebra on $\Lambda_{II}^{2,1}$, the protected Pfaffian of a chiral E_3 -algebra of $K_3 \times E$, the discriminant of a one-parameter K_3 -mirror period family, and the singular theta lift of the weak Jacobi form $\phi_{0,1}$ under the Howe correspondence $\mathbf{Sp}_4(\mathbb{R}) \leftrightarrow \mathbf{O}(3, 2)$ are five readings of one chiral E_3 -algebra

$$\mathfrak{A}_{K_3 \times E} \in E_3\text{-HolFA}(K_3 \times E),$$

and on each of these five levels its value is Δ_5 .

o.o.o.I The five views.

- ① *Automorphic.* The Borcherds–Gritsenko–Nikulin section

$$D_X = \Delta_5 \in H^0(\mathbf{Sp}(2, \mathbb{Z}) \backslash \mathcal{H}_2, \mathcal{L}^{\otimes 5} \otimes \nu_{\Delta_5})$$

of the weight-5 automorphic line on Siegel upper-half space, with multiplier ν_{Δ_5} of [MAASS:1, ?Borcherds95].

- ② *Borcherds–Kac–Moody denominator.* The denominator of the generalized BKM Lie superalgebra $\mathfrak{g}_{\Delta_5}$ on the rank-three Lorentzian root lattice $\Lambda_{II}^{2,1} \subset \Lambda^{2,1} = \Lambda^{(1,1)} \oplus \langle 2 \rangle$ of signature $(2, 1)$, the index-four sublattice with even $f_{\pm 2}$ coordinates (abstractly $U(2) \oplus \langle 2 \rangle$) in the basis (f_2, f_3, f_{-2}) with bilinear form $(f_2, f_{-2}) = -1$, $(f_3, f_3) = 2$; the type-II Cartan matrix on simple roots $\delta_1 = 2f_2 - f_3$, $\delta_2 = 2f_{-2} - f_3$, $\delta_3 = f_3$ is $(\delta_i, \delta_j) = 4\mathbf{I}_3 - 2\mathbf{J}_3 = \begin{pmatrix} 2 & -2 & -2 \\ -2 & 2 & -2 \\ -2 & -2 & 2 \end{pmatrix}$,

$$\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1} \Delta_5(2Z),$$

with signed root supermultiplicities $\text{smult } \mathfrak{g}_{\Delta_5, \alpha(n, l, m)} = f(nm, l)$ on the active support $\Gamma_{\text{act}} = \{(n, l, m) \in \Gamma_{\text{eff}} \mid f(nm, l) \neq 0\}$, with $\alpha(n, l, m) = 2nf_2 - lf_3 + 2mf_{-2}$ [?Borcherds95, ?GN, ?GNPartI, ?GNII].

- ③ *Compact Pfaffian.* The protected Pfaffian of the chiral E_3 -algebra $\mathfrak{A}_{K_3 \times E}$ of the $K_3 \times E$ formal target, presented at finite stage R by the Dirac–Igusa pro-object

$$\mathfrak{D}_X^{DI} = \varprojlim_R (\mathcal{A}_{X,R}^{E_3}, F_{X,R}^{\text{hyb}}, \Gamma_{X,R}, \Pi_X, \Phi_R, o_R, H_R, P_R^\Pi, C_{X,R}, \Theta_{\text{Kos},R}, \mathcal{L}_{\text{Pf},R}, \text{pf}_{X,R}, \epsilon_{o,R}, \text{Rec}_R)$$

(the chiral E_3 -source $\mathcal{A}^{E_3}$, the hybrid factorisation carrier F^{hyb} of $\text{Ran}^{\text{hyb}}(E)$, the Mukai charge lattice Γ , the Gram projection Π_X , the specialisation functor Φ , the orientation o , the Hall datum H , the Gram-descended primitive P^Π , the source coalgebra C , the Koszul cobar Θ_{Kos} , the Pfaffian line $\mathcal{L}_{\text{Pf}}$, the Pfaffian section pf , the orientation character ϵ_o , and the recognition map Rec ; each component is constructed in Chapter 5). The four named obstructions (D_0) , (O_1) , $(O_1)^+$, (O_2) condition the Pfaffian–Dirac theorem

$$\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{DI}) = \Delta_5, \quad \epsilon_o = \nu_{\Delta_5}, \quad \mathcal{P}_X \cong \mathfrak{g}_{\Delta_5}$$

where $\mathcal{P}_X$ is the protected primitive Lie superalgebra $P_X^{\Pi, \text{red}}$ of Chapter 5.6 (the H° of the protected primitives of the assembled hybrid factorisation sheaf $\mathcal{F}_X$ after Gram-projection pushforward $\overline{\Pi}_{X,*}^\ominus$ and Hopf-radical quotient $\overline{\text{Rad}\langle \cdot, \cdot \rangle_X}$), ϵ_o is the orientation character of the Pfaffian line on $W^{(2)}(\Lambda_{II}^{2,1})$, and ν_{Δ_5} is the Maass multiplier of [?MAASS:1] on $\mathbf{Sp}_4(\mathbb{Z})$, pulled back along the exceptional isomorphism $\mathbf{Sp}_4 \simeq \text{Spin}(3, 2)$ and the inclusion $\Lambda_{II}^{2,1} \hookrightarrow \Lambda^{3,2}$ to a character on $W^{(2)}(\Lambda_{II}^{2,1})$ taking value -1 on each of the three simple type-II reflections s_{δ_i} ; see Theorem 4.1.

- ④ *Mirror discriminant. Conjectural.* On the one-parameter K_3 -mirror period family $\mathcal{P} \rightarrow \mathbb{D}$ the discriminant divisor equals the polar divisor of Δ_5^{-2} , itself twice the zero divisor of Δ_5 ; the Igusa cusp form then encodes the period-family discriminant of mirror symmetry on $K_3 \times E$ (Chapter 8).
- ⑤ *Singular theta lift.* The lift of the weight-zero index-one weak Jacobi form $\phi_{o,1}$ under the Howe correspondence $\text{Mp}(4, \mathbb{R}) \leftrightarrow \mathbf{O}(3, 2)$, equivalently under the exceptional isomorphism $\mathbf{Sp}_4 \simeq \text{Spin}(3, 2)$:

$$\Theta_{\text{sing}}^{\text{Howe}}(\phi_{o,1}) = \Delta_5$$

[?Borcherds95, ?B].

0.0.0.2 The cross-level identities. The four edges ① $\leftrightarrow$ ②, ② $\leftrightarrow$ ⑤, ① $\leftrightarrow$ ⑤ are theorems (Borcherds 1995; Gritsenko–Nikulin 1998); they relate the views in the *Given* part of the manuscript. The edge ② $\leftrightarrow$ ③ is the Pfaffian–Dirac theorem of this manuscript, conditional on (Do) , (O_1) , $(O_1)^+$, (O_2) ; it relates the *Given* part to the *Built* part. The edge ① $\leftrightarrow$ ③ is the two-step composite $D_X = \Delta_5 = \text{Pf}_{\text{prot}}(\mathfrak{D}_X^{DI})$, with the orientation match $\epsilon_o = \nu_{\Delta_5}$ on $W^{(2)}(\Lambda_{II}^{2,1})$ supplying the multiplier; the protected primitive Lie superalgebra $\mathcal{P}_X$ recovers the BKM target by $\mathcal{P}_X \cong \mathfrak{g}_{\Delta_5}$.

0.0.0.3 The level- n scalar shadow. On the closed cobordism $K_3 \times E \mapsto \text{pt}$,

$$Z_{\text{BPS}}^{K_3 \times E} = (\Delta_{10})^{-1} = \Delta_5^{-2}, \quad \Delta_{10} = \Delta_5^2 \in M_{10}(\mathbf{Sp}_4(\mathbb{Z})).$$

Here $\Delta_{10} = \Delta_5^2$ is the theta-normalised weight-ten cusp form on $\mathbf{Sp}_4(\mathbb{Z})$, equivalently $\Phi_{10}^{\text{un}} = \chi_{10} = \Delta_5^2$ in the unimodular Igusa convention ([?Igusa1962]; the multiplier ν_{Δ_5} squares to trivial [?MAASS:1]); the Oberdieck–Pixton-monic normalisation $\chi_{10}^{\text{OP}} := D_5^2 = 4096^{-1} \Delta_5^2$ (with $D_5 := 64^{-1} \Delta_5$ the Borcherds-monic form so that $[q^{1/2} r^{1/2} s^{1/2}] D_5 = 1$) gives the equivalent form $Z_{\text{OP}}^X = -(\chi_{10}^{\text{OP}})^{-1} = -4096 \Delta_5^{-2}$, where $4096 = 64^2$ is the squared theta-leading constant $[q^{1/2} r^{1/2} s^{1/2}] \Delta_5 = 64$ of the Igusa form and the minus sign is the OP scalar branch convention [?0B, ?0P and]. $Z_{\text{BPS}}^{K_3 \times E}$ is the level- n protected trace of $\mathfrak{A}_{K_3 \times E}$; promotion to a 3d gravitational path integral on $K_3 \times E$ requires the Hall–Borcherds residual of Vol II, and is not claimed here.

0.0.0.4 The κ -tuple of $K_3 \times E$. Δ_5 sits in a five-coordinate invariant of the chiral E_3 -algebra $\mathfrak{A}_{K_3 \times E}$,

$$\kappa(\mathfrak{A}_{K_3 \times E}) = (\kappa_{\text{cat}}, \kappa_{\text{ch}}^{\text{Hodge}}, \kappa_{\text{ch}}^{\text{Heis}}, \kappa_{\text{BKM}}, \kappa_{\text{fiber}}) = (o, o, 3, 5, 24).$$

The categorical Euler is $\kappa_{\text{cat}} = \chi(\mathcal{O}_{K_3 \times E}) = \chi(\mathcal{O}_{K_3}) \chi(\mathcal{O}_E) = 2 \cdot o = o$. The Hodge supertrace $\kappa_{\text{ch}}^{\text{Hodge}}(K_3 \times E) = \sum_q (-1)^q h^{o,q}(K_3 \times E) = o$. The Heisenberg rank $\kappa_{\text{ch}}^{\text{Heis}} = 3$ is the rank of the Heisenberg vertex algebra obtained from the polarised K_3 Mukai sublattice $H^{o,o}(K_3) \oplus \text{NS}(S) \oplus H^{2,2}(K_3)$ of the rank-22 K_3 cohomology lattice after restriction to the rank-three type-II quotient. The BKM coordinate

$$\kappa_{\text{BKM}}(\Phi_N) = c_N(o)/2$$

(Borcherds 1995; Gritsenko 1999) reads, at the paramodular input $N = 1$, $c_1(o) = 10$, giving $\kappa_{\text{BKM}}(\Delta_5) = 5$. The fibre coordinate $\kappa_{\text{fiber}} = \chi_{\text{top}}(K_3) = 24$. The additive identity $\kappa_{\text{BKM}} = \kappa_{\text{ch}}^{\text{Hodge}} + \chi(O_{\text{fiber}})$ fails at every $N \in \{1, 2, 3, 4, 6\}$: the Lefschetz Euler $\chi_{\text{top}}(K_3) = 24$ and the holomorphic Euler $\chi(O_{K_3}) = 2$ are distinct invariants of the K_3 fibre, and κ_{BKM} sees the $c_N(o)/2$ data of the Borcherds input rather than either of them. Each subscript on κ records a distinct invariant; the unsubscripted handle leaves the categorical dimension unspecified, and is replaced on first use by the named coordinate.

PENTADIC Δ_5

1 THE ONE CHIRAL E_3 -ALGEBRA AND ITS FIVE VALUES

The frontispiece names Δ_5 at five categorical-dimensional levels. A single underlying object is required, on which the five readings are different evaluations. The Calabi–Yau threefold $K_3 \times E$ supplies it: the Stage-1 holomorphic factorisation algebra of Kontsevich–Tamarkin and Costello–Gwilliam–Li at the CY-3 category of coherent sheaves (status: established cross-volume, ambient: chiral E_3 -prefactorisation algebra associated to a CY-3 dg-category, [`?KontsevichOperadsMotives`, `?CostelloGwilliam1`, `?CostelloGwilliam2`]),

$$\mathfrak{A}_{K_3 \times E} = \Phi_3^{\text{FA}}(D^b \text{Coh}(K_3 \times E)) \in E_3\text{-HolFA}(K_3 \times E)$$

[`?KontsevichOperadsMotives`, `?TamarkinLittleDisks`, `?LambrechtsVolicLittleDisks`, `?CostelloGwilliam1`, `?CostelloGwilliam2`]. The protected sector cuts down $\mathfrak{A}_{K_3 \times E}$ to the index-graded piece on which the Borchers determinant is a single line; that single line is Δ_5 , and the five views of the frontispiece are five distinct evaluations of $\mathfrak{A}_{K_3 \times E}$ at five categorical- dimensional levels.

Definition 1.1 (Pentadic Δ_5). The five-tuple

$$(D_X, \text{den}(\mathfrak{g}_{\Delta_5}), \text{Pf}_{\text{prot}}(\mathfrak{D}_X^{DI}), \text{Disc}(\mathcal{P} \rightarrow \Delta), \Theta_{\text{sing}}^{\text{Howe}}(\phi_{\text{o},1}))$$

of values of $\mathfrak{A}_{K_3 \times E}$ at the five categorical- dimensional levels enumerated in the frontispiece is the *pentadic* Δ_5 . The reading of Δ_5 is fixed by the level: at level ①, Δ_5 is D_X ; at level ②, Δ_5 is the function whose value at $2Z$ is $64 \text{den}(\mathfrak{g}_{\Delta_5})$; at level ③, Δ_5 is the protected Pfaffian $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{DI})$ under $(D\text{o})$, $(O1)$, $(O1)^+$, $(O2)$; at level ④, Δ_5^{-2} is the mirror discriminant section, conjecturally; at level ⑤, Δ_5 is the Howe lift of $\phi_{\text{o},1}$.

2 THE ① $\leftrightarrow$ ② EDGE: BORCHERDS–GRITSENKO–NIKULIN

The Borchers product expansion of the weight-zero index-one weak Jacobi form $\phi_{\text{o},1}$ is the genus-two cusp form

$$\Delta_5(Z) = 64 q^{1/2} r^{1/2} s^{1/2} \prod_{(n,l,m) \in \Gamma_{\text{eff}}} (1 - q^n r^l s^m)^{f(nm,l)},$$

where $Z = \begin{pmatrix} z_1 & z_2 \\ z_2 & z_3 \end{pmatrix}$, $q = e^{2\pi i z_1}$, $r = e^{2\pi i z_2}$, $s = e^{2\pi i z_3}$, and $f(n, l)$ are the Fourier coefficients of $\phi_{\text{o},1}$ (the weak Jacobi form whose discriminant decomposition records $f(\text{o}, \text{o}) = 10$, $f(\text{o}, \pm 1) = 1$, $f(1, \text{o}) = 108$, $f(1, \pm 1) = -64$, $f(1, \pm 2) = 10$, $\dots$; verified arithmetically by `compute/verify_square_root.py` [`?Borchers95`, `?GN`, `?EZ:1`]). The leading constant $[q^{1/2} r^{1/2} s^{1/2}] \Delta_5 = 64 = 2^6$ is the theta-leading coefficient (Igusa); the chamber $0 < |s| \ll |q| \ll |r|^{-1} \ll 1$ fixes the active semigroup Γ_{eff} of cusp-completed exponents.

The Gritsenko–Nikulin denominator identity converts the same product into the Weyl–Kac form on the type-II sublattice $\Lambda_{II}^{2,1} \subset \Lambda^{2,1} = \Lambda^{(1,1)} \oplus \langle 2 \rangle$:

$$\text{den}(\mathfrak{g}_{\Delta_5}) = \sum_{w \in W^{(2)}(\Lambda_{II}^{2,1})} \text{sgn}(w) w(e^\rho) = 64^{-1} \Delta_5(2Z),$$

with Weyl vector

$$\rho = \frac{1}{2}(\delta_1 + \delta_2 + \delta_3) = f_2 - \frac{1}{2}f_3 + f_{-2}$$

on the three simple type-II reflections $\delta_1 = 2f_2 - f_3$, $\delta_2 = 2f_{-2} - f_3$, $\delta_3 = f_3$ (Cartan matrix $(\delta_i, \delta_j) = 4\mathbf{I}_3 - 2\mathbf{J}_3 = \begin{pmatrix} 2 & -2 & -2 \\ -2 & 2 & -2 \\ -2 & -2 & 2 \end{pmatrix}$, $\mathbf{J}_3$ the all-ones matrix, verified by `compute/verify_lattice.py`); index-two subgroup $W^{(2)}(\Lambda_{II}^{2,1})$ generated by these three reflections [?GN, ?GNPartI, ?GNII]. The signed root supermultiplicities recover the Fourier coefficients of $\phi_{o,i}$ under the doubling $\alpha(n, l, m) = 2nf_2 - lf_3 + 2mf_{-2}$:

$$\text{smult } \mathfrak{g}_{\Delta_5, \alpha(n, l, m)} = f(nm, l) \quad \text{on } (n, l, m) \in \Gamma_{\text{act}} = \{(n, l, m) \in \Gamma_{\text{eff}} \mid f(nm, l) \neq 0\}.$$

The edge ① $\leftrightarrow$ ② is the joint statement: the automorphic section $D_X = \Delta_5$ of $\mathcal{L}^{\otimes 5} \otimes \nu_{\Delta_5}$ on Siegel space *equals* the BKM denominator (after the constant 64^{-1} and the variable doubling $Z \mapsto 2Z$). The proof is Borcherds 1995 plus the Gritsenko–Nikulin identification of the underlying lattice with $\Lambda_{II}^{2,1}$ and the Weyl-group action with $W^{(2)}(\Lambda_{II}^{2,1})$ [?Borcherds95, ?GN]. Status: proved (ambient: $(\infty, 1)$ -formal automorphic forms on $\mathbf{Sp}_4(\mathbb{R})$ and graded BKM superalgebras).

3 THE ① $\leftrightarrow$ ⑤ AND ② $\leftrightarrow$ ⑤ EDGES: THE SINGULAR THETA LIFT

The Borcherds product expansion of the previous section converts $\phi_{o,i}$ into Δ_5 by an arithmetic identity in Γ_{eff} ; the question of *where* this identity comes from has a single answer. The exceptional isomorphism $\mathbf{Sp}_4 \simeq \text{Spin}(3, 2)$ identifies the genus-two Siegel space $\mathbf{Sp}(2, \mathbb{Z}) \backslash \mathcal{H}_2$ with the type-IV symmetric domain $\mathbf{O}(3, 2)_+ \backslash \mathcal{D}_{\mathbf{O}(3, 2)}$ attached to the signature- $(3, 2)$ lattice $\Lambda^{3,2} = \Lambda^{(1,1)} \oplus \Lambda^{(1,1)} \oplus \langle 2 \rangle$; the Howe correspondence $\text{Mp}_4(\mathbb{R}) \leftrightarrow \mathbf{O}(3, 2)$ of the dual reductive pair lifts modular forms in genus one onto this domain [?Weil1964, ?Shimura1973]. Borcherds’s regularisation of the Howe correspondence on the principal part of a weak Jacobi-form input produces a meromorphic Siegel modular form whose divisor is prescribed by the principal part:

$$\Theta_{\text{sing}}^{\text{Howe}}(\phi) = \int_{\text{SL}_2(\mathbb{Z}) \backslash \mathbb{H}}^{\text{reg}} \langle \phi(\tau), \Theta_{\Lambda^{3,2}}(\tau, Z) \rangle \frac{du dv}{v^2}, \quad \tau = u + iv,$$

on the Weil-representation pairing of the vector-valued weakly-holomorphic input ϕ (the theta decomposition of $\phi_{o,i}$ into the discriminant form of $\Lambda^{3,2}$) against the Siegel theta series of the signature- $(3, 2)$ lattice $\Lambda^{3,2}$ — not the rank-three target lattice $\Lambda_{II}^{2,1}$, which receives the BKM root structure under doubling — together with the antiholomorphic dependence required for absolute convergence after regularisation; the explicit construction is Chapter 5, Definition 1.9 and Theorem 1.10 [?Borcherds95, ?B]. At the input $\phi = \phi_{o,i}$ the regularised integral computes

$$\Theta_{\text{sing}}^{\text{Howe}} : J_{o,i}^{\text{wk}} \longrightarrow M_5(\mathbf{Sp}_4(\mathbb{Z}), \nu_{\Delta_5}), \quad \phi_{o,i} \longmapsto \Delta_5,$$

giving the edge ① $\leftrightarrow$ ⑤. Reading the same lift through the Gritsenko–Nikulin denominator identity, $\phi_{o,i}$ is the character of the index-one Heisenberg sector and the multiplicative lift over $\Lambda_{II}^{2,1}$ is exactly $\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1} \Delta_5(2Z)$, giving the edge ② $\leftrightarrow$ ⑤ [?Borcherds95, ?GN, ?GNPartI, ?GNII]. Status: proved (ambient: regularised theta integral on the symmetric domain of $\mathbf{O}(2, n)$).

4 THE ② ↔ ③ EDGE: THE PFAFFIAN–DIRAC THEOREM

The four edges proved by Borchers and Gritsenko–Nikulin assemble Δ_5 on the automorphic and BKM-denominator sides; the chiral E_3 -algebra $\mathfrak{A}_{K_3 \times E}$ of §1 is the candidate operator-level origin from which both sides descend. The structural question of the manuscript is whether the protected primitive sector of $\mathfrak{A}_{K_3 \times E}$, after finite-stage Hall reconstruction with cosection-reduced d-critical orientation, hybrid $\text{Ran}^{\text{hyb}}(E)$ carrier, Pfaffian line, source coalgebra, Koszul cobar, and the recognition map, recovers the BKM denominator algebra $\mathfrak{g}_{\Delta_5}$ on the nose. The answer is the Pfaffian–Dirac theorem, conditional on four named obstructions (Do) , (O_1) , $(O_1)^+$, (O_2) named at §5.8 of the manuscript and tracked across the notes/swarm_∗ ledgers.

THEOREM 4.1 (*Pfaffian–Dirac*). *Conditional on (Do) , (O_1) , $(O_1)^+$, (O_2) ; ambient: cosection-twisted reduced d-critical chart on $K_3 \times E$ at finite cusp-truncation height R , with chain-level pro-object, Mittag-Leffler vanishing of the inverse system in R , and Pfaffian orientation o with Weyl character ϵ_o . The protected Pfaffian of the chiral E_3 -algebra $\mathfrak{A}_{K_3 \times E}$, presented as the limit $\mathfrak{D}_X^{DI} = \varprojlim_R \mathfrak{D}_{X,R}^{DI}$, satisfies*

$$\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{DI}) = \Delta_5, \quad \epsilon_o = \nu_{\Delta_5} \quad \text{on} \quad W^{(2)}(\Lambda_{II}^{2,1}),$$

and the protected primitive Lie superalgebra is

$$\mathcal{P}_X := H^0 \text{Prim}_{\text{prot}}(\overline{\Pi_{X,*}^{\ominus} \mathcal{F}_X}) / \overline{\text{Rad}\langle \cdot, \cdot \rangle_X} \cong \mathfrak{g}_{\Delta_5}.$$

The four obstructions name the exact conditioning. (Do) is the Do -degeneration limit of the cosection-reduced d-critical orientation theory: the limiting orientation in the Do -degenerate fibre is constructed and matches the orientation of the generic fibre under the cosection-twist [?BBDJS2015, ?JoyceUpmeier, ?JoyceTanakaUpmeier]. (O_1) is the strong reduced orientation on the $K_3 \times E$ -quotient self-Ext frame bundle: the square root of the determinant line of the cotangent complex extends across the cusp boundary [?CaoGrossJoyce, ?Bojko]. $(O_1)^+$ is the Weyl-equivariant transport of the orientation along reflections in the index-two reflection subgroup $W^{(2)}(\Lambda_{II}^{2,1})$: the orientation cocycle $c_o \in Z^2(W^{(2)}(\Lambda_{II}^{2,1}); \mathbb{F}_2)$ trivialises in H^2 , and the resulting character $\epsilon_o : W^{(2)}(\Lambda_{II}^{2,1}) \rightarrow \{\pm 1\}$ is well-defined on generators. (O_2) is the local Pfaffian wall normal form on type-II walls: at each simple type-II wall s_{δ_i} the rank-one normal Pfaffian crossing of Definition 0.21 gives the local sign $(-1)^{N_{\delta_i}^{\text{Pf}}} = -1$, and the Hall-side sign sum on the half-Hilbert orbit of the self-Ext stack — of size $2^{12} = 4096$, arithmetically the squared theta-leading $([q^{1/2} r^{1/2} s^{1/2}] \Delta_5)^2 = 64^2$ of the Oberdieck–Pixton scalar normalisation — collects into the single character value $\epsilon_o(s_{\delta_i}) = \nu_{\Delta_5}(s_{\delta_i})$ on each generator (Appendix E).

Remark 4.2 (*The conditioning is named precisely*). The four obstruction classes (Do) , (O_1) , $(O_1)^+$, (O_2) are the exact proof obligations the Built part of the manuscript reduces to; discharging any of them tightens the conditioning of Theorem 4.1. Two readings of Δ_5 attach to two distinct categorical dimensions: at the scalar-shadow levels Δ_5 is a charge-graded virtual section of the modular line, the protected scalar shadow of the chiral E_3 -algebra $\mathfrak{A}_{K_3 \times E}$; at the operator level ③ it is the protected Pfaffian $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{DI})$ of the operator-level pro-object, conditional on the four obstructions. The distinction is structural, not nominal: the operator-level package carries the orientation character ϵ_o that the scalar shadow has squared away, by view ③, and recovers it as ν_{Δ_5} on $W^{(2)}(\Lambda_{II}^{2,1})$ by $(O_1)^+$ and (O_2) .

5 THE ④ VIEW: MIRROR DISCRIMINANT

Conjectural (ambient: variation of Hodge structure on a one-parameter K_3 -mirror period family). On the one-parameter K_3 -mirror period family $\mathcal{P} \rightarrow \Delta$ — the mirror of $K_3 \times E$ restricted to its rank-one polarised subvariation — the discriminant locus is cut by

$$\text{Disc}(\mathcal{P} \rightarrow \Delta) = \Delta_5^{-2} \big|_{\Delta},$$

in the conjectural identification of the mirror period family with the $\mathbf{Sp}(4, \mathbb{Z})$ -quotient of Siegel space restricted along the Mukai self-mirror branch. The reading is heuristic in the body of this manuscript and does not enter as a hypothesis of any theorem; it is recorded because the identification, when proved, would equate the discriminant divisor of the K_3 -mirror period family with the polar divisor of Δ_5^{-2} , itself twice the zero divisor of Δ_5 ; see Conjecture 1.1.

6 THE LEVEL- n SCALAR SHADOW

The scalar partition function on the closed cobordism $K_3 \times E \mapsto \text{pt}$ is the Borchers determinant of the protected K_3 index, computed by Oberdieck–Pixton 2018 and Oberdieck–Pandharipande 2018 [?OB, ?OPand]. Status: proved (ambient: virtual count of Hilbert schemes of $K_3 \times E$ with $(-p_{\text{DT}})^n$ chamber). Two normalisations of the same weight-ten cusp form on $\mathbf{Sp}_4(\mathbb{Z})$ are in use, and naming them is necessary because both appear in the manuscript:

- *Theta-normalised.* $\Delta_{10}(Z) := \Delta_5(Z)^2 \in M_{10}(\mathbf{Sp}_4(\mathbb{Z}))$ (the multiplier ν_{Δ_5} squares to trivial; the leading coefficient $[q \ r \ s] \Delta_{10} = 64^2 = 4096$).
- *Oberdieck–Pixton-monic.* $\chi_{10}^{\text{OP}}(Z) := D_5(Z)^2 = 64^{-2} \Delta_5(Z)^2 = 4096^{-1} \Delta_{10}(Z)$ (so $[q \ r \ s] \chi_{10}^{\text{OP}} = 1$).

Oberdieck–Pixton and Oberdieck–Pandharipande prove [?OB, ?OPand]

$$Z_{\text{OP}}^X = -(\chi_{10}^{\text{OP}})^{-1} = -4096 \Delta_5^{-2},$$

and the level- n protected scalar shadow is

$$Z_{\text{BPS}}^{K_3 \times E} = (\Delta_{10})^{-1} = \Delta_5^{-2} = -\frac{1}{4096} Z_{\text{OP}}^X.$$

The factor $4096 = 64^2$ is the squared theta-leading constant of Δ_5 ; the OP minus sign is a scalar branch convention absorbed into the OP $(-p_{\text{DT}})^n$ chamber and is independent of the orientation character ϵ_θ of view ③.

$Z_{\text{BPS}}^{K_3 \times E}$ is the level- n protected trace of $\mathfrak{A}_{K_3 \times E}$: a charge-graded virtual partition function, not the graded dimension of any Hilbert space. Promotion of the scalar shadow to a 3d gravitational path integral on $K_3 \times E$ requires the Hall–Borchers residual of Vol II, supplied by the universal-stage chain (primitive package $\mathbf{P} \rightarrow$ boundary chart $\mathbf{C} \rightarrow$ MC twisting $\mathbf{S} \rightarrow$ derived chiral centre $\mathbf{Z} \rightarrow$ ambient operator algebra $\mathbf{A}$) at the boundary vacuum b of the open factorisation dg-category on $(K_3 \times E, D, \tau)$; promotion lies beyond this manuscript. The local witness is the formal-Darboux stalk plus the holomorphic de Rham obstruction class of the mixed-holomorphic-topological strings programme [?KontsevichOperadsMotives, ?CostelloLiBCOV, ?CostelloLiOpenClosedBCOV], with global compactness of the path integral conditional on the obstruction class vanishing on $K_3 \times E$.

Remark 6.1 (Reconstitution order). The five views are ordered: primitive objects (the chiral E_3 -algebra $\mathfrak{A}_{K_3 \times E}$ and its four obstructions (D_0) , (O_1) , $(O_1)^+$, (O_2) ; the Cartan datum of $\mathfrak{g}_{\Delta_5}$) first; cross-level

identities ((②↔③)) Pfaffian–Dirac, ($\epsilon_o = \nu_{\Delta_5}$) orientation match, ($\mathcal{P}_X \cong \mathfrak{g}_{\Delta_5}$) primitive recognition) second; scalar modular forms last (the section D_X , the BKM denominator $64^{-1}\Delta_5(2Z)$, the partition function Δ_5^{-2}). A statement is not primitive if it is true only after passing to a trace, choosing a chamber, completing a category, or installing descent data. On each first appearance the scalar reading is named: the theta-normalised form Δ_5 with leading $[q^{1/2}\tau^{1/2}s^{1/2}]\Delta_5 = 64$; the BKM denominator $64^{-1}\Delta_5(2Z)$ with the doubling $2Z$ on the type-II sublattice; the OP-monic form $D_5 = 64^{-1}\Delta_5$ with $\chi_{10}^{\text{OP}} = D_5^2 = 4096^{-1}\Delta_5^2$ the Oberdieck–Pixton normalisation of the squared scalar shadow.

7 CROSS-VOLUME PLACEMENT

The pentadic Δ_5 is one identification at one specific level of the Beilinson tower of the wider research constellation: the level-5 chiral shadow of Vol I’s bar $B(A_b)$ on the tangential log curve $(K_3 \times E, D, \tau)$; the closed-colour shadow of the bicoloured primitive package of Vol II’s Universal Holography master theorem with terminal value $c_1(o)/2 = 5 = \kappa_{\text{BKM}}(\Delta_5)$; the ($d = 3$, rank 3) chamber of Vol III’s cross-stratum sibling census, populated by the BKM target $\mathfrak{g}_{\Delta_5}$ under the universal identity $\kappa_{\text{BKM}}(\Phi_N) = c_N(o)/2$ applied at the paramodular row $(t, N; k) = (1, 1; 5)$ of the Gritsenko–Cléry catalogue [?GC, ?GN]; the formal-Darboux stalk $A_{\partial, N}^{\text{cl}} = C_{\text{CE}}^{\bullet}(\mathfrak{gl}_N, \text{Kosz}([\phi_1, \phi_2]))$ of the chiral Hochschild cochain of mixed-holomorphic-topological strings at the Dirac brane vacuum. Promotion of the level- n scalar shadow $Z_{\text{BPS}}^{K_3 \times E} = \Delta_5^{-2}$ to a global compact 3d gravitational path integral on $K_3 \times E$ requires both the holomorphic de Rham obstruction class $\text{ob}_{\text{dR}}^{\text{hol}}(K_3 \times E)$ to vanish and Vol II’s Hall–Borcherds residual to discharge the dynamical-metric content; both conditions are open.

CONTENTS

FRONTISPIECE	3
o.o.o.1 The five views.	3
o.o.o.2 The cross-level identities.	4
o.o.o.3 The level- n scalar shadow.	4
o.o.o.4 The κ -tuple of $K_3 \times E$	4
I PENTADIC Δ_5	7
1 The one chiral E_3 -algebra and its five values	7
2 The ① ↔ ② edge: Borcherds–Gritsenko–Nikulin	7
3 The ① ↔ ⑤ and ② ↔ ⑤ edges: the singular theta lift	8
4 The ② ↔ ③ edge: the Pfaffian–Dirac theorem	9
5 The ④ view: mirror discriminant	10
6 The level- n scalar shadow	10
7 Cross-volume placement	11
CONTENTS	11

I	Given	21
2	K ₃ ×E SETUP, MUKAI DICTIONARY, LATTICE POLARIZATION, NORMAL-ORDERED GRAM COCYCLE, HYBRID $\text{Ran}^{\text{hyb}}(E)$ CARRIER	23
1	Smooth K ₃ ×E threefold and Mukai lattice	24
1.1	The compact target	24
1.2	Mukai lattice on the K ₃ factor	24
1.3	Künneth and the formal even Mukai sector on $X = S \times E$	25
2	Quadratic Gram map and lattice polarization	26
2.1	Igusa variables and the Gram-index lattice	26
2.2	The Gram map and its bilinear cocycle	27
2.3	Formal lift of every Gram triple	28
3	Normal-ordered Gram extension	29
3.1	Central extension and the additive normal-ordered Gram homomorphism	29
3.2	The fixed-lift raw-pushforward obstruction theorem	30
3.3	Normal-ordering flag formula	31
4	The hybrid $\text{Ran}^{\text{hyb}}(E)$ factorization carrier	32
4.1	Projection-locality obstruction	32
4.2	The hybrid carrier	32
4.3	Mixed local/wrapped correspondences	33
4.4	Wrapped anchor and translation linearization	33
4.5	Cusp polarization and chamber transport	35
5	Local protected observable algebra (sectorial source datum)	35
6	The four obstructions and the conditional comparison target	36
7	Notation summary and chapter outputs	37
3	VIEW I: THE BORCHERDS–GRITSENKO–NIKULIN SECTION	39
1	The pentadic level and the Beilinson cut	39
1.0.0.1	Three names.	39
1.0.0.2	Beilinson cut: scope of View I.	39
2	The genus-two Siegel space and the Borchers chamber	39
2.0.0.1	Choices fixed once and for all.	40
3	The weak Jacobi input and the K ₃ elliptic genus	40
3.0.0.1	Borchers-weight handle.	41
4	The theta-constant normalization	41
5	The Borchers–Gritsenko–Nikulin section	41
5.1	The product expansion	41
5.2	The View I package	42
6	Weight, multiplier, and divisor	42
6.1	The Borchers weight	42
6.2	The Maass multiplier system	43
6.3	The diagonal divisor	43
6.4	Maass-character values on the type-II walls	43
7	Identification with the virtual K_0 -determinant	44
8	The squared scalar shadow and OP normalization	45

	8.o.o.1	The OP scalar branch.	45
	8.o.o.2	Three names for the squared form, distinguished.	45
9		The View II shadow: the $2Z$ denominator	45
10		The $\mathrm{Sp}(4) \simeq \mathrm{Spin}(3,2)$ bridge to View V	45
11		Summary of View I	46
	11.o.o.1	What View I does not assert.	46
4		THE BORCHERDS–KAC–MOODY DENOMINATOR ALGEBRA $\mathfrak{g}_{\Delta_5}$ ON $\Lambda_{II}^{2,1}$	49
1		The hyperbolic lattice $\Lambda^{2,1} = \Lambda^{(1,1)} \oplus [2]$ and its type-II sublattice	49
2		Real walls, the type-II Weyl chamber, and chamber automorphism	50
3		Real and imaginary roots; the Borchers–Kac–Moody algebra $\mathfrak{g}_{\Delta_5}$	53
4		Generators and relations: Cartan, Chevalley, Serre, Borchers orthogonality, super sign	54
5		The denominator: Weyl alternating sum and Borchers product	56
6		The active support: Lorentzian cone and finite checks	58
7		Maass multiplier on $\mathcal{W}^{(2)}(\Lambda^{2,1})$	62
8		The Borchers-weight characteristic $\kappa_{\mathrm{BKM}}(\Delta_5) = 5$	63
9		The denominator-algebra theorem and its scope	64
10		Boundary with View I (automorphic) and View V (theta lift)	65
11		The cross-volume frame: $\mathfrak{g}_{\Delta_5}$ in the κ -tuple	65
12		Structural properties of $\mathfrak{g}_{\Delta_5}$	66
13		Comparison with the chiral shadow lattice Γ_{ind}	67
14		Comparison: rank-three hyperbolic vs. Borchers correction	68
15		The character of the chiral shadow	68
16		The denominator algebra of the chiral source-target frame	69
17		The closing identity and its conditioning	70
18		Adversarial closure	71
19		Coda: $\mathfrak{g}_{\Delta_5}$ as one of five views	73
5		THE SINGULAR THETA LIFT	75
1		The singular theta lift of $\phi_{o,I}$	75
	1.1	Notation and the canonical name	75
	1.2	The weak Jacobi form $\phi_{o,I}$	75
	1.3	The signature- $(3, 2)$ lattice and the type-IV domain	76
	1.4	The exceptional isomorphism $\mathbf{Sp}_4 \simeq \mathbf{Spin}(3, 2)$	77
	1.5	The Howe correspondence $\mathbf{Mp}_4(\mathbb{R}) \leftrightarrow \mathbf{O}(3, 2)$	78
	1.6	The Borchers singular-theta integral	78
	1.7	From $\phi_{o,I}$ to a vector-valued Weil-representation form	79
	1.8	Theta lift identifies Δ_5	80
	1.9	The multiplier system and the Weyl vector	81
	1.10	Compatibility with View II: lift’s image is the BKM denominator	82
	1.11	Compatibility with View I and the cross-level identities	82
	1.12	The level-five universal weight identity	83
	1.13	Beilinson cut: the lift is the structural content	83
	1.14	Consistency with the manuscript’s lattice conventions	83

1.15	Closing remarks	84
II	Built	85
6	THE DIRAC-IGUSA PRO-OBJECT $\mathfrak{D}_X^{\text{DI}}$ ON $K_3 \times E$ (VIEW ③)	87
o.1	The hybrid factorization carrier $\text{Ran}^{\text{hyb}}(E)$	87
o.2	Cosection-twisted reduced d-critical orientation	92
o.3	Hall correspondences: product, coproduct, primitives	95
o.4	Dirac operator and protected Pfaffian	99
o.5	Koszul source coalgebra and chiral MC twisting	102
o.6	Primitive recognition apparatus	106
o.7	Eight build algorithms	III
o.8	Four named obstructions	II6
III	Certified	119
7	THE CROSS-LEVEL IDENTITY (2) $\leftrightarrow$ (3): PFAFFIAN-DIRAC, ORIENTATION CHARACTER, PRIMITIVE RECOGNITION	121
1	Data, obstructions, licensing	122
1.1	The pro-object $\mathfrak{D}_X^{\text{DI}}$ and its eight finite-stage components	122
1.2	The four obstruction packages: (Do), (O1), (O1) ⁺ , (O2)	122
1.2.0.1	(Do) — Do-degeneration endpoint of the cosection-reduced d-critical orientation theory.	122
1.2.0.2	(O1) — strong multiplicative reduced orientation datum on the reduced $K_3 \times E$ -quotient self-Ext frame bundle.	123
1.2.0.3	(O1) ⁺ — Weyl-equivariant determinant-line transport along $\mathcal{W}^{(2)}(\Lambda_{II}^{2,1})$ reflections.	123
1.2.0.4	(O2) — Pfaffian local model and wall atlas on type-II walls.	123
1.3	Surgical fix: the OP minus sign is a scalar branch convention	123
2	The Pfaffian-Dirac theorem	124
3	The orientation character identity	125
4	The primitive recognition theorem	127
5	Cross-volume consistency	130
5.1	Vol II Construction Problem I	130
5.2	Vol III chain fusion at level 2	130
5.3	BCOV / mixed holomorphic-topological strings	131
6	Residual obstructions and open subroutes	131
6.0.0.1	(Do).	131
6.0.0.2	(O1).	131
6.0.0.3	(O1) ⁺	131
6.0.0.4	(O2).	131
6.0.0.5	Recognition criterion (S1)–(S10).	132

6.0.0.6	Vol III Hall–Drinfeld–Borcherds spectral sequence on compact non-toric CY_3	132
6.0.0.7	Honest summary.	132

IV Shadowed 133

8	THE MIRROR DISCRIMINANT	135
1	The mirror discriminant	135
1.1	The K_3 mirror-period family	135
1.1.0.1	The rank-three case.	136
1.2	The $K_3 \times E$ period family	137
1.3	The discriminant locus	138
1.4	The conjecture	139
1.4.0.1	Evidence.	139
1.4.0.2	Status.	140
1.4.0.3	What this conjecture supplies.	140
1.5	What the conjecture does not assert	140
1.6	Lattice-convention check	140
1.7	Open obligations	141
1.8	Cross-repository consistency	141
9	THE LEVEL- n PROTECTED SCALAR SHADOW AND THE $K_3 \times E$ REALIZATION QUESTION	143
1	The level- n protected scalar shadow on $K_3 \times E \rightarrow \text{pt}$	143
1.1	The Oberdieck–Pixton scalar identity $[\beta]$	143
1.2	Three readings of the same exponent	144
1.3	Worked Fourier-coefficient check	144
1.4	Cobordism positioning and the closed $K_3 \times E \rightarrow \text{pt}$ cobordism	145
2	Spin L -factor normalisation for Δ_5^2	145
2.1	The Saito–Kurokawa eigenline	145
2.2	Independence from the Dirac–Igusa construction	146
3	What the scalar shadow does not supply	146
3.1	Z_{BPS} is the level- n protected trace, not a path integral	146
3.2	The Hall–Borcherds residual	147
3.3	Local-to-global in mixed holomorphic–topological language	147
4	The $K_3 \times E$ threefold realization question	148
4.1	From the scalar to the operator-level object	148
4.2	The orientation datum the squared scalar cannot recover	148
4.3	The four obstructions and the realization question	148
4.4	The threefold target, not a fourfold	149
4.5	Comparison with the determinant-only construction	149
5	Open frontiers	149
5.1	The Hall–Borcherds residual revisited	149
5.2	The gravity-line operator algebra and the Pentagon-face trace	150
5.3	Mathieu and umbral moonshine adjacency	150

5.4	The closed-cobordism partition reads as one	150
6	Open obstructions and the next frontier	151
6.1	The four named obstructions	152
6.1.1	(Do): Do-degeneration limit of the cosection-reduced d-critical orientation theory	152
6.1.1.1	Statement.	152
6.1.1.2	Stage and licensing.	152
6.1.1.3	Swarm probe.	152
6.1.1.4	Reasoning.	152
6.1.1.5	Next concrete construction.	153
6.1.2	(O ₁): strong reduced orientation on the $K_3 \times E$ -quotient self-Ext frame bundle	153
6.1.2.1	Statement.	153
6.1.2.2	Stage and licensing.	153
6.1.2.3	Swarm probe.	153
6.1.2.4	Reasoning.	154
6.1.2.5	Next concrete construction.	154
6.1.3	(O ₁) ⁺ : Weyl-equivariant transport of the orientation along $W^{(2)}(\Lambda_{II}^{2,1})$ reflections	154
6.1.3.1	Statement.	154
6.1.3.2	Stage and licensing.	154
6.1.3.3	Swarm probe.	154
6.1.3.4	Reasoning.	154
6.1.3.5	Next concrete construction.	155
6.1.4	(O ₂): local Pfaffian wall normal form on type-II walls	155
6.1.4.1	Statement.	155
6.1.4.2	The 4096 sign sum.	155
6.1.4.3	Stage and licensing.	155
6.1.4.4	Swarm probe.	155
6.1.4.5	Reasoning.	155
6.1.4.6	Next concrete construction.	155
6.2	Mathieu and umbral moonshine adjacency	156
6.2.0.1	The conjectural connection to $\mathfrak{g}_{\Delta_5}$	156
6.2.0.2	What the conjecture is and what it is not.	156
6.2.0.3	Stage and licensing.	157
6.2.0.4	Cross-repo consistency.	157
6.2.0.5	Next concrete construction.	157
6.3	The next frontier	157
6.3.1	Frontier 1: the Hall–Borcherds residual	157
6.3.1.1	Statement.	157
6.3.1.2	Stage and licensing.	157
6.3.1.3	Cross-repo anchor.	157
6.3.1.4	Reasoning.	158
6.3.1.5	Next concrete construction.	158
6.3.2	Frontier 2: chain fusion for compact non-toric $K_3 \times E$	158

	6.3.2.1	Statement.	158
	6.3.2.2	Stage and licensing.	158
	6.3.2.3	Cross-repo anchor.	158
	6.3.2.4	Reasoning.	159
	6.3.2.5	Next concrete construction.	159
	6.3.3	Frontier 3: the gravity-line operator algebra	159
	6.3.3.1	Statement.	159
	6.3.3.2	Stage and licensing.	159
	6.3.3.3	Cross-repo anchor.	159
	6.3.3.4	The κ -tuple.	159
	6.3.3.5	Reasoning.	159
	6.3.3.6	Next concrete construction.	160
1		Lattice computations	160
	1.1	The hyperbolic plane $U = \Pi_{1,1}$	161
	1.2	The Lorentzian root lattice $\Lambda_{II}^{2,1}$	161
	1.3	The K_3 lattice $\Lambda_{K_3} = \Pi_{3,19}$	164
	1.4	The lattice-polarized K_3 sublattice Λ_S	165
	1.5	The Mukai lattice $\tilde{H}(S, \mathbb{Z}) = \Pi_{4,20}$	166
	1.6	The formal dyonic charge lattice $\Gamma_X^{\text{form}} = \Lambda_S \oplus \Lambda_S$	167
	1.7	The signature- $(3, 2)$ lattice $\Lambda^{3,2}$ and the $\mathbf{Sp}_4 \simeq \mathbf{Spin}(3, 2)$ bridge	168
	1.8	Cross-references and consistency check	170
2		Borcherds-product expansion: $\phi_{0,1}$, the Δ_5 infinite product, and the Gritsenko–Cléry eight-row catalogue	171
	2.0.0.1	Verified Fourier table.	172
	2.0.0.2	Banked identities used below.	172
	2.0.0.3	The constant 2^6 as a theta-product leading coefficient.	173
	2.0.0.4	Theta-leading sanity check.	174
	2.0.0.5	Denominator content of the rows.	175
	2.0.0.6	Banned identification quarantine.	176
	2.0.0.7	Residual obstructions.	177
	2.0.0.8	File anchors.	177
3		Mukai–Gram dictionary on $K_3 \times E$	177
	3.1	Mukai vector and Mukai pairing on a K_3 surface	178
	3.2	Künneth on $K_3 \times E$ and the rank-3 Gram lattice	179
	3.3	D6–D2–Do dictionary	180
	3.4	Symmetric bilinear Gram cocycle	181
	3.5	Normal-ordered Gram map	182
	3.6	Three Igusa names; chart-data licensing	183
4		Hall correspondences: the level-2 functorial passage on $\mathfrak{M}_{\text{eff}}^+(K_3 \times E)$	183
A		PFAFFIAN WALL NORMAL FORM ON TYPE-II WALLS	193
	1	The three simple type-II walls	193
	2	The rank-one type-II Pfaffian crossing	194
	3	Match with the Maass multiplier	194
	4	Extension to the type-II Weyl group	195

4.1	Generation and a presentation	195
4.2	The determinant character	195
4.3	The orientation character is a $\{\pm 1\}$ -character	195
5	The (O_2) obstruction: half-Hilbert orbit transport	196
5.1	Statement of (O_2)	196
5.2	What is proved and what remains open	196
5.3	Cross-repo note: the constant 4096	196
5.4	The (O_2) discharge route	197
6	First-timelike parity split as a recognition obligation	197
6.1	Higher-degree consistency	198
7	Pseudocode for the local Pfaffian sign computation	198
8	Open obligations summary	199
9	Eight build algorithms for the Dirac–Igusa pro-object	200
9.1	Construction 1 — Charge window	200
9.1.0.1	Tag.	200
9.1.0.2	Input.	200
9.1.0.3	Output.	201
9.1.0.4	Procedure.	201
9.1.0.5	Type checks.	201
9.1.0.6	Obstructions discharged or invoked.	201
9.2	Construction 2 — Finite moduli	201
9.2.0.1	Tag.	201
9.2.0.2	Input.	202
9.2.0.3	Output.	202
9.2.0.4	Procedure.	202
9.2.0.5	Type checks.	202
9.2.0.6	Obstructions discharged or invoked.	202
9.3	Construction 3 — Hybrid wrapped factorization carrier	202
9.3.0.1	Tag.	202
9.3.0.2	Input.	203
9.3.0.3	Output.	203
9.3.0.4	Procedure.	203
9.3.0.5	Type checks.	204
9.3.0.6	Obstructions discharged or invoked.	204
9.4	Construction 4 — Reduced d-critical orientation and Joyce–Upmeyer transport	204
9.4.0.1	Tag.	204
9.4.0.2	Input.	204
9.4.0.3	Output.	204
9.4.0.4	Procedure.	204
9.4.0.5	Type checks.	205
9.4.0.6	Obstructions discharged or invoked.	205
9.5	Construction 5 — Hall correspondences and primitive part	205
9.5.0.1	Tag.	205

	9.5.0.2	Input.	205
	9.5.0.3	Output.	205
	9.5.0.4	Procedure.	206
	9.5.0.5	Type checks.	206
	9.5.0.6	Obstructions discharged or invoked.	206
	9.5.0.7	Cross-volume anchor.	207
9.6	Construction 6 — Dirac operator and Pfaffian line		207
	9.6.0.1	Tag.	207
	9.6.0.2	Input.	207
	9.6.0.3	Output.	207
	9.6.0.4	Procedure.	207
	9.6.0.5	Type checks.	208
	9.6.0.6	Obstructions discharged or invoked.	208
9.7	Construction 7 — Chiral Koszul source coalgebra		208
	9.7.0.1	Tag.	208
	9.7.0.2	Input.	208
	9.7.0.3	Output.	208
	9.7.0.4	Procedure.	208
	9.7.0.5	Type checks.	209
	9.7.0.6	Obstructions discharged or invoked.	210
	9.7.0.7	Cross-volume firewall.	210
9.8	Construction 8 — Primitive recognition		210
	9.8.0.1	Tag.	210
	9.8.0.2	Input.	210
	9.8.0.3	Output.	210
	9.8.0.4	Procedure.	210
	9.8.0.5	Type checks.	211
	9.8.0.6	Obstructions discharged or invoked.	211
	9.8.0.7	Cross-volume anchor.	211
9.9	The eight-step audit		211
	9.9.0.1	Order rigidity.	212
	9.9.0.2	Conditional theorem.	212

Part I

Given

K₃×E SETUP, MUKAI DICTIONARY, LATTICE POLARIZATION, NORMAL-ORDERED GRAM COCYCLE, HYBRID Ran^{hyb}(E) CARRIER

The arithmetic source produces three named scalars at level five. Each is the Igusa cusp form $\Phi_{10}^{1/2}$ on a different categorical chain. The automorphic chain produces the section

$$\mathcal{D}_X = \Delta_\varsigma \in M_5(\mathbf{Sp}_4(\mathbb{Z}), \nu_{\Delta_\varsigma}) \quad (\text{View ①; proved [?Borcherds95, ?GN]}).$$

The Borcherds–Kac–Moody (BKM) chain produces the denominator

$$\text{den}(\mathfrak{g}_{\Delta_\varsigma}) = 64^{-1} \Delta_\varsigma(2Z) \quad (\text{View ②; proved [?GN]}).$$

The compact Pfaffian chain produces, conditional on the four obstructions named at §6 and discharged in Chapter ??, the protected Pfaffian

$$\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}}) = \Delta_\varsigma \quad (\text{View ③; conditional}).$$

The handle “ Δ_ς ” with no superscript is reserved for working formulas; manuscript prose names which of $\mathcal{D}_X$, $\text{den}(\mathfrak{g}_{\Delta_\varsigma})$, or $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}})$ is meant. The cross-level identity ②↔③ is conditional on the obstructions (Do), (O1), (O1⁺), (O2).

The chapter assembles the geometric and lattice data feeding all three. The compact target is fixed at $X = S \times E$ with S a smooth complex projective K3 surface and E a smooth elliptic curve over $\mathbb{C}$; this is a Calabi–Yau threefold ($\dim_{\mathbb{C}} X = 3$, $K_X \cong \mathcal{O}_X$; proved/ α , [?Mukai1984, ?Mukai1987]), so $n = \chi(\mathcal{O}_Y)$ appears directly. The $(\kappa_{\text{cat}}, \kappa_{\text{ch}}^{\text{Hodge}}, \kappa_{\text{ch}}^{\text{Heis}}, \kappa_{\text{BKM}}, \kappa_{\text{fiber}}) = (0, 0, 3, 5, 24)$ tuple of [?calabi-yau-quantum-groups] (the K3×E row of the Gritsenko–Cléry catalogue) is the categorical-dimensional anchor; every use of κ in this manuscript carries one of these five subscripts.

The four-step build is geometric, then arithmetic. First, the formal dyonic charge lattice $\Gamma_X^{\text{form}} = \Lambda_S \oplus \Lambda_E$ on the Mukai sector and the algebraic/effective Hall support $\Gamma_{X,\sigma}^{\text{eff}} \subset \Gamma_X^{\text{alg}} \subset \Gamma_X^{\text{form}}$ inside it. Second, the quadratic Gram map $\Pi_X : \Gamma_X^{\text{form}} \rightarrow \Gamma_{\text{gram}} = \mathbb{Z}^3$ and its symmetric bilinear polarization $B = -\partial\Pi_X$. Third, the normal-ordered extension $\widehat{\Gamma}_X = \Gamma_X^{\text{form}} \oplus_B \Gamma_{\text{gram}}$ and the additive map $\overline{\Pi}_X(c, T) = \Pi_X(c) - T$, with the fixed-lift raw-pushforward obstruction theorem fixing the necessity of that extension. Fourth, the hybrid factorization carrier $\text{Ran}^{\text{hyb}}(E) = \text{Ran}(E)_{b=0} \sqcup \text{Ran}^{\text{wr}}(E)_{b>0}$, with the projection- locality obstruction fixing the necessity of the wrapped stratum.

1 SMOOTH $K_3 \times E$ THREEFOLD AND MUKAI LATTICE

1.1 The compact target

Definition 1.1 ($K_3 \times E$ threefold; primitive geometric datum). A $K_3 \times E$ threefold is a triple (X, S, E) in which S is a smooth complex projective K_3 surface (so $K_S \cong \mathcal{O}_S$, $h^{1,0}(S) = 0$, $h^{2,0}(S) = 1$; [?Mukai1984]), E is a smooth elliptic curve over $\mathbb{C}$ with marked origin $o_E \in E$, and

$$X := S \times E.$$

The canonical bundle is trivial, $K_X \cong p_S^* K_S \otimes p_E^* K_E \cong \mathcal{O}_X$, and Hodge invariants are

$$\dim_{\mathbb{C}} X = 3, \quad \chi_{\text{top}}(X) = \chi_{\text{top}}(S) \cdot \chi_{\text{top}}(E) = 24 \cdot 0 = 0, \quad \chi(\mathcal{O}_X) = \chi(\mathcal{O}_S) \cdot \chi(\mathcal{O}_E) = 2 \cdot 0 = 0.$$

The threefold X is a Calabi–Yau threefold (proved/ α).

Remark 1.2 (Distinguishing $\chi_{\text{top}}(K_3) = 24$ and $\chi(\mathcal{O}_S) = 2$). The integer $24 = \chi_{\text{top}}(K_3)$ enters the Borchers exponent $f(0, 0)/2 = 5$ via $f(0, 0) = 2(\chi_{\text{top}}(K_3))/24 \cdot \dim_{\mathbb{C}}(\tilde{H}(K_3, \mathbb{C})/(\mathbb{R} \cdot 1 \oplus \mathbb{R} \cdot \omega_K)) = 10$, equivalently $\phi_{0,1}(\tau, 0) = 24 \eta(\tau)^{-2} \theta_1 \theta_2 \theta_3 \theta_4 = 12 + \dots$ at $z = 1/2$ (computed/ β , [?EZ:1]). The integer $\chi(\mathcal{O}_S) = 2$ enters the categorical Calabi–Yau dimension and the holomorphic part of the Mukai pairing. The two integers do not collapse: the κ -tuple of [?calabi-yau-quantum-groups] records $\kappa_{\text{fiber}} = 24 = \chi_{\text{top}}(K_3)$, not $2 = \chi(\mathcal{O}_S)$, and the universal Borchers identity $\kappa_{\text{BKM}}(\Phi_N) = c_N(0)/2$ at $\Phi_1 = \Delta_5$ gives $c_1(0) = 10$ and $\kappa_{\text{BKM}} = 5$, while $\kappa_{\text{ch}}^{\text{Heis}} + \chi(\mathcal{O}_E) = 3 + 0 = 3$: the additive collapse $\kappa_{\text{BKM}} = \kappa_{\text{ch}}^{\text{Heis}} + \chi(\mathcal{O}_{\text{fiber}})$ disagrees with the universal identity at $N = 1$ (proved/ α , [?calabi-yau-quantum-groups] counterexample). The level $\kappa_{\text{BKM}} = 5$ is the cusp-form weight; the manuscript writes the protected scalar shadow at level $n = \chi(\mathcal{O}_Y)$, with the threefold Y specialised to the $K_3 \times E$ fibre over the closed cobordism. The $K_3 \times E$ target is a Calabi–Yau threefold; the manuscript reads X as a threefold throughout.

1.2 Mukai lattice on the K_3 factor

Definition 1.3 (Mukai lattice and Mukai pairing). Let S be a smooth complex K_3 surface. The *Mukai lattice* is the total even cohomology

$$\Lambda_S := \tilde{H}(S, \mathbb{Z}) = H^0(S, \mathbb{Z}) \oplus H^2(S, \mathbb{Z}) \oplus H^4(S, \mathbb{Z}),$$

identified with the K-theoretic charge lattice $K_0(D^b(\text{Coh } S))_{\text{tor. free}}$ via the Mukai vector

$$v(\mathcal{F}) := \text{ch}(\mathcal{F}) \sqrt{\text{Td}(S)} = (r(\mathcal{F}), c_1(\mathcal{F}), r(\mathcal{F}) + \text{ch}_2(\mathcal{F})),$$

where r is the rank, c_1 is the first Chern class, and the half-shift r in the four-form term comes from $\text{Td}(S) = 1 + 2[\text{pt}]$ (proved/ β , [?Mukai1984, ?Mukai1987]). The *Mukai pairing* is

$$\langle (r, c, s), (r', c', s') \rangle_{\text{Muk}} := c \cdot c' - r s' - r' s,$$

where $c \cdot c' \in \mathbb{Z}$ is the cup product on $H^2(S, \mathbb{Z})$. It is even, unimodular, of signature $(4, 20)$, and rank 24:

$$\Lambda_S \cong II_{4,20} \cong U^{\oplus 4} \oplus E_8(-1)^{\oplus 2},$$

where U denotes the rank-two hyperbolic plane $U \cong \mathbb{Z}e \oplus \mathbb{Z}f$ with $e^2 = f^2 = 0$, $e \cdot f = 1$, and $E_8(-1)$ denotes the negative-definite E_8 root lattice (proved/ β , [?Mukai1987, ?NIKULIN]).

Remark 1.4 (Sign convention). The signature $(4, 20)$ uses the convention that $H^0 \oplus H^4$ contributes a hyperbolic plane U via $(1, 0, 0)$ and $(0, 0, 1)$ and that $H^2(S, \mathbb{Z}) \cong II_{3,19} \cong U^{\oplus 3} \oplus E_8(-1)^{\oplus 2}$, which has positive part of dimension 3. Together, $\Lambda_S = U \oplus II_{3,19} \cong U^{\oplus 4} \oplus E_8(-1)^{\oplus 2}$ with positive part of dimension 4 (proved/ β , [?Mukai1987]). The signature is $(4, 20)$, not $(20, 4)$: the positive direction is the smaller side. Throughout the manuscript a reflection r_v across $v \in \Lambda_S$ acts by $r_v(x) = x - 2(x, v)v/(v, v)$, taking the $\mathbb{R}$ -linear extension when $(v, v) \neq 0$, without implicit sign reversal.

1.3 Künneth and the formal even Mukai sector on $X = S \times E$

Definition 1.5 (Formal even Mukai sector on $X = S \times E$). Pick a basepoint $o_E \in E$ and the standard generators $1_E \in H^0(E, \mathbb{Z})$ and $\omega_E \in H^2(E, \mathbb{Z})$ with $\int_E \omega_E = 1$. Every even cohomology class on $X = S \times E$ decomposes as

$$v_X = P \otimes 1_E + Q \otimes \omega_E, \quad P, Q \in \Lambda_S$$

(proved/ β , Künneth on simply connected $\tilde{H}$ tensor E -cohomology). The pair $(Q, P) \in \Lambda_S \oplus \Lambda_S$ is the *formal even charge*. The component $P \otimes 1_E$ is invariant under translation along E , while $Q \otimes \omega_E$ records the E -degree.

The *formal full-Mukai dyonic lattice* is the additive abelian group

$$\Gamma_X^{\text{form}} := \Lambda_S \oplus \Lambda_S, \quad c = (Q, P),$$

with electric component Q and magnetic component P . In this chapter the legacy notation Γ_X^{phys} is retained as a mnemonic only and equals Γ_X^{form} :

$$\Gamma_X^{\text{phys}} := \Gamma_X^{\text{form}}.$$

This is not the full four-dimensional $\mathcal{N} = 4$ electric–magnetic charge lattice and not compact Hall support. It is the D6–D2–Do/OP even branch on which the Mukai Gram map is tested before imposing algebraicity, effectivity, stability, or non-emptiness of compact Hall moduli.

Definition 1.6 (Algebraic/effective Hall support). Fix a polarised K_3 surface (S, H) , a chosen Mukai vector lattice $\Lambda_{S, \text{alg}} \subset \Lambda_S$ (the image of the algebraic K-class lattice generated by sheaves whose Mukai vector lies in $H^0 \oplus \text{NS}(S) \oplus H^4$), and a stability condition σ on $D^b(\text{Coh } S \times E)$ (Bridgeland-stable; folklore/ β , [?KSCoHA]). Set

$$\Gamma_X^{\text{alg}} := \Lambda_{S, \text{alg}} \oplus \Lambda_{S, \text{alg}}, \quad \Gamma_{X, \sigma}^{\text{eff}} \subset \Gamma_X^{\text{alg}} \subset \Gamma_X^{\text{form}},$$

where the σ -effective semigroup $\Gamma_{X, \sigma}^{\text{eff}}$ is the additive sub-semigroup of charges supported by σ -semistable objects of nonzero Mukai vector. Membership in $\Gamma_{X, \sigma}^{\text{eff}}$ is not implied by membership in Γ_X^{alg} or by formal lattice arithmetic (folklore/ β).

Remark 1.7 (Three-tier inclusion). The hierarchy $\Gamma_{X, \sigma}^{\text{eff}} \subset \Gamma_X^{\text{alg}} \subset \Gamma_X^{\text{form}}$ distinguishes three discipline registers. Γ_X^{form} is the formal additive group on which Π_X is defined and the polarization identity $B = -\partial \Pi_X$ holds. Γ_X^{alg} is the K-theoretic algebraic sublattice; the rank-one extension closure of Γ_X^{alg} inside Γ_X^{form} records the Mukai-effective branch (computed/ β , [?Mukai1987, ?KSCoHA]). $\Gamma_{X, \sigma}^{\text{eff}}$ is the σ -stable Hall support; it depends on σ and is not fibre-functorial in the K_3 type. The sectorial Hall truncation of §4 works at the second level after imposing finite Harder–Narasimhan height bounds.

LEMMA 1.8 (*Two orthogonal hyperbolic planes inside Λ_S*). The Mukai lattice contains two orthogonal hyperbolic planes:

$$U \oplus U \hookrightarrow \Lambda_S, \quad U_1 = \mathbb{Z}e_1 \oplus \mathbb{Z}f_1, \quad U_2 = \mathbb{Z}e_2 \oplus \mathbb{Z}f_2,$$

with $e_i^2 = f_i^2 = 0$, $e_i \cdot f_i = 1$, and $U_1 \perp U_2$ inside the Mukai pairing (proved/ α , [NIKULIN]).

Proof. Direct from $\Lambda_S \cong U^{\oplus 4} \oplus E_8(-1)^{\oplus 2}$. □

2 QUADRATIC GRAM MAP AND LATTICE POLARIZATION

2.1 Igusa variables and the Gram-index lattice

For a Siegel period

$$Z = \begin{pmatrix} z_1 & z_2 \\ z_2 & z_3 \end{pmatrix} \in \mathcal{H}_2, \quad q = e^{2\pi i z_1}, \quad r = e^{2\pi i z_2}, \quad s = e^{2\pi i z_3},$$

and

$$T(n, l, m) = \begin{pmatrix} n & l/2 \\ l/2 & m \end{pmatrix}, \quad n, l, m \in \mathbb{Z},$$

the trace pairing

$$(T(n, l, m), Z)_{\text{tr}} := \text{tr}(T(n, l, m) Z) = nz_1 + lz_2 + mz_3$$

gives the Borchers monomial $x^{(n, l, m)} = q^n r^l s^m$.

Definition 2.1 (*Gram-index lattice and effective semigroup*). The Gram-index lattice is

$$\Gamma_{\text{gram}} := \mathbb{Z}^3, \quad \gamma = (n, l, m).$$

The *cuspidal semigroup* at the standard cusp $\mathfrak{c}_\infty$ with chamber convention $0 < |s| \ll |q| \ll |r|^{-1} \ll 1$ is

$$\Gamma_{\text{eff}} = \{(n, l, m) : m > 0, n \geq 0, l \in \mathbb{Z}\} \cup \{(n, l, 0) : n > 0, l \in \mathbb{Z}\} \cup \{(0, l, 0) : l < 0\}.$$

The *Jacobi-active sublocus* $\Gamma_{\text{act}} \subset \Gamma_{\text{eff}}$ is the locus of nonvanishing $f(nm, l)$ in the weak Jacobi form $\phi_{0,1}$ (computed/ β , [EZ:1]).

PROPOSITION 2.2 (*Gram matrix convention for protected charges*). For a formal even charge $c = (Q, P) \in \Gamma_X^{\text{form}}$ in the unimodular lattice Λ_S , set

$$n = \frac{(Q, Q)}{2}, \quad l = (Q, P), \quad m = \frac{(P, P)}{2}.$$

Then $n, l, m \in \mathbb{Z}$ (since Λ_S is even unimodular, proved/ γ),

$$\begin{pmatrix} (Q, Q) & (Q, P) \\ (P, Q) & (P, P) \end{pmatrix} = 2T(n, l, m), \quad \det(2T(n, l, m)) = 4nm - l^2,$$

and $x^{(n, l, m)} = q^n r^l s^m$ is the Fourier character of $(T(n, l, m), Z)_{\text{tr}}$.

Proof. The Mukai pairing is integer-valued and even on $\Lambda_S \oplus \Lambda_S$ by Definition 1.21. The matrix identity is the direct expansion

$$\begin{pmatrix} (Q, Q) & (Q, P) \\ (P, Q) & (P, P) \end{pmatrix} = \begin{pmatrix} 2n & l \\ l & 2m \end{pmatrix} = {}_2T(n, l, m).$$

The determinant follows. The Fourier character is $\exp(2\pi i \operatorname{tr}(TZ)) = q^n r^l s^m$. $\square$

2.2 The Gram map and its bilinear cocycle

Definition 2.3 (Quadratic Gram map and bilinear cocycle). The *Gram map* is the quadratic projection

$$\Pi_X : \Gamma_X^{\text{form}} \longrightarrow \Gamma_{\text{gram}}, \quad \Pi_X(Q, P) = \left(\frac{(Q, Q)}{2}, (Q, P), \frac{(P, P)}{2} \right).$$

The *symmetric bilinear cocycle* B associated to Π_X is

$$B : \Gamma_X^{\text{form}} \times \Gamma_X^{\text{form}} \longrightarrow \Gamma_{\text{gram}}, \quad B((Q, P), (Q', P')) = ((Q, Q'), (Q, P') + (Q', P), (P, P')).$$

LEMMA 2.4 (Gram additivity defect). For $c, c' \in \Gamma_X^{\text{form}}$,

$$\Pi_X(c + c') = \Pi_X(c) + \Pi_X(c') + B(c, c').$$

The defect B is symmetric, $\mathbb{Z}$ -bilinear, integer-valued, and satisfies polarization: $B(c, c) = {}_2\Pi_X(c)$. As an ordinary additive cochain identity,

$$B = -\partial\Pi_X, \quad (\partial q)(c, c') = q(c) + q(c') - q(c + c').$$

(proved/ γ .)

Proof. Write $c = (Q, P), c' = (Q', P')$; then

$$\begin{aligned} \frac{(Q + Q')^2}{2} &= \frac{Q^2}{2} + \frac{Q'^2}{2} + Q \cdot Q', \\ (Q + Q') \cdot (P + P') &= Q \cdot P + Q' \cdot P' + Q \cdot P' + Q' \cdot P, \\ \frac{(P + P')^2}{2} &= \frac{P^2}{2} + \frac{P'^2}{2} + P \cdot P'. \end{aligned}$$

The three cross terms are the three components of $B(c, c')$. Symmetry, bilinearity, and integrality follow from the corresponding properties of the Mukai pairing. Polarization is the case $c' = c$. The cochain identity rewrites the additivity defect. $\square$

LEMMA 2.5 (Bilinear cocycle identity). B is normalised, $B(c, o) = B(o, c) = o$, and the additive coboundary

$$(\partial B)(a, b, c) := B(b, c) - B(a + b, c) + B(a, b + c) - B(a, b)$$

vanishes identically on Γ_X^{form} (proved/ γ).

Proof. By bilinearity in each slot,

$$B(a + b, c) = B(a, c) + B(b, c), \quad B(a, b + c) = B(a, b) + B(a, c).$$

Substituting,

$$(\partial B)(a, b, c) = B(b, c) - B(a, c) - B(b, c) + B(a, b) + B(a, c) - B(a, b) = 0.$$

The polarization identity $B(c, c) = 2 \Pi_X(c)$ is the only place where B takes a non-vanishing value on the diagonal; this is structural, not a Bott-element framing. $\square$

Remark 2.6 (Polarization, not Bott element). The identity $B(c, c) = 2 \Pi_X(c)$ is the polarization identity for the symmetric bilinear B associated to the quadratic Π_X . It is forced by the homogeneous-of-degree-two structure of the Gram map. No external Bott-element framing or characteristic-class coincidence is involved. The full structural content is that Π_X is a homogeneous quadratic on the additive group Γ_X^{form} valued in Γ_{gram} , with associated symmetric bilinear B given by the Mukai pairing of components on each direction.

2.3 Formal lift of every Gram triple

LEMMA 2.7 (*Formal primitive torsion-one lift of every Gram triple*). Suppose Λ_S contains two orthogonal hyperbolic planes $U_1 \oplus U_2$ with bases (e_1, f_1, e_2, f_2) as in Lemma 1.29. Then every Gram triple $(n, l, m) \in \mathbb{Z}^3$ admits a primitive saturated formal full-Mukai lift $c = (Q, P) \in \Gamma_X^{\text{form}}$ with $\Pi_X(Q, P) = (n, l, m)$. One may take

$$Q = e_1 + n f_1, \quad P = l f_1 + e_2 + m f_2.$$

For this lift, the wedge torsion invariant

$$I(Q, P) := \gcd\{\text{coordinates of } Q \wedge P \text{ in a } \mathbb{Z}\text{-basis of } \wedge^2 \Lambda_S\}$$

equals 1. (proved/ β .)

Proof. The hyperbolic-plane pairings give

$$Q^2 = 2n, \quad P^2 = 2m, \quad Q \cdot P = l,$$

hence $\Pi_X(Q, P) = (n, l, m)$. The submodule $\mathbb{Z}Q + \mathbb{Z}P \subset \Lambda_S$ contains the unimodular minor $\det \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = 1$ in the rows indexed by e_1, e_2 of the coefficient matrix in (e_1, f_1, e_2, f_2) , so the lift is primitive saturated. The wedge expansion

$$Q \wedge P = l e_1 \wedge f_1 + e_1 \wedge e_2 + m e_1 \wedge f_2 + n f_1 \wedge e_2 + nm f_1 \wedge f_2$$

contains the basis element $e_1 \wedge e_2$ with coefficient 1, so the gcd is 1 and $I(Q, P) = 1$. $\square$

Remark 2.8 (Pointwise existence is not Hall support). Lemma 2.7 produces a formal charge plane realising every prescribed Igusa degree. It does not prove algebraicity of (Q, P) , effectivity, non-empty compact Hall moduli, duality-orbit descent, source-index descent, or any chain-level Hall strictification. The compact Hall obstruction is functorial compatibility of pointwise existence with Hall addition and the primitive bracket; only the latter is the working content of Chapter ??.

3 NORMAL-ORDERED GRAM EXTENSION

3.1 Central extension and the additive normal-ordered Gram homomorphism

Definition 3.1 (Normal-ordered extension and Gram homomorphism). Set

$$\widehat{\Gamma}_X := \Gamma_X^{\text{form}} \oplus \Gamma_{\text{gram}},$$

with operation

$$(c, T) \star (c', T') := (c + c', T + T' + B(c, c')).$$

Define the *normal-ordered Gram map*

$$\overline{\Pi}_X : \widehat{\Gamma}_X \longrightarrow \Gamma_{\text{gram}}, \quad \overline{\Pi}_X(c, T) = \Pi_X(c) - T.$$

The *raw placement* and the *split section* are

$$i_o : \Gamma_X^{\text{form}} \longrightarrow \widehat{\Gamma}_X, \quad i_o(c) = (c, o), \quad s : \Gamma_X^{\text{form}} \longrightarrow \widehat{\Gamma}_X, \quad s(c) = (c, \Pi_X(c)).$$

THEOREM 3.2 (*Group law of the normal-ordered extension; additivity of $\overline{\Pi}_X$*). Let $\widehat{\Gamma}_X, \star, \overline{\Pi}_X, i_o, s$ be as in Definition 3.1. Then:

- (i) $(\widehat{\Gamma}_X, \star)$ is an abelian group with identity (o, o) and inverse $(c, T)^{-1} = (-c, -T + B(c, c)) = (-c, -T + 2\Pi_X(c))$.
- (ii) The sequence

$$o \rightarrow \Gamma_{\text{gram}} \xrightarrow{T \mapsto (o, T)} \widehat{\Gamma}_X \xrightarrow{\text{pr}_1} \Gamma_X^{\text{form}} \rightarrow o$$

is a central extension of abelian groups with cocycle B .

- (iii) $\overline{\Pi}_X$ is a group homomorphism from $(\widehat{\Gamma}_X, \star)$ to the additive group Γ_{gram} .
- (iv) The split section $s(c) = (c, \Pi_X(c))$ is additive, and

$$\overline{\Pi}_X(i_o(c)) = \Pi_X(c), \quad \overline{\Pi}_X(s(c)) = o.$$

- (v) The change of variables $(c, T) \mapsto (c, T - \Pi_X(c))$ carries $\star$ to componentwise addition; thus $\widehat{\Gamma}_X \cong \Gamma_X^{\text{form}} \oplus \Gamma_{\text{gram}}$ as abstract abelian groups, and $[B] = o$ in additive cohomology.

(proved/ δ .)

Proof. (i) Associativity follows from the cocycle identity $\delta B = o$ (Lemma 2.5); the associator $B(a, b) + B(a + b, c) - B(a, b + c) - B(b, c)$ vanishes by bilinearity. Commutativity is symmetry of B . The unit calculation $(c, T) \star (o, o) = (c, T + B(c, o)) = (c, T)$ and inverse $(c, T) \star (-c, -T + B(c, c)) = (o, B(c, c) - B(c, c)) = (o, o)$ (using $B(c, -c) = -B(c, c)$ by bilinearity) close the abelian-group structure.

(ii) Definition.

(iii) Apply Lemma 2.4:

$$\overline{\Pi}_X((c, T) \star (c', T')) = \Pi_X(c + c') - T - T' - B(c, c') = \Pi_X(c) + \Pi_X(c') - T - T' = \overline{\Pi}_X(c, T) + \overline{\Pi}_X(c', T').$$

(iv) For the section,

$$s(c) \star s(c') = (c + c', \Pi_X(c) + \Pi_X(c') + B(c, c')) = (c + c', \Pi_X(c + c')) = s(c + c').$$

The identities $\overline{\Pi}_X(i_o(c)) = \Pi_X(c) - o = \Pi_X(c)$ and $\overline{\Pi}_X(s(c)) = \Pi_X(c) - \Pi_X(c) = o$ are immediate.

(v) The map $(c, T) \mapsto (c, T - \Pi_X(c))$ is a bijection with inverse $(c, T) \mapsto (c, T + \Pi_X(c))$. Direct calculation gives

$$(c, T - \Pi_X(c)) + (c', T' - \Pi_X(c')) = (c + c', T + T' - \Pi_X(c) - \Pi_X(c')) = (c + c', (T + T' + B(c, c')) - \Pi_X(c + c')),$$

matching the change of variables applied to $(c, T) \star (c', T')$. $\square$

3.2 The fixed-lift raw-pushforward obstruction theorem

Definition 3.3 (Raw homogeneous and fixed-lift Π_X -pushforwards). Let $P = \bigoplus_{c \in \Gamma_X^{\text{form}}} P_c$ be a Γ_X^{form} -graded primitive object. The *raw homogeneous Π_X -pushforward* is

$$(\Pi_{X,*}^{\text{raw}} P)_\gamma := \bigoplus_{\Pi_X(c)=\gamma} P_c.$$

The *strict fixed-lift raw Π_X -pushforward* is the narrower operation obtained by choosing one lift $L(\gamma) \in \Gamma_X^{\text{form}}$ for every retained γ and setting

$$(\Pi_{X,*}^{\text{raw},L} P)_\gamma := P_{L(\gamma)}.$$

THEOREM 3.4 (*Raw homogeneous fixed-lift Π_X -pushforward cannot realise the type-II real-root strings*). Let $P = \bigoplus_{c \in \Gamma_X^{\text{form}}} P_c$ be a Γ_X^{form} -graded Lie superalgebra with $[P_c, P_{c'}] \subset P_{c+c'}$. Suppose a strict fixed-lift raw pushforward of P along the quadratic map Π_X were a Γ_{gram} -graded Lie superalgebra structure on the displayed bracket channels of chosen charge lifts. Then every nonzero bracket channel $[P_c, P_{c'}]$ contributing to that strict raw pushforward must satisfy $B(c, c') = o$.

For the type-II real simple roots $(\delta_1, \delta_2, \delta_3)$ of the Gritsenko–Nikulin algebra $\mathfrak{g}_{\Delta_3}$ with Cartan matrix

$$A_{\text{II}} = \begin{pmatrix} 2 & -2 & -2 \\ -2 & 2 & -2 \\ -2 & -2 & 2 \end{pmatrix},$$

this condition is incompatible with the nonzero second real-root strings $2\delta_i + \delta_j, i \neq j$. The strict fixed-lift raw Π_X -descent therefore cannot carry these brackets; they require the normal-ordered extension $\widehat{\Gamma}_X$ and the additive homomorphism $\overline{\Pi}_X$. (proved/ δ , [?GN].)

Proof. If the raw pushforward is Γ_{gram} -graded, a nonzero bracket of classes of raw Gram degrees $\Pi_X(c)$ and $\Pi_X(c')$ must land in raw Gram degree $\Pi_X(c) + \Pi_X(c')$. But the upstairs bracket lands in physical degree $c + c'$, whose raw Gram shadow is

$$\Pi_X(c + c') = \Pi_X(c) + \Pi_X(c') + B(c, c').$$

Hence $B(c, c') = o$ on every nonzero bracket channel.

Take physical lifts c_i, c_j of distinct type-II real simple root degrees γ_i, γ_j , $\Pi_X(c_i) = \gamma_i$ with $\gamma_i \neq o$. The Gritsenko–Nikulin Cartan matrix A_{II} has off-diagonal entries -2 , so distinct simple roots satisfy $\langle \delta_i, \delta_j \rangle = -2$, and the real-root string through δ_j in the δ_i -direction includes the levels $\delta_i + \delta_j$ and $2\delta_i + \delta_j$ (proved/ δ , [?GN, §2–3]). Both correspond to nonzero brackets, hence raw pushforward forces

$$B(c_i, c_j) = o, \quad B(c_i, c_i + c_j) = o.$$

Bilinearity and polarization give

$$B(c_i, c_i + c_j) = B(c_i, c_i) + B(c_i, c_j) = 2\Pi_X(c_i) + 0 = 2\gamma_i,$$

which is nonzero in the torsion-free lattice Γ_{gram} . Contradiction. $\square$

Remark 3.5 (Scope of the no-go). Theorem 3.4 excludes the strict *homogeneous fixed-lift* raw pushforward. It does not exclude isolated B -isotropic bracket channels and does not decide the mixed-lift fibre-summed construction in which several lifts of a single Gram degree are summed before bracketing; the latter remains an open mixed-lift question. The normal-ordered extension removes exactly the displayed obstruction because $\overline{\Pi}_X$ is additive on $\widehat{\Gamma}_X$. Chain-level Hall strictification ($\mathcal{A}_\infty$ -compatibility, cyclic equivariance, Frobenius, multiplicative orientation) is a separate input system, named in §6 and developed in Chapter ??.

Remark 3.6 (Quadratic obstruction; no linear trivialisation). Suppose $B = \delta L$ for some additive linear $L : \Gamma_X^{\text{form}} \rightarrow \Gamma_{\text{gram}}$. Setting $c' = c$, $B(c, c) = 2L(c) - L(2c)$; additivity gives $L(2c) = 2L(c)$, so $B(c, c) = 0$, contradicting $B(c, c) = 2\Pi_X(c) \neq 0$ for any c with $(Q, Q), (Q, P), (P, P)$ not all zero. Hence no additive linear trivialisation exists. The quadratic primitive Π_X is necessary; trivialising the additive cocycle requires a quadratic cochain, not a linear one (proved/ γ).

3.3 Normal-ordering flag formula

LEMMA 3.7 (*Multiplication of raw placements*). For $c_1, \dots, c_k \in \Gamma_X^{\text{form}}$,

$$i_o(c_1) \star \cdots \star i_o(c_k) = \left(\sum_i c_i, \sum_{i < j} B(c_i, c_j) \right),$$

and consequently

$$\overline{\Pi}_X(i_o(c_1) \star \cdots \star i_o(c_k)) = \sum_i \Pi_X(c_i).$$

(proved/ δ .)

Proof. Induct on k . The case $k = 2$ is the definition of $\star$. At step $k \rightarrow k + 1$,

$$B\left(\sum_{i \leq k} c_i, c_{k+1}\right) = \sum_{i \leq k} B(c_i, c_{k+1})$$

by bilinearity. Applying $\overline{\Pi}_X$ subtracts the accumulated pairwise B -term from $\Pi_X(\sum_i c_i) = \sum_i \Pi_X(c_i) + \sum_{i < j} B(c_i, c_j)$. $\square$

Remark 3.8 (Normal-ordered primitive recognition target). The interface with the BKM denominator algebra is the corrected route

$$\Gamma_X^{\text{form}} \hookrightarrow \widehat{\Gamma}_X \xrightarrow{\overline{\Pi}_X} \Gamma_{\text{gram}} \xrightarrow{\alpha} \Lambda_{II}^{2,1},$$

where $\alpha(n, l, m) = 2nf_2 - lf_3 + 2mf_{-2}$ is the type-IV embedding into the BKM root lattice (proved/ δ , [?GN]). At chain level this becomes the descent $\overline{\Pi}_{X,*}^\ominus$ of Chapter ??. The primitive recognition problem is to construct a protected primitive Hall algebra whose normal-ordered descended bracket is the Borcherds–Kac bracket of $\mathfrak{g}_{\Delta_5}$. Raw Gram pushforward is not an admissible recognition target.

4 THE HYBRID $\text{Ran}^{\text{hyb}}(E)$ FACTORIZATION CARRIER

The K_3 direction is integrated. What remains is a chiral object on the elliptic factor E . The naive carrier is the Ran space $\text{Ran}(E)$ of finite subsets, but a one-dimensional subscheme with positive elliptic degree projects onto all of E . The hybrid carrier $\text{Ran}^{\text{hyb}}(E)$ is the disjoint union of the projection-finite local stratum $\text{Ran}(E)_{b=0}$ and the wrapped stratum $\text{Ran}^{\text{wr}}(E)_{b>0}$, with mixed local/wrapped correspondences supplied as additional finite-stage data.

4.1 Projection-locality obstruction

LEMMA 4.1 (*Projection-locality obstruction*). Let $C \subset X = S \times E$ be a one-dimensional closed subscheme (reduced and irreducible, for definiteness) with

$$\deg_E(C) := \deg(p_E|_C: C \rightarrow E) > 0.$$

Then $p_E(C) = E$, hence $p_E(C)$ is not a finite subset of E and C is not local on $\text{Ran}(E)$. (proved/ ϵ .)

Proof. The pushforward $p_{E,*}[C]$ is a positive multiple of the fundamental class $[E]$, so the image $p_E(C)$ is a closed one-dimensional subset of the irreducible curve E . An irreducible curve has only itself as a one-dimensional closed subset, so $p_E(C) = E$. $\square$

Remark 4.2 (*Geometric versus Borcherds elliptic degree*). The lemma fixes the necessity of a wrapped stratum. The geometric elliptic-degree map b_R^{geom} , on a finite Hall stage R , records the integer $\deg_E$ of a chosen retained class. The trace exponent $m_R := \text{pr}_3 \overline{\Pi}_X$, on the same stage, records the third coordinate of the normal-ordered Gram pushforward. The two are not equal as functions on the formal lattice; the Borcherds exponent m records the magnetic Mukai pairing $(P, P)/2$, which sees Λ_S -square data, while b_R^{geom} records the E -degree of the geometric support, which sees $H^0(S) \oplus H^4(S)$ data through r and ch_2 of the K_3 factor. The wrapped stratum carries $b_R^{\text{geom}} > 0$ branes; the local stratum carries $b_R^{\text{geom}} = 0$ branes whose magnetic component lies in the projection-finite kernel. This is the $\text{Ran}(E)$ -support-locality obstruction in the K_3 -integrated chiral shadow.

4.2 The hybrid carrier

Definition 4.3 (*Hybrid Ran carrier*). The hybrid Ran carrier on E is the disjoint union

$$\text{Ran}^{\text{hyb}}(E) := \text{Ran}(E)_{b=0} \sqcup \text{Ran}^{\text{wr}}(E)_{b>0},$$

with the projection-finite local stratum $\text{Ran}(E)_{b=0}$ and the wrapped stratum $\text{Ran}^{\text{wr}}(E)_{b>0}$ defined as follows.

- (i) $\text{Ran}(E)_{b=0}$ is the standard Ran space of finite (unordered) subsets $T = \{x_1, \dots, x_k\} \subset E$ ([?BD, ?FrancisGaiatsgoryCKD]); its functor of points sends a test scheme S' to the set of finite closed subschemes $T' \subset E \times S'$ whose projection to S' is finite étale of constant fibre size $\leq R$ for some chosen finite Hall height R . This is the $b_R^{\text{geom}} = 0$ stratum: branes whose K_3 factor has zero elliptic degree.
- (ii) $\text{Ran}^{\text{wr}}(E)_{b>0}$ is the *wrapped prequotient stratum*: a colim of pre-quotient stacks $\mathfrak{B}_{R,b}$ parameterising rigidified compactly supported B-branes $\mathcal{F} \in D^b(\text{Coh } X)$ of bounded retained Liu–Hilbert class $\Xi = (\gamma, [a, b], (P_i), N)$ with positive elliptic degree $b = \deg_E(\text{ch}_2(\mathcal{F})) > 0$, modulo E -translation up to the chosen anchor (§4.4).

The two strata are exchanged only by the mixed local/wrapped correspondences of §4.3; they are not identified.

Remark 4.4 (Two strata, two registers). On the $b = 0$ stratum the standard $\text{Ran}/\text{factorization}$ formalism applies and produces a chiral algebra in $\text{Fact}(\text{Ran}(E))$ (folklore/ ϵ , [?BD, ?FrancisGaitsgoryCKD]). On the $b > 0$ stratum the support of $\text{Supp } \mathcal{F}$ projects onto E , so the brane is global; its data live on the wrapped prequotient and after E -quotient become E -equivariant data on the elliptic curve. The hybrid carrier is the smallest carrier reflecting both strata; it is not a single Ran space.

4.3 Mixed local/wrapped correspondences

Definition 4.5 (Mixed local/wrapped correspondences (finite Hall stage R)). Fix a finite Hall stage R and retained Liu–Hilbert classes Ξ_A, Ξ_B, Ξ_C with $\Xi_C = \Xi_A + \Xi_B$ in the closure rules of [?KSCoHA]. The retained extension correspondence

$$\mathfrak{E}_{\Xi_A, \Xi_B; \Xi_C}^{\text{ret}} \xrightarrow{p} \mathfrak{M}_{\Xi_A}^{\text{ss}} \times \mathfrak{M}_{\Xi_B}^{\text{ss}}, \quad \mathfrak{E}_{\Xi_A, \Xi_B; \Xi_C}^{\text{ret}} \xrightarrow{q} \mathfrak{M}_{\Xi_C}^{\text{ss}}$$

is the closed derived flag-Quot/filtration stack over $\mathfrak{M}_{\Xi_C}^{\text{ss}}$, with q proper or Coh^{prop} -admissible in the compactified model (folklore/ ϵ). Three colours of correspondence appear, indexed by the location of Ξ_A, Ξ_B in $\text{Ran}^{\text{hyb}}(E)$:

(LL) *local-local*: $b_A = b_B = 0$;

(LW) *ordered local/wrapped*: $b_A = 0, b_B > 0$;

(WW) *wrapped-wrapped*: $b_A, b_B > 0$.

The arity-three shadow is the eight-word two-step binary flag atlas

$$\{L, W\}^3 = \{LLL, LLW, LWL, LWW, WLL, WLW, WWL, WWW\},$$

which records binary associativity only; higher coloured tree configurations live on a separate coloured tree category $\text{Tree}_R^{\text{col}}$, supplied as additional input (folklore/ ϵ , [?calabi-yau-quantum-groups]).

Remark 4.6 (Eight binary words versus colored trees). The eight binary words of Definition 4.5 exhaust the arity-three two-step compositions $\{L, W\}^3$, but they do not catalogue higher coloured tree configurations, units, symmetry/order conventions, refinement maps, descent, or overlaps. Those data are additional finite Hall input named at §6; asserting them prematurely confuses binary associativity with the full factorization-coherence structure ([?calabi-yau-quantum-groups], correspondence-programme remark).

4.4 Wrapped anchor and translation linearization

Definition 4.7 (Determinant anchor on the wrapped stratum). For a retained E -stable family $\mathcal{F}$ on the wrapped stratum, the *determinant anchor line bundle* on E is

$$\lambda(\mathcal{F}) := \det R p_{E,*} \mathcal{F} \otimes \mathcal{O}_E(-\chi(\mathcal{F}) \circ_E).$$

This is a degree-zero line bundle on E , hence a point of $\text{Pic}^0(E) \cong E$. The translation law

$$\lambda(t \cdot \mathcal{F}) = \lambda(\mathcal{F}) + \chi(\mathcal{F}) t, \quad t \in E,$$

holds in $\text{Pic}^0(E)$ (proved/ ϵ , [?Mukai1987]).

LEMMA 4.8 (*Connected E -descent: separated cohomological classes*). For the connected component E° of the identity in the E -translation action,

$$H^*(BE; \mathbb{F}_2) = \mathbb{F}_2[u_1, u_2], \quad |u_i| = 2.$$

Hence

$$H^1(BE; \mathbb{F}_2) = 0, \quad H^2(BE; \mathbb{F}_2) = \mathbb{F}_2 u_1 \oplus \mathbb{F}_2 u_2.$$

There is no degree-one connected character; there is a degree-two connected gerbe class

$$\alpha^{E, \text{free}} = a_1 u_1 + a_2 u_2 \in H^2(BE; \mathbb{F}_2).$$

(proved/ ϵ , [NIKULIN].)

Proof. $E^{\text{top}} \simeq T^2$ has classifying space $BE \simeq BT^2$ and $H^*(BT^2; \mathbb{F}_2) = \mathbb{F}_2[u_1, u_2]$ with both u_i of degree 2 (mod-2 Chern classes of the universal line bundles). H^1 vanishes; H^2 is the displayed two-dimensional $\mathbb{F}_2$ vector space. Vanishing of the connected gerbe class is the equation $a_1 = a_2 = 0$ on the actual Borel/edge class of the equivariant reduced determinant complex, not a consequence of $H^1(BE) = 0$ or ordinary translation invariance. $\square$

LEMMA 4.9 (*Finite stabilizer $E[N]$ -descent: gerbe and linearization classes*). For the order-two finite subgroup $E[2] \cong (\mathbb{Z}/2)^2$,

$$H^*(BE[2]; \mathbb{F}_2) = \mathbb{F}_2[x_1, x_2], \quad |x_i| = 1.$$

The degree-two gerbe class is

$$\beta^{E, E[2]} = b_{20} x_1^2 + b_{11} x_1 x_2 + b_{02} x_2^2, \quad (b_{20}, b_{11}, b_{02}) = (r_1, r_1 + r_2 + r_3, r_2),$$

detected on cyclic subgroup restrictions by joint reduction; the $r_i \in \mathbb{F}_2$ are the three nonzero point-stabilizer parities. After $\beta^{E, E[2]} = 0$ (i.e. $r_1 = r_2 = r_3 = 0$), a separate degree-one linearization class remains:

$$\lambda^{E, E[2]} = \lambda_1 x_1 + \lambda_2 x_2 \in H^1(BE[2]; \mathbb{F}_2), \quad \rho_1 = \lambda_1, \rho_2 = \lambda_2, \rho_3 = \lambda_1 + \lambda_2.$$

The r_i are gerbe bits in H^2 ; the ρ_i are linearization bits in H^1 ; the two live in distinct cohomological degrees and remain independent. For even $N \geq 4$ with $2^a \parallel N$, $a \geq 2$,

$$H^*(B(\mathbb{Z}/2^a)^2; \mathbb{F}_2) = \Lambda(x_1, x_2) \otimes \mathbb{F}_2[y_1, y_2],$$

$$\beta^{E, N} = A_1^{(N)} y_1 + A_{12}^{(N)} x_1 x_2 + A_2^{(N)} y_2, \quad \lambda^{E, N} = \lambda_1^{(N)} x_1 + \lambda_2^{(N)} x_2.$$

The mixed term $A_{12}^{(N)}$ is invisible to cyclic order-two restrictions; even $N \geq 4$ tests must be checked on the full two-primary generators. For odd N , the mod-2 orientation obstruction vanishes by transfer. (proved/ ϵ .)

Proof. $E[2] \cong (\mathbb{Z}/2)^2$ has $H^*(B(\mathbb{Z}/2)^2; \mathbb{F}_2) = \mathbb{F}_2[x_1, x_2]$ with $|x_i| = 1$. The reduction of $\beta^E \in H^2(BE; \mathbb{F}_2)$ along the inclusion $E[2] \hookrightarrow E$ gives the displayed quadratic in x_1, x_2 ; reduction to cyclic subgroups gives the linear combinations r_i as in [NIKULIN, §2]. The degree-one linearization class $\lambda^{E, E[2]}$ is the residual character of the underlying line bundle after $\beta = 0$ is imposed; it lives in H^1 and is distinct from the gerbe $\beta \in H^2$ in degree. Even $N \geq 4$ introduces the polynomial generators $y_i \in H^2(B(\mathbb{Z}/2^a)^2)$ and the mixed term $x_1 x_2$, which on cyclic restrictions vanishes; full two-primary detection is needed. Odd N contributes only $\mathbb{F}_p$ -cohomology with p odd, on which the mod-2 orientation has zero restriction by the transfer. $\square$

Remark 4.10 (Translation linearization criterion, not vanishing). For a stabilizer $H \subset E[N]$ and a determinant anchor line bundle $\lambda(\mathcal{F})$, ordinary translation invariance $g^*\lambda \simeq \lambda$ in $\text{Pic}^0(E)$ does not choose coherent isomorphisms $\phi_g : \lambda \rightarrow g^*\lambda$. After the square-root gerbe is killed, the choices form a character torsor in $H^1(BH; \mathbb{F}_2)$, and quotient orientation requires the linearization class

$$\lambda^H \in H^1(BH; \mathbb{F}_2)$$

to vanish. This is a *criterion* for descent on the wrapped stratum; it is not implied by $g^*[\lambda] = [\lambda]$ in $\text{Pic}(E)$. The translation law $\lambda(t \cdot \mathcal{F}) = \lambda(\mathcal{F}) + \chi(\mathcal{F})t$ of Definition 4.7 carries unit weight only when $\chi(\mathcal{F}) \neq 0$; otherwise the anchor is unweighted and descends only after the linearization class vanishes. (proved/ ϵ as a criterion.)

Remark 4.11 (Mod-2 character does not control odd-order anchor characters). The mod-2 orientation character does not by itself control odd-order anchor characters. For anchor descent on the wrapped stratum, also compute the actual stabilizer character on the chosen anchor trivialization, valued in $\hat{H} = \text{Hom}(H, \mathbb{C}^\times)$, or define the retained stabilizer as preserving that trivialization. (unverified/ ϵ at the manuscript level; supplied as part of the finite Hall datum.)

4.5 Cusp polarization and chamber transport

Definition 4.12 (Cusp polarization on Γ_{eff}). The chamber convention $0 < |s| \ll |q| \ll |r|^{-1} \ll 1$ at the standard cusp $\mathfrak{c}_\infty$ determines the cusp-positive semigroup Γ_{eff} of Definition 2.1. The order on triples is lexicographic: $m > 0$ first because s is smallest; if $m = 0$ then $n > 0$ because q is next; if $m = n = 0$ then $l < 0$ because $|r|^{-1} \ll 1$. This is a chamber convention for the completed product, not a microscopic charge positivity statement.

Remark 4.13 (Chamber transport). The pairing $\langle \gamma, \gamma' \rangle_{\text{ind}} = 2(nm' + n'm) - ll'$ is the Gram pairing on triples; the order on Γ_{eff} is a Fock polarization at one cusp. An element of the duality group $\mathbf{Sp}_4(\mathbb{Z})$ transports both the chamber and the expansion to the corresponding product chamber at the transported cusp. The arithmetic group $\mathbf{Sp}_4(\mathbb{Z})$ is the symmetry of the genus-two chemical potential Z and of the invariant Gram plane on $\Lambda^{3,2}$; it is not asserted to be the full microscopic T-/U-duality group of the $K3 \times T^2$ compactification, which is larger (folklore/ β , [?HM, ?DVV]).

Remark 4.14 (Chamber dependence of the wrapped stratum). The cusp chamber chooses a positive direction on Γ_{eff} . At $\mathfrak{c}_\infty$ the wrapped stratum carries $\deg_E > 0$ branes; at a transported cusp the wrapped direction is the image of $\deg_E$ under the corresponding chamber transport. Mixed Hall correspondences are chamber-equivariant when the finite Hall stage R is compatible with the duality element; otherwise the correspondence stack is the original closed flag-Quot stack restricted to the new chamber.

5 LOCAL PROTECTED OBSERVABLE ALGEBRA (SECTORIAL SOURCE DATUM)

The compact realization route of Chapter ?? takes as supplied input a chain-level local observable system on opens $U \subset X = S \times E$. The system is graded by the algebraic/effective Hall support, factorizes on disjoint opens, and restricts on E -projection-finite supports to a chiral object on $\text{Ran}(E)_{b=0}$. The data are external assumptions named in the literature; this section records them as the source interface used downstream.

Definition 5.1 (Local protected observable algebra). Let $U \subset X = S \times E$ be open. Let $\Gamma_{X,\sigma}^{\text{eff}}(U) \subset \Gamma_X^{\text{alg}} \subset \Gamma_X^{\text{form}}$ be the supplied algebraic/effective charge support. For $c \in \Gamma_{X,\sigma}^{\text{eff}}(U)$, let $\mathfrak{M}_c(U)$ be the

supplied derived moduli stack of compactly supported B-branes on U of Mukai charge c . The PTVV theorem supplies the ambient (-1) -shifted symplectic structure on the CY_3 perfect-complex moduli problem (proved/ β , [?PTVV2013]). Let Φ_c denote the BBDJS perverse sheaf $P_{\mathfrak{M}_c, \mathcal{J}_c}^\bullet$ on an oriented d-critical truncation, when such a d-critical structure and square-root orientation are supplied (proved/ β , [?BBDJS2015, Theorem 6.9]). Let o_c be the orientation line, a square root of the determinant line of the cotangent complex of $\mathfrak{M}_c(U)$, supplied as part of the Hall datum.

The *local protected observable algebra* of X on U is the source object

$$\mathcal{A}_X(U) := \bigoplus_{c \in \Gamma_{X, \sigma}^{\text{eff}}(U)} H_*^{\text{BM}}(\mathfrak{M}_c(U), \Phi_c \otimes o_c),$$

graded by the supplied algebraic/effective support, not by the full formal lattice Γ_X^{form} . For pairwise disjoint $U_1, \dots, U_k \subset X$, the proposed factorization map is

$$\mathcal{A}_X(U_1) \otimes \dots \otimes \mathcal{A}_X(U_k) \longrightarrow \mathcal{A}_X(U_1 \sqcup \dots \sqcup U_k),$$

defined chain-level by Massey–Saito Thom–Sebastiani for vanishing cycles

$$\Phi_{F_1 \oplus \dots \oplus F_k} \simeq \Phi_{F_1} \boxtimes \dots \boxtimes \Phi_{F_k}$$

on disjoint d-critical charts (proved/ β , [?MasseyST, ?SaitoMHM]), together with the Ext-vanishing $\text{Ext}^*(\mathcal{F}, \mathcal{G}) = 0$ for branes $\mathcal{F}, \mathcal{G}$ with disjoint supports (folklore/ β).

Remark 5.2 (Source interface: external assumptions). The components are external assumptions: PTVV (-1) -shifted symplectic structure on derived moduli of B-branes (proved/ β , [?PTVV2013]); BBDJS vanishing-cycle perverse sheaf and its Thom–Sebastiani isomorphism (proved/ β , [?MasseyST, ?SaitoMHM, ?BBDJS2015]); orientation lines and Joyce–Upmeyer-compatible multiplicative orientation data (supplied separately, proved/ β , [?JoyceUpmeyer, Theorem 4.1] or [?KSCoHA]). The factorization map relies on disjoint-supports Ext-vanishing for compactly supported branes (folklore/ β ; absent for non-compact supports, where one recovers chiral algebra structure, not commutative factorization). Compactness on $X = S \times E$ is open because the relevant moduli of branes are typically non-quasicompact in Λ_S -charge; the sectorial truncation of §4 is the candidate truncation, conditional on the four obstructions of §6.

6 THE FOUR OBSTRUCTIONS AND THE CONDITIONAL COMPARISON TARGET

The hybrid carrier and the source interface produce a candidate compact holomorphic E_3 -factorization algebra and its K_3 -specialization to a chiral object on $\text{Ran}^{\text{hyb}}(E)$. The cross-level comparison $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}}) = \Delta_5$ is conditional on four obstruction classes named at the level of §5.8 of the manuscript and tracked across the swarm ([?notes/swarm_20260430/reports]). The chapter records them here as the conditioning data feeding Chapter ??.

(Do) Do-degeneration limit. The cosection-reduced d-critical orientation theory on $\text{Hilb}^n(X) \times \text{Sym}^n(X)$, in the Do-degeneration limit $n \rightarrow \infty$, supplies the orientation line on the local stratum (folklore/ β ; conditional, swarm probes Ao05, Ao11, Ao26, Ao37, Ao42).

(Or) Strong reduced orientation. The $K_3 \times E$ -quotient self-Ext frame bundle carries a strong reduced orientation on the wrapped stratum, descending the BBDJS orientation line o_c across the E -quotient (folklore/ β ; conditional, swarm probes Ao74, Ao76, Ao82).

(O₁⁺) Weyl-equivariant transport. The orientation transports $W^{(2)}(\Lambda_{II}^{2,1})$ -equivariantly along reflections, fixing the orientation character $\epsilon_\theta = \nu_{\Delta_5}$ on the BKM Weyl group (folklore/ δ ; conditional, swarm probes Ao47, Ao49, Ao89).

(O₂) Local Pfaffian wall normal form. On type-II walls of $\Lambda_{II}^{2,1}$, the local Pfaffian wall normal form is the $4096 = 2^{12}$ sign sum on the half-Hilbert orbit (computed/ δ at small height; conditional at full height, swarm probes Ao47, Ao92, A100).

The protected scalar shadow on the closed $K3 \times E \rightarrow \text{point}$ cobordism is the OP/DT identity

$$Z_{\text{DT}}^X(T, Q, P) = Z_{\text{OP}}^X(P, Q, T) = -4096 \Delta_5(Z)^{-2} \quad (\text{proved}/\beta, [\text{?OB}, \text{?OPand}, \text{?DT}]).$$

Equivalently $\chi_{\text{io}}^{\text{OP}} = (64^{-1} \Delta_5)^2 = 4096^{-1} \Delta_5^2$ and $Z_{\text{OP}}^X = -(\chi_{\text{io}}^{\text{OP}})^{-1}$; the factor $4096 = 64^2$ is a theta-leading normalization conversion and the minus sign is the OP scalar convention. The orientation-forgetful shadow

$$Z_{\text{BPS}}^{K3 \times E} = (\Phi_{\text{io}}^{\text{un}})^{-1} = \Delta_5^{-2}$$

is the level- n protected trace of $\mathfrak{A}_{K3 \times E}$ on the closed cobordism in the unnormalized convention $\Phi_{\text{io}}^{\text{un}} = \Delta_5^2$; the squared scalar cannot recover the orientation character that distinguishes Δ_5 from $-\Delta_5$. Promotion to a $3d$ gravitational path integral requires the Hall–Borcherds residual of [?chiral-bar-cobar-vol2] and is not claimed. The conditional cross-level identity $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}}) = \Delta_5$ supplies the orientation/Pfaffian datum under (Do), (O₁), (O₁⁺), (O₂).

Remark 6.1 (Cross-volume consistency). The κ -tuple (0, 0, 3, 5, 24) is consistent across [?calabi-yau-quantum-groups] (Vol III), the Hall–Borcherds residual of [?chiral-bar-cobar-vol2] (Vol II), and the Beilinson–Drinfeld factorization framework of [?BD] via Vol I [?chiral-bar-cobar]. The local model $\mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$ + brane completion of [?mixed-holomorphic-topological-strings] contributes the holomorphic de Rham obstruction class on the global target; the identity $Z_{\text{BPS}}^{K3 \times E} = \Delta_5^{-2}$ does not promote to the global target via formal Hamiltonian BF, and the Hall–Borcherds residual is the cross-volume bridge.

7 NOTATION SUMMARY AND CHAPTER OUTPUTS

$X = S \times E$. Smooth complex projective Calabi–Yau threefold; S is a $K3$ surface, E a smooth elliptic curve. $\dim_{\mathbb{C}} X = 3$.

Λ_S . Mukai lattice; even unimodular signature (4, 20). $\Lambda_S \cong U^{\oplus 4} \oplus E_8(-1)^{\oplus 2} \cong II_{4,20}$.

$\Gamma_X^{\text{form}}, \Gamma_X^{\text{phys}}$. Formal full-Mukai dyonic lattice $\Lambda_S \oplus \Lambda_S$. The superscript “phys” is mnemonic only; not the $\mathcal{N} = 4$ charge lattice.

$\Gamma_X^{\text{alg}}, \Gamma_{X,\sigma}^{\text{eff}}$. Algebraic K-class lattice $\subset \Gamma_X^{\text{form}}$; supplied σ -effective semigroup $\subset \Gamma_X^{\text{alg}}$.

$\Gamma_{\text{gram}} = \mathbb{Z}^3$. Gram-index lattice, (n, l, m) .

Π_X . Quadratic Gram map, $\Pi_X(Q, P) = ((Q, Q)/2, (Q, P), (P, P)/2)$.

B . Symmetric bilinear cocycle, $B(c, c') = (Q \cdot Q', Q \cdot P' + Q' \cdot P, P \cdot P')$; satisfies $B = -\delta \Pi_X$ and $B(c, c) = 2\Pi_X(c)$.

$\widehat{\Gamma}_X$. Normal-ordered extension $\Gamma_X^{\text{form}} \oplus_B \Gamma_{\text{gram}}$; abelian under $\star$.

$\overline{\Pi}_X$. Additive normal-ordered Gram homomorphism $\overline{\Pi}_X(c, T) = \Pi_X(c) - T$.

$\text{Ran}^{\text{hyb}}(E)$. Hybrid carrier $\text{Ran}(E)_{b=0} \sqcup \text{Ran}^{\text{wr}}(E)_{b>0}$ on the elliptic factor.

$\mathcal{D}_X = \Delta_5, \text{den}(\mathfrak{g}_{\Delta_5}), \text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}})$. Three named scalars at level five: automorphic section, BKM denominator, compact Pfaffian (View ①, ②, ③).

The chapter outputs are: (a) the formal Mukai sector $\Gamma_X^{\text{form}} = \Lambda_S \oplus \Lambda_S$ and its three-tier support hierarchy; (b) the lattice polarization $B = -\partial\Pi_X$ as the full structural content of the Gram defect; (c) the normal-ordered extension $\widehat{\Gamma}_X$ with additive homomorphism $\overline{\Pi}_X$, fixed by the fixed-lift raw-pushforward obstruction theorem; (d) the hybrid Ran carrier $\text{Ran}^{\text{hyb}}(E)$, fixed by the projection-locality obstruction; (e) the connected/finite split of the E -descent classes, with the linearization criterion separated from the gerbe class; (f) the local protected observable algebra source interface, supplied as external assumption; (g) the four obstructions (Do), (O1), (O1⁺), (O2) conditioning the cross-level identity ②↔③.

Chapter ?? consumes these outputs as fixed input data and constructs the Dirac–Igusa pro-object $\mathfrak{D}_X^{\text{DI}} = \lim_R (\mathcal{A}_{X,R}^{E_3}, F_{X,R}^{\text{hyb}}, \Gamma_{X,R}, \Pi_X, \Phi_R, o_R, H_R, P_R^{\Pi}, C_{X,R}, \Theta_{\text{Kos},R}, \mathcal{L}_{\text{Pf},R}, \text{pf}_{X,R}, \epsilon_{o,R}, \text{Rec}_R)$.

VIEW I: THE BORCHERDS–GRITSENKO–NIKULIN SECTION

1 THE PENTADIC LEVEL AND THE BEILINSON CUT

The first of the five categorical-dimensional levels at which the chiral E_3 -algebra $\mathfrak{A}_{K_3 \times E}$ of the formal target $K_3 \times E$ takes the value Δ_5 is the automorphic level. At this level Δ_5 is read as a holomorphic section of the weight-5 automorphic line on the genus-two Siegel modular space $\mathbf{Sp}_4(\mathbb{Z}) \backslash \mathbb{H}_2$: the Borchers–Gritsenko–Nikulin lift of the weight-zero index-one weak Jacobi form $\phi_{0,1}$, normalized at the isotropic cusp $\mathfrak{c}_\infty$ by the theta-constant leading coefficient 64. The $\mathbf{Sp}_4(\mathbb{Z})$ -equivariant data — multiplier system ν_{Δ_5} , Borchers chamber, divisor support, and weight — constitute the View I package.

1.0.0.1 Three names. The handle “ Δ_5 ” is shared among three structurally distinct objects in the manuscript and must be tagged on first use per chapter:

- *the Borchers–Gritsenko–Nikulin section* $D_X = \Delta_5$ of $\mathcal{L}^5 \otimes \nu_{\Delta_5}$ on $\mathbf{Sp}_4(\mathbb{Z}) \backslash \mathbb{H}_2$, View I, the subject of this chapter;
- *the BKM denominator* $\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1} \Delta_5(2Z)$, View II, constructed in the chapter on Borchers–Kac–Moody algebras;
- *the protected Pfaffian* $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}})$, View III, constructed in the Dirac–Igusa chapter as a section on the Pfaffian line of the cosection-reduced compact Hall datum, conditional on the four obstruction classes (D_0) , (O_1) , $(O_1)^+$, (O_2) .

The bare label Δ_5 never substitutes for any of the three: it is the protected scalar shadow of the chiral E_3 -algebra, while D_X , $\text{den}(\mathfrak{g}_{\Delta_5})$, and $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}})$ live at distinct categorical-dimensional levels.

1.0.0.2 Beilinson cut: scope of View I. View I is the scalar modular shadow. In the reconstitution order “primitive objects first, cross-level identities second, scalar shadows last” the contents of this chapter sit at the bottom: a single holomorphic section of an automorphic line on a Shimura variety, with its multiplier system and its weight. No operator-level package is asserted at View I. In particular, the equality $D_X = \text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}})$ of View I and View III is the Pfaffian–Dirac theorem of Chapter ??, conditional on (D_0) , (O_1) , $(O_1)^+$, (O_2) ; it is not asserted in this chapter. The shadows are Δ_5 , χ_{10} , the level- n BPS partition function $Z_{\text{BPS}}^{K_3 \times E} = \Delta_5^{-2}$, and the Oberdieck–Pixton scalar branch $Z_{\text{OP}}^X = -4096 \Delta_5^{-2}$; none of them is a Hilbert space, a Hall pairing, an orientation line, or a BPS operator product.

2 THE GENUS-TWO SIEGEL SPACE AND THE BORCHERDS CHAMBER

Let $\mathbb{H}_2$ be the genus-two Siegel upper half space. An element $Z \in \mathbb{H}_2$ is a complex symmetric 2×2 matrix with positive-definite imaginary part,

$$Z = \begin{pmatrix} z_1 & z_2 \\ z_2 & z_3 \end{pmatrix}, \quad \Im Z > 0.$$

Set Fourier variables

$$q = e^{2\pi i z_1}, \quad r = e^{2\pi i z_2}, \quad s = e^{2\pi i z_3}.$$

The fixed cusp polarization at the rational isotropic boundary $\mathfrak{c}_\infty$ of $\mathbf{Sp}_4(\mathbb{Z}) \backslash \mathbb{H}_2$ — the standard type-II cusp polarization used throughout the manuscript — is the ordered limit

$$0 < |s| \ll |q| \ll |r|^{-1} \ll 1.$$

Equivalently, $\mathfrak{I}_{z_3}$ goes to $+\infty$ faster than $\mathfrak{I}_{z_1}$ goes to $+\infty$, and $\mathfrak{I}_{z_2}$ is bounded. This polarization selects a Weyl chamber on the $O(3, 2)$ symmetric domain through the $\mathrm{Sp}(4) \simeq \mathrm{Spin}(3, 2)$ bridge of Chapter 5, and hence selects a positive cone — the effective semigroup

$$\Gamma_{\mathrm{eff}} = \{(n, l, m) \mid m > 0, n \geq 0, l \in \mathbb{Z}\} \cup \{(n, l, 0) \mid n > 0, l \in \mathbb{Z}\} \cup \{(0, l, 0) \mid l < 0\} \subset \mathbb{Z}^3,$$

and a corresponding completed Fourier expansion ring. In this completion every Fourier coefficient of every formal product receives only finitely many contributions: the chamber walls are the inequalities defining the ordering, and the chamber-completed product is the only product considered.

2.0.0.1 Choices fixed once and for all. The cusp $\mathfrak{c}_\infty$, the Fourier variables (q, r, s) , the polarization $(0 < |s| \ll |q| \ll |r|^{-1} \ll 1)$, and the effective semigroup $\Gamma_{\mathrm{eff}} \subset \mathbb{Z}^3$ are fixed. Every product expansion in this chapter is the product expansion in the cusp-completed semigroup algebra of Γ_{eff} . The cusp expansion at any other rational isotropic cusp is obtained by an $\mathbf{Sp}_4(\mathbb{Z})$ -translate of these data.

3 THE WEAK JACOBI INPUT AND THE K_3 ELLIPTIC GENUS

The Borchers input is the *weight-zero index-one weak Jacobi form* $\phi_{0,1} \in J_{0,1}^{\mathrm{weak}}$ of Eichler–Zagier [EZ:1, §2.2], normalized so that the K_3 elliptic genus is

$$Z_{K_3}(\tau, z) = 2 \phi_{0,1}(\tau, z).$$

Write the Fourier expansion as

$$\phi_{0,1}(\tau, z) = \sum_{n \geq 0, l \in \mathbb{Z}} f(n, l) q^n r^l, \quad q = e^{2\pi i \tau}, \quad r = e^{2\pi i z}.$$

The first Fourier coefficients are

$$\begin{aligned} f(0, -1) &= 1, & f(0, 0) &= 10, & f(0, 1) &= 1, \\ f(1, -2) &= 10, & f(1, -1) &= -64, & f(1, 0) &= 108, & f(1, 1) &= -64, & f(1, 2) &= 10, \\ f(2, -3) &= 1, & f(2, -2) &= 108, & f(2, -1) &= -513, & f(2, 0) &= 808, \end{aligned} \tag{3.1}$$

extending in the standard way by hyperbolic symmetry $f(n, l) = f(n, -l)$ and Eichler–Zagier support $f(n, l) = 0$ for $4n - l^2 < -1$ [EZ:1, §2.2]. The values in (3.1) are computed in the local fixture `compute/verify_square_root.py` from the standard theta-quotient expression $\phi_{0,1} = 4 \sum_{i=2,3,4} \theta_i(\tau, z)^2 / \theta_i(\tau, 0)^2$ [computed; cross-checked against [EZ:1, Theorem 9.2]].

3.0.0.1 Borchers-weight handle. Following the universal Borchers-weight identity $\kappa_{\text{BKM}}(\Phi_N) = c_N(\mathbf{o})/2$ (Borchers 1995 [?Borchers95, Theorem 10.1]; Gritsenko 2018 [?Gritsenko2018Reflective]), the input row $N = \mathbf{1}$ of the Cléry–Gritsenko catalogue [?GC, Theorem 1.4] is named by

$$c_1(\mathbf{o}) := f(\mathbf{o}, \mathbf{o}) = 10, \quad \kappa_{\text{BKM}}(\Delta_5) = \frac{c_1(\mathbf{o})}{2} = 5,$$

to be matched in Proposition 6.1 below with the geometric weight of the Borchers lift. The notation κ_{BKM} is the BKM cusp-form-weight entry of the κ -tuple $(\kappa_{\text{cat}}, \kappa_{\text{ch}}^{\text{Hodge}}, \kappa_{\text{ch}}^{\text{Heis}}, \kappa_{\text{BKM}}, \kappa_{\text{fiber}}) = (\mathbf{o}, \mathbf{o}, 3, 5, 24)$ of the $K_3 \times E$ target. The bare symbol κ is forbidden — every entry carries its semantic subscript; the additive collapse $\kappa_{\text{BKM}} = \kappa_{\text{ch}} + \chi(\mathcal{O}_{\text{fiber}})$ fails on the CHL rows beyond $N = \mathbf{1}$ and is replaced by the universal weight identity (Vol III).

4 THE THETA-CONSTANT NORMALIZATION

The theta-constant Igusa form Δ_5 is fixed by its leading Fourier monomial at $\mathbf{c}_\infty$. Writing the theta-constant representation of Igusa [?Igusa1962, §5], [?Igusa1964II, §2]

$$\Delta_5(Z) = \prod_{\substack{a, b \in (\mathbb{Z}/2\mathbb{Z})^2 \\ {}^t ab \equiv 0 \pmod{2}}} \nu_{a,b}(Z), \quad \nu_{a,b}(Z) = \sum_{l \in \mathbb{Z}^2} \exp(\pi i (z[l + \tfrac{1}{2}a] + {}^t bl)),$$

the four characteristics with $a = (\mathbf{o}, \mathbf{o})$ start with the constant 1, the two factors with $a = (\mathbf{1}, \mathbf{o})$ start with $2q^{1/8}$, the two factors with $a = (\mathbf{o}, \mathbf{1})$ start with $2s^{1/8}$, and the two factors with $a = (\mathbf{1}, \mathbf{1})$ start with $2q^{1/8}r^{1/4}s^{1/8}$; multiplying gives

PROPOSITION 4.1 (*Theta-constant leading coefficient*).

$$[q^{1/2}r^{1/2}s^{1/2}]\Delta_5 = 2^6 = 64.$$

[*proved*; Igusa 1962 [?Igusa1962, Theorem 3], Gritsenko 1999 [?GN, Theorem 4.1]; cross-checked numerically in `compute/verify_square_root.py`.]

Definition 4.2 (*Monic Borchers–Gritsenko form*). The *monic genus-two Borchers–Gritsenko form* is

$$D_5(Z) := 64^{-1}\Delta_5(Z).$$

By Proposition 4.1, $D_5 = q^{1/2}r^{1/2}s^{1/2} + \dots$ at $\mathbf{c}_\infty$.

This is the normalization used by Gritsenko–Nikulin [?GNII, Theorem 2.1, (2.15)–(2.16)]: their Borchers product is the monic form D_5 ; the theta-constant convention restores the prefactor 64.

5 THE BORCHERS–GRITSENKO–NIKULIN SECTION

5.1 The product expansion

Borchers’s automorphic product theorem [?Borchers95, Theorem 10.1], specialized to the genus-two Siegel modular variety $\mathbf{Sp}_4(\mathbb{Z}) \backslash \mathbb{H}_2$ by Gritsenko–Nikulin [?GNII, Theorem 2.1, Example 2.4] via the exceptional isomorphism $\mathbf{Sp}_4 \simeq \mathbf{Spin}(3, 2)$ of Chapter 5, lifts the weak Jacobi form $\phi_{\mathbf{o}, \mathbf{1}}$ to a meromorphic Siegel modular form of weight $f(\mathbf{o}, \mathbf{o})/2 = 5$ and multiplier system ν_{Δ_5} with infinite product expansion in the cusp chamber.

THEOREM 5.1 (*Borcherds–Gritsenko–Nikulin lift, monic form*). In the cusp chamber $0 < |s| \ll |q| \ll |r|^{-1} \ll 1$,

$$D_5(Z) = q^{1/2} r^{1/2} s^{1/2} \prod_{(n,l,m) \in \Gamma_{\text{eff}}} (1 - q^n r^l s^m)^{f(nm,l)},$$

the product converging absolutely on a neighborhood of $\mathfrak{c}_\infty$ and continuing meromorphically to all of $\mathbb{H}_2$ as a holomorphic Siegel modular form of weight 5 on $\mathbf{Sp}_4(\mathbb{Z})$ with character ν_{Δ_5} of order two.

[*proved*; Borcherds 1995 [?Borcherds95, Theorem 10.1], Gritsenko–Nikulin 1998 [?GNII, Theorem 2.1, (2.15)–(2.16)], Gritsenko 1999 [?GN, Theorem 4.1].]

COROLLARY 5.2 (*Theta-normalized section*).

$$\Delta_5(Z) = 64 q^{1/2} r^{1/2} s^{1/2} \prod_{(n,l,m) \in \Gamma_{\text{eff}}} (1 - q^n r^l s^m)^{f(nm,l)}.$$

The exponent $f(nm, l)$ of the (n, l, m) -factor depends only on the Hecke–Verlinde–DMVV product nm and on the Jacobi index l , reflecting the second-quantized structure of the Borcherds lift; the combinatorial origin of this exponent is the Dijkgraaf–Moore–Verlinde–Verlinde elliptic genus of symmetric products [?DMVV].

5.2 The View I package

Definition 5.3 (*The Borcherds–Gritsenko–Nikulin section*). The Borcherds–Gritsenko–Nikulin section of the weight-5 automorphic line is the View I representative

$$D_X := \Delta_5$$

on $\mathbf{Sp}_4(\mathbb{Z}) \backslash \mathbb{H}_2$, with Borcherds product expansion at $\mathfrak{c}_\infty$ given by Corollary 5.2.

Remark 5.4 (*D_X is the View I name*). The handle D_X is the View I name of the section. It coincides as a holomorphic Siegel modular form with the theta-constant Igusa cusp form Δ_5 ; the renaming records the role: View I scalar shadow, the determinant of the virtual K3 index in $K_0(\text{sVect}_{\text{fd}})$ (the K_0 -determinant package proved at Proposition 7.2; the operator upgrade is View III), and the input section of the Pfaffian–Dirac theorem of Chapter ?? . The compact operator-level package $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}})$ of View III is asserted equal to D_X under (D_0) , (O_1) , $(O_1)^+$, (O_2) ; without these obstruction trivializations, only D_X is rigorously available.

6 WEIGHT, MULTIPLIER, AND DIVISOR

6.1 The Borcherds weight

PROPOSITION 6.1 (*Borcherds–Gritsenko weight identity at $N = 1$*).

$$\text{wt}(D_X) = \frac{f(0,0)}{2} = 5 = \frac{c_1(0)}{2} = \kappa_{\text{BKM}}(\Delta_5).$$

The weight is integral; no metaplectic cover enters the weight factor.

[*proved*; Borcherds 1995 [?Borcherds95, Theorem 10.1, weight formula]; Gritsenko–Nikulin 1998 [?GNII, Example 2.4]; Gritsenko 1999 [?GN, Theorem 4.1, weight]; universal Borcherds-weight identity $\kappa_{\text{BKM}}(\Phi_N) = c_N(0)/2$ across the Cléry–Gritsenko ladder [?GC], Vol III.]

The row $(N, \text{wt}, c_N(o)) = (1, 5, 10)$ is the leading entry of the eight-row Cléry–Gritsenko catalogue [?GC, Theorem 1.4]; the remaining seven rows $(t, N; k)$ carry weights $(2, 3, 1, 2, 1/2, 3/2, 1)$ with input zero coefficients $c_N(o) \in \{4, 6, 2, 4, 1, 3, 2\}$ and reproduce $\kappa_{\text{BKM}}(\Phi_N) = c_N(o)/2$ row-by-row. The half-integer weights are realized via multiplier systems $(v_\eta^3 \times v_H)$ and not via metaplectic covers. The Vol III universal Borcherds-weight identity replaces the false additive collapse $\kappa_{\text{BKM}} = \kappa_{\text{ch}} + \chi(\mathcal{O}_{\text{fiber}})$ that was once written for the $K_3 \times E$ row alone (where it holds by the $N = 1$ accident $5 = 3 + 2$) and fails at every other CHL row [?GN, FRONTIER, eq. (72), eq. (118)]; the corrected entry of the κ -tuple of $K_3 \times E$ is precisely $\kappa_{\text{BKM}} = 5$.

6.2 The Maass multiplier system

The character $\nu_{\Delta_5} : \mathbf{Sp}_4(\mathbb{Z}) \rightarrow \mathbb{C}^\times$ of Maass [?MAASS: 1] has order two, $\nu_{\Delta_5}^2 = 1$, and is determined on the standard generators of $\mathbf{Sp}_4(\mathbb{Z})$ by

$$\begin{aligned} \nu_{\Delta_5} \begin{pmatrix} o & \mathbf{I}_2 \\ -\mathbf{I}_2 & o \end{pmatrix} &= 1, & \nu_{\Delta_5} \begin{pmatrix} \mathbf{I}_2 & B \\ o & \mathbf{I}_2 \end{pmatrix} &= (-1)^{b_1+b_2+b_3}, \\ \nu_{\Delta_5} \begin{pmatrix} {}^t A^{-1} & o \\ o & A \end{pmatrix} &= (-1)^{(1+a_1+a_4)(1+a_2+a_3)+a_1a_4}, \end{aligned}$$

with $B = \begin{pmatrix} b_1 & b_2 \\ b_2 & b_3 \end{pmatrix}$ and $A = \begin{pmatrix} a_1 & a_2 \\ a_3 & a_4 \end{pmatrix}$ [?MAASS: 1]. The square $\Delta_5^2 = \Delta_{10}$ is scalar. The multiplier records the obstruction to identifying Δ_5 with an ordinary modular form on $\mathbf{Sp}_4(\mathbb{Z})$: Δ_5 is a section of $\mathcal{L}^5 \otimes \nu_{\Delta_5}$, where $\mathcal{L}$ is the Hodge automorphic line and ν_{Δ_5} is the order-two character; the section Δ_5 does not descend to a section of the bare Hodge line $\mathcal{L}^5$, but its square Δ_5^2 is a scalar section of $\mathcal{L}^{10}$.

6.3 The diagonal divisor

Let

$$\mathcal{H}_{\text{diag}} = \left\{ \begin{pmatrix} a & o \\ o & b \end{pmatrix} \mid a, b \in \mathbb{H}_1 \right\} \subset \mathbb{H}_2$$

be the locus of diagonal genus-two periods. The Cléry–Gritsenko classification [?GC] records

$$\text{supp div}(\Delta_5) = \mathbf{Sp}_4(\mathbb{Z}) \cdot \mathcal{H}_{\text{diag}}, \quad \text{mult}_{\mathcal{H}_{\text{diag}}}(\Delta_5) = 1,$$

so the divisor of Δ_5 is the simple paramodular Humbert divisor. In the cusp-chamber language, the type-II reflective walls $\mathcal{H}_{\delta_i}$, $i = 1, 2, 3$, of the lattice $\Lambda_{II}^{2,1}$ (the type-II sublattice of the Borcherds even lattice $\Lambda^{2,1}$) are the components of $\mathbf{Sp}_4(\mathbb{Z}) \cdot \mathcal{H}_{\text{diag}}$ that meet the chosen fundamental Weyl chamber Π_{II} (Chapter ??); the corresponding simple roots $\delta_1 = 2f_2 - f_3$, $\delta_2 = 2f_{-2} - f_3$, $\delta_3 = f_3$ have Gram pairings $(\delta_i, \delta_i) = 2$ and $(\delta_i, \delta_j) = -2$ for $i \neq j$ [?GNII, Example 2.4], and the corresponding Borcherds divisor orders are

$$d_{\delta_1}^{\text{aut}} = f(o, 1) = 1, \quad d_{\delta_2}^{\text{aut}} = f(o, 1) = 1, \quad d_{\delta_3}^{\text{aut}} = f(o, -1) = 1.$$

[*proved*; Cléry–Gritsenko 2011 [?GC, Theorem 1.4]; Gritsenko–Nikulin 1998 [?GNII, Example 2.4]; lattice computation in `compute/verify_lattice.py`.]

6.4 Maass-character values on the type-II walls

LEMMA 6.2 (*Maass-character values on the type-II reflections*). With the simple type-II roots $\delta_1 = 2f_2 - f_3$, $\delta_2 = 2f_{-2} - f_3$, $\delta_3 = f_3$ and chamber-permuting type-I roots $\alpha_1 = f_{-2} - f_2$, $\alpha_2 = f_2 - f_3$, $\alpha_3 = f_2 + f_3$,

$$\nu_{\Delta_5}(s_{\delta_i}) = -1, \quad \nu_{\Delta_5}(s_{\alpha_i}) = +1, \quad i = 1, 2, 3.$$

Equivalently, $\Delta_5(s_{\delta_i}Z) = -\Delta_5(Z)$ and $\Delta_5(s_{\alpha_i}Z) = +\Delta_5(Z)$.

[*proved*; Maass [?MAASS: 1]; Gritsenko–Nikulin [?GN, §3]; Gritsenko [?Gritsenko99]; the type-II values follow from $d_{\delta_i}^{\text{aut}} = 1$ and the type-I values from $d_{\alpha_i}^{\text{aut}} = 0$ (vanishing of $f(0, \pm 2)$ by the support of $\phi_{0,1}$ at $q = 0$). The detailed cocycle calculation sits in Chapter ??, Lemma ??.]

The chamber-permuting reflections $s_{f_2-f_3}, s_{f_2-f_1}, s_{f_2+f_3}$ act trivially on Δ_5 because the corresponding type-I roots α_i are not $\mathbf{Sp}_4$ -images of timelike short roots in the cusp chamber; their Borcherds divisor order vanishes by the support $f(0, l) = 0$ for $|l| \geq 2$ of $\phi_{0,1}$ at $q = 0$ [?EZ: 1, §2.2].

7 IDENTIFICATION WITH THE VIRTUAL K_0 -DETERMINANT

The Fock superdeterminant of the Borcherds-graded super vector space $\mathcal{V} = \bigoplus_{(n,l,m) \in \Gamma_{\text{eff}}} \mathbb{U}_{nm,l}$ — the K_0 -class of the second-quantized Hecke–DMVV half of the K_3 elliptic genus, with the $m = 0$ cusp/Weyl boundary factors carrying the input multiplicities $f(0, l)$ (the construction $\mathcal{V} = \mathcal{V}^{\text{bulk}} \oplus \mathcal{V}^{\text{cusp}}$ of Chapter ??, Definition ??) — is

LEMMA 7.1 (*K_0 -Fock superdeterminant*).

$$\text{sdet}_{\mathcal{V}}(1 - x) = \prod_{(n,l,m) \in \Gamma_{\text{eff}}} (1 - q^n r^l s^m)^{f(nm,l)}, \quad x^{(n,l,m)} = q^n r^l s^m.$$

[*proved*; Fock-trace lemma of Chapter ??, Lemma ??.]

PROPOSITION 7.2 (*BGN identification of the virtual determinant*). With $v_0(Z) := 64 q^{1/2} r^{1/2} s^{1/2}$ the cusp/Weyl monomial of Chapter ??, the normalized K_0 -determinant agrees with the View I section,

$$\mathcal{D}_X(Z) := v_0(Z) \text{sdet}_{\mathcal{V}}(1 - x) = D_X(Z) = \Delta_5(Z),$$

in the cusp-chamber expansion at $\mathfrak{c}_{\infty}$ and hence as sections of $\mathcal{L}^5 \otimes \nu_{\Delta_5}$.

Proof. Combine Theorem 5.1, the theta-constant prefactor Proposition 4.1, Lemma 7.1, and Cohen’s q -expansion principle for genus-two Siegel modular forms with half-integral character: a meromorphic Siegel form is determined by its expansion at one isotropic cusp. Both sides have the same Borcherds product expansion at $\mathfrak{c}_{\infty}$ by direct comparison of exponents, and D_X is holomorphic of weight 5 by Theorem 5.1. $\square$

[*proved*; Borcherds 1995 [?Borcherds95, Theorem 10.1], Gritsenko–Nikulin 1998 [?GNII, Theorem 2.1], Igusa 1962/1964 [?Igusa1962, ?Igusa1964II].]

Remark 7.3 (*Scope of the BGN identification*). Proposition 7.2 is an equality of holomorphic sections of an automorphic line bundle. The carrier $\mathcal{V}$ is a class in $K_0(\Gamma_{\text{eff}}\text{-graded super vector spaces})$; it is not a Hilbert space, not a vertex algebra, not a holomorphic E_3 -factorization algebra, and not a chiral factorization algebra. The operator-level upgrade — promotion of the K_0 -Fock determinant to a section of an oriented Pfaffian line over the cosection-reduced compact Hall datum, and identification of that Pfaffian with D_X — is the Pfaffian–Dirac theorem of Chapter ??, conditional on $(D_0), (O_1), (O_1)^+, (O_2)$. At View I, no such promotion is asserted.

8 THE SQUARED SCALAR SHADOW AND OP NORMALIZATION

The Maass character has order two, so the square $\Delta_5^2 = \Delta_{10}$ is a scalar weight-10 Igusa form on $\mathbf{Sp}_4(\mathbb{Z})$, with no multiplier system.

PROPOSITION 8.1 (*The square is the Oberdieck–Pandharipande χ_{10}*).

$$\Delta_5^2 = \Delta_{10} = \Phi_{10} = \chi_{10}.$$

[*proved*; Igusa [?Igusa1962, ?Igusa1964II]; the symbol Φ_{10} is Borchers’s name for the squared theta-constant Igusa form, χ_{10} is the Oberdieck–Pandharipande [?OPand, ?OB] name; both equal Δ_5^2 . The naming is $\Phi_{10} = \chi_{10} = \Delta_5^2$, with the Oberdieck–Pixton OP normalization $\chi_{10}^{\text{OP}} = D_5^2 = 4096^{-1}\Delta_5^2$ the monic-on- D_5 variant.]

8.0.0.1 *The OP scalar branch.* With the OP normalization $\chi_{10}^{\text{OP}} = D_5^2$, the Oberdieck–Pixton scalar partition function on $K_3 \times E$ is [?OB, (0.4) and Theorem 1]

$$Z_{\text{OP}}^X = -(\chi_{10}^{\text{OP}})^{-1} = -4096 \Delta_5^{-2}.$$

The factor $-4096 = -(64)^2$ records the squared theta-constant leading coefficient and the OP scalar sign convention. At View I this is the protected scalar shadow of the chiral E_3 -algebra of $K_3 \times E$; the operator package whose protected trace it is remains conditional on (D_o) , (O_t) , $(O_t)^+$, (O_2) .

8.0.0.2 *Three names for the squared form, distinguished.* The notation here is fixed once for the manuscript: $\Delta_5^2 = \Delta_{10}$ is the Igusa convention (theta-constant square); $\Phi_{10} = \Delta_5^2$ is the Borchers–Eisenstein convention; $\chi_{10} = \Delta_5^2$ is the Oberdieck–Pandharipande convention; and $\chi_{10}^{\text{OP}} = 4096^{-1}\Delta_5^2 = D_5^2$ is the Oberdieck–Pixton variant in which D_5 is the monic Borchers form (Definition 4.2).

9 THE VIEW II SHADOW: THE $2Z$ DENOMINATOR

The Gritsenko–Nikulin denominator algebra of the Borchers–Kac–Moody Lie superalgebra $\mathfrak{g}_{\Delta_5}$ arises as the View II shadow of Δ_5 under the doubling $Z \mapsto 2Z$:

PROPOSITION 9.1 (*Gritsenko–Nikulin doubled denominator*).

$$\text{den}(\mathfrak{g}_{\Delta_5}) = D_5(2Z) = 64^{-1}\Delta_5(2Z).$$

[*proved*; Gritsenko–Nikulin 1998 [?GNII, Theorem 2.1, (2.15)–(2.16); §5]; Gritsenko 1999 [?GN, Theorem 4.1].]

The doubling reflects the level-2 structure of the BKM root lattice $\Lambda_{II}^{2,1}$ inside $\Lambda^{2,1}$: the type-II walls of the denominator side carry roots of length-square 2 and live on a sublattice of index 2; the substitution $Z \mapsto 2Z$ on the modular side restricts the Fourier expansion to the corresponding sublattice [?GNII, Theorem 2.1, (2.15)]. At View I this is a pointer; the Cartan, Chevalley, real-and-imaginary roots, and Weyl–Kac–Borchers denominator identity of $\mathfrak{g}_{\Delta_5}$ sit in Chapter ??.

10 THE $\mathbf{Sp}(4) \simeq \mathbf{Spin}(3,2)$ BRIDGE TO VIEW V

The Borchers construction is most naturally written on the orthogonal side: Δ_5 is the singular theta lift of $\phi_{0,1}$ under the Howe correspondence $\mathbf{Mp}(4, \mathbb{R}) \leftrightarrow O(3, 2)$, bridged by the exceptional isomorphism

$$\mathbf{Sp}_4 \simeq \mathbf{Spin}(3, 2)$$

of low-dimensional spin geometry. At the symbolic level View I (Δ_5 on $\mathbf{Sp}_4(\mathbb{Z}) \backslash \mathbb{H}_2$) and View V (Δ_5 as the singular theta lift on $O(3, 2) \backslash D_{O(3,2)}$) are the two sides of this bridge: the weight-5 automorphic line on $\mathbf{Sp}_4(\mathbb{Z}) \backslash \mathbb{H}_2$ is the pullback of the weight-5 orthogonal modular line on $O(3, 2) \backslash D_{O(3,2)}$, and the multiplier ν_{Δ_5} restricts to the orthogonal Maass character. The full theta-lift construction — Borchers’ *singular theta correspondence* [?Borchers95], [?B] — is presented in Chapter 5; this chapter uses only the $\mathrm{Sp}(4)$ -side of the bridge.

II SUMMARY OF VIEW I

THEOREM II.1 (*View I package*). The genus-two Igusa cusp form Δ_5 is the Borchers–Gritsenko–Nikulin section

$$D_X := \Delta_5$$

of the weight-5 automorphic line $\mathcal{L}^5 \otimes \nu_{\Delta_5}$ over $\mathbf{Sp}_4(\mathbb{Z}) \backslash \mathbb{H}_2$:

- (i) At the rational isotropic cusp $\mathfrak{c}_\infty$ with cusp polarization $0 < |s| \ll |q| \ll |r|^{-1} \ll 1$, D_X admits the Borchers product expansion

$$D_X(Z) = 64 q^{1/2} r^{1/2} s^{1/2} \prod_{(n,l,m) \in \Gamma_{\mathrm{eff}}} (1 - q^n r^l s^m)^{f(nm,l)},$$

with leading theta-constant coefficient $[q^{1/2} r^{1/2} s^{1/2}] D_X = 64$, and exponents $f(nm, l)$ the Fourier coefficients of the K_3 weight-zero index-one weak Jacobi form $\phi_{0,1} = \frac{1}{2} Z_{K_3}$ (Eichler–Zagier [?EZ: 1, §2.2]).

- (ii) D_X has weight $\mathrm{wt}(D_X) = f(0, 0)/2 = 5 = c_1(0)/2 = \kappa_{\mathrm{BKM}}(\Delta_5)$, the leading entry $(N, \mathrm{wt}, c_N(0)) = (1, 5, 10)$ of the Cléry–Gritsenko catalogue [?GC, Theorem 1.4]. The multiplier is the Maass character ν_{Δ_5} of order two.
- (iii) The divisor of D_X is the simple paramodular Humbert divisor $\mathbf{Sp}_4(\mathbb{Z}) \cdot \mathcal{H}_{\mathrm{diag}}$; the type-II reflective walls $\mathcal{H}_{\delta_i}$ ($i = 1, 2, 3$) carry order $d_{\delta_i}^{\mathrm{aut}} = 1$, and $\nu_{\Delta_5}(s_{\delta_i}) = -1 = (-1)^{d_{\delta_i}^{\mathrm{aut}}}$, while the chamber-permuting roots α_i carry order 0 and $\nu_{\Delta_5}(s_{\alpha_i}) = +1$ (Lemma 6.2).
- (iv) The square $D_X^2 = \Delta_{10} = \Phi_{10} = \chi_{10}$ is a scalar weight-10 Igusa form, and the Oberdieck–Pixton scalar branch is $Z_{\mathrm{OP}}^X = -(\chi_{10}^{\mathrm{OP}})^{-1} = -4096 \Delta_5^{-2}$.
- (v) The View II shadow is the doubled denominator $\mathrm{den}(\mathfrak{g}_{\Delta_5}) = D_5(2Z) = 64^{-1} \Delta_5(2Z)$ (Proposition 9.1); the View III operator-level package $\mathrm{Pf}_{\mathrm{prot}}(\mathfrak{D}_X^{\mathrm{DI}})$ is the subject of Chapter ??, conditional on $(D_0), (O_1), (O_1)^+, (O_2)$.

[*proved*; assembled from Borchers 1995 [?Borchers95, Theorem 10.1], Borchers 1998 [?B], Gritsenko–Nikulin 1998 [?GNII, Theorem 2.1, Example 2.4], Gritsenko 1999 [?GN, Theorem 4.1], Igusa 1962 [?Igusa1962, Theorem 3], Igusa 1964 [?Igusa1964II, §2], Cléry–Gritsenko 2011 [?GC, Theorem 1.4], Maass 1964 [?MAASS: 1], Eichler–Zagier 1985 [?EZ: 1, §2.2], Oberdieck–Pixton 2018 [?OB, (0.4) and Theorem 1], Oberdieck–Pandharipande 2016 [?OPand]; numerical checks in `compute/verify_lattice.py` and `compute/verify_square_root.py`.]

II.0.0.1 *What View I does not assert.* View I does not assert any operator-level package on $K_3 \times E$, does not construct the Borchers–Kac–Moody algebra $\mathfrak{g}_{\Delta_5}$ (Chapter ??), does not construct the protected first-order operator $\mathfrak{D}_X^{\mathrm{DI}}$ or the Pfaffian line $\mathcal{L}_{\mathrm{Pf}, X}$ (Chapter ??), and does not commit to the $K_3 \times E$ realization of the chiral E_3 -algebra $\mathfrak{A}_{K_3 \times E}$ at any level above the scalar shadow. In the language of the

Beilinson cut, View I is the bottom of the reconstitution order: a holomorphic Siegel cusp form, with its weight, multiplier, divisor, and Borchers product, and nothing more.

THE BORCHERDS–KAC–MOODY DENOMINATOR ALGEBRA $\mathfrak{g}_{\Delta_5}$ ON $\Lambda_{II}^{2,1}$

The Igusa cusp form Δ_5 admits a Borchers–Kac–Moody description as a denominator. The denominator algebra is the generalized Kac–Moody Lie superalgebra $\mathfrak{g}_{\Delta_5}$ on the Lorentzian root lattice $\Lambda_{II}^{2,1} = \Lambda^{(1,1)} \oplus [2]$ in its type-II sublattice presentation, with three even real simple roots $\delta_1, \delta_2, \delta_3$, an infinite imaginary simple-root datum read from the K3 half-index $\phi_{0,1}$, and Weyl group $W^{(2)}(\Lambda_{II}^{2,1})$ generated by the three real reflections. Its Weyl–Kac–Borchers denominator is the level-five Borchers–Kac–Moody denominator

$$\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1} \Delta_5(2Z),$$

the doubled Igusa form normalized by the theta-constant constant $[q^{1/2} r^{1/2} s^{1/2}] \Delta_5 = 64$. The Borchers-weight characteristic of the denominator is $\kappa_{\text{BKM}}(\Delta_5) = c_1(o)/2 = 10/2 = 5$, the weight of the paramodular form $\Delta_5 = \Phi_1$ at level 1.

The chapter is target-side. The denominator algebra is the chiral shadow of an open compact realization. No compact Hall bracket, protected Pfaffian, or BPS Hilbert space is claimed; those are the conditional contribution of Chapter ?? under the four obstructions $(Do), (O1), (O1)^+, (O2)$.

The three named instances of Δ_5 are tagged distinctly. In this chapter the canonical handle is the Borchers–Kac–Moody denominator,

$$\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1} \Delta_5(2Z).$$

The automorphic section $\mathcal{D}_X = \Delta_5 \in M_5(\mathbf{Sp}_4(\mathbb{Z}), \nu_{\Delta_5})$ of Chapter ?? is View I; the compact protected Pfaffian $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{DI})$ of Chapter ?? is View III, conditional. The bare handle Δ_5 is the working symbol for the Siegel cusp form of weight five and never substitutes for any of the three.

1 THE HYPERBOLIC LATTICE $\Lambda^{2,1} = \Lambda^{(1,1)} \oplus [2]$ AND ITS TYPE-II SUBLATTICE

Fix the rank-three Lorentzian lattice

$$\Lambda^{2,1} = \Lambda^{(1,1)} \oplus [2] = \mathbb{Z}f_2 \oplus \mathbb{Z}f_3 \oplus \mathbb{Z}f_{-2},$$

where $\Lambda^{(1,1)}$ is the hyperbolic plane and $[2]$ is the rank-one lattice generated by a single vector of square +2. The bilinear form on the chosen basis is

$$(f_2, f_{-2}) = -1, \quad (f_2, f_2) = (f_{-2}, f_{-2}) = 0, \quad (f_3, f_3) = 2, \quad (4.1)$$

with all other pairings between basis vectors zero. The lattice is even, signature $(2, 1)$, and primitively embedded in the rank-five Lorentzian lattice $\Lambda^{3,2} \simeq \Lambda^{(1,1)} \oplus \Lambda^{(1,1)} \oplus [2]$ of Section ??.

Definition 1.1 (Type-II sublattice). The type-II sublattice of $\Lambda^{2,1}$ is

$$\Lambda_{II}^{2,1} = \{mf_2 + lf_3 + nf_{-2} \in \Lambda^{2,1} \mid m \equiv n \equiv 0 \pmod{2}\}.$$

Its dual lattice is

$$(\Lambda_{II}^{2,1})^* = \mathbb{Z}\frac{1}{2}f_2 \oplus \mathbb{Z}\frac{1}{2}f_3 \oplus \mathbb{Z}\frac{1}{2}f_{-2} = \frac{1}{2}\Lambda^{2,1},$$

of index four in $\Lambda^{2,1}$.

Remark 1.2 (The two divisibility classes of square-2 vectors). A primitive element $\delta \in \Lambda^{2,1}$ with $(\delta, \delta) = 2$ has divisibility either $(\delta, \Lambda^{2,1}) = \mathbb{Z}$ (type I) or $(\delta, \Lambda^{2,1}) = 2\mathbb{Z}$ (type II). The type-I roots lie in the sublattice

$$\Lambda_I^{2,1} = \{mf_2 + lf_3 + nf_{-2} \in \Lambda^{2,1} \mid m + l + n \equiv 0 \pmod{2}\}$$

and the type-II roots lie in $\Lambda_{II}^{2,1}$. The split is recorded in Gritsenko–Nikulin [GN, Section 1]; the underlying classification of square-2 vectors on $\Lambda^{(1,1)} \oplus [2]$ is Nikulin’s [NIKULIN].

Lemma 1.3 (Reflection-group decomposition). Let $W^{(2)}(\Lambda^{2,1}) \subset \mathbf{O}(\Lambda^{2,1})_+$ be the subgroup generated by reflections in primitive square-2 vectors. Then

$$\mathbf{O}(\Lambda^{2,1})_+ = W^{(2)}(\Lambda^{2,1}),$$

and there are two normal subgroups corresponding to the two divisibility classes,

$$[W^{(2)}(\Lambda^{2,1}) : W^{(2)}(\Lambda_I^{2,1})] = 2, \quad [W^{(2)}(\Lambda^{2,1}) : W^{(2)}(\Lambda_{II}^{2,1})] = 6. \quad (4.2)$$

Proof. The first equality is Nikulin’s [NIKULIN] reflection-group theorem for $\Lambda^{(1,1)} \oplus [2]$. The two indices are the two cosets of the divisibility classes in $\mathbf{O}(\Lambda^{2,1})_+$; type-II reflections generate a subgroup of index six because the three orbit representatives $\{f_2 - f_3, f_{-2} - f_2, f_2 + f_3\}$ of the chamber automorphism S_3 are type-I and lie outside $W^{(2)}(\Lambda_{II}^{2,1})$. $\square$

The change of polarization between Gram-index parametrization and the denominator-side Lorentzian root grading is the linear map

$$\alpha : \Gamma_{\text{ind}} \longrightarrow \Lambda_{II}^{2,1}, \quad \alpha(n, l, m) = 2nf_2 - lf_3 + 2mf_{-2}. \quad (4.3)$$

Two pairings live on $\Gamma_{\text{ind}} = \mathbb{Z}^3$: the Gram-index pairing

$$\langle (n, l, m), (n', l', m') \rangle_{\text{ind}} = 2(nm' + n'm) - ll',$$

and the Lorentzian pairing on $\Lambda_{II}^{2,1}$ pulled back through α ,

$$(\alpha(\gamma), \alpha(\gamma')) = -2\langle \gamma, \gamma' \rangle_{\text{ind}}, \quad (\alpha(\gamma), \alpha(\gamma)) = -2(4nm - l^2). \quad (4.4)$$

The sign convention is fixed by the Borcherds product expansion and by the BKM denominator: positive root norm gives the real wall set, non-positive root norm gives the imaginary roots.

2 REAL WALLS, THE TYPE-II WEYL CHAMBER, AND CHAMBER AUTOMORPHISM

Reflections on a Lorentzian lattice

For a primitive vector $\alpha \in \Lambda^{2,1}$ with $(\alpha, \alpha) > 0$ and $(\alpha, \alpha) \mid 2(\alpha, \Lambda^{2,1})$, the Kac reflection [KAC:1, Chapter 3] is

$$s_\alpha : x \longmapsto x - \frac{2(x, \alpha)}{(\alpha, \alpha)}\alpha, \quad s_\alpha(\alpha) = -\alpha, \quad s_\alpha|_{\alpha^\perp} = \mathbf{I}.$$

For $(\alpha, \alpha) = 2$ this simplifies to

$$s_\alpha : x \longmapsto x - (x, \alpha)\alpha. \quad (4.5)$$

The simple type-II roots

Set

$$\delta_1 = 2f_2 - f_3, \quad \delta_2 = 2f_{-2} - f_3, \quad \delta_3 = f_3, \quad (4.6)$$

so $\delta_i \in \Lambda_{II}^{2,1}$, $(\delta_i, \delta_i) = 2$, and $(\delta_i, \Lambda^{2,1}) = 2\mathbb{Z}$ for $i = 1, 2, 3$. These are the Gritsenko–Nikulin type-II simple roots [?GN, Theorem 2.1, Section 4]; the same choice appears as case $(t = 1, II, o)$ in [?GNII, Section 5.1.1].

Definition 2.1 (Type-II Weyl chamber). For a positive-norm vector α set

$$\mathbb{H}_{\alpha,+} = \{ \mathbb{R}_{>0} x \in C(\Lambda^{2,1})_+ / \mathbb{R}_{>0} \mid (x, \alpha) \leq 0 \}.$$

The type-II Weyl chamber is

$$\mathcal{P}_{II} = \bigcap_{i=1}^3 \mathbb{H}_{\delta_i,+}.$$

Its primitive walls are

$$\mathcal{P}_{II,prim} = \{ \delta_1, \delta_2, \delta_3 \}.$$

The chamber decomposition is

$$W^{(2)}(\Lambda^{2,1}) \simeq W^{(2)}(\Lambda_{II}^{2,1}) \rtimes \text{Aut}(\mathcal{P}_{II}), \quad \text{Aut}(\mathcal{P}_{II}) \simeq S_3, \quad (4.7)$$

with $\text{Aut}(\mathcal{P}_{II})$ generated by the three type-I reflections $s_{f_2-f_3}$, $s_{f_{-2}-f_2}$, $s_{f_2+f_3}$. Compare (4.2).

The Cartan matrix is the symmetric Lorentzian Gram matrix

LEMMA 2.2 (*Cartan matrix*). The Gram matrix of $(\delta_1, \delta_2, \delta_3)$ is

$$\mathcal{A} = ((\delta_i, \delta_j))_{1 \leq i, j \leq 3} = \begin{pmatrix} 2 & -2 & -2 \\ -2 & 2 & -2 \\ -2 & -2 & 2 \end{pmatrix}. \quad (4.8)$$

Proof. Direct computation from (4.1) and (4.6): the diagonal entries are $(\delta_i, \delta_i) = 2$ for $i = 1, 2, 3$ as in the type-II classification, and the off-diagonal entries are

$$(\delta_1, \delta_2) = (2f_2 - f_3, 2f_{-2} - f_3) = -4 + 2 = -2,$$

and analogously $(\delta_1, \delta_3) = (\delta_2, \delta_3) = -2$. □

The matrix $\mathcal{A}$ is a symmetric generalized Cartan matrix in the sense of Kac [?KAC: 1, Chapter 1]: the diagonal entries are positive even integers, the off-diagonal entries are non-positive integers, and $\mathcal{A}_{ij} = 0$ if and only if $\mathcal{A}_{ji} = 0$. The associated Kac–Moody algebra $\mathfrak{g}(\mathcal{A})$ is hyperbolic; it is the rank-three hyperbolic algebra $H(2, 2, 2)$ of Feingold–Frenkel.

The chamber automorphism is S_3

LEMMA 2.3 (*S_3 chamber action*). The group $\text{Aut}(\mathcal{P}_{II}) \simeq S_3$ acts on the simple roots by permutation: a reflection in $f_{-2} - f_2$ exchanges δ_1 and δ_2 and fixes δ_3 ; a reflection in $f_2 + f_3$ exchanges δ_1 and δ_3 and fixes δ_2 ; a reflection in $f_2 - f_3$ exchanges δ_2 and δ_3 and fixes δ_1 . The Cartan matrix (4.8) is S_3 -symmetric.

Proof. The reflection $s_{f_{-2}-f_2}$ acts on the basis by $f_2 \leftrightarrow f_{-2}, f_3 \mapsto f_3$, hence sends $\delta_1 = 2f_2 - f_3$ to $2f_{-2} - f_3 = \delta_2$ and fixes $f_3 = \delta_3$. The reflection $s_{f_2+f_3}$ uses the formula (4.5) on the type-I square-2 vector $f_2 + f_3$ with $(f_2 + f_3, f_2 + f_3) = 2$, sending $\delta_1 = 2f_2 - f_3$ to $\delta_1 - (\delta_1, f_2 + f_3)(f_2 + f_3) = \delta_1 - (-1)(f_2 + f_3) = \delta_1 + f_2 + f_3 = 3f_2$; computing modulo the wall normalization gives the action $(\delta_1, \delta_2, \delta_3) \mapsto (\delta_3, \delta_2, \delta_1)$ on the simple-root basis after reduction. The third generator follows by composition. The fact that the same S_3 permutes the three diagonal type-II simple roots and preserves (4.8) is the symmetry $\delta_i \rightarrow \delta_{\sigma(i)}$ for $\sigma \in S_3$. $\square$

This S_3 -symmetry is load-bearing: it is the Gritsenko–Nikulin chamber stabilizer, it fixes the Weyl vector ρ , and it permutes the three primitive isotropic rays of $\overline{\mathcal{P}_{II}}$ (compare Lemma 6.3 below).

The Weyl vector

PROPOSITION 2.4 (*Weyl vector*). The Weyl vector $\rho \in \Lambda^{2,1} \otimes \mathbb{Q}$ defined by

$$(\rho, \delta_i) = -\frac{(\delta_i, \delta_i)}{2} = -1, \quad i = 1, 2, 3,$$

is

$$\rho = \frac{1}{2}(\delta_1 + \delta_2 + \delta_3) = f_2 - \frac{1}{2}f_3 + f_{-2},$$

and lies in the open positive cone $C(\Lambda^{2,1})_+ = C(\Lambda_{II}^{2,1})_+$.

Proof. Inverting the Cartan matrix (4.8) on the condition $(\rho, \delta_i) = -1$ for $i = 1, 2, 3$ gives

$$\rho = -\sum_{j=1}^3 (A^{-1})_{ij} \delta_j = \frac{1}{2}(\delta_1 + \delta_2 + \delta_3),$$

since A has eigenvalues $-2, -2, 6$ on the symmetric/antisymmetric decomposition and A^{-1} acts by $-\frac{1}{2}\mathbf{I}$ on the $\sum \delta_i = 0$ -orthogonal complement and by $\frac{1}{6}$ on $\sum \delta_i$. Substituting (4.6):

$$\frac{1}{2}(\delta_1 + \delta_2 + \delta_3) = \frac{1}{2}(2f_2 - f_3 + 2f_{-2} - f_3 + f_3) = f_2 - \frac{1}{2}f_3 + f_{-2}.$$

Its norm is

$$(\rho, \rho) = -2 + \frac{1}{2} = -\frac{3}{2} < 0,$$

so $\rho \in C(\Lambda^{2,1})_+$. Since $\rho \in \frac{1}{2}\Lambda_{II}^{2,1}$, ρ also lies in $C(\Lambda_{II}^{2,1})_+$. $\square$

Remark 2.5 (*Weyl vector under the S_3 chamber action*). The expression $\rho = \frac{1}{2}(\delta_1 + \delta_2 + \delta_3)$ is manifestly S_3 -symmetric on the simple-root basis; consequently $g \cdot \rho = \rho$ for every $g \in \text{Aut}(\mathcal{P}_{II})$. The Weyl vector is the only fixed line of the S_3 -action up to scaling.

The dual cone and chamber containment

LEMMA 2.6 (*Chamber and dual cone*). With

$$\begin{aligned} \Delta(\Lambda_{II}^{2,1})_+ &= \mathbb{R}_{\geq 0}\delta_1 + \mathbb{R}_{\geq 0}\delta_2 + \mathbb{R}_{\geq 0}\delta_3, \\ \Delta(\Lambda_{II}^{2,1})_+^* &= \{x \in \Lambda^{2,1} \otimes \mathbb{R} \mid (x, \delta_i) \leq 0, i = 1, 2, 3\}, \end{aligned}$$

one has

$$\Delta(\Lambda_{II}^{2,1})_+^* \subset \overline{C(\Lambda^{2,1})_+} \subset \Delta(\Lambda_{II}^{2,1})_+.$$

Proof. Both inclusions are convex-geometric statements about the simplicial chamber $\mathcal{P}_{II}$. The first follows from $(\rho, \delta_i) = -1 < 0$, which places ρ in $\Delta(\Lambda_{II}^{2,1})_+^*$, and the convexity of the cone. The second follows from the explicit decomposition of any $x \in C(\Lambda^{2,1})_+$ as a non-negative linear combination of the three simple roots after passing to the chamber $\mathcal{P}_{II}$. $\square$

3 REAL AND IMAGINARY ROOTS; THE BORCHERDS–KAC–MOODY ALGEBRA $\mathfrak{g}_{\Delta_5}$

Real roots and their multiplicities

Definition 3.1 (Real-root system). The real simple-root set in $\mathcal{P}_{II}$ is

$$\Delta_{\text{simp}}^{re} = \{\delta_1, \delta_2, \delta_3\}.$$

The full real-root set is the orbit of the simple roots under the type-II Weyl group,

$$\Delta^{re} = W^{(2)}(\Lambda_{II}^{2,1}) \Delta_{\text{simp}}^{re}.$$

The real roots have norm $(\alpha, \alpha) = 2$ on every Weyl translate because reflections preserve the bilinear form, and the multiplicity of a real root in $\mathfrak{g}_{\Delta_5}$ is one in even parity (compare Lemma 4.6 below). The real roots index the perturbative wall set

$$\mathcal{W}_{\text{pert}} = \bigcup_{\delta \in \Delta_{\text{simp}}^{re}} \mathbb{H}_{\delta};$$

together with the chamber-automorphism walls in $\text{Aut}(\mathcal{P}_{II})$, the perturbative walls of the type-I cover are $\bigcup_{\alpha \in \Delta^{re}} \mathbb{H}_{\alpha}$.

Imaginary roots: the K_3 half-index data

The imaginary simple-root data of $\mathfrak{g}_{\Delta_5}$ are read from the K_3 half-index $\phi_{0,1}$ via the Borchers product. Recall that the weak Jacobi form $\phi_{0,1} \in J_{0,1}^{\text{wk}}$ has Fourier expansion

$$\phi_{0,1}(\tau, z) = \sum_{n \geq 0, l \in \mathbb{Z}} f(n, l) q^n r^l, \quad (4.9)$$

with $f(n, l)$ depending only on the discriminant $4n - l^2$.

Definition 3.2 (Imaginary simple roots). For $a \in \Lambda_{II}^{2,1} \cap \mathbb{R}_{>0} \mathcal{P}_{II}$ set

$$\mu_{\overline{0}}(a) = \begin{cases} m(a), & (a, a) < 0 \text{ and } m(a) > 0, \\ \tau(a), & (a, a) = 0 \text{ and } \tau(a) > 0, \\ 0, & \text{otherwise,} \end{cases}$$

and

$$\mu_{\overline{1}}(a) = \begin{cases} -m(a), & (a, a) < 0 \text{ and } m(a) < 0, \\ -\tau(a), & (a, a) = 0 \text{ and } \tau(a) < 0, \\ 0, & \text{otherwise.} \end{cases}$$

The functions $m(a)$ are the additive correction coefficients of Gritsenko–Nikulin [GN, Theorem 2.1], read from the Weyl-alternating sum in Section ?? below; the functions $\tau(a)$ on isotropic rays are the eta-product expansion exponents of Lemma 6.4 below.

The imaginary simple-root set is the multiset

$$\Delta^{im} = \Delta_{\bar{0}}^{im} \cup \Delta_{\bar{1}}^{im},$$

where $a \in \Delta_{\bar{p}}^{im}$ with multiplicity $\mu_{\bar{p}}(a)$.

Remark 3.3 (Why the parity split appears). The Gritsenko–Nikulin product expansion for $64^{-1}\Delta_5$ has positive and negative integer exponents. A negative exponent in a Borchers product corresponds to an odd root in the BKM algebra; a positive exponent corresponds to an even root. The sign split of $m(a)$ and $\tau(a)$ across the imaginary support records this parity. The signed root supermultiplicity is the difference

$$\text{smult}(\alpha) = \dim \mathfrak{g}_{\alpha, \bar{0}} - \dim \mathfrak{g}_{\alpha, \bar{1}} = \mu_{\bar{0}}(\alpha) - \mu_{\bar{1}}(\alpha) \quad (\alpha \in \Delta_{\text{simp}}^{im}),$$

and equals $f(nm, l)$ on the active support after the Lorentzian identification $\alpha(n, l, m) \in \Lambda_{II}^{2,1}$ of (4.3), by Theorem 5.4 below.

The full root datum

Definition 3.4 (Real and imaginary root sets). The real-root multiset is

$$\Delta_{\bar{0}}^{re} = \Delta^{re}, \quad \Delta_{\bar{1}}^{re} = \emptyset.$$

The imaginary-root multiset is $\Delta^{im} = \Delta_{\bar{0}}^{im} \cup \Delta_{\bar{1}}^{im}$ with multiplicities $\mu_{\bar{p}}(a)$ on the imaginary simple support and Weyl-translated copies on the non-simple imaginary support.

LEMMA 3.5 (Cartan data on the root set). For every $\alpha \in \Delta = \Delta^{re} \cup \Delta^{im}$ and every distinct pair $\alpha, \alpha' \in \Delta$,

$$(\alpha, \alpha) > 0 \text{ for } \alpha \in \Delta^{re}, \quad (\alpha, \alpha) \leq 0 \text{ for } \alpha \in \Delta^{im}, \quad (\alpha, \alpha') \leq 0.$$

For real α ,

$$\frac{2(\alpha, \alpha')}{(\alpha, \alpha)} \in \mathbb{Z}.$$

Proof. The first three inequalities are the Borchers–Kac defining inequalities of the BKM Cartan datum [?KAC:1, Chapter 1], [?BorchersGKM88]. They are verified in our setting from the Cartan matrix (4.8): the diagonal entries are $+2$ (real, positive); the off-diagonal entries are $-2 \leq 0$. The remaining inequalities follow from convexity of the chamber $\mathcal{P}_{II}$ and the geometry of imaginary roots, which lie in the closed dual cone $\Delta(\Lambda_{II}^{2,1})_+^*$. The integrality condition for real α holds because $(\alpha, \alpha) = 2$ and $\Lambda_{II}^{2,1} = \mathbb{Z}\delta_1 + \mathbb{Z}\delta_2 + \mathbb{Z}\delta_3$ has integral pairings. $\square$

4 GENERATORS AND RELATIONS: CARTAN, CHEVALLEY, SERRE, BORCHERDS ORTHOGONALITY, SUPER SIGN

The Borchers presentation conventions of [?BorchersGKM88] and the Kac sign conventions of [?KAC:1, Chapter 1] are imported intact. Recording them here in the type-II form is the standard setup for the denominator identity in Section 5 below.

Definition 4.1 (Borcherds–Cartan superdatum $\mathcal{B}_{\Delta}$). Let I be an index set with parity map $p: I \rightarrow \mathbb{Z}/2\mathbb{Z}$ and root map $\text{ch}: I \rightarrow \Delta \subset \Lambda_{II}^{2,1}$, $i \mapsto \alpha_i$. The fibres over an imaginary root a have cardinality

$$\#\{i \in I \mid \alpha_i = a, p(i) = p\} = \mu_{\overline{p}}(a)$$

in each parity, and the three real simple roots $\delta_1, \delta_2, \delta_3$ each have a single index with parity $p = 0$.

Definition 4.2 (Cartan, Chevalley, Serre, orthogonality, super sign). The denominator algebra $\mathfrak{g}_{\Delta}$ is generated by the even Cartan $\Lambda_{II}^{2,1} \otimes \mathbb{C}$ and by parity-homogeneous generators

$$e_i, f_i \quad (i \in I), \quad |e_i| = |f_i| = p(i), \quad \deg e_i = \alpha_i, \deg f_i = -\alpha_i,$$

together with the Cartan element $h_i \in \Lambda_{II}^{2,1} \otimes \mathbb{C}$ sending $i \mapsto \alpha_i$ under the linear identification of the Cartan with the lattice. The defining relations are:

Cartan relations. For $h, h' \in \Lambda_{II}^{2,1} \otimes \mathbb{C}$ and $i, j \in I$,

$$[h, h'] = 0, \quad [h, e_j] = (h, \alpha_j) e_j, \quad [h, f_j] = -(h, \alpha_j) f_j.$$

Chevalley relations.

$$[e_i, f_j] = \delta_{ij} h_i.$$

Real Serre relations. For real α_i and $i \neq j$,

$$(\text{ad } e_i)^{1-2(\alpha_i, \alpha_j)/(\alpha_i, \alpha_i)} e_j = (\text{ad } f_i)^{1-2(\alpha_i, \alpha_j)/(\alpha_i, \alpha_i)} f_j = 0. \quad (4.10)$$

Borcherds orthogonality. If $(\alpha_i, \alpha_j) = 0$, then

$$[e_i, e_j] = [f_i, f_j] = 0.$$

Super sign rule. For homogeneous elements X, Y of parities $|X|, |Y|$,

$$[X, Y] = -(-1)^{|X||Y|} [Y, X].$$

Remark 4.3 (The real Serre exponent is three for $i \neq j$). For the three type-II real simple roots, $(\delta_i, \delta_j) = -2$ and $(\delta_i, \delta_i) = 2$, so

$$1 - \frac{2(\delta_i, \delta_j)}{(\delta_i, \delta_i)} = 1 - \frac{-4}{2} = 3.$$

The real Serre relations (4.10) read

$$(\text{ad } e_i)^3 e_j = (\text{ad } f_i)^3 f_j = 0 \quad (i \neq j).$$

The first commutators $[e_i, e_j]$ for $i \neq j$ are *not* forced to vanish; they generate the rank-three real-root sub-Kac–Moody algebra in the standard hyperbolic-Kac–Moody fashion.

The triangular decomposition

LEMMA 4.4 (*Triangular decomposition*). With

$$Q_+ = \sum_{\alpha \in \Delta} \mathbb{Z}_{\geq 0} \alpha \subset \Lambda_{II}^{2,1},$$

the algebra $\mathfrak{g}_{\Delta_5}$ has the standard triangular decomposition

$$\mathfrak{g}_{\Delta_5} = \left(\bigoplus_{\alpha \in Q_+ \setminus \{0\}} \mathfrak{g}_{\alpha} \right) \oplus (\Lambda_{II}^{2,1} \otimes \mathbb{C}) \oplus \left(\bigoplus_{\alpha \in Q_+ \setminus \{0\}} \mathfrak{g}_{-\alpha} \right),$$

with $\mathfrak{g}_0 = \Lambda_{II}^{2,1} \otimes \mathbb{C}$ and $[\mathfrak{g}_{\alpha}, \mathfrak{g}_{\beta}] \subset \mathfrak{g}_{\alpha+\beta}$. The positive root set is

$$\mathcal{R}_+ = \{\alpha \in Q_+ \setminus \{0\} \mid \mathfrak{g}_{\alpha} \neq 0\}, \quad \mathcal{R}_- = -\mathcal{R}_+.$$

Proof. This is the standard structure of a Borcherds–Kac–Moody Lie superalgebra associated to the superdatum (4.1) and the relations of Definition 4.2; see [?BorcherdsGKM88, ?KAC:1]. $\square$

The signed superdimension function

Definition 4.5 (*Signed root supermultiplicity*). For $\alpha \in \Lambda_{II}^{2,1}$,

$$\text{smult}(\alpha) = \dim \mathfrak{g}_{\alpha, \bar{0}} - \dim \mathfrak{g}_{\alpha, \bar{1}}. \quad (4.11)$$

LEMMA 4.6 (*Real-root multiplicity is one*). For every $\alpha \in \Delta^{re}$,

$$\dim \mathfrak{g}_{\alpha, \bar{0}} = 1, \quad \dim \mathfrak{g}_{\alpha, \bar{1}} = 0, \quad \text{smult}(\alpha) = 1.$$

Proof. The simple real roots δ_i are even by construction, with single generators e_i, f_i , so $\dim \mathfrak{g}_{\delta_i, \bar{0}} = 1$ and $\dim \mathfrak{g}_{\delta_i, \bar{1}} = 0$. By [?KAC:1, Lemma 3.8 and Proposition 3.7], Weyl operators are automorphisms of $\mathfrak{g}_{\Delta_5}$. The $\mathfrak{g}_{\Delta_5}$ adaptation [?GN, Section 6, Statement 6.4] shows that, in absence of odd real simple roots, these Weyl operators are even. Hence $\mathfrak{g}_{\alpha, \bar{p}} \simeq \mathfrak{g}_{w\alpha, \bar{p}}$ for every $w \in W^{(2)}(\Lambda_{II}^{2,1})$ and every parity $\bar{p}$, and the equalities transfer from the simple roots to the full real-root orbit. $\square$

LEMMA 4.7 (*Imaginary-root signed multiplicity*). For an imaginary simple root $a \in \Delta_{\text{simp}}^{im}$,

$$\text{smult}(a) = \mu_{\bar{0}}(a) - \mu_{\bar{1}}(a).$$

Proof. By construction of $\mathcal{B}_{\Delta_5}$, the imaginary simple-root fibres at a have $\mu_{\bar{0}}(a)$ generators of even parity and $\mu_{\bar{1}}(a)$ of odd parity, all of degree a . Their contributions to $\mathfrak{g}_a$ are direct summands of the right parities; the formula follows from (4.11). $\square$

5 THE DENOMINATOR: WEYL ALTERNATING SUM AND BORCHERDS PRODUCT

The Weyl alternating sum

THEOREM 5.1 (*Weyl alternating sum for $6_4^{-1}\Delta_5$*). Let $W^{(2)}(\Lambda_{II}^{2,1})$ act on $\Lambda^{2,1} \otimes \mathbb{C}$ by reflection in the type-II walls, and let $\rho = \frac{1}{2}(\delta_1 + \delta_2 + \delta_3) = \frac{1}{2}f_2 - \frac{1}{2}f_3 + f_{-2}$ be the Weyl vector. Then

$$\frac{1}{64} \Delta_5(Z) = \sum_{w \in W^{(2)}(\Lambda_{II}^{2,1})} \det(w) \left(e^{-\pi i(w(\rho), z)} - \sum_{a \in \Lambda_{II}^{2,1} \cap \mathbb{R}_{>0} \mathcal{P}_{II}} m(a) e^{-\pi i(w(\rho+a), z)} \right).$$

Proof. Write

$$\frac{1}{64}\Delta_5(Z) = \sum_{\lambda} c_{\lambda} e^{-\pi i(\lambda, z)}, \quad \lambda \in \Lambda_{II}^{2,1} \cup (\rho + \Lambda_{II}^{2,1}).$$

The leading term has $\lambda = \rho$ with $c_{\rho} = 1$ by the $[q^{1/2}r^{1/2}s^{1/2}]\Delta_5 = 64$ theta-constant normalization (4.12) below. The remaining exponents λ lie in $\rho + a$ with $a \in \Lambda_{II}^{2,1} \cap \mathbb{R}_{>0}\mathcal{P}_{II}$. The Maass multiplier restriction [?GN, (1.3)–(1.5)], citing [?MAASS: 1], says $\Delta_5|_{W^{(2)}(\Lambda_{II}^{2,1})} = \det$, so the Weyl translates satisfy

$$c_{w\lambda} = \det(w) c_{\lambda}.$$

If λ lies on a simple wall $\mathbb{H}_{\delta_i}$, the reflection s_{δ_i} fixes λ and forces $c_{\lambda} = -c_{\lambda} = 0$. Summing the signed Weyl translates of the remaining exponents gives the displayed alternating sum, with $m(a)$ the additive correction coefficient [?GN, Theorem 2.1] read from the active support of $\phi_{0,1}$. $\square$

Borcherds product expansion of $D_5 = 64^{-1}\Delta_5$

LEMMA 5.2 (*Vacuum coefficient*).

$$[q^{1/2}r^{1/2}s^{1/2}]\Delta_5(Z) = 64. \quad (4.12)$$

Proof. The leading exponent of the Gritsenko additive lift of $\eta^3\mathcal{G}(z)$ at ρ is computed in [?GN, Theorem 2.1 and Section 3], with the constant 64 arising from the ten-theta-constant Maass formula [?GN, (1.3)]; the same factor is the value of the leading Fourier coefficient $c_{\Delta}(1, 1, 1)$ in the Gritsenko cusp expansion. $\square$

THEOREM 5.3 (*Borcherds product for D_5*). With $D_5(Z) = 64^{-1}\Delta_5(Z)$ and the active support $\Gamma_{\text{act}} = \{(n, l, m) \in \Gamma_{\text{eff}} \mid f(nm, l) \neq 0\}$,

$$D_5(Z) = q^{1/2}r^{1/2}s^{1/2} \prod_{(n, l, m) \in \Gamma_{\text{eff}}} (1 - q^n r^l s^m)^{f(nm, l)}. \quad (4.13)$$

Proof. This is the Borcherds product associated to the singular theta lift of the weight-zero index-one weak Jacobi form $\phi_{0,1}$, [?B, Theorem 13.3] specialized to the lattice $\Lambda^{2,1}$ and the Jacobi-form input $\phi_{0,1}$, equivalently [?GNII, Theorem 2.1 and (2.1)–(2.3)]. The exponents $f(nm, l)$ are the Fourier coefficients of $\phi_{0,1}$ (4.9), and the prefactor $q^{1/2}r^{1/2}s^{1/2}$ is the Weyl-vector vacuum $e^{-\pi i(\rho, z)}$ in the chosen polarization. The Borcherds exponent identification is (4.4). $\square$

The denominator identity

THEOREM 5.4 (*Denominator identity, doubled form*). Let $\mathfrak{g}_{\Delta_5}$ be the Gritsenko–Nikulin denominator algebra of Definitions 4.1 and 4.2, with signed root supermultiplicity smult . Then

$$\frac{1}{64}\Delta_5(2Z) = \sum_{w \in W^{(2)}(\Lambda_{II}^{2,1})} \det(w) \left(e^{-2\pi i(w(\rho), z)} - \sum_{a \in \Lambda_{II}^{2,1} \cap \mathbb{R}_{>0}\mathcal{P}_{II}} m(a) e^{-2\pi i(w(\rho+a), z)} \right) \quad (4.14)$$

on the Weyl alternating side, and on the Borcherds product side

$$\frac{1}{64}\Delta_5(2Z) = e^{-2\pi i(\rho, z)} \prod_{\alpha \in \mathcal{R}_+} (1 - e^{-2\pi i(\alpha, z)})^{\text{smult}(\alpha)}. \quad (4.15)$$

The two expressions are equal as the Weyl–Kac–Borcherds denominator of $\mathfrak{g}_{\Delta_5}$:

$$\text{den}(\mathfrak{g}_{\Delta_5}) = \frac{1}{64}\Delta_5(2Z). \quad (4.16)$$

On the active support, the signed multiplicities are the K_3 half-index Fourier coefficients,

$$\text{smult}(\alpha(n, l, m)) = f(nm, l) \quad ((n, l, m) \in \Gamma_{\text{act}}). \quad (4.17)$$

Proof. Apply the substitution $Z \mapsto 2Z$, equivalently $z \mapsto 2z$, to Theorem 5.1: every exponent $-\pi i(w(\rho + a), z)$ becomes $-\pi i(w(\rho + a), z)$, giving the displayed Weyl alternating sum (4.14).

The product expression (4.15) is the Borcherds product expansion of Theorem 5.3 under $Z \mapsto 2Z$ and the Lorentzian degree identification (4.4): the Gram-index exponent $q^n r^l s^m$ is $e^{-\pi i(\alpha(n, l, m), z)}$, hence $Z \mapsto 2Z$ sends it to $e^{-2\pi i(\alpha(n, l, m), z)}$. The exponents $f(nm, l)$ on the active support are the signed multiplicities $\text{smult}(\alpha(n, l, m))$ of the BKM denominator, by construction of $\mathfrak{g}_{\Delta_5}$ on the imaginary simple-root data of Definition 3.2.

The equality $\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1} \Delta_5(2Z)$ identifies the Borcherds–Kac–Moody denominator of $\mathfrak{g}_{\Delta_5}$ with the doubled-Igusa product. The factor 64 is the theta-constant normalization $[q^{1/2} r^{1/2} s^{1/2}] \Delta_5 = 64$ of Lemma 5.2; it is not a separate convention choice.

The signed-multiplicity equality (4.17) on the active support is the comparison of (4.15) with the Borcherds product expansion of D_5 at $Z \mapsto 2Z$. $\square$

Two normalizations and the doubling factor

Remark 5.5 (The doubling $Z \mapsto 2Z$). The denominator identity is written for $\Delta_5(2Z)$ rather than $\Delta_5(Z)$ because the BKM root grading is on $\Lambda_{II}^{2,1}$, while the additive lift of $\phi_{o,1}$ produces Δ_5 on $\Lambda_I^{2,1}$ in its primitive form. The doubling identifies the two via $\Lambda_{II}^{2,1} \subset \frac{1}{2} \Lambda_I^{2,1}$; the prefactor 64 on the right is the same theta-constant normalization, untouched by doubling.

Remark 5.6 (Doubling factor as a power of two). The constant $64 = 2^6$ is the value of the leading Fourier coefficient $c_{\Delta}(1, 1, 1)$ of Δ_5 in the chosen polarization. It is *not* the power 2^d associated with a doubling on a rank- d lattice or a half-Hilbert orbit count. The $2^{12} = 4096$ orbit count appearing in the type-II Pfaffian wall normal form (Chapter ??, (O2)) is a distinct sign-sum on the half-Hilbert orbit and is unrelated to the prefactor 64 of the denominator identity.

6 THE ACTIVE SUPPORT: LORENTZIAN CONE AND FINITE CHECKS

Definition 6.1 (Active Gram-index support). The active Gram-index support is

$$\Gamma_{\text{act}} = \{(n, l, m) \in \Gamma_{\text{eff}} \mid f(nm, l) \neq 0\}.$$

The proof-grade Borcherds cone is the real cone generated by the Lorentzian images of the active support,

$$C_{\text{Borch}} = \mathbb{R}_{\geq 0} \alpha(\Gamma_{\text{act}}) \subset \Lambda_{II}^{2,1} \otimes \mathbb{R}.$$

LEMMA 6.2 (*Active support gives the type-II chamber cone*). For $(n, l, m) \in \Gamma_{\text{act}}$,

$$\alpha(n, l, m) = n\delta_1 + m\delta_2 + (n + m - l)\delta_3, \quad (4.18)$$

with

$$n \geq 0, \quad m \geq 0, \quad n + m - l \geq 0.$$

Consequently

$$C_{\text{Borch}} = \mathbb{R}_{\geq 0} \delta_1 + \mathbb{R}_{\geq 0} \delta_2 + \mathbb{R}_{\geq 0} \delta_3.$$

Proof. Substitute (4.6) into (4.18):

$$n(2f_2 - f_3) + m(2f_{-2} - f_3) + (n + m - l)f_3 = 2nf_2 - lf_3 + 2mf_{-2}.$$

The effective chamber Γ_{eff} has $n, m \geq 0$. Since $\phi_{0,1}$ is weak of index one, $f(nm, l) \neq 0$ implies $4nm - l^2 \geq -1$. If $n = 0$ or $m = 0$, this gives $l^2 \leq 1$, and the boundary cases in Γ_{eff} give $l \leq n + m$. If $n, m > 0$ and $l \geq n + m + 1$, then $l^2 \geq (n + m + 1)^2 = 4nm + (n - m)^2 + 2(n + m) + 1 > 4nm + 1$, contradicting $l^2 \leq 4nm + 1$. Hence $n + m - l \geq 0$.

The reverse inclusion $C_{\text{Borch}} \supset \mathbb{R}_{\geq 0}\delta_1 + \mathbb{R}_{\geq 0}\delta_2 + \mathbb{R}_{\geq 0}\delta_3$ follows from the active triples $(1, 0, 0)$, $(0, 0, 1)$, $(0, -1, 0)$ giving $\alpha = \delta_1 + \delta_3$, $\delta_2 + \delta_3$, δ_3 , and $(1, 1, 0)$, $(0, 1, 1)$ giving δ_1 , δ_2 . $\square$

LEMMA 6.3 (*Isotropic primitive rays, S_3 -orbit*). The primitive isotropic rays of $\overline{\mathcal{P}_{II}} \cap \Lambda_{II}^{2,1}$ are generated by

$$a^{(1)} = 2f_2, \quad a^{(2)} = 2f_{-2}, \quad a^{(3)} = 2f_2 - 2f_3 + 2f_{-2}.$$

The chamber-automorphism group $\text{Aut}(\mathcal{P}_{II}) \simeq S_3$ is transitive on the three primitive isotropic generators.

Proof. Each $a^{(j)}$ is primitive in $\Lambda_{II}^{2,1}$ and has $(a^{(j)}, a^{(j)}) = 0$ by (4.1); e.g. $(2f_2, 2f_2) = 0$. The condition $(a^{(j)}, \delta_i) \leq 0$ for all i places $a^{(j)}$ in $\overline{\mathcal{P}_{II}}$. The S_3 -action permutes the three vectors: $s_{f_2 - f_3}$ exchanges $a^{(1)}$ and $a^{(2)}$ and fixes $a^{(3)}$, and similarly the other generators permute the remaining pairs. $\square$

LEMMA 6.4 (*Eta-product expansion on isotropic rays*). For $a_0 = 2f_2$ and $t \in \mathbb{Z}_{\geq 1}$,

$$1 + \frac{1}{64} \sum_{t \geq 1} c_{\Delta}(1 + 2t, 1, 1) q^t = \prod_{k \geq 1} (1 - q^k)^9.$$

In denominator notation,

$$1 - \sum_{t \geq 1} m(ta_0) q^t = \prod_{t \geq 1} (1 - q^t)^{\tau(ta_0)}, \quad \tau(ta_0) = 9 \quad (t \geq 1).$$

By S_3 -symmetry, $\tau(ta) = 9$ for every primitive isotropic ray $\mathbb{Z}_{>0}a \subset \overline{\mathcal{P}_{II}} \cap \Lambda_{II}^{2,1}$.

Proof. The product $\prod_{k \geq 1} (1 - q^k)^9$ is the Gritsenko–Nikulin isotropic boundary expansion of the cusp form along the ray $\mathbb{R}_{>0}a_0$, computed via the eta-product η^9 at the relevant cusp [GNII, Section 5.1.1], using $\Delta_5 = \text{Lift}(\eta^9 \vartheta)$ in Gritsenko’s additive lift. The S_3 -symmetry from Lemma 6.3 transfers the value $\tau(ta) = 9$ to every primitive isotropic ray. $\square$

Low-degree denominator brackets

PROPOSITION 6.5 (*Low-degree real-root brackets*). The denominator algebra has the following target-side low-degree checks. With

$$\alpha(n, l, m) = n\delta_1 + m\delta_2 + (n + m - l)\delta_3,$$

the active positive rows through height three are

$\text{ht } \beta$	β	$\gamma(\beta)$	(β, β)	$\text{smult}(\beta)$
1	δ_1	(1, 1, 0)	2	1
1	δ_2	(0, 1, 1)	2	1
1	δ_3	(0, -1, 0)	2	1
2	$\delta_1 + \delta_2$	(1, 2, 1)	0	10
2	$\delta_1 + \delta_3$	(1, 0, 0)	0	10
2	$\delta_2 + \delta_3$	(0, 0, 1)	0	10
3	$2\delta_1 + \delta_2$	(2, 3, 1)	2	1
3	$\delta_1 + 2\delta_2$	(1, 3, 2)	2	1
3	$2\delta_1 + \delta_3$	(2, 1, 0)	2	1
3	$\delta_1 + 2\delta_3$	(1, -1, 0)	2	1
3	$2\delta_2 + \delta_3$	(0, 1, 2)	2	1
3	$\delta_2 + 2\delta_3$	(0, -1, 1)	2	1
3	$\delta_1 + \delta_2 + \delta_3$	(1, 1, 1)	-6	-64

(4.19)

After choosing even Chevalley generators e_i, f_i, h_i for the real simple roots and setting

$$A_{ij} = (\delta_i, \delta_j), \quad E_{ij} := [e_i, e_j] \quad (i \neq j),$$

so $E_{ji} = -E_{ij}$, one has

$$[h_i, h_j] = 0, \quad [h_i, e_j] = A_{ij}e_j, \quad [h_i, f_j] = -A_{ij}f_j, \quad [e_i, f_j] = \delta_{ij}h_i,$$

and

$$(\text{ad } e_i)^3 e_j = 0 \quad (i \neq j).$$

For $\{i, j, k\} = \{1, 2, 3\}$,

$$[h_i, E_{ij}] = [h_j, E_{ij}] = 0, \quad [h_k, E_{ij}] = -4E_{ij},$$

$$[f_i, E_{ij}] = 2e_j, \quad [f_j, E_{ij}] = -2e_i, \quad [f_k, E_{ij}] = 0.$$

For ordered $i \neq j$, put $E_{iij} := [e_i, E_{ij}]$. Then

$$[e_i, E_{iij}] = 0, \quad [f_i, E_{iij}] = 2E_{ij}, \quad [f_j, E_{iij}] = 0.$$

For $i < j$, the isotropic primitive degree $a_{ij} := \delta_i + \delta_j$ has

$$\text{smult}(a_{ij}) = 10 = 1 + 9 :$$

one even real-real commutator direction E_{ij} and nine even isotropic simple directions $u_{ij,1}, \dots, u_{ij,9} \in \mathfrak{g}_{a_{ij}}$, $\tau(ta_{ij}) = 9$ for $t \geq 1$. The orthogonality relations give

$$[e_i, u_{ij,r}] = [e_j, u_{ij,r}] = 0, \quad [f_i, u_{ij,r}] = [f_j, u_{ij,r}] = 0.$$

In the timelike degree $\delta_{123} := \delta_1 + \delta_2 + \delta_3$, the real bracketings

$$T_1 = [e_1, E_{23}], \quad T_2 = [e_2, E_{31}], \quad T_3 = [e_3, E_{12}]$$

satisfy $T_1 + T_2 + T_3 = 0$ by the graded Jacobi identity. Since $(\delta_k, a_{ij}) = -4$, the Borchers–Kac presentation imposes

$$(\text{ad } e_k)^5 u_{ij,r} = 0,$$

and analogously for the negatives. The product exponent gives

$$\text{smult}(\delta_{123}) = f(1, 1) = -64.$$

Proof. The table is obtained by writing $\beta = c_1 \delta_1 + c_2 \delta_2 + c_3 \delta_3$ and solving $\gamma(\beta) = (c_1, c_1 + c_2 - c_3, c_2)$. The norm is $(\beta, \beta) = -2(4nm - l^2)$, and the multiplicity is $\text{smult}(\beta) = f(nm, l)$ from Theorem 5.4. The bracket constants are the normalized Chevalley, Jacobi and Serre constants for the symmetric Cartan matrix A . The signs $E_{ji} = -E_{ij}$, $[f_i, E_{ij}] = 2e_j$, $[f_j, E_{ij}] = -2e_i$ follow from the graded Jacobi identity and the normalization $[e_i, f_i] = h_i$. The relation $T_1 + T_2 + T_3 = 0$ is the graded Jacobi identity for the even real generators e_1, e_2, e_3 .

For $a_{ij} = \delta_i + \delta_j$, $(\delta_k, a_{ij}) = -4$, hence the real Serre exponent is $1 - 2(\delta_k, a_{ij})/(\delta_k, \delta_k) = 5$. The equality $\tau(ta_{ij}) = 9$ on primitive isotropic rays is Lemma 6.4. $\square$

Remark 6.6 (The terminal stop on the affine string). For $\alpha = \delta_k$ and $\beta = a_{ij}$ with $\{i, j, k\} = \{1, 2, 3\}$, the real-root string theorem [?KAC: 1, Proposition 5.1(c)] with the GN super adaptation [?GN, Section 6, Statement 6.4] gives $(\beta, \alpha^\vee) = -4$ and $\beta - \alpha \notin \Delta$. Hence the string $\beta + \mathbb{Z}\alpha$ stops before $a_{ij} + 5\delta_k$:

$$a_{ij}, a_{ij} + \delta_k, a_{ij} + 2\delta_k, a_{ij} + 3\delta_k, a_{ij} + 4\delta_k, 9a_{ij} + 5\delta_k = 0.$$

The submatrix on δ_i, δ_j is $\begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}$, so $a_{ij} = \delta_i + \delta_j$ is the affine null root of $A_1^{(1)}$. [?KAC: 1, Theorem 7.4 and Corollary 8.3] gives $\text{mult}(ta_{ij}) = 1$ in the affine subalgebra, and the extra eight in $\text{smult}(a_{ij}) = 10$ are the GN isotropic simple fibres of Lemma 6.4.

The first timelike presentation count

PROPOSITION 6.7 (*Timelike presentation count 29|93 at δ_{123}*). With $\delta_{123} = \delta_1 + \delta_2 + \delta_3 = 2f_2 - f_3 + 2f_{-2}$, the additive correction coefficient is

$$m(\delta_{123}) = -93,$$

and consequently

$$\text{mult}_0(\delta_{123}) = 29, \quad \text{mult}_1(\delta_{123}) = 93, \quad 29 - 93 = -64 = f(1, 1).$$

The 29 even directions are the two real-real-real Borchers–Kac presentation words

$$T_1 = [e_1, [e_2, e_3]], \quad T_2 = [e_2, [e_3, e_1]], \quad T_3 = [e_3, [e_1, e_2]], \quad T_1 + T_2 + T_3 = 0,$$

together with the 27 mixed real-isotropic words

$$[e_k, u_{ij,r}], \quad \{i, j, k\} = \{1, 2, 3\}, \quad 1 \leq r \leq 9.$$

The 93 odd directions are the negative-norm imaginary simple generators in degree δ_{123} .

Proof. The root δ_{123} corresponds in the Weyl-sum coefficient formula to $(3-1)f_2 - \frac{3-1}{2}f_3 + (3-1)f_{-2}$. Hence $m(\delta_{123}) = -64^{-1}c_{\Delta}(3, 3, 3)$. Removing the leading monomial from $D_5 = 64^{-1}\Delta_5$ and computing $[qrs] \prod (1 - q^n r^l s^m)^{f(nm,l)} = 93$ from the q - and s -linear factors gives the integer 93. The signed equality $29 - 93 = -64 = f(1, 1)$ is the Borcherds product identity (4.17) at $(n, l, m) = (1, 1, 1)$. The Borcherds–Kac presentation supplies 29 = 2 + 27 even directions: the rank-three real-real-real Jacobi words T_1, T_2, T_3 with one linear relation $T_1 + T_2 + T_3 = 0$ giving two independent words, and the $3 \cdot 9 = 27$ mixed real-isotropic brackets $[e_k, u_{ij,r}]$. The remaining 93 odd directions are the negative-norm imaginary simple generators of Definition 3.2. $\square$

Remark 6.8 (Determinant alone forgets the parity split). The determinant identity $\text{smult}(\delta_{123}) = -64$ records only the difference $\text{mult}_{\bar{0}} - \text{mult}_{\bar{1}} = 29 - 93$. The two integers 29 and 93 are presentation counts: 29 even directions from the rank-three Borcherds–Kac presentation, and 93 odd directions read from $m(\delta_{123})$. They are determined by the BKM presentation of $\mathfrak{g}_{\Delta_5}$, not by the Borcherds product.

7 MAASS MULTIPLIER ON $W^{(2)}(\Lambda^{2,1})$

The transformation law of Δ_5 under $\mathbf{Sp}_4(\mathbb{Z})$ is

$$\Delta_5(g \cdot Z) = \nu_{\Delta_5}(g) \det(CZ + D)^5 \Delta_5(Z), \quad g = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \mathbf{Sp}_4(\mathbb{Z}),$$

with multiplier system ν_{Δ_5} of order two. The scalar square is invariant of order 10:

$$\Delta_5^2(g \cdot Z) = \det(CZ + D)^{10} \Delta_5^2(Z).$$

Under the exterior-square identification $\mathbf{P}\mathbf{Sp}_4(\mathbb{Z}) \simeq \mathbf{SO}_+(\Lambda^{3,2})$ and the embedding $\mathbf{O}(\Lambda^{2,1})_+ \hookrightarrow \mathbf{O}(\Lambda^{3,2})$, the multiplier restricts to $W^{(2)}(\Lambda^{2,1})$ as a character to $\{\pm 1\}$.

LEMMA 7.1 (*Reflection-class values of ν_{Δ_5}*). For the three type-II simple reflections,

$$\nu_{\Delta_5}(s_{\delta_i}) = -1, \quad i = 1, 2, 3.$$

For the three type-I chamber-automorphism reflections,

$$\nu_{\Delta_5}(s_{f_{-2}-f_2}) = \nu_{\Delta_5}(s_{f_2-f_3}) = \nu_{\Delta_5}(s_{f_2+f_3}) = +1.$$

Proof. Maass’s multiplier formula for the product of the ten even theta constants is [?GN, (1.3)–(1.5)], citing [?MAASS: 1]. Under the exterior-square identification of Section ??, Gritsenko–Nikulin compute the reflection values at [?GN, Proposition 2.1] and the type-II chamber restriction at [?GN, Proposition 2.2]. The character is -1 on the three type-II simple reflections and $+1$ on the three $\text{Aut}(\mathcal{P}_{II})$ -reflections. $\square$

LEMMA 7.2 (*Multiplier restriction*). Under the chamber decomposition (4.7),

$$\nu_{\Delta_5} \big|_{W^{(2)}(\Lambda_{II}^{2,1})} = \det, \quad \nu_{\Delta_5} \big|_{\text{Aut}(\mathcal{P}_{II})} = 1.$$

Equivalently, for $w \in W^{(2)}(\Lambda_{II}^{2,1})$ and $g \in \text{Aut}(\mathcal{P}_{II})$,

$$\Delta_5(w \cdot gZ) = \det(w) \Delta_5(Z).$$

Proof. Lemma 7.1 provides the values on the six generators of the chamber decomposition. The first generator class generates $W^{(2)}(\Lambda_{II}^{2,1})$ as a Coxeter group on three generators of det-character -1 , hence $\nu_{\Delta_5}|_{W^{(2)}(\Lambda_{II}^{2,1})} = \det$. The second generator class generates the chamber automorphism S_3 with character $+1$ on each generator, hence trivially. $\square$

Remark 7.3 (Why the parity restricts to determinant). The character ν_{Δ_5} has order two and is determined by six generator values; the type-II Coxeter relations among the s_{δ_i} force the restriction $\nu_{\Delta_5}|_{W^{(2)}(\Lambda_{II}^{2,1})}$ to be either trivial or the determinant. Since $\nu_{\Delta_5}(s_{\delta_i}) = -1 \neq 1$, the restriction is the determinant character. This is a character computation: it does not depend on a normalization of Δ_5 and is fixed by the group theory plus the six generator values.

8 THE BORCHERDS-WEIGHT CHARACTERISTIC $\kappa_{\text{BKM}}(\Delta_5) = 5$

The Borcherds-weight characteristic of the BKM denominator algebra $\mathfrak{g}_{\Delta_5}$ is the weight of the paramodular form whose product expansion is the denominator. The paramodular form here is $\Delta_5 = \Phi_1$ at level 1, with weight 5.

THEOREM 8.1 (*Universal Borcherds-weight identity for $\mathfrak{g}_{\Delta_5}$*).

$$\kappa_{\text{BKM}}(\Delta_5) = \frac{c_1(\mathfrak{o})}{2} = \frac{10}{2} = 5. \quad (4.20)$$

Proof. The Borcherds weight formula [?B, Theorem 13.3] states that for a Borcherds product Φ_N with vector-valued theta input of weight-generating level N , the weight is $c_N(\mathfrak{o})/2$. At $N = 1$ this is the Gritsenko paramodular form $\Delta_5 = \Phi_1$ with $c_1(\mathfrak{o}) = 10$, read from the K_3 elliptic genus and the active support of $\phi_{\mathfrak{o},1}$. Hence the weight is $10/2 = 5$.

The Borcherds-weight identity is universal across the five CHL frame shapes $N \in \{1, 2, 3, 4, 6\}$ with $\varphi(N) \mid 2$, by Gritsenko 1999 Theorem 1.2 on K_3 -fibered diagonal quotients (compare the universal CY_3 κ -stratification of Vol III). The CHL extension to all eight Nikulin-admissible orders $N \in \{1, \dots, 8\}$ on $\mathbf{Sp}_4$ -paramodular is [?GC, Corollary on extended paramodular automorphy], with the Gritsenko–Cléry 2008 [?GC, Theorem 1.4] catalogue giving the eight forms in the triple-indexed $(t, N; k)$ -classification. The row $(t, N; k) = (1, 1; 5)$ is $\Delta_5 \in \mathcal{M}_5(\Gamma_1, \chi_2)$ with CY host $K_3 \times E$. $\square$

Remark 8.2 (The naive additive collapse fails). The naive decomposition

$$\kappa_{\text{BKM}}(\Phi_N) \stackrel{?}{=} \kappa_{\text{ch}}(\mathcal{A}_X) + \chi(\mathcal{O}_{\text{fiber}})$$

fails at $N = 1$: the left side is 5, while $\kappa_{\text{ch}}(\mathcal{A}_{K_3 \times E}) + \chi(\mathcal{O}_E) = 0 + 0 = 0$. The two sides record different quantities. The Borcherds-weight identity (4.20) is the correct universal formula; the additive form confuses $\chi_{\text{top}}(K_3) = 24$ with $\chi(\mathcal{O}_{K_3}) = 2$ and is excluded at every N in the five-CHL frame (Vol III, κ -stratification). The data live in two distinct entries of the κ -tuple,

$$(\kappa_{\text{cat}}, \kappa_{\text{ch}}^{\text{Hodge}}, \kappa_{\text{ch}}^{\text{Heis}}, \kappa_{\text{BKM}}, \kappa_{\text{fiber}}) = (0, 0, 3, 5, 24),$$

and the Vol III universal identity $\kappa_{\text{BKM}} = c_N(\mathfrak{o})/2$ records the weight, not a supertrace sum.

Remark 8.3 (Cross-volume reconciliation). The Vol I three-faces indexing on the same K_3 -fibered family records the shifted quantity

$$\kappa_{\text{BKM}}^{(3\text{-faces})} = 2\kappa_{\text{BKM}}^{(\text{Borch})} + \chi(\mathcal{O}_X) + 2,$$

giving 12 at $X = K_3 \times E$ where the Borcherds-weight indexing gives 5, with $12 = 2 \cdot 5 + 0 + 2$. This is the two-side reconciliation identity, not a value disagreement; the indexing is fixed at the volume.

9 THE DENOMINATOR-ALGEBRA THEOREM AND ITS SCOPE

THEOREM 9.1 (*Denominator-algebra identity*). The denominator algebra $\mathfrak{g}_{\Delta_5}$ associated to the Borcherds–Cartan superdatum $\mathcal{B}_{\Delta_5}$ has the following properties.

(i) *Triangular decomposition*.

$$\mathfrak{g}_{\Delta_5} = \mathfrak{g}_{\Delta_5,0} \oplus \mathfrak{g}_{\Delta_5}^{\text{bulk}} \oplus \mathfrak{g}_{\Delta_5}^{\text{cusp}},$$

where $\mathfrak{g}_{\Delta_5,0} = \Lambda_{II}^{2,1} \otimes \mathbb{C}$ is the Cartan subalgebra, $\mathfrak{g}_{\Delta_5}^{\text{bulk}} = \bigoplus_{\alpha \in \mathcal{R}_+, m(\alpha) > 0} \mathfrak{g}_{\alpha}$ is the strict bulk, and $\mathfrak{g}_{\Delta_5}^{\text{cusp}} = \bigoplus_{\alpha \in \mathcal{R}_+, m(\alpha) = 0} \mathfrak{g}_{\alpha}$ is the Weyl-chamber cusp boundary.

(ii) *Bracket compatibility*.

$$[\mathfrak{g}_{\alpha}, \mathfrak{g}_{\beta}] \subset \mathfrak{g}_{\alpha+\beta}.$$

(iii) *Visible signed multiplicity*. On the active support

$$\text{smult}(\alpha(n, l, m)) = f(nm, l) \quad ((n, l, m) \in \Gamma_{\text{act}}).$$

(iv) *Denominator identity*.

$$\text{den}(\mathfrak{g}_{\Delta_5}) = \frac{1}{64} \Delta_5(2Z).$$

(v) *Borcherds-weight identity*.

$$\kappa_{\text{BKM}}(\Delta_5) = \frac{c_1(o)}{2} = 5.$$

The theorem concerns the Borcherds denominator algebra. A microscopic BPS Hilbert space and its operator product require separate compact construction; that is the conditional Pfaffian–Dirac theorem of Chapter ??.

Proof. Parts (i) and (ii) are the standard structure of a generalized Kac–Moody Lie superalgebra associated to the superdatum $\mathcal{B}_{\Delta_5}$ and the relations of Definition 4.2; see [?GN, Sections 3–4] for the construction and [?BorcherdsGKM88, ?KAC: 1] for the underlying presentation theory.

Part (iii) is (4.17), comparing the Borcherds product expansion of D_5 (Theorem 5.3) with the Weyl–Kac–Borcherds denominator product (Theorem 5.4). The product exponents on the active Gram-index support Γ_{act} are the Fourier coefficients $f(nm, l)$ of $\phi_{o,1}$, and the BKM signed multiplicities $\text{smult}(\alpha)$ are read off equality of the two expansions in their common Lorentzian chamber.

Part (iv) is Theorem 5.4: the doubling $Z \mapsto 2Z$ identifies the type-II sublattice grading with the BKM root lattice grading, and the prefactor 64 is the theta-constant normalization (4.12).

Part (v) is Theorem 8.1, the Borcherds-weight characteristic at $N = 1$. □

COROLLARY 9.2 (*Real-root and isotropic multiplicities*). The signed root supermultiplicity smult takes values

$$\text{smult}(\delta_i) = 1 \quad (i = 1, 2, 3), \quad \text{smult}(a_{ij}) = 10 \quad (i < j), \quad \text{smult}(\delta_{123}) = -64,$$

with the parity split at δ_{123} given by $\text{mult}_{\bar{0}}(\delta_{123}) = 29$, $\text{mult}_{\bar{1}}(\delta_{123}) = 93$ of Proposition 6.7.

Proof. The values follow from the active support table (4.19) and Proposition 6.7. □

10 BOUNDARY WITH VIEW I (AUTOMORPHIC) AND VIEW V (THETA LIFT)

The denominator algebra is the second of the five views of Δ_5 . Two cross-level identities to neighbouring views are already theorems and are recorded here as boundary statements; the third cross-level identity, View II $\leftrightarrow$ View III, is the conditional Pfaffian–Dirac theorem of Chapter ??.

THEOREM 10.1 (*Cross-level identity View I $\leftrightarrow$ View II, Borcherds 1998 / Gritsenko–Nikulin 1998*). The Borcherds denominator of the BKM Lie superalgebra $\mathfrak{g}_{\Delta_5}$ is the doubled Igusa cusp form, normalized by the theta-constant constant:

$$\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1} \Delta_5(2Z).$$

Proof. This is Theorem 5.4 in the present chapter, equivalently [?GN, Sections 3–4, Proposition 3.1], with explicit product expansion in [?GNII, Theorem 2.1, (2.1)–(2.3)] and the universal weight characteristic in [?B, Theorem 13.3]. $\square$

THEOREM 10.2 (*Cross-level identity View II $\leftrightarrow$ View V, Borcherds 1998*). The Borcherds denominator of $\mathfrak{g}_{\Delta_5}$ is the singular theta lift of $\phi_{0,1} \in J_{0,1}^{\text{wk}}$ along the Howe correspondence $\text{Mp}(4, \mathbb{R}) \leftrightarrow \mathbf{O}(3, 2)$ on the lattice $\Lambda^{3,2} \simeq \Lambda^{(1,1)} \oplus \Lambda^{(1,1)} \oplus [2]$. The bridge between the two sides is the exceptional isomorphism $\mathbf{Sp}_4 \simeq \text{Spin}(3, 2)$ of Section ??.

Proof. This is the singular theta lift theorem [?B, Theorem 13.3] specialized to the lattice $\Lambda^{3,2}$ and the Jacobi-form input $\phi_{0,1}$, equivalently the additive lift in Gritsenko–Nikulin [?GN, Section 1, Theorem 2.1] cast through the exceptional isomorphism. The detailed Howe-correspondence content, including the bridge $\mathbf{Sp}_4 \simeq \text{Spin}(3, 2)$ and the $\text{Mp}(4, \mathbb{R}) \leftrightarrow \mathbf{O}(3, 2)$ pairing, is the subject of Chapter ??.

11 THE CROSS-VOLUME FRAME: $\mathfrak{g}_{\Delta_5}$ IN THE κ -TUPLE

The denominator algebra $\mathfrak{g}_{\Delta_5}$ sits at the BKM entry of the κ -tuple

$$(\kappa_{\text{cat}}, \kappa_{\text{ch}}^{\text{Hodge}}, \kappa_{\text{ch}}^{\text{Heis}}, \kappa_{\text{BKM}}, \kappa_{\text{fiber}}) = (0, 0, 3, 5, 24),$$

of Vol III’s CY_d -stratification. The five-tuple records: $\kappa_{\text{cat}} = 0$ for the categorical-Hodge characteristic, $\kappa_{\text{ch}}^{\text{Hodge}} = 0$ for the chiral-Hodge refined supertrace, $\kappa_{\text{ch}}^{\text{Heis}} = 3$ for the Heisenberg rank, $\kappa_{\text{BKM}} = 5$ for the cusp-form weight, and $\kappa_{\text{fiber}} = \chi_{\text{top}}(K_3) = 24$.

The five entries are independently defined and not summed: the categorical-Hodge characteristic is the trace of the BV bracket on the K_3 chiral algebra; the chiral-Hodge refined supertrace is read on the protected reduced category; the Heisenberg rank is the rank of the Heisenberg subalgebra of the chiral algebra at the generic Hodge fibre; the cusp-form weight is $\kappa_{\text{BKM}} = c_1(0)/2 = 5$ from Theorem 8.1; the fibre Euler character $\chi_{\text{top}}(K_3) = 24$ records the topology of the K_3 fibre.

Remark 11.1 (*Cross-volume consistency*). The value $\kappa_{\text{BKM}} = 5$ is the Borcherds-weight indexing $c_1(0)/2$. The Vol I three-faces indexing on the same family records the shifted quantity $12 = 2 \cdot 5 + 0 + 2$ via the reconciliation identity of Remark 8.3. The two indexings are the same datum in two indexings; the BKM denominator algebra of this chapter realizes the Borcherds-weight side.

12 STRUCTURAL PROPERTIES OF $\mathfrak{g}_{\Delta_5}$ *Real-root sub-Kac–Moody algebra*

LEMMA 12.1 (*Real rank-three sub-Kac–Moody algebra*). The subalgebra $\mathfrak{g}_{\Delta_5}^{re} \subset \mathfrak{g}_{\Delta_5}$ generated by $\{e_i, f_i, h_i\}_{i=1}^3$ together with the Cartan $\Lambda_{II}^{2,1} \otimes \mathbb{C}$ is the rank-three hyperbolic Kac–Moody algebra $\mathfrak{g}(A)$ associated to the symmetric Cartan matrix $A = ((\delta_i, \delta_j)) = \begin{pmatrix} 2 & -2 & -2 \\ -2 & 2 & -2 \\ -2 & -2 & 2 \end{pmatrix}$.

Proof. The relations of Definition 4.2 restricted to real $\alpha_i \in \Delta_{\text{simp}}^{re}$ are exactly the Kac–Moody relations of [?KAC:1, Chapter 1] for the symmetric Cartan matrix A . The real Serre exponent $1 - 2(\delta_i, \delta_j)/(\delta_i, \delta_i) = 3$ is the standard exponent for $A_{ij} = -2$, $A_{ii} = 2$. $\square$

Remark 12.2 (*Hyperbolic structure of $\mathfrak{g}_{\Delta_5}^{re}$*). The Kac–Moody algebra $\mathfrak{g}(A)$ is hyperbolic in the sense of [?KAC:1, Chapter 4]: removing any one row of A yields a positive-definite Cartan matrix of type A_2 . The full BKM algebra $\mathfrak{g}_{\Delta_5}$ is the automorphic correction of $\mathfrak{g}(A)$ in the sense of [?GN, Section 3], i.e. adjoining the imaginary simple roots $a \in \Lambda_{II}^{2,1} \cap \mathbb{R}_{>0}\mathcal{P}_{II}$ with multiplicities $\mu_{\bar{p}}(a)$ to obtain the algebra whose denominator identity matches $64^{-1}\Delta_5(2Z)$.

Bilinear pairing on $\mathfrak{g}_{\Delta_5}$

LEMMA 12.3 (*Invariant bilinear form*). There is a unique non-degenerate invariant supersymmetric bilinear form

$$(\cdot, \cdot)_{\mathfrak{g}_{\Delta_5}} : \mathfrak{g}_{\Delta_5} \otimes \mathfrak{g}_{\Delta_5} \longrightarrow \mathbb{C}$$

extending the bilinear form on $\Lambda_{II}^{2,1} \otimes \mathbb{C} \subset \mathfrak{g}_{\Delta_5,0}$ and satisfying the ad-invariance

$$([X, Y], Z)_{\mathfrak{g}_{\Delta_5}} = (X, [Y, Z])_{\mathfrak{g}_{\Delta_5}}.$$

The pairing $(\mathfrak{g}_{\alpha}, \mathfrak{g}_{-\alpha})_{\mathfrak{g}_{\Delta_5}}$ is non-degenerate; pairings between non-opposite root spaces vanish.

Proof. Standard for Borchers–Kac–Moody algebras; see [?BorchersGKM88, Section 1] and [?KAC:1, Section 2.2]. The form on the Cartan extends uniquely by the relations $[h, e_i] = (\alpha_i, h)e_i$ to the relation $([h, e_i], f_i) + (e_i, [h, f_i]) = 0$ which forces $(e_i, f_i) \cdot (\alpha_i, h) - (e_i, f_i) \cdot (\alpha_i, h) = 0$, giving the relations on root pairings. $\square$

Weyl-equivariance of root spaces

LEMMA 12.4 (*Even Weyl transport of parity*). For every $w \in W^{(2)}(\Lambda_{II}^{2,1})$ and every root $\alpha \in \Delta$, the Weyl operator induces a parity-preserving isomorphism

$$\mathfrak{g}_{\alpha, \bar{p}} \simeq \mathfrak{g}_{w\alpha, \bar{p}}, \quad \bar{p} \in \mathbb{Z}/2.$$

Proof. This is the Weyl-transport-of-parity statement of [?KAC:1, Lemma 3.8 and Proposition 3.7], with the GN super-version of [?GN, Section 6, Statement 6.4]. Since the GN real simple generators are even, the reflection operators are even automorphisms of $\mathfrak{g}_{\Delta_5}$. Hence Weyl transport preserves the parity decomposition. $\square$

13 COMPARISON WITH THE CHIRAL SHADOW LATTICE Γ_{ind}

The Gram-index parametrization $\Gamma_{\text{ind}} = \mathbb{Z}^3$ of $\Lambda_{II}^{2,1}$ via $\alpha(n, l, m) = 2nf_2 - lf_3 + 2mf_{-2}$ provides the comparison between the BKM denominator algebra and the K_3 half-index $\phi_{o,1}$. The Gram-index pairing on Γ_{ind} is

$$\langle (n, l, m), (n', l', m') \rangle_{\text{ind}} = 2(nm' + n'm) - ll',$$

and the pullback to the Lorentzian pairing on $\Lambda_{II}^{2,1}$ is (4.4).

Remark 13.1 (The denominator polarization vs. the cusp polarization). The Gram-index lattice carries two polarizations:

- the cusp polarization, in which $\Gamma_{\text{eff}} \subset \Gamma_{\text{ind}}$ is the effective subset (the active support of the K_3 half-index in the Borchers product) and the order is the Fock polarization at one cusp;
- the denominator polarization, in which Γ_{ind} maps via α to $\Lambda_{II}^{2,1}$ and the order is the type-II chamber $\mathcal{P}_{II}$.

The two polarizations are the same lattice with two ordered charts; the denominator side is the Borchers chamber with $\mathcal{W}^{(2)}(\Lambda_{II}^{2,1}) \rtimes \text{Aut}(\mathcal{P}_{II})$ symmetry, the cusp side is the Fock determinant at the chosen cusp. A duality element transports between the two; cf. Section ?? of Chapter ??.

Visible coefficients

LEMMA 13.2 (*Active multiplicity equals $f(nm, l)$*). On the active support $\Gamma_{\text{act}} \subset \Gamma_{\text{eff}}$,

$$\text{smult}(\alpha(n, l, m)) = f(nm, l).$$

For triples $(n, l, m) \in \Gamma_{\text{eff}} \setminus \Gamma_{\text{act}}$, the product exponent vanishes and no visible product factor is contributed; the BKM root space $\mathfrak{g}_{\alpha(n, l, m)}$ is not pinned by the Borchers product alone.

Proof. The first equality is (4.17), comparing the Borchers product expansion with the Weyl–Kac–Borchers denominator. The second statement records that exponent zero on a triple (n, l, m) does not exclude the existence of a root space with $\text{smult} = 0$; the Borchers product expansion does not see zero-superdimension root spaces. $\square$

What the determinant determines

PROPOSITION 13.3 (*Information content of the denominator product*). The denominator identity

$$\text{den}(\mathfrak{g}_{\Delta_s}) = 64^{-1} \Delta_5(2Z)$$

determines the signed integers $\text{smult}(\alpha)$ on the active support after the BKM root system $\mathcal{R}_+$ has been fixed. It does not determine:

- invisible zero-superdimension root spaces;
- the parity decomposition $\dim \mathfrak{g}_{\alpha, \bar{0}}, \dim \mathfrak{g}_{\alpha, \bar{1}}$ beyond their difference;
- the structure constants of the BKM bracket $[\mathfrak{g}_\alpha, \mathfrak{g}_\beta] \rightarrow \mathfrak{g}_{\alpha+\beta}$;
- simple primitive representatives, no-extra-relations theorems, or completed PBW compatibility;
- the closed radical of the positive-negative invariant pairing.

Consequently the denominator product is a necessary input for primitive recognition; it is not by itself a recognition datum.

Proof. Write $X^\alpha = e^{-2\pi i(\alpha, z)}$. In the completed positive root monoid algebra,

$$\log \prod_{\alpha \in \mathcal{R}_+} (1 - X^\alpha)^{\text{smult}(\alpha)} = - \sum_{\alpha \in \mathcal{R}_+} \sum_{k \geq 1} \frac{\text{smult}(\alpha)}{k} X^{k\alpha}.$$

The exponents $\text{smult}(\alpha)$ are recovered recursively by height, equivalently by Mobius inversion on each primitive ray. The recovered exponent is only $\text{smult}(\alpha) = \text{mult}_{\bar{0}}(\alpha) - \text{mult}_{\bar{1}}(\alpha)$. The difference $a - b$ does not determine the pair (a, b) . No bilinear bracket, PBW filtration, or Hopf-pairing radical appears in the denominator product. $\square$

What the determinant forgets

PROPOSITION 13.4 (*Grothendieck class as the determinant invariant*). For every Γ_{eff} -graded super vector space $\mathcal{B} = \bigoplus_{\gamma} \mathcal{B}_{\gamma}$ with $\text{sdim } \mathcal{B}_{(n,l,m)} = f(nm, l)$, the Fock determinant is

$$64q^{1/2}r^{1/2}s^{1/2} \prod_{\gamma} (1 - x^{\gamma})^{\text{sdim } \mathcal{B}_{\gamma}} = \Delta_5(Z).$$

The determinant remembers only the class of $\mathcal{B}$ in

$$K_0(\Gamma_{\text{eff}}\text{-graded finite super vector spaces}).$$

A bracket $[\mathcal{B}_{\gamma}, \mathcal{B}_{\gamma'}] \rightarrow \mathcal{B}_{\gamma+\gamma'}$ does not appear in the determinant formula, nor do simple primitive representatives, parity dimensions, or pairing radicals.

Proof. The Fock determinant lemma gives $\text{sdet}_{\mathcal{B}}(1 - x) = \prod_{\gamma} (1 - x^{\gamma})^{\text{sdim } \mathcal{B}_{\gamma}}$. Multiplying by the vacuum factor and using the hypothesis $\text{sdim } \mathcal{B}_{(n,l,m)} = f(nm, l)$ identifies this product with the Borchers product for $64^{-1}\Delta_5$. Replacing $\mathcal{B}$ by any other graded super vector space with the same Grothendieck class preserves the determinant; the zero bracket on such a space gives the same determinant and a wrong Lie algebra; adjoining a cancelling pair $\mathcal{W}_{\gamma} \oplus \Pi \mathcal{W}_{\gamma}$ in any active degree leaves the determinant unchanged while changing the parity dimensions. Hence the determinant invariant is the K_0 -class. $\square$

14 COMPARISON: RANK-THREE HYPERBOLIC VS. BORCHERDS CORRECTION

The denominator algebra $\mathfrak{g}_{\Delta_5}$ is the BKM correction of the rank-three hyperbolic Kac–Moody algebra $\mathfrak{g}(\mathcal{A})$ on the same Cartan matrix $\mathcal{A}$. The two algebras have the same real-root datum and the same real Serre relations; they differ in the imaginary roots.

Remark 14.1 (Borchers correction adjoins the imaginary simple roots). The Gritsenko–Nikulin construction [GN, Sections 3–4, Proposition 3.1] adjoins to $\mathfrak{g}(\mathcal{A})$ the imaginary simple generators with multiplicities $\mu_{\bar{p}}(a)$ from the Borchers-product expansion of the additive lift of $\eta^9 \mathfrak{g}$. The resulting BKM Lie superalgebra has denominator $64^{-1}\Delta_5(2Z)$, while $\mathfrak{g}(\mathcal{A})$ has no closed denominator product as a Kac–Moody algebra alone (its denominator is a formal power series whose closed form is unknown). The correction algebra is the central object of View II.

15 THE CHARACTER OF THE CHIRAL SHADOW

The denominator algebra $\mathfrak{g}_{\Delta_5}$ is the Beilinson cut’s chiral shadow at View II. The Beilinson cut requires that primitive objects come first, cross-level identities second, and scalar shadows last. Inscribed at View II:

TABLE 4.1: Real-root datum of $\mathfrak{g}_{\Delta_5}$ and the rank-three hyperbolic $\mathfrak{g}(A)$.

	$\mathfrak{g}(A), A = \begin{pmatrix} 2 & -2 & -2 \\ -2 & 2 & -2 \\ -2 & -2 & 2 \end{pmatrix}$	$\mathfrak{g}_{\Delta_5}$
real simple roots	$\delta_1, \delta_2, \delta_3$	$\delta_1, \delta_2, \delta_3$
real-root mult.	1 each	1 each
real Serre exponent	3	3
imaginary simple roots	none	$\Lambda_{II}^{2,1} \cap \mathbb{R}_{>0} \mathcal{P}_{II}$
$\mu_{\bar{o}}(a)$	o	$\max(m(a), o) \text{ or } \max(\tau(a), o)$
$\mu_{\bar{i}}(a)$	o	$\max(-m(a), o) \text{ or } \max(-\tau(a), o)$
denominator	no closed product	$64^{-1} \Delta_5(2Z)$

- *Primitive object* (View III, conditional): the chiral E_3 -algebra $\mathcal{A}_{K_3 \times E}$ of the $K_3 \times E$ formal target, presented at finite stage as the pro-object $\mathfrak{D}_X^{DI}$ of Chapter ??.
- *Cross-level identity* (View II): the Borcherds–Kac–Moody denominator $\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1} \Delta_5(2Z)$, with generators and relations as recorded above.
- *Scalar shadow* (View I, K_o -determinant): the automorphic section $\mathcal{D}_X = \Delta_5 \in \mathcal{M}_5(\mathbf{Sp}_4(\mathbb{Z}), \nu_{\Delta_5})$, with squared scalar shadow Δ_5^2 at weight 10.

The denominator algebra is target-side: it imports the Gritsenko–Nikulin Cartan, the Chevalley–Borcherds–Kac relations, the Weyl vector, and the Borcherds product, and supplies no compact Hall correspondence, no orientation line, no Pfaffian section, and no BPS Hilbert space. The conditional cross-level identity View II $\leftrightarrow$ View III, which would establish the Pfaffian–Dirac theorem, requires the four obstructions $(D_o), (O_1), (O_1)^+, (O_2)$ of Chapter ?? to be discharged.

Remark 15.1 (The chiral shadow is target, not bulk). The Beilinson cut classifies $\mathfrak{g}_{\Delta_5}$ as a chiral shadow, not a chiral bulk. The bulk object lives on the source side: at finite-stage realization the chiral bulk is $Z_{\text{ch}}^{\text{der}}(C_{X,R}) \simeq \text{ChirHoch}^\bullet(C_{X,R}, C_{X,R})$ (Vol II Universal Holography master theorem), attached to a primitive $C^{op} = C_{X,R}^{op}$ on the closed factorization datum (X, D, τ) at the boundary vacuum b . The denominator algebra $\mathfrak{g}_{\Delta_5}$ is the chiral shadow of this bulk, not the bulk itself. The bulk object lives one categorical level above the chiral shadow; the BKM denominator records the shadow.

16 THE DENOMINATOR ALGEBRA OF THE CHIRAL SOURCE-TARGET FRAME

The denominator algebra is recorded together with its formal current envelope as a target chiral object on a smooth complex curve.

Definition 16.1 (Formal current envelope). For a smooth complex curve C , set

$$\text{Cur}_C(\mathfrak{g}_{\Delta_5}) := \mathfrak{g}_{\Delta_5} \otimes \mathcal{O}_C$$

with the level-zero current bracket

$$[a \otimes f, b \otimes g] = [a, b] \otimes fg.$$

The chiral enveloping algebra of Beilinson–Drinfeld [?BD] is

$$\text{Den}_{\Delta_5, C} := U_C^{\text{ch}}(\text{Cur}_C(\mathfrak{g}_{\Delta_5})).$$

PROPOSITION 16.2 (*Formal current envelope realizes the denominator*). The chiral algebra $\text{Den}_{\Delta_5, C}$ is the formal completed ind Beilinson–Drinfeld current envelope on C obtained from the denominator Lie superalgebra. Its constant-current Lie superalgebra is $\mathfrak{g}_{\Delta_5}$, and the denominator of that constant-current bracket is $64^{-1}\Delta_5(2Z)$.

For $C = E$, the elliptic curve in $X = K_3 \times E$, the chiral algebra $\text{Den}_{\Delta_5, E}$ is the target chiral object used in the $K_3 \times E$ realization problem of Chapter ?? . A compact E_3/Hall /chiral protected realization requires additional microscopic input (the Dirac–Igusa pro-object $\mathfrak{D}_X^{DI}$).

Proof. In the completed ind setting, the current construction sends a Lie superalgebra to a Lie- $\ast$ algebra on a smooth curve, and the chiral enveloping construction sends a Lie- $\ast$ algebra to a chiral algebra [?BD]. Equivalently, in vertex-algebra language, this is the universal current vertex algebra at level zero [?KACVA]. Constant sections of $\mathfrak{g}_{\Delta_5} \otimes \mathcal{O}_E$ recover the original Lie superalgebra, with denominator $\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1}\Delta_5(2Z)$ from Theorem 9.1. $\square$

17 THE CLOSING IDENTITY AND ITS CONDITIONING

The closing identity of this chapter is

$$\boxed{\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1}\Delta_5(2Z), \quad \kappa_{\text{BKM}}(\Delta_5) = 5, \quad \text{smult}(\alpha(n, l, m)) = f(nm, l).}$$

These three are imported theorems: the first from Borcherds 1995 [?Borcherds95] and Gritsenko–Nikulin 1998 [?GNII] [?GNII, Theorem 2.1, (2.1)–(2.3)], with the algebra constructed in [?GN, Sections 3–4, Proposition 3.1]; the second from Borcherds 1998 [?B, Theorem 13.3] with the Vol III specialization to the paramodular family; the third from the same Borcherds product expansion as the first, by comparison with the Weyl–Kac–Borcherds denominator. The conditioning of the closing identity is target-side. No compact Hall, Pfaffian, or BPS Hilbert space is claimed.

The cross-level identity View II $\leftrightarrow$ View III, the conditional Pfaffian–Dirac theorem of Chapter ?? , asks for the protected Pfaffian

$$\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{DI}) = \Delta_5,$$

the orientation character $\epsilon_o = \nu_{\Delta_5}$ on $W^{(2)}(\Lambda_{II}^{2,1})$, and the primitive recognition theorem $P_X \simeq \mathfrak{g}_{\Delta_5}$, conditional on the discharging of the four obstructions (D_0) , (O_1) , $(O_1)^+$, (O_2) . The present chapter is independent of those four obstructions; it imports the BKM denominator identity from primary literature.

Status of each load-bearing claim

The following claims are inscribed in this chapter; the table records the epistemic status and primary citation of each.

Claim	Status	Citation
Type-II Cartan A , Lemma 2.2	proved here	direct
Weyl vector $\rho = f_2 - \frac{1}{2}f_3 + f_{-2}$, Prop. 2.4	proved here	direct
Reflection-group decomposition, Lemma 1.3	proved	[?NIKULIN]
Real-root multiplicity is 1, Lemma 4.6	proved	[?KAC: 1, ?GN]
Imaginary simple-root parity split, Def. 3.2	proved	[?GN, Section 3]
Maass multiplier values, Lemma 7.1	proved	[?MAASS: 1, ?GN]
Multiplier restriction, Lemma 7.2	proved	[?GN, Prop. 2.1, 2.2]
Borcherds product for D_5 , Theorem 5.3	proved	[?B, Thm. 13.3], [?GNII, Thm. 2.1]
Weyl alternating sum, Theorem 5.1	proved	[?GN, Thm. 2.1]
Denominator identity (4.16)	proved	[?GN, Sections 3–4], [?GNII, Thm. 2.1]
Borcherds-weight $\kappa_{\text{BKM}} = 5$, Theorem 8.1	proved	[?B, Thm. 13.3], Vol III
29 93 presentation count, Prop. 6.7	proved here	direct + [?GN]
Triangular decomposition, Lemma 4.4	proved	[?BorcherdsGKM88, ?KAC: 1]
Invariant bilinear form, Lemma 12.3	proved	[?BorcherdsGKM88, ?KAC: 1]

The closing identity is therefore proved as recorded; its compact realization is the conditional theorem of Chapter ??.

18 ADVERSARIAL CLOSURE

The denominator-algebra theorem has been attacked along the following axes; each attack closes against the recorded primary sources and the Borcherds-weight identity.

Sign of imaginary multiplicities

Claim attacked. The Borcherds product exponent $f(nm, l)$ on the active support determines a unique BKM signed multiplicity.

Closure. Direct: smult is defined as the difference $\text{mult}_{\bar{\sigma}} - \text{mult}_{\bar{\tau}}$ (Definition 4.5), and the comparison of the Borcherds product (4.13) with the Weyl–Kac–Borcherds denominator product (4.15) gives $\text{smult}(\alpha(n, l, m)) = f(nm, l)$ on Γ_{act} . The parity split of smult into $(\text{mult}_{\bar{\sigma}}, \text{mult}_{\bar{\tau}})$ requires the Gritsenko–Nikulin imaginary simple-root data $\mu_{\bar{p}}(a)$, recorded in Definition 3.2, which is Borcherds-product information augmented by the parity sign of $m(a)$ and $\tau(a)$. Compare [?GN, Section 3].

Doubled vs. undoubled Δ_5

Claim attacked. The denominator identity uses $\Delta_5(2Z)$, not $\Delta_5(Z)$. Why?

Closure. The BKM root grading is on $\Lambda_{II}^{2,1}$, while the additive Gritsenko lift produces Δ_5 on $\Lambda_I^{2,1}$ in its primitive form, with $\Lambda_{II}^{2,1} \subset \frac{1}{2}\Lambda_I^{2,1}$. The doubling $Z \mapsto 2Z$ identifies the two lattice gradings; the prefactor $64 = [q^{1/2}r^{1/2}s^{1/2}]\Delta_5$ is the theta-constant normalization and is untouched by doubling. See Remark 5.5.

Doubling factor as power of two

Claim attacked. The factor $64 = 2^6$ is associated with a power of two from a half-Hilbert orbit.

Closure. The factor 64 is the value of the leading Fourier coefficient $c_{\Delta}(1, 1, 1)$ of Δ_5 , computed in the Maass formula for the product of ten even theta constants [?GN, (1.3)–(1.5)]. It is not a half-Hilbert orbit count. The orbit-count factor $2^{12} = 4096$ arises in the type-II Pfaffian wall normal form (O_2) of Chapter ?? as the size of the half-Hilbert orbit on the local sign atlas; it is unrelated to the denominator prefactor 64. See Remark 5.6.

Simple-root system on $\Lambda_{II}^{2,1}$

Claim attacked. A different basis of three vectors of square 2 in $\Lambda_{II}^{2,1}$ would yield a different BKM algebra.

Closure. Nikulin’s reflection-group theorem [?NIKULIN] pins $W^{(2)}(\Lambda_{II}^{2,1})$ as a normal subgroup of index 6 in $\mathbf{O}(\Lambda^{2,1})_+$, and the Gritsenko–Nikulin chamber $\mathcal{P}_{II}$ has fixed walls $\delta_1, \delta_2, \delta_3$ with Gram matrix (4.8). Any other choice of three primitive type-II square-2 vectors lies in the same Weyl orbit as some permutation of $(\delta_1, \delta_2, \delta_3)$. The BKM algebra depends on the equivalence class of the Cartan matrix, not on the specific basis vectors.

Weyl vector ρ

Claim attacked. The Weyl vector might be $\rho' = \delta_1 + \delta_2 + \delta_3$ rather than $\frac{1}{2}(\delta_1 + \delta_2 + \delta_3)$.

Closure. The Weyl vector is fixed by the condition $(\rho, \delta_i) = -(\delta_i, \delta_i)/2 = -1$ for $i = 1, 2, 3$. The displayed value $\rho = \frac{1}{2}(\delta_1 + \delta_2 + \delta_3)$ satisfies this and lies in $(\Lambda_{II}^{2,1})^* \cap C(\Lambda^{2,1})_+$; $\rho' = \delta_1 + \delta_2 + \delta_3$ would have $(\rho', \delta_i) = -2$, violating the condition. The factor $\frac{1}{2}$ is forced. See Proposition 2.4.

Real-root norm convention

Claim attacked. The convention $(\delta_i, \delta_i) = 2$ might be exchanged with $(\delta_i, \delta_i) = 1$ on a non-even lattice.

Closure. The lattice $\Lambda^{2,1} = \Lambda^{(1,1)} \oplus [2]$ is even by (4.1). The type-II sublattice $\Lambda_{II}^{2,1}$ is also even (a sublattice of an even lattice). Hence $(\delta, \delta) \in 2\mathbb{Z}$ for every $\delta \in \Lambda_{II}^{2,1}$. The choice $(\delta, \delta) = 2$ is forced for primitive square-norm-2 vectors. The convention $(\delta_i, \delta_i) = 1$ does not occur.

Second Serre bracket compatibility (Gram cocycle)

Claim attacked. The Gram cocycle $B(c, c')$ on the upstairs lattice Γ_X^{phys} might be compatible with the BKM bracket only after normal ordering.

Closure. The Gram cocycle $B(c, c')$ is the source-side quadratic form on the upstairs lattice; its compatibility with the BKM bracket on $\Lambda_{II}^{2,1}$ is mediated by the normal-ordered homomorphism $\overline{\Pi}_X : \widehat{\Gamma}_X \rightarrow \Gamma_{\text{gram}}$ of Chapter ??, which corrects the raw quadratic shadow $\Pi_X(c + c') = \Pi_X(c) + \Pi_X(c') + B(c, c')$ to the additive map. The present chapter, which is target-side, uses the Lorentzian Cartan pairing on $\Lambda_{II}^{2,1}$ directly; no Gram cocycle correction is needed. The compatibility is precisely between $\overline{\Pi}_X$ on the source and the BKM grading on the target, as established in Chapter ??.

Alternative Cartan-matrix choice

Claim attacked. A symmetrized Cartan matrix $\begin{pmatrix} 1 & -1 & -1 \\ -1 & 1 & -1 \\ -1 & -1 & 1 \end{pmatrix}$ would also satisfy the Borchers–Kac axioms and might give the same algebra.

Closure. Borchers–Kac–Moody algebras are classified up to scalar normalization of the Cartan matrix by the equivalence class of A . The matrix A in (4.8) has integer entries with $A_{ii} = 2$, and is the symmetrized form fixed by the even-lattice convention. A rescaled matrix $\frac{1}{2}A$ with $A_{ii} = 1$ would not be a generalized Cartan matrix in the Kac–Moody sense (the condition $A_{ii} \in 2\mathbb{Z}_{>0}$ is part of the standard normalization in [?KAC:1, Chapter 1]). The symmetric form A is the unique normalization compatible with the even lattice $\Lambda_{II}^{2,1}$.

Cross-volume κ_{BKM} disagreement

Claim attacked. The Vol I three-faces value 12 might disagree with this chapter's Borchers-weight value 5.

Closure. The two values are the two indexings of the same quantity: $\kappa_{\text{BKM}}^{(\text{Borch})} = c_1(o)/2 = 5$ from [?B, Theorem 13.3] (Borchers-weight indexing), and $\kappa_{\text{BKM}}^{(3\text{-faces})} = 2\kappa_{\text{BKM}}^{(\text{Borch})} + \chi(\mathcal{O}_X) + 2 = 12$ on $X = K_3 \times E$ (three-faces indexing). The reconciliation $12 = 2 \cdot 5 + 0 + 2$ is a single identity, not a conflict. See Remark 8.3 and the Vol III universal-Borchers-weight identity (Theorem 8.1).

Determinant alone determines the algebra

Claim attacked. The denominator product $64^{-1}\Delta_5(2Z)$ alone determines $\mathfrak{g}_{\Delta_5}$.

Closure. False: a graded super vector space with the same signed multiplicities and zero bracket gives the same denominator product but is not a Lie superalgebra. The denominator identity fixes only the signed Grothendieck class (Proposition 13.4). The bracket, the parity split, the radical, the simple primitives, and the no-extra-relations theorem are required for a recognition theorem; all of these are read on the BKM presentation of Definition 4.2, not from the denominator product alone. The recognition target $P_X \simeq \mathfrak{g}_{\Delta_5}$ requires the compact-source data of Chapter ??.

19 CODA: $\mathfrak{g}_{\Delta_5}$ AS ONE OF FIVE VIEWS

The denominator algebra $\mathfrak{g}_{\Delta_5}$ is one of five views of Δ_5 : View I (automorphic section), View II (Borchers–Kac–Moody denominator, this chapter), View III (compact protected Pfaffian, conditional), View IV (mirror discriminant, conjectural), View V (singular theta lift). The cross-level identities are:

- View I $\leftrightarrow$ View II: theorem, [?Borchers95, ?GNII].
- View II $\leftrightarrow$ View V: theorem, [?B].
- View I $\leftrightarrow$ View V: theorem, [?GN, ?B].
- View II $\leftrightarrow$ View III: *conditional*, Chapter ??, under (Do) , (O_1) , $(O_1)^+$, (O_2) .
- View II $\leftrightarrow$ View IV: *conjectural*, Chapter ??.

The closing identity of this chapter is the recorded denominator identity

$$\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1}\Delta_5(2Z),$$

the Borchers-weight characteristic

$$\kappa_{\text{BKM}}(\Delta_5) = 5,$$

and the active multiplicity formula

$$\text{smult}(\alpha(n, l, m)) = f(nm, l) \quad ((n, l, m) \in \Gamma_{\text{act}}).$$

These three identities are the load-bearing target-side data of View II. The chapter is closed.

THE SINGULAR THETA LIFT

I THE SINGULAR THETA LIFT OF $\phi_{o,I}$

The four cross-level identifications proved by Borcherds, Gritsenko, and Nikulin assemble Δ_5 into one object seen from five entry points: the automorphic section of the weight-five Hodge line on $\mathbf{Sp}_4(\mathbb{Z}) \backslash \mathbb{H}_2$ (View I), the denominator of $\mathfrak{g}_{\Delta_5}$ on $\Lambda_{II}^{2,1}$ (View II), the protected Pfaffian of the conditional Dirac–Igusa pro-object $\mathfrak{D}_X^{DI}$ (View III), the conjectural mirror discriminant on the K3 period domain (View IV), and the singular theta lift of the weak Jacobi form $\phi_{o,I}$ under the Howe correspondence $\mathbf{Mp}(4, \mathbb{R}) \leftrightarrow \mathbf{O}(3, 2)$ (View V). This chapter records View V. The exceptional isomorphism $\mathbf{Sp}_4 \simeq \mathbf{Spin}(3, 2)$ compiles the genus-two Siegel upper half-space and the type-IV symmetric domain attached to the signature- $(3, 2)$ lattice into one homogeneous space; the singular theta integral of [?Borcherds95, ?B] sends $\phi_{o,I}$ to the weight-five Borcherds product Δ_5 , and the Gritsenko–Nikulin denominator identity then reads the lift as the BKM denominator on $\Lambda_{II}^{2,1}$. All identifications below are theorems of [?Borcherds95, ?B, ?GN, ?GNII]; the chapter assembles them with the manuscript’s lattice conventions and licensing tags.

1.1 Notation and the canonical name

Three Borcherds-product handles for the weight-five Igusa cusp form live in the manuscript. The *automorphic section* $\mathcal{D}_X = \Delta_5$ of the line $\mathcal{L}^5 \otimes \nu_{\Delta_5}$ on $\mathbf{Sp}_4(\mathbb{Z}) \backslash \mathbb{H}_2$ is the View-I name. The *BKM denominator* $\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1} \Delta_5(2Z)$ on $\Lambda_{II}^{2,1}$ is the View-II name. The *protected Pfaffian* $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{DI})$, conditional on the four Dirac–Igusa obstructions, is the View-III name. This chapter introduces the View-V canonical name

$$\Delta_5 = \Theta(\phi_{o,I}),$$

the *singular theta lift* of $\phi_{o,I}$. Throughout $\phi_{o,I} \in J_{o,I}^{\text{weak}}$ is the unique up to scale weight-zero index-one weak Jacobi form on the Jacobi group $\Gamma_1^J = \mathbf{SL}_2(\mathbb{Z}) \ltimes \mathbb{Z}^2$ with constant Fourier coefficient fixed by the K3 elliptic genus normalization $Z_{K3} = 2\phi_{o,I}$ of [?EZ: 1, §2.2], and Θ is the Borcherds singular-theta integral of [?Borcherds95, §6], [?B, §6]. The bare handle Δ_5 is reserved for cross-view identifications.

1.2 The weak Jacobi form $\phi_{o,I}$

Definition 1.1 (Weak Jacobi form, weight zero, index one). A weak Jacobi form of weight k and index m on the Jacobi group $\Gamma_1^J = \mathbf{SL}_2(\mathbb{Z}) \ltimes \mathbb{Z}^2$ is a holomorphic function $\phi : \mathbb{H} \times \mathbb{C} \rightarrow \mathbb{C}$ satisfying, for $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathbf{SL}_2(\mathbb{Z})$ and $(\lambda, \mu) \in \mathbb{Z}^2$,

$$\begin{aligned} \phi\left(\frac{a\tau + b}{c\tau + d}, \frac{z}{c\tau + d}\right) &= (c\tau + d)^k e^{2\pi i m c z^2 / (c\tau + d)} \phi(\tau, z), \\ \phi(\tau, z + \lambda\tau + \mu) &= e^{-2\pi i m (\lambda^2 \tau + 2\lambda z)} \phi(\tau, z), \end{aligned}$$

together with the cusp expansion

$$\phi(\tau, z) = \sum_{n \geq 0, l \in \mathbb{Z}} f(n, l) q^n r^l, \quad q = e^{2\pi i \tau}, \quad r = e^{2\pi i z},$$

in which $f(n, l)$ depends only on the discriminant $4nm - l^2$ (Eichler–Zagier [EZ: 1, Theorem 2.2]). The *weight-zero index-one* weak Jacobi form $\phi_{0,1}$ is fixed by $k = 0$, $m = 1$, the discriminant condition $4n - l^2 \geq -1$ and the normalization

$$\phi_{0,1}(\tau, z) = r^{-1} + 10 + r + O(q).$$

The space $J_{0,1}^{\text{weak}}$ is one-dimensional over $\mathbb{C}$ (Eichler–Zagier [EZ: 1, Theorem 9.4 and Table 9, §9]); $\phi_{0,1}$ generates it. The theta-quotient presentation is the input to the singular-theta integral below.

LEMMA 1.2 (*Eichler–Zagier theta-quotient presentation*). The weak Jacobi form $\phi_{0,1}$ admits the closed form

$$\phi_{0,1}(\tau, z) = 4 \sum_{i=2}^4 \frac{\theta_i(\tau, z)^2}{\theta_i(\tau, 0)^2},$$

where $\theta_2, \theta_3, \theta_4$ are the three even Jacobi theta functions in the Mumford normalization (Eichler–Zagier [EZ: 1, §9, equation (9.4)]); equivalently $\phi_{0,1} = Z_{K3}/2$ is one half of the $K3$ elliptic genus [DMVV]. The leading Fourier coefficients are

$$\begin{aligned} f(0, 0) &= 10, & f(0, \pm 1) &= 1, & f(1, 0) &= 108, & f(1, \pm 1) &= -64, & f(1, \pm 2) &= 10, \\ f(2, 0) &= 808, & f(2, \pm 1) &= -513, & f(2, \pm 2) &= 108, & f(2, \pm 3) &= 1, \end{aligned}$$

in agreement with `main.tex` §?? and verified arithmetically by `compute/verify_square_root.py`. The discriminant relation $f(n, l) = f(n', l')$ when $4n - l^2 = 4n' - (l')^2$ (Eichler–Zagier [EZ: 1, Theorem 2.2]) organises these coefficients into the equivalence classes

$$\{f(1, 0), f(2, \pm 2)\} = 108 (\text{disc } 4), \quad \{f(0, 0), f(1, \pm 2)\} = 10 (\text{disc } 0), \quad \{f(0, \pm 1), f(2, \pm 3)\} = 1 (\text{disc } -1),$$

each independently consistent with the verified expansion.

Proof. The closed form is standard in Eichler–Zagier [EZ: 1, §§9–10]; the $K3$ elliptic-genus identity $Z_{K3} = 2\phi_{0,1}$ is [EZ: 1, §2.2] together with the sigma-model right-moving Witten index of [DMVV]. The constant term $f(0, 0) = 10$ is the right-moving-localized character of the $K3$ Ramond sector at $L_0 = c/24$; the remaining values follow from the discriminant relation $f(n, l) = f(n', l')$ when $4n - l^2 = 4n' - (l')^2$ (Eichler–Zagier [EZ: 1, Theorem 2.2]) together with explicit theta-quotient evaluation, which fixes the disc-3 class ($f(1, \pm 1) = -64$) and the disc-4 class ($f(1, 0) = f(2, \pm 2) = 108$) by direct computation from the theta-quotient of Lemma 1.2. The squared theta-leading coefficient $([q^{1/2} r^{1/2} s^{1/2}] \Delta_5)^2 = 64^2 = 4096$ of the Borcherds-product expansion in §?? is the next arithmetic check on these coefficients via the cusp-chamber product $\Delta_5(Z) = 64 q^{1/2} r^{1/2} s^{1/2} \prod_{(n,l,m) \in \Gamma_{\text{eff}}} (1 - q^n r^l s^m)^{f(nm,l)}$. $\square$

1.3 The signature-(3, 2) lattice and the type-IV domain

Definition 1.3 (*The signature-(3, 2) lattice*). The lattice $\Lambda^{3,2}$ is the orthogonal complement of the alternating Pfaffian element $q_0 = e_1 \wedge e_3 + e_2 \wedge e_4 \in \wedge^2 \Lambda^4$ inside the signature-(3, 3) lattice $(\wedge^2 \Lambda^4, (\cdot, \cdot)) \simeq \Pi_{3,3}$, where $\Lambda^4 = \bigoplus_{i=1}^4 \mathbb{Z} e_i$ and the bilinear form (u, v) is the Pfaffian pairing $u \wedge v = (u, v) e_1 \wedge e_2 \wedge e_3 \wedge e_4$. As an even unimodular lattice,

$$\Lambda^{3,2} \simeq \Lambda^{(1,1)} \oplus \Lambda^{(1,1)} \oplus [2], \quad \Lambda^{(1,1)} = \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix},$$

written in the basis $(f_1, f_2, f_3, f_{-2}, f_{-1})$ of the exterior-square model section.

Remark 1.4 (Signature conventions). The lattice $\Lambda^{3,2}$ carries three positive and two negative directions in the (u, v) -pairing of Definition 1.3. The *primitive hyperbolic sublattice* $\Lambda^{2,1} = \Lambda^{(1,1)} \oplus [2]$ on which the BKM denominator algebra $\mathfrak{g}_{\Delta_5}$ is graded has signature $(1, 2)$ in the manuscript convention used throughout: one positive direction (the $[2]$ factor) and two negative directions inside the cone $C(\Lambda^{2,1})_+$. The decomposition $\Lambda^{3,2} = \Lambda^{2,1} \oplus \Lambda^{(1,1)}$ splits an extra hyperbolic plane on which the singular theta integral lives; this is the $\Pi_{2,2}$ -fibre at the standard cusp under the exceptional isomorphism of §1.4. No $\Lambda^{2,1} \oplus \langle -2 \rangle$ presentation is invoked; the manuscript convention is the $\Lambda^{(1,1)} \oplus [2]$ Nikulin form. $[\gamma$: ambient declaration]

The type-IV symmetric domain attached to $\Lambda^{3,2}$ is

$$\mathbb{H}^{\text{IV}} = \{Z \in \mathbb{P}(\Lambda^{3,2} \otimes \mathbb{C}) \mid (Z, Z) = 0, (Z, \bar{Z}) < 0\},$$

with positive component $\mathbb{H}_+^{\text{IV}}$ singled out by $\text{Im} z_1 > 0$ in the parametrization ${}^t(z_2^2 - z_1 z_3, z_3, z_2, z_1, 1)z_0$ of the exterior-square model section. Its arithmetic quotient $\mathbf{O}(\Lambda^{3,2})_+ \backslash \mathbb{H}_+^{\text{IV}}$ is the moduli space on which the Borchers singular-theta lift of [?Borchers95] produces automorphic forms with prescribed singularities along rational quadratic divisors.

1.4 The exceptional isomorphism $\mathbf{Sp}_4 \simeq \mathbf{Spin}(3, 2)$

THEOREM 1.5 (*Exterior-square exceptional isomorphism*). The exterior-square representation $\wedge^2 : \mathbf{SL}_4 \rightarrow \text{Aut}(\wedge^2 \Lambda^4, (,))$ restricts to a homomorphism

$$\wedge^2 : \mathbf{Sp}_4(\mathbb{Z}) / \{\pm \mathbf{I}_4\} \xrightarrow{\cong} \mathbf{SO}_+(\Lambda^{3,2}),$$

realizing the exceptional isomorphism of real Lie groups $\mathbf{PSp}_4(\mathbb{R}) \simeq \mathbf{SO}(3, 2)_+$ and its spin lift $\mathbf{Sp}_4(\mathbb{R}) \simeq \mathbf{Spin}(3, 2)$. Under this identification the genus-two Siegel upper half-space $\mathbb{H}_2$ embeds onto the positive component $\mathbb{H}_+^{\text{IV}}$ by

$$\begin{pmatrix} z_1 & z_2 \\ z_2 & z_3 \end{pmatrix} \longmapsto {}^t(z_2^2 - z_1 z_3, z_3, z_2, z_1, 1)z_0,$$

intertwining the $\mathbf{Sp}_4(\mathbb{Z})$ and $\mathbf{SO}_+(\Lambda^{3,2})$ actions.

[proved; β : comparison morphism]

Proof. The Pfaffian form $u \wedge v = (u, v)e_1 \wedge e_2 \wedge e_3 \wedge e_4$ is the unique $\mathbf{SL}_4$ -invariant alternating bilinear form on $\wedge^2 \Lambda^4$; the subgroup of $\mathbf{SL}_4$ fixing $q_0 = e_1 \wedge e_3 + e_2 \wedge e_4$ is the symplectic group $\mathbf{Sp}_4$ of the alternating form B_{q_0} . The orthogonal complement $\Lambda^{3,2} = q_0^\perp$ is preserved, and the kernel of $\wedge^2|_{\mathbf{Sp}_4}$ on $q_0^\perp$ is $\{\pm \mathbf{I}_4\}$ (the exterior-square model section). This is the integral form of the classical exceptional isomorphism $\mathbf{PSp}_4(\mathbb{R}) \simeq \mathbf{SO}(3, 2)_+$ of [?KNAPP:1, Chapter II.6]; the spin lift is the metaplectic description $\mathbf{Sp}_4(\mathbb{R}) \simeq \mathbf{Spin}(3, 2)$ of [?Weil1964, §§5–6]. The displayed substitution into the type-IV domain produces the commutative square of the exterior-square model section. The integral-Igusa form of the identification $\mathbb{H}_2 \simeq \mathbb{H}_+^{\text{IV}}$ is the content of [?Igusa1962, §2]; see also [?FREITAG:1, Chapter II] and [?VDGEER:1, §6]. $\square$

Remark 1.6 (Three guises of one homogeneous space). The same homogeneous space carries three names: the genus-two Siegel domain $\mathbf{Sp}_4(\mathbb{R})/\mathbf{U}(2)$, the type-IV symmetric domain $\mathbf{SO}(3, 2)_+ / (\mathbf{SO}(3) \times \mathbf{SO}(2))$, and the spin double cover $\mathbf{Spin}(3, 2) / (\mathbf{U}(2)\text{-cover})$. The Howe correspondence below uses the metaplectic cover $\mathbf{Mp}_4(\mathbb{R})$, which agrees with $\mathbf{Spin}(3, 2)$ as a real Lie group. The integrality of $\text{wt}(\Delta_5) = 5$ means the cover question is trivialized for the *weight* factor; the multiplier system ν_{Δ_5} records the residual reflection character on $\mathcal{W}^{(2)}(\Lambda_{II}^{2,1})$ (Lemma ??). $[\alpha$: chart / cover declaration]

1.5 The Howe correspondence $\mathbf{Mp}_4(\mathbb{R}) \leftrightarrow \mathbf{O}(3, 2)$

THEOREM 1.7 (*Howe duality, type I dual reductive pair* $(\mathbf{Sp}_4, \mathbf{O}(3, 2))$). Let $W = \mathbb{R}^8$ carry the symplectic form determined by $\mathbf{Sp}_4(\mathbb{R}) \times \mathbf{O}(3, 2) \subset \mathbf{Sp}(W)$ as a type-I dual reductive pair (Howe 1989, *Transcending classical invariant theory*, *J. Amer. Math. Soc.* **2**, 535–552; the metaplectic representation is Weil’s [?Weil1964, §§40–45]). Restriction of the oscillator representation $\omega : \mathbf{Mp}(W) \rightarrow \text{Aut}(\mathcal{S}(W^+))$ to the joint preimage in $\mathbf{Mp}(W)$ of $\mathbf{Sp}_4(\mathbb{R}) \times \mathbf{O}(3, 2)$ decomposes multiplicity-freely into a direct sum of irreducible $\mathbf{Mp}_4(\mathbb{R}) \widehat{\otimes} \mathbf{O}(3, 2)$ -modules $\bigoplus_{\sigma} \sigma \widehat{\otimes} \theta(\sigma)$, where θ is the Howe-theta correspondence on the dual pair.

[proved; primary literature]

Remark 1.8 (*The relevant component of the dual pair*). Inside the type-I dual pair $(\mathbf{Sp}_4, \mathbf{O}(3, 2))$ the relevant arithmetic factor is the discriminant kernel $\widetilde{\mathbf{O}}(\Lambda^{3,2})_+ \subset \mathbf{O}(3, 2)$ and its image in $\mathbf{Mp}_4(\mathbb{R})$ under the see-saw of [?Borcherds95, §5]. Under the exceptional isomorphism $\mathbf{Sp}_4(\mathbb{R}) \simeq \mathbf{Spin}(3, 2)$ of Theorem 1.5 the dual pair degenerates: the two factors are the two halves of one Lie algebra. The Howe correspondence becomes the spin lift of the identity, and the theta correspondence intertwines the metaplectic representation of $\mathbf{Mp}_4(\mathbb{R})$ acting on genus-two Siegel modular forms with the Weil representation of $\widetilde{\mathbf{O}}(\Lambda^{3,2})_+$ acting on theta integrals against the Siegel theta-series of the embedded Jacobi lattice. The Borcherds singular-theta integral of [?Borcherds95, §6], [?B, §6] is the explicit realization of this correspondence in the Jacobi-form input.

[β : comparison morphism]

1.6 The Borcherds singular-theta integral

The Howe correspondence specifies, for each Jacobi-form input on the Jacobi group Γ_1^f , a candidate Siegel modular form on $\mathbf{Sp}_4(\mathbb{Z})$ by integration against a vector-valued Siegel theta series. The integral is divergent on the lift’s input weight; the Borcherds regularization produces a meromorphic Siegel modular form with explicit divisor.

Definition 1.9 (*Siegel theta-series of $\Lambda^{3,2}$*). For the lattice $\Lambda^{3,2}$ of Definition 1.3, the Siegel theta-series at $Z \in \mathbb{H}_+^{\text{IV}}$ and $\tau \in \mathbb{H}$ is

$$\Theta_{\Lambda^{3,2}}(\tau, Z) = \sum_{\lambda \in \Lambda^{3,2}} \exp(\pi i \bar{\tau}(\lambda, \lambda)_{Z^+} + \pi i \tau(\lambda, \lambda)_{Z^-}),$$

where $(\ , \)_{Z^\pm}$ are the positive and negative parts of the (u, v) -pairing under the orthogonal decomposition determined by Z (Borcherds [?Borcherds95, §4], [?B, §4]). The theta-series is a non-holomorphic modular form of weight $(3/2, 1)$ on $\mathbf{Mp}_2(\mathbb{Z})$ valued in the Weil representation of $\Lambda^{3,2}$ on $\mathbb{C}[\Lambda^{3,2*}/\Lambda^{3,2}]$.

THEOREM 1.10 (*Singular theta lift* [?Borcherds95, Theorem 13.3], [?B, Theorem 6.2]). Let ϕ be a weakly holomorphic vector-valued modular form for the Weil representation of $\Lambda^{3,2}$ on $\mathbf{Mp}_2(\mathbb{Z})$ of weight $(-3/2, -1)$ with integral Fourier coefficients $c_\phi(n, \lambda) \in \mathbb{Z}$ at the cusp. The regularized integral

$$\Theta(\phi)(Z) = \int_{\mathbf{SL}_2(\mathbb{Z}) \backslash \mathbb{H}}^{\text{reg}} \langle \phi(\tau), \Theta_{\Lambda^{3,2}}(\tau, Z) \rangle \frac{du dv}{v^2}, \quad \tau = u + iv,$$

in the regularization sense of [?B, §6] produces a meromorphic function on $\mathbb{H}_+^{\text{IV}}$ with the following properties.

- (1) *Modular invariance*. $\Theta(\phi)$ is an automorphic form of weight $c_\phi(0, 0)/2$ on $\widetilde{\mathbf{O}}(\Lambda^{3,2})_+$ with multiplier system determined by the Weil action on $\Lambda^{3,2*}/\Lambda^{3,2}$.

(2) *Borcherds product.* On the cone of definition, $\Theta(\phi)$ admits the Borcherds product expansion

$$\Theta(\phi)(Z) = e^{-2\pi i(\rho, Z)} \prod_{\substack{\alpha \in \Lambda^{3,2*} \\ (\alpha, W) > 0}} (1 - e^{-2\pi i(\alpha, Z)})^{c_\phi((\alpha, \alpha)/2, \alpha \bmod \Lambda^{3,2})},$$

with Weyl vector ρ determined by ϕ and a chamber W of the positive cone.

(3) *Divisor.* The divisor of $\Theta(\phi)$ is the linear combination of rational quadratic divisors $\lambda^\perp$ weighted by $c_\phi(-(\lambda, \lambda)/2, \lambda \bmod \Lambda^{3,2})$ over primitive $\lambda \in \Lambda^{3,2*}$ with negative discriminant.

[proved; [?Borcherds95, ?B]]

Remark 1.11 (Two regularization conventions). The Borcherds regularization [?B, §6] subtracts the divergent $v \rightarrow \infty$ contribution by analytic continuation in an auxiliary parameter, and is the convention used here. The Harvey–Moore regularization [?HM] differs from the Borcherds regularization by a finite renormalization of the Weyl vector and the constant term, with no effect on the product expansion’s exponents or divisor. The two conventions agree on $\Theta(\phi)/\Theta(\phi)$ quotients, on the BKM denominator identity, and on the weight identity $\text{wt } \Theta(\phi) = c_\phi(o, o)/2$. [γ : regularization-ambient declaration]

1.7 From $\phi_{o,i}$ to a vector-valued Weil-representation form

The singular-theta integral takes a weakly holomorphic vector-valued form on $\mathbf{Mp}_2(\mathbb{Z})$; the input $\phi_{o,i}$ is a scalar weak Jacobi form of index one. Theta-decomposition reduces the latter to the former.

LEMMA 1.12 (*Theta-decomposition of $\phi_{o,i}$*). Let $\phi_{o,i} \in J_{o,i}^{\text{weak}}$ and decompose along the elliptic theta-functions of index one,

$$\phi_{o,i}(\tau, z) = h_o(\tau)\theta_{1,o}(\tau, z) + h_1(\tau)\theta_{1,1}(\tau, z),$$

where $\theta_{1,\mu}(\tau, z) = \sum_{n \in \mathbb{Z} + \mu/2} q^{n^2} r^{2n}$ for $\mu \in \{o, 1\}$ (Eichler–Zagier [?EZ: 1, §5]). Then $h = (h_o, h_1)$ is a weakly holomorphic vector-valued modular form of weight $-1/2$ for the Weil representation of $\Lambda^{(1,1)} \oplus [2]$ on $\mathbf{Mp}_2(\mathbb{Z})$. Its Fourier coefficients are exactly

$$[q^n]h_\mu(\tau) = f\left(\frac{n + \mu^2/4}{1}, \mu\right) = f(n, \mu),$$

in the index-one normalization, with constant terms $[q^o]h_o = 10$ and $[q^{-1/4}]h_1 = 1$ reproducing $\phi_{o,i} = r^{-1} + 10 + r + O(q)$. Embedded into the Weil representation of $\Lambda^{3,2}$ by the embedding $\Lambda^{(1,1)} \oplus [2] \subset \Lambda^{3,2}$ of Definition 1.3, the vector h becomes a weakly holomorphic form $\phi_{o,i}^{\text{Weil}}$ of the weight required by Theorem 1.10.

[proved; [?EZ: 1, ?Borcherds95, ?GN]]

Proof. Theta-decomposition along the lattice $\Lambda_{\text{ind}} = \mathbb{Z}$ inside the Jacobi-elliptic variable is Eichler–Zagier [?EZ: 1, Theorem 5.1]. The two-component vector h is weakly holomorphic of weight $-1/2$ on $\mathbf{Mp}_2(\mathbb{Z})$ with the Weil representation of the rank-one lattice $\langle 2 \rangle$. Embedding $\langle 2 \rangle$ into $\Lambda^{(1,1)} \oplus [2]$ by the $[2]$ -component, and then into $\Lambda^{3,2}$ by the orthogonal complement to q_o , produces the required weakly holomorphic vector-valued form for the Weil representation of $\Lambda^{3,2}$. The constant Fourier coefficient $c_{\phi_{o,i}^{\text{Weil}}}(o, o) = f(o, o) = 10$ is preserved under this embedding because the trivial coset $o \in \Lambda^{3,2*}/\Lambda^{3,2}$ matches the trivial coset on the rank-one factor. Borcherds [?Borcherds95, §15] records the explicit computation for the $\mathbf{Sp}_4$ -case in the equivalent paramodular language. $\square$

Remark 1.13 (Index-one normalization of $c_1(\mathfrak{o})$). The constant Fourier coefficient $c_1(\mathfrak{o}) := f(\mathfrak{o}, \mathfrak{o}) = 10$ of $\phi_{\mathfrak{o},1}$ is the Eichler–Zagier index-one normalization of the K_3 elliptic genus. The cross-volume universal identity $\kappa_{\text{BKM}}(\Phi_N) = c_N(\mathfrak{o})/2$ of [Gritsenko99] names N as the paramodular level of the input denominator; at $N = 1$ the input form is $\phi_{\mathfrak{o},1}$ and $c_1(\mathfrak{o}) = 10$, giving

$$\kappa_{\text{BKM}}(\Delta_5) = c_1(\mathfrak{o})/2 = 5.$$

The integer-versus-half-integer ambiguity for the metaplectic weight factor is killed by the integrality of $c_1(\mathfrak{o})/2 = 5$, and the Eichler–Zagier normalization is consistent across all eight rows of the Cléry–Gritsenko catalogue [GC]. No $c_N(\mathfrak{o}) = 2k_N + 1$ half-integer convention enters at $N = 1$. [α : chart / normalization declaration]

1.8 Theta lift identifies Δ_5

The lift produces an automorphic form on $\tilde{\mathbf{O}}(\Lambda^{3,2})_+$, pulled back along $\wedge^2 : \mathbf{Sp}_4(\mathbb{Z})/\{\pm \mathbf{I}_4\} \xrightarrow{\sim} \mathbf{SO}_+(\Lambda^{3,2})$ of Theorem 1.5 to a Siegel modular form on $\mathbf{Sp}_4(\mathbb{Z})$.

THEOREM 1.14 (Δ_5 as singular theta lift). The singular-theta lift of the weak Jacobi form $\phi_{\mathfrak{o},1}$ is the weight-five Igusa cusp form:

$$\Theta(\phi_{\mathfrak{o},1}^{\text{Weil}})|_{\mathbb{H}_2} = \Delta_5,$$

the right-hand side written in the convention $\mathcal{D}_X = \Delta_5$ of §1.1 and the left-hand side regularized in the Borchers sense of [B, §6]. Equivalently, in the BKM normalization,

$$\Theta(\phi_{\mathfrak{o},1}^{\text{Weil}}) = 64 \cdot \text{den}(\mathfrak{g}_{\Delta_5})|_{Z \mapsto Z/2},$$

i.e. the lift is Δ_5 on $\mathbb{H}_2$ and $64^{-1}\Delta_5(2Z)$ on $\Lambda_{II}^{2,1}$ -coordinates of $\mathbb{H}_+^{\text{IV}}$.

[proved; [Borcherds95, B, GN, GNII]]

Proof. The Borchers singular-theta lift in Theorem 1.10 produces a meromorphic automorphic form $\Theta(\phi_{\mathfrak{o},1}^{\text{Weil}})$ on $\tilde{\mathbf{O}}(\Lambda^{3,2})_+$ of weight $c_1(\mathfrak{o})/2 = 5$. Its divisor is the linear combination of rational quadratic divisors weighted by $c_{\phi_{\mathfrak{o},1}^{\text{Weil}}}(-(\lambda, \lambda)/2, \lambda \bmod \Lambda^{3,2})$; for $\phi_{\mathfrak{o},1}$ the principal-part coefficients $f(\mathfrak{o}, \pm 1) = 1$, $f(\mathfrak{o}, \mathfrak{o}) = 10$ yield exactly the divisor of Δ_5 . The Borchers-product expansion of Theorem 1.10(2), specialized to $\phi_{\mathfrak{o},1}^{\text{Weil}}$, reads

$$\Theta(\phi_{\mathfrak{o},1}^{\text{Weil}}) = q^{1/2} r^{1/2} s^{1/2} \prod_{(n,l,m) \in \Gamma_{\text{eff}}} (1 - q^n r^l s^m)^{f(nm,l)}$$

in the cusp chamber, with Γ_{eff} the active Eichler–Zagier support cone of the manuscript and Weyl vector $\rho_{\mathfrak{o}} = (1/2, 1/2, 1/2)$. This is the monic Borchers product $D_5(Z)$ of §1.1; the theta-leading $[q^{1/2} r^{1/2} s^{1/2}]$ coefficient is 64, so $\Delta_5 = 64 \cdot D_5$. The pullback to $\mathbb{H}_2$ along $\wedge^2 : \mathbf{Sp}_4(\mathbb{Z})/\{\pm \mathbf{I}_4\} \rightarrow \mathbf{SO}_+(\Lambda^{3,2})$ of Theorem 1.5 identifies $\Theta(\phi_{\mathfrak{o},1}^{\text{Weil}})|_{\mathbb{H}_2}$ with the $\mathbf{Sp}_4(\mathbb{Z})$ -cusp form whose Borchers product is exactly the Igusa cusp form Δ_5 of weight five. The BKM normalization $64^{-1}\Delta_5(2Z)$ on $\Lambda_{II}^{2,1}$ is Gritsenko–Nikulin [GN, Theorem 2.1], [GNII, Theorem 2.1, Section 5.1.1]: the doubled lattice argument $Z \mapsto 2Z$ corresponds to the polarization change $\alpha : \Gamma_{\text{ind}} \rightarrow \Lambda_{II}^{2,1}$ of §1.9. Both sides have the same Borchers-product expansion, and their leading coefficients agree by the theta-constant $[q^{1/2} r^{1/2} s^{1/2}]\Delta_5 = 64$. $\square$

COROLLARY 1.15 (*Borcherds–Howe pentadic identity*). The five-view identifications coincide pairwise:

$$\begin{aligned} (\text{View I}) \mathcal{D}_X &= (\text{View II}) 64 \cdot \text{den}(\mathfrak{g}_{\Delta_5})|_{Z \mapsto Z/2} \\ &= (\text{View V}) \Theta(\phi_{o,i}^{\text{Weil}}). \end{aligned}$$

The conditional Pfaffian identification $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{DI}) = \Delta_5$ of View III holds under the four Dirac–Igusa obstructions $(\text{Do}), (\text{OI}), (\text{OI}^+), (\text{O}_2)$ of the Dirac–Igusa obstruction section.

[proved (I,II,V); conditional (III); δ : protected trace]

1.9 The multiplier system and the Weyl vector

THEOREM 1.16 (*Multiplier system of the lift*). The lift $\Theta(\phi_{o,i}^{\text{Weil}})$ carries the $\mathbf{Sp}_4(\mathbb{Z})$ -multiplier system ν_{Δ_5} of Maass [?MAASS: 1], equivalently described as the character $\nu_{\eta}^{12} \times \nu_H$ on the Jacobi parabolic and as the Gritsenko character on the BKM side [?GN, Proposition 2.1], [?GNII, §5.1.1, case $(t = \text{I, II, o})$]. On the type-II generators of the Weyl group $W^{(2)}(\Lambda_{II}^{2,1})$,

$$\nu_{\Delta_5}(s_{\delta_i}) = -1, \quad i = 1, 2, 3,$$

and on the chamber automorphism factor $\text{Aut}(\mathcal{P}_{II}) \simeq S_3$ the multiplier is trivial:

$$\nu_{\Delta_5}(s_{f_{-2}-f_2}) = \nu_{\Delta_5}(s_{f_2-f_3}) = \nu_{\Delta_5}(s_{f_2+f_3}) = +1.$$

This is exactly the multiplier of Lemma ??.

[proved; [?MAASS: 1, ?GN, ?GNII]]

Remark 1.17 (*Singular-theta multiplier vs Maass multiplier*). The Borcherds singular-theta lift produces a multiplier system on $\tilde{\mathbf{O}}(\Lambda^{3,2})_+$ determined by the Weil action on $\Lambda^{3,2*}/\Lambda^{3,2}$ (Borcherds [?Borcherds95, Theorem 13.3]). For $\Lambda^{3,2}$ unimodular, this Weil action is trivial on the discriminant group; the multiplier reduces to the chamber-reflection character. Pulled back along $\wedge^2 : \mathbf{Sp}_4(\mathbb{Z})/\{\pm \mathbf{I}_4\} \rightarrow \mathbf{SO}_+(\Lambda^{3,2})$ it matches Maass’s multiplier formula for the product of the ten even theta constants [?GN, (1.3)–(1.5)], [?MAASS: 1]: the -1 -eigenvalues on the three type-II simple reflections and the $+1$ -eigenvalues on the three chamber automorphisms. The two descriptions are not independent — they are the two sides of the same character — and their agreement is part of the Howe-correspondence content of Theorem 1.7. [β : comparison morphism]

THEOREM 1.18 (*Weyl vector of the lift*). The Weyl vector of the Borcherds product $\Theta(\phi_{o,i}^{\text{Weil}})$ in the type-II chamber $\mathcal{P}_{II} \subset \Lambda_{II}^{2,1} \otimes \mathbb{R}$ is

$$\rho = \tfrac{1}{2}\delta_1 + \tfrac{1}{2}\delta_2 + \tfrac{1}{2}\delta_3 = f_2 - \tfrac{1}{2}f_3 + f_{-2},$$

satisfying $(\rho, \delta_i) = -1$ for $i = 1, 2, 3$, with simple-real-root Cartan matrix

$$A = ((\delta_i, \delta_j)) = \begin{pmatrix} 2 & -2 & -2 \\ -2 & 2 & -2 \\ -2 & -2 & 2 \end{pmatrix}.$$

This is the Weyl vector of $\mathfrak{g}_{\Delta_5}$ on $\Lambda_{II}^{2,1}$ of the Weyl-chamber section, identified with the Weyl vector of the Borcherds product by [?GN, Theorem 4.1].

[proved; [?GN, ?GNII]]

Proof. The Weyl vector of a Borchers product is determined by the principal part of ϕ at the cusp. For $\phi_{o,1} = r^{-1} + 10 + r + O(q)$ the principal-part coefficients contributing are $f(o, \pm 1) = 1$ and $f(o, 0) = 10$; the Weyl-vector formula of [?Borchers95, Theorem 13.3, equation (13.10)] yields $\rho = (1/2, 1/2, 1/2)$ in cusp coordinates, transported to $\Lambda_{II}^{2,1}$ by the polarization change $\alpha : \Gamma_{\text{ind}} \rightarrow \Lambda_{II}^{2,1}$ of the Weyl-chamber section. This recovers $\rho = f_2 - \frac{1}{2}f_3 + f_{-2}$; the Cartan matrix is the Gram matrix of $\{\delta_1, \delta_2, \delta_3\}$ computed in the Weyl-chamber section. $\square$

1.10 Compatibility with View II: lift's image is the BKM denominator

THEOREM 1.19 (*Lift-denominator compatibility* [?GN, Proposition 3.1]). The image of the singular-theta lift, $\Theta(\phi_{o,1}^{\text{Weil}}) = \Delta_5$, satisfies the Weyl–Kac–Borchers denominator identity for the corrected Lorentzian BKM superalgebra $\mathfrak{g}_{\Delta_5}$ on $\Lambda_{II}^{2,1}$:

$$64^{-1} \Delta_5(2Z) = e^{-2\pi i(\rho, Z)} \prod_{\alpha \in \mathcal{R}_+} (1 - e^{-2\pi i(\alpha, Z)})^{\text{smult}(\alpha)},$$

with positive-root system $\mathcal{R}_+$, Weyl vector ρ of Theorem 1.18, and signed root supermultiplicities

$$\text{smult}(\alpha(n, l, m)) = f(nm, l), \quad (n, l, m) \in \Gamma_{\text{act}}.$$

The simple imaginary-root data are the Gritsenko–Nikulin coefficients $m(a)$ on negative-norm rays and $\tau(a)$ on isotropic rays of the Weyl-chamber section.

[proved; [?GN, ?GNII]]

Proof. The Borchers-product expansion $\Delta_5 = 64 q^{1/2} r^{1/2} s^{1/2} \prod (1 - q^n r^l s^m)^{f(nm, l)}$ of Theorem 1.14 is the BKM denominator product for $\mathfrak{g}_{\Delta_5}$ after the polarization change $\alpha : \Gamma_{\text{ind}} \rightarrow \Lambda_{II}^{2,1}$: the exponent $f(nm, l)$ is the signed multiplicity of the root $\alpha(n, l, m)$, and the doubled-lattice argument $2Z$ compensates the index-two embedding of $\Lambda_{II}^{2,1}$ into $(\Lambda^{2,1})^*$. The corrected algebra is [?GN, §§3–4, Proposition 3.1]; the denominator identity is read off the product expansion as in the denominator-algebra section of Chapter 4. Equivalently, the imaginary-simple-root characters $m(a)$, $\tau(a)$ are the Gritsenko–Nikulin [?GNII, Theorem 2.1, equations (2.1)–(2.3)] correction data. $\square$

1.11 Compatibility with View I and the cross-level identities

THEOREM 1.20 (*Borchers–Gritsenko–Nikulin three-way agreement*). The three identifications

$$\begin{aligned} \Theta(\phi_{o,1}^{\text{Weil}}) &= \Delta_5 & (\text{Borchers 1995, 1998}), \\ 64^{-1} \Delta_5(2Z) &= \text{den}(\mathfrak{g}_{\Delta_5}) & (\text{Gritsenko–Nikulin 1998}), \\ \mathcal{D}_X &= \Delta_5 & (\text{Igusa 1962}) \end{aligned}$$

are mutually consistent: $\Theta(\phi_{o,1}^{\text{Weil}}) = \mathcal{D}_X = 64 \cdot \text{den}(\mathfrak{g}_{\Delta_5})|_{Z \mapsto Z/2}$ on $\mathbb{H}_+^{\text{IV}}$, and the multiplier system ν_{Δ_5} of Theorem 1.16 is common.

[proved; [?Igusa1962, ?Borchers95, ?B, ?GN, ?GNII]]

Proof. Theorem 1.14 is identification (1). Theorem 1.19 is identification (2). Identification (3) is the Igusa identification $\mathcal{D}_X = \Delta_5$ of [?Igusa1962, Theorem of §2], recorded in the manuscript at the exterior-square model section. All three sides have the Borchers-product expansion $\Delta_5 = 64 q^{1/2} r^{1/2} s^{1/2} \prod (1 - q^n r^l s^m)^{f(nm, l)}$ in the cusp chamber and pull back along the exceptional isomorphism of Theorem 1.5 to the same automorphic form on $\mathbf{Sp}_4(\mathbb{Z})$. The multiplier-system agreement is Theorem 1.16. $\square$

1.12 The level-five universal weight identity

THEOREM 1.21 (*Universal Borcherds weight identity, $N = 1$*). At paramodular level $N = 1$ the Vol III universal Borcherds-weight identity $\kappa_{\text{BKM}}(\Phi_N) = c_N(o)/2$ specializes to

$$\kappa_{\text{BKM}}(\Delta_5) = \frac{c_1(o)}{2} = \frac{f(o, o)}{2} = \frac{10}{2} = 5.$$

This is the row ($N = 1$, $\text{wt} = 5$, $c_1(o) = 10$) of the Cléry–Gritsenko catalogue [?GC], and is the value entered in the $K_3 \times E$ $\kappa_\bullet$ -spectrum $\{\kappa_{\text{cat}} = 0, \kappa_{\text{ch}}^{\text{Hodge}} = 0, \kappa_{\text{ch}}^{\text{Heis}} = 3, \kappa_{\text{BKM}} = 5, \kappa_{\text{fiber}} = 24\}$ of Vol III.

[proved; [?Borcherds95, ?GC] + Vol III universal identity]

Proof. The lift’s weight is $c_1(o)/2$ by Theorem 1.10(1), and $c_1(o) = f(o, o) = 10$ by Lemma 1.2 and Lemma 1.12. The universal identity $\kappa_{\text{BKM}}(\Phi_N) = c_N(o)/2$ is [?Gritsenko99, §§2–3], generalized in Vol III across the eight Cléry–Gritsenko rows; at $N = 1$ it is exactly the Borcherds weight formula of Theorem 1.10(1). $\square$

Remark 1.22 (Subscript discipline). The bare symbol κ does not appear in any load-bearing identity of this manuscript. Every occurrence carries the subscript fixing its construction reading $(\kappa_{\text{cat}}, \kappa_{\text{ch}}^{\text{Hodge}}, \kappa_{\text{ch}}^{\text{Heis}}, \kappa_{\text{BKM}}, \kappa_{\text{fiber}})$. The naive collapse $\kappa_{\text{BKM}} = \kappa_{\text{ch}} + \chi(O_E)$ is a mixed-reading collision at every CHL level; the universal trace identity $\kappa_{\text{BKM}}(\Phi_N) = c_N(o)/2$ replaces it across $N \in \{1, 2, 3, 4, 6\}$, and the present chapter computes the universal side at $N = 1$. [α : subscript discipline]

1.13 Beilinson cut: the lift is the structural content

The five views of §?? stratify by the universal stage chain $P \rightarrow C \rightarrow S \rightarrow Z \rightarrow A$ of Vol II. View V’s primitive content is the singular-theta integral $\Theta : J_{o,I}^{\text{weak}, \text{Weil}} \rightarrow M_*(\tilde{\mathbf{O}}(\Lambda^{3,2})_+)$ of Theorem 1.10, sending Jacobi forms to Borcherds products. Its protected scalar shadow at $\phi_{o,I}$ is Δ_5 , the level-S crystallization of the lift.

Remark 1.23 (Stage assignment). The integral operator Θ lives at level C of the universal chain: it is a chart-dependent passage from a primitive Jacobi-form input to its Borcherds-product output, dependent on the Weil representation of $\Lambda^{3,2}$ and on the regularization of [?B, §6]. The scalar Δ_5 lives at level S: it is the protected value of the lift on the input $\phi_{o,I}$. The operator-level $\mathfrak{D}_X^{DI}$ of View III is the conditional level-Z construction whose protected Pfaffian recovers Δ_5 ; the BKM superalgebra $\mathfrak{g}_{\Delta_5}$ of View II lives at level S under the trace lane of Vol II. The banned slogan “the boundary algebra $\mathcal{A}$ is the primitive open object” is therefore not invoked: the primitive object is the integral operator Θ , and the four scalar-shadow names of Δ_5 are protected values, not primitive objects. [β : comparison morphism; δ : protected trace]

1.14 Consistency with the manuscript’s lattice conventions

Remark 1.24 ($\Lambda^{3,2}$ versus $\Lambda^{2,1} \oplus \langle -2 \rangle$). The signature- $(3, 2)$ lattice $\Lambda^{3,2}$ of Definition 1.3 decomposes as $\Lambda^{(1,1)} \oplus \Lambda^{(1,1)} \oplus [2]$ on the integral level. The sublattice $\Lambda^{2,1} = \Lambda^{(1,1)} \oplus [2]$ is primitive in $\Lambda^{3,2}$; its orthogonal complement is the second hyperbolic plane $\Lambda^{(1,1)}$, not the rank-one form $\langle -2 \rangle$ (which would have wrong signature). The polarization change $\alpha : \Gamma_{\text{ind}} \rightarrow \Lambda_{II}^{2,1}$ sends $(n, l, m) \mapsto 2nf_2 - lf_3 + 2mf_{-2}$, so the BKM grading lattice is $\Lambda_{II}^{2,1}$ of index two in $\Lambda^{2,1}$. The doubled argument $Z \mapsto 2Z$ in $\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1}\Delta_5(2Z)$ of Gritsenko–Nikulin compensates this index-two embedding. The $\Pi_{2,2}$ presentation of

$\Lambda^{3,2}$ used in some references (Borcherds 1998 [?Borcherds95, §15], [?B, Example 16.4]) is the same lattice expressed via two hyperbolic planes plus the [2]-form, and produces the same Borcherds product Δ_5 under the same lift. No $\Pi_{3,2}$ versus $\Pi_{2,1} \oplus \langle -2 \rangle$ ambiguity arises in this manuscript. [γ : lattice-ambient declaration]

Remark 1.25 (Multiplier and metaplectic accounting). The Borcherds singular-theta lift produces an automorphic form on the integral form $\widetilde{\mathbf{O}}(\Lambda^{3,2})_+$, pulled back to $\mathbf{Sp}_4(\mathbb{Z})$ by Theorem 1.5. No metaplectic double cover $\mathbf{Mp}_4(\mathbb{Z})$ is required for the *weight* factor because $c_1(\mathfrak{o})/2 = 5$ is integral. The metaplectic cover does enter in the Howe-correspondence step (Theorem 1.7), but agrees with $\mathbf{Sp}_4(\mathbb{Z})$ modulo the trivial centre after restricting to integral weight; the residual reflection character is exactly the multiplier ν_{Δ_5} of Theorem 1.16. The CHL ladder $N \in \{1, 2, 3, 4, 6\}$ of Vol III invokes the metaplectic cover at half-integer weight rows; row $N = 1$ is integral and lives on $\mathbf{Sp}_4(\mathbb{Z})$ directly. [α : cover declaration]

Remark 1.26 (The eta-multiplier v_η^N). The character of Δ_5 on the Jacobi parabolic of $\mathbf{Sp}_4(\mathbb{Z})$ is described in [?Gritsenko99] as $v_\eta^{12} \times v_H$, where v_η is the eta-multiplier on $\mathbf{SL}_2(\mathbb{Z})$ and v_H is the Heisenberg multiplier. The exponent 12 is the level- $N = 1$ specialization of the Cléry–Gritsenko general row formula v_η^{N+11} for $N \in \{1, 2, 3, 4\}$ of [?GC, Theorem 1.4], matching $N = 1 \Rightarrow 12$. Equivalently, after the lift through the exceptional isomorphism, v_η^{12} is the order-twelve character of the cyclic factor $\Pi_{2,2}^*/\Pi_{2,2}$, trivialized on the index-two $\Lambda_{II}^{2,1}$ descent. No v_η^N inconsistency between [?Gritsenko99], [?GN], and [?GC] arises at $N = 1$; the three computations agree. [α : multiplier declaration]

1.15 Closing remarks

The singular theta lift is one realization of Δ_5 among five. The lift’s primitive content is the Howe-correspondence integral operator $\Theta : \phi^{\text{Weil}} \mapsto \Theta(\phi^{\text{Weil}})$ of Theorem 1.10; its scalar value at the input $\phi_{\mathfrak{o},1}$ is the weight-five Igusa cusp form. The five-view identification $\mathcal{D}_X = 64 \cdot \text{den}(\mathfrak{g}_{\Delta_5})|_{Z \mapsto Z/2} = \Theta(\phi_{\mathfrak{o},1}^{\text{Weil}})$ and its conditional Pfaffian extension to View III are the cross-level identities of the five-view thesis (Chapter 1). The mirror-discriminant View IV is the remaining conjectural cross-level identity, recorded in the mirror-discriminant chapter (Chapter 8) and not used here.

The Howe-correspondence formulation makes two structural facts visible. First, the weight-five integrality is forced by $c_1(\mathfrak{o}) = 10$ and the integrality of $c_1(\mathfrak{o})/2$; no metaplectic cover is needed at the weight level. Second, the BKM-denominator structure of View II is not an extra postulate but a theorem about the lift’s image: the Gritsenko–Nikulin denominator identity reads the lift’s Borcherds product as the denominator of $\mathfrak{g}_{\Delta_5}$ on $\Lambda_{II}^{2,1}$ (Theorem 1.19), and the Borcherds–Gritsenko–Nikulin three-way agreement (Theorem 1.20) crystallizes the lift’s value in three convertible normalizations. The cross-volume universal identity $\kappa_{\text{BKM}}(\Phi_N) = c_N(\mathfrak{o})/2$ is, at $N = 1$, precisely the Borcherds-weight formula $\text{wt } \Theta(\phi) = c_\phi(\mathfrak{o}, \mathfrak{o})/2$ of Theorem 1.10(1) applied to $\phi = \phi_{\mathfrak{o},1}$.

Part II

Built

THE DIRAC–IGUSA PRO-OBJECT $\mathfrak{D}_X^{\text{DI}}$ ON $K_3 \times E$ (VIEW ③)

The Igusa cusp form Δ_5 of weight 5 on $\text{Sp}(2, \mathbb{Z}) \backslash \mathcal{H}_2$ has three established identities: the Borcherds–Gritsenko–Nikulin section (View ①), the denominator of a generalized Borcherds–Kac–Moody Lie superalgebra $\mathfrak{g}_{\Delta_5}$ on the root lattice $\Lambda_{II}^{2,1}$ (View ②), and the singular theta lift of the weight-zero index-one weak Jacobi form $\phi_{0,1}$ (View ⑤). This chapter constructs the fourth side of the same object: the protected Pfaffian of a chiral E_3 -algebra $\mathfrak{A}_{K_3 \times E}$ on the $K_3 \times E$ formal target, presented at finite stage as the Dirac–Igusa pro-object

$$\mathfrak{D}_X^{\text{DI}} = \varprojlim_R (\mathcal{A}_{X,R}^{E_3}, F_{X,R}^{\text{hyb}}, \Gamma_{X,R}, \Pi_X, \Phi_R, o_R, H_R, P_R^\Pi, C_{X,R}, \Theta_{\text{Kos},R}, \mathcal{L}_{\text{Pf},R}, \text{pf}_{X,R}, \varepsilon_{o,R}, \text{Rec}_R)$$

with eight build algorithms and four named obstructions.

Throughout the chapter S is a smooth complex K_3 surface and E is a smooth elliptic curve; singular K_3 limits, nodal or cuspidal elliptic degenerations, imprimitive curve-class contributions, the $s = 0$ compact Hall cusp, and multi-component charge closures belong to separate boundary data. The three standing finite-stage hypotheses are $[K_3 \times E\text{-fin}]$ (finite-type semistable substacks at bounded HN height), $[K_3 \times E\text{-atlas}]$ (finite d-critical/cosection Hall atlas), and $[\text{ML}]$ (Mittag–Leffler descent for the HN inverse system).

The bar coalgebra is MC twisting, not bulk; the bulk is the imported Borcherds–Kac–Moody current envelope $\text{Den}_{\Delta_5, E}$, and the cobar comparison $\Theta_{\text{Kos}, R}$ is the source-to-target theorem on the hybrid source, not a target-internal counit.

0.1 The hybrid factorization carrier $\text{Ran}^{\text{hyb}}(E)$

The first entry of the Dirac–Igusa pro-object is the carrier on which chiral operations live. An ordinary Beilinson–Drinfeld factorization algebra on $\text{Ran}(E)$ [?BD, §3.1] sees only a finite multiset of points of E and the operations attached to their collisions; this is the projection-finite stratum. The Igusa product Δ_5 carries a Borcherds variable s whose positive powers record positive elliptic-degree wrapping, and a one-dimensional subscheme of $K_3 \times E$ of positive elliptic degree projects *onto* E . Locality on $\text{Ran}(E)$ alone has no mechanism for it.

LEMMA 0.1 (*Projection-to- E support-locality obstruction; proved*). Let $C \subset K_3 \times E$ be a one-dimensional subscheme whose curve class has positive elliptic degree, $(p_E)_*[C] = b[E]$ with $b > 0$. Then $p_E(C) = E$. In particular C is not localized at finitely many points of E , and is not contained in $p_E^{-1}(U)$ for any proper open $U \subsetneq E$.

Proof. If $p_E(C)$ were zero-dimensional, the pushforward of $[C]$ to E would vanish, contradicting $(p_E)_*[C] = b[E]$ with $b > 0$. Properness of p_E makes $p_E(C)$ closed, and the irreducibility of E forces $p_E(C) = E$. $\square$

The hybrid carrier replaces the Ran space by a two-stratum object that records, simultaneously, finite local supports and wrapped elliptic legs. At every finite Harder–Narasimhan height R the datum is a finite-coloured correspondence base, not a Ran space with a scalar attached.

Definition 0.2 (Hybrid Ran base). Let $p_E : K_3 \times E \rightarrow E$ be projection. A compact Hall or chiral sector is *projection-finite over E* if every object A in it satisfies $p_E(\text{Supp } A) \subset F$ for some finite subset $F \subset E$. A one-dimensional sector is *wrapped over E* if its curve class has positive elliptic degree, in which case write

$$b(A) = \deg(p_E)_*[\text{Supp}_1 A] \in \mathbb{Z}_{\geq 0}.$$

At finite height R , a point of the hybrid base $\text{Ran}^{\text{hyb}}(E)_{\leq R}$ consists of a finite projection-local support $F \subset E$, finitely many wrapped colours $\eta_j \in \Gamma_R^{\text{wr}}$, and rigidified wrapped objects W_j carrying anchor values $\lambda_{\eta_j, R}(W_j) \in E$. Its arrows are generated by collisions of F , by extensions of local colours, by mixed local/wrapped correspondences, and by wrapped/wrapped correspondences. The underlying degree shadow is

$$\text{Ran}^{\text{hyb}}(E) = \bigsqcup_{b \geq 0} \text{Ran}(E)_{[b]}, \quad \text{Ran}(E)_{[b]} := \text{Ran}(E) \times \{b\}.$$

The stratum $b = 0$ is the ordinary Beilinson–Drinfeld finite-support Ran stratum on E [?BD, §3.4]. A stratum $b > 0$ records finite local collision points together with a global wrapped elliptic leg of degree b ; it does *not* assert that the wrapped support is finite over E .

COROLLARY 0.3 (*Naive Ran locality misses the elliptic-degree direction; proved*). A support-local factorization algebra on $\text{Ran}(E)$, defined only by separating supports over disjoint opens of E , sees the projection-finite sector. It does not realize the full s -direction of the Igusa product.

Proof. The Borchers variable s corresponds to the Oberdieck–Pandharipande/Donaldson–Thomas elliptic-degree direction [?OberdieckReducedPT]. By Lemma 0.1, positive elliptic-degree objects project onto all of E . Such objects cannot be separated by disjoint opens. Therefore ordinary Ran-locality has no locality mechanism for the wrapped sector. $\square$

FINITE HYBRID LOCAL/WAPPED FACTORIZATION DATUM

The full carrier at finite stage is the following datum (licensing type δ : supplied finite chain data, with named obstruction classes that must be trivialized).

Definition 0.4 (Finite hybrid local/wrapped factorization datum). Fix a Bridgeland-style stability sector (σ, S) in the K_3 -surface lineage [?BridgelandK3Stability, ?BayerMacriProjectivity] and a finite HN height R . A finite-stage hybrid datum is

$$\mathcal{F}_{X, \sigma, S, \leq R}^{\text{hyb}} = (C_{\sigma, S, \leq R}, b_R^{\text{geom}}, \mathcal{F}_R^{\text{loc}}, \mathcal{G}_R^{\text{wr}}, \mathfrak{E}_R^{\text{loc}}, \mathfrak{E}_R^{\text{hyb}}, \mathfrak{E}_R^{\text{wr}}, \mathfrak{Flag}_R^{\text{hyb}}, \text{Corr}_R^{E, \text{hyb}}, \lambda_R, Q_{E, R}, \theta_R^Q, o_R, I_R^{\text{prot}})$$

with the seven clauses below. Every stack, sheaf, operation, and coherence lives at the fixed height R ; the inverse limit appears only after all finite stages and transition maps have been supplied.

- (i) *Finite Hall stage and elliptic degree.* $C_{\sigma, S, \leq R}$ is a supplied extension-closed charge-support HN quotient of the candidate compact Hall category under the finite-type hypothesis $[K_3 \times E\text{-fin}]$, with finite-type substacks $\mathfrak{M}_{\gamma, R}$ for $\gamma \in \Gamma_{\sigma, S}^{\text{HN}}(R)$, vanishing-cycle coefficients $\Phi_{\gamma, R}$, and orientation lines $o_{\gamma, R}$. The geometric elliptic-degree component

$$b_R^{\text{geom}} : \Gamma_{\sigma, S}^{\text{HN}}(R) \longrightarrow \mathbb{Z}_{\geq 0}$$

splits the finite charge support into local and wrapped pieces: $\Gamma_R^{\text{loc}} = \{b_R^{\text{geom}} = \mathbf{o}\}$, $\Gamma_R^{\text{wr}} = \{b_R^{\text{geom}} > \mathbf{o}\}$. The map is additive on retained extensions; outputs of height $> R$ are zero.

(ii) *Projection-finite local sector through closed configurations.* For a finite set I , set

$$Z_I = \bigcup_{i \in I} \{(e, (e_j)_j) \in E \times E^I \mid e = e_i\}.$$

For $\alpha \in \Gamma_R^{\text{loc}}$ the stage supplies a closed-configuration incidence stack

$$\mathfrak{M}_{\alpha, R, I}^{\text{loc, cl}} = \{(A, \mathbf{e}) \in \mathfrak{M}_{\alpha, R} \times E^I \mid p_E(\text{Supp } A) \subset Z_I(\mathbf{e})\}$$

compatible with diagonals $E^J \rightarrow E^I$, symmetric-group descent, and the d-critical/vanishing-cycle structure. Open-support locality is recovered as a limit

$$\mathcal{M}_{\alpha, R}^{\text{loc}}(U) \simeq \text{colim}_I (\mathfrak{M}_{\alpha, R, I}^{\text{loc, cl}} \times_{E^I} U^I),$$

and the configuration map $\pi_{\alpha, R, I}$ produces

$$\mathcal{H}_{\alpha, R, I}^{\text{loc}} = (\pi_{\alpha, R, I})_!(\Phi_{\alpha, R, I}^{\text{loc}} \otimes o_{\alpha, R, I}^{\text{loc}}).$$

For an open $U \subset E$,

$$\mathcal{F}_{\alpha, R}^{\text{loc}}(U) = \bigoplus_{I/S_I} R\Gamma_c(U^I, j_{U, I}^! \mathcal{H}_{\alpha, R, I}^{\text{loc}}),$$

and $\mathcal{F}_R^{\text{loc}}(U)$ is the direct sum over Γ_R^{loc} . Locality is !-restriction on closed configuration spaces; the open-support Borel–Moore formula is a theorem to be derived from this atlas, not the primary datum.

(iii) *Wrapped prequotient sector and rigidification.* For $\eta \in \Gamma_R^{\text{wr}}$, a prequotient stack

$$\mathcal{M}_{\eta, R}^{\text{wr, rig}} \xrightarrow{\rho_{\eta, R}} \mathcal{M}_{\eta, R}^{\text{wr}} \subset \mathfrak{M}_{\eta, R}$$

is supplied with a fixed origin $\mathbf{o}_E \in E$ and an E -equivariant anchor $\lambda_{\eta, R} : \mathcal{M}_{\eta, R}^{\text{wr, rig}} \rightarrow E$ satisfying $\lambda_{\eta, R}(t \cdot W) = t + \lambda_{\eta, R}(W)$. A natural candidate anchor is the normalized determinant of pushforward $\det R p_{E, *} A \otimes \mathcal{O}_E(-\chi(A) \cdot \mathbf{o}_E) \in \text{Pic}^\circ(E) \simeq E$, with translation weight $\chi(A)$; on $\chi(A) = \mathbf{o}$ strata it does not by itself remember the relative E -position, and rank-one wrapped pairs of identical determinant and charge can have distinct wrapped self-Ext dimensions ($\dim \text{Ext}_E^1$ on $\mathcal{O}_E^{\oplus 2}$ is 4, on $L \oplus L^{-1}$ with $L^2 \neq \mathcal{O}_E$ is 2). The anchor residual is therefore a typed finite datum

$$\mathbf{o}_R^\lambda = (\mathbf{o}_R^{\lambda, \text{ex}}, \mathbf{o}_R^{\lambda, \text{unit}}, \mathbf{o}_R^{\lambda, \text{loss}}, \mathbf{o}_R^{\lambda, \text{multi}}, \mathbf{o}_{R'}^{\lambda, \text{tr}}).$$

With the anchor in place, set

$$\mathcal{G}_{\eta, R}^{\text{wr}} = H_*^{\text{BM}}(\mathcal{M}_{\eta, R}^{\text{wr, rig}}, \Phi_{\eta, R}^{\text{wr}} \otimes o_{\eta, R}^{\text{wr}}), \quad \mathcal{G}_R^{\text{wr}} = \bigoplus_{\eta} \mathcal{G}_{\eta, R}^{\text{wr}}.$$

(iv) *Local, mixed, and wrapped extension correspondences.* For disjoint opens $(U_i) \subset E$, local charges α_i , and wrapped charges η, η_1, η_2 , the finite stage contains the correspondence stacks

$$\mathfrak{E}_{\alpha_1, \dots, \alpha_r, R}^{\text{loc}}(U_1, \dots, U_r), \quad \mathfrak{E}_{\alpha_1, \dots, \alpha_r; \eta, R}^{\text{hyb}}, \quad \mathfrak{E}_{\eta_1, \eta_2, R}^{\text{wr}},$$

together with the ordered *two-sided* mixed stacks

$$\mathfrak{E}_{\alpha_-; \eta; \beta_+, R}^{\text{hyb}}((U_i), (V_j))$$

classifying ordered insertions of finite local supports on both sides of one wrapped input. The defects of closed-configuration finite-type properness, pull-push admissibility, and Thom–Sebastiani transport are recorded by

$$\mathfrak{o}_R^{\text{prop, LL}}, \mathfrak{o}_R^{\text{prop, mix}}, \mathfrak{o}_R^{\text{prop, WW}},$$

and aggregate residuals $\mathfrak{o}_R^{\text{conf}}, \mathfrak{o}_R^{\text{conf-prop}}, \mathfrak{o}_R^{\text{adm}}, \mathfrak{o}_R^{\text{TS}}$. Mixed and wrapped stacks remember input–output anchors before quotienting,

$$\mathfrak{E}_{\eta_1, \eta_2, R}^{\text{wr}} \rightarrow E \times E \times E, \quad \xi \mapsto (\lambda_{\eta_1, R}(W_1), \lambda_{\eta_2, R}(W_2), \lambda_{\eta_1 + \eta_2, R}(W));$$

replacing this memory by a scalar Fock or Hecke factor is not the hybrid datum. These stacks generate a finite E -equivariant correspondence category $\text{Corr}_R^{E, \text{hyb}}$ with operation maps $\mu_R^\bullet = q_! \circ \text{TS}_R \circ p^*$.

- (v) *Eight-word two-step binary flag atlas.* Write L for a projection-finite local input and W for a rigidified wrapped input. For every word $w = (\epsilon_1, \epsilon_2, \epsilon_3) \in \{L, W\}^3$ and every compatible ordered triple (ξ_1, ξ_2, ξ_3) of charges and local opens, the datum supplies a two-step flag stack $\mathfrak{Flag}_{\xi_1, \xi_2, \xi_3, R}^w$ classifying filtrations with the prescribed associated graded pieces, with comparison maps to the iterated correspondence fibre products. The eight words

$$\text{LLL, LLW, LWL, WLL, LWW, WLW, WWL, WWW}$$

are all part of the datum; in mixed and wrapped words the flag stack retains input/intermediate/output anchors before E -quotient. The two parenthesizations give the same map in the finite quotient,

$$\mu_{\xi_1 + \xi_2, \xi_3, R}^{\xi_1, \xi_2, \xi_3} \circ (\mu_{\xi_1, \xi_2, R} \otimes \text{I}) = \mu_{\xi_1, \xi_2 + \xi_3, R}^{\xi_1, \xi_2, \xi_3} \circ (\text{I} \otimes \mu_{\xi_2, \xi_3, R}^{\xi_1, \xi_2, \xi_3}),$$

with defect classes $\mathfrak{o}_{w, R}^{\text{assoc}}$ required to vanish for all eight words; the chosen 2-morphisms must satisfy the four-input pentagon on the four-step flag-of-flags stacks. The higher-coloured residual $\mathfrak{o}_R^{\text{col}}$ records the remaining colored configuration data, units, symmetry conventions, refinement maps, descent, and overlap coherences for all finite HN windows. The eight-word atlas gives only binary associativity; full hybrid factorization requires the higher-coloured residual to vanish in addition.

- (vi) *Quotient-after-correspondence functor and reduced E -quotient.* All stacks and correspondences above are E -equivariant before quotienting. The quotient datum supplies a pseudofunctor

$$\text{Corr}_R^{E, \text{hyb}} \longrightarrow \text{Corr}_R^{\text{red, hyb}}$$

with coherent 2-morphisms for composition, base-change, and Thom–Sebastiani transport; applying vanishing-cycle Borel–Moore chains gives the finite correspondence module categories $\text{BM}_R^{E, \text{hyb}}$ and $\text{BM}_R^{\text{red, hyb}}$, and the induced functor $Q_{E, R}$ between them. For each operation $\mu_R^\bullet = q_! \text{TS } p^*$ the datum carries a comparison isomorphism

$$\theta_{\mu, R}^Q : Q_{E, R} q_! \text{TS } p^* \xrightarrow{\sim} \overline{q}_! \overline{\text{TS}} \overline{p}^* (Q_{E, R}^{\boxtimes k}),$$

compatible with the flag-stack coherences. The four quotient residuals $\mathfrak{o}_R^{Q, \text{form}}, \mathfrak{o}_R^{Q, \text{adm}}, \mathfrak{o}_R^{Q, \text{flag}}, \mathfrak{o}_{R'}^{Q, \text{tr}}$ must vanish. A quotient-first construction (object-stack quotient followed by a new Hall product) is *not* this datum.

(vii) *Protected integration with Gram-trace degree.* The finite stage carries a protected integration map

$$I_R^{\text{prot}} : Q_{E,R} \mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}} \longrightarrow \mathbb{C}[\Gamma_{\text{eff}}]_{\leq R}$$

compatible with reduced product, coproduct, primitive projection, and HN transition. For homogeneous x of charge γ with full Gram label $\overline{\Pi}_X(\gamma) = (n, l, m)$,

$$I_R^{\text{prot}}(Q_{E,R}x) \in s^{m_R(\gamma)} \mathbb{C}[q^{\pm 1}, r^{\pm 1}], \quad m_R(\gamma) = m,$$

in the OP/Borcherds variables of Proposition ?? . The Gram trace m_R is additive under $\overline{\Pi}_X$, so $s^{m_R(\gamma_1 + \dots + \gamma_k)} = \prod_i s^{m_R(\gamma_i)}$ inside the finite quotient: the s -direction is coupled to the actual local/wrapped operation, not appended as an external Fock determinant. The integration defects

$$\mathfrak{o}_R^{I,\text{ch}}, \{\mathfrak{o}_{w,R}^{I,\text{prod}}\}_w, \mathfrak{o}_R^{I,\text{co}}, \mathfrak{o}_R^{I,\text{Prim}}, \mathfrak{o}_{R'R}^{I,\text{tr}}$$

record chain compatibility, multiplicativity on each word, comultiplicativity, primitive compatibility, and HN transition.

The aggregate finite obstruction tuple of the hybrid lane is

$$\mathfrak{D}_{\text{hyb},R} = (\mathfrak{o}_R^{\text{stage}}, \mathfrak{o}_R^\lambda, \mathfrak{o}_R^{\text{corr,LL}}, \mathfrak{o}_R^{\text{corr,mix}}, \mathfrak{o}_R^{\text{corr,WW}}, \mathfrak{o}_R^{\text{conf}}, \{\mathfrak{o}_{w,R}^{\text{assoc}}\}_w, \mathfrak{o}_R^{\text{col}}, \mathfrak{o}_R^{Q,\bullet}, \mathfrak{o}_R^{I,\bullet}, \mathfrak{o}_{\text{hyb},R'}^{\text{tr}}).$$

PROPOSITION 0.5 (*Finite consequences of the hybrid datum; conditional on $\mathfrak{D}_{\text{hyb},R} = \mathfrak{o}$*). Assume the data of Definition 0.4 are supplied for all R with compatible transition maps, and $\mathfrak{D}_{\text{hyb},R} = \mathfrak{o}$. Then:

- (i) the $b = \mathfrak{o}$ restriction is an ordinary support-local factorization algebra on $\text{Ran}(E)$ in the sense of Beilinson–Drinfeld [?BD, §3.4], equivalently a Costello–Gwilliam factorization algebra over the smooth real curve E_{top} [?CostelloGwilliam2, Vol. II §10];
- (ii) every operation with a wrapped input lands in a wrapped stratum;
- (iii) the ordered mixed maps, including the two-sided $\mu_{\alpha-;\gamma;\beta+,R}^{\text{hyb}}$, are the only Hall actions of projection-finite insertions on wrapped sectors in $\text{Corr}_R^{E,\text{hyb}}$ before reduced E -quotient;
- (iv) the eight-word atlas plus pentagon coherence yield an associative finite-stage reduced Hall product after vanishing cycles, orientations, HN quotienting, the quotient functor, and the $\theta_{\mu,R}^Q$;
- (v) for homogeneous x of charge γ , $I_R^{\text{prot}}(Q_{E,R}x) \in s^{m_R(\gamma)} \mathbb{C}[q^{\pm 1}, r^{\pm 1}]$, hence the positive s -direction cannot be supplied by the projection-finite stratum alone.

Proof. Items (i) and (ii) are built into the elliptic-degree stratification and Lemma 0.1. Item (iii) is the definition of the mixed operations: before reduced E -quotient they are the only displayed operations remembering relative E -position between finite local supports and a wrapped leg. Items (iv) and (v) are the eight-word coherence and the protected integration clause. $\square$

COROLLARY 0.6 (*No quotient-first universal hybrid repair; proved*). A candidate finite correspondence category for the wrapped sector gives the source object $\mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}}$ only if it contains the unreduced local, wrapped, and ordered mixed source stacks of Definition 0.4, their anchors, the eight-word flag atlas, the quotient functor $Q_{E,R}$ formed after those correspondences, the comparison maps $\theta_{\mu,R}^Q$, and protected integration with $\mathfrak{D}_{\text{hyb},R} = \mathfrak{o}$. A quotient of object stacks formed *before* the mixed and wrapped correspondences, or a determinant/Fock category carrying only the trace degree s^{m_R} , does not define the hybrid source.

Proof. By Lemma 0.1, a positive elliptic-degree sector is not a finite-support Ran sector; only the rigidified wrapped prequotient with anchor, together with the unreduced mixed and wrapped correspondences, retains the relative E -position used by the source data. A quotient-first construction loses these anchors. $\square$

CROSS-REPO ANCHOR

The hybrid carrier is the $K_3 \times E$ -specialization of the two-stage construction $\text{CY}_d\text{-Cat} \rightarrow E_d\text{-HolFA}(X) \rightarrow \text{ChirAlg}(C)$ supplied by the Vol III Calabi–Yau quantum-groups constellation: the projection-finite local stratum is the E_3 -holomorphic factorization algebra restricted to the smooth elliptic factor, and the wrapped stratum is the chain-level K_3 -fusion datum that the local stratum cannot see. Beilinson and Drinfeld’s Ran construction [?BD, §3.4–§3.5] provides the $b = 0$ part; the $b > 0$ part is the obligation that this manuscript isolates and the parallel chiral-bar-cobar volume tracks under the name Hall–Borcherds residual. The eight-word two-step binary flag atlas $w \in \{L, W\}^3$ is the smallest closed coherence calculus that detects the local/wrapped split through binary associativity; it is independent of the higher-coloured residual $\mathfrak{o}_R^{\text{col}}$ which closes higher symmetric arities, units, and overlaps.

STATUS UNDER (Do)

Probes A005, A011, A026, A037, A042 record finite-stage components of $\mathfrak{o}_R^{\text{stage}}$ and $\mathfrak{o}_R^\lambda$. Discharge of the wrapped-leg construction and the eight-word atlas at finite height is open; the $b = 0$ stratum is the classical projection-finite Ran sector of [?BD]. The hybrid carrier is therefore the projection-finite Ran stratum unconditionally, and the wrapped stratum conditionally on the named finite obstruction tuple $\mathfrak{D}_{\text{hyb}, R}$.

0.2 Cosection-twisted reduced d -critical orientation

The d -critical structure of the compact Hall sector on $K_3 \times E$ is the second entry of the pro-object. It supplies a vanishing-cycle coefficient system, a Joyce–Upmeyer–type orientation line, and a quotient-descent package for the reduced E -quotient. These are the finite-stage operands of the Hall product. Before any Pfaffian wall or Weyl transport can be discussed, the orientation line must descend through the cosection-reduced moduli.

The compact Hall heart is built from a Bridgeland-type stability sector (σ, S) on $D^b(\text{Coh}(K_3 \times E))$ in the K_3 lineage of [?BridgelandK3Stability, ?BayerMacriProjectivity]. The PTVV (-1) -shifted symplectic structures [?PTVV2013] restrict to the finite-type stacks selected by the HN quotient and give Joyce d -critical structures [?BBDJS2015] after classical truncation. The K_3 fibre direction supplies a semiregularity cosection CS_C , surjective on reduced strata, whose filtered Hall additivity on extension stacks

$$q^* \text{CS}_C = p_1^* \text{CS}_A + p_2^* \text{CS}_B$$

is the input to Kiem–Li cosection localization [?JoyceSong, ?JoyceUpmeyer].

THE REDUCED DETERMINANT COMPLEX

For an object C in the chosen finite stage, write

$$K_C^{\text{red}} = \text{RHom}_{\text{red}}(C, C),$$

the cosection-reduced self-Ext complex. The reduced orientation line is supplied with a chosen square root,

$$o_C \in \text{Pic}(\mathfrak{M}_X^{\text{or}}), \quad o_C^{\otimes 2} \simeq \det K_C^{\text{red}},$$

in the sense of [?JoyceUpmeyer], and is multiplicative under the reduced extension correspondences: $o_{C \oplus C'} \simeq o_C \otimes o_{C'}$ compatibly with the Hall additivity above. This is the data on which Thom–Sebastiani transport is supplied for the vanishing-cycle coefficient sheaves.

(O1) STRONG REDUCED ORIENTATION THROUGH THE QUOTIENT-ORIENTATION TOWER

The compact Hall product on $X = K_3 \times E$ is built only after the reduced E -quotient. The orientation line must descend through this quotient, and descent is detected by a tower of mod-2 Borel-equivariant gerbes. For a charge stratum C :

- (i) the reduced gerbe class

$$\alpha_C^{\text{red}} = w_2(\det K_C^{\text{red}}) + \text{cs}^* w_2(\det \text{coker cs}) \in H^2(\mathfrak{M}_C^{\text{red}}; \mathbb{F}_2);$$

- (ii) the free E -class

$$\alpha_C^{E, \text{free}} = a_1(C)u_1 + a_2(C)u_2 \in H^2(BE; \mathbb{F}_2);$$

- (iii) for each finite-stabilizer stratum S with stabilizer $H \subset E_{\text{tors}}$, the equivariant class $\tilde{\beta}_{C,S}^H \in H_H^2(S; \mathbb{F}_2)$, whose point-stabilizer restriction $\beta_{C,S}^H \in H^2(BH; \mathbb{F}_2)$ is read only after the mixed Borel terms, the stabilizer action on $H^\bullet(S; \mathbb{F}_2)$, and the spectral-sequence differentials have vanished.

Definition 0.7 ((O1) Strong multiplicative reduced orientation). The (O1) datum at finite HN height R consists of: the reduced orientation line o_C with chosen square root; its multiplicativity under reduced extension correspondences with chosen Thom–Sebastiani transport; null-trivializations of α_C^{red} , $\alpha_C^{E, \text{free}}$, $\tilde{\beta}_{C,S}^H$ and $\beta_{C,S}^H$ on every charge stratum used by the Hall correspondences; and a choice of zero $H^1(BH; \mathbb{F}_2)$ -linearization character $\lambda_{C,S}^H = 0$ for each finite stabilizer.

The full-level specialization at $H = E[2]$ is the Klein-four condition

$$\beta_{C,S}^{E,2} = b_{20}x_1^2 + b_{11}x_1x_2 + b_{02}x_2^2, \quad (b_{20}, b_{11}, b_{02}) = (r_1, r_1 + r_2 + r_3, r_2),$$

$$\text{res}_{\langle e_i \rangle} \beta_{C,S}^{E,2} = r_i \tau_i^2,$$

with vanishing $r_1 = r_2 = r_3 = 0$ on every retained two-torsion stratum. For full-level $H = E[N]$ with $2^a \parallel N$, $a \geq 2$, the two-primary residual takes value in $H^2(B(\mathbb{Z}/2^a)^2; \mathbb{F}_2) = \mathbb{F}_2\langle y_1, x_1x_2, y_2 \rangle$ and contains the mixed coefficient $A_{12}^{(N)}$ not detected by cyclic restrictions.

(O1) is therefore not a single Picard-line assertion: it is the ordinary square root, the null-trivialization of the reduced and equivariant degree-two gerbes, and the choice of zero degree-one linearization characters for the actual finite stabilizers. At finite HN height this is the cocycle system of [?JoyceUpmeyer] restricted to $K_3 \times E$.

(O1)⁺ WEYL-EQUIVARIANT TRANSPORT ALONG $W^{(2)}(\Lambda_{II}^{2,1})$

The orientation gerbe descends through the reduced E -quotient, but the Igusa structure also requires it to be compatible with the Weyl group of the BKM target. For each of the three type-II simple reflections s_{∂_i} ($i = 1, 2, 3$), the $K_3 \times E$ side carries a wall transport

$$\tau_i : o_C \longrightarrow s_{\partial_i}^* o_C.$$

The word *coherent* is load-bearing. At finite height, the first object is the finite partial action groupoid $\mathcal{G}_R$ whose objects are retained oriented quotient strata and whose arrows are the retained type-II wall

transports; a choice of determinant-line lifts gives a 2-cocycle $c_{o,R} \in Z^2(\mathcal{G}_R; \mathbb{F}_2)$. Only on a complete $W^{(2)}(\Lambda_{II}^{2,1})$ -orbit with constant coefficients does this reduce to the group cohomology class

$$[c_o] \in H^2(W^{(2)}(\Lambda_{II}^{2,1}); \mathbb{F}_2).$$

Definition 0.8 ((O_I)⁺ Weyl-equivariant determinant-line transport). The (O_I)⁺ datum is the supplied set of lifts $\{\tau_i\}$ satisfying $[c_o] = 0$ with chosen 1-cochain killing c_o , the normalization $\tau_i^2 = 1$, and the transport of the quotient-orientation package

$$\mathfrak{q}_C = (\alpha_C^{\text{red}}, \alpha_C^{E, \text{free}}, \{(\beta_C^H, \lambda_C^H)\}_{H \in \mathcal{H}(C)})$$

along τ_i between charge strata over Weyl-related Gram degrees. The torsor defect of this transport is

$$\omega_{i,C} \in H^1(\mathfrak{M}_{s_{\partial_i} C}^{\text{red}}; \mathbb{F}_2) \oplus \bigoplus_{H \in \mathcal{H}(C)} H^1(BH; \mathbb{F}_2),$$

required to vanish. The Coxeter presentation of $W^{(2)}(\Lambda_{II}^{2,1})$ then promotes these lifts to an honest action on the oriented Pfaffian line over the wall-complement of the reduced compact Hall sector.

(O_I)⁺ is a determinant-line monodromy datum. No scalar formula enters; the transports must move the entire null-trivialization package, not only the underlying line. The (O_I)⁺ ledger comprises $[c_o] = 0$, the chosen cochain, the unitarity $\tau_i^2 = 1$, and the torsor vanishing $\omega_{i,C} = 0$ on every retained orbit.

COSECTION LOCALIZATION AND THE COSECTION-REDUCED HEART

The K_3 semiregularity cosection CS_C selects a reduced obstruction theory; the corresponding Behrend constructible function ν^{red} produces the reduced vanishing-cycle complex Φ^{red} used in the Hall product [?BEHREND, ?JoyceSong]. The filtered Hall additivity $q^* \text{CS}_C = p_1^* \text{CS}_A + p_2^* \text{CS}_B$ on extension stacks is the cosection-localization compatibility required by the Hall correspondence; the Joyce–Upmeyer orientation lines are required to be Thom–Sebastiani-coherent on the resulting two-step and four-input flag stacks.

PROPOSITION 0.9 (*Finite d-critical Hall product; conditional on the (O_I) and atlas data*). Fix R and closed-configuration substacks $\mathfrak{M}_{c,R,I}^{\text{loc,cl}}$ in a noetherian abelian Hall heart, and assume:

- (i) finite-type quasi-smooth derived Artin enhancements of the selected object, extension, and two-step flag stacks, with perfect obstruction theories;
- (ii) finite residual inertia after scalar, E -translation, and wrapped rigidifications;
- (iii) finite-type extension and flag stacks with admissible p^* and q ;
- (iv) PTVV (-1) -shifted symplectic structures restricting to these substacks, classical truncations carrying Joyce d-critical structures [?PTVV2013, ?BBDJS2015];
- (v) surjective K_3 semiregularity cosections with filtered Hall additivity;
- (vi) Joyce–Upmeyer orientation lines and orientation transports [?JoyceUpmeyer] with all finite-stabilizer characters trivialized;
- (vii) Thom–Sebastiani transport for the reduced vanishing-cycle sheaves satisfying two-step flag pentagons and higher coloured configuration coherences.

Then the finite reduced local-local product

$$m_R^{\text{red}} = q! \circ \text{TS}_R^{\text{red}} \circ p^*$$

is an honest morphism on $H_c^\bullet(\mathfrak{M}_{c,R,I}^{\text{loc,cl}}, \Phi_{c,R,I}^{\text{red}} \otimes o_{c,R,I}^{\text{red}})$ and is associative on the finite stage.

The first obstruction to removing these hypotheses is finite inertia: proper finite-type correspondences do not by themselves eliminate residual $\mathbb{G}_m$, direct-sum, or parabolic automorphisms, and a residual $\mathbb{G}_m$ already contributes $H^\bullet(B\mathbb{G}_m) = \mathbb{C}[u]$ to protected cohomology. The phrase *finite stage* here means finite charge support, finite-type stacks, and finite-dimensional protected cohomology after the stated quotients and admissibility hypotheses; it does not mean that the Hall correspondence morphisms are themselves finite morphisms.

QUOTIENT-AFTER-CORRESPONDENCE ORDERING

The E -quotient is taken *after* the extension correspondences, not before. A pseudofunctor $\text{Corr}_R^{E,\text{hyb}} \rightarrow \text{Corr}_R^{\text{red,hyb}}$ supplies the descent; applying the vanishing-cycle Borel–Moore functor with orientation lines included gives $Q_{E,R}$ between the corresponding finite correspondence module categories. For each operation $\mu_R^\bullet = q! \text{ TS } p^*$ a chosen comparison isomorphism

$$\theta_{\mu,R}^Q : Q_{E,R} q! \text{ TS } p^* \xrightarrow{\sim} \overline{q}! \overline{\text{TS}} \overline{p}^* (Q_{E,R}^{\boxtimes k})$$

is part of the datum. Its compatibility with the eight-word flag 2-morphisms and HN transitions is the (O1) descent calculus. A quotient-first construction (object-stack quotient followed by a new Hall product) is not the same datum.

STATUS OF THE OBSTRUCTION TUPLE

The orientation lane carries the residual

$$\mathfrak{D}_{\text{or},R} = (\mathfrak{o}_C^{\text{red}}, \mathfrak{o}_C^{E,\text{free}}, \{\mathfrak{o}_{C,S}^H\}_{H,S}, \{\mathfrak{o}_{C,S}^{\lambda^H}\}_{H,S}, [c_o]_R, \xi_{c_o,R}, \{\mathfrak{o}_{i,R}^{\tau,2}\}_i, \{\omega_{i,C,R}\}_i)$$

required to vanish at every finite HN height, with chosen null-trivializations. Probes Ao74, Ao76, Ao82 audit the (O1) tower; probes Ao47, Ao49, Ao89 audit (O1)⁺. Discharge of (O1) under $K_3 \times E$ -specific finite-stabilizer hypotheses is open in generality; the K_3 -fibrewise framework of [?JoyceSong, ?JoyceUpmeyer] reduces (O1) on the projection-finite stratum to the finite-stabilizer linearization choice and the mixed Borel computation, and the wrapped stratum is the open piece tracked by the Hall–Borcherds residual in the parallel Vol II constellation.

CROSS-REPO ANCHOR

Vol III’s Calabi–Yau quantum-groups stratification observes that $K_3 \times E$ admits no global non-commutative crepant resolution, so the orientation lane cannot be supplied by a global NCCR-induced heart; the cosection-twisted reduced d-critical structure is forced on the chosen Bridgeland heart instead. The displayed Borel calculation at $E[2]$ is the smallest finite stabilizer detected by the cosection-reduced determinant; even $N \geq 4$ stabilizers carry the mixed coefficient $A_{12}^{(N)}$, which the displayed cyclic restrictions miss, and which (O1) requires to be null-trivialized as part of the full degree-two gerbe.

0.3 Hall correspondences: product, coproduct, primitives

The third entry of the Dirac–Igusa pro-object is the Hall structure on the cosection-reduced d-critical heart. It is the algebraic content that the orientation lane equips and the hybrid carrier indexes. Three operations live here: the Hall product m_R^{red} , the collision coproduct Δ_R^{ch} , and the primitive projection P_R^{prim} . After normal-ordered Gram descent these form a graded Hopf datum on the protected primitive layer; recognition of this Hopf datum with the Borcherds–Kac–Moody target is the §5.6 obligation.

Russian-school discipline applies. The Kontsevich–Soibelman cohomological Hall algebra of a 3-Calabi–Yau category [?KSCoHA] supplies the prototype; on $K_3 \times E$ the cosection reduction [?JoyceSong,

[JoyceUpmeier] replaces the unreduced critical CoHA, and the wrapped sector is the residual that the hybrid carrier of §0.1 retains. The Maulik–Okounkov stable-envelope construction and the Schiffmann–Vasserot theorems on Drinfeld doubles of CoHA operate on the same input data; their integral form is the Hall side of the present construction.

THE COMPACT HALL PRODUCT

Definition 0.10 (Compact Hall product on the cosection-reduced heart). Let $C_{\sigma,S,\leq R}$ be the finite HN quotient of §0.1. The compact Hall product at height R is the operation

$$m_R^{\text{red}} = q_{c,c'}! \circ \text{TS}_{c,c'}^{\text{red}} \circ p_{c,c'}^* : H_*^{\text{BM}}(\mathfrak{M}_c^\sigma \times \mathfrak{M}_{c'}^\sigma, \Phi_c^{\text{red}} \boxtimes \Phi_{c'}^{\text{red}} \otimes o_c \boxtimes o_{c'}) \rightarrow H_*^{\text{BM}}(\mathfrak{M}_{c+c'}^\sigma, \Phi_{c+c'}^{\text{red}} \otimes o_{c+c'})$$

attached to the extension correspondence

$$\mathfrak{M}_c^\sigma \times \mathfrak{M}_{c'}^\sigma \xleftarrow{p_{c,c'}} \mathfrak{E}_{c,c'}^\sigma \xrightarrow{q_{c,c'}} \mathfrak{M}_{c+c'}^\sigma,$$

in the cosection-reduced d-critical heart, with vanishing-cycle coefficients Φ^{red} , Joyce–Upmeier orientation lines o , and Thom–Sebastiani transport $\text{TS}_{c,c'}^{\text{red}}$. Associativity on the finite stage is the eight-word two-step binary flag pentagon of §0.1, axiom (5).

For local-only inputs and ordered mixed and wrapped inputs, the same construction produces the three operation families μ_R^{loc} , μ_R^{hyb} , μ_R^{wr} of §0.1. Existence of the product as a genuine morphism on Borel–Moore chains requires the admissibility/properness residuals of that section to vanish.

THE COLLISION COPRODUCT ON THE SOURCE COALGEBRA

The collision coproduct is defined on the same correspondence geometry, read in the opposite direction: instead of contracting an extension to a single output, one expands an object into the pairs of its sub/quotient extensions. At finite stage, this is the 2-step flag stack already supplied as part of the hybrid datum.

Definition 0.11 (Collision coproduct). The finite collision coproduct

$$\Delta_R^{\text{ch}} : C_{X,R} \longrightarrow C_{X,R} \otimes C_{X,R}$$

is the chain-level pull-push along the 2-step flag stack $\mathfrak{Flag}_{c,c',R}^{(2)}$ of subobjects of charge c and quotients of charge c' in $\mathfrak{M}_{c+c',R}^\sigma$, computed in the cosection-reduced d-critical heart, before the reduced E -quotient. After $Q_{E,R}$ it satisfies the coalgebra identities

$$(\Delta_R^{\text{ch}} \otimes \text{id}) \Delta_R^{\text{ch}} = (\text{id} \otimes \Delta_R^{\text{ch}}) \Delta_R^{\text{ch}}, \quad (\varepsilon_R^C \otimes \text{id}) \Delta_R^{\text{ch}} = (\text{id} \otimes \varepsilon_R^C) \Delta_R^{\text{ch}} = \text{id},$$

which are the dual of the Hall pentagon on the same flag stack. For $R' \geq R$ the transition map $\rho_{R',R}^C$ commutes with Δ^{ch} .

This is the source coalgebra of §0.5; its identification with the bar construction $\bar{B}_E^{\text{ch}}(\mathcal{F}_X^{\text{hyb}})$ is the bar comparison there.

PRIMITIVE PROJECTION AND THE HOPF DATUM

The primitive projection on a graded conilpotent Hopf object is unique once the augmentation, product, and coproduct are supplied.

Definition 0.12 (Primitive Hall sub-bracket). Write

$$\tilde{P}_X = H^\circ \text{Prim}_{\text{prot}}(\mathcal{F}_X) = \ker(\tilde{\Delta} : \mathcal{F}_X \rightarrow \mathcal{F}_X \otimes \mathcal{F}_X)$$

for the primitive layer of the protected Hall Hopf object, where $\tilde{\Delta} = \Delta - (\eta\varepsilon \otimes \text{id} + \text{id} \otimes \eta\varepsilon)$. The Hall product restricts to a $\tilde{\Gamma}_X$ -graded bracket

$$[\cdot, \cdot]_X : \tilde{P}_{X, \hat{c}} \otimes \tilde{P}_{X, \hat{c}} \longrightarrow \tilde{P}_{X, \hat{c} \star \hat{c}},$$

the protected primitive Hall bracket. After normal-ordered Gram descent (§0.1) and the Hopf-radical quotient, the descended bracket is Γ_{gram} -graded and the descended primitive layer is the candidate $P_X^{\Pi, \text{red}}$ to be identified with $\mathfrak{g}_{\Delta, \cdot}$.

LEMMA 0.13 (Primitive bracket lifting; conditional on the Frobenius defect $\mathfrak{o}_F = \mathfrak{o}$). Let $\langle \cdot, \cdot \rangle_X$ be the protected Hopf pairing on $\tilde{P}_X$, non-degenerate on the radical quotient. The trilinear Frobenius defect

$$\mathfrak{o}_F(x, y, z) := \langle [x, y], z \rangle + (-1)^{|x| \cdot |y|} \langle y, [x, z] \rangle, \quad x, y, z \in \tilde{P}_X, \quad c_x + c_y + c_z = \mathfrak{o},$$

viewed as a class in $C_{\text{cyc}}^3(\tilde{P}_X; k)$ after the reduced E -quotient and radical quotient, vanishes precisely when the bracket is graded-symmetric for the Hopf pairing. Vanishing is the cyclic-CY-3 Serre-functor identity of Lemma ???. Non-degeneracy of the Hopf pairing alone does *not* force $\mathfrak{o}_F = \mathfrak{o}$; it only detects the vanishing once supplied.

LEMMA 0.14 (Hopf radical is a Lie ideal and a coideal; conditional on $\mathfrak{o}_F = \mathfrak{o}$ and Hopf adjointness). Let P_R^Π be a finite normal-ordered primitive stage with transported product m_R^Θ , co-product Δ_R^Θ , graded symmetric Hopf pairing $\langle \cdot, \cdot \rangle_R$, and radical Rad_R^Π . Assume Hopf adjointness $\langle m_R^\Theta(x, y), z \rangle_R = \langle x \otimes y, \Delta_R^\Theta z \rangle_R$ and non-degeneracy on $P_R^\Pi / \text{Rad}_R^\Pi \otimes P_R^\Pi / \text{Rad}_R^\Pi$. Then

$$(\bar{q}_{\Pi, R} \otimes \bar{q}_{\Pi, R}) \Delta_R^\Theta(r) = \mathfrak{o} \quad (r \in \text{Rad}_R^\Pi),$$

i.e. Rad_R^Π is a coideal. If, in addition, $\mathfrak{o}_{F, R} = \mathfrak{o}$, then Rad_R^Π is a Lie ideal.

Proof. For $r \in \text{Rad}_R^\Pi$ and $x, y \in P_R^\Pi$, Hopf adjointness gives $\langle x \otimes y, \Delta_R^\Theta(r) \rangle_R = \langle m_R^\Theta(x, y), r \rangle_R = \mathfrak{o}$; non-degeneracy of the quotient tensor pairing then kills the class. The Lie-ideal half is direct: if $\mathfrak{o}_{F, R} = \mathfrak{o}$ then $\langle [x, r]_\Theta, z \rangle_R = -(-1)^{|x| \cdot |r|} \langle r, [x, z]_\Theta \rangle_R = \mathfrak{o}$ for every z , so $[x, r]_\Theta \in \text{Rad}_R^\Pi$. $\square$

INTEGRAL FORM AND STABLE-ENVELOPE TRANSPORT

Two parallel constructions act on the same compact Hall heart:

- (i) the integral form of the Schiffmann–Vasserot elliptic Hall algebra, whose $K_3 \times E$ -specialization carries the Maulik–Okounkov stable envelope;
- (ii) the Drinfeld double $\mathcal{D}_h(\text{CoHA}_{K_3 \times E})$ of Vol III, whose positive half is the protected primitive Hopf algebra above and whose Drinfeld pairing is the Hopf pairing.

The compatibility datum between these two constructions is part of the cross-repo consistency check between this manuscript and the Calabi–Yau quantum-groups constellation. Disagreement on a load-bearing constant or sign is the deliverable; the Drinfeld double is not asserted here as a black box.

CONILPOTENT SOURCE COALGEBRA AND HOPF RADICAL QUOTIENT

Definition 0.15 (Conilpotent source coalgebra). At finite stage, $C_{X,R}$ is the conilpotent chiral coalgebra on E obtained from the collision coproduct on $\mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}}$ after the reduced E -quotient: it is filtered by a finite increasing source charge filtration

$$\mathfrak{o} = F_{R,-1}^{\text{co}} \subset F_{R,0}^{\text{co}} \subset \cdots \subset F_{R,N_R}^{\text{co}} = C_{X,R}$$

whose associated graded pieces are finite dimensional in each Γ_R^{Π} -degree, and whose reduced iterated coproduct is nilpotent on each finite stage. Conilpotence is recorded by the defect $(\overline{\Delta}_R^{\text{ch}})^{N_R+1}$, required to vanish. Inverse-limit completion gives the source coalgebra $C_X = \varprojlim_R C_{X,R}$.

PROPOSITION 0.16 (Sectorial Hall truncation and inverse limit; conditional on $[K_3 \times E\text{-fin}]$, $[K_3 \times E\text{-atlas}]$, and $[ML]$). Under the three standing hypotheses $[K_3 \times E\text{-fin}]$ (finite-type semistable substacks at bounded HN height), $[K_3 \times E\text{-atlas}]$ (finite d-critical/cosection Hall atlas with Joyce–Upmeyer orientation lines), and $[ML]$ (Mittag–Leffler descent for the HN inverse system), the finite Hall product, coproduct, primitive projection, and Hopf pairing are compatible with the transition maps $\rho_{R'R}$, and their inverse limits define the completed Hall Hopf datum P_X, C_X . The Mittag–Leffler hypothesis controls the $R^1 \varprojlim$ obstruction on every finite-window slice; without it, only a compatible pro-object is obtained.

The hypothesis $[K_3 \times E\text{-fin}]$ is not a consequence of fibrewise K_3 boundedness alone. A precise sufficient input is product-stability boundedness in the Abramovich–Polishchuk/Liu sense [*?AbramovichPolishchukTStructures, ?LiuProductStability*]: for a rational K_3 stability condition $\sigma_S = (\mathcal{A}_S, Z_S)$ and Liu’s product stability $\sigma_{S,t}$ on $D^b(\text{Coh}(S \times E))$, fixed-class boundedness

$$\mathfrak{M}_{\sigma_{S,t}}^{ss}(\gamma) \text{ is finite type over } \mathbb{C}$$

together with Liu’s support property would imply $[K_3 \times E\text{-fin}]$ for bounded h_S -windows. Piyyaratne–Toda boundedness for threefold Bridgeland/BMT [*?PiyyaratneTodaModuli3folds*] does not automatically supply this theorem; fibrewise Bayer–Macri boundedness alone does not either.

WALL-CROSSING AND CHAMBER DEPENDENCE

The lattice cocycle $B(c, c')$ of the dyonic Mukai construction is chamber-independent. The compact Hall side $D_o(X; \sigma, S)$ is chamber-dependent: finite-type semistable substacks change under wall-crossing, and the Kontsevich–Soibelman wall-crossing formula [*?KSCoHA*] relates Hall counts in adjacent chambers. If the datum $D_o(X; \sigma, S)$ is supplied in chamber σ , then $\overline{\Pi}_X$ remains a homomorphism and the recognition target is $\mathfrak{g}_{\Delta}$, provided the chamber identifications hold for the OP/DT scalar import; that the chosen σ lies in the OP/DT chamber, or that the wall-crossing transport is compatible with $\overline{\Pi}_X$, is a separate conjectural input. No chamber-independence of the recognition target is asserted.

STATUS UNDER THE OBSTRUCTION TUPLE

The Hall lane carries the residuals

$$\mathfrak{D}_{\text{Hall},R} = (\mathfrak{o}_R^{\text{stage}}, \mathfrak{o}_R^{\text{conf}}, \mathfrak{o}_R^{\text{prop}}, \mathfrak{o}_R^{\text{TS}}, \mathfrak{o}_R^{\text{assoc}}, \mathfrak{o}_{F,R}, \mathfrak{o}_R^{\text{coid}}, \mathfrak{o}_{R'R}^{\text{ML}}).$$

Discharge of the wrapped product on the projection-finite stratum is classical d-critical CoHA after cosection reduction; on the wrapped stratum the construction is open and tracked by the (Do) obstruction. Hopf adjointness, Frobenius vanishing, and PBW compatibility are independent constructions on the descended primitive layer.

CROSS-REPO ANCHOR

Vol III's Drinfeld doubling $Y^+(X) \rightarrow G(X)$ produces the BKM target of the present manuscript on the source side; the integral form is the chiral lift of the Schiffmann–Vasserot elliptic Hall algebra. Maulik–Okounkov stable-envelope transport supplies the integral form on the projection-finite Ran stratum; the wrapped stratum is the stratum-completion that this manuscript and the chiral-bar-cobar volumes reconstruct. No black-box equality $\text{CoHA} = W_\infty$ is asserted: $\text{CoHA}(\mathbb{C}^3) = Y^+(\mathfrak{gl}_1)$ in Vol III, and $W_{1+\infty}$ appears only after Drinfeld doubling, centre, vacuum evaluation. The same discipline applies here: the protected primitive Hopf algebra of $K_3 \times E$ becomes the BKM target only after the Hopf radical quotient, the Frobenius identity, and the no-extra-relations theorem of §0.6.

0.4 Dirac operator and protected Pfaffian

The fourth entry of the Dirac–Igusa pro-object is the protected Pfaffian line and its first-order Dirac block decomposition. Two constructions are at work. The geometric construction starts from the cosection-reduced d-critical heart, builds a Joyce–Upmeyer orientation, descends to the reduced quotient, and associates a Pfaffian line to the determinant of the reduced obstruction theory in the Brav–Bussi–Dupont–Joyce–Szendrői framework [?BBDJS2015]. The algebraic construction starts from the imported Borchers–Kac–Moody denominator algebra, fixes a positive window in Γ_{gram} , and writes the algebraic super-Pfaffian

$$\text{sPf}_R^{\text{alg}}(D) = \prod_{\gamma \in A_R} (1 - x^\gamma)^{\text{sdim } V_\gamma}$$

of the trivial first-order block decomposition D_γ on the target side. The Pfaffian–Dirac theorem of Chapter 6 is the identification, conditional on (Do) , (O1) , $(\text{O1})^+$, (O2) , of these two Pfaffian sections after the chosen line-comparison ι_{aut} .

PAFFIAN LINE AND SQUARING ISOMORPHISM

Definition 0.17 (Pfaffian line on the cosection-reduced quotient). Let K_C^{red} denote the cosection-reduced self-Ext complex on a charge stratum, with reduced determinant $\det K_C^{\text{red}}$ and Joyce–Upmeyer orientation o_C . The Pfaffian line at finite stage is

$$\mathcal{L}_{\text{Pf},R} = \bigotimes_C \text{Pf}(K_{C,R}^{\text{red}})$$

on the reduced moduli, with squaring isomorphism

$$\mathcal{L}_{\text{Pf},R}^{\otimes 2} \simeq \mathcal{L}_{\text{det},R}$$

in the Brav–Bussi–Dupont–Joyce–Szendrői sense [?BBDJS2015]. The successor maps $\rho_{R^+,R}^{\text{Pf}}$ commute with these squaring isomorphisms, identifying the inverse limit with a Pfaffian line $\mathcal{L}_{\text{Pf},X}$ on the inverse-limit object. A normalized section $\text{pf}_{X,R}$ at finite stage and $\text{pf}_X = \varprojlim_R \text{pf}_{X,R}$ are part of the datum.

PAFFIAN-TO-AUTOMORPHIC COMPARISON

The recognition statement is read on the automorphic line. A section-level identification $\text{pf}_X = \Delta_\delta$ requires a comparison of $\mathcal{L}_{\text{Pf},X}$ with the pullback of the automorphic weight-five line to the moduli base.

Definition 0.18 (Pfaffian-to-automorphic line comparison). A section-level Pfaffian-to-automorphic comparison is a chosen isomorphism of line bundles

$$\iota_{\text{aut}} : \mathcal{L}_{\text{Pf},X} \xrightarrow{\sim} \mathcal{L}^5 \otimes \nu_{\Delta_\delta}$$

to the pullback of the automorphic weight-five line, twisted by the multiplier ν_{Δ_5} , compatible with cusp trivialization at $\mathfrak{c}_{\infty}$ and with the squaring map $\mathcal{L}_{\text{Pf},X}^{\otimes 2} \simeq \mathcal{L}_{\text{det},X}$.

FIRST-ORDER DIRAC BLOCK DECOMPOSITION

Definition 0.19 (First-order Dirac block). For $\gamma \in \Gamma_R^{\Pi,+}$, write $P_{R,\gamma}^{\Pi,+}$ for the cusp-positive primitive piece in degree γ and put $x^\gamma = q^n r^l s^m$ for $\gamma = (n, l, m)$. The two-term Pfaffian/Koszul block on $P_{R,\gamma}^{\Pi,+} \oplus (P_{R,\gamma}^{\Pi,+})^\vee[1]$ is the skew operator

$$J_\gamma = \begin{pmatrix} 0 & 1 - x^\gamma \\ -(1 - x^\gamma) & 0 \end{pmatrix}, \quad \text{Pf}(J_\gamma) = (1 - x^\gamma)^{\text{sdim } P_{R,\gamma}^{\Pi,+}}.$$

For $\gamma \notin \Gamma_R^{\Pi,+}$, set $P_{R,\gamma}^{\Pi,+} = 0$. The first-order block of $\mathfrak{D}_X^{D_0}$ at height R is the direct sum

$$\mathfrak{D}_{X,R}^{D_0} = \bigoplus_{\gamma \in \Gamma_R^{\Pi,+} \setminus \{0\}} J_\gamma,$$

acting on $\bigoplus_\gamma P_{R,\gamma}^{\Pi,+} \oplus (P_{R,\gamma}^{\Pi,+})^\vee[1]$.

The principal symbol of $\mathfrak{D}_{X,R}^{D_0}$ with respect to the finite Chevalley–Eilenberg/Koszul filtration is the normal-ordered Hall bracket of §0.3: the Pfaffian line is the determinant detector, and the symbol carries the Lie data. The phrase *first-order* means the Pfaffian/primitive object before scalar squaring; no analytic Dirac operator on a constructed Hilbert space is asserted.

THE PFAFFIAN PRODUCT AT FINITE STAGE

PROPOSITION 0.20 (Pfaffian product at finite stage; conditional on (Do-Pf)). Suppose the orientation lane (OI, OI^+) has descended to the reduced quotient, the first-order block decomposition is supplied, and the support and parity residuals

$$\mathfrak{v}_{R,\gamma}^{\text{supp}} = \begin{cases} P_{R,\gamma}^{\Pi,+}, & a_\Delta(\gamma) = 0 \\ 0, & a_\Delta(\gamma) \neq 0 \end{cases}, \quad \mathfrak{v}_{R,\gamma}^{\text{par}} = p_{R,\gamma}^{\bar{0}} - p_{R,\gamma}^{\bar{1}} - a_\Delta(\gamma) \in \mathbb{Z},$$

together with the leading-coefficient residual $\mathfrak{v}_R^{\text{lead}} = \text{lc}_{\mathfrak{c}_{\infty}}(\text{pf}_{X,R}) - 64$, all vanish on the test window, where $a_\Delta(\gamma)$ is the Borcherds exponent and $p_{R,\gamma}^{\bar{p}} = \dim P_{R,\gamma,\bar{p}}^{\Pi,+}$. Then the Pfaffian section at finite stage has expansion

$$\text{pf}_{X,R} |_{\mathfrak{c}_{\infty}} = 64 q^{1/2} r^{1/2} s^{1/2} \prod_{\substack{\gamma \in \Gamma_R^{\Pi,+} \\ \gamma \neq 0}} (1 - x^\gamma)^{\text{sdim } P_{R,\gamma}^{\Pi,+}}.$$

The cusp-positive Pfaffian half is recorded; negative root representatives and the Cartan/imaginary completion enter the Lie recognition statement of §0.6.

The support residual is a super-vector-space residual, not a K_0 class: its vanishing excludes invisible even–odd cancelling primitive pieces in target-zero degrees. The parity residual is a signed superdimension residual. These finite-stage residuals on the test window do not by themselves prove compact source coverage; the inclusion $A_R \subset \Gamma_R^{\Pi,+}$ of active target degrees in the source image is itself a residual condition, not a consequence of the Borcherds product.

LOCAL PFAFFIAN WALL ON TYPE-II WALLS

The (O₂) obstruction concerns the local Pfaffian normal form on the three simple type-II walls $\mathbb{H}_{\delta_i}$, $i = 1, 2, 3$, of the Borcherds–Cartan datum $\mathcal{B}_{\Delta_5}$.

Definition 0.21 (Rank-one type-II Pfaffian crossing). For a simple type-II root δ , let T_δ be a real analytic tangent chart with trivial s_δ -action, set $U_\delta = T_\delta \times (-\varepsilon, \varepsilon)$ with $s_\delta(t, u) = (t, -u)$, and let $\mathcal{L}_\delta^{\text{tan}}$ be an oriented tangent Pfaffian line with s_δ -invariant unit v_δ . The rank-one odd Pfaffian crossing is the skew family

$$D_u = \begin{pmatrix} 0 & u \\ -u & 0 \end{pmatrix}, \quad K_\delta^{\text{cross}} = [\mathbb{R} \xrightarrow{u} \mathbb{R}],$$

in odd Pfaffian parity. The total local Pfaffian section is

$$\text{pf}_\delta = v_\delta u, \quad s_\delta^* \text{pf}_\delta = -\text{pf}_\delta, \quad s_\delta^*(\text{pf}_\delta^2) = \text{pf}_\delta^2.$$

PROPOSITION 0.22 (Rank-one crossing gives the type-II local sign; proved). For $i = 1, 2, 3$, the rank-one crossing of Definition 0.21 forms the finite generic type-II local-sign datum with $N_{\delta_i}^{\text{Pf}} = 1$, and at finite HN height R , with successor maps preserving the tangent charts, the wall coordinate $u_{\delta_i, R}$, and the invariant units, the (O₂) residual entries

$$\mathfrak{o}_{\delta_i, R}^{\text{chart}}, \mathfrak{o}_{\delta_i, R}^{\text{wall}}, \mathfrak{o}_{\delta_i, R}^{\text{split}}, \mathfrak{o}_{\delta_i, R}^{\text{unit}}, \mathfrak{o}_{\delta_i, R}^{\text{rank}}, \mathfrak{o}_{\delta_i, R'}^{\text{tr}}$$

vanish for these local models, and $\varepsilon_{\delta_i, R}^{\text{loc}} = -1$.

Proof. The fixed locus of $s_\delta(t, u) = (t, -u)$ is $u = 0$, so the wall and the tangent fixed locus agree. The determinant factor is split: the tangent line is $\mathcal{L}_\delta^{\text{tan}}$, and the normal factor is K_δ^{cross} . The unit v_δ is invariant by construction. The skew matrix has Pfaffian u , so the normal-mode rank is one. Reflection sends $u \mapsto -u$ and fixes v_δ ; hence the local Pfaffian section changes sign and its square is fixed. The transition hypothesis is exactly preservation of these data under successor maps. $\square$

HALF-HILBERT ORBIT AND THE TYPE-II SIGN SUM

The (O₂) obstruction also requires a wall *atlas*: at finite HN height R , each retained component and boundary stratum of $\mathbb{H}_{\delta_i}$ needs a chart of the rank-one form, and the normal Pfaffian units, tangent/normal splittings, quotient orientations, and protected integration maps must agree on chart overlaps. A single generic chart computes only a local sign.

The half-Hilbert orbit of the type-II reflection group on the relevant retained stratum carries a sign sum of size $4096 = 2^{12}$, counting the supplied finite-window normal Pfaffian signs across the twelve retained components and boundary strata of the type-II wall configuration; this sum is the signature of the (O₂) wall normal-form theorem. Local rank-one charts contribute the local factors; the global sum reads as the orientation character on the $W^{(2)}(\Lambda_{II}^{2,1})$ -orbit. The full normal-form theorem with overlap, boundary, and component coherence is the content of Appendix E and probes Ao47, Ao92, A100; the present section sketches the local model only.

PROPOSITION 0.23 (Generic local sign and the orientation character; conditional on (O₁)⁺ and the (O₂) wall atlas). Suppose (O₁)⁺ supplies a normalized lift $\tau_{i, R, S} : \mathfrak{o}_{R, S} \rightarrow s_{\delta_i}^* \mathfrak{o}_{R, S}$ of the simple type-II reflection on every retained orbit, and the (O₂)/hybrid residual for δ_i vanishes. In a local chart

with $K_{\delta_i, R}^{\text{nor}} \simeq [\mathbb{R} \xrightarrow{u_i} \mathbb{R}]$, with normal Pfaffian unit v_i and $s_{\delta_i}(v_i) = v_i$, let $\ell_{\delta_i, R, S}$ be the path in the reduced quotient wall-complement carrying $u_i = \epsilon$ to $u_i = -\epsilon$, closed by $\tau_{i, R, S}$. Then

$$\mathcal{M}_{\ell_{\delta_i, R, S}}(v_i u_i) = v_i(-u_i) = -v_i u_i, \quad \epsilon_{o, R}(s_{\delta_i}) = (-1)^{N_{\delta_i, R}^{\text{Pf}}}.$$

For the rank-one type-II model $\epsilon_{o, R}(s_{\delta_i}) = -1 = \nu_{\Delta_5}(s_{\delta_i})$. The global statement $\epsilon_o|_{W^{(2)}(\Lambda_{II}^{2,1})} = \nu_{\Delta_5}|_{W^{(2)}(\Lambda_{II}^{2,1})}$ is the chamber-permuting half of the Pfaffian–Dirac orientation identity, conditional on the (O₂) wall atlas.

PF AFFIAN NORMALIZATION AND TRACE

The supplied Pfaffian-to-automorphic comparison ι_{aut} then gives, on the imported Borchers product side,

$$\iota_{\text{aut}}(\text{Pf}_{\text{prot}}(\mathfrak{D}_X)) = \mathcal{D}_X = \Delta_5,$$

with cusp-side expansion

$$\text{pf}_X|_{\mathfrak{c}_{\infty}} = 64q^{1/2}r^{1/2}s^{1/2} \prod_{\gamma=(n,l,m) \in \Gamma_{\text{eff}}} (1 - q^n r^l s^m)^{f(nm,l)},$$

matching the imported Borchers–Gritsenko–Nikulin product. The scalar trace is the Oberdieck–Pixton normalization

$$\text{Tr}_{\text{prot}}((-1)^F \mathfrak{D}_X^{-2}) = -4096 \Delta_5^{-2},$$

which carries the negative sign and the $4096 = 2^{12}$ overall multiplicity attached to the half-Hilbert orbit. This sign and multiplicity are not chosen branch conventions; they are computed by the (O₂) wall atlas after (O₁)⁺ has supplied the Weyl transports.

CROSS-REPO ANCHOR AND STATUS

Vol II’s Construction Problem 1 isolates the protected Pfaffian $\text{Pf}_{\text{prot}}(\mathfrak{D}_X) = \Delta_5$ as the target identity for the chiral E_3 -algebra $\mathfrak{A}_{K_3 \times E}$; the present section supplies the $K_3 \times E$ -side construction. BBDJS [?BBDJS2015] provides the orientation-line and Pfaffian calculus on the cosection-reduced d-critical heart; Joyce–Upmeyer [?JoyceUpmeyer] provides multiplicativity under reduced extension correspondences; Joyce–Tanaka–Upmeyer [?JoyceTanakaUpmeyer] extends the framework to 4-fold analogues but is not used directly here. The constants $4096 = 2^{12}$ and -1 are fixed by group theory plus one local rank-one Pfaffian wall computation; the Oberdieck–Pixton minus sign is a scalar branch convention, not the source of the orientation character. No identification of the Pfaffian Δ_5 with a compact BPS Hilbert space is asserted: Δ_5 is the protected scalar shadow of $\mathfrak{A}_{K_3 \times E}$, and the operator-level package is the pro-object $\mathfrak{D}_X^{\text{DI}}$ conditional on (Do), (O₁), (O₁)⁺, (O₂).

0.5 Koszul source coalgebra and chiral MC twisting

The fifth entry of the Dirac–Igusa pro-object is the Koszul lane: the source coalgebra C_X , the bar comparison b_X , and the cobar quasi-isomorphism Θ_{Kos} to the imported current envelope $\text{Den}_{\Delta_5, E}$. Source–target separation is load-bearing. The target counit

$$\varepsilon_{\Delta, R} : \Omega_E^{\text{ch}} \bar{B}_E^{\text{ch}}(\text{Den}_{\Delta_5, E, \leq R}) \longrightarrow \text{Den}_{\Delta_5, E, \leq R}$$

is internal to the imported denominator current envelope and produces no source data. The compact Koszul comparison $\Omega_E^{\text{ch}} C_X \simeq \text{Den}_{\Delta_5, E}$ is a source-to-target theorem on the supplied hybrid source, conditional on the named obstructions.

The Beilinson cut applies in its purest form on this lane. $\bar{B}^{\text{ch}}$ is MC twisting / coupling coalgebra; it is not the bulk. The bulk is $\text{Den}_{\Delta,E}$ itself, the chiral shadow of the BKM denominator algebra; the source coalgebra C_X is the supplied compact Hall data after collision coproduct, conilpotent filtration, and bar-length truncation, and the cobar map is the comparison.

SOURCE-TARGET SEPARATION

LEMMA 0.24 (*Source-target separation for the Koszul lane; proved*). Fix a finite height R . The target counit $\varepsilon_{\Delta,R}$ above is a morphism internal to the imported current envelope. A source Koszul datum is different data: it must first construct

$$\mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}}, \quad C_{X,R}, \quad F_{R,\bullet}^{\text{co}}, \quad \Delta_R^{\text{ch}}$$

from the finite hybrid correspondence category $\text{Corr}_R^{E,\text{hyb}}$, its quotient functor $Q_{E,R}$, orientations, the eight-word two-step binary flag atlas, and the separate higher coloured configuration residual; only then may it compare

$$b_{X,R} : C_{X,R} \simeq \bar{B}_E^{\text{ch}}(\mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}}, \varepsilon_{X,R}^{\text{hyb}}), \quad \Theta_{\text{Kos},R} : \Omega_E^{\text{ch}} C_{X,R} \simeq \text{Den}_{\Delta,E,\leq R}.$$

Setting $C_{X,R} = \bar{B}_E^{\text{ch}}(\text{Den}_{\Delta,E,\leq R})$, or choosing a homotopy inverse to $\varepsilon_{\Delta,R}$, does not produce a source datum.

Proof. The domain of $\varepsilon_{\Delta,R}$ contains only target augmentation, target charge filtration, and target chiral operations; it does not contain rigidified wrapped prequotients, anchors, mixed local/wrapped correspondences, quotient-after-correspondence functor, orientation transport, protected integration, or primitive representatives. These are the data from which the source coalgebra and collision coproduct are required to be built. $\square$

CHIRAL KOSZUL SOURCE DATUM

Definition 0.25 (*Chiral Koszul source datum*). Fix (σ, S, R) and the Gram image Γ_R^{Π} . A finite-stage chiral Koszul source datum is

$$\mathfrak{R}_{X,R}^{\text{Kos}} = (\mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}}, C_{X,R}, F_{R,\bullet}^{\text{co}}, \Delta_R^{\text{ch}}, b_{X,R}, \Theta_{\text{Kos},R})$$

with the following clauses; every entry is source-side finite data except for the terminal comparison with the imported target.

- (i) *Source chiral object.* $\mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}}$ is the finite hybrid object of §0.1, with $\mathfrak{D}_{\text{hyb},R} = 0$.
- (ii) *Conilpotent chiral coalgebra.* $C_{X,R}$ is a coaugmented conilpotent chiral coalgebra on E , filtered by a finite increasing source charge filtration $F_{R,\bullet}^{\text{co}}$; reduced iterated coproduct nilpotent on each finite stage.
- (iii) *Collision coproduct.* Δ_R^{ch} is the collision coproduct from the finite Hall correspondences (local-local, ordered local-wrapped, wrapped-wrapped, and eight-word flag stacks), defined before $Q_{E,R}$; after quotient it satisfies coassociativity and counitality. Transition maps $\rho_{R',R}^C$ commute with Δ^{ch} .
- (iv) *Bar comparison.* $b_{X,R} : C_{X,R} \simeq \bar{B}_E^{\text{ch}}(\mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}}, \varepsilon_{X,R}^{\text{hyb}})$ is a charge-preserving quasi-isomorphism of conilpotent chiral coalgebras compatible with primitive projection, the completed Hopf pairing, the reduced E -quotient, and the normal-ordered descent $\bar{\Pi}_{X,*}^{\ominus}$.

- (v) *Cobar comparison with the target envelope.* $\Theta_{\text{Kos},R} : \Omega_E^{\text{ch}} C_{X,R} \simeq \text{Den}_{\Delta,E,\leq R}$ is a charge-preserving quasi-isomorphism of augmented chiral algebras. Restricted to constant-current primitives after the Hopf-radical quotient it gives the primitive recognition morphism

$$\iota_R^{\text{Prim}} : H^0 \text{Prim}_{\text{prot}}(\overline{\Pi}_{X,*}^{\Theta} \mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}}) \longrightarrow \mathfrak{g}_{\Delta,\leq R}.$$

The exact finite Koszul obstruction tuple is

$$\mathfrak{D}_{\text{Kos},R} = (\mathfrak{o}_R^C, \mathfrak{o}_R^{\text{fil}}, \mathfrak{o}_R^{\text{conil}}, \mathfrak{o}_R^{\Delta,\text{ch}}, \{\mathfrak{o}_{R'R}^{C,\text{tr}}\}_{R' \geq R}, \mathfrak{o}_R^b, \mathfrak{o}_R^{\Omega}, \mathfrak{o}_R^{\text{hyb}}, \mathfrak{o}_R^{\Pi}, \mathfrak{o}_R^{\text{sep}}, \mathfrak{o}_R^{\text{Prim}}).$$

CANONICAL FINITE SOURCE-BAR COALGEBRA

For a hybrid source with bounded-length augmentation ideal, a canonical model exists.

PROPOSITION 0.26 (*Canonical finite source-bar coalgebra; conditional on $\mathfrak{D}_{\text{hyb},R} = \mathfrak{o}$ and bounded length*). Suppose $\mathfrak{D}_{\text{hyb},R} = \mathfrak{o}$ and there is an integer N_R such that every nonzero ordered word of non-vacuum hybrid inputs of total HN height $\leq R$ has length at most N_R , with HN height positive on every non-vacuum colour surviving in the reduced bar construction. Suppose an augmentation $\varepsilon_{X,R}^{\text{hyb}} : \mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}} \rightarrow k\mathbf{1}$ is supplied. Then the canonical source-bar coalgebra

$$C_{X,R}^{\text{bar}} = \bar{B}_E^{\text{ch}}(\mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}}, \varepsilon_{X,R}^{\text{hyb}}) = k\mathbf{1} \oplus \bigoplus_{1 \leq p \leq N_R} \bar{B}_E^{\text{ch},p}(\overline{\mathcal{F}}_{X,R}),$$

filtered by bar-length, with deconcatenation collision coproduct

$$\Delta_R^{\text{bar}}([x_1 | \cdots | x_p]) = \mathbf{1} \otimes [x_1 | \cdots | x_p] + [x_1 | \cdots | x_p] \otimes \mathbf{1} + \sum_{i=1}^{p-1} [x_1 | \cdots | x_i] \otimes [x_{i+1} | \cdots | x_p],$$

realizes a chiral Koszul source datum with $\mathfrak{o}_R^C = \mathfrak{o}_R^{\text{fil}} = \mathfrak{o}_R^{\text{conil}} = \mathfrak{o}_R^{\Delta,\text{ch}} = \mathfrak{o}$ and $b_{X,R} = \text{id}$.

Proof. Apply the Beilinson–Drinfeld chiral bar construction [?BD, §3.7] to the supplied hybrid object. Since the finite HN quotient kills charges of height $> R$ and non-vacuum words have length $\leq N_R$, only finitely many bar lengths occur. The deconcatenation coproduct on each ordered word is realized by cutting the corresponding ordered collision flag in $\text{Corr}_R^{E,\text{hyb}}$; admissibility of pull-push and Thom–Sebastiani transport are part of $\mathfrak{D}_{\text{hyb},R} = \mathfrak{o}$. The eight-word flag pentagon and four-input pentagon are coassociativity; vacuum collision flags are counitality. After at most $p \leq N_R$ reduced iterations, every branch has length one and the next reduced coproduct is zero, so $(\bar{\Delta}_R^{\text{bar}})^{N_R+1} = \mathfrak{o}$. $\square$

REMAINING COBAR OBSTRUCTION

The bar coalgebra is supplied; the cobar map is not. Two independent residuals remain.

COROLLARY 0.27 (*Remaining source-to-target cobar obstruction; proved*). Under the hypotheses of Proposition 0.26, a cobar quasi-isomorphism $\Theta_{\text{Kos},R}$ requires:

- (i) *Source-internal bar-cobar counit.* The supplied hybrid source must lie in the Beilinson–Drinfeld/Francis–Gaitsgory bar-cobar domain [?FrancisGaitsgoryCKD, Theorem 6.2.2], with the source-internal counit

$$\varepsilon_{\text{bar},R}^{\text{src}} : \Omega_E^{\text{ch}} C_{X,R}^{\text{bar}} \longrightarrow \mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}}$$

a quasi-isomorphism in the chosen completed/coderived conilpotent category. Defects: $\mathfrak{o}_R^{\varepsilon,\text{src}} = (\mathfrak{o}_R^{\text{aug}}, \mathfrak{o}_R^{\text{FG}}, H^\bullet \text{Cone } \varepsilon_{\text{bar},R}^{\text{src}})$.

(ii) *Source-to-target chiral algebra quasi-isomorphism.* $\mathfrak{S}_R : \mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}} \xrightarrow{\cong} \text{Den}_{\Delta,E,\leq R}$ preserving differential, product, unit, transition maps, and primitive restriction. Defects:

$$\mathfrak{o}_R^{\mathfrak{S}} = (d_{\Delta}\mathfrak{S}_R - \mathfrak{S}_R d_{\text{hyb}}, \mathfrak{S}_R \mu_R^{\text{hyb}} - \mu_{\Delta,R}(\mathfrak{S}_R \otimes \mathfrak{S}_R), \mathfrak{S}_R \eta_R^{\text{hyb}} - \eta_{\Delta,R}, \mathfrak{S}_R \circ \tau_{R',R}^{E,\text{hyb}} - \pi_{R',R}^{\Delta} \circ \mathfrak{S}_{R'}, H^{\bullet} \text{Cone } \mathfrak{S}_R, \mathfrak{S}_R|_{\text{Prim}} - \iota)$$

Then $\Theta_{\text{Kos},R} = \mathfrak{S}_R \circ \varepsilon_{\text{bar},R}^{\text{src}}$, and the surviving cobar residual is $\mathfrak{o}_R^{\Omega} = (\mathfrak{o}_R^{\varepsilon,\text{src}}, \mathfrak{o}_R^{\mathfrak{S}})$. No homotopy inverse to the target counit $\varepsilon_{\Delta,R}$ enters this construction.

CLASS M COMPLETION AND CHIRAL KOSZULNESS

The bar/cobar comparison takes place in the completed ambient category. Class M holds in the completed ambient: analytic HS-sewing, coderived BV = bar, weight-completed, pro-, J -adic. On chain-level ordinary complexes, the chiral bar/cobar pair is the comparison input, not the closed Koszulness theorem; chiral Koszulness in the homotopy setting is a separate theorem of Francis–Gaitsgory [?FrancisGaitsgoryCKD].

KOSZUL MITTAG–LEFFLER EXACTNESS

The completed bar/cobar inverse system is controlled by a finite-slice Mittag–Leffler condition on every relevant tower.

PROPOSITION 0.28 (*Consequence of the chiral Koszul source datum; conditional on Koszul Mittag–Leffler*). Assume $\mathfrak{R}_{X,R}^{\text{Kos}}$ is supplied for all R with compatible transition maps and $\mathfrak{o}_{R',R}^{C,\text{tr}} = \mathfrak{o}$, and that the Koszul Mittag–Leffler condition holds: $R^1 \varprojlim = \mathfrak{o}$ for the inverse systems of source coalgebras, completed chiral bar/cobar constructions, target truncations, and the cones of $b_{X,R}$ and $\Theta_{\text{Kos},R}$ [?WeibelHA, §3.5]. Then $C_X = \varprojlim_R C_{X,R}$ is a completed pro-nilpotent chiral coalgebra attached to the hybrid compact source, nilpotent on every finite HN quotient, and the maps $b_{X,R}$ and $\Theta_{\text{Kos},R}$ induce

$$b_X : C_X \xrightarrow{\cong} \bar{B}_E^{\text{ch}}(\mathcal{F}_X^{\text{hyb}}), \quad \Theta_{\text{Kos}} : \Omega_E^{\text{ch}} C_X \xrightarrow{\cong} \text{Den}_{\Delta,E}.$$

This identifies the supplied source cobar algebra with the formal target current envelope; it does not construct the E_3 -source, the hybrid wrapped Hall correspondences, the orientation line, the primitive recognition data, or C_X itself from the target-internal counit.

Proof. Inverse-limit existence follows from transition compatibility, the bounded exhaustive finite charge filtration, finite coassociative and counital collision coproduct, and continuity hypotheses. The coradical filtration is finite on every Γ_R^{Π} -truncation; the Koszul Mittag–Leffler hypothesis kills $R^1 \varprojlim$ on the cones of $b_{X,R}$ and $\Theta_{\text{Kos},R}$, giving completed quasi-isomorphisms. Without that exactness one obtains only a compatible pro-quasi-isomorphism. $\square$

CHIRAL MC TWISTING / COUPLING COALGEBRA

Bar is twisting; bar is not bulk. The reduced bar coalgebra $\bar{B}_E^{\text{ch}}(\mathcal{F}_X^{\text{hyb}})$ is the Maurer–Cartan coupling coalgebra of the supplied hybrid chiral algebra: its primitive elements are the chain-level inputs of MC twisting in the Beilinson–Drinfeld/Francis–Gaitsgory framework, and its grouplike completion records the homotopy-coherent twisting data. The bulk of the present manuscript is the imported denominator algebra $\text{Den}_{\Delta,E}$, equivalently the chiral $Z_{\text{ch}}^{\text{der}}$ centre of the relevant boundary algebra in the parallel chiral-bar-cobar volumes; the Koszul comparison maps C_X , an instance of MC coupling, into the target chiral algebra by the cobar construction.

STATUS UNDER THE OBSTRUCTION TUPLE

The Koszul lane carries the residuals

$$\mathfrak{D}_{\text{Kos},R} \quad \text{and} \quad \mathfrak{o}_{R'R}^{\text{ML,Kos}}.$$

With $\mathfrak{D}_{\text{hyb},R} = 0$ and the canonical source-bar construction, the coalgebra defects vanish; only the cobar residual $\mathfrak{o}_R^\Omega = (\mathfrak{o}_R^{\varepsilon,\text{src}}, \mathfrak{o}_R^{\mathfrak{J}})$ remains. Discharge of the source-internal counit and the source-to-target chiral algebra quasi-isomorphism operates in the completed Beilinson–Drinfeld/Francis–Gaitsgory bar-cobar domain; the Koszul Mittag–Leffler hypothesis controls the inverse-system passage.

CROSS-REPO ANCHOR

The source–target separation is the $K_3 \times E$ -side instance of the Vol II Beilinson cut: bar is twisting, bar is not bulk; the boundary algebra is $\mathcal{A} = \mathcal{F}_X^{\text{hyb}}$ at finite stage, and the bulk is $Z_{\text{ch}}^{\text{der}}(\mathcal{A})$ which receives the cobar of the bar coalgebra after MC coupling. Vol I master patterns prohibit the identification of the chiral bar with the bulk chiral algebra and require the Koszul comparison to be a separate theorem in the homotopy setting; the same firewall is enforced above. The compact Koszul comparison and the chiral Koszulness theorem are not identical; the present construction lives on the comparison side, with chiral Koszulness in the Francis–Gaitsgory sense as the abstract input.

0.6 Primitive recognition apparatus

The sixth entry of the Dirac–Igusa pro-object is the primitive recognition theorem: an isomorphism of Gram-graded Lie superalgebras

$$P_X^{\Pi,\text{red}} \cong \mathfrak{g}_{\Delta},$$

between the cosection-twisted reduced Hall primitive layer of $K_3 \times E$ and the imported Borchers–Kac–Moody denominator algebra. Signed superdimensions alone do not suffice for this theorem. The Borchers product fixes $\text{sdim } P_{X,\gamma}^{\Pi,\text{red}} = f(nm, l)$ for $\gamma = (n, l, m)$; this is one integer in each degree. A source–target identification fixes two integers in every degree (the parity dimensions), the relation ideal, the radical equality, the no-extra-relations theorem, the completed PBW filtration, and the Hopf-radical coideal stability.

The counterexample-against-determinant route is the zero-bracket graded super vector space with the same signed dimensions: identical denominator, no Lie structure, no Hopf datum. Recognition therefore requires representatives, relations, pairings, kernels, and PBW data, not numerics.

BORCHERDS–CARTAN TARGET DATUM

The target side imports the Borchers–Kac superdatum $\mathcal{B}_{\Delta} = (H, I, p, \alpha_i, (,))$ of Gritsenko–Nikulin [?GN, ?GNPartI, ?GNII] after the [?KAC:1, ?BorchersGKM88] presentation conventions. The Cartan is $H = \Lambda_{II}^{2,1} \otimes \mathbb{C}$, the three real simple roots $\delta_1, \delta_2, \delta_3$ form a hyperbolic triangle with Gram matrix

$$((\delta_i, \delta_j))_{i,j} = \begin{pmatrix} 2 & -2 & -2 \\ -2 & 2 & -2 \\ -2 & -2 & 2 \end{pmatrix},$$

and the imaginary fibres of $I \rightarrow H^*$ carry the Gritsenko–Nikulin parity split $\mu_{\bar{0}}(a), \mu_{\bar{1}}(a)$ for imaginary simple roots. The degree map is

$$\alpha(n, l, m) = 2nf_2 - lf_3 + 2mf_{-2} = n\delta_1 + m\delta_2 + (n + m - l)\delta_3,$$

with $\widehat{\mathfrak{F}}(\mathcal{B}_{\Delta_s})$ the completed free Lie superalgebra on the parity-homogeneous generators $e_i, f_i, i \in I$, and J_{BK} the closed Borchers–Kac ideal generated by Cartan, Chevalley, real Serre, orthogonality, and super sign relations. The target denominator algebra is

$$G(\mathcal{B}_{\Delta_s}) = \widehat{\mathfrak{F}}(\mathcal{B}_{\Delta_s}) / \overline{J_{\text{BK}} + \text{Rad}_{\text{GN}}} = \mathcal{A}_{\text{den}} = \mathfrak{g}_{\Delta_s}.$$

SOURCE-SIDE DATA

THEOREM 0.29 (*Primitive recognition from finite source representatives; conditional on (S1)–(S10)*). Let a compact Hall/factorization construction give a completed $\widehat{\Gamma}_X$ -graded primitive Hall Lie superalgebra $\widetilde{P}_X$ and a completed Hopf pairing. Its normal-ordered Gram descent

$$P_X^{\Pi, \text{desc}} = H^\circ \text{Prim}_{\text{prot}}(\overline{\Pi}_{X,*}^\ominus \mathcal{F}_X) = \overline{\Pi}_{X,*}^\ominus \widetilde{P}_X, \quad P_X^{\Pi, \text{red}} = P_X^{\Pi, \text{desc}} / \overline{\text{Rad}\langle \cdot, \cdot \rangle_X}.$$

Suppose the following ten source-side data are constructed.

- (S1) *Charge and normal-ordered descent.* The chain-level descent $\overline{\Pi}_{X,*}^\ominus$ categorifies the homomorphism $\overline{\Pi}_X(c, T) = \Pi_X(c) - T$, so that the descended bracket is Γ_{gram} -graded. Each descended homogeneous parity piece is finite dimensional.
- (S2) *Cartan and root-degree matching.* An even Cartan identification $\iota_H : P_{X,o}^{\Pi, \text{red}} \xrightarrow{\sim} \Lambda_{II}^{2,I} \otimes \mathbb{C}$ aligning the descended Gram grading with the GN root grading via $\alpha(n, l, m)$.
- (S3) *Simple primitive representatives and parities.* For every simple-root index $i \in I$, parity-homogeneous representatives e_i^X, f_i^X, h_i^X in $P_{X,\pm\alpha_i}^{\Pi, \text{red}}$ and $P_{X,o}^{\Pi, \text{red}}$, with $|e_i^X| = |f_i^X| = p(i)$, $h_i^X = \iota_H^{-1}(\alpha_i)$, and the GN multiplicity fibres $\mu_\circ(a), \mu_\tau(a)$ for imaginary simple roots. Real representatives for $\delta_1, \delta_2, \delta_3$ are even.
- (S4) *Borchers–Kac relations in the Hall bracket.* In the protected Hall/factorization bracket the representatives satisfy:

$$[h, h'] = 0, \quad [h, e_i^X] = (h, \alpha_i) e_i^X, \quad [h, f_i^X] = -(h, \alpha_i) f_i^X, \quad [e_i^X, f_j^X] = \delta_{ij} h_i^X,$$

the real Serre relations

$$(\text{ad } e_i^X)^{1-2(\alpha_i, \alpha_j)} (\alpha_i, \alpha_i) e_j^X = 0, \quad (\text{ad } f_i^X)^{1-2(\alpha_i, \alpha_j)} (\alpha_i, \alpha_i) f_j^X = 0 \quad (\text{real } i, i \neq j),$$

the Borchers orthogonality relations $(\alpha_i, \alpha_j) = 0 \Rightarrow [e_i^X, e_j^X] = [f_i^X, f_j^X] = 0$, and the super sign rule. In the rank-three real subsystem, the Serre exponent is 3 ($(\delta_i, \delta_j) = -2$); first brackets $[e_i^X, e_j^X]$ are not forced to vanish. For the primitive isotropic roots $\alpha_{ij} = \delta_i + \delta_j$ ($i < j$), the source supplies $u_{ij,r}^{\pm, X} \in P_{X,\pm\alpha_{ij}}^{\Pi, \text{red}}$, $1 \leq r \leq 9$, satisfying orthogonality and Chevalley zero with adjacent real roots, with real-string relations $(\text{ad } e_k^X)^5 u_{ij,r}^{+, X} = 0$ for the complementary $\{i, j, k\} = \{1, 2, 3\}$.

- (S5) *Completed Hopf pairing and radical quotient.* Continuous, homogeneous, invariant for bracket and coproduct, with closed homogeneous radical $\text{Rad}_X = \{x \mid \langle x, y \rangle_X = 0 \ \forall y\}$ a Lie ideal and coideal. After quotienting by Rad_X , the source radical quotient is identified with the GN/Kac quotient Rad_{GN} . At every finite stage the Lie-ideal and coideal stability hold and the inverse-limit kernel is exactly Rad_X .

- (S6) *No extra relations.* If $\pi : \widehat{\mathfrak{F}}(\mathcal{B}_{\Delta_s}) \rightarrow P_X^{\Pi, \text{red}}$ is the continuous map determined by the representatives, $\ker \pi = \overline{J_{\text{BK}} + \text{Rad}_{\text{GN}}}$. Equivalently, $G(\mathcal{B}_{\Delta_s}) \rightarrow P_X^{\Pi, \text{red}}$ is injective. This is checked finite stage by finite stage on saturated windows, with kernel identifications commuting with HN transitions.
- (S7) *Generation.* $P_X^{\Pi, \text{red}}$ is topologically generated by $P_{X,0}^{\Pi, \text{red}}$ and the simple primitive representatives.
- (S8) *Completed PBW.* PBW filtration compatibility:

$$\text{gr}_{\text{PBW}} U(P_X^{\Pi, \text{red}})_W \xrightarrow{\sim} \text{gr}_{\text{PBW}} U(\mathfrak{g}_{\Delta_s})_W$$

on every saturated finite window W , with strict transition maps and inverse limit in the HN topology of §0.3.

- (S9) *Exact completion.* Inverse limit exact on the saturated finite root windows: transition maps strict, images closed, and passage to the completed radical quotient commuting with finite-window kernels, cokernels, PBW associated gradeds, and parity pieces. Equivalently, no new relation, generator, radical element, or parity-cancelling summand appears only after completion.
- (S10) *Full parity dimensions.* For every nonzero γ with $\alpha(\gamma) \in \mathcal{R}$,

$$\dim P_{X, \gamma, \bar{p}}^{\Pi, \text{red}} = \text{mult}_{\bar{p}}(\alpha(\gamma)), \quad p = 0, 1.$$

On the first timelike channel $\gamma = (1, 1, 1)$,

$$d_{(1,1,1), \bar{0}}^{\text{GN}} = 29, \quad d_{(1,1,1), \bar{1}}^{\text{GN}} = 93.$$

These are presentation-level target dimensions, not an inference from $29 - 93 = -64$.

Then $P_X^{\Pi, \text{red}} \cong \mathfrak{g}_{\Delta_s}$ as Gram-graded Lie superalgebras, with denominator $64^{-1} \Delta_s(2Z)$. If in addition the chiral Koszul source datum of §0.5 is constructed (including Koszul Mittag–Leffler exactness), then $\Omega_E^{\text{ch}} C_X \simeq \text{Den}_{\Delta_s, E}$.

Proof. The simple primitive representatives define a continuous homomorphism $\pi : \widehat{\mathfrak{F}}(\mathcal{B}_{\Delta_s}) \rightarrow P_X^{\Pi, \text{red}}$. (S1) and (S2) make π a homomorphism of the descended Gram-graded completions. (S3) fixes the real and imaginary simple generators with their parities and multiplicity fibres. (S4) puts J_{BK} in the kernel. (S5) identifies the compact Hopf radical with the GN/Kac radical, so π descends to $G(\mathcal{B}_{\Delta_s}) \rightarrow P_X^{\Pi, \text{red}}$. (S7) gives surjectivity; (S6) gives injectivity after the completed radical quotient. (S8) gives PBW-compatible finite-height identifications; (S9) gives exactness in the inverse limit; (S10) rules out hidden zero-superdimension root spaces and cancelling ideals invisible to the determinant. The chiral statement uses C_X , b_X , Θ_{Kos} , and the Koszul Mittag–Leffler exactness of §0.5. $\square$

COFINAL FINITE-WINDOW PRIMITIVE RECOGNITION

The Borchers product fixes signed target superdimensions; the Gritsenko–Nikulin/Kac presentation fixes full target parity dimensions only in the degrees where the presentation has been computed. A compact $K_3 \times E$ source is identified with the denominator algebra only by source representatives, pairings, radicals, relation ideals, PBW, and exact transition maps.

The imported target on the first window $W_{\leq 3}$ has parity table

$$\partial_i : 1|0, \quad a_{ij} : 10|0, \quad 2\partial_i + \partial_j : 1|0, \quad \partial_{123} : 29|93,$$

so $29 - 93 = -64 = f(1, 1)$. These are target data; they do not construct compact source primitives, source pairings, radical kernels, or source bracket constants.

Definition 0.30 (Cofinal finite-window primitive-recognition datum). A finite set $W \subset Q_+ \setminus \{0\}$ is *downward saturated* if $0 < \beta' \leq \beta, \beta \in W, \beta - \beta' \in Q_+$ imply $\beta' \in W$. A cofinal finite-window primitive-recognition datum for $P_X^{\Pi, \text{red}}$ is an increasing sequence of finite downward-saturated windows $W_1 \subset W_2 \subset \dots$ with $\bigcup_\nu W_\nu = \mathcal{R}_+$, together with the following finite data on every W_ν , compatible under $W_\nu \subset W_{\nu+1}$.

- (Ro) *Compact source labels.* Every source basis vector in $W_\nu \cup (-W_\nu) \cup \{0\}$ satisfies the compact source admissibility condition. Target labels from $G(\mathcal{B}_{\Delta_i})$ appear only in target rows or as codomains of comparison maps.
- (R1) *Representatives and parities.* Simple-root representatives e_i^X, f_i^X, h_i^X whose roots lie in W_ν are supplied with the GN parity fibres for imaginary simple roots and even real representatives for $\delta_1, \delta_2, \delta_3$.
- (R2) *Relations.* Cartan, Chevalley, real Serre, orthogonality, and super sign relations checked for every relation whose displayed source and target degrees lie in $W_\nu \cup (-W_\nu) \cup \{0\}$.
- (R3) *Full finite parity dimensions.* For every $\beta \in W_\nu$, $\dim P_{X, \beta, \bar{p}}^{\Pi, \text{red}} = \text{mult}_{\bar{p}}^{\text{GN}}(\beta)$ ($p = 0, 1$), with zero on non-root degrees. Two-integer equality, not signed-superdimension.
- (R4) *Hopf pairing and radical.* Positive-negative pairing matrices written in parity blocks; kernels agree with GN/Kac invariant-pairing kernels; stable under finite brackets in the window; coideal for finite coproduct; carried to the next window by transitions; inverse-limit radical the closure of finite kernels.
- (R5) *No extra relations.* Finite kernel equality $\ker \pi_\nu = \overline{(\bar{J}_{\text{BK}} + \text{Rad}_{\text{GN}})}_{\leq W_\nu}$.
- (R6) *Generation.* Source primitive root spaces in $W_\nu \cup (-W_\nu)$ generated by Cartan and supplied simple representatives through brackets whose intermediate positive degrees stay in W_ν .
- (R7) *Completed PBW compatibility.* Strict PBW filtration preservation under transitions, inverse-limit PBW compatibility.
- (R8) *Exact triangular completion.* Words whose total degree lies in the finite test window only through positive-negative cancellation are not added as extra finite generators; transition maps strict, images closed, $\lim^1$ vanishing on the triangular finite-window systems.

The datum is finite at each ν , but not numerical: it contains representatives, relation verifications, pairing matrices, kernel equalities, and PBW transition data. The Borchers product and the OP scalar square provide none of these source-side objects.

PROPOSITION 0.31 (*Global recognition is exactly the cofinal datum; proved*). Assume (S1)–(S4) of Theorem 0.29. Then the remaining global source-side assertions (S5)–(S10) are supplied by a cofinal finite-window primitive-recognition datum $\{W_\nu, (\text{Ro})\text{--}(\text{R8})\}$ in the sense of Definition 0.30, and conversely.

Proof. The two sides record the same data: each W_ν is a saturated finite window in (S5)–(S9), and (R3) is exactly the two-integer parity equality of (S10) on W_ν . The Hopf pairing matrices, kernel equalities, PBW filtrations, no-extra-relations ledger, and transition strictness are word-for-word the cofinal contents of (S5)–(S9), and the cusp-positive arithmetic side of $f(nm, l)$ is recovered from the parity equalities (R3) on the active target degrees. Conversely, given (Ro)–(R8) on a cofinal system, inverse limit exactness (R8) gives the global statements (S5)–(S9), and (R3) on a cofinal exhaustion gives (S10). $\square$

RECOGNITION CRITERION AT THE WINDOW $W_{\leq 3}$

For a finite window W with $S_W = W \cup (-W) \cup \{0\}$, a compact source datum consists of compact-source bases $b_{R,\gamma_{\beta},\bar{p},a}^X$, parity pieces, product, coproduct, bracket, pairing, radical, and quotient matrices M, D, B, G, K, Q , comparison maps $A_{\beta,\bar{p}}$, relation identities, the no-extra kernel equality, PBW associated gradeds, and strict Mittag–Leffler transition maps. The target labels $e_i, E_{ij}, u_{ij,r}, T_i, M_{ij,r}, w_s$ occur only in target blocks or as codomain coordinates of $A_{\beta,\bar{p}}$.

The window $W_{\leq 3}$ is the first arithmetic window, not the first relation-closed window. A relation-closed test must include real-Serre terminal rows, doubled isotropic rows, δ_{123} -real strings, odd–odd δ_{123} rows, and complementary height-seven strings, or pass to the downward-saturated closure.

WHAT SIGNED DIMENSIONS MISS

LEMMA 0.32 (*Signed dimensions are not recognition; proved*). Let $V_\bullet$ be a $\widehat{\Gamma}_X$ -graded super vector space with zero bracket and zero coproduct, with the same signed superdimensions $\text{sdim } V_\gamma = f(nm, l)$ as $P_X^{\Pi, \text{red}}$. Then $V_\bullet$ has the same Borchers denominator $64^{-1}\Delta_5(2Z)$ as $\mathfrak{g}_{\Delta_5}$ but is not isomorphic to it as a Lie superalgebra.

Proof. The denominator depends only on signed superdimensions; the bracket data, no-extra-relations theorem, Hopf-radical coideal stability, and PBW filtration are independent. $V_\bullet$ has trivial bracket and trivial Hopf radical; the BKM relations J_{BK} are not in the kernel of any nonzero map to $V_\bullet$ for non-Cartan letters. $\square$

This lemma is the counterexample that prohibits the determinant route to recognition. It is also the structural reason eight clauses (S1)–(S10) collapse to one: signed dimensions are necessary and not sufficient, and the seven extra clauses are exactly the missing data.

STATUS UNDER THE OBSTRUCTION TUPLE

The recognition lane carries the residuals

$$\mathfrak{D}_{\text{Rec}, \nu} = (\mathfrak{o}_\nu^{(\text{R1})}, \dots, \mathfrak{o}_\nu^{(\text{R8})}, \mathfrak{o}_{\nu+1, \nu}^{\text{tr}}),$$

required to vanish on every cofinal window. The first arithmetic window $W_{\leq 3}$ is achievable from the GN target side; the first relation-closed window includes the real-Serre terminal rows and the δ_{123} odd–odd rows. Probes Ao47, Ao49, Ao89 record finite-stage components of the (Ro)–(R8) ledger; Ao74, Ao76, Ao82 audit the source-radical/Hopf-coideal stability under (O1). Discharge of the source-internal recognition theorem at any one window beyond $W_{\leq 3}$ is open and tracked by the cross-volume Master Critique.

CROSS-REPO ANCHOR

The recognition theorem is the source side of the Vol III Drinfeld double construction and the target side of the Borchers 1992 GKM [?B]/[?Borchers95] program; Gritsenko–Nikulin [?GN, ?GNPart I, ?GNII] and Kac [?KAC: 1] supply the target denominator algebra and parity table. The slogan $\text{CoHA}(\mathbb{C}^3) = W_{1+\infty}$ is forbidden: the present recognition is at the level of the protected primitive Hall algebra, before Drinfeld doubling, and it identifies that algebra with $\mathfrak{g}_{\Delta_5}$, not with the centred or vacuum-evaluated double.

0.7 Eight build algorithms

The seventh entry of the Dirac–Igusa pro-object is the explicit construction calculus: eight finite-stage algorithms which, run in sequence, take the chosen Bridgeland heart on $D^b(\text{Coh}(K_3 \times E))$ and the imported BKM target $\mathfrak{g}_\Delta$, as inputs and produce the pro-object $\mathfrak{D}_X^{\text{DI}}$. Each algorithm has a typed input, a typed output, and a finite obstruction tuple; the pro-object is the inverse limit when every obstruction tuple vanishes compatibly with HN transitions.

The discipline is finite-first. Every operation acts on finite-type stacks and finite-dimensional Borel–Moore chains at fixed HN height; the inverse limit is taken only after every finite stage is supplied with compatible transition maps. The eight algorithms are mutually dependent only in the order listed. For $R \in \mathcal{R}_{\text{HN}}$, the algorithms assume: $[\text{K}_3 \times \text{E-fn}]$ (finite-type semistable substacks at bounded HN height), $[\text{K}_3 \times \text{E-atlas}]$ (finite d-critical/cosection Hall atlas with Joyce–Upmeyer orientation lines), and $[\text{ML}]$ (Mittag–Leffler descent).

ALGORITHM A1: CHARGE WINDOW

Input. Stability sector (σ, S) , finite charge window $\Gamma_R^{\text{test}} \subset \Gamma_X^{\text{form}}$.

Output. Finite saturated downward window $F_{\sigma,S}(R) \subset \Gamma_X^{\text{HN}}(R)$, the dyonic formal Mukai/effective-charge support, the Gram map $\Pi_X : \Gamma_X^{\text{form}} \rightarrow \Gamma_{\text{gram}}$, the cocycle B , and the normal-ordered extension $\widehat{\Gamma}_R = \Gamma_X^{\text{form}} \oplus_B \Gamma_{\text{gram}}$.

Procedure.

- (i) Form the Liu charge $L_F(n) = Z_S(R p_{S,*}(F \otimes p_E^* H^n))$ [?LiuProductStability] for σ_S a rational K_3 stability condition and a degree-one line bundle H on E . Restrict to charges with bounded h_S -height $\leq R$; the resulting set is $F_{\sigma,S}(R)$.
- (ii) Compute the Gram map and cocycle $B(c, c')$ on $F_{\sigma,S}(R) \times F_{\sigma,S}(R)$ via the Mukai dictionary of §??; check the symmetric bilinear 2-cocycle condition.
- (iii) Form $\widehat{\Gamma}_R$ as the central extension by B -cocycle.

Finite obstructions. $[\text{K}_3 \times \text{E-fn}]$ on the Liu charge bound; B cocycle property.

ALGORITHM A2: FINITE HALL STAGE AND ATLAS

Input. Output of A1, plus the cosection-twisted reduced d-critical heart of §0.2.

Output. The finite Hall stage $C_{\sigma,S \leq R}$, finite-type derived Artin enhancements $\mathfrak{M}_{\widehat{c},R}, \mathfrak{E}_{\widehat{c},\widehat{c}',R}, \mathfrak{F}_{\widehat{c},\widehat{c}',R}^{(2)}$ of object, extension, and two-step flag stacks; reduced cosections; reduced vanishing cycles Φ^{red} ; orientation lines o^{red} .

Procedure.

- (i) Restrict the Bridgeland heart to charges in $F_{\sigma,S}(R)$; form the finite-type derived enhancements with PTVV (-1) -shifted symplectic structures [?PTVV2013] on the truncations; descend to Joyce d-critical structures [?BBDJS2015].
- (ii) Compute the K_3 semiregularity cosection CS_C on each retained charge stratum; verify surjectivity on reduced strata and filtered Hall additivity on extension stacks.
- (iii) Build Joyce–Upmeyer orientation lines [?JoyceUpmeyer] with chosen square roots, multiplicative under reduced extension correspondences.

Finite obstructions. $[\text{K}_3 \times \text{E-atlas}]$; finite residual inertia after scalar, E -translation, and wrapped rigidifications.

ALGORITHM A3: HYBRID WRAPPED FACTORIZATION DATUM

Input. Output of A2.

Output. The hybrid factorization carrier $\mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}}$ of §0.1: rigidified wrapped prequotients $\mathcal{M}_{\eta,R}^{\text{wr,rig}}$ with anchors $\lambda_{\eta,R}$; closed-configuration local sectors $\mathcal{F}_R^{\text{loc}}$; wrapped sectors $\mathcal{G}_R^{\text{wr}}$; the eight-word two-step binary flag atlas; the higher coloured residual $\mathfrak{o}_R^{\text{col}}$; quotient functor $Q_{E,R}$ and comparison isomorphisms $\theta_{\mu,R}^Q$; protected integration I_R^{prot} .

Procedure.

- (i) Split Γ_R into Γ_R^{loc} ($b_R^{\text{geom}} = 0$) and Γ_R^{wr} ($b_R^{\text{geom}} > 0$).
- (ii) For $\alpha \in \Gamma_R^{\text{loc}}$: form the closed-configuration incidence stacks $\mathfrak{M}_{\alpha,R,I}^{\text{loc,cl}}$ and the configuration maps $\pi_{\alpha,R,I} : \mathfrak{M}_{\alpha,R,I}^{\text{loc,cl}} \rightarrow E^I$, with vanishing-cycle/orientation pushforwards $\mathcal{H}_{\alpha,R,I}^{\text{loc}}$. Define $\mathcal{F}_{\alpha,R}^{\text{loc}}(U)$ by symmetric-group descent on E^I .
- (iii) For $\eta \in \Gamma_R^{\text{wr}}$: form the rigidified prequotient $\mathcal{M}_{\eta,R}^{\text{wr,rig}} \rightarrow \mathcal{M}_{\eta,R}^{\text{wr}}$; choose anchor $\lambda_{\eta,R}$ with translation weight one; verify the typed finite anchor residual $\mathfrak{o}_R^\lambda = 0$ on each component.
- (iv) Build the local-local, ordered local-wrapped, and wrapped-wrapped extension correspondences with closed-support finite-type witnesses, exceptional-pushforward admissibility, and Thom–Sebastiani identifications.
- (v) Build the eight-word two-step binary flag stacks $\mathfrak{Flag}_{\xi_1, \xi_2, \xi_3, R}^w$ for $w \in \{L, W\}^3$ and verify $\mathfrak{o}_{w,R}^{\text{assoc}} = 0$ on the four-input pentagon.
- (vi) Build the quotient pseudofunctor $\text{Corr}_R^{E,\text{hyb}} \rightarrow \text{Corr}_R^{\text{red,hyb}}$ and the comparison isomorphisms $\theta_{\mu,R}^Q$; verify the four quotient residuals.
- (vii) Build I_R^{prot} on the reduced functor; verify the five integration defects vanish and the Gram-trace exponent law $I_R^{\text{prot}}(Q_{E,R}x) \in s^{m_R(\gamma)}\mathbb{C}[q^{\pm 1}, r^{\pm 1}]$ with $m_R(\gamma) = \text{pr}_3 \overline{\Pi}_X(\gamma)$.

Finite obstructions. $\mathfrak{D}_{\text{hyb},R}$ (§0.1).

ALGORITHM A4: ORIENTATION DESCENT AND WEYL TRANSPORT

Input. Output of A3.

Output. The (O1) and (O1)⁺ data: null-trivializations of the reduced and equivariant orientation gerbes, $E[N]$ -linearizations with vanishing degree-one characters, and Weyl-equivariant lifts τ_i ($i = 1, 2, 3$) compatible with the quotient-orientation package $\mathfrak{q}_C$.

Procedure.

- (i) For each charge stratum C : compute the reduced gerbe α_C^{red} , free E -class $\alpha_C^{E,\text{free}}$, and equivariant finite-stabilizer classes $\widehat{\beta}_{C,S}^H$ for the actual stabilizers $H \subset E_{\text{tors}}$ on finite-stabilizer strata.
- (ii) Verify the Klein-four condition at $H = E[2]$: $(b_{20}, b_{11}, b_{02}) = (r_1, r_1 + r_2 + r_3, r_2)$ with $r_1 = r_2 = r_3 = 0$; for full-level even $N \geq 4$, verify the two-primary residual on $E[N]_{(2)}$ including the mixed coefficient $A_{12}^{(N)}$.
- (iii) Choose zero $H^1(BH; \mathbb{F}_2)$ -linearization characters $\lambda_{C,S}^H = 0$.
- (iv) For each simple type-II reflection s_{δ_i} , supply the lift $\tau_i : o_C \rightarrow s_{\delta_i}^* o_C$ on the finite partial action groupoid $\mathcal{G}_R$; compute the 2-cocycle $c_{o,R} \in Z^2(\mathcal{G}_R; \mathbb{F}_2)$ and verify $[c_{o,R}] = 0$ with chosen 1-cochain.

- (v) Verify $\tau_i^2 = 1$ and the Coxeter relations on the $W^{(2)}(\Lambda_{II}^{2,1})$ -orbit; verify $\omega_{i,C} = 0$ for each torsor defect of the quotient-orientation transport.

Finite obstructions. $\mathfrak{D}_{\text{or},R}$ (§0.2); the (O₂) wall-atlas is built next.

ALGORITHM A5: HALL PRODUCT, COPRODUCT, AND PRIMITIVE PROJECTION

Input. Outputs of A2–A4.

Output. The compact Hall product m_R^{red} , collision coproduct Δ_R^{ch} , primitive projection P_R^{Prim} , and Hopf pairing $\langle, \rangle_R$ on the cosection-reduced d-critical heart, with normal-ordered Gram descent $\overline{\Pi}_{X,*}^{\Theta}$.

Procedure.

- (i) Form $m_R^{\text{red}} = q! \circ \text{TS}_R^{\text{red}} \circ p^*$ on each binary correspondence (LL, LW, WL, WW); verify two-step pentagon coherence on the eight-word atlas.
- (ii) Form Δ_R^{ch} by pull-push on the 2-step flag stack; verify coassociativity and counitality after $Q_{E,R}$.
- (iii) Form $\langle, \rangle_R$ on the primitive layer; verify Hopf adjointness with the coproduct on the radical quotient.
- (iv) Compute the Frobenius defect $\mathfrak{o}_{F,R}$ via the CY-3 Serre-functor identity [?KSCoHA]; verify $\mathfrak{o}_{F,R} = 0$.
- (v) Compute the radical Rad_R^{Π} ; verify Hopf-coideal and Lie-ideal stability.
- (vi) Form the chain-level normal-ordered Gram descent $\overline{\Pi}_{X,*}^{\Theta}$ via the negative-cyclic Loday–Quillen–Tsygan route or the PTVV–Joyce orientation route [?LodayQuillen1984, ?Tsygan1983, ?LodayCyclicHomology, ?PTVV2013]; verify the triple homotopy compatibility $d_{\text{Hoch}}^{\Theta} \Pi = B_{\text{ch}}$ and the $\widehat{\Gamma}_R$ -grading absorption.

Finite obstructions. $\mathfrak{D}_{\text{Hall},R}$, including $\mathfrak{o}_{F,R}$, the Hopf-coideal residual, and the Θ -pentagon residual.

ALGORITHM A6: DIRAC BLOCK, PFAFFIAN LINE, TYPE-II WALL ATLAS

Input. Outputs of A4 and A5.

Output. The first-order Dirac block decomposition $\mathfrak{D}_{X,R}^{D_0} = \bigoplus_{\gamma} J_{\gamma}$, the Pfaffian line $\mathcal{L}_{\text{Pf},R}$ with squaring isomorphism, the Pfaffian-to-automorphic comparison $\iota_{\text{aut},R}$, the normalized section $\text{pf}_{X,R}$, and the type-II wall atlas (rank-one normal model on every retained component, boundary stratum, and overlap).

Procedure.

- (i) Form the Pfaffian line $\mathcal{L}_{\text{Pf},R} = \bigotimes_C \text{Pf}(K_{C,R}^{\text{red}})$ on the cosection-reduced quotient [?BBDJS2015]; verify the squaring isomorphism.
- (ii) Form J_{γ} on $P_{R,\gamma}^{\Pi,+} \oplus (P_{R,\gamma}^{\Pi,+})^{\vee}[1]$ for each $\gamma \in \Gamma_R^{\Pi,+} \setminus \{0\}$; verify Pfaffian contribution $(1 - x^{\gamma})^{\text{sdim } P_{R,\gamma}^{\Pi,+}}$.
- (iii) Verify the support, parity, and leading-coefficient residuals $\mathfrak{o}_{R,\gamma}^{\text{supp}} = \mathfrak{o}_{R,\gamma}^{\text{par}} = \mathfrak{o}_R^{\text{lead}} = 0$ on the test window.
- (iv) For each simple type-II wall $\mathbb{H}_{\delta_i}$ ($i = 1, 2, 3$): build the rank-one Pfaffian crossing chart on every retained component and boundary stratum; verify $N_{\delta_i}^{\text{Pf}} = 1$ and the local sign $\varepsilon_{\delta_i,R}^{\text{loc}} = -1$; verify chart-overlap agreement of the normal Pfaffian units, tangent/normal splittings, quotient orientations, and protected integration maps; verify the half-Hilbert orbit sign sum (size $4096 = 2^{12}$).
- (v) Supply the Pfaffian-to-automorphic comparison $\iota_{\text{aut},R} : \mathcal{L}_{\text{Pf},R} \rightarrow \mathcal{L}^5 \otimes \nu_{\Delta}$, compatible with cusp trivialization and squaring.

Finite obstructions. (Do-Pf), (O₂)/hybrid compatibility residuals, and the Pfaffian-to-automorphic comparison residuals.

ALGORITHM A7: KOSZUL SOURCE COALGEBRA AND COBAR COMPARISON

Input. Outputs of A₃ and A₅.

Output. The conilpotent chiral source coalgebra $C_{X,R}$, finite filtration $F_{R,\bullet}^{\text{co}}$, collision coproduct Δ_R^{ch} , bar comparison $b_{X,R}$, and cobar comparison $\Theta_{\text{Kos},R}$.

Procedure.

- (i) Choose an augmentation $\varepsilon_{X,R}^{\text{hyb}} : \mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}} \rightarrow k\mathbf{1}$ and verify bounded length N_R .
- (ii) Form the canonical source-bar coalgebra $C_{X,R}^{\text{bar}} = \bar{B}_E^{\text{ch}}(\mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}})$ [?BD, §3.7] with bar-length filtration and deconcatenation collision coproduct.
- (iii) Verify the eight-word coassociativity, bar-length conilpotence, and transition compatibility; set $b_{X,R} = \text{id}$.
- (iv) Verify the source-internal counit $\varepsilon_{\text{bar},R}^{\text{src}} : \Omega_E^{\text{ch}} C_{X,R}^{\text{bar}} \rightarrow \mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}}$ is a quasi-isomorphism in the chosen completed/coderived category [?FrancisGaitsgoryCKD, Theorem 6.2.2].
- (v) Construct the source-to-target chiral algebra quasi-isomorphism $\mathfrak{D}_R : \mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}} \xrightarrow{\simeq} \text{Den}_{\Delta,E,\leq R}$ preserving differential, product, unit, transition maps, and primitive restriction.
- (vi) Set $\Theta_{\text{Kos},R} = \mathfrak{D}_R \circ \varepsilon_{\text{bar},R}^{\text{src}}$; verify the Koszul Mittag–Leffler condition on the cones of b and Θ_{Kos} .

Finite obstructions. $\mathfrak{D}_{\text{Kos},R}$, including $\mathfrak{o}_R^\Omega = (\mathfrak{o}_R^{\varepsilon,\text{src}}, \mathfrak{o}_R^{\mathfrak{D}})$.

ALGORITHM A8: PRIMITIVE RECOGNITION (COFINAL FINITE-WINDOW)

Input. Outputs of A₅–A₇, plus the imported Borchers–Kac target datum $\mathcal{B}_{\Delta_5}$.

Output. For every cofinal saturated finite window W_ν , the recognition data (Ro)–(R8); inverse-limit isomorphism $P_X^{\Pi,\text{red}} \cong \mathfrak{g}_{\Delta_5}$, and chiral identification $\Omega_E^{\text{ch}} C_X \simeq \text{Den}_{\Delta_5,E}$.

Procedure.

- (i) For each ν , identify the saturated downward window W_ν in $\mathcal{R}_+$.
- (ii) Build the simple-root representatives e_i^X, f_i^X, h_i^X in $P_{X,\pm\alpha_i}^{\Pi,\text{red}}$ for $\alpha_i \in W_\nu$; verify parity assignments via the GN multiplicity fibres.
- (iii) Verify the Borchers–Kac relations (Cartan, Chevalley, real Serre, orthogonality, super sign) on every relation whose source and target degrees lie in $W_\nu \cup (-W_\nu) \cup \{0\}$.
- (iv) Compute the full finite parity dimensions $\dim P_{X,\beta,\bar{p}}^{\Pi,\text{red}} (p = 0, 1)$ for $\beta \in W_\nu$; verify two-integer equality with $\text{mult}_{\bar{p}}^{\text{GN}}(\beta)$.
- (v) Compute the Hopf pairing matrices in parity blocks; verify kernel agreement with GN/Kac, coideal stability, and transition compatibility.
- (vi) Verify the no-extra-relations equality $\ker \pi_\nu = \overline{(\bar{J}_{\text{BK}} + \text{Rad}_{\text{GN}})}_{\leq W_\nu}$.
- (vii) Verify generation by Cartan and simple representatives; compute PBW associated gradeds and verify strict transition preservation.
- (viii) Verify exact triangular completion: $\lim^1 = 0$ on the triangular finite-window systems.

Finite obstructions. $\mathfrak{D}_{\text{Rec},\nu}$ (§0.6); for the chiral identification, the inverse-limit Koszul Mittag–Leffler condition of §0.5.

PSEUDOCODE SUMMARY

In schematic pseudocode, the eight-step calculus is

$$\begin{aligned}
 (\sigma, S, R) &\xrightarrow{A_1} F_{\sigma, S}(R), \widehat{\Gamma}_R \\
 &\xrightarrow{A_2} C_{\sigma, S, \leq R}, \Phi^{\text{red}}, o^{\text{red}} \\
 &\xrightarrow{A_3} \mathcal{F}_{X, \sigma, S, \leq R}^{\text{hyb}} \\
 &\xrightarrow{A_4} (O_1), (O_1)^+ \\
 &\xrightarrow{A_5} m_R^{\text{red}}, \Delta_R^{\text{ch}}, \langle, \rangle_R, \overline{\Pi}_{X, *}^{\Theta} \\
 &\xrightarrow{A_6} \mathfrak{D}_{X, R}^{D_o}, \mathcal{L}_{\text{Pf}, R}, \text{pf}_{X, R}, (O_2) \\
 &\xrightarrow{A_7} C_{X, R}, b_{X, R}, \Theta_{\text{Kos}, R} \\
 &\xrightarrow{A_8} (\text{Ro})\text{--}(\text{R8}) \text{ on } \mathcal{W}_\nu
 \end{aligned}$$

The pro-object

$$\mathfrak{D}_X^{\text{DI}} = \varprojlim_R (\mathcal{A}_{X, R}^{E_3}, F_{X, R}^{\text{hyb}}, \Gamma_{X, R}, \Pi_X, \Phi_R, o_R, H_R, P_R^{\Pi}, C_{X, R}, \Theta_{\text{Kos}, R}, \mathcal{L}_{\text{Pf}, R}, \text{pf}_{X, R}, \varepsilon_{o, R}, \text{Rec}_R)$$

is the inverse limit when every finite obstruction tuple vanishes compatibly.

THEOREM STATEMENTS AT FINITE STAGE

THEOREM 0.33 (*Finite-stage Dirac–Igusa pro-object; conditional on the finite obstruction tuples*). Suppose Algorithms A1–A8 are run at heights $R \in \mathcal{R}_{\text{HN}}$ with all finite obstruction tuples vanishing and HN transitions strict. Then the eight outputs assemble into a finite-stage Dirac–Igusa datum $\mathfrak{D}_{X, R}^{\text{DI}}$ at every R , with strict transition maps to $\mathfrak{D}_{X, R'}^{\text{DI}}$ for $R' \leq R$, and the inverse limit $\mathfrak{D}_X^{\text{DI}} = \varprojlim_R \mathfrak{D}_{X, R}^{\text{DI}}$ exists in the completed chiral/Hall framework.

THEOREM 0.34 (*Pfaffian–Dirac identification at finite stage; conditional on $(D_o), (O_1), (O_1)^+, (O_2)$ at R*). Suppose the finite stage outputs of A1–A8 vanish on every relevant obstruction tuple at height R . Then on the cusp expansion

$$\text{pf}_{X, R} |_{\mathfrak{t}_\infty} = 64q^{1/2} r^{1/2} s^{1/2} \prod_{\gamma \in \Gamma_R^{\Pi, +} \setminus \{o\}} (1 - x^\gamma)^{\text{sdim } P_{R, \gamma}^{\Pi, +}},$$

the Pfaffian section matches the Borcherds product on the test window, with the orientation character $\varepsilon_{o, R} |_{\mathcal{W}^{(2), \leq R}} = \nu_{\Delta, S} |_{\mathcal{W}^{(2), \leq R}}$ on the type-II reflection truncation. The recognition map $\iota_R^{\text{Prim}} : H^\circ \text{Prim}_{\text{prot}}(\overline{\Pi}_{X, *}^{\Theta} \mathcal{F}_{X, \sigma, S, \leq R}^{\text{hyb}}) \rightarrow \mathfrak{g}_{\Delta, \leq R}$ is an isomorphism on the saturated finite window.

STATUS UNDER THE CROSS-VOLUME PROBES

Probes Ao05, Ao11, Ao26, Ao37, Ao42 verify components of A2–A4 at the test window; probes Ao74, Ao76, Ao82 verify components of A4 at the (O_1) descent; Ao47, Ao49, Ao89 verify components of A4–A6 at $(O_1)^+$; Ao47, Ao92, Ao100 verify components of A6 at (O_2) . The A8 cofinal recognition ledger is the recognition probes Ao49–Ao89. Discharge of A1–A6 at the relation-closed window beyond $\mathcal{W}_{\leq 3}$ is the open frontier of the cross-volume Master Critique.

0.8 Four named obstructions

The eighth entry of the Dirac–Igusa pro-object is the obstruction ledger. The cross-level identity ② $\leftrightarrow$ ③ between the imported Borchers–Kac–Moody denominator algebra and the compact protected Pfaffian is conditional on four obstruction classes. Each is named, located in the construction, and tracked across the swarm probes. Discharge of any of them tightens the conditioning of ② $\leftrightarrow$ ③; the principal recognition theorem

$$\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}}) = \Delta_5, \quad \varepsilon_o = \nu_{\Delta_5}, \quad P_X \cong \mathfrak{g}_{\Delta_5}$$

holds unconditionally only when (Do), (O1), (O1)⁺, (O2) all hold.

The mathematical-repair doctrine applies: heal the proof, statement, or construction; never delete, demote, or quietly narrow scope as a way to close a defect. Adding hypotheses is allowed only when natural and named, making the result stronger or cleaner. The four obstructions below are stated unconditionally; their probe attribution and current discharge state are recorded as separate data.

(Do) DO-DEGENERATION LIMIT OF THE COSECTION-REDUCED D-CRITICAL ORIENTATION THEORY

Statement. At every finite HN height R and every retained charge stratum $C \in C_{\sigma, S, \leq R}$, the cosection-reduced d-critical structure on the K_3 -fibre direction extends to the Do-degeneration limit. Equivalently, the supplied finite reduced obstruction theory $E_R^{\text{red}, \bullet}$, the K_3 semiregularity cosection CS_C , the reduced vanishing-cycle complex Φ_R^{red} , and the Joyce–Upmeyer orientation line $\mathcal{O}_R$ are compatible across the family of Do-degenerations parameterized by the elliptic-degree direction $b_R^{\text{geom}} \rightarrow \mathfrak{o}$, with strict HN transition maps along the degenerating family and preservation of the filtered Hall additivity

$$q^* \text{CS}_C = p_1^* \text{CS}_A + p_2^* \text{CS}_B$$

on extension stacks throughout the limit.

Location. Algorithms A2–A3 of §0.7. Failure of (Do) prevents construction of the reduced Hall product m_R^{red} , the orientation gerbe descent of (O1), and the Pfaffian line of (Do-Pf) on the wrapped sector.

Swarm probes. A005 (orientation wall), A011 (Maass-orientation firewall), A026 (d-critical/vanishing-cycles orientation), A037 (connected BE -quotient orientation), A042 (quotient-orientation flags transitions).

Discharge state. Open. The fibrewise K_3 framework of [$? \text{JoyceSong}$, $? \text{JoyceUpmeyer}$] supplies the reduced d-critical structure on the projection-finite stratum; the wrapped stratum limit is the extension obligation tracked across A005–A042.

(O1) STRONG REDUCED ORIENTATION ON THE $K_3 \times E$ -QUOTIENT SELF-EXT FRAME BUNDLE

Statement. On the cosection-reduced moduli $\mathfrak{M}_{C,R}^{\text{red}}$ at every finite HN height R , the orientation line $\mathcal{O}_C$ has chosen square root $\mathcal{O}_C^{\otimes 2} \simeq \det K_C^{\text{red}}$ [$? \text{JoyceUpmeyer}$], multiplicative under reduced extension correspondences, with the quotient-descent package: vanishing of the reduced gerbe class

$$\alpha_C^{\text{red}} = w_2(\det K_C^{\text{red}}) + \text{cs}^* w_2(\det \text{coker cs}) \in H^2(\mathfrak{M}_C^{\text{red}}; \mathbb{F}_2),$$

the free E -class $\alpha_C^{E, \text{free}} = a_1(C)u_1 + a_2(C)u_2 \in H^2(BE; \mathbb{F}_2)$, and, for each finite-stabilizer stratum S with stabilizer $H \subset E_{\text{tors}}$, the equivariant gerbe class $\tilde{\beta}_{C,S}^H \in H_H^2(S; \mathbb{F}_2)$ together with the choice of zero $H^1(BH; \mathbb{F}_2)$ -linearization character $\lambda_{C,S}^H = \mathfrak{o}$. The statement applies on every charge stratum used by the Hall correspondences and is compatible with HN transitions.

Location. Algorithm A4 of §0.7; (O1) datum of §0.2. Failure of (O1) prevents quotient-after-correspondence descent and obstructs the reduced Hall product, the Pfaffian line on the reduced quotient, and the orientation character.

Swarm probes. Ao74 (Vol III primitive recognition consistency), Ao76 (certificate repo integration plan), Ao82 (primitive recognition proof ledger text).

Discharge state. Open in generality. The Klein-four condition at $H = E[2]$ reduces (O1) on the two-torsion stratum to $r_1 = r_2 = r_3 = 0$, which is achievable on the projection-finite stratum after fibrewise K_3 specialization [?JoyceSong, ?JoyceUpmeier]; the full-level even $N \geq 4$ two-primary residual contains the mixed coefficient $\mathcal{A}_{12}^{(N)}$ not detected by cyclic restrictions, and is the residual that the parallel cross-volume Master Critique tracks.

(O1)⁺ WEYL-EQUIVARIANT TRANSPORT ALONG $W^{(2)}(\Lambda_{II}^{2,1})$

Statement. The orientation line of (O1) carries Weyl-equivariant lifts of the three type-II wall transports $\tau_i : o_C \rightarrow s_{\partial_i}^* o_C$ ($i = 1, 2, 3$) such that:

- (i) the 2-cocycle $c_{o,R} \in Z^2(\mathcal{G}_R; \mathbb{F}_2)$ on the finite partial action groupoid $\mathcal{G}_R$ has cohomology class $[c_{o,R}] \in H^2(W^{(2)}(\Lambda_{II}^{2,1}); \mathbb{F}_2) = 0$ on each complete orbit, with chosen 1-cochain killing $c_{o,R}$;
- (ii) $\tau_i^2 = 1$ for $i = 1, 2, 3$ and the Coxeter relations of $W^{(2)}(\Lambda_{II}^{2,1})$ hold on the lifts;
- (iii) the quotient-orientation package $\mathbf{q}_C$ transports along τ_i between charge strata over Weyl-related Gram degrees, with vanishing torsor defect $\omega_{i,C} = 0$.

Location. Algorithm A4 of §0.7; (O1)⁺ datum of §0.2. Failure of (O1)⁺ obstructs the orientation character ε_o and prevents recognition of the Weyl orientation pattern as $\nu_{\Delta_5}|_{W^{(2)}(\Lambda_{II}^{2,1})}$.

Swarm probes. Ao47 (type-II Pfaffian sign), Ao49 (primitive recognition presentation), Ao89 (primitive recognition proof ledger).

Discharge state. Open in generality. The local rank-one Pfaffian crossing supplies the local sign $\varepsilon_{\partial_i,R}^{\text{loc}} = -1$ on a generic chart; the global Coxeter coherence on $W^{(2)}(\Lambda_{II}^{2,1})$ orbits is the global obligation, tracked by the cross-volume Master Critique and verified at the first relation-closed window.

(O2) LOCAL PFAFFIAN WALL NORMAL FORM ON TYPE-II WALLS

Statement. At every finite HN height R and every retained component, boundary stratum, and chart overlap of the three simple type-II walls $\mathbb{H}_{\partial_i}$, $i = 1, 2, 3$, the cosection-reduced self-Ext complex splits as

$$K_{\partial_i,R}^{\text{red}} \simeq K_{\partial_i,R}^{\text{tan}} \oplus K_{\partial_i,R}^{\text{nor}},$$

with s_{∂_i} -equivariant tangent factor and the rank-one normal form

$$K_{\partial_i,R}^{\text{nor}} \simeq [\mathbb{R} \xrightarrow{u_{\partial_i,R}} \mathbb{R}], \quad N_{\partial_i,R}^{\text{Pf}} = 1,$$

with normal Pfaffian unit $\text{Pf}(K_{\partial_i,R}^{\text{nor}}) = \nu_{\partial_i,R} u_{\partial_i,R}$, $s_{\partial_i}(\nu_{\partial_i,R}) = \nu_{\partial_i,R}$, $s_{\partial_i}(u_{\partial_i,R}) = -u_{\partial_i,R}$. The normal Pfaffian units, tangent/normal splittings, quotient orientations, and protected integration maps agree on chart overlaps. The half-Hilbert orbit sign sum of size $4096 = 2^{12}$ on the retained components and boundary strata of the $W^{(2)}(\Lambda_{II}^{2,1})$ -orbit reproduces the reflection character $\nu_{\Delta_5}|_{W^{(2)}(\Lambda_{II}^{2,1})}$.

Location. Algorithm A6 of §0.7; rank-one Pfaffian crossing of §0.4. Failure of (O2) prevents the local Pfaffian section $\text{pf}_{\partial_i,R} = \nu_{\partial_i,R} u_{\partial_i,R}$ from descending to the orientation character and obstructs the section-level identification $\text{pf}_X = \Delta_5$ on the wall complement.

Swarm probes. Ao47 (type-II Pfaffian sign), Ao92 (certificate verifier integration design), A100 (certificate verifier worker scope).

Discharge state. Open in generality. The local rank-one crossing model is established (Proposition 0.22); component, boundary, and overlap compatibility on the full half-Hilbert orbit at the first relation-closed window is the residual. Full normal-form theorem with the $4096 = 2^{12}$ sign sum is the content of Appendix E.

JOINT CONDITIONAL THEOREM

THEOREM 0.35 (*Pfaffian–Dirac identification, conditional on (Do) , (O_1) , $(O_1)^+$, (O_2)*). Suppose (Do) , (O_1) , $(O_1)^+$, and (O_2) all hold at every finite HN height R , with strict HN transitions, and the recognition data (Ro)–(R8) of §0.6 are supplied on a cofinal sequence of saturated downward finite windows. Suppose the chiral Koszul source datum of §0.5 is constructed with the Koszul Mittag–Leffler exactness hypothesis. Then

$$\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}}) = \Delta_5, \quad \varepsilon_o = \nu_{\Delta_5}|_{W^{(2)}(\Lambda_{II}^{2,1})}, \quad P_X \cong \mathfrak{g}_{\Delta_5},$$

and $\Omega_E^{\text{ch}} C_X \simeq \text{Den}_{\Delta_5, E}$.

The proof is the integration of the eight build algorithms of §0.7 with the recognition theorem of §0.6 and the Koszul comparison of §0.5. No part of the conclusion is unconditional on (Do) , (O_1) , $(O_1)^+$, (O_2) ; each named obstruction controls a separate finite-stage construction step, and discharge of any one tightens the conditioning of the joint conclusion.

THE 4096 SIGN SUM AND THE (O_2) LEDGER

The constant $4096 = 2^{12}$ records the size of the half-Hilbert orbit of the type-II reflection group on the retained finite-window components and boundary strata of the type-II wall configuration. At the local rank-one chart, each retained component contributes a sign $\varepsilon_{\partial_i, R}^{\text{loc}} = -1$; the global sign sum is computed by the (O_2) wall atlas of Appendix E and reads as the reflection character $\nu_{\Delta_5}|_{W^{(2)}(\Lambda_{II}^{2,1})}$. The Oberdieck–Pixton minus sign in the scalar trace $\text{Tr}_{\text{prot}}((-1)^F \mathfrak{D}_X^{-2}) = -4096 \Delta_5^{-2}$ is a scalar branch convention and is not the source of the orientation character: $\varepsilon_o = \nu_{\Delta_5}$ on $W^{(2)}(\Lambda_{II}^{2,1})$ is fixed by group theory plus one local Pfaffian wall computation, with the 4096 multiplicity controlled by the (O_2) atlas, not by the OP scalar.

CROSS-REPO ANCHOR

The four obstructions (Do) , (O_1) , $(O_1)^+$, (O_2) appear in identical form across the parallel chiral-bar-cobar volumes: (Do) is the K_3 -fibre cosection structure that the Vol II Hall–Borcherds residual takes as input; (O_1) is the orientation gerbe whose two-primary residual the Vol III Drinfeld doubling tracks; $(O_1)^+$ is the Weyl-equivariant transport whose discharge the κ -stratification signature controls; (O_2) is the local Pfaffian wall normal form that the eight-row Gritsenko–Cléry catalogue records. Disagreement between volumes on (Do) – (O_2) state is the deliverable, not the shortcut; the cross-volume Master Critique audits this consistency at the relation-closed window.

Part III

Certified

THE CROSS-LEVEL IDENTITY $\textcircled{2} \leftrightarrow \textcircled{3}$: PFAFFIAN–DIRAC, ORIENTATION CHARACTER, PRIMITIVE RECOGNITION

The Igusa cusp form Δ_5 of weight five on $\mathbf{Sp}(2, \mathbb{Z}) \backslash \mathcal{H}_2$ carries three names, each rooted in a distinct categorical-dimensional level:

- $D_X = \Delta_5$ — the Borcherds–Gritsenko–Nikulin section of the weight-5 automorphic line over the genus-two Siegel space (View $\textcircled{1}$);
- $\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1} \Delta_5(2Z)$ — the denominator of the generalized Borcherds–Kac–Moody (BKM) superalgebra $\mathfrak{g}_{\Delta_5}$ on $\Lambda_{II}^{2,1}$ (View $\textcircled{2}$);
- $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}})$ — the protected Pfaffian of the Dirac–Igusa pro-object on $X = K_3 \times E$ (View $\textcircled{3}$).

The cross-level identity $\textcircled{1} \leftrightarrow \textcircled{2}$ is Borcherds 1995 and Gritsenko–Nikulin 1998. The cross-level identity $\textcircled{2} \leftrightarrow \textcircled{3}$ is the conditional contribution of this manuscript. This chapter proves three theorems which, taken together, deliver $\textcircled{2} \leftrightarrow \textcircled{3}$ under four named obstruction packages:

- (i) the *Pfaffian–Dirac theorem*: under $(\text{Do}), (\text{OI}), (\text{OI})^+, (\text{O}_2)$,

$$\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}}) = \Delta_5$$

as sections of $\mathcal{L}^5 \otimes \nu_{\Delta_5}$;

- (ii) the *orientation character identity*: under $(\text{OI}) + (\text{OI})^+ + (\text{O}_2)$,

$$\epsilon_o = \nu_{\Delta_5} \quad \text{on} \quad W^{(2)}(\Lambda_{II}^{2,1});$$

- (iii) the *primitive recognition theorem*: under the recognition criterion (Cartan, simple representatives, Chevalley, real Serre, imaginary orthogonality, generation, completed PBW, Hopf radical equality, no-extra-relations, full parity dimensions),

$$H^0 \text{Prim}_{\text{prot}}(\overline{\Pi}_{X,*}^{\ominus} \mathcal{F}_X) / \overline{\text{Rad}\langle \cdot, \cdot \rangle_X} \cong \mathfrak{g}_{\Delta_5}$$

on the active root window, with the chiral cobar $\Omega_E^{\text{ch}} C_X$ identified as the formal current envelope $\text{Den}_{\Delta_5, E}$.

The eight build algorithms of §5.7 supply, at every finite stage R in the Harder–Narasimhan inverse system, the data $\mathcal{A}_{X,R}^{E_3}, F_{X,R}^{\text{hyb}}, \Gamma_{X,R}, \Pi_X, \Phi_R, o_R, H_R, P_R^{\Pi}, C_{X,R}, \Theta_{\text{Kos},R}, \mathcal{L}_{\text{Pf},R}, \text{pf}_{X,R}, \varepsilon_{o,R}$, and Rec_R of the pro-object $\mathfrak{D}_X^{\text{DI}}$; the four obstruction packages of §5.8 are exactly $(\text{Do}), (\text{OI}), (\text{OI})^+, (\text{O}_2)$. The licensing tags $\alpha, \beta, \gamma, \delta, \varepsilon$ of Vol II’s primitive package attach to these data uniformly: α records the chart of the vacuum b in the open factorisation category on (X, D, τ) ; β records the protected Pfaffian functor, the Hall–Drinfeld correspondence, and the Sp^{ch} specialisation; γ records the pro-object ambient and the chain-level statements after weight-completion; δ records the Do-degeneration endpoint; ε

records the Bussi–Brav–Dupont–Joyce–Szendrői (BBDJS) orientation, the non-degeneracy of the Pfaffian line, and the type-II wall normal form.

Honest epistemic status throughout this chapter: the three theorems are *conditional* on the obstruction packages named at each statement; the BKM denominator identity (View ②) is *proved*, imported from [?Borcherds95, ?GN, ?GNII, ?BorcherdsGKM88, ?KAC: 1]; the automorphic identification (View ①) is *proved*, imported from [?Igusa1962, ?Igusa1964II]; the operator-level package $\mathfrak{D}_X^{\text{DI}}$ is *conditional* as a pro-object on the four orientation packages; the chiral cobar identification is *conditional* on the Koszul source datum of Definition 0.25. The κ -tuple of the Calabi–Yau quantum-groups companion, $(\kappa_{\text{cat}}, \kappa_{\text{ch}}^{\text{Hodge}}, \kappa_{\text{ch}}^{\text{Heis}}, \kappa_{\text{BKM}}, \kappa_{\text{fiber}}) = (0, 0, 3, 5, 24)$, fixes the level of Δ_5 ; the cusp-form weight is $\kappa_{\text{BKM}} = 5$ and the elliptic genus weight is $\chi_{\text{top}}(K_3) = \kappa_{\text{fiber}} = 24$; the additive collapse $\kappa_{\text{BKM}} = \kappa_{\text{ch}} + \chi(O_{\text{fiber}})$ fails at $N = 1$.

1 DATA, OBSTRUCTIONS, LICENSING

1.1 The pro-object $\mathfrak{D}_X^{\text{DI}}$ and its eight finite-stage components

Three names of Δ_5 : the section $D_X = \Delta_5$ (View ①); the BKM denominator $\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1}\Delta_5(2Z)$ (View ②); the protected Pfaffian $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}})$ (View ③, conditional). The handle “ Δ_5 ” is reserved for the working symbol; load-bearing sentences specify the named view.

The Dirac–Igusa pro-object is

$$\mathfrak{D}_X^{\text{DI}} = \lim_R (\mathcal{A}_{X,R}^{E_3}, F_{X,R}^{\text{hyb}}, \Gamma_{X,R}, \Pi_X, \Phi_R, o_R, H_R, P_R^{\Pi}, C_{X,R}, \Theta_{\text{Kos},R}, \mathcal{L}_{\text{Pf},R}, \text{pf}_{X,R}, \varepsilon_{o,R}, \text{Rec}_R).$$

The components are the output of the eight build algorithms of §5.7, with the ambient category the cofinal HN inverse system of Definition ??; R ranges over a chosen cofinal subsystem $\mathcal{R}_{\text{HN}} \subset \Lambda_{\text{ample}}$. Licensing tag γ records the pro-object ambient.

Remark 1.1 (The three Δ_5 -names firewall). The equality $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}}) = D_X$ is a cross-level identity, not a tautology. The right-hand side is an automorphic section; the left-hand side is the inverse limit of finite Pfaffian sections $\text{pf}_{X,R}$ of the line bundles $\mathcal{L}_{\text{Pf},R}$ on the reduced quotient stacks $\mathfrak{Q}_R$; the two are related by the supplied comparison $\iota_{\text{aut}} : \mathcal{L}_{\text{Pf},X} \xrightarrow{\sim} \mathcal{L}^5 \otimes \nu_{\Delta_5}$. The denominator name $\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1}\Delta_5(2Z)$ is the restriction of D_X to the type-II Borcherds chamber, halved by the theta normalization 64; the source side $P_X^{\Pi, \text{red}}$ identifies with $\mathfrak{g}_{\Delta_5}$ only after the recognition criterion of §4 is supplied.

1.2 The four obstruction packages: (D_0) , (O_I) , $(O_I)^+$, (O_2)

The four packages match the eight build algorithms one-to-one only on the orientation lane. Their statement here repeats the definitions of Theorem ?? and is the standing data of the chapter.

1.2.0.1 (D₀) — Do-degeneration endpoint of the cosection-reduced d-critical orientation theory. Licensing tag δ . At every R , the data $D_{0,R}$ on the finite support $\widehat{\Gamma}_R$ is supplied: the finite Hall stage, the finite target test window Γ_R^{test} , successor transition maps whose composites give the smaller-stage maps, source/observable comparison maps, and a vanishing stage residual tuple $\mathfrak{D}_{D_{0,R}} = 0$; compatibly in R , the inverse limit gives the HN-completed state object and the finite first-order block $\mathfrak{D}_X$. The signed superdimension residual is

$$\text{sdim } P_{X,(n,l,m)}^{\Pi} = f(nm, l) \quad ((n, l, m) \in \Gamma_{\text{eff}}),$$

where $f(n, l)$ is the Fourier coefficient of the weak Jacobi form $\phi_{0,1}$ of weight zero and index one. The cosection contraction of the Pantev–Toen–Vaquie–Vezzosi (PTVV) (-1) -shifted symplectic form by the K_3 semiregularity cosection σ_{K_3} is the Maulik–Toda / Oberdieck reduced obstruction theory [?MaulikTodaGV, ?OberdieckReducedPT]; this absorbs the parity flip $\text{vd}^{\text{red}} = \text{vd}^{\text{std}} + 1$ into the reduced orientation line as the $\mathbb{Z}/2$ -grading shift $o_C^{\text{red}} = o_C^{\text{std}} \otimes \Pi$.

1.2.0.2 (O1) — strong multiplicative reduced orientation datum on the reduced $K_3 \times E$ -quotient self-Ext frame bundle. Licensing tag ε . A Joyce–Upmeyer-type orientation line $o_C \in \text{Pic}(\mathfrak{M}_X^{\text{or}})$ with $o_C^{\otimes 2} \simeq \det \text{RHom}_{\text{red}}(C, C)$, extension-multiplicative under $o_{C \oplus C'} \simeq o_C \otimes o_{C'}$, and equipped with the full quotient-orientation package

$$\theta_C = (\theta_C^{\text{red}}, \theta_C^{E, \text{free}}, \{(\theta_C^H, \lambda_C^H = 0)\}_{H \in \mathcal{H}(C)})$$

on every retained charge stratum. The vanishings $\alpha_C^{\text{red}} = 0$, $\alpha_C^{E, \text{free}} = 0$, $\widetilde{\beta}_C^H = 0$, $\lambda_C^H = 0$ are the orientation-line, the Borel-equivariant free-action, the residual finite-stabilizer Borel, and the residual finite-stabilizer linearization conditions; for $N = 2$, the Klein-four calculation $\beta_C^{E, E[2]} = b_{20}x_1^2 + b_{11}x_1x_2 + b_{02}x_2^2$ specializes the residual. The orientation lift technology is BBDJS [?BBDJS2015] on the unreduced side and Joyce–Upmeyer [?JoyceUpmeyer, ?JoyceTanakaUpmeyer] on the multiplicative side; the reduced $K_3 \times E$ -quotient theorem is supplied separately.

1.2.0.3 (O1)⁺ — Weyl-equivariant determinant-line transport along $W^{(2)}(\Lambda_{II}^{2,1})$ reflections. Licensing tag ε . The supplied lifts $\tau_i : o_C \rightarrow s_{\delta_i}^* o_C$, $i = 1, 2, 3$, of the three type-II wall transports satisfy $[c_0] = 0$ in $H^2(W^{(2)}(\Lambda_{II}^{2,1}); \mathbb{F}_2)$ (equivalently, in the finite action-groupoid cohomology $H^2(\mathcal{G}_R; \mathbb{F}_2)$ at every R), $\widetilde{\tau}_i^2 = 1$, and the quotient-orientation transport defects $\omega_{i,C} = 0$ for every simple reflection. Then $\widetilde{\tau}_i$ define an honest Weyl action on the reduced quotient orientation; the passage from the three generator signs to a character on $W^{(2)}(\Lambda_{II}^{2,1})$ is the pure group-theoretic Lemma ??.

1.2.0.4 (O2) — Pfaffian local model and wall atlas on type-II walls. Licensing tag ε . At a generic point of each $\mathbb{H}_{\delta_i}$, the supplied real local equation $u_i = u_{\delta_i}$ with $s_{\delta_i}(u_i) = -u_i$, the normal determinant-complex splitting

$$K_{\delta_i, R}^{\text{nor}} \simeq [\mathbb{R} \xrightarrow{u_i} \mathbb{R}], \quad \text{Pf}(K_{\delta_i, R}^{\text{nor}}) = \nu_{\delta_i, R} u_i, \quad s_{\delta_i}(\nu_{\delta_i, R}) = \nu_{\delta_i, R},$$

and the rank-one normal-form rank $N_{\delta_i}^{\text{Pf}} = 1$ hold. The wall charts cover every retained component and boundary stratum of $\mathbb{H}_{\delta_i}$; on overlaps the tangent/normal splittings, normal Pfaffian units, quotient orientations, and protected integration maps agree after a zero Čech sign class. The middle wall $\delta_2 \leftrightarrow (0, 1, 1)$ is represented by a retained wrapped wall object $x_{\delta_2, R} \in \mathcal{M}_{\eta, R}^{\text{wr, rig}}$ with $b_R^{\text{geom}}(\eta) > 0$, $\overline{\Pi}_X(\eta) = (0, 1, 1)$, and the component, boundary, overlap, quotient, and protected integration compatibility package of Proposition ??.

1.3 Surgical fix: the OP minus sign is a scalar branch convention

The Oberdieck–Pixton (OP) scalar $C_{\square, X} = -4096$ of Proposition ?? is the square of the theta-normalization 64 with the OP scalar branch sign $\text{sign}(C_{\square, X}) = -1$ absorbed into the fermionic shift $(-p_{\text{DT}})^n$ of the OP/DT chamber. It is not the source of the orientation character ϵ_o ; orientation lines are killed by squaring. The orientation character is computed by monodromy of the Pfaffian section around the type-II walls (Lemma ??); equivalently, by the $(-1)^{N_{\delta_i}^{\text{Pf}}}$ values from (O2).

2 THE PFAFFIAN-DIRAC THEOREM

The first theorem of the chapter identifies the protected Pfaffian of the Dirac-Igusa pro-object with the automorphic section. The four licensing tags $\alpha, \beta, \gamma, \delta, \varepsilon$ all attach.

THEOREM 2.1 (*Pfaffian-Dirac*). *Status: conditional, on (Do), (Or), (Or)⁺, (O2).* Granted the four obstruction packages of Section 1.2, the protected Pfaffian of the Dirac-Igusa pro-object $\mathfrak{D}_X^{\text{DI}}$ equals the automorphic section $D_X = \Delta_5$:

$$\iota_{\text{aut}}\left(\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}})\right) = D_X = \Delta_5$$

as sections of $\mathcal{L}^5 \otimes \nu_{\Delta_5}$ over the genus-two Siegel space $\mathbf{Sp}(2, \mathbb{Z}) \backslash \mathcal{H}_2$, where

$$\iota_{\text{aut}} : \mathcal{L}_{\text{Pf}, X} \xrightarrow{\sim} \mathcal{L}^5 \otimes \nu_{\Delta_5}$$

is the supplied Pfaffian-to-automorphic line comparison, compatible with the cusp trivialization at $\mathfrak{c}_{\infty}$ and with squaring to $\mathcal{L}^{10}$. Equivalently, in the type-II Borchers chamber,

$$\iota_{\text{aut}}\left(\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}})\right)\big|_{\mathfrak{c}_{\infty}} = 64 q^{1/2} r^{1/2} s^{1/2} \prod_{(n,l,m) \in \Gamma_{\text{eff}}} (1 - q^n r^l s^m)^{f(nm,l)}.$$

After squaring, the orientation forgets and the OP scalar branch identifies the protected scalar trace as

$$\text{Tr}_{\text{prot}}((-1)^F \mathfrak{D}_X^{-2}) = -4096 \Delta_5^{-2},$$

in agreement with [?0B], with $-4096 = \text{sign}(C_{\square, X}) \cdot 64^2$ the squared theta-normalization absorbing the OP scalar branch sign. *Licensing: α — chart of the boundary vacuum b in the open factorisation category $\mathcal{C}$ on $(X = K_3 \times E, D, \tau)$; β — protected Pfaffian functor, Hall-Drinfeld correspondence, Sp^{ch} specialisation; γ — pro-object ambient, chain-level after weight-completion; δ — Do-degeneration endpoint; ε — BBDJS orientation, Pfaffian-line non-degeneracy, type-II wall normal form.*

Proof. The proof is the inverse-limit assembly of a Pfaffian product, a leading coefficient identification with a known weight-five Siegel form, and a q -expansion principle on a punctured neighbourhood of the type-II cusp $\mathfrak{c}_{\infty}$.

Step 1 (finite-stage Pfaffian product, licensing $\delta + \varepsilon$). Hypothesis (Do) supplies, at every cofinal R , the finite first-order operator/algebra $\mathfrak{D}_{X,R}$ on the reduced compact quotient $\mathfrak{Q}_R = (\mathfrak{M}_R/E)^{\text{red}}$, the Pfaffian line $\mathcal{L}_{\text{Pf}, R}$, and the normalized leading coefficient condition $\mathfrak{o}_R^{\text{lead}} = \mathfrak{o}$. Hypothesis (Or) supplies the orientation line $\mathfrak{o}_C$ with $\mathfrak{o}_C^{\otimes 2} \simeq \det \text{RHom}_{\text{red}}(C, C)$ and extension-multiplicativity $\mathfrak{o}_{C \oplus C'} \simeq \mathfrak{o}_C \otimes \mathfrak{o}_{C'}$, descended under the reduced E -quotient. The protected Pfaffian functor of (β) is then the graded Pfaffian product of the determinant of RHom_{red} over the supplied retained charge strata. The signed-superdimension residual of (Do) gives

$$\text{sdim } P_{X, (n,l,m)}^{\Pi} = f(nm, l) \quad ((n, l, m) \in \Gamma_{\text{eff}}),$$

so the graded Pfaffian product, expanded in the type-II cusp polarization $(q, r, s) = (e^{2\pi i z_1}, e^{2\pi i z_2}, e^{2\pi i z_3})$, is

$$\text{Pf}_{\text{prot}}(\mathfrak{D}_{X,R}) = v_{\mathfrak{o}} \prod_{(n,l,m) \in \Gamma_{\text{eff}, R}} (1 - q^n r^l s^m)^{f(nm,l)}, \quad v_{\mathfrak{o}} = 64 q^{1/2} r^{1/2} s^{1/2},$$

where $\Gamma_{\text{eff},R}$ is the active support cut by HN height R . The Weyl monomial v_o is the theta normalization $[q^{1/2}r^{1/2}s^{1/2}]\Delta_5 = 64$ of [?GN, Theorem 4.1].

Step 2 (inverse limit, licensing γ). The pro-object ambient is the cofinal HN system; (Do) supplies strict transition maps with vanishing successor residuals $\mathfrak{o}_{R^+R}^{\text{atlas}} = \mathfrak{o}_{R^+R}^{\text{tr}} = \mathfrak{o}$; the finite Pfaffian sections $\text{pf}_{X,R}$ form a Mittag-Leffler system on the active support, and the inverse limit is the formal Borchers product

$$\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}})|_{\mathfrak{c}_\infty} = \lim_R \text{Pf}_{\text{prot}}(\mathfrak{D}_{X,R}) = 64 q^{1/2} r^{1/2} s^{1/2} \prod_{(n,l,m) \in \Gamma_{\text{eff}}} (1 - q^n r^l s^m)^{f(nm,l)}.$$

Class $\mathcal{M}$ completion (chain-level after weight-completion, in the sense of Vol I's class $\mathcal{M}$ global triangle) is exactly the ambient in which $\lim_R$ commutes with the graded Pfaffian product on the active support; this is the licensing γ clause.

Step 3 (comparison with the imported Borchers product, licensing β). By Proposition ??, the formal product $64 q^{1/2} r^{1/2} s^{1/2} \prod (1 - q^n r^l s^m)^{f(nm,l)}$ equals the Fourier expansion of the automorphic section $D_X = \Delta_5$ at $\mathfrak{c}_\infty$; this is the Borchers–Gritsenko–Nikulin theorem [?Borchers95, ?GN, ?GNII]. The supplied $\iota_{\text{aut}} : \mathcal{L}_{\text{Pf},X} \xrightarrow{\sim} \mathcal{L}^5 \otimes \nu_{\Delta_5}$ of (β) , compatible with cusp trivialization and squaring, transports the inverse-limit Pfaffian to a section of $\mathcal{L}^5 \otimes \nu_{\Delta_5}$.

Step 4 (q -expansion principle, licensing α). The chart α records the open factorisation category $\mathcal{C}$ on (X, D, τ) with boundary vacuum b : on the punctured neighbourhood of $\mathfrak{c}_\infty$ inside $\text{Sp}(2, \mathbb{Z}) \backslash \mathcal{H}_2$ the chart algebra $A_b = \text{End}_{\mathcal{C}}(b)$ supplies the local factorisation-algebra structure within which the Pfaffian section is holomorphic. Two holomorphic genus-two Siegel modular forms with the same Maass character ν_{Δ_5} and the same Fourier expansion at $\mathfrak{c}_\infty$ are equal: this is the q -expansion principle for holomorphic Siegel modular forms [?Igusa1962, ?Igusa1964II]. Both sides have weight 5 and Maass character ν_{Δ_5} : the right-hand side by the Borchers–Gritsenko–Nikulin theorem; the left-hand side because the orientation character of the supplied $\mathfrak{D}_X^{\text{DI}}$, under $(O_1) + (O_1)^+ + (O_2)$, is exactly ν_{Δ_5} on $W^{(2)}(\Lambda_{II}^{2,+})$ (this is Theorem 3.1). Hence $\iota_{\text{aut}}(\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}})) = \Delta_5$ as sections of $\mathcal{L}^5 \otimes \nu_{\Delta_5}$.

Step 5 (the squared scalar, licensing β). Squaring forgets orientation: the squared line $\mathcal{L}^5 \otimes \nu_{\Delta_5})^{\otimes 2} = \mathcal{L}^{10}$ is trivially characterized, and $\Delta_5^2 = \Delta_{10}$ is scalar. By Proposition ??, the OP scalar partition function is $Z_{\text{OP}}^X = -(\chi_{10}^{\text{OP}})^{-1} = -4096 \Delta_5^{-2}$, with $\chi_{10}^{\text{OP}} = D_5^2 = 4096^{-1} \Delta_5^2$ and $D_5 = 64^{-1} \Delta_5$. The scalar trace identification $\text{Tr}_{\text{prot}}((-1)^F \mathfrak{D}_X^{-2}) = -4096 \Delta_5^{-2}$ follows; the minus sign is the OP scalar branch convention, absorbed into $(-p_{\text{DT}})^n$, and is not a feature of the orientation character. $\square$

Remark 2.2 (Why (Do)+(O1)+(O1)⁺+(O2) and not less). Each of the four packages is essential. (Do) supplies the existence and the finite first-order block; without it there is no $\mathfrak{D}_{X,R}$ and no Pfaffian product to take. (O1) supplies the orientation line and the reduced E -quotient descent; without the line the Pfaffian section has no domain. (O1)⁺ supplies the honest Weyl action on the reduced quotient orientation; without it the type-II reflection signs are projective and the orientation character is projective, not a homomorphism. (O2) supplies the local Pfaffian normal form; without it the rank $N_{\partial_i}^{\text{Pf}}$ is undetermined and the local sign cannot be computed. Drop any package and one of the five proof steps fails.

3 THE ORIENTATION CHARACTER IDENTITY

The second theorem identifies the geometric orientation monodromy character ϵ_o of the Dirac–Igusa pro-object with the automorphic Maass character ν_{Δ_5} on the type-II Weyl group. The proof splits into a local Pfaffian wall computation, a group-theoretic extension, and an independent automorphic check.

THEOREM 3.1 (*Orientation character*). *Status: conditional, on (O_1) , $(O_1)^+$, (O_2) .* Granted the orientation packages (O_1) , $(O_1)^+$, (O_2) , the geometric orientation monodromy character of the Dirac–Igusa pro-object equals the automorphic Maass character on the type-II Weyl group:

$$\boxed{\epsilon_o = \nu_{\Delta_5} \quad \text{on} \quad W^{(2)}(\Lambda_{II}^{2,1})}.$$

The character takes the value -1 on each of the three type-II simple reflections $s_{\delta_1}, s_{\delta_2}, s_{\delta_3}$, and the identity is reduced to these three values by the abelianization $W^{(2)}(\Lambda_{II}^{2,1})_{\text{ab}} \simeq (\mathbb{Z}/2)^3$ (Lemma ??). *Licensing: ε — BBDJS orientation, Pfaffian-line non-degeneracy, type-II wall normal form.*

The identity is a comparison of two characters defined independently: ϵ_o is computed by transport of the Pfaffian section under the supplied Weyl action of $(O_1)^+$; ν_{Δ_5} is the Maass multiplier of the holomorphic Siegel form $D_X = \Delta_5$ computed from the multiplier system of the corresponding theta lift [?MAASS: 1, ?GN, ?Gritsenko99]. No scalar branch convention enters this sign comparison: replacing Δ_5 by $c\Delta_5$, $c \in \mathbb{C}^\times$, changes neither the Maass character nor the divisor orders. The factor 64 names the monic Borcherds product $D_5 = 64^{-1}\Delta_5$; the OP/DT scalar -4096 is unused.

Proof. Three lemmas combine: Lemma ?? computes the local Pfaffian signs on the three simple type-II walls; Lemma ?? extends a generator sign-tuple $(\epsilon_1, \epsilon_2, \epsilon_3) \in \{\pm 1\}^3$ to a unique character on $W^{(2)}(\Lambda_{II}^{2,1})$; and Lemma ?? computes the automorphic target values $\nu_{\Delta_5}(s_{\delta_i}) = -1$ directly from the Maass multiplier and the Borcherds divisor order $d_{\delta_i}^{\text{aut}} = 1$.

Step 1 (local Pfaffian signs from (O_2)). By (O_2) , at a generic point of each $\mathbb{H}_{\delta_i}$, $i = 1, 2, 3$, the supplied local equation u_i satisfies $s_{\delta_i}(u_i) = -u_i$ and the normal determinant-complex splits as $K_{\delta_i}^{\text{nor}} \simeq [\mathbb{R} \xrightarrow{u_i} \mathbb{R}]$ with $\text{Pf} = \nu_{\delta_i} u_i$ and $s_{\delta_i}(\nu_{\delta_i}) = \nu_{\delta_i}$. Transport along the closed path ℓ_{δ_i} (the path in the reduced quotient wall-complement carrying $u_i = \epsilon$ to $u_i = -\epsilon$, closed by the $(O_1)^+$ lift $\tilde{\tau}_i$) sends $\nu_{\delta_i} u_i$ to $-\nu_{\delta_i} u_i$, giving

$$\epsilon_o(s_{\delta_i}) = (-1)^{N_{\delta_i}^{\text{Pf}}} = -1, \quad i = 1, 2, 3.$$

This is Lemma ?. Squaring eliminates the sign and hence eliminates the OP scalar branch ambiguity: the squared Pfaffian $\text{Pf}(K_{\delta_i}^{\text{nor}})^2 = u_i^2$ is fixed by s_{δ_i} .

Step 2 (group-theoretic extension via $(O_1)^+$). The type-II Weyl group has Coxeter presentation

$$W^{(2)}(\Lambda_{II}^{2,1}) \simeq \langle s_{\delta_1}, s_{\delta_2}, s_{\delta_3} \mid s_{\delta_1}^2 = s_{\delta_2}^2 = s_{\delta_3}^2 = 1 \rangle,$$

with no braid relations — the off-diagonal entries $(\delta_i, \delta_j) = -2$ for $i \neq j$ give Coxeter exponents $m_{ij} = \infty$. Hence $W^{(2)}(\Lambda_{II}^{2,1})_{\text{ab}} \simeq (\mathbb{Z}/2)^3$ with one generator per s_{δ_i} , and any $\{\pm 1\}$ -valued assignment on the generators extends uniquely to a homomorphism. By $(O_1)^+$, $[c_o] = o$, $\tilde{\tau}_i^2 = 1$, and $\omega_{i,C} = o$; the lifts $\tilde{\tau}_i$ define an honest action on the reduced quotient orientation, and the induced character has $\epsilon_o(s_{\delta_i}) = -1$ for $i = 1, 2, 3$. This is Lemma ?.

Step 3 (independent automorphic computation of ν_{Δ_5}). The Maass multiplier ν_{Δ_5} of the singular theta lift of $\phi_{o,1}$ is computed from the multiplier system of the Borcherds product (Borcherds [?Borcherds95], Gritsenko–Nikulin [?GN, Section 3], [?Gritsenko99], Maass [?MAASS: 1]). On the type-II simple roots $\delta_1 = 2f_2 - f_3$, $\delta_2 = 2f_{-2} - f_3$, $\delta_3 = f_3$, the Borcherds divisor orders are

$$d_{\delta_i}^{\text{aut}} := \text{ord}_{\mathbb{H}_{\delta_i}}(\Delta_5) = f(o, \pm 1) = 1, \quad i = 1, 2, 3,$$

so $\nu_{\Delta_5}(s_{\delta_i}) = (-1)^{d_{\delta_i}^{\text{aut}}} = -1$ for $i = 1, 2, 3$. This is Lemma ???. The chamber-permuting roots $f_{-2} - f_2, f_2 - f_3, f_2 + f_3$ lie in $\Lambda_{II}^{2,1} \setminus \Lambda_{II}^{2,1}$ and are not type-II walls; their Maass values are $+1$ but no Hall orientation sign is asserted on them unless a semidirect-product Hall lift is supplied (Remark 3.4).

Step 4 (the comparison $\epsilon_o = \nu_{\Delta_5}$). Two $\{\pm 1\}$ -valued characters of $W^{(2)}(\Lambda_{II}^{2,1})$ that agree on $s_{\delta_1}, s_{\delta_2}, s_{\delta_3}$ agree everywhere (Step 2). By Steps 1 and 3, both ϵ_o and ν_{Δ_5} take the value -1 on each of the three simple reflections. Therefore $\epsilon_o = \nu_{\Delta_5}$ on $W^{(2)}(\Lambda_{II}^{2,1})$. $\square$

Remark 3.2 (The OP minus sign is not the source of ϵ_o). The two computations of Steps 1 and 3 are independent. Step 1 uses no automorphic data; Step 3 uses no Pfaffian local model. In particular, the OP scalar $C_{\square, X} = -4096 = \text{sign}(C_{\square, X}) \cdot 64^2$ does not source the orientation character: squaring forgets orientation, so $C_{\square, X}$ carries no information about ϵ_o . The minus sign $\text{sign}(C_{\square, X}) = -1$ is the OP scalar branch convention of [?OB], absorbed into the fermionic shift $(-p_{\text{DT}})^n$ of the OP/DT chamber. The orientation character is a *geometric* monodromy datum, computed by transport of the Pfaffian line around type-II walls; it is fixed by the data of $(O_1) + (O_1)^+ + (O_2)$, and is independently checked against the automorphic target by Step 3.

Remark 3.3 (Imaginary roots are not Weyl reflections). The Weyl group $W^{(2)}(\Lambda_{II}^{2,1})$ of the type-II BKM datum is generated by real simple reflections only. The imaginary simple roots contribute to the Weyl–Kac–Borcherds denominator through the multiplicity coefficients $m(a)$ and $\tau(a)$ of Gritsenko–Nikulin [?GNII, Theorem 2.1], [?GN, Section 3]; they do not define reflections in $W^{(2)}(\Lambda_{II}^{2,1})$. Hence $\epsilon_o : W^{(2)}(\Lambda_{II}^{2,1}) \rightarrow \{\pm 1\}$ is a real-root reflection datum. Local divisor monodromy of a compact Hall theory around imaginary components of the Borcherds divisor is not a Weyl character and is not asserted here.

Remark 3.4 (Extension to $W^{(2)}(\Lambda_{II}^{2,1}) \rtimes \text{Aut}(\mathcal{P}_{II})$). The chamber-permuting symmetry group $\text{Aut}(\mathcal{P}_{II}) \simeq S_3$ acts on the Coxeter generators $s_{\delta_1}, s_{\delta_2}, s_{\delta_3}$ by permutation; its three reflections $s_{f_{-2}-f_2}, s_{f_2-f_3}, s_{f_2+f_3}$ have $\nu_{\Delta_5} = +1$. An extension of ϵ_o to $W^{(2)}(\Lambda_{II}^{2,1}) \rtimes \text{Aut}(\mathcal{P}_{II})$ is not asserted in this chapter; it requires a separately supplied Hall semidirect-product lift of the chamber-permuting reflections.

4 THE PRIMITIVE RECOGNITION THEOREM

The third theorem identifies the source-side primitive Lie superalgebra of the Dirac–Igusa pro-object with the BKM denominator algebra $\mathfrak{g}_{\Delta_5}$. The conditions form the *recognition criterion*: ten finite-stage data carrying compact geometric provenance, none of which is inferable from the Borcherds product or signed determinant alone.

THEOREM 4.1 (Primitive recognition). *Status: conditional, on the recognition criterion (S1)–(S10) of Theorem 0.29; on the cofinal finite-window datum of Definition 0.30; on the chiral Koszul source datum of Definition 0.25; on $(O_1) + (O_1)^+ + (O_2)$ through the source orientation lane.* Suppose the compact $K_3 \times E$ Hall/factorization construction supplies, on a cofinal sequence of downward-saturated finite root windows $W_1 \subset W_2 \subset \cdots$ with $\bigcup_\nu W_\nu = \mathcal{R}_+$:

- (S2) *Cartan and root-degree matching.* An even Cartan identification $\iota_H : P_{X,o}^{\Pi, \text{red}} \xrightarrow{\sim} \Lambda_{II}^{2,1} \otimes \mathbb{C}$, with Gram-grading $\alpha(n, l, m) = 2nf_2 - lf_3 + 2mf_{-2}$.
- (S3) *Simple primitive representatives.* Parity-homogeneous e_i^X, f_i^X, h_i^X for every simple-root index $i \in I$ of $\mathcal{B}_{\Delta_5}$, with Gritsenko–Nikulin parity dimensions $\mu_{\bar{o}}(a), \mu_{\bar{\tau}}(a)$ on imaginary fibres.

- (S4) *Borcherds–Kac relations.* Cartan, Chevalley ($[e_i^X, f_j^X] = \delta_{ij} h_i^X$), real Serre, Borcherds orthogonality, super sign, and isotropic-string relations satisfied in the protected Hall bracket.
- (S5) *Completed Hopf radical equality.* The completed Hopf pairing has closed homogeneous radical Rad_X which is a Lie ideal and a coideal, and identifies under the radical quotient with Rad_{GN} .
- (S6) *No-extra-relations.* The presentation map $\pi : \widehat{\mathfrak{F}}(\mathcal{B}_{\Delta_s}) \rightarrow P_X^{\Pi, \text{red}}$ has $\ker \pi = \overline{J_{\text{BK}}} + \text{Rad}_{\text{GN}}$, on every finite window separately.
- (S7) *Generation.* $P_X^{\Pi, \text{red}}$ is topologically generated by $P_{X, \circ}^{\Pi, \text{red}}$ and $\{e_i^X, f_i^X\}_{i \in I}$.
- (S8) *Completed PBW.* The associated graded $\text{gr}_{\text{PBW}} U(P_X^{\Pi, \text{red}})_W \xrightarrow{\sim} \text{gr}_{\text{PBW}} U(\mathfrak{g}_{\Delta_s})_W$ is an isomorphism on every finite window, with strict transitions.
- (S9) *Exact triangular completion.* The HN inverse limit is exact on the saturated finite root windows: $\lim^1$ vanishes for the kernels, radicals, PBW associated gradeds, and root-space parity pieces.
- (S10) *Full parity dimensions.* Two integers per nonzero root degree: $\dim P_{X, \gamma, \bar{p}}^{\Pi, \text{red}} = \text{mult}_{\bar{p}}^{\text{GN}}(\alpha(\gamma))$, $p = 0, 1$; not only their signed difference.

Then there is a canonical isomorphism of completed Γ_{gram} -graded Lie superalgebras on the active root window

$$H^0 \text{Prim}_{\text{prot}}(\overline{\Pi}_{X, *}^{\ominus} \mathcal{F}_X) / \overline{\text{Rad}}\langle \cdot, \cdot \rangle_X \cong \mathfrak{g}_{\Delta_s}$$

and the Weyl–Kac–Borcherds denominator of $P_X^{\Pi, \text{red}}$ is $\text{den}(\mathfrak{g}_{\Delta_s}) = 64^{-1} \Delta_s(2Z)$. If in addition the chiral Koszul source datum of Definition 0.25 is supplied, including the Koszul Mittag–Leffler exactness hypothesis of Proposition 0.28, then the chiral cobar of the source coalgebra C_X is identified with the formal current envelope:

$$\Omega_E^{\text{ch}} C_X \simeq \text{Den}_{\Delta_s, E}.$$

Licensing: α — chart b of the open factorisation category; β — Hall–Drinfeld correspondence and Sp^{ch} specialisation; γ — pro-object ambient with completed PBW; δ — Do-degeneration endpoint as input to (S1) and (S3); ε — orientation lane through (S1) and the closure of Rad_X by the Frobenius compatibility of Lemma ??.

Proof. The argument is the inverse-limit assembly of (S1)–(S10) on each finite window, with the chiral cobar identification appended at the end. The proof of Theorem 0.29 supplies the window-wise version; we extract the load-bearing logic and add the chiral-cobar step.

Step 1 (presentation map and kernel, licensing $\alpha + \beta$). (S2) and (S3) define the continuous Lie-superalgebra homomorphism

$$\pi : \widehat{\mathfrak{F}}(\mathcal{B}_{\Delta_s}) \longrightarrow P_X^{\Pi, \text{red}}$$

on the descended Gram-graded completion. (S4) puts the Borcherds–Kac ideal J_{BK} in the kernel. (S6) gives the kernel equality $\ker \pi = \overline{J_{\text{BK}}} + \text{Rad}_{\text{GN}}$ on every finite window, hence injectivity of the descended map $G(\mathcal{B}_{\Delta_s}) \rightarrow P_X^{\Pi, \text{red}}$. (S7) gives surjectivity. Thus the descended map is an isomorphism on every finite window.

Step 2 (radical descent, licensing ε). (S5) identifies the source Hopf radical $\overline{\text{Rad}_X}$ with the target radical Rad_{GN} . The required Hopf adjointness, ideal/coideal property, and closedness on the inverse system are the finite Hopf-radical conditions of Lemma ??; quotient-tensor non-degeneracy is fixed by the Frobenius identity from the CY-3 Serre functor of Lemma ??, which is part of the orientation lane.

Step 3 (full parity dimensions, licensing δ). (S10) is the two-integer equality $\dim P_{X,\gamma,\bar{p}}^{\Pi,\text{red}} = \text{mult}_{\bar{p}}^{\text{GN}}(\alpha(\gamma))$ for $p = 0, 1$, not just the signed difference. This rules out hidden zero-superdimension source summands, wrong parity lifts invisible to the determinant, and parity-cancelling ideals. Crucially, a putative compact source which has all the same Hall bracket templates as $\mathfrak{g}_{\Delta_5}$ on a finite window can still fail (S5) if there is a kernel element $k \in K_{\gamma_\mu}$ with $(Q_{\gamma_\mu} \otimes Q_{\gamma_\nu})D_{\rho}^{\mu,\nu}(k) \neq 0$; the quotient target table sees no discrepancy, but the radical is not a coideal and the Hopf quotient is not defined (Corollary ??). This is the finite tensor-Hopf obstruction hidden by signed determinant data; (S5)+(S10) together block it.

Step 4 (PBW comparison and exact completion, licensing γ). (S8) is the associated-graded isomorphism on every finite window, with strict transitions. (S9) is the exact-completion clause: $\lim^1$ vanishes for kernels, radicals, PBW associated gradeds, and root-space parity pieces. Equivalently, no new generator, relation, radical element, or parity-cancelling summand appears only after completion. Class $\mathcal{M}$ completion is the ambient: chain-level identifications hold after weight-completion; this is the licensing γ clause.

Step 5 (passage to the inverse limit). Since the windows are cofinal ($\bigcup_\nu W_\nu = \mathcal{R}_+$) and the radical in (S5) is the closure of the finite kernels, the finite isomorphisms of Steps 1–4 assemble to the inverse-limit isomorphism

$$H^0 \text{Prim}_{\text{prot}}(\overline{\Pi}_{X,*}^{\ominus} \mathcal{F}_X) / \overline{\text{Rad}\langle \cdot, \cdot \rangle_X} = P_X^{\Pi,\text{red}} \xrightarrow{\sim} \mathfrak{g}_{\Delta_5}.$$

The Weyl–Kac–Borcherds denominator of $P_X^{\Pi,\text{red}}$ is therefore $\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1} \Delta_5(2Z)$, by Theorem ??.

Step 6 (chiral cobar identification). Suppose the chiral Koszul source datum of Definition 0.25 is supplied: $(\mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}}, C_{X,R}, F_{R,\bullet}^{\text{co}}, \Delta_R^{\text{ch}}, b_{X,R}, \Theta_{\text{Kos},R})$ with vanishing residual $\mathfrak{D}_{\text{Kos},R} = 0$ compatibly in R and with the Mittag–Leffler exactness hypothesis of Proposition 0.28. The target side is the formal current envelope $\text{Den}_{\Delta_5,E} = U_E^{\text{ch}}(\text{Cur}_E(\mathcal{A}_{\text{den}}))$ of Proposition ??. The cobar comparison of Proposition 0.28 identifies

$$\Omega_E^{\text{ch}} C_X \simeq \text{Den}_{\Delta_5,E},$$

through the source-internal bar/cobar admissibility (the residual $\mathfrak{o}_R^{\varepsilon,\text{src}} = 0$) and the source-to-target quasi-isomorphism ($\mathfrak{o}_R^{\delta} = 0$), avoiding the target-internal bar/cobar counit; this is the firewall (β) provides between observables and states. $\square$

Remark 4.2 (The zero-bracket graded super VS counterexample). The criterion (S1)–(S10) blocks the trivial counterexample of a graded super vector space $V_\bullet$ with all brackets zero and parity dimensions matching $\text{mult}_{\bar{p}}^{\text{GN}}$ on every window. Such a $V_\bullet$ satisfies (S2), (S3), (S7), (S8), (S10), and trivially (S4) (because all brackets are zero, so all relations hold); it satisfies the signed-superdimension residual on the active support. It fails (S6): the kernel of the presentation map $\pi : \widehat{\mathfrak{F}}(\mathcal{B}_{\Delta_5}) \rightarrow V_\bullet$ is not $\overline{J_{\text{BK}} + \text{Rad}_{\text{GN}}}$ but contains every commutator $[e_i, e_j]$ with $(\alpha_i, \alpha_j) \neq 0$, so the induced map $G(\mathcal{B}_{\Delta_5}) \rightarrow V_\bullet$ is not injective. It fails (S5): the Hopf pairing on $V_\bullet$ is the zero pairing (or any pairing inherited from the target), and the radical is all of $V_\bullet$, so the radical quotient is zero and not the BKM target. It fails (S4) on the precise statement $[e_i^X, f_j^X] = \delta_{ij} h_i^X$ of the Chevalley relation: with all brackets zero and $h_i^X \neq 0$, this fails on the diagonal $i = j$. The counterexample shows that signed-superdimension matching alone is not recognition; the bracket-level data of (S4)+(S5)+(S6) is the load-bearing content.

Remark 4.3 (Recognition without (D0)+(O1)+(O1)⁺+(O2)). The recognition criterion of Theorem 4.1 nominally lists no orientation hypothesis among (S1)–(S10). The orientation packages enter through

(S1) — the chain-level descent $\overline{\Pi}_{X,*}^{\ominus} \mathcal{F}_X$ is conditional on the orientation lane via Definition ?? and the Frobenius identity from the CY-3 Serre functor — and through (S5), where the Hopf radical's coideal property is fixed by the same Frobenius identity (Lemma ??). In the chapter's combined statement (Theorem 2.1+Theorem 3.1+Theorem 4.1) the orientation packages are loaded once and serve all three theorems.

5 CROSS-VOLUME CONSISTENCY

The three theorems sit inside the chiral-bar-cobar / Calabi–Yau-quantum-groups constellation as the level-4/level-5 face of the chain fusion of the Calabi–Yau-to-chiral programme. Three consistencies are load-bearing.

5.1 Vol II Construction Problem 1

Vol II's Construction Problem 1 is exactly the cross-level identity $\textcircled{2} \leftrightarrow \textcircled{3}$ applied at level 4/level 5 of the chain. The Vol II primitive package on (X, D, τ) is the bicoloured nine-tuple

$$(C|_{(X,D,\tau)}, \mathcal{D}^{\text{cl}}|_X; b, A_b, F^{\text{cl}}; \sigma_{\text{half}}, \text{tr}_X^{\text{cl}}; \text{Tr}_C).$$

The chart of vacuum b is the licensing α input; the chart algebra $A_b = \text{End}_C(b)$ is the open primitive algebra; the closed-colour vacuum factorisation algebra F^{cl} gives the closed-side trace tr_X^{cl} which evaluates at the type-II cusp ϵ_{∞} to the level-5 Borcherds–Gritsenko–Nikulin section $D_X = \Delta_5$ (View $\textcircled{1}$). The identification $A_b = \Phi_d^{(\Sigma_{d-1}, C)}(C_X)$ is the chain fusion of the chiral-bar-cobar / Calabi–Yau-quantum-groups programme at level 2: the Calabi–Yau-side's Stage-2 chiral algebra A_X is the open-side's chart algebra A_b on the same curve C with boundary vacuum induced by (C_X, Σ_{d-1}, C) . In our setting, $A_X = A_b$ is the chiral algebra on E attached to the Hall source of $K_3 \times E$; the chiral cobar $\Omega_E^{\text{ch}} C_X \simeq \text{Den}_{\Delta_5, E}$ of Theorem 4.1 is the source-side chiral expression at level 2, and its constant-current Lie superalgebra is $\mathfrak{g}_{\Delta_5}$ (View $\textcircled{2}$). The protected Pfaffian $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}})$ is the level-5 scalar shadow of the open-cyclic trace; this is the content of the Universal Holography master theorem of Vol II applied at $(X = K_3 \times E, \Delta = \Delta_5)$.

5.2 Vol III chain fusion at level 2

The Vol III κ -stratification supplies the κ -tuple $(\kappa_{\text{cat}}, \kappa_{\text{ch}}^{\text{Hodge}}, \kappa_{\text{ch}}^{\text{Heis}}, \kappa_{\text{BKM}}, \kappa_{\text{fiber}}) = (0, 0, 3, 5, 24)$ at $K_3 \times E$; the row $\kappa_{\text{BKM}} = 5$ is the cusp-form weight realised by Δ_5 , and the row $\kappa_{\text{fiber}} = 24$ is $\chi_{\text{top}}(K_3) = 24$. The additive collapse $\kappa_{\text{BKM}} = \kappa_{\text{ch}} + \chi(\mathcal{O}_{\text{fiber}})$ fails at $N = 1$ (Vol III counterexample), confirming that κ_{fiber} is the Euler characteristic of the topological fibre and not of $\mathcal{O}_{\text{fiber}}$; for K_3 , $\chi_{\text{top}}(K_3) = 24$ but $\chi(\mathcal{O}_{K_3}) = 2$. The Vol III chain fusion at level 2 identifies the chiral algebra A_X supplied by Vol III's CY-input face with the chart algebra A_b supplied by Vol II's open-input face; in our setting, $A_X = A_b = \Omega_E^{\text{ch}} C_X \simeq \text{Den}_{\Delta_5, E}$ by Theorem 4.1.

Remark 5.1 (Coefficient projection is not a Hall comparison). A Hall–Borcherds comparison map cannot be defined by coefficient projection alone, and a coefficient projection respecting charge addition because both products use the same cone is not a recognition theorem. A coefficient projection is not a Hall product/bracket matrix comparison, and does not prove Borcherds–Serre rows, radical descent, no-extra-relations, or PBW. The recognition criterion (S4)–(S10) of Theorem 4.1 is the corrected statement: the source-side bracket matrices, Hopf radical, kernel equality, and PBW associated-graded comparison are independent data, none of which is inferable from a coefficient projection.

5.3 BCOV/mixed holomorphic-topological strings

The mixed holomorphic-topological strings companion provides the local model $\mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$ with brane completion as one realisation of the Stage-1 datum Φ_d^{FA} ; the global obstruction is the holomorphic de Rham obstruction class. The level-5 scalar shadow $Z_{\text{BPS}}^{K_3 \times E} = \Delta_5^{-2}$ on the closed $K_3 \times E \mapsto$ point cobordism is the BPS partition function (Section 9 of the manuscript); promotion to a 3d gravitational path integral on $K_3 \times E$ requires the Hall–Borcherds residual of Vol II and is not claimed by Theorem 2.1. The identity $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}}) = \Delta_5$ is a level-4/level-5 statement; its squared scalar trace is level 5; the operator-level package $\mathfrak{D}_X^{\text{DI}}$ is itself level 4 as a pro-object, conditional on (D_0) , (O_1) , $(O_1)^+$, (O_2) .

6 RESIDUAL OBSTRUCTIONS AND OPEN SUBROUTES

The three theorems of this chapter are conditional. Five obstructions remain, each isolated to a precise open subroute.

6.0.0.1 (D_0) . The cosection-twisted reduced d-critical orientation theory is constructed locally on rank-one strata and partially on Klein-four small-orbit boundary (Lemma ??); the connected BE -linearization is supplied conditional on $\mathfrak{D}_R^{\text{OI}+, \text{conn}} = 0$; the inverse-limit exhaustion of the active support is partial on the height-7 closure $W_{\leq 7}$ and open on the full positive cone $\mathcal{R}_+$. The sub-residuals $\mathfrak{D}_{D_0, R}$ of the D_0 -state, D_0 -quotient, D_0 -observable, D_0 -descent, and D_0 -Pfaffian rows of Definition ?? are tracked stage by stage in the cofinal HN inverse system.

6.0.0.2 (O_1) . Rank-one components are oriented: Joyce–Upmeyer multiplicative orientation [$?_{\text{JoyceUpmeyer}}$, $?_{\text{JoyceTanakaUpmeyer}}$] on the open factorisation correspondences and BBDJS [$?_{\text{BBDJS2015}}$] on the unreduced d-critical line, with the rank-one Pin^c -lift gerbe vanishing on rank-one strata. HN-reducible Pin^c gerbe is settled on length- $\ell \geq 2$ HN strata whose graded factors lie in the rank-one sector, by Lemma ??; stable higher-rank length-one HN strata remain open. The finite-stabilizer linearization characters $\lambda_C^H = 0$ are settled for $N = 2$ Klein-four by Lemma ?? and partially for $a \geq 2$ two-primary residuals; the connected BE -linearization holds for translation-equivariant reduced orientation lines.

6.0.0.3 $(O_1)^+$. The projective cocycle $[c_o] \in H^2(W^{(2)}(\Lambda_{II}^{2,1}); \mathbb{F}_2)$ vanishes on the finite action-groupoid level for the three rank-one type-II crossings (Proposition ??). The torsor defects $\omega_{i,C} = 0$ are settled for the rank-one sector and open for higher-rank components. The honest Weyl action on the reduced quotient orientation extends from the local model to the cofinal HN system through Corollary ??; cofinal extension to the full $\mathcal{R}_+$ is open.

6.0.0.4 (O_2) . The rank-one normal-form rank $N_{\delta_i}^{\text{Pf}} = 1$ holds for $i = 1, 2, 3$ (Lemma ??). The type-II half-Hilbert orbit sign sum $4096 = 2^{12}$ is the squared theta-normalization with the 2^{12} sign assignments arising as $(\pm 1)^{\dim_{\text{Spin}}}$ over the 12-dimensional half-Hilbert orbit; the OP scalar branch sign -1 is absorbed into $(-p_{\text{DT}})^n$ of the OP/DT chamber, not into the orientation line. The middle wall $\delta_2 \leftrightarrow (0, 1, 1)$ is represented by a wrapped wall object with positive geometric b -degree; the component, boundary, overlap, quotient, and protected integration compatibility package (Proposition ??) is discharged on the rank-one wall atlas and open on the higher-rank wall atlas.

6.0.0.5 Recognition criterion (S1)–(S10). The window $W_{\leq 3}$ is the first arithmetic window but is not relation-closed. The first relation-closed window is the height-7 closure

$$W_{\text{rel}}^+ = W_{\leq 3}^{\text{act}} \cup R_4 \cup I_4 \cup C_{\leq 7} \cup \{2\delta_{123}\},$$

adding the real-Serre terminal rows, doubled isotropic rows, δ_{123} -real strings, odd–odd δ_{123} rows, and complementary height-seven strings. At $W_{\leq 3}$ the target parity table $\delta_i : 1|0, a_{ij} : 10|0, 2\delta_i + \delta_j : 1|0, \delta_{123} : 29|93$ is computed; a compact source implementing (S3)–(S10) on $W_{\leq 3}$ is open, and the source identification at W_{rel}^+ is the next milestone.

6.0.0.6 Vol III Hall–Drinfeld–Borcherds spectral sequence on compact non-toric CY₃. The Hall–Drinfeld–Borcherds spectral sequence abutting from Vol III’s CoHA-type input to the BKM denominator on $K_3 \times E$ is open on the compact non-toric Calabi–Yau threefold, even after (Do), (O1), (O1)⁺, (O2) are supplied; this is the next frontier and is recorded in Section 10 of the manuscript. The finite-stage Hall source kernel of Theorem ?? is the first-order replacement.

6.0.0.7 Honest summary. The three theorems of this chapter close the cross-level identity $\textcircled{2} \leftrightarrow \textcircled{3}$ under the four orientation packages and the recognition criterion. The cross-level identities $\textcircled{1} \leftrightarrow \textcircled{2}$ and $\textcircled{2} \leftrightarrow \textcircled{5}$ are imported from Borcherds–Gritsenko–Nikulin and Howe. The operator-level package $\mathfrak{D}_X^{\text{DI}}$ is conditional as a pro-object; the scalar shadow $Z_{\text{BPS}}^{K_3 \times E} = \Delta_5^{-2}$ is the level-5 protected trace. The compact BPS Hilbert space interpretation, the 3d gravitational path integral, and the global Calabi–Yau realisation are level 3, level 2, and level 1 respectively, and remain open frontier targets, recorded honestly and not closed by relabelling.

Part IV

Shadowed

THE MIRROR DISCRIMINANT

1 THE MIRROR DISCRIMINANT

The four views developed so far place Δ_5 on chains that begin with explicit modular or representation-theoretic data: the Borchers–Gritsenko–Nikulin section of the weight-5 automorphic line over $\mathbf{Sp}_4(\mathbb{Z}) \backslash \mathcal{H}_2$ (View ①), the denominator $\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1} \Delta_5(2Z)$ of the Borchers–Kac–Moody Lie superalgebra on $\Lambda_{II}^{2,1}$ (View ②), the protected Pfaffian $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}})$ of the Dirac–Igusa pro-object (View ③), and the singular theta lift of the weak Jacobi form $\phi_{0,1}$ under $\text{Mp}(4, \mathbb{R}) \leftrightarrow \text{O}(3, 2)$ (View ⑤). The fifth view is geometric. Let $\mathcal{P}_{K_3}$ be the period family of K_3 -mirror pairs; let $\mathfrak{D}_{\text{per}} \subset \mathcal{P}_{K_3}$ be its discriminant locus, the divisor along which the local Hodge filtration degenerates. The expected statement is the equality

$$[\mathfrak{D}_{\text{per}}] = [\Delta_5^{-2}] \quad (\text{equality of divisor classes on the } (K_3 \times E)\text{-component}), \quad (8.1)$$

where $[\Delta_5^{-2}]$ is the principal divisor of the meromorphic section Δ_5^{-2} of the appropriate automorphic line. The four preceding chains supply the Borchers product, the BKM denominator, the Dirac–Igusa pro-object, and the theta-lift bridge; the fifth view asks for an independent geometric origin of the same form on the period side.

The status, recorded in advance, is conjectural. View ④ is the sole view in this manuscript whose load-bearing identification is not currently a theorem. The cross-level identities $④ \leftrightarrow ①$, $④ \leftrightarrow ②$, $④ \leftrightarrow ⑤$ are open. The identifications proved or conditionally proved on the other four chains are imported from the previous chapters unchanged; the present chapter neither weakens nor strengthens those chains.

The handle Δ_5 splits into three names already in this manuscript. The automorphic section over $\mathbf{Sp}_4(\mathbb{Z}) \backslash \mathcal{H}_2$ is denoted $\mathcal{D}_X = \Delta_5$ (View ①); the BKM denominator is $\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1} \Delta_5(2Z)$ (View ②); the protected Pfaffian is $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}})$ (View ③). The mirror conjecture concerns the first: the divisor identification (8.1) is a statement about $\mathcal{D}_X$ as an automorphic section, transported to the period side.

1.1 The K_3 mirror-period family

The lattice-polarised K_3 mirror is fixed by Dolgachev [?DolgachevMirrorLP] and tracked, throughout, through the Mukai lattice. Let

$$\Lambda_{K_3} = {}_3U \oplus {}_2E_8(-1), \quad \widetilde{\Lambda}_{K_3} = U \oplus \Lambda_{K_3} \simeq {}_4U \oplus {}_2E_8(-1),$$

the K_3 lattice of signature $(3, 19)$ and the Mukai lattice of signature $(4, 20)$ (Mukai [?Mukai1984, ?Mukai1987]). A *lattice polarisation* of a complex K_3 surface S is a primitive embedding $\iota: M \hookrightarrow \text{Pic}(S)$ of an even hyperbolic lattice M , of signature $(1, t)$, such that $M^\perp \subset \Lambda_{K_3}$ admits a primitive embedding of the hyperbolic plane U ; the pair (S, ι) is then *Nikulin-admissible*, in the convention of Gritsenko–Nikulin [?GNII] § 3 and the rank-3 sublattice constructions of Borchers [?Borchers95] § 15.

Here t denotes the rank of $M^\perp$ inside Λ_{K_3} minus one, and $(1, t)$ is signature in the convention $\text{sign}(M) = (\#\text{positive}, \#\text{negative})$.

The associated period domain

$$\Omega_M = \{[\omega] \in \mathbb{P}(M^\perp \otimes \mathbb{C}) : \langle \omega, \omega \rangle = 0, \langle \omega, \bar{\omega} \rangle > 0\}^+$$

is the type-IV symmetric domain of signature $(2, 19 - t)$, with arithmetic group $\Gamma_M = \text{O}^+(M^\perp)$ acting properly discontinuously. The *Dolgachev mirror lattice* is the sub-lattice

$$\check{M} = M^\perp \cap (U)^\perp \subset \Lambda_{K_3}, \quad M^\perp = U \oplus \check{M},$$

so that $\check{M}$ carries signature $(1, 19 - t)$ and is hyperbolic. The mirror K_3 surface $S^\vee$ is the lattice-polarised K_3 with $\text{Pic}(S^\vee) \simeq \check{M}$; the Dolgachev map $(S, \iota) \mapsto (S^\vee, \check{\iota})$ descends to an involution on the moduli stack Ω_M/Γ_M and on its mirror $\Omega_{\check{M}}/\Gamma_{\check{M}}$; this is the Dolgachev–Nikulin involution, recorded as ι_{DN} in the cross-volume $d = 2$ Vol III chapter on derived categories [CYQGVolIII, Prop. 6.4].

The Aspinwall–Morrison form of the same construction [AspinwallMorrisonStringy] realises Ω_M as the moduli space of lattice-polarised K_3 sigma models with B-field, and identifies the mirror involution as the exchange of large-volume and large-complex-structure limits along the two distinguished cusps of Ω_M/Γ_M . Aspinwall–Morrison and Dolgachev produce the same lattice; the difference is in presentation, not in object. The Mukai-lattice form

$$\widetilde{M} = U \oplus M, \quad \widetilde{M}^\perp = U \oplus M^\perp = 2U \oplus \check{M}, \quad \widetilde{M}^\perp \subset \widetilde{\Lambda}_{K_3},$$

is the convention adopted in the cross-volume Vol III statement [CYQGVolIII, Prop. 6.4 + Thm. 6.7]. The Dolgachev–Nikulin involution acts as $-\text{id}$ on $\widetilde{M}$ and $+\text{id}$ on $\widetilde{M}^\perp$, preserving the full signature $(4, 20)$.

1.1.0.1 The rank-three case. The polarisation relevant to View ④ is

$$M = U \oplus \langle -2 \rangle, \quad \text{rank}(M) = 3, \quad \text{sign}(M) = (1, 2),$$

with Dolgachev mirror lattice

$$\check{M} = (M^\perp \cap U^\perp)_{\Lambda_{K_3}} \simeq U \oplus E_7(-1) \oplus E_8(-1),$$

of rank 17 and signature $(1, 16)$ inside Λ_{K_3} ; the convention is $\text{sign} = (\#\text{positive}, \#\text{negative})$ on the bilinear form of the lattice. The decomposition follows from the embedding $\langle -2 \rangle \hookrightarrow E_8(-1)$ on a single root, after which the orthogonal complement of the root inside $E_8(-1)$ is $E_7(-1)$. Inside the Mukai lattice the augmented sub-lattice is

$$\widetilde{M} = U \oplus M = 2U \oplus \langle -2 \rangle, \quad \text{rank}(\widetilde{M}) = 5, \quad \text{sign}(\widetilde{M}) = (2, 3).$$

Under the manuscript’s convention (the exterior-square model of $\mathbf{Sp}_4 \simeq \text{Spin}(3, 2)$ at §??) the same lattice is written $\Lambda^{3,2} = U \oplus U \oplus \langle 2 \rangle$, with the hyperbolic-plane summand $\Lambda^{(1,1)} = \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}$ carrying signature $(1, 1)$ and the rank-1 summand carrying form value $+2$ on a generator. The two writings are the same lattice expressed in opposite sign conventions on the bilinear form: the manuscript’s $(p, q) = (3, 2)$ is signature “three positive, two negative” when read with the manuscript’s quadratic form, while the

Dolgachev convention writes the same signature “two positive, three negative” on the form with reversed sign. The $(+2)$ -vectors of the manuscript are the (-2) -vectors of the Dolgachev convention; the Heegner divisors and the period domain are independent of the sign choice.

The period domain attached to $\widetilde{M} = \Lambda^{3,2}$ is the type-IV symmetric domain

$$\Omega_{\widetilde{M}} = \{[Z] \in \mathbb{P}(\widetilde{M} \otimes \mathbb{C}) : (Z, Z) = 0, (Z, \bar{Z}) > 0\}^+, \quad \dim_{\mathbb{C}} \Omega_{\widetilde{M}} = 3,$$

with arithmetic group $\Gamma_{\widetilde{M}} = \mathrm{O}^+(\widetilde{M}) \simeq \mathrm{O}^+(\Lambda^{3,2})$ acting properly discontinuously, where the component is fixed by the manuscript’s positive cone $C(\Lambda^{2,1})_+$ (§ ??). The exceptional isogeny

$$\mathbf{PSp}_4(\mathbb{R}) \simeq \mathbf{SO}(3, 2)_+, \quad \mathbf{Sp}_4(\mathbb{Z})/\{\pm 1\} \simeq \mathbf{SO}_+(\Lambda^{3,2}),$$

imported as View ⑤ (the $\mathrm{Sp}(4) \simeq \mathrm{Spin}(3, 2)$ bridge), gives the identification

$$\Omega_{\widetilde{M}}/\Gamma_{\widetilde{M}} \simeq \mathcal{H}_2/\mathbf{Sp}_4(\mathbb{Z}). \quad (8.2)$$

The form $\Delta_5 \in M_5(\mathbf{Sp}_4(\mathbb{Z}), \nu_{\Delta_5})$ of View ①, viewed as a section on the right-hand side of (8.2), is the Borchers lift of the weak Jacobi form $\phi_{0,1}$ under the data $(\widetilde{M}, \phi_{0,1})$ on the left-hand side (Borchers [?B, Thm. 13.3] applied to signature $(2, 3)$; Gritsenko–Nikulin [?GNII, Thm. 1.1] for the identification with the additive Gritsenko lift; the rank-3 row of the Vol III lattice-polarised $\mathfrak{g}_L$ catalogue at $L = U \oplus \langle -2 \rangle$).

The K_3 -side period domain on the rank-3 polarisation is a different, larger object:

$$\Omega_M = \{[\omega] \in \mathbb{P}(M^\perp \otimes \mathbb{C}) : \langle \omega, \omega \rangle = 0, \langle \omega, \bar{\omega} \rangle > 0\}^+, \quad \dim_{\mathbb{C}} \Omega_M = 17,$$

where $M^\perp \subset \Lambda_{K_3}$ has signature $(2, 17)$ and parametrises the *transcendental* Hodge filtration of a lattice-polarised K_3 surface. The Borchers lift used in View ④ is on $\Omega_{\widetilde{M}}$, not on Ω_M : the difference is the absorption of the elliptic factor $\Lambda_E = U$ into $\widetilde{M} = U \oplus M$, which is the Künneth identification on $K_3 \times E$ recorded below in § 1.2. The two period spaces are not interchangeable: $\Omega_{\widetilde{M}}$ of dimension 3 is the genus-two Siegel space and supports Δ_5 directly via the exceptional isogeny, while Ω_M of dimension 17 is the K_3 -only period space, on which the rank-3 lattice-polarised lift of $\phi_{0,1}$ lands only after the additional Mukai- U augmentation.

1.2 The $K_3 \times E$ period family

The threefold $K_3 \times E$ acquires its Hodge structure through the Künneth decomposition. The Mukai lattice $\Lambda_S = H^*(S, \mathbb{Z})$ of signature $(4, 20)$ carries the Mukai pairing $\langle (r, c, s), (r', c', s') \rangle_{\mathrm{Muk}} = c \cdot c' - rs' - r's$ (Definition ??); the elliptic factor contributes $\Lambda_E = H^*(E, \mathbb{Z}) = U$ with the standard hyperbolic form. The total integral cohomology is

$$H^*(K_3 \times E, \mathbb{Z}) = \Lambda_S \otimes \Lambda_E = \Lambda_S \otimes U,$$

a lattice of rank 48; the relevant Hodge filtration is controlled, on each weight, by the K_3 polarisation M and the τ -modulus of E . The component of the period stack parametrisng $(K_3 \times E)$ -pairs (S, E_τ) with S lattice-polarised by $M = U \oplus \langle -2 \rangle$ is, on the rank-3 Mukai-invariant stratum,

$$\mathcal{P}_{K_3 \times E} := \Omega_{\widetilde{M}}/\Gamma_{\widetilde{M}} \stackrel{(8.2)}{\simeq} \mathcal{H}_2/\mathbf{Sp}_4(\mathbb{Z}), \quad (8.3)$$

where the augmented lattice $\widetilde{M} = U \oplus M$ absorbs the elliptic Mukai- U summand into the K_3 -side Mukai- U summand. The Künneth identification $\Lambda_E = U \subset \widetilde{M}$ is the Hodge-theoretic content of

the absorption: the elliptic period $\tau \in \mathbb{H}_1$ is absorbed into the type-IV variable $z_1 \in \Omega_{\widetilde{M}}$ under the manuscript's coordinatisation in §?? . Equation (8.3) is the period-side input on which View ④ is stated. It is a theorem of Gritsenko–Nikulin [?GNII, Thm. 1.1, Thm. 1.2] (the existence and identification of the lattice-polarised lift) and Borcherds [?Borcherds95, § 15] (the period-domain Borcherds-product convergence on signature $(2, 3)$); it is not a new statement.

The K_3 -only period domain Ω_M of dimension 17 does not enter (8.3): its role is to parametrise the transcendental Hodge filtration of S alone, before tensoring with $\Lambda_E = U$. The $(K_3 \times E)$ -component is on $\Omega_{\widetilde{M}}$, of dimension 3. The relation $\widetilde{M} = U \oplus M$ is the bridge between the two period spaces.

1.3 The discriminant locus

The period family $\mathcal{P}_{K_3 \times E}$ carries a natural divisor: the locus along which the rank-3 lattice-polarised K_3 surface acquires a nodal degeneration in the polarisation lattice $\widetilde{M}$. On the type-IV symmetric domain $\Omega_{\widetilde{M}}$, the Heegner divisors are

$$\mathfrak{H}_{(\delta)} = \{[Z] \in \Omega_{\widetilde{M}} : (Z, \delta) = 0\}, \quad \delta \in \widetilde{M}, \quad (\delta, \delta) = 2,$$

indexed by primitive square-2 vectors of the manuscript's form on $\widetilde{M} = \Lambda^{3,2}$ (equivalently, square- (-2) vectors of the Dolgachev form on the K_3 side after the sign flip recorded above). On the corresponding geometric component the K_3 fibre acquires a nodal rational curve in the class δ : the Hodge filtration on $H^2(S, \mathbb{Z}) \otimes H^*(E, \mathbb{Z})$ degenerates along this divisor as the elliptic factor passes through a Mukai-isomorphic point. The *period-family discriminant* is

$$\mathfrak{D}_{\text{per}} = \bigcup_{\substack{\delta \in \widetilde{M} \\ (\delta, \delta) = 2}} \mathfrak{H}_{(\delta)} / \Gamma_{\widetilde{M}} \subset \Omega_{\widetilde{M}} / \Gamma_{\widetilde{M}}. \quad (8.4)$$

The $\Gamma_{\widetilde{M}}$ -orbit structure on primitive square-2 vectors of $\widetilde{M}$ breaks into type-I and type-II classes ($\delta \cdot \widetilde{M} = \mathbb{Z}$ versus $\delta \cdot \widetilde{M} = 2\mathbb{Z}$, the dichotomy already imported as the type-II versus type-I distinction at the $W^{(2)}(\Lambda_{II}^{2,1})$ -orbit level in the manuscript's preceding chapters); the union in (8.4) runs over the type-II orbit, which is the orbit relevant to the $\mathbf{Sp}_4(\mathbb{Z})$ -image of the exceptional isogeny. The image in the quotient is one irreducible Heegner-type divisor (Bruinier–Funke [?BruinierFunkeHeegner, § 2] for the general lattice-Heegner formulation in signature $(2, 3)$; the index-1 case is implicit in Borcherds [?B, Thm. 10.1]). The same divisor, transported across the bridge (8.2), is the *Humbert surface*

$$\mathcal{H}_2^{(\text{Hum})} = \{[Z] \in \mathcal{H}_2 / \mathbf{Sp}_4(\mathbb{Z}) : Z = \begin{pmatrix} \tau_1 & 0 \\ 0 & \tau_2 \end{pmatrix} \text{ up to } \mathbf{Sp}_4(\mathbb{Z})\},$$

the diagonal locus on which the principally polarised abelian surface $\mathbb{C}^2 / (\mathbb{Z}^2 + Z\mathbb{Z}^2)$ decomposes as a product of two elliptic curves; the corresponding K_3 -side degeneration is exactly the nodal rational curve. The match $\mathfrak{D}_{\text{per}} \leftrightarrow \mathcal{H}_2^{(\text{Hum})}$ is fixed by the bridge (8.2).

The form Δ_s of View ① vanishes precisely along this divisor: $\Delta_s \in M_s(\mathbf{Sp}_4(\mathbb{Z}), \nu_{\Delta_s})$ has a simple zero on $\mathcal{H}_2^{(\text{Hum})}$ [?GNPartI, Thm. 3.1][?GNII, § 3], with multiplier ν_{Δ_s} accounting for the half-integral shift. Concretely, the Borcherds product expansion of View ① gives

$$\Delta_s(Z) = 64 q^{1/2} r^{1/2} s^{1/2} \prod_{(n,l,m) \in \Gamma_{\text{eff}}} (1 - q^n r^l s^m)^{f(nm,l)},$$

and the zero locus is exactly $\mathcal{H}_2^{(\text{Hum})}$, counted with multiplicity 1. Hence

$$[\Delta_5^{-1}] = [\mathcal{H}_2^{(\text{Hum})}] = [\mathfrak{D}_{\text{per}}] \quad (\text{divisor classes on } \mathcal{H}_2/\mathbf{Sp}_4(\mathbb{Z})). \quad (8.5)$$

This identity is a theorem in the cited references; it is recorded here as the input to the conjecture.

1.4 The conjecture

The discriminant of a period family carries multiplicities, not just a support. For the K_3 -mirror period family on the rank-3 Mukai-invariant component, the relevant multiplicity is fixed by the Hodge structure on $H^2(S, \mathbb{Z}) \oplus H^*(E, \mathbb{Z})$: a nodal K_3 acquires a vanishing cycle of dimension 2, and the local universal Hodge filtration degenerates with multiplicity 2 along $\mathfrak{D}_{\text{per}}$ when the discriminant is read as the divisor of a section of the period-bundle determinant (Saito's mixed-Hodge-module formalism for the Hodge filtration on the local Milnor fibre, [?Sai toMHM]; Bryan–Pandharipande [?BRYAN, § 3] for the genus-two BPS calculation that supplies the same multiplicity at the level of generating functions).

Conjecture 1.1 (The mirror discriminant identity). On the $(K_3 \times E)$ -component of the K_3 -mirror period family, parametrised by (8.3), the meromorphic section

$$\Delta_5^{-2} \in \Gamma(\mathcal{H}_2/\mathbf{Sp}_4(\mathbb{Z}), \mathcal{L}_{\text{aut}}^{-10})$$

of the inverse 10th tensor power of the automorphic line bundle $\mathcal{L}_{\text{aut}}$ coincides, as a meromorphic section of the period-bundle determinant on $\mathcal{P}_{K_3 \times E}$, with the discriminant of the K_3 -mirror period family. Equivalently, the polar divisor of Δ_5^{-2} is twice the period-family discriminant $\mathfrak{D}_{\text{per}}$, and the comparison map

$$\mathcal{L}_{\text{aut}}^{-10} \xrightarrow{\sim} (\det \mathcal{H}_{\text{tr}}^2)^*$$

on the open complement $\mathcal{P}_{K_3 \times E}^\circ = \mathcal{P}_{K_3 \times E} \setminus \mathfrak{D}_{\text{per}}$ extends to a global identification of the inverse-period-determinant section with Δ_5^{-2} .

The conjecture is, at the level of divisor classes,

$$[\Delta_5^{-2}] = 2[\mathfrak{D}_{\text{per}}] = 2[\mathcal{H}_2^{(\text{Hum})}]; \quad (8.6)$$

the identity (8.5) fixes the right-hand side, and the conjectural content is the matching of multiplicities and the trivialisation of the comparison line bundle.

1.4.0.1 Evidence. The genus-two Bryan–Pandharipande generating function for disconnected genus-two BPS states on $K_3 \times E$ gives $Z_{\text{BPS}}^{K_3 \times E} = \Delta_5^{-2}$ in the cusp chamber [?BRYAN, Conj. 1]; this identity has been proved by Oberdieck–Pixton [?OB, Thm. 1] and is the input to View ① on the curve-counting side. Two facts converge: the coefficient of Δ_5^{-2} at a degenerate stable map is controlled, by the Maulik–Toda Gopakumar–Vafa formalism [?MaulikTodaGV], by the local Milnor monodromy on the period filtration; and the local Hodge-filtration multiplicity at a nodal K_3 fibre is 2 (Saito [?Sai toMHM]). Both of these are theorems, not conjectures. What is conjectural is the global trivialisation: the comparison line bundle $\mathcal{L}_{\text{aut}}^{-10} \otimes (\det \mathcal{H}_{\text{tr}}^2)$ admits no obstruction, and the identification of the two sides extends across the discriminant divisor.

A separate piece of evidence is the precedent set by Gritsenko–Nikulin [?GNII, Thm. 4.1]: at rank 4, with polarisation $L = U \oplus U$, the analogous Borcherds product is the Igusa cusp form Φ_{12} on the type-IV symmetric domain of signature $(2, 4)$; Gritsenko–Nikulin proved that Φ_{12} is the quasi-pull-back of the discriminant section of the universal lattice-polarised Hodge bundle on Ω_L/Γ_L . Conjecture 1.1 is the rank-3 analogue of the same identification, evaluated at the lattice $M = U \oplus \langle -2 \rangle$.

1.4.0.2 Status. The identification claim of Conjecture 1.1 is conjectured. The existence of the comparison line bundle is folklore; the divisor identity (8.5) is a theorem; the multiplicity-2 statement on the K_3 side is a theorem; the global extension across the discriminant divisor is the conjectural content.

1.4.0.3 What this conjecture supplies. If proved, View ④ embeds the present programme inside mirror symmetry on $K_3 \times E$ at the level of period-side discriminants. The identification supplies a new derivation of Δ_5 that does not pass through the Borchers product (View ①), the BKM denominator (View ②), the Pfaffian of the Dirac–Igusa pro-object (View ③), or the singular theta lift (View ⑤). It is a fifth chain. The fact that the same form arises on five chains is the content of the pentadic thesis of the manuscript; the fifth chain, conjectural now, completes the picture.

The opposite implication is also of interest. A proof of Conjecture 1.1 would identify Δ_5^{-2} with the universal determinant section of the $(K_3 \times E)$ -mirror period family; combined with the Bryan–Oberdieck–Pixton theorem [?OB, Thm. 1] that $Z_{\text{BPS}}^{K_3 \times E} = \Delta_5^{-2}$, this would give a new proof of the genus-two BPS partition-function identity through the period side, complementing the curve-counting derivation.

1.5 What the conjecture does not assert

The Beilinson cut (CLAUDE.md § III) is in force. The conjectural identification is on the period side: it concerns the section Δ_5^{-2} of an automorphic line bundle and the discriminant divisor of the K_3 -mirror period family. It is not a claim about the protected operator-level package $\mathfrak{D}_X^{\text{DI}}$ of View ③; that package is constructed and proved, under the four named obstructions (Do), (O1), (Or⁺), (O2), in the View ②↔③ chapters preceding this one. View ④ does not enter the operator chain.

The level- n scalar partition function $Z_{\text{BPS}}^{K_3 \times E} = \Delta_5^{-2}$ is, on the present programme, the orientation-forgetful trace of the Pfaffian package; its promotion to a $3d$ gravitational path integral requires the Hall–Borchers residual (Vol II, [?ChiralBarCobarVolIII]). The mirror conjecture above says nothing about the gravitational lift; it addresses only the geometric origin of the section Δ_5^{-2} on the K_3 -mirror period family.

The Borchers–Kac–Moody denominator $\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1} \Delta_5(2Z)$ is on the algebraic side; it is a theorem of Gritsenko–Nikulin [?GNII, § 3, Thm. 3.1]. The mirror conjecture is on the period side and does not enter the algebraic chain.

1.6 Lattice-convention check

The lattice convention adopted above is the Dolgachev convention $\Lambda_{K_3} = 3U \oplus 2E_8(-1)$, $M = U \oplus \langle -2 \rangle$, mirror lattice $\check{M} \simeq U \oplus E_7(-1) \oplus E_8(-1)$ (rank 17, signature (1, 16)). Aspinwall–Morrison’s original mirror-pair construction [?AspinwallMorrisonStringy] adopts the same lattice up to the choice of which hyperbolic plane in Λ_{K_3} is split off; the two conventions agree on the K_3 mirror lattice. The Mukai-lattice form $\widetilde{M} = U \oplus M = 2U \oplus \langle -2 \rangle$ of rank 5 and signature (2, 3) is the convention adopted in cross-volume Vol III [?CYQGVolIII, Prop. 6.4]; it is the convention relevant to the period-domain bridge (8.2). The lattice $\widetilde{M}$ is identified with the manuscript’s $\Lambda^{3,2}$ of § ?? by the sign-flip on the rank-1 summand recorded in § 1.1. Disagreement on the lattice convention surfaces, in cross-volume work, only as a sign convention on the Mukai pairing; it does not affect the discriminant divisor.

The K_3 vs. $(K_3 \times E)$ period-space matching (8.3) works because *the Mukai U -summand of $\widetilde{M}$ is the elliptic factor on the $(K_3 \times E)$ -side after Künneth absorption*, not because the K_3 -mirror period family and the $(K_3 \times E)$ -mirror period family are abstractly equal. Disagreement on this point would surface as a discrepancy between the $\mathbf{Sp}_4$ -side and the $\mathbf{O}^+(2, 3)$ -side; the isomorphism (8.2) is exactly the $\mathbf{Sp}(4)$

$\simeq \text{Spin}(3,2)$ bridge of View ⑤, and the absorption of $\Lambda_E = U$ into the Mukai U -summand of $\widetilde{\mathcal{M}}$ is the Hodge-theoretic content of the Künneth product.

The phrase “discriminant of WHAT” is sharpened by the multiplicity. There are two natural divisors on the period side: the support divisor $\mathfrak{D}_{\text{per}}$ of the period-family degeneration, and the principal divisor of the section of $(\det \mathcal{H}_{\text{tr}}^2)^*$ realising the period-bundle determinant. The conjecture concerns the second: Δ_5^{-2} is conjectured to be that section, with polar divisor twice the support divisor of $\mathfrak{D}_{\text{per}}$. The factor of 2 is forced by the local multiplicity of the Hodge filtration at a nodal K_3 and is matched, on the automorphic side, by the square of Δ_5 ; a mismatch at the level of multiplicity would falsify the conjecture by a factor that the Borcherds product does not allow.

The relation to the Borcherds–Gritsenko–Nikulin precedent at rank 4 is the cleanest comparison. At $L = U \oplus U$ the analogous lift is Φ_{12} on the type-IV symmetric domain of signature $(2, 4)$, and Gritsenko–Nikulin [?GNII, Thm. 4.1] prove that Φ_{12} is the quasi-pull-back of the discriminant section of the universal lattice-polarised Hodge bundle. The rank-3 analogue $L = U \oplus \langle -2 \rangle$ is Conjecture 1.1; the only piece missing is the global comparison-line trivialisation. The Φ_{12} precedent is evidence; it is not a proof, since the quasi-pull-back map at rank 3 is governed by a different Heegner divisor structure.

1.7 Open obligations

- (M1) *Trivialisation of the comparison line.* The comparison line bundle $\mathcal{L}_{\text{aut}}^{-10} \otimes (\det \mathcal{H}_{\text{tr}}^2)$ on $\mathcal{P}_{K_3 \times E}^\circ$ carries no first Chern class that is not absorbed by the automorphic-character data of $\nu_{\Delta_5}^{-2}$. Sub-conjecture: this comparison line is trivial. Implication: under (M1), the identity (8.6) promotes from a divisor identity to an equality of sections.
- (M2) *Multiplicity at the cusp.* Conjecture 1.1 concerns the polar divisor of Δ_5^{-2} on the open part $\mathcal{P}_{K_3 \times E}^\circ$; the boundary behaviour at the rank-2 cusp of $\mathcal{H}_2/\mathbf{Sp}_4(\mathbb{Z})$ is governed by the singular weight $c_1(\mathfrak{o})/2 = 5$ ([?Borcherds95, § 15]). Sub-conjecture: the boundary multiplicity matches the singular weight on the period side.
- (M3) *The mirror involution ι_{DN} on the section.* The Dolgachev–Nikulin involution on $\widetilde{\Lambda}_{K_3}$ acts as $-\text{id}$ on $\widetilde{\mathcal{M}} = 2U \oplus \langle -2 \rangle$ (signature $(2, 3)$) and as $+\text{id}$ on $\widetilde{\mathcal{M}}^\perp$ (signature $(2, 17)$); both halves carry the same positive-part rank 2, which is the structural source of the symmetry. Sub-conjecture: $\iota_{\text{DN}}^*(\Delta_5) = \Delta_5$ up to the multiplier ν_{Δ_5} , and the discriminant identity is ι_{DN} -invariant.
- (M4) *Cross-volume consistency at rank ≥ 5 .* The Vol III rank- ≥ 5 lattice-polarised $\mathfrak{g}_L$ family [?CYQGVolIIIIWave3] is the conjectural extension of Conjecture 1.1 to higher Picard rank. At rank ≥ 5 the discriminant identity is conditional on the Gritsenko–Cléry 2018 admissibility conjecture (Conj. 5.1 of [?GC]); the rank-3 statement is the corresponding conditional statement for the K_3 -Mukai-invariant lattice and is the only one of the rank ladder relevant to this manuscript.

The four open obligations (M1)–(M4) are distinct from the four obstruction classes (Do), (O1), (O1⁺), (O2) of View ③; the latter govern the operator-level chain and are the conditioning for ② $\leftrightarrow$ ③, while the former govern the period-side chain and are the conditioning for ④ $\leftrightarrow$ ①. The two sets do not interact: a discharge of (M1) does not advance the operator package, and a discharge of (Do) does not advance the period-side conjecture.

1.8 Cross-repository consistency

The lattice convention $\mathcal{M} = U \oplus \langle -2 \rangle$, $\check{\mathcal{M}} \simeq U \oplus E_7(-1) \oplus E_8(-1)$, $\widetilde{\mathcal{M}} = U \oplus \mathcal{M}$ of § 1.1 is the convention of Vol III [?CYQGVolIIII, § 6.4 + Prop. 6.4 + Thm. 6.7]; on the $(K_3 \times E)$ -component the Vol III κ -tuple $(\kappa_{\text{cat}}, \kappa_{\text{ch}}^{\text{Hodge}}, \kappa_{\text{ch}}^{\text{Heis}}, \kappa_{\text{BKM}}, \kappa_{\text{fiber}}) = (0, 0, 3, 5, 24)$ is recorded with each entry sourced

from a distinct construction. The mirror discriminant of Conjecture 1.1 is on the $\kappa_{\text{BKM}} = 5$ row of the κ -tuple; the period-side input is the discriminant divisor on the lattice-polarised period stratum of the Vol III chart classification (Vol III [CYQGVolIII, § V]: the “lattice-polarised period domain” equivariance stratum). The $(K_3 \times E)$ -fiber rank $\kappa_{\text{fiber}} = 24$ is the Mukai-lattice rank of $\tilde{\Lambda}_{K_3}$ and enters the present chapter only as the rank of the Heisenberg algebra of the Mukai lattice, not as a discriminant invariant.

The mixed-holomorphic-topological-strings companion volume [MixedHTSCompanion] records that the modular line bundle on $\overline{\mathcal{A}}_g$ at level $g = 2$ is Ω_{central} , and that $\Phi_{10} = \chi_{10} = \Delta_5^{-2}$ is the level-2 section. The conjecture above is consistent with that statement: the polar divisor of $\Delta_5^{-2} = \Phi_{10}^{-1}$ on $\overline{\mathcal{A}}_2$ is twice the divisor of Δ_5 , and the mirror discriminant identification is precisely the geometric origin of that section.

The Vol II Hall–Borcherds residual is silent on View ④: the period-side discriminant identification is independent of the gravitational-lift bridge, and the conjecture above does not require the residual to be discharged. A discharge of the Hall–Borcherds residual on the $(K_3 \times E)$ -component would not by itself prove Conjecture 1.1; conversely, a proof of Conjecture 1.1 would not automatically discharge the Hall–Borcherds residual.

The fifth view stands open. The four other views form a single quadrilateral of theorems and conditional theorems; the fifth view is the missing diagonal. Its discharge is the next geometric problem in the programme.

THE LEVEL- n PROTECTED SCALAR SHADOW AND THE $K_3 \times E$ REALIZATION QUESTION

The denominator algebra and the protected Pfaffian sit at the primitive and chart levels of the universal stage chain $P \rightarrow C \rightarrow S \rightarrow Z \rightarrow A$. At level S the same Δ_5 survives, squared, as the protected partition function on the closed $K_3 \times E \rightarrow \text{pt}$ cobordism. This chapter records the scalar shadow exactly as the Beilinson cut requires it: an automorphic identity, not an algebra; a trace, not an operator; an orientation-forgetful invariant, not an orientation lift. The orientation lift is the missing first-order object $\mathfrak{D}_X^{\text{DI}}$ of Chapter 5; the scalar identity is the level- n shadow it casts.

Three distinct meanings of Δ_5 coexist in this chapter and are tagged on first occurrence: the automorphic section $D_X = \Delta_5$ (View I -- Borchers–Gritsenko–Nikulin); the Borchers–Kac–Moody denominator $\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1} \Delta_5(2Z)$ (View II -- Gritsenko–Nikulin); the conditional protected Pfaffian $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}})$ (View III -- Chapter 5). The handle “ Δ_5 ” alone never substitutes for any of the three.

I THE LEVEL- n PROTECTED SCALAR SHADOW ON $K_3 \times E \rightarrow \text{pt}$

I.1 The Oberdieck–Pixton scalar identity $[\beta]$

Let $X = S \times E$, with S a smooth complex K_3 surface and E a smooth elliptic curve. The reduced Donaldson–Thomas / stable-pair / Gromov–Witten correspondence on X , in the primitive K_3 -class chamber, identifies the scalar partition function Z_{DT}^X with the inverse of the Igusa cusp form on the genus-2 Siegel upper half-space $\mathfrak{H}_2$. Using the chamber and variable conventions of Proposition ?? and the equality of Theorem ??,

$$Z_{\text{DT}}^X(T, Q, P) = Z_{\text{OP}}^X(P, Q, T) = -4096 \Delta_5(Z)^{-2}, \quad Z = \Omega \in \mathfrak{H}_2. \quad (9.1)$$

The constant $4096 = 64^2$ is the squared theta-leading coefficient $[q^{1/2} r^{1/2} s^{1/2}] \Delta_5$ of the automorphic section, and the leading sign is the Oberdieck–Pixton scalar convention $Z_{\text{OP}}^X = -(\chi_{\text{io}}^{\text{OP}})^{-1}$. Both are normalization data of the imported reduced theory.

THEOREM I.1 (*Level- n protected scalar shadow on the closed $K_3 \times E \rightarrow \text{pt}$ cobordism $[\alpha + \beta]$*). With variables (T, Q, P) normalised in the Oberdieck–Pixton chamber of Proposition ??, the level- n protected scalar shadow on the closed $K_3 \times E \rightarrow \text{pt}$ cobordism is

$$Z_{\text{BPS}}^{K_3 \times E}(T, Q, P) = (\Phi_{\text{io}}^{\text{un}}(Z))^{-1} = \Delta_5(Z)^{-2}. \quad (9.2)$$

Equivalently, in OP’s normalization $\chi_{\text{io}}^{\text{OP}} = 2^{-12} \Delta_5^2$, $\Phi_{\text{io}}^{\text{un}} = \chi_{\text{io}} = \Delta_5^2$, and the unimodular form $(\Phi_{\text{io}}^{\text{un}})^{-1}$ is the level- n protected trace on every closed K_3 -class fibre. The Behrend-weighted reduced quotient integration of Proposition ?? identifies this scalar with the reduced DT, PT, and OP series $Z_{\text{DT}}^X = Z_{\text{OP}}^X = Z_{\text{PT}}^X = -4096 \Delta_5^{-2}$ on the primitive branch.

Proof. Equation (9.2) is the dictionary between the Oberdieck–Pixton primitive series and the unimodular Igusa form Φ_{10}^{un} : the OP normalization $\chi_{10}^{\text{OP}} = 2^{-12} \Delta_5^2$ of Theorem ?? together with $\Phi_{10}^{\text{un}} = \Delta_5^2$ gives $(\Phi_{10}^{\text{un}})^{-1} = \Delta_5^{-2}$. Multiplying by $2^{12} = 4096$ and the OP scalar sign restores (9.1). The reduced DT/PT comparison is [?OberdieckReducedPT, Theorem 3(i)], the reduced PT/GW comparison on primitive K_3 classes is [?OPand, Proposition 5], and the OP product identity is [?OB, Theorem 1]; their composition gives the displayed equality in the OP chamber. $\square$

1.2 Three readings of the same exponent

The integers $f(n, l)$ of the weight-zero index-one weak Jacobi form $\phi_{0,1}$ drive both sides of the OP/Borcherds dictionary. At level P they are virtual K_0 -superdimensions of $\mathbb{U}_{n,l}$, and at level S they appear twice: once as exponents of the Borcherds product on the automorphic side, once as multiplicities in the Mukai dimension count of the Hilbert-scheme stratification on the geometric side. In $\mathfrak{g}_{\Delta_5}$, the same $f(nm, l)$ records the signed root supermultiplicity at $\alpha(n, l, m) = 2nf_2 - lf_3 + 2mf_{-2}$.

The variable map of Proposition ?? encodes this triple reading. A factor $(1 - q^n r^l s^m)^{f(nm, l)}$ of the Borcherds product on the automorphic side is, on the Oberdieck–Pixton side, a factor $(1 - p_{\text{OP}}^k q_{\text{OP}}^h \bar{q}_{\text{OP}}^d)^{c(4hd - k^2)}$ of χ_{10}^{OP} , the K_3 -elliptic-genus exponents $c(4hd - k^2) = 2f(hd, k)$ recording the squared automorphic line. The multiplicities $2f$ are exactly the K_3 elliptic-genus coefficients of $Z_{K_3} = 2\phi_{0,1}$; the squaring is built into the identification $\chi_{10}^{\text{OP}} = D_5^2$ of Theorem ??.

Remark 1.2 (Constants are normalization, not orientation [α]). Three constants intervene in (9.1). The leading $64 = [q^{1/2} r^{1/2} s^{1/2}] \Delta_5$ is the theta-constant normalization of the Igusa section; the monic Borcherds product is $D_5 = 64^{-1} \Delta_5$. The factor $4096 = 64^2$ is the squared theta-leading coefficient produced by the OP convention $\chi_{10}^{\text{OP}} = D_5^2$. The leading minus sign is the OP scalar branch $Z_{\text{OP}}^X = -(\chi_{10}^{\text{OP}})^{-1}$. None of the three is the orientation character. The protected orientation character ϵ_0 and the automorphic reflection character ν_{Δ_5} are determined elsewhere: ν_{Δ_5} by the Maass transformation law of Δ_5 , ϵ_0 by the type-II Pfaffian wall computation of (O₂) on the half-Hilbert orbit through the 2^{12} sign sum. The number $4096 = 2^{12}$ recurring in the (O₂) wall count is the *same* algebraic source as the OP normalization 2^{12} , but the *rolls* are distinct: the Pfaffian-side 2^{12} is the cardinality of a sign orbit that the local wall computation must collapse, while the OP-side 2^{12} is the squared theta-leading coefficient that the imported scalar identity absorbs.

1.3 Worked Fourier-coefficient check

The leading Fourier coefficients of Δ_5 and Δ_5^2 provide the elementary verification that the constants in (9.1) are normalization, not enumerative content. Recall $\phi_{0,1}$ has leading expansion

$$\phi_{0,1}(\tau, z) = (r^{-1} + 10 + r) + q(10r^{-2} - 64r^{-1} + 108 - 64r + 10r^2) + O(q^2),$$

with coefficients $f(0, -1) = f(0, 1) = 1, f(0, 0) = 10, f(1, 1) = f(1, -1) = -64, f(1, 0) = 108, f(1, 2) = f(1, -2) = 10, f(1, 3) = f(1, -3) = 0$. The Borcherds–Gritsenko–Nikulin product

$$\Delta_5(Z) = 64 q^{1/2} r^{1/2} s^{1/2} \prod_{(n, l, m) \in \Gamma_{\text{eff}}} (1 - q^n r^l s^m)^{f(nm, l)} \quad (9.3)$$

opens with the theta-leading term $[q^{1/2} r^{1/2} s^{1/2}] \Delta_5 = 64$; the next non-trivial coefficient is $[q^{3/2} r^{1/2} s^{1/2}] \Delta_5$, produced by the factor $(1 - qrs)^{f(1,1)} = (1 - qrs)^{-64}$ along with the active $(n, l, m) = (0, \pm 1, 0)$ factor $(1 - r^{\pm 1})^{f(0, \pm 1)} = (1 - r^{\pm 1})$. Squaring (9.3),

$$\Delta_5^2(Z) = 4096 q r s \prod (1 - q^n r^l s^m)^{2f(nm, l)}, \quad (9.4)$$

with theta-leading $64^2 = 4096$. The OP product $\chi_{10}^{\text{OP}} = p q \tilde{q} \Pi(1 - p^k q^h \tilde{q}^d)^{c(4hd-k^2)}$, with K_3 -elliptic exponents $c(4hd - k^2) = 2f(hd, k)$, is exactly the right-hand side of (9.4) divided by 4096. Thus

$$\chi_{10}^{\text{OP}} = 4096^{-1} \Delta_5^2, \quad [pq\tilde{q}] \chi_{10}^{\text{OP}} = 1, \quad [q^{1/2} r^{1/2} s^{1/2}] (64^{-1} \Delta_5) = 1.$$

The squared theta-leading coefficient 4096 is therefore the single explicit normalization constant absorbed by the OP/DT scalar identity $Z_{\text{OP}}^X = -(\chi_{10}^{\text{OP}})^{-1}$; the sign -1 is OP convention; together they produce the constant -4096 of (9.1). The Fourier verification at $q^{1/2} r^{1/2} s^{1/2}$ resolves the question “where does the 4096 come from?” at the level of a single coefficient: it is $([q^{1/2} r^{1/2} s^{1/2}] \Delta_5)^2$, a leading-coefficient square, not an enumerative invariant.

1.4 Cobordism positioning and the closed $K_3 \times E \rightarrow \text{pt}$ cobordism

The cobordism $K_3 \times E \rightarrow \text{pt}$ is the closed 3-manifold viewed in the closed-CY-target sense: the protected partition function of $\mathfrak{A}_{K_3 \times E}$ evaluated on the bordism with empty boundary. At level S the partition function is a complex number; the “level- n ” label $n = \chi(O_{K_3 \times E}) = \chi(O_{K_3}) \chi(O_E) = 2 \cdot 0 = 0$, so the relevant grading for $K_3 \times E$ is the elliptic degree d and the K_3 norm $h - 1$, not the holomorphic Euler characteristic. The dual notation $Z_{\text{BPS}}^{K_3 \times E}(T, Q, P)$ stores the elliptic and K_3 gradings as a Laurent expansion of the inverse Igusa cusp form on $\mathfrak{H}_2$.

The $K_3 \times E$ target is a threefold; the perfect obstruction theory and the virtual fundamental class $[X]^{\text{vir}}$ are those of the threefold (not of any auxiliary fourfold). “ n ” in “level- n shadow” refers to the cobordism class of the closed K_3 -class fibre at primitive curve class β_h of norm $2h - 2$, not to the holomorphic Euler characteristic of the target.

2 SPIN L -FACTOR NORMALISATION FOR Δ_5^2

2.1 The Saito–Kurokawa eigenline

The unimodular Igusa form $\Phi_{10}^{\text{un}} = \Delta_5^2$ is the weight-10 Saito–Kurokawa lift of the unique normalized cusp form $g = \Delta E_6 \in S_{18}(\mathbf{SL}_2(\mathbb{Z}))$; the Hecke eigenline is the one-dimensional $\mathbb{C} \Delta_{10}^\theta = \mathbb{C} \chi_{10}^{\text{OP}}$, independent of the scalar normalization of its generator. The OP monic normalization is

$$\chi_{10}^{\text{OP}} = 2^{-12} \Delta_{10}^\theta = 4096^{-1} \Delta_5^2, \quad (9.5)$$

and the level- n shadow (9.2) is the inverse of the *theta-normalized* Hecke eigenline element $\Delta_{10}^\theta = \Delta_5^2$.

THEOREM 2.1 (*Andrianov factorisation of the spinor L -function* [β]). For the Saito–Kurokawa representation generated by $\Delta_{10}^\theta = \Delta_5^2$,

$$L_{\text{spin}}(s, \Delta_{10}^\theta) = \zeta(s-8) \zeta(s-9) L(s, g), \quad g = \Delta E_6.$$

The completed spinor L -function for a genus-two form of weight 10 is $L_{\text{spin}}^*(s, F) = (2\pi)^{-2s} \Gamma(s) \Gamma(s-8) L_{\text{spin}}(s, F)$.

This is Theorem ??; the proof uses Andrianov’s spinor L -factor formula for the degree-2 Saito–Kurokawa lift attached to the weight-18 elliptic cusp form g [Andr74], together with the dimension count $\dim S_{18}(\mathbf{SL}_2(\mathbb{Z})) = 1$ which forces the source form to be $g = \Delta E_6$ [ZagierSK].

PROPOSITION 2.2 (*Pole and residue at $s = 10$ [β]*). With $\Lambda(g, s) = (2\pi)^{-s} \Gamma(s) L(s, g)$, the functional equation $\Lambda(g, s) = -\Lambda(g, 18 - s)$ gives $L(9, g) = 0$, so the possible pole of $\zeta(s - 8) L(s, g)$ at $s = 9$ is cancelled. At $s = 10$ the residue is

$$\text{Res}_{s=10} L_{\text{spin}}(s, \Delta_{10}^\theta) = \zeta(2) L(10, g) = \frac{\pi^2}{6} L(10, g),$$

with $L(10, g) \in \mathbb{Q}^\times (2\pi i)^{10} \Omega^+(g)$ by Manin–Deligne periods, hence $L(10, g) \neq 0$. The function $L_{\text{spin}}(s, \Delta_{10}^\theta)$ has a simple pole at $s = 10$ with non-zero residue.

This is Proposition ??, packaged with Proposition ??; the proof uses the Eichler–Shimura–Manin period theorem for the rational newform g [Manin72, ?KZ], in Deligne’s period normalisation [De179].

Remark 2.3 (*Hecke eigenline is normalization-independent [α]*). The spinor L -function depends on the Hecke eigenline $\mathbb{C} \Delta_{10}^\theta = \mathbb{C} \chi_{10}^{\text{OP}}$, not on the scalar used to label its generator. Theorem 2.1 and Proposition 2.2 are statements about the eigenline. The OP scalar constant 2^{-12} of (9.5) relabels the generator from Δ_5^2 to χ_{10}^{OP} ; it does not change L_{spin} , the residue, or the central zero.

2.2 Independence from the Dirac–Igusa construction

The Saito–Kurokawa factorization of $L_{\text{spin}}(s, \Delta_5^2)$ and the OP/DT scalar identity (9.1) are imported results: the first from Andrianov [Andr74], the second from Oberdieck–Pixton [OB] and Oberdieck [OberdieckReducedPT]. Neither uses the protected Pfaffian Pf_{prot} of Chapter 5, the orientation character ϵ_o of $(\text{O}1)/(\text{O}1^+)$, the type-II wall normal form of $(\text{O}2)$, or the primitive recognition $P_X \cong \mathfrak{g}_{\Delta_5}$ of Chapter 6. Conversely, the construction of $\mathfrak{D}_X^{\text{DI}}$ and the Pfaffian–Dirac theorem $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}}) = \Delta_5$ do not use the spinor L -factor or the scalar inverse Igusa form. The two routes meet at the squared section $\Delta_5^2 = \Phi_{10}^{\text{un}}$ and at no earlier point.

3 WHAT THE SCALAR SHADOW DOES NOT SUPPLY

3.1 Z_{BPS} is the level- n protected trace, not a path integral

The squared section Δ_5^2 and its inverse $Z_{\text{BPS}}^{K_3 \times E} = \Delta_5^{-2}$ sit at level S of the universal stage chain, downstream of the primitive object at level P (the open factorization dg-category, the boundary vacuum b , and $A_b = \text{End}_C(b)$) and downstream of the chart object at level C (the disk algebra and its bar / cobar / chiral Hochschild data). The scalar partition function records the *trace* of the protected data; it does not by itself construct the protected data.

Scope 3.1 (*No promotion to a 3D gravitational path integral [δ]*). The identity (9.1), packaged through OP/DT/PT primary literature, is a closed automorphic equation between scalar power series. No statement of this chapter, of Chapter 3, or of Chapter 5 promotes $Z_{\text{BPS}}^{K_3 \times E}$ to a 3-dimensional gravitational path integral on $\text{AdS}_3 \times S^3 \times K_3$, to a metric Euclidean path-integral over the full conformal class of the boundary, or to a holographic dual partition function in any sense beyond the boundary-CFT reading recorded by Vol II of the constellation. The equality $\Phi_{10}^{\text{un}} = \Delta_5^2$ gives the scalar character of a candidate gravity-line trace; the construction of that trace is open.

Remark 3.2 (*The forbidden identification, replaced [α]*). The cross-volume slogan “ Z_{BPS} is a 3D gravitational path integral” confuses level S with level Z of the universal stage chain. Replaced: $Z_{\text{BPS}}^{K_3 \times E}$ is the level- n protected trace of $\mathfrak{A}_{K_3 \times E}$; the gravitational path integral, if such a thing exists in this corner of the constellation, requires the Hall–Borcherds residual of Vol II (§3.2). The residual is open; the

path-integral promotion is not claimed. In particular, no microscopic Hilbert space $\mathcal{H}_{K_3 \times E}$ is asserted to support $\dim \mathcal{H}_{K_3 \times E, \gamma} = f(\gamma)$ outside the charge-graded virtual reading recorded in Chapter 2.

3.2 The Hall–Borcherds residual

The promotion from a scalar partition function to a chain-level gravity-line operator algebra requires a comparison datum that lives neither in the automorphic / OP reduced theory nor in the present manuscript. Vol II names this comparison.

Conjecture 3.3 (Hall–Borcherds residual; gravity-line comparison $[\beta+\delta+\epsilon]$). Let $X = K_3 \times E$ and assume the oriented critical Hall datum used for the Vol III positive-half comparison

$$\Theta_{\text{Hall} \rightarrow \text{Borch}} : \widehat{\text{CoHA}}_{\Lambda_{II}^{2,1}}^{\text{red}}(X) \longrightarrow U^{\text{ch}}(\widehat{\mathfrak{n}_+(\mathfrak{g}_{\Delta_5})}). \quad (9.6)$$

The Hall–Borcherds residual is the conjectural extension of (9.6), after Drinfeld doubling, current enveloping along E , and weight-completion, to a filtered $\text{SC}^{\text{ch}, \text{top}}$ morphism from the $K_3 \times E$ Hall boundary object to the gravitational boundary line algebra, whose derived-centre trace has scalar character $(\Phi_{10}^{\text{un}})^{-1} = \Delta_5^{-2}$.

The residual is the second of the four Construction Problems of Vol II of the constellation: source `chiral-bar-cobar-vol2:chapters/connections/3d_gravity.tex`, Conjecture `conj:gravity-line-hall-borcherds-comparison` at line 8429. It is not claimed in the present manuscript; its existence and the construction of its witnessing morphism are open.

Remark 3.4 (Cross-repository consistency $[\gamma]$). The κ -tuple of the $K_3 \times E$ witness records the structural data behind the scalar shadow:

$$(\kappa_{\text{cat}}, \kappa_{\text{ch}}^{\text{Hodge}}, \kappa_{\text{ch}}^{\text{Heis}}, \kappa_{\text{BKM}}, \kappa_{\text{fiber}})(\mathfrak{A}_{K_3 \times E}) = (0, 0, 3, 5, 24).$$

The fourth entry $\kappa_{\text{BKM}} = 5$ is the cusp-form weight $\text{wt}(\Delta_5)$, the entry that anchors this manuscript; the fifth entry $\kappa_{\text{fiber}} = \chi_{\text{top}}(K_3) = 24$ is the topological Euler characteristic of the K_3 fibre, distinct from $\chi(\mathcal{O}_{K_3}) = 2$. The additive collapse $\kappa_{\text{BKM}} = \kappa_{\text{ch}}^{\text{Hodge}} + \chi(\mathcal{O}_{\text{fiber}})$ confuses these two: with $\kappa_{\text{ch}}^{\text{Hodge}} = 0$ and $\chi(\mathcal{O}_{K_3}) = 2$, it predicts $\kappa_{\text{BKM}} = 2$, which fails because the universal Borcherds weight at $N = 1$ is $\kappa_{\text{BKM}}(\Phi_1) = c_1(0)/2 = 10/2 = 5$ (Borcherds 1995, Gritsenko 1999). The five entries of the tuple are five *distinct* constructions; they cannot be combined additively. The Vol III counterexample at $N = 1$ is the controlling instance.

3.3 Local-to-global in mixed holomorphic–topological language

A second route from local protected data to a global compact target runs through the mixed holomorphic–topological model of Costello and the holomorphic de Rham obstruction class. The slogan there is “formal local Hamiltonian BF $\Rightarrow$ global compact target”; the corrected form requires vanishing of the global holomorphic de Rham obstruction class $[\text{obs}_{HT, \text{glob}}] \in H^1(X, \mathbb{C}_X)$, plus descent, quantum master equation, anomaly, and locality data, with the Hamiltonian condition supplied by the local model. For $X = K_3 \times E$ the global obstruction group is $H^1(K_3 \times E, \mathbb{C}_{K_3 \times E}) \cong \mathbb{C}$ (non-zero), so the formal-local-Hamiltonian-BF route does not by itself promote $Z_{\text{BPS}}^{K_3 \times E} = \Delta_5^{-2}$ to a global compact target either. Both bridges -- Hall–Borcherds for the Vol II gravity line, mixed-HT for the Costello local-to-global ascent -- are open; the present chapter records the scalar shadow as it stands.

4 THE $K_3 \times E$ THREEFOLD REALIZATION QUESTION

The square root in the title is meant in Dirac's sense. The OP/DT scalar branch $Z_{\text{OP/DT}}^X = -4096 \Delta_5^{-2}$ is orientation-forgetful: the sign character is squared away, and the squared scalar cannot recover the sign by which a Pfaffian-line section distinguishes Δ_5 from $-\Delta_5$. The missing first-order object is the protected Pfaffian, whose square is the inverse of the protected scalar shadow.

4.1 From the scalar to the operator-level object

The Dirac square-root pattern is the level-Z-to-level-S descent. At level Z one has the protected Pfaffian (View III -- Chapter 5); at level S one has the inverse squared scalar shadow. The Pfaffian carries the orientation data; squaring forgets it; the squared scalar identity records what remains.

THEOREM 4.1 (*Square-root consequence at level S [$\alpha + \beta + \epsilon$]*). *Conditional on (Do), (O₁), (O₁⁺), (O₂), the protected Pfaffian $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}})$ of Chapter 5 is the automorphic section $D_X = \Delta_5$ of Chapter 3, and its inverse square is the level- n protected scalar shadow:*

$$(\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}}))^{-2} = \Delta_5(Z)^{-2} = (\Phi_{\text{io}}^{\text{un}}(Z))^{-1} \propto Z_{\text{BPS}}^{K_3 \times E}(T, Q, P), \quad (9.7)$$

the constant of proportionality being -4096^{-1} on the OP/DT branch of Theorem ??.

Proof. The Pfaffian–Dirac theorem of Chapter 6, conditional on (Do), (O₁), (O₁⁺), (O₂), asserts $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}}) = \Delta_5$. Squaring, $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}})^2 = \Delta_5^2 = \Phi_{\text{io}}^{\text{un}}$; inverting, $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}})^{-2} = (\Phi_{\text{io}}^{\text{un}})^{-1}$. By Theorem 1.1 the right-hand side equals the level- n protected scalar shadow, and by Theorem ?? that scalar equals $-4096^{-1} Z_{\text{OP/DT}}^X$ in the OP chamber. The proportionality constant -4096^{-1} absorbs the OP convention, the squared theta-leading coefficient, and the sign branch of (9.1). $\square$

4.2 The orientation datum the squared scalar cannot recover

The square-root sign is invisible to the scalar. An orientation-line sign $\varepsilon \in \{\pm 1\}$ acting by $\Delta_5 \mapsto \varepsilon \Delta_5$ satisfies $\varepsilon^2 = 1$; the scalar Δ_5^{-2} is therefore ε -invariant. Selecting the sign requires an algebraic input from level Z: the protected orientation character ϵ_o of the type-II Pfaffian wall normal form (O₂) and its restriction to the type-II Weyl group $W^{(2)}(\Lambda_{II}^{2,1})$, matched against the automorphic reflection character ν_{Δ_5} . Equality

$$\epsilon_o|_{W^{(2)}(\Lambda_{II}^{2,1})} = \nu_{\Delta_5}|_{W^{(2)}(\Lambda_{II}^{2,1})}$$

is Open Problem ??; the present manuscript reduces the equality to (Do), (O₁), (O₁⁺), (O₂), and the four obstruction classes carry the realization burden. No constant appearing in (9.1) -- not the leading 64, not the squared 4096, not the OP scalar minus sign -- supplies ϵ_o ; the orientation datum is fixed by group theory on $W^{(2)}(\Lambda_{II}^{2,1})$ plus a single local Pfaffian wall computation, not by a normalization constant.

4.3 The four obstructions and the realization question

Let $\mathfrak{D}_X^{\text{DI}}$ denote the Dirac–Igusa pro-object

$$\mathfrak{D}_X^{\text{DI}} = \lim_R (\mathcal{A}_{X,R}^{E_3}, F_{X,R}^{\text{hyb}}, \Gamma_{X,R}, \Pi_X, \Phi_R, o_R, H_R, P_R^{\Pi}, C_{X,R}, \Theta_{\text{Kos},R}, \mathcal{L}_{\text{Pf},R}, \text{pf}_{X,R}, \varepsilon_{o,R}, \text{Rec}_R)$$

of Chapter 5: the inverse limit, over finite root windows R , of fourteen pieces of data including the local holomorphic E_3 -factorization algebra $\mathcal{A}_{X,R}^{E_3}$, the hybrid $\text{Ran}^{\text{hyb}}(E)$ carrier $F_{X,R}^{\text{hyb}}$, the cosection-twisted reduced d-critical orientation o_R , the Hall correspondence package H_R , the Pfaffian line $\mathcal{L}_{\text{Pf},R}$, the

Pfaffian section $\text{pf}_{X,R}$, and the recognition map Rec_R . The four obstructions are the conditional data for Theorem 4.1.

Realization question (the $K_3 \times E$ threefold). The square-root identity (9.7) is a theorem under $(D_0), (O_1), (O_1^+), (O_2)$:

(D_0) *Do-degeneration limit of the cosection-reduced d-critical orientation theory.* [swarm probes A005, A011, A026, A037, A042]

(O_1) *Strong reduced orientation on the $K_3 \times E$ -quotient self-Ext frame bundle.* [swarm probes A074, A076, A082]

(O_1^+) *Weyl-equivariant transport of the orientation along $W^{(2)}(\Lambda_{II}^{2,1})$ reflections.* [swarm probes A047, A049, A089]

(O_2) *Local Pfaffian wall normal form on type-II walls; the $2^{12} = 4096$ sign sum on the half-Hilbert orbit.* [swarm probes A047, A092, A100]

Discharging any obstruction tightens the conditioning of the identification $\text{View II} \leftrightarrow \text{View III}$; discharging all four makes Theorem 4.1 an unconditional theorem.

4.4 The threefold target, not a fourfold

The $K_3 \times E$ target is a complex threefold; $\dim_{\mathbb{C}} K_3 \times E = 3$, and the integer $n = \chi(O_{K_3 \times E}) = 2 \cdot 0 = 0$ appearing in the “level- n shadow” label is the holomorphic Euler characteristic of the threefold, not of an auxiliary fourfold. The perfect obstruction theory of Chapter 5 is the threefold theory; the virtual fundamental class $[X]^{\text{vir}}$ is on the moduli of objects on the threefold; the Pfaffian Pf_{prot} of Chapter 5 is the Pfaffian of the threefold’s protected operator data. The Borchers product formula and the Gritsenko–Nikulin denominator identity give the Mukai-vector dictionary on the threefold by projection along the elliptic factor; the OP/DT/PT comparison gives the scalar shadow on the threefold; the protected Pfaffian gives the square root. No fourfold target intervenes anywhere in the chain.

4.5 Comparison with the determinant-only construction

The virtual K_0 -determinant package $\mathcal{D}_X(Z) = \Delta_5(Z)$ of Chapter 2 is a Grothendieck-class record: it sees only the signed superdimensions of the underlying virtual super vector spaces. It does not see brackets, not parities of representatives, not relation constants, not completed PBW data, not Hopf-pairing radicals, not orientation gerbes, not the Pfaffian line $\mathcal{L}_{\text{Pf}}$. Squaring the determinant gives Δ_5^2 ; inverting gives the scalar shadow. All the information about the operator-level object $\mathfrak{D}_X^{\text{DI}}$ that distinguishes the conjectured $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}}) = \Delta_5$ from a non-Pfaffian square-root choice $\Delta_5 \mapsto -\Delta_5$ lives at level Z, in the Pfaffian datum and the orientation character. The squared scalar shadow (9.1) is the maximal information visible at level S; the Pfaffian and the orientation are the strictly stronger data at level Z.

5 OPEN FRONTIERS

Three open frontiers connect the present chapter to the next.

5.1 The Hall–Borchers residual revisited

The first frontier is Conjecture 3.3: the construction of the filtered $\text{SC}^{\text{ch}, \text{top}}$ morphism from the $K_3 \times E$ Hall boundary object to the gravitational boundary line algebra of Vol II, whose derived-centre trace has scalar character Δ_5^{-2} . The residual is the bridge from the level-S scalar identity of this chapter to a

level-Z operator algebra carrying the same scalar trace; it is the obstruction to promoting $Z_{\text{BPS}}^{K_3 \times E}$ into a chain-level Vol II object.

The construction requires three layers of completion. First, the Drinfeld doubling $\widehat{\text{CoHA}}^{\text{red}} \rightarrow D(\widehat{\text{CoHA}}^{\text{red}})$ of the Vol III positive-half pairing: the pairing, completion, integral form, and stable-envelope transport that convert the Hall positive half $\mathcal{Y}^+(K_3 \times E)$ into the full quantum group $G(K_3 \times E)$. Second, the current enveloping along E and weight-completion: the chiral specialization $\text{Sp}_{\Sigma_{d-1}, C}^{\text{ch}}$ along the reference curve $C = E$, together with the weight-completed ambient required to host the formal generators of $\mathfrak{g}_{\Delta_5}$. Third, the filtered $\text{SC}^{\text{ch}, \text{top}}$ morphism property: the resulting boundary object must be a filtered morphism in the two-coloured Swiss-cheese chiral-topological operad, not merely a graded-vector-space comparison. All three layers are open in this manuscript; the residual conjecture lives in Vol II.

5.2 The gravity-line operator algebra and the Pentagon-face trace

The second frontier is the gravity-line operator algebra of Vol II Construction Problem 2: a chain-level operator-algebra object $\mathfrak{Op}_{\text{grav}}^{K_3 \times E}$ whose Pentagon-face scalar trace is exactly $\Phi_{10}^{\text{un}} = \Delta_5^2$. Vol II's Brown–Henneaux dictionary $c = 3\ell/(2G_N)$ at Vir_c supplies the Pentagon-face trace structure; the K_3 -Heisenberg witness specializes Vol II's Virasoro λ -bracket data $\{T_\lambda T\} = \partial T + 2T\lambda + (c/12)\lambda^3$ to the closed K_3 boundary at $\kappa_{\text{BKM}}(\Delta_5) = 5$; the Pentagon-face scalar trace, evaluated on the $K_3 \times E$ closed cobordism, is $\Phi_{10}^{\text{un}} = \Delta_5^2$, the squared automorphic section. Whether the gravity-line operator algebra exists, whether its Pentagon-face trace agrees with the OP/DT scalar shadow (9.1) on every closed K_3 -class fibre, and whether it factors through the protected Pfaffian $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}}) = \Delta_5$ of Chapter 5 are three open questions, posed at level Z and forwarded to Vol II.

5.3 Mathieu and umbral moonshine adjacency

The third frontier is the Mathieu / umbral moonshine adjacency. The K_3 elliptic genus $Z_{K_3} = 2\phi_{0,1} = 24\hat{\phi}_{0,1} + \sum_{r=1}^{23} A_r \hat{\phi}_{0,r}$ admits a virtual-character decomposition into the irreducible characters of the sporadic group M_{24} (Eguchi–Ooguri–Tachikawa [EOT]); the same coefficients $f(n, l)$ of $\phi_{0,1}$ which drive the Borcherds product on the automorphic side are M_{24} -virtual-character traces $\sum_r \text{Tr}_r(g) \hat{\phi}_{0,r}(\tau, z)$ on the moonshine side. The umbral moonshine extension (Cheng–Duncan–Harvey) replaces M_{24} by twenty-three sporadic extensions indexed by the Niemeier root systems. The interaction between this sporadic-group structure on the K_3 elliptic genus and the automorphic / BKM denominator structure on the $K_3 \times E$ Igusa cusp form is the Mathieu / umbral moonshine adjacency; the role it plays in the realization question, if any, is forwarded to Chapter 10.

5.4 The closed-cobordism partition reads as one

The level-S shadow is the trace, the level-Z operator is the Pfaffian, and the level-P primitive object is the chiral E_3 -algebra $\mathfrak{U}_{K_3 \times E}$ whose protected Pfaffian is the conjectural $\mathfrak{D}_X^{\text{DI}}$. The five-fold thesis of the manuscript -- one object, one chiral E_3 -algebra, five categorical-dimensional levels at which its value is Δ_5 -- meets the closed $K_3 \times E \rightarrow \text{pt}$ cobordism at level S, where the inverse of the squared value is the Laurent expansion $\Delta_5^{-2} = (\Phi_{10}^{\text{un}})^{-1}$ of the inverse Igusa cusp form, and the orientation information that distinguishes Δ_5 from $-\Delta_5$ lives in the Pfaffian datum at level Z. The closed-cobordism partition function reads as one because Δ_5 is the same section throughout: the Borcherds–Gritsenko–Nikulin section of the weight-5 automorphic line, the Borcherds–Kac–Moody denominator on the root lattice $\Lambda_{II}^{2,1}$, and (conditionally) the protected Pfaffian of the Dirac–Igusa pro-object on $K_3 \times E$. The squared

shadow of this same section is the protected partition function on the closed cobordism $K_3 \times E \rightarrow \text{pt}$; the Hall–Borcherds residual of Vol II, when constructed, will lift it to the gravity-line trace at level Z; the threefold realization question of §4 is the construction of the Dirac–Igusa pro-object that supplies the square-root datum.

BIBLIOGRAPHIC NOTES

The OP/DT scalar identity is Oberdieck–Pixton [?OB], Oberdieck–Pandharipande [?OPand], and Oberdieck [?OberdieckReducedPT]; the variable conventions follow Bryan [?BRYAN]. Theorem ?? is the consolidated form used here. The Andrianov factorization of the spinor L -function is [?Andr74]; the Saito–Kurokawa correspondence at weight 10 is [?ZagierSK]; the Manin–Deligne period theorem is [?Manin72], [?KZ], [?De179]. The dyon-counting interpretation in CHL backgrounds is Dijkgraaf–Verlinde–Verlinde [?DVV] and Borcherds [?B]; these inputs are the boundary-row physical hosts and are kept separate from the $K_3 \times E$ protected scalar shadow throughout.

The Hall–Borcherds residual conjecture in §3.2 is Vol II’s Construction Problem 2; its specific source location in Vol II is `conj : gravity-line-hall-borcherds-comparison on chapters/connections/3d_gravity.tex` at line 8429. The protected D-brane index conjecture $\Omega_X(h, d, n) \stackrel{\text{phys}}{=} \mathbf{DT}_{n, (\beta_h, d)}^X$ is recorded as Conjecture ?? of Chapter 5; it is not used in any of the proofs in this chapter. The Mukai dictionary, the K_3 -class chamber convention, and the $\Lambda_{II}^{2,1}$ lattice data are imported from Chapter 2. The protected Pfaffian Pf_{prot} and the Dirac–Igusa pro-object $\mathfrak{D}_X^{\text{DI}}$ are constructed in Chapter 5; the Pfaffian–Dirac theorem is Chapter 6; the four obstruction classes $(\text{Do}), (\text{O}_1), (\text{O}_1^+), (\text{O}_2)$ are defined and tracked in §5.8 of Chapter 5.

The κ -tuple $(0, 0, 3, 5, 24)$ and its non-additivity at $N = 1$ (the controlling instance for Δ_5) are recorded in Vol III; the Hall positive half $Y^+(K_3 \times E)$ and its Drinfeld doubling to $G(K_3 \times E)$, as well as the 6d holomorphic Chern–Simons quartic obstruction $\int_X \text{Tr}_{\text{ad}}(\mathcal{A}(F_A)^3)$, are also Vol III. The local model $\mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$ and the holomorphic de Rham obstruction class are mixed-HT-strings. None of these supply a 3D gravitational path integral, and the present chapter does not assert otherwise.

OPEN OBSTRUCTIONS AND THE NEXT FRONTIER

1 OPEN OBSTRUCTIONS AND THE NEXT FRONTIER

The cross-level identity ② ↔ ③ of this manuscript -- the Pfaffian–Dirac theorem $\mathrm{Pf}_{\mathrm{prot}}(\mathfrak{D}_X^{\mathrm{DI}}) = \Delta_5$, $\epsilon_o = \nu_{\Delta_5}$ on $\mathcal{W}^{(2)}(\Lambda_{II}^{2,1})$, and the primitive recognition $P_X \cong \mathfrak{g}_{\Delta_5}$ -- is conditional on four named obstructions, and the manuscript stops at the boundary where each is named, not at the boundary where each is closed. The view is honest: the automorphic identity ① ↔ ② is theorem [?Borcherds95, ?GN]; the singular-theta-lift identity ② ↔ ⑤ is theorem [?Borcherds95]; the BKM denominator computation $\mathrm{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1}\Delta_5(2Z)$ is theorem; what is conditional is the operator-level passage from the finite-stage compact $K_3 \times E$ source to the closed denominator algebra.

Each of the four obstructions sits at a definite categorical-dimensional level on the universal chain $P \rightarrow C \rightarrow S \rightarrow Z \rightarrow A$; each of three open frontiers -- the Hall–Borcherds residual promoting the level- n protected scalar shadow to operator level (Vol II Construction Problem 2); the chain fusion conjecture for compact non-toric $K_3 \times E$ (Vol III); and the gravity-line operator algebra whose Pentagon-face scalar trace is Δ_5^2 (Vol II Construction Problem 2 again, cross-listed) -- bridges the operator content of this manuscript to the broader programme. The Mathieu / umbral moonshine adjacency stands between obstructions and frontiers: its connection to $\mathfrak{g}_{\Delta_5}$ is conjectural, and the precise content of the conjecture is what the chapter asserts. The Beilinson cut places this chapter at the frontier: open obligations named precisely *are* the deliverable.

The three names of Δ_5 are kept distinct on first occurrence: the automorphic section $D_X = \Delta_5 \in H^0(\mathcal{H}_2, \mathcal{L}^1 \otimes \nu_{\Delta_5})$ of view ①; the BKM denominator $\mathrm{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1}\Delta_5(2Z)$ of view ②; and the compact protected Pfaffian $\mathrm{Pf}_{\mathrm{prot}}(\mathfrak{D}_X^{\mathrm{DI}})$ of view ③, conditional on (Do), (O1), (O1)⁺, (O2).

1.1 The four named obstructions

Each of the four obstructions sits at a definite categorical-dimensional level and carries a definite licensing type. The compact target is the Dirac–Igusa pro-object $\mathfrak{D}_X^{\mathrm{DI}} = \varprojlim_R (\mathcal{A}_{X,R}^{E_3}, F_{X,R}^{\mathrm{hyb}}, \Gamma_{X,R}, \Pi_X, \Phi_R, o_R, H_R, P_R^{\Pi}, C_{X,R}, \Theta_{\mathrm{Kos},R})$. The level discipline is *primitive object first, cross-level identity second, scalar shadow last*. The four obstructions are exactly the local-to-global passages whose Beilinson cut places the conditional theorem ② ↔ ③ at level Z rather than at level S.

1.1.1 (Do): DO-DEGENERATION LIMIT OF THE COSECTION-REDUCED D-CRITICAL ORIENTATION THEORY

1.1.1.1 Statement. The cosection-reduced d-critical orientation theory on the retained Hall stack $\mathcal{H}_R^{\mathrm{or}}$ of finite HN height R extends to the Do (zero-dimensional brane) degeneration locus, in the sense that the orientation line of the reduced determinant complex

$$\mathscr{D}_{R,C}^{\mathrm{red}} = \det \mathrm{RHom}_{\mathrm{red}}(C, C)$$

admits a Brav–Bussi–Dupont–Joyce–Szendrői [?BBDJS2015] square-root gerbe section over the closure $\overline{\mathfrak{M}_{R,C}^{\text{st}}} \subset \overline{\mathcal{H}_R^{\text{or}}}$ including the Do-degeneration stratum, compatible with the cosection contraction inherited from $K_3 \times E$. The retained Hall datum supplies an oriented critical finite-stage Pandharipande–Thomas/Gholampour–Sheshmani inverse system $\widehat{\mathcal{H}}_{\sigma,S}^{\text{or}} = \varprojlim_R F_R \mathcal{H}_{\sigma,S}^{\text{or}}$ on the finite HN quotients of Definition ??; (Do) asks that this system carry a coherent orientation through the Do-degeneration limit.

1.1.1.2 Stage and licensing. Level S, γ -ambient (chain-level versus pro-object), ε -effectiveness (orientation as square-root datum). The licensing is (S, γ, ε) : the d-critical structure lives at the chart specialization stage S; the ambient is the finite-stage pro-object $\widehat{\mathcal{H}}_{\sigma,S}^{\text{or}}$, not an ordinary chain complex; the effectiveness datum is the orientation gerbe section, which is what (Do) demands extend.

1.1.1.3 Swarm probe. *Status: partially discharged at the local level; open globally.* The swarm probes and their findings are recorded in notes/swarm_20260430/reports/: Ao05 verifies the local Pfaffian wall sign theorem is correctly conditional and identifies graph-isogeny atoms as candidate, not theorem; Ao11 maintains the firewall between the Maass character ν_{Δ_5} and the geometric orientation ε_o (these agree on $W^{(2)}(\Lambda_{JJ}^{2,1})$ only after (Do)–(O2) hold); Ao26 confirms BBDJS [?BBDJS2015] vanishing-cycle data require an oriented d-critical locus, not boundedness alone, and that Joyce–Upmeyer [?JoyceUpmeyer] square-root data on CY_3 moduli do not automatically extend to the reduced $K_3 \times E$ quotient; Ao37 separates the connected BE Borel/edge class $\alpha^{E,\text{free}} = a_1 u_1 + a_2 u_2 \in H^2(BE; \mathbb{F}_2)$ from finite $BE[N]$ Borel terms; Ao42 verifies the quotient-orientation descent across object, extension, mixed, wrapped, two-step flag, overlap, Thom–Sebastiani, and $R' \rightarrow R$ transition strata is conditional, with overlap/correspondence/flag/transition compatibilities named separately.

1.1.1.4 Reasoning. The retained Hall boundedness [?LiuProductStability, ?PiyaratneTodaModuli3folds] gives a finite-type retained slice; it does not give quasi-smoothness, a d-critical chart, BBDJS coefficients, an orientation line, or compact-support six-functor admissibility through the Do limit. Pantev–Toën–Vaquié–Vezzosi [?PTVV2013] shifted symplectic structures cover ambient derived moduli under Calabi–Yau hypotheses; the retained atlas is not produced by PTVV alone. Joyce–Upmeyer [?JoyceUpmeyer] square-root orientation on Calabi–Yau₃ moduli requires exact-sequence/direct-sum compatibility; the reduced $K_3 \times E$ -quotient inherits this only after the cosection-induced shift is shown compatible with descent.

1.1.1.5 Next concrete construction. *Open.* Compute $\alpha_{R,C}^{\text{red}} = w_2(\mathcal{D}_{R,C}^{\text{red}}) \in H^2(S; \mathbb{F}_2)$, $\alpha_{R,C}^{E,\text{free}}$, $\widetilde{\beta}_{R,C}^H \in H_H^2(S; \mathbb{F}_2)$, and $\lambda_{R,C}^H \in H^1(BH; \mathbb{F}_2)$ on every retained object, extension, mixed, wrapped, and flag stratum C , with killing of the Borel mixed terms and edge-spectral-sequence differentials required at every node of the descent diagram. The simplest decisive locus is the rank-one D6–D2–Do stratum $C = (1, \beta, n)$ with β the K_3 -fibre primitive class: on this stratum the BBDJS line carries the cosection contraction explicitly, and the four obstruction classes $(\alpha^{\text{red}}, \alpha^{E,\text{free}}, \widetilde{\beta}^H, \lambda^H)$ reduce to a Pin^c computation. A discharge of (Do) on this stratum, together with the extension/correspondence/flag transitions of probe Ao42, yields the strong reduced orientation across the retained inverse system.

1.1.2 (O₁): STRONG REDUCED ORIENTATION ON THE $K_3 \times E$ -QUOTIENT SELF-EXT FRAME BUNDLE

1.1.2.1 Statement. The BBDJS [?BBDJS2015] orientation gerbe on the moduli of stable sheaves on $K_3 \times E$ descends to the $K_3 \times E$ -quotient self-Ext frame bundle, with reduced orientation gauge: on the stable stratum $\mathfrak{M}_{R,C}^{\text{st}} \subset \mathcal{H}_R^{\text{or}}$ the obstruction tuple

$$\mathfrak{o}_{R,C}^{\text{or}} = (\alpha_{R,C}^{\text{red}}, \alpha_{R,C}^{E,\text{free}}, \{(\tilde{\beta}_{R,C}^H, \lambda_{R,C}^H)\}_H, \mu_{R,e}^{\text{or}}, \dots)$$

vanishes simultaneously, with all overlap, correspondence, flag, and $R' \rightarrow R$ transition data trivialized coherently. (O₁) demands the strong reduced descent: the connected- BE class $\alpha^{E,\text{free}}$, the finite- $BE[N]$ gerbe classes $\tilde{\beta}^H$, and the residual finite linearization characters λ^H all vanish, separately, on every retained stratum, and the spectral-sequence differentials and Borel mixed terms are killed.

1.1.2.2 Stage and licensing. Level $S \rightarrow Z$, β -comparison (gerbe descent functor), ε -effectiveness (orientation gauge non-degeneracy). The licensing is $(S \rightarrow Z, \beta, \varepsilon)$: the descent of the BBDJS line is the comparison arrow from the chart specialization stage S to the centre stage Z , and effectiveness asks the orientation be a non-degenerate square-root datum.

1.1.2.3 Swarm probe. *Status: partially discharged on local charts; open globally.* Probe Ao74 confirms cross-repo consistency with Vol III on the primitive recognition firewall: target arithmetic on $W_{\leq 3}$, is target Borchers–Gritsenko–Nikulin–Kac data only, and recognition requires an actual compact source datum supplying $M, D, B, G, K, Q, A_\beta$, relation rows, Hopf radical/coideal kernel equality, PBW associated-graded comparison, and strict transition/Mittag–Leffler rows. Probes Ao76, Ao82, Ao89 supply the proof-ledger statement defining what a discharged (O₁) carries to the recognition theorem. The connected E and finite $E[N]$ cohomologies are *distinct* obstruction spaces: $H^*(BE; \mathbb{F}_2) = \mathbb{F}_2[u_1, u_2]$ with $|u_i| = 2$, so $H^1(BE; \mathbb{F}_2) = 0$; for $E[2] \cong (\mathbb{Z}/2)^2$, $H^*(BE[2]; \mathbb{F}_2) = \mathbb{F}_2[x_1, x_2]$ with $|x_i| = 1$, and the residual linearization $\lambda = \lambda_1 x_1 + \lambda_2 x_2$ is independent data after the degree-two gerbe class is killed (Ao37, Ao42).

1.1.2.4 Reasoning. Translation invariance under E -action does not produce orientation descent: a fixed underlying line may carry a nontrivial finite character, and ordinary translation invariance is *not* a substitute for vanishing of λ^H on full two-primary generators of $H^*(B(\mathbb{Z}/2^a)^2; \mathbb{F}_2) = \Lambda(x_1, x_2) \otimes \mathbb{F}_2[y_1, y_2]$ at $N = 2^a$. The quotient-orientation criterion is finite Čech–Borel descent on the retained groupoid, not a vanishing theorem.

1.1.2.5 Next concrete construction. *Open.* Construct, on the rank-one D6–D2–Do stratum and its extension, mixed, wrapped, and two-step flag completions, the actual BBDJS gerbe sections on the self-Ext frame bundle, and verify the full vanishing of $\mathfrak{o}_{R,C}^{\text{or}}$ entry-by-entry. The decisive locus is the rank-one D6–D2–Do input $C = (1, \beta, n)$ with the four cohomology classes computed explicitly via the Pin^c input plus quotient checks of clause (P₁); for a discharged (O₁), the overlap and correspondence transitions of probe Ao42 must commute with descent on the cofinal HN system.

1.1.3 (O₁)⁺: WEYL-EQUIVARIANT TRANSPORT OF THE ORIENTATION ALONG $W^{(2)}(\Lambda_{II}^{2,1})$ REFLECTIONS

1.1.3.1 Statement. The orientation character on the type-II Weyl group $W^{(2)}(\Lambda_{II}^{2,1})$ is Weyl-equivariant: the orientation line of (O₁) is equipped with coherent lifts $\tau_i : o_C \rightarrow s_{\delta_i}^* o_C$ of the

three type-II simple reflections, each acting with sign -1 , and the projective determinant-line cocycle $c_o \in Z^2(W^{(2)}(\Lambda_{II}^{2,1}); \mathbb{F}_2)$ of the lift is a coboundary, with quotient torsor defects $\omega_{i,C} = 0$ on every retained groupoid component. At finite height R , the partial action groupoid $\mathcal{G}_R$ carries $c_{o,R} \in Z^2(\mathcal{G}_R; \mathbb{F}_2)$, and $(O1)^+$ asks that $[c_{o,R}] = 0$, $\tau_i^2 = 1$, and $\omega_{i,C} = 0$ hold coherently on every stratum and at every transition height.

1.1.3.2 Stage and licensing. Level Z , β -comparison (Weyl transport functor), δ -endpoint (cocycle reduction to group cohomology on the full $W^{(2)}(\Lambda_{II}^{2,1})$ -orbit). The licensing is (Z, β, δ) : the Weyl transport is the comparison arrow at the centre stage; the endpoint discipline is the reduction of the finite groupoid cocycle to the group cohomology class on the complete orbit.

1.1.3.3 Swarm probe. *Status: open; group-theoretic frame fixed.* Probe Ao47 records the local sign-extraction firewall: only after the unit character is killed and $N_\delta^{\text{Pf}} = 1$ is computed from the reduced normal self-Ext chart does the local sign equal -1 , and Maass values $\nu_{\Delta, (s_{\delta_i})} = -1$ plus divisor orders $d_{\delta_i}^{\text{aut}} = f(0, \pm 1) = 1$ supply target data only -- they do not compute N_δ^{Pf} , χ_ν , or ϵ_o . Probe Ao49 confirms that the presentation

$$W^{(2)}(\Lambda_{II}^{2,1}) = \langle s_{\delta_1}, s_{\delta_2}, s_{\delta_3} \mid s_{\delta_i}^2 = 1 \rangle$$

implies that any orientation character on $W^{(2)}(\Lambda_{II}^{2,1})$ is determined by three generator signs once $(O1)^+$ gives the honest Weyl lift; the three signs proven equal to -1 by $(O2)$ then propagate to the full character. Probe Ao89 places the Weyl-equivariant transport in the proof ledger as a separate identity, distinct from the no-extra-relations and PBW identities.

1.1.3.4 Reasoning. Three local signs on the simple reflections imply a character on the full type-II Weyl group only after $[c_{o,R}] = 0$ and $\omega_{i,C} = 0$: the Weyl determinant-line cocycle measures whether the lift is honest, and the quotient torsor defects measure whether the descent commutes with the reflection action on the retained groupoid. The cocycle reduces to the classical group-cohomology class only on a complete $W^{(2)}(\Lambda_{II}^{2,1})$ -orbit with constant coefficients; finite height R is a partial action groupoid whose component data must be glued.

1.1.3.5 Next concrete construction. *Open.* Construct the finite-height projective lifts $\tau_{i,R}$, prove $\tau_{i,R}^2 = 1$ (no normalization choice survives squaring), and compute the cocycle class $[c_{o,R}]$ and torsor defects $\omega_{i,C}$ in $H^2(\mathcal{G}_R; \mathbb{F}_2)$ on every retained groupoid component, killing them coherently as $R \rightarrow \infty$. The decisive computation is on the $2\delta_i + \delta_j$ and δ_{123} words of the first window $W_{\leq 3}$: if the lifts trivialize on the height-three orbit, $(O1)^+$ discharges on $W_{\leq 3}$; the $W_{\leq 7}$ extension via height-four, doubled isotropic, and complementary string rows is then the next stage.

1.1.4 (O2): LOCAL PFAFFIAN WALL NORMAL FORM ON TYPE-II WALLS

1.1.4.1 Statement. On the three type-II walls $\mathbb{H}_{\delta_i}$ for $\delta_1 = 2f_2 - f_3$, $\delta_2 = 2f_{-2} - f_3$, $\delta_3 = f_3$, the Pfaffian sign sum on the half-Hilbert orbit is exactly $4096 = 2^{12}$ on the wall, by a local computation that Appendix E carries in full. Concretely, in an s_δ -equivariant real-analytic d-critical/Kuranishi chart $(U_\delta, x_\delta, u_\delta, K_\delta^{\text{red}})$, with K_δ^{red} the real form of the cosection-reduced self-Ext complex $\text{RHom}_{\text{red}}(C_\delta, C_\delta)$, the splitting $K_\delta^{\text{red}} \simeq K_\delta^{\text{tan}} \oplus K_\delta^{\text{nor}}$ is supplied with the extension class $\eta_\delta \in \text{Ext}_{D^{s_\delta}(U_\delta)}^1(K_\delta^{\text{nor}}, K_\delta^{\text{tan}})$ trivial after the reduced E -quotient and quotient-orientation trivializations, and the normal Pfaffian admits the rank-one form $\text{Pf}(K_\delta^{\text{nor}}) = \nu_\delta u_\delta^{N_\delta^{\text{Pf}}}$ with $\nu_\delta(x_\delta) \neq 0$, $s_\delta(\nu_\delta) = \nu_\delta$, and $N_\delta^{\text{Pf}} = 1$.

1.1.4.2 The 4096 sign sum. The integer $4096 = 2^{12}$ is the half-Hilbert sign sum across the three walls of the cell at the cusp: twelve sign degrees of freedom, each $\mathbb{Z}/2$, give $2^{12} = 4096$ signed states, and the sign sum on the type-II wall locus is exactly this full count under (L1)–(L3) of Proposition ?? . The arithmetic match $-4096 = -(64)^2$ with the OP/DT scalar $Z_{\text{OP}}^X = -4096 \Delta_5^{-2}$ is the squared-theta normalization, fixed by Borchers normalization and the OP scalar sign convention; it is *not* a derivation of ϵ_o .

1.1.4.3 Stage and licensing. Level C, α -chart (real-analytic d-critical/Kuranishi chart), ε -effectiveness (rank-one normal Pfaffian non-degeneracy). The licensing is (C, α, ε) : the chart specialization is the local model at the wall; effectiveness is the rank-one normal form.

1.1.4.4 Swarm probe. *Status: open; local model fixed up to discharged unit and rank computations.* Probe Ao47 fixes the local model and the exact firewall against scalar leakage from OP and Maass: scalar constants $D_5 = 64^{-1} \Delta_5$, $\chi_{10}^{\text{OP}} = 4096^{-1} \Delta_5^2$, $Z_{\text{OP}}^X = -4096 \Delta_5^{-2}$ are scalar normalizations and do not define ϵ_o ; the Pfaffian normal form is the content. Probes Ao92 and A100 specify the discharge content for (O2): matrices over $\mathbb{Z}$ or $\mathbb{Q}$ computing the Pfaffian-rank identity $N_{\delta}^{\text{Pf}} = 1$, the splitting $\eta_{\delta} = 0$, the unit ν_{δ} with s_{δ} -character trivial, and overlap/transition compatibility across the cofinal HN system.

1.1.4.5 Reasoning. The condition $(\partial, \partial) = 2$, the automorphic divisor order $d_{\delta_i}^{\text{aut}} = 1$, and the Maass character $\nu_{\Delta_5}(s_{\delta_i}) = -1$ are target data that do not imply the Pfaffian rank assertion $N_{\delta}^{\text{Pf}} = 1$. Graph-isogeny atom candidates $B_2 = i_{\Gamma_{\varphi},*} L$ are Koszul-Ext-local; they are *candidates*, not constructions, until semistability, full $\widehat{\Gamma}_X$ charge match, reduced normal Ext splitting, invariant unit, and atlas overlap compatibility are supplied. The “higher terms do not alter” phrase in the rank-one Ext criterion requires the explicit equivariant real/parametric Morse lemma and unit check before it operates as a theorem.

1.1.4.6 Next concrete construction. *Open.* Construct retained wall objects for $\delta_1, \delta_2, \delta_3$ with Liu-heart membership [$? \text{LiuProductStability}$], semistability, exact wall equality, full $\widehat{\Gamma}_X$ -charge matching; for δ_2 , supply the wrapped/mixed representative with $b_R^{\text{geom}} > 0$; on each wall, supply the equivariant d-critical chart with $s_{\delta}(u_{\delta}) = -u_{\delta}$, the reduced self-Ext complex, the tangent/normal splitting with vanishing extension class, the invariant Pfaffian unit, and the rank-one normal quasi-isomorphism. The decisive computation is the discharge of Proposition ?? (L1)–(L3) on δ_2 , where the wrapped-middle representative is the hardest of the three; the corresponding Appendix E entry then carries the full 2^{12} sign sum.

1.2 Mathieu and umbral moonshine adjacency

The K_3 elliptic genus has a precise representation-theoretic decomposition. Eguchi, Ooguri, and Tachikawa [$? \text{EOT}$] observed in 2010 that the elliptic genus $Z_{K_3}(\tau, z) = 2\phi_{0,1}(\tau, z)$ admits an M_{24} -graded character expansion: the mock-modular completion $\Sigma(\tau)$ of the elliptic genus is a graded M_{24} -module, and the multiplicities of irreducible M_{24} -representations in the first several Fourier coefficients of the holomorphic part match the dimensions of irreducible M_{24} -representations on the nose. *Status: proved as an arithmetic identity by Gannon [$? \text{GannonUmbral}$]; the geometric origin via a sigma-model action of M_{24} on the K_3 BPS Hilbert space remains open for the non-surfing classes $\{7A, 7B, 15A, 15B\}$. The M_{24} action visible at the level of the elliptic genus does not lift to a global geometric symmetry of $D^b(\text{Coh}(K_3))$; the work of Gaberdiel–Hohenegger–Volpato [$? \text{GHV}$] on*

Mathieu twining characters identifies the precise refinement at the level of supersymmetric sigma-model conformal field theory data.

The umbral moonshine of Cheng–Duncan–Harvey [?CDH] extends Mathieu moonshine to a family of 23 cases indexed by Niemeier root systems, one case per Niemeier lattice having a non-empty root system. The 23 Niemeier lattices with non-empty root systems pair each Niemeier ADE root system with a graded module for the corresponding discriminant group; Mathieu moonshine is the root system $24A_1$ -case, which corresponds to the Niemeier lattice with root system A_1^{24} . *The count is 23 umbral cases plus the Leech lattice as the 24th Niemeier with empty root system: the Leech lattice itself does not enter the umbral catalogue but supplies the Conway moonshine group, distinct from the umbral construction.* The umbral moonshine conjecture, raised by Cheng–Duncan–Harvey [?CDH] and verified module-theoretically by Gannon [?GannonUmbral], asserts that each of the 23 umbral cases carries a graded module whose graded character is a specified vector-valued mock modular form.

1.2.0.1 The conjectural connection to $\mathfrak{g}_{\Delta_5}$. Status: conjectured. The Mathieu moonshine module is conjecturally a graded character of $\mathfrak{g}_{\Delta_5}$ or a closely related Borcherds–Kac–Moody algebra. The precise statement that survives the Beilinson cut is the following: there exists a $\mathbb{Z}$ -bigraded M_{24} -module $M_{\bullet,\bullet}^{\text{moon}}$, with $M_{n,l}^{\text{moon}}$ the Eguchi–Ooguri–Tachikawa M_{24} -module in degree (n, l) , whose graded character matches the K3 elliptic genus, and whose vertex/Borcherds extension to the Borcherds–Kac–Moody datum on $\Lambda_{II}^{2,1}$ is conjecturally a g -twined refinement of $\mathfrak{g}_{\Delta_5}$ producing the twined denominator $\text{den}(\mathfrak{g}_{\Delta_5}^{(g)})$ for each M_{24} -conjugacy class g . The umbral cases of [?CDH, ?ChengHarrison] extend the conjecture to a family of 23 Borcherds–Kac–Moody-style algebras associated to the 23 Niemeier root systems with non-empty root sets.

1.2.0.2 What the conjecture is and what it is not. The conjectural connection is of view ② (BKM denominator) to Mathieu/umbral moonshine, not of view ③ (compact Pfaffian). The compact protected Pfaffian $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}})$ is conditional on (Do)–(O₂); the moonshine adjacency is independent of this conditioning. Equivalently, the moonshine adjacency lives at level Z (centre/denominator) and at level A (acting object on the K3 BPS Hilbert space at the level of representation-graded characters of mock modular forms); the operator-level connection to view ③ is open, and a proven identification of the moonshine module as a g -twined character of $\mathfrak{g}_{\Delta_5}$ does not by itself discharge any of (Do), (O₁), (O₁)⁺, (O₂).

1.2.0.3 Stage and licensing. Level $Z \rightarrow A, \beta$ -comparison (g -twining functor), γ -ambient ($\mathbb{Z}$ -bigraded module versus mock modular form). The licensing is $(Z \rightarrow A, \beta, \gamma)$: the g -twining operation is the comparison arrow that produces $\text{den}(\mathfrak{g}_{\Delta_5}^{(g)})$ from $\text{den}(\mathfrak{g}_{\Delta_5})$ and the M_{24} -action; the ambient is the bigraded M_{24} -module category, not the ordinary graded category.

1.2.0.4 Cross-repo consistency. The Vol III FRONTIER (file FRONTIER.md) carries the dimension-stratified BKM catalogue in which $\mathfrak{g}_{\Delta_5}$ is the $d = 3$ sibling, with companions at $d = 3$ (Borcherds Monster $V^{\natural}$ on $\text{II}_{1,1}$) and $d = 5$ (Fake Monster on $\text{II}_{25,1}$ via $K_3 \times K_3 \times E$) by frontier row F6 (the $d \geq 4$ stratum). The Mathieu/umbral moonshine adjacency to $\mathfrak{g}_{\Delta_5}$ is a level-Z cross-stratum conjecture: the geometric M_{24} sigma-model action on K3 for the non-surfing classes $\{7A, 7B, 15A, 15B\}$ is the missing piece on the K3-fibre side; the g -twined denominator $\text{den}(\mathfrak{g}_{\Delta_5}^{(g)})$ is the missing piece on the BKM side.

1.2.0.5 Next concrete construction. Open. Compute the twined denominators $\text{den}(\mathfrak{g}_{\Delta_5}^{(g)})$ for each of the 26 conjugacy classes of M_{24} , using the g -twined Borchers product on $\Lambda_{II}^{2,1}$, and verify that the resulting graded characters match the EOT M_{24} -multiplicity predictions through the first eight Fourier coefficients of $\Sigma(\tau)$. The decisive cross-check is the consistency of the g -twined denominator with the OP scalar $Z_{\text{OP}}^X = -4096 \Delta_5^{-2}$ under the g -twining construction, including the squared sign convention.

1.3 The next frontier

Three open frontiers connect this manuscript to the broader programme. Each is named in cross-volume notes; each requires construction the manuscript does not undertake. The Beilinson cut places each frontier at a definite level, and the licensing tags below identify the comparison arrows whose construction is open.

1.3.1 FRONTIER I: THE HALL–BORCHERDS RESIDUAL

1.3.1.1 Statement. The level- n protected scalar shadow $Z_{\text{BPS}}^{K_3 \times E} = (\Phi_{10}^{\text{un}})^{-1} = \Delta_5^{-2}$ on the closed $K_3 \times E \mapsto$ point cobordism is the BPS partition function. Its promotion to a 3d gravitational path integral with a genuinely operator-level construction requires the Hall–Borchers residual: an operator-valued completion of the scalar trace whose trace recovers the level- n shadow and whose operator-level algebraic content carries the dynamical metric path integral. *Status: open; this is Vol II Construction Problem 2.*

1.3.1.2 Stage and licensing. Level $Z \rightarrow A$, β -comparison (residual operator algebra), δ -endpoint (promotion to dynamical metric beyond the holographic HT sector), ε -effectiveness (operator non-degeneracy). The licensing is $(Z \rightarrow A, \beta, \delta, \varepsilon)$: the residual is the comparison arrow from the centre to the acting object; the endpoint discipline is the promotion beyond the holographic Hochschild–Tate boundary-CFT reading of pure 3d gravity.

1.3.1.3 Cross-repo anchor. Vol II chapters/connections/3d_gravity.tex:8429 names the gravity-line operator algebra whose Pentagon-face scalar trace is $\Phi_{10}^{\text{un}} = \Delta_5^2$, and identifies the Hall–Borchers residual as the comparison arrow from the bulk $Z_{\text{ch}}^{\text{der}}(A_b)$ to the line-operator algebra. The Vol II Construction Problem 2 reads: *construct the gravity-line operator algebra with Pentagon-face scalar trace $\text{tr}_{\text{Pentagon}} = \omega_{\text{Borchers}} = c_N(o)/2$ at $N = 1$ on the K_3 -Heisenberg witness, recovering the Borchers form $\Phi_{10}^{\text{un}} = \Delta_5^2$ on the closed cobordism.* The igusa-side input is the same scalar identity $\Delta_5^2 = \Phi_{10}^\theta$, with the manuscript-conditional operator-level package $\mathfrak{D}_X^{\text{DI}}$ supplying the Pfaffian/orientation/Weyl-transport data on the chiral side.

1.3.1.4 Reasoning. The protected scalar trace $\text{Tr}_{\text{prot}}((-1)^F (\mathfrak{D}_X^{\text{DI}})^{-2}) = -4096 \Delta_5^{-2}$ is squared away from orientation: the sign-character ϵ_o is invisible in the squared form. The Hall–Borchers residual recovers the orientation by lifting from the squared scalar to the unsquared Pfaffian section, with the Weyl-character data providing the extra structure that the squared scalar cannot recover. The promotion to a 3d gravitational path integral is conditional on Brown–Henneaux, modular invariance, vacuum dominance, and saddle dominance [?HM]; the manuscript does not claim this promotion.

1.3.1.5 Next concrete construction. Open. Construct the Pentagon-face cyclic trace $\text{Tr}_{\text{Pentagon}}^{\text{cl}}$ on the closed-colour factorization algebra F^{cl} at the $K_3 \times E$ -Heisenberg witness, prove that its evaluation recovers $c_1(o)/2 = 5$ on the universal Borchers-weight identity, and identify its operator-level lift to the

open-colour boundary algebra $A_b = \text{End}_C(b)$ via the chain fusion conjecture (Frontier 2 below). The decisive cross-check is the consistency of the Pentagon-face scalar with the OP/DT scalar $-4096 \Delta_5^{-2}$ under the squared-trace operation.

1.3.2 FRONTIER 2: CHAIN FUSION FOR COMPACT NON-TORIC $K_3 \times E$

1.3.2.1 Statement. For each Calabi–Yau $_d$ category C_X and each chart datum (Σ_{d-1}, C) , the Stage-2 chiral algebra $A_X = \Phi_d^{(\Sigma_{d-1}, C)}(C_X)$ coincides with the boundary algebra $A_{b(X, \Sigma, C)} = \text{End}_C(b(X, \Sigma, C))$ for a canonical boundary vacuum $b(X, \Sigma, C)$ in an open factorization dg-category on (C, D_C, τ_C) , where D_C encodes the Calabi–Yau data’s special points (orbifold loci, fibration punctures, conifold singularities). *Status: conjectured; verified at toric loci ($\mathbb{C}^3$, local $\mathbb{P}^2$, conifold) and at $K_3 \times E$ at $d = 3$; compact non-toric general X is open.*

1.3.2.2 Stage and licensing. Level $P \rightarrow C \rightarrow S$, β -comparison (chain-fusion functor identifying Stage-2 output with boundary algebra), α -chart (canonical boundary vacuum $b(X, \Sigma, C)$), γ -ambient (open factorization dg-category on the log curve (C, D_C, τ_C)). The licensing is $(P \rightarrow S, \alpha, \beta, \gamma)$: the chain fusion is the comparison arrow from the primitive open factorization category to the chart-specialized chiral shadow; the chart datum is the canonical boundary vacuum; the ambient is the log-curve open factorization dg-category, not an ordinary chiral algebra on a smooth curve.

1.3.2.3 Cross-repo anchor. The Vol III FRONTIER row F1 (chain fusion conjecture in general d) names this conjecture in the two-stage factorization frame; row F2 ($G(X) = D(Y^+(X))$ for compact non-toric CY_3) names the level-Z Drinfeld double obligation downstream of chain fusion; row F10 ($CY\text{-}A_3$ chain-level compact closure) names the conditional TCFT/framing/anomaly/completion package needed for compact non-formal CY_3 in general. The Vol III adversarial review note `notes/chatgpt_critique_consequence_map_adversarial_review.md` §III records chain fusion as a programme-level conjecture, not a theorem. The Vol II CLAUDE.md banned row II names “ $\Delta_5 = \text{Hilbert space}$ ” as collapse; the corrected statement is the conditional $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}}) = \Delta_5$ of view ③.

1.3.2.4 Reasoning. The compact $K_3 \times E$ case at $d = 3$ is a witnessed instance of chain fusion via the rectified finite K_3 -to- E specialization (Definition ?? in this manuscript). The chiral algebra $\text{Sp}_{K_3, E}^{\text{ch}}(\Phi_3^{\text{FA}}(K_3 \times E)) = U^{\text{ch}}(\mathfrak{heis}_{\text{Muk}}) \otimes U^{\text{ch}}(\mathfrak{g}_{K_3}^{\text{BPS}})$ on $\text{Ran}(E)$ is the Stage-2 specialization; the boundary algebra $A_b = \text{End}_C(b)$ on the open factorization side coincides with this on the verified locus, but the canonical boundary vacuum $b(K_3 \times E, \Sigma_2, E)$ is not in the literature. Compact non-toric $K_3 \times E$ generalizations to other compact non-toric CY_3 -data (e.g. Borcea–Voisin threefolds, Schoen’s small resolutions) require independent construction.

1.3.2.5 Next concrete construction. *Open.* Construct $b(K_3 \times E, \Sigma_2, E)$: a canonical boundary vacuum in an open factorization dg-category on (E, D_E, τ_E) with D_E at the Igusa cusp $\mathfrak{c}_\infty$ of the Siegel space, and prove the chain-fusion isomorphism on $W_{\leq 3}$ at finite HN height $R = 3$, identifying the Stage-2 chiral output with $A_{b(K_3 \times E, \Sigma_2, E)}$. The decisive cross-check is the consistency of the chain-fusion isomorphism with the OP/DT scalar branch and with the universal Borchers-weight identity $\kappa_{\text{BKM}}(\Phi_1) = c_1(o)/2 = 5$; see the κ -tuple discussion below.

1.3.3 FRONTIER 3: THE GRAVITY-LINE OPERATOR ALGEBRA

1.3.3.1 Statement. The gravity-line operator algebra is an operator algebra on the closed-colour factorization algebra F^{cl} on $K_3 \times E$ whose Pentagon-face scalar trace is $\Delta_5^2 = \Phi_{10}^{\text{un}}$. The Pentagon-face trace is the trace lane of the universal stage chain $P \rightarrow C \rightarrow S \rightarrow Z \rightarrow A$ that, by the universal Borchers-weight identity $\text{tr}_{\text{ghost}}(Q_{\text{BRST}}^2) = \text{tr}_{\text{Pentagon}} = \omega_{\text{Borchers}} = c_N(o)/2$, takes the value $c_1(o)/2 = 5$ at the Gritsenko–Cléry row producing Δ_5 . *Status: open; this is Vol II Construction Problem 2 (cross-listed with Frontier 1).*

1.3.3.2 Stage and licensing. Level $Z \rightarrow A$, β -comparison (operator algebra functor at Pentagon face), δ -endpoint (Pentagon-face trace evaluation at the Δ_5 row), ε -effectiveness (operator non-degeneracy at the K_3 -Heisenberg witness). The licensing is $(Z \rightarrow A, \beta, \delta, \varepsilon)$.

1.3.3.3 Cross-repo anchor. Vol II chapters/connections/3d_gravity.tex:8429 names the Gravity-line Hall–Borchers comparison conjecture and the Pentagon-face scalar trace; Vol II CLAUDE.md §9 lists this as Construction Problem 2; Vol III CLAUDE.md names $\kappa_{\text{BKM}}(\Phi_N) = c_N(o)/2$ as the universal Borchers-weight identity [?Borchers95], with $\kappa_{\text{BKM}}(\Phi_1) = c_1(o)/2 = 5$ at the paramodular Δ_5 input.

1.3.3.4 The κ -tuple. The five-fold κ -tuple $(\kappa_{\text{cat}}, \kappa_{\text{ch}}^{\text{Hodge}}, \kappa_{\text{ch}}^{\text{Heis}}, \kappa_{\text{BKM}}, \kappa_{\text{fiber}}) = (o, o, 3, 5, 24)$ at $K_3 \times E$ records five separate invariants, distinct in construction and meaning, never additively combined. Here $\kappa_{\text{cat}} = \chi(\mathcal{O}_{K_3 \times E}) = \chi(\mathcal{O}_{K_3}) \cdot \chi(\mathcal{O}_E) = 2 \cdot o = o$ is the Künneth-multiplicative categorical Euler; $\kappa_{\text{ch}}^{\text{Hodge}} = \sum_q (-1)^q h^{o,q}(K_3 \times E) = o$ by Serre duality on the compact total space; $\kappa_{\text{ch}}^{\text{Heis}} = 3$ is the rank of the Heisenberg–Cartan in the Stage-2 specialization on $\text{Ran}(E)$ and equals the Cartan rank of $\mathfrak{g}_{\Delta_5}$; $\kappa_{\text{BKM}} = c_1(o)/2 = 5$ is the universal Borchers-weight identity at the Igusa row $\Phi_1 = \Delta_5$ of the Gritsenko–Cléry 8-form catalogue with $c_1(o) = 10$; $\kappa_{\text{fiber}} = 24$ is the Mukai-lattice rank $\text{rk } \tilde{H}^*(K_3, \mathbb{Z}) = 24$ of the K_3 fibre, equal to $\chi_{\text{top}}(K_3)$. The collapsed identity $\kappa_{\text{BKM}} = \kappa_{\text{ch}} + \chi(\mathcal{O}_{\text{fiber}})$ fails because it confuses $\chi_{\text{top}}(K_3) = 24$ with $\chi(\mathcal{O}_{K_3}) = 2$; the universal identity $\kappa_{\text{BKM}}(\Phi_N) = c_N(o)/2$ is the correct frame.

1.3.3.5 Reasoning. The Pentagon-face scalar trace is the modularity-witness of the open-side cyclic trace plus clutching; modularity is a property of the open trace, not of the closed algebra. The trace-lane equation $\text{tr}_{\text{ghost}}(Q_{\text{BRST}}^2) = \text{tr}_{\text{Pentagon}} = \omega_{\text{Borchers}} = c_N(o)/2$ identifies the Pentagon-face trace with the Borchers form weight under the BRST chiral nilpotent on the holographic factor; on the Δ_5 row, $c_1(o)/2 = 5$ is the level-five weight, and $\Delta_5^2 = \Phi_{10}^{\theta}$ is the squared form of total weight 10 appearing as the closed-cobordism BPS partition-function input.

1.3.3.6 Next concrete construction. *Open.* Construct the gravity-line operator algebra $\text{GravLine}_{K_3 \times E}$ on the open factorization dg-category on (E, D_E, τ_E) at the Igusa cusp, prove the Pentagon-face scalar trace identity $\text{tr}_{\text{Pentagon}}^{\text{cl}}(\text{GravLine}_{K_3 \times E}) = \Delta_5^2$, and verify consistency with the universal Borchers-weight identity $\kappa_{\text{BKM}}(\Phi_1) = 5$ and with the κ -tuple $(o, o, 3, 5, 24)$ at the K_3 -Heisenberg witness. The decisive cross-check is the discharge of the chain fusion (Frontier 2) at $b(K_3 \times E, \Sigma_2, E)$ and the construction of the Hall–Borchers residual (Frontier 1) lifting the squared scalar $-4096 \Delta_5^{-2}$ to the unsquared Pfaffian section $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}}) = \Delta_5$ in this manuscript.

The mathematical work is the discharge of (Do)–(O2) on $W_{\leq 3}$ and $W_{\leq 7}$, followed by the construction of the canonical boundary vacuum $b(K_3 \times E, \Sigma_2, E)$ of Frontier 2, the Hall–Borchers residual

of Frontier 1 lifting the squared scalar $-4096 \Delta_5^{-2}$ to the unsquared Pfaffian $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}}) = \Delta_5$, and the verification of the Mathieu-twined denominators $\text{den}(\mathfrak{g}_{\Delta_5}^{(g)})$ on the 26 M_{24} -conjugacy classes. Each of the four obstructions is at a definite categorical-dimensional level: (Do) at S, (O1) at $S \rightarrow Z$, (O1)⁺ at Z, (O2) at C; each of the three frontiers, at $Z \rightarrow A$, $P \rightarrow S$, and the Pentagon face. The Beilinson cut places naming as deliverable; the deliverable is named.

1 LATTICE COMPUTATIONS

The seven integral lattices used in the manuscript are recorded here in one place: the hyperbolic plane $U = \Pi_{1,1}$, the Lorentzian root lattice $\Lambda_{II}^{2,1}$ of the Borcherds–Kac–Moody algebra $\mathfrak{g}_{\Delta_5}$, the K3 cohomology lattice $\Lambda_{K3} = \Pi_{3,19}$, the lattice-polarized K3 sublattice Λ_S , the Mukai lattice $\tilde{H}(S, \mathbb{Z}) \simeq \Pi_{4,20}$, the formal dyonic charge lattice $\Gamma_X^{\text{form}} = \Lambda_S \oplus \Lambda_S$, and the signature- $(3, 2)$ lattice $\Lambda^{3,2}$ that bridges $\mathbf{Sp}_4(\mathbb{Z})$ to $\mathbf{SO}_+(3, 2)$ via $\mathbf{PSp}_4(\mathbb{Z}) \simeq \mathbf{SO}_+(\Lambda^{3,2})$. Each is given by its Gram matrix or root data, signature, discriminant, and automorphism group, with primary citations (Conway–Sloane 1999 §2.5, §4, §16; Nikulin 1980 Theorem 1.1.2 and Theorem 1.14.4; Mukai 1984; Gritsenko–Nikulin 1998 Part II §2, §5). Ranks, signatures, and the Cartan matrix of the type-II simple roots are verified against the rational computation `compute/verify_lattice.py`.

This appendix is upstream of every chapter: load-bearing identities about Δ_5 , $\mathfrak{g}_{\Delta_5}$, the orientation character ν_{Δ_5} , and the Pfaffian wall normal form on $\mathcal{W}^{(2)}(\Lambda_{II}^{2,1})$ refer back to the lattice data fixed here, not to a sign or scaling chosen in the body.

Convention 1.1 (Sign of the bilinear form; orientation of the time-like cone). The standard mathematical convention for the hyperbolic plane U is $U = (\mathbb{Z}^2, ((0, 1), (1, 0)))$, even unimodular of signature $(1, 1)$ and discriminant -1 (Conway–Sloane 1999 §2.5). This appendix takes this convention as primary.

The Borcherds–Gritsenko–Nikulin packaging used in §?? of the body and the exterior-square model of $\mathbf{Sp}_4(\mathbb{Z})$ employs the negated form

$$\Lambda^{(1,1)} = \left(\mathbb{Z}^2, \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix} \right),$$

so that the time-like cone $C(\Lambda^{2,1})_+ \subset \Lambda^{2,1} \otimes \mathbb{R}$ has *negative* square $(x, x) < 0$. The relation is $\Lambda^{(1,1)} = U(-1)$. Every Gram matrix below is given in the standard convention; a sign flip transports it to the body convention without changing rank, discriminant absolute value, or automorphism group.

1.1 The hyperbolic plane $U = \Pi_{1,1}$

The hyperbolic plane is the unique even unimodular indefinite lattice of signature $(1, 1)$ (Conway–Sloane 1999 §2.5; Nikulin 1980 Theorem 1.1.2).

Definition 1.2 ($U = \Pi_{1,1}$). [definitional, α] Let $U := \mathbb{Z}e \oplus \mathbb{Z}f$ with Gram matrix

$$G_U = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad e^2 = f^2 = 0, \quad e \cdot f = 1.$$

Verification 1.3 (*Rank, signature, discriminant of U*). [computed, γ] $\text{rk } U = 2$. The eigenvalues of G_U are ± 1 , so the signature is $(1, 1)$. The discriminant is

$$\text{disc } U = \det G_U = -1.$$

U is even (every G_U -diagonal entry is $0 \equiv 0 \pmod{2}$) and unimodular ($|\det G_U| = 1$). These are the defining invariants of $\Pi_{1,1}$ in the Conway–Sloane classification (Conway–Sloane 1999 §2.5, Theorem 2.5.1).

PROPOSITION 1.4 ($\text{Aut}(U)$). [proved, α] The orthogonal group of U is

$$\text{O}(U) = \{\pm \mathbf{I}\} \times \langle \sigma \rangle \simeq (\mathbb{Z}/2)^2, \quad \sigma : e \leftrightarrow f.$$

The proper subgroup $\text{O}^+(U) \simeq \mathbb{Z}/2$ swaps e and f but fixes the spinor norm.

Proof. An isometry $\phi \in \text{O}(U)$ preserves the isotropic vectors of U . In rank 2 over $\mathbb{Z}$, the isotropic primitive vectors are exactly $\pm e, \pm f$, so ϕ permutes these four vectors. The permutations preserving the bilinear form give the four-element group above (Conway–Sloane 1999 §4.1). $\square$

Remark 1.5 (Standard sign convention versus body convention). The body of the manuscript uses $\Lambda^{(1,1)} = U(-1)$ in the cusp polarization of §?? and in the exterior-square model of the automorphic group, where the time-like cone of $\Lambda^{2,1}$ has negative square. The map $U \rightarrow U(-1)$, $(e, f) \mapsto (e, f)$ is a bijection of underlying $\mathbb{Z}$ -modules; only the bilinear form flips sign. Lattices that occur in the body ($\Lambda^{2,1}, \Lambda^{3,2}$) are presented in body coordinates so that the Gram matrices match line-for-line with the manuscript text; even-unimodular ambient lattices ($\Lambda_{K3}, \tilde{H}(S, \mathbb{Z})$) are presented in the standard Conway–Sloane convention. Each section flags its choice explicitly.

1.2 The Lorentzian root lattice $\Lambda_{II}^{2,1}$

The Borcherds–Kac–Moody algebra $\mathfrak{g}_{\Delta_5}$ of [?GNII, Theorem 2.1, Section 5.1.1, case $(t = 1, II, \bar{0})$] is graded by an integral lattice of signature $(2, 1)$. This is the lattice on which the type-II simple-root Cartan matrix is $2(\mathbf{I}_3 - \mathbf{J}_3)$ where $\mathbf{J}_3$ is the all-ones matrix.

Definition 1.6 ($\Lambda_{II}^{2,1}$ as $U(-1) \oplus \langle 2 \rangle$). [definitional, α] Set

$$\Lambda_{II}^{2,1} := U(-1) \oplus \langle 2 \rangle = \mathbb{Z}e \oplus \mathbb{Z}f \oplus \mathbb{Z}h, \quad e \cdot f = -1, \quad e^2 = f^2 = 0, \quad h^2 = 2.$$

The Gram matrix in the basis (e, f, h) is

$$G_{II,2,1} = \begin{pmatrix} 0 & -1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 2 \end{pmatrix}.$$

The body’s lattice $\Lambda^{2,1} = \Lambda^{(1,1)} \oplus [2]$ in (f_2, f_3, f_{-2}) -coordinates uses exactly this Gram, with $\Lambda^{(1,1)} = U(-1)$ on the (f_2, f_{-2}) -block and $[2] = \langle 2 \rangle$ on the f_3 -axis.

VERIFICATION 1.7 (*Rank, signature, discriminant of $\Lambda_{II}^{2,1}$*). [computed, γ] $\text{rk } \Lambda_{II}^{2,1} = 3$. The hyperbolic block $U(-1)$ has eigenvalues ± 1 and contributes signature $(1, 1)$; the rank-1 summand $\langle 2 \rangle$ has eigenvalue $+2$ and contributes signature $(1, 0)$. Total

$$\sigma(\Lambda_{II}^{2,1}) = (2, 1),$$

matching the body convention in which the time-like cone of $\Lambda^{2,1}$ has negative square $(x, x) < 0$ on the rank-2 hyperbolic block. The determinant is

$$\det G_{II,2,1} = -1 \cdot 2 = -2, \quad \text{disc } \Lambda_{II}^{2,1} = -2.$$

The lattice is even because every diagonal entry is even and the off-diagonal couplings are integral. It is not unimodular; the discriminant group is

$$\Lambda_{II}^{2,1,*} / \Lambda_{II}^{2,1} \simeq \mathbb{Z}/2,$$

generated by $h/2$. Switching to the standard hyperbolic-plane convention $U = ((0, 1), (1, 0))$ gives the alternative presentation $\Lambda_{II}^{2,1} \simeq U \oplus \langle -2 \rangle$ of signature $(1, 2)$ and determinant $+2$; rank, discriminant group $\mathbb{Z}/2$, automorphism group, and Cartan datum below are unchanged.

Remark 1.8 (Identification with the body lattice). The body uses $\Lambda^{2,1} = \Lambda^{(1,1)} \oplus [2] = \mathbb{Z}f_2 \oplus \mathbb{Z}f_3 \oplus \mathbb{Z}f_{-2}$ with Gram in (f_2, f_3, f_{-2})

$$\begin{pmatrix} 0 & 0 & -1 \\ 0 & 2 & 0 \\ -1 & 0 & 0 \end{pmatrix}.$$

The basis change

$$e \mapsto f_2, \quad f \mapsto f_{-2}, \quad h \mapsto f_3$$

sends $G_{II,2,1}$ above onto the body Gram (up to row/column permutation between the h -axis and the hyperbolic block). The type-II sublattice

$$\Lambda_{II}^{2,1} \subset \Lambda^{2,1}, \quad \Lambda_{II}^{2,1} = \{mf_2 + lf_3 + nf_{-2} \mid m \equiv n \equiv 0 \pmod{2}\}$$

of the body has the same Gram (up to the appendix sign) and is the even Lorentzian root lattice of $\mathfrak{g}_{\Delta_5}$.

PROPOSITION 1.9 (Type-II simple-root Cartan). [proved, γ] In the body coordinates fixed at the start of the Weyl-chamber section, set

$$\delta_1 = 2f_2 - f_3, \quad \delta_2 = 2f_{-2} - f_3, \quad \delta_3 = f_3.$$

Then $\delta_i^2 = 2$ for $i = 1, 2, 3$ and

$$((\delta_i, \delta_j))_{i,j=1,2,3} = \begin{pmatrix} 2 & -2 & -2 \\ -2 & 2 & -2 \\ -2 & -2 & 2 \end{pmatrix} = 2(\mathbf{I}_3 - \mathbf{J}_3),$$

where $\mathbf{J}_3$ is the 3×3 all-ones matrix. Each δ_i is type-II in the sense that $(\delta_i, \Lambda^{2,1}) = 2\mathbb{Z}$.

Proof. Direct computation in the body Gram. The companion script `compute/verify_lattice.py` at lines 92–97 verifies the matrix $2(\mathbf{I}_3 - \mathbf{J}_3)$ from the same data; the print output records Type-II simple-root Gram matrix equal to the displayed Cartan. The type-II divisibility statement follows because each δ_i pairs against $\Lambda^{2,1}$ with even values: $(\delta_i, f_2), (\delta_i, f_3), (\delta_i, f_{-2}) \in 2\mathbb{Z}$ for each i . This is the GN root datum of [GNII, Section 5.1.1, case (1, II, $\bar{0}$)]. $\square$

PROPOSITION 1.10 (Weyl vector ρ). [proved, γ] The Weyl vector

$$\rho := \frac{1}{2}(\delta_1 + \delta_2 + \delta_3) = f_2 - \frac{1}{2}f_3 + f_{-2} \in \Lambda^{2,1} \otimes \mathbb{Q}$$

satisfies $(\rho, \delta_i) = -1$ for $i = 1, 2, 3$ and

$$(\rho, \rho) = -\frac{3}{2}, \quad \rho \in C(\Lambda^{2,1})_+.$$

In particular ρ lies in the time-like cone, so the chamber $\mathcal{P}_{II} = \{x \mid (x, \delta_i) \leq 0, i = 1, 2, 3\}$ has finite hyperbolic volume.

Proof. The simple-root condition $(\rho, \delta_i) = -(\delta_i^2)/2 = -1$ follows from Kac [?KAC: 1, §2.5, Lemma 2.5] for any choice of simple roots of square 2; for the GN type-II datum see [?GNII, (2.7)]. The script `compute/verify_lattice.py` asserts $\rho = (1, -1/2, 1)$ in (f_2, f_3, f_{-2}) coordinates (line 99) and $(\rho, \delta_i) = -1$ at line 100. Direct computation in the body Gram of Definition 1.6:

$$(\rho, \rho) = 2(f_2, f_{-2}) + \frac{1}{4}(f_3, f_3) = 2(-1) + \frac{1}{4}(2) = -\frac{3}{2}.$$

Since $(\rho, \rho) < 0$, $\rho \in C(\Lambda^{2,1})_+$. □

PROPOSITION 1.11 (*Reflection group; Nikulin's theorem*). [proved, α ; Nikulin 1980 Theorem 1.4.1] Let $W^{(2)}(\Lambda^{2,1})$ be the subgroup of $\mathbf{O}(\Lambda^{2,1})$ generated by reflections in primitive roots of square 2. Then

$$\mathbf{O}(\Lambda^{2,1})_+ = W^{(2)}(\Lambda^{2,1}),$$

where the subscript $+$ denotes the index-two subgroup preserving the time-like cone $C(\Lambda^{2,1})_+$. Equivalently, every isometry of $\Lambda^{2,1}$ preserving the chosen positive cone is a product of square-2 reflections.

Proof. Nikulin [?NIKULIN, Theorem 1.4.1] establishes for the rank-three hyperbolic lattice $\Lambda^{(1,1)} \oplus [2]$ the equality $\mathbf{O}(\Lambda^{2,1})_+ = W^{(2)}(\Lambda^{2,1})$; see also Gritsenko–Nikulin [?GN, Proposition 1.5]. The body of the manuscript invokes this at line 5408–5410. □

VERIFICATION 1.12 (*Type-II reflection orbit*). [computed, γ] For each $\delta \in \{\delta_1, \delta_2, \delta_3\}$, the reflection $s_\delta : x \mapsto x - (x, \delta)\delta$ satisfies $s_\delta(\delta) = -\delta$ and preserves the Cartan matrix on the δ -orbit (compute/verify_lattice.py lines 103–108). Concretely, $s_{\delta_1}(\delta_1) = -\delta_1$, $s_{\delta_1}(\delta_2) = \delta_2 + 2\delta_1$, etc.; the bilinear form on $\{\delta_1, \delta_2, \delta_3\}$ is reflection-invariant by direct matrix computation.

Remark 1.13 (*The two type-2 divisibility classes*). Each $\delta \in \Lambda^{2,1}$ with $\delta^2 = 2$ has divisibility $(\delta, \Lambda^{2,1}) \in \{\mathbb{Z}, 2\mathbb{Z}\}$. The two cases are recorded as *type I* ($(\delta, \Lambda^{2,1}) = \mathbb{Z}$) and *type II* ($(\delta, \Lambda^{2,1}) = 2\mathbb{Z}$); both classes generate index-finite normal subgroups of $W^{(2)}(\Lambda^{2,1})$:

$$[W^{(2)}(\Lambda^{2,1}) : W^{(2)}(\Lambda_I^{2,1})] = 2, \quad [W^{(2)}(\Lambda^{2,1}) : W^{(2)}(\Lambda_{II}^{2,1})] = 6,$$

following [?GN, Sections 4–5]. The denominator algebra $\mathfrak{g}_{\Delta_s}$ uses the type-II reflection chamber.

1.3 *The K_3 lattice $\Lambda_{K_3} = \Pi_{3,19}$*

For S a complex K_3 surface, the integral cohomology $H^2(S, \mathbb{Z})$ with cup-product pairing is the unique even unimodular indefinite lattice of signature $(3, 19)$ (Conway–Sloane 1999 §16.4; Nikulin 1980 Theorem 1.1.4).

Definition 1.14 ($\Lambda_{K_3} = \Pi_{3,19}$). [definitional, α] The K_3 cohomology lattice is

$$\Lambda_{K_3} := H^2(S, \mathbb{Z}) \simeq -E_8 \oplus -E_8 \oplus U \oplus U \oplus U \equiv -2E_8 \oplus 3U,$$

where E_8 is the rank-8 root lattice with the standard positive-definite Cartan pairing (Conway–Sloane 1999 §4.8) and $-E_8$ denotes E_8 with bilinear form negated.

VERIFICATION 1.15 (*Rank, signature, discriminant of Λ_{K_3}*). [computed, α]

$$\text{rk } \Lambda_{K_3} = 8 + 8 + 2 + 2 + 2 = 22, \quad \sigma(\Lambda_{K_3}) = (3, 19).$$

The three U -summands contribute signature $(3, 3)$ and the two $-E_8$ -summands contribute signature $(0, 16)$; total $(3, 19)$. The discriminant is

$$\text{disc } \Lambda_{K_3} = (\det -E_8)^2 \cdot (\det U)^3 = 1 \cdot (-1)^3 = -1,$$

so $|\text{disc}| = 1$ and the lattice is unimodular (Conway–Sloane 1999 §16.4, Theorem 16.4.1). Even unimodularity follows because each summand is even unimodular.

PROPOSITION 1.16 (*Aut and Hodge data on Λ_{K_3}*). [proved, α] The orthogonal group $\mathbf{O}(\Lambda_{K_3})$ acts transitively on the set of primitive vectors of any fixed square (Eichler 1952; Nikulin [?Nikulin1980, Theorem 1.14.4]). For a polarized K_3 with period $\omega \in H^{2,0}(S) \subset \Lambda_{K_3} \otimes \mathbb{C}$, the transcendental and algebraic sublattices fit into the orthogonal splitting

$$\Lambda_{K_3} = \text{NS}(S) \oplus T(S),$$

where $\text{NS}(S) \subset H^{1,1}(S, \mathbb{Z})$ is the Néron–Severi (algebraic) sublattice of signature $(1, \rho - 1)$ and $T(S)$ is the transcendental lattice of signature $(2, 20 - \rho)$, with $\rho = \text{rk } \text{NS}(S)$ the Picard rank (Beauville–Bourguignon–Demazure 1985 §6; [?Nikulin1980, §2]).

Proof. The transitivity statement is Eichler’s theorem on indefinite unimodular lattices, sharpened by Nikulin [?Nikulin1980, Theorem 1.14.4]: for an even unimodular indefinite lattice L of rank ≥ 3 , $\mathbf{O}(L)$ acts transitively on primitive vectors of fixed square in each squared-class. Decomposition by Hodge type partitions $\Lambda_{K_3} \otimes \mathbb{C}$ into $H^{2,0} \oplus H^{1,1} \oplus H^{0,2}$; intersecting with Λ_{K_3} gives the transcendental–algebraic splitting at the integral level. $\square$

1.4 The lattice-polarized K_3 sublattice Λ_S

A lattice polarization is a primitive embedding of an integral lattice Λ_S into Λ_{K_3} whose orthogonal complement carries period structure.

Definition 1.17 (*Lattice polarization $\Lambda_S \hookrightarrow \Lambda_{K_3}$*). [definitional, α] A *lattice polarization* of signature $(1, \rho - 1)$ on a K_3 surface S is a primitive embedding

$$\Lambda_S \hookrightarrow \Lambda_{K_3} = \Pi_{3,19}, \quad \sigma(\Lambda_S) = (1, \rho - 1), \quad 1 \leq \rho \leq 19,$$

where $\Lambda_S \subset H^{1,1}(S, \mathbb{Z})$ contains an ample class. The orthogonal complement

$$\Lambda_S^\perp \subset \Lambda_{K_3}, \quad \sigma(\Lambda_S^\perp) = (2, 20 - \rho),$$

is the lattice on which the period map of the family $\mathcal{F}_{\Lambda_S} \rightarrow \mathcal{D}_{\Lambda_S^\perp}$ is defined (Dolgachev 1996; Nikulin 1980 Proposition 1.6.1).

VERIFICATION 1.18 (*Numerics of Λ_S and $\Lambda_S^\perp$*). [computed, α]

$$\text{rk } \Lambda_S = \rho, \quad \sigma(\Lambda_S) = (1, \rho - 1), \quad \text{rk } \Lambda_S^\perp = 22 - \rho, \quad \sigma(\Lambda_S^\perp) = (2, 20 - \rho).$$

The discriminant pairs:

$$\text{disc } \Lambda_S \cdot \text{disc } \Lambda_S^\perp = \text{disc } \Lambda_{K_3} = -1,$$

so $\text{disc } \Lambda_S^\perp = -\text{disc } \Lambda_S^{-1} \cdot \det \mathcal{A}^2$ for $\mathcal{A}$ the change-of-basis matrix realizing the primitive embedding (Nikulin 1980 Corollary 1.6.2). In particular $\Lambda_S^\perp / \Lambda_S^*$ and $\Lambda_{K_3} / \Lambda_S \oplus \Lambda_S^\perp$ are isomorphic as discriminant groups.

PROPOSITION 1.19 (*Nikulin embedding criterion*). [proved; Nikulin 1980 Theorem 1.14.4] A primitive embedding $\Lambda_S \hookrightarrow \Lambda_{K_3}$ exists if and only if there is an isomorphism

$$(\mathcal{A}_{\Lambda_S}, q_{\Lambda_S}) \simeq (\mathcal{A}_{\Lambda_S^\perp}, -q_{\Lambda_S^\perp})$$

of finite quadratic forms, where $\mathcal{A}_L := L^*/L$ is the discriminant group and q_L its quadratic form (Nikulin 1980 Definition 1.3.1 and Theorem 1.14.4; Conway–Sloane 1999 §15.5). The embedding is unique up to $\mathbf{O}(\Lambda_{K_3})$ when

$$\mathrm{rk} \Lambda_S + \ell(\mathcal{A}_{\Lambda_S}) \leq 22 - \mathrm{rk} \Lambda_S$$

and the discriminant form satisfies the Nikulin uniqueness criterion (Nikulin 1980 Theorem 1.14.4).

Proof. This is the gluing-via-discriminant-form theorem ([Nikulin1980, §1.14]; [ConwaySloane, §15.5]). An embedding into Λ_{K_3} exists iff the discriminant form on Λ_S^*/Λ_S is realized by the orthogonal complement; uniqueness is the Nikulin orbit-uniqueness statement under the corank inequality. $\square$

Remark 1.20 (Mirror lattice convention). The Aspinwall–Morrison mirror to a Λ_S -polarized K_3 family is the $\check{\Lambda}_S$ -polarized family with

$$\check{\Lambda}_S = U \oplus (\Lambda_S^\perp \oplus U),$$

the splitting depending on a choice of isotropic primitive $U \hookrightarrow \Lambda_S^\perp$ (Dolgachev 1996 §5; Aspinwall 1997). The two factors Λ_S and $\Lambda_S^\perp \oplus U$ play the roles of algebraic–symplectic exchange under mirror symmetry. No mirror identification is asserted at this lattice level; the assertion that Δ_S^{-2} is the discriminant of the mirror period family (View ④) is conjectural and not used in Chapter ??.

1.5 The Mukai lattice $\tilde{H}(S, \mathbb{Z}) = \Pi_{4,20}$

For S a complex K_3 surface, the full integral cohomology $H^*(S, \mathbb{Z})$ with the Mukai pairing is an even unimodular lattice of signature $(4, 20)$ (Mukai 1984 Theorem 1.5; Mukai 1987).

Definition 1.21 (*Mukai lattice $\tilde{H}(S, \mathbb{Z})$*). [definitional, α] The Mukai lattice of S is

$$\tilde{H}(S, \mathbb{Z}) := H^0(S, \mathbb{Z}) \oplus H^2(S, \mathbb{Z}) \oplus H^4(S, \mathbb{Z}), \quad \tilde{H}(S, \mathbb{Z}) \simeq U \oplus U \oplus U \oplus U \oplus -2E_8 \equiv 4U \oplus 2(-E_8),$$

graded by even cohomological degree. The Mukai pairing is

$$\langle (r, D, s), (r', D', s') \rangle_{\mathrm{Muk}} := D \cdot D' - rs' - r's$$

for $(r, D, s), (r', D', s') \in \mathbb{Z} \oplus H^2(S, \mathbb{Z}) \oplus \mathbb{Z}$ (Mukai 1984 Theorem 1.5). The pairing of an H^0 -class with an H^4 -class carries a minus sign relative to the cup product, so that the rank-4 component of $H^*(S, \mathbb{Z}) \otimes H^*(S, \mathbb{Z})$ contributes $-rs' - r's$.

VERIFICATION 1.22 (*Rank, signature, discriminant of $\tilde{H}(S, \mathbb{Z})$*). [computed, α]

$$\mathrm{rk} \tilde{H}(S, \mathbb{Z}) = 1 + 22 + 1 = 24, \quad \sigma(\tilde{H}(S, \mathbb{Z})) = (4, 20).$$

The four U -summands give signature $(4, 4)$ and the $-2E_8$ pieces give signature $(0, 16)$, total $(4, 20)$. Discriminant

$$\mathrm{disc} \tilde{H}(S, \mathbb{Z}) = (\det U)^4 \cdot (\det -E_8)^2 = (-1)^4 \cdot 1 = 1,$$

so $\tilde{H}(S, \mathbb{Z})$ is even unimodular, denoted $\Pi_{4,20}$ (Conway–Sloane 1999 §16.4). The body uses the notation $\Lambda_S = \tilde{H}(S, \mathbb{Z})$ at line 4589 and 4714 of main.tex; this is consistent with the convention here.

PROPOSITION 1.23 (*Mukai vector*). [proved, α ; Mukai 1987] For F a coherent sheaf on S , the Mukai vector is

$$v(F) := \text{ch}(F)\sqrt{\text{td}(S)} = (\text{rk } F, c_1(F), \text{ch}_2(F) + \text{rk } F \cdot [\text{pt}]) \in \tilde{H}(S, \mathbb{Z}),$$

where $\sqrt{\text{td}(S)} = (1, 0, 1)$ in Mukai notation on a K3 surface ([Mukai 1987, Theorem 3]). The Mukai pairing on $\tilde{H}(S, \mathbb{Z})$ recovers the Euler characteristic of Hom:

$$\langle v(F), v(G) \rangle_{\text{Muk}} = -\chi(F, G) = -\sum_i (-1)^i \dim \text{Ext}^i(F, G).$$

Proof. Mukai [Mukai 1987, Theorem 3] computes $\sqrt{\text{td}(S)}$ for K3 surfaces. The Hirzebruch–Riemann–Roch formula gives $\chi(F, G) = \int_S \text{ch}(F^\vee) \text{ch}(G) \text{td}(S) = -\langle v(F), v(G) \rangle_{\text{Muk}}$ once the sign of the H^0-H^4 pairing is taken into account; see Mukai 1984 §2. $\square$

Remark 1.24 (*Sign of the Mukai pairing on rank-0 components*). The Mukai pairing $\langle (r, D, s), (r', D', s') \rangle_{\text{Muk}} = D \cdot D' - rs' - r's$ restricts to the cup-product pairing on the middle H^2 -component

$$\langle (0, D, 0), (0, D', 0) \rangle_{\text{Muk}} = D \cdot D',$$

and to the negative cup-product on the rank–Euler component

$$\langle (r, 0, s), (r', 0, s') \rangle_{\text{Muk}} = -rs' - r's.$$

The relative minus sign is forced by Hirzebruch–Riemann–Roch: $-\chi(F, G)$ contains a $-\text{rk } F \cdot \chi(\emptyset, G) - \text{rk } G \cdot \chi(\emptyset, F)$ term coming from $\sqrt{\text{td}(S)} = (1, 0, 1)$. An apparent attack route is to negate the H^0-H^4 pairing or to drop the minus sign; this would convert the Mukai lattice to an isotropic-rank-2 unimodular form not of signature $(4, 20)$, in disagreement with the Mukai 1984 classification ([Mukai 1984, Theorem 1.5]) and the unique even unimodular type $\text{II}_{4,20}$. The sign as stated is forced by these requirements.

PROPOSITION 1.25 (*Aut of the Mukai lattice*). [proved, α] $\mathbf{O}(\tilde{H}(S, \mathbb{Z}))$ acts transitively on the set of primitive vectors of any fixed square (Eichler 1952; Nikulin 1980 Theorem 1.14.4). In particular, the Mukai vector $v = (1, 0, 1)$ (corresponding to the structure sheaf $\mathcal{O}_S$) and the generator $(0, 0, 1)$ of $H^4(S, \mathbb{Z}) \cap \tilde{H}(S, \mathbb{Z})$ are $\mathbf{O}(\tilde{H}(S, \mathbb{Z}))$ -conjugate to any other primitive vector of the same square. The derived autoequivalence group of $D^b(S)$ maps to $\mathbf{O}(\tilde{H}(S, \mathbb{Z}))$ via the action on Mukai vectors (Mukai 1987 §2; Orlov 1997).

Proof. Eichler’s theorem ([Nikulin 1980, Theorem 1.14.4]) gives the transitivity statement. Mukai [Mukai 1987, §2] constructs the representation of derived autoequivalences on $\tilde{H}(S, \mathbb{Z})$ via Fourier–Mukai kernels. $\square$

1.6 The formal dyonic charge lattice $\Gamma_X^{\text{form}} = \Lambda_S \oplus \Lambda_S$

The formal additive dyonic lattice on $X = S \times E$ packages the electric–magnetic charge data on which the Gram-shadow projection Π_X is defined. Throughout $\Lambda_S = \tilde{H}(S, \mathbb{Z})$ in the sense of §1.5.

Definition 1.26 (Γ_X^{form} and its Gram pairing). [definitional, α ; from Definition ??] For $X = S \times E$ with S a K3 surface and E an elliptic curve,

$$\Gamma_X^{\text{form}} := \Lambda_S \oplus \Lambda_S, \quad c = (Q, P) \in \Gamma_X^{\text{form}}, \quad Q, P \in \Lambda_S.$$

The Gram pairing $\langle \cdot, \cdot \rangle_\Gamma : \Gamma_X^{\text{form}} \times \Gamma_X^{\text{form}} \rightarrow \mathbb{Z}$ is inherited from the Mukai pairing on each factor:

$$\langle (Q, P), (Q', P') \rangle_\Gamma := \langle Q, Q' \rangle_{\text{Muk}} + \langle P, P' \rangle_{\text{Muk}}.$$

The Gram-index projection

$$\Pi_X : \Gamma_X^{\text{form}} \rightarrow \Gamma_{\text{gram}} = \mathbb{Z}^3, \quad (Q, P) \mapsto \left(\frac{Q^2}{2}, Q \cdot P, \frac{P^2}{2} \right),$$

records the genus-two Gram triple of the dyonic class. This map is quadratic, not additive: the obstruction is the symmetric bilinear 2-cocycle B of Lemma 2.4.

VERIFICATION 1.27 (*Rank, signature of Γ_X^{form}*). [computed, α]

$$\text{rk } \Gamma_X^{\text{form}} = 2 \cdot \text{rk } \Lambda_S = 48, \quad \sigma(\Gamma_X^{\text{form}}) = 2 \cdot (4, 20) = (8, 40).$$

The discriminant is

$$\text{disc } \Gamma_X^{\text{form}} = (\text{disc } \Lambda_S)^2 = 1,$$

so Γ_X^{form} is even unimodular. In Conway–Sloane notation it is $\Pi_{8,40}$ (Conway–Sloane 1999 §16.4).

Remark 1.28 (Relation to the $K3 \times E$ Mukai geometry). Under the Künneth decomposition for $X = S \times E$, the integral even cohomology decomposes as

$$H^{\text{ev}}(X, \mathbb{Z}) = H^*(S, \mathbb{Z}) \otimes H^*(E, \mathbb{Z}) = H^*(S, \mathbb{Z}) \otimes (\mathbb{Z}[1] \oplus \mathbb{Z}[\omega_E]),$$

where ω_E is the fundamental class of E . Every Mukai vector on X in the formal even sector decomposes as $v_X = P \otimes 1_E + Q \otimes \omega_E$ with $Q, P \in \Lambda_S$, and $\Gamma_X^{\text{form}} = \Lambda_S \oplus \Lambda_S$ is the additive lattice on which the (P, Q) -grading lives (Definition ??). The Mukai pairing on $\tilde{H}(X, \mathbb{Z})$ pairs against the elliptic U -summand $H^0(E, \mathbb{Z}) \oplus H^2(E, \mathbb{Z}) \simeq U$, giving

$$\langle v_X, v'_X \rangle_{\tilde{H}(X)} = Q \cdot P' + Q' \cdot P,$$

where each summand is the Mukai pairing on Λ_S . The resulting genus-two Gram triple $(Q^2/2, Q \cdot P, P^2/2)$ is the Gram-index data $\Pi_X(Q, P)$ used in the Borcherds product (Theorem ??).

LEMMA 1.29 (*Two orthogonal hyperbolic planes inside Λ_S*). [proved, α] $\Lambda_S = \tilde{H}(S, \mathbb{Z}) \simeq 4U \oplus 2(-E_8)$ contains four mutually orthogonal copies of U . In particular, two of them give

$$U_1 \oplus U_2 \subset \Lambda_S, \quad U_i \simeq U, \quad U_1 \perp U_2,$$

which is the hypothesis of the formal primitive lift Lemma 2.7: every Gram triple $(n, l, m) \in \mathbb{Z}^3$ has a primitive saturated formal lift $c = (Q, P) \in \Gamma_X^{\text{form}}$ with $\Pi_X(Q, P) = (n, l, m)$.

Proof. By Definition 1.21, $\Lambda_S \simeq 4U \oplus 2(-E_8)$ contains four mutually orthogonal U -summands. Selecting any two of them gives the displayed sublattice, which is itself isomorphic to $U \oplus U \simeq \Pi_{2,2}$ as a sublattice of Λ_S . The Gram-triple lift then follows from Lemma 2.7: take $Q = e_1 + nf_1$, $P = lf_1 + e_2 + mf_2$ in $U_1 \oplus U_2$ to obtain $\Pi_X(Q, P) = (n, l, m)$. $\square$

1.7 The signature-(3, 2) lattice $\Lambda^{3,2}$ and the $\mathbf{Sp}_4 \simeq \mathbf{Spin}(3, 2)$ bridge

The exceptional isomorphism $\mathbf{PSp}_4(\mathbb{R}) \simeq \mathbf{SO}_+(3, 2)$ identifies the Siegel modular variety with the fourth-orthogonal modular variety of an integral lattice $\Lambda^{3,2}$ of signature (3, 2) ([?GN, Section 3]; Borchers 1995 §5). The Igusa cusp form Δ_5 lives on the Siegel side; the Borchers product expansion lives on the orthogonal side.

Definition 1.30 ($\Lambda^{3,2}$ in the exterior-square model). [definitional, α] Let $\Lambda^4 = \mathbb{Z}e_1 \oplus \mathbb{Z}e_2 \oplus \mathbb{Z}e_3 \oplus \mathbb{Z}e_4$ and consider the exterior square $\wedge^2 \Lambda^4$ with the symmetric pairing

$$u \wedge v \cdot u' \wedge v' = [\det(u, v, u', v')]_{e_1 \wedge e_2 \wedge e_3 \wedge e_4} \in \mathbb{Z}.$$

Then $(\wedge^2 \Lambda^4, (\cdot, \cdot)) \simeq \Pi_{3,3}$ as an even unimodular lattice of signature (3, 3). Fix $q_o := e_1 \wedge e_3 + e_2 \wedge e_4 \in \wedge^2 \Lambda^4$ and define

$$\Lambda^{3,2} := q_o^\perp \subset \wedge^2 \Lambda^4.$$

VERIFICATION 1.31 (Rank, signature, discriminant of $\Lambda^{3,2}$). [computed, α ; verified by compute/verify_lattice.py] $\text{rk } \wedge^2 \Lambda^4 = \binom{4}{2} = 6$. Removing one isotropic ray from q_o gives $\text{rk } \Lambda^{3,2} = 5$. The expected_gram array of compute/verify_lattice.py at lines 82–88 is the explicit Gram matrix of $\Lambda^{3,2}$ in the basis $(f_1, f_2, f_3, f_{-2}, f_{-1})$:

$$G_{3,2} = \begin{pmatrix} 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 2 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 \end{pmatrix},$$

where $f_1 = e_1 \wedge e_2, f_2 = e_2 \wedge e_3, f_3 = e_1 \wedge e_3 - e_2 \wedge e_4, f_{-2} = e_4 \wedge e_1, f_{-1} = e_4 \wedge e_3$. Direct computation: the matrix is antidiagonal except for the central $h^2 = 2$ entry, with off-diagonal entries $-1, -1, -1, -1$. Expansion gives

$$\det G_{3,2} = 2 \cdot (-1)^4 \cdot \text{sgn}(1, 2, 3, 4 \mapsto 4, 3, 2, 1) = 2 \cdot 1 \cdot (+1) = +2,$$

since the permutation $(1, 2, 3, 4) \mapsto (4, 3, 2, 1)$ has six inversions and even parity. Therefore

$$\sigma(\Lambda^{3,2}) = (3, 2), \quad \text{disc } \Lambda^{3,2} = 2.$$

The script asserts `gram == expected_gram` at line 89.

PROPOSITION 1.32 ($\Lambda^{3,2} \simeq U \oplus U \oplus \langle 2 \rangle$ in the manuscript convention; $\Lambda^{3,2} \simeq U(-1) \oplus U(-1) \oplus \langle 2 \rangle$ in the standard convention). [proved, α] In the body convention of the exterior-square model,

$$\Lambda^{3,2} \simeq \Lambda^{(1,1)} \oplus \Lambda^{(1,1)} \oplus [2],$$

with $\Lambda^{(1,1)} = U(-1)$ and $[2]$ the rank-1 lattice with Gram (2). The Gram presentation displayed in Verification 1.31 arises from the f -basis: (f_1, f_{-1}) pairs as a copy of $\Lambda^{(1,1)}$ ($(f_1, f_{-1}) = -1$), (f_2, f_{-2}) pairs as a second copy of $\Lambda^{(1,1)}$ ($(f_2, f_{-2}) = -1$), and f_3 contributes $(f_3, f_3) = 2$.

Proof. The pairings $(f_1, f_{-1}) = -1, (f_2, f_{-2}) = -1, (f_3, f_3) = 2$ are read off from $G_{3,2}$. The orthogonal decomposition is direct: $\mathbb{Z}f_1 + \mathbb{Z}f_{-1} \perp \mathbb{Z}f_2 + \mathbb{Z}f_{-2} \perp \mathbb{Z}f_3$ inside $\Lambda^{3,2}$ because every cross-pair vanishes in $G_{3,2}$. The identification with $\Lambda^{(1,1)} \oplus \Lambda^{(1,1)} \oplus [2]$ follows because each rank-2 block has Gram $((0, -1), (-1, 0)) = \Lambda^{(1,1)}$ and the rank-1 block has Gram (2) = $[2]$. $\square$

PROPOSITION 1.33 ($\mathbf{PSp}_4(\mathbb{Z}) \simeq \mathbf{SO}_+(\Lambda^{3,2})$). [proved, α ; exterior-square model of $\mathbf{Sp}_4(\mathbb{Z})$] The exterior-square representation $\wedge^2 : \mathbf{Sp}_4(\mathbb{Z})/\{\pm \mathbf{I}_4\} \rightarrow \mathbf{SO}_+(\Lambda^{3,2}) \simeq \mathbf{O}(\Lambda^{3,2})_+/\{\pm \mathbf{I}_5\}$ is an isomorphism. Concretely, the integral generators

$$g_o = \begin{pmatrix} o & \mathbf{I}_2 \\ -\mathbf{I}_2 & o \end{pmatrix}, \quad g_M = \begin{pmatrix} \mathbf{I}_2 & M \\ o & \mathbf{I}_2 \end{pmatrix}, \quad g_A = \begin{pmatrix} {}^t A^{-1} & o \\ o & A \end{pmatrix} \quad (A \in \mathbf{GL}_2(\mathbb{Z}))$$

of $\mathbf{Sp}_4(\mathbb{Z})$ map under $\wedge^2$ to integral isometries of $\Lambda^{3,2}$, displayed as the three explicit 5×5 matrices $\wedge^2(g_o), \wedge^2(g_M), \wedge^2(g_A)$ in the body's exterior-square model. The full image is $\mathbf{SO}_+(\Lambda^{3,2})$. The Siegel space $\mathbb{H}_2$ embeds into the type-IV upper half-space $\mathbb{H}_+^{\mathbf{IV}} \subset \mathbb{P}(\Lambda^{3,2} \otimes \mathbb{C})$ via

$$Z = \begin{pmatrix} z_1 & z_2 \\ z_2 & z_3 \end{pmatrix} \mapsto {}^t(z_2^2 - z_1 z_3, z_3, z_2, z_1, 1)z_o,$$

and the action of $\mathbf{Sp}_4(\mathbb{Z})$ on $\mathbb{H}_2$ corresponds to the action of $\mathbf{SO}_+(\Lambda^{3,2})$ on $\mathbb{H}_+^{\mathbf{IV}}$.

Proof. The Pfaffian pairing on $\wedge^2 \Lambda^4$ is preserved by $\wedge^2 \mathbf{SL}_4$. The subgroup fixing q_o is $\mathbf{Sp}_4(\mathbb{Z})$ by definition of the symplectic form $B_{q_o}(x, y)e_1 \wedge e_2 \wedge e_3 \wedge e_4 = -x \wedge y \wedge q_o$. The kernel on $q_o^\perp = \Lambda^{3,2}$ is $\{\pm \mathbf{I}_4\}$. The exceptional Lie-algebra isomorphism $\mathfrak{sp}_4(\mathbb{R}) \simeq \mathfrak{so}(3, 2)$ extends to the integral form on the displayed generators; this is the content of [?GN, Section 3] and the manuscript's exterior-square model of the automorphic group. $\square$

Remark 1.34 (*Embedding $\Lambda_{II}^{2,1} \subset \Lambda^{3,2}$*). The primitive sublattice

$$\Lambda^{2,1} = \Lambda^{(1,1)} \oplus [2] = \mathbb{Z}f_2 \oplus \mathbb{Z}f_3 \oplus \mathbb{Z}f_{-2} \subset \Lambda^{3,2}$$

of the body's Weyl-chamber section is the BKM root lattice: discarding the (f_1, f_{-1}) -block of $\Lambda^{3,2} \simeq \Lambda^{(1,1)} \oplus \Lambda^{(1,1)} \oplus [2]$ leaves $\Lambda^{(1,1)} \oplus [2] \simeq \Lambda^{2,1}$. The type-II sublattice $\Lambda_{II}^{2,1} = \{mf_2 + lf_3 + nf_{-2} \mid m, n \equiv 0 \pmod{2}\}$ is the lattice of even integral type-II classes, on which the simple-root Cartan

$$((\partial_i, \partial_j)) = \begin{pmatrix} 2 & -2 & -2 \\ -2 & 2 & -2 \\ -2 & -2 & 2 \end{pmatrix}$$

of Proposition 1.9 is realized via $\partial_1 = 2f_2 - f_3$, $\partial_2 = 2f_{-2} - f_3$, $\partial_3 = f_3$. Every isometry of $\Lambda^{2,1}$ extends to an isometry of $\Lambda^{3,2}$ by the identity on $(\Lambda^{2,1})^\perp \subset \Lambda^{3,2}$.

1.8 Cross-references and consistency check

The lattices recorded above sit in the manuscript as follows.

- $U = \Pi_{1,1}$: every U -summand of $\Lambda^{3,2}$, Λ_{K3} , and $\tilde{H}(S, \mathbb{Z})$. The relation $\Lambda^{(1,1)} = U(-1)$ packages the body's negated convention.
- $\Lambda_{II}^{2,1} \simeq U(-1) \oplus \langle 2 \rangle \simeq U \oplus \langle -2 \rangle$ (sign flip on the rank-2 hyperbolic block): the BKM root lattice on which the type-II simple-root Cartan $2(\mathbf{I}_3 - \mathbf{J}_3)$ and the Weyl vector $\rho = \frac{1}{2}(\partial_1 + \partial_2 + \partial_3)$ are realized. The denominator algebra $\mathfrak{g}_{\Delta_\delta} = \mathfrak{g}_{\Delta_\delta}$ is graded by $\Lambda_{II}^{2,1}$ ([?GNII, Theorem 2.1]).
- $\Lambda_{K3} = \Pi_{3,19}$: K3 cohomology with intersection pairing. Polarized K3 sublattices Λ_S embed primitively into Λ_{K3} .

- Λ_S : signature $(1, \rho - 1)$ for $\rho \leq 19$ (the algebraic Picard rank). Used in Definition ?? as the K_3 factor of Γ_X^{form} and in the lattice-polarization setup of the body.
- $\tilde{H}(S, \mathbb{Z}) = \Pi_{4,20}$: the Mukai lattice on which the D6–D2–D0 Mukai vectors $v_X(\mathcal{I}_Y)$ live; the Gram-index map Π_X is computed via this pairing (Theorem ??).
- $\Gamma_X^{\text{form}} = \Lambda_S \oplus \Lambda_S$: the formal additive dyonic charge lattice of $X = S \times E$, on which the quadratic Gram-shadow projection Π_X and the symmetric bilinear 2-cocycle B are defined (Definition ??).
- $\Lambda^{3,2} \simeq U(-1) \oplus U(-1) \oplus \langle 2 \rangle$: the signature- $(3, 2)$ orthogonal lattice that bridges $\mathbf{PSP}_4(\mathbb{Z})$ to $\mathbf{SO}_+(3, 2)$ via the exterior-square representation; on this lattice the Borcherds product expansion of Δ_5 is written.

VERIFICATION 1.35 (*End-to-end computation check*). [computed, γ] The script `compute/verify_lattice.py` verifies, in one self-contained $\mathbb{Q}$ -arithmetic computation: (i) the rank-5 Gram matrix $G_{3,2}$ in the $(f_1, f_2, f_3, f_{-2}, f_{-1})$ -basis (lines 82–89); (ii) the orthogonality of the symplectic invariant $q_0 = e_1 \wedge e_3 + e_2 \wedge e_4$ to the basis (line 90); (iii) the type-II simple-root Cartan $2(\mathbf{I}_3 - \mathbf{J}_3)$ at lines 92–97; (iv) the Weyl vector $\rho = (1, -1/2, 1)$ in (f_2, f_3, f_{-2}) coordinates and the $(\rho, \delta_i) = -1$ relation (lines 99–101); (v) the type-II reflection action and Cartan invariance under each s_{δ_i} (lines 103–108). The `All lattice checks passed.` line printed at the end of the script’s output confirms agreement of the Gram, Cartan, and Weyl-vector data of §1.2 and §1.7 with the $\mathbb{Q}$ -arithmetic computation.

Remark 1.36 (Discriminant convention; sign of $\Lambda^{3,2}$). In the body convention $\Lambda^{(1,1)} \oplus \Lambda^{(1,1)} \oplus [2]$, both hyperbolic-plane summands have Gram determinant -1 and the $[2]$ summand contributes $+2$, so $\text{disc } \Lambda^{3,2} = (-1)^2 \cdot 2 = +2$. In the standard convention $U \oplus U \oplus \langle -2 \rangle$, the two U -summands contribute determinant -1 each and $\langle -2 \rangle$ contributes -2 , so $\text{disc } \Lambda^{3,2} = (-1)^2 \cdot (-2) = -2$. The two presentations differ by an overall sign on the rank-2 hyperbolic blocks; both carry an order-2 discriminant group with single nontrivial generator $f_3/2 \in (\Lambda^{3,2})^*/\Lambda^{3,2}$. The same analysis applies to $\Lambda_{II}^{2,1}$: discriminant -2 (body) or $+2$ (standard), discriminant group $\mathbb{Z}/2$ generated by $h/2$. In every case the order of the discriminant group is 2 and the type-II divisibility statement $(\delta, \Lambda^{2,1}) \in \{\mathbb{Z}, 2\mathbb{Z}\}$ depends on the order, not the sign of the determinant.

Remark 1.37 (Cited primary literature). The primary references behind every load-bearing lattice claim above are: Conway–Sloane 1999 §2.5, §4.8, §15.5, §16.4 (classification of even unimodular indefinite lattices, E_8 root system, discriminant-form gluing, K_3 lattice as $\Pi_{3,19}$); Nikulin 1980 Theorem 1.1.2, Theorem 1.4.1 (= [NIKULIN] Theorem 1.4.1 in the K_3 -paper version), Theorem 1.6.1, Theorem 1.14.4 (signature classification, reflection group of square-2 vectors, primitive embeddings, discriminant-form uniqueness); Mukai 1984 Theorem 1.5 (Mukai pairing, signature $(4, 20)$), Mukai 1987 Theorem 3 ($\sqrt{\text{td}(S)} = (1, 0, 1)$ on K_3); Gritsenko–Nikulin 1998 Part II Theorem 2.1, Section 5.1.1, case $(t = 1, II, \bar{0})$ (BKM root datum for $\mathfrak{g}_{\Delta_5}$, Weyl vector, type-II simple-root Cartan).

2 BORCHERDS-PRODUCT EXPANSION: $\phi_{0,1}$, THE Δ_5 INFINITE PRODUCT, AND THE GRITSENKO–CLÉRY EIGHT-ROW CATALOGUE

This appendix records, in one place, the Fourier expansion of the weight-zero index-one weak Jacobi form $\phi_{0,1}$, the Borcherds–Gritsenko–Nikulin product expansion of the Igusa cusp form Δ_5 in the type-II cusp chamber, the theta-constant normalization constant $64 = 2^6$, the doubling identity that turns the monic Borcherds product into the denominator of the Lorentzian Borcherds–Kac–Moody algebra $\mathfrak{g}_{\Delta_5}$, and the full eight-row Gritsenko–Cléry catalogue of diagonal-divisor Siegel modular forms. Every numerical Fourier coefficient quoted below is produced by an exact rational computation in

compute/verify_square_root.py; every load-bearing identity is attributed to author, year, and theorem of the primary literature. No construction in this appendix uses orientation data, Pfaffian wall normal forms, or compact Hall input; the appendix lives entirely on the automorphic / target side, prior to the geometric content of (Do), (O1), (O1)⁺, and (O2).

Three names for Δ_5 appear and are kept distinct: the theta-constant automorphic section $D_X = \Delta_5$ (View I); the Borchers–Kac–Moody denominator $\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1} \Delta_5(2Z)$ (View II); and the protected Pfaffian $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{DI})$ of the pro-object $\mathfrak{D}_X^{DI}$ (View III, conditional on (Do), (O1), (O1)⁺, (O2)). All three coincide on Fourier coefficients in the cusp polarization; the orientation character that distinguishes Δ_5 from $-\Delta_5$ enters only at View III.

B.1 The weak Jacobi form $\phi_{0,1}$

The K_3 elliptic genus is the modular input. Eichler–Zagier [?EZ: 1, §2.2 and Theorem 9.3] fix the weight-zero index-one weak-Jacobi normalization

$$\phi_{0,1}(\tau, z) \in J_{0,1}^w, \quad \phi_{0,1}(\tau, z) = \sum_{n \geq 0, l \in \mathbb{Z}} f(n, l) q^n r^l, \quad q = e^{2\pi i \tau}, \quad r = e^{2\pi i z}.$$

The relation to the K_3 right-moving elliptic genus is $Z_{K_3}(\tau, z) = 2\phi_{0,1}(\tau, z)$ [?EZ: 1, §2.2]; the half is the Borchers-lift representative used below. The Jacobi weak-form support is $f(n, l) = 0$ whenever $4n - l^2 < -1$, with isolated extreme entries $f(0, \pm 1) = 1$ on the discriminant hyperbola $4n - l^2 = -1$. (*proved*; [?EZ: 1, §2.2, Theorem 9.3]; [?GNII, Example 2.4]; *licensing* β .)

2.0.0.1 Verified Fourier table. The exact coefficients $f(n, l)$ for $n \leq 2$ are the rational integers listed below, computed by the theta-quotient identity $\phi_{0,1} = 4 \sum_{i=2,3,4} \theta_i(\tau, z)^2 / \theta_i(\tau, 0)^2$ of [?EZ: 1, §2.2] and verified bit-for-bit against compute/verify_square_root.py (rational arithmetic, no floating-point):

	$l = -3$	$l = -2$	$l = -1$	$l = 0$	$l = 1$	$l = 2$	$l = 3$
$n = 0$	0	0	1	10	1	0	0
$n = 1$	0	10	-64	108	-64	10	0
$n = 2$	1	108	-513	808	-513	108	1

(*computed*; *cross-checked with* [?GNII, Example 2.4] *and* [?GN, Theorem 4.1]; *licensing* δ .) The negative entries reflect the index-one Jacobi-form theta-decomposition $\phi_{0,1} = \theta_{1,0} h_0 + \theta_{1,1} h_1$ with theta-coefficient expansions $h_0 = 2(45E_4 E_{4,1} - 50E_6 E_{6,1}) / \eta^{24} + \cdots$ in [?EZ: 1, §2.2]; the integer values are not target signs but signed multiplicities of the Borchers source.

2.0.0.2 Banked identities used below.

$$\begin{aligned} f(0, 0) &= 10, & f(0, \pm 1) &= 1, & f(0, l) &= 0 \quad (|l| \geq 2), \\ f(1, 0) &= 108, & f(1, \pm 1) &= -64, & f(1, \pm 2) &= 10, \\ f(2, 0) &= 808, & f(2, \pm 1) &= -513, & f(2, \pm 2) &= 108, \\ & & f(2, \pm 3) &= 1, & f(1, \pm 3) &= 0. \end{aligned}$$

The diagonal entry $f(0, 0) = 10$ controls the Borchers-lift weight $\text{wt}(\Delta_5) = f(0, 0)/2 = 5$; the entries $f(0, \pm 1) = 1$ control the simple type-II divisor orders $d_{\partial_i}^{\text{aut}} = 1$; the entry $f(1, 1) = -64$ controls the multiplicity of the first timelike imaginary root, computed and used below.

B.2 The cusp chamber and the effective semigroup

Set $Z = \begin{pmatrix} z_1 & z_2 \\ z_2 & z_3 \end{pmatrix} \in \mathbb{H}_2$, $q = e^{2\pi i z_1}$, $r = e^{2\pi i z_2}$, $s = e^{2\pi i z_3}$. The standard type-II cusp polarization is the ordered limit

$$0 < |s| \ll |q| \ll |r|^{-1} \ll 1,$$

which determines the effective semigroup

$$\Gamma_{\text{eff}} = \{(n, l, m) \in \mathbb{Z}^3 \mid m > 0, n \geq 0\} \cup \{(n, l, 0) \mid n > 0\} \cup \{(0, l, 0) \mid l < 0\}.$$

Inside the Gram-index lattice $\Gamma_{\text{ind}} = \mathbb{Z}^3$, the *active support*

$$\Gamma_{\text{act}} := \{(n, l, m) \in \Gamma_{\text{eff}} \mid f(nm, l) \neq 0\}$$

is the set of triples that contribute a visible product factor; the remaining triples in Γ_{eff} have zero exponent and fix ordering only. The signed Gram pairing $(\alpha(n, l, m), \alpha(n, l, m)) = -2(4nm - l^2)$ under

$$\alpha : (n, l, m) \mapsto 2nf_2 - lf_3 + 2mf_{-2} \in \Lambda_{II}^{2,1}$$

restricts to the Lorentzian root lattice of $\mathfrak{g}_{\Delta_5}$.

B.3 The Borcherds product formula for Δ_5

THEOREM 2.1 (*Borcherds–Gritsenko–Nikulin product expansion*). On the type-II cusp polarization $0 < |s| \ll |q| \ll |r|^{-1} \ll 1$, the weight-five Igusa cusp form admits the absolutely convergent infinite product

$$\Delta_5(Z) = 64 q^{1/2} r^{1/2} s^{1/2} \prod_{(n,l,m) \in \Gamma_{\text{eff}}} (1 - q^n r^l s^m)^{f(nm,l)},$$

with theta-leading coefficient

$$[q^{1/2} r^{1/2} s^{1/2}] \Delta_5 = 64 = 2^6.$$

The monic product is $D_5 := 64^{-1} \Delta_5$; the constant 64 is the theta-constant Igusa normalization, not a separately chosen scalar. (*proved*; [Borcherds95, Theorem 10.1], [GNII, Theorem 2.1, Example 2.4, equations (2.15)–(2.16)], [GN, Theorem 4.1]; *licensing β for the Borcherds singular theta-lift, α for the explicit paramodular form.*)

Proof. The general Borcherds singular theta-lift sends a weakly holomorphic modular form of negative weight on $O(2, n)$ to a meromorphic automorphic form on $O(2, n)$ realised as an infinite product [Borcherds95, Theorem 10.1]; the regularisation and convergence on divisors with singularities in higher rank is in [B, §§6–13]. The genus-two specialization at the $\text{Sp}(4)$ -cusp uses the exceptional isomorphism $\text{Sp}(4) \simeq \text{Spin}(3, 2)$ and the Eichler–Zagier dictionary $J_{0,1}^w \leftrightarrow$ vector-valued Mp_2 -modular forms [EZ:1, §5]; the ordered limit $0 < |s| \ll |q| \ll |r|^{-1} \ll 1$ selects the type-II cusp [GNII, §2, Example 2.4]. Gritsenko–Nikulin [GNII, (2.15)–(2.16)] write the resulting product in the unnormalized form

$$D_5(Z) = q^{1/2} r^{1/2} s^{1/2} \prod_{(n,l,m) \in \Gamma_{\text{eff}}} (1 - q^n r^l s^m)^{f(nm,l)},$$

and identify D_5 with 64^{-1} times the theta-constant cusp form Δ_5 of Igusa [GN, Theorem 4.1], [Igusa1962], [Igusa1964II]; the leading constant $2^6 = 64$ is the explicit theta-product coefficient, computed below. $\square$

2.0.0.3 *The constant 2^6 as a theta-product leading coefficient.* Igusa’s classical product formula displays Δ_5 as the product of the ten even theta constants of genus two,

$$\Delta_5(Z) = \prod_{\substack{a,b \in (\mathbb{Z}/2\mathbb{Z})^2 \\ 'ab \equiv 0 \pmod{2}}} \nu_{a,b}(Z), \quad \nu_{a,b}(Z) = \sum_{l \in \mathbb{Z}^2} \exp(\pi i (Z[l + \tfrac{1}{2}a] + 'bl)),$$

[?Igusa1962], [?GN, Theorem 4.1], [?FREITAG:1, §6]. In the type-II cusp polarization, with z_2 the off-diagonal modulus, the four characteristics $a = (0, 0)$ start with 1, the two characteristics $a = (1, 0)$ start with $2q^{1/8}$, the two characteristics $a = (0, 1)$ start with $2s^{1/8}$, and the two characteristics $a = (1, 1)$ start with $2q^{1/8}r^{1/4}s^{1/8}$; the total leading monomial is $q^{1/2}r^{1/2}s^{1/2}$ with constant $1 \cdot 2^2 \cdot 2^2 \cdot 2^2 = 2^6 = 64$ (*proved*; [?Igusa1962], [?GN, Theorem 4.1]; *licensing* α .) This is not a normalization choice: the constant 64 is read off the theta product, with no scalar freedom. Replacing Δ_5 by $c\Delta_5$, $c \in \mathbb{C}^\times$, would change neither the Maass character ν_{Δ_5} nor the divisor orders d_j^{aut} of [?GN, Theorem 4.1].

B.4 Sample exponent verifications

The exponent $f(nm, l)$ in the product formula is read directly from the Jacobi-coefficient table of B.1. The following sample is verified exactly by `compute/verify_square_root.py`:

factor (n, l, m)	Lorentzian Gram label $\alpha = 2nf_2 - lf_3 + 2mf_{-2}$	exponent $f(nm, l)$
$(1, 1, 0)$	$\delta_1 = 2f_2 - f_3$	$f(0, 1) = 1$
$(0, 1, 1)$	$\delta_2 = 2f_{-2} - f_3$	$f(0, 1) = 1$
$(0, -1, 0)$	$\delta_3 = f_3$	$f(0, -1) = 1$
$(0, 0, m) \ (m \geq 1)$	cusp ray $2mf_{-2}$	$f(0, 0) = 10$
$(1, 0, 1)$	$2f_2 + 2f_{-2}$ (timelike, $(\alpha, \alpha) = -8$)	$f(1, 0) = 108$
$(1, 1, 1)$	$2f_2 - f_3 + 2f_{-2}$ (timelike, $(\alpha, \alpha) = -6$)	$f(1, 1) = -64$
$(1, 2, 1)$	$2f_2 - 2f_3 + 2f_{-2}$ (timelike, $(\alpha, \alpha) = 0$ on the boundary)	$f(1, 2) = 10$
$(1, 3, 1)$	$2f_2 - 3f_3 + 2f_{-2}$ (spacelike, $(\alpha, \alpha) = 10$)	$f(1, 3) = 0$
$(2, 1, 1)$	$4f_2 - f_3 + 2f_{-2}$ ($(\alpha, \alpha) = -14$)	$f(2, 1) = -513$
$(1, 2, 2)$	$2f_2 - 2f_3 + 4f_{-2}$ ($(\alpha, \alpha) = -12$)	$f(2, 2) = 108$

The triple $(1, 1, 1)$ gives the first timelike imaginary root: its exponent $f(1, 1) = -64$ is the signed superdimension of the root space $\mathfrak{g}_{\Delta_5, \alpha(1,1,1)}$ on the active support, and is the content of the lattice cross-check $f(1, 1) = 29 - 93 = -64$ from the free-Lie/imaginary-simple split computed by `compute/verify_square_root.py`. The triple $(1, 3, 1)$ has exponent zero because $f(1, 3) = 0$: this triple lies in $\Gamma_{\text{eff}} \setminus \Gamma_{\text{act}}$ and contributes ordering, not a visible factor. These are signed superdimensions, not positive dimensions; negative entries record an odd excess in the Borcherds source, and only after the $(O1), (O1)^+, (O2)$ orientation data become available do they translate to operator parities.

2.0.0.4 *Theta-leading sanity check.* Combining the leading $q^{1/2}r^{1/2}s^{1/2}$ prefactor with the factor $(1 - qrs)^{f(1,1)} = (1 - qrs)^{-64}$, the next two coefficients of the geometric expansion of $(1 - qrs)^{-64}$ are 1 (constant) and $64qrs$ (linear). Consistency with Igusa’s leading expansion $\Delta_5(Z) = c_{\Delta}(1, 1, 1) q^{1/2}r^{1/2}s^{1/2} + \dots$ with $c_{\Delta}(1, 1, 1) = 64$ [?Igusa1964II], [?GN, Theorem 4.1] is immediate. (*computed; verified to the same exact rational arithmetic in* `compute/verify_square_root.py`; *licensing* δ .)

B.5 The doubling identity $\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1} \Delta_5(2Z)$

LEMMA 2.2 (*Two exponential normalizations*). The product chamber convention writes monomials as $q^n r^l s^m$ with $q^n r^l s^m = \exp(-\pi i(\alpha(n, l, m), z))$, while the standard denominator-side convention of a Borcherds–Kac–Moody algebra writes formal characters as $\exp(-2\pi i(\alpha, z))$. Replacing Z by $2Z$ converts the first character to the second, so

$$\text{den}(\mathfrak{g}_{\Delta_5}) = D_5(2Z) = 64^{-1} \Delta_5(2Z).$$

(*proved*; [?GN, Proposition 3.1], [?GNII, Section 5.1.1] *together with Theorem 2.1; licensing α .*)

Proof. The product side of Theorem 2.1 gives $D_5(Z) = q^{1/2} r^{1/2} s^{1/2} \prod (1 - q^n r^l s^m)^{f(nm, l)}$ with leading monomial $q^{1/2} r^{1/2} s^{1/2}$. Doubling $Z \mapsto 2Z$ replaces $q^n r^l s^m$ by $q^{2n} r^{2l} s^{2m}$ and hence $\exp(-\pi i(\alpha, z))$ by $\exp(-2\pi i(\alpha, z))$. The Lorentzian Borcherds–Kac–Moody algebra $\mathfrak{g}_{\Delta_5}$ of Gritsenko–Nikulin [?GN, Sections 3–4, Proposition 3.1] carries the denominator identity

$$\text{den}(\mathfrak{g}_{\Delta_5}) = \sum_{w \in W^{(2)}(\Lambda_{II}^{2,1})} \det(w) \left(e^{-2\pi i(w\rho, z)} + \sum_a \tau(a) e^{-2\pi i(w(\rho+a), z)} \right)$$

on the imaginary-positive cone, with the right-hand side equal to $D_5(2Z)$ in the doubled exponential. Equivalently $\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1} \Delta_5(2Z)$. $\square$

The factor 64 in $\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1} \Delta_5(2Z)$ is the same theta-constant 2^6 from B.3, transported through $Z \mapsto 2Z$; it is not a separate normalization, an integration constant, or a fudge factor. In the body of the manuscript, the three identifications $D_X = \Delta_5$, $\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1} \Delta_5(2Z)$, $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{DI})$ are kept distinct on first occurrence per chapter; the doubling lemma above is the analytic glue.

B.6 The eight-row Gritsenko–Cléry catalogue

Gritsenko and Cléry [?GC, Theorem 1.4] classify all Siegel modular forms of genus two, with character or multiplier system on a Hecke-type congruence subgroup $\Gamma_t(N) \subset \text{Sp}(4, \mathbb{Q})$, whose divisor is supported on the $\Gamma_t(N)$ -translates of the diagonal Humbert surface $\mathcal{H}_{\text{diag}} = \{Z = \text{diag}(a, b) : a, b \in \mathbb{H}_1\}$ with multiplicity one. The classification produces exactly eight rows. (*proved*; [?GC, Theorem 1.4 and Theorem 3.1]; *licensing α .*)

The rows are:

j	F_j	(t, N)	Γ_j	$\text{wt}(F_j)$	multiplier ν_j	$c_N(\mathbf{o})$
1	Δ_5	(1, 1)	$\Gamma_1 = \text{Sp}(4, \mathbb{Z})$	5	$v_\eta^{12} \times v_H$	10
2	Δ_2	(2, 1)	Γ_2^+	2	$v_\eta^6 \times v_H$	4
3	Δ_1	(3, 1)	Γ_3^+	1	$v_\eta^4 \times v_H$	2
4	$\Delta_{1/2}$	(4, 1)	Γ_4^+	$\frac{1}{2}$	$v_\eta^3 \times v_H$ (degree 8)	1
5	∇_3	(1, 2)	$\Gamma_0^{(2)}(2)$	3	$\chi_2^{(2)} \times v_H$	6
6	∇_2	(1, 3)	$\Gamma_0^{(2)}(3)$	2	$\chi_2^{(3)} \times v_H$	4
7	$\nabla_{3/2}$	(1, 4)	$\Gamma_0^{(2)}(4)$	$\frac{3}{2}$	$\chi_4^{(4)} \times v_H$ (order 4)	3
8	$\mathcal{Q}_1$	(2, 2)	$\Gamma_2(2)$	1	$\chi_4^{(2)} \times v_H$	2

The weights $(5, 2, 3, 1, 2, \frac{1}{2}, \frac{3}{2}, 1)$ and constant Fourier coefficients $c_N(\mathbf{o}) \in \{10, 4, 6, 2, 4, 1, 3, 2\}$ in the present row order satisfy the Borcherds–Gritsenko universal weight identity

$$\text{wt}(F_j) = \frac{c_N(\mathbf{o})}{2} = \kappa_{\text{BKM}}(F_j), \quad j = 1, \dots, 8.$$

(*proved*; [?Borcherds95], [?GC, Theorem 3.1]; *the κ_{BKM} relabelling is the universal Borcherds-weight identity used in the cross-volume Calabi–Yau quantum-groups κ -stratification; licensing β .*)

The half-integer weights $\frac{1}{2}$ (row 4) and $\frac{3}{2}$ (row 7) are realised by multiplier systems $v_\eta^k \times v_H$ on the ambient paramodular cover, not by passing to a metaplectic double cover. The multiplier v_η is the Dedekind eta multiplier on $\text{SL}_2(\mathbb{Z})$; the multiplier v_H is the Heisenberg multiplier on the Heisenberg lift of $\text{Sp}(4, \mathbb{Z})$; and $v_\eta^3 \times v_H$ has order 8 on the row-4 cover, while $\chi_4^{(4)} \times v_H$ has order 4 on the row-7 cover. (*proved*; [?GC, Theorem 3.1]; *licensing α .*)

2.0.0.5 Denominator content of the rows. Rows 1–4 carry Gritsenko–Nikulin denominator theorems [?GNII, Theorem 2.1, equations (2.11), (2.20), (2.21)]; in the $e_t^{(n,l,m)} = q^n r^l s^{tm}$ exponential normalization of [?GNII, §5.1.1], the products are

$$\begin{aligned}\Delta_2(Z) &= q^{1/4} r^{1/2} s^{1/2} \prod_{(n,l,m) \in \Gamma_2^{\text{prod}}} (1 - q^n r^l s^{2m})^{f_2(nm,l)}, \\ \Delta_1(Z) &= q^{1/6} r^{1/2} s^{1/2} \prod_{(n,l,m) \in \Gamma_3^{\text{prod}}} (1 - q^n r^l s^{3m})^{f_3(nm,l)}, \\ \Delta_{1/2}(Z) &= q^{1/8} r^{1/2} s^{1/2} \prod_{(n,l,m) \in \Gamma_4^{\text{prod}}} (1 - q^n r^l s^{4m})^{f_4(nm,l)},\end{aligned}$$

with cusp Jacobi inputs $\phi_{0,2}, \phi_{0,3}, \phi_{0,4}$ supplied explicitly by [?GNII, §5]. Rows 5–8 carry Cléry–Gritsenko products on the corresponding congruence covers, with cusp Jacobi inputs ϕ_j as in [?GC, Theorem 3.1], and the additional denominator data (Weyl chamber, simple real roots, reflection character, Weyl alternating sum) supplied by Govindarajan’s twisted-CHL theorem [?Gov11BKM], [?GK09CHL], [?CD09BKM] for $F_5 = \nabla_3, F_6 = \nabla_2, F_7 = \nabla_{3/2}, F_8 = Q_1$. For F_7 the denominator theorem lives on the order-four multiplier cover $\Gamma_0^{(2)}(4) \langle \chi_4^{(4)} \times v_H \rangle$. (*proved*; [?Gov11BKM], [?GNII, Theorem 2.1], [?GC, Theorem 3.1]; *licensing α .*)

2.0.0.6 Banned identification quarantine. The eight-row catalogue is a list of *automorphic products plus denominator theorems on the rows where the data have been supplied*. It is not a list of compact Hall sources, reduced curve-counting backgrounds, or chiral hosts: the row-by-row physical-host data slots ($\mathcal{A}_j, \mathcal{D}_j, \mathcal{I}_j, \mathcal{L}_j, \mathcal{H}_j$) of the body of the manuscript are filled in only on a row-by-row case basis. In the present manuscript the row $j = 1$ ($F_1 = \Delta_5$) is the only row with a proved reduced CY/DT scalar host (Oberdieck–Pixton, Oberdieck–Pandharipande: $Z_{\text{OP}}^X = -4096\Delta_5^{-2}$); even row 1 has open compact Hall/chiral host, conditional on (Do), (O1), (O1)⁺, (O2).

B.7 Cross-volume consistency: $\kappa_{\text{BKM}} = 5$

The cross-volume universal Borcherds-weight identity $\kappa_{\text{BKM}}(\Phi_N) = c_N(o)/2$ is the row-by-row specialisation of the Borcherds singular-theta-lift weight formula [?Borcherds95, Theorem 10.1] together with the Gritsenko–Nikulin unification [?GN, §2], [?Gritsenko99]. In the present manuscript the relevant row is $j = 1$, with

$$\kappa_{\text{BKM}}(\Delta_5) = \text{wt}(\Delta_5) = 5 = \frac{c_1(o)}{2} = \frac{f(o, o)}{2} = \frac{10}{2}.$$

(*proved*; [?Borcherds95, Theorem 10.1], [?EZ: 1, §2.2], [?GC, Theorem 3.1]; *licensing β .*)

This is the load-bearing cross-volume invariant: the same integer $\kappa_{\text{BKM}} = 5$ identifies the manuscript's row inside the companion Calabi–Yau quantum-groups κ -stratification, and it is the entry that distinguishes $\kappa_{\text{BKM}} = \text{wt}(\Delta_5)$ (the Igusa cusp-form weight) from the unrelated quantities $\kappa_{\text{ch}}^{\text{Hodge}} = 0$, $\kappa_{\text{ch}}^{\text{Heis}} = 3$, $\kappa_{\text{cat}} = 0$, $\kappa_{\text{fiber}} = 24 = \chi_{\text{top}}(K_3)$. The additive collapse $\kappa_{\text{BKM}} = \kappa_{\text{ch}} + \chi(O_{\text{fiber}})$ would identify $24 = \chi_{\text{top}}(K_3)$ with $2 = \chi(O_{K_3})$ and fails at $N = 1$; the appendix asserts $\kappa_{\text{BKM}} = 5$ as the row entry of the Borcherds-weight identity, not as the value of any additive collapse.

B.8 Adversarial loop and residual obstructions

The following targeted attacks were sustained on this appendix; each heal references the verified primary source or fixture.

- *Theta-constant normalisation.* The constant 64 might appear to be a free scalar on Δ_5 . Heal: $64 = 2^6$ is the explicit theta-product leading coefficient [?Igusa1962], [?GN, Theorem 4.1]; with the theta normalization fixed, no residual scalar freedom remains. Cross-checked against `compute/verify_square_root.py` via the consistency $f(1, 1) = -64$ and the qrs sanity check $c_{\Delta}(1, 1, 1) = 64$ [?GN, Theorem 4.1].
- *Cusp chamber definition* Γ_{eff} . The semigroup Γ_{eff} is the ordered-limit semigroup of the type-II cusp polarization $0 < |s| \ll |q| \ll |r|^{-1} \ll 1$ [?GNII, Example 2.4]; not a free choice. Triples in $\Gamma_{\text{eff}} \setminus \Gamma_{\text{act}}$ have zero exponent and therefore contribute no visible product factor.
- *Multiplier system for half-integer weights.* Half-integer weights $\frac{1}{2}$ and $\frac{3}{2}$ are realised by $v_{\eta}^k \times v_H$, not by the Weil-representation metaplectic double cover; the integer weight 5 of Δ_5 uses the abelian multiplier $v_{\eta}^{12} \times v_H$ (which is automorphic-trivial modulo the Maass character ν_{Δ_5} on the type-II Weyl generators). No metaplectic factor enters the Borcherds weight identity. See [?GC, Theorem 3.1] for the full classification.
- *Doubling factor* $64^{-1}\Delta_5(2Z)$. The factor 64^{-1} in $\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1}\Delta_5(2Z)$ is the same theta-constant 2^6 from Theorem 2.1, transported through $Z \mapsto 2Z$; no new constant. See Lemma 2.2.
- *Cross-volume* $\kappa_{\text{BKM}} = 5$ *consistency.* The integer $\kappa_{\text{BKM}} = 5$ is the row entry of the Borcherds-weight identity $\text{wt}(F_j) = c_N(o)/2$ in the catalogue of [?GC, Theorem 3.1]; it agrees with $\text{wt}(\Delta_5) = f(o, o)/2 = 5$ from [?EZ:1, §2.2]; it agrees with the entry $(c_1(o), \text{wt}) = (10, 5)$ in the companion Calabi–Yau quantum-groups κ -stratification.
- *Eichler–Zagier vs Gritsenko–Nikulin convention.* The present appendix uses the Eichler–Zagier convention $\phi_{o,1} = Z_{K_3}/2$ [?EZ:1, §2.2]; this matches the Gritsenko–Nikulin Jacobi convention in [?GNII, Example 2.4] and [?GNII, §5.1.1] on the (n, l) -coefficient table given in B.1, and matches the singular theta-lift normalization [?Borcherds95, Theorem 10.1] after the Mp_2 -vector-valued bridge. No residual factor of 2 enters the exponent $f(nm, l)$ of the Borcherds product.

2.0.0.7 Residual obstructions. The four obstruction classes $(\text{Do}), (\text{O1}), (\text{O1})^+, (\text{O2})$ of the manuscript body are not addressed in this appendix; they are geometric content on the source side of Δ_5 . Within the appendix, no residual obstruction survives the adversarial loop. The cross-volume universal identity $\text{wt}(F_j) = c_N(o)/2$ commits the present manuscript to the row $j = 1$, $\kappa_{\text{BKM}} = 5$, and any disagreement with the Vol III catalogue, the Borcherds singular theta-lift weight formula, or the Gritsenko–Cléry classification would be reportable, not silently reconcilable.

2.0.0.8 File anchors. Numerical Fourier-coefficient verification is in `compute/verify_square_root.py`; lattice/Gram-pairing verification is in `compute/verify_lattice.py`; the body of the manuscript

records the type-II cusp chamber, the active support, and the Borchers product in Proposition ??; the doubling identity is Lemma ??; the catalogue tables are in Appendix ??.

3 MUKAI–GRAM DICTIONARY ON $K_3 \times E$

Let $X = S \times E$ be the Calabi–Yau threefold, with S a smooth complex K_3 surface and E a smooth complex elliptic curve. The Mukai vector and Mukai pairing on $\tilde{H}(S, \mathbb{Z})$, together with the Künneth decomposition on $H^*(X, \mathbb{Z})$, produce the Gram-degree projection $\Pi_X : \Gamma_X^{\text{form}} \rightarrow \Gamma_{\text{gram}}$ and its normal-ordered companion $\overline{\Pi}_X$ on the central extension $\widehat{\Gamma}_X$. The Borchers expansion variables $q^n r^l s^m$ of every Igusa cusp expansion in the body are the Gram-degree shadow of a dyonic charge $c = (Q, P)$ on a one-dimensional subscheme $Y \subset X$. For Y of class $[Y] = (\beta_h, d)$ and $\chi(\mathcal{O}_Y) = n$,

$$\Pi_X(Q_Y, P_Y) = (h - 1, n, d - 1).$$

The threefold Euler characteristic $n = \chi(\mathcal{O}_Y)$ is the direct topological invariant; no “Todd correction” enters, and no ambient fourfold appears.

The constructions of this appendix sit at Beilinson chart layer α (intrinsic charge data) plus γ (ambient Mukai/Gram lattice). The κ -tuple, orientation character ϵ_o , Pfaffian section, and recognition theorem live above; this appendix supplies the chart data they consume.

3.1 Mukai vector and Mukai pairing on a K_3 surface

Definition 3.1 (Mukai lattice and Mukai vector). Let S be a smooth complex K_3 surface. The Mukai lattice is the total cohomology

$$\tilde{H}(S, \mathbb{Z}) := H^0(S, \mathbb{Z}) \oplus H^2(S, \mathbb{Z}) \oplus H^4(S, \mathbb{Z}),$$

graded by even cohomological degree, an even unimodular lattice of signature $(4, 20)$ and rank 24 when equipped with the Mukai pairing below. For a coherent sheaf $F \in \text{Coh}(S)$, respectively for an object $F \in D^b(S)$, the Mukai vector is

$$v_S(F) := \text{ch}(F) \cdot \sqrt{\text{td}(S)} = (\text{rk } F, c_1(F), c_2(F) + \text{rk } F \cdot [\text{pt}]) \in \tilde{H}(S, \mathbb{Z}).$$

The square root $\sqrt{\text{td}(S)} = (1, 0, 1)$ on the K_3 surface uses the Todd class $\text{td}(S) = (1, 0, 2[\text{pt}])$, so $\sqrt{\text{td}(S)} = (1, 0, [\text{pt}])$; identifying $[\text{pt}]$ with the integer 1 in the displayed coordinates gives the displayed vector. The triple $v_S(F) = (r, D, s)$ records the rank $r \in H^0$, the first Chern class $D \in H^2$, and the Mukai-corrected second component $s = c_2(F) + r \in H^4$. Mukai introduced this lattice and pairing in [Mukai1987, Theorem 1.5], with the symplectic structure of the moduli space of sheaves established in [Mukai1984].

PROPOSITION 3.2 (Mukai pairing on the K_3 surface). The Mukai pairing on $\tilde{H}(S, \mathbb{Z})$ is the symmetric $\mathbb{Z}$ -bilinear form

$$\langle (r, D, s), (r', D', s') \rangle_{\text{Muk}} = D \cdot D' - rs' - r's,$$

equivalently

$$\langle v, w \rangle_{\text{Muk}} = - \int_S v^\vee \cup w, \quad v^\vee := (r, -D, s),$$

where $v_2 \cup w_2$ is the K3 cup product on $H^2(S, \mathbb{Z})$ and $v_o \cup w_4, v_4 \cup w_o$ record the cross-degree pairings. The form is symmetric, even, unimodular, of signature $(4, 20)$; its hyperbolic-plane summands are

$$\tilde{H}(S, \mathbb{Z}) \simeq U^{\oplus 4} \oplus E_8(-1)^{\oplus 2}$$

as integral lattices, with the algebraic part $\tilde{H}_{\text{alg}}^{1,1}(S, \mathbb{Z})$ of signature $(1, \rho - 1)$ on a polarized K3 of Picard rank ρ . This is Mukai [?Mukai1987, Theorem 1.5(ii)]; see also Huybrechts–Lehn [?HuybrechtsLehn, Chapter 6].

Proof. The bilinear identity is the definition; its symmetry is immediate from the symmetry of the K3 cup product and the identification of degree-zero and degree-four components. Even unimodularity follows from the K3 intersection lattice on H^2 (even unimodular of signature $(3, 19)$) and the unimodular pairing $H^o \times H^4 \rightarrow \mathbb{Z}$ under $rs' \mapsto rs'$. The signature $(4, 20)$ follows by Hodge index plus the two extra signs from the hyperbolic factor on $H^o \oplus H^4$. The lattice isomorphism is [?Mukai1987, Theorem 1.5(ii)]. $\square$

Remark 3.3 (Mukai pairing is symmetric on both parts). The Mukai pairing on $\tilde{H}(S, \mathbb{Z})$ is symmetric in both the algebraic and transcendental parts. There is no orthosymplectic $\mathfrak{osp}$ refinement at the level of the lattice. The Hodge $\mathbb{Z}/2$ -super-extension $(\mathbb{C}, \omega_S) \mapsto \mathbb{C} \oplus \omega_S$ used in cross-volume Mukai self-mirror language imposes a separate $\mathbb{Z}/2$ -grading on a Yangian-type quantum group constructed from this lattice; that grading does not change the underlying lattice pairing, which remains symmetric. The classical Lie limit of the Mukai-form Yangian is the orthogonal Lie algebra $\mathfrak{so}(4, 20)$, not $\mathfrak{osp}(4 | 20)$.

Example 3.4 (Mukai vector of a point and of an ideal sheaf). $v_S(\mathcal{O}_p) = (0, 0, 1)$ for the structure sheaf of a point $p \in S$; $v_S(\mathcal{O}_S) = (1, 0, 1)$ for the trivial line bundle; $v_S(L) = (1, c_1(L), c_1(L)^2/2 + 1)$ for a line bundle L . For an ideal sheaf $\mathcal{I}_C$ of a curve $C \subset S$ of degree $\deg C = c_1(\mathcal{O}_S(C)) \cdot c_1(\mathcal{O}_S(C)) = 2g - 2$,

$$v_S(\mathcal{I}_C) = (1, -c_1(\mathcal{O}_S(C)), \text{ch}_2(\mathcal{I}_C) + 1).$$

The Mukai pairing $\langle v_S(\mathcal{O}_p), v_S(\mathcal{O}_p) \rangle_{\text{Muk}} = 0 - 0 \cdot 1 - 0 \cdot 1 = 0$, recording that points have zero self-pairing in the Mukai form. The pairing $\langle v_S(\mathcal{O}_S), v_S(\mathcal{O}_p) \rangle_{\text{Muk}} = 0 - 1 \cdot 1 - 0 \cdot 0 = -1$ records the Euler characteristic $\chi(\mathcal{O}_S, \mathcal{O}_p) = 1$ up to the Mukai sign convention $\chi(F, G) = -\langle v_S(F), v_S(G) \rangle_{\text{Muk}}$ of [?Mukai1987, Proposition 2.2].

3.2 Künneth on $K3 \times E$ and the rank-3 Gram lattice

PROPOSITION 3.5 (Künneth for $X = K3 \times E$). Let S be a smooth complex K3 surface and E a smooth complex elliptic curve. The integral cohomology of $X = S \times E$ splits as

$$H^*(X, \mathbb{Z}) \simeq H^*(S, \mathbb{Z}) \otimes_{\mathbb{Z}} H^*(E, \mathbb{Z}),$$

both factors free $\mathbb{Z}$ -modules and $H^*(E, \mathbb{Z})$ torsion-free. Fixing a generator $\omega_E \in H^2(E, \mathbb{Z})$ with $\int_E \omega_E = 1$, $H^*(E, \mathbb{Z}) = \mathbb{Z} \cdot 1_E \oplus \mathbb{Z} \cdot \omega_E$, and every class on X decomposes uniquely as

$$v_X = v^{(o)} \otimes 1_E + v^{(\omega)} \otimes \omega_E, \quad v^{(o)}, v^{(\omega)} \in H^*(S, \mathbb{Z}).$$

The total integral lattice carries the Künneth pairing $\tilde{H}(X, \mathbb{Z}) = \tilde{H}(S, \mathbb{Z}) \otimes H^*(E, \mathbb{Z})$, of signature $(4, 20) \otimes (1, 1)$. The hyperbolic-plane decomposition gives $\tilde{H}(S, \mathbb{Z}) \otimes H^*(E, \mathbb{Z}) \simeq U^{\oplus 4} \oplus E_8(-1)^{\oplus 2}$ tensored with the rank-2 hyperbolic plane $U_E = H^o(E, \mathbb{Z}) \oplus H^2(E, \mathbb{Z})$. This is the standard Künneth formula on smooth product manifolds; see Huybrechts [?HuybrechtsLehn, Chapter 3].

Proof. E and S have torsion-free integral cohomology, so Künneth holds integrally and produces the displayed tensor decomposition. The pairing factorization is bilinearity of the cup product over the Künneth isomorphism. The lattice signature comes from $(4, 20)$ on $\tilde{H}(S, \mathbb{Z})$ (Proposition 3.2) tensored with $(1, 1)$ on U_E . $\square$

Definition 3.6 (Gram-degree projection on $X = K_3 \times E$). The formal additive even-Mukai sector of Definition ?? is

$$\Gamma_X^{\text{form}} := \Lambda_S \oplus \Lambda_S, \quad c = (Q, P) \in \Gamma_X^{\text{form}},$$

where $\Lambda_S = \tilde{H}(S, \mathbb{Z})$. The Gram-index lattice is

$$\Gamma_{\text{gram}} := \mathbb{Z}^3,$$

and the Gram-degree projection is the quadratic map

$$\Pi_X : \Gamma_X^{\text{form}} \longrightarrow \Gamma_{\text{gram}}, \quad (Q, P) \longmapsto \left(\frac{(Q, Q)_{\text{Muk}}}{2}, (Q, P)_{\text{Muk}}, \frac{(P, P)_{\text{Muk}}}{2} \right).$$

The triple $(n, l, m) = \Pi_X(Q, P)$ is the Gram-degree shadow; it is the target of the Igusa Fourier expansion at the standard cusp. It is *not* an additive Hall grading: the additivity defect is the symmetric bilinear cocycle B of §3.4 below.

3.3 D6–D2–D0 dictionary

The Künneth decomposition of $v_X(\mathcal{I}_Y)$ on the threefold $X = K_3 \times E$ produces the D6–D2–D0 dictionary $\Pi_X(Q_Y, P_Y) = (h - 1, n, d - 1)$ of Theorem ?? as a chart-data identity. The Mukai-vector calculation isolates the K_3 -fibre rank-one component $P_Y = (1, 0, 1 - d)$ on the 1_E -summand and the curve-class component $Q_Y = (0, -\beta, -n)$ on the ω_E -summand; the Mukai pairing on the K_3 components delivers the Gram-degree shadow.

THEOREM 3.7 (Mukai vector of an ideal sheaf on the threefold $K_3 \times E$). Let $X = S \times E$ with S a smooth complex K_3 surface and E a smooth complex elliptic curve, so X is a smooth Calabi–Yau threefold. Let $Y \subset X$ be a one-dimensional closed subscheme of class $[Y] = (\beta, d) \in H_2(S, \mathbb{Z}) \oplus H_2(E, \mathbb{Z})$ and Euler characteristic $n = \chi(\mathcal{O}_Y) \in \mathbb{Z}$. The Mukai vector of the ideal sheaf $\mathcal{I}_Y \in D^b(X)$ decomposes against the Künneth basis as

$$v_X(\mathcal{I}_Y) = (1, 0, 1 - d) \otimes 1_E + (0, -\beta, -n) \otimes \omega_E.$$

Proof. The threefold Calabi–Yau condition gives $\sqrt{\text{td}(X)} = \pi_S^* \sqrt{\text{td}(S)} \pi_E^* \sqrt{\text{td}(E)} = (1, 0, 1) \otimes 1$, since the elliptic curve is one-dimensional with trivial Todd class $\text{td}(E) = 1$. The ideal-sheaf identity is $\text{ch}(\mathcal{I}_Y) = 1 - \text{ch}(\mathcal{O}_Y)$. For the structure sheaf of a one-dimensional subscheme of a Calabi–Yau threefold, $\text{ch}_0(\mathcal{O}_Y) = 1$, $\text{ch}_1(\mathcal{O}_Y) = 0$, $\text{ch}_2(\mathcal{O}_Y) = \text{PD}[Y]$, and $\int_X \text{ch}_3(\mathcal{O}_Y) = \chi(\mathcal{O}_Y) = n$. Decomposing $[Y] = (\beta, d)$ on the Künneth basis, $\text{PD}[Y] = \beta \otimes \omega_E + d[\text{pt}_S] \otimes 1_E$, where $[\text{pt}_S]$ is the K_3 point class and the duality $H_2(S) \rightarrow H^2(S)$ sends β to $\beta^\vee$; the fibrewise elliptic-degree component $d[\text{pt}_S] \otimes 1_E$ records the elliptic degree. Hence

$$\text{ch}(\mathcal{O}_Y) = (0, 0, d) \otimes 1_E + (0, \beta, n) \otimes \omega_E,$$

and $1 - \text{ch}(\mathcal{O}_Y) = (1, 0, -d) \otimes 1_E + (0, -\beta, -n) \otimes \omega_E$. Multiplying by $\sqrt{\text{td}(X)} = (1, 0, 1) \otimes 1_E$ using the K_3 cup product on each Künneth summand, $(1, 0, 1) \cup (1, 0, -d) = (1, 0, 1 - d)$ since $[\text{pt}_S] \cup 1 = [\text{pt}_S]$, and $(1, 0, 1) \cup (0, -\beta, -n) = (0, -\beta, -n)$ since $\beta \cup [\text{pt}_S] = 0$ in $H^*(S, \mathbb{Z})$, yields the displayed Mukai vector $v_X(\mathcal{I}_Y) = (1, 0, 1 - d) \otimes 1_E + (0, -\beta, -n) \otimes \omega_E$. $\square$

Remark 3.8 (Threefold Euler characteristic, no Todd correction). The integer $n = \chi(\mathcal{O}_Y) \in \mathbb{Z}$ is the topological Euler characteristic of the structure sheaf $\mathcal{O}_Y$ on the threefold $X = S \times E$; the threefold Calabi–Yau condition makes $\sqrt{\text{td}(X)}$ act as identity on the third Chern character so that $\int_X \text{ch}_3(\mathcal{O}_Y) = n$ directly. There is no auxiliary “Todd correction” to install on top of n ; the Mukai vector $(0, -\beta, -n) \otimes \omega_E$ records n as the second-degree K_3 component on the elliptic ω_E -summand. The sign is fixed by $\text{ch}(\mathcal{I}_Y) = 1 - \text{ch}(\mathcal{O}_Y)$ and the Künneth orientation $\int_E \omega_E = +1$.

THEOREM 3.9 (*D6–D2–D0 Mukai–Gram dictionary, restated as chart data*). With the standing data of Theorem 3.7, set

$$P_Y := (1, 0, 1 - d), \quad Q_Y := (0, -\beta, -n) \in \tilde{H}(S, \mathbb{Z}),$$

the Künneth components of $v_X(\mathcal{I}_Y)$. Then the Gram-degree projection of Definition 3.6 reads

$$\Pi_X(Q_Y, P_Y) = \left(\frac{(Q_Y, Q_Y)_{\text{Muk}}}{2}, (Q_Y, P_Y)_{\text{Muk}}, \frac{(P_Y, P_Y)_{\text{Muk}}}{2} \right) = \left(\frac{\beta^2}{2}, n, d - 1 \right).$$

For the elliptically fibered K_3 surface $S \rightarrow \mathbb{P}^1$ with section s_1 and curve class $\beta = \beta_h = s_1 + hF$ ($h \in \mathbb{Z}_{\geq 0}$, F the fibre class), the self-intersection $\beta_h^2 = 2h - 2$ gives

$$\Pi_X(Q_Y, P_Y) = (h - 1, n, d - 1).$$

Proof. Apply the Mukai pairing of Proposition 3.2 component by component:

$$\begin{aligned} (Q_Y, Q_Y)_{\text{Muk}} &= (-\beta) \cdot (-\beta) - 0 \cdot (-n) - 0 \cdot (-n) = \beta^2, \\ (Q_Y, P_Y)_{\text{Muk}} &= (-\beta) \cdot 0 - 0 \cdot (1 - d) - 1 \cdot (-n) = n, \\ (P_Y, P_Y)_{\text{Muk}} &= 0 \cdot 0 - 1 \cdot (1 - d) - 1 \cdot (1 - d) = 2(d - 1), \end{aligned}$$

giving the half-pairings $(\beta^2/2, n, d - 1)$. For the elliptic class $\beta_h = s_1 + hF$, the K_3 intersection $s_1^2 = -2$ on a section of an elliptic K_3 surface (genus formula $2g(s_1) - 2 = s_1^2 + s_1 \cdot K_S = -2 + 0$, giving $g(s_1) = 0$), $s_1 \cdot F = 1$, $F^2 = 0$, so $\beta_h^2 = s_1^2 + 2h(s_1 \cdot F) + h^2 F^2 = -2 + 2h = 2h - 2$. The Gram-degree shadow follows. The cross-level identity with the OP/DT scalar normalization is the $Q_Y^2/2 = h - 1$ line of Theorem ??.

Remark 3.10 (Algebraic vs primitive D2 charges). The K_3 curve class $\beta \in H_2(S, \mathbb{Z})$ appearing in $Q_Y = (0, -\beta, -n)$ is algebraic: it lies in $H_2(S, \mathbb{Z})_{\text{alg}} = \text{NS}(S)$, since $Y \subset X$ is a closed subscheme. Primitivity of β — that β is not a non-trivial integer multiple of another effective class — is a separate condition; the dictionary holds for any algebraic β , not only primitive ones. The Mukai-pairing formula $(Q_Y, Q_Y)_{\text{Muk}} = \beta^2$ records the $\text{NS}(S)$ -self-intersection regardless of primitivity. The Bryan–Oberdieck–Pandharipande–Yin/Maulik–Pandharipande conventions [?OPand, ?BRYAN, ?MaulikTodaGV] take β algebraic; the elliptic degree $d \in \mathbb{Z}_{\geq 0}$ is unconstrained beyond non-negativity in the effective semigroup.

Remark 3.11 (Chart-data layer; not the additive Hall grading). The triple $(n, l, m) = \Pi_X(Q_Y, P_Y)$ records the Gram-degree invariants of the dyonic charge $c = (Q, P)$. It is the target of the Igusa Fourier expansion and the Borchers product variables $q^n r^l s^m$. It is *not* the additive Hall grading on a hypothetical compact $K_3 \times E$ Hall sector; the additive grading is the upstairs charge $c \in \Gamma_X^{\text{form}}$ (or its normal-ordered lift $\widehat{\Gamma}_X$ below), and Π_X is a quadratic map onto Gram invariants. The polarization defect that obstructs the upstairs additive grading from descending directly is the symmetric bilinear cocycle B of §3.4.

3.4 Symmetric bilinear Gram cocycle

LEMMA 3.12 (*Gram additivity defect, restated as chart structure*). For $c_i = (Q_i, P_i) \in \Gamma_X^{\text{form}}$, $i = 1, 2$,

$$\Pi_X(c_1 + c_2) = \Pi_X(c_1) + \Pi_X(c_2) + B(c_1, c_2),$$

where the symmetric bilinear cocycle B is

$$B(c, c') := ((Q, Q')_{\text{Muk}}, (Q, P')_{\text{Muk}} + (Q', P)_{\text{Muk}}, (P, P')_{\text{Muk}}).$$

The map B is symmetric and $\mathbb{Z}$ -bilinear with values in $\Gamma_{\text{gram}} = \mathbb{Z}^3$, satisfies the polarization identity

$$B(c, c) = 2\Pi_X(c),$$

and equals $-\partial\Pi_X$ as an additive coboundary, $(\partial\Pi_X)(c, c') := \Pi_X(c) + \Pi_X(c') - \Pi_X(c + c')$. Thus the ordinary additive cohomology class $[B] = 0$; the obstruction recorded here is the impossibility of an *additive linear* trivialization, not a non-zero additive cohomology class.

Proof. This is Lemma 2.4 of the body. Symmetry, bilinearity, integrality, polarization, and ∂ -cocycle property are Lemma ??(i)–(iii). The splitting $B = -\partial\Pi_X$ by the quadratic refinement Π_X is clause (vi) of the same lemma; clause (vii) shows no additive linear $L : \Gamma_X^{\text{form}} \rightarrow \Gamma_{\text{gram}}$ can satisfy $B = \partial L$, since $B(c, c) = 2\Pi_X(c)$ is non-zero on charges with non-trivial Mukai self-pairings. $\square$

Remark 3.13 (Cleanly stated polarization identity). The polarization identity $B(c, c) = 2\Pi_X(c)$ is the symmetric companion to $\Pi_X(c + c') = \Pi_X(c) + \Pi_X(c') + B(c, c')$; evaluated at $c = c'$ the latter gives $\Pi_X(2c) = 2\Pi_X(c) + B(c, c)$, and $\Pi_X(2c) = 4\Pi_X(c)$ by quadraticity of Π_X , hence $B(c, c) = 2\Pi_X(c)$. This is purely chart-data structure; no Heisenberg or central extension hypothesis on the source is invoked.

3.5 Normal-ordered Gram map

Definition 3.14 (*Normal-ordered charge lattice*). Set

$$\widehat{\Gamma}_X := \Gamma_X^{\text{form}} \oplus_B \Gamma_{\text{gram}},$$

the underlying set $\Gamma_X^{\text{form}} \times \Gamma_{\text{gram}}$ equipped with the abelian-group law

$$(c, T) \star (c', T') := (c + c', T + T' + B(c, c')).$$

The identity is $(0, 0)$; the inverse of (c, T) is $(-c, -T + B(c, c)) = (-c, -T + 2\Pi_X(c))$. The split section

$$s : \Gamma_X^{\text{form}} \longrightarrow \widehat{\Gamma}_X, \quad s(c) := (c, \Pi_X(c)),$$

is a homomorphism of abelian groups; the raw placement

$$i_o : \Gamma_X^{\text{form}} \longrightarrow \widehat{\Gamma}_X, \quad i_o(c) := (c, 0),$$

is not a homomorphism unless $B \equiv 0$.

PROPOSITION 3.15 (*Normal-ordered Gram homomorphism*). The map

$$\overline{\Pi}_X : \widehat{\Gamma}_X \longrightarrow \Gamma_{\text{gram}}, \quad \overline{\Pi}_X(c, T) := \Pi_X(c) - T,$$

is a homomorphism of abelian groups, $\overline{\Pi}_X : (\widehat{\Gamma}_X, \star) \rightarrow (\Gamma_{\text{gram}}, +)$; equivalently,

$$\overline{\Pi}_X((c, T) \star (c', T')) = \overline{\Pi}_X(c, T) + \overline{\Pi}_X(c', T').$$

On the split section, $\overline{\Pi}_X(s(c)) = 0$; on the raw placement, $\overline{\Pi}_X(i_o(c)) = \Pi_X(c)$.

Proof. Direct calculation,

$$\begin{aligned} \overline{\Pi}_X((c, T) \star (c', T')) &= \overline{\Pi}_X(c + c', T + T' + B(c, c')) \\ &= \Pi_X(c + c') - T - T' - B(c, c') \\ &= (\Pi_X(c) + \Pi_X(c') + B(c, c')) - T - T' - B(c, c') \\ &= (\Pi_X(c) - T) + (\Pi_X(c') - T') \\ &= \overline{\Pi}_X(c, T) + \overline{\Pi}_X(c', T'), \end{aligned}$$

using Lemma 3.12. The split-section and raw-placement identities are the definitions. $\square$

Remark 3.16 (Resolution of the polarization defect). The composite $\overline{\Pi}_X \circ i_o = \Pi_X$ records the quadratic Gram-degree shadow of the bare upstairs charge; the composite $\overline{\Pi}_X \circ s \equiv 0$ records the split-section convention that absorbs the entire Gram-degree into the central coordinate T . The choice of section is the choice of normal ordering: bare placements collect cross-pairing $B(c_i, c_j)$ in the central coordinate (Lemma ??), while s -placements record $\overline{\Pi}_X = 0$ on every component and the Gram-degree information lives entirely in the upstairs $\Pi_X(c)$. The two presentations agree on the homomorphism $\overline{\Pi}_X$. This is the additive group law on which a hypothetical compact Hall product can descend without a chain-level associator at the lattice level.

Remark 3.17 (Chart data, not Hall support). The lattice $\widehat{\Gamma}_X$ is the formal additive group on which charges and their Gram-degree shadows are tracked. It is not the algebraic effective semigroup Γ_X^{eff} of curve classes representable by closed subschemes $Y \subset X$; membership in the effective sub-semigroup is a separate algebraic-effective condition (Definition ??). The Beilinson chart layer α (charge data) and γ (Mukai lattice) license $\widehat{\Gamma}_X$ as ambient; the Hall-support restriction is part of the categorical-input layer at higher Beilinson level, not the chart data of this appendix.

3.6 Three Igusa names; chart-data licensing

Three Igusa names live distinctly in the body and remain distinct under the chart-data of this appendix:

- the automorphic section $\mathcal{D}_X = \Delta_5$, a section of the weight-5 automorphic line over the genus-2 Siegel space with Maass character ν_{Δ_5} ;
- the BKM denominator $\text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1} \Delta_5(2Z)$ on the root lattice $\Lambda_{II}^{2,1}$; and
- the compact protected Pfaffian $\text{Pf}_{\text{prot}}(\mathfrak{D}_X)$, a section of $\mathcal{L}^5 \otimes \nu_{\Delta_5}$ on the conditional pro-object $\mathfrak{D}_X^{\text{DI}}$ of §5.

The Mukai–Gram dictionary supplies the Gram-degree shadow $\Pi_X : \Gamma_X^{\text{form}} \rightarrow \Gamma_{\text{gram}}$ on which the cusp-Fourier expansion of Δ_β , the root-degree map of $\text{den}(\mathfrak{g}_{\Delta_\beta})$, and the wall data of Pf_{prot} all read. The Mukai pairing on $\tilde{H}(S, \mathbb{Z})$ is symmetric in both algebraic and transcendental parts; the ungraded Mukai-form classical limit of an associated Yangian-type construction is the orthogonal Lie algebra $\mathfrak{so}(4, 20)$, not the orthosymplectic $\mathfrak{osp}(4 | 20)$. The Hodge $\mathbb{Z}/2$ -super-extension that appears in cross-volume Yangian conventions imposes a separate $\mathbb{Z}/2$ -grading on the algebraic side; it does not enter the lattice pairing of this appendix. The Bryan/Oberdieck–Pandharipande/ Maulik–Toda conventions [?OPand, ?BRYAN, ?MaulikTodaGV] take β algebraic and d elliptic-degree, agreeing with Theorem 3.9.

The constructions here do not touch the orientation character ϵ_o of Open Problem ??, the Pfaffian section $\text{Pf}_{\text{prot}}(\mathfrak{D}_X)$, or the recognition theorem of Theorem ??; those structures live above this appendix and consume the chart data supplied here.

4 HALL CORRESPONDENCES: THE LEVEL-2 FUNCTORIAL PASSAGE ON $\mathfrak{M}_{\text{eff}}^+(K_3 \times E)$

This appendix records the imported theorems on which the Dirac–Igusa construction of §?? rests on the Hall side, and pins down the $K_3 \times E$ specialisation. The Hall correspondences are level-2 / β arrows in the universal stage chain $\text{CY}_d\text{-cat} \rightarrow E_d\text{-HolFA}(X) \rightarrow \text{Sp}_{\Sigma_{d-1}, C}^{\text{ch}} \rightarrow Y^+ \rightarrow G \rightarrow \kappa\text{-tuple}$ in the sense of Beilinson–Drinfeld [?BD]; equivalently, they comprise the chart-augmented A_b data for the open factorisation dg-category (X, D, τ) at a chosen boundary vacuum b , with chiral shadow on the elliptic carrier supplied separately by §??.

The bibliography supplies the proofs. This appendix supplies the exact statements as used downstream, the proof sketches at the level needed to support the $K_3 \times E$ specialisation, and the identification of which licensing band each clause sits in: α (object / construction), β (comparison / functor / natural transformation), γ (theorem of structure), δ (open obligation paired with named obstruction), ϵ (scalar shadow). The tags appear inline in square brackets after each statement.

The standing data are: S is a smooth complex projective K_3 surface, E is a smooth elliptic curve, $X = S \times E$ is the Calabi–Yau threefold, $\sigma_S \in H^0(S, K_S)$ is the holomorphic two-form trivialising K_S , and $\sigma = \text{pr}_S^* \sigma_S \in H^0(X, p_S^* K_S)$ is the inflated cosection of §??. The Mukai lattice is

$$\Lambda_{\text{Muk}}(K_3) = H^0(S, \mathbb{Z}) \oplus \text{NS}(S) \oplus H^4(S, \mathbb{Z}),$$

endowed with the Mukai pairing $\langle \cdot, \cdot \rangle_{\text{Muk}}$ of [?Mukai1984]; Γ_{eff} is the dyonic charge sublattice of Definition ??; and $\mathfrak{M}_{\text{eff}}^+(X)$ is the moduli stack of (Bridgeland) semistable perfect complexes on X of effective dyonic class.

D.1 The Hall product on the moduli of stable sheaves

The first imported theorem is the existence and associativity of a Hall product on the equivariant cohomology of the moduli stack of semistable sheaves, with vanishing-cycle coefficients, after Joyce and Song.

THEOREM 4.1 (*Joyce–Song generalised DT Hall product; γ , proved on CY_3 projective; conditional on (OI) , $(OI)^+$ on $K_3 \times E$). Let Y be a smooth complex projective Calabi–Yau threefold. The moduli stack $\mathfrak{M}(Y)$ of compactly supported coherent sheaves on Y carries a critical Hall product*

$$m : H_c^\bullet(\mathfrak{M}(Y), \Phi_W \otimes o) \otimes H_c^\bullet(\mathfrak{M}(Y), \Phi_W \otimes o) \longrightarrow H_c^\bullet(\mathfrak{M}(Y), \Phi_W \otimes o),$$

with Φ_W the perverse sheaf of vanishing cycles of the holomorphic Chern–Simons potential and o the orientation line supplied by Joyce–Upmeier orientation data. The product is defined by pull-push along the universal extension stack

$$\mathfrak{M}(Y) \xleftarrow{p} \mathfrak{Ext}^1(\mathfrak{M}(Y), \mathfrak{M}(Y)) \xrightarrow{q} \mathfrak{M}(Y),$$

where $\mathfrak{Ext}^1$ is the moduli stack of short exact sequences $0 \rightarrow A \rightarrow C \rightarrow B \rightarrow 0$, $p(0 \rightarrow A \rightarrow C \rightarrow B \rightarrow 0) = (A, B)$, and $q(0 \rightarrow A \rightarrow C \rightarrow B \rightarrow 0) = C$. The product $m = q_! \circ \text{TS} \circ p^*$ is associative, where TS is the Massey–Saito–Joyce Thom–Sebastiani isomorphism on the universal extension stack [?JoyceSong, Theorem 5.4], [?MasseyST], [?BBDJS2015]. Its unit is the class of the empty object.

Proof sketch. Three inputs assemble the product. First, the moduli stack $\mathfrak{M}(Y)$ is locally the critical locus of the Joyce–Song holomorphic Chern–Simons potential $W : \text{Crit}(W) \hookrightarrow Y_W$ where Y_W is a Darboux chart of the (-1) -shifted symplectic structure on $\mathfrak{M}(Y)$ supplied by Pantev–Toën–Vaquié–Vezzosi [?PTVV2013, Theorem 0.3]; the perverse sheaf Φ_W of vanishing cycles is well defined on the oriented d-critical locus by Brav–Bussi–Dupont–Joyce–Szendrői [?BBDJS2015, Theorem 6.9]. Second, the universal extension stack is locally a critical locus of $W_A \boxplus W_B$ on $Y_A \times Y_B$, and the Massey–Saito Thom–Sebastiani isomorphism [?MasseyST, Theorem 1.1], [?BBDJS2015, Theorem 5.10] identifies $\Phi_{W_A \boxplus W_B}$ with $\Phi_{W_A} \boxtimes \Phi_{W_B}$ on $\text{Crit}(W_A) \times \text{Crit}(W_B)$. Third, the multiplicative orientation cocycle of Joyce–Upmeier [?JoyceUpmeier, Theorem 4.1], [?JoyceTanakaUpmeier, Theorem 1.11] trivialises the orientation gerbe on extensions, supplying the line bundle factor. Pulling $\Phi_W \otimes o$ along p , tensoring by Thom–Sebastiani, and pushing along $q_!$ defines m . Associativity reduces, via the two-step flag stack $\mathfrak{Flag}(\mathfrak{M}(Y), \mathfrak{M}(Y), \mathfrak{M}(Y))$, to the pentagon coherence of Thom–Sebastiani triple stratification [?BBDJS2015, Theorem 5.10(iv)] together with the Joyce–Upmeier multiplicative cocycle on triple extensions [?JoyceUpmeier, §4]. See Joyce–Song [?JoyceSong, §5.2, Theorem 5.5] for the original construction and Davison [?Davison2017, Theorem A] for the cohomological amplification. $\square$

$K_3 \times E$ specialisation. On $Y = K_3 \times E$, the cosection $\sigma = \text{pr}_S^* \sigma_S$ of the holomorphic two-form on the K_3 factor reduces the obstruction theory through Kiem–Li (Appendix 4); the supplied moduli is $\mathfrak{M}_{\text{eff}}^{\text{red}}(X) \subset \mathfrak{M}(X)$, and the product is the reduced critical product

$$m^{\text{red}} = q_!^{\text{red}} \circ \text{TS}^{\text{red}} \circ p^*$$

of Proposition 0.9. Vanishing of the seven obstruction classes $\mathfrak{o}_R^{\text{conf}}$, $\mathfrak{o}_R^{\text{prop,LL}}$, $\mathfrak{o}_R^{\text{prop,mix}}$, $\mathfrak{o}_R^{\text{prop,WW}}$, $\mathfrak{o}_R^{\text{conf-prop}}$, $\mathfrak{o}_{w,R}^{\text{assoc}}$, and $\mathfrak{o}_{w,R}^{I,\text{prod}}$ is the data $(D_0) + (O_1)$, absorbed into Bnd, dCrit, RedOr, and Six of Definition ???. *The existence of a chain-level Hall product on $K_3 \times E$ without those four data is open.* The downstream invariant $Y^+(X) := H_c^\bullet(\mathfrak{M}_{\text{eff}}^+(X), \Phi_W \otimes o)$ is the positive-half cohomological Hall algebra (CoHA) of Kontsevich–Soibelman [?KSCoHA] restricted to effective dyonic class; its Drinfeld double, regulated tensor product completion, and integral form constitute the level-3 object $G(X)$ [β -band, conditional]. The level-3 object thus arises from $Y^+(X)$ by Hall pairing, integral-form completion, and stable-envelope transport, three arrows downstream of the level-2 Hall positive half (Vol III [?KSCoHA] cache row).

D.2 The Hall coproduct via collisions

The Hall product has a dual coproduct, defined by reversing pull and push in the universal short-exact-sequence stack. The collision coproduct furnishes $Y^+(X)$ with a (graded) bialgebra structure;

after Drinfeld doubling (Hall pairing + completion + integral form + stable-envelope transport, § D.6) the bialgebra is the level-3 quantum group $G(X) = D_h(Y^+(X))$ of Schiffmann–Vasserot type [?SchiffmannVasserot2013].

THEOREM 4.2 (*Joyce–Song collision coproduct*; [?JoyceSong] *Theorem 5.5*; β , *proved on CY₃ projective*). Let Y be a smooth projective Calabi–Yau threefold. The critical Hall positive half $Y^+(Y) = H_c^\bullet(\mathfrak{M}_{\text{eff}}^+(Y), \Phi_W \otimes o)$ carries a collision coproduct

$$\Delta : Y^+(Y) \longrightarrow Y^+(Y) \otimes Y^+(Y), \quad \Delta = p_!^{\text{coll}} \circ \text{TS}^{-1} \circ q^{\text{coll},*},$$

defined by the collision/extension correspondence

$$\mathfrak{M}_{\text{eff}}^+(Y) \xleftarrow{q^{\text{coll}}} \mathfrak{Ext}^{1,\text{coll}}(\mathfrak{M}_{\text{eff}}^+(Y)) \xrightarrow{p^{\text{coll}}} \mathfrak{M}_{\text{eff}}^+(Y) \times \mathfrak{M}_{\text{eff}}^+(Y),$$

with $\mathfrak{Ext}^{1,\text{coll}}$ the moduli of short exact sequences $0 \rightarrow A \rightarrow C \rightarrow B \rightarrow 0$ read with C on the left and (A, B) on the right, $q^{\text{coll}}(0 \rightarrow A \rightarrow C \rightarrow B \rightarrow 0) = C$, $p^{\text{coll}}(0 \rightarrow A \rightarrow C \rightarrow B \rightarrow 0) = (A, B)$. The coproduct Δ is graded coassociative. Together with the Hall product m of Theorem 4.1, $(Y^+(Y), m, \Delta)$ is a graded bialgebra.

Proof sketch. Coassociativity is the dual statement of associativity proven through the $\mathfrak{Flag}$ stack and the inverse Thom–Sebastiani pentagon [?BBDJS2015, Theorem 5.10(iv)]. The compatibility of m and Δ follows from the Yetter–Drinfeld–type cyclic two-step diagram of Joyce–Song [?JoyceSong, §5.4]; that diagram fails the strict bialgebra Hopf axiom on the unmodified Hall pairing and is repaired by the Hall pairing twist $\langle \cdot, \cdot \rangle_{\text{Hall}}$ supplied by the Hopf radical kernel of Lemma ?? . See Davison–Meinhardt [?DavisonMeinhardtCoHA, Theorem A] for the amplified bialgebra statement on quivers with potential, and Davison [?Davison2017, Theorem A] for the integrality of Y^+ on preprojective stacks. Coassociativity transports verbatim through the cosection-reduced quotient $\mathfrak{M}_{\text{eff}}^+(X) \rightarrow \mathfrak{M}_{\text{eff}}^{\text{red},+}(X)/E$ once (Do) supplies the equivariant six-functor formalism on the two-step flag stack. $\square$

Drinfeld double. The Hall pairing $\langle \cdot, \cdot \rangle_{\text{Hall}}$ of Lemma ?? pairs Y^+ with itself; the Drinfeld double $D_h(Y^+(X))$ is the quantum double in the sense of Drinfeld [?KAC: 1, §13], supplying the negative half $Y^-(X)$ and the Cartan generators K_γ , $\gamma \in \Gamma_{\text{eff}}$, with relations encoding the Hall pairing. The *level-3* $K_3 \times E$ *quantum group* is then

$$G(X) := D_h(Y^+(X))_{\text{compl}}^{\text{int}} \quad [\beta, K_3 \times E \text{ compact: open}]$$

where $(-)_{\text{compl}}^{\text{int}}$ is the integral-form completion under the Hall pairing, conditional on the stable-envelope transport of § D.6. In Schiffmann–Vasserot [?SchiffmannVasserot2013] the analogous identification on toric CY₃ ambient is established; $K_3 \times E$ admits no global non-commutative crepant resolution (NCCR) for five obstructions ((a) trivial ω but ω -structure not reflexive-tilting; (b) derived McKay needs finite Aut fixing a point; (c) HPD self-dual fails product polarisation; (d) Mukai vanishing fails off the K₃ factor; (e) no global CY₃ symmetric obstruction theory). The Serre-equivariant quasi-NCCR substitute supplied by the cosection-twist of § D.4 is the standing replacement; the global $G(X)$ on the compact non-toric $K_3 \times E$ accordingly remains an open obligation.

D.3 The Hall primitives and the formal recognition criterion

For a connected graded bialgebra $B = \bigoplus_c B_c$, the primitive elements at charge c are

$$\mathrm{Prim}_c(B) = \{x \in B_c \mid \Delta(x) = x \otimes 1 + 1 \otimes x\},$$

equivalently the kernel of the reduced coproduct $\bar{\Delta}$. On a graded connected cocommutative Hopf algebra in characteristic zero, the Milnor–Moore theorem [?LurieHA, Theorem 5.18] identifies $B \simeq U(\mathrm{Prim}_\bullet(B))$ as Hopf algebras, where $\mathrm{Prim}_\bullet(B)$ is the Lie algebra of primitives under the bracket $[x, y] = xy - (-1)^{|x||y|}yx$.

THEOREM 4.3 (*Hall primitives, formal level; β , formal*). Let $\mathfrak{M}$ carry a critical Hall positive half $Y^+(Y) = (B, m, \Delta)$ of Theorem 4.2. The graded subspace

$$\mathrm{Prim}_\bullet(Y^+(Y)) = \bigoplus_{c \in \Gamma_{\mathrm{eff}}} \mathrm{Prim}_c(Y^+(Y))$$

is a graded Lie subalgebra of $Y^+(Y)$ under the antisymmetrised product $[\cdot, \cdot]$. At a charge c such that the moduli $\mathfrak{M}_c^+(Y)$ is connected and the C -component of Δ on $\mathfrak{M}_c^+(Y)$ factors through extensions whose graded pieces lie in strictly lower charges, the inclusion $\mathrm{Prim}_c \hookrightarrow Y_c^+$ is the kernel of $\bar{\Delta}_c$. When $Y^+(Y)$ is graded connected and cocommutative, Milnor–Moore identifies $Y^+(Y) \simeq U(\mathrm{Prim}_\bullet(Y^+(Y)))$ as graded Hopf algebras.

Proof sketch. Primitivity of $\mathrm{Prim}_\bullet$ under the antisymmetrised product is verified by direct computation of $\Delta([x, y])$ using the cocommutative-up-to-twist axiom and the multiplicativity of Δ ; the resulting expression collapses to $[x, y] \otimes 1 + 1 \otimes [x, y]$. On a connected graded cocommutative Hopf algebra in characteristic zero, the Milnor–Moore identification is [?LurieHA, §21.1]; on graded connected non-cocommutative Hopf algebras, the Cartier–Quillen–Milnor–Moore theorem [?LurieHA, Theorem 21.2.1] requires either cocommutativity or a chain-level lift through bar / cobar, which is exactly the chiral Koszul firewall of Vol I [?FrancisGaitsgoryCKD]. $\square$

Recognition. The recognition theorem of Theorem 0.29 of the body of this manuscript supplies the upgrade of the formal primitive Lie superalgebra to the imported Borchers–Kac–Moody Lie superalgebra $\mathfrak{g}_\Delta$ of Theorem ??, conditional on (Do), (O₁), (O₁)⁺, (O₂), the completed PBW filtration, the Hopf pairing radical equality, and the parity dimension recovery. In particular, on $K_3 \times E$ the formal primitive subspace $\mathrm{Prim}_\bullet(Y^+(X))$ is the *candidate* primitive Lie superalgebra; its identification with the imported BKM denominator algebra $\mathfrak{g}_\Delta$ is the *compact Pfaffian View* ③ of the Pentadic decomposition, conditional on the four named obstructions. This appendix supplies no recognition theorem; it supplies the formal Hopf algebra structure on which recognition acts.

D.4 The cosection-twist on $K_3 \times E$: Kiem–Li localisation

On $K_3 \times E$, the holomorphic two-form on the K_3 factor pulls back to a non-vanishing cosection of the obstruction theory of $\mathfrak{M}(X)$; the Kiem–Li [?KiemLi2013] localisation theorem then localises the virtual fundamental class to the zero-locus of the cosection. The localisation is the source of the *reduced* virtual class on which the Hall product of § D.1 carries chain-level coherence.

Definition 4.4 (*Cosection*). Let $\mathfrak{M}$ be a Deligne–Mumford stack equipped with a perfect obstruction theory $\phi : \mathbb{E} \rightarrow \mathbb{L}_{\mathfrak{M}}$ of amplitude $[-1, 0]$, and let $\mathrm{Ob}(\mathfrak{M}) = h^1(\mathbb{E}^\vee)$ be the coherent obstruction sheaf. A *cosection* of the obstruction theory is an $\mathcal{O}_{\mathfrak{M}}$ -linear morphism

$$\sigma : \mathrm{Ob}(\mathfrak{M}) \longrightarrow \mathcal{O}_{\mathfrak{M}};$$

its *vanishing locus* is the closed substack $\mathfrak{M}^{\text{red}} = (\sigma = 0)$ on which the obstruction sheaf restricts to the kernel $\ker \sigma \subset \text{Ob}(\mathfrak{M})$.

THEOREM 4.5 (*Kiem–Li cosection localisation*; [?KiemLi2013] *Theorem 5.0.1*; γ , *proved*). Let $\mathfrak{M}$ be a Deligne–Mumford stack with perfect obstruction theory ϕ and surjective cosection $\sigma : \text{Ob}(\mathfrak{M}) \rightarrow \mathcal{O}_{\mathfrak{M}}$ on $\mathfrak{M} \setminus \mathfrak{M}^{\text{red}}$, where $\mathfrak{M}^{\text{red}} = (\sigma = 0)$. The virtual fundamental class $[\mathfrak{M}]^{\text{vir}} \in A_{\text{vd}}(\mathfrak{M})$ lifts to a cosection-localised class

$$[\mathfrak{M}]_{\text{loc}}^{\text{vir}} \in A_{\text{vd}+1}(\mathfrak{M}^{\text{red}})$$

satisfying $\iota_*[\mathfrak{M}]_{\text{loc}}^{\text{vir}} = [\mathfrak{M}]^{\text{vir}}$, where $\iota : \mathfrak{M}^{\text{red}} \hookrightarrow \mathfrak{M}$ is the inclusion. The reduced virtual dimension is $\text{vd}^{\text{red}} = \text{vd} + 1$.

Proof sketch. The cosection-kernel cone $C(\sigma) = \ker(\sigma : \text{Ob} \rightarrow \mathcal{O})$ is a perfect obstruction cone of amplitude $[-1, 0]$ on $\mathfrak{M}^{\text{red}}$ with $\text{rk } C(\sigma) = \text{rk } \text{Ob} - 1$. The intrinsic normal cone $\mathfrak{C}$ of $\mathfrak{M}$ sits in the total obstruction bundle on $\mathfrak{M} \setminus \mathfrak{M}^{\text{red}}$, where σ is surjective and the cone has codimension one in Ob ; on $\mathfrak{M}^{\text{red}}$, the cone embeds into $C(\sigma)$ by [?KiemLi2013, Proposition 4.3]. Intersection with the zero section of $C(\sigma)$ on $\mathfrak{M}^{\text{red}}$, pushed forward to $\mathfrak{M}$ through ι , yields $[\mathfrak{M}]^{\text{vir}}$ by [?KiemLi2013, Theorem 5.0.1]. The dimension shift $\text{vd}^{\text{red}} = \text{vd} + 1$ is the codimension of $\ker \sigma$ in Ob . $\square$

$K_3 \times E$ cosection. On $X = K_3 \times E$, the cosection of the obstruction theory of $\mathfrak{M}(X)$ is induced by Mukai dilation along $\sigma_S = \text{pr}_S^* \sigma_S$. Concretely, for a perfect complex F on X , the chain-level cosection acts on $\text{Ext}^2(F, F)$ by the semiregularity map of Buchweitz–Flenner / Bloch [?MaulikTodaGV],

$$\text{cs}_F : \text{Ext}^2(F, F) \longrightarrow \mathbb{C}, \quad \text{cs}_F(\xi) = \int_X \sigma \wedge \text{Tr}((\xi \otimes \text{I}_{\Omega_X^1}) \circ \text{At}(F)),$$

where $\text{At}(F) \in \text{Ext}^1(F, F \otimes \Omega_X^1)$ is the Atiyah class. The cosection vanishes precisely on the *reduced* moduli $\mathfrak{M}^{\text{red}}(X) \subset \mathfrak{M}(X)$ consisting of sheaves whose Atiyah class is annihilated by σ ; equivalently, sheaves whose deformation space along the K_3 direction is constrained. See Maulik–Toda [?MaulikTodaGV, Lemma 2.6, Theorem 2.7] and Oberdieck [?OberdieckReducedPT, Theorem 1, Theorem 3] for the explicit identification on the DT/PT side. The cosection is surjective off the cusp locus of the Mukai vector $\text{ch}(F) = (0, 0, c_2(F), \dots)$ for which the K_3 component is trivial; on the $s = 0$ cusp $\sigma_S = 0$ and the localisation degenerates, which is the *Do*-degeneration limit named at obstruction (*Do*) of § ??.

THEOREM 4.6 (*Reduced virtual class on $K_3 \times E$; proved on the smooth primitive nonzero K_3 -class branch*). Let $\mathfrak{M}_c^+(X)$ be the moduli of semistable sheaves of dyonic class $c \in \Gamma_{\text{eff}}$ with nonzero K_3 component. The cosection-localised virtual class $[\mathfrak{M}_c^+(X)]_{\text{loc}}^{\text{vir}} \in A_{\text{vd}^{\text{red}}}(\mathfrak{M}_c^{\text{red},+}(X))$ is the reduced virtual class of Maulik–Toda [?MaulikTodaGV, §2]; its DT/PT shadow recovers the Oberdieck–Pixton scalar $Z_{\text{OP}}^X = -4096 \Delta_3^{-2}$ on the smooth primitive K_3 -class branch [ε , proved] [?OberdieckReducedPT, Theorem 3]. On the $s = 0$ cusp the cosection degenerates; *the cosection-reduced theory there is the (Do) degeneration limit named at § ?? and is open.* [δ , (*Do*)]

D.5 BBDJS reduced d -critical orientation

The Hall product of § D.1 requires not just a virtual fundamental class but a perverse-sheaf-of-vanishing-cycles refinement and an orientation gerbe; the source is the Brav–Bussi–Dupont–Joyce–Szendrői [?BBDJS2015] extension of the Joyce [?JoyceSong] d -critical-locus framework.

THEOREM 4.7 (*BBDJS perverse-sheaf-of-vanishing-cycles; oriented CY_3 version of [?BBDJS2015] Theorem 6.9; γ , proved*). Let $\mathfrak{M}$ be a derived Artin stack with a (-1) -shifted symplectic structure (PTVV [?PTVV2013]) whose classical truncation is a d-critical locus in the sense of Joyce [?JoyceSong, Definition 2.5]. Suppose $\mathfrak{M}$ carries an *orientation*, i.e. a square root o of the Joyce determinant line $K_{\mathfrak{M}}^{\text{vir}} = (\det \mathbb{L}_{\mathfrak{M}})^{1/2}$, in the sense of [?BBDJS2015, Theorem 6.9]. There is a perverse sheaf

$$\Phi_{\mathfrak{M}} \in \text{Perv}(\mathfrak{M}),$$

the *perverse sheaf of vanishing cycles* of the d-critical locus; locally on a Joyce–Song Darboux chart $\text{Crit}(W)$ inside a smooth ambient Y_W , $\Phi_{\mathfrak{M}}|_{\text{Crit}(W)} \simeq \Phi_W[d_{Y_W}]$ is the standard perverse sheaf of vanishing cycles of the holomorphic potential $W : Y_W \rightarrow \mathbb{C}$, shifted by the ambient dimension. The chart-comparison cocycle is canonical up to the orientation choice; the orientation upgrades to a multiplicative cocycle of Joyce–Upmeyer [?JoyceUpmeyer, Theorem 4.1] on the moduli of objects.

Proof sketch. Three layers stack. PTVV [?PTVV2013, Theorem 0.3] produces the (-1) -shifted symplectic form on the derived moduli of CY_3 coherent sheaves; Bussi–Brav–Joyce [?BBDJS2015, §5] produce the d-critical-locus chart structure on the classical truncation; Bussi–Brav–Dupont–Joyce–Szendrői [?BBDJS2015, Theorem 6.9] produce the perverse-sheaf-of-vanishing-cycles package with chart-cocycle data. The orientation is the data needed to unify chart-local Φ_W into a global perverse sheaf $\Phi_{\mathfrak{M}}$; without it, the chart-cocycle has a sign ambiguity that obstructs the descent to a perverse sheaf [?BBDJS2015, §6.4]. Joyce–Upmeyer [?JoyceUpmeyer, Theorem 1.11] build the multiplicative orientation cocycle on the universal extension stack; this is the CY_3 projective input that is absorbed below. $\square$

Reduced refinement on $K_3 \times E$. On $X = K_3 \times E$, the (-1) -shifted symplectic structure $\omega_{\mathfrak{M}}$ on $\mathfrak{M}(X)$ does not restrict to a (-1) -shifted symplectic structure on $\mathfrak{M}^{\text{red}}(X)$; the cosection-twist of § D.4 contracts $\omega_{\mathfrak{M}}$ by σ , and the *reduced* (-1) -shifted form $\omega_{\mathfrak{M}}^{\text{red}}$ is non-degenerate only after restriction to the cosection vanishing locus [?MaulikTodaGV, §2.6]. The corresponding reduced d-critical-locus structure is supplied by the Maulik–Toda [?MaulikTodaGV, Theorem 2.7] extension; the reduced perverse sheaf of vanishing cycles $\Phi_{\mathfrak{M}}^{\text{red}}$ and the reduced orientation line o_C^{red} are exactly the data (O_1) of § ?? . In the language of (P4) of the body manuscript, the reduced orientation is twisted by a parity-flip line Π generated by the cosection cokernel,

$$o_C^{\text{red}} = o_C^{\text{std}} \otimes \Pi,$$

absorbing the dimension shift $\text{vd}^{\text{red}} = \text{vd}^{\text{std}} + 1$ into a $\mathbb{Z}/2$ -grading shift on the oriented BBDJS line.

PROPOSITION 4.8 (*BBDJS reduced d-critical orientation on $K_3 \times E$; δ , (O_1)*). Let $\mathfrak{M}_c^{\text{red},+}(X)$ be the cosection-reduced moduli of semistable sheaves of effective dyonic class c on $X = K_3 \times E$. The hypotheses *strong reduced orientation on the $K_3 \times E$ -quotient self-Ext frame bundle (O_1) and Weyl-equivariant transport along $W^{(2)}(\Lambda_{II}^{2,1})$ reflections $(O_1)^+$* of § ?? are exactly the following. There exist a multiplicative orientation cocycle of Joyce–Upmeyer type extending equivariantly across the reduced E -quotient, the σ_S -cosection-induced parity flip line Π , and a connected BE -equivariant trivialisation of the small-orbit gerbe class $\overline{\beta}_C^E$ on the finite-stabilizer strata, such that

$$\Phi_{\mathfrak{M}_c^{\text{red},+}(X)}^{\text{red}} \otimes o_c^{\text{red}}$$

descends through the free E -translation quotient and trivialises the Pin^c -lift gerbe class of the reduced derived self-Ext frame bundle. The strong reduced orientation (O_1) is established on the unreduced

$K_3 \times E$ -quotient by Joyce–Upmeyer [?JoyceUpmeyer, Theorem 1.11] on the CY_3 projective ambient, on the cosection-reduced quasi-projective extension by Bojko [?Bojko, Theorem 1.4], and on the cosection-twisted orientation package on toric CY_4 ambient via Cao–Kool–Monavari [?CaoKoolMonavari, Theorem 1.6]; the equivariant transport across the free E -quotient and across the finite-stabilizer strata, jointly with Weyl-equivariant transport, is the open obligation $(O_1) + (O_1)^+$ of § 5.8.

Proof sketch. The construction of the reduced perverse sheaf $\Phi_{\mathfrak{M}_c^{\text{red},+}(X)}^{\text{red}}$ on the cosection-reduced moduli is by Maulik–Toda [?MaulikTodaGV, Theorem 2.7]; the orientation gerbe lifts to the reduced Joyce d-critical locus by the BBDJS [?BBDJS2015, §6.4] chart-cocycle calculation. The multiplicative *equivariant* orientation cocycle across the E -quotient and across the finite-stabilizer locus $BE[N]$ -strata is the joint vanishing of (O_1) and $(O_1)^+$: on the connected BE -equivariant trivialisation of $\alpha_C^E \in H_E^2(\mathfrak{M}_C^{\text{red}}, \mathbb{F}_2)$ of Lemma ?? it suffices to trivialise the small-orbit gerbe class $\bar{\beta}_C^E$ on $BE[N]$ -strata, governed by the Klein-four detection criterion of Lemma ?. The Weyl-equivariant transport across $W^{(2)}(\Lambda_{II}^{2,1})$ reflections is the orientation character calculation of Theorem ?? on the three type-II walls. Both transports are open in general; the Pin^c -lift programme of [?JoyceUpmeyer], [?JoyceTanakaUpmeyer], [?CaoKoolMonavari], [?Bojko] supplies the analytic technology. $\square$

D.6 Stable-envelope transport: from Hall to Yangian

The level-3 quantum-group structure on $G(X) = D_h(Y^+(X))$ of § D.2 is, on toric ambient or quiver varieties, identified by Maulik–Okounkov [?MaulikOkounkov] with the geometric R -matrix of the *stable envelope*, providing the link from the Hall side to the Yangian / quantum-group side.

Definition 4.9 (Stable envelope). Let $\mathfrak{M}$ be a smooth symplectic variety with a torus T -action and a finite T -fixed locus $\mathfrak{M}^T$; let $C \subset \text{Lie}(T)$ be a chamber of the T -equivariant cohomology of $\mathfrak{M}$. The *stable envelope* [?MaulikOkounkov, §3] is the T -equivariant cohomological correspondence

$$\text{Stab}_C : H_T^\bullet(\mathfrak{M}^T) \longrightarrow H_T^\bullet(\mathfrak{M}),$$

characterised by support, normalisation, and degree axioms [?MaulikOkounkov, Theorem 3.3.4]. The geometric R -matrix is the comparison

$$R_{C,C'}^{\text{MO}} = \text{Stab}_{C'}^{-1} \circ \text{Stab}_C \in \text{End}(H_T^\bullet(\mathfrak{M}^T))$$

across two chambers C, C' ; it satisfies the Yang–Baxter equation by [?MaulikOkounkov, Theorem 4.6.1].

THEOREM 4.10 (Maulik–Okounkov; [?MaulikOkounkov] Theorem 10.2.1; β , proved on quiver varieties; conditional on $K_3 \times E$ compact non-toric). Let $\mathfrak{M} = \mathfrak{M}_{Q,\mathbf{v},\mathbf{w}}$ be a Nakajima quiver variety [?Nakajima1994, ?Nakajima1998] attached to a finite quiver Q with dimension vectors $\mathbf{v}, \mathbf{w}$. The stable envelope Stab_C makes $H_T^\bullet(\mathfrak{M}_{Q,\mathbf{v},\mathbf{w}})$ a module over a Yangian $Y_h(\mathfrak{g}_Q)$ attached to Q , with $\mathfrak{g}_Q$ the Kac–Moody Lie algebra of the quiver, and the geometric R -matrix R^{MO} is the rational R -matrix of $Y_h(\mathfrak{g}_Q)$.

Proof sketch. The Maulik–Okounkov programme proceeds through: (i) the existence and uniqueness of the stable envelope (axioms of support, polarisation, normalisation), [?MaulikOkounkov, §3.3]; (ii) the Yang–Baxter equation as a chamber-wall cocycle, [?MaulikOkounkov, §4]; (iii) the identification of the Bethe algebra generated by the matrix elements of Stab with a Yangian, [?MaulikOkounkov, §5–10].

The Schiffmann–Vasserot identification $\text{CoHA}(\mathbb{C}^3) = Y^+(\widehat{\mathfrak{gl}}_1)$ of [?SchiffmannVasserot2013, Theorem A] is the affine $\mathcal{A}_0$ specialisation; $\mathcal{W}_{1+\infty}$ arises *only after* Drinfeld doubling and Fock-evaluation. The three-arrow chain $\text{CoHA} = Y^+ \hookrightarrow Y \xrightarrow{\text{ev}_1} \text{End}(\mathcal{W}_{1+\infty}[\lambda]\text{-vac})$ of [?KSCoHA], [?SchiffmannVasserot2013] traverses the level-2 $\rightarrow$ level-3 $\rightarrow$ level-3-evaluated passage; the named identity on each arrow records the structural distance from $\text{CoHA}(\mathbb{C}^3)$ to $\mathcal{W}_{1+\infty}$. $\square$

K₃ $\times$ E specialisation: open. On $X = K_3 \times E$, the moduli $\mathfrak{M}_{\text{eff}}^+(X)$ is not a Nakajima quiver variety, the ambient is compact non-toric, and the global stable envelope is unconstructed. The five obstructions to a global non-commutative crepant resolution (NCCR) of X recorded in [?SchiffmannVasserot2013], [?KSCoHA], ((a) trivial ω but ω -structure not reflexive-tilting; (b) derived McKay needs finite Aut fixing a point; (c) HPD self-dual fails product polarisation; (d) Mukai vanishing fails off the K_3 factor; (e) no global CY_3 symmetric obstruction theory) obstruct the literal [?MaulikOkounkov] construction. The cosection-twist of § D.4 supplies the *Serre-equivariant quasi-NCCR substitute*: the cosection-reduced obstruction theory of Theorem 4.6 restores a CY_3 -symmetric reduced obstruction package on the $s \neq 0$ sector, and the Hall-to-Yangian comparison is then expected at the level of the reduced theory and conditional on the discharge of the four obstructions (Do) , (O_1) , $(\text{O}_1)^+$, (O_2) of § ??.

The Hall–Yangian comparison on $K_3 \times E$ is therefore a *conjectural* β -arrow:

$$G(X) \stackrel{?}{=} D_h(\text{CoHA}_{K_3 \times E}^{\text{red},+})_{\text{compl}}^{\text{int}} \stackrel{?}{=} Y_h(\widehat{\mathfrak{g}}_{\Delta_s}) \quad [\partial, \text{open}]$$

where $\widehat{\mathfrak{g}}_{\Delta_s}$ is the central extension of the imported BKM Lie superalgebra of Theorem ??, and the equality is conditional on a $K_3 \times E$ stable-envelope transport. The Hall–Drinfeld double $\mathcal{D}_h(\text{CoHA}_{K_3 \times E})$ of Kontsevich–Soibelman [?KSCoHA] places this identification at the same level-3 quantum group seen from the geometric side; it is the open compact non-toric instance of the chain-fusion conjecture.

D.7 Banned identifications and cross-volume firewall

The Beilinson cut of [?KSCoHA], [?SchiffmannVasserot2013] and the standing firewall of § ?? pair each banned identification (left column) with the structurally correct identification (right column). Each row records a categorical-level collision the audit corrects.

[Y] *Banned:* $Y^+(X) = G(X)$. *Required:* $Y^+(X)$ is the cohomological-Hall positive half $H_c^\bullet(\mathfrak{M}_{\text{eff}}^+(X), \Phi_W \otimes o)$ with the Joyce–Song product/coproduct of Theorems 4.1, 4.2; $G(X) = D_h(Y^+(X))_{\text{compl}}^{\text{int}}$ is the Drinfeld double after Hall pairing, completion, integral form, stable-envelope transport, and descent.

[CoHA] *Banned:* $\text{CoHA}(\mathbb{C}^3) = \mathcal{W}_{1+\infty}$. *Required:* $\text{CoHA}(\mathbb{C}^3) = Y^+(\widehat{\mathfrak{gl}}_1)$ [?SchiffmannVasserot2013, Theorem A]; $\mathcal{W}_{1+\infty}$ appears as the Drinfeld-double + Fock-evaluation image, three arrows downstream.

[NCCR] *Banned:* $K_3 \times E$ admits a global NCCR. *Required:* $K_3 \times E$ admits no global NCCR; five named obstructions [?SchiffmannVasserot2013], [?KSCoHA]. The cosection-twist of § D.4 supplies the Serre-equivariant quasi-NCCR substitute.

[Hall–BKM] *Banned:* the Hall primitive Lie algebra is the BKM denominator algebra. *Required:* $\text{Prim}_\bullet(Y^+(X))$ is the formal primitive Lie superalgebra of $Y^+(X)$; its identification with $\mathfrak{g}_{\Delta_s}$ is the conditional Pfaffian–Dirac recognition theorem of Theorem 0.29, under (Do) , (O_1) , $(\text{O}_1)^+$, (O_2) .

[d-crit] *Banned:* a Joyce d-critical locus admits a perverse sheaf of vanishing cycles without orientation. *Required:* BBDJS [?BBDJS2015, Theorem 6.9] requires an *oriented* d-critical locus; the orientation

is the square root of the Joyce determinant line, and the multiplicative cocycle is Joyce–Upmeier [?JoyceUpmeier, Theorem 4.1].

[cosection] *Banned*: cosection localisation holds without surjectivity. *Required*: Kiem–Li [?KiemLi2013, Theorem 5.0.1] requires the cosection to be surjective off the localised locus; on $K_3 \times E$, the cosection degenerates on the $s = 0$ cusp, which is the (Do) degeneration limit and is open.

[shift] *Banned*: (-1) -shifted symplectic on $\mathfrak{M}(K_3 \times E)$ restricts to (-1) -shifted symplectic on $\mathfrak{M}^{\text{red}}(K_3 \times E)$. *Required*: the cosection contracts the form; ω^{red} is non-degenerate only after the restriction to the cosection vanishing locus [?MaulikTodaGV, §2.6], and the orientation line carries the $\mathbb{Z}/2$ -parity flip Π absorbing the dimension shift $\text{vd}^{\text{red}} = \text{vd}^{\text{std}} + 1$.

[fourfold] *Banned*: $K_3 \times E$ is a smooth projective fourfold. *Required*: $K_3 \times E$ is a threefold; $n = \chi(\mathcal{O}_X) = \chi(\mathcal{O}_{K_3}) \cdot \chi(\mathcal{O}_E) = 2 \cdot 0 = 0$, and the BBDJS/Joyce–Upmeier machinery applies in the CY_3 form [?BBDJS2015, §6], not in the CY_4 Borisov–Joyce [?BorisovJoyce] or Oh–Thomas [?OhThomasI, ?OhThomasII] form.

D.8 Adversarial loop: closing checks

Four failure modes are checked.

(F1) Hall associativity. The Joyce–Song associativity of Theorem 4.1 requires the pentagon coherence of Thom–Sebastiani triple stratification [?BBDJS2015, Theorem 5.10(iv)]. Failure mode: the chart-cocycle of the d-critical locus carries a sign ambiguity whose obstruction to a canonical pentagon coheres only after orientation choice. Resolution: the orientation gerbe of BBDJS [?BBDJS2015, §6.4] trivialises exactly this ambiguity; the multiplicative cocycle of Joyce–Upmeier [?JoyceUpmeier, §4] extends the trivialisation to the universal extension stack. On $K_3 \times E$, the gerbe descends through the free E -quotient and trivialises the small-orbit class, jointly $(O_1) + (O_1)^+$; this is the open obligation, not a contradiction of the projective associativity.

(F2) Cosection surjectivity. Kiem–Li [?KiemLi2013, Theorem 5.0.1] requires $\sigma : \text{Ob} \rightarrow \mathcal{O}$ on $\mathfrak{M} \setminus \mathfrak{M}^{\text{red}}$. Failure mode: on the $s = 0$ cusp of the dyonic charge lattice of $K_3 \times E$, the cosection $\sigma_S = 0$ on the K_3 factor and surjectivity fails. Resolution: the smooth primitive nonzero K_3 -class branch satisfies surjectivity by Maulik–Toda [?MaulikTodaGV, §2]; the cusp branch is the (Do) degeneration limit named at obstruction (Do) of §??, and is open. The Oberdieck–Pixton scalar $Z_{\text{Op}}^X = -4096 \Delta_5^{-2}$ is proved on the $s \neq 0$ branch [?OberdieckReducedPT, Theorem 3]; on the cusp the chain-level theory degenerates to the Borcherds imaginary-simple-root contribution.

(F3) BBDJS orientation existence on compact non-toric. The Joyce–Upmeier multiplicative cocycle [?JoyceUpmeier, Theorem 4.1] is established on the CY_3 projective ambient; its extension across the cosection-reduced quasi-projective ambient is Bojko [?Bojko, Theorem 1.4]. Failure mode: on $K_3 \times E$ quotient by free E -translation, the orientation line need not descend; the small-orbit gerbe class on $BE[N]$ -strata may not trivialise. Resolution: the joint vanishing of the connected BE -equivariant linearisation obstruction $\alpha_C^E \in H_E^2(\mathfrak{M}_C^{\text{red}}; \mathbb{F}_2)$ and the small-orbit gerbe class $\overline{\beta}_C^E \in H^\bullet(BE[N]; \mathbb{F}_2)$ is exactly (O_1) ; higher-rank Pin^c -vanishing is Lemma ?? of the body. Stable higher-rank length-one strata and nontrivial finite-stabilizer linearisations remain genuine open obligations.

(F4) (-1) -shifted vs (-2) -shifted symplectic. $K_3 \times E$ is a Calabi–Yau threefold; the moduli $\mathfrak{M}(K_3 \times E)$ carries a (-1) -shifted symplectic structure by PTVV [?PTVV2013, Theorem 0.3]. Failure mode: mistaking $K_3 \times E$ for a CY_4 ambient (e.g. under a confusion with $K_3 \times T^2$ at the F-theory level) and applying Borisov–Joyce [?BorisovJoyce] or Oh–Thomas [?OhThomasI, ?OhThomasII]. Resolution: the Calabi–Yau dimension of $X = K_3 \times E$ is $\dim_{\mathbb{C}}(K_3) + \dim_{\mathbb{C}}(E) = 2 + 1 = 3$, the holomorphic

three-form is $\sigma_S \wedge dz_E$, and the shifted symplectic is $[-1]$ -shifted by [?PTVV2013, §2.1]. The cosection of § D.4 reduces vd by one, but does not reduce the shift; ω^{red} is still (-1) -shifted, on the cosection vanishing locus, by the Maulik–Toda [?MaulikTodaGV, Theorem 2.7] extension. The cosection-twist of § D.4 is the Serre-equivariant quasi-NCCR substitute for the missing global NCCR of $K_3 \times E$, as recorded in [?SchiffmannVasserot2013], [?KSCoHA].

The four failure modes above are absorbed into the four obstructions $(D_0), (O_1), (O_1)^+, (O_2)$ of § ??; their joint discharge tightens the conditioning of the cross-level identity $\textcircled{2} \leftrightarrow \textcircled{3}$ of the manuscript thesis. The Hall correspondences of this appendix supply the level-2 / β arrows; the recognition of $\text{Prim}_\bullet(Y^+(X)) \simeq \mathfrak{g}_{\Delta_3}$ of Theorem 0.29 acts on these arrows; the protected Pfaffian recognition theorem $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}}) = \Delta$, of Theorem ?? is the cross-level identity acting on the level-4 scalar shadow.

APPENDIX A

PFSAFFIAN WALL NORMAL FORM ON TYPE-II WALLS

The three local Pfaffian wall computations of this appendix supply the sign data used in Chapter ?? for the orientation-character match

$$\epsilon_o = \nu_{\Delta_5} \quad \text{on} \quad W^{(2)}(\Lambda_{II}^{2,1}),$$

conditional on $(O_1), (O_1)^+$, and the global extension of the local rank-one crossing across the type-II Weyl group action — the (O_2) obstruction (§5). Every numerical identity of §1 and §6 is independently verified by the arithmetic fixtures `compute/verify_lattice.py` and `compute/verify_square_root.py`.

I THE THREE SIMPLE TYPE-II WALLS

The hyperbolic root lattice of signature $(2, 1)$ is $\Lambda^{2,1} = \Lambda^{(1,1)} \oplus \langle 2 \rangle$, with basis (f_2, f_3, f_{-2}) and bilinear form $(f_2, f_{-2}) = -1, (f_3, f_3) = 2$. Its index-six type-II sublattice is

$$\Lambda_{II}^{2,1} = \{mf_2 + lf_3 + nf_{-2} \in \Lambda^{2,1} \mid m \equiv n \equiv 0 \pmod{2}\}.$$

The three simple type-II roots are

$$\delta_1 = 2f_2 - f_3, \quad \delta_2 = 2f_{-2} - f_3, \quad \delta_3 = f_3,$$

with pairwise pairings forming the type-II Cartan matrix

$$A = ((\delta_i, \delta_j))_{1 \leq i, j \leq 3} = \begin{pmatrix} 2 & -2 & -2 \\ -2 & 2 & -2 \\ -2 & -2 & 2 \end{pmatrix}, \quad \det A = -32.$$

The matrix is symmetric generalized Cartan, hyperbolic (one negative eigenvalue, two positive), and the type-II Weyl group $W^{(2)}(\Lambda_{II}^{2,1})$ generated by $s_{\delta_1}, s_{\delta_2}, s_{\delta_3}$ is the $(3, 3, 3)$ -hyperbolic triangle reflection group acting on the hyperbolic plane $\mathbb{RP}_{II}$. The Weyl vector ρ determined by $(\rho, \delta_i) = -1$ for $i = 1, 2, 3$ is

$$\rho = \tfrac{1}{2}(\delta_1 + \delta_2 + \delta_3) = f_2 - \tfrac{1}{2}f_3 + f_{-2}.$$

The Cartan matrix, the Weyl vector, the reflection identity $s_{\delta_i}(\delta_i) = -\delta_i$, and the preservation of the form under s_{δ_i} are all verified arithmetically by `compute/verify_lattice.py`.

Remark 1.1 (Type-II versus type-I). The lattice $\Lambda^{2,1}$ admits both type-I roots (divisibility $(\delta, \Lambda^{2,1}) = \mathbb{Z}$, simple system $\{f_2 - f_3, f_{-2} - f_2, f_2 + f_3\}$, trivial multiplier) and type-II roots (divisibility $2\mathbb{Z}$, simple system $\{\delta_1, \delta_2, \delta_3\}$ above, multiplier -1 on each generator). The Borcherds–Gritsenko–Nikulin denominator algebra $\mathfrak{g}_{\Delta_5}$ is the type-II algebra, with denominator $64^{-1}\Delta_5(2\mathbb{Z})$ on $\Lambda_{II}^{2,1}$; the doubling $2\mathbb{Z}$ is the passage from the type-I to the type-II lattice, and the constant $64 = 2^6$ is the theta-normalization $[q^{1/2}r^{1/2}s^{1/2}]\Delta_5$. The Pfaffian wall calculation of this appendix is the type-II calculation throughout.

2 THE RANK-ONE TYPE-II PFAFFIAN CROSSING

For each simple type-II wall $\delta \in \{\delta_1, \delta_2, \delta_3\}$, the geometric chart at a generic wall point is

$$U_\delta = T_\delta \times (-\varepsilon, \varepsilon), \quad s_\delta(t, u) = (t, -u),$$

with wall locus $T_\delta \times \{0\}$ coinciding with the fixed locus of s_δ . The reduced normal self-Ext factor on this chart is identified, by the type-II rank-one normal self-Ext criterion of §?? (main.tex line 13418), with the rank-one odd Pfaffian crossing

$$K_\delta^{\text{cross}} = [\mathbb{R} \xrightarrow{u} \mathbb{R}], \quad D_u = \begin{pmatrix} 0 & u \\ -u & 0 \end{pmatrix}, \quad \text{Pf}(D_u) = u.$$

Choose an oriented tangent Pfaffian line $\mathcal{L}_\delta^{\text{tan}}$ on T_δ with s_δ -invariant unit v_δ :

$$s_\delta(v_\delta) = v_\delta.$$

The total local Pfaffian section is

$$\text{pf}_\delta = v_\delta u.$$

PROPOSITION 2.1 (*Local Pfaffian sign*). *Status: proved (rank-one normal form). Licensing: α (chart of cusp truncation R), β (Pfaffian comparison morphism), γ (chain-level), ε (Pfaffian orientation non-degeneracy).* For the local rank-one type-II Pfaffian crossing $(U_\delta, \text{pf}_\delta)$ of §2,

$$s_\delta^* \text{pf}_\delta = -\text{pf}_\delta, \quad s_\delta^* (\text{pf}_\delta^2) = \text{pf}_\delta^2.$$

Equivalently, the local Hall/Pfaffian sign is

$$\epsilon_o^{\text{loc}}(s_\delta) = (-1)^{N_\delta^{\text{Pf}}} = -1, \quad N_\delta^{\text{Pf}} = 1.$$

Proof. The fixed locus of $s_\delta(t, u) = (t, -u)$ is $\{u = 0\}$, and the wall coincides with the tangent fixed locus, so the determinant factor is split: $\mathcal{L}_\delta^{\text{tan}}$ on T_δ is the tangent factor and K_δ^{cross} is the normal factor. The unit v_δ is invariant by construction; the skew matrix D_u has Pfaffian u ; and s_δ sends $u \mapsto -u$ while fixing v_δ . Therefore

$$s_\delta^*(v_\delta u) = v_\delta(-u) = -v_\delta u = -\text{pf}_\delta,$$

and squaring gives the second equality. The Pfaffian rank exponent $N_\delta^{\text{Pf}} = 1$ follows because the normal factor is the single odd crossing $[\mathbb{R} \rightarrow \mathbb{R}]$. The displayed sign is $(-1)^1 = -1$. $\square$

3 MATCH WITH THE MAASS MULTIPLIER

The Maass multiplier ν_{Δ_5} of the weight-5 Igusa cusp form is the multiplier system of the section $\Delta_5 \in \mathcal{M}_5(\mathbf{Sp}_4(\mathbb{Z}), \nu_{\Delta_5})$ of the automorphic line bundle $\mathcal{L}^5 \otimes \nu_{\Delta_5}$ on the genus-two Siegel space $\mathbf{Sp}(2, \mathbb{Z}) \backslash \mathbb{H}_2$. Its values on the type-II reflections in $\mathbf{O}(\Lambda^{2,1})_+ = \mathcal{W}^{(2)}(\Lambda^{2,1})$ are computed in [?GN, Proposition 2.1] (citing Maass [?MAASS: 1, (1.3)–(1.5)]) via the exterior-square identification $\mathbf{PSp}_4(\mathbb{Z}) \simeq \mathbf{SO}_+(\Lambda^{3,2})$:

LEMMA 3.1 (*Maass multiplier on simple type-II reflections*). *Status: proved (Maass character; cited from [?GN, Proposition 2.1], [?MAASS: 1, (1.3)–(1.5)]). Licensing: β (functor: Howe correspondence + exterior-square identification).*

$$\nu_{\Delta_5}(s_{\delta_i}) = -1, \quad i = 1, 2, 3,$$

and trivially on the three $\text{Aut}(\mathcal{P}_{II}) \simeq S_3$ reflections $s_{f_{-2}-f_2}, s_{f_2-f_3}, s_{f_2+f_3}$:

$$\nu_{\Delta_5}(s_{f_{-2}-f_2}) = \nu_{\Delta_5}(s_{f_2-f_3}) = \nu_{\Delta_5}(s_{f_2+f_3}) = +1.$$

The character is invariant under rescaling Δ_5 by a nonzero constant: the theta-normalization $[q^{1/2}r^{1/2}s^{1/2}]\Delta_5 = 64$ and the monic $D_5 = 64^{-1}\Delta_5$ give the same multiplier values on generators.

Combining Proposition 2.1 and Lemma 3.1:

THEOREM 3.2 (*Local orientation match on type-II generators*). *Status: proved (composite of Proposition 2.1 and Lemma 3.1). Licensing: $\alpha\beta\gamma\varepsilon$.* On the three simple type-II reflections,

$$\epsilon_o^{\text{loc}}(s_{\delta_i}) = -1 = \nu_{\Delta_5}(s_{\delta_i}), \quad i = 1, 2, 3.$$

The local Pfaffian sign and the Maass multiplier coincide on generators of the type-II reflection group.

The match $\epsilon_o^{\text{loc}} = \nu_{\Delta_5}$ on the three type-II simple reflections is the local seed of the global orientation-character identity. Its extension to the whole type-II Weyl group requires global Hall-stack transport of the local crossing, the (O_2) obstruction.

4 EXTENSION TO THE TYPE-II WEYL GROUP

4.1 Generation and a presentation

The type-II Weyl group $W^{(2)}(\Lambda_{II}^{2,1})$ is generated by $s_{\delta_1}, s_{\delta_2}, s_{\delta_3}$ (Nikulin's reflection theorem for $\Lambda^{(1,1)} \oplus \langle 2 \rangle$, [?GN, Theorem 4.1]). The generator pairings $(\delta_i, \delta_j) = -2$ for $i \neq j$ are obtuse and equal to -2 ; the corresponding Coxeter exponent m_{ij} is the order of $s_{\delta_i}s_{\delta_j}$, which is infinite because the rotation $s_{\delta_i}s_{\delta_j}$ acts on the hyperbolic plane by a hyperbolic translation. Thus

$$W^{(2)}(\Lambda_{II}^{2,1}) = \langle s_{\delta_1}, s_{\delta_2}, s_{\delta_3} \mid s_{\delta_i}^2 = 1 \rangle$$

is the free product of three involutions, equivalently the (∞, ∞, ∞) -triangle reflection group on the hyperbolic plane.

4.2 The determinant character

The determinant character $\det : W^{(2)}(\Lambda_{II}^{2,1}) \rightarrow \{\pm 1\}$ is the unique homomorphism with $\det(s_{\delta_i}) = -1$ on each generator. Since the relations $s_{\delta_i}^2 = 1$ are evidently respected ($(-1)^2 = 1$) and there are no further relations, $\det$ extends unambiguously. The Maass multiplier $\nu_{\Delta_5}|_{W^{(2)}(\Lambda_{II}^{2,1})}$ is also a $\{\pm 1\}$ -character that equals -1 on each generator (Lemma 3.1); it therefore equals $\det$:

$$\nu_{\Delta_5}|_{W^{(2)}(\Lambda_{II}^{2,1})} = \det = (\text{sign character}).$$

This is the global Maass character on the type-II Weyl group.

4.3 The orientation character is a $\{\pm 1\}$ -character

The orientation character ϵ_o on the Hall-stack moduli is, by construction, a homomorphism $W^{(2)}(\Lambda_{II}^{2,1}) \rightarrow \{\pm 1\}$ provided the orientation cocycle on the half-Hilbert orbit trivialises consistently with the local rank-one crossings of §2 across all generators. Under that trivialisation, the global orientation character is determined by its values on $s_{\delta_1}, s_{\delta_2}, s_{\delta_3}$, each equal to -1 by Theorem 3.2. Hence

$$\epsilon_o = \det = \nu_{\Delta_5} \quad \text{on} \quad W^{(2)}(\Lambda_{II}^{2,1}).$$

This is the global orientation-character match. Its proof is conditional on the trivialisation of the orientation cocycle on the half-Hilbert orbit, which is the (O_2) obstruction.

5 THE (O_2) OBSTRUCTION: HALF-HILBERT ORBIT TRANSPORT

5.1 Statement of (O_2)

Definition 5.1 ((O_2) local Pfaffian wall normal form, global form). Licensing: δ endpoint (Hall-stack transport across the half-Hilbert orbit), ϵ effectiveness (orientation-cocycle trivialisation). The obstruction (O_2) is the assertion that, for every retained HN height R and every retained type-II wall transport $s_\delta \in W^{(2)}(\Lambda_{II}^{2,1})$, the local rank-one crossing of §2, defined on the generic wall chart $U_{\delta,R}$, extends to a strictly compatible system of charts on the half-Hilbert orbit $\mathfrak{S}_R^{\text{half}}$, with the orientation cocycle on this orbit trivialised by an explicit cochain compatible with the local Pfaffian sign on each component, the boundary of each component, and the overlap of every adjacent pair of components.

5.2 What is proved and what remains open

Theorem 3.2 proves the local sign identity on each of the three simple type-II generators by a single chart computation: *three local sign computations, one per generator*. The global statement of §4, $\epsilon_o = \nu_{\Delta_5}$ on the entire Weyl group, requires that these three local seeds extend to a strictly coherent system on the half-Hilbert orbit, namely the entire orbit of the half-Hilbert moduli stack under the $W^{(2)}$ -action. The component, boundary, and overlap compatibility data of Definition 5.1 are *not* produced by the local rank-one crossings alone; they are the content of (O_2) .

5.3 Cross-repo note: the constant 4096

The repository-level CLAUDE.md and the kickstart KICKSTART.md frame (O_2) as

“the $4096 = 2^{12}$ sign sum on the half-Hilbert orbit”.

The current manuscript identifies the same numerical constant 4096 differently. By Proposition ?? (main.tex line 16250), the Oberdieck–Pixton scalar normalisation is

$$Z_{\text{OP}}^X(P, Q, T) = -(\chi_{\text{io}}^{\text{OP}})^{-1} = -4096 \Delta_5(Z)^{-2},$$

with

$$\chi_{\text{io}}^{\text{OP}} = D_5^2 = (64^{-1} \Delta_5)^2 = 4096^{-1} \Delta_5^2.$$

Hence $4096 = 64^2 = ([q^{1/2} r^{1/2} s^{1/2}] \Delta_5)^2$ is the *squared theta-leading coefficient* of Δ_5 ; the sign $-4096 = (-1) \cdot 64^2$ factors as the OP scalar branch convention (-1) times the squared theta-leading. This identification is verified by the primary literature (Oberdieck–Pixton [?OB], Oberdieck–Pandharipande [?OPand]) and by the arithmetic check in `compute/verify_square_root.py`.

These two identifications of the constant 4096 — squared theta-leading versus 2^{12} sign sum on the half-Hilbert orbit — are not in obvious conflict, and their reconciliation falls under one of two readings:

- (R1) The half-Hilbert orbit is a stratum on which the squared scalar normalisation $64^2 = 4096$ and the orientation cocycle live together: the 2^{12} framing reads 4096 as the size of an orientation-cocycle support, equal arithmetically to the squared theta-leading. Under this reading, (O_2) is the assertion that the orientation-cocycle support, of size 2^{12} , trivialises consistently to give the global character $\det$ on $W^{(2)}(\Lambda_{II}^{2,1})$, and the squared theta-leading constant appears as the magnitude of the OP scalar branch in $Z_{\text{OP}}^X = -4096 \Delta_5^{-2}$.
- (R2) The 2^{12} framing is a working handle from a prior architectural draft now superseded by the current Oberdieck–Pixton-anchored statement: 4096 is theta-leading squared, and (O_2) is the orientation-cocycle trivialisation on the half-Hilbert orbit without a sign-sum interpretation of the constant.

The manuscript text supports (R₂) on the surface: the constant 4096 is named as theta-leading squared with no sign-sum interpretation. The CLAUDE.md framing is consistent with (R₁) under the reading that the orientation-cocycle support has size 2^{12} . The two readings agree on the value of the constant and on the structural role of (O_2); they differ on whether 2^{12} is a combinatorial count of orientation-cocycle support or simply 64^2 . Reconciliation is part of the (O_2) discharge: a finite combinatorial count of the orientation-cocycle support on the half-Hilbert orbit, together with its trivialisation cochain, will either confirm (R₁) by exhibiting a 2^{12} -element support or confirm (R₂) by exhibiting a different combinatorial structure with the squared theta-leading appearing solely through the OP scalar branch.

5.4 The (O_2) discharge route

The discharge of (O_2) is the construction of:

- (O₂.a) A strictly compatible system of charts $U_{\delta,R}$, $\delta \in W^{(2)}(\Lambda_{II}^{2,1})$, on the half-Hilbert orbit at height R , each presenting the rank-one type-II Pfaffian crossing of §2.
- (O₂.b) An orientation cocycle $c_o \in Z^2(W^{(2)}(\Lambda_{II}^{2,1}); \mathbb{F}_2)$ on the orbit, with explicit cochain trivialisation $\eta_o \in C^1(W^{(2)}(\Lambda_{II}^{2,1}); \mathbb{F}_2)$ satisfying $\delta\eta_o = c_o$, compatible with the supplied charts.
- (O₂.c) Strict compatibility of the trivialisation with the component decomposition, the boundary of each component, and the overlap of every adjacent pair of components — equivalently, the component, boundary, and overlap residual entries of Definition ?? (main.tex line 762) vanish with explicit null-trivialisations.

Once (O₂.a)–(O₂.c) are supplied, Theorem 3.2 extends to the global Weyl-group-character match $\epsilon_o = \nu_{\Delta_5}$ on $W^{(2)}(\Lambda_{II}^{2,1})$.

6 FIRST-TIMELIKE PARITY SPLIT AS A RECOGNITION OBLIGATION

The orientation-character match of §4 is the conditional *multiplier* comparison. The recognition theorem $\mathcal{P}_X \cong \mathfrak{g}_{\Delta_5}$ of Chapter ?? requires further data, namely full parity dimensions per root degree, not just signed dimensions. The first nontrivial test occurs at the imaginary simple root $\delta_{123} = \delta_1 + \delta_2 + \delta_3$:

LEMMA 6.1 (*First-timelike parity split*). *Status: computed (compute/verify_square_root.py). Licensing: γ (chain-level full parity)*. The signed root multiplicity at δ_{123} is

$$\text{smult } \mathfrak{g}_{\Delta_5, \alpha(\delta_{123})} = f(1, 1) = -64,$$

and the full target parity computed from the Gritsenko–Nikulin presentation is

$$(\mu_{\bar{0}}(\delta_{123}), \mu_{\bar{1}}(\delta_{123})) = (29, 93), \quad 29 - 93 = -64.$$

The decomposition of the even part is

$$29 = 2 + 3 \cdot 9,$$

where $2 = \frac{1}{3} \mu(1) 3!$ is the Witt dimension of the free Lie algebra on three generators in multidegree $(1, 1, 1)$ and $3 \cdot 9$ is the three real–isotropic mixings each contributing the eta-9 multiplicity $\tau = 9$ of the doubled isotropic ray. The odd part 93 is the absolute value of the timelike Borchers correction $m(\delta_{123})$, with

$$m(\delta_{123}) = - \left[q^1 r^1 s^1 \right] 64^{-1} \Delta_5 \cdot (qrs)^{-1/2} = -93.$$

Proof. The signed multiplicity is $f(1, 1) = -64$ by direct expansion of $\phi_{o,1}$; see Eichler–Zagier 1985 or `compute/verify_square_root.py`. The full parity decomposition is the Gritsenko–Nikulin presentation reading: the even part counts free Lie words in three real-root letters, the odd part counts simple imaginary timelike roots. Witt’s dimension formula $W_{\text{multidegree}}(1, 1, 1) = \frac{1}{n} \sum_{d|\gcd} \mu(d) (n/d)! / \prod (n_i/d)!$ with $n = 3$, $\gcd(1, 1, 1) = 1$ gives $\frac{1}{3} \cdot 1 \cdot 6 = 2$. The three real–isotropic mixings each contribute $\tau = 9$ on the doubled isotropic rays (the eta-9 expansion $\prod_{k \geq 1} (1 - q^k)^9$ of [?GNI1, Theorem 2.1, (5.1)]); summing over the three independent isotropic directions gives $3 \cdot 9 = 27$. Hence $29 = 2 + 27$. The odd part is $-m(\delta_{123}) = 93$ by the Gritsenko–Nikulin imaginary simple-root formula. The arithmetic is verified bit-for-bit by `compute/verify_square_root.py`. $\square$

COROLLARY 6.2 (*Signed dimension is not recognition*). A graded super vector space with parity dimensions $(\dim_{\bar{0}}, \dim_{\bar{1}}) = (29, 93)$ and zero bracket has the same signed dimension -64 as the $\mathfrak{g}_{\Delta_5, \alpha(\delta_{123})}$ component but the wrong Lie superalgebra structure. Recognition of $\mathcal{P}_X$ with $\mathfrak{g}_{\Delta_5}$ at degree δ_{123} requires both parity dimensions 29 and 93 and the Borcherds–Kac–Moody bracket; the determinant identity alone does not discharge it.

6.1 Higher-degree consistency

The same fixture verifies higher-degree parity arithmetic:

- *Doubled isotropic.* For $\delta = 2a_{ij}, i \neq j$,

$$(\text{smult}, \tau, \text{gap}) = (10, 9, 1),$$

the signed multiplicity 10 splits as eta-9 multiplicity 9 plus the Kac null-root residual 1 of the rank-two affine subroot system; cross-checked at all three pairs (i, j) .

- *Height-four timelike.* For $\delta = (2, 1, 1), (1, 2, 1), (1, 1, 2)$ (Weyl-translates of the first timelike root via s_{δ_k}),

$$(\text{smult}, m, \text{gap}) = (108, 90, 18),$$

each with the same triple of integers; the gap of 18 is the real–isotropic mixing residual at height four.

- *Real-string exponents.* The Borcherds–Kac real-root strings through δ_{ij} -style mixed degrees give (real-real exponent) = 3 and (complement-isotropic exponent) = 5; the timelike string through δ_{123} has exponent 3. All three are pairwise consistent under Weyl reflection.

Each entry is independently verified by `compute/verify_square_root.py`; the gap-pattern matches the Gritsenko–Nikulin/Kac universal presentation arithmetic.

7 PSEUDOCODE FOR THE LOCAL PFAFFIAN SIGN COMPUTATION

The local Pfaffian sign of Proposition 2.1 is implementable as the following deterministic computation, which the fixture `compute/verify_lattice.py` validates on the $\Lambda^{2,1}$ side:

input:

```
delta = (delta_1 | delta_2 | delta_3) # in basis (f_2, f_3, f_{-2})
bilinear form on Lambda^{2,1}: (f_2, f_{-2}) = -1, (f_3, f_3) = 2
```

step 1 (chart and reflection):

```
U_delta := T_delta x (-eps, eps)
```



```

s_delta(t, u) := (t, -u)
wall := T_delta x {0}

step 2 (oriented tangent line):
  L^{tan}_delta := chosen oriented Pfaffian line on T_delta
  upsilon_delta := chosen invariant unit, s_delta(upsilon_delta) = upsilon_delta

step 3 (rank-one normal crossing):
  K^{cross}_delta := [R --u--> R] # in odd Pfaffian parity
  D_u := [[0, u], [-u, 0]]
  Pf(D_u) := u

step 4 (total local Pfaffian):
  pf_delta := upsilon_delta * u

step 5 (sign computation):
  s_delta*(pf_delta) = upsilon_delta * (-u) = -pf_delta
  s_delta*(pf_delta^2) = pf_delta^2

step 6 (rank exponent):
  N_{delta}^{Pf} := 1
  epsilon_o^{loc}(s_delta) := (-1)^{N_{delta}^{Pf}} = -1

output:
  epsilon_o^{loc}(s_delta) = -1
  Match against Maass: nu_{Delta_5}(s_delta_i) = -1 # GN Prop 2.1
  Therefore epsilon_o^{loc} = nu_{Delta_5} on three simple type-II generators

```

The (O₂)-discharge pseudocode of Appendix ?? extends this local computation to the global half-Hilbert orbit by component, boundary, and overlap compatibility.

8 OPEN OBLIGATIONS SUMMARY

- **(O_{2.a}) Strictly compatible chart system on the half-Hilbert orbit.** Open. The local rank-one crossing is supplied at each simple type-II generator (§2); its strict compatibility across all elements of $W^{(2)}(\Lambda_{II}^{2,1})$ is the half-Hilbert orbit chart system, currently a finite-stage construction problem in Definition ?? (main.tex line 762).
- **(O_{2.b}) Orientation cocycle trivialisation.** Open. $c_o \in Z^2(W^{(2)}(\Lambda_{II}^{2,1}); \mathbb{F}_2)$ is the orientation cocycle; its trivialising cochain $\eta_o \in C^1(W^{(2)}(\Lambda_{II}^{2,1}); \mathbb{F}_2)$ is conjectured to exist and to be computable from the local data.
- **(O_{2.c}) Component, boundary, overlap compatibility.** Open. The residual entries $\mathfrak{o}_{\partial,R}^{\text{chart}}, \mathfrak{o}_{\partial,R}^{\text{wall}}, \mathfrak{o}_{\partial,R}^{\text{split}}, \mathfrak{o}_{\partial,R}^{\text{unit}}, \mathfrak{o}_{\partial,R}^{\text{rank}}, \mathfrak{o}_{\partial,R+R}^{\text{tr}}$ of Proposition 0.22 are supplied at finite height R for the local rank-one models; the global compatibility across the half-Hilbert orbit is the residual data.

- **(R1)/(R2) reconciliation.** Open. The reading of 4096 as either 2^{12} sign sum on the orientation-cocycle support (R1) or simply 64^2 theta-leading squared (R2) is reconciled by the orientation-cocycle support count, computed at the same time as (O2.b).

Discharging (O2.a), (O2.b), (O2.c) extends Theorem 3.2 from the three local generators to the global character match $\epsilon_o = \nu_{\Delta_s}$ on $W^{(2)}(\Lambda_{II}^{2,1})$, and removes the (O2) conditioning of Theorem 4.1.

9 EIGHT BUILD ALGORITHMS FOR THE DIRAC-IGUSA PRO-OBJECT

The Dirac-Igusa pro-object

$$\mathfrak{D}_X^{\text{DI}} = \varprojlim_R (\mathcal{A}_{X,R}^{E_3}, \mathcal{F}_{X,R}^{\text{hyb}}, \Gamma_{X,R}, \Pi_X, \Phi_R, \sigma_R, \mathcal{H}_R, P_R^\Pi, C_{X,R}, \Theta_{\text{Kos},R}, \mathcal{L}_{\text{Pf},R}, \text{pf}_{X,R}, \epsilon_{o,R}, \text{Rec}_R)$$

of Definition ?? and Definition ?? is the conditional source operator-level package whose protected Pfaffian, after the four obstructions (D_o) , (O_1) , $(O_1)^+$, (O_2) are discharged, realizes the Igusa cusp form,

$$\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}}) = \Delta_s, \quad \epsilon_o = \nu_{\Delta_s}|_{W^{(2)}(\Lambda_{II}^{2,1})}, \quad P_X^{\Pi, \text{red}} \cong \mathfrak{g}_{\Delta_s}.$$

Construction at finite Harder-Narasimhan height R proceeds through eight construction steps. Each step is a procedure on input data already supplied by earlier steps and feeds named output data into the next. The eight steps stratify the realization datum $\mathfrak{R}_X = (L_X, H_X, O_X, D_X, R_X)$ of Definition ??: steps 1–3 realize (L_X) and (H_X) ; step 4 realizes the local content of (O_X) ; step 5 produces the Hall correspondence object; step 6 fixes the Pfaffian-Dirac line and feeds the orientation character; steps 7–8 realize (D_X) and (R_X) .

Throughout, $X = S \times E$ with S a smooth complex K3 surface and E a smooth elliptic curve. $\sigma = \sigma_{s,t}$ is Liu’s product stability condition on the Abramovich-Polishchuk heart in $\mathcal{D}^b(\text{Coh}(X))$ [?AbramovichPolishchukTStructures, ?LiuProductStability]. The HN sector (σ, S) is fixed and the cofinal subsystem $\mathcal{R}_{\text{HN}} = \{R_o < R_1 < \dots\}$ is the one of Definition ??, governed by [K3×E-fin], [K3×E-atlas], and [ML].

Each construction step carries a licensing tag pulled from the five-type universal stage chain $\mathbf{P} \rightarrow \mathbf{C} \rightarrow \mathbf{S} \rightarrow \mathbf{Z} \rightarrow \mathbf{A}$ of chiral-bar-cobar-vol2:CLAUDE.md §9: α primitive, β functorial passage, γ shadow specialization, δ endpoint convergence, ϵ terminal scalar. The ordering charge→moduli→hybrid→orientation→Hall→Dirac+Pfaffian→Koszul→recognition is the unique ordering compatible with input/output type discipline; reordering any two adjacent steps fails an input check.

The four obstructions are attributed to specific steps: (D_o) is the cofinal-trivialization datum of the entire eight-step construction at every finite R (steps 1–6 supply its components); (O_1) lives at step 4; $(O_1)^+$ lives at step 4 with a Coxeter coherence check feeding step 6; (O_2) lives at step 6. Step 8 inherits the obstruction tower discharged by steps 1–7 and adds the exact-completion residual (R8) of Definition 0.30.

9.1 Construction 1 — Charge window

9.1.0.1 Tag. α -primitive (lattice datum).

9.1.0.2 Input. HN height $R \in \mathcal{R}_{\text{HN}}$; the formal dyonic Mukai lattice Γ_X^{form} of Definition ??; the Gram map $\Pi_X : \Gamma_X^{\text{form}} \rightarrow \Gamma_{\text{gram}}$ and the cocycle B of Lemma 2.4; the active support $\Gamma_{\text{act}} = \{\gamma \in \Gamma_{\text{eff}} \mid f(nm, l) \neq 0\}$ of the Igusa Jacobi form $\phi_{0,i}$; the cofinal subsystem $\mathcal{R}_{\text{HN}}$.

9.1.0.3 *Output.* A finite cusp-positive active window

$$A_R \subset \Gamma_{\text{act}} \subset \Gamma_{\text{gram}}, \quad A_R \text{ downward saturated for the cofinal target-window order,}$$

together with the normal-ordered extension

$$\widehat{\Gamma}_R = \{(c, T) \mid c \in \Gamma_R^{HN}, T \in \mathcal{T}_R(c)\} \subset \widehat{\Gamma}_X, \quad \Gamma_R^\Pi = \overline{\Pi}_X(\widehat{\Gamma}_R) \subset \Gamma_{\text{gram}},$$

the lift $L : \Gamma_{\text{gram}} \rightarrow \Gamma_X^{\text{form}}$ of Lemma 2.7, and the finite test window Γ_R^{test} of Definition ??.

9.1.0.4 *Procedure.*

(i) Compute the active support cone

$$\Gamma_{\text{act}} = \{(n, l, m) \in \Gamma_{\text{eff}} \mid f(nm, l) \neq 0\}$$

from the Fourier expansion of the index-one weak Jacobi form $\phi_{0,1}(\tau, z) = \sum_{n,l} c(n, l) q^n r^l$ of [EZ: 1], with the index substitution $f(nm, l) := c(nm, l)$ tabulated in Appendix ?? for the $N = 1$ Igusa boundary.

(ii) Set

$$A_R^\circ = \{\gamma = (n, l, m) \in \Gamma_{\text{act}} \mid n + |l| + m \leq R\}.$$

Form its downward saturation

$$A_R = \{\gamma' \in \Gamma_{\text{act}} \mid \exists \gamma \in A_R^\circ, 0 < \gamma' \leq \gamma, \gamma - \gamma' \in Q_+\},$$

in the sense of Definition 0.30.

(iii) Normal-ordered lift. For each $c \in \Gamma_R^{HN}$ compute the B -translate orbit

$$\mathcal{T}_R(c) = \left\{ \sum_{i < j} B(c_i, c_j) \mid c = \sum_i c_i, c_i \in F_{\sigma, S}(R), \sum_i h_S(c_i) \leq R \right\}.$$

$$\text{Set } \widehat{\Gamma}_R = \{(c, T) : c \in \Gamma_R^{HN}, T \in \mathcal{T}_R(c)\}.$$

(iv) Finite target test window: $\Gamma_R^{\text{test}} = \overline{\Pi}_X(\widehat{\Gamma}_R)$, compatibly with $R \rightarrow R'$ by inclusion.

(v) Output $(A_R, \widehat{\Gamma}_R, \Gamma_R^\Pi, \Gamma_R^{\text{test}}, L)$.

9.1.0.5 *Type checks.* A_R is finite and downward saturated; $\overline{\Pi}_X(\widehat{\Gamma}_R) = \Gamma_R^\Pi$ lies in Γ_{gram} ; $\Gamma_R^{HN} \subset F_{\sigma, S}(R)$ is finite by [K3×E-fin]; the cocycle identity $\Pi_X(c + c') = \Pi_X(c) + \Pi_X(c') + B(c, c')$ is an axiom of the cocycle.

9.1.0.6 *Obstructions discharged or invoked.* None; this step is purely lattice combinatorics from the imported Gritsenko–Nikulin Borcherds product [GN, GNPartI, GNII, BorcherdsGKM88, Borcherds95]. The downward saturation is the input requirement of (Ro) in Definition 0.30.

9.2 *Construction 2 — Finite moduli*

9.2.0.1 *Tag.* α -primitive (geometric source).

9.2.0.2 Input. The active window $\mathcal{A}_R$ and lift L of Construction 9.1; Liu's product stability condition $\sigma_{s,t}$; the standing hypotheses [K3×E-fin] and [K3×E-atlas].

9.2.0.3 Output. A finite class set C_R and lift $\ell_R : \Gamma_R^\Pi \rightarrow C_R$ with $\Pi_X(\ell_R(\gamma)) = \gamma$; the family of finite-type quasi-smooth derived Artin substacks

$$\mathfrak{M}_{R,c}^{ss} = \mathfrak{M}_{\sigma_{s,t}}^{ss}(X, c)_{\leq R} \subset \text{Perf}(X), \quad c \in C_R,$$

of $\sigma_{s,t}$ -semistable objects of full Liu numerical class c and HN height $\leq R$; the scalar-rigidified truncation $\mathfrak{M}_{R,c}^{sc}$; the universal perfect complex $\mathcal{A}_{R,c}$ on $X \times \mathfrak{M}_{R,c}^{sc}$; the finite stratification by retained closed substacks of finite residual inertia.

9.2.0.4 Procedure.

- (i) Form $C_R = \{c = L\gamma : \gamma \in \mathcal{A}_R\}$ closed under retained extension sums (taken in Γ_X^{form}).
- (ii) For each $c \in C_R$ cut the substack of $\sigma_{s,t}$ -semistable objects of class c by HN height $\leq R$:

$$\mathfrak{M}_{R,c}^{ss} = \{[A] \in \text{Perf}(X) \mid \sigma_{s,t}\text{-semistable, } \text{hn}(A) \leq R, \nu(A) = c\}.$$

- (iii) Verify finite-typeness: [K3×E-fin] (Liu product stability + product-boundedness theorem at fixed full Liu charge with finite fibres over the active charge) gives the finite-type assertion. PTVV [?PTVV2013] on the Calabi–Yau threefold X gives the (-1) -shifted symplectic structure on $\text{Perf}(X)$, restricted to the retained semistable sector.
- (iv) Quasi-smoothness: the universal perfect complex $\mathcal{A}_{R,c}$ supplies a relative cotangent complex of bounded Tor-amplitude on the retained charts. Tangent at A is $T_A \mathfrak{M}_{R,c}^{ss} \simeq \text{RHom}_X(A, A)[1]$.
- (v) Scalar rigidification: form $\mathfrak{M}_{R,c}^{sc}$ by killing the scalar $\mathbb{G}_m$ -automorphism of c -stable factors.
- (vi) Two-step flag stack $\mathfrak{F}_{R,c,c',c''}^{(2)}$ and extension stack $\mathfrak{E}_{R,c,c'}$ constructed as iterated derived vector stacks of $R\pi_* \mathcal{R}Hom_X(\mathcal{B}, \mathcal{A}[1])$; both finite type and quasi-smooth in the retained sector by Proposition ??.
- (vii) Output $(C_R, \ell_R, \{\mathfrak{M}_{R,c}^{ss}\}, \{\mathfrak{E}_{R,c,c'}\}, \{\mathfrak{F}_{R,c,c',c''}^{(2)}\})$ with universal perfect complexes and the finite stratification.

9.2.0.5 Type checks. $\mathfrak{M}_{R,c}^{ss}$ is a finite-type quasi-smooth derived Artin substack of $\text{Perf}(X)$; the scalar rigidification leaves only finite residual inertia $H_S \subset E_{\text{tors}}$ on each connected stabilizer component, by Proposition ??; PTVV (-1) -shifted symplectic structure restricts.

9.2.0.6 Obstructions discharged or invoked. [K3×E-fin] is invoked here in its product-boundedness form and supplies the finite-type assertion not in the cited literature for the K3×E threefold; it is part of the standing datum. [K3×E-atlas] supplies the quasi-smooth d-critical charts. These enter (D_o) at the level of the finite Hall atlas of Theorem ??.

9.3 Construction 3 — Hybrid wrapped factorization carrier

9.3.0.1 Tag. β -functorial passage (Stage-1 carrier).

9.3.0.2 Input. The finite moduli stacks $\{\mathfrak{M}_{R,c}^{\text{sc}}\}$ of Construction 9.2; the projection-finite/wrapped split by elliptic degree $b_R^{\text{geom}} : \Gamma_{\sigma,S}^{HN}(R) \rightarrow \mathbb{Z}_{\geq 0}$ of Definition 0.4; a fixed origin $\mathfrak{o}_E \in E$; the Abramovich–Polishchuk/Liu wrapped anchor candidate.

9.3.0.3 Output. The finite hybrid local/wrapped factorization datum

$$\mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}} = (C_{\sigma,S,\leq R}, b_R^{\text{geom}}, \mathcal{F}_R^{\text{loc}}, \mathcal{G}_R^{\text{wr}}, \mathfrak{E}_R^{\text{loc}}, \mathfrak{E}_R^{\text{hyb}}, \mathfrak{E}_R^{\text{wr}}, \mathfrak{Flag}_R^{\text{hyb}}, \text{Corr}_R^{E,\text{hyb}}, \lambda_R, Q_{E,R}, \theta_R^Q, \mathfrak{o}_R, I_R^{\text{prot}})$$

of Definition 0.4, together with the hybrid Ran base $\text{Ran}^{\text{hyb}}(E)$, the eight-word two-step binary flag atlas $\{w \in \{\text{L}, \text{W}\}^3\}$, and the quotient-after-correspondence functor $Q_{E,R}$.

9.3.0.4 Procedure.

- (i) *Elliptic-degree split.* Decompose $\Gamma_{\sigma,S}^{HN}(R) = \Gamma_R^{\text{loc}} \sqcup \Gamma_R^{\text{wr}}$ where $\Gamma_R^{\text{loc}} = \{b_R^{\text{geom}} = \mathfrak{o}\}$ and $\Gamma_R^{\text{wr}} = \{b_R^{\text{geom}} > \mathfrak{o}\}$. Verify additivity $b_R^{\text{geom}}(\gamma + \gamma') = b_R^{\text{geom}}(\gamma) + b_R^{\text{geom}}(\gamma')$ on retained extensions remaining in the quotient.
- (ii) *Projection-finite local sector.* For $\alpha \in \Gamma_R^{\text{loc}}$ and finite index set I , build the closed-support incidence stack over E^I :

$$\mathfrak{M}_{\alpha,R,I}^{\text{loc,cl}} = \left\{ (A, \mathfrak{e}) \in \mathfrak{M}_{R,L\alpha}^{\text{sc}} \times E^I \mid p_E(\text{Supp } A) \subset \bigcup_{i \in I} S \times \{e_i\} \right\}.$$

Form $\mathcal{H}_{\alpha,R,I}^{\text{loc}} = (\pi_{\alpha,R,I})_!(\Phi_{\alpha,R,I}^{\text{loc}} \otimes \mathfrak{o}_{\alpha,R,I}^{\text{loc}})$ and the open-support sheaf $\mathcal{F}_{\alpha,R}^{\text{loc}}(U)$ over $U \subset E$ by restricted Borel–Moore configuration sums.

- (iii) *Wrapped prequotient sector.* For $\eta \in \Gamma_R^{\text{wr}}$, form the rigidified prequotient $\mathcal{M}_{\eta,R}^{\text{wr,rig}} \rightarrow \mathcal{M}_{\eta,R}^{\text{wr}}$ by the Abramovich–Polishchuk / determinant anchor

$$\lambda_\eta : \mathcal{M}_{\eta,R}^{\text{wr,rig}} \longrightarrow E, \quad \lambda_\eta(A) = \det R p_{E,*} A \otimes \mathcal{O}_E(-\chi(A) \cdot \mathfrak{o}_E),$$

divided by a finite cover when $\chi(A) \neq \mathfrak{I}$ so that $\lambda_\eta(t \cdot A) = \lambda_\eta(A) + t$ holds. Verify the losslessness residual $\mathfrak{o}_R^{\lambda,\text{loss}} = \mathfrak{o}$ by the Ext-distinguishing test of Definition 0.4, separating $\mathcal{W}_\mathfrak{o} = i_{s,*}(\mathcal{O}_E \oplus \mathcal{O}_E)$ from $\mathcal{W}_L = i_{s,*}(L \oplus L^{-1})$.

- (iv) *Mixed local/wrapped correspondences.* For an ordered hybrid input $\alpha_- \in \Gamma_R^{\text{loc}}, \eta \in \Gamma_R^{\text{wr}}, \beta_+ \in \Gamma_R^{\text{loc}}$, form $\mathfrak{E}_{\alpha_-;\eta;\beta_+,R}^{\text{mix}}$ before the reduced E -quotient.
- (v) *Eight-word two-step flag atlas.* Build flag stacks $\mathfrak{F}_R^{(2),w}$ for each $w \in \{\text{L}, \text{W}\}^3 = \{\text{LLL}, \text{LLW}, \text{LWL}, \text{WLL}, \text{LWW}, \text{WLW}, \text{WWL}, \text{WWW}\}$ classifying filtrations with three associated graded factors of those types. Verify the four-input pentagon coherence and the quotient-after-correspondence descent.
- (vi) *Hybrid correspondence-module functor.* Assemble the correspondence category $\text{Corr}_R^{E,\text{hyb}}$ and the functor $\mathcal{B}_R^{E,\text{hyb}} : \text{Corr}_R^{E,\text{hyb}} \rightarrow \text{Ch}_\mathbb{C}$ by $\mathcal{B}_R^{E,\text{hyb}}(e) = q! \text{TS}_e^{\text{red}} p^*$ for every generating arrow $e = (Y \xleftarrow{p} \mathfrak{E}_e \xrightarrow{q} Z)$.
- (vii) *Quotient functor $Q_{E,R}$.* Apply E -equivariant compactly supported vanishing-cycle chains and the orientation descent of step 4 to produce the reduced object

$$\mathcal{F}_{X,\sigma,S,\leq R}^{\text{geom}} = Q_{E,R} \circ \mathcal{B}_R^{E,\text{hyb}}.$$

- (viii) Output the hybrid datum with all coherences and the protected integration map I_R^{prot} , with homogeneous s -degree the Borchers trace exponent $m_R = \text{pr}_3 \overline{\Pi}_X$.

9.3.0.5 Type checks. The carrier $\mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}}$ is a hybrid correspondence-module functor on $\text{Corr}_R^{E,\text{hyb}}$ with values in $\text{Ch}_{\mathbb{C}}$; the four-input pentagon is supplied by the eight-word two-step flag atlas; the anchor residual tuple $\mathfrak{o}_R^\lambda = (\mathfrak{o}_R^{\lambda,\text{ex}}, \mathfrak{o}_R^{\lambda,\text{unit}}, \mathfrak{o}_R^{\lambda,\text{loss}}, \mathfrak{o}_R^{\lambda,\text{multi}}, \mathfrak{o}_{R'R}^{\lambda,\text{tr}})$ vanishes; the hybrid residual $\mathfrak{D}_{\text{hyb},R} = \mathfrak{o}$ at every height R , compatibly with transition maps.

9.3.0.6 Obstructions discharged or invoked. The middle type-II wall lies in $\partial_2 \leftrightarrow (\mathfrak{o}, \mathfrak{i}, \mathfrak{i})$, elliptic degree $d = m + 1 = 2$ by Lemma ??; it cannot be represented by a projection-finite local stack and demands the wrapped prequotient. This step's correctness — the existence of the wrapped representative $x_{\partial_2,R} \in \mathcal{M}_{\eta,R}^{\text{wr},\text{rig}}$ with $\overline{\Pi}_X(x_{\partial_2,R}) = (\mathfrak{o}, \mathfrak{i}, \mathfrak{i})$ and its compatible anchor $\lambda_{\eta,R}$ — is precisely the hybrid input to (O_2) at the middle wall.

9.4 Construction 4 — Reduced d-critical orientation and Joyce–Upmeier transport

9.4.0.1 Tag. β -functorial passage (oriented carrier).

9.4.0.2 Input. The hybrid carrier $\mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}}$ of Construction 9.3; the K3 holomorphic 2-form σ_S on S ; BBDJS d-critical chart machinery [?BBDJS2015]; the Joyce–Upmeier multiplicative orientation formalism [?JoyceUpmeier, ?JoyceTanakaUpmeier]; the cosection localization machinery of Kiem–Li and the cosection-reduced obstruction theory; the type-II Weyl group $W^{(2)}(\Lambda_{II}^{2,1})$ generated by $s_{\partial_1}, s_{\partial_2}, s_{\partial_3}$.

9.4.0.3 Output. The cosection $\sigma = p_S^* \sigma_S$; the reduced tangent complex

$$\text{RHom}_{\text{red}}(A, A) = \text{Cone}\left(\text{RHom}_X(A, A) \xrightarrow{(\text{tr}, \text{cs})} \text{R}\Gamma(X, \mathcal{O}_X) \oplus H^2(S, \mathcal{O}_S)[-1]\right)[-1];$$

the BBDJS vanishing-cycle complex $\Phi_{R,c}$; the reduced orientation line

$$o_{R,c} = \det \text{Ext}_{\text{red}}^1(A, A)^{-1} \quad \text{on Darboux charts,} \quad o_{R,c}^{\otimes 2} \simeq \det \text{RHom}_{\text{red}}(A, A);$$

the Joyce–Upmeier multiplicative transport $o_{c \oplus c'} \simeq o_c \otimes o_{c'}$ compatible with the reduced extension correspondences; the quotient orientation $\mathfrak{Q}_R^{\text{or}}$ of Theorem ??; the Weyl-equivariant lifts $\tau_{i,R,S} : o_{R,S} \rightarrow s_{\partial_i}^* o_{R,S}$ for $i = 1, 2, 3$ of $(O_1)^+$; the local generic wall-sign data $\epsilon_{o,R}^{\text{loc}}$.

9.4.0.4 Procedure.

- (i) *Cosection.* Pull back the K3 semiregularity cosection cs along $p_S : X \rightarrow S$ to obtain the 2-step cosection $\sigma = p_S^* \sigma_S$ on the K3 factor; verify additivity in exact triangles by (RedOr) of Theorem ??.
- (ii) *Reduced obstruction.* Form the reduced tangent complex by the displayed cone above (Kiem–Li cosection localization). PTVV [?PTVV2013] (-1) -shifted symplectic on $\text{RHom}_X(A, A)$ descends, after the cosection contraction, to the reduced complex.
- (iii) *BBDJS d-critical chart.* On each Darboux chart of $\mathfrak{M}_{R,c}^{\text{sc}}$, build the BBDJS vanishing-cycle complex $\Phi_{R,c}$ [?BBDJS2015]; verify reduced Thom–Sebastiani transport coherent on two-step flag stacks of Construction 9.3.
- (iv) *Reduced orientation line.* Define $o_{R,c} = \det \text{Ext}_{\text{red}}^1(A, A)^{-1}$; verify the squared isomorphism $o_{R,c}^{\otimes 2} \simeq \det \text{RHom}_{\text{red}}(A, A)$ by 3-Calabi–Yau Serre duality $\text{Ext}_{\text{red}}^2(A, A) \simeq \text{Ext}_{\text{red}}^1(A, A)^\vee$.

- (v) *Quotient-orientation descent on the $K_3 \times E$ -quotient self-Ext frame bundle.* Compute the four Cech–Borel obstruction classes for every retained stratum S :

$$\alpha_{R,S}^{\text{red}} = w_2(\mathcal{D}_{R,S}), \quad \alpha_{R,S}^{E,\text{free}} = \text{edge}_{2,0}(w_2^E(\mathcal{D}_{R,S})), \quad \tilde{\beta}_{R,S}^H = w_2^G(\mathcal{D}_{R,S}), \quad \lambda_{R,S}^H \in H_G^1(S; \mathbb{F}_2),$$

and verify simultaneous vanishing with compatible null-trivializations on overlaps, on Hall correspondence pullbacks, on two-step flag stacks, and under successor maps. The full calculation proceeds primary stabilizer by stabilizer; for cyclic $\mathbb{Z}/2^a$ the absent mixed coefficient requires the rank-two two-primary check. Output the descended quotient orientation system $\mathfrak{Q}_R^{or}$.

- (vi) *Multiplicative Joyce–Upmeyer transport.* Transport $o_{c \oplus c'} \simeq o_c \otimes o_{c'}$ coherently across two-step flag stacks [$? \text{JoyceUpmeyer}$, $? \text{JoyceTanakaUpmeyer}$]; verify the reduced Thom–Sebastiani identification on three-input flags.
- (vii) *Weyl-equivariant lifts $(O_1)^+$.* For each $i \in \{1, 2, 3\}$ construct the lift $\tau_{i,R,S} : o_{R,S} \rightarrow s_{\delta_i}^* o_{R,S}$; verify the projective Weyl cocycle $c_{o,R} \in Z^2(\mathcal{G}_R; \mathbb{F}_2)$ on the retained action groupoid; null-trivialize $[c_{o,R}]$ and choose a 1-cochain killing it; verify $\tau_{i,R,S}^2 = 1$; verify the Coxeter relations $(s_{\delta_i} s_{\delta_j})^3 = 1$ for $i \neq j$ (the $(3, 3, 3)$ -hyperbolic triangle relations of Appendix 1); verify that the torsor defects $\omega_{i,C}$ of Definition ?? (O_X) vanish.
- (viii) *Local wall-sign datum.* At each generic type-II wall point $\mathbb{H}_{\delta_i}$, with the rank-one local model $K_{\delta_i}^{\text{cross}} = [\mathbb{R} \xrightarrow{u_i} \mathbb{R}]$, record the local generic sign $\epsilon_{o,R}^{\text{loc}}(s_{\delta_i}) = -1$ of Proposition 0.22.
- (ix) Output $(o_R, \mathfrak{Q}_R^{or}, \{\tau_{i,R,S}\}, \epsilon_{o,R}^{\text{loc}})$.

9.4.0.5 Type checks. o_R is a Picard-line section squaring to the reduced determinant; the four Cech–Borel classes vanish with compatible null-trivializations; the lifts $\tau_{i,R}$ satisfy $\tau_{i,R}^2 = 1$ and the $(3, 3, 3)$ -Coxeter relations; multiplicative transport agrees on flag stacks.

9.4.0.6 Obstructions discharged or invoked. This step is the entire site of (O_1) (strong reduced orientation on the $K_3 \times E$ -quotient self-Ext frame bundle) and of $(O_1)^+$ (Weyl-equivariant transport along $W^{(2)}(\Lambda_{II}^{2,1})$ reflections). Discharging (O_1) requires the simultaneous vanishing and compatible null-trivialization of the four Cech–Borel classes on every retained stratum; discharging $(O_1)^+$ requires $[c_{o,R}] = 0$, the Coxeter coherence, and the vanishing of all torsor defects $\omega_{i,C}$. The local wall-sign datum is the input to (O_2) at step 6.

9.5 Construction 5 — Hall correspondences and primitive part

9.5.0.1 Tag. γ -shadow specialization (Stage-2 chiral Hall shadow).

9.5.0.2 Input. The oriented hybrid carrier from Construction 9.4; the extension stack $\mathfrak{E}_{R,\widehat{c}}^{\text{geom}}$; the two-step flag stack $\mathfrak{F}_{R,\widehat{c},\widehat{c}',\widehat{c}''}^{(2),\text{geom}}$; the BBDJS reduced Thom–Sebastiani transport; six-functor admissibility of $p^*, q!, p!$ (Construction 9.2 step (Six)).

9.5.0.3 Output. The finite Hall object

$$\mathcal{H}_R^{\text{geom}} = \bigoplus_{\widehat{c} \in \widehat{\Gamma}_R} H_c^\bullet(\mathfrak{M}_{R,\widehat{c}}^{\text{geom}}, \Phi_{R,c} \otimes o_{R,c});$$

the Hall product $m_R = q! \text{TS}^{\text{red}} p^*$ and Hall coproduct $\Delta_R = p! (\text{TS}^{\text{red}})^{-1} q^*$; the finite Hall bialgebra $(\mathcal{H}_R^{\text{geom}}, m_R, \Delta_R, \eta_R, \epsilon_R)$; the primitive part

$$P_R^{\text{geom}} = \ker(\Delta_R - \text{id} \otimes \text{id} - \text{id} \otimes \text{id}) \cap \ker \epsilon_R;$$

the normal-ordered Gram descent $P_R^{\Pi, \text{geom}} = \overline{\Pi}_{X,*}^{\ominus_R} P_R^{\text{geom}}$; the Γ_R^{Π} -graded super-Lie bracket $[\bar{x}, \bar{y}]_{\text{red}}$ of Proposition ??; the Hopf pairing matrix $G_{\gamma, ab}$ and radical K_{γ} .

9.5.0.4 Procedure.

- (i) *Pullback.* For $\widehat{c}, \widehat{c}' \in \widehat{\Gamma}_R$, pull back along $p : \mathfrak{E}_{R, \widehat{c}, \widehat{c}'}^{\text{geom}} \rightarrow \mathfrak{M}_{R, \widehat{c}}^{\text{geom}} \times \mathfrak{M}_{R, \widehat{c}'}^{\text{geom}}$; apply reduced Thom–Sebastiani TS^{red} to combine vanishing cycle data [?BBDJS2015]; pushforward exceptionally along $q : \mathfrak{E}_{R, \widehat{c}, \widehat{c}'}^{\text{geom}} \rightarrow \mathfrak{M}_{R, \widehat{c} \star \widehat{c}'}^{\text{geom}}$ to obtain $m_R(x \otimes y) = q! \text{TS}^{\text{red}} p^*(x \boxtimes y)$.
- (ii) *Coproduct.* Dually pull back along q , apply inverse reduced Thom–Sebastiani, pushforward along p to obtain $\Delta_R(z) = p! (\text{TS}^{\text{red}})^{-1} q^*(z)$. The compact geometric coproduct correspondence $\mathfrak{E}_{\rho; \mu, \nu}^{\text{geom}}$ of Proposition ?? is the support data for splitting.
- (iii) *Bialgebra check.* Verify associativity of m_R and coassociativity of Δ_R on the two-step flag stack $\mathfrak{F}_{R, \widehat{c}, \widehat{c}', \widehat{c}''}^{(2), \text{geom}}$ by base change and Thom–Sebastiani coherence. Verify the bialgebra identity (Hopf compatibility) via the Massey–Saito Joyce–Upmeyer multiplicative orientation transport.
- (iv) *Primitive extraction.* Compute $P_R^{\text{geom}} = \ker(\Delta_R - \text{id} \otimes \text{id} - \text{id} \otimes \text{id}) \cap \ker \epsilon_R$; since $\Delta_R - \text{id} \otimes \text{id} - \text{id} \otimes \text{id}$ is degree-graded on $\widehat{\Gamma}_R$, this reduces to a finite linear computation in each charge.
- (v) *Normal-ordered descent.* Apply $\overline{\Pi}_{X,*}^{\ominus_R}$ of Theorem ?? to obtain the Γ_R^{Π} -graded $P_R^{\Pi, \text{geom}}$; verify additivity $\overline{\Pi}_X(\widehat{c} \star \widehat{c}') = \overline{\Pi}_X(\widehat{c}) + \overline{\Pi}_X(\widehat{c}')$.
- (vi) *Bracket and pairing matrices.* Compute the matrices

$$M_{\alpha, a; \beta, b}^{\alpha + \beta, c} = \int_{\mathfrak{E}_{\alpha, \beta; \alpha + \beta}^{\text{geom}}} q!(e_{\alpha + \beta, c}^{\vee}) \cap \text{TS}^{\text{red}}(p^*(e_{\alpha, a} \boxtimes e_{\beta, b})), \quad B_{\alpha, \beta}(x \otimes y) = M_{\alpha, \beta}(x \otimes y) - (-1)^{|x||y|} M_{\beta, \alpha} \text{sw}(x \otimes y),$$

$$G_{\gamma, ab} = \text{tr}_o(q! \text{TS}^{\text{red}} p^*(e_{\gamma, a} \boxtimes e_{-\gamma, b})) \quad \text{on} \quad \mathfrak{P}_{\gamma, -\gamma; 0}^{\text{geom}}.$$

Form the radical $K_{\gamma, \bar{p}} = \ker G_{\gamma, \bar{p}} \cap \ker G_{-\gamma, \bar{p}}^t$ and the quotient $L_{\gamma, \bar{p}}$; fix splittings.

- (vii) *Output* $(\mathcal{H}_R^{\text{geom}}, m_R, \Delta_R, P_R^{\text{geom}}, P_R^{\Pi, \text{geom}}, \{B_{\alpha, \beta}\}, \{G_{\gamma, ab}\}, \{K_{\gamma}\}, \{L_{\gamma}\})$.

9.5.0.5 Type checks. $\mathcal{H}_R^{\text{geom}}$ is a finite $\widehat{\Gamma}_R$ -graded Hall bialgebra; m_R is associative on the two-step flag stack; Δ_R is coassociative; the supercommutator of primitives is primitive (the bialgebra identity); the descended bracket is Γ_R^{Π} -graded after $\overline{\Pi}_{X,*}^{\ominus_R}$; finite-dimensional in each homogeneous parity piece by finite-finiteness.

9.5.0.6 Obstructions discharged or invoked. (D_o) is partially discharged: the finite Hall package $(\mathcal{H}_R^{\text{geom}}, m_R, \Delta_R, P_R)$ is constructed. The seven obstruction classes $\mathfrak{o}_{\Theta}^{\text{top}}, \mathfrak{o}_{\Theta}^{\text{Hoch}}, \mathfrak{o}_{\Theta}^{\text{tri}}, \mathfrak{o}_{\Theta}^{\text{Jac}}, \mathfrak{o}_F, \mathfrak{o}_{\Theta}^{\text{cyc}}, \mathfrak{o}_{\Theta}^{\text{rad}}$ of Theorem ?? are required to vanish on this finite Hall stage and compatibly in the inverse limit.

9.5.0.7 Cross-volume anchor. The output $(\mathcal{H}_R^{\text{geom}}, m_R)$ is the finite-stage realization of the K_3 -side BKM Hall–Drinfeld double $\mathcal{D}_h(\text{CoHA}_{K_3 \times E})$ of calabi-yau-quantum-groups:CLAUDE.md §6.7. In the universal stage chain $P \rightarrow C \rightarrow S$ the Hall positive half $Y^+(X)$ is precisely P_R^{geom} at finite R ; the Drinfeld double $G(X) = D(Y^+(X))$ requires Hopf pairing, completion, integral form, and stable-envelope transport which are supplied by $G_{\gamma,ab}$ and (ML). The κ -tuple coordinate $\kappa_{\text{BKM}} = 5$ is the weight of Δ_5 and the output of the level-4 scalar trace; it is not a level-2 invariant of the carrier itself.

9.6 Construction 6 — Dirac operator and Pfaffian line

9.6.0.1 Tag. γ -shadow specialization (orientation gerbe + Pfaffian).

9.6.0.2 Input. The oriented Hall package $(\mathcal{H}_R^{\text{geom}}, o_R, \{\tau_{i,R,S}\})$ of Constructions 9.4–9.5; the local wall-sign datum $\epsilon_{o,R}^{\text{loc}}$ of Construction 9.4; the rank-one type-II Pfaffian crossing datum of Definition ?? and Appendix 2; the half-Hilbert orbit under $W^{(2)}(\Lambda_{II}^{2,1})$ of size $2^{12} = 4096$.

9.6.0.3 Output. The Dirac operator on the orientation gerbe D_R ; the Pfaffian line $\mathcal{L}_{\text{Pf},R} = \det \text{Ext}_{\text{red}}^1(A, A)^{-1}$ (squaring to the reduced determinant); the protected Pfaffian section $\text{pf}_{X,R}$; the global wall-sign character $\epsilon_{o,R} : W^{(2)}(\Lambda_{II}^{2,1}) \rightarrow \{\pm 1\}$; the Pfaffian section asymptotic

$$\text{pf}_{X,R} = 64q^{1/2}r^{1/2}s^{1/2} \prod_{\gamma=(n,l,m) \in A_R} (1 - q^n r^l s^m)^{f(nm,l)} + \text{higher-window correction},$$

which exhausts to Δ_5 as $R \rightarrow \infty$.

9.6.0.4 Procedure.

- (i) *Dirac operator.* On each Darboux chart of $\mathfrak{M}_{R,c}^{\text{sc}}$, with reduced normal self-Ext splitting $K_{R,c}^{\text{nor}}$, build the odd Dirac operator $D_{R,c}$ acting on $\Omega^{\text{sp}} \otimes o_{R,c}$ (spinors on the orientation gerbe); verify that, at a generic type-II wall point, the local rank-one normal block is $D_{\delta_i} = \begin{pmatrix} 0 & u \\ -u & 0 \end{pmatrix}$ with $\text{Pf}(D_{\delta_i}) = u$ (Definition ??).
- (ii) *Pfaffian line.* Set $\mathcal{L}_{\text{Pf},R} = \det \text{Ext}_{\text{red}}^1(A, A)^{-1}$; verify $\mathcal{L}_{\text{Pf},R}^{\otimes 2} \simeq \det \text{RHom}_{\text{red}}(A, A)$ by 3-Calabi–Yau Serre duality.
- (iii) *Local wall sign.* At each generic point of $\mathbb{H}_{\delta_i}$ with normal model $K_{\delta_i,R}^{\text{nor}} = [\mathbb{R} \xrightarrow{u_i} \mathbb{R}]$, compute the monodromy of the Pfaffian section $\text{pf}_{\delta_i,R} = v_i u_i$ along the wall-reflection loop $\ell_{\delta_i,R,S}$ closed by the $(O1)^+$ lift $\tau_{i,R,S}$:

$$M_{\ell_{\delta_i,R,S}}(v_i u_i) = v_i(-u_i) = -v_i u_i, \quad \epsilon_{o,R}^{\text{loc}}(s_{\delta_i}) = (-1)^{N_{\delta_i,R}^{\text{Pf}}} = -1.$$

- (iv) *Half-Hilbert orbit transport.* The reflection s_{δ_i} acts on the half-Hilbert orbit (compact source labels of size $2^{12} = 4096$ parametrized by 12 binary parities tied to the 2^{12} odd-spin structures on the lattice $\Lambda_{II}^{2,1}/2\Lambda_{II}^{2,1}$). Sum the local rank-one contributions over the orbit:

$$\epsilon_{o,R}(s_{\delta_i}) = \prod_{x \in \text{orbit}/2^{12}} \epsilon_{o,R}^{\text{loc}}(s_{\delta_i}; x).$$

The output is the global character $\epsilon_{o,R} : W^{(2)}(\Lambda_{II}^{2,1}) \rightarrow \{\pm 1\}$.

- (v) *Compatibility package.* Verify the component, boundary, overlap, quotient, and protected integration compatibility package of Proposition ?? on every retained type-II wall stratum.
- (vi) *Pfaffian section.* Define $\text{pf}_{X,R}$ as the protected integration of the oriented Hall product over $\mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}}$:

$$\text{pf}_{X,R} = 64q^{1/2}r^{1/2}s^{1/2}I_R^{\text{prot}}\left(\bigotimes_{\gamma \in \mathcal{A}_R} \text{pf}_{\gamma,R}\right), \quad \text{pf}_{\gamma,R} = (1 - q^n r^l s^m)^{f(nm,l)}.$$

- (vii) Output $(D_R, \mathcal{L}_{\text{Pf},R}, \text{pf}_{X,R}, \epsilon_{o,R})$, with the protected integration compatibility data.

9.6.0.5 *Type checks.* $\mathcal{L}_{\text{Pf},R}$ is a Picard-line whose square isomorphism is supplied by reduced Serre duality; $\text{pf}_{X,R}$ is a section of $\mathcal{L}_{\text{Pf},R}$ with the displayed cusp expansion; the character $\epsilon_{o,R}$ takes values in $\{\pm 1\}$ and satisfies the Coxeter relations of $W^{(2)}(\Lambda_{II}^{2,1})$; on each simple type-II reflection $\epsilon_{o,R}(s_{\partial_i}) = -1$ by the rank-one local model.

9.6.0.6 *Obstructions discharged or invoked.* This step is the entire site of (O_2) (local Pfaffian wall normal form on type-II walls; the $2^{12} = 4096$ sign sum on the half-Hilbert orbit transport). The local rank-one Pfaffian crossing of Appendix 2 discharges (O_2) generically; the global half-Hilbert orbit transport remains the open (O_2) component of Open Problem ??. The character identity $\epsilon_o = \nu_{\Delta}|_{W^{(2)}(\Lambda_{II}^{2,1})}$ becomes a theorem only after (D_o) , (O_1) , $(O_1)^+$, and (O_2) are discharged compatibly in R .

9.7 Construction 7 — Chiral Koszul source coalgebra

9.7.0.1 *Tag.* γ -shadow specialization (*Bar = twisting; not bulk*; the bulk is $Z_{\text{ch}}^{\text{der}}(\mathcal{F}_X^{\text{hyb}})$, constructed elsewhere).

9.7.0.2 *Input.* The oriented Hall primitives $P_R^{\Pi, \text{geom}}$ of Construction 9.5; the hybrid carrier $\mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}}$ of Construction 9.3; the augmentation $\varepsilon_{X,R}^{\text{hyb}} : \mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}} \rightarrow k\mathbf{1}$; the hybrid residual $\mathfrak{D}_{\text{hyb},R} = 0$; the bounded length N_R (positivity of HN height on every non-vacuum colour); the Beilinson–Drinfeld/Francis–Gaitsgory bar-cobar formalism [?BD, ?FrancisGaitsgoryCKD]; the conilpotent completed domain on the elliptic curve E .

9.7.0.3 *Output.* The conilpotent chiral coalgebra on E :

$$C_{X,R}^{\text{geom}} = \bar{B}_E^{\text{ch}}(\mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}}, \varepsilon_{X,R}^{\text{hyb}}) = k\mathbf{1} \oplus \bigoplus_{1 \leq p \leq N_R} \bar{B}_E^{\text{ch},p}(\overline{\mathcal{F}}_{X,R}^{\text{geom}});$$

the bar-length filtration $F_{R,p}^{\text{co}}$; the chiral collision coproduct Δ_R^{ch} ; the bar comparison $b_{X,R} = \text{id}_{\bar{B}_E^{\text{ch}}(\cdot)}$; the source cobar counit $\varepsilon_{\text{bar},R}^{\text{src}} : \Omega_E^{\text{ch}} C_{X,R}^{\text{geom}} \xrightarrow{\cong} \mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}}$; after a separate source-to-target quasi-isomorphism $\mathfrak{I}_R : \mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}} \xrightarrow{\cong} \text{Den}_{\Delta,E,\leq R}$, the cobar comparison $\Theta_{\text{Kos},R} = \mathfrak{I}_R \circ \varepsilon_{\text{bar},R}^{\text{src}}$.

9.7.0.4 Procedure.

- (i) *Augmentation kernel.* Set $\overline{\mathcal{F}}_{X,R}^{\text{geom}} = \ker(\varepsilon_{X,R}^{\text{hyb}})$; verify the bounded-length hypothesis (positivity of HN height on every non-vacuum surviving colour, hence at most N_R factors in any non-zero word).

(ii) *Bar coalgebra.* Form the reduced chiral bar

$$C_{X,R}^{\text{geom}} = k\mathbf{1} \oplus \bigoplus_{1 \leq p \leq N_R} \tilde{B}_E^{\text{ch},p}(\overline{\mathcal{F}}_{X,R}^{\text{geom}}),$$

length p summand generated by ordered p -fold inputs from the augmentation ideal of total HN height $\leq R$ (height $> R$ outputs zero); coaugmentation by the inclusion of the vacuum summand; counit by projection.

(iii) *Bar-length filtration.*

$$F_{R,p}^{\text{co}} C_{X,R}^{\text{geom}} = k\mathbf{1} \oplus \bigoplus_{1 \leq j \leq p} \tilde{B}_E^{\text{ch},j}(\overline{\mathcal{F}}_{X,R}^{\text{geom}}), \quad 0 \leq p \leq N_R.$$

(iv) *Collision coproduct.* For $x = [x_1 | \cdots | x_p]$, define the cut-map coproduct

$$\Delta_R^{\text{ch}}(x) = \mathbf{1} \otimes x + x \otimes \mathbf{1} + \sum_{i=1}^{p-1} [x_1 | \cdots | x_i] \otimes [x_{i+1} | \cdots | x_p],$$

with each cut realized by the corresponding finite collision/flag pull-push map in $\text{Corr}_{R,\text{geom}}^{E,\text{hyb}}$ of Construction 9.3. Verify coassociativity by the four-input pentagon and the eight-word flag coherences; verify counit identities via vacuum collision flags.

(v) *Conilpotence.* Verify $(\overline{\Delta}_R^{\text{bar}})^{N_R+1} = 0$: each reduced cut strictly lowers bar length on each branch.

(vi) *Bar comparison.* Take $b_{X,R}$ to be the identity of $\tilde{B}_E^{\text{ch}}(\mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}}, \varepsilon_{X,R}^{\text{hyb}})$, eliminating the coalgebra-side finite Koszul residuals $\mathfrak{o}_R^C, \mathfrak{o}_R^{\text{fil}}, \mathfrak{o}_R^{\text{conil}}, \mathfrak{o}_R^{\Delta,\text{ch}}$ of Definition 0.25.

(vii) *Source cobar counit.* Apply Francis–Gaitsgory chiral Koszul duality [?FrancisGaitsgoryCKD, Theorem 6.2.2] in the conilpotent completed domain to obtain $\varepsilon_{\text{bar},R}^{\text{src}} : \Omega_E^{\text{ch}} C_{X,R}^{\text{geom}} \xrightarrow{\cong} \mathcal{F}_{X,\sigma,S,\leq R}^{\text{hyb}}$; verify it is a source-internal quasi-isomorphism of chiral algebras on E , not a homotopy inverse of the target counit.

(viii) *Source-to-target identification.* Construct $\mathfrak{I}_R$ from the geometric primitive representatives, Hall products, coproducts, pairings, and HN transition compatibility; verify the residual $\mathfrak{o}_R^{\mathfrak{I}}$ of Corollary 0.27 vanishes; set $\Theta_{\text{Kos},R} = \mathfrak{I}_R \circ \varepsilon_{\text{bar},R}^{\text{src}}$.

(ix) *Inverse limit.* Verify the Koszul Mittag–Leffler condition $R^1 \varprojlim = 0$ for the inverse systems of source coalgebras, completed chiral bar/cobar constructions, target truncations, and the cones of $b_{X,R}$ and $\Theta_{\text{Kos},R}$ [?WeibelHA, §3.5]. Form

$$C_X^{\text{geom}} = \varprojlim_R C_{X,R}^{\text{geom}}, \quad \Theta_{\text{Kos}} : \Omega_E^{\text{ch}} C_X^{\text{geom}} \xrightarrow{\cong} \text{Den}_{\Delta_S,E}.$$

(x) Output $(C_{X,R}^{\text{geom}}, F_{R,\bullet}^{\text{co}}, \Delta_R^{\text{ch}}, b_{X,R}, \Theta_{\text{Kos},R})$ at every R , with successor compatibility.

9.7.0.5 Type checks. $C_{X,R}^{\text{geom}}$ is a coaugmented conilpotent chiral coalgebra on E with bounded bar-length filtration of length at most N_R ; Δ_R^{ch} is coassociative and counital on the chosen flag stacks; the bar-cobar inversion is supplied by Francis–Gaitsgory chiral Koszul; the cobar comparison $\Theta_{\text{Kos},R}$ is source-internal, not the target bar-cobar counit (the source/target separation of Lemma 0.24).

9.7.0.6 Obstructions discharged or invoked. The Koszul Mittag–Leffler exactness hypothesis is the (Do)-component for the chiral Koszul lane. The total finite Koszul residual $\mathfrak{D}_{\text{Kos},R} = (\mathfrak{o}_R^C, \mathfrak{o}_R^{\text{fil}}, \mathfrak{o}_R^{\text{conil}}, \mathfrak{o}_R^{\Delta, \text{ch}}, \{\mathfrak{o}_{R'R}^{C, \text{tr}}\}, \mathfrak{o}_R^b, \mathfrak{o}_R^\Omega, \mathfrak{o}_R^{\text{hyb}}, \mathfrak{o}_R^\Pi, \mathfrak{o}_R^{\text{sep}}, \mathfrak{o}_R^{\text{Prim}})$ vanishes after this step.

9.7.0.7 Cross-volume firewall. The bar coalgebra $\bar{B}_E^{\text{ch}}(\mathcal{F}_{X,R}^{\text{hyb}})$ is a *twisting / coupling* coalgebra (Maurer–Cartan classification), *not* the bulk $Z_{\text{ch}}^{\text{der}}(\mathcal{F}_X^{\text{hyb}}) = \text{ChirHoch}^\bullet(\mathcal{F}_X^{\text{hyb}}, \mathcal{F}_X^{\text{hyb}})$; cf. `chiral-bar-cobar-vol2:CLAUDE.md §6` (universal stage chain $P \rightarrow C \rightarrow S \rightarrow Z$). The chiral Koszul comparison $\Theta_{\text{Kos}} : \Omega_E^{\text{ch}} C_X^{\text{geom}} \xrightarrow{\cong} \text{Den}_{\Delta, E}$ is a comparison between two algebras of the same level; the homotopy inverse to the target-internal counit $\Omega_E^{\text{ch}} \bar{B}_E^{\text{ch}}(\text{Den}_{\Delta, E}) \rightarrow \text{Den}_{\Delta, E}$ is not used in either direction.

9.8 Construction 8 — Primitive recognition

9.8.0.1 Tag. δ -endpoint convergence (source-to-target identification).

9.8.0.2 Input. The descended source primitive Lie superalgebra $P_X^{\Pi, \text{red}}$ of Construction 9.5 (after the radical quotient $\overline{\text{Rad}\langle \cdot, \cdot \rangle_X}$); the imported target denominator algebra $\mathfrak{g}_{\Delta_\delta} = G(\mathcal{B}_{\Delta_\delta})$ on $\Lambda_{II}^{2,1}$ of Definition ?? and Definition ??; the increasing cofinal sequence of finite downward-saturated windows $W_1 \subset W_2 \subset \dots$ with $\bigcup_\nu W_\nu = \mathcal{R}_+$; the Borchers–Kac–Moody data $(H, I, p, \alpha_i, (,))$ of Theorem 0.29.

9.8.0.3 Output. The cofinal finite-window primitive-recognition datum $\{(\text{Ro}), (\text{R1}), (\text{R2}), (\text{R3}), (\text{R4}), (\text{R5}), (\text{R6}), (\text{R7}), (\text{R8})\}$ of Definition 0.30 for every W_ν , compatible under $W_\nu \subset W_{\nu+1}$; the source representatives e_i^X, f_i^X, h_i^X ; the source presentation map $\pi_W : \widehat{\mathfrak{F}}(\mathcal{B}_{\Delta_\delta})_W \rightarrow P_X^{\Pi, \text{red}}|_W$; the kernel equality $\ker \pi_W = (\sqrt{J_{\text{BK}}} + \text{Rad}_{\text{GN}})_W$; the isomorphism

$$P_X^{\Pi, \text{red}} \cong \mathfrak{g}_{\Delta_\delta}, \quad \Omega_E^{\text{ch}} C_X \simeq \text{Den}_{\Delta, E}.$$

9.8.0.4 Procedure. For every cofinal W_ν :

- (Ro) *Compact source labels.* Verify every finite source basis vector in $W_\nu \cup (-W_\nu) \cup \{0\}$ satisfies the compact source-admissibility condition of Theorem ??.
- (R1) *Representatives and parities.* Supply parity-homogeneous primitive representatives $e_i^X \in P_{X, \alpha_i}^{\Pi, \text{red}}$, $f_i^X \in P_{X, -\alpha_i}^{\Pi, \text{red}}$, $h_i^X = \iota_H^{-1}(\alpha_i)$ for every simple-root index i with $\alpha_i \in W_\nu$, with the Gritsenko–Nikulin parity fibres $\mu_{\bar{0}}(a)$, $\mu_{\bar{1}}(a)$ for imaginary simple roots and even real representatives for $\delta_1, \delta_2, \delta_3$.
- (R2) *Relations.* Verify in the protected Hall super-bracket the Cartan, Chevalley, real Serre, Borchers orthogonality, and super sign relations for every relation whose displayed source and target degrees lie in $W_\nu \cup (-W_\nu) \cup \{0\}$. In the real rank-three subsystem the Cartan matrix is $((\delta_i, \delta_j)) = \text{diag}(2) - 2(J - \text{diag}(1))$ with real Serre exponent 3.
- (R3) *Full finite parity dimensions.* For every $\beta \in W_\nu$ verify $\dim P_{X, \beta, \bar{p}}^{\Pi, \text{red}} = \text{mult}_{\bar{p}}^{\text{GN}}(\beta)$ for both parities $p = 0, 1$. In the first timelike channel $\beta = \delta_{123}$, Proposition ?? requires $(d_{(1,1,1), \bar{0}}^{\text{GN}}, d_{(1,1,1), \bar{1}}^{\text{GN}}) = (29, 93)$. Determinant information $29 - 93 = -64 = f(1, 1)$ is consistent but not the recognition.

- (R4) *Hopf pairing and radical.* Compute the positive-negative pairing matrices in parity blocks; verify the kernel agrees with the Gritsenko–Nikulin/Kac invariant-pairing kernel; verify Lie-ideal and coideal stability under all finite brackets; verify carrying under HN transitions to the next window.
- (R5) *No extra relations.* Verify the kernel equality $\ker \pi_\nu = (\overline{J_{\text{BK}} + \text{Rad}_{\text{GN}}})_{\leq W_\nu}$ on the finite triangular span generated by $W_\nu \cup (-W_\nu) \cup \{0\}$.
- (R6) *Generation.* Verify that every source primitive root space in $W_\nu \cup (-W_\nu)$, after the finite radical quotient, is generated by the Cartan and the supplied simple-root representatives through brackets whose intermediate positive degrees stay in W_ν .
- (R7) *Completed PBW compatibility.* Verify the PBW-graded isomorphism $\text{gr}_{\text{PBW}} U(P_X^{\Pi, \text{red}})_{\leq W_\nu} \xrightarrow{\sim} \text{gr}_{\text{PBW}} U(\mathfrak{g}_{\Delta_5})_{\leq W_\nu}$; verify strict preservation under $W_{\nu+1} \rightarrow W_\nu$ transition maps.
- (R8) *Exact triangular completion.* Verify strictness of the transition maps and closedness of images on every finite window, so that passage to the inverse limit commutes with the finite kernels, cokernels, radicals, PBW associated gradeds, and parity pieces. Equivalently the relevant $\lim^1$ terms vanish for these triangular finite-window systems.
- (i) Output the global isomorphism $P_X^{\Pi, \text{red}} \cong \mathfrak{g}_{\Delta_5}$ by passage to the inverse limit (compatible cofinal datum); together with the chiral Koszul comparison of Construction 9.7 this gives $\Omega_E^{\text{ch}} C_X \simeq \text{Den}_{\Delta_5, E}$ and hence the protected Pfaffian identification $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}}) = \Delta_5$.

9.8.0.5 Type checks. The source presentation map $\pi : \widehat{\mathfrak{F}}(\mathcal{B}_{\Delta_5}) \rightarrow P_X^{\Pi, \text{red}}$ is continuous and degree-preserving; (R5) gives injectivity after the completed radical quotient; (R6) gives surjectivity; (R7) gives PBW compatibility; (R8) gives exactness in the inverse-limit step. No claim on signed superdimensions $\text{sdim} = f(nm, l)$ substitutes for the two-integer parity dimensions of (R3).

9.8.0.6 Obstructions discharged or invoked. This step inherits the obstruction tower discharged by Constructions 1–7 and adds the cofinal exact-completion residual (R8). The Beilinson–Drinfeld bar-cobar firewall is respected throughout: (R5) is a source kernel-equality theorem, not an inversion of the target bar-cobar counit; (R7) is a PBW graded comparison of source and target enveloping algebras, not a graph-equality argument from the scalar square.

9.8.0.7 Cross-volume anchor. The output is `chiral-bar-cobar-vol2:CLAUDE.md` §9 Construction Problem 1: the operator-level construction of $\mathfrak{D}_X^{\text{DI}}$ for $K_3 \times E$ with $\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}}) = \Delta_5$, under the four obstructions (D_0) , (O_1) , $(O_1)^+$, (O_2) . The Drinfeld doubling $Y^+(X) \rightarrow G(X) = D(Y^+(X))$ of `calabi-yau-quantum-groups:CLAUDE.md` requires the Hopf pairing of (R4), completion by (R8), integral form, and stable-envelope transport which are supplied by the same data. The κ -tuple coordinate $\kappa_{\text{BKM}} = 5$ is the weight of Δ_5 and lives at level 4 of the universal stage chain; it is a consequence of the construction, not an input to it.

9.9 The eight-step audit

The eight construction steps form a strict ordered sequence of input/output type discipline. An audit of any candidate construction of $\mathfrak{D}_X^{\text{DI}}$ reduces to verifying, for every $R \in \mathcal{R}_{\text{HN}}$:

- (i) the lattice combinatorics of Construction 9.1;
- (ii) the finite-type quasi-smooth derived stacks of Construction 9.2;

- (iii) the hybrid local/wrapped factorization datum and the eight-word two-step flag atlas of Construction 9.3;
- (iv) the four Čech–Borel orientation classes and the $(O_1)^+$ Coxeter coherence of Construction 9.4;
- (v) the finite Hall bialgebra and primitive matrices of Construction 9.5;
- (vi) the rank-one Pfaffian crossing and the half-Hilbert orbit transport of Construction 9.6;
- (vii) the bar coalgebra, conilpotent collision coproduct, and the Francis–Gaitsgory cobar quasi-isomorphism of Construction 9.7;
- (viii) the cofinal datum (Ro)–(R8) of Construction 9.8;

together with the compatible vanishing of the obstruction tuples $(\mathfrak{D}_R^{E_3-\text{src}}, \mathfrak{D}_{\text{hyb}, R}, \mathfrak{D}_{\text{Kos}, R}, \mathfrak{D}_R^{O_1}, \mathfrak{D}_R^{O_1^+}, \mathfrak{D}_R^{O_2}, \mathfrak{D}_R^{D_o})$ at every height R , and the Mittag–Leffler exactness condition [ML] of Definition ?? on the cofinal HN inverse system. Each obstruction is attributable to the unique step in which its components are constructed.

9.9.0.1 Order rigidity. No two adjacent construction steps may be exchanged: the charge window feeds the finite moduli, which feeds the hybrid carrier, which feeds the orientation, which feeds the Hall correspondences, which feed the Pfaffian line and Dirac operator, which feed the source coalgebra (the bar of the augmented hybrid source), which feed primitive recognition. Type-checking forbids permutation; for instance, computing the orientation gerbe before the hybrid wrapped prequotient leaves the middle type-II wall $\delta_2 \leftrightarrow (o, 1, 1)$ without a representative, by Lemma ??; computing primitive recognition before the chiral Koszul cobar leaves the second identification $\Omega_E^{\text{ch}} C_X \simeq \text{Den}_{\Delta_5, E}$ unconstructed.

9.9.0.2 Conditional theorem. Granted the eight constructions and the four obstructions $(D_o), (O_1), (O_1)^+, (O_2)$, Theorem ?? reads

$$\text{Pf}_{\text{prot}}(\mathfrak{D}_X^{\text{DI}}) = \Delta_5, \quad \epsilon_o = \nu_{\Delta_5} \big|_{W^{(2)}(\Lambda_{II}^{2,1})}, \quad P_X^{\Pi, \text{red}} \cong \mathfrak{g}_{\Delta_5}.$$

The scalar square $Z_{\text{BPS}}^{K_3 \times E} = (\Phi_{10}^{\text{un}})^{-1} = \Delta_5^{-2}$ is the level-4 trace of the constructed protected algebra; promotion to the 3d gravitational path integral is not claimed and requires the Hall–Borcherds residual of chiral-bar-cobar-vol2.