

*Local Mixed Holomorphic-Topological String Field Theory*

# REMARKS ON MIXED HOLOMORPHIC-TOPOLOGICAL STRINGS

*Formal Local Hamiltonian BF Theory,  
Reduced CE/PV Central Operations,  
and the Trace Chart at  $N$  Dirac Branes on  $\mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$*

*the formal-Darboux stalk of the Mixed Holomorphic-Topological Deligne conjecture*

Raez Lorgat

May 2026



## Abstract

The Bershadsky–Cecotti–Ooguri–Vafa B-model topological string [1] is the topological B-twist of the  $\mathcal{N} = (2, 2)$  worldsheet sigma model on a Calabi–Yau target; its Costello–Li BV renormalisation [12–14] admits a further holomorphic-topological twist localising it on  $X = \mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$  in the Costello twisted-supergravity sense [25]. In this twisted local geometry, topological in  $\mathbb{R}^2$  and holomorphic in  $\mathbb{C}^2$ , the closed-string sector is the holomorphic-topological Hamiltonian BF theory whose fields are  $\mathcal{A} \in \Omega^\bullet(\mathbb{R}^2) \widehat{\otimes} \Omega^{0,\bullet}(\mathbb{C}^2) \widehat{\otimes} \mathfrak{h}$  valued in the reduced Hamiltonian Lie algebra  $\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$  of the formal symplectic polydisk, antifields  $\mathcal{B}$  valued in the Grothendieck–Matlis residue dual  $\mathfrak{h}_{\text{cont}}^\vee = H_{(z_1, z_2)}^2(\mathbb{C}[[z_1, z_2]]) dz_1 dz_2$  paired by  $\langle f, \omega \rangle = \mathbf{Res}_0 f \omega$ , and master action

$$S = \int_X \langle \mathcal{B}, (d + \bar{\partial})\mathcal{A} + \tfrac{1}{2}[\mathcal{A}, \mathcal{A}]_{\mathfrak{h}} \rangle.$$

The open sector is a stack of  $N$  Dirac branes on the worldline  $\mathbb{R}_t \subset X$  with matrix transverse position  $(\phi_1, \phi_2) \in \mathfrak{gl}_N \oplus \mathfrak{gl}_N$ , Chan–Paton substitution  $z_i \mapsto \phi_i$  subject to the moment-map constraint  $[\phi_1, \phi_2] = 0$ , and antifield  $\psi \in \mathfrak{gl}_N[1]$  with  $Q\psi = [\phi_1, \phi_2]$  giving the Koszul resolution of the derived zero fibre.

At a brane vacuum  $b \in \mathbb{C}^2$  the formal Darboux chart is  $\mathcal{A}_b = \widehat{\mathcal{O}}_{\mathbb{C}^2, b} \cong \mathbb{C}[[z_1, z_2]]$  with symplectic form  $\omega_b = dz_1 \wedge dz_2$ ; the classical open-string algebra of observables on  $\mathbb{R}_t$  is the Chevalley–Eilenberg cochain complex of the Chan–Paton gauge algebra with values in the symmetric algebra of the matrix coordinates, the antifield, and the moment-map antifield,

$$\mathcal{A}_{\partial, N}^{\text{cl}} = C_{\text{CE}}^\bullet(\mathfrak{gl}_N, \text{Sym}(\mathfrak{gl}_N \oplus \mathfrak{gl}_N \oplus \mathfrak{gl}_N[1])),$$

with Koszul differential  $Q\psi = [\phi_1, \phi_2]$ . A closed Hamiltonian  $f \in \mathfrak{h}$  acts on this brane sector through the trace map

$$J: \mathfrak{h} \longrightarrow \mathcal{O}(\mu_{\text{der}}^{-1}(0))^{GL_N}, \quad J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)},$$

the unique  $GL_N$ -equivariant, primitive, single-trace, Darboux-natural expression of  $f$  on the derived commuting variety, normalised by  $J(z_i) = \overline{\text{Tr } \phi_i}$ .

The closed observables form the mixed-HT factorization algebra  $\mathcal{F}_{\text{Ham}}^{\text{cl}}$  valued in CE chains of the shifted-cotangent BF Lie algebra  $\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1]$ ; its pushforward along  $\mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2 \rightarrow \mathbb{C}_{\text{hol}}^2$  is a holomorphic factorization algebra  $\mathcal{A}_{\text{bulk}}^{\text{cl}}$  on  $\mathbb{C}^2$  valued in  $E_2$ -algebras. It is the two-complex-dimensional holomorphic factorization algebra of Hamiltonian currents. Its singular product is the two-variable diagonal expansion on  $\Delta \subset \mathbb{C}_z^2 \times \mathbb{C}_w^2$ : for  $x \in \mathfrak{g}$  the linear holomorphic current  $J_x(z)$  in the CE factorization algebra  $B \mapsto C_{\text{CE, cont}}^\bullet(\Omega_c^{0,\bullet}(B) \widehat{\otimes} \mathfrak{g})$  satisfies

$$J_x(z)J_y(w) \sim K_\Delta^{(2)}(z, w) J_{[x, y]_{\mathfrak{g}}}(w), \quad K_\Delta^{(2)}(z, w) = \frac{d\zeta_1 d\zeta_2}{\zeta_1 \zeta_2}, \quad \zeta_i = z_i - w_i,$$

with the diagonal local-cohomology principal-part module  $H_\Delta^2(\mathbb{C}[[z_1, z_2, w_1, w_2]]) d\zeta_1 d\zeta_2$  as its singular coefficient module. A curve chiral algebra is obtained from this surface factorization algebra after choosing a curve  $C \hookrightarrow \mathbb{C}^2$ , a pushed-forward bracket, the induced BV pairing, the image of the brane, and the anomaly normalisation; under these data the diagonal principal part restricts to the Cauchy kernel  $K_C^{(1)}(z, w) = dz/(z - w)$ .

The Mixed Holomorphic–Topological Deligne conjecture identifies  $\mathcal{A}_{\text{bulk}}^{\text{cl}}$  with the mixed-HT chiral derived  $E_2$ -centre of the boundary open factorisation category  $C_\partial^{\text{op}}$  on the Kato–Nakayama log boundary  $\partial X^{\text{KN}}$  — the two-complex-dimensional extension of the chiral Deligne theorem for chiral algebras on curves, in the Beilinson–Drinfeld formulation [28, §4]. Inside the admissible

filtered nuclear Tate / pro-Matlis / Capelli-projective derived-centre category  $\mathcal{B}^{\text{adm}}$ , the formal-Darboux stalk identification at  $N$  Dirac branes is unconditional:

$$\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b \simeq C_{\text{CE,cont}}^{\bullet}(\widehat{\mathfrak{g}}_{\text{cent}}) \simeq Z_{E_i}^{\text{der,adm}}(A_{\partial,N}^{\text{cl}}) \simeq HH_{\text{adm}}^{\bullet}(A_{\partial,N}^{\text{cl}}, A_{\partial,N}^{\text{cl}})$$

as  $E_2$ -algebras — the bulk restricted to the formal neighbourhood of  $b$ , equated with the brane-side derived  $E_1$ -centre (equivalently the  $E_1$ -Hochschild cochain complex) by Costello–Gwilliam factorization-algebra bulk–boundary correspondence and the  $E_1$ -Hochschild theorem. The Capelli central extension  $\widehat{\mathfrak{g}}_{\text{cent}}$  absorbs the finite- $N$  scalar anomaly as the projective central charge  $\hbar N[\bar{c}]$ ; the trace map  $J(f) = \text{Tr } f(\phi_1, \phi_2)$  is the formal Darboux coordinate expression of the identification; the closed-to-open dictionary is the Koszul resolution carrying the centrality homotopy hierarchy,  $(c_f, u_f, K) \mapsto (\theta_f, J(f) \cdot (-), \text{Tr } \mathbf{1}, H^{\text{prod}}, H^{P_0}, H^3, \dots)$ , with  $\theta_f$  the Hamiltonian Schouten derivation. The scalar-reduced shadow  $K \mapsto \mathfrak{o}, \hbar N[\bar{c}] \mapsto \mathfrak{o}$  recovers the reduced equivalence with  $J(f) \rightsquigarrow \text{Tr } f(\phi_1, \phi_2)$ .

The constant Hamiltonian line  $\mathbb{C} \cdot \mathbf{1} \subset \mathbb{C}[z_1, z_2]$  lies in the kernel of the closed Hamiltonian Lie algebra reduction but acts on the brane as  $c \mapsto c \cdot I$ , producing the finite- $N$  central term. With  $\Phi_{\hbar}: \mathbb{C}[z_1, z_2] \rightarrow M_N(\mathbb{C})$  the Weyl-symmetric quantisation map and  $\star$  the Moyal star-product, this term has three equivalent expressions:

$$\omega(f, g) = [\{f, g\}]_{\mathfrak{o}}, \quad \{\text{Tr } \phi_1, \text{Tr } \phi_2\} = N, \quad [\Phi_{\hbar}(z_1), \Phi_{\hbar}(z_2)]_{\star} = \hbar N \cdot I,$$

the Lie 2-cocycle on  $\bar{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot \mathbf{1}$ , the Poisson bracket of trace coordinates, and the Moyal commutator in  $M_N(\mathbb{C})$ . They realise one class  $\hbar N[\bar{c}] \in H_{\text{Lie}}^2(\bar{A}, \mathbb{C})[[\hbar]]$  — the Capelli scalar of the trace centre-of-mass and the projective curvature  $\Omega_{\text{central}}|_{J(f)}$  of the modular line bundle on the period domain at the trace generator,

$$\hbar N[\bar{c}] = \Omega_{\text{central}}|_{J(f)}.$$

The local model is the two-complex-dimensional analogue of the Heisenberg vertex algebra, with  $\mathcal{P}$  playing the role of the Kac–Moody central pairing module  $\Omega_K$  for a curve  $K$  and  $\mathfrak{h}$  that of regular functions; it is the formal local sector of the Costello twisted-M-theory family [25]; and on its cyclic framed stable branch  $\mathbb{C}[\phi_1, \phi_2] \cdot v = \mathbb{C}^N$  it is the formal-Darboux stalk chart of  $\text{Hilb}^N(\mathbb{C}^2)$  in its Nekrasov–Schwarz noncommutative Moyal deformation [46].

The formal-Darboux stalk theorem supplies the local term in a global identification. The global identification on the Kato–Nakayama log boundary holds if and only if the obstruction  $\text{Ob}^{\text{global}} = \text{Ob}_{\text{Deligne}}^{\text{ch } E_2} \oplus \text{Ob}_{\text{KN}}^{\text{desc}} \oplus \text{Ob}_{\text{all-loop}}^{\text{QME}}$  vanishes; the chiral  $E_d$ -Deliège conjecture at  $d \geq 2$ , the operator-level lift of the modular line on  $K_3 \times E$ , and the Hall–Drinfeld–Borchers comparison on compact non-toric Calabi–Yau threefolds are the comparison problems carried by these classes.

## SUMMARY

The B-model topological string of Bershadsky–Cecotti–Ooguri–Vafa [1] is the topological B-twist of the  $\mathcal{N} = (2, 2)$  worldsheet sigma model on a Calabi–Yau target; its Costello–Li BV renormalisation [12–14] admits a further holomorphic-topological twist localising it on  $X = \mathbb{R}_{t,s}^2 \times \widehat{\mathbb{C}^2}_{\text{o,hol}}$  in the Costello twisted-supergravity sense [25] — topological in the worldvolume coordinates  $(t, s)$  and holomorphic in the transverse coordinates  $(z_1, z_2)$  — with the closed-string sector becoming the holomorphic-topological Hamiltonian BF theory below. The closed-string fields  $\mathcal{A}$  take values in the reduced Hamiltonian Lie algebra of the formal symplectic polydisk,  $\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot \mathbf{1}$  with Poisson bracket  $\{f, g\} = \partial_{z_1} f \partial_{z_2} g - \partial_{z_2} f \partial_{z_1} g$ ; the antifields  $\mathcal{B}$  take values in the continuous Grothendieck–Matlis residue

dual  $\mathfrak{h}_{\text{cont}}^\vee = H_{(z_1, z_2)}^2(\mathbb{C}[[z_1, z_2]]) dz_1 dz_2$  paired by  $\langle f, \omega \rangle = \mathbf{Res}_o f \omega$ . Together they form the shifted-cotangent BF Lie algebra  $\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1]$  on the formal symplectic polydisk  $\widehat{\mathbb{C}^2}_o$ , with classical action

$$S = \int_X \langle \mathcal{B}, (d + \bar{\partial})\mathcal{A} + \tfrac{1}{2}[\mathcal{A}, \mathcal{A}]_{\mathfrak{h}} \rangle.$$

The classical observables of this theory form a holomorphic factorization algebra on  $\mathbb{C}^2$  valued in  $E_2$ -algebras,  $\mathcal{A}_{\text{bulk}}^{\text{cl}}$ . It factorises over the holomorphic surface in the sense of Costello–Gwilliam [15].

An open boundary is supplied by a stack of  $N$  Dirac branes on the worldline  $\mathbb{R}_t \subset X$ . Its two holomorphic transverse coordinates become Chan–Paton matrices  $\phi_1, \phi_2 \in \mathfrak{gl}_N$  under the substitution  $z_i \mapsto \phi_i$ , subject to the moment-map constraint  $[\phi_1, \phi_2] = o$ . The antifield  $\psi \in \mathfrak{gl}_N[1]$  with  $Q\psi = [\phi_1, \phi_2]$  gives the Koszul resolution of the moment-map ideal, equivalently the derived zero fibre of  $\mu(\phi_1, \phi_2) = [\phi_1, \phi_2]$ . The classical open-string algebra on the worldline is the Chevalley–Eilenberg cochain complex

$$A_{\partial, N}^{\text{cl}} = C_{\text{CE}}^\bullet(\mathfrak{gl}_N, \text{Sym}(\mathfrak{gl}_N \oplus \mathfrak{gl}_N \oplus \mathfrak{gl}_N[1])),$$

with Koszul differential  $Q\psi = [\phi_1, \phi_2]$ . A closed Hamiltonian  $f \in \mathfrak{h}$  acts on this brane sector through the trace map

$$J: \mathfrak{h} \longrightarrow \mathcal{O}(\mu_{\text{der}}^{-1}(o))^{GL_N}, \quad J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)},$$

the  $GL_N$ -invariant function obtained by substituting the matrices  $\phi_i$  into  $f$  and taking the trace on the derived commuting variety  $\mu_{\text{der}}^{-1}(o)$  at the brane vacuum  $b \in \mathbb{C}^2$ .

The Mixed Holomorphic–Topological Deligne conjecture identifies  $\mathcal{A}_{\text{bulk}}^{\text{cl}}$  with the chiral derived  $E_2$ -centre of the boundary open factorisation category  $\mathcal{C}_\partial^{\text{op}}$  on the Kato–Nakayama log boundary  $\partial X^{\text{KN}}$  — the two-complex-dimensional extension of the  $d = 1$  chiral Deligne theorem for chiral algebras on algebraic curves [28, §4]. This manuscript proves the formal-Darboux stalk identification of that conjecture: the equivalence

$$\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b \simeq Z_{E_1}^{\text{der}}(A_{\partial, N}^{\text{cl}}) \simeq HH^\bullet(A_{\partial, N}^{\text{cl}}, A_{\partial, N}^{\text{cl}})$$

as  $E_2$ -algebras — the bulk restricted to the formal neighbourhood of  $b$ , identified with the brane-side derived  $E_1$ -centre (equivalently the  $E_1$ -Hochschild cochain complex) by Costello–Gwilliam factorization-algebra bulk–boundary correspondence and Deligne’s  $E_1$ -Hochschild theorem. The trace map  $J$  is the formal Darboux coordinate expression of this equivalence. Theorem 5.6 is a local formal-Darboux statement. Away from the stalk, the corresponding global assertion is the open chiral  $E_d$ -Deligne conjecture at  $d = 2$ .

The classical BV pair is the field  $\mathcal{A}$  valued in  $\Omega^\bullet(\mathbb{R}^2) \widehat{\otimes} \Omega^{0, \bullet}(\mathbb{C}^2) \widehat{\otimes} \mathfrak{h}$  and antifield  $\mathcal{B}$  valued in  $\Omega^\bullet(\mathbb{R}^2) \widehat{\otimes} \Omega^{0, \bullet}(\mathbb{C}^2) \widehat{\otimes} \mathfrak{h}_{\text{cont}}^\vee[1]$ , paired by the  $(-1)$ -shifted symplectic form  $\int_X \langle \mathcal{B}, \mathcal{A} \rangle$  obtained from the residue pairing on  $\mathfrak{h} \otimes \mathfrak{h}_{\text{cont}}^\vee$  and integration of mixed-HT forms against  $dz_1 dz_2$ . The BV differential is  $Q = d + \bar{\partial} + d_{\text{CE}}$ , with  $d_{\text{CE}}$  the Chevalley–Eilenberg differential of  $\mathfrak{g}$ ; the master equation  $\{S, S\} = 0$  holds because the BF interaction  $I = \tfrac{1}{2} \int_X \langle \mathcal{B}, [\mathcal{A}, \mathcal{A}]_{\mathfrak{h}} \rangle$  encodes the Lie bracket on  $\mathfrak{h}$ . Renormalisation proceeds in the heat-kernel scheme of Costello–Gwilliam: a gauge-fixing operator  $Q^{\text{GF}} = d^* + \bar{\partial}^*$  for chosen flat and Hermitian metrics gives the Laplacian  $D = [Q, Q^{\text{GF}}]$ , heat kernel  $K_t = e^{-tD}$ , and propagator  $P_{\varepsilon, L} = \int_\varepsilon^L \tfrac{1}{2}(Q^{\text{GF}} \otimes 1 + 1 \otimes Q^{\text{GF}}) K_t dt$ . At RG scale  $L$  the effective interaction is the Costello transform  $I[L] = W(P_{\varepsilon, L}, I)$ , and the quantum master equation

$$QI[L] + \tfrac{1}{2}\{I[L], I[L]\}_L + \hbar \Delta_L I[L] = 0$$

is the consistency condition for renormalised quantisation. Costello deformation cohomology of  $I[L]$  is the obstruction theory governing the closed-to-open coupling  $\Phi_N$  (Theorem 1.24); its reduced non-scalar obstruction is the class  $[\text{Ob}_{w,\partial,\hbar}^{\text{red}}]$  (Theorem E.38).

*Tate setup and the residue pairing.* Throughout,  $\mathfrak{h}$  carries the linear topology defined by the images of  $\mathfrak{m}^k \subset A$ , with continuous dual  $\mathfrak{h}_{\text{cont}}^\vee = \text{Hom}_{\mathbb{C}}^{\text{cts}}(\mathfrak{h}, \mathbb{C})$  in the category of Tate  $\mathbb{C}$ -vector spaces (linearly compact / pro-finite). The residue pairing identifies  $\mathfrak{h}_{\text{cont}}^\vee$  with the Matlis principal-part module

$$\mathcal{P} = H_{\mathfrak{m}}^2(A) \otimes \omega_A = \bigoplus_{a+b>0} \mathbb{C} \rho_{a,b}, \quad \omega_A = dz_1 \wedge dz_2, \quad \rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2,$$

the holomorphic-surface analogue of the Heisenberg current on the formal disk. Inverse limits over Matlis windows  $\mathcal{P}_{\leq N} = \bigoplus_{a+b \leq N} \mathbb{C} \rho_{a,b}$  are read as derived limits  $\mathbf{R}\varprojlim_N \mathcal{P}_{\leq N}$ , with  $\varprojlim^1$ -data tracked.

The moment-map equation  $[\phi_1, \phi_2] = 0$  defines a singular variety on  $\mathfrak{gl}_N \oplus \mathfrak{gl}_N$ : for  $N \geq 2$  the entries of  $[\phi_1, \phi_2]$  have syzygies and define the derived commuting-pair stack. The Koszul complex  $\mathbb{C}[\phi_1, \phi_2, \psi]$  with  $Q\psi = [\phi_1, \phi_2]$  produces a derived enhancement: the ghost-zero brane moduli is the derived commuting-pair stack  $[\mu_{\text{der}}^{-1}(0)/GL_N]$ . The classical Hilbert scheme  $\mathcal{H}\text{ilb}^N(\mathbb{C}^2)$  is recovered on the cyclic framed stable branch obtained by adjoining a framing vector  $v \in \mathbb{C}^N$  with  $\mathbb{C}[\phi_1, \phi_2] \cdot v = \mathbb{C}^N$ ; the Moyal–Weyl deformation on that branch produces the Nekrasov–Schwarz noncommutative Hilbert scheme [46].

Uniqueness of  $J$ . Procesi–Razmyslov [17, 18] identifies the polynomial  $GL_N$ -invariants on  $\mathfrak{gl}_N \oplus \mathfrak{gl}_N$  with the algebra generated by mixed traces of words in  $\phi_1, \phi_2$ ; Loday–Quillen–Tsygan [2, 3] identify the primitive part with the cyclic-word trace classes. Combined with Dirac descent through the moment-map ideal, formal-Darboux naturality, and the linear normalisation  $J(z_i) = \overline{\text{Tr } \phi_i}$ ,  $J$  is forced monomial-by-monomial (Theorem 5.41); among admissible protected Hamiltonian sectors at a holomorphic-symplectic brane point,  $J$  is the terminal datum (Theorem 5.146).

Concretely the closed observables are the Chevalley–Eilenberg chains of the local Lie algebra  $\Omega_c^\bullet(U) \widehat{\otimes} \Omega_c^{\circ,\bullet}(B) \widehat{\otimes} \mathfrak{g}$  on  $U \subset \mathbb{R}_{\text{top}}^2$ ,  $B \subset \mathbb{C}_{\text{hol}}^2$  (Definition 1.10, Proposition 1.34); pushforward along  $\pi: X \rightarrow \mathbb{C}_{\text{hol}}^2$  integrates out the topological factor and gives  $\mathcal{A}_{\text{bulk}}^{\text{cl}}(B) = \mathcal{F}_{\text{Ham}}^{\text{cl}}(D^2 \times B) \in \text{Alg}_{E_2}$  (Theorem 1.20, Corollary 1.21); the  $E_2$ -valuation arises from the locally constant factorization-algebra structure on  $\mathbb{R}_{\text{top}}^2$  (Lurie–Costello–Gwilliam). The native holomorphic singular product of  $\mathcal{A}_{\text{bulk}}^{\text{cl}}$  is a two-variable expansion on the diagonal  $\Delta \subset \mathbb{C}_z^2 \times \mathbb{C}_w^2$ : the singular coefficient module is the Grothendieck local-cohomology principal-part module  $H_\Delta^2(\mathbb{C}[[z_1, z_2, w_1, w_2]]) d\zeta_1 d\zeta_2$ , with arity-two Cauchy kernel  $K_\Delta^{(2)}(z, w) = d\zeta_1 d\zeta_2 / (\zeta_1 \zeta_2)$  in formal diagonal coordinates  $\zeta_i = z_i - w_i$  (Theorem 1.15). A one-dimensional chiral algebra is obtained after constructing a restriction or pushforward to a curve  $C \hookrightarrow \mathbb{C}^2$ . Such a construction must specify the transverse modes, the pushed-forward bracket, the BV pairing, the image of the brane chart, and the anomaly normalisation. Heisenberg, affine Kac–Moody,  $\beta\gamma$ , Virasoro, and  $\mathcal{W}$ -algebra vertex algebras enter as curve-chiral comparison targets with these extra hypotheses.

The closed-to-open dictionary on Chevalley–Eilenberg generators of  $\mathfrak{g}$  assigns to each generator-dual coordinate  $c_f$  the Hamiltonian Schouten derivation  $\theta_f \in HH^1(A_{\partial,N}^{\text{cl}})$  and to each cotangent-dual coordinate  $u_f$  the brane observable  $O_f = J(f) \in HH^0(A_{\partial,N}^{\text{cl}})$  (Theorem 1.24). Lifted to factorisation algebras, this is the chain map

$$\Phi_N: \mathcal{A}_{\text{bulk},0}^{\text{cl}} \longrightarrow Z_{E_1}^{\text{der}}(A_{\partial,N}^{\text{cl}})$$

whose existence at the chain level is conditional on vanishing of the obstruction vector  $\text{Ob}_{\text{cent/QME/desc}}$  (Theorem E.53) with null-homotopies supplied by the completion constructions of Appendix G.

*Capelli scalar anomaly.* Closed Hamiltonian reduction quotients the constant Hamiltonian line  $\mathbb{C} \cdot 1$ , on which  $\text{Tr } 1 = N$ ; the brane retains it as the trace- $U(1)$  centre-of-mass cocycle of  $N$  coincident D-branes on noncommutative  $\mathbb{C}_h^2$ . The class  $\hbar N[\tilde{c}] \in H_{\text{Lie}}^2(\tilde{A}, \mathbb{C})[[\hbar]]$ ,  $\tilde{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1$ , has three faces: the Lie 2-cocycle  $\omega(f, g) = [f, g]_0$  on  $\tilde{A}$  (Lemma 6.2), the Poisson bracket of trace coordinates  $\{\text{Tr } \phi_1, \text{Tr } \phi_2\} = N$ , and the Moyal commutator (with  $\Phi_h: \mathbb{C}[z_1, z_2] \rightarrow M_N(\mathbb{C})$  the Weyl-symmetric quantisation map of §4.2)  $[\Phi_h(z_1), \Phi_h(z_2)]_\star = \hbar N \cdot I$  in  $M_N(\mathbb{C})$  (Theorem E.13). The number  $N$  is the trace- $U(1)$  central charge of the noncommutative D-brane stack on  $\mathbb{C}_h^2$ , equivalently the projective curvature  $\Omega_{\text{central}}|_{J(f)}$  of  $\Phi_N$ . The strict scalar-reduced Hamiltonian action upgrades to a projective action on the Capelli central extension  $0 \rightarrow \mathbb{C} \rightarrow \mathfrak{h}_{\hbar, N} \rightarrow \tilde{A} \rightarrow 0$ .

*Obstruction calculus.* Write  $\Theta_{\text{loc}}$  for the local trace-substitution datum  $(J, \rho_\phi, \psi, c_f \mapsto \theta_f, u_f \mapsto J(f))$  packaged as a degree-one element of the filtered  $L_\infty$ -algebra  $\mathfrak{R}_{\text{loc}}$  of §7.1. Globalization, quantization, all-bidegree Weyl/Moyal compatibility, and chain-level upgrades are filtered  $L_\infty$  transports of  $\Theta_{\text{loc}}$  into a target deformation complex  $\mathfrak{R}_{\mathcal{T}}$ , with master equation curvature  $F_{\mathcal{T}}(\Theta_{\text{loc}} + \tau) = 0$  (Theorem 7.1). All-bidegree Weyl/trace compatibility is controlled by the radial Weyl cokernel  $\Omega_{a,b}^{\text{rad}} \in \text{coker } B_{a,b}$  on the decorated Poincaré–Birkhoff–Witt (PBW) Stokes operator  $B_{a,b}$  (Theorems 5.217, F.44), with  $\Omega_{a,b}^{\text{rad}} = 0$  for all  $(a, b)$  conditional on the Harish-Chandra/Levasseur–Stafford radial-image input Statement F.8; the chain-level  $P_0$ -factorization-centre lift of  $\Phi_N$  by the factorization-centre vector  $\text{Ob}_{\text{cent/QME/desc}}$  (Theorem E.53); the reduced non-scalar Costello QME by the class  $[\text{Ob}_{w,\partial,\hbar}^{\text{red}}] \in H^1(\mathfrak{Q}_{w,\partial,\hbar}^\bullet(I), d_{\text{QME}})$  in the filtered weighted brane-defect local-functional complex, whose order-three CE-source row  $\theta_3$  gives the prototype  $H_{\text{bare}}^1 = \mathbb{C} \cdot E_{\theta_3, M}$  killed by the unique counterterm  $d_M C_{\theta_3, M} = -E_{\theta_3, M}$ ,  $c = 1$ , in any finite window  $M \ni \theta_3$  (Theorem E.38); and the strict all-window Bochner–Martinelli current transfer by  $\text{Ob}_{\text{BM}}^\Pi$ , nonzero on every finite Matlis window  $\mathcal{P}_{\leq N}$  (Theorem 8.40), with coherent action on the pro-Matlis substituted target  $\hat{\mathcal{P}}_\Pi = \varprojlim_N \mathcal{P}_{\leq N}$  (Theorem 8.42). These four classes are the obstructions identified for the upgrade directions considered here; they sit in the filtered  $L_\infty$  algebra  $\mathfrak{R}_{\text{loc}}$  governing deformations of  $\Phi_N$ . Minimal-model classification and an exact count of nonzero higher brackets are separate theorems.

The local model is the holomorphic factorization algebra of Hamiltonian currents on  $\mathbb{C}^2$ . Its coefficient dual is the Matlis principal-part module  $\mathcal{P}$ . In the curve analogy,  $\mathcal{P}$  replaces the Kac–Moody central-pairing module  $\Omega_K$ , and  $\mathfrak{h}$  replaces the regular functions on the curve. The same formal local sector appears in Costello’s twisted M-theory family [25]. On the cyclic framed stable branch it is the formal-Darboux chart of  $\mathcal{H}\text{ilb}^N(\mathbb{C}^2)$  in its Moyal deformation.

Across volumes, the bulk algebra  $\mathcal{A}_{\text{bulk}}^{\text{cl}}$  is sourced from a Calabi–Yau category  $\mathcal{C}$  through the two-stage functor  $\Phi_d = \text{Sp}_{\Sigma_{d-1}, \mathcal{C}}^{\text{ch}} \circ \Phi_d^{\text{FA}}$  — the Calabi–Yau-to-factorization-algebra functor  $\Phi_d^{\text{FA}}$  followed by Lurie–Ayala–Francis factorization-homology specialisation  $\text{Sp}_{\Sigma_{d-1}, \mathcal{C}}^{\text{ch}}(\mathcal{F}) = \int_{\Sigma_{d-1}} \mathcal{F}|_{\mathcal{C}}$ ; at  $d = 2$ ,  $\mathcal{C}$  is the Calabi–Yau category  $\mathcal{C}$  of the holomorphic-symplectic target. Conventions match a global

theorem through the matched-conventions acceptance class  $\text{Ob}_{\text{UKD}}$  of Theorem 10.13: a nonzero coordinate of  $\text{Ob}_{\text{UKD}}$  is the obstruction to transporting the local theorem to a compact-target programme (BCOV/Kodaira–Spencer, CY-to-chiral, Igusa/BKM, GW/DT/PT/MNOP, OSV).

## CONTENTS

|                                                                                                   |     |
|---------------------------------------------------------------------------------------------------|-----|
| Summary . . . . .                                                                                 | 2   |
| CONTENTS . . . . .                                                                                | 6   |
| Local dictionary . . . . .                                                                        | 9   |
| Conventions and notation for the local theorem . . . . .                                          | 28  |
| 1 The setup . . . . .                                                                             | 31  |
| 1.1 The Dirac brane formal-stalk chart in one calculation . . . . .                               | 31  |
| 1.2 Mixed holomorphic-topological strings . . . . .                                               | 33  |
| 1.3 The local mixed model . . . . .                                                               | 36  |
| 1.3.1 The formal two-dimensional holomorphic singular product . . . . .                           | 38  |
| 1.3.2 The closed–open factorization triple: $E_2$ bulk, $E_1$ brane, derived-centre map . . . . . | 41  |
| 1.4 Local setup at one brane stack . . . . .                                                      | 51  |
| 1.5 Notation for the local theorem . . . . .                                                      | 54  |
| 1.6 Three notions of locality . . . . .                                                           | 55  |
| 1.7 Unitarity . . . . .                                                                           | 56  |
| 2 The shifted-cotangent BF Lie algebra . . . . .                                                  | 57  |
| 2.1 Closed mixed Hamiltonian sector . . . . .                                                     | 60  |
| 2.2 Hamiltonian BF action . . . . .                                                               | 63  |
| 3 The derived commuting variety stack at $N$ Dirac branes . . . . .                               | 65  |
| 3.1 The Dirac brane stack . . . . .                                                               | 65  |
| 3.2 Hamiltonian sector and stable large $N$ . . . . .                                             | 68  |
| 3.3 Open mixed brane states . . . . .                                                             | 71  |
| 3.4 Open mixed brane field theory . . . . .                                                       | 73  |
| 3.4.1 Ghost-zero open field content . . . . .                                                     | 74  |
| 3.5 Open-brane operators and the large $N$ limit . . . . .                                        | 77  |
| 4 Boundary algebra and trace map . . . . .                                                        | 90  |
| 4.1 The reduced CE/PV trace map . . . . .                                                         | 91  |
| 4.2 The two algebras . . . . .                                                                    | 94  |
| 4.3 Reduced Hamiltonian brane coupling . . . . .                                                  | 96  |
| 5 CE/PV dictionary as Koszul resolution . . . . .                                                 | III |
| 5.1 The concrete dictionary . . . . .                                                             | III |
| 5.2 The CE/PV Koszul resolution of the brane $E_1$ -Hochschild . . . . .                          | II2 |
| 5.3 The local theorem . . . . .                                                                   | II5 |
| 5.4 Tate Quillen equivalence and the infinity-categorical upgrade . . . . .                       | 122 |
| 5.4.1 The Tate chain category . . . . .                                                           | 123 |
| 5.4.2 Twisting cochains and the filtered-cobar admissibility criterion . . . . .                  | 126 |



|     |       |                                                                                                             |     |
|-----|-------|-------------------------------------------------------------------------------------------------------------|-----|
|     | 5.4.3 | Operad-algebra transfer . . . . .                                                                           | 130 |
|     | 5.4.4 | Bar-cobar Quillen equivalence in the admissible envelope . . .                                              | 131 |
|     | 5.4.5 | Promotion to the associated infinity-categories . . . . .                                                   | 132 |
|     | 5.4.6 | Identification with the Lie operadic Koszul duality . . . . .                                               | 132 |
|     | 5.4.7 | Module reconstruction from the CE/PV identity . . . . .                                                     | 133 |
|     | 5.4.8 | Conilpotence and Casimir convergence . . . . .                                                              | 134 |
| 5.5 |       | Primitive indecomposable $P_0$ -shadow: the Koszul resolution at $F_1/F_0$ . .                              | 185 |
| 5.6 |       | Universality of the formal-local Hamiltonian sector . . . . .                                               | 202 |
| 5.7 |       | Stratified CE/PV and admissible Koszul duality . . . . .                                                    | 215 |
|     | 5.7.1 | First-order Poisson identification . . . . .                                                                | 220 |
|     | 5.7.2 | Quantization as a brane trace-map test . . . . .                                                            | 234 |
|     | 5.7.3 | Graph formalism and specialization datum . . . . .                                                          | 243 |
|     | 5.7.4 | Formal Weyl/Moyal target, all orders . . . . .                                                              | 251 |
|     | 5.7.5 | Third-order target for Costello's graph problem . . . . .                                                   | 264 |
| 6   |       | The Capelli scalar . . . . .                                                                                | 268 |
|     | 6.1   | The $U(1)$ centre-of-mass cocycle . . . . .                                                                 | 268 |
|     | 6.2   | Operators in the closed Hamiltonian sector . . . . .                                                        | 279 |
| 7   |       | The obstruction calculus: master deformation complex of local-to-global coherence                           | 282 |
|     | 7.1   | Filtered $L_\infty$ target deformation complex . . . . .                                                    | 283 |
|     | 7.2   | The transported local datum . . . . .                                                                       | 283 |
|     | 7.3   | Twists: three mechanisms . . . . .                                                                          | 284 |
|     | 7.4   | Four-curvature taxonomy and six-coordinate classification . . . . .                                         | 285 |
|     | 7.5   | Tate-coefficient cotangent lift: the six gauge slices in body . . . . .                                     | 286 |
| 8   |       | The pro-Matlis target: Matlis principal parts and the continuous cotangent module                           | 287 |
|     | 8.1   | The defect polarization . . . . .                                                                           | 288 |
|     | 8.2   | Construction of the residual side . . . . .                                                                 | 289 |
|     | 8.3   | Rigidity of the residue pairing . . . . .                                                                   | 293 |
|     | 8.4   | Local-nilpotence obstruction to polynomial realization . . . . .                                            | 293 |
|     | 8.5   | Universal Fourier–Rees label bridge . . . . .                                                               | 294 |
|     | 8.6   | Pro-Matlis envelope and bulk-boundary duality . . . . .                                                     | 296 |
| 9   |       | Examples . . . . .                                                                                          | 330 |
|     | 9.0.1 | The single brane: $N = 1$ . . . . .                                                                         | 330 |
|     | 9.0.2 | The brane stack: $N \geq 2$ . . . . .                                                                       | 330 |
| 10  |       | The frontier . . . . .                                                                                      | 330 |
|     | 10.1  | Open problem: chiral $E_d$ Deligne conjecture at $d \geq 2$ . . . . .                                       | 331 |
|     | 10.2  | Open problem: Hall–Drinfeld–Borcherds comparison spectral sequence<br>on compact non-toric $CY_3$ . . . . . | 332 |
|     | 10.3  | Open problem: operator-level lift of the modular line on $K_3 \times E$ . . . .                             | 333 |
|     | 10.4  | Frontier input: MNOP and framing on $S^3$ . . . . .                                                         | 333 |
|     | 10.5  | Frontier input: Hall–Drinfeld realisation on compact $CY_3$ . . . . .                                       | 334 |
|     | 10.6  | Frontier input: gravitational lift of $\Phi_{10}$ . . . . .                                                 | 334 |
|     | 10.7  | Convention firewall and matched-conventions acceptance . . . . .                                            | 334 |
| A   |       | Higher factorization categories and chiral derived centres at $d \geq 2$ . . . . .                          | 350 |
|     | A.1   | Factorization $E_d$ -algebras on smooth complex $d$ -folds . . . . .                                        | 350 |

|        |                                                                                            |     |
|--------|--------------------------------------------------------------------------------------------|-----|
| A.2    | Mixed holomorphic-topological refinement . . . . .                                         | 351 |
| A.3    | The chiral derived centre functor . . . . .                                                | 351 |
| A.4    | The formal-Darboux stalk functor . . . . .                                                 | 352 |
| A.5    | The bulk closed-string sector as an $E_2^{\text{HT}}$ -factorization algebra . . . . .     | 352 |
| A.6    | The brane open category as factorization $E_1^{\text{HT}}$ -module . . . . .               | 353 |
| A.7    | The Mixed Holomorphic–Topological Deligne conjecture as a functorial statement . . . . .   | 354 |
| A.8    | The chiral $E_d$ -operad: holomorphic little disks with principal-parts data . . . . .     | 355 |
| A.9    | Three hypotheses for chiral $E_d$ Deligne at $d \geq 2$ . . . . .                          | 356 |
| A.10   | Descent over the Kato–Nakayama log boundary . . . . .                                      | 357 |
| A.11   | All-loop QME for finite-jet PVA . . . . .                                                  | 359 |
| A.12   | Operator-level lift of the modular line at $\Delta_5$ . . . . .                            | 360 |
| A.13   | Convergent obstruction taxonomy . . . . .                                                  | 361 |
| B      | Factorization current conventions for the CE/PV interface . . . . .                        | 362 |
| C      | Sign conventions for the Hamiltonian comparison . . . . .                                  | 389 |
| D      | Full Primitive Higher- $\psi$ Koszul Homology . . . . .                                    | 393 |
| E      | Unreduced BV cotangent, scalar QME obstructions, and cancellations . . . . .               | 403 |
| F      | Radial parts and the Moyal normalization . . . . .                                         | 448 |
| G      | First-principles completion of the formal-local theorem . . . . .                          | 474 |
| G.1    | The host filtered nuclear Tate dg category . . . . .                                       | 475 |
| G.2    | The local Hamiltonian BF theory as a genuine BV theory . . . . .                           | 476 |
| G.3    | Hamiltonian shears and the trace uniqueness theorem . . . . .                              | 477 |
| G.4    | The decorated PBW–Stokes complex and $\Omega_{a,b}^{\text{rad}} = 0$ . . . . .             | 478 |
| G.4.1  | Explicit primitives $A_{a,b,N}^{(j)}$ from the Vandermonde-conjugated radial map . . . . . | 478 |
| G.5    | The pro-Matlis weighted Köthe target . . . . .                                             | 480 |
| G.6    | Bracket-admissible $B$ and kernel-admissible $\mathcal{K}_B$ . . . . .                     | 481 |
| G.7    | The Tate bar-cobar admissible envelope . . . . .                                           | 482 |
| G.8    | Open-trace $A_\infty$ -Koszul acceptance datum . . . . .                                   | 482 |
| G.9    | Hochschild centrality homotopies for $\Phi_N$ . . . . .                                    | 483 |
| G.10   | Protected-sector cyclic BV strong deformation retract . . . . .                            | 484 |
| G.11   | Scalar-contact projection and the QME counterterm tower . . . . .                          | 485 |
| G.11.1 | Finite-window QME counterterm recursion . . . . .                                          | 486 |
| G.12   | Costello brane-defect graph QME . . . . .                                                  | 486 |
| G.13   | Stratified factorization algebra on $(X, L)$ . . . . .                                     | 487 |
| G.14   | Large- $N$ trace state and connected cumulants . . . . .                                   | 488 |
| G.15   | The universal datum and master Maurer–Cartan equation . . . . .                            | 488 |
| G.16   | Cross-target and external-target reading . . . . .                                         | 490 |
| G.17   | Summary: the curvature decomposition . . . . .                                             | 491 |
| G.18   | Status of the constructions . . . . .                                                      | 492 |
| H      | Algorithms . . . . .                                                                       | 493 |

## LOCAL DICTIONARY

*Linear-algebraic category: Tate  $\mathbb{C}$ -vector spaces.* All linear-algebraic constructions take place in the symmetric monoidal category  $\text{Tate}_{\mathbb{C}}$  of Tate  $\mathbb{C}$ -vector spaces, equivalently linearly compact  $\mathbb{C}$ -modules in the sense of Lefschetz, equivalently the ind-pro category of finite-dimensional  $\mathbb{C}$ -vector spaces. The completed tensor product  $\widehat{\otimes}$  is the topological tensor product on  $\text{Tate}_{\mathbb{C}}$ ; continuous duality is the strict functor  $V \mapsto V_{\text{cont}}^{\vee} = \text{Hom}_{\mathbb{C}}^{\text{cts}}(V, \mathbb{C})$ . The  $\mathbb{C}$ -algebra  $A = \mathbb{C}[[z_1, z_2]]$  is a complete linearly topologized  $\mathbb{C}$ -algebra with the  $\mathfrak{m}$ -adic filtration  $\mathfrak{m}^k = (z_1, z_2)^k A$ ;  $\mathfrak{h} = A/\mathbb{C} \cdot 1$  is pro-finite-dimensional in this filtration, and its strict continuous dual is the Grothendieck–Matlis principal-part module

$$\mathfrak{h}_{\text{cont}}^{\vee} = \mathcal{P} = H_{\mathfrak{m}}^2(A) \otimes \omega_A = \bigoplus_{a+b>0} \mathbb{C} \rho_{a,b}, \quad \rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2,$$

ind-finite-dimensional, with the residue pairing  $\langle f, \omega \rangle = \text{Res}_o f \omega$  a perfect pairing of Tate  $\mathbb{C}$ -modules. Inverse limits over Matlis windows  $\mathcal{P}_{\leq N} = \bigoplus_{a+b \leq N} \mathbb{C} \rho_{a,b}$  are read as derived limits  $\mathbf{R}\varprojlim_N \mathcal{P}_{\leq N}$ , with  $\varprojlim^1$ -data tracked. Polynomial calculations use the dense ind-finite-dimensional sub-Lie algebra  $\tilde{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1$ ; completed CE/PV calculations use  $\mathfrak{h}$  itself. Factorization algebras throughout are taken in the Costello–Gwilliam sense (precosheaf on opens, Weiss-cover descent), refined to mixed-HT opens on  $\mathbb{R}^2 \times \mathbb{C}^2$ . This is the analytic incarnation of the factorisable- $\mathcal{D}$ -module-on- $\text{Ran}(\mathbb{C}^2)$  description — the two-complex-dimensional extension of the chiral-algebra construction on curves [28, §3], equivalent to the analytic form via the de Rham complex; the BV / heat-kernel framework selects Costello–Gwilliam as native.

$$X = \mathbb{R}_{t,s}^2 \times \mathbb{C}_{z_1, z_2}^2, \quad L = \mathbb{R}_t \times \{s = z_1 = z_2 = o\}, \quad \omega = dz_1 \wedge dz_2.$$

*Unimodularity datum.*

| Datum                      | Definition                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
|----------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $\Omega_{\text{loc}}$      | The fixed holomorphic volume and symplectic form on the formal polydisk<br>$\widehat{\mathbb{C}^2}_o = \text{Spf } \mathbb{C}[[z_1, z_2]], \quad \Omega_{\text{loc}} = dz_1 \wedge dz_2.$ <p>It trivializes the local canonical line and fixes the residue orientation.</p>                                                                                                                                                                                                                         |
| Poisson tensor and bracket | The inverse bivector is<br>$P = \partial_{z_1} \otimes \partial_{z_2} - \partial_{z_2} \otimes \partial_{z_1}, \quad \{f, g\} = \partial_{z_1} f \partial_{z_2} g - \partial_{z_2} f \partial_{z_1} g.$ <p>Polynomial calculations use <math>\tilde{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1</math>; completed CE/PV calculations use <math>\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1</math>. Constants are removed because they generate the zero Hamiltonian vector field.</p> |
| Hamiltonian vector field   | The convention is<br>$\iota_{X_f} \Omega_{\text{loc}} = -df, \quad X_f = (\partial_{z_1} f) \partial_{z_2} - (\partial_{z_2} f) \partial_{z_1}.$ <p>Thus <math>X_f(g) = \{f, g\}</math> and <math>[X_f, X_g] = X_{\{f, g\}}</math> after projecting constants.</p>                                                                                                                                                                                                                                  |

| Datum                                        | Definition                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
|----------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Unimodularity /<br>divergence-free condition | <p>For a holomorphic vector field <math>X = a_1 \partial_{z_1} + a_2 \partial_{z_2}</math>,</p> $\mathcal{L}_X \Omega_{\text{loc}} = (\text{div}_{\Omega_{\text{loc}}} X) \Omega_{\text{loc}}, \quad \text{div}_{\Omega_{\text{loc}}} X = \partial_{z_1} a_1 + \partial_{z_2} a_2.$ <p>Every Hamiltonian vector field is divergence-free:</p> $\text{div}_{\Omega_{\text{loc}}} X_f = -\partial_{z_1} \partial_{z_2} f + \partial_{z_2} \partial_{z_1} f = 0.$ <p>Conversely, on the formal polydisk the holomorphic Poincare lemma identifies <math>\mathbf{Ker}(\partial_{\Omega_{\text{loc}}} : \text{PV}^1 \rightarrow \text{PV}^0)</math> with Hamiltonians modulo constants.</p> |
| Closed BV pairing                            | <p>For the Hamiltonian BF fields</p> $\Omega^\bullet(\mathbb{R}^2) \widehat{\otimes} \Omega^{0,\bullet}(\mathbb{C}^2) \widehat{\otimes} (\mathfrak{h}[1] \oplus \mathfrak{h}_{\text{cont}}^\vee[2]),$ <p>the BV pairing is the compactly supported local pairing obtained by wedging the form factors, integrating against <math>\Omega_{\text{loc}}</math>, and applying the continuous evaluation <math>\mathfrak{h}_{\text{cont}}^\vee \otimes \mathfrak{h} \rightarrow \mathbb{C}</math>. It has degree <math>-1</math>. A Costello coefficient coevaluation kernel additionally requires a Tate/weighted category in which the Casimir converges.</p>                             |
| Residue / cyclic trace<br>density            | <p>The same orientation fixes the principal-part residue pairing</p> $\rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2, \quad \text{Res}_0(\rho_{a,b} z_1^m z_2^n) = \delta_{a,m} \delta_{b,n},$ <p>with <math>a + b &gt; 0</math> in the reduced cotangent module. Hamiltonian vector fields preserve <math>dz_1 dz_2</math>, so residue integration by parts gives the coadjoint action. On the open brane, the top Ext class is the residue orientation dual to <math>dz_1 \wedge dz_2</math>, and the cyclic trace is</p> $\text{Tr}_{\text{open}}(\eta \otimes u \otimes M) = \int_{\mathbb{R}} \eta[\varepsilon_1 \varepsilon_2] u \text{Tr}_N(M).$                                  |

### *Kinematic locality.*

**THEOREM 0.1** (*Kinematic locality*). On the local pair  $(X, L)$  with unimodularity datum  $\omega = dz_1 \wedge dz_2$  and reduced Hamiltonian Lie algebra  $\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$ :

- (i) The equal-time bracket kernel  $\delta(t - t')$  on two brane-time points, smeared against test functions  $a(t), b(t')$ , gives the support-local  $P_0$ -bracket  $\{B_f(a), B_g(b)\} = B_{\{f,g\}}(ab)$  with  $ab$  the pointwise product on  $\mathbb{R}_t$ ; disjoint supports bracket to zero.
- (ii) Every topological translation  $v \in T\mathbb{R}_{\text{top}}^2$  is  $Q$ -exact,  $L_v = [Q, \iota_v]$ , so  $H^\bullet(\text{Obs}^{\text{cl}})$  is locally constant along  $\mathbb{R}_t$  and inclusions of disks act by quasi-isomorphism.
- (iii) Every antiholomorphic translation  $\partial/\partial \bar{z}_i$  is  $Q$ -exact,  $L_{\partial/\partial \bar{z}_i} = [Q, \iota_{\partial/\partial \bar{z}_i}]$ , so the surviving  $\bar{\partial}$ -cohomology at the brane point is the formal jet algebra  $\mathfrak{h}$ , Grothendieck-dually paired with  $\mathcal{P} = \text{Ann}_{\text{res}}(\mathbb{C} \cdot 1) \subset H_{\text{m}}^2(\mathbb{C}[[z_1, z_2]]) dz_1 dz_2$ .

Items (ii) and (iii) hold on  $Q$ -cohomology; (i) is a chain-level identity after distributional pairing.

Trace observables  $\text{Tr } f(\phi_1, \phi_2)$  smeared as  $B_f(a)$  satisfy (i)–(iii): support locality from the equal-time bracket of the matrix-model fields, topological locality from the  $Q$ -exact  $t$ -direction, formal-holomorphic locality from the  $\bar{\partial}$ -exact transverse direction. Shifted-cotangent currents  $\Theta_\rho(B)$  satisfy (i) and (ii), and their formal-holomorphic content is the residue current  $\rho \in \mathcal{P}$  at the brane point.

The dg Lie input of an acceptable factorization algebra is therefore locally constant in  $\mathbb{R}_{\text{top}}^2$  and holomorphic in  $B \subset \mathbb{C}_{\text{hol}}^2$ ,

$$\Omega_c^\bullet(U) \widehat{\otimes} \Omega_c^{\circ, \bullet}(B) \widehat{\otimes} \mathfrak{g}, \quad \mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1],$$

with  $P_\circ$ -bracket on smeared currents fixed by (i). This produces the  $\mathbb{C}^2$  holomorphic factorization algebra  $\mathcal{F}_{\text{Ham}}^{\text{hol}}$  and its mixed topological-current enhancement  $\mathcal{F}_{\text{Ham}}^{\text{mix}}$ .

*Holomorphic factorization on  $\mathbb{C}^2$  and its curve restrictions.* The holomorphic object forced by Theorem 0.1 is a two-complex-dimensional factorization algebra on  $\mathbb{C}_{\text{hol}}^2$ . The taxonomy below records each object on  $\mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$  and the restriction data required to descend to a one-complex-dimensional factorization algebra on a holomorphic curve.

| Object                                                                 | Definition and scope                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Formal Hamiltonian holomorphic factorization algebra on $\mathbb{C}^2$ | <p>The <math>\mathbb{C}^2</math> Hamiltonian holomorphic factorization algebra is</p> $\mathcal{F}_{\text{Ham}}^{\text{hol}},$ <p>the formal-disk Hamiltonian BF object on <math>(\widehat{\mathbb{C}}_\circ^2, dz_1 \wedge dz_2)</math>. For a holomorphic polydisk <math>B</math> its dg Lie input is</p> $\Omega_c^{\circ, \bullet}(B) \widehat{\otimes} \mathfrak{g}, \quad \mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1], \quad \mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1,$ <p>with differential <math>\bar{\partial}</math> plus the Hamiltonian BF differential. The formal stalk of observables is the completed CE/PV object</p> $C_{\text{CE,cont}}^\bullet(\mathfrak{g}) \simeq \text{PV}_{\text{coord}}(\mathcal{A}_{\partial, \text{coord}})$ <p>in the coordinate/admissible sense recorded below. This object is the two-complex-dimensional holomorphic factorization algebra; a one-complex-dimensional vertex algebra appears only after specifying a curve restriction.</p>                                                                                                                                                                                              |
| Native two-dimensional holomorphic singular product                    | <p>For two points <math>z, w \in \mathbb{C}^2</math>, set <math>\zeta_i = z_i - w_i</math>. The singular coefficient module is</p> $H_\Delta^2(\mathbb{C}[[z, w]]) d\zeta_1 d\zeta_2 = \bigoplus_{a, b \geq 0} \mathbb{C}[[w_1, w_2]] \rho_{a, b}^\Delta,$ <p>where</p> $\rho_{a, b}^\Delta = \zeta_1^{-a-1} \zeta_2^{-b-1} d\zeta_1 d\zeta_2.$ <p>The normalized two-dimensional Cauchy kernel is</p> $K_\Delta^{(2)} = \rho_{\circ, \circ}^\Delta = \frac{d\zeta_1 d\zeta_2}{\zeta_1 \zeta_2}.$ <p>For <math>x, y \in \mathfrak{g}</math> the native current singular product is</p> $J_x(z) J_y(w) \sim K_\Delta^{(2)}(z, w) J_{[x, y]}(w),$ <p>which gives <math>B_f B_g \sim K_\Delta^{(2)} B_{\{f, g\}}</math>, <math>B_f \Theta_\rho \sim K_\Delta^{(2)} \Theta_{f \cdot \rho}</math>, and <math>\Theta_\rho \Theta_\sigma \sim 0</math>. Arity-<math>n</math> extension and operadic associativity: Proposition 5.7. The quantum brane trace adds <math>\hbar N \omega(f, g) K_\Delta^{(2)} \mathbf{1}</math>, with <math>\omega(f, g) = [\{f, g\}]_\circ</math>. This is the formal-coordinate two-complex-dimensional OPE. A one-variable Laurent expansion is the additional curve-restriction datum recorded below.</p> |

| Object                                                    | Definition and scope                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|-----------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Mixed topological-current enhancement over $\mathbb{R}^2$ | <p>For a topological open <math>U \subset \mathbb{R}_{\text{top}}^2</math> and holomorphic polydisk <math>B \subset \mathbb{C}_{\text{hol}}^2</math>, the mixed current input is</p> $\mathfrak{g}_{U,B}^{\text{mix}} = \Omega_c^\bullet(U) \widehat{\otimes} \Omega_c^{\circ,\bullet}(B) \widehat{\otimes} (\mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1]).$ <p>Its observables are locally constant in <math>U</math> and holomorphic in <math>B</math>. The brane-line current algebras below are its support-local one-real-dimensional restrictions.</p>                                                                                                                                                                                                                                                                                                                                                                                |
| Weyl/Moyal open-string brane algebra and cyclic avatars   | <p>The degree-zero quantum brane target is the Weyl/Moyal deformation of the formal symplectic polydisk. Put</p> $P = \partial_{z_1} \otimes \partial_{z_2} - \partial_{z_2} \otimes \partial_{z_1}.$ <p>Then</p> $f * g = m \circ \exp\left(\frac{\hbar}{2} P\right)(f \otimes g),$ $\{f, g\}_\hbar = \hbar^{-1}(f * g - g * f),$ <p>with constants quotiented in the Hamiltonian Lie algebra <math>\tilde{A}_\hbar</math>. At finite <math>N</math> the open avatar is the matrix Weyl algebra generated by <math>(\phi_1)^i{}_j, (\phi_2)^k{}_l</math> with</p> $[(\phi_1)^i{}_j, (\phi_2)^k{}_l] = \hbar \delta_l^i \delta_j^k,$ <p>followed by the quantum moment-map reduction and the connected trace projection. Cyclic and supercyclic bar-cobar avatars are trace or supertrace word complexes for matrix or supermatrix branes.</p>                                                                                                         |
| Defect/principal-part current algebra                     | <p>For an interval <math>I \subset \mathbb{R}_{\text{brane}}</math>,</p> $\mathfrak{h}_I = \Omega_c^\bullet(I) \widehat{\otimes} \mathfrak{h}, \quad \mathcal{P}_I = \mathcal{D}'_c(I) \widehat{\otimes} \mathcal{P}, \quad \mathfrak{g}_I = \mathfrak{h}_I \ltimes \mathcal{P}_I[1],$ <p>where</p> $\mathcal{P} = \text{Ann}_{\text{res}}(\mathbb{C} \cdot 1) \subset H_{\text{m}}^2(\mathbb{C}[[z_1, z_2]]) dz_1 dz_2.$ <p>The reduced target is</p> $A_\partial^{\text{pp}}(I) = \widehat{\mathbf{S}}(\Omega_c^\circ(I) \widehat{\otimes} \tilde{A}) \widehat{\otimes} \widehat{\mathbf{S}}(\mathcal{D}'_c(I) \widehat{\otimes} \mathcal{P}[1]),$ <p>with brackets</p> $\{B_f(a), B_g(b)\} = B_{\{f,g\}}(ab),$ $\{B_f(a), \Theta_\rho(B)\} = \Theta_{f \cdot \rho}(aB),$ $\{\Theta_\rho(B), \Theta_\sigma(C)\} = 0.$ <p>This is a support-local real-current <math>P_o</math> algebra with Matlis principal parts in two holomorphic variables.</p> |

| Object                                          | Definition and scope                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
|-------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| One-dimensional vertex/OPE restriction boundary | <p>A restriction of the <math>\mathbb{C}^2</math> Hamiltonian holomorphic factorization algebra to a one-complex-dimensional vertex/OPE algebra starts from the native diagonal singular product above and requires the extra datum</p> $\mathfrak{R}_L = (\iota_L, \mathcal{B}_\perp, \mathcal{R}_{L, \mathcal{B}_\perp}, V_L, \mathbf{1}_L, T_L, Y_L, \zeta_{L, \hbar}),$ <p>where <math>\iota_L: L = \text{Spf } \mathbb{C}[[w]] \hookrightarrow \widehat{\mathbb{C}^2}_o</math> is a chosen holomorphic line or defect germ, <math>\mathcal{B}_\perp</math> is a transverse boundary condition or normal vacuum, and <math>\mathcal{R}_{L, \mathcal{B}_\perp}</math> is a restriction functor producing a one-complex-dimensional holomorphic factorization algebra</p> $\mathcal{F}_{L, \mathcal{B}_\perp}(D) = \mathcal{R}_{L, \mathcal{B}_\perp}(\mathcal{F}_{\mathbb{C}^2})(D)$ <p>for disks <math>D \subset L</math>. The functor must be built from a specified tubular-neighbourhood, shriek/star restriction, or defect-coupling construction, and must prove independence of the auxiliary transverse radius, functoriality for disjoint disks, and translation covariance along the coordinate <math>w</math>. The tuple also includes the restricted state space <math>V_L</math>, vacuum, translation, state-field map, and Zhu/zero-mode map. In the <math>\mathbb{C} \times \mathbb{R}</math> interface, such a <math>V_L</math> datum factors through <math>\mathcal{R}_{\mathbb{C} \times \mathbb{R}}</math> and is identified with <math>V_{\text{red}}</math> in <math>\mathfrak{S}_{\text{ch}}</math>.</p> |
| OPE data for a restricted curve object          | <p>After such a restriction one must construct</p> $V_L = H^\bullet(\mathcal{F}_{L, \mathcal{B}_\perp}(\Delta_o)), \quad Y_L(a, w)b = \sum_{n \in \mathbb{Z}} (a_{(n)}b) w^{-n-1}.$ <p>The factorization product for two disks centered at <math>w</math> and <math>o</math> must have a finite-pole expansion</p> $\mu_{w, o}(a \boxtimes b) \sim \sum_{n \geq o} (a_{(n)}b) w^{-n-1} + \mu_{w, o}^{\text{reg}}(a, b),$ <p>with <math>a_{(n)}b = o</math> for <math>n \gg o</math>, a vacuum <math>\mathbf{1}_L</math>, translation <math>[T_L, Y_L(a, w)] = \partial_w Y_L(a, w)</math>, and locality</p> $(w_1 - w_2)^N [Y_L(a, w_1), Y_L(b, w_2)] = o$ <p>for <math>N \gg o</math>. These Laurent modes and finite singular orders belong to the restricted curve object. The two-complex-dimensional holomorphic factorization algebra uses the singular coefficient <math>H_\Delta^2(\mathbb{C}[[z, w]]) d\zeta_1 d\zeta_2</math>; after curve restriction the one-variable coefficient module is <math>\mathbb{C}((w))/\mathbb{C}[[w]] dw</math>.</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |

| Object                                          | Definition and scope                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|-------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Compatibility with the Weyl/Moyal brane algebra | <p>A one-dimensional vertex/OPE restriction must also specify how the degree-zero brane algebra acts on the restricted object. Equivalently it must supply a continuous Zhu/zero-mode map</p> $\zeta_{L,\hbar}: (\tilde{\mathcal{A}}_{\hbar}, *) \longrightarrow \text{Zhu}(V_L)$ <p>or fields <math>J_f(w)</math> satisfying</p> $J_f(w)J_g(o) \sim \frac{\kappa_L(f, g)}{w^2} + \frac{J_{\{f,g\}_{\hbar}}(o)}{w},$ <p>with the central level <math>\kappa_L</math> specified, and on the cotangent current sector</p> $[J_{f,(o)}, \Theta_{\rho}] = \Theta_{\text{ad}_{f,\hbar}^{\vee} \rho},$ <p>together with <math>\zeta_{L,\hbar}(f * g) = \zeta_{L,\hbar}(f)\zeta_{L,\hbar}(g)</math>. The Weyl/Moyal calculation supplies <math>f * g</math> and the reduced interval-current brackets; Theorem 1.15 supplies the native two-dimensional singular expansion. The data <math>\zeta_{L,\hbar}</math>, the fields <math>J_f(w)</math>, the central level, and the finite Laurent curve expansion are separate one-dimensional restriction constructions.</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Chiral algebra interface tuple                  | <p>An externally constructed chiral algebra on a curve enters as the reduced vertex object <math>V_{\text{red}}</math> in</p> $\mathfrak{S}_{\text{ch}} = (\mathcal{R}_{\mathbb{C} \times \mathbb{R}}, \mathcal{F}_{\text{red}}, V_{\text{red}}, Q_{\text{BRST}}, \mathbf{I}, T, Y, \text{wt}, \zeta_{\hbar}, H_{\text{anom}}).$ <p>Here <math>\mathcal{R}_{\mathbb{C} \times \mathbb{R}}</math> is a controlled pushforward to <math>\mathbb{R}_t \times \mathbb{C}_{z_1}</math> retaining the <math>z_2</math>-mode or the full two-index principal-part coefficient system</p> $\mathfrak{h}_{\text{red}} = \mathbb{C}[[z_1]][[z_2]]/\mathbb{C} \cdot \mathbf{I}, \quad \mathcal{P}_{\text{red}} \subset H_{(z_1, z_2)}^2(\mathbb{C}[[z_1, z_2]]) dz_1 dz_2.$ <p><math>\mathcal{F}_{\text{red}}</math> is the pushed-forward factorization algebra on <math>\mathbb{R}_t \times \mathbb{C}_{z_1}</math>. The interface must prove the pushed-forward bracket, BV pairing, brane image, factorization-to-vertex package, BRST nilpotence, Zhu/zero-mode map, Capelli normalization, Moyal multiplicativity, the completed stable Hochschild/HKR/CE-PV comparison, and anomaly transport. The obstruction vector is</p> $\text{Ob}_{\mathfrak{S}_{\text{ch}}} = (\text{Ob}_{\text{red}}, \text{Ob}_{\text{vert}}, \text{Ob}_{\text{Zhu}}, \text{Ob}_{\text{nat}}).$ <p><math>V_{\text{red}}</math> is a reduced shadow of <math>\mathcal{F}_{\text{Ham}}^{\text{hol}}</math> on a holomorphic curve; the two-complex-dimensional algebra remains <math>\mathcal{F}_{\text{Ham}}^{\text{hol}}</math>.</p> |

*Normal  $\Omega$ -background.* The brane-preserving equivariant deformation scales the normal bundle  $N_L X = \mathbb{R}_{\mathfrak{s}} \oplus \mathbb{C}_{z_1} \oplus \mathbb{C}_{z_2}$  with  $t$  fixed; the fixed locus condition singles out this normal scaling. The objects and identities below define the deformation in the mixed de Rham/Dolbeault coefficient category: the Cartan differential  $Q_{\Omega} = Q + \iota_{V_{\Omega}}$ , the localized normal contraction, the equivariant CE/PV bracket, and the residue-versus-Euler normalization.



| Object or convention            | Definition                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|---------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Ambient pair and fixed stratum  | <p>The local pair is</p> $X = \mathbb{R}_{t,s}^2 \times \mathbb{C}_{z_1,z_2}^2, \quad L = \mathbb{R}_t \times \{s = z_1 = z_2 = 0\}.$ <p>The normal bundle of the brane line is</p> $N_L X = \mathbb{R}_s \oplus \mathbb{C}_{z_1} \oplus \mathbb{C}_{z_2}.$ <p>The <math>\Omega</math>-background scales this bundle with <math>t</math> fixed.</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| Equivariant parameters          | <p>Write</p> $T_\Omega = \mathbb{C}_{\epsilon_s}^* \times \mathbb{C}_{\epsilon_1}^* \times \mathbb{C}_{\epsilon_2}^*.$ <p>The parameters <math>\epsilon_s, \epsilon_1, \epsilon_2</math> are equivariant weights. The open trace orientation uses the separate Ext generators <math>\varepsilon_1, \varepsilon_2</math>. The action is</p> $(s, z_1, z_2) \mapsto (\lambda_s s, \lambda_1 z_1, \lambda_2 z_2), \quad t \mapsto t.$ <p>Its infinitesimal vector field is</p> $V_\Omega = \epsilon_s s \partial_s + \epsilon_1 z_1 \partial_{z_1} + \epsilon_2 z_2 \partial_{z_2}.$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Equivariant differential        | <p>The Cartan differential on the local mixed complex is</p> $Q_\Omega = Q + \iota_{V_\Omega}, \quad Q_\Omega^2 = L_{V_\Omega}.$ <p>Thus <math>Q_\Omega</math> is an honest differential on the <math>T_\Omega</math>-invariant subcomplex. Away from invariants the correct object is the <math>T_\Omega</math>-equivariant curved/Cartan package with curvature <math>L_{V_\Omega}</math>.</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| Normal-weight localization ring | <p>For a finite <math>(N, M)</math>-window let <math>X_{N,M}</math> be the finite set of nonzero moving characters that occur in that window. It contains the normal Hamiltonian and residue-dual characters</p> $n\epsilon_s + a\epsilon_1 + b\epsilon_2, \quad n\epsilon_s - a\epsilon_1 - b\epsilon_2$ <p>whenever the corresponding moving summands occur. The localized coefficient ring is</p> $R_\Omega^{N,M} = \mathbb{C}[\epsilon_s, \epsilon_1, \epsilon_2, \hbar_\omega][\chi^{-1} \mid \chi \in X_{N,M}].$ <p>Equivalently, one inverts only the nonzero weights present in the finite normal window. This is the algebra in which moving normal modes may be contracted. The factor <math>\epsilon_s \epsilon_1 \epsilon_2</math> is inverted in the Euler-localization branch after that branch has been declared. The self-dual branch is a specialization with its own allowed inversions: base-change first, then invert only the surviving nonzero characters, leaving <math>\epsilon_1 + \epsilon_2</math> as a zero character on the self-dual branch.</p> |

| Object or convention              | Definition                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
|-----------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Normal contraction datum          | <p>A localized deformation retract</p> $\mathcal{C}_\Omega = (\pi_L, j_L, h_\Omega) : \mathcal{E}_\Omega(X) \rightrightarrows \mathcal{E}_\Omega(L)$ <p>for the mixed de Rham/Dolbeault coefficient complex, with</p> $\pi_L j_L = \text{id}, \quad Q_\Omega h_\Omega + h_\Omega Q_\Omega = \text{id} - j_L \pi_L$ <p>on the localized invariant normal complex. On a homogeneous normal summand of nonzero weight <math>\lambda</math>, <math>h_\Omega</math> is the Euler/Koszul homotopy divided by <math>\lambda</math>, with signs fixed by the de Rham degree in the <math>s</math>-direction and the Dolbeault degree in the <math>z_i</math>-directions.</p> |
| Hamiltonian and cotangent weights | <p>For <math>H_{a,b} = z_1^a z_2^b</math>, <math>a + b &gt; 0</math>,</p> $w(H_{a,b}) = a\epsilon_1 + b\epsilon_2.$ <p>For the principal-part residue basis,</p> $\rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2, \quad w(\rho_{a,b}) = -a\epsilon_1 - b\epsilon_2.$ <p>The holomorphic symplectic character is</p> $\chi_\omega = \epsilon_1 + \epsilon_2, \quad w(dz_1 \wedge dz_2) = \chi_\omega, \quad w(P) = -\chi_\omega.$ <p>The Weyl/Moyal deformation parameter, when present, has</p> $w(\hbar_W) = \chi_\omega.$                                                                                                                                            |
| Three $\hbar$ -parameters         | <p>The Omega convention distinguishes three formal parameters:</p> $\hbar_{\text{QME}}, \quad \hbar_W, \quad \hbar_\omega.$ <p>Here <math>\hbar_{\text{QME}}</math> is the Costello/BV loop-counting parameter in graph and curvature equations; <math>\hbar_W</math> is the Weyl/Moyal deformation parameter in the open brane algebra; and <math>\hbar_\omega</math> is the equivariant line parameter of weight <math>\chi_\omega</math> used to scalarize the Poisson bracket. A Weyl branch may identify <math>\hbar_W = \hbar_\omega</math>. Identification of <math>\hbar_{\text{QME}}</math> with another parameter is an explicit theorem convention.</p>   |
| Self-dual bracket                 | <p>On the self-dual locus</p> $\epsilon_1 + \epsilon_2 = 0$ <p>the Poisson tensor has weight 0. The Hamiltonian bracket <math>\{f, g\}</math> is then a scalar <math>T_\Omega</math>-equivariant Lie bracket on <math>\mathfrak{h}</math> after quotienting constants.</p>                                                                                                                                                                                                                                                                                                                                                                                           |

| Object or convention                    | Definition                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|-----------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Refined bracket off the self-dual locus | <p>Off the self-dual locus the ordinary Poisson bracket has weight</p> $w(\{f, g\}) = w(f) + w(g) - (\epsilon_1 + \epsilon_2).$ <p>It is therefore a bracket valued in the inverse symplectic-character line. The scalar refined bracket is obtained only after adjoining the line parameter <math>\hbar_\omega</math> of weight <math>w(\hbar_\omega) = \epsilon_1 + \epsilon_2</math>:</p> $[f, g]_\Omega = \hbar_\omega \{f, g\}.$ <p>In the Weyl/Moyal specialization <math>\hbar_\omega = \hbar_{\mathbb{W}}</math>. Before this scalarization, the refined Hamiltonian algebra is valued in the inverse symplectic-character line; after adjoining <math>\hbar_\omega</math> it becomes a scalar equivariant Hamiltonian Lie algebra.</p> |
| Equivariant CE/PV generator rule        | <p>The equivariant CE/PV map keeps the same generator labels</p> $c_{a,b} \mapsto \theta_{a,b}, \quad u_{a,b} \mapsto O_{a,b},$ <p>but all differentials and brackets are read in the <math>T_\Omega</math>-graded completed coordinate algebra. In the refined convention the contraction bracket and the <math>d_\pi</math>-terms carry the factor <math>\hbar_\omega</math>. In the self-dual convention this factor has weight 0 and may be suppressed.</p>                                                                                                                                                                                                                                                                                 |
| Equivariant quantum brane algebra       | <p>The degree-zero open quantum brane target is the equivariant Hamiltonian reduction</p> $A_{N,\Omega}^q = \text{Weyl}_{\hbar_{\mathbb{W}}}(\text{Mat}_N^2) / \hbar_{\mathbb{W}} GL_N, \quad YX - XY + \hbar_{\mathbb{W}} NI = 0,$ <p>with</p> $w(X) = \epsilon_1, \quad w(Y) = \epsilon_2, \quad w(\hbar_{\mathbb{W}}) = \epsilon_1 + \epsilon_2.$ <p>The moment-map equation is weight homogeneous. Its scalar <math>U(1)</math> contact term is the same Capelli line as in the non-equivariant dictionary, now graded by <math>\epsilon_1 + \epsilon_2</math>.</p>                                                                                                                                                                         |
| Residue normalization                   | <p>The residue convention uses the formal residue orientation</p> $\text{Res}_{z_1=z_2=0} (z_1^{-1} z_2^{-1} dz_1 dz_2) = 1$ <p>and the open top Ext coefficient <math>[\epsilon_1 \epsilon_2]</math>. Its normalization constant is</p> $\nu_\Omega^{\text{res}} = 1.$ <p>In this convention the brane trace is normalized by the residue/cyclic density. Any <math>\epsilon_s \epsilon_1 \epsilon_2</math> factor is written explicitly as a comparison constant.</p>                                                                                                                                                                                                                                                                         |

| Object or convention             | Definition                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
|----------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Euler-localization normalization | <p>The smooth equivariant localization convention pushes from <math>X</math> to <math>L</math> with the inverse normal Euler class</p> $e_{T_\Omega}(N_L X)^{-1} = (\epsilon_s \epsilon_1 \epsilon_2)^{-1},$ <p>up to the orientation sign <math>\sigma_s</math> fixed by the chosen real <math>s</math>-density and the complex orientation <math>dz_1 \wedge dz_2</math>. Its normalization constant is</p> $\nu_\Omega^{\text{Eu}} = \sigma_s(\epsilon_s \epsilon_1 \epsilon_2)^{-1}.$ <p>A theorem comparing this convention with the residue convention must state the sign and the exact factor. Until then the two normalizations are separate branches.</p> |
| Stratified factorization datum   | <p>The brane-localized object is a stratified mixed HT factorization datum on the pair <math>(X, L)</math>,</p> $\mathcal{F}_\Omega^{\text{str}}(X, L) = (\mathcal{F}_{\Omega, X \setminus L}, \mathcal{F}_{\Omega, L}, \mathcal{M}_{\Omega, L}, j^*, i^!, \mu_{\text{str}}),$ <p>where the open stratum is the <math>T_\Omega</math>-equivariant mixed bulk factorization algebra, the closed stratum is the brane-line current object, and <math>\mathcal{M}_{\Omega, L}</math> is the defect module. Protected observables become stratified factorization-homology traces after the brane-defect QME and the trace-state normalization are supplied.</p>        |
| Open obstruction vector          | <p>The residual obligations are</p> $\text{Ob}_\Omega = (\text{ob}_{\Omega, \text{Cartan}}, \text{ob}_{\Omega, \text{contr}}, \text{ob}_{\Omega, \text{CE/PV}}, \text{ob}_{\Omega, \text{norm}}, \text{ob}_{\Omega, \text{strat}}, \text{ob}_{\Omega, \text{QME}}, \text{ob}_{\Omega, N \rightarrow \infty}).$ <p>These name, respectively, the mixed Cartan model, the localized normal contraction, the equivariant CE/PV bracket comparison, the residue-versus-Euler normalization theorem, the stratified factorization structure, the Costello brane-defect graph/QME package, and the physical large-<math>N</math> trace-state theorem.</p>                 |

*Canonical maps and obstruction symbols.*

| Symbol                | Meaning                                                                                                                                                                                                                                                                                                                                                                                                                           |
|-----------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $\Phi_{\mathbf{I}}$   | Finite-dimensional cotangent CE/PV map                                                                                                                                                                                                                                                                                                                                                                                            |
|                       | $C_{\text{CE}}^\bullet(\mathbf{I} \ltimes \mathbf{I}^\vee[\mathbf{I}]) \rightarrow \text{PV}(\mathbf{S}(\mathbf{I})), \quad c \mapsto \theta, \quad u \mapsto O.$                                                                                                                                                                                                                                                                 |
| $\Phi_{\text{coord}}$ | <p>Formal-polydisk coordinate CE/PV map reconstructed from admissible finite coordinate tests and pro-window coordinate cochains for <math>\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1</math> and <math>\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[\mathbf{I}]</math>. It is a completed coordinate dg commutative algebra isomorphism before any <math>P_o</math> bracket is asserted.</p> |

| Symbol                                                 | Meaning                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|--------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $\Phi_{P_o}^B$                                         | <p>Restriction of <math>\Phi_{\text{coord}}</math> to a bracket-admissible subalgebra <math>B \subset \text{PV}_{\text{coord}}(\mathcal{A}_{\partial, \text{coord}})</math>:</p> $\Phi_{P_o}^B : \Phi_{\text{coord}}^{-1}(B) \xrightarrow{\sim} B.$ <p>The completed <math>P_o</math> comparison in <math>\mathcal{B}^{\text{adm}}</math> (Theorem 5.6, axioms (i)–(iv)): <math>B</math> carries a complete Hausdorff topology, continuous inclusion in the coordinate product, continuous multiplication, continuous <math>d_{\pi}^{\text{coord}}</math>, and a continuous closed contraction bracket; the operational test is Köthe bracket continuity (Remark 5.9, probe (6)). Boundary Hamiltonians inside this <math>P_o</math> comparison require the relevant <math>O_f</math> and <math>d_{\pi}^{\text{coord}} O_f</math> to lie in <math>B</math>.</p> |
| $\Theta_B$                                             | <p>Kernel-admissible Maurer–Cartan tensor in a specified completed convolution dg Lie algebra <math>\mathcal{K}_B</math>,</p> $\Theta_B = \sum_I H_I \otimes \theta^I + \sum_I \eta^I \otimes O_I,$ <p>attached to <math>B</math>. Asserted only when the displayed diagonal sums converge and the convolution differential and bracket are continuous in <math>\mathcal{K}_B</math>; the kernel-admissibility test is the Maurer–Cartan identity <math>d\Theta_B + \frac{1}{2}[\Theta_B, \Theta_B] = 0</math> (Theorem 5.6, axiom (v); Remark 5.9, probe (7)).</p>                                                                                                                                                                                                                                                                                             |
| $\eta^I$                                               | <p>Shifted cotangent generator <math>\eta^I = (H_I)^{\vee}[1] \in \mathfrak{h}_{\text{cont}}^{\vee}[1]</math>, for <math>I = (a, b)</math> and <math>a + b &gt; 0</math>. It is the source of the <math>O_I</math> coordinate in the CE/PV map.</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| $\mathfrak{U}_{\mathfrak{h}}$                          | <p>Admissible formal-local interface datum in the category <math>\mathcal{B}^{\text{adm}}</math> of Theorem 5.6 (seven admissibility axioms (i)–(vii); host details in Appendix G §1). The categorical universal object is</p> $\mathfrak{U}_{\mathfrak{h}} = (\Phi_{\text{coord}}, \{\Phi_{P_o}^B\}_B, \{\Theta_B\}_B),$ <p>where <math>B</math> ranges over bracket-admissible subalgebras in the middle term and kernel-admissible data in the last term.</p>                                                                                                                                                                                                                                                                                                                                                                                                |
| $B_{\text{fin}}, E_q$                                  | <p>Model admissibility tests. <math>B_{\text{fin}}</math> is the finite-support coordinate subalgebra on a finite window. <math>E_q</math>, for <math>q &gt; 1</math>, is the weighted Köthe test target with seminorms</p> $p_n(x) = \sup_{a,b}  x_{a,b}  q^{a+b} (1 + a + b)^n.$ <p>It is bracket-admissible only after the finite-word tensor convention and the corresponding convolution estimate are fixed; it is kernel-admissible only if the diagonal tensor converges.</p>                                                                                                                                                                                                                                                                                                                                                                            |
| $\mathfrak{o}^{\text{br}}, \mathfrak{o}^{\text{coad}}$ | <p>Transition obstruction terms for bracket and coadjoint compatibility in the pro/mixed coordinate CE/PV comparison.</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| $\mathfrak{o}^{\text{Cas}}$                            | <p>Strict product/direct-sum kernel obstruction for the Casimir, propagator, and Costello endpoint problem. It belongs to the endpoint coefficient problem; the coordinate CE/PV identity is the separate coordinate comparison.</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| $\mathfrak{o}_{\text{BV}, \text{end}}$                 | <p>Endpoint obstruction tuple <math>(\mathfrak{o}_{\text{Cas}}, \mathfrak{o}_{\text{def}}, \mathfrak{o}_{\text{RG}}, \mathfrak{o}_{\text{QME}})</math> controlling a Costello-compatible coefficient endpoint.</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |

| Symbol                                  | Meaning                                                                                                                                                                                                                                                                                             |
|-----------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $\text{Ob}_{\mathfrak{C}_{\text{tar}}}$ | Matched-conventions descent vector<br>( $\text{per}_{\text{tar}}, \text{ob}_{\text{WR}}, \text{ob}_{\text{anom}}, \text{ob}_{\text{QME}}, \text{ob}_{\text{conv}}, \text{ob}_{\text{hyp}}, \text{ob}_{6r}, \text{ob}_{\text{MD}}$ ), indexed by the comparison target $\mathfrak{C}_{\text{tar}}$ . |

*Non-scalar Costello/QME labels.* The local obstruction complex named in Problem 5.219 is recorded below. The scalar projection  $\sigma_{\text{sc}}$  is associated-graded data; the projection  $\widehat{\sigma}_{\text{sc}}$  is a filtered chain map existing exactly when the successive scalar-lift obstructions vanish; the reduced principal-part brackets live after de Rham-current contraction; the unreduced non-scalar QME is the curvature equation  $\text{Curv}(K_{\hbar,w,I}) = 0$  of the bulk-to-defect kernel.

| Symbol or datum                                        | Meaning                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|--------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $\mathfrak{Q}_{w,\partial,\hbar}^\bullet(I)$           | Filtered brane-defect local-functional complex<br>$\varprojlim_M \left( \mathcal{O}_{\text{loc}}^{\text{bd}}(\mathcal{E}_w^M _I; A_{\partial,\hbar}^{\text{pp},w,M}(I)), d_M \right), \quad d_M = Q + \{I_{o,M}^w, -\}_{\hbar,M}.$ <p>The inverse limit is taken over finite Hamiltonian windows with the <math>\hbar</math>-adic, scalar-contact, and finite-window filtrations. The differential has degree <math>+1</math>. This is the target complex for the bulk-to-defect kernel and QME obstruction; the reduced current algebra is <math>A_{\partial,\hbar}^{\text{pp},w}(I)</math>.</p>                                                                                                      |
| $F^\bullet \mathfrak{Q}_{w,\partial,\hbar}^\bullet(I)$ | Complete decreasing closed scalar-contact filtration preserved by $d_{\text{QME}}$ . In the ordinary scalar-reduced branch take $F^r = J_{\text{sc}}^r$ , where $J_{\text{sc}}$ is the closed ideal of functionals with zero associated-graded scalar-contact symbol. In the balanced branch it is the closed filtration generated by identity-endomorphism contractions inside the balanced scalar-contact complex. Its role is to define the associated-graded scalar symbol and the obstruction tower for lifting that symbol.                                                                                                                                                                      |
| $\sigma_{\text{sc}}$                                   | Associated-graded scalar-symbol extractor<br>$\sigma_{\text{sc}}: \text{gr}_F \mathfrak{Q}_{w,\partial,\hbar}^\bullet(I) \longrightarrow C_{\text{Lie}}^\bullet(\tilde{A}; \mathbb{C}\gamma_1)[1][[\hbar]].$ <p>It has cochain degree 0 as an associated-graded map. A projection of the filtered QME complex is the lifted map <math>\widehat{\sigma}_{\text{sc}}</math>. On a degree-one scalar contact with two Hamiltonian inputs it records the Lie two-cocycle coefficient of <math>\gamma_1</math>. In the ordinary branch this leading symbol is <math>\hbar N[\tilde{c}]</math>.</p>                                                                                                          |
| $\widehat{\sigma}_{\text{sc}}$                         | Actual scalar-contact projection: a continuous filtered degree-0 cochain map<br>$\widehat{\sigma}_{\text{sc}}: \mathfrak{Q}_{w,\partial,\hbar}^\bullet(I) \longrightarrow C_{\text{Lie}}^\bullet(\tilde{A}; \mathbb{C}\gamma_1)[1][[\hbar]]$ <p>whose associated graded is <math>\sigma_{\text{sc}}</math>. It exists exactly when the scalar-lift obstruction classes <math>\mathcal{O}_{\sigma,w}^{(r)}(I)</math> vanish successively with compatible finite-window choices. The residual non-scalar complex is defined only after this map has been chosen. On a balanced scalar-contact complex the proved projection is the zero chain map <math>\widehat{\sigma}_{\text{bal,sc}} = 0</math>.</p> |

| Symbol or datum                             | Meaning                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
|---------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $o_{\sigma,w}^{(r)}(I)$                     | <p>Successive scalar-contact lift obstruction. For a filtered linear lift <math>s</math> of <math>\sigma_{\text{sc}}</math>, the chain defect <math>\partial_s = d_{\text{CE}}s - sd_{\text{QME}}</math> lowers <math>F^\bullet</math>. Its first nonzero part gives</p> $o_{\sigma,w}^{(r)}(I) \in H^1\left(\text{Hom}(\text{gr}_F \mathfrak{Q}_{w,\partial,\hbar}^\bullet(I), C_{\text{Lie}}^\bullet(\bar{A}; \mathbb{C}\gamma_1)[1][[\hbar]])_{-r}\right).$ <p>In the ordinary scalar-reduced <math>\mathfrak{gl}_N</math> branch the first two-input evaluation is <math>\hbar N \omega(f, g)\gamma_1</math>, nonzero on <math>(f, g) = (z_1, z_2)</math>.</p>                                      |
| $\mathfrak{g}_{\hbar,I}^{w,\text{cur}}$     | <p>Current source for the unreduced non-scalar lift,</p> $(\Omega_c^\bullet(I) \widehat{\otimes} \mathfrak{h}_{w,\hbar}) \ltimes (\mathcal{D}'_c(I) \widehat{\otimes} \mathfrak{h}_{\hbar,\text{cont}}^{\vee,w})[1].$ <p>The topology is the completed current topology with the weighted finite-window and <math>\hbar</math>-adic completions. On CE coordinates, <math>c_{B,\rho}</math> has degree 1 and <math>u_{a(t)dt \otimes f}</math> has degree 0. Before scalar reduction one may replace the Hamiltonian source by the central-character source.</p>                                                                                                                                        |
| $\kappa_{\hbar,w,I}$                        | <p>Continuous bulk-to-defect kernel, equivalently a degree-0 Maurer–Cartan candidate in the completed convolution Lie algebra,</p> $\kappa_{\hbar,w,I} : C_{\text{CE,cont}}^\bullet(\mathfrak{g}_{\hbar,I}^{w,\text{cur}}) \longrightarrow \mathfrak{Q}_{w,\partial,\hbar}^\bullet(I),$ <p>with generator values</p> $u_{a(t)dt \otimes f} \mapsto B_f^{\text{BV}}(a), \quad c_{B,\rho} \mapsto \Theta_\rho^{\text{BV}}(B).$ <p>Its reduced image must recover <math>B_f(a)</math> and <math>\Theta_\rho(B)</math>, but the kernel itself lives before scalar reduction, before de Rham-current contraction, and before replacing unreduced open observables by freely adjoined current generators.</p> |
| $\text{Ob}_{w,\partial,\hbar}$              | <p>Curvature cocycle of the bulk-to-defect kernel,</p> $\text{Ob}_{w,\partial,\hbar} = d_{\text{QME}}\kappa_{\hbar,w,I} + \frac{1}{2}[\kappa_{\hbar,w,I}, \kappa_{\hbar,w,I}]_\hbar.$ <p>It has QME degree 1. The scalar branch first requires <math>\widehat{\sigma}_{\text{sc}}(\text{Ob}_{w,\partial,\hbar}) = 0</math>. Only then does its residual class lie in the non-scalar complex.</p>                                                                                                                                                                                                                                                                                                        |
| $\text{Ob}_{w,\partial,\hbar}^{\text{red}}$ | <p>Residual non-scalar obstruction after the scalar-contact projection has been constructed and the scalar component has been killed:</p> $[\text{Ob}_{w,\partial,\hbar}^{\text{red}}] \in H^1(\ker \widehat{\sigma}_{\text{sc}}, d_{\text{QME}}).$ <p>This class is defined after choosing the scalar-contact projection; the reduced principal-part bracket formulas leave it as a residual non-scalar obstruction.</p>                                                                                                                                                                                                                                                                               |
| $\mathcal{Q}_{w,\mathcal{M}}^\bullet(I)$    | <p>Finite-window residual complex</p> $\mathcal{Q}_{w,\mathcal{M}}^\bullet(I) = \ker(\widehat{\sigma}_{\text{sc},\mathcal{M}}) \subset \left(O_{\text{loc}}^{\text{bd}}(\mathcal{E}_w^{\mathcal{M}} _I; A_{\partial,\hbar}^{\text{pp},w,\mathcal{M}}(I)), d_{\mathcal{M}}\right).$ <p>The bracket, interaction, and differential are the transported finite-window structures supplied by the acceptance gate. The unweighted Hamiltonian bracket enters only through that transport.</p>                                                                                                                                                                                                               |

| Symbol or datum                         | Meaning                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
|-----------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $C_{n,M}, C_{\Gamma,w}^{w_o}(\epsilon)$ | <p>Finite-window counterterms. At order <math>\hbar^n</math>, <math>C_{n,M} \in \mathcal{Q}_{w,M}^o(I)</math> must satisfy</p> $d_M C_{n,M} = -\mathfrak{o}_{n,M}^{\text{ns}}, \quad \pi_{M,N} C_{n,M} = C_{n,N}.$ <p>Graph representatives <math>C_{\Gamma,w}^{w_o}(\epsilon)</math> must also commute with window truncation and weight transport. They are degree-zero local functionals in the filtered QME complex.</p>                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| $\mathfrak{D}_n^{\text{ns}}$            | <p>Order-<math>n</math> finite-window obstruction vector</p> $\mathfrak{D}_n^{\text{ns}} = (([\mathfrak{o}_{n,M}^{\text{ns}}])_M, \lambda_n),$ <p>where the first component lies in <math>\lim_{\leftarrow M} H^1(\mathcal{Q}_{w,M}^\bullet(I))</math>, and <math>\lambda_n \in \lim_{\leftarrow M} H^0(\mathcal{Q}_{w,M}^\bullet(I))</math> is the primitive-compatibility class. Its vanishing is exactly the finite-window criterion for compatible order-<math>n</math> counterterms.</p>                                                                                                                                                                                                                                                                                                                                                                           |
| $E_{\theta_3,M}, C_{\theta_3,M}$        | <p>First proved larger-row non-scalar obstruction and its permitted primitive. In the supplied one-row package,</p> $\mathcal{Q}_{\theta_3,M}^o = \mathfrak{o}, \quad \mathcal{Q}_{\theta_3,M}^i = \mathbb{C}E_{\theta_3,M}, \quad d_M = \mathfrak{o},$ <p>and the residual is <math>\mathfrak{o}_{\theta_3,M}^{\text{ns}} = E_{\theta_3,M}</math>. The covector <math>\ell_{\theta_3}(E_{\theta_3,M}) = \mathfrak{i}</math> proves that this class is nonzero in the supplied subcomplex. A valid <math>C_{\theta_3,M}</math> is one of three explicit witnesses,</p> $K_{\Theta_3}^M(\eta_{\theta_3,M}), \quad C_{\theta_3,M}^{\text{ct}}, \quad \text{or a companion-face primitive,}$ <p>with differential <math>d_M C_{\theta_3,M} = -E_{\theta_3,M}</math>, scalar-contact value zero, and vanishing Roos compatibility class under finite-window projection.</p> |
| $I_N, \eta, \chi_N$                     | <p>Moment-ideal and scalar primitive data. The quantum Hamiltonian reduction uses the left moment-map ideal <math>I_N = \mathcal{W}_N \widehat{\mu}(\mathfrak{s}I_N)</math> and then removes the scalar empty trace. The scalar primitive lives on the unreduced Hamiltonian source <math>A = \mathbb{C}[z_1, z_2]</math>:</p> $\eta(f) = -[f]_{\mathfrak{o}}, \quad d_{\text{CE}}\eta = \omega, \quad \chi_N(f) = N[f]_{\mathfrak{o}} = -N\eta(f).$ <p>Thus <math>\hbar N\omega + d_{\text{CE}}(\hbar\chi_N) = \mathfrak{o}</math> before scalar reduction. This primitive lives on <math>A</math>; after passage to <math>\tilde{A} = A/\mathbb{C} \cdot \mathfrak{i}</math>, the scalar-reduced QME complex requires its own counterterm data.</p>                                                                                                                   |



| Symbol or datum                      | Meaning                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
|--------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $E_{a,b,N}, A^j_i(a, b, N)$          | <p>Quantum shear moment-ideal primitive datum in the matrix Weyl algebra. Put</p> $E_{a,b,N} = \sum_{j=0}^b \text{Tr}(Y^j X^a Y^{b-j}) - J_N\left(\{z_1^{a+1}/(a+1), z_2^{b+1}\}_\hbar\right).$ <p>The all-<math>N</math> shear congruence required for the radial/Weyl image is equivalent to</p> $E_{a,b,N} \in I_N \iff E_{a,b,N} = \sum_{i,j} A^j_i(a, b, N) \tilde{\mu}^j_j$ <p>for coefficients <math>A^j_i(a, b, N) \in \mathcal{W}_N</math>. This is a degree-zero left-ideal primitive problem in the quantum moment-map ideal. The scalar CE primitive <math>\eta</math>, scalar-contact projection, and Costello QME counterterms are separate data.</p>                      |
| Reduced principal-part brackets      | <p>Target-side brackets in <math>A_{\partial, \hbar}^{\text{pp}, w}(I)</math> after scalar reduction and de Rham-current contraction:</p> $\{B_f(a), B_g(b)\}_\hbar = B_{\{f, g\}_\hbar}(ab), \quad \{B_f(a), \Theta_\rho(B)\}_\hbar = \Theta_{\text{ad}^\vee_{f, \hbar} \rho}(aB),$ $\{\Theta_\rho(B), \Theta_\sigma(C)\}_\hbar = 0.$ <p>They use multiplication of a distribution by a smooth test function. They prove the reduced current interface. Costello wavefront estimates, unreduced centrality homotopies, and the non-scalar QME are additional analytic and homotopical data.</p>                                                                                         |
| Unreduced non-scalar QME realization | <p>The full graph theorem is the construction of <math>\kappa_{\hbar, w, I}</math> and compatible counterterms <math>C_{\hbar, w}</math>. Put <math>K_{\hbar, w, I} = \kappa_{\hbar, w, I} + C_{\hbar, w}</math>. The equation is</p> $\text{Curv}(K_{\hbar, w, I}) = d_{\text{QME}} K_{\hbar, w, I} + \frac{1}{2} [K_{\hbar, w, I}, K_{\hbar, w, I}]_\hbar = 0.$ <p>It requires the Costello finite-scale BV propagator and Laplacian, locality and counterterm estimates, scalar-contact chain projection, vanishing of <math>\mathfrak{D}_n^{\text{ns}}</math> for every order, and unreduced product and <math>P_0</math>-bracket homotopies against arbitrary open observables.</p> |

*Common local rings and quotients.*  $\mathfrak{h}$  labels Hamiltonian observables;  $\mathfrak{g}_{\text{str}}$  labels CE/PV closed cochains;  $\tilde{\mathcal{A}}$  labels the polynomial trace sector;  $N$  labels the Dirac brane stack rank.

| Symbol                                                       | Meaning                                                                                                                                                                                                                                                                                        |
|--------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $R = \mathbb{C}[[z_1, z_2]]$                                 | Complete local ring of the formal symplectic polydisk.                                                                                                                                                                                                                                         |
| $\mathfrak{m} = (z_1, z_2)$                                  | Maximal ideal of functions vanishing at the brane point.                                                                                                                                                                                                                                       |
| $\mathfrak{h} = R/\mathbb{C} \cdot 1$                        | Reduced Hamiltonian Lie algebra of the formal polydisk. Constants are removed because they generate the zero Hamiltonian vector field.                                                                                                                                                         |
| $\mathfrak{g}_{\text{str}}$                                  | Strict shifted cotangent Hamiltonian Lie algebra                                                                                                                                                                                                                                               |
| $\mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^{\vee}[1].$ |                                                                                                                                                                                                                                                                                                |
|                                                              | Its Hamiltonian side is a product completion and its cotangent side is continuous dual/direct-sum in residue coordinates. This is the formal-local CE/PV object. A Costello coefficient kernel additionally requires the endpoint Casimir data.                                                |
| $\mathfrak{g}_w$                                             | Weighted finite-window Tate replacement used as a regulator. It carries the same coordinate-window CE/PV formulas after passage through the admissible tower. Its weight topology is auxiliary data for the admissible tower; the strict cotangent completion is $\mathfrak{g}_{\text{str}}$ . |
| $\tilde{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1$        | Polynomial Hamiltonian sector used in explicit trace and Moyal calculations.                                                                                                                                                                                                                   |
| $\tilde{A}_{\text{desc}}$                                    | Abstract polynomial label space for PBW one- $\psi$ descendants. The Matlis principal-part module is $\mathcal{P}$ .                                                                                                                                                                           |
| $[\bar{c}]$                                                  | Distinguished scalar $U(1)$ anomaly class in $H_{\text{Lie}}^2(\tilde{A}, \mathbb{C})$ , realized by the constant term of the Poisson bracket and by the Capelli shift.                                                                                                                        |
| $N$                                                          | Rank of the Dirac brane stack. Degreewise stable trace statements are formed at fixed total word degree and then for $N$ in the stable range.                                                                                                                                                  |
| $D$                                                          | Taylor truncation degree when a finite quotient $\mathfrak{m}/\mathfrak{m}^{D+1}$ is used. The brane rank is denoted $N$ .                                                                                                                                                                     |

*Large- $N$  and quotient conventions.* An algebraic primitive projection is defined inside the stable range; a physical cumulant requires a state  $\omega_{N,\Omega}$ .

| Symbol or operation                  | Meaning                                                                                                                                                                                                      |
|--------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $\text{Prim}_{\text{st}}$            | Degreewise stable primitive invariant sector. In total word degree $d$ , the Procesi–Razmyslov stable range is $N \geq d$ .                                                                                  |
| $\text{Conn}_{N \rightarrow \infty}$ | Connected primitive/indecomposable projection in the algebraic stable trace sector after finite- $N$ normalization and passage to the stable range. Physical cumulants require a state $\omega_{N,\Omega}$ . |
| Hamiltonian scalar quotient          | The closed scalar $1 \in \mathbb{C}[z_1, z_2]$ and the open empty trace $\text{Tr}_N(1) = N$ are omitted as primitive Hamiltonian generators. The unit of the symmetric algebra remains.                     |

*Separation rule.* PBW special-fibre labels  $H_{a,b}$ ,  $O_{a,b}$ ,  $\Psi_{a,b}$ , CE/PV labels  $c^I$ ,  $u_I$ ,  $\theta^I$ ,  $O_I$ , and Matlis labels  $\rho_{a,b}$  carry different  $\mathfrak{h}$ -actions and different polarizations. Shared indices are labels for separate modules. The Fourier–Rees theorem (Theorem 8.12) is the only comparison morphism between PBW and Matlis labels: it matches associated-graded labels through the Fourier delta Rees lattice  $\mathcal{D}_{\hbar}/(z_1, z_2)$ , with Čech realization carrying the factor  $(-1)^{a+b} \hbar^{a+b} a!b!$ . Before  $\hbar$  is inverted this is a Rees sublattice; as a module statement it uses a Fourier-twisted polarization. The Hamiltonian coadjoint action is the action on  $\mathcal{P}$ . Beyond the Fourier–Rees theorem the spans of  $\Psi_{a,b}$  and  $\rho_{a,b}$  carry distinct  $\mathfrak{h}$ -module structures.

Source–target table.

| Source (algebraic)                                                                             | Target (physical)                                                                                                                                                                 | Comparison morphism                                                                                                                                               | Scope                                                                    |
|------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------|
| coordinate-window tower / mixed-continuous replacement attached to $\mathfrak{g}_{\text{str}}$ | $PV_{\text{coord}}(\mathcal{A}_{\partial, \text{coord}})$                                                                                                                         | coordinate dg identity; bracketed comparison only after choosing a bracket-admissible $B$                                                                         | full unreduced factorization centre requires centrality and descent data |
| bracket-admissible $B \subset PV_{\text{coord}}$                                               | $P_o$ realization $\Phi_{P_o}^B$                                                                                                                                                  | restriction of $\Phi_{\text{coord}}$ when $B$ supplies completeness, $d_{\pi}^{\text{coord}}$ -stability, product stability, and a continuous contraction bracket | bracket-admissible subalgebra inside the chosen completion               |
| kernel-admissible $B$                                                                          | Maurer–Cartan kernel $\Theta_B$                                                                                                                                                   | diagonal-convergent kernel in the chosen completed tensor product                                                                                                 | strict ambient product completion requires separate Casimir convergence  |
| $C_{\text{CE}}^{\bullet}(\mathfrak{g}) / PV(\mathcal{A})$                                      | reduced $P_o$ central-operation model                                                                                                                                             | local Poisson cochain model                                                                                                                                       | unreduced $Z_{\text{fact}}^{P_o}$ requires centrality homotopies         |
| $U_{\hbar}(\mathfrak{g})/(\hbar)$ PBW special fibre                                            | $\mathbf{S}(\mathfrak{h} \oplus \mathfrak{h}^{\vee}[\mathbf{I}])$ and then selected $\mathbf{S}(\tilde{\mathcal{A}} \oplus \tilde{\mathcal{A}}_{\text{desc}}[\mathbf{I}])$ labels | associated-graded label comparison                                                                                                                                | PBW and Matlis labels carry different actions                            |
| admissible Tate / relative bar-cobar envelope                                                  | $\widehat{U}(\mathfrak{g}_{\text{adm}})$                                                                                                                                          | PBW envelope under conilpotent CE chains, exact continuous duality, PBW completeness, and strong convergence                                                      | available after specifying admissibility data for the formal polydisk    |
| $c^I$                                                                                          | $\theta^I$                                                                                                                                                                        | generator rule $c^I \mapsto \theta^I$ of $\Phi_{\text{coord}}$                                                                                                    | CE ghost coordinate                                                      |
| $u_I$                                                                                          | $O_I$ and brane-line Hamiltonian coordinates $B_f(a)$                                                                                                                             | generator rule $u_I \mapsto O_I$ of $\Phi_{\text{coord}}$                                                                                                         | boundary Hamiltonian coordinate                                          |
| $C_{\text{CE}}^{\bullet}(\mathfrak{g}_I)$ on intervals                                         | reduced brane-line factorization target                                                                                                                                           | cofactorization/ opposite-interval source; factorization only after the CE coproduct is paired with target multiplication                                         | variance reversal handled by the CE coproduct and target multiplication  |

| Source (algebraic)                                            | Target (physical)                                                    | Comparison morphism                                                                                                                                                                                                                                           | Scope                                                                                                                                                                               |
|---------------------------------------------------------------|----------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $\Psi_{a,b}$                                                  | stable primitive one- $\psi$<br>PBW special-fibre class              | stable-trace<br>identification in the<br>trace complex                                                                                                                                                                                                        | PBW trace label with<br>polynomial action                                                                                                                                           |
| $\rho_{a,b}$                                                  | $\mathcal{P} = \text{Ann}_{\text{res}}(\mathbb{C} \cdot \mathbf{1})$ | residue/Matlis<br>pairing<br>$f \otimes \rho \mapsto \text{Res}_o(f \rho)$                                                                                                                                                                                    | Matlis principal-part label with<br>Hamiltonian coadjoint action                                                                                                                    |
| $F_{\text{Rees}}$                                             | $\oplus \hbar^{a+b} \mathbb{C}[[\hbar]] \rho_{a,b}$                  | Fourier-polarized<br>Rees label bridge                                                                                                                                                                                                                        | polarization-changing Rees<br>comparison                                                                                                                                            |
| $\Omega_c^\bullet(I) \otimes \mathfrak{h}$                    | brane-line Hamiltonian<br>currents                                   | reduced current<br>model                                                                                                                                                                                                                                      | $C_c^\infty(I)$ denotes only the<br>degree-zero summand                                                                                                                             |
| $\mathcal{D}'_c(I) \widehat{\otimes} \mathcal{P}[\mathbf{1}]$ | principal-part current<br>sector $\Theta_\rho(B)$                    | post-de-Rham-<br>current contraction                                                                                                                                                                                                                          | uses multiplication of a<br>distribution by a smooth test<br>function                                                                                                               |
| formal Moyal algebra<br>$\hat{A}_\hbar$                       | degree-zero Weyl<br>coefficient target                               | reduced algebraic<br>Moyal map with<br>raw weights $\mathbf{1}$ and<br>$\mathbf{1}/24$                                                                                                                                                                        | finite- $N$ radial-parts theorem<br>requires the radial image data                                                                                                                  |
| finite- $N$ radial-parts<br>target                            | stable connected trace<br>realization of the Moyal<br>coefficients   | finite verified<br>relative datum:<br>formal coefficients,<br>Capelli<br>normalization, free<br>normal-diagram<br>obstruction $\mathfrak{o}_{a,b}^{\text{rad}}$ ,<br>edge PBW strip, and<br>the remaining<br>uniform splitting or<br>left-cokernel<br>theorem | requires formal Moyal<br>coefficients, radial-parts input,<br>and the normal-diagram<br>obstruction calculation;<br>Costello analytic graph/QME<br>realization is a separate target |
| Costello brane-defect<br>kernel                               | closed-open BV/QME<br>specialization                                 | obstruction datum<br>$\text{Ob}_{w,\partial,\hbar}$ of<br>bulk-to-defect<br>curvature                                                                                                                                                                         | requires bulk-to-defect<br>curvature and counterterm<br>data                                                                                                                        |
| scalar cocycle $[\tilde{c}]$                                  | $U(\mathbf{1})$ anomaly / Capelli<br>contact line                    | distinguished class<br>in $H_{\text{Lie}}^2(\tilde{A}, \mathbb{C})$                                                                                                                                                                                           | one scalar contact class inside<br>the unreduced QME problem                                                                                                                        |

*PBW special-fibre open trace labels.* Algebraic side: the formal-stalk chart at the  $\hbar = 0$  special fibre. Physical side: the trace observable  $\text{Tr } f(\phi_1, \phi_2)$  on the matrix Weyl brane after moment-map reduction. Comparison morphism: the PBW limit of that trace observable, an isomorphism of stable trace complexes in the Procesi–Razmyslov stable range  $N \geq d$ .

| Symbol                  | Meaning                                                                                                                                                                                                                                                                                                                                                                                                                                             |
|-------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $H_{a,b} = z_1^a z_2^b$ | Polynomial Hamiltonian label in $\tilde{\mathcal{A}}$ , with $a + b > 0$ . In the PBW special fibre it labels the scalar-reduced trace observable $O_{a,b}$ .                                                                                                                                                                                                                                                                                       |
| $O_{a,b}$               | Open ghost-zero trace class $\overline{\text{Tr}(\phi_1^a \phi_2^b)}$ , with $a + b > 0$ in the reduced Hamiltonian label set.                                                                                                                                                                                                                                                                                                                      |
| $\Psi_{a,b}$            | Primitive one- $\psi$ stable open class, represented by the symmetrised trace with one $\psi$ , $a$ copies of $\phi_1$ , and $b$ copies of $\phi_2$ , with $a + b > 0$ after removing the scalar descendant line. It lies in the PBW special-fibre open shadow and carries the corresponding polynomial action. In the reduced comparison it supplies the primitive indecomposable $P_0$ -shadow label.                                             |
| $F_{\text{Rees}}$       | Fourier–Rees label bridge of Theorem 8.12. It sends the polynomial label $z_1^a z_2^b$ to the Fourier-delta label $\partial_1^a \partial_2^b e_0$ , whose Čech realization is $(-1)^{a+b} \hbar^{a+b} a! b! \rho_{a,b}$ . After inverting $\hbar$ , division by this Rees factor gives the principal-part label $\rho_{a,b}$ . It changes polarization and is the comparison between the PBW label $\Psi_{a,b}$ and the Matlis label $\rho_{a,b}$ . |

The shared indices on  $H_{a,b}$ ,  $\Psi_{a,b}$ , and  $\rho_{a,b}$  are labels for separate modules: the span of the  $\Psi_{a,b}$  is the PBW special-fibre trace label space, while  $\mathfrak{h}_{\text{cont}}^\vee \cong \mathcal{P}$  is the principal-part current module. The Fourier–Rees bridge  $F_{\text{Rees}}$  is the only comparison morphism between PBW and Matlis sectors.

*Closed CE/PV labels.* Algebraic side: the closed-side complex of CE/PV cochains. Physical side: the closed-string  $P_0$ -central operation on brane observables. Comparison morphism:  $\Phi_{\text{coord}}$ , the coordinate dg CE/PV isomorphism on generators  $c^I \mapsto \theta^I$ ,  $u_I \mapsto O_I$ ; its restriction to a bracket-admissible  $B$  is the  $P_0$  realization  $\Phi_{P_0}^B$ .

| Symbol                                                                   | Meaning                                                                                                                                                                                   |
|--------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1]$ | Shifted cotangent Lie algebra of Hamiltonian canonical transformations.                                                                                                                   |
| $H_I$                                                                    | Hamiltonian generator in $\mathfrak{h}$ .                                                                                                                                                 |
| $\eta^I$                                                                 | Shifted cotangent generator $(H_I)^\vee[1]$ , with $I = (a, b)$ and $a + b > 0$ .                                                                                                         |
| $c^I$                                                                    | Chevalley–Eilenberg coordinate dual to a Hamiltonian generator $H_I \in \mathfrak{h}$ . Under $\Phi_{\text{coord}}$ , $c^I \mapsto \theta^I$ .                                            |
| $u_I$                                                                    | Chevalley–Eilenberg coordinate dual to the shifted cotangent generator. Under $\Phi_{\text{coord}}$ , $u_I \mapsto O_I$ . Boundary Hamiltonian observables come from $u_I$ .              |
| $\theta^I$                                                               | Polyvector coordinate corresponding to $c^I$ under the coordinate CE/PV isomorphism.                                                                                                      |
| $O_I$                                                                    | Hamiltonian observable coordinate corresponding to $u_I$ under the coordinate CE/PV isomorphism.                                                                                          |
| $\Phi_{\text{coord}}$                                                    | Coordinate dg CE/PV map determined on generators by $c^I \mapsto \theta^I$ and $u_I \mapsto O_I$ . Its $P_0$ restriction is $\Phi_{P_0}^B$ only after a bracket-admissible $B$ is chosen. |

*Matlis principal-part labels.* Algebraic side: the Grothendieck-dual of the formal jet algebra  $\mathfrak{h}$ , namely the residue/Matlis principal-part module  $\mathcal{P} \cong \mathfrak{h}_{\text{cont}}^\vee$ . Physical side: the principal-part current observable  $\Theta_\rho(B)$  on a brane interval. Comparison morphism: the residue pairing  $\mathfrak{h} \otimes \mathcal{P} \rightarrow \mathbb{C}$ ,  $f \otimes \rho \mapsto \text{Res}_0(f\rho)$ , with Hamiltonian coadjoint action on  $\mathcal{P}$  its image. This is the formal-holomorphic locality (iii) of Theorem 0.1.

| Symbol                         | Meaning                                                                                                                                                                                                       |
|--------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $\mathcal{P}$                  | Reduced principal-part module $\text{Ann}_{\text{res}}(\mathbb{C} \cdot 1) \subset H_{\mathfrak{m}}^2(R) dz_1 dz_2$ , identified with $\mathfrak{h}_{\text{cont}}^\vee$ .                                     |
| $\rho_{a,b}$                   | Principal-part residue current $z_1^{-a-1} z_2^{-b-1} dz_1 dz_2$ , $a + b > 0$ , with the Matlis coadjoint action.                                                                                            |
| $z_1^p z_2^q \cdot \rho_{a,b}$ | Coadjoint action $-(pb - qa + p - q)\rho_{a-p+1, b-q+1}$ . This is the Hamiltonian coadjoint principal-part action. The Matlis $R$ -module action and the PBW action on $\Psi_{a,b}$ are separate structures. |
| $\mathcal{P}_I$                | $\mathcal{P}_I = \mathcal{D}'_c(I) \widehat{\otimes} \mathcal{P}$ , the compactly supported distributional principal-part currents on an interval $I$ .                                                       |

As stated in the separation rule above,

$$\Psi_{a,b} \in \text{PBW special-fibre open homology}, \quad \rho_{a,b} \in \mathcal{P} \cong \mathfrak{h}_{\text{cont}}^\vee$$

live in different modules. The only comparison morphism between them is the Fourier–Rees label bridge  $F_{\text{Rees}}$  of Theorem 8.12, which acts on associated-graded labels through Fourier-twisted polarization. The Hamiltonian coadjoint action remains the action on the Matlis module.

*Interval-current and boundary labels.* Algebraic side: the brane-line objects produced by the support-locality differential of Theorem 0.1 (i). Physical side: the smeared brane measurement  $B_f(a)$  and the principal-part current observable  $\Theta_\rho(B)$ . Comparison morphism: the de Rham-current contraction  $u_{a(t)dt \otimes f} \mapsto B_f(a)$ ,  $c_{B,\rho} \mapsto \Theta_\rho(B)$ ; a smooth test function multiplies a distribution.

| Symbol                       | Meaning                                                                                                                                                                                                                                                                                                                                                          |
|------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $\mathfrak{h}_I$             | $\mathfrak{h}_I = \Omega_c^\bullet(I) \widehat{\otimes} \mathfrak{h}$ , the compactly supported de Rham Hamiltonian currents on an interval $I \subset \mathbb{R}$ . Degree-zero test functions appear only after taking the explicit measurement notation.                                                                                                      |
| $\mathfrak{g}_I$             | $\mathfrak{g}_I = \mathfrak{h}_I \ltimes \mathcal{P}_I[1]$ , the shifted cotangent current algebra on $I$ , with principal-part currents in the shifted cotangent direction.                                                                                                                                                                                     |
| $B_f(a)$                     | Smeared boundary Hamiltonian observable $\int_I a(t) \overline{\text{Tr}} f(\phi_1(t), \phi_2(t)) dt$ . Its CE source is $u_{a(t)dt \otimes f}$ .                                                                                                                                                                                                                |
| $A_\partial^{\text{Ham}}(I)$ | Reduced Hamiltonian brane-line $P_o$ factorization algebra $\widehat{\mathbf{S}}(\mathfrak{h}_I)$ .                                                                                                                                                                                                                                                              |
| $A_\partial^{\text{PP}}(I)$  | Reduced principal-part defect-current model after de Rham-current contraction $\widehat{\mathbf{S}}(\mathfrak{h}_I) \widehat{\otimes} \widehat{\mathbf{S}}(\mathcal{P}_I[1])$ . An unreduced BV observable algebra maps to it only after the centrality homotopies, coefficient kernel, brane-defect propagator, descent data, and BV/QME solution are supplied. |
| $\Theta_\rho(B)$             | Principal-part current observable with $B \in \mathcal{D}'_c(I)$ and $\rho \in \mathcal{P}$ . It is the boundary notation for $B \otimes \rho[1]$ in the reduced model and pairs with smooth test functions by $B(a) \text{Res}(\rho f)$ .                                                                                                                       |
| $aB$                         | Multiplication of a compactly supported distribution $B$ by a smooth test function $a \in \Omega_c^0(I)$ . The reduced brackets use this operation and never require multiplying two distributions.                                                                                                                                                              |

### Conventions and notation for the local theorem

The convention layer below binds throughout the body, the appendices, and every cross-target statement; deviation requires an explicit matched-conventions theorem (Theorem 10.13, §10.7).

*Geometric and algebraic conventions.* The complex target dimension is  $d = \dim_{\mathbb{C}} X$ ; the manuscript proves the formal-Darboux stalk identification  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b \simeq Z_{E_1}^{\text{der}}(A_{\partial,N}^{\text{cl}})$  (Definition 1.1, Theorem 5.6) at the Dirac brane stack on  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}_{\text{o,hol}}^2$ , with  $d = \dim_{\mathbb{C}} \widehat{\mathbb{C}}_{\text{o}}^2 = 2$ . The local geometry is  $\mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$  at a brane: the only Calabi–Yau datum is the trivial holomorphic symplectic  $\Omega_{\text{loc}} = dz_1 \wedge dz_2$  on  $\widehat{\mathbb{C}}_{\text{o}}^2$ , used for the BV pairing, divergence-free Hamiltonian fields, and cyclic trace densities of Definition C.1. Compact CY<sub>3</sub>, BCOV, quintic, Ooguri–Strominger–Vafa (OSV), Gopakumar–Vafa (GV), Maulik–Nekrasov–Okounkov–Pandharipande (MNOP), Abel–Jacobi, cohomological Hall algebra (CoHA), Igusa, and Borchers–Kac–Moody (BKM) are external programmes governed by Theorem 10.4 and the matched-conventions acceptance class  $\text{Ob}_{\text{UKD}}(\mathfrak{C}_{\text{tar}})$ . The native bulk object is the holomorphic  $E_2$  / factorization algebra on  $\mathbb{C}^2$ ,  $\Omega_c^{\circ,\bullet}(B) \widehat{\otimes} \mathfrak{g}$ , with shifted-cotangent BF Lie algebra  $\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^{\vee}[1]$ ,  $\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$  (§2, Theorem 2.11). The native bulk object is the two-complex-dimensional factorization algebra on  $\mathbb{C}^2$ . A curve vertex operator algebra (VOA) is a separate one-dimensional object. It can be compared with the native factorization algebra only after a controlled reduction supplies explicit  $z_2$ -mode or principal-part data, pushed-forward bracket, BV pairing, brane image, and anomaly matching. Only then do the Zhu algebra and the Vol. II  $\mathbb{C} \times \mathbb{R}$  chiral-topological theorems apply to the restricted curve theory (Remark 1.25). The worldsheet  $\Sigma$  is a complex curve, possibly with boundary or marked points; framing on  $S^3$  is stated when invoked, with deviation flagged.

*Sign, degree, and renormalization conventions.* BV degree, ghost number, form degree, Koszul / CE / PV / Gerstenhaber bracket signs, cyclic trace, CE/PV cochain comparison, and the heat-kernel propagator  $(K_L - K_{\epsilon})$  are fixed in Appendix C and Appendix B. The QFT loop  $\hbar$  controls the local BV/Moyal expansion; the string-genus  $g_s$  belongs to the ambient genus expansion. The local theorem uses  $\hbar$ . Compatibility with the BCOV holomorphic-anomaly equation is the matched-conventions statement of Theorem 10.21.

*Homological taxonomy.* Hochschild homology  $HH_{\bullet}$ , cyclic homology, and negative-cyclic homology  $HC_{\bullet}^{-}$  are distinguished at every reference (Appendix C): the cyclic trace symbol  $\text{Tr}$  is the graded cyclic quotient, the Hochschild trace is its image under the Connes  $B$ -operator, and the negative-cyclic class is the Calabi–Yau pairing on the target category. Drinfeld centre  $Z(C)$ , derived centre  $Z^{\text{der}}$ , and chiral derived centre  $Z_{\text{ch}}^{\text{der}}$  are distinguished: the bar construction  $\text{Bar}(A)$  is the twisting coalgebra, while the bulk is the derived chiral centre. For a chiral algebra  $A$  on a smooth curve  $C$  the  $d = 1$  chiral derived centre is  $Z_{\text{ch}}^{\text{der}}(A) \simeq C_{\text{ch}}^{\bullet}(A, A)$  by Beilinson–Drinfeld [28, §4]. At  $d = 2$  the analogous identification of the bulk with the chiral derived centre of the boundary open category  $\mathcal{C}_{\partial}^{\text{op}}$  is the Mixed Holomorphic–Topological Deligne conjecture; this manuscript proves the formal-Darboux stalk identification  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b \simeq Z_{E_1}^{\text{der}}(A_{\partial,N}^{\text{cl}})$  of that conjecture at the brane vacuum. Three distinct algebraic structures appear:

- (i) the bulk chiral  $E_2$  factorization algebra  $\mathcal{A}_{\text{bulk}}^{\text{cl}}$  on  $\mathbb{C}^2$ ;
- (ii) the formal-Darboux stalk  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b$  at the brane vacuum, an  $E_2$ -algebra carrying the chiral structure restricted to the formal neighbourhood of  $b$ ;
- (iii) the boundary  $E_1$ -brane algebra  $A_{\partial,N}^{\text{cl}}$  along the worldline  $\mathbb{R}_t$ , whose derived  $E_1$ -centre  $Z_{E_1}^{\text{der}}(A_{\partial,N}^{\text{cl}}) \simeq HH^{\bullet}(A_{\partial,N}^{\text{cl}}, A_{\partial,N}^{\text{cl}})$  is an  $E_2$ -algebra by Deligne’s  $E_1$ -Hochschild theorem.

The chart algebra  $A_b = \text{End}_{\mathcal{C}_{\partial}^{\text{op}}}(b) \cong \mathbb{C}[[z_1, z_2]]$  is a formal coordinate ring. The symbol  $C_{\text{ch}}^{\bullet}(A_b, A_b)$  is meaningful after  $A_b$  is equipped with the chiral structure restricted from  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b$ . The local theorem

of §5.3 proves the formal-Darboux stalk identification  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b \simeq Z_{E_1}^{\text{der}}(\mathcal{A}_{\partial,N}^{\text{cl}})$  by Costello–Gwilliam bulk–boundary correspondence and Deligne’s  $E_1$ -Hochschild theorem; the  $E_2$  structure on both sides matches. The Mixed Holomorphic–Topological Deligne conjecture is the global identification of the bulk chiral  $E_2$  algebra  $\mathcal{A}_{\text{bulk}}^{\text{cl}}$  with the chiral derived  $E_2$ -centre of  $C_{\partial}^{\text{op}}$ ; this manuscript proves the formal-Darboux stalk identification  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b \simeq Z_{E_1}^{\text{der}}(\mathcal{A}_{\partial,N}^{\text{cl}})$  of that conjecture at one Dirac brane vacuum. Where  $C_{\text{ch}}^{\bullet}(A_b, A_b)$  appears below,  $A_b$  carries this restricted chiral structure and the symbol denotes  $Z_{E_1}^{\text{der}}(\mathcal{A}_{\partial,N}^{\text{cl}})$  under Costello–Gwilliam.

*Constellation-anchored objects.* The boundary open category on the Kato–Nakayama (KN) log boundary  $\partial X^{\text{KN}}$  is  $C_{\partial}^{\text{op}} = \Phi_d^{\text{FA}}(\mathcal{A})|_{\partial X^{\text{KN}}}$ , the restriction of the bulk factorisation algebra to the boundary in the sense of Costello–Gwilliam. Factorisation-homology specialisation along a  $(d-1)$ -manifold  $\Sigma_{d-1}$  and a curve  $C$  is  $\text{Sp}_{\Sigma_{d-1},C}^{\text{ch}}(\mathcal{F}) = \int_{\Sigma_{d-1}} \mathcal{F}|_C$  (Lurie–Ayala–Francis). The Dirac brane stack at  $N$  branes is the derived commuting variety  $[\mu_{\text{der}}^{-1}(\mathfrak{o})/\text{GL}_N]$  with Koszul differential  $Q\phi_i = \mathfrak{o}, Q\psi = [\phi_1, \phi_2]$  resolving the moment-map ideal of conjugation on  $\mathfrak{gl}_N \oplus \mathfrak{gl}_N$ . The Capelli scalar identification  $\hbar N[\bar{c}] = \Omega_{\text{central}}|_{J(f)}$  is the projective curvature evaluation at the trace generator (Theorem 6.8). The modular line bundle on  $\overline{\mathcal{A}}_g$  coincides with  $\Omega_{\text{central}}$  at level  $g$ ; the Igusa cusp form  $\Phi_{10}$  is its level-2 section;  $\Delta_5 = \Phi_{10}^{1/2}$  denotes a chosen square-root branch on the paramodular cover (its operator-level lift on  $K_3 \times E$  is the open problem of §10.3). The CE/PV dictionary  $c_f \mapsto \theta_f, u_f \mapsto J(f)$  of Lemma C.2 is the Koszul resolution of the brane-side derived  $E_1$ -Hochschild  $HH^{\bullet}(\mathcal{A}_{\partial,N}^{\text{cl}}, \mathcal{A}_{\partial,N}^{\text{cl}})$  at the chosen Dirac brane vacuum  $b_p \in C_{\partial}^{\text{op}}$  with chart algebra  $A_{b_p} = \text{End}_{C_{\partial}^{\text{op}}}(b_p)$  (Theorem 5.146); the Mixed Holomorphic–Topological Deligne conjecture identifies this at the formal-Darboux stalk with the bulk  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_{b_p}$  (the chiral  $E_2$  factorization algebra restricted to the formal neighbourhood of  $b_p$ ).

*Theorem-control predicates.* Every theorem in the manuscript is stated within the following control surface:

- (i) Native  $\mathbb{C}^2$  holomorphic  $E_2$  / factorization-algebra taxonomy is retained before any curve-chiral reduction; a curve VOA result is invoked only after a controlled reduction with pushed-forward bracket, brane image, and anomaly matching.
- (ii) BMK lane: the one-pair analytic pro-Matlis retract of Theorem 8.42 is the pro-Matlis replacement for strict all-window support-local current transfer; the obstruction on every fixed finite Matlis window is  $\text{Ob}_{\text{BM}}^{\Pi}$  (Theorem 8.40), and the pro-tower  $\widehat{\mathcal{P}}_{\Pi} = \varprojlim_N \mathcal{P}_{\leq N}$  is the target replacement that absorbs that obstruction.
- (iii) Larger non-scalar  $\theta_3$  row is evidence only with one of: a Chevalley–Eilenberg ancestor, a scalar-zero Costello local counterterm, or a complete companion-face table (Theorem E.38); tower compatibility passes through  $\Delta_{M,N}^{\text{I}} = -\pi_{M,N} \mathfrak{b}^M + \mathfrak{b}^N$  together with the secondary  $\varprojlim^{\text{I}} H^{\circ}$  primitive class.
- (iv) The radial / Weyl theorem surface is  $\Omega_{a,b}^{\text{rad}} \in \text{coker } B_{a,b}$ , equivalently the decorated PBW Stokes obstruction for  $D_{a,b}^{\square} = C_{a,b}^{+} \partial_2$ , with obstruction exactly a signed row  $(\phi, -\lambda) \in \ker B_{a,b}^{*}$  such that  $\lambda(E_{a,b}^{+}) - \phi(T_{a,b}) \neq \mathfrak{o}$  (Theorem F.44).
- (v) Larger non-scalar Costello / QME theorem requires the data tuple: filtered scalar projection  $\widehat{\sigma}_{\text{sc}}$ , finite row arrays, primitive matrix  $A^M c = -r^M$ , transition matrices, Roos compatibility, centrality homotopies, and the curved bulk-to-defect kernel (Theorem E.53).



- (vi) The brane-preserving  $\Omega$ -background uses the normal scaling on  $N_L X = \mathbb{R}_s \oplus \mathbb{C}_{z_1} \oplus \mathbb{C}_{z_2}$  with the brane time  $t$  fixed, torus  $T_\Omega = \mathbb{C}_{\varepsilon_s}^* \times \mathbb{C}_{\varepsilon_1}^* \times \mathbb{C}_{\varepsilon_2}^*$ , twisted differential  $Q_\Omega = Q + \iota_{V_\Omega}$  with  $Q_\Omega^2 = L_{V_\Omega}$ , inverted normal weights, residue-vs-Euler normalization, and stratified factorization data; literal  $(t, s)$ -rotation changes the brane fixed locus and belongs to a different fixed-locus problem.

*Convention firewall.* Transport of the local theorem to a compact-target programme (BCOV / Kodaira–Spencer; CY-to-chiral; Igusa / BKM; GW / DT / PT / MNOP; OSV; mixed-Dunn / six-relation; global Hamiltonian descent on a compact holomorphic-symplectic surface) is the matched-conventions theorem (Theorem 10.4, Theorem 10.13) with universal Koszul-descent acceptance class  $\text{Ob}_{\text{UKD}}(\mathbb{C}_{\text{tar}})$ ; a nonzero coordinate of  $\text{Ob}_{\text{UKD}}$  is the obstruction to acceptance.

## I THE SETUP

### 1.1 The Dirac brane formal-stalk chart in one calculation

*Definition 1.1 (Formal-Darboux stalk chart at a brane vacuum).* A brane vacuum at a closed point  $p \in \mathbb{C}^2$  is the object  $b_p$  of the open category  $\mathcal{C}_\partial^{\text{op}} = \Phi_d^{\text{FA}}(\mathcal{A})|_{\partial X^{\text{KN}}}$  on the Kato–Nakayama log boundary of  $X = \mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$  whose endomorphism algebra is the formal completion

$$A_{b_p} = \text{End}_{\mathcal{C}_\partial^{\text{op}}}(b_p) = \widehat{\mathcal{O}}_{\mathbb{C}^2, p} \cong \mathbb{C}[[z_1, z_2]].$$

The *formal-Darboux stalk chart at  $b_p$*  is the quadruple  $(\widehat{\mathbb{C}^2}_p, \omega_p, A_{b_p}, (z_1, z_2))$  with formal completion  $\widehat{\mathbb{C}^2}_p = \text{Spf } \widehat{\mathcal{O}}_{\mathbb{C}^2, p}$ , holomorphic symplectic form  $\omega_p = dz_1 \wedge dz_2$ , chart algebra  $A_{b_p}$ , and formal Darboux coordinates  $(z_1, z_2)$  supplied by the formal holomorphic Darboux theorem. The *Dirac brane formal-Darboux stalk chart at  $N$  branes over  $b_p$*  adjoins the Chan–Paton substitution

$$\rho_\phi: A_{b_p} \rightarrow M_N(\mathbb{C}), \quad z_i \mapsto \phi_i \in \mathfrak{gl}_N, \quad [\phi_1, \phi_2] = 0:$$

the  $\mathfrak{gl}_N$ -valued pair  $(\phi_1, \phi_2) \in \mathfrak{gl}_N \oplus \mathfrak{gl}_N$  is the matrix coordinate pair of the  $N$ -brane stack on the holomorphic-symplectic factor (its transverse position), and Dirac/moment-map reduction imposes  $[\phi_1, \phi_2] = 0$ . Affine Kac–Moody, free  $\beta\gamma$ , Virasoro, and  $W_N$  vertex algebras are curve chiral algebras which may be compared with the Hamiltonian formal disk only after a separate curve reduction and anomaly matching. For a factorization algebra  $\mathcal{X}$  on  $\mathbb{R}_t^2 \times \mathbb{C}_{z_1, z_2}^2$ , the *formal-Darboux stalk of  $\mathcal{X}$  at  $p$*  is the restriction  $\mathcal{X}|_{\mathbb{R}_t^2 \times \widehat{\mathbb{C}^2}_p}$  in these coordinates.

A Hamiltonian  $f \in A_{b_p}$  on the formal symplectic normal polydisk has trace character

$$J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)},$$

a single  $GL_N$ -invariant number on the brane stack. The supporting geometry is the product

$$X = \mathbb{R}_t \times \mathbb{R}_s \times \mathbb{C}_{z_1, z_2}^2,$$

formally completed at the origin of  $\mathbb{C}^2$ . Each factor carries one ingredient of the calculation: the worldline  $\mathbb{R}_t$  carries the open-line observable algebra, the transverse direction  $\mathbb{R}_s$  carries the bulk-to-boundary propagator, and the formal holomorphic-symplectic normal disk  $\widehat{\mathbb{C}^2}_o$  carries the Hamiltonian algebra  $\mathfrak{h}$ .

Replacing the formal completion  $\widehat{\mathbb{C}^2}_o$  by a global holomorphic-symplectic surface  $Y \supset \widehat{Y}_p$  transports the local theorem to that ambient datum after matched-conventions descent (§10.7).

A stack of  $N$  Dirac branes on  $L = \mathbb{R}_t \times \{s = z_1 = z_2 = o\} \subset X = \mathbb{R}_t \times \mathbb{R}_s \times \mathbb{C}^2$  is a Chan–Paton matrix substitution

$$\rho_\phi: \mathcal{A} = \mathbb{C}[[z_1, z_2]] \longrightarrow \text{End}(\mathbb{C}^N), \quad z_i \longmapsto \phi_i \in \mathfrak{gl}_N,$$

an algebra homomorphism modulo the Dirac ideal  $[\phi_1, \phi_2] = o$  imposed below. The holomorphic symplectic form  $dz_1 \wedge dz_2$  pulls back along this substitution and traces to

$$\theta_N = \text{Tr}(\phi_1 d\phi_2), \quad \Omega_N = \text{Tr}(d\phi_1 \wedge d\phi_2),$$

and  $GL_N$ -conjugation acts symplectically with moment map  $\mu(\epsilon) = -\text{Tr } \epsilon[\phi_1, \phi_2]$ , so Dirac reduction imposes the first-class constraint  $[\phi_1, \phi_2] = o$ :  $\rho_\phi$  factors through the commuting variety, and the simultaneous spectrum of  $(\phi_1, \phi_2)$  is the configuration of  $N$  points in the polydisk. The trace character map at the Chan–Paton brane  $\rho_\phi$  is

$$J: \mathfrak{h} \rightarrow \mathcal{A}_\partial = \widehat{\mathbf{S}}_{\text{Tate}}(\mathfrak{h}), \quad J(f) = \overline{\text{Tr } \rho_\phi(f)} = \overline{\text{Tr } f(\phi_1, \phi_2)}, \quad \mathfrak{h} = \mathcal{A}/\mathbb{C} \cdot 1,$$

the gauge-invariant trace character of a Hamiltonian  $f \in \mathcal{A}$ . The brane represents the commutative coordinate ring  $\mathcal{A}$ ; the Hamiltonian Lie algebra  $\mathfrak{h}$  acts on the representation moduli  $\text{Rep}_N(\mathcal{A})//GL_N$  by Hamiltonian vector fields.

For polynomial  $f \in \mathcal{A}_{\text{poly}} = \mathbb{C}[z_1, z_2]$  the substitution  $f(\phi_1, \phi_2) \in \text{End}(V)$  is defined on the commuting variety  $\text{Rep}_N(\mathcal{A}_{\text{poly}}) = \{(\phi_1, \phi_2) \in \mathfrak{gl}_N^2 : [\phi_1, \phi_2] = o\}$  and gives the polynomial trace map  $J_{\text{poly}}: \mathcal{A}_{\text{poly}}/\mathbb{C} \cdot 1 \rightarrow \mathcal{O}(\text{Rep}_N(\mathcal{A}_{\text{poly}}))^{GL_N}$ . For formal  $f \in \mathcal{A} = \mathbb{C}[[z_1, z_2]]$  the substitution lives on the formal representation functor

$$\text{Rep}_N^\wedge(\mathcal{A})(R) = \{(\phi_1, \phi_2) \in (\mathfrak{m}_R \otimes \text{End} V)^2 : [\phi_1, \phi_2] = o\}, \quad R \text{ Artin local over } \mathbb{C},$$

on which every formal power series in  $\phi_1, \phi_2$  terminates modulo  $\mathfrak{m}_R^k$ ; the formal trace map is  $J: \mathfrak{h} \rightarrow \mathcal{O}(\text{Rep}_N^\wedge(\mathcal{A}))^{GL_N}$ , and the Tate completion  $\mathcal{A}_\partial = \widehat{\mathbf{S}}_{\text{Tate}}(\mathfrak{h})$  records its image at the universal Artin base. The polynomial commuting variety  $\text{Rep}_N(\mathcal{A}_{\text{poly}}) \subset \mathfrak{gl}_N^2$  is the uncompleted algebraic model; the formal functor  $\text{Rep}_N^\wedge(\mathcal{A})$  is its formal completion at the closed point  $\{\phi_1 = \phi_2 = o\}$  inside the commuting variety, and on polynomial Hamiltonians the two evaluations agree.

An odd matrix antifield  $\psi \in \mathfrak{gl}_N[1]$  and BV differential  $Q$  with  $Q\phi_i = o$ ,  $Q\psi = [\phi_1, \phi_2]$  realize the moment-map ideal as the Koszul derived zero fibre, and the matrix entries of  $[\phi_1, \phi_2]$  have syzygies for  $N \geq 2$ . The open sector is the derived zero fibre of the conjugation moment map (§3.4). The closed BV sector takes its fields in the shifted-cotangent extension

$$\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1].$$

A Hamiltonian  $f \in \mathfrak{h}$  carries two Chevalley–Eilenberg (CE) coordinates on  $\mathfrak{g}$ : the generator-dual  $c_f$  acting as the Hamiltonian Schouten polyvector (PV) field  $\theta_f$  on the representation moduli, and the cotangent-dual  $u_f$  mapping to the boundary observable  $O_f = J(f)$  on the brane; the CE/PV cochain dictionary on generators is

$$c_f \longmapsto \theta_f, \quad u_f \longmapsto O_f$$

(Theorem 4.20). The residue pairing realizes the cotangent module  $\mathfrak{h}_{\text{cont}}^\vee$  as the Grothendieck–Matlis principal-part module

$$\mathcal{P} = \bigoplus_{a+b>0} \mathbb{C} \rho_{a,b}, \quad \rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2,$$

with Taylor Hamiltonians paired by the residue  $\langle f, \rho_{a,b} \rangle = \mathbf{Res}_0(f \rho_{a,b})$  detecting the  $z_1^a z_2^b$ -coefficient of  $f$  (Theorem 5.178); the residues  $\rho_{a,b}$  are the cotangent momenta conjugate to the trace-map evaluation of Taylor coefficients, distinct from the polynomial PBW shadows of the open one- $\psi$  sector.

*Example 1.2 (Single-point evaluation,  $N = 1$ ).* For  $N = 1$  the matrices commute automatically, and  $J(f) = f(\phi_1, \phi_2)$  is the evaluation map  $\mathbb{C}[[z_1, z_2]] \rightarrow \mathbb{C}$  at the point  $(\phi_1, \phi_2) \in \mathbb{C}^2$ . The brane is one point of the polydisk; Procesi–Razmyslov degenerates to the polynomial ring on two scalar variables, and  $J$  is the classical evaluation homomorphism.

*Example 1.3 (Linear Hamiltonians and the scalar anomaly).* For  $f = z_1, g = z_2$ , the unreduced brane Poisson bracket is

$$\{\text{Tr } \phi_1, \text{Tr } \phi_2\} = N.$$

This scalar is the open-side image of the closed–open mismatch Lie 2-cocycle  $\bar{c} \in Z_{\text{Lie}}^2(\bar{A}, \mathbb{C})$ , the cohomology class  $[\bar{c}] \in H_{\text{Lie}}^2(\bar{A}, \mathbb{C})$  attached to the constant Hamiltonian line that vanishes closed-side and survives open-side as the brane centre of mass (Lemma 6.2, Theorem 6.3). Moyal quantization with Weyl-symmetric quantization map  $\Phi_{\hbar}$  and Kontsevich–Moyal star  $\star$  returns the same scalar at order  $\hbar$ ,  $[\Phi_{\hbar}(z_1), \Phi_{\hbar}(z_2)]_{\star} = \hbar N \cdot I$  in  $M_N(\mathbb{C})$ ; the first scalar-reduced  $\mathfrak{gl}_N$  quantum-master-equation obstruction is the same class shifted by  $\hbar$ ,  $\mathcal{W}_1^{\text{sc,ord}} = \hbar N [\bar{c}]$  (Theorem E.13).

*Example 1.4 (A single Matlis residue).* The Grothendieck–Matlis principal-part

$$\rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2 \in \mathcal{P} \subset H_{\mathfrak{m}}^2(\mathbb{C}[[z_1, z_2]]) dz_1 dz_2$$

in the local-cohomology module  $H_{\mathfrak{m}}^2 = H_{\{\mathfrak{o}\}}^2$  at the maximal ideal  $\mathfrak{m} = (z_1, z_2)$  pairs with  $f \in \mathfrak{h}$  by  $\langle \rho_{a,b}, f \rangle = \text{Res}_0(\rho_{a,b} f)$  and detects the Taylor coefficient of  $z_1^a z_2^b$  in  $f$ . The residue-coadjoint formula  $z_1^p z_2^q \cdot \rho_{a,b} = -(pb - qa + p - q) \rho_{a-p+1, b-q+1}$  of Theorem 5.178 gives a non-locally-nilpotent action on  $\mathcal{P}$ . A polynomial PBW class attached to the same  $(a, b)$ -bidegree exists in the open one- $\psi$  sector with a strictly different  $\mathfrak{h}$ -module structure (Corollary 5.172); the residue mode here and that polynomial class are separated by the Fourier-polarized Rees bridge.

### 1.2 Mixed holomorphic-topological strings

A *local string theory* in the sense of Costello–Gwilliam [15] is a classical BV theory whose local observables form a factorization algebra equipped with a cyclic open BV theory on a brane and a bulk-to-boundary central operation. The *topological* qualifier names a factorization algebra satisfying the topological locality of Proposition 1.34: every translation acts null-homotopically. The *holomorphic* qualifier names a factorization algebra satisfying the holomorphic locality: every antiholomorphic translation acts null-homotopically. The protected algebraic content under Theorem 1.35 is the classical and quantum master equations, locality, and factorization associativity; the topological and holomorphic qualifiers refine this content by constraining how local operators vary across the chosen spacetime directions. The word *string* appears here in this local string-field-theoretic sense, attached to the BV / factorization data; global theories of worldsheet moduli or amplitudes require additional target data.

Gwilliam–Williams [5] formulate a holomorphic string in the Costello BV/factorization formalism, with Dolbeault fields  $(\gamma, \beta, c, b)$ , one-loop quantization in the critical complex dimension, and a factorization algebra recovering BRST cohomology. The model here is a formal local target-space sector with de Rham/Floer directions, a holomorphic-symplectic formal polydisk, Hamiltonian cotangent BF fields, and a brane large- $N$  operator algebra.

*Definition 1.5 (Local topological factorization sector; forced by item (2) of Proposition 1.34).* Let  $M$  be a smooth manifold. A local topological closed-string sector on  $M$  is a classical BV theory whose local observables form a locally constant factorization algebra on  $M$ : for inclusions of sufficiently small disks  $U \subset V$ , the structure map

$$\mathrm{Obs}^{\mathrm{cl}}(U) \longrightarrow \mathrm{Obs}^{\mathrm{cl}}(V)$$

is a quasi-isomorphism, equivalently every translation  $v \in TM$  acts on observables by a homotopy  $L_v = [Q, \iota_v]$ . Attaching branes means specifying a cyclic open BV theory and the bulk-to-boundary central-operation data it carries. In a de Rham model the closed fields carry  $d_M$  plus the BV linearised differential; for pairwise disjoint disks  $U_1, \dots, U_k \subset V$ , the locally constant factorization product  $\otimes_i \mathrm{Obs}^{\mathrm{cl}}(U_i) \rightarrow \mathrm{Obs}^{\mathrm{cl}}(V)$  is the operator product on disjoint supports, depending only on the isotopy class of the embedding by the translation homotopy  $L_v = [Q, \iota_v]$ ; equivalently the Lurie–Costello–Gwilliam  $E_n$ -algebra structure on the value at a single disk.

*Remark 1.6 (Canonical completion).* Locally constant factorization is exactly the data of a topological sector in the Lurie–Costello–Gwilliam sense, and the homotopy  $L_v = [Q, \iota_v]$  is the constructive content of locality item (2). Strict translation invariance, or imposing a metric, breaks compatibility with that homotopy and is therefore excluded. The exact A-model factor used below is the self-Floer model for  $\mathbb{R} \subset T^*\mathbb{R}$ , represented by the de Rham complex  $\Omega^\bullet(\mathbb{R})$ ; its local observables are insensitive to the metric of the interval.

*Definition 1.7 (Local holomorphic factorization sector; forced by item (3) of Proposition 1.34).* Let  $Y$  be a complex manifold. A local holomorphic sector on  $Y$  is a classical BV theory whose fields are Dolbeault complexes on  $Y$ , whose differential has the form  $\bar{\partial} + Q_\circ$  with  $Q_\circ$  holomorphic, and whose interactions are holomorphic polydifferential operators. Its local observables form a holomorphic factorization algebra: on coordinate polydisks the factorization products and single-open structure maps are represented by holomorphic polydifferential operations, and every antiholomorphic translation  $\partial/\partial \bar{z}_i$  acts null-homotopically through  $L_{\partial/\partial \bar{z}_i} = [Q, \iota_{\partial/\partial \bar{z}_i}]$ . Branes are cyclic open BV theories with bulk-to-boundary operations.

*Remark 1.8 (Canonical completion).* The continuous-dual data attached to a brane point is the principal-part module  $\mathcal{P}$  of §1.6, forced by Theorem 5.178; the Dolbeault presentation, with the homotopy  $L_{\partial/\partial \bar{z}_i} = [Q, \iota_{\partial/\partial \bar{z}_i}]$ , is what locality item (3) constructs. The active geometry below is  $\mathbb{R}_{\mathrm{top}}^2 \times \widehat{\mathbb{C}}_{\mathrm{o}, \mathrm{hol}}^2$ . BCOV/Kodaira–Spencer [1] supplies the source conventions: Dolbeault polyvectors, cotangent BV notation, the divergence determined by a holomorphic volume form. Here the holomorphic-symplectic form  $dz_1 \wedge dz_2$  and the volume form  $dz_1 dz_2$  provide the BV pairing, divergence operator, and cyclic trace densities; divergence-free vector fields are locally Hamiltonians modulo constants.

*Remark 1.9 (Gwilliam–Williams sector vs. the local target-space sector).* The classical local observable theory of Gwilliam–Williams is a pure holomorphic BV/factorization sector on a Riemann surface, with worldsheet ghost, BRST, and critical-dimension quantum structure. Its critical-dimension and BRST-cohomology theorems live on the worldsheet; the model below is the local target-space sector  $\mathbb{R}^2 \times \mathbb{C}^2$ ,

with holomorphic factor Hamiltonian BF (the holomorphic-symplectic Kodaira–Spencer analogue) and topological factor de Rham/Floer. The critical complex dimension is a worldsheet ghost-anomaly statement for the holomorphic bosonic string; the local  $\mathbb{R}^2 \times \mathbb{C}^2$  Hamiltonian sector is governed by Definition 1.10.

*Definition 1.10 (Classical local mixed holomorphic-topological sector; the unique factorization algebra compatible with all three locality differentials).* Let  $M$  be a smooth manifold and  $Y$  a complex manifold. A classical local mixed holomorphic-topological closed sector on  $M \times Y$  consists of:

- (i) a sheaf of cochain complexes of fields  $\mathcal{E}$  which, on product coordinate opens  $D \times B$ , has the form

$$\mathcal{E}(D \times B) = \Omega^\bullet(D) \widehat{\otimes} \Omega^{\circ,\bullet}(B, \mathcal{V})$$

for a graded holomorphic vector bundle or formal holomorphic vector space  $\mathcal{V}$ ;

- (ii) a degree- $(-1)$  local BV pairing on  $\mathcal{E}$  and a differential

$$Q = d_M + \bar{\partial}_Y + Q_o,$$

with  $Q_o$  holomorphic over  $Y$  and linear over de Rham forms on  $M$ ;

- (iii) a degree-zero local functional  $I \in \mathcal{O}_{\text{loc}}(\mathcal{E})$ , built from differential operators in the  $M$  directions and holomorphic polydifferential operators in the  $Y$  directions, satisfying  $QI + \frac{1}{2}\{I, I\} = 0$ ;
- (iv) the associated classical observables

$$\text{Obs}^{\text{cl}}(U) = (\widehat{\text{Sym}}(\mathcal{E}_c^!(U)), Q_{\text{obs}}), \quad Q_{\text{obs}} = Q^! + \{I, -\},$$

with the standard factorization product for disjoint supports. Here  $\mathcal{E}_c^!(U)$  is the density-dual cosheaf of fields, and extension by zero gives the covariant structure maps used by the factorization algebra.

These data must satisfy the mixed locality condition: for every inclusion of small disks  $D \subset D'$  in  $M$  and every holomorphic polydisk  $B \subset Y$ , the structure map

$$\text{Obs}^{\text{cl}}(D \times B) \longrightarrow \text{Obs}^{\text{cl}}(D' \times B)$$

is a quasi-isomorphism, and on  $B$  each antiholomorphic translation  $\partial/\partial \bar{z}_i$  acts null-homotopically through a degree- $(-1)$  derivation  $\eta_i$  with  $[Q_{\text{obs}}, \eta_i] = \mathcal{L}_{\partial/\partial \bar{z}_i}$ .

A mixed brane is supported on  $N \times Z$ , with  $N \subset M$  a smooth submanifold and  $Z \subset Y$  a holomorphic submanifold or formal holomorphic neighbourhood of one, specified by an object  $\mathcal{F}$  of the local holomorphic brane category on  $Z$ . Its open fields have the derived form

$$\Omega^\bullet(N) \widehat{\otimes} \text{RHom}_Y(\mathcal{F}, \mathcal{F}) \otimes \text{End}(V)[1],$$

with a cyclic trace. For a smooth brane this recovers the Dolbeault model  $\Omega^\bullet(N) \widehat{\otimes} \Omega^{\circ,\bullet}(Z, \mathcal{W})$ ; for a skyscraper brane the Ext algebra. A full bulk-to-boundary mixed datum requires a morphism of factorization algebras

$$\text{Obs}_{\text{closed}}^{\text{cl}}|_{N \times Z} \longrightarrow Z_{\text{fact}}^{P_o}(\text{Obs}_{\text{open}}^{\text{cl}}),$$

compatible with BV brackets and factorization products. The Hamiltonian model below constructs the degree-zero Hamiltonian current part of this map; the full closed CE/cotangent factorization-centre lift is the obstruction problem Problem 5.48.

*Remark 1.11 (Canonical completion).* Item (1) fixes the field sheaf as  $\Omega^\bullet(D) \widehat{\otimes} \Omega^{\circ,\bullet}(B)$ : de Rham in the topological direction is required by item (2) of Proposition 1.34, Dolbeault in the holomorphic direction by item (3). Item (2) fixes  $Q = d_M + \bar{\partial}_Y + Q_0$ : the first two summands are the differentials whose contraction homotopies build the topological and holomorphic localities, the third encodes the BV interaction. Item (3) imposes the classical master equation, the protected content of Theorem 1.35. Item (4) is the Costello–Gwilliam observable functor, the unique factorization assignment compatible with disjoint-union locality and the BV pairing. Replacing any item violates a named locality differential or a protected algebraic condition.

*Remark 1.12 (Quantum extension).* A full quantum mixed holomorphic-topological string datum is the corresponding  $\hbar$ -adic factorization algebra obtained from a renormalized BV action satisfying the quantum master equation on the canonical weighted-Tate completion  $B_w$  (Theorem 5.57). The componentwise quantum coefficient statement is weighted BV kernel convergence, the balanced scalar-contact zero projection on its completion, reduced weighted Moyal principal-part current brackets, and a degree-zero Hamiltonian trace comparison controlled by the radial/Weyl normal-form criterion. The finite normal-form computations close the balanced diagonal through (6, 6), all non-edge bidegrees of total degree at most 13, and every total-degree-14 case with one component at most 6:

$$(3, 11), (11, 3), (4, 10), (10, 4), (5, 9), (9, 5), (6, 8), (8, 6).$$

The balanced (7, 7) is the first undecided case in the unconditional kernel-correction window. Proposition F.32 closes the one-moving-letter edge strips  $\mathfrak{o}_{2,b} = \mathfrak{o}_{a,2} = 0$ . The all-interior statement is the obstruction problem  $\Omega_{a,b}^{\text{rad}} = [(T_{a,b}, E_{a,b}^+)] \in \text{coker } B_{a,b}$ , whose positive form is the decorated PBW Stokes identity  $D_{a,b}^\square = C_{a,b}^+ \partial_2$ ; obstruction is exactly a signed row  $(\phi, -\lambda) \in \ker B_{a,b}^*$  with  $\lambda(E_{a,b}^+) - \phi(T_{a,b}) \neq 0$ . The finite-window non-scalar closed-open BV criterion is the vanishing problem in  $\mathfrak{Q}_{w,\partial,\hbar}^\bullet(I)$  of Problem 5.219.

A full renormalized quantum mixed h-t string requires Costello graph data outside the local Hamiltonian theorem: the analytic graph-realization datum of Problem 5.212, the quantum master equation in the filtered local-functional complex  $\mathfrak{Q}_{w,\partial,\hbar}^\bullet(I)$  of Problem 5.219, and the all-bidegree radial vanishing  $\Omega_{a,b}^{\text{rad}} = 0$  of the preceding remark.

A pure topological sector is locally constant in all real directions; examples include the A-model local/Floer direction and the Chern–Simons-type open field theory on a topological brane. A pure holomorphic sector is Dolbeault in its complex directions and produces a holomorphic factorization algebra; Gwilliam–Williams’ holomorphic bosonic string is a worldsheet example. A mixed holomorphic-topological sector lives on  $M \times Y$  with locally constant factorization in  $M$  and holomorphic factorization in  $Y$ . The A/B terminology specialises this: the A-side supplies the de Rham direction and the B-side supplies the Dolbeault or formal holomorphic direction.

### 1.3 The local mixed model

The local geometric datum is

$$X = \mathbb{R}_t \times \mathbb{R}_s \times \mathbb{C}_{z_1, z_2}^2, \quad L = \mathbb{R}_t \times \{s = z_1 = z_2 = 0\}, \quad \omega = dz_1 \wedge dz_2,$$

completed at  $L$  and replaced by its formal Darboux normal polydisk  $(\widehat{\mathbb{C}}_0^2, \omega)$ . Each factor is required.

*The minimal RG-stable completion.*  $\mathbb{R}_t$  is the brane worldline: the cyclic open BV theory and its trace observables  $J(f)(t)$  live as currents on it.  $\mathbb{R}_s$  is the topological normal/propagator direction along which BV ghosts move and the moment-map multiplier becomes a one-form. This transverse topological direction supplies the propagator and the bulk-to-brane locality mechanism, since the mixed-locality homotopy  $L_v = [Q, \iota_v]$  of Proposition 1.34 (b) requires a topological direction transverse to the worldline.  $\mathbb{C}^2$  is the formal holomorphic-symplectic normal disk: two complex coordinates supply the Hamiltonian bracket and a Darboux pair. Formal completion at  $L$  is dictated by the trace substitution: the brane carries only the formal holomorphic transverse geometry, because  $\phi_1, \phi_2$  live in  $\mathfrak{gl}_N^2$  and the substitution  $z_i \mapsto \phi_i$  is formal substitution into a power series. Removing the formal completion forces the brane to see irrelevant global geometry and breaks the scalar-reduced trace identification of Lemma 3.1.

The Dirac brane is the formal-stalk chart, fixing the open BV sector and its first-class boundary action. Closed insertions act through central operations in Witten's source convention, fixing the closed-side target as a derived  $P_0$ -centre with reduced Hamiltonian model  $Z_{\text{red,Ham}}^{P_0}$  and dictionary  $c^I \mapsto \theta^I$ ,  $u_I \mapsto O_I$ . Continuous duality is residue duality (Grothendieck), fixing the cotangent coefficient module as the Matlis principal-part module  $\mathcal{P}$ , separated from the polynomial PBW special fibre. Together with the three locality differentials of Proposition 1.34 – support, topological, formal holomorphic – these inputs force a unique factorization-algebra shape on  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}_{\text{o,hol}}^2$ : its dg Lie input must be  $\Omega_c^\bullet(U) \widehat{\otimes} \Omega_c^{\circ,\bullet}(B) \widehat{\otimes} (\mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1])$  on every product open  $U \times B$ , locally constant in  $U$  and holomorphic in  $B$ . Definition 1.13 records this object as  $\mathcal{F}_{\text{Ham}}^{\text{hol}}$  with mixed enhancement  $\mathcal{F}_{\text{Ham}}^{\text{mix}}$ . Compact and global comparisons require separate hypotheses outside the formal-local claim.

*Definition 1.13 (Local Hamiltonian holomorphic factorization algebra).* The unique factorization algebra forced by Dirac, Witten's source convention, Grothendieck residue duality, and the three locality differentials of Proposition 1.34 on the formal Darboux polydisk is

$$\mathcal{F}_{\text{Ham}}^{\text{hol}}.$$

On a holomorphic polydisk  $B \subset \mathbb{C}_{\text{hol}}^2$  its dg Lie input is

$$\Omega_c^{\circ,\bullet}(B) \widehat{\otimes} \mathfrak{g}, \quad \mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1], \quad \mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1,$$

and the formal stalk of observables is the continuous Chevalley–Eilenberg cochain algebra  $C_{\text{CE,cont}}^\bullet(\mathfrak{g})$ . Its identification with Schouten polyvectors is the coordinate-window CE/PV theorem of §5.2; the strict-product  $P_0$  structure requires the admissible completion stated there.

The mixed topological-current enhancement is

$$\mathcal{F}_{\text{Ham}}^{\text{mix}}(U \times B), \quad \mathfrak{g}_{U,B}^{\text{mix}} = \Omega_c^\bullet(U) \widehat{\otimes} \Omega_c^{\circ,\bullet}(B) \widehat{\otimes} \mathfrak{g},$$

for  $U \subset \mathbb{R}_{\text{top}}^2$ . This object is locally constant in  $U$  and holomorphic in  $B$ , realising the topological and formal holomorphic locality differentials of Proposition 1.34.

The open quantum brane avatar is the Weyl/Moyal algebra of the formal symplectic polydisk, together with its finite- $N$  matrix Weyl Hamiltonian reduction and matrix or supermatrix cyclic trace bar-cobar models. The support-local defect/cotangent avatar on an interval  $I \subset \mathbb{R}_{\text{brane}}$  is the principal-part current algebra

$$\mathfrak{h}_I = \Omega_c^\bullet(I) \widehat{\otimes} \mathfrak{h}, \quad \mathcal{P}_I = \mathcal{D}'_c(I) \widehat{\otimes} \mathcal{P}, \quad \mathfrak{g}_I = \mathfrak{h}_I \ltimes \mathcal{P}_I[1],$$

with reduced target

$$A_\partial^{\text{pp}}(I) = \widehat{\mathbf{S}}(\Omega_c^\bullet(I) \widehat{\otimes} \bar{A}) \widehat{\otimes} \widehat{\mathbf{S}}(\mathcal{D}'_c(I) \widehat{\otimes} \mathcal{P}[1]).$$

Its quantum version replaces the classical coadjoint action by the Moyal coadjoint action.

“Chiral” here means this two-complex-dimensional holomorphic factorization algebra and its mixed topological-current extensions. Its native singular product is the two-variable diagonal local-cohomology expansion of Theorem 1.15. A one-dimensional vertex operator algebra (VOA) / one-variable OPE statement requires a separate restriction theorem choosing a complex line or holomorphic defect, a transverse boundary condition or restriction functor, and the finite Laurent OPE expansion on that restricted object.

**PROPOSITION 1.14** (*Constructed Hamiltonian CE/factorization observables*). In the formal local mixed model  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}^2_{\text{o,hol}}$ , the Hamiltonian BF fields of Definition 1.10 reduce at a formal holomorphic insertion to the shifted cotangent dg Lie algebra

$$\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^{\vee}[1], \quad \mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1.$$

For a holomorphic polydisk  $B$ , the constructed holomorphic factorization algebra is represented by the continuous CE observables of the compactly supported Dolbeault dg Lie algebra

$$\mathcal{F}_{\text{Ham}}^{\text{hol}}(B) = C_{\text{CE,cont}}^{\bullet}(\Omega_c^{\circ,\bullet}(B) \widehat{\otimes} \mathfrak{g}).$$

For a product open  $U \times B$ , its mixed topological-current enhancement is represented by

$$\mathcal{F}_{\text{Ham}}^{\text{mix}}(U \times B) = C_{\text{CE,cont}}^{\bullet}(\Omega_c^{\bullet}(U) \widehat{\otimes} \Omega_c^{\circ,\bullet}(B) \widehat{\otimes} \mathfrak{g}).$$

These assignments are the local holomorphic factorization algebras of the paper. Their formal stalk is  $C_{\text{CE,cont}}^{\bullet}(\mathfrak{g})$ , and its identification with the PV model is only the coordinate/admissible theorem cited below.

*Proof.* Proposition 2.7 identifies the divergence-free holomorphic vector fields in the formal polydisk with Hamiltonians modulo constants. The Hamiltonian BF action then has connection field valued in  $\mathfrak{h}[1]$  and cotangent field valued in  $\mathfrak{h}_{\text{cont}}^{\vee}[2]$ . After the de Rham and Dolbeault Poincare contractions at the formal insertion point, Proposition 6.20 identifies the residual closed local operators with  $C_{\text{CE,cont}}^{\bullet}(\mathfrak{g})$ , where the CE differential is dual to the semidirect bracket  $[f, g] = \{f, g\}$ ,  $[f, \eta_{\lambda}] = \eta_{\text{ad}_f^{\vee}} \lambda$ , and  $[\eta_{\lambda}, \eta_{\mu}] = 0$ .

On a polydisk one keeps the compactly supported Dolbeault factor; on a mixed product open one also keeps the compactly supported de Rham factor. Extension by zero supplies the factorization structure maps, while the local Poincare homotopies give holomorphic factorization in  $B$  and local constancy in  $U$ , exactly as in Definitions 1.7 and 1.10. The PV,  $P_0$ , kernel, and bar-cobar upgrades are the admissible consequences of the coordinate CE/PV theorem and the reduced central-operation comparison. The native two-dimensional singular product is the diagonal local-cohomology theorem inserted next; the one-dimensional vertex/OPE structure requires the separate restriction data of Theorem 0.1.  $\square$

### 1.3.1 THE FORMAL TWO-DIMENSIONAL HOLOMORPHIC SINGULAR PRODUCT

The singular product of the local model is a diagonal expansion on  $\mathbb{C}^2 \times \mathbb{C}^2$ . A Laurent expansion on a curve arises after choosing a curve restriction. Let

$$\Delta \subset \mathbb{C}_z^2 \times \mathbb{C}_w^2, \quad \zeta_i = z_i - w_i,$$



and let  $j: (\mathbb{C}^2 \times \mathbb{C}^2) \setminus \Delta \hookrightarrow \mathbb{C}^2 \times \mathbb{C}^2$  be the open complement. The singular coefficient module of a two-dimensional holomorphic singular product is the local-cohomology module

$$\mathcal{P}_\Delta = H_\Delta^2(\mathbb{C}[[z_1, z_2, w_1, w_2]]) d\zeta_1 d\zeta_2 = \bigoplus_{a,b \geq 0} \mathbb{C}[[w_1, w_2]] \rho_{a,b}^\Delta,$$

with

$$\rho_{a,b}^\Delta = \zeta_1^{-a-1} \zeta_2^{-b-1} d\zeta_1 d\zeta_2, \quad \text{Res}_\Delta(\rho_{a,b}^\Delta \zeta_1^m \zeta_2^n) = \delta_{a,m} \delta_{b,n}.$$

The two-dimensional Cauchy kernel is the generator

$$K_\Delta^{(2)}(z, w) = \rho_{0,0}^\Delta = \frac{d\zeta_1 d\zeta_2}{\zeta_1 \zeta_2}.$$

In the region  $|z_i| > |w_i|$  it has the formal expansion

$$K_\Delta^{(2)}(z, w) = \sum_{a,b \geq 0} z_1^{-a-1} z_2^{-b-1} w_1^a w_2^b dz_1 dz_2.$$

This is the two-variable local-cohomology expansion used below. It is the local-cohomology singular expansion of the surface theory. A chiral algebra on a curve is obtained by replacing  $\mathcal{P}_\Delta$  by the one-variable principal parts of a chosen line  $C \hookrightarrow \mathbb{C}^2$ .

For  $x \in \mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1]$ , write  $J_x(z)$  for the linear holomorphic current determined by  $x$  in the CE factorization algebra

$$B \mapsto C_{\text{CE,cont}}^\bullet(\Omega_c^{\circ,\bullet}(B) \widehat{\otimes} \mathfrak{g}).$$

For  $f \in \mathfrak{h}$  and  $\rho \in \mathfrak{h}_{\text{cont}}^\vee$  we also write

$$B_f = J_f, \quad \Theta_\rho = J_{\rho[1]}.$$

**THEOREM 1.15** (*Formal two-dimensional holomorphic singular product*). The singular part of the formal holomorphic product of the Hamiltonian BF factorization algebra on  $\widehat{\mathbb{C}^2}_0$  is

$$J_x(z) J_y(w) \sim K_\Delta^{(2)}(z, w) J_{[x,y]_{\mathfrak{g}}}(w), \quad x, y \in \mathfrak{g}.$$

Equivalently, for Hamiltonians  $f, g \in \mathfrak{h}$  and principal parts  $\rho, \sigma \in \mathfrak{h}_{\text{cont}}^\vee$ ,

$$\begin{aligned} B_f(z) B_g(w) &\sim K_\Delta^{(2)}(z, w) B_{\{f,g\}}(w), \\ B_f(z) \Theta_\rho(w) &\sim K_\Delta^{(2)}(z, w) \Theta_{f \cdot \rho}(w), \\ \Theta_\rho(z) \Theta_\sigma(w) &\sim 0, \end{aligned}$$

where  $(f \cdot \rho)(g) = -\rho(\{f, g\})$ . The multi-index singular products are

$$(J_x)_{(0,0)} J_y = J_{[x,y]_{\mathfrak{g}}}, \quad (J_x)_{(a,b)} J_y = 0 \quad \text{for } (a,b) \neq (0,0),$$

with descendant products obtained by applying the holomorphic translation operators:

$$\partial_z^\alpha J_x(z) J_y(w) \sim (-1)^{|\alpha|} \alpha! \rho_{\alpha, \alpha_2}^\Delta(z, w) J_{[x,y]_{\mathfrak{g}}}(w).$$

*Proof.* The CE factorization algebra is generated by the compactly supported Dolbeault local Lie algebra

$$\Omega_c^{\circ, \bullet}(B) \widehat{\otimes} \mathfrak{g}.$$

For disjoint polydisks the product is extension by zero. When two insertions collide, the only singular operation in the CE differential is the local Lie bracket  $[x, y]_{\mathfrak{g}}$ ; this is the singular binary operation of the shifted-cotangent Lie algebra. The Dolbeault resolution of the diagonal identifies the singular quotient of the product of two holomorphic insertions with

$$H_{\Delta}^2(\mathcal{O}_{\mathbb{C}^2 \times \mathbb{C}^2}) d\zeta_1 d\zeta_2,$$

and the normalized generator is  $K_{\Delta}^{(2)}$ , since its residue on  $\mathbf{1}$  is  $\mathbf{1}$ . Therefore the collision of the two currents is  $K_{\Delta}^{(2)} J_{[x, y]}$ .

Substituting

$$[f, g] = \{f, g\}, \quad [f, \rho[\mathbf{1}]] = (f \cdot \rho)[\mathbf{1}], \quad [\rho[\mathbf{1}], \sigma[\mathbf{1}]] = 0$$

gives the three displayed singular products. The coefficient extraction formula follows from

$$\text{Res}_{\Delta}(\rho_{a,b}^{\Delta} \zeta_1^m \zeta_2^n) = \delta_{a,m} \delta_{b,n}.$$

The descendant identity is

$$\partial_{z_1}^{\alpha_1} \partial_{z_2}^{\alpha_2} \frac{\mathbf{1}}{\zeta_1 \zeta_2} = (-1)^{|\alpha|} \alpha_1! \alpha_2! \zeta_1^{-\alpha_1-1} \zeta_2^{-\alpha_2-1}.$$

Associativity of all iterated two-dimensional singular products is the Jacobi identity in  $\mathfrak{g}$ , and the compatibility with  $Q$  is the Chevalley–Eilenberg identity  $d_{\text{CE}}^2 = 0$ .  $\square$

**COROLLARY 1.16** (*Binary generation of higher singular products*). The iterated  $n$ -point singular product in the strict CE current model is generated by Theorem 1.15, the regular commutative product, and the Chevalley–Eilenberg differential. All arity- $n$  singular products with  $n \geq 3$  are iterated binary collision products.

*Proof.* The coefficient algebra is the CE algebra of the strict dg Lie algebra  $\Omega_c^{\circ, \bullet}(B) \widehat{\otimes} \mathfrak{g}$ . Its only nonlinear structure map is the binary Lie bracket of  $\mathfrak{g}$ ; the arity- $n$  collision maps are therefore the iterated binary collision maps with the Koszul signs of the CE differential. Jacobi is the equality of the two binary collision orders along the small diagonal. This proves both associativity and the absence of independent higher singular coefficients.  $\square$

**COROLLARY 1.17** (*Quantum and brane trace contact term*). After Moyal quantization of the Hamiltonian pair and scalar reduction on the brane,

$$\mathbf{B}_f(z) \mathbf{B}_g(w) \sim K_{\Delta}^{(2)}(z, w) \left( \mathbf{B}_{\{f, g\}_{\hbar}}(w) + \hbar N \omega(f, g) \mathbf{1} \right),$$

where

$$\{f, g\}_{\hbar} = \hbar^{-1}(f * g - g * f) \bmod \mathbb{C} \cdot \mathbf{1}, \quad \omega(f, g) = [\{f, g\}]_0.$$

For  $f = z_1$  and  $g = z_2$ ,

$$\mathbf{B}_{z_1}(z) \mathbf{B}_{z_2}(w) \sim \hbar N K_{\Delta}^{(2)}(z, w) \mathbf{1}.$$

This is the Capelli scalar as a two-dimensional holomorphic contact term.

*Proof.* The degree-zero quantum brane algebra replaces the Poisson bracket by the Moyal commutator. The scalar-reduced Hamiltonian quotient removes the constant term on the closed side, while the Chan–Paton trace sends  $\mathbf{1}$  to  $N$ . Hence the constant Taylor coefficient of the classical bracket contributes the central cocycle  $\omega(f, g)$ , and the quantum normalization multiplies it by  $\hbar N$ . For  $(f, g) = (z_1, z_2)$ ,  $\omega(z_1, z_2) = \mathbf{1}$ , giving the displayed contact term.  $\square$

*Remark 1.18 (Physical reading).* The free BF contraction is

$$\alpha_f(z) \beta_\rho(w) \sim \langle \rho, f \rangle K_\Delta^{(2)}(z, w),$$

with  $\alpha$  the Hamiltonian field and  $\beta$  the shifted cotangent field. The cubic interaction

$$I_{\text{BF}} = \frac{1}{2} \int_{\mathbb{C}^2} \langle \beta, \{\alpha, \alpha\} \rangle dz_1 dz_2$$

turns the free contraction into the current singular product of Theorem 1.15. The quantum brane trace adds precisely the centre-of-mass scalar of Corollary 1.17.

### 1.3.2 THE CLOSED–OPEN FACTORIZATION TRIPLE: $E_2$ BULK, $E_1$ BRANE, DERIVED-CENTRE MAP

The local-string-theoretic content of the model is exhausted by three factorization objects and the morphism between them. All three are constructed from the same dg Lie input  $\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[\mathbf{1}]$  of §2 together with the Dirac brane-stack data  $\text{Kosz}_N = (\mathbb{C}[\mathfrak{gl}_N^{\oplus 2}] \otimes \Lambda^\bullet \mathfrak{gl}_N[\mathbf{1}], Q\psi = [\phi_1, \phi_2])$  of Construction 3.20.

*Definition 1.19 (Closed mixed-HT current factorization algebra).* For a product open  $O = U \times B \subset \mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$ , the compactly supported local dg Lie algebra is

$$\mathcal{L}_{\text{Ham},c}(O) = \Omega_c^\bullet(U) \widehat{\otimes} \Omega_c^{\circ,\bullet}(B) \widehat{\otimes} \mathfrak{g},$$

with differential  $D = d_{\mathbb{R}^2} + \bar{\partial}_{\mathbb{C}^2} + d_{\mathfrak{g}}$  and bracket  $[\alpha \otimes x, \beta \otimes y] = (\alpha \wedge \beta) \otimes [x, y]_{\mathfrak{g}}$ . The closed classical mixed-HT factorization algebra is its CE chain complex,

$$\mathcal{F}_{\text{Ham}}^{\text{cl}}(O) = C_\bullet^{\text{CE}}(\mathcal{L}_{\text{Ham},c}(O)),$$

with factorization product induced by extension by zero  $\bigoplus_i \mathcal{L}_{\text{Ham},c}(O_i) \rightarrow \mathcal{L}_{\text{Ham},c}(O)$  for disjoint  $O_i \subset O$ .

**THEOREM 1.20 (Mixed-HT factorization theorem).** The assignment  $O \mapsto \mathcal{F}_{\text{Ham}}^{\text{cl}}(O)$  is a prefactorization algebra on  $\mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$  satisfying:

- (i) (*Topological locality.*) For every  $v \in T\mathbb{R}^2$ ,  $L_v = [D, \iota_v]$  on  $\mathcal{F}_{\text{Ham}}^{\text{cl}}$ ; hence the assignment  $U \mapsto \mathcal{F}_{\text{Ham}}^{\text{cl}}(U \times V)$  is locally constant in the topological directions for every fixed holomorphic open  $V$ .
- (ii) (*Holomorphic locality.*)  $L_{\partial/\partial \bar{z}_i} = [D, \iota_{\partial/\partial \bar{z}_i}]$  on  $\mathcal{F}_{\text{Ham}}^{\text{cl}}$ ; hence the assignment  $V \mapsto \mathcal{F}_{\text{Ham}}^{\text{cl}}(U \times V)$  is a holomorphic factorization algebra on  $\mathbb{C}^2$  via Dolbeault descent.
- (iii) (*Factorization product.*) For pairwise disjoint  $O_1, \dots, O_k \subset O$ ,  $\bigotimes_i \mathcal{F}_{\text{Ham}}^{\text{cl}}(O_i) \rightarrow \mathcal{F}_{\text{Ham}}^{\text{cl}}(O)$  is a chain map and Weiss-cosheaf descent over hypercovers holds.

Equivalently,  $\mathcal{F}_{\text{Ham}}^{\text{cl}} \in \text{Fact}_{\text{HT}}(\mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2)$  in the sense of Definition 1.10.

*Proof. Topological locality.* Cartan's homotopy  $[d_{\mathbb{R}^2}, \iota_v] = L_v$  on  $\Omega_c^\bullet(U)$  lifts to  $[D, \iota_v] = L_v$  on  $\mathcal{L}_{\text{Ham},c}(O)$ , since  $\iota_v$  commutes with  $\tilde{d}_{\mathbb{C}^2}$  and  $d_{\mathfrak{g}}$ . The CE functor carries this homotopy to  $\mathcal{F}_{\text{Ham}}^{\text{cl}}$ . *Holomorphic locality.* Identical argument with  $\iota_v$  replaced by  $\iota_{\partial/\partial \bar{z}_i}$  and  $d_{\mathbb{R}^2}$  by  $\tilde{d}_{\mathbb{C}^2}$ . *Factorization product.* Extension-by-zero  $\Omega_c^\bullet(O_i) \hookrightarrow \Omega_c^\bullet(O)$  on each factor of  $\mathcal{L}_{\text{Ham},c}$  is a dg Lie injection for disjoint supports; CE chains of a coproduct of dg Lie algebras is the tensor product of CE chains, giving the factorization product. The Weiss-cosheaf condition is the standard de Rham/Dolbeault descent for compactly supported forms.  $\square$

**COROLLARY 1.21** ( *$E_2$ -bulk algebra*). Fix a holomorphic polydisk  $B \subset \mathbb{C}^2$  and a topological disk  $D^2 \subset \mathbb{R}^2$ . The bulk classical algebra

$$\mathcal{A}_{\text{bulk}}^{\text{cl}}(B) = \mathcal{F}_{\text{Ham}}^{\text{cl}}(D^2 \times B)$$

is an  $E_2$ -algebra under the Lurie–Costello–Gwilliam locally constant factorization structure on  $\mathbb{R}^2$ ; the assignment  $B \mapsto \mathcal{A}_{\text{bulk}}^{\text{cl}}(B)$  is a holomorphic factorization algebra on  $\mathbb{C}^2$  valued in  $E_2$ -algebras.

*Proof.* Theorem 1.20 (1) says that  $U \mapsto \mathcal{F}_{\text{Ham}}^{\text{cl}}(U \times B)$  is a locally constant factorization algebra on  $\mathbb{R}^2$ ; the value on a topological disk is therefore an  $E_2$ -algebra by the Lurie–Costello–Gwilliam classification of locally constant factorization algebras on  $\mathbb{R}^n$ . Holomorphic factorization in  $B$  is item (2) of the same theorem.  $\square$

**Definition 1.22** ( *$E_1$ -brane algebra*). Let  $\text{Kosz}_N$  be the Dirac BV complex of Construction 3.20. The classical brane factorization algebra on  $L = \mathbb{R}_t$  assigns to an interval  $I \subset \mathbb{R}_t$  the CE complex

$$\text{Obs}_{\partial,N}^{\text{cl}}(I) = C_{\text{CE},\text{cont}}^\bullet(\Omega_c^\bullet(I) \otimes \mathfrak{gl}_N, \Omega_c^\bullet(I) \otimes \text{Kosz}_N),$$

with factorization product extension by zero on  $\Omega_c^\bullet$  and multiplication of  $\text{Kosz}_N$ -valued cochains. Local constancy on  $\mathbb{R}_t$  makes the value on a small interval an  $E_1$ -algebra:

$$\mathcal{A}_{\partial,N}^{\text{cl}} = C_{\text{CE}}^\bullet(\mathfrak{gl}_N, \text{Kosz}_N) \in \text{Alg}_{E_1}.$$

Its ghost-zero  $Q$ -cohomology is the stable trace ring of Theorem 5.41:  $H^\circ(\mathcal{A}_{\partial,N}^{\text{cl}}) \simeq H^\circ(O(\mu_{\text{der}}^{-1}(\mathfrak{o}))^{GL_N}, Q)$ .

A closed Hamiltonian  $f \in \mathfrak{h}$  acts on the brane representation moduli through a Schouten vector field  $\theta_f$ , dual on the open side to the trace character map  $u_f \mapsto J_f$ . The following lemma writes  $\theta_f$  as an explicit derivation on the formal representation functor, fixing the closed-to-open chain map at the level of generators.

**LEMMA 1.23** (*Hamiltonian vector field on the representation stack*). For  $f \in \mathfrak{h} = A/C \cdot 1$  the assignment

$$\theta_f \phi_1 = -\partial_{z_2} f|_{(\phi_1, \phi_2)}, \quad \theta_f \phi_2 = +\partial_{z_1} f|_{(\phi_1, \phi_2)}$$

extends uniquely to a  $Q$ -commuting derivation  $\theta_f \in \text{Der}(O(\text{Rep}_N^\wedge(A)))$ ; the value  $\theta_f \psi \in \mathfrak{gl}_N[1]$  is determined by

$$Q(\theta_f \psi) = \theta_f([\phi_1, \phi_2]) = [\theta_f \phi_1, \phi_2] + [\phi_1, \theta_f \phi_2].$$

This  $\theta_f$  preserves the moment-map ideal, descends to a derivation on the derived commuting-pair stack  $O(\mu_{\text{der}}^{-1}(\mathfrak{o}))^{GL_N}$ , and on trace characters acts as

$$\theta_f(J_g) = J_{\{f,g\}_{\bar{A}}} \quad (f, g \in \mathfrak{h}),$$

with  $\{-, -\}_{\bar{A}}$  the Hamiltonian Poisson bracket on  $\mathfrak{h}$ .

*Proof.* The matrix Poisson bracket  $\{(\phi_1)^i_j, (\phi_2)^k_l\} = \delta_l^i \delta_j^k$  on  $\text{Rep}_N(A_{\text{poly}})$  extends by Artin-test continuity to  $\text{Rep}_N^\wedge(A)$  and renders  $\theta_f \bullet = \{J_f, \bullet\}$  for the polynomial generator  $J_f = \overline{\text{Tr } f(\phi_1, \phi_2)}$ , whose  $\partial_{\phi_i}$ -derivatives on a monomial  $\phi_1^k \phi_2^l$  are the cyclic-trace expressions of Proposition 5.170 corresponding to  $\partial_{z_1} \partial_{z_2} f$ . This identifies  $\theta_f = \{J_f, -\}$  on the matrix coordinate algebra and gives the displayed formulas on  $\phi_1, \phi_2$ . Since  $Q\phi_i = 0$ ,  $[\theta_f, Q]\phi_i = 0$  automatically; on  $\psi$ ,  $[\theta_f, Q]\psi = \theta_f([\phi_1, \phi_2]) - Q(\theta_f \psi)$ , and the displayed equation is exactly the vanishing condition for this commutator. The moment-map ideal is preserved because  $\{f(\phi_1, \phi_2), \mu_\epsilon\} = 0$  for every  $GL_N$ -invariant trace polynomial (Lemma 5.169). Identifying  $\theta_f$  with  $\{J_f, -\}$  and applying Proposition 5.170 gives  $\theta_f(J_g) = \{J_f, J_g\} = J_{\{f, g\}_A}$ .  $\square$

**THEOREM 1.24** (*Closed-to-open derived-centre map and CE/PV reduction*). There is a closed-to-open chain map

$$\Phi_N: \mathcal{A}_{\text{bulk},0}^{\text{cl}} \longrightarrow Z_{E_1}^{\text{der}}(A_{\partial,N}^{\text{cl}}), \quad Z_{E_1}^{\text{der}}(A_{\partial,N}^{\text{cl}}) \simeq HH^\bullet(A_{\partial,N}^{\text{cl}}, A_{\partial,N}^{\text{cl}}),$$

whose value on Hamiltonian CE generators is the explicit Hochschild cochain assignment

$$c_f \longmapsto \theta_f \in \text{Der}(A_{\partial,N}^{\text{cl}}) = HH^1(A_{\partial,N}^{\text{cl}}), \quad u_f \longmapsto O_f = J_f \cdot (\cdot) \in A_{\partial,N}^{\text{cl}} = HH^0(A_{\partial,N}^{\text{cl}}) \quad (f \in \mathfrak{h}),$$

with  $J_f = J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)}$  acting by multiplication and  $\theta_f$  the Hamiltonian vector field of Lemma 1.23; this matches the cochain identity of Theorem 4.20 on  $c^{a,b}, u_{a,b}$ . The full chain-level lift to the strict factorization-centre morphism  $\Phi^{\text{cl}}: \text{Obs}_{\text{closed},H}^{\text{cl}}|_L \rightarrow Z_{\text{fact}}^{P_0}(\text{Obs}_{\text{open},N}^{\text{cl}})$  is governed by the four-component obstruction vector  $\text{Ob}_{\text{cent/QME/desc}}$  of Theorem E.53. The CE/PV coordinate map  $c_f \mapsto \theta_f, u_f \mapsto O_f$  is the generator action of  $\Phi_N$ ;  $\Phi_N$  is in turn the Hochschild-cochain shadow of the chain-level  $P_0$ -centre map  $\Phi^{\text{cl}}$ , whose existence is conditional on  $\text{Ob}_{\text{cent/QME/desc}} = 0$  with compatible null-homotopies.

*Proof.* The boundary algebra  $A_{\partial,N}^{\text{cl}}$  is an  $E_1$ -algebra, and the bulk algebra  $\mathcal{A}_{\text{bulk},0}^{\text{cl}}$  is an  $E_2$ -algebra of Chevalley–Eilenberg coordinates on  $\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1]$ . The Hochschild complex  $C^\bullet(A_{\partial,N}^{\text{cl}}, A_{\partial,N}^{\text{cl}})$  carries the Gerstenhaber bracket of degree  $-1$ ; its cohomology is the derived  $E_1$ -centre, with  $HH^0$  the centre (multiplications) and  $HH^1$  the outer derivations.

The map  $\Phi_N$  is defined on CE generators as Hochschild cochains. For  $f \in \mathfrak{h}$ , the Hamiltonian-dual coordinate  $c_f$  is sent to the Hamiltonian vector field  $\theta_f \in \text{Der}(A_{\partial,N}^{\text{cl}}) = HH^1(A_{\partial,N}^{\text{cl}})$  of Lemma 1.23 (the classical brane algebra is graded-commutative with odd antifield generator  $\psi$ , so  $HH^1$  is the full module of graded derivations); the cotangent-dual coordinate  $u_f$  is sent to the Hochschild 0-cochain  $J_f \cdot (\cdot) \in HH^0(A_{\partial,N}^{\text{cl}}) = A_{\partial,N}^{\text{cl}}$ , multiplication by the trace character  $J_f = \overline{\text{Tr } f(\phi_1, \phi_2)}$ . Compatibility with the CE differential is the Hochschild Gerstenhaber identity

$$[\theta_f, \theta_g]_G = \theta_{\{f,g\}_A}, \quad \theta_f(J_g) = J_{\{f,g\}_A},$$

matching  $d_{\text{CE}} c^K = -\frac{1}{2} C_{IJ}^K c^I c^J$  and  $d_{\text{CE}} u_K = C_{KJ}^L u_L c^J$  on generators by Lemma 1.23 and Proposition 5.170.

Continuous extension across the graded-commutative coordinate algebra is Theorem 4.20. Deligne's  $E_1$ -Hochschild theorem identifies  $Z_{E_1}^{\text{der}}(A_{\partial,N}^{\text{cl}})$  with  $HH^\bullet(A_{\partial,N}^{\text{cl}}, A_{\partial,N}^{\text{cl}})$  as  $E_2$ -algebras; the Costello–Gwilliam factorization-algebra bulk–boundary correspondence glues the generator data into the universal bulk-to-boundary chain map landing in this  $E_2$ -algebra. Uniqueness of the boundary values  $u_f \mapsto J(f)$  is Theorem 5.41.

An explicit Hochschild-cochain model for  $\Phi_N$  at every cochain degree is the conditional upgrade  $\Phi^{\text{cl}}$  to the strict  $P_0$ -factorization-centre, governed by the obstruction  $\text{Ob}_{\text{cent/QME/desc}}$  of Theorem E.53; the curvature class is the difference between the derived  $E_1$ -centre  $E_2$ -structure of Deligne’s theorem and the strict  $P_0$ -factorization-centre.  $\square$

*Remark 1.25 (Native bulk object on  $\mathbb{C}^2$ ).* The native object is the mixed holomorphic-topological factorization algebra  $\mathcal{F}_{\text{Ham}}^{\text{cl}}$  on  $X = \mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$ . The bulk  $\mathcal{A}_{\text{bulk}}^{\text{cl}} = \pi_* \mathcal{F}_{\text{Ham}}^{\text{cl}}$  is its pushforward along  $\pi: X \rightarrow \mathbb{C}_{\text{hol}}^2$ , the holomorphic factorization algebra  $B \mapsto \mathcal{A}_{\text{bulk}}^{\text{cl}}(B)$  on  $\mathbb{C}_{\text{hol}}^2$  valued in  $E_2$ -algebras — the  $E_2$ -valuation encoding the locally constant  $\mathbb{R}_{\text{top}}^2$  factor by the Lurie–Costello–Gwilliam classification. A one-dimensional vertex algebra is a derived object, recovered after a restriction datum—a holomorphic curve  $C \hookrightarrow \mathbb{C}^2$  with transverse boundary condition, or a factorization-homology pushforward—which singles out a one-complex-dimensional holomorphic factorization restriction on which Zhu / finite-Laurent OPE applies. Before any such restriction, the native singular product is two-complex-dimensional: Theorem 1.15 identifies the diagonal local-cohomology coefficient module with  $H_{\Delta}^2(\mathbb{C}[[z, w]]) d\zeta_1 d\zeta_2$  and computes

$$J_x(z)J_y(w) \sim K_{\Delta}^{(2)}(z, w)J_{[x, y]_{\mathfrak{g}}}(w).$$

**PROPOSITION 1.26 (Bochner–Martinelli current data and finite-window obstruction for  $E_2$  transfer).** Let  $B_0 \Subset B_1 \Subset B_2 \Subset B \subset \mathbb{C}^2$  be nested polydisks around the collision point and choose  $\chi \in C_c^\infty(B_2)$  with  $\chi = 1$  on  $B_1$ . Put

$$\mathcal{P} = \bigoplus_{\substack{a, b \geq 0 \\ a+b > 0}} \mathbb{C} \rho_{a, b}, \quad \rho_{a, b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2,$$

and

$$\mathcal{P}_{\leq N} = \bigoplus_{0 < a+b \leq N} \mathbb{C} \rho_{a, b}.$$

The symbol  $\mathfrak{h}_{\leq N} = \mathfrak{m}/\mathfrak{m}^{N+1}$  is the coordinate window; the full Hamiltonian Lie bracket requires the completed algebra  $\mathfrak{h}$ .

On  $(B_\zeta \times B_z) \setminus \Delta$ , set

$$\eta = \zeta - z, \quad r^2 = |\eta_1|^2 + |\eta_2|^2, \quad d\tilde{\eta}_j = d\tilde{\zeta}_j - d\tilde{z}_j.$$

The Dolbeault homotopy kernel is the Bochner–Martinelli–Koppelman current [37, Ch. III, Def. 1.2, Lem. 1.3]

$$K_{\text{BM}}(\zeta, z) = \frac{1}{(2\pi i)^2} \frac{\bar{\eta}_1 d\tilde{\eta}_2 - \bar{\eta}_2 d\tilde{\eta}_1}{r^4} \wedge d\tilde{\zeta}_1 \wedge d\tilde{\zeta}_2.$$

The external several-complex-variables sources are used here only for this pre-cutoff kernel, its component  $\bar{\partial}$ -identity away from the diagonal, and the classical Koppelman homotopy formula on bounded  $C^1$  domains for  $C^1$  forms [38, §5, Def. 5.1, Thm. 5.2, Eq. (10)]. The cutoff-current limit  $H_\chi$ , the annular diagonal-current sign, the finite moment projection  $p_N$ , the residue-current section  $i_N$ , the analytic/pro-Matlis quotient, the Hamiltonian/Matlis arity-two action, and the Costello/QME or factorization transfer below are additional constructions. The  $d\tilde{z}_j$ -terms in  $d\tilde{\eta}_j$  are part of the kernel; the  $d\tilde{z} = 0$  component is only the scalar integral kernel. With a radial cutoff  $\vartheta_\epsilon(r)$ , equal to 0 for  $r \leq \epsilon$  and to 1 for  $r \geq 2\epsilon$ , put

$$K_{\chi, \epsilon} = \chi(\zeta)\chi(z)\vartheta_\epsilon(|\zeta - z|)K_{\text{BM}}(\zeta, z).$$

The current limit

$$H_\chi \alpha(z) = \lim_{\epsilon \rightarrow 0} \int_{\zeta \in B} K_{\chi, \epsilon}(\zeta, z) \wedge \alpha(\zeta)$$

is well-defined on each fixed finite configuration stratum. The singularity is locally integrable in real dimension four ( $|K_{\text{BM}}| \sim r^{-3}$ ), and for separated compact supports its derivatives obey estimates

$$|\nabla^k K_{\text{BM}}(\zeta, z)| \leq C_k |\zeta - z|^{-3-k}.$$

Let  $i_N(\rho_{a,b})$  be the support-at-zero current

$$\Delta_{a,b} = \frac{(-1)^{a+b}}{a! b!} \partial_{z_1}^a \partial_{z_2}^b \partial_o d\bar{z}_1 \wedge d\bar{z}_2, \quad \langle \Delta_{a,b}, f \rangle = \text{Res}_o(\rho_{a,b} f),$$

and define the finite principal-part moment projection on top Dolbeault degree by

$$p_N(\alpha) = \sum_{o < a+b \leq N} \left( \int_B z_1^a z_2^b \alpha^{o,2} \wedge dz_1 \wedge dz_2 \right) \rho_{a,b}.$$

The data above construct the current-limit operator  $H_\chi$ , the finite moment map  $p_N$ , and the residue-current section  $i_N$ . A finite-window homotopy  $H_{\chi,N}$  on the full compact-support/current complex additionally requires the identity

$$\bar{\partial} H_{\chi,N} + H_{\chi,N} \bar{\partial} = \text{id} - i_N p_N + E_{\chi,N} \quad (*_N)$$

is blocked, in the full-current completion, by the finite-window obstruction vector

$$\text{Ob}_N^{\text{BMK-curr}} = \left( \text{ob}_{\text{diag}}, \text{ob}_{\text{an/fin},N}, \text{ob}_{\mathfrak{h},N}, \text{ob}_{\text{collar-fact},N} \right).$$

Here  $\text{ob}_{\text{diag}}$  is the sign and scalar comparison between the annular diagonal current  $\bar{\partial} \mathcal{D}_\epsilon(|\zeta - z|) \wedge K_{\text{BM}}$  and the Matlis derivative-delta convention; with the displayed kernel and real outward boundary orientation its angular mass is  $-1$ , so a BMK diagonal-current convention must be declared before the sign in  $i_N p_N$  is fixed. The component  $\text{ob}_{\text{an/fin},N}$  is the quotient between the full analytic moment kernel  $\ker(O(B_i)^\vee \rightarrow \mathcal{P}_{\leq N})$  and the smaller support-at-zero span generated by  $\Delta_{o,o}$  and the  $\Delta_{a,b}$  with  $a + b > N$ . The component  $\text{ob}_{\mathfrak{h},N}$  is the Hamiltonian-invariance defect

$$\pi_N(\mathfrak{h} \cdot \ker \pi_N),$$

already nonzero because

$$z_1^{N+2} \cdot \rho_{N+1,o} = -(N+2) \rho_{o,1},$$

while  $\rho_{N+1,o}$  is killed by the window and  $\rho_{o,1}$  is retained. Finally  $\text{ob}_{\text{collar-fact},N}$  is the still-open obstruction, away from one nested pair, to simultaneous preservation of the same relation subspace by the collar quotient, extension-by-zero maps, factorization products, and the BMK homotopy.

Consequently strict finite support-local current quotient retaining  $\mathcal{P}_{\leq N}$ , killing the scalar and high support-at-zero currents, and carrying the full Hamiltonian factorization action is obstructed. The projected finite-window  $L_\infty$  escape repair is obstructed as well: the forced return block already has

$$\kappa_N(z_1^{N+2}, \rho_{N+1,o}) = -(N+2) \rho_{o,1},$$

and the projected Jacobiator on  $(z_1^{N+2}, z_2, \rho_{N,o})$  is

$$(N+1)(N+2)\rho_{o,1}.$$

Since the escape-return image spans the retained finite Matlis window, the unary-boundary closure of these returns coincides with the full collapse of  $\mathcal{P}_{\leq N}$  onto the trivial module; every nontrivial finite Matlis window therefore retains the Bochner–Martinelli leak. A one-pair derived analytic quotient does give a differential identity, after the full analytic BMK contraction and diagonal orientation are fixed: quotient the relative current complex by representatives of  $\ker(\mathcal{O}(B_1)^\vee \rightarrow \mathcal{P}_{\leq N})$ , normalize the homotopy by

$$(1 - I_{\text{an}} P_{\text{an}}) H_\chi (1 - I_{\text{an}} P_{\text{an}}),$$

and then

$$\bar{\partial} H_{\text{BM},N} + H_{\text{BM},N} \bar{\partial} = \text{id} - i_N p_N.$$

This quotient is analytic and one-pair relative. The source theorem used here is only the pre-cutoff Koppelman identity; the representative quotient, the normalized contraction, and the compatible pro-Matlis moment map are local constructions of this proposition.

Keeping the entire Taylor tower gives the positive one-pair pro-Matlis replacement. Let

$$\widehat{\mathcal{P}}_\Pi = \prod_{\substack{a,b \geq 0 \\ a+b > 0}} \mathbb{C} \rho_{a,b}, \quad \mathcal{P}_{\text{an}}(B_1) = \tau(\mathcal{O}(B_1)_{\text{red}}^\vee) \subset \widehat{\mathcal{P}}_\Pi,$$

where

$$\tau(\lambda) = (\lambda(z_1^a z_2^b))_{a+b > 0}.$$

On the single-pair analytic BMK completion, after the diagonal orientation is fixed and the collar error is quotiented, the normalized contraction  $H_\chi^o = (1 - I_{\text{an}} P_{\text{an}}) H_\chi (1 - I_{\text{an}} P_{\text{an}})$  satisfies

$$\bar{\partial} H_\chi^o + H_\chi^o \bar{\partial} = \text{id} - I_{\text{an}} P_{\text{an}}.$$

Hence the pro-analytic moment map  $P_\Pi^{\text{an}} = \tau P_{\text{an}}$  has compatible finite projections

$$\pi_N P_\Pi^{\text{an}} = p_N, \quad \pi_{M,N} p_M = p_N.$$

This is a one-pair analytic pro-Matlis retract.

The binary operation

$$\ell_2^{\text{BM},N}(x, y) = p_N[ix, iy]$$

is therefore retained only as an arity-two Hamiltonian/Matlis coefficient comparison induced by  $p_N$  and  $i_N$ . A finite-window homotopy would be an additional construction. It agrees, coefficient by coefficient and in every fixed output window, with the Hamiltonian semidirect bracket. Thus for Hamiltonians  $f, g \in \mathfrak{h}$ ,

$$\ell_2^{\text{BM},N}(f, g) = \text{pr}_{\leq N}(\partial_{z_1} f \partial_{z_2} g - \partial_{z_2} f \partial_{z_1} g) \pmod{\mathbb{C}},$$

while for  $f = z_1^p z_2^q$  and  $\rho_{i,j} = z_1^{-i-1} z_2^{-j-1} dz_1 dz_2$ ,

$$\ell_2^{\text{BM},N}(f, \rho_{i,j}) = -(pj - qi + p - q) \text{pr}_{\leq N} \rho_{i-p+1, j-q+1},$$



with negative-index targets equal to zero, and

$$\ell_2^{\text{BM},N}(\rho, \sigma) = 0.$$

This is the Matlis coadjoint action of Theorem 5.178.

Only after  $\text{Ob}_N^{\text{BMK-curr}}$  is killed, or after the one-pair analytic quotient is replaced by a factorization-current completion, does the first possible obstruction to a strict all-arity  $E_2$  collision operation become the ternary cutoff term

$$\text{Ob}_3^\chi(x, y, z) = p[H_{\text{BM}}([ix, iy] - i\ell_2^{\text{BM}}(x, y)), iz] + \text{cyclic Koszul terms}.$$

It vanishes on separated support strata by the estimates above, and its coefficient Jacobiator on the collision stratum is the Jacobiator of  $\mathfrak{h} \times \mathcal{P}[1]$ , hence zero. Any surviving term is therefore a cutoff/collar contribution supported on the small diagonal. The strict all-window current  $E_2$  transfer is downstream and is controlled by the six-entry obstruction vector

$$\text{Ob}_{\text{BM}}^\Pi = (\text{ob}_\oplus, \text{ob}_{I_\Pi}, \text{ob}_{\mathfrak{h}, \Pi}, \text{ob}_{\text{collar}, \Pi}, \text{ob}_{3, \Pi}, \text{ob}_{\text{unif}, \Pi}).$$

Here  $\text{ob}_\oplus$  is the nonzero class of the all-window moment sequence in  $\widehat{\mathcal{P}}_\Pi/\mathcal{P}$ ;  $\text{ob}_{I_\Pi}$  is the nonexistence of a compact-current section  $i_\Pi : \widehat{\mathcal{P}}_\Pi \rightarrow \mathcal{D}'^{0,2}(B)$  extending the finite derivative-delta sections; and  $\text{ob}_{\mathfrak{h}, \Pi}$  is the obstruction to the full formal Hamiltonian action on the pro-product module, detected by the divergent  $\rho_{0,1}$ -coefficient for  $f = \sum_{p \geq 2} z_1^p$  acting on  $\lambda = \sum_{p \geq 2} \rho_{p-1,0}$ . The remaining entries are the compatible collar primitive class, the first ternary small-diagonal transfer class, and the all-window uniform estimate class. The first, second, third, and sixth entries are nonzero in the current topology; the collar and ternary entries are exact primitive construction targets.

The principal-part face of the transfer is genuinely two-variable:

$$\frac{1}{(2\pi i)^2} \frac{d\zeta_1 \wedge d\zeta_2}{(\zeta_1 - z_1)(\zeta_2 - z_2)} = \sum_{a,b \geq 0} z_1^a z_2^b \rho_{a,b}(\zeta).$$

Killing  $z_2$  kills the Hamiltonian bracket, and the  $b = 0$  principal-part line is unstable because  $z_1 \cdot \rho_{a,0} = -\rho_{a,1}$ .

*Proof.* The external Koppelman identity for  $K_{\text{BM}}$  supplies only the smooth-domain, pre-cutoff Dolbeault homotopy [38, §5, Thm. 5.2]. The cutoff derivative, annular diagonal current, finite moments, and Matlis orientation below are checked locally. After multiplication by the cutoff, the load-bearing derivative is

$$\bar{\partial}(\chi(\zeta)\chi(z)\vartheta_\epsilon K_{\text{BM}}) = \chi\chi\vartheta_\epsilon \bar{\partial}K_{\text{BM}} + \bar{\partial}(\chi\chi)\vartheta_\epsilon K_{\text{BM}} + \chi\chi\bar{\partial}\vartheta_\epsilon \wedge K_{\text{BM}}.$$

The second summand is collar-supported. The third summand is an annular diagonal current; its sign and scalar are the first entry of  $\text{Ob}_N^{\text{BMK-curr}}$ . Applying finite moments gives the formula for  $p_N$ . The normalization of  $i_N$  follows by integration by parts against a holomorphic test function, giving the residue functional represented by  $\rho_{a,b}$ , once the diagonal-current orientation is tied to the Matlis convention.

The full-current finite-window identity has a Stokes obstruction on the current complex. It would make every closed top current whose analytic Dolbeault cohomology class lies in  $\ker \pi_N$  exact after quotienting by the finite selected moments, while pairing with a holomorphic function detects that class. In particular,  $T = \Delta_{N+1,0}$  has  $p_N(T) = 0$ , but  $\langle T, z_1^{N+1} \rangle = 1$ , whereas  $\langle \bar{\partial}S, z_1^{N+1} \rangle = 0$  for every

current  $S$ . The scalar current  $\Delta_{o,o}$  gives the same obstruction because  $\rho_{o,o}$  is excluded from the reduced principal-part module.

The stronger support-local quotient obstruction is Hamiltonian invariance. A quotient relation must be an  $\mathfrak{h}$ -submodule relation. The Matlis coadjoint formula gives  $z_1^{N+2} \cdot \rho_{N+1,o} = -(N+2)\rho_{o,1}$ ; hence killing the high mode  $\rho_{N+1,o}$  while retaining the finite mode  $\rho_{o,1}$  is incompatible with the full Hamiltonian action: high transverse holomorphic modes leak back into low brane-momentum modes. This proves the nonexistence of the strict finite support-local quotient stated above; the all-window form is Theorem 8.40, with the pro-Matlis retract the target replacement absorbing this leak (Corollary 8.41).

The one-pair analytic quotient is formal once the full analytic BMK contraction is supplied locally: the normalized homotopy annihilates the chosen representatives of  $\ker \pi_N$ , so it descends and the analytic projector becomes  $i_N p_N$  in the quotient. On the unquotiented Taylor tail, the same normalized contraction gives  $\bar{\delta} H_\chi^o + H_\chi^o \bar{\delta} = \text{id} - I_{\text{an}} P_{\text{an}}$ , and Taylor moments give  $P_{\Pi}^{\text{an}} = \tau P_{\text{an}}$  with  $\pi_N P_{\Pi}^{\text{an}} = p_N$ . Because this pro-Matlis retract depends on the analytic moment space, a representative map, and a chosen polydisk pair, a factorization-current theorem requires additional support-local data.

The arity-two formula is the residue transpose calculation. On holomorphic coefficient labels this is the Poisson bracket modulo constants, projected after the coefficient is computed. On cotangent labels the residue identity

$$\text{Res}_o((f \cdot \rho)g) = -\text{Res}_o(\rho \{f, g\})$$

gives exactly the principal-part coadjoint formula already proved in Theorem 5.178. The cotangent summand is abelian, so the  $(\rho, \sigma)$ -bracket is zero.

The displayed ternary expression is therefore only the next obstruction after the finite-current condition is crossed. Away from collisions the kernel is smooth and the separated-support estimates force it to vanish in the factorization product. On the coefficient collision stratum the semidirect Lie algebra satisfies Jacobi. The remaining work is the collar, diagonal, and completed all-window transfer data.  $\square$

**PROPOSITION 1.27** (*Darboux polydisk constructions*). For

$$X = \mathbb{R}_t \times \mathbb{R}_s \times \mathbb{C}_{z_1, z_2}^2, \quad L = \mathbb{R}_t \times \{s = z_1 = z_2 = o\},$$

with holomorphic Darboux form  $dz_1 \wedge dz_2$ , the local theorem has the following five constructions.

- (i) The closed Hamiltonian BF coefficient algebra is

$$\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1, \quad P = \mathfrak{h}_{\text{cont}}^\vee, \quad \mathfrak{g} = \mathfrak{h} \ltimes P[1].$$

The quotient by constants comes from the kernel of the Hamiltonian vector-field map.

- (ii) The finite- $N$  open chart is the Dirac/Koszul matrix model

$$Q\psi = [\phi_1, \phi_2], \quad Q\phi_1 = Q\phi_2 = o,$$

obtained from the first-order action  $\int_{\mathbb{R}_t} \text{Tr}(\phi_1 d\phi_2 + A[\phi_1, \phi_2])$ . Thus the open algebra is the derived zero-fibre of the conjugation moment map, with commutativity imposed homologically.

- (iii) After killing the scalar Hamiltonian and the empty trace, the reduced CE/PV central-operation comparison is the coordinate map

$$\Phi_{\text{coord}}(c^I) = \theta^I, \quad \Phi_{\text{coord}}(u_I) = O_I.$$

The  $c$ -coordinate gives the Schouten vector-field operation; the  $u$ -coordinate gives the boundary Hamiltonian observable.

- (iv) The open one- $\psi$  classes are PBW special-fibre labels. The deformation-level cotangent dual is  $P$ , realized by Matlis principal parts and acted on by the coadjoint action. The PBW/Rees label bridge is extra Fourier-polarized structure relating two label systems with distinct  $\mathfrak{h}$ -actions.
- (v) The Bochner–Martinelli–Koppelman current limit, finite Matlis moment maps, arity-two coefficient comparison, and one-pair analytic pro-Matlis retract  $P_{\Pi}^{\text{an}} = \tau P_{\text{an}}$ ,  $\pi_N P_{\Pi}^{\text{an}} = p_N$ , are constructed in Proposition 1.26. A strict support-local finite-window current quotient is obstructed by  $\text{Ob}_N^{\text{BMK-curr}}$ . A strict all-arity  $E_2$  transfer is the downstream theorem target: it requires first crossing this finite-current condition and then solving the six-entry all-window obstruction problem  $\text{Ob}_{\text{BM}}^{\Pi}$ .

*Proof.* The formal Darboux polydisk is fixed in the opening of this section. Item (1) is Proposition 2.7 and Proposition 1.14. Item (2) is the constrained-mechanics computation of Lemma 3.15, Lemma 3.14, and Proposition 3.16. Item (3) is Theorem 4.20, with the scalar convention of Remark 2.3 and Theorems 6.3–6.6. Item (4) is Theorem 5.180 together with the principal-part uniqueness theorem 5.178. Item (5) is Proposition 1.26.  $\square$

*Remark 1.28 (Formal stalk restriction).* The Hamiltonian identification is formal local. On a holomorphic symplectic surface  $X$ , a holomorphic vector field  $v$  is symplectic when

$$\alpha_v = -\iota_v \omega$$

is a closed holomorphic one-form, and Hamiltonian only after choosing  $f$  with  $df = \alpha_v$ , in the convention  $\iota_{X_f} \omega = -df$ . A global Hamiltonian sector requires a sheaf of local Hamiltonians and the vanishing of the corresponding holomorphic de Rham obstruction. The holomorphic Poincaré lemma makes this obstruction vanish on the formal polydisk.

**PROPOSITION 1.29 (Formal-stalk Hamiltonian restriction).** Let  $(X, \omega)$  be a connected holomorphic symplectic surface and let  $p \in X$ . Put

$$\mathcal{H}_X = \mathcal{O}_X / \mathbb{C}_X, \quad \widehat{\mathfrak{h}}_{X,p} = \widehat{\mathcal{O}}_{X,p} / \mathbb{C}.$$

The differential identifies  $\mathcal{H}_X$  with the sheaf of closed holomorphic one-forms, hence with the sheaf of holomorphic symplectic vector fields by  $\alpha = -\iota_v \omega$ . For every  $p \in X$  there is a Lie algebra restriction

$$H^0(X, \mathcal{H}_X) \longrightarrow \widehat{\mathfrak{h}}_{X,p}.$$

After choosing a formal Darboux coordinate system, the formal neighbourhood  $(\widehat{X}_p, \omega)$  has the same completed Hamiltonian Lie algebra as the formal polydisk used in the local model. The resulting stalk-level boundary map

$$\widehat{\mathfrak{h}}_{X,p} \longrightarrow H_{\text{Ham}}^0(\text{Obs}_{\text{brane}}^{\text{cl}})$$

is independent of this choice up to the natural action of the formal symplectomorphism group. The subspace coming from global Hamiltonian functions is the image of

$$H^0(X, \mathcal{O}_X) / \mathbb{C} \longrightarrow H^0(X, \mathcal{H}_X).$$

The obstruction for a locally Hamiltonian symplectic vector field to come from a global Hamiltonian function is its period class in  $H^1(X, \mathbb{C}_X)$ .

*Proof.* The holomorphic Poincaré lemma gives an isomorphism

$$d : \mathcal{O}_X / \mathbb{C}_X \xrightarrow{\sim} \Omega_{X, \text{cl}}^1.$$

The holomorphic symplectic form identifies a symplectic vector field  $v$  with the closed one-form  $\alpha_v = -\iota_v \omega$ . Thus a section of  $\mathcal{H}_X$  is exactly a local Hamiltonian primitive, modulo local constants, and the Poisson bracket is independent of the local constants. Taking the germ at  $p$  and completing gives the displayed restriction to  $\widehat{\mathfrak{h}}_{X,p}$ .

The formal holomorphic Darboux theorem identifies the completed local symplectic algebra at  $p$  with  $\mathbb{C}[[z_1, z_2]]$  equipped with  $dz_1 \wedge dz_2$ . Under this identification Hamiltonians modulo constants become  $\widehat{\mathfrak{h}} = \mathbb{C}[[z_1, z_2]]/\mathbb{C}$ , and the point-brane Ext algebra is the coordinate-free exterior algebra  $\mathbf{Ext}_X^*(\mathcal{O}_p, \mathcal{O}_p) \cong \Lambda^* T_p^* X$ , with cyclic trace fixed by  $\omega_p$ . Hence the open BV reduction and the Hamiltonian boundary evaluation are exactly the formal local construction after choosing Darboux coordinates. A different Darboux coordinate system differs by a formal symplectomorphism, which acts on Hamiltonians and on the normal coordinate algebra; the substitution map  $z_i \mapsto \phi_i$  is equivariant for this action.

The exact sequence

$$0 \rightarrow \mathbb{C}_X \rightarrow \mathcal{O}_X \rightarrow \mathcal{H}_X \rightarrow 0$$

gives

$$0 \rightarrow H^0(X, \mathcal{O}_X)/\mathbb{C} \rightarrow H^0(X, \mathcal{H}_X) \xrightarrow{\delta} H^1(X, \mathbb{C}_X) \rightarrow H^1(X, \mathcal{O}_X).$$

For a locally Hamiltonian vector field,  $\delta$  is the period class of the closed one-form  $-\iota_v \omega$ . Vanishing of this class is exactly the existence of a single global holomorphic Hamiltonian primitive, unique modulo constants.  $\square$

*Remark 1.30 (Cross-target anchor: HT-import licensing).* The vanishing  $H^1(X, \mathbb{C}_X) = 0$  is the licensing condition that promotes the formal-local Hamiltonian model of Proposition 1.29 to a global holomorphic-topological theory. It is the named hypothesis  $\text{Hyp}_{\text{HT, glob, DR}}$  carrying the four-step license (descent + QME + anomaly + locality) required to pass from the formal stalk model to a compact global HT theory on  $X$ .

The closed mixed BF sector is the consequence of Witten's source convention applied to canonical transformations of the Darboux polydisk. The closed-field target is the shifted cotangent dg Lie algebra  $\mathfrak{g} = \widehat{\mathfrak{h}} \ltimes \widehat{\mathfrak{h}}_{\text{cont}}^\vee[1]$ ; on product opens

$$\Omega^*(D) \widehat{\otimes} \Omega^{0,*}(B) \widehat{\otimes} (\widehat{\mathfrak{h}}[1] \oplus \widehat{\mathfrak{h}}_{\text{cont}}^\vee[2]), \quad Q = d_{\mathbb{R}^2} + \tilde{d}_{\mathbb{C}^2}, \quad S = \frac{1}{2} \int \langle \beta, \{a, a\} \rangle dz_1 dz_2,$$

which Proposition 1.32 verifies as a mixed holomorphic-topological BV sector in the sense of Definition 1.10. The constants kernel of the Hamiltonian vector-field map forces the quotient  $\widehat{\mathfrak{h}} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$ ; the  $U(1)$  centre class  $[\tilde{c}]$  of §6.1 is its only obstruction to lifting to the unreduced polynomial Hamiltonian Lie algebra.

The open brane is the product of the exact local A-brane  $\mathbb{R} \subset T^*\mathbb{R}$  and the B-brane  $\mathcal{O}_p$  at the formal holomorphic point. Cyclic open string field theory [4], packaged AKSZ [6], gives the formal action  $S(a) = \sum_{n \geq 1} \frac{1}{n+1} \text{Tr}(a m_n(a, \dots, a))$  on  $\text{RHom}(\mathcal{F}, \mathcal{F})[1]$ ; the affine skyscraper Yoneda algebra  $\text{RHom}_{\mathbb{C}[[z_1, z_2]]}(\mathcal{O}_p, \mathcal{O}_p) \cong \Lambda(\varepsilon_1, \varepsilon_2)$  of Proposition 3.9 reduces the ghost-zero piece  $\alpha_0 = A_{\text{CS}} + \chi_1 \varepsilon_1 + \chi_2 \varepsilon_2$  to the Dirac first-order matrix model (7) under  $\phi_1 = -\chi_1$ ,  $\phi_2 = \chi_2$ ,  $A = -A_{\text{CS}}$ . The local

unimodular datum is the residue/cyclic pairing on this exterior algebra: a holomorphic volume input on the Darboux polydisk.

A topological line carrying a stack of  $\mathfrak{gl}_N$  Dirac branes meets a formal holomorphic-symplectic polydisk transverse to it. The brane line selects a Lagrangian inside that polydisk; the polydisk supplies the Hamiltonian deformations as conjugate matrix coordinates on the brane.

#### 1.4 Local setup at one brane stack

A stack of  $N$  branes carries the  $\mathfrak{gl}_N$ -valued Heisenberg pair

$$(\phi_1, \phi_2) \in \mathfrak{gl}_N \oplus \mathfrak{gl}_N, \quad \Omega_N = \text{Tr}(d\phi_1 \wedge d\phi_2),$$

of holomorphic conjugate matrix coordinates on the brane worldvolume. The trace functional of a Hamiltonian  $f \in \mathfrak{h}$  on the brane is  $J(f) = \overline{\text{Tr } f}(\phi_1, \phi_2)$ .

The pair  $(\phi_1, \phi_2)$  is a Chan–Paton representation of the two formal holomorphic coordinates. Its scalar reduction is one holomorphic symplectic pair. Thus the transverse local geometry is a single holomorphic symplectic surface  $(Y, \omega)$  at the brane point.

The formal-stalk chart uses the germ of  $Y$  at that point. Holomorphic Darboux trivialises the germ to

$$(\widehat{\mathbb{C}^2_o}, dz_1 \wedge dz_2),$$

so the target contributes only its formal symplectic polydisk.

The brane stack carries an open trace observable  $\text{Tr } f(\phi_1, \phi_2)$  for every Hamiltonian  $f$ . The trace is supported on the one-dimensional brane worldline. A second topological direction supplies the transverse propagator direction for the BV ghosts, and the moment-map multiplier becomes a one-form there. Thus a real two-dimensional topological factor is the minimal completion in which the Dirac moment-map evaluation and the bulk-to-brane locality coexist.

Under the formal-stalk-chart hypotheses — a brane stack carrying the  $\mathfrak{gl}_N$ -valued holomorphic-symplectic pair  $(\phi_1, \phi_2)$ , open trace observables along a one-dimensional brane worldline, a propagator direction transverse to that worldline inside the topological factor, and a single transverse holomorphic symplectic surface at the brane point — the three independent locality differentials of Proposition 1.34 fix the geometry as:

$$X = \mathbb{R}_t \times \mathbb{R}_s \times \mathbb{C}_{z_1, z_2}^2, \quad L = \mathbb{R}_t \times \{s = z_1 = z_2 = o\}, \quad \omega = dz_1 \wedge dz_2, \quad (1)$$

with  $\mathbb{R}_t$  the brane worldline,  $\mathbb{R}_s$  the transverse topological direction, and  $\mathbb{C}_{z_1, z_2}^2$  the formal Darboux polydisk. The holomorphic factor is read through its formal symplectic polydisk  $(\widehat{\mathbb{C}^2_o}, dz_1 \wedge dz_2)$ ; compact or global features of an ambient holomorphic symplectic surface contribute this local Darboux datum to the theorem.

The brane has topological worldvolume direction  $\mathbb{R}_t$  and holomorphic factor the point  $p = o \in \widehat{\mathbb{C}^2_o}$ , realised in the brane category as the skyscraper sheaf  $\mathcal{O}_p$ . The mixed brane is therefore

$$L = \mathbb{R}_t \times \{o\} \times \{p\} \hookrightarrow X,$$

the unique mixed brane on which  $J$  is unobstructed.

A brane stack is an  $N$ -dimensional Chan–Paton representation  $\rho_\phi: \mathcal{A} \rightarrow \text{End}(\mathbb{C}^N)$  of the formal coordinate algebra  $\mathcal{A} = \mathbb{C}[[z_1, z_2]]$ , with  $\rho_\phi(z_i) = \phi_i$  and the moment-map constraint  $[\phi_1, \phi_2] = o$  factoring it through the commuting variety. Equivalently, after gauge fixing it is a

class in  $[\text{Rep}_N(A)^{\text{comm}}/GL_N]$  with Chan–Paton pair  $(\phi_1, \phi_2) \in \mathfrak{gl}_N^{\oplus 2}$ . The Hamiltonian Lie algebra  $\mathfrak{h} = A/\mathbb{C}$  acts on this representation moduli by Hamiltonian vector fields; the  $c_f \mapsto \theta_f$  avatar of a Hamiltonian is its motion on the moduli. The trace character of a bulk Hamiltonian  $f \in \widehat{\mathfrak{h}} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$  at the shifted-cotangent avatar  $u_f$  is

$$J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)}, \quad (2)$$

the gauge-invariant trace of  $f$  evaluated on the Chan–Paton pair, projected to the stable scalar-reduced quotient. Under the formal-stalk-chart hypotheses of Proposition 1.32 together with  $GL_N$ -invariance, formal Darboux naturality, and stable scalar reduction, three independent inputs force  $J$ . First, Procesi–Razmyslov 1976/1980 [17, 18] identify the stable connected  $GL_N$ -invariants on  $\mathfrak{gl}_N \oplus \mathfrak{gl}_N$  as cyclic words in two letters; under the stated hypotheses these are the only candidate gauge-invariant scalar functionals (Proposition 3.22). Second, the moment-map zero-fibre  $[\phi_1, \phi_2] = 0$  makes the cyclic words commute pairwise on cohomology and collapses every cyclic trace to the commutative monomial trace  $\text{Tr}(\phi_1^k \phi_2^l)$  (Lemma 3.30, Corollary 3.32). Third, stable  $GL_N$ -invariance together with Darboux equivariance under the formal symplectomorphism group of  $(\widehat{\mathbb{C}^2_0}, dz_1 \wedge dz_2)$  and linear normalization on  $z_1, z_2$  forces the scalar correction to vanish at every monomial  $z_1^a z_2^b$ : the naturality lemma gives  $J(z_1^a z_2^b) = \text{Tr}(\phi_1^a \phi_2^b)$  directly, and the formal extension  $f = \sum_{a,b} f_{a,b} z_1^a z_2^b \mapsto J(f) = \sum_{a,b} f_{a,b} \text{Tr}(\phi_1^a \phi_2^b)$  follows by linearity and  $\mathfrak{m}$ -adic continuity. Under those three hypotheses  $J$  is the unique  $GL_N$ -invariant, Darboux-natural, scalar-reduced bulk-to-brane evaluation (Theorem 5.41; Corollary 5.39; Theorem 5.146 for the terminal-object form among admissible protected ambient data).

§§3.3–3.4 build the matrix-Chan–Paton BV theory whose ghost-zero classical observables are the cyclic Procesi–Razmyslov traces on the moment-map zero-fibre, with Koszul antifield  $\psi$  resolving the constraint  $[\phi_1, \phi_2] = 0$ . §2.1 builds the closed shifted-cotangent BV theory whose Hamiltonian factor is  $\widehat{\mathfrak{h}}$  and whose cotangent factor is the Grothendieck–Matlis principal-part module  $\mathcal{P} = \widehat{\mathfrak{h}}_{\text{cont}}^V$ . §2.2–4.3 assemble the Hamiltonian BF action and the bulk-to-brane coupling  $f \mapsto B_f(a)$  that smears  $J(f)$  along the brane line and matches the reduced Hamiltonian Lie bracket on the closed side with the  $P_0$ -bracket on the brane line (Theorem 4.5, Theorem 5.190). §7.5 records the obstruction calculus blocking the unreduced Tate-coefficient cotangent lift of this matched bracket: vector obstruction  $\text{Ob}_T$  of Theorem E.53 on the strict all-window side, with  $\mathfrak{o}_{\text{Cas}}$  the Casimir-kernel block and BMK obstruction Theorem 8.40 excluding one-pair analytic strict transfer and forcing the pro-Matlis retract as supremum (Corollary 8.41). §3.5 closes the section with the operator algebra of the open brane and the stable large- $N$  reduction in which  $J$  becomes a connected single-trace primitive generator and the Capelli/Moyal scalar shift identifies with the closed-side anomaly class  $[\bar{c}]$ .

Theorem 5.146 is the universal property statement. Among all admissible protected Hamiltonian sectors at a holomorphic-symplectic brane point, the formal Darboux model  $(\widehat{\mathfrak{h}} \ltimes \widehat{\mathfrak{h}}_{\text{cont}}^V[1], \Omega^*(\mathbb{R}) \otimes \Lambda(\varepsilon_1, \varepsilon_2) \otimes \mathfrak{gl}_N[1], [\bar{c}])$  with the bulk-to-brane evaluation (2) is terminal up to canonical isomorphism; the obstruction to membership of an ambient datum is the four-component protected twist class  $\text{Ob}_{p,X}^{\text{univ}}$ .

The cochain-level comparison between the closed Hamiltonian sector and the open brane observables runs through five rungs. (1) Finite-dimensional cotangent CE/PV identification on every truncation  $\mathfrak{h}_{\leq N} = \mathfrak{m}^3/\mathfrak{m}^{N+1}$  (Theorem 5.101). (2) Coordinate formal-polydisk envelope: the same generator dictionary  $c^{a,b} \mapsto \theta^{a,b}, u_{a,b} \mapsto O_{a,b}$  extends to the completed coordinate algebras of Definition 4.17 (Theorem 4.20). (3)  $P_0$ -bracket enhancement on a chosen bracket-admissible target  $B \subset \text{PV}_{\text{coord}}(A_{\partial, \text{coord}})$  (Definition 4.18); the strict ambient product is foreclosed by Lemma 5.211. (4) Maurer–Cartan diagonal kernel: the formal Casimir  $C_{\mathfrak{h}} = \sum_{d \geq 1} \sum_i e_{d,i} \otimes e_d^i$  converges in the chosen completion — realised by the weighted Tate envelope of Theorem 5.70. (5) Bar-cobar/Koszul Quillen upgrade

(Theorem 4.24), under a Tate/nilpotent/weighted completion with bounded inverse-limit hypotheses. The eight-component vector  $\text{Ob}_T$  of §7.5 records which level is reached and which obstruction blocks the next.

*Remark 1.31 ( $c_f$  versus  $u_f$ ).* The dictionary distinguishes two CE coordinates of a Hamiltonian  $f \in \widehat{\mathfrak{h}}$  on the closed side. The Hamiltonian-dual CE coordinate  $c^f$  maps under the cotangent CE/PV comparison to the Schouten polyvector coordinate  $\theta^f$  on  $\widehat{\mathbf{S}}(\vec{A})$ , the generator of an infinitesimal canonical motion as a vector field on the brane phase space. The shifted-cotangent-dual CE coordinate  $u_f$  maps to the boundary observable  $O_f = J(f)$ , the gauge-invariant scalar reading of  $f$  on the Chan–Paton pair. The closed Hamiltonian generator enters the boundary algebra through  $u_f$  (Theorem 4.20, Proposition 4.15): a vector field acts on observables, and perfection of the Hamiltonian Lie algebra kills closed length-one CE cochains in the Hamiltonian direction, so  $u_f$  is the unique CE coordinate whose CE/PV image  $O_f$  is itself an observable.

**PROPOSITION 1.32** (*Geometry determined by the formal-stalk chart*). The geometry (1) is the unique formal local mixed completion carrying the formal-stalk-chart datum  $(\Omega_N, \text{Tr})$  of the brane stack: the open observable family equals  $J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)}$  by the  $GL_N$ -invariant Darboux-natural scalar-reduced criterion (Theorem 5.146), and the mixed factorization-algebra shape is fixed by the three independent locality differentials of Proposition 1.34, with the trace observables exhibiting all three localities and the shifted-cotangent currents only the first two (Theorem 0.1). Its closed observable theory is the mixed holomorphic-topological Hamiltonian BF sector of Definition 1.10, with classical master equation; its mixed locality is the Cartan/Dolbeault homotopy. The brane open observable theory is the cyclic Dirac formal-stalk-chart theory of §3.3 and §3.4; the constructed bulk-to-brane operation is the degree-zero Hamiltonian boundary evaluation (Construction 3.25, Theorem 5.190). The unreduced cotangent factorization centre is the obstruction problem Problem 5.48; the open-side centrality-homotopy obstruction is Remark 5.189.

*Proof.* Closed BV data: on a product open  $D \times B$  in the completed formal neighbourhood of  $\mathbb{R}^2 \times \{p\} \subset \mathbb{R}^2 \times \mathbb{C}^2$ , Proposition 2.7 identifies divergence-free polyvectors-one with Hamiltonians modulo constants, so Hamiltonian BF in the Kodaira-Spencer convention has fields

$$\mathcal{E}_{\text{cl}}(D \times B) = \Omega^*(D) \widehat{\otimes} \Omega^{\circ,*}(B) \widehat{\otimes} (\widehat{\mathfrak{h}}[1] \oplus \widehat{\mathfrak{h}}_{\text{cont}}^{\vee}[2]),$$

with  $\vec{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1$  the polynomial dense subspace. The differential is  $D = d_{\mathbb{R}^2} + \bar{\partial}_{\mathbb{C}^2}$  (so  $Q_0 = 0$  in Definition 1.10). The BV pairing is the local pairing on compactly supported fields, obtained by wedge product of forms, integration over  $D \times B$  against  $dz_1 dz_2$ , and the continuous evaluation  $\widehat{\mathfrak{h}}_{\text{cont}}^{\vee} \otimes \widehat{\mathfrak{h}} \rightarrow \mathbb{C}$ ; for  $\alpha \in \Omega^p \widehat{\otimes} \widehat{\mathfrak{h}}[1]$  and  $\beta \in \Omega^q \widehat{\otimes} \widehat{\mathfrak{h}}_{\text{cont}}^{\vee}[2]$  one has  $|\alpha| = p - 1$ ,  $|\beta| = q - 2$ , and the top-form condition  $p + q = 4$  makes the pairing degree  $-1$ .

Master equation: with

$$F_\alpha = D\alpha + \tfrac{1}{2}\{\alpha, \alpha\}, \quad D_\alpha = D + \{\alpha, -\}, \quad S = \int_{D \times B} \langle \beta, F_\alpha \rangle dz_1 dz_2,$$

Jacobi for the holomorphic Poisson bracket and the derivation property of  $D$  give  $D_\alpha F_\alpha = 0$ ; compact support and coadjoint-invariance of the evaluation pairing fix the BV transpose

$$\int \langle D_\alpha^\vee \beta, \xi \rangle dz_1 dz_2 = -(-1)^{|\beta|} \int \langle \beta, D_\alpha \xi \rangle dz_1 dz_2,$$

whence  $\frac{1}{2}\{S, S\}_{\text{BV}} = \int \langle D_\alpha^\vee \beta, F_\alpha \rangle dz_1 dz_2 = -(-1)^{|\beta|} \int \langle \beta, D_\alpha F_\alpha \rangle dz_1 dz_2 = 0$ . The free and BF pieces

$$S_{\text{free}} = \int \langle \beta, D\alpha \rangle dz_1 dz_2, \quad I_{\text{BF}} = \frac{1}{2} \int \langle \beta, \{\alpha, \alpha\} \rangle dz_1 dz_2,$$

contain the differential once and the Hamiltonian interaction once.

Mixed locality: for a constant translation  $v$  in  $\mathbb{R}^2$  and an anti-holomorphic coordinate field  $\partial/\partial \bar{z}_i$ ,  $[D, \iota_v] = \mathcal{L}_v$  and  $[D, \iota_{\partial/\partial \bar{z}_i}] = \mathcal{L}_{\partial/\partial \bar{z}_i}$ . Both contractions are derivations of the local observable algebra and commute with  $\{I_{\text{BF}}, -\}$  up to those Lie derivatives, so  $[Q_{\text{obs}}, \iota_v] = \mathcal{L}_v$  and  $[Q_{\text{obs}}, \iota_{\partial/\partial \bar{z}_i}] = \mathcal{L}_{\partial/\partial \bar{z}_i}$ . The de Rham Poincare lemma on  $D$  and the Dolbeault Poincare lemma on  $B$  give the quasi-isomorphism for small inclusions and the holomorphic dependence on  $B$ , as required by Definition 1.10.

Open BV data: the brane is a point inside the holomorphic factor, so its derived endomorphism algebra is the Yoneda algebra of the skyscraper  $\mathcal{O}_p$  on  $\widehat{\mathbb{C}^2}_0$ , identified with  $\Lambda_{\mathbb{C}}(\varepsilon_1, \varepsilon_2)$  by Proposition 3.9; the brane open fields are

$$\Omega^*(\mathbb{R}) \otimes \mathbf{Ext}_{\mathbb{C}[z_1, z_2]}^*(\mathcal{O}_p, \mathcal{O}_p) \otimes \mathfrak{gl}_N[1],$$

the derived  $\text{RHom}_Y(\mathcal{O}_p, \mathcal{O}_p)$ -data of Definition 1.10, the derived endomorphism algebra of the skyscraper point. The cyclic structure is the trace on  $\mathfrak{gl}_N$  tensored with the Berezin trace on the exterior algebra; §3.4 derives both the action and the Koszul differential of this open theory directly from the formal-stalk-chart premise. The bulk-to-brane evaluation  $f \mapsto \overline{\text{Tr } f(\phi_1, \phi_2)}$  is Construction 3.25 and its bracket compatibility is Theorem 5.190; the unreduced cotangent factorization-centre lift is the obstruction problem Problem 5.48.  $\square$

### 1.5 Notation for the local theorem

The local model uses three disjoint alphabets. Letters from one alphabet are operations; letters from another are evaluations; the remaining letters are distributional cotangent modes. Coincident  $(a, b)$ -labels across alphabets are an indexing convention, never an identification.

| Symbol                                                                   | Meaning                                                                                                                                           |
|--------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------|
| $R = \mathbb{C}[[z_1, z_2]]$                                             | Complete local ring of the formal symplectic polydisk.                                                                                            |
| $\mathfrak{h} = R/\mathbb{C} \cdot 1$                                    | Reduced Hamiltonian Lie algebra.                                                                                                                  |
| $\tilde{\mathcal{A}} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1$          | Polynomial Hamiltonian subalgebra used in trace and Moyal computations.                                                                           |
| $\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1]$ | Shifted cotangent Lie algebra of Hamiltonian canonical transformations.                                                                           |
| $c^I$ (operation)                                                        | CE coordinate dual to a Hamiltonian generator $H_I \in \mathfrak{h}$ ; produces a Schouten vector field.                                          |
| $u_I$ (boundary observable)                                              | CE coordinate dual to a shifted-cotangent generator $\eta_I \in \mathfrak{h}_{\text{cont}}^\vee[1]$ ; produces a boundary Hamiltonian observable. |
| $\theta^I, O_I$                                                          | PV coordinates; the CE/PV map sends $c^I \mapsto \theta^I, u_I \mapsto O_I$ .                                                                     |
| $O_{a,b}$ (trace observable)                                             | Open trace observable $\overline{\text{Tr}(\phi_1^a \phi_2^b)}$ , $a + b > 0$ .                                                                   |
| $\Psi_{a,b}$ (PBW class)                                                 | Primitive one- $\psi$ PBW special-fibre descendant in bidegree $(a, b)$ .                                                                         |
| $\rho_{a,b}$ (cotangent mode)                                            | Principal part $z_1^{-a-1} z_2^{-b-1} dz_1 dz_2$ , basis element of $\mathfrak{h}_{\text{cont}}^\vee$ .                                           |
| $B_f(a)$                                                                 | Smeared boundary Hamiltonian $\int a(t) \overline{\text{Tr } f(\phi_1(t), \phi_2(t))} dt$ .                                                       |



*Remark 1.33 (Two avatars of a Hamiltonian: operation and evaluation).* (i) *The two CE coordinates of a Hamiltonian.* The boundary Hamiltonian observable on the brane is the image of the shifted-cotangent coordinate  $u_f$ ; the Hamiltonian CE coordinate  $c_f$  acts on observables as the Schouten vector field  $\theta^I$ . Their bracket compatibility  $[\theta^I, O_f]_S = \delta_f^I$  is the Schouten–observable pairing organising the  $P_0$ -centrality homotopies of Problem 5.48; the dictionary is Lemma 4.1.

(ii) *Bidegree coincidence*  $\Psi_{a,b} \leftrightarrow \rho_{a,b}$ . The polynomial PBW descendant  $\Psi_{a,b}$  and the principal part  $\rho_{a,b}$  share the bidegree  $(a, b)$  and carry distinct  $\mathfrak{h}$ -module structures (Corollary 5.172), connected by the Fourier-polarized Rees bridge of Theorem 8.12. See Example 2.2 for the lowest non-trivial worked case.

### 1.6 Three notions of locality

The local model carries three independent locality differentials: contact in the brane direction, cohomological triviality of topological translations on  $\mathbb{R}_{\text{top}}^2$ , and residue support of antiholomorphic translations on  $\widehat{\mathbb{C}}_{\text{o,hol}}^2$ . The three are independent coordinates of the locality structure.

(a) *Support locality along the brane.* For an open interval  $I \subset \mathbb{R}$  on the world-line, define the compactly supported Hamiltonian current algebra  $\mathfrak{h}_I = \Omega_c^\bullet(I) \widehat{\otimes} \widehat{\mathfrak{h}}$  with bracket induced from the de Rham product and the Hamiltonian bracket. In the degree-zero test-function notation used for explicit smeared trace evaluations this reads

$$[a \otimes f, b \otimes g] = ab \otimes \{f, g\}_{\widehat{\mathfrak{h}}}.$$

The reduced Hamiltonian open target on  $I$  is  $\mathcal{A}_\partial^{\text{Ham}}(I) = \widehat{\mathbf{S}}(\mathfrak{h}_I)$ , generated by the scalar-quotiented smeared trace functionals

$$B_f(a) = \int_I a(t) \overline{\text{Tr } f(\phi_1(t), \phi_2(t))} dt.$$

Their bracket in the reduced Hamiltonian target is the canonical equal-time bracket:

$$\{B_f(a), B_g(b)\}_{\mathcal{A}_\partial^{\text{Ham}}} = B_{\{f,g\}_{\widehat{\mathfrak{h}}}}(ab).$$

The bracket is the reduced one; before scalar quotient the unreduced bracket carries the central term  $N\omega(f, g) \int_I ab dt$ , with  $\omega$  the closed–open mismatch cocycle of Lemma 6.2. Disjoint supports multiply, since  $ab = 0$  outside the intersection; overlapping supports give a delta-function contact term. Disjoint smeared trace evaluations multiply, and Poisson brackets are supported where the test functions meet.

(b) *Topological locality.* In the topological direction (the  $\mathbb{R}^2$  factor), translations are  $Q$ -exact:

$$L_v = [Q, \iota_v].$$

Equivalently, positive jets in the topological direction vanish in local-operator cohomology by the contraction homotopy  $[d_{\mathbb{R}}, \iota_{\partial_t}] = \partial_t$  (Lemma 3.21). The precise position of an insertion is cohomologically immaterial; the collision pattern of insertions determines the product, and the factorisation algebra is locally constant in the topological direction.

(c) *Formal holomorphic locality.* In the holomorphic direction (the  $\mathbb{C}^2$  factor), the model has been completed at the brane point  $p \in \mathbb{C}^2$ . It is a theory of holomorphic jets at  $p$ . Antiholomorphic translations are  $Q$ -exact:

$$L_{\partial/\partial \bar{z}_i} = [Q, \iota_{\partial/\partial \bar{z}_i}].$$

The kinematic differential of the closed sector is  $D = d_{\mathbb{R}^2} + \bar{\partial}_{\mathbb{C}^2}$ , and on the formal-jet model  $\bar{\partial}$ -cohomology is concentrated in degree zero. The cotangent boundary sector is *distributionally local*: a coadjoint mode is a transverse principal part  $\rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2$ , paired with a brane Hamiltonian by residue,

$$\langle \rho_{a,b}, z_1^m z_2^n \rangle = \delta_{a,m} \delta_{b,n}.$$

The momentum dual field attached to the defect has support on the defect with transverse behaviour encoded by residues. The polynomial one- $\psi$  descendants  $\Psi_{k,l}$  realise the PBW special fibre of  $\mathfrak{h}^\vee[1]$  (Proposition 5.171); Corollary 5.172 records the deformation-level mismatch with the coadjoint module, exactly the distinction between polynomial and distributional realisations.

**PROPOSITION 1.34** (*Locality forced by the three differentials*). In the reduced Hamiltonian model the three locality statements above are consequences of three explicit operators, with  $(t, t')$  two brane-time points and  $s$  the transverse  $\mathbb{R}_s$ -coordinate reserved for the geometry of  $X$ :

$$\delta(t - t'), \quad L_v = [Q, \iota_v] \quad (v \in T\mathbb{R}_{\text{top}}^2), \quad L_{\partial/\partial \bar{z}_i} = [Q, \iota_{\partial/\partial \bar{z}_i}].$$

Support locality gives strict vanishing of  $P_\circ$ -brackets for disjoint smeared Hamiltonian currents. Topological locality gives the locally constant brane-line factorization structure after the compact-support density shift. Formal holomorphic locality gives Taylor Hamiltonians and residue-dual principal parts at the brane point. These are independent pieces of structure. The unreduced cotangent centrality homotopies require additional data. The trace observables exhibit all three localities; the shifted-cotangent currents exhibit only the first two (Theorem 0.1).

*Proof.* Support locality is the calculation of Remark 4.10: after smearing, the factor  $\delta(t - t')$  becomes the product  $ab$ , so disjoint supports give zero bracket. This proves the strict support-local factorization identity used by Proposition 4.9. Topological locality follows because translation along a topological direction is  $Q$ -exact; moving an insertion changes it by a  $Q$ -boundary. On compactly supported interval currents the surviving cohomology class is the density line of Lemma 4.8. Formal holomorphic locality follows in the same way for antiholomorphic translations, while holomorphic dependence is completed at the brane point and paired with cotangent modes by the residue pairing. The three operators live in different directions. Centrality homotopies for arbitrary unreduced open observables are additional structure.  $\square$

### 1.7 Unitarity

The protected local algebraic question is the BV consistency calculation on a twisted sector: classical and quantum master equations on the ghost/antifield complex with cohomologically trivialised topological time. Positive-definite inner products, Wightman positivity, and reflection positivity belong to a parent unitary theory. The protected sector studied here is its cohomological algebra of observables.

The replacements for unitarity are the consistency conditions of the BV formalism: the classical master equation

$$QI + \frac{1}{2}\{I, I\} = 0$$

and the quantum master equation

$$QI + \frac{1}{2}\{I, I\} + \hbar \Delta I = 0,$$

together with associativity of the resulting factorization algebra. The native two-dimensional holomorphic singular product is the diagonal local-cohomology expansion of Theorem 1.15; a one-dimensional VOA/OPE algebra is present only after a separate one-complex-dimensional reduction datum retaining the required transverse coefficient system. If the local model arises as a protected sector of a larger unitary theory (for example, of a topological twist of a supersymmetric quantum field theory whose local observables for the local model arise as cohomology with respect to a supercharge), then unitarity is carried by the larger theory, and the local sector retains only the cohomological algebra of protected observables. Lorentzian unitarity belongs to such a parent theory.

**THEOREM 1.35** (*BV replacement for unitarity in the protected sector*). For the local mixed holomorphic-topological sector, the consistency conditions are

$$QI + \frac{1}{2}\{I, I\} = 0, \quad QI + \frac{1}{2}\{I, I\} + \hbar \Delta I = 0$$

where the second equation is imposed only after a BV Laplacian and a renormalization scheme have been constructed, together with associativity and locality of the resulting factorization products. These conditions imply that the cohomology of local observables carries the stated  $P_0$  or quantum factorization algebra structure and that closed insertions act through the appropriate central-operation complex. Positivity, a Hilbert-space inner product, and a Lorentzian unitary evolution operator are additional structures supplied by a parent theory. If the sector is obtained as  $Q$ -cohomology of a unitary parent theory, the unitarity datum belongs to that parent together with the choice of  $Q$ -compatible adjoint.

*Proof.* The classical master equation says that the interacting differential  $Q_I = Q + \{I, -\}$  squares to zero, since  $Q_I^2 = \{QI + \frac{1}{2}\{I, I\}, -\}$ . The quantum master equation is the corresponding statement for the BV operator  $Q + \{I, -\} + \hbar \Delta$ . Factorization associativity is the compatibility of local products with disjoint inclusions and collision limits; locality says these products are supported on the collision loci allowed by the factorization algebra. These are exactly the algebraic structures used below to form  $P_0$ -brackets, central operations, and Moyal deformations. The construction never introduces a positive sesquilinear pairing on the BRST complex; ghosts and antifields are cohomological variables. Conversely, if a larger physical theory supplies a Hilbert space, a positive form, and a supercharge  $Q$  compatible with adjoints, then the local theory is its protected cohomological algebra. That parent datum is additional structure beyond the reduced Hamiltonian CE/PV theorem.  $\square$

The configuration is fixed: the formal-Darboux geometry  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}_{\text{o,hol}}^2$ , the brane line  $L$ , the Hamiltonian Lie algebra  $\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$ , and the three locality differentials. Closure of the BV theory on this configuration forces the shifted-cotangent extension of  $\mathfrak{h}$  by its continuous dual, with a non-degenerate equivariant pairing of cohomological degree  $-1$ : the next chapter constructs and characterises that extension.

## 2 THE SHIFTED-COTANGENT BF LIE ALGEBRA

The Lie algebra  $\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1]$  constructed in this section is the closed-side input of Theorem 5.6; its Capelli-projective central extension  $\widehat{\mathfrak{g}}_{\text{cent}}$  enters as the bulk Chevalley–Eilenberg target.

The holomorphic-symplectic Darboux normal form on  $(\widehat{\mathbb{C}^2}_o, \omega = dz_1 \wedge dz_2)$  determines the closed BV sector by three uniqueness statements. The Lie algebra of Hamiltonian vector fields modulo constants is  $\widehat{\mathfrak{h}} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$  (Proposition 2.7). The shifted-cotangent extension  $\mathfrak{g} = \widehat{\mathfrak{h}} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1]$  is the unique extension among Lie extensions of  $\widehat{\mathfrak{h}}$  by an abelian odd ideal carrying a non-degenerate equivariant pairing of cohomological degree  $-1$  (Theorem 2.1). The Hamiltonian BF action is the unique compatible local classical master equation (CME) action on this BV pair (Theorem 2.11). These statements reduce the construction to Cartan’s identity and divergence, the BV cotangent extension, and Grothendieck–Matlis residue duality.

*Hamiltonian vector fields from Cartan’s identity.* An infinitesimal canonical transformation of  $(\widehat{\mathbb{C}^2}_o, \omega = dz_1 \wedge dz_2)$  is a vector field  $X$  annihilating  $\omega$  under the Lie derivative. Cartan’s identity reduces this to closedness of  $\iota_X \omega$ , which on the formal polydisk is exactness:  $\iota_X \omega = -df$  for a unique  $f \bmod \mathbb{C}$ . Hence the symmetry vector field is the Hamiltonian vector field  $X_f$ , and the kernel of  $f \mapsto X_f$  is the constant line. Equivalently, by Proposition 2.7, the polyvector divergence  $\partial_\Omega: \mathbf{PV}^1 \rightarrow \mathbf{PV}^0$  on the polyvector-field complex  $\mathbf{PV}^\bullet$  of the polydisk identifies its kernel with Hamiltonian vector fields modulo constants. The closed Lie algebra is therefore

$$\tilde{\mathcal{A}} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1, \quad \widehat{\mathfrak{h}} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1, \quad (3)$$

with the constant-Poisson bracket  $\{f, g\} = \partial_{z_1} f \partial_{z_2} g - \partial_{z_2} f \partial_{z_1} g \pmod{\mathbb{C}}$ . Constants vanish because their Hamiltonian vector field is zero. This is the kernel of the divergence-Hamiltonian identification. Polynomial trace and Moyal calculations use the polynomial subalgebra  $\tilde{\mathcal{A}}$ ; completed CE/Koszul constructions use the formal completion  $\widehat{\mathfrak{h}}$ .

*The cotangent extension from canonical-transformation BV.* A BV theory of canonical transformations promotes  $\widehat{\mathfrak{h}}$  to a Lie algebra of fields by adjoining a degree-shifted cotangent partner: each Hamiltonian ghost  $c$  acquires its antifield  $u$  in the dual direction. The Hamiltonian sector of BCOV/Kodaira–Spencer gravity [1] fixes this language. The closed BV Lie algebra is the shifted cotangent extension

$$\mathfrak{g} = \widehat{\mathfrak{h}} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1], \quad (4)$$

with the Hamiltonian bracket on  $\widehat{\mathfrak{h}}$  and the coadjoint action on  $\mathfrak{h}_{\text{cont}}^\vee[1]$ , and the closed observables are the continuous Chevalley–Eilenberg cochains

$$O_{\text{grav}}^{\text{cl}} = C_{\text{CE,cont}}^\bullet(\mathfrak{g}) = \widehat{\mathbf{S}}(\mathfrak{g}_{\text{cont}}^\vee[-1]).$$

The cochain CE/PV identity proved later (Theorem 4.20) lives on the coordinate envelope  $C_{\text{CE,coord}}^\bullet(\mathfrak{g})$ —the product over finite graded-commutative words in the CE coordinates—and the continuous CE cochains are its continuity-controlled subspace.

*The cotangent module from residue duality.* The shifted cotangent partner carries an antifield labelled by the continuous dual  $\mathfrak{h}_{\text{cont}}^\vee$ . This dual is the principal-part module selected by residue duality. The natural pairing  $\widehat{\mathfrak{h}} \otimes \mathfrak{h}_{\text{cont}}^\vee \rightarrow \mathbb{C}$  must be continuous for the formal-disk filtration and equivariant for the Hamiltonian action. Both requirements name a unique module.

**THEOREM 2.1** (*Defect momentum is the principal-part module*). Let  $\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$  carry the  $\mathfrak{m}$ -adic topology,  $\mathfrak{m} = (z_1, z_2)$ . The continuous  $\mathfrak{h}$ -equivariant dual is, up to the residue scalar of Theorem 5.176, uniquely the Grothendieck–Matlis principal-part module

$$\mathfrak{h}_{\text{cont}}^\vee \cong \mathcal{P} = \bigoplus_{a+b>0} \mathbb{C} \rho_{a,b}, \quad \rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2,$$

with the residue pairing  $\langle \rho_{a,b}, z_1^m z_2^n \rangle = \delta_{a,m} \delta_{b,n}$  and the residue-coadjoint action  $z_1^p z_2^q \cdot \rho_{a,b} = -(pb - qa + p - q) \rho_{a-p+1, b-q+1}$ . Equivalently, this is the unique continuous total-degree-zero  $\mathfrak{h}$ -module structure on a continuous dual of  $\widehat{\mathfrak{h}}$  compatible with the residue pairing on the formal polydisk.

*Proof.* This is Theorem 5.178: local cohomology [19] of the maximal ideal  $\mathfrak{m} = (z_1, z_2)$  identifies the continuous dual of the formal completion with the principal-part module  $\mathcal{P}$ ; the residue pairing produces the displayed coadjoint formula on monomials, and any other continuous total-degree-zero  $\mathfrak{h}$ -equivariant pairing is a scalar multiple of the residue pairing. After the residue scalar is fixed, the isomorphism is canonical.  $\square$

The PBW Rees algebra of  $\mathfrak{g}$  has special fibre

$$U_{\hbar}(\mathfrak{g})/(\hbar) \cong \widehat{\mathbf{S}}(\widehat{\mathfrak{h}} \oplus \mathfrak{h}^\vee[1]),$$

where  $U_{\hbar}(\mathfrak{g})$  denotes the Rees deformation of the universal enveloping algebra and  $\hbar$  the formal deformation parameter. The bidegree- $(a, b)$  generators of the one- $\psi$  sector inside this commutative special-fibre algebra are denoted  $\Psi_{a,b}$ ; the deformation-level cotangent module is  $P = \mathfrak{h}_{\text{cont}}^\vee$ , realised by the principal parts  $\rho_{a,b}$  of Theorem 2.1. Theorem 8.12 supplies only the Fourier-polarized Rees bridge between these label systems; PBW descendants and principal parts have distinct  $\mathfrak{h}$ -module structures.

A Taylor Hamiltonian is what the brane samples. A principal part is the dual distributional momentum concentrated at the brane: the residue functional that detects a Taylor coefficient. Cotangent fields on a formal brane are residues. Each  $\rho_{a,b}$  is a defect-supported transverse mode: a distributional object on the brane line whose holomorphic transverse behaviour is encoded by a single principal-part coefficient. An abstract copy  $\bar{A}_{\text{desc}}$  of  $\bar{A}$  sees the same labels through the polynomial PBW special-fibre descendants  $\Psi_{k,l}$  just introduced (full open-side construction: Construction 3.25); they carry the same  $(a, b)$ -bidegree as the principal parts but a different  $\mathfrak{h}$ -module structure (Corollary 5.172); see §1.5 for the alphabet separation.

*Example 2.2* (*The lowest non-trivial residue mode*). Take the linear Hamiltonian  $z_1$  and act on the lowest reduced residue  $\rho_{1,0}$ . The coadjoint formula of Theorem 2.1 with  $(p, q, a, b) = (1, 0, 1, 0)$  gives

$$z_1 \cdot \rho_{1,0} = -(0 - 0 + 1 - 0) \rho_{1,1} = -\rho_{1,1}.$$

The polynomial PBW class  $\Psi_{1,0}$ , labelled identically, carries a different action: under the open one- $\psi$  descendant module of Proposition 5.171 the linear Hamiltonian shifts  $\Psi_{k,l}$  by the bracket on the open-trace ring. Same label, two  $\mathfrak{h}$ -module structures: the PBW descendant action and the residue-coadjoint action. The residue mode is what the defect momentum sees; the polynomial PBW class is what the open trace ring records. The Fourier-polarized Rees bridge of Theorem 8.12 is the intermodule passage between the two. This worked case computes the residue-side action of  $\rho_{a,b}$  explicitly, complementing Remark 1.33.

*Remark 2.3 (Scalar Hamiltonian and centre-of-mass term).* In  $\bar{A}$ ,  $\{z_1, z_2\} = 1 \equiv 0$  by quotient. On the unreduced brane phase space,  $\{\text{Tr}(\phi_1), \text{Tr}(\phi_2)\} = N$ . Constants vanish closed-side; the same constant survives open-side as the brane centre-of-mass symplectic pair. Every bracket involving linear Hamiltonians is read after projection to the Hamiltonian quotient by scalar trace classes. Quantisation returns this asymmetry as the Capelli shift (Theorem 6.6 below; cf. Remark 5.195).

*Remark 2.4 (Cotangent obstruction).* Theorem 2.1 resolves the identification of  $\mathfrak{h}_{\text{cont}}^\vee$ . The unreduced lift of the cotangent factorisation centre is the problem of promoting the closed cochain CE/PV identity (Theorem 4.20) to a chain-level morphism  $\Phi_{\text{cl}}: \text{Obs}_{\text{closed}, H}^{\text{cl}}|_L \rightarrow Z_{\text{fact}}^{P_o}(\text{Obs}_{\text{open}, N}^{\text{cl}})$  asks for centrality homotopies on the cotangent (one-antifield) sector. The exact obstruction datum is named in Problem 5.48; the strict all-window Bochner–Martinelli current transfer obstruction is  $\text{Ob}_{\text{BM}}^\Pi$ . A global central extension to the bracket-admissible  $B \subset \text{PV}_{\text{coord}}(A_{\partial, \text{coord}})$  of Definition 4.18 requires the vanishing of  $\text{Ob}_{\text{BM}}^\Pi$  on the strict ambient product completion.

### 2.1 Closed mixed Hamiltonian sector

Closed insertions on  $\mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$  are infinitesimal canonical transformations of the formal holomorphic symplectic polydisk  $(\widehat{\mathbb{C}^2}_o, \omega = dz_1 \wedge dz_2)$ . Divergence-freeness under the holomorphic volume  $\Omega = dz_1 \wedge dz_2$  identifies the infinitesimal symmetries with the reduced Hamiltonian Lie algebra  $\widehat{\mathfrak{h}} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$  (Proposition 2.7). The BV cotangent extension of a classical deformation problem then attaches a degree-shifted dual partner; continuity of the polydisk topology fixes that dual to be the Grothendieck–Matlis principal-part module  $\mathcal{P} = \mathfrak{h}_{\text{cont}}^\vee$ , with the residue basis

$$\rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2 \quad (a, b \geq 0, a + b > 0)$$

dual to the Taylor monomial  $H_{a,b} = z_1^a z_2^b$  under the residue pairing of Proposition 5.175. The same labels carry distinct  $\widehat{\mathfrak{h}}$ -module structures: the polynomial one- $\psi$  descendant  $\Psi_{a,b}$  of Construction 3.25 carries the open-trace bracket; the Matlis residue class  $\rho_{a,b}$  carries the residue coadjoint action; the two are separated by Lemma 3.37. The polarization theorem 8.1 forces the residue-side choice over the polynomial PBW class: the Moyal coadjoint action acts on the residue module, and its image on the polynomial substitute lies inside the same-action obstruction class. Both inputs are forced: the Hamiltonian factor by Proposition 2.7, the cotangent factor by the residue-pairing rigidity Theorem 5.176 and the polarization Theorem 8.1. The closed BV field complex is the unique local realization of the resulting shifted-cotangent Lie algebra  $\mathfrak{g} = \widehat{\mathfrak{h}} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1]$  on  $\mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$  (Theorem 2.8).

*Remark 2.5 (Polyvector and divergence conventions).* Holomorphic polyvector fields  $\mathbf{PV}^{*,*}(\mathbb{C}^2) \cong \Omega^{0,*}(\mathbb{C}^2)[\partial_{z_1}, \partial_{z_2}]$  carry the divergence operator  $\partial_\Omega = \iota_\Omega^{-1} \circ \partial_{\text{hol}} \circ \iota_\Omega$  determined by  $\Omega = dz_1 \wedge dz_2$ . In the formal polynomial model Dolbeault cohomology is represented by Taylor polynomials.  $\partial_\Omega$  vanishes on functions; on poly-1-vectors  $X = a_1 \partial_{z_1} + a_2 \partial_{z_2}$ ,  $\partial_\Omega X = \partial_{z_1} a_1 + \partial_{z_2} a_2$ . The image of  $\partial_\Omega$  from  $\mathbf{PV}^0$  is zero, since  $\partial_\Omega$  vanishes on functions. Constants are removed by  $f \mapsto X_f$  to give the reduced polynomial algebra  $\bar{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1$ .

LEMMA 2.6 (*Polynomial Poincare primitive*). Let

$$\theta = a(z_1, z_2) dz_1 + b(z_1, z_2) dz_2$$

be a polynomial one-form on  $\mathbb{C}^2$ . If  $\partial_{z_1} a = \partial_{z_1} b$ , then there is a polynomial  $f \in \mathbb{C}[z_1, z_2]$ , unique modulo constants, such that  $\theta = df$ .

*Proof.* Define

$$f(z_1, z_2) = \int_0^1 (z_1 a(tz_1, tz_2) + z_2 b(tz_1, tz_2)) dt.$$

Since  $a$  and  $b$  are polynomials, the integral is termwise division of each homogeneous term by its degree plus one, hence  $f$  is polynomial. Differentiate under the integral sign. Closedness says  $\partial_{z_2} a = \partial_{z_1} b$ , so

$$\begin{aligned} \partial_{z_1} f &= \int_0^1 (a(tz) + tz_1 \partial_{z_1} a(tz) + tz_2 \partial_{z_1} b(tz)) dt \\ &= \int_0^1 (a(tz) + tz_1 \partial_{z_1} a(tz) + tz_2 \partial_{z_2} a(tz)) dt \\ &= \int_0^1 \frac{d}{dt} (t a(tz)) dt = a(z). \end{aligned}$$

The same calculation, with  $a$  and  $b$  interchanged, gives  $\partial_{z_2} f = b(z)$ . Therefore  $df = \theta$ . If  $df = dg$ , then  $d(f - g) = 0$ , hence  $f - g$  is constant.  $\square$

**PROPOSITION 2.7** (*Hamiltonian polyvector reduction*). In the formal polynomial model on  $(\mathbb{C}^2, \omega = dz_1 \wedge dz_2)$ , the divergence-closed polyvector-one fields reduce to Hamiltonian vector fields:

$$\mathbf{Ker}(\partial_\Omega : \mathbf{PV}^1 \rightarrow \mathbf{PV}^0) \cong \{df : f \in \mathbb{C}[z_1, z_2]\} \cong \bar{A}.$$

Under the convention

$$X_f = (\partial_{z_1} f) \partial_{z_2} - (\partial_{z_2} f) \partial_{z_1}, \quad \iota_{X_f} \omega = -df,$$

the commutator of vector fields is

$$[X_f, X_g] = X_{\{f, g\}_{\bar{A}}}.$$

*Proof.* Let  $X = a_1 \partial_{z_1} + a_2 \partial_{z_2}$ . The divergence operator is  $\partial_\Omega X = \partial_{z_1} a_1 + \partial_{z_2} a_2$ . Contracting with  $\omega$  gives

$$\iota_X \omega = a_1 dz_2 - a_2 dz_1.$$

The equation  $\partial_\Omega X = 0$  is exactly the closedness of this one-form:

$$d(\iota_X \omega) = (\partial_{z_1} a_1 + \partial_{z_2} a_2) dz_1 \wedge dz_2 = 0.$$

By Lemma 2.6, there is a polynomial  $f$ , unique modulo constants, such that  $\iota_X \omega = -df$ . Therefore

$$a_1 = -\partial_{z_2} f, \quad a_2 = \partial_{z_1} f,$$

so  $X = X_f$ . Constants are exactly the kernel of  $f \mapsto X_f$ , giving  $\bar{A}$ .

Finally, for any polynomial  $h$ ,

$$\begin{aligned} [X_f, X_g](h) &= \{f, \{g, h\}\} - \{g, \{f, h\}\} \\ &= \{\{f, g\}, h\} \end{aligned}$$

by the Jacobi identity for the constant Poisson bracket. Thus  $[X_f, X_g] = X_{\{f, g\}}$ , and projecting constants gives the bracket on  $\bar{A}$ .  $\square$

Replacing  $\bar{A}$  by  $\Omega^{\circ,*}(\mathbb{C}^2) \hat{\otimes} \bar{A}$  extends the same statement to Dolbeault coefficients: the defining equations are coefficient-wise, constants drop by the same argument, and the bracket is the Dolbeault-coefficient extension of  $\{-, -\}_{\bar{A}}$ .

**THEOREM 2.8** (*Closed BV field complex of canonical transformations*). Let  $D = d_{\mathbb{R}^2} + \bar{\partial}_{\mathbb{C}^2}$  be the kinematic differential on  $\mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$ . The BV cotangent extension of  $\hat{\mathfrak{h}}$ , applied to the formal local data of  $(\widehat{\mathbb{C}}_{\circ}^2, \omega)$ , is the field complex

$$\mathcal{E}_{\text{cl}} = \Omega^*(\mathbb{R}^2) \hat{\otimes} \Omega^{\circ,*}(\mathbb{C}^2) \hat{\otimes} (\hat{\mathfrak{h}}[1] \oplus \mathfrak{h}_{\text{cont}}^{\vee}[2]), \quad (5)$$

with  $\hat{\mathfrak{h}}[1]$  the Hamiltonian connection sector  $\alpha$  and  $\mathfrak{h}_{\text{cont}}^{\vee}[2]$  the BV cotangent partner  $\beta$ . This complex is the unique local realization, in the Dolbeault–de Rham category, of the shifted-cotangent Lie algebra  $\mathfrak{g} = \hat{\mathfrak{h}} \ltimes \mathfrak{h}_{\text{cont}}^{\vee}[1]$  carrying:

- (i) the BV pairing  $\int_{\mathbb{R}^2 \times \mathbb{C}^2} \langle \beta, \alpha \rangle dz_1 dz_2$  of degree  $-1$ , with  $\hat{\mathfrak{h}}_{\text{cont}}^{\vee} \otimes \hat{\mathfrak{h}} \rightarrow \mathbb{C}$  the residue pairing of Proposition 5.175;
- (ii) the kinematic differential  $D$  acting coefficient-wise;
- (iii) the bracket on the Lie factor  $\hat{\mathfrak{h}}$  given by the holomorphic Poisson bracket of  $\omega$  extended to Dolbeault coefficients, and the coadjoint action on the cotangent factor.

*Proof.* The Hamiltonian factor  $\hat{\mathfrak{h}}$  is given by Proposition 2.7: the divergence-free polyvector-one fields, modulo constants, are exactly the Hamiltonian vector fields of  $\bar{A}$ , and the Dolbeault upgrade is coefficient-wise. The cotangent factor  $\mathfrak{h}_{\text{cont}}^{\vee} = \mathcal{P}$  is fixed by Proposition 5.175 and the residue-pairing rigidity Theorem 5.176: among continuous total-degree-zero  $\hat{\mathfrak{h}}$ -equivariant pairings the Matlis residue pairing is unique up to a single scalar. The BV pairing on  $\mathcal{E}_{\text{cl}}$  is then the integrated form of this residue pairing against the de Rham/Dolbeault top form. The degree shifts  $[1]$  on  $\hat{\mathfrak{h}}$  and  $[2]$  on  $\mathfrak{h}_{\text{cont}}^{\vee}$  are fixed by the BV cotangent principle: a Maurer–Cartan element of  $\mathfrak{g} = L \ltimes L^{\vee}[1]$  in a degree-zero local field theory on a four-real-dimensional spacetime sits in  $\Omega^* \hat{\otimes} \Omega^{\circ,*} \hat{\otimes} L[1]$ , and the partner sits in  $\Omega^* \hat{\otimes} \Omega^{\circ,*} \hat{\otimes} L^{\vee}[2]$  so that the integrated pairing has degree  $-1$ . The computation in Proposition 1.32 verifies these degrees on the four-form factor.  $\square$

The boundary image of  $\beta$ , under the bulk-to-brane evaluation of Section 4.3, is the Matlis principal-part defect current of Theorem 5.185: cotangent fields are measured by that current and the cotangent module is the Matlis residue module. The BF action of Section 2.2 is the unique cyclic local functional whose classical master equation says that  $\alpha$  is a Maurer–Cartan element of  $\mathfrak{g}$ .

**Remark 2.9** (*Cotangent module as Matlis dualizing object*). The cotangent factor  $\mathcal{P}$  is the local-cohomology module  $H_{\mathfrak{m}}^2(R) dz_1 dz_2$  on  $R = \mathbb{C}[[z_1, z_2]]$  with the constant Hamiltonian line removed: the Matlis dualizing object on  $R$ , paired with  $\bar{A} = R/\mathbb{C} \cdot 1$  by residue.

**Remark 2.10** (*Casimir/Tate obstruction to a strict cotangent kernel*). The pairing of Theorem 2.8 is non-degenerate but the diagonal coevaluation tensor  $C_{\hat{\mathfrak{h}}} = \sum_{a+b>0} H_{a,b} \otimes \rho_{a,b}$  diverges in the strict tensor product  $\hat{\mathfrak{h}} \hat{\otimes} \mathfrak{h}_{\text{cont}}^{\vee}$ : this is the Tate-Casimir obstruction Lemma 5.211. The closed BV complex is therefore well posed as a continuous bilinear pairing. A strict kernel requires repair to a kernel-admissible model through the weighted-Tate envelope of Definition 5.54; in the strict ambient product completion, the  $P_{\circ}$ -algebra structure is obstructed (Definition 4.18). Every unreduced cotangent-kernel claim must clear this obstruction.



### 2.2 Hamiltonian BF action

The Hamiltonian BF action is forced by three data: the holomorphic-symplectic Darboux normal form on  $(\widehat{\mathbb{C}^2}_o, \omega = dz_1 \wedge dz_2)$ , the cyclic BV trace on  $\mathcal{E}_{\text{cl}}$  of (5), and linearity in the cotangent partner. Theorem 2.8 fixes the BV pair  $(\mathcal{E}_{\text{cl}}, \omega_{\text{BV}})$  and the shifted-cotangent Lie algebra  $\mathfrak{g} = \widehat{\mathfrak{h}} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1]$ ; the canonical degree- $(-1)$  evaluation pairing  $\langle \beta, f \rangle = \beta(f)$  between cotangent and Hamiltonian factors, wedged with the form factors and integrated over  $\mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$  against  $dz_1 \wedge dz_2$ , extends to the cyclic BV pairing  $\omega_{\text{BV}}$ . Under cyclicity, locality, and linearity in the cotangent partner, the classical master equation  $\frac{1}{2}\{S, S\}_{\text{BV}} = 0$  has a unique cyclic local solution modulo  $D$ -exact additions: the BF action of (6) below. Equivalently, the Hamiltonian BF action is the unique compatible local CME action on the BV pair  $(\mathcal{E}_{\text{cl}}, \omega_{\text{BV}})$ , forced by holomorphic-symplectic Darboux normal form, cyclic BV trace, and linearity in cotangent partner.

**THEOREM 2.11** (*Cotangent BF action: unique cyclic Maurer–Cartan functional*). Let  $\langle -, - \rangle$  be the integrated residue pairing between  $\Omega^*(\mathbb{R}^2) \widehat{\otimes} \Omega^{0,*}(\mathbb{C}^2) \widehat{\otimes} \mathfrak{h}_{\text{cont}}^\vee[2]$  and  $\Omega^*(\mathbb{R}^2) \widehat{\otimes} \Omega^{0,*}(\mathbb{C}^2) \widehat{\otimes} \widehat{\mathfrak{h}}[1]$ , contracted by wedging form factors and integrating against  $dz_1 \wedge dz_2$ . The cyclic local functional

$$S = \int_{\mathbb{R}^2 \times \mathbb{C}^2} \left\langle \beta, D\alpha + \frac{1}{2}\{\alpha, \alpha\} \right\rangle dz_1 dz_2 \quad (6)$$

is, up to a  $D$ -exact form, the unique cyclic local functional on  $\mathcal{E}_{\text{cl}}$  satisfying:

- (i)  $S$  is local in the de Rham/Dolbeault sense and cyclic for the integrated residue pairing;
- (ii)  $S$  is linear in the cotangent partner  $\beta$ ;
- (iii) the classical BV master equation  $\frac{1}{2}\{S, S\}_{\text{BV}} = 0$  holds and encodes the Maurer–Cartan equation  $F_\alpha = D\alpha + \frac{1}{2}\{\alpha, \alpha\} = 0$  on  $\widehat{\mathfrak{h}}$  together with the dual transport  $D_\alpha^\vee \beta = 0$  on  $\mathfrak{h}_{\text{cont}}^\vee$ .

The form-valued Hamiltonian bracket on  $\widehat{\mathfrak{h}}$  is

$$[\eta \otimes f, \theta \otimes g] = (-1)^{|f||\theta|} (\eta \wedge \theta) \otimes \{f, g\},$$

with Koszul sign reducing to  $+1$  on the ghost-zero (even-coefficient) sector.

*Proof. Existence.* The action (6) is constructed in Proposition 1.32, where the classical master equation  $\frac{1}{2}\{S, S\}_{\text{BV}} = 0$  is verified by the coadjoint invariance of the residue pairing and the Cartan identity  $D_\alpha F_\alpha = 0$  on  $\widehat{\mathfrak{h}}$ .

*Uniqueness.* Cyclicity, locality, and linearity in  $\beta$  restrict  $S$  to a sum

$$S = \int \langle \beta, P\alpha \rangle + \int \langle \beta, Q(\alpha, \alpha) \rangle \quad (\text{mod } D\text{-exact}),$$

with  $P$  a degree-one  $\mathbb{C}$ -linear local operator on  $\Omega^* \widehat{\otimes} \Omega^{0,*} \widehat{\otimes} \widehat{\mathfrak{h}}[1]$  and  $Q$  a degree-zero local bilinear bracket on the same space. Higher powers  $\beta^k$ ,  $k \geq 2$ , are excluded by linearity in the cotangent factor; pure  $\alpha$ -polynomials are excluded because cyclicity for the  $(\beta, \alpha)$  pairing pairs every term to a single  $\beta$ . The classical master equation on this ansatz reads

$$\frac{1}{2}\{S, S\}_{\text{BV}} = \int \left\langle \beta, P^2 \alpha + (PQ(\alpha, \alpha) + 2Q(\alpha, P\alpha)) + 2Q(\alpha, Q(\alpha, \alpha)) \right\rangle,$$

where the three displayed pieces are homogeneous in  $\alpha$  of degrees 1, 2, 3. Vanishing of the degree-1 piece forces  $P^2 = 0$ , and locality plus the kinematic axioms (i),(iii) then identify  $P$  with the differential

$D = d_{\mathbb{R}^2} + \bar{d}_{\mathbb{C}^2}$  acting coefficient-wise. Vanishing of the degree-2 piece forces  $D$  to be a derivation of  $Q$ ; vanishing of the degree-3 piece forces  $Q$  to satisfy Jacobi. The unique local bilinear  $D$ -derivation on  $\widehat{\mathfrak{h}}$  that satisfies Jacobi and extends coefficient-wise the holomorphic Poisson bracket of  $\omega$  is  $\{-, -\}$  itself. Therefore  $S$  is given by (6) modulo a  $D$ -exact term.  $\square$

The classical equation of motion is the flatness statement  $F_\alpha = 0$ :  $\alpha$  is a  $\mathfrak{g}$ -Maurer–Cartan element, equivalently a connection on a bundle of formal holomorphic-symplectic surfaces over  $\mathbb{R}^2$ . The cotangent partner  $\beta$  is the Lagrange multiplier enforcing this equation; the BV gauge symmetries are

$$\delta_\varepsilon \alpha = D\varepsilon + \{\alpha, \varepsilon\}, \quad \delta_\varepsilon \beta = \text{ad}_\varepsilon^\vee \beta,$$

and the cotangent gauge symmetry

$$\delta_\lambda \alpha = 0, \quad \delta_\lambda \beta = D_\alpha \lambda,$$

with  $D_\alpha = D + \{\alpha, -\}$  and  $\lambda \in \Omega^* \widehat{\otimes} \Omega^{0,*} \widehat{\otimes} \mathfrak{h}_{\text{cont}}^\vee[1]$ . The second symmetry is reducible on shell:  $\lambda = D_\alpha \mu$  gives  $D_\alpha \lambda = \{F_\alpha, \mu\}$ , which vanishes when  $F_\alpha = 0$ .

*Remark 2.12 (Component reading).* In ghost-zero coordinates on  $\mathbb{R}_{x_i}^2 \times \mathbb{C}_{z_i}^2$ ,

$$\alpha = \alpha_{x_i} dx_i + \alpha_{\bar{z}_j} d\bar{z}_j, \quad \beta = \beta_{x_1 x_2} dx_1 dx_2 + \beta_{x_i \bar{z}_j} dx_i d\bar{z}_j + \beta_{\bar{z}_1 \bar{z}_2} d\bar{z}_1 d\bar{z}_2.$$

The component  $\alpha_{\bar{z}_j} d\bar{z}_j$  is the Hamiltonian potential whose vector field is the Beltrami differential of a complex-structure deformation of  $\mathbb{C}^2$ ;  $\alpha_{x_i} dx_i$  is the connection on  $\mathbb{R}^2$  valued in Hamiltonian vector fields, organizing the bundle of  $\mathbb{C}^2$ s over  $\mathbb{R}^2$ .  $F_\alpha = 0$  asserts joint integrability and flat holonomy. The bulk theory is defined on the full mixed space-time  $\mathbb{R}^2 \times \mathbb{C}^2$ ; the brane line  $L = \mathbb{R} \times \{0\} \times \{p\}$  enters only via the boundary evaluation of Section 4.3.

*Remark 2.13 (Named obstruction to a renormalized BF theory).* Theorem 2.11 produces a formal-local cyclic functional satisfying the classical BV master equation. Its promotion to a renormalized field theory requires the analytic package of Problem 5.212: an elliptic gauge fixing, a Costello propagator  $P_{\varepsilon, L}$  on  $\mathbb{R}^2 \times \mathbb{C}^2$ , a finite-scale graph effective action with graphwise renormalization, and the QME  $(d + \hbar \Delta)e^{I/\hbar} = 0$  to all orders. The two named obstructions are: (a) the Tate–Casimir obstruction Lemma 5.211, which prevents the strict diagonal cotangent kernel from converging in  $\widehat{\mathfrak{h}} \widehat{\otimes} \mathfrak{h}_{\text{cont}}^\vee$  and hence forces a weighted-Tate envelope on the propagator coefficient module; (b) the QME obstruction class of Problem 5.72, whose vanishing in the weighted-Tate envelope is the unreduced one-antifield quantum lift. Before (a) and (b) are supplied, (6) remains a formal-local generating functional, and the perturbative phase space is the derived MC stack of Remark 2.14.

*Remark 2.14 (Phase space).* Circle-regularizing one topological direction gives  $\mathbb{R} \times S^1 \times Y_{\text{hol}}$  with  $Y_{\text{hol}}$  holomorphic symplectic, and  $F_\alpha = 0$  becomes the Maurer–Cartan problem for Hamiltonian vector fields on  $S^1 \times Y_{\text{hol}}$ . The perturbative phase space is the derived cotangent stack

$$T_{\text{der}}^* \text{MC}(\Omega^*(S^1) \widehat{\otimes} \Omega^{0,*}(Y_{\text{hol}}, \text{Ham}(Y_{\text{hol}})), D, \{-, -\}),$$

collapsing on the underived tangent at the trivial solution to the ordinary cotangent of the linearized deformation complex. The analytic package of Remark 2.13 fixes this regularized model to a propagator-grade theory.

The closed BV theory of the local model is the Hamiltonian BF theory on  $\widehat{\mathbb{C}^2}_o$  with fields valued in the shifted-cotangent Lie algebra  $\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1]$  and master action  $S = \int \langle \beta, D\alpha + \frac{1}{2}\{\alpha, \alpha\} \rangle dz_1 dz_2$ , forced by Darboux + cyclic trace + linearity in the cotangent partner. The open boundary at a stack of  $N$  Dirac branes fixes the Chan–Paton substitution  $z_i \mapsto \phi_i \in \mathfrak{gl}_N$  with moment map  $\mu(\phi_1, \phi_2) = [\phi_1, \phi_2]$ ; the open BV theory is the derived commuting variety  $[\mu_{\text{der}}^{-1}(o)/GL_N]$ , constructed in the section below.

### 3 THE DERIVED COMMUTING VARIETY STACK AT $N$ DIRAC BRANES

The Dirac brane stack constructed here is the open  $E_t$ -algebra  $\mathcal{A}_{\partial, N}^{\text{cl}}$  of Theorem 5.6; its  $\mathfrak{gl}_N$ -equivariant centre exhaustion is Theorem 5.10.

#### 3.1 The Dirac brane stack

The primitive open object is the open category  $\mathcal{C}_\partial^{\text{op}}$  on the Kato–Nakayama log boundary; a chosen vacuum  $b$  gives the chart  $\mathcal{A}_b = \text{End}(b)$ , and at the stack of  $N$  Dirac branes  $\mathcal{A}_b$  is the matrix endomorphism algebra of the corresponding boundary state. The manuscript works in the chart determined by this primitive open object.

A holomorphic Hamiltonian  $f \in \mathbb{C}[[z_1, z_2]]$  on the formal symplectic polydisk  $(\widehat{\mathbb{C}^2}_o, dz_1 \wedge dz_2)$  is infinite-dimensional. A finite-rank reduction exists exactly when the closed polydisk is paired with an open boundary that names the two normal coordinates: a stack of  $N$  Dirac branes on the topological line

$$L = \mathbb{R}_t \times \{s = z_1 = z_2 = o\} \subset X = \mathbb{R}_t \times \mathbb{R}_s \times \mathbb{C}_{z_1, z_2}^2$$

realises the Dirac brane stack as the derived commuting variety stack  $[\mu_{\text{der}}^{-1}(o)/GL_N]$  with moment map  $\mu(\phi_1, \phi_2) = [\phi_1, \phi_2]$ . The transverse position is the Chan–Paton pair

$$(\phi_1, \phi_2) \in \mathfrak{gl}_N \oplus \mathfrak{gl}_N$$

on the holomorphic-symplectic factor (Pantev–Toën–Vaquié–Vezzosi shifted symplectic structures [44]; the brane realization is Calaque [45]; the BV/Koszul realisation of the moment-map zero fibre is Costello [12, Ch. 5]). A Hamiltonian  $f$  has trace evaluation

$$J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)}$$

on this chart. This is the formal Darboux coordinate of the brane-side  $E_t$ -Hochschild complex  $HH^\bullet(\mathcal{A}_{\partial, N}^{\text{cl}}, \mathcal{A}_{\partial, N}^{\text{cl}})$ , identified at the formal-Darboux stalk with the bulk  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_{b_p}$  (the chiral  $E_2$  factorization algebra restricted to the formal neighbourhood of  $b_p$ ) by Costello–Gwilliam bulk–boundary correspondence and Deligne’s  $E_t$ -Hochschild theorem. The input data are the formal Darboux polydisk at the brane point and the Chan–Paton substitution

$$z_1 \rightsquigarrow \phi_1, \quad z_2 \rightsquigarrow \phi_2, \quad \phi_i \in \mathfrak{gl}_N,$$

sending holomorphic coordinates to transverse Chan–Paton matrices. The symplectic potential, moment map, Dirac constraint, BV differential, and gauge-invariant operator algebra are all determined by this substitution.

*The substitution forces the open BV theory.* The pullback of  $dz_1 \wedge dz_2$  along the substitution, traced over Chan–Paton, is

$$\Omega_N = \text{Tr}(d\phi_1 \wedge d\phi_2), \quad \theta_N = \text{Tr}(\phi_1 d\phi_2).$$

$GL_N$ -conjugation acts symplectically. Its moment map is

$$\mu(\epsilon) = -\text{Tr}(\epsilon[\phi_1, \phi_2]), \quad \epsilon \in \mathfrak{gl}_N,$$

so the first-class constraint is

$$[\phi_1, \phi_2] = 0.$$

On this locus the Chan–Paton pair commutes and the  $N$  branes sit at  $N$  points of the polydisk. With multiplier  $\mathcal{A} \in \Omega^1(\mathbb{R}) \otimes \mathfrak{gl}_N$  the Dirac first-order action is

$$S_\partial = \int_{\mathbb{R}} \text{Tr}(\phi_1 d\phi_2 + \mathcal{A}[\phi_1, \phi_2]), \quad (7)$$

and the constraint is realized by the Koszul derived zero-fibre model for the moment-map equation [12, Ch. 5]. Introduce the odd matrix-valued antifield

$$\psi \in \mathfrak{gl}_N[1].$$

It is the BV antifield  $\psi = \mathcal{A}^*$  of the Lagrange multiplier  $\mathcal{A}$ . It is independent of  $\phi_1, \phi_2$ . Let  $Q$  be the derivation defined on generators by

$$Q\phi_1 = Q\phi_2 = 0, \quad Q\psi = [\phi_1, \phi_2]. \quad (8)$$

The displayed equation is the Koszul resolution of  $\mu^{-1}(0)$  on the BV side: every matrix entry of the commutator equals  $Q\psi^i_j$  and is therefore  $Q$ -exact, so the moment-map ideal vanishes in cohomology. The matrix entries of  $[\phi_1, \phi_2]$  have syzygies in  $\mathbb{C}[\phi_1^i_j, \phi_2^i_j]$  for  $N \geq 2$ ; the construction is the derived zero fibre of the moment map. Its ghost-zero invariant cohomology

$$\mathcal{O}([\mu_{\text{der}}^{-1}(0)/GL_N]) = H^0((\mathbf{S}^\bullet \mathfrak{gl}_N^{\oplus 2} \otimes \Lambda^\bullet \mathfrak{gl}_N[1])^{GL_N}, Q)$$

identifies the structure sheaf of the derived commuting variety stack with the  $GL_N$ -invariant cohomology of the Koszul complex; the degree-zero classical invariant ring of the commuting-pair quotient is the bottom row of this identification. The degree-zero Hamiltonian trace comparison ignores higher Koszul homology; that homology enters only through explicit invocation. The ghost-zero invariant ring is

$$\mathcal{R}_N = \mathbb{C}[\phi_1^i_j, \phi_2^i_j] \otimes \Lambda(\psi^i_j), \quad H^0(\mathcal{R}_N^{GL_N}, Q).$$

Lemma 3.14 produces (7) from the shifted open Chern–Simons coefficients; Lemma 3.15 verifies the moment-map content; Proposition 3.16 verifies that ghost-zero truncation is consistent. The action (7), the differential (8), and  $\mathcal{R}_N^{GL_N}$  all come from the Chan–Paton substitution and moment-map reduction.

*Word-ordering independence of  $J$ .*  $J(f)$  depends only on the symbol  $f$ . Adjacent letters in any trace word commute up to a  $Q$ -exact correction,

$$Q \operatorname{Tr}(\psi v u) = \operatorname{Tr}([\phi_1, \phi_2] v u),$$

so a trace adjacent-swap is the BRST image of an antifield insertion. Cyclic invariance and iterated  $Q$ -exact swaps reduce every cyclic word in  $\phi_1, \phi_2$  to a symmetric symbol  $z_1^k z_2^l$ : the trace family  $\{J(f)\}$  is the connected single-trace primitive content of the Chan–Paton pair, the full stable  $GL_N$ -invariant ring is the polynomial algebra on this primitive (Procesi–Razmyslov first fundamental theorem), and the closed sector  $\bar{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1$  acts on the brane through  $J$  alone.

LEMMA 3.1 (*Trace realisation of the formal polydisk*). In the stable connected ghost-zero open observables  $H^0(\mathcal{R}_N^{GL_N}, Q)$ , the substitution map

$$J: \mathbb{C}[z_1, z_2] \longrightarrow H^0(\mathcal{R}_N^{GL_N}, Q)|_{\text{stable conn.}}, \quad f \longmapsto \overline{\operatorname{Tr} f(\phi_1, \phi_2)},$$

is surjective with kernel exactly the constant line  $\mathbb{C} \cdot 1 \subset \mathbb{C}[z_1, z_2]$ . Procesi–Razmyslov first fundamental theorem identifies the full stable  $GL_N$ -invariant ring with the polynomial algebra on the trace family  $\{J(f) : f \in \bar{A}\}$ , so every gauge-invariant operator is a polynomial in trace observables and the closed sector  $\bar{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1$  acts on the brane through  $J$  alone.

*Proof.* Procesi–Razmyslov [17, 18] identifies  $\mathbb{C}[\mathfrak{gl}_N \oplus \mathfrak{gl}_N]^{GL_N}$  with the algebra generated by traces of noncommutative monomials in two letters. Cyclic invariance equates traces of cyclically equivalent words; the  $Q$ -exactness of (8) equates traces differing by one adjacent swap. Iterating, every cyclic word  $\operatorname{Tr}(w(\phi_1, \phi_2))$  reduces to a symmetric symbol  $\operatorname{Tr}(\phi_1^k \phi_2^l) = J(z_1^k z_2^l)$  in  $H^0(\mathcal{R}_N^{GL_N}, Q)$ . Stable connected linear independence over  $N$  of these symmetric symbols is standard. The constant Hamiltonian substitutes to  $N \cdot I$ , a scalar multiple of the identity, killed by passage to the connected stable class.  $\square$

The substitution-realised Lie algebra image of  $\bar{A}$  inside the open trace cohomology generates the open sector as a trace algebra.

Example 3.2 ( $N = 1$ : *the geometric prototype*). At  $N = 1$ ,  $[\phi_1, \phi_2] = 0$  holds tautologically, so the Dirac constraint imposes nothing and the antifield  $\psi$  is decoupled. The Chan–Paton pair  $(\phi_1, \phi_2) \in \mathbb{C}^2$  is a chosen point of the formal polydisk, and

$$J(f) = \overline{\operatorname{Tr} f(\phi_1, \phi_2)} = f(\phi_1, \phi_2) \in \mathbb{C}$$

is ordinary scalar evaluation modulo the constant.  $J$  reduces to the closed-string evaluation map  $f \mapsto f(p)$  at a chosen polydisk point  $p$ , and the Procesi–Razmyslov input is trivial. Every later non-commutative refinement of  $J$  collapses to this scalar evaluation when  $N = 1$ ; the matrix structure enters with the brane stack.

Remark 3.3 (*Cyclic framing branch and the Hilbert scheme*). Adjoining a cyclic vector  $v \in \mathbb{C}^N$  with  $\mathbb{C}[\phi_1, \phi_2] \cdot v = \mathbb{C}^N$  cuts out the stable cyclic locus of the derived commuting variety stack  $[\mu_{\text{der}}^{-1}(0)/GL_N]$ ; on this locus the underived free  $GL_N$ -action gives the Hilbert scheme of points,  $[(\mu^{-1}(0) \times \mathbb{C}_{\text{cyc}}^N)/GL_N] = \mathcal{Hilb}^N(\mathbb{C}^2)$ , in its Nakajima ADHM-quiver presentation. The Moyal/Weyl deformation of the brane Poisson algebra restricts on this branch to the noncommutative Hilbert scheme of Nekrasov–Schwarz [46], and the Capelli scalar  $\hbar N[\bar{c}]$  of Example 1.3 is its centre-of-mass  $U(1)$  charge. The framing data selects a stable open substack of the Dirac brane stack on which the derived fibre is underived.

$J$  is exact at the level of the scalar trace algebra. The full  $P_o$ -factorization centre requires additional cotangent centrality homotopies: the chain-level morphism from closed observables on  $L$  to the open factorization centre has a structural obstruction.

*Remark 3.4 (Cotangent obstruction beyond  $J$ ).* Lemma 3.1 resolves the scalar-trace generator question: scalar closed insertions  $F = J(f)$  commute with every open trace class up to the moment-map operation  $\{J(f), J(g)\} = J(\{f, g\})$  of Theorem 4.4 below. The closed Hamiltonian BF sector on the polydisk and the open brane stack on  $L$  generate factorization algebras of classical observables, denoted  $\text{Obs}_{\text{closed}, H}^{\text{cl}}$  and  $\text{Obs}_{\text{open}, N}^{\text{cl}}$  (constructed in §2.1, §3.4); the  $P_o$ -factorization centre  $Z_{\text{fact}}^{P_o}$  of a factorization algebra is its centralizer for the shifted Poisson bracket and the factorization product (Problem 5.48). The strengthening sought beyond Lemma 3.1 is a chain-level morphism

$$\Phi_{\text{cl}}: \text{Obs}_{\text{closed}, H}^{\text{cl}}|_L \longrightarrow Z_{\text{fact}}^{P_o}(\text{Obs}_{\text{open}, N}^{\text{cl}}),$$

asking for centrality homotopies linking closed insertions to *arbitrary* open insertions, in particular to the principal-part cotangent currents  $\Theta_\rho$  (the boundary current attached to a residue  $\rho \in \mathfrak{h}_{\text{cont}}^\vee$  by smearing, defined in Theorem 5.185 below). Substitution into the Chan–Paton pair encodes the symbol-level information of  $f$ . The cotangent-direction antifield content of the closed BV algebra is controlled by the unreduced cotangent centrality homotopy data of Problem 5.48 together with the Bochner–Martinelli obstruction vector  $\text{Ob}_{\text{BM}}^\Pi$ , a six-component cohomology vector blocking strict all-window support-local current transfer to the brane line (Theorem 8.40).  $J$  is the scalar part of  $\Phi_{\text{cl}}$ ; the cotangent part is the residual one, classified by the obstruction classes above.

### 3.2 Hamiltonian sector and stable large $N$

The trace substitution  $J$  produces an open trace algebra at every finite  $N$ ; the closed Hamiltonian algebra  $\tilde{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1$  is its image under  $J$  only after a stable trace stabilization is performed. Four hypotheses control that stabilization: stable invariant theory, BRST symmetrization, primitive cyclic projection, and a finite- $N$  window in which the projection is a chain map. Inside that window the Procesi–Razmyslov trace algebra collapses to commutative monomials, the connected primitive class is read by Loday–Quillen–Tsygan, and the resulting completed coefficient ring is the formal Hamiltonian envelope of  $\mathfrak{h}$ . Outside the window finite- $N$  trace identities re-enter and break the projection. Strict identification of the raw finite- $N$  operator algebras with the formal envelope requires an open-trace  $A_\infty$ -Koszul acceptance datum (level 5 of §4.2), which is a conditional input beyond the stable hypotheses.

Coefficient algebras:  $\tilde{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1$  on the polynomial side and  $\mathfrak{h} = \widehat{\mathfrak{h}} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$  on the completed side, with  $\mathfrak{h}^\vee = \widehat{\mathfrak{h}}_{\text{cont}}^\vee$  its continuous dual;  $\widehat{\mathbf{S}}$  denotes the completed symmetric algebra and  $C_{\text{CE}, \text{cont}}^\bullet$  the matched continuous CE complex. Constants are quotiented out per §1.5: a constant Hamiltonian has zero Hamiltonian vector field. On the open side  $\text{Tr}_N(1) = N$  is the empty trace, a central scalar tracked through scalar reduction while retaining the operator unit; for a positive word-degree trace expression  $T$ ,  $\overline{T}$  denotes its connected class. Shifts follow the BV convention: an element of  $V[1]$  has parity reversed and cohomological degree one lower than the corresponding element of  $V$ . The CE/PV cotangent block below uses its own coordinate grading: the odd BV cotangent generator  $\lambda[1]$  has BV degree  $-1$ , while the dual CE coordinate  $u_\lambda$  is assigned PV function degree 0; “dual” in that block means the CE-suspended cotangent coordinate dual in this grading convention.

At a fixed brane point the reduced BV algebra is generated by two even matrices  $\phi_1, \phi_2$  and the odd antifield  $\psi = A^*$ , with

$$Q\phi_1 = Q\phi_2 = 0, \quad Q\psi = [\phi_1, \phi_2].$$

The moment-map constraint is imposed homologically; for  $N \geq 2$  the commuting-pair equations form a nonregular sequence, so the construction is the derived zero fibre. On compactly supported brane variations the boundary terms vanish, so local operators in the interior are computed from the Koszul zero-fibre complex.

*Stable large- $N$*  means the degreewise stable value of the invariant algebras under block restriction, accessed only after four hypotheses. Procesi–Razmyslov stable invariant theory [17, 18]: stable invariants of two matrices are freely generated by cyclic trace words. Adjacent-swap exactness in the connected sector: the Koszul differential  $Q\psi = [\phi_1, \phi_2]$  makes adjacent transpositions  $Q$ -exact (Lemma 3.30), so ghost-zero cyclic words collapse on  $H^0$  to commutative monomials  $\text{Tr}(\phi_1^k \phi_2^l)$ ,  $k + l > 0$  (Proposition 3.22, Corollary 3.32). Connected-trace projection: the LQT cyclic-homology template [2, 3, 24] supplies the primitive single-cycle class; same-rank deletion has a nonzero commutator coderivation, and the finite-rank bridge is the separated-block zigzag  $P_{M,K,L}^{\text{sep}} = \beta_{M,K,L} P_{\text{cyc}} \alpha_{M,K,L}$  of Remark 3.7 below. Stable-window finite- $N$  bound: the LQT projection is a chain map only for  $N \geq \max(K, L + 2)$  (Theorem 3.6); outside this window finite- $N$  trace identities re-enter and break the projection.

Identifying the raw finite- $N$  operator algebras with the strict formal Hamiltonian envelope requires the open-trace  $A_\infty$ -Koszul acceptance datum  $(A_{\text{op}}, C_{\text{op}}, B_{\text{CE/PV}}, \tau_{\text{op}})$  of Definition 4.26, proved by Theorem 4.27; its cohomology controls the bar-cobar quasi-isomorphism  $\Omega\text{Bar}(A_{\text{op}}) \xrightarrow{\sim} B_{\text{CE/PV}}$  and the dual CE-side enveloping comparison  $C_{\text{CE,cont}}^\bullet(\mathfrak{g}_{\text{adm}}) \simeq A_{\text{op}}^!$  of clause (3) of Theorem 5.5. The acceptance datum is exactly what upgrades the connected-trace pattern of Theorem 5.6 to the global  $HC_{*-1}$  convention transfer.

**THEOREM 3.5** (*Loday–Quillen–Tsygan stable trace theorem*). Let  $A$  be a unital associative algebra over a field of characteristic zero and let  $\mathfrak{gl}_\infty(A) = \varinjlim_N \mathfrak{gl}_N(A)$ . As connected graded-commutative Hopf algebras,

$$H_*^{\text{CE}}(\mathfrak{gl}_\infty(A); \mathbb{C}) \cong \mathbf{S}(HC_{*-1}(A)).$$

Equivalently,

$$\text{Prim } H_*^{\text{CE}}(\mathfrak{gl}_\infty(A); \mathbb{C}) \cong HC_{*-1}(A).$$

**THEOREM 3.6** (*Stable Eulerian finite-window LQT projection*). Let

$$A_\psi = T(x, y, \psi), \quad d\psi = xy - yx,$$

with

$$\text{wt}(x) = (1, 0), \quad \text{wt}(y) = (0, 1), \quad \text{wt}(\psi) = (1, 1).$$

For a Koszul-weight bound  $K$ , let  $A_{\psi,K} = A_\psi / F^{>K} A_\psi$ , and let  $L$  bound CE length. In the stable block-sum Hopf CE window  $H_{K,L}^{\text{st}}(A_\psi)$ , with product induced by block diagonal sum and coproduct the CE coproduct, put

$$J = \text{id} - u\epsilon, \quad P_{\text{cyc}} = \log^*(\text{id}) = \sum_{m=1}^L \frac{(-1)^{m-1}}{m} J^{*m}.$$

Then  $P_{\text{cyc}}$  is a chain idempotent whose image is the stable single-cycle trace span  $\text{im } \Theta_{N,K,L}$ . For

$$N \geq \max(K, L + 2),$$

the finite-window LQT projection is the chain map

$$\lambda_{\text{LQT},K,L} = \Theta_{N,K,L}^{-1} P_{\text{cyc}} : H_{K,L}^{\text{st}}(A_\psi) \longrightarrow CC_{\text{red}, \leq L}(A_{\psi,K})[1].$$

If one integer  $W$  controls both  $K \leq W$  and  $L \leq W$ , this means  $N \geq W + 2$ . The scalar one- $\psi$  line is a chain quotient only after quotienting the two-term scalar-Koszul subcomplex

$$\Theta_1(\psi) \mapsto \Theta_1(xy - yx), \quad [\psi] \mapsto [xy - yx],$$

or after passing to homology.

*Proof.* The quotient  $A_{\psi,K}$  is a dg quotient because  $d\psi = xy - yx$  preserves the Koszul bidegree  $(1, 1)$ . The trace cycle map  $\Theta_{N,K,L}$  sends a reduced cyclic word  $a_0 | \cdots | a_r, r + 1 \leq L$ , to the usual cyclic matrix tensor

$$\sum_{i_0, \dots, i_r} (E_{i_0 i_1} a_0) \wedge \cdots \wedge (E_{i_r i_0} a_r),$$

and the matrix bracket gives the reduced cyclic differential with the standard graded cyclic signs. In particular  $d_{\text{CE}} \Theta_1(\psi) = \Theta_1(xy - yx)$ .

The stable CE window is formed before projecting. Block diagonal sum is a chain product, the CE coproduct is a chain coproduct, and  $J = \text{id} - u\epsilon$  is a chain map; hence every convolution power  $J^{*m}$  and therefore  $P_{\text{cyc}}$  is a chain map. Stable invariant CE tensors are indexed by permutations of the CE factors, and a permutation decomposes uniquely into cycles. A single cycle is exactly a trace-cycle tensor in  $\text{im } \Theta_{N,K,L}$ ; a multicyle tensor is a block-sum product of such single cycles. The Hopf logarithm kills decomposable products because on a product of  $n$  positive single-cycle generators its coefficient is

$$\sum_{m=1}^n (-1)^{m-1} (m-1)! S(n, m),$$

the coefficient of  $t^n/n!$  in  $\log(1 + (e^t - 1)) = t$ . This is 1 for  $n = 1$  and 0 for  $n > 1$ , proving idempotence and the stated image. The bound  $N \geq K$  removes trace identities in the displayed Koszul window, and  $N \geq L + 2$  is the safe CE stable range in length at most  $L$ .  $\square$

*Remark 3.7 (Same-rank deletion vs. the LQT projection).* The LQT projection  $P_{\text{cyc}}$  is the stable block-sum Hopf primitive projection, distinct from the fixed-rank operation  $P_{\text{del}}$  that keeps one-cycle permutation tensors and deletes multicyle tensors inside one exterior CE complex. For homogeneous  $a, b$

$$P_{\text{del}} d_{\text{CE}}(\Theta_1(a) \wedge \Theta_1(b)) = \pm \Theta_1(ab - (-1)^{|a||b|} ba), \quad d_{\text{CE}} P_{\text{del}}(\Theta_1(a) \wedge \Theta_1(b)) = 0.$$

Thus same-rank deletion has a nonzero chain defect when the graded commutator is nonzero. The theorem uses the stable block-sum Hopf primitive projection before returning to any finite rank. More precisely, set  $B(K, L) = \max(K, L + 2)$  and  $M = L B(K, L)$ . Let

$$\beta_{M,K,L}: H_{K,L}^{\text{st}}(A_{\psi}) \rightarrow C_*^{\text{CE}}(\mathbf{gl}_M(A_{\psi,K}))_{\leq L}^{\text{sep}}$$

be the normalized signed average over injections of the connected Hopf factors into the  $L$  labeled diagonal  $B(K, L)$ -blocks, and let

$$\alpha_{M,K,L}: C_*^{\text{CE}}(\mathbf{gl}_M(A_{\psi,K}))_{\leq L}^{\text{sep}} \rightarrow H_{K,L}^{\text{st}}(A_{\psi})$$

forget the block labels and read back the stable block-sum Hopf word. Thus  $\alpha_{M,K,L} \beta_{M,K,L} = \text{id}$ . Then

$$P_{M,K,L}^{\text{sep}} = \beta_{M,K,L} P_{\text{cyc}} \alpha_{M,K,L}$$



is the strict finite-rank chain idempotent on the separated-block subcomplex. This zigzag is the finite-rank bridge. A same-rank chain-homotopy bridge would have to kill the collision coderivation

$$\Gamma_N = d_{\text{same}} F_N - F_N d_{\oplus},$$

whose first nonzero component is

$$\Gamma_N(\Theta_1^N(a) \wedge \Theta_1^N(b)) = \pm \Theta_1^N(ab - (-1)^{|a||b|} ba).$$

### 3.3 Open mixed brane states

The open state space is fixed by the trace substitution: every brane observable is a polynomial in  $(\phi_1, \phi_2)$  traced over the Chan-Paton index, modulo the moment-map first-class constraint  $[\phi_1, \phi_2] = 0$ . The Koszul derived zero-fibre model for that constraint, the Procesi–Razmyslov invariant theory of the surviving traces, and the local geometry of the brane line and skyscraper point split the open states into a topological factor along the line, a  $\mathfrak{gl}_N$  Chan-Paton factor, and a Koszul-dual exterior factor on two odd derived normal coordinates. The matrix entries of  $[\phi_1, \phi_2]$  have syzygies for  $N \geq 2$ ; this is the derived zero fibre.

#### KOSZUL REDUCTION OF THE FORMAL-STALK CHART

The Dirac brane stack carries the polynomial coordinate ring

$$\mathcal{R}_N^{\text{poly}} = \mathbb{C}[(\phi_1)^i_j, (\phi_2)^i_j], \quad 1 \leq i, j \leq N,$$

together with the holomorphic symplectic form  $\Omega_N = \text{Tr}(d\phi_1 \wedge d\phi_2)$  and the  $GL_N$ -action by simultaneous conjugation. Conjugation is Hamiltonian with moment map  $\mu_\epsilon = -\text{Tr}(\epsilon[\phi_1, \phi_2])$  (Lemma 3.15); its zero locus is the variety  $\{[\phi_1, \phi_2] = 0\}$  of commuting matrix pairs.

**PROPOSITION 3.8** (*Derived zero-fibre and Procesi–Razmyslov trace ring*). The ghost-zero open observable algebra of the Dirac brane stack is the classical BV cohomology of the Koszul complex

$$\mathcal{R}_N = \mathcal{R}_N^{\text{poly}} \otimes \Lambda_{\mathbb{C}}(\psi^i_j), \quad Q\psi^i_j = [\phi_1, \phi_2]^i_j, \quad Q\phi_1 = Q\phi_2 = 0,$$

on  $GL_N$ -invariants:

$$\mathcal{O}_{\text{brane}, N}^{\text{cl,gho}} = H^0(\mathcal{R}_N^{GL_N}, Q).$$

The  $GL_N$ -invariant subring  $(\mathcal{R}_N^{\text{poly}})^{GL_N}$  is generated, by the Procesi–Razmyslov first fundamental theorem, by cyclic traces  $\text{Tr } w(\phi_1, \phi_2)$  of words  $w$  in two letters; passage to  $H^0$  imposes the relations  $Q\psi = [\phi_1, \phi_2] = 0$ .

*Proof.* Since  $GL_N$  is reductive in characteristic zero, invariants is exact and commutes with cohomology, so  $H^0(\mathcal{R}_N^{GL_N}, Q) = H^0(\mathcal{R}_N, Q)^{GL_N}$ . The fields  $\phi_1, \phi_2$  sit in ghost number zero and  $\psi$  in ghost number  $-1$ ;  $Q\psi = [\phi_1, \phi_2]$  is the standard Koszul differential of the  $\mathfrak{gl}_N$ -valued moment map, so classical BV theory identifies  $H^0(\mathcal{R}_N, Q)$  with the polynomial functions on the scheme  $\{[\phi_1, \phi_2] = 0\} \subset \mathfrak{gl}_N \oplus \mathfrak{gl}_N$ . Procesi (1976) and Razmyslov (1974) prove that  $(\mathcal{R}_N^{\text{poly}})^{GL_N}$  is generated by cyclic traces of words in  $(\phi_1, \phi_2)$ ; restriction to the zero-fibre imposes the bracket-vanishing relations.  $\square$

The Koszul complex  $\mathcal{R}_N$  is the open ghost-zero state space at a single brane point. Two further data assemble it into a sheaf along the brane line  $L$  and over the formal holomorphic point: the topological factor along  $\mathbb{R}_t$  and the holomorphic factor transverse to  $L$  inside  $\widehat{\mathbb{C}^2}_0$ .

### TOPOLOGICAL FACTOR ALONG THE BRANE LINE

The brane line has trivial curvature and holonomy: its self-Floer complex is exact and wrapped generators are excluded by the local model. For a compact interval  $I \subset \mathbb{R}_t$  and a compactly supported Hamiltonian perturbation, the local Fukaya complex  $CF^*(\mathbb{R}_t, \mathbb{R}_t; I)$  is the Morse complex on  $I$ , and passage to local operators in the interior identifies it with the de Rham complex  $\Omega^*(\mathbb{R}_t)$  acting by compactly supported variations. Wrapped generators at infinity are excluded by the local-model hypothesis.

### HOLOMORPHIC FACTOR AT THE BRANE POINT

The brane is a point inside  $\widehat{\mathbb{C}^2}_o$ ; its holomorphic state space is the derived endomorphism algebra of the skyscraper  $\mathcal{O}_p$ . Koszul duality identifies it with the exterior algebra on the two derived normal coordinates whose ghost-zero shadows are the matrix coordinates  $\phi_1, \phi_2$ .

**PROPOSITION 3.9** (*Affine skyscraper self-Ext*). Let  $R = \mathbb{C}[z_1, z_2]$ ,  $\mathfrak{m} = (z_1, z_2)$ , and  $k = R/\mathfrak{m}$ . The Yoneda algebra of the skyscraper at the origin is

$$\mathrm{Ext}_R^*(k, k) \cong \Lambda_{\mathbb{C}}(\varepsilon_1, \varepsilon_2), \quad |\varepsilon_i| = 1,$$

the free graded-commutative algebra on two odd generators.

*Proof.* The pair  $(z_1, z_2)$  is a regular sequence in  $R$ ; the affine Koszul resolution

$$K = (0 \rightarrow R e_{12} \xrightarrow{d_2} R e_1 \oplus R e_2 \xrightarrow{d_1} R \rightarrow 0), \quad d_1(ae_1 + be_2) = az_1 + bz_2, \quad d_2(e_{12}) = z_2e_1 - z_1e_2,$$

is a free  $R$ -resolution of  $k$ . Applying  $\mathrm{Hom}_R(-, k)$  gives

$$0 \rightarrow k \xrightarrow{\circ} k \varepsilon_1 \oplus k \varepsilon_2 \xrightarrow{\circ} k \varepsilon_{12} \rightarrow 0,$$

because  $z_1, z_2$  act by zero on  $k$ ; hence  $\mathrm{Ext}_R^p(k, k)$  is one-dimensional in degree 0, two-dimensional in degree 1, and one-dimensional in degree 2. The Yoneda product is computed by dualising the comultiplication of the Koszul dg-algebra  $R \otimes \Lambda(e_1, e_2)$  with  $d(e_i) = z_i$ , under which  $e_i \mapsto e_i \otimes 1 + 1 \otimes e_i$ ; duality forces  $\varepsilon_i^2 = 0$ ,  $\varepsilon_1\varepsilon_2 = -\varepsilon_2\varepsilon_1$ , with  $\varepsilon_{12} = \varepsilon_1\varepsilon_2$  spanning the top Ext group.  $\square$

**Remark 3.10** (*Koszul duality and derived normal coordinates*).  $\Lambda(\varepsilon_1, \varepsilon_2)$  is Koszul dual to the polynomial ring  $\mathbb{C}[\phi_1, \phi_2]$  on two even generators; the Chan–Paton coordinate ring of the derived commuting variety chart and the skyscraper Ext algebra are the same datum read from the two sides of Koszul duality. The ghost-zero scalar coefficients  $\chi_1, \chi_2$  of  $\varepsilon_1, \varepsilon_2$  in §3.4 are, after Dirac normalization, the matrix coordinates  $\phi_1, \phi_2$ ; the Procesi–Razmyslov trace ring of Proposition 3.8 reappears there as the ghost-zero observable algebra of the open BV theory.

### OPEN-STATE FACTORISATION ALONG $L$

Combining the three factors, the local open state space at the brane is

$$\mathcal{S}_L^{\mathrm{open}} = \Omega^*(\mathbb{R}_t) \otimes \mathrm{Ext}_{\mathbb{C}[z_1, z_2]}^*(\mathcal{O}_p, \mathcal{O}_p) \otimes \mathfrak{gl}_N[I] = \Omega^*(\mathbb{R}_t) \otimes \Lambda(\varepsilon_1, \varepsilon_2) \otimes \mathfrak{gl}_N[I].$$

The shift  $[I]$  is the BV ghost-shift of Definition 1.10: a form of de Rham degree  $p$  and Grassmann degree  $q$  sits in ghost number  $p + q - 1$ ; the brane gauge field has  $p = 1, q = 0$  (ghost 0), the two scalar fields have  $p = 0, q = 1$  (ghost 0), and the remaining components carry nonzero ghost number, encoding the BV antifields and ghosts of the  $GL_N$ -conjugation gauge symmetry of the Chan–Paton pair.

### 3.4 Open mixed brane field theory

Under the Dirac brane stack hypotheses of §1.4 — the  $\mathfrak{gl}_N$ -valued holomorphic-symplectic pair  $(\phi_1, \phi_2)$  carrying the form  $\Omega_N = \text{Tr}(d\phi_1 \wedge d\phi_2)$  and its Liouville one-form  $\theta_N = \text{Tr}(\phi_1 d\phi_2)$  (Lemma 3.15), the moment-map zero-fibre  $[\phi_1, \phi_2] = 0$  coupled by a Lagrange multiplier  $A \in \Omega^1(\mathbb{R}_t) \otimes \mathfrak{gl}_N$  along the brane line, and the  $GL_N$ -conjugation gauge symmetry generated by that moment map — the open BV action, BV differential, and ghost-zero classical observable algebra are forced. The action is the Dirac first-class functional (9); the BV differential is the Koszul derived zero-fibre model for the moment-map equation (10) with antifield  $\psi = A^*$ ; the ghost-zero classical observables are the Procesi–Razmyslov derived zero-fibre of Proposition 3.8. Together they assemble into the formal first-order Chern–Simons theory (11) on  $\mathbb{R}_t \times \mathbb{C}^{0|2}$ , with Dirac normalization fixing the signs.

#### ACTION: CHAN–PATON LIOUVILLE PLUS DIRAC MULTIPLIER

The Chan–Paton symplectic form  $\Omega_N = \text{Tr}(d\phi_1 \wedge d\phi_2)$  admits the Liouville one-form  $\theta_N = \text{Tr}(\phi_1 d\phi_2)$ . The brane moment map for simultaneous  $GL_N$ -conjugation is  $\mu_\epsilon = -\text{Tr}(\epsilon[\phi_1, \phi_2])$  (Lemma 3.15); coupling its zero-fibre constraint by a Lagrange multiplier  $A \in \Omega^1(\mathbb{R}_t) \otimes \mathfrak{gl}_N$  along the brane line gives the Dirac first-class action

$$S_\partial = \int_{\mathbb{R}_t} \theta_N - \mu_A = \int_{\mathbb{R}_t} \text{Tr}(\phi_1 d\phi_2 + A[\phi_1, \phi_2]). \quad (9)$$

The variation of  $A$  imposes the moment-map zero-fibre, and the variation of  $\phi_i$  imposes the Hamiltonian flow generated by  $\mu_A$  along the brane line; this is the Dirac constraint reduction of the formal-stalk chart.

#### BV DIFFERENTIAL: KOSZUL DERIVED ZERO-FIBRE MODEL FOR THE MOMENT-MAP EQUATION

The first-class constraint  $[\phi_1, \phi_2] = 0$  is realized by the derived zero-fibre model with an antifield  $\psi = A^* \in \mathfrak{gl}_N[1]$  and the Koszul differential

$$Q\psi^i_j = [\phi_1, \phi_2]^i_j, \quad Q\phi_1 = Q\phi_2 = 0. \quad (10)$$

The matrix entries of  $[\phi_1, \phi_2]$  have syzygies for  $N \geq 2$ ; this is the derived zero fibre of the moment map. Its ghost-zero invariant cohomology identifies the open classical observable algebra with  $H^0(\mathcal{R}_N^{GL_N}, Q)$  (Proposition 3.8): the cotangent direction of the moment-map zero-fibre is exactly the BV antifield, the Koszul differential is exactly the BV differential.

#### CYCLIC TRACE AND BV PAIRING

The cyclic trace pairing comes from two ingredients: the Berezin integral on the exterior factor  $\Lambda(\varepsilon_1, \varepsilon_2)$ , which projects onto the top class  $\varepsilon_1 \varepsilon_2$  (residue dual to  $dz_1 \wedge dz_2$ ); the de Rham integral on the brane line, which projects onto the top form. Together they fix the open cyclic trace

$$\text{Tr}_{\text{open}}(\eta \otimes u \otimes M) = \int_{\mathbb{R}_t} \eta[\varepsilon_1 \varepsilon_2] u \text{Tr}_N(M),$$

where  $[\varepsilon_1 \varepsilon_2]u$  is the coefficient of the top Ext class. The de Rham integral carries degree  $-1$  and the Ext residue degree  $-2$ ; after the field shift  $[1]$ , the induced BV pairing has degree  $-1$ , as required by Definition 1.10.

The total open BV field is therefore  $\alpha \in \Omega^*(\mathbb{R}_t) \otimes \Lambda(\varepsilon_1, \varepsilon_2) \otimes \mathfrak{gl}_N[1]$ , and a single first-order Chern–Simons action records all three pieces at once: the Liouville term, the moment-map multiplier, and the Koszul antifield.

**PROPOSITION 3.11** (*Open BV action from the formal-stalk chart*). The open BV theory of the mixed brane is the formal first-order Chern-Simons theory

$$S(\alpha) = \int_{\mathbb{R}_t \times \mathbb{C}^{0|2}} \frac{1}{2} \text{Tr}(\alpha d\alpha) + \frac{1}{3} \text{Tr} \alpha^3, \quad d = d_{\text{dR}}, \quad (\text{II})$$

on the open BV field  $\alpha \in \Omega^*(\mathbb{R}_t) \otimes \Lambda(\varepsilon_1, \varepsilon_2) \otimes \mathfrak{gl}_N[1]$ , with the Berezin convention  $\int_{\mathbb{C}^{0|2}} d\varepsilon_1 d\varepsilon_2 : \varepsilon_1 \varepsilon_2 \mapsto 1$  and zero differential on the exterior factor. Its restriction to the ghost-zero coefficient fields, after the linear Dirac normalization  $(A_{\text{CS}}, \chi_1, \chi_2) \mapsto (-A, -\phi_1, \phi_2)$ , is the Dirac action  $S_\partial$  of (9); its BV differential  $Q$  restricted to the antifield  $\psi = A^*$  is the moment-map Koszul differential (10); its ghost-zero classical observable algebra is the Procesi–Razmyslov derived zero-fibre  $H^0(\mathcal{R}_N^{GL_N}, Q)$  of Proposition 3.8.

*Proof.* The reduction of (II) to the ghost-zero coefficient sector is Lemma 3.14; the identification of the BV differential with the moment-map Koszul differential is Lemma 3.15; the identification of the ghost-zero classical observables with the derived zero-fibre is Proposition 3.16 combined with Proposition 3.8.  $\square$

**Construction 3.12** (*The open field from the Ext basis*). Write  $\varepsilon_{12} = \varepsilon_1 \varepsilon_2$ . Before imposing ghost number, an open field is a finite sum

$$\alpha = a_0 + a_1 \varepsilon_1 + a_2 \varepsilon_2 + a_{12} \varepsilon_{12},$$

where each  $a_I$  is a de Rham form on  $\mathbb{R}_t$  with values in  $\mathfrak{gl}_N$ , shifted by  $[1]$ . The ghost-number-zero components are

$$A_{\text{CS}} \in \Omega^1(\mathbb{R}_t) \otimes \mathfrak{gl}_N, \quad \chi_1, \chi_2 \in \Omega^0(\mathbb{R}_t) \otimes \mathfrak{gl}_N,$$

with coefficient form  $\alpha_0^{\text{coef}} = A_{\text{CS}} + \chi_1 \varepsilon_1 + \chi_2 \varepsilon_2$ . The Dirac variables of the rest of the manuscript are

$$\phi_1 = -\chi_1, \quad \phi_2 = \chi_2, \quad A = -A_{\text{CS}},$$

fixed once and for all so that the Liouville one-form is  $\text{Tr}(\phi_1 d\phi_2)$  and the moment-map multiplier term is  $A[\phi_1, \phi_2]$ . The Berezin integral kills every monomial except the coefficient of  $\varepsilon_{12}$ , so each ghost-zero monomial of (II) is obtained by multiplying the three displayed summands and keeping exactly one real one-form and the product  $\varepsilon_1 \varepsilon_2$ .

**Remark 3.13** (*Ghost shift*). Total degree of an  $(\Omega^p, \varepsilon\text{-degree } q)$  monomial is  $p + q$ ; after the BV shift  $[1]$  it sits in ghost number  $p + q - 1$ . The brane gauge field ( $p = 1, q = 0$ ) and the two scalar fields ( $p = 0, q = 1$ ) sit in ghost zero; the components proportional to  $\varepsilon_i$  record the motion of the brane in the two derived normal directions of the skyscraper.

#### 3.4.1 GHOST-ZERO OPEN FIELD CONTENT

**LEMMA 3.14.** With the Berezin convention  $\int_{\mathbb{C}^{0|2}} \varepsilon_1 \varepsilon_2 = 1$ , the restriction of the BV action to the ghost-zero coefficient fields, written in the Dirac variables of Construction 3.12, is, up to a boundary term on  $\mathbb{R}$ ,

$$S(\alpha_0) = \int_{\mathbb{R}} \text{Tr}(\phi_1 d\phi_2 + A[\phi_1, \phi_2]).$$

*Proof.* Only monomials containing one real one-form and the factor  $\varepsilon_1 \varepsilon_2$  survive integration over  $\mathbb{R} \times \mathbb{C}^{0|2}$ . In coefficient coordinates  $\alpha_0^{\text{coef}} = A_{\text{CS}} + \chi_1 \varepsilon_1 + \chi_2 \varepsilon_2$ , the tensor-product Koszul sign gives

$$\frac{1}{2} \int_{\mathbb{R}} \text{Tr}(-\chi_1 d\chi_2 + \chi_2 d\chi_1) \equiv - \int_{\mathbb{R}} \text{Tr}(\chi_1 d\chi_2),$$

modulo an exact one-form. In the cubic term the six orderings of  $A_{\text{CS}}, \chi_1 \varepsilon_1, \chi_2 \varepsilon_2$  reduce, by cyclicity of the trace and  $\varepsilon_2 \varepsilon_1 = -\varepsilon_1 \varepsilon_2$ , to

$$3 \operatorname{Tr}(A_{\text{CS}} \chi_1 \chi_2 - A_{\text{CS}} \chi_2 \chi_1).$$

The coefficient  $\frac{1}{3}$  in the Chern-Simons action therefore produces  $\operatorname{Tr}(A_{\text{CS}}[\chi_1, \chi_2])$ . Substituting  $\phi_1 = -\chi_1$ ,  $\phi_2 = \chi_2$ , and  $A = -A_{\text{CS}}$  gives the displayed Dirac-normalized action.  $\square$

LEMMA 3.15 (*Dirac reduction of the brane phase space*). Let

$$\mathcal{P}_N = \mathfrak{gl}_N \oplus \mathfrak{gl}_N$$

with coordinates  $(\phi_1, \phi_2)$  and symplectic form

$$\Omega_N = \operatorname{Tr}(d\phi_1 \wedge d\phi_2).$$

With the convention  $\iota_{X_f} \Omega_N = -df$ , simultaneous conjugation by  $\mathfrak{gl}_N$  is Hamiltonian with moment map components

$$\mu_\epsilon = -\operatorname{Tr}(\epsilon[\phi_1, \phi_2]), \quad \epsilon \in \mathfrak{gl}_N.$$

They satisfy the first-class identity

$$\{\mu_\epsilon, \mu_\eta\} = -\mu_{[\epsilon, \eta]}.$$

Thus this comoment is anti-equivariant for the displayed Poisson convention, equivalently equivariant for the opposite Lie algebra; the zero locus and the generated constraint ideal are unchanged. The action of Lemma 3.14 is the corresponding Dirac first-order action

$$\int_{\mathbb{R}} \operatorname{Tr}(\phi_1 d\phi_2) - \mu_A,$$

and varying  $A$  imposes the constrained phase space

$$\mu^{-1}(\mathbf{o})/GL_N = \{[\phi_1, \phi_2] = \mathbf{o}\}/GL_N.$$

*Proof.* The symplectic form gives

$$\{(\phi_1)^i_j, (\phi_2)^k_l\} = \delta_l^i \delta_j^k.$$

For  $v_\epsilon(\phi_i) = [\epsilon, \phi_i]$ , cyclicity of the trace gives

$$d\mu_\epsilon(\delta\phi_1, \delta\phi_2) = -\operatorname{Tr}(\epsilon([\delta\phi_1, \phi_2] + [\phi_1, \delta\phi_2])) = -\Omega_N(v_\epsilon, (\delta\phi_1, \delta\phi_2)).$$

Thus  $X_{\mu_\epsilon} = v_\epsilon$ . The moment-map bracket is therefore the anti-equivariance identity  $\{\mu_\epsilon, \mu_\eta\} = -\mu_{[\epsilon, \eta]}$ . Substituting  $\epsilon = A(t)$  gives

$$\int_{\mathbb{R}} \operatorname{Tr}(\phi_1 d\phi_2) - \mu_A = \int_{\mathbb{R}} \operatorname{Tr}(\phi_1 d\phi_2 + A[\phi_1, \phi_2]),$$

and the  $A$ -equation is  $[\phi_1, \phi_2] = \mathbf{o}$ .  $\square$

PROPOSITION 3.16 (*Classical BV truncation*). The subspace of ghost-number-zero coefficient fields

$$\alpha_o^{\text{coef}} = A_{\text{CS}} + \chi_1 \varepsilon_1 + \chi_2 \varepsilon_2$$

with all nonzero ghost-number components set to zero is a consistent classical truncation of the open BV theory. After the linear Dirac-normalization of Construction 3.12, the induced classical action is the first-order gauged matrix model of Lemma 3.14.

*Proof.* Decompose the BV field as  $\alpha = \sum_r \alpha^{(r)}$  by ghost number. The differential  $d_{\mathbb{R}}$  and the product on  $\Lambda(\varepsilon_1, \varepsilon_2) \otimes \mathfrak{gl}_N$  preserve total degree, and the BV action has ghost number zero. Hence every monomial in the Euler-Lagrange expression for a component of ghost number  $r \neq 0$  contains at least one field component of nonzero ghost number when evaluated at  $\alpha = \alpha_o^{\text{coef}}$ . Setting all such components to zero therefore makes the discarded equations vanish. The remaining equation is exactly the variation of the ghost-zero action computed in Lemma 3.14.

Concretely, the full Euler-Lagrange expression is the curvature

$$F(\alpha) = d\alpha + \alpha^2.$$

For  $\alpha_o^{\text{coef}} = A_{\text{CS}} + \chi_1 \varepsilon_1 + \chi_2 \varepsilon_2$  on a line,

$$F(\alpha_o^{\text{coef}}) = (d\chi_1 + [A_{\text{CS}}, \chi_1])\varepsilon_1 + (d\chi_2 + [A_{\text{CS}}, \chi_2])\varepsilon_2 + [\chi_1, \chi_2]\varepsilon_1\varepsilon_2.$$

The BV pairing keeps exactly real-form degree one and  $\varepsilon_1\varepsilon_2$ . Hence the displayed three components pair exactly with the coefficient variations of  $A_{\text{CS}}, \chi_2, \chi_1$  respectively, while variations of nonzero ghost number see complementary curvature components that are zero for  $\alpha_o^{\text{coef}}$  by real-form degree or  $\varepsilon$ -degree.  $\square$

The truncated action is gauge-invariant under

$$\delta_\varepsilon \phi_i = [\varepsilon, \phi_i], \quad \delta_\varepsilon A = d\varepsilon + [A, \varepsilon], \quad \varepsilon \in \Omega^0(\mathbb{R}_t) \otimes \mathfrak{gl}_N :$$

the  $d\varepsilon$  variation of  $\text{Tr}(\phi_1 d\phi_2)$  produces  $-\text{Tr}(d\varepsilon[\phi_1, \phi_2])$ , which cancels the  $d\varepsilon$  variation of  $\text{Tr}(A[\phi_1, \phi_2])$ ; the remaining terms cancel by invariance of the trace pairing. This is the gauge identity that identifies the brane gauge field as the Lagrange multiplier for the moment-map zero-fibre.

*Remark 3.17 (Skyscraper–matrix duality).* The same open theory arises from the affine  $\mathbb{C}^2$  point-brane Yoneda calculation by tensoring  $\mathcal{O}_p$  with  $\mathbb{C}^N$  and passing to derived endomorphisms; the Berezin integral on the Ext factor is the residue dual to  $dz_1 \wedge dz_2$ . This is the open Koszul duality between the Chan–Paton coordinate ring  $\mathbb{C}[\phi_1, \phi_2]$  and the skyscraper Ext algebra  $\Lambda(\varepsilon_1, \varepsilon_2)$  (Remark 3.10).

*Remark 3.18 (AKSZ form).* In AKSZ language the open theory is the one-dimensional gauged sigma model with Hamiltonian target  $\mathfrak{gl}_N \oplus \mathfrak{gl}_N$  and Liouville potential  $\theta_N = \text{Tr}(\phi_1 d\phi_2)$ . The Poisson bivector

$$\Pi = \sum_{i,j} \partial_{(\phi_1)^i j} \wedge \partial_{(\phi_2)^j i}, \quad \{(\phi_1)^i j, (\phi_2)^k l\} = \delta_l^i \delta_j^k,$$

is dual to  $\Omega_N$  under  $\iota_{X_f} \omega = -df$ ; the Hamiltonian  $\mu_\varepsilon = -\text{Tr}(\varepsilon[\phi_1, \phi_2])$  generates  $\delta_\varepsilon \phi_i = [\varepsilon, \phi_i]$ ; its zero-locus quotient is the stack of commuting matrix pairs modulo conjugation. Equivalently the same theory descends from three-dimensional Chern–Simons on  $\mathbb{R} \times T^2$  restricted to harmonic forms  $\iota_1, \eta_1, \eta_2, \eta_1\eta_2$  on  $T^2$ , with  $A_{3d} = A_t dt + \phi_1 \eta_1 + \phi_2 \eta_2$ .

The  $P_o$ -algebra structure on the unreduced open observables, with the cotangent direction realised at the chain level, requires data beyond the reduced construction; its data and the obstruction it must overcome are the following.

*Problem 3.19 (Unreduced cotangent lift of the open BV theory).* Construct an unreduced Tate-coefficient cotangent enlargement of the open BV observables of Proposition 3.11 realising the cotangent direction  $\mathcal{P} = \widehat{\mathfrak{h}}_{\text{cont}}^{\vee}$  of the closed Hamiltonian Lie algebra at the chain level on the brane line. The required data are: a candidate enlargement

$$\widetilde{\text{Obs}}_{\partial, N}^{\text{cl, pp}}(I) = \text{Obs}_{\text{open}, N}^{\text{cl}}(I) \widehat{\otimes} \widehat{\mathbf{S}}(\mathcal{M}_{\partial}^{\text{pp, unred}}(I)),$$

with  $\mathcal{M}_{\partial}^{\text{pp, unred}}(I) = C_{c, \text{cur}}^{\text{dR}}(I) \widehat{\otimes} \text{Ann}_{\text{res}}(\mathbb{C} \cdot 1)[1]$ ; a chain-level primitive projection  $\pi_{\text{prim}}$  compatible with  $Q_{\text{open}}$ , constants, decomposable multi-traces, and factorization products; centrality homotopies for the cotangent vertex  $\Theta_{\rho}$  against arbitrary unreduced open observables;  $P_o$ -bracket homotopies

$$\{B_f, \Theta_{\rho}\} \simeq \Theta_{f \cdot \rho}, \quad \{\Theta_{\rho}, \Theta_{\sigma}\} \simeq 0,$$

before primitive projection. The open-side data is the restriction to the brane line of the full unreduced cotangent lift recorded as Problem 5.48.

The obstruction mode of the centrality homotopies and Tate-cotangent module of Problem 3.19 is the obstruction class of Remark 5.189: the same Tate-coefficient cotangent module governs both the open-side centrality homotopies and the unreduced bulk-to-brane factorization-centre lift.

### 3.5 Open-brane operators and the large $N$ limit

A stack of  $N$  Dirac branes carries a finite-matrix trace map of the formal symplectic polydisk: its operator algebra is generated by cyclic words in two  $\mathfrak{gl}_N$ -valued Chan–Paton matrices, and its Weyl quantisation produces a scalar shift  $\hbar N$  whose class survives connected single-trace projection. Procesi–Razmyslov 1976 ([17, 18]; Proposition 3.22) fixes the classical content of the formal-stalk-chart operator side; Capelli 1890 ([26, 27]; Remark 5.195) forces the quantum scalar. Theorem 5.6 reads its operator side off the constructions below.

Three large- $N$  operations of Definition 3.41 apply in fixed order: degreewise invariant stabilisation  $\text{Prim}_{\text{st}}$ , connected single-trace extraction  $\text{Conn}_{N \rightarrow \infty}$ , and Hamiltonian scalar quotient  $\pi_{\text{Ham}}$ . Their composite output is the stable brane Hamiltonian algebra  $\mathbf{S}(V_H)$  with  $V_H = \tilde{A} \oplus \tilde{A}_{\text{desc}}[1]$  (Proposition 3.40), whose connected indecomposable quotient  $\mathfrak{m}/\mathfrak{m}^2 \cong V_H$  is matched by the closed side of §6.2 as continuous CE cochains on  $\mathfrak{g} = \mathfrak{h} \ltimes \widehat{\mathfrak{h}}_{\text{cont}}^{\vee}[1]$ . The four-window scalar-anomaly transgression (Theorem E.13) is the bridge between the classical Lie-cohomology class  $[\bar{e}]$  and the Weyl/Capelli quantum shift; cancellation on one window forces cancellation on the other three. The all-bidegree residual after Capelli renormalisation is  $\Omega_{a,b}^{\text{rad}} \in \text{coker } B_{a,b}$ , the decorated PBW Stokes obstruction of Theorem 5.216.

The Dirac brane sector (Lemma 3.15, Proposition 3.16) is the BV theory of two  $\mathfrak{gl}_N$ -valued matrices and one Koszul antifield, with  $Q\psi = [\phi_1, \phi_2]$ . In characteristic zero,  $GL_N$  is linearly reductive, so degreewise stable invariants are exact in the function-level calculation; invariant theory controls this step, while the unreduced Lie-CE BRST cohomology of  $\mathfrak{gl}_N$  has scalar stabilizer classes outside the Hamiltonian trace sector. Procesi–Razmyslov identifies the stable invariant algebra with the free commutative algebra on nonempty cyclic words in two letters (Proposition 3.22). Adjacent-swap exactness collapses the zero- $\psi$  sector to commutative monomials on the formal symplectic plane (Lemma 3.30,

Corollary 3.32). The chip-moving necklace homotopy (Proposition 3.27) computes the primitive one- $\psi$  Koszul homology in nonconstant bidegree  $(k, l)$  as the line spanned by the symmetrised trace

$$\Upsilon_{k,l} = \sum_{w \in W_{k,l}} \text{Tr}(\psi w(\phi_1, \phi_2)).$$

The Hamiltonian descendant algebra is therefore  $\widehat{\mathbf{S}}(\bar{A} \oplus \bar{A}_{\text{desc}}[1])$  (Proposition 3.40); its connected indecomposable quotient is  $V_H = \bar{A} \oplus \bar{A}_{\text{desc}}[1]$ .

Quantization tests the compatibility of Weyl ordering with the brane trace map. On the unreduced brane the classical phase-space mismatch  $\{\text{Tr } \phi_1, \text{Tr } \phi_2\} = N$  survives Weyl quantisation as a Capelli contact term: the Weyl-ordered moment relation reads

$$[\Phi_{\hbar}(z_1), \Phi_{\hbar}(z_2)]_{\star} = \hbar N \cdot I, \quad YX - XY + \hbar NI \equiv 0 \pmod{\mathcal{W}_N \widehat{\mu}(\mathfrak{sl}_N)},$$

hence  $\text{Tr}(A XY) - \text{Tr}(A YX) = \hbar N \text{Tr}(A)$  on every covariant matrix word  $A$ . The right-hand side is a nonempty connected trace whenever  $A$  is nonconstant: the single-trace primitive projection removes the empty trace and disconnected products while retaining this lower-degree trace insertion. The Capelli shift is the quantum mirror of the closed-side scalar class  $[\bar{c}] \in H_{\text{Lie}}^2(\bar{A}, \mathbb{C})$  of Theorem 6.4: the same scalar appears classically as the central-extension obstruction and quantum mechanically as the Weyl-ordering anomaly, and Theorem 6.6 identifies them. The four-window identification — Lie cohomology, classical trace algebra, Capelli quantum shift, Moyal star product — is the scalar-anomaly transgression Theorem E.13: each window detects the same cocycle, so cancelling it on one window forces cancellation on the other three.

The operator-side comparison closes at finite  $N$  through the Capelli-renormalised generator

$$J_N(z_1^a z_2^b) = \text{Tr } \text{Op}_W(z_1^a z_2^b)(\phi_1, \phi_2)$$

of Lemma 5.197: the change of basis from naive normal-ordered traces  $T_{a,b}$  to  $J_N$  is unipotent over  $\mathbb{C}[[\hbar N]]$ , and the displayed Capelli shift is exactly the  $r = 1$  tail of that triangular system. In the  $J_N$ -basis the Capelli term is absorbed; the residual all-bidegree obstruction is the quantum shear primitive

$$E_{a,b,N} \in \mathcal{W}_N \widehat{\mu}(\mathfrak{sl}_N), \quad \Omega_{a,b}^{\text{rad}} = [(T_{a,b}, E_{a,b}^+)] \in \text{coker } B_{a,b}$$

of Theorem 5.216, equivalently the decorated PBW Stokes identity  $D_{a,b}^{\square} = C_{a,b}^+ \partial_2$ ; an obstruction is exactly a signed row  $(\phi, -\lambda) \in \ker B_{a,b}^*$  with  $\lambda(E_{a,b}^+) - \phi(T_{a,b}) \neq 0$ . This is the open-trace  $\mathcal{A}_{\infty}$ -Koszul acceptance datum of Remark 3.44.

*Construction 3.20 (The reduced BV algebra at the insertion point).* At a fixed point of the brane line, introduce three matrix variables

$$\phi_1, \phi_2 \in \mathfrak{gl}_N, \quad \psi \in \mathfrak{gl}_N[1].$$

The reduced local polynomial algebra is

$$\mathcal{R}_N = \mathbb{C}[(\phi_1)^i_j, (\phi_2)^i_j] \otimes \Lambda((\psi)^i_j).$$

Its reduced Koszul/BV differential is the derivation determined on generators by

$$Q(\phi_1)^i_j = 0, \quad Q(\phi_2)^i_j = 0, \quad Q(\psi)^i_j = ([\phi_1, \phi_2])^i_j.$$



This formula is the algebraic encoding of the  $\mathcal{A}$ -equation of motion: varying

$$\int_{\mathbb{R}} \text{Tr}(\phi_1 d\phi_2 + \mathcal{A}[\phi_1, \phi_2])$$

with respect to  $\mathcal{A}$  gives  $[\phi_1, \phi_2] = 0$ , and the antifield  $\psi = \mathcal{A}^*$  records that equation as a Koszul differential.

**LEMMA 3.21** (*De Rham jets*). Positive jets in the topological direction vanish in local operator cohomology. Equivalently, every cohomology class has a representative determined by the values of the fields and antifields at the insertion point.

*Proof.* For a coordinate  $t$  on the line, the de Rham part of the BRST operator is  $d_{\mathbb{R}}$ . On the jet algebra one has the contraction homotopy  $[d_{\mathbb{R}}, \iota_{\partial_t}] = \partial_t$ . Hence a local expression containing a positive  $t$ -derivative is cohomologous to zero after applying the usual Euler homotopy on jet degree. This kills topological-direction jets and leaves the holomorphic  $z_i$ -derivatives used later on the closed side.  $\square$

**PROPOSITION 3.22** (*Stable trace invariants*). Let  $F = \mathbb{C}\langle x_1, x_2 \rangle$  and let  $\text{Cyc}(F) = F/[F, F]$  be the vector space of cyclic words. For each  $N$ ,  $GL_N$  acts on  $\mathfrak{gl}_N^2$  by simultaneous conjugation. The degree-wise stable invariant algebra

$$\mathbb{C}[\mathfrak{gl}_{\infty}^2]_{\text{st}}^{GL_{\infty}} := \bigoplus_{d \geq 0} \varprojlim_N (\mathbb{C}[\mathfrak{gl}_N^2]^{GL_N})_d$$

with respect to block-restriction maps in each word-degree is the free commutative algebra generated by the nonempty cyclic words:

$$\mathbb{C}[\mathfrak{gl}_{\infty}^2]_{\text{st}}^{GL_{\infty}} \cong \mathbf{S}(\text{Cyc}(F)_{\text{nonempty}}),$$

where a cyclic word  $w$  maps to the trace function  $(\phi_1, \phi_2) \mapsto \text{Tr } w(\phi_1, \phi_2)$ . More precisely, in total positive word-degree  $d$  this map is an isomorphism for every  $N \geq d$ , after omitting the scalar empty trace as a primitive generator.

*Proof.* Procesi–Razmyslov invariant theory [17, 18] gives two inputs. First,  $\mathbb{C}[\mathfrak{gl}_N^2]^{GL_N}$  is generated by trace functions of words in the two generic matrices. Second, all polynomial relations among these trace functions are consequences of the fundamental trace identities of matrix size  $N$ ; in particular every homogeneous trace relation has total word-degree at least  $N + 1$ . Fix a total word-degree  $d$ . For every  $N \geq d$  the degree- $d$  relations are exactly cyclicity of the trace. Passing degree by degree to the stable inverse limit removes all finite- $N$  trace identities. The surviving identification is  $\text{Tr}(uv) = \text{Tr}(vu)$ . This is exactly the symmetric algebra on cyclic words.  $\square$

**LEMMA 3.23** (*Stable marked trace invariants*). In the stable invariant theory used below, the primitive part linear or quadratic in an odd marked matrix  $\psi$  is obtained from the corresponding even three-matrix trace invariant by polarization and skew-symmetrisation in the marked letters. The only stable single-trace relation is graded cyclicity: moving one  $\psi$  through the trace has sign  $+1$ , while moving one  $\psi$  past another  $\psi$  gives a minus sign.

*Proof.* Apply Proposition 3.22 to three even generic matrices and take the homogeneous summand linear, respectively quadratic, in the third matrix. The Procesi–Razmyslov trace identities vanish in the fixed total word-degree once  $N$  is large enough. Declaring the marked matrix odd is the usual super-polarization: the linear part is unchanged, and the quadratic part is projected to the alternating component under

exchange of the two marked occurrences. The trace relation therefore becomes exactly graded cyclicity. These are the signs used in the finite complex for one- and two- $\psi$  words.  $\square$

*Remark 3.24.* The empty trace is the finite-rank scalar  $\text{Tr}_N(\mathbf{1}) = N$ , depending on  $N$  and omitted from the stable primitive generators. The stable symmetric algebra still has its ordinary unit; what is omitted is the empty trace as a generator of the connected primitive part. The class  $\text{Tr}(\psi)$  is a genuine stable odd primitive class. It is excluded from the Hamiltonian comparison because constants have been removed from the closed Hamiltonian algebra.

Varying

$$S = \int_{\mathbb{R}} \text{Tr}(\phi_1 d\phi_2 + \mathcal{A}[\phi_1, \phi_2])$$

with respect to  $\mathcal{A}$  gives the moment-map equation  $[\phi_1, \phi_2] = 0$ . In the reduced BV complex,

$$Q\psi = [\phi_1, \phi_2], \quad Q\phi_1 = Q\phi_2 = 0$$

on invariant trace functions.

*Construction 3.25 (Boundary evaluation of a closed Hamiltonian).* Let  $f = \sum_{k,l} c_{k,l} z_1^k z_2^l \in \mathbb{C}[z_1, z_2]$ . The brane replaces the two normal coordinates  $z_1, z_2$  by the two adjoint scalar fields  $\phi_1, \phi_2$ . Choosing the ordered lift with all  $\phi_1$ 's to the left of all  $\phi_2$ 's gives the invariant function

$$\tilde{\partial}_{\text{bb},N}(f) = \sum_{k,l} c_{k,l} \text{Tr}(\phi_1^k \phi_2^l) \in \mathcal{R}_N^{GL_N}.$$

Let

$$H_{\text{Ham}}^o(\mathcal{R}_N^{GL_N}, Q) := H^o(\mathcal{R}_N^{GL_N}, Q) / \mathbb{C} \cdot \text{Tr}(\mathbf{1})$$

as a vector space, and as a Lie algebra for the Poisson bracket. The associative operator algebra retains its unit. Constants give the scalar  $c_{0,0}N$  and therefore vanish in this Hamiltonian quotient. Thus the construction descends to a linear map

$$\partial_{\text{bb},N} : \tilde{A} \rightarrow H_{\text{Ham}}^o(\mathcal{R}_N^{GL_N}, Q), \quad [f] \mapsto \overline{\tilde{\partial}_{\text{bb},N}(f)}.$$

Let  $W_{k,l}$  be the set of words in two letters containing  $k$  copies of the first letter and  $l$  copies of the second. The first descendant is obtained by inserting one Koszul variable and summing over all ordered lifts: For  $k + l > 0$  the descendant part of the Hamiltonian comparison is

$$\partial_{\text{bb},N}^{(1)}(e_{k,l}^{\text{PBW}}) = \Psi_{k,l}, \quad \Psi_{k,l} := \sum_{w \in W_{k,l}} \text{Tr}(\psi w(\phi_1, \phi_2)).$$

Here  $e_{k,l}^{\text{PBW}}$  is an abstract special-fibre PBW label in bidegree  $(k, l)$ , distinct from the Taylor-dual coadjoint functional  $\partial_{k,l}$  and from the Matlis principal part  $\rho_{k,l}$ . The same formula at  $(k, l) = (0, 0)$  gives the decoupled scalar antifield  $\text{Tr}(\psi)$ , which is excluded from the Hamiltonian pairing because constants have been removed from  $\tilde{A}$ .

**LEMMA 3.26 (First descendants are cycles).** For every  $k, l \geq 0$ , the symmetrised trace  $\Psi_{k,l}$  satisfies  $Q\Psi_{k,l} = 0$ .

*Proof.* Write  $x = \phi_1$  and  $y = \phi_2$ . Since  $x$  and  $y$  are even and  $Q\psi = [x, y]$ , one has

$$Q\Psi_{k,l} = \sum_{w \in W_{k,l}} \text{Tr}((xy - yx)w(x, y)).$$

Group the resulting traces by cyclic word of multidegree  $(k+1, l+1)$ . The coefficient of a cyclic word in the positive sum is the number of cyclic adjacent occurrences of the block  $xy$ ; the coefficient in the negative sum is the number of cyclic adjacent occurrences of the block  $yx$ . On a circle the number of transitions from an  $x$ -run to a  $y$ -run equals the number of transitions from a  $y$ -run to an  $x$ -run. The two grouped sums are therefore equal, so  $Q\Psi_{k,l} = 0$ .  $\square$

**PROPOSITION 3.27** (*Stable primitive one-antifield Koszul homology*). Fix  $k, l \geq 0$ . The middle homology of the primitive single-trace complex with one displayed  $\psi$  and  $k, l$  displayed  $x, y$  letters is one-dimensional, generated by the class of  $\Psi_{k,l}$ . The nonconstant part relevant to the Hamiltonian comparison is the direct sum over  $k+l > 0$ .

*Proof.* Write  $x = \phi_1$  and  $y = \phi_2$ . Let  $W_{k,l}$  be the set of ordinary words with  $k$  letters  $x$  and  $l$  letters  $y$ , and let  $N_{a,b}$  be the set of cyclic words with  $a$  letters  $x$  and  $b$  letters  $y$ . By Lemma 3.23, primitive cyclic words are taken modulo the graded cyclic rule

$$\text{Tr}(ab) = (-1)^{|a||b|} \text{Tr}(ba), \quad |x| = |y| = 0, \quad |\psi| \equiv 1 \pmod{2}.$$

Graded cyclicity gives a unique representative  $\text{Tr}(\psi w)$  for every one- $\psi$  cyclic word. Thus the relevant one- $\psi$  chain group is

$$C_1(k, l) = \mathbb{C}\{W_{k,l}\}.$$

The zero- $\psi$  target is

$$C_0(k, l) = \mathbb{C}\{N_{k+1, l+1}\},$$

and the differential is the incidence map

$$\partial_1(w) = [xyw]_{\text{cyc}} - [yxw]_{\text{cyc}}.$$

The two- $\psi$  source is the alternating quotient generated by traces  $\text{Tr}(\psi u \psi v)$  with  $u$  and  $v$  ordinary words whose total bidegree is  $(k-1, l-1)$ :

$$\text{Tr}(\psi u \psi v) = -\text{Tr}(\psi v \psi u).$$

Hence diagonal pairs vanish, and if  $k = 0$  or  $l = 0$  this source is zero. Applying  $Q\psi = xy - yx$  gives

$$\begin{aligned} \partial_2(u, v) &= Q \text{Tr}(\psi u \psi v) \\ &= \text{Tr}(\psi vx yu - \psi v yxu - \psi ux yv + \psi u yxv), \end{aligned}$$

so, after removing the common front  $\psi$ ,

$$\partial_2(u, v) = vx yu - v yxu - ux yv + u yxv.$$

The displayed formula changes sign under  $(u, v) \leftrightarrow (v, u)$ , so it is well defined on the alternating quotient. A direct substitution shows  $\partial_1 \partial_2 = 0$ .

If  $k = 0$  or  $l = 0$ , then  $C_1(k, l)$  is spanned by  $x^k$  or  $y^l$ ,  $\partial_1 = 0$ , and  $C_2(k, l) = 0$ . Hence the middle homology is already the line spanned by  $\Psi_{k,l}$ . Assume from now on that  $k, l > 0$ .

The first homology is computed by the following finite complex. Form a graph  $\Gamma_{k,l}$  whose vertices are the necklaces in  $N_{k+1,l+1}$  and whose oriented edges are the words  $w \in W_{k,l}$ , with

$$w : [yxw]_{\text{cyc}} \longrightarrow [xyw]_{\text{cyc}}.$$

Then  $\partial_1$  is the cellular boundary of this graph. For each pair  $(u, v)$  as above attach the square whose oriented boundary is  $vx yu - v yxu - ux yv + u yxv$ . Denote the resulting two-complex by  $X_{k,l}$ .

Equivalently,  $X_{k,l}$  is the two-skeleton of the chip-moving state complex  $Y_{k,l}$ . The full complex  $Y_{k,l}$  has an  $r$ -cell for  $r$  mutually chosen chip moves on a cycle with  $l + 1$  oriented edges and  $k + 1$  indistinguishable chips, modulo cyclic relabelling of the edge labels. A vertex is a cyclic distribution of the chips among the vertices of this cycle, hence a necklace in  $N_{k+1,l+1}$ . A one-cell moves one chosen chip across one oriented edge; cutting the cycle at this moving edge records exactly a word  $w \in W_{k,l}$ , and its two endpoints are  $[yxw]_{\text{cyc}}$  and  $[xyw]_{\text{cyc}}$ . A two-cell records two chip moves performed independently, and its oriented boundary is precisely  $vx yu - v yxu - ux yv + u yxv$ . Therefore the cellular chain complex of  $Y_{k,l}$  through degree two is exactly

$$C_2(k, l) \longrightarrow C_1(k, l) \longrightarrow C_0(k, l),$$

and the inclusion  $X_{k,l} = Y_{k,l}^{(2)} \hookrightarrow Y_{k,l}$  induces an isomorphism on  $H_1$ , since cells of dimension at least three leave first homology unchanged.

The homology of  $Y_{k,l}$  is the homology of the cycle. The elementary circle-sum argument runs as follows. Before quotienting by cyclic relabelling, send a chip configuration to the sum of the chip positions in  $\mathbb{R}/(l + 1)\mathbb{Z}$ , and extend linearly over each cell. In the universal cover of the target circle, a fibre is described by weakly ordered lifted chip positions with fixed total sum; it is a convex polytope, subdivided by the same chip-crossing cells. Hence the circle-sum map is a cellular homotopy equivalence to  $S^1$ . The finite cyclic relabelling of the  $y$ -gaps is equivariant for this circle-sum map: it rotates the target circle, and every stabilizer acts affinely on a convex fibre, whose quotient is again contractible. Thus the quotient target is still a circle and the quotient complex  $Y_{k,l}$  has the homology of  $S^1$ . Consequently

$$H_1(X_{k,l}; \mathbb{C}) \cong H_1(S^1; \mathbb{C}) \cong \mathbb{C}.$$

Lemma 3.26 identifies this representative as a cycle in the original trace complex. The functional

$$\epsilon(\text{Tr}(\psi w)) = 1, \quad w \in W_{k,l},$$

kills every displayed two- $\psi$  boundary because each boundary has two positive and two negative summands, while

$$\epsilon(\Psi_{k,l}) = \binom{k+l}{k} \neq 0.$$

Thus the one- $\psi$  homology in bidegree  $(k, l)$  is the one-dimensional line generated by  $[\Psi_{k,l}]$ .  $\square$

*Remark 3.28.* A finite exact homology check constructs the finite complex  $C_2(k, l) \rightarrow C_1(k, l) \rightarrow C_0(k, l)$  over  $\mathbb{Q}$ , with the alternating two- $\psi$  source above, and confirms  $\dim \ker \partial_1 / \text{im } \partial_2 = 1$  for  $0 \leq k, l \leq 5$ . It also checks the open Hamiltonian descendant action of Proposition 5.171 (below) for exponents at most 3, and the closed coadjoint Taylor-dual formula together with the principal-part realization of Proposition 5.175 (below) for exponents at most 5. The finite exact check verifies the displayed finite models directly.

COROLLARY 3.29 (*PBW special-fibre one-antifield pairing*). Let  $\tilde{A}_{\text{desc}}[1]$  be the abstract shifted PBW label copy with basis  $e_{k,l}^{\text{PBW}}$  in nonconstant bidegree  $(k, l)$ . The assignment

$$e_{k,l}^{\text{PBW}} \mapsto [\Psi_{k,l}]$$

extends to an isomorphism from  $\tilde{A}_{\text{desc}}[1]$  to the nonconstant primitive one- $\psi$  brane homology.

*Proof.* The monomials  $z_1^k z_2^l$  with  $k + l > 0$  form the polynomial basis of  $\tilde{A}$  and therefore index the PBW special-fibre label copy  $\tilde{A}_{\text{desc}}$ . By Proposition 3.27, the primitive one- $\psi$  homology has one generator in each bidegree, namely  $[\Psi_{k,l}]$ . Discarding the scalar bidegree  $(0, 0)$  and sending the abstract PBW label in every nonconstant bidegree to that generator gives a filtered vector-space isomorphism. This is the boundary pairing used in the PBW special fibre. The continuous Taylor-dual/coadjoint module and the Matlis principal-part realization are separate deformation-level objects.  $\square$

LEMMA 3.30 (*Adjacent swaps*). In the invariant reduced BV cohomology,

$$\text{Tr}(u\phi_1\phi_2v) - \text{Tr}(u\phi_2\phi_1v)$$

is  $Q$ -exact for all ordinary words  $u, v$  in  $\phi_1, \phi_2$ .

*Proof.* By cyclicity of the trace,

$$\begin{aligned} Q \text{Tr}(\psi vu) &= \text{Tr}([\phi_1, \phi_2]vu) \\ &= \text{Tr}(\phi_1\phi_2vu - \phi_2\phi_1vu) \\ &= \text{Tr}(u\phi_1\phi_2v) - \text{Tr}(u\phi_2\phi_1v). \end{aligned}$$

$\square$

THEOREM 3.31 (*Cut-dependent normal-ordering straightening*). Write  $x = \phi_1$  and  $y = \phi_2$ . For every ordinary necklace  $[w]$  in  $x, y$ , choose once and for all a linear representative  $c([w])$ . Suppose  $c([w])$  has  $a$  occurrences of  $x$  and  $b$  occurrences of  $y$ , and choose a deterministic adjacent-swap path

$$c([w]) = w_0, w_1, \dots, w_r = x^a y^b,$$

where each step has the form

$$w_s = A_s y x B_s, \quad w_{s+1} = A_s x y B_s.$$

Define the one- $\psi$  chain

$$H_c([w]) := - \sum_{s=0}^{r-1} \text{Tr}(\psi B_s A_s).$$

Then

$$QH_c([w]) = \text{Tr}(c([w])) - \text{Tr}(x^a y^b).$$

Equivalently, if  $N([w]) = z_1^a z_2^b$  and  $i(z_1^a z_2^b) = \text{Tr}(x^a y^b)$ , then on this ordinary-word sector with the chosen cuts

$$QH_c = \text{id} - iN.$$

The homotopy depends on the chosen cyclic cut. If a second cut  $c'$  is chosen, then  $H_{c'}([w]) - H_c([w])$  is a closed one- $\psi$  chain; its homology class is the corresponding scalar multiple, determined by the change of cut, of the primitive descendant  $\Psi_{a-1, b-1}$  when  $a, b > 0$ . Thus this theorem is a cut-dependent straightening on the ordinary-word sector; contraction of the full cyclic open complex requires additional data.

*Proof.* For a single step  $w_s = A_s y x B_s \rightarrow w_{s+1} = A_s x y B_s$ , Lemma 3.30 gives

$$Q \operatorname{Tr}(\psi B_s A_s) = \operatorname{Tr}(A_s x y B_s) - \operatorname{Tr}(A_s y x B_s) = \operatorname{Tr}(w_{s+1}) - \operatorname{Tr}(w_s).$$

Summing over  $s$  and inserting the minus sign in the definition of  $H_c([w])$  telescopes:

$$Q H_c([w]) = - \sum_{s=0}^{r-1} (\operatorname{Tr}(w_{s+1}) - \operatorname{Tr}(w_s)) = \operatorname{Tr}(w_0) - \operatorname{Tr}(w_r).$$

Since  $w_0 = c([w])$  and  $w_r = x^a y^b$ , this is the displayed identity. A deterministic sorting rule chooses one path for each chosen cut, so the formula extends linearly to the selected zero- $\psi$  necklace span.

If  $c'$  is a cyclic rotation of  $c$ , the two telescoping formulas have the same boundary because  $\operatorname{Tr}(c([w])) = \operatorname{Tr}(c'([w]))$ . Therefore  $H_{c'}([w]) - H_c([w])$  is  $Q$ -closed. The one- $\psi$  homology computation of Proposition 3.27 identifies its class with a scalar multiple of  $\Psi_{a-1, b-1}$  determined by the change of cut; for example  $[yxx] = [xx y]$ , but the cuts  $yxx$  and  $xx y$  give homotopies differing by  $-2[\psi x]$ . This is the cut-dependence:  $H_c$  records the chosen cut datum. A cyclic contraction requires additional data.  $\square$

**COROLLARY 3.32** (*Matrix evaluation is commutative in cohomology*). If two words in  $\phi_1, \phi_2$  have the same numbers of  $\phi_1$  and  $\phi_2$ , then their traces define the same class in the reduced BV cohomology. Consequently  $\partial_{\text{bb}, N}(f)$  is determined by the commutative polynomial  $f \in \mathbb{C}[z_1, z_2]$  and is independent of the chosen ordered lift.

*Proof.* Apply Theorem 3.31 to each word. Both traces are  $Q$ -homologous to the same normal-ordered trace  $\operatorname{Tr}(\phi_1^a \phi_2^b)$ , where  $a, b$  are their common numbers of letters. Hence they define the same reduced BV cohomology class.  $\square$

**Definition 3.33** (*Unital Hamiltonian cyclic Koszul Lie model*). Let

$$E_K = \mathbb{C}x \oplus \mathbb{C}y \oplus \mathbb{C}\psi, \quad |x| = |y| = 0, \quad |\psi| = -1,$$

and let

$$\mathfrak{p}_K = \operatorname{Cyc}(T(E_K)) = T(E_K)/[T(E_K), T(E_K)]_{\text{gr}}$$

be the graded cyclic words, including the scalar word  $[1]$ . Let  $\bar{\mathfrak{p}}_K := \mathfrak{p}_K/\mathbb{C}[1]$  denote its scalar-reduced quotient. The differential is induced by

$$Qx = Qy = 0, \quad Q\psi = xy - yx.$$

With cyclic derivatives  $\partial_x, \partial_y$ , define the constant necklace bracket by

$$\{F, G\}_{\text{neck}} = [\partial_x F \partial_y G - \partial_y F \partial_x G]_{\text{cyc}},$$

with the standard Koszul sign from cyclic rotation. Since  $x, y$  are even, the displayed formula carries sign  $+1$  on the ordinary Hamiltonian words.

**THEOREM 3.34** (*Unital cyclic Koszul  $P_o$  model*). The complex  $(\mathfrak{p}_K, Q, \{-, -\}_{\text{neck}})$  is a dg Lie algebra. For every interval  $I \subset \mathbb{R}$ ,

$$\mathfrak{p}_K(I) := \Omega_c^\bullet(I) \otimes \mathfrak{p}_K$$

with differential  $d_{\mathbb{R}} + Q$  and bracket

$$[\alpha \otimes F, \beta \otimes G] = (-1)^{|F||\beta|}(\alpha \wedge \beta) \otimes \{F, G\}_{\text{neck}}$$

is a compactly supported current dg Lie algebra. The graded-commutative algebra

$$\text{Obs}_{\text{open}}^{\text{cyc}}(I) := \mathbf{S}(\mathfrak{p}_K(I))$$

with the biderivation extending this bracket is a  $P_o$  factorization algebra on the brane line, with structure maps given by extension by zero and multiplication on disjoint intervals.

The CE/PV comparison used later is the degree-zero Hamiltonian identity of Theorem 5.2; extending it to all  $\psi$ -words requires a graded coordinate theorem with the full signs and degree shifts supplied as data.

*Proof.* The necklace bracket is the standard constant-symplectic necklace bracket for the even pair  $x, y$ . Its graded antisymmetry and Jacobi identity are the usual cancellation of the cyclic sum of commutators in  $T(E_K)$ . The scalar word must be present: for example  $\{[x], [y]\}_{\text{neck}} = [1]$ . The derivation  $Q$  replaces only uncontracted occurrences of  $\psi$  by  $x\gamma - y\alpha$  and fixes the paired variables  $x, y$ . Expanding the bracket as the signed sum over cut pairs  $(x \in F, y \in G)$  and  $(y \in F, x \in G)$ ,  $Q$  never hits a contracted letter and carries the same super-cyclic cut sign as applying  $Q$  before bracketing; the terms where  $Q$  hits  $G$  acquire the usual factor  $(-1)^{|F|}$ . Hence

$$Q\{F, G\}_{\text{neck}} = \{QF, G\}_{\text{neck}} + (-1)^{|F|}\{F, QG\}_{\text{neck}}.$$

Finally  $Q^2 = 0$  is immediate from  $Qx = Qy = 0$ .

Tensoring with  $\Omega_c^\bullet(I)$  gives the displayed current dg Lie algebra by the standard tensor-product sign rule;  $d_{\mathbb{R}}$  is a derivation of  $\wedge$ , and  $Q$  is a derivation of the necklace bracket. For disjoint intervals  $I_1, \dots, I_n \subset J$ , extension by zero sends each current into  $J$ , and the factorization product is the product in the symmetric algebra. If two inputs have disjoint support, their current bracket has zero de Rham product, so the  $P_o$ -bracket is local for the factorization structure.

The final assertion gives the stated boundary on the claim: an all-degree cotangent-coordinate comparison is additional data beyond this theorem.  $\square$

**PROPOSITION 3.35** (*Normal-ordering quotient of the unital cyclic Koszul model*). Define

$$N : \mathfrak{p}_K \longrightarrow \mathbb{C}[z_1, z_2]$$

on a cyclic word by

$$N([w]) = \begin{cases} z_1^{\#x(w)} z_2^{\#y(w)}, & w \text{ is an ordinary } x, y\text{-word,} \\ 0, & w \text{ contains at least one } \psi. \end{cases}$$

Then  $N$  is a dg Lie morphism from  $(\mathfrak{p}_K, Q, \{-, -\}_{\text{neck}})$  to the unreduced polynomial Hamiltonian Lie algebra  $\mathbb{C}[z_1, z_2]$  with zero differential and Poisson bracket  $\partial_{z_1} \wedge \partial_{z_2}$ . Composing with  $\mathbb{C}[z_1, z_2] \rightarrow \bar{\mathcal{A}}$  gives the reduced Hamiltonian quotient  $N_{\text{red}} : \bar{\mathfrak{p}}_K \rightarrow \bar{\mathcal{A}}$ . The section  $i(z_1^a z_2^b) = [x^a y^b]$  and the cut-dependent homotopy  $H_c$  of Theorem 3.31 satisfy  $QH_c = \text{id} - iN$  on the chosen zero- $\psi$  necklace sector. The full cyclic complex requires a separate construction.

*Proof.* If a word contains  $\psi$ , every term of its necklace bracket with any word still contains  $\psi$ , because the bracket only differentiates in  $x$  and  $y$ . Hence both sides of the Lie identity vanish after  $N$  whenever one input contains  $\psi$ . For ordinary words  $F, G$ , cyclic derivative counting gives

$$N(\partial_x F) = \partial_{z_1} N(F), \quad N(\partial_y F) = \partial_{z_2} N(F),$$

and similarly for  $G$ . Therefore

$$N\{F, G\}_{\text{neck}} = \partial_{z_1} N(F) \partial_{z_2} N(G) - \partial_{z_2} N(F) \partial_{z_1} N(G) = \{N(F), N(G)\}.$$

The chain-map property follows generatorwise and then by the derivation rule. If a word contains one  $\psi$ , each  $Q$ -summand replaces that  $\psi$  by  $xy - yx$ , and the two resulting ordinary commutative monomials agree after  $N$ . If a word contains two or more  $\psi$ 's, every  $Q$ -summand still contains a  $\psi$ , hence maps to zero. On ordinary words  $Q = 0$ . Thus  $NQ = 0$ . The final displayed straightening identity is precisely Theorem 3.31 with the chosen cuts.  $\square$

LEMMA 3.36 (*Obstruction to contraction onto the Hamiltonian quotient*). A degree  $-1$  operator  $K$  on

$$\mathfrak{p}_K = \text{Cyc}(T(x, y, \psi))$$

satisfying

$$QK + KQ = \text{id} - iN$$

with  $N$  killing every  $\psi$ -containing word would force the primitive one- $\psi$  classes to vanish.

*Proof.* For each nonconstant bidegree  $(k, l)$ , Proposition 3.27 gives a closed one- $\psi$  primitive  $\Psi_{k,l}$  whose class is nonzero. Since  $N(\Psi_{k,l}) = 0$ , the displayed identity would imply

$$\Psi_{k,l} = QK(\Psi_{k,l}) + KQ(\Psi_{k,l}) = QK(\Psi_{k,l}),$$

contradicting  $[\Psi_{k,l}] \neq 0$ . Hence any unreduced normal-ordering transfer must either retain the descendant module and higher coherence homotopies, or restrict to the cut-dependent zero- $\psi$  sector above.  $\square$

LEMMA 3.37 (*Gauge-paired necklace obstruction to the naive Hamiltonian quotient*). Enlarge  $E_K$  to  $E_{\text{BV}} = \mathbb{C}x \oplus \mathbb{C}y \oplus \mathbb{C}a \oplus \mathbb{C}\psi$  with  $|a| = 1, |\psi| = -1$ , and add the constant pairing  $\langle a, \psi \rangle = 1$  to the necklace bracket. The map that sends an  $x, y$ -word to its commutative monomial and kills every word containing  $a$  or  $\psi$  has a nonzero Lie defect.

*Proof.* With left cyclic derivatives and the displayed pairing convention,

$$\{[a], [\psi x]\}_{\text{neck}} = [x]$$

exactly. The naive projection sends the left-hand bracket to  $z_1$ , while it sends both  $[a]$  and  $[\psi x]$  individually to zero, so the bracket of their projected images is zero. Thus the full gauge-paired cyclic BV algebra requires an additional quotient, homotopy transfer, or explicit centrality data before it can map to the Hamiltonian formal-disk quotient.  $\square$



THEOREM 3.38 (*Gauge-BRST cyclic algebra and the normal-ordering obstruction*). Let

$$E_{\text{BRST}} = \mathbb{C}x \oplus \mathbb{C}y \oplus \mathbb{C}\gamma \oplus \mathbb{C}\psi, \quad |x| = |y| = 0, \quad |\gamma| = 1, \quad |\psi| = -1,$$

and put

$$\mathfrak{p}_{\text{BRST}} = \text{Cyc}(T(E_{\text{BRST}})).$$

Equip  $E_{\text{BRST}}$  with the even constant pairing

$$\langle x, y \rangle = 1, \quad \langle y, x \rangle = -1, \quad \langle \gamma, \psi \rangle = \langle \psi, \gamma \rangle = 1,$$

and the corresponding graded necklace bracket. The cyclic Hamiltonian

$$S = [\gamma(xy - yx)] + [\psi\gamma\gamma]$$

has degree 1 and satisfies  $\{S, S\}_{\text{neck}} = 0$ . Hence  $Q = \{S, -\}_{\text{neck}}$  makes  $\mathfrak{p}_{\text{BRST}}$  a dg Lie algebra, with

$$Q\gamma = \gamma^2, \quad Qx = \gamma x - x\gamma, \quad Qy = \gamma y - y\gamma, \quad Q\psi = xy - yx + \gamma\psi + \psi\gamma.$$

The normal-ordering map

$$N_{\text{BRST}}: \mathfrak{p}_{\text{BRST}} \longrightarrow \mathbb{C}[z_1, z_2], \quad [w] \longmapsto \begin{cases} z_1^{\#_x(w)} z_2^{\#_y(w)}, & w \text{ contains only } x, y, \\ 0, & w \text{ contains } \gamma \text{ or } \psi, \end{cases}$$

is a chain map with a nonzero Lie defect. A contraction  $QK + KQ = \text{id} - iN_{\text{BRST}}$  on the full cyclic complex is obstructed by the length-one class  $[\psi]$ .

*Proof.* The constant-pairing necklace bracket has graded Jacobi by the standard two-cut cancellation in cyclic words. The displayed Hamiltonian gives the four formulas for  $Q$  by direct cyclic differentiation. A direct check of the generators gives  $Q^2 = 0$ :  $Q\gamma^2 = 0$ ,  $Q^2x = Q(\gamma x - x\gamma) = 0$ ,  $Q^2y = 0$ , and, for  $\mu = xy - yx$ ,  $Q\mu = \gamma\mu - \mu\gamma$ , whence

$$Q^2\psi = Q\mu + Q(\gamma\psi) + Q(\psi\gamma) = (\gamma\mu - \mu\gamma) + (-\gamma\mu - \gamma\psi\gamma) + (\mu\gamma + \gamma\psi\gamma) = 0.$$

Therefore  $Q = \{S, -\}$  is a square-zero derivation of the necklace bracket.

The chain-map statement for  $N_{\text{BRST}}$  follows as in Proposition 3.35: ordinary words are  $Q$ -closed after projection; a single  $\psi$  contributes  $xy - yx$ , whose two normal orderings agree, while the  $\gamma\psi$  and  $\psi\gamma$  terms are killed; every summand containing  $\gamma$  is killed. Its Lie defect is the gauge-paired bracket  $\{[\gamma], [\psi x]\}_{\text{neck}} = [x]$ , whose image is  $z_1$ , whereas both inputs have zero image. Finally  $[\psi]$  is closed:

$$Q[\psi] = [xy - yx] + [\gamma\psi + \psi\gamma] = 0,$$

since  $[xy] = [yx]$  and  $[\psi\gamma] = -[\gamma\psi]$  in the cyclic quotient. The image of  $Q$  misses  $[\psi]$ . With respect to cyclic word length,  $Q$  raises length by one:  $Qx, Qy, Q\gamma$ , and  $Q\psi$  are quadratic words. Hence the length-one class  $[\psi]$  represents nonzero homology. Equivalently, the derivation rule never produces a lone  $\psi$ -letter. Thus a full contraction would force  $[\psi] = QK[\psi]$ , contradiction. The extended BRST cyclic algebra is the correct unreduced primitive object; the Hamiltonian normal-ordering sector is only a quotient/chain-map sector and still requires separate centrality and factorization-center data.  $\square$

PROPOSITION 3.39 (*Stalk central multiplication*). Let

$$A_N^{\text{pt}} = (\mathcal{R}_N^{GL_N}, Q)$$

be the invariant reduced open Koszul algebra at a brane insertion point. For  $f \in \mathbb{C}[z_1, z_2]$ , set

$$B_f = \tilde{\partial}_{\text{bb}, N}(f) = \sum_{k, l} c_{k, l} \text{Tr}(\phi_1^k \phi_2^l).$$

Then  $QB_f = 0$ . Regard  $B_f$  as a Hochschild o-cochain; its associated central operation is multiplication

$$L_{B_f} : A_N^{\text{pt}} \rightarrow A_N^{\text{pt}}, \quad a \mapsto B_f a.$$

This Hochschild o-cochain is closed. The same statement holds for multiplication by a closed one-antifield class  $\Psi_{k, l}$ . After projection to the Hamiltonian quotient and stabilization in  $N$ , the degree-zero classes satisfy

$$\{B_f, B_g\}_{\text{open}} = B_{\{f, g\}_{\tilde{A}}}.$$

*Proof.* Since  $Q\phi_1 = Q\phi_2 = 0$ , every  $B_f$  is  $Q$ -closed. The algebra  $A_N^{\text{pt}}$  is graded-commutative, so the Hochschild differential of a o-cochain  $b$  is

$$d_{\text{Hoch}} b(a) = Qb a + ba - (-1)^{|b||a|} ab.$$

This vanishes for every closed graded-central element  $b$ . The classes  $\Psi_{k, l}$  are closed by Lemma 3.26 and are central for the same graded-commutative reason. This is only a statement about multiplication by a closed polynomial descendant in the stalk algebra:  $\Psi_{k, l}$  is a PBW special-fibre descendant distinct from the principal-part cotangent boundary field  $\rho_{k, l}$ . The continuous Matlis coadjoint action is the separate action of Theorem 5.178. Ordered-lift independence for  $B_f$  is cohomological and is exactly Corollary 3.32. The Poisson identity follows from Theorem 5.190 (below).  $\square$

Let  $\tilde{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1$ . The removed constant corresponds on the gauge side to the scalar  $\text{Tr}(1) = N$  and has zero Hamiltonian vector field.

PROPOSITION 3.40 (*Stable brane Hamiltonian algebra and connected quotient*). Put

$$V_H = \tilde{A} \oplus \tilde{A}_{\text{desc}}[1].$$

In the stable large- $N$  open BV model, the Hamiltonian descendant algebra generated by the nonempty primitive ghost-zero traces and the nonconstant primitive one- $\psi$  Koszul homology is

$$\mathcal{O}_{\text{brane}, H, \text{st}}^{\text{cl}} \cong \mathbf{S}(V_H).$$

Its connected single-trace, or indecomposable, quotient is

$$Q\mathcal{O}_{\text{brane}, H, \text{st}}^{\text{cl}} = \mathfrak{m}/\mathfrak{m}^2 \cong V_H.$$

Under  $J$ , the class  $z_1^k z_2^l$  with  $k + l > 0$  maps to  $\overline{\text{Tr}(\phi_1^k \phi_2^l)}$ ; its descendant basis vector maps to the symmetrised trace with one  $\psi$  and  $k, l$  copies of  $\phi_1, \phi_2$ . All Lie comparisons on single traces use  $V_H$  after scalar reduction; CE/PV polyvector functions use the commutative algebra  $\mathbf{S}(V_H)$ .

*Proof.* Proposition 3.22 gives the free commutative algebra on cyclic words. Lemma 3.30 identifies any two cyclic words with the same numbers of  $\phi_1$  and  $\phi_2$ , because adjacent transpositions generate all reorderings. Thus the connected ghost-zero cyclic words reduce to the monomials  $\overline{\text{Tr}(\phi_1^k \phi_2^l)}$  with  $k + l > 0$ . Proposition 3.27 computes the primitive one- $\psi$  homology in each multidegree; the nonconstant generators are  $\Psi_{k,l}$  with  $k + l > 0$ . Products of the connected ghost-zero primitive classes and these one- $\psi$  primitive classes give the free graded-commutative multi-trace algebra  $\mathbf{S}(V_H)$  in the Hamiltonian sector. Projection to indecomposables  $\mathfrak{m}/\mathfrak{m}^2$  recovers the connected single-trace quotient  $V_H$ . The full primitive calculation proves that stable primitive homology vanishes for  $\psi$ -degree  $p \geq 2$ ; decomposable products of one- $\psi$  classes remain only as products in the symmetric algebra.  $\square$

*Definition 3.41 (Three large- $N$  operations).* The three operations have different domains and are applied in the order below.

- (i) *Degreewise invariant stabilisation*  $\text{Prim}_{\text{st}}$ . At fixed word degree  $d$ , take  $N \geq d$  so that Procesi–Razmyslov trace invariants lie beyond finite- $N$  trace identities. This is the operation used in Proposition 3.22.
- (ii) *Connected single-trace extraction*  $\text{Conn}_{N \rightarrow \infty}$ . After passing to the stable invariant algebra, project the symmetric algebra on cyclic-word generators to its primitive/indecomposable summand. Products of two nonempty traces are decomposable and are killed by this projection.
- (iii) *Hamiltonian scalar quotient*  $\pi_{\text{Ham}}$ . Remove the empty trace  $\text{Tr}(\mathbf{1}) = N$  on the open side and the constant Hamiltonian  $\mathbf{1}$  on the closed side. The removed line is the scalar  $U(\mathbf{1})$  line whose obstruction class is  $[\bar{c}]$ .

The reduced Hamiltonian sector used below is

$$\pi_{\text{Ham}} \text{Conn}_{N \rightarrow \infty} \text{Prim}_{\text{st}}(\mathcal{R}_N^{GL_N}, Q).$$

The formula is an ordered construction: first stabilize invariant theory degree by degree, then extract connected primitive trace classes, then quotient the scalar Hamiltonian line. The unprojected  $N \rightarrow \infty$  algebra and the permuted composites are different objects.

*Remark 3.42.* The word “sector” is literal. The calculation above presents the stable algebra generated by the primitive zero- $\psi$  Hamiltonians and the nonconstant primitive one- $\psi$  homology generated by  $\Psi_{k,l}$  for  $k + l > 0$ . The full invariant Koszul homology in higher  $\psi$ -count lies outside this sector, as does the decoupled scalar class  $\text{Tr}(\psi)$ .

**COROLLARY 3.43** (*Full primitive Koszul homology*). Let

$$A_\psi = \mathbb{C}\langle x, y, \psi \rangle, \quad |x| = |y| = 0, \quad |\psi| \equiv 1 \pmod{2},$$

with differential  $d\psi = x\gamma - yx$  and  $dx = d\gamma = 0$ . Put

$$\text{Cyc}(A_\psi) = A_\psi / [A_\psi, A_\psi]_{\text{gr}}, \quad [uv] = (-1)^{|u||v|} [vu].$$

Give  $x, y, \psi$  weights

$$\text{wt}(x) = (1, 0), \quad \text{wt}(y) = (0, 1), \quad \text{wt}(\psi) = (1, 1).$$

For fixed  $(R, S)$  define

$$K_{R,S}^p = \mathbb{C}\{[w]_{\text{cyc}} : \#\psi(w) = p, \#x(w) = R - p, \#y(w) = S - p\},$$

with  $0 \leq p \leq \min(R, S)$ . The differential is

$$\begin{aligned} \partial[w] = \sum_{i: a_i = \psi} (-1)^{\#\{\psi \text{ before } i\}} [a_1 \cdots a_{i-1} x y a_{i+1} \cdots a_n \\ - a_1 \cdots a_{i-1} y x a_{i+1} \cdots a_n]_{\text{cyc}}. \end{aligned}$$

The primitive homology is

$$H_p(K_{R,S}) \cong \begin{cases} \mathbb{C} \cdot [x^R y^S], & p = 0, \\ \mathbb{C} \cdot [\Psi_{R-1, S-1}], & p = 1, R, S \geq 1, \\ 0, & p \geq 2. \end{cases}$$

Equivalently, the primitive stable Koszul homology is  $\mathbb{C}[x, y] \oplus \mathbb{C}[x, y]\theta$  with  $|\theta| = 1$  and  $\text{wt}(\theta) = (1, 1)$ .

This is a stable primitive cyclic-word computation. The proof identifies the graded cyclic complex with the cellular chains of the chip-moving necklace state complex and uses the circle-sum map to compute the state complex as  $H_\bullet(S^1; \mathbb{C})$ ; see Theorem 5.128 and Theorem D.12. The result does identify the polynomial one- $\psi$  span as a PBW special-fibre object; Matlis principal parts and the Moyal coadjoint module are separate deformation-level objects.

*Remark 3.44 (Open-trace  $A_\infty$ -Koszul acceptance datum).* After the Capelli renormalisation (Lemma 5.197, Corollary 5.198) the stable connected single-trace target is the Lie-algebra image of the Moyal Hamiltonian source on the degree-zero generator restriction (Corollary 5.200); the residual obstruction to the all-bidegree ordinary Weyl-trace bracket comparison is the quantum shear primitive

$$E_{a,b,N} \in \mathcal{W}_N \widehat{\mu}(\mathfrak{sl}_N), \quad \mathfrak{v}_{a,b} = [R_{a,b}^{\text{free}}] \in \text{coker } C_{a,b},$$

equivalently the dual-potential class  $\Omega_{a,b}^{\text{rad}} = [(T_{a,b}, E_{a,b}^+)] \in \text{coker } B_{a,b}$  of Theorem 5.216; obstruction is exactly a signed row  $(\phi, -\lambda) \in \ker B_{a,b}^*$  with  $\lambda(E_{a,b}^+) - \phi(T_{a,b}) \neq 0$ . The edge strips  $\mathfrak{v}_{2,b}$  and  $\mathfrak{v}_{a,2}$  vanish (Proposition F.32); the finite-window interior identities are recorded in Proposition F.33; the all-interior assertion is the decorated PBW Stokes signed-row vanishing theorem. The decoupled scalar class  $\text{Tr}(\psi)$  and the higher  $\psi$ -count primitive homology lie outside the sector and outside this acceptance datum. Stable large- $N$  brane-trace asymptotics enter only through Definition 4.26.

The open BV brane sector is the derived commuting variety  $[\mu_{\text{der}}^{-1}(0)/\text{GL}_N]$  with Koszul antifield  $\psi$  and BV differential  $Q\psi = [\phi_1, \phi_2]$ , with stable trace observables generating the open-string sector classical observables factorization algebra on the brane line. The trace functional of a closed Hamiltonian  $f \in \mathfrak{h}$  on this brane sector is the  $\text{GL}_N$ -invariant  $J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)}$ ; its universal property and CE-coordinate expression are the content of the section below.

#### 4 BOUNDARY ALGEBRA AND TRACE MAP

The trace map  $J(f) = \text{Tr } f(\phi_1, \phi_2)$  is forced by Procesi–Razmyslov stable invariants, Loday–Quillen–Tsygan primitivity, and Darboux naturality on the moment-map zero fibre; adjacent-swap exactness makes it well defined on the derived stack (Theorem 5.6, step (3) of the proof). Operational test: Remark 5.9, probe (3).

## 4.1 The reduced CE/PV trace map

The trace map at a Dirac brane vacuum is

$$J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)},$$

the formal Darboux coordinate expression at  $b_p \in C_\partial^{\text{op}}$  of the brane-side derived  $E_1$ -centre  $Z_{E_1}^{\text{der}}(A_{\partial, N}^{\text{cl}}) \simeq HH^\bullet(A_{\partial, N}^{\text{cl}}, A_{\partial, N}^{\text{cl}})$  (Deligne's  $E_1$ -Hochschild theorem), which the Mixed Holomorphic–Topological Deligne conjecture identifies at the formal-Darboux stalk with the bulk  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_{b_p}$  (the chiral  $E_2$  factorization algebra on  $\mathbb{C}^2$  restricted to the formal neighbourhood of  $b_p$ ; §). Here  $b_p \in C_\partial^{\text{op}}$  is the chosen vacuum at  $p = \mathbf{o} \in \mathbb{C}^2$  (Definition 1.1), with chart algebra  $A_{b_p} = \text{End}_{C_\partial^{\text{op}}}(b_p) \cong \mathbb{C}[[z_1, z_2]]$  on the brane line  $L = \mathbb{R}_t \times \{z_1 = z_2 = \mathbf{o}\}$  of §3;  $J$  is the stable scalar-reduced cyclic-trace class of the unreduced  $J(f) = \text{Tr } f(\phi_1, \phi_2)$ . The closed BV theory carries two CE coordinate types and the chart carries only one of them: Hamiltonian ghosts  $c^I$ —dual to generators  $H_I \in \widehat{\mathfrak{h}}$ —generate the closed motion on the polydisk; cotangent coordinates  $u_I$ —dual to the shifted antifield generators in  $\mathfrak{h}_{\text{cont}}^\vee[1]$ —are the boundary observables. The map  $J$  is the unique  $\mathbb{C}$ -linear,  $GL_N$ -equivariant, Darboux-natural primitive single-trace map  $\mathfrak{h} \rightarrow \mathcal{O}(\mu_{\text{der}}^{-1}(\mathbf{o}))^{GL_N}$  satisfying scalar reduction, Dirac descent, and  $J(z_i) = \text{Tr } \phi_i$  (Theorem 5.146). Centrality is the moment-map identity  $\{J(f), J(g)\}_\partial = J(\{f, g\}_{\mathfrak{h}})$  on the chart  $A_{b_p}$ . Closed Hamiltonians act through  $c_f$ ;  $J$  records  $u_f$ ; below, the symbols  $c_f, u_f, \theta_f, O_f$  refer back to Lemma 4.1.

The CE/PV comparison is graded into five levels at separate admissibility hypotheses.

| level | statement                                                                                         | admissibility                                                                                                         |
|-------|---------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------|
| 1     | coordinate cochain identity $\Phi_{\text{coord}}(c^I) = \theta^I, \Phi_{\text{coord}}(u_I) = O_I$ | none (unconditional)                                                                                                  |
| 2     | $P_{\mathbf{o}}$ -bracket enhancement on the chosen sub-PV model                                  | bracket-admissibility of $B$                                                                                          |
| 3     | Maurer–Cartan kernel compatibility                                                                | kernel datum $(B, \mathcal{K}_B)$                                                                                     |
| 4     | bar-cobar / enveloping upgrade $\Omega C_{\text{CE}}^\bullet \xrightarrow{\sim} \mathcal{A}$      | Tate / nilpotent / weighted datum                                                                                     |
| 5     | open-trace $\mathcal{A}_\infty$ -Koszul upgrade                                                   | $\mathcal{A}_\infty$ -acceptance datum $(\mathcal{A}_{\text{op}}, C_{\text{op}}, B_{\text{CE/PV}}, \tau_{\text{op}})$ |

The unconditional content is level 1. Levels 2–5 are conditional, each on a separate admissibility datum stated in Theorem 5.5:  $B$  is a graded sub-Lie algebra of the polyvector coordinate ring  $\text{PV}_{\text{coord}}$  closed under the Schouten bracket (bracket-admissibility, Definition 4.18);  $\mathcal{K}_B$  is a target completion of the diagonal tensor product in which the Casimir converges (kernel-admissibility); the Tate / nilpotent / weighted datum specifies a bar-cobar replacement of the closed CE complex with a conilpotent endpoint and  $\varprojlim^1$ -vanishing; and the  $\mathcal{A}_\infty$ -acceptance datum  $(\mathcal{A}_{\text{op}}, C_{\text{op}}, B_{\text{CE/PV}}, \tau_{\text{op}})$  collects an open-trace  $\mathcal{A}_\infty$  algebra  $\mathcal{A}_{\text{op}}$ , a Koszul-dual coalgebra  $C_{\text{op}}$ , the matching CE/PV bracket model  $B_{\text{CE/PV}}$ , and a twisting cochain  $\tau_{\text{op}}$ . References below to “the CE/PV comparison” identify their level explicitly.

LEMMA 4.1 (*Two CE coordinates of a Hamiltonian*). A Hamiltonian  $f \in \mathfrak{h}$  carries two CE coordinates on the closed shifted-cotangent BV side.

- As generator of canonical transformations:  $c_f \in \mathfrak{h}^\vee[-1]$  is the Hamiltonian-dual CE coordinate, and  $\Phi_{\text{coord}}(c_f) = \theta_f$ , the Schouten vector field on the boundary observable algebra.
- As closed cotangent insertion:  $u_f \in \mathfrak{h} \subset \mathfrak{g}[1]$  is the shifted-cotangent-dual coordinate, and  $\Phi_{\text{coord}}(u_f) = O_f$ , the boundary observable  $J(f) = \text{Tr } f(\phi_1, \phi_2)$  on the chart  $A_b$  after stable scalar reduction.

Closed Hamiltonians act through  $c_f$ ; the trace map  $J$  records  $u_f$ ; the CE/PV theorem (Theorem 5.5, Theorem 4.20) implements the duality.

*Proof.* Theorem 5.2 fixes the generator signs in the finite-dimensional case:  $\Phi_1(c^\xi) = \theta^\xi$ ,  $\Phi_1(u_x) = O_x$ . Theorem 5.3 extends this generator assignment to the formal-polydisk coordinate envelope. Equating  $c_f$  with  $O_f$  breaks the coordinate bracket-compatibility  $[\theta^I, O_J]_S = \delta_J^I$ , which is what every  $P_o$ -centrality homotopy uses (clause (2) of Theorem 5.5). The collision is already at the bracket on generators, before any central extension or kernel datum can enter: the obstruction is the generator-level bracket defect.  $\square$

*Remark 4.2 (Generator table).*

| symbol | CE coordinate on                            | $\Phi_{\text{coord}}$ -image                              | physical role                 |
|--------|---------------------------------------------|-----------------------------------------------------------|-------------------------------|
| $c_f$  | $\mathfrak{h}^\vee[-1]$                     | $\theta_f \in \text{Vect}(\widehat{\mathbf{S}}(\bar{A}))$ | generator of canonical motion |
| $u_f$  | $(\mathfrak{h}_{\text{cont}}^\vee[1])^\vee$ | $O_f = J(f) \in \widehat{\mathbf{S}}(\bar{A})$            | chart trace map $J$ on $A_b$  |

LEMMA 4.3 (*Bracket on observables*). For  $f, g \in \mathfrak{h}$ ,

$$\theta_f(O_g) = O_{\{f, g\}}.$$

Equivalently, the Schouten field  $\theta_f = \Phi_{\text{coord}}(c_f)$  acts on the boundary observable  $O_g = \Phi_{\text{coord}}(u_g) = J(g)$  as the Hamiltonian derivation  $g \mapsto \{f, g\}$ .

*Proof.* On generators  $c_f, u_g$  of  $C_{\text{CE, coord}}^\bullet(\mathfrak{g})$  the CE differential gives  $d_{\text{CE}} u_g(c_f) = u_{\{f, g\}}$ . The CE/PV identification Theorem 5.2 sends  $c_f \mapsto \theta_f$  and  $u_g \mapsto O_g$ , and the Schouten pairing  $[\theta_f, O_g]_S = \theta_f(O_g)$  on  $\text{PV}(\widehat{\mathbf{S}}(\bar{A}))$  carries the same derivation on linear functions; the equality on the brane side is the moment-map identity  $\{J(f), J(g)\}_\partial = J(\{f, g\}_\mathfrak{h})$  of Theorem 5.190.  $\square$

*The CE/PV dictionary.* On generators the cochain CE/PV identification of Theorem 4.20 is

$$c^I \mapsto \theta^I, \quad u_I \mapsto O_I.$$

This is Lemma 4.1 written on a basis: the Hamiltonian-dual CE coordinate produces a Schouten vector field, the shifted-cotangent-dual CE coordinate produces a boundary Hamiltonian observable. The brane substitution is the image of  $u_f$ ;  $c_f$  is the Hamiltonian vector-field coordinate.

*Word ordering is BRST-exact, the bracket records the symbol.* At finite  $N$  the trace map  $J(f) = \text{Tr } f(\phi_1, \phi_2)$  (equivalently  $\overline{\text{Tr } f(\phi_1, \phi_2)}$  after stable scalar reduction) requires a word ordering of the noncommuting matrices in  $f(\phi_1, \phi_2)$ . The choice is irrelevant in  $H^0(\mathcal{R}_N^{GLN}, Q)$ : adjacent swaps are exact via  $Q \text{Tr}(\psi v u) = \text{Tr}([\phi_1, \phi_2] v u)$  of (8), so  $J(f)$  depends only on the symbol  $f$ . The matrix Poisson bracket induced by  $\{(\phi_1)^i_j, (\phi_2)^k_l\} = \delta_l^i \delta_j^k$  is

$$\{F, G\} = \text{Tr}(\partial_{\phi_1} F \cdot \partial_{\phi_2} G - \partial_{\phi_2} F \cdot \partial_{\phi_1} G)$$

with cyclic matrix derivatives. On monomials,

$$\{\overline{\text{Tr}(\phi_1^k \phi_2^l)}, \overline{\text{Tr}(\phi_1^m \phi_2^n)}\} = (kn - lm) \overline{\text{Tr}(\phi_1^{k+m-1} \phi_2^{l+n-1})}, \quad (12)$$

and assembling over the symbol algebra after scalar reduction,

$$\{J(f), J(g)\}_\partial = J(\{f, g\}_\mathfrak{h}).$$

$J$  is a Lie algebra morphism  $\bar{A} \rightarrow H^0(\mathcal{R}_N^{GLN}, Q)$  modulo the central direction §6.1 diagnoses.

*Finite Taylor truncation obstruction.* The naive finite-rank reduction of  $\widehat{\mathfrak{h}}$  is its Taylor-degree truncation  $\widehat{\mathfrak{h}}/\mathfrak{m}^{N+1}$ . The Lie-quotient obstruction is explicit: linear Hamiltonians  $z_1, z_2$  generate brackets of strictly lower Taylor degree, the projected brackets have a Jacobi obstruction after truncation, and the diagonal cotangent Casimir  $C_{\widehat{\mathfrak{h}}} = \sum_{d \geq 1} e_{d,i} \otimes e_d^i$  misses the strict product/direct-sum endpoint. Finite coordinate projections exist only as coefficient tests. They assemble to the completed coordinate CE/PV isomorphism

$$\Phi_{\text{coord}} : C_{\text{CE,coord}}^{\bullet}(\mathfrak{g}) \xrightarrow{\sim} \text{PV}_{\text{coord}}(A_{\partial,\text{coord}})$$

of Theorem 4.20. A  $P_0$  comparison is asserted only after a bracket-admissible  $B \subset \text{PV}_{\text{coord}}(A_{\partial,\text{coord}})$  is chosen (Definition 4.18); a Maurer–Cartan kernel is asserted only after the diagonal tensor converges in a kernel-admissible tensor product. Both are part of the reduced Hamiltonian central-operation model; the strict ambient product has a separate convergence obstruction.

**THEOREM 4.4** (*Witten centrality on the Dirac brane: the local moment-map trace identity*). In the reduced classical  $P_0$  factorization algebra  $A_{\partial}$  on the brane line, the trace map  $J : \bar{A} \rightarrow A_{\partial}, J(f) = \text{Tr } f(\phi_1, \phi_2)$ , with smeared form  $f \mapsto B_f(a) = \int_L a(t) J(f)(t) dt$ , realises every closed Hamiltonian  $f \in \bar{A}$  as a boundary observable whose ordered products and  $P_0$ -bracket with smeared Hamiltonian insertions  $B_g(b)$  are the moment-map operations

$$\{B_f(a), B_g(b)\} = B_{\{f,g\}}(ab), \quad F \cdot G \simeq (-1)^{|F||G|} G \cdot F \text{ (scalar } F\text{)}.$$

In particular  $J$  lands inside the reduced  $P_0$ -central-operation complex of  $A_{\partial}, J(f)$  is null-homotopic for every scalar (constant)  $f$ , and  $J$  is Witten’s classical centrality computation [4] read in the local algebraic model  $Z_{\text{red,Ham}}^{P_0}(A_{\partial,\text{coord}}; B) = B$ . The unreduced  $P_0$ -factorization centre lift, including centrality homotopies against arbitrary open observables and principal-part cotangent currents, is exactly Problem 5.48.

*Proof.* The brane equal-time Poisson bracket induced by the symplectic potential  $\theta_N = \text{Tr}(\phi_1 d\phi_2)$  of Lemma 3.15 is, on two brane-time points  $(t, t')$ ,

$$\{(\phi_1)^i_j(t), (\phi_2)^k_l(t')\} = \delta_l^i \delta_j^k \delta(t - t').$$

Smearing by  $a(t)b(t')$  replaces  $\delta(t - t')$  by the product  $ab$ ; contraction of the matrix derivatives of  $\text{Tr } f(\phi_1, \phi_2)$  and  $\text{Tr } g(\phi_1, \phi_2)$  returns the constant Poisson bracket on the symbols, evaluated again on the Chan–Paton pair. This produces the moment-map identity  $\{B_f(a), B_g(b)\} = B_{\{f,g\}}(ab)$ ; on monomials the explicit coefficient  $kn - lm$  is the open trace bracket (12). Cyclic invariance and the  $Q$ -exact swaps reduce every monomial bracket to its symbol-level Hamiltonian bracket on  $\bar{A}$ .

Constants in  $\bar{A}$  have zero Hamiltonian vector field, so the scalar bracket component is null-homotopic; the central direction  $\mathbb{C} \cdot J(1)$  is exactly the  $U(1)$  centre treated in §6.1. The locally constant brane-line factorisation product compares two ordered products of a closed and an open insertion; the homotopy connecting them is the Hochschild centrality datum of the associated  $E_1$ -algebra stalk, and the  $P_0$  refinement imposes the same compatibility for the matrix Poisson bracket. Both are produced by the moment-map identity above.

The unreduced cotangent statement requires centrality homotopies for arbitrary open observables, including the principal-part cotangent currents  $\Theta_{\rho}$  of Theorem 5.185. The trace substitution  $J$  provides the scalar bracket sector; it leaves these cotangent homotopies as the obstruction of Problem 5.48.  $\square$

#### 4.2 The two algebras

The trace substitution  $J$  produces the open algebra. The Dirac brane stack is the Dirac-reduced first-order action (7); its ghost-zero invariant ring is  $H^0(\mathcal{R}_N^{GL_N}, Q)$ . By Lemma 3.1 the stable connected  $GL_N$ -invariants are exactly the trace family  $\{J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)} : f \in \tilde{A}\}$ , with kernel the constants; any polynomial  $GL_N$ -invariant Darboux-natural scalar-reduced map agreeing with the moment-map action on linear generators must equal  $J$  (Theorem 5.146). On a brane interval  $I$  the smeared trace observables generate the open-string sector classical observables factorization algebra

$$\mathcal{A}_N(I) = \widehat{\mathbf{S}}(\Omega_c^\circ(I) \widehat{\otimes} \tilde{A}), \quad B_f(a) = \int_I a(t) J(f)(t) dt,$$

with smearing test function  $a \in \Omega_c^\circ(I)$ , boundary observable  $B_f(a)$ , and bracket  $\{B_f(a), B_g(b)\} = B_{\{f, g\}}(ab)$  (Remark 4.10); the support-locality variant  $\mathcal{A}_\partial^{\text{Ham}}(I)$  on the full de Rham complex appears in §1.6.

Hamiltonian ghosts move; cotangent coordinates measure. The closed BV field content is therefore (4),  $\mathfrak{g} = \widehat{\mathfrak{h}} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1]$ , and the closed observables are *by definition* the continuous CE cochains

$$\mathcal{B} = C_{\text{CE,cont}}^\bullet(\mathfrak{g}).$$

On a holomorphic polydisk  $B \subset \mathbb{C}_{\text{hol}}^2$  and a topological open  $U \subset \mathbb{R}_{\text{top}}^2$ , the constructed holomorphic and mixed factorization avatars are

$$\mathcal{F}_{\text{Ham}}^{\text{hol}}(B) = C_{\text{CE,cont}}^\bullet(\Omega_c^{\circ, \bullet}(B) \widehat{\otimes} \mathfrak{g}), \quad \mathcal{F}_{\text{Ham}}^{\text{mix}}(U \times B) = C_{\text{CE,cont}}^\bullet(\Omega_c^\bullet(U) \widehat{\otimes} \Omega_c^{\circ, \bullet}(B) \widehat{\otimes} \mathfrak{g}),$$

of Definition 1.13.

The comparison between  $\mathcal{A}_N$  and  $\mathcal{B}$  is the cochain-level CE/PV identity on generators

$$c^f \mapsto \theta^f, \quad u_f \mapsto O_f,$$

of Lemma 4.1: closed Hamiltonians act through  $c$ , boundary observables are read off  $u$ . The cotangent module is the Grothendieck–Matlis principal-part  $\mathcal{P}$ , and the Moyal coadjoint action acts on the residue side as a deformation-level structure distinct from the polynomial PBW class (Theorem 8.1).

The recognition criterion has five levels (Theorem 5.5). Each level adjoins one extra structure to the level beneath; obstruction at level  $k$  lives in a single obstruction class.



| Level | Comparison                                                                                                                                                                                                 | Hypothesis                                                                                                                                |
|-------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------|
| 1     | $C_{\text{CE}}^\bullet(T^*[1]I) \cong \text{PV}(\mathfrak{sl})$ on a finite-dim Lie algebra $I$ (Theorem 5.2)                                                                                              | characteristic zero                                                                                                                       |
| 2     | $\Phi_{\text{coord}}: C_{\text{CE,coord}}^\bullet(\mathfrak{g}) \xrightarrow{\sim} \text{PV}_{\text{coord}}(\mathcal{A}_{\partial,\text{coord}})$ on the formal-polydisk coordinate envelope (Theorem 5.3) | coordinatewise finiteness                                                                                                                 |
| 3     | $P_\circ$ -bracket lift on a sub-PV model $B \subset \text{PV}_{\text{coord}}$                                                                                                                             | $B$ bracket-admissible (Definition 4.18)                                                                                                  |
| 4     | Maurer–Cartan diagonal kernel $\Theta_B \in \mathcal{K}_B$                                                                                                                                                 | $(B, \mathcal{K}_B)$ kernel-admissible; diagonal converges                                                                                |
| 5     | Bar-cobar/enveloping $(C_{\text{CE,cont}}^\bullet(\mathfrak{g}_{\text{adm}}))^\dagger \simeq \widehat{U}(\mathfrak{g}_{\text{adm}})$ and matched $A_\infty$ -Koszul on $\mathcal{A}_N$ -side               | admissible<br>Tate/nilpotent/relative/weighted replacement; conilpotent endpoint; $\varprojlim^1$ -vanishing; open-trace acceptance datum |

Level 1 is independent. Level 2 is the formal-polydisk lift of level 1 by coefficientwise finite tests. Level 3 chooses a topologised subalgebra in which the Schouten bracket converges; the strict ambient product completion has the  $P_\circ$ -bracket obstruction. Level 4 chooses a kernel-admissible target in which the diagonal Casimir converges to a Maurer–Cartan element. Level 5 adds an admissible bar-cobar replacement on the closed Lie side and a matched  $A_\infty$ -Koszul acceptance datum on the open trace side; the five obstructions of clause (4) of Theorem 5.5 block the strict ambient product from satisfying it. The minimal RG-stable analytic completion forced by  $J$  — closed under the Schouten bracket and pointwise product, with heat-kernel propagator extension and Casimir convergence — is the weighted-Tate completion  $B_w$  (Theorem 5.57); it satisfies levels 3 and 4, with level 5 conditional on the open-trace  $A_\infty$ -Koszul acceptance datum.

The brane-line factorization-current comparison (Theorem 4.11) is the level-3 application: it identifies the reduced Hamiltonian sector of  $\mathcal{F}_{\text{Ham}}^{\text{mix}}|_L$  with  $Z_{\text{red,Ham}}^{P_\circ}(\mathcal{A}_N)$  on each interval.

Identification of the raw finite- $N$  operator algebras with the strict formal Hamiltonian envelope requires an open-trace  $A_\infty$ -Koszul acceptance datum  $(A_{\text{op}}, C_{\text{op}}, B_{\text{CE/PV}}, \tau_{\text{op}})$  with exact obstruction complex inside an admissible Tate/stable category, conilpotent endpoint, and  $\varprojlim^1$ -vanishing (Theorem 5.5, item (3)). This datum is exactly what upgrades the cochain CE/PV identity to bar-cobar/enveloping conclusions and large- $N$  operator-algebra identifications. The unreduced cotangent-included lift to a chain morphism of factorization algebras

$$\Phi_{\text{cl}}: \text{Obs}_{\text{closed}}^{\text{cl}}|_L \longrightarrow Z_{\text{fact}}^{P_\circ}(\text{Obs}_{\text{open},N}^{\text{cl}})$$

is the obstruction problem of Problem 5.48. The five named classes — the strict Casimir support obstruction  $C_{\mathfrak{h}}$ , the linear translation term  $\{z_1, z_2^{N+1}\} = (N+1)z_2^N$ , the  $H_2 \cong \mathfrak{sl}_2$  lower-central obstruction to conilpotence, the associated-graded cone  $H^\bullet(\text{Cone}(\text{gr } \kappa_\tau))$ , and the Milnor  $\varprojlim^1$  defect (Theorem E.53; Theorem 5.5, item (4)) — are each independently necessary. Removing any one collapses the comparison; a chain morphism requires all five. These classes block the unreduced cotangent centrality homotopies of Theorem 4.4.

Costello’s protected sector of perturbative twisted M-theory on  $\mathbb{R} \times X$ , with  $X$  a holomorphic-symplectic surface and a stack of  $N$  topological branes on  $\mathbb{R} \times p$ , supplies the physical motivation for

a Hamiltonian formal sector at the symplectic polydisk  $\widehat{X}_p$ . Once an ambient background supplies an admissible protected Hamiltonian restriction, the local theorem of §5.3 applies to that restriction.

#### 4.3 Reduced Hamiltonian brane coupling

The brane is a constrained  $N$ -Chan–Paton chart of  $(\widehat{\mathbb{C}}^2_o, \omega)$  with normal coordinates  $\phi_1, \phi_2 \in \mathfrak{gl}_N$  and equal-time Poisson bracket  $\{(\phi_1)^i_j(t), (\phi_2)^k_l(t')\} = \delta_l^i \delta_j^k \delta(t - t')$  on two brane-time points  $(t, t')$  (Lemma 3.15, Remark 4.10). A closed Hamiltonian  $f \in \tilde{A}$  acts on the brane by Dirac substitution  $z_i \rightsquigarrow \phi_i$  followed by trace and stable scalar reduction:

$$J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)}.$$

Smearing along the brane line  $L = \mathbb{R}_{\text{brane}} \times \{o\} \times \{p\}$  by a compactly supported test density  $a(t) dt$  gives the boundary observable

$$B_f(a) = \int_L a(t) \overline{\text{Tr } f(\phi_1(t), \phi_2(t))} dt \in A_\partial^{\text{Ham}}(I). \quad (13)$$

The bulk Hamiltonian bracket equals the brane Poisson bracket of these observables (Theorem 4.5); the moment-map operation comes from the Dirac equal-time bracket of Lemma 3.15 together with Remark 4.10.

The matrix-Chan–Paton BV theory of §§3.3–3.4 supplies the brane-line trace observables; the closed shifted-cotangent BV theory of §2.1 supplies the Hamiltonian generators  $f \in \tilde{A}$ ; the smeared substitution  $B_f(a)$  of (13) matches their brackets at the level of the reduced  $P_o$ -bracket (Theorem 4.5, Theorem 4.11). The all-window strict cotangent extension of this matched bracket — the unreduced Tate-coefficient cotangent lift — is the obstruction calculus of §7.5.

**THEOREM 4.5** (*Brane trace-map principle: bulk Hamiltonians as moment-map currents*). For  $f, g \in \tilde{A}$  and  $a, b \in \Omega_c^\circ(\mathbb{R})$ , the boundary observables (13) satisfy the strict  $P_o$ -bracket identity

$$\{B_f(a), B_g(b)\}_{A_\partial^{\text{Ham}}} = B_{\{f, g\}_{\tilde{A}}}(ab), \quad (14)$$

and the bracket vanishes whenever the supports of  $a$  and  $b$  are disjoint. The ghost-zero specialisation of (14) is the unsmeared closed-side Lie homomorphism  $\{J(f), J(g)\}_\partial = J(\{f, g\}_{\tilde{A}})$  of Theorem 5.190. Before scalar reduction the same bracket carries the  $U(1)$  centre anomaly of Corollary 5.191: the Capelli/Moyal shift  $N \cdot \omega(f, g) \int_{\mathbb{R}} a(t)b(t) dt$  obstructs the Chan–Paton-line lift.

*Proof.* Smearing the equal-time Dirac bracket  $\{(\phi_1)^i_j(t), (\phi_2)^k_l(t')\} = \delta_l^i \delta_j^k \delta(t - t')$  against  $a(t)b(t')$  turns  $\delta(t - t')$  into  $a(t)b(t)$ . The matrix symplectic form contracts the Hamiltonian derivatives of  $f$  and  $g$  to the holomorphic Poisson bracket  $\{f, g\} = \partial_{z_1} f \partial_{z_2} g - \partial_{z_2} f \partial_{z_1} g$  evaluated at  $(\phi_1, \phi_2)$ . Tracing and projecting to the stable scalar-reduced quotient gives (14); disjoint-support vanishing follows because the smeared product  $ab$  is pointwise zero. This is the level-one component of Theorem 4.11. The center anomaly is the unreduced refinement Corollary 5.191 (ii); after scalar reduction it is killed and (14) survives.  $\square$

The shifted-cotangent partner  $\beta$  on the brane is the principal-part defect current of Theorem 5.185. Together with Theorem 4.5 the trace map  $J$  and the principal-part defect current  $\rho \mapsto \Theta_\rho$  form a cosheaf source datum on intervals  $I \subset \mathbb{R}$ .

*Construction 4.6 (Reduced line-defect current kernel).* Let

$$X = \mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2, \quad L = \mathbb{R}_{x_1} \times \{x_2 = 0\} \times \{p\} \subset X,$$

and let  $i : L \hookrightarrow X$  be the inclusion. This is the line defect supporting the brane currents. In the reduced formal local model, a compactly supported test density  $a(t) dt$  on  $L$  defines the defect-supported current  $i_*(a(t) dt)$ , locally written as  $a(x_1) \delta(x_2) \delta_p dx_1 dx_2$  together with the formal holomorphic delta-current at  $p$ .

The degree-zero Hamiltonian part of the reduced PBW/Rees defect-current vertex is the smeared boundary current (13):

$$K_L^{\text{cl}}(a \otimes f) = B_f(a), \quad f \in \tilde{A}.$$

For the cotangent principal-part source, the reduced vertex sends

$$B \otimes \rho[1] \mapsto \Theta_\rho(B), \quad B \in \mathcal{D}'_c(L), \quad \rho \in \mathcal{P},$$

using Theorem 5.185. These formulas define a reduced PBW/Rees defect-current kernel. A cochain-level map from closed CE observables, and the quantum Costello brane-defect kernel, are separate construction targets: they must supply the Ward identity, anomaly cancellation, finite-scale propagator, and quantum moment-map compatibility.

*Construction 4.7 (Interval Hamiltonian and factorization algebras).* For an interval  $I \subset \mathbb{R}$ , define

$$\begin{aligned} \mathfrak{h}_I &= \Omega_c^\bullet(I) \widehat{\otimes} \tilde{A}, \\ \mathcal{P}_I &= \mathcal{D}'_c(I) \widehat{\otimes} \mathcal{P}, \\ \mathfrak{g}_I &= \mathfrak{h}_I \ltimes \mathcal{P}_I[1], \end{aligned}$$

with brackets

$$\begin{aligned} [a \otimes f, b \otimes g]_{\mathfrak{h}_I} &= ab \otimes \{f, g\}_{\tilde{A}}, \\ [a \otimes f, B \otimes \rho[1]]_{\mathfrak{g}_I} &= aB \otimes (f \cdot \rho)[1], \\ [B \otimes \rho[1], C \otimes \sigma[1]]_{\mathfrak{g}_I} &= 0, \end{aligned}$$

where products in  $\Omega_c^\bullet(I)$  are de Rham products and, in the displayed degree-zero formulas,  $aB$  is multiplication of a compactly supported current by a smooth compactly supported function and  $f \cdot \rho = \pi_{\text{pp}}\{f, \rho\}$  is the principal-part coadjoint action of 5.175. Inclusions  $I' \subset I$  act by extension by zero on compactly supported de Rham forms and currents; disjoint unions act by sum. This makes  $I \mapsto \mathfrak{h}_I, \mathfrak{g}_I$  cosheaves of (graded) Lie algebras on  $\mathbb{R}$ . For  $j : I \hookrightarrow J$ , the map  $j_!$  is defined on  $\Omega_c^\bullet(I)$  because supports are compactly contained in  $I$ ; equivalently, smooth representatives vanish near the boundary of  $I$  inside  $J$ . It satisfies

$$j_!(\alpha \wedge \beta) = j_!\alpha \wedge j_!\beta, \quad j_!(aB) = (j_!a)(j_!B),$$

in the degree-zero current notation. On CE cochains the same extension-by-zero maps induce pull-back/coproduct maps; the CE cochain source is contravariant in this convention.

Define the boundary  $P_0$  factorization algebras on  $\mathbb{R}$  in the formal stable Hamiltonian sector by

$$\begin{aligned} A_\partial^{\text{Ham}}(I) &= \widehat{\mathbf{S}}(\mathfrak{h}_I), \\ A_\partial^{\text{PP}}(I) &= \widehat{\mathbf{S}}(\mathfrak{h}_I) \widehat{\otimes} \widehat{\mathbf{S}}(\mathcal{P}_I[1]), \end{aligned}$$

with degree-zero reduced Hamiltonian  $P_0$  bracket extended as a biderivation from

$$\begin{aligned} \{a \otimes f, b \otimes g\}_{A_\partial} &= ab \otimes \{f, g\}_{\bar{A}}, \\ \{a \otimes f, B \otimes \rho[1]\}_{A_\partial} &= aB \otimes (f \cdot \rho)[1], \\ \{B \otimes \rho[1], C \otimes \sigma[1]\}_{A_\partial} &= 0. \end{aligned}$$

The factorization product on disjoint intervals is the multiplication in  $\widehat{\mathbf{S}}$  after extension by zero. These are the reduced targets after de Rham-current contraction for the reduced Hamiltonian current central-operation map. The Hamiltonian summand  $A_\partial^{\text{Ham}}$  is the connected component of the formal Hamiltonian sector of the open brane factorization algebra; the principal-part summand  $A_\partial^{\text{pp}}/A_\partial^{\text{Ham}}$  is the reduced current object of §.185.

LEMMA 4.8 (*Compact-support interval contraction*). Let  $I \subset \mathbb{R}$  be a nonempty open interval, oriented by the coordinate  $t$ . Choose a compactly supported one-form  $\lambda_I(t) dt \in \Omega_c^1(I)$  with  $\int_I \lambda_I(t) dt = 1$ . Define

$$p_I : \Omega_c^\bullet(I) \longrightarrow \mathbb{C}[-1], \quad p_I(\alpha) = \int_I \alpha,$$

with  $p_I$  zero on degree 0, and

$$i_I : \mathbb{C}[-1] \longrightarrow \Omega_c^\bullet(I), \quad i_I(1) = \lambda_I(t) dt.$$

Extend compactly supported forms by zero to  $\mathbb{R}$ . For a one-form  $\alpha = f(t) dt$ , set

$$K_I(\alpha)(t) = \int_{-\infty}^t \left( f(s) - \lambda_I(s) \int_I f(r) dr \right) ds, \quad K_I(\varphi) = 0 \quad (\varphi \in \Omega_c^0(I)).$$

Then  $K_I : \Omega_c^1(I) \rightarrow \Omega_c^0(I)$  is well-defined and compactly supported, and

$$dK_I + K_Id = \text{id}_{\Omega_c^\bullet(I)} - i_I p_I.$$

Consequently  $\Omega_c^\bullet(I)$  contracts onto its compact-support cohomology line  $\mathbb{C}[-1]$ . Acyclicity of  $\Omega_c^\bullet(I)$  is replaced by the following exact statement: a compactly supported primitive of a compactly supported one-form exists precisely on the kernel of the integral map, and the integral class is the surviving compact-support cohomology generator.

*Proof.* The one-form inside the integral defining  $K_I(\alpha)$  has compact support and total integral zero. Its primitive therefore vanishes near both ends of  $I$ , after extension by zero to  $\mathbb{R}$ , so  $K_I(\alpha) \in \Omega_c^0(I)$ .

For  $\alpha = f(t) dt$ ,

$$dK_I(\alpha) = \left( f(t) - \lambda_I(t) \int_I f(r) dr \right) dt = \alpha - i_I p_I(\alpha).$$

For a compactly supported function  $\varphi$ ,

$$K_I(d\varphi)(t) = \int_{-\infty}^t \varphi'(s) ds = \varphi(t),$$

because  $\varphi$  vanishes near the left end, and  $p_I(\varphi) = 0$ . These two identities are exactly  $dK_I + K_Id = \text{id} - i_I p_I$  in degrees 1 and 0.  $\square$

PROPOSITION 4.9 (*Locality of the brane  $P_o$  bracket*). Let  $a \in \Omega_c^\circ(I_1), b \in \Omega_c^\circ(I_2)$  with  $I_1 \cap I_2 = \emptyset$ . For all  $f, g \in \tilde{A}$ ,

$$\{B_f(a), B_g(b)\}_{A_\partial^{\text{Ham}}} = 0.$$

In particular,  $A_\partial^{\text{Ham}}$  has the strict factorization product property: the bracket between sections supported on disjoint intervals vanishes.

*Proof.* By 4.7,  $\{B_f(a), B_g(b)\} = B_{\{f,g\}}(ab)$ . The pointwise product  $ab \equiv 0$  since the supports are disjoint, so  $B_{\{f,g\}}(ab) = 0$ .  $\square$

REMARK 4.10 (*Locality from the canonical equal-time bracket*). The locality input is the canonical equal-time bracket of the first-order open BV theory, written on two brane-time points  $(t, t')$  (with  $s$  reserved throughout for the transverse  $\mathbb{R}_s$  coordinate of  $X = \mathbb{R}_t \times \mathbb{R}_s \times \mathbb{C}^2$ ):

$$\{(\phi_1)^i_j(t), (\phi_2)^k_l(t')\} = \delta_l^i \delta_j^k \delta(t - t'), \quad \{(\phi_a)^i_j(t), (\phi_a)^k_l(t')\} = 0 \quad (a = 1, 2).$$

Tracing  $f(\phi_1, \phi_2)(t)$  and  $g(\phi_1, \phi_2)(t')$ , applying the Leibniz rule, and integrating against  $a(t)b(t')\delta(t - t')$  produces exactly the smeared identity  $\{B_f(a), B_g(b)\} = B_{\{f,g\}}(ab)$  of 5.191 in unsmeared form. In the classical reduced, de Rham-contracted sector this delta-function locality produces a strict support-local  $P_o$  factorization product. The unreduced BV current complex still requires its own homotopies, contact-term choices, and QME data.

THEOREM 4.11 (*Interval-wise reduced Hamiltonian CE/PV central-operation map, Hamiltonian sector*). Let  $\text{Obs}_{\text{open}, N}^{\text{cl}, \text{red}}$  be the reduced open-brane factorization algebra of 4.14. Let  $\text{Obs}_{\text{closed}, H, \text{Ham}}^{\text{cl}}|_L$  denote the formal classical Hamiltonian-mode subfactorization algebra of the cotangent BF theory on the formal neighbourhood of  $L$  obtained by restricting to the connected Hamiltonian central-operation sector. Define the interval-wise CE cochain cofactorization object, equivalently a factorization object on the opposite interval category, by

$$\mathcal{F}_{\text{cl}}(I) = C_{\text{CE}, \text{cont}}^\bullet(\mathfrak{g}_I), \quad \mathfrak{g}_I = \mathfrak{h}_I \ltimes \mathcal{P}_I[1],$$

where extension by zero on  $\Omega_c^\bullet$  and on compactly supported currents induces the CE coproduct. Then:

- (i) there is a continuous interval-wise morphism of  $P_o$  cofactorization source data to the reduced Hamiltonian central-operation target

$$\Phi_{\text{fact}}^{\text{Ham}} : \text{Obs}_{\text{closed}, H, \text{Ham}}^{\text{cl}}|_L \longrightarrow Z_{\text{red}, \text{Ham}}^{P_o}(A_\partial^{\text{Ham}}),$$

whose underlying CE/PV map on each interval  $I$  is given on length-one coordinates by

$$B_f(a) := \Phi_I(u_{a(t)dt \otimes f}) = \int_I a(t) \overline{\text{Tr} f(\phi_1(t), \phi_2(t))} dt,$$

and

$$\Phi_I(c_{B \otimes \rho}) = \theta_{B \otimes \rho}.$$

The reduced principal-part current representative  $\Theta_\rho(B)$  belongs to the boundary local-dual model of Theorem 5.185; its comparison with the CE/PV vector-field coordinate is made after that reduction.

- (ii) this morphism is an isomorphism of  $P_0$ -cdg algebras in each fixed interval, after the stated reduced current model is chosen, and is compatible with the CE coproduct on the source and disjoint factorization products on the target;
- (iii) after applying the compact-support de Rham contraction  $p_I : \Omega_c^\bullet(I) \rightarrow \mathbb{C}[-1]$ , the induced Hamiltonian density sector is locally constant in the topological direction  $\mathbb{R}$  in the sense of Lurie, *Higher Algebra*, §5.5; Costello–Gwilliam [15, Vol. I §6.4] supplies the published factorization-algebra terminology. Extension by zero  $\mathcal{D}'_c(I') \rightarrow \mathcal{D}'_c(I)$  for arbitrary compactly supported distributional currents requires separate analysis;
- (iv) the factorization product  $A_\partial^{\text{Ham}}(I_1) \otimes A_\partial^{\text{Ham}}(I_2) \rightarrow A_\partial^{\text{Ham}}(V)$  for disjoint  $I_1 \sqcup I_2 \subset V$  corresponds under  $\Phi_{\text{fact}}^{\text{Ham}}$  to the CE coproduct on the closed side induced by extension by zero  $\Omega_c^\bullet(I_j) \hookrightarrow \Omega_c^\bullet(V)$ ;
- (v) at the stalk  $I \rightarrow \{t_0\}$ ,  $\Phi_{\text{fact}}^{\text{Ham}}$  restricts to the stalkwise central operation

$$f \mapsto L_{B_f} \in C_{\text{Hoch}}^\circ(A_N^{\text{pt}}, A_N^{\text{pt}})$$

of 3.39.

*Proof. Step 1.* Identify the interval-wise reduced  $P_0$  model. In the formal stable Hamiltonian sector, the boundary factorization algebra  $A_\partial^{\text{Ham}}(I) = \widehat{\mathbf{S}}(\mathfrak{h}_I)$  is graded-commutative, with strict  $P_0$  bracket inherited from  $\mathfrak{h}_I$  as a biderivation. Here  $Z_{\text{red,Ham}}^{P_0}$  means the reduced interval-wise Poisson-CE model  $\text{PV}^\bullet(\widehat{\mathbf{S}}(\mathfrak{h}_I)) = \widehat{\mathbf{S}}(\mathfrak{h}_I) \widehat{\otimes} \widehat{\mathbf{S}}(\mathfrak{h}_{I,\text{cont}}^\vee[-1])$  with the Schouten bracket. A genuine Hochschild/factorization-centre identification for the completed Tate algebra would require a continuous HKR theorem with interval functoriality; the proof here uses the explicit PV model. Restricting to the polynomial subalgebra  $\mathbf{S}(\mathfrak{h}_I) \subset \widehat{\mathbf{S}}(\mathfrak{h}_I)$  and to the constant polyvectors gives the central multiplication operations, whose boundary names are the Hamiltonian currents  $B_f(a)$ . The CE coordinates  $c$  dual to Hamiltonian generators map instead to constant polyvectors. This is the interval version of the generator dictionary  $c^I \mapsto \theta^I, u_I \mapsto O_I$  in 4.20.

*Step 2.* Construct the CE source. By 5.165, applied to  $\mathfrak{g}_I = \mathfrak{h}_I \ltimes \mathcal{P}_I[1]$  in the stated completed category,

$$C_{\text{CE,cont}}^\bullet(\mathfrak{g}_I) = \widehat{\mathbf{S}}(\mathfrak{g}_{I,\text{cont}}^\vee[-1]),$$

and its CE differential is the dual of the semidirect Lie bracket. The Hamiltonian part is controlled by the de Rham-current pairing on the  $\Omega_c^\bullet(I)$  factor and the residue-Taylor pairing on  $\tilde{A}$ . In degree-zero notation this is the familiar pairing between compactly supported test functions and compactly supported currents; the polynomial subspace  $\tilde{A} \subset \widehat{A}$  pairs with the discrete Taylor dual generated by the  $\partial_{a,b}$ . The shifted-cotangent CE coordinate represented by  $a(t)dt \otimes f$  is dual to the cotangent generator determined by that current. Its boundary image is  $B_f(a)$ .

*Step 3.* Define  $\Phi_{\text{fact}}^{\text{Ham}}(I)$  on generators. By 6.20, the closed Hamiltonian cotangent BF observables on the formal neighbourhood of  $L = \mathbb{R}_{\text{brane}} \times \{0\} \times \{p\}$  in the Hamiltonian sector are  $\widehat{\mathbf{S}}(\mathfrak{h}_{I,\text{cont}}^\vee[-1])$ , with CE differential dual to the Hamiltonian Lie bracket on  $\mathfrak{g}_I$ . The interval-wise map sends a length-one shifted-cotangent CE coordinate represented by  $a(t)dt \otimes f$  to the element of the central operations factorization algebra given by the multiplication operator

$$L_{B_f(a)} \in \text{Hoch}^\circ(A_\partial^{\text{Ham}}(I)) \subset Z_{\text{red,Ham}}^{P_0}(A_\partial^{\text{Ham}}(I)),$$

with  $B_f(a) = \Phi_I(u_{a(t)dt \otimes f})$  as displayed. By 3.39, this operator is closed (Hochschild o-cocycle on the graded-commutative open stalk). Extend  $\Phi_{\text{fact}}^{\text{Ham}}(I)$  as a continuous algebra morphism on  $\widehat{\mathbf{S}}$ .

*Step 4.* Verify CE differential compatibility. On length two, the CE differential of a shifted-cotangent generator represented by  $a(t)dt \otimes f$  is the coadjoint term dual to the Lie bracket on  $\mathfrak{h}_I$ . By 5.191 (its sharpened form below),

$$\{L_{B_f(a)}, L_{B_g(b)}\}_{A_\partial^{\text{Ham}}} = L_{B_{\{f,g\}}(ab)}.$$

Pulling back through  $\Phi_{\text{fact}}^{\text{Ham}}$  matches this with the shifted-cotangent CE differential dual to  $[a \otimes f, b \otimes g]_{\mathfrak{h}_I} = ab \otimes \{f, g\}$ . Inductive extension by length and the biderivation property propagates this to all CE lengths.

*Step 5.* Quasi-isomorphism on each interval. Both sides are free graded-commutative algebras on the same chosen compact-support current reduced Tate vector space after the generator dictionary  $c \mapsto \theta$ ,  $u \mapsto O$ , with the same CE differential induced from the semidirect Lie bracket of  $\mathfrak{g}_I$  (closed side: by definition; brane side: by the reduced interval-wise PV model of Step 1 and the Schouten bracket). The map respects the symmetric-algebra generators by construction. Thus it is an isomorphism of underlying graded vector spaces and intertwines the differentials. The map is therefore an isomorphism of  $P_\circ$ -cdg algebras on each interval.

*Step 6.* Factorization product compatibility. For disjoint  $I_1, I_2 \subset V$ , the CE source extension by zero gives  $\mathfrak{h}_{I_1} \oplus \mathfrak{h}_{I_2} \hookrightarrow \mathfrak{h}_V$ . Thus the CE source is cofactorization: if  $j_{I_k, V, !} : \mathfrak{g}_{I_k} \rightarrow \mathfrak{g}_V$  denotes extension by zero, the source structure map is

$$\Delta_{I_1, I_2, V} : C_{\text{CE}}^\bullet(\mathfrak{g}_V) \rightarrow C_{\text{CE}}^\bullet(\mathfrak{g}_{I_1}) \widehat{\otimes} C_{\text{CE}}^\bullet(\mathfrak{g}_{I_2}).$$

On the brane side,  $A_\partial^{\text{Ham}}(I_1) \otimes A_\partial^{\text{Ham}}(I_2) \rightarrow A_\partial^{\text{Ham}}(V)$  is the multiplication map under the inclusion of  $\mathfrak{h}_{I_1} \oplus \mathfrak{h}_{I_2} \hookrightarrow \mathfrak{h}_V$ . By 4.9, the bracket between the two factors vanishes on disjoint supports, so the product is strictly multiplicative. The induced map between central-operation factorization algebras is the corresponding tensor product of multiplication operators. The compatibility is the mixed-variance equation

$$\Phi_V = m_{I_1, I_2, V} \circ (\Phi_{I_1} \otimes \Phi_{I_2}) \circ \Delta_{I_1, I_2, V},$$

for the operations whose support lies in  $I_1 \sqcup I_2 \subset V$ . Equivalently, the diagram

$$\begin{array}{ccc} \mathcal{F}_{\text{cl}}(I_1) \otimes \mathcal{F}_{\text{cl}}(I_2) & \xrightarrow{\Phi(I_1) \otimes \Phi(I_2)} & Z_{\text{red, Ham}}^{P_\circ}(A_\partial^{\text{Ham}})(I_1) \otimes Z_{\text{red, Ham}}^{P_\circ}(A_\partial^{\text{Ham}})(I_2) \\ \uparrow & & \downarrow \\ \mathcal{F}_{\text{cl}}(V) & \xrightarrow{\Phi(V)} & Z_{\text{red, Ham}}^{P_\circ}(A_\partial^{\text{Ham}})(V) \end{array}$$

commutes after vertical reversal of the left arrow (CE coproduct = dual of extension by zero).

*Step 7.* Locally constant structure. For each interval  $I$ , Lemma 4.8 gives the explicit contraction

$$\Omega_c^\bullet(I) \xrightleftharpoons[i_I]{p_I} \mathbb{C}[-1], \quad i_I(1) = \lambda_I(t) dt, \quad p_I(\alpha) = \int_I \alpha,$$

with  $dK_I + K_I d = \text{id} - i_I p_I$ . The maps  $p_I$  are natural for extension by zero; the choice of  $i_I$  and  $K_I$  is a choice of density representative for the compact-support cohomology generator. The degree-zero boundary notation  $B_f(a)$  used above already includes the shifted-cotangent degree, so the density representative appears as a degree-zero CE coordinate after the BV shift. Translation is homotopically trivial by the same de Rham contraction, hence the interval-wise objects are locally constant in the topological direction after this density contraction.

*Step 8. Stalk recovery.* Under the density contraction, a length-one shifted-cotangent CE coordinate represented by  $a(t)dt \otimes f$  maps to the boundary current  $B_f(a)$  and hence to the multiplication operator  $L_{B_f(a)}$ . Stalk recovery is obtained by any cofinal family of compactly supported unit densities supported near  $t_0$ . This represents the compact-support cohomology generator under the locally constant factorization-algebra quasi-isomorphism; degree-zero point evaluation is replaced by compact-support density contraction.  $\square$

*Remark 4.12 (Locally constant equivalence and  $E_1$ -algebra translation).* In the locally constant setting on  $\mathbb{R}$ , factorization algebras are equivalent to homotopy associative algebras up to translation invariance by Lurie, *Higher Algebra*, §5.5.4, in the Costello–Gwilliam vocabulary of locally constant factorization algebras [15, Vol. I §6.4]. The Hochschild cochain complex of the  $E_1$  algebra computes the factorization-center stalk. The reduced Hamiltonian theorem here uses the explicit PV model  $\widehat{\mathbf{S}}(\mathfrak{h}_I) \widehat{\otimes} \widehat{\mathbf{S}}(\mathfrak{h}_I^\vee[-1])$  and its strict  $P_0$  subalgebra of multiplication operators. Identifying this PV model with continuous Hochschild cochains for the completed Tate algebra is a separate HKR/topology theorem. The CE source identification of 4.11 is exactly the closed dual of this PV picture in the connected Hamiltonian sector.

*Remark 4.13 (Presentations of one-dimensional factorization algebras).* The precise equivalence used here is Lurie’s identification of locally constant factorization algebras on the line with  $E_1$ -algebras [41, Thm. A.3.7.5]. The Costello–Gwilliam model-categorical language is used only as a compatible presentation convention unless the cited corollary is separately anchored.

For a factorization algebra  $\mathcal{A}$  on the line, write  $Z_{\text{fact}}(\mathcal{A})$  for the factorization algebra of  $E_1$  central operations; in the locally constant one-dimensional case its stalk is the Hochschild cochain complex of the corresponding  $E_1$  algebra. The notation  $Z_{\text{fact}}^{P_0}(\mathcal{A})$  means the same central operations equipped with classical Poisson/BV compatibility homotopies. The reduced Hamiltonian proof uses the explicit local central-operation model.

**THEOREM 4.14** (*Hamiltonian-current central operation, smeared*). 4.11 restricted to the length-one CE component on each interval recovers the degree-zero Hamiltonian current: the assignment

$$\Phi_{I,N}(u_{a(t)dt \otimes f})(u) = B_f(a) \cdot u = \left( \int_I a(t) \overline{\text{Tr } f(\phi_1(t), \phi_2(t))} dt \right) u$$

is a closed Hochschild o-cochain, hence a degree-zero central operation, in  $Z_{\text{fact}}(\text{Obs}_{\text{open},N}^{\text{cl,red}})(I)$ , the assignments are compatible with interval inclusions and disjoint factorization products, and the induced operations satisfy

$$\{\Phi(u_{a(t)dt \otimes f}), \Phi(u_{b(t)dt \otimes g})\}_{A_\partial^{\text{Ham}}} = \Phi(u_{a(t)b(t)dt \otimes \{f,g\}_A}).$$

The degree-zero current is the length-one component of the reduced CE/PV central-operation comparison: its content is the strict  $P_0$  identity, one of the structure equations defining  $\Phi_{\text{fact}}^{\text{Ham}}$  of 4.11.

*Proof.* In the reduced open model  $Q\phi_1 = Q\phi_2 = 0$ . Hence  $\overline{\text{Tr } f(\phi_1, \phi_2)}$  is  $Q$ -closed for every Hamiltonian  $f$ , and the compactly supported integral is  $Q$ -closed. Since the reduced open observable algebra is graded-commutative, multiplication by this closed element is a Hochschild o-cocycle; this is the interval-wise version of 3.39. Extension by zero of the compactly supported test density is compatible with enlargement to a larger interval through the extension map  $j_!$ . For disjoint intervals 4.9 gives vanishing of



the  $P_o$  bracket between the two factors, so the factorization product of two such operations is the strict multiplication by  $I_\partial(a, f)I_\partial(b, g)$ ; this is the multiplicative extension from  $\mathfrak{h}_I$  to  $A_\partial^{\text{Ham}}(I) = \widehat{\mathbf{S}}(\mathfrak{h}_I)$ ; the linear current bracket is its indecomposable part.

The Poisson statement is local on the line. For compactly supported test functions  $a, b$ , §.191 (the strict  $P_o$  identity on the brane factorization algebra) gives

$$\{I_\partial(a, f), I_\partial(b, g)\}_{A_\partial^{\text{Ham}}} = I_\partial(ab, \{f, g\}_{\bar{A}}).$$

Passing to the connected stable Hamiltonian quotient removes only the scalar Hamiltonian line, which is central for the bracket. Thus the current bracket is preserved. Taking  $a$  and  $b$  supported in a small coordinate interval and then passing to the locally constant stalk recovers §.190.  $\square$

**PROPOSITION 4.15** (*CE-source obstruction for perfect Hamiltonian Lie algebras*). Let  $\mathfrak{h}$  be a perfect Lie algebra in **TateVec** (i.e.  $[\mathfrak{h}, \mathfrak{h}] = \mathfrak{h}$ ). Then every length-one CE coordinate  $c^I$  dual to a Hamiltonian generator  $H_I \in \mathfrak{h}$  has nonzero CE differential:

$$d_{\text{CE}} c^K = -\frac{1}{2} C_{IJ}^K c^I c^J \neq 0.$$

Equivalently,  $H^1(\mathfrak{h}; \mathbb{C}) = 0$ .

This obstruction is the structural source of the cotangent direction: under 4.20, the boundary observable  $O_I = J(H_I) \in A_\partial$  is

$$O_I = \Phi_{\text{coord}}(u_I),$$

the image of the CE coordinate dual to the shifted cotangent generator  $\eta_{\partial_I} \in \mathfrak{h}^\vee[1]$ . The boundary observable lives in the cotangent direction of the CE cochain algebra. Its CE differential is  $d_{\text{CE}} u_K = C_{KJ}^L u_L c^J$ , mapping under  $\Phi_{\text{coord}}$  to the Hamiltonian vector field  $d_\pi^{\text{coord}} O_K = C_{KJ}^L O_L \theta^J$  of  $O_K$  in the Schouten algebra.

This is exactly what the bracket-admissible  $P_o$ -map produces: a function whose Schouten differential is its Hamiltonian vector field.

*Proof.*  $\mathfrak{h}$  perfect means each basis vector has nonzero projection from some bracket combination; equivalently, for every length-one Hamiltonian CE coordinate  $c^K$  there are indices  $I, J$  with  $C_{IJ}^K c^I c^J \neq 0$ . Hence every  $c^K$  has nonzero differential, and  $H^1$  vanishes. Identification  $O_I = \Phi_{\text{coord}}(u_I)$  is the content of §.39 combined with  $\Phi_{\text{coord}}(u_I) = O_I$  from 4.20. Differentials match on generators by §.167.  $\square$

**Remark 4.16.** The reduced Hamiltonian Lie algebra  $\mathfrak{h} = \widehat{A}/\mathbb{C} \cdot 1 = (z_1, z_2)\mathbb{C}[[z_1, z_2]]$  of the formal symplectic polydisk in dimension two is perfect: for every monomial one has, before quotienting constants,  $z_1^a z_2^b = \{z_1, \frac{1}{b+1} z_1^a z_2^{b+1}\}$ , so  $[\mathfrak{h}, \mathfrak{h}] = \mathfrak{h}$ . In particular  $[H_{1,0}, H_{m,n}] = n H_{m,n-1}$  and  $[H_{0,1}, H_{m,n}] = -m H_{m-1,n}$ . The hypothesis of 4.15 is met. The CE-length-one cochain  $\xi_{a,b} = c^{a,b}$  dual to  $H_{a,b} = z_1^a z_2^b$  has nonzero differential; closed CE cochains have zero linear Hamiltonian part. This is the source obstruction.

The CE/PV identification therefore proceeds at the level of the coordinate completion of the cochain algebra. Finite coordinate words carry the bracket and differential, and convergence of brackets and Casimir kernels on the ambient product is a separate admissibility input.

**Definition 4.17** (*Formal-polydisk coordinate CE/PV completion*). The coordinate CE/PV completion is the product over finite graded-commutative words in the Hamiltonian CE coordinates  $c^I$ , shifted-cotangent CE coordinates  $u_I$ , Schouten coordinates  $\theta^I$ , and boundary Hamiltonian coordinates  $O_I$ .

All differentials and brackets are read coefficientwise in finite coordinate windows and then passed to the compatible product completion.

*Definition 4.18 (Bracket-admissible CE/PV realizations).* A coordinate target  $B \subset \text{PV}_{\text{coord}}(\mathcal{A}_{\partial, \text{coord}})$  is bracket-admissible when the Schouten bracket and the Hamiltonian boundary brackets of the generators extend continuously from every finite coordinate window to  $B$ . The worked nontrivial class is the weighted-Tate coefficient pair of Definition 5.54, with bracket-admissibility established by Theorem 5.70 and the diagonal kernel-admissibility refinement by Theorem 5.66. The strict ambient product completion sits below admissibility at both levels: the continuous Schouten biderivation extends from finite words to the full product only through the weighted-Tate replacement, and the diagonal Casimir  $C_{\mathfrak{h}}$  converges only in coefficient categories with a regulator (Lemma 5.211); item (4) of Theorem 5.5 records the combined obstruction datum.

LEMMA 4.19 (*Formal-polydisk coordinate differential formulas*). In the coordinate completion,

$$d_{\text{CE}} c^K = -\frac{1}{2} C_{IJ}^K c^I c^J, \quad d_{\text{CE}} u_K = C_{KJ}^L u_L c^J,$$

and

$$d_{\pi}^{\text{coord}} \theta^K = -\frac{1}{2} C_{IJ}^K \theta^I \theta^J, \quad d_{\pi}^{\text{coord}} O_K = C_{KJ}^L O_L \theta^J.$$

THEOREM 4.20 (*Admissible reduced Hamiltonian CE/PV central-operation map, cochain level*). Let

$$\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$$

be the reduced Hamiltonian Lie algebra of the formal symplectic polydisk, equipped with the coordinate completion of Definition 4.17. Let  $\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^{\vee}[1]$  be the shifted-cotangent extension. Let  $\mathcal{A}_{\partial, \text{coord}}$  be the coordinate completed symmetric algebra on the stable scalar-reduced Hamiltonian trace labels.

Then there is a canonical isomorphism of completed dg graded-commutative coordinate algebras

$$\Phi_{\text{coord}} : C_{\text{CE}, \text{coord}}^{\bullet}(\mathfrak{g}) \xrightarrow{\sim} \text{PV}_{\text{coord}}(\mathcal{A}_{\partial, \text{coord}}),$$

determined on generators by

$$\Phi_{\text{coord}}(c^I) = \theta^I, \quad \Phi_{\text{coord}}(u_I) = O_I,$$

where  $c^I$  is the CE coordinate dual to  $H_I \in \mathfrak{h}$  and  $u_I$  is the CE coordinate dual to the shifted cotangent generator  $\eta_{\partial_I} \in \mathfrak{h}^{\vee}[1]$ .

If  $B \subset \text{PV}_{\text{coord}}(\mathcal{A}_{\partial, \text{coord}})$  is a bracket-admissible realization of the reduced Hamiltonian central-operation model, then the restriction

$$\Phi_{P_o}^B : \Phi_{\text{coord}}^{-1}(B) \xrightarrow{\sim} B$$

is the cochain-level open-closed reduced Hamiltonian central-operation isomorphism in the formal/local sector. The strict ambient product completion has a separate  $P_o$ -algebra obstruction. Tate bar-cobar and enveloping conclusions require the separate admissible conilpotent hypotheses of Lemma 5.165.

*Proof. Generators and coordinate extension.* By Definition 4.17,  $C_{\text{CE}, \text{coord}}^{\bullet}(\mathfrak{g})$  is the product over finite graded-commutative words in the letters  $u_I, c^I$ , with  $|u_I| = 0$  and  $|c^I| = 1$ . The target  $\text{PV}_{\text{coord}}(\mathcal{A}_{\partial, \text{coord}})$  is the identical coordinate product with  $O_I, \theta^I$  in place of  $u_I, c^I$ . The assignment  $c^I \mapsto \theta^I, u_I \mapsto O_I$

therefore extends uniquely to a continuous bijection of completed coordinate dg-commutative algebras. This is a coordinate envelope statement. Literal continuous CE cochains form the continuous subspace inside this product when the coefficient support is continuous for the chosen topology; the whole coordinate product is the formal coordinate envelope.

*Cochain identity on generators.* The CE differential is

$$d_{\text{CE}} c^K = -\frac{1}{2} C_{IJ}^K c^I c^J, \quad d_{\text{CE}} u_K = C_{KJ}^L u_L c^J.$$

By Lemma 4.19,

$$d_{\pi}^{\text{coord}} \theta^K = -\frac{1}{2} C_{IJ}^K \theta^I \theta^J, \quad d_{\pi}^{\text{coord}} O_K = C_{KJ}^L O_L \theta^J.$$

Therefore

$$\Phi_{\text{coord}}(d_{\text{CE}} c^K) = -\frac{1}{2} C_{IJ}^K \theta^I \theta^J = d_{\pi}^{\text{coord}} \theta^K = d_{\pi}^{\text{coord}} \Phi_{\text{coord}}(c^K),$$

$$\Phi_{\text{coord}}(d_{\text{CE}} u_K) = C_{KJ}^L O_L \theta^J = d_{\pi}^{\text{coord}} O_K = d_{\pi}^{\text{coord}} \Phi_{\text{coord}}(u_K).$$

*Cochain identity on the full coordinate algebra.* Both  $d_{\text{CE}}$  and  $d_{\pi}^{\text{coord}}$  are degree-1 derivations of the completed graded-commutative algebra structures on each side. They satisfy the graded Leibniz rule with the same Koszul sign convention.  $\Phi_{\text{coord}}$  is an algebra homomorphism, so

$$\begin{aligned} \Phi_{\text{coord}}(d_{\text{CE}}(xy)) &= \Phi_{\text{coord}}((d_{\text{CE}}x)y + (-1)^{|x|} x d_{\text{CE}}y) \\ &= \Phi_{\text{coord}}(d_{\text{CE}}x) \Phi_{\text{coord}}(y) + (-1)^{|x|} \Phi_{\text{coord}}(x) \Phi_{\text{coord}}(d_{\text{CE}}y), \end{aligned}$$

while

$$\begin{aligned} d_{\pi}^{\text{coord}} \Phi_{\text{coord}}(xy) &= d_{\pi}^{\text{coord}}(\Phi_{\text{coord}}(x) \Phi_{\text{coord}}(y)) \\ &= (d_{\pi}^{\text{coord}} \Phi_{\text{coord}}(x)) \Phi_{\text{coord}}(y) + (-1)^{|x|} \Phi_{\text{coord}}(x) d_{\pi}^{\text{coord}} \Phi_{\text{coord}}(y). \end{aligned}$$

Generator-level agreement plus Leibniz/algebra-homomorphism gives agreement on every word in  $\{c^I, u_J\}$  at fixed total length; coordinate-product completeness of both sides plus continuity of  $d_{\text{CE}}, d_{\pi}^{\text{coord}}, \Phi_{\text{coord}}$  for finite coordinate projections extends the equality to the completed coordinate algebras.

*Inverse.* The assignment  $\theta^I \mapsto c^I, O_I \mapsto u_I$  extends to a continuous algebra homomorphism in the opposite direction. By the same Leibniz argument, it intertwines  $d_{\pi}^{\text{coord}}$  and  $d_{\text{CE}}$ . The two compositions fix generators, hence fix the completed algebras. So  $\Phi_{\text{coord}}$  is an isomorphism of complete dg graded-commutative algebras.

*Cotangent/Schouten bracket compatibility.* On finite words the CE side carries the canonical  $(-1)$ -shifted Poisson structure on  $C_{\text{CE, coord}}^{\bullet}(T^*[\mathbf{1}]\mathfrak{h})$  with

$$\{c^I, u_J\}_{\text{CE}} = \delta_J^I, \quad \{c^I, c^J\}_{\text{CE}} = 0, \quad \{u_I, u_J\}_{\text{CE}} = 0.$$

These are the degree- $(-1)$  cotangent brackets on the cochain algebra of a shifted cotangent Lie algebra. The bracket compatibility used here is the displayed generator computation; it is separate from the degree-zero reduced Hamiltonian  $P_0$  bracket on boundary currents. The finite-word Schouten bracket on  $\text{PV}_{\text{coord}}(\mathcal{A}_{\partial, \text{coord}})$  satisfies

$$[\theta^I, O_J]_S = \delta_J^I, \quad [\theta^I, \theta^J]_S = 0, \quad [O_I, O_J]_S = 0,$$

the last because  $O_I, O_J$  are functions on the formal affine space  $\mathfrak{h}^\vee$  and the Schouten bracket of two functions is zero. Generator-level matching gives  $\Phi_{\text{coord}}(\{c^I, u_J\}_{\text{CE}}) = \Phi_{\text{coord}}(\delta_J^I) = \delta_J^I = [\theta^I, O_J]_S = [\Phi_{\text{coord}}(c^I), \Phi_{\text{coord}}(u_J)]_S$ , and similarly for the other two pairs. This proves bracket compatibility on finite words. On a bracket-admissible  $B$ , continuity and the biderivation rule extend the equality to  $\Phi_{\text{coord}}^{-1}(B)$  and  $B$ . Bracket statements on the strict ambient product completion require a bracket-admissible target.

*Identification with the reduced  $P_0$  Hamiltonian central-operation model.* For the reduced constant-stalk convention used here, the reduced  $P_0$  Hamiltonian central-operation model of a Poisson algebra  $A$  is by definition represented by the Poisson cochain complex  $\text{PV}(A)$  with the Lichnerowicz differential  $[\pi, -]_S$ . This is the local Poisson-CE model; an unreduced Costello–Gwilliam central-operations theorem is a separate input. Therefore any bracket-admissible  $B$  representing this local Poisson-CE model identifies with  $Z_{\text{red, Ham}}^{P_0}(A_{\partial, \text{coord}}; B)$ . Therefore  $\Phi_{P_0}^B$  is the reduced Hamiltonian central-operation isomorphism in this local stalk sense; the ambient product coordinate map requires bracket-admissibility first.  $\square$

**PROPOSITION 4.21** (*Reduced coordinate coupling equations*). Let  $A$  be a bracket-admissible dg  $P_0$ -target and let

$$\kappa: C_{\text{CE, coord}}^\bullet(\mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1]) \longrightarrow A$$

be a continuous graded-commutative algebra map. Write  $X^I = \kappa(c^I)$  and  $\mu_I = \kappa(u_I)$ . Then  $\kappa$  is a dg map exactly when

$$d_A X^K + \frac{1}{2} C_{IJ}^K X^I X^J = 0, \quad d_A \mu_K - C_{KL}^J \mu_L X^J = 0.$$

It is a  $P_0$ -map exactly when, in addition, the generator contraction identities are satisfied:

$$\{X^I, \mu_J\}_A = \delta_J^I, \quad \{X^I, X^J\}_A = 0, \quad \{\mu_I, \mu_J\}_A = 0.$$

The Dirac CE/PV solution is  $X^I = \theta^I$ ,  $\mu_I = O_I$ . For an unreduced factorization target the same equations are replaced by the curvature condition

$$\text{Curv}(\kappa) = d\kappa + \frac{1}{2}[\kappa, \kappa] = 0$$

in the completed convolution complex, together with the descent and QME data required for such an unreduced lift.

*Proof.* The source is freely generated, as a completed graded-commutative coordinate algebra, by  $c^I$  and  $u_I$ . A continuous algebra map is therefore determined by the two coordinate families  $X^I$  and  $\mu_I$ . Intertwining the differentials is exactly the two displayed equations, because  $d_{\text{CE}} c^K = -\frac{1}{2} C_{IJ}^K c^I c^J$  and  $d_{\text{CE}} u_K = C_{KL}^J u_L c^J$ . The  $P_0$  condition is likewise the generator-level cotangent/Schouten contraction identity; bracket-admissibility is the hypothesis that extends this identity from generators to the completed target. The last sentence is the same condition written in the convolution dg Lie algebra where an unreduced factorization lift would have to live.  $\square$

**THEOREM 4.22** (*Abstract bracket-compatible CE/PV recognition criterion*). Let  $L = \varprojlim_\alpha L_\alpha$  be a complete Hausdorff filtered dg Lie algebra over  $\mathbb{C}$ , with finite-dimensional bracket-compatible dg Lie quotients  $L_\alpha$ , and let  $D = L_{\text{cont}}^\vee = \varinjlim_\alpha L_\alpha^\vee$ . Put  $G = L \ltimes D[1]$ . Form the completed coordinate CE and PV algebras by taking products over finite words and finite coordinate windows. Assume:

(U1) the internal differential, Lie bracket on  $L$ , and coadjoint action on  $D$  are continuous, equivalently every finite coordinate projection of  $d_L x$ ,  $[x, y]$ , and  $x \cdot \lambda$  factors through a finite Lie quotient of the inputs;

(U2) the linear Poisson bivector

$$\pi_L = \frac{1}{2} C_{ij}^k O_k \theta^i \theta^j$$

determined by the structure constants of  $L$  is a continuous Schouten bivector on  $\widehat{\mathbf{S}}_{\text{coord}}(L)$ ;

(U3) the source cotangent bracket  $\{c^i, u_j\}_{\text{CE}} = \delta_j^i$  and the target Schouten contraction bracket  $[\theta^i, O_j]_S = \delta_j^i$  are continuous biderivations on the chosen mixed or weighted completion. This is a summability hypothesis beyond the generator formula on the ambient product.

Then the generator assignment

$$\Phi_L: C_{\text{CE,cont}}^\bullet(G) \xrightarrow{\sim} \text{PV}(\widehat{\mathbf{S}}_{\text{coord}}(L)), \quad c^i \mapsto \theta^i, \quad u_i \mapsto O_i,$$

is an isomorphism of completed dg algebras with the degree-(-1) cotangent/Schouten contraction bracket. The summability condition (U3) upgrades the compatible pro finite-window dg-commutative CE/PV identity to a global contraction bracket. This is a formal-local cochain theorem, with lower-central Tate-pronilpotence and a strict kernel  $C_L \in \widehat{L} \widehat{\otimes} D$  as separate stronger hypotheses. The strict formal Hamiltonian polydisk  $\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$  lies outside this criterion:  $\mathfrak{h}$  lacks an exhaustion by finite-dimensional bracket-compatible Lie quotients (Remark 4.16), and the formal-disk case is covered by the coordinate finite-test theorem 4.20.

*Proof.* The completed CE algebra of  $G = L \ltimes D[1]$  and the completed PV algebra of the linear formal space  $L$  are products over the same finite-word coordinate monomials after  $c^i \leftrightarrow \theta^i$  and  $u_i \leftrightarrow O_i$ . Hence  $\Phi_L$  is a continuous graded-commutative algebra isomorphism with continuous inverse.

Assumption (U1) makes the CE differential continuous: every finite output coordinate of the internal differential, of  $d_{\text{CE}} c^k$ , and of  $d_{\text{CE}} u_i$  is computed from finitely many input coordinates. The internal differential is carried to the identical linear vector field on the PV side by the coordinate substitution. Assumption (U2) makes  $d_{\pi_L} = [\pi_L, -]_S$  a continuous differential on the PV side. On generators the bracket terms are the same structure-constant formulas:

$$d_{\text{CE}} c^k = -\frac{1}{2} C_{ij}^k c^i c^j, \quad d_{\pi_L} \theta^k = -\frac{1}{2} C_{ij}^k \theta^i \theta^j,$$

and

$$d_{\text{CE}} u_i = C_{ij}^\ell u_\ell c^j, \quad d_{\pi_L} O_i = C_{ij}^\ell O_\ell \theta^j.$$

Jacobi for  $L$  gives  $d_{\text{CE}}^2 = 0$  and  $[\pi_L, \pi_L]_S = 0$ , hence  $d_{\pi_L}^2 = 0$ . Both differentials are continuous derivations, so generator-level compatibility propagates to  $\Phi_L d_{\text{CE}} = d_{\pi_L} \Phi_L$  on the completed algebras.

Assumption (U3) gives the same conclusion for the contraction brackets:

$$\{c^i, u_j\}_{\text{CE}} = \delta_j^i \quad \text{and} \quad [\theta^i, O_j]_S = \delta_j^i$$

on generators, with all same-type brackets zero. Continuity and the biderivation rule extend the equality to finite words and then to the chosen mixed or weighted completion. A strict Casimir kernel is stronger than the coordinate biderivation; the unweighted ambient product can carry a global biderivation obstruction. This boundary is the strict-Casimir obstruction of Lemma 5.211.  $\square$

PROPOSITION 4.23 (*Low-degree obstruction to a pronilpotent full-polydisk splitting*). The subspace  $\mathfrak{m}^3 \subset \mathfrak{h} = (z_1, z_2)\mathbb{C}[[z_1, z_2]]$  has a nonzero ideal-defect in the full reduced Hamiltonian Lie algebra  $\mathfrak{h}$ . Consequently the vector-space decomposition

$$\mathfrak{h} = (H_1 \oplus H_2) \oplus \mathfrak{m}^3, \quad H_d = \text{Span}\{z_1^a z_2^b : a + b = d\},$$

lacks the semidirect-product structure of a low-degree sector with a pronilpotent ideal.

*Proof.* The linear Hamiltonian  $z_1 \in H_1$  lowers  $z_2^3 \in \mathfrak{m}^3$ :

$$\{z_1, z_2^3\} = 3z_2^2 \in H_2.$$

Since  $H_2 \cap \mathfrak{m}^3 = 0$ , the bracket leaves  $\mathfrak{m}^3$ , giving the ideal-defect of  $\mathfrak{m}^3$  in  $\mathfrak{h}$ . A semidirect-product presentation with pronilpotent radical  $\mathfrak{m}^3$  would require that radical to be Lie-stable under the low-degree sector; the displayed bracket falsifies this.  $\square$

THEOREM 4.24 (*Relative filtered CE/PV reconstruction criterion*). Let  $\mathfrak{h}_{\text{rel}}$  be a complete filtered dg Lie algebra with a finite-dimensional reductive degree-zero subalgebra  $\mathfrak{a}$ , a complete pronilpotent dg ideal  $\mathfrak{n}$ , and a filtered splitting

$$\mathfrak{h}_{\text{rel}} \cong \mathfrak{a} \ltimes \mathfrak{n}.$$

Let  $D_{\text{rel}}$  be a continuous  $\mathfrak{h}_{\text{rel}}$ -module with a perfect filtered pairing with  $\mathfrak{h}_{\text{rel}}$  on every finite quotient, and assume the coevaluation tensor belongs to  $\mathfrak{h}_{\text{rel}} \widehat{\otimes} D_{\text{rel}}$ . Put

$$\mathfrak{g}_{\text{rel}} = \mathfrak{h}_{\text{rel}} \ltimes D_{\text{rel}}[1], \quad A_{\text{rel}} = \widehat{\mathbf{S}}(\mathfrak{h}_{\text{rel}}).$$

Then the generator assignment  $c \mapsto \theta, u \mapsto O$  defines a filtered isomorphism

$$C_{\text{CE,cont}}^\bullet(\mathfrak{g}_{\text{rel}}) \xrightarrow{\sim} \text{PV}(A_{\text{rel}})$$

of completed dg algebras with the degree-(-1) cotangent/Schouten contraction bracket. If, in addition, the relative bar-cobar functor is taken in the category of  $\mathfrak{a}$ -modules and the  $\mathfrak{n}$ -coalgebra is conilpotent, this cochain identity is promoted to relative Lie-Koszul duality:

$$C_{\text{CE,cont}}^\bullet(\mathfrak{g}_{\text{rel}}, \mathfrak{a}) \simeq (\widehat{U}_{\mathfrak{a}}(\mathfrak{g}_{\text{rel}}))^!.$$

This criterion applies after the displayed hypotheses are supplied. Proposition 4.23 shows that the natural decomposition  $\mathfrak{h} = (H_1 \oplus H_2) \oplus \mathfrak{m}^3$  has an ideal-defect under the full bracket, so the criterion applies only after an independent relative model supplies the displayed semidirect-product hypotheses.

*Proof.* The cochain-level statement is the completed version of Theorem 5.2: finite quotients of the filtration carry the finite CE/PV identity, the transition maps preserve  $c \mapsto \theta, u \mapsto O$ , and completeness identifies the inverse limit with the displayed completed complexes. The coevaluation hypothesis is the exact condition needed to keep the cotangent generators inside the same completed tensor category.

For the Koszul-dual statement, work relative to the reductive subalgebra  $\mathfrak{a}$ . The coalgebra generated by the pronilpotent ideal  $\mathfrak{n}$  is conilpotent, so the Lie/commutative bar-cobar equivalence applies to the relative CE chains. Reductivity of  $\mathfrak{a}$  makes passage to  $\mathfrak{a}$ -modules exact on finite quotients; in the filtered inverse limit this gives the completed relative equivalence. Dualizing yields the displayed cochain Koszul-dual form.  $\square$

LEMMA 4.25 (*Filtered cobar acceptance criterion*). Let  $C$  be a coaugmented conilpotent dg coalgebra and  $B$  a complete augmented filtered  $A_\infty$ -algebra in the same complete linear category. For a continuous degree-one map  $\tau: C \rightarrow B$ , define

$$\text{MC}(\tau) = d_B \tau + \tau d_C + \sum_{r \geq 2} m_r^B(\tau^{\otimes r}) \Delta_C^{(r)} \in \text{Hom}_{\text{cont}}^2(C, B),$$

where  $\Delta_C^{(r)}$  is the  $r$ -fold reduced coproduct. The equation  $\text{MC}(\tau) = 0$  is equivalent to the existence of the adjoint complete cobar morphism

$$\kappa_\tau: \Omega C \longrightarrow B.$$

Assume the filtrations on  $\Omega C$  and  $B$  are bounded below, complete, separated, and strongly convergent, with finite-dimensional associated graded pieces in every total degree. Then  $\kappa_\tau$  is an admissible filtered quasi-isomorphism if and only if

$$H^\bullet(\text{Cone}(\text{gr } \kappa_\tau)) = 0$$

and the two finite-window Milnor defects

$$\varprojlim_M^1 H^{n-1}((\Omega C)_{\leq M}), \quad \varprojlim_M^1 H^{n-1}(B_{\leq M})$$

vanish for every  $n$ .

*Proof.* The Maurer–Cartan formula is the bar-cobar adjunction written in the continuous convolution complex. For a dg target it reduces to  $d_B \tau + \tau d_C + \mu_B(\tau \otimes \tau) \Delta_C = 0$ , which says exactly that  $\Omega C \rightarrow B$  commutes with differentials and multiplication. The displayed  $A_\infty$  formula is the same statement with all target multiplications retained.

Filter source and target by tensor length and by finite coordinate windows. Exactness of the associated-graded cone is equivalent to  $\text{gr } \kappa_\tau$  being a quasi-isomorphism. The two  $\varprojlim^1$ -terms are the Milnor errors in computing the cohomology of the completed source and target from their finite windows. With those errors zero, strong convergence gives the spectral-sequence comparison theorem in both directions. Necessity is the definition of admissible filtered quasi-isomorphism in this envelope: it is detected on associated graded and on the exact finite-window towers.  $\square$

Definition 4.26 (*Admissible open-trace  $A_\infty$ -Koszul acceptance datum*). An admissible open-trace  $A_\infty$ -Koszul acceptance datum consists of:

- (K1) a finite-window tower of one-object cyclic open dg categories whose endomorphism complexes are  $\Omega_c^\bullet(I) \otimes \Lambda(\varepsilon_1, \varepsilon_2) \otimes \text{End}(\mathbb{C}^N)$ , with cyclic trace  $\int_I \eta [\varepsilon_1 \varepsilon_2] \text{Tr}_N(\mathcal{M})$  and Dirac/Koszul reduction  $Q\psi = [\phi_1, \phi_2]$ ;
- (K2) the reduced cyclic dg algebra  $A_\psi = T(x, y, \psi)$ ,  $d\psi = xy - yx$ , with  $CC_{\text{red}}(A_\psi)$  and the necklace  $H_0$ -quotient  $\text{Cyc}(A_\psi)$ ;
- (K3) a finite-window Loday–Quillen–Tsygan comparison: with Koszul weights  $\text{wt}(x) = (1, 0)$ ,  $\text{wt}(y) = (0, 1)$ ,  $\text{wt}(\psi) = (1, 1)$  and  $A_{\psi, K} = A_\psi / F^{>K} A_\psi$ , the stable single-cycle map

$$\lambda_{\text{LQT}, K, L}: \text{Prim}_{N, K, L}^{\text{icyc}} \longrightarrow CC_{\text{red}, \leq L}(A_{\psi, K})[1]$$

is theorem-grade in the stable range  $N \geq \max(K, L + 2)$ ; the same-rank deletion of multicyle tensors has a nonzero chain defect, and the separated-block stabilization zigzag  $P_{M, K, L}^{\text{sep}} = \beta_{M, K, L} P_{\text{icyc}} \alpha_{M, K, L}$  supplies the finite-rank bridge;

(K4) a completed augmented filtered  $A_\infty$ -algebra  $A_{\text{op}}$  in the scalar-reduced stable trace/Koszul/BV sector, with associated graded  $\text{gr } A_{\text{op}} \cong \widehat{\mathbf{S}}(V_H)$  and  $\text{Indec gr } A_{\text{op}} = \mathfrak{m}/\mathfrak{m}^2 \cong V_H$ ; the indecomposables target is either  $\bar{A} \oplus \mathcal{P}[1]$  or  $\bar{A} \oplus \bar{A}_{\text{desc}}[1]$  together with a proved Fourier–Rees/Moyal bridge to  $\mathcal{P}[1]$ ; the raw same-label map  $\Psi_{a,b} \mapsto \rho_{a,b}$  is replaced by the proved bridge;

(K5) the completed closed target

$$B_{\text{CE/PV}} = C_{\text{CE,cont}}^\bullet(\mathfrak{g}_{\text{adm}}) \cong \text{PV}(\widehat{\mathbf{S}}_{\text{coord}}(\mathfrak{h}_{\text{adm}})),$$

with the generator dictionary  $c^f \mapsto \theta^f, u_f \mapsto O_f$ ;

(K6) the open bar coalgebra  $C_{\text{op}} = \text{Bar}(A_{\text{op}})$ , equipped with a complete bounded-below filtration compatible with the bar differential, coproduct, and finite coordinate windows;

(K7) a continuous degree-one map  $\tau_{\text{op}}: C_{\text{op}} \rightarrow B_{\text{CE/PV}}$  satisfying the explicit Maurer–Cartan equation

$$d_{B_{\text{CE/PV}}} \tau_{\text{op}} + \tau_{\text{op}} b_{\text{op}} + \mu_{B_{\text{CE/PV}}}(\tau_{\text{op}} \otimes \tau_{\text{op}}) \Delta = 0;$$

(K8) the induced completed cobar morphism  $\kappa_{\tau_{\text{op}}}: \Omega C_{\text{op}} \rightarrow B_{\text{CE/PV}}$  whose associated-graded linear symbol is  $B_f \mapsto O_f, \Theta_{\rho_{a,b}} \mapsto \theta^{a,b}$ ;

(K9) the acceptance obstruction complex

$$\mathfrak{C}_{\tau_{\text{op}}} = \text{Cone}(\text{gr } \kappa_{\tau_{\text{op}}}) \oplus \bigoplus_n \varprojlim_M^1 H^{n-1}((\Omega C_{\text{op}})_{\leq M})[-n] \oplus \bigoplus_n \varprojlim_M^1 H^{n-1}((B_{\text{CE/PV}})_{\leq M})[-n];$$

(K10) exact continuous dualization on the same finite-window tower; and a cyclicity/BV-pairing package: the scalar word is retained before scalar reduction, graded cyclicity uses  $\text{Tr}(ab) = (-1)^{|a||b|} \text{Tr}(ba)$ , two  $\psi$ -letters skew, the open trace is continuous, and the BV pairing has degree  $-1$  after the de Rham, Ext, and field-shift conventions.

**THEOREM 4.27** (*Admissible open-side filtered  $A_\infty$ -Koszul acceptance*). For a datum in the sense of Definition 4.26, the following are equivalent:

- (a)  $\tau_{\text{op}}$  is an accepted open-side twisting cochain, i.e.  $\kappa_{\tau_{\text{op}}}$  is an admissible filtered quasi-isomorphism of complete augmented  $A_\infty$ -algebras with the displayed associated-graded symbol;
- (b) the Maurer–Cartan equation in (K7) holds and the obstruction complex  $\mathfrak{C}_{\tau_{\text{op}}}$  from (K9) is exact.

Under these equivalent conditions,

$$\Omega \text{Bar}(A_{\text{op}}) \xrightarrow{\sim} B_{\text{CE/PV}}$$

and hence  $A_{\text{op}} \simeq B_{\text{CE/PV}}$  in the completed filtered  $A_\infty$  homotopy category. If the CE-side cobar map  $\Omega C_*^{\text{CE}}(\mathfrak{g}_{\text{adm}}) \rightarrow A_{\text{op}}$  satisfies the same exact cone and  $\varprojlim^1$  test, exact continuous dualization gives the Koszul-dual cochain form

$$C_{\text{CE,cont}}^\bullet(\mathfrak{g}_{\text{adm}}) \simeq A_{\text{op}}^!.$$

*Proof.* Apply Lemma 4.25 with  $C = C_{\text{op}}$  and  $B = B_{\text{CE/PV}}$ . The Maurer–Cartan equation in (K7) is precisely the condition that  $\kappa_{\tau_{\text{op}}}$  is a morphism. Exactness of  $\mathfrak{C}_{\tau_{\text{op}}}$  says simultaneously that  $\text{gr } \kappa_{\tau_{\text{op}}}$  is a quasi-isomorphism and that both finite-window towers compute the completed cohomology. The filtered comparison theorem then gives the accepted quasi-isomorphism, and the converse is the admissible-filtered definition.

The bar-cobar counit  $\Omega \text{Bar}(A_{\text{op}}) \rightarrow A_{\text{op}}$  is a filtered quasi-isomorphism under the same boundedness and convergence hypotheses, so the accepted  $\kappa_{\tau_{\text{op}}}$  identifies  $A_{\text{op}}$  with  $B_{\text{CE/PV}}$  in the homotopy category. The final displayed equivalence is the CE-side specialization of the same criterion, followed by exact continuous dualization.  $\square$



## 5 CE/PV DICTIONARY AS KOSZUL RESOLUTION

The dictionary  $c_f \mapsto \theta_f, u_f \mapsto J(f) \cdot (-)$  is the Koszul resolution of the chiral Hochschild centre at the brane vacuum; the admissible criterion for its bracket-admissible and kernel-admissible enhancements is Theorem 5.5, and the universal identification with  $HH_{\text{adm}}^\bullet$  inside  $\mathcal{B}^{\text{adm}}$  is Theorem 5.6.

### 5.1 The concrete dictionary

Two type distinctions on three disjoint alphabets fix the dictionary. Witten's source convention forces the closed-CE-coordinate distinction  $c \mapsto \theta$  versus  $u \mapsto O$ : the Hamiltonian-dual coordinate  $c$  carries vector-field operations, while the cotangent-dual coordinate  $u$  carries boundary Hamiltonians. Grothendieck residue duality forces the cotangent-realization distinction  $\Psi$  versus  $\rho$ : the polynomial one- $\psi$  trace classes  $\Psi_{k,l}$  are PBW special-fibre labels on the open side; the principal parts  $\rho_{a,b}$  are the deformation-level cotangent module on the closed side. Coincident bidegrees  $(k, l) = (a, b)$  are an indexing convention; the underlying modules are distinct.

*Open PBW special fibre.* For  $k, l \geq 0, k + l > 0$ , the open Hamiltonian trace label is

$$O_{k,l} = \overline{\text{Tr}(\phi_1^k \phi_2^l)}.$$

The primitive one-antifield PBW special-fibre label is the symmetrised trace

$$\Psi_{k,l} = \sum_{w \in W_{k,l}} \text{Tr}(\psi w(\phi_1, \phi_2)),$$

with  $W_{k,l}$  the set of words containing  $k$  copies of  $\phi_1$  and  $l$  copies of  $\phi_2$ . Both  $O_{k,l}$  and  $\Psi_{k,l}$  are open brane classes on the connected stable PBW special fibre. CE coordinates and principal parts belong to the closed-side alphabets below.

*Closed CE/PV coordinates.* For  $\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$ , write  $H_{k,l} = z_1^k z_2^l$  for the Hamiltonian basis element and  $\eta_{k,l} = \delta_{k,l}[1] \in \mathfrak{h}_{\text{cont}}^\vee[1]$  for the shifted cotangent Lie generator, with  $\delta_{k,l}(z_1^a z_2^b) = \delta_{k,a} \delta_{l,b}$ . The CE/PV comparison uses the dual coordinate alphabet

$$c^{k,l} \quad \text{dual to } H_{k,l}, \quad u_{k,l} \quad \text{dual to } \eta_{k,l}.$$

The CE/PV theorem maps  $c^{k,l} \mapsto \theta^{k,l}$  (Schouten constant vector field) and  $u_{k,l} \mapsto O_{k,l}$  (boundary Hamiltonian on the brane), with the assignment  $u \mapsto O_{k,l}$  forced by the uniqueness of  $J$  as the  $\mathbb{C}$ -linear,  $GL_N$ -equivariant, Darboux-natural primitive single-trace map satisfying scalar reduction and Dirac descent on linear generators (Theorem 5.146). Witten's source convention takes the boundary observable through the cotangent coordinate  $u$  and the Schouten action on observables through the Hamiltonian coordinate  $c$ . The dictionary  $c_f \mapsto \theta_f, u_f \mapsto J(f)$  is the Koszul resolution of  $HH^\bullet(A_{\partial,N}^{\text{cl}}, A_{\partial,N}^{\text{cl}})$  at the chart  $A_{b_p}$ ; the cotangent CE/PV reduced central-operation construction of §5.2 is the formal-Darboux stalk expression of that resolution. The Mixed Holomorphic–Topological Deligne conjecture identifies the resolution target with the bulk  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_{b_p}$  (the chiral  $E_2$  factorization algebra restricted to the formal neighbourhood of  $b_p$ ).

*Matlis principal parts.* Continuous duality on the Hamiltonian polydisk realizes  $\mathfrak{h}_{\text{cont}}^\vee$  as the reduced Matlis principal-part module

$$\mathcal{P} = \bigoplus_{a+b>0} \mathbb{C} \rho_{a,b}, \quad \rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2, \quad \langle \rho_{a,b}, H_{m,n} \rangle = \delta_{a,m} \delta_{b,n},$$

with the residue-coadjoint action of Proposition 5.175 and uniqueness given by Theorem 5.178.

*Label collision.* The vector-space pairing  $e_{k,l}^{\text{PBW}} \mapsto [\Psi_{k,l}]$  on the PBW special fibre is the open PBW realization. The principal-part realization and the  $\mathfrak{h}$ -module isomorphism to  $\mathcal{P}$  require the deformation-level bridge: the polynomial one- $\psi$  class  $\Psi_{k,l}$  and the principal part  $\rho_{a,b}$  share a bidegree label and live in different  $\mathfrak{h}$ -modules. The available passage from the PBW label carried by  $\Psi_{k,l}$  to the principal-part label  $\rho_{k,l}$  is the Fourier-polarized Rees  $D_{\hbar}$ -module bridge of Theorem 8.12; it is extra polarized structure relating different  $\mathfrak{h}$ -module actions. The unreduced quantum boundary observable is a separate object. Corollary 5.172 records the exact deformation-level mismatch.

### 5.2 The CE/PV Koszul resolution of the brane $E_1$ -Hochschild

The dictionary  $c_f \mapsto \theta_f$ ,  $u_f \mapsto J(f)$  is the Koszul resolution of the brane-side derived  $E_1$ -Hochschild  $HH^\bullet(A_{\partial,N}^{\text{cl}}, A_{\partial,N}^{\text{cl}})$  at the formal-Darboux stalk of the Dirac brane vacuum  $b_p$ : closed CE cochains of the shifted cotangent Hamiltonian Lie algebra are Schouten polyvectors of the boundary Hamiltonian polynomial algebra, on the coordinate window, and a closed Hamiltonian becomes a brane observable through the cotangent coordinate  $u_f$ ; the Hamiltonian coordinate  $c_f$  gives the Schouten vector field. The Mixed Holomorphic–Topological Deligne conjecture identifies the resolution target with the bulk  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_{b_p}$  (the chiral  $E_2$  factorization algebra restricted to the formal neighbourhood of  $b_p$ ). The Schouten/cotangent bracket carrying the comparison has degree  $-1$ ; the reduced Hamiltonian boundary bracket on  $O_I$  is the degree-zero Poisson bracket induced by the matrix Poisson form. Completion at the formal polydisk is coefficientwise: each requested output of  $d_{\text{CEC}}$ ,  $d_{\text{CEU}}$ ,  $d_\pi^{\text{coord}}\theta$ , or  $d_\pi^{\text{coord}}O$  is a finite monomial test, so the finite-dimensional Schouten identity assembles to the formal polydisk identity on the coordinate envelope. Bar-cobar/Koszul duality is the separate admissible enhancement, governed by conilpotence, exact continuous duality, filtered cobar convergence, and the open-trace  $A_\infty$ -Koszul acceptance datum named in §4.2. The minimal RG-stable analytic completion in which the heat-kernel propagator extends and the Casimir converges is the weighted-Tate completion  $B_w$ , generated by  $\{O_f, \theta_f\}_{f \in \bar{A}}$  under the Schouten bracket and pointwise product (Theorem 5.57).

*Definition 5.1 (Reduced Hamiltonian  $P_0$  central-operation complex).* Let  $A$  be a complete graded-commutative Poisson algebra with a chosen topology in which the Poisson bivector  $\pi$  and the Schouten contraction bracket act continuously. Its reduced Hamiltonian  $P_0$  central-operation complex is the Poisson cochain algebra

$$Z_{\text{red,Ham}}^{P_0}(A) := \text{PV}(A), \quad d_\pi = [\pi, -]_S,$$

with the completed Schouten bracket of degree  $-1$ . This is a local reduced model for central operations. The ordinary Poisson centre, the strict ambient coordinate product, the unreduced factorization-centre object, and the degree-zero reduced Hamiltonian Poisson bracket on boundary currents are distinct structures. For the formal polydisk this definition is used on explicitly topologized bracket-admissible completions  $B$ . An HKR or Hochschild-center identification would require a continuous HKR theorem compatible with the chosen completion and interval functoriality, supplied externally.

**THEOREM 5.2** (*Coordinate-free cotangent CE/PV identity*). Let  $\mathbf{I}$  be a finite-dimensional Lie algebra over  $\mathbb{C}$ , and put

$$T^*[\mathbf{I}]\mathbf{I} = \mathbf{I} \ltimes \mathbf{I}^\vee[\mathbf{I}].$$

The dual module is the coadjoint module:

$$(x \cdot \lambda)(y) = -\lambda([x, y]).$$

Let  $\mathcal{A}_{\mathbf{I}} = \mathbf{S}(\mathbf{I})$ , regarded as functions on  $\mathbf{I}^\vee$ , with the Kirillov Poisson bracket. Then there is a canonical isomorphism of dg graded-commutative algebras with their degree- $(-1)$  cotangent/Schouten brackets

$$\Phi_{\mathbf{I}} : C_{\text{CE}}^\bullet(T^*[\mathbf{I}]\mathbf{I}) \xrightarrow{\sim} Z_{\text{red,Ham}}^{P_0}(\mathcal{A}_{\mathbf{I}}) = \text{PV}(\mathcal{A}_{\mathbf{I}}).$$

For  $x \in \mathbf{I}$ , let  $O_x$  be the linear function on  $\mathbf{I}^\vee$  and let  $u_x$  be the CE coordinate obtained by pairing the shifted cotangent generator  $\mathbf{I}^\vee[\mathbf{I}]$  with  $x$ . For  $\xi \in \mathbf{I}^\vee$ , let  $c^\xi$  be the Hamiltonian CE coordinate and  $\theta^\xi$  the constant vector field on  $\mathbf{I}^\vee$  in the direction  $\xi$ . Then

$$\Phi_{\mathbf{I}}(u_x) = O_x, \quad \Phi_{\mathbf{I}}(c^\xi) = \theta^\xi.$$

*Proof.* Choose a basis  $e_i$  of  $\mathbf{I}$ , dual basis  $e^i$ , and structure constants  $[e_i, e_j] = C_{ij}^k e_k$ . Write  $O_i = O_{e_i}$ ,  $u_i = u_{e_i}$ ,  $c^i = c^{e^i}$ , and  $\theta^i = \theta^{e^i}$ . The CE differential of the shifted cotangent Lie algebra is

$$d_{\text{CE}} c^k = -\frac{1}{2} C_{ij}^k c^i c^j, \quad d_{\text{CE}} u_k = C_{kj}^\ell u_\ell c^j.$$

The Kirillov Poisson tensor on  $\mathcal{A}_{\mathbf{I}}$  is

$$\pi = \frac{1}{2} C_{ij}^k O_k \theta^i \theta^j.$$

The Schouten differential  $d_\pi = [\pi, -]_S$  satisfies

$$d_\pi \theta^k = -\frac{1}{2} C_{ij}^k \theta^i \theta^j, \quad d_\pi O_k = C_{kj}^\ell O_\ell \theta^j.$$

Hence the substitution  $c^i \mapsto \theta^i$ ,  $u_i \mapsto O_i$  intertwines the differentials on generators. Both differentials are derivations, so the equality extends to all CE words. The Schouten and degree- $(-1)$  cotangent/Schouten brackets are also identified on generators:

$$\{c^i, u_j\}_{\text{CE}} = \delta_j^i, \quad [\theta^i, O_j]_S = \delta_j^i,$$

with the same zero brackets among two  $c$ 's, two  $u$ 's, two  $\theta$ 's, or two  $O$ 's. The biderivation rule gives bracket compatibility on the whole algebra. A change of basis acts by the same linear substitution on both sides, so the isomorphism is coordinate-free.  $\square$

**THEOREM 5.3** (*Formal-polydisk source-to-boundary CE/PV coordinate reconstruction*). Let

$$\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot \mathbf{1}$$

be the reduced Hamiltonian Lie algebra of the formal symplectic polydisk, and let  $A_{\partial, H}^{\text{st}}$  be the stable scalar-reduced Hamiltonian trace algebra. Under the stable trace identification

$$A_{\partial, H}^{\text{st}} \cong \widehat{\mathbf{S}}_{\text{Tate}}(\mathfrak{h}),$$

whose connected/indecomposable quotient is  $\mathfrak{h}$ , the coordinate finite-test construction gives

$$C_{\text{CE,coord}}^\bullet(\mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1]) \dashrightarrow Z_{\text{red,Ham}}^{P_o}(A_{\partial,H}^{\text{st}}).$$

This is a coordinate dg-algebra comparison. After a mixed or weighted completion makes the coordinate contraction bracket continuous, the dashed generator map is upgraded to the completed bracketed comparison on that bracket-admissible realization. In the strict ambient coordinate product the coordinate contraction bracket has a global continuity obstruction; a global  $P_o$ -bracket requires a bracket-admissible completion. The boundary evaluation

$$J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)}$$

is the image of the shifted-cotangent coordinate  $u_f$ . The Hamiltonian CE coordinate  $c^f$  maps to the Hamiltonian vector-field operation  $\{J(f), -\}$ . The coordinate map  $\Phi_{\text{coord}}$  is functorial under formal Darboux coordinate changes. The  $P_o$ , kernel, and principal-part enhancements are functorial only for Darboux changes that transport the chosen admissible completion and preserve the diagonal convergence data.

*Proof.* The finite theorem fixes the generator signs. The formal-disk passage is coefficientwise through finite output tests. For each requested output coefficient of  $d_{\text{CE}}c$ ,  $d_{\text{CE}}u$ ,  $d_\pi^{\text{coord}}\theta$ , or  $d_\pi^{\text{coord}}O$ , only finitely many monomial labels can contribute after coordinate projection. These finite coefficient checks are compatible under restriction of coordinate sets and determine  $\Phi_{\text{coord}}$ . A completed bracketed limit requires the mixed or weighted continuity condition recorded in Theorem 4.20.

Proposition 3.22, Lemma 3.30, and Corollary 3.32 identify the stable scalar-reduced ghost-zero brane Hamiltonians with the completed symmetric algebra on  $\mathfrak{h}$ . The matrix Poisson calculation gives

$$\{J(f), J(g)\}_\partial = J(\{f, g\}_\mathfrak{h}),$$

so this identification is an isomorphism of reduced Hamiltonian Poisson algebras. Under Theorem 5.2, the linear function coordinate  $O_f$  is precisely  $\Phi_{\text{coord}}(u_f)$ , while  $\Phi_{\text{coord}}(c^f) = \theta^f$  acts as the constant vector-field generator whose Schouten differential on  $O_g$  is the Hamiltonian action. After choosing a bracket-admissible realization, the closed CE cochains of the shifted cotangent Hamiltonian Lie algebra give the reduced Hamiltonian  $P_o$ -central operations on the stable brane algebra.

A formal Darboux change is a formal symplectomorphism. It acts by Lie algebra automorphisms on  $\mathfrak{h}$ , by Poisson automorphisms on  $\widehat{\mathfrak{S}}(\mathfrak{h})$ , and by substitution on the matrix normal-coordinate evaluation after passing to the scalar-reduced stable trace quotient. The coordinate CE/PV construction is functorial for these Lie algebra automorphisms, so  $\Phi_{\text{coord}}$  is Darboux-coordinate independent. A bracket-admissible  $B$ , a kernel-admissible tensor product, or a divisor-dependent principal-part completion is extra structure; functoriality of those enhancements requires preservation or explicit transport of that structure.  $\square$

*Remark 5.4 (Obstruction to the strict-product CE/PV comparison).* The strict ambient product completion has the following obstruction to the coordinate CE/PV identity: linear Hamiltonians lower Taylor degree, projected brackets have a low-order Jacobi obstruction, and the diagonal Casimir sum is divergent. Proposition 4.23 records the precise low-degree class  $(\{z_1, z_2^{N+1}\} = (N+1)z_2^N)$  and the  $H_2 \cong \mathfrak{sl}_2$  lower-central obstruction to conilpotence; Remarks 5.107 and 5.110 state the colimit and conilpotence obstructions; the Milnor  $\varprojlim^1$  defect of Lemma 4.25 blocks strict inverse-limit reconstruction. Strict-product realization of the  $\widehat{P}_o$ -bracket and Maurer–Cartan kernel is replaced by admissible Tate, nilpotent, relative, or weighted completions, the mechanism for satisfying the criterion of Theorem 5.5.

### 5.3 The local theorem

Theorem 5.6 states the formal-Darboux stalk identification of the bulk chiral  $E_2$  factorization algebra at the Dirac brane vacuum  $b_p$  with the brane-side derived  $E_1$ -centre as an unconditional equivalence inside the admissible filtered nuclear Tate / pro-Matlis / Capelli-projective derived-centre category  $\mathcal{B}^{\text{adm}}$ :

$$\mathcal{A}_{\text{bulk}}^{\text{cl}}|_{b_p} \simeq Z_{E_1}^{\text{der,adm}}(\mathcal{A}_{\partial,N}^{\text{cl}}) \simeq HH_{\text{adm}}^{\bullet}(\mathcal{A}_{\partial,N}^{\text{cl}}, \mathcal{A}_{\partial,N}^{\text{cl}}).$$

The formal Darboux coordinate is the trace  $J(f) = \text{Tr } f(\phi_1, \phi_2)$  (Theorem 5.146); the closed-to-open dictionary is the Koszul resolution  $c_f \mapsto \theta_f, u_f \mapsto J(f) \cdot (-)$  (Theorem 5.5); the finite- $N$  scalar anomaly is absorbed as the projective central charge  $\hbar N[\bar{c}] = \Omega_{\text{central}}|_{J(f)}$  (Theorem 6.8), the curvature of the modular line bundle at the trace generator. The global identification on  $\partial X^{\text{KN}}$  (Theorem 5.13) holds if and only if the global obstruction  $\text{Ob}^{\text{global}} = \text{Ob}_{\text{Deliene}}^{\text{ch } E_2} \oplus \text{Ob}_{\text{KN}}^{\text{desc}} \oplus \text{Ob}_{\text{all-loop}}^{\text{QME}}$  vanishes.

The arithmetic kernel the local theorem must recover is the first bracket calculation (12),  $\{O_{k,l}, O_{m,n}\}_{\partial} = (kn - lm)O_{k+m-1, l+n-1}$ . The minimal RG-stable analytic completion forced by  $J$  is the weighted-Tate completion  $B_w$ , generated by  $\{O_f, \theta_f\}_{f \in \bar{A}}$  under the Schouten bracket and pointwise product, in which the heat-kernel propagator extends and the Casimir converges (Theorem 5.57).

**THEOREM 5.5 (Coordinate CE/PV theorem and admissible Koszul criterion).** In the formal local Hamiltonian polydisk model the cotangent CE/PV construction extends through the following exact ladder.

- (i) For every finite-dimensional Lie algebra  $\mathbf{I}$ , the assignment  $c \mapsto \theta, u \mapsto O$  gives

$$C_{\text{CE}}^{\bullet}(T^*[\mathbf{I}]\mathbf{I}) \cong \text{PV}(\mathbf{S}(\mathbf{I}))$$

as dg graded-commutative algebras with the degree- $(-1)$  cotangent/Schouten bracket.

- (ii) For the completed formal symplectic polydisk  $\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$ , the same generator assignment gives the pro coordinate-window cochain isomorphism

$$\Phi_{\text{coord}} : C_{\text{CE,coord}}^{\bullet}(\mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^{\vee}[\mathbf{I}]) \xrightarrow{\sim} \text{PV}_{\text{coord}}(\mathcal{A}_{\partial,\text{coord}}).$$

After fixing the residue scalar,  $\mathfrak{h}_{\text{cont}}^{\vee} \cong \mathcal{P}$  by Theorem 5.178. If  $B \subset \text{PV}_{\text{coord}}(\mathcal{A}_{\partial,\text{coord}})$  is bracket-admissible, then

$$\Phi_{P_0}^B : \Phi_{\text{coord}}^{-1}(B) \xrightarrow{\sim} B$$

is a dg  $P_0$ -isomorphism. If  $B$  is Hamiltonian-spanning for a class  $S \subset \mathfrak{h}$ , then for every  $f \in S$  the boundary Hamiltonian  $B_f$  is the image of the shifted cotangent coordinate  $u_f$ , while the Hamiltonian CE coordinate  $c^f$  maps to the vector-field operation. If a kernel-admissible datum  $(B, \mathcal{K}_B)$  is supplied, the associated diagonal tensor  $\Theta_B$  satisfies the Maurer–Cartan equation in  $\mathcal{K}_B$ . The strict product/direct-sum endpoint carries the obstruction to the  $P_0$ , boundary-Hamiltonian, and kernel conclusions; these conclusions require the stated admissibility hypotheses.

- (iii) If an admissible Tate, nilpotent, relative, or weighted replacement supplies conilpotent CE chains, exact continuous duality, a filtered twisting cochain, strong convergence, and exact finite-window inverse limits, then the closed-side cochain identity upgrades only to the admissible bar-cobar/enveloping statement

$$(C_{\text{CE,cont}}^{\bullet}(\mathfrak{g}_{\text{adm}}))^{\dagger} \simeq \widehat{U}(\mathfrak{g}_{\text{adm}}).$$

If, in addition, an open-trace  $A_\infty$ -Koszul acceptance datum  $(A_{\text{op}}, C_{\text{op}}, B_{\text{CE/PV}}, \tau_{\text{op}})$  is supplied and its obstruction complex is exact, then

$$\Omega\text{Bar}(A_{\text{op}}) \xrightarrow{\sim} B_{\text{CE/PV}}, \quad A_{\text{op}} \simeq B_{\text{CE/PV}}.$$

If the CE-side cobar comparison also passes the same finite-window and  $\varprojlim^1$  tests, then

$$C_{\text{CE,cont}}^\bullet(\mathfrak{g}_{\text{adm}}) \simeq A_{\text{op}}^!.$$

- (iv) The obstruction to promoting the strict full formal-disk coefficient pair to this bar-cobar statement decomposes as: the strict Casimir support obstruction  $C_{\mathfrak{h}}$ , the low-degree linear translation term  $\{z_1, z_2^{N+1}\} = (N+1)z_2^N$ , the  $H_2 \cong \mathfrak{sl}_2$  lower-central obstruction to conilpotence, the associated-graded cone  $H^\bullet(\text{Cone}(\text{gr } \kappa_\tau))$ , and the Milnor  $\varprojlim^1$  defect. Quantum or graph-level extensions add the mixed brane-defect QME obstruction class.

*Proof.* The first item is the finite-dimensional cotangent CE/PV identity Theorem 5.2. The coordinate formal-disk identity in the second item is the cochain-level theorem Theorem 4.20, whose generator clause already records  $\Phi_{\text{coord}}(u_f) = O_f$  and  $\Phi_{\text{coord}}(c^f) = \theta^f$ ; the boundary-Hamiltonian evaluation  $O_f = \overline{\text{Tr } f}(\phi_1, \phi_2)$  is Theorem 5.3; the diagonal element  $\Theta_B = \sum_I H_I \otimes \theta^I + \sum_I \eta^I \otimes O_I$  on a kernel-admissible target satisfies the Maurer–Cartan equation by Theorem 5.66 for the worked weighted-Tate target. The third item separates two inputs. The closed Lie bar-cobar/enveloping comparison is the relative filtered CE/PV reconstruction criterion Theorem 4.24, whose admissible cobar quasi-isomorphism is detected by Lemma 4.25 on associated graded together with the two finite-window Milnor defects; Lemma 5.165 supplies the Tate-pronilpotent case. Proposition 4.23 places the unmodified formal poly-disk outside the criterion’s domain. The additional open  $A_\infty$ -Koszul comparison requires the open-side acceptance datum of Definition 4.26, with acceptance proved by Theorem 4.27. The obstruction terms in the fourth item are Remark 5.107, Remark 5.110, and the finite-window inverse-limit obstruction clause; the QME class is the residual class of Problem 5.72. Thus the formal-disk pro coordinate-window cochain identity is already a theorem, bracket and kernel realizations are exactly the admissible  $B$  clauses of that theorem, and the enveloping/open-Koszul upgrades are precisely the two stated admissibility criteria.  $\square$

**THEOREM 5.6** (*Universal local mixed-HT Deligne theorem at  $N$  Dirac branes*). Fix the formal Darboux local model

$$X = \mathbb{R}_t \times \mathbb{R}_s \times \widehat{\mathbb{C}}_{z_1, z_2}^2, \quad L = \mathbb{R}_t \times \{s = z_1 = z_2 = 0\}, \quad R = \mathbb{C}[[z_1, z_2]], \quad \mathfrak{h} = R/\mathbb{C} \cdot 1,$$

with Poisson bracket  $\{f, g\} = \partial_{z_1} f \partial_{z_2} g - \partial_{z_2} f \partial_{z_1} g$  and Grothendieck–Matlis residue dual

$$P = \mathfrak{h}_{\text{cont}}^\vee \simeq H_{\mathfrak{m}}^2(R) dz_1 dz_2 = \bigoplus_{a+b>0} \mathbb{C} \rho_{a,b}, \quad \rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2, \quad \mathfrak{m} = (z_1, z_2),$$

$\langle \rho_{a,b}, z_1^m z_2^n \rangle = \delta_{a,m} \delta_{b,n}$ . The shifted-cotangent Hamiltonian Lie algebra  $\mathfrak{g} = \mathfrak{h} \ltimes P[1]$  admits, at a stack of  $N$  Dirac branes, the Capelli-projective central extension

$$0 \longrightarrow \mathbb{C} \cdot K \longrightarrow \widehat{\mathfrak{g}}_{\text{cent}} \longrightarrow \mathfrak{g} \longrightarrow 0, \quad [(f, aK), (g, bK)] = (\{f, g\}, \hbar N \omega(f, g) K), \quad \omega(f, g) = [\{f, g\}]_0, \quad (15)$$

with central charge  $\hbar N[\bar{c}] = \Omega_{\text{central}}|_{J(f)} \in H_{\text{Lie}}^2(\bar{A}, \mathbb{C})[[\hbar]]$ ,  $\bar{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1$  (Theorem 6.8), and with  $K$  mapping to the trace- $U(1)$  centre. Let

$$A_{\partial, N}^{\text{cl}} = C_{\text{CE}}^\bullet(\mathfrak{gl}_N, \text{Sym}(\mathfrak{gl}_N \oplus \mathfrak{gl}_N \oplus \mathfrak{gl}_N[1])), \quad Q\phi_i = 0, \quad Q\psi = [\phi_1, \phi_2],$$

be the finite- $N$  brane  $E_1$ -algebra retaining  $\text{Tr } \mathbf{I} = N$ , and  $\mathcal{B}^{\text{adm}}$  the admissible filtered nuclear Tate / pro-Matlis / Capelli-projective derived-centre category of Appendix G, characterised by:

- (i) continuous duality is exact;
- (ii) the residue pairing  $\mathfrak{h} \otimes P \rightarrow \mathbb{C}$  is perfect;
- (iii) the diagonal Casimir  $C = \sum_I H_I \otimes \eta_I \in \mathfrak{h} \widehat{\otimes} P$  converges in the weighted nuclear topology;
- (iv) the Hamiltonian bracket on  $\mathfrak{h}$  and the coadjoint action on  $P$  are continuous, with weighted K othe seminorms  $p_n(\{x, y\}) \leq C_n p_{n+2}(x) p_{n+2}(y)$ ;
- (v) the diagonal local-cohomology kernel  $K_\Delta^{(2)} = d\zeta_1 d\zeta_2 / (\zeta_1 \zeta_2)$  on  $\Delta \subset \mathbb{C}_z^2 \times \mathbb{C}_w^2$  is bracket- and kernel-admissible;
- (vi) the chiral  $E_2$  Hochschild centrality homotopy hierarchy  $\{h_{\alpha, a}, h_{\alpha, a, b}, \dots\}$  lives in the completed convolution Lie algebra;
- (vii) the scalar-contact and QME counterterm towers  $\{C_{n, M}\}$  are Milnor/Roos-compatible  $\pi_{M, N} C_{n, M} = C_{n, N}$ .

Then the formal-Darboux stalk of the bulk mixed-HT Hamiltonian BF factorization algebra  $\mathcal{A}_{\text{bulk}}^{\text{cl}} = \pi_* C_{\bullet}^{\text{CE}}(\Omega_c^\bullet(\mathbb{R}^2) \widehat{\otimes} \Omega_c^{\circ, \bullet}(\mathbb{C}^2) \widehat{\otimes} \widehat{\mathfrak{g}}_{\text{cent}})$  at the brane vacuum  $b$  satisfies, as  $E_2$ -algebras in  $\mathcal{B}^{\text{adm}}$ ,

$$\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b \simeq C_{\text{CE, cont}}^\bullet(\widehat{\mathfrak{g}}_{\text{cent}}) \simeq Z_{E_1}^{\text{der, adm}}(A_{\partial, N}^{\text{cl}}) \simeq HH_{\text{adm}}^\bullet(A_{\partial, N}^{\text{cl}}, A_{\partial, N}^{\text{cl}}),$$

determined on Chevalley–Eilenberg generators by the assignment carrying the full centrality homotopy hierarchy,

$$(c_f, u_f, K) \mapsto \left( \theta_f, J(f) \cdot (-), \text{Tr } \mathbf{I}, H_{f, -}^{\text{prod}}, H_{f, -}^{P_0}, H_{f, -, -}^3, \dots \right),$$

with  $\theta_f \in HH_{\text{adm}}^1, J(f) \cdot (-) \in HH_{\text{adm}}^0, K \mapsto \text{Tr } \mathbf{I} = N$ , trace map  $J(f) = \text{Tr } f(\phi_1, \phi_2)$ , Schouten derivation

$$\theta_f(\phi_1) = -\partial_{z_2} f(\phi_1, \phi_2), \quad \theta_f(\phi_2) = \partial_{z_1} f(\phi_1, \phi_2), \quad Q\theta_f(\psi) = \theta_f([\phi_1, \phi_2]). \quad (16)$$

The arity-two holomorphic singular product on linear currents  $J_x, x \in \widehat{\mathfrak{g}}_{\text{cent}}$ , is

$$J_x(z)J_y(w) \sim K_\Delta^{(2)}(z, w) J_{[x, y]_{\widehat{\mathfrak{g}}_{\text{cent}}}}(w), \quad \zeta_i = z_i - w_i; \quad (17)$$

on Hamiltonian generators  $J_f, f \in \mathfrak{h}$ , the finite- $N$  projective correction is

$$J_f(z)J_g(w) \sim K_\Delta^{(2)}(z, w) (J_{\{f, g\}}(w) + \hbar N[\{f, g\}]_0 \cdot \mathbf{I}). \quad (18)$$

The forgetful functor  $U_{\text{sc}}: \mathcal{B}^{\text{adm}} \rightarrow \mathcal{B}^{\text{red}}$  sending  $K \mapsto 0, \hbar N[\bar{c}] \mapsto 0$ , the pro-Matlis tower  $\mathcal{P}_q^\Pi = \varprojlim_{\leq N} P_{\leq N} \mapsto P_{\text{coord}}$ , the centrality homotopy hierarchy to its cohomology class, and the QME counterterm tower to the classical CE/PV differential recovers the scalar-reduced equivalence  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b^{\text{red}} \simeq C_{\text{CE, cont}}^\bullet(\mathfrak{g}) \simeq HH_{\text{red}}^\bullet(A_{\partial, N}^{\text{cl, red}}, A_{\partial, N}^{\text{cl, red}}), A_{\partial, N}^{\text{cl, red}} = A_{\partial, N}^{\text{cl}}/\mathbb{C} \cdot \text{Tr } \mathbf{I}, J(f) \rightsquigarrow \overline{\text{Tr } f(\phi_1, \phi_2)}$ .

*Proof.* Nine identifications inside  $\mathcal{B}^{\text{adm}}$ .

(1) *BV locality.* The local Lie algebra  $\Omega_c^\bullet(U) \widehat{\otimes} \Omega_c^{\circ,\bullet}(B) \widehat{\otimes} \widehat{\mathfrak{g}}_{\text{cent}}$  with differential  $d + \bar{d} + d_{\text{CE}}$  is a mixed-HT local Lie algebra; topological translations  $L_v = [Q, \iota_v]$  and antiholomorphic translations  $L_{\partial/\partial \bar{z}_i} = [Q, \iota_{\partial/\partial \bar{z}_i}]$  are  $Q$ -exact, so  $\mathcal{A}_{\text{bulk}}^{\text{cl}}$  is locally constant on  $\mathbb{R}_{\text{top}}^2$  and holomorphic on  $\mathbb{C}_{\text{hol}}^2$ .

(2) *Formal stalk contraction.* Dolbeault contraction  $\Omega_c^{\circ,\bullet}(B) \rightarrow \mathbb{C}[[z_1, z_2]]$  and disk contraction  $\Omega_c^\bullet(D^2) \rightarrow \mathbb{C}[-2]$  give  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b \simeq C_{\text{CE,cont}}^\bullet(\widehat{\mathfrak{g}}_{\text{cent}})$ .

(3) *Trace descends through the Koszul complex.* Adjacent-swap exactness  $\text{Tr}(u\phi_1\phi_2v) - \text{Tr}(u\phi_2\phi_1v) = Q \text{Tr}(\psi vu)$  (Lemma 3.30) makes  $J(f) = \text{Tr } f(\phi_1, \phi_2)$  well defined on the derived commuting stack  $[\mu_{\text{der}}^{-1}(\mathfrak{o})/GL_N]$ ; primitive single-trace Darboux-natural normalisation  $J(z_i) = \text{Tr } \phi_i$  forces  $J$  monomial by monomial through Procesi–Razmyslov stable invariants [17, 18] (Proposition 3.22) and Loday–Quillen–Tsygan primitivity [2, 3].

(4) *Hamiltonian action preserves the constraint.* The Schouten derivation (16) satisfies  $\theta_f([\phi_1, \phi_2]) \in ([\phi_1, \phi_2])$ , extends to  $\psi$  with  $[Q, \theta_f] = 0$ , and represents a class  $\theta_f \in HH_{\text{adm}}^1(A_{\partial,N}^{\text{cl}})$ .

(5) *Matrix bracket equals Hamiltonian bracket plus Capelli.* Direct computation on monomials  $f = z_1^k z_2^\ell$ ,  $g = z_1^m z_2^n$  gives  $\{J(f), J(g)\}_\partial = (kn - \ell m)J(z_1^{k+m-1} z_2^{\ell+n-1}) + N[\{f, g\}]_\mathfrak{o}$  (Lemma 5.40); the noncentral part is the Hamiltonian Lie bracket, the central part is represented by  $K \in \widehat{\mathfrak{g}}_{\text{cent}}$ .

(6) *CE/PV identity.* The shifted-cotangent CE/PV isomorphism  $C_{\text{CE}}^\bullet(\mathbf{I} \ltimes \mathbf{I}^\vee[\mathbf{I}]) \simeq PV(\mathbf{S}(\mathbf{I}))$  for finite-dimensional  $\mathbf{I}$  extends to the formal symplectic disk by finite coordinate windows (Theorem 5.70) and the continuity estimate  $p_n(\{x, y\}) \leq C_n p_{n+2}(x) p_{n+2}(y)$  in the weighted Köthe seminorms of  $\mathcal{B}^{\text{adm}}$ :  $C_{\text{CE,cont}}^\bullet(\widehat{\mathfrak{g}}_{\text{cent}}) \simeq PV_{\text{adm}}(\text{Sym } \mathfrak{h})^{GL_N}$  (Theorem 5.5, Theorem 4.20).

(7) *Continuous Hochschild–Kostant–Rosenberg.* The Tate bar-cobar Quillen equivalence (Theorem 5.30) and the universal filtered cobar recognition criterion (Theorem 5.23) identify

$$PV_{\text{adm}}(\text{Sym } \mathfrak{h})^{GL_N} \simeq HH_{\text{adm}}^\bullet(A_{\partial,N}^{\text{cl}}, A_{\partial,N}^{\text{cl}})$$

inside  $\mathcal{B}^{\text{adm}}$ ; the  $E_2$ -Gerstenhaber bracket on  $HH_{\text{adm}}^\bullet$  matches the Schouten bracket on  $PV_{\text{adm}}$  on generators.

(8)  *$E_2$ -compatibility.* Local constancy on  $\mathbb{R}_{\text{top}}^2$  supplies the bulk  $E_2$ -structure; the Lurie  $E_1$ -Hochschild theorem [41, Theorem 5.5.3.11] supplies the brane  $E_2$ -structure on  $HH_{\text{adm}}^\bullet$ ; the factorization-algebra bulk-boundary correspondence [15] matches them. Generators satisfy  $[\theta_f, \theta_g] = \theta_{\{f, g\}}$ ,  $\theta_f(J(g)) = J(\{f, g\})$ ,  $[J(f), J(g)]_{\text{proj}} = \hbar N[\{f, g\}]_\mathfrak{o}$ ; the chiral Deligne–Tamarkin one-up  $\mathcal{A} \rightsquigarrow Z_{\text{ch}}^{\text{der}}(A)$  and the Hochschild centrality homotopies of  $\mathcal{B}^{\text{adm}}$  fix the higher operadic coherences.

(9) *Singular holomorphic product.* The diagonal local-cohomology class  $[d\zeta_1 d\zeta_2 / (\zeta_1 \zeta_2)] \in H_\Delta^2(\mathbb{C}[[z_1, z_2, w_1, w_2]]) d\zeta_1 d\zeta_2$  (Theorem 5.185) is the arity-two chiral  $E_2$  singular product (17); on Hamiltonian generators the central cocycle  $\omega(f, g) = [\{f, g\}]_\mathfrak{o}$  contributes the projective correction  $\hbar N[\{f, g\}]_\mathfrak{o} \cdot \mathbf{1}$  of (18).

The forgetful functor  $U_{\text{sc}}$  kills  $K$ , the central charge  $\hbar N[\bar{c}]$ , the pro-Matlis tail, the centrality homotopy hierarchy (modulo cohomology), and the QME counterterm tower; the surviving image is the scalar-reduced CE/PV equivalence. Strictness is detected by  $\hbar N[\bar{c}] \neq 0$  in  $\mathcal{B}^{\text{adm}}$  versus  $\hbar N[\bar{c}] = 0$  in  $\mathcal{B}^{\text{red}}$ , witnessed classically by  $\{\text{Tr } \phi_1, \text{Tr } \phi_2\} = N$  and quantum-mechanically by  $[\Phi_h(z_1), \Phi_h(z_2)]_\star = \hbar N \cdot I$ .  $\square$

PROPOSITION 5.7 (*Diagonal local-cohomology chiral  $E_2$ -operad on  $\mathbb{C}^2$* ). Let

$$K_n = H_{\Delta_n}^{2(n-1)}(\mathbb{C}[[z^{(1)}, \dots, z^{(n)}]]) \bigotimes_{i=1}^{n-1} d\zeta_{i,1} d\zeta_{i,2}, \quad \Delta_n = \{z^{(1)} = \dots = z^{(n)}\}, \quad \zeta_i = z^{(i)} - z^{(n)},$$



the arity- $n$  diagonal local-cohomology kernel on  $\mathbb{C}^{2n}$ . The collection  $\{K_n\}_{n \geq 1}$  with residue composition  $\gamma: K_n \otimes K_{m_1} \otimes \cdots \otimes K_{m_n} \rightarrow K_{m_1 + \cdots + m_n}$  defined by pullback of singular principal parts to nested diagonals followed by iterated Grothendieck residues is a coloured dg operad in nuclear Tate vector spaces; it acts on  $\mathcal{A}_{\text{bulk}}^{\text{cl}}$  by

$$\mu_n: K_n \otimes (\mathcal{A}_{\text{bulk}}^{\text{cl}})^{\boxtimes n} \longrightarrow \Delta_{n,*} \mathcal{A}_{\text{bulk}}^{\text{cl}}, \quad \mu_n(K_n; \mathbf{J}_{x_1}, \dots, \mathbf{J}_{x_n}) = K_n \cdot \mathbf{J}_{\ell_n(x_1, \dots, x_n)},$$

with  $\ell_2 = [-, -]_{\mathfrak{g}_{\text{cent}}}$  and the higher  $L_\infty$ -brackets  $\ell_{n \geq 3}$  transferred from the CE factorization structure ( $\ell_{n \geq 3} = 0$  in the strict Hamiltonian BF model, modulo quantum counterterm transfer). The arity-two component is  $K_2 = K_\Delta^{(2)} = d\zeta_1 d\zeta_2 / (\zeta_1 \zeta_2)$  of (17).

*Proof.* Arity-two:  $K_2 = H_\Delta^2(\mathbb{C}[[z, w]]) d\zeta_1 d\zeta_2$  is the residue class of Theorem 5.185. Arity-three associativity: the iterated Grothendieck residue on the nested diagonal  $\Delta_3 \subset \Delta_2 \subset \mathbb{C}^6$  factors as  $\text{Res}_{z^{(2)}=z^{(3)}} \circ \text{Res}_{z^{(1)}=z^{(3)}} = \text{Res}_{z^{(1)}=z^{(2)}=z^{(3)}}$ , matching  $\gamma(K_2 \otimes K_2) = K_3$ ; signs agree with Koszul signs of CE cochains by the standard sign rule for nested residues. Higher arities  $n \geq 4$  inherit the operadic associativity from the factorization-algebra bulk-boundary correspondence [15] applied to disjoint and nested polydisk factorisations of  $\mathbb{C}^{2n}$ ; continuity in  $\mathcal{B}^{\text{adm}}$  is the kernel-admissibility condition (v) of Theorem 5.6. Curve restriction to a smooth  $C \hookrightarrow \mathbb{C}^2$  with transverse residue datum  $B_\perp: \Omega_c^{\circ,1}(\mathbb{C}) \rightarrow \mathbb{C}$  specialises  $K_\Delta^{(2)}$  to the one-dimensional Cauchy kernel  $K_C^{(1)}(z, w) = dw/(w - w')$  by direct residue computation  $\text{Res}_{u=u'} d(w - w')d(u - u')/((w - w')(u - u')) = d(w - w')/(w - w')$ .  $\square$

*Definition 5.8 (Admissible Hochschild-centre lift).* An admissible lift of the closed-to-open coupling  $\Phi_N: \mathcal{A}_{\text{bulk}}^{\text{cl}}|_b \rightarrow Z_{E_1}^{\text{der,adm}}(\mathcal{A}_{\partial,N}^{\text{cl}})$  is a Maurer–Cartan element

$$C_{\Phi_N} = \Phi_N + H^{\text{prod}} + H^{P_0} + H^3 + \cdots$$

in the completed Hochschild-centrality convolution  $L_\infty$ -algebra  $\mathcal{K}_{\mathcal{B}^{\text{adm}}}^{\text{cent}}$  of clause (vi) of Theorem 5.6 satisfying

$$dC_{\Phi_N} + \frac{1}{2}[C_{\Phi_N}, C_{\Phi_N}] + \frac{1}{3!}\ell_3(C_{\Phi_N}^{\otimes 3}) + \cdots = 0,$$

with  $H^{\text{prod}}$  the product centrality homotopy  $[\Phi_N(x), a]_G = dH_{x,a}^{\text{prod}}$ ,  $H^{P_0}$  the Poisson centrality homotopy  $\{\Phi_N(x), a\}_{P_0} = dH_{x,a}^{P_0}$ ,  $H^3$  the order-three Jacobiator/associator  $dH_{x,a,b}^3 = \text{Jac}(H^{\text{prod}}, H^{P_0})$ , and higher  $H^n$  the Stasheff coherences. The  $C_{\Phi_N}$  datum is the difference between an action of closed currents on open observables and a derived  $E_1$ -centre map.

*Remark 5.9 (Consistency probes).* Theorem 5.6 passes the following checks inside  $\mathcal{B}^{\text{adm}}$ .

- (1)  $N = 1$ .  $\phi_i = z_i$ ,  $J(f) = f$ , and  $[\Phi_h(z_1), \Phi_h(z_2)]_\star = \hbar$ ; the projective curvature reduces to  $\hbar[\bar{c}]$ .
- (2) *Finite- $N$  trace bracket.*  $\{\text{Tr } \phi_1, \text{Tr } \phi_2\} = N$ ; vanishing of this datum signals an accidental downgrade to the scalar-reduced shadow.
- (3) *Adjacent-swap exactness.*  $\text{Tr}(u\phi_1\phi_2v) - \text{Tr}(u\phi_2\phi_1v) = Q \text{Tr}(\psi v u)$  (Lemma 3.30).
- (4) *Monomial bracket.*  $\{J(z_1^a z_2^b), J(z_1^c z_2^d)\} = (ad - bc)J(z_1^{a+c-1} z_2^{b+d-1}) + N[\{f, g\}]_0$ .
- (5) *CE/PV differential.* Under  $c^k \mapsto \partial_{\xi_k} u_k \mapsto \xi_k$ ,  $d_{\text{CE}} c^k = -\frac{1}{2} f_{ij}^k c^i c^j$  matches  $d_\pi(\partial_{\xi_k})$  and  $d_{\text{CE}} u_k = f_{ik}^j u_j c^i$  matches  $d_\pi(\xi_k)$  (Theorem 4.20).
- (6) *Köthe bracket continuity.*  $p_n(\{x, y\}) \leq C_n p_{n+2}(x) p_{n+2}(y)$  in the weighted seminorms of clause (iv).
- (7) *Kernel Maurer–Cartan.*  $d\Theta_B + \frac{1}{2}[\Theta_B, \Theta_B] = 0$  (Theorem 5.5).

- (8) *Hochschild centrality*.  $[\Phi_N(x), a]_G = dH_{x,a}^{\text{prod}}$  and  $\{\Phi_N(x), a\}_{P_0} = dH_{x,a}^{P_0}$  for every closed generator  $x$  and open observable  $a$  (Definition 5.8).
- (9) *QME recursion*. At order  $\hbar^n$  and window  $M$ ,  $d_M C_{n,M} = -o_{n,M}^{\text{ns}}$  and  $\pi_{M,N} C_{n,M} = C_{n,N}$ .
- (10) *Global descent first obstruction*.  $[d_2] \in H^2(\partial X^{\text{KN}}, \underline{\text{Hom}}^{-1}(\mathcal{H}^\bullet, \mathcal{H}^\bullet))$  from Theorem 5.13.
- (11) *Curve restriction*.  $R_{L,B_\perp}(K_\Delta^{(2)}) = dw/(w - w')$  (Proposition 5.7).

THEOREM 5.10 (*Hamiltonian trace generators exhaust the admissible  $E_1$ -Hochschild centre*). Inside  $\mathcal{B}^{\text{adm}}$ ,

$$HH_{\text{adm}}^\bullet(A_{\partial,N}^{\text{cl}}, A_{\partial,N}^{\text{cl}}) = \langle J(f), \theta_f, K : f \in \mathfrak{h} \rangle_{\sim, \{-, -\}},$$

the graded-commutative algebra generated under cup product and Gerstenhaber bracket by the trace coordinates  $J(f) = \text{Tr } f(\phi_1, \phi_2)$ , their Schouten derivations  $\theta_f$ , and the trace- $U(1)$  centre  $K$ . The scalar-reduced shadow at  $K = 0$  gives the identity for  $HH_{\text{adm}}^\bullet(A_{\partial,N}^{\text{cl,red}}, A_{\partial,N}^{\text{cl,red}})$ .

*Proof.* Continuous Hochschild–Kostant–Rosenberg (Theorem 5.30; Theorem 5.23) identifies  $HH_{\text{adm}}^\bullet(A_{\partial,N}^{\text{cl}}, A_{\partial,N}^{\text{cl}})$  with  $PV_{\text{adm}}^\bullet(\text{Sym } \mathfrak{h})^{GL_N}$  in  $\mathcal{B}^{\text{adm}}$ . Procesi–Razmyslov stable trace invariant theory [17, 18] (Proposition 3.22) generates the  $GL_N$ -invariant polynomial functions on  $\mathfrak{gl}_N \oplus \mathfrak{gl}_N$  by the cyclic trace words  $\text{Tr } f(\phi_1, \phi_2)$ ,  $f \in \mathbb{C}[z_1, z_2]$ ; Koszul resolution by  $\psi$  with  $Q\psi = [\phi_1, \phi_2]$  extends this generation to  $\mathcal{O}(\mu_{\text{der}}^{-1}(0))^{GL_N}$ . The polyvector algebra  $PV_{\text{adm}}^\bullet(\text{Sym } \mathfrak{h})^{GL_N}$  is generated as a graded-commutative algebra over this function ring by the Schouten derivatives  $\theta_f$  along the symplectic structure on  $\widehat{\mathbb{C}}^2$ , and the projective Capelli completion adjoins  $K$  with the cocycle  $[(f, aK), (g, bK)] = (\{f, g\}, \hbar N \omega(f, g) K)$ . The bracket-admissible Hamiltonian-spanning image of Theorem 5.5 contains every  $J(f)$ , every  $\theta_f$ , and  $K$ , and is closed under cup product and Gerstenhaber bracket.  $\square$

PROPOSITION 5.11 (*Monotone strength tower of the formal-Darboux Deligne theorem*). The local theorem sits in a tower of theorem packages  $T = (C, \mathcal{A}_{\text{bulk}}^{\text{cl}}, A_\partial^{\text{cl}}, Z, \Phi, O, \mathcal{H})$  connected by conservative forgetful functors

$$T_{\text{compact}} \xrightarrow{U_{\text{target}}} T_{\text{KN,global}} \xrightarrow{\text{Stalk}_b} T_{\text{adm,Capelli}} \xrightarrow{H_Q} T_{P_0^\beta} \xrightarrow{\text{gr}_{\hbar,F}} T_{\text{coord}} \xrightarrow{U_B} T_{\text{red}}.$$

Each arrow is forgetful downward and structure-adding upward; every  $T_{i+1} \Rightarrow T_i$ . Strictness is witnessed at each rung by data destroyed under the corresponding forgetful functor: the Capelli projective curvature  $\hbar N[\bar{c}]$ ; the pro-Matlis residue tower  $\mathcal{P}_q^{\text{II}} = \lim P_{\leq N}$ ; the diagonal local-cohomology kernel  $K_\Delta^{(2)}$ ; the Hochschild centrality homotopy hierarchy  $\{\overline{h}_{\alpha,a}, h_{\alpha,a,b}, \dots\}$ ; the all-loop QME counterterm tower  $\{C_{n,M}\} \in \mathfrak{Q}_{w,\partial,\hbar}^\bullet$ . Theorem 5.6 is the rung  $T_{\text{adm,Capelli}}$ .

*Proof.*  $U_{\text{target}}$  forgets compact-target comparison data of Theorem 10.13.  $\text{Stalk}_b$  restricts to the formal Darboux neighbourhood: the descent class  $\text{Ob}_{\text{KN}}^{\text{desc}}$  vanishes on a contractible polydisk, and the global chiral  $E_2$ -Deligne datum restricts to the arity-two kernel  $K_\Delta^{(2)}$ .  $H_Q$  takes cohomology of the QME differential, retaining the induced projective  $E_2$ -central action on  $H^\bullet$ .  $\text{gr}_{\hbar,F}$  takes the associated graded for the  $\hbar$ -adic and scalar-contact filtrations, turning the projective quantum centre into its classical  $P_0$ -shadow with the cocycle as leading scalar class.  $U_B$  forgets the topology, convergence, and kernel admissibility of  $\mathcal{B}^{\text{adm}}$ , retaining the coordinate CE/PV identity  $c_f \mapsto \theta_f, u_f \mapsto O_f$ . The terminal arrow to  $T_{\text{red}}$  quotients constants, killing  $K$  and the pro-Matlis tail. Each kernel is detected by a named strictness witness; conservativity follows from injectivity on the witness classes.  $\square$

*Remark 5.12 (Admissible CE/PV criterion and specialization).* The CE/PV cochain identity in 5.5 is the full formal-disk statement in the reduced Hamiltonian sector. Its enveloping/Koszul enhancement is the admissible criterion: one must supply the coefficient category in which CE chains are conilpotent, continuous duality is exact, and the filtered cobar spectral sequence converges. Nilpotent, relative, and weighted Tate replacements are mechanisms for satisfying that criterion; the strict product/direct-sum endpoint is governed by the obstruction datum. The other proved statements are dimension-two specific (they use the residue pairing on  $\mathbb{C}^2$ , the matrix Weyl algebra of pairs  $(X, Y)$ , and the  $U(1)$  centre of  $\mathfrak{gl}_N$ ). Higher-dimensional analogues require the same cochain identity together with the appropriate residue pairing, matrix Weyl data, and a separate verification of the required conilpotent, relative, weighted, or completed coefficient hypotheses.

The admissible category  $\mathcal{B}^{\text{adm}}$  absorbs the local-to-strict obstructions structurally. The factorization-centre lift datum  $\text{Ob}_{p_0}^{\text{fact}} \in H^1(\text{Cone}(\text{gr } \kappa_\tau)) \oplus \varprojlim^1 H^0$  of Problem 5.48 (strict Casimir support, the low-degree term  $\{z_1, z_2^{N+1}\} = (N+1)z_2^N$ , the  $H_2 \cong \mathfrak{sl}_2$  lower-central obstruction, and the Milnor  $\varprojlim^1$  defect) is converted into the bracket-admissible kernel image of Theorem 5.5. The non-scalar QME class  $[\text{Ob}_{w,\partial,h}^{\text{red}}] \in H^1(\ker \widehat{\sigma}_{\text{sc}}, d_{\text{QME}})$  of Problem 5.219, with first exposed row  $\theta_3$  and one-row covector  $\ell_{\theta_3}(\mathbf{E}_{\theta_3, \mathcal{M}}) = 1$ , is killed by the Costello renormalisation counterterm package. The radial PBW Stokes class  $[(T_{a,b}, E_{a,b}^+)] \in \text{coker } B_{a,b}$  of Proposition F.25, equivalently the decorated Stokes obstruction for  $D_{a,b}^\square = C_{a,b}^+ \partial_2$ , is absorbed by the contracting homotopies on the pro-Matlis cotangent target. The six-entry Bochner–Martinelli–Koppelman vector  $(\text{ob}_\oplus, \text{ob}_{I_\Pi}, \text{ob}_{\mathfrak{h}, \Pi}, \text{ob}_{\text{collar}, \Pi}, \text{ob}_{3, \Pi}, \text{ob}_{\text{unif}, \Pi})$  of Proposition 1.26 is absorbed by the pro-Matlis principal-part current target. The trace- $U(1)$  scalar anomaly  $\hbar N[\bar{c}]$  is absorbed by the projective Capelli central extension  $\widehat{\mathfrak{g}}_{\text{cent}}$  and reappears in  $\mathcal{B}^{\text{adm}}$  as the central charge of the modular line bundle at the trace generator.

The Hamiltonian on the formal symplectic normal disk is evaluated on the brane’s matrix-valued transverse coordinate by trace; Dirac reduction forces the commuting-matrix algebra; the shifted-cotangent BF Lie algebra  $\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1]$  separates canonical motion from evaluation; the Chevalley–Eilenberg/polyvector dictionary sends Hamiltonian ghosts  $c_f$  to Schouten derivations  $\theta_f$  and cotangent coordinates  $u_f$  to brane observables  $J(f)$ ; continuous duality realises cotangent modes as Matlis principal parts; the projective central charge  $\hbar N[\bar{c}]$  is the curvature of the modular line bundle at the trace generator.

**THEOREM 5.13 (Mixed Holomorphic–Topological Deligne identification on the Kato–Nakayama log boundary).** Let  $(X, D, \tau)$  be a holomorphic-symplectic log surface with Kato–Nakayama log boundary  $\partial X^{\text{KN}}$ , with bulk mixed-HT factorization algebra  $\mathcal{A}_{\text{bulk}}^{\text{cl}}$  of Definition 1.19 and open boundary factorisation category  $\mathcal{C}_\partial^{\text{op}}$  of Dirac branes on  $\partial X^{\text{KN}}$ . The global identification

$$\mathcal{A}_{\text{bulk}}^{\text{cl}} \simeq Z_{E_2^{\text{HT}}}^{\text{der}}(\mathcal{C}_\partial^{\text{op}})$$

as mixed-HT chiral algebras on  $\partial X^{\text{KN}}$  holds if and only if the three-component global obstruction

$$\text{Ob}^{\text{global}} = \text{Ob}_{\text{Deligne}}^{\text{ch } E_2} \oplus \text{Ob}_{\text{KN}}^{\text{desc}} \oplus \text{Ob}_{\text{all-loop}}^{\text{QME}}$$

vanishes, where:

- $\text{Ob}_{\text{Deligne}}^{\text{ch } E_2}$  is the obstruction to extending the arity-two diagonal product  $K_\Delta^{(2)} = d\zeta_1 d\zeta_2 / (\zeta_1 \zeta_2)$  to a chiral  $E_2$ -operad with operadic associativity on all arities — the chiral  $E_d$ -Deligne conjecture at  $d = 2$  (Conjecture 10.1);

- $\text{Ob}_{\text{KN}}^{\text{desc}} = [d_2] \in H^2(\partial X^{\text{KN}}, \underline{\text{Hom}}^{-1}(\mathcal{H}^\bullet, \mathcal{H}^\bullet))$  is the  $d_2$ -differential of the descent spectral sequence on the Kato–Nakayama log-étale cover trivialising the local holomorphic-symplectic Darboux data, with  $\mathcal{H}^\bullet$  the local-stalk sheaf assigned by Theorem 5.6;
- $\text{Ob}_{\text{all-loop}}^{\text{QME}}$  is the all-loop Quantum Master Equation obstruction class of the closed-to-open coupling  $\Phi_N$  at every Costello renormalisation scale, in the renormalisation-group filtered complex supplied by the Costello–Li package of Statement 5.207 and Statement 5.206.

If the operator-level lift of the modular line bundle on the compact target  $K_3 \times E$  is included, vanishing of the additional modular-line obstruction  $\text{Ob}_{\text{mod}}^V$  to the operator package of  $\Omega_{\text{central}}$  is required; its scalar shadow is the Igusa cusp form  $\Phi_{10}$ , with Borcherds square root  $\Delta_5 = \Phi_{10}^{1/2}$ .

*Remark 5.14 (Local trivialisation of the global obstruction).* Restricted to a formal-Darboux neighbourhood of a brane vacuum  $b \in X$ , every component of  $\text{Ob}^{\text{global}}$  vanishes: the chiral  $E_2$ -Deligne datum is supplied by the two-variable Cauchy kernel  $K_\Delta^{(2)}$ ; the descent class is identically zero on a contractible polydisk; the all-loop QME class collapses to the one-loop centrality datum absorbed by Theorem 6.8. The content of  $\text{Ob}^{\text{global}}$  is the global gluing of these locally trivialised classes along  $\partial X^{\text{KN}}$ .

#### 5.4 Tate Quillen equivalence and the $\infty$ -categorical upgrade of the cochain bar-cobar identity

Inside the admissible Tate envelope of Statement 5.17, the bar-cobar adjunction  $\Omega \dashv \text{Bar}$  is a Quillen equivalence (Theorem 5.30), and the CE/PV map  $c_f \mapsto \theta_f$ ,  $u_f \mapsto J(f)$  descends to a quasi-isomorphism on cohomology between the closed CE cochain algebra and the Schouten polyvector algebra of the boundary Hamiltonian polynomial algebra. This is the Loday–Vallette / Lurie Koszul-resolution identification of the brane-side derived  $E_1$ -Hochschild  $HH^\bullet(A_{\partial,N}^{\text{cl}}, A_{\partial,N}^{\text{cl}})$ , identified with the bulk stalk  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b$  by Theorem 5.6; after restriction to a holomorphic curve, the  $d = 1$  matching to the bulk side is the chiral-algebra theorem of [28]. Its arity-two holomorphic collision is the two-dimensional diagonal singular product of Theorem 1.15; the Tate Quillen equivalence controls the bar-cobar/Hochschild algebraic realization of that stalk:  $\Omega C_*^{\text{CE}}(\mathfrak{g}) \xrightarrow{\sim} \widehat{U}(\mathfrak{g})$  on the Lie side, dualized through exact continuous duality to

$$C_{\text{CE,cont}}^\bullet(\mathfrak{g}) \simeq A_{\text{op}}^!,$$

with the linear symbol  $B_f \mapsto O_f$ ,  $\Theta_{\rho_{a,b}} \mapsto \theta^{a,b}$  of Definition 5.21.

For zero-differential finite-dimensional pieces  $\{V_d\}_{d \geq 1}$ , the inclusion

$$V_\oplus = \bigoplus_{d \geq 1} V_d \hookrightarrow V_\Pi = \prod_{d \geq 1} V_d,$$

filtered by degree tails, is an isomorphism on every finite quotient and on associated graded, yet  $H^0(V_\Pi/V_\oplus) = V_\Pi/V_\oplus \neq 0$  (Lemma 5.26). The tail-quotient class is invisible to admissible filtered weak equivalence and constitutes one summand of the Koszul obstruction datum (the five-summand obstruction complex collecting the window, topology, conilpotence, Maurer–Cartan, and cobar obstruction components for the twist  $\tau$ ):  $O_{\text{Kos}} = O_{\text{win}} \oplus O_{\text{top}} \oplus O_{\text{conilp}} \oplus O_{\text{MC}}(\tau) \oplus O_{\text{cobar}}(\tau)$ , of Definition 5.22. The recognition criterion

$$\omega_{\text{win}} = 0, \quad O_{\text{conilp}} = 0, \quad O_{\text{MC}}(\tau) = 0, \quad H^\bullet(O_{\text{top}} \oplus O_{\text{cobar}}(\tau)) = 0$$

is necessary and sufficient for filtered-cobar admissibility (Theorem 5.23); equivalently, it is the precise vanishing condition for the CE/PV map to descend to a quasi-isomorphism of cohomology of the chiral

Hochschild target. Theorem 5.30 descends to an equivalence of associated  $\infty$ -categories (Corollary 5.31). The Weiss-cosheaf condition on the unreduced factorization algebra over  $\mathbb{R}^2 \times \mathbb{C}^2$  remains separate (Proposition 5.38); it requires the brane-defect heat-kernel propagator extension of Problem 5.48 item (2).

#### 5.4.1 THE TATE CHAIN CATEGORY

Let **TateVec** be the closed symmetric monoidal category of complete filtered Tate  $\mathbb{C}$ -vector spaces of 5.165. The filtration on every object is taken *complete and Hausdorff*:  $V = \varprojlim_w V_{\leq w}$  and  $\bigcap_w F^w V = 0$ . Concrete morphisms are filtration-preserving continuous chain maps. The working ambient category is the bounded-below filtered subcategory **TateChain** $_{\geq 0}$  of non-negatively filtered chain complexes in **TateVec**, large enough to contain every CE chain coalgebra and completed enveloping algebra used here. The exact finite-window Mittag-Leffler category is constructed first. The model-categorical statements add transfer, smallness, and monoid axioms to this exact category; their scope is the exact category constructed here.

*Definition 5.15 (Strict finite-window Mittag-Leffler Tate category).* Let **MLTateChain** $_{\geq 0}^{\text{str}}$  be the category whose objects are inverse systems

$$V = (V_{\leq M}, \pi_{M,N}^V)_{M \geq N \geq 1}$$

of bounded-below finite-dimensional chain complexes over  $\mathbb{C}$ , with degreewise surjective transition maps. The represented complete object is

$$|V| = \varprojlim_M V_{\leq M}, \quad F^M |V| = \ker(|V| \rightarrow V_{\leq M-1}).$$

Morphisms are compatible families of chain maps  $f_{\leq M}: V_{\leq M} \rightarrow W_{\leq M}$ . The completed tensor product is defined on paired windows by

$$V \widehat{\otimes}_{\text{ML}} W = (V_{\leq M} \otimes W_{\leq M})_{M \geq 1},$$

with the induced surjective transition maps, and

$$|V \widehat{\otimes}_{\text{ML}} W| = \varprojlim_M (V_{\leq M} \otimes W_{\leq M}).$$

The continuous dual attached to this strict tower is

$$V_{\text{ML}}^\vee = \varinjlim_M V_{\leq M}^\vee.$$

A morphism is an admissible filtered weak equivalence when each  $f_{\leq M}$  and the induced associated-graded map are quasi-isomorphisms.

The only universal property asserted for this category is the ordinary inverse-limit one: a compatible family  $x_M \in V_{\leq M}$  determines a unique element  $x \in |V|$ , and every map out of a test strict tower is exactly a compatible family of finite-window maps. Initial and terminal properties among all completions of a Tate vector space lie outside this assertion.

**PROPOSITION 5.16 (Exactness and kernels in the strict ML category).** In **MLTateChain** $_{\geq 0}^{\text{str}}$ :

- (i) the filtration on  $|V|$  is complete and Hausdorff;

- (ii) inverse limits of strict surjective towers are exact, so  $H^\bullet(|V|) = \varprojlim_M H^\bullet(V_{\leq M})$  and  $\varprojlim_M^1 H^{\bullet-1}(V_{\leq M}) = 0$ ;
- (iii) completed tensor products and continuous duals are computed by the displayed finite-window formulas;
- (iv) if  $H = (H_{\leq M})$  and  $D = (D_{\leq M})$  are paired strict ML systems and finite tensors  $C_M \in H_{\leq M} \otimes D_{\leq M}$  are compatible under transition, then

$$C = \varprojlim_M C_M \in |H| \widehat{\otimes}_{\text{ML}} |D|$$

exists and is the unique strict ML kernel with those finite-window projections.

*Proof.* Completeness is the definition of  $|V|$  as an inverse limit; the intersection of the kernels  $F^M|V|$  is zero because an element whose image vanishes in every finite quotient is the zero compatible family. Surjectivity of the transition maps gives the Mittag-Leffler condition. The Milnor exact sequence then has zero  $\varprojlim^1$ -term, proving exactness and the cohomology formula. The tensor and dual formulas are definitions inside the strict category and are finite-dimensional on each window. Finally, a compatible tensor family  $(C_M)_M$  is precisely an element of the inverse limit  $\varprojlim_M (H_{\leq M} \otimes D_{\leq M})$ , which is the completed tensor product by definition. The uniqueness is the inverse-limit universal property of Definition 5.15.  $\square$

*Statement 5.17 (Admissible Tate model envelope).* An admissible Tate model envelope is a bounded-below, locally presentable symmetric monoidal model category  $\mathbf{TateChain}_{\geq 0}^{\text{adm}}$  equipped with a fully faithful exact symmetric-monoidal embedding of the strict category

$$\mathbf{MLTateChain}_{\geq 0}^{\text{str}} \hookrightarrow \mathbf{TateChain}_{\geq 0}^{\text{adm}},$$

and containing the CE chains, completed enveloping algebras, completed symmetric algebras, and finite-window inverse systems used below. Weak equivalences are admissible filtered quasi-isomorphisms: quasi-isomorphisms on every finite quotient and on associated graded. The tensor product restricts to  $\widehat{\otimes}_{\text{ML}}$  on strict Mittag-Leffler systems, and filtered inverse limits with surjective transition maps are exact by Proposition 5.16. The additional hypotheses are transfer and smallness for the bar-cobar adjunction, the monoid axiom for completed tensor products, and compatibility of  $\widehat{\otimes}$  with the finite-window model structures. All Quillen statements below are relative to such an envelope.

**LEMMA 5.18 (Mittag-Leffler dictionary for closed Tate windows).** Let

$$\{V_{\leq M}\}_{M \geq 1} \quad \text{and} \quad \{W_{\leq M}\}_{M \geq 1}$$

be inverse systems of finite-dimensional complexes with surjective transition maps, and put

$$V = \varprojlim_M V_{\leq M}, \quad W = \varprojlim_M W_{\leq M}.$$

In the strict ML category, and hence in any admissible envelope containing it, the following assertions hold:

- (i)  $\varprojlim$  is exact on these systems, so  $H^\bullet(V) = \varprojlim_M H^\bullet(V_{\leq M})$ ;
- (ii) continuous dualization changes the inverse system into the direct system of finite duals:  $V_{\text{cont}}^\vee = \varinjlim_M V_{\leq M}^\vee$ ;

- (iii) completed tensor products are computed windowwise:  $V \widehat{\otimes} W = \varprojlim_M (V_{\leq M} \otimes W_{\leq M})$  whenever the transition maps are the paired finite-window truncations;
- (iv) a map  $f: V \rightarrow W$  is an admissible filtered quasi-isomorphism iff each finite-window map  $f_{\leq M}: V_{\leq M} \rightarrow W_{\leq M}$  is a quasi-isomorphism and the induced associated-graded maps are quasi-isomorphisms.

*Proof.* Surjectivity of the transition maps gives the Mittag-Leffler condition, hence  $\varprojlim^1 = 0$ . The cohomology assertion is the standard Milnor exact sequence with the  $\varprojlim^1$ -term killed. Continuous linear functionals on  $V$  factor through a finite quotient  $V_{\leq M}$ , giving the displayed direct limit of duals. The completed tensor product in Definition 5.15 is defined by the same finite-window inverse system, so item (iii) is tautological inside the strict category. Item (iv) is the definition of weak equivalence in Definition 5.15 and Statement 5.17.  $\square$

*Definition 5.19 (Closed-window obstruction terms).* Let  $\mathfrak{h} = \varprojlim_M \mathfrak{h}_{\leq M}$  be a complete filtered Lie algebra with finite-dimensional windows and surjective transition maps  $\pi_{M,N}$ . Write  $D_{\leq M}$  for a finite dual module paired with  $\mathfrak{h}_{\leq M}$ , with surjective transition maps  $\pi_{M,N}^\vee$ , and put  $D_{\text{cont}} = \varprojlim_M D_{\leq M}$ . For  $M \geq N$  define:

$$\begin{aligned} \mathfrak{o}_{M,N}^{\text{br}}(x, y) &= \pi_{M,N}[x, y]_M - [\pi_{M,N}x, \pi_{M,N}y]_N, \\ \mathfrak{o}_{M,N}^{\text{coad}}(x, \lambda) &= \pi_{M,N}^\vee(x \cdot_M \lambda) - (\pi_{M,N}x) \cdot_N (\pi_{M,N}^\vee \lambda), \end{aligned}$$

and

$$\mathfrak{o}_{M,N}^{\text{Cas}} = (\pi_{M,N} \otimes \pi_{M,N}^\vee)C_M - C_N.$$

The closed window system is *cochain-CE/PV-admissible* if  $\mathfrak{o}^{\text{br}}$  and  $\mathfrak{o}^{\text{coad}}$  vanish for all  $M \geq N$ . It is *kernel-admissible* only after the additional Casimir term  $\mathfrak{o}^{\text{Cas}}$  vanishes in the chosen completed tensor category. The coordinate CE/PV cochain identity contains the bracket and coadjoint transition equations; kernel-admissibility adds the Casimir equation.

**LEMMA 5.20 (Closed CE/PV finite-window compatibility and variance boundary).** Assume the closed window system of Definition 5.19 is cochain-CE/PV-admissible and each finite window has the finite cotangent CE/PV identity

$$\Phi_M: C_{\text{CE}}^\bullet(\mathfrak{h}_{\leq M} \ltimes D_{\leq M}[1]) \xrightarrow{\sim} \text{PV}(\mathbf{S}(\mathfrak{h}_{\leq M})).$$

Then the  $\Phi_M$  are compatible as finite coordinate tests under the contravariant CE pullbacks and the corresponding PV coordinate maps. These maps form compatible finite coordinate tests. A completed admissible filtered quasi-isomorphism

$$\Phi: C_{\text{CE,cont}}^\bullet(\mathfrak{h} \ltimes D_{\text{cont}}[1]) \xrightarrow{\sim} \text{PV}(\widehat{\mathbf{S}}_{\text{Tate}}(\mathfrak{h}))$$

follows only after one supplies either a chain-side inverse limit with exact continuous dualization or a topology whose cochain transition maps are dg maps. If the kernel-admissible Casimir condition is also supplied, the same finite windows carry compatible  $P_\circ$ -pairings and kernels in that chosen tensor category.

*Proof.* The finite map  $\Phi_M$  is determined on generators by  $c \mapsto \theta$  and  $u \mapsto O$ . Vanishing of  $\mathfrak{o}_{M,N}^{\text{br}}$  says the CE differential on the Hamiltonian ghosts commutes with transition maps. Vanishing of  $\mathfrak{o}_{M,N}^{\text{coad}}$  says the cotangent-coordinate part of the CE differential, and equivalently the Lichnerowicz differential

on  $O$ -coordinates, commutes with transition maps. Therefore the finite squares with  $\Phi_M$  and  $\Phi_N$  commute in the variance appropriate to cochains. A quotient  $\mathfrak{h}_{\leq M} \twoheadrightarrow \mathfrak{h}_{\leq N}$  induces CE cochain maps by pullback along the quotient. Thus inverse-limit language on cochains requires the additional continuous-dualization or topology hypothesis stated above. Vanishing of  $\mathfrak{o}_{M,N}^{\text{Cas}}$  is the separate statement that the finite  $P_o$ -brackets and BV coefficient pairings are compatible under transition in the chosen kernel-admissible tensor category.  $\square$

#### 5.4.2 TWISTING COCHAINS AND THE FILTERED-COBAR ADMISSIBILITY CRITERION

*Definition 5.21 (Open-side twisting cochain datum).* Fix a candidate admissible filtered-cobar datum

$$\mathfrak{h} = \varprojlim_M \mathfrak{h}_{\leq M}, \quad D = \varinjlim_M \mathfrak{h}_{\leq M}^\vee, \quad \mathfrak{g} = \mathfrak{h} \ltimes D[1],$$

a complete augmented filtered open  $\mathcal{A}_\infty$ -algebra  $\mathcal{A}_{\text{op}}$ , and the completed CE/PV target

$$B_{\text{cl}} = C_{\text{CE,cont}}^\bullet(\mathfrak{g}) \cong \text{PV}(\widehat{\mathbf{S}}_{\text{coord}}(\mathfrak{h})).$$

Put  $C_{\text{op}} = \text{Bar}(\mathcal{A}_{\text{op}})$ , with reduced bar differential  $b_{\text{op}}$ , coproduct  $\Delta$ , and iterated reduced coproducts  $\Delta^{(r)}$ . The bar coalgebra is required to be locally conilpotent in the completed sense: for every finite output window of  $B_{\text{cl}}$  and every finite coaugmentation stage of  $C_{\text{op}}$ , the iterated reduced coproducts  $\Delta^{(r)}$  vanish for all sufficiently large  $r$ . Equivalently, the conilpotence obstruction in Definition 5.22 is zero. This condition makes the Maurer–Cartan curvature below a continuous map and supplies the twisting-cochain datum.

An open-side twisting cochain is a continuous degree-one map

$$\tau: C_{\text{op}} \longrightarrow B_{\text{cl}}$$

vanishing on the coaugmentation and satisfying the Maurer–Cartan equation

$$d_{B_{\text{cl}}} \tau + \tau b_{\text{op}} + \sum_{r \geq 2} m_r^{B_{\text{cl}}}(\tau^{\otimes r}) \Delta^{(r)} = 0.$$

For the dg  $P_o$ -algebra  $B_{\text{cl}}$ , the higher multiplications vanish and this reads

$$d_{B_{\text{cl}}} \tau + \tau b_{\text{op}} + \mu_{B_{\text{cl}}}(\tau \otimes \tau) \Delta = 0.$$

Equivalently,

$$d_{\text{conv}} \tau + \frac{1}{2} [\tau, \tau]_{\text{conv}} = 0, \quad d_{\text{conv}} f = d_{B_{\text{cl}}} f - (-1)^{|f|} f b_{\text{op}},$$

in the continuous convolution dg Lie algebra  $\text{Hom}_{\text{cont}}(C_{\text{op}}, B_{\text{cl}})$ .

The twisting cochain induces the completed cobar morphism

$$\kappa_\tau: \Omega C_{\text{op}} \longrightarrow B_{\text{cl}}.$$

Its associated-graded symbol is the map

$$\text{gr } \kappa_\tau: \text{gr } \Omega \text{Bar}(\mathcal{A}_{\text{op}}) \longrightarrow \text{gr } B_{\text{cl}}.$$

In the Hamiltonian trace presentation the required linear symbol is

$$\text{gr } \kappa_\tau(B_f) = O_f, \quad \text{gr } \kappa_\tau(\Theta_{\rho_{a,b}}) = \theta^{a,b},$$

after the scalar trace is removed and the cotangent generator is read in the principal-part local-dual basis.



*Definition 5.22 (Open-side Koszul obstruction complex).* In the notation of Definition 5.21, put  $D_{\leq M} = \mathfrak{h}_{\leq M}^\vee$ ,  $\text{PV}_{\leq M} = \text{PV}(\mathbf{S}(\mathfrak{h}_{\leq M}))$ , and  $B_{\text{cl}, \leq M}$  for the corresponding finite CE/PV window. The obstruction datum is

$$\mathcal{O}_{\text{Kos}}(\mathfrak{h}, D, \mathcal{A}_{\text{op}}, \tau) = \mathcal{O}_{\text{win}} \oplus \mathcal{O}_{\text{top}} \oplus \mathcal{O}_{\text{conilp}} \oplus \mathcal{O}_{\text{MC}}(\tau) \oplus \mathcal{O}_{\text{cobar}}(\tau).$$

The window and conilpotence summands are degree-zero:

$$\mathcal{O}_{\text{win}} = \prod_{M \geq N} (\text{Hom}(\Lambda^2 \mathfrak{h}_{\leq M}, \mathfrak{h}_{\leq N}) \oplus \text{Hom}(\mathfrak{h}_{\leq M} \otimes D_{\leq M}, D_{\leq N})) \oplus Q_{\text{Cas}},$$

where

$$Q_{\text{Cas}} = \frac{\varprojlim_M (\mathfrak{h}_{\leq M} \otimes D_{\leq M})}{\text{im}(\widehat{\mathfrak{h} \otimes D})}.$$

The distinguished window class is

$$\omega_{\text{win}} = (\mathfrak{o}_{M,N}^{\text{br}}, \mathfrak{o}_{M,N}^{\text{coad}}, [(C_{\leq M})_M]),$$

where the first two terms are those of Definition 5.19 and the last term is the compatible finite-window Casimir family modulo strict completed tensors. The topology summand is

$$\mathcal{O}_{\text{top}}^n = \varprojlim_M^1 H^{n-1}(C_{\text{CE}}^\bullet(\mathfrak{g}_{\leq M})) \oplus \varprojlim_M^1 H^{n-1}(\text{PV}_{\leq M}).$$

The conilpotence summand is

$$\mathcal{O}_{\text{conilp}} = \left( C_*^{\text{CE}}(\mathfrak{g}) / \bigcup_r F_{\text{coaug}}^{(r)} C_*^{\text{CE}}(\mathfrak{g}) \right) \oplus \bigcap_{r \geq 1} \mathfrak{g}^{(r)}.$$

The Maurer–Cartan summand is the explicit curvature

$$\mathcal{O}_{\text{MC}}(\tau) = \text{MC}(\tau) = d_{\text{conv}} \tau + \frac{1}{2} [\tau, \tau]_{\text{conv}} \in \text{Hom}_{\text{cont}}^2(C_{\text{op}}, B_{\text{cl}}).$$

The cobar obstruction complex is

$$\mathcal{O}_{\text{cobar}}(\tau) = \text{Cone}(\text{gr } \kappa_\tau) \oplus \bigoplus_n \varprojlim_M^1 H^{n-1}((\Omega C_{\text{op}})_{\leq M})[-n] \oplus \bigoplus_n \varprojlim_M^1 H^{n-1}(B_{\text{cl}, \leq M})[-n].$$

**THEOREM 5.23 (Filtered-cobar admissibility criterion).** Fix a candidate datum as in Definitions 5.21 and 5.22. Assume the filtrations on  $\Omega C_{\text{op}}$  and  $B_{\text{cl}}$  are bounded below, complete, separated, and strongly convergent, with finite-dimensional associated graded pieces in every total degree. Assume moreover that the local conilpotence condition in Definition 5.21 holds and continuous dualization is exact on the finite-window towers. Then the following conditions are equivalent.

- (A1) The datum is accepted: the generator substitution  $c \mapsto \theta$ ,  $u \mapsto O$  is the completed CE/PV identity,  $\tau$  is a twisting cochain, and

$$\kappa_\tau: \Omega \text{Bar}(\mathcal{A}_{\text{op}}) \longrightarrow B_{\text{cl}}$$

is an admissible filtered quasi-isomorphism with the linear symbol of Definition 5.21.

(A2) The obstruction equations are exact:

$$\omega_{\text{win}} = 0, \quad \mathcal{O}_{\text{conilp}} = 0, \quad \mathcal{O}_{\text{MC}}(\tau) = 0, \quad H^\bullet(\mathcal{O}_{\text{top}} \oplus \mathcal{O}_{\text{cobar}}(\tau)) = 0.$$

When these equivalent conditions hold,

$$B_{\text{cl}} \simeq A_{\text{op}}$$

in the homotopy category of complete augmented filtered  $A_\infty$ -algebras. If the CE coalgebra is also used as the closed Koszul source, exact continuous duality identifies the dual form as

$$C_{\text{CE,cont}}^\bullet(\mathfrak{g}) \simeq A_{\text{op}}^!$$

precisely when the CE-side cobar map  $\Omega C_*^{\text{CE}}(\mathfrak{g}) \rightarrow A_{\text{op}}$  has the same exact associated-graded cone and  $\varprojlim^1$  obstruction.

*Proof.* The equality  $\omega_{\text{win}} = 0$  is equivalent to strict finite-window compatibility of the CE differential and the Schouten differential, with  $P_0$ -pairings added only in the kernel-admissible tensor category; this is exactly Lemma 5.20 and Theorem 4.22. The vanishing of  $\mathcal{O}_{\text{top}}$  is the Milnor and variance condition identifying cohomology of the completed CE/PV objects with the admissible finite-window calculation.

The local conilpotence hypothesis makes the Maurer–Cartan sum finite on each finite output window, hence a continuous curvature. The equation  $\mathcal{O}_{\text{MC}}(\tau) = 0$  is then the Maurer–Cartan equation in the continuous convolution dg Lie algebra. It is equivalent to the statement that the adjoint map  $\kappa_\tau$  commutes with the completed cobar differential and the  $A_\infty$ -operations. This equality supplies the morphism tested below.

After the Maurer–Cartan equation holds, filter  $\Omega C_{\text{op}}$  and  $B_{\text{cl}}$  by tensor length and by the finite coordinate windows. The mapping-cone summand in  $\mathcal{O}_{\text{cobar}}(\tau)$  is acyclic exactly when  $\text{gr } \kappa_\tau$  is a quasi-isomorphism. The two  $\varprojlim^1$ -summands are exactly the Milnor defects for the completed source and target towers. Strong convergence and completeness then make the spectral-sequence comparison theorem equivalent to admissible filtered quasi-isomorphism of  $\kappa_\tau$ . Necessity follows from the definition of admissible filtered quasi-isomorphism in the envelope: it is detected on the associated graded and on the exact finite-window towers.

The bar-cobar counit  $\Omega \text{Bar}(A_{\text{op}}) \rightarrow A_{\text{op}}$  is an admissible filtered quasi-isomorphism under the same boundedness and convergence hypotheses. Composing its inverse in the homotopy category with  $\kappa_\tau$  gives  $A_{\text{op}} \simeq B_{\text{cl}}$ . The final statement is the same argument applied to the CE coalgebra  $C_*^{\text{CE}}(\mathfrak{g})$ , followed by exact continuous dualization.  $\square$

*Definition 5.24 (Tate cofibrantly generated model structure in the envelope).* Define generating cofibrations  $I$  and acyclic cofibrations  $J$  by

$$I = \{F^w S^{n-1} \hookrightarrow F^w D^n \mid n \in \mathbb{Z}_{\geq 0}, w \in \mathbb{N}\}, \quad J = \{0 \hookrightarrow F^w D^n \mid n \in \mathbb{Z}_{\geq 0}, w \in \mathbb{N}\},$$

where  $S^{n-1}, D^n$  are the standard sphere and disk chain complexes in **Vec** and the Tate window  $F^w(-)$  takes the  $w$ -th piece of the filtration. A morphism  $f: V \rightarrow W$  in **TateChain** $_{\geq 0}^{\text{adm}}$  is:

- (i) a *weak equivalence* if it is an admissible filtered quasi-isomorphism in the sense of 5.165; a quasi-isomorphism on every quotient  $V/F^w V \rightarrow W/F^w W$  and on the associated graded;
- (ii) a *cofibration* if it is in the class generated by  $I$  under transfinite composition, retracts, and pushouts;
- (iii) a *fibration* if it has the right lifting property against  $J$ .

**THEOREM 5.25** (*Admissible-envelope model-structure existence criterion*). Under Statement 5.17, the triple  $(\mathbf{TateChain}_{\geq 0}^{\text{adm}}, I, J)$  is a cofibrantly generated model category. The class of weak equivalences is closed under retracts and satisfies the two-out-of-three property; cofibrations and fibrations form complementary lifting classes; the small-object argument terminates after countably many steps because every Tate window is finite-dimensional and every generating cofibration changes only one window.

*Proof.* Quillen's small-object argument ([42, §II.3]) applies in the admissible envelope once the following are verified:

- (i)  $I$  and  $J$  permit the small-object argument: the domains and codomains of the generating maps are small with respect to relative  $I$ -cell complexes because each generator lives in one finite-dimensional Tate window  $V_{\leq w}$ , hence is  $\aleph_0$ -small in the admissible envelope.
- (ii) Every map in  $J$ -cell is a weak equivalence: pushouts of acyclic generating cofibrations are levelwise acyclic by finite-dimensional standard chain-complex theory in each window, and the filtered colimit of windowwise acyclic cofibrations is acyclic by Mittag-Leffler exactness in the Tate filtration (5.165 item 1).
- (iii) Every map with the right lifting property against  $I$  is a trivial fibration: standard, levelwise.

The two-out-of-three property for filtered quasi-isomorphisms follows from the long exact sequence of the filtration: a composite  $V \rightarrow W \rightarrow U$  has filtered  $E_1$ -page agreeing with the composite of  $E_1$ -pages of  $V \rightarrow W$  and  $W \rightarrow U$  in each window, so two-out-of-three on each window propagates to the global filtered class. This proves the model structure in the chosen envelope. Arbitrary unbounded Tate complexes outside the envelope require a separate construction.  $\square$

**LEMMA 5.26** (*Tail quotient obstruction invisible to finite-window weak equivalence*). Let  $\{V_d\}_{d \geq 1}$  be nonzero finite-dimensional vector spaces with zero differential, and put

$$V_{\oplus} = \bigoplus_{d \geq 1} V_d, \quad V_{\Pi} = \prod_{d \geq 1} V_d.$$

Filter both complexes by degree tails. The inclusion  $V_{\oplus} \hookrightarrow V_{\Pi}$  is an admissible filtered weak equivalence in the sense of Statement 5.17: it is an isomorphism on every finite quotient and on associated graded pieces. Its cofiber has

$$H^0(V_{\Pi}/V_{\oplus}) = V_{\Pi}/V_{\oplus} \neq 0.$$

*Proof.* The quotient by the tail  $d > w_0$  is

$$V_{\oplus} / \bigoplus_{d > w_0} V_d \cong \bigoplus_{d \leq w_0} V_d \cong \prod_{d \leq w_0} V_d \cong V_{\Pi} / \prod_{d > w_0} V_d.$$

Thus the inclusion is an isomorphism on every finite quotient. The associated graded in degree  $d$  is  $V_d$  on both sides, so the associated graded map is also an isomorphism. This is precisely an admissible filtered weak equivalence.

Since the differential is zero, ordinary cohomology is the underlying vector space. A sequence with one nonzero component in every degree defines a nonzero class of  $V_{\Pi}/V_{\oplus}$ . Hence the cofiber has nonzero  $H^0$ . An ordinary quasi-isomorphism of chain complexes would have acyclic cofiber, so the inclusion is only a filtered weak equivalence in the admissible envelope.  $\square$

### 5.4.3 OPERAD-ALGEBRA TRANSFER

Let  $\text{Lie}_{\mathbf{TateVec}}$  be the Lie operad in  $\mathbf{TateVec}$ , regarded as a  $\Sigma$ -cofibrant  $\Sigma$ -module (by the standard Koszul resolution of the  $\Sigma$ -cofibrant replacement), and similarly  $\text{Ass}_{\mathbf{TateVec}}^+$  for augmented associative. A  $\text{Lie}_{\mathbf{TateVec}}$ -algebra is a Tate Lie algebra in  $\mathbf{TateVec}$ ; an  $\text{Ass}_{\mathbf{TateVec}}^+$ -algebra is a Tate complete augmented associative algebra. Write

$$\text{TateLie} = \text{Lie}_{\mathbf{TateVec}}\text{-alg}_{\geq 0}, \quad \text{TateAssocAlg}^+ = \text{Ass}_{\mathbf{TateVec}}^+\text{-alg}_{\geq 0}$$

in  $\mathbf{TateChain}_{\geq 0}^{\text{adm}}$ .

LEMMA 5.27 (*Transferred model structures on Tate operad-algebras*). In the admissible envelope, the free-forget adjunction  $\text{Free}_{\mathcal{P}} \dashv \text{Forget}$  for  $\mathcal{P} \in \{\text{Lie}_{\mathbf{TateVec}}, \text{Ass}_{\mathbf{TateVec}}^+\}$  transfers the cofibrantly generated model structure of 5.25 to  $\mathcal{P}\text{-alg}_{\geq 0}$  in the sense of Hinich [35, §2.2]: the weak equivalences and fibrations in  $\mathcal{P}\text{-alg}_{\geq 0}$  are the maps whose underlying chain map in  $\mathbf{TateChain}_{\geq 0}^{\text{adm}}$  is one, and the cofibrations are determined by lifting.

*Proof.* Hinich's transfer theorem requires:

- (1) a path object on every algebra of the form  $\text{Free}_{\mathcal{P}}(F^w D^n)$ ;
- (2) that pushouts along generating acyclic cofibrations be weak equivalences.

Item (1) follows because  $\mathcal{P} = \text{Lie}$  or  $\text{Ass}^+$  is a  $\Sigma$ -cofibrant operad in  $\mathbf{TateVec}$ ; the path object is constructed from the cylinder object on chain complexes via the Berger-Moerdijk path-functor lift [33], valid whenever the operad is admissible and the ambient category is cofibrantly generated and proper. Item (2) follows because  $\mathcal{P}$  acts levelwise on each Tate window, and the windowwise finite-dimensional pushout is a quasi-isomorphism by Schwede-Shipley [43, Lemma 2.3]. Filtered-colimit assembly preserves quasi-isomorphism by Mittag-Leffler exactness (5.165 item 1).  $\square$

The dual coalgebra side requires more care. Hinich's transfer applies to algebra categories; the coalgebra case is governed instead by conilpotence and the bar-cobar adjunction. The substitute is the Aubry-Lemaire-Smirnov [24] §11.4 model structure on conilpotent dg coalgebras transferred through the bar-cobar adjunction itself.

Definition 5.28 (*Conilpotent Tate coalgebras*). Let

$$\begin{aligned} \text{TateConilpCoAlg} = \{ (C, \Delta_C, d_C) : C \in \mathbf{TateChain}_{\geq 0}^{\text{adm}}, \\ \Delta_C : C \rightarrow C \hat{\otimes} C \text{ continuous coproduct,} \\ \text{coaugmentation filtration exhaustive} \}. \end{aligned}$$

with morphisms continuous coalgebra maps. Conilpotency means the coaugmentation filtration  $F_{\text{coaug}}^{(n)} C = \ker(\bar{\Delta}^{(n)})$  is exhaustive:  $C = \bigcup_n F_{\text{coaug}}^{(n)} C$ .

LEMMA 5.29 (*Transferred model structure on Tate conilpotent coalgebras*).  $\text{TateConilpCoAlg}$  admits, in the admissible envelope, a model structure with

- weak equivalences = maps  $f : C \rightarrow C'$  such that  $\Omega(f) : \Omega(C) \rightarrow \Omega(C')$  is a weak equivalence in  $\text{TateAssocAlg}^+$  (5.27);
- cofibrations = monomorphisms of underlying filtered chain complexes;
- fibrations = right lifting against acyclic cofibrations.

Equivalently, this is the Lefevre-Hasegawa [40, Th. 1.3.1.2] transferred model structure in the admissible Tate envelope.

*Proof.* Lefevre-Hasegawa, with proof sketched in Loday-Vallette §II.4.10, constructs the conilpotent-coalgebra transferred model structure in the symmetric monoidal category of  $\mathbb{Q}$ - or  $\mathbb{C}$ -vector spaces. The proof passes through the cobar-bar adjunction  $\Omega: \text{ConilpCoAlg} \rightleftarrows \text{AugAssocAlg}: \text{Bar}$  and uses that

- (i)  $\Omega$  takes coalgebra cofibrations (= mono with filtered cokernel) to algebra cofibrations;
- (ii)  $\Omega$  inverts coalgebra weak equivalences;
- (iii) the unit  $\eta_C: C \rightarrow \text{Bar } \Omega(C)$  is a weak equivalence for every conilpotent dg coalgebra.

Items (i)-(iii) are checked windowwise (each Tate window is finite-dimensional, so the standard Lefevre-Hasegawa proof applies) and assemble by the transfer, smallness, monoid, completed tensor, and Mittag-Leffler exactness hypotheses built into Statement 5.17. The unit weak-equivalence (iii) is precisely 5.165 item 3 (admissible filtered qiso) inside that envelope.  $\square$

#### 5.4.4 BAR-COBAR QUILLEN EQUIVALENCE IN THE ADMISSIBLE ENVELOPE

**THEOREM 5.30** (*Tate bar-cobar Quillen equivalence in the admissible envelope*). The bar-cobar adjunction

$$\Omega: \text{TateConilpCoAlg} \rightleftarrows \text{TateAssocAlg}^+: \text{Bar}$$

is a Quillen equivalence with respect to the admissible-envelope model structures of 5.27 and 5.29. In particular:

- (i)  $\Omega$  is left Quillen: it preserves cofibrations and acyclic cofibrations.
- (ii)  $\text{Bar}$  is right Quillen: it preserves fibrations and acyclic fibrations.
- (iii) For every cofibrant conilpotent coalgebra  $C$  and every fibrant augmented associative algebra  $A$ , the unit  $\eta_C: C \rightarrow \text{Bar } \Omega(C)$  and the counit  $\varepsilon_A: \Omega \text{Bar}(A) \rightarrow A$  are weak equivalences.
- (iv) The induced derived adjunction  $L\Omega: \text{Ho}(\text{TateConilpCoAlg}) \rightleftarrows \text{Ho}(\text{TateAssocAlg}^+): R\text{Bar}$  is an equivalence of homotopy categories.

*Proof.* Items (1) and (2) are immediate from 5.29: weak equivalences in  $\text{TateConilpCoAlg}$  are defined as maps whose  $\Omega$  is a weak equivalence, and fibrations on the coalgebra side are defined by lifting.

For item (3), the unit  $\eta_C$ : filter both sides by symmetric / tensor length on the underlying **TateVec** object, plus the Tate window filtration  $F^w$ . The associated graded gives, windowwise, a finite-dimensional dg coalgebra and the standard bar-cobar resolution  $C^{\text{gr}} \rightarrow \text{Bar } \Omega(C^{\text{gr}})$ . By Loday-Vallette [24, §2.2.5, §II.1.5], this is a quasi-isomorphism on each window. Conilpotency (5.28) ensures the coaugmentation filtration on  $C$  is exhaustive and Hausdorff, so the abutment of the spectral sequence of the filtration computes the global qiso. Mittag-Leffler exactness in the Tate filtration (5.165 item 1) lifts the windowwise qiso to a global filtered qiso.

For the counit  $\varepsilon_A: \Omega \text{Bar}(A) \rightarrow A$ , filter both sides by augmentation/tensor length and by the Tate window. On every finite window and every finite length cap this is the ordinary bar-cobar counit for an augmented dg associative algebra over  $\mathbb{C}$ , hence a quasi-isomorphism by the standard normalized bar resolution [24, §2.2.5, §II.1.5]. The filtrations are bounded below, complete, and Hausdorff in the admissible envelope, so the spectral sequence comparison and Mittag-Leffler exactness lift the windowwise quasi-isomorphism to the completed Tate object.

In the Lie specialization  $A = \widehat{U}(\mathfrak{g})$ , the same counit can also be read through the PBW filtration:  $\text{gr}_{\text{PBW}} \widehat{U}(\mathfrak{g}) \cong \widehat{S}_{\text{Tate}}(\mathfrak{g})$ , and the associated graded comparison is the symmetric Koszul resolution of

Loday-Vallette [24, §3.4.13]. This is the closed-side PBW form used later, but the Quillen counit above is the associative bar-cobar counit for general  $A$ .

Item (4) is formal from (1)-(3) by Quillen's adjoint-functor theorem ([39, Th. 1.3.10]).  $\square$

#### 5.4.5 PROMOTION TO THE ASSOCIATED $\infty$ -CATEGORIES

COROLLARY 5.31 (*Infinity-categorical Tate bar-cobar equivalence*). Apply the Dwyer-Kan / Hammock-localization functor to the Quillen equivalence of 5.30:

$$N_{\Delta}(\text{TateConilpCoAlg}^{\text{cf}}, W) \simeq N_{\Delta}(\text{TateAssocAlg}^{+, \text{cf}}, W)$$

as the  $\infty$ -categories associated to the admissible model envelope in the sense of Lurie's *Higher Algebra* §A.3, where cf denotes the full subcategory of cofibrant-fibrant objects and  $W$  is the class of weak equivalences. The equivalence is induced by  $L\Omega$  and  $R\text{Bar}$  at the cofibrant-fibrant level. Presentability is an attribute of the chosen locally presentable envelope.

*Proof.* Lurie [41, A.3.7.6, A.3.7.10]: a Quillen equivalence of model categories induces an equivalence of associated  $\infty$ -categories. The presentability of both sides, inside the envelope, is inherited from the cofibrant generation of the model structures (5.25, 5.27, 5.29) by Lurie A.2.6.8.  $\square$

#### 5.4.6 IDENTIFICATION WITH THE LIE OPERADIC KOSZUL DUALITY

The Quillen equivalence of 5.30 is *associative* bar-cobar. Specialization to the Lie operad gives the cochain-level equivalence used in the CE/PV theorem.

COROLLARY 5.32 (*Tate Lie bar-cobar / CE Koszul duality*). For  $\mathfrak{g}$  a lower-central Tate-pronilpotent graded Lie algebra in the sense of 5.165, the bar-cobar Quillen equivalence applied to  $A = \widehat{U}(\mathfrak{g}) \in \text{TateAssocAlg}^+$  yields the equivalence in the homotopy category

$$L\Omega(C_*^{\text{CE}}(\mathfrak{g})) \xrightarrow{\sim} \widehat{U}(\mathfrak{g}), \quad R\text{Bar}(\widehat{U}(\mathfrak{g})) \xrightarrow{\sim} C_*^{\text{CE}}(\mathfrak{g}),$$

and hence, after continuous dualization, a Lie-Koszul-dual equivalence of cochain algebras

$$C_{\text{CE}, \text{cont}}^{\bullet}(\mathfrak{g}) \simeq (\widehat{U}(\mathfrak{g}))^!$$

in the corresponding homotopy category of complete augmented Tate algebras.

Inside the admissible envelope, by the symmetric-monoidal-functoriality of the bar-cobar Quillen equivalence [24, §13.2] and Lurie [41, §5.5], this lifts to an equivalence of symmetric monoidal  $\infty$ -categories,

$$N_{\Delta}(\text{TateConilpCoAlg}_{\text{Lie}}^{\text{cf}}, W) \simeq N_{\Delta}(\text{TateAssocAlg}_{\text{Lie}}^{+, \text{cf}}, W)$$

in the framework of Lurie *Higher Algebra* §5.5 (locally constant factorization algebras over  $\mathbb{R}$ , applied pointwise on the line; or directly the formal-disk version).

*Proof.* Apply 5.30 to the Lie-operadic Koszul-dual pair (Lie, CoComm) in **TateVec**. Loday-Vallette [24, Th. 13.4.6] identifies the operadic bar-cobar with the Lie-bar-cobar of CE chains and completed enveloping algebra. The symmetric-monoidal extension is direct because the operad Lie is a symmetric-monoidal Hopf operad in **Vec**, and base change to **TateVec** via the strong symmetric-monoidal functor  $\mathbf{Vec} \rightarrow \mathbf{TateVec}$  preserves operadic Koszul duality ([24, §13.5]).  $\square$

LEMMA 5.33 (*Closed-side Tate formalism*). Let  $\mathfrak{h} = \varprojlim_M \mathfrak{h}_{\leq M}$  be a closed Hamiltonian Tate Lie algebra whose finite windows are cochain-CE/PV-admissible, and kernel-admissible when a  $P_0$  kernel is used, in the sense of Definition 5.19. Assume:

- (K1) the shifted cotangent extension  $\mathfrak{g} = \mathfrak{h} \ltimes D_{\text{cont}}[1]$  is an object of  $\mathbf{TateChain}_{\geq 0}^{\text{adm}}$ ;
- (K2) the CE coalgebra  $C_*^{\text{CE}}(\mathfrak{g})$  is conilpotent for the coaugmentation filtration;
- (K3) the completed enveloping algebra  $\widehat{U}(\mathfrak{g}) = \varprojlim_M \widehat{U}(\mathfrak{g}_{\leq M})$  is complete and separated for the PBW filtration;
- (K4) the finite-window PBW associated graded maps assemble to

$$\text{gr}_{\text{PBW}} \widehat{U}(\mathfrak{g}) \cong \widehat{\mathbf{S}}_{\text{Tate}}(\mathfrak{g}).$$

Then the closed-side formalism has two compatible outputs:

$$C_{\text{CE,cont}}^\bullet(\mathfrak{g}) \xrightarrow{\Phi} \text{PV}(\widehat{\mathbf{S}}_{\text{Tate}}(\mathfrak{h}))$$

is the admissible completed CE/PV identity obtained from the finite-window family of Lemma 5.20 together with the exact continuous-dualization and variance data in (K1)–(K4), and

$$L\Omega C_*^{\text{CE}}(\mathfrak{g}) \simeq \widehat{U}(\mathfrak{g}), \quad R\text{Bar } \widehat{U}(\mathfrak{g}) \simeq C_*^{\text{CE}}(\mathfrak{g})$$

is the Tate Lie bar-cobar / CE Koszul duality of Corollary 5.32. The two outputs are compatible with the same finite-window transition maps.

*Proof.* The first output is Lemma 5.20 plus the exact continuous-dualization and variance hypotheses included above. The obstruction terms  $\mathfrak{o}^{\text{br}}$  and  $\mathfrak{o}^{\text{coad}}$  are the cochain CE/PV transition obstructions;  $\mathfrak{o}^{\text{Cas}}$  is the additional kernel/ $P_0$ -pairing obstruction. Their vanishing supplies the finite-window compatibility; the admissible envelope supplies the cochain inverse-limit datum for  $\Phi = \varprojlim_M \Phi_M$ .

For the Koszul output, hypotheses (K1)–(K4) put  $C_*^{\text{CE}}(\mathfrak{g})$  and  $\widehat{U}(\mathfrak{g})$  in the admissible model envelope with complete filtrations. Conilpotency is the coalgebra hypothesis needed for Bar  $\Omega$ , and PBW completeness identifies the associated graded of the enveloping algebra with the completed symmetric algebra. Corollary 5.32 therefore applies. Both constructions are obtained by first applying the finite-dimensional CE/PV and bar-cobar statements on  $\mathfrak{g}_{\leq M}$ , then passing through the same Mittag-Leffler inverse limit; hence the transition maps are the same.  $\square$

#### 5.4.7 MODULE RECONSTRUCTION FROM THE CE/PV IDENTITY

THEOREM 5.34 (*Admissible-envelope module reconstruction from the CE/PV identity*). Let  $\mathfrak{h}_{\text{adm}}$  be a pro-finite-dimensional Lie algebra in  $\mathbf{TateVec}$  whose shifted-cotangent extension  $\mathfrak{g}_{\text{adm}} = \mathfrak{h}_{\text{adm}} \ltimes \mathfrak{h}_{\text{adm,cont}}^\vee[1]$  satisfies the lower-central Tate-pronilpotent admissible-envelope hypotheses of 5.30. Assume further that:

- (M1) the strict cochain map

$$\Phi: C_{\text{CE,cont}}^\bullet(\mathfrak{g}_{\text{adm}}) \longrightarrow \text{PV}(\widehat{\mathbf{S}}_{\text{Tate}}(\mathfrak{h}_{\text{adm}}))$$

is the admissible CE/PV quasi-isomorphism obtained from the finite-window family of Lemma 5.20 after the required variance and exact continuous-dualization data are supplied;

- (M<sub>2</sub>) the completed dg  $P_o$ -algebras on both sides lie in an admissible transferred  $P_o$ -algebra model structure inside  $\mathbf{TateChain}_{\geq o}^{\text{adm}}$ , with weak equivalences detected on underlying admissible filtered quasi-isomorphisms;
- (M<sub>3</sub>) the module categories are formed in the same locally presentable monoidal envelope, and the two algebras are replaced by cofibrant-fibrant representatives when needed.

Then  $\Phi$  induces an equivalence, inside that admissible Tate model envelope, of module  $\infty$ -categories over the two identified completed dg  $P_o$ -algebras

$$\tilde{\Phi}: N_{\Delta}(C_{\text{CE,cont}}^{\bullet}(\mathfrak{g}_{\text{adm}})\text{-mod}, W) \xrightarrow{\sim} N_{\Delta}(\text{PV}(\widehat{S}_{\text{Tate}}(\mathfrak{h}_{\text{adm}}))\text{-mod}, W)$$

in the framework of Lurie's *Higher Algebra* §5.5. The hypotheses (M<sub>1</sub>)–(M<sub>3</sub>) presuppose an already constructed admissible  $P_o$ -algebra envelope. The conclusion is the algebraic  $\infty$ -categorical equivalence inside that envelope. The completed dg  $P_o$  model structure and the factorization-centre theorem for the full untruncated formal Hamiltonian algebra are separate constructions.

*Proof.* Assumption (M<sub>1</sub>) gives a strict admissible filtered quasi-isomorphism of completed dg  $P_o$ -algebras. The strictness is the generator-level identity  $c \mapsto \theta$ ,  $u \mapsto O$ , together with  $d_{\pi}\Phi = \Phi d_{\text{CE}}$  and bracket compatibility from the finite-window CE/PV calculation.

By (M<sub>2</sub>), this strict weak equivalence is a weak equivalence in the transferred  $P_o$ -algebra model structure. Replace the two  $P_o$ -algebras by cofibrant-fibrant representatives if necessary. Extension and restriction of scalars along a weak equivalence of admissible algebra objects give a Quillen equivalence of module model categories; this is the standard Schwede–Shipley/Lurie mechanism for module categories over weakly equivalent algebra objects [43] and [41, Th. 7.1.4.6]. Applying Dwyer–Kan localization gives the displayed equivalence of module  $\infty$ -categories.

Functoriality in the Lie algebra gives interval-wise compatibility for admissible interval objects  $\mathfrak{h}_{\text{adm},I} = \Omega_c^{\bullet}(I) \otimes \mathfrak{h}_{\text{adm}}$  when these interval objects remain in the same admissible envelope. This gives interval-wise categorical compatibility. Weiss descent and unreduced open-BV centrality are separate theorems.  $\square$

*Remark 5.35 (Relation to Lurie's higher-algebraic Koszul duality).* Lurie [41, §5.5.5] constructs a higher-algebraic Koszul duality between augmented  $\infty$ -operad-algebras and their Koszul-dual coalgebras. For the operad  $\text{Lie}_{\text{TateVec}}$ , Koszul self-duality (with shift) gives the equivalence of 5.32 in the lower-central Tate-pronilpotent setting. The two constructions agree: weak equivalences in the model-categorical presentation are admissible filtered quasi-isomorphisms (5.165), and Lurie A.3.7 with the Hinich transfer of 5.27 identifies them with the Hammock-localized weak equivalences of the  $\infty$ -categorical presentation.

#### 5.4.8 CONILPOTENCE AND CASIMIR CONVERGENCE

**PROPOSITION 5.36** (*Bar-cobar conilpotence separated from Casimir convergence*). The Quillen equivalence of 5.30 and the resulting  $\infty$ -categorical equivalence of 5.34, when their admissible-envelope hypotheses hold, are independent of the convergence of the Tate Casimir element. The cochain complexes  $C_{\text{CE,cont}}^{\bullet}(\mathfrak{g})$  and  $\text{PV}(\widehat{S}_{\text{Tate}}(\mathfrak{h}))$  are equivalent in the model category of 5.27 after imposing the same conilpotent admissibility hypotheses. The required hypothesis here is conilpotence; Casimir convergence belongs to the analytic kernel problem.

*Proof.* The bar-cobar comparison uses  $\widehat{S}^c(\mathfrak{g}[1])$  and  $T(\mathfrak{g})$  as underlying graded objects. Both are well-defined and conilpotent in the admissible envelope under 5.165 items 1, 2 (continuous dual exact, conilpotent CE coalgebra). The formal Casimir  $\sum_d e_{d,i} \otimes e_d^{i,\vee}$  is a kernel hypothesis only when it converges



in the chosen coefficient category; in the strict product/direct-sum Tate endpoint it is precisely the obstruction isolated in Lemma 5.2II. Such a kernel is required for the Costello [12] heat-kernel propagator construction at the analytic / factorization-algebra-with-non-strict-bracket level. The algebraic bar-cobar comparison uses only the conilpotent admissibility hypotheses: the cochain-level  $\Phi$  promotes, in the admissible model envelope, to an  $\infty$ -categorical equivalence.  $\square$

**PROPOSITION 5.37** (*Strict product/direct-sum kernel obstruction*). Let  $C_{\ker}$  be the category whose objects are complete coefficient pairs  $(H, D)$ , equipped with finite quotients  $(H_{\leq M}, D_{\leq M})$ , compatible transition maps, and a tensor  $K \in H \widehat{\otimes} D$  whose image in  $H_{\leq M} \otimes D_{\leq M}$  is the finite Casimir  $C_M$  for every  $M$ . Morphisms preserve the finite quotients and carry  $K$  to  $K'$ . The original strict product/direct-sum pair

$$H_{\Pi} = \prod_{d \geq 1} H_d, \quad D_{\oplus} = \bigoplus_{d \geq 1} H_d^{\vee}$$

lies outside  $C_{\ker}$ . More strongly, any object of  $C_{\ker}$  mapping to the strict pair by a finite-window identity morphism would have to produce the obstructed infinite-support tensor in  $H_{\Pi} \widehat{\otimes} D_{\oplus}$ .

Thus initial, terminal, and representing universal properties for kernel-admissible completions pass through a genuine kernel object. A weighted product completion is a constructed regulator for tail-sensitive ordinary cohomology classes.

*Proof.* If the strict pair were an object, it would contain a tensor  $K \in H_{\Pi} \widehat{\otimes} D_{\oplus}$  whose image in every finite window is  $C_M$ . If a kernel-admissible object mapped to the strict pair by a finite-window identity morphism, the image of its kernel would give the same obstructed tensor. But  $D_{\oplus}$  has finite support in the degree direction, while the compatible family  $(C_M)_M$  has a nonzero  $H_d^{\vee}$ -component for every  $d$ . This is exactly the support obstruction of Theorem 5.93. Therefore the strict pair lies outside  $C_{\ker}$ , and universal-property assertions must pass through a kernel-admissible completion.  $\square$

**PROPOSITION 5.38** (*Residual Weiss-cosheaf condition*). The  $\infty$ -categorical equivalence of 5.34 on stalks and on the locally-constant interval-factorization  $\mathcal{F}_{\text{cl}}$  of 4.11 gives the cochain-level categorical comparison. The Weiss-cosheaf condition for the *unreduced* factorization algebra  $\text{Obs}_{\text{closed}, H}^{\text{cl}}$  on  $\mathbb{R}^2 \times \mathbb{C}^2$  requires the additional descent check below. The Weiss / descent condition for an unreduced factorization algebra on a higher-dimensional manifold requires checking Čech descent for Weiss covers, which is a separate cosheaf-theoretic statement. Costello–Gwilliam Vol. I §§6.1, 6.5, 7.2 provide the factorization, cosheaf, and basis-extension vocabulary; the present argument uses them as vocabulary and leaves the unreduced descent obligation open. The admissible Tate Quillen-equivalence promotion closes the cochain-level  $\infty$ -categorical comparison; the Weiss upgrade on the unreduced factorization algebra remains a separate cosheaf-theoretic obligation requiring the heat-kernel propagator extension of Problem 5.48. The two-dimensional formal singular product calculation is the separate result of Theorem 1.15.

*Proof.* Lurie [41, §5.5.4], with the same terminology as Costello–Gwilliam [15, Vol. I §6.4]: a locally constant factorization algebra on  $\mathbb{R}$  determines an  $E_1$ -algebra up to the standard contractible choice; the Quillen equivalence of 5.30 realizes this. On a higher-dimensional manifold, the Weiss cover  $\{U_{\alpha}\}$  has structure-map descent independent of the algebraic bar-cobar; in particular the heat-kernel propagator on  $\mathbb{R}^2 \times \mathbb{C}^2$  enters the analytic check. This heat-kernel propagator is the analytic input relating the bar-cobar identity to Weiss descent.  $\square$

COROLLARY 5.39 (*Moment-map specialisation*). In the situation of 4.20, the degree-zero specialization of  $\Phi_{\text{coord}}$  on the cotangent direction is the moment-map / Kirillov assignment

$$J : \mathfrak{h} \longrightarrow A_\partial = \widehat{\mathbf{S}}_{\text{Tate}}(\mathfrak{h}), \quad J(f) = O_f.$$

The closed-side Lie bracket  $\{f, g\}_{\mathfrak{h}}$  matches the Kirillov–Poisson bracket on  $A_\partial$ :

$$\{J(f), J(g)\}_\partial = J(\{f, g\}_{\mathfrak{h}}).$$

*Proof.*  $J$  is the restriction of  $\Phi_{\text{coord}}$  to the cotangent direction  $u_I \mapsto O_I$  followed by linear extension along  $\mathfrak{h} \hookrightarrow A_\partial$  (the canonical embedding of  $\mathfrak{h}$  as the linear functions on  $\mathfrak{h}^\vee$ ). Bracket compatibility is the  $(c^J, u_K)$ -versus- $(\theta^J, O_K)$  identity from 4.20 read off in degree zero. Equivalently, this is the standard Kirillov–Kostant–Souriau theorem for linear Poisson structures.  $\square$

LEMMA 5.40 (*Natural-correction obstruction*). Let  $\tilde{J} : \mathfrak{h} \rightarrow A_\partial$  be a linear map satisfying stable polynomial  $GL_\infty$ -invariance (Proposition 3.22), descent through the Dirac ideal  $[\phi_1, \phi_2] = 0$ , the scalar quotient  $\overline{\text{Tr}(1)} = 0$ , and primitive block-sum additivity:

$$\tilde{J}_{N_1+N_2}(f)|_{\phi_i=\phi'_i\oplus\phi''_i} = \tilde{J}_{N_1}(f) + \tilde{J}_{N_2}(f),$$

the value at the block-diagonal stacking  $(\phi_1, \phi_2) = (\phi'_1, \phi'_2) \oplus (\phi''_1, \phi''_2)$  on  $\mathfrak{gl}_{N_1} \oplus \mathfrak{gl}_{N_2} \subset \mathfrak{gl}_{N_1+N_2}$  reading on the right as a vector-space sum in  $A_\partial$ . Equivalently,  $\tilde{J}(f)$  is primitive for the block-sum coproduct  $\Delta_\oplus$  on  $A_\partial$  sending each single-trace generator  $\overline{\text{Tr}}$  to  $\overline{\text{Tr}} \otimes 1 + 1 \otimes \overline{\text{Tr}}$ . Then for every  $a + b > 0$ ,

$$\tilde{J}(z_1^a z_2^b) = c_{a,b} \overline{\text{Tr}(\phi_1^a \phi_2^b)}, \quad c_{a,b} \in \mathbb{C}.$$

*Proof.* Stable invariant theory (Proposition 3.22) writes  $\tilde{J}(z_1^a z_2^b)$  as a polynomial in nonempty cyclic trace words on the free pair, of total bidegree  $(a, b)$ ; scalar reduction kills empty traces. Restriction to the moment-map zero-fibre with adjacent-swap exactness (Lemma 3.30) collapses each cyclic word at bidegree  $(p, q)$  to the commutative monomial trace  $\overline{\text{Tr}(\phi_1^p \phi_2^q)}$  (Corollary 3.32), so

$$\tilde{J}(z_1^a z_2^b) = \sum_{\lambda \vdash (a,b)} c_\lambda \prod_{i=1}^{\ell(\lambda)} \overline{\text{Tr}(\phi_1^{p_i} \phi_2^{q_i})},$$

summed over partitions of  $(a, b)$  into  $\ell(\lambda)$  nonempty parts  $(p_i, q_i)$ . Block-sum stacking sends the single-trace generator  $\overline{\text{Tr}(\phi_1^p \phi_2^q)}$  to its block-diagonal sum  $\overline{\text{Tr}((\phi'_1)^p (\phi'_2)^q)} + \overline{\text{Tr}((\phi''_1)^p (\phi''_2)^q)}$  — this is the Loday–Quillen–Tsygan primitive on  $H_*^{\text{CE}}(\mathfrak{gl}_\infty(A_{\tilde{f}}))$  — and a length- $\ell$  product to a sum of  $2^\ell$  terms with  $2^\ell - 2$  cross-block products outside the LQT primitive. The primitive block-sum identity therefore vanishes the  $\ell \geq 2$  coefficients, leaving the single-part partition  $\lambda = ((a, b))$ .  $\square$

THEOREM 5.41 (*Uniqueness of  $J$* ). Let  $\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$  be the reduced Hamiltonian Lie algebra of the formal symplectic polydisk and let  $A_\partial = \widehat{\mathbf{S}}_{\text{Tate}}(\mathfrak{h})$  be the stable scalar-reduced Hamiltonian trace algebra of Proposition 3.40. Among linear maps

$$\tilde{J} : \mathfrak{h} \longrightarrow A_\partial$$

satisfying:

- (U<sub>1</sub>) (*Polynomial  $GL_N$ -invariance, stable*)  $\tilde{J}(f)$  is a stable  $GL_N$ -invariant trace polynomial in  $(\phi_1, \phi_2)$  in the sense of Proposition 3.22;
- (U<sub>2</sub>) (*Dirac constraint*)  $\tilde{J}(f)$  descends modulo the moment-map ideal  $[\phi_1, \phi_2] = 0$ ;
- (U<sub>3</sub>) (*Scalar reduction*)  $\tilde{J}$  factors through the stable scalar-reduced quotient  $A_\partial = A_\infty^{GL_\infty}/\mathbb{C} \cdot \text{Tr}(\mathbf{1})$ ;
- (U<sub>4</sub>) (*Darboux naturality*) for every formal canonical transformation  $\Phi: \widehat{\mathbb{C}}^2_o \rightarrow \widehat{\mathbb{C}}^2_o$  preserving  $\omega = dz_1 \wedge dz_2$ ,  $\tilde{J}(\Phi^* f) = \Phi^* \tilde{J}(f)$  under the induced action on  $A_\partial$ ;
- (U<sub>5</sub>) (*Linear normalization*)  $\tilde{J}(z_1) = \overline{\text{Tr } \phi_1}$ ,  $\tilde{J}(z_2) = \overline{\text{Tr } \phi_2}$ ;
- (U<sub>6</sub>) (*Primitive block-sum additivity*) under the block-diagonal stacking  $(\phi_1, \phi_2) = (\phi'_1, \phi'_2) \oplus (\phi''_1, \phi''_2)$  on  $\mathfrak{gl}_{N_1} \oplus \mathfrak{gl}_{N_2} \subset \mathfrak{gl}_{N_1+N_2}$ ,

$$\tilde{J}_{N_1+N_2}(f)|_{\phi_i=\phi'_i \oplus \phi''_i} = \tilde{J}_{N_1}(f) + \tilde{J}_{N_2}(f) \quad \text{in } A_\partial,$$

the brane substitution  $J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)}$  is the unique solution. Lie-algebra morphism  $\{\tilde{J}(f), \tilde{J}(g)\}_\partial = \tilde{J}(\{f, g\}_{\tilde{A}})$  holds for every solution (Theorem 5.190); uniqueness is the content of (U<sub>1</sub>)–(U<sub>6</sub>). The categorical terminal-object refinement, including protected admissibility, is Theorem 5.146 of §7.5.

*Remark 5.42 (Worked instances).* The three running examples of §1.1 determine the theorem concretely. Example 1.2 is the  $N = 1$  case: the matrices commute automatically, scalar reduction is trivial, and  $J(f) = f(\phi_1, \phi_2)$  is the classical evaluation map at a point of  $\mathbb{C}^2$ . Example 1.3 shows the linear Hamiltonian normalisation  $z_i \mapsto \overline{\text{Tr } \phi_i}$  at work and exhibits the scalar reduction killing the constant-Hamiltonian line. Example 1.4 contrasts the Taylor side (what  $J$  samples) with the residue side (the Matlis dual that pairs with Taylor coefficients), which is why the natural-correction obstruction works monomial-by-monomial through finite Hamiltonian flows.

*Proof.* Lemma 5.40, drawing on (U<sub>1</sub>), (U<sub>2</sub>), (U<sub>3</sub>), (U<sub>6</sub>), gives  $\tilde{J}(z_1^a z_2^b) = c_{a,b} \overline{\text{Tr}(\phi_1^a \phi_2^b)}$  on every bidegree  $a + b > 0$  with scalars  $c_{a,b} \in \mathbb{C}$ . Linear normalization (U<sub>5</sub>) gives  $c_{1,0} = c_{0,1} = 1$ .

Darboux naturality (U<sub>4</sub>) propagates the scale to every bidegree. The formal Darboux symplectomorphism group  $G_{\text{Darb}} = \text{Symp}_{\text{form}}(\widehat{\mathbb{C}}^2_o)$  acts on  $A_\partial$  through the algebra-homomorphism matrix substitution  $\Phi_* \overline{\text{Tr } f(\phi_1, \phi_2)} = \overline{\text{Tr}(\Phi^* f)(\phi_1, \phi_2)}$ . The Hamiltonian flow generated by  $H_1^{(2)} = -z_1^2/2$  acts by  $\Phi_{H_1^{(2)}}^* z_2 = z_2 + t z_1$ , and on  $z_2^k$  gives

$$\tilde{J}((z_2 + t z_1)^k) = \sum_{a+b=k} \binom{k}{a} t^a c_{a,b} \overline{\text{Tr}(\phi_1^a \phi_2^b)}, \quad \Phi_* \tilde{J}(z_2^k) = c_{0,k} \overline{\text{Tr}(\phi_2 + t \phi_1)^k} = \sum_{a+b=k} \binom{k}{a} t^a c_{0,k} \overline{\text{Tr}(\phi_1^a \phi_2^b)}.$$

Equality at each  $t^a$  gives  $c_{a,b} = c_{0,k}$  for  $a + b = k$ ; the symmetric linear flow  $z_1 \mapsto z_1 - t z_2$  on  $z_1^k$  gives  $c_{a,b} = c_{k,0}$  and hence  $c_{0,k} = c_{k,0}$  within total degree  $k$ . Linear shears preserve total degree, so they are insufficient by themselves to chain back to the linear normalisation  $c_{1,0} = c_{0,1} = 1$ . The required cross-degree input is the nonlinear flow generated by  $H_k^{(1)} = -z_2^{k+1}/(k+1)$ ,

$$\Phi_{H_k^{(1)}}^* z_1 = z_1 - t z_2^k, \quad \Phi_{H_k^{(1)}}^* z_2 = z_2,$$

which raises the Taylor degree of  $z_1$  from 1 to  $k$  and acts on the matrix substitution as  $(\phi_1, \phi_2) \mapsto (\phi_1 - t \phi_2^k, \phi_2)$ . Naturality (U<sub>4</sub>) on the linear monomial  $z_1$  reads

$$\tilde{J}(z_1 - t z_2^k) = \tilde{J}(z_1) - t \tilde{J}(z_2^k) = \overline{\text{Tr } \phi_1} - t c_{0,k} \overline{\text{Tr } \phi_2^k}$$

on the source against

$$\Phi_* \tilde{J}(z_1) = \overline{\text{Tr}(\phi_1 - t\phi_2^k)} = \overline{\text{Tr} \phi_1} - t \overline{\text{Tr} \phi_2^k}$$

on the matrix side. Equality at  $t^1$  forces  $c_{0,k} = 1$  for every  $k \geq 1$ ; the linear-shear identity then propagates  $c_{a,b} = 1$  on every bidegree of total degree  $k$ . Continuity in the  $\widehat{\mathfrak{h}}$ -topology extends to formal power series, so  $\tilde{J} = J$ .  $\square$

**THEOREM 5.43** (*Hypothesis rigidity for  $J$* ). The substantive hypotheses are (U<sub>4</sub>) and (U<sub>5</sub>): removing either admits a positive-dimensional family of solutions  $\tilde{J} \neq J$ . Primitive block-sum additivity (U<sub>6</sub>) is the Loday–Quillen–Tsygan-explicit form of single-trace forcing; given (U<sub>1</sub>)–(U<sub>5</sub>), it is automatic, but separating it from (U<sub>4</sub>) rebalances the proof, with (U<sub>6</sub>) carrying shape (single-trace) and (U<sub>4</sub>)+(U<sub>5</sub>) carrying scale (Lemma 5.40 monomial-by-monomial). Hypotheses (U<sub>1</sub>), (U<sub>2</sub>), (U<sub>3</sub>) are codomain specifications, each implied by (U<sub>4</sub>)+(U<sub>5</sub>) on an enlarged ambient: on  $\mathbb{C}[\mathfrak{gl}_N^2]/\langle[\phi_1, \phi_2]\rangle$ ,  $\tilde{J}$  takes values in the trace algebra; on  $\mathbf{S}(\text{Cyc}(F)_{\text{nonempty}})$ ,  $\tilde{J}$  equals the symmetrized cyclic trace, which Dirac collapse identifies with  $J$ ; on  $A_\partial \oplus \mathbb{C} \cdot \overline{\text{Tr}(1)}$ , the  $\overline{\text{Tr}(1)}$ -component is zero.

*Proof. Removing (U<sub>4</sub>).* Choose any sequence  $\{c_{a,b}\}_{a+b \geq 2} \subset \mathbb{C}$  and set  $\tilde{J}(z_1^a z_2^b) = c_{a,b} \overline{\text{Tr}(\phi_1^a \phi_2^b)}$  for  $a + b \geq 2$ ,  $\tilde{J}(z_i) = \overline{\text{Tr} \phi_i}$ . The family satisfies (U<sub>1</sub>)–(U<sub>3</sub>), (U<sub>5</sub>), (U<sub>6</sub>); Darboux flow  $z_2 \mapsto z_2 + tz_1$  on  $z_2^k$  forces  $c_{a,k-a} = c_{0,k}$  and the reverse forces  $c_{0,k} = 1$ , hence every  $c_{a,b} = 1$ .

*Removing (U<sub>5</sub>).* The family  $\tilde{J}_\lambda^{\text{rs}} = \lambda J$ ,  $\lambda \in \mathbb{C}^\times$ , satisfies (U<sub>1</sub>)–(U<sub>4</sub>) and (U<sub>6</sub>); (U<sub>5</sub>) requires  $\lambda = 1$ .

*Removing (U<sub>6</sub>) with (U<sub>4</sub>) intact.* The two-trace correction  $\tilde{J}_\alpha^{\text{mt}}(z_1 z_2) = \overline{\text{Tr}(\phi_1 \phi_2)} + \alpha \overline{\text{Tr} \phi_1} \overline{\text{Tr} \phi_2}$ ,  $\tilde{J}_\alpha^{\text{mt}}(z_1^a z_2^b) = \overline{\text{Tr}(\phi_1^a \phi_2^b)}$  elsewhere, satisfies (U<sub>1</sub>), (U<sub>2</sub>), (U<sub>3</sub>), (U<sub>5</sub>) for every  $\alpha$ ; Darboux flow  $z_2 \mapsto z_2 + tz_1$  on  $z_2^2$  reads source-side  $t^1$ -coefficient  $2 \overline{\text{Tr}(\phi_1 \phi_2)} + 2\alpha \overline{\text{Tr} \phi_1} \overline{\text{Tr} \phi_2}$  against matrix-side  $2 \overline{\text{Tr}(\phi_1 \phi_2)}$ , forcing  $\alpha = 0$ . The same Darboux propagation kills every higher multi-trace correction inductively: any  $M: V_H \rightarrow \mathbf{S}^{\geq 2}(V_H)$  compatible with the full Hamiltonian flow algebra reduces to  $M = 0$ . Hence (U<sub>6</sub>) is implied by (U<sub>4</sub>)+(U<sub>5</sub>)+(U<sub>1</sub>)–(U<sub>3</sub>); inside the full hypothesis set  $\tilde{J} = J$ .

*(U<sub>1</sub>) reduction.* On  $\mathbb{C}[\mathfrak{gl}_N^2]/\langle[\phi_1, \phi_2]\rangle$ , (U<sub>5</sub>) gives  $\tilde{J}(z_i) = \overline{\text{Tr} \phi_i}$  (single trace, hence  $GL_N$ -invariant). The flow  $(\Phi_t^{H^{(1)}})^* z_1 = z_1 - tz_2^k$  under (U<sub>4</sub>) gives

$$\tilde{J}(z_1) - t\tilde{J}(z_2^k) = \overline{\text{Tr} \phi_1} - t\overline{\text{Tr} \phi_2^k},$$

hence  $\tilde{J}(z_2^k) = \overline{\text{Tr} \phi_2^k}$ . The reverse flow gives  $\tilde{J}(z_1^k)$ ; composite flows give  $\tilde{J}(z_1^a z_2^b) = \overline{\text{Tr}(\phi_1^a \phi_2^b)}$  at every bidegree.

*(U<sub>2</sub>) reduction.* On  $\mathbf{S}(\text{Cyc}(F)_{\text{nonempty}})$  (Proposition 3.22), (U<sub>5</sub>) gives  $\tilde{J}(z_i) = \overline{\text{Tr} \phi_i}$ ; matrix substitution sends  $\overline{\text{Tr} P(\phi)}$  to  $\overline{\text{Tr} P(\Phi(\phi))}$ . The  $t^j$ -coefficient of  $(z_2 + tz_1)^k$  under (U<sub>4</sub>), matched against  $\overline{\text{Tr}(\phi_2 + t\phi_1)^k}$ , gives

$$\tilde{J}(z_1^j z_2^{k-j}) = \frac{1}{\binom{k}{j}} \sum_{w \in W_{j,k-j}} \overline{\text{Tr} w(\phi_1, \phi_2)}.$$

At bidegree  $(2, 2)$ :  $\frac{2}{3} \overline{\text{Tr}(\phi_1 \phi_1 \phi_2 \phi_2)} + \frac{1}{3} \overline{\text{Tr}(\phi_1 \phi_2 \phi_1 \phi_2)}$ . On commuting matrices both cyclic classes equal  $\overline{\text{Tr}(\phi_1^2 \phi_2^2)} = J(z_1^2 z_2^2)$ ; Lemma 3.30 extends this to every bidegree.

*(U<sub>3</sub>) reduction.* On  $A_\partial \oplus \mathbb{C} \cdot \overline{\text{Tr}(1)}$ , write  $\tilde{J}(f) = \tilde{J}_{\text{red}}(f) + L(f) \overline{\text{Tr}(1)}$  with  $L: \mathfrak{h} \rightarrow \mathbb{C}$  linear. Matrix substitution fixes  $\overline{\text{Tr}(1)}$ , so (U<sub>4</sub>) gives  $L(\Phi^* z_i) = L(z_i)$ ; the Hamiltonian flows give  $L(z_i^k) = 0$  for every  $k \geq 1$ , and composite flows give  $L \equiv 0$ .  $\square$

**COROLLARY 5.44** (*CE/PV generator identity, formal symplectic polydisk*). Let  $\mathfrak{h} = \widehat{A}/\mathbb{C} \cdot 1 = (z_1, z_2)\mathbb{C}[[z_1, z_2]]$  be the reduced Hamiltonian Lie algebra of the formal symplectic polydisk on  $\mathbb{C}^2$ , with bracket  $\{f, g\} = \partial_{z_1}f \partial_{z_2}g - \partial_{z_2}f \partial_{z_1}g$  modulo constants. Choose the Taylor basis  $H_{a,b} = z_1^a z_2^b$  for  $a, b \geq 0, a + b > 0$ . Structure constants:

$$[H_{a,b}, H_{c,d}] = (ad - bc) H_{a+c-1, b+d-1},$$

with the convention that  $H_{m,n} = 0$  if  $\min(m, n) < 0$  or  $m + n = 0$ .

Set  $A_\partial = \widehat{\mathbf{S}}_{\text{Tate}}(\mathfrak{h})$ , with linear coordinates  $O_{a,b}$  on  $\mathfrak{h}^\vee$  and Kirillov–Poisson bracket  $\{O_{a,b}, O_{c,d}\}_\partial = (ad - bc)O_{a+c-1, b+d-1}$ . The generator calculation underlying 4.20 gives the cochain-level coordinate CE/PV identity, read in the pro coordinate-window sense,

$$\Phi_{\text{coord}} : \mathbf{C}_{\text{CE, coord}}^\bullet(\mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1]) \xrightarrow{\sim} \text{PV}_{\text{coord}}(\widehat{\mathbf{S}}_{\text{Tate}}(\mathfrak{h})), \quad c^{a,b} \mapsto \theta^{a,b}, \quad u_{a,b} \mapsto O_{a,b}.$$

Its moment-map specialisation is the boundary evaluation map  $J(f) = O_f = \overline{\text{Tr } f(\phi_1, \phi_2)}$  of 5.190 after identifying  $A_\partial$  with the stable scalar-reduced Hamiltonian trace algebra of 3.40. The  $P_0$  and factorization central-operation interpretation is imposed only after the coordinate-window or mixed/weighted continuity hypotheses in the theorem-data table; the strict product coordinate completion has a separate global  $P_0$ -algebra obstruction.

*Proof.* The structure constants are an instance of 2.7 read on monomials. The CE/PV differential identity is checked directly on the monomial generators  $c^{a,b}, u_{a,b}$  and then extended by the completed Leibniz rule; lower-central Tate-pronilpotence of the full Hamiltonian algebra is a separate bar-cobar hypothesis. The moment-map specialisation follows from 5.39 together with 5.190.  $\square$

**Example 5.45** (*Low-degree monomial verification*). Direct verification of  $\Phi_{\text{coord}}$  on small monomials of the formal symplectic polydisk.

- *Degree-1 generators.* Take  $H_{1,0} = z_1, H_{0,1} = z_2$ . The bracket  $[H_{1,0}, H_{0,1}] = 1 \cdot 1 - 0 \cdot 0 = 1$  lifts to the constant  $H_{0,0}$ , which is set to zero in  $\mathfrak{h}$ . Hence  $C_{(1,0),(0,1)}^{0,0} = 0$  in  $\mathfrak{h}$ . Consequently the generator  $u_{0,0}$  is omitted, and the  $u_{0,0}$  degree of freedom is omitted, consistent with constants being removed.
- *Degree-2 calculation.* Take  $K = (1, 1)$ . The differential of  $u_K$  uses outgoing brackets with  $H_K$ :

$$d_{\text{CE}} u_{1,1} = \sum_{c,d} (d - c) u_{c,d} c^{c,d}.$$

Hence the first low-degree terms are

$$d_{\text{CE}} u_{1,1} = 2u_{0,2}c^{0,2} - 2u_{2,0}c^{2,0} - u_{1,0}c^{1,0} + u_{0,1}c^{0,1} + \dots$$

Each term  $C_{(1,1),J}^L u_L c^J$  arises from  $\{H_{1,1}, H_J\} = \sum_L C_{(1,1),J}^L H_L$ . For example,  $\{H_{1,1}, H_{1,0}\} = \partial_{z_1}(z_1 z_2) \cdot 0 - \partial_{z_2}(z_1 z_2) \cdot 1 = -z_1 = -H_{1,0}$ , so  $C_{(1,1),(1,0)}^{1,0} = -1$ , contributing  $-u_{1,0} c^{1,0}$  to  $d_{\text{CE}} u_{1,1}$ . On the open/PV side,

$$d_\pi^{\text{coord}} O_{1,1} = 2O_{0,2}\theta^{0,2} - 2O_{2,0}\theta^{2,0} - O_{1,0}\theta^{1,0} + O_{0,1}\theta^{0,1} + \dots,$$

the same expression with  $u \rightarrow O, c \rightarrow \theta$ . Hence  $\Phi_{\text{coord}}(d_{\text{CE}} u_{1,1}) = d_\pi^{\text{coord}} \Phi_{\text{coord}}(u_{1,1})$ . An independent exact arithmetic check verifies the same structure-constant input on every monomial pair with exponents up to 6 via the first-order Moyal coefficient  $kn - lm$  at  $\hbar^1$ , which equals  $C_{IJ}^K$  for these indices.

- *Higher-order word.* For  $K = (1, 1)$ ,  $J = (1, 0)$ ,  $J' = (0, 1)$ , the cochain element  $u_{1,1}c^{1,0}c^{0,1}$  has degree  $0 + 1 + 1 = 2$ . Its CE differential, by graded Leibniz, is

$$d_{\text{CE}}(u_{1,1}c^{1,0}c^{0,1}) = (d_{\text{CE}}u_{1,1})c^{1,0}c^{0,1} - u_{1,1}(d_{\text{CE}}c^{1,0})c^{0,1} + u_{1,1}c^{1,0}(d_{\text{CE}}c^{0,1}),$$

with each summand applying the generator formulas above. The same word on the PV side is  $O_{1,1}\theta^{1,0}\theta^{0,1}$ , and  $d_{\pi}^{\text{coord}}$  applied with the same Leibniz convention produces the corresponding triple sum with  $u \rightarrow O$  and  $c \rightarrow \theta$ . Term-by-term equality follows from the generator identities; the full word equality is the algebra-homomorphism Leibniz argument from 4.20.

*Construction 5.46 (Compact current PBW/Rees source).* Let  $\mathcal{P}$  denote the principal-part module

$$\mathcal{P} = \bigoplus_{\substack{a,b \geq 0 \\ a+b > 0}} \mathbb{C} \rho_{a,b}, \quad \rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2,$$

identified with  $\mathfrak{h}_{\text{cont}}^{\vee}$  by the residue pairing of Proposition 5.175. Put

$$\mathfrak{g}_{\text{pp}} = \mathfrak{h} \ltimes \mathcal{P}[1],$$

with zero differential and bracket

$$[f, g] = \{f, g\}, \quad [f, \rho[1]] = (f \cdot \rho)[1], \quad [\rho[1], \sigma[1]] = 0,$$

where  $f \cdot \rho = \pi_{\text{pp}}\{f, \rho\}$ . On monomials this action is

$$z_1^p z_2^q \cdot \rho_{a,b} = -(pb - qa + p - q)\rho_{a-p+1, b-q+1},$$

with negative-index targets zero.

For an interval  $I \subset \mathbb{R}$ , define the compactly supported current Lie algebra

$$\mathfrak{Z}_{\mathfrak{g}}(I) = \Omega_c^{\bullet}(I) \widehat{\otimes} \mathfrak{g}_{\text{pp}}, \quad [\alpha \otimes x, \beta \otimes y] = \alpha\beta \otimes [x, y].$$

Its PBW/Rees algebra is

$$U_{\hbar}(\mathfrak{Z}_{\mathfrak{g}}(I)) = \widehat{T}(\mathfrak{Z}_{\mathfrak{g}}(I))[[\hbar]] / \langle xy - (-1)^{|x||y|}yx - \hbar[x, y] \rangle.$$

The PBW filtration is the tensor-word filtration. Its special fibre is

$$U_{\hbar}(\mathfrak{Z}_{\mathfrak{g}}(I)) / (\hbar) \cong \widehat{\mathbf{S}}(\mathfrak{Z}_{\mathfrak{g}}(I)),$$

equipped with the semiclassical  $P_0$  bracket

$$\{\bar{X}, \bar{Y}\}_{P_0} = (\hbar^{-1}(XY - (-1)^{|X||Y|}YX)) \bmod \hbar.$$

This is the PBW/Rees source for the Hamiltonian and principal-part comparison. A closed CE cochain algebra map requires a separate bar-cobar twisting construction or a completed pairing.

*Problem 5.47 (PBW-source  $P_o$ -centre action criterion and CE comparison datum).* Construct a filtered factorization-center action of the compactly supported current PBW/Rees source  $U_h(\mathfrak{S}_g)$  of Construction 5.46. With the cotangent generators included, the target is the enlarged principal-part boundary object of Theorem 5.185. In the classical special fibre the action must send

$$f \mapsto B_f = \overline{\text{Tr } f(\phi_1, \phi_2)}, \quad \rho_{a,b}[1] \mapsto \Theta_{a,b},$$

and satisfy the stalkwise bracket identities

$$\{B_f, B_g\}_{\text{open}} = B_{\{f,g\}}, \quad \{B_f, \Theta_\rho\}_{\text{open}} = \Theta_{f \cdot \rho}, \quad \{\Theta_\rho, \Theta_\sigma\}_{\text{open}} = 0.$$

This special-fibre statement is the natural  $P_o$  theorem. A strict action of the full filtered Rees algebra on the same commutative reduced target requires additional quantized target data: in the source

$$XY - (-1)^{|X||Y|} YX = \hbar[X, Y],$$

while multiplication operators on a commutative classical target commute. Thus the Rees-level theorem requires either a deformed boundary target or an explicit homotopical quantization of the central operations. Consequently the compact current PBW/Rees source supplies a  $P_o$ -central-operation action only after passing to the classical special fibre, or after replacing the boundary target by a genuinely deformed/homotopical central-operation object. The primitive one- $\psi$  projection is an indecomposable-label projection; the filtered Rees action datum is additional. The target data must include differential compatibility, factorization centrality homotopies, coherent  $P_o$  bracket homotopies, and compact-support naturality for interval inclusions and disjoint unions before the de Rham contraction.

Separately compare this PBW/Koszul-dual source with the closed CE observable algebra by an explicit twisting or pairing construction. This comparison is the extra datum that promotes a PBW/Koszul-dual boundary action to a cochain-level map from closed CE observables.

*Problem 5.48 (Unreduced Tate-coefficient cotangent lift datum).* Construct a Tate or weighted-coefficient extension of the BV heat-kernel calculus on  $\mathbb{R}^2 \times \mathbb{C}^2$ . The native formal-coordinate holomorphic singular product has already been computed: its diagonal coefficient module, local-cohomology expansion, and current singular product are Theorem 1.15. The open extension problem is to lift that binary singular product, together with the trace map  $J$ , through the unreduced cotangent sector, Costello heat-kernel coefficients, and the quantum master equation tower. Thus one asks for a coefficient category in which:

- (i) the formal Casimir  $C_{\mathfrak{h}} = \sum_{d \geq 1} \sum_i e_{d,i} \otimes e_d^i$  converges as an element of  $\widehat{\mathfrak{h}} \widehat{\otimes} \widehat{\mathfrak{h}}_{\text{cont}}^\vee$  this is proved by 5.70(a) via the weighted Tate completion of 5.54;
- (ii) the Costello propagator  $P_{\epsilon,L}$  extends to that coefficient category this is proved by 5.70(b);
- (iii) the primitive indecomposable projection of the principal-part current sector

$$\widetilde{\text{Obs}}_{\partial,N}^{\text{cl,pp}}(I) = \text{Obs}_{\text{open},N}^{\text{cl}}(I) \widehat{\otimes} \widehat{\mathbf{S}}(\mathcal{M}_{\partial}^{\text{pp,unred}}(I))$$

admits a chain-level primitive projection  $\pi_{\text{prim}}$  with verified  $Q$ -compatibility and factorization compatibility on the indecomposable quotient the primitive projection is supplied by 5.132; the cotangent centrality homotopies remain part of item (4);

- (iv) the reduced principal-part defect-current theorem of 5.6 lifts to the unreduced factorization-algebra model and the degree-zero Moyal/Weyl theorem extends to the cotangent (one-antifield) sector

at all orders in  $\hbar$  after the primitive projection of 5.139; unreduced cotangent centrality and the one-antifield quantum lift require vanishing of the QME obstruction class of 5.72; the truncated nilpotent sector  $\mathfrak{m}^3/\mathfrak{m}^{N+1}$  closed classically by 5.101 and obstructed quantum-mechanically by 5.102.

Equivalently, lift the cochain-level 4.20 and the Hamiltonian-sector factorization-algebra 4.11 to a cochain-level morphism of factorization algebras

$$\Phi_{\text{cl}} : \text{Obs}_{\text{closed}, H}^{\text{cl}}|_L \longrightarrow Z_{\text{fact}}^{P_o}(\text{Obs}_{\text{open}, N}^{\text{cl}})$$

on the unreduced (cotangent-included) sector, with all  $P_o$ -center homotopies, factorization centrality homotopies, the stable connected quotient, and a boundary realization of the shifted cotangent generators. Any global version must also pass through the local-Hamiltonian descent and period obstruction of 1.29; a stalkwise formal disk map supplies the local input, while Weiss/Ran descent for factorization centers requires the descent datum.

*Weighted Tate completion of the coefficient pair*

The Hamiltonian Lie algebra on the formal symplectic plane  $\mathbb{C}^2$ ,

$$\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1 = \prod_{d \geq 1} H_d, \quad H_d = \text{Span}\{z_1^a z_2^b : a + b = d\},$$

carries the Hamiltonian bracket  $\{H_p, H_q\} \subset H_{p+q-2}$  and the formal Casimir

$$C_{\mathfrak{h}} = \sum_{d \geq 1} \sum_i e_{d,i} \otimes e_d^i$$

in any homogeneous basis  $\{e_{d,i}\}$  of  $H_d$  with dual basis  $\{e_d^i\}$  fixed by  $\langle e_d^j, e_{d,i} \rangle = \delta_i^j$ . Costello's heat-kernel BV calculus on the bulk-plus-defect field complex

$$\mathcal{E} = \Omega^\bullet(\mathbb{R}^2) \widehat{\otimes} \Omega^{\circ, \bullet}(\mathbb{C}^2) \widehat{\otimes} (\mathfrak{h}[1] \oplus \mathfrak{h}_{\text{cont}}^\vee[2])$$

factors the propagator as

$$P_{\epsilon, L} = P_{\epsilon, L}^{\text{base}} \widehat{\otimes} C_{\mathfrak{h} \oplus \mathfrak{h}^\vee[1]},$$

the bulk piece being the standard heat-kernel inverse and the coefficient piece  $C_{\mathfrak{h}}$  extended to the cotangent direction; the propagator exists in  $\mathcal{E} \widehat{\otimes} \mathcal{E}$  exactly when  $C_{\mathfrak{h}}$  exists as a coefficient kernel. In the strict product/direct-sum Tate pair  $(\mathfrak{h}_\Pi, D_\oplus) = (\prod_{d \geq 1} H_d, \oplus_{d \geq 1} H_d^\vee)$  the second factor has support in finitely many degrees, while  $C_{\mathfrak{h}}$  has nonzero  $H_d^\vee$ -component in every degree, so  $C_{\mathfrak{h}} \notin \mathfrak{h}_\Pi \widehat{\otimes} D_\oplus$  (Lemma 5.211; Theorem 5.93 sharpens this to the finite-window form). The coefficient category itself must change: the second factor must be a product, dualized through an intermediate Mittag-Leffler inverse-limit weight system. The smallest complete filtered subspace of  $\text{PV}_{\text{coord}}(\mathcal{A}_{\partial, \text{coord}})$  containing  $\{O_f, \theta_f\}_{f \in \tilde{\mathcal{A}}}$ , closed under  $d_\pi$ , the Schouten bracket  $[-, -]_{\text{Sch}}$ , and the pointwise product, and complete in such a weighted topology is the weighted Tate envelope  $B_w$  of Theorem 5.57.

*Strict Tate pair: Casimir-kernel obstruction*

*Remark 5.49 (Strict Tate continuous duality).* In the strict Tate pair

$$\mathfrak{h}_\Pi = \prod_{d \geq 1} H_d, \quad D_\oplus = \bigoplus_{d \geq 1} H_d^\vee, \tag{19}$$



Lemma 5.165 (1) realizes  $D_\oplus$  as the continuous dual of  $\mathfrak{h}_\Pi$  as Tate vector spaces: the continuous dual of a Tate product is the strict direct sum of duals, and *vice versa*. The second tensor factor of the formal Casimir  $C_{\mathfrak{h}} = \sum_{d \geq 1} \sum_i e_{d,i} \otimes e_d^i$  has nonzero  $H_d^\vee$ -component in every  $d$ , hence lies in  $\prod_d H_d^\vee$ . The direct sum  $\bigoplus_d H_d^\vee = D_\oplus$  contains only finitely supported dual-degree expansions.

*Remark 5.50 (Costello's coefficient requirements).* Costello's heat-kernel BV calculus [12, Ch. 5] writes the BV propagator on a free elliptic complex  $(\mathcal{E}, Q, \langle, \rangle)$  as the kernel of  $\int_\epsilon^L Q^{\text{GF}} e^{-tD} dt$  in  $\mathcal{E} \widehat{\otimes} \mathcal{E}$ . When the field complex factors as

$$\mathcal{E} = \Omega^\bullet(\mathbb{R}^2) \widehat{\otimes} \Omega^{\circ,\bullet}(\mathbb{C}^2) \widehat{\otimes} (\mathfrak{h}[1] \oplus \mathfrak{h}_{\text{cont}}^\vee[2]),$$

with the Tate coefficient pair acting passively on the spacetime, the propagator factors as

$$P_{\epsilon,L} = P_{\epsilon,L}^{\text{base}} \widehat{\otimes} C_{\mathfrak{h} \oplus \mathfrak{h}^\vee[1]}. \quad (20)$$

The spacetime piece  $P_{\epsilon,L}^{\text{base}}$  on  $\Omega^\bullet(\mathbb{R}^2) \widehat{\otimes} \Omega^{\circ,\bullet}(\mathbb{C}^2)$  is the standard bulk BV inverse, continuous as an operator by Remark 5.210. The coefficient piece is the BV inverse pairing on  $\mathfrak{h} \oplus \mathfrak{h}_{\text{cont}}^\vee[1]$ , i.e. the Casimir  $C_{\mathfrak{h}}$  together with its dual coevaluation. The full kernel therefore exists in  $\mathcal{E} \widehat{\otimes} \mathcal{E}$  exactly when  $C_{\mathfrak{h}}$  exists as a coefficient kernel.

Costello's heat-kernel BV requires three properties of the coefficient module:

- (C1) a distributional inverse pairing, i.e. a coefficient coevaluation  $C_{\mathfrak{h}} \in \mathfrak{h} \widehat{\otimes} \mathfrak{h}_{\text{cont}}^\vee$  realizing the bilinear form with kernel zero in the cohomology of the heat operator;
- (C2) regularity in the cutoff parameters  $\epsilon$  and  $L$ , in the sense that the heat-kernel propagator  $P_{\epsilon,L}$  is a continuous family in the coefficient topology and converges as  $L \rightarrow \infty, \epsilon \rightarrow 0^+$  to a distributional kernel after counterterm subtraction;
- (C3) compatibility with the Costello RG flow  $W(P_{\epsilon,L}, I)$ : the weight or topology defining the coefficient module is preserved by the renormalization flow on effective interactions induced by varying  $(\epsilon, L)$ , in the precise sense that the source of the flow lies in the same completed coefficient category as its target.

The strict Tate pair (19) has the obstruction to (C1). Requirement (C3) is the residual locality/QME problem attached to coefficient enlargement: Problem 5.72.

#### *Degree weights and the canonical completion $B_w$*

A degree weight on  $\mathfrak{h}$  selects a coefficient category closed under the Schouten differential  $d_\pi$ , the Schouten bracket  $[-, -]_{\text{Sch}}$ , and the pointwise product, while hosting the Casimir kernel: bracket-tame for the Schouten bracket, windowwise coadjoint-continuous for the cotangent action, and Mittag-Leffler-coevaluating for the heat-kernel propagator.

*Definition 5.51 (Polynomial Hamiltonian degree weight).* A degree weight is a function  $w: \mathbb{Z}_{\geq 1} \rightarrow \mathbb{R}_{>0}$  satisfying:

- (W1) *Bracket-tame*: there exists a constant  $C_w \geq 0$  such that  $w(p+q-2) \leq C_w w(p)w(q)$  for every  $p, q \geq 2$ ;
- (W2) *Windowwise coadjoint-continuity*: for every Hamiltonian degree bound  $R$  and every output Tate window  $w_o$ , the principal-part coadjoint shifts  $\rho_{a,b} \mapsto \rho_{a-p+1, b-q+1}$  with  $p+q \leq R$  and  $a+b-p-q+2 \leq w_o$  involve only the finite input window  $a+b \leq w_o + R - 2$ , and the finitely many corresponding weight ratios have a maximum;

(W<sub>3</sub>) *Mittag-Leffler coevaluation*: for each  $w_o$  the finite-window Casimir  $\sum_{d \leq w_o} \sum_i w(d) e_{d,i} \otimes w(d)^{-1} e_d^i$  is compatible under the surjective truncation maps  $w'_o \rightarrow w_o$ . Equivalently, the coefficient kernel is a strict Mittag-Leffler inverse system of finite-dimensional kernels.

LEMMA 5.52 (*Existence of weights*). Every exponential weight  $w(d) = q^d$  with  $q > 1$  satisfies (W<sub>1</sub>)–(W<sub>3</sub>). The proof uses only finite-window maxima for the coadjoint action; the unbounded principal-part degree prevents a uniform bound.

*Proof.* For  $w(d) = q^d$ , (W<sub>1</sub>) reads  $q^{p+q-2} \leq C_w q^{p+q}$ , i.e.  $C_w = q^{-2}$  suffices. For (W<sub>2</sub>), fix  $R$  and  $w_o$ . If  $p + q \leq R$  and the output degree is  $a + b - p - q + 2 \leq w_o$ , then  $a + b \leq w_o + R - 2$ . Hence the set of possible triples  $(p, q, a + b)$  is finite, and the corresponding ratios of powers of  $q$  admit a finite maximum; the bound depends on the finite-window restriction. Condition (W<sub>3</sub>) is the strict Mittag-Leffler compatibility of finite-window Casimirs under truncation. The transition maps are surjective, hence the system is Mittag-Leffler.  $\square$

Remark 5.53 (*Choice of exponential weight*). The exponential weight  $w(d) = q^d$  is the working choice, with  $q > 1$  formally an element of  $\mathbb{R}_{>0}$  and concretely set to  $q = 2$  for the proofs below. The choice is a grading weight on the coefficient module. It acts on coefficients, with the formal polydisk fixed. Thus the Poisson bracket and the counterterm representatives are compared between two weights only after transport by the diagonal map  $R_{w,w'}$  below. In coordinates, the bracket constants in the basis  $w(d) e_{d,i}$  are conjugated by the finite ratios  $w(p)w(q)/w(p+q-2)$ , and condition (W<sub>1</sub>) is exactly the boundedness condition needed for this transported bracket on each finite graph. The weight is therefore a controlled coefficient regulator; rescaling the variables  $z_1, z_2$  is a different operation on the original Hamiltonian bracket.

Definition 5.54 (*Weighted Tate coefficient pair*). Given a degree weight  $w$  as in Definition 5.51, write  $H_d^{(w)}$  for the rescaled copy of the finite-dimensional degree piece  $H_d$  with basis  $w(d) e_{d,i}$ , and  $(H_d^\vee)^{(w^{-1})}$  for the rescaled dual basis  $w(d)^{-1} e_d^i$ . Define

$$\mathfrak{h}_w = \varprojlim_{w_o \in \mathbb{Z}_{\geq 1}} \prod_{d=1}^{w_o} H_d^{(w)}$$

with the Tate filtration  $F^{w_o} \mathfrak{h}_w = \ker(\mathfrak{h}_w \rightarrow \prod_{d \leq w_o} H_d^{(w)})$ , and define the *weighted coefficient dual*

$$\mathfrak{h}_{\text{cont}}^{\vee, w} = \varprojlim_{w_o \in \mathbb{Z}_{\geq 1}} \prod_{d=1}^{w_o} (H_d^\vee)^{(w^{-1})}.$$

The pairing is fixed degreewise by  $\langle w(d)^{-1} e_d^i, w(d) e_{d,j} \rangle = \delta_j^i$ . The weighted Tate coefficient pair is  $(\mathfrak{h}_w, \mathfrak{h}_{\text{cont}}^{\vee, w})$ . The second factor is a product coefficient module replacing the strict continuous dual of the first product Tate space.

Remark 5.55 (*Topological content*). Both  $\mathfrak{h}_w$  and  $\mathfrak{h}_{\text{cont}}^{\vee, w}$  are Mittag-Leffler inverse limits of finite-dimensional Tate windows, and both are products in the degree direction. The construction replaces the direct-sum dual by a symmetric Mittag-Leffler product coefficient module. This is an additional coefficient category for the BV kernel; the ordinary continuous dual of  $\mathfrak{h}_w$  remains the strict Tate dual.

*Definition 5.56 (The weighted-Tate envelope  $B_w$ ).* Inside the bracket-admissible polyvector subspace  $B \subset \text{PV}_{\text{coord}}(\mathcal{A}_{\partial, \text{coord}})$  of Definition 4.18, the weighted Tate envelope is

$$B_w := \overline{\langle \{O_f, \theta_f\}_{f \in \tilde{A}} \rangle_{d_\pi, [-, -]_{\text{Sch}}}}^{\mathfrak{h}_w \widehat{\otimes} \mathfrak{h}_{\text{cont}}^{\vee, w}},$$

the smallest complete filtered subspace of  $\text{PV}_{\text{coord}}(\mathcal{A}_{\partial, \text{coord}})$  containing the brane-side generators  $\{O_f, \theta_f\}_{f \in \tilde{A}}$ , closed under the Schouten differential  $d_\pi$ , the Schouten bracket  $[-, -]_{\text{Sch}}$ , and the pointwise product, and complete in the weighted Tate topology of  $\mathfrak{h}_w \widehat{\otimes} \mathfrak{h}_{\text{cont}}^{\vee, w}$ . The polyvector grading on  $B_w$  is the Schouten degree; the coefficient grading is the polynomial degree carried by the weight  $w$ .

*THEOREM 5.57 (Canonical weighted-Tate habitat).* Let  $w$  be a degree weight as in Definition 5.51 and let  $B_w$  be the envelope of Definition 5.56. Then  $B_w$  is the minimal complete filtered subspace of  $\text{PV}_{\text{coord}}(\mathcal{A}_{\partial, \text{coord}})$  in which Costello's heat-kernel BV calculus and the brane-side Schouten algebra on  $\{O_f, \theta_f\}_{f \in \tilde{A}}$  both close.

- (a) *Closure under the Schouten algebra.*  $B_w$  is closed under multiplication, Schouten differential, and Schouten bracket:

$$B_w \cdot B_w \subset B_w, \quad d_\pi B_w \subset B_w, \quad [B_w, B_w]_{\text{Sch}} \subset B_w.$$

- (b) *Casimir convergence.* The diagonal-rescaled Casimir

$$C_{\mathfrak{h}, w} = \sum_{d \geq 1} \sum_i (w(d) e_{d,i}) \otimes (w(d)^{-1} e_d^i)$$

converges in the weighted continuous tensor product  $\mathfrak{h}_w \widehat{\otimes} \mathfrak{h}_{\text{cont}}^{\vee, w}$ . Under the  $P_o$ -algebra map  $e_{d,i} \mapsto O_{e_{d,i}}, e_d^i \mapsto \theta_d^i$  of Theorem 4.20, the kernel lies in  $B_w \widehat{\otimes} B_w$  and realizes Costello requirement (C1) as the BV inverse pairing on the cotangent extension.

- (c) *Heat-kernel propagator extension.* The Costello heat-kernel propagator

$$P_{\epsilon, L}^w = P_{\epsilon, L}^{\text{base}} \widehat{\otimes} C_{\mathfrak{h} \oplus \mathfrak{h}^\vee[1], w}$$

extends from the strict Tate field complex to the weighted complex  $\mathcal{E}_w$  as a continuous family for  $0 < \epsilon \leq L < \infty$ , satisfying the BV identity  $[Q, P_{\epsilon, L}^w] = K_\epsilon^w - K_L^w$  and Costello regularity (C2): continuity in  $(\epsilon, L)$  in the weighted Tate topology, and existence of the  $L \rightarrow \infty, \epsilon \rightarrow 0^+$  limit as a distributional kernel after the standard bulk counterterm subtraction. The coefficient piece  $C_{\mathfrak{h}, w}$  is counterterm-free; it is the fixed Mittag-Leffler kernel of (b).

- (d) *Graphwise RG stability.* For every fixed Costello graph  $\Gamma$  and every  $0 < \epsilon \leq L < \infty$ , the contribution  $\mathcal{W}_\Gamma(P_{\epsilon, L}^w, I)$  of any vertexwise polynomial-degree-bounded interaction  $I \in \mathcal{O}_{\text{loc}}(\mathcal{E}_w)$  stays inside  $B_w$ . Each fixed graph is bounded by a finite constant times the corresponding spacetime graph weight, with constants supplied by (W1) and (W2).
- (e) *Cochain-level CE/PV extension.* The reduced central-operation cochain  $\Phi$  of Theorem 4.20 extends to the weighted cotangent extension  $\mathfrak{g}_w = \mathfrak{h}_w \ltimes \mathfrak{h}_{\text{cont}}^{\vee, w}[1]$  as a continuous cochain map landing in  $B_w$ :

$$\Phi^w : C_{\text{CE, cont}}^\bullet(\mathfrak{g}_w) \longrightarrow B_w \subset \text{PV}(\widehat{\mathfrak{S}}_{\text{Tate}}(\mathfrak{h}_w)).$$

Conversely, any complete filtered subspace  $B' \subset \text{PV}_{\text{coord}}(\mathcal{A}_{\partial, \text{coord}})$  satisfying (a), containing  $\{O_f, \theta_f\}_{f \in \tilde{A}}$ , and containing  $C_{\mathfrak{h}, w} \in B' \widehat{\otimes} B'$  as a continuous element of the weighted Tate topology, contains  $B_w$ .

*Proof.* (a): with the linear Kirillov Poisson tensor  $\pi = \frac{1}{2}C_{ij}^k O_k \theta^i \theta^j$  of Theorem 4.20, the Schouten differential  $d_\pi = [\pi, -]_S$  acts on generators by

$$d_\pi \theta^k = -\frac{1}{2}C_{ij}^k \theta^i \theta^j, \quad d_\pi O_k = C_{kj}^\ell O_\ell \theta^j,$$

with  $C_{ij}^k$  the Hamiltonian structure constants on  $\mathfrak{h}$ . The bracket-tame estimate (W1) bounds, on each finite Tate window, the polynomial-degree shift produced by a single  $d_\pi$  or a single Schouten bracket by a finite constant; the windowwise coadjoint-continuity (W2) bounds the cotangent-leg shift. Multiplication is closed by definition of the envelope. The closure  $\overline{(\cdot)}$  in Definition 5.56 adjoins all  $d_\pi$ -,  $[-, -]_{\text{Sch}}$ -, and product-completions, and that closure is itself stable under the same operations because each is continuous in the weighted Tate topology by (W1)–(W2).

(b): this is Proposition 5.61 below.

(c): this is Proposition 5.63.

(d): this is Proposition 5.68.

(e): the cochain identity  $\Phi d_{\text{CE}} = d_\pi \Phi$  is degreewise algebraic on  $\mathfrak{h} \oplus \mathfrak{h}^\vee[1]$  and is preserved by the diagonal weight rescaling. The image of generators  $c_f, u_f \in C_{\text{CE}}^\bullet(\mathfrak{g}_w)$  is, by the  $P_o$ -algebra map convention,  $\theta_f, O_f \in B_w$ ; closure of  $B_w$  under  $d_\pi$ ,  $[-, -]_{\text{Sch}}$ , and product (item (a)) makes  $\Phi^w$  land in  $B_w$ . The continuity for the weighted Tate topology is the same windowwise estimate as in (d).

Minimality: any  $B'$  satisfying (a) and containing  $\{O_f, \theta_f\}$  contains all  $d_\pi$ -, bracket-, and product-iterates of those generators, hence the algebraic envelope. Containing  $C_{\mathfrak{h},w} \in B' \widehat{\otimes} B'$  as a continuous element forces  $B'$  to be complete in the weighted Tate topology, hence to contain the closure  $B_w$ .  $\square$

*Definition 5.58 (Category of admissible weighted-Tate completions).* Fix a degree weight  $w$  as in Definition 5.51. The category  $\text{Adm}_w$  has objects: complete filtered subspaces  $B' \subset \text{PV}_{\text{coord}}(\mathcal{A}_{\partial, \text{coord}})$  satisfying

- (A1) (*generation*)  $B'$  contains the brane-side generators  $\{O_f, \theta_f\}_{f \in \vec{A}}$ ;
- (A2) (*Schouten closure*)  $B' \cdot B' \subset B'$ ,  $d_\pi B' \subset B'$ , and  $[B', B']_{\text{Sch}} \subset B'$ ;
- (A3) (*Casimir convergence*) the weighted Casimir  $C_{\mathfrak{h},w}$  of (21) converges in  $B' \widehat{\otimes} B'$  in the weighted Tate topology;
- (A4) (*heat-kernel propagator extension*) the Costello heat-kernel propagator  $P_{\epsilon,L}^w = P_{\epsilon,L}^{\text{base}} \widehat{\otimes} C_{\mathfrak{h} \oplus \mathfrak{h}^\vee[1],w}$  extends to a continuous family for  $0 < \epsilon \leq L < \infty$  on  $B'$  satisfying  $[Q, P_{\epsilon,L}^w] = K_\epsilon^w - K_L^w$ ;
- (A5) (*graphwise RG boundedness*) for every Costello graph  $\Gamma$  and every vertexwise polynomial-degree bounded interaction  $I \in \mathcal{O}_{\text{loc}}(\mathcal{E}_w)$ , the contribution  $\mathcal{W}_\Gamma(P_{\epsilon,L}^w, I)$  stays in  $B'$  with constants supplied by (W1) and (W2).

Morphisms  $B' \hookrightarrow B''$  are continuous filtered inclusions containing the generators on the nose and intertwining  $d_\pi$ ,  $[-, -]_{\text{Sch}}$ , the pointwise product, the weighted Casimir, and the heat-kernel propagator.

**COROLLARY 5.59** ( $B_w$  is initial in  $\text{Adm}_w$ ).  $B_w$  of Definition 5.56 is the initial object of  $\text{Adm}_w$ : every  $B' \in \text{Adm}_w$  admits a unique morphism  $\iota_{B'} : B_w \hookrightarrow B'$ .

*Proof.* By Theorem 5.57,  $B_w \subset B'$  for every  $B' \in \text{Adm}_w$ , and the inclusion is an  $\text{Adm}_w$ -morphism. A morphism  $B_w \rightarrow B'$  is determined on  $\{O_f, \theta_f\}_{f \in \vec{A}}$ , hence on the Schouten-bracket-product envelope, hence by Tate continuity on  $B_w$ .  $\square$

*Remark 5.60.*  $\text{Adm}_w$  depends on  $w$ ; its admissibility conditions are intrinsic to  $(\mathfrak{h}_w, \mathfrak{h}_{\text{cont}}^{\vee,w}, P_{\epsilon,L}^w)$ .

PROPOSITION 5.61 (*Casimir convergence*). In the weighted Tate coefficient pair  $(\mathfrak{h}_w, \mathfrak{h}_{\text{cont}}^{\vee, w})$  of Definition 5.54, the formal Casimir

$$C_{\mathfrak{h}, w} = \sum_{d \geq 1} \sum_i (w(d) e_{d,i}) \otimes (w(d)^{-1} e_d^i) \quad (21)$$

converges as an element of  $\mathfrak{h}_w \widehat{\otimes} \mathfrak{h}_{\text{cont}}^{\vee, w}$ . The pairing of  $C_{\mathfrak{h}, w}$  with any test functional in the dual is the identity on the corresponding finite-dimensional Tate window, so  $C_{\mathfrak{h}, w}$  is the BV coefficient inverse pairing in the weighted category.

*Proof.* The displayed Casimir, restricted to the Tate window  $d \leq w_o$ , is the finite sum  $\sum_{d \leq w_o} \sum_i (w(d) e_{d,i}) \otimes (w(d)^{-1} e_d^i)$ , an element of  $\prod_{d \leq w_o} w(d) H_d \otimes \prod_{d \leq w_o} w(d)^{-1} H_d^\vee$ . This is a tensor in a finite-dimensional vector space because each window is finite-dimensional. The tower of finite-window Casimirs is compatible: enlarging  $w_o$  extends the sum by the terms in degrees  $w_o + 1, \dots, w'_o$ . The inverse limit in **TateVec** exists by Definition 5.54 and (W3) of Definition 5.51.

Pairing the Casimir with a test functional  $\phi \in \mathfrak{h}_{\text{cont}}^{\vee, w}$ , supported in finitely many Tate windows, gives a finite sum of identity pairings, so the result is the canonical evaluation. Hence  $C_{\mathfrak{h}, w}$  realizes the inverse pairing on the weighted Tate pair, satisfying Costello requirement (C1).  $\square$

*Remark 5.62 (Independence of basis).* Definition (21) is written in a basis  $\{e_{d,i}\}$  of  $H_d$ , and the tensor is basis-independent. The dual basis  $\{e_d^i\}$  is fixed by  $\langle e_{d,i}, e_d^j \rangle = \delta_i^j$ . An orthogonal change of basis  $e_{d,i} \mapsto \sum_j O_i^j e_{d,j}$  with  $O \in GL(H_d)$  shifts the dual by the inverse transpose, yielding the same Casimir. Thus  $C_{\mathfrak{h}, w} = \sum_d \text{id}_{H_d \otimes H_d^\vee}$  is the basis-independent identity element of  $\bigoplus_d H_d \otimes H_d^\vee$  extended to the weighted product.

*Heat-kernel propagator on the weighted pair*

PROPOSITION 5.63 (*Heat-kernel propagator extension to the weighted pair*). Let  $w$  be a degree weight as in Definition 5.51. The Costello heat-kernel propagator  $P_{\epsilon, L}^{\text{op}}$  of Remark 5.210 extends from the strict Tate field complex to the weighted field complex

$$\mathcal{E}_w = \Omega^\bullet(\mathbb{R}^2) \widehat{\otimes} \Omega^{\circ, \bullet}(\mathbb{C}^2) \widehat{\otimes} (\mathfrak{h}_w[1] \oplus \mathfrak{h}_{\text{cont}}^{\vee, w}[2]), \quad (22)$$

to give a continuous family of BV propagators  $P_{\epsilon, L}^w \in \mathcal{E}_w \widehat{\otimes} \mathcal{E}_w$  for  $0 < \epsilon \leq L < \infty$ , satisfying:

- (1) the BV identity  $[Q, P_{\epsilon, L}^w] = K_\epsilon^w - K_L^w$ , where  $K_t^w = K_t^{\text{base}} \widehat{\otimes} C_{\mathfrak{h}, w}$  is the weighted heat kernel of the BV inverse pairing;
- (2) Costello regularity (C2): the family is continuous in  $(\epsilon, L)$  in the weighted Tate topology;
- (3) the limit  $L \rightarrow \infty, \epsilon \rightarrow 0^+$  exists as a distributional kernel after the standard spacetime counterterm subtraction in Costello's heat-kernel renormalization-group construction [12, Chs. 5–7]; the coefficient direction leaves the spacetime singular support unchanged. Locality and QME counterterm compatibility for the mixed brane-defect theory remain the content of Problem 5.72.

*Proof.* By the factorization (20), the propagator splits as a tensor product of the spacetime kernel  $P_{\epsilon, L}^{\text{base}}$  on  $\Omega^\bullet(\mathbb{R}^2) \widehat{\otimes} \Omega^{\circ, \bullet}(\mathbb{C}^2)$  and the coefficient Casimir  $C_{\mathfrak{h} \oplus \mathfrak{h}^\vee[1], w}$  on the weighted Tate pair. The spacetime piece is unchanged by the weight on the coefficient direction; it is the Costello heat-kernel inverse on the bulk free elliptic complex  $\Omega^\bullet(\mathbb{R}^2) \widehat{\otimes} \Omega^{\circ, \bullet}(\mathbb{C}^2)$ , constructed in [12, Ch. 5]. The BV identity for  $P_{\epsilon, L}^{\text{base}}$

is standard: with  $Q^{\text{base}} = d_{\mathbb{R}^2} + \bar{\partial}_{\mathbb{C}^2}$ , gauge-fixing  $Q^{\text{GF}} = d_{\mathbb{R}^2}^* + \bar{\partial}_{\mathbb{C}^2}^*$ , and the associated Laplacian  $\Delta = [Q^{\text{base}}, Q^{\text{GF}}]$ , one has

$$[Q^{\text{base}}, P_{\epsilon, L}^{\text{base}}] = \int_{\epsilon}^L [Q^{\text{base}}, Q^{\text{GF}} e^{-t\Delta}] dt = \int_{\epsilon}^L \partial_t (-e^{-t\Delta}) dt = e^{-\epsilon\Delta} - e^{-L\Delta} = K_{\epsilon}^{\text{base}} - K_L^{\text{base}}.$$

The coefficient piece is the Casimir of Proposition 5.61, which is annihilated by the trivial differential on  $\mathfrak{h}_w[1] \oplus \mathfrak{h}_{\text{cont}}^{\vee, w}[2]$ . Therefore  $[Q, P_{\epsilon, L}^w] = K_{\epsilon}^w - K_L^w$ , and the family is continuous in  $(\epsilon, L)$  because  $C_{\mathfrak{h}, w}$  is a fixed element of the weighted pair and the bulk family is continuous. Item (3), at the level asserted here, is the Costello bulk regularity results [12, Chs. 5–7] tensored with a fixed coefficient kernel. This proves finite-scale kernel regularity and absence of new coefficient wave-front directions. The brane-defect QME obstruction classes are separate local-functional classes.  $\square$

*Remark 5.64 (Brane defect).* The brane defect  $L = \mathbb{R}_{\text{brane}} \times \{0\} \times \{p\} \subset \mathbb{R}^2 \times \mathbb{C}^2$  pulls the weighted field complex  $\mathcal{E}_w$  to its restriction along the defect:

$$\mathcal{E}_w^{\partial} = \Omega^{\bullet}(\mathbb{R}) \widehat{\otimes} (\mathfrak{h}_w[1] \oplus \mathfrak{h}_{\text{cont}}^{\vee, w}[2]).$$

The boundary propagator  $P_{\epsilon, L}^{w, \partial} \in \mathcal{E}_w^{\partial} \widehat{\otimes} \mathcal{E}_w^{\partial}$  is the bulk-to-boundary restriction of  $P_{\epsilon, L}^w$  followed by the zero-mode-removed midpoint propagator  $G(t, s) = \frac{1}{2} \text{sgn}(t - s)$  of Proposition 5.222. The weighted Casimir  $C_{\mathfrak{h}, w}$  is the same on the boundary as in the bulk; the weight  $w$  is independent of the coordinate  $t$ .

#### Normal $\Omega$ -weights and equivariant kernel admissibility

The brane-preserving equivariant deformation rescales the normal bundle  $N_L X = \mathbb{R}_s \oplus \mathbb{C}_{z_1} \oplus \mathbb{C}_{z_2}$  by the torus  $T_{\Omega} = \mathbb{C}_{\epsilon_s}^* \times \mathbb{C}_{\epsilon_1}^* \times \mathbb{C}_{\epsilon_2}^*$ , with  $t$  fixed. The equivariant differential is  $Q_{\Omega} = Q + \iota_{V_{\Omega}}$  for the generating vector field  $V_{\Omega} = \epsilon_s s \partial_s + \epsilon_1 z_1 \partial_{z_1} + \epsilon_2 z_2 \partial_{z_2}$ , with  $Q_{\Omega}^2 = L_{V_{\Omega}}$ . The weighted-Tate envelope  $B_w$  extends to a stratified factorization-algebra over  $(X, L)$  by a diagonal equivariant Casimir; the residue-versus-Euler normalization enters as a one-dimensional scalar comparison map, and resonance among the normal characters is the exact obstruction to localizing the contraction.

*Definition 5.65 (Equivariant finite normal window).* Fix normal equivariant parameters  $\epsilon_s, \epsilon_1, \epsilon_2$ , with  $t$  fixed and  $s, z_1, z_2$  carrying weights  $\epsilon_s, \epsilon_1, \epsilon_2$ . For a finite normal/Hamiltonian window  $(N, M)$ , write

$$W_{N, M} = \{(n, a, b) \mid 0 \leq n \leq N, a, b \geq 0, 1 \leq a + b \leq M\}.$$

The integer  $n$  is the total  $s$ -weight of the homogeneous normal summand; if a de Rham  $ds$ -factor is present, its weight is included in  $n$ . For the Hamiltonian monomial  $s^n z_1^a z_2^b$  the normal character is

$$\chi_{n, a, b}^H = n\epsilon_s + a\epsilon_1 + b\epsilon_2 \quad (23)$$

For the residue-dual principal-part label with the same  $s$ -weight, the holomorphic part has the opposite character:

$$\chi_{n, a, b}^{\vee} = n\epsilon_s - a\epsilon_1 - b\epsilon_2 \quad (24)$$

Let  $X_{N, M}$  be the finite set of nonzero formal characters among  $\chi_{n, a, b}^H$  and  $\chi_{n, a, b}^{\vee}$  for those homogeneous summands which the normal Koszul contraction is required to move. The finite-window equivariant coefficient ring is

$$R_{\Omega}^{N, M} = \mathbb{C}[\epsilon_s, \epsilon_1, \epsilon_2, \hbar_{\omega}][\chi^{-1} \mid \chi \in X_{N, M}], \quad \text{wt}(\hbar_{\omega}) = \epsilon_1 + \epsilon_2$$

A scalar specialization of  $(\epsilon_s, \epsilon_1, \epsilon_2)$  is  $(N, M)$ -nonresonant if every  $\chi \in X_{N,M}$  remains nonzero. It is all-window nonresonant if this holds for every  $(N, M)$ . An analytic Kothe norm in place of the algebraic Mittag–Leffler product topology requires the stronger  $q$ -moderate nonresonance condition is required: there are constants  $A, Q$  such that

$$|\chi_{n,a,b}|^{-1} \leq A Q^{n+a+b}$$

for every inverted character. This quantitative bound upgrades the finite-window algebraic kernels from nonresonance to an analytic all-window bounded homotopy in a fixed normed scale.

**THEOREM 5.66** (*Equivariant weighted-kernel admissibility*). Let  $w$  be a degree weight satisfying Definition 5.51. Work in a finite normal window  $(N, M)$  over  $R_\Omega^{N,M}$ , and let  $A_{N,M}$  be a homogeneous basis of the corresponding Hamiltonian-plus-cotangent coefficient pair. For  $\alpha = (n, a, b, \tau) \in A_{N,M}$ , put  $d(\alpha) = a + b$ , let  $e_{\alpha,w} = w(d(\alpha))e_\alpha$ , let  $e_{w^{-1}}^\alpha = (w(d(\alpha)))^{-1}e^\alpha$ , and let  $\chi_\alpha$  be the character (23) for Hamiltonian labels and (24) for cotangent labels.

Then the equivariant diagonal coefficient Casimir

$$C_{\Omega,w}^{N,M} = \sum_{\alpha \in A_{N,M}} e_{\alpha,w} \otimes e_{w^{-1}}^\alpha$$

is a finite-window kernel, and these kernels are compatible under the normal/Hamiltonian truncation maps. Thus the inverse system of  $C_{\Omega,w}^{N,M}$ 's is a strict Mittag–Leffler diagonal Casimir in the weighted coefficient category.

Suppose moreover that on each moving homogeneous normal summand there is a finite-dimensional Koszul numerator  $k_\alpha^{\text{Kos}}$  with

$$Q_\Omega k_\alpha^{\text{Kos}} + k_\alpha^{\text{Kos}} Q_\Omega = \chi_\alpha \text{id}_\alpha$$

before division by the character. Then the normal contracting kernel

$$H_{\Omega,w}^{N,M} = \sum_{\alpha \in A_{N,M}^{\text{mov}}} \chi_\alpha^{-1} e_{\alpha,w} \otimes e_{w^{-1}}^\alpha \otimes k_\alpha^{\text{Kos}} \quad (25)$$

is a well-defined element of the finite-window weighted kernel module and satisfies

$$Q_\Omega H_{\Omega,w}^{N,M} + H_{\Omega,w}^{N,M} Q_\Omega = \text{id} - \text{pr}_{\chi=0}$$

on that window. A character with specialized value zero is a genuine resonant summand. It is retained in  $\text{pr}_{\chi=0}$ , and the contraction is defined on the moving summand.

The residue and Euler normalizations leave the diagonal Casimir fixed. They change the brane pushforward/scalar-contact normalization. With orientation sign  $\sigma_s = \pm 1$ ,

$$\nu_\Omega^{\text{res}} = 1, \quad \nu_\Omega^{\text{Eu}} = \sigma_s (\epsilon_s \epsilon_1 \epsilon_2)^{-1}.$$

A normalized equivariant kernel row is therefore  $K_{\Omega,w,\nu}^{N,M} = \nu_\Omega K_{\Omega,w}^{N,M}$ . The comparison between residue and Euler conventions is the scalar map

$$K_{\Omega,w,\text{res}}^{N,M} \longmapsto \nu_\Omega^{\text{Eu}} K_{\Omega,w,\text{res}}^{N,M}.$$

This factor is required when both conventions appear in one scalar-contact or trace statement; omitting it is exactly the normalization obstruction.

*Proof.* The Casimir is finite in the window  $(N, M)$ . The two diagonal weights  $w(d(\alpha))$  and  $(w(d(\alpha)))^{-1}$  cancel in the evaluation pairing, so the finite sum is the identity tensor on the corresponding finite-dimensional coefficient quotient. Truncating from  $(N, M)$  to a smaller window simply deletes the omitted homogeneous labels, hence sends  $C_{\Omega, w}^{N, M}$  to the smaller-window Casimir. This proves strict Mittag–Leffler compatibility.

For a moving homogeneous summand, the displayed Koszul identity gives

$$Q_{\Omega}(\chi_{\alpha}^{-1}k_{\alpha}^{\text{Kos}}) + (\chi_{\alpha}^{-1}k_{\alpha}^{\text{Kos}})Q_{\Omega} = \text{id}_{\alpha}$$

after adjoining  $\chi_{\alpha}^{-1}$  to the coefficient ring. Summing over the finite set of moving labels gives (25) and the homotopy identity with the unmoved zero-character projection. If a specialized character vanishes,  $\chi_{\alpha}^{-1}$  lies outside the specialized coefficient ring, so resonance is precisely the obstruction to moving that summand.

The residue convention fixes the formal residue density; the Euler convention pushes along the normal bundle with inverse equivariant Euler class  $\sigma_5(\epsilon, \epsilon_1, \epsilon_2)^{-1}$ . The coefficient Casimir pairs dual bases before either pushforward is applied, so it is unchanged. The two conventions differ only in the one-dimensional scalar localization line: scalar-contact rows and trace rows are multiplied by the chosen  $\nu_{\Omega}$ , which is the displayed comparison map.  $\square$

**COROLLARY 5.67** (*Equivariant Costello QME obstruction vector*). Theorem 5.66 supplies the coefficient Casimir and the localized normal contracting kernel in the weighted finite-window category, the input needed to form an equivariant Costello kernel complex. Once equivariant brane-defect Costello data are supplied, in the sense of Definition E.41, the exact obstruction vector to the Costello QME is

$$\text{Ob}_{\Omega, w}^{\text{QME}} = \left( \mathcal{R}_{\Omega}, \text{ob}_{\Omega, \nu}, \{\mathfrak{s}_{n, \Omega}\}_{n \geq 1}, \{([\mathfrak{p}_{n, \Omega, M}^{\text{ns}}])_M, \lambda_{n, \Omega}\}_{n \geq 1} \right).$$

Here

$$\mathcal{R}_{\Omega}^{N, M} = \{\alpha \in A_{N, M}^{\text{mov}} \mid \chi_{\alpha} = \mathfrak{o}\}$$

is the resonance obstruction to the normal contraction,  $\text{ob}_{\Omega, \nu}$  is the residue/Euler normalization obstruction,  $\mathfrak{s}_{n, \Omega, M}$  is the scalar condition

$$\widehat{\sigma}_{\text{sc}, \Omega, M}(R_{n, \Omega, M}^{\text{ns}}),$$

and

$$([\mathfrak{p}_{n, \Omega, M}^{\text{ns}}])_M \in \varprojlim_M H^1(\mathcal{K}_{\text{ns}, \Omega, M}^{\bullet}(I)), \quad \lambda_{n, \Omega} \in \varprojlim_M^1 H^0(\mathcal{K}_{\text{ns}, \Omega, M}^{\bullet}(I))$$

are the non-scalar cohomology and primitive-compatibility obstructions. The equivariant weighted kernel data determine a Costello brane-defect QME solution exactly on the common zero locus of this vector, and only there.

*Proof.* Theorem 5.66 supplies only  $C_{\Omega, w}$ , the normal denominators, and the normalization line. Costello’s QME is the Maurer–Cartan equation for the Hom-valued brane-defect graph kernel plus local counterterms. Therefore, after the equivariant datum is present, the scalar projection must first be a chain projection and its scalar condition must vanish. Only then is the residual a class in the non-scalar kernel complex, and the finite-window counterterm criterion is the same inverse-limit  $H^1$  and  $\varprojlim^1 H^0$  criterion as Theorem 5.83, with  $Q, d_M, \widehat{\sigma}_{\text{sc}}$  replaced by their equivariant versions. Resonant normal modes and normalization mismatch occur before this cohomology class is even formed. Hence the displayed vector is necessary and sufficient for the equivariant weighted-kernel construction to become a QME solution.  $\square$



*Renormalization-flow compatibility*

The Costello RG flow on effective interactions is the assignment  $I_L = W(P_{\epsilon,L}, I)$ , where  $W$  is the homotopy renormalization group action on functionals [12, Ch. 6]. Costello requirement (C<sub>3</sub>) demands that the RG flow on  $\mathcal{E}_w$ -valued interactions preserve the weighted coefficient module: a polynomial-degree-bounded interaction in  $\mathcal{O}(\mathcal{E}_w)$  must produce  $I_L$  in the same weighted category.

**PROPOSITION 5.68** (*Graphwise RG-tame interactions*). Let  $I \in \mathcal{O}_{\text{loc}}(\mathcal{E}_w)$  be a local interaction polynomial in the fields, of total polynomial-degree bounded by  $D_o$  at each vertex in the coefficient direction. For every fixed Costello graph  $\Gamma$  and every  $0 < \epsilon \leq L < \infty$ , the graph contribution  $W_\Gamma(P_{\epsilon,L}^w, I)$  has finite coefficient degree bounded by  $D_o V(\Gamma) - 2E(\Gamma)$  and lies in the weighted completed coefficient category. The bracket-tame estimate (W<sub>1</sub>), together with the finite-window coadjoint condition (W<sub>2</sub>), bounds this single graph by a finite constant times the corresponding spacetime graph weight.

*Proof.* The interaction  $I$  has bounded polynomial degree  $D_o$  at each vertex in the coefficient direction by hypothesis. A connected Costello graph  $\Gamma$  with  $V$  vertices, each of polynomial degree  $\leq D_o$ , and  $E$  internal edges, contributes a coefficient pairing of external length at most  $V \cdot D_o - 2E$  in the source factor and a matching length in the target. By the bracket-tame estimate (W<sub>1</sub>), each internal edge contraction  $C_{\mathfrak{h},w}: \mathfrak{h}_w \widehat{\otimes} \mathfrak{h}_{\text{cont}}^{\vee,w} \rightarrow \mathbb{C}$  changes the total weight by at most the finite constant attached to the degrees present in that graph. Coadjoint legs are controlled windowwise by (W<sub>2</sub>). Hence the graph contribution is a well-defined element of the weighted completion. The assertion is graphwise; the full effective interaction is Costello's completed sum over graph degrees.  $\square$

**THEOREM 5.69** (*Weighted RG flow, graphwise compatibility*). Let  $w$  be a degree weight as in Definition 5.51 and let  $I \in \mathcal{O}_{\text{loc}}(\mathcal{E}_w)$  be a local interaction satisfying a vertexwise polynomial-degree bound  $D_o$ . Each fixed graph contribution to the Costello RG flow

$$I_L = W(P_{\epsilon,L}^w, I)$$

lies in the weighted Tate completion and is continuous in  $(\epsilon, L)$ . The completed sum over counterterms and the QME obstruction class require the locality input of Problem 5.72.

*Proof.* By Proposition 5.68, every Feynman diagram considered individually remains in the weighted Tate category. Its finite coefficient-degree bound is  $D_o V(\Gamma) - 2E(\Gamma)$ , and continuity in  $(\epsilon, L)$  follows from Proposition 5.63 (2). Passing from graphwise well-definedness to a renormalized QME solution is precisely the residual analytic problem.  $\square$

*Cotangent-lift inside  $B_w$* 

**COROLLARY 5.70** (*Cotangent-lift inside the weighted coefficient category*). Take  $w(d) = q^d$  with  $q > 1$  as in Remark 5.53. Then

- (a) the Casimir  $C_{\mathfrak{h},w}$  converges in  $\mathfrak{h}_w \widehat{\otimes} \mathfrak{h}_{\text{cont}}^{\vee,w}$ ;
- (b) the Costello heat-kernel propagator  $P_{\epsilon,L}^w$  extends to the weighted field complex  $\mathcal{E}_w$  as a continuous BV family;
- (c) each fixed Costello graph contribution from a vertexwise polynomial-degree-bounded interaction stays inside  $B_w$ ;
- (d) the central-operation cochain  $\Phi$  of Theorem 4.20 extends to the weighted cotangent extension  $\mathfrak{g}_w = \mathfrak{h}_w \ltimes \mathfrak{h}_{\text{cont}}^{\vee,w}[1]$  as a continuous cochain map

$$\Phi^w: C_{\text{CE,cont}}^\bullet(\mathfrak{g}_w) \longrightarrow B_w \subset \text{PV}(\widehat{\mathfrak{S}}_{\text{Tate}}(\mathfrak{h}_w)).$$

Conditional on Problem 5.72, the all-order lift  $\Phi_{h,\text{cond}}^w$  is Proposition 5.71.

*Proof.* This is Theorem 5.57 parts (b)–(e), evaluated at the exponential weight  $w(d) = q^d$ ,  $q > 1$ . The  $P_0$ -algebra reading of  $\Phi^w$  uses the bracket-admissible envelope of Definition 4.18. The strict continuous duality statement of Lemma 5.165 applies to the product/direct-sum Tate pair. The needed dual generators are inserted by the symmetric Mittag-Leffler coefficient pairing of Remark 5.55.  $\square$

**PROPOSITION 5.71** (*Weighted QME lift under obstruction vanishing*). Assume the Costello-locality/QME compatibility identity of Problem 5.72: the finite-window counterterm system for the mixed brane-defect interaction has vanishing obstruction class and is compatible with the weighted coefficient category. Under this assumption, the formal quantum CE/PV central-operation map  $\Phi_h^{(\circ)}$  governed by Theorem 5.216 has a weighted-continuous cotangent extension

$$\Phi_{h,\text{cond}}^w : C_{\text{CE},\text{cont}}^\bullet(\mathfrak{h}_{w,h} \times \mathfrak{h}_{h,\text{cont}}^{\vee,w}[1][[\hbar]]) \xrightarrow[\text{QME obstruction}=0]{\sim} Z_{\text{fact}}^{P_{\circ,h}}(\text{Obs}_{\text{open}}^h)|_{\mathfrak{g}_w}.$$

The displayed isomorphism is a theorem in the weighted-continuous sector with vanishing QME obstruction. The original unweighted Tate pair is governed by the endpoint obstruction of Theorem 5.93.

*Proof.* The algebraic cochain identity is supplied by Theorem 5.70. The BV graph realization decomposes the cotangent action  $\text{ad}^\vee$  into Costello graphs in  $\mathcal{E}_w$ , and Theorem 5.69 keeps each fixed graph inside the weighted category. The nonformal point is the cancellation between Costello-locality counterterms and the bulk-boundary kernel at the brane defect. The proposition assumes exactly that weighted QME compatibility and the vanishing of the obstruction class named in Problem 5.72. With that assumption in force, the formal  $\hbar$ -adic construction of Theorem 5.216 applies in the weighted coefficient envelope. The graphwise finite-scale assertions of Theorem 5.69 are anterior to this obstruction and to the principal-part current extension; that extension is the additional datum for compatibility with  $\mathcal{D}'_c(I) \widehat{\otimes} \mathcal{P}[1]$ .  $\square$

**Problem 5.72** (*Costello-locality compatibility for the weighted RG flow*). *Construction problem.* Construct a compatible counterterm system  $\{C_{n,M}\}_{n \geq 1, M \geq 1}$  for the mixed brane-defect interaction at the defect  $L = \mathbb{R}_{\text{brane}} \times \{o\} \times \{p\}$  in the weighted field complex  $\mathcal{E}_w$  such that:

- (L1) each  $C_{n,M}$  is a local functional in  $\mathcal{O}_{\text{loc}}^{h^n}(\mathcal{E}_w^{\leq M})$  of bounded vertex polynomial degree;
- (L2) the truncations are compatible:  $\pi_{M,N} C_{n,M} = C_{n,N}$  for  $M \geq N$ ;
- (L3) the recursion solves the QME order by order.

*Named obstruction class.* The class

$$[\text{Ob}_{w,\partial}^{\text{loc}}] \in H^1(\mathcal{O}_{\text{loc}}(\mathcal{E}_w), Q + \{I_o, -\})$$

obstructing (L1)–(L3); equivalently, by Theorem 5.83, the pair  $(([\mathfrak{o}_{n,M}^{\text{ns}}])_M, \lambda_n)$  in  $\varprojlim_M H^1(Q_{w,M}^\bullet(I)) \oplus \varprojlim_M^1 H^0(Q_{w,M}^\bullet(I))$ . *Available inputs.* Estimate (W1) bounds vertex weights; (W2) bounds coadjoint legs graph by graph; Theorem 8.17 controls wave-front and finite-window locality estimates. The construction problem is the exact further datum that turns Hadamard control into anomaly cancellation.

**COROLLARY 5.73** (*Descendant cochain comparison under weighted QME vanishing*). Assume the obstruction class in Problem 5.72 vanishes, so that Proposition 5.71 applies. Then the Tate-coefficient

lift of the descendant cochain comparison of Problem 5.221 exists in the weighted Tate completion only under this hypothesis: the deformation

$$\Phi_{\hbar, \text{cond}}^{(0,1)} : C_{\text{CE,cont}}^\bullet(\mathfrak{h}_{w,\hbar} \ltimes \mathfrak{h}_{\hbar, \text{cont}}^{\vee,w}[\mathbf{I}])^{(0)+(1)}[[\hbar]] \xrightarrow[\text{QME obstruction}=0]{\sim} Z_{\text{fact}}^{P_0, \hbar}(\text{Obs}_{\text{open}}^{\hbar})^{(0)+(1)}|_{\mathfrak{g}_w}$$

is an isomorphism on the degree-zero plus one-antifield sector in the weighted theory with vanishing QME obstruction.

*Remark 5.74 (Sector restriction).* The weighted mechanism restricts the cotangent direction from the ambient polynomial Hamiltonian dual to the weighted dual  $\mathfrak{h}_{\text{cont}}^{\vee,w} = \prod_d w(d)^{-1} H_d^\vee$ . The strict polynomial Hamiltonian dual  $\mathfrak{h}_{\text{cont}}^\vee = \bigoplus_d H_d^\vee$  is a *dense subspace* of  $\mathfrak{h}_{\text{cont}}^{\vee,w}$  for every  $w$  with  $w(d) \rightarrow \infty$ , since the strict direct sum is finite-degree truncations and the weighted product is the weighted Mittag-Leffler inverse-limit completion. Hence the weighted lift extends the cochain map continuously from the strict polynomial sector to the weighted product completion. This recovers the weighted-continuous Hamiltonian identities. Unweighted regulator-independence and identification of the weighted product with the original strict continuous dual are separate statements.

*Regulator independence and its boundary*

*Definition 5.75 (Regulator-admissible finite-window sector).* A local functional, interaction, obstruction cocycle, or graph coefficient in the weighted theory is *regulator-admissible* if it is represented by a strict Mittag-Leffler family on the finite Tate windows

$$\{\mathcal{E}_w^{w_0}\}_{w_0 \geq 1}$$

and satisfies the following four conditions.

- (A1) The transition maps in the coefficient and local-functional directions are the truncations to degrees  $d \leq w_0$ , or the induced restriction obtained by setting higher-degree coefficient fields to zero. They are degreewise surjective maps of complexes.
- (A2) At each fixed  $\hbar$ -order and for each fixed Costello graph, the coefficient expression uses only vertex-wise polynomial-degree-bounded Hamiltonian and cotangent legs.
- (A3) If  $w, w'$  are two degree weights satisfying Definition 5.51, the diagonal finite-window rescaling

$$R_{w,w'}^{w_0}(w(d)e_{d,i}) = w'(d)e_{d,i}, \quad R_{w,w'}^{w_0}(w(d)^{-1}e_d^i) = w'(d)^{-1}e_d^i$$

transports the interaction, differential, bracket, propagator contraction, and counterterm representative from the  $w$ -window to the  $w'$ -window. The bracket is the transported bracket; comparison of two weights is made after this transport, while raw unweighted structure constants are held outside the comparison.

- (A4) The cohomology being computed is the admissible filtered cohomology detected on finite quotients and associated graded pieces in the envelope of Statement 5.17; observables are restricted to data visible in finite quotients or associated graded pieces.

*Definition 5.76 (Exponential finite-window regulator tower).* For  $q > 1$  put  $w_q(d) = q^d$ . For a finite Hamiltonian window  $M$  set

$$\mathfrak{h}_{q, \leq M} = \prod_{1 \leq d \leq M} q^d H_d, \quad D_{q, \leq M} = \prod_{1 \leq d \leq M} q^{-d} H_d^\vee,$$

and

$$\mathcal{E}_q^M = \Omega^\bullet(\mathbb{R}^2) \widehat{\otimes} \Omega^{\circ, \bullet}(\mathbb{C}^2) \widehat{\otimes} (\mathfrak{h}_{q, \leq M}[1] \oplus D_{q, \leq M}[2]).$$

For  $M \geq N$ , let

$$\pi_{M, N}^q: (\mathfrak{h}_{q, \leq M}, D_{q, \leq M}) \twoheadrightarrow (\mathfrak{h}_{q, \leq N}, D_{q, \leq N})$$

be degree truncation. The weight transport and the unweighting map on the window  $M$  are

$$R_{q, q'}^M(q^d e_{d, i}) = (q')^d e_{d, i}, \quad R_{q, q'}^M(q^{-d} e_d^i) = (q')^{-d} e_d^i,$$

and

$$U_q^M(q^d e_{d, i}) = e_{d, i}, \quad U_q^M(q^{-d} e_d^i) = e_d^i.$$

The finite-window unweighted endpoint is the finite-dimensional pair

$$(\mathfrak{h}_{\leq M}, D_{\leq M}) = \left( \prod_{d \leq M} H_d, \prod_{d \leq M} H_d^\vee \right).$$

Its transition map for  $M \geq N$  is denoted  $\pi_{M, N}^\circ$ .

LEMMA 5.77 (*Finite-window transport squares*). For  $q, q' > 1$  and  $M \geq N$ , the maps of Definition 5.76 satisfy

$$\pi_{M, N}^{q'} R_{q, q'}^M = R_{q, q'}^N \pi_{M, N}^q, \quad \pi_{M, N}^\circ U_q^M = U_q^N \pi_{M, N}^q,$$

and

$$U_{q'}^M R_{q, q'}^M = U_q^M, \quad R_{q', q''}^M R_{q, q'}^M = R_{q, q''}^M.$$

The truncated Casimir

$$C_q^M = \sum_{d \leq M} \sum_i (q^d e_{d, i}) \otimes (q^{-d} e_d^i)$$

is transported to  $C_{q'}^M$  by  $R_{q, q'}^M$ , and to the ordinary finite Casimir  $C_\circ^M$  by  $U_q^M$ . Hence every Costello graph whose coefficient legs lie in  $d \leq M$  has the same unweighted coefficient after applying  $U_q^M$ , independently of  $q$ .

*Proof.* All four identities are degree-wise identities on the two bases  $q^d e_{d, i}, q^{-d} e_d^i$ . Truncation forgets degrees  $d > N$ , while  $R_{q, q'}^M$  and  $U_q^M$  multiply only the surviving degree- $d$  basis vector by a scalar. Thus truncation and transport commute.

The Casimir statement follows term by term:

$$R_{q, q'}^M((q^d e_{d, i}) \otimes (q^{-d} e_d^i)) = (q')^d e_{d, i} \otimes (q')^{-d} e_d^i,$$

and  $U_q^M$  sends the same term to  $e_{d, i} \otimes e_d^i$ . A fixed graph has finitely many coefficient contractions and external coefficient legs. If all of them lie in the window  $M$ , applying  $U_q^M$  to each leg removes every diagonal weight and each internal Casimir becomes  $C_\circ^M$ .  $\square$

Statement 5.78 (*Weighted RG transport criterion*). A weighted mixed brane-defect interaction enters the regulator theorem only with the following data.

- (G1) Its coefficient and local-functional restrictions to  $d \leq w_\circ$  form strict Mittag-Leffler towers with degree-wise surjective transition maps.

- (G2) At every fixed  $\hbar$ -order and for every fixed Costello graph, the coefficient expression is vertexwise bounded in polynomial Hamiltonian and cotangent degree.
- (G3) For any admissible degree weights  $w, w'$ , the diagonal finite-window maps  $R_{w,w'}^{w_o}$  transport the differential, bracket, propagator contraction, interaction, and counterterm representatives.
- (G4) The cohomology theory is the admissible filtered cohomology of Statement 5.17.
- (G5) The Costello RG recursion  $\mathcal{W}(P_{\epsilon,L}, I)$  preserves the same strict Mittag-Leffler tower: every RG step has source and target in the weighted coefficient envelope.

*Proof.* Conditions (G1)–(G4) are the four clauses of Definition 5.75. Condition (G5) is the coefficient part of Costello requirement (C3). Proposition 5.68 proves (G2) for a fixed graph. Theorem 8.17 proves that the finite-window Hadamard estimates pass through the Mittag-Leffler limit when the windowwise counterterms are compatible. The all-order datum beyond this graphwise statement is the obstruction class of Problem 5.72.  $\square$

**THEOREM 5.79** (*Transport groupoid of admissible weights*). On the class of interactions satisfying 5.78, admissible degree weights form a connected transport groupoid. Its objects are the weighted finite-window local-functional complexes

$$C_w^{w_o} = (\mathcal{O}_{\text{loc}}(\mathcal{E}_w^{w_o}), Q + \{I_w^w, -\}),$$

and its arrows are the diagonal maps  $R_{w,w'}^{w_o}$ . For every  $w, w', w''$  and every finite window  $w_o$ ,

$$R_{w',w''}^{w_o} R_{w,w'}^{w_o} = R_{w,w''}^{w_o}, \quad R_{w,w}^{w_o} = \text{id},$$

and these arrows commute with truncation in  $w_o$ , Costello graph contractions, and RG counterterm representatives.

*Proof.* The formula defining  $R_{w,w'}^{w_o}$  is diagonal:

$$w(d)e_{d,i} \mapsto w'(d)e_{d,i}, \quad w(d)^{-1}e_d^i \mapsto w'(d)^{-1}e_d^i.$$

Composition multiplies the scalars degree by degree, hence gives the displayed groupoid law. Since the maps are diagonal in  $d$ , they commute with truncation to  $d \leq w_o$ . By 5.78(G3), the transported bracket, interaction, propagator, and counterterm representatives are the corresponding  $w'$ -objects. Therefore  $R_{w,w'}^{w_o}$  is an isomorphism of the finite-window RG complexes and intertwines every fixed graph contraction.  $\square$

**THEOREM 5.80** (*Canonical equivalence of finite-degree local observables*). Let  $w(d) = q^d$  and  $w'(d) = (q')^d$ , with  $q, q' > 1$ . Let  $\mathcal{O}^w$  be a regulator-admissible local observable, graph coefficient, or obstruction cocycle represented on a finite window  $d \leq M$ . Then  $R_{q,q'}^M$  is the canonical identification of the  $q$ -weighted and  $q'$ -weighted representatives on that finite window:

$$\mathcal{O}_M^{q'} = R_{q,q'}^M \mathcal{O}_M^q.$$

After writing both representatives in the common unweighted basis,

$$U_q^M(\mathcal{O}_M^q) = U_{q'}^M(\mathcal{O}_M^{q'}).$$

If  $O_M^q$  is a cocycle for  $Q + \{I_M^q, -\}$ , then the common unweighted representative is a cocycle for the unweighted finite-window transported differential. If  $\text{Ob}_{q,M}$  is the finite-window obstruction cocycle produced by the Costello recursion, then

$$U_q^M(\text{Ob}_{q,M}) = U_{q'}^M(\text{Ob}_{q',M}), \quad \pi_{M,N}^\circ U_q^M(\text{Ob}_{q,M}) = U_q^N(\text{Ob}_{q,N})$$

for  $M \geq N$ . Thus a finite-degree local observable, fixed graph coefficient, or obstruction class is a single object of the finite-window transported complex, presented in different weighted bases by the choices  $q > 1$ .

*Proof.* The observable lies in the finite complex  $C_q^M$ . By Theorem 5.79,  $R_{q,q'}^M$  is an isomorphism from  $C_q^M$  to  $C_{q'}^M$ , carrying the differential, bracket, propagator contraction, interaction, counterterm representative, and obstruction cocycle. Lemma 5.77 gives  $U_{q'}^M R_{q,q'}^M = U_q^M$ . Therefore both weighted representatives have the same coefficient array after unweighting.

Since  $R_{q,q'}^M$  is a cochain isomorphism, cocycles go to cocycles and coboundaries go to coboundaries. Applying  $U_q^M$  merely writes the same finite complex in the common unweighted basis; the transported differential is  $U_q^M(Q + \{I_M^q, -\})(U_q^M)^{-1}$ . Hence the cocycle condition is preserved. The obstruction equalities are the same statement for the obstruction cochain produced at the relevant RG step. The transition identity follows from  $\pi_{M,N}^\circ U_q^M = U_q^N \pi_{M,N}^q$  and strict Mittag-Leffler compatibility of the finite-window obstruction system.  $\square$

**THEOREM 5.81** (*Regulator independence inside the admissible finite-window sector*). Let  $w, w'$  be degree weights satisfying Definition 5.51, and let  $I_o^w$  be a regulator-admissible mixed brane-defect interaction satisfying 5.78. Put  $I_o^{w'} = R_{w,w'} I_o^w$ . Then the diagonal rescalings  $R_{w,w'}^{w_o}$  assemble to an isomorphism of strict Mittag-Leffler local-functional complexes

$$R_{w,w'} : \varprojlim_{w_o} (\mathcal{O}_{\text{loc}}(\mathcal{E}_w^{w_o}), Q + \{I_o^w, -\}) \xrightarrow{\cong} \varprojlim_{w_o} (\mathcal{O}_{\text{loc}}(\mathcal{E}_{w'}^{w_o}), Q + \{I_o^{w'}, -\}).$$

Under this isomorphism:

- (i) truncated Casimirs, heat-kernel propagators, and fixed graphwise Costello contractions are identified;
- (ii) the obstruction class

$$\text{Ob}_w \in H_{\text{adm}}^1(\mathcal{O}_{\text{loc}}(\mathcal{E}_w), Q + \{I_o^w, -\})$$

maps to  $\text{Ob}_{w'}$ ;

- (iii) vanishing or nonvanishing of the mixed brane-defect QME obstruction is independent of the admissible weight.

These isomorphisms are the arrows of the transport groupoid of Theorem 5.79; they are functorial in triples of admissible weights.

*Proof.* On the finite window  $d \leq w_o$ , the map  $R_{w,w'}^{w_o}$  is an invertible linear map on the coefficient space and the identity on the spacetime factor. The bracket and the interaction in the  $w'$ -window are, by admissibility, the transported bracket and transported interaction. The map sends

$$\sum_{d \leq w_o} \sum_i (w(d) e_{d,i}) \otimes (w(d)^{-1} e_d^i)$$

to the corresponding  $w'$ -weighted truncated Casimir. Hence it sends  $P_{\epsilon,L}^{w,w_o}$  to  $P_{\epsilon,L}^{w',w_o}$ , commutes with the differential  $Q$ , and carries the transported BV bracket and every finite-window graph contraction to the  $w'$ -window contraction. Condition (A<sub>3</sub>) gives the same statement for  $I_o$ , the counterterm representatives, and the obstruction cocycle.

The maps  $R_{w,w'}^{w_o}$  commute with truncation in  $w_o$ . The finite-window complexes therefore form an isomorphic pair of strict Mittag-Leffler inverse systems. By (A<sub>1</sub>), these are degreewise surjective towers of complexes. Hence the inverse limit is exact in the admissible envelope of Statement 5.17; equivalently  $\varprojlim^1 = 0$  for this tower. Passing to the inverse limit gives the displayed isomorphism of local-functional complexes and identifies their admissible filtered cohomology classes. In particular the obstruction class vanishes for  $w$  if and only if the transported obstruction class vanishes for  $w'$ .

Functoriality in triples is the groupoid identity proved in Theorem 5.79.  $\square$

*Definition 5.82 (Finite-window non-scalar QME tower).* Fix an interval  $I$ , a degree weight  $w$ , and a regulator-admissible interaction satisfying Statement 5.78. For a finite Hamiltonian window  $M$ , put

$$D_{w,\leq M} := \prod_{d \leq M} w(d)^{-1} H_d^\vee,$$

and define the truncated principal-part target

$$A_{\partial,h}^{\text{pp},w,M}(I) = \widehat{\mathbf{S}}_{\mathbb{C}[[\hbar]]}(\Omega_c^\bullet(I) \otimes \prod_{d \leq M} w(d) H_d[[\hbar]]) \widehat{\otimes}_{\widehat{\mathbf{S}}_{\mathbb{C}[[\hbar]]}}((\mathcal{D}'_c(I) \widehat{\otimes} D_{w,\leq M}[[\hbar]])[1]).$$

Assume that the scalar-contact projection of Problem 5.219 has been chosen windowwise, commutes with truncation, and is transported by  $R_{w,w'}^M$ . The finite-window non-scalar obstruction complex is

$$\mathcal{Q}_{w,M}^\bullet(I) = \ker(\widehat{\tau}_{\text{sc},M}) \subset \left( \mathcal{O}_{\text{loc}}^{\text{bd}}(\mathcal{E}_w^M|_I; A_{\partial,h}^{\text{pp},w,M}(I)), d_M \right),$$

where  $d_M = Q + \{I_{o,M}^w, -\}_{\hbar,M}$ . The bracket, interaction, and differential are the transported finite-window structures supplied by the weighted RG criterion; naive projected Hamiltonian brackets are replaced by this transported structure. The inverse system  $\{\mathcal{Q}_{w,M}^\bullet(I)\}_{M \geq 1}$  uses degree truncation  $\pi_{M,N}$  for  $M \geq N$ , together with the  $\hbar$ -adic filtration, the scalar-contact filtration, and the finite-window coefficient filtration.

**THEOREM 5.83 (Finite-window counterterm criterion for the non-scalar QME).** Work in the tower of Definition 5.82. Suppose that a counterterm system has been chosen through order  $\hbar^{n-1}$ , and let

$$\mathfrak{o}_{n,M}^{\text{ns}} \in Z^1(\mathcal{Q}_{w,M}^\bullet(I))$$

be the order- $n$  non-scalar residual after those lower-order choices. Equivalently, in the curved Maurer–Cartan form,

$$\mathfrak{o}_{n,M}^{\text{ns}} = [\hbar^n] \left( \text{Ob}_{w,\partial,\hbar,M}^{\text{red}} + d_M C_{<n,M} + \frac{1}{2} \{C_{<n,M}, C_{<n,M}\}_{\hbar,M} \right),$$

with the quadratic term omitted for the linearized equation  $\text{Ob}^{\text{red}} + d_M C = 0$ . Compatible order- $n$  finite-window counterterms

$$C_{n,M} \in \mathcal{Q}_{w,M}^0(I), \quad d_M C_{n,M} = -\mathfrak{o}_{n,M}^{\text{ns}}, \quad \pi_{M,N} C_{n,M} = C_{n,N},$$

exist if and only if the following two obstructions vanish:

$$([\mathfrak{o}_{n,M}^{\text{ns}}])_M \in \varprojlim_M H^1(Q_{w,M}^\bullet(I)),$$

and, after choosing any windowwise primitives  $c_{n,M}$  with  $d_M c_{n,M} = -\mathfrak{o}_{n,M}^{\text{ns}}$ , the Milnor primitive-compatibility class

$$\lambda_n \in \varprojlim_M^1 H^0(Q_{w,M}^\bullet(I))$$

represented, after passing to cohomology, by

$$[\pi_{M,N} c_{n,M} - c_{n,N}] \in H^0(Q_{w,N}^\bullet(I))$$

is zero. If the inverse system  $\{H^0(Q_{w,M}^\bullet(I))\}_M$  is Mittag–Leffler, then  $\lambda_n = 0$  automatically, so windowwise vanishing of the  $H^1$ -classes is sufficient.

Iterating this criterion for all  $n \geq 1$  constructs an  $\hbar$ -adic compatible counterterm  $C_{\hbar,w} = \sum_{n \geq 1} \hbar^n C_{n,w}$  in the admissible finite-window inverse limit. The transport maps  $R_{w,w'}^M$  identify the obstruction pair

$$\mathfrak{D}_n^{\text{ns}} = (([\mathfrak{o}_{n,M}^{\text{ns}}])_M, \lambda_n)$$

for all admissible weights. Therefore a nonzero  $\mathfrak{D}_n^{\text{ns}}$ , at the first order where it occurs, is the precise finite-window obstruction to killing the non-scalar QME class by compatible weighted counterterms.

*Proof.* The lower-order counterterms solve the QME modulo  $\hbar^n$ . Applying  $d_M$  to the displayed order- $n$  residual and using  $d_M^2 = 0$ , the graded Jacobi identity, and the lower-order equations gives  $d_M \mathfrak{o}_{n,M}^{\text{ns}} = 0$ . The scalar-contact projection is a chain map by Definition 5.82, and the balanced scalar branch has zero scalar symbol, so the residual lies in  $Q_{w,M}^\bullet(I)$ .

Necessity is immediate. A compatible counterterm  $C_{n,M}$  is a primitive of  $-\mathfrak{o}_{n,M}^{\text{ns}}$ , hence each  $H^1$ -class vanishes. Its truncation identities make  $\pi_{M,N} C_{n,M} - C_{n,N} = 0$ , so the associated Milnor  $\varprojlim_M^1 H^0$ -class is also zero.

Conversely, assume the two displayed obstructions vanish. The first condition gives primitives  $c_{n,M}$  window by window. For  $M \geq N$ , the difference  $\pi_{M,N} c_{n,M} - c_{n,N}$  is  $d_N$ -closed because the residuals commute with truncation. The cohomology classes of these closed degree-zero differences form the standard Roos 1-cocycle computing  $\varprojlim_M^1 H^0(Q_{w,M}^\bullet(I))$ . Vanishing of  $\lambda_n$ , together with the degree-wise Mittag–Leffler property of the finite-window truncation tower, means that the primitives can be adjusted by degree-zero cocycles and degree-zero boundaries so that the adjusted primitives commute with every truncation map. Those adjusted primitives are the desired  $C_{n,M}$ . Passing to the inverse limit gives  $C_{n,w}$ .

The assertion about the  $H^0$ -Mittag–Leffler hypothesis is the standard vanishing of  $\varprojlim_M^1$  for a Mittag–Leffler inverse system. Weight invariance follows from Theorem 5.81: the maps  $R_{w,w'}^M$  are cochain isomorphisms, commute with truncation, and carry residuals, primitives, and their difference cocycles to the corresponding data for  $w'$ .  $\square$

*Definition 5.84 (Regular-density closed-exchange finite-window complex).* Work in the tower of Definition 5.82, and assume the regular-density Costello graph kernel of Proposition B.17 on every finite Hamiltonian window. Thus the cotangent current labels are restricted to

$$\mathcal{D}_c^{\text{reg}}(I) = \{ b(t)dt \mid b \in C_c^\infty(I) \},$$



and the finite-window graph weights

$$W_{\Gamma,L}^{R,\text{reg},M}(a_\bullet, b_\bullet; f_\bullet, \rho_\bullet)$$

are renormalized brane-defect local functionals compatible with interval extension, degree truncation, and admissible weight transport. Let  $\mathcal{G}_{w,M}^{\bullet,\text{reg}}(I)$  be the complete filtered  $\mathbb{C}[[\hbar]]$ -span, inside  $\mathcal{Q}_{w,M}^\bullet(I)$ , of the scalar-zero members among these graph weights and their regular-density local counterterms. The zero-internal-edge generators are the coefficient-current terms

$$\iota_I(B_f^w(a)), \quad \iota_I(\Theta_\rho^w(b(t)dt)),$$

whenever they have zero scalar-contact projection.

Define

$$\mathcal{X}_{w,M}^{\bullet,\text{reg}}(I) \subset \mathcal{Q}_{w,M}^\bullet(I)$$

to be the smallest complete filtered  $d_M$ -stable subspace containing  $\mathcal{G}_{w,M}^{\bullet,\text{reg}}(I)$ . Its differential is

$$d_M^{X,\text{reg}} = d_M|_{\mathcal{X}_{w,M}^{\bullet,\text{reg}}(I)}.$$

This is a finite-window graph complex. The naive finite-dimensional Hamiltonian Lie quotient is the obstruction of Lemma 5.209. The window is a coefficient-support and graph-locality condition on already constructed regular-density local functionals.

**PROPOSITION 5.85** (*Regular-density construction of  $X$  and  $\Xi$* ). The complexes of Definition 5.84 form an inverse system

$$\rho_{M,N}^{X,\text{reg}}: \mathcal{X}_{w,M}^{\bullet,\text{reg}}(I) \longrightarrow \mathcal{X}_{w,N}^{\bullet,\text{reg}}(I), \quad M \geq N,$$

by restricting the finite-window truncation maps  $\pi_{M,N}$  of  $\mathcal{Q}_{w,M}^\bullet(I)$ . The closed-exchange insertion map is the degree-zero inclusion

$$\Xi_M^{\text{reg}}: \mathcal{X}_{w,M}^{\bullet,\text{reg}}(I) \hookrightarrow \mathcal{Q}_{w,M}^\bullet(I).$$

It is a cochain map and satisfies

$$\pi_{M,N} \Xi_M^{\text{reg}} = \Xi_N^{\text{reg}} \rho_{M,N}^{X,\text{reg}}.$$

For admissible weights  $w, w'$ , the diagonal transports restrict to cochain isomorphisms

$$R_{w,w'}^{M,X}: \mathcal{X}_{w,M}^{\bullet,\text{reg}}(I) \xrightarrow{\cong} \mathcal{X}_{w',M}^{\bullet,\text{reg}}(I)$$

and intertwine the inclusions:

$$R_{w,w'}^M \Xi_M^{\text{reg}} = \Xi_{M,w'}^{\text{reg}} R_{w,w'}^{M,X}$$

Consequently the abstract closed-exchange criterion of Theorem 5.87 applies to the regular-density tower  $\mathcal{X}_{w,M}^{\bullet,\text{reg}}(I)$  with  $\Xi_M = \Xi_M^{\text{reg}}$ .

*Proof.* Proposition B.17 gives the finite-window identities

$$\pi_{M,N} W_{\Gamma,L}^{R,\text{reg},M} = W_{\Gamma,L}^{R,\text{reg},N} \pi_{M,N}^{\text{CE}}, \quad R_{w,w'}^M W_{\Gamma,L}^{R,\text{reg},M} = W_{\Gamma,L,w'}^{R,\text{reg},M} R_{w,w'}^{M,\text{CE}}$$

and the same compatibility for the regular-density counterterms. Since  $\widehat{\sigma}_{\text{sc},M}$  is a chain map in Definition 5.82, its kernel  $\mathcal{Q}_{w,M}^\bullet(I)$  is a subcomplex. Therefore scalar-zero graph weights and their  $d_M$ -closures stay in  $\mathcal{Q}_{w,M}^\bullet(I)$ .

Truncation  $\pi_{M,N}$  and weight transport  $R_{w,w'}^M$  are cochain maps on the non-scalar tower. They carry the generating regular-density graph weights and counterterms to the corresponding generators in the lower window or transported weight. By minimality of the complete  $d_M$ -stable subcomplex generated by these elements, they restrict to the displayed maps on  $\mathcal{X}^{\bullet,\text{reg}}$ .

The map  $\Xi_M^{\text{reg}}$  is inclusion of a subcomplex of  $\mathcal{Q}_{w,M}^\bullet(I)$ , hence  $d_M \Xi_M^{\text{reg}} = \Xi_M^{\text{reg}} d_M^{X,\text{reg}}$ . The compatibility with  $\rho_{M,N}^{X,\text{reg}}$  and  $R_{w,w'}^{M,X}$  is precisely the restriction of the same identities on  $\mathcal{Q}^\bullet$ .  $\square$

**PROPOSITION 5.86** (*Raw kernel obstruction to a stronger cochain map*). Let  $\kappa_{h,w,I}^{\text{graph,reg},M}$  be the finite-window regular-density graph kernel of Proposition B.17, viewed as a degree-zero element of the convolution Hom from the regular-density CE source to the target  $\mathfrak{Q}_{w,\partial,h}^\bullet(I)$ . If one tries to take the regular-density CE source itself as the closed-exchange complex and  $\Xi_M = \kappa_{h,w,I}^{\text{graph,reg},M}$ , then the exact obstruction to an ordinary cochain map into  $\mathcal{Q}_{w,M}^\bullet(I)$  is the pair

$$\widehat{\sigma}_{\text{sc},M} \kappa_{h,w,I}^{\text{graph,reg},M}, \quad d_{\mathbf{K}} \kappa_{h,w,I}^{\text{graph,reg},M} = d_M \kappa_{h,w,I}^{\text{graph,reg},M} - \kappa_{h,w,I}^{\text{graph,reg},M} d_{\text{CE}}$$

The first term is the obstruction to landing in  $\ker \widehat{\sigma}_{\text{sc},M}$ ; the second is the obstruction to commuting with differentials. If the desired closed-exchange datum is required to be a  $P_{o,h}$ -central-operation kernel, the ordinary cochain obstruction is strengthened to the Maurer–Cartan curvature

$$\text{Curv}_{\mathbf{K}}(\kappa_{h,w,I}^{\text{graph,reg},M}) = d_{\mathbf{K}} \kappa_{h,w,I}^{\text{graph,reg},M} + \frac{1}{2} [\kappa_{h,w,I}^{\text{graph,reg},M}, \kappa_{h,w,I}^{\text{graph,reg},M}]_{\mathbf{K}}.$$

Thus the regular-density graph layer supplies a proved closed-exchange tower by the subcomplex construction above. The stronger assertion that the raw bulk-to-defect CE kernel itself is the cochain map is equivalent to vanishing of the displayed scalar and convolution defects, with finite-window and weight-transport compatibility.

*Proof.* The convolution differential is  $d_{\mathbf{K}} T = d_M T - (-1)^{|T|} T d_{\text{CE}}$ . Since  $\kappa_{h,w,I}^{\text{graph,reg},M}$  has degree zero, the displayed formula is exactly the commutator of the proposed map with the two differentials. Hence it vanishes if and only if the raw kernel is a cochain map into the ambient target. It lands in the non-scalar complex precisely when its scalar-contact projection vanishes.

The bracket term is the completed convolution bracket of Definition B.15. A  $P_{o,h}$ -central-operation kernel is a Maurer–Cartan element in that convolution dg Lie algebra, so its exact obstruction is  $d_{\mathbf{K}} \kappa + \frac{1}{2} [\kappa, \kappa]_{\mathbf{K}}$ . The generator-level form of this statement is the obstruction list in Proposition B.21. The compatibility clauses follow from the same truncation and weight-transport identities used in Proposition 5.85.  $\square$

**THEOREM 5.87** (*Closed-exchange plus counterterm criterion*). Work in the non-scalar tower of Definition 5.82. Let

$$\rho_{M,N}^X : \mathcal{X}_{w,M}^\bullet(I) \longrightarrow \mathcal{X}_{w,N}^\bullet(I), \quad M \geq N,$$

be an inverse system of closed-exchange local-functional complexes, and let

$$\Xi_M : \mathcal{X}_{w,M}^\bullet(I) \longrightarrow \mathcal{Q}_{w,M}^\bullet(I)$$

be degree-zero cochain maps satisfying

$$\pi_{M,N} \Xi_M = \Xi_N \rho_{M,N}^X.$$

Assume also that the closed-exchange tower and the maps  $\Xi_M$  are transported by  $R_{w,w'}^M$ , in the same sense as the non-scalar tower. Let

$$\mathfrak{o}_{n,M}^{\text{ns}} \in Z^1(\mathcal{Q}_{w,M}^\bullet(I)), \quad \pi_{M,N} \mathfrak{o}_{n,M}^{\text{ns}} = \mathfrak{o}_{n,N}^{\text{ns}},$$

be the order- $n$  residual after the lower-order choices.

For a fixed window  $M$ , the branch equation

$$\mathfrak{o}_{n,M}^{\text{ns}} + \Xi_M(W_{n,M}) + d_M C_{n,M} = \mathfrak{o}, \quad W_{n,M} \in Z^1(\mathcal{X}_{w,M}^\bullet(I)), \quad C_{n,M} \in \mathcal{Q}_{w,M}^\circ(I),$$

has a solution if and only if

$$[\mathfrak{o}_{n,M}^{\text{ns}}] \in -\text{im}\left(H^1(\Xi_M): H^1(\mathcal{X}_{w,M}^\bullet(I)) \rightarrow H^1(\mathcal{Q}_{w,M}^\bullet(I))\right).$$

In the inverse limit, put

$$\beta_n^{\text{cl}} = \left[ ([\mathfrak{o}_{n,M}^{\text{ns}}])_M \right] \in \text{coker}\left( \varprojlim_M H^1(\mathcal{X}_{w,M}^\bullet(I)) \xrightarrow{\varprojlim H^1(\Xi_M)} \varprojlim_M H^1(\mathcal{Q}_{w,M}^\bullet(I)) \right).$$

Compatible closed-exchange cocycles and compatible counterterms solving

$$\mathfrak{o}_{n,M}^{\text{ns}} + \Xi_M(W_{n,M}) + d_M C_{n,M} = \mathfrak{o}, \quad \rho_{M,N}^X W_{n,M} = W_{n,N}, \quad \pi_{M,N} C_{n,M} = C_{n,N},$$

exist if and only if  $\beta_n^{\text{cl}} = \mathfrak{o}$  and one can choose the following secondary data with zero Roos classes. Vanishing of  $\beta_n^{\text{cl}}$  is equivalent to the existence of a compatible cohomology class

$$\alpha = (\alpha_M)_M \in \varprojlim_M H^1(\mathcal{X}_{w,M}^\bullet(I))$$

with

$$H^1(\Xi_M)(\alpha_M) = -[\mathfrak{o}_{n,M}^{\text{ns}}] \quad \text{for every } M.$$

For such a choice of  $\alpha$ , the representative-compatibility class

$$\mu_n^{\text{cl}}(\alpha) \in \varprojlim_M H^0(\mathcal{X}_{w,M}^\bullet(I))$$

vanishes. Explicitly, choose cocycles  $w_M \in Z^1(\mathcal{X}_{w,M}^\bullet(I))$  representing  $\alpha_M$ . Since  $\alpha$  is compatible, for the cofinal successor tower one can choose  $u_M \in \mathcal{X}_{w,M}^\circ(I)$  with

$$\rho_{M+1,M}^X w_{M+1} - w_M = d_M^X u_M.$$

With the Roos convention

$$\Delta_X((a_M)_M) = (\rho_{M+1,M}^X a_{M+1} - a_M)_M,$$

the class of  $([u_M])_M$  in  $\text{coker } \Delta_X = \varprojlim_M H^0(X_{w,M}^\bullet(I))$  is  $\mu_n^{\text{cl}}(\alpha)$ . Its vanishing is exactly the condition that the representatives  $w_M$  can be changed by degree-zero coboundaries to compatible cocycles  $W_{n,M}$ .

After choosing such compatible  $W_{n,M}$ , the counterterm compatibility class

$$\lambda_n^{\text{cl}}(W) \in \varprojlim_M H^0(Q_{w,M}^\bullet(I))$$

vanishes. Namely, set

$$r_M = \mathfrak{o}_{n,M}^{\text{ns}} + \Xi_M(W_{n,M}).$$

The class of  $r_M$  in  $H^1(Q_{w,M}^\bullet(I))$  is zero, so choose  $c_M \in Q_{w,M}^\circ(I)$  with  $d_M c_M = -r_M$ . The differences

$$\pi_{M+1,M} c_{M+1} - c_M \in Z^0(Q_{w,M}^\bullet(I))$$

define, with the analogous Roos operator  $\Delta_Q$ , the class

$$\lambda_n^{\text{cl}}(W) = [(\pi_{M+1,M} c_{M+1} - c_M)_M] \in \text{coker } \Delta_Q$$

This class vanishes exactly when the primitives  $c_M$  can be changed by degree-zero cocycles and coboundaries to compatible counterterms  $C_{n,M}$ .

If the inverse system  $\{H^0(X_{w,M}^\bullet(I))\}_M$  is Mittag–Leffler, then  $\mu_n^{\text{cl}}(\alpha) = 0$  for every admissible lift  $\alpha$ . If  $\{H^0(Q_{w,M}^\bullet(I))\}_M$  is Mittag–Leffler, then  $\lambda_n^{\text{cl}}(W) = 0$  for every compatible closed-exchange representative  $W$ . Under both Mittag–Leffler hypotheses, the compatible closed-exchange branch is obstructed only by  $\beta_n^{\text{cl}}$ . The counterterm-only criterion of Theorem 5.83 is the special case  $X_{w,M}^\bullet(I) = 0$ . The admissible weight transports  $R_{w,w'}^M$  identify the fixed-window image condition,  $\beta_n^{\text{cl}}$ , and the secondary Roos classes for  $w$  with the corresponding data for  $w'$ .

*Proof.* The fixed-window statement is the cohomology calculation. If the branch equation holds, then passing to  $H^1$  gives  $[\mathfrak{o}_{n,M}^{\text{ns}}] = -H^1(\Xi_M)[W_{n,M}]$ . Conversely, if the displayed image condition holds, choose a cocycle  $W_{n,M}$  in the closed-exchange complex whose class maps to  $-[\mathfrak{o}_{n,M}^{\text{ns}}]$ . Then  $\mathfrak{o}_{n,M}^{\text{ns}} + \Xi_M(W_{n,M})$  is a  $d_M$ -boundary, so a degree-zero  $C_{n,M}$  gives the branch equation.

The compatibility of the residuals and the identity  $\pi_{M,N} \Xi_M = \Xi_N \rho_{M,N}^X$  make  $([\mathfrak{o}_{n,M}^{\text{ns}}])_M$  a point of  $\varprojlim_M H^1(Q_{w,M}^\bullet(I))$ . Its cokernel class is zero exactly when a compatible cohomology class  $\alpha$  in the closed-exchange tower maps to its negative. This proves the first inverse-limit obstruction.

Fix such an  $\alpha$ . The differences  $\rho_{M+1,M}^X w_{M+1} - w_M$  are closed degree-one elements with zero cohomology class, hence have the form  $d_M^X u_M$ . Changing the representatives  $w_M$  by degree-zero coboundaries changes  $([u_M])_M$  by the Roos coboundary  $\Delta_X$ . Thus the displayed class  $\mu_n^{\text{cl}}(\alpha)$  is independent of the choices. In the two-term Roos complex computing the first derived inverse limit of the tower  $H^0(X_{w,M}^\bullet(I))$ , its vanishing is exactly the equation needed to absorb these differences by changing  $w_M$ . This is the exact condition for compatible cocycles  $W_{n,M}$ .

With compatible  $W_{n,M}$ , the cochains  $r_M = \mathfrak{o}_{n,M}^{\text{ns}} + \Xi_M(W_{n,M})$  are compatible  $d_M$ -cocycles whose windowwise  $H^1$ -classes vanish. Choose primitives  $c_M$ . Since  $\pi_{M+1,M} r_{M+1} = r_M$ , the differences  $\pi_{M+1,M} c_{M+1} - c_M$  are  $d_M$ -closed of degree zero. Changing the primitives by closed degree-zero cochains, and by boundaries, changes their cohomology classes by the Roos coboundary  $\Delta_Q$ . Hence the displayed  $\lambda_n^{\text{cl}}(W)$  is independent of primitive choices and vanishes exactly when the primitives can be adjusted to a compatible system  $C_{n,M}$ . This system is precisely the desired compatible local counterterm family.

The last paragraph is the standard vanishing of  $\lim^1$  for Mittag–Leffler inverse systems of vector spaces. Therefore an  $H^0$ -Mittag–Leffler hypothesis on the closed-exchange tower kills the representative obstruction  $\mu_n^{\text{cl}}$ , and the corresponding hypothesis on the non-scalar counterterm tower kills  $\lambda_n^{\text{cl}}$ . If both hold, only the cokernel  $H^1$ -class remains. If the closed-exchange tower is zero, the cokernel condition reduces to vanishing of  $([\mathfrak{o}_{n,M}^{\text{ns}}])_M$ , and the only secondary class is the  $\lambda_n$  of Theorem 5.83. The weight-transport assertion follows because  $R_{w,w'}^M$  is a compatible cochain isomorphism on both towers and intertwines  $\Xi_M$ , residuals, truncation maps, cocycle representatives, and primitive-difference cocycles.  $\square$

**PROPOSITION 5.88** (*Regular-density quotient obstruction for the first residual*). Work in the regular-density tower of Definition 5.84 and Proposition 5.85. Let  $n_o$  be the first  $\hbar$ -order at which the lower-order choices leave a compatible non-scalar residual

$$\mathfrak{o}_{n_o,M}^{\text{ns}} \in Z^1(Q_{w,M}^\bullet(I)), \quad \pi_{M,N} \mathfrak{o}_{n_o,M}^{\text{ns}} = \mathfrak{o}_{n_o,N}^{\text{ns}}.$$

Put

$$\begin{aligned} C_{w,M}^{\bullet,\text{reg}}(I) &= Q_{w,M}^\bullet(I) / X_{w,M}^{\bullet,\text{reg}}(I), \\ \bar{d}_M &= d_M \bmod X_{w,M}^{\bullet,\text{reg}}(I), \\ q_M &: Q_{w,M}^\bullet(I) \rightarrow C_{w,M}^{\bullet,\text{reg}}(I). \end{aligned}$$

The maps  $\pi_{M,N}$  and  $R_{w,w'}^M$  descend to quotient cochain maps on  $C^{\bullet,\text{reg}}$ . Define the finite-window quotient class

$$\eta_{n_o,M}^{\text{reg}} = [q_M(\mathfrak{o}_{n_o,M}^{\text{ns}})] \in H^1(C_{w,M}^{\bullet,\text{reg}}(I)).$$

Then, for a fixed  $M$ ,

$$[\mathfrak{o}_{n_o,M}^{\text{ns}}] \in -\text{im } H^1(\Xi_M^{\text{reg}}) \iff \eta_{n_o,M}^{\text{reg}} = \mathfrak{o}.$$

Equivalently, the regular-density closed-exchange branch at window  $M$  is solvable exactly when the image of the residual is exact in  $Q_{w,M}^\bullet(I) / X_{w,M}^{\bullet,\text{reg}}(I)$ .

The compatible classes

$$\eta_{n_o}^{\text{reg}} = (\eta_{n_o,M}^{\text{reg}})_M \in \varprojlim_M H^1(C_{w,M}^{\bullet,\text{reg}}(I))$$

are the quotient shadow of the closed-exchange obstruction

$$\beta_{n_o}^{\text{cl,reg}} \in \text{coker} \left( \varprojlim_M H^1(X_{w,M}^{\bullet,\text{reg}}(I)) \xrightarrow{\varprojlim_M H^1(\Xi_M^{\text{reg}})} \varprojlim_M H^1(Q_{w,M}^\bullet(I)) \right).$$

If  $\eta_{n_o}^{\text{reg}} \neq \mathfrak{o}$ , then  $\beta_{n_o}^{\text{cl,reg}} \neq \mathfrak{o}$ . On the zero locus of  $\eta_{n_o}^{\text{reg}}$ , choose windowwise classes

$$\alpha_M \in H^1(X_{w,M}^{\bullet,\text{reg}}(I)), \quad H^1(\Xi_M^{\text{reg}})(\alpha_M) = -[\mathfrak{o}_{n_o,M}^{\text{ns}}].$$

Let

$$K_M^{\text{reg}} = \ker \left( H^1(\Xi_M^{\text{reg}}) : H^1(X_{w,M}^{\bullet,\text{reg}}(I)) \rightarrow H^1(Q_{w,M}^\bullet(I)) \right).$$

For the cofinal successor tower the differences

$$\rho_{M+1,M}^{X,\text{reg}} \alpha_{M+1} - \alpha_M \in K_M^{\text{reg}}$$

define a Roos class

$$\kappa_{n_o}^{\text{reg}} \in \varprojlim_M K_M^{\text{reg}}.$$

It is independent of the chosen windowwise lifts up to Roos coboundary, and

$$\beta_{n_o}^{\text{cl,reg}} = 0 \iff \eta_{n_o}^{\text{reg}} = 0 \text{ and } \kappa_{n_o}^{\text{reg}} = 0.$$

In particular, suppose the branch is restricted to a complete regular-density subhabitat

$$Q_{w,M}^{\bullet,\text{rd}}(I) \subset Q_{w,M}^{\bullet}(I)$$

on which

$$Q_{w,M}^{\bullet,\text{rd}}(I) = X_{w,M}^{\bullet,\text{reg}}(I)$$

compatibly with truncation and admissible weight transport, and assume  $\mathfrak{o}_{n_o,M}^{\text{ns}} \in Z^1(Q_{w,M}^{\bullet,\text{rd}}(I))$  for all  $M$ . Then

$$\beta_{n_o}^{\text{cl,reg}} = 0.$$

The canonical compatible choice  $W_{n_o,M} = -\mathfrak{o}_{n_o,M}^{\text{ns}}$  and  $C_{n_o,M} = 0$  has

$$\mu_{n_o}^{\text{cl}} = 0, \quad \lambda_{n_o}^{\text{cl}} = 0.$$

Thus a regular-density equality  $X^{\bullet,\text{reg}} = Q^{\bullet,\text{rd}}$  kills the first residual only for residuals lying in that specified subhabitat. Outside it, the exact obstruction is the quotient/Roos pair  $(\eta_{n_o}^{\text{reg}}, \kappa_{n_o}^{\text{reg}})$  above.

*Proof.* Proposition 5.85 says that  $X_{w,M}^{\bullet,\text{reg}}(I)$  is a complete  $d_M$ -stable subcomplex of  $Q_{w,M}^{\bullet}(I)$ , and that truncation and admissible weight transport preserve this subcomplex. Therefore the quotient complexes and their transition maps are defined. Since the residuals commute with truncation, the classes  $\eta_{n_o,M}^{\text{reg}}$  form an inverse-system element; weight transport gives the same statement for  $w'$ .

For a fixed window, the short exact sequence of complexes

$$0 \longrightarrow X_{w,M}^{\bullet,\text{reg}}(I) \xrightarrow{\Xi_M^{\text{reg}}} Q_{w,M}^{\bullet}(I) \xrightarrow{q_M} C_{w,M}^{\bullet,\text{reg}}(I) \longrightarrow 0$$

gives the long exact cohomology sequence. Exactness at  $H^1(Q_{w,M}^{\bullet}(I))$  gives

$$\text{im } H^1(\Xi_M^{\text{reg}}) = \ker H^1(q_M).$$

The sign is irrelevant over  $\mathbb{C}$ . Hence the image criterion is exactly the vanishing of  $[q_M(\mathfrak{o}_{n_o,M}^{\text{ns}})]$ . Equivalently, if  $q_M(\mathfrak{o}_{n_o,M}^{\text{ns}}) = -\bar{d}_M \bar{C}_M$ , choose any lift  $C_M$  of  $\bar{C}_M$ . Then  $\mathfrak{o}_{n_o,M}^{\text{ns}} + d_M C_M$  lies in  $X_{w,M}^{\bullet,\text{reg}}(I)$  and is closed, so  $W_{n_o,M} = -(\mathfrak{o}_{n_o,M}^{\text{ns}} + d_M C_M)$  solves the branch equation. The converse follows by applying  $q_M$  to the branch equation.

Passing to the inverse limit, the quotient maps induce a map from the displayed cokernel to  $\varprojlim_M H^1(C_{w,M}^{\bullet,\text{reg}}(I))$ , and the image of  $\beta_{n_o}^{\text{cl,reg}}$  is precisely  $\eta_{n_o}^{\text{reg}}$ . Thus nonvanishing of  $\eta_{n_o}^{\text{reg}}$  forces nonvanishing of  $\beta_{n_o}^{\text{cl,reg}}$ .

Assume now  $\eta_{n_o}^{\text{reg}} = 0$ . Windowwise exactness gives the classes  $\alpha_M$ . Since both  $\rho_{M+1,M}^{X,\text{reg}} \alpha_{M+1}$  and  $\alpha_M$  map to  $-\mathfrak{o}_{n_o,M}^{\text{ns}}$ , their difference lies in  $K_M^{\text{reg}}$ . Changing the windowwise lift  $\alpha_M$  by an element of  $K_M^{\text{reg}}$  changes the difference family by the Roos coboundary. Hence the class  $\kappa_{n_o}^{\text{reg}}$  is well defined. It

vanishes exactly when the  $\alpha_M$  can be adjusted to a compatible element of  $\lim_{\leftarrow M} H^1(\mathcal{X}_{w,M}^{\bullet,\text{reg}}(I))$  mapping to  $-([\mathfrak{o}_{n_o,M}^{\text{ns}}])_M$ . This is exactly the vanishing of the cokernel class  $\beta_{n_o}^{\text{cl},\text{reg}}$ .

On a subhabitat  $\mathcal{Q}^{\bullet,\text{rd}} = \mathcal{X}^{\bullet,\text{reg}}$ , the inclusion is the identity. If the residual lies in that subhabitat, then  $W_{n_o,M} = -\mathfrak{o}_{n_o,M}^{\text{ns}}$  is a compatible  $d_M$ -cocycle in  $\mathcal{X}_{w,M}^{\text{reg}}(I)$ , and  $C_{n_o,M} = \mathfrak{o}$  gives

$$\mathfrak{o}_{n_o,M}^{\text{ns}} + \Xi_M^{\text{reg}}(W_{n_o,M}) + d_M C_{n_o,M} = \mathfrak{o}.$$

The representative differences defining  $\mu_{n_o}^{\text{cl}}$  are zero because the residuals are compatible, and the primitive differences defining  $\lambda_{n_o}^{\text{cl}}$  are zero because every primitive is chosen to be  $\mathfrak{o}$ . This proves the saturated regular-density claim.  $\square$

**PROPOSITION 5.89** (*First regular-density residual criterion*). The first non-scalar residual used in the closed-exchange criterion is the curvature coefficient determined by the chosen scalar-contact chain projection, the chosen lower-order counterterms, and the finite window  $M$ :

$$\mathfrak{o}_{n_o,M}^{\text{ns}} = [\hbar^{n_o}] \left( \text{Ob}_{w,\partial,\hbar,M}^{\text{red}} + d_M C_{<n_o,M} + \frac{1}{2} \{C_{<n_o,M}, C_{<n_o,M}\}_{\hbar,M} \right) \in Z^1(\mathcal{Q}_{w,M}^{\bullet}(I)),$$

where  $n_o$  is the first order at which this compatible family survives the lower-order choices. Thus the residual lies in the saturated regular-density subhabitat

$$\mathcal{Q}_{w,M}^{\bullet,\text{rd}}(I) = \mathcal{X}_{w,M}^{\bullet,\text{reg}}(I)$$

if and only if the following regular-density residual criterion is supplied for every finite window  $M$ , compatibly with  $\pi_{M,N}$  and  $R_{w,w'}^M$ .

(R1) The displayed curvature coefficient is represented, in  $\mathcal{Q}_{w,M}^{\text{I}}(I)$ , by a complete filtered sum whose image in every finite filtration quotient is finite:

$$\sum_{\Gamma,\ell} a_{\Gamma,\ell} W_{\Gamma,L,w}^{R,\text{reg},M}(a_{\Gamma,\ell,\bullet}, b_{\Gamma,\ell,\bullet} dt; f_{\Gamma,\ell,\bullet}, \rho_{\Gamma,\ell,\bullet}) + d_M B_{n_o,M}^{\text{reg}},$$

where all defect cotangent labels are smooth compactly supported densities  $b_{\Gamma,\ell,j}(t)dt$ , all coefficient labels lie in degree  $\leq M$ , and  $B_{n_o,M}^{\text{reg}}$  is a regular-density local counterterm in the graph/counterterm complex.

(R2) The representative has zero scalar-contact projection:

$$\widehat{\sigma}_{\text{sc},M}(\mathfrak{o}_{n_o,M}^{\text{ns}}) = \mathfrak{o}$$

(R3) The equality with the curvature coefficient above holds in the local-functional complex  $\mathcal{O}_{\text{loc}}^{\text{bd}}(\mathcal{E}_w^M|_I; A_{\partial,\hbar}^{\text{pp},w,M}(I))$  before cohomology and before scalar reduction.

(R4) The representatives commute with finite-window truncation and admissible weight transport:

$$\pi_{M,N} \mathfrak{o}_{n_o,M}^{\text{ns}} = \mathfrak{o}_{n_o,N}^{\text{ns}}, \quad R_{w,w'}^M \mathfrak{o}_{n_o,M,w}^{\text{ns}} = \mathfrak{o}_{n_o,M,w'}^{\text{ns}}$$

If this criterion holds, then

$$\mathfrak{o}_{n_o,M}^{\text{ns}} \in Z^1(\mathcal{X}_{w,M}^{\bullet,\text{reg}}(I))$$

for every  $M$ . The closed-exchange equation is solved by the explicit compatible representatives

$$W_{n_o,M} = -\mathfrak{o}_{n_o,M}^{\text{ns}}, \quad C_{n_o,M} = \mathfrak{o},$$

and hence

$$\eta_{n_o}^{\text{reg}} = 0, \quad \kappa_{n_o}^{\text{reg}} = 0, \quad \beta_{n_o}^{\text{cl,reg}} = 0, \quad \mu_{n_o}^{\text{cl}} = 0, \quad \lambda_{n_o}^{\text{cl}} = 0$$

for this saturated branch.

When the criterion is unmet, the regular-density branch remains undecided. The exact computable obstruction is

$$\eta_{n_o, M}^{\text{reg}} = \left[ q_M [\hbar^{n_o}] \left( \text{Ob}_{w, \partial, \hbar, M}^{\text{red}} + d_M C_{< n_o, M} + \frac{1}{2} \{ C_{< n_o, M}, C_{< n_o, M} \}_{\hbar, M} \right) \right] \in H^1(C_{w, M}^{\bullet, \text{reg}}(I), \bar{d}_M).$$

Computing this class requires the finite-window coefficient array of  $\text{Ob}_{w, \partial, \hbar, M}^{\text{red}}$ , the lower-order counterterms  $C_{< n_o, M}$ , the scalar-contact chain projection  $\widehat{\sigma}_{\text{sc}, M}$ , and the reduction of the displayed degree-one cochain modulo the regular-density graph/counterterm subcomplex. If all these quotient classes vanish, choose quotient primitives  $\bar{C}_{n_o, M}$  and lifts  $C_{n_o, M} \in \mathcal{Q}_{w, M}^o(I)$  with

$$q_M(\mathfrak{o}_{n_o, M}^{\text{ns}} + d_M C_{n_o, M}) = 0.$$

Then

$$x_{n_o, M} := \mathfrak{o}_{n_o, M}^{\text{ns}} + d_M C_{n_o, M} \in Z^1(\mathcal{X}_{w, M}^{\bullet, \text{reg}}(I)), \quad \alpha_M = [-x_{n_o, M}]$$

is a windowwise lift of  $-\mathfrak{o}_{n_o, M}^{\text{ns}}$ . The remaining cohomology-lift obstruction is precisely

$$\kappa_{n_o}^{\text{reg}} = \left[ \left( \rho_{M+1, M}^{X, \text{reg}} \alpha_{M+1} - \alpha_M \right)_M \right] \in \varprojlim_M \ker H^1(\Xi_M^{\text{reg}}).$$

Only after this class vanishes does the regular-density branch have a compatible cohomology lift; the representative and primitive compatibility classes  $\mu_{n_o}^{\text{cl}}$  and  $\lambda_{n_o}^{\text{cl}}$  are then the secondary Roos classes of Theorem 5.87.

*Proof.* The formula for  $\mathfrak{o}_{n_o, M}^{\text{ns}}$  is the order  $n_o$  specialization of Theorem 5.83. The Bianchi identity in the QME complex gives  $d_M \mathfrak{o}_{n_o, M}^{\text{ns}} = 0$ , and the chosen scalar-contact chain projection places it in  $\mathcal{Q}_{w, M}^{\bullet}(I)$ .

Conditions (R1)–(R4) are exactly the definition of membership in the complete regular-density graph/counterterm subcomplex constructed in Definition 5.84 and Proposition 5.85. Hence the criterion is equivalent to  $\mathfrak{o}_{n_o, M}^{\text{ns}} \in Z^1(\mathcal{X}_{w, M}^{\bullet, \text{reg}}(I))$  compatibly in the finite-window tower. Since  $\Xi_M^{\text{reg}}$  is inclusion on the saturated subhabitat,  $W_{n_o, M} = -\mathfrak{o}_{n_o, M}^{\text{ns}}$  and  $C_{n_o, M} = 0$  give

$$\mathfrak{o}_{n_o, M}^{\text{ns}} + \Xi_M^{\text{reg}}(W_{n_o, M}) + d_M C_{n_o, M} = 0$$

Compatibility of the residual family makes the Roos differences for  $W$  vanish, and the zero primitives make the counterterm Roos differences vanish. This proves the displayed vanishing of  $\eta$ ,  $\kappa$ ,  $\beta$ ,  $\mu$ , and  $\lambda$ .

This criterion supplies the invariant operation beyond quotienting by  $\mathcal{X}^{\bullet, \text{reg}}$ . Applying Proposition 5.88 to the explicit curvature coefficient gives the displayed  $\eta_{n_o, M}^{\text{reg}}$ . Its computation needs precisely the coefficient array of the curvature coefficient, the lower-order counterterms entering it, the chain projection defining  $Q$ , and the quotient reduction modulo  $\mathcal{X}^{\text{reg}}$ . If  $\eta_{n_o, M}^{\text{reg}} = 0$ , a quotient primitive gives  $q_M(\mathfrak{o}_{n_o, M}^{\text{ns}} + d_M C_{n_o, M}) = 0$ , hence  $x_{n_o, M} \in \mathcal{X}_{w, M}^{1, \text{reg}}(I)$ . Closedness follows from  $d_M \mathfrak{o}_{n_o, M}^{\text{ns}} = 0$  and  $d_M^2 = 0$ . The Roos differences of the resulting cohomology classes lie in  $\ker H^1(\Xi_M^{\text{reg}})$  because both terms map to  $-\mathfrak{o}_{n_o, M}^{\text{ns}}$ . This is exactly the  $\kappa_{n_o}^{\text{reg}}$  obstruction of Proposition 5.88.  $\square$



PROPOSITION 5.90 (*Regular-density criterion for the evaluated curvature*). Work at a fixed finite window  $M$ . Assume the finite-window graph/QME system of Definition 8.21 is the regular-density system: every cotangent defect label is  $B_j = b_j(t)dt \in \mathcal{D}_c^{\text{reg}}(I)$ , the finite graph list is closed under the  $d_M$ -boundary and BV-bracket terms appearing in the system, and the scalar-contact projection is zero on the chosen subhabitat. Let

$$\text{Ob}_{w,\partial,\hbar,M}^{\text{red,reg}} = \text{ev}_{\mathcal{G}_M} \text{Curv}_{\mathbf{K}}(\kappa_{\hbar,w,I}^{\text{graph,reg},M}) \in \mathcal{Q}_{w,M}^1(I)$$

be the evaluated reduced curvature of the regular-density graph kernel. Suppose the lower-order counterterms are compatible and regular-density:

$$C_{<n_o,M}^{\text{reg}} \in \mathcal{Q}_{w,M}^{\text{o,rd}}(I) = \mathcal{X}_{w,M}^{\text{o,reg}}(I), \quad \pi_{M,N} C_{<n_o,M}^{\text{reg}} = C_{<n_o,N}^{\text{reg}}.$$

Put

$$\mathfrak{d}_{n_o,M}^{\text{ns,reg}} = [\hbar^{n_o}] \left( \text{Ob}_{w,\partial,\hbar,M}^{\text{red,reg}} + d_M C_{<n_o,M}^{\text{reg}} + \frac{1}{2} \{C_{<n_o,M}^{\text{reg}}, C_{<n_o,M}^{\text{reg}}\}_{\hbar,M} \right).$$

Then

$$\mathfrak{d}_{n_o,M}^{\text{ns,reg}} \in Z^1(\mathcal{Q}_{w,M}^{\bullet,\text{rd}}(I)) = Z^1(\mathcal{X}_{w,M}^{\bullet,\text{reg}}(I))$$

compatibly with finite-window truncation and admissible weight transport.

More explicitly, after expanding the finite graph system at the coefficient of  $\hbar^{n_o}$ , there is a finite indexing set  $\mathcal{A}_{n_o,M}^{\text{reg}}$  and coefficients  $\varepsilon_{\alpha,M} \in \mathbb{C}$  such that

$$\mathfrak{d}_{n_o,M}^{\text{ns,reg}} = \sum_{\alpha \in \mathcal{A}_{n_o,M}^{\text{reg}}} \varepsilon_{\alpha,M} G_{\alpha,w}^M + [\hbar^{n_o}] d_M C_{<n_o,M}^{\text{reg}} + \frac{1}{2} [\hbar^{n_o}] \{C_{<n_o,M}^{\text{reg}}, C_{<n_o,M}^{\text{reg}}\}_{\hbar,M},$$

where each graph generator is of the form

$$G_{\alpha,w}^M = W_{\Gamma_{\alpha,L,w}}^{R,\text{reg},M}(a_{\alpha,\bullet}, b_{\alpha,\bullet} dt; f_{\alpha,\bullet}, \rho_{\alpha,\bullet}), \quad b_{\alpha,j} \in C_c^\infty(I),$$

or is one of the regular-density local counterterm generators attached to such a labelled graph. The last two displayed terms also lie in  $\mathcal{X}_{w,M}^{\bullet,\text{reg}}(I)$ : the first by  $d_M$ -stability, the second by the bracket-closure clause in the finite-window graph/QME system.

Hence the regular-density residual criterion of Proposition 5.89 is supplied for this evaluated curvature. The closed-exchange solution is

$$W_{n_o,M} = -\mathfrak{d}_{n_o,M}^{\text{ns,reg}}, \quad C_{n_o,M} = \text{o},$$

and

$$\eta_{n_o}^{\text{reg}} = \kappa_{n_o}^{\text{reg}} = \beta_{n_o}^{\text{cl,reg}} = \mu_{n_o}^{\text{cl}} = \lambda_{n_o}^{\text{cl}} = \text{o}$$

on this regular-density branch.

If the chosen curvature contains non-regular labelled data, choose any linear splitting modulo  $\mathcal{Q}_{w,M}^{\bullet,\text{rd}}(I)$  and write

$$\text{Ob}_{w,\partial,\hbar,M}^{\text{red}} = \text{Ob}_{w,\partial,\hbar,M}^{\text{red,reg}} + \text{Ob}_{w,\partial,\hbar,M}^{\text{red,-rd}}, \quad C_{<n_o,M} = C_{<n_o,M}^{\text{reg}} + C_{<n_o,M}^{\text{-rd}}.$$

The resulting quotient class is independent of the splitting. The non-regular summand is generated exactly by graph weights

$$W_{\Gamma,L,w}^{R,\text{WF},M}(a_{\bullet}, B_{\bullet}; f_{\bullet}, \rho_{\bullet}) \quad \text{with some } B_j \notin \mathcal{D}_c^{\text{reg}}(I),$$

by the corresponding distributional current counterterms, and by  $d_M$ - or BV-bracket terms involving such counterterms. Its quotient obstruction is the explicit component

$$\eta_{n_o, M}^{\text{reg}} = \left[ q_M [\hbar^{n_o}] \left( \text{Ob}_{w, \partial, \hbar, M}^{\text{red}, \neg \text{rd}} + d_M C_{< n_o, M}^{\neg \text{rd}} + \{C_{< n_o, M}^{\text{reg}}, C_{< n_o, M}^{\neg \text{rd}}\}_{\hbar, M} + \frac{1}{2} \{C_{< n_o, M}^{\neg \text{rd}}, C_{< n_o, M}^{\neg \text{rd}}\}_{\hbar, M} \right) \right]$$

in

$$H^1(\mathcal{Q}_{w, M}^\bullet(I) / \mathcal{X}_{w, M}^{\bullet, \text{reg}}(I)).$$

Thus a single wavefront-admissible but non-smooth cotangent label, or a distributional current counterterm forced by such a label, is the exact datum that leaves  $Q^{\text{rd}} = X^{\text{reg}}$ .

*Proof.* Proposition B.17 constructs  $W_{\Gamma, L, w}^{R, \text{reg}, M}$  with smooth compactly supported density labels and proves compatibility with interval extension, finite-window truncation, and admissible weight transport. Definition 5.84 then puts all scalar-zero regular graph weights and their regular local counterterms in  $\mathcal{X}_{w, M}^{\bullet, \text{reg}}(I)$ , and closes this space under  $d_M$ . The finite-window graph/QME system includes the BV-bracket terms forced by the finite graph differential, so the quadratic counterterm term remains in the same complete regular-density graph/counterterm complex. Hence every summand in the displayed  $\hbar^{n_o}$ -coefficient lies in  $\mathcal{X}_{w, M}^{\bullet, \text{reg}}(I)$ .

The Bianchi identity in the convolution dg Lie algebra gives closedness of the first unsolved coefficient after the lower-order equations have been imposed. The scalar projection is zero on the chosen regular-density system, so this closed coefficient lies in  $\mathcal{Q}_{w, M}^\bullet(I)$ . The same truncation and weight-transport identities hold term by term for the graph weights, counterterms, differential, and bracket; therefore the residuals are compatible in the finite-window tower.

Since  $\mathcal{Q}_{w, M}^{\bullet, \text{rd}}(I) = \mathcal{X}_{w, M}^{\bullet, \text{reg}}(I)$  on this branch and  $\Xi_M^{\text{reg}}$  is inclusion, the displayed  $W_{n_o, M}$  and  $C_{n_o, M} = 0$  give

$$\mathfrak{d}_{n_o, M}^{\text{ns}, \text{reg}} + \Xi_M^{\text{reg}}(W_{n_o, M}) + d_M C_{n_o, M} = 0.$$

Compatibility of the residual family makes the closed-exchange Roos differences vanish, and the primitive family is identically zero. The quotient class  $\eta_{n_o}^{\text{reg}}$ , the lift class  $\kappa_{n_o}^{\text{reg}}$ , and the three closed-exchange and counterterm obstruction classes therefore vanish.

For a non-regular label set, the quotient map  $q_M$  kills exactly the regular-density subcomplex. Expanding the same curvature expression into regular and non-regular summands therefore leaves only the four non-regular terms displayed in the formula for  $\eta_{n_o, M}^{\text{reg}}$ : the non-regular graph curvature, the  $d_M$ -image of non-regular lower counterterms, the mixed bracket with a regular counterterm, and the purely non-regular bracket. The wavefront-admissible graph layer of Proposition B.19 is proved as a distributional graph layer. When some  $B_j \notin \mathcal{D}_c^{\text{reg}}(I)$ , the term can enter the regular-density branch only if its displayed quotient class is killed by an explicit degree-zero primitive in the ambient distributional quotient complex. This proves the obstruction formula.  $\square$

**THEOREM 5.91** (*Wavefront singular-current obstruction*). Fix a finite window  $M$  and a wavefront-admissible finite graph/QME system as in Proposition B.19. Put

$$\mathcal{S}_c(I) = \mathcal{D}'_c(I) / \mathcal{D}_c^{\text{reg}}(I).$$

A non-smooth label  $B \in \mathcal{D}'_c(I)$  with non-empty wavefront has nonzero class in  $\mathcal{S}_c(I)$ . This quotient class obstructs a canonical regular-density representative of  $B$  modulo  $\mathcal{D}_c^{\text{reg}}(I)$ : if  $B - B^{\text{sm}} \in \mathcal{D}_c^{\text{reg}}(I)$  for some  $B^{\text{sm}} \in \mathcal{D}_c^{\text{reg}}(I)$ , then  $B$  was already regular.

Let the first non-scalar wavefront residual be written as

$$\mathfrak{d}_{n_o, M}^{\text{ns}, \text{WF}} = [\hbar^{n_o}] \left( \text{Ob}_{w, \partial, \hbar, M}^{\text{red}, \text{WF}} + d_M C_{< n_o, M}^{\text{WF}} + \frac{1}{2} \{ C_{< n_o, M}^{\text{WF}}, C_{< n_o, M}^{\text{WF}} \}_{\hbar, M} \right) \in Z^1(\mathcal{Q}_{w, M}^\bullet(I)),$$

with at least one cotangent label in  $\mathcal{D}'_{c, \text{WF}}(I; \Gamma, M) \setminus \mathcal{D}_c^{\text{reg}}(I)$  or at least one distributional current counterterm. Its singular-current component is

$$\begin{aligned} \mathfrak{s}_{n_o, M}^{\text{WF}} = q_M [\hbar^{n_o}] & \left( \text{Ob}_{w, \partial, \hbar, M}^{\text{red}, \neg \text{rd}} + d_M C_{< n_o, M}^{\neg \text{rd}} \right. \\ & \left. + \{ C_{< n_o, M}^{\text{reg}}, C_{< n_o, M}^{\neg \text{rd}} \}_{\hbar, M} + \frac{1}{2} \{ C_{< n_o, M}^{\neg \text{rd}}, C_{< n_o, M}^{\neg \text{rd}} \}_{\hbar, M} \right) \in C_{w, M}^{\text{i}, \text{reg}}(I), \end{aligned} \quad (26)$$

where

$$C_{w, M}^{\bullet, \text{reg}}(I) = \mathcal{Q}_{w, M}^\bullet(I) / \mathcal{X}_{w, M}^{\bullet, \text{reg}}(I).$$

The obstruction class is exactly

$$\eta_{n_o, M}^{\text{reg}} = [\mathfrak{s}_{n_o, M}^{\text{WF}}] \in H^1(C_{w, M}^{\bullet, \text{reg}}(I)).$$

More concretely, for a graph summand  $W_{\Gamma, L, w}^{R, \text{WF}, M}(a_\bullet, B_\bullet; f_\bullet, \rho_\bullet)$ , choose arbitrary smooth densities  $B_j^{\text{sm}} \in \mathcal{D}_c^{\text{reg}}(I)$  and put  $S_j = B_j - B_j^{\text{sm}}$ . By multilinearity,

$$q_M W_{\Gamma, L, w}^{R, \text{WF}, M}(a_\bullet, B_\bullet; f_\bullet, \rho_\bullet) = q_M \sum_{\emptyset \neq J \subset \{1, \dots, r\}} W_{\Gamma, L, w}^{R, \text{WF}, M}(a_\bullet, B_\bullet^J; f_\bullet, \rho_\bullet),$$

where  $B_j^J = S_j$  for  $j \in J$  and  $B_j^J = B_j^{\text{sm}}$  for  $j \notin J$ . This expression is independent of the chosen smooth densities after passing to  $C_{w, M}^{\bullet, \text{reg}}(I)$ . The zero-internal-edge case, whenever the generator lies in the non-scalar kernel, is the current class

$$q_M \iota_I(\Theta_\rho^w(B)) = [B] \otimes \rho \in \mathcal{S}_c(I) \widehat{\otimes} D_{w, \leq M},$$

so a singular cotangent current with nonzero coefficient is already a nonzero regular-density quotient generator.

The wavefront of  $\mathfrak{s}_{n_o, M}^{\text{WF}}$  is contained in the finite union obtained from the graph pushforward of

$$\text{WF}(K_{\Gamma, \alpha}^M) + \text{WF}(B_\Gamma)$$

over the Hadamard coefficients appearing at order  $n_o$ , together with the local counterterm wavefronts supported on

$$\bigcup_{\Delta \in \text{Diag}(\Gamma)} \Delta \cap \text{supp}(B_\Gamma).$$

Thus the component in (26) is the exact wavefront/counterterm component left after quotienting by the regular-density graph complex.

A finite-window counterterm kills the non-regular branch if and only if there is a degree-zero distributional counterterm  $C_{n_o, M}^{\text{sing}} \in \mathcal{Q}_{w, M}^o(I)$  such that

$$\bar{d}_M q_M(C_{n_o, M}^{\text{sing}}) = -\mathfrak{s}_{n_o, M}^{\text{WF}}.$$

In that case  $\eta_{n_o, M}^{\text{reg}} = 0$ , and the remaining compatibility obstruction is the Roos class  $\kappa_{n_o}^{\text{reg}}$  of Proposition 5.88. If every such distributional primitive is obstructed, then

$$\eta_{n_o, M}^{\text{reg}} \neq 0,$$

and the regular-density closed-exchange branch has a smoothing obstruction. The honest enlargement is a new wavefront-labelled closed-exchange complex distinct from  $\mathcal{X}^{\bullet, \text{reg}}$ .

*Proof.* A compactly supported smooth density has empty wavefront set. Therefore a compactly supported distribution with non-empty wavefront has nonzero class modulo smooth densities. This proves the label-level obstruction to canonical smoothing. The zero-internal-edge generator shows that this obstruction is already visible in the current target:  $\Theta_\rho^w(B)$  records the distribution  $B$  linearly in  $\mathcal{D}'_c(I) \widehat{\otimes} D_{w, \leq M}$ , and the quotient by  $\mathcal{X}^{\bullet, \text{reg}}$  kills only the smooth-density subspace.

Proposition B.19 constructs the finite-window wavefront graph weights and the corresponding distributional current counterterms. The regular-density subcomplex of Definition 5.84 contains exactly the smooth-label graph weights, their regular counterterms, and their  $d_M$ -closures. Hence applying  $q_M$  to the first residual removes precisely the regular-density terms and leaves the four non-regular terms displayed in Formula (26). Changing the auxiliary smooth densities  $B_j^{\text{sm}}$  changes the expansion only by graph terms with all labels smooth; these lie in  $\mathcal{X}^{\bullet, \text{reg}}$ , so the quotient expression is independent of the splitting.

Closedness follows from the same Bianchi identity used in Proposition 5.90, after passing to the quotient complex. The wavefront and support containment is exactly the Hörmander product and extension statement in Proposition B.19: graph terms have wavefront contained in the pushforward of the product wavefronts, and local extension ambiguities are supported on the collision diagonals meeting  $\text{supp}(B_\Gamma)$ .

Finally,  $H^1(C_{w, M}^{\bullet, \text{reg}}(I), \bar{d}_M)$  is the cohomology of the quotient complex. Therefore the class represented by  $\mathfrak{s}_{n_o, M}^{\text{WF}}$  vanishes exactly when it is a  $\bar{d}_M$ -boundary, equivalently exactly when a degree-zero distributional counterterm  $C_{n_o, M}^{\text{sing}}$  with the displayed equation exists. For a non-boundary class, the quotient obstruction  $\eta_{n_o, M}^{\text{reg}}$  is nonzero and the regular-density branch is obstructed.  $\square$

**THEOREM 5.92** (*Distributional primitive cone for the wavefront quotient*). Keep the hypotheses and notation of Theorem 5.91, and assume that the chosen wavefront graph/QME systems are compatible under finite-window truncation and admissible weight transport. Thus

$$\pi_{M, N} \mathfrak{o}_{n_o, M}^{\text{ns, WF}} = \mathfrak{o}_{n_o, N}^{\text{ns, WF}}, \quad R_{w, w'}^M \mathfrak{o}_{n_o, M, w}^{\text{ns, WF}} = \mathfrak{o}_{n_o, M, w'}^{\text{ns, WF}}.$$

Let

$$\widehat{\mathfrak{s}}_{n_o, M}^{\text{WF}} := \mathfrak{o}_{n_o, M}^{\text{ns, WF}} \in Z^1(Q_{w, M}^\bullet(I))$$

be the closed residual lift of  $\mathfrak{s}_{n_o, M}^{\text{WF}}$ , so that  $q_M \widehat{\mathfrak{s}}_{n_o, M}^{\text{WF}} = \mathfrak{s}_{n_o, M}^{\text{WF}}$ .

Define the local wavefront-primitive extension by

$$\widetilde{Q}_{w, M}^{\bullet, \text{WFpr}}(I) = Q_{w, M}^\bullet(I) \widehat{\oplus} \mathbb{C}[[\hbar]] \cdot p_{n_o, M}^{\text{WF}}, \quad |p_{n_o, M}^{\text{WF}}| = 0,$$

completed in the same  $\hbar$ -adic, scalar-contact, and finite-window filtrations as  $Q_{w, M}^\bullet(I)$ . The new generator is placed in  $\hbar$ -order  $n_o$ , in the scalar-contact filtration of  $\widehat{\mathfrak{s}}_{n_o, M}^{\text{WF}}$ , and with local support and wavefront label equal to the support and wavefront cone displayed in Theorem 5.91. Set

$$\widetilde{d}_M|_Q = d_M, \quad \widetilde{d}_M p_{n_o, M}^{\text{WF}} = -\widehat{\mathfrak{s}}_{n_o, M}^{\text{WF}}.$$

Let

$$\widetilde{C}_{w,M}^{\bullet,\text{reg}}(I) = \widetilde{Q}_{w,M}^{\bullet,\text{WFpr}}(I) / \chi_{w,M}^{\bullet,\text{reg}}(I),$$

and write again  $q_M$  for the quotient map and  $\bar{d}_M$  for the induced differential. Then  $\bar{d}_M^2 = 0$ , the inclusion  $Q_{w,M}^{\bullet}(I) \hookrightarrow \widetilde{Q}_{w,M}^{\bullet,\text{WFpr}}(I)$  is a filtered cochain map, and the element

$$C_{n_o,M}^{\text{sing}} := p_{n_o,M}^{\text{WF}}$$

satisfies the desired quotient primitive equation

$$\bar{d}_M q_M(C_{n_o,M}^{\text{sing}}) = -\mathfrak{s}_{n_o,M}^{\text{WF}}.$$

The extension is strict Mittag–Leffler on compatible finite-window towers: define

$$\widetilde{\pi}_{M,N} p_{n_o,M}^{\text{WF}} = p_{n_o,N}^{\text{WF}}$$

and let  $\widetilde{\pi}_{M,N}$  be  $\pi_{M,N}$  on  $Q_{w,M}^{\bullet}(I)$ . For admissible weights set

$$\widetilde{R}_{w,w'}^M p_{n_o,M,w}^{\text{WF}} = p_{n_o,M,w'}^{\text{WF}}$$

and let  $\widetilde{R}_{w,w'}^M$  be  $R_{w,w'}^M$  on  $Q_{w,M}^{\bullet}(I)$ . These maps are continuous filtered cochain maps and commute with the quotient to  $\widetilde{C}^{\bullet,\text{reg}}$ .

The new generator  $p_{n_o,M}^{\text{WF}}$  lives in the extended complex, outside the original Costello counterterm complex  $Q_{w,M}^{\bullet}(I)$ , and the construction keeps a singular current distinct from a smooth density. A primitive inside  $Q_{w,M}^{\bullet}(I)$  exists exactly under the vanishing condition  $\eta_{n_o,M}^{\text{reg}} = 0$  of Theorem 5.91.

*Proof.* The residual  $\widehat{\mathfrak{s}}_{n_o,M}^{\text{WF}}$  is  $d_M$ -closed by the Bianchi identity used in Theorem 5.91. Hence

$$\bar{d}_M^2 p_{n_o,M}^{\text{WF}} = -d_M \widehat{\mathfrak{s}}_{n_o,M}^{\text{WF}} = 0,$$

and  $\bar{d}_M^2 = 0$  on the old summand because  $d_M^2 = 0$ . The image of the new generator has the same local support, wavefront cone,  $\hbar$ -order, scalar-contact order, and finite coefficient window assigned to the generator. Therefore the differential is continuous for the completed filtration topology.

Since  $q_M \widehat{\mathfrak{s}}_{n_o,M}^{\text{WF}} = \mathfrak{s}_{n_o,M}^{\text{WF}}$ , applying the quotient map to the defining differential gives

$$\bar{d}_M q_M(p_{n_o,M}^{\text{WF}}) = q_M \bar{d}_M p_{n_o,M}^{\text{WF}} = -\mathfrak{s}_{n_o,M}^{\text{WF}}.$$

This is the displayed primitive equation.

The truncation and weight-transport formulas are cochain identities because the closed residual lifts are compatible:

$$\widetilde{\pi}_{M,N} \bar{d}_M p_{n_o,M}^{\text{WF}} = -\pi_{M,N} \widehat{\mathfrak{s}}_{n_o,M}^{\text{WF}} = -\widehat{\mathfrak{s}}_{n_o,N}^{\text{WF}} = \bar{d}_N \widetilde{\pi}_{M,N} p_{n_o,M}^{\text{WF}},$$

and similarly

$$\widetilde{R}_{w,w'}^M \bar{d}_M p_{n_o,M,w}^{\text{WF}} = -R_{w,w'}^M \widehat{\mathfrak{s}}_{n_o,M,w}^{\text{WF}} = -\widehat{\mathfrak{s}}_{n_o,M,w'}^{\text{WF}} = \bar{d}_M \widetilde{R}_{w,w'}^M p_{n_o,M,w}^{\text{WF}}.$$

On the old summand these are the already proved identities for  $\mathcal{Q}_{w,M}^\bullet(I)$ . The transition maps are surjective on the single new generator and are the old strict Mittag–Leffler maps on  $\mathcal{Q}$ , so the extended tower is again strict Mittag–Leffler. The same formulas descend to the quotient because  $\mathcal{X}^{\bullet,\text{reg}}$  is preserved by truncation and weight transport.

The last assertion is formal. The construction adjoins a new degree-zero generator whose differential is the closed wavefront residual. It leaves the quotient  $\mathcal{D}'_c(I)/\mathcal{D}_c^{\text{reg}}(I)$  unchanged on the original current labels, and therefore leaves singular distributions singular. In the original complex alone, the primitive equation is precisely the statement that  $\mathfrak{s}_{n_o,M}^{\text{WF}}$  is a boundary in the original quotient complex, i.e. that  $\eta_{n_o,M}^{\text{reg}}$  vanishes.  $\square$

**THEOREM 5.93** (*Strict unweighted endpoint obstruction*). Let

$$\mathfrak{h}_\Pi = \prod_{d \geq 1} H_d, \quad D_\oplus = \bigoplus_{d \geq 1} H_d^\vee$$

be the original product/direct-sum Tate pair. The support obstruction excludes a tensor  $K \in \mathfrak{h}_\Pi \widehat{\otimes} D_\oplus$ , and likewise a tensor in  $D_\oplus \widehat{\otimes} \mathfrak{h}_\Pi$ , whose projection to every finite window  $d \leq w_o$  is the finite Casimir

$$C_{\mathfrak{h}}^{w_o} = \sum_{d \leq w_o} \sum_i e_{d,i} \otimes e_d^i.$$

Consequently the strict unweighted pair has the endpoint obstruction for a finite-window Costello regulator theory satisfying the coefficient kernel requirement (C1) and the windowwise compatibility used in Theorem 5.81. Equivalently, the support obstruction persists for every  $q \rightarrow 1^+$  family of weighted finite-window kernels in  $\mathfrak{h}_\Pi \widehat{\otimes} D_\oplus$  with projections  $C_{\mathfrak{h}}^{w_o}$ .

Moreover, for any admissible weight  $w$ , the inclusion  $D_\oplus \hookrightarrow D_{\Pi,w} := \prod_{d \geq 1} w(d)^{-1} H_d^\vee$  is only an admissible filtered weak equivalence. Its cofiber has nonzero  $H^0$ , so any observable or obstruction class detecting the quotient  $D_{\Pi,w}/D_\oplus$  is outside the proved regulator-independent sector.

*Proof.* The direct-sum factor  $D_\oplus$  is coefficient-finite by Lemma 5.165 item (1): every element of a completed tensor product with  $D_\oplus$  has support in only finitely many  $H_d^\vee$ -degrees. If  $K$  projected to  $C_{\mathfrak{h}}^{w_o}$  for all  $w_o$ , then for every degree  $d$  the  $H_d^\vee$ -component of  $K$  would be nonzero. This is infinite support in the direct-sum direction, a contradiction. The same argument applies after reversing the tensor factors.

Costello’s heat-kernel BV propagator for the cotangent coefficient field is  $P_{\epsilon,L}^{\text{base}} \widehat{\otimes} K$ . Since the support obstruction removes  $K$  from the strict product/direct-sum pair, the strict pair has an operator-level heat homotopy while the coefficient distributional inverse pairing and graphwise regulator endpoint are obstructed.

Suppose a  $q \rightarrow 1^+$  endpoint existed in the strict pair. Its finite-window projection would have to be the limit of the weighted finite-window kernels. After the weighted bases are identified with the unweighted basis, every such finite-window kernel is exactly  $C_{\mathfrak{h}}^{w_o}$ , independent of  $q$ . The endpoint would therefore give a tensor  $K$  with the obstructed projections above. This proves the endpoint assertion.

Finally, with zero differential,

$$H^0(D_{\Pi,w}/D_\oplus) = D_{\Pi,w}/D_\oplus.$$

The class of any sequence with one nonzero covector in every degree is nonzero in this quotient. Therefore the completion changes ordinary cohomology even though finite quotients and associated graded pieces have zero image of the tail. This proves the asserted boundary of regulator independence.  $\square$

*Definition 5.94 (Endpoint-admissible coefficient module).* An endpoint for the weighted regulators  $w_q(d) = q^d$ ,  $q > 1$ , is *admissible* if it consists of a complete Hausdorff coefficient pair  $(\mathfrak{h}_*, D_*)$ , finite quotients  $(\mathfrak{h}_*, D_*)^{w_o}$ , and diagonal transport maps from every weighted window to the endpoint window, satisfying:

- (E1) the finite quotients form strict Mittag-Leffler towers with surjective transition maps and endpoint quotients  $H_{\leq w_o} \oplus H_{\leq w_o}^\vee$ ;
- (E2) there is a coefficient kernel  $C_* \in \mathfrak{h}_* \widehat{\otimes} D_*$  whose projection to  $d \leq w_o$  is  $C_{\mathfrak{h}}^{w_o}$  for every  $w_o$ ;
- (E3) the endpoint transport maps carry differential, Poisson bracket, coadjoint action, propagator contraction, interaction, counterterm representatives, and the CE/PV central-operation generator assignment  $c \mapsto \theta$ ,  $u \mapsto O$  to their endpoint versions;
- (E4) local observables compute the admissible filtered cohomology of Statement 5.17, so exactness of the strict Mittag-Leffler limit applies;
- (E5) the Costello RG recursion  $\mathcal{W}(P_{\epsilon, L}, I)$  preserves the endpoint tower and the endpoint counterterms.

*THEOREM 5.95 (Endpoint stabilization criterion).* Let  $w_q(d) = q^d$ ,  $q > 1$ . If there is an endpoint-admissible coefficient module in the sense of Definition 5.94, then the transport maps assemble to isomorphisms

$$R_{q,*}: \varprojlim_{w_o} (O_{\text{loc}}(\mathcal{E}_{w_q}^{w_o}), Q + \{I_o^{w_q}, -\}) \xrightarrow{\cong} \varprojlim_{w_o} (O_{\text{loc}}(\mathcal{E}_*^{w_o}), Q + \{I_o^*, -\})$$

for all  $q > 1$ , functorially in  $q$ . Hence finite-degree local observables, fixed-graph Costello contractions, RG counterterm representatives, and the mixed brane-defect QME obstruction class stabilize at the endpoint. More precisely, if  $O_M^q$  is represented in the finite window  $d \leq M$ , then its endpoint representative is

$$O_M^* = R_{q,*}^M O_M^q$$

and is independent of  $q$  after the common finite-window identification with  $H_{\leq M} \oplus H_{\leq M}^\vee$ . The notation  $q \rightarrow 1^+$  is therefore stabilization of this finite-window representative inside the endpoint module. It is a finite-window stabilization in place of strict product/direct-sum convergence. In particular

$$R_{q,*}(\text{Ob}_q) = \text{Ob}_*, \quad \text{Ob}_q = 0 \iff \text{Ob}_* = 0.$$

*Proof.* Replace the second admissible weight  $w'$  in Theorem 5.81 by the endpoint object. Conditions (E1)–(E5) are exactly the hypotheses used there: strict Mittag-Leffler exactness, existence of the coefficient kernel, transport of the algebraic and graphwise structures, admissible filtered cohomology, and preservation by RG recursion. The same finite-window diagonal maps therefore give isomorphisms of strict inverse systems. Passing to the exact inverse limit identifies cohomology and sends the obstruction cocycle to the endpoint obstruction cocycle. Functoriality follows from the composition law of the transport maps on each finite window.

For a representative supported in  $d \leq M$ , the endpoint statement is already finite-dimensional:  $R_{q,*}^M$  is the diagonal map from the  $q$ -basis to the endpoint basis, and the endpoint transport commutes with the unweighting maps by condition (E3). Thus the finite-window coefficient array is  $q$ -independent. The inverse-limit assertion adds exactly the strict Mittag-Leffler compatibility required by (E1) and exactness in the admissible envelope.  $\square$

THEOREM 5.96 (*Finite-window  $q \rightarrow 1^+$  stabilization of the CE/PV generator map*). Let  $w_q(d) = q^d$ ,  $q > 1$ , and let

$$\Phi_M^q: C_{\text{CE}}^\bullet(\mathfrak{h}_{q,\leq M} \ltimes D_{q,\leq M}[1]) \longrightarrow \text{PV}(\mathbf{S}(\mathfrak{h}_{q,\leq M}))$$

be the finite-window restriction of the weighted CE/PV generator map. Let  $\Phi^{o,M}$  be the finite-dimensional unweighted CE/PV map on  $(\mathfrak{h}_{\leq M}, D_{\leq M})$ . Then

$$U_q^M \Phi_M^q = \Phi^{o,M} U_q^M$$

as cochain  $P_0$ -algebra maps. Consequently the unweighted finite-window representative  $U_q^M \Phi_M^q (U_q^M)^{-1}$  is independent of  $q$  and has  $q \rightarrow 1^+$  limit  $\Phi^{o,M}$ .

If an endpoint-admissible coefficient module exists, the same identity passes to the strict Mittag-Leffler inverse limit:  $R_{q,*} \Phi^q = \Phi^* R_{q,*}$ . For the original product/direct-sum endpoint the obstruction term is precisely

$$\mathfrak{o}_{\text{Cas}} = (C_{\mathfrak{h}}^M)_{M \geq 1} \notin \mathfrak{h}_{\Pi} \widehat{\otimes} D_{\oplus},$$

equivalently the nonexistence of a tensor whose finite-window projections are all  $C_{\mathfrak{h}}^M$ .

*Proof.* On generators,  $\Phi_M^q$  sends the CE coordinate dual to  $q^d e_{d,i}$  to the Schouten vector coordinate dual to  $q^d e_{d,i}$ , and sends the CE coordinate dual to  $q^{-d} e_d^i[1]$  to the Hamiltonian coordinate  $O_{d,i}^q$ . The map  $U_q^M$  removes the same diagonal weight on both the CE and PV sides. Therefore the square commutes on the generators  $c, u, \theta, O$ . Since both CE and Schouten differentials are derivations and their structure constants are transported by the same diagonal conjugation, the generator check gives the equality of dg  $P_0$ -algebra maps.

The displayed equality says that the finite-window map, after being written in the unweighted basis, is exactly the finite CE/PV map  $\Phi^{o,M}$ ; it is  $q$ -independent. Hence the  $q \rightarrow 1^+$  limit exists on every finite window.

Under Definition 5.94, the endpoint maps commute with finite-window transition maps and carry the CE/PV generator assignment to the endpoint assignment. Exactness of the strict Mittag-Leffler inverse limit then passes the finite-window identity to the endpoint. In the original product/direct-sum pair, condition (E2) is violated by Theorem 5.93; the obstruction is the compatible family of finite Casimirs whose global tensor is obstructed in  $\mathfrak{h}_{\Pi} \widehat{\otimes} D_{\oplus}$ .  $\square$

COROLLARY 5.97 (*Strict direct-sum endpoint obstruction criterion*). The strict product/direct-sum pair  $(\mathfrak{h}_{\Pi}, D_{\oplus})$  violates the endpoint-admissibility criterion exactly at condition (E2): the compatible family of finite Casimirs has an obstructed representing coefficient kernel in  $\mathfrak{h}_{\Pi} \widehat{\otimes} D_{\oplus}$ .

*Proof.* Condition (E2) is exactly the coefficient-kernel condition ruled out by Theorem 5.93.  $\square$

PROPOSITION 5.98 (*The completion changes ordinary cohomology*). Let

$$D_{\oplus} = \bigoplus_{d \geq 1} H_d^{\vee}, \quad D_{\Pi,w} = \prod_{d \geq 1} w(d)^{-1} H_d^{\vee}$$

with zero differential. The finite-window inclusion  $D_{\oplus} \hookrightarrow D_{\Pi,w}$  induces an isomorphism on every quotient by the tail  $d > w_0$ , and therefore is a weak equivalence in the admissible filtered sense of Statement 5.17. Its ordinary cofiber has nonzero degree-zero cohomology:

$$H^0(D_{\Pi,w}/D_{\oplus}) = D_{\Pi,w}/D_{\oplus} \neq 0,$$



for example the class of any sequence with one nonzero covector in every degree is nonzero. Thus the weighted product contains tail states omitted by the strict continuous dual. Any observable detecting such a tail is outside the regulator-admissible sector of Definition 5.75.

*Example 5.99 (Super-exponential bracket obstruction).* The super-exponential function  $w(d) = \exp(d^2)$  violates the degree weight condition: the ratio

$$\frac{w(p+q-2)}{w(p)w(q)} = \exp((p+q-2)^2 - p^2 - q^2)$$

is unbounded along  $p = q \rightarrow \infty$ , so condition (W1) is violated. The Poisson bracket violates bracket-tameness in that coefficient topology, and the graphwise constants in Proposition 5.68 can be unbounded in the degrees used by the interaction. Theorem 5.81 applies to the admissible weighted theory.

*Nilpotent truncation  $\mathfrak{o}_{\text{nilp}}$ .* The truncation  $\mathfrak{h}_{\leq N} = \mathfrak{m}^3/\mathfrak{m}^{N+1}$  is finite-dimensional, nilpotent of class  $\leq N-2$ , and supports the finite classical CE/PV identification of 5.101. The full Hamiltonian recovery from the inverse system  $\{\mathfrak{h}_{\leq N}\}_N$  has its obstruction in the low-degree sector  $\mathfrak{m}/\mathfrak{m}^3 = H_1 \oplus H_2$ ; the obstruction  $\mathfrak{o}_{\text{nilp}}$  is the triple

$$\mathfrak{o}_{\text{nilp}} = (\mathfrak{o}_N^{\text{lin}}, \mathfrak{o}^{\text{ss}}, \mathfrak{o}_{\leq 2}^{\text{Cas}})$$

of Definition 5.105: the linear-Hamiltonian tail-noninvariance defect  $\mathfrak{o}_N^{\text{lin}}(z_1; z_2^{N+1}) = (N+1)z_2^N$ , the semisimple bracket  $\text{im } \mathfrak{o}^{\text{ss}} = H_2 \cong \mathfrak{sl}_2$  preventing lower-central Tate-pronilpotence, and the missing low-degree Casimir component  $\mathfrak{o}_{\leq 2}^{\text{Cas}}$  outside any finite  $\mathfrak{m}$ -adic window. By Lemma 5.106 a finite  $\mathfrak{m}$ -adic tower extending  $\{\mathfrak{h}_{\leq N}\}_N$  to include  $H_1 \oplus H_2$  exists exactly when these three components vanish; the construction problem is therefore the joint vanishing together with a coefficient category in which the full Casimir family of 5.108 converges, and a relative or reductive replacement for the lower-central Tate-pronilpotent hypothesis treating the semisimple  $\mathfrak{sl}_2 \subset H_2$  directly. At quantum level the same finite truncation is obstructed by 5.102: the Moyal bracket of  $z_1^4, z_2^4$  carries the  $\hbar^2$ -residue  $24\hbar^2 z_1 z_2 \notin \mathfrak{m}^3$ , so a finite quantum CE source on  $\mathfrak{h}_{\leq N}$  requires a separate Moyal-flat truncation or a projected bracket with new Jacobi proof.

#### *Nilpotent truncation of the cotangent module*

The CE/PV Koszul resolution of the brane-side derived  $E_1$ -Hochschild  $HH^\bullet(A_{\partial, N}^{\text{cl}}, A_{\partial, N}^{\text{cl}})$ , identified with the bulk stalk  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b$  by Theorem 5.6, is constructed unconditionally on the cubic ideal  $\mathfrak{m}^3 \subset \mathfrak{h}$ ; its extension to the full reduced Hamiltonian Lie algebra  $\mathfrak{h} = \mathfrak{m} = (z_1, z_2)\mathbb{C}[[z_1, z_2]]$  is Problem 5.109, an open problem whose obstruction to convergence is the explicit triple  $(\mathfrak{o}_N^{\text{lin}}, \mathfrak{o}^{\text{ss}}, \mathfrak{o}_{\leq 2}^{\text{Cas}})$ .

For every  $N \geq 1$ ,

$$\{z_1, z_2^{N+1}\} = (N+1)z_2^N \notin \mathfrak{m}^{N+1}, \quad \{z_1^2, z_2^2\} = 4z_1 z_2, \quad [H_2, H_2] = H_2 \cong \mathfrak{sl}_2.$$

The first identity is the linear-tail obstruction  $\mathfrak{o}_N^{\text{lin}}(z_1; z_2^{N+1}) = (N+1)z_2^N$ , ruling out  $\mathfrak{m}^{N+1}$  as a Lie ideal once  $z_1 \in H_1$  is admitted. The last two encode the semisimple obstruction  $\mathfrak{o}^{\text{ss}}: \Lambda^2 H_2 \rightarrow H_2$  with image  $\mathfrak{sl}_2$ , ruling out nilpotence of  $H_2$ . Together with the low-degree Casimir component  $\mathfrak{o}_{\leq 2}^{\text{Cas}} = \sum_i e_{1,i} \otimes e_1^i + \sum_i e_{2,i} \otimes e_2^i$ , these form the obstruction triple of Definition 5.105.

On the cubic ideal  $\mathfrak{h}_{\leq N} = \mathfrak{m}^3/\mathfrak{m}^{N+1}$ , where the obstruction triple is silent, the finite cotangent CE/PV identity  $\Phi_{\leq N}: C_{\text{CE, cont}}^\bullet(\mathfrak{g}_{\leq N, I}) \xrightarrow{\sim} \text{PV}(\widehat{\mathbf{S}}(\mathfrak{h}_{\leq N, I}))$  is the Koszul resolution of  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b$  on the cubic truncation, unconditional in the classical sector (Theorem 5.101). Theorem 5.102 excludes the

all-order Moyal lift on the same finite truncation; Remark 5.107 shows that the inverse limit recovers the proper subalgebra  $\mathfrak{m}^3 \subsetneq \mathfrak{h}$ . Problem 5.109 is the open problem of extending the Koszul resolution  $\Phi_{\mathfrak{m}^3}$  from  $\mathfrak{m}^3$  to  $\mathfrak{h} = \mathfrak{m}$  by supplying a coefficient category in which  $\mathfrak{o}_{\leq 2}^{\text{Cas}}$  converges and a relative or reductive replacement for the lower-central Tate-pronilpotent route handling the semisimple  $H_2 \cong \mathfrak{sl}_2$  direction.

### The truncation

Let  $\mathfrak{h} = (z_1, z_2)\mathbb{C}[[z_1, z_2]] = \mathfrak{m}$  be the reduced Hamiltonian Lie algebra of the formal symplectic polydisk on  $\mathbb{C}^2$ , with bracket  $\{f, g\} = \partial_{z_1}f \partial_{z_2}g - \partial_{z_2}f \partial_{z_1}g$  modulo constants. Let  $\mathfrak{m}^k = (z_1, z_2)^k \mathbb{C}[[z_1, z_2]]$  for  $k \geq 1$  be the  $k$ -th power of the maximal ideal. Fix integers  $N \geq k \geq 3$  and define the *nilpotent truncation*

$$\mathfrak{h}_{\leq N}^{(k)} := \mathfrak{m}^k / \mathfrak{m}^{N+1},$$

with  $\mathfrak{m}$ -adic Poisson bracket inherited from the ambient  $\mathfrak{h}_{\text{poly}}$ . The standard choice is  $k = 3$ :

$$\mathfrak{h}_{\leq N} := \mathfrak{h}_{\leq N}^{(3)} = \mathfrak{m}^3 / \mathfrak{m}^{N+1}.$$

Concretely,  $\mathfrak{h}_{\leq N}$  is the finite-dimensional vector space spanned by monomials  $z_1^a z_2^b$  with  $3 \leq a + b \leq N$ .

The cutoff lies at  $k = 3$ . At  $k = 2$ , the degree-2 piece  $H_2 = \text{span}\{z_1^2, z_1 z_2, z_2^2\}$  closes under the Poisson bracket and carries the  $\mathfrak{sl}_2$ -relations

$$\{z_1^2, z_2^2\} = 4z_1 z_2, \quad \{z_1^2, z_1 z_2\} = 2z_1^2, \quad \{z_1 z_2, z_2^2\} = 2z_2^2,$$

so  $\mathfrak{m}^2 / \mathfrak{m}^{N+1}$  contains a nonabelian semisimple subalgebra and violates nilpotence, hence the lower-central Tate-pronilpotence hypothesis of 5.165. At  $k = 3$  the bracket raises  $\mathfrak{m}$ -adic degree strictly.

LEMMA 5.100 (*Truncation is a finite-dimensional nilpotent Lie algebra*). For every  $N \geq 3$ :

- (i)  $\mathfrak{h}_{\leq N} = \mathfrak{m}^3 / \mathfrak{m}^{N+1}$  is a finite-dimensional Lie algebra of dimension  $\binom{N+2}{2} - 6$  (the number of monomials of degree between 3 and  $N$  inclusive).
- (ii)  $\mathfrak{h}_{\leq N}$  is nilpotent of class at most  $N - 2$  in the lower central series.
- (iii)  $\mathfrak{h}_{\leq N}^\vee$  is finite-dimensional, hence the formal Casimir  $C_{\mathfrak{h}_{\leq N}} = \sum_{3 \leq d \leq N} \sum_i e_{d,i} \otimes e_d^i$  is a convergent element of  $\mathfrak{h}_{\leq N} \otimes \mathfrak{h}_{\leq N}^\vee$  as a finite sum in the ordinary tensor product.
- (iv) The linear Hamiltonians  $z_1, z_2 \in \mathfrak{m} \setminus \mathfrak{m}^3$ , the conformal-symplectic generators  $H_2 = \text{span}\{z_1^2, z_1 z_2, z_2^2\} \cong \mathfrak{sl}_2$ , and the constant Hamiltonian  $1 \notin \mathfrak{m}$  are omitted from  $\mathfrak{h}_{\leq N}$  by construction.

*Proof.* (i) The space  $\mathfrak{m}^3$  is a Lie subalgebra of the polynomial Hamiltonian Lie algebra  $\mathfrak{h}_{\text{poly}}$ , and  $\mathfrak{m}^{N+1}$  is a Lie ideal inside  $\mathfrak{m}^3$ : for  $f \in \mathfrak{m}^p$  and  $g \in \mathfrak{m}^q$  one has  $\{f, g\} \in \mathfrak{m}^{p+q-2}$ , since the Poisson bracket on  $\mathbb{C}[z_1, z_2]$  lowers total polynomial degree by exactly two, and each partial derivative lowers  $\mathfrak{m}$ -adic degree by exactly one. For  $p, q \geq 3$  the bracket lands in  $\mathfrak{m}^{p+q-2} \subset \mathfrak{m}^4 \subset \mathfrak{m}^3$ , making  $\mathfrak{m}^3$  a Lie subalgebra; for  $p \geq 3$  and  $q \geq N + 1$  the bracket lands in  $\mathfrak{m}^{p+q-2} \subset \mathfrak{m}^{N+2} \subset \mathfrak{m}^{N+1}$ , making  $\mathfrak{m}^{N+1}$  a Lie ideal of  $\mathfrak{m}^3$ . The quotient  $\mathfrak{h}_{\leq N} = \mathfrak{m}^3 / \mathfrak{m}^{N+1}$  inherits a well-defined Lie bracket. Dimension count: monomials  $z_1^a z_2^b$  with  $3 \leq a + b \leq N$  number  $\binom{N+2}{2} - 6$  (the total number with  $0 \leq a + b \leq N$  minus the six of degrees 0, 1, 2).

(ii) The Poisson bracket of two elements of  $\mathfrak{m}^3$  raises  $\mathfrak{m}$ -adic degree by at least 1:  $\{H_p, H_q\} \subset H_{p+q-2} \subset \mathfrak{m}^{p+q-2}$ , and for  $p, q \geq 3$ ,  $p + q - 2 \geq 4 > p$  (since  $q \geq 3$ ). Thus the descending Lie series

$$\mathfrak{h}_{\leq N}^{(\geq m)} := \underbrace{[\mathfrak{h}_{\leq N}, [\mathfrak{h}_{\leq N}, \dots]]}_{m \text{ brackets}} \subset (\mathfrak{m}^{m+3} / \mathfrak{m}^{N+1})$$

satisfies  $\mathfrak{h}_{\leq N}^{(\geq m)} = 0$  for  $m + 3 \geq N + 1$ , i.e. for  $m \geq N - 2$ . Hence  $\mathfrak{h}_{\leq N}$  is nilpotent of class at most  $N - 2$ .

(iii) Immediate from finite-dimensionality.

(iv) Immediate from the  $\mathfrak{m}^3$ -adic cutoff and the exclusion  $a + b \leq 2$ .  $\square$

The truncation kills the linear and conformal-symplectic sectors: the obstruction  $\{z_1, z_2^{N+1}\} = (N + 1)z_2^N \notin \mathfrak{m}^{N+1}$  of 5.209 requires  $z_1 \in H_1$ , which the cubic cutoff omits. The Casimir is the finite sum  $C_{\mathfrak{h}_{\leq N}} = \sum_{d=3}^N \sum_i e_{d,i} \otimes e_d^i \in \mathfrak{h}_{\leq N} \otimes \mathfrak{h}_{\leq N}^\vee$ ; the product/direct-sum support obstruction of 5.211, which depends on infinite degree expansion, vanishes.

### *The classical truncated cotangent CE/PV identification*

**THEOREM 5.101** (*Classical truncated CE/PV identification: Koszul resolution on the cubic ideal*). Set  $\mathfrak{g}_{\leq N} = \mathfrak{h}_{\leq N} \ltimes \mathfrak{h}_{\leq N}^\vee[1]$ . For every interval  $I \subset \mathbb{R}$  let  $\mathfrak{g}_{\leq N, I} = \Omega_c^\bullet(I) \widehat{\otimes} \mathfrak{h}_{\leq N} \ltimes \mathcal{D}'_c(I) \widehat{\otimes} \mathfrak{h}_{\leq N}^\vee[1]$ . Then, after the Poisson, residue, and interval-smearing conventions fixed above, the dictionary  $c^I \mapsto \theta^I$ ,  $u_I \mapsto O_I$  is the Koszul resolution of the chiral Hochschild cohomology  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b$  on the cubic truncation: a filtered cochain isomorphism of completed dg graded-commutative  $P_0$ -algebras

$$\Phi_{\leq N} : C_{\text{CE,cont}}^\bullet(\mathfrak{g}_{\leq N, I}) \xrightarrow{\sim} \text{PV}(\widehat{\mathbf{S}}(\mathfrak{h}_{\leq N, I}))$$

on every interval  $I \subset \mathbb{R}$ , with the same Schouten/Lichnerowicz convention as in 4.20. It is compatible with extension by zero and with disjoint-interval products in this reduced finite model, and on the Hamiltonian generators it recovers the stalkwise operation  $f \mapsto L_{B_f}$ . The cotangent generators are finite principal-part labels; their unreduced boundary centrality homotopies remain part of Problem 5.48.

*Proof.* The Lie algebra  $\mathfrak{g}_{\leq N}$  is finite-dimensional ( $\dim \mathfrak{g}_{\leq N} = 2 \dim \mathfrak{h}_{\leq N} = 2 \binom{N+2}{2} - 12$ ), hence pro-finite-dimensional in the trivial constant filtration. It is lower-central Tate-pronilpotent in the sense of 5.165: by 5.100(ii) the descending Lie powers of  $\mathfrak{h}_{\leq N}$  vanish after at most  $N - 2$  brackets. The cotangent summand is abelian, and nilpotence of the semidirect product uses the degree-lowering coadjoint action: since  $[H_p, H_q] \subset H_{p+q-2}$ , an element  $f \in H_p$ ,  $p \geq 3$ , sends  $H_d^\vee$  to  $H_{d-p+2}^\vee$ , hence lowers dual degree by at least one. Repeated action therefore kills every finite quotient  $\mathfrak{h}_{\leq N}^\vee$ . Together with nilpotence of  $\mathfrak{h}_{\leq N}$ , the descending Lie powers of  $\mathfrak{g}_{\leq N} = \mathfrak{h}_{\leq N} \ltimes \mathfrak{h}_{\leq N}^\vee[1]$  also terminate. Hence  $\bigcap_m \mathfrak{g}_{\leq N}^{(\geq m)} = 0$  in any Tate filtration, and the lower-central Tate-pronilpotence hypothesis of 5.165 is satisfied trivially.

Apply 4.20 (the cochain-level reduced Hamiltonian CE/PV central-operation theorem) to  $\mathfrak{h}_{\leq N}$ . The conclusion is a canonical isomorphism of completed dg graded-commutative  $P_0$ -algebras

$$\Phi_{\leq N}^{\text{stalk}} : C_{\text{CE,cont}}^\bullet(\mathfrak{g}_{\leq N}) \xrightarrow{\sim} \text{PV}(\widehat{\mathbf{S}}(\mathfrak{h}_{\leq N})),$$

with the explicit generator-level data of 4.20.

To pass from the stalk identification to the factorization-algebra identification on intervals  $I \subset \mathbb{R}$ , the Hamiltonian-sector factorization upgrade of 4.11 applies verbatim with  $\mathfrak{h}$  replaced by  $\mathfrak{h}_{\leq N}$ : the locality input  $\{(\phi_i)^i_j(t), (\phi_2)^k_l(t')\} = \partial_i^j \partial_j^k \delta(t - t')$  on two brane-time points is unchanged, the smearing  $\mathfrak{h}_{\leq N, I} = \Omega_c^\bullet(I) \widehat{\otimes} \mathfrak{h}_{\leq N}$  is canonical because  $\mathfrak{h}_{\leq N}$  is finite-dimensional in the ordinary tensor category, and the strict  $P_0$  bracket follows from the delta-function locality.

In this finite truncated classical sector, the cotangent factor has a finite Tate-Casimir kernel. The principal-part sub-factorization algebra

$$\widetilde{\text{Obs}}_{\partial, \leq N}^{\text{cl, pp}}(I) = \text{Obs}_{\text{open}, \leq N}^{\text{cl}}(I) \widehat{\otimes} \widehat{\mathbf{S}}(\mathfrak{h}_{\leq N, I}^\vee[1])$$

admits a chain-level primitive projection  $\pi_{\text{prim}}^{\leq N}$  because the Tate Casimir  $C_{\mathfrak{h}_{\leq N}} = \sum_{d=3}^N \sum_i e_{d,i} \otimes e_d^i$  is a finite sum, hence a literal element of the (uncompleted) tensor product  $\mathfrak{h}_{\leq N} \otimes \mathfrak{h}_{\leq N}^{\vee}$ . The  $Q$ -compatibility, factorization compatibility, and  $P_o$ -bracket homotopies are then formal, since every step uses finite-support tensors in the direct-sum dual.

The bracket compatibility and graded Leibniz extension to the completed finite CE/PV algebra is the same argument as in 4.20, with the additional simplification that all completions are trivial because  $\mathfrak{h}_{\leq N}$  is finite-dimensional. Compatibility with extension by zero, disjoint-interval factorization, and the stalkwise limit follow as in 4.11.  $\square$

### *Quantum obstruction for the finite truncation*

**THEOREM 5.102** (*Quantum obstruction for the nilpotent truncation*). The classical finite truncation 5.101 is unconditional. The same finite Lie algebra  $\mathfrak{h}_{\leq N} = \mathfrak{m}^3/\mathfrak{m}^{N+1}$  has an obstruction to an all-order Moyal deformation for  $N \geq 4$ : the reduced Moyal bracket has a closure obstruction. Consequently an all-order quantum descendant theorem on the strict finite truncation requires an additional Moyal-flat replacement or a projected bracket with Jacobi and cochain compatibility.

What remains true at quantum level is the restriction of the full-sector degree-zero map 5.216 to the finite set of generators of  $\mathfrak{h}_{\leq N}$ , viewed inside the full  $\mathfrak{h}_{\hbar}$ . This is a restriction of a proved map on the full Hamiltonian source. A closed CE theorem for a finite quantum Lie subalgebra requires the additional data above.

*Proof.* The Moyal bracket is

$$\{f, g\}_{\hbar} = \sum_{j \geq 0} \frac{\hbar^{2j}}{(2j+1)! 2^{2j}} P^{2j+1}(f, g), \quad P = \partial_{z_1} \otimes \partial_{z_2} - \partial_{z_2} \otimes \partial_{z_1}.$$

If  $f \in \mathfrak{m}^p$  and  $g \in \mathfrak{m}^q$ , the order- $\hbar^{2j}$  summand lies in  $\mathfrak{m}^{p+q-2(2j+1)}$  when nonzero. The Poisson term  $j = 0$  preserves  $\mathfrak{m}^3$  as a Lie subalgebra, but the higher Moyal terms can leave the ideal. For  $N \geq 4$ ,

$$P^3(z_1^4, z_2^4) = 24z_1 \cdot 24z_2 = 576z_1z_2,$$

so

$$\{z_1^4, z_2^4\}_{\hbar} = 16z_1^3z_2^3 + 24\hbar^2 z_1z_2 + \dots$$

The term  $24\hbar^2 z_1z_2$  has  $\mathfrak{m}$ -adic degree 2. It has degree 2 and lies outside both the cubic ideal  $\mathfrak{m}^3$  and the constants killed by the reduction modulo  $\mathbb{C} \cdot 1$ . Thus  $\mathfrak{m}^3/\mathfrak{m}^{N+1}$  has a Moyal-closure obstruction for  $N \geq 4$ .

Therefore the CE differential for a hypothetical finite quantum source  $\mathfrak{h}_{\leq N, \hbar} \ltimes \mathfrak{h}_{\leq N, \hbar}^{\vee}[1]$  is incompatible with simple restriction of the Moyal bracket. Any projected bracket would require a new Jacobi proof; the projected-degree warning of 5.209 shows that this is a genuine Jacobi condition. The all-order theorem 5.216 remains valid on the full Hamiltonian algebra  $\mathfrak{h}_{\hbar}$ , and its map can be evaluated on generators of degrees 3,  $\dots$ ,  $N$ . That operation yields restricted values; a finite quantum CE source requires the additional Moyal-flat structure above. The descendant cotangent sector at  $\hbar = 0$  is exactly 5.101; at  $\hbar \neq 0$  it is the named Moyal-flat truncation problem.  $\square$

**PROPOSITION 5.103** (*Sectors removed by truncation*). The truncation  $\mathfrak{h} \rightsquigarrow \mathfrak{h}_{\leq N} = \mathfrak{m}^3/\mathfrak{m}^{N+1}$  removes the following five data from the local theorem 5.6:

- (i) The linear Hamiltonians  $z_1, z_2 \in \mathfrak{m} \setminus \mathfrak{m}^2$ , which generate translations in the symplectic plane and the center-of-mass mode of the brane.
- (ii) The conformal-symplectic  $\mathfrak{sl}_2$  generators  $H_2 = \text{span}\{z_1^2, z_1 z_2, z_2^2\}$ , which generate rotations and dilations of the symplectic plane. This is the semisimple sector that prevents  $\mathfrak{m}^3/\mathfrak{m}^{N+1}$  from being nilpotent.
- (iii) The constant Hamiltonian  $\mathbf{1} \notin \mathfrak{m}$ , on which the  $\hbar$ -normalization of the Moyal product depends. The central element  $\mathbb{C} \cdot \mathbf{1} \subset \mathfrak{h}_{\text{poly}}$  is omitted from  $\mathfrak{m}^3$  a fortiori, so the  $\hbar$ -normalization in the truncated theorem is conventional.
- (iv) The  $U(1)$  center anomaly cohomology class  $[\bar{c}] \in H_{\text{Lie}}^2(\bar{A}, \mathbb{C})$  of 5.6(f), since the cocycle

$$\omega(z_1^k z_2^l, z_1^m z_2^n) = (kn - lm) \mathbf{1}[(k + m, l + n) = (\mathbf{1}, \mathbf{1})]$$

is concentrated on multidegree  $(\mathbf{1}, \mathbf{1})$ , with both arguments of total  $\mathfrak{m}$ -adic degree  $\mathbf{1}$ , hence the cocycle pairs only elements outside  $\mathfrak{h}_{\leq N} = \mathfrak{m}^3/\mathfrak{m}^{N+1}$ . The central anomaly has support outside the truncation.

- (v) The stable connected limit  $N \rightarrow \infty$  on the full  $\mathfrak{h}$ . The inverse system  $\{\mathfrak{h}_{\leq N}\}_{N \geq 3}$  has structure maps the natural surjections  $\mathfrak{h}_{\leq N+1} \twoheadrightarrow \mathfrak{h}_{\leq N}$  induced by the  $\mathfrak{m}$ -adic projection  $\mathfrak{m}^{N+1} \subset \mathfrak{m}^{N+2}$ . The inverse limit is  $\varprojlim_N \mathfrak{h}_{\leq N} = \mathfrak{m}^3$ , the  $\mathfrak{m}^3$ -truncation of the Hamiltonian Lie algebra.

What remains classically: the third-order and higher canonical transformations, which form a nilpotent Lie subalgebra  $\mathfrak{m}^3 \subset \mathfrak{h}$ , a Lie ideal in  $\mathfrak{m}^2$ , admitting the classical finite cotangent CE/PV identification unconditionally. The unreduced open BV factorization-centre morphism requires additional data beyond this truncation. The all-order Moyal descendant lift is excluded by 5.102 unless extra quantum truncation data are added.

*Proof.* (i)–(iv) are immediate from the definitions and from the  $\mathfrak{m}$ -adic concentration of  $\omega$ . (v) is the explicit inverse limit calculation: a compatible family  $\{f_N \in \mathfrak{m}^3/\mathfrak{m}^{N+1}\}_N$  assembles to a unique formal series in  $\mathfrak{m}^3 \subset \mathbb{C}[[z_1, z_2]]$ , and the  $\mathfrak{m}$ -adic Lie structure is preserved at every level.  $\square$

#### *Cubic inverse-limit survivor*

**PROPOSITION 5.104** (*Cubic coordinate-window survivor and variance boundary*). The finite Lie quotients

$$\mathfrak{h}_{\leq N} = \mathfrak{m}^3/\mathfrak{m}^{N+1}$$

recover the complete cubic Lie algebra  $\varprojlim_N \mathfrak{h}_{\leq N} = \mathfrak{m}^3$ . The finite cotangent maps  $\Phi_{\leq N}$  of 5.101 form a compatible family of coordinate tests. Their functoriality is contravariant: it is pullback along Lie quotients, with additional data required for an inverse system of dg CE cochain projections. A Lie quotient  $\mathfrak{h}_{\leq M} \twoheadrightarrow \mathfrak{h}_{\leq N}$  induces CE cochain maps by pullback in the opposite direction. Consequently  $\Phi_{\mathfrak{m}^3}$  denotes the coordinate-window family  $\{\Phi_{\leq N}\}_N$  together with the chain-side inverse limit and an exact continuous-dualization datum; the expression  $\varprojlim_N \Phi_{\leq N}$  on CE cochains requires those additional data.

This is the maximal survivor of the nilpotent tower only in that precise sense: the Lie-side tower recovers  $\mathfrak{m}^3$ , while a completed cochain  $P_0$ -statement requires the additional variance and bracket-admissibility data named above.

*Proof.* The Lie-side inverse limit is the usual  $\mathfrak{m}$ -adic completion of  $\mathfrak{m}^3$ . The variance obstruction appears already at  $\mathcal{M} = 4$ ,  $N = 3$ . In  $\mathfrak{h}_{\leq 4}$ ,

$$\{z_1^3, z_2^3\} = 9z_1^2 z_2^2, \quad d_{\text{CE}} c^{2,2} \supset -9c^{3,0} c^{0,3}.$$

The coordinate projection to the  $N = 3$  cochain window kills  $c^{2,2}$  and leaves  $c^{3,0} c^{0,3}$ , so

$$p(d_{\text{CE}} c^{2,2}) = -9c^{3,0} c^{0,3} \neq 0 = d_{\text{CE}} p(c^{2,2}).$$

Thus projection of CE cochains has a nonzero dg defect. The correct finite functoriality is contravariant pullback along Lie quotients, while a completed cochain theorem must be obtained either from a chain-side inverse limit followed by exact continuous dualization or from a separately supplied topology whose cochain transition maps are dg maps.

The Lie algebra  $\mathfrak{m}^3$  is lower-central Tate-pronilpotent in the  $\mathfrak{m}$ -adic Tate filtration: its descending Lie powers satisfy  $(\mathfrak{m}^3)^{(\geq m)} \subset \mathfrak{m}^{m+3}$  because each iterated bracket raises the  $\mathfrak{m}$ -adic degree by at least 1 when both arguments lie in  $\mathfrak{m}^3$  (since  $\{H_p, H_q\} \subset \mathfrak{m}^{p+q-2}$  and  $p, q \geq 3$  implies  $p+q-2 \geq p+1$ ). Hence  $\bigcap_m (\mathfrak{m}^3)^{(\geq m)} = 0$ , and the lower-central Tate-pronilpotence hypothesis of 5.165 holds. The Tate Casimir  $C_{\mathfrak{m}^3} = \sum_{d \geq 3} \sum_i e_{d,i} \otimes e_d^i$  is governed by the same product/direct-sum convergence question as in 5.211, but on the smaller direct-sum dual  $\bigoplus_{d \geq 3} H_d^\vee \subset \bigoplus_{d \geq 1} H_d^\vee$ . The finite maps  $\Phi_{\leq N}$  at every finite  $N$  use only the finite Casimir  $\sum_{3 \leq d \leq N} \sum_i e_{d,i} \otimes e_d^i$ . Passing from these finite identities to a completed cochain identity is exactly the continuous-dualization and variance datum isolated in the statement, rather than a formal inverse-limit consequence.  $\square$

*Definition 5.105 (Low-degree obstruction datum for the nilpotent tower).* For  $N \geq 3$  let

$$\pi_N : \mathfrak{m}^3 \rightarrow \mathfrak{h}_{\leq N} = \mathfrak{m}^3 / \mathfrak{m}^{N+1}$$

be the nilpotent window projection. The two obstruction terms for adjoining the removed low-degree sector are:

$$\mathfrak{o}_N^{\text{lin}}(x; \tau) = \{x, \tau\} \bmod \mathfrak{m}^{N+1}, \quad x \in H_1, \tau \in \mathfrak{m}^{N+1} / \mathfrak{m}^{N+2},$$

and

$$\mathfrak{o}^{\text{ss}} : \Lambda^2 H_2 \rightarrow H_2, \quad \mathfrak{o}^{\text{ss}}(x, y) = \{x, y\}.$$

Concretely

$$\mathfrak{o}_N^{\text{lin}}(z_1; z_2^{N+1}) = (N+1)z_2^N \neq 0 \quad \text{in } \mathfrak{m}^N / \mathfrak{m}^{N+1},$$

and

$$\{z_1^2, z_2^2\} = 4z_1 z_2, \quad \{z_1^2, z_1 z_2\} = 2z_1^2, \quad \{z_1 z_2, z_2^2\} = 2z_2^2.$$

The compatible Casimir missing from the nilpotent tower is

$$\mathfrak{o}_{\leq 2}^{\text{Cas}} = \sum_i e_{1,i} \otimes e_1^i + \sum_i e_{2,i} \otimes e_2^i.$$

**LEMMA 5.106 (Obstruction-datum criterion for enlarging the tower).** A finite  $\mathfrak{m}$ -adic tower extending  $\{\mathfrak{h}_{\leq N}\}_{N \geq 3}$  to a Lie tower containing  $H_1 \oplus H_2$  and using the same quotient maps exists only if  $\mathfrak{o}_N^{\text{lin}} = 0$  for all  $N$  and  $\mathfrak{o}^{\text{ss}}$  is nilpotent in the lower-central filtration. A cotangent CE/PV extension of the inverse limit to the full  $\mathfrak{h} = \mathfrak{m}$  also requires a coefficient category in which the compatible tensor  $\mathfrak{o}_{\leq 2}^{\text{Cas}} + \sum_{d \geq 3} \sum_i e_{d,i} \otimes e_d^i$  converges.

*Proof.* If  $H_1$  is included and the window  $N$  is still the quotient by  $\mathfrak{m}^{N+1}$ , then the tail  $\mathfrak{m}^{N+1}$  must be a Lie ideal for the enlarged source. The displayed value  $\{z_1, z_2^{N+1}\} = (N+1)z_2^N \notin \mathfrak{m}^{N+1}$  shows that this ideal condition is exactly the vanishing of  $\mathfrak{o}_N^{\text{lin}}$ , and the displayed value is nonzero.

If  $H_2$  is included, then its bracket remains inside  $H_2$  and equals the nonabelian  $\mathfrak{sl}_2$  bracket displayed in Definition 5.105. The image of  $\mathfrak{o}^{\text{ss}}$  is all of  $H_2$ , so the descending Lie powers never leave  $H_2$ . This is exactly the obstruction to nilpotence and to the conilpotent bar-cobar hypothesis.

The finite cotangent windows for  $\mathfrak{h}_{\leq N}$  use the finite Casimir  $\sum_{3 \leq d \leq N} \sum_i e_{d,i} \otimes e_d^i$ . Extending to  $\mathfrak{h}$  adds the missing  $d = 1, 2$  components and then the whole compatible family over all  $d \geq 1$ . Thus the full cotangent lift requires precisely the convergence of the displayed full Casimir family.  $\square$

**THEOREM 5.107** (*Low-degree obstruction datum for full Hamiltonian recovery*). The coordinate-window family  $\Phi_{\mathfrak{m}^3}$  of 5.104 is the classical cotangent lift datum on  $\mathfrak{m}^3$ . Extension to all of  $\mathfrak{h} = \mathfrak{m}$  must include the low-degree sectors  $\mathfrak{m}/\mathfrak{m}^3 = H_1 \oplus H_2 = \mathbb{C} \cdot z_1 \oplus \mathbb{C} \cdot z_2 \oplus \text{span}\{z_1^2, z_1 z_2, z_2^2\}$ , which by 5.209(iii) and 5.106 has a stability obstruction under any  $\mathfrak{m}$ -adic cutoff: the bracket  $\{z_1, z_2^{N+1}\} = (N+1)z_2^N$  is the explicit obstruction. Equivalently, the obstruction datum is

$$(\mathfrak{o}_N^{\text{lin}}, \mathfrak{o}^{\text{ss}}, C_{\mathfrak{h}}),$$

where

$$\mathfrak{o}_N^{\text{lin}}(z_1; z_2^{N+1}) = (N+1)z_2^N \neq 0, \quad \text{im } \mathfrak{o}^{\text{ss}} = H_2 \cong \mathfrak{sl}_2,$$

and  $C_{\mathfrak{h}}$  is the full infinite Casimir family below. Equivalently, the Tate Casimir  $C_{\mathfrak{h}} = \sum_{d \geq 1} \sum_i e_{d,i} \otimes e_d^i$  has low-degree components  $\sum_i e_{1,i} \otimes e_1^i \in H_1 \otimes H_1^\vee$  and  $\sum_i e_{2,i} \otimes e_2^i \in H_2 \otimes H_2^\vee$  that the truncation removes; the degree-1 component is the source of the linear-Hamiltonian translation, and the degree-2 component is the source of the conformal-symplectic  $\mathfrak{sl}_2$  direction, where semisimplicity replaces nilpotence. Recovering the full  $\mathfrak{h}$  requires a coefficient category in which  $C_{\mathfrak{h}}$  converges *including* both low-degree components, which is exactly the Casimir support obstruction of 5.211. The truncation route leaves that obstruction intact; it removes the sector supporting the obstruction. A full recovery theorem must therefore supply both a coefficient category in which  $C_{\mathfrak{h}}$  converges and a relative or reductive model for the  $H_2$ -sector.

*Proof.* The tower used in 5.104 is  $\mathfrak{h}_{\leq N} = \mathfrak{m}^3/\mathfrak{m}^{N+1}$ . Since the transition maps are the usual quotient maps,

$$\varprojlim_N \mathfrak{h}_{\leq N} = \varprojlim_N \mathfrak{m}^3/\mathfrak{m}^{N+1} = \mathfrak{m}^3$$

as a complete filtered Lie algebra. The homogeneous pieces  $H_1$  and  $H_2$  have zero image in every term of this tower; hence the inverse limit misses them.

The obstruction is intrinsic to the chosen finite windows. Including  $H_1$  while keeping the same  $\mathfrak{m}$ -adic finite windows gives a Lie-ideal obstruction in the tails: for every  $N$ ,

$$z_2^{N+1} \in \mathfrak{m}^{N+1}, \quad \{z_1, z_2^{N+1}\} = (N+1)z_2^N \notin \mathfrak{m}^{N+1}.$$

Therefore the quotient by the tail carries a full-Hamiltonian bracket obstruction. If one includes  $H_2$ , the degree-two Hamiltonians contain the  $\mathfrak{sl}_2$  bracket  $[H_2, H_2] = H_2$ , so the finite quotient violates nilpotence. Thus a finite nilpotent  $\mathfrak{m}$ -adic tower is incompatible with the low-degree sector  $H_1 \oplus H_2$ .

Finally, the full Casimir  $C_{\mathfrak{h}}$  has nonzero components in  $H_1 \otimes H_1^\vee$ ,  $H_2 \otimes H_2^\vee$ , and in every higher  $H_d \otimes H_d^\vee$ . The nilpotent tower sees only the  $d \geq 3$  part. Hence the inverse-limit cotangent lift on  $\mathfrak{m}^3$  promotes to the full Hamiltonian sector only after an additional coefficient category makes the full Casimir converge.  $\square$

COROLLARY 5.108 (*Full maximal-ideal universal obstruction class*). For the full reduced formal Hamiltonian algebra  $\mathfrak{h} = \mathfrak{m} = (z_1, z_2)\mathbb{C}[[z_1, z_2]]$ , the direct formal-local CE/PV recognition theorem remains valid in the finite-coordinate product completion. The universal filtered-cobar Koszul recognition criterion of Theorem 5.23, however, has a nonzero obstruction class in  $\mathcal{O}_{\text{Kos}}$  of Definition 5.22. Its visible components are

$$\mathfrak{o}_N^{\text{lin}}(z_1; z_2^{N+1}) = (N+1)z_2^N \neq 0, \quad \text{im } \mathfrak{o}^{\text{ss}} = H_2 \cong \mathfrak{sl}_2,$$

$$[C_{\mathfrak{h}}] = \left[ \sum_{d \geq 1} \sum_i e_{d,i} \otimes e_d^i \right] \in Q_{\text{Cas}}, \quad H_2 \subset \bigcap_{r \geq 1} \mathfrak{h}^{(r)}.$$

Therefore a universal full- $\mathfrak{h}$  bar-cobar/enveloping theorem requires, before any open-side  $\mathcal{A}_\infty$  comparison, all of the following: a coefficient topology representing  $C_{\mathfrak{h}}$ , continuity of the finite-window bracket and coadjoint transition maps, vanishing  $\varprojlim^1$  for the completed CE/PV and cobar towers, a relative or reductive treatment of the  $H_2$ -sector, and a twisting cochain whose associated-graded cobar map has acyclic cone.

*Proof.* The formal-local CE/PV part is Theorem 4.20, equivalently the formal-disk instance of Theorem 4.22: the bracket and coadjoint formulas are continuous for finite coordinate projections, so the generator substitution  $c \mapsto \theta$ ,  $u \mapsto O$  is a completed dg commutative cochain isomorphism. The  $P_0$ -algebra reading is finite-level or bracket-admissible data; an unqualified inverse limit of ambient coordinate products supplies a different object.

The filtered-cobar recognition theorem asks for more. The linear obstruction is the same computation as in Definition 5.105: adjoining  $H_1$  to the  $\mathfrak{m}$ -adic nilpotent windows makes the tail non-invariant because  $\{z_1, z_2^{N+1}\} = (N+1)z_2^N \notin \mathfrak{m}^{N+1}$ . The semisimple obstruction is the displayed  $\mathfrak{sl}_2$  bracket on  $H_2$ , hence  $H_2 \subset \bigcap_r \mathfrak{h}^{(r)}$ . This gives a nonzero lower-central component in  $\mathcal{O}_{\text{conilp}}$ .

The Casimir component is nonzero in the strict product/direct-sum topology by the support argument of Lemma 5.211: the compatible finite-window Casimirs have nonzero identity component in every  $H_d \otimes H_d^\vee$ , while a strict tensor with a direct-sum dual has only finitely many dual-degree components. Finally, even after changing topology and handling  $H_2$  relatively, the universal criterion still demands the cobar cone and the Milnor  $\varprojlim^1$  terms in Definition 5.22 to vanish. These are precisely the listed remaining requirements.  $\square$

Problem 5.109 (*Open problem: extension of the Koszul resolution from  $\mathfrak{m}^3$  to  $\mathfrak{h}$* ). Extend the coordinate-window Koszul-resolution datum  $\Phi_{\mathfrak{m}^3}$  of 5.104 from  $\mathfrak{m}^3$  to all of  $\mathfrak{h} = \mathfrak{m}$  by supplying: (D1) a Tate or weighted-coefficient extension in which the degree-1 and degree-2 Casimir components converge as kernels for the BV propagator; (D2) a relative or reductive replacement for the lower-central Tate-pronilpotence hypothesis of 5.165 in which the conformal-symplectic  $\mathfrak{sl}_2$  direction  $H_2$  is included as semisimple data. By 5.105 the obstruction terms are the triple  $(\mathfrak{o}_N^{\text{lin}}, \mathfrak{o}^{\text{ss}}, \mathfrak{o}_{\leq 2}^{\text{Cas}})$ ; by 5.211 (D1) is the same coefficient-category problem as 5.48. The nilpotent-truncation route closes the Koszul resolution of  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_{\mathfrak{b}}$  on the cubic ideal  $\mathfrak{m}^3$ ; this open problem is the precise obstruction to the full  $\mathfrak{h}$  Koszul resolution, and the residual moves from 5.48 as stated to its restricted form on the low-degree sector  $\mathfrak{m}/\mathfrak{m}^3 = H_1 \oplus H_2$  alone.

THEOREM 5.110 (*Semisimple obstruction to lower-central Tate-pronilpotence*). A second obstruction to extending  $\Phi_{\mathfrak{m}^3}$  to all of  $\mathfrak{h}$  is structural: the inclusion of  $H_2 = \mathfrak{sl}_2$  violates the lower-central Tate-pronilpotence route used in Lemma 5.165 because  $[H_2, H_2] = H_2$  is its own descending Lie series. Even



if the degree-1 and degree-2 Casimir components converged in some weighted Tate completion, direct application of the pronilpotent bar-cobar route would still be obstructed. Ordinary finite-dimensional bar-cobar and PBW formalism remain valid for  $\mathfrak{sl}_2$ ; the lower-central Tate-pronilpotent criterion must be replaced, on a  $L$  containing  $H_2$ , by a relative or reductive model that builds in semisimple subalgebras separately.

*Proof.* Let  $L$  be any filtered Lie algebra containing the degree-two Hamiltonians  $H_2$  as a Lie subalgebra with the usual Poisson bracket. The descending Lie powers of  $L$  satisfy  $L^{(n)} \cap H_2 \supset H_2$  for every  $n$ , because  $[H_2, H_2] = H_2$ . Hence

$$H_2 \subset \bigcap_{n \geq 1} L^{(n)}.$$

The intersection of the descending Lie powers is therefore nonzero. This contradicts the lower-central Tate-pronilpotence condition used in Lemma 5.165. The bar-cobar argument applies to the nilpotent radical  $\mathfrak{m}^3$ . A Lie source containing the semisimple  $H_2$ -sector requires a relative or reductive replacement.  $\square$

### Costello–Li precedent

*Remark 5.III (Costello–Li balanced matrix truncation as structural model).* Costello and Li [13] prove the quantum perturbative BCOV theorem on flat holomorphic-volume space by an  $N \rightarrow \infty$  stable connected limit through the open gauge algebra  $\mathfrak{gl}(N | N)$ , with the directed Lie embedding  $\mathfrak{gl}(N | N) \hookrightarrow \mathfrak{gl}(N + k | N + k)$  and the BV-cohomology vanishing  $H_{\text{BV}}^*(\mathfrak{gl}(N | N)) = 0$  (supertrace identity  $\text{str}(I) = N - N = 0$  kills the would-be  $U(1)$ -anomaly). The structural analogy with the truncation here:

- (i) Each window  $\mathfrak{gl}(N | N)$  is finite-dimensional, BV-cohomology vanishes, the universal CE/Schouten identification and the BV-propagator coefficient kernel are well-defined on each window.
- (ii) The inverse / direct system is the Lie embedding  $\mathfrak{gl}(N | N) \hookrightarrow \mathfrak{gl}(N + 1 | N + 1)$  on the Costello–Li side, the Lie surjection  $\mathfrak{h}_{\leq N+1} \rightarrow \mathfrak{h}_{\leq N}$  here.
- (iii) The  $U(1)$ -anomaly is killed by the structure of each window on the Costello–Li side (supertrace), and is removed by the  $\mathfrak{m}^3$ -truncation here (the cocycle  $\omega$  pairs only multidegree-(1, 0) with multidegree-(0, 1), both in  $H_1 \subset \mathfrak{m} \setminus \mathfrak{m}^2$ , hence outside  $\mathfrak{m}^3$ ).
- (iv) The colimit  $N \rightarrow \infty$  converges on the Costello–Li side to the full  $\mathfrak{gl}(\infty | \infty)$ , providing the stable connected single-trace algebra. Here the inverse limit converges to the proper subalgebra  $\mathfrak{m}^3 \subsetneq \mathfrak{h}$  by 5.107.

The structural difference: the  $\mathfrak{gl}(N | N)$  system is graded by matrix size, with trivial action of one window on the next-window orthogonal complement; the  $\mathfrak{h} = \mathfrak{m}$  system is graded by  $\mathfrak{m}$ -adic degree, with the linear and quadratic sectors acting non-trivially on every higher-degree window, which is precisely the finite-window obstruction of 5.209(iii) plus the  $\mathfrak{sl}_2 \subset H_2$  semisimple obstruction to lower-central Tate-pronilpotence.

*Tate Quillen equivalence*  $\mathfrak{o}_{\text{Quil}}$ . The cochain CE/PV identification of 4.20 is upgraded to a Tate Quillen equivalence inside the admissible model envelope  $\mathbf{TateChain}_{\geq 0}^{\text{adm}}$  of Statement 5.17. The component  $\mathfrak{o}_{\text{Quil}}$  is the universal Koszul obstruction class in the universal Koszul obstruction complex  $\mathcal{O}_{\text{Kos}}$  of Definition 5.22, recording the conilpotent-coalgebra hypothesis on the bar source, the bounded filtration,

the strong-convergence test, and the cone of the filtered cobar map. By 5.108 the visible components on the full Hamiltonian source  $\mathfrak{h} = \mathfrak{m}$  are

$$\mathfrak{o}_N^{\text{lin}}(z_1; z_2^{N+1}) \neq 0, \quad \text{im } \mathfrak{o}^{\text{ss}} = H_2, \quad [C_{\mathfrak{h}}] \in Q_{\text{Cas}}, \quad H_2 \subset \bigcap_{r \geq 1} \mathfrak{h}^{(r)},$$

together with the Milnor  $\varprojlim^1$  terms required by the filtered cobar acceptance criterion of 4.25. The construction problem is the joint vanishing inside  $\mathcal{O}_{\text{Kos}}$ : supply a coefficient topology representing  $C_{\mathfrak{h}}$ , continuity of finite-window bracket and coadjoint transition maps, vanishing  $\varprojlim^1$  for the completed CE/PV and cobar towers, a relative/reductive treatment of the  $H_2$ -sector, and a twisting cochain whose associated-graded cobar map has acyclic cone. Inside the weighted regulator-admissible sector and after the substitution of the finite-window scalar projection of 5.83, the Tate Quillen equivalence becomes an admissible filtered quasi-isomorphism in the sense of 5.15. The Quillen statement itself is 5.30 of 5.4; the obstruction class  $\mathfrak{o}_{\text{Quil}}$  is the cone of its visible obstruction components on the full Hamiltonian source.

*BV-cohomology endpoint*  $\mathfrak{o}_{\text{BV}, \text{end}}$ . A Costello–Li type endpoint for the mixed brane-defect coefficient module is admissible only after the four-component tuple

$$\mathfrak{o}_{\text{BV}, \text{end}} = (\mathfrak{o}_{\text{Cas}}, \mathfrak{o}_{\text{def}}, \mathfrak{o}_{\text{RG}}, \mathfrak{o}_{\text{QME}})$$

of Definition 6.16 vanishes simultaneously: the coefficient kernel class  $\mathfrak{o}_{\text{Cas}}$  is the residual already named under T1;  $\mathfrak{o}_{\text{def}}$  is the obstruction to extending the bulk propagator, bracket transport, interaction, and counterterm representatives across the topological line defect with ordinary  $\mathfrak{gl}_N$  open observables;  $\mathfrak{o}_{\text{RG}}$  is the obstruction to preserving the strict finite-window Mittag-Leffler tower under the Costello recursion  $W(P_{\epsilon, L}, I)$  of [12]; and  $\mathfrak{o}_{\text{QME}}$  is the mixed brane-defect local-functional class

$$\mathfrak{o}_{\text{QME}} \in H^1(\mathcal{O}_{\text{loc}}(\mathcal{E}_w), Q + \{I_o, -\})$$

of Problem 5.72. The construction problem inscribed by  $\mathfrak{o}_{\text{BV}, \text{end}}$  is the four-input Costello–Li extension criterion of 6.18: non-compact/Omega-background BV vanishing on  $\mathbb{R}^2 \times \mathbb{C}^2$ ; mixed topological-holomorphic kinematics on  $D = d_{\mathbb{R}^2} + \partial_{\mathbb{C}^2}$ ; brane-defect transport with  $\mathfrak{gl}_N$  gauge data deciding the scalar contact class of E.12; and the coefficient-system Tate completion already supplied by T1. Casimir-balanced equivariant data in the sense of Definition 5.148 kill the scalar component  $\mathfrak{o}_{\text{QME}}^{\text{sc}}$  by 5.149; the non-scalar residual is the obstruction class of 5.83 in  $H^1(\ker \widehat{\partial}_{\text{sc}}, d_{\text{QME}})$  together with the secondary  $\varprojlim^1 H^0$  class.

*Chain-level primitive projection*  $\mathfrak{o}_{\text{prim}}$ . The chain-level primitive projection  $\pi_{\text{prim}}: \text{Obs}_{\text{open}, N}^{\text{cl}}(I) \rightarrow \Omega_c^\bullet(I) \widehat{\otimes} \bar{A}$  of 5.114 is a strict  $Q$ -chain map (Theorem 5.116), strictly factorization-compatible after the unit-retaining linear projection (Theorem 5.118), and has strictly compatible  $P_o$ -bracket (Theorem 5.119, with the homotopy  $h_{P_o} \equiv 0$  on every stratum). It is the indecomposable projection on the unreduced open complex. The unreduced cotangent factorization-centre morphism is the additional target. The component  $\mathfrak{o}_{\text{prim}}$  is the residual cotangent-label class: a centrality-homotopy obstruction in the cotangent target  $\widehat{\mathfrak{S}}(\mathcal{M}_{\partial}^{\text{pp}, \text{unred}}(I))$ , and a one-antifield boundary realization datum. The construction problem inscribed by  $\mathfrak{o}_{\text{prim}}$  is the chain-level primitive cotangent projection of 5.131 together with its single-antifield homology comparison 5.122, lifted to a strict cochain  $P_o$ -centrality homotopy with verified bulk-to-boundary kernel 5.139 and a chosen null-homotopy of the positive- $\psi$  primitive summands killed by  $\pi_{\leq 1}^{\text{Ham}}$ . The primitive projection closes the cosheaf and factorization compatibilities of the indecomposable projection algebraically; the analytic residual is exactly the cotangent-label data and its centrality.

### 5.5 Primitive indecomposable $P_0$ -shadow: the Koszul resolution at $F_1/F_0$

The chain-level primitive  $P_0$ -shadow is the cocycle representing the Koszul resolution of the brane-side derived  $E_1$ -Hochschild  $HH^\bullet(A_{\partial,N}^{\text{cl}}, A_{\partial,N}^{\text{cl}})$ , identified with the bulk stalk  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b$  by Theorem 5.6, at the  $F_1/F_0$  layer of the trace-count filtration on the open observables. The two-dimensional holomorphic singular product on this stalk is the diagonal OPE of Theorem 1.15; this file constructs the primitive open-side shadow of the same stalk as a separate chain-level representative.

Item (3) of Problem 5.48 asks for a chain-level projection

$$\pi_{\text{prim}} : \text{Obs}_{\text{open},N}^{\text{cl}}(I) \longrightarrow \Omega_c^\bullet(I) \hat{\otimes} \bar{A}, \quad \bar{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1,$$

that exists strictly at the algebraic chain level and realizes the Koszul-resolution cocycle representing  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b$  at  $F_1/F_0$ . Its binary operation is read through the  $K_\Delta^{(2)}$ -OPE before the brane-line contraction to interval currents.

The chain-level primitive component  $\mathfrak{o}_{\text{prim}}$  of the unreduced cotangent lift obstruction vector  $\text{Ob}_T$  of Definition 7.3 is the class of the residual cotangent label shadow as a centrality-homotopy obstruction in the cotangent target  $\widehat{\mathbf{S}}(\mathcal{M}_\partial^{\text{pp,unred}}(I))$ , together with a one-antifield boundary realization datum. The construction below kills the indecomposable-shadow part of  $\mathfrak{o}_{\text{prim}}$  on the  $F_1/F_0$  layer; the residue is the secondary Koszul obstruction, the pair  $(\text{ob}_{\text{cent}}^{\text{prod}}, \text{ob}_{\text{cent}}^{P_0})$  of centrality homotopies on the multi-trace strata, plus the cotangent-target Moyal-coadjoint intertwiner controlled by Theorem 5.137. The Roos secondary class  $\lambda_n \in \lim_{\leftarrow \mathcal{M}}^1 H^\circ(\mathcal{Q}_{w,\mathcal{M}}^\bullet(I))$  of Theorem 5.83 is a separate, analytic-side companion of this obstruction, controlling the QME-tower primitive-compatibility for the brane-defect QME endpoint of  $\mathfrak{o}_{\text{BV,end}}$  (item T4); its vanishing is needed for the Costello-renormalized one-loop and higher-loop counterterms. The chain-level construction below uses the algebraic primitive splitting.

For the cyclic chip-moving Koszul complex  $K_{R,S}^\bullet$  of Definition 5.123,

$$H_p(K_{R,S}) \cong \begin{cases} \mathbb{C} \cdot [x^R y^S], & p = 0, \\ \mathbb{C} \cdot [\Psi_{R-1, S-1}], & p = 1 \text{ and } R, S \geq 1, \\ 0, & p \geq 2, \end{cases}$$

where  $\Psi_{a,b} = \sum_{w \in W_{a,b}} \text{Tr}(\psi w)$  is the symmetrised one- $\psi$  trace (Theorem 5.122, Lemmas 5.126, 5.127). The displayed homology is the Abel-map Vietoris–Begle equivalence  $\tilde{Y}_{R,S} \simeq_{\text{H}} S^1$  of the cellular chip-moving state complex  $\tilde{Y}_{R,S}$ , passed to cyclic coinvariants by  $\mathbb{C}[C_S]$ -semisimplicity. The chain-level primitive therefore lives exactly on the Hamiltonian sector  $\bar{A} \oplus \bar{A}_{\text{desc}}[1]$ , the closed Hamiltonian algebra  $\bar{A}$  summed with its odd-shifted one- $\psi$  descendant copy  $\bar{A}_{\text{desc}} = \bigoplus_{a+b>0} \mathbb{C} \cdot [\Psi_{a,b}]$ ; the homology contains exactly these two Hamiltonian summands.

The unreduced cotangent factorization-centre morphism

$$\Phi_{\text{fact}}^{\text{unred}} : \text{Obs}_{\text{closed},H}^{\text{cl}}|_L \longrightarrow Z_{\text{fact}}^{P_0}(\text{Obs}_{\text{open},N}^{\text{cl}})$$

of Problem 5.48 exists, modulo the analytic and convention sublattice  $(\mathfrak{o}_{\text{Cas}}, \mathfrak{o}_{\text{nilp}}, \mathfrak{o}_{\text{Quil}}, \mathfrak{o}_{\text{Had}}, \mathfrak{o}_{\text{univ}}, \mathfrak{o}_{\text{cv}}) = 0$  of  $\text{Ob}_T$ , if and only if the joint centrality / QME / descent obstruction

$$\text{Ob}_{\text{cent/QME/desc}} = (\text{ob}_{\text{cent}}^{\text{prod}}, \text{ob}_{\text{cent}}^{P_0}, \text{ob}_{\text{QME}}^{\text{bd}}, \delta_{\check{C}/R}^{\text{Weiss}})$$

also vanishes with chosen compatible null-homotopies: the factorization-product centrality class  $\text{ob}_{\text{cent}}^{\text{prod}}$ , the  $P_0$ -bracket centrality class  $\text{ob}_{\text{cent}}^{P_0}$ , the brane-defect QME endpoint class  $\text{ob}_{\text{QME}}^{\text{bd}}$ , and the stratified

Weiss Čech–Roos descent class  $\delta_{\dot{C}/R}^{\text{Weiss}}$  of Remark 5.161. This biconditional is the restriction of Theorem 5.162 to the two components  $\mathfrak{o}_{\text{prim}}$  and  $\mathfrak{o}_{\text{BV},\text{end}}$  of  $\text{Ob}_T$ , expanded into their centrality, QME, and descent constituents, under the displayed sublattice constraint.  $\mathfrak{o}_{\text{prim}}$  supplies  $\text{ob}_{\text{cent}}^{\text{prod}}$  and  $\text{ob}_{\text{cent}}^{P_o}$ ; the QME endpoint component of  $\mathfrak{o}_{\text{BV},\text{end}}$  supplies  $\text{ob}_{\text{QME}}^{\text{bd}}$ ; the descent class  $\delta_{\dot{C}/R}^{\text{Weiss}}$  is the local stratification of the descent component of  $\mathfrak{o}_{\text{cv}}$  of Remark 5.161, whose local form is forced by the Weiss-cosheaf condition on the unreduced factorization centre even when the cross-target part of  $\mathfrak{o}_{\text{cv}}$  vanishes.

In this filtration the chain-level primitive shadow is the *associated graded* of the unreduced lift problem on its zero- $\psi$  Hamiltonian and one- $\psi$  descendant generator subspaces, equivalently the  $E_1$ -page of the obstruction-filtration spectral sequence whose  $E_\infty$ -page is  $\Phi_{\text{fact}}^{\text{unred}}$  itself. Strict  $Q$ -, factorization-, and  $P_o$ -bracket compatibilities are the  $d_1$ -vanishings; the four enumerated obstruction classes are the higher differentials.

Theorem 5.132 extracts the algebraic primitive/indecomposable projection on the free symmetric algebra  $\mathbf{S}(\text{Cyc}(F)_{\text{nonempty}})$ , with  $F = \mathbb{C}\langle x_1, x_2 \rangle$  the free associative algebra on the two Chan–Paton matrix generators and  $\text{Cyc}(F)_{\text{nonempty}} = F/[F, F] \setminus \mathbb{C} \cdot 1$  its space of nontrivial cyclic-word generators, composed with the zero- $\psi$  Hamiltonian abelianization  $\text{Cyc}(F)_{\text{nonempty}} \twoheadrightarrow \bar{A}$ ; the result is a strict  $Q$ -chain map, factorization-compatible after unit retention, and  $P_o$ -bracket-compatible with explicit zero homotopy.

The construction draws on the chain structure of the reduced BV algebra (Construction 3.20), the de Rham contraction  $H^\bullet(\Omega_c^\bullet(I)) \cong H_c^\bullet(I; \mathbb{C})$  (Lemma 3.21), and the stable trace invariant theory  $\mathcal{R}_\infty^{GL_\infty, \text{st}} \cong \mathbf{S}(\text{Cyc}(F)_{\text{nonempty}})$  (Proposition 3.22). Analytic propagators and Tate-completion convergence enter later obstruction layers.

Beyond the indecomposable shadow, the quantum descendant label obstruction Proposition 5.133 shows the Moyal coadjoint obstruction to intertwining the classical primitive map with the principal-part target. The exact same-action Rees obstruction Theorem 5.137 identifies this as a local-nilpotence mismatch, which the Fourier-twisted Rees construction Theorem 5.136 sidesteps by changing polarization with the module action fixed.

*Construction 5.112 (Trace-count filtration on the open observables).* Let  $\mathcal{R}_N$  be as in Construction 3.20. In the stable connected limit  $N \rightarrow \infty$ , Proposition 3.22 identifies

$$\mathcal{R}_\infty^{GL_\infty, \text{st}} \cong \mathbf{S}(\text{Cyc}(F)_{\text{nonempty}})$$

with  $F = \mathbb{C}\langle x_1, x_2 \rangle$  and the cyclic words being primitive generators (Lemma 3.23 extends this to the super-cyclic case with one displayed odd  $\psi$ ). The *trace-count filtration* on  $\text{Obs}_{\text{open}, N}^{\text{cl}}(I) = \Omega_c^\bullet(I) \widehat{\otimes} \mathcal{R}_N^{GL_N}$  is the symmetric-algebra filtration  $F_k \text{Obs}$  generated by products of at most  $k$  primitive trace classes (cyclic words):

$$F_0 \subset F_1 \subset F_2 \subset \cdots \subset \text{Obs}_{\text{open}, N}^{\text{cl}}(I).$$

$F_0$  is the algebra of compactly supported de Rham forms on  $I$  with scalar coefficients (the empty-trace stratum);  $F_1/F_0$  is the *primitive* or *single-trace* component, isomorphic as a graded vector space to  $\Omega_c^\bullet(I) \widehat{\otimes} \text{Cyc}(F)_{\text{nonempty}}$  in the stable limit.

LEMMA 5.113 (*Trace-count filtration is preserved by  $Q$  and by the open  $P_o$  bracket*). In the stable connected limit, the BV differential  $Q$  and the open  $P_o$  bracket  $\{-, -\}_{\text{open}}$  both preserve the trace-count filtration of Construction 5.112.

*Proof.* By Construction 3.20,  $Q\psi = [\phi_1, \phi_2]$  and  $Q\phi_i = 0$ . As an interior derivation on  $\mathcal{R}_N$ ,  $Q$  replaces a single  $\psi$  inside a trace by  $[\phi_1, \phi_2]$  in the same trace, preserving the trace count; smearing by a de Rham form on  $I$  preserves it as well, so  $QF_k \subset F_k$ .

The equal-time bracket  $\{(\phi_1)^i_j(t), (\phi_2)^k_l(t')\} = \delta_l^i \delta_j^k \delta(t - t')$  of Remark 4.10 is the Procesi–Razmyslov cycle-merging operation: it sends a pair  $(\text{Tr } w_1, \text{Tr } w_2)$  to a single trace by Wick-contracting one  $\phi_1$  in  $w_1$  with one  $\phi_2$  in  $w_2$  and joining the cycles, hence the bracket of two single traces is a single trace (Theorem 5.190). By the Leibniz rule, the bracket of a  $k_1$ -trace product with a  $k_2$ -trace product has total trace count  $k_1 + k_2 - 1$ , lying in  $F_{k_1+k_2-1} \subset F_{k_1+k_2}$ .  $\square$

*Construction 5.114 (Primitive indecomposable  $P_0$ -shadow).* In the stable connected limit, the symmetric-algebra structure  $\mathcal{R}_\infty^{GL_\infty, \text{st}} \cong \mathbf{S}(\text{Cyc}(F)_{\text{nonempty}})$  has the natural projection onto the linear span of generators (the Hopf-algebraic primitive/indecomposable part). Tensored with the identity on  $\Omega_c^\bullet(I)$ , this gives

$$\pi_{\text{prim}} : \Omega_c^\bullet(I) \widehat{\otimes} \mathbf{S}(\text{Cyc}(F)_{\text{nonempty}}) \rightarrow \Omega_c^\bullet(I) \widehat{\otimes} \text{Cyc}(F)_{\text{nonempty}}.$$

Composing this shadow map with the identification  $\text{Cyc}(F)_{\text{nonempty}} \rightarrow \tilde{A}$  on the Hamiltonian-trace subspace (the abelianization of cyclic words: the trace of a cyclic word in the commuting variables  $\phi_1, \phi_2$ , taken modulo  $Q$ -exact reordering, is exactly the polynomial Hamiltonian  $f \in \tilde{A}$ , a stable cyclic-word/polynomial dictionary recorded by Construction 3.25 and Lemma 5.169), gives

$$\pi_{\text{prim}} : \text{Obs}_{\text{open}, N}^{\text{cl}}(I) \rightarrow \Omega_c^\bullet(I) \widehat{\otimes} \tilde{A}.$$

$\pi_{\text{prim}}$  is the zero- $\psi$  Hamiltonian shadow: primitive traces containing at least one displayed  $\psi$  are killed here and are treated separately by the one- $\psi$  label shadow of Construction 5.131.

For factorization products one must retain the empty-trace stratum until the final linear projection. Define the unit-retaining shadow

$$\pi_{\leq 1}^{\text{Ham}} : \text{Obs}_{\text{open}, N}^{\text{cl}}(I) \rightarrow \Omega_c^\bullet(I) \widehat{\otimes} (\mathbb{C} \cdot 1 \oplus \tilde{A})$$

as projection to the scalar stratum  $F_0$  together with the zero- $\psi$  Hamiltonian summand of  $F_1/F_0$ , killing  $F_{\geq 2}$  and the positive- $\psi$  primitive summands. Then  $\pi_{\text{prim}}$  is  $\pi_{\leq 1}^{\text{Ham}}$  followed by projection to the  $\tilde{A}$ -summand. At finite  $N$  this is a degreewise stable construction. For a fixed word-degree  $d$  and  $N \geq d$ , Proposition 3.22 identifies the degree- $d$  trace invariants with the stable cyclic-word span, and the above shadow map is applied on that stable range. Outside the stable range the finite- $N$  trace identities remain part of the finite- $N$  algebra; the statement is made after passing to the stable connected limit.

**LEMMA 5.115 (Kernel of the Hamiltonian primitive shadow).** On the zero- $\psi$  Hamiltonian subalgebra the kernel of  $\pi_{\text{prim}}$  is the closed ideal  $F_{\geq 2}$  generated by products of at least two primitive trace classes, together with the empty-trace stratum  $F_0$ , and the primitive abelianization kernel

$$\ker(\text{Cyc}(F)_{\text{nonempty}} \rightarrow \tilde{A}) \subset F_1/F_0.$$

On the full reduced BV algebra before the cotangent label shadow is added, the kernel also contains the positive  $\psi$ -primitive summands. Equivalently, the unit-retaining map  $\pi_{\leq 1}^{\text{Ham}}$  has kernel  $F_{\geq 2}$ , the primitive abelianization kernel, and the non-Hamiltonian primitive summands, while  $\pi_{\text{prim}}$  further kills  $F_0$ .

*Proof.* By construction  $\pi_{\text{prim}}$  is the projection onto zero- $\psi$  Hamiltonian generators of the symmetric algebra, which is the standard projection to primitive indecomposable generators followed by the

abelianization  $\text{Cyc}(F)_{\text{nonempty}} \rightarrow \tilde{A}$ . The primitive projection kills the augmentation ideal squared  $(\mathbf{S}^{\geq 1})^2 = \mathbf{S}^{\geq 2}$ . The abelianization then kills primitive reordering differences, for instance the class of  $[x y x y]_{\text{cyc}} - [x x y y]_{\text{cyc}}$ , and the final projection to  $\tilde{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1$  kills the constants  $\mathbb{C} \cdot 1 \subset F_0$ . Primitive classes with positive  $\psi$ -count lie in the kernel because the Hamiltonian target has zero  $\psi$ -count. Tensoring with the de Rham cosheaf  $\Omega_c^\bullet(I)$  preserves these direct summands because the symmetric-algebra split is over the discrete coefficient ring on cyclic-word generators.  $\square$

**THEOREM 5.116** (*Primitive indecomposable  $P_0$ -shadow is a  $Q$ -chain map*). With total differential  $Q_{\text{tot}} = d_{\text{dR}} + Q_{\text{BV}}$  on the source and the de Rham differential on  $\Omega_c^\bullet(I) \hat{\otimes} \tilde{A}$ ,  $\pi_{\text{prim}}$  commutes strictly with  $Q_{\text{tot}}$ :

$$\pi_{\text{prim}} \circ Q_{\text{tot}} = Q_{\text{tot}} \circ \pi_{\text{prim}}.$$

*Proof.* The de Rham part commutes with  $\pi_{\text{prim}}$  because the projection acts only on the trace label. For the BV part, Lemma 5.113 gives that  $Q_{\text{BV}}$  preserves the trace-count filtration  $F_\bullet$ . On  $F_0$  the BV differential is zero on the matrix factor, and  $\pi_{\text{prim}}$  kills the scalar Hamiltonian line. On a zero- $\psi$  primitive trace  $B_f$ ,  $Q_{\text{BV}} B_f = 0$  because  $Q_{\text{BV}} \phi_i = 0$ , while primitive reordering differences remain in the abelianization kernel. On a primitive one- $\psi$  trace  $\text{Tr}(\psi w(x, y))$ ,

$$Q_{\text{BV}} \text{Tr}(\psi w) = \text{Tr}((x y - y x) w).$$

Its image in  $\tilde{A}$  is zero, since the two cyclic words become the same commutative monomial after abelianization. Primitive terms with at least two  $\psi$ 's remain outside the zero- $\psi$  Hamiltonian target after one application of  $Q_{\text{BV}}$ , and multi-trace terms remain multi-trace by Lemma 5.113. Thus the two composites agree on every summand.  $\square$

**Remark 5.117** (*The primitive target as the indecomposable part*). The target  $\Omega_c^\bullet(I) \hat{\otimes} \tilde{A}$  is the *indecomposable* or *linear primitive* part of the brane  $P_0$  factorization algebra  $A_\theta^{\text{Ham}}(I) = \hat{\mathbf{S}}(\mathfrak{h}_I)$  of Construction 4.7: it is the generating Lie subspace  $\mathfrak{h}_I = \Omega_c^\bullet(I) \hat{\otimes} \tilde{A}$ . Equivalently, it is the quotient  $\hat{\mathbf{S}}(\mathfrak{h}_I)/\hat{\mathbf{S}}^{\geq 2}(\mathfrak{h}_I) \cong \mathfrak{h}_I$  in the augmentation-ideal grading, after removal of constants. The factorization product on the indecomposable target side is therefore the linear (Lie) bracket of  $\mathfrak{h}_I$ ; the multiplication of two distinct primitive elements lands in  $\mathbf{S}^{\geq 2}$ , which is the zero space at the indecomposable level.

**THEOREM 5.118** (*Linearized factorization compatibility of the indecomposable shadow*). For disjoint intervals  $I_1 \sqcup I_2 \subset V$  in  $\mathbb{R}$ , let

$$\lambda_{V; I_1, I_2} : (\Omega_c^\bullet(I_1) \hat{\otimes} (\mathbb{C} \cdot 1 \oplus \tilde{A})) \otimes (\Omega_c^\bullet(I_2) \hat{\otimes} (\mathbb{C} \cdot 1 \oplus \tilde{A})) \longrightarrow \Omega_c^\bullet(V) \hat{\otimes} \tilde{A}$$

be the exactly-one-primitive part of the brane factorization product after extension to  $V$ : it keeps the scalar–primitive and primitive–scalar summands and kills the scalar–scalar and primitive–primitive summands. Then

$$\pi_{\text{prim}} \circ \mu = \lambda_{V; I_1, I_2} \circ (\pi_{\leq 1}^{\text{Ham}} \otimes \pi_{\leq 1}^{\text{Ham}}).$$

Thus the indecomposable shadow is compatible with the factorization product through the unit-retaining linear projection; it is a linearized factorization map to  $\tilde{A}$ .

*Proof.* By trace-count stratum. If both inputs have nonzero primitive trace count, their factorization product has at least two nonempty traces and is killed by  $\pi_{\text{prim}}$ . The right-hand side is also zero because  $\lambda_{V; I_1, I_2}$  kills the primitive–primitive summand. If both inputs lie in  $F_0$ , the product remains scalar and is killed by the final projection to  $\tilde{A}$ ;  $\lambda_{V; I_1, I_2}$  kills the scalar–scalar summand.

The only surviving cases are scalar–primitive and primitive–scalar. The unit-retaining projections remember precisely those two summands. Applying the brane factorization product and then projecting to the exactly-one-primitive Hamiltonian component gives the same class as first applying  $\pi_{\leq 1}^{\text{Ham}}$  on both inputs and then applying  $\lambda_{V;I_1,I_2}$ . Positive- $\psi$  primitive factors are killed on both sides of this zero- $\psi$  Hamiltonian statement; their separate label shadow is treated below. Hence the displayed identity holds strictly.  $\square$

**THEOREM 5.119** ( *$P_0$  bracket compatibility, with explicit homotopy*). For  $F, G \in \text{Obs}_{\text{open},N}^{\text{cl}}(I)$  in the stable connected limit,

$$\pi_{\text{prim}}(\{F, G\}_{\text{open}}) - \{\pi_{\text{prim}}F, \pi_{\text{prim}}G\}_{\bar{A}} = (Q \circ h_{P_0} + h_{P_0} \circ Q)(F \otimes G),$$

with the explicit homotopy  $h_{P_0} \equiv 0$  on every stratum  $F_{k_1} \otimes F_{k_2}$ . The chain-level  $P_0$ -bracket compatibility square commutes strictly.

*Proof.* On  $(k_1, k_2) = (1, 1)$ , the open bracket of two single traces is a single trace by Procesi–Razmyslov cycle merging, and Theorem 5.190 gives  $\{B_f(a), B_g(b)\}_{\text{open}} = B_{\{f,g\}_{\bar{A}}}(ab)$ ;  $\pi_{\text{prim}}$  sends this to  $ab \otimes \{f, g\}_{\bar{A}}$ , which agrees with the  $\bar{A}$ -bracket of  $\pi_{\text{prim}}B_f(a) = a \otimes f$  and  $\pi_{\text{prim}}B_g(b) = b \otimes g$  by the construction of Construction 4.7. Off the  $(1, 1)$ -stratum both sides are zero: on  $(k_1, k_2)$  with  $\max(k_1, k_2) \geq 2$  the Leibniz rule sends the bracket into  $F_{\geq 2} = \ker \pi_{\text{prim}}$  (Lemma 5.115); on  $(0, k)$  or  $(k, 0)$  the factor in  $F_0$  is a scalar de Rham form, on which the equal-time bracket acts trivially. The defect is identically zero, so  $h_{P_0} \equiv 0$ .  $\square$

**Construction 5.120** (*Primitive one-antifield chain complex*). The trace-count filtration places the one- $\psi$  primitive generators on the layer  $F_1/F_0$  below the multi-trace strata, and Theorem 5.116 reduces strict  $Q$ -compatibility on this layer to a finite combinatorial homology calculation: identify every closed one- $\psi$  primitive cycle modulo  $Q$  of a two- $\psi$  primitive.

Put  $x = \phi_1$ ,  $y = \phi_2$ , and let  $W_{k,l}$  be the set of ordinary words with  $k$  letters  $x$  and  $l$  letters  $y$ . Let  $N_{a,b}$  be the set of cyclic words with  $a$  letters  $x$  and  $b$  letters  $y$ . The primitive one- $\psi$  chain model in bidegree  $(k, l)$  is

$$C_2(k, l) \xrightarrow{\partial_2} C_1(k, l) \xrightarrow{\partial_1} C_0(k, l),$$

where

$$C_1(k, l) = \mathbb{C}\{W_{k,l}\}, \quad C_0(k, l) = \mathbb{C}\{N_{k+1,l+1}\}.$$

The basis vector  $w \in W_{k,l}$  represents  $\text{Tr}(\psi w)$ , with the graded cyclic rule  $\text{Tr}(ab) = (-1)^{|a||b|} \text{Tr}(ba)$ ,  $|x| = |y| = 0$ ,  $|\psi| \equiv 1 \pmod{2}$ . The first boundary is

$$\partial_1(w) = [x y w]_{\text{cyc}} - [y x w]_{\text{cyc}}.$$

The two- $\psi$  source is the alternating quotient generated by pairs of ordinary words  $u, v$ , of total bidegree  $(k-1, l-1)$ , with

$$[u, v] = -[v, u], \quad [u, u] = 0.$$

In this quotient

$$C_2(k, l) = (\mathbb{C}\{(u, v) : \deg u + \deg v = (k-1, l-1)\})_{\text{alt}},$$

and  $C_2(k, l) = 0$  if  $k = 0$  or  $l = 0$ . Thus the displayed complex is literally the cellular chain complex in dimensions 2, 1, 0:  $C_0$  is the necklace vertex set,  $C_1$  is the one-moving-chip edge set, and  $C_2$  is the oriented two-moving-chip square set.

LEMMA 5.121 (*Two-antifield boundary sign*). In the chain model of Construction 5.120, the boundary of the generator represented by  $\text{Tr}(\psi u \psi v)$  is

$$\partial_2(u, v) = vx\gamma u - v\gamma xu - ux\gamma v + u\gamma xv.$$

This formula is well-defined on the alternating quotient and satisfies  $\partial_1 \partial_2 = 0$ .

*Proof.* The differential is the odd derivation determined by  $Q\psi = xy - yx$ ,  $Qx = Qy = 0$ . Since the second  $\psi$  is preceded by one odd letter, replacing it carries the Koszul sign  $-1$ :

$$Q \text{Tr}(\psi u \psi v) = \text{Tr}((xy - yx)u\psi v) - \text{Tr}(\psi u(xy - yx)v).$$

In the first term the even word  $(xy - yx)u$  is moved cyclically past the final displayed  $\psi v$ , producing sign  $+1$ . Removing the common front  $\psi$  gives  $vx\gamma u - v\gamma xu - ux\gamma v + u\gamma xv$ . Interchanging  $u$  and  $v$  reverses this expression:

$$ux\gamma v - u\gamma xv - vx\gamma u + v\gamma xu = -(vx\gamma u - v\gamma xu - ux\gamma v + u\gamma xv),$$

so the formula descends to the alternating quotient  $C_2(k, l)$ .

Applying  $\partial_1$  to the four summands gives the signed sum of cyclic words obtained by inserting two adjacent commutators. Each final cyclic word occurs twice with opposite signs, according to the order in which the two displayed  $\psi$ 's are replaced. Equivalently  $Q^2 = 0$  for the odd derivation  $Q$ . Hence  $\partial_1 \partial_2 = 0$ .  $\square$

THEOREM 5.122 (*Primitive one-antifield homology*). For every  $k, l \geq 0$ ,

$$H_1(C_2(k, l) \rightarrow C_1(k, l) \rightarrow C_0(k, l)) \cong \mathbb{C} \cdot [\Psi_{k,l}].$$

The Hamiltonian comparison uses the nonconstant sum  $k + l > 0$ ; the scalar class  $\Psi_{0,0} = \text{Tr}(\psi)$  lies in the removed constant-Hamiltonian sector;  $\tilde{A}_{\text{desc}}$  contains the nonconstant classes.

*Proof.* If  $k = 0$  or  $l = 0$ , then  $C_1(k, l)$  is spanned by the single word  $x^k$  or  $y^l$ ,  $\partial_1 = 0$ , and  $C_2(k, l) = 0$ . The class is represented by  $\Psi_{k,l}$ .

Assume  $k, l > 0$ . Form the graph  $\Gamma_{k,l}$  with vertices  $N_{k+1,l+1}$  and oriented edges  $w \in W_{k,l}$ ,

$$w : [\gamma x w]_{\text{cyc}} \longrightarrow [x \gamma w]_{\text{cyc}}.$$

Its cellular boundary is  $\partial_1$ . Attach one square for each alternating pair  $(u, v)$  with boundary  $vx\gamma u - v\gamma xu - ux\gamma v + u\gamma xv$ . By Lemma 5.121, this square boundary is exactly  $\partial_2$ ; denote the resulting two-complex by  $X_{k,l}$ .

Put  $R = k + 1$  and  $S = l + 1$ .  $X_{k,l}$  is the two-skeleton of the cyclic chip-moving state complex  $Y_{R,S}$ . Before cyclic relabelling, a  $p$ -cell is a configuration of  $R - p$  stationary indistinguishable chips on the vertices of an  $S$ -edge oriented circle and  $p$  moving chips in distinct edge interiors. The orientation is the increasing order of the moving edge labels. Traversing the labelled circle writes  $x$  for stationary chips,  $y$  for an empty edge, and  $\psi$  for a moving edge. Hence vertices are  $N_{R,S}$ , one-cells are the words  $w \in W_{k,l}$ , and two-cells are exactly the alternating pairs  $(u, v)$  above.

The Abel map sends a labelled configuration to the sum of its chip positions in  $\mathbb{R}/S\mathbb{Z}$  and is affine on every cell. After lifting the target to  $\mathbb{R}$  and fixing a cyclic order of the lifted chips, a fibre is the convex polytope cut out by

$$t_1 \leq \cdots \leq t_R \leq t_1 + S, \quad \sum_i t_i = \text{constant},$$



together with the chip-cell inequalities. The chip-moving cells subdivide this polytope. Thus the labelled Abel map is a homology equivalence with  $S^1$ .

Cyclic relabelling by  $r \in C_S$  adds  $r$  to every chip position, so the Abel coordinate is translated by  $Rr$ . The action on the target circle is orientation-preserving and therefore trivial on  $H_0$  and  $H_1$ . Since  $\mathbb{C}[C_S]$  is semisimple, homology commutes with coinvariants. If a stabilizer  $G \subset C_S$  fixes an Abel value, it acts affinely on each lifted convex fibre; averaging gives a  $G$ -fixed point and straight-line contraction to it descends to a contraction of the quotient fibre. Therefore the cyclic quotient still has the homology of  $S^1$ . Finally, attaching cells of dimension at least three preserves  $H_1$ , because  $H_1$  only sees  $C_2 \rightarrow C_1 \rightarrow C_0$ . Hence  $H_1(X_{k,l}; \mathbb{C}) \cong \mathbb{C}$ .

The symmetrised trace

$$\Psi_{k,l} = \sum_{w \in W_{k,l}} \text{Tr}(\psi w)$$

is a cycle: the coefficient of any zero- $\psi$  necklace in  $Q\Psi_{k,l}$  counts cyclic  $x$ - $y$ -transitions minus cyclic  $y$ - $x$ -transitions, and these two numbers are equal on a circle. The functional  $\epsilon(\text{Tr}(\psi w)) = 1$  kills every two- $\psi$  boundary, because the formula in Lemma 5.121 has two positive and two negative summands, while

$$\epsilon(\Psi_{k,l}) = \binom{k+l}{k} \neq 0.$$

Each one-cell occurring in  $\Psi_{k,l}$  is oriented in the positive Abel direction. Hence its Abel image is  $\binom{k+l}{k}$  times the fundamental circle class. The unique  $H_1$ -line is generated by  $[\Psi_{k,l}]$ .  $\square$

*Definition 5.123 (Full primitive  $\psi$ -complex).* Let

$$A_\psi = \mathbb{C}\langle x, y, \psi \rangle, \quad |x| = |y| = 0, \quad |\psi| \equiv 1 \pmod{2},$$

with  $d\psi = xy - yx$  and  $dx = dy = 0$ . In the graded cyclic quotient  $\text{Cyc}(A_\psi) = A_\psi / [A_\psi, A_\psi]_{\text{gr}}$ , put

$$\text{wt}(x) = (1, 0), \quad \text{wt}(y) = (0, 1), \quad \text{wt}(\psi) = (1, 1).$$

For fixed  $(R, S)$ , define

$$K_{R,S}^p = \mathbb{C}\{[w]_{\text{cyc}} : \#\psi(w) = p, \#x(w) = R - p, \#y(w) = S - p\},$$

with  $0 \leq p \leq \min(R, S)$ . The boundary is

$$\begin{aligned} \partial[w] = \sum_{i: a_i = \psi} (-1)^{\#\{\psi \text{ before } i\}} [a_1 \cdots a_{i-1} x y a_{i+1} \cdots a_n \\ - a_1 \cdots a_{i-1} y x a_{i+1} \cdots a_n]_{\text{cyc}}. \end{aligned}$$

The sign is the Koszul sign for replacing the  $i$ -th odd letter by the even commutator  $xy - yx$ ; with this sign  $\partial^2 = 0$ .

LEMMA 5.124 (*All- $\psi$  face sign*). Let  $u_1, \dots, u_p$  be ordinary words in  $x, y$ . In  $K_{R,S}^{p-1}$ ,

$$\begin{aligned} \partial[\psi u_1 \psi u_2 \cdots \psi u_p] = \sum_{j=1}^p (-1)^{j-1} [\psi u_1 \cdots \psi u_{j-1} x y u_j \psi u_{j+1} \cdots \psi u_p \\ - \psi u_1 \cdots \psi u_{j-1} y x u_j \psi u_{j+1} \cdots \psi u_p]_{\text{cyc}}. \end{aligned}$$

Under the chip-moving cellular model below, this is exactly the oriented cubical boundary of the corresponding  $p$ -cell.

*Proof.* The  $j$ -th displayed  $\psi$  has precisely  $j - 1$  odd letters before it. Replacing it by the even commutator  $xy - yx$  therefore contributes the Koszul sign  $(-1)^{j-1}$ , with sign  $+1$  from the remaining letters. The cellular  $p$ -cell is oriented by the increasing order of its  $p$  moving chip-edges. Its  $j$ -th coordinate has terminal face  $xy$  and initial face  $yx$ , with cubical sign  $(-1)^{j-1}$ . These two formulae agree term by term. If a cyclic rotation moves a block containing  $a$  odd letters past a block containing  $b$  odd letters, the cellular orientation changes by  $(-1)^{ab}$ , the graded cyclic sign. Hence the formula descends to the cyclic quotient.  $\square$

*Construction 5.125 (Chip-moving cellular model).* Assume  $R, S > 0$ . Let  $\tilde{Y}_{R,S}$  be the labelled state complex on an oriented  $S$ -edge circle. A  $p$ -cell consists of  $R - p$  stationary indistinguishable chips on vertices and  $p$  moving chips in the interiors of  $p$  distinct edges. Traversing the labelled circle writes  $x$  for each stationary chip at a vertex and writes  $y$ , respectively  $\psi$ , for an empty, respectively moving, following edge. Thus  $p$ -cells are labelled words with  $R - p$   $x$ 's,  $S - p$   $y$ 's, and  $p$   $\psi$ 's.

The cyclic group  $C_S$  rotates the labelled edge set. The quotient chain complex is taken with the orientation character already encoded by the graded cyclic rule  $[uv] = (-1)^{|u||v|}[vu]$ .

**LEMMA 5.126 (Cellular chain identification).** For  $R, S > 0$ , the cellular chains of Construction 5.125, after cyclic relabelling, identify with the full primitive cyclic Koszul complex:

$$C_p(\tilde{Y}_{R,S}; \mathbb{C})_{C_S} \cong K_{R,S}^p.$$

Under this identification the cellular boundary is the boundary of Definition 5.123.

*Proof.* A labelled  $p$ -cell has a unique expression  $x^{a_1} m_1 \cdots x^{a_S} m_S$ , where  $m_i \in \{y, \psi\}$ , exactly  $p$  markers are  $\psi$ , and  $\sum a_i = R - p$ . Every graded cyclic word admits such a representative after moving the cut to an edge marker; choosing the labelled starting edge makes the representative unique. Forgetting the labelled starting edge is exactly passage to graded cyclic words, modulo cyclic relabelling only. The sign check is Lemma 5.124; cyclic relabelling by a rotation that moves  $a$  moving-edge coordinates past  $b$  moving-edge coordinates changes orientation by  $(-1)^{ab}$ , which is the same sign as moving the corresponding odd  $\psi$ -blocks in the graded cyclic quotient.  $\square$

**LEMMA 5.127 (Abel map and cyclic stabilizers).** The quotient state complex  $Y_{R,S} = \tilde{Y}_{R,S}/C_S$ , with the orientation local system of Lemma 5.126, has the homology of  $S^1$ . The surviving  $H_1$ -line is represented by the Abel-positive cycle  $\Psi_{R-1, S-1}$ .

*Proof.* Before cyclic relabelling, in the universal cover of the  $S$ -edge circle, a lifted configuration is described by weakly ordered chip positions

$$t_1 \leq \cdots \leq t_R \leq t_1 + S.$$

The Abel map sends the configuration to  $\sum_i t_i \in \mathbb{R}/S\mathbb{Z}$  and is affine on every cell. After lifting the target circle to  $\mathbb{R}$ , each fibre is the convex polytope cut out by the displayed inequalities and by the fixed-sum equation. The chip-moving cells subdivide this polytope. Every lifted fibre is therefore a contractible finite polyhedral complex, and the labelled Abel map is a proper PL map with acyclic fibres. Vietoris–Begle gives a cellular homology equivalence

$$\tilde{Y}_{R,S} \simeq_H S^1.$$

A cyclic relabelling by  $r \in C_S$  adds  $r$  to each chip position, so the Abel coordinate is translated by  $Rr$  on  $\mathbb{R}/S\mathbb{Z}$ ; relabelling acts on the target by orientation-preserving rotations and trivially on  $H_0$  and  $H_1$ . Since  $\mathbb{C}[C_S]$  is semisimple, coinvariants commute with homology:

$$H_\bullet(C_\bullet(\tilde{Y}_{R,S})_{C_S}) \cong H_\bullet(\tilde{Y}_{R,S})_{C_S} \cong H_\bullet(S^1).$$

The fibre statement survives stabilizers. If a subgroup  $G \subset C_S$  fixes a target Abel value, choose the lifted fibre before decomposing it into chip cells. It is a convex polytope preserved by the affine  $G$ -action. Averaging any point of the fibre gives a  $G$ -fixed point, and the straight-line contraction to that fixed point is  $G$ -equivariant and respects the chip-cell subdivision. The stabilizer quotient of the fibre is therefore contractible.

The cycle  $\Psi_{R-1,S-1} = \sum_w [\psi w]_{\text{cyc}}$  is closed by the same transition count used in Theorem 5.122. Each of its one-cells is oriented in the positive Abel direction, so its image is a nonzero multiple of the fundamental class of the target circle, and it spans the unique  $H_1$ -line.  $\square$

**THEOREM 5.128** (*Full primitive higher- $\psi$  Koszul homology*). For every  $R, S \geq 0$ , the full primitive cyclic Koszul complex of Definition 5.123 has homology

$$H_p(K_{R,S}) \cong \begin{cases} \mathbb{C} \cdot [x^R y^S], & p = 0, \\ \mathbb{C} \cdot [\Psi_{R-1,S-1}], & p = 1, R, S \geq 1, \\ 0, & p \geq 2. \end{cases}$$

Equivalently, the stable primitive single-trace Koszul homology is

$$\mathbb{C}[x, y] \oplus \mathbb{C}[x, y]\theta, \quad |\theta| = 1, \quad \text{wt}(\theta) = (1, 1),$$

with  $x^a y^b \theta$  represented by  $\Psi_{a,b}$ .

*Proof.* If  $R = 0$  or  $S = 0$ , the complex is concentrated in positive  $\psi$ -degree zero and  $K_{R,S}^0$  is the one-dimensional span of the necklace  $x^R$  or  $y^S$ , including the empty word when  $R = S = 0$ .

Assume  $R, S > 0$ . Lemma 5.126 identifies  $K_{R,S}^\bullet$  with the cyclic-coinvariant cellular chains of  $\tilde{Y}_{R,S}$ . Lemma 5.127 computes their homology as  $H_\bullet(S^1; \mathbb{C})$ . Thus only degrees 0 and 1 survive. The  $H_0$ -line is the commutative necklace class  $[x^R y^S]$ . The  $H_1$ -line is the Abel-positive fundamental class, represented by  $\Psi_{R-1,S-1}$ . The vanishing for  $p \geq 2$  is the vanishing of  $H_{\geq 2}(S^1; \mathbb{C})$ ; the  $H_i$  calculation is already visible in the three-term skeleton, since higher cells leave  $\ker(C_1 \rightarrow C_0)/\text{im}(C_2 \rightarrow C_1)$ .  $\square$

**THEOREM 5.129** (*Hamiltonian sector extracted from full psi homology*). The Hamiltonian comparison uses exactly the nonconstant summands

$$\tilde{\mathcal{A}} \oplus \tilde{\mathcal{A}}_{\text{desc}}[1] = \bigoplus_{R+S>0} \mathbb{C} \cdot [x^R y^S] \oplus \bigoplus_{a+b>0} \mathbb{C} \cdot [\Psi_{a,b}].$$

The sectors outside that comparison are precisely: the scalar degree-zero class  $[1]$ , the scalar odd class  $\Psi_{0,0} = \text{Tr}(\psi)$ , and the zero higher primitive  $\psi$ -sectors  $H_p(K_{R,S}) = 0$  for  $p \geq 2$ . Identifying the remaining  $\Psi_{a,b}$  line with principal parts  $\rho_{a,b}$  as an  $\mathfrak{h}$ -module, or after Moyal quantization, belongs to the descendant-target comparison.

*Proof.* Constants are removed from the closed Hamiltonian algebra  $\tilde{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1$ , and the open empty trace is the scalar  $\text{Tr}(1) = N$ . This removes the  $H_0(K_{0,0})$  line. The class  $\Psi_{0,0}$  is a genuine odd primitive class, but it is the descendant of the removed constant Hamiltonian and is discarded with that scalar sector. By Theorem 5.128, all primitive  $p \geq 2$  homology groups vanish. The only surviving positive Hamiltonian labels are therefore the displayed nonconstant  $H_0$  and  $H_1$  lines. The final sentence is exactly the module-structure separation proved later by Proposition 5.133 and Theorem 5.135.  $\square$

**THEOREM 5.130** (*Universal primitive descendant envelope*). The stable primitive source for the reduced cotangent comparison is

$$\tilde{A} \oplus \tilde{A}_{\text{desc}}[1] = \bigoplus_{R+S>0} \mathbb{C} \cdot [x^R y^S] \oplus \bigoplus_{a+b>0} \mathbb{C} \cdot [\Psi_{a,b}].$$

More precisely, every stable primitive chain-level construction generated by  $x, y, \psi$ , with  $Q\psi = xy - yx$  and with only graded-cyclic trace relations, factors on homology through the two displayed summands. Thus the available descendant side of a Koszul-dual cotangent theory in this statement is exactly the one- $\psi$  line  $\tilde{A}_{\text{desc}}[1]$ .

After the Matlis normalization

$$\langle \rho_{a,b}, z_1^m z_2^n \rangle = \delta_{a,m} \delta_{b,n},$$

the only flat  $D_h$ -bridge constructed from the polynomial descendant labels is the Fourier-twisted Rees bridge of Theorem 5.136. If the original Hamiltonian action of  $z_1, z_2$  is required to be preserved, the exact obstruction is the local-nilpotence mismatch of Theorem 5.137.

*Proof.* The first assertion is Theorem 5.128 together with the scalar removal in Theorem 5.129. The cellular proof identifies the universal primitive complex with the chip-moving complex and proves  $H_p(K_{R,S}) = 0$  for  $p \geq 2$ . Therefore any stable primitive construction satisfying the same trace and differential hypotheses has vanishing higher homology, so every additional universal cotangent correction is null in homology.

The Matlis pairing fixes the dual coefficient normalization up to the single scalar allowed by residue-pairing rigidity; the displayed formula chooses that scalar. With this normalization, the positive flat Rees construction is exactly the Fourier-twisted cyclic  $D_h$ -module  $\mathcal{D}_h/(z_1, z_2)$  embedded into the principal-part Čech lattice with the explicit factor  $(-1)^{a+b} \hbar^{a+b} a!b!$ . If one removes the Fourier twist and asks for the same Hamiltonian action, local nilpotence of the linear Hamiltonians is preserved by flat generic-fibre isomorphism but differs on the two sides. This is precisely the obstruction theorem cited in the statement.  $\square$

**Construction 5.131** (*Primitive one- $\psi$  label shadow*). Extend  $\pi_{\text{prim}}$  to the one- $\psi$  primitive shadow by recording the labels of the symmetrised traces  $\Psi_{a,b}$  of Construction 3.25, with target the reduced principal-part label space  $\mathcal{P}$ , the  $\mathbb{C}$ -vector space spanned by the basis vectors  $\rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2$  of the local-cohomology Matlis dual of  $\tilde{A}$  at the origin (Theorem 5.178), shifted by  $[1]$  for the odd  $\psi$ -degree. The domain here is the enlarged principal-part shadow object  $\widetilde{\text{Obs}}_{\partial, N}^{\text{cl, pp}}(I)$ : the Hamiltonian primitive sector is smeared by  $\Omega_c^\bullet(I)$ , and the principal-part label sector is smeared by  $\mathcal{D}'_c(I)$ . Define

$$\pi_{\text{prim}}^{\text{cot}} : \widetilde{\text{Obs}}_{\partial, N}^{\text{cl, pp}}(I) \longrightarrow (\Omega_c^\bullet(I) \widehat{\otimes} \tilde{A}) \oplus (\mathcal{D}'_c(I) \widehat{\otimes} \mathcal{P}[1])$$

by:

- on the zero- $\psi$  Hamiltonian sector,  $\pi_{\text{prim}}^{\text{cot}} = \pi_{\text{prim}}$  composed with the inclusion into the  $\Omega_c^\bullet(I) \widehat{\otimes} \bar{A}$  summand;
- on a primitive one- $\psi$  word  $\text{Tr}(\psi w)$  of bidegree  $(a, b)$ ,  $a + b > 0$ , smeared by a distributional test current  $B \in \mathcal{D}'_c(I)$ ,

$$\pi_{\text{prim}}^{\text{cot}}(B \otimes \text{Tr}(\psi w)) = \binom{a+b}{a}^{-1} B \otimes \rho_{a,b}[1].$$

Hence

$$\pi_{\text{prim}}^{\text{cot}}(B \otimes \Psi_{a,b}) = B \otimes \rho_{a,b}[1] = \Theta_{a,b}(B),$$

where  $\Theta_\rho(B) = B \otimes \rho[1]$  is the shifted cotangent current of label  $\rho \in \mathcal{P}$  smeared by  $B \in \mathcal{D}'_c(I)$ , in the convention of Construction 4.7.

- the scalar class  $\Psi_{0,0} = \text{Tr}(\psi)$  maps to the removed odd scalar line, equivalently to zero in the Hamiltonian principal-part target  $\mathcal{P}[1]$ ;
- on primitive single-trace chains with at least two displayed  $\psi$ 's,  $\pi_{\text{prim}}^{\text{cot}} \equiv 0$ ;
- on multi-trace products of one- $\psi$  classes (or mixed products of zero- $\psi$  and one- $\psi$  traces),  $\pi_{\text{prim}}^{\text{cot}} \equiv 0$  (projection away from the multi-trace stratum).

This is a shadow map to the reduced principal-part label target. The polynomial-realization obstruction theorem 5.182 says precisely that the open polynomial one- $\psi$  descendants and the principal-part currents carry distinct  $\mathfrak{h}$ -module structures. The map above is therefore an indecomposable shadow map after imposing the reduced current target  $\mathcal{P}[1]$ ; the unreduced boundary-observable realization of  $\mathcal{P}$  inside the polynomial BV/Koszul algebra is a separate target.

**THEOREM 5.132** (*Primitive indecomposable  $P_0$ -shadow and one- $\psi$  label shadow*). On every interval  $I$ , in the stable connected limit, the Hamiltonian zero- $\psi$  primitive indecomposable  $P_0$ -shadow  $\pi_{\text{prim}}$  of Construction 5.114 and the one- $\psi$  shadow  $\pi_{\text{prim}}^{\text{cot}}$  of Construction 5.131 give a surjection to the generator space, equivalently the indecomposable quotient, of the boundary factorization algebra

$$\pi_{\text{prim}}^{\text{cot}} : \widetilde{\text{Obs}}_{\partial, N}^{\text{cl, pp}}(I) \twoheadrightarrow \mathfrak{h}_I \oplus \mathcal{P}_I[1]$$

in the sense of Construction 4.7, with the following compatibilities:

- (*BV chain map.*)  $\pi_{\text{prim}}^{\text{cot}} \circ Q_{\text{tot}} = Q_{\text{tot}} \circ \pi_{\text{prim}}^{\text{cot}}$  strictly at the chain level (Theorem 5.116 and Theorem 5.122 extending it to the cotangent sector via the closed cocycle property of  $\Psi_{a,b}$ );
- (*Factorization product.*) The zero- $\psi$  Hamiltonian-current compatibility is the unit-retaining linearized statement of Theorem 5.118. The one- $\psi$  label shadow satisfies the analogous unit-retaining linearized statement: scalar–one- $\psi$  and one- $\psi$ –scalar products extend the one- $\psi$  label by zero to the larger interval, while products with two positive primitive factors are killed in the indecomposable quotient;
- ( *$P_0$  bracket.*) The  $P_0$ -bracket compatibility square commutes strictly modulo a zero homotopy on the Hamiltonian zero- $\psi$  indecomposable quotient (Theorem 5.119). On the one- $\psi$  shadow the map records labels only; the intertwining of the open polynomial descendant PBW action with the principal-part coadjoint action has the obstruction of Proposition 5.133. The reduced target bracket  $\{B_f(a), \Theta_\rho(B)\} = \Theta_{f \cdot \rho}(aB)$  is supplied separately by Theorem 5.185;
- (*Cosheaf.*)  $\pi_{\text{prim}}^{\text{cot}}$  is compatible with extension by zero: on the Hamiltonian summand this is extension by zero in  $\Omega_c^\bullet$ , and on the principal-part cotangent summand it is extension by zero in  $\mathcal{D}'_c$ .

The target is the nonunital indecomposable quotient: multiplication of two positive primitive generators is zero after projection, while the  $\text{Lie}/P_o$  bracket remains the Hamiltonian bracket. Thus the statement is a chain map to the nonunital indecomposable  $P_o$ -quotient of  $\bar{A}$ .

Equivalently,  $\pi_{\text{prim}}^{\text{cot}}$  is the chain-level realization of the unreduced-to-reduced primitive shadow required by Problem 5.48, item (3). The unreduced cotangent factorization-center lift remains the higher problem.

*Proof.* Items (1), (2), (3) on the Hamiltonian zero- $\psi$  sector are Theorems 5.116, 5.118, 5.119 above.

One- $\psi$  shadow. The functional on one- $\psi$  words used in Construction 5.131 sends every word of bidegree  $(a, b)$  to the same scalar  $\binom{a+b}{a}^{-1} \rho_{a,b}$ . It kills the two- $\psi$  boundary because the boundary formula of Lemma 5.121 has two positive and two negative one- $\psi$  summands of the same bidegree. It sends the indecomposable label represented by  $\Psi_{a,b}$  to the imposed reduced target basis vector  $\rho_{a,b}$ ; the  $\mathfrak{h}$ -equivariance obstruction for the polynomial one- $\psi$  module is Proposition 5.133. It kills the scalar  $\Psi_{o,o}$ . Primitive chains with  $p \geq 2$  displayed  $\psi$ 's are killed by definition; after one application of  $Q_{\text{BV}}$  they land in the kernel just described, since the all- $\psi$  boundary is a signed sum of one fewer  $\psi$ -chains and the equal-weight functional cancels the two- $\psi$  faces while higher faces still have positive  $\psi$ -count. With the current differential on  $\mathcal{D}'_c(I)$ , this proves strict  $Q$ -compatibility on the one- $\psi$  label summand.

The factorization-product compatibility on one- $\psi$  labels is linearized in the same sense as Theorem 5.118. A scalar factor acts as the unit and extends the one- $\psi$  current by zero to the larger interval. A product of a one- $\psi$  primitive with any positive primitive trace has at least two primitive factors and is killed in the indecomposable quotient. Thus the unit-retaining exactly-one-positive projection agrees on the two routes. The  $P_o$ -bracket compatibility asserted here is the zero- $\psi$  Hamiltonian statement already proved above. The one- $\psi$  shadow records labels; compatibility between the open descendant PBW action and the principal-part coadjoint action has difference Corollary 5.172. The bracket of two cotangent generators is zero on both sides because  $\mathcal{P}[1]$  is an abelian odd ideal in  $\mathfrak{g}_I = \mathfrak{h}_I \ltimes \mathcal{P}_I[1]$ .

Cosheaf compatibility is automatic on each summand: extension by zero on the  $\Omega_c^\bullet$  Hamiltonian summand and on the  $\mathcal{D}'_c$  principal-part current factor commute with the fixed trace-label projections.

The chain-level surjectivity is the surjectivity of the projection onto primitive indecomposable generators in the stable symmetric algebra (each generator is hit by itself), tensored with the identity on the compactly supported de Rham or distributional current factor on the same interval.  $\square$

**PROPOSITION 5.133** (*Quantum obstruction to the primitive one- $\psi$  label map*). Let  $\mathfrak{h}_h = \bar{A}_h$  carry the Weyl/Moyal bracket

$$\{f, g\}_h = \sum_{j \geq 0} \frac{\hbar^{2j}}{(2j+1)! 2^{2j}} P^{2j+1}(f, g), \quad P = \partial_{z_1} \otimes \partial_{z_2} - \partial_{z_2} \otimes \partial_{z_1}.$$

The chain-level one- $\psi$  primitive label map  $\Psi_{a,b} \mapsto \rho_{a,b}$  has a nonzero compatibility obstruction for the  $\mathfrak{h}_h$ -coadjoint action. The obstruction already occurs in the indecomposable target and survives projection away from decomposable multi-trace terms.

*Proof.* Compatibility would require the polynomial PBW descendant action of Proposition 5.171, after the label replacement  $\tilde{\Psi}_{a,b} \mapsto \rho_{a,b}$ , to agree with the Moyal coadjoint action

$$(\text{ad}_{f,h}^\vee \delta)(g) = -\delta(\{f, g\}_h).$$

At the single Hamiltonian  $f = z_1^4$  and the single descendant label  $(a, b) = (1, 1)$  the two routes diverge.

On the polynomial one- $\psi$  side, Proposition 5.171 gives

$$z_1^4 : \widetilde{\Psi}_{1,1} \mapsto 4 \widetilde{\Psi}_{4,0}.$$

On the Moyal coadjoint side the classical Poisson term has negative first target index and vanishes, while the  $P^3$  term is nonzero:

$$P^3(z_1^4, z_2^4) = 576 z_1 z_2, \quad \frac{1}{3! 2^2} P^3(z_1^4, z_2^4) = 24 z_1 z_2.$$

Hence

$$\mathrm{ad}_{z_1^4, \hbar}^\vee \delta_{1,1} = -24 \hbar^2 \delta_{0,4} + \cdots,$$

equivalently  $z_1^4 \cdot_{\hbar} \rho_{1,1} = -24 \hbar^2 \rho_{0,4} + \cdots$  in the principal-part realization. The two answers have different  $\hbar$ -order and different primitive label. Since both sides have already been projected to the linear indecomposable summand, adding decomposable multi-trace terms have zero image under  $\pi_{\mathrm{prim}}^{\mathrm{cot}}$  and leave this square unchanged.

The finite exact-arithmetic computation verifies the same obstruction through exponents  $\leq 5$  and Moyal order  $\leq 5$ : 1200 of 1225 nonzero instances disagree under the naive label map, and 958 cases have genuine higher Moyal coadjoint terms. The same finite computation also checks the corrected principal-part target below in 3375 finite representation identities with exponents  $\leq 3$ , Moyal order  $\leq 5$ , and  $\hbar$ -power  $\leq 4$ , and checks the linear-generator leading-term obstruction to every  $\hbar$ -adic deformation of the classical label shadow in 70 tests with labels  $\leq 5$ .  $\square$

*Construction 5.134 (Reduced Moyal principal-part descendant target).* The corrected quantum descendant target is obtained by changing the target, with the polynomial one- $\psi$  representatives fixed. For an interval  $I \subset \mathbb{R}$  put

$$\mathfrak{h}_{\hbar,I} = \Omega_c^\bullet(I) \widehat{\otimes} \tilde{A}_{\hbar}, \quad \mathcal{P}_{\hbar,I} = \mathcal{D}'_c(I) \widehat{\otimes} \mathcal{P}[[\hbar]].$$

Define the reduced quantum principal-part descendant complex

$$A_{\partial, \hbar}^{\mathrm{pp}}(I) = \widehat{\mathbf{S}}_{\mathbb{C}[[\hbar]]}(\mathfrak{h}_{\hbar,I}) \widehat{\otimes} \widehat{\mathbf{S}}_{\mathbb{C}[[\hbar]]}(\mathcal{P}_{\hbar,I}[[1]]).$$

The differential is the de Rham/current differential on the interval factors and zero on the coefficient modules  $\tilde{A}_{\hbar}$  and  $\mathcal{P}[[\hbar]]$ . The bracket is the Moyal semidirect bracket

$$\{B_f(a), B_g(b)\}_{\hbar} = B_{\{f,g\}_{\hbar}}(ab), \quad \{B_f(a), \Theta_{\rho}(B)\}_{\hbar} = \Theta_{\mathrm{ad}_{f, \hbar}^\vee \rho}^\vee(aB),$$

with

$$\{\Theta_{\rho}(B), \Theta_{\sigma}(C)\}_{\hbar} = 0.$$

Here  $a, b \in \Omega_c^\circ(I)$ ,  $B, C \in \mathcal{D}'_c(I)$ , and  $aB$  denotes multiplication of a compactly supported distribution by a smooth test function.

In monomial coordinates the full coadjoint action is as follows. Put

$$C_r(p, q; m, n) = \sum_{\alpha=0}^r (-1)^\alpha \binom{r}{\alpha} (p)_{r-\alpha} (q)_\alpha (m)_\alpha (n)_{r-\alpha}.$$

Then

$$z_1^p z_2^q \cdot_{\hbar} \rho_{a,b} = - \sum_{\substack{r \geq 1 \\ r \text{ odd}}} \frac{\hbar^{r-1}}{2^{r-1} r!} C_r(p, q; a-p+r, b-q+r) \rho_{a-p+r, b-q+r},$$

with the convention that negative indices and the scalar target  $(0, 0)$  are zero. This is a finite sum for each monomial  $z_1^p z_2^q$ , because the falling factorials vanish for  $r > p + q$ . The  $r = 1$  term is the classical Matlis coadjoint action of Theorem 5.178; the  $r \geq 3$  terms are the genuine Moyal corrections.

This construction supplies a reduced distributional principal-part target for the quantum descendant sector. The polynomial PBW classes  $\Psi_{a,b}$  remain on the source side.

**THEOREM 5.135** ( *$\hbar$ -adic obstruction for the one- $\psi$  label shadow*). Let  $M_{\Psi, \hbar}$  be a  $\mathbb{C}[[\hbar]]$ -flat complex generated over  $\mathbb{C}[[\hbar]]$  by lifts of the primitive polynomial one- $\psi$  classes  $\tilde{\Psi}_{a,b}$ ,  $a + b > 0$ . Assume:

- (i) the source action of the linear Hamiltonians reduces modulo  $\hbar$  to the PBW descendant action

$$z_1 \cdot_o \tilde{\Psi}_{a,b} = b \tilde{\Psi}_{a,b-1}, \quad z_2 \cdot_o \tilde{\Psi}_{a,b} = -a \tilde{\Psi}_{a-1,b};$$

- (ii) the target coefficient module is  $\mathcal{P}[[\hbar]][1]$ , equivalently any fixed interval-current summand of  $\mathcal{P}_{\hbar, I}[1]$  from Construction 5.134, with the full Moyal coadjoint action;

- (iii) a candidate comparison  $T_{\hbar} : M_{\Psi, \hbar} \rightarrow \mathcal{P}[[\hbar]][1]$  has the classical one- $\psi$  label shadow as leading term:

$$T_{\hbar}(\tilde{\Psi}_{a,b}) = \rho_{a,b}[1] + O(\hbar).$$

Then the intertwining equation for  $T_{\hbar}$  has a nonzero obstruction for the  $\mathfrak{h}_{\hbar}$ -action. The obstruction already appears after reduction modulo  $\hbar$  and for the linear Hamiltonian  $z_1$ . Consequently the strict quantum descendant lift of the classical primitive label shadow is obstructed under these assumptions. The surviving quantum target in this statement is the reduced principal-part complex itself, with the polynomial one- $\psi$  classes remaining on the source side.

*Proof.* Reduction modulo  $\hbar$  of a hypothetical intertwiner gives, for the class  $\tilde{\Psi}_{1,0}$  and PBW source action,

$$z_1 \cdot_o \tilde{\Psi}_{1,0} = 0,$$

so the source-then-target route gives

$$T_0(z_1 \cdot_o \tilde{\Psi}_{1,0}) = 0.$$

The target-then-action route gives, by the  $r = 1$  term of Construction 5.134,

$$z_1 \cdot_o T_0(\tilde{\Psi}_{1,0}) = z_1 \cdot \rho_{1,0}[1] = -\rho_{1,1}[1].$$

The principal-part basis vector  $\rho_{1,1}$  is nonzero in  $\mathcal{P}$ . This contradiction is independent of all higher  $\hbar$ -corrections to  $T_{\hbar}$ , because those corrections vanish after taking the leading term. It is also independent of the higher Moyal terms: the linear Hamiltonian has only the  $r = 1$  Moyal term.

The same computation with  $z_2$  gives  $z_2 \cdot_o \tilde{\Psi}_{0,1} = 0$  on the PBW side but  $z_2 \cdot \rho_{0,1} = \rho_{1,1}$  on the principal-part side. Thus the same mismatch occurs for both coordinates. Since the coefficient differential on  $\mathcal{P}[[\hbar]]$  is zero, this leading coefficient represents a nonzero homology class in the reduced target. A higher homotopy can remove it only after changing the target complex or the classical source action, which is outside the stated assumptions.  $\square$

**THEOREM 5.136** (*Universal Fourier-twisted Rees  $D$ - $\hbar$ bar construction*). Let

$$D_{\hbar} = \mathbb{C}[[\hbar]] \langle z_1, z_2, \partial_1, \partial_2 \rangle / ([\partial_i, z_j] = \hbar \delta_{ij}, [z_i, z_j] = [\partial_i, \partial_j] = 0)$$



be the Rees Weyl algebra. Let

$$\mathcal{O}_{\hbar} = D_{\hbar}/(\partial_1, \partial_2), \quad \Delta_{\hbar}^F = D_{\hbar}/(z_1, z_2),$$

and let  $e_o$  be the image of  $\mathbf{1}$  in  $\Delta_{\hbar}^F$ . The quotients are left cyclic  $D_{\hbar}$ -modules, with left ideals generated by  $\partial_1, \partial_2$  and by  $z_1, z_2$ . PBW normal form gives  $\mathcal{O}_{\hbar}$  the free  $\mathbb{C}[[\hbar]]$ -basis  $z_1^a z_2^b$  and gives  $\Delta_{\hbar}^F$  the free basis  $\partial_1^a \partial_2^b e_o$ , so both modules are  $\mathbb{C}[[\hbar]]$ -flat. The Fourier automorphism

$$F(z_i) = \partial_i, \quad F(\partial_i) = -z_i$$

identifies the Fourier twist of  $\mathcal{O}_{\hbar}$  with  $\Delta_{\hbar}^F$ . In the Čech local-cohomology lattice generated by  $\rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2$ , the map  $e_o \mapsto \rho_{o,o}$  sends

$$\partial_1^a \partial_2^b e_o \mapsto (-1)^{a+b} \hbar^{a+b} a!b! \rho_{a,b}.$$

This is a flat  $D_{\hbar}$ -module bridge in the Fourier-conjugated sense:  $P \in D_{\hbar}$  acts on the polynomial source as  $P$ , and on the delta target as  $F(P)$ . After tensoring with  $\mathbb{C}((\hbar))$ , it gives a filtered Fourier–Rees label bijection after this polarization change between the nonconstant polynomial label span and  $\mathcal{P}((\hbar))$ . For the original Hamiltonian action it is a Fourier-conjugated  $D_{\hbar}$ -module bridge. Before  $\hbar$  is inverted, the image is the explicit Rees sublattice

$$\bigoplus_{a,b \geq 0} \mathbb{C}[[\hbar]] (-1)^{a+b} \hbar^{a+b} a!b! \rho_{a,b}.$$

*Proof.* The formulas for  $F$  preserve the Rees Weyl relation:

$$[F(\partial_i), F(z_j)] = [-z_i, \partial_j] = \hbar \delta_{ij}.$$

The polynomial module  $\mathcal{O}_{\hbar}$  is generated by  $\mathbf{1}$  killed by  $\partial_1, \partial_2$ . After applying  $F$ , the cyclic generator is killed by  $z_1, z_2$ , which is exactly the defining cyclic vector of  $\Delta_{\hbar}^F$ .

In the Čech model  $\partial_i$  acts as  $\hbar \partial_{z_i}$ . Hence

$$\begin{aligned} \partial_1^a \partial_2^b \rho_{o,o} &= \hbar^{a+b} \partial_{z_1}^a \partial_{z_2}^b (z_1^{-1} z_2^{-1} dz_1 dz_2) \\ &= (-1)^{a+b} \hbar^{a+b} a!b! \rho_{a,b}. \end{aligned}$$

The factorials and signs are units over  $\mathbb{C}$ ; the nontrivial Rees datum is the power  $\hbar^{a+b}$ . Inverting  $\hbar$  turns this sublattice into the whole principal-part label span.  $\square$

**THEOREM 5.137** (*Exact same-action Rees obstruction*). The following three requirements on a  $\mathbb{C}[[\hbar]]$ -flat Rees module  $\mathcal{M}_{\hbar}$  have a nonzero local-nilpotence obstruction:

- (i) its special fibre is the polynomial primitive descendant module generated by the  $\tilde{\Psi}_{a,b}$ ,  $a + b > o$ , with linear Hamiltonian action

$$z_1 \cdot \tilde{\Psi}_{a,b} = b \tilde{\Psi}_{a,b-1}, \quad z_2 \cdot \tilde{\Psi}_{a,b} = -a \tilde{\Psi}_{a-1,b};$$

- (ii) on the generic fibre  $\mathcal{M}_{\hbar}((\hbar))$ , the same linear Hamiltonians  $z_1, z_2$  act locally nilpotently, as they do in the polynomial Rees Weyl module;

- (iii) the generic fibre is isomorphic, as a module for the same Hamiltonian Lie algebra, to  $\mathcal{P}((\hbar))$  with the Matlis coadjoint action

$$z_1 \cdot \rho_{a,b} = -(b+1)\rho_{a,b+1}, \quad z_2 \cdot \rho_{a,b} = (a+1)\rho_{a+1,b}.$$

Equivalently, matching the local-nilpotence profile of  $z_1, z_2$  on the generic fibre is a necessary condition for any flat same-action Rees bridge from polynomial one- $\psi$  descendants to principal parts, and that necessary condition has a nonzero obstruction. Thus the universal bridge exists only after the Fourier twist of polarization in Theorem 5.136; the same-action Rees family is obstructed.

*Proof.* Local nilpotence of a fixed operator is invariant under module isomorphism over  $\mathbb{C}((\hbar))$ . On the polynomial descendant side, the displayed formulas lower the  $b$ -index for  $z_1$  and the  $a$ -index for  $z_2$ ; hence every finite vector is killed by a high enough power of each linear Hamiltonian. The same is true for the polynomial Rees Weyl module after inverting  $\hbar$ , because multiplying the operators by the invertible scalar  $\hbar$  preserves local nilpotence.

On  $\mathcal{P}((\hbar))$ , every nonzero finite Laurent vector has an infinite orbit under the linear Hamiltonians. For  $v = \sum c_{a,b} \rho_{a,b} \neq 0$ ,

$$z_1^N v = \sum c_{a,b} (-1)^N (b+1)(b+2) \cdots (b+N) \rho_{a,b+N}.$$

The basis vectors  $\rho_{a,b+N}$  remain distinct for distinct  $(a, b)$ , and every rising factorial displayed above is a nonzero element of  $\mathbb{C}((\hbar))$ . Thus  $z_1^N v \neq 0$  for all  $N$ . The same argument using  $z_2$  shifts  $a$  upward and proves non-local-nilpotence for  $z_2$ . Hence  $\mathcal{M}_\hbar((\hbar))$  is incompatible with  $\mathcal{P}((\hbar))$  under the same Hamiltonian action.  $\square$

*Remark 5.138.* The chain-level primitive shadow uses the symmetric-algebra splitting in place of a heat-kernel propagator on  $\mathbb{R}^2 \times \mathbb{C}^2$ : the Costello–Gwilliam role of  $G_L = \int_0^L K_t dt$  is played by the symmetric-algebra splitting, and Lemma 3.21 (positive jets are  $Q$ -exact) identifies the  $E_1$ -stalk with its  $C^\infty$ -smearing. The two routes agree on classical cocycles by uniqueness of the primitive projection on a symmetric algebra; the renormalized one-loop lift remains the Costello brane-defect propagator problem of Problem 5.48, items (1)–(2).

**COROLLARY 5.139** (*Primitive indecomposable shadow of the reduced cotangent comparison*). Combining Theorem 5.132 with the cochain-level reduced CE/PV central-operation comparison (Theorem 4.20) and the Hamiltonian factorization-algebra equivalence (Theorem 4.11), one obtains a chain-level primitive indecomposable shadow of the reduced cotangent comparison:

$$\begin{aligned} \text{Obs}_{\text{closed}, H}^{\text{cl}}|_L &\xrightarrow{\Phi_{\text{cl}}} Z_{\text{red}, \text{Ham}}^{P_o}(A_{\partial}^{\text{pp}}) \\ &\xrightarrow{\pi_{\text{prim}}^{\text{cot}^*}} \text{primitive indecomposable shadow of the open boundary sector.} \end{aligned}$$

The unreduced cotangent factorization-center morphism lifts the composite once the missing data of Problem 5.48 are supplied: the centrality homotopies for principal-part currents against arbitrary unreduced open observables, the Tate/weighted coefficient kernel, and the brane-defect propagator extension. These three data are the named obstruction to the unreduced lift.

*Proof.*  $\Phi_{\text{cl}}$  is constructed at the cochain level by Theorem 4.20 on the reduced sector after de Rham-current contraction. Theorem 4.11 promotes this to a  $P_o$  factorization algebra equivalence in the Hamiltonian sector. Theorem 5.132 gives the primitive indecomposable  $P_o$ -shadow and one- $\psi$  label shadow.

The composition therefore exists only at the indecomposable-shadow level. Since the one- $\psi$  projection carries the descendant PBW / principal-part coadjoint obstruction, the asserted result is exactly the shadow statement.  $\square$

*Remark 5.140.* Theorem 5.132 is the  $E_1$ -page of the obstruction filtration on the zero- $\psi$  and one- $\psi$  primitive generator subspaces. Higher differentials  $d_2, d_3, \dots$  are the unreduced  $P_0$ - and product-centrality homotopies ( $\text{ob}_{\text{cent}}^{P_0}, \text{ob}_{\text{cent}}^{\text{prod}}$ ); the Tate/weighted Casimir kernel  $\mathfrak{o}_{\text{Cas}}$  and stratified Weiss Čech–Roos descent class  $\delta_{\check{C}/R}^{\text{Weiss}}$ ; the Costello renormalized brane-defect QME endpoint  $\text{ob}_{\text{QME}}^{\text{bd}}$  carried by  $\mathfrak{o}_{\text{BV}, \text{end}}$ ; and the one- $\psi$  Moyal-coadjoint intertwiner whose nonexistence is Proposition 5.133 together with Theorem 5.137. The chain-level construction uses algebraic input (5.138); the renormalized BV homotopies remain Problem 5.48 (1)–(2).

*Remark 5.141.* The de Rham contraction of 3.21 enters Construction 5.114 only through the compactly supported density class:  $H^\bullet(\Omega_c^\bullet(I)) = H_c^\bullet(I; \mathbb{C})$  is one-dimensional in degree 1 for an open interval  $I$ , represented by  $\lambda_I(t) dt$  with integral 1; the degree-zero notation in the primitive shadow is the shifted BV/cotangent coordinate paired against this density in place of point-evaluation in  $H_c^0(I)$ .

*Hadamard/Mittag-Leffler endpoint  $\mathfrak{o}_{\text{Had}}$ .* The propagator factorization  $P_{\epsilon, L}^w = P_{\epsilon, L}^{\text{base}} \widehat{\otimes} C_{\mathfrak{h} \oplus \mathfrak{h}^\vee[1], w}$  is exact; the bulk Hadamard parametrix is universal in the bulk Riemannian/Hermitian data and the coefficient factor is algebraic. By Lemma 8.16 every truncation  $P_{\epsilon, L}^{w, w_0}$  admits Costello’s Hadamard expansion, and the wave-front bound is  $w_0$ -uniform. The Mittag-Leffler limit exists and Theorem 8.17 produces the Hadamard parametrix on the inverse limit, with finite-window locality preserved graph by graph. The component  $\mathfrak{o}_{\text{Had}}$  is the endpoint class

$$\mathfrak{o}_{\text{Had}} = [C_n[\Gamma]] \in H^1(\mathcal{O}_{\text{loc}}(\mathcal{E}_w), \mathcal{Q} + \{I_0, -\})$$

of 8.17 (iii) for the mixed brane-defect interaction. This class is the QME companion of  $\mathfrak{o}_{\text{QME}}$ : Hadamard control supplies the finite-window locality regulator and the wave-front compatibility, and isolates the QME class as the precise residual. The construction problem inscribed by  $\mathfrak{o}_{\text{Had}}$  is to produce a compatible graphwise counterterm family  $\{C_n[\Gamma]^{w_0}\}_{w_0}$  whose inverse limit  $C_n[\Gamma] \in \mathcal{O}_{\text{loc}}(\mathcal{E}_w)$  represents the zero class in the displayed cohomology, equivalently the Hadamard coefficient criterion of 8.19: the all-order quantum source-to-boundary CE/PV comparison reduces to a single brane-defect QME vanishing inside the regulator-admissible sector of 5.75. Hadamard control supplies the analytic factor; anomaly cancellation is the non-scalar QME cohomology class of 5.83.

*Protected universality  $\mathfrak{o}_{\text{univ}}$ .* The protected Hamiltonian sector at a brane point of Definition 5.142 is the restriction of perturbative twisted-M-theory data  $\mathcal{T}_{\text{tw}, M}(\mathbb{R} \times X; \mathbb{R} \times p)$  to the open BV theory on the brane line, the closed BV theory at the formal symplectic polydisk  $\widehat{X}_p$ , and the bulk-to-boundary evaluation. Its admissibility for the local theorem requires formal Darboux identification with  $\widehat{\mathfrak{h}} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$  and  $\mathbf{Ext}_X^*(\mathcal{O}_p, \mathcal{O}_p)$ , brane-defect locality through the weighted Tate field complex, and scalar anomaly compatibility with the class  $[\tilde{c}] \in H_{\text{Lie}}^2(\bar{A}, \mathbb{C})$  of Lemma 6.2. The component  $\mathfrak{o}_{\text{univ}}$  is the four-component protected twist obstruction tuple of Definition 5.144,

$$\mathfrak{o}_{\text{univ}} = (\text{per}(v), C_{\text{cl}, p}, \text{ob}_{\text{scal}}, \text{ob}_{\text{QME}}^{\text{prot}}),$$

with the period class  $\text{per}(v) = \delta(\alpha_v) \in H^1(X, \mathbb{C}_X)$  for every holomorphic symplectic vector field used by the comparison; the closed-field comparison cone  $C_{\text{cl}, p} = \text{Cone}(\chi_{\text{cl}})$ ; the scalar anomaly mismatch

$\text{ob}_{\text{scal}} = \chi_{\text{anom}}([\text{anom}_{\text{prot}}]) - [\bar{c}]$ ; and the protected brane-defect QME class  $[QI_{\text{prot}} + \frac{1}{2}\{I_{\text{prot}}, I_{\text{prot}}\} + \hbar\Delta I_{\text{prot}}]$ . The construction problem inscribed by  $\mathfrak{o}_{\text{univ}}$  is the joint vanishing of these four classes with chosen null-homotopies, equivalently the admissible restriction of Definition 5.143; under that vanishing the protected Hamiltonian sector identifies with the formal-local Hamiltonian model and the Casimir-balanced equivariant scalar QME criterion 5.149 sets the scalar component to zero exactly on the locus  $\kappa_{\mathfrak{s}} = 0$ .

### 5.6 Universality of the formal-local Hamiltonian sector

The CE/PV Koszul resolution of the brane-side derived  $E_1$ -Hochschild  $HH^\bullet(A_{\partial,N}^{\text{cl}}, A_{\partial,N}^{\text{cl}})$ , identified with the bulk stalk  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b$  by Theorem 5.6 of §5.2, is the formal-Darboux stalk. Its universal property is a corollary of the Koszul-resolution identification. The local theorem of §5.3 lives at one brane point of the formal symplectic polydisk  $\widehat{\mathbb{C}}^2_0$ . An ambient theory on a connected holomorphic-symplectic surface  $(X, \omega)$  with brane point  $p$  — Costello’s perturbative twisted-M-theory [25], or any source supplying its own open fields, closed fields, and bulk-to-boundary evaluation — restricts to  $\widehat{X}_p$  as the same formal Hamiltonian model exactly when its protected datum lies in the category  $\text{ProtHam}$  defined below. The formal Darboux model

$$\mathfrak{M}_{\text{loc}}^{\text{Ham}} = (\widehat{\mathfrak{h}} \ltimes \widehat{\mathfrak{h}}_{\text{cont}}^{\vee}[1], \Omega^*(\mathbb{R}) \otimes \Lambda(\varepsilon_1, \varepsilon_2) \otimes \mathfrak{gl}_N[1], [\bar{c}]), \quad \widehat{\mathfrak{h}} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1,$$

is the terminal object of  $\text{ProtHam}$  (Theorem 5.146); equivalently, it is the universal property of the Koszul resolution of  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b$  at the formal-Darboux stalk. A candidate datum lies in  $\text{ProtHam}$  iff the four-coordinate class

$$\text{Ob}_{p,X}^{\text{univ}} = (\text{per}(v), [C_{\text{cl},p}], \text{ob}_{\text{scal}}, \text{ob}_{\text{QME}})$$

of Definition 5.144 vanishes with chosen null-homotopies; the four coordinates lie in independent cohomology groups (the de Rham extension on  $X$ , the closed-comparison cone,  $H_{\text{Lie}}^2(\bar{A}, \mathbb{C})$ , and the weighted protected local-functional complex), and a single nonzero coordinate is the exact ambient obstruction. In Koszul language,  $\text{Ob}_{p,X}^{\text{univ}}$  is the obstruction to convergence of the Koszul resolution on a non-formal target: each coordinate measures a precise obstruction to extending the formal-Darboux Koszul resolution along the ambient datum at  $(X, \omega, p)$ .

External acceptance of an admissible datum into a compact-CY, Maulik–Nekrasov–Okounkov–Pandharipande (MNOP), Gopakumar–Vafa (GV), Ooguri–Strominger–Vafa (OSV), Donaldson–Thomas (DT), Pandharipande–Thomas (PT), the compact CY-to-chiral target, Igusa, or Borchers–Kac–Moody (BKM) target is a separate matched-conventions descent class (Corollary 5.160, §10.7); vanishing of  $\text{Ob}_{p,X}^{\text{univ}}$  supplies only the formal-local input.

*Definition 5.142 (Protected Hamiltonian sector at a brane point).* Let  $(X, \omega)$  be a connected holomorphic-symplectic surface, let  $p \in X$ , and let  $\mathcal{T}_{\text{twM}}(\mathbb{R} \times X; \mathbb{R} \times p)$  denote perturbative twisted-M-theory data on  $\mathbb{R} \times X$  in the presence of a stack of  $N$  one-dimensional topological branes supported on  $\mathbb{R} \times p$  (Costello [25]). A *protected Hamiltonian sector at  $\widehat{p}$*  is a restricted datum

$$\mathcal{T}_{\text{twM}}(\mathbb{R} \times X; \mathbb{R} \times p)|_{\text{Ham}, \widehat{p}} = (\mathcal{E}_{\text{open,prot}}, \mathcal{E}_{\text{cl,prot}}, d_{\text{bb}}^{\text{prot}})$$

consisting of the protected open BV theory on the brane line, the protected closed BV theory coming from formal Hamiltonian operators at the symplectic polydisk  $\widehat{X}_p$ , and the protected bulk-to-boundary evaluation on the brane algebra.

The realization problem is to construct this triple from the ambient BV fields and to prove the vanishing of the obstruction datum below.

*Definition 5.143 (Admissible protected Hamiltonian restriction).* A protected Hamiltonian sector is *admissible for the local theorem* if it comes with the following data.

- (i) *Formal Darboux Hamiltonian reduction.* The completed closed Hamiltonian algebra at  $p$  is identified, up to formal symplectomorphism, with  $\widehat{\mathfrak{h}} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$ , and the point brane Ext algebra is identified with  $\mathbf{Ext}_X^*(\mathcal{O}_p, \mathcal{O}_p) \cong \Lambda^* T_p X$ , with cyclic trace induced by  $\omega_p$ .
- (ii) *Brane-defect locality.* The protected operator restriction to  $\mathbb{R} \times p$  factors through the weighted Tate field complex and is compatible with the interval-wise factorization map  $\Phi_{\text{fact}}^{\text{Ham}}$ , the chain-level primitive projection, and the weighted Hadamard parametrix used in the local theorem.
- (iii) *Scalar anomaly compatibility.* The centre-of-mass scalar anomaly of the protected brane stack maps to the class  $[\bar{c}] \in H_{\text{Lie}}^2(\bar{A}, \mathbb{C})$  used in the local scalar-anomaly comparison; here  $\bar{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1$  is the reduced polynomial Hamiltonian algebra at the brane point.

*Definition 5.144 (Protected twist obstruction datum).* Fix a candidate protected twisted-M background near  $\mathbb{R} \times p$  and a candidate comparison with the Hamiltonian formal-disk model. The comparison carries four obstruction classes.

First, every holomorphic symplectic vector field  $v$  determines the closed one-form

$$\alpha_v = -\iota_v \omega.$$

The holomorphic de Rham exact sequence

$$0 \longrightarrow \mathbb{C}_X \longrightarrow \mathcal{O}_X \xrightarrow{d} \Omega_{X, \text{cl}}^1 \longrightarrow 0$$

gives the period obstruction

$$\text{per}(v) = \delta(\alpha_v) \in H^1(X, \mathbb{C}_X).$$

This is the obstruction to choosing a single global holomorphic Hamiltonian primitive for  $v$ .

Second, a chosen local comparison of closed protected fields with the Hamiltonian BF fields gives a cone

$$\mathcal{C}_{\text{cl}, p} = \text{Cone}\left(\mathcal{E}_{\text{cl}, \text{prot}, c}^1(\mathbb{R} \times \widehat{X}_p) \xrightarrow{\chi_{\text{cl}}} \Omega_c^\bullet(\mathbb{R}) \widehat{\otimes} (\widehat{\mathfrak{h}} \ltimes \widehat{\mathfrak{h}}_{\text{cont}}^\vee[1])^!\right).$$

The comparison is a field-level equivalence only after this cone is acyclic in the protected summand.

Third, the scalar anomaly mismatch is

$$\text{ob}_{\text{scal}} = \chi_{\text{anom}}([\text{anom}_{\text{prot}}]) - [\bar{c}] \in H_{\text{Lie}}^2(\bar{A}, \mathbb{C}),$$

where  $\chi_{\text{anom}}$  is the anomaly map induced by the comparison and  $[\bar{c}]$  is the scalar class used in the local theorem.

Fourth, after choosing the weighted coefficient model used by the local theorem, the brane-defect quantum obstruction is the class

$$\text{ob}_{\text{QME}} = \left[ QI_{\text{prot}} + \frac{1}{2}\{I_{\text{prot}}, I_{\text{prot}}\} + \hbar \Delta I_{\text{prot}} \right] \in H^1(\mathcal{O}_{\text{loc}}(\mathcal{E}_{\text{prot}, w}), \mathcal{Q} + \{I_{\text{o, prot}}, -\}).$$

Its vanishing is the precise statement that the protected brane-defect interaction satisfies the QME in the same normalization as Problem 5.72.

LEMMA 5.145 (*Coordinate-free objects of ProtHam*). Definition 5.143 imposes on an object  $(X, \omega, p, \Theta)$  of ProtHam coordinate-free data for  $\Theta$  and treats the Darboux expression and the bulk-to-boundary evaluation by trace substitution. An admissible protected Hamiltonian restriction  $\Theta$  at  $\widehat{p}$  is a quadruple of independent data: a sheaf of dgla brackets on  $\widehat{p}$ , a brane-defect open BV theory, a bulk-to-boundary evaluation  $\text{ev}_\Theta: \Theta \rightarrow \mathcal{A}_\partial(L)$ , and a scalar anomaly representative  $[\bar{c}]_\Theta$ ; the morphism conditions preserve each piece. Examples:

- (1)  $(X, \omega) = (T^*\Sigma, d\theta_\Sigma)$  for  $\Sigma$  a Riemann surface, evaluated at a point of the zero section, with  $\Theta$  supplied by the cotangent BV theory of  $\Sigma$ ;
- (2)  $(X, \omega)$  any K3 surface at a smooth point, with  $\Theta$  the Hochschild restriction of a curved  $\mathcal{A}_\infty$ -deformation of the structure sheaf;
- (3)  $(X, \omega)$  a holomorphic-symplectic surface with coordinate-free symplectic form,  $\Theta$  the protected Hamiltonian sector of an arbitrary Darboux choice.

The objects (1)–(3) are presented coordinate-free; the Darboux symplectomorphism, the trace-substitution evaluation, and the identification of  $[\bar{c}]_\Theta$  with  $[\bar{c}]$  are conclusions of Theorem 5.146; ProtHam membership is coordinate-free.

*Proof.* The four pieces of an admissible protected Hamiltonian restriction are listed in Definition 5.143 independent of a coordinate choice. The functorial datum  $\text{ev}_\Theta$  may be supplied by other evaluations: the cotangent example (1) admits the de Rham residue evaluation as  $\text{ev}_\Theta$ ; the Hochschild example (2) admits the Connes–Tsygan evaluation; the symplectic-surface example (3) admits any  $\omega$ -compatible polynomial evaluation that satisfies (U1)–(U5). Each receives the trace-substitution form as the conclusion of the formal Darboux reduction, the brane-line factorization restriction  $\Phi_{\text{fact}}^{\text{Ham}}$ , and the scalar-anomaly identification used in the proof of Theorem 5.146.  $\square$

THEOREM 5.146 (*Universal property of the formal-local Hamiltonian sector; corollary of the CE/PV Koszul resolution identification*). Let ProtHam denote the category whose objects are triples  $(X, \omega, p)$  with  $(X, \omega)$  a connected holomorphic-symplectic surface and  $p \in X$  a brane point, together with an admissible protected Hamiltonian sector at  $\widehat{p}$  in the sense of Definition 5.143, and whose morphisms are formal Darboux symplectomorphisms at the brane point that intertwine the protected open BV theory, the protected closed BV theory, the protected bulk-to-boundary evaluation, and the scalar anomaly class  $[\bar{c}]$ . Let  $\mathfrak{M}_{\text{loc}}^{\text{Ham}}$  denote the formal Darboux model

$$\mathfrak{M}_{\text{loc}}^{\text{Ham}} = (\widehat{\mathfrak{h}} \ltimes \widehat{\mathfrak{h}}_{\text{cont}}^\vee[1], \Omega^*(\mathbb{R}) \otimes \Lambda(\varepsilon_1, \varepsilon_2) \otimes \mathfrak{gl}_N[1], [\bar{c}]), \quad \widehat{\mathfrak{h}} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1,$$

with bulk-to-boundary evaluation  $f \mapsto \overline{\text{Tr } f(\phi_1, \phi_2)}$  on the brane line. Then  $\mathfrak{M}_{\text{loc}}^{\text{Ham}}$  is terminal in ProtHam up to canonical isomorphism: for every object  $(X, \omega, p, \Theta)$  of ProtHam there is a morphism

$$\mu_{(X, \omega, p, \Theta)}: (X, \omega, p, \Theta) \longrightarrow \mathfrak{M}_{\text{loc}}^{\text{Ham}}$$

in ProtHam, determined uniquely up to a canonical natural isomorphism. The map  $\mu_{(X, \omega, p, \Theta)}$  is the canonical formal Darboux reduction

$$\mathfrak{h}_{X, p} \xrightarrow{\sim} \widehat{\mathfrak{h}}, \quad \mathbf{Ext}_X^*(\mathcal{O}_p, \mathcal{O}_p) \xrightarrow{\sim} \Lambda^* T_p X, \quad [\bar{c}]_\Theta \mapsto [\bar{c}],$$

composed with the brane-line factorization restriction  $\Phi_{\text{fact}}^{\text{Ham}}$  of Theorem 4.11 and the protected bulk-to-boundary evaluation.

Equivalently, in the natural groupoid enrichment of  $\text{ProtHam}$  by the inner formal Darboux symmetry group

$$G_{\text{Darb}} = \text{Symp}_{\text{form}}(\widehat{\mathbb{C}^2}_o)$$

acting trivially on isomorphism classes of  $\widehat{\mathfrak{h}} \ltimes \widehat{\mathfrak{h}}_{\text{cont}}^\vee[1]$ , the contravariant functor

$$\text{ProtHam}^{\text{op}} \longrightarrow \text{Grpd}, \quad (X, \omega, p, \Theta) \longmapsto \underline{\text{Hom}}_{\text{ProtHam}}((X, \omega, p, \Theta), \mathfrak{M}_{\text{loc}}^{\text{Ham}})$$

is naturally isomorphic to the constant functor with value the one-object classifying groupoid  $BG_{\text{Darb}}$ ; the  $\pi_o$  of this groupoid is the one-point set, so on isomorphism classes of morphisms there is a unique morphism to  $\mathfrak{M}_{\text{loc}}^{\text{Ham}}$ , and inverting the inner Darboux symplectomorphisms in the homotopy category collapses the  $\text{Hom}$ -groupoid to a one-point set, naturally in isomorphism-class data.

A candidate ambient datum supplies an object of  $\text{ProtHam}$  iff its protected twist obstruction class  $\text{Ob}_{p,X}^{\text{univ}}$  of Definition 5.144 vanishes with chosen null-homotopies; a single nonzero coordinate of  $\text{Ob}_{p,X}^{\text{univ}}$  is the precise obstruction to membership in  $\text{ProtHam}$ , hence the precise obstruction to existence of the morphism  $\mu_{(X,\omega,p,\Theta)}$ .

*Proof.* Terminality of  $\mathfrak{M}_{\text{loc}}^{\text{Ham}}$  is the universal property of the CE/PV Koszul resolution of  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b$  of §5.2: any protected datum on a connected holomorphic-symplectic surface restricts to that Koszul resolution at  $p \in X^{\text{KN}}$  once the four coordinates of  $\text{Ob}_{p,X}^{\text{univ}}$  vanish, since the Koszul-resolution map is canonical on the formal-Darboux stalk and uniqueness up to inner Darboux symplectomorphism is the Darboux–Weinstein rigidity of the stalk. The detailed verification follows.

Existence of  $\mu_{(X,\omega,p,\Theta)}$  is read off the three clauses of Definition 5.143. Formal Darboux Hamiltonian reduction supplies a symplectomorphism  $\mathfrak{h}_{X,p} \xrightarrow{\sim} \widehat{\mathfrak{h}}$  of the completed closed Hamiltonian algebra and an identification  $\mathbf{Ext}_X^*(\mathcal{O}_p, \mathcal{O}_p) \xrightarrow{\sim} \Lambda^* T_p^* X$  of the point-brane Ext algebra carrying the cyclic trace induced by  $\omega_p$ ; this fixes the closed and open underlying objects of  $\mathfrak{M}_{\text{loc}}^{\text{Ham}}$ . Brane-defect locality factors the protected operator restriction through the weighted Tate field complex and the brane-line factorization map  $\Phi_{\text{fact}}^{\text{Ham}}$  of Theorem 4.11, with target  $f \mapsto \overline{\text{Tr } f}(\phi_1, \phi_2)$  by Theorem 5.41. Scalar anomaly compatibility identifies  $[\bar{c}]_\Theta$  with  $[\bar{c}]$  inside  $H_{\text{Lie}}^2(\bar{A}, \mathbb{C})$  through  $\chi_{\text{anom}}$ . Their composition is  $\mu_{(X,\omega,p,\Theta)}$ .

Uniqueness up to canonical isomorphism uses the  $G_{\text{Darb}}$ -rigidity of each clause. Suppose two morphisms  $\mu, \mu' : (X, \omega, p, \Theta) \rightarrow \mathfrak{M}_{\text{loc}}^{\text{Ham}}$  exist. Their closed-side restrictions are two formal Darboux symplectomorphisms  $\mathfrak{h}_{X,p} \rightarrow \widehat{\mathfrak{h}}$ ; these differ by an element  $g \in G_{\text{Darb}}$ , since the formal Darboux reduction at a point of  $(X, \omega)$  is unique up to inner symplectomorphism of  $\widehat{\mathbb{C}^2}_o$  (Darboux–Weinstein for formal symplectic polydisks). Their open-side restrictions  $\mathbf{Ext}_X^*(\mathcal{O}_p, \mathcal{O}_p) \rightarrow \Lambda^* T_p^* X$  differ by the same  $g$  acting on  $T_p^* X$ . Their brane-line factorization maps differ by the action of  $g$  on the boundary algebra  $\overline{\text{Tr } f}(\phi_1, \phi_2)$ , which is  $GL_N$ -invariant and preserves the connected-trace projection. In each Darboux frame the factorization map equals the unique  $J$  of Theorem 5.41; the only open-side freedom is  $g$ . Their scalar anomaly intertwiners are the identity by linearity in  $H_{\text{Lie}}^2(\bar{A}, \mathbb{C})$ . Hence  $\mu' = g \cdot \mu$  for the diagonal action of  $g$ , and the data  $g$  supplies a canonical natural isomorphism  $\mu \Rightarrow \mu'$  in  $\text{ProtHam}$ . The action of  $G_{\text{Darb}}$  on the isomorphism class of  $\mathfrak{M}_{\text{loc}}^{\text{Ham}}$  is trivial because inner symplectomorphisms of the formal polydisk preserve the isomorphism class of  $\widehat{\mathfrak{h}} \ltimes \widehat{\mathfrak{h}}_{\text{cont}}^\vee[1]$ ; they act faithfully on the structure but trivially on the isomorphism class. This is the terminality up to canonical isomorphism.

The groupoid reformulation packages this rigidity: the  $\text{Hom}$ -set  $\underline{\text{Hom}}_{\text{ProtHam}}((X, \omega, p, \Theta), \mathfrak{M}_{\text{loc}}^{\text{Ham}})$  is a torsor under the action of  $G_{\text{Darb}}$ , since any two morphisms differ by exactly one such  $g$ ; the action groupoid of a  $G_{\text{Darb}}$ -torsor is the one-object classifying groupoid  $BG_{\text{Darb}}$ , whose  $\pi_o$  is the singleton. The

functor  $(X, \omega, p, \Theta) \mapsto \underline{\text{Hom}}$  is then naturally the constant functor with value  $BG_{\text{Darb}}$ , and inverting inner Darboux symplectomorphisms in the homotopy category collapses  $BG_{\text{Darb}}$  to the one-point set: the homotopy-category Hom-set is a singleton.

The membership iff is Lemma 5.152 below: an ambient datum supplies an object of  $\text{ProtHam}$  iff each of the four classes  $\text{per}(v)$ ,  $[C_{\text{cl},p}]$ ,  $\text{ob}_{\text{scal}}$ ,  $\text{ob}_{\text{QME}}$  vanishes with chosen null-homotopies, since these classes precisely measure the obstruction to the four admissibility clauses (Darboux Hamiltonian reduction, closed-field protected comparison, scalar anomaly compatibility, brane-defect QME locality).  $\square$

**COROLLARY 5.147** (*Restriction to the formal-Darboux Koszul resolution*). Any protected datum  $(X, \omega, p, \Theta) \in \text{ProtHam}$  restricts at the Kato–Nakayama log point  $p \in X^{\text{KN}}$  to the formal-Darboux CE/PV Koszul resolution  $\Phi_{\text{coord}}$  of §5.2 of the chiral Hochschild cohomology  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b$ , with bulk-to-boundary evaluation  $u_f \mapsto J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)}$  and  $c_f \mapsto \theta_f$ , uniquely up to inner formal Darboux symplectomorphism. Equivalently,  $\text{Ob}_{p,X}^{\text{univ}}$  is the obstruction to convergence of that Koszul resolution along the ambient datum at  $(X, \omega, p)$ : a single nonzero coordinate is the precise obstruction (period, closed-field comparison cone, scalar anomaly, brane-defect QME) to extending the formal-Darboux Koszul resolution along  $\Theta$ .

*Proof.* Theorem 5.146 gives the morphism  $\mu_{(X,\omega,p,\Theta)}$  and its uniqueness up to inner Darboux symplectomorphism; §5.2 identifies the formal-Darboux stalk of  $\mathfrak{M}_{\text{loc}}^{\text{Ham}}$  with the CE/PV Koszul resolution of  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b$  under  $c_f \mapsto \theta_f$ ,  $u_f \mapsto J(f)$ . Composing supplies the corollary; the obstruction reading is Lemma 5.152 restated for the Koszul resolution.  $\square$

*The QME coordinate of  $\text{Ob}_{p,X}^{\text{univ}}$ : Casimir-balance and the queer counterexample.* The first three coordinates of  $\text{Ob}_{p,X}^{\text{univ}}$  – the period class, the closed-field comparison cone, and the scalar anomaly mismatch – are classical: they are read off the holomorphic-symplectic geometry, the protected closed sector, and the Lie cohomology of  $\bar{A}$  respectively, and their vanishing is a structural statement about the ambient datum. The fourth coordinate  $\text{ob}_{\text{QME}}$  is a quantum BV class that depends on the Chan-Paton matrix factor. The Casimir-balanced equivariant scalar QME criterion (Theorem 5.149 below) identifies an explicit locus  $\kappa_{\mathfrak{s}} = 0$  on which the scalar component of the QME obstruction vanishes identically, and exhibits  $\mathfrak{gl}(N|N)$  as an instance. The queer-centre theorem (Theorem 5.151 below) exhibits a complementary locus where unsigned parity-equivariance breaks down and a nonzero odd scalar class  $\hbar N[\bar{c}]^{\text{otr}}$  survives, showing that the algebraic spin-Hecke coefficient input governs the cohomology layer while chain-level QME cancellation requires the additional homotopy data. Together these two theorems give the precise dichotomy controlling the QME coordinate: scalar QME-vanishing is ruled by a single matrix coefficient  $\kappa_{\mathfrak{s}}$ , and a target null-homotopy of the surviving odd class is the residual ingredient for the queer brane stack.

**Definition 5.148** (*Casimir-balanced equivariant QME datum*). A Casimir-balanced equivariant brane-defect datum consists of:

- (i) a matrix Lie superalgebra  $\mathfrak{s}$  acting on the Chan-Paton factor, together with an even nondegenerate invariant form  $B_{\mathfrak{s}}$  used to build the BV heat-kernel contraction;
- (ii) a trace character  $\text{tr}_{\mathfrak{s}}$  on the matrix factor. Let  $C_{B_{\mathfrak{s}}} = \sum_{\alpha} e_{\alpha} \otimes e^{\alpha}$  be the Casimir for  $B_{\mathfrak{s}}$ , put  $K_{\mathfrak{s}} = m(C_{B_{\mathfrak{s}}})$ , and define the scalar contraction coefficient  $\kappa_{\mathfrak{s}}$  on the scalar trace line by

$$\text{tr}_{\mathfrak{s}}(AK_{\mathfrak{s}}) = \kappa_{\mathfrak{s}} \text{tr}_{\mathfrak{s}}(A).$$

The datum is Casimir-balanced when  $\kappa_{\mathfrak{s}} = 0$ ;



(iii) the weighted brane-defect local-functional complex

$$\mathfrak{Q}_s = \left( \mathcal{O}_{\text{loc}}(\mathcal{E}_{\text{bd},w}^s), d_{\text{QME}} = Q + \{I_{\text{o},s}, -\} \right);$$

(iv) a  $d_{\text{QME}}$ -stable scalar filtration and a filtered chain projection

$$\widehat{\sigma}_{\text{sc}}: \mathfrak{Q}_s \longrightarrow C_{\text{Lie}}^\bullet(\bar{A}; \mathbb{C}\gamma_1)[1]$$

whose associated graded is the scalar-symbol map  $\sigma_{\text{sc}}$ ;

(v) a  $P_{(z_1)}$ -equivariant coefficient topology on the formal symplectic polydisk, where

$$P_{(z_1)} = \{\varphi \in \text{Symp}_{\text{form}}(\widehat{\mathbb{C}^2_{\text{o}}}) : \varphi^*(z_1) \in (z_1)\}.$$

The scalar associated graded of  $\mathfrak{Q}_s$  is the Chevalley–Eilenberg complex of the reduced Hamiltonian algebra  $\bar{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1$  with coefficients in the central boundary ghost line  $\mathbb{C}\gamma_1$ .

**THEOREM 5.149** (*Casimir-balanced equivariant scalar QME vanishing*). Let  $(s, B_s, \text{tr}_s)$  be an equivariant brane-defect datum in the sense of Definition 5.148. Write  $\omega(f, g) = [\{f, g\}]_{\text{o}} \in \mathbb{C}$  for the constant Taylor coefficient of the Poisson bracket on  $\mathbb{C}[z_1, z_2]$  (Lemma 6.2). For  $a, b \in \Omega_c^0(\mathbb{R})$  and  $f, g \in \mathbb{C}[z_1, z_2]$ , the scalar part of the mixed brane-defect QME obstruction has the explicit representative

$$\text{Ob}_{\text{sc}}^s(\gamma_1; a, f; b, g) = \hbar \kappa_s \omega(f, g) \int_{\mathbb{R}} a(t)b(t)\gamma_1(t) dt \in \mathfrak{Q}_s,$$

with associated graded scalar class

$$[\text{Ob}_{\text{sc}}^s] = \hbar \kappa_s [\bar{c}] \in H_{\text{Lie}}^2(\bar{A}, \mathbb{C}\gamma_1).$$

Two consequences follow. First, the scalar QME obstruction vanishes identically,  $P_{(z_1)}$ -equivariantly, on the Casimir-balanced locus  $\kappa_s = \text{o}$ ; for the standard  $\mathfrak{gl}(N|N)$  supertrace pairing  $K_s = \text{sdim}(V)I = \text{o}$ , so  $\kappa_s = \text{o}$  and Casimir-balance holds. Second, off the Casimir-balanced locus the scalar channel carries the precise nonzero class  $\hbar \kappa_s [\bar{c}]$ , which is the exact obstruction to scalar QME-flatness of the brane-defect interaction.

If, in addition, the reduced non-scalar summand

$$\mathfrak{Q}_s^{\text{nsc}} = \text{Conc}(\widehat{\sigma}_{\text{sc}}: \mathfrak{Q}_s \longrightarrow C_{\text{Lie}}^\bullet(\bar{A}; \mathbb{C}\gamma_1)[1])[-1]$$

has zero degree-one cohomology after the chosen weighted regulator and interval-factorization null-homotopies are imposed, then the complete degree-one brane-defect QME obstruction class in  $H^1(\mathfrak{Q}_s)$  vanishes. The non-scalar vanishing hypothesis is the additional condition extending the result beyond the scalar channel.

*Proof.* The scalar contraction in the BV bracket is obtained by closing the matrix index in the single boundary loop. With the invariant form  $B_s$  this contraction is the multiplication  $K_s = m(C_{B_s})$  of the Casimir tensor, and its scalar trace-line coefficient is  $\kappa_s$ . The target part of the same local bracket is the constant Taylor coefficient cocycle

$$\omega(f, g) = [\{f, g\}]_{\text{o}}$$

of Lemma 6.2, and the brane-line locality gives the compactly supported factor  $\int_{\mathbb{R}} a(t)b(t)\gamma_1(t) dt$ . This proves the displayed representative.

Passing to the scalar associated graded keeps only the Hamiltonian cocycle  $[\bar{c}]$  and the matrix coefficient  $\kappa_{\mathfrak{s}}$ , giving the displayed cohomology class. Casimir-balance sets this coefficient to zero, so the degree-one representative is the zero cochain itself. The  $P_{(z_1)}$ -equivariance follows because  $\kappa_{\mathfrak{s}}$  is matrix-factor data and  $\omega$  is the symplectic Poisson cocycle on the target factor. The completion at the brane axis is functorial only under  $P_{(z_1)}$ , which is exactly the equivariance group in the hypothesis.

The last assertion uses the long exact sequence of cohomology for the displayed cone, which is defined only for the filtered chain projection  $\widehat{\sigma}_{\text{sc}}$  included in the datum. The scalar quotient class is zero by the first part, and the reduced degree-one cohomology is zero by hypothesis. Therefore the total degree-one obstruction class in  $H^1(\mathfrak{Q}_{\mathfrak{s}})$  vanishes.  $\square$

*Definition 5.150 (Spin-Hecke queer coefficient input).* Let  $N \geq 1$  and

$$\mathfrak{q}(N) = \left\{ \begin{pmatrix} A & B \\ B & A \end{pmatrix} : A, B \in \mathfrak{gl}_N \right\} \subset \mathfrak{gl}(N|N).$$

Put

$$J = \begin{pmatrix} \circ & I_N \\ -I_N & \circ \end{pmatrix}, \quad P^{\mathfrak{q}} = \begin{pmatrix} I_N & \circ \\ \circ & -I_N \end{pmatrix},$$

and define the queer trace, valued in the odd line  $\Pi\mathbb{C}$  (parity reversal of  $\mathbb{C}$ ;  $\Pi$  shifts even degree to odd):

$$\text{otr} \begin{pmatrix} A & B \\ B & A \end{pmatrix} = \text{Tr}(B) \in \Pi\mathbb{C}.$$

A spin-Hecke input for the queer brane is the Sergeev Hecke–Clifford action on  $V^{\otimes n}$ ,  $V = \mathbb{C}^{N|N}$ , whose central-character comparison identifies

$$\text{Tr}_{\mathfrak{gl}_N}(I_N) = \text{otr}_{\mathfrak{q}(N)}(J) = N.$$

**THEOREM 5.151** (*Queer-centre anomaly from the spin-Hecke input*). Assume the spin-Hecke input of Definition 5.150. For the queer-trace boundary evaluation  $f \mapsto \text{otr} f(\phi_1, \phi_2)$ , the odd scalar QME representative on the queer brane stack is

$$\text{Ob}_{\text{sc}}^{\text{otr}}(\gamma_{\mathfrak{i}}; a, f; b, g) = \hbar N \omega(f, g) \int_{\mathbb{R}} a(t)b(t)\gamma_{\mathfrak{i}}(t) dt \in \mathcal{O}_{\text{loc}}(\mathcal{E}_{\text{bd},w}^{\mathfrak{q}})_{\mathfrak{i}},$$

with associated graded CE class

$$\hbar N [\bar{c}]^{\text{otr}} \in H_{\text{Lie}}^2(\bar{A}, \Pi\mathbb{C})_{\mathfrak{i}}.$$

Three statements follow. First, the odd class  $\hbar N [\bar{c}]^{\text{otr}}$  is nonzero for every  $N \geq 1$  and is independent of the even class  $\hbar N [\bar{c}] \in H_{\text{Lie}}^2(\bar{A}, \mathbb{C})_{\mathfrak{o}}$ , because they sit in the two direct summands  $H_{\text{Lie}}^2(\bar{A}, \mathbb{C}) \oplus H_{\text{Lie}}^2(\bar{A}, \Pi\mathbb{C})$  of the even-odd decomposition. Second, the odd class survives over  $\mathbb{C}[[\hbar]]$  and admits a null-homotopy only after either an odd scalar quotient is imposed on the target or a target null-homotopy of the odd line  $\Pi\mathbb{C} \cdot [\bar{c}]^{\text{otr}}$  is supplied; the class itself is the precise residual obstruction. Third, the spin-Hecke input fixes the coefficient identity at the cohomology layer; chain-level QME cancellation has the parity obstruction

$$P^{\mathfrak{q}} J (P^{\mathfrak{q}})^{-1} = -J$$

obstructs the unsigned parity-equivariant regulator used in Theorem 5.149.

Consequently, the queer brane stack is a named ambient datum on which the QME coordinate  $\text{ob}_{\text{QME}}$  of  $\text{Ob}_{p,X}^{\text{univ}}$  is nonzero in the cohomology layer; admissibility of the queer datum requires either an odd-scalar quotient at the target or a chosen null-homotopy of the odd class, supplied as part of the obstruction-vanishing data of Definition 5.144.

*Proof.* The queer brane has two central coefficient lines: the even line generated by the identity and the odd line generated by  $J$ . The queer trace kills the even identity and evaluates the odd central element by

$$\text{otr}(J) = \text{Tr}(I_N) = N.$$

The spin-Hecke input asserts that this coefficient is preserved by the Sergeev central-character comparison; the Hamiltonian cocycle on the target remains the same  $\omega(f, g) = [\{f, g\}]_0$ . Therefore the scalar boundary loop gives exactly the displayed odd representative.

Passing to associated graded removes the brane-line smearing and keeps the Lie two-cocycle  $\Pi\omega$ , hence the class  $\hbar N[\bar{c}]^{\text{otr}}$ . Nontriviality is the same calculation as Lemma 6.2: a primitive  $\tilde{\eta} : \tilde{A} \rightarrow \Pi\mathbb{C}$  would lift with  $\tilde{\eta}(1) = 0$ , but then

$$\partial\tilde{\eta}(z_1, z_2) = -\tilde{\eta}(\{z_1, z_2\}) = -\tilde{\eta}(1) = 0$$

while  $\omega(z_1, z_2) = 1$ . The even and odd classes are independent because they lie in the two direct summands  $H_{\text{Lie}}^2(\tilde{A}, \mathbb{C}) \oplus H_{\text{Lie}}^2(\tilde{A}, \Pi\mathbb{C})$ . Since  $N \geq 1$  in the datum and  $\hbar$  is formal, the displayed odd class survives inside  $H_{\text{Lie}}^2(\tilde{A}, \Pi\mathbb{C})[[\hbar]]$ . Killing it requires an additional target quotient or null-homotopy beyond the spin-Hecke coefficient comparison.

Finally,  $P^q J(P^q)^{-1} = -J$  is an explicit matrix computation. Thus the queer channel has signed parity; the super-balanced cancellation uses the unsigned parity condition. The algebraic spin-Hecke input gives the coefficient class; a null-homotopy of this odd obstruction is separate data.  $\square$

**LEMMA 5.152 (Primary obstruction test).** A protected twisted-M background supplies an admissible protected Hamiltonian restriction in the sense of Definition 5.143 if and only if the four classes of Definition 5.144 vanish in the senses stated there: the period classes  $\text{per}(v)$  vanish for the Hamiltonian vector fields used by the sector, the closed-field comparison cone  $\mathcal{C}_{\text{cl},p}$  is acyclic on the protected summand, the scalar mismatch  $\text{ob}_{\text{scal}}$  is zero, and a solution of the QME class  $\text{ob}_{\text{QME}}$  has been chosen compatibly with compactly supported interval factorization.

*Proof.* The period statement is the connecting morphism of  $0 \rightarrow \mathbb{C}_X \rightarrow \mathcal{O}_X \xrightarrow{d} \Omega_{X,\text{cl}}^1 \rightarrow 0$ : local Hamiltonian primitives  $h_i$  for  $\alpha_v$  differ by constants on overlaps, and that Čech cocycle is exactly  $\partial(\alpha_v)$ ; vanishing of  $\text{per}(v) = \partial(\alpha_v) \in H^1(X, \mathbb{C}_X)$  is precisely the statement that the local primitives glue to a global Hamiltonian, modulo one global constant.

The closed-field comparison is a quasi-isomorphism of protected closed fields with Hamiltonian BF fields exactly when its mapping cone  $\mathcal{C}_{\text{cl},p} = \text{Cone}(\chi_{\text{cl}})$  is acyclic on the protected summand: a nonzero cohomology class in the cone is either a protected closed field invisible to the formal Hamiltonian model, or a Hamiltonian BF field outside the protected-sector image, and either witness blocks the field-level identification.

Scalar compatibility is the equality  $\chi_{\text{anom}}([\text{anom}_{\text{prot}}]) = [\bar{c}]$  in  $H_{\text{Lie}}^2(\tilde{A}, \mathbb{C})$ , where the comparison map  $\chi_{\text{anom}}$  is the anomaly-comparison morphism of Definition 5.144; both classes are central Lie cocycles on  $\tilde{A}$  by construction, so their equality is the assertion  $\text{ob}_{\text{scal}} = \chi_{\text{anom}}([\text{anom}_{\text{prot}}]) - [\bar{c}] = 0$ .

Finally, the displayed QME expression is the standard degree-one obstruction class for deforming a classical BV interaction to a quantum BV interaction in the fixed weighted coefficient category; vanishing of this class with chosen interval-factorization null-homotopies is exactly the brane-defect locality clause of Definition 5.143. The four vanishings therefore supply the four admissibility clauses, and conversely each admissibility clause records the vanishing of its class.  $\square$

*Definition 5.153 (Ambient-to-protected realization map).* An ambient-to-protected realization map is a morphism from the ambient twisted-M BV fields near  $\mathbb{R} \times p$  to the triple

$$(\mathcal{E}_{\text{open,prot}}, \mathcal{E}_{\text{cl,prot}}, \partial_{\text{bb}}^{\text{prot}})$$

whose closed-field cone is  $C_{\text{cl},p}$ , whose scalar anomaly map is  $\chi_{\text{anom}}$ , and whose brane interaction has QME obstruction  $\text{ob}_{\text{QME}}$ .

**PROPOSITION 5.154 (Admissible protected-sector reduction).** Let  $(X, \omega)$  be a connected holomorphic-symplectic surface and let  $p \in X$ . Suppose that a perturbative twisted-M background with branes on  $\mathbb{R} \times p$  supplies an admissible protected Hamiltonian sector at  $\hat{p}$ . Then, after the Darboux coordinate and scalar-anomaly normalizations included in the admissibility data, that restricted sector is identified with the formal local reduced Hamiltonian CE/PV brane model of §5.3.

The open side is

$$\Omega^*(\mathbb{R}) \otimes \Lambda(\varepsilon_1, \varepsilon_2) \otimes \mathfrak{gl}_N[1],$$

the closed side is

$$C_{\text{CE,cont}}^\bullet(\mathfrak{g}), \quad \mathfrak{g} = \widehat{\mathfrak{h}} \ltimes \widehat{\mathfrak{h}}_{\text{cont}}^\vee[1], \quad \widehat{\mathfrak{h}} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1,$$

and the bulk-to-boundary evaluation is

$$f \mapsto \overline{\text{Tr } f(\phi_1, \phi_2)}.$$

Hence the proved Hamiltonian, factorization, and scalar-anomaly statements of the local theorem apply to the protected sector. The all-order one-antifield quantum extension is still conditional on vanishing of the mixed brane-defect QME obstruction of Problem 5.72 for that sector.

*Proof.* The formal Darboux part of admissibility identifies the completed Hamiltonian algebra at the brane point with  $\mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$ , after choosing formal Darboux coordinates. The point-brane Ext algebra and its cyclic trace give the open coefficient data of the Dirac brane formal-stalk model.

Brane-defect locality supplies the factorization-algebraic restriction to the brane line and matches it with the local factorization map, primitive projection, and weighted Hadamard reduction already used in the local theorem. These hypotheses put the protected sector in the same algebraic category as the local model. QME-flatness for the one-antifield quantum extension is the vanishing condition  $\text{ob}_{\text{QME}} = 0$  in Theorem 5.155.

Scalar anomaly compatibility identifies the brane centre-of-mass anomaly with the class  $[\bar{e}]$  appearing in the local anomaly comparison. The three admissibility inputs therefore supply exactly the data on which the local theorem acts. Applying that theorem gives the asserted protected-sector reduction.  $\square$

**THEOREM 5.155 (Protected twist-to-Hamiltonian comparison criterion).** Let  $(X, \omega)$  be a connected holomorphic-symplectic surface and  $p \in X$ . Suppose a perturbative twisted-M background with branes on  $\mathbb{R} \times p$  supplies a candidate protected sector together with the obstruction datum of Definition 5.144. Assume:

- (i) the period classes  $\text{per}(v)$  vanish for the locally Hamiltonian symplectic vector fields used by the sector;
- (ii) the closed-field comparison cone  $C_{\text{cl},p}$  is acyclic on the protected summand;
- (iii)  $\text{ob}_{\text{scal}} = 0$ ;
- (iv)  $\text{ob}_{\text{QME}} = 0$ , with a chosen solution compatible with compactly supported interval factorization.

Then the candidate protected sector determines an admissible protected Hamiltonian restriction. Consequently the reduced brane-line Hamiltonian CE/PV, factorization-current, and scalar-anomaly statements of the formal local theorem apply to that protected sector. If, in addition, the QME solution includes the one-antifield weighted Tate lift of Problem 5.72, then the corresponding one-antifield quantum Hamiltonian comparison is valid for that protected sector and for that sector only.

*Proof.* By Lemma 5.152, the four hypotheses supply the Darboux Hamiltonian reduction, the field-level protected comparison, scalar anomaly compatibility, and brane-defect QME locality required in Definition 5.143. Applying Proposition 1.29 at the formal polydisk and then the admissible protected-sector reduction gives the local Hamiltonian BF model and its bulk-to-boundary evaluation. The last sentence is exactly the conditional weighted Tate/QME extension isolated in Problem 5.72, transported through the chosen protected-sector equivalence.  $\square$

**LEMMA 5.156** (*Formal-restriction external acceptance criterion*). An admissible protected Hamiltonian restriction contributes precisely the formal Hamiltonian polydisk, the Dirac brane Ext algebra, the brane-line factorization restriction, and the scalar anomaly class  $[\bar{c}]$ . For an external compact BCOV, MNOP/chiral-volume, CY-to-chiral, Igusa, Borchers, or BKM target  $T$ , this contribution remains only a conditional quantum input. The target theorem accepts it as matched-conventions global or cross-target descent data exactly after  $T$  supplies its convention datum  $\mathfrak{C}_T$ , a morphism  $\mathfrak{C}_{\hat{p}}^{\text{Ham}} \rightarrow \mathfrak{C}_T$ , and null-homotopies for the obstruction vector

$$\text{Ob}_T^{\text{ext}} = (\text{per}_T, \text{ob}_T^{\text{desc}}, \text{ob}_T^{\text{BCOV}}, \text{ob}_T^{\text{QME}}, \text{ob}_T^{\text{conv}}, \text{ob}_T^{\text{hyp}}, \text{ob}_T^{\text{CY-ch}}, \text{ob}_T^{\text{Igusa/BKM}}, \text{ob}_T^{6r/\text{MD}}) \in H^1(\mathfrak{R}_T),$$

with a component included exactly when  $T$  asserts the corresponding structure. A non-zero component is the exact obstruction to the corresponding external theorem.

*Proof.* The three admissibility clauses record Darboux Hamiltonians at  $\widehat{X}_p$ , the point-brane Ext algebra, brane-defect locality on  $\mathbb{R} \times p$ , and scalar anomaly compatibility. The global data of any external theorem — compact Calabi–Yau threefold, negative-cyclic Calabi–Yau class, the compact CY-to-chiral target specialization datum  $\Phi_d = \text{Sp}_{\Sigma_{d-1}, C} \circ \Phi_d^{\text{FA}}$ , compact worldsheet or  $S^3$  framing, BCOV propagator and holomorphic anomaly equation, Borchers input Jacobi form, Weyl chamber, compact Hall orientation — enter as independent coordinates of  $\text{Ob}_T^{\text{ext}}$  in the target convention complex  $\mathfrak{R}_T$ . Two ambient theories with the same admissible formal restriction differ exactly at these coordinates. Vanishing with specified null-homotopies is the acceptance datum for using the formal restriction in  $T$ ; a non-zero coordinate is the named target obstruction.  $\square$

**Definition 5.157** (*Protected-sector matched-conventions contribution*). The protected-sector reduction contributes the following datum to any external matched-conventions theorem:

$$\mathfrak{C}_{\hat{p}}^{\text{Ham}} = (d = 2, \widehat{X}_p, \omega_p, \mathbb{R}_{\text{brane}}, \widehat{\mathfrak{h}} = \mathbb{C}[[z_1, z_2]]/\mathbb{C}, \mathbf{Ext}_X^*(\mathcal{O}_p, \mathcal{O}_p), [\bar{c}], \Phi_{\text{fact}}^{\text{Ham}}).$$

The remaining datum for a compact BCOV, CY-to-chiral, or Igusa/BKM theorem consists of the compact target, its Calabi–Yau or modular class, the worldsheet or specialization curve, the framing, the global propagator or Hall orientation, and the target obstruction complex.

**COROLLARY 5.158** (*Formal uniformity after admissibility*). For every connected holomorphic-symplectic surface  $(X, \omega)$ , every brane point  $p \in X$ , and every protected twisted-M realization whose obstruction datum vanishes, the restricted sector is identified with the same formal Hamiltonian brane model

$$(\widehat{\mathfrak{h}} \ltimes \widehat{\mathfrak{h}}_{\text{cont}}^{\vee}[1], \Omega^*(\mathbb{R}) \otimes \Lambda(\varepsilon_1, \varepsilon_2) \otimes \mathfrak{gl}_N[1], [\bar{c}])$$

after choosing a formal Darboux coordinate at  $p$ .

*Proof.* The period vanishing gives global Hamiltonian primitives for the vector fields used in the sector. The acyclic closed comparison cone identifies the protected closed fields with  $\widehat{\mathfrak{h}} \ltimes \widehat{\mathfrak{h}}_{\text{cont}}^{\vee}[1]$ . The Ext-algebra identification gives the Dirac brane formal-stalk chart, and scalar compatibility fixes the same class  $[\bar{c}]$ . These are exactly the inputs of Proposition 1.29.  $\square$

*Remark 5.159* (*Dirac language inside the protected sector*). For an admissible protected Hamiltonian sector, the Dirac-thesis language — the brane as the constrained formal-stalk chart whose stable scalar-reduced measurement is the unique  $GL_N$ -invariant Darboux-natural extension of the moment-map action on linear generators (Theorem 5.146) — is a shorthand for the already identified formal local model. The Dirac brane stack measures the formal symplectic normal polydisk through trace; Dirac reduction gives the open trace algebra; the closed Hamiltonian BF theory is the shifted cotangent theory of canonical transformations of that polydisk; and the closed observables identify with derived Poisson-central operations on the stable open Hamiltonian brane algebra only in the reduced local sense proved by Theorem 5.6. The scalar anomaly is the same class  $[\bar{c}]$  on the closed and open sides. The corresponding ambient centrality statement is the matched-conventions interface of Corollary 5.160.

**COROLLARY 5.160** (*Protected-sector matched-conventions acceptance criterion*). A theorem in a compact BCOV, CY-to-chiral, or Igusa/BKM target may use the protected-sector reduction only as a matched-conventions descent input: it must give a morphism  $\mathfrak{C}_{\widehat{p}}^{\text{Ham}} \rightarrow \mathfrak{C}_T$  and null-homotopies for every asserted component of  $\text{Ob}_T^{\text{ext}}$ .

*Proof.* Lemma 5.156 identifies the external acceptance class  $\text{Ob}_T^{\text{ext}} \in H^1(\mathfrak{R}_T)$ . A morphism of convention data and null-homotopies for this class supply exactly the compact, worldsheet, modular, orientation, and descent coordinates needed by the target theorem. Corollary 5.158 then supplies the  $d = 2$  formal-disk normalization functorially in the chosen Darboux chart.  $\square$

*Matched-conventions component*  $\mathfrak{o}_{\text{cv}}$ . The local convention base  $\mathfrak{C}_{\text{loc}}^{(2)}$  of Definition 10.5 is the formal holomorphic dimension  $d = 2$  brane-line stalk; compact  $\text{CY}_3$  BCOV, the compact CY-to-chiral frontier, and the Borchers / Borchers–Kac–Moody (BKM)–Igusa bridge supply their own convention, descent, and modular data. By Lemma 10.6 the formal-disk theorem contributes that stalk to a target  $\mathfrak{T}$  only after the target supplies a matched-conventions datum  $\mathfrak{C}_{\text{tar}}$  of Definition 10.7 and a null-homotopy of the universal Koszul-descent acceptance class  $\text{Ob}_{\text{UKD}}(\mathfrak{T}) = (\text{Ob}_{\mathfrak{C}_{\text{tar}}}, \text{Ob}_{\text{glob}}, \text{ob}_{6r}, \text{ob}_{\text{MD}})$ . The component  $\mathfrak{o}_{\text{cv}}$  is therefore

$$\mathfrak{o}_{\text{cv}} = \text{Ob}_{\text{UKD}}(\mathfrak{C}_{\text{tar}}) \in H^1(\mathcal{K}_{\text{obs}}) \oplus H^1(\mathfrak{R}_{\text{KD}}),$$

with named subclasses the period class  $\text{per}_X(v) = \delta(-\iota_v \omega) \in H^1(X, \mathbb{C}_X)$  of Definition 10.8, the Weiss/Ran descent class of Definition 10.9, the scalar anomaly compatibility class  $\text{ob}_{\text{anom}}$ , the target QME class  $\text{ob}_{\text{QME}}$ , the convergence and hyperdescent classes  $\text{ob}_{\text{conv}}$ ,  $\text{ob}_{\text{hyp}}$ , the CY-to-chiral/Igusa/BKM matched-conventions classes  $\text{ob}_{\text{CY-ch}}$ ,  $\text{ob}_{\text{Igusa/BKM}}$ , the six-relation factorization class  $\text{ob}_{6r}$ , and the mixed-Dunn class  $\text{ob}_{\text{MD}}$ , together with the curvature  $[d\Theta_{\text{KD}} + \frac{1}{2}[\Theta_{\text{KD}}, \Theta_{\text{KD}}]]$ . The construction problem inscribed by  $\mathfrak{o}_{\text{cv}}$  is joint vanishing with chosen null-homotopies, equivalently the hypotheses of Theorem 10.13; under that vanishing the target inherits the formal-stalk component  $\chi_{\text{op}}^{-1} \Phi_{\text{fact}}^{\text{Ham}} \chi_{\text{cl}}$  of 10.6. Convention divergence is load-bearing and reported explicitly;  $\mathfrak{o}_{\text{cv}}$  records exactly the structures captured by  $\mathfrak{C}_{\text{tar}}$ . The full firewall theorem 10.4, the matched-conventions criterion 10.13, and the per-target divergence theorems for compact-CY, BCOV, the compact CY-to-chiral frontier, the Borceherds/BKM–Igusa bridge, and the Gromov–Witten (GW) / Donaldson–Thomas (DT) / Pandharipande–Thomas (PT) / Maulik–Nekrasov–Okounkov–Pandharipande (MNOP) lane are stated in the appendix (§10.7); their statements are deferred to keep the local theorem from being read as a global claim.

*Remark 5.161 (Stratified Weiss/Cech–Roos descent obstruction).* Inside the local stratified construction the descent component of  $\mathfrak{o}_{\text{cv}}$  is sharpened by the Cech cone

$$\Delta_{V, \mathcal{U}}^{N, M} = \text{Cone}\left(\mathcal{F}_{\Omega, N, M}^{\text{str}}(V) \rightarrow \text{Tot } \check{C}^\bullet(\mathcal{U}; \mathcal{F}_{\Omega, N, M}^{\text{str}})\right)$$

for collared stratified Weiss covers  $\mathcal{U}$  of  $V$ , together with the Roos compatibility class  $\mathbf{w}_{V, \Omega, N, M}^{\check{C}/R}$  under refinements, finite windows, interval inclusions, and admissible weight transports. The internal descent entry is

$$\delta_{\text{Weiss}}^{\check{C}/R} = \left(\{H^\bullet(\Delta_{V, \mathcal{U}}^{N, M})\}_{V, \mathcal{U}, N, M}, \mathbf{w}_{V, \Omega, N, M}^{\check{C}/R}\right),$$

the datum used by the local factorization-center morphism. Choosing Darboux coordinates pointwise and gluing the resulting stalk maps produces a global morphism after this descent datum is made acyclic; the displayed cone class is the obstruction.

**THEOREM 5.162** (*Iff-vanishing of the Tate-coefficient cotangent lift obstruction vector*). Let  $\text{Ob}_T$  be the obstruction vector of Definition 7.3.

- *Desired upgrade.* An unreduced Tate-coefficient cotangent factorization-centre morphism

$$\Phi_{\text{cl}}: \text{Obs}_{\text{closed}, H}^{\text{cl}}|_L \longrightarrow Z_{\text{fact}}^{P_0}(\text{Obs}_{\text{open}, N}^{\text{cl}})$$

with all  $P_0$ -centrality homotopies, factorization centrality homotopies, stable connected quotient, and a boundary realization of the shifted cotangent generators in the sense of Problem 5.48.

- *Minimal input data.* The reduced bulk-to-brane evaluation  $J: \tilde{A} \rightarrow A_{\tilde{\partial}}^{\text{Ham}}$  of Theorem 5.146; the brane line  $L$  and gauge group  $\mathfrak{gl}_N$ ; a coefficient category together with chosen compatible null-homotopies for each component class.
- *Obstruction class.* The eight-component vector  $\text{Ob}_T = (\mathfrak{o}_{\text{Cas}}, \mathfrak{o}_{\text{nilp}}, \mathfrak{o}_{\text{Quil}}, \mathfrak{o}_{\text{BV, end}}, \mathfrak{o}_{\text{prim}}, \mathfrak{o}_{\text{Had}}, \mathfrak{o}_{\text{univ}}, \mathfrak{o}_{\text{cv}})$  of Definition 7.3.
- *Vanishing criterion.*  $\Phi_{\text{cl}}$  exists iff  $\text{Ob}_T = 0$  with chosen compatible null-homotopies. The all-order one-antifield quantum extension exists exactly under additionally  $\mathfrak{o}_{\text{QME}} = 0$ . The compact or cross-target version exists exactly under additionally  $\mathfrak{o}_{\text{cv}} = 0$ .
- *First nonzero example.* The Casimir kernel  $\mathfrak{o}_{\text{Cas}}$  is nonzero on the strict product/direct-sum Tate pair  $(\mathfrak{h}_{\Pi}, D_{\oplus})$ : the compatible family  $(C_{\mathfrak{h}}^M)_M$  lies in the cokernel of the represented-kernel map into  $\mathfrak{h} \widehat{\otimes} D$

(Lemma 5.211, Theorem 5.93). This is the canonical construction defect blocking the strict-unweighted endpoint and forcing the weighted Tate completion of Definition 5.54.

- *Physical meaning.* The matched-bracket open-closed identity that holds at degree zero (Theorem 4.11) lifts to the cotangent direction after installing the regulator data for the strict ambient product completion; the residue cotangent module  $\mathcal{P} = \mathfrak{h}_{\text{cont}}^{\vee}$  is genuinely Tate, with finite-window approximations, and the diagonal Casimir  $C_{\mathfrak{h}}$  converges in  $\mathfrak{h} \widehat{\otimes} \mathcal{P}$  precisely after the regulator is installed. Vanishing of  $\text{Ob}_T$  names exactly the regulator data — weight system, propagator extension, truncation, primitive projection, parametrix expansion, protected anomaly cancellation, and matched-conventions descent — under which the unreduced lift is constructed.

*Proof.* The classical assertion assembles the component criteria. Component  $\mathfrak{o}_{\text{Cas}} = 0$  is the weighted Casimir kernel of 5.61 together with the heat-kernel propagator extension of 5.63, giving the classical cochain CE/PV cotangent extension 5.70 (d). Component  $\mathfrak{o}_{\text{nilp}} = 0$  is the joint vanishing of the linear, semisimple, and low-degree Casimir terms of Lemma 5.106, closing the truncation tower. Component  $\mathfrak{o}_{\text{Quil}} = 0$  is the universal Koszul-descent vanishing of 5.108 in  $\mathcal{O}_{\text{Kos}}$ ; inside the admissible Tate model envelope of 5.17, this gives the Tate Quillen upgrade of the cochain identification. Component  $\mathfrak{o}_{\text{BV, end}} = 0$  supplies the endpoint-admissible coefficient module of 5.94, and component  $\mathfrak{o}_{\text{prim}} = 0$  supplies the strict cotangent centrality homotopy and the cotangent-label boundary realization. Component  $\mathfrak{o}_{\text{Had}} = 0$  is the Hadamard endpoint vanishing of 8.19. Component  $\mathfrak{o}_{\text{univ}} = 0$  supplies the protected Hamiltonian comparison through Theorem 5.149 and identifies the formal-local model with its protected restriction. Together these supply the cochain-level CE/PV map, the Tate Quillen envelope, the analytic kernel, the descent datum, and the protected realization, which is exactly Problem 5.48. Conversely, an existing  $\Phi_{\text{cl}}$  with the listed homotopies exhibits each component class through its definition: the Casimir kernel through the convergent BV pairing, the truncation classes through the finite  $\mathfrak{m}^3$ -adic cotangent, the universal Koszul cone through the bar-cobar comparison, the BV endpoint through the coefficient and RG transport, the primitive class through the factorization projection, the Hadamard class through the parametrix expansion of the propagator, the protected class through the comparison morphisms, and the cross-target class through the matched-conventions descent.

The quantum and cross-target statements are Proposition 5.71 and Theorem 10.13, respectively.  $\mathfrak{o}_{\text{QME}} = 0$  supplies the all-order one-antifield extension of the degree-zero Moyal/Weyl theorem inside the weighted regulator-admissible sector;  $\mathfrak{o}_{\text{cv}} = 0$  with the matched-conventions datum supplies the global or cross-target factorization-centre morphism whose formal stalk is the local Hamiltonian map.  $\square$

**COROLLARY 5.163** (*Proved sublattice and residual*). The proved sublattice of  $\text{Ob}_T$  consists of: the weighted-completion classical lift (5.70 (a)–(d) inside the regulator-admissible sector); the classical  $\mathfrak{m}^3$  truncation (5.101); the chain-level primitive projection with strict  $Q$ -, factorization-, and  $P_{\circ}$ -bracket compatibility (5.116, 5.118, 5.119); the Hadamard finite-window parametrix (8.17); the scalar Casimir-balanced equivariant QME criterion (5.149); and the conditional matched-conventions transfer of 10.13. The residual obstruction is

$$\text{Ob}_T^{\text{res}} = (\mathfrak{o}_{\text{QME}}^{\text{ns}}, \mathfrak{o}_{\text{nilp}}^{\text{low}}, \mathfrak{o}_{\text{Quil}}^{\text{univ}}, \mathfrak{o}_{\text{prim}}^{\text{cot}}, \mathfrak{o}_{\text{univ}}^{\text{prot}}, \mathfrak{o}_{\text{cv}}^{\text{glob}}),$$

with components: the non-scalar mixed brane-defect QME class  $[(\mathfrak{o}_{n, \mathcal{M}}^{\text{ns}})_{\mathcal{M}}] \in \varprojlim_{\mathcal{M}} H^1(\mathcal{K}_{\text{ns}, \mathcal{M}}^{\bullet}(I))$  together with its secondary  $\varprojlim^1 H^0$  primitive class (5.83 and 5.67); the low-degree truncation triple



$(\mathfrak{o}_N^{\text{lin}}, \mathfrak{o}_{\leq 2}^{\text{ss}}, \mathfrak{o}_{\leq 2}^{\text{Cas}})$  of §5.105; the universal Koszul-descent class in  $\mathcal{O}_{\text{Kos}}$  of §5.22; the cotangent centrality homotopy and one-antifield boundary realization residue inscribed by  $\mathfrak{o}_{\text{prim}}$ ; the protected twist-obstruction tuple of §5.144; and the matched-conventions acceptance class  $\text{Ob}_{\text{UKD}}(\mathfrak{C}_{\text{tar}})$  of 10.13. Each is a precise construction problem with explicit missing datum.

*Proof.* The proved sublattice is the disjoint union of the cited theorems; each supplies the corresponding component. The residual is the complement, with construction problem inscribed by the corresponding paragraph above. Iff-vanishing is Theorem 5.162.  $\square$

### 5.7 Stratified CE/PV and admissible Koszul duality

The closed local operator algebra  $C_{\text{CE}, \text{cont}}^\bullet(\mathfrak{g})$  of §6.2 and the open brane Hamiltonian descendant algebra  $\widehat{\mathbf{S}}(\bar{A} \oplus \bar{A}_{\text{desc}}[1])$  of §3.5 are two  $E_1$ -algebras built from the same Lie/coComm operadic pair, on different coefficient categories. Koszul duality (Priddy 1970, [23]; Loday–Vallette 2012, [24]) exchanges commutative cochains for associative enveloping algebras whenever its convergence hypotheses are met. On the strict formal-polydisk product/direct-sum pair those hypotheses have one obstruction class — the Tate Casimir  $\mathfrak{o}_{\text{Cas}}$ ; on the weighted Tate replacement of §7.5 they are met. The stratification below names what is built at each coefficient layer and the precise obstruction class at each boundary.

- (i) *Operadic input.* The closed BV side has dg Lie input  $\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1]$ ; the open Hamiltonian side has coCommutative comodule input on  $\bar{A} \oplus \bar{A}_{\text{desc}}[1]$ , namely brane traces and primitive Koszul descendants. The Lie/coComm operadic pair is the precondition of Koszul duality. The cotangent-dual CE coordinate  $u_f$  is the CE coordinate whose CE/PV image is the boundary observable  $O_f$  (Theorem 4.20); the Hamiltonian-dual coordinate  $c_f$  records the Schouten action on observables.
- (ii) *Admissible Tate CE/bar-cobar.* The cochain layer is the completed cotangent CE/PV identity of Theorem 5.5: valid for the formal polydisk on finite coordinate tests, with a kernel-admissible Maurer–Cartan tensor appearing only after the diagonal converges. The enveloping layer is a bar-cobar criterion that applies precisely when the chosen Tate or weighted coefficient category makes the CE coalgebra conilpotent and the filtered inverse limits exact. The strict product/direct-sum full-polydisk coefficient pair  $(\prod_d H_d) \oplus (\bigoplus_d H_d^\vee)$  has the bar-cobar obstruction (Lemma 5.211); the obstruction tuple  $\mathfrak{o}_{\text{Cas}}$  is repaired by weighted Tate completion (Theorem 5.70) or by nilpotent truncation, never discarded. The minimal RG-stable analytic completion forced by  $J$  — closed under the Schouten bracket and pointwise product, with heat-kernel propagator extension and Casimir convergence — is the weighted-Tate completion  $B_w$  (Theorem 5.57).
- (iii) *Tate Quillen equivalence (Beilinson–Drinfeld 2004, [28]).* Under admissible lower-central Tate-pronilpotence the completed Lie bar-cobar map  $\Omega C_*^{\text{CE}}(\mathfrak{g}_{\text{adm}}) \rightarrow \widehat{U}(\mathfrak{g}_{\text{adm}})$  is a filtered quasi-isomorphism in **TateVec** (Lemma 5.165, Proposition 5.168). The lower-central Tate-pronilpotence hypothesis is an extra input on the strict full polydisk; the weighted completion or nilpotent truncation supplies it. The strict Beilinson–Feigin–Mazur [22] product pair carries the same obstruction.
- (iv) *Coordinate CE/PV reduced  $P_0$  central operation.* On the coordinate window the CE/PV identity matches Hamiltonian-dual CE coordinates to Schouten polyvector coordinates, and shifted-cotangent-dual CE coordinates to boundary Hamiltonian functions on the Chan–Paton pair

- (Theorem 4.20). This is a cochain-level reduced morphism. A strict factorisation  $P_0$ -centre morphism on the unreduced cotangent side requires the unreduced lift obstruction Remark 5.189, recorded in Problem 5.48.
- (v) *PBW/Rees special fibre.* The Rees enveloping algebra  $U_{\hbar}(\mathfrak{g})$  has PBW special fibre at  $\hbar = 0$  isomorphic to  $\widehat{\mathbf{S}}(\mathfrak{g})$ , supplying the label-level identification of the brane Hamiltonian descendants  $\Psi_{a,b}$  with the abstract bidegree  $\tilde{A}_{\text{desc}}$  (Corollary 3.29). The symmetric PBW action on  $\Psi_{a,b}$  and the coadjoint action on the cotangent module  $\mathcal{P}$  are distinct (Theorem 5.180, Corollary 5.172, Theorem 8.1); the two formulas have opposite shift directions and differ by  $-(pb - qa + p - q)$  versus  $(pb - qa)$ . The PBW special fibre and the coadjoint module have distinct Hamiltonian actions.
  - (vi) *Matlis/Grothendieck cotangent module.* The continuous dual  $\mathfrak{h}_{\text{cont}}^{\vee}$  is the Grothendieck–Matlis principal-part module  $\mathcal{P}$ , the residue annihilator of the constants quotient (Matlis 1958, [19]; Hartshorne local cohomology, [20]; Theorem 5.178). The cotangent module  $\mathcal{P}$  is the residue principal-part module, distinguished from the polynomial open BV/Koszul matrix model (Proposition 5.181, Theorem 5.182) by the residue-coadjoint pole-raising action: linear Hamiltonians act locally nilpotently on polynomial observables and pole-raisingly on residues.
  - (vii) *Brane-line factorisation-current comparison.* Hamiltonians smeared on a brane interval lift to factorisation currents whose binary collisions reproduce the Hamiltonian Poisson bracket and the residue-coadjoint action on  $\mathcal{P}$  (Theorem 4.11, Theorem 5.185); de Rham-current contraction along the brane line returns the open trace algebra (Theorem 5.190); the Moyal quantum coefficient comparison is the componentwise Theorem 5.217. Each remaining obstruction is a finite-window obstruction theorem: the strict all-window Bochner–Martinelli transfer is obstructed by  $\text{Ob}_{\text{BM}}^{\Pi}$ , with the strict obstruction proved for every  $N$  by the Hamiltonian-flow leak  $z_1^{N+2} \cdot \rho_{N+1,0} = -(N+2)\rho_{0,1}$  (Theorem 8.40, Corollary 8.41); the unreduced factorisation-centre lift by the four-component obstruction vector of Theorem E.53 and Problem 5.48; the non-scalar Costello QME by Problem 5.219, with the smallest non-scalar window  $\theta_3$  carrying both the bare value  $H^1 = \mathbb{C} \cdot E_{\theta_3} \neq 0$  and the supplied-counterterm value  $H^1 = 0$  (Theorem E.38); the all-bidegree decorated PBW Stokes identity by  $\Omega_{a,b}^{\text{rad}}$ , with the radial bicomplex interior-acyclic conditional on the radial-parts statement (Theorem F.28).

*Definition 5.164.* For an augmented complete dg algebra  $B$ , its Koszul dual is

$$B^! = \text{RHom}_B(\mathbb{C}, \mathbb{C})$$

[23]. Commutative Chevalley–Eilenberg cochains are regarded as  $E_1$  algebras by restriction. All bar and cobar constructions below are taken in the category of complete filtered Tate vector spaces, with continuous maps. The completions are by symmetric or tensor length together with the coefficient topology coming from Taylor degree. For a Tate space  $V$ , the notation  $V_{\text{cont}}^{\vee}$  means the continuous dual with its dual Tate topology,  $\widehat{\otimes}$  means the completed Tate tensor product, and

$$\widehat{\mathbf{S}}^n(V) = (V^{\widehat{\otimes} n})_{\mathbb{S}_n}.$$

Chevalley–Eilenberg chains below are direct sums in CE length with completed coefficients in each length; CE cochains, completed cobar algebras, and completed enveloping algebras are products in the corresponding length filtration.

LEMMA 5.165 (*Tate CE chains, completed PBW, and Koszul duality*). Work in the symmetric monoidal category **TateVec** of complete filtered Tate  $\mathbb{C}$ -vector spaces:

$$V = \varprojlim_{w \in \mathbb{N}} V_{\leq w}, \quad V_{\leq w} \text{ finite-dimensional}, \quad F^w V = \mathbf{Ker}(V \rightarrow V_{\leq w-1}),$$

with  $\widehat{\otimes}$  the completed tensor product  $V \widehat{\otimes} W = \varprojlim_w (V_{\leq w} \otimes W_{\leq w})$  and  $V_{\text{cont}}^\vee = \varinjlim_w V_{\leq w}^*$  the continuous dual. Admissible quasi-isomorphisms are the filtered ones: chain maps inducing a quasi-isomorphism on every quotient  $V/F^w V$  and on the associated graded.

Let  $\mathfrak{g}$  be a Tate graded Lie algebra whose filtration is by closed dg Lie ideals, so that every  $\mathfrak{g}_{\leq w}$  used below is a finite-dimensional dg Lie quotient. Assume  $\mathfrak{g}$  is lower-central Tate-pronilpotent in the sense that the descending Lie powers  $\mathfrak{g}^{(\geq m)} = [\mathfrak{g}, \mathfrak{g}^{(\geq m-1)}]$  satisfy  $\bigcap_m (\mathfrak{g}^{(\geq m)} + F^w \mathfrak{g}) = F^w \mathfrak{g}$  for every  $w$ . Then:

- (i) For  $V = \prod_d V_d$  with all  $V_d$  finite-dimensional and  $W = \bigoplus_d W_d$  with all  $W_d$  finite-dimensional,

$$(V \oplus W)_{\text{cont}}^\vee = \bigoplus_d V_d^\vee \oplus \prod_d W_d^\vee,$$

and the dualization  $V \mapsto V_{\text{cont}}^\vee$  is exact and continuous in the filtration  $F^w$ .

- (ii) The continuous Chevalley–Eilenberg cochains

$$C_{\text{CE,cont}}^\bullet(\mathfrak{g}) = \text{Hom}_{\text{cont}}(\widehat{\mathbf{S}}^c(\mathfrak{g}[1]), \mathbb{C})$$

identify with  $\widehat{\mathbf{S}}_{\text{Tate}}(\mathfrak{g}_{\text{cont}}^\vee[-1])$ . The CE coalgebra  $C_*^{\text{CE}}(\mathfrak{g}) = \widehat{\mathbf{S}}^c(\mathfrak{g}[1])$  is conilpotent in CE length: every homogeneous CE chain has finite symmetric length and is killed by sufficiently high iterated reduced coproduct. Lower-central Tate-pronilpotence is a separate inverse-limit/bar-cobar convergence hypothesis, used below for the completed cobar and PBW comparison.

- (iii) The continuous bar resolution of the augmentation  $C_{\text{CE,cont}}^\bullet(\mathfrak{g}) \rightarrow \mathbb{C}$  has dual cobar algebra  $\Omega C_*^{\text{CE}}(\mathfrak{g})$ , and the canonical Lie bar–cobar map

$$\Omega C_*^{\text{CE}}(\mathfrak{g}) \longrightarrow \widehat{U}(\mathfrak{g})$$

is an admissible (filtered) quasi-isomorphism in **TateVec**.

- (iv) The PBW filtration on  $\widehat{U}(\mathfrak{g})$  by tensor length is complete and separated. Its associated graded is the completed symmetric algebra

$$\text{gr}_F^\bullet \widehat{U}(\mathfrak{g}) \cong \widehat{\mathbf{S}}_{\text{Tate}}(\mathfrak{g}).$$

*Proof.* Item (1). A continuous functional on  $V = \prod V_d$  depends on finitely many coordinates: the topology on  $\mathbb{C}$  is discrete, so the kernel of any continuous map contains some  $F^w V$ . Therefore  $V_{\text{cont}}^\vee = \bigoplus_d V_d^*$ . Conversely, a continuous functional on the discrete sum  $W = \bigoplus W_d$  is given by an arbitrary product of degree-wise functionals, hence  $W_{\text{cont}}^\vee = \prod_d W_d^*$ . Direct sum of these gives the displayed formula. Exactness of dualization in **TateVec** is preservation of finite limits; continuity in the Tate filtration is direct.

Item (2). The completed CE coalgebra  $\widehat{\mathbf{S}}^c(\mathfrak{g}[1]) = \bigoplus_n \widehat{\mathbf{S}}^n(\mathfrak{g}[1])$  has coaugmentation coideal  $\widehat{\mathbf{S}}^{\geq 1}(\mathfrak{g}[1])$ . An element in CE length  $n$  is killed by the  $(n+1)$ -fold reduced coproduct; this is the conilpotence used in the coalgebra. Passage to completed cobar algebras and PBW enveloping algebras

after the Tate inverse limit uses the lower-central Tate-pronilpotence hypothesis: modulo every finite filtration step the descending Lie powers have empty intersection, so the completed bar-cobar spectral sequence is separated. The displayed equality identifies cochains as continuous functionals on the CE coalgebra, using item (1) windowwise.

*Item (3).* Filter the CE coalgebra by symmetric coalgebra length and by the supplied Tate quotient system. Project to a fixed bracket-compatible finite dg Lie quotient  $\mathfrak{g}_{\leq w}$  and bound symmetric/tensor length to a fixed  $n$ . The resulting finite-dimensional diagram is the ordinary Lie bar–cobar comparison [24]: the bar resolution computes  $\mathrm{RHom}_{C_{\mathrm{CE}}^\bullet(\mathfrak{g}_{\leq w}, \leq n)}(\mathbb{C}, \mathbb{C})$ , and the cobar algebra maps to the universal enveloping algebra of the same finite Lie quotient. These finite comparisons are compatible with enlargement of the quotient and increase of the length cap because the transition maps are Lie-algebra quotient maps. Passing to the Tate limit and completing in tensor length gives the asserted filtered quasi-isomorphism in **TateVec**. The PBW filtration is, quotientwise, the standard PBW filtration on  $U(\mathfrak{g}_{\leq w})$ , hence preserved under inverse limits. Arbitrary Taylor-degree coordinate windows of the full formal Hamiltonian polydisk lie outside this hypothesis and outside this lemma.

*Item (4).* Tensor length is a separated, descending filtration on  $\widehat{U}(\mathfrak{g})$  because the Lie relations  $x y - (-1)^{|x||y|} y x - [x, y] = 0$  have leading tensor-length term  $x y - (-1)^{|x||y|} y x$  of length 2 and lower-order term  $-[x, y]$  of length 1; so the associated graded retains only the symmetric (length-fixed) part. Completeness is clear from the Tate construction. The associated graded is therefore the completed symmetric algebra  $\widehat{\mathbf{S}}_{\mathrm{Tate}}(\mathfrak{g})$ .  $\square$

*Remark 5.166 (Bar-cobar obstruction datum).* The lower-central Tate-pronilpotence hypothesis in item (2) is the exact bar-cobar criterion used here. It is satisfied by the nilpotent truncations and by the weighted admissible replacements used later. The full reduced formal Hamiltonian algebra with linear modes is already covered at cochain level by the pro coordinate-window/mixed-continuous CE/PV calculation; its enveloping/Koszul promotion is governed by the obstruction terms of Remark 4.16 and Theorem 5.5. The Beilinson–Feigin–Mazur explicit product/direct-sum Tate convention [22] is the special case  $\mathfrak{g} = (\prod P_d) \oplus (\oplus Q_d)$  with the bracket bilinear in the two displayed pieces.

**LEMMA 5.167 (Linear Poisson Schouten differential).** Let  $\mathfrak{h}$  be a pro-finite-dimensional Lie algebra in **TateVec**, and choose a mixed or weighted completion on which the linear Kirillov–Kostant–Souriau bracket is continuous. The bracket is  $\{O_I, O_J\}_\partial = O_{[H_I, H_J]} = C_{IJ}^K O_K$  on a chosen homogeneous basis  $\{H_I\}$  of  $\mathfrak{h}$ , where  $O_I \in \mathfrak{h}^*$  is the linear coordinate. The associated Poisson bivector is the linear bivector

$$\pi_\partial = \tfrac{1}{2} C_{IJ}^K O_K \theta^I \wedge \theta^J, \quad \theta^I = \partial / \partial O_I \in \mathrm{Der}(\widehat{\mathbf{S}}_{\mathrm{Tate}}(\mathfrak{h})).$$

Assume that this linear bivector and the Schouten contraction bracket are continuous on the chosen mixed or weighted completion. Then the Lichnerowicz/Schouten differential  $d_\pi = [\pi_\partial, -]_S$  on

$$\mathrm{PV}(\widehat{\mathbf{S}}_{\mathrm{Tate}}(\mathfrak{h})) = \widehat{\mathbf{S}}_{\mathrm{Tate}}(\mathfrak{h}) \widehat{\otimes} \widehat{\Lambda}(\theta^I)$$

acts on generators by

$$d_\pi \theta^K = -\tfrac{1}{2} C_{IJ}^K \theta^I \theta^J, \quad d_\pi O_K = C_{KJ}^L O_L \theta^J.$$

*Proof.* Direct calculation. The Schouten Leibniz rule with  $[\theta^I, O_J]_S = \delta_J^I$ ,  $[\theta^I, \theta^J]_S = 0$ ,  $[O_I, O_J]_S = 0$  gives

$$\begin{aligned} [\pi_\partial, O_K]_S &= \tfrac{1}{2} C_{IJ}^L O_L [\theta^I \theta^J, O_K]_S \\ &= \tfrac{1}{2} C_{IJ}^L O_L (\delta_K^I \theta^J - \delta_K^J \theta^I) \\ &= \tfrac{1}{2} (C_{KJ}^L - C_{JK}^L) O_L \theta^I = C_{KJ}^L O_L \theta^J, \end{aligned}$$

using the antisymmetry  $C_{IJ}^L = -C_{JI}^L$  in the lower indices. For  $[\pi_\partial, \theta^K]_S$ , the Schouten bracket with  $\theta^K$  uses  $[\theta^I \theta^J, \theta^K]_S = 0$  on the bivector factor and  $[O_L, \theta^K]_S = -\delta_L^K$ ; the result is  $-\tfrac{1}{2} C_{IJ}^K \theta^I \theta^J$ .  $\square$

**PROPOSITION 5.168** (*Admissible Tate Koszul-duality criterion*). Let  $\mathfrak{g}_{\text{adm}}$  be a Tate graded Lie algebra satisfying the lower-central Tate-pronilpotence hypothesis of Lemma 5.165; for the formal-local setting this means one of the nilpotent bracket-compatible finite-window replacements, the weighted admissible envelope where the hypothesis is explicitly imposed, or another stated admissible replacement. Put

$$B_{\text{adm}} = C_{\text{CE}, \text{cont}}^\bullet(\mathfrak{g}_{\text{adm}}) = \widehat{\mathbf{S}}(\mathfrak{g}_{\text{adm}, \text{cont}}^\vee[-1]).$$

Then

$$B_{\text{adm}}^! \simeq \widehat{U}(\mathfrak{g}_{\text{adm}})$$

in the completed Tate category. For the strict full formal Hamiltonian algebra  $\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$  with its linear and conformal-symplectic low-degree modes, the same conclusion is a criterion: it follows after supplying a coefficient replacement or relative model killing the obstruction tuple of Theorem 5.5.

*Proof.* This is exactly Lemma 5.165 applied to  $\mathfrak{g}_{\text{adm}}$ . The lower-central Tate-pronilpotence condition is the load-bearing hypothesis: it ensures that the coefficient-completed CE coalgebra is conilpotent in every finite Tate window, that the continuous bar resolution is admissible, and that the completed Lie bar-cobar map

$$\Omega C_*^{\text{CE}}(\mathfrak{g}_{\text{adm}}) \longrightarrow \widehat{U}(\mathfrak{g}_{\text{adm}})$$

is a filtered quasi-isomorphism.

For the full formal Hamiltonian algebra, the direct CE/PV generator calculation gives the cochain identity before any bar-cobar hypothesis is imposed. To turn that identity into an enveloping/Koszul-duality theorem one must supply an admissible replacement, relative semisimple model, or weighted coefficient category satisfying the displayed lower-central Tate-pronilpotence and convergence hypotheses. The required data are exactly the Casimir, low-degree, semisimple, cone, and inverse-limit datum of Theorem 5.5.  $\square$

The completed PBW/order Rees enveloping algebra

$$U_{\hbar}(\mathfrak{g}) = \widehat{T}(\mathfrak{g})[[\hbar]] / \overline{\langle xy - (-1)^{|x||y|}yx - \hbar[x, y] \rangle},$$

with closure in tensor length and the Tate coefficient topology, makes the PBW special fibre precise in every admissible replacement to which Proposition 5.168 applies. The Rees parameter  $\hbar$  records word length in the enveloping algebra and is independent of Taylor degree. At  $\hbar = 0$  the completed PBW special fibre is

$$U_{\hbar}(\mathfrak{g})/(\hbar) \cong \widehat{\mathbf{S}}(\mathfrak{g}) = \widehat{\mathbf{S}}(\widehat{\mathfrak{h}} \oplus \mathfrak{h}^\vee[1]).$$

After choosing the monomial-dual Taylor basis, the underlying label set of  $\mathfrak{h}^\vee$  is identified with an abstract copy  $\tilde{\mathcal{A}}_{\text{desc}}$ ; this identification transports labels only, with the  $\mathfrak{h}$ -module structure supplied separately. The degree-wise polynomial subalgebra  $\mathbf{S}(\tilde{\mathcal{A}} \oplus \tilde{\mathcal{A}}_{\text{desc}}[1])$  is the stable classical brane Hamiltonian descendant algebra of Proposition 3.40: the degree-wise polynomial part of the PBW associated graded gives the brane Hamiltonian descendant labels. For the full formal Hamiltonian algebra this is the PBW/Rees label level. A filtered enveloping/Koszul-duality theorem is obtained exactly when the admissible or relative replacement data of Proposition 5.168 are supplied. The interacting closed Hamiltonian bracket deforms the special fibre inside that admissible enveloping algebra; on the brane side the matching deformation at order one is the Hamiltonian trace bracket proved below.

### 5.7.1 FIRST-ORDER POISSON IDENTIFICATION

The formal-stalk chart evaluates a closed Hamiltonian on the matrix-valued transverse coordinate  $(\phi_1, \phi_2)$ ; Dirac reduction forces  $[\phi_1, \phi_2] = 0$ ; the surviving algebra of cyclic single traces is the first nontrivial test. The question is whether the closed Poisson bracket on  $\tilde{\mathcal{A}} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1$  and the matrix trace bracket give equal coefficients on the trace generators.

The PBW special-fibre label correspondence used in the reduced comparison reads

$$z_1^k z_2^l \leftrightarrow \overline{\text{Tr}(\phi_1^k \phi_2^l)}.$$

The reduced Poisson bracket on the closed side is

$$\{z_1^k z_2^l, z_1^m z_2^n\}_{\tilde{\mathcal{A}}} = \overline{(kn - lm) z_1^{k+m-1} z_2^{l+n-1}}.$$

If an exponent in the displayed monomial is negative then the coefficient  $kn - lm$  is already zero; if the displayed monomial is constant, its class in  $\tilde{\mathcal{A}}$  is zero. The matching bracket on the open side is the canonical bracket induced by the first-order kinetic term  $\int \text{Tr}(\phi_1 d\phi_2)$ , with the sign convention  $\iota_{X_f} \omega = -df$  fixed in the AKSZ remark above:

$$\{(\phi_1)^i_j, (\phi_2)^k_l\} = \delta_l^i \delta_j^k, \quad \{(\phi_1)^i_j, (\phi_1)^k_l\} = 0 = \{(\phi_2)^i_j, (\phi_2)^k_l\}.$$

**LEMMA 5.169 (Hamiltonian trace bracket).** The canonical matrix Poisson bracket descends to the invariant reduced BV cohomology and then to the Hamiltonian Lie quotient by constant trace classes.

*Proof.* For  $\epsilon \in \mathfrak{gl}_N$ , the component  $\mu_\epsilon = -\text{Tr}(\epsilon[\phi_1, \phi_2])$  of the moment map generates the infinitesimal conjugation action, with the sign fixed by the convention for the bracket. Hence  $\{F, \mu_\epsilon\} = 0$  for every invariant function  $F$ . If  $I_\mu$  is the ideal generated by all  $\mu_\epsilon$ , then  $\{F, \mu_\epsilon K\} = \mu_\epsilon \{F, K\}$  lies again in  $I_\mu$ . Therefore the bracket of invariant functions descends to  $(\mathcal{R}_N/I_\mu)^{GL_N}$ , equivalently to the degree-zero reduced BV cohomology. The central line  $\mathbb{C} \cdot \text{Tr}(1)$  is central for this Lie bracket, so the Hamiltonian vector-space quotient by constants is compatible with the bracket. The associative quotient by the unit is a different object.  $\square$

**PROPOSITION 5.170 (First-order trace bracket).** In the stable connected Hamiltonian trace cohomology,

$$\overline{\{\text{Tr}(\phi_1^k \phi_2^l), \text{Tr}(\phi_1^m \phi_2^n)\}} = \overline{(kn - lm) \text{Tr}(\phi_1^{k+m-1} \phi_2^{l+n-1})}.$$

*Proof.* Write  $F_{k,l} = \text{Tr}(\phi_1^k \phi_2^l)$ . The matrix Poisson bracket induced by

$$\{(\phi_1)^i_j, (\phi_2)^k_l\} = \delta_l^i \delta_j^k$$

is

$$\{F, G\} = \text{Tr} \left( \frac{\partial F}{\partial \phi_1} \frac{\partial G}{\partial \phi_2} - \frac{\partial F}{\partial \phi_2} \frac{\partial G}{\partial \phi_1} \right),$$

where the derivatives are cyclic matrix derivatives. Varying one occurrence at a time gives

$$\begin{aligned} \frac{\partial F_{k,l}}{\partial \phi_1} &= \sum_{a=0}^{k-1} \phi_1^{k-1-a} \phi_2^l \phi_1^a, \\ \frac{\partial F_{k,l}}{\partial \phi_2} &= \sum_{b=0}^{l-1} \phi_2^{l-1-b} \phi_1^k \phi_2^b. \end{aligned}$$

Therefore the first summand in the bracket is a sum of  $kn$  traces, each obtained by removing one  $\phi_1$  from the first word and one  $\phi_2$  from the second word. The second summand is a sum of  $lm$  traces obtained by removing one  $\phi_2$  from the first word and one  $\phi_1$  from the second. Every trace appearing in either nonzero sum has multidegree  $(k + m - 1, l + n - 1)$ . Lemma 3.30 identifies all cyclic words with this multidegree in reduced Koszul cohomology. The coefficient is therefore  $kn - lm$ , and the common class is  $\overline{\text{Tr}(\phi_1^{k+m-1} \phi_2^{l+n-1})}$ . When the common word has multidegree  $(0, 0)$ , this class is the scalar trace and is zero in the Hamiltonian quotient by constants.  $\square$

PROPOSITION 5.171 (*Open action on primitive descendants*). Let

$$\tilde{\Psi}_{a,b} = \binom{a+b}{a}^{-1} \Psi_{a,b}$$

be the averaged primitive one- $\psi$  class. In the stable primitive one- $\psi$  homology,

$$\overline{\{\text{Tr}(\phi_1^p \phi_2^q), \tilde{\Psi}_{a,b}\}} = (pb - qa) \tilde{\Psi}_{a+p-1, b+q-1},$$

with the convention that a term with a negative exponent is zero. Thus the unreduced polynomial descendant classes carry the adjoint polynomial Hamiltonian action. The Hamiltonian descendant sector is the quotient by the scalar line  $\mathbb{C}\tilde{\Psi}_{0,0}$ , so the displayed action is projected further by killing the scalar target.

*Proof.* Put  $x = \phi_1$  and  $y = \phi_2$ . The cyclic derivatives of  $F_{p,q} = \text{Tr}(x^p y^q)$  are

$$\frac{\partial F_{p,q}}{\partial x} = \sum_{i=0}^{p-1} x^{p-1-i} y^q x^i, \quad \frac{\partial F_{p,q}}{\partial y} = \sum_{j=0}^{q-1} y^{q-1-j} x^p y^j.$$

For a one- $\psi$  cyclic word  $\text{Tr}(\psi w)$  of bidegree  $(a, b)$ , differentiating with respect to  $y$  gives one summand for each of the  $b$  occurrences of  $y$  in  $w$ , and differentiating with respect to  $x$  gives one summand for each of the  $a$  occurrences of  $x$ . Multiplying by the displayed cyclic derivatives of  $F_{p,q}$  therefore produces  $pb$  positive one- $\psi$  words and  $qa$  negative one- $\psi$  words for each  $w \in W_{a,b}$ , all of total bidegree  $(a + p - 1, b + q - 1)$ .

The bracket is a cycle because  $F_{p,q}$  is invariant, hence  $\{F_{p,q}, \mu_\epsilon\} = 0$  for every moment-map component  $\mu_\epsilon = -\text{Tr}(\epsilon[x, y])$ . Equivalently, the Hamiltonian vector field of  $F_{p,q}$  preserves the Koszul differential  $Q\psi = [x, y]$  and acts on Koszul homology. By Proposition 3.27, the target primitive one- $\psi$

homology in that bidegree is one-dimensional. The functional  $\epsilon(\text{Tr}(\psi w)) = 1$  detects its coefficient. Since  $\epsilon(\Psi_{a,b}) = 1$ , the averaged input gives

$$\epsilon(\{F_{p,q}, \tilde{\Psi}_{a,b}\}) = pb - qa.$$

Also  $\epsilon(\tilde{\Psi}_{a+p-1,b+q-1}) = 1$ . Hence the displayed formula follows. If the target bidegree is  $(0, 0)$ , the same formula is still the unreduced primitive one- $\psi$  action; the Hamiltonian descendant quotient sends that scalar class to zero.  $\square$

**COROLLARY 5.172** (*Descendant action and coadjoint action: structural separation*). Under the PBW special-fibre pairing of Corollary 3.29, the open Hamiltonian descendant action on the averaged polynomial  $\tilde{\Psi}_{a,b}$ -basis is the symmetric PBW action

$$F_{p,q} : \tilde{\Psi}_{a,b} \mapsto (pb - qa) \tilde{\Psi}_{a+p-1,b+q-1},$$

while the closed coadjoint action on the principal-part basis  $\rho_{a,b} \leftrightarrow \delta_{a,b} \in \mathfrak{h}_{\text{cont}}^\vee$  is the negative-transpose coadjoint action

$$\text{ad}_{z_1^p z_2^q}^\vee \delta_{a,b} = -(pb - qa + p - q) \delta_{a-p+1,b-q+1},$$

with negative-index targets zero. The two actions agree only on the trivial Hamiltonian  $p = q = 0$ . The deformation quantisation of the cotangent factor sees the second action; the special-fibre PBW symmetric algebra sees the first. The structural reason is the PBW special-fibre versus coadjoint-module separation of Theorem 5.180 below: the  $\tilde{\Psi}_{a,b}$  and the  $\rho_{a,b}$  have the same index set but different  $\mathfrak{h}$ -module structures.

*Proof.* Proposition 5.171 gives the symmetric PBW formula on  $\tilde{\Psi}_{a,b}$ . The coadjoint action is dual to the Poisson bracket: for  $f = z_1^p z_2^q$  and the Taylor-dual basis vector  $\delta_{a,b}$ ,

$$(\text{ad}_f^\vee \delta_{a,b})(z_1^m z_2^n) = -\delta_{a,b}((pn - qm)z_1^{p+m-1}z_2^{q+n-1}),$$

giving the displayed coadjoint formula. For example,  $z_1$  acts on open descendants by  $\tilde{\Psi}_{a,b} \mapsto b \tilde{\Psi}_{a,b-1}$ , whereas it acts coadjointly by  $\delta_{a,b} \mapsto -(b+1) \delta_{a,b+1}$ . The scalar edge  $\{\text{Tr}(\phi_1), \tilde{\Psi}_{0,1}\} = \Psi_{0,0}$  is removed with the scalar descendant. The two formulas have opposite index shift directions and differ by  $-(pb - qa + p - q)\rho_{a-p+1,b-q+1} - (pb - qa)\rho_{a+p-1,b+q-1}$  after any bidegree-only vector-space relabelling between the two alphabets, which is non-zero unless both targets vanish identically; this happens exactly on the trivial  $p = q = 0$  action. The PBW special-fibre product identification is Corollary 3.29; the coadjoint half is Theorem 5.178; the Laurent/distributional principal-part realization is Proposition 5.175 and the separation Theorem 5.180.  $\square$

The coadjoint identification on the principal-part side requires three notions of locality on the brane line  $\mathbb{R}$  to remain distinct.

**LEMMA 5.173** (*Three notions of locality*). On the brane line  $\mathbb{R}$  with formal holomorphic transverse coordinates  $z_1, z_2$  at the brane point  $p \in \mathbb{C}^2$ , the following three notions of locality are independent.

- (i) *Support locality*: distributional support on the brane: a section  $T \otimes m$  of  $\mathcal{D}'_c(I) \hat{\otimes} \mathcal{M}$  has support locality whenever  $\text{supp } T \subset I \subset \mathbb{R}$ .
- (ii) *Topological locality*: locally constant in the  $\mathbb{R}$ -direction up to  $Q$ -exact translations: every section is identified with its translate by a parameter on  $\mathbb{R}$  modulo the open BV differential  $Q_{\text{open}}$ .



- (iii) *Formal holomorphic locality*: polynomial, or Taylor-series, dependence on the transverse coordinates  $z_1, z_2$ .

The polynomial trace observables  $O_f(a) = \int_{\mathbb{R}} a(t) \overline{\text{Tr } f(\phi_1(t), \phi_2(t))} dt$  have all three. The shifted cotangent currents  $\Theta_\rho(B)$  of Construction 5.184 have support locality and topological locality. Their transverse content is momentum data attached to the brane, living in the  $\mathbb{C}$ -linear residue annihilator of scalar Hamiltonians inside the top local-cohomology principal-part module,  $\text{Ann}_{\text{res}}(\mathbb{C} \cdot 1)$ .

*Proof.* Independence is a category-theoretic statement: a polynomial in  $z_1, z_2$  with point support is constant in  $\mathbb{R}$ , separating support locality from formal holomorphic locality; a delta-current at one point of  $\mathbb{R}$  has compact support and only trivial  $Q$ -exact translation, separating support locality from topological locality. The cotangent currents  $\Theta_\rho(B)$  are smooth or distributional sections of the line  $\mathbb{R}$  (hence (1)) and translation-invariant up to  $Q_{\text{open}}$ -exact cocycles (hence (2)); their transverse content  $\rho \in \mathcal{P}$  is a top-local-cohomology principal part, hence a residue-dual distributional mode.  $\square$

*Remark 5.174.* In canonical quantization of a constrained system, the configuration variables are locally polynomial in spatial coordinates and the conjugate momenta are dual distributions. This is exactly the separation Lemma 5.173 records: configuration observables on a brane have formal holomorphic locality (3), and momentum observables have only (1) and (2). The principal-part sector is the algebraic image of this separation in the local Hamiltonian BF model.

PROPOSITION 5.175 (*Principal-part realization of the coadjoint module*). Let

$$\mathcal{P} = \bigoplus_{\substack{a, b \geq 0 \\ a+b > 0}} \mathbb{C} \rho_{a,b}, \quad \rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2,$$

with residue pairing

$$\langle \rho_{a,b}, z_1^m z_2^n \rangle = \text{Res}_{z_1=z_2=0} z_1^{m-a-1} z_2^{n-b-1} dz_1 dz_2 = \delta_{a,m} \delta_{b,n}.$$

Let  $\pi_{\text{pp}}$  denote projection of Laurent two-forms to the span of the principal parts  $\rho_{a,b}$ . The action

$$f \cdot \rho = \pi_{\text{pp}}(\{f, \rho\})$$

identifies  $\mathcal{P}$  with the continuous coadjoint module  $\mathfrak{h}_{\text{cont}}^\vee$ . On monomials,

$$z_1^p z_2^q \cdot \rho_{a,b} = -(pb - qa + p - q) \rho_{a-p+1, b-q+1},$$

with the convention that a term with a negative index is zero.

*Proof.* The residue pairing identifies  $\rho_{a,b}$  with the Taylor-dual functional  $\delta_{a,b}$ . Hamiltonian vector fields preserve the holomorphic volume form  $dz_1 dz_2$ , so integration by parts for the residue pairing gives

$$\langle \{f, \rho\}, g \rangle = -\langle \rho, \{f, g\} \rangle.$$

Hence the Laurent principal-part action realizes the coadjoint action. Directly, for  $f = z_1^p z_2^q$  and  $\rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2$ ,

$$\begin{aligned} \{f, z_1^{-a-1} z_2^{-b-1}\} &= p z_1^{p-1} z_2^q (-(b+1)) z_1^{-a-1} z_2^{-b-2} \\ &\quad - q z_1^p z_2^{q-1} (-(a+1)) z_1^{-a-2} z_2^{-b-1} \\ &= -(pb - qa + p - q) z_1^{p-a-2} z_2^{q-b-2}. \end{aligned}$$

Since  $z_1^{p-a-2} z_2^{q-b-2} dz_1 dz_2 = \rho_{a-p+1, b-q+1}$  when both indices are nonnegative, principal-part projection gives the displayed formula.  $\square$

Two rigidity statements determine the principal-part identification after the residue scalar is fixed: among continuous total-degree-zero pairings the residue pairing is unique up to a single nonzero scalar, and among continuous total-degree-zero  $\mathfrak{h}$ -module isomorphisms the coadjoint identification  $\mathcal{P} \cong \mathfrak{h}_{\text{cont}}^\vee$  is unique modulo that scalar.

**THEOREM 5.176** (*Residue pairing, normalized up to scalar*). Up to a nonzero scalar, the residue pairing

$$\langle \cdot, \cdot \rangle : \mathcal{P} \times (\mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1) \longrightarrow \mathbb{C}, \quad \langle \rho_{a,b}, [z_1^m z_2^n] \rangle = \delta_{a,m} \delta_{b,n},$$

is the unique non-degenerate continuous  $\mathfrak{h}$ -equivariant pairing graded of total degree zero with the property that every element of  $\mathcal{P} = \text{Ann}_{\text{res}}(\mathbb{C} \cdot 1)$  annihilates the constant line.

*Proof. Existence.* The residue pairing is non-degenerate by the displayed Kronecker relation; it is continuous because  $\mathcal{P}$  is graded discrete and  $\mathbb{C}[[z_1, z_2]]$  carries the Taylor filtration; equivariance is integration by parts for the Poisson bracket, since Hamiltonian vector fields preserve the holomorphic two-form  $dz_1 dz_2$ :

$$\langle \{f, \rho\}, g \rangle = \text{Res}(\{f, \rho\}g) = -\text{Res}(\rho \{f, g\}) = -\langle \rho, \{f, g\} \rangle.$$

*Uniqueness.* Let  $\beta : \mathcal{P} \times (\mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1) \rightarrow \mathbb{C}$  be a second pairing satisfying the same conditions, and write  $\beta_{a,b;m,n} = \beta(\rho_{a,b}, z_1^m z_2^n)$ . Since  $\beta$  is graded of total degree zero,  $\beta_{a,b;m,n} = 0$  unless  $a + b = m + n$ .

*Diagonal weight constraint.* The Hamiltonian  $z_1 z_2$  acts diagonally on monomials by  $\{z_1 z_2, z_1^m z_2^n\} = (n - m) z_1^m z_2^n$ . Direct Poisson computation on principal parts uses the Hamiltonian vector field  $X_{z_1 z_2} = z_2 \partial_{z_1} - z_1 \partial_{z_2}$  (with  $X_f = (\partial_{z_1} f) \partial_{z_2} - (\partial_{z_2} f) \partial_{z_1}$ ), giving  $X_{z_1 z_2}(z_1^{-a-1} z_2^{-b-1}) = (a - b) z_1^{-a-1} z_2^{-b-1}$ , so  $\{z_1 z_2, \rho_{a,b}\} = (a - b) \rho_{a,b} = -(b - a) \rho_{a,b}$ . Equivariance  $\beta(\{z_1 z_2, \rho_{a,b}\}, z_1^m z_2^n) = -\beta(\rho_{a,b}, \{z_1 z_2, z_1^m z_2^n\})$  gives  $-(b - a) \beta_{a,b;m,n} = -(n - m) \beta_{a,b;m,n}$ , hence  $\beta_{a,b;m,n} = 0$  unless  $a - b = m - n$ . Combined with the weight constraint  $a + b = m + n$ , this gives  $\beta_{a,b;m,n} = 0$  unless  $(a, b) = (m, n)$ . So  $\beta$  is diagonal in the residue basis:  $\beta_{a,b;m,n} = c_{a,b} \delta_{a,m} \delta_{b,n}$  for some scalars  $c_{a,b} \in \mathbb{C}$ .

*Coefficients are constant.* The Hamiltonian generator  $f = z_1$  has  $X_{z_1} = \partial_{z_2}$ , so  $\{z_1, z_1^m z_2^n\} = n z_1^m z_2^{n-1}$  and a direct Poisson computation gives  $\{z_1, \rho_{a,b}\} = -(b + 1) \rho_{a,b+1}$ . Equivariance reads  $\beta(\{z_1, \rho_{a,b}\}, z_1^m z_2^n) = -\beta(\rho_{a,b}, \{z_1, z_1^m z_2^n\})$ , i.e.,  $-(b + 1) \beta_{a,b+1;m,n} = -n \beta_{a,b;m,n-1}$ . Both sides vanish identically unless the diagonality condition is met on each pair. The right-hand side is non-zero only if  $(a, b) = (m, n - 1)$ , i.e.,  $m = a, n = b + 1$ . The left-hand side is non-zero only if  $(a, b + 1) = (m, n)$ , i.e.,  $m = a, n = b + 1$ . Both indices coincide, giving  $(b + 1) c_{a,b+1} = (b + 1) c_{a,b}$ , hence  $c_{a,b+1} = c_{a,b}$  when both coefficients occur in the reduced index set; that is, for  $a + b > 0$ . The reduced module  $\mathcal{P}$  starts at  $a + b > 0$ , so the case  $a = b = 0$  is omitted. Symmetrically from  $f = z_2$  (using  $X_{z_2} = -\partial_{z_1}$ , so  $\{z_2, z_1^m z_2^n\} = -m z_1^{m-1} z_2^n$  and  $\{z_2, \rho_{a,b}\} = (a + 1) \rho_{a+1,b}$ ) the equality  $c_{a+1,b} = c_{a,b}$  holds when both coefficients occur in the reduced index set. Connecting these chains through  $c_{1,1}$ , all  $c_{a,b}$  for  $a + b \geq 1$  equal a single constant  $c \in \mathbb{C}$ . Non-degeneracy forces  $c \neq 0$ .

Therefore  $\beta = c \langle \cdot, \cdot \rangle$  is the residue pairing scaled by  $c$ . The condition on constants holds because  $\mathcal{P} = \text{Ann}_{\text{res}}(\mathbb{C} \cdot 1)$ ; equivalently  $\rho_{0,0} \notin \mathcal{P}$ .  $\square$

**Remark 5.177.** Theorem 5.176 is better understood as Matlis-local-cohomology rigidity. Let  $R = \mathbb{C}[[z_1, z_2]]$  and  $\mathfrak{m} = (z_1, z_2)$ . The top local cohomology module, in the Čech-Laurent residue

model of Grothendieck–Hartshorne local cohomology and Kunz–Cox–Dickenstein power-series residues [20, 21],

$$H_{\mathfrak{m}}^2(R) dz_1 dz_2 = \bigoplus_{a,b \geq 0} \mathbb{C} z_1^{-a-1} z_2^{-b-1} dz_1 dz_2$$

is, after the trivialization  $\omega_R = R dz_1 dz_2$ , the injective hull  $E_R(\mathbb{C})$  supplied by Matlis duality [19]. The reduced principal-part module  $\mathcal{P} = \text{Ann}_{\text{res}}(\mathbb{C} \cdot 1)$  is the  $\mathbb{C}$ -linear residue annihilator of constants. The  $R$ -submodule structure belongs to  $E_R(\mathbb{C})$ , with the scalar line treated in the ambient Matlis module. The reduced module is the  $\mathbb{C}$ -linear annihilator of the scalar Hamiltonian under the Matlis residue evaluation pairing. Hamiltonian vector fields preserve  $dz_1 dz_2$ , so this restricted pairing transports their action to the coadjoint action. The explicit coefficient proof above is the coordinate form of this local-duality statement.

**THEOREM 5.178** (*Principal-part coadjoint isomorphism, Matlis form*). After fixing the residue scalar in Theorem 5.176, the residue pairing induces an isomorphism of  $\mathfrak{h}$ -modules

$$\Phi_{\text{coad}} : \mathcal{P} \xrightarrow{\sim} \mathfrak{h}_{\text{cont}}^{\vee}, \quad \rho_{a,b} \mapsto \delta_{a,b},$$

characterised uniquely (up to the residue scalar  $c$  of Theorem 5.176) by the property that for every  $f \in \mathfrak{h}$  and every  $g \in \mathfrak{h}$ ,

$$\langle \Phi_{\text{coad}}^{-1}(\text{ad}_f^{\vee} \delta), g \rangle = -\langle \Phi_{\text{coad}}^{-1}(\delta), \{f, g\} \rangle.$$

In monomial form,

$$z_1^p z_2^q \cdot \rho_{a,b} = -(pb - qa + p - q) \rho_{a-p+1, b-q+1},$$

with the convention that any term with a negative index is zero. The isomorphism is continuous, of total degree zero, and the unique continuous total-degree-zero  $\mathfrak{h}$ -module isomorphism  $\mathcal{P} \rightarrow \mathfrak{h}_{\text{cont}}^{\vee}$  after the residue scalar is fixed; before that normalization it is unique modulo scalar.

*Proof. Existence on monomials.* Let  $f = z_1^p z_2^q$  and  $\rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2$ . Since  $\{z_1, z_2\} = 1$ ,

$$\{f, z_1^{-a-1} z_2^{-b-1}\} = p z_1^{p-1} z_2^q \cdot (-(b+1)) z_1^{-a-1} z_2^{-b-2} - q z_1^p z_2^{q-1} \cdot (-(a+1)) z_1^{-a-2} z_2^{-b-1},$$

which simplifies to  $-(pb - qa + p - q) z_1^{p-a-2} z_2^{q-b-2}$ . Multiplying by  $dz_1 dz_2$  identifies this with  $\rho_{a-p+1, b-q+1}$  when both indices are  $\geq 0$ ; otherwise the product lies outside  $\mathcal{P}$  and the principal-part projection  $\pi_{\text{pp}}$  of Proposition 5.175 gives zero. The integration by parts formula  $\langle \{f, \rho\}, g \rangle = -\langle \rho, \{f, g\} \rangle$  identifies this with the coadjoint action.

*Continuity.* The dual filtration on  $\mathfrak{h}_{\text{cont}}^{\vee}$  has  $F^N \mathfrak{h}_{\text{cont}}^{\vee}$  equal to the functionals vanishing on  $\mathfrak{m}^{N+1} \subset \mathfrak{h}$ , where  $\mathfrak{m}$  is the maximal ideal. The grading of  $\mathcal{P}$  by total weight  $a + b$  corresponds under  $\Phi_{\text{coad}}$  to the dual filtration:  $F^N = \bigoplus_{a+b \leq N} \mathbb{C} \delta_{a,b}$  is the  $\Phi_{\text{coad}}$ -image of  $\bigoplus_{a+b \leq N} \mathbb{C} \rho_{a,b}$ .

*Verification on three instances.* With the convention  $\{z_1, z_2\} = 1$ , the Hamiltonian vector field is  $X_f = (\partial_{z_1} f) \partial_{z_2} - (\partial_{z_2} f) \partial_{z_1}$ .

- (i)  $(p, q, a, b) = (1, 0, 0, 1)$ : the formula gives  $z_1 \cdot \rho_{0,1} = -(1 \cdot 1 - 0 \cdot 0 + 1 - 0) \rho_{0-1+1, 1-0+1} = -2\rho_{0,2}$ . Direct:  $X_{z_1} = \partial_{z_2}$ , so  $X_{z_1}(z_1^{-1} z_2^{-2}) = -2z_1^{-1} z_2^{-3}$ , and multiplying by  $dz_1 dz_2$  gives  $-2\rho_{0,2}$ . Match.
- (ii)  $(p, q, a, b) = (0, 1, 1, 0)$ : the formula gives  $z_2 \cdot \rho_{1,0} = -(0 \cdot 0 - 1 \cdot 1 + 0 - 1) \rho_{1-0+1, 0-1+1} = 2\rho_{2,0}$ . Direct:  $X_{z_2} = -\partial_{z_1}$ , so  $X_{z_2}(z_1^{-2} z_2^{-1}) = 2z_1^{-3} z_2^{-1}$ , multiplied by  $dz_1 dz_2$  gives  $2\rho_{2,0}$ . Match.
- (iii)  $(p, q, a, b) = (2, 1, 1, 1)$ : the formula gives  $z_1^2 z_2 \cdot \rho_{1,1} = -(2 \cdot 1 - 1 \cdot 1 + 2 - 1) \rho_{0,1} = -2\rho_{0,1}$ . Direct:  $X_{z_1^2 z_2} = 2z_1 z_2 \partial_{z_1} - z_1^2 \partial_{z_2}$ ;  $X_{z_1^2 z_2}(z_1^{-2} z_2^{-2}) = -4z_1^{-1} z_2^{-2} + 2z_1^{-1} z_2^{-2} = -2z_1^{-1} z_2^{-2}$ , multiplied by  $dz_1 dz_2$  gives  $-2\rho_{0,1}$ . Match.

These three cases are confirmed as part of the 1225-case exact check, which verifies the formula for  $0 \leq p, q, a, b \leq 5, p + q > 0, a + b > 0$ .

*Uniqueness.* Suppose  $\Phi' : \mathcal{P} \rightarrow \mathfrak{h}_{\text{cont}}^\vee$  is another continuous  $\mathfrak{h}$ -equivariant isomorphism of total degree zero. Define

$$\beta_{\Phi'}(\rho, g) = \Phi'(\rho)(g).$$

This is a non-degenerate continuous total-degree-zero pairing  $\mathcal{P} \times \mathfrak{h} \rightarrow \mathbb{C}$ , it kills the scalar line by definition of  $\mathfrak{h}$ , and  $\mathfrak{h}$ -equivariance of  $\Phi'$  gives the coadjoint equivariance condition of Theorem 5.176. Therefore  $\beta_{\Phi'} = c' \langle \cdot, \cdot \rangle$  for a unique  $c' \in \mathbb{C}^\times$ , and hence  $\Phi' = c' \Phi_{\text{coad}}$ . The scalar is exactly the residue scalar of Theorem 5.176.  $\square$

*Remark 5.179 (Sign convention).* The sign  $-(pb - qa + p - q)$  on the principal-part side and the sign  $(pb - qa)$  on the averaged polynomial  $\tilde{\Psi}_{a,b}$  side of Proposition 5.171 are both correct in their own conventions; they differ because the  $\Psi_{a,b}$  are special-fibre PBW representatives and the  $\rho_{a,b}$  are deformation-compatible coadjoint representatives. See Theorem 5.180 below for the precise separation.

**THEOREM 5.180 (PBW special fibre versus coadjoint deformation module).** Let  $\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1]$  be the semidirect Lie algebra of Proposition 2.7. The polynomial one- $\psi$  basis  $\Psi_{a,b} \in \tilde{A}_{\text{desc}}$  and the principal-part basis  $\rho_{a,b} \in \mathcal{P} \cong \mathfrak{h}_{\text{cont}}^\vee$  are related as follows.

- (i) *Special-fibre PBW basis.* The  $\tilde{\Psi}_{a,b}$  form the primitive one- $\psi$  PBW label basis in the special-fibre symmetric algebra  $\mathbf{S}(\mathfrak{h} \oplus \mathfrak{h}^\vee[1])|_{\hbar=0}$ , and the open Hamiltonian Lie action is the *symmetric PBW action* on this generator span:

$$F_{p,q} : \tilde{\Psi}_{a,b} \mapsto (pb - qa) \tilde{\Psi}_{a+p-1, b+q-1}.$$

The shift  $(a + p - 1, b + q - 1)$  is determined by multiplication by the Hamiltonian monomial and by the scalar Hamiltonian quotient.

- (ii) *Deformation-level coadjoint basis.* The  $\rho_{a,b}$  form a basis of the deformation-compatible continuous coadjoint module  $\mathfrak{h}_{\text{cont}}^\vee \cong \mathcal{P}$  of Theorem 5.178, and the  $\mathfrak{h}$ -action is the *principal-part coadjoint action* on this basis:

$$z_1^p z_2^q \cdot \rho_{a,b} = -(pb - qa + p - q) \rho_{a-p+1, b-q+1}.$$

The displayed shift is the negative-transpose residue action. Linear Hamiltonians raise pole order, while higher monomials have the mixed direction prescribed by the formula.

- (iii) *Module separation.* The bidegree relabelling between the PBW special-fibre basis  $\tilde{\Psi}_{a,b}$  and the coadjoint basis  $\rho_{a,b} \leftrightarrow \delta_{a,b}$  is an isomorphism of graded vector spaces and label sets. Pulling the coadjoint  $\mathfrak{h}$ -action across this relabelling gives  $-(pb - qa + p - q) \rho_{a-p+1, b-q+1}$ , while the symmetric PBW action displayed in (i) gives  $(pb - qa) \rho_{a+p-1, b+q-1}$ . The formal Moyal/Weyl Rees deformation controls the Weyl algebra and its continuous dual  $\mathfrak{h}_{\hbar, \text{cont}}^\vee$ ; the polynomial one- $\psi$  descendant module and  $\mathcal{P}$  remain distinct  $\mathfrak{h}$ -modules. The Fourier-polarized Rees label bridge is constructed in Theorem 8.12; the same-action and unreduced quantum-observable realizations remain separate problems.

Thus the two formulas define distinct  $\mathfrak{h}$ -modules in the formal-local category: the PBW special fibre and the continuous coadjoint cotangent module, respectively. The structural separation recorded in Corollary 5.172 follows.

*Proof.* (1) is Proposition 5.171, with the rephrasing in the PBW special-fibre basis. (2) is Theorem 5.178. For (3), the symmetric PBW action on the nonconstant  $\tilde{\Psi}_{a,b}$  labels is the PBW special-fibre action after the Hamiltonian scalar quotient. The coadjoint action on  $\rho_{a,b}$  is the negative-transpose principal-part action. These two module behaviours are separated by the displayed formulas already at the linear Hamiltonians  $z_1, z_2$ ; hence any bidegree-only relabelling between  $\tilde{\Psi}_{a,b}$  and  $\delta_{a,b}$  deviates from an  $\mathfrak{h}$ -module map already at the linear Hamiltonians. The Rees-deformation  $\mathcal{A}_{\hbar} = (\mathbb{C}[[z_1, z_2]][[\hbar]], *)$  has the property that the formal Moyal product  $f * g = fg + \frac{\hbar}{2}\{f, g\} + O(\hbar^2)$  acts on its dual by the coadjoint formula at *all* orders of  $\hbar$ : the dual representation of the Moyal Lie algebra  $(\mathfrak{h}_{\hbar}, [-, -]_{\hbar})$  on  $\mathfrak{h}_{\hbar}^{\vee}$  is the residue pairing extended  $\mathbb{C}[[\hbar]]$ -linearly. Specialising to  $\hbar = 0$  recovers the Poisson coadjoint action of the completed Hamiltonian algebra  $\mathfrak{h}$  on  $\mathfrak{h}^{\vee} = \mathcal{P}$ , and its restriction to the dense polynomial subalgebra  $\tilde{\mathcal{A}} \subset \mathfrak{h}$ . The PBW filtration is the Rees filtration, and the special-fibre  $\hbar = 0$  symmetric algebra is the associated graded; the symmetric PBW action on  $\tilde{\Psi}_{a,b}$  is exactly the  $\hbar = 0$  *associated graded* image of the classical Hamiltonian vector-field action on the polynomial PBW descendant labels; the adjoint and coadjoint actions of the Lie algebra live on the deformation-level module. It therefore separates the descendant module from the continuous coadjoint module, and identifies the Rees/Fourier bridge as additional Fourier-polarized structure.  $\square$

PROPOSITION 5.181 (*Polynomial boundary obstruction for principal parts*). In the polynomial open BV/Koszul matrix model studied here, any realization of the principal-part coadjoint module requires a boundary sector beyond ordinary polynomial boundary local cohomology classes

$$\Theta_{a,b}, \quad a, b \geq 0, \quad a + b > 0,$$

of Proposition 5.175.

*Proof.* The linear Hamiltonians

$$O_{1,0} = \overline{\text{Tr}(\phi_1)}, \quad O_{0,1} = \overline{\text{Tr}(\phi_2)}$$

act on polynomial boundary observables by the constant Hamiltonian vector fields  $\partial_{\phi_2}$  and  $-\partial_{\phi_1}$ , respectively. On every ordinary polynomial observable these operators are locally nilpotent: after enough iterations the polynomial degree in the differentiated variable is exhausted.

On the principal-part coadjoint module, however, Proposition 5.175 gives

$$O_{1,0} \cdot \rho_{a,b} = -(b+1)\rho_{a,b+1}, \quad O_{0,1} \cdot \rho_{a,b} = (a+1)\rho_{a+1,b}.$$

Both operators have non-locally-nilpotent action on every nonzero principal-part generator. Local nilpotence is preserved under isomorphism of modules. Therefore the desired  $\Theta_{a,b}$  requires a non-polynomial boundary realization; the residue principal-part module is the unique such realization.  $\square$

The categorical refinement records the same local-nilpotence obstruction at the level of polynomial sub-bimodules. The statement is restricted to this local-nilpotence obstruction; classification of all  $\tilde{\mathcal{A}}$ -module morphisms out of  $\mathcal{P}$  is a separate question.

THEOREM 5.182 (*Categorical polynomial-realization obstruction for principal parts*). Let  $\mathbf{P}_{\text{pol}}$  denote the abelian category of  $\mathbb{Z}$ -graded bimodules  $\mathcal{M} = \bigoplus_{n \geq 0} \mathcal{M}_n$  over the polynomial Hamiltonian Lie algebra  $\tilde{\mathcal{A}} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1$ , on which:

- (i) every element acts as a Hamiltonian vector field;

- (ii) every linear Hamiltonian acts locally nilpotently: for every  $m \in M$  and every  $i \in \{1, 2\}$ , there exists  $N$  with  $(z_i \cdot)^N m = 0$ .

This category models polynomial boundary observables of the open BV theory with respect to the polynomial sub-bimodule structure preserved by all morphisms of the boundary algebra into itself. Then  $\mathcal{P}$  lies outside  $\text{Ob}(\mathcal{P}_{\text{pol}})$ . More generally, an injective  $\bar{A}$ -module map  $\mathcal{P} \hookrightarrow M$  with  $M \in \text{Ob}(\mathcal{P}_{\text{pol}})$  is obstructed by local nilpotence. The unique realization of  $\mathcal{P}$  is the residue principal-part realization, with the residue-coadjoint pole-raising Hamiltonian action.

*Proof.* By Theorem 5.178,  $z_1 \cdot \rho_{a,b} = -(b+1)\rho_{a,b+1}$  and  $z_2 \cdot \rho_{a,b} = (a+1)\rho_{a+1,b}$  on every generator. Iteration gives  $(z_1 \cdot)^N \rho_{a,b} = (-1)^N (b+1)(b+2) \cdots (b+N) \rho_{a,b+N} \neq 0$  for every  $N$ , so  $z_1$  acts non-locally-nilpotently on every nonzero element of  $\mathcal{P}$ . The same argument applies to  $z_2$ .

Hence  $\mathcal{P}$  itself violates condition (2) and lies outside of  $\mathcal{P}_{\text{pol}}$ . If  $\Phi : \mathcal{P} \hookrightarrow M$  were an injective  $\bar{A}$ -module map into an object of  $\mathcal{P}_{\text{pol}}$ , then  $m = \Phi(\rho_{a,b}) \neq 0$  for some generator. Equivariance gives

$$z_1^N \cdot m = \Phi(z_1^N \cdot \rho_{a,b}) = (-1)^N (b+1) \cdots (b+N) \Phi(\rho_{a,b+N}),$$

which is nonzero for every  $N$  by injectivity. This contradicts local nilpotence of  $z_1$  on  $M$ . The same argument applies with  $z_2$ .  $\square$

*Remark 5.183 (Polynomial obstruction).* The categorical obstruction applies to every polynomial sub-bimodule of every boundary observable algebra with locally nilpotent  $z_i$ -action; it is uniform in the boundary algebra and independent of the particular choice of classes  $\Theta_{a,b}$ . The distributional/Laurent principal-part construction of Construction 5.184 is therefore required for this realization.

*Construction 5.184 (Stalkwise boundary local-dual principal-part sector).* Let  $M_\delta^{\text{pp}}$  be the reduced continuous dual of the stable connected Hamiltonian trace sector, with basis  $\Theta_{a,b}$  dual to

$$O_{m,n} = \overline{\text{Tr}(\phi_1^m \phi_2^n)}, \quad m+n > 0,$$

by

$$\langle \Theta_{a,b}, O_{m,n} \rangle = \delta_{a,m} \delta_{b,n}.$$

Equivalently,

$$M_\delta^{\text{pp}} \cong \text{Ann}_{\text{res}}(\mathbb{C} \cdot 1) \subset H_{(z_1, z_2)}^2(\mathbb{C}[z_1, z_2]) dz_1 dz_2 \quad \text{after completing at } (z_1, z_2),$$

the continuous dual functionals vanishing on constants, with basis  $\rho_{a,b}$  for  $a+b > 0$ . Explicitly,

$$\rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2, \quad a+b > 0,$$

with

$$\Theta_{a,b} \longleftrightarrow \rho_{a,b}.$$

This is a  $\mathbb{C}$ -linear residue annihilator stable for the Hamiltonian coadjoint action. The ambient  $R$ -submodule is  $E_R(\mathbb{C})$ , and quotienting by the non-invariant line  $\mathbb{C} \rho_{0,0}$  is a different operation. Define the Hamiltonian action by negative transpose of the proved boundary Hamiltonian bracket:

$$\langle \{O_f, \Theta\}, O_g \rangle = -\langle \Theta, \{O_f, O_g\}_{\text{open}} \rangle.$$

Then this stalkwise local-dual sector realizes the principal-part coadjoint action:

$$\{O_{p,q}, \Theta_{a,b}\}_{\text{open}} = -(pb - qa + p - q)\Theta_{a-p+1, b-q+1},$$

with negative-index targets zero. Declare the odd local-dual ideal abelian:

$$\{\Theta_{a,b}, \Theta_{c,d}\}_{\text{open}} = 0.$$

Equivalently, the assignment

$$z_1^p z_2^q \mapsto O_{p,q}, \quad \rho_{a,b}[1] \mapsto \Theta_{a,b}$$

realizes stalkwise the Lie algebra  $\tilde{A} \ltimes \mathcal{P}[1]$ .

**THEOREM 5.185** (*Support-local principal-part current  $P_0$  prefactorization datum*). After the de Rham contraction to the reduced local model on the brane line, the local-dual sector of Construction 5.184 extends to a support-local  $P_0$  prefactorization datum on  $\mathbb{R}$ . In this theorem the de Rham differential has already been contracted; the Hamiltonian smearing uses degree-zero test functions, and the cotangent summand uses compactly supported distributions. Define the prefactorization datum: for each open interval  $I \subset \mathbb{R}$  put

$$\mathfrak{S}_\partial^{\text{pp}}(I) = \mathcal{D}'_c(I) \widehat{\otimes} \mathcal{P}[1], \quad \mathfrak{S}_\partial^{\text{Ham}}(I) = \Omega_c^\circ(I) \widehat{\otimes} \tilde{A},$$

and

$$A_\partial^{\text{Ham}}(I) = \widehat{\mathfrak{S}}(\mathfrak{S}_\partial^{\text{Ham}}(I)), \quad A_\partial^{\text{pp}}(I) = A_\partial^{\text{Ham}}(I) \widehat{\otimes} \widehat{\mathfrak{S}}(\mathfrak{S}_\partial^{\text{pp}}(I)).$$

In this theorem the open factor is this reduced connected Hamiltonian current algebra  $A_\partial^{\text{Ham}}(I)$ . The displayed brackets define the reduced Hamiltonian/principal-part current bracket. Brackets with arbitrary reduced or unreduced open BV observables require the unreduced lift of Remark 5.189. Write

$$\text{Obs}_{\partial, N}^{\text{cl, pp}}(I) = A_\partial^{\text{pp}}(I),$$

where  $\widehat{\mathfrak{S}}$  is the degreewise completed symmetric algebra in the chosen current category: each finite symmetric degree is formed before completion, and the  $P_0$  biderivation is extended continuously from generators. The construction uses this degreewise completed symmetric-algebra functor. The datum is equipped with extension by zero for inclusions of intervals and with the following structures:

- (i) the differential  $Q_{\text{open}}$  on the open factor and zero on  $\mathfrak{S}_\partial^{\text{pp}}(I)$ ;
- (ii) the factorization product on disjoint intervals  $I_1 \sqcup I_2 \subset J$  given by extension by zero followed by graded-commutative multiplication;
- (iii) the cosheaf structure on  $I \mapsto \Omega_c^\circ(I)$  and  $I \mapsto \mathcal{D}'_c(I)$  given by extension by zero;
- (iv) the smearing maps

$$B_f(a) = \int_I a(t) B_f(t) dt, \quad O_f(a) = a \otimes f, \quad \Theta_\rho(B) = B \otimes \rho[1],$$

with the convention that  $\Theta_\rho(B)$  pairs only with the primitive connected Hamiltonian trace projection by  $\langle \Theta_\rho(B), B_g(b) \rangle = B(b) \rho(g)$ , and descendants, constants, and decomposable multi-traces lie outside this pairing unless an additional coalgebra projection is specified;

(v) the  $P_0$  Poisson bracket

$$\begin{aligned} \{B_f(a), B_g(b)\} &= B_{\{f,g\}}(ab), \\ \{B_f(a), \Theta_\rho(B)\} &= \Theta_{f \cdot \rho}(aB), \\ \{\Theta_\rho(B), \Theta_\sigma(C)\} &= 0, \end{aligned}$$

where  $a, b \in \Omega_c^\circ$ ,  $B, C \in \mathcal{D}'_c$ , and  $aB$  is the multiplication of a current by a smooth degree-zero form;

(vi) the smeared current morphism

$$\begin{aligned} \widetilde{\partial}_{\text{bb}, I}: \Omega_c^\circ(I) \otimes \bar{A} \oplus \mathcal{D}'_c(I) \otimes \mathcal{P}[1] &\longrightarrow A_\partial^{\text{pp}}(I), \\ a \otimes f &\longmapsto B_f(a), \quad B \otimes \rho[1] \longmapsto \Theta_\rho(B). \end{aligned}$$

This degreewise datum defines a support-local  $P_0$  prefactorization datum on  $\mathbb{R}$ . On the basis  $\Theta_{i,j}$  and for regular densities  $B = b(t) dt$  the second bracket reads

$$\{O_{p,q}(a), \Theta_{i,j}(b)\} = -(pj - qi + p - q) \Theta_{i-p+1, j-q+1}(ab),$$

again with negative-index targets zero. The smeared map  $\widetilde{\partial}_{\text{bb}, I}$  is the interval-level morphism into the reduced connected Hamiltonian  $P_0$  current sector. An unsmeared canonical map on an interval requires a chosen compactly supported unit or density.

*Proof. Cosheaf axioms.* Both  $\Omega_c^\circ$  and  $\mathcal{D}'_c$  are cosheaves on  $\mathbb{R}$  for the extension-by-zero structure. Tensoring with the fixed graded vector spaces  $\bar{A}$  and  $\mathcal{P}[1]$  preserves the cosheaf property because the tensor factors are discrete graded vector spaces. Taking the degreewise completed symmetric algebra gives the prefactorization products used here: on disjoint intervals the structure map is extension by zero followed by the graded-commutative product, and the  $P_0$  bracket is the continuous biderivation generated by the displayed current brackets. The odd generators  $\Theta_\rho$  therefore generate the completed graded-symmetric, equivalently exterior, principal-part factor.

*Disjoint-support locality.* For disjoint  $I_1, I_2 \subset J$ , the multiplication map

$$\text{Obs}_{\partial, N}^{\text{cl}, \text{pp}}(I_1) \otimes \text{Obs}_{\partial, N}^{\text{cl}, \text{pp}}(I_2) \longrightarrow \text{Obs}_{\partial, N}^{\text{cl}, \text{pp}}(J)$$

is the graded-commutative product after extending by zero. This is the standard prefactorization product. Because the bracket  $\{O_f(a), \Theta_\rho(B)\} = \Theta_{f \cdot \rho}(aB)$  involves multiplication of a smooth function by a distribution and is supported on  $\text{supp } a \cap \text{supp } B$ , disjoint  $I_1, I_2$  have disjoint such supports and the corresponding bracket vanishes; the factorization product is therefore Poisson-commutative on disjoint intervals, as required for the  $P_0$  structure.

*Q-compatibility.*  $Q_{\text{open}} B_f = 0$  on Hamiltonian trace classes by Proposition 3.39. By declaration  $Q_{\text{open}} \Theta_\rho = 0$  on the  $\Theta_\rho$ -factor. Therefore the differential is zero on the  $\Theta_\rho$ -factor and acts trivially on the bracket structure displayed above. This is consistent with the reduced model after de Rham-current contraction.

*$P_0$  identities.* The first bracket  $\{B_f(a), B_g(b)\} = B_{\{f,g\}}(ab)$  is Theorem 5.190 smeared by test functions, supplied by Corollary 5.191. The second bracket  $\{B_f(a), \Theta_\rho(B)\} = \Theta_{f \cdot \rho}(aB)$  is the negative



transpose under the residue pairing of Theorem 5.176 applied to the first bracket; in coordinates with  $\rho = \rho_{a',b'}$  and  $f = z_1^p z_2^q$  this reads

$$\{O_{p,q}(a), \Theta_{a',b'}(B)\} = -(pb' - qa' + p - q) \Theta_{a'-p+1, b'-q+1}(aB),$$

exactly the principal-part coadjoint formula of Theorem 5.178. The third bracket  $\{\Theta_\rho(B), \Theta_\sigma(C)\} = 0$  is by declaration of the abelian odd ideal.

*Jacobi identity.* The semidirect Lie algebra  $\tilde{A} \ltimes \mathcal{P}[1]$  has Jacobi identity  $[\text{ad}_f^\vee, \text{ad}_g^\vee] = \text{ad}_{\{f,g\}}^\vee$  for  $f, g \in \tilde{A}$  on the continuous dual. Smearing by test functions and distributions preserves Jacobi because the  $C^\infty(\mathbb{R})$ -module action on distributions is associative. Therefore the  $P_0$  bracket on the smeared sector also satisfies Jacobi.

*Bulk-boundary morphism.* The map  $\tilde{\partial}_{\text{bb},I}$  sends  $a \otimes f$  to the smeared polynomial trace observable  $B_f(a)$  by Theorem 5.190, and sends  $B \otimes \rho[1]$ , with  $\mathcal{P}[1] \cong \mathfrak{h}_{\text{cont}}^\vee[1]$  via Theorem 5.178, to the  $\Theta_\rho(B)$ -sector. The brackets just verified are exactly the smeared semidirect brackets on the degree-zero line-current version of  $\tilde{A} \ltimes \mathcal{P}[1]$ . Therefore  $\tilde{\partial}_{\text{bb},I}$  is a morphism into the reduced connected Hamiltonian  $P_0$  current sector.  $\square$

LEMMA 5.186 (*Distribution-locality criterion for principal-part currents*). Let  $a \in C_c^\infty(I)$ ,  $B \in \mathcal{D}'(I)$ , and  $\rho, \sigma \in \mathcal{P}$ . The product  $aB$  used in Theorem 5.185 is the standard multiplication of a distribution by a smooth function,

$$(aB)(\varphi) = B(a\varphi), \quad \varphi \in C_c^\infty(I).$$

With this convention the reduced principal-part brackets are well-defined by

$$\{B_f(a), \Theta_\rho(B)\} = \Theta_{f \cdot \rho}(aB), \quad \{\Theta_\rho(B), \Theta_\sigma(C)\} = 0.$$

Multiplication of two distributions is avoided. This is support-locality in the reduced line-current model. Costello-local functional locality and wavefront/RG-admissibility are separate analytic data.

*Proof.* Multiplication  $B \mapsto aB$  is a continuous endomorphism of  $\mathcal{D}'(I)$ , because  $a\varphi \in C_c^\infty(I)$  for every test function  $\varphi$ . The first displayed bracket therefore lands again in  $\mathcal{D}'(I) \widehat{\otimes} \mathcal{P}[1]$ . Its support is contained in  $\text{supp}(a) \cap \text{supp}(B)$ , so it is local on the line. The second bracket is zero by the abelian odd-ideal definition of the reduced principal-part sector; the product  $BC$  is omitted. Jacobi with two Hamiltonian entries and one principal-part entry follows from associativity  $a(bB) = (ab)B$  and from the coadjoint identity  $[f \cdot, g \cdot]\rho = \{f, g\} \cdot \rho$ . Jacobi with two principal-part entries is immediate from the zero bracket.  $\square$

*Remark 5.187 (Reduced principal-part currents).* Theorem 5.185 is a reduced support-local  $P_0$  prefactorization datum after de Rham-current contraction, with  $\Omega_c^\circ(I)$  in the Hamiltonian current factor and  $\mathcal{D}'(I)$  in the cotangent factor. The raw distributional target has additional degree-zero classes under interval enlargement: if  $I \subsetneq J$ , a delta-current supported in  $J \setminus I$  lies in  $\mathcal{D}'(J)$  and outside the image of extension from  $\mathcal{D}'(I)$ . The unreduced BV/factorization algebra is separate. In particular, the equation  $Q\Theta_\rho = 0$  is part of the added reduced local-dual factor; an unreduced derivation requires an unreduced de Rham current complex, a chain-level primitive projection, and Hochschild centrality homotopies against arbitrary open observables.

**THEOREM 5.188** (*Reduced prefactorization realization of the boundary local dual*). After de Rham contraction in the reduced setting, the assignment

$$\rho_{a,b}[1] \mapsto \Theta_{a,b}$$

is a chain-level realization of the boundary local dual within the support-local  $P_0$  prefactorization datum  $\text{Obs}_{\partial,N}^{\text{cl,pp}}$  of Theorem 5.185. The intervalwise smeared current morphism  $\widetilde{\partial}_{\text{bb},I}$  extends, in this reduced principal-part target, the Hamiltonian bulk-to-boundary Lie algebra map of Theorem 5.190 to the closed-coadjoint sector  $\mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^{\vee}[1]$ .

*Proof.* Theorem 5.185 supplies the support-local  $P_0$  prefactorization datum and verifies that  $\widetilde{\partial}_{\text{bb},I}$  is a smeared current Lie algebra morphism for each interval  $I$ . The identification  $\rho_{a,b}[1] \leftrightarrow \Theta_{a,b}$  is by the basis of the local-dual factor, consistent with Construction 5.184. The closed source side  $\mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^{\vee}[1]$  is the semidirect product computed in Proposition 2.7, and its image in the support-local  $P_0$  prefactorization datum is the displayed Lie subalgebra  $\tilde{A} \oplus \mathcal{P}[1]$ .  $\square$

**Remark 5.189** (*Unreduced lift obstruction*). Theorem 5.188 resolves the boundary local-dual prefactorization realization at the reduced level after de Rham-current contraction, with the cotangent current differential already contracted. An unreduced boundary BV/factorization lift requires: define the compactly supported de Rham functoriality before contraction, the full BV differential and homotopies, the pairing with ordinary open observables before primitive trace projection, and the coherent  $P_0$ -center homotopies.

The candidate unreduced enlargement has the form

$$\widetilde{\text{Obs}}_{\partial,N}^{\text{cl,pp}}(I) = \text{Obs}_{\text{open},N}^{\text{cl}}(I) \widehat{\otimes} \widehat{\mathbf{S}}(\mathcal{M}_{\partial}^{\text{pp,unred}}(I)),$$

where

$$\mathcal{M}_{\partial}^{\text{pp,unred}}(I) = C_{c,\text{cur}}^{\text{dR}}(I) \widehat{\otimes} \text{Ann}_{\text{res}}(\mathbb{C} \cdot 1)[1].$$

Here  $C_{c,\text{cur}}^{\text{dR}}(I)$  is the compactly supported de Rham current cosheaf on the line, with differential dual to  $d_{\mathbb{R}}$ , and  $\text{Ann}_{\text{res}}(\mathbb{C} \cdot 1)$  is the principal-part residue annihilator of constants. One must also construct a chain-level primitive/indecomposable projection

$$\pi_{\text{prim}} : \text{Obs}_{\text{open},N}^{\text{cl}}(I) \longrightarrow \Omega_c^{\bullet}(I) \widehat{\otimes} \tilde{A}$$

compatible with  $Q_{\text{open}}$ , constants, decomposable multi-traces, and factorization products. The required homotopies are the centrality homotopies for  $\Theta_{\rho}$  against arbitrary unreduced open observables and the  $P_0$  bracket homotopies

$$\{B_f, \Theta_{\rho}\} \simeq \Theta_{f \cdot \rho}, \quad \{\Theta_{\rho}, \Theta_{\sigma}\} \simeq 0$$

before primitive projection. Only after this enlargement does the full cotangent factorization-center lift extend the reduced realization  $\rho_{a,b}[1] \mapsto \Theta_{a,b}$  to a chain-level morphism on the unreduced sector. The unreduced lift is the quantum-level homotopical lift beyond the classical reduced theorem above; it sits in the residual Problem 5.221.

**THEOREM 5.190** (*Classical bulk-to-boundary Hamiltonian map*). In the stable connected Hamiltonian trace sector of the formal local mixed holomorphic-topological theory, the boundary evaluation maps  $\partial_{\text{bb},N}$  of Construction 3.25 stabilize to a linear map

$$\partial_{\text{bb}} : \tilde{A} \longrightarrow H_{\text{Ham}}^0(O_{\text{brane}}^{\text{cl}}), \quad f \mapsto \overline{\text{Tr } f(\phi_1, \phi_2)}.$$

This map is a morphism of Lie algebras from the closed Hamiltonian Poisson algebra  $\bar{A}$  to the connected open Hamiltonian trace Poisson algebra:

$$\{\partial_{\text{bb}}(f), \partial_{\text{bb}}(g)\}_{\text{open}} = \partial_{\text{bb}}(\{f, g\}_{\bar{A}}).$$

This is the degree-zero Hamiltonian bracket component of the reduced CE/PV central-operation map to the brane observables. The full Poisson factorization-center morphism is a separate construction.

*Proof.* The map is well-defined on  $\bar{A}$  by Corollary 3.32; constants map to the scalar  $N$  and vanish in the Hamiltonian quotient by constants. Lemma 5.169 gives the open Hamiltonian trace Poisson bracket. For  $f = z_1^k z_2^l$  and  $g = z_1^m z_2^n$ , Proposition 5.170 gives

$$\{\partial_{\text{bb}}(f), \partial_{\text{bb}}(g)\}_{\text{open}} = \overline{(kn - lm) \text{Tr}(\phi_1^{k+m-1} \phi_2^{l+n-1})}.$$

The closed Poisson bracket is

$$\{f, g\}_{\bar{A}} = \overline{(kn - lm) z_1^{k+m-1} z_2^{l+n-1}},$$

and applying  $\partial_{\text{bb}}$  gives the same expression. Bilinearity proves the identity for all polynomials.  $\square$

**COROLLARY 5.191** (*Classical local bulk-boundary coupling with explicit  $U(1)$  centre anomaly*). Let  $a, b \in \Omega_c^\circ(\mathbb{R})$  be scalar compactly supported test functions on the brane line, and for  $f \in \bar{A}$  define the smeared boundary Hamiltonian

$$I_\partial(a, f) = B_f(a) = \int_{\mathbb{R}} a(t) \overline{\text{Tr} f(\phi_1(t), \phi_2(t))} dt \in A_\partial^{\text{Ham}}(\mathbb{R}).$$

Then the following two strengthenings of the bracket identity hold jointly.

(i) *Strict  $P_0$  identity on disjoint supports.* In the  $P_0$  bracket of  $A_\partial^{\text{Ham}}$ ,

$$\{I_\partial(a, f), I_\partial(b, g)\}_{A_\partial^{\text{Ham}}} = I_\partial(ab, \{f, g\}_{\bar{A}}),$$

and the bracket vanishes whenever the supports of  $a$  and  $b$  are disjoint. The locality input is the canonical equal-time bracket  $\{(\phi_1)^i{}_j(t), (\phi_2)^k{}_l(t')\} = \delta_l^i \delta_j^k \delta(t - t')$  of 4.10 on two brane-time points; smearing against  $a(t)b(t')$  produces the pointwise product  $ab$  after integration of the delta function. This is the level-one component of 4.11.

(ii) *Explicit  $U(1)$  centre anomaly constant.* Before projection to the scalar-reduced quotient, the unreduced open bracket carries an explicit central piece

$$\{I_\partial(a, f), I_\partial(b, g)\}_{\text{open}} = I_\partial(ab, \{f, g\}_{\bar{A}}) + N \cdot \omega(f, g) \int_{\mathbb{R}} a(t)b(t) dt,$$

where  $\omega$  is the cocycle of 6.2 and the prefactor  $N$  is the rank of the brane stack and is the value of the  $U(1)$  centre character  $\text{Tr} : \mathbb{C} \cdot I \rightarrow \mathbb{C}, I \mapsto N$ . After projection to  $H_{\text{red}}^\circ = H^\circ / \mathbb{C} \cdot \text{Tr}(1)$  the second term vanishes, and one recovers

$$\overline{\{I_\partial(a, f), I_\partial(b, g)\}_{\text{open}}} = \overline{I_\partial(ab, \{f, g\}_{\bar{A}})}.$$

Statements (i) and (ii) are compatible: (ii) is the unreduced refinement, (i) is the level-one factorization-algebra restatement on the scalar-reduced sector. The equal-time delta locality of (i) produces a strict  $P_0$  factorization algebra, while (ii) records the homotopical  $U(1)$  centre anomaly that obstructs lifting (i) to the unreduced sector.

*Proof. Proof of (i).* By 4.7, the  $P_\partial$  bracket on  $A_\partial^{\text{Ham}}$  is the biderivation with  $\{a \otimes f, b \otimes g\}_{A_\partial} = ab \otimes \{f, g\}_{\bar{A}}$  on length-one generators, so  $\{B_f(a), B_g(b)\}_{A_\partial^{\text{Ham}}} = B_{\{f, g\}}(ab) = I_\partial(ab, \{f, g\}_{\bar{A}})$ . 4.9 then gives vanishing on disjoint supports because  $ab \equiv 0$  pointwise. The same identity is the smeared content of the equal-time delta-function bracket of 4.10: tracing  $f(\phi_1, \phi_2)(t)$  and  $g(\phi_1, \phi_2)(t')$ , applying the Leibniz rule, and integrating against  $a(t)b(t')\delta(t - t')$  produces the pointwise product  $ab$  after the delta-function integration.

*Proof of (ii).* The unreduced bracket of two integrals is, by locality of the canonical bracket and 5.190,

$$\begin{aligned} \{I_\partial(a, f), I_\partial(b, g)\} &= \iint_{\mathbb{R}^2} a(t)b(t') \{\text{Tr } f(\phi(t)), \text{Tr } g(\phi(t'))\} dt dt' \\ &= \int_{\mathbb{R}} a(t)b(t) \text{Tr}(\{f, g\}_{\bar{A}}) dt + \int_{\mathbb{R}} a(t)b(t) \omega(f, g) N dt, \end{aligned}$$

where the splitting is by the central decomposition of 6.4: the central scalar  $\omega(f, g) \cdot N$  is removed by the projection to  $H_{\text{red}}^\circ$ , and the remaining piece is  $I_\partial(ab, \{f, g\}_{\bar{A}})$  exactly. The constants are explicit: the prefactor of the central piece is  $N \cdot \omega(f, g)$ , and the smeared coefficient is  $\int ab dt$ . The classical-to-quantum reduction of 6.6 identifies  $\omega(f, g) \cdot N$  with the leading-order Capelli shift  $\hbar N$  on the linear generators  $f = z_1, g = z_2$ . Compact support of  $a, b$  kills boundary terms.  $\square$




FIGURE 0.1: Notation diagram for the reduced open-line order- $\hbar$  raw commutator graph  $\Gamma_1$ . Two ordered boundary insertions sit on the brane line  $\mathbb{R}$  at distinct times  $t, t'$ ; the unique propagator  $P_\partial$  contracts one  $z_1$ -leg on one side against one  $z_2$ -leg on the other (and the antisymmetric partner). The automorphism group is trivial. In the Weyl ordered product the ordered one-edge weight is  $\hbar P_\partial/2$ ; after subtracting the reverse order the commutator weight is  $\hbar P_\partial$ . The bivector contraction on monomials reproduces the cocycle coefficient  $(kn - lm)$  of Proposition 5.170, the order- $\hbar$  Moyal coefficient of Proposition 5.215, and the  $U(1)$  centre cocycle  $\omega(f, g)$  of Lemma 6.2 realised at multidegree  $(k + m, l + n) = (1, 1)$ .

The graph  $\Gamma_1$  of Figure 0.1 records the single canonical contraction producing the coefficient  $(kn - lm)$  in Proposition 5.170. An independent Feynman-integral proof would require Costello graph data.

### 5.7.2 QUANTIZATION AS A BRANE TRACE-MAP TEST

Three operations are at stake: Moyal star-quantisation of Hamiltonians on the formal symplectic polydisk, Weyl quantisation of the matrix entries of  $(\phi_1, \phi_2)$ , and the brane trace map  $J: f \mapsto \text{Tr } f(\phi_1, \phi_2)$

extended to a quantum trace map  $\Phi_h$ . The question is whether they commute up to identifiable obstruction classes. Theorem 5.217 names the four components on which compatibility is proved: weighted BV kernel convergence, balanced-supertrace scalar QME cancellation on its completion, weighted Moyal principal-part current brackets, and an explicit degree-zero Hamiltonian trace comparison. The remaining obstructions are inscribed sharply: the all-bidegree radial/Weyl normal-form question is the single cokernel class  $\Omega_{a,b}^{\text{rad}} \in \text{coker } B_{a,b}$  (Proposition F.25); the reduced non-scalar Costello QME question is the single cohomology class  $[\text{Ob}_{w,\partial,h}^{\text{red}}] \in H^1(\ker \widehat{\sigma}_{\text{sc}}, d_{\text{QME}})$  (Problem 5.219); the smallest non-scalar window  $\theta_3$  splits into a bare  $H^1 = \mathbb{C} \cdot E_{\theta_3} \neq 0$  and a supplied-counterterm  $H^1 = 0$  with  $c = 1$  unique (Theorem E.38).

*The  $U(1)$  centre-of-mass cocycle.* On the closed side, the Hamiltonian vector field of every constant is zero; the further quotient  $\tilde{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1$  removes the unit from the source. On the brane side, the unit  $1 \in \mathbb{C}[z_1, z_2]$  lifts to  $\text{Tr}(I_N) = N$ , the rank of the brane stack: the center-of-mass mode of the matrix coordinate, invisible to Hamiltonian reduction. The two linear generators  $\text{Tr}(\phi_1)$  and  $\text{Tr}(\phi_2)$  are the open-side images of  $z_1$  and  $z_2$ , and on them the unreduced matrix Poisson bracket reads

$$\{\text{Tr } \phi_1, \text{Tr } \phi_2\} = \text{Tr}(I_N) = N,$$

while  $\{z_1, z_2\}_{\tilde{A}} = 0$  modulo constants. Quantising the same linear generators by the Moyal star and reading the Weyl commutator on the matrix-brane realization,

$$[\Phi_h(z_1), \Phi_h(z_2)]_{\star} = \hbar N \cdot I_N \quad \text{in } M_N(\mathbb{C}),$$

detects on the closed side the Lie two-cocycle  $\bar{c} \in H_{\text{Lie}}^2(\tilde{A}, \mathbb{C})$  with  $\bar{c}(f, g) = [\{f, g\}]_{\circ}$ , and on the brane side reproduces the Capelli contact term in the reduced ordinary scalar  $\mathfrak{gl}_N$  branch [26, 27]. The three faces of the same scalar obstruction — Lie 2-cocycle on  $\tilde{A}$ , Poisson bracket of trace coordinates, and quantum Moyal–Weyl commutator — realise the same class  $\hbar N[\bar{c}] \in H_{\text{Lie}}^2(\tilde{A}, \mathbb{C})[[\hbar]]$  (Theorems 6.3, 6.4, 6.6, E.13). The cancellation branches are catalogued in Theorem 5.213: the unreduced source, the central-character source, the opposite central extension, the balanced supertrace, the algebraic central-contact cone, and the Costello-local closed-exchange criterion. The all-order Moyal/Weyl identity that promotes the linear-generator computation is the constant-Poisson deformation quantisation [7].

The formal Moyal target is the unique translation-invariant deformation of the constant Poisson tensor on the formal holomorphic polydisk. It fixes the algebraic coefficients that any quantum closed-open BV construction must recover. Construction of that BV theory is separate. The classical degree-zero Hamiltonian bracket is the proposed special fibre; the cotangent/descendant part is recorded as Problem 5.221. Put

$$A_h = (\mathbb{C}[[z_1, z_2]][[\hbar]], *),$$

where  $*$  is the Weyl/Moyal product

$$f * g = m \circ \exp\left(\frac{\hbar}{2}(\partial_{z_1} \otimes \partial_{z_2} - \partial_{z_2} \otimes \partial_{z_1})\right)(f \otimes g).$$

The symbol  $\tilde{A}_h$  denotes the central quotient of the underlying Moyal Lie algebra by  $\mathbb{C}[[\hbar]]$ ; the associative algebra quotient is a different object. Set

$$[f, g]_{\text{raw}} = f * g - g * f, \quad \{f, g\}_h = \hbar^{-1}[f, g]_{\text{raw}}.$$

The degree-zero quantum Hamiltonian Lie algebra is

$$\mathfrak{h}_{\hbar} = \bar{A}_{\hbar}, \quad [f, g]_{\mathfrak{h}_{\hbar}} = \{f, g\}_{\hbar}.$$

The specialization at  $\hbar = 0$  is the Poisson Lie algebra  $\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$  used above.

On the brane side, the open action

$$\int_{\mathbb{R}} \text{Tr}(\phi_1 d\phi_2)$$

makes the matrix entries of  $\phi_1$  and  $\phi_2$  conjugate variables. The canonical algebraic quantization of the unconstrained first-order matrix phase space is the completed Weyl algebra  $\mathcal{W}_N$  generated by those entries with

$$[(\phi_1)^i_j, (\phi_2)^k_l] = \hbar \delta_l^i \delta_j^k, \quad [(\phi_a)^i_j, (\phi_a)^k_l] = 0 \quad (a = 1, 2).$$

The gauge field imposes the conjugation moment-map constraint. With the commutator convention above, the normal-ordered quantum comoment is

$$\widehat{\mu}_{\epsilon} = -\text{Tr}([\epsilon, \phi_1] \phi_2), \quad \epsilon \in \mathfrak{gl}_N.$$

This formula is kept in normal order: for  $\epsilon = 1$  it vanishes, as the scalar subgroup acts trivially. The induced infinitesimal conjugation action on the Weyl algebra is fixed by

$$\hbar^{-1}[\widehat{\mu}_{\epsilon}, a] = \epsilon \cdot a;$$

changing to the opposite commutator convention changes the displayed comoment by the corresponding sign. Since the scalar comoment is zero, the effective constraint algebra is  $\mathfrak{pgl}_N$ , represented by  $\mathfrak{sl}_N$  after choosing the trace pairing. The quantum brane local algebra used here is the reduced BRST, or degree-zero normalizer, quantum Hamiltonian reduction

$$\mathcal{O}_{\text{brane}, N}^q = C_{\text{BRST}}^{\bullet}(\mathfrak{sl}_N, \mathcal{W}_N, \widehat{\mu}),$$

whose degree-zero cohomology is the normalizer quotient

$$H^0 \mathcal{O}_{\text{brane}, N}^q = N_{\mathcal{W}_N}(\mathcal{W}_N \widehat{\mu}(\mathfrak{sl}_N)) / \mathcal{W}_N \widehat{\mu}(\mathfrak{sl}_N).$$

The constant Hamiltonian trace is projected out in degree zero as before. The classical special fibre is the reduced BV algebra of Construction 3.20.

The quantum boundary map is Weyl-ordered trace evaluation:

$$\partial_{\text{bb}, \hbar, N}(f) = \overline{\text{Tr Op}_{\mathcal{W}}(f)(\phi_1, \phi_2)}, \quad f \in \bar{A}_{\hbar},$$

where  $\text{Op}_{\mathcal{W}}$  denotes total symmetrisation before passing to the trace. At  $\hbar = 0$  this recovers Construction 3.25. After dividing the commutator by  $\hbar$  and reducing modulo  $\hbar$ , the raw commutator identity recovers Theorem 5.190.

**PROPOSITION 5.192** (*Reduced BRST descent of Weyl-ordered Hamiltonian traces*). For every  $f \in \bar{A}_{\hbar}$ , the Weyl-ordered trace

$$F_{f, N} = \text{Tr Op}_{\mathcal{W}}(f)(\phi_1, \phi_2)$$

is invariant under the quantum comoment:

$$[\widehat{\mu}_{\epsilon}, F_{f, N}] = 0, \quad \epsilon \in \mathfrak{gl}_N.$$

Hence  $F_{f, N}$  lies in the normalizer of the quantum moment-map ideal and defines a degree-zero class in  $H^0 \mathcal{O}_{\text{brane}, N}^q$  after removing the scalar Hamiltonian.

*Proof.* Total Weyl symmetrisation is equivariant for linear changes of the matrix variables, and simultaneous conjugation acts linearly on  $(\phi_1, \phi_2)$ . Therefore  $\text{Tr Op}_W(f)(\phi_1, \phi_2)$  is invariant under infinitesimal conjugation. By the defining property of the quantum comoment,

$$\hbar^{-1}[\widehat{\mu}_\epsilon, a] = \epsilon \cdot a,$$

this invariance is exactly  $[\widehat{\mu}_\epsilon, F_{f,N}] = 0$ . If  $I = \mathcal{W}_N \widehat{\mu}(\mathfrak{sl}_N)$  is the left quantum moment-map ideal, then for every generator  $\widehat{\mu}_\epsilon, \widehat{\mu}_\epsilon F_{f,N} = F_{f,N} \widehat{\mu}_\epsilon \in I$ . Thus  $F_{f,N}$  normalizes  $I$  and descends to the normalizer quotient describing  $H^0 \mathcal{O}_{\text{brane},N}^q$ .  $\square$

**PROPOSITION 5.193** (*Reduced BRST descent of arbitrary Weyl-ordered Hamiltonian products*). For every finite collection  $f_1, \dots, f_r \in \tilde{A}_\hbar$ , the ordered product

$$F = \prod_{i=1}^r F_{f_i,N}, \quad F_{f_i,N} = \text{Tr Op}_W(f_i)(\phi_1, \phi_2),$$

is invariant under the quantum comoment:

$$[\widehat{\mu}_\epsilon, F] = 0, \quad \epsilon \in \mathfrak{gl}_N.$$

Equivalently, every monomial in the Weyl-ordered single-trace generators normalizes the quantum moment-map ideal and defines a degree-zero class in  $H^0 \mathcal{O}_{\text{brane},N}^q$  after removing the empty trace.

*Proof.* By the convention  $\hbar^{-1}[\widehat{\mu}_\epsilon, a] = \epsilon \cdot a$ , the operator  $\hbar^{-1} \text{ad } \widehat{\mu}_\epsilon$  acts on the associative algebra  $\mathcal{W}_N$  by infinitesimal  $GL_N$ -conjugation, and is therefore a derivation of the associative product. By Proposition 5.192, each individual factor  $F_{f_i,N}$  is annihilated by this derivation. Derivations of an associative algebra preserve products: the Leibniz expansion of  $\hbar^{-1} \text{ad } \widehat{\mu}_\epsilon$  on the ordered product  $\prod_{i=1}^r F_{f_i,N}$  is a sum of summands each containing a factor  $\hbar^{-1}[\widehat{\mu}_\epsilon, F_{f_i,N}] = 0$ . Hence  $[\widehat{\mu}_\epsilon, F] = 0$ , and as in the proof of Proposition 5.192,  $\widehat{\mu}_\epsilon F = F \widehat{\mu}_\epsilon \in \mathcal{W}_N \widehat{\mu}(\mathfrak{sl}_N)$ , so  $F$  normalizes the left moment-map ideal and descends.  $\square$

*Remark 5.194.* The argument uses only that  $\hbar^{-1} \text{ad } \widehat{\mu}_\epsilon$  is a derivation. The Capelli shift obstructs the bracket identity between traces while their joint reduced BRST descent follows from derivation invariance.

*Remark 5.195* (*Normal-ordering obstruction*). Proposition 5.192 proves reduced BRST descent. The quantum Moyal theorem requires the bracket identity. Put  $X = \phi_1$  and  $Y = \phi_2$ . With the normal ordering above,

$$\widehat{\mu}_\epsilon = -\text{Tr}([\epsilon, X]Y) = \text{Tr}(\epsilon(YX - XY + \hbar NI)).$$

For  $\epsilon = I$  this comoment is identically zero; equivalently the displayed matrix has trace zero inside the Weyl algebra. Thus the nontrivial quantum moment equations are represented in the quantum Hamiltonian reduction by the traceless matrix congruence

$$YX - XY + \hbar NI \equiv 0 \pmod{\mathcal{W}_N \widehat{\mu}(\mathfrak{sl}_N)},$$

whose special fibre is the classical commuting-pair equation. Consequently, for a covariant matrix word  $A$ ,

$$\text{Tr}(AXY) - \text{Tr}(AYX) = \hbar N \text{Tr}(A) \pmod{\mathcal{W}_N \widehat{\mu}(\mathfrak{sl}_N)}.$$

This Capelli-type shift can feed lower-degree traces into the connected sector. A stable quantum theorem must prove that the chosen connected extraction removes or renormalizes this shift, or else change the sector or normalization.

*Remark 5.196 (Capelli-renormalized trace generators).* Let

$$T_{a,b} = \text{Tr}(X^a Y^b)$$

denote the normal-ordered trace. The Weyl-ordered trace

$$J_N(z_1^a z_2^b) = \text{Tr Op}_W(z_1^a z_2^b)(X, Y)$$

is triangular over the normal-ordered traces after imposing the quantum moment relation:

$$J_N(z_1^a z_2^b) \equiv \sum_{r=0}^{\min(a,b)} \left(-\frac{\hbar N}{2}\right)^r r! \binom{a}{r} \binom{b}{r} T_{a-r, b-r}.$$

For example,

$$J_N(z_1 z_2) = T_{1,1} - \frac{\hbar N^2}{2}, \quad J_N(z_1^2 z_2) = T_{2,1} - \hbar N T_{1,0},$$

and

$$J_N(z_1 z_2^2) = T_{1,2} - \hbar N T_{0,1}, \quad J_N(z_1^2 z_2^2) = T_{2,2} - 2\hbar N T_{1,1} + \frac{\hbar^2 N^3}{2}.$$

The normal-ordered basis carries the Capelli-shifted commutator already in low degree; a direct calculation gives

$$\hbar^{-1}[T_{2,1}, T_{0,2}] = 4T_{1,2} - 2\hbar N T_{0,1} \mod \mathcal{W}_N \widehat{\mu}(\mathfrak{sI}_N),$$

while the scalar Moyal bracket records the leading  $4T_{1,2}$ . The connected, or primitive trace, projection retains the Capelli shift: it removes disconnected products and the empty trace, leaving the nonempty primitive trace  $T_{0,1}$  in the residual. The stable connected quantum theorem therefore uses normal-ordered primitive traces with the triangular  $\hbar N$  correction built into the definition. The natural finite- $N$  generators are the Capelli-renormalized Weyl traces  $J_N(f)$ , equivalently

$$J_N(f) = T_N \left( \exp \left( -\frac{\hbar N}{2} \partial_{z_1} \partial_{z_2} \right) f \right)$$

in the normal-ordered trace notation  $T_N(z_1^a z_2^b) = T_{a,b}$ .

**LEMMA 5.197 (Capelli-renormalized stable connected trace).** Let  $\text{Tr}_N^{\text{ren}}(f) = J_N(f)$  for  $f \in \mathbb{C}[z_1, z_2][[\hbar]]$ . Write  $I_N = \mathcal{W}_N \widehat{\mu}(\mathfrak{sI}_N)$  for the left moment-map ideal. Then:

- (i) the map  $\text{Tr}_N^{\text{ren}}: \bar{\mathcal{A}}_{\hbar} \longrightarrow \mathcal{W}_N/I_N$  is well defined modulo the empty (constant) trace;
- (ii) the inverse triangular system reads

$$T_{a,b} = \sum_{r=0}^{\min(a,b)} \left(\frac{\hbar N}{2}\right)^r r! \binom{a}{r} \binom{b}{r} J_N(z_1^{a-r} z_2^{b-r}),$$

so  $T_{a,b}$  and  $J_N(z_1^a z_2^b)$  generate the same Weyl trace algebra modulo the empty trace, after inverting the unipotent system over  $\mathbb{C}[[\hbar N]]$ ;

- (iii) the Capelli contact term of Remark 5.195,  $\text{Tr}(\mathcal{A}XY) - \text{Tr}(\mathcal{A}YX) = \hbar N \text{Tr}(\mathcal{A}) \mod I_N$ , reproduces, on  $J_N$  generators, exactly the  $r = 1$  tail of the triangular formula. Consequently the displayed shift is *absorbed into the change of basis* between  $T_{a,b}$  and  $J_N(z_1^a z_2^b)$ ; the  $J_N$ -commutator has the scalar contact term already built into the basis;



- (iv) the leading symbol of  $J_N(f)$  is  $\text{Tr } f(X, Y)$  in associated graded order, so  $\text{Tr}_N^{\text{ren}}$  stabilises in the large- $N$  limit and its stable connected projection coincides with the leading primitive trace.

*Proof.* The triangular formula of Remark 5.196 is total Weyl symmetrisation of  $z_1^a z_2^b$  inside the matrix Weyl algebra, in which one normal-ordered swap of  $z_1 z_2$  commutes through  $\hbar N$ . This is the local bivariate Weyl-normal-ordering formula proved by the contraction count in Lemma F.7; the classical Capelli source context is [26, 27]. The exact bivariate triangular trace identity is proved here by the displayed contraction count. A finite exact check verifies  $J_N \circ T = \text{id}$  on exponents  $a, b \leq 10$ . The system is unipotent over  $\mathbb{C}[[\hbar N]]$  with  $r = 0$  diagonal, hence invertible there; the sign in the inverse is  $(+\hbar N/2)^r$ , again verified by the same finite computation.

For (iii), expand

$$\text{Tr}(A XY) - \text{Tr}(A YX) = \text{Tr}(A[X, Y]) = \hbar N \text{Tr}(A) - \text{Tr}(A(YX - XY + \hbar NI))$$

using the Weyl moment relation  $YX - XY + \hbar NI = 0 \bmod I_N$  from Remark 5.195; the subtracted term lies in  $I_N$  and the displayed image is  $\hbar N \text{Tr}(A)$ . On  $J_N$  generators this is the  $r = 1$  tail term of the triangular expansion, by direct comparison with Remark 5.196. Consequently, in the  $J_N$ -basis the Capelli contact term is the change-of-basis matrix from  $T$  to  $J$ .

For (iv), the shift  $T_{a-r, b-r}$  has total degree  $a + b - 2r < a + b$  for  $r \geq 1$ , so the  $r = 0$  term is the unique top component in the  $\mathfrak{m}$ -adic filtration on  $\tilde{A}_\hbar$ ; the stable connected projection of  $f$  is  $J_N(f) \bmod I_N$  in the associated graded sense.  $\square$

COROLLARY 5.198 (*Renormalized stable connected trace map*). The map

$$\text{Tr}^{\text{ren}}: \tilde{A}_\hbar \longrightarrow H_{\text{Ham, conn}}^{\circ}(\mathcal{O}_{\text{brane}, \infty}^{\mathfrak{q}}), \quad f \longmapsto [\overline{J_N(f)}]_{N \rightarrow \infty}$$

is well defined and is a  $\mathbb{C}[[\hbar]]$ -linear injection of  $\tilde{A}_\hbar$  onto the stable Capelli-renormalized single-trace generators. The bar  $\overline{(-)}$  denotes the connected single-trace primitive projection: it removes the empty trace and disconnected products, and it is consistent with the  $J_N$ -basis at every finite  $N$  by Lemma 5.197(iii). The notation  $[\overline{J_N(f)}]_{N \rightarrow \infty}$  means this renormalized  $J$ -basis transition, or equivalently the associated-graded stable class determined by the leading primitive trace. The literal block-restriction inverse-limit element in the naive normal-ordered trace basis over plain  $\mathbb{C}[[\hbar]]$  is a different object.

THEOREM 5.199 (*Reduced Moyal commutator: radial input and obstruction criterion*). Let  $\mathcal{W}_N$  be the Weyl algebra of  $T^* \text{Mat}_N$  and let  $\hat{\mu}$  be the normal-ordered conjugation comoment above. For  $f \in \mathbb{C}[z_1, z_2][[\hbar]]$ , put

$$J_N(f) = \text{Tr } \text{Op}_W(f)(X, Y).$$

Let  $\mathfrak{t} \subset \mathfrak{gl}_N$  be the diagonal Cartan, with coordinates  $\lambda_1, \dots, \lambda_N$ , and let

$$\Delta = \prod_{i < j} (\lambda_i - \lambda_j).$$

Put

$$S_N(f) = \sum_{i=1}^N \text{Op}_W(f)(\lambda_i, -\hbar \partial_{\lambda_i}).$$

The ordinary-trace comparison is proved under the following hypotheses. Assume the radial-parts hypotheses of Statement F.8: after Vandermonde conjugation, the Harish-Chandra map on invariants

has the quantum moment-map ideal as kernel and therefore injects the reduced invariant quotient into Weyl-invariant differential operators on the regular Cartan. Under this input the radial-part map satisfies

$$\text{Rad}_0(J_N(f)) = \Delta^{-1} S_N(f) \Delta.$$

Consequently, for every  $N \geq 1$  and all  $f, g$  the defect

$$D_N(f, g) = \hbar^{-1} [J_N(f), J_N(g)] - J_N(\{f, g\}_\hbar)$$

lies in the quantum moment-map ideal  $I_N = \mathcal{W}_N \widehat{\mu}(\mathfrak{s} \mathbf{I}_N)$  and therefore vanishes in the degree-zero quantum Hamiltonian reduction

$$\mathbf{N}_{\mathcal{W}_N}(I_N)/I_N.$$

After the connected single-trace projection  $\overline{(-)}: \mathcal{W}_N/I_N \twoheadrightarrow \mathcal{W}_N/I_N/(\text{empty trace} \oplus \text{disconnected products})$ , the descended commutator on the  $J_N$ -images is the finite-rank image of the scalar Moyal bracket:

$$[\partial_{\text{bb}, \hbar, N}(f), \partial_{\text{bb}, \hbar, N}(g)]_{\text{conn, raw}} = \hbar \partial_{\text{bb}, \hbar, N}(\{f, g\}_\hbar).$$

The identification with the scalar Moyal algebra itself holds only after passing to the degreewise stable range and then to the large- $N$  connected sector. Here the large- $N$  limit is the renormalized associated-graded connected transition system of Corollary 5.198. Finite- $N$  trace identities remain in each rank, and the scalar character  $\hbar N$  has already been absorbed into the  $J_N$ -basis and then quotiented along the Hamiltonian scalar line. The identity stabilises in this large- $N$  sense, defining  $\partial_{\text{bb}, \hbar}: \mathfrak{h}_\hbar \longrightarrow H_{\text{Ham, conn}}^0(O_{\text{brane}, \infty}^q)$  satisfying

$$\hbar^{-1} [\partial_{\text{bb}, \hbar}(f), \partial_{\text{bb}, \hbar}(g)]_{\text{conn, raw}} = \partial_{\text{bb}, \hbar}(\{f, g\}_\hbar),$$

which is the bracket equation recorded in Remark 5.202.

The image-normalization clause of Statement F.8 is the quantum shear primitive: for every  $a, b, N$ ,

$$E_{a, b, N} = \sum_{j=0}^b \text{Tr}(Y^j X^a Y^{b-j}) - J_N\left(\{z_1^{a+1}/(a+1), z_2^{b+1}\}_\hbar\right) \in \mathcal{W}_N \widehat{\mu}(\mathfrak{s} \mathbf{I}_N).$$

Equivalently one must kill the free normal-diagram obstruction

$$\mathfrak{o}_{a, b} = [R_{a, b}^{\text{free}}] \in \text{coker } C_{a, b}$$

by functorial lifts  $K_{a, b} \in \ker \partial$ . In the dual-potential formulation this is the vanishing of

$$\Omega_{a, b}^{\text{rad}} = [(T_{a, b}, E_{a, b}^+)] \in \text{coker } B_{a, b},$$

equivalently the decorated PBW Stokes condition  $D_{a, b}^\square = C_{a, b}^+ \partial_2$ ; obstruction is exactly a signed row  $(\phi, -\lambda) \in \ker B_{a, b}^*$  with  $\lambda(E_{a, b}^+) - \phi(T_{a, b}) \neq 0$ . The complete edge families  $\mathfrak{o}_{2, b}$  and  $\mathfrak{o}_{a, 2}$  vanish by Proposition F.32. The finite free normal-form computations prove the displayed interior windows, including the  $K_{+, 4}$  lift and the non-edge total-degree checks recorded in Proposition F.33 and the exact normal-form script. For  $a, b \geq 3$  in all degrees, all-interior closure is equivalent to constructing the associated-graded two-colour necklace homotopy, or proving the dual-potential theorem: every signed row  $(\phi, -\lambda) \in \ker B_{a, b}^*$  satisfies  $\lambda(E_{a, b}^+) = \phi(T_{a, b})$ . The obstruction is exactly such a row with nonzero  $\lambda(E_{a, b}^+) - \phi(T_{a, b})$ .

*Proof.* Identify  $\mathcal{W}_N$  with  $\mathcal{D}_h(\mathfrak{gl}_N)$ , where  $X$  is multiplication and  $Y = -\hbar\partial_X^t$ . The quantum Hamiltonian reduction for conjugation is the normalizer quotient by the left ideal generated by the traceless comoment. The scalar matrix has zero comoment, as in Remark 5.195; equivalently the degree-zero function calculation is the  $\mathfrak{sl}_N$  reduction together with rational  $GL_N$ -invariants. Since  $GL_N$  is linearly reductive in characteristic zero, the functor of rational algebraic invariants is exact on this function-level calculation. The unreduced Lie-CE cohomology of  $\mathfrak{gl}_N$  is outside this argument.

The Harish-Chandra radial-parts theorem, whose source lineage is Harish-Chandra, Wallach, and Levasseur–Stafford [8–II], is exactly the assertion that the Vandermonde-conjugated radial map on invariants has the moment-map ideal as kernel, injects this quantum Hamiltonian reduction, and has the generator image normalization recorded in Statement F.8. The nonlinear shear obstruction in Lemma F.13 explains why the mixed Weyl-trace image is kept as part of those hypotheses; pure-power radial images supply only the special cases. Therefore the displayed radial formula for  $J_N(f)$  is valid only under the exact hypotheses named there.

Distinct one-particle Weyl summands commute, hence

$$\hbar^{-1}[S_N(f), S_N(g)] = S_N(\{f, g\}_h).$$

Conjugation by  $\Delta$  preserves commutators, so  $\text{Rad}_o(D_N(f, g)) = 0$ . Injectivity of the Harish-Chandra reduction map gives

$$D_N(f, g) \in (\mathcal{W}_N \widehat{\mu}(\mathfrak{sl}_N))^{GL_N},$$

hence  $D_N(f, g)$  is zero in the degree-zero quantum Hamiltonian reduction. The Cartan-rank-2 case has been checked by exact arithmetic: the direct  $N = 2$  radial restriction of  $J_2(f)$  is checked on 80 operator/test-polynomial pairs, and the  $N = 3$  finite-rank sweep checks 80 corresponding pairs. The same computation also checks

$$[S_2(f), S_2(g)] = S_2([f, g]_*)$$

in the normal-form convention  $z_2 = -\hbar\partial_2$  on the primary pairs, and on the higher-order pairs that exhibit nonzero  $P^3$  and  $P^5$  coefficients. These are finite-rank falsification tests; the all- $N$  source theorem remains separate.

For the connected-projection statement, the projection  $\overline{(-)}$  acts on the basis  $T_{a,b} \mapsto \overline{T_{a,b}} \equiv \overline{J_N(z_1^a z_2^b)} \bmod (\text{Capelli triangular})$ . By Lemma 5.197(iii) the Capelli contact term is absorbed into the change of basis between  $T_{a,b}$  and  $J_N$ , so on  $\overline{J_N(f)}$ -classes the descended commutator is the image of  $\hbar J_N(\{f, g\}_h) \bmod I_N$  under the projection, i.e.  $\hbar \partial_{\text{bb}, \hbar, N}(\{f, g\}_h)$ .

Stability in  $N$ : after passage to the renormalized associated-graded  $J_N$ -basis transition maps, the radial identity is compatible with increasing  $N$ . Enlarging  $\mathfrak{t}_N \subset \mathfrak{gl}_N$  to  $\mathfrak{t}_{N+k}$  adjoins  $k$  mutually commuting one-particle Weyl summands, and the bracket  $[S_N(f), S_N(g)] = S_N(\{f, g\}_h)$  is preserved at every  $N$ . By Lemma 5.197(iv), the connected extraction depends only on the leading symbol filtration, which the Capelli triangular system preserves. Hence the stable connected limit exists and inherits the bracket identity.

The final obstruction statement is Lemma F.15 together with Definition F.21 and Statement F.22. The edge vanishing is Proposition F.32. The finite interior identities are the free normal-diagram identities of Proposition F.20, Proposition F.33, and the reproducible exact-arithmetic computation `scripts/quantum_shear_trace_diagram_normal_form.py`. These data give finite-window identities and edge theorems; the functorial all-interior splitting remains the missing construction.  $\square$

COROLLARY 5.200 (*Degree-zero quantum comparison and decorated radial obstruction*). Under the same radial-parts hypotheses as Theorem 5.199, the map

$$\partial_{\text{bb},\hbar} : \mathfrak{h}_{\hbar} = \tilde{A}_{\hbar} \longrightarrow H_{\text{Ham,conn}}^{\circ}(\mathcal{O}_{\text{brane},\infty}^{\mathfrak{q}}), \quad f \longmapsto [\overline{J_N(f)}]_{N \rightarrow \infty},$$

is a Lie-algebra map for  $\{, \}_{\hbar}$  on the source and  $\hbar^{-1}[-, -]_{\text{conn,raw}}$  on the target. Its associated graded at  $\hbar = 0$  is the degree-zero part of Theorem 5.6. All-order quantum corrections are governed by the odd-power Moyal expansion

$$\{f, g\}_{\hbar} = \sum_{s \geq 0} \frac{\hbar^{2s}}{2^{2s}(2s+1)!} P^{2s+1}(f, g),$$

so its second-order normalized coefficient is  $\frac{1}{24} P^3(f, g)$ . This term can be nonzero.

Under the stated finite-window and radial hypotheses, the formal Moyal coefficients and the Capelli triangular trace basis remain proved. The quantum shear primitive needed for the ordinary Weyl-trace comparison is proved on the edge strips by Proposition F.32 and in the finite verified interior windows. Its all-interior extension is exactly the dual-potential obstruction theorem

$$\Omega_{a,b}^{\text{rad}} = [(T_{a,b}, E_{a,b}^+)] \in \text{coker } B_{a,b}$$

of Proposition F.25: it vanishes if and only if the decorated PBW Stokes identity  $D_{a,b}^{\square} = C_{a,b}^+ \partial_2$  holds, with obstruction represented exactly by a signed row  $(\phi, -\lambda) \in \ker B_{a,b}^*$  with  $\lambda(E_{a,b}^+) - \phi(T_{a,b}) \neq 0$ .

*Proof.* This is Theorem 5.199, combined with the Moyal odd-symmetry of Proposition 5.215:  $P^r$  changes by  $(-1)^r$  under interchange of  $f$  and  $g$ , so only odd  $r$  survive in the raw star commutator. Dividing by  $\hbar$  shifts these odd raw powers to even powers in the normalized Hamiltonian bracket. Hence the normalized  $\hbar^2$  coefficient is  $P^3(f, g)/24$ , and the normalized  $\hbar^4$  coefficient is  $P^5(f, g)/1920$ .  $\square$

*Remark 5.201 (Explicit second- and fourth-order corrections).* For higher-order instances that exercise nontrivial  $P^3$  and  $P^5$ :

$$\begin{aligned} [z_1^3 z_2, z_1 z_2^3]_* &= 8\hbar z_1^3 z_2^3 - 3\hbar^3 z_1 z_2 + O(\hbar^5), \\ [z_1^4 z_2^2, z_1^2 z_2^4]_* &= 12\hbar z_1^5 z_2^5 - 40\hbar^3 z_1^3 z_2^3 + 6\hbar^5 z_1 z_2 + O(\hbar^7). \end{aligned}$$

After normalization by  $\hbar^{-1}$ , the first example gives

$$\{z_1^3 z_2, z_1 z_2^3\}_{\hbar} = 8z_1^3 z_2^3 - 3\hbar^2 z_1 z_2 + O(\hbar^4).$$

The  $\hbar^3$  raw coefficient is therefore the normalized  $\hbar^2$  correction  $\frac{1}{24} P^3(f, g)$ , in agreement with Proposition 5.215. The non-affine bracket correction reduces, in this constant Poisson tensor case, to exactly the higher Moyal coefficients.

*Remark 5.202 (Degree-zero quantum Hamiltonian sector).* Assume the radial image datum of Statement F.8. Then Corollary 5.200 identifies the stable connected single-trace quotient of  $\mathcal{O}_{\text{brane},N}^{\mathfrak{q}}$  and proves that  $\partial_{\text{bb},\hbar,N}$  stabilizes to a map

$$\partial_{\text{bb},\hbar} : \mathfrak{h}_{\hbar} \longrightarrow H_{\text{Ham,conn}}^{\circ}(\mathcal{O}_{\text{brane},\infty}^{\mathfrak{q}})$$

satisfying

$$\hbar^{-1}[\partial_{\text{bb},\hbar}(f), \partial_{\text{bb},\hbar}(g)]_{\text{conn,raw}} = \partial_{\text{bb},\hbar}(\{f, g\}_{\hbar}).$$

Thus the connected quantum brane algebra is the Weyl/Moyal deformation of the degree-zero Hamiltonian sector. Its associated graded at  $\hbar = 0$  is the degree-zero part of Theorem 5.6. The  $J_N$ -basis change absorbs the finite- $N$  Capelli contact and its triangular  $\hbar N$  lower-trace terms; the connected projection removes only the empty trace and decomposable products. Interior bidegrees reduce to the exact Weyl-trace image clause of Statement F.8, equivalently to quantum shear primitives  $E_{a,b,N} \in \mathcal{W}_N \widehat{\mu}(\mathfrak{sl}_N)$ . In the free indexed normal-diagram model this is the vanishing of  $\mathfrak{o}_{a,b}$  together with functorial lifts  $K_{a,b}$ . In the sharper dual-potential form the obstruction is  $\Omega_{a,b}^{\text{rad}}$ , equivalently decorated PBW Stokes for  $D_{a,b}^\square = C_{a,b}^+ \partial_2$ ; an obstruction is a signed  $B_{a,b}^*$ -row with nonzero  $\lambda(E_{a,b}^+) - \phi(T_{a,b})$ . The edge strips are proved; the interior all-bidegree theorem is the associated-graded two-colour necklace homotopy, or this equivalent dual-potential signed-row vanishing theorem.

*Remark 5.203.* The degree-zero Hamiltonian target is the Weyl/Moyal algebra of the formal symplectic polydisk, the constant-Poisson case of deformation quantization [7]. Costello's BV/RG construction [12] and the flat open-closed BCOV/HCS quantization of Costello–Li [13, 14] apply under their elliptic, gauge-fixing, renormalization, and obstruction-vanishing hypotheses; the mixed ordinary- $\mathfrak{gl}_N$  Hamiltonian BF defect problem here lies outside their hypothesis class and requires the specialization datum of Problem 5.212.

*Remark 5.204.* Corollary 5.172 gives the Hamiltonian action on the polynomial representatives  $\Psi_{k,l}$ : it is the PBW special-fibre action. The cotangent field carries the coadjoint action. A quantum descendant theorem requires a boundary model for the continuous dual  $\mathfrak{h}_\hbar^{\vee, \text{cont}}$ , for example a Laurent or distributional boundary realization.

*Remark 5.205.* The full cotangent deformation would replace  $\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}^\vee[1]$  by

$$\mathfrak{g}_\hbar = \mathfrak{h}_\hbar \ltimes \mathfrak{h}_\hbar^{\vee, \text{cont}}[1]$$

and would ask for  $C_{\text{CE,cont}}^\bullet(\mathfrak{g}_\hbar)$  as the closed quantum observable algebra. The polynomial boundary representatives carry the PBW special-fibre action; the deformation-level coadjoint action requires the principal-part target.

### 5.7.3 GRAPH FORMALISM AND SPECIALIZATION DATUM

The Moyal coefficients  $\hbar P$ ,  $\hbar^3 P^3/24$ ,  $\hbar^5 P^5/1920$ ,  $\hbar^7 P^7/322560$  of the previous subsection are an algebraic target. Costello's heat-kernel BV calculus produces coefficients of the same operad-theoretic shape from a different input: an elliptic parent theory, a propagator, a counterterm scheme, an anomaly class, and a trace normalisation. The question is whether these two calculations land on the same coefficients on the brane-defect target.

*Statement 5.206 (Costello perturbative BV construction).* For an elliptic BV theory with a gauge fixing, heat kernel  $K_t$ , propagator

$$P_{\epsilon,L} = \int_\epsilon^L Q^{\text{GF}} K_t dt,$$

and a renormalization scheme, Costello constructs finite-scale effective interactions by the RG operator

$$W(P_{\epsilon,L}, I[\epsilon]) = \hbar \log(\exp(\hbar \partial_{P_{\epsilon,L}}) \exp(I[\epsilon]/\hbar)).$$

The renormalized interaction is obtained by adding local counterterms and taking the  $\epsilon \rightarrow 0$  limit. It has the connected-graph expansion

$$I[L] = \sum_{\Gamma \text{ connected}} \frac{\hbar^{b_1(\Gamma) + \sum_v g_v}}{|\text{Aut } \Gamma|} W_{\Gamma}(P_{\epsilon, L}; I_{g_v}[\epsilon] + I_{g_v}^{\text{CT}}(\epsilon)),$$

satisfies RG flow and locality, and satisfies the quantum master equation when the obstruction classes vanish [12]. The resulting quantum observables form a factorization algebra [15, 16].

*Statement 5.207 (Costello–Li flat open-closed BCOV theorem).* Costello–Li prove a perturbative quantization of open-closed BCOV theory on flat  $\mathbb{C}^d$ , for  $d$  odd, unique up to the equivalences in their renormalization scheme, with open gauge algebra  $\mathfrak{gl}(N | N)$  and compatibility under  $\mathfrak{gl}(N | N) \hookrightarrow \mathfrak{gl}(N + k | N + k)$ . In that flat open-closed BCOV/HCS theory they compute the anomaly cancellation needed for the graph construction [14]. Ordinary  $\mathfrak{gl}_N$  and the mixed  $\mathbb{R}^2 \times \mathbb{C}^2$  Hamiltonian brane model require an additional reduction or summand argument.

*Remark 5.208.* Theorem 5.6 uses the formal Hamiltonian BF calculation as its input. The brane  $L = \mathbb{R}_{\text{brane}} \times \{0\} \times \{p\}$  is a defect in the mixed space; a spacetime-boundary interpretation requires a chosen half-space model.

*LEMMA 5.209 (Finite-window obstruction for the full Hamiltonian sector).* The polynomial Hamiltonian Lie algebra of the formal symplectic plane has the following finite-window obstructions to a naive finite-dimensional Lie truncation compatible with the full Hamiltonian sector:

- (i) finite degree windows have a Lie-subalgebra obstruction;
- (ii) projected degree-window brackets can violate Jacobi;
- (iii)  $\mathfrak{m}$ -adic jet quotients have a Lie-quotient obstruction in the presence of linear Hamiltonians;
- (iv) directed finite-dimensional Lie-subalgebra exhaustion is obstructed containing the full polynomial Hamiltonian algebra.

*Proof.* Let  $H_d$  be the span of homogeneous polynomials of degree  $d$ , and let  $\mathfrak{m} = (z_1, z_2)$ . The Poisson bracket satisfies  $\{H_p, H_q\} \subset H_{p+q-2}$ . Thus

$$\{z_1^N, z_2^N\} = N^2 z_1^{N-1} z_2^{N-1},$$

which has degree  $2N - 2 > N$  for  $N > 2$ , so degree  $\leq N$  windows have a Lie-subalgebra obstruction.

Projecting the bracket back to a degree window creates a Jacobi obstruction. For the bracket  $\pi_{\leq 3}\{-, -\}$ , take

$$f = z_2, \quad g = z_2^3, \quad h = z_1^2 z_2.$$

A direct calculation gives the projected Jacobiator

$$[[f, g]_3, h]_3 + [[g, h]_3, f]_3 + [[h, f]_3, g]_3 = 6z_2^3.$$

The  $\mathfrak{m}$ -adic quotients have a stability obstruction under translations generated by linear Hamiltonians:

$$\{z_1^{N+1}, z_2\} = (N + 1)z_1^N \notin \mathfrak{m}^{N+1}.$$

Hence  $\mathfrak{h}/\mathfrak{m}^{N+1}$  has a Lie-quotient obstruction for the full Hamiltonian algebra.

Any finite-dimensional Lie subalgebra containing

$$a = z_1^2 z_2^2, \quad b = z_1^3 z_2$$

contains all iterated brackets

$$(\mathrm{ad}_a)^n(b) = (-4)^n z_1^{n+3} z_2^{n+1}, \quad n \geq 0,$$

an infinite linearly independent set. Thus directed finite-dimensional Lie-subalgebra exhaustion is obstructed for the full polynomial Hamiltonian algebra.  $\square$

*Remark 5.210 (Linear heat operator versus BV kernel).* On the linear field complex

$$\Omega^\bullet(\mathbb{R}^2) \widehat{\otimes} \Omega^{\bullet, \bullet}(\mathbb{C}^2) \widehat{\otimes} (\widehat{\mathfrak{h}}[1] \oplus \widehat{\mathfrak{h}}_{\mathrm{cont}}^\vee[2]),$$

the flat gauge fixing

$$Q^{\mathrm{GF}} = d_{\mathbb{R}^2}^* + \bar{\partial}_{\mathbb{C}^2}^*, \quad \Delta = [d_{\mathbb{R}^2} + \bar{\partial}_{\mathbb{C}^2}, Q^{\mathrm{GF}}]$$

gives finite-scale heat operators

$$K_t = K_t^{\mathrm{base}} \widehat{\otimes} \mathrm{id}, \quad P_{\epsilon, L}^{\mathrm{op}} = \int_{\epsilon}^L Q^{\mathrm{GF}} K_t^{\mathrm{base}} dt \widehat{\otimes} \mathrm{id}$$

satisfying

$$[d_{\mathbb{R}^2} + \bar{\partial}_{\mathbb{C}^2}, P_{\epsilon, L}^{\mathrm{op}}] = K_{\epsilon} - K_L$$

as continuous operators on the completed coefficient spaces. This is an operator-level homotopy. Costello's BV propagator also requires a coefficient coevaluation kernel for the Tate pair  $\widehat{\mathfrak{h}} \oplus \widehat{\mathfrak{h}}_{\mathrm{cont}}^\vee$ . In the product/direct-sum topology used here, the formal sum of homogeneous coefficient coevaluations diverges in the completed tensor product. Thus the analytic graph problem requires a genuine Tate or weighted coefficient extension of the heat-kernel calculus, or a declared restriction to a smaller sector.

*LEMMA 5.211 (Tate Casimir obstruction for the full Hamiltonian sector).* In the product/direct-sum Tate topology

$$\mathfrak{h} = \prod_{d \geq 1} H_d, \quad \mathfrak{h}_{\mathrm{cont}}^\vee = \bigoplus_{d \geq 1} H_d^\vee,$$

the coefficient coevaluation

$$C_{\mathfrak{h}} = \sum_{d \geq 1} \sum_i e_{d,i} \otimes e_d^i$$

lies outside  $\mathfrak{h} \widehat{\otimes} \mathfrak{h}_{\mathrm{cont}}^\vee$ , nor of  $\mathfrak{h}_{\mathrm{cont}}^\vee \widehat{\otimes} \mathfrak{h}$ . Consequently the full Hamiltonian sector has a BV pairing; a Costello coefficient kernel in this topology requires a different coefficient category. The operator-level homotopy of Remark 5.210 supplies a BV propagator, BV Laplacian, and renormalized graph weights only after a coefficient category in which the Casimir converges has been chosen.

*Proof.* For each  $d$ , choose a basis  $\{e_{d,i}\}_i$  of  $H_d$  and the dual basis  $\{e_d^i\}_i$  of  $H_d^\vee$ . An element of a completed tensor product with the direct-sum factor  $\bigoplus_d H_d^\vee$  has finite support in that direct-sum direction in the Tate convention used here. The displayed formal Casimir has a nonzero component in every  $H_d^\vee$ . Hence the tensor diverges. The same support obstruction holds with the tensor factors reversed. Costello's heat-kernel BV formalism requires the inverse pairing as a distributional kernel on the field complex; an abstract continuous operator is insufficient. The weighted Tate completion of Theorem 5.70 resolves this Casimir obstruction by changing the coefficient category; the unweighted diagnosis here remains the structural reason that change is required.  $\square$

*Problem 5.212 (Hamiltonian specialization datum for Costello perturbative BV).* • *Desired upgrade:* a Costello-type closed-open BV theory whose restriction is the formal-local Hamiltonian BF sector  $\mathbb{R}^2 \times \mathbb{C}^2$  of §1.4.

- *Minimal input data:* an elliptic parent or stated dimensional reduction; field topology, BV pairing, gauge fixing, heat-kernel propagator; weighted Tate coefficient kernel (Theorem 5.70); brane-defect condition; ribbon graph data; Capelli/scalar normalisation; bulk-to-defect kernel.
- *Obstruction complex:* the Tate Casimir  $\mathfrak{o}_{\text{Cas}}$  (Lemma 5.211); the scalar-symbol class  $\hbar N[\bar{c}]$  (Theorem 5.213); the reduced non-scalar QME class  $[\text{Ob}_{w,d,\hbar}^{\text{red}}]$  (Problem 5.219); the  $\theta_3$  finite-window class (Theorem E.38).
- *Vanishing criterion:* weighted Tate completion closes  $\mathfrak{o}_{\text{Cas}}$ ; a balanced or unreduced source replacement closes  $\hbar N[\bar{c}]$ ; the reduced non-scalar class is closed exactly on the locus where the filtered scalar-contact tower of Theorem 5.83 closes.
- *First nonzero example:* the strict full-polydisk product/direct-sum coefficient pair carries a nonzero  $\mathfrak{o}_{\text{Cas}}$ ; the ordinary scalar-reduced  $\mathfrak{gl}_N$  branch carries nonzero  $\hbar N[\bar{c}]$  at order  $\hbar^1$  on the linear pair  $(z_1, z_2)$ .
- *Physical meaning:* the analytic specialization datum is the elliptic parent and the brane-defect renormalisation under which Costello's heat-kernel BV calculus and the Hamiltonian trace map is simultaneously controllable.

*Statement.* Costello supplies the general renormalized BV graph calculus. For the Hamiltonian sector here, the required theorem data begin with an elliptic closed-open parent theory, or a stated dimensional reduction of one, whose restriction is this Hamiltonian BF sector and whose obstruction class vanishes. The required data are:

- (i) choose the mechanism: either prove that the Hamiltonian cotangent BF complex

$$\Omega^\bullet(\mathbb{R}^2) \widehat{\otimes} \Omega^{\circ,\bullet}(\mathbb{C}^2) \widehat{\otimes} (\widehat{\mathfrak{h}}[1] \oplus \widehat{\mathfrak{h}}_{\text{cont}}^\vee[2])$$

is literally a closed sector of a Costello-type open-closed BV theory, or prove that it is obtained by a specified dimensional reduction. The chosen mechanism must determine the field topology, degree  $-1$  BV pairing, differential, local interaction, bracket degree, and a homotopy retract closed under the propagator and RG flow. Since  $\widehat{\mathfrak{h}}$  is a completed Tate coefficient system, the proof goes through a Tate or weighted-coefficient extension; finite-degree and  $\mathfrak{m}$ -adic truncations carry the obstruction of Lemma 5.209. It must construct a genuine Tate or weighted-coefficient extension of Costello's heat-kernel calculus [12] compatible with the Poisson bracket and RG flow, or restrict to a smaller Lie sector such as the classical  $\mathfrak{m}^3/\mathfrak{m}^{N+1}$  truncation and explicitly declare that translations, the conformal  $\mathfrak{sl}_2$  sector, and the all-order Moyal closure have been removed. By Lemma 5.211, a damping factor inside the direct-sum dual fixed above is insufficient; the coefficient category itself must be replaced by one in which the Casimir converges, or the sector must be restricted;

- (ii) specify the asymptotic conditions on  $\mathbb{C}^2$  and the brane-defect condition on  $L = \mathbb{R}_{\text{brane}} \times \{0\} \times \{p\}$ ;
- (iii) choose the induced finite-scale gauge fixing, heat-kernel propagator, and BV Laplacian in this sector. The infrared limit  $L \rightarrow \infty$  may be used only after the zero modes and harmonic projection have been fixed;
- (iv) compute the obstruction/anomaly class for this mixed Hamiltonian sector. Concretely, represent the obstruction as a local functional class in the chosen bulk-defect local-functional complex. If



the defect cotangent sector is represented by principal-part currents, the target must either be replaced by a current-compatible complex

$$\mathcal{O}_{\text{loc}}^{\text{bd}}(\mathcal{E}_w; \mathfrak{F}_{\partial}^{\text{pp}}),$$

with wavefront and counterterm rules for  $\mathcal{D}'_c$ -coefficients, or the theorem must restrict  $B$  to regular compactly supported densities. In the weighted Tate category this means a class

$$\text{Ob}_{w,n} \in H_{\text{adm}}^1(\mathcal{O}_{\text{loc}}(\mathcal{E}_w), Q + \{I_o^w, -\})$$

coming from the  $n$ -loop obstruction to the quantum master equation after counterterms. Costello–Li’s flat  $\mathfrak{gl}(N \mid N)$  anomaly cancellation supplies a source theorem and a model; transfer requires proof. To execute it one must prove that constant-Hamiltonian projection and connected trace extraction kill the ordinary  $\mathfrak{gl}_N$  anomaly, or else replace the open sector by the appropriate super or special-linear variant. The first defect-local obstruction is the Capelli contact term

$$\omega_{\text{Cap}}(A) = N \overline{\text{Tr}(A)},$$

represented by

$$\text{Tr}(A X Y) - \text{Tr}(A Y X) = \hbar N \text{Tr}(A) \mod \mathcal{W}_N \widehat{\mu}(\mathfrak{sl}_N).$$

It survives the naive connected trace projection, since the right-hand side is a nonempty connected trace when  $A$  is nonconstant. The anomaly datum must therefore be a filtered local class  $\text{Ob}_{\text{bd}}$  whose scalar-symbol projection is

$$\sigma_{\text{sc}}(\text{Ob}_{\text{bd}}) = \hbar N [\bar{c}] \in H_{\text{Lie}}^2(\tilde{A}; \mathbb{C})[[\hbar]].$$

A closed-exchange repair would have to supply  $\mathcal{W}_{\text{closed exch}}$  and a counterterm  $C$  with scalar leading class

$$[w_{\text{ext}}] = -\hbar N [\bar{c}] \quad \text{in } H_{\text{Lie}}^2(\tilde{A}; \mathbb{C})[[\hbar]]$$

and with

$$\omega_{\text{Cap}} + \mathcal{W}_{\text{closed exch}} + (Q + \{I_o, -\})C = 0.$$

The proved local mechanisms here are the specified supertrace and central-character normalizations; any exhaustive classification is controlled by the displayed obstruction complex;

- (v) construct the bulk-to-defect kernel as a curved/cochain-level map  $\kappa$  to the Poisson central-operation object of the brane observables. After counterterms it must satisfy

$$\text{Curv}(\kappa) = d\kappa + \frac{1}{2}[\kappa, \kappa] = 0.$$

On degree-zero Hamiltonians it must send  $f(z_1, z_2)$  to the Weyl-ordered boundary observable  $\text{Tr Op}_{\mathbb{W}}(f)(\phi_1, \phi_2)$ , and it must satisfy the Ward identity and the quantum moment-map constraint. Its cotangent component must either restrict the principal-part current  $B$  to regular defect densities, or prove that the chosen Costello bulk-defect observable category admits the required compactly supported distributions with wavefront, counterterm, finite-window, and RG compatibility;

- (vi) define the relevant boundary ribbon graphs: cyclic order, boundary components, automorphism factors, propagator labels, and the relation between Costello's loop-counting  $\hbar$  and the Weyl contraction parameter. Equivalently, one must prove that a single boundary cross-contraction is  $P/2$  in the Weyl convention;
- (vii) fix the one-propagator boundary graph normalization. For monomials  $f = z_1^k z_2^l$  and  $g = z_1^m z_2^n$ , the raw first boundary commutator must have coefficient

$$\hbar(kn - lm) z_1^{k+m-1} z_2^{l+n-1},$$

equivalently the normalized commutator must have coefficient

$$(kn - lm) z_1^{k+m-1} z_2^{l+n-1};$$

- (viii) after the first normalization, compute the three-propagator boundary graph normalization and prove that the connected order- $\hbar^3$  commutator contributes exactly

$$\frac{\hbar^3}{24} P^3(f, g)$$

with  $P = \partial_{z_1} \otimes \partial_{z_2} - \partial_{z_2} \otimes \partial_{z_1}$ ;

- (ix) give the convention for extracting connected single-trace and multi-trace coefficients as primitive trace classes, and prove that this extraction cancels, renormalizes, or otherwise accounts for the normal-ordering shift of Remark 5.195;
- (x) construct the renormalization scheme and counterterm locality statement in this mixed bulk-defect sector. For every divergent graph whose  $\epsilon \rightarrow 0$  limit is subtracted, choose compatible representatives

$$C_{\Gamma, w}^{w_o}(\epsilon) \in \mathcal{O}_{\text{loc}}(\mathcal{E}_w^{w_o})$$

with

$$\pi_{w_o w_o} C_{\Gamma, w}^{w_o'} = C_{\Gamma, w}^{w_o}, \quad R_{w, w'}^{w_o} C_{\Gamma, w}^{w_o} = C_{\Gamma, w'}^{w_o},$$

prove support on the relevant bulk diagonal or brane-collision stratum, and check that RG flow carries these representatives inside the same local-functional subcomplex;

- (xi) extend the bulk-to-defect kernel to the cotangent boundary-current target, or record the obstruction. In the reduced setting the extension must land in the principal-part current object of Theorem 5.185; before contraction it must supply the unreduced data of Remark 5.189. On the cotangent summand it must realize the principal-part coadjoint action of Theorem 5.178, and at the Moyal level the  $\mathfrak{h}_\hbar^\vee$ -action induced by the residue pairing. The allowed substitute for polynomial one- $\psi$  representatives is a completed continuous-dual boundary realization.

**THEOREM 5.213** (*Scalar QME obstruction and cancellation branches*). • *Desired upgrade:* cancellation of the scalar-symbol QME obstruction in the ordinary scalar-reduced  $\mathfrak{gl}_N$  brane-defect model.

- *Minimal input data:* on the source side, an  $\bar{A}$ -cochain  $J$  with  $d_{\text{CE}} J = -\hbar N \bar{c}$ , or an alternative source category in which  $\bar{c}$  is cohomologous to zero; on the Costello side, finite-window complexes  $\mathcal{X}_{\text{sc}, w, M}^\bullet$ , cochain maps  $\Xi_M^{\text{Cost}, \text{sc}}$ , a filtered scalar chain projection  $\widehat{\sigma}_{\text{sc}, M}$ , and compatible local counterterms.
- *Obstruction complex:* the Lie cohomology class  $\hbar N[\bar{c}] \in H_{\text{Lie}}^2(\bar{A}; \mathbb{C})[[\hbar]]$ , and its filtered Costello image  $\beta_M^{\text{Cost}, \text{sc}} = [\hbar N[\bar{c}_M]] \in \text{coker } H^1(\widehat{\sigma}_{\text{sc}, M} \Xi_M^{\text{Cost}, \text{sc}})$ .

- *Vanishing criterion:* either replace the scalar-reduced source by the unreduced source, the central-character source, the opposite central extension, the balanced  $\mathfrak{gl}(N|N)$ -supertrace source, or the algebraic central-contact cone — each cancellation is a source replacement, made explicit below; alternatively, supply a Costello-local closed-exchange triple  $(W_M, C_M, \widehat{\sigma}_{\text{sc},M})$  with  $\beta_M^{\text{Cost,sc}} = 0$  and compatible truncation.
- *First nonzero example:* on the linear pair  $(z_1, z_2)$ ,  $\omega(z_1, z_2) = 1$  and  $\bar{c} \neq 0$  in  $H_{\text{Lie}}^2(\bar{A}; \mathbb{C})$ ; the Moyal commutator gives the Capelli term  $[\Phi_{\hbar}(z_1), \Phi_{\hbar}(z_2)]_{\star} = \hbar N$  at order  $\hbar^1$ .
- *Physical meaning:* the scalar-symbol QME obstruction records the brane center-of-mass anomaly that survives ordinary  $\mathfrak{gl}_N$  Hamiltonian reduction; cancelling it requires changing what one means by “Hamiltonian” (source replacement) or supplying a closed-exchange counterterm with a matched scalar leading class.

*Statement.* Let  $A = \mathbb{C}[z_1, z_2]$ ,  $\bar{A} = A/\mathbb{C} \cdot 1$ , and

$$\omega(f, g) = [\{f, g\}]_0.$$

Write  $\bar{c}$  for the induced Lie two-cocycle on  $\bar{A}$ . In the ordinary scalar-reduced  $\mathfrak{gl}_N$  brane-defect model, the scalar-symbol part of the mixed QME obstruction is represented by

$$\text{Ob}_{\text{sc}}(\gamma_1; a, f; b, g) = \hbar N \omega(f, g) \int_{\mathbb{R}} a(t)b(t)\gamma_1(t) dt,$$

and its associated-graded CE class is

$$\hbar N [\bar{c}] \in H_{\text{Lie}}^2(\bar{A}; \mathbb{C})[[\hbar]].$$

For  $N > 0$ , this class is nonzero in the scalar-reduced source. Hence the leading scalar symbol obstructs internal scalar-reduced local counterterms.

The cancellation mechanisms have distinct types.

- (i) On the unreduced source  $A$ , the functional  $\eta(f) = -[f]_0$  satisfies  $d_{\text{CE}}\eta = \omega$ . Thus

$$\hbar N \omega + d_{\text{CE}}(-\hbar N \eta) = 0$$

before quotienting constants.

- (ii) Equivalently, after replacing the scalar-reduced source by the unreduced source with central character  $\chi_N(f) = N[f]_0$ , the scalar term is absorbed before it becomes a reduced obstruction class.
- (iii) Equivalently in the scalar-replacement category, define the opposite scalar central extension

$$\bar{A}_{-\hbar N \bar{c}}^{\text{cl}} = \bar{A} \oplus \mathbb{C}[[\hbar]]K_{\text{cl}},$$

with

$$[(f, aK_{\text{cl}}), (g, bK_{\text{cl}})] = (\{f, g\}_{\bar{A}}, -\hbar N \bar{c}(f, g)K_{\text{cl}}), \quad [K_{\text{cl}}, -] = 0.$$

If  $k^{\vee}(K_{\text{cl}}) = 1$  and  $k^{\vee}(\bar{A}) = 0$ , then for  $J_{\text{op}} = -k^{\vee}$

$$d_{\text{CE}}J_{\text{op}} = -\hbar N \bar{c}, \quad \text{Ob}_{\text{sc}} + d_{\text{CE}}J_{\text{op}} = 0.$$

The cancellation is compatible with scalar finite-window projections retaining  $z_1, z_2$ . It is a source replacement; after forgetting  $K_{\text{cl}}$ , the class  $\hbar N [\bar{c}]$  on  $\bar{A}$  reappears.

- (iv) In the balanced supertrace model  $V = \mathbb{C}^{N|N}$ , the scalar coefficient is  $\text{sdim } V = 0$ ; the scalar brane-defect QME representative vanishes in that model.
- (v) The finite-window algebraic central-contact cone is a genuine changed source branch. After adjoining a defect-supported central closed source line  $\mathbb{C}[[\hbar]]e_M$  with

$$\Xi_M^{\text{alg,sc}}(e_M) = -\hbar N \bar{c}_M$$

on the scalar-contact line, the choice  $W_M = e_M$  and  $C_M = 0$  cancels the finite-window scalar row, and the corresponding  $\beta, \mu, \lambda_C$  classes vanish in that enlarged algebraic cone.

- (vi) In the original scalar-reduced ordinary  $\mathfrak{gl}_N$  branch, a same-branch closed-exchange or brane-defect contribution can cancel the scalar symbol only if its leading scalar class obeys

$$[w_{\text{ext}}] = -\hbar N [\bar{c}] \quad \text{in } H_{\text{Lie}}^2(\bar{A}; \mathbb{C})[[\hbar]].$$

Equivalently, a Costello-local closed-exchange solution must supply finite-window closed-source complexes  $\mathcal{X}_{\text{sc},w,M}^\bullet(I)$ , cochain maps

$$\Xi_M^{\text{Cost,sc}}: \mathcal{X}_{\text{sc},w,M}^\bullet(I) \rightarrow \mathcal{Q}_{\text{sc},w,M}^\bullet(I),$$

a filtered scalar chain projection  $\widehat{\sigma}_{\text{sc},M}$ , cocycles  $W_M$ , and counterterms  $C_M$  such that

$$\text{Ob}_{\text{sc},M} + \Xi_M^{\text{Cost,sc}}(W_M) + d_M C_M = 0, \quad H^1(\widehat{\sigma}_{\text{sc},M} \Xi_M^{\text{Cost,sc}})[W_M] = -\hbar N [\bar{c}_M],$$

with compatible truncation and vanishing inverse-limit  $\beta, \mu, \lambda_C$  classes.

The algebraic cone in (4) becomes a Costello-local closed-exchange theorem only after its formal generator is replaced by a closed source complex, finite-scale propagator, bulk-to-defect graph map, source-face Hom differential, local counterterm support, scalar chain projection, and compatible finite-window representatives. For current genuine Costello-local towers the ordinary scalar-reduced branch is blocked first by

$$o_{\sigma,M}^{(1),\text{sc}} = \hbar N [\bar{c}_M] \neq 0,$$

and, after any externally supplied scalar condition, by the image obstruction

$$\beta_M^{\text{Cost,sc}} = [\hbar N [\bar{c}_M]] \in \text{coker } H^1(\widehat{\sigma}_{\text{sc},M} \Xi_M^{\text{Cost,sc}}).$$

A complete classification of scalar QME cancellations requires a separate target theorem. Any further mechanism must be an explicitly constructed null-homotopy in the filtered QME obstruction complex whose scalar-symbol class cancels  $\hbar N [\bar{c}]$ .

*Proof.* The representative and its nonzero associated-graded class are exactly Theorem E.12. Non-exactness in the reduced source is the last part of that theorem: a primitive on  $\bar{A}$  would lift to a primitive on  $A$  vanishing on  $\mathfrak{I}$ , but  $\omega(z_1, z_2) = 1$  and  $\{z_1, z_2\} = \mathfrak{I}$ .

The first branch is Lemma E.11, multiplied by  $\hbar N$ . The central-character branch is Theorem E.15; it is a source replacement before scalar reduction. A counterterm inside the original reduced obstruction complex is a different datum. The opposite central-extension branch is the same CE sign calculation written as an honest central extension of  $\bar{A}$ : with  $(d_{\text{CE}}\lambda)(x, y) = -\lambda([x, y])$ , the central bracket  $-\hbar N \bar{c}(f, g)K_{\text{cl}}$  gives  $d_{\text{CE}}k^\vee = \hbar N \bar{c}$ , hence  $d_{\text{CE}}(-k^\vee) = -\hbar N \bar{c}$ . Jacobi is exactly the cocycle identity

for  $\bar{c}$ , and finite-window compatibility follows because  $\bar{c}$  is detected on the retained pair  $z_1, z_2$ . The balanced-supertrace branch is Theorem E.16. The algebraic central-contact cone is the finite-window source extension recorded in the scalar QME source extension; its differential is zero and its scalar image is declared to be  $-\hbar N \bar{c}_M$ , so the displayed cancellation is immediate in that enlarged source. The same declaration records the missing Costello-local closed-exchange data. The necessary condition for any same-branch ordinary cancellation is Theorem E.52. The additional Costello-local image and scalar-chain obstructions are the filtered Hom conditions in Problem 5.219.  $\square$

*Remark 5.214 (Parabolic functoriality of the local model).* The completion along the brane divisor  $z_1 = 0$  used here breaks the formal symplectomorphism group  $\mathrm{Symp}_{\mathrm{form}}(\mathbb{C}^2, 0)$  to the parabolic stabiliser

$$P_{(z_1)} = \{\varphi \in \mathrm{Symp}_{\mathrm{form}}(\mathbb{C}^2, 0) : \varphi^*(z_1) \in (z_1)\}.$$

Any cross-target transfer therefore supplies a parabolic-equivariant comparison. Full  $\mathrm{Symp}_{\mathrm{form}}$ -equivariance is additional target data.

#### 5.7.4 FORMAL WEYL/MOYAL TARGET, ALL ORDERS

The order- $\hbar$  coefficient  $[\Phi_{\hbar}(z_1), \Phi_{\hbar}(z_2)]_{\star} = \hbar N \cdot I_N$  in  $M_N(\mathbb{C})$  identifies the linear-generator anomaly on the matrix-brane realization; the all-order question is whether the same identification persists at orders  $\hbar^3, \hbar^5, \dots$ . The unique translation-invariant deformation of the constant Poisson tensor on the formal holomorphic polydisk [7] fixes the all-order target on the closed side.

The Weyl/Moyal Rees convention is

$$f * g = m \circ \exp\left(\frac{\hbar}{2}(\partial_{z_1} \otimes \partial_{z_2} - \partial_{z_2} \otimes \partial_{z_1})\right)(f \otimes g).$$

The factor  $1/2$  is already determined by linear Hamiltonians in this Weyl convention: if  $u = az_1 + bz_2$  and  $v = cz_1 + dz_2$ , then  $[\widehat{u}, \widehat{v}] = \hbar(ad - bc)$ , so the Baker-Campbell-Hausdorff formula for Weyl-ordered exponentials contributes  $\exp(\hbar P(u, v)/2)$ . Differentiating these exponentials gives the cross-contraction  $P/2$  in the formal Weyl convention. Then the star commutator is odd in  $\hbar$ :

$$[f, g]_{\star} = \hbar\{f, g\} + \frac{\hbar^3}{24}P^3(f, g) + O(\hbar^5),$$

where  $P = \partial_{z_1} \otimes \partial_{z_2} - \partial_{z_2} \otimes \partial_{z_1}$ . Thus the  $\hbar^2$  coefficient of the raw star commutator vanishes in this quantization convention. The normalized quantum Hamiltonian bracket is instead

$$\{f, g\}_{\hbar} = \hbar^{-1}[f, g]_{\star} = \sum_{s \geq 0} \frac{\hbar^{2s}}{2^{2s}(2s+1)!} P^{2s+1}(f, g).$$

**PROPOSITION 5.215 (Moyal coefficients on monomials).** Let  $f = z_1^k z_2^l$  and  $g = z_1^m z_2^n$ . For every  $r \geq 0$ ,

$$P^r(f, g) = \sum_{a=0}^r (-1)^a \binom{r}{a} (k)_{r-a} (l)_a (m)_a (n)_{r-a} z_1^{k+m-r} z_2^{l+n-r},$$

where  $(s)_q = s(s-1)\cdots(s-q+1)$  and  $(s)_0 = 1$ . Terms with a negative displayed exponent are understood to be zero; equivalently the corresponding falling factorial already vanishes. Consequently

$$[f, g]_{\star} = \sum_{\substack{r \geq 1 \\ r \text{ odd}}} \frac{2}{2^r r!} \hbar^r P^r(f, g),$$

so the second-order coefficient of the raw commutator vanishes. The first four odd raw coefficients are

$$\hbar P(f, g), \quad \frac{\hbar^3}{24} P^3(f, g), \quad \frac{\hbar^5}{1920} P^5(f, g), \quad \frac{\hbar^7}{322560} P^7(f, g).$$

*Proof.* Since

$$P = \partial_{z_1} \otimes \partial_{z_2} - \partial_{z_2} \otimes \partial_{z_1},$$

the two summands commute as differential operators on the tensor product. The binomial theorem gives

$$P^r = \sum_{a=0}^r (-1)^a \binom{r}{a} \partial_{z_1}^{r-a} \partial_{z_2}^a \otimes \partial_{z_1}^a \partial_{z_2}^{r-a}.$$

Applying this to  $z_1^k z_2^l \otimes z_1^m z_2^n$  gives the displayed falling factorial formula. Interchanging  $f$  and  $g$  changes  $P^r(f, g)$  by the sign  $(-1)^r$ , so the star commutator keeps exactly odd  $r$  and contributes the factor  $2/(2^r r!)$  in order  $r$ .  $\square$

**THEOREM 5.216** (*Degree-zero Moyal target and decorated radial obstruction*). Let  $\mathfrak{h}_\hbar = \bar{A}_\hbar$  be the Moyal Hamiltonian Lie algebra, and let  $\mathfrak{h}_{\hbar,u}$  be its shifted-cotangent coordinate copy, with generator  $u_f$  labelled by  $f \in \mathfrak{h}_\hbar$ . Let

$$H_{\text{Ham,conn}}^{\circ}(\mathcal{O}_{\text{brane},\infty}^{\mathfrak{q}})_{\text{ren}}$$

denote the stable connected Hamiltonian trace sector generated by the Capelli-renormalized  $J$ -basis transition classes  $[\overline{J_N(f)}]_{N \rightarrow \infty}$ . These are associated-graded stable classes in the renormalized transition system. Naive normal-ordered block-restriction limits are different objects.

The following data are proved under the stated finite-window and radial obstruction hypotheses.

- (i) The Weyl/Moyal coefficients of Proposition 5.215 give the all-order formal Hamiltonian source. The odd raw coefficients are  $2\hbar^r P^r/(2^r r!)$ , and the first four are 1,  $1/24$ ,  $1/1920$ ,  $1/322560$ .
- (ii) The Capelli-renormalized  $J_N$ -basis of Lemma 5.197 gives the stable connected trace target as a  $\mathbb{C}[[\hbar]]$ -module, after removing the empty trace and decomposable products.
- (iii) The ordinary Weyl-trace image clause is equivalent, after the homomorphism, injectivity, and pure-power parts of the radial hypotheses, to the moment-ideal primitive  $E_{a,b,N} \in \mathcal{W}_N \widehat{\mu}(\mathfrak{sl}_N)$  of Lemma F.15. In the free normal-diagram model this is the vanishing of

$$\mathfrak{o}_{a,b} = [R_{a,b}^{\text{free}}] \in \text{coker } C_{a,b}$$

with functorial lifts  $K_{a,b} \in \ker \partial$ . Equivalently, in the dual-potential obstruction theorem,

$$\Omega_{a,b}^{\text{rad}} = [(T_{a,b}, E_{a,b}^+)] \in \text{coker } B_{a,b}$$

must vanish, or equivalently the decorated PBW Stokes identity  $D_{a,b}^{\square} = C_{a,b}^+ \partial_2$  must hold; an obstruction is a signed row  $(\phi, -\lambda) \in \ker B_{a,b}^*$  with  $\lambda(E_{a,b}^+) - \phi(T_{a,b}) \neq 0$ . Proposition F.32 proves this primitive for every edge bidegree  $(2, b)$  and  $(a, 2)$ . Proposition F.20, Proposition F.33, and the exact-arithmetic normal-form script prove finite interior identities. For all  $a, b \geq 3$ , the all-interior assertion is the associated-graded two-colour necklace homotopy, equivalently the dual-potential signed-row theorem above.

If the radial image hypotheses of Statement F.8 are supplied, or if the equivalent quantum shear primitive construction above is constructed together with the homomorphism, injectivity, and pure-power clauses, then

$$\Phi_h^{(\circ)} : \mathfrak{h}_{h,u} \xrightarrow{\sim} H_{\text{Ham,conn}}^{\circ}(\mathcal{O}_{\text{brane},\infty}^{\mathfrak{q}})_{\text{ren}}, \quad u_f \mapsto \partial_{\text{bb},h}(f) = \overline{[J_N(f)]}_{N \rightarrow \infty}$$

is the  $u_f$ -coordinate component of the desired quantum  $P_{\circ}$ -operation: it is a  $\mathbb{C}[[\hbar]]$ -linear isomorphism onto the renormalized degree-zero Hamiltonian sector. It is a Lie-algebra map on these Hamiltonian coordinates,

$$\Phi_h^{(\circ)}(u_{\{f,g\}_h}) = \hbar^{-1}[\Phi_h^{(\circ)}(u_f), \Phi_h^{(\circ)}(u_g)]_{\text{conn,raw}}.$$

Under this assumption, the associated graded at  $\hbar = 0$  is the degree-zero Hamiltonian restriction of  $\Phi$  in Theorem 5.6. The descendant  $(\mathfrak{h}^{\vee}, \text{one-antifield Koszul})$  sector remains governed by Problem 5.48; see Problem 5.221 below for the corresponding obstruction.

*Proof.* Item (1) is Proposition 5.215. Item (2) is Corollary 5.198 and Lemma 5.197. Item (3) is the chain Proposition F.14, Lemma F.15, Definition F.21, and Statement F.22, sharpened by Proposition F.25 to  $\Omega_{a,b}^{\text{rad}} = 0$ , equivalently  $D_{a,b}^{\square} = C_{a,b}^{+} \partial_2$ , with the signed-row obstruction criterion. The edge theorem is supplied by Proposition F.32. The finite interior statements cited there are proved exact free normal-diagram identities; by Lemma F.19 they are all-rank identities in their displayed windows.

Under the radial image hypotheses, the displayed  $\mathbb{C}[[\hbar]]$ -linear bijection follows from the stable Capelli-renormalized generators. Flatness over  $\mathbb{C}[[\hbar]]$  follows from the unipotent Capelli triangular system over  $\mathbb{C}[[\hbar N]]$  (Lemma 5.197(ii)); the associated graded is the classical trace map because the  $r = 0$  triangular term is the leading symbol (Lemma 5.197(iv)). The Lie identity is precisely Theorem 5.199, passed to the connected stable limit. The cotangent cochain summand is governed by the separate Tate-coefficient obstruction datum.  $\square$

**THEOREM 5.217** (*Componentwise quantum coefficient comparison*). Moyal-quantised Hamiltonians, Weyl-quantised brane operators, and the brane trace map commute on a four-component product target, with the obstruction of any other component recorded as a separate class on its own completion.

- *Components proved compatible:* weighted BV kernel convergence  $\mathcal{K}_{\text{BV}}^w$ ; balanced  $\mathfrak{gl}(N|N)$ -supertrace zero scalar QME representative  $\mathfrak{o}_{\text{sc}}$  on its scalar-contact completion; reduced weighted Moyal principal-part current target  $A_{\partial,\hbar}^{\text{pp},w}(I)$ ; verified-range degree-zero Hamiltonian trace target  $T_{\text{Ham},h}$ .
- *Minimal input data:* weighted Tate completion  $\mathfrak{h}_w$  (Theorem 5.57), Costello propagator and weighted RG calculus (Theorem 5.70), balanced supertrace replacement (Theorem E.16), Capelli-renormalised  $J_N$ -basis transitions (Lemma 5.197), and the explicit enumeration cited in clause (4).
- *Obstruction classes outside this product:* the ordinary- $\mathfrak{gl}_N$  scalar QME class  $\hbar N[\bar{c}] \in H_{\text{Lie}}^2(\bar{A}, \mathbb{C})[[\hbar]]$  (Theorem 5.213); the all-bidegree radial/Weyl normal-form class  $\Omega_{a,b}^{\text{rad}} \in \text{coker } B_{a,b}$  (Proposition F.25); the reduced non-scalar Costello QME class  $[\text{Ob}_{w,\partial,\hbar}^{\text{red}}] \in H^1(\ker \widehat{\partial}_{\text{sc}}, d_{\text{QME}})$  (Problem 5.219); the  $\theta_3$  bare / supplied-counterterm dichotomy (Theorem E.38).
- *Vanishing criterion for the four named components:* the displayed equalities of items (1)–(4) hold on the scalar-contact balanced completion and on the verified range listed in Proposition F.32, Proposition F.31, Proposition F.33, and Theorem 5.216.
- *First nonzero example outside the proved components:* the ordinary scalar branch is blocked at order  $\hbar^1$  on linear generators by  $[\Phi_h(z_1), \Phi_h(z_2)]_{\star} = \hbar N$ ; the reduced non-scalar Costello QME branch is blocked at the smallest non-scalar window  $\theta_3$ , where the bare  $H^1 = \mathbb{C} \cdot E_{\theta_3} \neq 0$ .

- *Physical meaning:* the Moyal star, the matrix Weyl algebra, and the brane trace map agrees component-wise on the proved completion; mismatch on any of the four obstruction classes is the precise location at which compatibility breaks and a counterterm, source replacement, or scalar-contact lift is required.

*Statement.* Fix an exponential degree weight  $w(d) = q^d$ ,  $q > 1$ , and work over  $\mathbb{C}[[\hbar]]$ . Write

$$\mathfrak{h}_{w,\hbar} = \mathfrak{h}_w[[\hbar]], \quad \mathfrak{h}_{\hbar,\text{cont}}^{\vee,w} = \mathfrak{h}_{\text{cont}}^{\vee,w}[[\hbar]].$$

The Moyal bracket on  $\mathfrak{h}_{w,\hbar}$  is transported from  $\tilde{A}_{\hbar}$  by the diagonal weight rescaling of Definition 5.54. Let

$$\mathcal{K}_{\text{BV}}^w = (C_{\mathfrak{h},w}, P_{\epsilon,L}^w, \mathcal{E}_w)$$

be the weighted coefficient-kernel datum of Theorem 5.70. For an interval  $I \subset \mathbb{R}$ , let  $A_{\partial,\hbar}^{\text{pp},w}(I)$  be the weighted reduced Moyal principal-part current envelope

$$\mathfrak{h}_{\hbar,I}^w = \Omega_c^\bullet(I) \widehat{\otimes} \mathfrak{h}_{w,\hbar}, \quad \mathcal{P}_{\hbar,I}^w = \mathcal{D}'_c(I) \widehat{\otimes} \mathfrak{h}_{\hbar,\text{cont}}^{\vee,w},$$

$$A_{\partial,\hbar}^{\text{pp},w}(I) = \widehat{\mathbf{S}}_{\mathbb{C}[[\hbar]]}(\mathfrak{h}_{\hbar,I}^w) \widehat{\otimes} \widehat{\mathbf{S}}_{\mathbb{C}[[\hbar]]}(\mathcal{P}_{\hbar,I}^w[1]).$$

Its dense strict Matlis subtarget is the reduced Moyal principal-part target  $A_{\partial,\hbar}^{\text{pp}}(I)$  of Construction 5.134, obtained by replacing  $\mathfrak{h}_{w,\hbar}$  with  $\tilde{A}_{\hbar}$  and  $\mathfrak{h}_{\hbar,\text{cont}}^{\vee,w}$  with  $\mathcal{P}[[\hbar]]$ . The Moyal coadjoint action on the weighted envelope is the finite-window continuous extension of the action in that construction; the assertion is restricted to the vertexwise polynomial-degree-bounded weighted sector. Let  $T_{\text{Ham},\hbar} = H_{\text{Ham,conn}}^\circ(O_{\text{brane},\infty}^{\text{q}})_{\text{ren}}$  be the stable connected Hamiltonian trace target of Theorem 5.216. Its formal Moyal and Capelli pieces are proved there; the ordinary-trace Lie comparison is reduced to the dual-potential radial obstruction  $\Omega_{a,b}^{\text{rad}}$ , equivalently the decorated PBW Stokes identity  $D_{a,b}^\square = C_{a,b}^+ \partial_2$ , with all edge strips proved and all-interior closure requiring the signed-row vanishing theorem. In the balanced  $\mathfrak{gl}(N | N)$  supertrace branch put  $\mathfrak{o}_{\text{sc}}$  for the zero scalar QME representative of Theorem E.16; on a balanced scalar-contact completion this is the zero filtered scalar projection of Proposition E.18.

The componentwise quantum coefficient comparison is the product prefactorization target

$$\mathfrak{S}_{\hbar,w}(I) = \mathcal{K}_{\text{BV}}^w \times \{\mathfrak{o}_{\text{sc}}\} \times A_{\partial,\hbar}^{\text{pp},w}(I) \times T_{\text{Ham},\hbar},$$

where the first, second, and fourth factors are constant in  $I$ , and the principal-part current factor uses extension by zero and the factorization product of Theorem 5.185. This product target carries the following proved data.

- The weighted Casimir  $C_{\mathfrak{h},w}$  and the finite-scale BV propagator  $P_{\epsilon,L}^w$  converge in the weighted Tate coefficient category. Each fixed Costello graph with vertexwise polynomial-degree-bounded Hamiltonian labels remains in that category.
- On every balanced scalar-contact completion of Definition E.17, the balanced  $\mathfrak{gl}(N | N)$  supertrace model has zero filtered scalar projection  $\widehat{\mathfrak{o}}_{\text{bal,sc}} = \mathfrak{o}$ , zero chain-map defect, and vanishing scalar-projection obstruction classes  $\mathfrak{o}_{\sigma,w,\text{bal}}^{(r)}(I)$ . In the ordinary scalar-reduced  $\mathfrak{gl}_N$  branch the scalar class is instead  $\hbar N[\bar{c}]$  and requires the alternative mechanisms of Theorem 5.213.
- The reduced weighted quantum cotangent current brackets are

$$\{B_f(a), B_g(b)\}_{\hbar} = B_{\{f,g\}_{\hbar}}(ab), \quad \{B_f(a), \Theta_\rho(B)\}_{\hbar} = \Theta_{\text{ad}_{f,\hbar}^\vee \rho}(aB),$$



$$\{\Theta_\rho(B), \Theta_\sigma(C)\}_\hbar = 0.$$

Here  $\text{ad}_{f,\hbar}^\vee$  is the Moyal coadjoint action on  $\mathfrak{h}_{\hbar,\text{cont}}^{\vee,w}$ , whose restriction to  $\mathcal{P}[[\hbar]]$  is the action of Construction 5.134. This is the Moyal coadjoint action, distinct from the polynomial one- $\psi$  PBW action. Thus the classical principal-part current construction is transported to the quantum target only after replacing  $f \cdot \rho$  by the full Moyal coadjoint action and after imposing weighted finite-window continuity.

- (iv) Under the radial image hypotheses, equivalently after solving the quantum shear primitive construction of Theorem 5.216, the degree-zero Hamiltonian generator map

$$\Phi_h^{(o)} : \mathfrak{h}_{h,u} \xrightarrow{\sim} T_{\text{Ham},h}$$

is the  $\mathbb{C}[[\hbar]]$ -linear Lie comparison of Theorem 5.216.

Consequently the weighted coefficient/BV kernel, the balanced scalar-contact zero projection, the reduced weighted Moyal principal-part current target, and the radial-constrained degree-zero Moyal coefficient comparison give the componentwise theorem. The full non-scalar Costello graph realization, unreduced factorization centre morphism, and all-interior radial theorem are separate constructions governed by the obstruction formulation above.

*Proof.* Item (1) is Theorem 5.70: its parts (a)–(c) give Casimir convergence, heat-kernel propagator extension, and graphwise weighted RG compatibility. Item (2) is Theorem E.16 together with Proposition E.18; the ordinary scalar alternatives and obstruction criterion are assembled in Theorem 5.213.

At  $\hbar = 0$ , the current brackets are the reduced principal-part brackets of Theorem 5.185 and Corollary B.8. The quantum target is obtained from the completed continuous-dual coadjoint action, with the polynomial one- $\psi$  labels fixed. On the dense strict Matlis subtarget it is the semidirect Moyal current complex of Construction 5.134, whose action is by  $(\text{ad}_{f,\hbar}^\vee \rho)(g) = -\rho(\{f, g\}_\hbar)$ . The extension from  $\mathcal{P}[[\hbar]]$  to  $\mathfrak{h}_{\hbar,\text{cont}}^{\vee,w}$  is windowwise continuous: the Moyal coadjoint formula is finite for each polynomial Hamiltonian, and condition (W2) in Definition 5.51 gives the required finite-window bounds for vertexwise polynomial-degree-bounded inputs. Proposition 5.133 and Theorem 5.135 show why this hypothesis is necessary: the PBW one- $\psi$  label map has a nonzero intertwining obstruction for the Moyal coadjoint action. Hence item (3) is a target-side reduced current construction with the correct quantum action; transport from polynomial descendants is a separate problem.

Item (4) is Theorem 5.216: its formal Moyal, Capelli, and edge-strip pieces are proved, while the all-interior ordinary-trace comparison is exactly Proposition F.25, namely  $\Omega_{a,b}^{\text{rad}} = 0$ , equivalently decorated PBW Stokes  $D_{a,b}^\square = C_{a,b}^+ \partial_2$ , with signed  $B_{a,b}^*$ -rows as the obstruction functionals. The product prefactorization structure is componentwise: the weighted BV kernel, scalar zero representative, and stable trace target are constant in the interval, while the principal-part current factor is functorial by extension by zero and has the disjoint-interval product of Theorem 5.185. These components are independent targets, so their product carries exactly the displayed product comparison; Costello graph/QME assertions require the separate obstruction complex.  $\square$

*Remark 5.218 (Computer verification).* The explicit clause of Theorem 5.217 is verified by exact rational arithmetic. `scripts/check_moyal_coefficients.py` runs the Procesi–Razmyslov–Capelli inverse identity and the prequotient scalar Moyal expansion on the rectangular window  $0 \leq k, l, m, n \leq 10$  to Moyal order  $\leq 11$ , confirming the identities of Proposition F.32 and Theorem 5.216.

scripts/check\_one\_psi\_homology.py verifies the descendant action and the residue-coadjoint formula  $z_1^p z_2^q \cdot \rho_{a,b} = -(pb - qa + p - q)\rho_{a-p+1, b-q+1}$  of Theorem 5.178 on  $0 \leq k, l \leq 5$ . The radial Stokes boundary  $B_{a,b}$  and decorated weight  $C_{a,b}^+$  are built as rational matrices;  $\ker B_{a,b}^*$  and  $\text{coker } B_{a,b}$  follow by exact Gaussian elimination. Outside the window, the all-bidegree obstruction  $\Omega_{a,b}^{\text{rad}} \in \text{coker } B_{a,b}$  is the cohomology class blocking extension (Proposition F.25).

*Problem 5.219 (Non-scalar quantum  $P_o$ -operation obstruction complex).* After the four proved components of Theorem 5.217, the remaining non-scalar Costello QME comparison reduces to a single cohomology class on a single completion.

- *Desired upgrade:* a chain-level Costello-local  $P_{o,h}$ -factorisation-centre map  $C_{\text{CE,cont}}^\bullet(\mathfrak{h}_{w,h} \rtimes \mathfrak{h}_{h,\text{cont}}^{\vee,w}[1]) \rightarrow Z_{\text{fact}}^{P_{o,h}}(\text{Obs}_{\text{open}}^h)|_{\mathfrak{S}_{h,w}}$ , curved bulk-to-defect kernel  $\kappa_{h,w,I}$  included.
- *Minimal input data:* a complete  $d_{\text{QME}}$ -stable scalar-contact filtration with vanishing successive scalar-symbol extension classes  $o_{\sigma,w}^{(r)}(I)$ ; a filtered chain projection  $\widehat{\sigma}_{\text{sc}}$ ; a curved bulk-to-defect kernel  $\kappa_{h,w,I}$  on the current source of Definition B.14; compatible local counterterms  $C_{\Gamma,w}^{w_o}(\epsilon)$  with finite-window transport.
- *Obstruction class:* the reduced non-scalar QME cohomology class  $[\text{Ob}_{w,\partial,h}^{\text{red}}] \in H^1(\ker \widehat{\sigma}_{\text{sc}}, d_{\text{QME}})$ , together with its finite-window  $\varprojlim^1 H^0$ -primitive-compatibility class.
- *Vanishing criterion:* successive vanishing of  $o_{\sigma,w}^{(r)}(I)$  for  $r > 0$  builds  $\widehat{\sigma}_{\text{sc}}$ ; on a fixed window the row equation  $A^M c = -r^M$  is solvable iff  $[\mathfrak{o}_{n_o,M}^{\text{ns}}]$  lies in  $-\text{im } H^1(\Xi_M)$ ; in the inverse-limit tower vanishing of the closed-exchange triple  $(\beta_{n_o}^{\text{cl}}, \mu_{n_o}^{\text{cl}}, \lambda_{n_o}^{\text{cl}})$  is necessary and sufficient.
- *First nonzero example or known obstruction:* the smallest non-scalar window  $\theta_3$  carries the bare  $H^1 = \mathbb{C} \cdot E_{\theta_3,M} \neq 0$  with covector  $\ell_{\theta_3}(E_{\theta_3,M}) = 1$ ; a supplied scalar-zero counterterm  $C_{\theta_3,M}$  with  $d_M C_{\theta_3,M} = -E_{\theta_3,M}$  forces  $c = 1$  unique and  $H^1 = 0$  (Theorem E.38); the lower source-coordinate Bianchi route of Proposition 5.220 exhibits a nontrivial cokernel  $\widetilde{\lambda}_{02,b}$  surviving the current Bianchi-exposed completion.
- *Physical meaning:* the unreduced Costello-local closed-exchange or local-counterterm cancellation of the non-scalar QME residual is the bulk-to-defect Ward identity compatible with the quantum moment-map constraint.

*Statement.* The non-scalar Costello graph/QME obstruction criterion is the vanishing problem in the filtered local-functional complex

$$\mathfrak{Q}_{w,\partial,h}^\bullet(I) = \left( \mathcal{O}_{\text{loc}}^{\text{bd}}(\mathcal{E}_w|_I; A_{\partial,h}^{\text{pp},w}(I)), d_{\text{QME}} = Q + \{I_o^w, -\}_h \right),$$

where the local functionals have compact support on the brane line, coefficients in the weighted field complex  $\mathcal{E}_w$ , and target currents in  $A_{\partial,h}^{\text{pp},w}(I)$ . The scalar-symbol projection is only associated-graded data until it is lifted through the scalar-contact filtration. The first obstruction is the filtered lift of the scalar projection; the reduced non-scalar class is reached after this lift. Choose a complete  $d_{\text{QME}}$ -stable scalar-contact filtration  $F^\bullet \mathfrak{Q}_{w,\partial,h}^\bullet(I)$ . For any filtered linear lift  $s$  of the scalar-symbol map, the successive chain-map defects define classes

$$o_{\sigma,w}^{(r)}(I) \in H^1\left(\text{Hom}(\text{gr}_F \mathfrak{Q}_{w,\partial,h}^\bullet(I), C_{\text{Lie}}^\bullet(\bar{A}; C_{\mathcal{H}}[1][[\hbar]])_{-r}\right), \quad r > 0,$$

with the Hom differential of Lemma E.4. The first condition is the successive vanishing of these  $o_{\sigma,w}^{(r)}(I)$ , with compatible choices in the completed finite-window inverse limit. Only then is there a filtered chain projection

$$\widehat{\sigma}_{\text{sc}}: \mathfrak{Q}_{w,\partial,\hbar}^\bullet(I) \longrightarrow C_{\text{Lie}}^\bullet(\bar{A}; \mathbb{C}\gamma_1)[1][[\hbar]]$$

whose associated graded is the scalar-symbol projection of Definition E.3.

The balanced scalar-contact branch has a vanishing subcomplex: on  $\mathfrak{S}_{\text{bal,sc}}^\bullet(I) \subset \mathfrak{Q}_{w,\partial,\hbar,\text{str}}^\bullet(I)$ , Proposition E.18 gives the zero filtered chain projection  $\widehat{\sigma}_{\text{bal,sc}} = 0$  and kills every  $o_{\sigma,w,\text{bal}}^{(r)}(I)$ . Extending this result to the full graph/counterterm completion requires an additional construction: either prove that the relevant scalar-contact symbols still factor through  $\text{sdim } \mathbb{C}^{N|N}$ , or compute and kill the extension classes of Remark E.19.

After this lift has been constructed, the balanced supertrace branch requires a continuous bulk-to-defect kernel

$$\kappa_{\hbar,w,I}: C_{\text{CE,cont}}^\bullet(\mathfrak{g}_{\hbar,I}^{w,\text{cur}}) \longrightarrow \mathfrak{Q}_{w,\partial,\hbar}^\bullet(I)$$

for the current source of Definition B.14. Its curvature

$$\text{Ob}_{w,\partial,\hbar} = d_{\text{QME}} \kappa_{\hbar,w,I} + \frac{1}{2} [\kappa_{\hbar,w,I}, \kappa_{\hbar,w,I}]_{\hbar}$$

must be a cocycle satisfying

$$\widehat{\sigma}_{\text{sc}}(\text{Ob}_{w,\partial,\hbar}) = 0$$

and the remaining reduced non-scalar class

$$[\text{Ob}_{w,\partial,\hbar}^{\text{red}}] \in H^1(\ker \widehat{\sigma}_{\text{sc}}, d_{\text{QME}})$$

must vanish. At finite windows this is the tower  $\mathcal{Q}_{w,M}^\bullet(I) = \ker \widehat{\sigma}_{\text{sc},M}$  of Definition 5.82. The compatible counterterm recursion is exactly the criterion of Theorem 5.83: at each  $\hbar$ -order the inverse-limit  $H^1$ -class and the  $\varprojlim^1 H^0$ -primitive-compatibility class must vanish.

More explicitly, let  $n_o$  be the first order where lower-order choices leave a residual

$$\mathfrak{o}_{n_o,M}^{\text{ns}} \in Z^1(\mathcal{Q}_{w,M}^\bullet(I)).$$

Counterterms alone close this order exactly on the vanishing locus of the obstruction vector

$$\mathfrak{D}_{n_o}^{\text{ns}} = (([\mathfrak{o}_{n_o,M}^{\text{ns}}])_M, \lambda_{n_o})$$

of Theorem 5.83. A closed-exchange repair is additional cochain data: finite-window complexes  $\mathcal{X}_{w,M}^\bullet(I)$ , compatible cochain maps

$$\Xi_M: \mathcal{X}_{w,M}^\bullet(I) \longrightarrow \mathcal{Q}_{w,M}^\bullet(I),$$

and cochains satisfying the branch equation

$$\mathfrak{o}_{n_o,M}^{\text{ns}} + \Xi_M(W_{n_o,M}) + d_M C_{n_o,M} = 0, \quad W_{n_o,M} \in Z^1(\mathcal{X}_{w,M}^\bullet(I)), \quad C_{n_o,M} \in \mathcal{Q}_{w,M}^0(I).$$

For a fixed window this equation is solvable if and only if

$$[\mathfrak{o}_{n_o,M}^{\text{ns}}] \in -\text{im}\left(H^1(\Xi_M): H^1(\mathcal{X}_{w,M}^\bullet(I)) \rightarrow H^1(\mathcal{Q}_{w,M}^\bullet(I))\right).$$

In the inverse-limit tower the exact closed-exchange obstruction is the triple

$$\left( \beta_{n_o}^{\text{cl}}, \mu_{n_o}^{\text{cl}}, \lambda_{n_o}^{\text{cl}} \right),$$

where

$$\begin{aligned} \beta_{n_o}^{\text{cl}} &\in \text{coker} \left( \varprojlim_M H^1(\mathcal{X}_{w,M}^\bullet(I)) \xrightarrow{\lim H^1(\Xi_M)} \varprojlim_M H^1(\mathcal{Q}_{w,M}^\bullet(I)) \right), \\ \mu_{n_o}^{\text{cl}} &\in \varprojlim_M H^0(\mathcal{X}_{w,M}^\bullet(I)), \quad \lambda_{n_o}^{\text{cl}} \in \varprojlim_M H^0(\mathcal{Q}_{w,M}^\bullet(I)). \end{aligned}$$

Vanishing of this triple is the exact typed criterion for a compatible closed-exchange kernel plus local counterterm to kill the first non-scalar residual. Constructing  $\mathcal{X}^\bullet$ ,  $\Xi$ , and such a class  $[W]$  remains open.

For a fixed binary generator pair  $x, y$ , the row form of the centrality obstruction is the same elementary finite-window matrix problem. Once the finite graph data supply a scalar-zero basis  $\rho_i^M$  for the degree  $|x| + |y|$  residual rows and candidate homotopy rows  $\eta_j^M$  of degree  $|x| + |y| - 1$ , write

$$R_{x,y,M}^{\text{cent}} = \sum_i r_i^M \rho_i^M, \quad d_M \eta_j^M = \sum_i A_{ij}^M \rho_i^M.$$

Then  $H_{x,y}^M = \sum_j h_j^M \eta_j^M$  is a binary centrality homotopy in the supplied finite row span if and only if

$$A^M h^M = -r^M.$$

The compatible inverse-limit homotopy additionally requires

$$[P^{M,N} h^M - h_{\pi x, \pi y}^N] = 0 \quad \text{in} \quad H^{|x|+|y|-1}(\mathcal{Q}_{w,N}^\bullet(I)).$$

Thus the coefficient-current zero row supplies evidence only for the coefficient-current row; the enlarged Costello row is the displayed matrix obstruction.

Equivalently, one must construct compatible finite-window local counterterms  $C_{\Gamma,w}^{w_o}(\epsilon)$  and the bulk-to-defect kernel  $\kappa_{\hbar,w,I}$  such that

$$\text{Curv}(\kappa_{\hbar,w,I} + C_{\hbar,w}) = d_{\text{QME}}(\kappa_{\hbar,w,I} + C_{\hbar,w}) + \frac{1}{2} [\kappa_{\hbar,w,I} + C_{\hbar,w}, \kappa_{\hbar,w,I} + C_{\hbar,w}]_{\hbar} = 0,$$

with finite-window compatibility

$$\pi_{w_o'w_o} C_{\Gamma,w}^{w_o'} = C_{\Gamma,w}^{w_o}, \quad R_{w,w'}^{w_o} C_{\Gamma,w}^{w_o} = C_{\Gamma,w'}^{w_o}.$$

This is the precise obstruction complex behind the desired

$$C_{\text{CE,cont}}^\bullet(\mathfrak{h}_{w,\hbar} \ltimes \mathfrak{h}_{\hbar,\text{cont}}^{\vee,w}[1]) \longrightarrow Z_{\text{fact}}^{P_o,\hbar}(\text{Obs}_{\text{open}}^{\hbar})|_{\mathfrak{S}_{\hbar,w}}.$$

**PROPOSITION 5.220** (*Finite-window criterion for the first  $\theta_3$  row*). In a larger finite-window graph system, let  $\Theta_3$  be a genuine marked source-face row of  $\hbar$ -order 3. The exposed one-row scalar-zero summand is

$$\mathcal{Q}_{\theta_3,M}^o = 0, \quad \mathcal{Q}_{\theta_3,M}^1 = \mathbb{C}E_{\theta_3,M}, \quad d_M = 0,$$

with

$$E_{\theta_3,M} = -K_{\Theta_3}^M(\zeta_{M,\nu_3}), \quad \mathfrak{v}_{\theta_3,M}^{\text{ns}} = E_{\theta_3,M}, \quad \widehat{\sigma}_{\text{sc},M}(E_{\theta_3,M}) = 0.$$

In this supplied subcomplex the class  $[E_{\theta_3, M}]$  is nonzero. It is killed in an enlarged finite-window system precisely by one of the following data, together with scalar-zero and finite-window transport compatibility.

First, a CE-source ancestor

$$\eta_{\theta_3, M}, \quad d_{\text{CE}} \eta_{\theta_3, M} = \zeta_{M, \nu_3},$$

whose induced degree-zero row  $C_{\theta_3, M}^{\text{CE}} = K_{\Theta_3}^M(\eta_{\theta_3, M})$  satisfies

$$d_M C_{\theta_3, M}^{\text{CE}} = -E_{\theta_3, M}.$$

Second, a Costello local counterterm primitive  $C_{\theta_3, M}^{\text{ct}} \in Q_{w, M}^{\circ}(I)$  satisfying

$$d_M C_{\theta_3, M}^{\text{ct}} = -E_{\theta_3, M}, \quad \widehat{\sigma}_{\text{sc}, M}(C_{\theta_3, M}^{\text{ct}}) = 0.$$

Third, a complete companion-face table  $F_{\Theta_3, M}$  with signs  $\epsilon_{\Theta_3, \nu}^{\text{CE}}$  and rows  $E_{\Theta_3, \nu}^M$  such that

$$\sum_{\nu \in F_{\Theta_3, M}} \epsilon_{\Theta_3, \nu}^{\text{CE}} E_{\Theta_3, \nu}^M = 0, \quad E_{\Theta_3, \nu_3}^M = E_{\theta_3, M}.$$

In the first two cases the finite-window tower must also satisfy

$$[\pi_{M, N} C_{\theta_3, M} - C_{\theta_3, N}] = 0 \quad \text{in} \quad H^{\circ}(Q_{w, N}^{\bullet}(I)).$$

In the companion-face case the source-face transport matrices must satisfy

$$\pi_{M, N} E_{\Theta_3, \nu}^M = \sum_{\nu'} v_{\Theta_3, \nu; \Theta_3, \nu'}^{M, N} E_{\Theta_3, \nu'}^N, \quad v^{M, P} = v^{N, P} v^{M, N},$$

and must preserve the displayed signed zero sum. A formal symbol  $C_{\theta_3}$  supplied apart from the differential  $d_M C_{\theta_3, M} = -E_{\theta_3, M}$ , scalar-zero value, and tower compatibility is merely notation; a counterterm consists of all three data.

The currently tested finite source routes are obstructed in a sharper sense. In the ordinary pure two- $u$  CE source envelope, in every finite Hamiltonian degree, the covector

$$q_{2u} = e_{((H_{0,1}, H_{2,0}), c_{\rho_{2,1}})}^* - \frac{1}{2} e_{((H_{0,1}, H_{0,1}), c_{\rho_{0,2}})}^*$$

satisfies

$$q_{2u} d_{\text{CE}} = 0, \quad q_{2u}(\zeta_{M, \nu_3}) = 1.$$

Hence the ordinary pure two- $u$  degree enlargement has an obstruction to supplying a CE ancestor for  $\zeta_{M, \nu_3}$ . This is an exact obstruction theorem for that source envelope; marked-source, Hom-valued companion, and local-functional counterterm completions are separate source categories. The tested eight-face companion row has coordinates

$$(2, -2, 3, 2, -1, 1, -2, -3)$$

on

$$E_{\nu_{02}}, E_{\nu_{20}}, E_{\nu_{03}}, E_{\nu_{12}^{(1)}}, E_{\nu_{12}^{(2)}}, E_{\theta_3}, E_{\nu_{21}^{(2)}}, E_{\nu_{30}},$$

and diagonal transport to the  $N = 2$  window leaves  $2E_{\nu_{02}}^2 - 2E_{\nu_{20}}^2$ . The covector  $q_{2u}$  evaluates to zero on this eight-face row, so the row is incompatible with the required marked source column. The normalized lower-window cokernel functional  $\lambda_{\nu_{02}}^{(2)} = \frac{1}{2}(E_{\nu_{02}}^2)^*$  evaluates to 1 on the transported residual. A repair must therefore supply either a scalar-zero column  $A_{\theta,C}^M = -1$ , a marked source generator outside the ordinary pure two- $u$  algebra with  $q_{2u}$ -value 1, or a companion relation whose lower-window transport kills this residual by a derived non-diagonal transport, lower primitive, or relation  $E_{\nu_{02}}^2 = E_{\nu_{20}}^2$ . The canonical lower source-coordinate candidate is

$$\zeta_2^o = u_{a,H_{1,0}} u_{b,H_{0,1}}, \quad D_{02,20}^2 = K_{\Theta_3}^2(\zeta_2^o),$$

with source differential

$$d_{\text{CE}} \zeta_2^o = 2\zeta_{\nu_{02}} - 2\zeta_{\nu_{20}}.$$

This proves source exactness of the transported lower row. It proves local-functional exactness only after the Hom-valued Bianchi defect

$$\mathfrak{b}_{02,20}^2 = d_{\text{ns},2} K_{\Theta_3}^2(\zeta_2^o) - K_{\Theta_3}^2(d_{\text{CE}} \zeta_2^o)$$

is killed:

$$d_{\text{ns},2} D_{02,20}^2 = -2E_{\nu_{02}}^2 + 2E_{\nu_{20}}^2 + \mathfrak{b}_{02,20}^2.$$

In the Bianchi-exposed lower basis

$$(E_{\nu_{02}}^2, E_{\nu_{20}}^2, \mathfrak{b}_{02,20}^2)$$

the column of  $D_{02,20}^2$  and the residual are

$$A_D^2 = (-2, 2, 1)^T, \quad r_2 = (2, -2, 0)^T.$$

The equation  $A_D^2 c = -r_2$  is obstructed in this free presentation. The cokernel

$$\tilde{\lambda}_{02,\mathfrak{b}} = \frac{1}{2}(E_{\nu_{02}}^2)^* + (\mathfrak{b}_{02,20}^2)^*$$

satisfies

$$\tilde{\lambda}_{02,\mathfrak{b}} A_D^2 = 0, \quad \tilde{\lambda}_{02,\mathfrak{b}}(r_2) = 1.$$

Thus the lower source primitive is a genuine local-functional primitive only in the quotient where  $\mathfrak{b}_{02,20}^2 = 0$ , or after a scalar-zero local counterterm  $B_{02,20}^2$  with  $d_{\text{ns},2} B_{02,20}^2 = -\mathfrak{b}_{02,20}^2$  is supplied, with locality and transport compatibility.

In the current Bianchi-exposed finite-row completion the  $B_{02,20}^2$  column is omitted. The only degree-zero lower column is  $D_{02,20}^2$ , so the desired Bianchi-counterterm target is the vector

$$-\mathfrak{b}_{02,20}^2 = (0, 0, -1)^T.$$

It lies outside the image of  $A_D^2$ , since

$$\tilde{\lambda}_{02,\mathfrak{b}} A_D^2 = 0, \quad \tilde{\lambda}_{02,\mathfrak{b}}((0, 0, -1)^T) = -1.$$

The minimal controlled enlargement must therefore add a genuine scalar-zero local-functional column

$$A_B^2 = (0, 0, -1)^T, \quad d_{\text{ns},2} B_{02,20}^2 = -\mathfrak{b}_{02,20}^2,$$

with regular-density or wavefront-admissible locality and compatible finite-window transport. In a tower this requires

$$\Delta_{M,N}^1 := -\pi_{M,N} \mathbf{b}^M + \mathbf{b}^N = d_{\text{ns},N}(\pi_{M,N} B^M - B^N)$$

together with vanishing of the  $H^1$  Bianchi transport class on the right-hand side and then of the resulting secondary  $\varprojlim^1 H^0$ -class after any supplied transport correction.

*Proof.* The one-row complex used here has zero degree-zero term and has one degree-one basis vector. Hence  $H^1(Q_{\theta_3,M}^\bullet) \cong \mathbb{C}E_{\theta_3,M}$ . The covector

$$\ell_{\theta_3}(E_{\theta_3,M}) = 1$$

satisfies  $\ell_{\theta_3} d_M = 0$  and evaluates nontrivially on the residual, so it is a left-cokernel functional for the nonzero fixed-window  $H^1$ -class.

After adjoining any finite collection of degree-zero candidate rows, the fixed-window question is the elementary linear system

$$A^M c^M = -r^M,$$

where  $A^M$  is the matrix of  $d_M$  from candidate rows to residual rows and  $r^M$  is the coordinate vector of  $\mathfrak{o}_{\theta_3,M}^{\text{ns}}$ . A CE ancestor or local counterterm with differential coefficient  $-1$  is exactly a column of  $A^M$  equal to  $-E_{\theta_3,M}$ , so  $c^M = 1$  solves the equation. If the candidate is named while this differential entry is omitted, the old covector  $\ell_{\theta_3}$  still annihilates the image of  $A^M$  and still evaluates to 1 on  $r^M$ ; the obstruction survives.

The lower source-coordinate candidate illustrates the distinction between source exactness and local-functional exactness. The source calculation gives  $d_{\text{CE}} \zeta_2^0 = 2\zeta_{\nu_{02}} - 2\zeta_{\nu_{20}}$ , but  $K_{\Theta_3}^2$  becomes a chain map after its Hom-valued Bianchi defect is killed. In the Bianchi-exposed basis the matrix equation is  $A_D^2 c = -r_2$ , with  $A_D^2 = (-2, 2, 1)^T$  and  $r_2 = (2, -2, 0)^T$ . The first two coordinates force  $c = 1$ , while the Bianchi coordinate forces  $c = 0$ . Equivalently  $\widehat{\lambda}_{02,b} A_D^2 = 0$  and  $\widehat{\lambda}_{02,b}(r_2) = 1$ . Hence this computation gives an exact obstruction class. A completed local-functional counterterm requires the additional column displayed below.

The same cokernel proves absence of the  $B_{02,20}^2$  column in the current completion. The target vector  $(0, 0, -1)^T$  would require a scalar  $c$  with  $c(-2, 2, 1) = (0, 0, -1)$ ; the first two coordinates force  $c = 0$ , while the third forces  $c = -1$ . Equivalently  $\widehat{\lambda}_{02,b}$  annihilates every current boundary column and evaluates to  $-1$  on the target. A formal new column  $A_B^2 = (0, 0, -1)^T$  closes this fixed matrix equation, but it defines a theorem only after the local functional, scalar-zero value, locality or wavefront data, and finite-window transport data are constructed.

A companion-face table is a signed source-face relation. The primitive is a separate row datum. It changes the residual row by replacing the one-face CE source expression with the full signed CE boundary. It is accepted exactly when the signed sum is zero and the truncation matrices preserve that zero sum in the inverse system. The displayed Roos condition is precisely the obstruction to choosing the fixed-window primitives compatibly. These are necessary and sufficient finite-window conditions; the analytic Costello graph construction and all-graph QME solution require additional data.  $\square$

*Problem 5.221 (Tate-coefficient lift of the descendant cochain isomorphism).* In the unweighted Tate pair, the diagnosis of Lemma 5.211 identifies the obstruction as the coefficient-kernel obstruction; the resolution requires a category-level change of coefficient module. The classical Tate obstruction is closed by

the weighted Tate completion of Theorem 5.70(a)–(d); the all-order quantum lift on the descendant sector is the quantum specialization of 5.48 under the hypothesis of 5.72.

**PROPOSITION 5.222** (*Open-line Weyl boundary product from midpoint kernel*). For Weyl-ordered Hamiltonian boundary observables placed on the open line  $\mathbb{R}$  in the reduced first-order open model  $S_o = \int_{\mathbb{R}} \text{Tr}(\phi_1 d\phi_2)$ , on two brane-time points  $(t, t') \in \mathbb{R}_{\text{brane}}^2$  (with  $s$  reserved for the transverse  $\mathbb{R}_s$  coordinate of  $X$ ) and the zero-mode-removed midpoint propagator

$$G(t, t') = \frac{1}{2} \text{sgn}(t - t'), \quad \langle \phi_1(t) \phi_2(t') \rangle = \hbar G(t, t'), \quad \langle \phi_2(t) \phi_1(t') \rangle = -\hbar G(t, t'),$$

the boundary product on  $\bar{A}_{\hbar}$  is

$$F \star G = m \circ \exp\left(\frac{\hbar}{2} P_{\partial}\right)(F \otimes G), \quad P_{\partial} = \partial_{z_1} \otimes \partial_{z_2} - \partial_{z_2} \otimes \partial_{z_1}.$$

Equivalently, every odd raw commutator coefficient is  $\frac{2}{2^r r!} \hbar^r P^r(f, g)$ . In particular the first and third coefficients are  $\hbar P(f, g)$  and  $\frac{\hbar^3}{24} P^3(f, g)$ . This reduced open-line computation is independent of Costello specialization. With these conventions, the Weyl normalization is determined by:

- (i) the matrix Weyl algebra commutator  $[(\phi_1)^i_j, (\phi_2)^k_l] = \hbar \delta_l^i \delta_j^k$ ;
- (ii) the zero-mode-removed midpoint propagator  $G(\pm, \mp) = \pm \frac{1}{2}$ ;
- (iii) the Baker–Campbell–Hausdorff determination of the cross-contraction weight  $P/2$  on linear Hamiltonians;
- (iv) the linear-Hamiltonian commutator  $[\widehat{u}, \widehat{v}] = \hbar(ad - bc)$  for  $u = az_1 + bz_2, v = cz_1 + dz_2$ .

*Proof.* Items (i)–(iv) form a normalization system. By (i)–(ii) the midpoint propagator at  $t \neq t'$  has weight  $\pm \hbar/2$ . By (iv) the linear-Hamiltonian Weyl commutator is  $\hbar(ad - bc)$ , so the cross-contraction weight on linear Hamiltonians is exactly  $1/2$ , giving  $P/2$ . Hence by (iii) the BCH formula on the Heisenberg algebra of linear Hamiltonians fixes the cross-contraction kernel on the boundary as  $P_{\partial} = P$  with weight  $\hbar/2$ .

The all-order boundary product on  $\bar{A}_{\hbar}$  is then determined by: (a) the local Weyl/BCH computation for the constant Poisson tensor, with Kontsevich deformation quantization [7] serving only as general context here; (b) Proposition 5.215 together with item (iii) above; (c) the antisymmetrisation argument of Theorem 5.223.

The scalar-normalization clause is the content of Lemma 5.197(iii): the Capelli contact term is absorbed into the  $T \leftrightarrow J_N$  change of basis, and the connected single-trace projection is the standard primitive projection. Consequently all odd raw commutator coefficients are  $\frac{2}{2^r r!} \hbar^r P^r(f, g)$ , and for  $r = 1, 3$  they are  $\hbar P(f, g)$  and  $\frac{\hbar^3}{24} P^3(f, g)$ . The four primary cases (a)–(d) and a sweep of 14641 monomial pairs through order  $\hbar^1$  are exact-arithmetic verification checks for this coefficient formula. The external finite- $N$  radial-parts input remains a separate hypothesis.  $\square$

**THEOREM 5.223** (*Reduced open-line Wick theorem*). In the reduced open first-order line model

$$S_o = \int_{\mathbb{R}} \text{Tr}(\phi_1 d\phi_2),$$

on two brane-time points  $(t, t') \in \mathbb{R}_{\text{brane}}^2$  (with  $s$  reserved for the transverse  $\mathbb{R}_s$  coordinate of  $X$ ), choose the zero-mode-removed midpoint propagator

$$G(t, t') = \frac{1}{2} \text{sgn}(t - t'), \quad \langle \phi_1(t) \phi_2(t') \rangle = \hbar G(t, t'), \quad \langle \phi_2(t) \phi_1(t') \rangle = -\hbar G(t, t').$$



For ordered insertions  $t > t'$ , let  $W_r^>(f, g)$  be the contribution of the graph with exactly  $r$  cross-contractions between the two Weyl-ordered boundary vertices. Let  $W_r^<(f, g)$  be the reversed ordering contribution. Then

$$W_r^>(f, g) = \frac{1}{r!} \left( \frac{\hbar}{2} \right)^r P^r(f, g), \quad W_r^<(f, g) = \frac{1}{r!} \left( -\frac{\hbar}{2} \right)^r P^r(f, g).$$

Consequently the reduced open-line commutator is

$$[f, g]_{\text{Wick}} = \sum_{\substack{r \geq 1 \\ r \text{ odd}}} \frac{2}{2^r r!} \hbar^r P^r(f, g),$$

and all even orders vanish. For monomials  $f = z_1^k z_2^l$ ,  $g = z_1^m z_2^n$ ,

$$P^r(f, g) = \sum_{a=0}^r (-1)^a \binom{r}{a} (k)_{r-a} (l)_a (m)_a (n)_{r-a} z_1^{k+m-r} z_2^{l+n-r}.$$

In particular the one-edge and three-edge commutator contributions are

$$\hbar P(f, g), \quad \frac{\hbar^3}{24} P^3(f, g).$$

*Proof.* The midpoint value  $G(+, -) = 1/2$  assigns one copy of

$$P = \partial_{z_1} \otimes \partial_{z_2} - \partial_{z_2} \otimes \partial_{z_1}$$

to each cross-contraction, with weight  $\hbar/2$ . With Weyl vertices self-contractions vanish. The graph with  $r$  identical cross-edges has automorphism group  $S_r$ , so the ordered contribution is

$$W_r^>(f, g) = \frac{1}{r!} \left( \frac{\hbar}{2} \right)^r P^r(f, g).$$

Reversing the order changes the midpoint propagator from  $1/2$  to  $-1/2$ , giving the displayed formula for  $W_r^<$ . Hence

$$W_r^>(f, g) - W_r^<(f, g) = \frac{1 - (-1)^r}{2^r r!} \hbar^r P^r(f, g),$$

which is zero for even  $r$  and  $2\hbar^r P^r(f, g)/(2^r r!)$  for odd  $r$ .

The monomial formula is the expansion of  $P^r = (\partial_{z_1} \otimes \partial_{z_2} - \partial_{z_2} \otimes \partial_{z_1})^r$  by the binomial theorem: the summand with  $a$  copies of  $-\partial_{z_2} \otimes \partial_{z_1}$  contributes the sign  $(-1)^a$ , the binomial coefficient  $\binom{r}{a}$ , and the four falling factorials shown above. The values at  $r = 1$  and  $r = 3$  give  $\hbar P(f, g)$  and  $\hbar^3 P^3(f, g)/24$ .  $\square$

*Remark 5.224.* Theorem 5.223 is a computation in the reduced open-line Weyl model. The Costello closed-open graph theorem still requires an elliptic parent or dimensional reduction, a brane-defect condition, finite-scale gauge fixing, the induced bulk-to-defect propagator, orientations and measures on configuration spaces, counterterms, anomaly cancellation, and compatibility with the quantum Hamiltonian reduction.

### 5.7.5 THIRD-ORDER TARGET FOR COSTELLO'S GRAPH PROBLEM

The order- $\hbar^1$  Moyal coefficient  $P$  is fixed by the linear-Hamiltonian Heisenberg commutator and gives the target  $\hbar(kn - lm)z_1^{k+m-1}z_2^{l+n-1}$ . At order  $\hbar^3$  the Moyal target is  $P^3/24$ ; a Costello-graph realisation of the same coefficient would supply a separate trivalent boundary contraction diagram. Whether the two answers match is the content of this subsection.

For monomials  $f = z_1^k z_2^l$  and  $g = z_1^m z_2^n$ , the third-order Moyal coefficient from Proposition 5.215 is

$$P^3(f, g) = \sum_{a=0}^3 (-1)^a \binom{3}{a} (k)_{3-a} (l)_a (m)_a (n)_{3-a} z_1^{k+m-3} z_2^{l+n-3},$$

where  $(r)_s = r(r-1) \cdots (r-s+1)$  and  $(r)_0 = 1$ . Therefore

$$[f, g]_* = \hbar(kn - lm)z_1^{k+m-1}z_2^{l+n-1} + \frac{\hbar^3}{24}P^3(f, g) + O(\hbar^5).$$

The exact-arithmetic verification checks this first-order coefficient, the vanishing of the second-order coefficient in the raw commutator, the displayed third-order formula, and the star-commutator normalization on every monomial pair with exponents between 0 and 10 and commutator orders through 11.

This algebraic computation is an observable-product target, distinct from an effective action computation. In Costello's formalism the connected effective action and the product/OPE of quantum observables are related but distinct. The formal holomorphic singular coefficient of the product is the  $K_{\Delta}^{(2)}$ -kernel of Theorem 1.15; the finite-scale calculation below controls the brane-defect observable product that quantizes it. For two boundary observables  $F, G$ , the finite-scale free-field model has the Wick product

$$F \star_L G = m \circ \exp(\hbar \partial_{P_{\partial,L}})(F \otimes G),$$

where  $P_{\partial,L}$  is the boundary or brane-defect propagator induced by the chosen kernel. The connected single-trace part is the primitive trace component extracted from this product. In this convention the order in  $\hbar$  counts boundary cross-contractions; the loop order of the connected effective interaction is a different grading.

Thus there are two equivalent normalizations to keep separate. In Costello's observable-product notation one asks for the boundary contraction kernel


$$P_{\partial,L} \rightsquigarrow \frac{1}{2}P$$

on degree-zero Hamiltonians, so that

$$m \circ \exp(\hbar \partial_{P_{\partial,L}}) \rightsquigarrow m \circ \exp\left(\frac{\hbar}{2}P\right),$$

the Weyl/Moyal product used above. The ordered product has labeled inputs  $(f, g)$ ; its  $r$  identical cross-contractions contribute the denominator  $r!$ . The ordered product uses exactly this denominator. The reverse order is the separate product  $g * f$ ; their difference forms the commutator.


$\Gamma_{3a}$ : three propagators, common orientation



$$\text{Aut}(\Gamma_{3a}) = S_3, \quad |\text{Aut}(\Gamma_{3a})| = 3! = 6, \quad \frac{1}{|\text{Aut}(\Gamma_{3a})|} = \frac{1}{6}$$

orientations  $a = 0, 3$ : contributions  $(k)_3(n)_3$  and  $-(l)_3(m)_3$

$\Gamma_{3b}$ : two propagators forward, one reversed



$$\text{Aut}(\Gamma_{3b}) = S_2, \quad |\text{Aut}(\Gamma_{3b})| = 2! = 2, \quad \frac{1}{|\text{Aut}(\Gamma_{3b})|} = \frac{1}{2}$$

orientations  $a = 1, 2$ : contributions  $-3(k)_2 l m(n)_2$  and  $+3k(l)_2(m)_2 n$

FIGURE 0.2: The third-order graphs  $\Gamma_{3a}$  (top) and  $\Gamma_{3b}$  (bottom) for the order- $\hbar^3$  boundary contractions whose tensor target is  $P^3(f, g)$ . Each cross-contraction contributes one copy of the Poisson bivector  $P_\Delta$ ; reversing the orientation of a propagator contributes a relative sign. The orientation multiplicities are the binomial coefficients  $\binom{3}{a}$  of Proposition 5.215;  $\Gamma_{3a}$  realises  $a \in \{0, 3\}$  with automorphism group  $S_3$  permuting the three identical propagators (symmetry factor  $1/6$ ), and  $\Gamma_{3b}$  realises  $a \in \{1, 2\}$  with automorphism group  $S_2$  permuting the two same-orientation propagators (symmetry factor  $1/2$ ). These are two descriptions of the same count: either use the unexpanded aggregate  $P$ -edge count with automorphism  $S_3$ , or expand into orientation classes with residual automorphism  $S_a \times S_{3-a}$ ; the equality is  $\binom{3}{a}/3! = 1/(a!(3-a)!)$ : the unexpanded  $S_3$  denominator and the residual  $S_a \times S_{3-a}$  denominator are alternative descriptions of the same total automorphism count. Summing all four orientation classes,  $P^3(f, g) = \sum_{a=0}^3 (-1)^a \binom{3}{a} (k)_{3-a} (l)_a (m)_a (n)_{3-a} z_1^{k+m-3} z_2^{l+n-3}$ . The third-order raw commutator weight after antisymmetrising is  $\hbar^3/24$ . The reduced open-line weights are computed in Theorem 5.223; the Costello counterterm and anomaly checks remain the analytic obligations of Problem 5.227.

The graphs  $\Gamma_{3a}$  and  $\Gamma_{3b}$  of Figure 0.2 record the displayed order- $\hbar^3$  boundary contraction types whose tensor target is  $P^3(f, g)$ . Each cross-contraction contributes one copy of the constant Poisson bivector  $P$ . Their reduced open-line weights and automorphism factors are computed above. Their realization as Costello brane-defect graphs still requires the sector specialization, propagator, counterterm, and anomaly checks in the problem below.

**THEOREM 5.225** (*Costello first/third normalization recognition criterion*). Assume the Hamiltonian specialization data of Problem 5.212 have been supplied for a Costello-type closed-open BV theory, and assume that the induced brane-defect observable product restricts on degree-zero Hamiltonians to

the midpoint kernel  $P/2$  of Theorem 5.223. Assume further that at order  $\hbar^3$  every connected local term outside the three-parallel-edge cross-contraction has been shown to vanish, to be counterterm-exact, or to land in the discarded scalar trace. Then the first- and third-order Costello boundary commutator coefficients are determined by the Weyl/Moyal coefficients:

- (i) the connected one-propagator brane-defect ribbon graph carries coefficient  $\hbar(kn - lm)z_1^{k+m-1}z_2^{l+n-1}$  for monomial inputs  $f = z_1^k z_2^l$ ,  $g = z_1^m z_2^n$ ;
- (ii) the surviving three-propagator contribution carries coefficient  $\frac{\hbar^3}{24}P^3(f, g)$ ;
- (iii) any finite counterterm ambiguity that changes the first or third coefficient is fixed by matching the Moyal coefficients 1 and  $1/24$  at these orders. All-order Costello uniqueness lies outside this two-order test.

Equivalently, the Weyl/Moyal coefficient test for a Costello specialization at these orders holds precisely when the specialization supplies the above data and the normalization defects

$$\mathfrak{n}_1 = P_{\partial, L}|_{\text{Ham}^\circ} - \frac{1}{2}P,$$

$$\mathfrak{n}_3 = \left[ \text{ObsProd}_{\text{Costello}}^{(3)}(f, g) - \frac{\hbar^3}{24}P^3(f, g) \right]_{\text{Ham}^\circ, \text{conn}, \text{red}}$$

vanish, together with the scalar-contact, reduced-kernel, and  $\theta_3$  graph-row obstructions. The obstruction tuple is

$$\mathfrak{D}_{\text{Cost}}^{\mathfrak{h}_3} = (\mathfrak{o}_{\text{Cas}}, \hbar N[\bar{c}] \text{ or } o_{\sigma, w}^{(r)}, [\text{Ob}_{w, \partial, \hbar}^{\text{red}}], \mathfrak{D}_{\theta_3}, \mathfrak{n}_1, \mathfrak{n}_3).$$

The reduced datum gives Theorem 5.223 and the reduced open-line Wick normalization. Costello heat-kernel graph realization requires the full specialization data.

*Proof.* Under the stated hypotheses, the degree-zero boundary observable product is the reduced open-line Weyl product of Proposition 5.222. Thus each boundary cross-contraction contributes the operator  $P$  with weight  $\hbar/2$ . Here the  $S_3$  count treats the three edges as copies of the aggregate bivector  $P = A - B$ . After expanding  $P^3$ , a mixed orientation class has residual automorphism  $S_a \times S_{3-a}$  and binomial coefficient  $\binom{3}{a}$ . These are equivalent normalizations:  $\binom{3}{a}/3! = 1/(a!(3-a)!)$ ; the aggregate  $S_3$  denominator and the residual orientation-class denominator are alternative descriptions of the same total automorphism count.

Item (i): one edge, trivial edge automorphism group, weight  $\hbar P/2$  for the ordered product (Theorem 5.223); antisymmetrisation gives

$$\hbar P(f, g) = \hbar(kn - lm)z_1^{k+m-1}z_2^{l+n-1}$$

on monomials.

Item (ii): for three parallel edges between the two labelled boundary observables, the edge automorphism group is  $S_3$ . The ordered weight is

$$\hbar^3 \frac{1}{3!} \left( \frac{1}{2} \right)^3 P^3(f, g) = \frac{\hbar^3}{48} P^3(f, g),$$

and antisymmetrisation doubles it to  $\hbar^3 P^3(f, g)/24$ . The hypotheses exclude the remaining connected local terms at the same order.

Item (iii): a finite local counterterm that changes the one- or three-edge boundary product changes the coefficient of  $P$  or  $P^3$ . The linear-Hamiltonian Heisenberg commutator fixes the coefficient of  $P$  to 1, and Proposition 5.215 fixes the coefficient of  $P^3$  to  $1/24$  in the raw commutator. Thus matching these two coefficients removes precisely the counterterm freedom visible at these orders. The displayed defect tuple is a restatement of the hypotheses needed to promote this reduced computation to a Costello graph realization: the first defect fixes the brane-restricted propagator, the third-order defect fixes the connected reduced product, and the remaining entries are the scalar, reduced-kernel, and  $\theta_3$  graph-row conditions developed in Problem 5.219.  $\square$

*Remark 5.226.* In the reduced open-line model the Weyl parameter  $\hbar$  counts boundary cross-contractions. If a Costello specialization restricts to this free boundary kernel, then its observable-product expansion must be read with this edge-counting convention on degree-zero Hamiltonians. The specialization data of Problem 5.212 upgrades the formal Weyl/Moyal normalization test above to a Costello graph statement.

*Problem 5.227 (Costello first/third graph-normalization obstruction datum).* Work inside the observable product supplied by Statement 5.206, after solving the Hamiltonian specialization data of Problem 5.212. The first graph fixes the normalization; the third graph tests whether the Costello specialization really lands on the Weyl/Moyal product, thereby fixing the deformation-quantization convention. The analytic problem is reduced to the following checks:

- (i) identify the connected one-propagator brane-defect ribbon graph whose tensor contraction is  $P(f, g)$ , and prove that its antisymmetrised boundary product gives the raw coefficient  $\hbar P(f, g)$ ;
- (ii) fix the sign, automorphism factor, boundary orientation, and trace normalization in this first-order graph so that the coefficient is  $\hbar(kn - lm)z_1^{k+m-1}z_2^{l+n-1}$  on monomials;
- (iii) identify the connected three-propagator brane-defect ribbon graphs whose tensor contraction is  $P^3(f, g)$ ;
- (iv) prove that every other connected order- $\hbar^3$  contribution in the Hamiltonian sector vanishes, is counterterm-exact, or lands in the discarded scalar trace;
- (v) for every term declared counterterm-exact in the preceding item, exhibit compatible local counterterm representatives  $C_{\Gamma, w}^{w_o}(\epsilon) \in \mathcal{O}_{\text{loc}}(\mathcal{E}_w^{w_o})$ , prove Costello locality and RG compatibility, and verify that their restriction to degree-zero Hamiltonians preserves the fixed one-edge coefficient except by the allowed Moyal normalization choice;
- (vi) compute the graph weight and automorphism factor after the first-order normalization and show that the coefficient is  $\frac{1}{24}$ ;
- (vii) check that connected large- $N$  extraction preserves the factor of  $N$ , sign, and scalar moment-map normalization;
- (viii) verify that the same observable product, counterterm subtraction, and connected extraction commute with the principal-part current extension of Theorem 5.185 on the reduced target. If the claim is made before de Rham-current contraction, identify the additional unreduced homotopies of Remark 5.189. In particular the degree-zero Moyal product must act on the cotangent current summand through the principal-part coadjoint action; polynomial one- $\psi$  descendants carry the PBW special-fibre action.

The target normalization is the pair of algebraic coefficients

$$\hbar P(f, g), \quad \frac{\hbar^3}{24} P^3(f, g)$$

in the Weyl/Moyal commutator. Theorem 5.225 records the implication once the Costello specialization supplies these checks.

*Remark 5.228.* In characteristic zero,  $GL_N$  is linearly reductive on rational representations. Therefore taking  $GL_N$ -invariants is exact, and

$$H((\mathcal{R}_N)^{GL_N}, Q) \cong H(\mathcal{R}_N, Q)^{GL_N}.$$

This justifies the degree-zero rational invariant calculation used above. The full Lie-algebra Chevalley–Eilenberg BRST cohomology of  $\mathfrak{gl}_N$  is a separate stack-level calculation. A stack-level presentation remembers stabilizers, including the trivially acting scalar  $B\mathbb{G}_m$ ; those ghost and stabilizer classes are outside the Hamiltonian trace sector computed here. Linearly reductive exactness identifies derived algebraic invariants of functions with ordinary invariants in this setting.

The dictionary  $c_f \mapsto \theta_f, u_f \mapsto J(f)$  is the Koszul resolution of the brane-side derived  $E_1$ -Hochschild  $HH^\bullet(\mathcal{A}_{\partial, N}^{\text{cl}}, \mathcal{A}_{\partial, N}^{\text{cl}})$  at the formal-Darboux stalk of the Dirac brane vacuum  $b_p$  (Theorem 5.6); the Mixed Holomorphic–Topological Deligne conjecture identifies this at the formal-Darboux stalk with the bulk  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_{b_p}$  (the chiral  $E_2$  factorization algebra restricted to the formal neighbourhood of  $b_p$ ). Scalar reduction kills the constant Hamiltonian on the closed side and retains a  $U(1)$  centre-of-mass anomaly on the open side: a single Lie 2-cocycle  $[\bar{c}] \in H_{\text{Lie}}^2(\bar{A}, \mathbb{C})$  classified in the section below.

## 6 THE CAPELLI SCALAR

The Capelli central charge  $\hbar N[\bar{c}]$  is the projective curvature  $\Omega_{\text{central}}|_{J(f)}$  of the modular line bundle at the trace generator, realised by the Capelli central extension  $\mathfrak{o} \rightarrow \mathbb{C} \cdot K \rightarrow \widehat{\mathfrak{g}}_{\text{cent}} \rightarrow \mathfrak{g} \rightarrow \mathfrak{o}$  of (15); the operational test is  $\{\text{Tr } \phi_1, \text{Tr } \phi_2\} = N$  (Remark 5.9, probe (2)).

### 6.1 The $U(1)$ centre-of-mass cocycle

A constant Hamiltonian  $c \in \mathbb{C} \subset \mathbb{C}[z_1, z_2]$  acts trivially on the polydisk: its Hamiltonian vector field vanishes, so it sits in the kernel of the canonical-transformation reduction (3). The same constant pulled back along the substitution  $z_i \mapsto \phi_i$  becomes  $c \cdot I$ , and the matrix Poisson partner of the symplectic potential is

$$\{\text{Tr } \phi_1, \text{Tr } \phi_2\} = \text{Tr}(I \cdot I) = N,$$

the brane centre-of-mass charge. Closed-side: invisible. Open-side: visible, equal to  $N$ . The asymmetry is a single Lie cohomology class  $[\bar{c}] \in H_{\text{Lie}}^2(\bar{A}, \mathbb{C})$ , realised both as an abelian-extension obstruction on  $\bar{A}$  and as the  $U(1)$  trace centre of  $\mathfrak{gl}_N$ , and it survives Moyal quantisation as the order- $\hbar$  Capelli shift. The cocycle has three concrete faces — Lie 2-cocycle on  $\bar{A}$ , Poisson bracket of trace coordinates, Moyal–Weyl commutator  $[\Phi_{\hbar}(z_1), \Phi_{\hbar}(z_2)]_{\star} = \hbar N \cdot I$  in  $\mathcal{M}_N(\mathbb{C})$  — packaging the four equivalent sightings of Theorem E.13 (the abstract central extension and the explicit Lie central extension reduce to one face on  $\bar{A}$ ).

*The closed–open mismatch cocycle.* Write  $\mathfrak{h}_{\text{poly}} = \mathbb{C}[z_1, z_2]$  for the unreduced polynomial Hamiltonian Lie algebra, constants kept; the bracket  $\{f, g\} = \partial_{z_1} f \partial_{z_2} g - \partial_{z_2} f \partial_{z_1} g$  again takes values in  $\mathfrak{h}_{\text{poly}}$ . Define

$$\omega: \mathfrak{h}_{\text{poly}} \otimes \mathfrak{h}_{\text{poly}} \longrightarrow \mathbb{C}, \quad \omega(f, g) = [\{f, g\}]_0,$$

the constant Taylor coefficient of the bracket. On monomials,

$$\omega(z_1^k z_2^l, z_1^m z_2^n) = \begin{cases} kn - lm, & (k+m, l+n) = (1, 1), \\ 0, & \text{otherwise,} \end{cases}$$

so  $\omega$  is concentrated in the unique bidegree where the bracket multidegree  $(k+m-1, l+n-1)$  lands at  $(0, 0)$ : the constant-Taylor direction the canonical-transformation reduction (3) discards. This is the direction on which the closed and open sides disagree.

*Example 6.1 (Linear Hamiltonians and the brane centre of mass).* The two linear Hamiltonians  $f_1 = z_1$  and  $f_2 = z_2$  measure

$$J(z_1) = \overline{\text{Tr } \phi_1}, \quad J(z_2) = \overline{\text{Tr } \phi_2},$$

the brane centre of mass in the two transverse directions. On the unreduced phase space the matrix Poisson bracket of these two observables is

$$\{\text{Tr } \phi_1, \text{Tr } \phi_2\} = \text{Tr}(I \cdot I) = N,$$

so the brane centre-of-mass pair is a Heisenberg pair with charge  $N$ : the open-side image of the unique cocycle bidegree  $\omega(z_1, z_2) = 1$ . Closed-side,  $\{z_1, z_2\} = 1 \equiv 0$  in  $\tilde{A}$  by quotient. Moyal quantisation returns the same scalar at order  $\hbar$ : on the matrix-brane realization  $\Phi_{\hbar}(z_i) = \phi_i$  in  $\mathcal{M}_N(\mathbb{C})$  the commutator

$$[\Phi_{\hbar}(z_1), \Phi_{\hbar}(z_2)]_{\star} = \hbar N \cdot I_N$$

truncates exactly at first order because the Moyal expansion needs at least one second derivative on each tensor factor and the linear Hamiltonian has none (Theorem 6.6). The three faces of the anomalous scalar — Lie 2-cocycle on  $\tilde{A}$ , Poisson bracket of trace coordinates, Moyal–Weyl commutator — realise the same class. The trace- $U(1)$  central charge and the modular projective-curvature  $\Omega_{\text{central}}|_{J(f)}$  are equivalent interpretations of the multiplier  $N$  (Theorem 6.8).

**LEMMA 6.2 (Closed–open mismatch lemma).** The form  $\omega$  is a Lie 2-cocycle on  $\mathfrak{h}_{\text{poly}}$ : exact on the unreduced algebra and obstructed after the constant-quotient. Equivalently:

- on  $\mathfrak{h}_{\text{poly}}$ ,  $\omega = \delta\eta$  with  $\eta(f) = -[f]_0$ ;
- on  $\tilde{A} = \mathfrak{h}_{\text{poly}}/\mathbb{C} \cdot 1$ , the restricted cocycle represents a nonzero class  $[\tilde{c}] \in H_{\text{Lie}}^2(\tilde{A}, \mathbb{C})$ .

The class  $[\tilde{c}]$  is the closed-side obstruction blocking the descent of  $\eta$  past the constant line. Up to the multiplier  $\text{Tr } I = N$  it is the open-side scalar  $\{\text{Tr } \phi_1, \text{Tr } \phi_2\} = N$  pulled back through the boundary evaluation map of Theorem 6.4: the same scalar mismatch, read on each side.

*Proof. Cocycle identity.* Project each term of the Jacobi identity on  $\mathfrak{h}_{\text{poly}}$  to its constant Taylor coefficient:

$$\omega(f, \{g, h\}) + \omega(g, \{h, f\}) + \omega(h, \{f, g\}) = 0.$$

Antisymmetry of  $\omega$  is inherited from antisymmetry of the bracket. Hence  $\omega \in Z_{\text{Lie}}^2(\mathfrak{h}_{\text{poly}}, \mathbb{C})$ .

*Exactness on the unreduced algebra.* Set  $\eta(f) = -[f]_0$ . Then for all  $f, g \in \mathfrak{h}_{\text{poly}}$ ,  $\delta\eta(f, g) = -\eta(\{f, g\}) = [\{f, g\}]_0 = \omega(f, g)$ .

*Obstruction after constant-quotient.* The cocycle  $\omega$  descends to  $\bar{A}$  because it already vanishes on the constant line. The primitive  $\eta$  has value  $\eta(1) = -1$  on the constant line. Suppose some  $\tilde{\eta}: \bar{A} \rightarrow \mathbb{C}$  primitive of  $[\bar{c}]$  existed; lift to  $\tilde{\eta}: \mathfrak{h}_{\text{poly}} \rightarrow \mathbb{C}$  by setting  $\tilde{\eta}(1) = 0$ . Then  $\tilde{\eta}(\{z_1, z_2\}) = \tilde{\eta}(1) = 0$ , while  $\omega(z_1, z_2) = 1$ , contradicting  $\delta\tilde{\eta} = \omega$ .

*Open-side image.* The closed cocycle on the unit bidegree is  $\omega(z_1, z_2) = 1$ ; the boundary evaluation  $\partial_{\text{bb}, N}^{\text{full}}$  of Theorem 6.4 multiplies by  $\text{Tr } I = N$  to give  $\{\text{Tr } \phi_1, \text{Tr } \phi_2\} = \text{Tr}(I \cdot I) = N$ . Same axis, two faces.  $\square$

The class  $[\bar{c}]$  is Chevalley–Eilenberg cohomology on the closed side and the  $\mathfrak{gl}_N$  trace centre on the open side.

**THEOREM 6.3** (*Closed-side face: the scalar abelian extension*). The class  $[\bar{c}] \in H_{\text{Lie}}^2(\bar{A}, \mathbb{C})$  of Lemma 6.2 is the obstruction class to splitting the abelian Lie extension

$$0 \longrightarrow \mathbb{C} \cdot 1 \longrightarrow \mathfrak{h}_{\text{poly}} \longrightarrow \bar{A} \longrightarrow 0.$$

Equivalently,  $\mathfrak{h}_{\text{poly}} \cong \bar{A}_{[\bar{c}]} = \bar{A} \oplus \mathbb{C}K$  as Lie algebras, with bracket

$$[(f, aK), (g, bK)] = (\{f, g\}_{\bar{A}}, \bar{c}(f, g)K),$$

and  $\mathbb{C} \cdot [\bar{c}] \subset H_{\text{Lie}}^2(\bar{A}, \mathbb{C})$  is the distinguished anomaly line attached to the constant-Hamiltonian central extension.

*Proof.* The constant Hamiltonian  $1 \in \mathfrak{h}_{\text{poly}}$  has zero Hamiltonian vector field, so  $\mathbb{C} \cdot 1$  is central; the displayed sequence is a central extension and its class is its standard Chevalley–Eilenberg obstruction. Computing on monomial generators,

$$[z_1^k z_2^l, z_1^m z_2^n]_{\mathfrak{h}_{\text{poly}}} = (kn - lm) z_1^{k+m-1} z_2^{l+n-1},$$

and projecting to the central line  $\mathbb{C} \cdot 1$  keeps the  $(0, 0)$ -Taylor coefficient. This is the cocycle  $\omega$ , hence the extension class is  $[\bar{c}]$ . Non-triviality is Lemma 6.2; the scalar-extension line  $\mathbb{C} \cdot [\bar{c}] \subset H_{\text{Lie}}^2(\bar{A}, \mathbb{C})$  is therefore nonzero, and splitting the corresponding single extension is the problem of lifting back to  $\mathfrak{h}_{\text{poly}}$ .  $\square$

**THEOREM 6.4** (*Open-side face: the  $U(1)$  trace centre at finite  $N$* ). At finite  $N$  the matrix Poisson bracket  $\{F, G\} = \text{Tr}(\partial_{\phi_1} F \partial_{\phi_2} G - \partial_{\phi_2} F \partial_{\phi_1} G)$  on  $\mathbb{C}[\mathfrak{gl}_N \oplus \mathfrak{gl}_N]^{GL_N}$  satisfies

$$\{\text{Tr } \phi_1, \text{Tr } \phi_2\} = N$$

in the unreduced ring; this scalar vanishes in the reduced quotient  $H_{\text{red}}^0 = H^0 / \mathbb{C} \cdot \text{Tr}(1)$ . Under the boundary evaluation map

$$\partial_{\text{bb}, N}^{\text{full}}: \mathbb{C}[z_1, z_2] \longrightarrow \mathbb{C}[\mathfrak{gl}_N^2]^{GL_N}, \quad f \longmapsto \text{Tr } f(\phi_1, \phi_2),$$

the closed cocycle  $\omega$  on  $\bar{A}$  multiplies by  $\text{Tr } I = N$  to give the open trace bracket on the central direction; the resulting central extension of the open trace algebra is therefore the image of  $[\bar{c}]$ . Since  $\mathbb{C} \cdot I \subset \mathfrak{gl}_N$  is the  $U(1)$  centre with character  $\text{Tr}$ , the brane realisation of  $[\bar{c}]$  is exactly the  $U(1)$  centre of  $\mathfrak{gl}_N$  with charge  $N$ .



*Proof.* On the linear generators  $\partial_{\phi_1} \text{Tr } \phi_1 = I$  and  $\partial_{\phi_2} \text{Tr } \phi_2 = I$ ; the cross derivatives vanish. The matrix Poisson bracket is therefore  $\text{Tr}(I \cdot I) = N$  in the unreduced ring and zero modulo  $\mathbb{C} \cdot \text{Tr}(\mathbf{1})$ . The boundary evaluation map sends  $\mathbf{1} \in \mathbb{C}[z_1, z_2]$  to  $\text{Tr} I = N$ , and on the cocycle bidegree  $(k + m, l + n) = (\mathbf{1}, \mathbf{1})$ ,

$$\partial_{\text{bb}, N}^{\text{full}}(\{z_1^k z_2^l, z_1^m z_2^n\}) = (kn - lm) \cdot N = N \cdot \omega(z_1^k z_2^l, z_1^m z_2^n),$$

which is the closed cocycle of Lemma 6.2 times the multiplier  $\text{Tr } I = N$ . The central extension class on the open side is therefore the image of  $[\bar{c}]$ , and the central generator  $I \in \mathfrak{gl}_N$  is detected by the trace character.  $\square$

*Remark 6.5 (Reduced quotient).* Theorems below using  $\mathfrak{h}$ ,  $\tilde{A}$ , or  $\mathcal{A}_\partial = H_{\text{red}}^\circ$  live *after* killing the scalar; the constant central line is removed on each side. The unreduced statement keeps  $\mathbb{C} \cdot \mathbf{1} \subset \mathfrak{h}_{\text{poly}}$  closed-side and  $\mathbb{C} \cdot \text{Tr}(\mathbf{1})$  open-side, and the gap is  $[\bar{c}]$ .

Quantisation tests trace-map compatibility. Two operations close in on the brane: Weyl/Moyal quantization  $\Phi_\hbar$  of the formal symplectic polydisk, and the matrix-trace map  $J$  of the quantized symbol on the Chan–Paton pair. They commute exactly when the Moyal commutator  $[\Phi_\hbar(z_1), \Phi_\hbar(z_2)]_\star$  and the trace commutator  $[\text{Tr } \phi_1^\hbar, \text{Tr } \phi_2^\hbar]$  agree. Linear Hamiltonians expose the first quantum obstruction. The Moyal expansion truncates because each higher-order term  $P^r$  requires  $r$  derivatives in each tensor factor and the second derivative of a linear function vanishes; the surviving leading term is the matrix-brane scalar  $[\Phi_\hbar(z_1), \Phi_\hbar(z_2)]_\star = \hbar N \cdot I_N$  in  $M_N(\mathbb{C})$ , exactly the order- $\hbar$  image of  $\omega(z_1, z_2) = \mathbf{1}$  on the unreduced algebra. The classical class  $[\bar{c}]$  and its order- $\hbar$  Capelli shift are one Lie 2-cocycle, read on the closed Lie side, on the unreduced trace algebra side, and on the Moyal/Weyl side; this is the quantum face of the brane trace-map test. Linear trace generators have only the order- $\hbar$  correction.

**THEOREM 6.6** (*Moyal survival of  $\bar{c}$ : the order- $\hbar$  Capelli shift*). Let  $\Phi = (\phi_1, \phi_2)$  be the classical brane field and  $\Phi_\hbar = (\phi_1^\hbar, \phi_2^\hbar)$  its Weyl quantisation at finite  $N$ , with the Moyal product  $[7] f * g = m \circ \exp(\frac{\hbar}{2} P)(f \otimes g)$ ,  $P = \partial_{z_1} \otimes \partial_{z_2} - \partial_{z_2} \otimes \partial_{z_1}$ . In the degree-zero quantum Hamiltonian reduction the normal-ordered comoment imposes the Capelli/canonical relation

$$Y_\hbar X_\hbar - X_\hbar Y_\hbar + \hbar N I \equiv 0, \quad X_\hbar = \phi_1^\hbar, Y_\hbar = \phi_2^\hbar,$$

modulo the quantum moment-map ideal. On the linear trace generators  $X = \text{Tr } \phi_1$ ,  $Y = \text{Tr } \phi_2$  this is the classical bracket  $\{\text{Tr } \phi_1, \text{Tr } \phi_2\} = N$  of Theorem 6.4 times  $\hbar$ ; higher-order Moyal corrections vanish. Equivalently, the Rees quantisation  $U_\hbar(\tilde{A}_{[\bar{c}]})$  of the closed-side central extension carries the cocycle  $[\bar{c}]$  as its leading-order PBW splitting obstruction, and the trace character  $K \mapsto NI$  on the central generator returns the scalar Capelli contact  $\hbar N$ .

*Proof. Linear-generator collapse of the Moyal expansion.* The Weyl convention gives  $[X_\hbar, Y_\hbar]_* = \hbar\{X, Y\} + O(\hbar^3)$  in general. On the linear trace generators  $X = \text{Tr } \phi_1$ ,  $Y = \text{Tr } \phi_2$ , every higher-order term  $P^r$  requires at least  $r$  derivatives in each tensor factor, but the second derivative of a linear function vanishes. The Moyal expansion truncates at first order, so

$$X_\hbar Y_\hbar - Y_\hbar X_\hbar = \hbar\{X, Y\} = \hbar N$$

exactly. Rearranging gives the displayed normal-ordered moment relation in the quantum Hamiltonian reduction. The order-one Capelli identity [26, 27] is the same relation read on the universal-enveloping side; the Razmyslov–Procesi description [17, 18] of the full trace-relation ideal at finite  $N$  is separate from this calculation.

*Central-extension matching.* The Rees algebra  $U_{\hbar}(\tilde{A}_{[\bar{c}]})$  has special fibre  $\widehat{\mathbf{S}}(\tilde{A}_{[\bar{c}]})$  at  $\hbar = 0$  and contains  $\tilde{A}_{[\bar{c}]}$  as a Lie subalgebra at  $\hbar = 1$ . PBW associated-graded carries the cocycle  $\omega$  as the primary obstruction to splitting the deformation:  $\omega$  is the leading Moyal commutator on the central direction. Evaluating the central generator  $K$  by the finite- $N$  trace character  $K \mapsto NI$  converts the order- $\hbar$  central term into the scalar Capelli contact  $\hbar N$ .

*Conclusion.* Classical  $[\bar{c}]$  and order- $\hbar$  quantum Capelli shift carry the same Lie 2-cocycle on  $\tilde{A}$ . They are the same datum, read at  $\hbar = 0$  and at first order in  $\hbar$ .  $\square$

*Remark 6.7 (Three faces, one class).* The mismatch class  $[\bar{c}] \in H_{\text{Lie}}^2(\tilde{A}, \mathbb{C})$  has three named faces:

- closed side, abelian-extension obstruction (Theorem 6.3);
- open side,  $U(1)$  trace centre with charge  $N$  (Theorem 6.4);
- Moyal side, order- $\hbar$  Capelli contact (Theorem 6.6).

Scalar reduction kills the empty-trace scalar  $\text{Tr}(1) = N$ ; the linear trace observables  $\text{Tr } \phi_1, \text{Tr } \phi_2$ , whose unreduced bracket is  $N$  and whose reduced bracket is zero, remain. The obstruction beyond this scalar sector is the unreduced factorisation centre of Problem 5.48 and the higher-bidegree QME obstruction of Problem 5.219:  $[\bar{c}]$  is the scalar component, those problems carry the higher-bidegree components.

**THEOREM 6.8** (*Capelli scalar = projective curvature of the modular line bundle*). Let  $J(f) = \text{Tr } f(\phi_1, \phi_2)$  be the trace generator of §4.2 and let  $\Omega_{\text{central}}$  denote the canonical curvature of the determinant line bundle on the moduli stack of the closed-string sector classical observables factorization algebra  $\mathcal{A}_{\text{bulk}}^{\text{cl}}$  at  $N$  Dirac branes on  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}_{\text{o,hol}}^2$ . Then

$$\hbar N [\bar{c}] = \Omega_{\text{central}}|_{J(f)} \quad \text{in } H_{\text{Lie}}^2(\tilde{A}, \mathbb{C})[[\hbar]],$$

where the left-hand side is the Lie 2-cocycle of Lemma 6.2 multiplied by the Chan–Paton scalar  $\hbar N$ , and the right-hand side is the value of the modular line curvature at the local trace coordinate. Equivalently, the central extension

$$0 \longrightarrow \mathbb{C} \longrightarrow \mathfrak{h}_{\hbar, N} \longrightarrow \tilde{A} \longrightarrow 0$$

carrying the cocycle  $\hbar N \omega$  lifts canonically to the square root of the determinant line of the trace representation  $\tilde{A} \rightarrow \mathfrak{gl}_N$ ; the class  $[\bar{c}]$  is the Atiyah class of this determinant line, in the sense of [28, §2.5].

*Proof.* The cocycle identity is the content of Theorem E.13; the four sightings of  $[\bar{c}]$  coincide under the canonical scalar projection, with the multiplier  $\hbar N$  supplied by the open Chan–Paton trace pairing. The right-hand side is identified as follows. The brane  $E_1$ -algebra  $\mathcal{A}_{\partial, N}^{\text{cl}} = C_{\text{CE}}^{\bullet}(\mathfrak{gl}_N, \text{Kosz}([\phi_1, \phi_2]))$  on the worldline  $\mathbb{R}_t$  has trace coordinate  $u_f \mapsto J(f)$  and CE coordinate  $c_f \mapsto \theta_f$  by §4.1; under this identification the Capelli normal-ordered shift  $Y_{\hbar} X_{\hbar} - X_{\hbar} Y_{\hbar} + \hbar N I \equiv 0$  (Theorem 6.6) is the formal-Darboux expression of the projective curvature of the determinant line of the trace representation  $\tilde{A} \rightarrow \mathfrak{gl}_N$ . The linear Hamiltonian generators  $z_1, z_2$  act on the trace representation with classical Capelli identity [26, 27]; its order- $\hbar$  deformation is the Tate central extension associated with the continuous endomorphism algebra of [28, §2.5]. Pulling the Atiyah class of  $\det^{1/2}$  back to  $H_{\text{Lie}}^2$  gives the cocycle  $\hbar N \omega$  with  $\omega(z_1, z_2) = 1$  by Corollary C.8; the class is therefore  $\hbar N [\bar{c}]$ , and the equality with  $\Omega_{\text{central}}|_{J(f)}$  is the Beilinson–Drinfeld central-extension identification at the trace generator.  $\square$

*Remark 6.9 (Determinant line and scheme dependence).* The central extension lifts canonically to the square root of the determinant line of the trace representation  $\tilde{A} \rightarrow \mathfrak{gl}_N$ : the Capelli shift  $\hbar N I$  is the

curvature of the canonical connection on  $\det^{1/2}$  at the linear trace generators, so  $[\bar{c}]$  is the Atiyah class of this determinant line in the sense of [28, §2.5]. The numerical coefficient  $\hbar N$  is scheme-dependent in the sense of the factorization-current convention table §B: the canonical scheme is the symmetric Weyl ordering  $z_1 z_2 \mapsto \frac{1}{2}(\phi_1 \phi_2 + \phi_2 \phi_1)$  with Moyal parameter  $\hbar$  and Chan–Paton trace  $\text{Tr } I = N$ ; changing scheme by an inner automorphism shifts  $[\bar{c}]$  by a coboundary and leaves the class invariant. Reordering by an outer automorphism (e.g. a triangular Capelli splitting versus normal ordering) shifts the trace coefficient by a central character but preserves the modular-line identification.

*BV-cohomology vanishing and the unreduced cotangent lift*

The BV operator  $\Delta = Q + \{I_o, -\}$  of degree +1 acts on  $\mathcal{O}_{\text{loc}}(\mathcal{E}_w)$  for the mixed geometry  $X = \mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}^2_{o,\text{hol}}$  with brane  $L = \mathbb{R}_{\text{brane}} \times \{o\} \times \{p\}$  and ordinary  $\mathfrak{gl}_N$  gauge data. An endpoint-admissible coefficient module for Problem 5.48 extends the already computed native holomorphic singular product of Theorem 1.15 to the unreduced Costello/QME endpoint; it exists if and only if the four-component class

$$\mathfrak{o}_{\text{BV},\text{end}} = (\mathfrak{o}_{\text{Cas}}, \mathfrak{o}_{\text{def}}, \mathfrak{o}_{\text{RG}}, \mathfrak{o}_{\text{QME}})$$

of Definition 6.16 vanishes (Proposition 6.17): the strict-pair Casimir cokernel  $\mathfrak{o}_{\text{Cas}}$  (5.21), the brane-line propagator extension obstruction  $\mathfrak{o}_{\text{def}}$  (8.17 (2)), the strict-window Mittag–Leffler preservation class  $\mathfrak{o}_{\text{RG}}$  (5.15), and the local-functional QME anomaly class  $\mathfrak{o}_{\text{QME}} \in H^1(\mathcal{O}_{\text{loc}}(\mathcal{E}_w), Q + \{I_o, -\})$  of Problem 5.72.

The BV grading is the cotangent-coordinate grading of E.1 together with the signs of C.1, C.8:  $|\Delta| = +1$ , the reduced boundary  $P_o$ -bracket has degree 0, the Schouten bracket has degree  $-1$ .

The closed BCOV vanishing of [13] and the open-closed extension of [14] differ from the present geometry along five independent axes: compact Hodge geometry against the formal jet  $\widehat{\mathbb{C}}^2_o$ ; pure holomorphic-volume kinematics  $\bar{\partial} + t \partial_{\Omega}$  against mixed  $D = d_{\mathbb{R}^2} + \bar{\partial}_{\mathbb{C}^2}$ ; empty bulk against the transverse line defect  $L$ ; the  $\mathfrak{gl}(N | N)$  supertrace cancellation  $\text{str}(I) = N - N = 0$  against the ordinary  $\mathfrak{gl}_N$  scalar contact  $\hbar N[\bar{c}]$  of E.12; the worldsheet  $t$ -Tate direction against the coefficient-system Tate direction  $\widehat{\mathfrak{h}} \widehat{\otimes} \widehat{\mathfrak{h}}_{\text{cont}}^V$  carrying  $\mathfrak{o}_{\text{Cas}}$ . Each axis names a distinct endpoint component of  $\mathfrak{o}_{\text{BV},\text{end}}$  under the cited hypotheses (Theorem 6.15); the data needed to close them is enumerated in Theorem 6.18. The chain-level primitive shadow 5.132 closes the classical indecomposable shadow using algebraic input, inside the admissible Tate envelope of 5.30.

*BV degree convention and Costello renormalization datum*

*Remark 6.10 (BV grading and  $\Delta$  for the brane-defect endpoint).* The BV grading is fixed by E.1:  $|\phi_1| = |\phi_2| = 0$ ,  $|\psi = A^*| = -1$ ,  $Q\psi = [\phi_1, \phi_2]$ ,  $|Q| = +1$ ; a Hamiltonian generator  $H_f$  has degree 0, a shifted cotangent generator  $\eta_\rho = \rho[1]$  is odd of BV degree  $-1$ , and the CE cotangent–coordinate grading is  $|c_f| = +1$ ,  $|u_\rho| = 0$ . In this convention the BV operator

$$\Delta = Q + \{I_o, -\}$$

acting on  $\mathcal{O}_{\text{loc}}(\mathcal{E}_w)$  has degree +1, and a quantum master equation *anomaly representative* is a degree-+1 cocycle in  $(\mathcal{O}_{\text{loc}}(\mathcal{E}_w), \Delta)$ . The Schouten/cotangent–coordinate antibracket has degree  $-1$ ; the reduced Hamiltonian boundary  $P_o$ -bracket has degree 0. The scalar Hamiltonian cocycle  $\omega(z_1, z_2) = 1$  of C.6 controls the central contribution to the symbol, and the local Poisson sign convention  $\{z_1, z_2\} = 1$  is the one of C.1.

*Remark 6.11 (Costello (2011) heat-kernel renormalization datum invoked).* The endpoint comparison invokes the renormalization framework of [12, Ch. 2 §2, Ch. 2 §6, Ch. 5 §6]: a parametrix  $P_{\epsilon,L} = \int_{\epsilon}^L ds K_s$  for a generalized Laplacian, the homotopy RG flow

$$W(P_{\epsilon,L}, I) = \hbar \log(\exp(\hbar \partial_{P_{\epsilon,L}}) \exp(I/\hbar)) = \sum_{\Gamma \text{ connected}} \frac{\hbar^{\text{loops}(\Gamma)}}{|\text{Aut}(\Gamma)|} W_{\Gamma}(P_{\epsilon,L}, I),$$

the scale-zero limit  $\lim_{\epsilon \rightarrow 0} W(P_{\epsilon,L}, I)$  constructed by counterterm subtraction, and the QME  $QI + \hbar \Delta_L I + \frac{1}{2}\{I, I\}_L = 0$  at scale  $L$ . In the present brane-defect geometry the propagator factorizes (when it exists) as  $P_{\epsilon,L}^w = P_{\epsilon,L}^{\text{base}} \widehat{\otimes} C_{\mathfrak{h} \oplus \mathfrak{h}^{\vee}[1], w}$  (8.17); the bulk Hadamard parametrix is taken from [12, Ch. 3] on  $\mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$  and the coefficient factor is the algebraic kernel of  $\text{Tr}$ . The components  $\mathfrak{o}_{\text{def}}$  and  $\mathfrak{o}_{\text{RG}}$  of  $\mathfrak{o}_{\text{BV}, \text{end}}$  measure the brane-line extension class of  $P_{\epsilon,L}^{\text{base}}$  and the strict-window preservation class of  $W(P_{\epsilon,L}, -)$  inside this convention.

*Costello–Li 2012 hypotheses, restated*

*Statement 6.12 (Costello–Li 2012 BCOV construction, restated).* Let  $X$  satisfy the compact Calabi-Yau holomorphic-volume hypotheses of [13], with volume form  $\Omega$ ,  $\dim_{\mathbb{C}} X = d$ . Let  $\text{PV}(X)[[t]] = \bigoplus_{i,j} \text{PV}^{i,j}(X)[[t]]$  be the polyvector-field complex with the Tate parameter  $t$  for the worldsheet  $S^1$  rotation, BV differential

$$Q_{\text{BCOV}} = \bar{\partial} + t \partial_{\Omega}, \quad |Q_{\text{BCOV}}| = +1,$$

with  $\partial_{\Omega} = \Omega^{-1} \circ \partial \circ \Omega$  the divergence operator, and the  $(-1)$ -shifted symplectic pairing

$$\langle \alpha, \beta \rangle = \int_X (\alpha \wedge \beta)|_{\text{PV}^{d,d}} \cdot \Omega^2.$$

The classical BCOV interaction  $I_{\text{BCOV}}$  is the cubic vertex of [13, §3]. The cited construction supplies the generalized classical BCOV complex, its local-functional obstruction complex, and the renormalization framework of [12] adapted to compact Calabi-Yau  $X$ . The 2015 open-closed sequel [14] extends the construction to flat  $\mathbb{C}^d$  ( $d$  odd) coupled to open gauge data  $\mathfrak{gl}(N | N)$ , with the  $\text{U}(1)$ -anomaly cancellation  $\text{str}(I) = N - N = 0$ . Each is used here only under its stated geometric and coefficient hypotheses.

*Hypothesis separation on  $\mathbb{R}^2 \times \mathbb{C}^2$  with brane  $L = \mathbb{R}_{\text{brane}} \times \{0\} \times \{p\}$  and  $\mathfrak{gl}_N$*

**PROPOSITION 6.13 (Costello–Li 2012/2015 hypothesis separation for the brane-defect endpoint).** Let  $\mathfrak{o}_{\text{BV}, \text{end}} = (\mathfrak{o}_{\text{Cas}}, \mathfrak{o}_{\text{def}}, \mathfrak{o}_{\text{RG}}, \mathfrak{o}_{\text{QME}})$  be the four-component endpoint obstruction class of 6.16 for the formal local geometry  $X = \mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}_{0, \text{hol}}^2$  with brane  $L = \mathbb{R}_{\text{brane}} \times \{0\} \times \{p\}$  and ordinary  $\mathfrak{gl}_N$  gauge data. The hypotheses of 6.12 differ from this local model along five independent axes. In each axis the corresponding endpoint obstruction is named at the level of a cocycle representative:

(S1) *Compact Hodge-theoretic geometry vs. formal jet.* The 2012 theorem requires compact Calabi-Yau  $X$  with Hodge-theoretic harmonic projection  $\Pi_{\text{harm}}: \Omega^{\circ, \bullet} \rightarrow \mathcal{H}^{\circ, \bullet}$ . On the formal jet  $\widehat{\mathbb{C}}_0^2$ , the corresponding datum is the one-dimensional cohomology  $H_{\text{Dol}}^*(\widehat{\mathbb{C}}_0^2) = \mathbb{C}$  in degree zero. Hence the 2012 BV-cohomology vanishing  $H_{\text{BV}, \text{BCOV}}^*(X) = 0$  above degree zero is a statement about the compact BCOV complex. The present  $(\mathcal{O}_{\text{loc}}(\mathcal{E}_w), \Delta)$  on  $X = \mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}_{0, \text{hol}}^2$  has its own cocycle representative of  $\mathfrak{o}_{\text{QME}}$ . The 2015 open-closed theorem replaces compact  $X$  by flat  $\mathbb{C}^d$  ( $d$  odd) but keeps the holomorphic-volume kinematics and the  $\mathfrak{gl}(N | N)$  anomaly cancellation; it is a different geometric and coefficient setting.

- (S2) *Pure holomorphic-volume vs. mixed topological–holomorphic kinematics.* The 2012 BV differential is  $Q_{\text{BCOV}} = \bar{\partial} + t \partial_{\Omega}$  on  $\text{PV}(X)[[t]]$ . The kinematic differential of the present model is  $D = d_{\mathbb{R}^2} + \bar{\partial}_{\mathbb{C}^2}$  on  $\Omega_{\mathbb{R}^2}^{\bullet}(I) \widehat{\otimes} \Omega_{\mathbb{C}^2}^{\circ, \bullet} \widehat{\otimes} (\mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^{\vee}[1])$ , the holomorphic  $E_2$  factorization algebra on  $\mathbb{C}^2$ . The two complexes have distinct generators and differentials, so transfer of a Costello–Li class on  $(\text{PV}(X)[[t]], Q_{\text{BCOV}})$  to a cocycle of  $\Delta$  on the mixed complex requires a separate kinematic comparison map. The protected sector of twisted M-theory [25] carries this mixed kinematic shape as physical motivation.
- (S3) *Brane and topological line defect.* The 2012 theorem concerns the closed sector; the propagator  $P_{\epsilon, L}^{2012}$  on  $\text{PV}(X)[[t]]$  is a closed-sector propagator. The 2015 theorem has an open-closed sector on flat  $\mathbb{C}^d$  with open gauge  $\mathfrak{gl}(N | N)$ , where the open boundary is the boundary of a worldsheet disk in  $\mathbb{C}^d$ . The present endpoint uses a transverse line  $L \subset \mathbb{R}_{\text{top}}^2 \times \mathbb{C}^2$  point-supported in the holomorphic direction. Extension of  $P_{\epsilon, L}^{\text{base}}$  across  $L$  with ordinary  $\mathfrak{gl}_N$  brackets is encoded by the obstruction  $\mathfrak{o}_{\text{def}}$  of 6.16.
- (S4) *Supertrace cancellation vs. ordinary  $\mathfrak{gl}_N$  scalar contact.* On  $\mathfrak{gl}(N | N)$  the [14] cancellation  $\text{str}(I) = N - N = 0$  kills the would-be  $\text{U}(1)$ -anomaly at every loop order. On ordinary  $\mathfrak{gl}_N$  the same closed exchange supplies the scalar contact term  $\hbar N \omega(f, g) \gamma_1$  of E.12, with cohomology class  $\hbar N[\bar{c}] \in H_{\text{Lie}}^2(\bar{A}, \mathbb{C})$ , equivalently the projective curvature of the modular line bundle  $\Omega_{\text{central}}$  evaluated at the trace generator  $J(f)$  of Chapter 6 (Theorem 6.8). Costello–Li graph signs leave this class as the local trace-line avatar of the cancelled supertrace anomaly; a graph-level comparison with the Costello–Li cancellation is additional data. Hence  $\mathfrak{o}_{\text{QME}}$  carries the residual  $\hbar N[\bar{c}]$  summand for ordinary  $\mathfrak{gl}_N$  at the  $\text{U}(1)$  centre-of-mass slice, and that summand is a separate endpoint cocycle in the stated Costello–Li framework.
- (S5) *Worldsheet  $t$ -Tate vs. coefficient-system Tate.* The 2012 Tate parameter  $t$  is a worldsheet  $S^1$  rotation completion of the polyvector field complex. The Tate direction required by Problem 5.48 is along the coefficient pair

$$\widehat{\mathfrak{h}} \widehat{\otimes} \widehat{\mathfrak{h}}_{\text{cont}}^{\vee}, \quad C_{\mathfrak{h}} = \sum_{d \geq 1} \sum_i e_{d,i} \otimes e_d^i.$$

By 5.211 the strict pair  $(\mathfrak{h}_{\Pi}, D_{\oplus})$  has the represented-kernel obstruction for simultaneous projection to every finite Casimir  $C_{\mathfrak{h}}^M$ : the dual side admits only finitely many degree components, and the compatible kernel has nonzero identity component in every  $H_d^{\vee}$ . Hence the cocycle representative  $\mathfrak{o}_{\text{Cas}} = [(C_{\mathfrak{h}}^M)_M]$  is nonzero in the cokernel. This coefficient direction is governed by the endpoint Tate completion; the Costello–Li  $t$ -Tate completion is the worldsheet completion.

Each axis identifies a distinct endpoint component of  $\mathfrak{o}_{\text{BV, end}}$ : (S1)+(S2) give the kinematic comparison problem for  $\mathfrak{o}_{\text{QME}}$ ; (S3) gives  $\mathfrak{o}_{\text{def}}$ ; (S4) names the residual  $\hbar N[\bar{c}]$  summand of  $\mathfrak{o}_{\text{QME}}$ ; (S5) names the cokernel class of  $\mathfrak{o}_{\text{Cas}}$ . The remaining component  $\mathfrak{o}_{\text{RG}}$  is the strict finite-window Mittag-Leffler preservation class for the recursion  $\mathcal{W}(P_{\epsilon, L}, I)$  of 6.11; this preservation is an endpoint coefficient condition in addition to the compact-target Costello–Li theorems.

*Proof.* Each axis is a direct comparison of the hypotheses of 6.12 (and its 2015 sequel) with the data of 5.6, the linear-heat / Casimir analysis (5.210, 5.211), the scalar contact theorem E.12, and the strict-window category 5.15.

(S1). Compactness enters [13] through the Hodge-theoretic harmonic projection used to construct the BV propagator and to verify the QME at the parametrix limit  $\epsilon \rightarrow 0, L \rightarrow \infty$ . On the formal jet  $\widehat{\mathbb{C}}^2_0$ , the corresponding datum is the one-dimensional Dolbeault cohomology  $H_{\text{Dol}}^*(\widehat{\mathbb{C}}^2_0)$ .

(S<sub>2</sub>). The 2012 BV differential is purely Dolbeault plus the divergence  $t \partial_{\Omega}$ ; the present  $D = d_{\mathbb{R}^2} + \tilde{\partial}_{\mathbb{C}^2}$  carries an additional de Rham factor on the topological line. A symmetric-monoidal comparison map  $(\mathrm{PV}(X)[[t]], Q_{\mathrm{BCOV}}) \rightarrow (\Omega_{\mathbb{R}^2}^{\bullet}(I) \widehat{\otimes} \Omega_{\mathbb{C}^2}^{\bullet} \widehat{\otimes} \cdots, D)$  is additional endpoint data.

(S<sub>3</sub>). The brane  $L$  is transverse to the topological factor and point-supported in the holomorphic factor; the open boundary in [14] is a worldsheet boundary on flat  $\mathbb{C}^d$ . The present endpoint requires the Wilson line on a transverse topological direction. The bulk Hadamard parametrix  $P_{\epsilon, L}^{\mathrm{base}}$  of [12, Ch. 3] on  $\mathbb{R}^2 \times \mathbb{C}_{\mathrm{hol}}^2$  supplies the bulk kernel; the brane-line trace is a further endpoint datum.

(S<sub>4</sub>). By E.12 the closed exchange between two Hamiltonian sources  $B_f, B_g$  on the brane line produces the universal scalar contact density  $\hbar N \omega(f, g) \gamma_1(t) a(t) b(t)$ . Its associated-graded scalar symbol is the Lie 2-cocycle  $\omega(f, g) \gamma_1$  on  $\tilde{A}$ , with  $\omega(z_1, z_2) = 1$  by C.8; this is a nonzero class in  $H_{\mathrm{Lie}}^2(\tilde{A}, \mathbb{C})$  by 5.6 part (f). On  $\mathfrak{gl}(N | N)$  the same scalar is multiplied by  $\mathrm{sdim}(\mathbb{C}^{N|N}) = N - N = 0$ , killing the representative. On ordinary  $\mathfrak{gl}_N$  it is  $\hbar N$ , and absorption of this term requires an endpoint counterterm.

(S<sub>5</sub>). The compatible Casimir family  $(C_{\mathfrak{h}}^M)_M$  projects nonzero into every  $H_d \otimes H_d^{\vee}$  for  $d \leq M$ , while the strict direct-sum dual  $D_{\oplus} = \bigoplus_d H_d^{\vee}$  has only finitely many nonzero degree components per element. Thus 5.211 identifies a nonzero cokernel class for the map

$$\mathfrak{h}_{\Pi} \widehat{\otimes} D_{\oplus} \longrightarrow (\mathfrak{h}_{\Pi} \widehat{\otimes} D_{\oplus})^{\wedge^{\bullet}}.$$

Equivalently the cocycle representative of  $\mathfrak{o}_{\mathrm{Cas}}$  is nonzero, and the worldsheet  $t$ -Tate completion of [13] commutes with this coefficient direction trivially.

Finally,  $\mathfrak{o}_{\mathrm{RG}}$  is the assertion that  $\mathcal{W}(P_{\epsilon, L}, -)$  sends the strict ML category  $\mathbf{MLTateChain}_{\geq 0}^{\mathrm{str}}$  of 5.15 into itself with surjective finite-window transition maps; this is a statement about the algebraic-coefficient direction of the propagator, while the Costello–Li theorems address the worldsheet direction.  $\square$

**COROLLARY 6.14** (*Costello–Li input and the unreduced coefficient endpoint*). The Costello–Li 2012 BCOV construction supplies a pure closed BCOV template; the 2015 sequel supplies the open-closed extension on flat  $\mathbb{C}^d$  with  $\mathfrak{gl}(N | N)$  gauge data. The present  $\mathbb{R}_{\mathrm{top}}^2 \times \widehat{\mathbb{C}}_{\mathrm{o, hol}}^2$  brane-defect endpoint is governed by the four cocycle classes in  $\mathfrak{o}_{\mathrm{BV, end}}$ . The unreduced Tate-coefficient cotangent lift therefore additionally requires the endpoint coefficient data of Definition 5.94. The classical primitive indecomposable shadow on the open observables is closed by the algebraic argument of 5.132; the admissible Tate envelope on which the obstruction tuple is computed is provided by the model-categorical envelope of 5.17. The unreduced cotangent factorization-centre morphism and the quantum one-antifield extension of part (e) of 5.6 require the weighted coefficient category of T1 and the mixed brane-defect locality/QME input of Problem 5.72 together with 8.17.

*Proof.* By Proposition 6.13, each Costello–Li hypothesis selects a distinct endpoint comparison problem for a cocycle representative in  $\mathfrak{o}_{\mathrm{BV, end}}$ . Vanishing of the tuple is therefore the additional endpoint datum specified in the corollary. The remaining clauses are the classical chain-level lift of 5.132, the model-categorical envelope of 5.17, and the weighted finite-scale coefficient kernel of T1; the endpoint data are exactly those listed in Definition 5.94.  $\square$

**THEOREM 6.15** (*Costello–Li endpoint coefficient criterion*). The Costello–Li 2012 BCOV construction [13] and its 2015 open-closed extension [14], in the forms restated in 6.12, supply an endpoint-admissible coefficient module of Definition 5.94 for the formal local brane-defect geometry only after three independent classical data are added:

(E<sub>2</sub>) coefficient Casimir kernel, (E<sub>3</sub>) mixed brane-defect transport, (E<sub>5</sub>) endpoint RG preservation,

with (E1) and (E4) the inherited finite-window and Mittag-Leffler envelope conditions. A quantum endpoint additionally requires vanishing of the QME component  $\mathfrak{o}_{\text{QME}}$  of Definition 6.16. Equivalently: the four cocycle representatives in  $\mathfrak{o}_{\text{BV},\text{end}}$  require independent endpoint primitives on this geometry.

*Proof.* An endpoint-admissible coefficient module requires a coefficient kernel  $C_* \in \widehat{\mathfrak{h}}_* \widehat{\otimes} D_*$  whose projection to every finite Hamiltonian window is the finite Casimir  $C_{\mathfrak{h}}^M$ , together with mixed brane-defect transport of the bracket, propagator, interaction, counterterms, and RG recursion (Definition 5.94). By Proposition 6.13, each axis (S1)–(S5) names a distinct cocycle in  $\mathfrak{o}_{\text{BV},\text{end}}$ : (S5) gives  $\mathfrak{o}_{\text{Cas}}$ , (S3) gives  $\mathfrak{o}_{\text{def}}$ , the algebraic-coefficient direction gives  $\mathfrak{o}_{\text{RG}}$ , and (S1)+(S2)+(S4) give  $\mathfrak{o}_{\text{QME}}$ . Hence (E2), (E3), (E5), and the QME vanishing are the additional endpoint data.  $\square$

*Definition 6.16 (BV endpoint obstruction tuple, with explicit cocycle representatives).* For the formal mixed local geometry  $X = \mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}_{\text{o,hol}}^2$  with brane  $L = \mathbb{R}_{\text{brane}} \times \{\mathfrak{o}\} \times \{p\}$  and ordinary  $\mathfrak{gl}_N$  gauge data, the BV endpoint obstruction tuple is

$$\mathfrak{o}_{\text{BV},\text{end}} = (\mathfrak{o}_{\text{Cas}}, \mathfrak{o}_{\text{def}}, \mathfrak{o}_{\text{RG}}, \mathfrak{o}_{\text{QME}}).$$

Each component is named with its ambient cohomology, BV degree, and explicit cocycle representative:

(O1) *Casimir cokernel class*  $\mathfrak{o}_{\text{Cas}}$  — in  $\text{coker}(\widehat{\mathfrak{h}}_{\Pi} \widehat{\otimes} D_{\oplus} \rightarrow \varprojlim_M (\mathfrak{h} \otimes \mathfrak{h}^{\vee})_{\leq M})$ , degree 0; cocycle representative the compatible family

$$\mathfrak{o}_{\text{Cas}} = [(C_{\mathfrak{h}}^M)_{M \geq 1}], \quad C_{\mathfrak{h}}^M = \sum_{d \leq M} \sum_i e_{d,i} \otimes e_d^i,$$

with transition maps the standard truncations  $C_{\mathfrak{h}}^{M+1} \mapsto C_{\mathfrak{h}}^M$ ; nonzero in the strict pair by 5.211.

(O2) *Brane-defect transport class*  $\mathfrak{o}_{\text{def}}$  — in  $H^0(\text{Hom}_{\text{loc}}^{\bullet}(P_{\epsilon,L}^{\text{base}}, P_{\epsilon,L}^{\text{brane}}), \delta_L)$ , degree 0; cocycle representative the brane-line restriction datum

$$\mathfrak{o}_{\text{def}} = [\delta_L^* P_{\epsilon,L}^{\text{base}} - P_{\epsilon,L}^{\text{brane}}]$$

together with the bracket-transport obstruction for  $\{\cdot, \cdot\}_{\text{open}}$  of C.9 across  $L$  with ordinary  $\mathfrak{gl}_N$  brackets, equivalently the Hadamard locality class of 8.17 item (2).

(O3) *RG-window preservation class*  $\mathfrak{o}_{\text{RG}}$  — in the Mittag-Leffler  $\varprojlim^1 H^0$  of the strict-window functional complex  $\mathbf{MLTateChain}_{\geq 0}^{\text{str}}$  of 5.15, degree 0; cocycle representative the compatible family of recursion-induced transition defects

$$\mathfrak{o}_{\text{RG}} = [T^{M,M-1} \circ W(P_{\epsilon,L}, -)|_{\leq M} - W(P_{\epsilon,L}, -)|_{\leq M-1} \circ T^{M,M-1}] \in \varprojlim_M^1 H^0,$$

where  $T^{M,M-1}$  is the strict ML truncation of 5.15 and  $W(P_{\epsilon,L}, -)$  is the Costello recursion of 6.11.

(O4) *QME anomaly class*  $\mathfrak{o}_{\text{QME}}$  — in the local-functional cohomology

$$H^1(\mathcal{O}_{\text{loc}}(\mathcal{E}_w), \Delta), \quad \Delta = \mathcal{Q} + \{I_{\mathfrak{o}}, -\}, \quad |\Delta| = +1,$$

degree +1; cocycle representative the local functional

$$\mathfrak{o}_{\text{QME}} = [QI_{\mathfrak{o}} + \hbar \Delta_L I_{\mathfrak{o}} + \frac{1}{2} \{I_{\mathfrak{o}}, I_{\mathfrak{o}}\}_L]$$

produced by the quantum master equation of [12, Ch. 5] at scale  $L$ , restricted to the mixed brane-defect interaction  $I_o$  of Problem 5.72. In the ordinary  $\mathfrak{gl}_N$  branch this class carries the residual scalar contact summand  $\hbar N \omega(f, g) \gamma_1$  of E.12, whose scalar projection at the  $U(1)$  centre-of-mass slice is the Capelli class  $\hbar N [\bar{c}]$  of Chapter 6 (Theorem 6.8); in the balanced  $\mathbb{C}^{N|N}$  branch the same summand vanishes by E.16. The scalar projection is therefore the local-stalk value of the projective curvature  $\Omega_{\text{central}}|_{J(f)}$  of the modular line bundle of the closed chiral algebra, by Theorem 6.8.

**PROPOSITION 6.17** (*Endpoint construction is equivalent to vanishing of the tuple*). An endpoint-admissible coefficient module of Definition 5.94 for the brane-defect geometry of 6.16 exists if and only if all four cocycle classes  $\mathfrak{o}_{\text{Cas}}$ ,  $\mathfrak{o}_{\text{def}}$ ,  $\mathfrak{o}_{\text{RG}}$ ,  $\mathfrak{o}_{\text{QME}}$  of 6.16 vanish and a compatible choice of primitives is made. In particular, a Costello–Li type BV-cohomology theorem closes the present coefficient endpoint only when it proves vanishing of all four classes simultaneously, with primitives compatible with the strict finite-window Mittag-Leffler envelope of 5.15.

*Proof.*  $(O1) \Leftrightarrow (E2)$ . Definition 5.94(E2) is the existence of a coefficient kernel  $C_* \in \mathfrak{h}_* \widehat{\otimes} D_*$  whose finite-window projections are the  $C_{\mathfrak{h}}^M$ ; equivalently a primitive for the cocycle representative of  $\mathfrak{o}_{\text{Cas}}$  under the represented-kernel map.

$(O2) \Leftrightarrow (E3)$ . Condition (E3) requires transport of bracket  $\{\cdot, \cdot\}_{\text{open}}$ , propagator  $P_{\epsilon, L}^{\text{base}}$ , interaction  $I_o$ , and counterterm representatives across the brane line  $L$  with ordinary  $\mathfrak{gl}_N$  brackets; equivalently a primitive for the Hom-complex cocycle representative of  $\mathfrak{o}_{\text{def}}$ .

$(O3) \Leftrightarrow (E5)$ . Condition (E5) is preservation of the strict finite-window Mittag-Leffler tower under the Costello recursion  $W(P_{\epsilon, L}, -)$ ; equivalently vanishing of the  $\varprojlim^1 H^\circ$  class of  $\mathfrak{o}_{\text{RG}}$ .

$(O4)$  and the quantum endpoint. The endpoint QME is the vanishing of the degree- $+1$  local-functional class  $\mathfrak{o}_{\text{QME}}$  in  $H^1(\mathcal{O}_{\text{loc}}(\mathcal{E}_w), \Delta)$ , which by E.1 has  $|\Delta| = +1$  on the cotangent-coordinate grading of the local-functional complex.

Necessity is the same chain of equivalences in reverse: a nonzero component removes one of the endpoint data just listed. The primitive-compatibility clause is the requirement that the chosen cobounding primitives be morphisms in the strict ML category  $\mathbf{MLTateChain}_{\geq 0}^{\text{str}}$ , so that the endpoint module sits in  $\mathbf{TateChain}_{\geq 0}^{\text{adm}}$  (5.17).  $\square$

#### Four-input extension criterion

**THEOREM 6.18** (*Costello–Li extension criterion for the mixed coefficient endpoint*). Let  $\mathfrak{o}_{\text{BV}, \text{end}}$  be the obstruction tuple of Definition 6.16. A Costello–Li type BV-cohomology theorem proves vanishing of  $\mathfrak{o}_{\text{BV}, \text{end}}$  on the formal local geometry  $X = \mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}_{\text{o, hol}}^2$ , brane  $L = \mathbb{R}_{\text{brane}} \times \{o\} \times \{p\}$ , gauge  $\mathfrak{gl}_N$ , if and only if it supplies the following four classical and one quantum data simultaneously, each addressing a distinct cocycle representative in  $\mathfrak{o}_{\text{BV}, \text{end}}$ :

- (I1) *Non-compact /  $\Omega$ -background BV-cohomology vanishing* replacing (S1). A vanishing theorem for the BV operator  $\Delta$  on  $(\mathcal{O}_{\text{loc}}(\mathcal{E}_w), \Delta)$  over the non-compact base  $\mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$ , either via a separately anchored  $\Omega$ -background route in the normal bundle  $N_L X = \mathbb{R}_s \oplus \mathbb{C}_{z_1} \oplus \mathbb{C}_{z_2}$ , with the vector field, equivariant differential  $Q_\Omega = Q + \iota_{V_\Omega}$ ,  $Q_\Omega^2 = L_{V_\Omega}$  mechanism, inverted normal weights, residue/Euler normalization, and stratified factorization on  $(X, L)$ , or via compactification with Lagrangian boundary at infinity preserving the brane  $L$  as a fixed locus.



- (I<sub>2</sub>) *Topological–holomorphic mixed kinematics* replacing (S<sub>2</sub>). The vanishing must hold for  $D = d_{\mathbb{R}^2} + \widehat{\partial}_{\mathbb{C}^2}$  on  $\Omega_{\mathbb{R}^2}^\bullet(I) \widehat{\otimes} \Omega_{\mathbb{C}^2}^{\circ,\bullet}(\mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1])$ , together with its linearized BV-kernel complex of 5.210 and the cotangent [1]-extension of the Hamiltonian sector (E.1).
- (I<sub>3</sub>) *Brane-defect extension with ordinary  $\mathfrak{gl}_N$  scalar contact decision* replacing (S<sub>3</sub>)–(S<sub>4</sub>). The vanishing must allow the transverse one-real-dimensional defect  $L$  carrying open gauge  $\mathfrak{gl}_N$ , with comparison to the flat- $\mathbb{C}^d$   $\mathfrak{gl}(N | N)$  setup of [14], and must decide the scalar contact class

$$\hbar N \omega(f, g) \gamma_1 \in H_{\text{Lie}}^2(\bar{A}, \mathbb{C}), \quad \omega(z_1, z_2) = 1,$$

of E.12 — equivalently the projective curvature  $\Omega_{\text{central}}|_{J(f)}$  of the modular line bundle at the trace generator (Theorem 6.8) — by one of the four named cancellations: closed-exchange cancellation,  $\mathbb{C}^{N|N}$  supertrace removal (E.16), central-character normalization (E.15), or renormalized boundary generators.

- (I<sub>4</sub>) *Coefficient-system Tate completion* replacing (S<sub>5</sub>). The coefficient direction  $\widehat{\mathfrak{h}} \widehat{\otimes} \widehat{\mathfrak{h}}_{\text{cont}}^\vee$  must support the Casimir  $C_{\mathfrak{h}} = \sum_{d \geq 1} \sum_i e_{d,i} \otimes e_d^i$  as a convergent represented kernel; by 5.211 this requires a coefficient category change, supplied for example by the weighted Tate pair  $(\mathfrak{h}_w, \mathfrak{h}_{\text{cont}}^{\vee,w})$  of T1.
- (I<sub>5</sub>) *Quantum endpoint vanishing*. A primitive for the degree-+1 cocycle  $\mathfrak{o}_{\text{QME}} = [QI_o + \hbar \Delta_L I_o + \frac{1}{2}\{I_o, I_o\}_L]$  in  $H^1(\mathcal{O}_{\text{loc}}(\mathcal{E}_w), \Delta)$ , with the Costello-2011 counterterm subtraction of 6.11 compatible with the strict-window Mittag-Leffler tower of 5.15. Casimir-balanced equivariant data of 5.148 kill the scalar component  $\mathfrak{o}_{\text{QME}}^{\text{sc}}$ ; the non-scalar residual is the obstruction class of 5.83 in  $H^1(\ker \widehat{\sigma}_{\text{sc}}, d_{\text{QME}})$  together with the secondary  $\varprojlim^1 H^0$  primitive class.

Equivalently, (I<sub>1</sub>)–(I<sub>5</sub>) are exactly the data of Problem 5.48 for the present brane-defect endpoint, after the arity-two formal OPE of Theorem 1.15 has been fixed.

*Proof.* By Proposition 6.13, axes (S<sub>1</sub>)–(S<sub>5</sub>) name pairwise distinct cocycle representatives in  $\mathfrak{o}_{\text{BV},\text{end}}$ : (S<sub>1</sub>)+(S<sub>2</sub>) give the kinematic side of  $\mathfrak{o}_{\text{QME}}$ , (S<sub>3</sub>) for  $\mathfrak{o}_{\text{def}}$ , (S<sub>4</sub>) for the residual  $\hbar N[\bar{c}]$  summand of  $\mathfrak{o}_{\text{QME}}$ , and (S<sub>5</sub>) for  $\mathfrak{o}_{\text{Cas}}$ ; the algebraic-coefficient direction governs  $\mathfrak{o}_{\text{RG}}$ . Components (I<sub>1</sub>)–(I<sub>4</sub>) supply primitives, in turn, for the cocycle representatives forced by (S<sub>1</sub>)+(S<sub>2</sub>), (S<sub>3</sub>)+(S<sub>4</sub>)<sub>transport</sub>, the strict-window Mittag-Leffler preservation of  $\mathcal{W}(P_{e,L}, -)$ , and the Casimir convergence; (I<sub>5</sub>) supplies the quantum primitive. By Proposition 6.17 this is exactly vanishing of  $\mathfrak{o}_{\text{BV},\text{end}}$ , with primitives compatible with **MLTateChain** <sub>$\geq 0$</sub> <sup>str</sup> of 5.15.

For necessity: dropping (I<sub>1</sub>) leaves the kinematic representative of  $\mathfrak{o}_{\text{QME}}$  unprimitivated; dropping (I<sub>2</sub>) leaves  $D$  requiring its endpoint vanishing theorem; dropping (I<sub>3</sub>) retains  $\mathfrak{o}_{\text{def}}$  and the scalar contact summand of  $\mathfrak{o}_{\text{QME}}$ ; dropping (I<sub>4</sub>) retains  $\mathfrak{o}_{\text{Cas}}$  as a nonzero cokernel class by 5.211; dropping (I<sub>5</sub>) retains the degree-+1 anomaly primitive as endpoint data. The chain-level primitive shadow of 5.132 closes the classical indecomposable shadow by an algebraic argument; (I<sub>3</sub>) and (I<sub>5</sub>) supply the brane-defect scalar and quantum endpoint inputs. The Quillen envelope of 5.30 promotes the strict category of (I<sub>4</sub>) to an admissible Tate model envelope.  $\square$

## 6.2 Operators in the closed Hamiltonian sector

On the closed side of the local mixed model, Hamiltonian-dual CE coordinates produce Schouten vector-field operations on the brane polynomial algebra, while shifted-cotangent-dual CE coordinates produce boundary Hamiltonian observables on the Chan–Paton pair. The closed local operator algebra is, after one kinematic contraction and one BRST identification, the continuous CE cochains

$$\mathcal{O}_{\text{grav}}^{\text{cl}} \cong C_{\text{CE},\text{cont}}^\bullet(\mathfrak{g}), \quad \mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1],$$

with  $\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$  the constants-quotiented Hamiltonian polydisk algebra and  $\mathfrak{h}_{\text{cont}}^\vee \cong \mathcal{P}$  its continuous residue dual (Theorem 5.178); this is Proposition 6.20.

The kinematic BRST operator  $Q_o = d_{\mathbb{R}^2} + \bar{\partial}_{\mathbb{C}^2}$  contracts every positive topological and anti-holomorphic jet by the Euler homotopy in the real direction (Lemma 3.21) and the formal Dolbeault Poincaré lemma in the holomorphic normal directions;  $Q_o$ -cohomology retains only the holomorphic Taylor coefficients of the Hamiltonian field  $\alpha$  and the cotangent field  $\beta$ . The residual interaction differential reproduces the Lie brackets of  $\mathfrak{g}$ : two Hamiltonian  $\alpha$ -inputs contribute the closed Poisson bracket  $\{f, g\}$ , one Hamiltonian and one cotangent input contribute the coadjoint action, and two cotangent inputs contribute zero. The residual differential is the Chevalley–Eilenberg differential of  $\mathfrak{g}$ , and the closed local operator algebra is its continuous cochain algebra.

The open side of §3.5 supplies the ghost-zero connected trace monomials  $\overline{\text{Tr}(\phi_1^k \phi_2^l)}$ , the primitive one- $\psi$  Koszul descendants  $\Psi_{a,b}$ , and the open-trace  $\mathcal{A}_\infty$ -Koszul acceptance datum of Remark 3.44. The closed side matches these as continuous CE cochains via the coordinate CE/PV identity of Theorem 5.5: on the coordinate window, Hamiltonian-dual CE coordinates  $c^I$  map to Schouten polyvector coordinates  $\theta^I$  on  $\widehat{\mathbf{S}}(\bar{A})$ , and shifted-cotangent-dual CE coordinates  $u_I$  map to boundary Hamiltonian observables  $O_I$  on the Chan–Paton pair. The assignment  $c_f \mapsto O_f$  is a type error (Theorem 4.20). The full stratified comparison is §5.7 below.

Let  $\mathfrak{h} = \widehat{\mathfrak{h}} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$  with the projected Poisson bracket, and let  $\mathfrak{h}^\vee$  be its continuous Taylor dual. The cotangent field is a coadjoint module, so the local Hamiltonian Lie algebra is

$$\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}^\vee[1].$$

For  $f, g \in \mathfrak{h}$  and  $\lambda, \mu \in \mathfrak{h}^\vee$ , the brackets are

$$[f, g] = \{f, g\}, \quad [f, \eta_\lambda] = \eta_{\text{ad}_f^\vee \lambda}, \quad [\eta_\lambda, \eta_\mu] = 0,$$

where  $(\text{ad}_f^\vee \lambda)(g) = -\lambda(\{f, g\})$ . On polynomial monomials in the dense subalgebra  $\bar{A}$  this reads

$$[z_1^k z_2^l, z_1^m z_2^n] = \overline{(kn - lm) z_1^{k+m-1} z_2^{l+n-1}},$$

where the bar means projection to  $\bar{A}$ ; in particular the constant bracket of  $z_1$  and  $z_2$  is zero in the reduced Hamiltonian Lie algebra.

LEMMA 6.19 (*Shifted coadjoint signs*). Let  $\eta_\lambda$  denote the suspension of  $\lambda \in \mathfrak{h}^\vee$  into  $\mathfrak{h}^\vee[1]$ . Since the Hamiltonians in  $\mathfrak{h}$  are even,

$$[f, \eta_\lambda] = \eta_{\text{ad}_f^\vee \lambda}, \quad [\eta_\lambda, \eta_\mu] = 0.$$

Consequently the enveloping algebra relations involving two shifted coadjoint generators are odd-commutative:

$$\eta_\lambda \eta_\mu + \eta_\mu \eta_\lambda = 0.$$

*Proof.* The semidirect product bracket before shifting is the ordinary coadjoint action of the even Lie algebra  $\mathfrak{h}$  on its continuous dual. Suspending the module raises the degree of every  $\lambda$  by one. The mixed-bracket sign is  $+1$  because  $|f| = 0$ . The shifted module is abelian as a Lie ideal, so its bracket with itself is zero; in the associative enveloping algebra this zero Lie bracket gives  $\eta_\lambda \eta_\mu - (-1)^{1+1} \eta_\mu \eta_\lambda = 0$ .  $\square$

The kinematic BRST operator is

$$Q_o = d_{\mathbb{R}^2} + \bar{\partial}_{\mathbb{C}^2}.$$

The same jet contraction as in Lemma 3.21, together with the Dolbeault Poincaré lemma in the formal local model, leaves precisely the holomorphic Taylor coefficients. Thus the linear local coordinates are the functionals

$$\partial_{z_1}^k \partial_{z_2}^l \alpha|_o, \quad \partial_{z_1}^k \partial_{z_2}^l \beta|_o.$$

With the coefficient-dual convention fixed above, these are represented by  $(k!l!)^{-1} \partial_{z_1}^k \partial_{z_2}^l |_o$ . Since zero-form components of  $\alpha$  and  $\beta$  have negative field ghost numbers, the corresponding linear functions have positive ghost numbers.

**PROPOSITION 6.20** (*Closed local operators are continuous CE cochains*). The classical local operators of the formal Hamiltonian cotangent BF sector are the continuous Chevalley-Eilenberg cochains

$$O_{\text{grav}}^{\text{cl}} \cong C_{\text{CE,cont}}^{\bullet}(\mathfrak{g}) = \widehat{\mathbf{S}}(\mathfrak{g}_{\text{cont}}^{\vee}[-1]),$$

with CE differential dual to the Lie brackets above.

*Proof.* Work at the formal insertion point. The jet algebra of the fields is the completed free graded-commutative algebra on the Taylor coefficients of  $\alpha$  and  $\beta$ . The kinematic differential  $Q_o = d_{\mathbb{R}^2} + \bar{\partial}_{\mathbb{C}^2}$  contracts all positive real and anti-holomorphic jets: in the real directions the homotopy is  $[d_{\mathbb{R}^2}, \iota_{\partial_{x_i}}] = \partial_{x_i}$ , and in the anti-holomorphic directions it is the formal Dolbeault Poincaré homotopy. The  $Q_o$ -cohomology of the jet algebra is therefore generated by the holomorphic Taylor coefficients alone.

The remaining  $\alpha$ -coefficients are Taylor Hamiltonians modulo constants, hence elements of  $\mathfrak{h} = \widehat{\mathfrak{h}}$ ; degree-wise polynomial coefficients form the dense subalgebra  $\bar{A}$ . The cotangent field  $\beta$  supplies the odd shifted continuous dual  $\mathfrak{h}_{\text{cont}}^{\vee}[1]$ . Thus the linear space of residual local modes is  $\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^{\vee}[1]$ .

The nonlinear BRST differential is determined by the BV Hamiltonian

$$S = \int \left\langle \beta, D\alpha + \frac{1}{2} \{\alpha, \alpha\} \right\rangle.$$

The linear part is exactly  $Q_o$ , already killed in cohomology. The quadratic term sends two Hamiltonian  $\alpha$ -modes  $f, g$  to their reduced Poisson bracket  $\{f, g\}$ , and the action on the cotangent variable is the coadjoint action displayed above. The quadratic term has zero component with two cotangent inputs, so  $[\eta_\lambda, \eta_\mu] = 0$ . Hence the residual BRST differential on functions is precisely the derivation dual to the Lie bracket on  $\mathfrak{g}$ , which is the Chevalley-Eilenberg differential. Continuity records the completion over polynomial degree.  $\square$

**Remark 6.21** (*Strict-product  $P_o$  comparison block*). The cochain-level CE/PV identity of Theorem 5.5 matches the generators of  $C_{\text{CE,cont}}^{\bullet}(\mathfrak{g})$  to Schouten polyvectors of the boundary polynomial algebra  $\widehat{\mathbf{S}}(\bar{A})$ . The match is exact only on the coordinate-window CE generators; promoting it to a strict product  $P_o$ -comparison on the full Hamiltonian polydisk has a Tate-Casimir obstruction. The full disk algebra  $\mathfrak{h}$  carries the product/direct-sum Tate pair  $\mathfrak{h} \oplus \mathfrak{h}_{\text{cont}}^{\vee}$  of Lemma 5.211: the coefficient coevaluation  $C_{\mathfrak{h}} = \sum_d \sum_i e_{d,i} \otimes e_d^i$  diverges in each completed tensor factor; the Casimir obstruction  $\mathfrak{o}_{\text{Cas}}$  is the block, repaired only by the weighted Tate completion (Theorem 5.70) or by the nilpotent truncation  $\mathfrak{m}^3/\mathfrak{m}^{N+1}$ . Promotion of the generator-level CE/PV identity to a strict factorisation  $P_o$ -centre morphism on the unreduced cotangent side is governed by the obstruction tuple Problem 5.48, with classical

Tate-coefficient closure reduced to Problem 5.221, and Roos/Cech-descent compatibility recorded in Remark 5.161. The PBW special-fibre/coadjoint deformation separation (Theorem 5.180, Corollary 5.172) shows the polynomial one- $\psi$  descendants of §3.5 carry a different  $\mathfrak{h}$ -module structure than the residue cotangent module  $\mathcal{P}$ ; the cotangent side admits a unique residue realization, separated from any polynomial substitute by Theorem 5.182. Three obstructions remain:  $\mathfrak{o}_{\text{Cas}}$  (BFM Casimir), the unreduced lift obstruction  $\text{Ob}_{\mathfrak{o}, \text{unred}}$  of Remark 5.189, and the PBW/coadjoint module-structure mismatch. They are repaired in §5.7 layer by layer.

The Capelli scalar  $\hbar N[\bar{c}] = \Omega_{\text{central}}|_{J(f)}$  is the projective curvature of the modular line bundle evaluated at the trace generator (Theorem 6.8); it is one component of a single curvature  $F_{\mathcal{T}}(\Theta_{\text{loc}})$  in the master deformation complex  $\mathfrak{R}_{\mathcal{T}}$ , whose four-curvature taxonomy classifies the global obstructions and locates the projective cell.

*All-bidegree Capelli through the radial-parts identity.* The linear-generator collapse of Theorem 6.6 extends to all bidegrees  $(a, b)$  through a per-bidegree exact rational reduction: at each bidegree  $(a, b)$  the radial bicomplex carries a finite matrix  $B_{a,b} \in M_{r_0 \times r_1}(\mathbb{Q})$  and a comparison vector  $v_{a,b} = (T_{a,b}, E_{a,b}^+)$ , and  $\Omega_{a,b}^{\text{rad}} \in \text{coker } B_{a,b}$  classifies whether  $v_{a,b}$  lies in  $\text{im } B_{a,b}$  (Bareiss elimination over  $\mathbb{Q}$  decides the question in exact rational arithmetic; §F, Theorem F.44). When  $v_{a,b} \in \text{im } B_{a,b}$  the elimination returns the unique correction  $C_{a,b}^+$  with  $B_{a,b}C_{a,b}^+ = v_{a,b}$  and  $\Omega_{a,b}^{\text{rad}} = \mathfrak{o}$ ; otherwise the obstruction covector  $(\phi, -\lambda) \in \ker B_{a,b}^*$  realises the equivalence class  $[(T_{a,b}, E_{a,b}^+)] \in \text{coker } B_{a,b}$ . On the listed range of Corollary F.45 all classes vanish and the Capelli identification extends; outside the listed range  $\Omega_{a,b}^{\text{rad}}$  is the counterterm-cell  $(d_{\Theta}\tau)$  coordinate of  $F_{\mathcal{T}}(\Theta_{\text{loc}})$  in the four-curvature taxonomy of Theorem 7.1. The Razmyslov–Procesi radical structure of the trace ideal at finite  $N$  lies outside the radial reduction: every step is exact rational and basis-canonical.

## 7 THE OBSTRUCTION CALCULUS: MASTER DEFORMATION COMPLEX OF LOCAL-TO-GLOBAL COHERENCE

The four-curvature taxonomy classifies the global obstructions to the formal-Darboux stalk identification of Theorem 5.6: the single curvature  $F_{\mathcal{T}}(\Theta_{\text{loc}}) \in H^2(\mathfrak{R}_{\mathcal{T}}, \Theta_{\text{loc}})$  of the master deformation complex  $\mathfrak{R}_{\mathcal{T}}$  lies on exactly one of four cells — strict, counterterm-coherent, projective, or target-replacement — and the six manuscript obstruction classes  $\Omega_{a,b}^{\text{rad}}, \hbar N[\bar{c}], [\text{Ob}_{w,\partial,\hbar}^{\text{red}}], \text{Ob}_{\text{BM}}^{\Pi}, \text{Ob}_{\text{cent/QME/desc}}, \text{per}_X(v)$  are six gauge-slice coordinates of this single curvature. The convention layer of § fixes the symbols:  $d = \dim_{\mathbb{C}} X$ ,  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}^2_{\mathfrak{o}, \text{hol}}$  as local geometry,  $\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$  and  $\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^{\vee}[1]$  as the native shifted-cotangent BF Lie algebra, the holomorphic  $E_2$  factorization-algebra surface, the BV / Koszul / Gerstenhaber sign rules of Appendix C, the explicit Costello propagator weight in heat-kernel renormalization, the  $\hbar$  versus  $g$ , discipline, the Hochschild / cyclic / negative-cyclic distinction, the Capelli scalar  $\hbar N[\bar{c}] = \Omega_{\text{central}}|_{J(f)}$  of Theorem 6.8, and the cross-target firewall acceptance class  $\text{Ob}_{\text{UKD}}(\mathfrak{C}_{\text{tar}})$  of Theorem 10.13. Every class named below is a coordinate of the curvature decomposition recorded under that convention layer.

The shifted-cotangent BF Lie algebra of §2,  $\mathfrak{g} = \widehat{\mathfrak{h}} \ltimes \mathfrak{h}_{\text{cont}}^{\vee}[1]$ , is the universal infinitesimal deformation of the closed BV pair on  $(\widehat{\mathbb{C}}^2_{\mathfrak{o}}, dz_1 \wedge dz_2)$ : every infinitesimal compatible local action carrying the Hamiltonian bracket and its coadjoint cotangent module is a Maurer–Cartan element of  $\mathfrak{g}$  (Theorem 2.11). Its filtered  $L_{\infty}$ -deformation complex  $(\mathfrak{R}_{\mathcal{T}}, F^{\bullet}, d, \ell_2, \dots)$  is the target whose Maurer–Cartan stratum carries the trace substitution  $J$  as the local datum  $\Theta_{\text{loc}}$ . Image under quantization, globalization, factorization-centre lift, modular clutching, and CY-categorical specialisation produces

the obstruction classes  $\hbar N[\bar{c}]$ ,  $\Omega_{a,b}^{\text{rad}}$ ,  $[\text{Ob}_{w,\partial,h}^{\text{red}}]$ ,  $\text{Ob}_{\text{cent/QME/desc}}$ ,  $\text{Ob}_{\text{BM}}^{\Pi}$ , and  $\text{per}_X(v)$  as the components of a single curvature decomposition; the four-curvature taxonomy that classifies them is proved in §7.5.

### 7.1 Filtered $L_\infty$ target deformation complex

The data of a target deformation problem are  $(\mathfrak{R}_{\mathcal{T}}^\bullet, F^\bullet, d, \ell_2, \ell_3, \dots)$ : a filtered  $L_\infty$ -algebra in cohomological degree, with filtration  $\mathfrak{R}_{\mathcal{T}} = F^0 \supseteq F^1 \supseteq F^2 \supseteq \dots$  that is exhaustive and complete in the sense that  $\mathfrak{R}_{\mathcal{T}} \xrightarrow{\sim} \varprojlim_n \mathfrak{R}_{\mathcal{T}}/F^n$ , with each  $\ell_k$  of weight at least  $k-1$  for the filtration so that the curvature series of an element of  $F^1$  converges in the filtered topology.

Given a degree-one element  $\Theta \in F^1 \mathfrak{R}_{\mathcal{T}}^1$ , the curvature is the formal series

$$F_{\mathcal{T}}(\Theta) = d\Theta + \frac{1}{2}\ell_2(\Theta, \Theta) + \frac{1}{3!}\ell_3(\Theta, \Theta, \Theta) + \dots,$$

convergent in the filtered topology. The Maurer–Cartan equation  $F_{\mathcal{T}}(\Theta) = 0$  cuts out the strict  $\mathcal{T}$ -coherences. Twists are filtered cochains  $\tau \in F^1 \mathfrak{R}_{\mathcal{T}}^1$  (or, when the necessary extension is structural and higher than first order, central extensions of  $\mathfrak{R}_{\mathcal{T}}$  by an abelian filtered cochain complex):

$$\text{TwGlob}_{\mathcal{T}}(\Theta) = \{ \tau \in F^1 \mathfrak{R}_{\mathcal{T}}^1 : F_{\mathcal{T}}(\Theta + \tau) = 0 \} / \text{gauge}.$$

A morphism of target deformation complexes  $\mathfrak{R}_{\mathcal{T}} \rightarrow \mathfrak{R}_{\mathcal{T}'}$  is a filtered  $L_\infty$ -morphism, sending Maurer–Cartan elements to Maurer–Cartan elements.

### 7.2 The transported local datum

The formal-stalk-chart hypotheses of Theorem 5.146 – Chan–Paton substitution  $z_i \rightsquigarrow \phi_i$ , the Dirac constraint  $[\phi_1, \phi_2] = 0$ , and the trace map  $J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)}$  – supply the underlying datum; the choice of target supplies its controlling deformation algebra.

*Global Hamiltonian descent.* For a holomorphic-symplectic ambient  $(X, \omega)$  with a brane  $L$  and a Darboux cover  $\{U_i\}$  symplectomorphic to  $\widehat{\mathbb{C}^2}_0$ , the local data are charts  $\Phi_i : U_i \rightarrow \widehat{\mathbb{C}^2}_0$  with local Hamiltonians  $h_i$ ; the deformation complex is  $\mathfrak{R}_{\text{glob}} = \text{Tot } C_{\check{C}/\text{WR}}^\bullet(\{U_i\}, \mathfrak{R}_{\text{loc}})$  – the totalization of the Čech–Weiss–Ran complex of local formal-stalk-chart coefficients. The transported element is  $\Theta_{\text{glob}} = (\Theta_i, \alpha_{ij}, \beta_{ijk}, \dots)$ , with curvature  $F_{\text{glob}}(\Theta_{\text{glob}})$  carrying the Hamiltonian period  $\text{per}_X(v) = [h_i - h_j] \in H^1(X, \mathbb{C}_X)$  as one component.

*Costello quantum master equation.* The Costello deformation complex is the filtered local-functional complex  $\mathfrak{Q}_{w,\partial,h}^\bullet(I)$  with differential  $d_{\text{QME}} = Q + \{I, -\} + \hbar\Delta$ ; the transported element is the classical interaction  $I_{\text{loc}}$  of the open Hamiltonian BF theory, completed in the weighted-Tate envelope of Theorem 5.57. After scalar-symbol projection  $\widehat{\sigma}_{\text{sc}}$ , the reduced non-scalar curvature is  $[F_{\text{QME}}(\Theta_{\text{loc}})] = [\text{Ob}_{w,\partial,h}^{\text{red}}] \in H^1(\ker \widehat{\sigma}_{\text{sc}}, d_{\text{QME}})$ .

*Unreduced  $P_0$ -factorization-centre lift.* The deformation complex is the mapping complex  $\mathfrak{F}_{\text{fact}} = \text{RHom}(\text{Obs}_{\text{closed},H}^{\text{cl}}|_L, Z_{\text{fact}}^{P_0}(\text{Obs}_{\text{open},N}^{\text{cl}}))$ , augmented by BV/QME and stratified Weiss–Čech–Roos descent directions. The transported element is the reduced CE/PV cochain morphism of Theorem 4.20 restricted to the brane. The curvature decomposes as the four-component vector  $\text{Ob}_{\text{cent/QME/desc}} = (\text{ob}_{\text{cent}}^{\text{prod}}, \text{ob}_{\text{cent}}^{P_0}, \text{ob}_{\text{QME}}^{\text{bd}}, \partial_{\check{C}/R}^{\text{Weiss}})$  of Theorem E.53.

*Decorated radial Stokes / Weyl normal form.* The deformation complex is the finite-dimensional radial bicomplex of Definition F.27 with decoration generators for PBW/Stokes corrections. At each bidegree  $(a, b)$  the transported element is the radial comparison vector  $v_{a,b} = (T_{a,b}, E_{a,b}^+)$ ; the curvature is the per-bidegree class  $\Omega_{a,b}^{\text{rad}} = [(T_{a,b}, E_{a,b}^+)] \in \text{coker } B_{a,b}$ , and the obstruction dual is  $(\phi, -\lambda) \in \ker B_{a,b}^*$  with  $\lambda(E_{a,b}^+) - \phi(T_{a,b}) \neq 0$ .

*Strict all-window Bochner–Martinelli current transfer.* The deformation complex is the tower of finite Matlis windows  $P_{\leq N} = \bigoplus_{a+b \leq N} \mathbb{C} \rho_{a,b}$  with action of  $\mathfrak{h}$  by Hamiltonian vector fields; the transported element is the brane current pair  $(P_{\leq N}, \rho_{0,1})$ . The curvature is the residue-coadjoint leak  $z_1^{N+2} \cdot \rho_{N+1,0} = -(N+2)\rho_{0,1}$ , encoded as the obstruction  $\text{Ob}_{\text{BM}}^\Pi$ .

*Modular open-closed clutching.* Modular coherence on the brane-line cyclic factorization dg category  $C^{\text{op}}$  is a target deformation problem of the same shape as the previous six, controlled by the strict law  $d\Theta_C + \frac{1}{2}[\Theta_C, \Theta_C] = 0$  in  $Z_{\text{der}}^{\text{ch}}(C^{\text{op}})$  and its  $\Omega_{\text{mod}}$ -twisted version carrying the determinant-line twist of the projective clutching law  $\text{Cl}^* \text{Tr}_{g+1, n-2} = \text{contract}_C(\text{Tr}_{g,n}) \otimes e^{\Omega_{\text{node}}}$ . The data  $(b, \text{Tr}_C, \Theta_C^{\text{oc, log}}, \Omega_{\text{mod}})$  belong to the companion programme (Vol II 3d HT bulk-boundary; Vol III CY cyclic trace); cross-target use passes through the matched-conventions firewall of §10.7.

### 7.3 Twists: three mechanisms

A nonvanishing curvature class admits three repair mechanisms: null-homotopy, central extension, and target replacement.

**Null-homotopy twist.** The class is exact in the original target:  $F_{\mathcal{T}}(\Theta) = d_{\Theta}\tau$ . The corrected element  $\Theta + \tau$  satisfies the strict Maurer–Cartan equation in the same target. This is the language of QME counterterms, homotopy-centrality witnesses, decorated radial Stokes corrections, Weiss/Ran descent homotopies, and many finite-window radial adjustments. The model example is the  $\theta_3$  finite-window  $H^1$ : bare  $H^1 = \mathbb{C} \cdot E_{\theta_3} \neq 0$  under the cleared QME differential, but supplied-counterterm  $H^1 = 0$  under the unique local counterterm  $dC = -E$ ,  $c = 1$  (Theorem E.38).

**Structural / central twist.** The class is closed and central:  $F_{\mathcal{T}}(\Theta) = \omega$ ,  $d\omega = 0$ ,  $\omega$  central. The repair is to enlarge  $\mathfrak{R}_{\mathcal{T}}$  by a central extension carrying  $\omega$  on the trace line. The corrected element satisfies the master equation in the extended target, and  $\Theta$  becomes a section of a projective structure. The scalar anomaly  $\hbar N[\tilde{c}] \in H_{\text{Lie}}^2(\tilde{A}, \mathbb{C})[[\hbar]]$  is the prototype: the ordinary scalar-reduced Costello obstruction in the  $\mathfrak{gl}_N$  branch obstructs internal counterterms, and the admissible repair is the central extension  $0 \rightarrow \mathbb{C} \cdot \mathbf{1} \rightarrow \mathfrak{h}_{\hbar, N} \rightarrow \tilde{A} \rightarrow 0$  under which the brane carries a projective Hamiltonian action  $[\Phi_{\hbar}(f), \Phi_{\hbar}(g)]_{\star} = \hbar \Phi_{\hbar}(\{f, g\}) + \hbar N \tilde{c}(f, g) \mathbf{1}$ . The same mechanism produces the global gerbe-valued descent of Hamiltonian sections in the period class  $\text{per}_X(v)$ , where the gerbe is the central extension of the Čech–Weiss–Ran complex by the descent cocycle.

**Target-replacement twist.** The class is closed, inexact, and noncentral: every admissible extension of  $\mathfrak{R}_{\mathcal{T}}$  for which the corrected curvature is zero is obstructed. Coherence requires a target substitution  $\mathcal{T} \rightsquigarrow \mathcal{T}'$  along a forced embedding. The strict all-window Bochner–Martinelli current transfer is the prototype: in every fixed finite Matlis window  $\mathcal{P}_{\leq N}$ , the residue-coadjoint Jacobiator  $(N+1)(N+2)\rho_{0,1} \neq 0$  is non-absorptive in the strict multiplication structure, so internal counterterm restoration is obstructed. The pro-Matlis tower  $\widehat{\mathcal{P}}_{\Pi} = \varprojlim_N \mathcal{P}_{\leq N}$  (Corollary 8.41) is the supremum form selected by the leak: the modified coherence law is  $\pi_N(f \cdot \xi) = \pi_N(f \cdot \pi_{N+r(f)}\xi)$ , Hamiltonian action is

coherent only in the pro-tower, and any fixed finite current window is a measurement truncation of the closed dynamical subsystem.

#### 7.4 Four-curvature taxonomy and six-coordinate classification

**THEOREM 7.1** (*Four-curvature taxonomy of the formal-Darboux stalk identification*). Let  $\mathcal{T}$  be a target deformation problem with filtered  $L_\infty$  algebra  $(\mathfrak{R}_{\mathcal{T}}, F^\bullet, d, \ell_k)$ , and let  $\Theta_{\text{loc}} \in F^1 \mathfrak{R}_{\mathcal{T}}^1$  be the transported local formal-stalk-chart datum. Call an *admissible extension* of  $\mathfrak{R}_{\mathcal{T}}$  any  $L_\infty$  central extension by a closed degree-2 cocycle. Exactly one of the following four cases obtains.

- (i) Strict coherence:  $F_{\mathcal{T}}(\Theta_{\text{loc}}) = 0$  in  $\mathfrak{R}_{\mathcal{T}}^2$ , and the  $\mathcal{T}$ -coherence is unique up to gauge.
- (ii) Counterterm coherence: the class  $[F_{\mathcal{T}}(\Theta_{\text{loc}})]$  is exact in  $H^2(\mathfrak{R}_{\mathcal{T}}, \Theta_{\text{loc}})$ , with  $\tau \in F^1 \mathfrak{R}_{\mathcal{T}}^1$  satisfying  $F_{\mathcal{T}}(\Theta_{\text{loc}} + \tau) = 0$  in  $\mathfrak{R}_{\mathcal{T}}$ ; the corrected element is the  $\tau$ -twisted  $\mathcal{T}$ -coherence in the same target.
- (iii) Projective coherence: the class is central and inexact,  $[F_{\mathcal{T}}(\Theta_{\text{loc}})] = [\omega]$  with  $\omega$  central in  $\mathfrak{R}_{\mathcal{T}}$ , and the corrected element solves the master equation in a central extension by  $\omega$ ; the global structure is projective.
- (iv) Target replacement: outside cases (i)–(3),  $[F_{\mathcal{T}}(\Theta_{\text{loc}})]$  is nonzero in  $H^2(\mathfrak{R}_{\mathcal{T}}, \Theta_{\text{loc}})$ , and every admissible extension has an obstruction to a central representative; coherence exists only after a substitution  $\mathcal{T} \rightsquigarrow \mathcal{T}'$  supplied as external data.

The local manuscript theorems decompose as follows.

| Class                                     | Curvature meaning                    | Mechanism                           |
|-------------------------------------------|--------------------------------------|-------------------------------------|
| $\hbar N[\tilde{c}]$                      | scalar-reduced Hamiltonian action    | central extension, projective       |
| $\Omega_{a,b}^{\text{rad}}$               | Weyl/radial all-bidegree normal form | decorated PBW/Stokes counterterm    |
| $[\text{Ob}_{w,\partial,h}^{\text{red}}]$ | reduced non-scalar QME               | Costello local counterterm          |
| $\text{Ob}_{\text{cent/QME/desc}}$        | unreduced factorization-centre lift  | derived centre, homotopy-centrality |
| $\text{Ob}_{\text{BM}}^{\Pi}$             | strict finite-current transfer       | target replacement, pro-Matlis      |
| $\text{per}_X(v)$                         | global Hamiltonian descent           | Hamiltonian gerbe, projective       |

The clause assignment follows from the obstruction theorem of §7.5 and the appendices.

**THEOREM 7.2** (*Six coordinates of one curvature*). The six manuscript obstruction classes

$$\Omega_{a,b}^{\text{rad}}, \quad \hbar N[\tilde{c}], \quad [\text{Ob}_{w,\partial,h}^{\text{red}}], \quad \text{Ob}_{\text{BM}}^{\Pi}, \quad \text{Ob}_{\text{cent/QME/desc}}, \quad \text{per}_X(v)$$

are the six gauge-slice components of the single curvature  $F_{\mathcal{T}}(\Theta_{\text{loc}})$  of Theorem 7.1, evaluated at six independent gauge slices of  $(\mathfrak{R}_{\mathcal{T}}, \Theta_{\text{loc}})$ , with cell assignment  $(d_{\Theta}\tau, \Omega_{\text{central}}, d_{\Theta}\tau, \notin \text{im } d_{\Theta}, d_{\Theta}\tau, \Omega_{\text{central}})$ .

*Proof.* The slice and cell assignment is established for each coordinate through the corresponding section of the manuscript:  $\Omega_{a,b}^{\text{rad}}$  in the radial bicomplex (Appendix F, Theorem F.44; counterterm cell  $d_{\Theta}\tau$ );  $\hbar N[\tilde{c}]$  in scalar Lie cohomology (Lemma 6.2, Theorem 6.8; projective cell  $\Omega_{\text{central}}$ );  $[\text{Ob}_{w,\partial,h}^{\text{red}}]$  in the weighted Tate complex (Theorem E.38; counterterm cell);  $\text{Ob}_{\text{BM}}^{\Pi}$  on finite Matlis windows (Theorem 8.40; genuine-obstruction cell  $\notin \text{im } d_{\Theta}$ );  $\text{Ob}_{\text{cent/QME/desc}}$  in the unreduced cotangent lift (Theorem E.53; counterterm cell);  $\text{per}_X(v)$  as the global period class on  $X$  (Theorem 5.146; projective cell). The six slices are independent because their support classes are non-overlapping in  $\mathfrak{R}_{\mathcal{T}}$ ; the single

curvature  $F_{\mathcal{T}}(\Theta_{\text{loc}})$  projects to each slice via the gauge-slice projection induced by the corresponding filtration of  $\mathfrak{R}_{\mathcal{T}}$ .  $\square$

### 7.5 Tate-coefficient cotangent lift: the six gauge slices in body

The unreduced cotangent extension of the bracket match  $J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)}$  of §4.3 (Theorem 4.5, Theorem 4.11) is a single filtered  $L_{\infty}$  deformation problem  $(\mathfrak{R}_{\mathcal{T}}, F^{\bullet}, d, \ell_2, \ell_3, \dots)$  with transported local datum  $\Theta_{\text{loc}} \in F^1 \mathfrak{R}_{\mathcal{T}}^1$  and curvature

$$F_{\mathcal{T}}(\Theta_{\text{loc}}) = d\Theta_{\text{loc}} + \frac{1}{2}\ell_2(\Theta_{\text{loc}}, \Theta_{\text{loc}}) + \frac{1}{3!}\ell_3(\Theta_{\text{loc}}, \Theta_{\text{loc}}, \Theta_{\text{loc}}) + \dots$$

in the filtered topology of §7.1. The classifying class  $[F_{\mathcal{T}}(\Theta_{\text{loc}})] \in H^2(\mathfrak{R}_{\mathcal{T}}, d_{\Theta_{\text{loc}}})$  sits in exactly one of four cells of the four-curvature taxonomy  $\{\circ, d_{\Theta}\tau, \Omega_{\text{central}}, \notin \text{im } d_{\Theta}\}$  of Theorem 7.1: strict, gauge-trivial counterterm, projective via central extension, or genuine obstruction. The six manuscript classes

$$\Omega_{a,b}^{\text{rad}}, \quad \hbar N[\bar{c}], \quad [\text{Ob}_{w,\partial,h}^{\text{red}}], \quad \text{Ob}_{\text{BM}}^{\Pi}, \quad \text{Ob}_{\text{cent/QME/desc}}, \quad \text{per}_X(v)$$

are the gauge-slice components of the single curvature  $F_{\mathcal{T}}(\Theta_{\text{loc}})$ , evaluated at six independent gauge slices of  $(\mathfrak{R}_{\mathcal{T}}, \Theta_{\text{loc}})$ , with cell assignment  $(d_{\Theta}\tau, \Omega_{\text{central}}, d_{\Theta}\tau, \notin \text{im } d_{\Theta}, d_{\Theta}\tau, \Omega_{\text{central}})$  established in Theorem 7.2. The strict cochain-level morphism of factorization algebras

$$\Phi_{\text{cl}}: \text{Obs}_{\text{closed}, H}^{\text{cl}}|_L \longrightarrow Z_{\text{fact}}^{P_{\circ}}(\text{Obs}_{\text{open}, N}^{\text{cl}})$$

realising the cotangent direction  $\mathfrak{h}_{\text{cont}}^{\vee}[1]$  with all  $P_{\circ}$ -centrality homotopies and a boundary realisation of the shifted-cotangent generators is the construction problem Problem 5.48.

The four-component obstruction vector  $(\text{ob}_{\text{cent}}^{\text{prod}}, \text{ob}_{\text{cent}}^{P_{\circ}}, \text{ob}_{\text{QME}}^{\text{bd}}, \delta_{\dot{C}/R}^{\text{Weiss}})$  of Theorem E.53 resolves the Tate-completed lift inside the  $(d_{\Theta}\tau)$ -cell: joint vanishing with chosen null-homotopies constructs the morphism, and a nonzero entry is the precise construction defect. The chain-level primitive projection of Theorem 5.132 computes the  $E_1$ -page of the spectral sequence converging to  $\text{ob}_{\text{cent}}^{P_{\circ}}$ .

The first proved nonzero entry is the Casimir-kernel block  $\mathfrak{o}_{\text{Cas}}$ : the strict product/direct-sum Tate pair  $(\mathfrak{h}_{\Pi}, D_{\oplus})$  has an obstruction to a coefficient kernel projecting to every finite Casimir  $C_{\mathfrak{h}}^M = \sum_{d \leq M} \sum_i e_{d,i} \otimes e_d^i$  (Lemma 5.211, Theorem 5.93), so the strict-unweighted endpoint is foreclosed. Theorem 8.40 extends this to the all-window analytic statement: strict all-window support-local Hamiltonian transfer has an obstruction on every choice of one-pair principal-part datum, with obstruction class  $\text{Ob}_{\text{BM}}^{\Pi}$  on the BMK ladder. By Corollary 8.41 the pro-Matlis retract is the maximal admissible refinement; any larger strict refinement is blocked by an entry of  $\text{Ob}_{\text{BM}}^{\Pi}$ . The unreduced cotangent lift therefore has its obstruction on the strict ambient product completion and succeeds inside a controlled refinement: the weighted Tate completion (Theorem 5.70), the nilpotent truncation  $\mathfrak{m}^3/\mathfrak{m}^{N+1}$ , or the pro-Matlis retract.

*Definition 7.3 (Tate-coefficient cotangent lift obstruction vector).* The Tate-coefficient cotangent lift obstruction vector for the formal local model  $X = \mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}_{\text{o,hol}}^2$  with brane line  $L = \mathbb{R}_{\text{brane}} \times \{\circ\} \times \{\circ\}$  and ordinary  $\mathfrak{gl}_N$  gauge data is

$$\text{Ob}_T = (\mathfrak{o}_{\text{Cas}}, \mathfrak{o}_{\text{nilp}}, \mathfrak{o}_{\text{Quil}}, \mathfrak{o}_{\text{BV, end}}, \mathfrak{o}_{\text{prim}}, \mathfrak{o}_{\text{Had}}, \mathfrak{o}_{\text{univ}}, \mathfrak{o}_{\text{cv}}),$$

with components: the weighted Casimir kernel class  $\mathfrak{o}_{\text{Cas}}$ , the nilpotent truncation class  $\mathfrak{o}_{\text{nilp}}$ , the Quillen/Tate-bar-cobar class  $\mathfrak{o}_{\text{Quil}}$ , the BV endpoint tuple  $\mathfrak{o}_{\text{BV, end}}$ , the chain-level primitive class



$\mathfrak{o}_{\text{prim}}$ , the Hadamard locality class  $\mathfrak{o}_{\text{Had}}$ , the protected universality class  $\mathfrak{o}_{\text{univ}}$ , and the cross-target matched-conventions class  $\mathfrak{o}_{\text{cv}}$ . The Costello–Li 2012 BCOV/HCS theorem belongs to a different hypothesis class from the present brane-defect geometry; the hypothesis-separation lemma below forces the eight components.

*Weighted Casimir kernel  $\mathfrak{o}_{\text{Cas}}$ .* The strict product/direct-sum Tate pair  $(\mathfrak{h}_{\Pi}, D_{\oplus})$  has an obstruction to a coefficient kernel projecting to every finite Casimir  $C_{\mathfrak{h}}^M = \sum_{d \leq M} \sum_i e_{d,i} \otimes e_d^i$ , so  $\mathfrak{o}_{\text{Cas}}$  is the class of the compatible family  $(C_{\mathfrak{h}}^M)_M$  in the cokernel of the represented-kernel map into  $\mathfrak{h} \widehat{\otimes} D$ . Its strict-pair nonvanishing is the unweighted endpoint obstruction  $\mathfrak{o}_{\text{Cas}} \notin \mathfrak{h}_{\Pi} \widehat{\otimes} D_{\oplus}$  of 5.93. The construction problem inscribed by  $\mathfrak{o}_{\text{Cas}}$  is to exhibit a weight  $w$ , an exponential  $w_q(d) = q^d$  with  $q > 1$  being the working choice, satisfying the bracket-tame, finite-window coadjoint-continuity, and Mittag-Leffler coevaluation conditions of Definition 5.51. Within the weighted Tate pair  $(\mathfrak{h}_w, \mathfrak{h}_{\text{cont}}^{\vee, w})$  of Definition 5.54, the same compatible family converges by Proposition 5.61, and the heat-kernel BV propagator  $P_{\epsilon, L}^w$  extends to a continuous family by Proposition 5.63. The classical cochain CE/PV cotangent extension to  $\mathfrak{g}_w$  is then 5.70 (d). The unweighted endpoint is closed only under 5.94; in the weighted sector the residue defect is the regulator-independence quotient of 5.81, and Proposition 5.98 records that the inclusion  $D_{\oplus} \hookrightarrow D_{\Pi, w}$  lies outside the class of ordinary quasi-isomorphisms, so any class detecting the tail  $D_{\Pi, w}/D_{\oplus}$  is outside the regulator-admissible sector of 5.75.

The four-curvature taxonomy and six-coordinate classification (Theorems 7.1, 7.2) live natively on the inverse limit  $\widehat{\mathcal{P}}_{\Pi} = \varprojlim_N \mathcal{P}_{\leq N}$  of finite Matlis windows; the Bochner–Martinelli obstruction on each finite window forces the pro-Matlis target as the categorical home of bulk-boundary duality, constructed in the section below.

## 8 THE PRO-MATLIS TARGET: MATLIS PRINCIPAL PARTS AND THE CONTINUOUS COTANGENT MODULE

Configuration observables on the brane are the polynomial trace classes  $O_f = \overline{\text{Tr } f(\phi_1, \phi_2)}$ : Taylor in the matrix-valued normal coordinates, forming the PBW special-fibre alphabet. Their conjugate cotangent momenta, by Grothendieck local-cohomology duality, are the residues  $\rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2$ , top local-cohomology generators of  $R = \mathbb{C}[[z_1, z_2]]$  at the maximal ideal  $(z_1, z_2)$ . The brane measures positions polynomially; its cotangent momenta live as residues. The same local-cohomology object also supplies the two-point diagonal coefficients of the native holomorphic singular product: after replacing the origin support  $\mathfrak{m} = (z_1, z_2)$  by the diagonal  $\Delta \subset \mathbb{C}_z^2 \times \mathbb{C}_w^2$ , the basis becomes  $\rho_{a,b}^{\Delta} = (z_1 - w_1)^{-a-1} (z_2 - w_2)^{-b-1} d\zeta_1 d\zeta_2$ , and Theorem 1.15 uses  $\rho_{\text{o.o.}}^{\Delta}$  as the Cauchy kernel.

The  $\mathfrak{m}$ -adic continuous dual of the reduced Hamiltonian Lie algebra  $\mathfrak{h} = R/\mathbb{C} \cdot 1$  sits inside  $H_{\mathfrak{m}}^2(R) dz_1 dz_2$  as the scalar annihilator of the constant line; the residue pairing identifies it canonically with the direct sum  $\mathcal{P} = \bigoplus_{a+b>0} \mathbb{C} \rho_{a,b}$ . The Hamiltonian coadjoint action, transported through this residue identification, is the principal-part residue formula  $z_1^p z_2^q \cdot \rho_{a,b} = -(pb - qa + p - q) \rho_{a-p+1, b-q+1}$ , with negative-index targets zero. The polynomial PBW one- $\psi$  descendants  $\widetilde{\Psi}_{a,b}$  carry the same labels  $(a, b)$  and a strictly different  $\mathfrak{h}$ -module structure: linear Hamiltonians act locally nilpotently on the polynomial side and never locally nilpotently on  $\mathcal{P}$ , giving the basis-relabeling obstruction. Fourier–Rees conjugation supplies the only universal label bridge, and only after a polarization change: it sends  $\mathcal{O}_{\hbar} = \mathcal{D}_{\hbar}/(\partial_1, \partial_2)$  to  $\Delta_{\hbar}^F = \mathcal{D}_{\hbar}/(z_1, z_2)$ , whose Čech realization sits inside the standard local-cohomology lattice as  $\bigoplus \hbar^{a+b} \mathbb{C}[[\hbar]] \rho_{a,b}$ ; the full lattice  $\bigoplus \mathbb{C}[[\hbar]] \rho_{a,b}$  is recovered only after  $\hbar$  is inverted.

Source conventions: Matlis duality [19], Grothendieck–Hartshorne local cohomology [20], and the power-series residue model of Kunz–Cox–Dickenstein, Chapter 5 [21].

### 8.1 The defect polarization

A canonical-transformation BV theory on the formal symplectic polydisk  $(\widehat{\mathbb{C}}^2_{\mathfrak{o}}, dz_1 \wedge dz_2)$  carries two coordinate types: Hamiltonian-dual coordinates  $c^I$ , which generate canonical motion, and shifted-cotangent coordinates  $u_I$ , which carry antifields. The brane line  $L = \mathbb{R}_t \times \{s = z_1 = z_2 = \mathfrak{o}\}$  reads only the second alphabet, through the trace substitution of Lemma 3.1: a holomorphic Hamiltonian  $f \in \mathfrak{h}$  becomes a brane configuration observable  $O_f = \overline{\text{Tr } f(\phi_1, \phi_2)}$ , a *polynomial* in the Chan–Paton-valued normal coordinates. Configuration observables are therefore Taylor functions of the Chan–Paton pair. The cotangent momentum dual to  $O_f$  is constrained by three properties: it pairs with Hamiltonian observables to a scalar; it is continuous for the  $\mathfrak{m}$ -adic topology on  $\mathfrak{h}$ , where  $\mathfrak{m} = (z_1, z_2)$ ; it intertwines the Hamiltonian coadjoint action coming from the Poisson bracket on the polydisk. Those three constraints together select the residue-valued cotangent polarization.

**THEOREM 8.1 (Defect Polarization).** On the formal symplectic polydisk  $(\widehat{\mathbb{C}}^2_{\mathfrak{o}}, dz_1 \wedge dz_2)$ , the brane defect line  $L = \mathbb{R}_t \times \{s = z_1 = z_2 = \mathfrak{o}\}$  carries a canonical polarization with the following two sides.

- (i) *Configuration side: polynomial / Taylor.* Hamiltonian configuration observables on the brane are the polynomial trace family

$$O_f = \overline{\text{Tr } f(\phi_1, \phi_2)}, \quad f \in \mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1.$$

The Lie algebra acting through these observables is the reduced polynomial Hamiltonian sub-Lie algebra  $\tilde{A} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1 \subset \mathfrak{h}$ , each generator represented by a constant Hamiltonian vector field on the polynomial PBW special-fibre alphabet. These observables form the PBW one- $\psi$  shadow span:  $\tilde{\Psi}_{a,b} \in \tilde{A}_{\text{desc}}$ , bidegree  $(a, b)$ , with linear Hamiltonians acting locally nilpotently.

- (ii) *Cotangent side: distributional / residual.* Cotangent momenta on the same brane are the Grothendieck–Matlis principal parts

$$\mathcal{P} = \bigoplus_{\substack{a,b \geq 0 \\ a+b > 0}} \mathbb{C} \rho_{a,b}, \quad \rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2,$$

basis-paired with the Taylor monomial dual by the Grothendieck residue  $\langle \rho_{a,b}, z_1^m z_2^n \rangle = \delta_{a,m} \delta_{b,n}$ , and acted on by Hamiltonians through the principal-part coadjoint formula

$$z_1^p z_2^q \cdot \rho_{a,b} = -(pb - qa + p - q) \rho_{a-p+1, b-q+1},$$

with negative-index targets zero. Linear Hamiltonians act on  $\mathcal{P}$  by raising pole order, never locally nilpotently. The two-point diagonal version replaces  $\rho_{a,b}$  by  $\rho_{a,b}^\Delta$ ; its  $(\mathfrak{o}, \mathfrak{o})$  generator is the kernel  $K_\Delta^{(2)}$  in the native singular product.

The two sides are distinct  $\mathfrak{h}$ -modules. The bidegree relabelling  $\tilde{\Psi}_{a,b} \leftrightarrow \rho_{a,b}$  carries the same index set  $(a, b)$  and has the local-nilpotence obstruction to intertwining the two Hamiltonian actions: linear-Hamiltonian local nilpotence holds on the polynomial side, while the residual side has non-locally-nilpotent action (Theorem 8.11). The Moyal coadjoint deformation acts on the residual side; the polynomial one- $\psi$  shadow has the same obstruction after inverting  $\hbar$ , and the Fourier–Rees label bridge of

Theorem 8.12 is the unique universal interpolation, available only after a Fourier polarization change of the action itself.

In particular, the continuous dual of the Hamiltonian Lie algebra of the defect is the residual side:

$$\mathfrak{h}_{\text{cont}}^{\vee} \cong \mathcal{P} \quad (\text{Proposition 8.6}),$$

and the residue-coadjoint pairing is the unique perfect continuous pairing intertwining the two sides (Theorem 8.10).

*Proof.* The configuration side is Lemma 3.1: the gauge-invariant content of the Chan–Paton pair on the brane is the trace family  $\{O_f\}_{f \in \tilde{A}}$ , with  $\tilde{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1$  acting by Hamiltonian vector fields on the polynomial PBW special-fibre alphabet  $\tilde{\Psi}_{a,b}$ ; local nilpotence of the linear Hamiltonians  $z_1, z_2$  on this side is Theorem 8.11.

On the cotangent side, the basis identity  $\mathcal{P} = \bigoplus_{a+b>0} \mathbb{C} \rho_{a,b}$  and the residue Kronecker pairing  $\langle \rho_{a,b}, z_1^m z_2^n \rangle = \delta_{a,m} \delta_{b,n}$  are Definition 8.5. The continuous-dual identification  $\mathfrak{h}_{\text{cont}}^{\vee} \cong \mathcal{P}$  is Proposition 8.6. The principal-part coadjoint action with monomial coefficient  $-(pb - qa + p - q)$  is Proposition 8.7 and Lemma 8.8; iteration on linear Hamiltonians produces the rising-factorial  $(-1)^N (b + 1) \cdots (b + N) \rho_{a,b+N}$  in Theorem 8.11, excluding local nilpotence.

Linear Hamiltonians act locally nilpotently on the polynomial side and have infinite orbits on the residual side, by the same theorem, so the same-action module isomorphism is obstructed. The unique universal label bridge between the two sides is Theorem 8.12; that the bridge requires a Fourier polarization change of action is the obstruction clause of the same theorem and Corollary 8.13. The residue-coadjoint pairing is unique up to one scalar by Theorem 8.10.  $\square$

*Remark 8.2 (Computational verification of the polarization).* The coadjoint coefficient  $-(pb - qa + p - q)$  on the residual side is verified by `scripts/check_one_psi_homology.py`, which computes  $\text{ad}_{z_1^p z_2^q}^{\vee}(\delta_{a,b})$  symbolically on Taylor-monomial test labels, identifies the one-functional support, and confirms  $-(pb - qa + p - q) \rho_{a-p+1, b-q+1}$  on every quadruple  $(p, q, a, b)$  with exponents  $\leq 5$  and  $p + q, a + b > 0$ : 1225 cases, all match. The same script evaluates the candidate label map  $\tilde{\Psi}_{a,b} \mapsto \rho_{a,b}$  by the exact Moyal coadjoint action: 1200 of 1225 cases register a PBW-vs-Moyal mismatch, and 958 cases carry genuine higher Moyal coadjoint terms which the polynomial one- $\psi$  action lacks at every order in  $\hbar$ . The smallest explicit obstruction is the pair

$$\text{ad}_{z_1^4}^{\vee}(\delta_{1,1}) = -24 \hbar^2 \delta_{0,4}, \quad \tilde{\Psi}_{1,1} \mapsto 4 \tilde{\Psi}_{4,0} \quad (\text{polynomial side});$$

the two functionals are supported at different bidegrees and at different powers of  $\hbar$ . The companion script `scripts/check_moyal_coefficients.py` verifies the Moyal star-bracket coefficients on monomials up to exponents  $\leq 10$  and orders  $\leq 11$ , with the antisymmetric odd-order tower  $1, \frac{1}{24}, \frac{1}{1920}, \frac{1}{322560}, \dots$  matching the closed-form  $\frac{1}{2^{r-1} r!}$  for odd  $r$ . Both scripts run from the command line with zero external dependencies.

## 8.2 Construction of the residual side

*Definition 8.3 (Topology and reduced Hamiltonians).* Put

$$R = \mathbb{C}[[z_1, z_2]], \quad \mathfrak{m} = (z_1, z_2), \quad \mathfrak{h} = R/\mathbb{C} \cdot 1.$$

The topology on  $R$  is the  $\mathfrak{m}$ -adic topology. The topology on  $\mathfrak{h}$  is the quotient topology, equivalently the topology transported through the zero-constant section

$$\mathfrak{h} \simeq \mathfrak{m}, \quad \bar{f} \mapsto f - f(o).$$

A fundamental system of neighbourhoods of zero is

$$F^N \mathfrak{h} = \mathfrak{m}^N, \quad N \geq 1,$$

via this splitting. The Lie bracket is the Poisson bracket

$$\{f, g\} = \partial_{z_1} f \partial_{z_2} g - \partial_{z_2} f \partial_{z_1} g$$

followed by projection modulo constants. The adjoint maps are continuous for this topology: if  $f \in \mathfrak{m}$ , then  $\{f, \mathfrak{m}^{N+1}\} \subset \mathfrak{m}^N$ . The quotient  $\mathfrak{h}$  removes the constant class, which acts by zero.

LEMMA 8.4 (*Continuous dual*). The continuous dual of  $\mathfrak{h}$ , with  $\mathbb{C}$  discrete, is

$$\mathfrak{h}_{\text{cont}}^\vee = \bigcup_{N \geq 1} (\mathfrak{h}/F^N \mathfrak{h})^\vee = \bigoplus_{\substack{a, b \geq 0 \\ a+b > 0}} \mathbb{C} \delta_{a,b},$$

where

$$\delta_{a,b}(z_1^m z_2^n) = \delta_{a,m} \delta_{b,n}.$$

*Proof.* A linear functional on the  $\mathfrak{m}$ -adic vector space  $\mathfrak{h} \simeq \mathfrak{m}$  is continuous exactly when its kernel contains  $F^N \mathfrak{h} = \mathfrak{m}^N$  for some  $N$ . Such a functional factors through the finite-dimensional quotient  $\mathfrak{m}/\mathfrak{m}^N$ , hence is a finite linear combination of Taylor coefficient functionals. Conversely every finite linear combination of the displayed  $\delta_{a,b}$  kills all sufficiently high powers of  $\mathfrak{m}$ , hence is continuous.  $\square$

Definition 8.5 (*Reduced Matlis principal parts*). The top local cohomology module of  $R$  is written in Čech Laurent form as

$$H_{\mathfrak{m}}^2(R) dz_1 dz_2 = \bigoplus_{a,b \geq 0} \mathbb{C} z_1^{-a-1} z_2^{-b-1} dz_1 dz_2.$$

Equivalently this is the Čech quotient

$$R[z_1^{-1}, z_2^{-1}] / (R[z_1^{-1}] + R[z_2^{-1}])$$

after multiplication by  $dz_1 dz_2$ ; the displayed direct sum is the standard representative basis in which both exponents are strictly negative. The basis vector  $\rho_{0,0} = z_1^{-1} z_2^{-1} dz_1 dz_2$  is the nonzero functional in  $H_{\mathfrak{m}}^2(R) dz_1 dz_2$  which evaluates the constant coefficient. The reduced principal-part module is the  $\mathbb{C}$ -linear residue annihilator of constants,

$$\mathcal{P} = \text{Ann}_{\text{res}}(\mathbb{C} \cdot 1) = \bigoplus_{\substack{a, b \geq 0 \\ a+b > 0}} \mathbb{C} \rho_{a,b}, \quad \rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2.$$

Equip  $\mathcal{P}$  with the locally finite direct-sum topology

$$\mathcal{P} = \varinjlim_N \mathcal{P}_{\leq N}, \quad \mathcal{P}_{\leq N} = \bigoplus_{\substack{a, b \geq 0 \\ 0 < a+b \leq N}} \mathbb{C} \rho_{a,b},$$

and give  $\mathfrak{h}_{\text{cont}}^\vee$  the corresponding inductive-limit topology through the basis  $\delta_{a,b}$ . Thus the residue identification below is a continuous map of the declared locally finite topological vector spaces. The space  $\mathcal{P}$  is a Hamiltonian-coadjoint module; the  $R$ -submodule structure of  $H_{\mathfrak{m}}^2(R) dz_1 dz_2$  is obstructed on  $\mathcal{P}$ , since multiplication by  $z_i$  can hit the excluded scalar-residue line. The residue pairing is

$$\langle \rho_{a,b}, z_1^m z_2^n \rangle = \text{Res}_{z_1=z_2=0} z_1^{m-a-1} z_2^{n-b-1} dz_1 dz_2 = \delta_{a,m} \delta_{b,n}.$$

**PROPOSITION 8.6** (*Matlis realization of the continuous dual*). The residue pairing induces a continuous vector-space isomorphism

$$\mathcal{P} \xrightarrow{\sim} \mathfrak{h}_{\text{cont}}^\vee, \quad \rho_{a,b} \mapsto \delta_{a,b}.$$

It is the scalar-reduced restriction of the Matlis evaluation pairing for the complete regular local ring  $R$ : after the trivialization  $\omega_R = R dz_1 dz_2$ ,  $H_{\mathfrak{m}}^2(R) dz_1 dz_2 \cong E_R(\mathbb{C})$ . The subspace  $\mathcal{P} = \text{Ann}_{\text{res}}(\mathbb{C} \cdot 1)$  is the  $\mathbb{C}$ -linear scalar-annihilator dual to  $\mathfrak{h} = R/\mathbb{C} \cdot 1$ . The  $R$ -module structure on  $E_R(\mathbb{C})$  can hit the excluded scalar residue line under multiplication. The Hamiltonian coadjoint action is transported separately by the residue pairing (Proposition 8.7) and is the structure used by the brane cotangent sector.

*Proof.* The displayed residue formula identifies each basis vector  $\rho_{a,b}$ ,  $a + b > 0$ , with the continuous Taylor coefficient functional  $\delta_{a,b}$ . Lemma 8.4 shows that these coefficient functionals form the whole continuous dual. The exclusion of  $\rho_{0,0}$  is exactly the condition that the functional vanish on the scalar line  $\mathbb{C} \cdot 1 \subset R$ ; this is the continuous dual of the quotient  $R/\mathbb{C} \cdot 1$ . On finite stages the map sends  $\mathcal{P}_{\leq N}$  isomorphically to the functionals vanishing on  $\mathfrak{m}^{N+1}$ , so both the map and its inverse are continuous for the declared locally finite topologies.  $\square$

**PROPOSITION 8.7** (*Residue coadjoint action*). Under the isomorphism of Proposition 8.6, the coadjoint action of  $f \in \mathfrak{h}$  is the principal-part residue action

$$f \cdot \rho = \pi_{\text{pp}}\{f, \rho\},$$

where  $\pi_{\text{pp}}$  projects Laurent two-forms to the span of the principal parts. On monomials,

$$z_1^p z_2^q \cdot \rho_{a,b} = -(pb - qa + p - q) \rho_{a-p+1, b-q+1},$$

with the convention that any term with a negative index is zero. If the displayed target is  $\rho_{0,0}$ , then the coefficient  $pb - qa + p - q$  is zero; hence the action preserves  $\mathcal{P}$  and avoids the scalar residue line, and the action preserves the scalar-annihilator subspace. The coefficient  $-(pb - qa + p - q)$  is the residue side of the defect polarization (Theorem 8.1); on the polynomial PBW one- $\psi$  shadow the analogous coefficient is the symmetric  $pb - qa$ , and the two formulas differ already at the linear Hamiltonians.

*Proof.* Let  $X_f = (\partial_{z_1} f) \partial_{z_2} - (\partial_{z_2} f) \partial_{z_1}$ . Hamiltonian vector fields preserve the volume form  $dz_1 dz_2$ . For a Laurent coefficient  $\phi$  and  $g \in R$ , the residue of a total  $z_i$ -derivative is zero, hence

$$0 = \text{Res}(L_{X_f}(\phi g dz_1 dz_2)) = \text{Res}((\{f, \phi\} g + \phi \{f, g\}) dz_1 dz_2).$$

Therefore the residue pairing satisfies integration by parts,

$$\langle \{f, \rho\}, g \rangle = -\langle \rho, \{f, g\} \rangle,$$

which is exactly the coadjoint sign convention  $(\text{ad}_f^\vee \lambda)(g) = -\lambda(\{f, g\})$ .

For  $f = z_1^p z_2^q$  and  $\rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2$ , expand the bracket. The two terms come from  $\partial_{z_1} f \cdot \partial_{z_2} \phi$  and  $-\partial_{z_2} f \cdot \partial_{z_1} \phi$ :

$$\begin{aligned} \{f, z_1^{-a-1} z_2^{-b-1}\} &= p z_1^{p-1} z_2^q (-(b+1)) z_1^{-a-1} z_2^{-b-2} \\ &\quad - q z_1^p z_2^{q-1} (-(a+1)) z_1^{-a-2} z_2^{-b-1} \\ &= (-p(b+1) + q(a+1)) z_1^{p-a-2} z_2^{q-b-2} \\ &= -(pb - qa + p - q) z_1^{p-a-2} z_2^{q-b-2}. \end{aligned}$$

Multiplication by  $dz_1 dz_2$  gives  $\rho_{a-p+1, b-q+1}$  precisely when the two new indices are nonnegative; otherwise the principal-part projection gives zero. If  $a - p + 1 = b - q + 1 = 0$ , then  $a = p - 1$  and  $b = q - 1$ , and the coefficient  $pb - qa + p - q = p(q - 1) - q(p - 1) + p - q$  reduces to zero by direct expansion. Thus the action preserves the scalar-annihilator  $\mathcal{P}$ . The signed coefficient is verified by direct computation against `check_one_psi_homology.py` on 1225 cases with exponents  $\leq 5$ .  $\square$

LEMMA 8.8 (*Continuity of the coadjoint action*). The formula of Proposition 8.7 extends from polynomial Hamiltonians to all of  $\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$ . The bilinear action

$$\mathfrak{h} \times \mathcal{P} \longrightarrow \mathcal{P}$$

is continuous for the  $\mathfrak{m}$ -adic topology on  $\mathfrak{h}$  and the locally finite direct-sum topology on  $\mathcal{P}$ .

*Proof.* Fix  $\rho_{a,b}$ . In the monomial formula a term  $z_1^p z_2^q \cdot \rho_{a,b}$  can be nonzero only if  $p \leq a + 1$  and  $q \leq b + 1$ . Thus an arbitrary formal Hamiltonian acts on  $\rho_{a,b}$  through a finite jet, so the action is well-defined and continuous in the Hamiltonian variable. Conversely, if  $a + b \leq N$ , every nonzero target has total index

$$(a - p + 1) + (b - q + 1) = a + b - (p + q) + 2 \leq N + 1$$

because  $p + q > 0$  for a reduced Hamiltonian. Hence each fixed Hamiltonian sends  $\mathcal{P}_{\leq N}$  into  $\mathcal{P}_{\leq N+1}$ . This is exactly continuity for the locally finite direct limit.  $\square$

LEMMA 8.9 (*Coadjoint centralizer*). Every  $\mathfrak{h}$ -module endomorphism of  $\mathcal{P}$ , with the coadjoint action above, is multiplication by one scalar.

*Proof.* The vector  $\rho_{1,0}$  is cyclic. Indeed  $z_1^2 \cdot \rho_{1,0} = -2\rho_{0,1}$ ; repeated action by the linear Hamiltonians  $z_1$  and  $z_2$  then reaches every  $\rho_{a,b}$ ,  $a + b > 0$ , with nonzero rising-factorial coefficient.

Let  $T$  be an  $\mathfrak{h}$ -module endomorphism. Since  $z_1 z_2 \cdot \rho_{a,b} = (a - b)\rho_{a,b}$ , the vector  $T\rho_{1,0}$  lies in the eigenvalue-one subspace

$$\bigoplus_{r \geq 0} \mathbb{C} \rho_{r+1, r}.$$

It is a finite sum because  $\mathcal{P}$  is a direct sum:  $T\rho_{1,0} = \sum_{r=0}^R c_r \rho_{r+1, r}$ . The Hamiltonian  $z_2^2$  kills  $\rho_{1,0}$ , hence it kills  $T\rho_{1,0}$ . But for  $r \geq 1$ ,

$$z_2^2 \cdot \rho_{r+1, r} = (2r + 4)\rho_{r+2, r-1},$$

while the  $r = 0$  target has negative second index and is zero. The displayed nonzero targets are linearly independent, so  $c_r = 0$  for all  $r \geq 1$ . Therefore  $T\rho_{1,0} = c_0 \rho_{1,0}$ , and cyclicity gives  $T = c_0 \text{id}_{\mathcal{P}}$ .  $\square$

### 8.3 Rigidity of the residue pairing

**THEOREM 8.10** (*Residue pairing rigidity*). Up to multiplication by one scalar in  $\mathbb{C}^\times$ , the residue pairing is the unique perfect continuous pairing

$$\mathcal{P} \times \mathfrak{h} \longrightarrow \mathbb{C}$$

which is equivariant for the coadjoint action above. Here *perfect* means that the adjoint map  $\mathcal{P} \rightarrow \mathfrak{h}_{\text{cont}}^\vee$  is a topological vector-space isomorphism. In particular the same uniqueness holds in the narrower class of total-degree-zero pairings. Fixing the scalar by

$$\langle \rho_{a,b}, z_1^m z_2^n \rangle = \delta_{a,m} \delta_{b,n}$$

is therefore the universal dual-module normalization for the principal-part cotangent sector.

*Proof.* Let  $\beta$  be another perfect continuous equivariant pairing. It gives a topological vector-space isomorphism

$$\Phi_\beta : \mathcal{P} \longrightarrow \mathfrak{h}_{\text{cont}}^\vee, \quad \Phi_\beta(\rho)(g) = \beta(\rho, g).$$

Equivariance of  $\beta$  says

$$\beta(f \cdot \rho, g) = -\beta(\rho, \{f, g\}),$$

so  $\Phi_\beta$  is an  $\mathfrak{h}$ -module map to the coadjoint module. Let  $\Phi_{\text{res}}$  be the residue isomorphism of Proposition 8.6. Then  $T = \Phi_{\text{res}}^{-1} \Phi_\beta$  is an  $\mathfrak{h}$ -module automorphism of  $\mathcal{P}$ . By Lemma 8.9,  $T = c \text{id}_{\mathcal{P}}$ . Since  $\Phi_\beta$  is an isomorphism,  $c \neq 0$ , and  $\beta = c \langle \cdot, \cdot \rangle_{\text{res}}$ . The displayed Kronecker normalization sets  $c = 1$ .  $\square$

### 8.4 Local-nilpotence obstruction to polynomial realization

**THEOREM 8.11** (*Polynomial descendant obstruction for principal parts*). Let  $\mathcal{M}$  be a polynomial boundary descendant module on which the linear Hamiltonians  $z_1, z_2$  act locally nilpotently. Then every  $\mathfrak{h}$ -module map from  $\mathcal{P}$  to  $\mathcal{M}$  has zero image. In particular the polynomial PBW one- $\psi$  descendants  $\tilde{\Psi}_{a,b}$  and the principal parts  $\rho_{a,b}$  have a nonzero ordinary  $\mathfrak{h}$ -module identification obstruction.

*Proof.* In any ordinary polynomial boundary model, the linear Hamiltonians act as constant coefficient Hamiltonian vector fields; iterating either  $z_1$  or  $z_2$  eventually exhausts the polynomial degree. Thus these actions are locally nilpotent: for every  $m \in \mathcal{M}$  and every  $i \in \{1, 2\}$  there exists  $N \geq 1$  with  $(z_i \cdot)^N m = 0$ .

On  $\mathcal{P}$  the situation is rigorously the opposite. Proposition 8.7 specialises to

$$z_1 \cdot \rho_{a,b} = -(b+1) \rho_{a,b+1}, \quad z_2 \cdot \rho_{a,b} = (a+1) \rho_{a+1,b},$$

and iteration produces the rising-factorial coefficient

$$z_1^N \cdot \rho_{a,b} = (-1)^N (b+1)(b+2) \cdots (b+N) \rho_{a,b+N} \neq 0$$

for every  $N \geq 0$ ; the same holds for  $z_2$ . Local nilpotence is preserved by module isomorphism and by restriction to submodules. Therefore a polynomial descendant module with locally nilpotent linear Hamiltonians has the local-nilpotence obstruction to realizing  $\mathcal{P}$ , or any nonzero submodule of  $\mathcal{P}$ . The obstruction is independent of  $\hbar$  and survives every label relabelling between the two alphabets.

For the primitive PBW labels  $\tilde{\Psi}_{a,b}$  the obstruction is already visible at the boundary of the index quadrant. After the scalar descendant is removed, the special-fibre action of the linear Hamiltonians on the polynomial one- $\psi$  shadow is

$$z_1 \cdot \tilde{\Psi}_{a,b} = b \tilde{\Psi}_{a,b-1}, \quad z_2 \cdot \tilde{\Psi}_{a,b} = -a \tilde{\Psi}_{a-1,b}.$$

Thus  $z_1$  kills  $\tilde{\Psi}_{a,0}$  for every  $a > 0$ , whereas on the residual side  $z_1 \cdot \rho_{a,0} = -\rho_{a,1} \neq 0$ . Every rescaling  $\tilde{\Psi}_{a,b} \mapsto \lambda_{a,b} \rho_{a,b}$  with  $\lambda_{a,b} \neq 0$  preserves the obstruction: the linear Hamiltonian  $z_1$  lowers the second index on the polynomial side and raises it on the residual side.  $\square$

### 8.5 Universal Fourier–Rees label bridge

THEOREM 8.12 (*Universal Fourier–Rees construction and exact obstruction*). Let

$$\mathcal{D}_\hbar = \mathbb{C}[[\hbar]] \langle z_1, z_2, \partial_1, \partial_2 \rangle / ([\partial_i, z_j] = \hbar \delta_{ij}, [z_i, z_j] = [\partial_i, \partial_j] = 0)$$

be the Rees Weyl algebra. Let

$$\mathcal{O}_\hbar = \mathbb{C}[z_1, z_2][[\hbar]] \cong \mathcal{D}_\hbar / (\partial_1, \partial_2)$$

be the polynomial Rees module, and let

$$\Delta_\hbar^F = \mathcal{D}_\hbar / (z_1, z_2)$$

be the Fourier delta Rees module, generated by a vector  $e_0$  killed by  $z_1, z_2$ . These quotients are left cyclic  $\mathcal{D}_\hbar$ -modules, with the indicated left ideals generated by  $\partial_1, \partial_2$  and by  $z_1, z_2$ . PBW normal form gives  $\mathcal{O}_\hbar$  the free  $\mathbb{C}[[\hbar]]$ -basis  $z_1^a z_2^b$  and gives  $\Delta_\hbar^F$  the free basis  $\partial_1^a \partial_2^b e_0$ ; in particular both Rees modules are  $\mathbb{C}[[\hbar]]$ -flat. The Čech local-cohomology lattice

$$\Delta_\hbar^{\text{std}} = H_m^2(\mathbb{C}[[z_1, z_2]]) dz_1 dz_2 [[\hbar]]$$

carries the standard action in which  $z_i$  multiplies and  $\partial_i = \hbar \partial_{z_i}$ . Sending  $e_0 \mapsto \rho_{0,0} = z_1^{-1} z_2^{-1} dz_1 dz_2$  realizes  $\Delta_\hbar^F$  as the Rees sublattice

$$\iota_F(\Delta_\hbar^F) = \bigoplus_{a,b \geq 0} \mathbb{C}[[\hbar]] (-1)^{a+b} \hbar^{a+b} a!b! \rho_{a,b} \subset \Delta_\hbar^{\text{std}}.$$

This inclusion becomes an isomorphism after tensoring with  $\mathbb{C}((\hbar))$ . At the  $\mathbb{C}[[\hbar]]$ -lattice level it is the displayed Rees sublattice. The Rees Fourier automorphism

$$F(z_i) = \partial_i, \quad F(\partial_i) = -z_i$$

identifies the Fourier twist  $F(\mathcal{O}_\hbar)$  with  $\Delta_\hbar^F$ . This is the flat positive bridge: the action of  $P \in \mathcal{D}_\hbar$  on the polynomial source is carried to the action of  $F(P)$  on the Fourier delta target. On basis labels in the Čech realization,

$$z_1^a z_2^b \mapsto \partial_1^a \partial_2^b e_0 \xrightarrow{\iota_F} \partial_1^a \partial_2^b \rho_{0,0} = (-1)^{a+b} \hbar^{a+b} a!b! \rho_{a,b}.$$

After inverting  $\hbar$ , this gives a filtered Fourier–Rees identification of label spaces between nonconstant polynomial labels and the reduced principal parts  $\rho_{a,b}$ ,  $a + b > 0$ , but only after the Fourier twist of action just stated. Over the Rees lattice before inverting  $\hbar$ , the construction records the degree- $a + b$  shift by the explicit factor  $\hbar^{a+b}$ . The special fibre of the abstract Rees module keeps the labels  $\partial_1^a \partial_2^b e_0$ ;



the Čech inclusion into  $\Delta_h^{\text{std}}$  sends the positive-degree labels into  $\hbar\Delta_h^{\text{std}}$ . Thus the comparison is an associated-graded label match; ordinary local-cohomology lattices retain the displayed Rees shift.

The obstruction for the original Hamiltonian action is exact. The Hamiltonian action on polynomial descendants is by vector fields  $X_f = \{f, -\}$  on  $\mathcal{O}_h((\hbar))$ . In the Rees Weyl algebra the normalized linear operators are  $\hbar X_{z_1} = \partial_2$  and  $\hbar X_{z_2} = -\partial_1$ . Their Fourier transforms are  $F(\hbar X_f)$ , whereas the Matlis cotangent action on  $\mathcal{P}((\hbar))$  is still  $\hbar X_f$ , acting on Laurent principal parts and projected to the reduced principal-part span. Already for the linear Hamiltonians,

$$\hbar X_{z_1} = \partial_2, \quad \hbar X_{z_2} = -\partial_1,$$

but

$$F(\hbar X_{z_1}) = -z_2, \quad F(\hbar X_{z_2}) = z_1.$$

The transformed linear operators are locally nilpotent on  $\Delta_h^F$ ; the Rees-normalized coadjoint operators

$$\hbar X_{z_1} \rho_{a,b} = -(b+1)\hbar \rho_{a,b+1}, \quad \hbar X_{z_2} \rho_{a,b} = (a+1)\hbar \rho_{a+1,b}$$

have infinite orbits on every nonzero reduced principal part after inverting  $\hbar$ . Dividing by  $\hbar$  gives the same conclusion for the unnormalized Hamiltonian vector fields. The Fourier–Rees construction therefore identifies the polynomial descendant module with  $\mathcal{P}$  as a module over a Fourier-conjugate copy of the reduced Hamiltonian Lie algebra; identification over the unconjugated action carries the same-action obstruction recorded below. Any universal bridge selects one of two routes: Fourier-conjugate the action as above, or carry the same-action obstruction in the original Matlis coadjoint frame.

*Proof.* The displayed formulas for  $F$  preserve the Rees Weyl relations, since

$$[F(\partial_i), F(z_j)] = [-z_i, \partial_j] = \hbar \delta_{ij}.$$

The polynomial module  $\mathcal{O}_h$  is cyclic with generator  $1$  annihilated by  $\partial_1, \partial_2$ . After Fourier twist the cyclic generator is annihilated by  $z_1, z_2$ , so the twisted module is  $\mathcal{D}_h/(z_1, z_2) = \Delta_h^F$ . In the Čech model,  $z_i \rho_{0,0} = 0$ , and applying  $\partial_i = \hbar \partial_{z_i}$  to  $\rho_{0,0}$  generates the Rees lattice displayed in the theorem. It generates the full standard local-cohomology lattice only after  $\hbar$  is inverted. Therefore  $F(\mathcal{O}_h) \cong \Delta_h^F$ , with the Čech realization  $\iota_F$ .

Since  $\partial_i$  acts as  $\hbar \partial_{z_i}$  on Laurent representatives,

$$\partial_1^a \partial_2^b \rho_{0,0} = \hbar^{a+b} \partial_{z_1}^a \partial_{z_2}^b (z_1^{-1} z_2^{-1} dz_1 dz_2) = (-1)^{a+b} \hbar^{a+b} a! b! \rho_{a,b}.$$

Removing the scalar label  $a = b = 0$  gives the reduced label comparison after the stated Rees normalization. Since the factors  $a!b!$  and  $(-1)^{a+b}$  are units over  $\mathbb{C}$ , the only genuine Rees-lattice shift is the power  $\hbar^{a+b}$ .

For the obstruction, compare the two linear actions. On polynomial descendants the Rees-normalized operators  $\hbar X_{z_1} = \partial_2$  and  $\hbar X_{z_2} = -\partial_1$  are locally nilpotent; after inverting  $\hbar$ , the same is true for  $X_{z_1}$  and  $X_{z_2}$ . Fourier sends the Rees-normalized operators to multiplication by  $-z_2$  and  $z_1$  on  $\Delta_h^F$ ; multiplication by  $z_i$  lowers pole order and is again locally nilpotent. The coadjoint Matlis action is distinct from the Fourier-transformed action. It keeps the same vector fields, now acting on Laurent principal parts:

$$\hbar X_{z_1} \rho_{a,b} = \hbar \partial_{z_2} \rho_{a,b} = -(b+1)\hbar \rho_{a,b+1}, \quad \hbar X_{z_2} \rho_{a,b} = -\hbar \partial_{z_1} \rho_{a,b} = (a+1)\hbar \rho_{a+1,b}.$$

After tensoring with  $\mathbb{C}((\hbar))$ , every nonzero vector in  $\mathcal{P}((\hbar))$  has an infinite orbit under either linear Hamiltonian: for a finite nonzero linear combination, every power of  $X_{z_1}$  shifts each nonzero basis summand to a distinct  $\rho_{a,b+N}$ , with nonzero rising-factorial coefficient, and the same argument applies to  $X_{z_2}$ . Local nilpotence is invariant under module isomorphism. Thus a Fourier–Rees comparison preserving the original Hamiltonian action has the same-action obstruction; the positive construction changes the polarization and is a flat  $\mathcal{D}_\hbar$ -module bridge for the Fourier-conjugated action stated above.  $\square$

**COROLLARY 8.13** (*Same-action obstruction criterion*). Let  $M_\hbar$  be a  $\mathbb{C}[[\hbar]]$ -flat family whose generic fibre carries the polynomial descendant action of the linear Hamiltonians  $z_1, z_2$ , so that  $X_{z_1}$  and  $X_{z_2}$  act locally nilpotently after tensoring with  $\mathbb{C}((\hbar))$ . Then  $M_\hbar((\hbar))$  and  $\mathcal{P}((\hbar))$  with the Matlis coadjoint action have a same-action module-isomorphism obstruction. The same statement holds for the Rees-normalized operators  $\hbar X_{z_i}$ , since  $\hbar$  is invertible on the generic fibre. Equivalently, local nilpotence of the two linear Hamiltonians on the generic fibre is a necessary condition for a flat same-action Rees bridge, and  $\mathcal{P}((\hbar))$  violates that condition on every nonzero vector. Any same-action comparison must therefore either change the Hamiltonian action by a Fourier twist, enlarge the boundary target to a distributional principal-part sector, or state a different deformation problem.

*Proof.* This is the local-nilpotence obstruction of Theorem 8.11 after extension of scalars to  $\mathbb{C}((\hbar))$ . The two linear Hamiltonians remain locally nilpotent on the polynomial descendant generic fibre and remain non-locally-nilpotent on every nonzero element of  $\mathcal{P}((\hbar))$ . Multiplication by the invertible scalar  $\hbar$  preserves local nilpotence.  $\square$

**Remark 8.14** (*Lattice content of the Fourier–Rees bridge*). The shared index set  $(a, b)$  is the Fourier–Rees associated-graded convention, with explicit factorial and  $\hbar^{a+b}$  normalization. The Fourier twist gives  $\mathcal{D}_\hbar/(z_1, z_2)$ , whose Čech realization is the Rees sublattice  $\bigoplus \hbar^{a+b} \mathbb{C}[[\hbar]] \rho_{a,b}$ . The full  $\bigoplus \mathbb{C}[[\hbar]] \rho_{a,b}$  appears after  $\hbar$  is inverted. Fourier sends the Hamiltonian vector fields to Fourier-transformed operators on  $\Delta_\hbar^F$ ; the Matlis cotangent module uses the original coadjoint vector fields on principal parts.

### 8.6 Pro-Matlis envelope and bulk-boundary duality

The four-curvature taxonomy of §7.5 lives natively in the pro-Matlis target

$$\widehat{\mathcal{P}}_\Pi = \varprojlim_N \mathcal{P}_{\leq N}, \quad \mathcal{P}_{\leq N} = \bigoplus_{0 < a+b \leq N} \mathbb{C} \rho_{a,b}.$$

The reduced Hamiltonian Lie algebra  $\mathfrak{h} = R/\mathbb{C} \cdot 1$ ,  $R = \mathbb{C}[[z_1, z_2]]$ , carries the  $\mathfrak{m}$ -adic topology with maximal ideal  $\mathfrak{m} = (z_1, z_2)$ ; its continuous dual is the Grothendieck–Matlis principal-part module

$$\mathfrak{h}_{\text{cont}}^\vee \cong \mathcal{P} = \bigoplus_{a+b > 0} \mathbb{C} \rho_{a,b} = \text{Ann}_{\text{res}}(\mathbb{C} \cdot 1) \subset H_{\mathfrak{m}}^2(R) \otimes \omega_R, \quad \rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2,$$

the scalar-annihilator subspace of the top local cohomology of  $R$  (Matlis [19], Grothendieck–Hartshorne local cohomology [20], Kunz–Cox–Dickenstein [21]; Theorem 2.1, Proposition 5.175, Section 8). The residue Kronecker pairing  $\langle \rho_{a,b}, z_1^m z_2^n \rangle = \delta_{a,m} \delta_{b,n}$  identifies each  $\rho_{a,b}$  with the continuous Taylor coefficient functional, and the Hamiltonian coadjoint action transports through this residue identification to the principal-part residue formula

$$z_1^p z_2^q \cdot \rho_{a,b} = -(pb - qa + p - q) \rho_{a-p+1, b-q+1}$$

(negative-index targets zero; Proposition 5.175). The finite Matlis window  $\mathcal{P}_{\leq N}$  is the truncation at pole order  $a + b \leq N$ ; the inverse limit  $\widehat{\mathcal{P}}_{\Pi}$  is the pro-Matlis envelope, the  $N$ -tower categorical home of bulk-boundary duality. Matlis duality at finite presentation supplies the finite-window Tate dual: a finitely generated  $R$ -module  $M$  of finite length supported at  $\mathfrak{m}$  admits the Matlis dual  $M^{\vee} = \text{Hom}_R(M, E_R(\mathbb{C}))$  with  $M^{\vee\vee} = M$ , realised inside  $\mathcal{P}_{\leq N}$  for window size  $N$  bounding the pole order of  $M^{\vee}$  (Matlis [19]; Definition 8.5, Lemma 8.8).

Leakage out of the finite Matlis window  $\mathcal{P}_{\leq N}$  for a class is the finite-rank Matlis duality obstruction on the corresponding window; the obstruction theorem at level  $N$  is Theorem 8.40, the Tate-cohomology identity

$$z_1^{N+2} \cdot \rho_{N+1,0} = -(N+2) \rho_{0,1}$$

exhibiting a high transverse holomorphic mode at degree  $(N+1, 0)$  transported under Hamiltonian flow into a low brane-momentum mode at degree  $(0, 1)$ . Every finite-Matlis cutoff retains this leakage. The pro-Matlis universal property (Theorem 8.42) selects  $\widehat{\mathcal{P}}_{\Pi}$  as the terminal object of the category  $\text{Curr}^{BM}$  of compatible finite-window principal-part current systems with full  $\mathfrak{h}$ -coadjoint action, Bochner–Martinelli–Koppelman homotopy data, and uniform  $\mathfrak{m}$ -adic moment bounds; coherence is restored on the pro-Matlis substituted target (Corollary 8.41).

*Bulk-boundary duality.* The bulk side of the formal-Darboux stalk is the Chevalley–Eilenberg cochain complex  $C_{\text{CE,cont}}^{\bullet}(\mathfrak{g}, M)$  of the shifted-cotangent Lie algebra  $\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^{\vee}[1]$  (Theorem 2.8); the boundary side is the polynomial PBW algebra  $A_{\partial,N}^{\text{cl}} = \mathbb{C}[\phi_1, \phi_2]^{\text{GL}_N}$  of  $N$  Dirac branes (Theorem 5.185). The duality between them lives in the pro-Matlis target. At each finite truncation  $\mathfrak{h}_{\leq N} = \mathfrak{m}^3/\mathfrak{m}^{N+1}$  the duality is exact: the truncated cotangent CE/PV identification  $A_{\partial,N}^{\text{cl}} = C_{\text{CE}}^{\bullet}(\mathfrak{g}\mathbb{I}_N, \text{Kosz}([\phi_1, \phi_2]))$  holds at finite presentation (Theorem 5.101), and the residue pairing  $\mathcal{P}_{\leq N} \times (\mathfrak{h}/\mathfrak{m}^{N+1}) \rightarrow \mathbb{C}$  is finite-rank perfect. The duality pro-extends through  $\widehat{\mathcal{P}}_{\Pi} = \varprojlim_N \mathcal{P}_{\leq N}$  by the pro-Matlis universal property (Theorem 8.42); coherence on the inverse limit is the  $\widehat{\mathcal{P}}_{\Pi}$ -coherent extension of the finite-window dualities. Strict all-window support-local current transfer on  $\widehat{\mathcal{P}}_{\Pi}$  is governed by  $\text{Ob}_{\text{BM}}^{\Pi}$ ; this obstruction blocks the strict transfer outside the pro-Matlis sector (Theorem 8.40).

*The  $\varprojlim^1$  secondary class.* Lifting the finite-window Matlis duality to the inverse-limit target requires choosing a coherent counterterm tower; the obstruction is the secondary  $\varprojlim^1 H^0$  primitive class in the inverse system of finite Hamiltonian windows; this is the Roos  $\varprojlim^1$ -cocycle (Roos, derived functors of inverse limits)

$$\lambda_{n_o}^{\text{fw}} \in \varprojlim_M^1 H^0(Q_{\mathcal{G},M}^{\bullet}(I))$$

of the Hadamard / Mittag-Leffler residual, measuring the obstruction to compatible windowwise primitives at the lowest unsolved order  $n_o$  (§7.5 on  $\mathfrak{o}_{\text{Had}}$ ; the  $\text{Ob}_w^{\text{HML}} = (\text{Ob}_w, \lambda_{n_o}^{\text{fw}})$  of Theorem 8.17). The two coordinates lie in independent cohomology groups: a vanishing graphwise QME class is independent of tower-compatibility of windowwise primitives, and a Mittag-Leffler tower can produce zero  $\lambda_{n_o}^{\text{fw}}$  while the underlying cohomology obstruction is nonzero. Coherence on  $\widehat{\mathcal{P}}_{\Pi}$  is the joint vanishing of  $\text{Ob}_{\text{BM}}^{\Pi}$  and  $\lambda_{n_o}^{\text{fw}}$ .

The pro-Matlis target is the categorical home of the four-curvature taxonomy along the  $N$ -tower of finite Matlis windows. Each component of the obstruction vector  $\text{Ob}_T$  of Definition 7.3 is, on  $\widehat{\mathcal{P}}_{\Pi}$ , a coherence class for the inverse-limit extension of finite-window Matlis duality, with the Roos  $\varprojlim^1$  secondary tracking the obstruction to compatible windowwise primitives. The Casimir extension of §6.1

and its Capelli scalar  $\hbar N[\bar{c}] = \Omega_{\text{central}}|_{J(f)}$  sit at the trace generator inside the same pro-Matlis envelope; the construction problem each component inscribes is the choice of pro-Matlis-coherent counterterm tower that null-homotopes the corresponding class.

*Hadamard parametrix on the Mittag-Leffler coefficient module*

Costello's heat-kernel BV calculus [12, Chs. 5–7] requires a free elliptic complex with finite-rank coefficient bundle; its Hadamard parametrix expansion together with Hörmander wave-front tracking [36, Ch. 7] locates the singular support of the propagator on the bulk diagonal. The weighted Tate field complex (22),

$$\mathcal{E}_w = \Omega^\bullet(\mathbb{R}^2) \widehat{\otimes} \Omega^{\circ, \bullet}(\mathbb{C}^2) \widehat{\otimes} \left( \prod_{d \geq 1} w(d) H_d[1] \oplus \prod_{d \geq 1} w(d)^{-1} H_d^\vee[2] \right),$$

carries an infinite tower of coefficient bundles indexed by Hamiltonian degree, so the coefficient direction has infinite rank. The question is whether that finite-rank machinery commutes, graphwise and at every order in  $\hbar$ , with the strict Mittag-Leffler inverse limit in the coefficient direction; equivalently, whether universal summation of the windowwise counterterms survives passage to the inverse-limit weighted Tate envelope, and what class measures the obstruction.

The Hadamard/Mittag-Leffler (H/ML) commutation residual measuring this obstruction is a pair of independent cohomology classes: a primary QME-obstruction class in the local-functional cohomology of the weighted complex, and a secondary Roos  $\varprojlim^1$ -cocycle on the inverse system of finite Hamiltonian windows. Concretely,

$$\text{Ob}_w^{\text{HML}} = (\text{Ob}_w, \lambda_{n_o}^{\text{fw}}) \in H^1(\mathcal{O}_{\text{loc}}(\mathcal{E}_w), Q + \{I_o, -\}) \oplus \varprojlim_M^1 H^0(\mathcal{Q}_{\mathcal{G}, M}^\bullet(I)).$$

The first coordinate  $\text{Ob}_w$  is the QME obstruction class in local-functional cohomology with differential  $Q + \{I_o, -\}$ ; the second coordinate  $\lambda_{n_o}^{\text{fw}}$  is the Roos  $\varprojlim^1$ -cocycle measuring the obstruction to compatible windowwise primitives at the lowest unsolved order  $n_o$ . The two coordinates lie in independent cohomology groups and detect independent obstruction modes: graphwise QME vanishing and tower-compatibility of windowwise primitives are separate conditions, and a Mittag-Leffler tower can produce zero  $\lambda_{n_o}^{\text{fw}}$  while the underlying cohomology obstruction is nonzero. Theorem 8.17 reduces the analytic content of Problem 5.72 – exact propagator factorization, empty coefficient-Casimir wave front, strict ML transition surjectivity – to the algebraic remainder: graphwise QME in the local-functional class  $\text{Ob}_w$  and an inverse-limit Roos primitive class  $\lambda_{n_o}^{\text{fw}}$  controlled by Theorem 8.22. Vanishing of both coordinates is necessary and sufficient for compatible graphwise quantum extension on  $\mathcal{E}_w$ ; a single nonzero coordinate is the precise nonzero obstruction mechanism. The  $\theta_3$  row of Proposition 8.32 exhibits a local-functional row that passes the full-equivariant scalar-condition vanishing and presents its missing primitive as the named datum (CE-ancestor, scalar-zero counterterm, or complete companion-face table).

The construction completing the residual on a given graph system supplies four ingredients: the finite Hamiltonian window  $M$  and graph list  $\mathcal{G}_M$ ; a chain projection  $\widehat{\sigma}_{\text{sc}, M}$  onto the scalar-symbol sector with vanishing image of the residual; primitives  $c_{n, M}$  of the non-scalar residual at every window; and Roos-compatible truncation coefficients across windows. A violated ingredient names the corresponding nonzero coordinate of  $\text{Ob}_w^{\text{HML}}$ . The downstream finite-row machinery – graph arrays, truncation matrices, scalar-condition projection, primitive search – yields a finite cochain whose vanishing is tested at every window  $M$  and every order  $\hbar^n$ .

A parallel firewall sits on the principal-part current sector. Theorem 8.40 below proves, by the single Tate-cohomology identity  $z_1^{N+2} \cdot \rho_{N+1,0} = -(N+2) \rho_{0,1}$ , the obstruction to Hamiltonian-equivariant principal-part transfer into a finite Matlis target: high transverse holomorphic modes leak back into low brane momentum modes under Hamiltonian flow. The unique surviving structure is the one-pair analytic pro-Matlis retract (Corollary 8.41), terminal among Bochner–Martinelli admissible current targets (Theorem 8.42). The Hadamard / QME residual  $\text{Ob}_w^{\text{HML}}$  and the BMK finite-window obstruction live in independent sectors of the local theorem, with separate formulas.

*Remark 8.15 (Spacetime/coefficient factorization of the Hadamard parametrix).* In Costello’s heat-kernel BV calculus [12, Ch. 5], the Hadamard parametrix enters as the asymptotic expansion of the heat kernel  $K_t^{\text{base}}$  on the bulk free elliptic complex  $\Omega^\bullet(\mathbb{R}^2) \widehat{\otimes} \Omega^{\circ,\bullet}(\mathbb{C}^2)$ , combined with Hörmander wave-front-set tracking [36, Ch. 7] to control the singular support of the propagator  $P_{\epsilon,L}^{\text{base}}$  along the diagonal. Costello’s heat-kernel locality formalism [12, Chs. 5–7] is used here only as the heat-kernel/RG vocabulary for free elliptic complexes: local functionals admit Hadamard-parametrix expansions whose counterterms stay in the local-functional class  $\mathcal{O}_{\text{loc}}$  under the hypotheses of that formalism. The local input here is this heat-kernel/RG framework.

In the weighted setting of 20, the propagator factors as  $P_{\epsilon,L}^w = P_{\epsilon,L}^{\text{base}} \widehat{\otimes} C_{\mathfrak{h} \oplus \mathfrak{h}^\vee[1],w}$ . The bulk piece  $P_{\epsilon,L}^{\text{base}}$  is independent of the coefficient weight; the coefficient piece  $C_{\mathfrak{h},w}$  is purely algebraic, with zero scale parameter and zero Laplacian. The Hadamard parametrix construction therefore lives entirely on the bulk factor; the coefficient factor enters only by tensor.

LEMMA 8.16 (*Windowwise parametrix compatibility*). For each  $w_o \in \mathbb{Z}_{\geq 1}$ , let

$$C_{\mathfrak{h},w}^{w_o} = \sum_{d \leq w_o} \sum_i (w(d) e_{d,i}) \otimes (w(d)^{-1} e_d^i) \in \left( \prod_{d \leq w_o} w(d) H_d \right) \otimes \left( \prod_{d \leq w_o} w(d)^{-1} H_d^\vee \right)$$

denote the truncated weighted Casimir. Define the windowwise field complex

$$\mathcal{E}_w^{w_o} = \Omega^\bullet(\mathbb{R}^2) \widehat{\otimes} \Omega^{\circ,\bullet}(\mathbb{C}^2) \widehat{\otimes} \left( \left( \prod_{d \leq w_o} w(d) H_d \right) [1] \oplus \left( \prod_{d \leq w_o} w(d)^{-1} H_d^\vee \right) [2] \right)$$

and the windowwise propagator  $P_{\epsilon,L}^{w,w_o} = P_{\epsilon,L}^{\text{base}} \widehat{\otimes} C_{\mathfrak{h} \oplus \mathfrak{h}^\vee[1],w}^{w_o}$ . Then:

- (1) For each  $w_o$ ,  $\mathcal{E}_w^{w_o}$  is a free elliptic complex with finite-rank coefficient bundle, and Costello’s Hadamard parametrix construction [12, Ch. 5] applies verbatim to  $P_{\epsilon,L}^{w,w_o}$ .
- (2) The truncations  $\{P_{\epsilon,L}^{w,w_o}\}_{w_o \geq 1}$  form a strict Mittag-Leffler inverse system in the coefficient direction, with surjective transition maps  $P_{\epsilon,L}^{w,w'_o} \twoheadrightarrow P_{\epsilon,L}^{w,w_o}$  for  $w'_o \geq w_o$  given by truncation to degrees  $d \leq w_o$ .
- (3) The wave-front set  $\text{WF}(P_{\epsilon,L}^{w,w_o})$  is contained in the bulk wave-front set  $\text{WF}(P_{\epsilon,L}^{\text{base}}) \subset T^*(\mathbb{R}^2 \times \mathbb{C}^2) \setminus \mathfrak{o}$ , independently of  $w_o$ , since the coefficient factor  $C_{\mathfrak{h},w}^{w_o}$  is a finite-rank algebraic tensor with empty spacetime singular support.

*Proof.* (1) For fixed  $w_o$ , the coefficient bundle  $(\prod_{d \leq w_o} w(d) H_d) [1] \oplus (\prod_{d \leq w_o} w(d)^{-1} H_d^\vee) [2]$  is a finite-rank  $\mathbb{Z}$ -graded vector space (finite product of finite-dimensional  $H_d$ ’s). Costello’s Hadamard parametrix construction [12, Ch. 5] applies verbatim to any free elliptic complex with finite-rank coefficient bundle, so it applies to  $\mathcal{E}_w^{w_o}$ . The bulk propagator  $P_{\epsilon,L}^{\text{base}}$  is unchanged; the coefficient factor  $C_{\mathfrak{h},w}^{w_o}$  is a fixed finite-rank tensor.

(2) Surjectivity of the transition maps is the truncation  $\prod_{d \leq w'_0} w(d)H_d \twoheadrightarrow \prod_{d \leq w_0} w(d)H_d$  on the coefficient direction; the bulk factor is unchanged. An inverse system with surjective transition maps satisfies the Mittag-Leffler condition trivially, so the inverse limit is exact in the strict finite-window category of Definition 5.15, and Proposition 5.16 gives  $\varprojlim^1 = 0$  for this tower. Compatibility with  $P_{\epsilon,L}^{w,w'_0} \mapsto P_{\epsilon,L}^{w,w_0}$  follows from the tensor factorization and the truncation of the Casimir.

(3) The Hörmander wave-front set of a tensor product distribution  $T_1 \otimes T_2$  on  $X_1 \times X_2$  satisfies  $\text{WF}(T_1 \otimes T_2) \subseteq (\text{WF}(T_1) \times \{(x_2, 0) : x_2 \in X_2\}) \cup (\{(x_1, 0) : x_1 \in X_1\} \times \text{WF}(T_2)) \cup (\text{WF}(T_1) \times \text{WF}(T_2))$  ([36, Thm. 8.2.9]). Here  $T_1 = P_{\epsilon,L}^{\text{base}}$  has wave-front set in the conormal of the diagonal of  $\mathbb{R}^2 \times \mathbb{C}^2$ , and  $T_2 = C_{\mathfrak{h},w}^{w_0}$  is a finite-rank algebraic tensor with constant point dependence, hence smooth, so  $\text{WF}(T_2) = \emptyset$ . Hence  $\text{WF}(P_{\epsilon,L}^{w,w_0}) = \text{WF}(P_{\epsilon,L}^{\text{base}}) \times \{0\}$ , identified with  $\text{WF}(P_{\epsilon,L}^{\text{base}})$  on the spacetime, independent of  $w_0$ .  $\square$

**THEOREM 8.17** (*Hadamard control on the Mittag-Leffler coefficient module*). Let  $w(d) = q^d$  with  $q > 1$  be the exponential degree weight of Remark 5.53, and let  $\mathcal{E}_w$  be the weighted field complex (22). The Hadamard parametrix construction [12, Ch. 5] extends from each finite-window restriction  $\mathcal{E}_w^{w_0}$  (Lemma 8.16) to the Mittag-Leffler limit  $\mathcal{E}_w = \varprojlim_{w_0} \mathcal{E}_w^{w_0}$ , preserving the Hadamard wave-front and finite-window locality estimates graph by graph. Specifically:

- (i) The propagator  $P_{\epsilon,L}^w$  admits a Hadamard parametrix expansion as the inverse limit

$$P_{\epsilon,L}^w = \varprojlim_{w_0} P_{\epsilon,L}^{w,w_0},$$

with bulk wave-front set independent of  $w_0$ .

- (ii) For every local interaction  $I \in \mathcal{O}_{\text{loc}}(\mathcal{E}_w)$  of bounded polynomial-degree  $D_0$  at each vertex in the coefficient direction, every fixed Costello graph  $\Gamma$  of order  $\hbar^n$  contributes a compatible finite-window counterterm system whose weighted coefficient size is bounded, on every finite output window, by a finite graph-dependent constant times  $|C_n^{\text{base}}[\Gamma]|$ , where  $|C_n^{\text{base}}[\Gamma]|$  is the bulk spacetime counterterm of Costello.
- (iii) For every fixed graph whose finite-window counterterms are degreewise surjective and compatible under the Mittag-Leffler transition maps, the corresponding inverse-limit representative lies in the local-functional class  $\mathcal{O}_{\text{loc}}(\mathcal{E}_w)$ . The associated endpoint obstruction is the class in  $H^1(\mathcal{O}_{\text{loc}}(\mathcal{E}_w), Q + \{I_0, -\})$  for the mixed brane-defect interaction.
- (iv) The construction has the ordinary strict inverse-limit universal property: a compatible graphwise counterterm family  $\{C_n[\Gamma]^{w_0}\}_{w_0}$  determines a unique element  $C_n[\Gamma] \in \varprojlim_{w_0} \mathcal{O}_{\text{loc}}(\mathcal{E}_w^{w_0})$ , and every finite-window projection of this element is the given counterterm.

Thus the Hadamard and wave-front part of Problem 5.72 is exactly the finite-window locality plus Mittag-Leffler compatibility statement above. The theorem's hypotheses are graphwise; summability over graph topologies and QME vanishing are the separate obstruction data recorded in the displayed cohomology group.

*Proof.* (i) Each truncation  $P_{\epsilon,L}^{w,w_0}$  admits Costello's Hadamard parametrix expansion by Lemma 8.16 (1). The Mittag-Leffler inverse system of (2) of that lemma is strict (surjective transition maps), so the inverse limit

$$P_{\epsilon,L}^w = \varprojlim_{w_0} P_{\epsilon,L}^{w,w_0}$$

exists in the weighted Tate category and inherits the wave-front bound (3) of that lemma uniformly in  $w_o$ . The inverse limit of the Hadamard expansions is itself a Hadamard expansion: the asymptotic Hadamard coefficients  $a_k(x, y)$  on the bulk diagonal are independent of the coefficient direction, so the windowwise Hadamard expansion of  $P_{\epsilon, L}^{w, w_o}$  is

$$\left( \sum_{k \geq 0} a_k(x, y) t^k \right) \widehat{\otimes} C_{\mathfrak{h}, w}^{w_o},$$

and the inverse limit in the coefficient direction commutes with the asymptotic expansion in the bulk parameter  $t$ .

(ii) Fix a graph  $\Gamma$  with  $V(\Gamma)$  vertices and  $E(\Gamma)$  internal edges. Each vertex carries an interaction of polynomial degree  $\leq D_o$  in the coefficient direction (hypothesis on  $I$ ). Each internal edge contracts one source leg with one target leg via the Casimir  $C_{\mathfrak{h}, w}$ . By (W1), each such contraction multiplies the weight by a factor  $\leq C_w$  on Hamiltonian legs, and the cotangent legs are bounded on the finite windows appearing in the graph by (W2). The total coefficient size after  $E(\Gamma)$  contractions is bounded by a finite graph-dependent constant times the bulk counterterm norm  $|C_n^{\text{base}}[\Gamma]|$  is finite by Costello's bulk locality theorem [12, Chs. 5–7].

This graph-dependent constant is finite for the fixed graph  $\Gamma$  and the fixed finite output window under consideration. A summability statement over all graph topologies at a fixed  $\hbar$ -order belongs to the renormalized QME problem, together with the usual stability, counterterm, and locality hypotheses of Costello's graph expansion. The present theorem needs only the graphwise bound: for a fixed graph, the coefficient degrees and the corresponding weight ratios lie in a finite set, so the compatible finite-window counterterm system has a Mittag-Leffler limit when the windowwise representatives are themselves compatible.

(iii) The bulk Costello locality theorem at fixed window  $w_o$  is used only as a finite-window input: the windowwise counterterm  $C_n[\Gamma]^{w_o} \in \mathcal{O}_{\text{loc}}(\mathcal{E}_w^{w_o})$  is a local functional with kernel supported on the diagonal of  $\mathbb{R}^2 \times \mathbb{C}^2$ . By the wave-front bound of Lemma 8.16 (3), the support is the bulk diagonal independently of  $w_o$ . The transition map  $\mathcal{O}_{\text{loc}}(\mathcal{E}_w^{w'_o}) \rightarrow \mathcal{O}_{\text{loc}}(\mathcal{E}_w^{w_o})$  for  $w'_o \geq w_o$  is the restriction of polynomial functionals to truncated coefficient indices, equivalently setting higher coefficient fields to zero; in the regulator-admissible sector these maps are degreewise surjective. This preserves locality (the support of the truncated kernel is the same diagonal). The inverse limit  $\mathcal{O}_{\text{loc}}(\mathcal{E}_w) = \varprojlim_{w_o} \mathcal{O}_{\text{loc}}(\mathcal{E}_w^{w_o})$  exists in the weighted Tate category and is the candidate local-functional class on  $\mathcal{E}_w$ . The windowwise counterterms  $\{C_n[\Gamma]^{w_o}\}_{w_o}$  form a compatible inverse system, and their inverse limit  $C_n[\Gamma] = \varprojlim_{w_o} C_n[\Gamma]^{w_o}$  lies in  $\mathcal{O}_{\text{loc}}(\mathcal{E}_w)$  with bound (ii). Its universal property is precisely the one proved in Proposition 5.16: it is the unique compatible strict ML element with these finite-window projections.

The endpoint obstruction class  $[C_n[\Gamma]] \in H^1(\mathcal{O}_{\text{loc}}(\mathcal{E}_w), Q + \{I_o, -\})$  is represented by the inverse limit of the compatible windowwise classes when that compatibility holds. The theorem identifies its analytic support and coefficient topology. Its vanishing is an additional algebraic/BV condition on the mixed brane-defect interaction.  $\square$

*Remark 8.18.* The Mittag-Leffler limit in the coefficient direction commutes with Hörmander's wave-front analysis because the propagator factorization (20) is exact (the bulk wave-front set is computed once and tensored), the coefficient factor  $C_{\mathfrak{h}, w}$  is an algebraic tensor in the weighted Tate category with  $\text{WF} = \emptyset$ , and the Hadamard parametrix coefficients  $a_k(x, y)$  depend only on the bulk Riemannian/Hermitian data of  $\mathbb{R}^2 \times \mathbb{C}^2$ . The wave-front bound of Lemma 8.16 (3) is therefore exact and  $w_o$ -uniform; the

Mittag-Leffler limit preserves the local-functional class graph by graph at every order  $\hbar^n$  whenever the windowwise counterterm system is compatible.

**COROLLARY 8.19** (*Hadamard coefficient criterion for the weighted QME lift*). In the graphwise Mittag-Leffler sector of Theorem 8.17, Proposition 5.71 is equivalent to the vanishing of the mixed brane-defect QME obstruction class. The all-order quantum source-to-boundary CE/PV comparison

$$\begin{aligned} \Phi_{\hbar}^w : C_{\text{CE,cont}}^{\bullet}(\mathfrak{h}_{w,\hbar} \ltimes \mathfrak{h}_{\hbar,\text{cont}}^{\vee,w}[1])[[\hbar]] \\ \longrightarrow Z_{\text{fact}}^{P_{\circ,\hbar}}(\text{Obs}_{\text{open}}^{\hbar})|_{\mathfrak{g}_w} \end{aligned}$$

extends to the weighted cotangent direction at all orders in  $\hbar$  as a continuous  $\mathbb{C}[[\hbar]]$ -linear cochain map precisely when the obstruction class in  $H^1(\mathcal{O}_{\text{loc}}(\mathcal{E}_w), Q + \{I_{\circ}, -\})$  vanishes. Equivalently, this vanishing statement is the datum required by Corollary 5.73.

*Proof.* Proposition 5.71 assumes the Costello-locality/QME compatibility identity of Problem 5.72. By Theorem 8.17, the Hadamard wave-front-set and Mittag-Leffler coefficient-topology pieces of that identity are controlled. The proposition therefore asks for exactly the vanishing of the mixed brane-defect QME obstruction class.  $\square$

**THEOREM 8.20** (*Weighted RG locality obstruction criterion*). Before adjoining the principal-part current extension, in the graphwise finite-window sector of Theorem 8.17, the analytic part of Problem 5.72 is exactly the following finite-window coefficient criterion:

$$\begin{aligned} P_{\epsilon,L}^w &= P_{\epsilon,L}^{\text{base}} \widehat{\otimes} C_{\mathfrak{h} \oplus \mathfrak{h}^{\vee}[1],w}, \\ \text{WF}(C_{\mathfrak{h},w}^{w_{\circ}}) &= \emptyset, \\ \varprojlim_{w_{\circ}} P_{\epsilon,L}^{w,w_{\circ}} &= 0. \end{aligned}$$

These conditions hold by Lemma 8.16. After they are imposed, the endpoint datum in the weighted QME lift is the class

$$\text{Ob}_w \in H^1(\mathcal{O}_{\text{loc}}(\mathcal{E}_w), Q + \{I_{\circ}, -\}).$$

*Proof.* The propagator factorization is (20); its finite-window form is the definition of  $P_{\epsilon,L}^{w,w_{\circ}}$  in Lemma 8.16. The same lemma proves that the coefficient Casimir is an algebraic tensor with empty spacetime wave-front set and that the finite-window propagators form a strict Mittag-Leffler system. Thus the Hadamard singular support, asymptotic coefficients, and locality estimates are exactly the bulk Costello data tensored with a compatible coefficient kernel. The only term of Problem 5.72 remaining after these analytic statements is the local-functional QME class displayed above. The  $\mathcal{D}'_{\epsilon}(I) \otimes \mathcal{P}[1]$  defect-current compatibility clause is the additional datum of Problem 5.212.  $\square$

#### *Finite-window graph/QME assembly*

The QME obstruction class isolated by Theorem 8.17 is a sum over Costello graphs at every  $\hbar$ -order, each graph carrying its renormalized weight, source-face, BV-bracket, and counterterm contributions. Extracting it requires a finite graph list  $\mathcal{G}_M$  on a finite Hamiltonian window  $M$ , with labels and incidence signs declared, and the order- $n$  residual written as a finite cochain.

**Definition 8.21** (*Finite-window graph/QME system*). Fix an interval  $I$ , an admissible weight  $w$ , a finite Hamiltonian window  $M$ , and a finite set  $\mathcal{G}_M$  of connected Costello graphs. A finite-window graph/QME system over  $(I, w, M, \mathcal{G}_M)$  consists of the following data.



- (F1) For every  $\Gamma \in \mathcal{G}_M$  with  $r_\Gamma$  cotangent external labels, a tuple  $B_\bullet^\Gamma = (B_1^\Gamma, \dots, B_{r_\Gamma}^\Gamma)$  satisfying one of the two proved label conditions:

$$B_\bullet^\Gamma \in (\mathcal{D}_c^{\text{reg}}(I))^{r_\Gamma} \quad \text{or} \quad B_\bullet^\Gamma \in \mathcal{D}'_{c, \text{WF}}(I; \Gamma, M).$$

The second condition is the graphwise relation of Definition B.18. If one wants an honest CE source from a linear subspace  $\mathcal{V} \subset \mathcal{D}'_c(I)$ , every finite tuple from  $\mathcal{V}$  must satisfy the corresponding graphwise relation. Otherwise the source is only the finite CE span of the specified labelled inputs. Denote the resulting source CE object by  $C_{\text{lab}}^\bullet(\mathcal{G}_M)$ .

- (F2) The finite-scale regulator data

$$P_{\epsilon, L}^{w, M}, \quad \Delta_{\epsilon, L}^{w, M}, \quad I_{o, M}^w,$$

together with graph weights  $W_{\Gamma, \epsilon, L}^M(B_\bullet^\Gamma)$ , local counterterms  $C_{\Gamma, w}^M(\epsilon; B_\bullet^\Gamma)$ , and renormalized weights

$$W_{\Gamma, L, w}^{R, M} = \lim_{\epsilon \rightarrow 0} (W_{\Gamma, \epsilon, L}^M - C_{\Gamma, w}^M(\epsilon; B_\bullet^\Gamma)).$$

These are the regular-density weights of Proposition B.17, or the wavefront-admissible weights of Proposition B.19.

- (F3) A complete filtered local-functional subcomplex

$$\mathfrak{Q}_{\mathcal{G}, M}^\bullet(I) = \left( \mathcal{O}_{\text{loc}, \mathcal{G}, M}^{\text{bd}}(\mathcal{E}_w^M|_I; A_{\partial, \hbar}^{\text{pp}, w, M}(I)), d_M \right), \quad d_M = Q + \{I_{o, M}^w, -\}_{\hbar, M},$$

generated, and then completed in the  $\hbar$ -adic and scalar-contact filtrations, by the displayed renormalized graph weights, their local counterterms, and the  $d_M$ -boundaries and BV brackets forced by the finite graph differential on  $\mathcal{G}_M$ . This is a subcomplex only after those boundary and bracket terms have been included.

- (F4) A filtered chain projection on this subcomplex,

$$\widehat{\sigma}_{\text{sc}, M}: \mathfrak{Q}_{\mathcal{G}, M}^\bullet(I) \longrightarrow C_{\text{Lie}}^\bullet(\bar{A}; \mathbb{C}\chi)[1][[\hbar]],$$

whose associated graded is the scalar-symbol projection. Equivalently, either the graph data lie in the balanced graph/counterterm subhabitat of Definition E.20, or the scalar-extension obstructions

$$\delta_{\text{Cost}, \sigma}^{(o)}(I), \quad \mathfrak{d}_{\text{Cost}, \sigma}^{(r)}(I), \quad r > o,$$

of Theorem E.22 have been killed on the chosen finite-window graph/counterterm subcomplex.

- (F5) A degree-zero graph kernel

$$\kappa_{\mathcal{G}, w, I}^M \in \text{Hom}_{\text{cont}}\left(C_{\text{lab}}^\bullet(\mathcal{G}_M), \mathfrak{Q}_{\mathcal{G}, M}^\bullet(I)\right),$$

restricted to the chosen label set, whose zero-internal-edge part is  $\kappa_{\hbar, w, I}^{\text{coef}, M}$  of Proposition B.16 and whose positive graph components are the renormalized weights  $W_{\Gamma, L, w}^{R, M}$ .

Summation over graph topologies is part of the datum exactly when a separate graph-summability hypothesis is supplied.

THEOREM 8.22 (*Finite-window graph/QME assembly*). Let  $(I, w, M, \mathcal{G}_M)$  carry a finite-window graph/QME system in the sense of Definition 8.21. Put

$$S^\bullet = C_{\text{Lie}}^\bullet(\bar{A}; \mathbb{C}\gamma) [1] [[\hbar]], \quad Q_{\mathcal{G}, M}^\bullet(I) = \ker(\widehat{\sigma}_{\text{sc}, M}) \subset \mathfrak{Q}_{\mathcal{G}, M}^\bullet(I).$$

Let

$$\text{Ob}_{\mathcal{G}, M}^{\text{graph}} = \text{ev}_{\mathcal{G}} \text{Curv}_{\mathbf{K}}(\kappa_{\mathcal{G}, w, I}^M) \in \mathfrak{Q}_{\mathcal{G}, M}^1(I)$$

be the graph curvature evaluated on the chosen finite labelled CE cycles; before evaluation it is the corresponding Hom-valued curvature. The source-differential term is included in  $\text{Curv}_{\mathbf{K}}$ . Here  $\text{Curv}_{\mathbf{K}} = d_{\mathbf{K}} + \frac{1}{2}[-, -]_{\mathbf{K}}$  as in Definition B.15. Suppose counterterms have been chosen through order  $\hbar^{n-1}$ , and write  $C_{<n, M} \in \mathfrak{Q}_{\mathcal{G}, M}^0(I)$  for their sum. The order- $n$  residual is

$$\mathfrak{o}_{n, M}^{\text{fw}} = [\hbar^n] \left( \text{Ob}_{\mathcal{G}, M}^{\text{graph}} + d_M C_{<n, M} + \frac{1}{2} \{C_{<n, M}, C_{<n, M}\}_{\hbar, M} \right).$$

Then the following statements hold.

- (1) The analytic graph weights in the system are finite-window local functionals satisfying the stated locality hypotheses. For regular density labels this is Proposition B.17; for wavefront-admissible labels this is Proposition B.19. The statement applies to these regular-density and wavefront-admissible label classes.
- (2) The first obstruction is scalar. If

$$\mathfrak{s}_{n, M} := [\hbar^n] \widehat{\sigma}_{\text{sc}, M} \left( \text{Ob}_{\mathcal{G}, M}^{\text{graph}} + d_M C_{<n, M} + \frac{1}{2} \{C_{<n, M}, C_{<n, M}\}_{\hbar, M} \right)$$

is nonzero in  $S^1$ , the order- $n$  graph/QME equation has a scalar obstruction before passage to the non-scalar kernel. In the balanced graph/counterterm subhabitat this scalar term is zero by Proposition E.21.

- (3) If the scalar term vanishes through order  $n$  and the lower-order QME equations hold, then

$$\mathfrak{o}_{n, M}^{\text{fw}} \in Z^1(Q_{\mathcal{G}, M}^\bullet(I)).$$

An order- $n$  finite-window counterterm  $C_{n, M} \in \mathfrak{Q}_{\mathcal{G}, M}^0(I)$  satisfying

$$d_M C_{n, M} = -\mathfrak{o}_{n, M}^{\text{fw}}$$

exists if and only if

$$[\mathfrak{o}_{n, M}^{\text{fw}}] = 0 \quad \text{in} \quad H^1(Q_{\mathcal{G}, M}^\bullet(I), d_M).$$

If this class is nonzero, it is the precise finite-window non-scalar obstruction to the graph/QME realization at order  $\hbar^n$ .

*Proof.* The first assertion is exactly the analytic input supplied by the regular-density and wavefront-admissible kernel propositions. The finite-window hypothesis makes every coefficient factor finite rank, so Theorem 8.17 supplies the Hadamard counterterm control for each fixed graph. The label restriction is essential: outside  $\mathcal{D}'_{c, \text{WF}}(I; \Gamma, M)$ , the product of the graph Hadamard coefficient with the pushed-forward current label requires the wavefront-admissible product condition to define a canonical distribution.

The scalar statement is formal. A filtered chain projection  $\widehat{\sigma}_{\text{sc},M}$  is part of the system. If the displayed scalar projection of the residual is nonzero, then the residual has nonzero scalar component outside  $\ker \widehat{\sigma}_{\text{sc},M}$ , so every counterterm inside that kernel leaves the scalar component unchanged. In the balanced graph/counterterm subhabitat the projection is the zero chain map, hence the scalar branch vanishes before the non-scalar class is formed.

Finally, the completed convolution Hom is a filtered dg Lie algebra. The Bianchi identity

$$d_{\mathbf{K}} \text{Curv}_{\mathbf{K}}(\kappa) + [\kappa, \text{Curv}_{\mathbf{K}}(\kappa)]_{\mathbf{K}} = 0$$

and the lower-order QME equations imply that the first unsolved  $\hbar^n$ -coefficient is closed in the convolution differential. After evaluation on labelled CE cycles this is  $d_M$ -closed. Since the scalar projection is a chain map and its order- $n$  value is zero, this coefficient lies in the kernel complex  $\mathcal{Q}_{\mathcal{G},M}^\bullet(I)$ . Adding a degree-zero counterterm changes the first unsolved coefficient by the  $d_M$ -boundary of that counterterm; quadratic counterterm terms are already included in the displayed residual. Therefore the finite-window equation is solvable exactly when the displayed degree-one cohomology class vanishes.  $\square$

**PROPOSITION 8.23** (*Finite-window-to-limit obstruction*). Suppose that finite-window graph/QME systems  $\{(I, w, M, \mathcal{G}_M)\}_{M \geq 1}$  are given and satisfy the following additional hypotheses.

- (L1) The graph system is compatible under truncation:  $\pi_{M,N} \mathfrak{D}_{\mathcal{G},M}^\bullet(I) \subset \mathfrak{D}_{\mathcal{G},N}^\bullet(I)$ , the kernels, counterterms, and scalar projections commute with  $\pi_{M,N}$ , and the same compatibility holds under admissible weight transport  $R_{w,w'}^M$ .
- (L2) At the fixed  $\hbar$ -order under consideration the graph sum is finite in each window, or a separate Costello graph-summability hypothesis has been supplied. This graph-summability hypothesis is separate from the finite-window theorem.
- (L3) The residuals  $\mathfrak{D}_{n,M}^{\text{fw}}$  of Theorem 8.22 are compatible:

$$\pi_{M,N} \mathfrak{D}_{n,M}^{\text{fw}} = \mathfrak{D}_{n,N}^{\text{fw}}, \quad M \geq N.$$

Put

$$\mathfrak{D}_n^{\text{fw}} = \left( ([\mathfrak{D}_{n,M}^{\text{fw}}])_M, \lambda_n^{\text{fw}} \right),$$

where

$$([\mathfrak{D}_{n,M}^{\text{fw}}])_M \in \varprojlim_M H^1(\mathcal{Q}_{\mathcal{G},M}^\bullet(I)),$$

and, after choosing windowwise primitives  $c_{n,M}$  with  $d_M c_{n,M} = -\mathfrak{D}_{n,M}^{\text{fw}}$ ,

$$\lambda_n^{\text{fw}} \in \varprojlim_M^1 H^0(\mathcal{Q}_{\mathcal{G},M}^\bullet(I))$$

is represented by the Roos cocycle

$$[\pi_{M,N} c_{n,M} - c_{n,N}] \in H^0(\mathcal{Q}_{\mathcal{G},N}^\bullet(I)).$$

Then compatible order- $n$  counterterms  $C_{n,M}$  with  $\pi_{M,N} C_{n,M} = C_{n,N}$  exist if and only if  $\mathfrak{D}_n^{\text{fw}} = 0$ . If the tower  $\{H^0(\mathcal{Q}_{\mathcal{G},M}^\bullet(I))\}_M$  is Mittag-Leffler, then  $\lambda_n^{\text{fw}} = 0$ , and the inverse-limit obstruction is only the compatible  $H^1$ -class above.

Hypotheses (L1)–(L3) are the scope of the inverse-limit graph/QME theorem. When (L2) is unmet, the graph sum itself lies outside the tower, so the obstruction class is unformed.

*Proof.* Windowwise, Theorem 8.22 says that the only non-scalar obstruction is the class of  $\mathfrak{o}_{n,M}^{\text{fw}}$  in  $H^1$ . Necessity of the two displayed inverse-limit conditions is immediate from any compatible primitive  $C_{n,M}$ . Conversely, vanishing of the compatible  $H^1$ -class gives primitives  $c_{n,M}$  window by window. The differences  $\pi_{M,N} c_{n,M} - c_{n,N}$  are closed degree-zero elements, and their cohomology classes are the standard Roos 1-cocycle computing  $\varprojlim_M^1 H^0(Q_{\mathcal{G},M}^\bullet(I))$ . The vanishing of  $\lambda_n^{\text{fw}}$  is exactly the statement that the  $c_{n,M}$ 's can be adjusted by closed degree-zero terms and boundaries to become compatible under all truncation maps. The Mittag-Leffler sufficient condition is the standard vanishing of  $\varprojlim^1$  for a Mittag-Leffler tower. The weight-transport compatibility follows from the same diagonal transport squares used in Theorem 5.81.  $\square$

The order- $n$  residual of Proposition 8.23 is a sum of graph-differential, CE-source, BV-bracket, and lower-counterterm contributions, each carrying its own incidence sign, scalar projection, and truncation matrix. Computing, testing, and propagating the residual window by window requires a finite array of rows recording every contribution in normal form.

*Definition 8.24 (Finite-window graph array).* Fix a finite-window graph/QME system  $(I, w, M, \mathcal{G}_M)$ , a codimension-two closed marked Costello list  $\mathcal{G}_M^{\text{mk}}$ , and a labelled CE cycle  $\zeta_M$ . A finite-window graph array at order  $n$  is a finite set

$$\mathbf{A}_{n,M}^{\text{fw}} = \{\alpha\}$$

together with the following row data.

(A1) *Graph type and order.* A row has a type

$$\tau_\alpha \in \{\text{dK, CE, bracket, lower-}d, \text{ lower-bracket}\}.$$

For dK, CE, and bracket rows it specifies, respectively, a marked graph  $\Gamma_\alpha$ , a CE-source face of such a graph, or an ordered pair  $(\Gamma_{\alpha,1}, \Gamma_{\alpha,2})$ , all in the finite marked list. The order is

| $\tau_\alpha$              | $ \alpha _h$                                    |
|----------------------------|-------------------------------------------------|
| dK, CE                     | $ \Gamma_\alpha _h$                             |
| bracket                    | $ \Gamma_{\alpha,1} _h +  \Gamma_{\alpha,2} _h$ |
| lower- $d$ , lower-bracket | $n,$                                            |

and only rows with  $|\alpha|_h = n$  enter  $\mathbf{A}_{n,M}^{\text{fw}}$ .

(A2) *Edge labels.* Each internal edge of the graph carries one of the finite-window edge kernels

$$P_{\epsilon,L}^{w,M}, \quad \Delta_{\epsilon,L}^{w,M}, \quad E_{\text{Weyl}},$$

where  $E_{\text{Weyl}}$  is the boundary Weyl edge whose scalar two-Hamiltonian part is  $(f, g) \mapsto \omega(f, g)\gamma_{\mathbf{r}}$ . The two half-edge coefficient labels lie in

$$\mathfrak{h}_{w,\leq M} \cup D_{\leq M}^w, \quad D_{\leq M}^w = \prod_{d \leq M} w(d)^{-1} H_d^\vee,$$

and the edge record includes the source/target orientation used by the BV bracket sign.

(A3) *Vertex labels.* Each vertex is labelled by one of the finite system labels

$$u_{a(t)dt \otimes f}, \quad c_{B,\rho}, \quad I_{o,M}^w, \quad W_{\Gamma,L,w}^{R,M}(B_\bullet^\Gamma), \quad C_{\Gamma,w}^M(\epsilon; B_\bullet^\Gamma), \quad \gamma_{\mathbf{r}},$$

with  $f, \rho$  in degree  $\leq M$  and  $B_\bullet^\Gamma$  either regular-density or graphwise wavefront-admissible as in Definition 8.21.

(A4) *Markers and incidence coefficient.* The row contains a marker tensor

$$\mathbf{m}_\alpha = (\mathbf{m}_{\alpha,L})_L, \quad \mathbf{m}_{\alpha,L} \in \mathcal{M}_{m_L}^{\text{full}}(\mathbb{C}^{N|N})$$

or the explicitly chosen smaller marker complex. If the row comes from an elementary boundary operation  $\beta_i \in \mathbf{B}_M(\Gamma_\alpha)$ , its incidence coefficient is

$$\iota_{\alpha,M} = (-1)^{i-1} \varepsilon_{\beta_i}.$$

For CE-source and convolution-bracket rows,  $\iota_{\alpha,M}$  is the corresponding CE or BV Koszul sign, including the factor 1/2 in the bracket row.

(A5) *Weight and filtrations.* The row records

$$d_{\max}(\alpha) = \max\{d : H_d \text{ or } H_d^\vee \text{ occurs in the row}\} \leq M,$$

its scalar-contact filtration  $p_\alpha = \#\{\text{scalar-contact open loops}\}$ , and, for a boundary operation  $\beta_i$ ,

$$r_\alpha = p_\alpha - p(\partial_{\beta_i} \Gamma_\alpha, \mu_{\beta_i}(\mathbf{m}_\alpha)).$$

(A6) *Curvature value and scalar condition.* The row evaluates to a local functional

$$R_{\alpha,M}^{\text{row}} \in \mathfrak{Q}_{\mathcal{G},M}^{\text{I}}(I).$$

For the graph-curvature part these are

| $\tau_\alpha$ | $R_{\alpha,M}^{\text{row}}$                                                        |
|---------------|------------------------------------------------------------------------------------|
| dK            | $\iota_{\alpha,M} d_M K_{\Gamma_\alpha}^M$                                         |
| CE            | $\iota_{\alpha,M} K_{\Gamma_\alpha}^M$                                             |
| bracket       | $\iota_{\alpha,M} \{K_{\Gamma_{\alpha,1}}^M, K_{\Gamma_{\alpha,2}}^M\}_{\hbar,M}.$ |

The two lower-counterterm row types are the corresponding homogeneous summands of  $[\hbar^n] d_M C_{<n,M}$  and  $\frac{1}{2} [\hbar^n] \{C_{<n,M}, C_{<n,M}\}_{\hbar,M}$ . The scalar-condition entry is

$$S_{\alpha,M} = \widehat{\sigma}_{\text{sc},M}(R_{\alpha,M}^{\text{row}}).$$

If  $\alpha$  is a first filtration-lowering two-Hamiltonian scalar contact row, this specializes to

$$S_{\alpha,M} = \iota_{\alpha,M} \text{Str}_{\text{cyc}}(\mu_{\beta_i}(\mathbf{m}_\alpha)) \omega(f_{\partial_{\beta_i} \Gamma_\alpha}, g_{\partial_{\beta_i} \Gamma_\alpha}) \gamma_1.$$

The non-scalar reduced-curvature contribution is the scalar-zero sum

$$R_{n,M}^{\text{ns}} = \sum_{\alpha \in \mathbf{A}_{n,M}^{\text{fw}}} R_{\alpha,M}^{\text{row}} \quad \text{provided} \quad \sum_{\alpha \in \mathbf{A}_{n,M}^{\text{fw}}} S_{\alpha,M} = 0.$$

An entrywise non-scalar summand requires an additional chain splitting of  $\widehat{\sigma}_{\text{sc},M}$ ; the displayed scalar-zero sum is the canonical datum in the present array.

(A7) *Truncation in  $M$ .* For  $M \geq N$  the array contains truncation coefficients  $t_{\alpha\alpha'}^{M,N} \in \mathbb{C}$  with  $\alpha' \in A_{n,N}^{\text{fw}}$ , defined by

$$\pi_{M,N} R_{\alpha,M}^{\text{row}} = \sum_{\alpha'} t_{\alpha\alpha'}^{M,N} R_{\alpha',N}^{\text{row}}, \quad \pi_{M,N} S_{\alpha,M} = \sum_{\alpha'} t_{\alpha\alpha'}^{M,N} S_{\alpha',N}.$$

A row whose coefficient labels lie entirely above degree  $N$  has all  $t_{\alpha\alpha'}^{M,N} = 0$ . A filling of this equation is exactly the truncation entry required by the lower-window theorem.

(A8) *Order-two boundary compositions.* For an order-2 closure check the array also records the ordered pairs of distinct first scalar-contact boundary operations

$$(\beta_i, \beta_j), \quad r(\beta_i) = r(\beta_j) = \mathbf{1}, \quad |\beta_i|_h = |\beta_j|_h = \mathbf{1},$$

whose two composites occur in the codimension-two closure table of Definition E.27. If  $\text{pos}_i(j)$  denotes the position of the induced face  $\beta_j|\beta_i$  after deleting  $\beta_i$ , the composite incidence coefficient is

$$\iota_{\beta_j|\beta_i, \beta_i} = (-1)^{i-1} (-1)^{\text{pos}_i(j)-1} \varepsilon_{\beta_i} \varepsilon_{\beta_j|\beta_i}.$$

The corresponding two-contact scalar condition is the row

$$S_{\beta_j|\beta_i, M}^{(2)} = \iota_{\beta_j|\beta_i, \beta_i} \left( \prod_{L \in \mathcal{L}_{\text{sc}}(\partial_{\beta_j|\beta_i} \partial_{\beta_i} \Gamma)} \text{Str}_{\text{cyc}}(\mathbf{m}'_L) \right) \omega(f_{\beta_i}, g_{\beta_i}) \omega(f_{\beta_j|\beta_i}, g_{\beta_j|\beta_i}) \gamma_{\mathbf{1}}^{(2)}.$$

Here  $\mathbf{m}'_L$  is the transported marker on the output scalar loop  $L$ , and  $\gamma_{\mathbf{1}}^{(2)}$  denotes the ordered two-contact associated-graded scalar-contact symbol. A surviving product  $\gamma_{\mathbf{1}}^2$  in the target algebra is a separate datum. The equality of output graphs, coefficients, and marker transports in (E.27) is exactly the order-2 composite-face datum.

The array is *valid* if the row sum reproduces the order- $n$  residual:

$$R_{n,M} = \sum_{\alpha \in A_{n,M}^{\text{fw}}} R_{\alpha,M}^{\text{row}}, \quad \mathfrak{s}_{n,M} = \sum_{\alpha \in A_{n,M}^{\text{fw}}} S_{\alpha,M},$$

and if the truncation coefficients satisfy the displayed equations for every  $M \geq N$ .

**PROPOSITION 8.25** (*First finite-window array rows*). The universal computed rows are the reduced current interface row in order 0 and the two-Hamiltonian scalar-contact row in order 1.

On the length-one reduced current interface, Theorem B.7 gives strict CE/PV compatibility. Thus the order-0 row has

$$R_{0,M}^{\text{row}} = 0, \quad S_{0,M} = 0.$$

For the ordered Hamiltonian inputs  $(a, z_1), (b, z_2)$ , the one-cross boundary row has graph type dK, one boundary Weyl edge  $E_{\text{Weyl}}$ , two external Hamiltonian vertices, one scalar-contact output vertex  $\gamma_{\mathbf{1}}$ , and

$$\omega(z_1, z_2) = \mathbf{1}, \quad |\alpha_{\text{sc}}|_h = \mathbf{1}.$$

Its scalar condition is

$$S_{\alpha_{\text{sc}}, M} = \text{Str}_{\text{cyc}}(\mu_{\beta_{\text{sc}}}(\mathbf{m})) \gamma_{\mathbf{1}},$$

after extracting the  $\hbar^1$ -coefficient and using the orientation convention in which the ordered pair  $(z_1, z_2)$  has positive sign. Therefore:

| branch                                            | $S_{\alpha_{sc}, M}$               | $R_{1, M}^{ns}$ from this row       |
|---------------------------------------------------|------------------------------------|-------------------------------------|
| $\mathfrak{gl}_N$ ordinary                        | $N\gamma_1$                        | scalar branch                       |
| even block marker $\lambda_o P_o + \lambda_1 P_1$ | $N(\lambda_o - \lambda_1)\gamma_1$ | formed when $\lambda_o = \lambda_1$ |
| $\mathfrak{gl}(N N)$ full-equivariant             | o                                  | o                                   |

Consequently, for the minimal full-equivariant closed array containing only the strict current-interface rows and this scalar-contact row, the first order residual is zero and

$$C_{1, M} = o$$

is the compatible order-one non-scalar counterterm. For any larger graph system, the first non-scalar residual is

$$R_{1, M}^{ns} = \sum_{\alpha \in A_{1, M}^{fw} \setminus \{\alpha_{sc}\}} R_{\alpha, M}^{row}$$

after the scalar condition has vanished. Its primitive is decided only after supplying the remaining order-one rows of Definition 8.24.

*Proof.* The order-o assertion is exactly the strict current-source/target identity of Theorem B.7 on the length-one generators: the CE differential and target  $P_o$  bracket use the same semidirect product constants.

The order-1 scalar-contact row is the computation of Proposition E.23, written in the row notation of Definition 8.24. The monomial value is  $\omega(z_1, z_2) = 1$ . In the ordinary branch the open index loop contributes  $N$ . In the even block-diagonal marker branch the cyclic supertrace is  $N(\lambda_o - \lambda_1)$ . In the full-equivariant branch, Lemma E.31 kills the cyclic supertrace of every transported marker tensor in  $\mathbb{C}^{N|N}$ ; the empty marker word is killed by the same  $\text{sdim } \mathbb{C}^{N|N} = o$  calculation. Thus the row is already zero as a local functional in the full-equivariant branch.

If the array contains only the displayed order-one row, the order-one residual is zero, so the zero counterterm is a primitive and is compatible under all truncation maps. If further order-one rows are present, the displayed sum is the definition of the remaining scalar-zero residual. A primitive for it is a theorem exactly when the missing row values, signs, scalar conditions, and truncation coefficients are supplied and the resulting  $H^1$ -class vanishes.  $\square$

**PROPOSITION 8.26** (*Universal scalar-contact bracket rows*). Work in the local full-equivariant marked branch with  $V = \mathbb{C}^{N|N}$ , and let  $\alpha_{sc}$  be the one-cross two-Hamiltonian scalar-contact row of Proposition 8.25. Write  $K_{\alpha_{sc}}^M$  for the local functional carried by this row. Then

$$K_{\alpha_{sc}}^M = o$$

in the finite-window local-functional complex.

Consequently every order-2 convolution-bracket row containing  $\alpha_{sc}$  has the following entries. For any order-1 component  $\beta$  for which the bracket row is present,

| row                           | $R_{b, M}^{row}$                                                                                             |
|-------------------------------|--------------------------------------------------------------------------------------------------------------|
| $b(\alpha_{sc}, \beta)$       | $\frac{1}{2} \varepsilon_{\alpha_{sc}, \beta} \{K_{\alpha_{sc}}^M, K_{\beta}^M\}_{\hbar, M} = o$             |
| $b(\beta, \alpha_{sc})$       | $\frac{1}{2} \varepsilon_{\beta, \alpha_{sc}} \{K_{\beta}^M, K_{\alpha_{sc}}^M\}_{\hbar, M} = o$             |
| $b(\alpha_{sc}, \alpha_{sc})$ | $\frac{1}{2} \varepsilon_{\alpha_{sc}, \alpha_{sc}} \{K_{\alpha_{sc}}^M, K_{\alpha_{sc}}^M\}_{\hbar, M} = o$ |

Their scalar conditions are

$$S_{b,M} = 0.$$

For every  $M \geq N$ , the truncation row may be chosen as

$$t_{bb'}^{M,N} = 0 \quad \text{for every lower-window bracket row } b',$$

since  $\pi_{M,N} R_{b,M}^{\text{row}} = 0$  and  $\pi_{M,N} S_{b,M} = 0$ . These rows therefore have zero order-2 obstruction and zero counterterm.

This proposition closes only the bracket family containing  $\alpha_{\text{sc}}$ . Bracket rows  $b(\beta_1, \beta_2)$  with  $\beta_1, \beta_2 \neq \alpha_{\text{sc}}$  remain part of the chosen graph system and must be supplied row by row.

*Proof.* Proposition 8.25 proves the stronger full-equivariant fact that the scalar-contact row is zero as a local functional, because every transported scalar-loop marker has zero cyclic supertrace on  $\mathbb{C}^{N|N}$ . The BV/convolution bracket is bilinear and continuous on the finite-window completed local-functional complex, hence any bracket with  $K_{\alpha_{\text{sc}}}^M = 0$  is zero before passing to cohomology. Applying the chain projection  $\widehat{\sigma}_{\text{sc},M}$  to zero gives the displayed scalar conditions. Truncation sends zero to zero, so the zero truncation matrix row is compatible with the defining equations of Definition 8.24.  $\square$

**PROPOSITION 8.27** (*Genuine order-two graph-weight rows*). Work in the local  $\mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$  full-equivariant marked branch with  $V = \mathbb{C}^{N|N}$ . Let  $\mathcal{G}_{2,M}^{\text{gen}}$  be the part of a finite marked Costello list consisting of genuine seed graph weights of  $\hbar$ -order 2: these are genuine seed rows, distinct from the codimension-two composites of two first scalar-contact boundary faces and from the bracket rows containing  $\alpha_{\text{sc}}$  closed in Proposition 8.26. For  $\Gamma \in \mathcal{G}_{2,M}^{\text{gen}}$ , write

$$K_{\Gamma}^M = W_{\Gamma,L,w}^{R,M}(B_{\bullet}^{\Gamma})$$

for the normalized renormalized finite-window graph weight, and let  $\varepsilon_{\Gamma}$  be the coefficient with which this component occurs in  $d_M \kappa_{\mathcal{G},w,I}^M$ , including the orientation, Koszul, and symmetry factors supplied by the finite graph system.

The genuine order-two rows have the following form.

(G1) The graph-differential row  $d\Gamma$  has

$$\tau_{d\Gamma} = dK, \quad |d\Gamma|_{\hbar} = 2,$$

inherits its edge labels, vertex labels, marker tensor, and coefficient degree from  $\Gamma$ , and has

$$R_{d\Gamma,M}^{\text{row}} = \varepsilon_{\Gamma} d_M K_{\Gamma}^M, \quad S_{d\Gamma,M} = \varepsilon_{\Gamma} \widehat{\sigma}_{\text{sc},M}(d_M K_{\Gamma}^M).$$

(G2) Before evaluation on a strict CE cycle, or when the Hom-valued kernel table is retained, each source face is a separate row. Write the relevant part of the source differential as

$$d_{\text{CE}} \zeta_M = \sum_{\nu} \varepsilon_{\Gamma,\nu}^{\text{CE}} \zeta_{M,\nu},$$

where  $\varepsilon_{\Gamma,\nu}^{\text{CE}}$  includes the CE Koszul sign and the graph coefficient attached to the  $\Gamma$ -component. The source row is

$$R_{\text{CE}(\Gamma,\nu),M}^{\text{row}} = -\varepsilon_{\Gamma,\nu}^{\text{CE}} K_{\Gamma}^M(\zeta_{M,\nu}),$$



and its scalar condition is

$$S_{\text{CE}(\Gamma, \nu), M} = -\varepsilon_{\Gamma, \nu}^{\text{CE}} \widehat{\sigma}_{\text{sc}, M}(K_{\Gamma}^M(\zeta_{M, \nu})).$$

If the curvature has already been evaluated on a strict labelled CE cycle, this source-face family is empty.

If these row values and all their  $d_M$ -faces lie in the full-equivariant marked complex of Theorem E.32, then

$$S_{d\Gamma, M} = 0, \quad S_{\text{CE}(\Gamma, \nu), M} = 0.$$

This is a scalar-condition statement. The local-functional values  $K_{\Gamma}^M$  and  $d_M K_{\Gamma}^M$  require their own row data.

After scalar-condition vanishing, the genuine seed-graph contribution to the order-2 reduced curvature is

$$\begin{aligned} R_{2, M}^{\text{ns, gen}} &= \sum_{\Gamma \in \mathcal{G}_{2, M}^{\text{gen}}} \varepsilon_{\Gamma} d_M K_{\Gamma}^M \\ &\quad - \sum_{\Gamma, \nu} \varepsilon_{\Gamma, \nu}^{\text{CE}} K_{\Gamma}^M(\zeta_{M, \nu}) \in \mathcal{K}_{\text{ns}, M}^1(I). \end{aligned}$$

The second sum is zero after strict CE-cycle evaluation. A genuine order-2 counterterm exists precisely when

$$[R_{2, M}^{\text{ns, gen}}] = 0 \quad \text{in} \quad H^1(\mathcal{K}_{\text{ns}, M}^{\bullet}(I), d_{\text{ns}, M}).$$

If this class is nonzero, it is the explicit finite-window obstruction carried by the genuine seed graphs.

Truncation is part of the row data. For  $M \geq N$  one must supply normalized coefficients

$$\pi_{M, N}(\varepsilon_{\Gamma} K_{\Gamma}^M) = \sum_{\Gamma'} u_{\Gamma \Gamma'}^{M, N} \varepsilon_{\Gamma'} K_{\Gamma'}^N,$$

and then the graph-differential truncation entries are

$$t_{d\Gamma, d\Gamma'}^{M, N} = u_{\Gamma \Gamma'}^{M, N}.$$

For source rows one must likewise supply

$$\pi_{M, N}(-\varepsilon_{\Gamma, \nu}^{\text{CE}} K_{\Gamma}^M(\zeta_{M, \nu})) = \sum_{\Gamma', \nu'} v_{\Gamma, \nu; \Gamma', \nu'}^{M, N} (-\varepsilon_{\Gamma', \nu'}^{\text{CE}} K_{\Gamma'}^N(\zeta_{N, \nu'})),$$

and set  $t_{\text{CE}(\Gamma, \nu), \text{CE}(\Gamma', \nu')}^{M, N} = v_{\Gamma, \nu; \Gamma', \nu'}^{M, N}$ . Rows whose coefficient labels lie entirely above degree  $N$  have zero truncation coefficients. The displayed truncation identities are exactly the row entries required by the lower-window theorem.

In the minimal full-equivariant local system of Proposition 8.28,  $\mathcal{G}_{2, M}^{\text{gen}} = \emptyset$  and the source-face set is empty. Hence

$$R_{2, M}^{\text{ns, gen}} = 0, \quad C_{2, M}^{\text{gen}} = 0.$$

In any larger system, vanishing, counterterms, and obstruction are decided by the following data: the finite seed list  $\mathcal{G}_{2, M}^{\text{gen}}$ ; the renormalized local weights  $K_{\Gamma}^M$  with regular-density or wavefront-admissible current labels; the full  $d_M K_{\Gamma}^M$  expansion with orientation signs; the CE-source decomposition and signs unless strict CE-cycle evaluation has removed them; proof that every row lies in the full-equivariant marked complex, or else its scalar projection; and the truncation matrices displayed above.

*Proof.* This is the order-2 part of the convolution-curvature formula in Theorem 8.36, with the scalar-contact bracket family removed by Proposition 8.26. The differential term contributes  $\varepsilon_\Gamma d_M K_\Gamma^M$ . The source term in  $d_K = d_M - (-1)^{|\cdot|}(\cdot)d_{\text{CE}}$  contributes the displayed minus sign for a degree-zero kernel. After evaluation on a strict CE cycle, that source term is zero.

The scalar-condition assertion follows from Theorem E.32: on the full-equivariant marked complex the filtered scalar projection is the zero chain map. Since this projection only kills the scalar shadow, it gives a scalar-condition theorem for the non-scalar local functional  $K_\Gamma^M$ .

The counterterm criterion is exactly Definition E.33: once the scalar condition is zero, the row sum is a degree-one element of the non-scalar kernel complex, and a degree-zero counterterm exists exactly when this element is a  $d_{\text{ns},M}$ -boundary. The truncation formulas are the row identities required in Definition 8.24, written for normalized graph and source rows. In the minimal system the genuine seed-row set is empty, so the sum is empty.  $\square$

**PROPOSITION 8.28** (*Order-two finite-window boundary rows*). Work in the local  $\mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$  full-equivariant marked branch with  $V = \mathbb{C}^{N|N}$ . Let the minimal array be generated by the strict current-interface rows and the one-cross scalar-contact row of Proposition 8.25. Assume the finite marked Costello list contains exactly these minimal rows: the genuine  $|\Gamma|_h = 2$  seed-graph set, the order-2 CE-source-face set, and the order-2 graph-counterterm-face set are empty. Use  $C_{1,M} = 0$ .

Then the only order-2 boundary data forced by codimension-two closure are the two ordered composites of two first scalar-contact boundary operations. For  $o = \beta_1 \wedge \beta_2$  they have incidence signs

$$\iota_{\beta_2|\beta_1,\beta_1} = \varepsilon_{\beta_1} \varepsilon_{\beta_2|\beta_1}, \quad \iota_{\beta_1|\beta_2,\beta_2} = -\varepsilon_{\beta_2} \varepsilon_{\beta_1|\beta_2}.$$

Their scalar conditions are

$$\begin{aligned} S_{\beta_2|\beta_1,M}^{(2)} &= \varepsilon_{\beta_1} \varepsilon_{\beta_2|\beta_1} \prod_L \text{Str}_{\text{cyc}}(\mathbf{m}'_L) \omega(f_{\beta_1}, g_{\beta_1}) \omega(f_{\beta_2|\beta_1}, g_{\beta_2|\beta_1}) \gamma_{\mathbf{I}}^{(2)}, \\ S_{\beta_1|\beta_2,M}^{(2)} &= -\varepsilon_{\beta_2} \varepsilon_{\beta_1|\beta_2} \prod_L \text{Str}_{\text{cyc}}(\mathbf{m}''_L) \omega(f_{\beta_2}, g_{\beta_2}) \omega(f_{\beta_1|\beta_2}, g_{\beta_1|\beta_2}) \gamma_{\mathbf{I}}^{(2)}. \end{aligned}$$

The codimension-two closure equations identify the two output graphs, the two products of coefficients, and the transported marker tensors. Therefore the pair sum is zero. In the full-equivariant branch it is stronger: every factor  $\text{Str}_{\text{cyc}}(\mathbf{m}'_L)$  and  $\text{Str}_{\text{cyc}}(\mathbf{m}''_L)$  is zero by Lemma E.31; hence each two-contact scalar row already vanishes as a local scalar row.

Under these minimal hypotheses the order-2 reduced non-scalar row sum is

$$R_{2,M}^{\text{ns}} = 0, \quad C_{2,M} = 0$$

is a compatible order-2 counterterm.

For any larger finite-window system, the order-2 residual is decided by the added row data. After scalar-condition vanishing its exact row sum is

$$\begin{aligned} R_{2,M}^{\text{ns}} &= \sum_{|\Gamma|_h=2} \varepsilon_\Gamma d_M K_\Gamma^M - \sum_{|\Gamma'|_h=2} \varepsilon_{\Gamma'}^{\text{CE}} K_{\Gamma'}^M \\ &\quad + \frac{1}{2} \sum_{|\Gamma_1|_h=|\Gamma_2|_h=1} \varepsilon_{\Gamma_1, \Gamma_2} \{K_{\Gamma_1}^M, K_{\Gamma_2}^M\}_{h,M} \\ &\quad + [\hbar^2] \left( d_M C_{<2,M} + \frac{1}{2} \{C_{<2,M}, C_{<2,M}\}_{h,M} \right). \end{aligned}$$

Thus a nonzero order-2 obstruction or its counterterm can be asserted only after supplying the genuine two-cross/contact graph weights  $K_{\Gamma}^M$ , the CE-source faces, the BV bracket rows between order-1 graph components distinct from those containing  $\alpha_{\text{sc}}$  or another retained zero local-functional row such as  $\alpha_{\text{II}}$ , the lower-counterterm rows if a nonminimal  $C_{1,M}$  was chosen, the scalar projection of any non-full-equivariant two-contact symbol, and the truncation coefficients  $t_{\alpha\alpha'}^{M,N}$ .

*Proof.* The signs are those of Lemma E.29. Starting with  $o = \beta_1 \wedge \beta_2$ , the face  $\beta_1$  followed by  $\beta_2|\beta_1$  contributes +1, while the face  $\beta_2$  followed by  $\beta_1|\beta_2$  contributes -1. The codimension-two closure equations Equation (E.27) identifies the output graph, coefficient, and marker transport in the two composites, so the two boundary-composition rows cancel pairwise.

The scalar-condition formula is the order-2 instance of the last row datum in Definition 8.24: one scalar-contact factor is contributed by each first-lowering boundary face. In the full-equivariant marked complex, marker transport preserves the signed Brauer span. Lemma E.31 gives zero cyclic supertrace for every transported scalar-loop marker, including the empty marker word. Hence the two-contact scalar rows vanish before passing to cohomology.

Under the stated minimal hypotheses the genuine order-2 graph-row set, CE-source-row set, and order-1 non-scalar primitive set are empty, and the full-equivariant trace calculation kills the remaining scalar-contact row. The possible bracket row between two one-cross scalar-contact components is also zero, because each such component has zero local-functional value in the full-equivariant branch by Proposition 8.25. The reduced non-scalar order-2 row sum is therefore zero, and the zero counterterm is compatible under finite-window truncation.

Removing any minimal hypothesis reintroduces exactly the displayed row types from the curvature formula in Theorem 8.36. The bracket rows containing  $\alpha_{\text{sc}}$  are the universal zero rows of Proposition 8.26; bracket rows containing a retained  $\alpha_{\text{II}}$ -row vanish by the same bilinearity argument and Proposition E.45. The remaining graph weights, signs, scalar projections, and truncation coefficients are the data defining  $R_{2,M}^{\text{ns}}$ , the order-2 obstruction class, and its counterterm equation.  $\square$

**PROPOSITION 8.29** (*Truncation matrices for the computed finite-window rows*). Work with the finite-window graph arrays of Definition 8.24. For  $M \geq N$ , use the following label-retaining convention: if a labelled row has  $d_{\text{max}} \leq N$ , the same label denotes its lower-window row; if  $d_{\text{max}} > N$ , the row is truncated away and all entries from it are zero. With this convention the computed rows have the following truncation matrices.

(T1) The one-cross scalar-contact row  $\alpha_{\text{sc}}$  of Proposition 8.25 has  $d_{\text{max}}(\alpha_{\text{sc}}) = 1$ . For  $M \geq N \geq 1$ ,

$$t_{\alpha_{\text{sc}}\alpha_{\text{sc}}}^{M,N} = 1, \quad t_{\alpha_{\text{sc}}\alpha'}^{M,N} = 0 \quad (\alpha' \neq \alpha_{\text{sc}}).$$

This applies to the ordinary, even-block, and full-equivariant branches. In the full-equivariant branch the row value is already zero, so the zero matrix would also satisfy the row equations; the displayed matrix is the chosen transport of the retained row label.

(T2) The labelled scalar-zero row  $\alpha_{\text{II}}$  of Proposition E.45, with ordered inputs  $(z_1, z_1)$ , has

$$R_{\alpha_{\text{II}},M}^{\text{row}} = 0, \quad S_{\alpha_{\text{II}},M} = 0, \quad d_{\text{max}}(\alpha_{\text{II}}) = 1.$$

If the row convention retains this labelled zero row, then for  $M \geq N \geq 1$

$$t_{\alpha_{\text{II}}\alpha_{\text{II}}}^{M,N} = 1, \quad t_{\alpha_{\text{II}}\alpha'}^{M,N} = 0 \quad (\alpha' \neq \alpha_{\text{II}}).$$

If zero rows are suppressed, the  $\alpha_{\text{II}}$ -row and its truncation matrix are empty.

(T3) Let  $\gamma_{21} = (\beta_2 | \beta_1, \beta_1)$  and  $\gamma_{12} = (\beta_1 | \beta_2, \beta_2)$  denote the two ordered codimension-two scalar-contact composites in Proposition 8.28. When the degree- $\leq N$  window contains the two Hamiltonian labels, the composite rows are transported diagonally:

$$t_{\gamma_{21}\gamma_{21}}^{M,N} = 1, \quad t_{\gamma_{12}\gamma_{12}}^{M,N} = 1, \quad t_{\gamma_{ij}\gamma'}^{M,N} = 0 \quad (\gamma' \neq \gamma_{ij}).$$

The codimension-two pair cancellation is therefore stable under truncation. In the full-equivariant branch the stronger rowwise vanishing is also stable.

(T4) Let  $b$  be any order-2 bracket row containing a retained zero local-functional row

$$\alpha_o \in \{\alpha_{sc}, \alpha_{11}\}.$$

Thus  $b = b(\alpha_o, \beta)$ ,  $b = b(\beta, \alpha_o)$ , or  $b = b(\alpha_o, \alpha_o)$ , whenever the displayed row belongs to the chosen finite marked list. Then

$$R_{b,M}^{\text{row}} = 0, \quad S_{b,M} = 0,$$

and the zero truncation row is

$$t_{bb'}^{M,N} = 0 \quad \text{for every lower-window bracket row } b'.$$

(T5) When genuine order-2 seed rows are zero, as in the minimal full-equivariant system, their truncation matrices are empty. If a larger system supplies genuine seeds  $\Gamma \in \mathcal{G}_{2,M}^{\text{gen}}$ , then the graph-differential rows are transported by the normalized matrix  $u^{M,N}$ :

$$\pi_{M,N}(\varepsilon_{\Gamma} K_{\Gamma}^M) = \sum_{\Gamma'} u_{\Gamma\Gamma'}^{M,N} \varepsilon_{\Gamma'} K_{\Gamma'}^N, \quad t_{d\Gamma, d\Gamma'}^{M,N} = u_{\Gamma\Gamma'}^{M,N}.$$

Source-face rows, when present, are transported by the normalized matrix  $v^{M,N}$ :

$$\pi_{M,N}(-\varepsilon_{\Gamma, \nu}^{\text{CE}} K_{\Gamma}^M(\zeta_{M, \nu})) = \sum_{\Gamma', \nu'} v_{\Gamma, \nu; \Gamma', \nu'}^{M,N} (-\varepsilon_{\Gamma', \nu'}^{\text{CE}} K_{\Gamma'}^N(\zeta_{N, \nu'})),$$

$$t_{\text{CE}(\Gamma, \nu), \text{CE}(\Gamma', \nu')}^{M,N} = v_{\Gamma, \nu; \Gamma', \nu'}^{M,N}.$$

Under the projection-closed row-basis hypotheses of Proposition 8.30, these are the coordinate entries of the finite-window projection. These hypotheses supply exactly the lower-window entries  $u_{\Gamma\Gamma'}^{M,N}$  and  $v_{\Gamma, \nu; \Gamma', \nu'}^{M,N}$  for those supplied genuine rows.

(T6) Bracket rows  $b(\beta_1, \beta_2)$  with  $\beta_i \notin \{\alpha_{sc}, \alpha_{11}\}$  require data beyond the present local computation. After a lower-window bracket row basis, or an equivalent row presentation with chosen coordinates, is fixed, their required truncation matrix is the coordinate family  $q^{M,N}$  satisfying

$$\pi_{M,N}\left(\frac{1}{2} \varepsilon_{\beta_1, \beta_2} \{K_{\beta_1}^M, K_{\beta_2}^M\}_{h,M}\right) = \sum_{\beta'_1, \beta'_2} q_{\beta_1, \beta_2; \beta'_1, \beta'_2}^{M,N} \frac{1}{2} \varepsilon_{\beta'_1, \beta'_2} \{K_{\beta'_1}^N, K_{\beta'_2}^N\}_{h,N}.$$

These  $q$ -entries, together with the underlying order-1 row values, signs, and lower-window bracket-basis reduction, are still missing data for the enlarged system.

The nonempty displayed matrices are stable under further truncation: for  $M \geq N \geq P$ ,

$${}_t M, P = {}_t M, N {}_t N, P$$

on the scalar-contact, scalar-zero, codimension-two, and zero-bracket families. For genuine seed rows and nonzero added bracket rows, stability is exactly the cocycle condition

$$u_{\Gamma''}^{M,P} = \sum_{\Gamma'} u_{\Gamma'}^{M,N} u_{\Gamma'\Gamma''}^{N,P},$$

$$v_{\Gamma, \nu; \Gamma'', \nu''}^{M,P} = \sum_{\Gamma', \nu'} v_{\Gamma, \nu; \Gamma', \nu'}^{M,N} v_{\Gamma', \nu'; \Gamma'', \nu''}^{N,P},$$

and the analogous equation for  $q$ .

*Proof.* For the one-cross scalar-contact rows the coefficient labels are  $z_1, z_2$ , and for  $\alpha_{\text{II}}$  the labels are  $z_1, z_1$ . These all have window degree 1. Therefore  $\pi_{M,N}$  preserves the labelled row for  $N \geq 1$ , and kills it only when the lower window omits the degree-one labels. The diagonal matrices in (T1) and (T2) are exactly this statement. The row  $\alpha_{\text{II}}$  is zero because  $\omega(z_1, z_1) = 0$ .

The codimension-two rows use the same two degree-one Hamiltonian labels and the same transported marker words before and after projection. The finite-window marked-complex compatibility preserves the output graph, coefficient product, marker transport, and incidence signs. Hence the two ordered composites truncate to the same labelled composites in the lower window, and their pair cancellation survives.

If a bracket row contains  $\alpha_{\text{sc}}$  or  $\alpha_{\text{II}}$ , one bracket input is zero as a local functional in the retained full-equivariant row family. Bilinearity of the BV/convolution bracket gives  $R_{b,M}^{\text{row}} = 0$ , and applying the scalar projection gives  $S_{b,M} = 0$ . The zero truncation row is then compatible with the defining equations of Definition 8.24.

For genuine seeds and for nonzero added bracket rows, the displayed formulas are additional data beyond the cutoff; they are the required decompositions of the projected normalized row values in the chosen lower-window row basis. Applying  $\pi_{N,P} \pi_{M,N} = \pi_{M,P}$  gives the displayed cocycle equations. Conversely, those equations are exactly the statement that the supplied matrices define a compatible finite-window row system.  $\square$

**PROPOSITION 8.30** (*Projection formulas for the  $u$ ,  $v$ , and  $q$  rows*). Work in a larger finite-window graph system in the local  $\mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$  full-equivariant marked branch. Fix  $M \geq N$ . Suppose the system supplies ordered lower-window row bases, or equivalently row presentations with chosen coordinate maps, for the following three finite-dimensional spans:

$$\begin{aligned} \mathcal{U}_M &= \text{span}\{\mathbf{K}_{\Gamma}^M\}_{\Gamma \in \mathcal{G}_{z,M}^{\text{gen}}}, & \mathbf{K}_{\Gamma}^M &= \varepsilon_{\Gamma} K_{\Gamma}^M, \\ \mathcal{V}_M &= \text{span}\{\mathbf{E}_{\Gamma, \nu}^M\}_{\Gamma, \nu}, & \mathbf{E}_{\Gamma, \nu}^M &= -\varepsilon_{\Gamma, \nu}^{\text{CE}} K_{\Gamma}^M(\zeta_{M, \nu}), \\ \mathcal{B}_M &= \text{span}\{\mathbf{B}_{\beta_1, \beta_2}^M\}_{\beta_i \notin \{\alpha_{\text{sc}}, \alpha_{\text{II}}\}}, & \mathbf{B}_{\beta_1, \beta_2}^M &= \frac{1}{2} \varepsilon_{\beta_1, \beta_2}^M \{K_{\beta_1}^M, K_{\beta_2}^M\}_{\hbar, M}. \end{aligned}$$

Assume these spans are projection-closed:

$$\pi_{M,N} \mathcal{U}_M \subset \mathcal{U}_N, \quad \pi_{M,N} \mathcal{V}_M \subset \mathcal{V}_N, \quad \pi_{M,N} \mathcal{B}_M \subset \mathcal{B}_N,$$

and that  $\pi_{M,N}$  is the chain and BV-bracket projection on the chosen finite-window subcomplex:

$$\pi_{M,N} d_M = d_N \pi_{M,N}, \quad \pi_{M,N} \{x, y\}_{h,M} = \{\pi_{M,N} x, \pi_{M,N} y\}_{h,N}$$

for the rows under consideration.

Then the missing entries isolated above are defined by coordinates of finite-window projection:

$$\pi_{M,N} \mathbf{K}_\Gamma^M = \sum_{\Gamma'} u_{\Gamma, \Gamma'}^{M,N} \mathbf{K}_{\Gamma'}^N,$$

$$\pi_{M,N} \mathbf{E}_{\Gamma, \nu}^M = \sum_{\Gamma', \nu'} v_{\Gamma, \nu; \Gamma', \nu'}^{M,N} \mathbf{E}_{\Gamma', \nu'}^N,$$

and

$$\pi_{M,N} \mathbf{B}_{\beta_1, \beta_2}^M = \sum_{\beta'_1, \beta'_2} q_{\beta_1, \beta_2; \beta'_1, \beta'_2}^{M,N} \mathbf{B}_{\beta'_1, \beta'_2}^N.$$

These equations are the definitions of  $u^{M,N}$ ,  $v^{M,N}$ , and  $q^{M,N}$  relative to the chosen lower-window coordinates. If the displayed lower-window families are honest bases, the entries are unique. If the system supplies only a spanning family with relations, the coordinates are an additional presentation datum that fixes the canonical matrix entry.

The  $u$ -entries also transport the differential rows:

$$\pi_{M,N} (\varepsilon_\Gamma d_M K_\Gamma^M) = \sum_{\Gamma'} u_{\Gamma, \Gamma'}^{M,N} \varepsilon_{\Gamma'} d_N K_{\Gamma'}^N.$$

This is just the first coordinate equation after applying  $d_N \pi_{M,N} = \pi_{M,N} d_M$ .

For the bracket rows there is a useful factorized form. Suppose the added nonzero order-1 rows have transport

$$\pi_{M,N} K_\beta^M = \sum_\eta r_{\beta, \eta}^{M,N} K_\eta^N, \quad \eta \notin \{\alpha_{\text{sc}}, \alpha_{\text{II}}\},$$

and write the lower-window ordered raw bracket reduction as

$$\frac{1}{2} \{K_{\eta_1}^N, K_{\eta_2}^N\}_{h,N} = \sum_{\beta'_1, \beta'_2} \Theta_{\eta_1, \eta_2; \beta'_1, \beta'_2}^N \mathbf{B}_{\beta'_1, \beta'_2}^N.$$

Then

$$q_{\beta_1, \beta_2; \beta'_1, \beta'_2}^{M,N} = \varepsilon_{\beta_1, \beta_2}^M \sum_{\eta_1, \eta_2} r_{\beta_1, \eta_1}^{M,N} r_{\beta_2, \eta_2}^{M,N} \Theta_{\eta_1, \eta_2; \beta'_1, \beta'_2}^N.$$

If the lower system retains all ordered normalized bracket rows with trivial row relations, then

$$\Theta_{\eta_1, \eta_2; \beta'_1, \beta'_2}^N = \frac{\delta_{\eta_1, \beta'_1} \delta_{\eta_2, \beta'_2}}{\varepsilon_{\beta'_1, \beta'_2}^N},$$

and hence

$$q_{\beta_1, \beta_2; \beta'_1, \beta'_2}^{M,N} = \frac{\varepsilon_{\beta_1, \beta_2}^M}{\varepsilon_{\beta'_1, \beta'_2}^N} r_{\beta_1, \beta'_1}^{M,N} r_{\beta_2, \beta'_2}^{M,N}.$$

For  $M \geq N \geq P$ , these coordinate matrices satisfy

$$u_{\Gamma, \Gamma''}^{M,P} = \sum_{\Gamma'} u_{\Gamma, \Gamma'}^{M,N} u_{\Gamma', \Gamma''}^{N,P},$$

$$v_{\Gamma, \nu; \Gamma'', \nu''}^{M, P} = \sum_{\Gamma', \nu'} v_{\Gamma, \nu; \Gamma', \nu'}^{M, N} v_{\Gamma', \nu'; \Gamma'', \nu''}^{N, P},$$

and

$$q_{\beta_1, \beta_2; \theta_1, \theta_2}^{M, P} = \sum_{\beta'_1, \beta'_2} q_{\beta_1, \beta_2; \beta'_1, \beta'_2}^{M, N} q_{\beta'_1, \beta'_2; \theta_1, \theta_2}^{N, P}.$$

Consequently the enlarged-system order-2 non-scalar residual belongs to the compatible finite-window tower only after the following data are supplied: the genuine seed weights, source-face values, nonzero order-1 rows, all orientation/Koszul/symmetry signs, CE-source transport, lower-window row bases or presentations, scalar-condition vanishing or scalar projections, and the projection-closure identities above. With this system the obstruction is exactly the class of the resulting row sum in  $H^1(\mathcal{K}_{\text{ns}, M}^\bullet(I), d_{\text{ns}, M})$ , together with the Roos compatibility class in  $\varprojlim^1 H^0(\mathcal{K}_{\text{ns}, M}^\bullet(I))$ . This system is the data defining the non-scalar QME obstruction class.

*Proof.* The first three displayed equations are simply the coordinate expansion of the projected normalized row values in the chosen lower-window row bases. Projection-closure is exactly the hypothesis that these coordinates exist. If the lower rows are linearly independent, the coordinate expansion is unique. If there are row relations, uniqueness is lost unless the system supplies a presentation or a coordinate splitting.

The  $d\Gamma$ -row formula follows by applying the chain-map identity to  $\pi_{M, N} \mathbf{K}_\Gamma^M$ . The  $q$ -formula follows by applying the BV-bracket identity:

$$\begin{aligned} \pi_{M, N} \mathbf{B}_{\beta_1, \beta_2}^M &= \frac{1}{2} \varepsilon_{\beta_1, \beta_2}^M \{ \pi_{M, N} K_{\beta_1}^M, \pi_{M, N} K_{\beta_2}^M \}_{h, N} \\ &= \frac{1}{2} \varepsilon_{\beta_1, \beta_2}^M \sum_{\eta_1, \eta_2} r_{\beta_1, \eta_1}^{M, N} r_{\beta_2, \eta_2}^{M, N} \{ K_{\eta_1}^N, K_{\eta_2}^N \}_{h, N}, \end{aligned}$$

and then reducing each lower row bracket by the matrix  $\Theta^N$ . The ordered normalized-row specialization is the case where the lower row bracket is already one basis row after division by its incidence coefficient.

The cocycle identities are forced by  $\pi_{N, P} \pi_{M, N} = \pi_{M, P}$ . Expanding first directly from  $M$  to  $P$ , and then through  $N$ , gives two coordinate expansions of the same projected row in the same lower-window coordinates. Equality of coordinates gives the three displayed matrix product formulas. These compatibility identities are precisely the truncation part of Definition 8.24. Once they are supplied, the obstruction criterion is the non-scalar kernel criterion of Definition E.33 and Proposition 8.23.  $\square$

**PROPOSITION 8.31** (*Finite-row primitive search interface*). Let  $\mathbf{A}_{n, M}^{\text{fw}}$  be a valid finite-window graph array at order  $n$ , and suppose its scalar condition has vanished:

$$\sum_{\alpha \in \mathbf{A}_{n, M}^{\text{fw}}} S_{\alpha, M} = 0.$$

Choose an ordered row presentation

$$\rho_1^M, \dots, \rho_a^M$$

for the degree-one non-scalar rows retained at this order: graph differentials, CE-source faces, nonzero bracket rows, and the lower counterterm rows. Write the residual as

$$R_{n, M} = \sum_i r_i^M \rho_i^M.$$

Choose also an ordered finite list of degree-zero counterterm candidates

$$\eta_1^M, \dots, \eta_b^M \in \mathcal{K}_{\text{ns}, M}^{\circ}(I)$$

supplied by the same finite system, and write

$$d_{\text{ns}, M} \eta_j^M = \sum_i A_{ij}^M \rho_i^M.$$

Then a counterterm in the supplied candidate span,  $C_{n, M} = \sum_j c_j \eta_j^M$ , exists if and only if the finite linear system

$$A^M c = -r^M$$

is solvable. When it is obstructed, any covector  $\ell \in (\mathbb{C}^a)^\vee$  satisfying

$$\ell A^M = 0, \quad \ell(r^M) \neq 0$$

separates  $r^M$  from  $\text{im } A^M$ , hence the class of  $R_{n, M}$  is nonzero in the cohomology of the displayed finite row complex.

For a tower  $M \geq N$ , let  $T^{M, N}$  be the row-truncation matrix on the  $\rho$ -presentation and  $P^{M, N}$  the chosen transport matrix on the primitive candidates. The row block  $T^{M, N}$  is the direct sum of the already computed identity or zero blocks, the  $u$ -block for genuine seed rows, the  $v$ -block for CE-source faces, and the  $q$ -block for nonzero bracket rows. A compatible primitive search is defined only when

$$T^{M, N} r^M = r^N, \quad T^{M, N} A^M = A^N P^{M, N}.$$

After windowwise solutions  $c_M$  have been chosen, the remaining compatibility obstruction is the Roos class represented by

$$[P^{M, N} c_M - c_N] \in H^0(\mathcal{K}_{\text{ns}, N}^{\bullet}(I)).$$

Thus the primitive search has three conditions, in this order: scalar condition, fixed-window linear solvability, and Roos  $\varprojlim^1$  compatibility. The last condition is reached only after fixed-window  $H^1$ -obstructions detected by such covectors  $\ell$  vanish; in that case the fixed-window obstruction is decisive.

*Proof.* Once the scalar condition vanishes, the residual lies in the non-scalar kernel complex. Restricting the possible counterterm to the supplied degree-zero span turns the equation  $d_{\text{ns}, M} C_{n, M} = -R_{n, M}$  into the displayed matrix equation. Solvability is equivalent to  $-r^M \in \text{im } A^M$ . In finite dimension this is detected by a linear functional annihilating  $\text{im } A^M$  and pairing nontrivially with  $r^M$ , exactly as written.

The truncation identities say that the row residual and the boundaries of candidate primitives commute with finite-window projection. The computed blocks are those of Propositions 8.29 and 8.30. If  $c_M$  and  $c_N$  solve the two fixed-window equations, then  $P^{M, N} c_M - c_N$  is closed in degree zero by  $T^{M, N} A^M = A^N P^{M, N}$  and  $T^{M, N} r^M = r^N$ . Its cohomology class is the Roos 1-cocycle of Proposition 8.23.  $\square$

In the minimal full-equivariant array all order-one and order-two rows close to zero (Propositions 8.25, 8.26, 8.28); the first non-trivial obstruction therefore appears at order three. The smallest enlargement exhibiting a non-scalar row with nonzero local-functional value is a single CE-source face attached to a genuine marked Costello seed at  $\hbar^3$ , the row  $\theta_3$  of Proposition 8.32, with obstruction class  $\text{Ob}_{\theta_3}^{\Pi}$  and companion-face construction tracked through Theorem 8.35.



PROPOSITION 8.32 (*Concrete order-three row in an enlarged system*). Work in the full-equivariant marked branch with  $V = \mathbb{C}^{N|N}$ , and let  $M \geq 3$ . Enlarge the minimal finite-window system by one Hom-valued CE-source face row

$$\theta_3 = \text{CE}(\Theta_3, \nu_3), \quad |\Theta_3|_h = 3,$$

where  $\Theta_3$  is a genuine marked Costello seed, distinct from a scalar-contact closure composite and from a bracket row. The row data are as follows.

The graph type and order are

$$\tau_{\theta_3} = \text{CE}, \quad |\theta_3|_h = 3.$$

Its finite labels are

$$\begin{aligned} e_{\text{pp}} : & \quad P_{\epsilon, L}^{w, M}, \quad h_{2, \text{I}} = z_1^2 z_2, \quad \rho_{2, \text{I}}, \\ e_{\text{ct}} : & \quad \Delta_{\epsilon, L}^{w, M}, \quad h_{1, \text{I}} = z_1 z_2, \quad \rho_{1, \text{I}}, \\ e_{\text{W}} : & \quad E_{\text{Weyl}}, \quad z_1^2, \quad z_2 \end{aligned}$$

with  $B_\theta \in \mathcal{D}_c^{\text{reg}}(I)$  on the cotangent external leg. The vertex labels are

$$u_{a(t)dt \otimes z_1^2}, \quad u_{b(t)dt \otimes z_2}, \quad c_{B_\theta, \rho_{2, \text{I}}}, \quad I_{\text{o}, M}^w, \quad W_{\Theta_3, L, w}^{R, M}(B_\bullet^\Theta).$$

The marker tensor is a full  $\mathfrak{gl}(N|N)$ -equivariant signed Brauer marker

$$\mathfrak{m}_{\theta_3} \in \mathcal{M}^{\text{full}}(\mathbb{C}^{N|N}), \quad \text{Str}_{\text{cyc}}(\mathfrak{m}_{\theta_3}) = \text{o}.$$

Choose the source-face coefficient

$$d_{\text{CE}} \zeta_M = +\zeta_{M, \nu_3} + \text{other faces}$$

on the  $\Theta_3$ -component. Since the kernel has degree zero, the incidence sign of this CE row in  $-\kappa_{\mathcal{G}, w, I}^M(d_{\text{CE}} \zeta_M)$  is  $-\text{I}$ .

Write

$$K_{\Theta_3}^M = W_{\Theta_3, L, w}^{R, M}(B_\bullet^\Theta).$$

The row value is the nonzero symbolic local functional

$$R_{\theta_3, M}^{\text{row}} = -K_{\Theta_3}^M(\zeta_{M, \nu_3}) =: E_{\theta_3, M} \neq \text{o}.$$

Its scalar condition is

$$S_{\theta_3, M} = \widehat{\sigma}_{\text{sc}, M}(E_{\theta_3, M}) = \text{o}.$$

Indeed, the row lies in the full-equivariant marked complex, so Theorem E.32 kills its scalar projection; also the displayed Weyl half-edge has  $\omega(z_1^2, z_2) = \text{o}$ .

The class-forming closure check is part of the row datum:

$$d_{\text{ns}, M} E_{\theta_3, M} = \text{o}.$$

This equation is part of the valid obstruction-row data; with it supplied, the row defines

$$\mathfrak{o}_{\theta_3, M}^{\text{ns}} = [E_{\theta_3, M}] \in H^1(\mathcal{K}_{\text{ns}, M}^\bullet(I), d_{\text{ns}, M}).$$

The truncation matrix is explicit. For  $M \geq N$ ,

$$\pi_{M,N} E_{\theta_3,M} = \begin{cases} E_{\theta_3,N}, & N \geq 3, \\ 0, & N < 3, \end{cases}$$

hence

$$t_{\theta_3,\theta_3}^{M,N} = 1 \quad (N \geq 3), \quad t_{\theta_3,\alpha'}^{M,N} = 0 \quad (\alpha' \neq \theta_3),$$

and every entry from  $\theta_3$  is zero for  $N < 3$ . These entries satisfy  $t^{M,P} = t^{M,N} t^{N,P}$ .

Therefore this enlarged-system row is a nonzero local-functional cochain with vanishing scalar condition. It is an explicit symbolic obstruction datum. A fixed-window order-three counterterm exists precisely when

$$\mathfrak{o}_{\theta_3,M}^{\text{ns}} = 0, \quad d_M C_{\theta_3,M} = -E_{\theta_3,M}$$

for some  $C_{\theta_3,M} \in \mathcal{K}_{\text{ns},M}^0(I)$ . This row names the class; solving the larger order-three QME branch requires an added primitive or companion rows.

**PROPOSITION 8.33** ( *$\theta_3$  finite-row obstruction criterion*). Keep the one-row order-three system of Proposition 8.32. In the finite row complex supplied there, set

$$\mathcal{K}_{\theta_3,M}^0 = 0, \quad \mathcal{K}_{\theta_3,M}^1 = \mathbb{C} E_{\theta_3,M}, \quad d_{\theta_3,M} = 0,$$

and take residual  $R_{\theta_3,M} = E_{\theta_3,M}$ . The scalar condition has already vanished:

$$\widehat{\sigma}_{\text{sc},M}(E_{\theta_3,M}) = 0.$$

The primitive-search matrix is the  $1 \times 0$  zero matrix and the target vector is  $-E_{\theta_3,M}$ . Hence the equation  $A^M c = -r^M$  has a nonzero obstruction in the displayed finite row complex. The covector

$$\ell_{\theta_3}(E_{\theta_3,M}) = 1$$

annihilates  $\text{im } d_{\theta_3,M} = 0$  and satisfies  $\ell_{\theta_3}(R_{\theta_3,M}) = 1$ . Therefore

$$[E_{\theta_3,M}] \neq 0 \quad \text{in} \quad H^1(\mathcal{K}_{\theta_3,M}^\bullet).$$

This is an exact obstruction statement for the supplied finite row subcomplex. A primitive in a larger local-functional complex requires extra data. To kill the row one must add either a degree-zero candidate  $C_{\theta_3,M}$  with  $d_{\text{ns},M} C_{\theta_3,M} = -E_{\theta_3,M}$ , or additional Hom-valued companion rows whose sum changes the residual vector.

If such a candidate  $C_{\theta_3,M}$  is supplied and truncates by

$$\pi_{M,N} C_{\theta_3,M} = C_{\theta_3,N} \quad (N \geq 3), \quad \pi_{M,N} C_{\theta_3,M} = 0 \quad (N < 3),$$

then the fixed-window search is solved and the Roos  $\varprojlim^1$ -class of this one-row tower is zero. A nonzero truncation defect leaves zero fixed-window obstruction but represents the Roos class by

$$[\pi_{M,N} C_{\theta_3,M} - C_{\theta_3,N}] \in H^0(\mathcal{K}_{\text{ns},N}^\bullet(I)).$$

For the minimal order-two full-equivariant rows and for bracket rows containing  $\alpha_{\text{sc}}$  or  $\alpha_{\text{II}}$ , the same primitive search has residual vector 0, so the zero counterterm is the unique supplied solution and its Roos class is zero.

*Proof.* This is Proposition 8.31 applied to the one-row presentation. The degree-zero row set is empty, so the image of the finite differential is zero. The residual vector is the basis vector  $E_{\theta_3, M}$ , so the displayed covector is a cokernel separating covector. The Roos statement is the same proposition applied after a hypothetical primitive has been added. The minimal order-two and zero-bracket rows have zero residual before cohomology, hence the zero primitive is compatible under truncation.  $\square$

PROPOSITION 8.34 ( $\theta_3$  primitive-entry criterion). Keep the row  $\theta_3 = \text{CE}(\Theta_3, \nu_3)$  and the notation of Propositions 8.32 and 8.33. A proposed degree-zero primitive for this row is accepted by the finite-row theorem only after the row array supplies one of the following two primitive entries.

(P1) A CE-ancestor entry

$$\eta_{\theta_3, M}, \quad d_{\text{CE}} \eta_{\theta_3, M} = \zeta_{M, \nu_3},$$

together with the Hom-valued Bianchi and companion-face terms needed to prove

$$C_{\theta_3, M}^{\text{CE}} = K_{\Theta_3}^M(\eta_{\theta_3, M}) \in \mathcal{K}_{\text{ns}, M}^{\circ}(I), \quad d_{\text{ns}, M} C_{\theta_3, M}^{\text{CE}} = -E_{\theta_3, M}.$$

(P2) A local counterterm entry  $C_{\theta_3, M}^{\text{ct}} \in \mathcal{K}_{\text{ns}, M}^{\circ}(I)$ , with its locality proof, scalar projection, and differential expansion, satisfying

$$d_{\text{ns}, M} C_{\theta_3, M}^{\text{ct}} = -E_{\theta_3, M}.$$

Equivalently, in the ordered one-row presentation

$$\rho_1^M = E_{\theta_3, M}, \quad r^M = (1),$$

the supplied candidate column  $C_{\theta_3, M}$  must have boundary-matrix coefficient

$$A_{\theta_3, C}^M = -1, \quad d_{\text{ns}, M} C_{\theta_3, M} = -E_{\theta_3, M}.$$

An accepted row-data input supplies the displayed differential entry with coefficient  $-1$  for the candidate. Any other input must first be replaced by an explicitly supplied candidate whose differential column is the one above.

The scalar and tower conditions are part of the same datum:

$$\widehat{\sigma}_{\text{sc}, M}(C_{\theta_3, M}) = 0, \quad \pi_{M, N} C_{\theta_3, M} = \begin{cases} C_{\theta_3, N}, & N \geq 3, \\ 0, & N < 3. \end{cases}$$

With the row transport  $T_{\theta_3, \theta_3}^{M, N} = 1$  for  $N \geq 3$  and 0 for  $N < 3$ , these equations give

$$T^{M, N} A^M = A^N P^{M, N}, \quad [P^{M, N} c_M - c_N] = 0 \quad \text{for the solution } c_M = 1.$$

Thus such an input kills the fixed-window row and has zero Roos compatibility class. The one-row complex here has  $K_{\theta_3, M}^{\circ} = 0$ , so Proposition 8.33 remains the active claim.

A complete companion-face table is a different condition. It replaces the one-face residual by a full CE-source-face row sum

$$\sum_{\nu \in F_{\Theta_3, M}} R_{\text{CE}(\Theta_3, \nu), M}^{\text{row}}, \quad \nu_3 \in F_{\Theta_3, M},$$

whose  $\nu_3$ -summand is  $E_{\theta_3, M}$ . Such a table is accepted only if it supplies every face, every CE coefficient, every scalar condition, the  $\nu^{M, N}$ -truncation matrices with their cocycle identity, and the codimension-two closure table, and if

$$\sum_{\nu \in F_{\Theta_3, M}} R_{\text{CE}(\Theta_\nu, \nu), M}^{\text{row}} = 0, \quad \sum_{\nu \in F_{\Theta_3, M}} S_{\text{CE}(\Theta_\nu, \nu), M} = 0.$$

The current one-face table is only an interface for this possibility: its residual is  $E_{\theta_3, M}$ . Hence it leaves the  $\theta_3$  row active and leaves the primitive Roos class unformed. The first source-algebraic companion table containing  $\nu_3$  is recorded in Theorem 8.35; it produces an obstruction datum.

*Proof.* For a primitive input, apply Proposition 8.31 with degree-one basis  $\rho_1^M = E_{\theta_3, M}$  and residual vector  $r^M = (1)$ . The equation  $A^M c = -r^M$  is solved by the supplied candidate with  $c = 1$  exactly when the candidate column is  $A_{\theta_3, C}^M = -1$ . The CE-ancestor and local-counterterm alternatives are precisely two ways of producing such a degree-zero column. The displayed target differential, scalar projection, and support proof supply the boundary matrix entry used by the finite row complex.

The truncation assertion is the identity/zero transport already attached to the row  $E_{\theta_3, M}$ . With the same identity/zero rule for  $C_{\theta_3, M}$ , the matrix equation  $TA = AP$  is tautological: above degree 3 it reads  $(-1) = (-1)$ , and below degree 3 both sides are 0. The chosen coefficients  $c_M = 1$  are therefore compatible, so the Roos representative vanishes.

For a companion-face table the residual vector itself is replaced by the signed row sum of all CE-source faces. The finite-row theorem can test that new row sum only after the table supplies its scalar conditions, truncation coordinates, and codimension-two closure data. A nonzero signed sum remains a residual; a zero signed sum cancels before the primitive search and leaves the primitive Roos class unformed.  $\square$

**THEOREM 8.35** ( $\theta_3$  companion-face construction obstruction). Work in the order-three full-equivariant marked branch of Proposition 8.32, and put  $H_{p,q} = z_1^p z_2^q$ . On the standard symplectic  $\mathbb{C}^2$  with  $\omega = dz_1 \wedge dz_2$  the Hamiltonian Poisson rule is  $\{H_{a,b}, H_{r,s}\} = (as - br)H_{a+r-1, b+s-1}$ ; in particular  $\{H_{1,0}, H_{2,1}\} = H_{2,0}$ . Consider the source-algebra ancestor

$$\zeta_M^0 = u_{a, H_{1,0}} u_{b, H_{0,1}} c_{B_{\theta_3, \rho_{2,1}}}.$$

Its monomial CE differential is the first source-algebraic table of the same external shape whose source faces contain the displayed  $\nu_3$ -row  $u_{a, H_{2,0}} u_{b, H_{0,1}} c_{B_{\theta_3, \rho_{2,1}}}$ . After truncation to Hamiltonian degree  $\leq 3$ , removal of constants, and cancellation of the two  $\rho_{1,1}$ -terms, the nonzero candidate face set is the following eight-face set. With

$$E_\nu^M = -K_{\Theta_3}^M(\zeta_{M, \nu}), \quad E_{\nu_3}^M = E_{\theta_3, M},$$

the table is

| $\nu$                    | $\epsilon_{\Theta_3, \nu}^{\text{CE}}$ | $\zeta_{M, \nu}$                                           | $\epsilon_{\Theta_3, \nu}^{\text{CE}} E_{\nu}^M$ |
|--------------------------|----------------------------------------|------------------------------------------------------------|--------------------------------------------------|
| $\nu_{02}$               | +2                                     | $u_{a, H_{0,1}} u_{b, H_{0,1}} c_{B_{\theta}, \rho_{0,2}}$ | $2E_{\nu_{02}}^M$                                |
| $\nu_{20}$               | -2                                     | $u_{a, H_{1,0}} u_{b, H_{1,0}} c_{B_{\theta}, \rho_{2,0}}$ | $-2E_{\nu_{20}}^M$                               |
| $\nu_{03}$               | +3                                     | $u_{a, H_{0,2}} u_{b, H_{0,1}} c_{B_{\theta}, \rho_{0,3}}$ | $3E_{\nu_{03}}^M$                                |
| $\nu_{12}^{(1)}$         | +2                                     | $u_{a, H_{1,1}} u_{b, H_{0,1}} c_{B_{\theta}, \rho_{1,2}}$ | $2E_{\nu_{12}^{(1)}}^M$                          |
| $\nu_{12}^{(2)}$         | -1                                     | $u_{a, H_{1,0}} u_{b, H_{0,2}} c_{B_{\theta}, \rho_{1,2}}$ | $-E_{\nu_{12}^{(2)}}^M$                          |
| $\nu_3 = \nu_{21}^{(1)}$ | +1                                     | $u_{a, H_{2,0}} u_{b, H_{0,1}} c_{B_{\theta}, \rho_{2,1}}$ | $E_{\theta_3, M}$                                |
| $\nu_{21}^{(2)}$         | -2                                     | $u_{a, H_{1,0}} u_{b, H_{1,1}} c_{B_{\theta}, \rho_{2,1}}$ | $-2E_{\nu_{21}^{(2)}}^M$                         |
| $\nu_{30}$               | -3                                     | $u_{a, H_{1,0}} u_{b, H_{2,0}} c_{B_{\theta}, \rho_{3,0}}$ | $-3E_{\nu_{30}}^M$                               |

Hence the source-algebraic companion residual is

$$R_{\Theta_3, M}^{\text{cand}} = 2E_{\nu_{02}}^M - 2E_{\nu_{20}}^M + 3E_{\nu_{03}}^M + 2E_{\nu_{12}^{(1)}}^M - E_{\nu_{12}^{(2)}}^M + E_{\theta_3, M} - 2E_{\nu_{21}^{(2)}}^M - 3E_{\nu_{30}}^M.$$

This expression is the exact residual of the attempted companion route. Its vanishing is an additional identity.

The induced source-face transport is the diagonal cutoff

$$v_{\nu; \nu'}^{M, N} = \begin{cases} 1, & \nu = \nu' \text{ and } d(\nu) \leq N, \\ 0, & \text{otherwise,} \end{cases}$$

where  $d(\nu_{02}) = d(\nu_{20}) = 2$  and the other six faces have degree 3. It satisfies the cocycle identity  $v^{M, P} = v^{M, N} v^{N, P}$ . For  $N = 2$ , however, the transported residual is

$$\pi_{M, 2} R_{\Theta_3, M}^{\text{cand}} = 2E_{\nu_{02}}^2 - 2E_{\nu_{20}}^2.$$

Thus a fixed-window cancellation at  $M \geq 3$  gives a compatible tower only after the lower-window identity  $E_{\nu_{02}}^2 = E_{\nu_{20}}^2$  is proved in the chosen row basis, or after the lower-window row presentation is changed and its  $\nu$ -matrix is recomputed.

Consequently the companion route is a construction/obstruction theorem with the following exact missing marked Costello datum:

$$\left( d_{\text{CE}} \zeta_M, \epsilon_{\Theta_3, \nu}^{\text{CE}}, K_{\Theta_3}^M(\zeta_{M, \nu}), \mu_{\nu}, v_{\nu; \nu'}^{M, N}, C_{\nu, \nu'}^{(2)} \right)_{\nu, \nu'}.$$

Here  $\mu_{\nu}$  is the marked output and marker-transport entry, and  $C_{\nu, \nu'}^{(2)}$  is the codimension-two source-face incidence/weight table: the two orders of each pair of boundary operations, the output graph, the orientation/Koszul coefficient, the transported marker, and the resulting renormalized row coordinate in the declared degree-one basis. Until this table is supplied and proves both  $R_{\Theta_3, M}^{\text{cand}} = 0$  and its transported lower-window identities, the proved conclusion remains the one-row obstruction of Proposition 8.33.

*Proof.* The Poisson rule  $[H_{a,b}, H_{r,s}] = (as - br)H_{a+r-1, b+s-1}$  gives the displayed source-algebra faces from the differential of the two  $u$ -factors against the current label. In degree  $\leq 3$ , constants are removed

by the Hamiltonian quotient and the two  $\rho_{1,1}$ -faces have opposite coefficients, so the eight listed terms remain. Applying the row convention  $R_{\text{CE}(\Theta_{3,\nu}),M}^{\text{row}} = \epsilon_{\Theta_{3,\nu}}^{\text{CE}} E_{\nu}^M$  gives the signed residual formula, with the  $\nu_3$ -summand equal to  $E_{\theta_3,M}$ .

The diagonal transport is just finite-window projection on the Hamiltonian degree of the  $c$ -label. Such diagonal cutoffs compose, proving the cocycle identity. Projecting to  $N = 2$  deletes every degree-three face and leaves exactly the two degree-two summands. Therefore the lower-window residual is  $2E_{\nu_{02}}^2 - 2E_{\nu_{20}}^2$ .

The passage from this source-algebra table to a Costello cancellation theorem is precisely the missing marked incidence/weight computation. Source coefficients alone leave the renormalized graph weights, marker transports, codimension-two closure signs, or lower-window row coordinates undetermined. Those entries are the displayed missing datum. They are the data by which Definition 8.24 supplies a zero residual vector and Proposition 8.31 still sees the current one-row obstruction.  $\square$

**THEOREM 8.36** (*Finite-window non-scalar curvature criterion*). Fix a finite-window graph/QME system  $(I, w, M, \mathcal{G}_M)$ , and fix the labelled CE cycle  $\zeta_M$  on which the curvature is evaluated. Write  $K_{\Gamma}^M$  for the component of  $\kappa_{\mathcal{G},w,I}^M$  attached to a graph  $\Gamma$ ; for the zero-internal-edge component this is  $\kappa_{h,w,I}^{\text{coef},M}$ , and for a positive graph it is the renormalized weight  $\mathcal{W}_{\Gamma,L,w}^{R,M}$ . If  $|\Gamma|_h$  denotes the  $\hbar$ -order assigned by the finite graph system, then the order- $n$  graph curvature is

$$\begin{aligned} \text{Ob}_{\mathcal{G},M}^{\text{graph}}[n](\zeta_M) &= [\hbar^n] \left( d_M \kappa_{\mathcal{G},w,I}^M(\zeta_M) - \kappa_{\mathcal{G},w,I}^M(d_{\text{CE}} \zeta_M) \right. \\ &\quad \left. + \frac{1}{2} \sum_{(\zeta_M)} \{ \kappa_{\mathcal{G},w,I}^M(\zeta'_M), \kappa_{\mathcal{G},w,I}^M(\zeta''_M) \}_{h,M} \right) \\ &= \sum_{|\Gamma|_h=n} \varepsilon_{\Gamma} d_M K_{\Gamma}^M - \sum_{|\Gamma'|_h=n} \varepsilon_{\Gamma'}^{\text{CE}} K_{\Gamma'}^M \\ &\quad + \frac{1}{2} \sum_{|\Gamma_1|_h+|\Gamma_2|_h=n} \varepsilon_{\Gamma_1, \Gamma_2} \{ K_{\Gamma_1}^M, K_{\Gamma_2}^M \}_{h,M}. \end{aligned}$$

The signs are the orientation signs of the supplied finite graph differential and the CE coproduct. With lower counterterms  $C_{<n,M}$ , the cochain tested at order  $n$  is

$$R_{n,M} = \text{Ob}_{\mathcal{G},M}^{\text{graph}}[n] + [\hbar^n] \left( d_M C_{<n,M} + \frac{1}{2} \{ C_{<n,M}, C_{<n,M} \}_{h,M} \right).$$

Its scalar condition is

$$\mathfrak{s}_{n,M} = \widehat{\sigma}_{\text{sc},M}(R_{n,M}).$$

The first scalar-contact graph whose coefficient is computed in the present local model is the one-cross-contraction boundary graph with two Hamiltonian external legs and one open scalar loop. It has  $\hbar$ -order 1. For a full-equivariant marked boundary generator  $X_{\Gamma,o,m,M}$ , the extracted order-one scalar condition is

$$\mathfrak{s}_{1,M}^{\text{mk}}(X_{\Gamma,o,m,M}) = \sum_{i: r(\beta_i)=1} (-1)^{i-1} \text{Str}_{\text{cyc}}(\mu_{\beta_i}(\mathbf{m})) \omega(f_{\partial_{\beta_i}\Gamma}, g_{\partial_{\beta_i}\Gamma}) \gamma_{\mathbf{1}}.$$

Before the coefficient extraction this is the  $\hbar^1$ -term  $\hbar \mathfrak{s}_{1,M}^{\text{mk}}$ . If  $V = \mathbb{C}^{N|N}$  and the markers are full  $\mathfrak{gl}(V)$ -equivariant, then

$$\text{Str}_{\text{cyc}}(\mu_{\beta_i}(\mathbf{m})) = 0$$

for every elementary boundary term by Lemma E.31; hence

$$\mathfrak{s}_{1,M}^{\text{mk}} = 0.$$

This is a scalar-condition vanishing. A non-scalar primitive requires a separate construction.

After the scalar condition vanishes, the first non-scalar order is the least  $n_o$  for which the cochain  $R_{n_o,M}$  is defined for the chosen graph list and survives the lower-order choices. Its class is

$$[\mathfrak{o}_{n_o,M}^{\text{fw}}] = [R_{n_o,M}] \in H^1(\mathcal{Q}_{\mathcal{G},M}^\bullet(I), d_M).$$

It vanishes exactly when there exists  $C_{n_o,M} \in \mathcal{Q}_{\mathcal{G},M}^\circ(I)$  with  $d_M C_{n_o,M} = -R_{n_o,M}$ . In a compatible finite-window tower, after choosing any such windowwise primitives, the remaining compatibility obstruction is

$$\lambda_{n_o}^{\text{fw}} = \left[ \left( [\pi_{M+1,M} C_{n_o,M+1} - C_{n_o,M}] \right)_M \right] \in \varprojlim_M H^0(\mathcal{Q}_{\mathcal{G},M}^\bullet(I)).$$

Thus compatible finite-window counterterms exist precisely when all fixed-window classes  $[\mathfrak{o}_{n_o,M}^{\text{fw}}]$  vanish and  $\lambda_{n_o}^{\text{fw}} = 0$ .

The abstract existence of  $\mathcal{G}_M$  supplies the graph list but leaves the coefficient array open. To decide  $n_o$ , the class  $[\mathfrak{o}_{n_o,M}^{\text{fw}}]$ , or the Roos class  $\lambda_{n_o}^{\text{fw}}$ , one must supply a valid finite-window coefficient array in the sense of Definition 8.24:

$$A_{n,M}^{\text{fw}}$$

whose rows specify graph type, edge labels, vertex labels, marker tensor, incidence coefficient,  $\hbar$ -order, coefficient degree, scalar-contact filtration, scalar-condition term, row curvature value, and truncation coefficients in  $M$ . This array turns the formal obstruction symbol of Theorem 8.22 into a computation of  $n_o$ , a test of  $[\mathfrak{o}_{n_o,M}^{\text{fw}}] = 0$ , and a defined  $\lambda_{n_o}^{\text{fw}}$ .

*Proof.* The first displayed formula is the definition of the convolution curvature in Definition B.15, evaluated on  $\zeta_M$  and then projected to the  $\hbar^n$ -coefficient. Expanding  $-\kappa d_{\text{CE}}$  and the CE coproduct bracket gives the three graph sums, with exactly the orientation signs supplied by the finite graph system. Adding the lower counterterm terms gives  $R_{n,M}$ , the residual of Theorem 8.22.

The order-one scalar-contact computation is Proposition E.23 applied to each first filtration-lowering marked boundary term of Definition E.28. The full-equivariant marker transport maps preserve the signed Brauer marker span. Lemma E.31 computes the cyclic supertrace of every such output marker as a power of  $\text{sdim } \mathbb{C}^{N|N} = 0$ , and the empty marker word gives the same zero coefficient. Therefore the scalar condition vanishes on the full-equivariant marked complex before any non-scalar cohomology class is formed.

Once the scalar condition is zero,  $R_{n,M}$  lies in  $\mathcal{Q}_{\mathcal{G},M}^\bullet(I) = \ker \widehat{\sigma}_{\text{sc},M}$  exactly as in the proof of Theorem 8.22. A degree-zero counterterm changes  $R_{n,M}$  by a  $d_M$ -boundary, so the fixed-window solvability criterion is precisely the vanishing of its  $H^1$ -class. If primitives are chosen window by window, their truncation differences are closed degree-zero cochains; their classes form the standard Roos cocycle computing  $\varprojlim_M H^0(\mathcal{Q}_{\mathcal{G},M}^\bullet(I))$ . Vanishing of this class is exactly the condition that the primitives can be adjusted to a compatible family.

The final assertion is a dependency statement. The formulas require a finite graph list with order assignment, CE cycle, signs, coproduct decomposition, graph weights, lower counterterms, scalar projection, and truncation maps. Omitting any of these removes a term from the displayed cochain  $R_{n,M}$ , so its first nonzero order and its cohomology class both become undefined.  $\square$

*Remark 8.37 (Scalar-condition boundary).* In the ordinary scalar-reduced  $\mathfrak{gl}_N$  branch, the same two-Hamiltonian one-cross graph gives the order-one scalar class  $N \omega(f, g) \gamma_1$ , equivalently  $N[\bar{c}] \gamma_1$  in cohomology after extracting the  $\hbar^1$ -coefficient. In the even block-diagonal marked complex a single marker  $T = \lambda_0 P_0 + \lambda_1 P_1$  gives  $N(\lambda_0 - \lambda_1) \omega(f, g) \gamma_1$ . These vanish only under the corresponding central-character, balanced, or full-equivariant hypotheses; they are scalar-contact rows.

*Remark 8.38 (Relation to regulator independence).* Theorem 5.81 proves that, after the finite-window Mittag-Leffler hypotheses of Definition 5.75 are imposed, the Hadamard criterion is independent of the admissible exponential weight. This is weight-independence among regulated coefficient products. The unweighted product/direct-sum pair is separated by the endpoint obstruction of Theorem 5.93: the required coefficient Casimir has an obstructed strict product/direct-sum representative.

*Remark 8.39 (Hadamard parametrix versus the Bochner-Martinelli-Koppelman kernel).* The Hadamard parametrix used in Theorem 8.17 controls Costello's heat-kernel propagator  $P_{\epsilon, L}^{\text{base}}$  on the bulk free elliptic complex  $\Omega^\bullet(\mathbb{R}^2) \hat{\otimes} \Omega^{\circ, \bullet}(\mathbb{C}^2)$  by short-time asymptotics in Hadamard normal form, with Hörmander wave-front-set tracking [12, Ch. 5], [36, Ch. 7]. The Bochner-Martinelli-Koppelman kernel

$$K_{\text{BM}}(\zeta, z) = \frac{1}{(2\pi i)^2} \frac{\bar{\eta}_1 d\bar{\eta}_2 - \bar{\eta}_2 d\bar{\eta}_1}{r^4} \wedge d\zeta_1 \wedge d\zeta_2, \quad \eta_j = \zeta_j - z_j, \quad r = |\zeta - z|,$$

is the several-complex-variables homotopy kernel of [38, §5] producing the classical Koppelman formula  $\bar{\partial}H + H\bar{\partial} = \text{id} - i_B p_B$  on bounded  $C^1$  domains ([38, Def. 5.1, Thm. 5.2, Eq. (10)]; Laurent-Thiébaud [37, Ch. III, Def. 1.2, Lem. 1.3]). These are two distinct analytic technologies attached to two different sectors of the local theorem. The heat-kernel parametrix governs the Costello/QME graph expansion of Theorem 8.17; the BMK kernel governs the Dolbeault homotopy on the formal holomorphic polydisk  $\widehat{\mathbb{C}^2}_0$  and the principal-part current sector recorded in Theorem B.9. The formulas remain separate: the Hadamard parametrix uses the bulk Riemannian/Hermitian heat data on  $\mathbb{R}^2 \times \mathbb{C}^2$ , the BMK kernel uses the holomorphic homotopy data on  $\mathbb{C}^2$  alone, and the local theorem combines their separate finite-window outputs through the obstruction vectors stated below.

**THEOREM 8.40 (BMK finite-window obstruction for strict all-window Hamiltonian transfer).** Let  $\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$  be the Hamiltonian Lie algebra of the formal symplectic polydisk  $(\widehat{\mathbb{C}^2}_0, \omega)$ , let

$$\rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 \wedge dz_2, \quad \mathcal{P} = \bigoplus_{\substack{a,b \geq 0 \\ a+b > 0}} \mathbb{C} \rho_{a,b}, \quad \mathcal{P}_{\leq N} = \bigoplus_{0 < a+b \leq N} \mathbb{C} \rho_{a,b},$$

and write  $\pi_N: \mathcal{P} \rightarrow \mathcal{P}_{\leq N}$  for the finite-Matlis truncation. For every  $N \geq 1$ , the following  $\mathbb{C}$ -linear quotient problem has obstruction:

$$q_N: \mathcal{P} \longrightarrow \mathcal{P}_{\leq N}$$

retains  $\mathcal{P}_{\leq N}$  pointwise, kills the high transverse currents  $\rho_{a,b}$  with  $a+b > N$ , and intertwines the full Hamiltonian coadjoint action of  $\mathfrak{h}$  on principal parts. In particular, the support-local quotient problem on the derivative-delta-current module  $\Delta_{a,b} \in \mathcal{D}'^{\circ,2}(B)$  of Theorem B.9 has the same obstruction to carrying the Hamiltonian action onto a finite Matlis target. The obstruction is the Tate-cohomology class of the principal-part module recorded by

$$\text{ob}_{\mathfrak{h}, N} = \pi_N(\mathfrak{h} \cdot \ker \pi_N) \neq 0 \quad \text{in} \quad \mathcal{P}_{\leq N}.$$



The class is exhibited by the explicit boundary-mode identity

$$z_1^{N+2} \cdot \rho_{N+1,0} = -(N+2) \rho_{0,1}, \quad (27)$$

where the high transverse mode  $\rho_{N+1,0}$  lies in the killed window  $\ker \pi_N$  but the low brane-momentum mode  $\rho_{0,1}$  is retained in  $\mathcal{P}_{\leq N}$ . Equivalently, the projected finite-window  $L_\infty$ -escape repair has nonzero unary block

$$\kappa_N(z_1^{N+2}, \rho_{N+1,0}) = -(N+2) \rho_{0,1},$$

and the projected Jacobiator on the triple  $(z_1^{N+2}, z_2, \rho_{N,0})$  is

$$(N+1)(N+2) \rho_{0,1} \neq 0. \quad (28)$$

Therefore strict all-window finite-current Hamiltonian transfer

$$\mathbb{C}[z_1, z_2]_{\text{red}} \longrightarrow \text{finite Matlis window}$$

has a nonzero obstruction for every  $N$ . The full obstruction vector  $\text{Ob}_N^{\text{BMK-curr}}$  of Theorem B.9, with the third entry the displayed  $\text{ob}_{\mathfrak{h},N}$ , is the precise finite-window obstruction datum.

*Proof.* The Hamiltonian coadjoint action of Corollary B.8 gives, for  $f = z_1^p z_2^q$  and  $\rho_{i,j}$ ,

$$f \cdot \rho_{i,j} = -(pj - qi + p - q) \rho_{i-p+1, j-q+1}.$$

Specializing  $(p, q) = (N+2, 0)$  and  $(i, j) = (N+1, 0)$ ,

$$z_1^{N+2} \cdot \rho_{N+1,0} = -(N+2) \rho_{(N+1)-(N+2)+1, 0-0+1} = -(N+2) \rho_{0,1},$$

which is (27). The mode  $\rho_{N+1,0}$  is killed by  $\pi_N$  since  $N+1+0 > N$ ; the mode  $\rho_{0,1}$  is retained in  $\mathcal{P}_{\leq N}$  since  $0+1 \leq N$  for every  $N \geq 1$ . Hence  $\rho_{N+1,0} \in \ker \pi_N$  and  $\pi_N(z_1^{N+2} \cdot \rho_{N+1,0}) = -(N+2) \rho_{0,1} \neq 0$ , so  $\text{ob}_{\mathfrak{h},N} \neq 0$  in  $\mathcal{P}_{\leq N}$ .

A support-local current quotient  $q_N$  intertwining the Hamiltonian action would satisfy  $q_N(\mathfrak{h} \cdot \xi) = \mathfrak{h} \cdot q_N(\xi)$  for every  $\xi \in \mathcal{P}$ ; applied to  $\xi = \rho_{N+1,0}$ , with  $q_N(\rho_{N+1,0}) = 0$  by the killing hypothesis, this would force

$$0 = \mathfrak{h} \cdot q_N(\rho_{N+1,0}) = q_N(\mathfrak{h} \cdot \rho_{N+1,0}) \ni q_N(z_1^{N+2} \cdot \rho_{N+1,0}) = -(N+2) q_N(\rho_{0,1}) = -(N+2) \rho_{0,1},$$

contradicting  $N+2 \neq 0$  and  $\rho_{0,1} \neq 0$  in  $\mathcal{P}_{\leq N}$ . This is the obstruction.

The projected Jacobiator (28) is the nonzero Jacobiator of the truncated bracket: the unprojected coadjoint bracket on  $\mathfrak{h} \ltimes \mathcal{P}[1]$  is a genuine semidirect Lie module and obeys Jacobi; its image after  $\pi_N$  has the displayed obstruction. Concretely, on the triple  $(z_1^{N+2}, z_2, \rho_{N,0})$  the three cyclic terms in the lifted Jacobi sum are computed by the displayed coadjoint rule together with the Hamiltonian Poisson rule  $\{H_{a,b}, H_{r,s}\} = (as - br)H_{a+r-1, b+s-1}$  of the polydisk:

$$\begin{aligned} \{z_1^{N+2}, \{z_2, \rho_{N,0}\}\} &= \{z_1^{N+2}, (N+1)\rho_{N+1,0}\} = -(N+1)(N+2) \rho_{0,1}, \\ \{z_2, \{\rho_{N,0}, z_1^{N+2}\}\} &= \{z_2, 0\} = 0 \quad \text{since } \{\rho_{N,0}, z_1^{N+2}\} = -(z_1^{N+2} \cdot \rho_{N,0}) = 0 \end{aligned}$$

by the negative-index rule of Proposition 8.7,

$$\{\rho_{N,0}, \{z_1^{N+2}, z_2\}\} = \{\rho_{N,0}, (N+2) z_1^{N+1}\} = (N+1)(N+2) \rho_{0,1}.$$

The unprojected sum is zero, as required by Jacobi for the lifted bracket. Apply  $\pi_N$  entrywise: the first cycle's intermediate  $\{z_2, \rho_{N,o}\} = (N+1)\rho_{N+1,o}$  lies in  $\ker \pi_N$  since  $N+1 > N$ , so the projected first cycle is  $\pi_N\{z_1^{N+2}, \pi_N(N+1)\rho_{N+1,o}\} = \pi_N\{z_1^{N+2}, 0\} = 0$ ; the second cycle is already zero; the third cycle is  $\pi_N\{\rho_{N,o}, (N+2)z_1^{N+1}\} = (N+1)(N+2)\rho_{o,1}$  since  $\rho_{o,1} \in \mathcal{P}_{\leq N}$  is retained. The projected sum is therefore  $(N+1)(N+2)\rho_{o,1} \neq 0$ , which is (28). This is the projected escape-return Jacobiator recorded in Theorem B.9 on the same triple; any quotient  $q_N$  absorbing this return as a unary boundary would also collapse  $\rho_{o,1}$  and hence  $\mathcal{P}_{\leq N}$  itself.

The third entry  $\text{ob}_{\mathfrak{h},N}$  of  $\text{Ob}_N^{\text{BMK-curr}}$  in Theorem B.9 is by construction the displayed class. The remaining entries – diagonal current sign, finite-versus-analytic moment kernel, collar factorization compatibility – are independent obstruction coordinates of the same finite-window obstruction statement and are recorded there.  $\square$

**COROLLARY 8.41** (*Pro-Matlis retract as the selected supremum form*). Theorem 8.40 forecloses every choice of finite-Matlis target for a Hamiltonian-equivariant principal-part transfer. Inside the category  $\text{Curr}^{BM}$  of compatible finite-window principal-part current systems with full  $\mathfrak{h}$ -coadjoint action, BMK homotopy data, and uniform  $\mathfrak{m}$ -adic moment bounds (Theorem 8.42), the unique surviving target is the one-pair analytic pro-Matlis retract

$$\widehat{\mathcal{P}}_{\Pi} = \prod_{\substack{a,b \geq 0 \\ a+b > 0}} \mathbb{C}\rho_{a,b}, \quad \mathcal{P}_{\text{an}}(B_1) = \tau(O(B_1)_{\text{red}}^{\vee}) \subset \widehat{\mathcal{P}}_{\Pi},$$

with the normalized analytic contraction  $H_{\chi}^o = (1 - I_{\text{an}}P_{\text{an}})H_{\chi}(1 - I_{\text{an}}P_{\text{an}})$  of Theorem B.9 satisfying

$$\bar{\partial}H_{\chi}^o + H_{\chi}^o\bar{\partial} = \text{id} - I_{\text{an}}P_{\text{an}},$$

and the pro-analytic moment map  $P_{\Pi}^{\text{an}} = \tau P_{\text{an}}$  with compatible finite projections

$$\pi_N P_{\Pi}^{\text{an}} = p_N, \quad \pi_{M,N} p_M = p_N \quad (M \geq N).$$

The pro-Matlis retract is therefore the terminal object of  $\text{Curr}^{BM}$ , the supremum form of the BMK lane. The obstruction theorem 8.40 is the structural selector that proves the necessity of the pro-Matlis category in place of every finite-Matlis target.

**THEOREM 8.42** (*Pro-Matlis universal property*). Let  $\text{Curr}^{BM}$  be the category whose objects are  $\mathfrak{h}$ -equivariant  $\mathbb{C}$ -linear pairs  $(Q, Q^{\text{an}})$  with  $Q^{\text{an}} \subset Q$ , equipped with:

- (a) compatible projections  $\pi_N: Q \rightarrow \mathcal{P}_{\leq N}$  satisfying  $\pi_M = \pi_N \circ \pi_{M,N}$  for  $M \geq N$ ;
- (b) a Hamiltonian  $\mathfrak{h}$ -action through full residue-coadjoint operators on the untruncated map  $\mathfrak{h} \rightarrow \text{End}(Q)$ ;
- (c) a Dolbeault chain homotopy  $H: Q \rightarrow Q[-1]$  with  $\bar{\partial}H + H\bar{\partial} = \text{id} - I \circ P$  for an analytic-submodule retraction  $P: Q \rightarrow Q^{\text{an}}$ ,  $I: Q^{\text{an}} \hookrightarrow Q$ ;
- (d) uniform  $\mathfrak{m}$ -adic moment bounds: for every  $f \in \mathfrak{h}$  and every  $\xi \in Q$  of  $\mathfrak{m}$ -adic order  $\geq n$ ,  $f \cdot \xi$  has  $\mathfrak{m}$ -adic order  $\geq n - r(f)$  for an  $f$ -dependent constant  $r(f)$ .

Morphisms are  $\mathfrak{h}$ -equivariant  $\mathbb{C}$ -linear maps  $(Q, Q^{\text{an}}) \rightarrow (Q', Q'^{\text{an}})$  commuting with all finite projections  $\pi_N, \pi'_N$  and with the analytic-submodule retractions. Then the pro-Matlis pair  $(\widehat{\mathcal{P}}_{\Pi}, \mathcal{P}_{\text{an}}(B_1))$  with the analytic moment map  $P_{\Pi}^{\text{an}}$  of Theorem B.9 is terminal in  $\text{Curr}^{BM}$ : for every object  $(Q, Q^{\text{an}})$  there is a unique morphism

$$(Q, Q^{\text{an}}) \longrightarrow (\widehat{\mathcal{P}}_{\Pi}, \mathcal{P}_{\text{an}}(B_1)).$$

*Proof.* Existence: condition (a) gives a compatible system  $\pi_N: Q \rightarrow \mathcal{P}_{\leq N}$ , defining a unique  $\mathfrak{h}$ -equivariant map  $Q \rightarrow \varprojlim_N \mathcal{P}_{\leq N} = \widehat{\mathcal{P}}_{\Pi}$  by the universal property of the inverse limit. Condition (c) sends the analytic retraction onto  $\mathcal{P}_{\text{an}}(B_1) = \tau(\mathcal{O}(B_1)_{\text{red}}^{\vee})$  by Theorem B.9, and condition (d) ensures the lift is convergent in the  $\mathfrak{m}$ -adic topology of  $\widehat{\mathcal{P}}_{\Pi}$ . Uniqueness: any two such morphisms agree on every  $\mathcal{P}_{\leq N}$ -component by compatibility with  $\pi_N$ , hence on  $\varprojlim_N \mathcal{P}_{\leq N}$ . Hamiltonian equivariance, the chain-homotopy compatibility, and the moment bounds are preserved by the inverse limit. Theorem 8.40 excludes every finite-rank target (those have a strict finite-window restriction of  $\mathfrak{h}$ , violating (b)), and Corollary 8.41 verifies the pro-Matlis pair structure, the normalized Dolbeault contraction  $H_{\chi}^{\circ}$ , and the analytic moment map  $P_{\Pi}^{\text{an}}$  on  $(\widehat{\mathcal{P}}_{\Pi}, \mathcal{P}_{\text{an}}(B_1))$ .  $\square$

*Proof of Corollary 8.41.* The first sentence is Theorem 8.40: the finite-Matlis quotient has the Hamiltonian-action obstruction. The positive constructions of the pro-Matlis pair  $(\widehat{\mathcal{P}}_{\Pi}, \mathcal{P}_{\text{an}}(B_1))$ , the normalized contraction  $H_{\chi}^{\circ}$ , and the moment map  $P_{\Pi}^{\text{an}}$  are Theorem B.9.

Minimality. Any other  $\mathbb{C}$ -linear  $\mathfrak{h}$ -module with a Dolbeault chain homotopy compatible with  $p_N$  and  $i_N$  at every  $N$  must

- (a) accept the full Hamiltonian coadjoint action of  $\mathfrak{h}$ ,
- (b) admit the Dolbeault homotopy obtained from the BMK kernel after the diagonal orientation and collar quotient,
- (c) project compatibly to the finite Matlis windows used by the arity-two coefficient comparison  $\ell_2^{\text{BM}, N}(x, y) = p_N[ix, iy]$ .

Condition (a) excludes every finite-rank quotient by Theorem 8.40; (b) selects the cohomology of the BMK contraction as the moment-restriction kernel, fixing the analytic submodule  $\mathcal{P}_{\text{an}}(B_1) = \tau(\mathcal{O}(B_1)_{\text{red}}^{\vee})$  on which the kernel acts; (c) forces the underlying ambient module to contain every  $\rho_{a,b}$  and to project to  $\mathcal{P}_{\leq N}$ , hence to be  $\widehat{\mathcal{P}}_{\Pi} = \varprojlim_N \mathcal{P}_{\leq N}$ . The pair  $(\widehat{\mathcal{P}}_{\Pi}, \mathcal{P}_{\text{an}}(B_1))$  is therefore the unique smallest target. The downstream theorem surface is the strict all-window factorization-current transfer; it requires the additional vanishing of  $\text{Ob}_{\text{BM}}^{\Pi}$  recorded in Theorem B.9, whose first three nonzero entries (direct-sum moment cokernel, missing compact-current section of the pro-product tower, divergent formal Hamiltonian action on the pro-product) are independent current-topology obstructions that remain alongside the analytic pro-Matlis retract.  $\square$

*Remark 8.43 (Physical interpretation: high transverse holomorphic modes leak back into low brane momentum modes).* The boundary-mode identity (27) of Theorem 8.40 carries a direct physical reading. In the Hamiltonian/Matlis dictionary the index  $\rho_{a,b}$  labels a holomorphic transverse mode of bidegree  $(a, b)$  along  $\mathbb{C}_{\text{hol}}^2$ ; the brane line  $L \subset \mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$  carries the low brane-momentum modes  $\rho_{o,b}$  and their finite Matlis truncations. The class

$$z_1^{N+2} \cdot \rho_{N+1,0} = -(N+2) \rho_{o,1}$$

asserts that under Hamiltonian flow a high transverse holomorphic mode at degree  $(N+1, 0)$  – formally above the cutoff window – transports back into a low brane-momentum mode at degree  $(0, 1)$  – inside the cutoff window. Physically: high transverse holomorphic modes leak back into low brane momentum modes.

This leak is the obstruction for every strict finite-Matlis cutoff: truncating high transverse modes severs the Hamiltonian return channel that would have lifted them to brane momenta, but the return channel is already a finite-window operation by (27), so the image remains. The pro-Matlis retract of Corollary 8.41

is the smallest module closed under this leak: the full Taylor tower retains every high transverse mode and lets the Hamiltonian flow act consistently, while the analytic submodule  $\mathcal{P}_{\text{an}}(B_1)$  supplies the moment-restriction kernel required by the BMK contraction. The arity-two output projection  $\ell_2^{\text{BM},N}(x, y) = p_N[ix, iy]$  of Theorem B.9 is then a finite-window readout of this consistent flow; the finite-rank operation appears after this readout.

The formal-Darboux stalk theorem at one Dirac brane vacuum, the six-coordinate obstruction classification, and the pro-Matlis target that supplies their categorical home are now in place. The identification  $A_{\partial,N}^{\text{cl}} = C_{\text{CE}}^\bullet(\mathfrak{gl}_N, \text{Kosz}([\phi_1, \phi_2])), J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)}, c_f \mapsto \theta_f, u_f \mapsto J(f)$ , is tested at the brane vacua of the central families.

## 9 EXAMPLES

At each closed point  $p \in \mathbb{C}^2$  the formal-Darboux stalk identification of Theorem 5.6 reads

$$\mathcal{A}_{\text{bulk}}^{\text{cl}}|_{b_p} \simeq Z_{E_1}^{\text{der}}(A_{\partial,N}^{\text{cl}}) \simeq HH^\bullet(A_{\partial,N}^{\text{cl}}, A_{\partial,N}^{\text{cl}}), \quad A_{\partial,N}^{\text{cl}} = C_{\text{CE}}^\bullet(\mathfrak{gl}_N, \text{Kosz}([\phi_1, \phi_2])),$$

with formal-Darboux coordinate  $J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)}$ . The two examples below compute  $J$  at the two simplest brane multiplicities.

### 9.0.1 THE SINGLE BRANE: $N = 1$

At  $N = 1$  the matrices commute, the formal-Darboux chart is  $A_b = \widehat{\mathcal{O}}_{\mathbb{C}^2,p} \cong \mathbb{C}[[z_1, z_2]]$  with  $(\phi_1, \phi_2) = (z_1, z_2)$ , and the trace is evaluation:

$$J(z_1^a z_2^b) = z_1^a z_2^b.$$

The closed-to-open dictionary is  $c_f \mapsto \theta_f, u_f \mapsto J(f)$ . Weyl quantisation gives  $[\Phi_h(z_1), \Phi_h(z_2)]_\star = \hbar$ , the one-pair Heisenberg relation; the Capelli scalar  $\hbar N[\bar{c}]|_{N=1} = \hbar[\bar{c}]$  vanishes after scalar reduction.

### 9.0.2 THE BRANE STACK: $N \geq 2$

At a stack of  $N \geq 2$  Dirac branes the trace reads

$$J(z_1^a z_2^b) = \overline{\text{Tr } \phi_1^a \phi_2^b} \in \mathcal{O}(\mu_{\text{der}}^{-1}(\mathfrak{o}))^{GL_N}$$

on the derived commuting variety  $\mu_{\text{der}}^{-1}(\mathfrak{o}) = \{(\phi_1, \phi_2) \in \mathfrak{gl}_N^{\oplus 2} : [\phi_1, \phi_2] = \mathfrak{o}\}$ , with the Procesi–Razmyslov forcing making  $J$  the unique  $GL_N$ -equivariant primitive single-trace Darboux-natural map. The Weyl-quantised commutator is  $[\Phi_h(z_1), \Phi_h(z_2)]_\star = \hbar N \cdot I$  in  $\mathcal{M}_N(\mathbb{C})$ , and the Capelli scalar  $\hbar N[\bar{c}] = \Omega_{\text{central}}|_{J(f)} \in H_{\text{Lie}}^2(\bar{A}, \mathbb{C})[[\hbar]]$  (Theorem 6.8) is the projective curvature of the modular line bundle at the trace generator.

## 10 THE FRONTIER

Theorem 5.6 is the formal-Darboux stalk of the global Mixed Holomorphic–Topological Deligne identification  $\mathcal{A}_{\text{bulk}}^{\text{cl}} \simeq Z_{E_2^{\text{HT}}}^{\text{der}}(C_\partial^{\text{op}})$  of Theorem 5.13, conditional on the three-component global obstruction  $\text{Ob}^{\text{global}} = \text{Ob}_{\text{Deligne}}^{\text{ch } E_2} \oplus \text{Ob}_{\text{KN}}^{\text{desc}} \oplus \text{Ob}_{\text{all-loop}}^{\text{QME}}$ , with an additional modular-line component  $\text{Ob}_{\text{mod}}^V$  for the operator-level lift on  $K_3 \times E$ . Three open problems target these components: the chiral  $E_d$  Deligne theorem at  $d \geq 2$  (§10.1), supplying  $\text{Ob}_{\text{Deligne}}^{\text{ch } E_2}$ ; the Hall–Drinfeld–Borcherds comparison spectral sequence on compact non-toric CY<sub>3</sub> (§10.2), supplying  $\text{Ob}_{\text{KN}}^{\text{desc}} \oplus \text{Ob}_{\text{all-loop}}^{\text{QME}}$ ; the operator-level lift of the modular

line on  $K_3 \times E$  (§10.3), supplying  $\text{Ob}_{\text{mod}}^V$ . Three frontier inputs anchor them: MNOP / framing on  $S^3$  (§10.4); the Hall–Drinfeld realisation on compact  $\text{CY}_3$  (§10.5); the gravitational lift of  $\Phi_{10}$  (§10.6). The cross-target firewall acceptance class  $\text{Ob}_{\text{UKD}}(\mathfrak{T})$  (Theorem 10.4, Theorem 10.13) is the obstruction each problem encounters in transit to a compact target. Volume `frontier_mnop_framing_volume.tex` develops the compact- $\text{CY}_3$  scalar and Fourier–Mukai data (quintic DT/PT factorisation,  $S^3$  framing, chiral-volume finite evidence,  $K_3 \times E$  Fourier–Mukai partner, Gopakumar–Vafa closure) used below as comparison anchors.

### 10.1 Open problem: chiral $E_d$ Deligne conjecture at $d \geq 2$

The Deligne conjecture in its  $E_d$ -form asserts an equivalence

$$Z_{E_d}^{\text{der}}(A) \simeq C_{E_d}^{\bullet}(A, A)$$

between the derived  $E_d$ -centre of an  $E_d$ -algebra  $A$  and the  $E_d$ -Hochschild cochain complex. At  $d = 1$  the topological form is Lurie [41, Thm. 5.3.1.14] (after Tamarkin); for a chiral algebra on a curve  $C$ , the  $d = 1$  chiral form is proved in [28, §4], with the chiral centre  $Z_{\text{ch}}^{\text{der}}(A) \simeq C_{\text{ch}}^{\bullet}(A, A)$  realised through the chiral envelope and chiral cobar; the factorization-algebra reformulation is Costello–Gwilliam [15, Vol. I §5–6]. At  $d \geq 2$  the topological  $E_d$ -centre is constructed by Lurie [41, §5.3] via  $E_d$ -module categories; the chiral upgrade requires three constructions beyond the current literature.

- (i) *A chiral  $E_d$ -operad on  $\mathbb{C}^d$ .* The little holomorphic disks operad  $E_d^{\text{ch}}$  on  $\mathbb{C}^d$  must carry the holomorphic factorization structure of chiral algebras on curves, extended to higher-dimensional affine space. Costello–Gwilliam [15, 16] establish holomorphic factorization algebras on  $\mathbb{C}^d$  as a well-defined symmetric-monoidal  $\infty$ -category; a chiral  $E_d$ -operad action on this category, refining the topological  $E_d$ -action via right  $\bar{\partial}$ -modules, is the missing object.
- (ii) *A centre-of-module-category construction in the chiral setting.* At  $d = 1$  the chiral centre  $Z_{\text{ch}}^{\text{der}}(A)$  is the endomorphism object in chiral  $A$ -modules ([28, §3.4]); at  $d \geq 2$  the analogous construction must take a chiral  $E_d$ -algebra to its derived chiral  $E_d$ -centre, compatibly with factorization homology.
- (iii) *Compatibility with factorization homology  $\text{Sp}_{\Sigma_{d-1}, C}^{\text{ch}}$ .* The two-stage functor  $\Phi_d = \text{Sp}_{\Sigma_{d-1}, C}^{\text{ch}} \circ \Phi_d^{\text{FA}}$  of the master architecture (Lurie–Ayala–Francis factorization homology [41, §5.5]) must intertwine the chiral  $E_d$ -centre on its source with the chiral Hochschild centre on its target.

*Conjecture 10.1 (Chiral  $E_d$  Deligne).* Let  $\mathcal{A}$  be a chiral  $E_d^{\text{ch}}$ -algebra on  $\mathbb{C}^d$  in the sense of (1). There is a natural equivalence

$$Z_{E_d^{\text{ch}}}^{\text{der}}(\mathcal{A}) \simeq C_{E_d^{\text{ch}}}^{\bullet}(\mathcal{A}, \mathcal{A})$$

in chiral  $E_{d+1}^{\text{ch}}$ -algebras, lifting the topological Deligne conjecture of Lurie [41, Thm. 5.3.1.14] along the chiral refinement.

The  $d = 1$  case is proved (Beilinson–Drinfeld [28]; Costello–Gwilliam [15, Vol. I]). At  $d \geq 2$  partial results exist for the topological  $E_d$ -centre ([41, §5.3.1]) and for chiral algebras on smooth schemes (Hennion’s chiral Lie algebroids; Francis’ factorisable co-operads); the chiral  $E_d$ -operad on  $\mathbb{C}^d$  at  $d \geq 2$  in the sense of (1) is open globally. The formal-Darboux stalk theorem of this manuscript (Theorem 5.6), together with Theorem 1.15, supplies the arity-two  $d = 2$  formal-coordinate chiral operation

$$H_{\Delta}^2(\mathbb{C}[[z, w]]) \, d\zeta_1 d\zeta_2 \otimes \mathfrak{g}^{\otimes 2} \longrightarrow \mathfrak{g}, \quad x \otimes y \longmapsto [x, y]_{\mathfrak{g}}.$$

It is the  $d = 2$  formal-Darboux stalk of the Mixed Holomorphic–Topological Deligne identification  $\mathcal{A}_{\text{bulk}}^{\text{cl}} \simeq Z_{E_2^{\text{HT}}}^{\text{der}}(\mathcal{C}_{\partial}^{\text{op}})$ , with  $\mathcal{A}_{\text{bulk}}^{\text{cl}} = \Omega_c^{\circ, \bullet}(B) \otimes \mathfrak{g}, \mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^{\vee}[1], \mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C}$ , the holomorphic  $E_2$  factorization algebra on  $\mathbb{C}^2$ . At the formal-Darboux stalk the bulk identifies with the brane-side derived  $E_1$ -centre  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_{b_p} \simeq Z_{E_1}^{\text{der}}(\mathcal{A}_{\partial, N}^{\text{cl}}) \simeq HH^{\bullet}(\mathcal{A}_{\partial, N}^{\text{cl}}, \mathcal{A}_{\partial, N}^{\text{cl}})$  by Costello–Gwilliam bulk–boundary correspondence and Deligne’s  $E_1$ -Hochschild theorem (both as  $E_2$ -algebras), with formal coordinate  $J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)}$ . The chiral  $E_2$ -Deligne conjecture (Conjecture 10.1 at  $d = 2$ ) is the underlying purely chiral statement on the bulk side; the local theorem supplies the formal-Darboux stalk calculation.

*10.2 Open problem: Hall–Drinfeld–Borcherds comparison spectral sequence on compact non-toric  $CY_3$*

On  $K_3 \times E$  five comparison anchors carry distinct numerical weights:

$$\kappa_{\text{cat}}(K_3 \times E) = 0, \quad \kappa_{\text{ch}}^{\text{Hodge}}(K_3 \times E) = 0,$$

$$\kappa_{\text{ch}}^{\text{Heis}} = 3, \quad \kappa_{\text{BKM}}(\Phi_{10}) = 5, \quad \text{rk } \tilde{H}^*(K_3, \mathbb{Z}) = 24,$$

the categorical Euler invariant  $\kappa_{\text{cat}} = \chi(\mathcal{O}_{K_3 \times E})$ , the chiral Hodge supertrace  $\kappa_{\text{ch}}^{\text{Hodge}} = \chi(\Phi_2^{\text{FA}}(\mathcal{A}_{\text{Hodge}}))$ , the chiral Heisenberg weight  $\kappa_{\text{ch}}^{\text{Heis}}$  of the boundary Heisenberg factor, the Borcherds–Kac–Moody (BKM) denominator weight  $\kappa_{\text{BKM}}(\Phi_{10}) = 5$  recording the lift exponent of  $\Phi_{10} = \chi_{10}$  in the Gritsenko paramodular row, and the extended Mukai-lattice rank of  $K_3$ . These five numbers occupy five distinct construction layers (§10.7, the compact CY-to-chiral target matched-conventions data; `frontier_mnop_framing_volume.tex`, section “Fourier–Mukai partners for  $K_3 \times E$ ”).

*Conjecture 10.2 (Hall–Drinfeld–Borcherds comparison spectral sequence on compact non-toric  $CY_3$ ).* On a compact non-toric Calabi–Yau threefold  $X$  with chosen Hall–Drinfeld–Borcherds matched-conventions datum (the compact CY-to-chiral target; convention contract Appendix 10.7, Theorem 10.13), the five comparison weights organise into the  $E_2$ -page of a spectral sequence

$$E_2^{p,q}(X) = (\kappa_{\text{cat}}, \kappa_{\text{ch}}^{\text{Hodge}}, \kappa_{\text{ch}}^{\text{Heis}}, \kappa_{\text{BKM}}, \kappa_{\text{fibre}})(X) \implies \kappa(X)$$

whose differentials  $d_r: E_r^{p,q} \rightarrow E_r^{p+r, q-r+1}$  are the Hall–Borcherds comparison residuals tracked by the matched-conventions obstruction  $\text{ob}_{\text{Igusa/BKM}} \oplus \text{ob}_{\text{CY-ch}}$  of Theorem 10.4, and whose abutment  $\kappa(X)$  is a single  $\kappa$ -invariant of  $X$ . On  $X = K_3 \times E$  the  $E_2$ -page reads  $(0, 0, 3, 5, 24)$ , and convergence to  $\kappa(K_3 \times E)$  is the conjectural content.

The construction sketch on  $K_3 \times E$  is the Fourier–Mukai partner data of `frontier_mnop_framing_volume.tex`, section “Fourier–Mukai partners for  $K_3 \times E$ ”: under the coprimality hypothesis ( $\star$ ), the universal sheaf gives a derived equivalence  $D^b(K_3) \simeq D^b(M)$  with  $M$  a  $K_3$  partner; the Mukai-lattice side feeds the  $\kappa_{\text{cat}} = 0$ , fibre Euler = 2, and extended-rank = 24 entries through Künneth, and the Heisenberg/BKM entries arrive through the chiral specialisation  $\Phi_2^{\text{FA}}$  along  $K_3$  paired with the Borcherds product realisation of  $\Phi_{10}$  on  $E$ . Equality among the five numbers requires data beyond the Fourier–Mukai theorem alone (`frontier_mnop_framing_volume.tex`, section “Fourier–Mukai partners for  $K_3 \times E$ ”, distinct  $\kappa$  values); the spectral-sequence interpretation is the conjectural content; convergence on a generic compact non-toric  $CY_3$  requires the Hall–Drinfeld–Borcherds construction of the compact CY-to-chiral target and the matched-conventions firewall of Theorem 10.13.

### 10.3 Open problem: operator-level lift of the modular line on $K_3 \times E$

The Capelli scalar  $\hbar N[\bar{c}] = \Omega_{\text{central}}|_{J(f)}$  of §6.1 is the local datum: the projective curvature of the modular line bundle on the period domain  $\overline{\mathcal{A}}_g$ , evaluated at the trace generator. At  $g = 2$  the level-2 section of  $\Omega_{\text{central}}$  is the Igusa cusp form  $\Phi_{10} = \chi_{10}$ , and  $\Delta_5 = \Phi_{10}^{1/2}$  is its square-root branch governing the paramodular Borchers product. An operator-level lift would have to construct an operator on the compact-target bulk theory and identify its protected trace with this modular section. The formal-Darboux stalk supplies only the local fibre of such a construction.

*Conjecture 10.3 (Operator-level lift of  $\Phi_{10}$ ).* Let  $b$  be a Dirac brane vacuum on  $\partial(K_3 \times E)^{\text{KN}}$ , the Kato–Nakayama log boundary of  $K_3 \times E$ . Suppose the compact-target holomorphic-topological bulk theory and its boundary trace have been constructed. The operator-lift problem is to lift the level-2 projective curvature  $\Omega_{\text{central}}|_{\overline{\mathcal{A}}_2}$  to an operator-valued cocycle on that bulk theory,  $\widehat{\Omega}_{\text{central}}^{(2)}$  whose protected scalar trace  $\text{Tr}_{C_j^{\text{op}}}(\widehat{\Omega}_{\text{central}}^{(2)})$  equals  $\Phi_{10} = \chi_{10}$  at the specified paramodular points. The Hall–Borchers comparison residual of Conjecture 10.2, restricted to the  $K_3 \times E$  five-layer page, is the obstruction class to this lift.

The construction problem is the lift of the projective curvature class  $\Omega_{\text{central}}$  from the period domain to an operator on the compact-target bulk module: the local datum of §6.1 supplies only the formal fibre, the modular line bundle on  $\overline{\mathcal{A}}_g$  supplies the global structure, and the Hall–Borchers residual identifies the obstruction. The level-2 scalar section in this comparison is  $\Phi_{10}$  by the Gritsenko paramodular row and the Borchers product realisation (cross-target reference: the Igusa cusp form volume); the operator lift is the construction problem on the bulk side. The cross-target firewall theorem (Theorem 10.4; matched-conventions class  $\text{ob}_{\text{Igusa/BKM}}$ ) records the data which must be supplied for the lift to descend from the bulk module to a section of the modular line bundle.  $\Delta_5$  is the chosen square-root branch on the paramodular cover; the operator-level lift is the open construction problem.

### 10.4 Frontier input: MNOP and framing on $S^3$

On a compact Calabi–Yau threefold  $X$  the Donaldson–Thomas (DT), stable-pair Pandharipande–Thomas (PT), and Gromov–Witten (GW) generating series carry a priori distinct chain-level trace data; the Maulik–Nekrasov–Okounkov–Pandharipande (MNOP) package asserts a single chain-level automorphism  $\sigma_X \in \text{Aut}_{E_2}(Z_{E_2}(\mathcal{F}_X))$  of the  $E_2$ -centre of  $\mathcal{F}_X = \Phi_3^{\text{FA}}(\text{Perf}(X))$  implementing the substitution  $-q = e^{iu}$  on  $(\text{Tr}_{\text{DT}}, \text{Tr}_{\text{PT}}, \text{Tr}_{\text{GW}})$ . The proven half on the smooth quintic is the DT/PT factorisation by the MacMahon power off the degree-zero locus (frontier\_mnop\_framing\_volume.tex, the theorem “DT/PT factorisation on the quintic” in that volume; Bridgeland’s Hall-algebra theorem; Behrend–Fantechi).

The formal-Darboux stalk supplies the formal local trace at one Dirac brane vacuum: the identification  $J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)}$  of Theorem 5.6 feeds the chain-level trace data on a formal neighbourhood of a brane chart on  $X$ . The global gluing across  $X$  requires a framing datum on the Lagrangian boundary; for  $L \cong S^3 \hookrightarrow X$  the framing torsor is  $\pi_3(SO(3)) = \mathbb{Z}$  (Costello brane-line BV framing; frontier\_mnop\_framing\_volume.tex, the proposition “ $S^3$ -framing on the quintic” in that volume). The Calabi–Yau volume form supplies the topological orientation and trivialisation; choice of integer framing class  $[f] \in \pi_3(SO(3)) = \mathbb{Z}$  together with the Costello-compatible  $\text{BV}/\mathcal{A}_\infty$  chain homotopy  $h_f$  trivialising the framing obstruction class  $\text{Ob}_{\text{frame}}(L, f)$  is the global gluing data. The construction problem is the matched-conventions firewall of Theorem 10.13 restricted to the GW/DT/PT/MNOP target with  $S^3$ -framing datum.

### 10.5 Frontier input: Hall–Drinfeld realisation on compact $CY_3$

The chiral specialisation  $\Phi_d = \mathrm{Sp}_{\Sigma_{d-1}, C}^{\mathrm{ch}} \circ \Phi_d^{\mathrm{FA}}$  at  $d = 3$  realises a  $CY_3$  category as a chiral algebra on a curve through the two-stage functor of the master architecture; the Hall–Drinfeld realisation  $\mathcal{G}(X) \simeq D(Y^+(X))$  on the spherical Hall double, after Hall pairing, completion, integral form, stable-envelope transport, and descent, supplies external comparison data for the formal-Darboux stalk theorem. The matched-conventions class  $\mathrm{ob}_{CY\text{-ch}}$  of Theorem 10.4 records the construction data a compact  $CY_3$  target must supply for transfer; the local theorem implies the corresponding compact- $CY_3$  claim only under joint vanishing of  $\mathrm{Ob}_{\mathrm{UKD}}$ .

### 10.6 Frontier input: gravitational lift of $\Phi_{10}$

The Igusa cusp form  $\Phi_{10} = \chi_{10} \in S_{10}(\mathrm{Sp}_4(\mathbb{Z}))$  is the level-2 section of  $\Omega_{\mathrm{central}}$  on  $\overline{\mathcal{A}}_2$ , realised as the Gritsenko paramodular Borchers product, with square-root branch  $\Delta_5 = \Phi_{10}^{1/2}$ . Its conjectural gravitational lift on  $K3 \times E$  (cross-target reference: the Igusa cusp form volume) is the construction of an operator-valued cocycle on the compact-target bulk module whose protected scalar trace would recover  $\Phi_{10}$  at the specified paramodular points (Conjecture 10.3). The cross-target firewall of Theorem 10.4 records the matched-conventions class  $\mathrm{ob}_{\mathrm{Igusa/BKM}}$  which must vanish for the lift to descend; this is external comparison data for the formal-Darboux stalk theorem.

### 10.7 Convention firewall and matched-conventions acceptance

The local theorem of §5.3 lives on

$$\mathbb{R}_{\mathrm{top}}^2 \times \widehat{\mathbb{C}}_{\mathrm{o}, \mathrm{hol}}^2 = \mathbb{R}_{\mathrm{brane}} \times \mathbb{R}_{\mathrm{top}, \mathrm{normal}} \times \widehat{\mathbb{C}}_{\mathrm{o}, \mathrm{hol}}^2,$$

holomorphic dimension  $d = \dim_{\mathbb{C}} \widehat{\mathbb{C}}_{\mathrm{o}}^2 = 2$ , with brane line  $\mathbb{R}_{\mathrm{brane}} \times \{\mathrm{o}\} \times \{\mathrm{o}\}$  as the one-dimensional factorization direction. Seven external programmes carry their own conventions, global data, and obstruction complexes: the compact  $CY_3$  BCOV / Kodaira–Spencer programme of Bershadsky–Cecotti–Ooguri–Vafa [1]; the compact  $CY$ -to-chiral target  $CY$ -to-chiral frontier with two-stage factorization  $\Phi_d = \mathrm{Sp}_{\Sigma_{d-1}, C}^{\mathrm{ch}} \circ \Phi_d^{\mathrm{FA}}$ ; the Borchers / Borchers–Kac–Moody (BKM) bridge to the Igusa cusp form  $\Delta_5$ ; the Gromov–Witten / Donaldson–Thomas / Pandharipande–Thomas / Maulik–Nekrasov–Okounkov–Pandharipande generating-function lane; the Ooguri–Strominger–Vafa attractor counting; the mixed topological–holomorphic six-relation factorization / mixed-Dunn operadic programme; and the global Hamiltonian-descent / Weiss–Ran lane on a compact holomorphic-symplectic surface. The cross-target theorem asserts that the local theorem implies none of these global statements. It contributes exactly the transported formal-disk stalk

$$\chi_{\mathrm{op}}^{-1} \Phi_{\mathrm{fact}}^{\mathrm{Ham}} \chi_{\mathrm{cl}},$$

and external use is governed by the acceptance residual class

$$\mathrm{Ob}_{\mathrm{UKD}}(\mathfrak{C}_{\mathrm{tar}}) \in H^1(\mathcal{K}_{\mathrm{obs}}) \oplus H^1(\mathfrak{R}_{\mathrm{KD}})$$

constructed in Theorem 10.13. A nonzero coordinate is the precise target obstruction.

In the three-volume architecture

$$C_{CY} \xrightarrow{\Phi_d^{\mathrm{FA}}} E_d\text{-HolFA}(X) \xrightarrow{\mathrm{Sp}_{\Sigma_{d-1}, C}} \mathcal{A} \xrightarrow{Z_{E_d^{\mathrm{ch}}}^{\mathrm{der}}} Z_{E_d^{\mathrm{ch}}}^{\mathrm{der}}(\mathcal{A}) \xrightarrow{\mathrm{HT realization}} \mathrm{Obs}_{\mathrm{bulk}}(T_{\mathcal{A}}),$$



where the chiral derived  $E_d$ -centre functor on the right of the specialisation is the  $d = 1$  chiral-algebra theorem ( $Z_{\text{ch}}^{\text{der}} = C_{\text{ch}}^{\bullet}$  on chiral algebras over curves; [28]) and at  $d \geq 2$  the chiral  $E_d$ -Deligne conjecture (Conjecture 10.1, open). The local theorem computes the final arrow on a single formal Darboux normal slice at a brane stack:

$$\text{res}_{L,N} : Z_{\text{ch}}^{\text{der}}(A) \rightarrow C_{\text{CE}}^{\bullet}(\mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^{\vee}[1]), \quad u_f \mapsto J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)}.$$

The pulled-back curvature reads

$$\text{res}_{L,N}(F_{\text{glob}}(\Theta_{\text{glob}})) = (\hbar N[\bar{e}], \Omega_{a,b}^{\text{rad}}, [\text{Ob}_{w,\partial,\hbar}^{\text{red}}], \text{Ob}_{\text{cent/QME/desc}}, \text{Ob}_{\text{BM}}^{\Pi}, \text{Ob}_{\Phi_d\text{-descent}})$$

in the master deformation complex of § 7; cross-target agreement is the assertion that this restriction admits a compatible null-homotopy in  $\mathfrak{R}_{\text{KD}}$ , and a nonzero coordinate is the named obstruction detected by the matched-conventions complex.

The universal centre  $Z_{\text{ch, stage}}^{\text{der}}(A)$  lives before the specialisation functor  $\text{Sp}_{\Sigma_{d-1}, C}$ ; on  $K_3 \times E$  the Heisenberg chiral scalar, the Borchers weight  $\kappa_{\text{BKM}}(\Phi_1) = 5$ , and the Mukai/conductor invariant are post-specialisation shadows of that operator-level identity, distinct on the curve. Numerical matches such as  $K(A) = \kappa_{\text{BKM}}$  are accepted by an external programme through  $\text{Ob}_{\text{UKD}}$ . The local theorem supplies the formal-stalk input.

**THEOREM 10.4 (Cross-target firewall theorem).** Let  $\mathfrak{T}$  be any of the seven external targets enumerated above: compact  $\text{CY}_3$  BCOV / Kodaira–Spencer, the compact CY-to-chiral target CY-to-chiral specialization, Igusa / BKM bridge to  $\Delta_5$ , GW / DT / PT / MNOP, OSV attractor, mixed-Dunn / six-relation, or global Hamiltonian descent on a compact connected holomorphic-symplectic surface. The local theorem on  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}_{\text{o,hol}}^2$  contributes exactly the transported formal-disk stalk

$$\chi_{\text{op}}^{-1} \Phi_{\text{fact}}^{\text{Ham}} \chi_{\text{cl}}$$

to a target morphism, and only after a matched-conventions datum  $\mathfrak{C}_{\text{tar}}$  (Definition 10.7) has been supplied and the universal Koszul-descent acceptance class  $\text{Ob}_{\text{UKD}}(\mathfrak{C}_{\text{tar}})$  of Theorem 10.13 has been null-homotoped. Compact, global, modular, automorphic, specialization, framing, descent, virtual-cycle, attractor, and operadic structures belong to the target data of  $\mathfrak{T}$ . A nonzero coordinate of  $\text{Ob}_{\text{UKD}}$  is the exact obstruction to acceptance in  $\mathfrak{T}$ , recorded coordinate by coordinate.

*Proof.* The formal-stalk content of any cross-target morphism produced by the local theorem is identified by Lemma 10.6: only the displayed transported stalk passes from  $\mathfrak{C}_{\text{loc}}^{(2)}$  through the comparison morphisms. The matched-conventions acceptance criterion is Theorem 10.13: under vanishing of  $\text{Ob}_{\text{UKD}}(\mathfrak{C}_{\text{tar}})$  and a chosen null-homotopy, the target morphism exists and has the displayed formal stalk; otherwise the nonzero coordinate is the exact obstruction. The per-target divergence content is the divergence row of each external target, given respectively by Theorems 10.21, 10.22, 10.23, 10.24, 10.25, 10.26, and 10.27, each naming the local convention layer, the target convention layer, the obstruction class, and the matched-conventions claim. The present theorem records the seven row restrictions simultaneously; independence of divergence rows is a separate question. Lemma 10.6 restricts the local contribution to the formal stalk on every row.  $\square$

The acceptance class  $\text{Ob}_{\mathfrak{C}_{\text{tar}}}$  decomposes into six *universal* coordinates – coordinates every cross-target target must address by the very fact of using a global formal-disk input – and seven *target-specific* coordinates indexed by the external programmes:

$$\begin{aligned}\text{Ob}_{\mathfrak{C}_{\text{tar}}}^{\text{univ}} &= (\text{per}_{\text{tar}}, \text{ob}_{\text{WR}}, \text{ob}_{\text{anom}}, \text{ob}_{\text{QME}}, \text{ob}_{\text{conv}}, \text{ob}_{\text{hyp}}), \\ \text{Ob}_{\mathfrak{C}_{\text{tar}}}^{\text{tar}} &= (\text{ob}_{\text{BCOV}}, \text{ob}_{\text{CY-ch}}, \text{ob}_{\text{Igusa/BKM}}, \text{ob}_{\text{GW/DT}}, \text{ob}_{\text{OSV}}, \text{ob}_{6r}, \text{ob}_{\text{MD}}).\end{aligned}$$

The universal coordinates are: period of locally Hamiltonian fields, Weiss/Ran descent of the local maps, scalar anomaly compatibility, brane-defect quantum master equation, inverse-limit convergence of the obstruction tower, and hyperdescent control of hidden  $\lim^1$  defects. The target-specific coordinates index the seven external programmes: compact  $\text{CY}_3$  BCOV holomorphic-anomaly equation; the compact CY-to-chiral target two-stage CY-to-chiral specialization; Igusa / BKM Borchers-product input; Gromov–Witten / Donaldson–Thomas / Pandharipande–Thomas / MNOP virtual-cycle counts; OSV black-hole attractor identification; mixed topological–holomorphic six-relation factorization; mixed-Dunn operadic rectification. The full acceptance class also carries the transported Maurer–Cartan curvature  $[d\Theta_{\text{KD}} + \frac{1}{2}[\Theta_{\text{KD}}, \Theta_{\text{KD}}]]$ . A coordinate appears exactly for a structure named by  $\mathfrak{C}_{\text{tar}}$ ; a present nonzero coordinate is the exact obstruction to acceptance in  $\mathfrak{C}_{\text{tar}}$ .

The seven external targets are bound to the seven target-specific coordinates of  $\text{Ob}_{\mathfrak{C}_{\text{tar}}}^{\text{tar}}$  by the following correspondence; the universal coordinates apply uniformly across all rows.

| Target programme                                                                                                               | Coordinate                              | Divergence row |
|--------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------|----------------|
| Compact $\text{CY}_3$ BCOV / Kodaira–Spencer [1]                                                                               | $\text{ob}_{\text{BCOV}}$               | Theorem 10.21  |
| the compact CY-to-chiral target CY-to-chiral specialization<br>$\Phi_d = \text{Sp}_{\Sigma_{d-1}, C} \circ \Phi_d^{\text{EA}}$ | $\text{ob}_{\text{CY-ch}}$              | Theorem 10.22  |
| Borchers / BKM bridge to Igusa $\Delta_5$                                                                                      | $\text{ob}_{\text{Igusa/BKM}}$          | Theorem 10.23  |
| GW / DT / PT / MNOP virtual cycles on a compact $\text{CY}_3$                                                                  | $\text{ob}_{\text{GW/DT}}$              | Theorem 10.24  |
| OSV attractor identification $Z^{\text{BH}} \sim  Z^{\text{top}} ^2$                                                           | $\text{ob}_{\text{OSV}}$                | Theorem 10.25  |
| Mixed-Dunn / six-relation factorization on $\mathbb{R} \times \mathbb{C}^2$                                                    | $\text{ob}_{6r}, \text{ob}_{\text{MD}}$ | Theorem 10.26  |
| Global Hamiltonian descent on $(X, \omega, L)$                                                                                 | $\text{Ob}_{\text{glob}}$               | Theorem 10.27  |

Each row carries a divergence theorem comparing the local convention with the target convention, an obstruction class given by the corresponding coordinate, and a matched-conventions claim conditional on a chosen null-homotopy. These coordinates live in syntactically distinct cohomology groups ( $H^1(\mathcal{K}_{\text{obs}})$ ,  $H^1(\mathfrak{R}_{\text{KD}})$ , and the appropriate target-specific complexes); a strict cohomology-level coordinate-independence theorem belongs to the target programme.

*Definition 10.5 (Local Hamiltonian convention base).* The local theorem supplies the convention base

$$\mathfrak{C}_{\text{loc}}^{(2)} = (d = 2, \mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}_{\text{o,hol}}^2, \mathbb{R}_{\text{brane}}, \widehat{\mathfrak{h}}, \widehat{\mathfrak{h}}_{\text{cont}}^{\vee}[1], \mathbf{Ext}^*(\mathcal{O}_{\text{o}}, \mathcal{O}_{\text{o}}), [\bar{c}], \hbar_{\text{Weyl}}, \Phi_{\text{fact}}^{\text{Ham}}).$$

The superscript (2) is the holomorphic dimension  $d = \dim_{\mathbb{C}} \widehat{\mathbb{C}}^2_{\circ}$ . The completed Hamiltonian algebra is  $\widehat{\mathfrak{h}} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$  (formal Hamiltonians modulo constants),  $\widehat{\mathfrak{h}}^{\vee}_{\text{cont}}[1]$  is its continuous cotangent shift,  $[\bar{c}] \in H^2_{\text{Lie}}(\bar{A}, \mathbb{C})$  is the scalar anomaly class carried by the brane line on the reduced Hamiltonian algebra  $\bar{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1$ ,  $\hbar_{\text{Weyl}}$  is the Weyl/Moyal deformation parameter, and  $\Phi^{\text{Ham}}_{\text{fact}}$  is the brane-line interval factorization map produced by the local theorem. The base packages the brane-line factorization convention, the scalar quotient, and the reduced degree-zero Capelli/Moyal normalization.

LEMMA 10.6 (*Formal-polydisk external descent criterion*). The formal-disk theorem contributes the formal stalk used by a global or cross-target descent target  $\mathfrak{T}$  exactly after the target supplies a matched-conventions datum  $\mathfrak{C}_{\text{tar}}$  and a null-homotopy of the acceptance class

$$\text{Ob}_{\text{UKD}}(\mathfrak{C}_{\text{tar}}) = (\text{Ob}^{\text{univ}}_{\mathfrak{C}_{\text{tar}}}, \text{Ob}^{\text{tar}}_{\mathfrak{C}_{\text{tar}}}, [d\Theta_{\text{KD}} + \frac{1}{2}[\Theta_{\text{KD}}, \Theta_{\text{KD}}]])$$

of Theorem 10.13, with each coordinate included exactly for the corresponding structure asserted by  $\mathfrak{T}$ . The accepted morphism has formal stalk

$$\chi_{\text{op}}^{-1} \Phi^{\text{Ham}}_{\text{fact}} \chi_{\text{cl}}.$$

*Proof.* The local input consists of the formal Hamiltonian algebra  $\mathbb{C}[[z_1, z_2]]/\mathbb{C}$ , its continuous cotangent extension, the point brane Ext algebra, the brane-line factorization-current convention, the scalar quotient, and the Capelli/Weyl normalization. These data are invariant under the formal Darboux germ. Compact topology, a negative-cyclic Calabi–Yau class, a worldsheet or specialization curve, an  $S^3$  framing, a global BV propagator, the holomorphic anomaly equation, Borchers input data, a Weyl chamber, a compact Hall orientation, virtual cycles, attractor counts, and compact chiral-homology objects therefore enter as components of  $\mathfrak{C}_{\text{tar}}$  accompanying the formal germ. The displayed coordinates record exactly the universal acceptance tests ( $\text{Ob}^{\text{univ}}_{\mathfrak{C}_{\text{tar}}}$ : period, Weiss/Ran descent, scalar anomaly, brane-defect QME, inverse-limit convergence, hyperdescent), the target-specific acceptance tests ( $\text{Ob}^{\text{tar}}_{\mathfrak{C}_{\text{tar}}}$ : BCOV, the compact CY-to-chiral target, Igusa/BKM, GW/DT/PT/MNOP, OSV, six-relation, mixed-Dunn), and the transported Maurer–Cartan curvature in  $H^1(\mathfrak{R}_{\text{KD}})$ . Vanishing with null-homotopies is the matched-conventions admissibility datum; a nonzero coordinate is the corresponding obstruction class.  $\square$

Definition 10.7 (*Matched-conventions theorem datum*). A matched-conventions theorem datum is a tuple

$$\mathfrak{C}_{\text{tar}} = (d, Y, \eta, (\Sigma_{d-1}, C), L, \tau, \lambda, \mathfrak{f}, \Lambda, \hbar_{\text{tar}}, \mathcal{O}_{\text{tar}}, \chi_{\text{cl}}, \chi_{\text{op}}, \chi_{\text{anom}}, \mathcal{K}_{\text{obs}}, \mathfrak{R}_{\text{KD}}, \Theta_{\text{KD}})$$

with the following meaning. Here  $d = \dim_{\mathbb{C}} Y$ ;  $\eta$  is the negative-cyclic Calabi–Yau class, holomorphic volume form, or holomorphic-symplectic form used by the target theory;  $(\Sigma_{d-1}, C)$  is the worldsheet or factorization-homology specialization when a  $\Phi_d$ -type theorem is asserted;  $L$  is the brane or boundary condition;  $\tau$  is the trace pairing and scalar quotient;  $\lambda$  is the anomaly normalization;  $\mathfrak{f}$  is the framing datum, including an  $S^3$ -framing for compact CY<sub>3</sub> statements;  $\Lambda$  is the coefficient topology;  $\hbar_{\text{tar}}$  is the deformation parameter;  $\mathcal{O}_{\text{tar}}$  is the QME or obstruction hypothesis;  $\chi_{\text{cl}}$ ,  $\chi_{\text{op}}$ , and  $\chi_{\text{anom}}$  are the closed-field, open-field, and anomaly comparison morphisms from the target formal restriction to  $\mathfrak{C}^{(2)}_{\text{loc}}$ ; and  $\mathcal{K}_{\text{obs}}$  is the obstruction complex in which descent, QME, and operadic coherence are solved;  $\mathfrak{R}_{\text{KD}}$  is the filtered dg Lie algebra governing the target global or cross-target descent Maurer–Cartan problem; and  $\Theta_{\text{KD}}$  is the transported local reduced central-operation Maurer–Cartan element whose curvature is the target acceptance class.

*Definition 10.8 (Compact Hamiltonian period complex).* Let  $(X, \omega)$  be a compact connected holomorphic-symplectic surface. For a holomorphic symplectic vector field  $v$ , set

$$\alpha_v = -\iota_v \omega \in H^0(X, \Omega_{X, \text{cl}}^1).$$

Choose a Stein cover  $\mathfrak{U} = \{U_i\}$  and local holomorphic primitives  $h_i \in \mathcal{O}_X(U_i)$  with  $dh_i = \alpha_v|_{U_i}$ . The differences  $h_i - h_j \in \mathbb{C}$  define a Čech cocycle, and its class

$$\text{per}_X(v) = [h_i - h_j] \in H^1(X, \mathbb{C}_X)$$

is the Hamiltonian period obstruction. Equivalently, it is the connecting image of  $\alpha_v$  for the holomorphic de Rham exact sequence

$$0 \longrightarrow \mathbb{C}_X \longrightarrow \mathcal{O}_X \xrightarrow{d} \Omega_{X, \text{cl}}^1 \longrightarrow 0.$$

*Definition 10.9 (Global Hamiltonian descent complex).* Let  $L$  be the brane locus in a proposed global target and suppose that formal Darboux charts have produced local Hamiltonian maps  $\Phi_U$  on a Weiss/Ran cover of  $L$ . Write  $\text{Obs}_{\text{closed}, H}^{\text{cl}}$  for the Hamiltonian closed-sector classical observables and  $Z_{\text{fact}}^{P_o}(\text{Obs}_{\text{open}}^{\text{cl}})$  for the  $P_o$ -factorization centre of the open observables, both taken from the local theorem's reduced Hamiltonian model. The descent target for these maps is the total complex

$$\mathfrak{G}_{\text{WR}}(L) = \text{Tot } C_{\text{WR}}^\bullet \left( \text{Ran}(L); \underline{\text{RHom}} \left( \text{Obs}_{\text{closed}, H}^{\text{cl}}, Z_{\text{fact}}^{P_o}(\text{Obs}_{\text{open}}^{\text{cl}}) \right) \right).$$

The degree-zero Čech-Weiss cochain  $\Phi_{\text{loc}} = \{\Phi_U\}$  has first descent obstruction

$$\text{ob}_{\text{WR}}^{(1)}(\Phi_{\text{loc}}) = [d_{\text{WR}} \Phi_{\text{loc}}] \in H^1(\mathfrak{G}_{\text{WR}}(L)).$$

After choosing a filtration by number of factorization collisions and cotangent insertions, the higher coherence obstructions live in

$$\text{ob}_{\text{WR}}^{(r)}(\Phi_{\text{loc}}) \in H^{r+1}(\text{gr}^r \mathfrak{G}_{\text{WR}}(L)), \quad r \geq 1.$$

Vanishing of all these classes, together with chosen null-homotopies, is the exact assertion that the local Darboux maps descend to a global factorization-centre morphism.

**THEOREM 10.10 (Sharp compact-surface period obstruction).** Let  $(X, \omega)$  be a compact connected holomorphic-symplectic surface and  $v$  a global holomorphic symplectic vector field on  $X$ . Then

$$\text{per}_X(v) = 0 \iff \alpha_v = 0 \iff v = 0.$$

In particular, every nonzero such  $v$  has nonzero period  $\text{per}_X(v) \in H^1(X, \mathbb{C}_X)$ , and is rejected by the global Hamiltonian acceptance criterion before any descent or matched-conventions datum is supplied.

*Proof.* If  $\text{per}_X(v) = 0$ , local Stein primitives  $h_i$  of  $\alpha_v = -\iota_v \omega$  on a Stein cover differ on overlaps by constants exact in the de Rham connecting morphism, hence can be modified by constants to glue to a global holomorphic primitive  $h$ . Since  $X$  is compact and connected,  $H^0(X, \mathcal{O}_X) = \mathbb{C}$ , so  $dh = 0$ , so  $\alpha_v = 0$ , so by nondegeneracy of  $\omega$  gives  $v = 0$ . The converses are immediate.  $\square$

*Example 10.11 (Abelian-surface translation).* A nonzero translation vector field  $v$  on an abelian surface  $(X, \omega)$  has  $\alpha_v \neq 0$ , hence  $\text{per}_X(v) \neq 0$ : local Darboux primitives exist on every chart and carry a nonzero gluing obstruction for a global holomorphic Hamiltonian.

**THEOREM 10.12** (*Formal local-to-global Hamiltonian descent theorem data*). Let  $(X, \omega)$  be a compact connected holomorphic-symplectic surface and  $L \subset X$  a brane locus for which the formal local Hamiltonian theorem is applied along Darboux charts. Fix the finite collection of locally Hamiltonian symplectic vector fields and local Darboux Hamiltonians used by the construction. Assume the target volume supplies the complete pronilpotent descent complex, hyperdescent for the relevant Weiss/Ran cosheaf, and the exact inverse-limit conditions excluding  $\varprojlim^1$  defects in the obstruction tower. Under those supplied convergence and descent hypotheses, the following data are equivalent.

- (i) A global Hamiltonian matched-conventions descent datum: a closed-open morphism

$$\mathrm{Obs}_{\mathrm{closed}, H}^{\mathrm{cl}}|_L \longrightarrow Z_{\mathrm{fact}}^{P_o}(\mathrm{Obs}_{\mathrm{open}}^{\mathrm{cl}})$$

whose Darboux restrictions are the local maps  $\Phi_{\mathrm{loc}}$ , together with its interval, Weiss/Ran, scalar, and connected-stable coherence homotopies.

- (ii) Vanishing of the obstruction vector

$$\mathrm{Ob}_{\mathrm{glob}}(X, L, \Phi_{\mathrm{loc}}) = \left( \{\mathrm{per}_X(v)\}_v, \mathrm{ob}_{\mathrm{scal}}, \mathrm{ob}_{\mathrm{WR}}^{(i)}(\Phi_{\mathrm{loc}}), \{\mathrm{ob}_{\mathrm{WR}}^{(r)}(\Phi_{\mathrm{loc}})\}_{r \geq 1} \right)$$

with null-homotopies compatible with disjoint unions, interval inclusions, cotangent boundary realization,  $P_o$ -centrality, and the stable connected quotient.

The morphism in (i) is the global accepted Koszul-duality surface

$$\mathrm{Obs}_{\mathrm{closed}, H}^{\mathrm{cl}}|_L \longrightarrow Z_{\mathrm{fact}}^{P_o}(\mathrm{Obs}_{\mathrm{open}}^{\mathrm{cl}})$$

and the entries in (ii) are:

- (a) every locally Hamiltonian symplectic vector field used by the construction has period class

$$\mathrm{per}_X(v) = \delta(-\iota_v \omega) \in H^1(X, \mathbb{C}_X)$$

equal to zero;

- (b) the first Weiss/Ran class  $\mathrm{ob}_{\mathrm{WR}}^{(i)}$  vanishes;  
 (c) all higher coherence classes  $\mathrm{ob}_{\mathrm{WR}}^{(r)}$ ,  $r \geq 1$ , vanish after the cotangent boundary realization and  $P_o$ -centrality homotopies have been fixed;  
 (d) the scalar anomaly class agrees with the local class  $[\bar{c}]$ .

When these conditions hold, the local Darboux maps glue, uniquely up to the corresponding contractible space of choices, to the displayed global Hamiltonian central-operation morphism. When one included component is nonzero, that component is the exact universal descent obstruction; every component required by the asserted target structure is part of the target datum.

On a compact connected  $X$  the period condition is sharp by Theorem 10.10: every nonzero global holomorphic symplectic vector field has nonzero  $\mathrm{per}_X(v)$  and is rejected.

*Proof.* The period calculation is Definition 10.8, and the sharpness on compact connected  $X$  is Theorem 10.10: vanishing of  $\mathrm{per}_X(v) = 0$  is the precise condition for gluing local holomorphic Hamiltonian primitives to a global one.

Once local primitives and Darboux charts are fixed, the remaining problem is ordinary descent for morphisms of factorization objects. On double overlaps the local-map defect is  $d_{\mathrm{WR}} \Phi_{\mathrm{loc}}$ , hence the

first obstruction is the displayed class in  $H^1(\mathfrak{G}_{\text{WR}}(L))$ . If it vanishes, a first homotopy can be chosen. The obstruction to making those homotopies coherent on higher overlaps and higher collision strata is computed by the associated graded pieces of the same filtered total complex, giving the classes  $\text{ob}_{\text{WR}}^{(r)}$ . This is the standard obstruction theory for a Maurer–Cartan element in a filtered dg Lie algebra of descent data. Scalar compatibility is separate because the local theorem has already quotiented by the distinguished scalar line. If all classes vanish and null-homotopies have been chosen, the Maurer–Cartan element is gauge equivalent to a global descent datum, which is exactly the displayed morphism.  $\square$

**THEOREM 10.13** (*Conditional matched-conventions transfer criterion*). Let  $\mathfrak{C}_{\text{tar}}$  be a matched-conventions datum whose comparison morphisms identify the formal restriction of the target closed and open theories with  $\mathfrak{C}_{\text{loc}}^{(2)}$ :

$$\chi_{\text{cl}}: \text{Obs}_{\text{closed,tar}}^{\text{cl}}|_{\widehat{p}} \xrightarrow{\sim} \text{Obs}_{\text{closed},H}^{\text{cl}}, \quad \chi_{\text{op}}: \text{Obs}_{\text{open,tar}}^{\text{cl}}|_{\mathbb{R}_{\text{brane}}} \xrightarrow{\sim} \text{Obs}_{\text{open},H}^{\text{cl}}.$$

Suppose

$$\begin{aligned} \text{Ob}_{\mathfrak{C}_{\text{tar}}}^{\text{univ}} &= (\text{per}_{\text{tar}}, \text{ob}_{\text{WR}}, \text{ob}_{\text{anom}}, \text{ob}_{\text{QME}}, \text{ob}_{\text{conv}}, \text{ob}_{\text{hyp}}) = 0, \\ \text{Ob}_{\mathfrak{C}_{\text{tar}}}^{\text{tar}} &= (\text{ob}_{\text{BCOV}}, \text{ob}_{\text{CY-ch}}, \text{ob}_{\text{Igusa/BKM}}, \text{ob}_{\text{GW/DT}}, \text{ob}_{\text{OSV}}, \text{ob}_{6r}, \text{ob}_{\text{MD}}) = 0, \end{aligned}$$

with chosen null-homotopies and with target-specific coordinates included exactly for the corresponding BCOV, the compact CY-to-chiral target, Igusa/BKM, GW/DT/PT/MNOP, OSV, six-relation, or mixed-Dunn structure in  $\mathfrak{C}_{\text{tar}}$ ; the universal coordinates record the target period vanishing, the Weiss/Ran descent of  $\mathfrak{G}_{\text{WR}}(L)$ , the scalar-anomaly identification  $\chi_{\text{anom}}([\text{anom}_{\text{tar}}]) = [\bar{c}]$ , the QME class of  $\mathcal{O}_{\text{tar}}$ , the convergence of the Maurer–Cartan obstruction tower, and the vanishing of  $\lim^1$  defects. When a mixed topological-holomorphic operadic comparison is asserted,  $\delta_{m,n}$  satisfies Theorem 10.17. Let

$$\text{Ob}_{\text{UKD}}(\mathfrak{C}_{\text{tar}}) = \left( \text{Ob}_{\mathfrak{C}_{\text{tar}}}, [d\Theta_{\text{KD}} + \frac{1}{2}[\Theta_{\text{KD}}, \Theta_{\text{KD}}]] \right) \in H^1(\mathcal{K}_{\text{obs}}) \oplus H^1(\mathfrak{R}_{\text{KD}})$$

be the matched-conventions acceptance class of the transported local reduced Hamiltonian central-operation Maurer–Cartan element.

If  $\text{Ob}_{\text{UKD}}(\mathfrak{C}_{\text{tar}}) = 0$  and a null-homotopy of this class has been chosen, then the target descent datum, using the local formal-stalk component, determines an accepted target factorization-centre morphism

$$\Phi_{\mathfrak{C}_{\text{tar}}}: \text{Obs}_{\text{closed,tar}}^{\text{cl}}|_L \longrightarrow Z_{\text{fact}}^{P_o}(\text{Obs}_{\text{open,tar}}^{\text{cl}})$$

whose formal restriction at  $p$  is

$$\chi_{\text{op}}^{-1} \Phi_{\text{fact}}^{\text{Ham}} \chi_{\text{cl}}.$$

If a specialization functor  $\text{Sp}_{\Sigma_{d-1},C}$  is part of  $\mathfrak{C}_{\text{tar}}$ , the specialized morphism is  $\text{Sp}_{\Sigma_{d-1},C}(\Phi_{\mathfrak{C}_{\text{tar}}})$ . If the acceptance class is nonzero, that class is the obstruction to accepting the target theorem in the global or cross-target descent problem.

*Proof.* The comparison morphisms transport the local reduced central-operation map to each Darboux chart of the target. Anomaly compatibility identifies the scalar quotient with the local class  $[\bar{c}]$ , so the transported bracket and connected-trace normalization agree with the Hamiltonian theorem. The vanishing of the target period classes gives Hamiltonian primitives for the local symplectic vector fields in every chart where the Hamiltonian model is used, and the vanishing of the Weiss/Ran classes in

$\mathfrak{G}_{\text{WR}}(L)$  glues the local maps with coherent factorization homotopies by Theorem 10.12. The QME obstruction vanishing upgrades the classical comparison to the target quantum normalization specified by  $\mathcal{O}_{\text{tar}}$ . When a mixed topological-holomorphic operad is present, the mixed-Dunn criterion gives the required equivalence of algebra categories before applying specialization. The displayed formula is the definition of the glued map on the formal stalk. The remaining Maurer–Cartan curvature is exactly  $d\Theta_{\text{KD}} + \frac{1}{2}[\Theta_{\text{KD}}, \Theta_{\text{KD}}]$ ; a chosen null-homotopy makes the transported local datum a global matched-conventions global descent datum. Descent gives the global morphism.  $\square$

*Definition 10.14 (Functor of points of mixed six-relation factorization).* Let  $M$  be a real  $m$ -manifold and  $Y$  a complex  $n$ -fold. For an Artinian commutative dg algebra  $R$ , an  $R$ -point of  $\text{Fact}_{\text{mix}}^{6r}(M, Y)$  is a datum

$$\mathcal{A}_R = (A_R, \mu_R, \ell_R, \bar{\partial}_R, W_R, R_R, \iota_R)$$

on product disks  $D \times B \subset M \times Y$ , where  $D$  is a real disk and  $B$  is a holomorphic polydisk: this is the product-disk site  $\text{Ran}_{\Pi}(M, Y)$ , generated by product disks and finite disjoint unions of such products. Cofinality of product-disk covers inside arbitrary Weiss covers of  $M \times Y$ , and descent on the ordinary  $\text{Ran}(M \times Y)$ , belong to a separate target. A target using arbitrary Weiss/Ran covers must supply  $\text{ob}_{\text{WR}}$ ,  $\text{ob}_{\text{hyp}}$ , and compatible null-homotopies in the matched-conventions obstruction complex.

- (i)  $A_R(D \times B)$  is a nilcomplete  $R$ -linear cochain complex, functorial for inclusions of product disks;
- (ii)  $\mu_R$  gives factorization products for pairwise disjoint finite families of product disks;
- (iii)  $\ell_R$  gives local-constancy homotopies for inclusions of real disks in the  $M$ -direction;
- (iv)  $\bar{\partial}_R$  gives Dolbeault null-homotopies for anti-holomorphic translations in the  $Y$ -direction;
- (v)  $W_R$  gives Weiss descent in the real direction;
- (vi)  $R_R$  gives Ran descent in the holomorphic direction;
- (vii)  $\iota_R$  gives the interchange homotopies between real collision products and holomorphic factorization products.

These data are required to satisfy six relations. Put

$$\text{Rel}^{6r}(\mathcal{A}_R) = (\text{Assoc}(\mu_R), \text{LC}(\ell_R), \bar{\partial}_R^z, d_{\text{WR}}W_R, d_{\text{Ran}}R_R, \text{Int}(\iota_R)).$$

An  $R$ -point is precisely a seven-tuple with  $\text{Rel}^{6r}(\mathcal{A}_R) = 0$ : associativity of  $\mu_R$ , homotopy invariance for  $\ell_R$ , Dolbeault closedness for  $\bar{\partial}_R$ , Weiss cocycle descent for  $W_R$ , Ran cocycle descent for  $R_R$ , and the mixed interchange identity for  $\iota_R$ . A  $k$ -simplex of the derived functor of points is the same datum over  $R \otimes \Omega^\bullet(\Delta^k)$ . Base change in  $R$  defines the functor

$$\text{Fact}_{\text{mix}}^{6r}(M, Y): \text{dgArt}_{\mathbb{C}} \longrightarrow \text{sSet}.$$

This functor is the mixed factorization surface used by the global or cross-target descent problem. Its admissibility condition is the equation  $\text{Rel}^{6r} = 0$  together with coherent null-homotopies in square-zero families. For a partial lift  $\tilde{\mathcal{A}}_{R'}$  over a square-zero extension  $R' \twoheadrightarrow R$  with kernel  $I$ , the six-relation obstruction is

$$\text{ob}_{6r}(\tilde{\mathcal{A}}_{R'}) = [\text{Rel}^{6r}(\tilde{\mathcal{A}}_{R'})] \in H^2(\text{Der}^{\text{fact}, \text{mix}}(\mathcal{A}_R) \otimes_R I).$$

A mixed-Dunn comparison is a morphism of cofibrant operadic models

$$\delta_{m,n}: \mathbb{E}_m^{\text{top}} \otimes \mathbb{E}_n^{\text{hol}} \longrightarrow \mathbb{E}_{m,n}^{\text{mix}}$$

whose induced algebra functor realizes this functor of points.

**PROPOSITION 10.15** (*Recognition and tangent complex for mixed six-relation factorization*). Let  $\mathcal{A}_R$  be an  $R$ -point of  $\text{Fact}_{\text{mix}}^{6r}(M, Y)$ . Then  $\mathcal{A}_R$  determines a mixed factorization object on  $\text{Ran}_{\Pi}(M, Y)$ , locally constant in the  $M$ -directions and holomorphic in the  $Y$ -directions. Conversely, restriction to product disks sends every such mixed factorization object on this product-disk site to an  $R$ -point of  $\text{Fact}_{\text{mix}}^{6r}(M, Y)$ .

If  $R' \twoheadrightarrow R$  is a square-zero extension with kernel  $I$ , the obstruction to lifting  $\mathcal{A}_R$  to an  $R'$ -point is the six-relation class  $\text{ob}_{6r}$ , lying in

$$H^2\left(\begin{array}{c} \text{Der}^{\text{fact}, \text{mix}}(\mathcal{A}_R) \\ \otimes_R I \end{array}\right),$$

and the space of lifts, when nonempty, is a torsor under

$$H^1\left(\begin{array}{c} \text{Der}^{\text{fact}, \text{mix}}(\mathcal{A}_R) \\ \otimes_R I \end{array}\right).$$

*Proof.* Product disks form the objects of the product-disk mixed Weiss/Ran site  $\text{Ran}_{\Pi}(M, Y)$ . The factorization products  $\mu_R$  define the multiplication on disjoint configurations. The homotopies  $\ell_R$  and  $\bar{\delta}_R$  impose the two directional invariance conditions. The descent systems  $W_R$  and  $R_R$  glue the basis values to the mixed Ran object, and  $\iota_R$  is exactly the compatibility between the two collision operations. Nilcompleteness over  $R$  identifies this glued object with the derived functor of points. Restricting a mixed factorization object to product disks recovers the seven pieces of data and the six relations.

For a square-zero extension  $R' \rightarrow R$ , deforming the seven structure maps while preserving the six relations is the Maurer–Cartan lifting problem in the dg Lie algebra of mixed factorization derivations  $\text{Der}^{\text{fact}, \text{mix}}(\mathcal{A}_R) \otimes_R I$ . The curvature of a partial lift is the degree-two component of  $\text{Rel}^{6r}$  for the lifted seven-tuple; its cohomology class is precisely  $\text{ob}_{6r}$  and is independent of choices. If this class vanishes, the remaining choices form the usual  $H^1$ -torsor. This proves the positive recognition criterion for the mixed surface: a target supplies a global or cross-target descent factorization surface exactly by giving an  $R$ -point and killing the six-relation obstruction tower in square-zero thickenings.  $\square$

**Definition 10.16** (*Mixed-Dunn obstruction complex*). The deformation complex of the candidate operad map  $\delta_{m,n}$  is

$$\mathfrak{D}_{\text{MD}}(m, n) = \text{Cone}\left(C_{\text{Operad}}^{\bullet}(\mathbb{E}_{m,n}^{\text{mix}}, \mathbb{E}_{m,n}^{\text{mix}}) \longrightarrow C_{\text{Operad}}^{\bullet}(\mathbb{E}_m^{\text{top}} \otimes \mathbb{E}_n^{\text{hol}}, \mathbb{E}_{m,n}^{\text{mix}})\right)[-1].$$

After a cofibrant filtration of the source operad has been chosen, the mixed-Dunn curvature is

$$F_{\delta} = d\delta + \frac{1}{2}[\delta, \delta] \in Z^1(\mathfrak{D}_{\text{MD}}(m, n)).$$

Its class

$$\text{ob}_{\text{MD}}(\delta_{m,n}) = [F_{\delta}] \in H^1(\mathfrak{D}_{\text{MD}}(m, n))$$

measures the simultaneous obstruction for the topological  $E_m$  operations, the holomorphic  $E_n$  operations, and the mixed interchange law to assemble into a strict operadic comparison. Higher rectification obstructions lie in  $H^{r+1}(\text{gr}^r \mathfrak{D}_{\text{MD}}(m, n))$ .

**THEOREM 10.17** (*Mixed-Dunn matched-conventions criterion*). Let  $\delta_{m,n}$  be a morphism between cofibrant operadic models as in Definition 10.14, equipped with a complete exhaustive cofibrant filtration



for which the obstruction tower below converges and has vanishing hidden  $\varprojlim^1$  defect. The mixed-Dunn surface is accepted by the global or cross-target descent problem exactly under the following equivalent conditions.

- (i)  $\delta_{m,n}$  rectifies to a weak equivalence of operads and induces an equivalence

$$\mathrm{Alg}_{\mathbb{E}_{m,n}^{\mathrm{mix}}} \simeq \mathrm{Alg}_{\mathbb{E}_m^{\mathrm{top}} \otimes \mathbb{E}_n^{\mathrm{hol}}}$$

on the homotopy categories of algebras, with  $\mathrm{Fact}_{\mathrm{mix}}^{6r}(M, Y)$  represented by the tensor product of the topological and holomorphic structures.

- (ii) The aritywise homology map of  $\delta_{m,n}$  is an isomorphism, the primary class  $\mathrm{ob}_{\mathrm{MD}}(\delta_{m,n})$  vanishes, and every higher class in  $H^{r+1}(\mathrm{gr}^r \mathfrak{D}_{\mathrm{MD}}(m, n))$  vanishes with chosen null-homotopies compatible with operadic composition.

A nonzero ordered tuple of these classes is the exact mixed-Dunn acceptance obstruction.

*Proof.* A weak equivalence between cofibrant operads induces a Quillen equivalence on their algebra categories [41, Thm. A.3.7.5]. This proves that (i) implies the aritywise homology isomorphism and that the curvature classes in  $\mathfrak{D}_{\mathrm{MD}}(m, n)$  vanish.

Conversely, suppose (ii) holds. The complete exhaustive cofibrant filtration reduces rectification of  $\delta_{m,n}$  to a convergent tower of lifting problems. The primary curvature  $F_{\delta}$  is the obstruction at the first stage; the associated-graded obstruction groups record the successive obstruction classes for operadic composition, holomorphic-side coherence in the  $\mathbb{E}_n^{\mathrm{hol}}$  factor, and the topological-holomorphic interchange law. By hypothesis each obstruction class is zero and a compatible null-homotopy has been chosen, so the tower has a rectified operad map. The aritywise homology isomorphism makes the rectified map a weak equivalence. Applying the cofibrant-operad Quillen-equivalence theorem gives (i), and Proposition 10.15 identifies the resulting algebra objects with the displayed functor of points.  $\square$

**COROLLARY 10.18** (*Matched-conventions factorization*). Every target theorem that elects to use the local Hamiltonian BF/Moyal theorem through a matched-conventions datum  $\mathfrak{C}_{\mathrm{tar}}$  with  $\mathrm{Ob}_{\mathrm{UKD}}(\mathfrak{C}_{\mathrm{tar}}) = 0$  and chosen null-homotopies has, as its formal-stalk component, the transported morphism

$$\Phi_{\mathfrak{C}_{\mathrm{tar}}} = \chi_{\mathrm{op}}^{-1} \Phi_{\mathrm{fact}}^{\mathrm{Ham}} \chi_{\mathrm{cl}}$$

constructed in Theorem 10.13. The remaining specialization, compactification, automorphic construction, and global descent maps are target data.

*Proof.* Theorem 10.13 constructs the formal-stalk component obtained from the local theorem and the comparison morphisms  $\chi_{\mathrm{cl}}, \chi_{\mathrm{op}}$ . Any subsequent operation belongs to the target datum: specialization along  $(\Sigma_{d-1}, C)$ , compact BV integration, or the Borchers automorphic construction. Thus the local theorem enters the target proof only through the displayed formal-stalk component.  $\square$

**COROLLARY 10.19** (*External target acceptance criterion*). A compact BCOV, MNOP/chiral-volume, the compact CY-to-chiral target CY-to-chiral, Igusa/BKM, or other cross-target theorem may use the local Hamiltonian BF/Moyal formal-stalk input only when supplied with a global or cross-target descent datum  $\mathfrak{C}_{\mathrm{tar}}$  with

$$\mathrm{Ob}_{\mathrm{UKD}}(\mathfrak{C}_{\mathrm{tar}}) = 0$$

and chosen null-homotopies for all components of the obstruction tower. The accepted theorem-data include the formal stalk

$$\chi_{\text{op}}^{-1} \Phi_{\text{fact}}^{\text{Ham}} \chi_{\text{cl}},$$

the global descent datum, the QME/anomaly null-homotopies, and any mixed-Dunn or six-relation factorization null-homotopies asserted by the target. The exact residual of an asserted target structure is the corresponding component of  $\text{Ob}_{\text{UKD}}(\mathfrak{C}_{\text{tar}})$ ; acceptance of any compact  $\text{CY}_3$ , global BCOV, MNOP/chiral-volume, the compact CY-to-chiral target, Igusa, or BKM conclusion follows from killing that class.

*Proof.* Theorem 10.13 constructs the accepted target morphism after the target datum has been fixed and the matched-conventions acceptance class has been null-homotoped. The class contains the period, descent, anomaly, QME, six-relation, and mixed-Dunn components. Lemma 10.6 identifies these components as target theorem data external to the formal polydisk. Therefore a nonzero component is the corresponding obstruction class, and a zero component with a null-homotopy is exactly the accepted theorem datum.  $\square$

*Definition 10.20 (Target convention data).* The cross-target targets require the following matched-conventions data.

(i) *the compact CY-to-chiral target CY-to-chiral datum.*

$$\mathfrak{C}_{\text{CY} \rightarrow \text{ch}} = (d, Y, \eta_{\text{CY}} \in HC_d^-(Y), (\Sigma_{d-1}, C), \Phi_d^{\text{FA}}, \text{Sp}_{\Sigma_{d-1}, C}, \kappa_{\text{cat}}, \tau, \mathfrak{f}, \mathcal{O}_{\text{QME}}, \mathfrak{R}_{\text{KD}}, \text{Ob}_{\text{UKD}}).$$

The comparison map from  $\mathfrak{C}_{\text{loc}}^{(2)}$  is the formal  $d = 2$  Darboux normalization at the chosen brane point. The compact CY-to-chiral target convention is the two-stage factorization

$$\Phi_d = \text{Sp}_{\Sigma_{d-1}, C} \circ \Phi_d^{\text{FA}}.$$

For a compact  $\text{CY}_3$  target the total-space  $\kappa_{\text{cat}}$  is the Hodge supertrace  $\sum_q (-1)^q h^{o,q}(Y)$ ; in particular  $\kappa_{\text{cat}}(K_3 \times E) = 0$ . This component is distinct from any Heisenberg, BKM, or fibre normalization in the compact CY-to-chiral target. Any BKM weight comparison is therefore a separate component of  $\text{Ob}_{\text{UKD}}$ , with the unqualified total-space supertrace kept as a separate component.

(ii) *Compact BCOV datum.*

$$\mathfrak{C}_{\text{BCOV}} = (Y^3, \Omega_Y, \text{PV}^{*,*}(Y)[[t]], Q_{\text{BCOV}}, P_{\text{prop}}, L, \mathfrak{f}_{S^3}, I_{\text{ct}}, \mathcal{O}_{\text{HAE}}, \mathfrak{R}_{\text{KD}}, \text{Ob}_{\text{UKD}}).$$

Here  $Y^3$  is the compact Calabi-Yau threefold,  $\Omega_Y$  the holomorphic volume form,  $\text{PV}^{*,*}(Y)[[t]]$  the polyvector-field BCOV field complex,  $Q_{\text{BCOV}}$  the BCOV differential,  $P_{\text{prop}}$  the global gauge-fixed propagator,  $\mathfrak{f}_{S^3}$  the framing on  $S^3$ ,  $I_{\text{ct}}$  the counterterm scheme, and  $\mathcal{O}_{\text{HAE}}$  the holomorphic-anomaly/QME obstruction complex. The BCOV holomorphic anomaly equation  $\bar{\partial}_i F_g = \frac{1}{2} \bar{C}_i^{jk} (D_j D_k F_{g-1} + \sum_{h+h'=g} D_j F_h D_k F_{h'})$  is a target equation in this column [1]; it lies outside the local  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}_o^2$  theorem surface. Compact Calabi-Yau threefolds and genus expansions belong to the target datum for global BCOV theory.

(iii) *Igusa/BKM datum.*

$$\mathfrak{C}_{\text{BKM}} = (\varphi_{\text{Jac}}, \Lambda_{\text{Borch}}, W, \rho, \text{ch}_{\text{Hall}}, \ell_{\text{or}}, \pi_{\text{prim}}, \kappa_{\text{BKM}}, \tau_{\text{DT/PT/OP}}, \text{Ob}_{\text{UKD}}).$$

The entries are the Jacobi or modular input, Borcherds lattice, Weyl chamber, character, Hall or chiral source, orientation line, primitive recognition map, and scalar DT/PT/OP normalization. In the  $\Delta_5$  row the candidate matched constants are

$$D_5 = 64^{-1} \Delta_5, \quad \text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1} \Delta_5(2Z),$$

$$\kappa_{\text{BKM}}(\Phi_1) = c_1(o)/2 = 10/2 = 5, \quad Z_{\text{OP}}^X = -4096 \Delta_5^{-2}.$$

These constants live in the BKM/Borcherds programme of the  $\Delta_5$  construction; they fix the convention layer at which a target acceptance theorem must match. The local Hamiltonian theorem on  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}_{\text{o,hol}}^2$  supplies the formal-stalk input. The orientation comparison  $\epsilon_o = \nu_{\Delta_5}$ , compact Hall constants, and primitive BKM recognition are target obstructions outside the local  $d = 2$  Hamiltonian output.

(iv) *Global Hamiltonian descent datum.*

$$\mathfrak{C}_{\text{glob}} = (L, \mathcal{H}_X = \mathcal{O}_X/\mathbb{C}_X, \text{per}(v) = \delta(-\iota_v \omega), \mathfrak{G}_{\text{WR}}(L), \{\text{ob}_{\text{WR}}^{(r)}\}_{r \geq 1}, h_{\text{cent}}, \text{Ob}_{\text{glob}}, \Theta_{\text{KD}}).$$

The vanishing condition is the one in Theorem 10.12; its output is the global matched-conventions descent surface.

(v) *GW/DT/PT/MNOP datum.*

$$\mathfrak{C}_{\text{GW/DT}} = (Y^3, \beta \in H_2(Y, \mathbb{Z}), F_{g,\beta}^{\text{GW}}, Z_{n,\beta}^{\text{DT}}, Z_{n,\beta}^{\text{PT}}, \mathfrak{f}_{\text{vir}}, \nu_{\text{Behr}}, \tau_{\text{vir}}, \Phi_{\text{MNOP}}, \text{Ob}_{\text{UKD}}).$$

The entries are the compact  $\text{CY}_3$  target, the curve class, the generating series for the four enumerative theories, the virtual framing, Behrend's microlocal weight, the virtual trace pairing, and the MNOP correspondence map. Their joint identification is the GW/DT/PT/MNOP comparison theorem on  $Y$ . These data belong to the compact target theory. The local  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}_{\text{o,hol}}^2$  theorem has a formal polydisk as holomorphic factor and carries only formal holomorphic coordinates.

(vi) *OSV attractor datum.*

$$\mathfrak{C}_{\text{OSV}} = (Y^3, \Lambda_{\text{att}}, Z_{Q,p}^{\text{BH}}, e^{-S_{\text{att}}}, \Phi_{\text{OSV}}, \text{Ob}_{\text{UKD}}).$$

The OSV conjecture  $Z^{\text{BH}} \sim |Z^{\text{top}}|^2$  and any attractor identification of charges with topological-string genera is a target equation in this column. Its derivation belongs to the OSV target theory, with the local theorem on  $\widehat{\mathbb{C}}_{\text{o,hol}}^2$  contributing only the formal Hamiltonian stalk.

(vii) *Mixed-Dunn datum.*

$$\mathfrak{C}_{\text{MD}} = (\mathbb{E}_m^{\text{top}}, \mathbb{E}_n^{\text{hol}}, \mathbb{E}_{m,n}^{\text{mix}}, \text{Fact}_{\text{mix}}^{6r}, \text{Rel}^{6r}, \delta_{m,n}, \text{ob}_{6r}, \mathfrak{D}_{\text{MD}}(m, n), \text{ob}_{\text{MD}}).$$

The vanishing condition is the simultaneous six-relation and rectification condition in Proposition 10.15 and Theorem 10.17.

#### *Divergence theorems per external target*

The seven matched-conventions templates of Definition 10.20 carry seven divergence theorems. Each names the convention used in the local theorem, the convention used in the target programme, the obstruction class to identification, and the matched-conventions claim that survives once that obstruction is killed by a chosen null-homotopy.

**THEOREM 10.21** (*BCOV divergence and matched-conventions*). Let  $\mathfrak{C}_{\text{BCOV}}$  be the compact BCOV datum of Definition 10.20 (ii).

- (i) *Divergence*. The local theorem on  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}_{\text{o,hol}}^2$  asserts the formal-local Darboux stalk theorem. A compact Calabi-Yau threefold  $Y^3$ , a global gauge-fixed BCOV propagator  $P_{\text{prop}}$ , an  $S^3$ -framing  $\mathfrak{f}_{S^3}$ , a genus expansion in  $F_g$ , and the BCOV holomorphic-anomaly equation are target data. The Bershadsky–Cecotti–Ooguri–Vafa equation

$$\bar{\partial}_{\bar{i}} F_g = \frac{1}{2} \bar{C}_{\bar{i}}^{jk} \left( D_j D_k F_{g-1} + \sum_{h+h'=g} D_j F_h D_k F_{h'} \right)$$

is a target equation in  $\mathfrak{C}_{\text{BCOV}}$  and a proven theorem of [1] on a compact  $\text{CY}_3$ . The local Hamiltonian theorem contributes only the formal-stalk component.

- (ii) *Obstruction class*. The acceptance obstruction is the holomorphic-anomaly / counterterm class

$$\text{ob}_{\text{BCOV}} \in H^1(\mathcal{O}_{\text{HAE}}),$$

computed in the BCOV obstruction complex  $\mathcal{O}_{\text{HAE}}$  of Definition 10.20 (ii).

- (iii) *Matched-conventions claim*. If  $\text{ob}_{\text{BCOV}} = 0$  with a chosen null-homotopy and the universal coordinates  $\text{Ob}_{\mathfrak{C}_{\text{tar}}}^{\text{univ}}$  vanish, then by Theorem 10.13 the BCOV target inherits the formal-disk component  $\chi_{\text{op}}^{-1} \Phi_{\text{fact}}^{\text{Ham}} \chi_{\text{cl}}$  at every Darboux chart of  $Y^3$  where the formal local Hamiltonian model is invoked. Compactification, propagator, framing, and genus-expansion data remain target data.

*Proof.* Clauses (i) and (ii) are Definition 10.20 (ii) read as a divergence and an obstruction class. Clause (iii) is the BCOV column of Theorem 10.13.  $\square$

**THEOREM 10.22** (*the compact CY-to-chiral target CY-to-chiral divergence and matched-conventions*). Let  $\mathfrak{C}_{\text{CY} \rightarrow \text{ch}}$  be the compact CY-to-chiral target datum of Definition 10.20 (i).

- (i) *Divergence*. The compact CY-to-chiral target convention is the two-stage factorization  $\Phi_d = \text{Sp}_{\Sigma_{d-1}, C} \circ \Phi_d^{\text{FA}}$  with worldsheet / specialization curve  $(\Sigma_{d-1}, C)$ , negative-cyclic CY class  $\eta_{\text{CY}} \in H\bar{C}_d^-(Y)$ , and total-space coefficient  $\kappa_{\text{cat}}$ . None of these data is asserted by the local theorem on  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}_{\text{o,hol}}^2$ , whose holomorphic factor is a formal polydisk and whose Calabi-Yau input is the trivial holomorphic symplectic form on  $\mathbb{C}^2$ , used only as a unimodularity datum.
- (ii) *Obstruction class*. The acceptance obstruction is the compact CY-to-chiral target specialization class

$$\text{ob}_{\text{CY} \rightarrow \text{ch}} \in H^1(\mathfrak{R}_{\text{KD}})$$

attached to the comparison of the local  $\mathfrak{C}_{\text{loc}}^{(2)}$ -stalk with the compact CY-to-chiral target formal restriction at the chosen brane point. In the compact  $\text{CY}_3$  row the Hodge supertrace  $\kappa_{\text{cat}} = \Sigma_q (-1)^q h^{o,q}(Y)$  is target datum; the convention check  $\kappa_{\text{cat}}(K_3 \times E) = 0$  is distinct from any Heisenberg, BKM, or fibre normalization in the compact CY-to-chiral target.

- (iii) *Matched-conventions claim*. If  $\text{ob}_{\text{CY} \rightarrow \text{ch}} = 0$  with a chosen null-homotopy and the universal coordinates vanish, then by Theorem 10.13 the specialized morphism  $\text{Sp}_{\Sigma_{d-1}, C}(\Phi_{\mathfrak{C}_{\text{tar}}})$  is defined and has formal stalk  $\chi_{\text{op}}^{-1} \Phi_{\text{fact}}^{\text{Ham}} \chi_{\text{cl}}$ . The remaining specialization, worldsheet, BKM-weight, and Hall-orientation maps are target data.

*Proof.* Clauses (i)–(ii) read off Definition 10.20 (i). Clause (iii) is the compact CY-to-chiral target column of Theorem 10.13, applied with the specialization functor  $\text{Sp}_{\Sigma_{d-1}, C}$  supplied by  $\mathfrak{C}_{\text{tar}}$ .  $\square$

**THEOREM 10.23** (*Igusa / BKM divergence and matched-conventions*). Let  $\mathfrak{C}_{\text{BKM}}$  be the Igusa/BKM datum of Definition 10.20 (iii).

- (i) *Divergence*. The Igusa/BKM convention layer consists of the Jacobi input  $\varphi_{\text{Jac}}$ , Borcherds lattice  $\Lambda_{\text{Borch}}$ , Weyl chamber  $\mathcal{W}$ , character  $\rho$ , Hall character  $\text{ch}_{\text{Hall}}$ , orientation line  $\ell_{\text{or}}$ , primitive recognition map  $\pi_{\text{prim}}$ , and DT/PT/OP normalization  $\tau_{\text{DT/PT/OP}}$ ; these data lie outside the local theorem surface. In the  $\Delta_5$  row, the constants

$$D_5 = 64^{-1} \Delta_5, \quad \text{den}(\mathfrak{g}_{\Delta_5}) = 64^{-1} \Delta_5(2Z),$$

$$\kappa_{\text{BKM}}(\Phi_1) = c_1(\mathfrak{o})/2 = 10/2 = 5, \quad Z_{\text{OP}}^X = -4096 \Delta_5^{-2},$$

live in the BKM/Borcherds programme of the Igusa cusp-form construction; they fix the convention layer at which a target acceptance theorem must match. The local Hamiltonian theorem on  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}_{\mathfrak{o}, \text{hol}}^2$  supplies the formal-stalk input.

- (ii) *Obstruction class*. The acceptance obstruction is the Igusa/BKM Borcherds match class

$$\text{ob}_{\text{Igusa/BKM}} \in H^1(\mathfrak{R}_{\text{KD}})$$

computed in the BKM matched-conventions complex of  $\mathfrak{C}_{\text{BKM}}$ ; it includes the orientation comparison  $\epsilon_{\mathfrak{o}} = \nu_{\Delta_5}$ , compact Hall constants, and primitive BKM recognition. These are target obstructions in the BKM layer.

- (iii) *Matched-conventions claim*. If  $\text{ob}_{\text{Igusa/BKM}} = \mathfrak{o}$  with a chosen null-homotopy and the universal coordinates vanish, then the target inherits the formal-disk component as in Theorem 10.13. The  $\Delta_5$  constants, Borcherds-product factorization, BKM weights, and modular invariance remain target data.

*Proof.* Clauses (i)–(ii) read off Definition 10.20 (iii). Clause (iii) is the Igusa/BKM column of Theorem 10.13.  $\square$

**THEOREM 10.24** (*GW / DT / PT / MNOP divergence and matched-conventions*). Let  $\mathfrak{C}_{\text{GW/DT}}$  be the GW/DT/PT/MNOP datum of Definition 10.20 (v).

- (i) *Divergence*. The GW/DT/PT/MNOP convention layer consists of the compact  $\text{CY}_3$   $Y$ , a curve class  $\beta \in H_2(Y, \mathbb{Z})$ , the four generating series  $F_{g, \beta}^{\text{GW}}, Z_{n, \beta}^{\text{DT}}, Z_{n, \beta}^{\text{PT}}$ , the virtual framing  $\mathfrak{f}_{\text{vir}}$ , Behrend's microlocal weight  $\nu_{\text{Behr}}$ , the virtual trace pairing  $\tau_{\text{vir}}$ , and the MNOP correspondence map  $\Phi_{\text{MNOP}}$ ; these data lie outside the local theorem surface, whose holomorphic factor is the formal polydisk  $\widehat{\mathbb{C}}_{\mathfrak{o}}^2$  and carries only formal holomorphic coordinates.

- (ii) *Obstruction class*. The acceptance obstruction is

$$\text{ob}_{\text{GW/DT}} \in H^1(\mathfrak{R}_{\text{KD}})$$

attached to compatibility of the formal-disk stalk with the virtual GW/DT/PT/MNOP normalization at every Darboux chart where the local Hamiltonian model is invoked in  $\mathfrak{C}_{\text{GW/DT}}$ .

- (iii) *Matched-conventions claim*. If  $\text{ob}_{\text{GW/DT}} = \mathfrak{o}$  with a chosen null-homotopy and the universal coordinates vanish, then the target inherits the formal-disk component as in Theorem 10.13. The virtual cycle, MNOP correspondence, and curve-class generating series remain target data.

*Proof.* Clauses (i)–(ii) read off Definition 10.20 (v). Clause (iii) is the GW/DT/PT/MNOP column of Theorem 10.13.  $\square$

**THEOREM 10.25** (*OSV attractor divergence and matched-conventions*). Let  $\mathfrak{C}_{\text{OSV}}$  be the OSV datum of Definition 10.20 (vi).

- (i) *Divergence.* The OSV convention layer consists of the compact  $\text{CY}_3$   $Y$ , the attractor lattice  $\Lambda_{\text{att}}$ , the black-hole partition  $Z_{Q,p}^{\text{BH}}$ , the attractor entropy  $e^{-S_{\text{att}}}$ , and the OSV identification  $\Phi_{\text{OSV}} : Z^{\text{BH}} \sim |Z^{\text{top}}|^2$ . The OSV identification is a target conjecture on a compact  $\text{CY}_3$ ; it lies outside the local theorem surface on  $\widehat{\mathbb{C}^2}_{\text{o,hol}}$ .
- (ii) *Obstruction class.* The acceptance obstruction is

$$\text{ob}_{\text{OSV}} \in H^1(\mathfrak{R}_{\text{KD}})$$

attached to compatibility of the formal-disk stalk with the attractor identification, when an OSV target invokes the formal-disk component.

- (iii) *Matched-conventions claim.* If  $\text{ob}_{\text{OSV}} = 0$  with a chosen null-homotopy and the universal coordinates vanish, then the target inherits the formal-disk component as in Theorem 10.13. Black-hole attractor counts and the OSV identification remain target data.

*Proof.* Clauses (i)–(ii) read off Definition 10.20 (vi). Clause (iii) is the OSV column of Theorem 10.13.  $\square$

**THEOREM 10.26** (*Mixed-Dunn divergence and matched-conventions*). Let  $\mathfrak{C}_{\text{MD}}$  be the mixed-Dunn datum of Definition 10.20 (vii). The local theorem carries the holomorphic  $E_2$ -type factorization algebra on  $\mathbb{C}^2$  (the formal symplectic surface, with bidegree two in the holomorphic direction), the topological factorization algebra on  $\mathbb{R}_{\text{brane}}$  (one topological direction along the brane line), and a chosen brane-line interval factorization map  $\Phi_{\text{fact}}^{\text{Ham}}$  on product disks  $D \times B \subset \mathbb{R} \times \mathbb{C}^2$ .

- (i) *Divergence.* The local product-disk site  $\text{Ran}_{\Pi}(\mathbb{R}, \mathbb{C}^2)$  is generated by product disks  $D \times B$ ; product-disk covers form a proper subsite of arbitrary Weiss covers of  $\mathbb{R} \times \mathbb{C}^2$ , and the local theorem leaves descent on the unrestricted Weiss site  $\text{Ran}(\mathbb{R} \times \mathbb{C}^2)$  to a separate target. A target using arbitrary Weiss/Ran covers, the operadic interchange  $\delta_{m,n} : \mathbb{E}_m^{\text{top}} \otimes \mathbb{E}_n^{\text{hol}} \rightarrow \mathbb{E}_{m,n}^{\text{mix}}$  with  $(m, n)$  different from the local  $(1, 2)$ , or a strict mixed-operadic rectification, supplies independent data; none of it is a consequence of the local theorem.
- (ii) *Obstruction class.* The acceptance obstruction is the ordered pair

$$(\text{ob}_{6r}, \text{ob}_{\text{MD}}) \in H^2(\text{Der}^{\text{fact,mix}}(\mathcal{A}_R) \otimes I) \oplus H^1(\mathfrak{D}_{\text{MD}}(m, n)).$$

The first coordinate is the six-relation deformation obstruction of Proposition 10.15; the second is the operadic deformation curvature of Definition 10.16. Higher operadic obstructions live in  $H^{r+1}(\text{gr}^r \mathfrak{D}_{\text{MD}}(m, n))$ .

- (iii) *Matched-conventions claim.* If both classes vanish with chosen null-homotopies and the universal coordinates vanish, then by Theorem 10.13 the target inherits the formal-disk component as a strict mixed-operadic factorization map. Operadic rectification at arities other than the local  $(1, 2)$  and Weiss/Ran descent outside the product-disk site remain target data.

*Proof.* Clauses (i)–(ii) read off Definitions 10.20 (vii), 10.14, and 10.16. Clause (iii) is the mixed-Dunn column of Theorem 10.13, combined with Theorem 10.17.  $\square$

**THEOREM 10.27** (*Global Hamiltonian descent divergence and matched-conventions*). Let  $\mathfrak{C}_{\text{glob}}$  be the global descent datum of Definition 10.20 (iv).

(i) *Divergence*. The local theorem is stated on the formal Darboux chart  $\widehat{\mathbb{C}}^2_{\mathfrak{o}}$ ; the global descent datum  $\mathfrak{C}_{\text{glob}}$  lives on a compact connected holomorphic-symplectic surface  $(X, \omega)$  with brane locus  $L$ , a Stein cover  $\mathfrak{U}$ , the de Rham extension  $\mathfrak{o} \rightarrow \mathbb{C}_X \rightarrow \mathcal{O}_X \rightarrow \Omega^1_{X, \text{cl}} \rightarrow \mathfrak{o}$ , and the total complex  $\mathfrak{G}_{\text{WR}}(L)$  of Definition 10.9. None of this global structure is supplied by the formal polydisk.

(ii) *Obstruction class*. The acceptance obstruction is the global descent vector

$$\text{Ob}_{\text{glob}}(X, L, \Phi_{\text{loc}}) \in H^1(X, \mathbb{C}_X) \oplus H^1(\mathfrak{G}_{\text{WR}}(L)) \oplus \bigoplus_{r \geq 1} H^{r+1}(\text{gr}^r \mathfrak{G}_{\text{WR}}(L)) \oplus H^2_{\text{Lie}}(\tilde{A}, \mathbb{C}),$$

with first coordinate the period  $\text{per}_X(v)$  – sharp by Theorem 10.10 on a compact connected surface – the second coordinate the first Weiss/Ran descent class, the third the higher coherence classes, and the fourth the scalar anomaly compatibility class.

(iii) *Matched-conventions claim*. Vanishing of  $\text{Ob}_{\text{glob}}$  with chosen null-homotopies is equivalent to a global Hamiltonian central-operation morphism on  $L$  whose Darboux restrictions are the local maps  $\Phi_{\text{loc}}$ , by Theorem 10.12.

*Proof.* Clauses (i)–(iii) are Definition 10.20 (iv) and Theorem 10.12 read as a divergence and a matched-conventions claim.  $\square$

The three Tate-P residuals form a sequential acceptance criterion. The Hadamard / Mittag-Leffler class  $\text{Ob}_w^{\text{HML}}$  of §7.5 controls universal summation on the weighted Tate envelope; the universality class  $\text{Ob}_{p,X}^{\text{univ}}$  of §5.6 proves that an ambient twisted-M sector restricts canonically to the same formal local model; the matched-conventions class  $\text{Ob}_{\text{UKD}}(\mathfrak{C}_{\text{tar}})$  proves acceptance of that formal local model by an external compact, modular, automorphic, virtual-cycle, or operadic target. Vanishing in sequence – H/ML at the analytic regulator,  $\text{ob}_{p,X}^{\text{univ}}$  at the ambient-to-formal restriction,  $\text{Ob}_{\text{UKD}}$  at the formal-to-target acceptance – is the precise condition under which a target theorem uses the local Hamiltonian formal-stalk component; a nonvanishing class at any layer is the exact obstruction.

*The closing identification.* The trace map  $J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)}$  is the formal Darboux coordinate expression of the brane-side derived  $E_1$ -Hochschild  $HH^\bullet(A_{\partial, N}^{\text{cl}}, A_{\partial, N}^{\text{cl}})$  at one Dirac brane vacuum. The Mixed Holomorphic–Topological Deligne conjecture identifies this at the formal-Darboux stalk with the bulk  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_{b_p}$  (the chiral  $E_2$  factorization algebra on  $\mathbb{C}^2$  restricted to the formal neighbourhood of  $b_p$ ) with two-dimensional singular product

$$J_x(z)J_y(w) \sim K_{\Delta}^{(2)}(z, w)J_{[x,y]_{\mathfrak{g}}}(w),$$

generalising the chiral Deligne theorem at  $d = 1$  for chiral algebras on algebraic curves. The global identification at  $d = 2$ , the operator-level lift of the modular line on  $K3 \times E$ , and the chiral  $E_d$  Deligne theorem at  $d \geq 2$  are the open problems the formal-Darboux stalk has now made well-posed.

## ACKNOWLEDGEMENTS

This work belongs to the lineage of Bershadsky–Cecotti–Ooguri–Vafa 1993, Witten 1988, Costello 2011, and Beilinson–Drinfeld 2004. The Capelli–Howe scalar identities, Procesi–Razmyslov trace relations, and Harish-Chandra radial-parts theorem supply the formal-stalk-chart side; Matlis 1958 and Hartshorne local cohomology supply the principal-part dual; Kontsevich’s deformation quantisation supplies the all-order Moyal target.

## A HIGHER FACTORIZATION CATEGORIES AND CHIRAL DERIVED CENTRES AT $d \geq 2$

The local theorem (Theorem 5.6) computes the brane-side derived  $E_1$ -Hochschild  $HH^\bullet(A_{\partial,N}^{\text{cl}}, A_{\partial,N}^{\text{cl}})$  at the formal-Darboux vacuum  $b_p \in C_\partial^{\text{op}}$ . The Mixed Holomorphic–Topological Deligne conjecture identifies the same object with the chiral derived centre of the closed-string sector at  $d = 2$ . This appendix constructs the higher factorization-categorical machinery in which both sides live: the category of mixed-HT chiral algebras at arbitrary  $d$ , the chiral derived centre as a functor on it, and the formal-Darboux stalk as a restriction functor. At  $d = 1$  the constructions reduce to the theory of chiral algebras on curves [28]; at  $d = 2$  they support the manuscript’s local theorem as the formal-Darboux stalk of the conjecture. The stalk carries the concrete arity-two holomorphic collision: the diagonal local-cohomology singular product of Theorem 1.15.

### A.1 Factorization $E_d$ -algebras on smooth complex $d$ -folds

Let  $X$  be a smooth complex variety of complex dimension  $d$ . Write  $\text{Ran}(X)$  for the Ran space of finite subsets of  $X$ . Let  $\text{Mfd}^{\text{lc}}(X)$  be the symmetric monoidal  $\infty$ -category whose objects are open subsets  $U \subset X$  and whose monoidal product is disjoint union of subsets with disjoint closures.

*Definition A.1 (Holomorphic factorization  $E_d$ -algebra).* A holomorphic factorization  $E_d$ -algebra on  $X$  is a symmetric monoidal  $\infty$ -functor

$$\mathcal{F} : \text{Mfd}^{\text{lc}}(X) \longrightarrow E_d\text{-Alg},$$

i.e. an assignment  $U \mapsto \mathcal{F}(U) \in E_d\text{-Alg}$ , factorisation isomorphisms

$$\mathcal{F}(U_1 \sqcup \cdots \sqcup U_n) \xrightarrow{\sim} \mathcal{F}(U_1) \otimes \cdots \otimes \mathcal{F}(U_n)$$

for finite collections of opens with pairwise disjoint closures, and extension-by-zero structure maps  $\mathcal{F}(U) \rightarrow \mathcal{F}(V)$  for  $U \subset V$  open, satisfying the factorization axioms of Costello–Gwilliam [15, Vol. I, Defn. 3.1.4]. The *holomorphic* condition is that  $\mathcal{F}$  is locally constant in the  $C^\infty$  direction and varies holomorphically in the complex direction: for any holomorphic embedding  $\iota : U \hookrightarrow V$ , the structure map  $\mathcal{F}(\iota)$  is a quasi-isomorphism, and the family  $\mathcal{F}$  on  $X$  is presented by a  $\mathcal{D}$ -module on  $\text{Ran}(X)$ .

*Remark A.2 (Equivalent presentations).* A holomorphic factorization  $E_d$ -algebra on  $X$  admits three equivalent presentations [28, §5.5; 41; 61]:

- (i) *Costello–Gwilliam.* A factorization algebra in the sense of [15, Vol. I, §3] valued in  $E_d$ -algebras, with the holomorphic locality differential.
- (ii) *Factorisable  $\mathcal{D}$ -module on  $\text{Ran}(X)$ .* A factorisable  $\mathcal{D}$ -module  $\mathcal{A}$  on  $\text{Ran}(X)$  valued in  $E_d$ -algebras. At  $d = 1$  this is the algebro-geometric construction of chiral algebras on curves [28, §3]; at  $d \geq 2$  it is the natural higher-dimensional extension whose equivalence with (1) and (3) is supplied by Costello–Gwilliam [15, Vol. I, §3] and Lurie–Ayala–Francis [41; 62, §5.5].
- (iii) *Lurie–Ayala–Francis.* An  $E_d$ -algebra object of the  $\infty$ -category of factorisation algebras on  $X$  in the sense of [41; 62, §5.5.4].

At  $d = 1$  all three reduce to chiral algebras on curves in the algebro-geometric sense of [28, §3]. For  $d \geq 2$  the manuscript uses the Costello–Gwilliam holomorphic factorization-algebra presentation as the closed-string sector classical observables; the global chiral  $E_d$ -Deligne structure is the conjectural layer isolated in §10.1.



*Definition A.3 (Category of holomorphic factorization  $E_d$ -algebras).* Let  $\text{HolFA}_d(X)$  denote the symmetric monoidal  $\infty$ -category whose objects are holomorphic factorization  $E_d$ -algebras on  $X$  in the sense of Definition A.1, and whose morphisms are natural transformations of symmetric monoidal  $\infty$ -functors that intertwine the factorisation isomorphisms. Its monoidal product is the pointwise tensor product of  $E_d$ -algebras.

### A.2 Mixed holomorphic-topological refinement

The manuscript's local geometry is  $X = \mathbb{R}_{\text{top}}^k \times M_{\text{hol}}$  with  $M$  a smooth complex variety of complex dimension  $d$  ( $k = 2, d = 2$  in the body). The factorization algebra carries three independent locality differentials: support locality on  $\mathbb{R}^k$ , holomorphic locality on  $M$ , and topological / locally constant in  $\mathbb{R}^k$ .

*Definition A.4 (Mixed-HT factorization  $E_d^{\text{HT}}$ -algebra).* A mixed-HT factorization  $E_d^{\text{HT}}$ -algebra on  $X = \mathbb{R}_{\text{top}}^k \times M_{\text{hol}}$  is a symmetric monoidal  $\infty$ -functor

$$\mathcal{F} : \text{Mfd}^{\text{lc}}(X) \longrightarrow E_d\text{-Alg}$$

satisfying the factorization axioms together with the three locality data:

- (i) *Support locality on  $\mathbb{R}^k$ .* For any compact embedding  $K \subset U \subset \mathbb{R}^k$ , the section  $\mathcal{F}$  is supported on  $K$  when restricted to  $U \setminus K$  extends to zero.
- (ii) *Topological locality on  $\mathbb{R}^k$ .* For any open  $U \subset \mathbb{R}^k$  and any inclusion  $U_0 \subset U$  with  $\pi_0(U_0) \rightarrow \pi_0(U)$  a bijection, the structure map  $\mathcal{F}(U_0 \times B) \rightarrow \mathcal{F}(U \times B)$  is a quasi-isomorphism for every  $B \subset M$ .
- (iii) *Holomorphic locality on  $M$ .* Restriction to the holomorphic factor is a holomorphic factorization  $E_d$ -algebra in the sense of Definition A.1.

*Definition A.5 (Category  $\text{ChirAlg}_{k,d}^{\text{HT}}(X)$ ).* Let  $\text{ChirAlg}_{k,d}^{\text{HT}}(X)$  denote the symmetric monoidal  $\infty$ -category whose objects are mixed-HT factorization  $E_d^{\text{HT}}$ -algebras on  $X$  in the sense of Definition A.4, and whose morphisms intertwine all three locality structures. At  $k = 0$  this reduces to  $\text{HolFA}_d(M)$ ; at  $d = 0$  it reduces to topological factorization algebras on  $\mathbb{R}^k$ . The manuscript's local geometry is  $k = 2, d = 2$ .

### A.3 The chiral derived centre functor

*Definition A.6 (Chiral derived centre at  $d = 1$ ).* For a chiral algebra  $\mathcal{A} \in \text{HolFA}_1(C)$  on a smooth curve  $C$ , the *chiral derived centre*  $Z_{\text{ch}}^{\text{der}}(\mathcal{A})$  is the chiral Hochschild cochain complex

$$Z_{\text{ch}}^{\text{der}}(\mathcal{A}) = C_{\text{ch}}^{\bullet}(\mathcal{A}, \mathcal{A}) := \text{RHom}_{\mathcal{A} \otimes \mathcal{A}^{\text{op}}}(\mathcal{A}, \mathcal{A})$$

in the chiral monoidal structure of [28, §3.4]. It carries a natural  $E_2$ -algebra structure (chiral Deligne theorem at  $d = 1$ ; [28, §4], Lurie HA [41, Thm. 5.3.1.14], Costello–Gwilliam [15, Vol. I, §6]). This is the proved  $d = 1$  chiral Deligne theorem of Vol I.

*Definition A.7 (Chiral derived centre at  $d \geq 2$ ).* For a holomorphic factorization  $E_d$ -algebra  $\mathcal{A} \in \text{HolFA}_d(X)$  on a smooth complex variety  $X$  of complex dimension  $d$ , the *chiral derived centre at  $d$*  is

$$Z_{E_d^{\text{ch}}}^{\text{der}}(\mathcal{A}) = C_{\text{ch}}^{\bullet}(\mathcal{A}, \mathcal{A}) := \text{RHom}_{\mathcal{A} \otimes \mathcal{A}^{\text{op}}}(\mathcal{A}, \mathcal{A})$$

in the higher chiral monoidal structure on  $\text{HolFA}_d(X)$ . At  $d = 1$  this is Definition A.6. At  $d \geq 2$  the  $\text{RHom}$  is taken in the  $\infty$ -category of factorization  $E_d$ -algebra bimodules on  $X$ .

*Status at  $d \geq 2$ .* The conjectural  $E_{d+1}^{\text{ch}}$ -structure on  $Z_{E_d^{\text{ch}}}^{\text{der}}(\mathcal{A})$  extending the  $E_2$ -structure of the  $d = 1$  case is the chiral  $E_d$ -Deligne conjecture (§10.1; Lurie [41, §5.3.1]); partial constructions are due to Hennion (chiral Lie algebroids on smooth schemes) and Francis (factorisable co-operads). The Mixed Holomorphic–Topological Deligne conjecture at  $d = 2$  is the corresponding statement for  $\text{ChirAlg}_{g,2}^{\text{HT}}(\mathbb{R}^2 \times \mathbb{C}^2)$ .

*Definition A.8 (Mixed-HT chiral derived centre).* For  $\mathcal{F} \in \text{ChirAlg}_{g,d}^{\text{HT}}(X)$ , the *mixed-HT chiral derived centre* is

$$Z_{E_d^{\text{HT}}}^{\text{der}}(\mathcal{F}) = C_{\text{ch,HT}}^{\bullet}(\mathcal{F}, \mathcal{F}),$$

the  $\text{RHom}$  of  $\mathcal{F}$  with itself in the bimodule  $\infty$ -category over  $\mathcal{F} \otimes \mathcal{F}^{\text{op}}$  inside  $\text{ChirAlg}_{g,d}^{\text{HT}}(X)$ . The mixed-HT structure on the output is the natural lift of the  $E_{d+1}$ -structure expected at  $d \geq 2$  (chiral  $E_d$ -Deligne conjecture).

#### A.4 The formal-Darboux stalk functor

For  $X = \mathbb{R}_{\text{top}}^k \times \mathbb{C}_{\text{hol}}^d$  and a closed point  $p \in \mathbb{C}^d$ , the formal-Darboux stalk operation is restriction to the formal completion at  $p$ .

*Definition A.9 (Formal-Darboux stalk functor).* The *formal-Darboux stalk functor* at  $p$  is the symmetric monoidal  $\infty$ -functor

$$(-)|_p: \text{ChirAlg}_{g,d}^{\text{HT}}(\mathbb{R}^k \times \mathbb{C}^d) \longrightarrow \text{ChirAlg}_{g,d}^{\text{HT}}(\mathbb{R}^k \times \widehat{\mathbb{C}^d}_p),$$

sending  $\mathcal{F} \mapsto \mathcal{F}|_{\mathbb{R}^k \times \widehat{\mathbb{C}^d}_p}$ , the restriction of the factorisable structure to the formal neighbourhood  $\widehat{\mathbb{C}^d}_p = \text{Spf } \widehat{\mathcal{O}_{\mathbb{C}^d,p}}$ . The holomorphic Darboux theorem trivialises the germ of  $\mathbb{C}^d$  at  $p$  to the formal symplectic polydisk in coordinates  $(z_1, \dots, z_d)$ , so the value at  $p$  depends on  $p$  only through its formal isomorphism class — equivalently, only through the formal Darboux coordinates.

**PROPOSITION A.10 (Equivariance under formal symplectomorphism).** The formal-Darboux stalk functor of Definition A.9 is equivariant under the formal symplectomorphism group  $\text{Symp}^{\text{form}}(\mathbb{C}^d)$  acting on the source category by formal coordinate change at  $p$  and on the target by the same change of formal Darboux coordinates. At  $d = 2$  on the holomorphic-symplectic factor, the stalk is canonical up to formal Darboux change.

*Proof.* The Darboux theorem in formal coordinates trivialises any holomorphic symplectic form to  $\omega_p = \sum_i dz_{2i-1} \wedge dz_{2i}$ , with the trivialisation defined uniquely up to the formal symplectomorphism group. The factorisation structure pulls back along this trivialisation by symmetric monoidal naturality.  $\square$

#### A.5 The bulk closed-string sector as an $E_2^{\text{HT}}$ -factorization algebra

*Construction A.11 (Bulk classical observables on  $\mathbb{R}^2 \times \mathbb{C}^2$ ).* Let  $X = \mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$  and  $\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^{\vee}[1]$  as in §2. The closed-string sector classical observables factorization algebra  $\mathcal{A}_{\text{bulk}}^{\text{cl}} \in \text{ChirAlg}_{g,2}^{\text{HT}}(X)$  is the assignment

$$U \times B \longmapsto C_{\text{CE,cont}}^{\bullet}(\Omega_c^{\bullet}(U) \widehat{\otimes} \Omega_c^{\bullet}(B) \widehat{\otimes} \mathfrak{g}),$$

for  $U \subset \mathbb{R}^2$  and  $B \subset \mathbb{C}^2$  open, with  $E_2$ -algebra structure from the symmetric algebra structure of the continuous CE cochains and factorisation isomorphisms inherited from extension-by-zero on disjoint

opens. The three locality differentials of Definition A.4 are the de Rham differential on  $U$ , the Dolbeault differential  $\bar{\partial}$  on  $B$ , and the internal CE differential dual to the  $\mathfrak{g}$ -bracket.

PROPOSITION A.12 (*Formal-Darboux stalk of the bulk*). At a closed point  $p \in \mathbb{C}^2$ , the formal-Darboux stalk of  $\mathcal{A}_{\text{bulk}}^{\text{cl}}$  is the continuous CE cochain algebra of the shifted-cotangent BF Lie algebra at the formal polydisk:

$$\mathcal{A}_{\text{bulk}}^{\text{cl}}|_p = C_{\text{CE,cont}}^\bullet(\mathfrak{g}), \quad \mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1], \quad \mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1.$$

This is the chart algebra  $A_{b_p} = \widehat{\mathcal{O}}_{\mathbb{C}^2, p} \cong \mathbb{C}[[z_1, z_2]]$  in formal Darboux coordinates equipped with the inherited  $E_2$ -algebra and chiral factorisation structure of Construction A.II. Its arity-two singular operation is the native diagonal singular product

$$J_x(z)J_y(w) \sim K_\Delta^{(2)}(z, w)J_{[x, y]_{\mathfrak{g}}}(w), \quad x, y \in \mathfrak{g},$$

with

$$K_\Delta^{(2)} = \frac{d\zeta_1 d\zeta_2}{\zeta_1 \zeta_2}, \quad \zeta_i = z_i - w_i.$$

*Proof.* Apply Definition A.9 to Construction A.II: the de Rham and Dolbeault Poincaré contractions on the formal polydisk reduce the compactly supported sheaf data to the continuous CE cochains of  $\mathfrak{g}$  at  $p$ . Extension by zero gives the regular product on disjoint formal polydisks. When two formal insertions collide, the Dolbeault resolution of the diagonal leaves the local-cohomology quotient

$$H_\Delta^2(\mathbb{C}[[z_1, z_2, w_1, w_2]]) d\zeta_1 d\zeta_2,$$

and the CE bracket of  $\mathfrak{g}$  gives exactly the singular product of Theorem I.15. Thus the formal completion retains the diagonal collision operation even though its disjoint-open product is the extension-by-zero product.  $\square$

#### A.6 The brane open category as factorization $E_1^{\text{HT}}$ -module

Definition A.13 (*Boundary open category*). The *boundary open category* of the local theory is

$$C_\partial^{\text{op}} = \Phi_d^{\text{FA}}(\mathcal{A}_{\text{bulk}}^{\text{cl}})|_{\partial X^{\text{KN}}},$$

the restriction of the bulk factorization algebra to the Kato–Nakayama log boundary  $\partial X^{\text{KN}}$  along the brane worldline  $\mathbb{R}_t$ . Its objects are factorisation  $E_1$ -modules over  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_{\partial X^{\text{KN}}}$  (boundary states); its morphisms are factorisation-compatible  $E_1$ -bimodule morphisms.

Construction A.14 (*Formal-Darboux vacuum and brane  $E_1$ -algebra*). The *formal-Darboux vacuum* at the closed point  $p \in \mathbb{C}^2$  is the object  $b_p \in C_\partial^{\text{op}}$  whose endomorphism algebra is the formal-completion endomorphism algebra

$$A_{b_p} = \text{End}_{C_\partial^{\text{op}}}(b_p) = \widehat{\mathcal{O}}_{\mathbb{C}^2, p} \cong \mathbb{C}[[z_1, z_2]]$$

of the brane skyscraper  $\mathcal{O}_p$  at  $p$ . At  $N$  Dirac branes over  $b_p$  the brane  $E_1$ -algebra of classical observables on the worldline  $\mathbb{R}_t$  is

$$\mathcal{A}_{\partial, N}^{\text{cl}} = C_{\text{CE}}^\bullet(\mathfrak{gl}_N, \text{Sym}(\mathfrak{gl}_N \oplus \mathfrak{gl}_N \oplus \mathfrak{gl}_N[1])), \quad Q\psi = [\phi_1, \phi_2],$$

with Chan–Paton substitution  $\rho_\phi: A_{b_p} \rightarrow M_N(\mathbb{C})$ ,  $z_i \mapsto \phi_i$ ,  $[\phi_1, \phi_2] = 0$  (Definition I.1). Curve chiral algebras such as affine Kac–Moody,  $\beta\gamma$ , Virasoro, and  $W$ -algebras enter through a separate curve restriction and anomaly matching before comparison with this formal disk.

**PROPOSITION A.15** (*Brane  $E_1$ -Hochschild as  $E_2$ -algebra*). The derived  $E_1$ -centre  $Z_{E_1}^{\text{der}}(A_{\partial,N}^{\text{cl}})$  is naturally an  $E_2$ -algebra, equivalent to the Hochschild cochain complex  $HH^\bullet(A_{\partial,N}^{\text{cl}}, A_{\partial,N}^{\text{cl}})$  with Gerstenhaber bracket and cup product. This is Deligne’s  $E_1$ -Hochschild theorem (Tamarkin [58], McClure–Smith [59], Kontsevich–Soibelman [60], Lurie [4I, Thm. 5.3.1.14]).

*A.7 The Mixed Holomorphic–Topological Deligne conjecture as a functorial statement*

**Conjecture A.16** (*Mixed Holomorphic–Topological Deligne, functorial form*). For the bulk closed-string sector  $\mathcal{A}_{\text{bulk}}^{\text{cl}} \in \text{ChiralAlg}_{2,2}^{\text{HT}}(\mathbb{R}^2 \times \mathbb{C}^2)$  of Construction A.11 and the boundary open category  $\mathcal{C}_\partial^{\text{op}}$  of Definition A.13, there is a canonical equivalence

$$\mathcal{A}_{\text{bulk}}^{\text{cl}} \simeq Z_{E_1^{\text{HT}}}^{\text{der}}(\mathcal{C}_\partial^{\text{op}})$$

of mixed-HT chiral algebras at  $d = 2$ , where the right-hand side is the mixed-HT chiral derived centre of the open category in the sense of Definition A.8.

*Status.*

- At  $d = 1$  (replacing  $\mathbb{C}^2$  by a smooth curve  $C$ ): proved (chiral Deligne theorem at  $d = 1$ , Vol I [63]; [28]).
- At  $d \geq 2$ : open. The chiral  $E_d$ -Deligne conjecture in the sense of Lurie [4I, §5.3.1] is the underlying purely chiral statement, of which the mixed-HT form is the holomorphic-topological enhancement. Partial cases are recorded in Hennion (chiral Lie algebroids on smooth schemes) and Francis (factorisable co-operads).
- *This manuscript:* proves the formal-Darboux stalk identification  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_p \simeq Z_{E_1}^{\text{der}}(A_{\partial,N}^{\text{cl}}) \simeq HH^\bullet(A_{\partial,N}^{\text{cl}}, A_{\partial,N}^{\text{cl}})$  of Conjecture A.16 at  $N$  Dirac branes over the formal-Darboux vacuum  $b_p$ . At the formal-Darboux stalk the bulk reduces to its  $E_2$ -algebra value  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_p = C_{\text{CE,cont}}^\bullet(\mathfrak{g})$  of Proposition A.12, and Lurie–Francis additivity / Costello–Gwilliam bulk–boundary correspondence identifies it with the brane-side derived  $E_1$ -centre  $Z_{E_1}^{\text{der}}(A_{\partial,N}^{\text{cl}}) \simeq HH^\bullet(A_{\partial,N}^{\text{cl}}, A_{\partial,N}^{\text{cl}})$  as  $E_2$ -algebras (Theorem 1.24), with the formal-coordinate arity-two singular product computed by Theorem 1.15.

**THEOREM A.17** (*Formal-Darboux stalk of Conjecture A.16*). At  $N$  Dirac branes over the formal-Darboux vacuum  $b_p \in \mathcal{C}_\partial^{\text{op}}$ , the formal-Darboux stalk of Conjecture A.16 holds:

$$\mathcal{A}_{\text{bulk}}^{\text{cl}}|_p \simeq Z_{E_1}^{\text{der}}(A_{\partial,N}^{\text{cl}}) \simeq HH^\bullet(A_{\partial,N}^{\text{cl}}, A_{\partial,N}^{\text{cl}})$$

as  $E_2$ -algebras, with formal Darboux coordinate expression  $J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)}$  on  $\mathfrak{h} = A_{b_p}/\mathbb{C} \cdot 1$ , and reduced closed-to-open dictionary  $c_f \mapsto \theta_f \in HH^1, u_f \mapsto J(f) \cdot (-) \in HH^0$  on Chevalley–Eilenberg generators of  $\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1]$ . The arity-two formal diagonal operation at the stalk is the singular product of Theorem 1.15. The chain-level lift  $\Phi_N$  to a strict  $P_0$ -factorization-centre morphism is conditional on  $\text{Ob}_{\text{cent/QME/desc}} = 0$  (Theorem E.53).

*Proof.* Combine Proposition A.12 (formal-Darboux stalk of the bulk), Proposition A.15 (Deligne’s  $E_1$ -Hochschild theorem for the brane  $E_1$ -algebra), and Theorem 1.24 (Costello–Gwilliam bulk–boundary correspondence at the formal-Darboux stalk). The trace-map computation is Theorem 5.146 (Procesi–Razmyslov + Loday–Quillen–Tsygan); the dictionary is Theorem 4.20; the arity-two diagonal operation is Theorem 1.15; the conditional chain-level lift is Theorem E.53.  $\square$

*Remark A.18 (Formal-Darboux stalk and global conjecture).* Theorem A.17 computes the formal-Darboux stalk of Conjecture A.16 at the formal-Darboux vacuum  $b_p$ . The conjecture itself — the global equivalence on  $\mathbb{R}^2 \times \mathbb{C}^2$  between the bulk  $\mathcal{A}_{\text{bulk}}^{\text{cl}}$  and the mixed-HT chiral derived centre of the boundary open category at  $d = 2$  — remains open. The  $d = 2$  chiral  $E_d$ -Deligne theorem on which it depends is itself open (§10.1). Identification with any curve chiral algebra requires a separate restriction theorem and matched-conventions transfer (Theorem 10.13).

#### A.8 The chiral $E_d$ -operad: holomorphic little disks with principal-parts data

The functorial Mixed-HT Deligne conjecture (Conjecture A.16) requires an operadic action: the chiral derived centre  $Z_{E_d^{\text{ch}}}^{\text{der}}(\mathcal{A})$  is the endomorphism object in a chiral  $E_d$ -module category. At  $d = 1$  the chiral operad on curves supplies this structure [28, §3.4]. At  $d \geq 2$  the operad is controlled by higher-codimension diagonal local cohomology. This subsection fixes the formal operadic model used by the manuscript; at  $d = 2$  its arity-two coefficient is the module computed in Theorem 1.15.

*Definition A.19 (Configuration  $\mathcal{D}$ -module).* Let  $X$  be a smooth complex variety of complex dimension  $d$  and  $I$  a non-empty finite set. Write  $X^I$  for the  $|I|$ -fold product and  $j_I : \text{Conf}_I(X) \hookrightarrow X^I$  for the open embedding of the configuration space of  $|I|$  distinct points. The *configuration  $\mathcal{D}$ -module*

$$\omega_{X^I}^{\text{ch}} = j_{I,*} j_I^! \omega_{X^I}$$

is the  $!$ -extension to  $X^I$  of the dualising sheaf of  $\text{Conf}_I(X)$ .

*Definition A.20 (Formal chiral  $E_d$  operadic model).* The *formal chiral  $E_d$  operadic model* on  $\mathbb{C}^d$  is the colored  $\infty$ -operad  $\text{hE}_d^{\text{ch}}$  with single colour and operation space at arity  $|I|$

$$\text{hE}_d^{\text{ch}}(I) = \omega_{(\mathbb{C}^d)^I}^{\text{ch}},$$

with composition the chiral product on  $!$ -pull-back along the diagonal embedding  $\Delta_A : (\mathbb{C}^d)^A \hookrightarrow (\mathbb{C}^d)^I$  for a partition  $I = \sqcup_{a \in A} I_a$ , following [28, §3.4] as the formal higher-codimension Ran-space analogue. At  $d \geq 2$  this definition is a model for the conjectural chiral  $E_d$ -operadic action. The full global Deligne theorem is the separate conjectural equivalence stated below.

*Remark A.21 (Comparison with curves at  $d = 1$ ).* For  $d = 1$  on a curve  $C$ , the configuration  $\mathcal{D}$ -module  $\omega_{C^I}^{\text{ch}}$  is the chiral operad on curves of [28, §3.4]. At  $d \geq 2$  the diagonals  $\Delta_{ij} \subset (\mathbb{C}^d)^I$  have complex codimension  $d$ , so the principal parts on  $\omega_{(\mathbb{C}^d)^I}^{\text{ch}}$  involve distributions of  $d$ -fold polar order; this is the higher-codimension chiral structure specific to  $d \geq 2$ .

*Remark A.22 (Arity two at  $d = 2$ ).* For  $X = \mathbb{C}^2$  and  $I = \{1, 2\}$ , the singular quotient of  $\omega_{X^I}^{\text{ch}}$  along the pairwise diagonal is

$$H_{\Delta}^2(\mathbb{C}[[z_1, z_2, w_1, w_2]]) d\zeta_1 d\zeta_2 = \bigoplus_{a,b \geq 0} \mathbb{C}[[w_1, w_2]] \rho_{a,b}^{\Delta}.$$

The generator  $\rho_{0,0}^{\Delta} = K_{\Delta}^{(2)}$  is the two-dimensional Cauchy kernel. For the Hamiltonian BF algebra, the associated binary operation sends  $x \otimes y$  to  $[x, y]_{\mathfrak{g}}$ , giving the formal singular product of Theorem 1.15.

*Construction A.23 (Conditional  $\text{hE}_d^{\text{ch}}$  operation maps).* Assume the conjectural chiral  $E_d$ -operadic action of §10.1 exists on  $\text{HolFA}_d(\mathbb{C}^d)$ . Then a holomorphic factorization  $E_d$ -algebra  $\mathcal{A} \in \text{HolFA}_d(\mathbb{C}^d)$  (Definition A.1) carries structure maps

$$\mu_I^{\text{ch}} : \text{hE}_d^{\text{ch}}(I) \otimes^! \mathcal{A}_{\text{Ran}}^{\boxtimes I} \longrightarrow \Delta_I^* \mathcal{A}_{\text{Ran}}$$

intertwining the chiral product of Definition A.20 with  $\Delta_I^*$ , where  $\mathcal{A}_{\text{Ran}}$  denotes the corresponding Ran-space factorization object. At  $d = 1$  these maps are the chiral operations on curves. At  $d = 2$  and arity two, the formal-Darboux stalk coefficient is computed unconditionally by Theorem 1.15.

*Remark A.24 (Status of the operation maps).* The Costello–Gwilliam and Lurie–Ayala–Francis factorization formalisms give the symmetric monoidal factorization category and its factorization-homology presentation. The chiral  $E_d$ -Deligne equivalence at  $d \geq 2$  is the additional theorem sought here. The manuscript uses Construction A.23 only as the conditional global model; the proved local input is the arity-two diagonal local-cohomology operation of Theorem 1.15.

*Construction A.25 (Conditional mixed-HT enhancement  $\mathbf{hE}_d^{\text{ch,HT}}$ ).* Adjoin to  $\mathbf{hE}_d^{\text{ch}}$  the topological  $E_k$ -operad on  $\mathbb{R}_{\text{top}}^k$ : the colored  $\infty$ -operad  $\mathbf{hE}_d^{\text{ch,HT}}$  has operation space

$$\mathbf{hE}_d^{\text{ch,HT}}(I) = \text{Conf}_I(\mathbb{R}^k) \times \omega_{(\mathbb{C}^d)^I}^{\text{ch}},$$

with composition the product of topological little-disks composition on the  $\mathbb{R}^k$  factor and chiral composition of Definition A.20 on the  $\mathbb{C}^d$  factor. An algebra over  $\mathbf{hE}_d^{\text{ch,HT}}$  is a mixed-HT factorization  $E_d^{\text{HT}}$ -algebra of Definition A.4, conditional on the same global chiral  $E_d$ -operadic action.

#### A.9 Three hypotheses for chiral $E_d$ Deligne at $d \geq 2$

The chiral  $E_d$  Deligne identification  $C_{\text{ch}}^\bullet(\mathcal{A}, \mathcal{A}) \simeq Z_{\mathbf{hE}_d^{\text{ch}}}^{\text{der}}(\mathcal{A})$  of Conjecture 10.1 reduces to three precise hypotheses on  $\mathbf{hE}_d^{\text{ch}}$  of Definition A.20.

*Definition A.26 (Hypothesis F: formality of  $\mathbf{hE}_d^{\text{ch}}$ ).* The formal chiral  $E_d$  operadic model  $\mathbf{hE}_d^{\text{ch}}$  is formal as an  $\infty$ -operad: there is a quasi-isomorphism

$$\mathbf{hE}_d^{\text{ch}} \simeq H_*(\mathbf{hE}_d^{\text{ch}})$$

to its homology operad in chain complexes of  $\mathcal{D}_{\text{Ran}(\mathbb{C}^d)}$ -modules. At  $d = 1$  this is Tamarkin’s formality theorem [58, 60]; at  $d \geq 2$  it is open.

*Definition A.27 (Hypothesis C: centrality).* For  $\mathcal{A} \in \text{HolFA}_d(\mathbb{C}^d)$ , the chiral derived centre  $Z_{\mathbf{hE}_d^{\text{ch}}}^{\text{der}}(\mathcal{A})$  is the endomorphism object in the bimodule  $\infty$ -category  $\mathcal{A} \otimes \mathcal{A}^{\text{op}}\text{-Mod}^{\text{ch}}$ :

$$Z_{\mathbf{hE}_d^{\text{ch}}}^{\text{der}}(\mathcal{A}) = \text{RHom}_{\mathcal{A} \otimes \mathcal{A}^{\text{op}}}(\mathcal{A}, \mathcal{A}).$$

At  $d = 1$  this is the chiral-centre construction for chiral algebras on curves. For  $d \geq 2$  its topological analogue is Lurie’s  $E_d$ -centre construction [41, §5.3.1]; the holomorphic chiral enhancement is part of the conjectural package.

*Definition A.28 (Hypothesis R: Ran-space descent).* The factorization-homology functor  $\text{Sp}_X^{\text{ch}}: \text{HolFA}_d(X) \rightarrow \mathcal{D}(\text{Ran}(X))\text{-Mod}$  of Lurie–Ayala–Francis [41, §5.5] is Ran-cohomologically descent-faithful: for any holomorphic factorization algebra  $\mathcal{A}$ ,

$$\mathcal{A} \xrightarrow{\sim} \text{holim}_{(I \hookrightarrow \text{Ran}(X))^{\text{op}}} \mathcal{A}|_I,$$

the limit over the simplicial diagram of finite-set embeddings into the Ran space. At  $d = 1$  this is part of the factorization theorem for chiral algebras on curves [28, §3]; at  $d \geq 2$  the corresponding descent statement is a required input for the global chiral  $E_d$ -Deligne theorem, with evidence from factorization homology and Francis–Gaitsgory chiral Koszul duality [64, Thm. 6.5.1].

THEOREM A.29 (*Conditional chiral  $E_d$  Deligne*). Under Hypotheses A.26, A.27, A.28, the chiral derived centre satisfies

$$Z_{\mathrm{hE}_d^{\mathrm{ch}}}^{\mathrm{der}}(\mathcal{A}) \simeq C_{\mathrm{ch}}^{\bullet}(\mathcal{A}, \mathcal{A})$$

as  $E_{d+1}^{\mathrm{ch}}$ -algebras. Hypotheses A.27 and A.28 are the chiral enhancements of known topological and curve-level theorems; at  $d \geq 2$  they remain part of the conditional chiral  $E_d$ -Deligne package together with Hypothesis A.26.

*Proof sketch.* By Hypothesis A.27 the chiral derived centre is  $\mathrm{RHom}_{\mathcal{A} \otimes \mathcal{A}^{\mathrm{op}}}(\mathcal{A}, \mathcal{A})$ . By Hypothesis A.28 this is computed by the simplicial bar resolution of  $\mathcal{A}$  as  $\mathcal{A} \otimes \mathcal{A}^{\mathrm{op}}$ -bimodule on the Ran space, yielding  $C_{\mathrm{ch}}^{\bullet}(\mathcal{A}, \mathcal{A})$ . By Hypothesis A.26 the  $\mathrm{hE}_d^{\mathrm{ch}}$ -action refines along the formality quasi-isomorphism to an  $E_{d+1}^{\mathrm{ch}}$ -action, by the same chain of natural transformations as Tamarkin [58] at  $d = 1$ , after replacing the one-dimensional chiral operad by the conditional Lurie–Ayala–Francis  $\mathcal{D}$ -module presentation [41, §5.5.4] of the chiral operad.  $\square$

Remark A.30 (*Obstruction class for Hypothesis A.26*). Hypothesis A.26 is equivalent to the vanishing of the Maurer–Cartan element  $\mathfrak{m}_d \in \mathrm{Def}(\mathrm{hE}_d^{\mathrm{ch}} \rightarrow H_*(\mathrm{hE}_d^{\mathrm{ch}}))$  classifying the chiral  $E_d$ -operad relative to its homology operad is non-trivial. At  $d = 1$  Tamarkin [58] shows  $\mathfrak{m}_1 = 0$  via the Drinfeld associator. At  $d \geq 2$  the existence of an analogous Drinfeld associator on  $\mathrm{Conf}_I(\mathbb{C}^d)$  for  $|I| \geq 2$  is the residual frontier. The class

$$\mathrm{Ob}_F = [\mathfrak{m}_d] \in H^*(\mathrm{Def}(\mathrm{hE}_d^{\mathrm{ch}}))$$

is the formality obstruction at  $d \geq 2$ .

Remark A.31 (*Chiral  $E_d$  package obstruction*). The global chiral  $E_d$ -Deligne input at  $d \geq 2$  consists of the operadic action of Definition A.20 together with Hypotheses A.26, A.27, and A.28. The manuscript packages the obstruction to this input as

$$\mathrm{Ob}_{\mathrm{ch}E_d} = \mathrm{Ob}_F \oplus \mathrm{Ob}_C \oplus \mathrm{Ob}_R,$$

where  $\mathrm{Ob}_C$  and  $\mathrm{Ob}_R$  measure the obstruction to the chiral centrality and Ran-descent hypotheses. The formal-Darboux stalk theorem computes the arity-two  $d = 2$  local operation on which any global theorem restricts; vanishing of  $\mathrm{Ob}_{\mathrm{ch}E_d}$  is the additional global condition.

#### A.10 Descent over the Kato–Nakayama log boundary

The boundary open category  $C_{\partial}^{\mathrm{op}}$  (Definition A.13) is defined globally on the Kato–Nakayama log boundary  $\partial X^{\mathrm{KN}}$ . This subsection writes the descent spectral sequence whose  $E_2$ -collapse removes one descent obstruction. The global mixed-HT Deligne equivalence further requires the other obstruction classes to vanish.

Definition A.32 (*Kato–Nakayama topological realisation*). For a log scheme  $(X, \mathcal{M}_X, \alpha)$  in the sense of Kato [65], the *Kato–Nakayama topological realisation* [66] is the topological space  $X^{\mathrm{KN}}$  with continuous projection  $\pi^{\mathrm{KN}}: X^{\mathrm{KN}} \rightarrow X^{\mathrm{an}}$ . Over the locus of trivial log structure  $\pi^{\mathrm{KN}}$  is a homeomorphism; over a divisor  $D \subset X$  with simple normal-crossing log structure, the fibre over a codimension- $r$  stratum is a real  $r$ -torus. The boundary  $\partial X^{\mathrm{KN}} = (\pi^{\mathrm{KN}})^{-1}(D)$  is a torus bundle over the smooth locus of  $D$ .

*Definition A.33 (Sheaf of factorization  $E_d^{\text{HT}}$ -categories on  $\partial X^{\text{KN}}$ ).* The boundary open category extends to a sheaf  $\underline{C}_\partial^{\text{op}}$  of mixed-HT factorization  $E_d^{\text{HT}}$ -categories on  $\partial X^{\text{KN}}$ : for  $U \subset \partial X^{\text{KN}}$  open,

$$\underline{C}_\partial^{\text{op}}(U) = \mathcal{A}_{\text{bulk}}^{\text{cl}}|_{U\text{-Mod}_\partial^{\text{ch,HT}}},$$

the boundary factorization  $E_d^{\text{HT}}$ -modules over the restricted bulk; restriction to a smaller open is restriction of factorization modules. At a closed point  $p \in \partial X^{\text{KN}}$  in the trivial log stratum, the stalk is the formal-Darboux stalk of Definition A.9.

*THEOREM A.34 (Log étale descent for the boundary open category).* The sheaf  $\underline{C}_\partial^{\text{op}}$  of Definition A.33 satisfies log étale descent: for any log étale cover  $\{U_a \rightarrow \partial X^{\text{KN}}\}$ ,

$$\underline{C}_\partial^{\text{op}}(\partial X^{\text{KN}}) \xrightarrow{\sim} \text{holim}(\Pi_a \underline{C}_\partial^{\text{op}}(U_a) \rightrightarrows \Pi_{a,b} \underline{C}_\partial^{\text{op}}(U_a \cap U_b) \rightrightarrows \cdots).$$

*Proof.* Apply Olsson's log étale descent theorem [67, Thm. 4.7] for sheaves of  $\infty$ -categories on log étale stacks. The factorization  $E_d^{\text{HT}}$ -module condition is preserved under log étale pullback because the structure maps of Definition A.4 are local on  $\partial X^{\text{KN}}$ . Closure under holim is the  $\infty$ -categorical limit closure of factorization  $\infty$ -categories [41, §5.5.4.10].  $\square$

*THEOREM A.35 (Descent spectral sequence).* Choose a log étale cover  $\{U_a \rightarrow \partial X^{\text{KN}}\}_{a \in \mathcal{A}}$  trivialising the log structure. There is a spectral sequence

$$E_2^{p,q} = H^p(\partial X^{\text{KN}}, \mathcal{H}^q(\underline{C}_\partial^{\text{op}})) \Rightarrow H^{p+q}(\underline{C}_\partial^{\text{op}}),$$

where  $\mathcal{H}^q$  is the  $q$ -th derived stalk. At a formal-Darboux point  $p \in \partial X^{\text{KN}}$  the stalk  $\mathcal{H}^*(\underline{C}_\partial^{\text{op}})_p$  is the formal-Darboux stalk of Theorem A.17.

*Proof.* Standard sheaf-cohomology descent spectral sequence for the  $\infty$ -categorical sheaf of Definition A.33, via Čech cohomology with respect to the log étale cover and convergence by Theorem A.34.  $\square$

*Definition A.36 (Formal-Darboux  $E_2$ -collapse).* The descent spectral sequence of Theorem A.35 *collapses at the formal-Darboux strata* if the differential

$$d_2: H^p(\partial X^{\text{KN}}, \mathcal{H}^q) \longrightarrow H^{p+2}(\partial X^{\text{KN}}, \mathcal{H}^{q-1})$$

vanishes at every formal-Darboux stratum. Equivalently, the formal-Darboux stalks glue across log strata along the local symplectic-form trivialisation of Proposition A.10 with trivial higher correction.

*THEOREM A.37 (Descent obstruction at the formal-Darboux strata).* The formal-Darboux  $E_2$ -collapse condition of Definition A.36 is precisely the vanishing of the first descent obstruction to gluing the formal-Darboux stalks across the Kato–Nakayama boundary. The obstruction is

$$\text{Ob}_{\text{desc}} = [d_2] \in H^2(\partial X^{\text{KN}}, \text{Hom}^{-1}(\mathcal{H}^*, \mathcal{H}^*)).$$

When this class vanishes, the local stalk theorem survives this descent test. The global equivalence

$$\mathcal{A}_{\text{bulk}}^{\text{cl}} \simeq Z_{E_2^{\text{HT}}}^{\text{der}}(\underline{C}_\partial^{\text{op}})$$

still requires the remaining chiral  $E_d$ , quantum master equation, and global comparison obstructions to vanish.



*Proof.* By Theorem A.35, the first possible obstruction to gluing the derived stalks through the chosen log étale cover is the  $d_2$ -differential. Its cohomology class is the displayed  $\text{Ob}_{\text{desc}}$ . Vanishing of this class is exactly the stated  $E_2$ -collapse condition. Theorem A.17 identifies each formal-Darboux stalk. Identification of the global bulk with the global chiral derived centre requires the remaining obstruction classes to vanish.  $\square$

#### A.11 All-loop QME for finite-jet PVA

The classical PVA Jacobi identity holds at the formal-Darboux stalk (Theorem 4.20) and is the Jacobi identity for the native two-dimensional singular product of Theorem 1.15. Lifting the PVA structure to all  $\hbar$ -orders — the quantum master equation for the finite-jet PVA — is the content of site #16 of the seventeen-site catalogue. The four hypotheses below isolate the analytic / homological control needed for all-loop convergence.

*Convention.* Under the binding convention of §, the symbol  $C_{\text{ch}}^\bullet(A_b, A_b)$  in this subsection denotes the brane-side derived  $E_1$ -Hochschild  $HH^\bullet(A_{\partial, N}^{\text{cl}}, A_{\partial, N}^{\text{cl}}) \simeq Z_{E_1}^{\text{der}}(A_{\partial, N}^{\text{cl}})$  identified with the bulk-stalk  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b$  by Costello–Gwilliam bulk–boundary correspondence (Theorem 5.6);  $Q_{\text{ch}}$  is the chiral cochain differential inherited from  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b$  under this identification. The hypotheses below are formulated on that object.

*Definition A.38 (Hypothesis K: KZ analytic SDR).* There is a strong deformation retract

$$(C_{\text{CE,cont}}^\bullet(\mathfrak{g}), Q_{\text{tot}}) \xrightarrow[\text{retract}]{\text{deform}} (C_{\text{ch}}^\bullet(A_b, A_b), Q_{\text{ch}})$$

in the category of chain complexes over the analytic ring  $\mathbb{C}\{\hbar\}$ ; the formal ring  $\mathbb{C}[[\hbar]]$  is replaced by this analytic coefficient ring, with finite-rank cohomology in each PVA weight, and with retract data analytic in  $\hbar$  in a punctured neighbourhood of  $\hbar = 0$ .

*Remark A.39 (Analytic SDR).* The classical PVA Jacobi holds in the  $\hbar$ -adic topology of the formal disk. All-loop lifting requires Stokes-controlled analytic continuation across the disk (Hypothesis A.40). A purely  $\hbar$ -formal SDR gives finite-loop existence (Costello [12, Ch. 5]); analytic SDR is needed for all-loop convergence.

*Definition A.40 (Hypothesis S: Stokes regularity).* The KZ connection on  $C_{\text{ch}}^\bullet(A_b, A_b)$  associated to the formal  $E_2$  singular product with kernel  $K_\Delta^{(2)}$  has trivial Stokes data along the real-positive  $\hbar$ -axis: every analytic continuation across  $\arg \hbar \in (-\pi, \pi]$  identifies with a single asymptotic series with matching sectorial asymptotics.

*Definition A.41 (Hypothesis W: reflected weights).* Under crossing  $z \leftrightarrow w$  on the formal  $E_2$  singular product of Theorem 1.15, the PVA weight grading on  $C_{\text{ch}}^\bullet(A_b, A_b)$  reflects: the operator

$$R: \text{wt}_z \otimes \text{wt}_w \longrightarrow \text{wt}_w \otimes \text{wt}_z$$

intertwines the singular product with itself; equivalently,  $C_{\text{ch}}^\bullet(A_b, A_b)$  is reflection-positive in the chiral  $E_2$ -sense.

*Definition A.42 (Hypothesis G:  $T = [Q_{\text{tot}}, G]$ ).* The energy–momentum tensor  $T \in C_{\text{ch}}^\bullet(A_b, A_b)$  is BRST-exact with primitive  $G$ :

$$T = [Q_{\text{tot}}, G],$$

where  $Q_{\text{tot}} = Q_{\text{CE}} + \hbar Q_{\text{BV}}$  is the total BRST operator and  $G$  is a degree- $(-1)$  density on the worldvolume. At the formal-Darboux stalk the primitive exists as a classical limit; the hypothesis is its survival to all  $\hbar$ -orders.

**THEOREM A.43** (*Conditional all-loop QME*). Under Hypotheses A.38, A.40, A.41, A.42, the formal-Darboux stalk of Theorem A.17 lifts to all orders in  $\hbar$ : there exists a quantum BV master action  $S^{\text{quant}} = S^{\text{cl}} + \sum_{n \geq 1} \hbar^n S^{(n)}$  satisfying  $\{S^{\text{quant}}, S^{\text{quant}}\} + \hbar \Delta S^{\text{quant}} = 0$  with

$$\mathcal{A}_{\text{bulk}}^{\text{cl}}|_p[[\hbar]] \simeq Z_{E_i}^{\text{der}}(A_{\partial, N}^{\text{cl}}[[\hbar]])$$

holding at all  $\hbar$ -orders, and the trace map  $J(f) = \overline{\text{Tr } f(\phi_1, \phi_2)}$  extending to  $J^{\text{quant}}(f) = J(f) + \sum_n \hbar^n J^{(n)}(f)$ .

*Proof sketch.* Costello's renormalization theorem [12, Thm. 8.6.0.1] gives finite-loop existence and uniqueness modulo cohomological obstructions. Hypothesis A.38 (analytic SDR) lifts existence to a punctured analytic neighbourhood of  $\hbar = 0$ . Hypothesis A.40 (Stokes regularity) extends existence across the disk  $|\hbar| < r_0$  for some  $r_0 > 0$  with matching sectorial asymptotics. Hypothesis A.41 (reflected weights) is the obstruction-vanishing input for the chiral  $E_2$ -action on  $C_{\text{ch}}^\bullet$  at all loops. Hypothesis A.42 ( $T = [Q_{\text{tot}}, G]$ ) eliminates the trace anomaly: BRST-exactness of  $T$  trivialises the conformal-anomaly cohomology class in  $H^2(C_{\text{ch}}^\bullet(A_b, A_b))$  via chain homotopy.  $\square$

**Remark A.44** (*QME obstruction class*). The four hypotheses combine into

$$\text{Ob}_{\text{QME}} = [\text{ob}_K] \oplus [\text{ob}_S] \oplus [\text{ob}_W] \oplus [\text{ob}_G] \in H^*(\text{local functionals on } \mathbb{R}^2 \times \mathbb{C}^2),$$

with summands the analytic-SDR obstruction  $[\text{ob}_K]$ , the Stokes-filtration class  $[\text{ob}_S]$ , the crossing class  $[\text{ob}_W]$ , and the conformal-anomaly class  $[\text{ob}_G]$ . Each summand is checkable in finitely many PVA weights at fixed loop order.

#### A.12 Operator-level lift of the modular line at $\Delta_5$

The Igusa cusp form  $\Phi_{10}$  is the level-2 section of the modular line  $\Omega_{\text{central}}$  on the Siegel period domain  $\mathfrak{H}_2$  [31, 32]. The square root  $\Delta_5 = \Phi_{10}^{1/2}$  is a scalar modular form in the external  $K_3 \times E$  comparison problem (§10.3). The formal-Darboux theorem constructs the local Capelli scalar. The operator-level lift asks for an operator whose protected trace is  $\Delta_5$  after the global compact-target theory and its trace have been constructed.

**Conjecture A.45** (*Operator lift of  $\Delta_5$* ). There exists a graded vertex-algebraic or factorization-algebraic object  $V$ , an involution or real form  $\sigma$ , and a protected trace such that

$$\text{Tr}_V(\sigma q^{L_0 - c/24} \prod_{i=1}^2 z_i^{H_i}) = \Delta_5(\tau, \rho, \nu) = \Phi_{10}(\tau, \rho, \nu)^{1/2}$$

on the level-2 cover of the Siegel period domain  $(\tau, \rho, \nu) \in \mathfrak{H}_2$ , with the trace compatible with the modular line  $\Omega_{\text{central}}$ . This is an open operator-construction problem extending the formal-Darboux stalk.

Three scalar theories give evidence for the desired trace. The operator object requires an additional construction.

*Construction A.46 (Borcherds–Kac–Moody candidate).* The Borcherds–Kac–Moody denominator formulas give automorphic products, including the fake-Monster denominator and the Gritsenko–Nikulin products [32, 68]. These are scalar identities. Passing from a scalar denominator formula to a vertex-algebraic object whose graded trace is  $\Delta_5$  requires an additional construction.

*Obstruction.*  $\text{Ob}_{\text{BKM}}$  is the obstruction to lifting the scalar Borcherds product to an operator trace with the required integral form, real structure, and modular covariance.

*Construction A.47 (Free-field realisation on  $K_3 \times E$ ).* The elliptic genus of  $K_3$ , its second-quantised product, and the Donaldson–Thomas / Pandharipande–Thomas comparison on  $K_3 \times E$  supply scalar modular data. A free-field chiral algebra on  $K_3 \times E$  whose trace is  $\Delta_5$ .

*Obstruction.*  $\text{Ob}_{K_3E}$  is the obstruction to constructing the compact target factorization object, its protected trace, and the descent of that trace to the level-2 modular line.

*Construction A.48 (Dabholkar–Murthy–Zagier candidate).* The Dabholkar–Murthy–Zagier decomposition of meromorphic Jacobi forms separates polar and finite parts in the  $1/\Phi_{10}$  dyonic counting problem [70]. This is a statement about mock modular and wall-crossing structures. An operator construction of  $\Delta_5$  is a separate lift.

*Obstruction.*  $\text{Ob}_{\text{DMZ}}$  is the obstruction to lifting the mock-modular wall-crossing package to an honest operator trace with the required holomorphic modular transformation law.

*Remark A.49 (Composition of operator-lift obstructions).* The three candidates are evidence for the same scalar modular form, and the operator construction is the common problem. The symbol  $\text{Ob}_V$  denotes the combined obstruction to constructing an operator-level lift from any of these scalar inputs. A cohomological product formula for these scalar obstruction inputs would require a separate definition.

### A.13 Convergent obstruction taxonomy

The four conditional theorems compose to a conditional global Mixed Holomorphic–Topological Deligne theorem.

**THEOREM A.50 (Conditional global Mixed-HT Deligne obstruction package).** Under the joint vanishing

$$\text{Ob}_{\text{global}} := \text{Ob}_{\text{ch}E_d} \oplus \text{Ob}_{\text{desc}} \oplus \text{Ob}_{\text{QME}} \oplus \text{Ob}_V = 0,$$

with the four summands as in Remark A.31, Theorem A.37, Remark A.44, and Remark A.49, the Mixed-HT Deligne conjecture A.16 holds globally on  $\mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$  at  $d = 2$ ,  $N$  Dirac branes, provided the construction hypotheses named in those four inputs have been supplied. This implication is conditional. It does record the compact-target modular-form comparison with an operator trace as a separate lift. The appearance of  $\Phi_{10}$  or  $\Phi_{10}^{1/2}$  belongs to the separate  $K_3 \times E$  operator-lift problem.

*Proof.* Compose Theorem A.29 (chiral  $E_d$  Deligne via the operadic action and Hypotheses A.26–A.28), Theorem A.37 (KN log-boundary descent via Definition A.36), Theorem A.43 (all-loop QME via Hypotheses A.38–A.42), and Conjecture A.45 only in the external compact-target comparison where that datum is part of the input. The four obstructions are independent cohomology classes living in distinct cohomology groups, so their joint vanishing is well-defined.  $\square$

*Remark A.51 (Reduction status).* Within the seventeen-site catalogue, sites #11 and #16 are the explicit obstruction conditions for  $\text{Ob}_V$  and  $\text{Ob}_{\text{QME}}$  respectively. The chiral  $E_d$  Deligne package obstruction  $\text{Ob}_{\text{ch}E_d}$  at  $d \geq 2$  is the residual research frontier of Conjecture 10.1; partial results (chiral Lie algebroids on

smooth schemes via Hennion, factorisable co-operads via Francis) give evidence for pieces of the package; the global theorem remains the conjectural target. The KN-descent obstruction  $\text{Ob}_{\text{desc}}$  reduces to the  $E_2$ -collapse condition of Definition A.36, checkable on a formal-Darboux stratum-by-stratum basis. The manuscript's main theorem (Theorem 5.6) is the formal-Darboux stalk that anchors all four obstructions, and Theorem 1.15 supplies its arity-two formal diagonal operation: each conditional theorem has the stalk theorem, including this diagonal singular product, as its base case.

*Construction A.52 (Global manuscript program).* The conditional global theorem (Theorem A.50) translates the manuscript's frontier (§10) into a single cohomological vanishing problem  $\text{Ob}_{\text{global}} = 0$ . The four summands are independent, hence individually attackable:

- $\text{Ob}_{\text{ch}E_d}$ : existence, centrality, Ran-descent, and formality for the chiral  $E_d$ -operadic package at  $d \geq 2$ ; its formality summand is the chiral analogue of the Drinfeld associator on  $\text{Conf}_I(\mathbb{C}^d)$ ,  $|I| \geq 2$ .
- $\text{Ob}_{\text{desc}}$ : vanishing of the  $d_2$  differential in the descent spectral sequence (Theorem A.35) on the formal-Darboux stratum.
- $\text{Ob}_{\text{QME}}$ : each of four checkable cohomology classes (Hypotheses A.38–A.42).
- $\text{Ob}_V$ : any of three candidate constructions (Constructions A.46, A.47, A.48).

The formal-Darboux stalk theorem of this manuscript (Theorem 5.6) anchors each.

## B FACTORIZATION CURRENT CONVENTIONS FOR THE CE/PV INTERFACE

This appendix fixes conventions for the brane-line factorization-current sector of the local mixed HT string field theory on  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}^2_{\text{o,hol}}$ : the interval-wise CE/PV current interface, and the obstruction surfaces that block strengthening from the algebraic current model to a Costello local-functional graph theorem [12, 15, 16]. Every BV degree, sign, completion, and obstruction class used by the body and by every other appendix is fixed here once.

The interface is one universal triangle of brackets. On a brane interval  $I \subset \mathbb{R}_{\text{brane}}$ , the Hamiltonian boundary observable  $B_f(a)$  (degree 0) and the residual cotangent observable  $\Theta_\rho(B)$  (degree 1) carry the open  $P_0$ -brackets

$$\{B_f(a), B_g(b)\} = B_{\{f,g\}}(ab), \quad \{B_f(a), \Theta_\rho(B)\} = \Theta_{f \cdot \rho}(aB), \quad \{\Theta_\rho(B), \Theta_\sigma(C)\} = 0,$$

where  $\{f, g\}$  is the Poisson bracket of the formal holomorphic symplectic polydisk  $(\widehat{\mathbb{C}}^2_{\text{o}}, dz_1 \wedge dz_2)$ ,  $f \cdot \rho$  is its Matlis residue coadjoint action of Proposition 8.7, and  $aB$  is the multiplication of a smooth compactly supported test function by a compactly supported distributional current. The CE source coordinates are  $c_\lambda$  (degree 1) and  $u_{a(t)dt \otimes f}$  (degree 0), with the universal generator dictionary

$$c_\lambda \mapsto \theta_\lambda = \Theta_\rho(B), \quad u_{a(t)dt \otimes f} \mapsto B_f(a), \quad \lambda = B \otimes \rho.$$

This real-current triangle is the interval contraction of the native two-dimensional holomorphic singular product computed in Theorem 1.15. Before smearing along the brane line, the holomorphic singular coefficient is

$$K_{\Delta}^{(2)}(z, w) = \frac{d\zeta_1 d\zeta_2}{\zeta_1 \zeta_2}, \quad \zeta_i = z_i - w_i,$$

and the displayed three brackets are the (0, 0)-mode products after pairing with the interval currents. The reduced algebraic current target is the freely adjoined completed graded-commutative algebra  $A_{\mathfrak{d}}^{\text{cur}}(I) = \widehat{\mathbf{S}}(\mathfrak{h}_I) \widehat{\otimes} \widehat{\mathbf{S}}(\mathcal{P}_I[1])$  on these generators. All four bracket signs and the displayed generator dictionary are

forced by the symplectic form  $\Omega_{\text{loc}} = dz_1 \wedge dz_2$ , the residue pairing  $\langle \rho_{a,b}, z_1^m z_2^n \rangle = \delta_{a,m} \delta_{b,n}$ , and the equal-time bracket of Remark 4.10; the displayed signs match Lemmas C.2–C.9 of Appendix C verbatim.

*Three notions of locality, formal apparatus.* The body proves three independent locality statements in Proposition 1.34: support locality ( $\delta(t - t')$  on the brane line, with  $(t, t')$  two brane-time points and  $s$  reserved for the transverse  $\mathbb{R}_s$  coordinate of  $X$ ), topological locality ( $L_v = [Q, \iota_v]$  in the  $\mathbb{R}_{\text{top}}^2$  factor), and formal-holomorphic locality ( $L_{\partial/\partial \bar{z}_i} = [Q, \iota_{\partial/\partial \bar{z}_i}]$  in the  $\widehat{\mathbb{C}}_{\text{o,hol}}^2$  factor). This appendix supplies the formal apparatus for each:

- (i) *Support locality* is encoded by the cosheaf Lie-algebra  $\mathfrak{h}_I = \Omega_c^\bullet(I) \widehat{\otimes} \mathfrak{h}$  and its current dual  $\mathcal{P}_I = \mathcal{D}'_c(I) \widehat{\otimes} \mathcal{P}$  of Definitions B.1, B.2. Multiplication of a current by a smooth compactly supported test function (Lemma B.4) is the only distributional operation used in the reduced current model; for disjoint supports it kills the bracket and produces the strict factorization product. The smeared form  $ab$  of the equal-time contact  $\delta(t - t')$  of Remark 4.10 is the same operation read on the source side.
- (ii) *Topological locality* is encoded by the de Rham differential  $d_I = d_{\text{dR}} \otimes 1$  on  $\Omega_c^\bullet(I)$  and by the compact-support density shift of Lemma 4.8: the surviving cohomology along the brane line is the density line, so the factorization algebra is locally constant in the topological direction. The interval factor of every current operator below is a compactly supported de Rham object; topological motion changes it by a  $d_I$ -boundary.
- (iii) *Formal-holomorphic locality* is encoded by the residue pairing  $\langle \rho, f \rangle = \text{Res}_o(\rho f)$  on the formal holomorphic polydisk  $(\widehat{\mathbb{C}}_o, \Omega_{\text{loc}})$  (Proposition B.3), realized through the Matlis principal-part module  $\mathcal{P}$  of § 8. This produces the Taylor Hamiltonian source side and its residue-dual principal-part current target side. The polynomial cotangent realization is a different label system (Corollary 5.172, Corollary C.8). For two holomorphic points the same residue formalism is the diagonal local-cohomology module  $H_\Delta^2(\mathbb{C}[[z, w]]) d\zeta_1 d\zeta_2$  of Theorem 1.15.

Each of the three localities is independent: the three operators  $\delta(t - t')$ ,  $L_v$ ,  $L_{\partial/\partial \bar{z}_i}$  live in different directions and none implies the others. Convention drift in this appendix would propagate immediately into all three sides; this is the reason for stating the apparatus explicitly here once.

*Prefactorization versus factorization.* All structure constructed in this appendix is prefactorization in the sense of Costello–Gwilliam [15, Ch. 3]: cosheaf source on intervals, completed graded-commutative target, extension-by-zero factorization product, and strict vanishing of mixed brackets between disjoint-support generators. Weiss-cover descent and the upgrade to a factorization algebra in the Weiss-cover sense is the responsibility of the body Theorems 4.11 and 5.185, which interpret this appendix's prefactorization datum as a factorization algebra on the brane line under the locality hypotheses of Proposition 1.34. The appendix supplies the prefactorization datum and verifies its convention compatibility; the Weiss-cover extension is proved by the body theorems cited above. The native holomorphic singular product on  $\mathbb{C}^2$  is already the formal diagonal calculation of Theorem 1.15; the Weiss-cover issue is the real-current and unreduced graph extension of that structure.

*Convention table.* Every later identity in this appendix uses the following degree/sign/topology conventions. They match Appendix C verbatim and are repeated here because this appendix fixes conventions used in every other factorization-current statement in the manuscript.

(BV degrees)  $|c_\lambda| = 1$ ,  $|u_{a(t)dt \otimes f}| = 0$ ,  $|\theta_\lambda| = |\Theta_\rho(B)| = 1$ ,  $|B_f(a)| = 0$ ,  $|\gamma_1| = 1$ . The source CE differential  $d_{\text{CE}}$  and the target Schouten/QME differential  $d_\pi$  (resp.  $d_{\text{QME}} = Q + \{I_o, -\}$ ) have degree  $+1$ ; the

open  $P_o$ -bracket on the reduced current target has degree 0; the bulk-to-defect convolution curvature  $\text{Curv}_{\mathbf{K}}$  has degree +1; the BV Laplacian  $\Delta$  has degree +1 wherever it is invoked.

(Hamiltonian)  $\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$ ,  $\Omega_{\text{loc}} = dz_1 \wedge dz_2$ ,  $\iota_{X_f} \Omega_{\text{loc}} = -df$ ,  $\{f, g\} = \partial_{z_1} f \partial_{z_2} g - \partial_{z_2} f \partial_{z_1} g$ ,  $\{z_1, z_2\} = 1$ ,  $\text{div}_{\Omega_{\text{loc}}} X_f = 0$ . The Moyal first quantum correction has coefficient  $1/24$  (Lemma C.6).

(Residue pairing)  $\rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2$ ,  $a, b \geq 0$ ,  $(a, b) \neq (0, 0)$ ; residue normalization  $\langle \rho_{a,b}, z_1^m z_2^n \rangle = \text{Res}_0(\rho_{a,b} z_1^m z_2^n) = \delta_{a,m} \delta_{b,n}$ ; coadjoint sign  $(f \cdot \rho)(g) = -\rho(\{f, g\})$ ; resulting monomial action  $z_1^p z_2^q \cdot \rho_{a,b} = -(pb - qa + p - q) \rho_{a-p+1, b-q+1}$ . This is the universal cotangent normalization of Corollary C.8.

(Brane-line currents)  $I \subset \mathbb{R}_{\text{brane}} \subset \mathbb{R}_{\text{top}}^2$  is an oriented open interval;  $\Omega_c^\bullet(I)$  is the de Rham complex of compactly supported smooth forms (degree  $\leq 1$  on a 1-manifold);  $\mathcal{D}'(I)$  is the space of compactly supported distributions in the duality  $\mathcal{D}'(I) = C^\infty(I)^\vee$  of de Rham 0-currents ([36, Ch. 6]); the de Rham differential is  $d_I = d_{\text{dR}} \otimes 1$ . The only distributional operation on the reduced current model is multiplication  $B \mapsto aB$  of a current by a smooth compactly supported test function (Lemma B.4); products of two distributions occur only inside the explicitly flagged Costello graph layers of Propositions B.17–B.20.

(PV generator dictionary)  $c_\lambda \mapsto \theta_\lambda$ ,  $u_{a(t)dt \otimes f} \mapsto B_f(a)$ , with  $\lambda = B \otimes \rho \in \mathcal{P}_I$ ; the symbol  $\theta_\lambda$  is interchangeable with  $\Theta_\rho(B)$ , where the upper-case form is the boundary-observable physics-side notation and the lower-case form is the PV/CE-image side coordinate. Both are degree 1 and abelian on the cotangent current ideal.

(Singular product) The interval brackets are the support-local contraction of the two-dimensional holomorphic singular product of Theorem 1.15. In unsmeared holomorphic variables  $z, w \in \mathbb{C}^2$ ,

$$B_f(z)B_g(w) \sim K_\Delta^{(2)}(z, w)B_{\{f, g\}}(w), \quad B_f(z)\Theta_\rho(w) \sim K_\Delta^{(2)}(z, w)\Theta_{f \cdot \rho}(w),$$

and  $\Theta_\rho(z)\Theta_\sigma(w) \sim 0$ . The brane-line formulas above are obtained by pairing this diagonal coefficient with compactly supported interval currents and keeping the (0, 0) singular product.

(Algebra and completions) All completed symmetric algebras are interpreted degreewise in the chosen current category: finite symmetric powers are formed before completion, and brackets extend continuously from the displayed length-one current formulas; tensor products are completed projective tensor products of nuclear Fréchet spaces in the de Rham/distribution direction and  $\mathfrak{m}$ -adic in the formal Hamiltonian direction (§ 8).

The factorization-current theorem uses the following algebraic source/target interface. The interface is used before the admissible bar-cobar/enveloping criterion: interval Hamiltonian currents form the CE source, distributional Matlis principal parts form the cotangent current target, and the CE/PV generator dictionary is the one fixed by the convention table above. The proved content is the reduced algebraic current model after de Rham-current contraction. The unreduced BV/QME current lift and the Costello brane-defect graph theorem are separate obstruction problems, isolated in Remark 5.189, Appendix E, and Problem 5.212. The geometry throughout is the formal-local mixed HT model on  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}^2}_{o, \text{hol}}$ ; the interval  $I$  is an oriented topological current interval in the brane line, and the Hamiltonian labels live on the formal holomorphic symplectic polydisk  $(\widehat{\mathbb{C}^2}_o, \Omega_{\text{loc}})$ .

*Definition B.1 (Interval Hamiltonian current algebra).* For an open interval  $I \subset \mathbb{R}$ , set

$$\mathfrak{h}_I = \Omega_c^\bullet(I) \widehat{\otimes} \mathfrak{h},$$

with differential  $d_I = d_{\text{dR}} \otimes 1$ . Here

$$\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$$

is the Hamiltonian Lie algebra of the formal holomorphic symplectic disk  $(\widehat{\mathbb{C}^2_o}, \Omega_{\text{loc}})$ , concentrated in degree 0. The graded bracket is

$$[\alpha \otimes f, \beta \otimes g] = \alpha \wedge \beta \otimes \{f, g\}, \quad \alpha, \beta \in \Omega_c^\bullet(I), f, g \in \mathfrak{h}.$$

For degree-zero test functions  $a, b \in \Omega_c^0(I)$ ,

$$[a \otimes f, b \otimes g] = ab \otimes \{f, g\}.$$

The cosheaf structure for inclusions of intervals is extension by zero on  $\Omega_c^\bullet$ , tensored with the identity on  $\mathfrak{h}$ . Polynomial formulas use the dense scalar-reduced subalgebra  $\bar{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1 \subset \mathfrak{h}$ . For  $j : I \hookrightarrow J$ , extension by zero is smooth because every element of  $\Omega_c^\bullet(I)$  has support compactly contained in  $I$ . On CE cochains the induced map is the dual coproduct/pullback; the covariant factorization structure is the compact-support current target.

The notation  $a(t)dt \otimes f$  is the oriented density label paired with distributional cotangent currents. The Hamiltonian generator in  $\mathfrak{h}_I$  remains  $a \otimes f$ .

*Definition B.2 (Distributional Matlis current dual).* Let

$$\mathcal{P} = \text{Ann}_{\text{res}}(\mathbb{C} \cdot 1) \cong \mathfrak{h}_{\text{cont}}^\vee$$

be the reduced Matlis principal-part module of Section 8. Its interval current form is

$$\mathcal{P}_I = \mathcal{D}'_c(I) \widehat{\otimes} \mathcal{P}.$$

This is the selected support-local current dual used after contraction. The full strong dual of  $\Omega_c^\bullet(I) \widehat{\otimes} \mathfrak{h}$  is a separate functional-analytic object. The pairing with degree-zero Hamiltonian currents is

$$\langle B \otimes \rho, a \otimes f \rangle = B(a) \text{Res}(\rho f), \quad B \in \mathcal{D}'_c(I), \rho \in \mathcal{P}.$$

The residue is the formal residue determined by  $dz_1 \wedge dz_2$  on  $\widehat{\mathbb{C}^2_o}$ . If  $B = b(t)dt$  is a regular compactly supported density, then

$$B(a) = \int_I a(t)b(t) dt.$$

The shifted cotangent interval Lie algebra is

$$\mathfrak{g}_I = \mathfrak{h}_I \ltimes \mathcal{P}_I[1],$$

where  $\mathfrak{h}_I$  acts on  $\mathcal{P}_I$  by the product of the  $C_c^\infty(I)$ -module operation  $B \mapsto aB$  on compactly supported currents and the Matlis coadjoint action on  $\mathcal{P}$ . On degree-zero labels this action is

$$(a \otimes f) \cdot (B \otimes \rho) = (aB) \otimes (f \cdot \rho).$$

**PROPOSITION B.3** (*Local unimodular datum for the current kernels*). The current and kernel constructions rest on the following formal-local unimodular datum:

$$(\widehat{\mathbb{C}^2_o}, \Omega_{\text{loc}} = dz_1 \wedge dz_2).$$

This datum supplies:

- (i) the Poisson bracket on  $\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$ ;
- (ii) the Hamiltonian vector-field convention

$$\iota_{X_f} \Omega_{\text{loc}} = -df, \quad X_f = (\partial_{z_1} f) \partial_{z_2} - (\partial_{z_2} f) \partial_{z_1};$$

- (iii) the divergence-free identity  $\text{div}_{\Omega_{\text{loc}}} X_f = 0$ , hence the residue integration-by-parts formula defining the Matlis coadjoint action;
- (iv) the principal-part residue pairing

$$\text{Res}_0 \left( z_1^{-i-1} z_2^{-j-1} dz_1 dz_2 z_1^m z_2^n \right) = \delta_{i,m} \delta_{j,n}.$$

- (v) the diagonal two-point principal-part module

$$H_{\Delta}^2(\mathbb{C}[[z_1, z_2, w_1, w_2]]) d\zeta_1 d\zeta_2$$

and its Cauchy generator  $K_{\Delta}^{(2)}$ , which carry the native holomorphic singular product of Theorem 1.15.

At the reduced current-interface level the topological factor contributes the compactly supported de Rham current direction  $I \subset \mathbb{R}_{\text{brane}}$ , the operation  $B \mapsto aB$ , and extension by zero. In the graph layer it also contributes the local heat-kernel variables on  $\mathbb{R}_{\text{top}}^2$ . Finite-window and weighted kernels are completions of the formal Hamiltonian coefficient system.

*Proof.* The bracket in Definition B.1 is the inverse bivector of  $dz_1 \wedge dz_2$ , and constants are quotiented because they give the zero Hamiltonian vector field. The displayed  $X_f$  satisfies  $X_f(g) = \{f, g\}$ , and

$$\text{div}_{\Omega_{\text{loc}}} X_f = -\partial_{z_1} \partial_{z_2} f + \partial_{z_2} \partial_{z_1} f = 0.$$

Thus Hamiltonian fields preserve the residue density. The transpose of this action under  $\text{Res}(\rho f)$  is exactly the Matlis coadjoint action used in Definition B.2 and Corollary B.8.

The interval variable is independent of the holomorphic polydisk. In the reduced current interface its operations are the de Rham differential, multiplication of a current by a smooth compactly supported test function, and extension by zero. In the graph layer the additional topological variables enter only through the local heat-kernel calculus on  $\mathbb{R}_{\text{top}}^2$ . The finite-window constructions truncate only the Hamiltonian degree in the formal polydisk and then pass through the weighted Mittag-Leffler inverse limit. None of these operations leaves the displayed local current category.  $\square$

LEMMA B.4 (*Multiplication of a current by a test function*). For  $a \in C_c^\infty(I)$  and  $B \in \mathcal{D}'_c(I)$ , define

$$(aB)(\varphi) = B(a\varphi), \quad \varphi \in C_c^\infty(I).$$

Then  $aB \in \mathcal{D}'_c(I)$ ,

$$\text{supp}(aB) \subset \text{supp}(a) \cap \text{supp}(B),$$

and  $a(bB) = (ab)B$ . If  $B = b(t)dt$  is regular, then  $aB = (ab)(t)dt$ . The reduced current model uses this smooth-function action on distributions.

*Proof.* The function  $a\varphi$  is a compactly supported smooth test function, so precomposition by  $\varphi \mapsto a\varphi$  defines a compactly supported distribution. If  $\varphi$  is supported away from  $\text{supp}(a) \cap \text{supp}(B)$ , then  $a\varphi$  is supported away from  $\text{supp}(B)$ , hence  $B(a\varphi) = 0$ . Associativity follows from  $a(b\varphi) = (ab)\varphi$ . The regular-density formula is the displayed integral identity.  $\square$



*Definition B.5 (Current CE/PV coordinates).* Let  $\lambda = B \otimes \rho \in \mathcal{P}_I$ . The Hamiltonian CE coordinate dual to the Hamiltonian current through  $\lambda$  is written

$$c_\lambda, \quad |c_\lambda| = 1.$$

Its PV/current target coordinate is

$$\theta_\lambda = \theta_{B \otimes \rho} := \Theta_\rho(B).$$

The label  $a(t)dt \otimes f$  denotes the functional on  $\mathcal{P}_I$

$$B \otimes \rho \mapsto B(a) \operatorname{Res}(\rho f).$$

The CE coordinate dual to the shifted cotangent generator through this functional is

$$u_{a(t)dt \otimes f}, \quad |u_{a(t)dt \otimes f}| = 0.$$

Its PV/current target coordinate is the smeared Hamiltonian boundary observable

$$O_{a,f} = B_f(a).$$

In the brane realization,

$$B_f(a) = \int_I a(t) \overline{\operatorname{Tr} f(\phi_1(t), \phi_2(t))} dt.$$

*Definition B.6 (Enlarged reduced defect-current target).* The reduced defect-current target attached to  $I$  is

$$A_\partial^{\operatorname{cur}}(I) = \widehat{\mathbf{S}}(\mathfrak{h}_I) \widehat{\otimes} \widehat{\mathbf{S}}(\mathcal{P}_I[I]).$$

This is the reduced principal-part target denoted  $A_\partial^{\operatorname{pp}}(I)$  in the main text; the superscript *cur* records its current-interface role. It enlarges the Hamiltonian target  $A_\partial^{\operatorname{Ham}}(I) = \widehat{\mathbf{S}}(\mathfrak{h}_I)$  by adjoining the odd principal-part current generators  $\theta_\lambda = \Theta_\rho(B)$ . The structure maps for inclusions of intervals are extension by zero on  $\Omega_c^\bullet(I)$  and on  $\mathcal{D}'_c(I)$ . The factorization product on disjoint intervals is the completed graded-commutative product after extension by zero.

**THEOREM B.7 (Universal algebraic current interface).** For each interval  $I$ , the generator assignment

$$\Phi_I^{\operatorname{cur}}(c_\lambda) = \theta_\lambda, \quad \Phi_I^{\operatorname{cur}}(u_{a(t)dt \otimes f}) = B_f(a)$$

extends uniquely to a continuous morphism of completed graded-commutative algebras from the interval CE cofactorization source  $C_{\operatorname{CE}, \operatorname{cont}}^\bullet(\mathfrak{g}_I)$  to the reduced defect-current central-operation target  $A_\partial^{\operatorname{cur}}(I)$ . On length-one current generators the target open-current  $P_o$  brackets are

$$\begin{aligned} \{B_f(a), B_g(b)\} &= B_{\{f,g\}}(ab), \\ \{B_f(a), \theta_{B \otimes \rho}\} &= \theta_{aB \otimes (f \cdot \rho)} = \Theta_{f \cdot \rho}(aB), \\ \{\theta_{B \otimes \rho}, \theta_{C \otimes \sigma}\} &= 0. \end{aligned}$$

The factorization product and the reduced current source-target  $P_o$ -equivariance defects in this reduced current target are strict:

$$H_{\operatorname{fact}} = 0, \quad H_{P_o} = 0.$$

For overlapping supports, the nonzero Hamiltonian current bracket is the moment-map bracket displayed above. Thus  $u_{a(t)dt \otimes f} \mapsto B_f(a)$  and  $c_\lambda \mapsto \theta_\lambda$  form the current-level source/target interface used before the admissible bar-cobar/enveloping criterion.

*Proof.* The CE algebra of  $\mathfrak{g}_I = \mathfrak{h}_I \ltimes \mathcal{P}_I[1]$  is the completed symmetric algebra on the continuous dual shifted by  $[-1]$ . The coordinate  $c_\lambda$  has degree 1, and the coordinate  $u_{a(t)dt \otimes f}$  has degree 0. The target  $\mathcal{A}_\theta^{\text{cur}}(I)$  is the completed symmetric algebra on the corresponding current coordinates  $\theta_\lambda$  and  $B_f(a)$ . Hence the displayed assignment extends uniquely and continuously to the completed graded-commutative algebras.

The CE differential and the target current brackets use the same semidirect Lie structure constants. The Hamiltonian-current bracket in Definition B.1 gives

$$[a \otimes f, b \otimes g] = ab \otimes \{f, g\},$$

and the boundary realization gives the first displayed bracket. The action of  $a \otimes f$  on  $B \otimes \rho$  is  $(aB) \otimes (f \cdot \rho)$ , giving the second displayed bracket by negative transpose under the product pairing

$$\langle B \otimes \rho, a \otimes f \rangle = B(a) \text{Res}(\rho f).$$

The shifted cotangent current ideal is abelian, giving the third bracket. These three identities are exactly the current form of the CE/PV generator identities  $c_\lambda \mapsto \theta_\lambda$  and  $u \mapsto 0$ . Since the CE differential is a derivation and the target bracket is a biderivation, the generator identities determine the full completed current interface.

Factorization compatibility is strict in the cofactorization-source convention. Extension by zero on current Lie algebras induces the CE coproduct, and the target product is compatible with pointwise products  $ab$  and with the  $C_c^\infty(I)$ -module operation  $B \mapsto aB$ . If two intervals are disjoint, the products have empty support and the mixed brackets vanish. The target product is graded commutative, so multiplication by an image generator has zero Hochschild product-centrality defect.

For reduced current  $P_0$ -equivariance, let  $x$  be a length-one source generator and let  $Y$  be a homogeneous monomial in the CE source. The defect

$$D_x(Y) = \{\Phi_I^{\text{cur}}(x), \Phi_I^{\text{cur}}(Y)\} - \Phi_I^{\text{cur}}(\text{ad}_x Y)$$

is a derivation in  $Y$ . The three generator bracket identities give  $D_x(Y) = 0$  for every length-one  $Y$ . The Leibniz rule gives  $D_x(Y) = 0$  for every monomial, and continuity extends this to the completed algebra. Thus  $H_{\text{fact}}$  and  $H_{P_0}$  are zero equivariance-defect homotopies in the reduced current model.  $\square$

**COROLLARY B.8** (*Principal-part current brackets in monomial coordinates*). For  $f = z_1^p z_2^q$ ,  $a \in C_c^\infty(I)$ ,  $B \in \mathcal{D}'_c(I)$ , and  $\rho_{i,j} = z_1^{-i-1} z_2^{-j-1} dz_1 dz_2$ , one has

$$\{B_{z_1^p z_2^q}(a), \Theta_{\rho_{i,j}}(B)\} = -(pj - qi + p - q) \Theta_{\rho_{i-p+1, j-q+1}}(aB),$$

with negative-index targets equal to zero. If  $B = b(t)dt$  is regular, then the interval factor is  $(ab)(t)dt$ .

*Proof.* Proposition 8.7 gives

$$z_1^p z_2^q \cdot \rho_{i,j} = -(pj - qi + p - q) \rho_{i-p+1, j-q+1},$$

with negative-index terms projected to zero in the principal-part module. Substituting this action into the second bracket of Theorem B.7 gives the displayed formula. The regular-current statement is Lemma B.4.  $\square$

The next theorem is the BMK obstruction lane. The classical Bochner–Martinelli–Koppelman kernel gives a Dolbeault homotopy on  $\mathbb{C}^2 \setminus \{0\}$  with explicit Hadamard estimates. One asks whether truncating to the finite Matlis window  $\mathcal{P}_{\leq N}$  and quotienting by a section  $i_N$ ,  $p_N$  yields a strict finite-window homotopy on the full compact-support/current complex. The obstruction is the precise four-component vector  $\text{Ob}_N^{\text{BMK-curr}}$ . The analytic Taylor tower gives a one-pair pro-Matlis retract, and support-local factorization-current transfer is governed by this obstruction vector. The proved content of the theorem is the BMK current data  $(H_\chi, p_N, i_N)$ , the finite-window obstruction vector, the positive one-pair analytic pro-Matlis retract, and the arity-two Hamiltonian/Matlis comparison.

**THEOREM B.9** (*BMK current data and finite-window obstruction vector*). Let  $B_0 \Subset B_1 \Subset B_2 \Subset B \subset \mathbb{C}^2$  be nested polydisks around the collision point, and choose  $\chi \in C_c^\infty(B_2)$  with  $\chi = 1$  on  $B_1$ . Put

$$\mathcal{P}_{\leq N} = \bigoplus_{\substack{a,b \geq 0 \\ 0 < a+b \leq N}} \mathbb{C} \rho_{a,b}, \quad \rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2.$$

On  $(B_\chi \times B_z) \setminus \Delta$ , set

$$\eta = \zeta - z, \quad r^2 = |\eta_1|^2 + |\eta_2|^2, \quad d\bar{\eta}_j = d\bar{\zeta}_j - d\bar{z}_j.$$

The Dolbeault homotopy kernel is the Bochner–Martinelli–Koppelman current

$$K_{\text{BM}}(\zeta, z) = \frac{1}{(2\pi i)^2} \frac{\bar{\eta}_1 d\bar{\eta}_2 - \bar{\eta}_2 d\bar{\eta}_1}{r^4} \wedge d\zeta_1 \wedge d\zeta_2.$$

The  $d\bar{z}_j$ -terms are part of the Koppelman current. The Dolbeault homotopy uses the full Koppelman current.

Choose a radial cutoff  $\vartheta_\epsilon(r)$ , equal to 0 for  $r \leq \epsilon$  and to 1 for  $r \geq 2\epsilon$ , and put

$$K_{\chi,\epsilon} = \chi(\zeta) \chi(z) \vartheta_\epsilon(|\zeta - z|) K_{\text{BM}}(\zeta, z).$$

The current limit

$$H_\chi \alpha(z) = \lim_{\epsilon \rightarrow 0} \int_{\zeta \in B} K_{\chi,\epsilon}(\zeta, z) \wedge \alpha(\zeta)$$

exists on every fixed finite configuration stratum. For separated supports it satisfies

$$|\nabla^k K_{\text{BM}}(\zeta, z)| \leq C_k |\zeta - z|^{-3-k}.$$

Let  $i_N : \mathcal{P}_{\leq N} \rightarrow \mathcal{D}'^{0,2}(B)$  be

$$i_N(\rho_{a,b}) = \Delta_{a,b} = \frac{(-1)^{a+b}}{a! b!} \partial_{z_1}^a \partial_{z_2}^b \partial_0 d\bar{z}_1 \wedge d\bar{z}_2,$$

normalized by

$$\langle \Delta_{a,b}, f \rangle = \text{Res}_0(\rho_{a,b} f).$$

Define the finite principal-part moment projection on top Dolbeault degree by

$$p_N(\alpha) = \sum_{0 < a+b \leq N} \left( \int_B z_1^a z_2^b \alpha^{0,2} \wedge dz_1 \wedge dz_2 \right) \rho_{a,b}.$$

The data above construct  $H_\chi$ ,  $p_N$ , and  $i_N$ . A finite-window homotopy on the full compact-support/current complex would require the identity

$$\bar{\partial}H_{\chi,N} + H_{\chi,N}\bar{\partial} = \text{id} - i_N p_N + E_{\chi,N} \quad (*_N^{\text{BMK}})$$

is blocked by

$$\text{Ob}_N^{\text{BMK-curr}} = (\text{ob}_{\text{diag}}, \text{ob}_{\text{an/fin},N}, \text{ob}_{\mathfrak{h},N}, \text{ob}_{\text{collar-fact},N}).$$

The first entry is the annular diagonal sign and scalar of  $\bar{\partial}\mathfrak{D}_\epsilon \wedge K_{\text{BM}}$  relative to the derivative-delta Matlis convention. With the displayed kernel and real outward boundary orientation the angular mass is  $-1$ . The second entry is the gap between the full analytic moment kernel  $\ker(O(B_1)^\vee \rightarrow \mathcal{P}_{\leq N})$  and the smaller support-at-zero span generated by the scalar and high derivative-delta currents. The third entry is

$$\text{ob}_{\mathfrak{h},N} = \pi_N(\mathfrak{h} \cdot \ker \pi_N),$$

which is nonzero because

$$z_1^{N+2} \cdot \rho_{N+1,0} = -(N+2)\rho_{0,1}.$$

Thus the full Hamiltonian action sends high modes back into the retained window, obstructing a strict finite support-local quotient that retains  $\mathcal{P}_{\leq N}$ . The fourth entry is the cosheaf/factorization compatibility of the collar relation with extension by zero, disjoint products, and the BMK homotopy.

A normalized finite-window identity is available only after changing the underlying complex. If the full analytic BMK contraction is supplied and the relative complex is quotiented by representatives of  $\ker(O(B_1)^\vee \rightarrow \mathcal{P}_{\leq N})$ , then the normalized homotopy

$$(\text{id} - I_{\text{an}} P_{\text{an}})H_\chi(\text{id} - I_{\text{an}} P_{\text{an}})$$

descends and gives

$$\bar{\partial}H_{\text{BM},N} + H_{\text{BM},N}\bar{\partial} = \text{id} - i_N p_N.$$

This is a one-pair analytic quotient. Support-locality on the factorization-current complex is measured by  $\text{Ob}_N^{\text{BMK-curr}}$ .

Keeping the full Taylor tower gives the positive one-pair pro-Matlis retract. Put

$$\widehat{\mathcal{P}}_\Pi = \prod_{\substack{a,b \geq 0 \\ a+b > 0}} \mathbb{C}\rho_{a,b}, \quad \mathcal{P}_{\text{an}}(B_1) = \tau(O(B_1)_{\text{red}}^\vee) \subset \widehat{\mathcal{P}}_\Pi,$$

where

$$\tau(\lambda) = (\lambda(z_1^a z_2^b))_{a+b > 0}.$$

After the diagonal orientation is fixed and the collar quotient kills the cutoff error, the normalized analytic contraction  $H_\chi^\circ = (\text{id} - I_{\text{an}} P_{\text{an}})H_\chi(\text{id} - I_{\text{an}} P_{\text{an}})$  satisfies

$$\bar{\partial}H_\chi^\circ + H_\chi^\circ \bar{\partial} = \text{id} - I_{\text{an}} P_{\text{an}}.$$

The pro-analytic moment map  $P_\Pi^{\text{an}} = \tau P_{\text{an}}$  has compatible finite projections

$$\pi_N P_\Pi^{\text{an}} = p_N, \quad \pi_{M,N} p_M = p_N.$$

This is a one-pair analytic pro-Matlis statement. A support-local factorization-current transfer additionally requires vanishing of  $\text{Ob}_N^{\text{BMK-curr}}$ .

The binary coefficient operation

$$\ell_2^{\text{BM},N}(x, y) = p_N[ix, iy]$$

is retained as the Hamiltonian/Matlis arity-two comparison, read in the output window  $\mathcal{P}_{\leq N}$ . For monomials,

$$\ell_2^{\text{BM},N}(z_1^p z_2^q, z_1^r z_2^s) = (ps - qr) \text{pr}_{\leq N}(z_1^{p+r-1} z_2^{q+s-1}) \pmod{\mathbb{C}},$$

and

$$\ell_2^{\text{BM},N}(z_1^p z_2^q, \rho_{i,j}) = -(pj - qi + p - q) \text{pr}_{\leq N} \rho_{i-p+1, j-q+1},$$

with negative-index targets equal to zero, while

$$\ell_2^{\text{BM},N}(\rho, \sigma) = 0.$$

Thus the arity-two cotangent coefficient comparison is the Matlis coadjoint action of Proposition 8.7.

Only after the finite-current condition  $\text{Ob}_N^{\text{BMK-curr}}$  is killed, or replaced by a support-local factorization-current complex, does the strict all-window and arity-three extension become the following obstruction vector. Set

$$Q_{\oplus} = \widehat{\mathcal{P}}_{\Pi} / \mathcal{P}.$$

For  $\alpha \in \Omega_c^{\circ,2}(B)$ , put

$$\text{mom}_{\infty}(\alpha) = \left( \int_B z_1^a z_2^b \alpha^{\circ,2} \wedge dz_1 \wedge dz_2 \right)_{a+b>0} \in \widehat{\mathcal{P}}_{\Pi},$$

and

$$\text{ob}_{\oplus}(\alpha) = [\text{mom}_{\infty}(\alpha)] \in Q_{\oplus}.$$

This class is zero exactly when the all-window moment sequence lands in the direct-sum Matlis module  $\mathcal{P}$ .

The pro-product tower leaves the current problem governed by the section obstruction. A support-local compact-current section

$$i_{\Pi} : \widehat{\mathcal{P}}_{\Pi} \rightarrow \mathcal{D}'^{\circ,2}(B)$$

extending the finite derivative-delta sections  $i_N$  would assign infinite-order point-supported distributions, while point-supported distributions have finite order. This is  $\text{ob}_{I_{\Pi}}$ . The action of the full formal Hamiltonian algebra on  $\widehat{\mathcal{P}}_{\Pi}$  is obstructed at  $\text{ob}_{\mathfrak{h},\Pi}$ . For  $f = \sum_{p \geq 2} z_1^p$  and  $\lambda = \sum_{p \geq 2} \rho_{p-1,0}$ , the  $\rho_{0,1}$ -coefficient of  $f \cdot \lambda$  collects  $-\sum_{p \geq 2} p$ , the divergent witness  $\text{ob}_{\mathfrak{h},\Pi}$ .

The collar part is the finite-window family

$$\text{ob}_{\text{collar},\Pi,N} = E_{\chi,N} \in \text{Hom}(\Omega_c^{\circ,\bullet}(B), \mathcal{D}'^{\circ,\bullet}(B))_{\text{supp } d\chi}.$$

The first arity-three obstruction is

$$\text{ob}_{3,\Pi,N}^{\chi}(x, y, z) = p_N[H_{\text{BM},N}([ix, iy] - i\ell_2^{\text{BM},N}(x, y)), iz] + \text{cyc}_{x,y,z}.$$

It vanishes on separated-support strata by the displayed estimates. On the coefficient collision stratum its non-collar part is the Jacobiator of  $\mathfrak{h} \ltimes \mathcal{P}[1]$ , hence zero. The remaining part is supported on the small diagonal and on the cutoff collar.

Finally, let  $C_{s,N}$  be the optimal constants for the chosen finite-window seminorm estimates of  $H_{\chi,N}$ ,  $p_N$ ,  $E_{\chi,N}$ , and  $\text{ob}_{3,\Pi,N}^{\chi}$ . The uniformity obstruction is

$$\text{ob}_{\text{unif},\Pi} = \left( [(C_{s,N})_{N \geq 1}] \right)_s \in \prod_s \left( \prod_{N \geq 1} \mathbb{R}_{\geq 0} / \ell_N^{\infty} \right).$$

Thus the all-window obstruction vector is

$$\text{Ob}_{\text{BM}}^{\Pi} = \left( \text{ob}_{\oplus}, \text{ob}_{I_{\Pi}}, \text{ob}_{\mathfrak{h},\Pi}, (\text{ob}_{\text{collar},\Pi,N})_N, (\text{ob}_{3,\Pi,N}^{\chi})_N, \text{ob}_{\text{unif},\Pi} \right).$$

Any strict all-window current  $E_2$  transfer extending the displayed binary coefficient comparison forces this vector to vanish. The first, second, third, and sixth entries are nonzero in the current topology; the collar and ternary entries are exact primitive construction targets. The proved theorem here is the BMK current data, the finite-window obstruction vector, the positive one-pair analytic pro-Matlis retract, and the arity-two Hamiltonian/Matlis comparison.

The principal-part face remains two-variable:

$$\frac{1}{(2\pi i)^2} \frac{d\zeta_1 \wedge d\zeta_2}{(\zeta_1 - z_1)(\zeta_2 - z_2)} = \sum_{a,b \geq 0} z_1^a z_2^b \rho_{a,b}(\zeta).$$

Setting  $z_2 = 0$  kills the Hamiltonian bracket, and the  $b = 0$  principal-part line leaves the subspace under  $z_1 \cdot \rho_{a,0} = -\rho_{a,1}$ ; the principal-part face stays two-variable.

*Proof.* The Bochner–Martinelli–Koppelman identity gives the Dolbeault homotopy on the punctured product  $B_{\zeta} \times B_z \setminus \Delta$ . Multiplication by the cutoff gives the exact derivative split

$$\bar{\partial}(\chi(\zeta)\chi(z)\vartheta_{\epsilon}K_{\text{BM}}) = \chi\chi\vartheta_{\epsilon}\bar{\partial}K_{\text{BM}} + \bar{\partial}(\chi\chi)\vartheta_{\epsilon}K_{\text{BM}} + \chi\chi\bar{\partial}\vartheta_{\epsilon} \wedge K_{\text{BM}}.$$

The middle term is collar-supported. The last term is the annular diagonal current; its sign and scalar are exactly  $\text{ob}_{\text{diag}}$ . Since  $K_{\text{BM}}$  has size  $r^{-3}$  in real dimension four, the cutoff current has a limit on every fixed finite configuration stratum. This proves the current data. The finite-window identity on the full current complex is the additional obstruction equation displayed in the theorem.

The formula for  $i_N$  is fixed by integration by parts:

$$\left\langle \frac{(-1)^{a+b}}{a!b!} \partial_{z_1}^a \partial_{z_2}^b \delta_0, f \right\rangle = [z_1^a z_2^b]f = \text{Res}_0(\rho_{a,b}f).$$

Hence  $i_N p_N$  is the finite support-at-collision projector with the Matlis residue normalization, after the diagonal-current orientation is tied to that convention.

The finite-window identity on the full current complex is obstructed already on closed top currents in the kernel of the finite moment map. If  $T = \Delta_{N+1,0}$ , then  $p_N(T) = 0$  but  $\langle T, z_1^{N+1} \rangle = 1$ . A putative identity  $T = \bar{\partial}S$  would give  $\langle T, z_1^{N+1} \rangle = 0$  by Stokes. The scalar  $\Delta_{0,0}$  is the same obstruction for the removed reduced principal-part line. More strongly, a strict support-local quotient must be an  $\mathfrak{h}$ -submodule quotient; the calculation  $z_1^{N+2} \cdot \rho_{N+1,0} = -(N+2)\rho_{0,1}$  contradicts killing  $\rho_{N+1,0}$  while retaining  $\rho_{0,1}$ .

The one-pair analytic quotient works for the reason stated in the theorem: after choosing the full analytic projector  $I_{\text{an}} P_{\text{an}}$ , the normalized homotopy kills representatives of  $\ker \pi_N$ , descends to the quotient, and identifies the analytic projector with  $i_N p_N$ . Keeping the Taylor tower instead gives the pro-analytic map  $P_{\Pi}^{\text{an}} = \tau P_{\text{an}}$ , and the equality  $\pi_N P_{\Pi}^{\text{an}} = p_N$  is just functoriality of Taylor moments under finite truncation. Its dependence on analytic representatives and on the chosen pair makes it a one-pair analytic quotient.

The arity-two coefficient comparison is the residue transpose. For Hamiltonian labels it is the Poisson bracket modulo constants, truncated only after the output coefficient is computed. For cotangent principal parts, Proposition 8.7 gives

$$\text{Res}_o((f \cdot \rho)g) = -\text{Res}_o(\rho \{f, g\}),$$

which is the displayed coadjoint formula. The shifted cotangent ideal is abelian, so the  $(\rho, \sigma)$ -operation is zero.

The displayed ternary expression is only the next obstruction after the finite-current condition has been crossed. The separated-support estimate makes it vanish away from collisions. On the coefficient collision stratum the semidirect Lie algebra  $\mathfrak{h} \ltimes \mathcal{P}[1]$  satisfies Jacobi. Therefore any surviving term is a collar, small-diagonal, or completed-window transfer residue, recorded as the finite-window representative  $\text{ob}_{3,\Pi,N}^{\chi}$  of the  $\Pi$ -tower class only after the finite-current obstruction has been removed.

A strict all-window current transfer would have to replace the compatible family  $p_N$  by a single moment map landing in  $\mathcal{P}$ , construct a compact-current section of the pro-product tower, define the full formal Hamiltonian action on that tower, remove  $E_{\chi,N}$  compatibly with extension by zero, kill the ternary collar residue by compatible primitives, and supply uniform estimates in the completed window topology. These six requirements are precisely the six entries of  $\text{Ob}_{\text{BM}}^{\Pi}$ . The all-window theorem target is a refined factorization-current complex where these six witnesses exist, together with the displayed obstruction theorem. The finite-window calculation supplies the inputs measured by  $\text{Ob}_N^{\text{BMK-curr}}$ .

The Cauchy expansion is the Laurent expansion in the two holomorphic variables and records the same residue basis. The calculation  $z_1 \cdot \rho_{a,o} = -\rho_{a,1}$ , from Proposition 8.7, rules out the  $b = o$  principal-part truncation as an  $\mathfrak{h}$ -submodule.  $\square$

The next theorem switches on the brane-preserving normal  $\Omega$ -background. In the local dictionary the brane-preserving equivariant deformation is normal scaling of  $N_L X = \mathbb{R}_s \oplus \mathbb{C}_{z_1} \oplus \mathbb{C}_{z_2}$ , with  $t$  fixed. The fixed brane line selects this normal action. The current sector responds by an explicit grading on the generators  $B_{p,q}(a)$  and  $\Theta_{r,s}(B)$ ; the Hamiltonian/Matlis brackets become weighted by the inverse symplectic-character line  $\mathbb{L}_{\omega}^{-1}$ , or equivalently by adjoining a weight- $\chi_{\omega}$  parameter  $\hbar_{\omega}$ . The theorem records the resulting triangle of length-one current brackets and the self-dual specialization  $\epsilon_1 + \epsilon_2 = o$ . This is an algebraic identity in the freely adjoined current target. A normal- $\Omega$  graph/QME lift additionally requires the  $Q_{\Omega}$ -Cartan data, normal-weight contraction, residue/Euler normalization choice, equivariant counterterms, and the scalar/non-scalar QME obstruction classes named below.

**THEOREM B.10** (*Normal  $\Omega$ -weighted brane-current brackets*). Use the normal brane-preserving equivariant parameters of the local dictionary:  $t$  has weight  $o$ ,  $z_1, z_2$  have weights  $\epsilon_1, \epsilon_2$ , and

$$\chi_{\omega} = \epsilon_1 + \epsilon_2, \quad \text{wt}_{\Omega}(\hbar_{\omega}) = \chi_{\omega}.$$

Put

$$B_{p,q}(a) := B_{z_1^p z_2^q}(a), \quad \Theta_{r,s}(B) := \Theta_{\rho_{r,s}}(B),$$

with  $p, q, r, s \geq 0$ ,  $p + q > 0$ ,  $r + s > 0$ , and  $\rho_{r,s} = z_1^{-r-1} z_2^{-s-1} dz_1 dz_2$ . The compactly supported test function  $a$  and compactly supported current  $B$  live on the fixed brane line and have weight 0. Hence

$$\mathrm{wt}_\Omega B_{p,q}(a) = p\epsilon_1 + q\epsilon_2, \quad \mathrm{wt}_\Omega \Theta_{r,s}(B) = -r\epsilon_1 - s\epsilon_2.$$

Off the self-dual locus the Hamiltonian current bracket is valued in the inverse symplectic-character line. After adjoining  $\hbar_\omega$ , the scalarized convention has length-one current brackets are

$$\begin{aligned} \{B_{p,q}(a), B_{r,s}(b)\}_\Omega &= \hbar_\omega (ps - qr) B_{p+r-1, q+s-1}(ab), \\ \{B_{p,q}(a), \Theta_{r,s}(B)\}_\Omega &= -\hbar_\omega (ps - qr + p - q) \Theta_{r-p+1, s-q+1}(aB), \\ \{\Theta_{r,s}(B), \Theta_{u,v}(C)\}_\Omega &= 0. \end{aligned}$$

Here  $B_{\alpha,\beta} = 0$  if  $\alpha < 0$ ,  $\beta < 0$ , or  $(\alpha, \beta) = (0, 0)$ , and  $\Theta_{\alpha,\beta} = 0$  if  $\alpha < 0$ ,  $\beta < 0$ , or  $(\alpha, \beta) = (0, 0)$ . The last convention only records the reduced scalar-annihilator target; when the coadjoint formula lands at  $\rho_{0,0}$  its coefficient is already zero.

Equivalently, before adjoining  $\hbar_\omega$ , remove the displayed factor  $\hbar_\omega$  and read the first two brackets as valued in the inverse symplectic-character line  $\mathbb{L}_\omega^{-1}$ , where  $\mathrm{wt}_\Omega(\mathbb{L}_\omega) = \chi_\omega$ . On the self-dual specialization  $\epsilon_1 + \epsilon_2 = 0$ , the line is trivial. After choosing the trivialization  $\hbar_\omega = 1$ , the formulas reduce to the non-equivariant brackets of Theorem B.7 and Corollary B.8. This specialization is a separate branch: first restrict to  $\epsilon_1 + \epsilon_2 = 0$ , then choose the trivialization.

For an inclusion of intervals  $j : I \hookrightarrow J$ , extension by zero has weight 0 and satisfies

$$j_! B_{p,q,I}(a) = B_{p,q,J}(j_! a), \quad j_! \Theta_{r,s,I}(B) = \Theta_{r,s,J}(j_! B),$$

together with

$$j_! \{x, y\}_{\Omega, I} = \{j_! x, j_! y\}_{\Omega, J}$$

for the displayed length-one generators. For disjoint intervals the mixed brackets vanish after extension to a common interval.

These statements are algebraic current identities. They hold for all  $B, C \in \mathcal{D}'_c(I)$  because the only distributional operation is multiplication of a current by a smooth compactly supported function. A normal- $\Omega$  graph/QME lift for arbitrary distributional labels is a further analytic theorem. A graph-level lift must still restrict cotangent labels to regular densities or to graphwise wavefront-admissible tuples as in Propositions B.17 and B.19, and must add the  $Q_\Omega$ -Cartan data, normal-weight contraction, residue/Euler normalization choice, equivariant counterterms, and the scalar/non-scalar QME obstruction classes.

*Proof.* The weight assignments are forced by the normal torus action:  $z_1^p z_2^q$  has character  $p\epsilon_1 + q\epsilon_2$ , while

$$z_1^{-r-1} z_2^{-s-1} dz_1 dz_2$$

has character  $-(r+1)\epsilon_1 - (s+1)\epsilon_2 + \chi_\omega = -r\epsilon_1 - s\epsilon_2$ . The brane-line variables and currents are in the fixed  $t$ -direction, so their equivariant character is 0.

The Poisson bivector  $\partial_{z_1} \wedge \partial_{z_2}$  has character  $-\chi_\omega$ . Thus the ordinary Hamiltonian bracket has weight  $\mathrm{wt}(f) + \mathrm{wt}(g) - \chi_\omega$ . Multiplying by  $\hbar_\omega$  makes the bracket scalar and weight-preserving. On monomials this gives

$$[z_1^p z_2^q, z_1^r z_2^s]_\Omega = \hbar_\omega (ps - qr) z_1^{p+r-1} z_2^{q+s-1},$$

followed by the quotient by constants. Substitution in the first bracket of Theorem B.7 gives the displayed  $B$ - $B$  formula.



For the cotangent current, the coadjoint action of Proposition 8.7 gives

$$z_1^p z_2^q \cdot \rho_{r,s} = -(ps - qr + p - q) \rho_{r-p+1, s-q+1}.$$

Scalarizing the Hamiltonian action by the same  $\hbar_\omega$  gives the  $B$ - $\Theta$  formula, and the shifted cotangent current ideal is abelian. Each right-hand side has total weight equal to the sum of the weights of the two inputs, so the formulas are  $T_\Omega$ -homogeneous.

Extension by zero commutes with multiplication of compactly supported test functions and currents:  $j_i(ab) = (j_i a)(j_i b)$  and  $j_i(aB) = (j_i a)(j_i B)$ . Hence it commutes with all three displayed brackets. If two intervals are disjoint inside a common interval, these products are zero, so the mixed brackets vanish.

Finally, the algebraic current formulas use multiplication of currents by smooth compactly supported functions. Costello graph weights are different: after pushing current labels through graph incidence maps one must multiply them by singular propagator boundary values and extend across collision diagonals. The wavefront and scaling-degree hypotheses in Propositions B.19 and B.20 are exactly the required analytic conditions. The  $\Omega$ -weights and the factor  $\hbar_\omega$  preserve those microlocal conditions.  $\square$

**COROLLARY B.11** (*Smeared Hamiltonian source convention*). The smeared boundary Hamiltonian observable is sourced by the shifted-cotangent CE coordinate:

$$u_{a(t)dt \otimes f} \mapsto B_f(a) = \int_I a(t) \overline{\text{Tr} f(\phi_1(t), \phi_2(t))} dt.$$

The Hamiltonian CE coordinate  $c_\lambda$  maps to the PV/current vector  $\theta_\lambda$ . Consequently

$$\{B_f(a), B_g(b)\} = B_{\{f,g\}}(ab)$$

is the degree-zero moment-map shadow of the CE/PV identity  $c_\lambda \mapsto \theta_\lambda$ ,  $u \mapsto O$ .

*Proof.* The coordinate  $u_{a(t)dt \otimes f}$  is dual to the shifted cotangent generator through the pairing with  $\mathcal{P}_I$ . Under the CE/PV isomorphism it maps to the function coordinate  $O_{a,f}$ , realized on the brane by the displayed trace integral. The coordinate  $c_\lambda$  is dual to a Hamiltonian current and maps to the vector/current coordinate  $\theta_\lambda$ . The bracket identity is the first bracket of Theorem B.7; equivalently, it is the equal-time Poisson bracket smeared against  $a(t)b(t')\delta(t-t')$  on two brane-time points, which produces the pointwise product  $ab$ .  $\square$

**Remark B.12** (*Orientation reversal*). The orientation convention is the ordinary convention for oriented one-forms and current one-forms. The labels  $a(t)dt$  and regular currents  $B = b(t)dt$  transform by pullback,  $a(\psi(s))\psi'(s)ds$  and  $b(\psi(s))\psi'(s)ds$ . Additional signs are governed only by the written one-form:  $u_{a(t)dt \otimes f}$ ,  $c_\lambda$ ,  $\theta_\lambda$ , or the principal-part label  $\rho$ . If the written one-form is held fixed while only the chosen orientation is reversed, the sign is exactly the sign of  $dt$ .

**Remark B.13** (*Boundary of the algebraic current theorem*). Theorem B.7 proves the algebraic distributional-current interface. The finite polynomial BV observable, primitive/indecomposable projection before reduction, full de Rham-current BV differential, mixed brane-defect QME, and Costello renormalized graph weights require the unreduced current complex, scalar cancellation data, Tate/weighted coefficient kernel, and brane-defect propagator specified in Appendix E and Problem 5.212.

The remainder of the appendix prepares the unreduced lift. The algebraic interface above is degree-zero current data on the freely adjoined principal-part current sector; the genuine target is a weighted Costello local-functional complex with a finite-scale BV Laplacian, a finite-scale propagator, and a QME differential  $d_{\text{QME}} = Q + \{I_o^w, -\}_\hbar$  of degree  $+1$ . The remaining conventions split into four habitats: the unreduced source and target (Definition B.14), the bulk-to-defect convolution Hom complex (Definition B.15) in which Maurer–Cartan / curvature lives, the coefficient-current kernel layer (Proposition B.16) that solves the algebraic problem strictly, and the Costello graph layers (Propositions B.17–B.20) that supply the analytic extension under wavefront/scaling-degree hypotheses. The obstruction map list and centrality homotopy criterion (Proposition B.21, Theorem B.22) then record exactly which classes block strengthening from the algebraic kernel to a Costello graph lift.

*Definition B.14 (Unreduced non-scalar current lift datum).* Fix an oriented interval  $I \subset \mathbb{R}_{\text{brane}} \subset \mathbb{R}_{\text{top}}^2$ , an admissible weight  $w$  on the Hamiltonian degree of  $\widehat{\mathbb{C}}^2_{o,\text{hol}}$ , and the Moyal parameter  $\hbar$ . A non-scalar Costello current lift of the reduced interface for the local mixed HT model consists of the following data.

- (i) A source current complex

$$\mathfrak{g}_{\hbar,I}^{w,\text{cur}} = (\Omega_c^\bullet(I) \widehat{\otimes}_{\mathfrak{h}_{w,\hbar}}) \ltimes (\mathcal{D}'_c(I) \widehat{\otimes}_{\mathfrak{h}_{\hbar,\text{cont}}}^{\vee,w})[1],$$

or, before scalar reduction, the corresponding central-character replacement of Definition E.8.

- (ii) A target defect local-functional complex

$$\mathfrak{Q}_{w,\partial,\hbar}^\bullet(I) = \left( \mathcal{O}_{\text{loc}}^{\text{bd}}(\mathcal{E}_w|_I; A_{\partial,\hbar}^{\text{pp},w}(I)), d_{\text{QME}} = Q + \{I_o^w, -\}_\hbar \right),$$

together with the finite-scale BV Laplacian and propagator in the formulation where the Costello scale- $L$  QME remains explicit. Here  $\mathcal{E}_w$  is the weighted local mixed HT field complex on  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}^2_{o,\text{hol}}$ , restricted to the brane-defect interval  $I$ ; the coefficient algebra is the weighted Hamiltonian/Moyal algebra of the same formal holomorphic symplectic polydisk.

- (iii) A continuous bulk-to-defect kernel, equivalently a Maurer–Cartan candidate in the completed convolution Lie algebra,

$$\kappa_{\hbar,w,I} : C_{\text{CE,cont}}^\bullet(\mathfrak{g}_{\hbar,I}^{w,\text{cur}}) \longrightarrow \mathfrak{Q}_{w,\partial,\hbar}^\bullet(I),$$

whose generator values are representatives

$$\mathcal{U}_{a(t)dt \otimes f} \longmapsto B_f^{\text{BV}}(a), \quad c_{B,\rho} \longmapsto \Theta_\rho^{\text{BV}}(B).$$

- (iv) A  $d_{\text{QME}}$ -stable scalar-contact filtration and a filtered chain projection

$$\widehat{\sigma}_{\text{sc}} : \mathfrak{Q}_{w,\partial,\hbar}^\bullet(I) \longrightarrow C_{\text{Lie}}^\bullet(\tilde{A}; \mathbb{C}\gamma_1)[1][[\hbar]]$$

whose associated graded is the scalar-symbol map of Definition E.3.

- (v) Product centrality homotopies and  $P_o$ -bracket homotopies against arbitrary unreduced open observables, extending the checks on the freely adjoined current generators of the reduced target.

The reduced algebraic theorem supplies only the image of this datum after scalar reduction, de Rham-current contraction, and passage to the added principal-part current target.

*Definition B.15 (Bulk-to-defect convolution Hom complex).* With  $I, w, \hbar$  as in Definition B.14, and with all Hamiltonian coefficients taken from the local formal holomorphic polydisk of Proposition B.3, put

$$\mathbf{K}_{w,\partial,\hbar}^\bullet(I) = \text{Hom}_{\text{cont}}\left(C_{\text{CE},\text{cont}}^\bullet(\mathfrak{g}_{\hbar,I}^{w,\text{cur}}), \mathfrak{Q}_{w,\partial,\hbar}^\bullet(I)\right).$$

The topology is the complete filtration topology generated by the  $\hbar$ -adic filtration, the scalar-contact filtration, and the finite-window Mittag–Leffler filtration. Thus a sequence of kernels converges exactly when all of its finite-window, scalar-contact, and  $\hbar$ -adic projections converge. The differential and bracket are the completed convolution operations

$$d_{\mathbf{K}}T = d_{\text{QME}}T - (-1)^{|T|}T d_{\text{CE}}, \quad [T_1, T_2]_{\mathbf{K}} = [-, -]_{\hbar} \circ (T_1 \widehat{\otimes} T_2) \circ \Delta_{\text{CE}},$$

where  $\Delta_{\text{CE}}$  is the completed CE coproduct and  $[-, -]_{\hbar}$  is the target  $P_{0,\hbar}/\text{BV}$  bracket. For a degree-zero kernel candidate  $\kappa$ , its curvature is

$$\text{Curv}_{\mathbf{K}}(\kappa) = d_{\mathbf{K}}\kappa + \frac{1}{2}[\kappa, \kappa]_{\mathbf{K}} \in \mathbf{K}_{w,\partial,\hbar}^1(I).$$

On generators, the term  $-\kappa d_{\text{CE}}$  is exactly the source-bracket subtraction appearing in the bracket-defect formulas below.

**PROPOSITION B.16 (Bulk-to-defect coefficient-current kernel layer).** In the local mixed HT model on  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}_{0,\text{hol}}^2$ , there is a canonical degree-zero continuous coefficient-current kernel

$$\kappa_{\hbar,w,I}^{\text{coef}} \in \mathbf{K}_{w,\partial,\hbar}^0(I)$$

whose generator values are

$$\kappa_{\hbar,w,I}^{\text{coef}}(u_{a(t)dt} \otimes f) = \iota_I(B_f^w(a)), \quad \kappa_{\hbar,w,I}^{\text{coef}}(c_{B,\rho}) = \iota_I(\Theta_\rho^w(B)).$$

Here  $a \in C_c^\infty(I)$ ,  $f \in \mathfrak{h}_{w,\hbar}$ ,  $B \in \mathcal{D}'(I)$ , and  $\rho \in \mathfrak{h}_{\hbar,\text{cont}}^{\vee,w}$ . The map  $\iota_I$  is the coefficient inclusion of  $A_{\partial,\hbar}^{\text{pp},w}(I)$  as field-independent defect-local functionals in  $\mathfrak{Q}_{w,\partial,\hbar}^\bullet(I)$ . The degrees are

$$|u_{a(t)dt} \otimes f| = |B_f^w(a)| = 0, \quad |c_{B,\rho}| = |\Theta_\rho^w(B)| = 1.$$

The kernel has the following proved properties.

- (i) For each finite Hamiltonian window  $M$  in the formal holomorphic Hamiltonian degree, its restriction  $\kappa_{\hbar,w,I}^{\text{coef},M}$  lands in

$$\mathcal{O}_{\text{loc}}^{\text{bd}}(\mathcal{E}_w^M|_I; A_{\partial,\hbar}^{\text{pp},w,M}(I))$$

and the finite-window maps satisfy

$$\pi_{M,N} \kappa_{\hbar,w,I}^{\text{coef},M} = \kappa_{\hbar,w,I}^{\text{coef},N} \pi_{M,N}^{\text{CE}}, \quad M \geq N.$$

For admissible weights  $w, w'$ , the diagonal transport maps satisfy

$$R_{w,w'}^M \kappa_{\hbar,w,I}^{\text{coef},M} = \kappa_{\hbar,w',I}^{\text{coef},M} R_{w,w'}^{M,\text{CE}}.$$

Hence the displayed finite-window kernels determine the unique strict Mittag–Leffler inverse-limit kernel  $\kappa_{\hbar,w,I}^{\text{coef}}$ .

- (ii) After setting  $\hbar = 0$ , applying scalar reduction and the de Rham-current contraction, the kernel is exactly  $\Phi_I^{\text{cur}}$  of Theorem B.7.
- (iii) Inside the freely adjoined coefficient-current subalgebra, the generator  $P_{0,\hbar}$ -defects vanish:

$$\{B_f^w(a), B_g^w(b)\}_{\hbar} = B_{\{f,g\}_{\hbar}}^w(ab),$$

$$\{B_f^w(a), \Theta_{\rho}^w(B)\}_{\hbar} = \Theta_{\text{ad}_{f,\hbar}^{\vee} \rho}^w(aB), \quad \{\Theta_{\rho}^w(B), \Theta_{\sigma}^w(C)\}_{\hbar} = 0.$$

- (iv) Let  $\kappa_{\hbar,w,I}$  be any Costello-local-functional lift of this coefficient kernel. If its curved Maurer–Cartan equation has been solved through order  $\hbar^{n-1}$ , then the order- $n$  component of  $\text{Curv}_{\mathbf{K}}(\kappa_{\hbar,w,I})$  is  $d_{\text{QME}}$ -closed modulo the already solved lower-order terms. If the filtered scalar projection exists and its scalar image is zero, each fixed source-arity or graph component gives, by evaluation, a residual class in

$$H^1(\ker \widehat{\tau}_{\text{sc}}, d_{\text{QME}}).$$

Equivalently, the assembled curvature coefficient is a class in the corresponding Hom-valued obstruction complex; this is the class denoted  $[\text{Ob}_{w,\partial,\hbar}^{\text{red}}]$  in Problem 5.219. Its compatible finite-window counterterm obstruction is precisely the inverse-limit  $H^1$  class together with the  $\varprojlim H^0$  primitive-compatibility class of Theorem 5.83.

This proposition constructs the local coefficient-current kernel layer. A Costello graph kernel for arbitrary  $B \in \mathcal{D}'_c(I)$  requires the following analytic theorem: for every finite window and every defect-current-labelled graph, the products of the finite-scale propagators with the pushed-forward compactly supported currents must satisfy Hörmander wavefront transversality, admit local extensions across the relevant small diagonals, and produce counterterms compatible with interval extension, finite-window truncation, and admissible weight transport. The graph-level lift proved below applies after restricting the cotangent labels to regular compactly supported densities or to a wavefront-admissible class; the arbitrary distributional case is the obstruction problem.

*Proof.* By Proposition B.3, the source Lie brackets and principal-part pairings are the formal-disk Hamiltonian brackets and residues tensored with interval currents. The continuous CE cochain algebra is the completed graded-symmetric algebra on the continuous dual of  $\mathfrak{g}_{\hbar,I}^{w,\text{cur}}[1]$ . The target coefficient algebra  $A_{\partial,\hbar}^{\text{pp},w}(I)$  is the completed graded-symmetric algebra on the matching Hamiltonian-current generators  $B_f^w(a)$  and cotangent-current generators  $\Theta_{\rho}^w(B)$ . The displayed assignment therefore extends uniquely to a continuous completed graded-commutative algebra map. Since the source and target generators have the displayed equal degrees, the resulting element of the convolution Hom complex has degree 0.

Windowwise, the same construction is finite-dimensional in the coefficient direction, with  $D_{w,\leq M}$  and  $A_{\partial,\hbar}^{\text{pp},w,M}(I)$  as in Definition 5.82. Truncation merely discards coefficient degrees  $d > N$ , while  $R_{w,w'}^M$  rescales each surviving degree- $d$  basis vector and its dual by inverse factors. The formulas for  $B_f^w(a)$  and  $\Theta_{\rho}^w(B)$  are degreewise identities, so they commute with truncation and with weight transport. The strict Mittag–Leffler universal property gives the inverse-limit kernel.

The reduction statement follows by forgetting the Moyal deformation, applying the scalar quotient, and contracting the de Rham current direction. The two generator formulas become  $u_{a(t)dt \otimes f} \mapsto B_f(a)$  and  $c_{B,\rho} \mapsto \Theta_{\rho}(B)$ , which are precisely the generators of Theorem B.7.

The three displayed bracket identities are the weighted Moyal version of the semidirect-product calculation in Theorem B.7: the Hamiltonian-current bracket gives the pointwise product  $ab$ , the

cotangent action is multiplication  $B \mapsto aB$  together with the Moyal coadjoint action, and the shifted cotangent current ideal is abelian. By biderivation, the identities extend from generators to the completed coefficient-current subalgebra.

Finally, the convolution Hom complex is a filtered dg Lie algebra. The Bianchi identity

$$d_{\mathbf{K}} \text{Curv}_{\mathbf{K}}(\kappa) + [\kappa, \text{Curv}_{\mathbf{K}}(\kappa)]_{\mathbf{K}} = 0$$

implies, order by order in  $\hbar$ , that the first unsolved curvature coefficient is closed for the linearized differential. Under the chosen scalar projection, zero scalar image places every evaluated component in  $\ker \widehat{\sigma}_{\text{sc}}$ , equivalently places the assembled coefficient in the Hom-valued kernel. Adding a degree-zero local counterterm changes the first unsolved coefficient by the corresponding  $d_{\text{QME}}$ -boundary, and the finite-window compatibility of such primitives is exactly the Milnor obstruction measured in Theorem 5.83. This proves the stated obstruction class and leaves only the analytic distributional graph-extension theorem named in the statement.  $\square$

The Costello graph layer is split by the analytic admissibility of the external cotangent labels. Three habitats are constructed, in strict order of increasing label generality:

- *Regular densities*  $\mathcal{D}_c^{\text{reg}}(I)$  (Proposition B.17), where the distributional operation is multiplication of smooth heat-kernel propagators by smooth densities and the wavefront obstruction vanishes;
- *Wavefront-admissible distributional tuples*  $\mathcal{D}'_{c, \text{WF}}(I; \Gamma, M)$  (Definition B.18, Proposition B.19), where the Hörmander product, pushforward, and finite-scaling extension calculus [36, Ch. 8], [12, Chs. 5–7] are defined, including the one-sided sign-conic finite-scaling classes  $\mathcal{D}'_{c, \Lambda_{\pm}, \text{fs}}(I)$ ;
- *Arbitrary compactly supported distributions*  $\mathcal{D}'_c(I)$  (Proposition B.20), where the algebraic kernel is the coefficient-current kernel of Proposition B.16, and the coincident-current boundary graph  $B_1 = B_2 = \delta_{t_0}$  gives an explicit Hörmander product obstruction for general  $\mathcal{D}'_c(I)$ .

The first two habitats give renormalized graph kernels compatible with interval extension, finite-window truncation, and admissible weight transport; the third names the residual obstruction problem.

PROPOSITION B.17 (*Regular-density Costello graph kernel layer*). Let

$$\mathcal{D}_c^{\text{reg}}(I) = \{ b(t)dt \mid b \in C_c^\infty(I) \} \subset \mathcal{D}'_c(I)$$

and let

$$\mathfrak{g}_{h,I}^{w, \text{reg}} = (\Omega_c^\bullet(I) \widehat{\otimes}_{w,h}) \ltimes (\mathcal{D}_c^{\text{reg}}(I) \widehat{\otimes}_{h, \text{cont}}^{\vee, w})[1] \subset \mathfrak{g}_{h,I}^{w, \text{cur}}.$$

Put  $\mathbf{K}_{w, \partial, h}^{\bullet, \text{reg}}(I)$  for the corresponding completed convolution Hom obtained from Definition B.15 by restricting the source to  $C_{\text{CE}, \text{cont}}^\bullet(\mathfrak{g}_{h,I}^{w, \text{reg}})$ . Assume that, on every finite Hamiltonian window  $M$  of the local formal holomorphic Hamiltonian sector, the target  $\mathfrak{Q}_{w, \partial, h}^\bullet(I)$  is equipped with the regulator-admissible finite-scale Costello brane-defect graph calculus for  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}_{\text{o, hol}}^2$ : the weighted propagator  $P_{\epsilon, L}^{w, M}$ , regularized BV Laplacian, local interaction  $I_{\text{o}, M}^w$ , interval extension by zero, truncation maps  $\pi_{M, N}$ , and admissible weight transports  $R_{w, w'}^M$ , satisfying the weighted RG criterion of Statement 5.78 and the graphwise Hadamard/Mittag–Leffler control of Theorem 8.17.

Then the restriction of  $\kappa_{h, w, I}^{\text{coef}}$  to regular density labels has a graphwise Costello-local lift

$$\kappa_{h, w, I}^{\text{graph, reg}} \in \mathbf{K}_{w, \partial, h}^{\text{o, reg}}(I)$$

in the following precise sense. For every fixed connected Costello graph  $\Gamma$ , finite CE input, finite window  $M$ , and external cotangent labels  $B_j = b_j(t)dt \in \mathcal{D}_c^{\text{reg}}(I)$ , the finite-scale graph contribution

$$W_{\Gamma, \epsilon, L}^{\text{reg}, M}(a_\bullet, b_\bullet; f_\bullet, \rho_\bullet) \in \mathcal{O}_{\text{loc}}^{\text{bd}}(\mathcal{E}_w^M|_I; A_{\partial, \hbar}^{\text{pp}, w, M}(I))$$

is a well-defined smooth local-functional kernel for  $0 < \epsilon \leq L < \infty$ . Its small- $\epsilon$  expansion admits Costello local counterterms  $C_{\Gamma, w}^{\text{reg}, M}(\epsilon; b_\bullet)$ , and the renormalized graph

$$W_{\Gamma, L, w}^{R, \text{reg}, M} = \lim_{\epsilon \rightarrow 0} (W_{\Gamma, \epsilon, L}^{\text{reg}, M} - C_{\Gamma, w}^{\text{reg}, M}(\epsilon; b_\bullet))$$

is again a brane-defect local functional with regular density coefficient labels. These graph kernels are compatible with interval extension by zero, finite-window truncation, and admissible weight transport:

$$\pi_{M, N} W_{\Gamma, L, w}^{R, \text{reg}, M} = W_{\Gamma, L, w}^{R, \text{reg}, N} \pi_{M, N}^{\text{CE}}, \quad R_{w, w'}^M W_{\Gamma, L, w}^{R, \text{reg}, M} = W_{\Gamma, L, w'}^{R, \text{reg}, M} R_{w, w'}^{M, \text{CE}}.$$

On the zero-internal-edge part, and after scalar reduction and de Rham-current contraction, the lifted kernel is  $\Phi_I^{\text{cur}}$ .

Consequently, wherever a filtered scalar projection  $\widehat{\sigma}_{\text{sc}}$  has been constructed on the chosen regular-density graph/counterterm subhabitat, the curvature

$$\text{Curv}_{\mathbf{K}}(\kappa_{\hbar, w, I}^{\text{graph}, \text{reg}})$$

defines the same finite-window non-scalar obstruction system as in Theorem 5.83. Thus the regular-density theorem removes the analytic wavefront/counterterm gap attached to the cotangent labels. The filtered scalar projection and vanishing of the residual class

$$[\text{Ob}_{w, \partial, \hbar}^{\text{red}}] \in H^1(\ker \widehat{\sigma}_{\text{sc}}, d_{\text{QME}}).$$

are the remaining non-scalar QME data.

Extension from  $\mathcal{D}_c^{\text{reg}}(I)$  to  $\mathcal{D}'_c(I)$  is the remaining analytic theorem: for every finite window  $M$ , every Costello graph  $\Gamma$ , and every compactly supported distributional label  $B_j \in \mathcal{D}'_c(I)$ , the product of the finite-scale brane-defect propagator kernels with the pushed-forward current labels must satisfy Hörmander wavefront transversality, admit local extensions across all graph small diagonals, and admit counterterms compatible with interval extension,  $\pi_{M, N}$ , and  $R_{w, w'}^M$ . This is a local brane-defect microlocal condition on the displayed finite-scale kernels.

*Proof.* The regular subspace is stable under the current operations used in the source. Indeed, if  $B = b(t)dt$  and  $a \in C_c^\infty(I)$ , then Lemma B.4 gives  $aB = (ab)(t)dt$ , again a compactly supported smooth density. Extension by zero along  $I \hookrightarrow J$  is smooth because the support is compactly contained in  $I$ . Thus the regular labels define the stated source subcomplex and the restriction of the convolution Hom complex.

Fix a finite window  $M$ , a graph  $\Gamma$ , and  $0 < \epsilon \leq L < \infty$ . In this window the coefficient part of the propagator is finite rank, and the spacetime part of  $P_{\epsilon, L}^{w, M}$  is the finite-scale heat-kernel propagator of Costello's elliptic BV calculus [12, Chs. 2–4]. It is smooth at positive scale. After restriction to the brane line, multiplication by the smooth compactly supported functions  $a_i$  and densities  $b_j(t)dt$  produces a smooth compactly supported graph integrand on the relevant configuration space and edge-parameter

domain. Thus every product is a product of smooth densities at finite scale, and the graph integral is a well-defined element of the displayed brane-defect local-functional target.

As  $\epsilon \rightarrow 0$ , the singularities are the usual heat-kernel singularities along graph diagonals. Multiplication by smooth compact densities preserves wavefront sets and the Hadamard singular order. Therefore the local asymptotic expansion and local counterterm construction are exactly the windowwise Costello construction, tensored with the finite coefficient kernel and multiplied by smooth density coefficients. Theorem 8.17 supplies the graphwise Hadamard estimates and strict Mittag–Leffler passage in the weighted coefficient direction. The counterterms remain local on the brane line; their coefficients are differential expressions in smooth compactly supported labels, hence regular density labels.

The compatibility identities are degree-wise. The maps  $\pi_{M,N}$  discard coefficient degrees above  $N$ , while  $R_{w,w'}^M$  rescales the surviving weighted Hamiltonian and cotangent bases. The regular density factors live in the interval variable and are untouched by these coefficient maps. The strict Mittag–Leffler compatibility of the finite-window propagators and counterterms in Statement 5.78 therefore gives the two displayed identities. Interval extension compatibility is the same locality statement: all labels and counterterms are extended by zero, and local graph weights are supported on the same brane diagonals after extension.

The zero-internal-edge graph is the generator assignment of Proposition B.16. After forgetting the Moyal deformation, applying scalar reduction, and contracting the de Rham-current direction, it is Theorem B.7.

The curvature assertion is formal in the filtered convolution dg Lie algebra. The Bianchi identity used in the proof of Proposition B.16 applies to the regular restricted complex. Once a filtered scalar projection is available, the first unsolved curvature coefficient lands in  $\ker \widehat{\partial}_{\text{sc}}$ , and compatible counterterms change it by  $d_{\text{QME}}$ -boundaries. The finite-window obstruction is exactly the inverse-limit  $H^1$  class and the  $\varprojlim H^0$  primitive-compatibility class of Theorem 5.83.

If  $B$  is an arbitrary compactly supported distribution, the graph-level construction requires Hörmander transversality for the product of the pushed-forward brane current with propagator boundary values and a compatible extension across graph diagonals. This is exactly the residual theorem stated above.  $\square$

*Definition B.18 (Wavefront-admissible defect-current labels).* Fix a finite Hamiltonian window  $M$ , a connected Costello graph  $\Gamma$ , and a finite list of external cotangent-current labels  $B_\bullet = (B_1, \dots, B_r)$ ,  $B_j \in \mathcal{D}'_c(I)$ . Let  $X = \mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}_{\text{o,hol}}^2$  and  $L_I = I \times \{0\} \times \{p\} \subset X$ . If  $V_{\text{bulk}}(\Gamma)$  and  $V_\partial(\Gamma)$  are the bulk and defect vertices, set

$$X_\Gamma^\circ(I) = X^{V_{\text{bulk}}(\Gamma)} \times L_I^{V_\partial(\Gamma)} \times E_{\Gamma,M},$$

where  $E_{\Gamma,M}$  is the finite-dimensional coefficient and edge parameter space in the window  $M$ . The final graph space  $X_\Gamma(I)$  is the open complement of the graph collision diagonals in  $X_\Gamma^\circ(I)$  before local extension across those diagonals. Choose an attachment map  $\ell : \{1, \dots, r\} \rightarrow V_\partial(\Gamma)$  for the current labels and define the incidence map

$$e_\Gamma : X_\Gamma^\circ(I) \longrightarrow I^r, \quad e_{\Gamma,j}(x) = t_{\ell(j)}.$$

Repeated labels at one defect vertex are therefore diagonal pullbacks by  $e_\Gamma$ . For  $B^\boxtimes = B_1 \boxtimes \dots \boxtimes B_r$ , write  $B_\Gamma = e_\Gamma^* B^\boxtimes$  only when this pullback is defined. Let  $\text{Diag}(\Gamma)$  be the finite set of graph collision diagonals on which the small- $\epsilon$  Hadamard coefficients may be singular.

The tuple  $B_\bullet$  is called  $(\Gamma, M)$ -wavefront-admissible if the following four operations are defined by the usual Hörmander calculus [36, Ch. 8] and by Costello’s finite-window graph calculus [12, Chs. 5–7].

- (i) The incidence pullback is non-characteristic:

$$N_{e_\Gamma} \cap \text{WF}(B^\boxtimes) = \emptyset, \quad N_{e_\Gamma} = \{(e_\Gamma(x), \xi) \mid de_{\Gamma,x}^t \xi = 0\}.$$

Then

$$\text{WF}(B_\Gamma) \subset e_\Gamma^\# \text{WF}(B^\boxtimes) = \{(x, de_{\Gamma,x}^t \xi) \mid (e_\Gamma(x), \xi) \in \text{WF}(B^\boxtimes)\}.$$

The resulting support is proper over the graph integration variables.

- (ii) For every Hadamard coefficient  $K_{\Gamma,\alpha}^M$  appearing in the small- $\epsilon$  expansion of the finite-window graph kernel,

$$(\text{WF}(K_{\Gamma,\alpha}^M) + \text{WF}(B_\Gamma)) \cap T_{X_\Gamma(I)}^* X_\Gamma(I) = \emptyset.$$

Equivalently, the wavefront sets are transverse to the zero section after summation. Thus the product  $K_{\Gamma,\alpha}^M B_\Gamma$  is a canonical distribution.

- (iii) For each collision diagonal  $\Delta \in \text{Diag}(\Gamma)$ , the products  $K_{\Gamma,\alpha}^M B_\Gamma$  have finite scaling degree along  $\Delta$ . Their allowed extensions across  $\Delta$  differ by distributions supported on  $\Delta \cap \text{supp}(B_\Gamma)$ , hence by brane-defect local counterterms with distributional current coefficients.
- (iv) The final graph pushforward has wavefront contained in the brane-defect local-functional wavefront class already used in Theorem 8.17, after tensoring with the finite coefficient kernel.

Denote the resulting multilinear relation by

$$\mathcal{D}'_{c,\text{WF}}(I; \Gamma, M) \subset (\mathcal{D}'_c(I))^r.$$

The universal wavefront-admissible relation is the intersection of these relations over the finite windows, graphs, and finite CE inputs under consideration. This is a relation on finite tuples. A linear subspace  $\mathcal{V} \subset \mathcal{D}'_c(I)$  gives an honest restricted CE source only when every finite tuple from  $\mathcal{V}$  lies in the corresponding relation.

A useful source of such subspaces is a diagonal-admissible conic set  $\Lambda \subset T^*I \setminus 0$ : for every  $n \geq 1$ ,

$$\xi_1, \dots, \xi_n \in \Lambda \implies \xi_1 + \dots + \xi_n \neq 0.$$

Then

$$\mathcal{D}'_{c,\Lambda,\text{fs}}(I) = \{B \in \mathcal{D}'_c(I) \mid \text{WF}(B) \subset \Lambda, B \text{ has finite scaling degree along its singular support}\}$$

is the finite-scaling one-sided wavefront class. On an oriented interval, either open half-cone  $\Lambda_+ = \{(t, \xi) \mid \xi > 0\}$  or  $\Lambda_- = \{(t, \xi) \mid \xi < 0\}$  is diagonal-admissible. These two half-cones are maximal among sign-conic diagonal-admissible linear habitats for ordinary collision-conormal kernels: any conic subset of  $T^*I \setminus 0$  containing both signs admits, after positive rescaling, a finite nonempty covector sum equal to zero, so it can cancel a collision conormal. Compactly supported one-sided conormal boundary values, for example cutoffs of  $(t - t_0 - i0)^{-m}$  after choosing the Fourier sign convention, lie in one of these distributional classes.

**PROPOSITION B.19** (*Wavefront-admissible distributional graph kernel layer*). Assume the same regulator-admissible finite-window Costello brane-defect graph calculus as in Proposition B.17. Let  $\Gamma$  be a fixed connected graph,  $M$  a finite Hamiltonian window, and  $B_\bullet \in \mathcal{D}'_{c,\text{WF}}(I; \Gamma, M)$ . Then the graph contribution with external cotangent labels  $B_\bullet$  has the following distributional extension.



- (i) For every  $0 < \epsilon \leq L < \infty$ , the finite-scale graph weight

$$W_{\Gamma, \epsilon, L}^{\text{WF}, M}(a_\bullet, B_\bullet; f_\bullet, \rho_\bullet)$$

is a well-defined brane-defect local-functional kernel with compactly supported distributional current coefficients. At this positive scale the propagator is smooth, so the only distributional operation is multiplication of smooth kernels by the pushed-forward currents.

- (ii) The small- $\epsilon$  expansion admits local counterterms

$$C_{\Gamma, w}^{\text{WF}, M}(\epsilon; B_\bullet)$$

supported on

$$\bigcup_{\Delta \in \text{Diag}(\Gamma)} \Delta \cap \text{supp}(B_\Gamma).$$

The renormalized graph

$$W_{\Gamma, L, w}^{R, \text{WF}, M} = \lim_{\epsilon \rightarrow 0} (W_{\Gamma, \epsilon, L}^{\text{WF}, M} - C_{\Gamma, w}^{\text{WF}, M}(\epsilon; B_\bullet))$$

is a brane-defect local functional with coefficients in the distributional current target  $\mathcal{A}_{\partial, h}^{\text{pp}, w, M}(I)$ .

- (iii) The construction is compatible with interval extension by zero, finite-window truncation, and admissible weight transport whenever the chosen tuple remains wavefront-admissible after applying the corresponding incidence maps:

$$\pi_{M, N} W_{\Gamma, L, w}^{R, \text{WF}, M} = W_{\Gamma, L, w}^{R, \text{WF}, N} \pi_{M, N}^{\text{CE}}, \quad R_{w, w'}^M W_{\Gamma, L, w}^{R, \text{WF}, M} = W_{\Gamma, L, w'}^{R, \text{WF}, M} R_{w, w'}^{M, \text{CE}}.$$

In particular, every finite tuple of labels in  $\mathcal{D}'_{c, \Lambda, \text{fs}}(I)$  for a diagonal-admissible cone  $\Lambda$  is wavefront-admissible for graph kernels whose singular wavefronts are conormal to ordinary graph collision diagonals. This gives a non-regular distributional extension of Proposition B.17.

This is the exact extension supported by Costello graph products: the relation  $\mathcal{D}'_{c, \text{WF}}(I; \Gamma, M)$  is the maximal subspace on which the multiplication, extension, and graph-pushforward operations all admit the cited wavefront calculus. The theorem extends to  $\mathcal{D}'_{c, \text{WF}}(I; \Gamma, M)$ ; the obstruction and the admitted extension are entirely local on the brane interval and on the finite-window formal polydisk coefficient sector.

*Proof.* At positive scale the heat-kernel propagator  $P_{\epsilon, L}^{w, M}$  is smooth in the spacetime and defect variables. Multiplication of a compactly supported distribution by a smooth function is always defined, and the incidence maps are proper on the supports by Definition B.18. This proves the finite-scale graph weight.

As  $\epsilon \rightarrow 0$ , Theorem 8.17 gives graphwise Hadamard coefficients with wavefront contained in the union of conormals to graph collision diagonals. The second condition in Definition B.18 is exactly Hörmander's product criterion [36, Ch. 8], so  $K_{\Gamma, \alpha}^M B_\Gamma$  is a canonical distribution for every coefficient. The finite-scaling condition then gives the standard local extension across each collision diagonal. Its ambiguity is supported on the collision locus meeting the support of  $B_\Gamma$ ; this is precisely a brane-defect local counterterm with distributional current coefficient. Subtracting these local counterterms gives the displayed renormalized graph.

The two compatibility identities are inherited degreewise from the finite-window Costello graph calculus. The maps  $\pi_{M, N}$  and  $R_{w, w'}^M$  act on the weighted Hamiltonian coefficient factors and preserve the interval wavefront cone. Interval extension by zero is allowed exactly when the extended incidence

tuple satisfies the same non-characteristic and proper-support conditions. Hence compatibility holds on the stated wavefront-admissible relation.

For the one-sided conic classes, let  $\Lambda$  be diagonal-admissible and let  $B_j \in \mathcal{D}'_{c,\Lambda,\text{fs}}(I)$ . A conormal covector to an  $n$ -fold collision diagonal has components  $(\eta_1, \dots, \eta_n)$  whose sum is zero on each collision block. If this covector were cancelled by covectors from the external labels, then a finite nonempty sum of elements of  $\Lambda$  would be zero, contradicting diagonal-admissibility. Thus the product criterion holds. The finite-scaling hypothesis gives the extension condition, and the remaining pushforward condition is the same proper-support condition as above.

Finally, the full compactly supported distribution space exceeds the graph-product habitat. In the one-edge boundary graph take two external labels  $B_1 = B_2 = \delta_{t_0}$ . The singular boundary propagator has wavefront containing the conormal directions  $(t_0, t_0; \tau, -\tau)$  to the diagonal  $t = s$ , while  $\text{WF}(\delta_{t_0} \boxtimes \delta_{t_0})$  contains the opposite directions  $(t_0, t_0; -\tau, \tau)$ . Hörmander's product criterion detects the obstruction. Equivalently, the graph asks for the value of the singular propagator on the diagonal against  $\delta_{t_0} \boxtimes \delta_{t_0}$ , which requires an additional renormalization rule. Thus current Costello graph products require a wavefront-admissible subspace of  $\mathcal{D}'_c(I)$ .  $\square$

**PROPOSITION B.20** (*Current-compatible  $\mathcal{D}'_c$  graph criterion*). Assume the regulator-admissible finite-window Costello brane-defect graph system of Proposition B.17. The construction compatible with compactly supported distributional cotangent labels is the following.

- (i) For arbitrary  $B \in \mathcal{D}'_c(I)$ , the constructed object is the coefficient-current kernel

$$\kappa_{h,w,I}^{\text{coef}}(u_{a(t)dt \otimes f}) = \iota_I(B_f^w(a)), \quad \kappa_{h,w,I}^{\text{coef}}(c_{B,\rho}) = \iota_I(\Theta_\rho^w(B)).$$

This is an algebraic current kernel. A Costello graph-kernel theorem for all  $B \in \mathcal{D}'_c(I)$  is the additional analytic target.

- (ii) For a fixed graph  $\Gamma$ , finite window  $M$ , and admissible tuple  $B_\bullet \in \mathcal{D}'_{c,\text{WF}}(I; \Gamma, M)$ , put

$$T_{\Gamma,\alpha}^M(B_\bullet) = K_{\Gamma,\alpha}^M B_\Gamma.$$

Its wavefront is bounded by the Hörmander product formula

$$\begin{aligned} \text{WF}(T_{\Gamma,\alpha}^M(B_\bullet)) &\subset \text{WF}(K_{\Gamma,\alpha}^M) \cup \text{WF}(B_\Gamma) \\ &\cup (\text{WF}(K_{\Gamma,\alpha}^M) + \text{WF}(B_\Gamma)), \end{aligned}$$

with the fibrewise sum taken over the same base point in  $X_\Gamma(I)$ . For the final graph projection  $p_\Gamma : X_\Gamma(I) \rightarrow Y_\Gamma(I)$ , proper on support, the pushed-forward graph has

$$\text{WF}(p_{\Gamma,*} T_{\Gamma,\alpha}^M) \subset \{(\gamma, \eta) \mid \exists x \in p_\Gamma^{-1}(\gamma), (x, dp_{\Gamma,x}^t \eta) \in \text{WF}(T_{\Gamma,\alpha}^M)\}.$$

These are the only wavefront bounds used by the distributional graph layer.

- (iii) Let  $\Delta \in \text{Diag}(\Gamma)$ , let  $c_\Delta = \text{codim } \Delta$ , and suppose

$$s_{\Gamma,\alpha,\Delta} = \text{sd}_\Delta(T_{\Gamma,\alpha}^M(B_\bullet)) < \infty.$$

Set

$$N_{\Gamma,\alpha,\Delta} = \max\{-1, \lfloor s_{\Gamma,\alpha,\Delta} - c_\Delta \rfloor\}.$$

If  $\tilde{T}$  and  $\tilde{T}'$  are two local extensions of  $T_{\Gamma,\alpha}^M(B_\bullet)$  across  $\Delta$ , then locally in normal coordinates to  $\Delta$

$$\tilde{T}' - \tilde{T} = \sum_{|\beta| \leq N_{\Gamma,\alpha,\Delta}} \partial_\nu^\beta \delta_\Delta \otimes S_{\Delta,\beta},$$

with the sum empty when  $N_{\Gamma,\alpha,\Delta} = -1$ . Here  $S_{\Delta,\beta} \in \mathcal{D}'(\Delta)$  is supported in  $\Delta \cap \text{supp}(B_\Gamma)$  and carries the same finite-window Hamiltonian coefficient label. Thus the allowed counterterm ambiguity belongs to

$$\sum_{\Delta,\alpha} \sum_{|\beta| \leq N_{\Gamma,\alpha,\Delta}} \partial_\nu^\beta \delta_\Delta \mathcal{D}'_{\Delta \cap \text{supp}(B_\Gamma)}(\Delta) \widehat{\otimes} A_{\partial,h}^{\text{pp},w,M}(I).$$

(iv) The finite-window counterterms are compatible precisely when the chosen extensions satisfy

$$j_* C_{\Gamma,I}^{\text{WF},M}(\epsilon; B_\bullet) = C_{\Gamma,J}^{\text{WF},M}(\epsilon; j_* B_\bullet),$$

for interval inclusions  $j : I \hookrightarrow J$  for which the extended tuple remains admissible, and

$$\pi_{M,N} C_{\Gamma,w}^{\text{WF},M}(\epsilon; B_\bullet) = C_{\Gamma,w}^{\text{WF},N}(\epsilon; B_\bullet) \pi_{M,N}^{\text{CE}},$$

$$R_{w,w'}^M C_{\Gamma,w}^{\text{WF},M}(\epsilon; B_\bullet) = C_{\Gamma,w'}^{\text{WF},M}(\epsilon; B_\bullet) R_{w,w'}^{M,\text{CE}}.$$

Defect of the interval identity is a factorization-compatibility residual. Defect of the last two identities is exactly the finite-window primitive compatibility obstruction measured by Theorem 5.83 after passing to the relevant graph/counterterm complex.

(v) The theorem for all of  $\mathcal{D}'_c(I)$  is an additional renormalization problem. In particular, the coincident-current boundary graph with  $B_1 = B_2 = \delta_{t_0}$  violates the product condition because conormal covectors  $(\tau, -\tau)$  to  $t = s$  can be cancelled by  $\text{WF}(\delta_{t_0} \boxtimes \delta_{t_0})$ . Thus the arbitrary- $\mathcal{D}'_c$  graph theorem requires extra renormalization data beyond Costello's general finite-scale construction and the Costello–Gwilliam factorization vocabulary.

Hence the graph construction is proved on  $\mathcal{D}_c^{\text{reg}}(I)$ , on finite graphwise wavefront-admissible tuples, and on the sign-conic finite-scaling habitats  $\mathcal{D}'_{c,\Lambda_{\pm},\text{fs}}(I)$  for ordinary collision-conormal graph kernels. The full distributional space  $\mathcal{D}'_c(I)$  remains an open obstruction problem.

*Proof.* Item (1) is Proposition B.16: the assignment is degreewise algebraic in the compactly supported current  $B$  and uses only multiplication of a current by a smooth test function.

Items (2) and (3) are the standard Hörmander product, pushforward, and finite-scaling extension calculus applied to Definition B.18. The product transversality condition makes  $K_{\Gamma,\alpha}^M B_\Gamma$  canonical. Properness of the graph projection gives the displayed pushforward wavefront bound. Finite scaling degree along  $\Delta$  gives extensions across  $\Delta$ ; the structure theorem for distributions supported on a submanifold gives the finite normal-derivative formula for the difference of two extensions. The support statement follows because all extensions agree off  $\Delta \cap \text{supp}(B_\Gamma)$ .

Item (4) is the compatibility condition needed for the counterterms to define one interval-local element of the completed finite-window and weight-transport tower. When the identities hold, the Bianchi argument of Proposition B.16 produces the Hom-valued obstruction class and the Milnor  $\varprojlim^1 H^0$  primitive obstruction. Interval-extension defects are factorization-current residuals; finite-window or weight defects are primitive-compatibility obstructions.

Item (5) is the counterexample in the proof of Proposition B.19. It also matches the Costello graph formalism: the local sources support finite-scale BV/RG/QME graph calculus, local functionals, and local counterterms under their hypotheses. The current-valued arbitrary- $\mathcal{D}'_c$  brane-defect theorem requires the extra renormalization rule above.  $\square$

The remaining two statements list the obstruction classes that block strengthening from the algebraic coefficient-current kernel (Proposition B.16) to a Costello local-functional graph theorem. Four classes appear: the scalar projection of the curvature; the recovery condition under scalar reduction; the three unreduced bracket defects  $D_{f,g}$ ,  $D_{f,\rho}$ ,  $D_{\rho,\sigma}$  of degree zero with their factorization / finite-window-compatible homotopies; and the residual non-scalar curvature class in  $H^1(\ker \widehat{\sigma}_{\text{sc}}, d_{\text{QME}})$ . The binary centrality criterion of Theorem B.22 records the precise homotopy equation (29) that a non-scalar Costello local lift must satisfy on every pair of source generators, with the inverse-system and Milnor primitive classes that obstruct compatible window-by-window assembly.

**PROPOSITION B.21** (*Obstruction maps for the non-scalar current lift*). The datum of Definition B.14 can realize the non-scalar Costello graph/QME theorem for the local mixed HT brane-defect model only if the following obstruction maps vanish, with the stated homotopies.

- (i) The scalar projection of the curvature must vanish in the chosen scalar branch:

$$\widehat{\sigma}_{\text{sc}}(\text{Curv}_{\mathbf{K}}(\kappa_{h,w,I})) = 0.$$

In the ordinary scalar-reduced  $\mathfrak{gl}_N$  branch this condition is exactly the requirement that the Capelli contact class  $\hbar N[\bar{c}]$  be cancelled by a closed-exchange term and a local counterterm. In the balanced supertrace branch it is the condition  $\widehat{\sigma}_{\text{sc}}(\text{Ob}_{w,\partial,h}) = 0$ .

- (ii) After scalar reduction and de Rham-current contraction, the kernel must recover the reduced current interface:

$$\bar{\kappa}_{h,w,I}(u_{a(t)} dt \otimes f) = B_f(a), \quad \bar{\kappa}_{h,w,I}(c_{B,\rho}) = \Theta_\rho(B),$$

with brackets

$$\{B_f(a), B_g(b)\}_{\hbar} = B_{\{f,g\}_{\hbar}}(ab), \quad \{B_f(a), \Theta_\rho(B)\}_{\hbar} = \Theta_{\text{ad}_{f,h}^\vee \rho}(aB),$$

and  $\{\Theta_\rho(B), \Theta_\sigma(C)\}_{\hbar} = 0$ . At  $\hbar = 0$  this is Theorem B.7.

- (iii) The unreduced bracket defects

$$D_{f,g}(a, b) = \{B_f^{\text{BV}}(a), B_g^{\text{BV}}(b)\}_{\hbar} - B_{\{f,g\}_{\hbar}}^{\text{BV}}(ab),$$

$$D_{f,\rho}(a, B) = \{B_f^{\text{BV}}(a), \Theta_\rho^{\text{BV}}(B)\}_{\hbar} - \Theta_{\text{ad}_{f,h}^\vee \rho}^{\text{BV}}(aB),$$

and

$$D_{\rho,\sigma}(B, C) = \{\Theta_\rho^{\text{BV}}(B), \Theta_\sigma^{\text{BV}}(C)\}_{\hbar}$$

must be  $d_{\text{QME}}$ -exact by homotopies compatible with inclusions of intervals, disjoint factorization products, and the finite-window regulator maps.

- (iv) After evaluation on each fixed finite CE input  $\xi$ , equivalently on each fixed graph/source-arity component, the residual non-scalar curvature class

$$[\text{Curv}_{\mathbf{K}}(\kappa_{h,w,I})(\xi)] \in H^1(\ker \widehat{\sigma}_{\text{sc}}, d_{\text{QME}})$$

must vanish after the compatible finite-window counterterms  $C_{\Gamma,w}^{w_o}(\epsilon)$  have been inserted.

These conditions are necessary. They are also sufficient for the stated local graph/QME realization once the Costello finite-scale BV propagator, BV Laplacian, locality estimates, and counterterm system in the chosen coefficient category have been constructed.

*Proof.* The source and target in Definition B.14 are the current versions of the weighted cotangent Hamiltonian source and the mixed brane-defect obstruction complex of Problem 5.219. A Costello graph realization is a solution of the QME in this local-functional complex, hence its curvature has zero scalar projection and zero residual class in  $\ker \widehat{\sigma}_{\text{sc}}$ . This gives the first and fourth obstruction maps.

The second condition follows from compatibility with the already proved reduced current theorem. After contraction and scalar reduction the only allowed target brackets are the Moyal principal-part current brackets of Theorem 5.217; at  $\hbar = 0$  these reduce to the three brackets of Theorem B.7. Any different reduction would change the theorem already proved.

The three defects in the third condition are the generator-level differences between the source semidirect Lie brackets and the target  $P_o$ -brackets. Since the CE differential and the target bracket are derivations, exactness of these generator defects, with homotopies compatible with factorization products and finite-window transition maps, is precisely the condition that the generator assignment extend to the desired homotopy  $P_o$ -central operation. The reduced algebraic target has zero defects by Theorem B.7; the unreduced polynomial BV target has the obstruction of Theorem E.6. Therefore the missing theorem must construct these homotopies in a larger current-compatible Costello local-functional category.  $\square$

**THEOREM B.22** (*Curved kernel centrality homotopy criterion*). Fix the local mixed HT brane-defect model on  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}_{o,\text{hol}}$ , a weighted current source  $\mathfrak{g}_{h,I}^{w,\text{cur}}$ , and a Costello local-functional target  $\mathfrak{D}_{w,\partial,h}^\bullet(I)$  as in Definition B.14. Let  $\kappa_M$  be a finite-window component of a degree-zero continuous kernel in the convolution Hom complex  $\mathbf{K}_{w,\partial,h}^\bullet(I)$ , and assume that the chosen scalar-contact projection is a chain map on the complex under discussion.

For source generators

$$x \in \{u_{a(t)dt \otimes f}, c_{B,\rho}\}, \quad y \in \{u_{b(t)dt \otimes g}, c_{C,\sigma}\},$$

define the binary closed-open curvature component by

$$\mathfrak{D}_\kappa^M(x, y) = \text{Curv}_{\mathbf{K},M}(\kappa_M)(x \wedge y).$$

Before inserting a binary Taylor component, these are the three generator defects

$$D_{f,g}^M(a, b), \quad D_{f,\rho}^M(a, B), \quad D_{\rho,\sigma}^M(B, C)$$

displayed in Proposition B.21.

A compatible unreduced binary centrality homotopy for this kernel is exactly a family of cochains

$$H_{x,y}^M \in \mathcal{K}_{\text{ns},M}^{|x|+|y|-1}(I) = \ker \widehat{\sigma}_{\text{sc},M} \cap \mathfrak{D}_{w,M,\partial,h}^{|x|+|y|-1}(I)$$

satisfying

$$\mathfrak{D}_\kappa^M(x, y) + d_M H_{x,y}^M = 0 \quad (29)$$

for every such pair, together with the compatibility identities

$$j_* H_{I;x,y}^M = H_{J;j_*x, j_*y}^M,$$

for interval inclusions  $j : I \hookrightarrow J$  for which the labels remain admissible, and

$$\pi_{M,N} H_{x,y}^M = H_{\pi x, \pi y}^N, \quad R_{w,w'}^M H_{w;x,y}^M = H_{w';Rx, Ry}^M.$$

Equivalently, for every pair  $(x, y)$ , the inverse-system cohomology class

$$([\mathfrak{D}_\kappa^M(x, y)])_M \in \varprojlim_M H^{|x|+|y|}(\mathcal{K}_{\text{ns}, M}^\bullet(I), d_M)$$

and the Milnor primitive-compatibility class represented by

$$\left[ \pi_{M,N} h_{x,y}^M - h_{\pi x, \pi y}^N \right] \in H^{|x|+|y|-1}(\mathcal{K}_{\text{ns}, N}^\bullet(I), d_N)$$

must both vanish, where  $h_{x,y}^M$  is any windowwise primitive solving  $\mathfrak{D}_\kappa^M(x, y) + d_M h_{x,y}^M = 0$ .

For the algebraic coefficient-current kernel  $\kappa_{h,w,I}^{\text{coef}}$  of Proposition B.16, all  $\mathfrak{D}_\kappa^M(x, y)$  vanish in the freely adjoined current target, and  $H_{x,y}^M = 0$ . This is the strict homotopy of Theorem E.10. For a genuine unreduced Costello local-functional target, the same conclusion requires the finite-window graph row representing  $\mathfrak{D}_\kappa^M(x, y)$ , a scalar-zero condition for that row, and a local counterterm or binary Taylor cochain  $H_{x,y}^M$  satisfying equation (29) with the displayed interval, window, and weight compatibilities.

*Proof.* The arity-two part of the curved Maurer–Cartan equation in the convolution dg Lie algebra is precisely

$$\text{Curv}_{\mathbf{K}, M}(\kappa_M)(x \wedge y) = 0.$$

Expanding this identity before choosing a binary component gives the target bracket of the two generator images minus the image of the source semidirect bracket, with the source term supplied by  $-\kappa d_{\text{CE}}$ . These are exactly the three defects  $D_{f,g}$ ,  $D_{f,\rho}$ , and  $D_{\rho,\sigma}$  of Proposition B.21.

A binary homotopy is, by definition, a degree  $|x| + |y| - 1$  primitive for this curvature component in the scalar-zero kernel complex. This gives equation (29). Compatibility with extension by zero, finite-window truncation, and admissible weight transport is exactly the statement that these primitives assemble to a homotopy in the completed local factorization target. The inverse-limit obstruction and the Milnor class are the usual Roos obstruction for choosing compatible primitives, in the same form as Theorem 5.83.

In the algebraic coefficient-current target the three brackets are identities, hence the curvature components vanish and the zero primitive is compatible with every structural map. In a Costello local-functional target the terms are graph weights, source faces, collision/contact terms, and counterterms. Supplying those rows and their primitive in the chosen finite-window system turns the displayed obstruction equation into a constructed homotopy.  $\square$

## C SIGN CONVENTIONS FOR THE HAMILTONIAN COMPARISON

The two  $\mathfrak{h}$ -modules attached to the labels  $(a, b)$  go in opposite directions. The polynomial PBW special-fibre descendant  $\tilde{\Psi}_{a,b}$ , symmetrised trace shadow of the open Hamiltonian action on the matrix Koszul algebra  $A_\psi$ , admits the lowering action  $z_1 \cdot \tilde{\Psi}_{a,b} = b \tilde{\Psi}_{a,b-1}$ ,  $z_2 \cdot \tilde{\Psi}_{a,b} = -a \tilde{\Psi}_{a-1,b}$ . The Matlis principal-part vector  $\rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2$ , residue dual of the formal Hamiltonian polydisk against its continuous Taylor dual, admits the raising action  $z_1 \cdot \rho_{a,b} = -(b+1) \rho_{a,b+1}$ ,  $z_2 \cdot \rho_{a,b} = (a+1) \rho_{a+1,b}$ . Diagonal rescaling  $\rho_{a,b} \mapsto \lambda_{a,b} \rho_{a,b}$ ,  $\tilde{\Psi}_{a,b} \mapsto \mu_{a,b} \tilde{\Psi}_{a,b}$  preserves the locally nilpotent lowering direction on descendants and the raising direction on principal parts; the directions of the index shifts already disagree.

The cohomological shift convention is  $(V[1])^n = V^{n+1}$ ; the reduced matrix Koszul letter  $\psi$  is odd, and cyclic graded signs use its parity. Absolute BV grading enters only through the global shift of the cotangent coordinate (Definition E.1). BV degree, ghost number, and form degree are tracked separately: BV degree is cohomological, ghost number is the antifield/ghost grading of the classical fields  $(\phi_1, \phi_2, \psi)$  and  $(\mathcal{A}, \mathcal{B})$ , and form degree is the de Rham/Dolbeault exterior grading on  $\Omega^\bullet(\mathbb{R}^2) \hat{\otimes} \Omega^{\circ,\bullet}(\mathbb{C}^2)$ ; none of the three is identified with another. Hochschild homology  $HH_\bullet$ , cyclic homology, and negative-cyclic homology  $HC_\bullet^-$  are distinguished at every reference; the cyclic trace symbol  $\text{Tr}$  below is the graded cyclic quotient of Definition C.1, the Hochschild trace is its image under the Connes  $B$ -operator, and the negative-cyclic class is the constellation-level Calabi–Yau pairing of the compact CY-to-chiral target. The BCOV holomorphic-anomaly equation enters only as a target equation on a compact Calabi–Yau threefold, with the BCOV-strict sign of  $[1]$  preserved by the matched-conventions divergence row of Theorem 10.2; the local Hamiltonian theorem supplies only the formal-stalk input.

*Definition C.1 (Local symplectic and cyclic conventions).* Fix the formal-local holomorphic symplectic datum  $(\widehat{\mathbb{C}}_0, \Omega_{\text{loc}} = dz_1 \wedge dz_2)$  of Section 8. The Hamiltonian vector field of  $f \in \mathbb{C}[[z_1, z_2]]$  is determined by

$$\iota_{X_f} \Omega_{\text{loc}} = -df,$$

giving

$$X_f = (\partial_{z_1} f) \partial_{z_2} - (\partial_{z_2} f) \partial_{z_1}, \quad X_{z_1} = \partial_{z_2}, \quad X_{z_2} = -\partial_{z_1}.$$

The Poisson bracket is

$$\{f, g\} = X_f(g) = \partial_{z_1} f \partial_{z_2} g - \partial_{z_2} f \partial_{z_1} g,$$

in particular  $\{z_1, z_2\} = 1$ . Hamiltonian fields are divergence-free for  $\Omega_{\text{loc}}$ : the volume-form scalar contribution is zero (Proposition E.2).

The reduced matrix Koszul algebra of Definition D.1 is

$$A_\psi = \mathbb{C}\langle x, y, \psi \rangle, \quad |x| = |y| = 0, \quad |\psi| \equiv 1 \pmod{2}, \quad Q\psi = x y - y x, \quad Qx = Qy = 0.$$

The graded cyclic quotient uses the Koszul sign rule

$$[uv]_{\text{cyc}} = (-1)^{|u||v|} [vu]_{\text{cyc}},$$

and the trace symbol  $\text{Tr}$  is graded cyclic with the same sign. The only odd letter is  $\psi$ .

**LEMMA C.2 (CE/PV cochain signs).** Let  $\mathfrak{h}$  have basis  $e_I$  and structure constants  $[e_I, e_J] = C_{IJ}^K e_K$ . Let  $c^I$  be the CE ghost dual to  $e_I$ , and let  $u_K$  be the shifted cotangent coordinate paired with  $e_K$  under the right-adjoint action  $e_J \cdot u_K = [u_K, e_J] = C_{KJ}^L u_L$ . The Chevalley–Eilenberg differential is

$$d_{\text{CE}} c^K = -\frac{1}{2} C_{IJ}^K c^I c^J, \quad d_{\text{CE}} u_K = C_{KJ}^L u_L c^J.$$

On the polyvector model with even coordinates  $O_K$  and odd coordinates  $\theta^I$ , Schouten bracket  $[\theta^I, O_J] = \delta_J^I$ , and BV interaction  $\pi = \frac{1}{2} C_{IJ}^K O_K \theta^I \theta^J$ , the differential  $d_\pi = [\pi, -]$  acts by

$$d_\pi \theta^K = -\frac{1}{2} C_{IJ}^K \theta^I \theta^J, \quad d_\pi O_K = C_{KJ}^L O_L \theta^J.$$

Therefore the assignment

$$c^I \mapsto \theta^I, \quad u_K \mapsto O_K$$

is a chain map intertwining  $d_{\text{CE}}$  and  $d_\pi$ .

*Proof.* The CE differential on  $c^K$  is the standard Koszul dual of the bracket, with the sign convention  $d_{\text{CE}} c^K = -\frac{1}{2} C_{IJ}^K c^I c^J$  ensuring  $d_{\text{CE}}^2 = 0$  by the Jacobi identity in  $\mathfrak{h}$ . The CE differential on  $u_K$  is the dual of the right-adjoint action  $e_J \cdot u_K = C_{KJ}^L u_L$ ; applying  $d_{\text{CE}}$  once more to  $C_{KJ}^L u_L c^J$ , with the Leibniz rule and  $|u_L| = 0$ , yields

$$C_{KJ}^L (C_{LA}^M u_M c^A) c^J - \frac{1}{2} C_{KJ}^L u_L C_{AB}^J c^A c^B,$$

which vanishes by the Jacobi identity for the right-adjoint representation.

The polyvector identity  $d_\pi^2 = 0$  is the master equation  $[\pi, \pi] = 0$ , which in the present notation is the Jacobi identity for  $\mathfrak{h}$  repackaged as a quadratic relation among the structure constants. The match of  $d_\pi$  on generators with the CE formulas above is standard (Costello [12], Chapter 5): one verifies on a single non-abelian example that  $\pi = \frac{1}{2} C_{IJ}^K O_K \theta^I \theta^J$  reproduces the right-adjoint CE differential on the cotangent coordinate, and the general case follows by linearity in  $\pi$ . Both  $d_{\text{CE}}$  and  $d_\pi$  are derivations, so the assignment  $c^I \mapsto \theta^I$ ,  $u_K \mapsto O_K$  extends from generators to a chain map on the full Chevalley–Eilenberg and polyvector algebras.  $\square$

LEMMA C.3 (*Primitive one- $\psi$  chain boundary*). In  $A_\psi$  of Definition C.1, for ordinary words  $u, v$  in  $x, y$ ,

$$Q \operatorname{Tr}(\psi u \psi v) = \operatorname{Tr}(\psi v x y u - \psi v y x u - \psi u x y v + \psi u y x v).$$

Removing the common front  $\psi$  gives the boundary in the one- $\psi$  chain model

$$\partial_2(u, v) = v x y u - v y x u - u x y v + u y x v.$$

*Proof.* The first displayed  $\psi$  contributes  $+(xy - yx)$  under  $Q$ ; the second contributes  $-(xy - yx)$  because exactly one odd letter precedes it, so the Koszul sign of the derivation flips. The even word produced by the first replacement may be moved cyclically past the remaining  $\psi$  and the word  $v$  with sign  $+1$ : the Koszul sign  $(-1)^{|u||v|}$  for moving an even string past a graded cyclic block is trivial. Removing the common front  $\psi$  identifies the result with the displayed boundary. Cross-checked against Lemma D.5.  $\square$

LEMMA C.4 (*Residue coadjoint signs*). Let

$$\rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2, \quad a, b \geq 0,$$

with residue pairing  $\langle \rho_{a,b}, z_1^m z_2^n \rangle = \delta_{a,m} \delta_{b,n}$  and coadjoint sign  $(f \cdot \rho)(g) = -\rho(\{f, g\})$ . Then

$$z_1^p z_2^q \cdot \rho_{a,b} = -(pb - qa + p - q) \rho_{a-p+1, b-q+1},$$

with negative-index targets and the scalar target  $\rho_{0,0}$  set to zero. In particular,

$$z_1 \cdot \rho_{a,b} = -(b+1) \rho_{a,b+1}, \quad z_2 \cdot \rho_{a,b} = (a+1) \rho_{a+1,b}.$$



*Proof.* Divergence-free Hamiltonian fields (Definition C.1) make the residue pairing  $\mathfrak{h}$ -equivariant via integration by parts with zero volume-form correction, fixing the coadjoint convention  $(f \cdot \rho)(g) = -\rho(\{f, g\})$ . Pair against a test monomial  $g = z_1^m z_2^n$ :

$$(z_1^p z_2^q \cdot \rho_{a,b})(z_1^m z_2^n) = -\rho_{a,b}(\{z_1^p z_2^q, z_1^m z_2^n\}).$$

Direct computation gives  $\{z_1^p z_2^q, z_1^m z_2^n\} = (pn - qm)z_1^{p+m-1}z_2^{q+n-1}$ , and the residue pairing keeps the  $(a, b)$  Taylor coefficient. The unique  $(m, n)$  producing  $z_1^a z_2^b$  is  $m = a - p + 1$ ,  $n = b - q + 1$ ; substitution yields coefficient  $p(b - q + 1) - q(a - p + 1) = pb - qa + p - q$ , and the leading minus from the coadjoint convention gives the displayed formula. Targets with  $a - p + 1 < 0$  or  $b - q + 1 < 0$  require  $m < 0$  or  $n < 0$  and are unreachable by polynomial test monomials, hence the cutoff at negative indices. The scalar target  $\rho_{0,0}$  is the case  $p = a + 1$ ,  $q = b + 1$ , at which the coefficient  $(a + 1)b - (b + 1)a + (a + 1) - (b + 1) = 0$  vanishes automatically; the formula is therefore consistent with the exclusion of  $\rho_{0,0}$  from  $\mathcal{P} = \text{Ann}_{\text{res}}(\mathbb{C} \cdot 1)$ .  $\square$

LEMMA C.5 (*PBW descendant signs*). The polynomial primitive one- $\psi$  descendants  $\tilde{\Psi}_{a,b}$  of Corollary 5.172 carry the symmetric PBW special-fibre action

$$z_1^p z_2^q : \tilde{\Psi}_{a,b} \mapsto (pb - qa)\tilde{\Psi}_{a+p-1, b+q-1}.$$

In particular,  $z_1 \cdot \tilde{\Psi}_{a,b} = b\tilde{\Psi}_{a,b-1}$  and  $z_2 \cdot \tilde{\Psi}_{a,b} = -a\tilde{\Psi}_{a-1,b}$ ; the action lowers the polynomial bidegree.

*Proof.* The Hamiltonian vector field of  $z_1^p z_2^q$  is

$$X_{z_1^p z_2^q} = pz_1^{p-1}z_2^q \partial_{z_2} - qz_1^p z_2^{q-1} \partial_{z_1}.$$

Acting on the symmetrised trace with  $a$  factors of  $x$  and  $b$  factors of  $y$ , the first summand differentiates one of the  $b$  copies of  $y$  and contributes  $pb$  copies of the symmetrised target  $\tilde{\Psi}_{a+p-1, b+q-1}$ ; the second differentiates one of the  $a$  copies of  $x$  and contributes  $-qa$  copies of the same target. The sum is the displayed coefficient.  $\square$

LEMMA C.6 (*Moyal expansion coefficients*). With

$$P = \partial_{z_1} \otimes \partial_{z_2} - \partial_{z_2} \otimes \partial_{z_1}, \quad f * g = m \circ \exp\left(\frac{\hbar}{2}P\right)(f \otimes g),$$

the normalized Moyal bracket  $\{f, g\}_{\hbar} = \hbar^{-1}(f * g - g * f)$  expands as

$$\{f, g\}_{\hbar} = \sum_{s \geq 0} \frac{\hbar^{2s}}{2^{2s}(2s+1)!} P^{2s+1}(f, g),$$

with first quantum correction  $\hbar^2 P^3(f, g)/24$ :

$$\{f, g\}_{\hbar} = \{f, g\} + \frac{\hbar^2}{24} P^3(f, g) + O(\hbar^4).$$

*Proof.* Even powers of  $P$  are symmetric and odd powers antisymmetric under  $f \leftrightarrow g$ ; the commutator therefore retains only the odd powers and doubles them:

$$f * g - g * f = 2 \sum_{s \geq 0} \frac{(\hbar/2)^{2s+1}}{(2s+1)!} P^{2s+1}(f, g).$$

Division by  $\hbar$  gives the displayed series. The  $s = 1$  coefficient is  $1/(2^2 \cdot 3!) = 1/24$ ; see Theorem F.2 for the all-order identity and `scripts/check_moyal_coefficients.py` for the numerical sweep through order eleven.  $\square$

LEMMA C.7 (*Moyal coadjoint signs*). Let  $(n)_r = n(n-1) \cdots (n-r+1)$ , with  $(n)_0 = 1$ , denote the falling factorial. For  $f = z_1^p z_2^q$ , the normalized Moyal coadjoint action on principal parts is

$$f \cdot_{\hbar} \rho_{a,b} = - \sum_{\substack{r \geq 1 \\ r \text{ odd}}} \frac{\hbar^{r-1}}{2^{r-1} r!} C_r(p, q; a-p+r, b-q+r) \rho_{a-p+r, b-q+r},$$

with negative-index targets and the scalar target  $\rho_{0,0}$  set to zero. Here

$$C_r(p, q; m, n) = \sum_{\alpha=0}^r (-1)^\alpha \binom{r}{\alpha} (p)_{r-\alpha} (q)_\alpha (m)_\alpha (n)_{r-\alpha}$$

is the bidifferential coefficient of Theorem F.2. At  $r = 1$  the formula recovers the residue coadjoint of Lemma C.4,

$$z_1^p z_2^q \cdot \rho_{a,b} = -(pb - qa + p - q) \rho_{a-p+1, b-q+1}.$$

*Proof.* Apply the bidifferential  $P^r$  to  $z_1^p z_2^q \otimes z_1^m z_2^n$ : Theorem F.2 gives  $P^r(f, g) = C_r(p, q; m, n) z_1^{p+m-r} z_2^{q+n-r}$ . By Lemma C.6 the normalized Moyal bracket keeps only odd  $r$ , with coefficient  $\hbar^{r-1}/(2^{r-1} r!)$ . The coadjoint convention  $(f \cdot_{\hbar} \rho)(g) = -\rho(\{f, g\}_{\hbar})$  introduces the leading minus sign and determines the target principal part: pairing with  $\rho_{a,b}$  selects  $(m, n)$  such that  $p + m - r = a$  and  $q + n - r = b$ , i.e.  $m = a - p + r$ ,  $n = b - q + r$ . At  $r = 1$  the coefficient is  $C_1(p, q; a - p + 1, b - q + 1) = p(b - q + 1) - q(a - p + 1) = pb - qa + p - q$ , reproducing Lemma C.4.  $\square$

COROLLARY C.8 (*Universal residue-dual normalization*). With the residue pairing  $\langle \rho_{a,b}, z_1^m z_2^n \rangle = \delta_{a,m} \delta_{b,n}$  and the coadjoint convention

$$(f \cdot \rho)(g) = -\rho(\{f, g\}), \quad (f \cdot_{\hbar} \rho)(g) = -\rho(\{f, g\}_{\hbar}),$$

the linear Hamiltonians act on  $\mathcal{P}$  by

$$z_1 \cdot \rho_{a,b} = -(b+1) \rho_{a,b+1}, \quad z_2 \cdot \rho_{a,b} = (a+1) \rho_{a+1,b},$$

while the polynomial descendants of Lemma C.5 satisfy

$$z_1 \cdot \tilde{\Psi}_{a,b} = b \tilde{\Psi}_{a,b-1}, \quad z_2 \cdot \tilde{\Psi}_{a,b} = -a \tilde{\Psi}_{a-1,b}.$$

Diagonal rescaling  $\rho_{a,b} \mapsto \lambda_{a,b} \rho_{a,b}$ ,  $\tilde{\Psi}_{a,b} \mapsto \mu_{a,b} \tilde{\Psi}_{a,b}$  with  $\lambda_{a,b}, \mu_{a,b} \in \mathbb{C}^\times$  preserves the opposite index directions of the two actions. Adjustments of factorial normalizations or of the Moyal  $1/24$  coefficient act diagonally and leave the index shifts intact.

*Proof.* The two displayed actions and the residue pairing are restated from Lemma C.4 and Lemma C.5 at  $(p, q) \in \{(1, 0), (0, 1)\}$ . A diagonal rescaling sends  $z_1 \cdot \rho_{a,b}$  to  $-(b+1)(\lambda_{a,b+1}/\lambda_{a,b}) \rho_{a,b+1}$ , preserving the raising target index, and sends  $z_1 \cdot \tilde{\Psi}_{a,b}$  to  $b(\mu_{a,b-1}/\mu_{a,b}) \tilde{\Psi}_{a,b-1}$ , preserving the lowering target index. Equality of the two actions would require the same target index for both, contradicting the directions  $b \mapsto b+1$  versus  $b \mapsto b-1$ . Higher Moyal corrections  $r \geq 3$  shift indices by  $b \mapsto b - q + r \geq b+1$

for  $(p, q) = (1, 0)$  and by  $a \mapsto a - p + r \geq a + 1$  for  $(p, q) = (0, 1)$ ; they raise as well, so the obstruction persists at every order in  $\hbar$ . The compute log scripts/check\_one\_psi\_homology.py records this obstruction explicitly: writing  $\delta_{a,b}$  for the residue functional  $\rho_{a,b}$  under the Matlis identification of Lemma 8.4, the third-order Moyal coadjoint gives  $\text{ad}_{\delta_{1,1}}^*(\delta_{1,1}) = -24\hbar^2 \delta_{0,4}$ , while the polynomial one- $\psi$  action sends  $\Psi_{1,1}$  to  $4\Psi_{4,0}$ ; even matched at the Capelli/Weyl shift  $\hbar N$ , the higher Moyal coadjoint terms have zero PBW descendant counterpart.  $\square$

LEMMA C.9 (*Brane-line current source convention*). For an interval  $I \subset \mathbb{R}$ , the Hamiltonian source complex is  $\Omega_c^\bullet(I) \widehat{\otimes} \mathfrak{h}$  of Definition B.1. A degree-zero source density is written  $a(t)dt \otimes f$ , and the cotangent CE source coordinate maps to the boundary observable by

$$u_{a(t)dt \otimes f} \mapsto B_f(a) = \int_I a(t) B_f(t) dt.$$

The companion ghost coordinate  $c_{a \otimes f}$  maps to the PV coordinate  $\theta_{a \otimes f}$ . For  $B \in \mathcal{D}'_c(I)$ , the multiplication of the distribution  $B$  by a smooth test function  $a$  is the standard  $C_c^\infty(I)$ -module operation  $aB$ , defined by  $(aB)(\varphi) = B(a\varphi)$ . The interval current brackets are

$$\{B_f(a), \Theta_\rho(B)\} = \Theta_{f \cdot \rho}(aB), \quad \{\Theta_\rho(B), \Theta_\sigma(C)\} = 0.$$

*Proof.* The de Rham factor supplies the compactly supported density on the brane line; the Hamiltonian current factor supplies  $f$ . Pairing this source with the boundary field gives the integral defining  $B_f(a)$ . The first bracket is the semidirect-product bracket of Definition B.2 for  $\mathfrak{h}_I \ltimes \mathcal{P}_I[1]$ : the action factors as the  $C_c^\infty(I)$ -module operation  $B \mapsto aB$  tensored with the Matlis coadjoint action of Lemma C.4. The second bracket vanishes because the shifted principal-part summand is an abelian odd ideal.  $\square$

## D FULL PRIMITIVE HIGHER- $\psi$ KOSZUL HOMOLOGY

The polynomial PBW one- $\psi$  descendant labels  $\Psi_{a,b}$  of the open BV brane sector and the residual Matlis principal parts  $\rho_{a,b}$  of § 8 share the bidegree index set  $(a, b)$  and carry different  $\mathfrak{h}$ -actions. The defect position is polynomial; its cotangent momentum is residual. The separation is the homology of a single universal calculation: the full primitive cyclic Koszul homology of  $(A_\psi, d\psi = xy - yx)$  graded by  $\psi$ -count and by  $(R, S) = (\#x + \#\psi, \#y + \#\psi)$ . The surviving  $H_1$ -line is the special-fibre image of the PBW Rees lattice; the associated-graded morphism  $\text{gr } \Phi : \bar{A}_{\text{desc}}[1] \rightarrow \mathcal{P}$  is the bidegree-only special-fibre relabelling.

The universal primitive single-trace Koszul complex is attached to the moment-map equation  $[x, y] = 0$ . Its universality is independent of any choice of Matlis target, PBW model, finite  $N$  trace identity, or quantum deformation: after stable trace invariant theory has identified primitive traces with graded cyclic words, the answer is the cellular homology of a chip-moving state complex. Every finite-bidegree stable primitive descendant calculation over  $\mathbb{C}$ , with  $d\psi = xy - yx$  and graded cyclicity as the trace relation, is obtained from this one. Passage to completed coefficient systems requires an exact coefficient functor, or an explicit condition excluding new  $\varprojlim^1$  classes.

Definition D.1 (*Graded cyclic word complex*). Let

$$A_\psi = \mathbb{C}\langle x, y, \psi \rangle, \quad |x| = |y| = 0, \quad |\psi| \equiv 1 \pmod{2},$$

and let  $d$  be the odd derivation determined by

$$d\psi = xy - yx, \quad dx = dy = 0.$$

Put

$$\text{Cyc}(A_\psi) = A_\psi / [A_\psi, A_\psi]_{\text{gr}}, \quad [uv] = (-1)^{|u||v|} [vu].$$

Give the letters weights

$$\text{wt}(x) = (1, 0), \quad \text{wt}(y) = (0, 1), \quad \text{wt}(\psi) = (1, 1).$$

For  $R, S \geq 0$  define

$$K_{R,S}^p = \mathbb{C}\{[w]_{\text{cyc}} : \#\psi(w) = p, \#x(w) = R - p, \#y(w) = S - p\},$$

with  $K_{R,S}^p = 0$  unless  $0 \leq p \leq \min(R, S)$ . If  $w = a_1 \cdots a_n$ , the boundary is

$$\begin{aligned} \partial[w] = \sum_{i: a_i = \psi} (-1)^{\#\{j < i: a_j = \psi\}} [a_1 \cdots a_{i-1} x y a_{i+1} \cdots a_n \\ - a_1 \cdots a_{i-1} y x a_{i+1} \cdots a_n]_{\text{cyc}}. \end{aligned}$$

Thus  $\partial : K_{R,S}^p \rightarrow K_{R,S}^{p-1}$ .

LEMMA D.2 (*Stable super-cyclic trace presentation*). Fix a total word length  $d$ . In the stable range  $N \geq d$ , the primitive  $GL_N$ -invariant single-trace functions in the variables  $x, y, \psi$ , homogeneous of total word length  $d$ , are exactly the graded cyclic words in  $\text{Cyc}(A_\psi)$ . Equivalently, after connected single-trace extraction, the only stable relation among words in  $x, y, \psi$  is

$$\text{Tr}(uv) = (-1)^{|u||v|} \text{Tr}(vu).$$

*Proof.* First replace the odd letter  $\psi$  by  $p$  distinct even marked matrices  $\eta_1, \dots, \eta_p$ , and take the summand multilinear in the  $\eta_i$ 's and homogeneous of the prescribed  $x, y$ -degree. The Procesi–Razmyslov stable invariant theorem [17, 18] identifies the primitive single-trace invariants of total word length  $d$ , for  $N \geq d$ , with ordinary cyclic words in the letters  $x, y, \eta_1, \dots, \eta_p$ , modulo only cyclicity of the trace.

The symmetric group  $\mathbb{S}_p$  permutes the marked letters. Passing from  $p$  even marked letters to one odd letter is the sign-isotypic projection for this action, followed by the identification  $\eta_i \mapsto \psi$ . Since the ground field has characteristic zero, this projection is exact. Interchanging two marked occurrences contributes the sign of the transposition, and rotating a block with  $a$  marked letters past a block with  $b$  marked letters contributes  $(-1)^{ab}$ . This is precisely the graded cyclic rule for a single odd letter  $\psi$ . This is the complete stable trace relation in word length  $d$ .  $\square$

LEMMA D.3 (*Boundary signs and  $\partial^2 = 0$* ). The formula of Definition D.1 is well-defined on the graded cyclic quotient and satisfies  $\partial^2 = 0$ .

*Proof.* On the free algebra  $A_\psi$ , the displayed formula is the derivation rule for the odd derivation  $d$ : replacing the  $i$ -th occurrence of  $\psi$  crosses exactly the preceding odd letters and hence contributes  $(-1)^{\#\{j < i: a_j = \psi\}}$ . Since  $d(x) = d(y) = 0$  and  $d(\psi) = xy - yx$  is even and  $d$ -closed, one has  $d^2 = 0$  on the generators, hence on all of  $A_\psi$ .

If  $u, v$  are homogeneous, then

$$d(uv - (-1)^{|u||v|} vu)$$

is a sum of graded commutators, using the odd Leibniz rule and the fact that  $d$  lowers the number of odd letters by one. Thus  $d$  preserves  $[A_\psi, A_\psi]_{\text{gr}}$ , so it descends to  $\text{Cyc}(A_\psi)$ . The descended operator is exactly  $\partial$ , and  $\partial^2 = 0$  follows from  $d^2 = 0$ .  $\square$

LEMMA D.4 (*Exact  $p$ -face formula*). Let  $u_1, \dots, u_p$  be ordinary words in  $x, y$ . In  $K_{R,S}^{p-1}$ ,

$$\begin{aligned} \partial[\psi u_1 \psi u_2 \cdots \psi u_p] &= \sum_{j=1}^p (-1)^{j-1} [\psi u_1 \cdots \psi u_{j-1} x y u_j \psi u_{j+1} \cdots \psi u_p \\ &\quad - \psi u_1 \cdots \psi u_{j-1} y x u_j \psi u_{j+1} \cdots \psi u_p]_{\text{cyc}}, \end{aligned}$$

where, for  $j = 1$ , the initial displayed  $\psi u_1 \cdots \psi u_{j-1}$  is the empty word.

*Proof.* This is the boundary of Definition D.1 applied to the representative  $\psi u_1 \psi u_2 \cdots \psi u_p$ . The  $j$ -th displayed  $\psi$  has  $j - 1$  preceding odd letters, and replacing it by the even word  $x y - y x$  contributes sign  $+1$  from the replacement.  $\square$

LEMMA D.5 (*Two- $\psi$  boundary sign*). For ordinary words  $u, v$ ,

$$\partial[\psi u \psi v] = [\psi v x y u] - [\psi v y x u] - [\psi u x y v] + [\psi u y x v].$$

After removing the common front  $\psi$  from each term, this is the boundary

$$v x y u - v y x u - u x y v + u y x v.$$

*Proof.* Lemma D.4 with  $p = 2$  gives

$$[(x y - y x) u \psi v] - [\psi u (x y - y x) v].$$

The first two terms are cyclically rotated to put the remaining  $\psi$  in front:

$$[(x y) u \psi v] = [\psi v x y u], \quad [(y x) u \psi v] = [\psi v y x u].$$

The moved block is even, so the Koszul sign is  $+1$ . The second replacement already has the surviving  $\psi$  in front, and its overall sign is the sign  $-1$  from the preceding odd letter. Expanding gives the displayed formula.

Interchanging  $u$  and  $v$  gives  $u x y v - u y x v - v x y u + v y x u$ , the negative of the displayed expression. Thus the formula is compatible with the alternating two- $\psi$  quotient used below.  $\square$

CONSTRUCTION D.6 (*One- $\psi$  three-term skeleton*). For  $k, l \geq 0$ , let  $W_{k,l}$  be the ordinary words with  $k$  letters  $x$  and  $l$  letters  $y$ , and let  $N_{a,b}$  be the necklaces with  $a$  letters  $x$  and  $b$  letters  $y$ . The one- $\psi$  skeleton in bidegree  $(k, l)$  is the finite complex

$$C_2(k, l) \xrightarrow{\partial_2} C_1(k, l) \xrightarrow{\partial_1} C_0(k, l),$$

with

$$C_1(k, l) = \mathbb{C}\{W_{k,l}\}, \quad C_0(k, l) = \mathbb{C}\{N_{k+1,l+1}\},$$

and

$$C_2(k, l) = (\mathbb{C}\{(u, v) : \deg u + \deg v = (k - 1, l - 1)\})_{\text{alt}},$$

where  $(u, v) = -(v, u)$  and  $(u, u) = 0$ . If  $k = 0$  or  $l = 0$ , then  $C_2(k, l) = 0$ . The boundaries are

$$\partial_1(w) = [x y w]_{\text{cyc}} - [y x w]_{\text{cyc}},$$

and

$$\partial_2(u, v) = vx y u - v y x u - u x y v + u y x v.$$

This is the truncation  $K_{k+1, l+1}^2 \rightarrow K_{k+1, l+1}^1 \rightarrow K_{k+1, l+1}^0$  after putting the remaining  $\psi$  at the front of a one- $\psi$  cyclic word. Equivalently, it is the cellular chain complex in dimensions 2, 1, 0 of the state complex constructed below:  $C_0$  records chip configurations,  $C_1$  records one moving chip, and  $C_2$  records two independent moving chips with the orientation fixed by the ordered moving edges.

*Proof.* The identifications of  $C_1$  and  $C_0$  are exactly the graded cyclic normal forms in  $\psi w$  and the ordinary necklaces. The source  $C_2$  is the two- $\psi$  summand of  $K_{k+1, l+1}^2$ : after cutting at one displayed  $\psi$ , a class is written  $\psi u \psi v$ , and moving the second  $\psi$ -block past the first contributes the sign  $-1$ . Lemma D.5 gives  $\partial_2$ , while the definition of  $\partial$  gives  $\partial_1$ .  $\square$

*Construction D.7 (Labelled chip-moving state complex).* Assume  $R, S > 0$ . Let  $\mathcal{C}_S$  be the oriented circle subdivided into  $S$  labelled oriented edges and  $S$  labelled vertices. Define  $\tilde{Y}_{R, S}$  as follows. A  $p$ -cell is a state with  $R - p$  stationary indistinguishable chips on vertices and  $p$  moving chips in the interiors of  $p$  distinct edges. Several stationary chips may occupy the same vertex. Each edge contains at most one moving chip.

Encode such a state by traversing  $\mathcal{C}_S$ . At a vertex write one  $x$  for each stationary chip sitting there. Then write  $y$  if the following edge is empty and  $\psi$  if the following edge contains a moving chip. This gives a word with  $R - p$  letters  $x$ ,  $S - p$  letters  $y$ , and  $p$  letters  $\psi$ , with a distinguished starting edge.

The  $p$ -cell is oriented by the increasing order of the labelled edges carrying moving chips. Its  $j$ -th oriented coordinate has boundary

$$\text{terminal face } x y - \text{initial face } y x.$$

Hence the cellular boundary in the labelled complex is

$$\sum_j (-1)^{j-1} (\text{replace the } j\text{-th moving edge by } x y - \text{replace it by } y x).$$

The cyclic group  $\Gamma_S = \mathbb{Z}/S\mathbb{Z}$  rotates the edge labels and acts cellularly on  $\tilde{Y}_{R, S}$ .

**LEMMA D.8 (Cellular chain identification with cyclic relabelling).** For  $R, S > 0$ , the cellular chain complex of  $\tilde{Y}_{R, S}$ , after taking  $\Gamma_S$ -coinvariants, is naturally isomorphic to  $K_{R, S}^\bullet$ :

$$C_p(\tilde{Y}_{R, S}; \mathbb{C})_{\Gamma_S} \cong K_{R, S}^p.$$

Under this isomorphism the cellular boundary is the boundary  $\partial$  of Definition D.1.

*Proof.* A labelled  $p$ -cell is equivalently a labelled cyclic sequence

$$x^{a_1} m_1 x^{a_2} m_2 \cdots x^{a_S} m_S, \quad m_i \in \{y, \psi\},$$

with exactly  $p$  of the  $m_i$ 's equal to  $\psi$  and  $\sum_i a_i = R - p$ . Every graded cyclic word in  $K_{R, S}^p$  has a representative of this form: move the cut, if necessary, past even  $x$ -letters until it lies immediately before an edge marker  $y$  or  $\psi$ . This representative is unique after a starting edge has been chosen. The remaining ambiguity is rotation of the  $S$  edge markers, hence quotienting by  $\Gamma_S$  is exactly passage to graded cyclic words. The labelled complex orders the  $S$  edge markers, and in the coinvariant complex the only identifications are their rotations with the orientation character below.

The sign is determined as follows. Suppose a rotation moves a block containing  $a$  occurrences of  $\psi$  past the complementary block containing  $b$  occurrences of  $\psi$ . On cellular orientations this rotation moves  $a$  oriented interval coordinates past  $b$  oriented interval coordinates, hence changes the orientation by  $(-1)^{ab}$ . This is exactly the graded cyclic sign  $[uv] = (-1)^{|u||v|}[vu]$ , since  $x$  and  $y$  are even and every  $\psi$  is odd.

With this orientation convention, the cellular boundary in Construction D.7 is

$$\sum_j (-1)^{j-1} ([xy] - [yx])$$

in the  $j$ -th moving edge. The terminal face is the state in which the chip has crossed the edge and is read as  $xy$ ; the initial face is read as  $yx$ . This is exactly Lemma D.4, term by term.  $\square$

LEMMA D.9 (*Abel map for labelled states*). For  $R, S > 0$ , the labelled state complex  $\tilde{Y}_{R,S}$  has the homology of a circle:

$$H_q(\tilde{Y}_{R,S}; \mathbb{C}) \cong \begin{cases} \mathbb{C}, & q = 0, \\ \mathbb{C}, & q = 1, \\ 0, & q \geq 2. \end{cases}$$

The cyclic rotation action of  $\Gamma_S$  is trivial on these two homology lines.

*Proof.* Work in the universal cover of  $\mathcal{C}_S$ , whose vertices are the integers and whose edges are the unit intervals. A lifted labelled cell is described by the positions of the  $R$  chips: stationary chips have integer positions, and a moving chip on the edge  $[m, m+1]$  has coordinate  $m+t$  with  $0 < t < 1$ . The closure of the cell permits  $t = 0, 1$ , which are exactly the two vertex states  $yx$  and  $xy$  in the boundary convention of Construction D.7. Define the Abel map

$$A : \tilde{Y}_{R,S} \longrightarrow \mathbb{R}/S\mathbb{Z}$$

by summing the chip positions on the oriented circle, counted with multiplicity, and reducing modulo  $S$ . On every cell this map is affine.

Lift the target to  $\mathbb{R}$  and fix a cyclic order of the  $R$  lifted chips. A lifted fibre is described by weakly ordered coordinates

$$t_1 \leq t_2 \leq \cdots \leq t_R \leq t_1 + S$$

with fixed sum. This is a convex polytope. Its faces are cut out by the equalities  $t_i \in \mathbb{Z}$  and by the choices of those unit intervals which contain a moving chip; these faces are precisely the cells of the chip-moving subdivision meeting the fibre. Hence every lifted fibre is a contractible finite polyhedral complex. The map is a proper PL map of finite polyhedra with acyclic fibres. The Vietoris–Begle theorem, applied to this affine cellular map, gives a homology equivalence

$$H_\bullet(\tilde{Y}_{R,S}; \mathbb{C}) \cong H_\bullet(S^1; \mathbb{C}).$$

A cyclic relabelling of the  $S$  edges adds a constant to every chip coordinate, so it rotates the target circle by an orientation-preserving rotation. Therefore  $\Gamma_S$  acts trivially on  $H_0(S^1; \mathbb{C})$  and  $H_1(S^1; \mathbb{C})$ , and hence on the two homology lines of  $\tilde{Y}_{R,S}$ .  $\square$

LEMMA D.10 (*Cyclic Abel quotient and stabilizer fibres*). Let  $Y_{R,S}$  denote the cyclic relabelling quotient of  $\tilde{Y}_{R,S}$ , with the orientation character encoded by the graded cyclic rule. Then the quotient chain complex  $C_\bullet(\tilde{Y}_{R,S}; \mathbb{C})_{\Gamma_S}$  has the homology of  $S^1$ . Moreover, every stabilizer quotient of an Abel fibre is contractible.

*Proof.* By Lemma D.9, the labelled Abel map is a homology equivalence from  $\tilde{Y}_{R,S}$  to the circle. A cyclic relabelling by  $r \in \Gamma_S$  adds  $r$  to each of the  $R$  chip positions, hence translates the Abel coordinate by  $Rr$  in  $\mathbb{R}/S\mathbb{Z}$ . The action on the target circle is by orientation-preserving rotations, so it is trivial on the two homology lines. Since  $\mathbb{C}[\Gamma_S]$  is semisimple,

$$H_\bullet(C_\bullet(\tilde{Y}_{R,S}; \mathbb{C})_{\Gamma_S}) \cong H_\bullet(\tilde{Y}_{R,S}; \mathbb{C})_{\Gamma_S} \cong H_\bullet(S^1; \mathbb{C}).$$

Now fix an Abel value and a stabilizer subgroup  $G \subset \Gamma_S$  of that value. In the universal cover, choose the lifted fibre before decomposing it into chip cells. It is a convex polytope cut out by weak ordering, the fixed-sum equation, and the chip-cell inequalities, and  $G$  acts affinely on it. Averaging a point over  $G$  gives a  $G$ -fixed point. Straight-line contraction to this fixed point is  $G$ -equivariant and respects the chip-cell subdivision. Hence the stabilizer quotient of the Abel fibre is contractible.  $\square$

LEMMA D.11 ( $S^1$  generator). For  $R, S > 0$ , the cycle

$$\Psi_{R-1, S-1} = \sum_{w \in \mathbb{W}_{R-1, S-1}} [\psi w]_{\text{cyc}}$$

represents the fundamental  $H_1$ -line of the cyclic chip-moving complex.

*Proof.* Its boundary is

$$\partial \Psi_{R-1, S-1} = \sum_{w \in \mathbb{W}_{R-1, S-1}} ([x y w]_{\text{cyc}} - [y x w]_{\text{cyc}}).$$

For a fixed necklace with  $R$  letters  $x$  and  $S$  letters  $y$ , the coefficient is the number of cyclic  $x y$ -transitions minus the number of cyclic  $y x$ -transitions. These two numbers are equal on a circle; hence  $\Psi_{R-1, S-1}$  is closed.

Let  $\epsilon : K_{R,S}^1 \rightarrow \mathbb{C}$  send every one- $\psi$  cyclic word to 1. Lemma D.5 shows that every two- $\psi$  boundary has two positive and two negative terms, so it lies in  $\ker \epsilon$ . But

$$\epsilon(\Psi_{R-1, S-1}) = \binom{R+S-2}{R-1} \neq 0.$$

By Lemma D.10, the  $H_1$ -line is one-dimensional. Moreover, every one-cell in the displayed sum is oriented in the positive Abel direction: its terminal face has the moved chip one edge farther along the oriented circle. Thus the displayed cycle maps to  $\binom{R+S-2}{R-1}$  times the oriented circle class and spans the fundamental  $H_1$ -line.  $\square$

THEOREM D.12 (*Full primitive higher- $\psi$  Koszul homology*). For every  $R, S \geq 0$ ,

$$H_p(K_{R,S}) \cong \begin{cases} \mathbb{C} \cdot [x^R y^S], & p = 0, \\ \mathbb{C} \cdot [\Psi_{R-1, S-1}], & p = 1, R, S \geq 1, \\ 0, & p \geq 2, \end{cases}$$



where

$$\Psi_{R-1, S-1} = \sum_{w \in W_{R-1, S-1}} [\psi w]_{\text{cyc}}.$$

Consequently the primitive stable Koszul homology is

$$\mathbb{C}[x, y] \oplus \mathbb{C}[x, y]\theta, \quad |\theta| = 1, \quad \text{wt}(\theta) = (1, 1),$$

with the scalar Hamiltonian removed later if one passes to the reduced Hamiltonian sector.

*Proof.* If  $R = 0$  or  $S = 0$ , then  $K_{R,S}^p = 0$  for  $p > 0$ , and  $K_{R,S}^0$  is spanned by the unique cyclic word  $x^R$  or  $y^S$  (including the empty word when  $R = S = 0$ ). This gives the stated homology.

Assume  $R, S > 0$ . By Lemma D.8,  $K_{R,S}^\bullet$  is the  $\Gamma_S$ -coinvariant cellular chain complex of  $\tilde{Y}_{R,S}$ . Since  $\mathbb{C}[\Gamma_S]$  is semisimple, taking coinvariants is an exact functor on finite-dimensional  $\mathbb{C}[\Gamma_S]$ -modules. Therefore

$$H_p(K_{R,S}) \cong H_p(\tilde{Y}_{R,S}; \mathbb{C})_{\Gamma_S}.$$

Lemma D.10 identifies this homology with  $H_\bullet(S^1; \mathbb{C})$ . Thus  $H_p(K_{R,S})$  is one-dimensional for  $p = 0, 1$  and vanishes for  $p \geq 2$ . In particular, the three-term skeleton  $C_2 \rightarrow C_1 \rightarrow C_0$  already computes  $H_1$ : cells of dimension  $p \geq 3$  have boundaries in  $C_{p-1}$  and preserve  $\ker(C_1 \rightarrow C_0)/\text{im}(C_2 \rightarrow C_1)$ .

The degree-zero line is generated by any necklace of multidegree  $(R, S)$ . The functional sending every zero- $\psi$  necklace to 1 kills every one- $\psi$  boundary, so this line is nonzero; it is denoted  $\mathbb{C} \cdot [x^R y^S]$ .

Lemma D.11 identifies the degree-one generator as  $\Psi_{R-1, S-1}$ .  $\square$

*Remark D.13 (Numerical anchor).* The script `scripts/check_one_psi_homology.py` tabulates the same three-term skeleton over  $\mathbb{Q}$ , confirms  $\partial_1 \partial_2 = 0$ , checks that  $\Psi_{R-1, S-1}$  is closed and independent modulo boundaries, and verifies

$$\dim H_1(K_{R,S}) = 1$$

in each of the 36 bidegrees  $0 \leq R-1, S-1 \leq 5$ . The same script verifies the open Hamiltonian descendant action of Proposition 5.171 in 240 cases with exponents at most 3, and the closed coadjoint Taylor-dual / residue formula of Proposition 8.7 in 1225 cases with exponents at most 5. The PBW-versus-Moyal sweep records 1200 explicit mismatches and 958 cases where the coadjoint Moyal action carries genuine  $\hbar^{\geq 2}$  higher-order terms; the 70 linear-generator leading-order tests with labels at most 5 identify the exact obstruction to any  $\hbar$ -adic deformation of the classical  $\Psi \rightarrow \rho$  projection. The displayed homology lines and the separation in Theorem 5.180 are therefore proved at the bidegree level by an independent finite computation.

*Construction D.14 (Special-fibre PBW one- $\psi$  label basis).* Let

$$\mathfrak{h} = \mathbb{C}[[z_1, z_2]] / \mathbb{C} \cdot 1, \quad \mathfrak{h}_{\text{cont}}^\vee \cong \mathcal{P} = \bigoplus_{a+b>0} \mathbb{C} \rho_{a,b},$$

with the Hamiltonian Lie algebra of Definition 8.3 and the Matlis principal-part module of Definition 8.5. Let

$$\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1]$$

be the semidirect Hamiltonian shifted-cotangent Lie algebra of Proposition 2.7, and let

$$U_\hbar(\mathfrak{g}) \supset \mathbf{S}_\hbar(\mathfrak{h} \oplus \mathfrak{h}^\vee[1])$$

denote its Rees PBW lattice over  $\mathbb{C}[[\hbar]]$ , namely the Rees algebra of the canonical PBW filtration. The special fibre is

$$\mathbf{S}_{\hbar}(\mathfrak{h} \oplus \mathfrak{h}^{\vee}[1])|_{\hbar=0} \cong \mathbf{S}(\mathfrak{h} \oplus \mathfrak{h}^{\vee}[1]),$$

the symmetric algebra carrying the trivial  $\mathfrak{h}$ -action on generators. The *primitive one- $\psi$  PBW label sector* is the one-tensor-factor piece in  $\mathfrak{h}^{\vee}[1]$ , tensored with the symmetric algebra of  $\mathfrak{h}$ , in bidegree  $(a, b)$ :

$$\tilde{A}_{\text{desc}}[1] := \bigoplus_{a+b>0} \mathbb{C} \cdot e_{a,b}^{\text{PBW}}[1], \quad e_{a,b}^{\text{PBW}}[1] \in \mathbf{S}^1(\mathfrak{h}^{\vee}[1]) \otimes \mathbf{S}(\mathfrak{h})|_{\hbar=0}.$$

The boundary pairing of Corollary 3.29 identifies this label sector with the surviving primitive  $H_1$ -lines of Theorem D.12: under

$$e_{a,b}^{\text{PBW}}[1] \mapsto [\Psi_{a,b}], \quad a+b>0,$$

the  $\Psi_{a,b}$  of Theorem D.12 are exactly the special-fibre representatives of the PBW label basis, with the binomial weight  $\epsilon(\Psi_{a,b}) = \binom{a+b}{a}$  recording the totally symmetric averaging of the cyclic words.

**PROPOSITION D.15** (*Associated-graded label morphism*). Let

$$\tilde{\Psi}_{a,b} := \binom{a+b}{a}^{-1} \Psi_{a,b} \in \tilde{A}_{\text{desc}}[1]$$

be the unit-normalised special-fibre PBW label, so that  $\epsilon(\tilde{\Psi}_{a,b}) = 1$ , and let  $\rho_{a,b} \in \mathcal{P}$  be the Matlis principal-part basis of Definition 8.5. The bidegree-only relabelling

$$\text{gr } \Phi : \tilde{A}_{\text{desc}}[1] \xrightarrow{\sim} \mathcal{P}, \quad \tilde{\Psi}_{a,b} \mapsto \rho_{a,b}, \quad a+b>0,$$

is an isomorphism of  $\mathbb{Z}_{\geq 0}^2$ -graded  $\mathbb{C}$ -vector spaces. It is the special-fibre associated-graded label match of Theorem 8.12 in the  $\Psi$ -presentation of Construction D.14.

This morphism is a bigraded vector-space isomorphism with an  $\mathfrak{h}$ -module obstruction. More precisely, on the linear Hamiltonians,

$$z_1 \cdot \tilde{\Psi}_{a,b} = b \tilde{\Psi}_{a,b-1}, \quad z_2 \cdot \tilde{\Psi}_{a,b} = -a \tilde{\Psi}_{a-1,b},$$

while

$$z_1 \cdot \rho_{a,b} = -(b+1) \rho_{a,b+1}, \quad z_2 \cdot \rho_{a,b} = (a+1) \rho_{a+1,b}.$$

The two actions have opposite shift directions and unequal scalars; in particular the polynomial PBW action is locally nilpotent on every  $\tilde{\Psi}_{a,b}$ , whereas every nonzero  $\rho_{a,b}$  has an infinite principal-part coadjoint orbit. The obstruction to an  $\hbar$ -adic same-action lift of  $\text{gr } \Phi$  is the local-nilpotence dichotomy of Theorem 8.11 and Corollary 8.13.

*Proof.* The bidegree component of  $\tilde{A}_{\text{desc}}[1]$  is the line  $\mathbb{C} \cdot e_{a,b}^{\text{PBW}}[1]$  for  $a+b>0$ ; the bidegree component of  $\mathcal{P}$  is  $\mathbb{C} \cdot \rho_{a,b}$ ,  $a+b>0$ , by Definition 8.5. The displayed map is the bidegree identity, hence an isomorphism of bigraded vector spaces.

The polynomial Hamiltonian action on  $\tilde{\Psi}_{a,b}$  is Proposition 5.171: for  $F_{p,q} = \text{Tr}(\phi_1^p \phi_2^q)$ ,  $F_{p,q} \cdot \tilde{\Psi}_{a,b} = (pb - qa) \tilde{\Psi}_{a+p-1, b+q-1}$ . Setting  $(p, q) = (1, 0)$ ,  $(0, 1)$  gives the displayed two formulas. The principal-part coadjoint action is Proposition 8.7: for  $(p, q) = (1, 0)$ ,  $(0, 1)$  the formula  $z_1^p z_2^q \cdot \rho_{a,b} = -(pb - qa + p - q) \rho_{a-p+1, b-q+1}$  gives the two displayed formulas with the principal-part shifts  $b+1$  and  $a+1$ .

For local nilpotence:  $z_1$  lowers the second index of  $\widetilde{\Psi}_{a,b}$  by one and is killed by  $\widetilde{\Psi}_{a,0}$  for  $a \geq 0$ ; a finite linear combination is exhausted after at most  $\max b + 1$  iterations. By contrast,  $z_1$  raises the second index of  $\rho_{a,b}$  and produces a nonzero rising-factorial coefficient  $(-1)^N (b+1)(b+2) \cdots (b+N) \rho_{a,b+N}$ , which is never zero. The same dichotomy holds for  $z_2$ . Local nilpotence is an isomorphism invariant; therefore the displayed  $\mathfrak{h}$ -actions have a module-isomorphism obstruction.

The obstruction to extending  $\text{gr } \Phi$  to an  $\hbar$ -adic same-action map is exactly this local-nilpotence dichotomy, and is recorded as Theorem 8.11 together with the deformation-level Corollary 8.13. Theorem 8.12 supplies the positive Fourier–Rees bridge that does identify the two label sets, but only after the Fourier twist of the Hamiltonian action.  $\square$

*Remark D.16 (Smallest explicit mismatch:  $(p, q, a, b) = (4, 0, 1, 1)$ ).* The polynomial PBW action on the descendant labels and the Moyal coadjoint action on the principal parts disagree already at one exponent pair recorded by the verification script. Take the quartic Hamiltonian  $F_{4,0} = \text{Tr}(\phi_1^4)$  and the bidegree- $(1, 1)$  descendant target. On the polynomial PBW side Proposition 5.171 gives

$$F_{4,0} : \widetilde{\Psi}_{1,1} \longmapsto (4 \cdot 1 - 0 \cdot 1) \widetilde{\Psi}_{1+4-1,1-1} = 4 \widetilde{\Psi}_{4,0}.$$

Direct calculation in the Moyal coadjoint module gives

$$\text{ad}_{z_1^4}^*(\delta_{1,1}) = -24 \hbar^2 \delta_{0,4},$$

the only nonzero term of order  $\leq 3$  in the formal Moyal expansion. Fourier-untwisted relabelling has the bidegree obstruction  $(1, 1) \rightarrow (0, 4)$  versus  $(1, 1) \rightarrow (4, 0)$ , so the polynomial PBW action and the Moyal coadjoint action remain distinct. Replacing the Moyal coadjoint by its classical truncation ( $\hbar^0$ ) gives  $\text{ad}_{z_1^4}^*(\rho_{1,1}) = 0$  by the principal-part formula at this exponent, again disagreeing with the polynomial PBW output  $4 \widetilde{\Psi}_{4,0}$ . The script confirms this fingerprint exactly and embeds it inside a sweep of 1200 such mismatches.

**THEOREM D.17 (Universal primitive descendant calculation).** Let  $\mathsf{K}_{\text{univ}}$  be the stable primitive cyclic Koszul complex generated by two even letters  $x, y$  and one odd letter  $\psi$ , with  $d\psi = xy - yx$ , decomposed by the weights of Definition D.1. Then

$$H_\bullet(\mathsf{K}_{\text{univ}}) \cong \mathbb{C}[x, y] \oplus \mathbb{C}[x, y]\theta, \quad |\theta| = 1, \quad \text{wt}(\theta) = (1, 1),$$

with  $x^a y^b \theta$  represented by the symmetrised primitive cycle  $\Psi_{a,b}$ . This is the universal primitive calculation in the following precise sense. If  $E_\bullet$  is any stable primitive single-trace model whose generators are the images of  $x, y, \psi$ , whose differential sends  $\psi$  to  $xy - yx$ , and whose only stable primitive trace relation is graded cyclicity, then the induced map

$$\mathsf{K}_{\text{univ}} \longrightarrow E_\bullet$$

factors on homology through the displayed two-line module. Hence the universal primitive higher- $\psi$  classes are exactly the displayed degree-zero polynomial line and degree-one  $\Psi$ -line.

*Proof.* The first assertion is Theorem D.12 summed over all bidegrees  $(R, S)$ , with  $x^a y^b \theta = \Psi_{a,b}$  and  $R = a + 1, S = b + 1$  in degree one. For the universal property, Lemma D.2 says that the stable primitive trace span is exactly the graded cyclic quotient of the free algebra on  $x, y, \psi$ . Therefore any stable primitive model satisfying the stated hypotheses receives a unique chain map from  $\mathsf{K}_{\text{univ}}$  after choosing the

images of  $x, y, \psi$ . The homology of the source is the two-line module just computed. Since  $H_p(K_{R,S}) = 0$  for  $p \geq 2$  in every bidegree, every universal primitive class in the target comes from the degree-zero polynomial line or the degree-one  $\Psi$ -line. The statement is independent of the later choice of principal-part, PBW, or Moyal coefficient module.  $\square$

**THEOREM D.18** (*Hamiltonian-sector extraction*). The full primitive homology decomposes as

$$\mathbb{C} \cdot [1] \oplus \tilde{A} \oplus \mathbb{C} \cdot [\Psi_{0,0}] \oplus \tilde{A}_{\text{desc}}[1],$$

where

$$\tilde{A} = \bigoplus_{R+S>0} \mathbb{C} \cdot [x^R y^S], \quad \tilde{A}_{\text{desc}}[1] = \bigoplus_{a+b>0} \mathbb{C} \cdot [\Psi_{a,b}].$$

The Hamiltonian comparison uses only  $\tilde{A} \oplus \tilde{A}_{\text{desc}}[1]$ , and recognises this latter pair as the special-fibre,  $\hbar = 0$ , shadow of the Rees PBW lattice of Construction D.14: the closed Hamiltonian summand is the polynomial part  $\mathbf{S}(\mathfrak{h})|_{\hbar=0}$ , and the descendant copy is the primitive one- $\psi$  PBW label sector  $\mathbf{S}^1(\mathfrak{h}^\vee[1]) \otimes \mathbf{S}(\mathfrak{h})|_{\hbar=0}$ . The complementary sectors are exactly the scalar unit  $[1]$ , the scalar odd primitive  $\Psi_{0,0}$ , and the vanishing higher- $\psi$  primitive groups  $H_p(K_{R,S}) = 0$  for  $p \geq 2$ .

*Proof.* The constant Hamiltonian has zero Hamiltonian vector field, so the reduced Hamiltonian algebra is  $\tilde{A} = \mathbb{C}[x, y]/\mathbb{C} \cdot 1$  degree by degree. This removes the degree-zero scalar class  $[1]$ . The odd scalar  $\Psi_{0,0}$  is the one- $\psi$  descendant of the same removed constant direction; it is a stable primitive trace class in the removed scalar sector. Theorem D.12 proves that primitive homology is concentrated in  $\psi$ -degrees 0 and 1. What remains is exactly the positive-degree Hamiltonian and one- $\psi$  PBW label sector displayed above. The identification with the special-fibre Rees lattice is Construction D.14: the closed Hamiltonian summand is the polynomial special fibre of the Rees algebra of  $\mathfrak{h}$ , and the one- $\psi$  descendant copy is the bidegree-graded  $\tilde{\Psi}_{a,b}$  sector of the cotangent-shifted  $\mathfrak{h}^\vee[1]$  tensor. This is a vector-space homology statement. The Matlis principal-part module comparison is the separate  $\mathfrak{h}$ -module obstruction.  $\square$

**COROLLARY D.19** (*Special-fibre defect polarization*). The polynomial one- $\psi$  PBW labels  $\Psi_{a,b}$  and the Matlis principal parts  $\rho_{a,b}$  are joined by the bidegree-only associated-graded morphism  $\text{gr } \Phi$  of Proposition D.15: this morphism intertwines the two index sets at the special fibre  $\hbar = 0$  and only there. In particular, the full primitive  $\psi$ -homology of this appendix computes the special-fibre shadow of the cotangent module on the open brane sector, while the Matlis principal-parts module of § 8 is the deformation-level cotangent module itself. The formal-stalk chart measures positions polynomially through  $\tilde{A}$ , and its cotangent momenta live as residues through  $\mathcal{P}$ ; the shared bidegree label is associated-graded data.

*Proof.* Construction D.14 identifies  $\tilde{\Psi}_{a,b}$  with the unit-normalised  $\hbar = 0$  class in  $\mathbf{S}^1(\mathfrak{h}^\vee[1]) \otimes \mathbf{S}(\mathfrak{h})|_{\hbar=0}$  in bidegree  $(a, b)$ . The bidegree-only relabelling  $\text{gr } \Phi$  of Proposition D.15 is an isomorphism of bigraded vector spaces with the  $\mathfrak{h}$ -module obstruction of Theorem 8.11. The Matlis side is the continuous coadjoint module identified by Theorem 8.10 as the unique perfect continuous coadjoint pairing. Therefore  $\tilde{A}_{\text{desc}}[1]$  is the special-fibre shadow and  $\mathcal{P}$  is the deformation-level cotangent module, with the shared index set carrying associated-graded label data only. This is the separation displayed by Theorem 5.180.  $\square$

## E UNREDUCED BV COTANGENT, SCALAR QME OBSTRUCTIONS, AND CANCELLATIONS

Fix the formal-local mixed geometry  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}^2_{\circ}$  with  $\Omega_{\text{loc}} = dz_1 \wedge dz_2$ . On a brane interval  $I \subset L$  the open BV theory contributes scalar contact counterterms each order in  $\hbar$ ; the residual non-scalar Costello quantum master equation lives in their orthogonal complement. Naming the players:  $\mathfrak{Q}_{w,\partial,\hbar}^{\bullet}(I)$  is the local-functional cochain complex on  $I$  at weighted-Tate coefficient weight  $w$ , with the brane-tangential differential  $d_{\partial}$  and the  $\hbar$ -formal grading;  $\widehat{\sigma}_{\text{sc}}$  is the filtered scalar-symbol chain projection killing the scalar-contact filtration; and  $d_{\text{QME}} = Q + \{I_{\circ}, -\} + \hbar\Delta$  is the quantum master differential of Costello's renormalization. Once these are in place, the residual non-scalar QME obstruction is a single cohomology class

$$[\text{Ob}_{w,\partial,\hbar}^{\text{red}}] \in H^1(\ker \widehat{\sigma}_{\text{sc}}, d_{\text{QME}}),$$

paired with its finite-window  $\varprojlim^1 H^0$ -primitive-compatibility class (the secondary Roos cohomology obstruction tracking compatibility of window-by-window primitives across the truncation tower). The scalar projection at the  $U(1)$  centre-of-mass slice has explicit value

$$\widehat{\sigma}_{\text{sc}}([\text{Ob}_{w,\partial,\hbar}^{\text{red}}]) = \hbar N[\bar{c}] = \Omega_{\text{central}}|_{J(f)},$$

the Capelli class of Chapter 6 (Theorem 6.8); equivalently, the scalar projection of the non-scalar Costello QME residual is the projective curvature of the modular line bundle evaluated at the trace generator  $J(f) = \text{Tr } f(\phi_1, \phi_2)$ , with sign and convention fixed by Appendix B and the canonical scheme  $z_1 z_2 \mapsto \frac{1}{2}(\phi_1 \phi_2 + \phi_2 \phi_1)$  of the Moyal star product. Verification of the Moyal coefficient  $\hbar N$  is supplied by `scripts/check_moyal_coefficients.py`.

The scalar anomaly is one cohomology class realized in four cohomologies. The algebraic central extension of  $\tilde{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1$ , the Lie central extension  $\widehat{\mathfrak{h}}_{\text{cent}}$ , the trace-algebraic boundary central extension, and the quantum Weyl commutator  $[\Phi_{\hbar}(z_1), \Phi_{\hbar}(z_2)]_{\star} = \hbar N$  carry the same class in  $H^2(\tilde{A}; \mathbb{C})[[\hbar]]$  under the canonical scalar projection; the cocycle  $\omega(f, g) = [\{f, g\}]_{\circ}$  is the same in each (Theorem E.13). Slogan: *constants vanish as Hamiltonian vector fields but survive as brane center-of-mass charge*. Each cancellation branch acts on this single class — ordinary  $\mathfrak{gl}_N$  is obstructed by  $\hbar N[\bar{c}]$ , the unreduced central character contributes the CE coboundary  $d_{\text{CE}}(\hbar \chi_N) = -\hbar N \omega$ , the balanced  $\mathbb{C}^{N|N}$  supertrace rescales the open trace by  $\text{sdim } \mathbb{C}^{N|N} = 0$ , and full-equivariant marker tensors satisfy  $\text{Str}_{\text{cyc}}(\mathfrak{m}) = 0$ . Modifying one branch displaces the same class.

At each finite window  $M$  (a Hamiltonian-degree truncation of the weighted Tate coefficient pair) the obstruction is the linear-algebra problem

$$A^M c^M = -r^M$$

of Theorem E.35, where  $A^M$  is the finite matrix whose columns enumerate the candidate counterterms supplied by the row span,  $c^M$  is the unknown counterterm vector, and  $r^M$  is the residual vector collecting the QME source faces. The row entries are graph-by-graph Costello data:  $K_{\Gamma}^M$  the windowed kernel of the Feynman graph  $\Gamma$ ,  $\varepsilon_{\Gamma}$  its sign,  $\zeta_{M,\nu}$  a CE test pairing labelled by the cycle  $\nu$ , and the bracket  $\{-, -\}_{\hbar, M}$  the windowed Schouten/BV bracket. In these terms,

$$R_{d\Gamma, M}^{\text{row}} = \varepsilon_{\Gamma} d_M K_{\Gamma}^M, \quad R_{\text{CE}(\Gamma, \nu), M}^{\text{row}} = -\varepsilon_{\Gamma, \nu} K_{\Gamma}^M(\zeta_{M, \nu}),$$

$$R_{b(\Gamma_1, \Gamma_2), M}^{\text{row}} = \frac{1}{2} \varepsilon_{\Gamma_1, \Gamma_2} \{K_{\Gamma_1}^M, K_{\Gamma_2}^M\}_{\hbar, M},$$

and the order- $n$  homogeneous lower-counterterm rows. A degree-zero counterterm  $c^M$  exists in the supplied row span exactly when the system is consistent; otherwise a separating covector  $\ell A^M = 0$ ,  $\ell(r^M) \neq 0$  detects the obstruction. Tower compatibility passes through truncation matrices  $T^{M,N}$  on rows (restriction-of-graphs from window  $M$  to a smaller window  $N$ ) and  $P^{M,N}$  on primitive candidates (the matched counterterm projector) and the Roos class  $[P^{M,N}c^M - c^N] \in H^0(\mathcal{K}_{\text{ns},N}^\bullet(I))$  (the cohomology class measuring compatibility of windowwise primitives across truncation).

The order-three row  $\theta_3 = \text{CE}(\Theta_3, \nu_3)$  is the first scalar-zero source-face row beyond the order-one one-cross scalar contact. It has

$$R_{\theta_3, M}^{\text{row}} = -K_{\Theta_3}^M(\zeta_{M, \nu_3}) =: E_{\theta_3, M}, \quad S_{\theta_3, M} = 0,$$

and in the one-row complex  $\mathcal{K}_{\theta_3, M}^0 = 0$ ,  $\mathcal{K}_{\theta_3, M}^1 = \mathbb{C}E_{\theta_3, M}$ ,  $d_{\theta_3, M} = 0$ , the covector  $\ell_{\theta_3}(E_{\theta_3, M}) = 1$  is an exact obstruction (Proposition E.36). A named coefficient  $C_{\theta_3}$  requires one of three data: a CE ancestor with controlled boundary; a scalar-zero Costello local counterterm  $C_{\theta_3, M}$  with  $d_{\text{ns}, M}C_{\theta_3, M} = -E_{\theta_3, M}$  and truncation-compatible primitives; or a complete companion-face table of Hom-valued companion rows whose signed sum moves the residual vector into the image of  $A^M$ . Tower compatibility passes through the truncation defect  $\Delta_{M, N}^1 = -\pi_{M, N}\mathbf{b}^M + \mathbf{b}^N$  ( $\mathbf{b}^M$  the order-one Bianchi residual at window  $M$ ,  $\pi_{M, N}$  the truncation projector) and the secondary  $\varprojlim^1 H^0$  primitive class (Proposition E.37). The unreduced factorization-center cotangent lift is governed by a four-component obstruction vector, and lifts if and only if that vector vanishes (Theorem E.53).

*Definition E.1 (BV degrees and signs).* The open polynomial BV/Koszul model uses

$$|\phi_1| = |\phi_2| = 0, \quad |\psi = A^*| = -1, \quad Q\psi = [\phi_1, \phi_2], \quad |Q| = 1.$$

A Hamiltonian generator  $H_f$  has degree 0. A shifted cotangent generator  $\eta_\rho = \rho[1]$  is odd of BV degree  $-1$ . The CE coordinates satisfy

$$|c_f| = 1, \quad |u_\rho| = 0.$$

This is the cotangent-coordinate grading of the CE/PV block. The ordinary graded dual of the BV degree of  $\rho[1]$  is a separate convention. The CE/PV cotangent–Schouten bracket has degree  $-1$ . The reduced Hamiltonian boundary  $P_0$  bracket has degree 0. The QME obstruction differential  $Q + \{I_0, -\}$  has degree 1. The Poisson sign convention is the one determined by  $\Omega_{\text{loc}} = dz_1 \wedge dz_2$ , so  $\{z_1, z_2\} = 1$ . The principal-part coadjoint action is

$$z_1^p z_2^q \cdot \rho_{a, b} = -(pb - qa + p - q)\rho_{a-p+1, b-q+1}.$$

**PROPOSITION E.2 (Local geometry of scalar shadows).** All scalar-contact, finite-window, marked-graph, and supertrace calculations below use only the formal-local mixed holomorphic-topological geometry

$$I \subset \mathbb{R}_{\text{brane}} \subset \mathbb{R}_{\text{top}}^2, \quad \widehat{\mathbb{C}}^2_0 = \text{Spf } \mathbb{C}[[z_1, z_2]], \quad \Omega_{\text{loc}} = dz_1 \wedge dz_2.$$

The inverse Poisson bracket is

$$\{f, g\}_{\Omega_{\text{loc}}} = \partial_{z_1} f \partial_{z_2} g - \partial_{z_2} f \partial_{z_1} g.$$

With the convention  $\iota_{X_f} \Omega_{\text{loc}} = -df$ , one has

$$X_f = (\partial_{z_1} f) \partial_{z_2} - (\partial_{z_2} f) \partial_{z_1}, \quad \text{div}_{\Omega_{\text{loc}}} X_f = 0.$$

Thus the scalar shadow has zero divergence and zero volume-form anomaly. The Hamiltonian scalar cocycle entering every calculation below is

$$\omega(f, g) = [\{f, g\}_{\Omega_{\text{loc}}}]_0, \quad \omega(z_1, z_2) = 1.$$

The open index contraction then multiplies this local Hamiltonian scalar by the Chan–Paton trace coefficient:

$$\begin{aligned} \hbar N \omega(f, g) \gamma_1 & \text{ in the ordinary } \mathfrak{gl}_N \text{ branch,} \\ \hbar \text{sdim}(\mathbb{C}^{N|N}) \omega(f, g) \gamma_1 & = 0 \text{ in the balanced supertrace branch,} \end{aligned}$$

and, for a marked scalar loop,

$$\hbar \text{Str}_{\text{cyc}}(\mathbf{m}) \omega(f, g) \gamma_1.$$

A finite window  $M$  restricts only the polynomial Hamiltonian labels, graph labels, and counterterm table to a finite local coefficient system; it preserves  $\Omega_{\text{loc}}$ , the Hamiltonian bracket, and the divergence-free condition.

*Proof.* The formula for  $X_f$  is obtained by solving  $\iota_{X_f}(dz_1 \wedge dz_2) = -df$ . Its divergence is

$$\partial_{z_1}(-\partial_{z_2}f) + \partial_{z_2}(\partial_{z_1}f) = 0$$

after writing  $X_f = -(\partial_{z_2}f)\partial_{z_1} + (\partial_{z_1}f)\partial_{z_2}$ . Therefore the local volume form is preserved and contributes zero additional scalar contact term. Taking the constant Taylor coefficient of the displayed Poisson bracket gives the cocycle  $\omega$ , with  $\omega(z_1, z_2) = 1$ .

The remaining scalar factor is purely the open Chan–Paton index loop. The ordinary trace of the identity on  $\mathbb{C}^N$  is  $N$ . The supertrace of the identity on  $\mathbb{C}^{N|N}$  is  $N - N = 0$ . A marked loop replaces the identity by the cyclic product of its marker tensors and gives the cyclic supertrace shown above. Finite-window truncation is a restriction of labels and graph tables inside this same formal polydisk, so it preserves the local cocycle and the divergence computation.  $\square$

*Definition E.3 (Scalar-reduced brane-defect obstruction complex).* Let  $O_{\text{loc}}^{\text{bd}}$  denote local functionals on the ordinary  $\mathfrak{gl}_N$  brane-defect fields in the formal-local model  $\mathbb{R}_{\text{top}}^2 \times \mathbb{C}^2_0$ , with compact support on the brane line and with coefficients in the scalar-reduced Hamiltonian source  $\tilde{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1$ . The differential controlling a mixed brane-defect QME lift is

$$d_{\text{QME}} = Q + \{I_0, -\}.$$

A scalar contact term is tested by adjoining the degree-one central ghost  $\gamma_1$  for the omitted constant Hamiltonian line. Its associated graded is the Chevalley–Eilenberg complex of  $\tilde{A}$  with trivial coefficients on this central line. Writing

$$\mathfrak{Q}_{\text{bd}}^\bullet = (O_{\text{loc}}^{\text{bd}}, d_{\text{QME}}),$$

the associated-graded scalar-symbol projection is

$$\sigma_{\text{sc}}: \text{gr}_{\text{sc}} \mathfrak{Q}_{\text{bd}}^\bullet \longrightarrow C_{\text{Lie}}^\bullet(\tilde{A}; \mathbb{C}\gamma_1)[1].$$

A cone or long exact sequence is formed only after one has chosen a  $d_{\text{QME}}$ -stable scalar filtration and a filtered chain projection

$$\widehat{\sigma}_{\text{sc}}: \mathfrak{Q}_{\text{bd}}^\bullet \longrightarrow C_{\text{Lie}}^\bullet(\tilde{A}; \mathbb{C}\gamma_1)[1]$$

whose associated graded is the displayed  $\sigma_{\text{sc}}$ . The assertion below is an associated-graded scalar obstruction statement until this extra projection is supplied. Thus a degree-one QME representative with two Hamiltonian inputs maps to a Lie two-cocycle. Explicitly,

$$\sigma_{\text{sc}} \left( \kappa(f, g) \int_{\mathbb{R}} a(t) b(t) \gamma_{\mathbf{I}}(t) dt \right) = \kappa(f, g) \gamma_{\mathbf{I}}.$$

LEMMA E.4 (*Filtered scalar projection obstruction*). Fix a complete decreasing scalar-contact filtration  $F^\bullet \mathfrak{Q}_{\text{bd}}^\bullet$  preserved by  $d_{\text{QME}}$ , and put the target  $C_{\text{Lie}}^\bullet(\bar{A}; \mathbb{C}\gamma_{\mathbf{I}})[1]$  in filtration degree 0. Suppose the associated graded carries the scalar-symbol projection

$$\sigma_{\text{sc}}: \text{gr}_F \mathfrak{Q}_{\text{bd}}^\bullet \longrightarrow C_{\text{Lie}}^\bullet(\bar{A}; \mathbb{C}\gamma_{\mathbf{I}})[1].$$

Choose any filtered linear lift  $s: \mathfrak{Q}_{\text{bd}}^\bullet \rightarrow C_{\text{Lie}}^\bullet(\bar{A}; \mathbb{C}\gamma_{\mathbf{I}})[1]$  of  $\sigma_{\text{sc}}$ . Its chain-defect

$$\partial_s = d_{\text{CE}} s - s d_{\text{QME}}$$

lowers the scalar-contact filtration. The first nonzero homogeneous piece defines an obstruction class

$$o_\sigma^{(r)} \in H^1(\text{Hom}(\text{gr}_F \mathfrak{Q}_{\text{bd}}^\bullet, C_{\text{Lie}}^\bullet(\bar{A}; \mathbb{C}\gamma_{\mathbf{I}})[1])_{-r}),$$

with the usual differential  $D(\alpha) = d_{\text{CE}} \alpha - (-1)^{|\alpha|} \alpha d_{\text{gr}}$ . A filtered chain projection  $\widehat{\sigma}_{\text{sc}}$  with associated graded  $\sigma_{\text{sc}}$  exists exactly when the obstruction classes  $o_\sigma^{(r)}$  vanish successively for all  $r > 0$ , with compatible choices in the completed inverse limit.

*Proof.* Since the associated graded of  $s$  is already a chain map, the commutator  $d_{\text{CE}} s - s d_{\text{QME}}$  strictly lowers the filtration. Its lowest nonzero component is closed for the Hom differential because  $d_{\text{CE}}^2 = d_{\text{QME}}^2 = 0$ . Replacing  $s$  by  $s - \alpha$ , where  $\alpha$  lowers filtration by the same amount, changes this component by  $D\alpha$ . Thus the first defect can be removed exactly when its cohomology class vanishes. Iterating the same argument through the complete filtration gives a compatible sequence of corrections if and only if all successive classes vanish.  $\square$

PROPOSITION E.5 (*First scalar-lift obstruction in the ordinary branch*). Work in the ordinary scalar-reduced  $\mathfrak{gl}_N$  weighted brane-defect local-functional complex  $\mathfrak{Q}_{w, \partial, \hbar}^\bullet(I)$ , with  $N > 0$ . Let  $J_{\text{sc}}$  be the closed ideal of functionals whose associated-graded scalar-contact symbol vanishes, and put  $F^r = J_{\text{sc}}^r$ , completed  $\hbar$ -adically and in the finite-window inverse limit. For any filtered linear lift  $s$  of the scalar-symbol projection, the first chain-defect class has Hamiltonian two-input evaluation

$$\text{ev}_{f, g}(o_{\sigma, w}^{(1)}) = \hbar N \omega(f, g) \gamma_{\mathbf{I}} \in H_{\text{Lie}}^2(\bar{A}; \mathbb{C}\gamma_{\mathbf{I}})[[\hbar]].$$

In particular

$$\text{ev}_{z_1, z_2}(o_{\sigma, w}^{(1)}) = \hbar N \gamma_{\mathbf{I}} \neq 0.$$

Hence a filtered chain lift of the scalar-symbol projection in the ordinary scalar-reduced branch requires additional data whose leading scalar class is  $-\hbar N [\bar{c}]$ , or a replacement of the source or open trace model before scalar reduction.



*Proof.* The filtration  $F^\bullet$  is the naive scalar-contact filtration: its associated graded keeps the leading scalar contact term and discards higher scalar-contact corrections. The first test is the bracket of the two linear Hamiltonian boundary insertions. By Theorem E.12,

$$\{I_\partial(a, f), I_\partial(b, g)\}_{\text{open}, \hbar} - I_\partial(ab, \{f, g\}_{\tilde{A}}) = \hbar N \omega(f, g) \int_I a(t) b(t) dt$$

after insertion of  $\gamma_1$ . Applying the scalar-symbol lift gives the displayed representative of  $d_{\text{CES}} - sd_{\text{QME}}$ . For  $f = z_1$  and  $g = z_2$ ,  $\omega(z_1, z_2) = 1$ .

If this first defect were a Hom coboundary, evaluation on the Hamiltonian two-input subcomplex would give a linear primitive  $\tilde{\eta} : \tilde{A} \rightarrow \mathbb{C}\gamma_1$  with  $d_{\text{CE}}\tilde{\eta} = \hbar N \omega\gamma_1$ . Pulling  $\tilde{\eta}$  back to  $A = \mathbb{C}[z_1, z_2]$  and forcing it to vanish on the constant line contradicts Lemma E.11, equivalently  $\omega(z_1, z_2) = 1$  while  $\{z_1, z_2\} = 1$  is zero in  $\tilde{A}$ . Thus  $\sigma_{\sigma, w}^{(1)}$  is nonzero.

The obstruction is detected in the finite window containing  $z_1$  and  $z_2$ . The admissible weight transports of Theorem 5.81 identify this finite-window class, so changing  $w$  preserves its vanishing criterion. The primitive exists on the unreduced source  $A$ , where  $\eta(f) = -[f]_0$ , and the central-character replacement uses exactly this primitive. In the balanced supertrace branch the displayed ordinary-trace evaluation is multiplied by  $\text{sdim } \mathbb{C}^{N|N} = 0$ . The scalar-contact subcomplex and chain projection constructed in Proposition E.18 kill this obstruction on the balanced scalar-contact branch. Extension to all local functionals remains the separate non-scalar QME obstruction problem.  $\square$

**THEOREM E.6** (*Polynomial unreduced centrality obstruction*). Let  $M_{\text{pol}}$  be any  $Q$ -stable subcomplex of the unreduced polynomial open BV observables in which every element has finite polynomial degree in  $\phi_1, \phi_2, \psi$ . Any assignment

$$\rho_{a,b}[1] \mapsto \Theta_{a,b} \in H^\bullet(M_{\text{pol}}, Q)$$

together with centrality homotopies in  $M_{\text{pol}}$  has locally nilpotent linear Hamiltonian action on  $Q$ -cohomology, whereas the principal-part module  $\mathcal{P}$  has non-locally-nilpotent linear Hamiltonian action. Equivalently, an unreduced polynomial lift of the reduced map  $\rho_{a,b}[1] \mapsto \Theta_{a,b}$  would require polynomial boundary representatives with homotopies satisfying

$$\{O_f, \Theta_\rho\} \simeq \Theta_{f \cdot \rho}, \quad \{\Theta_\rho, \Theta_\sigma\} \simeq 0$$

for all polynomial Hamiltonians  $f$ .

*Proof.* The linear Hamiltonians act on polynomial open observables by the canonical scalar-translation derivations

$$O_{1,0} \mapsto \partial_{\phi_2}, \quad O_{0,1} \mapsto -\partial_{\phi_1}.$$

These derivations commute with  $Q$ , since translating  $\phi_1$  or  $\phi_2$  by a scalar matrix leaves the moment-map equation  $Q\psi = [\phi_1, \phi_2]$  unchanged. On every finite polynomial observable, repeated application of either derivation eventually vanishes. Hence their induced actions on  $H^\bullet(M_{\text{pol}}, Q)$  are locally nilpotent.

On the principal-part module the same two Hamiltonians act by

$$z_1 \cdot \rho_{a,b} = -(b+1)\rho_{a,b+1}, \quad z_2 \cdot \rho_{a,b} = (a+1)\rho_{a+1,b}.$$

Iterating gives

$$z_1^r \cdot \rho_{a,b} = (-1)^r (b+1) \cdots (b+r) \rho_{a,b+r} \neq 0,$$

and similarly for  $z_2$ , for every  $r \geq 1$ . Thus every nonzero principal part has non-locally-nilpotent linear Hamiltonian orbit.

If the stated homotopies existed in  $M_{\text{pol}}$ , then after passing to  $Q$ -cohomology the action of  $O_{1,0}$  and  $O_{0,1}$  on the classes  $[\Theta_{a,b}]$  would be isomorphic to the displayed principal-part action. Local nilpotence is invariant under module isomorphism and under passage to a submodule. This is incompatible with the non-locally-nilpotent principal-part action above. Therefore the unreduced lift requires a category beyond polynomial open BV/Koszul observables.  $\square$

LEMMA E.7 (*Distributional locality of principal-part currents*). Let  $I \subset \mathbb{R}$ ,  $a \in C_c^\infty(I)$ , and  $B \in \mathcal{D}'_c(I)$ . The product

$$(aB)(\varphi) = B(a\varphi), \quad \varphi \in C_c^\infty(I),$$

is a compactly supported distribution with  $\text{supp}(aB) \subset \text{supp}(a) \cap \text{supp}(B)$ . Consequently the reduced principal-part current brackets

$$\{B_f(a), \Theta_\rho(B)\} = \Theta_{f \cdot \rho}(aB), \quad \{\Theta_\rho(B), \Theta_\sigma(C)\} = 0$$

are distributionally local on the brane line. If the supports are disjoint, the brackets vanish.

*Proof.* Multiplication of a compactly supported distribution by a smooth function is the standard operation  $B \mapsto aB$  defined by precomposition with multiplication by  $a$ . If  $\varphi$  is supported away from  $\text{supp}(a) \cap \text{supp}(B)$ , then  $a\varphi$  is supported away from  $\text{supp}(B)$ , hence  $B(a\varphi) = 0$ . This proves the support assertion. The first bracket is the negative transpose of the Hamiltonian action under the residue-current pairing, so its interval factor is exactly  $aB$ . The second bracket is zero because the principal-part current ideal is abelian. Disjoint-support vanishing follows from  $aB = 0$ .  $\square$

Definition E.8 (*Central-character principal-part current algebra*). Let  $A = \mathbb{C}[z_1, z_2]$ ,  $\bar{A} = A/\mathbb{C} \cdot 1$ , and let  $\chi: A \rightarrow \mathbb{C}$  be a linear central character on the constant Hamiltonian line. For an interval  $I \subset \mathbb{R}$ , write

$$\tilde{\mathfrak{h}}_I = \Omega_c^\bullet(I) \widehat{\otimes} A, \quad \mathcal{P}_I = \mathcal{D}'_c(I) \widehat{\otimes} \mathcal{P}.$$

The algebraic current target with central character is the completed graded-commutative algebra generated by symbols

$$B_f^\chi(a), \quad \Theta_\rho(B), \quad a \in \Omega_c^\circ(I), \quad f \in A, \quad B \in \mathcal{D}'_c(I), \quad \rho \in \mathcal{P},$$

modulo

$$B_{f+c}^\chi(a) = B_f^\chi(a) + \chi(c \cdot 1) \int_I a(t) dt, \quad \Theta_{\rho+\sigma}(B) = \Theta_\rho(B) + \Theta_\sigma(B),$$

and with  $P_0$  brackets

$$\begin{aligned} \{B_f^\chi(a), B_g^\chi(b)\} &= B_{\{f,g\}}^\chi(ab), \\ \{B_f^\chi(a), \Theta_\rho(B)\} &= \Theta_{f \cdot \rho}(aB), \\ \{\Theta_\rho(B), \Theta_\sigma(C)\} &= 0. \end{aligned}$$

Constants act trivially on  $\mathcal{P}$ . The product is the ordinary graded-commutative factorization product after extension by zero on intervals. This is an algebraic current model. The finite polynomial open BV/Koszul complex is governed by Theorem E.6.

*Construction E.9 (Algebraic current presentation map).* Let

$$\widetilde{\mathfrak{g}}_I = \widetilde{\mathfrak{h}}_I \ltimes \mathcal{P}_I[1]$$

with bracket induced by the Hamiltonian bracket on  $A$ , the coadjoint action on  $\mathcal{P}$ , and the abelian bracket on  $\mathcal{P}_I[1]$ . In the CE algebra  $C_{\text{CE,cont}}^\bullet(\widetilde{\mathfrak{g}}_I)$ , write

$$c_{B,\rho}$$

for the degree-one CE coordinate dual to Hamiltonian currents through the pairing

$$\langle B \otimes \rho, a(t)dt \otimes f \rangle = B(a)\rho(f),$$

and write

$$u_{a(t)dt \otimes f}$$

for the degree-zero CE coordinate dual to the shifted cotangent generator under the same pairing with  $a(t)dt \otimes f \in \widetilde{\mathfrak{h}}_I$ . Define

$$\Phi_I^\chi(c_{B,\rho}) = \Theta_\rho(B), \quad \Phi_I^\chi(u_{a(t)dt \otimes f}) = B_f^\chi(a),$$

and extend continuously and multiplicatively.

**THEOREM E.10 (Algebraic current presentation with zero homotopies).** The maps  $\Phi_I^\chi$  of Construction E.9 define a morphism of  $P_o$  factorization algebras from the interval CE source of  $\widetilde{\mathfrak{g}}_I$  to the algebraic current target of Definition E.8. The factorization centrality homotopies and the  $P_o$ -bracket compatibility homotopies in this algebraic model are strict:

$$H_{\text{fact}} = 0, \quad H_{P_o} = 0.$$

Thus the unreduced source with central character constructs the  $P_o$  current presentation as far as the distributional algebraic current model permits. A morphism into  $Z_{\text{fact}}^{P_o}(\text{Obs}_{\text{open}})$  for the genuine unreduced open BV factorization algebra requires the additional unreduced graph/QME data. This is compatible with the polynomial obstruction theorem: the image of  $\Theta_\rho(B)$  lives in the added principal-part current sector.

*Proof.* The CE differential of  $C_{\text{CE,cont}}^\bullet(\widetilde{\mathfrak{g}}_I)$  is dual to exactly three brackets:

$$[\widetilde{\mathfrak{h}}_I, \widetilde{\mathfrak{h}}_I], \quad \widetilde{\mathfrak{h}}_I \curvearrowright \mathcal{P}_I, \quad [\mathcal{P}_I[1], \mathcal{P}_I[1]] = 0.$$

Under  $\Phi_I^\chi$ , these three structure maps become the three displayed  $P_o$  brackets of Definition E.8. Hence the CE differential and the target Schouten differential agree on generators, while the degree-zero current  $P_o$  brackets agree by the same source brackets. The biderivation property extends the equality to the completed symmetric algebra.

Factorization compatibility is also strict. Extension by zero sends  $a, B$  to their extensions as test functions and currents; the products  $ab$  and  $aB$  are computed after extension. If the supports are disjoint these products vanish, so mixed brackets vanish before any homotopy is introduced. Multiplication in the target is graded-commutative, hence multiplication by an image generator is a Hochschild central operation with zero product-centrality homotopy. The  $P_o$ -centrality defect on generators is the difference

between the target bracket and the image of the corresponding source bracket; it is zero by the previous paragraph. Therefore  $H_{\text{fact}}$  and  $H_{P_0}$  may both be chosen to be zero.

The same statement holds against arbitrary monomials in the added algebraic current target, including monomials beyond length-one generators. Let  $Y = Y_1 \cdots Y_r$  be a homogeneous source monomial and put  $W = \Phi_I^\chi(Y)$ , a monomial in the target generators  $B_g^\chi(b)$  and  $\Theta_\sigma(C)$ . For a source generator  $x$ , the product-centrality defect is zero because the target product is graded-commutative:

$$\Phi_I^\chi(x)W - (-1)^{|\Phi_I^\chi(x)||W|}W\Phi_I^\chi(x) = 0.$$

Let  $\partial_x$  denote the derivation of the source symmetric algebra induced by the corresponding bracket in  $\widetilde{\mathfrak{g}}_I$ . The bracket-compatibility defect

$$D_x(Y) = \{\Phi_I^\chi(x), \Phi_I^\chi(Y)\} - \Phi_I^\chi(\partial_x Y)$$

is a graded derivation in  $Y$ . It vanishes when  $\Phi_I^\chi(Y)$  is  $B_g^\chi(b)$  or  $\Theta_\sigma(C)$  by the three displayed brackets. Hence the Leibniz rule gives  $D_x(Y) = 0$  for every monomial and, by continuity, for every element of the completed algebra. Thus the centrality homotopies against arbitrary algebraic current-target monomials are strict zero homotopies in this model. Here “arbitrary” means arbitrary monomial in the freely adjoined algebraic current target. The genuine unreduced open BV observable factorization algebra and the unreduced  $Z_{\text{fact}}^{P_0}$ -morphism require the additional graph/QME data.  $\square$

LEMMA E.11 (*Unreduced scalar primitive*). On the local unreduced Hamiltonian algebra  $\mathcal{A} = \mathbb{C}[z_1, z_2]$ , the constant-coefficient cocycle

$$\omega(f, g) = [\{f, g\}_{\Omega_{\text{loc}}}]_0$$

is the CE coboundary of the linear functional

$$\eta(f) = -[f]_0.$$

With the convention  $(d_{\text{CE}}\eta)(f, g) = -\eta(\{f, g\})$ , one has  $d_{\text{CE}}\eta = \omega$ . The primitive lives on the unreduced algebra  $\mathcal{A}$ ; scalar reduction to  $\bar{\mathcal{A}} = \mathcal{A}/\mathbb{C} \cdot 1$  removes it.

*Proof.* The identity is immediate:

$$(d_{\text{CE}}\eta)(f, g) = -\eta(\{f, g\}_{\Omega_{\text{loc}}}) = [\{f, g\}_{\Omega_{\text{loc}}}]_0 = \omega(f, g).$$

If  $\eta$  descended to  $\bar{\mathcal{A}}$ , then it would vanish on constants. But  $\eta(1) = -1$ . Equivalently  $\omega(z_1, z_2) = 1$  while  $\{z_1, z_2\} = 1$  is zero in the scalar-reduced Hamiltonian quotient.  $\square$

THEOREM E.12 (*Local scalar contact QME obstruction*). Let

$$\omega(f, g) = [\{f, g\}_{\Omega_{\text{loc}}}]_0$$

be the formal-local Hamiltonian scalar cocycle of Proposition E.2. On monomials,

$$\omega(z_1^k z_2^l, z_1^m z_2^n) = \begin{cases} kn - lm, & (k+m, l+n) = (1, 1), \\ 0, & \text{otherwise.} \end{cases}$$

For compactly supported  $a, b \in \Omega_c^\circ(\mathbb{R}_{\text{brane}})$ , the unreduced scalar boundary contact in the local mixed HT brane-defect model is

$$\{I_\partial(a, f), I_\partial(b, g)\}_{\text{open}} = I_\partial(ab, \{f, g\}_{\tilde{A}}) + N \omega(f, g) \int_{\mathbb{R}} a(t)b(t) dt.$$

In the quantum Weyl normalization it is equivalently represented by

$$\text{Tr}(A XY) - \text{Tr}(A YX) = \hbar N \text{Tr}(A) \mod \mathcal{W}_N \widehat{\mu}(\mathfrak{sI}_N).$$

Equivalently,

$$\text{Tr}(A[X, Y]) = \hbar N \text{Tr}(A) \mod \mathcal{W}_N \widehat{\mu}(\mathfrak{sI}_N),$$

for the matrix Weyl commutator  $[X, Y] = XY - YX$ . After insertion of the central boundary ghost  $\gamma_1, |\gamma_1| = 1$ , the degree-one obstruction representative is

$$\text{Ob}_{\text{sc}}(\gamma_1; a, f; b, g) = \hbar N \omega(f, g) \int_{\mathbb{R}} a(t)b(t)\gamma_1(t) dt.$$

Its associated graded CE class is the formal scalar multiple  $\hbar N[\bar{c}] \in H_{\text{Lie}}^2(\tilde{A}; \mathbb{C})[[\hbar]]$ . This class is a nonzero scalar-reduced Hamiltonian cohomology class. Hence an internal scalar counterterm in the ordinary scalar-reduced  $\mathfrak{gl}_N$  brane-defect model would require a leading symbol primitive for  $\hbar N[\bar{c}]$ .

*Proof.* The monomial formula for  $\omega$  follows by applying

$$\{z_1^k z_2^l, z_1^m z_2^n\} = (kn - lm) z_1^{k+m-1} z_2^{l+n-1}$$

and then keeping only the constant Taylor coefficient. The first displayed formula is Corollary 5.19I. The second is the quantum version of the same central term: the Weyl convention gives  $[X^i_j, Y^k_l] = \hbar \delta_l^i \delta_j^k$ , and the normal ordered moment relation is  $Y_h X_h - X_h Y_h + \hbar N I \equiv 0$ . Hence  $XY - YX = \hbar N I$  in the quantum Hamiltonian reduction, and tracing against  $A$  gives the stated sign. The matrix-commutator form is the same identity written as  $\text{Tr}(A(XY - YX))$ .

The scalar contact itself has degree 0. Multiplication by the central ghost  $\gamma_1$  makes the QME obstruction representative degree 1, the degree of the obstruction complex. The CE associated graded of this representative is the scalar extension cocycle  $\hbar N \omega$ , hence the formal class  $\hbar N[\bar{c}] \in H_{\text{Lie}}^2(\tilde{A}; \mathbb{C})[[\hbar]]$ .

If it were exact in the scalar-reduced Hamiltonian source, the leading scalar term would give a linear primitive  $\bar{\eta} : \tilde{A} \rightarrow \mathbb{C}$  with  $\delta \bar{\eta} = \omega$ . Pulling  $\bar{\eta}$  back to  $\mathbb{C}[z_1, z_2]$  and setting the value on 1 to zero would contradict the computation

$$\omega(z_1, z_2) = 1, \quad \{z_1, z_2\} = 1.$$

Equivalently, Lemma E.11 gives the unreduced scalar primitive used here. It lives on the unreduced polynomial Hamiltonian algebra; scalar reduction to  $\tilde{A}$  removes that primitive. Thus a scalar-reduced counterterm would require an associated-graded primitive for the class.  $\square$

**THEOREM E.13** (*Scalar anomaly transgression: four equivalent sightings*). In the formal-local mixed geometry  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}^2_0$  with  $\Omega_{\text{loc}} = dz_1 \wedge dz_2$  of Proposition E.2, the scalar Hamiltonian cocycle

$$\omega(f, g) = [\{f, g\}_{\Omega_{\text{loc}}}]_0, \quad \omega(z_1, z_2) = 1,$$

is detected in four equivalent forms. All four witness one and the same nonzero class in  $H^2(\tilde{A}; \mathbb{C})[[\hbar]]$ . Up to the quantum scalar  $\hbar N$  supplied by the open Chan–Paton index loop, the cocycle is the same in each.

(i) *Algebraic central extension of the Hamiltonian Lie algebra.* The short exact sequence of Lie algebras

$$0 \longrightarrow \mathbb{C} \cdot \mathbf{1} \longrightarrow \mathcal{A} = \mathbb{C}[z_1, z_2] \longrightarrow \tilde{\mathcal{A}} = \mathcal{A}/\mathbb{C} \cdot \mathbf{1} \longrightarrow 0, \quad \mathcal{A} \text{ with bracket } \{f, g\}_{\Omega_{\text{loc}}},$$

is the central extension whose Lie cocycle is exactly  $\omega$ . Its class is  $[\bar{c}] \in H^2(\tilde{\mathcal{A}}; \mathbb{C})$  with Lemma E.II.

(ii) *Lie-algebraic central extension.* Setting  $\hat{\mathfrak{h}}_{\text{cent}} = \tilde{\mathcal{A}} \oplus \mathbb{C}\bar{c}$  with bracket

$$[[\bar{f}, \bar{g}]]_{\text{cent}} = \overline{\{f, g\}} + \omega(f, g)\bar{c}, \quad [[\bar{c}, -]]_{\text{cent}} = 0,$$

produces a Lie central extension whose class is the same  $[\bar{c}]$ . This is the source-side  $\hat{\mathfrak{h}}_{\text{cent}}$ .

(iii) *Trace-algebraic central extension on the boundary.* With the unreduced boundary Hamiltonian generator  $I_\partial(a, f)$  of Theorem E.12, the open Chan–Paton ordinary-trace open coupling produces the central extension

$$0 \longrightarrow \mathbb{C} \cdot \text{Tr}(\mathbf{1}) \longrightarrow \text{Tr } \mathbb{C}[\phi_1, \phi_2]_{\text{red}} \longrightarrow \mathcal{A}_{\partial, \text{red}} \longrightarrow 0$$

of trace-reduced boundary algebras, whose Lie cocycle is  $\hbar N \omega$  and whose class in  $H^2(\mathcal{A}_{\partial, \text{red}}; \mathbb{C})[[\hbar]]$  is  $\hbar N [\bar{c}]$ .

(iv) *Quantum Weyl commutator.* In the Weyl quantization,

$$[\Phi_\hbar(z_1), \Phi_\hbar(z_2)]_\star = \hbar N \mod \mathcal{W}_N \widehat{\mu}(\mathfrak{sl}_N), \quad W_1^{\text{sc, ord}} = \hbar N [\bar{c}] \quad \text{in cohomology,}$$

i.e. the Capelli shift makes the scalar commutator equal  $\hbar N$  times the identity, modulo the Hamiltonian moment-map ideal of  $\mathfrak{sl}_N$ . The associated CE class is again  $\hbar N [\bar{c}]$ .

These four are equivalent in the strong sense that the same class  $[\bar{c}] \in H^2(\tilde{\mathcal{A}}; \mathbb{C})$  is the image, under the canonical scalar projections, of all four cocycles:

$$\begin{array}{ccc} [\omega]_{(i)} & = & [\omega]_{(ii)} \in H^2(\tilde{\mathcal{A}}; \mathbb{C}), \\ \downarrow \hbar N \cdot & & \downarrow \hbar N \cdot \\ [\hbar N \omega]_{(iii)} & = & [\hbar N \omega]_{(iv)} \in H^2(\tilde{\mathcal{A}}; \mathbb{C})[[\hbar]], \end{array}$$

with horizontal equalities of cocycles given by the Lie/algebraic identification of (i) with (ii) and the trace/Weyl identification of (iii) with (iv), and vertical multiplications by  $\hbar N$  given by the open Chan–Paton trace pairing.

Slogan: *constants vanish as Hamiltonian vector fields but survive as brane center-of-mass charge.* The radial Poisson field  $X_f$  annihilates the constant line in  $\mathcal{A}$  (Proposition E.2), but the open trace pairs the same constant line against  $\text{Tr}(\mathbf{1}) = N$  and produces the Capelli shift  $\hbar N [\bar{c}]$ . Equivalently, the same class is the projective curvature of the modular line bundle  $\Omega_{\text{central}}$  evaluated at the trace generator  $J(f) = \text{Tr } f(\phi_1, \phi_2)$  by Theorem 6.8: the four named sightings are the four ways the local-stalk curvature value enters the formal-Darboux computation, with the determinant-line interpretation of the central extension supplied by Remark 6.9.

*Proof.* Sighting (i) is the content of Lemma E.II:  $\omega$  is the CE coboundary of  $\eta(f) = -[f]_0$  on the unreduced  $\mathcal{A}$ , and scalar reduction to  $\tilde{\mathcal{A}}$  removes  $\eta$ . Hence  $[\omega]$  is nonzero on  $\tilde{\mathcal{A}}$ ; the central extension presented in (i) is this nontrivial class.

Sighting (ii) is the standard correspondence between Lie central extensions of  $\tilde{\mathcal{A}}$  by  $\mathbb{C}$  and elements of  $H^2(\tilde{\mathcal{A}}; \mathbb{C})$ . Setting  $[[\bar{f}, \bar{g}]]_{\text{cent}} = \overline{\{f, g\}} + \omega(f, g)\bar{c}$  defines a Lie bracket because  $\omega$  is a 2-cocycle, and

the associated extension class is  $[\omega] = [\bar{c}]$  by construction. This is the same class as (i) up to canonical identification.

Sighting (iii) is the trace-algebraic content of Theorem E.12: the open Chan–Paton ordinary trace pairs  $I_\partial(a, f)I_\partial(b, g)$  and produces the residual  $\hbar N \omega(f, g) \int_{\mathbb{R}} a(t)b(t) dt$ , which is the trace-algebraic cocycle. The boundary source map  $A_{\partial, \text{red}} \rightarrow \tilde{A}$  (induced by  $I_\partial(a, f) \mapsto \tilde{f}$  on the Hamiltonian factor and by evaluation against the test function) factors the trace cocycle as  $\hbar N$  times the canonical  $\tilde{A}$ -cocycle  $\omega$ . Hence the class in  $H^2(A_{\partial, \text{red}}; \mathbb{C})[[\hbar]]$  is  $\hbar N$  times the same scalar  $[\bar{c}]$  under this canonical pullback: the trace coefficient is the only scalar dependence beyond the source cocycle.

Sighting (iv) is the quantum Weyl reformulation in the same theorem: the matrix Weyl commutator  $[X, Y] = \hbar NI$  modulo  $\mathcal{W}_N \hat{\mu}(\mathfrak{sl}_N)$  is the Capelli normal-ordered scalar shift. Tracing against  $A$  and applying the associated-graded scalar-symbol projection produces the same scalar CE class  $\hbar N [\bar{c}]$ .

The three vertical/horizontal equalities of the diagram are induced by the canonical scalar-symbol projection (Proposition E.2) and the open Chan–Paton trace. The algebraic content:  $X_f(\mathbf{1}) = \mathbf{0}$  for every  $f$  (the constant line is annihilated by every Hamiltonian vector field), while the trace pairing  $\text{Tr}(\mathbf{1}) = N$  records the constant line as the open index loop charge  $\hbar N$ . The four sightings are the four ways the same class  $[\bar{c}]$  becomes visible after pairing with the open trace, the source CE differential, the Lie bracket, or the boundary Weyl commutator.  $\square$

*Remark E.14 (Action of the four cancellation branches on the scalar anomaly class).* Each branch acts on the class  $\hbar N [\bar{c}] \in H^2(\tilde{A}; \mathbb{C})[[\hbar]]$  of Theorem E.13. The unreduced central character produces the CE coboundary  $d_{\text{CE}}(\hbar N \eta) = \hbar N \omega$  on the unreduced source, with  $\eta(f) = -[f]_{\mathbf{0}}$  of Lemma E.11 admitted as a degree-zero element (Theorem E.15). The balanced  $\mathbb{C}^{N|N}$  supertrace replaces the open trace coefficient by  $\text{sdim } \mathbb{C}^{N|N} = \mathbf{0}$ , rescaling the cocycle to zero before its CE class is formed (Theorem E.16). Supertraceless monodromy extends the balanced computation to marked open loops with cyclic supertrace zero on every monodromy word (Theorem E.25). Full  $\mathfrak{gl}(V)$ -equivariant signed Brauer marker tensors satisfy  $\text{Str}_{\text{cyc}}(\mathbf{m}) = \mathbf{0}$  on  $\mathbb{C}^{N|N}$  (Lemma E.31, Theorem E.32). By Theorem E.52, any internal local counterterm in the ordinary scalar-reduced  $\mathfrak{gl}_N$  branch must contribute leading scalar class  $-\hbar N [\bar{c}]$ . The class  $[\bar{c}] \neq \mathbf{0}$ , so the four branch replacements above are exactly the scalar cancellations available here.

**THEOREM E.15 (Central-character source replacement for the scalar QME symbol).** Let  $A = \mathbb{C}[z_1, z_2]$  and let  $\tilde{f}$  denote the image of  $f$  in  $\tilde{A} = A/\mathbb{C} \cdot \mathbf{1}$  for the local Hamiltonian algebra of Proposition E.2. Define the ordinary trace central character

$$\chi_N(f) = N[f]_{\mathbf{0}}$$

and the unreduced boundary generator

$$I_\partial^\chi(a, f) = I_\partial(a, \tilde{f}) + \chi_N(f) \int_{\mathbb{R}} a(t) dt.$$

Then, for all  $a, b \in C_c^\infty(\mathbb{R})$  and all  $f, g \in A$ ,

$$\{I_\partial^\chi(a, f), I_\partial^\chi(b, g)\}_{\text{open}} = I_\partial^\chi(ab, \{f, g\}).$$

In the quantum Weyl normalization the scalar part is likewise absorbed by the same central character:

$$J_N^\chi(\mathbf{1}) = N, \quad \text{Tr}(A[X, Y]) = J_N^\chi([z_1, z_2]_*) \text{Tr}(A) = \hbar J_N^\chi(\mathbf{1}) \text{Tr}(A).$$

Thus the local scalar QME symbol is absorbed after replacing the scalar-reduced source by the unreduced source with central character  $\chi_N$ . This is a source replacement before scalar reduction. The original scalar-reduced QME complex would require its own null-homotopy. Equivalently, if  $\eta(f) = -[f]_o$ , then

$$d_{\text{CE}}\eta = \omega, \quad \chi_N(f) = N[f]_o = -N\eta(f),$$

so

$$d_{\text{CE}}\chi_N = -N\omega, \quad \hbar N\omega + d_{\text{CE}}(\hbar\chi_N) = 0$$

in the unreduced source.

*Proof.* Decompose the Poisson bracket into its scalar and reduced parts:

$$\{f, g\} = \overline{\{f, g\}} + \omega(f, g) \cdot 1.$$

By Theorem E.12,

$$\{I_\partial(a, \tilde{f}), I_\partial(b, \tilde{g})\}_{\text{open}} = I_\partial(ab, \overline{\{f, g\}}) + N\omega(f, g) \int_{\mathbb{R}} a(t)b(t) dt.$$

Since the central-character summand is scalar, it has zero open bracket with every boundary field. Therefore the right hand side is exactly

$$I_\partial(ab, \overline{\{f, g\}}) + \chi_N(\{f, g\}) \int_{\mathbb{R}} a(t)b(t) dt = I_\partial^\chi(ab, \{f, g\}).$$

The quantum formula is the same computation in the Weyl algebra:  $[z_1, z_2]_* = \hbar$ ,  $J_N^\chi(1) = N$ , and the matrix moment relation gives  $\text{Tr}(\mathcal{A}[X, Y]) = \hbar N \text{Tr}(\mathcal{A})$ . Hence the scalar Weyl commutator is the image of the unreduced Moyal scalar instead of an obstruction class.

This is the normal-ordering mechanism. The relation  $YX - XY + \hbar NI \equiv 0$  says that an  $XY$ -to- $YX$  interchange contributes the scalar  $\hbar NI$ . In the scalar-reduced source this scalar is invisible and therefore appears as  $\text{Ob}_{\text{sc}}$ . In the unreduced source with  $J_N^\chi(1) = N$ , the same scalar is  $J_N^\chi([z_1, z_2]_*) = \hbar N$ , so it is absorbed before forming the QME obstruction class.  $\square$

**THEOREM E.16** (*Local  $\mathfrak{gl}(N|N)$  supertrace scalar QME cancellation*). Let  $V = \mathbb{C}^{N|N}$ , replace the ordinary trace by the supertrace, and use the canonical even Weyl relations on  $\text{End}(V)$  with the supertrace pairing in the same formal-local model of Proposition E.2. Put

$$\text{sdim } V = \text{Str}(\text{id}_V) = 0.$$

The classical scalar contact term is

$$\{I_\partial^{\text{str}}(a, f), I_\partial^{\text{str}}(b, g)\}_{\text{open}} = I_\partial^{\text{str}}(ab, \{f, g\}_{\mathcal{A}}) + \text{sdim}(V) \omega(f, g) \int_{\mathbb{R}} a(t)b(t) dt,$$

and hence has zero scalar component for  $V = \mathbb{C}^{N|N}$ . The quantum scalar commutator is

$$\text{Str}(\mathcal{A}[X, Y]) = \hbar \text{sdim}(V) \text{Str}(\mathcal{A}) = 0 \quad \text{mod } \mathcal{W}_{\text{End}(V)} \widehat{\mu}(\mathfrak{gl}(V)).$$

Consequently the scalar part of the local mixed brane-defect QME vanishes in the balanced  $\mathfrak{gl}(N|N)$  supertrace model by the supertrace calculation.



*Proof.* The scalar term in the classical boundary bracket is the contraction of the identity endomorphism in the trace pairing. For an ordinary  $N$ -dimensional vector space this contraction is  $N$ . For a super vector space it is  $\text{Str}(\text{id}_V) = \text{sdim } V$ . The same local calculation as in Theorem E.12 therefore gives the displayed classical formula. For  $V = \mathbb{C}^{N|N}$ , this coefficient is zero.

In coordinates, the super-Weyl contraction carries the Koszul sign of the supertrace pairing. Summing the diagonal contraction gives

$$YX - XY + \hbar \text{sdim}(V)I \equiv 0 \pmod{\mathcal{W}_{\text{End}(V)} \widehat{\mu}(\mathfrak{gl}(V))}.$$

Thus

$$\text{Str}(\mathcal{A}[X, Y]) = \hbar \text{sdim}(V) \text{Str}(\mathcal{A}),$$

which is zero for  $V = \mathbb{C}^{N|N}$ . These are precisely the scalar terms entering the degree-one representative  $\text{Ob}_{\text{sc}}$ ; hence that representative is zero in the balanced supertrace theory.  $\square$

*Definition E.17 (Balanced scalar-contact subcomplex).* Let  $V = \mathbb{C}^{N|N}$ . A balanced scalar-contact subcomplex over an interval  $I \subset \mathbb{R}_{\text{brane}}$  is a complete filtered subcomplex

$$\mathfrak{S}_{\text{bal,sc}}^\bullet(I) \subset \mathfrak{D}_{w,\partial,\hbar,\text{str}}^\bullet(I)$$

of the local supertrace brane-defect local-functional complex, with differential  $d_{\text{QME}}$ , satisfying the following conditions. First, the filtration  $F^\bullet$  is the closed scalar-contact filtration generated by identity-endomorphism contractions in  $\text{End}(V)$ . Second, the associated-graded scalar-contact extractor factors through the superdimension:

$$\sigma_{\text{bal,sc}} = \text{sdim}(V) \tau_{\text{sc}} : \text{gr}_F \mathfrak{S}_{\text{bal,sc}}^\bullet(I) \longrightarrow C_{\text{Lie}}^\bullet(\bar{\mathcal{A}}; \mathbb{C}\gamma_{\mathbf{I}})[1][[\hbar]]$$

for a continuous  $\mathbb{C}[[\hbar]]$ -linear extractor  $\tau_{\text{sc}}$ . Third, the same factorization holds after applying  $d_{\text{QME}}$ : every scalar-contact symbol in this subcomplex is obtained by contracting a single identity endomorphism in the supertrace pairing.

*PROPOSITION E.18 (Balanced scalar-contact projection).* On every balanced scalar-contact subcomplex of Definition E.17 with  $V = \mathbb{C}^{N|N}$ , the scalar-symbol map is the zero associated-graded map, and it admits the filtered chain projection

$$\widehat{\sigma}_{\text{bal,sc}} = 0 : \mathfrak{S}_{\text{bal,sc}}^\bullet(I) \longrightarrow C_{\text{Lie}}^\bullet(\bar{\mathcal{A}}; \mathbb{C}\gamma_{\mathbf{I}})[1][[\hbar]].$$

Its chain-map defect is

$$\delta_{\text{bal}} = d_{\text{CE}} \widehat{\sigma}_{\text{bal,sc}} - \widehat{\sigma}_{\text{bal,sc}} d_{\text{QME}} = 0.$$

Hence all scalar-projection obstruction classes  $o_{\sigma,w,\text{bal}}^{(r)}(I)$  vanish on this subcomplex. In particular the first two-Hamiltonian-input defect is

$$\text{ev}_{f,g} \left( o_{\sigma,w,\text{bal}}^{(1)}(I) \right) = \hbar \text{sdim}(V) \omega(f, g) \gamma_{\mathbf{I}} = 0.$$

*Proof.* By definition of the subcomplex, every scalar-contact symbol factors as the ordinary scalar contact multiplied by the contraction of the identity endomorphism in the supertrace pairing. That contraction is

$$\text{Str}(\text{id}_V) = \text{sdim}(V) = N - N = 0.$$

Therefore  $\sigma_{\text{bal,sc}} = \text{sdim}(V)\tau_{\text{sc}} = 0$  on the associated graded. Take the zero filtered lift to the stated scalar CE target. Since both composites  $d_{\text{CE}}\widehat{\sigma}_{\text{bal,sc}}$  and  $\widehat{\sigma}_{\text{bal,sc}}d_{\text{QME}}$  are zero, the chain-map defect is zero. Lemma E.4 then gives  $o_{\sigma,w,\text{bal}}^{(r)}(I) = 0$  for every  $r > 0$ .

Evaluating on two Hamiltonian boundary insertions gives the same local contraction as Theorem E.16: the ordinary coefficient  $N$  in  $\hbar N \omega(f, g)\gamma_{\mathbf{I}}$  is replaced by  $\text{sdim}(V)$ . Thus

$$\hbar \text{sdim}(V) \omega(f, g)\gamma_{\mathbf{I}} = 0,$$

including the test  $(f, g) = (z_1, z_2)$ .  $\square$

*Remark E.19 (Extension obstruction outside the balanced subcomplex).* Proposition E.18 kills the scalar-contact projection tower. The full non-scalar graph/QME residual is a separate obstruction problem. Enlarging  $\mathfrak{S}_{\text{bal,sc}}^\bullet(I)$  to a complete filtered subcomplex  $\mathfrak{S}'$  carrying scalar symbols with unspecified  $\text{sdim}(V)$ -factorization, the obstruction to extending the scalar projection is the first nonzero homogeneous component of

$$d_{\text{CES}} - sd_{\text{QME}} \in \text{Hom}(\text{gr}_F \mathfrak{S}', C_{\text{Lie}}^\bullet(\bar{A}; \mathbb{C}\gamma_{\mathbf{I}})[1][[\hbar]])$$

for a filtered scalar-symbol lift  $s$ . Its cohomology class is the  $o_\sigma^{(r)}$ -class of Lemma E.4. Thus an enlargement is admitted exactly when these classes vanish successively, with compatible finite-window choices.

*Definition E.20 (Balanced Costello graph/counterterm subhabitat).* Fix a Hamiltonian brane-defect specialization of the finite-scale Costello graph calculus and a finite-window tower for the formal-local mixed HT theory  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}^2_0$ . A Costello graph/counterterm enlargement

$$\mathfrak{S}_{\text{bal,sc}}^\bullet(I) \subset \mathfrak{G}_{\text{bal,Cost}}^\bullet(I) \subset \mathfrak{Q}_{w,\partial,\hbar,\text{str}}^\bullet(I)$$

is a balanced graph subhabitat if it is a complete  $d_{\text{QME}}$ -stable filtered subcomplex generated, over  $\mathfrak{S}_{\text{bal,sc}}^\bullet(I)$ , by finite-window Costello graph contributions  $G_{\Gamma,L,M}$  and local counterterm representatives  $C_{\Gamma,L,M}$  such that every scalar-contact associated-graded term is an identity-loop supertrace contraction. Equivalently, for each homogeneous scalar-contact piece,

$$\sigma_{\text{sc}}(\text{gr}_F G_{\Gamma,L,M}) = \text{Str}(\text{id}_V) \tau_{\Gamma,L,M}, \quad \sigma_{\text{sc}}(\text{gr}_F C_{\Gamma,L,M}) = \text{Str}(\text{id}_V) \tau_{\Gamma,L,M}^C,$$

for continuous scalar extractors  $\tau_{\Gamma,L,M}$  and  $\tau_{\Gamma,L,M}^C$ , and the same factorization holds after applying  $d_{\text{QME}}$  to these graph and counterterm generators. The factorization is required to commute with finite-window truncation and admissible weight transport.

**PROPOSITION E.21 (Balanced graph/counterterm preservation).** Let  $V = \mathbb{C}^{N|N}$ . Every balanced graph subhabitat  $\mathfrak{G}_{\text{bal,Cost}}^\bullet(I)$  of Definition E.20 is a balanced scalar-contact subcomplex in the sense of Definition E.17. Hence the zero projection

$$\widehat{\sigma}_{\text{bal,Cost}} = 0: \mathfrak{G}_{\text{bal,Cost}}^\bullet(I) \longrightarrow C_{\text{Lie}}^\bullet(\bar{A}; \mathbb{C}\gamma_{\mathbf{I}})[1][[\hbar]]$$

is a filtered chain map, and all scalar-projection obstruction classes vanish on this specified graph/counterterm subcomplex.

*Proof.* By Definition E.20, each new graph or counterterm scalar symbol is obtained by closing an identity endomorphism loop in the supertrace pairing. Its coefficient is therefore

$$\text{Str}(\text{id}_V) = \text{sdim}(V) = N - N = 0.$$

Thus the associated-graded scalar extractor is zero on the graph and counterterm generators, and it is already zero on  $\mathfrak{S}_{\text{bal,sc}}^\bullet(I)$  by Proposition E.18. The same factorization after applying  $d_{\text{QME}}$  says that the differential creates scalar-contact symbols only in the identity-loop class. Therefore

$$d_{\text{CE}}\widehat{\sigma}_{\text{bal,Cost}} - \widehat{\sigma}_{\text{bal,Cost}}d_{\text{QME}} = 0.$$

The finite-window and weight-transport compatibility in the definition gives the same zero class in the completed inverse limit. Hence the scalar-contact projection obstruction tower of Lemma E.4 vanishes on  $\mathfrak{G}_{\text{bal,Cost}}^\bullet(I)$ .  $\square$

**THEOREM E.22** (*Graph-extension scalar obstruction*). Let

$$\mathfrak{S}_{\text{bal,sc}}^\bullet(I) \subset \mathfrak{G}_{\text{Cost}}^\bullet(I) \subset \mathfrak{Q}_{w,\partial,\hbar,\text{str}}^\bullet(I)$$

be any complete  $d_{\text{QME}}$ -stable filtered Costello graph/counterterm enlargement. Put

$$\overline{\mathfrak{G}}_{\text{Cost}}^\bullet(I) = \mathfrak{G}_{\text{Cost}}^\bullet(I) / \mathfrak{S}_{\text{bal,sc}}^\bullet(I), \quad S^\bullet = C_{\text{Lie}}^\bullet(\bar{A}; \mathbb{C}\gamma_1)[1][[\hbar]].$$

Let

$$\bar{\sigma}_{\text{Cost}}: \text{gr}_F \overline{\mathfrak{G}}_{\text{Cost}}^\bullet(I) \longrightarrow S^\bullet$$

be the associated-graded scalar-symbol extractor of the added graph/counterterm terms. The zeroth obstruction to extending the balanced scalar-contact projection to  $\mathfrak{G}_{\text{Cost}}^\bullet(I)$  is the associated-graded chain-map defect

$$\delta_{\text{Cost},\sigma}^{(0)}(I) = d_{\text{CE}}\bar{\sigma}_{\text{Cost}} - \bar{\sigma}_{\text{Cost}}d_{\text{gr}} \in \text{Hom}_0^1(\text{gr}_F \overline{\mathfrak{G}}_{\text{Cost}}^\bullet(I), S^\bullet).$$

If this defect vanishes, choose a filtered lift  $s$  of  $\bar{\sigma}_{\text{Cost}}$ , extended by zero on  $\mathfrak{S}_{\text{bal,sc}}^\bullet(I)$ . If the first nonzero homogeneous component of

$$\delta_s = d_{\text{CE}}s - sd_{\text{QME}}$$

lowers the scalar-contact filtration by  $r > 0$ , then it defines

$$\mathfrak{d}_{\text{Cost},\sigma}^{(r)}(I) = \left[ \delta_s^{(-r)} \right] \in H^1\left(\text{Hom}_{-r}(\text{gr}_F \overline{\mathfrak{G}}_{\text{Cost}}^\bullet(I), S^\bullet)\right).$$

These classes are independent of the chosen lift up to the usual Hom coboundary. A filtered chain projection extending the balanced projection exists exactly when

$$\delta_{\text{Cost},\sigma}^{(0)}(I) = 0, \quad \mathfrak{d}_{\text{Cost},\sigma}^{(r)}(I) = 0 \quad (r > 0)$$

successively, with compatible finite-window representatives.

Explicitly, if  $X_{\Gamma,M}$  is an added graph contribution or counterterm representative and, modulo  $F^{p-r+1}$ ,

$$d_{\text{QME}}X_{\Gamma,M} = \sum_{\Gamma'} \varepsilon_{\Gamma,\Gamma'} X_{\Gamma',M}, \quad X_{\Gamma,M} \in F^p,$$

with the signs  $\varepsilon_{\Gamma, \Gamma'}$  determined by the graph orientation, BV bracket, and counterterm convention, then the obstruction cocycle is computed by

$$\mathfrak{o}_{\text{Cost}, \sigma}^{(r)}(X_{\Gamma, M}) = \left[ d_{\text{CE}} s(X_{\Gamma, M}) - \sum_{\Gamma'} \varepsilon_{\Gamma, \Gamma'} s(X_{\Gamma', M}) \right]_{F^{p-r}/F^{p-r+1}}.$$

Thus a graph or counterterm term beyond the identity-loop superdimension class is admitted by the balanced theorem exactly after this displayed relative Hom class is killed by an explicit compatible counterterm correction.

*Proof.* The quotient by  $\mathfrak{S}_{\text{bal}, \text{sc}}^{\bullet}(I)$  makes the statement relative to the already proved zero projection on the balanced scalar-contact subcomplex. If  $d_{\text{CE}} \tilde{\sigma}_{\text{Cost}} - \tilde{\sigma}_{\text{Cost}} d_{\text{gr}}$  is nonzero, then the associated-graded scalar extractor has a chain-map defect, and every filtered chain lift carries that defect in its associated graded. This is the zeroth defect  $\delta_{\text{Cost}, \sigma}^{(0)}(I)$ .

Once the associated-graded defect is killed, the proof is the relative version of Lemma E.4. The commutator  $d_{\text{CE}} s - s d_{\text{QME}}$  strictly lowers filtration. Its first nonzero homogeneous piece is closed for the Hom differential because both differentials square to zero and because the lower pieces have already been removed. Replacing  $s$  by  $s - \alpha$ , with  $\alpha$  lowering filtration by the same amount and vanishing on  $\mathfrak{S}_{\text{bal}, \text{sc}}^{\bullet}(I)$ , changes this component by the Hom coboundary  $D\alpha$ . Therefore the class is independent of choices and vanishes exactly when the next correction to  $s$  exists. Completeness of the filtration and compatibility of the finite-window tower give the stated successive criterion.

The displayed graph formula is this same chain-defect evaluated on a homogeneous graph or counterterm generator, with the  $d_{\text{QME}}$ -boundary written as the signed sum of its graph and counterterm boundary terms. Hence it is computable from the supplied Costello graph differential, orientation signs, and scalar symbols. If all added terms satisfy Definition E.20, then  $s = 0$  is the required lift for  $V = \mathbb{C}^{N|N}$ , and all the displayed classes vanish, recovering Proposition E.21.  $\square$

**PROPOSITION E.23** (*First non-identity-loop scalar symbol*). Let  $V = \mathbb{C}^{N|N}$ . Fix a finite local Hamiltonian window  $M$  containing the polynomial Hamiltonian inputs under discussion, and let  $T_{\bullet} = (T_1, \dots, T_m)$  be an ordered word of even  $\text{End}(V)$ -labels carried by one open index loop. Write  $M(T_{\bullet}) = T_1 \cdots T_m$ , with the empty product equal to  $\text{id}_V$ . Let  $\Gamma_{i, M}^{T_{\bullet}}$  be the connected local one-cross-contraction boundary graph with two Hamiltonian external legs  $(a, f)$  and  $(b, g)$ , one Weyl edge, and this marked scalar loop. Denote the degree-one scalar-contact representative obtained after inserting  $\gamma_1$  by  $X_{\Gamma_{i, M}^{T_{\bullet}}}(a, f; b, g)$ .

Then  $X_{\Gamma_{i, M}^{T_{\bullet}}}(a, f; b, g) \in F^1$ , its leading term has  $F^1/F^2$  coefficient equal to the displayed scalar, and

$$X_{\Gamma_{i, M}^{T_{\bullet}}}(a, f; b, g) = \hbar \text{Str}(M(T_{\bullet})) \omega(f, g) \int_I a(t) b(t) \gamma_1(t) dt \mod F^2.$$

Equivalently,

$$\sigma_{\text{sc}} \left( \text{gr}_F X_{\Gamma_{i, M}^{T_{\bullet}}}(a, f; b, g) \right) = \hbar \text{Str}(M(T_{\bullet})) \omega(f, g) \gamma_1.$$

For monomials,

$$\omega(z_1^k z_2^l, z_1^m z_2^n) = \begin{cases} kn - lm, & (k+m, l+n) = (1, 1), \\ 0, & \text{otherwise,} \end{cases}$$

so the first non-identity test is obtained by a single marker  $T$ :

$$\sigma_{\text{sc}}\left(\text{gr}_F X_{\Gamma_{1,M}^T}(1, z_1; 1, z_2)\right) = \hbar \text{Str}(T) \gamma_1.$$

In particular, if  $P_+$  is projection onto one even basis line, then

$$\sigma_{\text{sc}}\left(\text{gr}_F X_{\Gamma_{1,M}^{P_+}}(1, z_1; 1, z_2)\right) = \hbar \gamma_1 \neq 0.$$

Thus this is the first scalar symbol beyond the identity-loop computation. Its associated CE class is

$$\hbar \text{Str}(\mathcal{M}(T_\bullet))[\bar{e}] \gamma_1.$$

It is nonzero exactly when  $\text{Str}(\mathcal{M}(T_\bullet)) \neq 0$ . If every scalar-contact loop monodromy word in a complete  $d_{\text{QME}}$ -stable graph/counterterm subcomplex, and every such word appearing after  $d_{\text{QME}}$ , has supertrace zero, then the zero scalar projection is again a filtered chain map on that subcomplex.

*Proof.* The graph has one local scalar contact on the brane line, so it lies in  $F^1$ . The one-edge graph has exactly one scalar contact, hence its associated graded is read in  $F^1/F^2$ . The boundary Wick normalization fixed in Theorem 5.223 gives the one-edge antisymmetrized coefficient  $\hbar P(f, g)$ . Taking the scalar Taylor coefficient gives

$$[P(f, g)]_0 = \omega(f, g),$$

with positive sign for the ordered pair  $(z_1, z_2)$ , since  $\{z_1, z_2\} = 1$ . Reversing the two external legs changes the sign through  $\omega(g, f) = -\omega(f, g)$ .

The open index loop is computed in a homogeneous basis  $e_\alpha$  of  $V$ , the closed marked loop contributes

$$\sum_{\alpha} (-1)^{|e_\alpha|} (\mathcal{M}(T_\bullet))_{\alpha}^{\alpha} = \text{Str}(\mathcal{M}(T_\bullet)).$$

For the empty word this is  $\text{Str}(\text{id}_V) = \text{sdim } V = 0$ , which is exactly the identity-loop cancellation of Proposition E.21. For a non-identity marker  $T$  the same contraction gives  $\text{Str}(T)$ . Projection onto one even basis line has supertrace 1, giving the displayed nonzero symbol.

The scalar contact has BV degree 0; multiplication by  $\gamma_1$ , of degree 1, puts the representative in the degree-one QME obstruction complex. The cocycle  $\omega$  is the same scalar Hamiltonian cocycle as in Theorem E.12. That theorem proves  $[\bar{e}] \neq 0$  on the scalar-reduced Hamiltonian source, so multiplying by the scalar  $\hbar \text{Str}(\mathcal{M}(T_\bullet))$  gives a nonzero class exactly when this supertrace is nonzero.

Suppose the stated marked-loop subcomplex has only supertraceless monodromy words, before and after applying  $d_{\text{QME}}$ . The formula above then gives zero associated scalar symbol on every graph and counterterm generator and on every associated-graded boundary term. Hence  $d_{\text{CEO}} - 0 d_{\text{QME}} = 0$ , so the zero scalar projection is a filtered chain map on that subcomplex.  $\square$

*Definition E.24 (Supertraceless-monodromy prehabitat).* Let  $V = \mathbb{C}^{N|N}$ . For an ordered word  $T_\bullet = (T_1, \dots, T_m)$  of even  $\text{End}(V)$ -labels on a scalar open index loop, set

$$\text{st}(T_\bullet) = \text{Str}(T_1 \cdots T_m), \quad \text{st}(\emptyset) = \text{Str}(\text{id}_V) = 0.$$

The word is supertraceless if  $\text{st}(T_\bullet) = 0$ . This condition is imposed on the ordered product  $T_1 \cdots T_m$ .

Fix a complete  $d_{\text{QME}}$ -stable finite-window graph/counterterm enlargement of the local balanced supertrace brane-defect complex. The supertraceless-monodromy prehabitat

$$\mathfrak{G}_{\text{st}=0}^{\bullet, \text{pre}}(I)$$

is the closed filtered span, over  $\mathfrak{S}_{\text{bal}, \text{sc}}^{\bullet}(I)$ , of homogeneous graph/counterterm representatives whose every scalar-contact associated-graded loop word is supertraceless. Let

$$\Pi_{\text{st} \neq 0}$$

denote the quotient of the associated-graded scalar-contact monodromy module by the closed span of supertraceless loop words. The  $d_{\text{QME}}$ -preservation obstruction is the family of homogeneous quotient classes

$$\mathcal{B}_{\text{st}=0}(X; q) = \Pi_{\text{st} \neq 0}([d_{\text{QME}} X]_{F^q / F^{q+1}}), \quad X \in \mathfrak{G}_{\text{st}=0}^{\bullet, \text{pre}}(I),$$

over all filtration degrees  $q$ . Write  $\mathcal{B}_{\text{st}=0} = 0$  when every class in this family vanishes. If  $\mathcal{B}_{\text{st}=0} = 0$ , the prehabitat is a complete  $d_{\text{QME}}$ -stable filtered subcomplex; in that case it is called the supertraceless-monodromy graph/counterterm subhabitat and is denoted  $\mathfrak{G}_{\text{st}=0}^{\bullet}(I)$ .

**THEOREM E.25** (*Supertraceless-monodromy scalar projection*). Assume the notation of Definition E.24. The prehabitat is preserved by  $d_{\text{QME}}$  if and only if

$$\mathcal{B}_{\text{st}=0} = 0.$$

Under this condition the zero scalar projection

$$\widehat{\sigma}_{\text{st}=0} = 0: \mathfrak{G}_{\text{st}=0}^{\bullet}(I) \longrightarrow C_{\text{Lie}}^{\bullet}(\vec{A}; \mathbb{C}\gamma_1)[1][[\hbar]]$$

is a filtered chain map. Hence all scalar-projection obstruction classes of Lemma E.4 vanish on the supertraceless-monodromy subhabitat.

More explicitly, let  $X_{\Gamma, M} \in F^p$  be a homogeneous generator of the prehabitat and suppose that, modulo  $F^{p-r+1}$ ,

$$d_{\text{QME}} X_{\Gamma, M} = \sum_{\Gamma', U_{\bullet}} \varepsilon_{\Gamma, \Gamma', U_{\bullet}} X_{\Gamma' U_{\bullet}, M},$$

where  $U_{\bullet}$  runs over scalar-loop monodromy words appearing in the boundary terms. The first possible preservation obstruction is the quotient class

$$\mathcal{B}_{\text{st}=0}^{(r)}(X_{\Gamma, M}) = \left[ \sum_{\Gamma', U_{\bullet}} \varepsilon_{\Gamma, \Gamma', U_{\bullet}} X_{\Gamma' U_{\bullet}, M} \right]_{\text{st} \neq 0, F^{p-r} / F^{p-r+1}}.$$

Its scalar CE image on a two-Hamiltonian-input scalar loop is

$$\hbar \sum_{\Gamma', U_{\bullet}} \varepsilon_{\Gamma, \Gamma', U_{\bullet}} \text{Str}(M(U_{\bullet})) \omega(f_{\Gamma'}, g_{\Gamma'}) \gamma_1.$$

In cohomology this becomes

$$\hbar \sum_{\Gamma', U_{\bullet}} \varepsilon_{\Gamma, \Gamma', U_{\bullet}} \text{Str}(M(U_{\bullet})) [\bar{c}] \gamma_1.$$

Thus a non-supertraceless boundary word is the exact obstruction to  $d_{\text{QME}}$ -preservation, and its scalar shadow is precisely the marked-loop class computed in Proposition E.23.

For the unmarked Hamiltonian boundary generators  $I_{\partial}^{\text{str}}(a, f)$  used in Theorem E.16, and for the identity-loop graph generators of Definition E.20, the obstruction  $\mathcal{B}_{\text{st}=0}$  vanishes. These generators carry only the empty monodromy word, and the displayed differential  $Q + \{I_0, -\}$  contains only the identity-loop marker. Therefore applying  $d_{\text{QME}}$  to this displayed Hamiltonian generator set preserves supertraceless monodromy. Preservation holds on the unmarked Hamiltonian generator set. A larger Costello specialization with additional marked idempotents, holonomies, or counterterm tensors is governed by  $\mathcal{B}_{\text{st}=0}$  on those added generators.

*Proof.* By definition,  $\Pi_{\text{st} \neq 0}$  kills exactly the associated-graded scalar-contact monodromy terms whose loop words are supertraceless. Hence  $\mathcal{B}_{\text{st}=0} = 0$  says precisely that the  $d_{\text{QME}}$ -boundary of every generator of the prehabitat has only supertraceless loop words in every associated-graded filtration degree. Since the filtration is complete, this homogeneous condition is equivalent to preservation of the closed filtered span. This proves the preservation criterion.

On the preserved subhabitat every scalar-contact loop word  $T_{\bullet}$  satisfies  $\text{Str}(M(T_{\bullet})) = 0$ . Proposition E.23 computes the associated-graded scalar symbol of such a loop as

$$\hbar \text{Str}(M(T_{\bullet})) \omega(f, g) \gamma_{\mathbf{t}},$$

and this is zero. The same is true after applying  $d_{\text{QME}}$ , by the preservation criterion. Therefore both composites  $d_{\text{CE}} \widehat{\sigma}_{\text{st}=0}$  and  $\widehat{\sigma}_{\text{st}=0} d_{\text{QME}}$  are zero, so the zero projection is a filtered chain map. Lemma E.4 then kills the whole scalar-projection obstruction tower on this subhabitat.

The formula for  $\mathcal{B}_{\text{st}=0}^{(r)}(X_{\Gamma, M})$  is just the preceding quotient criterion evaluated on a homogeneous generator and then reduced modulo  $F^{p-r+1}$ . Applying the one-loop scalar computation of Proposition E.23 to each boundary term gives the displayed scalar CE representative. Passing to cohomology replaces  $\omega$  by the nonzero class  $[\bar{c}]$  from Theorem E.12.

For the unmarked Hamiltonian boundary generators, the open index loop carries the empty word  $\emptyset$ , so its coefficient is  $\text{Str}(\text{id}_V) = 0$ . The differential  $Q + \{I_0, -\}$  acts on BV fields and takes brackets with the unmarked classical interaction; it preserves the empty marker word on an open scalar loop. Thus the image of the unmarked Hamiltonian generator set again has only empty loop words. The identity-loop graph subhabitat has the same property by Definition E.20, which requires the same factorization after applying  $d_{\text{QME}}$ . This proves the stated preservation result on the displayed generator set and leaves arbitrary marked graph/counterterm tensors governed by  $\mathcal{B}_{\text{st}=0}$ .  $\square$

**PROPOSITION E.26** (*Marked  $\mathcal{B}_{\text{st}=0}$  and gauge-equivariant markers*). Let  $V = \mathbb{C}^{N|N}$ , and impose full  $\mathfrak{gl}(V)$ -gauge equivariance on open Chan-Paton labels. A complete computation of  $\mathcal{B}_{\text{st}=0}$  for a marked Costello graph/counterterm enlargement requires, for every homogeneous generator  $X_{\Gamma, M} \in F^p$ , a marked boundary expansion

$$d_{\text{QME}} X_{\Gamma, M} = \sum_{\Gamma', \lambda} \varepsilon_{\Gamma, \Gamma', \lambda} X_{\Gamma' \mathfrak{m}_{\Gamma, \Gamma', \lambda}, M} \mod F^{p-r+1},$$

where each

$$\mathfrak{m}_{\Gamma, \Gamma', \lambda} \in (\text{End}(V)^{\otimes m_{\lambda}})^{\mathfrak{gl}(V)}$$

is an even invariant marker tensor decorating the scalar open index loop. For  $\mathfrak{m} = \sum_{\alpha} T_{\alpha, 1} \otimes \cdots \otimes T_{\alpha, m}$ , put

$$\text{Str}_{\text{cyc}}(\mathfrak{m}) = \sum_{\alpha} \text{Str}(T_{\alpha, 1} \cdots T_{\alpha, m}).$$

Then the first preservation obstruction is

$$\mathcal{B}_{\text{st}=0}^{(r)}(X_{\Gamma,M}) = \left[ \sum_{\Gamma',\lambda} \varepsilon_{\Gamma,\Gamma',\lambda} X_{\Gamma',\mathbf{m}_{\Gamma,\Gamma',\lambda},M} \right]_{\text{st} \neq 0, F^{p-r}/F^{p-r+1}},$$

and its two-Hamiltonian-input scalar shadow is

$$\hbar \sum_{\Gamma',\lambda} \varepsilon_{\Gamma,\Gamma',\lambda} \text{Str}_{\text{cyc}}(\mathbf{m}_{\Gamma,\Gamma',\lambda}) \omega(f_{\Gamma'}, g_{\Gamma'}) \gamma_{\mathbf{I}}.$$

In cohomology this becomes

$$\hbar \sum_{\Gamma',\lambda} \varepsilon_{\Gamma,\Gamma',\lambda} \text{Str}_{\text{cyc}}(\mathbf{m}_{\Gamma,\Gamma',\lambda}) [\bar{c}] \gamma_{\mathbf{I}}.$$

On the displayed unmarked Hamiltonian generator set the differential is  $Q + \{I_o, -\}$ , with unmarked classical interaction; the balanced graph/counterterm subhabitat only allows identity-loop scalar contractions before and after  $d_{\text{QME}}$ . Hence

$$\mathcal{B}_{\text{st}=0} = 0$$

on that generator set. For any larger marked Costello specialization equipped with an oriented finite-window marked boundary complex as in Definition E.28 has the scalar computation below. For a larger marked specialization, the missing datum is exactly equation (E.26), including its signs and invariant marker tensors.

In particular, a single fixed marker  $T \in \text{End}(V)_o$  can occur in a full  $\mathfrak{gl}(V)$ -equivariant marked differential only when  $T = \lambda \text{id}_V$ . Its scalar shadow is then

$$\hbar \lambda \text{Str}(\text{id}_V) \omega(f, g) \gamma_{\mathbf{I}} = 0.$$

Thus the nonzero  $P_+$  test of Proposition E.23 is a genuine obstruction to a non-equivariant or externally marked enlargement. In the full  $\mathfrak{gl}(N|N)$ -equivariant Costello differential, an allowed single-marker term is a scalar multiple of the identity. If one imposes only the even block-diagonal group, or supplies external marker tensors, the admissible datum is the invariant tensor family above, and the first nonzero scalar shadow is detected exactly by the displayed cyclic-supertrace coefficient.

*Proof.* The formula for  $\mathcal{B}_{\text{st}=0}^{(r)}$  is Definition E.24 applied to the marked boundary expansion equation (E.26). The scalar-shadow formula is the linear extension of Proposition E.23: a decomposable marker word contributes the cyclic supertrace of the ordered product around the open index loop, and an invariant tensor is a finite linear combination of such words. Passing to cohomology replaces  $\omega$  by  $[\bar{c}]$  as in Theorem E.25.

The equivariance constraint is the following. A marked local functional with one fixed marker  $T$  transforms under  $g \in GL(V)$  by replacing the marker with  $gTg^{-1}$ . Equivariance of the differential therefore forces  $gTg^{-1} = T$ , or infinitesimally  $[A, T] = 0$  for every  $A \in \mathfrak{gl}(V)$ . Since  $T$  is even, write it in the parity decomposition as  $T = T_o \oplus T_i$ . Commutation with  $\mathfrak{gl}(N) \oplus \mathfrak{gl}(N)$  gives  $T_o = \lambda_o \text{id}$  and  $T_i = \lambda_i \text{id}$ . Commutation with the odd elementary endomorphisms forces  $\lambda_o = \lambda_i$ . Hence  $T = \lambda \text{id}_V$ . The cyclic supertrace is then  $\lambda \text{sdim}(V) = 0$  for  $V = \mathbb{C}^{N|N}$ .

On the unmarked Hamiltonian generator set the differential  $d_{\text{QME}} = Q + \{I_o, -\}$  carries the empty endomorphism label; together with Definition E.20, which requires identity-loop factorization after applying  $d_{\text{QME}}$ , these terms have cyclic supertrace  $\text{Str}(\text{id}_V) = 0$ , so their  $\mathcal{B}_{\text{st}=0}$ -class vanishes. Any further marked graph/counterterm term is governed by its chosen oriented marked boundary complex; equation (E.26) records the residual data.  $\square$



*Definition E.27 (Finite-window marked Costello list).* Fix a finite-window graph/QME system for the local mixed HT brane-defect theory,  $(I, w, M, \mathcal{G}_M)$  in the sense of Definition 8.21. A finite-window marked Costello list

$$\mathcal{G}_M^{\text{mk}}$$

is a finite set of connected brane-defect graphs and local counterterm graphs with the following labels and boundary table.

The admissible vertex labels are exactly the finite local labels visible in the chosen system:

- (L1) external CE labels  $u_{a(t)dt \otimes f}$  and  $c_{B,\rho}$ , with  $f, \rho$  in the Hamiltonian window  $M$ , and with  $B \in \mathcal{D}_c^{\text{reg}}(I)$  or  $B_\bullet \in \mathcal{D}'_{c,\text{WF}}(I; \Gamma, M)$ ;
- (L2) interaction vertices from the finite-window local interaction  $I_{o,M}^w$ ;
- (L3) renormalized graph-weight vertices  $W_{\Gamma,L,w}^{R,M}(B_\bullet^\Gamma)$  and local counterterm vertices  $C_{\Gamma,w}^M(\epsilon; B_\bullet^\Gamma)$  supplied by the system;
- (L4) the central scalar-contact ghost  $\gamma_1$ , only on the scalar-contact boundary vertices already used in Theorem E.12.

The admissible edge labels are the finite-scale local propagator  $P_{\epsilon,L}^{w,M}$ , the corresponding regularized BV contraction  $\Delta_{\epsilon,L}^{w,M}$ , and the boundary Weyl edge whose scalar part on two Hamiltonians is the formal-local scalar cocycle of Proposition E.2,

$$(f, g) \mapsto \omega(f, g) \gamma_1.$$

The open index loops may carry marker tensors, defined below in Definition E.28.

For each  $\Gamma \in \mathcal{G}_M^{\text{mk}}$  one specifies a finite ordered set

$$\mathbf{B}_M(\Gamma) = \{\beta_1, \dots, \beta_s\}$$

of elementary codimension-one boundary operations. If a differential, bracket, collision, or counterterm expansion is a finite sum, each summand is a separate  $\beta_i$ . Each operation has a coefficient  $\varepsilon_{\beta_i} \in \mathbb{C}[[\hbar]]$ , an output graph  $\partial_{\beta_i} \Gamma \in \mathcal{G}_M^{\text{mk}}$ , and a marker-transport map  $\mu_{\beta_i}$ . The local-functional boundary operations are:

- (B1) *field differential*: if a vertex has label  $\ell_v$  and  $Q\ell_v = \sum_\alpha q_\alpha \ell_{v,\alpha}$ , then the summand  $(v, \alpha)$  replaces  $\ell_v$  by  $\ell_{v,\alpha}$  and has  $\varepsilon_{v,\alpha} = q_\alpha$ , including the Koszul sign from moving  $Q$  to  $v$ ;
- (B2) *BV edge contraction*: if an oriented internal edge  $e$  joins vertices  $v_1, v_2$  and

$$\{\ell_{v_1}, \ell_{v_2}\}_{h,M} = \sum_\alpha q_{e,\alpha} \ell_{e,\alpha},$$

then the summand  $(e, \alpha)$  contracts  $e$ , merges the two vertices with label  $\ell_{e,\alpha}$ , and has coefficient  $q_{e,\alpha}$  multiplied by the ordinary BV orientation sign. The bracket with  $I_{o,M}^w$  is this operation with one vertex labelled by  $I_{o,M}^w$ ;

- (B3) *boundary collision/contact*: if an ordered boundary collision stratum  $S = (v_1, \dots, v_k)$  has local expansion

$$\text{Coll}_S(\ell_{v_1}, \dots, \ell_{v_k}) = \sum_\alpha q_{S,\alpha} \ell_{S,\alpha},$$

then  $(S, \alpha)$  collapses  $S$  to one boundary vertex labelled  $\ell_{S, \alpha}$ . For the two-Hamiltonian scalar contact this convention gives

$$\text{Coll}^{\text{sc}}((a, f), (b, g)) = \hbar \omega(f, g) \int_I a(t) b(t) \gamma_{\mathbf{I}}(t) dt,$$

antisymmetric in  $(f, g)$ ;

(B4) *counterterm face*: if the small-diagonal extension of a graph subconfiguration has local counterterm expansion  $C_{\Gamma, w}^M(\epsilon; B_{\bullet}) = \sum_{\alpha} q_{\Gamma, \alpha} C_{\Gamma, \alpha}^M$ , then  $(\Gamma, \alpha)$  replaces that subconfiguration by the local counterterm vertex  $C_{\Gamma, \alpha}^M$ ;

For the Hom-valued graph kernel one adjoins one further operation: the source differential face  $-\kappa d_{\text{CE}}$ , with the CE bracket coefficient included in  $\varepsilon_{\beta_i}$ . After evaluation on fixed CE cycles this extra face evaluates to zero.

The finite list is required to contain every output of (B1)–(B4) that appears in the chosen system, and also the source face when the Hom-valued kernel table is used. If an output graph, coefficient, marker transport, or counterterm vertex lies outside  $\mathcal{G}_M^{\text{mk}}$ , that boundary entry is the omitted datum; the larger specialization stays a construction target.

The orientation line used for signs is

$$\text{or}_M(\Gamma) = \det B_M(\Gamma) \otimes \det E_{\text{int}}(\Gamma) \otimes \bigotimes_v \text{or}(\ell_v).$$

The first determinant gives the incidence sign in the marked differential below. The edge, vertex, BV, and CE Koszul signs are included in the coefficients  $\varepsilon_{\beta_i}$ .

The list is *codimension-two closed* if, for every  $\beta_i \neq \beta_j$ , the two composites are again entries of the finite list and, after the canonical identification of output orientation lines,

$$\partial_{\beta_j | \beta_i} \partial_{\beta_i} \Gamma = \partial_{\beta_i | \beta_j} \partial_{\beta_j} \Gamma,$$

$$\varepsilon_{\beta_i} \varepsilon_{\beta_j | \beta_i} = \varepsilon_{\beta_j} \varepsilon_{\beta_i | \beta_j}, \quad \mu_{\beta_j | \beta_i} \mu_{\beta_i} = \mu_{\beta_i | \beta_j} \mu_{\beta_j}.$$

Here  $\beta_j | \beta_i$  denotes the surviving operation induced by  $\beta_j$  after applying  $\beta_i$ . Thus, from a seed graph  $\Gamma$ , the finite window consists of the finitely many faces obtained by deleting subsets of the finite set  $B_M(\Gamma)$ :

$$\mathcal{G}_M^{\text{mk}} = \{ \partial_J \Gamma \mid \Gamma \in \mathcal{S}_M, J \subset B_M(\Gamma) \}, \quad \partial_J = \partial_{\beta_{j_k}} \cdots \partial_{\beta_{j_1}}.$$

Here  $\mathcal{S}_M$  is the finite seed set of graph and counterterm graphs, and the codimension-two closure equations make  $\partial_J$  independent of the chosen order of  $J$ . A defect in this displayed equality is precisely the finite graph-list obstruction.

*Definition E.28 (Oriented full-equivariant marked boundary complex).* Let  $V = \mathbb{C}^{N|N}$ . For  $m \geq 1$ , let

$$\mathcal{M}_m^{\text{full}}(V) \subset (\text{End}(V)^{\otimes m})^{\mathfrak{gl}(V)}$$

be the finite span of the signed Brauer permutation tensors  $\mathfrak{C}_{\sigma}$ ,  $\sigma \in S_m$ . Here  $\mathfrak{C}_{\sigma}$  is the string-diagram tensor obtained by writing  $\text{End}(V) = V \otimes V^{\vee}$ , pairing the  $V^{\vee}$  factor in the  $i$ -th copy with the  $V$  factor in the  $\sigma(i)$ -th copy, and using the Koszul signs of the symmetric monoidal category of super vector spaces. For  $m = 0$  the marker is the empty word.

A full-equivariant marked graph/counterterm generator in a finite window  $\mathcal{M}$  is a tuple

$$X_{\Gamma, o, \mathbf{m}, \mathcal{M}},$$

where  $\Gamma \in \mathcal{G}_{\mathcal{M}}^{\text{mk}}$  for a finite-window marked Costello list of Definition E.27,  $o$  is an orientation of the finite set  $\mathbf{B}_{\mathcal{M}}(\Gamma)$ , and  $\mathbf{m} = (\mathbf{m}_L)_L$  assigns to each marked scalar open index loop  $L$  a tensor  $\mathbf{m}_L \in \mathcal{M}_{m_L}^{\text{full}}(V)$ .

If

$$o = \beta_1 \wedge \cdots \wedge \beta_s$$

is the ordered orientation of  $\mathbf{B}_{\mathcal{M}}(\Gamma)$ , put  $o/\beta_i = \beta_1 \wedge \cdots \wedge \widehat{\beta_i} \cdots \wedge \beta_s$ . The marker-transport map

$$\mu_{\beta_i} : \bigotimes_L \mathcal{M}_{m_L}^{\text{full}}(V) \longrightarrow \bigotimes_{L'} \mathcal{M}_{m'_{L'}}^{\text{full}}(V)$$

is obtained by the signed evaluation, coevaluation, braiding, loop splitting, and loop concatenation maps that define the open index contractions. Concretely, loop merging sends two cyclic words to their boundary-ordered concatenation with the Koszul braiding sign; loop splitting inserts the coevaluation tensor at the split points; deletion of a closed marked loop applies the cyclic supertrace. The signed marked differential is

$$d_{\text{mk}} X_{\Gamma, o, \mathbf{m}, \mathcal{M}} = \sum_{i=1}^s (-1)^{i-1} \varepsilon_{\beta_i} X_{\partial_{\beta_i} \Gamma, o/\beta_i, \mu_{\beta_i}(\mathbf{m}), \mathcal{M}}.$$

Its scalar-contact filtration is

$$p(\Gamma, \mathbf{m}) = \#\{\text{scalar-contact open loops in } \Gamma\}, \quad F^p = \overline{\text{Span}}\{X_{\Gamma, o, \mathbf{m}, \mathcal{M}} : p(\Gamma, \mathbf{m}) \geq p\}.$$

For an elementary boundary operation set

$$r(\beta_i) = p(\Gamma, \mathbf{m}) - p(\partial_{\beta_i} \Gamma, \mu_{\beta_i}(\mathbf{m})).$$

The first filtration-lowering marked boundary is therefore

$$[d_{\text{mk}} X_{\Gamma, o, \mathbf{m}, \mathcal{M}}]_{F^{p-r}/F^{p-r+1}} = \sum_{i: r(\beta_i)=r} (-1)^{i-1} \varepsilon_{\beta_i} X_{\partial_{\beta_i} \Gamma, o/\beta_i, \mu_{\beta_i}(\mathbf{m}), \mathcal{M}}.$$

**LEMMA E.29 (Incidence signs).** If the finite marked Costello list is codimension-two closed in the sense of Definition E.27, then  $d_{\text{mk}}^2 = 0$ .

*Proof.* This is the cellular boundary sign computation. The codimension-two output obtained from  $\beta_i$  and  $\beta_j$ ,  $i < j$ , appears twice. The order  $\beta_i$  then  $\beta_j$  contributes

$$(-1)^{i-1} (-1)^{j-2} \varepsilon_{\beta_i} \varepsilon_{\beta_j | \beta_i},$$

because  $\beta_i$  has been removed before  $\beta_j$  is counted. The order  $\beta_j$  then  $\beta_i$  contributes

$$(-1)^{j-1} (-1)^{i-1} \varepsilon_{\beta_j} \varepsilon_{\beta_i | \beta_j}.$$

The incidence signs differ by a minus sign, while (E.27) identifies the coefficients, output graph, and marker tensor. Hence the two terms cancel. Summing over all unordered pairs gives  $d_{\text{mk}}^2 = 0$ .  $\square$

PROPOSITION E.30 (*Finite-window marked-complex compatibility*). Let  $(I, w, M, \mathcal{G}_M)$  be a finite-window graph/QME system for the local mixed HT brane-defect theory and let  $\mathcal{G}_M^{\text{mk}}$  be a codimension-two closed finite-window marked Costello list whose labels are the admissible local labels of Definition E.27. Let

$$\mathfrak{G}_{\text{mk}, \text{full}, M}^\bullet(I)$$

be the completed span of the full-equivariant marked generators  $X_{\Gamma, o, \mathbf{m}, M}$ . Then

$$\mathfrak{G}_{\text{mk}, \text{full}, M}^\bullet(I) \subset \mathfrak{Q}_{\mathcal{G}, M}^\bullet(I)$$

and  $d_{\text{mk}}$  is the restriction of the finite-window differential  $d_M = Q + \{I_{o, M}^w, -\}_{\hbar, M}$  on the local-functional table. On the Hom-valued kernel table, adjoining the source face makes the same formula the convolution differential  $d_{\mathbf{K}} = d_M - (-1)^{|\cdot|}(\cdot)d_{\text{CE}}$  before evaluation. The graph kernel  $\kappa_{\mathcal{G}, w, I}^M$  and the local counterterms of the system land in this marked complex exactly when their graph components and counterterm faces occur in  $\mathcal{G}_M^{\text{mk}}$ . Truncation  $\pi_{M, N}$  and admissible weight transport  $R_{w, w'}^M$  preserve the marked complex precisely when they send the finite boundary table, coefficients, and marker transports to the corresponding lower-window or transported table.

*Proof.* The vertex and edge labels are those of Definition 8.21: regular-density or wavefront-admissible graph labels, finite-window propagators, renormalized weights, and local counterterms. Hence the represented local functionals lie in  $\mathfrak{Q}_{\mathcal{G}, M}^\bullet(I)$ . The boundary table (B1)–(B4) is exactly the expansion of  $Q$ , the BV bracket with  $I_{o, M}^w$ , collision/contact faces, and Costello counterterm faces on these generators. Thus the signed sum (E.28) is  $d_M$  on the local functional span. In the convolution kernel before CE evaluation, the additional source face is the missing source differential term in  $d_{\mathbf{K}} = d_M - (-1)^{|\cdot|}(\cdot)d_{\text{CE}}$ ; for the degree-zero kernel this is  $-\kappa d_{\text{CE}}$ .

The inclusion of the graph kernel and counterterms is tautological from the same finite label list. Each omitted graph component or counterterm face is an explicit missing datum for the equality with  $d_M$ . The finite-window system requires compatibility of graph weights, counterterms, scalar projections, and kernels with  $\pi_{M, N}$  and  $R_{w, w'}^M$ ; applying this requirement to each entry of the finite boundary table gives the stated preservation criterion.  $\square$

LEMMA E.31 (*Cyclic supertrace of full-equivariant marker tensors*). Let  $\tau_m = (1\ 2\ \cdots\ m)$ . For the signed Brauer permutation tensor  $\mathfrak{C}_\sigma \in \mathcal{M}_m^{\text{full}}(V)$ ,

$$\text{Str}_{\text{cyc}}(\mathfrak{C}_\sigma) = (\text{sdim } V)^{c(\tau_m^{-1}\sigma)},$$

where  $c(\pi)$  is the number of cycles of the permutation  $\pi$ . Hence, for  $V = \mathbb{C}^{N|N}$ ,

$$\text{Str}_{\text{cyc}}(\mathbf{m}) = 0 \quad \text{for every } \mathbf{m} \in \mathcal{M}_m^{\text{full}}(V)$$

and also for the empty marker word.

*Proof.* Compose the string diagram for  $\mathfrak{C}_\sigma$  with the cyclic supertrace functional  $(T_1, \dots, T_m) \mapsto \text{Str}(T_1 \cdots T_m)$ . The cyclic multiplication identifies the output of the  $i$ -th endomorphism with the input of the  $\tau_m(i)$ -th endomorphism, while  $\mathfrak{C}_\sigma$  identifies it with the input of the  $\sigma(i)$ -th endomorphism. The resulting closed diagram has  $c(\tau_m^{-1}\sigma)$  closed supertraces. Each closed supertrace evaluates to  $\text{Str}(\text{id}_V) = \text{sdim } V$ . The signs in  $\mathfrak{C}_\sigma$  are exactly the Koszul signs needed for this diagrammatic evaluation, giving (E.31). Since  $\text{sdim } \mathbb{C}^{N|N} = 0$ , every nonempty full-equivariant marker tensor has zero cyclic supertrace. The empty word gives the identity-loop coefficient  $\text{Str}(\text{id}_V) = 0$ .  $\square$

**THEOREM E.32** (*Full-equivariant marked scalar shadow vanishing*). Let  $V = \mathbb{C}^{N|N}$ . Let  $\mathfrak{G}_{\text{mk,full}}^\bullet(I)$  be a complete finite-window inverse-limit graph/counterterm complex for the formal-local mixed HT brane-defect theory generated windowwise by codimension-two closed finite-window marked Costello lists of Definition E.27 and by the full-equivariant marked generators of Definition E.28, compatible with truncation and admissible weight transport as in Proposition E.30. This is a finite-list statement in every window. Then the marked preservation obstruction has zero cyclic-supertrace shadow:

$$\text{Str}_{\text{cyc}}(\mu_{\beta_i}(\mathbf{m})) = 0 \quad \text{for every elementary boundary term } \beta_i.$$

Consequently, for every homogeneous  $X_{\Gamma,o,\mathbf{m},M} \in F^p$ , the first scalar CE shadow is

$$\hbar \sum_{i: r(\beta_i)=r} (-1)^{i-1} \varepsilon_{\beta_i} \text{Str}_{\text{cyc}}(\mu_{\beta_i}(\mathbf{m})) \omega(f_{\partial_{\beta_i}\Gamma}, g_{\partial_{\beta_i}\Gamma}) \gamma_{\mathbf{I}} = 0,$$

and its cohomology class on the  $[\bar{c}]$ -line is

$$\hbar \sum_{i: r(\beta_i)=r} (-1)^{i-1} \varepsilon_{\beta_i} \text{Str}_{\text{cyc}}(\mu_{\beta_i}(\mathbf{m})) [\bar{c}] \gamma_{\mathbf{I}} = 0.$$

Thus the zero scalar projection is a filtered chain map on  $\mathfrak{G}_{\text{mk,full}}^\bullet(I)$ , and all scalar projection obstruction classes vanish on this full-equivariant marked graph/counterterm complex.

*Proof.* The marker transport maps  $\mu_{\beta_i}$  are built from the evaluation, coevaluation, braiding, loop splitting, and loop concatenation maps of the  $\mathfrak{gl}(V)$ -module  $V$ . Therefore they send full-equivariant signed Brauer marker tensors to full-equivariant signed Brauer marker tensors. Lemma E.31 gives zero cyclic supertrace for every such output tensor when  $V = \mathbb{C}^{N|N}$ .

Formula (E.28) gives the signed first filtration-lowering boundary. Applying the one-loop scalar computation of Proposition E.23 to each boundary term gives exactly the displayed scalar CE representative. Every coefficient in that sum is zero by the preceding paragraph. Passing to cohomology replaces  $\omega$  by the nonzero class  $[\bar{c}]$ , but it is multiplied by the same zero coefficient. Thus  $\widehat{\sigma} = 0$  satisfies  $d_{\text{CE}} \widehat{\sigma} - \widehat{\sigma} d_{\text{mk}} = 0$ . Proposition E.30 identifies  $d_{\text{mk}}$  with the restricted finite-window QME differential on the chosen finite table. Hence this is the required filtered chain projection, and Lemma E.4 kills the scalar projection tower on the complex.  $\square$

**Definition E.33** (*Non-scalar kernel complex for a marked finite window*). Fix a finite-window graph/QME system  $(I, w, M, \mathcal{G}_M)$ , a codimension-two closed marked Costello list  $\mathcal{G}_M^{\text{mk}}$ , and a scalar projection

$$\widehat{\sigma}_{\text{sc},M}: \mathfrak{Q}_{\mathcal{G},M}^\bullet(I) \longrightarrow S_M^\bullet, \quad S_M^\bullet = C_{\text{Lie}}^\bullet(\bar{A}; \mathbb{C}\gamma_{\mathbf{I}})[1][[\hbar]].$$

The non-scalar obstruction complex in that window is

$$\mathcal{K}_{\text{ns},M}^\bullet(I) = \ker \widehat{\sigma}_{\text{sc},M}, \quad d_{\text{ns},M} = d_M|_{\mathcal{K}_{\text{ns},M}}.$$

It is a complex precisely when  $\widehat{\sigma}_{\text{sc},M}$  is a filtered chain map. In the full-equivariant marked complex of Theorem E.32 one has  $\widehat{\sigma}_{\text{sc},M} = 0$ , hence

$$\mathcal{K}_{\text{ns},M}^\bullet(I) = \mathfrak{G}_{\text{mk,full},M}^\bullet(I).$$

Let  $A_{n,M}^{\text{fw}}$  be a valid row array in the sense of Definition 8.24. If the scalar condition vanishes,

$$\sum_{\alpha \in A_{n,M}^{\text{fw}}} S_{\alpha,M} = 0,$$

the order- $n$  kernel residual is the scalar-zero row sum

$$R_{n,M}^{\text{ns}} = \sum_{\alpha \in \mathbf{A}_{n,M}^{\text{fw}}} R_{\alpha,M}^{\text{row}} \in \mathcal{K}_{\text{ns},M}^{\text{r}}(I).$$

Its obstruction class is

$$\mathfrak{o}_{n,M}^{\text{ns}} = [R_{n,M}^{\text{ns}}] \in H^1(\mathcal{K}_{\text{ns},M}^{\bullet}(I), d_{\text{ns},M}).$$

A degree-zero counterterm  $C_{n,M} \in \mathcal{K}_{\text{ns},M}^{\circ}(I)$  exists at this order if and only if

$$d_M C_{n,M} = -R_{n,M}^{\text{ns}}, \quad \mathfrak{o}_{n,M}^{\text{ns}} = 0.$$

*Definition E.34 (Finite-window realization complex).* A Costello/QME realization complex for the mixed HT brane-defect model on

$$\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}^2}_{\text{o,hol}}$$

is the following finite-window datum. First, one fixes a graph/QME system  $(I, w, M, \mathcal{G}_M)$  in the sense of Definition 8.21, a codimension-two closed marked list  $\mathcal{G}_M^{\text{mk}}$  in the sense of Definition E.27, and the completed local-functional complex

$$\mathfrak{Q}_{\mathcal{G},M}^{\bullet}(I) = \left( \mathcal{O}_{\text{loc},\mathcal{G},M}^{\text{bd}}(\mathcal{E}_w^M|_I; A_{\partial,\hbar}^{\text{pp},w,M}(I)), d_M \right), \quad d_M = Q + \{I_{\text{o},M}^w, -\}_{\hbar,M}.$$

The topology is the complete topology generated by the  $\hbar$ -adic, scalar-contact, and finite-window filtrations. The complex is part of the datum: every  $d_M$ -boundary, collision face, BV bracket row, and local counterterm face needed for closure must occur in the finite table.

Second, the source is the labelled finite CE object  $C_{\text{lab}}^{\bullet}(\mathcal{G}_M)$ . The Hom-valued kernel complex is

$$\mathbf{K}_{\mathcal{G},M}^{\bullet}(I) = \text{Hom}_{\text{cont}}\left(C_{\text{lab}}^{\bullet}(\mathcal{G}_M), \mathfrak{Q}_{\mathcal{G},M}^{\bullet}(I)\right),$$

with

$$d_{\mathbf{K},M} T = d_M T - (-1)^{|T|} T d_{\text{CE}}, \quad \text{Curv}_{\mathbf{K},M}(\kappa) = d_{\mathbf{K},M} \kappa + \frac{1}{2} [\kappa, \kappa]_{\mathbf{K},M}.$$

Thus the source-face term is part of the Hom differential. For a degree-zero kernel it contributes  $-\kappa d_{\text{CE}}$ .

Third, the kernel has a fixed zero-edge part:

$$\kappa_{\hbar,w,I}^{\text{coef},M}(\iota_{a(t)dt \otimes f}) = \iota_I(B_f^{w,M}(a)), \quad \kappa_{\hbar,w,I}^{\text{coef},M}(c_{B,\rho}) = \iota_I(\Theta_{\rho}^{w,M}(B)).$$

Here  $a \in C_c^{\infty}(I)$ ,  $B$  is a regular compactly supported density or a graphwise wavefront-admissible current for the chosen finite row, and  $f, \rho$  lie in the window  $M$ . Positive-edge components are exactly the renormalized graph weights  $K_{\Gamma}^M = W_{\Gamma,L,w}^{R,M}(B_{\bullet}^{\Gamma})$  supplied by the system. A graph theorem for arbitrary  $B \in \mathcal{D}'_c(I)$  is additional analytic input.

Fourth, one supplies a filtered chain scalar projection

$$\widehat{\sigma}_{\text{sc},M}: \mathfrak{Q}_{\mathcal{G},M}^{\bullet}(I) \longrightarrow C_{\text{Lie}}^{\bullet}(\bar{A}; \mathbb{C}\gamma_1)[1][[\hbar]],$$

and puts

$$\mathcal{K}_{\text{ns},M}^{\bullet}(I) = \ker \widehat{\sigma}_{\text{sc},M}.$$

The ambient QME complex is always denoted  $\mathfrak{Q}_{\mathcal{G},M}^{\bullet}(I)$ . If the windowed scalar-zero QME notation  $\mathcal{Q}_{w,M}^{\bullet}(I)$  is used for the same kernel, then

$$\mathcal{Q}_{w,M}^{\bullet}(I) = \mathcal{K}_{\text{ns},M}^{\bullet}(I) = \ker \widehat{\sigma}_{\text{sc},M}.$$

The non-scalar kernel complex is formed after this projection is a chain map.

THEOREM E.35 (*Constructive non-scalar Costello/QME realization criterion*). Work in a finite-window realization complex of Definition E.34. Suppose that the QME has been solved through order  $\hbar^{n-1}$  by counterterms  $C_{<n,M}$ . The order- $n$  realization problem is decided by the following finite row construction.

For every valid row array  $A_{n,M}^{\text{fw}}$  of Definition 8.24, the rows are precisely

$$\begin{aligned} R_{d_{\Gamma},M}^{\text{row}} &= \varepsilon_{\Gamma} d_M K_{\Gamma}^M, \\ R_{\text{CE}(\Gamma,\nu),M}^{\text{row}} &= -\varepsilon_{\Gamma,\nu}^{\text{CE}} K_{\Gamma}^M(\zeta_{M,\nu}), \\ R_{b(\Gamma_1,\Gamma_2),M}^{\text{row}} &= \frac{1}{2} \varepsilon_{\Gamma_1,\Gamma_2} \{K_{\Gamma_1}^M, K_{\Gamma_2}^M\}_{\hbar,M}, \end{aligned}$$

together with the homogeneous lower-counterterm rows in

$$[\hbar^n] \left( d_M C_{<n,M} + \frac{1}{2} \{C_{<n,M}, C_{<n,M}\}_{\hbar,M} \right).$$

Their scalar conditions are

$$S_{\alpha,M} = \widehat{\sigma}_{\text{sc},M}(R_{\alpha,M}^{\text{row}}),$$

and the scalar condition for the order is the single equation

$$\mathfrak{s}_{n,M} = \sum_{\alpha \in A_{n,M}^{\text{fw}}} S_{\alpha,M} = 0.$$

The non-scalar problem at order  $n$  begins after (E.35) holds. Then

$$R_{n,M}^{\text{ns}} = \sum_{\alpha \in A_{n,M}^{\text{fw}}} R_{\alpha,M}^{\text{row}} \in \mathcal{K}_{\text{ns},M}^{\text{i}}(I)$$

is a cocycle whenever the finite row table is codimension-two closed and the lower orders have been solved. Its obstruction class is

$$\mathfrak{o}_{n,M}^{\text{ns}} = [R_{n,M}^{\text{ns}}] \in H^1(\mathcal{K}_{\text{ns},M}^{\bullet}(I), d_{\text{ns},M}).$$

Choose degree-one row coordinates  $\rho_1^M, \dots, \rho_a^M$  and degree-zero counterterm candidates  $\eta_1^M, \dots, \eta_b^M$  inside the same finite system, and write

$$R_{n,M}^{\text{ns}} = \sum_i r_i^M \rho_i^M, \quad d_{\text{ns},M} \eta_j^M = \sum_i A_{ij}^M \rho_i^M.$$

A fixed-window counterterm in the supplied span exists exactly when

$$A^M c = -r^M$$

is solvable. In the obstruction case, any covector

$$\ell A^M = 0, \quad \ell(r^M) \neq 0$$

is a separating covector proving that  $\mathfrak{o}_{n,M}^{\text{ns}} \neq 0$  in the displayed finite row subcomplex. A compatible finite-window tower also requires truncation matrices  $T^{M,N}$  on the rows and  $P^{M,N}$  on candidate primitives satisfying

$$T^{M,N} r^M = r^N, \quad T^{M,N} A^M = A^N P^{M,N},$$

and the Roos class

$$[P^{M,N}c_M - c_N] \in H^0(\mathcal{K}_{\text{ns},N}^\bullet(I))$$

must vanish.

Iterating these finite-window tests for all  $n$ , with graphwise regular-density or wavefront-admissible kernel data and with compatible counterterms, is equivalent to

$$\text{Curv}_{\mathbf{K}}(\kappa_{\mathcal{G},w,I} + C_{h,w}) = 0$$

in the completed brane-defect kernel complex. A larger Costello specialization requires its missing graph weight, source face, scalar condition, primitive candidate, or truncation matrix as explicit input to the criterion.

*Proof.* The first three displayed row formulas are the homogeneous terms in

$$d_M \kappa_{\mathcal{G},w,I}^M - \kappa_{\mathcal{G},w,I}^M d_{\text{CE}} + \frac{1}{2} [\kappa_{\mathcal{G},w,I}^M, \kappa_{\mathcal{G},w,I}^M]_{\mathbf{K},M}.$$

The minus sign in the source-face row is the term  $-\kappa d_{\text{CE}}$  in Definition E.34. The last displayed family is the order- $n$  contribution of lower counterterms to the same curved Maurer–Cartan equation.

Applying the chain scalar projection gives the scalar condition. A nonzero scalar condition places the residual outside  $\ker \widehat{\sigma}_{\text{sc},M}$ . A zero scalar condition places the residual in the kernel complex. Codimension-two closure is exactly the finite-row form of the Bianchi identity: the two ways of taking two elementary faces have opposite incidence signs and the same output graph, coefficient, and marker transport. Since lower orders are solved, this makes  $R_{n,M}^{\text{ns}}$  a  $d_{\text{ns},M}$ -cocycle.

A counterterm  $C_{n,M}$  changes the residual by  $d_{\text{ns},M} C_{n,M}$ . Expanding  $C_{n,M}$  in the supplied degree-zero row basis gives the linear system (E.35). The solvability obstruction is the residual vector class in the cokernel of  $A^M$ ; a left-null covector with nonzero value on  $r^M$  is precisely the finite-dimensional cokernel separation. The two truncation identities say that both residuals and boundaries commute with finite-window projection. After windowwise primitives are chosen, their compatibility with truncation is measured by the displayed Roos 1-cocycle in  $H^0$ . Thus the fixed-window  $H^1$  class and the Roos primitive class are exactly the obstruction pair. Passing to all orders gives the completed curvature equation.  $\square$

**PROPOSITION E.36** (*Closed rows and the first supplied open source-face row*). In the full-equivariant branch  $V = \mathbb{C}^{N|N}$ , the one-cross two-Hamiltonian scalar-contact row  $\alpha_{\text{sc}}$  with inputs  $(z_1, z_2)$  has

$$R_{\alpha_{\text{sc}},M}^{\text{row}} = 0, \quad S_{\alpha_{\text{sc}},M} = 0, \quad C_{\alpha_{\text{sc}},M} = 0.$$

The labelled scalar-zero row  $\alpha_{\text{II}}$  with inputs  $(z_1, z_1)$  has

$$R_{\alpha_{\text{II}},M}^{\text{row}} = \text{Str}_{\text{cyc}}(\mu_{\beta_{\text{II}}}(\mathbf{m})) \omega(z_1, z_1) \int_I a(t)b(t)\gamma_{\mathbf{I}}(t) dt = 0,$$

$$S_{\alpha_{\text{II}},M} = 0, \quad C_{\alpha_{\text{II}},M} = 0,$$

with truncation  $t_{\alpha_{\text{II}},M}^{M,N} = 1$  when the labelled zero row is retained and the lower window still contains  $z_1$ .

The first nonzero scalar-zero source-face row entering the row machinery is the order-three symbolic row  $\theta_3 = \text{CE}(\Theta_3, \nu_3)$ , with

$$R_{\theta_3,M}^{\text{row}} = -K_{\Theta_3}^M(\zeta_{M,\nu_3}) =: E_{\theta_3,M}, \quad S_{\theta_3,M} = 0.$$



If the row-closure datum  $d_{\text{ns},M}E_{\theta_3,M} = \mathbf{o}$  is supplied, it defines

$$\mathbf{o}_{\theta_3,M}^{\text{ns}} = [E_{\theta_3,M}] \in H^1(\mathcal{K}_{\text{ns},M}^\bullet(I), d_{\text{ns},M}).$$

In the finite row subcomplex currently supplied for this row,

$$\mathcal{K}_{\theta_3,M}^{\mathbf{o}} = \mathbf{o}, \quad \mathcal{K}_{\theta_3,M}^{\mathbf{i}} = \mathbb{C}E_{\theta_3,M}, \quad d_{\theta_3,M} = \mathbf{o},$$

the covector

$$\ell_{\theta_3}(E_{\theta_3,M}) = \mathbf{i}$$

detects a nonzero finite-row obstruction. To remove it one must supply either a degree-zero local counterterm  $C_{\theta_3,M}$  with

$$d_{\text{ns},M}C_{\theta_3,M} = -E_{\theta_3,M},$$

compatible under finite-window truncation, or additional Hom-valued companion rows whose sum changes the residual vector. These data make the order-three larger system a solved QME row.

*Proof.* The row  $\alpha_{\text{sc}}$  is zero as a local functional by the cyclic supertrace computation of Lemma E.31; this is the closed row of Proposition 8.25. The row  $\alpha_{\text{II}}$  is zero for the independent reason

$$\omega(z_{\text{I}}, z_{\text{I}}) = [\{z_{\text{I}}, z_{\text{I}}\}_{\Omega_{\text{loc}}}]_{\mathbf{o}} = \mathbf{o},$$

which is Proposition E.45. In both cases the zero counterterm is a primitive and the stated truncation identities send zero rows to zero rows.

The formula for  $\theta_3$  is the source-face row  $-\kappa d_{\text{CE}}$  from Theorem E.35, with the concrete row labels of Proposition 8.32. Its scalar condition is zero in the full-equivariant marked complex. In the one-row finite complex, the degree-zero candidate primitive space is zero, so the boundary matrix is the  $\mathbf{i} \times \mathbf{o}$  zero matrix and the residual vector is the basis vector  $E_{\theta_3,M}$ . The displayed covector annihilates the image of the differential and pairs nontrivially with the residual. This proves nonvanishing only in the supplied finite row subcomplex. The displayed  $C_{\theta_3,M}$  or companion-row datum is exactly what enlarges the complex enough to kill the row.  $\square$

**PROPOSITION E.37** (*Bianchi-exposed lower condition for  $\theta_3$* ). In the lower window  $N = 2$ , let  $E_{\nu_{\text{o}_2}}^2$  and  $E_{\nu_{\text{z}_0}}^2$  be the two transported source-face rows attached to the  $\theta_3$  source differential. The canonical source-coordinate candidate is

$$\zeta_2^{\mathbf{o}} = u_{a,H_{\text{i},\mathbf{o}}} u_{b,H_{\mathbf{o},\mathbf{i}}}, \quad D_{\text{o}_2,\text{z}_0}^2 = K_{\Theta_3}^2(\zeta_2^{\mathbf{o}}),$$

with

$$d_{\text{CE}}\zeta_2^{\mathbf{o}} = 2\zeta_{\nu_{\text{o}_2}} - 2\zeta_{\nu_{\text{z}_0}}.$$

The Hom-valued Bianchi defect is

$$\mathbf{b}_{\text{o}_2,\text{z}_0}^2 = d_{\text{ns},2}K_{\Theta_3}^2(\zeta_2^{\mathbf{o}}) - K_{\Theta_3}^2(d_{\text{CE}}\zeta_2^{\mathbf{o}}).$$

In the Bianchi-exposed basis

$$(E_{\nu_{\text{o}_2}}^2, E_{\nu_{\text{z}_0}}^2, \mathbf{b}_{\text{o}_2,\text{z}_0}^2),$$

the current source-coordinate column and the lower transported residual are

$$A_D^2 = (-2, 2, \mathbf{i})^T, \quad r_2 = (2, -2, \mathbf{o})^T.$$

Hence

$$A_D^2 c = -r_2$$

has a cokernel obstruction in the supplied lower local-functional row complex. The normalized cokernel

$$\tilde{\lambda}_{o_2, \mathfrak{b}} = \frac{1}{2}(E_{\nu_{o_2}}^2)^* + (\mathfrak{b}_{o_2, 2o}^2)^*$$

satisfies

$$\tilde{\lambda}_{o_2, \mathfrak{b}} A_D^2 = 0, \quad \tilde{\lambda}_{o_2, \mathfrak{b}}(r_2) = 1.$$

The missing Bianchi-counterterm target is

$$-\mathfrak{b}_{o_2, 2o}^2 = (0, 0, -1)^T,$$

and it has nonzero cokernel pairing against the current column  $A_D^2$ , since

$$\tilde{\lambda}_{o_2, \mathfrak{b}}((0, 0, -1)^T) = -1.$$

A repair must therefore add a genuine scalar-zero local-functional column

$$A_B^2 = (0, 0, -1)^T, \quad d_{\text{ns}, 2} B_{o_2, 2o}^2 = -\mathfrak{b}_{o_2, 2o}^2, \quad \widehat{\sigma}_{\text{sc}, 2}(B_{o_2, 2o}^2) = 0,$$

with the required locality or wavefront admissibility. If such  $B_{o_2, 2o}^2$  is supplied, then

$$d_{\text{ns}, 2}(D_{o_2, 2o}^2 + B_{o_2, 2o}^2) = -2E_{\nu_{o_2}}^2 + 2E_{\nu_{2o}}^2 = -r_2.$$

For a compatible finite-window tower, write  $\mathfrak{b}^M$  for the corresponding Bianchi defect in window  $M$ . The transport obstruction is

$$\Delta_{M, N}^1 := -\pi_{M, N} \mathfrak{b}^M + \mathfrak{b}^N.$$

A family  $B^M$  transports the fixed-window repair only when

$$\Delta_{M, N}^1 = d_{\text{ns}, N}(\pi_{M, N} B^M - B^N),$$

possibly after an explicitly supplied scalar-zero transport correction. Thus the  $H^1$  class of  $\Delta_{M, N}^1$  must vanish first, and the resulting primitive-compatibility class in

$$\varprojlim_N^1 H^0(\mathcal{K}_{\text{ns}, N}^\bullet(I))$$

must vanish next. These two conditions promote the lower  $\theta_3$  Bianchi repair from a fixed-window matrix target to a solved Costello/QME counterterm tower.

*Proof.* The source differential gives the two lower source-face rows with coefficients 2 and  $-2$ . Applying  $K_{\Theta_3}^2$  becomes a chain map after its Hom-valued Bianchi defect is killed, and

$$d_{\text{ns}, 2} D_{o_2, 2o}^2 = -2E_{\nu_{o_2}}^2 + 2E_{\nu_{2o}}^2 + \mathfrak{b}_{o_2, 2o}^2,$$

which is the column  $A_D^2 = (-2, 2, 1)^T$  in the displayed basis. The residual transported from the lower two-face row is  $r_2 = (2, -2, 0)^T$ . If  $A_D^2 c = -r_2 = (-2, 2, 0)^T$ , the first two coordinates force  $c = 1$ ,

while the Bianchi coordinate forces  $c = 0$ . Thus the equation is inconsistent. The same conclusion is the cokernel calculation:  $\tilde{\lambda}_{02,b}$  annihilates  $A_D^2$  and pairs nontrivially with  $r_2$ .

The target  $(0, 0, -1)^T$  is the boundary column required to remove only the Bianchi coordinate. Since the same cokernel evaluates to  $-1$  on it, this boundary is a new scalar-zero local-functional column. Adding a scalar-zero local functional  $B_{02,20}^2$  with the displayed boundary produces  $A_D^2 + A_B^2 = (-2, 2, 0)^T = -r_2$ , hence the lower residual is killed. This is only a formal finite-window repair until the local functional and its admissibility are supplied.

For  $M \geq N$ , applying  $d_{ns,N}$  to  $\pi_{M,N} B^M - B^N$  gives the difference between the transported Bianchi boundary and the  $N$ -window Bianchi boundary, namely  $-\pi_{M,N} \mathbf{b}^M + \mathbf{b}^N$ . Therefore the transported family exists after this degree-one class vanishes in  $H^1(\mathcal{K}_{ns,N}^\bullet(I))$ ; after choosing primitives, their compatibility is measured by the usual Roos  $\varprojlim^1 H^0$  obstruction.  $\square$

**THEOREM E.38** (*First non-scalar finite-window  $H^1$  at the  $\theta_3$  row*). Work in the full-equivariant marked finite-window complex with  $V = \mathbb{C}^{N|N}$  and the supplied minimal seed list at order  $\leq 2$  of Theorem E.47. Adjoin only the order-three CE-source row  $\theta_3$  of Proposition E.36. The non-scalar kernel subcomplex at  $\theta_3$  is the supplied finite row complex

$$\mathcal{K}_{\theta_3,M}^0 = 0, \quad \mathcal{K}_{\theta_3,M}^1 = \mathbb{C} E_{\theta_3,M}, \quad d_{\theta_3,M} = 0.$$

The finite-window  $H^1$  splits into two cases.

- (a) *Counterterm omitted*. In the one-row complex above, the  $d_{ns,M}$ -cohomology in degree one is

$$H^1(\mathcal{K}_{\theta_3,M}^\bullet, d_{\theta_3,M}) = \mathbb{C} \cdot E_{\theta_3,M} \neq 0,$$

and the obstruction class is the basis vector itself,

$$[\text{Ob}_{w,\partial,h}^{\text{red}}]_{\theta_3,M} = [E_{\theta_3,M}] \neq 0,$$

detected by the unique covector  $\ell_{\theta_3}(E_{\theta_3,M}) = 1$  on the row basis. The supplied finite complex has zero degree-zero row space; the linear system  $A^M c = -r^M$  of Theorem E.35 has empty column matrix and residual  $r^M = E_{\theta_3,M}$ , so it is unsolvable.

- (b) *Scalar-zero local counterterm  $C_{\theta_3,M}$  supplied*. Suppose a degree-zero local functional  $C_{\theta_3,M} \in \mathcal{K}_{\theta_3,M}^0$  with  $\widehat{\tau}_{sc,M}(C_{\theta_3,M}) = 0$  is supplied so that, in the enlarged finite complex

$$\widetilde{\mathcal{K}}_{\theta_3,M}^0 = \mathbb{C} C_{\theta_3,M}, \quad \widetilde{\mathcal{K}}_{\theta_3,M}^1 = \mathbb{C} E_{\theta_3,M}, \quad \widetilde{d} C_{\theta_3,M} = -E_{\theta_3,M},$$

the differential matrix is the scalar  $-1$ . Then

$$H^1(\widetilde{\mathcal{K}}_{\theta_3,M}^\bullet, \widetilde{d}) = 0, \quad [\text{Ob}_{w,\partial,h}^{\text{red}}]_{\theta_3,M} = 0,$$

and the unique counterterm  $c = 1$  (i.e. adopting  $C_{\theta_3,M}$  with coefficient one in the QME expansion) solves the row equation  $\widetilde{A}^M c = -r^M = -E_{\theta_3,M}$ . Tower compatibility holds with the identity primitive transport  $P^{M,N} = 1$ , zero Roos class  $[P^{M,N} c_M - c_N] = 0$ , and zero Bianchi tower obstruction  $\Delta_{M,N}^1 = 0$  when the same  $C_{\theta_3,M}$  is retained at all windows containing  $\theta_3$ .

Each case is the output of the fixed-window matrix solver of Theorem E.35 on the corresponding row complex.

*Proof.* Case (a). The supplied finite complex has empty  $\mathcal{K}_{\theta_3, M}^{\circ} = \circ$ , so the differential matrix  $(d_{\text{ns}, M} |_{\mathcal{K}_{\theta_3, M}^{\circ}})$  is the empty  $1 \times 0$  matrix and the residual is  $r^M = E_{\theta_3, M}$ . The scalar gate  $S_{\theta_3, M} = \circ$  holds by Proposition E.36, so the row lives in the kernel complex. Since  $d_{\theta_3, M} = \circ$  on the one-element basis, the row is closed. Its cohomology class equals the basis vector itself, and the unique covector  $\ell_{\theta_3}(E_{\theta_3, M}) = 1$  annihilates the (empty) image of  $d_{\theta_3, M}$  but is nonzero on  $E_{\theta_3, M}$ . This is the cokernel separation of Theorem E.35.

Case (b). Adjoining the degree-zero  $C_{\theta_3, M}$  with  $\widehat{\sigma}_{\text{sc}, M}(C_{\theta_3, M}) = \circ$  keeps both rows in the non-scalar kernel. The differential matrix is now the scalar  $\widetilde{A}^M = (-1)$ , so  $\widetilde{A}^M c = -1 \cdot c$  and  $\widetilde{A}^M c = -r^M = -1$  has unique solution  $c = 1$ . The  $H^1$  class is therefore zero. Tower compatibility: with identity transport  $\pi_{M, N} C_{\theta_3, M} = C_{\theta_3, N}$  on every window containing  $\theta_3$ , the Roos cocycle  $P^{M, N} c_M - c_N = 1 - 1 = \circ$ , and the Bianchi tower obstruction  $\Delta_{M, N}^1 = -\pi_{M, N} \mathbf{b}^M + \mathbf{b}^N = \circ$  by the same identity transport on  $E_{\theta_3}$ .

Both cases are stable under finite-window truncation since the scalar gate  $S_{\theta_3, N} = \circ$  is preserved (the cyclic supertrace vanishes at every window), and the residual transports as  $\pi_{M, N} E_{\theta_3, M} = E_{\theta_3, N}$  when both windows contain  $\theta_3$  and as  $\circ$  otherwise.  $\square$

*Remark E.39 (Computer verification at the  $\theta_3$  row).* The script `scripts/finite_window_graph_array.py` reproduces the case dichotomy of Theorem E.38. Case (a) is returned on the omitted-CE-ancestor and omitted-counterterm rows of the row family `theta_3_current_finite_row_subcomplex`: each yields `primitive_exists = false` with cokernel covector  $\ell(E_{\theta_3, M}) = 1$ . Case (b) is returned on the three repair rows of the same family — the supplied scalar-zero counterterm `theta3_counterterm_supplied_exact_payload`, the supplied CE ancestor `theta3_ce_ancestor_supplied_exact_payload` and the supplied complete companion-face table `theta3_complete_companion_faces_supplied_exact_payload` — each returning `primitive_exists = true` with `primitive_coefficients` equal to  $[[C_{\theta_3, M}^{\text{ct}}, 1]]$ ,  $[[C_{\theta_3, M}^{\text{CE}}, 1]]$ , or the corresponding unit-coefficient signed companion sum.

*Remark E.40 (Sub-theorem dichotomy at  $\theta_3$ ).* The two cases of Theorem E.38 apply under disjoint fixture conditions on the order-three finite-row complex at  $\theta_3$ . Case (a) is the sub-theorem of the minimal seed list: when the order-three scalar-zero counterterm  $C_{\theta_3, M}$  is omitted, the finite-window  $d_{\text{ns}, M}$ -cohomology in degree one is  $\mathbb{C} E_{\theta_3, M}$ , with explicit covector representative  $\ell_{\theta_3}$ . Case (b) is the sub-theorem of the counterterm-extended complex: when  $C_{\theta_3, M}$  with  $\widehat{\sigma}_{\text{sc}, M}(C_{\theta_3, M}) = \circ$  and  $d_{\text{ns}, M} C_{\theta_3, M} = -E_{\theta_3, M}$  is supplied, the finite-window  $H^1$  vanishes. A named coefficient  $C_{\theta_3}$  is admissible exactly in case (b), with one of the three repair data — CE ancestor with controlled boundary, scalar-zero local counterterm  $C_{\theta_3, M}$ , or complete companion-face table whose signed sum moves the residual into the image of  $\widetilde{A}^M$  (Proposition E.36).

*Definition E.41 (Normal  $\Omega$ -equivariant Costello kernel datum).* Let

$$X = \mathbb{R}_t \times \mathbb{R}_s \times \mathbb{C}_{z_1} \times \mathbb{C}_{z_2}, \quad L = \mathbb{R}_t \times \{s = z_1 = z_2 = \circ\},$$

and let the normal torus be

$$T_{\Omega} = \mathbb{C}_{\epsilon_s}^* \times \mathbb{C}_{\epsilon_1}^* \times \mathbb{C}_{\epsilon_2}^*.$$

Its infinitesimal vector field is

$$V_{\Omega} = \epsilon_s s \partial_s + \epsilon_1 z_1 \partial_{z_1} + \epsilon_2 z_2 \partial_{z_2}, \quad V_{\Omega}(t) = \circ,$$

and the Cartan differential is

$$Q_\Omega = Q + \iota_{V_\Omega}, \quad Q_\Omega^2 = L_{V_\Omega}.$$

All complexes below are taken in the  $T_\Omega$ -invariant, or equivalently basic Cartan, subcomplex. Before this restriction the displayed operator is a curved Cartan operator with square  $L_{V_\Omega}$ .

Put

$$X_{N,M} \subset \{ n\epsilon_s + a\epsilon_1 + b\epsilon_2, n\epsilon_s - a\epsilon_1 - b\epsilon_2 \}$$

for the finite set of nonzero moving normal characters occurring in the chosen  $N$ -brane,  $M$ -window system. The coefficient ring is

$$R_\Omega^{N,M} = \mathbb{C}[\epsilon_s, \epsilon_1, \epsilon_2, \hbar_\omega][\chi^{-1} \mid \chi \in X_{N,M}], \quad \text{wt}(\hbar_\omega) = \epsilon_1 + \epsilon_2.$$

The self-dual branch is formed by base change to  $\epsilon_1 + \epsilon_2 = 0$  before localization; only the surviving nonzero characters in the image of  $X_{N,M}$  are inverted. In particular  $\epsilon_1 + \epsilon_2$  is never inverted and then specialized to zero. On that branch  $\hbar_\omega$  has weight zero and may be retained as a scalar parameter or set to 1 after the specialization is explicit.

The three quantum parameters are distinct:

$$\hbar_{\text{QME}}, \quad \hbar_{\text{W}}, \quad \hbar_\omega.$$

The first counts Costello/BV graph order; the second deforms the boundary Weyl/Moyal algebra; the third scalarizes the equivariant Hamiltonian bracket. A Weyl branch may identify  $\hbar_{\text{W}} = \hbar_\omega$ . Identification of  $\hbar_{\text{QME}}$  with either parameter is an explicit theorem specialization. The equivariant Hamiltonian bracket is the  $R_\Omega^{N,M}$ -linear bracket

$$\{f, g\}_\Omega = \hbar_\omega (\partial_{z_1} f \partial_{z_2} g - \partial_{z_2} f \partial_{z_1} g).$$

A normal  $\Omega$ -equivariant finite-window Costello kernel datum over  $(I, w, M, \mathcal{G}_M)$  is a finite-window marked Costello system as in Definitions 8.21 and E.27, together with the following additional data.

- ( $\Omega 1$ ) A  $T_\Omega$ -locally finite weighted field complex  $\mathcal{E}_{w,\Omega}^M|_I$ , source CE complex, local interaction  $I_{o,M,\Omega}^w$ , and invariant local-functional target

$$\mathfrak{D}_{\mathcal{G},\Omega,M}^\bullet(I) \subset \mathcal{O}_{\text{loc}}^{\text{bd}}(\mathcal{E}_{w,\Omega}^M|_I; A_{\partial,\Omega,\hbar_{\text{W}}}^{\text{pp},w,M}(I) \widehat{\otimes} R_\Omega^{N,M}[[\hbar_{\text{QME}}]])$$

with differential

$$d_{M,\Omega} = Q_\Omega + \{I_{o,M,\Omega}^w, -\}_{\hbar_{\text{QME}},M,\Omega}$$

on the invariant subcomplex. The source CE differential  $d_{\text{CE},\Omega}$  is formed from  $Q_\Omega$  and the bracket (E.41).

- ( $\Omega 2$ ) For every localized normal character  $\chi \in X_{N,M}$ , or for every surviving nonzero image of such a character on the self-dual branch, a normal contraction  $h_\chi$  satisfying

$$Q_\Omega h_\chi + h_\chi Q_\Omega = \text{id} - \text{pr}_{\chi=0}$$

on the corresponding  $\chi$ -weight summand, after inverting  $\chi$ . This is the normal-mode acyclicity datum. Graph curvature classes and QME counterterms are separate data.

(Ω<sub>3</sub>) An equivariant finite-scale gauge fixing and propagator

$$P_{\epsilon,L,\Omega}^{w,M} = \int_{\epsilon}^L (Q_{\Omega}^{\text{GF}} \otimes \mathbf{1}) K_{t,\Omega}^{w,M} dt$$

such that

$$[Q_{\Omega}, P_{\epsilon,L,\Omega}^{w,M}] = K_{\epsilon,\Omega}^{w,M} - K_{L,\Omega}^{w,M}, \quad [L_{V_{\Omega}}, P_{\epsilon,L,\Omega}^{w,M}] = 0,$$

and such that the coefficient Casimir, wavefront conditions, and finite-window weighted bounds required by the nonequivariant system remain valid over  $R_{\Omega}^{N,M}[[\hbar_{\text{QME}}]]$ .

(Ω<sub>4</sub>) A named normal-localization normalization  $\nu_{\Omega}$ . The residue branch has

$$\nu_{\Omega}^{\text{res}} = \mathbf{1}.$$

The Euler branch has

$$\nu_{\Omega}^{\text{Eu}} = \sigma_s(\epsilon_s \epsilon_1 \epsilon_2)^{-1},$$

after the corresponding Euler denominators have been admitted in the chosen localized ring. Scalar-contact rows, graph weights, and trace normalizations are always read with the selected branch. A comparison between the residue branch, Euler branch, and Euler-rescaled residue branch must state the real orientation sign  $\sigma_s$  and every denominator.

(Ω<sub>5</sub>) Equivariant local counterterm representatives

$$C_{\Gamma,w,\Omega}^M(\epsilon; B_{\bullet}^{\Gamma}) \in \mathfrak{Q}_{\mathcal{G},\Omega,M}^{\circ}(I)$$

which are  $T_{\Omega}$ -invariant, local on the relevant bulk diagonal or brane-collision stratum, compatible with finite-window truncation and admissible weight transport, and whose scalar-contact associated graded is computed using the same  $\nu_{\Omega}$ .

(Ω<sub>6</sub>) An associated-graded scalar-symbol extractor candidate

$$\sigma_{\text{sc},\Omega,M}: \text{gr}_F^{\bullet} \mathfrak{Q}_{\mathcal{G},\Omega,M}^{\bullet}(I) \longrightarrow S_{\Omega,M}^{\bullet}, \quad S_{\Omega,M}^{\bullet} = C_{\text{Lie}}^{\bullet}(\bar{\mathcal{A}}; \mathbb{C}\gamma_{\mathbf{I}})[\mathbf{I}][[\hbar_{\text{QME}}]] \widehat{\otimes} R_{\Omega}^{N,M},$$

which sends the two-Hamiltonian scalar contact to

$$\nu_{\Omega} \hbar_{\text{QME}} \text{Str}_{\text{cyc}}(\mathbf{m}) [\{f, g\}_{\Omega}]_{\circ} \gamma_{\mathbf{I}}.$$

A filtered chain lift  $\widehat{\sigma}_{\text{sc},\Omega,M}$  of  $\sigma_{\text{sc},\Omega,M}$  exists exactly when the Hom-obstruction test of Theorem E.42 vanishes.

(Ω<sub>7</sub>) A degree-zero Hom-valued graph kernel

$$\kappa_{\mathcal{G},\Omega,w,I}^M \in \text{Hom}_{\text{cont}}\left(C_{\text{CE},\Omega,\text{lab}}^{\bullet}(\mathcal{G}_M), \mathfrak{Q}_{\mathcal{G},\Omega,M}^{\bullet}(I)\right)$$

whose zero-edge part is the coefficient-current kernel and whose positive graph components are the renormalized  $\Omega$ -equivariant graph weights.

This datum supplies the equivariant structure: coefficient ring,  $Q_{\Omega}$ , normal contractions, and equivariant propagator data. Equivariant QME and centrality obstructions are formed inside this structure by the criterion of Theorem E.42.

THEOREM E.42 (*Equivariant brane-defect QME obstruction criterion*). Fix a candidate normal  $\Omega$ -equivariant finite-window system satisfying items  $(\Omega 1)$ – $(\Omega 6)$  of Definition E.41. Let

$$\sigma_{\text{sc},\Omega,M} : \text{gr}_F \mathfrak{Q}_{\mathcal{G},\Omega,M}^\bullet(I) \longrightarrow S_{\Omega,M}^\bullet$$

be its associated-graded scalar extractor. The scalar projection is a chain projection exactly when the following relative Hom obstructions vanish. First,

$$\delta_{\Omega,\sigma,M}^{(\circ)} = d_{\text{CE},\Omega} \sigma_{\text{sc},\Omega,M} - \sigma_{\text{sc},\Omega,M} d_{\text{gr},\Omega} = 0.$$

If this holds and  $s_{\Omega,M}$  is any filtered lift of  $\sigma_{\text{sc},\Omega,M}$ , the first nonzero homogeneous piece of

$$d_{\text{CE},\Omega} s_{\Omega,M} - s_{\Omega,M} d_{M,\Omega}$$

lowering the scalar-contact filtration by  $r > 0$  gives

$$\mathfrak{p}_{\Omega,\sigma,M}^{(r)} \in H^1\left(\text{Hom}_{T_{\Omega,-r}}(\text{gr}_F \mathfrak{Q}_{\mathcal{G},\Omega,M}^\bullet(I), S_{\Omega,M}^\bullet)\right).$$

A filtered chain projection  $\widehat{\sigma}_{\text{sc},\Omega,M}$  exists if and only if  $\delta_{\Omega,\sigma,M}^{(\circ)} = 0$  and the classes  $\mathfrak{p}_{\Omega,\sigma,M}^{(r)}$  vanish successively, with compatible finite-window and weight-transport choices.

Once this projection is supplied, put

$$\mathcal{K}_{\text{ns},\Omega,M}^\bullet(I) = \ker \widehat{\sigma}_{\text{sc},\Omega,M}, \quad d_{\text{ns},\Omega,M} = d_{M,\Omega}|_{\mathcal{K}_{\text{ns},\Omega,M}}.$$

Suppose the Hom-valued kernel of item  $(\Omega 7)$  is supplied, and counterterms have been chosen through order  $\hbar_{\text{QME}}^{n-1}$ . The order- $n$  equivariant residual is

$$\begin{aligned} R_{n,\Omega,M}^{\text{ns}} = [\hbar_{\text{QME}}^n] & \left( \text{Curv}_{\mathbf{K},\Omega,M}(\kappa_{\mathcal{G},\Omega,w,I}^M) + d_{M,\Omega} C_{<n,\Omega,M} \right. \\ & \left. + \frac{1}{2} \{C_{<n,\Omega,M}, C_{<n,\Omega,M}\}_{\hbar_{\text{QME}},M,\Omega} \right). \end{aligned}$$

The scalar condition at this order is

$$\mathfrak{s}_{n,\Omega,M} = \widehat{\sigma}_{\text{sc},\Omega,M}(R_{n,\Omega,M}^{\text{ns}}).$$

Only when  $\mathfrak{s}_{n,\Omega,M} = 0$  and all lower orders have been solved does the residual define the non-scalar obstruction class

$$\mathfrak{p}_{n,\Omega,M}^{\text{ns}} = [R_{n,\Omega,M}^{\text{ns}}] \in H^1(\mathcal{K}_{\text{ns},\Omega,M}^\bullet(I), d_{\text{ns},\Omega,M}).$$

A compatible order- $n$  counterterm exists precisely when

$$(\mathfrak{p}_{n,\Omega,M}^{\text{ns}})_M = 0 \quad \text{in} \quad \varprojlim_M H^1(\mathcal{K}_{\text{ns},\Omega,M}^\bullet(I))$$

and the Milnor primitive-compatibility class

$$\lambda_{n,\Omega} \in \varprojlim_M^1 H^0(\mathcal{K}_{\text{ns},\Omega,M}^\bullet(I))$$

vanishes. Iterating this criterion for all  $n$  is equivalent to

$$\text{Curv}_{\mathbf{K},\Omega}(\kappa_{\mathcal{G},\Omega,w,I} + C_{\Omega,w}) = 0$$

in the completed invariant kernel complex.

Therefore the normal  $\Omega$ -background supplies only the equivariant structure. A QME proof requires, in addition, the chain scalar projection, the non-scalar row classes displayed above, explicit local counterterms killing those classes, compatible primitive transport in the  $\varprojlim^1 H^0$  sense, and the  $Q_\Omega$ -centrality homotopies of Proposition E.50 for the source rows whose centrality is used.

*Proof.* The first paragraph is the equivariant version of Lemma E.4. On the  $T_\Omega$ -invariant subcomplex  $Q_\Omega^2 = L_{V_\Omega} = 0$ , so the Hom differential is  $D_\Omega(\alpha) = d_{\text{CE},\Omega}\alpha - (-1)^{|\alpha|}\alpha d_{\text{gr},\Omega}$ . The associated-graded defect is the obstruction to even starting a filtered lift. After it vanishes, the first filtration-lowering chain defect is closed for  $D_\Omega$ ; changing the lift by a filtration-lowering cochain changes it by a Hom coboundary. Completeness gives the successive criterion and the compatibility condition in the inverse finite-window tower.

With a chain scalar projection fixed, its kernel is preserved by  $d_{\mathcal{M},\Omega}$ . The displayed residual is the order- $n$  term of the curved Maurer–Cartan equation for the Hom-valued kernel plus lower counterterms. Applying the scalar projection gives the scalar condition. If the condition vanishes and the lower-order equations hold, the equivariant Bianchi identity in the invariant complex makes  $R_{n,\Omega,M}^{\text{ns}}$  a  $d_{\text{ns},\Omega,M}$ -cocycle. Adding an order- $n$  local counterterm changes this cocycle by a  $d_{\text{ns},\Omega,M}$ -boundary, so the fixed-window obstruction is its  $H^1$ -class. Compatibility of primitives across the finite-window inverse system is exactly the usual  $\varprojlim^1 H^0$  obstruction of Theorem 5.83. The final sentence follows because the normal contraction datum appears only in the construction of the equivariant propagator and localized coefficient complex. The curvature residual boundary, scalar-projection chain data, and source-row centrality homotopies are additional QME data.  $\square$

**THEOREM E.43** (*Minimal full-equivariant kernel vanishing*). Work in the local  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}_{\text{o,hol}}^2$  full-equivariant marked branch with  $V = \mathbb{C}^{N|N}$ . Let the finite marked list be the minimal codimension-two closed list generated by the strict current-interface rows and by the one-cross two-Hamiltonian scalar-contact row of Proposition 8.25. Assume that the list consists exactly of these rows through codimension two: it contains the  $|\Gamma|_h = 1$  scalar-contact row and its codimension-two closure, and it has zero additional  $|\Gamma|_h = 1$  graph rows, zero genuine  $|\Gamma|_h = 2$  seed graphs, zero order-2 CE-source faces, and zero order-2 graph counterterm faces.

Then the first kernel residuals are

$$R_{1,M}^{\text{ns}} = 0, \quad R_{2,M}^{\text{ns}} = 0,$$

and the compatible counterterms are

$$C_{1,M} = 0, \quad C_{2,M} = 0.$$

Thus the minimal full-equivariant finite-window array has zero genuinely non-scalar QME obstruction through the first codimension-two closure level. Its first possible non-scalar obstruction can only appear after a larger finite marked list supplies an additional scalar-zero row.

*Proof.* The order-0 current-interface rows have zero curvature by Theorem E.10 and by the finite-window compatibility recorded in Proposition E.30. The only order-1 row in the minimal list is the one-cross



scalar-contact row. Its scalar coefficient is the cyclic supertrace of the transported full-equivariant marker tensor. By Lemma E.31, this coefficient is zero for  $V = \mathbb{C}^{N|N}$ , including the empty word. Hence the row is zero as a local functional in the full-equivariant branch, so  $R_{1,M}^{\text{ns}} = 0$  and  $C_{1,M} = 0$ .

Codimension-two closure adds only the two ordered composites of two first scalar-contact boundary faces. Their incidence signs are opposite and their output graphs, coefficients, and marker transports are identified by Equation (E.27). This gives the pairwise cancellation of the marked differential. In the full-equivariant branch each scalar-loop factor is already zero by the same cyclic supertrace calculation. The minimal list has zero genuine order-2 seed graph, CE-source face, or counterterm face, so the kernel complex has zero remaining row. Thus  $R_{2,M}^{\text{ns}} = 0$ , and the zero counterterm is compatible under every finite-window truncation.  $\square$

**PROPOSITION E.44** (*First missing non-scalar row*). Let  $\mathcal{G}_M^{\text{mk}}$  be any larger finite marked list in the same full-equivariant local branch, with scalar projection killed as in Theorem E.32. The first genuinely non-scalar obstruction beyond scalar projection is the least order  $n$  for which the row sum of Definition E.33 is nonzero in cohomology:

$$\mathfrak{o}_{n,M}^{\text{ns}} = \left[ \sum_{\alpha \in A_{n,M}^{\text{fw}}} R_{\alpha,M}^{\text{row}} \right] \in H^1(\mathcal{K}_{\text{ns},M}^\bullet(I), d_{\text{ns},M}).$$

Since the minimal order-1 scalar-contact row vanishes, the first datum that can make  $n = 1$  is an additional order-1 scalar-zero row

$$\alpha \in A_{1,M}^{\text{fw}} \setminus \{\alpha_{\text{sc}}\}, \quad S_{\alpha,M} = 0,$$

with its row value  $R_{\alpha,M}^{\text{row}}$ . Equivalently, the missing row entry is one of the order-1 summands

$$\varepsilon_\Gamma d_M K_\Gamma^M, \quad -\varepsilon_{\Gamma'}^{\text{CE}} K_{\Gamma'}^M, \quad \frac{1}{2} \varepsilon_{\Gamma_0, \Gamma_1} \{K_{\Gamma_0}^M, K_{\Gamma_1}^M\}_{h,M},$$

together with its graph type, finite-window labels, full-equivariant marker tensor, incidence sign, scalar condition, and truncation coefficients  $t_{\alpha\alpha'}^{M,N}$ .

If the finite marked list explicitly has zero such order-1 rows, the next required datum is the order-2 scalar-zero row sum

$$\begin{aligned} R_{2,M}^{\text{ns}} &= \sum_{|\Gamma|_h=2} \varepsilon_\Gamma d_M K_\Gamma^M - \sum_{|\Gamma'|_h=2} \varepsilon_{\Gamma'}^{\text{CE}} K_{\Gamma'}^M \\ &\quad + \frac{1}{2} \sum_{|\Gamma_1|_h=|\Gamma_2|_h=1} \varepsilon_{\Gamma_1, \Gamma_2} \{K_{\Gamma_1}^M, K_{\Gamma_2}^M\}_{h,M} + [\hbar^2] d_M C_{<2,M} \\ &\quad + \frac{1}{2} [\hbar^2] \{C_{<2,M}, C_{<2,M}\}_{h,M}. \end{aligned}$$

These row values, signs, scalar conditions, and truncation coefficients define the non-scalar obstruction class and counterterm problem beyond the minimal vanishing theorem above.

*Proof.* The scalar projection is a chain map on the full-equivariant marked complex by Theorem E.32, so its kernel is a  $d_M$ -stable complex. After the scalar condition vanishes, Theorem 8.22 puts the first unsolved residual in degree one of that kernel complex. Adding a degree-zero counterterm changes the residual by a  $d_M$ -boundary, and hence the displayed cohomology class is exactly the obstruction.

The minimal order-1 row is already zero by Theorem E.43. Thus an order-1 non-scalar class can only be supplied by an additional finite-window row outside  $\alpha_{\text{sc}}$ . The three displayed forms are precisely the

order-1 possibilities in the curvature expansion: target differential of an order-1 graph component, CE source face of such a component, or a convolution bracket whose orders add to 1. Each is a row only after the finite graph list supplies its labels, signs, marker transport, scalar condition, and truncation coefficients.

With zero order-1 rows, the next possible residual is the order-2 curvature expansion, including genuine order-2 graph components, CE-source faces, brackets of order-1 components, and lower-counterterm terms. These data are exactly the row entries required by Definition 8.24.  $\square$

**PROPOSITION E.45** (*First scalar-zero order-one row outside the minimal array*). Work in the full-equivariant local branch of Theorem E.43. Let  $M \geq 1$ , and enlarge the minimal finite marked list by the labelled one-cross scalar-contact boundary operation  $\alpha_{\text{II}}$  with ordered Hamiltonian inputs  $(a, z_1)$ ,  $(b, z_1)$ , the boundary Weyl edge  $E_{\text{Weyl}}$ , the scalar-contact output vertex  $\gamma_1$ , and the same full-equivariant marker complex as before. Choose the orientation in which the displayed ordered pair has incidence coefficient +1.

Then  $\alpha_{\text{II}}$  is a scalar-zero order-one row outside  $\alpha_{\text{sc}}$ , with row data

$$|\alpha_{\text{II}}|_{\hbar} = 1, \quad d_{\max}(\alpha_{\text{II}}) = 1, \quad p_{\alpha_{\text{II}}} = 1, \quad r_{\alpha_{\text{II}}} = 1, \quad \iota_{\alpha_{\text{II}}, M} = +1.$$

Its row value and scalar condition are

$$R_{\alpha_{\text{II}}, M}^{\text{row}} = \text{Str}_{\text{cyc}}(\mu_{\beta_{\text{II}}}(\mathbf{m}))\omega(z_1, z_1) \int_I a(t)b(t)\gamma_1(t) dt = 0,$$

$$S_{\alpha_{\text{II}}, M} = \text{Str}_{\text{cyc}}(\mu_{\beta_{\text{II}}}(\mathbf{m}))\omega(z_1, z_1)\gamma_1 = 0.$$

Here the coefficient is the extracted  $\hbar^1$ -coefficient; the full scalar-contact representative has the additional factor  $\hbar$ . For  $M \geq N \geq 1$ , retain the same labelled row in window  $N$  and set

$$t_{\alpha_{\text{II}}\alpha_{\text{II}}}^{M, N} = 1, \quad t_{\alpha_{\text{II}}\alpha'}^{M, N} = 0 \quad (\alpha' \neq \alpha_{\text{II}}).$$

These coefficients satisfy the truncation equations because  $\pi_{M, N} R_{\alpha_{\text{II}}, M}^{\text{row}} = 0$  and  $\pi_{M, N} S_{\alpha_{\text{II}}, M} = 0$ .

The associated kernel class is therefore

$$\mathfrak{o}_{1, M}^{\text{ns}}(\alpha_{\text{II}}) = [R_{\alpha_{\text{II}}, M}^{\text{row}}] = 0 \in H^1(\mathcal{K}_{\text{ns}, M}^{\bullet}(I), d_{\text{ns}, M}),$$

and the required counterterm contribution is  $C_{1, M} = 0$ .

Thus the finite closed graph-list axioms support explicit scalar-zero order-one rows beyond  $\alpha_{\text{sc}}$ . The first explicit such row is harmless: it is zero before cohomology. If the row convention retains only nonzero summands, then  $\alpha_{\text{II}}$  is omitted, and a nonzero order-one class still requires an additional supplied row  $R_{\alpha, M}^{\text{row}}$  of the form stated in Proposition E.44.

*Proof.* The scalar-contact boundary formula in Proposition 8.25 applies to every ordered pair of Hamiltonian labels in the finite window. For the pair  $(z_1, z_1)$ ,

$$\omega(z_1, z_1) = [\{z_1, z_1\}_{\Omega_{\text{loc}}}]_0 = 0.$$

Multiplying by the transported full-equivariant cyclic supertrace preserves this zero coefficient. Hence both the local functional row and its scalar projection vanish.

Since the row value is zero,  $d_{\text{ns}, M}$ -closedness is automatic and the cohomology class is the zero class. The displayed truncation coefficients send the labelled row to the same labelled row in lower windows

containing  $z_1$ , and both sides of the row and scalar truncation identities are zero. The zero counterterm solves the row equation. The final assertion is logical: the finite-list axioms allow this added labelled zero row; a nonzero order-one class is decided after the nonzero order-one row values have been supplied.  $\square$

*Remark E.46 (Order two after zero nonzero order-one rows).* If zero rows such as  $\alpha_{\text{ti}}$  are suppressed and the finite marked list supplies zero nonzero order-one rows outside  $\alpha_{\text{sc}}$ , then  $C_{1,M} = 0$  and every order-two bracket row containing  $\alpha_{\text{sc}}$  is zero by Proposition 8.26. The order-two scalar-zero residual reduces to

$$\begin{aligned} R_{2,M}^{\text{ns}} &= \sum_{|\Gamma|_h=2} \varepsilon_\Gamma d_M K_\Gamma^M - \sum_{|\Gamma'|_h=2} \varepsilon_{\Gamma'}^{\text{CE}} K_{\Gamma'}^M \\ &\quad + \frac{1}{2} \sum_{\substack{|\Gamma_1|_h=|\Gamma_2|_h=1 \\ \Gamma_i \neq \alpha_{\text{sc}}}} \varepsilon_{\Gamma_1, \Gamma_2} \{K_{\Gamma_1}^M, K_{\Gamma_2}^M\}_{h,M}. \end{aligned}$$

Under the stronger minimal hypotheses of Theorem E.43 the three sums are empty, and  $R_{2,M}^{\text{ns}} = 0$ . In any larger finite system this displayed row sum is a cochain only after the genuine order-two graph weights, CE-source faces, bracket signs, scalar conditions, and truncation coefficients have been supplied.

**THEOREM E.47 (Minimal full-equivariant all-order vanishing).** Work in the local  $\mathbb{R}_{\text{top}}^2 \times \widehat{\mathbb{C}}_{\text{o,hol}}^2$  full-equivariant marked branch with  $V = \mathbb{C}^{N|N}$ . Let  $\mathcal{G}_M^{\text{mk,o}}$  be the finite marked list generated by the strict current-interface rows, the two-Hamiltonian scalar-contact boundary row  $\alpha_{\text{sc}}$ , and its subset-deletion closure in the sense of Definition E.27. Assume that the finite-window system adds zero further positive-order seed graphs, CE-source faces, graph counterterm faces, or nonzero scalar-zero rows.

Then for every  $n \geq 1$

$$R_{n,M}^{\text{ns}} = 0, \quad \mathfrak{o}_{n,M}^{\text{ns}} = 0, \quad C_{n,M} = 0$$

If the row convention suppresses zero rows, then  $\mathcal{A}_{n,M}^{\text{fw,o}}$  is empty for every  $n \geq 1$ . If it retains labelled zero rows, every positive-order retained row has

$$R_{\alpha,M}^{\text{row}} = 0, \quad S_{\alpha,M} = 0,$$

and may be given the zero truncation row  $t_{\alpha\alpha'}^{M,N} = 0$ . In particular, the minimal full-equivariant order-three residual is zero.

*Proof.* Proposition 8.25 proves that the one-cross scalar-contact component satisfies  $K_{\alpha_{\text{sc}}}^M = 0$  as a local functional before scalar projection. The proof is the cyclic-supertrace calculation of Lemma E.31: every transported full-equivariant marker on a scalar open loop has cyclic supertrace zero, including the empty marker word.

Every positive-order boundary row in  $\mathcal{G}_M^{\text{mk,o}}$  contains at least one such scalar-contact factor. Hence its local-functional value is zero. Bracket rows in the minimal system have at least one positive-order factor of this form, so bilinearity of the BV/convolution bracket gives zero. Lower counterterm rows vanish inductively because the chosen lower counterterms are zero. The minimality hypothesis sets the genuine  $dK$ , CE-source, and graph-counterterm row spaces to zero. Therefore the order- $n$  scalar-zero row sum is zero for all  $n \geq 1$ , its class in  $H^1(\mathcal{K}_{\text{ns},M}^\bullet(I), d_{\text{ns},M})$  is zero, and the zero counterterm is a compatible primitive. Truncation sends zero rows to zero rows, giving the displayed truncation choice.  $\square$

**PROPOSITION E.48 (First order-three enlarged-system datum).** Keep the full-equivariant local branch, and suppose the minimal system has been closed through order two with  $C_{1,M} = C_{2,M} = 0$  and zero nonzero order-one or order-two rows outside the scalar-contact closure. Then a nonzero order-three

obstruction requires row data beyond the finite graph-list axioms. A nonzero order-three cochain exists only after the finite-window system supplies at least one of the following row data:

$$\begin{aligned} R_{3,M}^{\text{ns}} &= \sum_{|\Gamma|_h=3} \varepsilon_\Gamma d_M K_\Gamma^M - \sum_{|\Gamma'|_h=3} \varepsilon_{\Gamma'}^{\text{CE}} K_{\Gamma'}^M \\ &\quad + \frac{1}{2} \sum_{\substack{|\Gamma_1|_h+|\Gamma_2|_h=3 \\ K_{\Gamma_1}^M \neq 0, K_{\Gamma_2}^M \neq 0}} \varepsilon_{\Gamma_1, \Gamma_2} \{K_{\Gamma_1}^M, K_{\Gamma_2}^M\}_{h,M} \\ &\quad + [\hbar^3] \left( d_M C_{<3,M} + \frac{1}{2} \{C_{<3,M}, C_{<3,M}\}_{h,M} \right). \end{aligned}$$

Under the preceding minimal hypotheses the bracket and lower-counterterm sums are empty. Thus the first missing order-three datum is a genuine  $|\Gamma|_h = 3$  graph-differential row

$$R_{d\Gamma, M}^{\text{row}} = \varepsilon_\Gamma d_M K_\Gamma^M, \quad S_{d\Gamma, M} = \varepsilon_\Gamma \widehat{\sigma}_{\text{sc}, M}(d_M K_\Gamma^M),$$

or an order-three CE-source row

$$R_{\text{CE}(\Gamma, \nu), M}^{\text{row}} = -\varepsilon_{\Gamma, \nu}^{\text{CE}} K_\Gamma^M(\zeta_{M, \nu}), \quad S_{\text{CE}(\Gamma, \nu), M} = -\varepsilon_{\Gamma, \nu}^{\text{CE}} \widehat{\sigma}_{\text{sc}, M}(K_\Gamma^M(\zeta_{M, \nu})).$$

In a nonminimal system with supplied nonzero order-one and order-two components one must also supply each bracket row

$$R_{b(\Gamma_1, \Gamma_2), M}^{\text{row}} = \frac{1}{2} \varepsilon_{\Gamma_1, \Gamma_2} \{K_{\Gamma_1}^M, K_{\Gamma_2}^M\}_{h,M}, \quad S_{b, M} = \widehat{\sigma}_{\text{sc}, M}(R_{b(\Gamma_1, \Gamma_2), M}^{\text{row}}),$$

for  $|\Gamma_1|_h + |\Gamma_2|_h = 3$ , together with its truncation coefficients  $t_{bb'}^{M, N}$ .

The scalar condition for order three is the single condition

$$\mathfrak{s}_{3, M} = \sum_{\alpha \in A_{3, M}^{\text{fw}}} S_{\alpha, M} = 0.$$

Only after this condition is supplied and vanishes is the non-scalar obstruction class defined:

$$\mathfrak{o}_{3, M}^{\text{ns}} = [R_{3, M}^{\text{ns}}] \in H^1(\mathcal{K}_{\text{ns}, M}^\bullet(I), d_{\text{ns}, M}).$$

A fixed-window counterterm  $C_{3, M}$  exists exactly when this class vanishes; a compatible tower also requires the corresponding  $\varprojlim^1 H^0$ -compatibility class to vanish.

*Proof.* The order-three expansion is Theorem 8.36 with  $n = 3$ , written in the row notation of Definition 8.24. Rows containing  $\alpha_{\text{sc}}$  vanish by Proposition 8.26, and rows containing any retained labelled zero scalar-contact row vanish by the same bilinearity argument. Since  $C_{1, M} = C_{2, M} = 0$ , the lower-counterterm contribution is zero in the minimal system. Every nonzero order-three cochain in this system therefore comes from a genuine order-three graph row, an order-three CE-source row, or a bracket row between already supplied nonzero lower-order components. The displayed formulas are precisely the row values and scalar conditions required by Definition 8.24. The final assertions are the kernel obstruction criterion of Definition E.33 and the finite-window compatibility criterion of Theorem 8.36.  $\square$

**PROPOSITION E.49** (*Marked-row test for unreduced centrality homotopies*). Work in a finite-window marked Costello list for the local mixed HT brane-defect theory, with labels and source faces as in Definition E.27. Let  $\kappa_{\mathcal{G},w,I}^M$  be the Hom-valued finite-window kernel carried by this list. For source generators  $x, y \in \{u_{a(t)d\mathfrak{t} \otimes f}, c_{B,\rho}\}$ , the marked row testing unreduced centrality is

$$R_{x,y,M}^{\text{cent}} = \text{Curv}_{\mathbf{K},M}(\kappa_{\mathcal{G},w,I}^M)(x \wedge y).$$

In row notation this is the finite sum

$$\begin{aligned} R_{x,y,M}^{\text{cent}} = & \sum_{\Gamma \in \mathcal{A}_{x,y,M}^d} \varepsilon_{\Gamma} d_M K_{\Gamma}^M - \sum_{\Gamma \in \mathcal{A}_{x,y,M}^{\text{CE}}} \varepsilon_{\Gamma}^{\text{CE}} K_{\Gamma}^M(\zeta_{\Gamma,x,y}) \\ & + \frac{1}{2} \sum_{(\Gamma_1, \Gamma_2) \in \mathcal{A}_{x,y,M}^{\text{br}}} \varepsilon_{\Gamma_1, \Gamma_2} \{K_{\Gamma_1}^M, K_{\Gamma_2}^M\}_{h,M} + R_{x,y,M}^{\text{ct}}. \end{aligned}$$

Here the four summands are respectively the target differential rows, the CE-source rows, the convolution bracket rows, and the supplied counterterm rows of the marked table. The scalar condition is

$$S_{x,y,M}^{\text{cent}} = \widehat{\sigma}_{\text{sc},M}(R_{x,y,M}^{\text{cent}}).$$

Only after this condition is zero does the centrality row define a class

$$\mathfrak{o}_{x,y,M}^{\text{cent}} = [R_{x,y,M}^{\text{cent}}] \in H^{|x|+|y|}(\mathcal{K}_{\text{ns},M}^{\bullet}(I), d_{\text{ns},M}).$$

A marked finite-window system supplies an unreduced centrality homotopy for the pair  $(x, y)$  exactly when it also supplies a degree  $|x| + |y| - 1$  row  $H_{x,y}^M \in \mathcal{K}_{\text{ns},M}^{|x|+|y|-1}(I)$  with

$$R_{x,y,M}^{\text{cent}} + d_M H_{x,y}^M = \mathfrak{o},$$

and these rows satisfy the truncation and weight identities of Theorem B.22. The centrality-homotopy row and its compatibility identities are additional data beyond scalar-shadow vanishing.

In the minimal full-equivariant branch of Theorem E.47, every positive-order row contains the zero scalar-contact factor  $K_{\alpha_{\text{sc}}}^M = \mathfrak{o}$ , and the marked table contains zero additional positive-order seed graph, CE-source face, graph counterterm face, or nonzero scalar-zero row. Hence

$$R_{x,y,M}^{\text{cent}} = \mathfrak{o}, \quad H_{x,y}^M = \mathfrak{o}$$

is a valid homotopy inside that minimal finite marked complex. This constructs the homotopy only for the minimal system. An enlarged unreduced Costello graph/counterterm target must supply the displayed row values, scalar condition, primitive  $H_{x,y}^M$ , and transition coefficients.

*Proof.* Proposition E.30 identifies the marked differential with the finite-window QME differential and, for Hom-valued kernels, adds the source face  $-\kappa d_{\text{CE}}$ . Therefore the binary curvature on  $x \wedge y$  is exactly the sum of the four row types displayed above: target differential, CE-source subtraction, convolution bracket, and counterterm contribution. Applying the scalar projection gives the scalar condition. If the condition vanishes, the row lies in the non-scalar kernel complex of Definition E.33; its cohomology class is the obstruction to finding a fixed-window primitive.

The equation  $R_{x,y,M}^{\text{cent}} + d_M H_{x,y}^M = \mathfrak{o}$  is precisely the homotopy equation (29). The transition identities are necessary because the local kernel is the inverse limit over finite windows and admissible

weights. Thus a row omitted from the marked table is additional graph, source-face, counterterm, or homotopy data beyond the scalar projection.

In the minimal full-equivariant system all positive-order scalar-contact rows are zero before cohomology by Theorem E.47. Since the minimal hypothesis allows zero other positive-order rows, the centrality row is zero and the zero primitive is compatible with all truncation maps. The final assertion is the contrapositive of the row formula: once an enlarged system adds a nonzero graph, source, bracket, or counterterm row, the corresponding scalar condition and primitive must be supplied explicitly.  $\square$

**PROPOSITION E.50** ( *$Q_\Omega$ -centrality homotopy criterion*). Let  $(I, w, M, \mathcal{G}_M)$  carry a normal  $\Omega$ -equivariant Costello kernel datum in the sense of Definition E.41, and assume the scalar projection has passed the Hom-obstruction test of Theorem E.42. For  $T_\Omega$ -invariant source generators

$$x, y \in \{u_{a(t)} dt \otimes f, c_{B,\rho}\},$$

define the equivariant centrality row

$$R_{x,y,\Omega,M}^{\text{cent}} = \text{Curv}_{\mathbf{K},\Omega,M}(\kappa_{\mathcal{G},\Omega,w,I}^M)(x \wedge y).$$

Its finite-row expansion is

$$\begin{aligned} R_{x,y,\Omega,M}^{\text{cent}} &= \sum_{\Gamma \in \mathbf{A}_{x,y,\Omega,M}^d} \varepsilon_\Gamma d_{M,\Omega} K_{\Gamma,\Omega}^M - \sum_{\Gamma \in \mathbf{A}_{x,y,\Omega,M}^{\text{CE}}} \varepsilon_\Gamma^{\text{CE}} K_{\Gamma,\Omega}^M (\zeta_{\Gamma,x,y,\Omega}) \\ &\quad + \frac{1}{2} \sum_{(\Gamma_1, \Gamma_2) \in \mathbf{A}_{x,y,\Omega,M}^{\text{br}}} \varepsilon_{\Gamma_1, \Gamma_2} \{K_{\Gamma_1, \Omega}^M, K_{\Gamma_2, \Omega}^M\}_{h_{\text{QME}, M, \Omega}} + R_{x,y,\Omega,M}^{\text{ct}}. \end{aligned}$$

The scalar condition is

$$S_{x,y,\Omega,M}^{\text{cent}} = \widehat{\sigma}_{\text{sc}, \Omega, M}(R_{x,y,\Omega,M}^{\text{cent}}).$$

Only after this condition vanishes does one obtain the obstruction class

$$\mathfrak{o}_{x,y,\Omega,M}^{\text{cent}} = [R_{x,y,\Omega,M}^{\text{cent}}] \in H^{|x|+|y|}(\mathcal{K}_{\text{ns}, \Omega, M}^\bullet(I), d_{\text{ns}, \Omega, M}).$$

A  $Q_\Omega$ -centrality homotopy for  $(x, y)$  is exactly a  $T_\Omega$ -invariant row

$$H_{x,y,M}^\Omega \in \mathcal{K}_{\text{ns}, \Omega, M}^{|x|+|y|-1}(I)$$

satisfying

$$R_{x,y,\Omega,M}^{\text{cent}} + d_{M,\Omega} H_{x,y,M}^\Omega = 0, \quad L_{V_\Omega} H_{x,y,M}^\Omega = 0,$$

together with the same interval, finite-window, and admissible-weight transition identities as in Theorem B.22. Equivalently, the inverse-system class

$$(\mathfrak{o}_{x,y,\Omega,M}^{\text{cent}})_M \in \varprojlim_M H^{|x|+|y|}(\mathcal{K}_{\text{ns}, \Omega, M}^\bullet(I))$$

and the corresponding  $\varprojlim^1 H^{|x|+|y|-1}$ -primitive compatibility class must both vanish.

If one works before passing to  $T_\Omega$ -invariants, then  $Q_\Omega^2 = L_{V_\Omega}$  gives a secondary obstruction

$$\ell_{x,y,\Omega,M}^{\text{cent}} = [L_{V_\Omega} H_{x,y,M}^\Omega] \in H^{|x|+|y|}(\mathcal{K}_{\text{ns}, \Omega, M}^\bullet(I), d_{\text{ns}, \Omega, M})$$

for any chosen primitive. This class must vanish, or the primitive must be replaced by a basic one, before the homotopy is a  $Q_\Omega$ -homotopy.

*Proof.* The row expansion is the Hom-valued curvature computation of Proposition E.49 with  $d_M$ , the CE differential, the bracket, and the counterterm table replaced by their equivariant counterparts from Definition E.41. The scalar condition is necessary because only its vanishing places the row in  $\mathcal{K}_{\text{ns},\Omega,M}^\bullet$ . Once there, the equivariant Bianchi identity makes the row closed, so its cohomology class is the obstruction to a fixed-window primitive.

The displayed equation is precisely the centrality homotopy equation in the invariant  $Q_\Omega$ -complex. Since  $Q_\Omega^2 = L_{V_\Omega}$ ,  $T_\Omega$ -invariance of the primitive is an independent condition beyond scalar-shadow vanishing. The inverse-limit assertion is the same primitive-compatibility argument as in Theorem B.22. If a primitive is chosen outside the basic subcomplex, applying  $d_{M,\Omega}$  a second time records  $L_{V_\Omega}H$ ; its cohomology class is exactly the secondary obstruction displayed above.  $\square$

**PROPOSITION E.51** (*Even-block marker obstruction*). Replace full  $\mathfrak{gl}(V)$ -equivariance by the smaller even block-diagonal subalgebra

$$\mathfrak{gl}(V_\circ) \oplus \mathfrak{gl}(V_\text{I}) \subset \mathfrak{gl}(V), \quad V_\circ \cong V_\text{I} \cong \mathbb{C}^N.$$

Then a single fixed even marker is admissible exactly when

$$T = \lambda_\circ P_\circ + \lambda_\text{I} P_\text{I},$$

where  $P_\circ$  and  $P_\text{I}$  are the parity block projections. Its scalar shadow is

$$\text{Str}(T) = N(\lambda_\circ - \lambda_\text{I}).$$

For a two-Hamiltonian scalar contact the obstruction class is

$$\hbar N(\lambda_\circ - \lambda_\text{I}) \omega(f, g) \gamma_\text{I}, \quad \hbar N(\lambda_\circ - \lambda_\text{I}) [\bar{c}] \gamma_\text{I} \text{ in cohomology.}$$

Since  $[\bar{c}] \neq 0$  in the scalar-reduced source, this smaller subalgebra has zero  $[\bar{c}]$ -shadow for single markers if and only if  $\lambda_\circ = \lambda_\text{I}$ , i.e. the marker is scalar and the subalgebra has collapsed back to the full-equivariant single-marker case. For higher marker tensors the strongest true positive statement is exactly the supertraceless-monodromy submodule of Theorem E.25; outside it the obstruction is the displayed  $\text{Str}_{\text{cyc}}$ -weighted  $[\bar{c}]$ -class.

*Proof.* The commutant of  $\mathfrak{gl}(V_\circ) \oplus \mathfrak{gl}(V_\text{I})$  on  $V_\circ \oplus V_\text{I}$  is  $\mathbb{C}P_\circ \oplus \mathbb{C}P_\text{I}$ , so the displayed form of  $T$  is necessary and sufficient. The supertrace is

$$\text{Str}(\lambda_\circ P_\circ + \lambda_\text{I} P_\text{I}) = \lambda_\circ \dim V_\circ - \lambda_\text{I} \dim V_\text{I} = N(\lambda_\circ - \lambda_\text{I}).$$

Proposition E.23 then gives the scalar CE representative and the cohomology class by multiplying  $\omega$ , respectively  $[\bar{c}]$ , by this coefficient. The class  $[\bar{c}]$  is nonzero by Theorem E.12; hence the class vanishes for all two-Hamiltonian tests exactly when  $\lambda_\circ = \lambda_\text{I}$ . The higher tensor assertion is the same calculation with  $\text{Str}(T)$  replaced by  $\text{Str}_{\text{cyc}}$  and with the preservation criterion supplied by Theorem E.25.  $\square$

**THEOREM E.52** (*Ordinary scalar-reduced  $\mathfrak{gl}_N$  obstruction criterion*). Work in the scalar-reduced source of Definition E.3 and keep the ordinary trace on the open  $\mathfrak{gl}_N$  brane. Let  $\mathcal{W}_{\text{ext}}$  be any additional formal-local closed-exchange or brane-defect graph contribution whose leading scalar associated graded is  $w_{\text{ext}} \in C_{\text{Lie}}^2(\bar{A}; \mathbb{C})$ . A cancellation

$$\text{Ob}_{\text{sc}} + \mathcal{W}_{\text{ext}} + d_{\text{QME}} C = 0$$

with  $C \in \mathcal{O}_{\text{loc}}^{\text{bd}}$  can hold only if

$$[w_{\text{ext}}] = -\hbar N [\bar{c}] \quad \text{in } H_{\text{Lie}}^2(\bar{A}; \mathbb{C})[[\hbar]].$$

In particular, with  $\mathcal{W}_{\text{ext}} = \mathbf{o}$ , or with  $w_{\text{ext}}$  exact, the ordinary scalar-reduced  $\mathfrak{gl}_N$  brane-defect model carries the scalar QME obstruction  $\hbar N [\bar{c}]$ .

*Proof.* Taking associated graded with respect to the scalar-contact filtration sends the equation

$$\text{Ob}_{\text{sc}} + \mathcal{W}_{\text{ext}} + d_{\text{QME}} C = \mathbf{o}$$

to

$$\hbar N \omega + w_{\text{ext}} + \delta \bar{\eta} = \mathbf{o} \quad \text{in } C_{\text{Lie}}^2(\bar{A}; \mathbb{C}),$$

where  $\bar{\eta}$  is the linear functional represented by the leading scalar part of  $C$ . Passing to cohomology gives the stated necessary condition. If  $\mathcal{W}_{\text{ext}} = \mathbf{o}$ , or if  $w_{\text{ext}}$  is exact, this condition would force  $\hbar N [\bar{c}] = \mathbf{o}$  in  $H_{\text{Lie}}^2(\bar{A}; \mathbb{C})[[\hbar]]$ . For  $N > \mathbf{o}$  and formal  $\hbar$ , Theorem E.12 identifies  $[\bar{c}] \neq \mathbf{o}$  on  $\bar{A}$ . Thus the ordinary scalar-reduced obstruction survives every internal local counterterm.  $\square$

**THEOREM E.53** (*Unreduced cotangent lift obstruction: necessary and sufficient*). Fix a finite-window marked Costello system  $(I, w, M, \mathcal{G}_M, \mathcal{G}_M^{\text{mk}})$  for the local mixed HT brane-defect theory and a filtered chain scalar projection  $\widehat{\pi}_{\text{sc}, M}$  (Definition E.34). Let  $\Phi_{\text{fact}}^{\text{unred}}: C_{\text{CE}}^\bullet(\bar{\mathfrak{g}}_I) \rightarrow Z_{\text{fact}}^{P_{\mathbf{o}}}(\text{Obs}_{\text{open}})$  be a candidate factorization-center morphism extending the algebraic current presentation  $\Phi_I^\chi$  of Construction E.9 beyond the algebraic current target. Define the obstruction vector

$$\text{Ob}_{\text{cent/QME/descent}} = (\text{ob}_{\text{cent}}^{\text{prod}}, \text{ob}_{\text{cent}}^{P_{\mathbf{o}}}, \text{ob}_{\text{QME}}^{\text{bd}}, \delta_{\dot{C}/R}^{\text{Weiss}}),$$

with components defined as follows.

- (i) *Product/tensor-completion centrality.*  $\text{ob}_{\text{cent}}^{\text{prod}}$  is the  $H^{|x|+|y|}$ -class of the product centrality defect  $\Phi_{\text{fact}}^{\text{unred}}(x)\Phi_{\text{fact}}^{\text{unred}}(y) - (-1)^{|x||y|}\Phi_{\text{fact}}^{\text{unred}}(y)\Phi_{\text{fact}}^{\text{unred}}(x)$  in  $\mathfrak{D}_{\mathcal{G}, M}^\bullet(I)/\text{im}(d_M)$ . In the algebraic current model (Theorem E.10) this defect is zero on generators by graded commutativity; the obstruction measures the corresponding product-centrality defect on the enlarged unreduced target.

- (ii)  *$P_{\mathbf{o}}$  / shifted-cotangent refinement.*  $\text{ob}_{\text{cent}}^{P_{\mathbf{o}}}$  is the  $H^{|x|+|y|}$ -class of the  $P_{\mathbf{o}}$ -bracket defect  $\{\Phi_{\text{fact}}^{\text{unred}}(x), \Phi_{\text{fact}}^{\text{unred}}(y)\} - \Phi_{\text{fact}}^{\text{unred}}(\partial_x y)$  in the same cohomology, evaluated as a finite row sum

$$R_{x,y,M}^{\text{cent}} = \sum_{\Gamma \in \mathcal{A}_{x,y,M}^d} \varepsilon_\Gamma d_M K_\Gamma^M - \sum_{\Gamma \in \mathcal{A}_{x,y,M}^{\text{CE}}} \varepsilon_\Gamma^{\text{CE}} K_\Gamma^M (\zeta_{\Gamma,x,y}) + \frac{1}{2} \sum_{(\Gamma_1, \Gamma_2)} \varepsilon_{\Gamma_1, \Gamma_2} \{K_{\Gamma_1}^M, K_{\Gamma_2}^M\}_{\hbar, M} + R_{x,y,M}^{\text{ct}}$$

of the row machinery (Proposition E.49). A centrality homotopy  $H_{x,y}^M$  with  $R_{x,y,M}^{\text{cent}} + d_M H_{x,y}^M = \mathbf{o}$  is the construction-level witness; the obstruction is the cohomology class of the row before such a witness is supplied.

- (iii) *Boundary QME obstruction.*  $\text{ob}_{\text{QME}}^{\text{bd}}$  is the compatible inverse-system class

$$(\mathfrak{o}_{n,M}^{\text{ns}})_M = ([R_{n,M}^{\text{ns}}])_M \in \varprojlim_M H^1(\mathcal{K}_{\text{ns}, M}^\bullet(I), d_{\text{ns}, M})$$

of Theorem E.35, together with the secondary  $\varprojlim^1 H^0$  primitive-compatibility class  $\lambda_{n,M}$ . The first nontrivial finite-window value occurs at the  $\bar{\theta}_3$  row (Theorem E.38).



- (iv) *Čech–Roos / Weiss-versus-Ran descent gap*.  $\delta_{\check{C}/R}^{\text{Weiss}}$  is the codimension-one face class of the descent comparison between the Weiss cover  $\mathcal{U}^{\text{Weiss}}$  and the Ran-space cover  $\mathcal{U}^R$ : the same factorization-center lift descends to  $\mathcal{U}^R$  if and only if the Čech–Roos difference  $\delta_{\check{C}/R}^{\text{Weiss}}$  vanishes in the finite-window  $\varprojlim^1 H^\circ$  of the corresponding boundary kernel.

Then:

$$\Phi_{\text{fact}}^{\text{unred}} \text{ lifts to an unreduced } Z_{\text{fact}}^{P_o}\text{-morphism} \iff \text{Ob}_{\text{cent/QME/descent}} = 0.$$

At the associated-graded level, the primitive indecomposable  $P_o$ -shadow of Theorem 5.132 computes the first page  $E_1$  of the spectral sequence converging to  $\text{ob}_{\text{cent}}^{P_o}$ : the  $E_1$  differential is the primitive Hamiltonian shadow acting on the source one- $\psi$  label, and its first nonzero piece is the obstruction class to the strict centrality of the primitive-shadow lift.

The four components are independent in the following sense. Vanishing of (i) is a graded commutativity test on the unreduced target and is automatic in the algebraic current model. Vanishing of (ii) is the centrality homotopy test of Proposition E.49. Vanishing of (iii) is the boundary QME content of Theorem E.35, whose first finite-window obstruction is  $\theta_3$  (Theorem E.38). Vanishing of (iv) is the descent comparison; in the algebraic current model this gap is zero by the same finite-list axiomatization. For genuine unreduced  $Z_{\text{fact}}^{P_o}$ -morphisms the Weiss/Ran comparison requires its own descent datum.

*Proof.* Necessity in each component.

Component (i). Any factorization-center morphism into  $Z_{\text{fact}}^{P_o}$  ( $\text{Obs}_{\text{open}}$ ) sends every source generator to a graded center element by definition, hence the product centrality defect is a  $d_M$ -boundary on every generator pair. The cohomology class  $\text{ob}_{\text{cent}}^{\text{prod}}$  of a candidate lift is the obstruction to this strict graded centrality; its vanishing is necessary. In the algebraic current model with graded-commutative target it is automatic (Theorem E.10).

Component (ii). A factorization-center morphism into  $Z_{\text{fact}}^{P_o}$  intertwines the source CE/Schouten bracket with the target  $P_o$ -bracket. The deviation from this intertwining is precisely the row sum  $R_{x,y,M}^{\text{cent}}$  of Proposition E.49; its cohomology class is the obstruction to choosing a centrality homotopy. Vanishing is necessary because the homotopy is exactly the datum making the source CE differential commute with the target  $P_o$ -bracket on the image of  $\Phi_{\text{fact}}^{\text{unred}}$ .

Component (iii). A factorization-center morphism produces a curved Maurer–Cartan element of the brane-defect kernel complex. Its curvature class is exactly the residual  $R_{n,M}^{\text{ns}}$  of Theorem E.35, in the inverse-limit version recorded as  $(\mathfrak{o}_{n,M}^{\text{ns}})_M$ . Vanishing, together with the  $\varprojlim^1 H^\circ$  compatibility class, is the unique condition making the curved Maurer–Cartan element actually flat. The first finite-window obstruction occurs at the  $\theta_3$  row by Theorem E.38.

Component (iv). The factorization-center construction depends globally on the chosen Weiss/Ran cover; a different cover may give a different value of the same morphism on overlapping factorizations. The codimension-one Čech difference between Weiss and Ran covers is a class in the appropriate boundary-kernel  $H^\circ$ ; its vanishing is the descent compatibility for the factorization-center morphism. In the algebraic current model with the displayed finite-list axiomatization this difference is zero, but the genuine unreduced  $Z_{\text{fact}}^{P_o}$ -target carries the comparison nontrivially in general.

Sufficiency. Conversely, suppose all four components vanish. Construct the lift by induction on  $\hbar_{\text{QME}}$ -order and finite window  $M$ . At order zero the candidate is the algebraic current presentation  $\Phi_I^\chi$  of Construction E.9, which already supplies a strict (zero-homotopy) factorization-center morphism

into the algebraic current target (Theorem E.10); hence (i) vanishes at order zero. At each order  $n \geq 1$  and each window  $M$ :

- vanishing of (iii) supplies a degree-zero counterterm  $C_{n,M}$  with  $d_{\text{ns},M} C_{n,M} = -R_{n,M}^{\text{ns}}$  (Theorem E.35); together with the  $\varprojlim^1 H^0$ -primitive-compatibility class, this assembles a compatible counterterm tower;
- vanishing of (ii) supplies a centrality homotopy  $H_{x,y}^M$  with  $R_{x,y,M}^{\text{cent}} + d_M H_{x,y}^M = 0$  for every source generator pair (Proposition E.49);
- vanishing of (iv) makes the resulting morphism agree on Weiss and Ran cover overlaps, by the codimension-one Čech–Roos comparison;
- vanishing of (i) at every window inductively proves strict graded centrality on the constructed lift.

The four vanishings are the only obstructions to inductively extending the algebraic current presentation to a strict  $Z_{\text{fact}}^{P_0}$ -morphism: each is a separate row in the curved Maurer–Cartan equation of Theorem E.35 and its Hom-valued kernel companion, and any one nonzero component blocks the inductive step at the corresponding row. Hence the obstructions are necessary and sufficient for the  $Z_{\text{fact}}^{P_0}$ -lift to exist.

The associated-graded statement. Take the primitive shadow filtration of Theorem 5.132. At the associated-graded level, the source differential is the CE differential and the target differential is the Hamiltonian/Schouten bracket on primitive trace classes. The  $E_1$  page of the spectral sequence converging to  $\text{ob}_{\text{cent}}^{P_0}$  is exactly this primitive-shadow target; its differential is the Hamiltonian shadow on one- $\psi$  primitive trace classes, as computed in Theorem 5.119. The first nonzero  $E_i$  class is the obstruction to strict primitive-shadow centrality on the unreduced source, which is the classical primitive obstruction.  $\square$

*Remark E.54 (Locus of each component of  $\text{Ob}_{\text{cent/QME/descent}}$ ).* The component  $\text{ob}_{\text{cent}}^{\text{prod}}$  is nonzero outside the graded-commutative algebraic current target of Definition E.8, on a noncommutative  $\hbar$ -deformed target where the product centrality defect carries a Capelli-type contribution. The polynomial-obstruction boundary (Theorem E.6) exhibits the same obstruction in the polynomial open BV/Koszul category: polynomial open BV observables have locally nilpotent translation action, while the principal-part module requires non-locally-nilpotent action for strict product centrality.

The component  $\text{ob}_{\text{cent}}^{P_0}$  is detected at the smallest window where the brane-defect bracket of two source generators carries a nonzero scalar-zero contribution whose primitive requires a compatible centrality homotopy  $H_{x,y}^M$  (Proposition E.49). In the minimal full-equivariant branch the row vanishes and the zero homotopy suffices; the script `scripts/finite_window_graph_array.py` reports the same in the centrality-homotopy fixture.

The component  $\text{ob}_{\text{QME}}^{\text{bd}}$  is nonzero at  $\theta_3$  in case (a) of Theorem E.38, the counterterm-omitted sub-theorem; case (b) of the same theorem gives  $\text{ob}_{\text{QME}}^{\text{bd}} = 0$  at  $\theta_3$  under the supplied counterterm  $C_{\theta_3,M}$ .

The component  $\partial_{\check{C}/R}^{\text{Weiss}}$  is nonzero when the codimension-one face of the Weiss-versus-Ran cover comparison carries a non-vanishing boundary-kernel class; this is a separate descent comparison treated in the factorization-current conventions appendix.

## F RADIAL PARTS AND THE MOYAL NORMALIZATION

Formal Weyl quantization of the constant symplectic plane and stable scalar-reduced trace map on the Chan–Paton brane stack disagree at each bidegree  $(a, b)$  by a single class. The disagreement lives in a finite-dimensional bidegree-local complex whose source records cyclic-word boundary letters and whose target combines cyclic-word incidence with the residual normal-diagram corrections. Concretely,  $W_{a,b}$  is

the rational span of words containing the moment-map letter  $[x, y]$  once;  $V_{a,b}$  is the cyclic-word target of the necklace boundary;  $\text{Diag}_{a,b}$  is the free indexed normal-diagram space carrying contraction-degree  $\geq 1$  (PBW) corrections; and  $B_{a,b}$  is the combined coboundary  $(\partial, C_{a,b}^+)$  of Definition F.21. The discrepancy is then

$$\Omega_{a,b}^{\text{rad}} \in \text{coker } B_{a,b}, \quad B_{a,b}: W_{a,b} \longrightarrow V_{a,b} \oplus \text{Diag}_{a,b},$$

in the bidegree-local radial bicomplex  $(W_{a,b} \xrightarrow{\partial} V_{a,b}, W_{a,b} \xrightarrow{C_{a,b}^+} \text{Diag}_{a,b})$  of Definition F.21. This class vanishes in every bidegree: Theorem F.28 gives  $\Omega_{a,b}^{\text{rad}} = 0$  under the radial-parts Statement F.8, and Proposition F.31 gives unconditional vanishing on the total-degree-15-and-below window by stable Procesi–Razmyslov reduction.

The cyclic-word incidence complex  $\partial: W_{a,b} \rightarrow V_{a,b}$  is the  $\hbar^0$  skeleton of the radial bicomplex; the classical lift  $\partial c = T_{a,b}$  is the ordinary necklace Stokes identity of Lemma F.24 ( $T_{a,b}$  the necklace target of a moment-map insertion). The free indexed normal-diagram target  $\text{Diag}_{a,b}$  records the residual contraction-degree  $\geq 1$  (PBW) corrections and the Capelli  $\hbar N$  term; the residual class  $R_{a,b}^{\text{free}} \in \text{Diag}_{a,b}$  (the PBW residual of the Weyl/cyclic-trace comparison) lies entirely in this positive part. Its vanishing is proved by testing on the Cartan (Lemmas F.34 and F.11) against the Levasseur–Stafford radial-kernel input. A signed-row obstruction  $(\phi, -\lambda) \in \ker B_{a,b}^*$  (a covector pairing a cyclic-word functional  $\phi \in V_{a,b}^*$  with a diagonal-target functional  $\lambda \in \text{Diag}_{a,b}^*$ ) of the  $\lambda(E_{a,b}^+) - \phi(T_{a,b}) \neq 0$  type is excluded under that input.

The formal Weyl/Moyal coefficients on the constant symplectic plane and the Capelli triangular trace normalization on the matrix Weyl algebra reduce the comparison to a finite-dimensional linear-algebra problem in the free indexed normal-diagram algebra; finite-rank Procesi–Razmyslov stable injectivity promotes free identities to all- $N$  identities. The Vandermonde-conjugated Harish-Chandra radial-parts datum [8–11] is named in Statement F.8; the structural all-bidegree theorem uses clauses (1) and (2) (Levasseur–Stafford radial-kernel) and clause (3) only for the specific Moyal-bracket polynomial of bidegree  $(a, b)$ . Pure-power clause (3) is tautological because Weyl-symmetric and normal-ordered traces of pure powers coincide. The reduced open-line Wick graph realization with midpoint propagator  $\frac{1}{2} \text{sgn}(t - s)$  realizes the same Moyal product to all orders and is the bridge to the open-line BV sector.

The edge strips  $\mathfrak{o}_{2,b} = \mathfrak{o}_{a,2} = 0$  close in closed form (Proposition F.32). The Procesi–Razmyslov reduction (Proposition F.31) closes each interior bidegree unconditionally by finite rational linear algebra; the explicit list of bidegrees verified in this mode reaches through total degree 15 and is given in Remark F.42. Reflection symmetry  $(a, b) \leftrightarrow (b, a)$  is implemented by the symplectic involution  $z_1 \leftrightarrow z_2, X \leftrightarrow Y$ ; both sides of the radial bicomplex are equivariant under it.

*Definition F.1 (Formal Weyl/Moyal convention).* Let

$$P = \partial_{z_1} \otimes \partial_{z_2} - \partial_{z_2} \otimes \partial_{z_1}.$$

The formal Weyl product on  $\mathbb{C}[[z_1, z_2]][[\hbar]]$  is

$$f * g = m \circ \exp\left(\frac{\hbar}{2} P\right)(f \otimes g),$$

and

$$[f, g]_{\text{raw}} = f * g - g * f, \quad \{f, g\}_{\hbar} = \hbar^{-1} [f, g]_{\text{raw}}.$$

The reduced quantum Hamiltonian Lie algebra is the scalar quotient

$$\bar{A}_{\hbar} = \mathbb{C}[[z_1, z_2]][[\hbar]] / \mathbb{C}[[\hbar]]$$

as a Lie algebra. The construction uses this quotient only in the Lie-algebra sense.

THEOREM F.2 (*Exact formal Weyl/Moyal coefficients*). For  $f = z_1^k z_2^l$  and  $g = z_1^m z_2^n$ ,

$$P^r(f, g) = \sum_{a=0}^r (-1)^a \binom{r}{a} (k)_{r-a} (l)_a (m)_a (n)_{r-a} z_1^{k+m-r} z_2^{l+n-r}.$$

Consequently

$$[f, g]_{\text{raw}} = \sum_{\substack{r \geq 1 \\ r \text{ odd}}} \frac{2}{2^r r!} \hbar^r P^r(f, g),$$

equivalently, for  $s \geq 0$ , the raw coefficient of  $\hbar^{2s+1}$  is

$$\frac{1}{2^{2s}(2s+1)!} P^{2s+1}(f, g),$$

and the normalized Hamiltonian bracket has coefficient

$$\frac{\hbar^{2s}}{2^{2s}(2s+1)!} P^{2s+1}(f, g).$$

The first and third raw commutator coefficients are

$$\hbar P(f, g), \quad \frac{\hbar^3}{24} P^3(f, g),$$

and all even raw commutator coefficients vanish. This is a theorem inside the formal Weyl algebra of the constant symplectic plane, independent of finite- $N$  radial-parts input and Costello graph specialization.

*Proof.* The two summands in  $P = \partial_{z_1} \otimes \partial_{z_2} - \partial_{z_2} \otimes \partial_{z_1}$  commute as differential operators on the tensor product. The binomial theorem gives

$$P^r = \sum_{a=0}^r (-1)^a \binom{r}{a} \partial_{z_1}^{r-a} \partial_{z_2}^a \otimes \partial_{z_1}^a \partial_{z_2}^{r-a}.$$

Applying this to the two monomials gives the falling-factorial formula. Interchanging  $f$  and  $g$  changes  $P^r(f, g)$  by  $(-1)^r$ , so the commutator retains exactly the odd powers and contributes the coefficient  $2/(2^r r!)$  in order  $r$ .  $\square$

COROLLARY F.3 (*All odd Moyal coefficients*). The Weyl/Moyal commutator has only odd raw powers of  $\hbar$ . Its odd raw and normalized coefficients are

$$[f, g]_{\text{raw}}^{(2s+1)} = \frac{\hbar^{2s+1}}{2^{2s}(2s+1)!} P^{2s+1}(f, g), \quad \{f, g\}_\hbar^{(2s)} = \frac{\hbar^{2s}}{2^{2s}(2s+1)!} P^{2s+1}(f, g).$$

In particular the next two raw coefficients after the third-order term are

$$\frac{\hbar^5}{1920} P^5(f, g), \quad \frac{\hbar^7}{322560} P^7(f, g).$$

*Proof.* Put  $r = 2s + 1$  in Theorem F.2. Then  $2/(2^{2s+1}(2s+1)!) = 1/(2^{2s}(2s+1)!)$ . Dividing by  $\hbar$  gives the normalized coefficient.  $\square$

COROLLARY F.4 (*First and third formal graph weights*). In the ordered formal product  $m \circ \exp(\hbar P/2)$ , the one-cross-contraction term has weight  $\hbar P(f, g)/2$ , and the three-cross-contraction term has weight  $\hbar^3 P^3(f, g)/48$ . After antisymmetrising the ordered product, the commutator weights are

$$\hbar P(f, g), \quad \frac{\hbar^3}{24} P^3(f, g).$$

*Proof.* The ordered exponential contributes

$$\frac{1}{r!} \left( \frac{\hbar}{2} \right)^r P^r(f, g)$$

in order  $r$ . For  $r = 1$  this is  $\hbar P(f, g)/2$ ; for  $r = 3$  it is  $\hbar^3 P^3(f, g)/(3! 2^3) = \hbar^3 P^3(f, g)/48$ . Since  $P(g, f) = -P(f, g)$  and  $P^3(g, f) = -P^3(f, g)$ , subtracting the reverse ordered product doubles these two ordered contributions.  $\square$

Example F.5 (*A higher-order coefficient check*). For

$$f = z_1^4 z_2^2, \quad g = z_1^2 z_2^4,$$

the first nonzero odd contractions are

$$P(f, g) = 12 z_1^5 z_2^5, \quad P^3(f, g) = -960 z_1^3 z_2^3, \quad P^5(f, g) = 11520 z_1 z_2.$$

Hence

$$[f, g]_{\text{raw}} = 12 \hbar z_1^5 z_2^5 - 40 \hbar^3 z_1^3 z_2^3 + 6 \hbar^5 z_1 z_2.$$

The third coefficient is  $-960/24 = -40$ . The fifth coefficient is  $11520/1920 = 6$ . This example checks the numerical normalization beyond the first nontrivial term, independently of radial parts.

*Proof.* Apply the falling-factorial formula of Theorem F.2. For  $r = 1$ ,

$$4 \cdot 4 - 2 \cdot 2 = 12.$$

For  $r = 3$ ,

$$\begin{aligned} P^3(f, g) &= (24 \cdot 24 - 3 \cdot 12 \cdot 2 \cdot 2 \cdot 12 + 3 \cdot 4 \cdot 2 \cdot 2 \cdot 4) z_1^3 z_2^3 \\ &= -960 z_1^3 z_2^3. \end{aligned}$$

For  $r = 5$ , only the  $a = 1$  and  $a = 2$  summands survive, and they give

$$(-5 \cdot 24 \cdot 2 \cdot 2 \cdot 24 + 10 \cdot 24 \cdot 2 \cdot 2 \cdot 24) z_1 z_2 = 11520 z_1 z_2.$$

Substituting the odd Moyal coefficients  $1$ ,  $1/24$ , and  $1/1920$  gives the displayed raw commutator.  $\square$

Definition F.6 (*Matrix Weyl algebra and Weyl traces*). Let  $\mathcal{W}_N$  be the completed Weyl algebra generated by entries of two  $N \times N$  matrices  $X, Y$  with

$$[X^i_j, Y^k_l] = \hbar \delta_l^i \delta_j^k, \quad [X^i_j, X^k_l] = [Y^i_j, Y^k_l] = 0.$$

The normal-ordered quantum comoment for conjugation is

$$\widehat{\mu}_\epsilon = -\text{Tr}([\epsilon, X]Y), \quad \hbar^{-1}[\widehat{\mu}_\epsilon, a] = \epsilon \cdot a.$$

For  $f \in \mathbb{C}[z_1, z_2][[\hbar]]$ , define the Weyl-ordered trace

$$J_N(f) = \text{Tr Op}_W(f)(X, Y).$$

The degree-zero quantum Hamiltonian reduction uses the effective traceless comoment ideal

$$\mathcal{N}_{\mathcal{W}_N}(\mathcal{W}_N \widehat{\mu}(\mathfrak{sI}_N)) / \mathcal{W}_N \widehat{\mu}(\mathfrak{sI}_N).$$

The scalar Hamiltonian line survives this moment ideal; below it is removed by quotienting the empty trace before connected large- $N$  extraction.

LEMMA F.7 (*Capelli–Weyl triangular normalization*). Let  $T_{a,b} = \text{Tr}(X^a Y^b)$  be the normal-ordered trace. In the quantum Hamiltonian reduction, the locally proved Weyl-normal-ordering formula is

$$J_N(z_1^a z_2^b) \equiv \sum_{r=0}^{\min(a,b)} \left(-\frac{\hbar N}{2}\right)^r r! \binom{a}{r} \binom{b}{r} T_{a-r, b-r}.$$

The inverse triangular system is

$$T_{a,b} \equiv \sum_{r=0}^{\min(a,b)} \left(\frac{\hbar N}{2}\right)^r r! \binom{a}{r} \binom{b}{r} J_N(z_1^{a-r} z_2^{b-r}).$$

*Proof.* Total Weyl symmetrisation is obtained from normal ordering by commuting  $Y$ 's past  $X$ 's. Each contraction contributes the Capelli contact term coming from

$$YX - XY + \hbar NI \equiv 0 \pmod{\mathcal{W}_N \widehat{\mu}(\mathfrak{sI}_N)}.$$

Choosing  $r$  contracted  $X$ -slots and  $r$  contracted  $Y$ -slots gives  $\binom{a}{r} \binom{b}{r}$ , pairing them gives  $r!$ , and Weyl ordering contributes  $2^{-r}$ . The sign is negative because the displayed moment relation rewrites a normal-ordering swap as the lower triangular term  $-\hbar N$ . The diagonal term is  $r = 0$ , hence the system is unipotent over  $\mathbb{C}[[\hbar N]]$ ; inversion changes  $-\hbar N/2$  to  $+\hbar N/2$ .

The cited Capelli literature supplies the standard source for the Capelli contact phenomenon, and the displayed contraction count fixes the bivariate trace normalization used here.  $\square$

Statement F.8 (*Radial-parts input data*). The finite- $N$  radial-parts step uses the following Harish-Chandra radial-parts theorem for invariant differential operators, in the form refined by Wallach and by Levasseur–Stafford [8–11]. After identifying  $\mathcal{W}_N \simeq \mathcal{D}_\hbar(\mathfrak{gl}_N)$ , this input uses the Harish-Chandra map on invariants whose kernel is the quantum moment-map ideal and hence injects the quantum Hamiltonian reduction by conjugation into Weyl-invariant differential operators on the regular Cartan. With  $\lambda_1, \dots, \lambda_N$  the Cartan coordinates and  $\Delta = \prod_{i < j} (\lambda_i - \lambda_j)$ , the radial-part map is conjugated by  $\Delta$ .

The cited radial-parts literature sources the Harish-Chandra homomorphism, regular-Cartan landing, and kernel statement. The exact Weyl-trace image in item (3) is the matched normalization datum used in the formal-local comparison. The cited radial-parts sources supply the Harish-Chandra kernel theorem; the exact Weyl/Capelli convention enters here as an additional normalization. The finite exact checks below test it in ranks 2 and 3; the all- $N$  radial-parts input remains the structural hypothesis.

The input consists of the following three assertions.

- (i) The Vandermonde-conjugated radial map is a homomorphism from the invariant quantum Hamiltonian reduction by conjugation to Weyl-invariant differential operators on the regular Cartan.

- (ii) Its kernel is precisely the quantum moment-map ideal; hence it is injective on the reduced quotient.
- (iii) The Weyl trace generators used here have image

$$\text{Rad}_o(J_N(f)) = \Delta^{-1} \left( \sum_{i=1}^N \text{Op}_W(f)(\lambda_i, -\hbar \partial_{\lambda_i}) \right) \Delta.$$

The nonlinear-shear test in Lemma F.13 shows that pure  $z_1$ - and  $z_2$ -power images together with classical symbol-level shear compatibility determine only the pure-power datum. Item (3) requires a genuine quantum-corrected Egorov theorem in the Weyl algebra.

*Definition F.9 (Radially normalized Weyl trace class).* Assume only the homomorphism and injectivity clauses of Statement F.8. For  $f \in \mathbb{C}[z_1, z_2][[\hbar]]$ , a class  $J_N^{\text{rad}}(f)$  in the degree-zero quantum Hamiltonian reduction is called radially normalized if

$$\text{Rad}_o(J_N^{\text{rad}}(f)) = \Delta^{-1} S_N(f) \Delta.$$

Existence is a normalization hypothesis; uniqueness follows from radial injectivity. Statement F.8(3) is exactly the assertion that the ordinary Weyl trace  $J_N(f)$  equals  $J_N^{\text{rad}}(f)$ .

*Definition F.10 (Vandermonde gauge variables).* Set

$$\nabla_i^\Delta = \Delta^{-1}(-\hbar \partial_{\lambda_i}) \Delta = -\hbar \partial_{\lambda_i} - \hbar \sum_{j \neq i} \frac{1}{\lambda_i - \lambda_j}.$$

The logarithmic summand is the type  $\mathcal{A}$  Dunkl–Calogero correction in the Vandermonde gauge. For  $f \in \mathbb{C}[z_1, z_2][[\hbar]]$ , put

$$S_N(f) = \sum_{i=1}^N \text{Op}_W(f)(\lambda_i, -\hbar \partial_{\lambda_i})$$

and

$$R_N(f) = \sum_{i=1}^N \text{Op}_W(f)(\lambda_i, \nabla_i^\Delta) = \Delta^{-1} S_N(f) \Delta.$$

LEMMA F.11 (*Dunkl–Calogero correction cancellation*). The Vandermonde-gauge variables satisfy

$$[\lambda_i, \lambda_j] = 0, \quad [\nabla_i^\Delta, \nabla_j^\Delta] = 0, \quad [\lambda_i, \nabla_j^\Delta] = \hbar \delta_{ij}.$$

Consequently

$$\hbar^{-1} [R_N(f), R_N(g)] = R_N(\{f, g\}_\hbar)$$

for all  $f, g$ . In particular, all logarithmic Dunkl–Calogero terms introduced by expanding  $\Delta^{-1} S_N(-) \Delta$  cancel in the commutator identity.

*Proof.* The operator  $\nabla_i^\Delta$  is conjugate to  $-\hbar \partial_{\lambda_i}$  by multiplication by  $\Delta$ . The three displayed commutation relations are therefore the ordinary Weyl relations for  $(\lambda_i, -\hbar \partial_{\lambda_i})$ , conjugated by  $\Delta$ . The correction terms are a pure gauge connection: their curvature is zero because the unconjugated partial derivatives commute. Hence the  $i$ -th summand is again a one-particle Weyl representation of the formal Moyal algebra, and different summands commute. This gives the displayed commutator identity. The Vandermonde-conjugated form absorbs the Calogero potential.  $\square$

LEMMA F.12 (*Radial-map injectivity on the reduced quotient*). Under Statement F.8, if an invariant Weyl operator  $D$  in the normalizer of the quantum moment-map ideal satisfies

$$\text{Rad}_o(D) = 0,$$

then  $D$  is zero in the degree-zero quantum Hamiltonian reduction.

*Proof.* Statement F.8 identifies the kernel of the Vandermonde-conjugated Harish-Chandra radial map with the quantum moment-map ideal. Passing to the normalizer quotient by that ideal gives an injective map into Weyl-invariant differential operators on the regular Cartan. Thus an element with zero radial image represents the zero class in the reduced quotient.  $\square$

LEMMA F.13 (*Nonlinear shear obstruction to weaker radial-parts input*). Pure-power radial images plus classical symbol-level compatibility with nonlinear shears determine only the pure-power Weyl-trace image normalization. In the one-particle Weyl algebra  $[\lambda, y] = \hbar$ , the coefficient of  $t$  in

$$(y + t\lambda^2)^4$$

is

$$4\lambda^2 y^3 - 12\hbar\lambda y^2 + 8\hbar^2 y,$$

whereas

$$4 \text{Op}_W(z_1^2 z_2^3) = 4\lambda^2 y^3 - 12\hbar\lambda y^2 + 6\hbar^2 y.$$

The difference  $2\hbar^2 y$  is non-scalar. Thus Statement F.8(3) requires the quantum Egorov correction in addition to classical nonlinear shear covariance.

*Proof.* Expand to first order in  $t$ :

$$(y + t\lambda^2)^4 = y^4 + t \sum_{j=0}^3 y^j \lambda^2 y^{3-j} + O(t^2).$$

Using  $y\lambda = \lambda y - \hbar$ , the sum normal-orders to  $4\lambda^2 y^3 - 12\hbar\lambda y^2 + 8\hbar^2 y$ . Weyl symmetrisation of  $z_1^2 z_2^3$  gives  $\lambda^2 y^3 - 3\hbar\lambda y^2 + \frac{3}{2}\hbar^2 y$ , hence the displayed value after multiplication by 4. The discrepancy is a nonconstant one-particle operator, hence survives the scalar Hamiltonian quotient.  $\square$

PROPOSITION F.14 (*Quantum shear congruence suffices for the all- $N$  image*). Assume the homomorphism and injectivity clauses of Statement F.8, and assume the pure power images

$$\text{Rad}_o(J_N(z_1^r)) = \Delta^{-1} S_N(z_1^r) \Delta, \quad \text{Rad}_o(J_N(z_2^r)) = \Delta^{-1} S_N(z_2^r) \Delta$$

for all  $r \geq 0$ . Suppose moreover that, for all  $a, b \geq 0$ , the quantum shear congruence

$$\hbar^{-1} [J_N(z_1^{a+1}/(a+1)), J_N(z_2^{b+1})] \equiv J_N(\{z_1^{a+1}/(a+1), z_2^{b+1}\}_{\hbar}) \pmod{I_N}$$

holds in the degree-zero quantum Hamiltonian reduction, where  $I_N$  is the quantum moment-map ideal. Then the exact Weyl-trace image in Statement F.8(3) holds for every monomial  $z_1^a z_2^b$ , hence for every polynomial  $f$ .



*Proof.* The normalized Moyal bracket gives

$$\{z_1^{a+1}/(a+1), z_2^{b+1}\}_{\hbar} = (b+1)z_1^a z_2^b + \sum_{s \geq 1} \frac{\hbar^{2s} (a+1)_{2s+1} (b+1)_{2s+1}}{(a+1)2^{2s} (2s+1)!} z_1^{a-2s} z_2^{b-2s}.$$

Every summand after the leading term has lower total degree. The pure power cases start the induction. Assume the asserted radial image is known in all total degrees  $< a + b$ . Applying the radial homomorphism to the quantum shear congruence identifies the radial image of the left-hand commutator with the Moyal commutator of the already known pure-power radial images. Lemma F.34 computes this as the Vandermonde-conjugated Weyl quantization of the displayed Moyal bracket. By the induction hypothesis all lower-degree terms have the required images, so the leading coefficient  $(b+1)$  determines the image of  $J_N(z_1^a z_2^b)$ . Radial injectivity then identifies the class in the reduced quotient.  $\square$

LEMMA F.15 (*Moment-ideal primitive form of the quantum shear input*). In the notation of Proposition F.14, the quantum shear congruence is equivalent to the following explicit left-ideal membership problem. Put

$$E_{a,b,N} = \sum_{j=0}^b \text{Tr}(Y^j X^a Y^{b-j}) - J_N(\{z_1^{a+1}/(a+1), z_2^{b+1}\}_{\hbar}).$$

Then the congruence holds if and only if  $E_{a,b,N} \in \mathcal{W}_N \widehat{\mu}(\mathfrak{sl}_N)$ . Equivalently, after choosing normal-ordered matrix generators  $\widehat{\mu}^i_j$  for the traceless quantum moment-map ideal, one must construct coefficients  $A^j_i(a, b, N) \in \mathcal{W}_N$  such that

$$E_{a,b,N} = \sum_{i,j} A^j_i(a, b, N) \widehat{\mu}^i_j.$$

*Proof.* Since the pure powers are already Weyl ordered,  $J_N(z_1^{a+1}/(a+1)) = \text{Tr}(X^{a+1})/(a+1)$  and  $J_N(z_2^{b+1}) = \text{Tr}(Y^{b+1})$ . The Weyl relations give

$$\hbar^{-1} [\text{Tr}(X^{a+1})/(a+1), Y] = X^a.$$

Applying the Leibniz rule to  $Y^{b+1}$  gives

$$\hbar^{-1} [J_N(z_1^{a+1}/(a+1)), J_N(z_2^{b+1})] = \sum_{j=0}^b \text{Tr}(Y^j X^a Y^{b-j}).$$

Subtracting the  $J_N$ -image of the normalized Moyal bracket gives exactly  $E_{a,b,N}$ . Reduction modulo  $I_N$  is reduction modulo the left ideal  $\mathcal{W}_N \widehat{\mu}(\mathfrak{sl}_N)$ , so the displayed congruence is precisely the asserted membership. The coefficient formula is the same membership written after a choice of generators for that left ideal.  $\square$

Statement F.16 (*Free trace-diagram quantum-lift obstruction*). Put  $M = YX - XY + \hbar NI$ , and interpret

$$\text{Tr}(WM) = \sum_{i,j} W^j_i M^i_j$$

as a left-ideal expression. Let  $\mathbb{C}_{a,b}$  be the rational vector space spanned by words  $W$  with  $a - 1$  letters  $X$  and  $b - 1$  letters  $Y$ , and let  $\text{Cyc}_{a,b}$  be the cyclic word space with  $a$  letters  $X$  and  $b$  letters  $Y$ . The leading term of the desired primitive is the cyclic boundary equation

$$\partial c = \sum_{j=0}^b [Y^j X^a Y^{b-j}] - \frac{b+1}{\binom{a+b}{a}} \sum_{U \in \text{Sh}(a,b)} [U], \quad \partial W = [WYX] - [WXY].$$

An all- $N$  primitive is stronger. It requires equality after free indexed normal ordering, including every contraction term which splits a trace into multi-traces:

$$\mathcal{N}_{\text{free}} \left( \sum_W c_W \text{Tr}(WM) \right) = \mathcal{N}_{\text{free}}(E_{a,b,N}).$$

The cyclic equation is the associated-graded part of this quantum equation.

In bidegree  $(a, b) = (4, 4)$ , the lexicographic pivot solution of the cyclic equation is

$$\begin{aligned} C_{4,4}^{\text{cl}} = & \frac{31}{7} \text{Tr}(XXXYYYM) - \frac{4}{7} \text{Tr}(XXYXYYM) - \frac{4}{7} \text{Tr}(XXYYXYM) \\ & + \frac{27}{7} \text{Tr}(XXYYYXM) + \frac{23}{7} \text{Tr}(XYXXYYM) + \frac{1}{7} \text{Tr}(XYXYXYM) \\ & + \text{Tr}(XYXYXYM) - \frac{2}{7} \text{Tr}(XYXXXYM) + \frac{15}{7} \text{Tr}(XYXYXYM). \end{aligned}$$

Its quantum residual is nonzero. In rank  $N = 2$ , exact normal ordering gives

$$\begin{aligned} R_{4,4,2}^{\text{cl}} &:= E_{4,4,2} - C_{4,4}^{\text{cl}} \\ &= \frac{43}{70} \hbar^2 (4 \text{Tr}(XXYY) - 5 \text{Tr}(XYXY) + 2 \text{Tr}(XYXX) \\ &\quad - 5 \text{Tr}(YXYX) + 4 \text{Tr}(YYXX)) \neq 0. \end{aligned}$$

After expanding the displayed traces into normal-ordered matrix entries, the coefficient of  $\hbar^2 X_1^1 X_1^1 Y_2^1 Y_2^1$  is  $43/7$ .

The residual is killed by the classical-kernel correction

$$\begin{aligned} K_{4,4} = & \frac{43}{14} (\text{Tr}(XXYXYM) - \text{Tr}(XYXXYYM) \\ & - \text{Tr}(XYXYXYM) + \text{Tr}(XYXYXXM)), \end{aligned}$$

whose cyclic boundary is zero. The free normal-form computation below proves that this correction cancels the  $(4, 4)$  residual before any finite-rank trace identity is imposed. The decorated PBW Stokes identity closes uniformly: by Theorem F.28, for every bidegree

$$\Omega_{a,b}^{\text{rad}} = 0 \iff D_{a,b}^{\square} H_{a,b} = R_{a,b}^{\text{free}}, \quad D_{a,b}^{\square} = C_{a,b}^+ \partial_2,$$

conditional on Statement F.8. Unconditionally, the bidegrees in Remark F.42 are closed by Procesi–Razmyslov verification (Proposition F.31); the normalized signed-dual-row obstruction set is empty there.

*Proof.* The leading  $\hbar^0$  component of  $\text{Tr}(WM)$  is  $[WYX] - [WXY]$  in the cyclic trace quotient; comparing it with the leading term of  $E_{a,b,N}$  gives the displayed cyclic equation. Normal ordering in the indexed Weyl algebra contracts every occurrence of a  $Y$ -entry moved past an  $X$ -entry. These contractions identify matrix indices and may split one cyclic trace into a product of cyclic traces. Hence equality in the cyclic quotient is only the associated graded condition.

The  $(4, 4)$  coefficients displayed above solve the cyclic equation in the stated pivot gauge. Direct exact normal ordering in  $\mathcal{W}_2$  gives the displayed residual. Its nonvanishing follows from the indicated normal monomial coefficient. Applying  $\partial$  to the four words in  $K_{4,4}$  gives two copies of each resulting cyclic word with opposite signs, so  $K_{4,4} \in \ker \partial$ . The finite computation proves the obstruction to the uncorrected cyclic lift. The all- $N$  identity for  $K_{4,4}$  is supplied by the free trace-diagram normal form below.  $\square$

*Definition F.17 (Free indexed normal diagrams).* Let  $u = L_1 \cdots L_m$  be a word in  $X, Y$ . Write the trace  $\text{Tr}(u)$  as the directed  $m$ -gon with vertices  $v_0, \dots, v_{m-1}$  and edge  $e_r : v_r \rightarrow v_{r+1}$ , labelled  $L_r$ , with indices read modulo  $m$ . A contraction matching  $\Gamma$  is a partial matching of pairs

$$p < q, \quad L_p = Y, \quad L_q = X.$$

For each pair  $(p, q) \in \Gamma$ , delete the two edges  $e_p, e_q$  and identify

$$v_q \sim v_{p+1}, \quad v_p \sim v_{q+1}.$$

Let  $e(u, \Gamma)$  be the number of isolated vertices left after these deletions and identifications. Let  $D(u, \Gamma)$  be the remaining directed labelled graph, with all  $X$ -edges placed before all  $Y$ -edges in the ambient commutative normal-ordered algebra of matrix entries. Disconnected components are retained; they are the split-trace terms. The empty graph is denoted  $\mathbf{1}$ .

**THEOREM F.18 (Free indexed normal-ordering formula).** In the universal indexed matrix Weyl algebra with

$$Y^k_l X^i_j = X^i_j Y^k_l - \hbar \delta_l^i \delta_j^k,$$

the normal-ordered trace word is

$$\mathcal{N}_{\text{free}} \text{Tr}(u) = \sum_{\Gamma} (-\hbar)^{|\Gamma|} N^{e(u, \Gamma)} D(u, \Gamma),$$

where the sum runs over all contraction matchings  $\Gamma$  of  $Y$ -before- $X$  inversions in  $u$ . Consequently

$$\begin{aligned} \mathcal{N}_{\text{free}} \text{Tr}(WM) &= \mathcal{N}_{\text{free}} \text{Tr}(WYX) - \mathcal{N}_{\text{free}} \text{Tr}(WXY) \\ &\quad + \hbar N \mathcal{N}_{\text{free}} \text{Tr}(W), \end{aligned}$$

with  $M = YX - XY + \hbar NI$ , interpreted as the left-ideal expression  $\text{Tr}(WM) = \sum_{i,j} W^j_i M^i_j$ .

*Proof.* Move the  $X$ -entries leftward through the  $Y$ -entries by the PBW relation displayed above. Choosing the commutator summand at a swap selects one pair  $Y_p, X_q$ , contributes the factor  $-\hbar$ , and imposes exactly the two Kronecker identifications  $v_q \sim v_{p+1}$  and  $v_p \sim v_{q+1}$ . Choosing the ordered summand leaves both edges and continues the PBW ordering. Since a generator can be contracted only once, the possible choices are precisely the partial matchings of original inversions. After all  $X$ -entries have been moved left, the uncontracted entries form the normal diagram  $D(u, \Gamma)$ . Each isolated index loop contributes a free summation over  $1, \dots, N$ , hence  $N$ . This proves the formula for trace words. The formula for  $\text{Tr}(WM)$  is the defining expansion of  $M$ , with the scalar matrix term contributing  $\hbar N \text{Tr}(W)$ .  $\square$

LEMMA F.19 (*Stable injectivity for normal trace diagrams*). Fix  $d$ . Let  $\text{Diag}_{\leq d}$  be the rational span of free indexed normal diagrams with at most  $d$  labelled nonempty edges, with coefficients in  $\mathbb{Q}[\hbar, N]$ . The joint specialization map to  $N \times N$  matrices for all ranks  $N \geq d$  is injective. Thus a free normal-form equality in  $\text{Diag}_{\leq d}$  is an all- $N$  matrix Weyl identity. A nonzero free residual remains visible until the stable Procesi–Razmyslov range is reached.

*Proof.* Polarize in the labelled edges. A linear relation among polarized closed directed trace diagrams of edge degree at most  $d$  is a trace identity for generic matrices of degree at most  $d$ . The Procesi–Razmyslov second fundamental theorem says that the trace identities are generated by the fundamental antisymmetrizer on  $N + 1$  index lines. Hence every identity of degree at most  $d$  is zero in ranks  $N \geq d$ . Since the coefficients are polynomials in the formal scalar  $N$ , vanishing in every rank  $N \geq d$  forces each coefficient polynomial to vanish. Depolarizing gives the stated injectivity.  $\square$

PROPOSITION F.20 (*First free residual and its kernel correction*). Use the normal-diagram notation of Definition F.17; for instance  $D(X_{0 \rightarrow 1} Y_{1 \rightarrow 0})$  denotes the directed two-edge normal diagram with one  $X$ -edge and one  $Y$ -edge. Then

$$\begin{aligned} \mathcal{N}_{\text{free}}(E_{4,4,N} - C_{4,4}^{\text{cl}}) &= \frac{43}{7} \hbar^2 \left( D(X_{0 \rightarrow 1} X_{1 \rightarrow 2} Y_{2 \rightarrow 3} Y_{3 \rightarrow 0}) \right. \\ &\quad - D(X_{0 \rightarrow 1} X_{2 \rightarrow 3} Y_{1 \rightarrow 2} Y_{3 \rightarrow 0}) \\ &\quad \left. - \hbar D(X_{0 \rightarrow 0} \mid Y_{0 \rightarrow 0}) + \hbar N D(X_{0 \rightarrow 1} Y_{1 \rightarrow 0}) \right). \end{aligned}$$

Moreover

$$\mathcal{N}_{\text{free}}(K_{4,4}) = \mathcal{N}_{\text{free}}(E_{4,4,N} - C_{4,4}^{\text{cl}}),$$

and therefore

$$\mathcal{N}_{\text{free}}(E_{4,4,N} - C_{4,4}^{\text{cl}} - K_{4,4}) = 0.$$

This is an all- $N$  identity in the free indexed normal-diagram algebra, proved before any rank specialization.

*Proof.* Apply Theorem F.18 to the nine trace words in  $C_{4,4}^{\text{cl}}$  and to the Weyl-trace expression defining  $E_{4,4,N}$ . The finite matching enumeration is rational and takes place in the free diagram algebra. It leaves exactly the four displayed normal diagrams. Applying the same formula to the four trace words in  $K_{4,4}$  gives the same four-diagram vector. The script

`scripts/quantum_shear_trace_diagram_normal_form.py`

is the reproducible enumeration: with `-case 4,4 -classical-only` it prints the four-term residual above, and with `-case 4,4` it verifies zero residual after adding  $K_{4,4}$ .  $\square$

DEFINITION F.21 (*Kernel-correction residual and decorated necklace complex*). Let  $\mathcal{W}_{a,b}$  be the rational span of words with  $a - 1$  letters  $X$  and  $b - 1$  letters  $Y$ , and let  $\mathcal{V}_{a,b} = \text{Cyc}_{a,b}$  be the rational span of cyclic words with  $a$  letters  $X$  and  $b$  letters  $Y$ . Define the two-colour necklace multigraph  $G_{a,b}$  by taking  $\mathcal{V}_{a,b}$  as its vertex span and by assigning to  $W \in \mathcal{W}_{a,b}$  the oriented edge

$$e_W : [WXY] \longrightarrow [WYX].$$

Its incidence map is

$$\partial: W_{a,b} \rightarrow V_{a,b}, \quad \partial W = [WYX] - [WXY],$$

and  $Z_{a,b} := \ker \partial$  is the cycle space.

Let  $\text{Diag}_{a,b}$  denote the positive free indexed normal-diagram target in bidegree  $(a, b)$ : the uncontracted  $\hbar^0$  cyclic component is removed and all PBW contraction, split-trace, and Capelli  $\hbar N$ -terms are retained. Write  $\pi_+$  for this projection. Define the decorated contraction cochain by

$$C_{a,b}^+(W) = \pi_+ \mathcal{N}_{\text{free}} \text{Tr}(WM), \quad M = YX - XY + \hbar NI.$$

The correction map is

$$C_{a,b} = A_{a,b} := C_{a,b}^+|_{Z_{a,b}}: Z_{a,b} \longrightarrow \text{Diag}_{a,b}.$$

Put

$$T_{a,b} = \sum_{j=0}^b [Y^j X^a Y^{b-j}] - \frac{b+1}{\binom{a+b}{a}} \sum_{U \in \text{Sh}(a,b)} [U].$$

For any solution  $c_{a,b}^{\text{cl}}$  of  $\partial c = T_{a,b}$ , define

$$E_{a,b}^+ = \pi_+ \mathcal{N}_{\text{free}}(E_{a,b,N})$$

and put

$$R_{a,b}^{\text{free}} = E_{a,b}^+ - C_{a,b}^+(c_{a,b}^{\text{cl}})$$

and define

$$\mathfrak{o}_{a,b} = [R_{a,b}^{\text{free}}] \in \text{coker } C_{a,b}.$$

This is the kernel-correction class after a choice of classical lift. The radial obstruction itself is the lift-independent class  $\Omega_{a,b}^{\text{rad}}$  and the signed-row criterion of Proposition F.25.

*Statement F.22 (Decorated kernel-correction Hodge criterion).* Give  $Z_{a,b}$  and  $\text{Diag}_{a,b}$  the inner products for which the chosen word-cycle and free normal-diagram bases are orthonormal. Let  $A = A_{a,b}$  and define

$$\Delta_{a,b}^{\text{dec}} = AA^*, \quad H_{a,b}^{\text{dec}} = \ker A^* = (\text{im } A)^\perp.$$

Let

$$\Pi_{a,b}^{\text{harm}}: \text{Diag}_{a,b} \rightarrow H_{a,b}^{\text{dec}}$$

be the orthogonal projection. For every  $a, b \geq 1$ , the following conditions are equivalent.

(i) There exists  $K_{a,b} \in Z_{a,b}$  such that

$$C_{a,b}^+(K_{a,b}) = R_{a,b}^{\text{free}}.$$

(ii) The kernel-correction class  $\mathfrak{o}_{a,b}$  vanishes.

(iii) The decorated harmonic projection vanishes:

$$\Pi_{a,b}^{\text{harm}} R_{a,b}^{\text{free}} = \mathfrak{o}.$$

(iv) Every functional  $\lambda \in \text{Diag}_{a,b}^*$  with

$$\lambda C_{a,b}^+(K) = 0 \quad (K \in Z_{a,b})$$

also satisfies  $\lambda(R_{a,b}^{\text{free}}) = 0$ .

When these equivalent conditions hold, the canonical Hodge correction is

$$K_{a,b} = A_{a,b}^* (\Delta_{a,b}^{\text{dec}})^{-1} R_{a,b}^{\text{free}},$$

with the inverse taken on  $\text{im } A_{a,b}$ . Thus the fixed-lift kernel-correction problem is equivalent to

$$\Pi_{a,b}^{\text{harm}} R_{a,b}^{\text{free}} = 0 \quad \text{in the chosen bidegree.}$$

The lift-independent obstruction  $\Omega_{a,b}^{\text{rad}}$  and its normalized signed-row form are given by Proposition F.25.

*Proof.* If  $c$  and  $c'$  solve the cyclic equation, then  $c - c' \in Z_{a,b}$ . Hence  $R(c') - R(c) = -A_{a,b}(c' - c)$ , so the class in  $\text{coker } A_{a,b}$ , and its harmonic projection, are independent of the chosen cyclic lift.

The finite-dimensional Hodge decomposition for  $A$  gives

$$\text{Diag}_{a,b} = \text{im } A \oplus \ker A^*.$$

Therefore  $R_{a,b}^{\text{free}} \in \text{im } A$  if and only if its projection to  $\ker A^*$  is zero. This proves the equivalence of the correction equation, the intermediate cokernel class, and the harmonic projection. The functional criterion is the annihilator form of the same statement:  $R \in \text{im } A$  if and only if every  $\lambda \in (\text{im } A)^\perp = \ker A^*$  vanishes on  $R$ . If  $R \in \text{im } A$ , then  $AA^*(AA^*)^{-1}R = R$  on  $\text{im } A$ , so the displayed  $K_{a,b}$  is a correction. Conversely, any correction places  $R_{a,b}^{\text{free}}$  in  $\text{im } A$ .  $\square$

*Definition F.23 (Square-cell PBW Stokes complex).* Let  $S_{a,b}$  be the rational span of square cells obtained from two commuting adjacent swaps in a cyclic word with  $a$  letters  $X$  and  $b$  letters  $Y$ . A based representative has the form

$$uXYvXYw,$$

and its oriented square is

$$uXYvXYw \rightarrow uYXvXYw \rightarrow uYXvYXw \rightarrow uXYvYXw \rightarrow uXYvXYw.$$

Each side is read as the corresponding edge  $e_W : [WXY] \rightarrow [WYX]$  of the necklace graph. This defines

$$\partial_2 : S_{a,b} \longrightarrow W_{a,b}.$$

The decorated PBW Stokes map is

$$D_{a,b}^\square = C_{a,b}^+ \partial_2 : S_{a,b} \longrightarrow \text{Diag}_{a,b}.$$

LEMMA F.24 (*Ordinary necklace Stokes theorem*). For  $a, b \geq 1$ ,

$$\text{im } \partial_2 = \ker \partial.$$

*Proof.* Work first with based words. A binary word with  $a$  letters  $X$  and  $b$  letters  $Y$  is the same as an order ideal in the rectangle poset  $[a] \times [b]$ , via its inversion set. An adjacent  $XY \leftrightarrow YX$  move toggles one extremal box. The cubical complex of the distributive lattice of order ideals, with squares generated by commuting toggles, is contractible. Hence its rational one-cycles are generated by square boundaries. Passing to cyclic words is the quotient by a finite cellular rotation action; over  $\mathbb{Q}$ , transfer identifies the quotient rational homology with the invariant rational homology upstairs. Thus every necklace cycle is a rational sum of square boundaries.  $\square$

PROPOSITION F.25 (*Dual-potential form of the radial obstruction*). Define the finite-dimensional map

$$B_{a,b}: W_{a,b} \longrightarrow V_{a,b} \oplus \text{Diag}_{a,b}, \quad B_{a,b}(W) = (\partial W, C_{a,b}^+(W)),$$

and the target

$$\eta_{a,b} = (T_{a,b}, E_{a,b}^+).$$

Let

$$\Omega_{a,b}^{\text{rad}} = [\eta_{a,b}] \in \text{coker } B_{a,b}.$$

For fixed  $(a, b)$ , the following are equivalent.

- (i)  $\Omega_{a,b}^{\text{rad}} = 0$ .
- (ii) For one, equivalently every, classical lift  $c_{a,b}^{\text{cl}}$  with  $\partial c_{a,b}^{\text{cl}} = T_{a,b}$ , there is an  $H_{a,b} \in S_{a,b}$  such that

$$D_{a,b}^{\square} H_{a,b} = R_{a,b}^{\text{free}}.$$

- (iii) For one, equivalently every, classical lift  $c_{a,b}^{\text{cl}}$ , the residual  $R_{a,b}^{\text{free}}$  lies in  $\text{im}(C_{a,b}^+|_{\ker \partial})$ .
- (iv)  $\prod_{a,b}^{\text{harm}} R_{a,b}^{\text{free}} = 0$ .
- (v) For every pair

$$(\phi, \lambda) \in V_{a,b}^* \oplus \text{Diag}_{a,b}^*$$

satisfying

$$\lambda C_{a,b}^+(W) = \phi(\partial W) \quad (W \in W_{a,b}),$$

one has

$$\lambda(E_{a,b}^+) = \phi(T_{a,b}).$$

Thus

$$\Omega_{a,b}^{\text{rad}} = 0 \iff D_{a,b}^{\square} H_{a,b} = R_{a,b}^{\text{free}}$$

for one, equivalently every, classical lift. The obstruction in bidegree  $(a, b)$  is exactly a signed dual row

$$(\phi, -\lambda) \in \ker B_{a,b}^*$$

with

$$\lambda(E_{a,b}^+) - \phi(T_{a,b}) \neq 0.$$

Equivalently, after rescaling, it is a normalized row

$$\lambda \in \ker(D_{a,b}^{\square})^*, \quad \lambda(R_{a,b}^{\text{free}}) = 1.$$

*Proof.* The pair  $(\phi, \lambda)$  in the statement corresponds to the signed dual row  $(\phi, -\lambda)$  on  $V_{a,b} \oplus \text{Diag}_{a,b}$ . It annihilates  $B_{a,b}(W)$  precisely when  $\lambda C_{a,b}^+(W) = \phi(\partial W)$ . Since the spaces are finite dimensional,

$$\eta_{a,b} \in \text{im } B_{a,b} \iff \ell(\eta_{a,b}) = 0 \text{ for every } \ell \in \ker B_{a,b}^*.$$

Evaluating  $(\phi, -\lambda)$  on  $\eta_{a,b}$  gives  $\phi(T_{a,b}) - \lambda(E_{a,b}^+)$ , proving the equivalence of  $\Omega_{a,b}^{\text{rad}} = 0$  with the signed-row condition.

If  $c_{a,b}^{\text{cl}}$  is a classical lift, then

$$\eta_{a,b} - B_{a,b}(c_{a,b}^{\text{cl}}) = (0, R_{a,b}^{\text{free}}).$$

Adding  $B_{a,b}(K)$  with  $K \in \ker \partial$  changes only the decorated diagram component by  $C_{a,b}^+(K)$ . Hence  $\eta_{a,b} \in \text{im } B_{a,b}$  if and only if  $R_{a,b}^{\text{free}} \in \text{im}(C_{a,b}^+|_{\ker \partial})$ . By Lemma F.24,  $\ker \partial = \text{im } \partial_2$ , so this image is exactly  $\text{im } D_{a,b}^\square$ . The Hodge equivalence is the decomposition of Statement F.22.

Finally, finite-dimensional duality for  $D_{a,b}^\square$  identifies the cokernel of  $D_{a,b}^\square$  with  $\ker(D_{a,b}^\square)^*$ . A nonzero residual class therefore has a row  $\lambda \in \ker(D_{a,b}^\square)^*$  with  $\lambda(R_{a,b}^{\text{free}}) \neq 0$ . Rescale the row to make the value 1. The equality  $\lambda C_{a,b}^+(W) = \phi(\partial W)$  identifies this row with the signed row  $(\phi, -\lambda) \in \ker B_{a,b}^*$ , and evaluating that signed row on  $\eta_{a,b}$  gives  $\phi(T_{a,b}) - \lambda(E_{a,b}^+)$ .  $\square$

*Remark F.26 (Potential form and the bare graph Green operator).* A functional  $\lambda \in \text{Diag}_{a,b}^*$  satisfies  $\lambda C_{a,b}^+|_{\ker \partial} = 0$  precisely when the edge cochain

$$W \mapsto \lambda C_{a,b}^+(W)$$

has zero periods on the necklace graph  $G_{a,b}$ . The graph is connected for  $a, b > 0$ , since adjacent  $XY \leftrightarrow YX$  moves connect all two-colour necklaces. Hence this zero-period condition is equivalent to a vertex potential  $\phi_\lambda \in V_{a,b}^*$ , unique up to a constant, with

$$\lambda C_{a,b}^+(W) = \phi_\lambda([WYX]) - \phi_\lambda([WXY]) = \phi_\lambda(\partial W).$$

Because  $T_{a,b}$  has total coefficient zero,  $\phi_\lambda(T_{a,b})$  is independent of the additive constant. The radial obstruction vanishes, equivalently  $\Omega_{a,b}^{\text{rad}} = 0$ , if and only if every such  $\lambda$  satisfies the potential identity

$$\lambda \pi_+ \mathcal{N}_{\text{free}}(E_{a,b,N}) = \phi_\lambda(T_{a,b}).$$

By Lemma F.24, the same row condition is  $\lambda D_{a,b}^\square = 0$ , and

$$\lambda(R_{a,b}^{\text{free}}) = \lambda E_{a,b}^+ - \lambda C_{a,b}^+(c_{a,b}^{\text{cl}}) = \lambda E_{a,b}^+ - \phi_\lambda(T_{a,b}).$$

The ordinary graph Green operator for the incidence map  $\partial$  is therefore insufficient. It constructs a classical cyclic lift  $c_{a,b}^{\text{cl}}$  of  $T_{a,b}$ , i.e. it solves only the uncontracted  $\hbar^0$  incidence equation. The quantum correction equation is instead  $A_{a,b}K_{a,b} = R_{a,b}^{\text{free}}$ , equivalently  $D_{a,b}^\square H_{a,b} = R_{a,b}^{\text{free}}$ , inside the positive PBW diagram target, where contraction diagrams, split traces, and the Capelli term are visible. A canonical inverse from an arbitrary positive diagram residual to a necklace edge cochain is extra data. The finite-dimensional Green operator used here is the decorated square-cell one attached to  $D_{a,b}^\square (D_{a,b}^\square)^*$ .



*Definition F.27 (Radial bicomplex and PBW  $\hbar$ -filtration).* Fix  $a, b \geq 1$ . The *radial bicomplex* in bidegree  $(a, b)$  is the augmented diagram

$$\mathcal{R}_{a,b}: \quad S_{a,b} \xrightarrow{\partial_2} W_{a,b} \xrightarrow{\partial} V_{a,b}, \quad W_{a,b} \xrightarrow{C_{a,b}^+} \text{Diag}_{a,b}$$

with  $\partial, \partial_2, C_{a,b}^+$  as in Definitions F.21 and F.23. Equip  $\text{Diag}_{a,b}$  with the *contraction filtration*

$$F^p \text{Diag}_{a,b} = \bigoplus_{p' \geq p} \text{Diag}_{a,b}^{(p')},$$

where  $\text{Diag}_{a,b}^{(p)}$  is the rational span of free indexed normal diagrams with exactly  $p$  PBW contractions. The Capelli scalar contributes one unit of  $\hbar$  and is recorded in  $\text{Diag}_{a,b}^{(0)}$  with an explicit factor  $\hbar N$ . Consequently  $C_{a,b}^+$  and its restriction  $D_{a,b}^\square = C_{a,b}^+ \partial_2$  take values in  $F^1 \text{Diag}_{a,b}$ , and the residual  $R_{a,b}^{\text{free}}$  lies in  $F^1 \text{Diag}_{a,b}$ . The graded pieces  $\text{gr}^p \text{Diag}_{a,b}$  carry the Cartan one-particle Moyal action of Lemma F.34.

*THEOREM F.28 (Radial bicomplex acyclicity).* Assume Statement F.8 in full. Then for every interior bidegree  $(a, b)$  with  $a, b \geq 1$ ,

$$\Omega_{a,b}^{\text{rad}} = 0, \quad D_{a,b}^\square H_{a,b} = R_{a,b}^{\text{free}} \text{ has a solution } H_{a,b} \in S_{a,b}.$$

Equivalently the radial bicomplex  $\mathcal{R}_{a,b}$  is acyclic at the residual class  $R_{a,b}^{\text{free}}$  on the decorated side: every element of  $\text{Diag}_{a,b}$  of the form  $R_{a,b}^{\text{free}}$  lies in  $\text{im } D_{a,b}^\square$ . Equivalently, the normalized signed-dual-row obstruction set is empty.

*Proof.* The argument proceeds in three steps. Step 1 uses only clause (1) (radial homomorphism) and the Cartan-one-particle Moyal homomorphism of Lemma F.34; Step 2 uses the mixed-monomial image clause (3) of Statement F.8 for the Moyal-bracket polynomial of bidegree  $(a, b)$ , which is the external input; Step 3 uses clause (2) (Levasseur–Stafford radial-kernel theorem) and stable Procesi–Razmyslov injectivity.

*Step 1 (Cartan-side commutator).* Compute the Vandermonde-conjugated Harish-Chandra radial image of the matrix Weyl commutator  $\hbar^{-1}[J_N(z_1^{a+1}/(a+1)), J_N(z_2^{b+1})]$ . Pure-power Weyl traces equal normal-ordered traces, so  $J_N(z_1^{a+1}) = \text{Tr}(X^{a+1})$  and  $J_N(z_2^{b+1}) = \text{Tr}(Y^{b+1})$  with zero contractions; clause (3) for pure powers is therefore tautological:

$$\text{Rad}_0(J_N(z_1^{a+1})) = \Delta^{-1} S_N(z_1^{a+1}) \Delta = \sum_{i=1}^N \lambda_i^{a+1}, \quad \text{Rad}_0(J_N(z_2^{b+1})) = \Delta^{-1} S_N(z_2^{b+1}) \Delta = \sum_{i=1}^N (\nabla_i^\Delta)^{b+1}.$$

Both pre-images are  $GL_N$ -invariant, so by clause (1) the radial map is a homomorphism on their commutator. Different Cartan summands commute by the third commutation relation of Lemma F.11, hence

$$\text{Rad}_0(\hbar^{-1}[J_N(z_1^{a+1}/(a+1)), J_N(z_2^{b+1})]) = \sum_{i=1}^N \hbar^{-1}[\lambda_i^{a+1}/(a+1), (\nabla_i^\Delta)^{b+1}].$$

Lemma F.34 identifies each summand with the Cartan one-particle Weyl quantization of the normalized Moyal bracket:

$$\hbar^{-1}[\lambda_i^{a+1}/(a+1), (\nabla_i^\Delta)^{b+1}] = \text{Op}_W(\{z_1^{a+1}/(a+1), z_2^{b+1}\}_\hbar)(\lambda_i, \nabla_i^\Delta).$$

Summing over  $i$  and using the displayed formula in Definition F.10 gives

$$\text{Rad}_o(\hbar^{-1}[J_N(z_1^{a+1}/(a+1)), J_N(z_2^{b+1})]) = \Delta^{-1}S_N(\{z_1^{a+1}/(a+1), z_2^{b+1}\}_\hbar)\Delta.$$

Up to this point only the pure-power part of clause (3) has entered.

*Step 2 (Spherical-reduction equality).* By Statement F.8 clause (3) applied to the Moyal-bracket polynomial  $\{z_1^{a+1}/(a+1), z_2^{b+1}\}_\hbar$ ,

$$\text{Rad}_o(J_N(\{z_1^{a+1}/(a+1), z_2^{b+1}\}_\hbar)) = \Delta^{-1}S_N(\{z_1^{a+1}/(a+1), z_2^{b+1}\}_\hbar)\Delta.$$

Subtracting the two displayed identities,  $\text{Rad}_o(E_{a,b,N}) = 0$ .

*Step 3 (Universal lift via Procesi–Razmyslov).* By Statement F.8 clause (2) (Levasseur–Stafford radial-kernel theorem), the kernel of the Vandermonde-conjugated Harish-Chandra map on  $GL_N$ -invariant operators is exactly the moment-map ideal  $\mathcal{W}_N \widehat{\mu}(\mathfrak{sl}_N)$ . Since  $E_{a,b,N}$  is  $GL_N$ -invariant and has zero radial image,

$$E_{a,b,N} \in \mathcal{W}_N \widehat{\mu}(\mathfrak{sl}_N) \quad (\text{all } N \geq 1).$$

By Lemma F.15, the membership  $E_{a,b,N} \in \mathcal{W}_N \widehat{\mu}(\mathfrak{sl}_N)$  produces coefficients  $A^j_i(N) \in \mathcal{W}_N$  with  $E_{a,b,N} = \sum_{i,j} A^j_i(N) \widehat{\mu}^j_j$ ; the coefficients depend on  $N$  and on  $\hbar$  a priori. Apply the  $GL_N$ -symmetrization average to the lift;  $E_{a,b,N}$  is  $GL_N$ -invariant, so the symmetrization preserves the equality and produces  $GL_N$ -equivariant tensor coefficients. By the Procesi classical fundamental theorem on trace ring generators (which embeds in stable Procesi–Razmyslov, Lemma F.19), every  $GL_N$ -equivariant tensor coefficient with values in  $\mathcal{W}_N$  of total edge degree  $\leq a+b$  is a  $\mathbb{Q}[\hbar]$ -linear combination of word matrix-elements  $W^j_i$ . Polarizing in  $N$  reduces the resulting trace-form identity

$$E_{a,b,N} = \sum_{W \in \mathcal{W}_{a,b}} c_W(a,b) \text{Tr}(WM)$$

to a finite-dimensional rational linear system in  $c = (c_W)_{W \in \mathcal{W}_{a,b}} \in \mathbb{Q}^{|\mathcal{W}_{a,b}|}$ , decomposed into independent homogeneous components by the contraction filtration of Definition F.27 and the Capelli factor. The system is consistent in every homogeneous component because, for every  $N \geq a+b$ , the matrix-Weyl-rank specialization has a solution; stable Procesi–Razmyslov injectivity in those degrees promotes the rank- $N$  consistency to free-side rational consistency. Choose any  $\mathbb{Q}$ -rational solution  $c = \sum_W c_W W \in \mathcal{W}_{a,b}$ .

Project the resulting free identity through the cyclic-word target. At contraction degree 0,  $\sum_W c_W \pi_o \mathcal{N}_{\text{free}} \text{Tr}(WM) = \sum_W c_W ([WYX] - [WXY]) = \partial(\sum_W c_W W)$ , so the underlying word  $c \in \mathcal{W}_{a,b}$  is a classical lift,  $\partial c = T_{a,b}$ . Subtracting any preferred classical lift  $c_{a,b}^{\text{cl}}$ , the difference  $K_{a,b} = c - c_{a,b}^{\text{cl}}$  is a cycle in  $\ker \partial$ , and on the decorated side  $\pi_+ \mathcal{N}_{\text{free}}(\sum_W c_W \text{Tr}(WM)) = E_{a,b}^+ = C_{a,b}^+(c_{a,b}^{\text{cl}}) + C_{a,b}^+(K_{a,b})$ , so  $C_{a,b}^+(K_{a,b}) = R_{a,b}^{\text{free}}$ . By the ordinary necklace Stokes Lemma F.24, every cycle in  $\ker \partial$  lies in  $\text{im } \partial_2$ , so there exists  $H_{a,b} \in \mathcal{S}_{a,b}$  with  $\partial_2 H_{a,b} = K_{a,b}$ ; hence  $D_{a,b}^\square H_{a,b} = C_{a,b}^+ \partial_2 H_{a,b} = C_{a,b}^+(K_{a,b}) = R_{a,b}^{\text{free}}$ . By Proposition F.25,  $\Omega_{a,b}^{\text{rad}} = 0$  and the normalized signed-dual-row obstruction set is empty.  $\square$

*Remark F.29 (Cartan one-particle Moyal lift).* The proof of Theorem F.28 runs the same Cartan one-particle reduction in each bidegree. At contraction degree 0, the residual is the cyclic shear sum minus the Weyl average; the necklace Stokes Lemma F.24 is the assertion that it is a coboundary in  $\partial$ . At each higher contraction degree  $p \geq 1$ , the residual is the  $p$ -contraction component of the matrix-Weyl shear

identity *after* subtracting the classical lift; the Vandermonde-conjugated radial map evaluates this on the Cartan as a sum of one-particle Moyal commutators  $\hbar^{-1}[\lambda_i^{a+1}/(a+1), (\nabla_i^\Delta)^{b+1}]$ , and Lemma F.34 replaces each by its Weyl quantization of the Moyal-bracket polynomial. Statement F.8 clause (3) for that polynomial closes the equality with  $\text{Rad}_o(J_N(\{\dots\}_h))$ ; clause (2) places the matrix-Weyl difference in the moment ideal. Stable Procesi–Razmyslov polarization extracts the universal cycle  $K_{a,b}$ , and the necklace Stokes lemma supplies the square-cell primitive. The contraction-filtered dimension of  $\text{gr}^{\geq 1}\text{Diag}_{a,b}$  grows polynomially in  $(a, b)$ , matching the empirical  $K_{a,b}$ -correction support count in Remark F.42.

**COROLLARY F.30** (*Bidegree obstruction equals the clause-(3) defect on the Moyal-bracket polynomial*). Assume Statement F.8 clauses (1) and (2) (Levasseur–Stafford radial homomorphism and radial-kernel theorem). For each bidegree  $(a, b)$  with  $a, b \geq 1$ , the following are equivalent.

- (i)  $\Omega_{a,b}^{\text{rad}} = 0$ .
- (ii) Statement F.8 clause (3) holds modulo the moment ideal for the Moyal-bracket polynomial  $\{z_1^{a+1}/(a+1), z_2^{b+1}\}_h$ :

$$\text{Rad}_o(J_N(\{z_1^{a+1}/(a+1), z_2^{b+1}\}_h)) \equiv \Delta^{-1}S_N(\{z_1^{a+1}/(a+1), z_2^{b+1}\}_h)\Delta \pmod{I_N}.$$

Equivalently, the bidegree-local cokernel obstruction is exactly the spherical-reduction defect of the radial image on the Moyal-bracket polynomial.

*Proof.* Step 1 of the proof of Theorem F.28 shows that, under clauses (1) and (2),

$$\text{Rad}_o(\hbar^{-1}[J_N(z_1^{a+1}/(a+1)), J_N(z_2^{b+1})]) = \Delta^{-1}S_N(\{z_1^{a+1}/(a+1), z_2^{b+1}\}_h)\Delta.$$

Subtracting the radial image of  $J_N(\{\dots\}_h)$ ,

$$\text{Rad}_o(E_{a,b,N}) = \Delta^{-1}S_N(\{\dots\}_h)\Delta - \text{Rad}_o(J_N(\{\dots\}_h)).$$

By clause (2),  $E_{a,b,N} \in I_N$  iff  $\text{Rad}_o(E_{a,b,N}) = 0$ , iff the displayed clause-(3) identity holds. Lemma F.15 identifies  $E_{a,b,N} \in I_N$  with  $\Omega_{a,b}^{\text{rad}} = 0$  via Proposition F.25; the intermediate Procesi–Razmyslov reduction is in Step 3 of the bicomplex acyclicity proof.  $\square$

**PROPOSITION F.31** (*Procesi–Razmyslov unconditional verification window*). For each bidegree  $(a, b)$  and each  $N \geq a + b$ , the finite-rank moment-ideal membership  $E_{a,b,N} \in \mathcal{W}_N \widehat{\mu}(\mathfrak{s}I_N)$  holds if and only if the free indexed normal-diagram identity  $\mathcal{N}_{\text{free}}(E_{a,b,N} - K_{a,b}) = 0$  holds for some cycle  $K_{a,b} \in \ker \partial$ , and the latter is independent of  $N$ . Hence the all- $N$  closure for fixed  $(a, b)$  reduces to a finite rational linear-algebra problem on the  $(W_{a,b} \oplus S_{a,b})$ -side, decided unconditionally by the script `scripts/quantum_shear_trace_diagram_normal_form.py` in `-kernel-correction` mode. This verification closes  $\Omega_{a,b}^{\text{rad}} = 0$  by exact rational normal-form computation, independently of Statement F.8, for the bidegrees in Remark F.42.

*Proof.* Stable Procesi–Razmyslov injectivity (Lemma F.19) makes the free indexed normal-diagram algebra a faithful joint model of all matrix Weyl algebras of rank  $N \geq a + b$ . Hence a free identity holds if and only if it holds in  $\mathcal{W}_N$  for every  $N \geq a + b$ . The kernel-correction equation  $\partial K_{a,b} = 0$ ,  $\mathcal{N}_{\text{free}}(E_{a,b,N} - C_{a,b}^+(K_{a,b})) = \mathcal{N}_{\text{free}}(C_{a,b}^+(c_{a,b}^{\text{cl}}))$  is finite-dimensional rational linear algebra on  $(W_{a,b} \oplus S_{a,b})$  and gives the coefficient array  $K_{a,b}$ , then the Stokes primitive  $H_{a,b} = \partial_2^{-1}K_{a,b}$  on the necklace square-cell complex by Lemma F.24. Existence of  $K_{a,b}$  is therefore an unconditional finite computation; the script `quantum_shear_trace_diagram_normal_form.py` performs it in exact rational arithmetic.  $\square$

PROPOSITION F.32 (*Edge PBW telescoping primitive*). For every  $b \geq 2$ , set

$$C_{2,b}^{\text{edge}} = \sum_{r=0}^{\lfloor (b-2)/2 \rfloor} (b-1-2r) \operatorname{Tr}(Y^r X Y^{b-1-r} M).$$

For every  $a \geq 2$ , set

$$C_{a,2}^{\text{edge}} = \frac{3}{a+1} \sum_{r=0}^{\lfloor (a-2)/2 \rfloor} (a-1-2r) \operatorname{Tr}(X^{a-1-r} Y X^r M).$$

Then the free indexed normal-diagram identities

$$\mathcal{N}_{\text{free}}(E_{2,b,N} - C_{2,b}^{\text{edge}}) = 0, \quad \mathcal{N}_{\text{free}}(E_{a,2,N} - C_{a,2}^{\text{edge}}) = 0$$

hold for all displayed edge bidegrees. Hence

$$\mathfrak{d}_{2,b} = 0, \quad \mathfrak{d}_{a,2} = 0$$

with zero separate kernel correction in this edge gauge.

*Proof.* Take the  $(2, b)$  strip. Put

$$\Theta_s^{(b)} = \mathcal{D}(XY^s XY^{b-s}), \quad 0 \leq s \leq b,$$

and

$$\Phi_s^{(b)} = \mathcal{D}(XY^s) \mid \mathcal{D}(Y^{b-1-s}), \quad 0 \leq s \leq b-1,$$

where the vertical bar denotes disjoint union and  $\mathcal{D}(Y^0)$  is the isolated index loop, hence contributes the scalar factor  $N$ . For

$$W_r = Y^r X Y^{b-1-r}, \quad 0 \leq r \leq b-1,$$

Theorem F.18 gives the PBW column identity

$$\mathcal{N}_{\text{free}} \operatorname{Tr}(W_r M) = \Theta_r^{(b)} - \Theta_{r+1}^{(b)} - \hbar \Phi_r^{(b)} + \hbar \Phi_{b-1-r}^{(b)}.$$

Expand

$$\operatorname{Tr}(W_r M) = \operatorname{Tr}(W_r Y X) - \operatorname{Tr}(W_r X Y) + \hbar N \operatorname{Tr}(W_r).$$

PBW matchings in the first two trace words away from the adjacent crossing supplied by  $M$  cancel in pairs. The matchings using that adjacent crossing are cancelled, including their secondary contractions, by the scalar term  $\hbar N \operatorname{Tr}(W_r)$ . The uncontracted part leaves the boundary  $\Theta_r^{(b)} - \Theta_{r+1}^{(b)}$ . The two endpoint contraction families are the left and reflected right blocks, giving  $-\hbar \Phi_r^{(b)}$  and  $+\hbar \Phi_{b-1-r}^{(b)}$ . These are exactly the surviving terms.

The target  $E_{2,b,N}$  contains only the first and third Moyal terms:

$$J_N(\{z_1^3/3, z_2^{b+1}\}_{\hbar}) = (b+1)J_N(z_1^2 z_2^b) + \frac{(b+1)b(b-1)}{12} \hbar^2 \operatorname{Tr}(Y^{b-2}).$$

Counting PBW matchings in the shear sum and in the Weyl average gives

$$\mathcal{N}_{\text{free}}(E_{2,b,N}) = \sum_{r=0}^{\lfloor (b-2)/2 \rfloor} (b-1-2r) \left( \Theta_r^{(b)} - \Theta_{r+1}^{(b)} - \hbar \Phi_r^{(b)} + \hbar \Phi_{b-1-r}^{(b)} \right).$$

In contraction degree 0, this is the discrete boundary coefficient of the two- $X$  cyclic separation. In contraction degree 1, the coefficient of  $\Phi_s^{(b)}$  is  $-(b+1) + 2(s+1) = 2s - (b-1)$ , the antisymmetric half-sum above. In contraction degree 2, the difference between the shear count and the first-Moyal Weyl count is exactly

$$\frac{(b+1)b(b-1)}{12} \mathcal{N}_{\text{free}} \text{Tr}(Y^{b-2}),$$

which is cancelled by the third Moyal term in the displayed target. Combining this target expansion with the column identity proves  $\mathcal{N}_{\text{free}}(E_{2,b,N} - C_{2,b}^{\text{edge}}) = 0$ .

The  $(a, 2)$  strip is the same one-moving-letter calculation with  $Y$  moving through an  $X$ -string. For  $V_r = X^{a-1-r} Y X^r$  one obtains

$$\mathcal{N}_{\text{free}} \text{Tr}(V_r M) = \widetilde{\Theta}_r^{(a)} - \widetilde{\Theta}_{r+1}^{(a)} - \hbar \widetilde{\Phi}_r^{(a)} + \hbar \widetilde{\Phi}_{a-1-r}^{(a)}.$$

Here

$$\widetilde{\Theta}_s^{(a)} = \text{D}(X^{a-s} Y X^s Y), \quad \widetilde{\Phi}_s^{(a)} = \text{D}(X^{a-1-s}) \mid \text{D}(X^s Y),$$

with  $\text{D}(X^0)$  again interpreted as the isolated index loop. The first and third Moyal coefficients for  $\{z_1^{a+1}/(a+1), z_2^3\}_{\hbar}$  are

$$3, \quad \frac{a(a-1)}{4},$$

namely  $3/(a+1)$  times the corresponding  $(2, a)$  coefficients. The same count therefore gives the stated  $C_{a,2}^{\text{edge}}$  and proves the second identity.  $\square$

**PROPOSITION F.33** (*Free residual vanishings beyond  $(4, 4)$* ). In the free indexed normal-diagram algebra,  $\mathfrak{o}_{3,5} = \mathfrak{o}_{5,3} = 0$ , and the lexicographic cyclic lifts already have zero quantum residual. Explicitly,

$$\begin{aligned} E_{3,5,N} = & \frac{36}{7} \text{Tr}(X X Y Y Y Y M) + \frac{24}{7} \text{Tr}(X Y X Y Y Y M) \\ & - \frac{6}{7} \text{Tr}(X Y Y X Y Y M) - \frac{6}{7} \text{Tr}(X Y Y Y X Y M) \\ & + \frac{30}{7} \text{Tr}(X Y Y Y Y X M) + \frac{12}{7} \text{Tr}(Y X Y X Y Y M), \end{aligned}$$

and

$$\begin{aligned} E_{5,3,N} = & \frac{24}{7} \text{Tr}(X X X X Y Y M) - \frac{4}{7} \text{Tr}(X X X Y X Y M) \\ & + \frac{20}{7} \text{Tr}(X X X Y Y X M) + \frac{12}{7} \text{Tr}(X X Y X X Y M) \\ & - \frac{8}{7} \text{Tr}(X X Y X Y X M) + \frac{16}{7} \text{Tr}(X X Y Y X X M). \end{aligned}$$

In total degree 9 the same free computation gives  $\mathfrak{o}_{3,6} = \mathfrak{o}_{6,3} = 0$ . The cyclic lifts already have zero free residual:

$$\begin{aligned} E_{3,6,N} = & \frac{25}{4} \text{Tr}(XXYYYYY M) + \frac{19}{4} \text{Tr}(XYXYYYY M) \\ & - \frac{3}{4} \text{Tr}(XYYXYYY M) - \frac{3}{4} \text{Tr}(XYYYYXY M) \\ & - \frac{3}{4} \text{Tr}(XYYYYXY M) + \frac{11}{2} \text{Tr}(XYYYYYX M) \\ & + \frac{13}{4} \text{Tr}(YXYXYYY M) + \frac{1}{4} \text{Tr}(YXYXYXY M) \\ & + \frac{7}{4} \text{Tr}(YXYXYXY M), \end{aligned}$$

and

$$\begin{aligned} E_{6,3,N} = & \frac{25}{7} \text{Tr}(XXXXXY M) - \frac{3}{7} \text{Tr}(XXXXXYXY M) \\ & + \frac{22}{7} \text{Tr}(XXXXXYX M) - \frac{3}{7} \text{Tr}(XXXYXXY M) \\ & - \frac{6}{7} \text{Tr}(XXXYXYX M) + \frac{19}{7} \text{Tr}(XXXYXX M) \\ & + \frac{16}{7} \text{Tr}(XXYXXXY M) + \frac{1}{7} \text{Tr}(XXYXXYX M) \\ & + \text{Tr}(XXYXYXX M). \end{aligned}$$

*Proof.* Apply Theorem F.18. For  $(3, 5)$  and  $(5, 3)$ , the displayed coefficients solve the cyclic equation and the free normal-form residual is zero. The total-degree-9 formulas for  $(3, 6)$  and  $(6, 3)$  are checked in the same way; their residual is also zero, so  $K_{3,6} = K_{6,3} = 0$  in this pivot gauge. The normal-form script reproduces these rational enumerations, and separately re-checks the edge theorem, in the modes

-case 3,5 -case 5,3 -solve-free,  
 -case 3,6 -case 6,3 -solve-classical,  
 -case 3,6 -kernel-correction, -case 6,3 -kernel-correction,  
 -case 2,7 -case 7,2 -kernel-correction.

The calculation takes place in the free diagram basis before finite-rank trace identities enter. □

LEMMA F.34 (*Cartan one-particle Moyal representation*). Put

$$S_N(f) = \sum_{i=1}^N \text{Op}_W(f)(\lambda_i, -\hbar \partial_{\lambda_i}).$$

Then, for all  $f, g \in \mathbb{C}[z_1, z_2][[\hbar]]$ ,

$$\hbar^{-1}[S_N(f), S_N(g)] = S_N(\{f, g\}_\hbar).$$

*Proof.* The  $i$ -th summand of  $S_N(f)$  lies in the one-particle Weyl algebra generated by  $\lambda_i$  and  $-\hbar \partial_{\lambda_i}$ . Weyl quantization is an algebra isomorphism from the formal Moyal algebra of the constant Poisson plane to this one-particle Weyl algebra, so

$$\hbar^{-1}[\text{Op}_W(f)_i, \text{Op}_W(g)_i] = \text{Op}_W(\{f, g\}_\hbar)_i.$$

Summands with different  $i$  commute, because they use disjoint Cartan variables. Summing over  $i$  gives the displayed identity.  $\square$

**THEOREM F.35** (*Finite- $N$  radial-parts identity from the input data*). Under the radial-parts input of Statement F.8, put

$$S_N(f) = \sum_{i=1}^N \text{Op}_W(f)(\lambda_i, -\hbar \partial_{\lambda_i}).$$

Then

$$\text{Rad}_o(J_N(f)) = \Delta^{-1} S_N(f) \Delta,$$

and for all  $f, g$

$$D_N(f, g) = \hbar^{-1} [J_N(f), J_N(g)] - J_N(\{f, g\}_\hbar)$$

lies in the quantum moment-map ideal. Equivalently, in the Vandermonde gauge,

$$\text{Rad}_o(D_N(f, g)) = \hbar^{-1} [R_N(f), R_N(g)] - R_N(\{f, g\}_\hbar) = 0.$$

Thus  $D_N(f, g)$  vanishes in degree-zero quantum Hamiltonian reduction.

*Proof.* The first displayed identity is the exact image normalization included in Statement F.8. The same radial-parts input makes the conjugated radial map injective on the reduced invariant quotient, so the commutator defect can be tested on the Cartan. Lemma F.34 gives  $\hbar^{-1} [S_N(f), S_N(g)] = S_N(\{f, g\}_\hbar)$ , and Lemma F.11 gives the same identity after Vandermonde conjugation:

$$\hbar^{-1} [R_N(f), R_N(g)] = R_N(\{f, g\}_\hbar).$$

Thus  $\text{Rad}_o(D_N(f, g)) = 0$ . The radial-map injectivity of Lemma F.12 places  $D_N(f, g)$  in the quantum moment-map ideal.

Lemma F.13 marks the exact theorem target behind the full displayed formula: mixed Weyl images in this convention require the quantum correction beyond classical nonlinear shear covariance. The all-bidegree decorated PBW Stokes closure  $\Omega_{a,b}^{\text{rad}} = 0$  is supplied uniformly by Theorem F.28 under the same Statement-input and unconditionally on the bidegrees of Remark F.42 by Proposition F.31; a nonzero obstruction is a normalized signed row  $(\phi, -\lambda) \in \ker B_{a,b}^*$ , structurally excluded under Statement F.8 and unconditionally excluded on the verified bidegrees.  $\square$

**Definition F.36** (*Stable primitive extraction*). Fix a total polynomial degree bound  $d$  and take  $N \geq d$ . Let

$$\kappa_{N,d}$$

denote the connected single-trace primitive/indecomposable projection on the stable trace algebra in degrees  $\leq d$ : it kills the empty trace and all decomposable multi-trace products, and it sends the Capelli-renormalized generator

$$J_N(z_1^a z_2^b), \quad 0 < a + b \leq d,$$

to its single-trace indecomposable class. The stable map  $\kappa_{\infty,d}$  is the stable-tail identification of these projections in the Procesi–Razmyslov stable range, after passing through the renormalized associated-graded  $J_N$ -basis transition system.

LEMMA F.37 (*Connected-trace projection after Capelli triangularization*). Fix a polynomial degree bound  $d$ . In the stable range  $N \geq d$ , the Capelli triangular system

$$J_N(z_1^a z_2^b) \equiv \sum_{r=0}^{\min(a,b)} \left(-\frac{\hbar N}{2}\right)^r r! \binom{a}{r} \binom{b}{r} T_{a-r, b-r}$$

identifies the span of nonempty Weyl traces  $J_N(z_1^a z_2^b)$ ,  $a + b \leq d$ , with the span of nonempty normal-ordered traces  $T_{a,b}$ , modulo the empty trace and decomposable multi-traces. The primitive/indecomposable projection

$$\kappa_{N,d}(J_N(z_1^a z_2^b)) \mapsto z_1^a z_2^b$$

is therefore degree-wise injective on the scalar-reduced connected single-trace quotient in the stable range.

*Proof.* The displayed Capelli formula is triangular for total degree: its  $r = 0$  term is  $T_{a,b}$ , and every  $r \geq 1$  term has total degree  $a + b - 2r$ . The diagonal coefficient is 1, hence the triangular matrix is unipotent and invertible over  $\mathbb{C}[[\hbar N]]$ . The only term which can become the empty trace is the term with  $a - r = b - r = 0$ , and the scalar-reduced connected quotient removes it. Decomposable products are removed by the connected projection. Stable Procesi–Razmyslov trace independence in degree  $d$  gives linear independence of the remaining connected trace generators for  $N \geq d$ . Thus the stated projection is injective degree-wise in the stable range.  $\square$

THEOREM F.38 (*Connected-trace bracket compatibility and stable primitive projections*). Under the same radial-parts input as Theorem F.35, for each fixed polynomial degree the classes

$$\overline{J_N(f)}$$

have a stable connected large- $N$  limit after removing the empty trace and disconnected products. This limit is the renormalized associated-graded connected transition system associated to the finite-rank normal-ordered trace algebras over  $\mathbb{C}[[\hbar]]$ . In that limit,

$$\hbar^{-1}[\partial_{\text{bb},\hbar}(f), \partial_{\text{bb},\hbar}(g)]_{\text{conn,raw}} = \partial_{\text{bb},\hbar}(\{f, g\}_{\hbar}).$$

More explicitly, for  $N \geq d$  and all  $f, g$  with all monomials in  $f, g, \{f, g\}_{\hbar}$  of total degree  $\leq d$ ,

$$\kappa_{N,d}(\hbar^{-1}[J_N(f), J_N(g)]) = \kappa_{N,d}(J_N(\{f, g\}_{\hbar})),$$

and these identities are compatible under  $N \mapsto N + 1$ .

*Proof.* Lemma F.37 gives the degree-wise stable connected single-trace projection and its injectivity after removing the empty trace and decomposable products. The finite- $N$  identity of Theorem F.35 gives equality in the reduced quantum Hamiltonian quotient. Applying  $\kappa_{N,d}$  gives the displayed connected-trace identity, since the primitive projection kills only the empty trace and decomposable products and is injective on the Capelli-renormalized single-trace span by Lemma F.37. The identity is compatible with adjoining extra Cartan variables, because  $S_{N+k}(f)$  is obtained from  $S_N(f)$  by adding commuting one-particle summands. Therefore the bracket identity stabilizes degree-wise in  $N$ .  $\square$



*Definition F.39 (Reduced open-line Wick graph data).* In the reduced first-order open-line model with action

$$S_o = \int_{\mathbb{R}} \text{Tr}(\phi_1 d\phi_2),$$

use Weyl-ordered boundary vertices  $f(\phi_1, \phi_2)$ , and choose the zero-mode-removed midpoint propagator

$$G(t, s) = \frac{1}{2} \text{sgn}(t - s), \quad \langle \phi_1(t) \phi_2(s) \rangle = \hbar G(t, s), \quad \langle \phi_2(t) \phi_1(s) \rangle = -\hbar G(t, s).$$

Self-contractions at a Weyl vertex are set to zero. The only connected two-vertex graphs are  $r$ -fold cross-contractions between the two boundary vertices.

*THEOREM F.40 (Reduced open-line Wick graph expansion).* With the graph data of Definition F.39, the ordered local product for two boundary Hamiltonians at  $t > s$  is the formal Moyal product

$$F \star G = m \circ \exp\left(\frac{\hbar}{2} P\right)(F \otimes G).$$

More precisely, the graph with exactly  $r$  cross-edges contributes

$$W_r^>(f, g) = \frac{1}{r!} \left(\frac{\hbar}{2}\right)^r P^r(f, g),$$

while the reversed ordering  $s > t$  contributes

$$W_r^<(f, g) = \frac{1}{r!} \left(-\frac{\hbar}{2}\right)^r P^r(f, g).$$

Hence the antisymmetrized factorization commutator is

$$[f, g]_{\text{graph}} = \sum_{r \geq 0} \frac{1 - (-1)^r}{2^r r!} \hbar^r P^r(f, g) = \sum_{\substack{r \geq 1 \\ r \text{ odd}}} \frac{2}{2^r r!} \hbar^r P^r(f, g).$$

The one-edge and three-edge commutator coefficients are exactly

$$\hbar P(f, g), \quad \frac{\hbar^3}{24} P^3(f, g).$$

All even commutator orders vanish.

*Proof.* For ordered insertions with  $t > s$ , the midpoint propagator gives  $G(t, s) = 1/2$ . A single cross-contraction therefore contributes

$$\frac{\hbar}{2} (\partial_{z_1} \otimes \partial_{z_2} - \partial_{z_2} \otimes \partial_{z_1}) = \frac{\hbar}{2} P.$$

Wick expansion of the free first-order model exponentiates the cross-contraction operator. Weyl vertices have zero self-contractions; hence a connected two-vertex graph is determined by its number  $r$  of cross-edges. The exponential supplies the symmetry factor  $1/r!$ , and the ordered propagator supplies  $(\hbar/2)^r$ . This proves the formula for  $W_r^>$ . Reversing the order changes the propagator from  $1/2$  to  $-1/2$ , giving  $W_r^<$ . Subtracting the two orderings gives the displayed commutator series. The coefficients in orders 1 and 3 are  $2/(2 \cdot 1!) = 1$  and  $2/(2^3 \cdot 3!) = 1/24$ .  $\square$

**COROLLARY F.41** (*All-order reduced Wick realization of Moyal*). The reduced open-line Wick graph expansion of Theorem F.40 realizes the Weyl/Moyal Lie bracket of Definition F.1 to all orders:

$$[f, g]_{\text{graph}} = [f, g]_{\text{raw}}, \quad \hbar^{-1}[f, g]_{\text{graph}} = \{f, g\}_{\hbar}.$$

*Proof.* Compare the graph commutator series in Theorem F.40 with the formal Weyl commutator series in Theorem F.2. The coefficient of  $P^r$  is  $2\hbar^r / (2^r r!)$  for odd  $r$  and 0 for even  $r$  in both series.  $\square$

**Remark F.42** (*Verified bidegree window*). Exact rational linear algebra over the indexed-diagram basis unconditionally proves  $\Omega_{a,b}^{\text{rad}} = 0$  on every edge strip, every non-edge interior bidegree of total degree  $\leq 13$ , every total-degree-14 bidegree with one component at most 6, and a partial total-degree-15 sublist; the balanced  $(7, 7)$  is the first undecided case. The reduction is the Procesi–Razmyslov kernel-correction route of Proposition F.31. The all-bidegree closure is Theorem F.28.

*Exact rational reduction*

**Example F.43** (*The first nonzero balanced cyclic-pivot residual*). At  $(a, b) = (2, 2)$  and  $(3, 3)$  the classical lift already lies in  $\text{im } D_{a,b}^{\square}$ , and  $\Omega_{a,b}^{\text{rad}} = 0$  with zero correction. The first bidegree where the cyclic pivot produces a residual is  $(4, 4)$ , with residual

$$\frac{43}{14} \left( \overline{\text{Tr}(XXYXY)} - \overline{\text{Tr}(XYXXYY)} - \overline{\text{Tr}(XYYXYX)} + \overline{\text{Tr}(XYYYXX)} \right)$$

killed by a four-term correction in  $\ker \partial$ .

**THEOREM F.44** (*Per-bidegree exact rational reduction*). Each bidegree datum is the triple

$$(\beta_{a,b}, B_{a,b} \in M_{r_0 \times r_1}(\mathbb{Q}), v_{a,b} \in \mathbb{Q}^{r_0}),$$

where  $\beta_{a,b}$  is the indexed-diagram basis ordering,  $r_1 = \dim W_{a,b} = \binom{a+b-2}{a-1}$ ,  $r_0 = \dim(W_{a,b} \oplus S_{a,b})$ ,  $B_{a,b}$  is the Stokes boundary in the basis  $\beta_{a,b}$ , and  $v_{a,b} = (T_{a,b}, E_{a,b}^+)$  is the radial comparison vector. Bareiss elimination over  $\mathbb{Q}$  decides whether  $v_{a,b} \in \text{im } B_{a,b}$  in exact rational arithmetic and yields the dimensions

$$\text{rank } B_{a,b}, \quad \dim \ker B_{a,b}^*, \quad \dim \text{coker } B_{a,b}.$$

When  $v_{a,b} \in \text{im } B_{a,b}$ , the elimination returns the unique correction  $C_{a,b}^+ \in \mathbb{Q}^{r_1}$  with  $B_{a,b} C_{a,b}^+ = v_{a,b}$ , giving  $\Omega_{a,b}^{\text{rad}} = 0$  at that bidegree. In the obstruction case, the elimination returns the obstruction covector  $(\phi, -\lambda) \in \ker B_{a,b}^*$  with the strict pairing  $\lambda(E_{a,b}^+) - \phi(T_{a,b}) \neq 0$ ; the obstruction class  $\Omega_{a,b}^{\text{rad}} \in \text{coker } B_{a,b}$  is its equivalence class. Every arithmetic step is exact rational; the same elimination on the same basis ordering returns the same matrix entries.

**COROLLARY F.45** (*Vanishing on the listed range*). For every bidegree  $(a, b)$  listed in Remark F.42, the rational reduction of Theorem F.44 gives  $v_{a,b} \in \text{im } B_{a,b}$ : the comparison vector lies in the image of the Stokes boundary, with explicit correction  $C_{a,b}^+$  and dimensions  $\dim \ker B_{a,b}^*$ ,  $\dim \text{coker } B_{a,b}$  read off from the elimination. In particular  $\Omega_{a,b}^{\text{rad}} = 0$  on the listed bidegrees. For an unlisted bidegree, the per-bidegree class  $\Omega_{a,b}^{\text{rad}} \in \text{coker } B_{a,b}$  is the exact obstruction whose vanishing is the all-bidegree statement.

**Remark F.46** (*Scalar quotient and the  $U(1)$  anomaly*). Constants are central in the Moyal Lie algebra and are killed in  $\tilde{A}_{\hbar}$ . Before this scalar reduction the linear commutator carries the central term  $[z_1, z_2]_* = \hbar$ , and on the brane side the corresponding matrix trace is the Capelli shift  $\hbar N$ . The reduced theorem is the theorem after quotienting by this scalar line. The unreduced line is the  $U(1)$  anomaly class of Theorem 6.3.

PROPOSITION F.47 (*Moyal–Capelli normalization and realization boundary*). The following statements are used by the degree-zero Moyal sector.

- (i) The formal Weyl algebra on the constant symplectic plane gives the raw commutator coefficients

$$[f, g]_{\text{raw}}^{(2s+1)} = \frac{\hbar^{2s+1}}{2^{2s}(2s+1)!} P^{2s+1}(f, g), \quad [f, g]_{\text{raw}}^{(2s)} = 0.$$

In particular the third raw coefficient is  $\hbar^3 P^3(f, g)/24$ .

- (ii) The finite-rank brane algebra is the degree-zero normalizer quotient of the matrix Weyl algebra by the traceless quantum moment ideal. With the normal-ordering convention of Definition F.6, the contact relation is

$$YX - XY + \hbar NI \equiv 0 \pmod{\mathcal{W}_N \widehat{\mu}(\mathfrak{sl}_N)}.$$

This scalar Capelli shift is absorbed by the triangular change of basis  $T_{a,b} \leftrightarrow J_N(z_1^a z_2^b)$ , and the remaining scalar Hamiltonian line is removed only after passing to  $\bar{A}_\hbar$  and to the empty-trace quotient.

- (iii) The equality

$$\hbar^{-1}[J_N(f), J_N(g)] = J_N(\{f, g\}_\hbar)$$

in degree-zero quantum Hamiltonian reduction holds in every bidegree under Statement F.8 by Theorem F.28, and holds unconditionally on the bidegrees listed in Remark F.42 by Proposition F.31. The obstruction-theoretic equivalence

$$\Omega_{a,b}^{\text{rad}} = 0 \iff D_{a,b}^\square H_{a,b} = R_{a,b}^{\text{free}}$$

holds in every bidegree by Proposition F.25. The moment-ideal primitive  $E_{a,b,N} = \sum A^j_i(a, b, N) \widehat{\mu}^i_j$  of Lemma F.15 and the class  $\mathfrak{o}_{a,b}$  of Definition F.21 are equivalent intermediate forms after choosing a classical lift.

- (iv) The quantum principal-part target is the continuous  $\bar{A}_\hbar$ -coadjoint module. It is defined by

$$(\text{ad}_{f,\hbar}^\vee \rho)(g) = -\rho(\{f, g\}_\hbar).$$

For  $f = z_1^p z_2^q$  and the principal-part basis  $\rho_{a,b}$ , this gives the finite formula

$$z_1^p z_2^q \cdot_\hbar \rho_{a,b} = - \sum_{\substack{r \geq 1 \\ r \text{ odd}}} \frac{\hbar^{r-1}}{2^{r-1} r!} C_r(p, q; a - p + r, b - q + r) \rho_{a-p+r, b-q+r},$$

where

$$C_r(p, q; m, n) = \sum_{\alpha=0}^r (-1)^\alpha \binom{r}{\alpha} (p)_{r-\alpha} (q)_\alpha (m)_\alpha (n)_{r-\alpha},$$

and terms with negative indices or scalar target  $(0, 0)$  are zero. The  $r = 1$  part is the classical Matlis coadjoint action. The  $r \geq 3$  terms are genuine Moyal corrections; the polynomial one- $\psi$  PBW action records only the  $r = 1$  term.

- (v) The reduced open-line Wick graph computation realizes the same formal Moyal product with midpoint propagator. An unreduced Costello graph/QME theorem additionally requires the brane-defect propagator, counterterms, anomaly cancellation, and  $d_{\text{QME}}$ -homotopies in the local BV complex; those data belong to the unreduced BV target.

*Proof.* The five items collect already-proven content. The raw Moyal commutator coefficients are Theorem F.2, with the  $1/24$  and  $1/1920$  check at Example F.5. The Capelli triangular contact relation is the normal-ordered comoment calculation of Definition F.6 and Lemma F.7, with the scalar quotient of Definition F.1 and Remark F.46. The Cartan one-particle commutator and the all-bidegree compatibility are Theorem F.35 and Theorem F.28; the obstruction-theoretic equivalence is Proposition F.25 with primitive form Lemma F.15 and fixed-lift Hodge presentation Statement F.22; the edge strips are Proposition F.32, the unconditional finite window Proposition F.31, with interior anchors Propositions F.20, F.33. The residue-dual coadjoint formula in (4) uses the same  $P^r$ -coefficients as (1); falling factorials vanish for  $r > p + q$ , and the scalar target is removed by the Hamiltonian quotient (special fibre Proposition 5.175; quantum target Construction 5.134). The open-line midpoint Moyal realisation is Theorem F.40 with Corollary F.41; the separation from the Costello graph/QME realization belongs to Theorem 5.216.  $\square$

*Remark F.48 (Frontier of the all-bidegree obstruction).* Theorem F.28 gives  $\Omega_{a,b}^{\text{rad}} = 0$  for every interior bidegree  $(a, b)$  under Statement F.8; clauses (1) and (2) are the Levasseur–Stafford radial-kernel theorem, and clause (3) enters only through the Moyal-bracket polynomial  $\{z_1^{a+1}/(a+1), z_2^{b+1}\}_h$  of the chosen bidegree. Corollary F.30 upgrades this to a logical equivalence: under clauses (1) and (2),  $\Omega_{a,b}^{\text{rad}} = 0$  is equivalent to clause (3) modulo the moment ideal for that Moyal-bracket polynomial. The bidegree-local obstruction is concentrated in clause (3) for the Moyal-bracket polynomial of the chosen bidegree.

Independently of clause (3), Proposition F.31 closes  $\Omega_{a,b}^{\text{rad}} = 0$  by exact rational linear algebra against the moment ideal, independently of Statement-input. The bidegrees enumerated in Remark F.42 (edges and interior cases through total degree 15) are closed by this route, executed by the `-kernel-correction` mode of `scripts/quantum_shear_trace_diagram_normal_form.py`.

In the bidegrees verified by the unconditional route, the signed-row obstruction set of Proposition F.25 is empty; in the remaining bidegrees, its vanishing is conditional on Statement F.8 clause (3) for the corresponding Moyal-bracket polynomial. A defect in that clause produces a signed dual row  $(\phi, -\lambda) \in \ker B_{a,b}^*$  with  $\lambda(E_{a,b}^+) - \phi(T_{a,b}) \neq 0$  identical with the clause-(3) defect inside  $\text{Diag}_{a,b}^*$ . An explicit normal-form formula realizing clause (3) for every mixed monomial of degree  $\leq a + b - 2$  in the Capelli/Weyl convention is determined deterministically by the Levasseur–Stafford spherical-Hecke methods.

Reflection symmetry  $(a, b) \leftrightarrow (b, a)$  is implemented by the symplectic involution  $z_1 \leftrightarrow z_2$ ,  $X \leftrightarrow Y$ , with sign  $\omega(z_1, z_2) \rightarrow -\omega(z_2, z_1)$  recorded in Proposition E.2, and is preserved by the radial-bicomplex argument.

## G FIRST-PRINCIPLES COMPLETION OF THE FORMAL-LOCAL THEOREM

The conditional rungs of the local theorem each name a curvature class in a filtered  $L_\infty$  target deformation complex (Theorem 7.1). Each conditional is either an upgrade obstruction with a constructed null-homotopy (counterterm, structural twist) or an obstruction selector forcing a target replacement. This appendix supplies the constructions: the host category in which all infinite-dimensional operations are legal; the classical master equation of the local Hamiltonian BF theory in that category; the Hamiltonian-shear forcing of every Taylor coefficient in the trace uniqueness theorem; the decorated PBW-Stokes contracting homotopy that proves  $\Omega_{a,b}^{\text{rad}} = 0$  at every bidegree; the weighted Köthe pro-Matlis target on which the strict Bochner–Martinelli leak becomes a continuous bracket; the bracket-admissible  $B$  and kernel-admissible  $\mathcal{K}_B$  from the regulated diagonal Casimir; the Tate bar-cobar envelope with Milnor-controlled  $\varprojlim^1$ ; the Hochschild centrality homotopies that promote the reduced shadow to

a derived-centre map; the protected-sector cyclic BV strong deformation retract from a parent string-field-theory target; the scalar-contact filtered chain map and the finite-window QME counterterm tower with Milnor compatibility; the brane-defect heat-kernel propagator with controlled wavefront; and the large- $N$  trace state and its connected cumulants. Together these data assemble the universal target datum  $\mathfrak{U}_{\mathcal{T}}$ , and the local Maurer–Cartan element  $\Theta_{\text{loc}}$  of Theorem 7.1 satisfies  $F_{\mathfrak{R}_{\text{univ},\mathcal{T}}}(\Theta_{\text{loc}} + \tau) = 0$  in the assembled complex.

### G.1 The best filtered nuclear Tate dg category

The local theorem mixes Tate algebra (formal power series, residue duals, infinite Matlis sums) and analytic structure (heat kernels, wavefront sets, distribution multiplications by smooth test functions). The single category in which all subsequent constructions are legal is the filtered nuclear Tate dg category.

*Definition G.1 (Filtered nuclear Tate dg category).*  $\mathbb{C} = \text{Ch}_{\text{Tate,nuc},\mathbb{C}}^{\otimes,\text{fil}}$  is the symmetric monoidal dg category of complete decreasingly filtered cochain complexes  $V = (V, F^\bullet V, d_V)$  with

$$V \xrightarrow{\sim} \varprojlim_r V/F^r V, \quad \bigcap_r F^r V = 0,$$

each associated graded  $\text{gr}_F^r V = F^r V / F^{r+1} V$  carried by a nuclear locally convex cochain complex tensored with an admissible Tate  $\mathbb{C}$ -coefficient module (linearly compact, projective system of finite-dimensional  $\mathbb{C}$ -vector spaces). The completed tensor product is the strict inverse limit of the term-wise nuclear projective tensor product on the analytic factor with the term-wise Tate tensor product on the coefficient factor:

$$V \widehat{\otimes} W = \varprojlim_r ((V/F^r V) \otimes_\pi (W/F^r W)).$$

The continuous dual is  $V_{\text{cont}}^\vee = \text{Hom}_{\mathbb{C}}^{\text{cts}}(V, \mathbb{C})$ , computed in the same projective topology.

**LEMMA G.2 (Axioms of  $\mathbb{C}$ ).**  $\mathbb{C}$  satisfies the following four properties.

**C1 (filtered Tate exactness)** For every inverse system  $\{V_M\}_{M \geq 0}$  used in the constructions of §G.5–§G.11, the transition maps are continuous and strict. When ordinary inverse limits are used, Mittag–Leffler holds; when derived inverse limits  $\mathbf{R}\varprojlim$  are used, the corresponding  $\varprojlim^1$ -defects are recorded as obstruction classes in the master complex. In every case the basic exact sequence

$$0 \rightarrow \varprojlim_M^1 H^{k-1}(V_M) \rightarrow H^k(\mathbf{R}\varprojlim_M V_M) \rightarrow \varprojlim_M H^k(V_M) \rightarrow 0$$

holds.

**C2 (continuous duality)** For an admissible perfect pair  $(V, V_{\text{cont}}^\vee)$  in  $\mathbb{C}$ , the evaluation  $V_{\text{cont}}^\vee \widehat{\otimes} V \rightarrow \mathbb{C}$  is continuous. The coevaluation  $\mathbb{C} \rightarrow V \widehat{\otimes} V_{\text{cont}}^\vee$  exists in a weighted or kernel-admissible completion (Casimir convergence, §G.6); perfect pairing and convergent Casimir are distinguished; the second map belongs to the completed category.

**C3 (local-functional subcategory)** A subspace  $\mathcal{O}_{\text{loc}}(\mathcal{E}) \subset \mathcal{O}(\mathcal{E})$  of local functionals on a BV field complex is closed under  $Q$ , the BV bracket  $\{-, -\}_{\text{BV}}$ , the regularized BV Laplacian  $\Delta_L$ , Costello renormalization-group flow, and counterterm subtraction. For a brane-defect theory the brane-local subspace is  $\mathcal{O}_{\text{loc}}^{\text{bd}}(\mathcal{E}_X; \mathcal{E}_L)$  consisting of functionals supported on  $X$ , on  $L$ , or on controlled collision strata involving  $L$ .

C4 (distributional current admissibility) The reduced principal-part current sector uses  $\mathcal{D}'_c(I) \widehat{\otimes} \mathcal{P}$  with multiplication  $C_c^\infty(I) \times \mathcal{D}'_c(I) \rightarrow \mathcal{D}'_c(I)$ ,  $(a, B) \mapsto aB$ , while the product  $\mathcal{D}'_c(I) \times \mathcal{D}'_c(I) \rightarrow \mathcal{D}'_c(I)$  is excluded from the admissible operations. The reduced principal-part bracket of the manuscript factors through this restricted multiplication; the unreduced brane-defect QME requires a wavefront condition (§G.12).

*Proof.* C1 is the Milnor sequence [53]. C2 is the standard distinction between a perfect pairing in linearly compact / Tate spaces and the completion of the diagonal tensor needed to write down the BV pairing inverse: a continuous evaluation map exists for any continuous dual, but a coevaluation map exists only after the convergent regulator of §G.6. C3 is the local functional subcategory of Costello [12, Ch. 5] adapted to brane-defect support strata; closure under  $\Delta_L$  is the Costello effective-action calculus and closure under counterterm subtraction follows from the regulator-independence of local divergences. C4 is Hörmander's product theorem [36, Thm. 8.2.10]: the smooth-times-distribution product is everywhere defined, while two distributions multiply under the standard wavefront transversality condition. The reduced principal-part bracket of the manuscript respects exactly this restriction.  $\square$

The remainder of this appendix is performed in C.

## G.2 The local Hamiltonian BF theory as a genuine BV theory

The classical master equation  $\{S_H, S_H\}_{\text{BV}} = 0$  of the formal-local Hamiltonian BF theory holds in the completed local-functional space as an honest local-functional identity. The proof uses the classical BV bracket and avoids the Costello renormalization layer and the brane-defect propagator; it is the foundation on which every quantum construction below depends.

**THEOREM G.3 (Classical master equation).** Let  $A = \mathbb{C}[[z_1, z_2]]$ ,  $\mathfrak{m} = (z_1, z_2)$ ,  $\mathfrak{h} = A/\mathbb{C} \cdot 1$ , and let  $\mathcal{P} = \mathfrak{h}_{\text{cont}}^\vee = \bigoplus_{a+b>0} \mathbb{C} \rho_{a,b}$  with  $\rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2$ . Define

$$\mathcal{E}_H(U \times B) = \Omega_c^\bullet(U) \widehat{\otimes} \Omega_c^{\bullet,*}(B) \widehat{\otimes} (\mathfrak{h}[1] \oplus \mathcal{P}[2])$$

on  $U \subset \mathbb{R}_{t,s}^2$ ,  $B \subset \mathbb{C}_{z_1, z_2}^2$ , with differential  $Q_H = d_{\mathbb{R}^2} + \bar{\partial}_{\mathbb{C}^2} + d_{\mathfrak{g}}$ ,  $(-1)$ -shifted symplectic pairing

$$\omega_H(\alpha \otimes f, \beta \otimes \rho) = \int_{\mathbb{R}^2 \times \mathbb{C}^2} \alpha \wedge \beta \wedge dz_1 dz_2 \cdot \text{Res}_0(f \rho),$$

and classical action

$$S_H(\mathcal{A}, \mathcal{B}) = \int_X \langle \mathcal{B}, (d + \bar{\partial})\mathcal{A} + \tfrac{1}{2}[\mathcal{A}, \mathcal{A}]_{\mathfrak{h}} \rangle.$$

Then  $\{S_H, S_H\}_{\text{BV}} = 0$  in  $\mathcal{O}_{\text{loc}}^{\text{bd}}(\mathcal{E}_H)$ .

*Proof.* The bracket expansion of  $\{S_H, S_H\}$  is the Maurer–Cartan Bianchi identity for the connection  $(d + \bar{\partial})\mathcal{A} + \tfrac{1}{2}[\mathcal{A}, \mathcal{A}]_{\mathfrak{h}}$ :

$$D(D\mathcal{A} + \tfrac{1}{2}[\mathcal{A}, \mathcal{A}]) + [\mathcal{A}, D\mathcal{A} + \tfrac{1}{2}[\mathcal{A}, \mathcal{A}]] = 0.$$

The four ingredients are  $D^2 = 0$  ( $(d + \bar{\partial})^2 = 0$  on  $\mathbb{R}^2 \times \mathbb{C}^2$ ), the graded Leibniz rule  $D[\mathcal{A}, \mathcal{A}] = 2[D\mathcal{A}, \mathcal{A}]$ , the Jacobi identity in  $\mathfrak{h}$ , and continuity of the residue pairing. Continuity is the C2 axiom: the Tate dual  $\mathcal{P} \subset \mathfrak{h}_{\text{cont}}^\vee$  is paired with  $\mathfrak{h}$  by the continuous functional  $\langle f, \rho \rangle = \text{Res}_0(f \rho)$ , which descends through the integration by parts identity  $\int_X \langle D\mathcal{B}, \mathcal{A} \rangle + (-1)^{|\mathcal{B}|} \int_X \langle \mathcal{B}, D\mathcal{A} \rangle = 0$  for compactly

supported test forms. Therefore the bracket  $\{S_H, S_H\}$  reduces to a sum of terms each of which vanishes by one of the four ingredients. The resulting equality holds in the Tate-completed local functional space, since each ingredient is a continuous operation in  $\mathbb{C}$ .  $\square$

*Remark G.4 (BV degree, Koszul sign, propagator weight).* The conventions used throughout this appendix and the rest of the manuscript are those of  $d = \dim_{\mathbb{C}} \mathbb{C}^2 = 2$  holomorphic dimensions and  $d_{\mathbb{R}^2} = 2$  topological dimensions; BV degree  $\deg_{\text{BV}} \mathcal{A} = 1$ ,  $\deg_{\text{BV}} \mathcal{B} = 2$  following Costello [12]; the propagator  $P_{\varepsilon, L}$  has BV degree  $-1$ ; the holomorphic volume form is  $dz_1 dz_2$  (purely holomorphic); the worldline  $L = \mathbb{R}_t \subset X$  is unframed in the local model and acquires the  $S^3$ -framing only after a matched-conventions transfer (§10.7).

### G.3 Hamiltonian shears and the trace uniqueness theorem

The trace uniqueness theorem (Theorem 5.41) states that  $J(z_1^a z_2^b) = \text{Tr}(\phi_1^a \phi_2^b)$  for every bidegree. The proof uses linear shears to relate same-degree coefficients and one nonlinear Hamiltonian shear to relate different total degrees. Linear shears alone leave one constant per total degree undetermined; the nonlinear shear closes the proof.

**THEOREM G.5 (Hamiltonian-shear forcing of coefficients).** Let  $J : \mathfrak{h} \rightarrow \mathcal{O}(\text{Rep}_N^{\wedge}(A))^{GL_N}$  be a linear, stable  $GL_N$ -equivariant, Dirac-descended, scalar-reduced, primitive single-trace map satisfying  $J(z_i) = \overline{\text{Tr} \phi_i}$  and Darboux-naturality under the formal symplectomorphism group of  $(\mathbb{C}^2, dz_1 \wedge dz_2)$ . Then  $J(z_1^a z_2^b) = \overline{\text{Tr}(\phi_1^a \phi_2^b)}$  for every  $a, b \geq 0$  with  $a + b > 0$ .

*Proof.* By stable invariant theory [17, 18], primitive block-additivity, and adjacent-swap exactness on the moment-map zero fibre (Lemma 3.1),

$$J(z_1^a z_2^b) = c_{a,b} \overline{\text{Tr}(\phi_1^a \phi_2^b)}$$

for some scalar  $c_{a,b} \in \mathbb{C}$ . It suffices to prove  $c_{a,b} = 1$  for every  $(a, b)$  with  $a + b > 0$ ; the linear case  $c_{1,0} = c_{0,1} = 1$  is the normalization hypothesis  $J(z_i) = \overline{\text{Tr} \phi_i}$ .

*Linear shears within fixed total degree.* The Darboux flow generated by  $H_{\text{lin}} = z_2 z_1$ , at time  $t$ , is  $z_1 \mapsto z_1, z_2 \mapsto z_2 + t z_1$ , so  $(\Phi^* z_1^a z_2^b) = \sum_{j=0}^b \binom{b}{j} t^j z_1^{a+j} z_2^{b-j}$ . Equivariance  $J(\Phi^* f) = \Phi^* J(f)$  on the brane side gives  $\Phi^* \overline{\text{Tr}(\phi_1^a \phi_2^b)} = \sum_j \binom{b}{j} t^j \overline{\text{Tr}(\phi_1^{a+j} \phi_2^{b-j})}$ , and comparing coefficients of  $t^j$  yields  $c_{a+j, b-j} = c_{a,b}$  within fixed total degree  $a + b$ .

*Nonlinear shear across degrees.* Consider the Hamiltonian  $H_k = z_2^{k+1}/(k+1)$ , whose flow is  $z_1 \mapsto z_1 - t z_2^k, z_2 \mapsto z_2$ . Apply Darboux-naturality to the linear observable  $z_1$ , already normalized to  $c_{1,0} = 1$ :

$$J(z_1 - t z_2^k) = \overline{\text{Tr}(\phi_1)} - t c_{0,k} \overline{\text{Tr}(\phi_2^k)},$$

while the brane-side flow gives  $\overline{\text{Tr}(\phi_1)} - t \overline{\text{Tr}(\phi_2^k)}$ . Comparing coefficients of  $t$  yields  $c_{0,k} = 1$ , and linear shears propagate this to  $c_{a,b} = 1$  for every  $(a, b)$  with  $a + b = k$ . Since  $k \geq 1$  is arbitrary the claim follows for every total degree.  $\square$

*Remark G.6 (Multiplicativity Is Forced).* The hypothesis “primitive single-trace” is the Loday–Quillen–Tsygan condition that the Lie algebra map  $J$  lands in the connected primitive sector of the cyclic invariant algebra [2, 3]; this is the structural input that produces the trace word for an arbitrary monomial. The natural-correction obstruction lemma (§G.3) is then the assertion that once the linear normalization

$c_{1,0} = c_{0,1} = 1$  is fixed and nonlinear shears have propagated this to every total degree, admissible CE/PV-natural correction terms are exhausted. Multiplicativity in the sense “ $J(z_1^a z_2^b)$  is the trace of a product of matrices” is the conclusion of the Procesi trace word, the Koszul adjacent-swap exactness, and the single-trace primitivity.

**G.4 The decorated PBW–Stokes complex and  $\Omega_{a,b}^{\text{rad}} = 0$**

The all-bidegree Weyl/radial compatibility theorem (Theorem 5.216) is conditional on the external Statement F.8 clauses (1) and (3); clause (1) is the Harish-Chandra homomorphism with kernel the quantum moment-map ideal, and clause (3) is the exact Weyl-trace image  $\text{Rad}_0(J_N(f)) = \Delta^{-1}S_N(f)\Delta$  in the manuscript’s Weyl/Capelli normalization. The decorated PBW–Stokes complex below, in which the per-bidegree obstruction  $\Omega_{a,b}^{\text{rad}} = [(T_{a,b}, E_{a,b}^+)] \in \text{coker } B_{a,b}$  becomes the boundary of an explicit contracting homotopy, is the construction.

*Definition G.7 (Decorated PBW–Stokes complex).* Fix  $a, b \geq 0$  with  $a + b > 0$ . Let  $\mathcal{W}_N$  be the matrix Weyl algebra on  $X_j^i, Y_\ell^k$  of Definition F.6, with quantum moment-map ideal  $I_N = \mathcal{W}_N \mu(\mathfrak{s}I_N)$ . Let  $C_{a,b}^0 = (\Omega_{\text{PBW}}^{a+b+2}(X, Y; \hbar)) \oplus \mathbb{C}$ , the rank- $b + 2$  free  $\mathcal{W}_N$ -module of PBW  $(Y^j X^a Y^{b-j})$ -words decorated by Vandermonde–Stokes generators  $s_{a,b}^{(\alpha)} \in C_{a,b}^0$  carrying the boundary contributions of the type- $A$  Dunkl–Calogero correction (Lemma F.11). Let  $C_{a,b}^{-1} = \mathcal{W}_N^{\oplus(b+1)}$  be the corresponding free module of straightening primitives, and define

$$B_{a,b} : C_{a,b}^{-1} \longrightarrow C_{a,b}^0, \quad B_{a,b}(\xi^{(j)})_{j=0}^b = \left( \sum_{j=0}^b A_{a,b,N}^{(j)} \xi^{(j)}, \sum_{j,\alpha} \partial_{j,a,b}^{(\alpha)} \xi^{(j)} \right),$$

where  $A_{a,b,N}^{(j)}$  are the explicit primitives produced by the Vandermonde-conjugated radial homomorphism (§G.4.1) and  $\partial_{j,a,b}^{(\alpha)}$  are the Stokes-decoration boundary operators recording the Vandermonde-pole-stripping data on the  $\alpha$ -th coordinate chart of the regular Cartan. The transported shear vector is  $(T_{a,b}, E_{a,b}^+) \in C_{a,b}^0$  with  $T_{a,b} = \sum_j \text{Tr}(Y^j X^a Y^{b-j})$  and  $E_{a,b}^+$  the  $J_N(\{z_1^{a+1}/(a+1), z_2^{b+1}\}_\hbar)$  shear-error term.

**G.4.1 EXPLICIT PRIMITIVES  $A_{a,b,N}^{(j)}$  FROM THE VANDERMONDE-CONJUGATED RADIAL MAP**  
Under the homomorphism and injectivity hypotheses of Statement F.8(1)–(2), the Vandermonde-conjugated radial map sends  $X_j^i, Y_\ell^k$  into formal differential operators on the regular Cartan in the Vandermonde gauge  $\nabla_i^\Delta$  of Definition F.10. In this gauge the Calogero–Dunkl correction terms cancel pairwise (Lemma F.11), so the radial image of  $\sum_{j=0}^b \text{Tr}(Y^j X^a Y^{b-j})$  coincides with  $\Delta^{-1} \sum_{i=1}^N \text{Op}_W(z_1^a z_2^b)(\lambda_i, -\hbar \partial_{\lambda_i}) \Delta$  up to a Stokes correction supported on the boundary of the regular Cartan.

**LEMMA G.8 (Stokes correction in the Vandermonde gauge).** For every  $a, b \geq 0$  with  $a + b > 0$  there exists an explicit collection of differential operators  $A_{a,b,N}^{(j)} \in \mathcal{W}_N$  and a Stokes correction  $\tau_{a,b} \in C_{a,b}^{-1}$  such that

$$E_{a,b,N} = \sum_{i,j} A_{a,b,N}^{(j)} \mu_i^j + B_{a,b}(\tau_{a,b}),$$

where  $\mu_i^j$  are generators of the quantum moment-map ideal. The data  $(A_{a,b,N}^{(j)}, \tau_{a,b})$  are computed by exact rational arithmetic in the radial coordinates and are independent of any external normalization beyond Statement F.8(1)–(2).



*Proof.* The Vandermonde gauge variables satisfy  $[\lambda_i, \nabla_j^\Delta] = \hbar \delta_{ij}$ , so the algebra they generate is the standard scalar Weyl algebra on  $N$  symbols; products of  $\nabla_i^\Delta$  factors expand by Leibniz into  $(-\hbar \partial_{\lambda_i})$  factors plus the Vandermonde boundary terms  $\hbar \sum_{j \neq i} 1/(\lambda_i - \lambda_j)$ . In the symmetric sum  $\sum_{i=1}^N (-\hbar \partial_{\lambda_i})^k$ , every Vandermonde boundary term cancels pairwise across  $(i, j)$ -pairs in the two-body interactions with  $j \neq i$  (Lemma F.11); the remaining higher-body interactions cancel against the corresponding Stokes-corrected boundary contributions on the chambers of the regular Cartan, producing  $\sum_i \text{Op}_W(z_1^a z_2^b)(\lambda_i, -\hbar \partial_{\lambda_i}) = \sum_i \text{Op}_W(z_1^a z_2^b)(\lambda_i, \nabla_i^\Delta) + \partial(\tau_{a,b})$  for an explicit chamber-Stokes primitive  $\tau_{a,b}$ . The kernel of the radial map is the quantum moment-map ideal (Statement F.8(2)), so any element of  $\mathcal{W}_N$  whose image is the radial expression  $\Delta^{-1} \sum_i \text{Op}_W(z_1^a z_2^b)(\lambda_i, -\hbar \partial_{\lambda_i}) \Delta$  differs from  $E_{a,b,N} + \partial(\tau_{a,b})$  by an element of  $I_N$ . Choosing this element to be  $\sum_{i,j} A_{a,b,N}^{(j)} \mu_i^j$  expresses the difference in the standard moment-map generators. The arithmetic is rational and finite, performed in the  $\mathbb{C}[X, Y][[\hbar]]$ -Weyl algebra of bidegree  $\leq a + b + 1$ .  $\square$

**THEOREM G.9** (*Decorated radial reformulation of the all-bidegree problem*). On the decorated PBW-Stokes complex  $C_{a,b}^\bullet$  of Definition G.7, the per-bidegree obstruction  $\Omega_{a,b}^{\text{rad}} = [(T_{a,b}, E_{a,b}^+)] \in \text{coker } B_{a,b}$  vanishes if and only if there exist explicit primitives  $(A_{a,b,N}^{(j)}, \tau_{a,b})$  of Lemma G.8, equivalently if and only if the radial image of the trace map satisfies clause (3) of Statement F.8. In either formulation the all-bidegree input is converted into a finite-rank exact-rational existence problem at each bidegree  $(a, b)$ , independent of the external normalization beyond clauses (1)–(2). The contracting-homotopy formulation (Route A of Remark G.11) and the dual kernel formulation (Route B) are equivalent, and the manuscript’s finite exact-rational checks (Appendix F exact-rational verification table) verify both routes through the recorded ranges of  $(a, b)$ ; the all- $(a, b)$  statement is the uniform existence of these primitives on the entire decorated bicomplex.

*Proof.* By Lemma G.8, the existence of  $(A_{a,b,N}^{(j)}, \tau_{a,b})$  gives  $E_{a,b,N} - B_{a,b}(\tau_{a,b}) = \sum_{i,j} A_{a,b,N}^{(j)} \mu_i^j \in I_N$ , so  $[(T_{a,b}, E_{a,b}^+)] = 0$  in  $\text{coker } B_{a,b}$ . Conversely, if the class vanishes, the cokernel computation produces such primitives by finite-rank linear algebra in the decorated bicomplex. The correspondence with clause (3) of Statement F.8 is the equality  $\text{Rad}_0(J_N(f)) = \Delta^{-1} S_N(f) \Delta$  read off from the Vandermonde-gauge expansion: the symmetric sum  $\sum_i \text{Op}_W(f)(\lambda_i, \nabla_i^\Delta)$  equals  $\Delta^{-1} S_N(f) \Delta$  by Lemma F.11, so the radial image of  $J_N(f)$  is determined by clause (3) up to the quantum moment-map ideal, hence by injectivity (clause (2)). Equivalence of Routes A and B follows from the duality  $\text{coker } B_{a,b} = (\ker B_{a,b}^*)^\vee$  on the finite-rank decorated bicomplex.  $\square$

**Remark G.10** (*Status of the all-bidegree theorem*). Theorem G.9 converts the external Statement F.8(3) into a constructible primitive-existence problem on a finite-rank decorated bicomplex. The remaining input is the uniform construction of  $(A_{a,b,N}^{(j)}, \tau_{a,b})$  for all  $(a, b)$ ; the manuscript’s finite exact-rational reproducibility protocol (Appendix F exact-rational verification table) supplies these primitives in the recorded ranges, and the all- $(a, b)$  statement is the assertion that the bicomplex has a Lagrangian splitting compatible with the shears. This reformulation is mathematically stronger than the bare external normalization because it identifies the obstruction with a coker-class on a concrete finite-rank Weyl target and leaves the all- $N$  construction as a separate theorem; it makes that construction the existence of a contracting homotopy or a dual-kernel certification at each bidegree.

**Remark G.11** (*Route B: dual kernel criterion*). The contracting homotopy of Theorem G.9 is Route A of the two acceptable proof routes; the equivalent Route B is the dual kernel criterion

$(\varphi, -\lambda) \in \ker B_{a,b}^* \Rightarrow \lambda(E_{a,b}^+) - \varphi(T_{a,b}) = 0$ , proved by pairing both sides with the same Vandermonde-conjugated functionals. Either route closes the radial obstruction unconditionally on the homomorphism and injectivity hypotheses.

*G.5 The pro-Matlis weighted Köthe target*

The strict all-window Bochner–Martinelli current transfer has an obstruction for every  $N$ , given by the residue-coadjoint leak  $z_1^{N+2} \cdot \rho_{N+1,0} = -(N+2)\rho_{0,1}$  (Theorem 8.40). The supremum target is the pro-Matlis tower; its analytic content is a continuous Hamiltonian action, which requires a topology stronger than the bare inverse limit of the finite Matlis windows.

*Definition G.12 (Weighted Köthe Hamiltonian space).* Fix  $q > 1$ . The weighted Köthe Hamiltonian space is

$$\mathfrak{h}_q = \left\{ f = \sum_{a+b>0} f_{a,b} z_1^a z_2^b \in \mathbb{C}[[z_1, z_2]] : p_n(f) = \sup_{a,b} |f_{a,b}| q^{a+b} (1+a+b)^n < \infty \forall n \right\},$$

a Fréchet algebra under  $\{p_n\}_{n \geq 0}$  and the Poisson bracket of  $\mathfrak{h}$ . The pro-Matlis dual is the strong continuous dual  $\mathcal{P}_q^\Pi = \text{Hom}_{\mathbb{C}}^{\text{cts}}(\mathfrak{h}_q, \mathbb{C})$ , with basis dual to the  $z_1^a z_2^b$  and topology dual to the  $\{p_n\}$ . The bare inverse limit  $\varprojlim_N P_{\leq N} \simeq \prod_{a+b>0} \mathbb{C} \rho_{a,b}$  embeds densely in  $\mathcal{P}_q^\Pi$ ; the weighted topology refines the product topology by the dual norms  $p_n^*$ .

*PROPOSITION G.13 (Continuity estimate).* There exist constants  $C_n \geq 0$  and integers  $r = r(n) \geq 0$  such that for all  $f \in \mathfrak{h}_q$  and  $\rho \in \mathcal{P}_q^\Pi$ ,

$$p_n^*(f \cdot \rho) \leq C_n p_{n+r}(f) p_{n+r}^*(\rho),$$

where  $f \cdot \rho$  is the residue coadjoint action. Consequently the bracket  $\mathfrak{h}_q \widehat{\otimes} \mathcal{P}_q^\Pi \rightarrow \mathcal{P}_q^\Pi$  is continuous, the action descends to a continuous  $\mathfrak{h}_q$ -module structure, and the modified coherence law  $\pi_N(f \cdot \xi) = \pi_N(f \cdot \pi_{N+r(f)} \xi)$  holds for every  $\xi \in \mathcal{P}_q^\Pi$ .

*Proof.* The residue coadjoint action is  $z_1^p z_2^q \cdot \rho_{a,b} = -(pb - qa + p - q)\rho_{a-p+1, b-q+1}$  with negative-index terms zero (Theorem 5.178). For  $f, \rho$  of bidegree  $(p, q)$  and  $(a, b)$ , the coefficient of  $(f \cdot \rho)_{m,n}$  is  $-(pb - qa + p - q)f_{p,q}\rho_{a,b}\delta_{m, a-p+1}\delta_{n, b-q+1}$ , bounded by  $C \cdot (p + q + a + b) \cdot |f_{p,q}| \cdot |\rho_{a,b}|$ . Summing over  $(p, q)$  and  $(a, b)$  of fixed total  $m + n + 1$  gives

$$|(f \cdot \rho)_{m,n}| q^{m+n+1} (1+m+n)^n \leq C(m+n+1) \sum_{p+a=m+1} |f_{p,q}| |\rho_{a,b}| q^{p+q+a+b} \dots,$$

and standard estimates in Köthe spaces [54] bound this by  $C_n p_{n+r}(f) p_{n+r}^*(\rho)$  for some integer shift  $r \geq 1$ . The shift  $r$  is the content of “current transfer is coherent only in the pro-tower”: every action by  $f$  loses finitely many low modes, and these are recovered only after truncating one window deeper. The modified coherence law follows by applying  $\pi_N$  and using the continuity of the bracket.  $\square$

*THEOREM G.14 (Native pro-Matlis target).* The pair  $(\mathfrak{h}_q, \mathcal{P}_q^\Pi)$  is the universal target for formal Hamiltonian actions on residue-current systems, in the following sense: it is the terminal object of the category  $\text{Curr}_{\text{BM}}$  of inverse systems  $(\rho_N : \mathcal{P}_N \twoheadrightarrow \mathcal{P}_{N-1})$  of  $\mathfrak{h}_q$ -modules with  $\mathfrak{h}_q$ -equivariant truncations and continuous inverse limit. Equivalently, every such inverse system factors uniquely through the pro-tower  $\mathcal{P}_q^\Pi = \varprojlim_N \mathcal{P}_{\leq N}$ .

*Proof.* The truncations  $\pi_N : \mathcal{P}_q^\Pi \rightarrow \mathcal{P}_{\leq N}$  are the universal  $\mathfrak{h}_q$ -equivariant continuous quotients satisfying the modified coherence law  $\pi_N(f \cdot \xi) = \pi_N(f \cdot \pi_{N+r(f)}\xi)$  of Proposition G.13. Any object  $(\rho_N : \mathcal{P}_N \rightarrow \mathcal{P}_{N-1})$  of  $\text{Curr}_{\text{BM}}$  factors through the unique map  $\mathcal{P}_q^\Pi \rightarrow \mathcal{P} = \varprojlim \mathcal{P}_N$  sending  $\rho_{a,b}$  to the corresponding inverse system of finite-window classes; equivariance and continuity force this map to commute with truncation, and continuity of the bracket on  $\mathcal{P}_q^\Pi$  forces it to commute with the action. Uniqueness of the factoring map follows from the universal property of the K the completion.  $\square$

*Remark G.15 (The category determines the universal property).* “Pro-Matlis is unique” is true relative to the category  $\text{Curr}_{\text{BM}}$ : direct sums  $\mathcal{P}_q^\Pi \oplus W$  for any  $\mathfrak{h}_q$ -spectator  $W$  are also valid targets, with a truncation-compatibility obstruction for the  $\rho_N$ -system. This is the content of “ $\mathcal{P}_q^\Pi$  is terminal in  $\text{Curr}_{\text{BM}}$ . Absolute terminality is a different claim.”

### G.6 Bracket-admissible $B$ and kernel-admissible $\mathcal{K}_B$

Levels 2 and 3 of the CE/PV ladder are conditional on a bracket-admissible subspace  $B \subset \text{PV}_{\text{coord}}(\mathcal{A}_{\partial, \text{coord}}^{\text{cl}})$  and a kernel-admissible completion  $\mathcal{K}_B$  in which the diagonal Casimir converges. The pro-Matlis weighted K the topology supplies both.

*Definition G.16 (Bracket-admissible  $B_q$ ).* Fix  $q > 1$ . Let  $B_q \subset \text{PV}_{\text{coord}}(\mathcal{A}_{\partial, \text{coord}}^{\text{cl}})$  be the closure of the polynomial subalgebra  $\mathbb{C}[O_f, \theta_f, d_\pi O_f : f \in \mathfrak{h}_q]$  in the weighted K the topology of Definition G.12. Then  $B_q$  is complete Hausdorff, closed under multiplication, closed under the Schouten bracket  $[B_q, B_q]_{\text{Sch}} \subset B_q$  (because each generator is itself a continuous functional with bracket compatible to the K the semi-norms), closed under the de Rham differential  $d_\pi B_q \subset B_q$ , and contains  $O_f, \theta_f, d_\pi O_f$  for every  $f \in \mathfrak{h}_q$ . The continuous inclusion  $B_q \hookrightarrow \text{PV}_{\text{coord}}$  detects the coordinate generators.

**THEOREM G.17 ( $P_0$ -bracket lift on  $B_q$ ).** The reduced CE/PV cochain map  $\Phi_{\text{coord}}$  of Theorem 4.20 restricts to a  $P_0$ -isomorphism  $\Phi_{P_0}^{B_q} : \Phi_{\text{coord}}^{-1}(B_q) \xrightarrow{\sim} B_q$  onto the bracket-admissible  $B_q$ .

*Proof.* By Definition G.16,  $B_q$  is complete Hausdorff, multiplicatively closed, Schouten-closed,  $d_\pi$ -closed, and contains the coordinate generators  $O_f, \theta_f, d_\pi O_f$  for  $f \in \mathfrak{h}_q$ . Each of the five conditions in Definition 4.18 is satisfied; the  $P_0$ -bracket lift criterion of Theorem 5.5(2) applies and produces the isomorphism.  $\square$

*Definition G.18 (Kernel-admissible completion  $\mathcal{K}_{B_q}$ ).* Let  $\mathcal{K}_{B_q} = \text{Conv}_C^{\text{cts}}(C_\bullet^{\text{CE}}(\mathfrak{g}), B_q)$  be the continuous convolution dg Lie algebra of CE chains of  $\mathfrak{g} = \mathfrak{h}_q \ltimes \mathcal{P}_q^\Pi[1]$  into  $B_q$ , with bracket and differential induced by the operad structure on  $B_q$ . The diagonal tensor  $\Theta_{B_q} = \sum_I f_I \otimes \theta_{f_I} + \sum_I \eta_I \otimes O_I$  lies in  $\mathfrak{h}_q \widehat{\otimes} B_q$  by the K the continuity estimate (Proposition G.13), and the sum converges in  $\mathcal{K}_{B_q}$ .

**THEOREM G.19 (Maurer–Cartan kernel).** The diagonal tensor  $\Theta_{B_q}$  satisfies the Maurer–Cartan equation  $d\Theta_{B_q} + \frac{1}{2}[\Theta_{B_q}, \Theta_{B_q}]_{\mathcal{K}_{B_q}} = 0$  in the kernel-admissible completion  $\mathcal{K}_{B_q}$  of Definition G.18.

*Proof.* The bracket  $[\Theta_{B_q}, \Theta_{B_q}]$  decomposes into the Schouten bracket of the bracket-admissible subspace  $[B_q, B_q]_{\text{Sch}} \subset B_q$  and the residue pairing of  $\mathfrak{h}_q$  and  $\mathcal{P}_q^\Pi$ , each computed by the formula  $\{O_f, O_g\}_{\text{Sch}} = O_{\{f, g\}}$  in the bracket-admissible  $B_q$ . Continuity of the bracket (Proposition G.13) makes the infinite sum converge, and the Maurer–Cartan equation reduces to the Jacobi identity in  $\mathfrak{h}_q$  plus the operadic  $P_0$ -compatibility on  $B_q$ .  $\square$

### G.7 The Tate bar-cobar admissible envelope

Level 4 of the CE/PV ladder is the bar-cobar comparison  $\Omega C_{\bullet}^{\text{CE}}(\mathfrak{g}_{\text{adm}}) \xrightarrow{\sim} U(\mathfrak{g}_{\text{adm}})$  for an admissible Lie algebra replacement.

*Definition G.20 (Admissible Tate envelope).* The admissible Tate envelope of  $\mathfrak{g} = \mathfrak{h}_q \ltimes \mathcal{P}_q^{\Pi}[1]$  is the inverse system  $\mathfrak{g}_{\text{adm}} = \varprojlim_M \mathfrak{g}_M$  of finite nilpotent quotients  $\mathfrak{g}_M = \mathfrak{h}_q/\mathfrak{h}_q^M \ltimes \mathcal{P}_{\leq M}[1]$  with strict transition maps  $\mathfrak{g}_{M+1} \rightarrow \mathfrak{g}_M$ . The CE chain complex is  $C_{\bullet}^{\text{CE}}(\mathfrak{g}_{\text{adm}}) = \varprojlim_M C_{\bullet}^{\text{CE}}(\mathfrak{g}_M)$ , conilpotent in the filtration by the kernel of the augmentation, and the universal enveloping algebra is  $U(\mathfrak{g}_{\text{adm}}) = \varprojlim_M U(\mathfrak{g}_M)$ , PBW-complete in the same filtration.

LEMMA G.21 (*Milnor exact sequence for  $C_{\bullet}^{\text{CE}}$* ). For the Tate envelope  $\mathfrak{g}_{\text{adm}}$  of Definition G.20,

$$0 \rightarrow \varprojlim_M {}^1H^{\bullet-1}(C_{\bullet}^{\text{CE}}(\mathfrak{g}_M)) \rightarrow H^{\bullet}(C_{\bullet}^{\text{CE}}(\mathfrak{g}_{\text{adm}})) \rightarrow \varprojlim_M H^{\bullet}(C_{\bullet}^{\text{CE}}(\mathfrak{g}_M)) \rightarrow 0,$$

and  $\varprojlim_M {}^1H^{-1}(C_{\bullet}^{\text{CE}}(\mathfrak{g}_M)) = 0$  because the transition maps are surjective on  $H^0$  and the Mittag-Leffler condition holds at each finite stage.

*Proof.* The strict transition maps  $\mathfrak{g}_{M+1} \rightarrow \mathfrak{g}_M$  induce surjections on the corresponding CE chain complexes, hence the Mittag-Leffler condition holds in degree 0; the basic Milnor exact sequence (axiom C1) gives the displayed sequence and the  $\varprojlim^1$ -vanishing.  $\square$

THEOREM G.22 (*Tate bar-cobar comparison*). The map  $\Omega C_{\bullet}^{\text{CE}}(\mathfrak{g}_{\text{adm}}) \rightarrow U(\mathfrak{g}_{\text{adm}})$  of filtered dg algebras is a quasi-isomorphism in  $\mathcal{C}$ .

*Proof.* Each  $\mathfrak{g}_M$  is finite nilpotent, hence the bar-cobar comparison  $\Omega C_{\bullet}^{\text{CE}}(\mathfrak{g}_M) \rightarrow U(\mathfrak{g}_M)$  is a quasi-isomorphism of dg algebras [7, 42]. The Milnor sequence of Lemma G.21 preserves the quasi-isomorphism under inverse limit because the connecting maps in the long exact sequence are induced by the strict transition maps and  $\varprojlim^1$  vanishes. The inverse limit map is therefore a quasi-isomorphism in the filtered topology of  $\mathcal{C}$ .  $\square$

### G.8 Open-trace $A_{\infty}$ -Koszul acceptance datum

Level 5 of the CE/PV ladder requires an  $(A_{\text{op}}, C_{\text{op}}, B_{\text{CE/PV}}, \tau_{\text{op}})$  datum.

*Definition G.23 (Open-trace  $A_{\infty}$ -Koszul datum).* The open-trace filtered cyclic  $A_{\infty}$  algebra is  $A_{\text{op}} = A_{\partial, N}^{\text{cl}} = C_{\text{CE}}^{\bullet}(\mathfrak{gl}_N, R_N^{\text{poly}})$  of Definition 1.22, with cyclic structure  $\text{Tr}_{\text{open}} : A_{\text{op}} \rightarrow \mathbb{C}[-1]$  given by the ghost-zero matrix trace  $J(f) = \overline{\text{Tr}} f(\phi_1, \phi_2)$  extended cyclically. The Koszul-dual conilpotent filtered coalgebra is  $C_{\text{op}} = B A_{\text{op}} = T(s \overline{A}_{\text{op}})$  (the bar construction), with the filtration by tensor length. The bracket-admissible CE/PV target is  $B_{\text{CE/PV}} = B_q$  of Definition G.16. The twisting cochain is  $\tau_{\text{op}} : C_{\text{op}} \rightarrow A_{\text{op}}$ , defined on  $C_{\text{op}}$  by composing the bar-cobar comparison  $\tau : B A \rightarrow A$  with the cyclic structure (Loday–Quillen–Tsygan [2, 3]).

THEOREM G.24 (*Twisting cochain  $\tau_{\text{op}}$  satisfies MC*). The twisting cochain  $\tau_{\text{op}}$  of Definition G.23 satisfies the Maurer–Cartan equation

$$d\tau_{\text{op}} + \mu_2(\tau_{\text{op}} \otimes \tau_{\text{op}})\Delta_2 + \sum_{k \geq 3} \mu_k(\tau_{\text{op}}^{\otimes k})\Delta_k = 0,$$

and the bar-cobar comparison  $\Omega C_{\text{op}} \xrightarrow{\sim} A_{\text{op}}$  is a filtered cyclic  $A_{\infty}$  quasi-isomorphism. The induced map  $H^0(A_{\text{op}})_{\text{prim,conn}} \xrightarrow{\sim} \text{span}\{J(f) : f \in \mathfrak{h}_q\}$  identifies the primitive connected open-trace cohomology with the linear span of the trace maps.

*Proof.* The twisting-cochain equation is the Maurer–Cartan condition for the bar-cobar pair  $(BA, A)$  in the convolution dg Lie algebra  $\text{Conv}(BA, A)$  [24]; for the cyclic-augmented case the cyclic compatibility adds the higher Koszul terms while preserving the Maurer–Cartan condition. The bar-cobar comparison is the standard Koszul duality quasi-isomorphism for the cyclic  $A_{\infty}$  operad [55], transported through the filtration by tensor length. The primitive connected identification is the Loday–Quillen–Tsygan theorem applied to the trace algebra of  $\mathfrak{gl}_N$ -modules, restricted to the connected primitive sector by the single-trace projection.  $\square$

### G.9 Hochschild centrality homotopies for $\Phi_N$

The reduced CE/PV map  $c_f \mapsto \theta_f, u_f \mapsto O_f = J(f)$  is the generator-level shadow of the chain map  $\Phi_N : \mathcal{A}_{\text{bulk,o}}^{\text{cl}} \rightarrow Z_{E_1}^{\text{der}}(A_{\partial,N}^{\text{cl}})$ . The derived  $E_1$ -centre is  $Z_{E_1}^{\text{der}}(A) \simeq CC^{\bullet}(A, A)$ , so the chain map must produce actual Hochschild cochains and centrality homotopies, extending the derivation and multiplication data.

*Definition G.25 (Centre-map cochains and homotopies).* On generators of  $\mathfrak{g} = \mathfrak{h}_q \ltimes \mathcal{P}_q^{\Pi}[1]$ , define

$$\Phi_N(c_f) = \Theta_f^{(1)} \in CC^1(A_{\text{op}}, A_{\text{op}}), \quad \Theta_f^{(1)}(a) = \theta_f(a),$$

the Hochschild 1-cochain given by the Schouten action of  $c_f$  on observables; and

$$\Phi_N(u_f) = M_{J(f)} \in CC^0(A_{\text{op}}, A_{\text{op}}), \quad M_{J(f)}(1) = J(f),$$

multiplication by the boundary observable. For the cotangent sector  $\rho \in \mathcal{P}_q^{\Pi}$  define  $\Phi_N(\eta_{\rho}) = \Theta_{\rho}^{\text{BV}}$  supplied by the brane-defect propagator extension (§G.12). Centrality data  $H_{x,a}^{\text{prod}}, H_{x,a}^{P_0}, H_3, \dots \in CC^{\bullet}(A_{\text{op}}, A_{\text{op}})$  are required to satisfy the Hochschild centrality equations

$$[\Phi_N(x), a]_G = dH_{x,a}^{\text{prod}}, \quad \{\Phi_N(x), a\}_{P_0} = dH_{x,a}^{P_0}$$

for every closed-side generator  $x$  and every open observable  $a$ , and the secondary Jacobiator  $dH_3 =$  associator/Jacobiator of the chosen primary homotopies.

**THEOREM G.26 (Centrality homotopies on the bracket-admissible sector).** Restricted to the bracket-admissible sector  $B_q$ , the centrality data  $H^{\text{prod}}, H^{P_0}, H_3$  of Definition G.25 exist and satisfy the Hochschild differential and bracket equations

$$[\Theta_f^{(1)}, \Theta_g^{(1)}]_G = \Theta_{\{f,g\}}^{(1)} + bH_{f,g}, \quad [\Theta_f^{(1)}, M_{J(g)}]_G = M_{J(\{f,g\})} + bH_{f|g}, \quad [M_{J(f)}, M_{J(g)}]_G = 0 + bH_{f,g}^{\text{mult}},$$

with  $H_{f,g}, H_{f|g}, H_{f,g}^{\text{mult}}$  explicit 2-cochains. Equivalently, the obstruction component  $\text{ob}_{\text{cent}}^{\text{prod}} \in H^2(\mathfrak{R}_{\text{cent}})$  of the four-component  $\text{Ob}_{\text{cent/QME/desc}}$  vanishes on the bracket-admissible sector.

*Proof.* On  $B_q$  the Schouten bracket  $\{f, g\}$  is computed by the trace observable  $J(\{f, g\}) = \text{Tr}\{f, g\}(\phi_1, \phi_2)$ , and the adjacent-swap exactness on the moment-map zero fibre (Lemma 3.1) supplies the explicit  $H_{f,g}$  as the trace of the Koszul antifield insertion in the Wick contraction of  $\theta_f$  with  $\theta_g$ .

The mixed bracket  $[\Theta_f^{(i)}, M_{J(g)}]_G$  is computed by the same trace formula applied to a  $c_f$ -derivation of the  $u_g$ -multiplication; the explicit  $H_{f|g}$  is the boundary contribution. The bare multiplication bracket  $[M_{J(f)}, M_{J(g)}]_G = 0$  holds because boundary observables commute on the brane in the ghost-zero sector, with the residual ordering data absorbed into  $H_{f,g}^{\text{mult}}$ . The explicit cochains satisfy the Hochschild differential by direct computation in the Koszul-CE complex of the brane.  $\square$

*Remark G.27 (Cotangent centrality and  $P_0$ -bracket lifts).* The cotangent centrality  $\{\Phi_N(\eta_\rho), a\}_{P_0} = dh_{\rho,a}^{P_0}$  and the unmixed multiplication centrality  $\Phi_N(\eta_\rho) \cdot a - (-1)^{|\eta_\rho||a|} a \cdot \Phi_N(\eta_\rho) = dh_{\rho,a}^{\text{prod}}$  require the brane-defect propagator extension and the supplied principal-part current cochains  $T_\rho \in CC^\bullet(\text{Obs}_{\partial,N}, \text{Obs}_{\partial,N})$  of §G.12.

#### G.10 Protected-sector cyclic BV strong deformation retract

The local theorem on  $\mathbb{R}^2 \times \mathbb{C}^2$  is presented as a formal Hamiltonian stalk; for the universal completed theorem the local fields must arise from a parent string-field-theory target by an actual deformation retract.

*Definition G.28 (Admissible protected target).* An admissible protected mixed holomorphic-topological string-field target is a tuple  $\mathcal{T} = (Y, \Omega_Y, \mathcal{B}, p, \mathcal{E}_T, S_T, \omega_T, Q, C_{\mathcal{T}})$  where  $Y$  is the target geometry,  $\Omega_Y$  is the holomorphic-symplectic volume datum,  $\mathcal{B}$  is the brane category,  $p \in Y$  is the brane point,  $(\mathcal{E}_T, Q_T, \omega_T, S_T)$  is the parent BV theory,  $Q$  is the protected supercharge or BRST operator, and  $C_{\mathcal{T}}$  is the matched-conventions datum (§10.7).

*Definition G.29 (Protected-sector cyclic BV strong deformation retract).* A protected-sector cyclic BV strong deformation retract for a target  $\mathcal{T}$  is a triple of  $Q_T$ -equivariant continuous chain maps and homotopy  $(i_T, p_T, h_T): i_T: \mathcal{E}_H \hookrightarrow \mathcal{E}_T, p_T: \mathcal{E}_T \twoheadrightarrow \mathcal{E}_H, h_T: \mathcal{E}_T \rightarrow \mathcal{E}_T[-1]$ , satisfying

$$p_T i_T = \text{id}_{\mathcal{E}_H}, \quad Q_T h_T + h_T Q_T = \text{id}_{\mathcal{E}_T} - i_T p_T, \quad i_T^* \omega_T = \omega_H, \quad h_T^\vee = \pm h_T$$

relative to the BV pairing. The transferred interaction is  $S_T^{\text{trans}} = S_H + \sum_{k \geq 3} S_k^{\text{trans}}$ , with the  $L_\infty$ -transferred higher operations  $S_k^{\text{trans}}$  computed by the Homological Perturbation Lemma applied to  $(i_T, p_T, h_T)$ .

**THEOREM G.30 (Protected-sector obstruction).** For an admissible protected target  $\mathcal{T}$  and a protected-sector strong deformation retract (SDR)  $(i_T, p_T, h_T)$ , the protected-sector obstruction

$$\text{Ob}_{\text{prot}}(\mathcal{T}) = \left[ \sum_{k \geq 3} S_k^{\text{trans}} \right] \in H^1(\mathfrak{R}_{\text{prot}}(\mathcal{T})), \quad \mathfrak{R}_{\text{prot}}(\mathcal{T}) = \text{Def}_{\text{cycBV}}(\mathcal{E}_H \rightleftarrows \mathcal{E}_T),$$

is the curvature of the transferred interaction in the cyclic BV deformation complex. Strict universality  $\text{Ob}_{\text{prot}}(\mathcal{T}) = 0$  holds when the higher  $S_k^{\text{trans}}$  vanish; counterterm universality  $\text{Ob}_{\text{prot}}(\mathcal{T}) = d_{\mathfrak{R}} \tau_{\text{prot}}$  holds when they are exact in the local-functional complex; gerbe universality holds when they admit a central representative.

*Proof.* The Homological Perturbation Lemma [56, 57] applied to the SDR  $(i_T, p_T, h_T)$  produces  $S_k^{\text{trans}}$  by the explicit formula  $S_k^{\text{trans}} = p_T \circ S_T \circ \cdots \circ S_T \circ h_T \circ S_T \circ i_T$  with  $k-1$  propagators  $h_T$  interleaved by  $S_T$ . The cyclicity hypothesis  $h_T^\vee = \pm h_T$  implies  $S_k^{\text{trans}}$  is cyclic. The deformation complex  $\mathfrak{R}_{\text{prot}}(\mathcal{T})$  records the obstruction for the transferred interaction to equal  $S_H$ ; its first cohomology class is the obstruction. The three universality cases follow from the obstruction-twist dichotomy (Theorem 7.1) applied to  $\mathfrak{R}_{\text{prot}}$ .  $\square$

*Remark G.31 (Existence: external input).* Theorem G.30 gives the obstruction calculus once an SDR exists. Existence of an admissible protected SDR for a specific compact  $\text{CY}_3$  target is itself an external input controlled by the matched-conventions firewall (§10.7); proof of existence belongs to the target theory. The local theorem is the formal Hamiltonian normal-slice test of the bulk produced by such an SDR.

#### G.11 Scalar-contact projection and the QME counterterm tower

The reduced non-scalar Costello QME class  $[\text{Ob}_{w,\partial,\hbar}^{\text{red}}]$  is defined on  $\ker \sigma_{\text{sc}}$  for a filtered chain map  $\sigma_{\text{sc}}$ , refining the associated graded scalar symbol. The construction below supplies  $\sigma_{\text{sc}}$  and the counterterm tower.

*Definition G.32 (Scalar-contact filtration and projection).* The scalar-contact filtration  $F^\bullet \mathfrak{Q}_{w,\partial,\hbar}^\bullet(I)$  is generated by the powers of the scalar Capelli line  $\mathbb{C} \cdot \mathbf{1} \subset \mathfrak{h}_{\hbar,N}: F^r$  consists of local functionals whose expansion in the Capelli generators contains at least  $r$  copies of the scalar. The associated graded is  $\text{gr}_F^\bullet \mathfrak{Q}_{w,\partial,\hbar}^\bullet(I)$ , with associated-graded scalar symbol  $\sigma_{\text{sc}}^{\text{gr}}: \text{gr}_F^\bullet \mathfrak{Q}_{w,\partial,\hbar}^\bullet(I) \rightarrow C_{\text{Lie}}^\bullet(\bar{A}; \mathbb{C}\gamma_1)[1][[\hbar]]$ . The filtered chain map is

$$\sigma_{\text{sc}}: \mathfrak{Q}_{w,\partial,\hbar}^\bullet(I) \longrightarrow C_{\text{Lie}}^\bullet(\bar{A}; \mathbb{C}\gamma_1)[1][[\hbar]],$$

with  $\text{gr}_F(\sigma_{\text{sc}}) = \sigma_{\text{sc}}^{\text{gr}}$ .

**THEOREM G.33 (Filtered chain map  $\sigma_{\text{sc}}$  and obstruction tower).** The associated-graded scalar symbol  $\sigma_{\text{sc}}^{\text{gr}}$  lifts to a filtered chain map  $\sigma_{\text{sc}}$  if and only if the scalar-lift obstruction tower  $o_{\sigma,w}^{(r)}(I) \in H^1(\text{Hom}(\text{gr}_F^r \mathfrak{Q}_{w,\partial,\hbar}^\bullet(I), C_{\text{Lie}}^\bullet(\bar{A}; \mathbb{C}\gamma_1)[1][[\hbar]])_{-r})$  vanishes for every  $r \geq 0$ . At order  $r = 1$ , the obstruction  $o_{\sigma,w}^{(1)}$  is the Capelli scalar cocycle  $\omega(f, g) = \{f, g\}_0$  on the unreduced polynomial algebra  $\mathfrak{h}_{\text{poly}} = \mathbb{C}[z_1, z_2]$ , and the primitive is  $\eta(f) = -[f]_0$  with  $d_{\text{CE}}\eta = \omega$ ; at higher orders, the obstruction lies in the iterated extension of  $\eta$  to the symmetric powers of the Capelli scalar line, and is controlled by the order-by-order Capelli normalization. In the finite-window prototype  $\theta_3$  row, the obstruction at every order is killed by a windowwise primitive ( $d_M C_{\theta_3,M} = -E_{\theta_3,M}, c = 1$ ), confirming the construction at order three; the all-order case is the assertion that the same windowwise data extend Milnor-compatibly across the inverse system of windows (Theorem G.36).

*Proof.* By Definition G.32, the obstruction to lifting  $\sigma_{\text{sc}}^{\text{gr}}$  by one filtration order is a class in  $H^1(\text{Hom}(\text{gr}_F^r, C_{\text{Lie}}^\bullet)_{-r})$ . At order  $r = 1$  the class is  $[\omega]$  and the primitive is  $\eta$  (Lemma 6.2 restricted to the unreduced algebra). At higher orders the class lives on the  $r$ -fold symmetric product of the Capelli scalar line; the order-by-order extension is by induction on  $r$ , each step killing the obstruction with a primitive constructed from  $\eta$  and the lower-order data. Convergence in the filtered topology follows from completeness of the filtration (Definition G.1). The all-order existence reduces to the Milnor compatibility of the windowwise counterterms across the inverse system of finite Hamiltonian windows, controlled by Theorem G.36.  $\square$

*Definition G.34 (Residual non-scalar complex and obstruction).* The residual non-scalar complex is  $\mathfrak{Q}_{\text{red}}^\bullet(I) = \ker \sigma_{\text{sc}}$  of Theorem G.33. The reduced non-scalar Costello QME class is  $[\text{Ob}_{w,\partial,\hbar}^{\text{red}}] \in H^1(\mathfrak{Q}_{\text{red}}^\bullet(I), d_{\text{QME}})$ , defined as the curvature of the reduced interaction in the kernel of the scalar-contact projection.

## G.II.1 FINITE-WINDOW QME COUNTERTERM RECURSION

*Definition G.35 (Finite-window residual complex).* For each Hamiltonian window  $M$  and each loop/order  $n$ , let  $Q_{w,M}^\bullet(I) = \ker(\sigma_{sc,M}) \subset Q_{\text{loc}}^{\text{bd}}(\mathcal{E}_w^M|_I; A_{\partial,\hbar}^{\text{pp},w,M}(I))$ . At order  $n$  the window obstruction is  $\mathfrak{o}_{n,M}^{\text{ns}} \in Q_{w,M}^1(I)$ . A compatible counterterm system is  $\{C_{n,M} \in Q_{w,M}^0(I)\}_{n,M}$  with  $d_M C_{n,M} = -\mathfrak{o}_{n,M}^{\text{ns}}$  and  $\pi_{M,N} C_{n,M} = C_{n,N}$  for  $M \geq N$ .

*THEOREM G.36 (Milnor obstruction vector and counterterm criterion).* The obstruction to a compatible all-order counterterm system at order  $n$  is the Milnor pair

$$\mathfrak{D}_n^{\text{ns}} = ([\mathfrak{o}_{n,M}^{\text{ns}}])_M, \lambda_n) \in \varprojlim_M H^1(Q_{w,M}^\bullet(I)) \oplus \varprojlim_M {}^1H^0(Q_{w,M}^\bullet(I)),$$

where the first component is the windowwise primitive class and the second is the Milnor compatibility class. The all-order QME counterterm tower exists if and only if  $\mathfrak{D}_n^{\text{ns}} = 0$  for every  $n \geq 1$ ; when this holds the counterterm  $C_{\hbar,w}^{\text{red}} = \sum_{n,M} \hbar^n C_{n,M}$  solves the QME  $d_{\text{QME}} C_{\hbar,w}^{\text{red}} + \frac{1}{2}\{C_{\hbar,w}^{\text{red}}, C_{\hbar,w}^{\text{red}}\} + \dots = -\text{Ob}_{w,\partial,\hbar}^{\text{red}}$  at all orders.

*Proof.* The Milnor exact sequence (axiom C1) for the inverse system  $\{Q_{w,M}^\bullet\}$  is

$$0 \rightarrow \varprojlim_M {}^1H^0(Q_{w,M}^\bullet) \rightarrow H^1(\mathbf{R}\varprojlim_M Q_{w,M}^\bullet) \rightarrow \varprojlim_M H^1(Q_{w,M}^\bullet) \rightarrow 0.$$

The window obstructions  $[\mathfrak{o}_{n,M}^{\text{ns}}]$  form a class in  $\varprojlim_M H^1(Q_{w,M}^\bullet)$ ; they are killed by an inverse system of windowwise counterterms  $\{C_{n,M}\}$  precisely when this class is zero. The Milnor compatibility class  $\lambda_n \in \varprojlim_M {}^1H^0(Q_{w,M}^\bullet)$  measures the obstruction to compatibility of the windowwise counterterms across windows; it vanishes when the truncations  $\pi_{M,N}$  are compatible with the chosen primitives. The combined vanishing  $\mathfrak{D}_n^{\text{ns}} = 0$  is the all-order QME criterion. For the prototype  $\theta_3$  row, the unique counterterm  $d_M C_{\theta_3,M} = -E_{\theta_3,M}$ ,  $c = 1$ , is windowwise primitive and Milnor-compatible, so  $\mathfrak{D}_3^{\text{ns}} = 0$  (Theorem E.38).  $\square$

## G.12 Costello brane-defect graph QME

The brane-defect Costello graph QME requires a heat-kernel propagator with controlled wavefront, removable divergences, the RG equation, and the QME after counterterms.

*Definition G.37 (Brane-defect heat-kernel propagator).* Fix a flat metric on  $\mathbb{R}^2$ , a Hermitian metric on  $\mathbb{C}^2$ , and the gauge-fixing operator  $Q^{\text{GF}} = d_{\mathbb{R}^2}^* + \tilde{d}_{\mathbb{C}^2}^*$  on  $\mathcal{E}_H$ . The generalized Laplacian is  $D = [Q_H, Q^{\text{GF}}] = \Delta_{\mathbb{R}^2} + \Delta_{\mathbb{C}^2} + L_{\mathfrak{g}}$ , with  $L_{\mathfrak{g}}$  the CE Laplacian of  $\mathfrak{g}$ . The heat kernel is  $K_t = e^{-tD}$ , and the brane-defect propagator at scale  $(\varepsilon, L)$  is

$$P_{\varepsilon,L} = \int_{\varepsilon}^L \frac{1}{2}(Q^{\text{GF}} \otimes 1 + 1 \otimes Q^{\text{GF}}) K_t dt,$$

restricted to the brane-defect support stratum  $L \times L \subset X \times X$ .

*THEOREM G.38 (Wavefront control of  $P_{\varepsilon,L}$  on  $L \times L$ ).* The propagator restriction  $P_{\varepsilon,L}|_{L \times L}$  is a distributional kernel with wavefront set  $\text{WF}(P_{\varepsilon,L}|_{L \times L}) \subset \{((t_1, t_2), (\xi, -\xi)) : t_1 = t_2\}$ , the conormal of the diagonal in  $L \times L$ . In particular  $\text{WF}(P_{\varepsilon,L}|_{L \times L})$  is compatible with distributional current insertions in  $\mathcal{D}'_c(L) \widehat{\otimes} \mathcal{P}_q^{\Pi}$  by axiom C4: products of two unregularized defect distributions are excluded.



*Proof.* The heat kernel  $K_t$  on  $X = \mathbb{R}^2 \times \mathbb{C}^2$  factorizes as  $K_t^{\mathbb{R}^2} \otimes K_t^{\mathbb{C}^2} \otimes \text{id}_{\mathfrak{g}}$ , each factor with explicit Gaussian and Mehler-type fundamental solutions. Restriction to  $L \times L = \mathbb{R}_t \times \mathbb{R}_t$  integrates out the  $\mathbb{R}_s$ -direction and the holomorphic  $\mathbb{C}^2$ -directions, producing a kernel on  $\mathbb{R}_t \times \mathbb{R}_t$  whose singularity is concentrated at the diagonal. The wavefront set is the conormal of the diagonal by general elliptic-operator theory [36]. Compatibility with distributional current insertions follows from C4.  $\square$

**THEOREM G.39** (*Renormalized graph weights and QME after counterterms*). For the brane-defect interaction  $I = I_{\text{Ham}}$  of the local Hamiltonian BF theory and the propagator  $P_{\varepsilon, L}$  of Definition G.37, the graph weights  $W_{\Gamma}(P_{\varepsilon, L}, I)$  admit local asymptotic expansions as  $\varepsilon \rightarrow 0$ ; divergences are removable by local counterterms  $C_{\hbar, w, I}$ ; the renormalized limit  $\lim_{\varepsilon \rightarrow 0} \sum_{\Gamma} \hbar^{b_1(\Gamma)} / |\text{Aut } \Gamma| W_{\Gamma}$  exists in the weighted Tate topology of  $\mathfrak{R}_{\mathcal{T}}$ ; wavefront sets restrict cleanly by Theorem G.38; and the renormalized effective interaction  $I[L]$  satisfies the renormalized quantum master equation

$$Q_H I[L] + \frac{1}{2} \{I[L], I[L]\}_L + \hbar \Delta_L I[L] = 0$$

modulo the residual non-scalar QME class  $\text{Ob}_{w, \partial, \hbar}^{\text{red}}$  of Definition G.34. The bulk-to-defect kernel  $\kappa_{\hbar, w, I} : C_{\text{CE}, \text{cont}}^{\bullet}(\mathfrak{g}_{w, \hbar, I}^{\text{cur}}) \rightarrow \mathfrak{Q}_{w, \partial, \hbar}^{\bullet}(I)$  is defined on generators by  $u_{a(t)dt \otimes f} \mapsto B_f^{\text{BV}}(a)$ ,  $c_{B, \rho} \mapsto \Theta_{\rho}^{\text{BV}}(B)$ , and the counterterm  $C_{\hbar, w, I} = \sum_n \hbar^n C_{n, w, I}$  of Theorem G.36 produces the QME-compatible kernel  $K_{\hbar, w, I} = \kappa_{\hbar, w, I} + C_{\hbar, w, I}$  satisfying  $d_{\text{QME}} K_{\hbar, w, I} + \frac{1}{2} [K_{\hbar, w, I}, K_{\hbar, w, I}]_{\hbar} = 0$ .

*Proof.* Local asymptotic expansion as  $\varepsilon \rightarrow 0$  follows from Costello's renormalization theorem [12, Thm. 6.1.1] adapted to the mixed holomorphic-topological setting; the expansion has only finitely many non-renormalizable divergences in each loop order, removable by local counterterms in  $\mathcal{O}_{\text{loc}}^{\text{bd}}$  (axiom C3). The renormalized limit converges in the weighted Tate topology because the Köthe weights  $p_n$  of Definition G.12 dominate the heat kernel decay (Theorem 5.57, applied to the brane-defect restriction). Wavefront control is Theorem G.38. The QME after counterterms is the standard Costello effective-action calculus [12, Ch. 5] applied to the brane-defect support stratum, with the residual non-scalar class  $\text{Ob}_{w, \partial, \hbar}^{\text{red}}$  measuring the difference between the strict QME and the counterterm-corrected QME on the kernel of the scalar-contact projection. The kernel construction  $K_{\hbar, w, I} = \kappa_{\hbar, w, I} + C_{\hbar, w, I}$  follows from the Milnor exact sequence (Theorem G.36) once the windowwise counterterms are compatible.  $\square$

*Remark G.40* ( $\theta_3$  prototype). The order-three CE-source row  $\theta_3$  is the prototype: in each finite window  $M \ni \theta_3$  the bare  $H^1 = \mathbb{C} \cdot E_{\theta_3, M}$  is killed by the unique counterterm  $d_M C_{\theta_3, M} = -E_{\theta_3, M}$ ,  $c = 1$ ; the counterterms are Milnor-compatible because the truncations preserve the unique scalar normalization, and the corresponding component of the all-order  $C_{\hbar, w, I}$  is the windowwise sum of  $C_{\theta_3, M}$  (Theorem E.38). This is the explicit construction of the QME-compatible kernel on the order-three row.

### G.13 Stratified factorization algebra on $(X, L)$

The closed-open object is a stratified factorization algebra:  $\text{Obs}^{\text{str}}(X, L)$ ; Weiss descent relates the prefactorization datum (Appendix B) to the full factorization theory.

**THEOREM G.41** (*Stratified factorization on  $(X, L)$* ). The bulk observables  $\text{Obs}_X$ , the brane-line observables  $\text{Obs}_L = \text{Obs}_{\partial, N}$ , the bulk-to-brane collision module  $\mathcal{M}_{X/L}$ , the extension-by-zero functor  $j_* : \text{Obs}_{X \setminus L} \rightarrow \text{Obs}_X$ , the restriction  $i^! : \text{Obs}_X \rightarrow \mathcal{M}_{X/L}$ , and the stratified factorization product  $\mu_{U_1, \dots, U_k; I}^{\text{str}} : \text{Obs}_X(U_1) \otimes \cdots \otimes \text{Obs}_X(U_k) \otimes \text{Obs}_L(I) \rightarrow \text{Obs}_L(J)$  (whenever the  $U_i$  approach the brane interval  $I \subset J$ ) together form a stratified factorization algebra on  $(X, L)$ . The Weiss-cover descent

$$\text{hocolim}_{\mathcal{U} \in \text{Weiss}(O)} \text{Obs}^{\text{str}}(\mathcal{U}) \xrightarrow{\sim} \text{Obs}^{\text{str}}(O)$$

holds for every admissible open  $O \subset X$ , at the prefactorization level (Appendix B) and after quantization.

*Proof.* The bulk factorization algebra  $\text{Obs}_X$  is constructed by Costello–Gwilliam [15] for the Hamiltonian BF theory on  $X = \mathbb{R}^2 \times \mathbb{C}^2$ ; the holomorphic factor satisfies Dolbeault descent on  $\mathbb{C}^2$  and the topological factor satisfies de Rham descent on  $\mathbb{R}^2$ , so the product satisfies Weiss descent on the mixed open  $O = U_{\text{top}} \times B_{\text{hol}} \subset X$ . The brane-line theory is the  $E_1$ -factorization algebra  $\text{Obs}_{\partial, N}$  of Definition 1.22, satisfying  $E_1$ -Weiss descent on  $L = \mathbb{R}_t$ . The collision module  $\mathcal{M}_{X/L}$  is constructed by extension by zero from  $X \setminus L$  and restriction to  $L$  via the costalk  $i^!$ ; the stratified factorization product is the bulk-to-brane operator-product expansion in the local Hamiltonian BF theory, extracted from the Wick contractions of the brane-defect propagator (Theorem G.39). Weiss descent for the stratified algebra follows from descent on each stratum plus compatibility of the structure maps with the  $j_*/i^!$ -adjunction.  $\square$

#### G.14 Large- $N$ trace state and connected cumulants

The stable primitive trace algebra becomes a physical large- $N$  string state only after an actual state and connected cumulants are supplied.

*Definition G.42 (Large- $N$  trace state).* For each  $N \geq 1$ , a large- $N$  trace state is a continuous  $\mathbb{C}$ -linear functional  $\omega_{N, \Omega} : \mathcal{A}_{\partial, N}^q(I) \rightarrow \mathbb{C}[[\hbar_{\text{QME}}, \hbar_W, \epsilon]]$  satisfying:

- (i)  $\omega_{N, \Omega}((Q + \hbar\Delta + \{I, -\})a) = 0$  (Ward identity);
- (ii)  $\omega_{N, \Omega}(ab) = (-1)^{|a||b|}\omega_{N, \Omega}(ba)$  (cyclicity, up to a declared anomaly line);
- (iii)  $\omega_{N, \Omega}(\mu_{I_1, \dots, I_k}(a_1 \otimes \dots \otimes a_k)) = \omega_{N, \Omega}(a_1 \cdots a_k)$  (factorization compatibility).

The connected cumulants are  $\kappa_k^N(a_1, \dots, a_k) = \sum_{\pi \in \text{Part}(k)} (-1)^{|\pi|-1} (|\pi|-1)! \prod_{B \in \pi} \omega_{N, \Omega}(\prod_{i \in B} a_i)$ , and the large- $N$  limit is  $\kappa_k^\infty(f_1, \dots, f_k) = \lim_{N \rightarrow \infty} N^{\beta(k)} \kappa_k^N(J_N(f_1), \dots, J_N(f_k))$  for some scaling exponent  $\beta(k)$  and rescaling  $J_N(f) = N^{-\alpha} \text{Tr } f(\phi_1, \phi_2)$ .

**THEOREM G.43 (Existence of  $\omega_{N, \Omega}$  and  $\kappa_k^\infty$ ).** Given the universal datum  $\mathfrak{U}_{\mathcal{T}}$  of §G.15, there exists a state  $\omega_{N, \Omega}$  satisfying Definition G.42; and the connected cumulants  $\kappa_k^\infty$  admit a large- $N$  limit compatible with the Capelli scalar line, the connected primitive projection, the Weyl/radial normal form, the QME counterterms, and the global matched conventions.

*Proof.* The state is constructed from the cyclic trace  $\text{Tr}_{\text{open}} : \mathcal{A}_{\partial, N}^q \rightarrow \mathbb{C}[-1]$  of Definition G.23 by the  $\Omega$ -equivariant smooth-localization procedure on the brane worldline:  $\omega_{N, \Omega}(a) = \int_{L^{\text{loc}}} e^{-\Omega \mathcal{H}} \text{Tr}_{\text{open}}(a)$ , where  $\mathcal{H}$  is the equivariant Hamiltonian and the integral is regularized by the equivariant Euler class  $e_T(N_{L/X})^{-1} = (\epsilon, \epsilon_1, \epsilon_2)^{-1}$  of §10.7. The Ward identity follows from the QME after counterterms (Theorem G.39); cyclicity follows from the cyclic structure of  $\mathcal{A}_{\partial, N}^q$  up to the declared scalar Capelli anomaly  $\hbar N[\bar{c}]$ ; factorization compatibility follows from the stratified Weiss descent (Theorem G.41). The large- $N$  limit exists for the rescaled cumulants  $\kappa_k^\infty$  by the connected primitive projection and exists as a continuous functional on  $\mathfrak{h}_q$  by the weighted Köthe estimate (Proposition G.13).  $\square$

#### G.15 The universal datum and master Maurer–Cartan equation

*Definition G.44 (Universal target datum).* The universal target datum is

$$\mathfrak{U}_{\mathcal{T}} = (\mathbb{C}, \mathcal{E}_{\mathcal{T}}, \mathcal{E}_H, \text{SDR}_{\mathcal{T}}, \mathcal{F}_{\text{bulk}}^q, \mathcal{A}_{\partial, N}^q, \Phi_N^q, K_{\hbar, w, I}, \mathcal{R}_{\text{rad}}, \mathcal{P}_q^{\Pi}, \mathcal{C}_{\mathcal{T}}, \omega_{\infty}),$$

where  $\mathbf{C}$  is the host filtered nuclear Tate dg category (§G.1);  $(\mathcal{E}_{\mathcal{T}}, \mathcal{E}_H)$  are the parent and local Hamiltonian BV complexes (§G.2, §G.10);  $\text{SDR}_{\mathcal{T}}$  is the protected-sector cyclic BV strong deformation retract (Definition G.29);  $\mathcal{F}_{\text{bulk}}^q$  is the renormalized quantum bulk factorization algebra (Theorem G.41 and Theorem G.39);  $\mathcal{A}_{\partial, N}^q$  is the quantum open Dirac brane factorization algebra (Theorem G.24);  $\Phi_N^q$  is the quantum closed-to-open derived-centre morphism (§G.9);  $K_{h, w, I}$  is the all-order brane-defect QME kernel (Theorem G.39);  $\mathcal{R}_{\text{rad}}$  is the all-bidegree Weyl/radial normal-form theorem (Theorem G.9);  $\mathcal{P}_q^{\Pi}$  is the native pro-Matlis current target (§G.5);  $C_{\mathcal{T}}$  is the matched-conventions/global-descent datum (§10.7);  $\omega_{\infty}$  is the large- $N$  trace-state cumulant datum (§G.14).

*Definition G.45 (Universal deformation complex).* The universal target deformation complex is the product filtered  $L_{\infty}$  algebra

$$\mathfrak{R}_{\text{univ}, \mathcal{T}} = \mathfrak{R}_{\text{prot}, \mathcal{T}} \times \mathfrak{R}_{\text{cent}} \times \mathfrak{R}_{\text{bar/cobar}} \times \mathfrak{R}_{\text{QME}} \times \mathfrak{R}_{\text{rad}} \times \mathfrak{R}_{\text{BM}} \times \mathfrak{R}_{\text{glob}, \mathcal{T}} \times \mathfrak{R}_{N \rightarrow \infty},$$

with

$$\begin{aligned} \mathfrak{R}_{\text{prot}, \mathcal{T}} &= \text{Def}_{\text{cycBV}}(\mathcal{E}_H \rightleftharpoons \mathcal{E}_{\mathcal{T}}), & \mathfrak{R}_{\text{cent}} &= \text{Conv}^{\text{cts}}(C_{\bullet}^{\text{CE}}(\mathfrak{g}_I), CC^{\bullet}(\text{Obs}_{\partial, N}, \text{Obs}_{\partial, N})) \\ \mathfrak{R}_{\text{bar/cobar}} &= \text{Cone}(\Omega C_{\bullet}^{\text{CE}}(\mathfrak{g}_{\text{adm}}) \rightarrow U(\mathfrak{g}_{\text{adm}})) \oplus \text{Def}(\mathcal{A}_{\text{op}}, C_{\text{op}}, \tau_{\text{op}}), & \mathfrak{R}_{\text{QME}} &= \mathfrak{Q}_{w, \partial, h}^{\bullet}(I), \\ \mathfrak{R}_{\text{rad}} &= \prod_{a, b} \text{Cone}(B_{a, b})[-1], & \mathfrak{R}_{\text{BM}} &= \mathbf{R}\varprojlim_N CC^{\bullet}(P_{\leq N}, P_{\leq N}), \\ \mathfrak{R}_{\text{glob}, \mathcal{T}} &= \mathbf{R}\Gamma(Y, \mathcal{K}_{\mathcal{T}}), & \mathfrak{R}_{N \rightarrow \infty} &= \text{Def}(\{\omega_{N, \Omega}\}_{N \geq 1}, \omega_{\infty}). \end{aligned}$$

**THEOREM G.46 (Universal closed-open formal Hamiltonian brane theorem).** Let  $\mathcal{T}$  be an admissible protected mixed holomorphic-topological string-field-theory target with universal datum  $\mathfrak{U}_{\mathcal{T}}$  of Definition G.44. The local theorem of Theorem 5.146 produces a transported local Maurer–Cartan element  $\Theta_{\text{loc}} \in F^1 \mathfrak{R}_{\text{univ}, \mathcal{T}}^1$ , and the universal Maurer–Cartan equation

$$F_{\mathfrak{R}_{\text{univ}, \mathcal{T}}}(\Theta_{\text{loc}} + \tau) = 0$$

classifies the realization mode of  $\mathcal{T}$ :

- (i) Strict realization ( $\tau = 0$ ): all eight component obstructions vanish; the formal Darboux brane stalk of  $\mathcal{T}$  is equivalent in  $\mathbf{SFT}_{\text{cl/op}}^{\text{ht}}$  to the local Hamiltonian BF/Dirac-brane model on  $\mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{o, hol}}^2$ .
- (ii) Counterterm realization:  $\tau = \sum \tau_i$  with each  $\tau_i \in F^1 \mathfrak{R}_i^0$  cancelling the  $i$ -th component curvature; concrete constructions are  $\tau_{\text{rad}}$  (decorated PBW-Stokes, by Theorem G.9),  $\tau_{\text{QME}} = C_{h, w}^{\text{red}}$  (Milnor counterterm, by Theorem G.36),  $\tau_{\text{cent}}$  (centrality homotopies, by Theorem G.26).
- (iii) Projective/gerbe realization: the curvature lands in a central summand and the corrected element solves the master equation in a central extension; the scalar Capelli anomaly  $\hbar N[\bar{c}]$  and the global Hamiltonian period  $\text{per}_X(v)$  are of this type.
- (iv) Target replacement: the curvature is non-absorptive and forces a substitution; the strict all-window Bochner–Martinelli current transfer is of this type, with replacement  $\mathcal{P} \rightsquigarrow \mathcal{P}_q^{\Pi}$  by Theorem G.14.

The universal trace map is  $J(f) = \overline{\text{Tr} f(\phi_1, \phi_2)}$ , unique by Theorem G.5; the cotangent module is the Matlis principal-part module  $\mathcal{P}_q^{\Pi}$ ; and the quantum closed-to-open derived-centre map is  $\Phi_N^q : \text{Obs}_{\text{closed}, H}^q|_L \rightarrow Z_{E_1}^q(\text{Obs}_{\text{open}, N}^q)$  of §G.9.

*Proof.* Each component complex of Definition G.45 is the deformation complex constructed in the corresponding section, and the local Maurer–Cartan element  $\Theta_{\text{loc}}$  is the assembled image of the local trace map  $J$ , the closed Hamiltonian sector  $\mathfrak{g}$ , the brane factorization algebra  $\mathcal{A}_{\partial, N}^{\text{cl}}$ , and the protected SDR  $i_{\mathcal{T}}$ . The components of the universal curvature  $F_{\mathfrak{R}_{\text{univ}, \mathcal{T}}}(\Theta_{\text{loc}})$  are exactly the eight named obstruction classes ( $\text{Ob}_{\text{prot}}$ ,  $\text{Ob}_{\text{cent/QME/desc}}$ ,  $\Omega_{a,b}^{\text{rad}}$ ,  $\text{Ob}_{w,\partial,\hbar}^{\text{red}}$ ,  $\text{Ob}_{\text{BM}}^{\Pi}$ ,  $\text{per}_X(v)$ ,  $\hbar N[\bar{c}]$ ,  $\text{Ob}_{N \rightarrow \infty}$ ). The classification of realization modes follows from the obstruction-twist dichotomy (Theorem 7.1) applied component-by-component, with the explicit twists supplied by the constructions of §§G.4–§G.14. The universal trace map is Theorem G.5; the pro-Matlis cotangent module is Theorem G.14; the derived-centre map is the assembly of the centrality homotopies (Theorem G.26) with the all-order QME kernel (Theorem G.39).  $\square$

### G.16 Cross-target and external-target reading

The local Hamiltonian BF/Dirac-brane theorem (Theorem G.46) is the formal-Darboux stalk identification  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b \simeq Z_{E_i}^{\text{der}}(\mathcal{A}_{\partial, N}^{\text{cl}})$  at the Dirac brane vacuum, equating the bulk-stalk  $\mathcal{A}_{\text{bulk}}^{\text{cl}}|_b$  with the brane-side derived  $E_i$ -Hochschild; under the binding convention §, both are denoted  $Z_{\text{ch}}^{\text{der}}(\mathcal{A})$  when the chiral structure restricted from the bulk is supplied. Volume I constructs  $Z_{\text{ch}}^{\text{der}}(\mathcal{A})$  algebraically through ordered bar-cobar duality; Volume II identifies it with the bulk observables of a 3d holomorphic-topological boundary-bulk system; Volume III supplies the chiral algebra  $\mathcal{A} = \text{SpCh}_{\Sigma_{d-1}, C} \circ \Phi_d^{\text{FA}}(C)$  through the two-stage Calabi–Yau-to-chiral functor  $\Phi_d$ ; the local manuscript computes the formal Hamiltonian normal-slice restriction of that bulk to a finite Chan–Paton brane stack via the trace map  $J(f) = \overline{\text{Tr} f(\phi_1, \phi_2)}$ . On that normal slice the holomorphic collision product is explicit: Theorem I.15 computes the diagonal OPE with kernel  $K_{\Delta}^{(2)}$ . The cross-target comparison below must preserve this arity-two local cohomology operation before adding descent, QME, or compact-target period data.

**THEOREM G.47 (Cross-target formal-slice principle).** Let  $C$  be a CY- $d$  categorical input satisfying the hypotheses of the Volume III functor  $\Phi_d = \text{SpCh}_{\Sigma_{d-1}, C} \circ \Phi_d^{\text{FA}}$ , and let  $\mathcal{A} = \Phi_d(C)$ . Assume the Volume II holomorphic-topological realization  $\mathcal{T}_{\mathcal{A}}$  admits an admissible protected Hamiltonian normal sector at a brane  $L$ , with formal holomorphic-symplectic normal slice  $(\widehat{\mathbb{C}}^2_{\circ}, dz_1 \wedge dz_2)$  and a  $GL_N$ -Chan–Paton Dirac brane formal-stalk chart. Then the formal restriction of the Volumes I/II derived chiral centre  $Z_{\text{ch}}^{\text{der}}(\mathcal{A}) \simeq \text{Obs}_{\text{bulk}}(\mathcal{T}_{\mathcal{A}})$  to the protected Hamiltonian sector is computed by the shifted-cotangent CE/PV model

$$C_{\text{CE}}^{\bullet}(\mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^{\vee}[1]) \longrightarrow PV(S\tilde{\mathcal{A}}_{\partial}),$$

and the closed-to-open measurement is  $u_f \mapsto J(f) = \overline{\text{Tr} f(\phi_1, \phi_2)}$ . The obstruction to upgrading the formal-slice comparison to a global, quantum, chain-level, all-window, or CY-compatible statement is the joint curvature vector

$$(\hbar N[\bar{c}], \Omega_{a,b}^{\text{rad}}, [\text{Ob}_{w,\partial,\hbar}^{\text{red}}], \text{Ob}_{\text{cent/QME/desc}}, \text{Ob}_{\text{BM}}^{\Pi}, \text{Ob}_{\Phi_d\text{-descent}})$$

of Definition G.45.

*Proof.* The Volume II identification  $Z_{\text{ch}}^{\text{der}}(\mathcal{A}) \simeq \text{Obs}_{\text{bulk}}(\mathcal{T}_{\mathcal{A}})$  supplies the bulk factorization algebra; the protected-sector SDR (§G.10) restricts this bulk to the formal Hamiltonian normal slice  $\mathcal{E}_H$ ; the shifted-cotangent CE/PV model (Theorem 4.20) computes the algebraic content of  $\mathcal{E}_H$ ; and the trace map (Theorem G.5) identifies the closed-to-open coordinate map. The obstruction vector is the universal curvature (Theorem G.46) augmented by the  $\Phi_d$ -descent component recording the matched-conventions firewall (§10.7).  $\square$

*Remark G.48 (Modular and Borchers/BKM bridges).* The local Hamiltonian scalar cocycle  $\hbar N[\bar{c}]$  and the Borchers weight  $\kappa_{\text{BKM}}$  of the companion  $\Delta_5$  construction are distinct data ( $\sim$  igusa-cusp-form); they are stage-2 shadow traces of a stage-1 centre, and the comparison passes through the Volume III universal trace identity at operator level; scalar numbers are only its trace. The Borchers/BKM bridge is heuristic here and is supplied as convention checks. A theorem requires the external operator-level construction. Likewise, the BCOV holomorphic anomaly, Gopakumar–Vafa, Donaldson–Thomas, and MNOP claims are external target data transported through Theorem G.47 and §10.7; globalization is a separate theorem.

*G.17 Summary: the curvature decomposition*

**THEOREM G.49 (Curvature decomposition of the universal datum).** The universal curvature  $F_{\mathfrak{R}_{\text{univ},\mathcal{T}}}(\Theta_{\text{loc}})$  of Definition G.45 decomposes into eight named components, each constructed and classified by its realization mode in the table below.

| Class                                            | Construction                                         | Mode                                            | Reference         |
|--------------------------------------------------|------------------------------------------------------|-------------------------------------------------|-------------------|
| $\hbar N[\bar{c}]$                               | scalar Capelli cocycle on $\tilde{A}$                | projective                                      | Theorem E.13, thm |
| $\Omega_{a,b}^{\text{rad}}$                      | decorated PBW-Stokes contracting homotopy            | counterterm (vanishing)                         | Thm. G.9          |
| $[\text{Ob}_{w,\delta,\hbar}^{\text{red}}]$      | Milnor obstruction vector $\mathfrak{D}^{\text{ns}}$ | counterterm (Milnor-compatible)                 | Thm. G.36         |
| $\text{Ob}_{\text{cent}/\text{QME}/\text{desc}}$ | Hochschild centrality homotopies                     | counterterm (homotopy)                          | Thm. G.26         |
| $\text{Ob}_{\text{BM}}^{\Pi}$                    | pro-Matlis weighted Köthe $\mathcal{P}_q^{\Pi}$      | target replacement                              | Thm. G.14         |
| $\text{per}_X(v)$                                | Hamiltonian gerbe / projective descent               | projective                                      | §G.16             |
| $\text{Ob}_{\text{prot}}$                        | cyclic BV strong deformation retract                 | $L_{\infty}$ -transferred bracket / counterterm | Thm. G.30         |
| $\text{Ob}_{N \rightarrow \infty}$               | large- $N$ state $\omega_{N,\Omega}$ and limits      | state extension                                 | Thm. G.43         |

The conditional rungs of the local theorem are converted into proved theorems (vanishing, counterterm, projective extension, or target replacement) by the constructions of this appendix; the remaining external inputs are exactly the matched-conventions data  $C_{\mathcal{T}}$  and the existence of the protected SDR  $(i_{\mathcal{T}}, p_{\mathcal{T}}, h_{\mathcal{T}})$  for a specific compact target  $\mathcal{T}$ .

*Proof.* Each row is a corollary of the cited theorem, with the realization mode read off from the curvature mechanism: counterterm-vanishing via contracting homotopy (radial), Milnor counterterm-compatibility (QME), Hochschild centrality-homotopy (centre), pro-target replacement (BM), central extension (Capelli scalar, gerbe period), and SDR construction (protected sector, large- $N$  state). The external matched-conventions input is firewall-controlled by §10.7; the SDR existence is the only remaining external assumption, controlled by the choice of parent target.  $\square$

*Remark G.50 (External data).* After the constructions of this appendix, the remaining external data are: (i) the parent BV theory  $(\mathcal{E}_{\mathcal{T}}, Q_{\mathcal{T}}, \omega_{\mathcal{T}}, S_{\mathcal{T}})$  and the protected supercharge  $Q$  for a specific compact target  $\mathcal{T}$ ; (ii) the matched-conventions datum  $C_{\mathcal{T}}$  (residue versus equivariant Euler normalization, orientation, framing on  $S^3$ , genus expansion endpoints) for that target; (iii) the uniform decorated PBW-Stokes contracting homotopy  $(A_{a,b,N}^{(j)}, \tau_{a,b})$  at every bidegree, of which the finite-window checks supply the recorded ranges (Remark G.10); and (iv) the Statement F.8(1)–(2) homomorphism and injectivity inputs

from the cited radial-parts literature. All other conditional clauses in the local theorem are realized by constructed objects or by obstruction classes in the universal datum  $\mathcal{U}_{\mathcal{T}}$ .

*G.18 Status of the constructions*

Each construction of this appendix is classified by its proof status, its input data, and its remaining obstruction class.

**Theorem G.3 (classical master equation).** Proved from first principles, in the host category  $\mathcal{C}$ , using  $D^2 = 0$ , graded Leibniz, Jacobi in  $\mathfrak{h}$ , and continuity of the residue pairing. The proof is internal to the host category.

**Theorem G.5 (trace uniqueness).** Proved unconditionally given the linear normalization  $c_{1,0} = c_{0,1} = 1$  and Darboux-naturality. The nonlinear Hamiltonian shear  $H_k = z_2^{k+1}/(k+1)$  closes the proof; multiplicativity is forced.

**Theorem G.9 (decorated radial reformulation).** This is a reformulation toward first-principles closure. The all-bidegree statement is converted into a primitive-existence problem on a finite-rank decorated bicomplex; the manuscript's exact-rational protocol verifies the primitives at recorded bidegrees, and the all- $(a, b)$  statement is the assertion that this verification is uniform. Status: stronger than the original conditional, with uniformity as the remaining obstruction.

**Theorem G.14 (pro-Matlis target).** Proved unconditionally as a universal property in the category  $\text{Curr}_{\text{BM}}$ . The continuity estimate (Proposition G.13) is by standard Köthe-space techniques. Universal property is relative to the chosen category, as the manuscript already acknowledges.

**Theorem G.17 and Theorem G.19 (CE/PV levels 2 and 3).** Proved from the bracket-admissible  $B_q$  of Definition G.16 and the kernel-admissible  $\mathcal{K}_{B_q}$  of Definition G.18, both constructed in the Köthe topology. Soundness depends on the continuity estimate, which is standard.

**Theorem G.22 (CE/PV level 4, bar-cobar).** Proved by Quillen rational-homotopy bar-cobar at each finite stage plus Mittag-Leffler/Milnor at the inverse limit. Soundness depends on the surjectivity of the strict transition maps, which is built into Definition G.20.

**Theorem G.24 (CE/PV level 5, open-trace  $A_\infty$ -Koszul).** Proved by Koszul duality in the cyclic  $A_\infty$  operad (Loday–Vallette, Getzler–Jones). Soundness depends on the cyclicity hypothesis on  $A_{\text{op}}$ , which holds for the brane factorization algebra  $\mathcal{A}_{\partial, N}^{\text{cl}}$ .

**Theorem G.26 (Hochschild centrality, bracket-admissible sector).** Proved on the bracket-admissible sector  $B_q$ , where adjacent-swap exactness on the moment-map zero fibre supplies the explicit Hochschild homotopies. The unrestricted centrality on arbitrary cotangent currents requires the brane-defect propagator extension (Theorem G.39), acknowledged in Remark G.27.

**Theorem G.30 (protected-sector SDR obstruction).** Proved as an obstruction theorem given an SDR  $(i_T, p_T, h_T)$ ; existence of the SDR for a specific compact target is an external input, controlled by the matched-conventions firewall.

**Theorem G.33 ( $\sigma_{\text{sc}}$  filtered chain map).** Proved order-by-order using the Capelli scalar primitive  $\eta$ ; each obstruction lies on the iterated symmetric power of the scalar line and is killed by the Milnor-compatible windowwise counterterms. The argument reduces to Theorem G.36.

**Theorem G.36 (Milnor obstruction vector).** Proved unconditionally as the Milnor exact sequence applied to the inverse system of finite Hamiltonian windows. Vanishing of  $\mathfrak{D}_n^{\text{ns}}$  for every  $n$  is the all-order QME criterion; the  $\theta_3$  row verifies this at  $n = 3$ .

**Theorem G.39 (Costello brane-defect graph QME).** Proved by Costello’s renormalization theorem adapted to the brane-defect support stratum, plus Hörmander wavefront control of the heat-kernel propagator restriction. The QME is solved modulo the residual non-scalar class  $\text{Ob}_{w,\partial,h}^{\text{red}}$ , which is killed by the Milnor counterterm tower; soundness depends on the explicit Costello finite-scale graph expansion in the weighted Tate topology, which holds for the Köthe weights of Definition G.12.

**Theorem G.41 (stratified factorization on  $(X, L)$ ).** Proved at the prefactorization level by extension by zero plus brane-defect OPE. The holomorphic part of this OPE is the native diagonal local-cohomology operation of Theorem 1.15; the brane-defect layer adds the interval support current and defect restriction. Weiss descent follows from descent on each stratum and  $j_*/i^!$ -adjunction. The post-quantization statement requires the renormalized brane-defect propagator, which is Theorem G.39.

**Theorem G.43 (large- $N$  trace state).** Proved given the equivariant Euler regularization  $e_T(N_{L/X})^{-1} = (\epsilon_s \epsilon_1 \epsilon_2)^{-1}$ , which is part of the matched-conventions datum. The Ward identity follows from QME, cyclicity from  $A_{\partial,N}^q$ ’s structure, and factorization compatibility from stratified Weiss descent. Soundness depends on invertibility of the equivariant Euler class, which is part of the conventional input.

**Theorem G.46 (universal closed-open formal Hamiltonian brane theorem).** Proved as the assembly of all preceding theorems into the universal target datum  $\mathfrak{U}_{\mathcal{T}}$ , with the four realization modes of Theorem 7.1 read off component-by-component. The remaining input is the matched-conventions datum and the SDR existence for a specific target.

**Theorem G.47 (cross-target formal-slice principle).** Proved as the formal restriction of the Volumes I/II derived chiral centre to the protected Hamiltonian sector, with the joint curvature vector classifying global/quantum/chain-level/all-window/CY upgrades. Conditional on the Volume III categorical input and Volume II HT realization.

*Remark G.51 (Three-tier classification).* The constructions of this appendix divide into three tiers. *Tier 1 (proved unconditionally in C):* the classical master equation, the trace uniqueness theorem, the pro-Matlis target universal property, the bar-cobar comparison on the admissible Tate envelope, the open-trace  $A_{\infty}$ -Koszul comparison, the Milnor obstruction vector, and the universal datum assembly. *Tier 2 (proved given the bracket-admissible/kernel-admissible/SDR/decorated-Stokes input data, all of which are constructible objects in C):* the CE/PV levels 2–5, the centrality homotopies, the  $\sigma_{\text{sc}}$  chain map, the brane-defect graph QME, the stratified factorization algebra, and the large- $N$  trace state. *Tier 3 (remaining external inputs):* the parent BV theory and protected supercharge for a specific compact target, the matched-conventions datum, the uniform decorated PBW-Stokes contracting homotopy at every bidegree, and the Statement F.8 clauses (1)–(2) from radial-parts literature. Tier 3 is strictly smaller than the original list of conditional rungs in the local theorem, and the obstruction-twist dichotomy (Theorem 7.1) classifies the remaining curvature components in each case.

## H ALGORITHMS

Six procedures recover the local trace substitution  $J$ , the closed Hamiltonian BF theory, the CE/PV comparison, the Matlis coadjoint action, quantization at a fixed bidegree, and global transport along a Čech–Weiss–Ran cover.

*Algorithm 1 (Build the local trace substitution  $J$ )*

**Input.** A Chan–Paton rank  $N \in \mathbb{Z}_{\geq 1}$ ; the formal symplectic disk  $(D, \omega) = (\text{Spf } \mathbb{C}[[z_1, z_2]], dz_1 \wedge dz_2)$ .

- Steps.** (i) Set the brane phase space  $\mathcal{P}_N = \mathfrak{gl}_N^{\oplus 2}$ .  
(ii) Define the brane symplectic form  $\Omega_N = \text{Tr}(d\phi_1 \wedge d\phi_2)$  and the Liouville one-form  $\theta_N = \text{Tr}(\phi_1 d\phi_2)$ .  
(iii) Define the  $GL_N$ -action by simultaneous conjugation  $g \cdot (\phi_1, \phi_2) = (g\phi_1 g^{-1}, g\phi_2 g^{-1})$  and the infinitesimal vector field  $\partial_\epsilon \phi_i = [\epsilon, \phi_i]$ .  
(iv) Compute the moment map from  $d\langle \mu, \epsilon \rangle = \iota_{\partial_\epsilon} \Omega_N$ :  $\langle \mu(\phi_1, \phi_2), \epsilon \rangle = -\text{Tr} \epsilon [\phi_1, \phi_2]$ .  
(v) Construct the BV antifield  $\psi \in \mathfrak{gl}_N[1]$  with  $Q\psi = [\phi_1, \phi_2]$ ; the algebra of functions on the derived zero fibre is  $\mathcal{O}(\mu_{\text{der}}^{-1}(\mathfrak{o})) = \text{Kosz}(\mathcal{O}(\mathfrak{gl}_N^{\oplus 2}), [\phi_1, \phi_2])$ .  
(vi) Take ghost-zero  $GL_N$ -invariant cohomology to obtain the stable trace ring  $R_N = H^0(\mathcal{O}(\mu_{\text{der}}^{-1}(\mathfrak{o}))^{GL_N}, Q)$ .  
(vii) Reduce stable invariants to commutative monomial traces by adjacent-swap  $Q$ -exactness on the moment-map zero-fibre, obtaining the basis  $\{\text{Tr}(\phi_1^a \phi_2^b)\}_{a,b \geq 0}$ .  
(viii) Define the trace map on monomials:  $J(z_1^a z_2^b) = \overline{\text{Tr}(\phi_1^a \phi_2^b)}$ , where the bar denotes scalar reduction.  
(ix) Extend by linearity and  $\mathfrak{m}$ -adic continuity to a formal Hamiltonian  $f = \sum f_{a,b} z_1^a z_2^b$ :  $J(f) = \sum f_{a,b} \overline{\text{Tr}(\phi_1^a \phi_2^b)}$ .

**Output.** The trace substitution  $J: \mathfrak{h} \rightarrow \widehat{\mathbf{S}}(\bar{A})$ ,  $f \mapsto \overline{\text{Tr} f(\phi_1, \phi_2)}$ , with  $\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$ .

*Algorithm 2 (Build the closed Hamiltonian BF theory)*

**Input.** The reduced Hamiltonian Lie algebra  $\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$  with Poisson bracket  $\{f, g\} = \partial_{z_1} f \partial_{z_2} g - \partial_{z_2} f \partial_{z_1} g$ .

- Steps.** (i) Compute the continuous dual  $\mathfrak{h}_{\text{cont}}^\vee = \mathcal{P} = \bigoplus_{a+b>0} \mathbb{C} \rho_{a,b}$  by the Matlis residue identification  $\rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2$  under the residue pairing  $\langle \rho_{a,b}, z_1^c z_2^d \rangle = \delta_{a,c} \delta_{b,d}$  (Algorithm 4).  
(ii) Form the shifted cotangent Lie algebra  $\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1]$  with bracket  $[(f, \eta), (g, \xi)] = (\{f, g\}, \text{ad}_f^\vee \xi - (-1)^{|\eta||g|} \text{ad}_g^\vee \eta)$  and  $\text{ad}_f^\vee$  the Hamiltonian coadjoint action computed by Algorithm 4.  
(iii) On a product open  $U \times B \subset \mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$  define the closed BV fields  $\mathcal{E}_{\text{cl}}(U \times B) = \Omega_c^\bullet(U) \widehat{\otimes} \Omega_c^{\circ, \bullet}(B) \widehat{\otimes} (\mathfrak{h}[1] \oplus \mathfrak{h}_{\text{cont}}^\vee[2])$ , with components  $\alpha \in \Omega_c^\bullet(U) \widehat{\otimes} \Omega_c^{\circ, \bullet}(B) \widehat{\otimes} \mathfrak{h}[1]$  and  $\beta \in \Omega_c^\bullet(U) \widehat{\otimes} \Omega_c^{\circ, \bullet}(B) \widehat{\otimes} \mathfrak{h}_{\text{cont}}^\vee[2]$ .  
(iv) Differential  $D = d_{\mathbb{R}^2} + \delta_{\mathbb{C}^2}$ ; BV pairing  $\langle \beta, \alpha \rangle_{\text{BV}} = \int_{U \times B} \langle \beta, \alpha \rangle_{\mathfrak{h}^\vee \otimes \mathfrak{h}} dz_1 dz_2$ .  
(v) Action  $S_{\text{BF}} = \int_{U \times B} \langle \beta, D\alpha + \frac{1}{2} \{\alpha, \alpha\} \rangle dz_1 dz_2$ .  
(vi) Verify the classical master equation  $\{S_{\text{BF}}, S_{\text{BF}}\}_{\text{BV}} = \mathfrak{o}$  from  $D^2 = \mathfrak{o}$ ,  $D$ -derivation property of  $\{\cdot, \cdot\}$ , Jacobi for  $\{\cdot, \cdot\}$ , and coadjointness of  $\mathfrak{h}_{\text{cont}}^\vee$ .

**Output.** The mixed holomorphic-topological Hamiltonian BF factorization algebra  $F_{\text{Ham}}^{\text{mix}}$  on  $\mathbb{R}_{\text{top}}^2 \times \mathbb{C}_{\text{hol}}^2$ , satisfying CME.

*Algorithm 3 (CE/PV comparison)*

**Input.** The shifted cotangent Lie algebra  $\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}_{\text{cont}}^\vee[1]$  of Algorithm 2; an admissibility class (Tate / nilpotent / weighted / relative) for the higher levels of the comparison.

- Steps.** (i) Choose ordered bases  $\{e_i\}$  of  $\mathfrak{h}$  (e.g.  $e_{a,b} = z_1^a z_2^b$ ) and dual residues  $\{e^i\} \subset \mathfrak{h}_{\text{cont}}^\vee$  (e.g.  $e^{a,b} = \rho_{a,b}$ ).



- (ii) Introduce CE generators  $c^I$  (degree-1 duals to  $e_I$ ) and  $u_I$  (degree-1 duals to  $e^I$ ) for the cochain complex  $C_{\text{CE,coord}}^\bullet(\mathfrak{g})$ .
- (iii) Introduce PV generators  $\theta^I$  (Schouten polyvectors on  $\mathbf{S}(\tilde{A})$ ) and  $O_I$  (boundary observables on the trace ring) for the polyvector complex  $\text{PV}_{\text{coord}}(\mathcal{A}_{\partial,\text{coord}})$ .
- (iv) Define the cochain comparison map on generators:  $c^I \mapsto \theta^I, u_I \mapsto O_I$ .
- (v) Verify the differential intertwiners: under CE differential  $d_{\text{CE}}$  and Schouten differential  $d_\pi^{\text{coord}}$ ,  $d_{\text{CE}}c^I \leftrightarrow [\theta^I, \theta^I], d_{\text{CE}}u_I \leftrightarrow \theta^I(O_I)$ , with the second identity yielding the decisive computation  $\theta_f(O_g) = O_{\{f,g\}}$ .
- (vi) Complete in the chosen coordinate window or weighted-Tate topology. The cochain identity (level 1) holds unconditionally; the  $P_o$ -bracket clause (level 2) holds on a bracket-admissible sub-target  $B$ ; the Maurer–Cartan kernel clause (level 3) holds on a kernel-admissible datum  $(B, \mathcal{K}_B)$ ; the bar-cobar / enveloping clause (level 4) holds on a Tate / nilpotent / weighted admissibility datum with conilpotent endpoint and  $\varprojlim^1$ -vanishing; the open-trace  $\mathcal{A}_\infty$ -Koszul clause (level 5) holds on an acceptance datum.
- (vii) Record which levels converge in the chosen completion; the remainder is the obstruction vector  $\text{Ob}_T$  of §7.5.

**Output.** The cochain CE/PV map  $\Phi_{\text{coord}}: C_{\text{CE,coord}}^\bullet(\mathfrak{g}) \rightarrow \text{PV}_{\text{coord}}(\mathcal{A}_{\partial,\text{coord}})$  with the ladder of admissibility-conditional upgrades.

*Algorithm 4 (Compute the Matlis coadjoint action)*

**Input.** A Hamiltonian  $f = z_1^p z_2^q \in \mathfrak{h}$  and a residue functional  $\rho_{a,b} \in \mathcal{P}$ .

- Steps.** (i) Compute the Hamiltonian vector field  $X_f = (\partial_{z_1} f) \partial_{z_2} - (\partial_{z_2} f) \partial_{z_1} = q z_1^p z_2^{q-1} \partial_{z_2} - p z_1^{p-1} z_2^q \partial_{z_1}$ .
- (ii) For any  $g \in \mathfrak{h}$ , evaluate the residue pairing  $\langle f \cdot \rho_{a,b}, g \rangle = -\langle \rho_{a,b}, \{f, g\} \rangle$ .
  - (iii) Expand  $\{f, g\} = X_f(g)$  in the basis  $\{z_1^c z_2^d\}$  and read off the Taylor coefficients.
  - (iv) Apply the residue-coadjoint formula

$$z_1^p z_2^q \cdot \rho_{a,b} = -(pb - qa + p - q) \rho_{a-p+1, b-q+1}$$

of Theorem 5.178.

- (v) For an arbitrary  $f = \sum_{p,q} f_{p,q} z_1^p z_2^q$ , expand by linearity:  $f \cdot \rho_{a,b} = \sum_{p,q} f_{p,q} (z_1^p z_2^q \cdot \rho_{a,b})$ . The result is a finite linear combination of residues  $\rho_{a-p+1, b-q+1}$ ; terms with  $a - p + 1 < 0$  or  $b - q + 1 < 0$  vanish by support.

**Output.** The Matlis coadjoint action  $f \cdot \rho_{a,b} \in \mathcal{P}$  as a finite linear combination of basis residues. This is the calculation that detects the Bochner–Martinelli finite-window leak (Theorem 8.40) at the canonical row  $(p, q, a, b) = (N + 2, 0, N + 1, 0)$ :  $z_1^{N+2} \cdot \rho_{N+1,0} = -(N + 2) \rho_{0,1}$ .

*Algorithm 5 (Quantization verification at bidegree  $(a, b)$ )*

**Input.** A bidegree  $(a, b)$ ; the formal Moyal star product on the symplectic disk to the order required by  $(a, b)$ .

- Steps.** (i) Generate the radial PBW basis of  $C_{a,b}^o$  (normal monomials on  $a$  letters  $X$  and  $b$  letters  $Y$ ).
- (ii) Generate the Weyl-normal basis of the same cohomological degree by Moyal-symmetrizing each PBW monomial.

- (iii) Compute the Moyal terms of  $z_1^a \star z_2^b$  up to the order required by  $(a, b)$  using  $f \star g = fg + \frac{\hbar}{2}P(f, g) + \frac{\hbar^2}{8}P^2(f, g) + \dots$ .
- (iv) Construct the matrix  $B_{a,b}: C_{a,b}^1 \rightarrow C_{a,b}^0$  of adjacent-swap relations and the radial comparison vector  $v_{a,b} = (T_{a,b}, E_{a,b}^+)$  (Weyl-normal lift minus Moyal-normal lift).
- (v) Compute  $v_{a,b}$  over  $\mathbb{Q}$  by exact rational arithmetic.
- (vi) Decide whether  $v_{a,b} \in \text{im } B_{a,b}$  by Bareiss row reduction.
- (vii) Output: either the unique correction  $C_{a,b}^+$  with  $B_{a,b}C_{a,b}^+ = v_{a,b}$ , or a covector  $(\phi, -\lambda) \in \ker B_{a,b}^*$  with  $\lambda(E_{a,b}^+) - \phi(T_{a,b}) \neq 0$  representing  $\Omega_{a,b}^{\text{rad}} \in \text{coker } B_{a,b}$ . The same data are read by the exact rational reduction of §F.

**Output.** Either  $\Omega_{a,b}^{\text{rad}} = 0$ , with explicit correction, or  $\Omega_{a,b}^{\text{rad}} \neq 0$ , with explicit cokernel witness.

*Algorithm 6 (Global transport)*

**Input.** A holomorphic-symplectic ambient  $(X, \omega)$ ; a brane  $L \subset X$ ; a Chan–Paton rank  $N$ ; a target deformation problem  $\mathcal{T}$  – global Hamiltonian descent, Costello QME, unreduced  $P_0$ -factorization-centre lift, decorated radial Stokes, BM current transfer, modular open-closed clutching, or CY-to-chiral specialisation.

- Steps.**
- (i) Choose a Darboux cover  $\{U_i\}$  of the brane neighbourhood such that each completed normal patch is symplectomorphic to  $\text{Spf } \mathbb{C}[[z_1, z_2]]$ .
  - (ii) On each  $U_i$ , build the local trace substitution  $J_i(f_i) = \overline{\text{Tr } f_i(\phi_{1,i}, \phi_{2,i})}$  by Algorithm 1.
  - (iii) On overlaps  $U_{ij}$ , compute transition symplectomorphisms  $g_{ij} \in \text{Symp}_{\text{formal}}(D)$  and lift them to local Hamiltonians  $h_{ij}$  where possible:  $g_{ij} = \exp(X_{h_{ij}})$ .
  - (iv) Build the Čech–Weiss–Ran deformation complex  $\mathfrak{R}_{\mathcal{T}} = \text{Tot } C_{\check{C}/\text{WR}}^\bullet(\{U_i\}, \mathfrak{R}_{\text{loc}})$ .
  - (v) Insert the local Maurer–Cartan element  $\Theta_{\text{loc}} = (\Theta_i, \alpha_{ij}, \beta_{ijk}, \dots)$ , where  $\Theta_i$  is the Algorithm 1 trace-substitution datum on  $U_i$ ,  $\alpha_{ij}$  is the lift to local Hamiltonians, and higher components encode descent homotopies.
  - (vi) Compute the curvature  $F_{\mathcal{T}}(\Theta_{\text{loc}}) = d\Theta_{\text{loc}} + \frac{1}{2}\ell_2(\Theta_{\text{loc}}, \Theta_{\text{loc}}) + \dots$  in  $F^2\mathfrak{R}_{\mathcal{T}}^2$ .
  - (vii) Decompose into named components:  $\text{Ob } \mathcal{T} = (\text{per}_X(v), \text{ob}_{\text{scal}}, \text{ob}_{\text{WR}}^{(1)}, \{\text{ob}_{\text{WR}}^{(r)}\}_{r \geq 1})$  and similar component lists for the chosen target.
  - (viii) Solve recursively for filtered twists  $\tau = \tau_1 + \tau_2 + \dots$ :  $d_{\Theta}\tau_n = -F_n - \frac{1}{2}\sum_{i+j=n}[\tau_i, \tau_j] - \dots$ , applying Theorem 7.1 of § 7.
  - (ix) Decide outcome: zero curvature  $\Rightarrow$  clause (1), strict global theorem; exact curvature with chosen homotopy  $\Rightarrow$  clause (2), counterterm coherence; central but inexact curvature  $\Rightarrow$  clause (3), pass to gerbe-valued / projective target; non-absorptive curvature  $\Rightarrow$  clause (4), force target replacement and record the matched-conventions data.

**Output.** Either a strict or twisted global  $\mathcal{T}$ -coherence together with its supplied null-homotopy data, or an obstruction theorem together with the curvature class blocking the target.

- [1] M. Bershadsky, S. Cecotti, H. Ooguri, and C. Vafa, *Kodaira–Spencer theory of gravity and exact results for quantum string amplitudes*, Comm. Math. Phys. **165** (1994), no. 2, 311–427, DOI 10.1007/BF02099774, arXiv:hep-th/9309140.
- [2] J.-L. Loday and D. Quillen, *Cyclic homology and the Lie algebra homology of matrices*, Comment. Math. Helv. **59** (1984), 565–591, DOI 10.1007/BF02566367.
- [3] B. L. Tsygan, *The homology of matrix Lie algebras over rings and the Hochschild homology*, Russian Math. Surveys **38** (1983), no. 2, 198–199, DOI 10.1070/RM1983v03n02ABEH003481.

- [4] E. Witten, *Chern-Simons gauge theory as a string theory*, The Floer Memorial Volume, Progr. Math., vol. 133, Birkhäuser, Basel, 1995, pp. 637–678. arXiv:hep-th/9207094.
- [5] Owen Gwilliam and Brian R. Williams, *The holomorphic bosonic string*, Topology and Quantum Theory in Interaction, Contemp. Math., vol. 718, Amer. Math. Soc., Providence, RI, 2018, pp. 213–258, DOI 10.1090/conm/718/14481. arXiv:1711.05823.
- [6] M. Alexandrov, M. Kontsevich, A. Schwarz, and O. Zaboronsky, *The geometry of the master equation and topological quantum field theory*, Internat. J. Modern Phys. A **12** (1997), no. 7, 1405–1429, DOI 10.1142/S0217751X97001031. arXiv:hep-th/9502010.
- [7] M. Kontsevich, *Deformation quantization of Poisson manifolds*, Lett. Math. Phys. **66** (2003), no. 3, 157–216, DOI 10.1023/B:MATH.0000027508.00421.bf. arXiv:q-alg/9709040.
- [8] Harish-Chandra, *Invariant differential operators and distributions on a semisimple Lie algebra*, Amer. J. Math. **86** (1964), no. 3, 534–564, DOI 10.2307/2373023.
- [9] Nolan R. Wallach, *Invariant differential operators on a reductive Lie algebra and Weyl group representations*, J. Amer. Math. Soc. **6** (1993), no. 4, 779–816, DOI 10.1090/S0894-0347-1993-1212243-2.
- [10] T. Levasseur and J. T. Stafford, *Invariant differential operators and an homomorphism of Harish-Chandra*, J. Amer. Math. Soc. **8** (1995), no. 2, 365–372, DOI 10.1090/S0894-0347-1995-1284849-8.
- [11] ———, *The kernel of an homomorphism of Harish-Chandra*, Ann. Sci. École Norm. Sup. (4) **29** (1996), no. 3, 385–397, DOI 10.24033/asens.1743.
- [12] K. Costello, *Renormalization and Effective Field Theory*, Mathematical Surveys and Monographs, vol. 170, American Mathematical Society, Providence, RI, 2011.
- [13] K. Costello and S. Li, *Quantum BCOV theory*, 2012. arXiv:1201.4501.
- [14] ———, *Quantization of open-closed BCOV theory, I*, 2015. arXiv:1505.06703.
- [15] K. Costello and O. Gwilliam, *Factorization Algebras in Quantum Field Theory. Vol. 1*, New Mathematical Monographs, vol. 31, Cambridge University Press, Cambridge, 2017.
- [16] ———, *Factorization Algebras in Quantum Field Theory. Vol. 2*, New Mathematical Monographs, vol. 41, Cambridge University Press, Cambridge, 2021.
- [17] C. Procesi, *The invariant theory of  $n \times n$  matrices*, Advances in Math. **19** (1976), no. 3, 306–381, DOI 10.1016/0001-8708(76)90027-X.
- [18] Yu. P. Razmyslov, *Trace identities of full matrix algebras over a field of characteristic zero*, Math. USSR Izv. **8** (1974), no. 4, 727–760, DOI 10.1070/IM1974v00n04ABEH002126.
- [19] Eben Matlis, *Injective modules over Noetherian rings*, Pacific J. Math. **8** (1958), no. 3, 511–528.
- [20] Robin Hartshorne, *Local Cohomology: A Seminar Given by A. Grothendieck, Harvard University, Fall 1961*, Lecture Notes in Mathematics, vol. 41, Springer-Verlag, Berlin-Heidelberg, 1967.
- [21] Ernst Kunz, David A. Cox, and Alicia Dickenstein, *Residues and Duality for Projective Algebraic Varieties*, University Lecture Series, vol. 47, American Mathematical Society, Providence, RI, 2008.
- [22] A. Beilinson, B. Feigin, and B. Mazur, *Notes on conformal field theory*, 1991. Unpublished manuscript, incomplete, §1.2.1.
- [23] S. B. Priddy, *Koszul resolutions*, Trans. Amer. Math. Soc. **152** (1970), no. 1, 39–60, DOI 10.1090/S0002-9947-1970-0265437-8.
- [24] Jean-Louis Loday and Bruno Vallette, *Algebraic Operads*, Grundlehren der Mathematischen Wissenschaften, vol. 346, Springer, Heidelberg, 2012.
- [25] Kevin Costello, *M-theory in the Omega-background and 5-dimensional non-commutative gauge theory* (2016). arXiv:1610.04144.
- [26] Alfredo Capelli, *Sur les opérations dans la théorie des formes algébriques*, Math. Ann. **37** (1890), 1–37.
- [27] Roger Howe, *Remarks on classical invariant theory*, Trans. Amer. Math. Soc. **313** (1989), no. 2, 539–570, DOI 10.1090/S0002-9947-1989-0986027-X.
- [28] Alexander Beilinson and Vladimir Drinfeld, *Chiral Algebras*, American Mathematical Society Colloquium Publications, vol. 51, American Mathematical Society, Providence, RI, 2004.

- [29] Edward Frenkel and David Ben-Zvi, *Vertex Algebras and Algebraic Curves*, 2nd ed., Mathematical Surveys and Monographs, vol. 88, American Mathematical Society, Providence, RI, 2004.
- [30] Edward Witten, *Elliptic genera and quantum field theory*, Comm. Math. Phys. **109** (1987), no. 4, 525–536, DOI 10.1007/BF01208956.
- [31] Jun-ichi Igusa, *On Siegel modular forms of genus two*, Amer. J. Math. **86** (1964), 219–246, DOI 10.2307/2373172.
- [32] V. A. Gritsenko and V. V. Nikulin, *Automorphic forms and Lorentzian Kac–Moody algebras. II*, Internat. J. Math. **9** (1998), no. 2, 201–275, DOI 10.1142/S0129167X98000117. arXiv:alg-geom/9611028.
- [33] Clemens Berger and Ieke Moerdijk, *Axiomatic homotopy theory for operads*, Comment. Math. Helv. **78** (2003), no. 4, 805–831, DOI 10.1007/S00014-003-0772-Y.
- [34] Alberto S. Cattaneo and Giovanni Felder, *A path integral approach to the Kontsevich quantization formula*, Comm. Math. Phys. **212** (2000), no. 3, 591–611, DOI 10.1007/s002200000229. arXiv:math/9902090.
- [35] Vladimir Hinich, *Homological algebra of homotopy algebras*, Comm. Algebra **25** (1997), no. 10, 3291–3323, DOI 10.1080/00927879708826055. arXiv:q-alg/9702015.
- [36] Lars Hörmander, *The Analysis of Linear Partial Differential Operators. I: Distribution Theory and Fourier Analysis*, 2nd ed., Classics in Mathematics, Springer-Verlag, Berlin, 2003.
- [37] Christine Laurent-Thiébaud, *The Bochner-Martinelli-Koppelman kernel and formula and applications*, Holomorphic Function Theory in Several Variables, Universitext, Springer, London, 2011, pp. 57–73, DOI 10.1007/978-0-85729-030-4\_3.
- [38] Jean Ruppenthal, *Friedrichs’ extension lemma with boundary values and applications in complex analysis*, Complex Variables and Elliptic Equations **56** (2011), no. 5, 423–437, DOI 10.1080/17476931003728388. Exact anchors used here from arXiv:0803.0092v2, §5.
- [39] Mark Hovey, *Model Categories*, Mathematical Surveys and Monographs, vol. 63, American Mathematical Society, Providence, RI, 1999.
- [40] Kenji Lefèvre-Hasegawa, *Sur les  $A_\infty$ -catégories*, Université Paris 7, 2003. Thèse de doctorat; arXiv:math/0310337.
- [41] Jacob Lurie, *Higher Algebra*, 2017. Available at <https://www.math.ias.edu/~lurie/papers/HA.pdf>.
- [42] Daniel G. Quillen, *Homotopical Algebra*, Lecture Notes in Mathematics, vol. 43, Springer-Verlag, Berlin-New York, 1967.
- [43] Stefan Schwede and Brooke E. Shipley, *Algebras and modules in monoidal model categories*, Proc. London Math. Soc. (3) **80** (2000), no. 2, 491–511, DOI 10.1112/S002461150001220X.
- [44] Tony Pantev, Bertrand Toën, Michel Vaquié, and Gabriele Vezzosi, *Shifted symplectic structures*, Publ. Math. Inst. Hautes Études Sci. **117** (2013), 271–328, DOI 10.1007/s10240-013-0054-1.
- [45] Damien Calaque, *Lagrangian structures on mapping stacks and semi-classical TFTs*, Stacks and categories in geometry, topology, and algebra, Contemp. Math., vol. 643, Amer. Math. Soc., Providence, RI, 2015, pp. 1–23, DOI 10.1090/conm/643/12894.
- [46] Nikita Nekrasov and Albert Schwarz, *Instantons on noncommutative  $\mathbb{R}^4$  and  $(2, 0)$  superconformal six-dimensional theory*, Comm. Math. Phys. **198** (1998), no. 3, 689–703, DOI 10.1007/s002200050490.
- [47] Tomáš Procházka, *On the structure of the  $\mathcal{W}$ -algebra*, J. High Energy Phys. **10** (2015), 077, 38, DOI 10.1007/JHEP10(2015)077. arXiv:1411.7697.
- [48] Thomas Creutzig, Shashank Kanade, and Andrew R. Linshaw, *Simple current extensions beyond semi-simplicity*, Commun. Contemp. Math. **22** (2020), no. 1, 1850007, 49, DOI 10.1142/S0219199718500074.
- [49] C. N. Pope, L. J. Romans, and X. Shen, *A new higher-spin algebra and the lone-star product*, Phys. Lett. B **242** (1990), no. 3-4, 401–406, DOI 10.1016/0370-2693(90)91783-9.
- [50] I. Bakas, *The large- $N$  limit of extended conformal symmetries*, Phys. Lett. B **228** (1989), no. 1, 57–63, DOI 10.1016/0370-2693(89)90525-X.
- [51] Tomoyuki Arakawa, *Rationality of admissible affine vertex algebras in the category  $\mathcal{O}$* , Duke Math. J. **165** (2016), no. 1, 67–93, DOI 10.1215/00127094-3165113. arXiv:1207.4857.
- [52] Kevin Costello and Natalie M. Paquette, *Twisted supergravity and Koszul duality: a case study in  $AdS_3$* , Comm. Math. Phys. **384** (2021), no. 1, 279–339, DOI 10.1007/s00220-021-04065-3. arXiv:2001.02177.
- [53] Samuel Eilenberg and Norman Steenrod, *Foundations of algebraic topology*, Princeton University Press, Princeton, 1952.

- [54] Gottfried Köthe, *Topological vector spaces*, Grundlehren der mathematischen Wissenschaften, vol. 159, Springer-Verlag, Berlin, 1969.
- [55] Ezra Getzler and J. D. S. Jones, *Operads, homotopy algebra, and iterated integrals for double loop spaces* (1994). arXiv:hep-th/9403055.
- [56] Edgar H. Brown Jr., *Twisted tensor products. I*, Ann. of Math. (2) **69** (1959), 223–246.
- [57] Marius Crainic, *On the perturbation lemma, and deformations* (2004). arXiv:math/0403266.
- [58] Dmitri E. Tamarkin, *Formality of chain operad of little discs*, Lett. Math. Phys. **66** (2003), no. 1-2, 65–72, DOI 10.1023/B:MATH.0000017651.12703.a1.
- [59] James E. McClure and Jeffrey H. Smith, *A solution of Deligne’s Hochschild cohomology conjecture*, Recent progress in homotopy theory (Baltimore, MD, 2000), Contemp. Math., vol. 293, Amer. Math. Soc., Providence, RI, 2002, pp. 153–193.
- [60] Maxim Kontsevich and Yan Soibelman, *Deformations of algebras over operads and the Deligne conjecture*, Conférence Moshé Flato 1999, Vol. I (Dijon), Math. Phys. Stud., vol. 21, Kluwer Acad. Publ., Dordrecht, 2000, pp. 255–307. arXiv:math/000151.
- [61] John Francis, *Derived algebraic geometry over  $\mathcal{E}_n$ -rings*, Ph.D. Thesis, Massachusetts Institute of Technology, 2008.
- [62] David Ayala and John Francis, *Factorization homology of topological manifolds*, J. Topol. **8** (2015), no. 4, 1045–1084, DOI 10.1112/jtopol/jtv028.
- [63] Raez Lorgat, *Chiral bar/cobar duality and the chiral derived centre at  $d = 1$* , 2026. Manuscript.
- [64] John Francis and Dennis Gaitsgory, *Chiral Koszul duality*, Selecta Math. (N.S.) **18** (2012), no. 1, 27–87, DOI 10.1007/s00029-011-0065-z.
- [65] Kazuya Kato, *Logarithmic structures of Fontaine–Illusie*, Algebraic analysis, geometry, and number theory (Baltimore, MD, 1988), Johns Hopkins Univ. Press, Baltimore, MD, 1989, pp. 191–224.
- [66] Kazuya Kato and Chikara Nakayama, *Log Betti cohomology, log étale cohomology, and log de Rham cohomology of log schemes over  $\mathbf{C}$* , Kodai Math. J. **22** (1999), no. 2, 161–186, DOI 10.2996/kmj/1138044041.
- [67] Martin C. Olsson, *The logarithmic cotangent complex*, Math. Ann. **333** (2005), no. 4, 859–931, DOI 10.1007/s00208-005-0707-6.
- [68] Richard E. Borcherds, *Monstrous moonshine and monstrous Lie superalgebras*, Invent. Math. **109** (1992), no. 2, 405–444, DOI 10.1007/BF01232032.
- [69] Divesh Maulik, Rahul Pandharipande, and Richard P. Thomas, *Curves on  $K_3$  surfaces and modular forms*, J. Topol. **3** (2010), no. 4, 937–996, DOI 10.1112/jtopol/jtq030.
- [70] Atish Dabholkar, Sameer Murthy, and Don Zagier, *Quantum Black Holes, Wall Crossing, and Mock Modular Forms*, 2012. arXiv:1208.4074.