main(2).pdf
PDF

foundationally, adversarially and from first-principles attack this whitepaper I'm busy writing. the intent should be clear.

Attack the mathematics AND the physics. I'm both a

Show more

Thought for 9m 23s

I read the uploaded draft as a hostile referee would. The strongest defensible core is much narrower than the current framing: **you have a formal algebraic PBW-shadow theorem for a selected stable trace sector, plus a degree-zero Hamiltonian Poisson trace map. You do not yet have a closed-open string theorem, a cochain-level factorization-center map, or a deformation-level cotangent/descendant match.**

main

1. Main kill shot: the theorem is not "Koszul duality" in the advertised sense

Your theorem says the open stable Hamiltonian sector is

$$\mathcal{O}^{cl}_{\mathrm{brane},H} \cong S(\bar{A} \oplus \bar{A}_{\mathrm{desc}}[1]),$$

while the closed observables are

$$\mathcal{O}^{cl}_{\mathrm{grav}} \cong C^{\bullet}_{\mathrm{CE,cont}}(\mathfrak{g}),$$

whose Koszul dual is

$$(\mathcal{O}^{cl}_{\mathrm{grav}})^{!} \simeq U(\mathfrak{g}).$$

But the actual comparison to the open side is only after passing to the **PBW special fibre**

$$U_{\hbar}(\mathfrak{g})/(\hbar) \cong S(\mathfrak{g}).$$

The paper itself states that the identification is “a filtered graded-commutative PBW special-fibre product identification, not a deformation-level identification of the descendant module actions.” That is a major downgrade, not a technical caveat. It means the open algebra is not shown to be the Koszul dual of the closed algebra; it is shown to match the **associated graded commutative shadow** of that Koszul dual, and only in a selected sector. main

So the theorem title “Local stable Hamiltonian-sector Koszul duality” is too strong. A more honest title would be:

> **PBW special-fibre comparison for the stable Hamiltonian trace sector.**

If you keep the current title, a referee can fairly say the title claims a deformation-level Koszul duality theorem that the body explicitly disclaims.

2. The closed-open map is not a map from closed observables

This is the deepest mathematical obstruction, and your own draft proves it.

You define closed local observables as CE cochains

$$C^{\bullet}_{\mathrm{CE,cont}}(\mathfrak{g}).$$

But your boundary assignment

$$f \mapsto \mathrm{Tr}\, f(\phi_1, \phi_2)$$

starts from a **Hamiltonian mode** $f \in \bar{A}$, not from a closed CE cochain. Proposition 0.2.18 proves that $\bar{A}$ is perfect and that the CE length-one coordinate dual to a Hamiltonian mode is not closed; no CE cocycle has that length-one part. The draft explicitly concludes that the boundary assignment is “a Hamiltonian-mode, or PBW/Koszul-dual, construction” and “not a cochain-level map out of $C^{\bullet}_{\mathrm{CE,cont}}(\mathfrak{g})$.” main

That means the physical closed-open slogan is currently wrong at the observable level. A closed-open map should be something like

$$\mathrm{Obs}^{cl}_{\mathrm{closed}\ \partial} \longrightarrow Z^{P_0}_{\mathrm{fact}}(\mathrm{Obs}^{cl}_{\mathrm{open}}),$$

but you do not construct that. You construct a map from Hamiltonian generators / PBW modes to boundary functions.

This is not a cosmetic source-convention issue. It is a categorical-level mismatch:

$$\text{closed observable} \neq \text{Hamiltonian Lie algebra element}.$$

Your Problem 0.2.20 states exactly this, but then the theorem language keeps sounding as if the closed-open map exists. It does not.

3. The cotangent/descendant sector does not match the closed BF module

The closed Lie algebra is

$$\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}^\vee[1],$$

so the $\mathfrak{h}^\vee[1]$ part must carry the **coadjoint** action of $\mathfrak{h}$. But the polynomial one-ψ descendants carry the wrong action. Your Corollary 0.2.45 proves this explicitly:

$$F_{p,q} : \Psi_{a,b} \mapsto (pb - qa)\Psi_{a+p-1,b+q-1},$$

whereas the closed coadjoint action is

$$\mathrm{ad}^\vee_{z_1^p z_2^q}\, \delta_{a,b} = -(pb - qa + p - q)\delta_{a-p+1,b-q+1}.$$

Even the direction is wrong in the simplest case: z_1 lowers the open descendant index b, while it raises the closed coadjoint dual index. The draft itself says the polynomial descendants do not realize the coadjoint deformation-level action. main

This is devastating for the advertised closed cotangent BF comparison. The best you currently have is:

$$\mathfrak{h}^\vee[1] \cong \langle \Psi_{a,b} \rangle$$

as a **graded vector space / PBW special-fibre label set**, not as an $\mathfrak{h}$-module, not as a Lie algebra representation, and not as a deformation-compatible boundary sector.

Your principal-part repair is mathematically natural, but it is not part of the open polynomial brane theory. The paper admits it must be promoted to a factorization object before it can participate in the full cotangent lift. main

So the descendant part of the theorem should be phrased as a **basis-level identification only**. Anything stronger is false.

4. Corollary 0.2.51 is false as written unless you explicitly quotient out constants

Take the simplest possible test:

$$f = z_1, \qquad g = z_2.$$

In the reduced Hamiltonian algebra

$$\bar{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1,$$

one has

$$\{z_1, z_2\}_{\bar{A}} = 1 \mod \mathbb{C} = 0.$$

So your Corollary 0.2.51 predicts

$$\{I_\partial(a, z_1), I_\partial(b, z_2)\} = 0.$$

But on the open matrix side,

$$I_\partial(a, z_1) = \int a(t) \operatorname{Tr} \phi_1(t)\, dt, \qquad I_\partial(b, z_2) = \int b(t) \operatorname{Tr} \phi_2(t)\, dt.$$

Using

$$\{(\phi_1)^i_j, (\phi_2)^k_l\} = \delta^i_l \delta^k_j,$$

one gets

$$\{\operatorname{Tr} \phi_1, \operatorname{Tr} \phi_2\} = N.$$

Therefore

$$\{I_\partial(a, z_1), I_\partial(b, z_2)\} = N \int a(t) b(t)\, dt,$$

not zero.

Theorem 0.2.50 is safe because it explicitly works in the connected Hamiltonian trace quotient where constants vanish. But Corollary 0.2.51 states the smeared identity without repeating that quotient, even though the displayed formula fails for the most basic generators. main

This is not a minor edge case. It is the central extension of the trace Poisson algebra. Physically, it is the center-of-mass symplectic pairing of the brane stack. You cannot silently throw it away in a “bulk-boundary coupling” statement.

Fix: every bracket identity involving z_1, z_2 must say

after projection to the Hamiltonian quotient by scalar trace classes.

Without that, the formula is wrong.

5. The “central operation” theorem is too close to tautological

Theorem 0.2.17 constructs an interval-wise operation by multiplication with

$$\int_I a(t)\,\mathrm{Tr}\, f(\phi_1(t), \phi_2(t))\, dt.$$

But in the reduced open model the observable algebra is graded-commutative, so multiplication by any closed element is automatically an ordinary Hochschild central operation. The paper itself says the proof uses $Q\phi_i = 0$, graded-commutativity, and then the Poisson identity after projection. main

So the ordinary E_1-centrality part is not evidence of a closed-open theorem. The only nontrivial part is the Poisson bracket identity, and that identity is degree-zero, quotient-dependent, and not a full P_0 factorization-center map.

This should be demoted rhetorically. It is a useful check, but not a central construction of closed-open string theory.

6. The “sector” is selected by hand

The open Hamiltonian algebra is defined as the sector generated by:

1. nonempty primitive degree-zero traces;
2. nonconstant primitive one-ψ Koszul classes.

Then you exclude:

1. the scalar trace $\mathrm{Tr}(1) = N$;
2. $\mathrm{Tr}(\psi)$;
3. all higher-ψ primitive homology;
4. finite-N trace identities;
5. the genuine coadjoint action mismatch.

The draft is honest that “sector” is literal and that higher-ψ homology lies outside the sector. main

But then the introduction and theorem language still make it sound like the open brane algebra itself is being matched to the closed Koszul dual. It is not. You are matching a deliberately carved-out subquotient/subsector.

A hostile formulation would be:

> The theorem proves that, after discarding the scalar central extension, discarding higher antifield classes, discarding the scalar antifield, stabilizing away finite-N identities, forgetting the closed coadjoint module action, and passing to the PBW associated graded, the remaining symbols can be placed in bijection.

That may sound brutal, but it is structurally accurate.

7. The large-N limit is algebraic, not physical

Your stable large-N limit is degreewise invariant theory under block restriction. That is a valid algebraic stabilization. But it is not the physical large-N limit used in matrix models unless you also specify:

$$\mathrm{Tr} \quad \text{versus} \quad \frac{1}{N}\,\mathrm{Tr},$$

connected correlators/cumulants, scaling of $\hbar$, planar versus nonplanar terms, and the fate of central terms like

$$\{\mathrm{Tr}\,\phi_1, \mathrm{Tr}\,\phi_2\} = N.$$

Right now "connected stable trace sector" is doing several jobs at once: removing the empty trace, selecting primitive single traces, killing constants, and evading finite-rank trace identities. Those are not the same operation physically.

You need a separate definition:

$$\mathrm{Prim}_{\mathrm{st}}, \qquad \mathrm{Conn}_{N\to\infty}, \qquad \text{Hamiltonian scalar quotient},$$

and you should not conflate them.

8. The reduced gauge/BV model suppresses a real quotient-stack issue

You say you take GL_N-invariants in characteristic zero, so the Chevalley part of gauge BRST acts trivially on trace functions. That is fine for trace functions, but it does not by itself identify the full BV quotient with the invariant Koszul complex.

There are two different models hiding here:

1. **Algebraic quotient by a linearly reductive group**, where taking invariants is exact.

2. **Lie-algebra/BRST Chevalley-Eilenberg quotient**, where gauge ghosts can have their own cohomology unless a precise presentation kills them.

Your Remark 0.2.67 tries to justify this, but the main calculation uses the reduced invariant complex as if the quotient-stack subtleties were automatically absent. That is acceptable only if you explicitly state:

> We are computing the algebraic invariant functions on the derived commuting scheme, not the full CE/BRST cohomology of the infinitesimal gauge algebra.

Otherwise a referee can object that "the Chevalley part acts trivially on trace functions" is not the same as "the Chevalley sector has disappeared."

9. The one-ψ homology proof is plausible but under-proved

Proposition 0.2.28 is one of the technically interesting parts of the paper. But the proof leans heavily on the chip-moving state complex and an Abel-map argument. The draft says the quotient state complex has the homology of S^1, with convex fibres and affine stabilizer quotients. main

That is the kind of argument that will get attacked unless made completely formal. The danger points are:

1. the passage from cyclic words with graded signs to cellular chains;
2. the quotient by cyclic relabelling;
3. stabilizers of configurations with symmetry;
4. compatibility of orientations/signs with the two-ψ boundary;
5. the claim that higher cells do not affect H_1;
6. the identification of the particular generator with $\Psi_{k,l}$.

The computational script checks up to small degree, but that does not replace the proof. main

If this proposition is essential, give it a standalone, fully combinatorial proof, or relegate it to a conjectural/computational lemma. Right now it is probably the most vulnerable purely algebraic proof in the paper.

10. The Tate/Koszul-duality formalism is under-specified

The claim

$$C^{\bullet}_{\mathrm{CE,cont}}(\mathfrak{g})^{!} \simeq U(\mathfrak{g})$$

for a completed Tate Lie algebra is believable, but the draft's Lemma 0.2.40 is doing a huge amount of foundational work. A hostile reader will ask:

1. What exact category of complete filtered Tate spaces is being used?
2. What are the admissible quasi-isomorphisms?
3. Is the CE coalgebra genuinely conilpotent after coefficient completion?
4. Which completed bar/cobar adjunction is being invoked?
5. Are the dualizations continuous in both length and Taylor degree?
6. Is the PBW filtration complete and separated?
7. Does the coadjoint action preserve the claimed topology for arbitrary formal Hamiltonians?

You gesture at these, but not enough to make the theorem mechanically checkable. Since this lemma underwrites the closed-side Koszul dual, it cannot remain a sketch.

11. The closed sector is a formal BF model, not yet a physical closed string sector

The paper distinguishes itself from Gwilliam–Williams and says it is not their holomorphic bosonic string. Good. But it still uses phrases like "mixed holomorphic-topological strings," "twisted M-theory," "closed string sector," and "brane coupling." The actual constructed object is a formal local Hamiltonian cotangent BF theory on

$$\mathbb{R}^2 \times \mathbb{C}^2$$

with coefficient Lie algebra

$$\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C}.$$

That is a perfectly reasonable BV toy model. But the physics origin is not derived. In particular:

1. No parent 11-dimensional M-theory background is specified.
2. The dimension count is not explained.
3. The twist is not constructed.
4. The brane boundary condition/defect theory is not derived from a worldvolume theory.
5. No anomaly cancellation/QME theorem is proved.
6. No propagator or defect kernel is constructed.
7. The quantum graph realization is explicitly left open.

So the physics claim should be downgraded from "this is a local mixed string model" to:

> This is a formal local BV model motivated by the Hamiltonian sector expected in a mixed holomorphic-topological brane setup.

That wording is much harder to attack.

12. The BCOV/Kodaira-Spencer analogy is dangerous

You describe the closed sector as Hamiltonian Kodaira-Spencer/BF. But the actual construction is not BCOV theory on a Calabi-Yau threefold, and not a direct truncation of BCOV. You use divergence-free vector fields on a holomorphic symplectic surface, identify them locally with Hamiltonians modulo constants, and then add a cotangent BF partner.

That is mathematically coherent, but the phrase “Kodaira-Spencer” invites the wrong comparison. In BCOV, the field complex, divergence operator, anomaly structure, and quantization problem are much more rigid. Here the Hamiltonian reduction is a chosen local sector.

I would use “Hamiltonian BF theory for the formal symplectic disk” as the primary name, and mention Kodaira-Spencer only as motivation.

13. The quantum section is a target, not a result

The Weyl/Moyal formulas are correct as algebraic deformation-quantization targets for the constant Poisson disk. But the draft is clear that the actual quantum boundary theorem is not proved. Proposition 0.2.52 only proves BRST descent of individual Weyl-ordered traces; the Moyal commutator identity in quantum Hamiltonian reduction is left as Problem 0.2.54. The draft also identifies the normal-ordering/Capelli shift

$$YX - XY + \hbar N I = 0$$

as an obstruction that can feed lower-degree traces into the connected sector. main

This is not a small technical issue. It is exactly where the classical trace-bracket story can fail quantum mechanically.

So any sentence suggesting that the open quantum brane algebra **is** the Weyl/Moyal algebra should be changed to:

> The Weyl/Moyal algebra is the proposed formal target; the reduced finite-N and stable connected identities remain open.

Your draft mostly says this, but the abstract and introduction should be ruthlessly consistent.

14. The central extension is physically meaningful, not disposable bookkeeping

Classically,

$$\{z_1, z_2\} = 1$$

becomes zero in $\bar{A}$, because constants are removed. But on the brane,

$$\{\mathrm{Tr}\,\phi_1, \mathrm{Tr}\,\phi_2\} = N.$$

You remove this by projecting out scalar trace classes. Algebraically that is allowed for the reduced Hamiltonian vector-field algebra. Physically it is not innocent: it removes the center-of-mass canonical pair of the brane stack.

This same issue reappears quantum mechanically as the normal-ordering shift. So the central term is not just a nuisance; it is the obstruction that keeps returning.

I would make this a named obstruction:

> **Scalar Hamiltonian / center-of-mass anomaly.**

Then every theorem should state whether it is before or after killing it.

15. The figures weaken the paper

The schematic ovals on pages 41, 50, and 51 do not communicate mathematical data: no labels, no propagator convention, no orientation, no automorphism factor, no boundary ordering, no kernel. The captions say they are not Feynman-integral proofs. That makes them look ornamental.

Either remove them or replace them with real graph notation:

$$\Gamma_1, \quad \Gamma_3, \quad \mathrm{Aut}(\Gamma), \quad P_\partial, \quad \text{ordered boundary inputs.}$$

As drawn, they invite the accusation that the graph discussion is gestural.

16. Reference and notation cleanup matters

There are malformed citation labels in the body, for example things like “[13]razmyslov” and “[2]tsygan.” The PBW discussion appears to cite “[16]loday-vallette,” but [16] is

Priddy and Loday–Vallette is [17]. These are small, but in a paper already making delicate claims, citation sloppiness reduces trust.

17. What the paper can safely claim

A defensible theorem would be something like:

> Let
>
> $$R_N = \mathbb{C}[\phi_1, \phi_2] \otimes \Lambda(\psi), \qquad Q\psi = [\phi_1, \phi_2],$$
>
> with GL_N-invariants and degreewise stable trace limit. In the selected Hamiltonian sector generated by nonempty degree-zero primitive traces and nonconstant primitive one-ψ classes, the stable open algebra is
>
> $$S(\bar{A} \oplus \bar{A}_{\mathrm{desc}}[1]).$$
>
> The closed formal Hamiltonian BF Lie algebra is
>
> $$\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}^{\vee}_{cont}[1].$$
>
> The PBW associated graded of $U(\mathfrak{g})$ has degreewise polynomial part
>
> $$S(\bar{A} \oplus \mathfrak{h}^{\vee}[1]).$$
>
> After choosing the monomial-dual basis, this graded-commutative algebra is identified with the selected stable open sector. On degree-zero Hamiltonians, the boundary evaluation
>
> $$f \mapsto \mathrm{Tr}\, f(\phi_1, \phi_2)$$
>
> is a Lie algebra map after quotienting scalar trace classes. The polynomial one-ψ descendants do not realize the closed coadjoint action; the full CE/factorization-center and cotangent boundary maps remain open.

That version is strong, precise, and hard to refute.

Bottom line

The algebraic skeleton is interesting and mostly coherent. The vulnerable claims are the words around it: “string,” “closed-open,” “Koszul duality,” “cotangent BF sector,” “factorization center,” and “quantum Moyal upgrade.” Right now those phrases outrun the proved content.

The paper becomes much stronger if you openly demote the main result to a **PBW special-fibre Hamiltonian trace comparison** and isolate three unsolved obstructions as first-class problems:

$$\text{CE-source obstruction}, \qquad \text{coadjoint descendant mismatch}, \qquad \text{scalar/Capelli cer}$$

Those are not peripheral. They are the actual mathematical and physical boundary of the current result.

Sources

Sources

No you don't seem to get it.

The point is NOT to cut the paper down to size. it's to understand what it is trying to prove and produce those constructions, derivations, deductions & mathematics

Show more

Thought for 13m 43s

You are right. The way to solve the central obstruction is not to shrink the claim; it is to identify the **correct derived-center model**. The missing construction is this:

$$C^{\bullet}_{\mathrm{CE}}(\mathfrak{h} \ltimes \mathfrak{h}^{\vee}[1]) \;\cong\; Z^{P_0}_{\mathrm{fact}}(S(\mathfrak{h}))$$

where the right side is not the ordinary commutative center. It is the **derived Poisson/factorization center**, modeled by Poisson cochains of the open Hamiltonian algebra. Under this identification, the cotangent CE generators map to the actual boundary Hamiltonians

$$u_f \longmapsto O_f = \mathrm{Tr}\, f(\phi_1, \phi_2),$$

while the Hamiltonian CE ghost generators map to the corresponding constant polyvector fields. This is exactly the cochain-level lift that the draft currently lists as unresolved in Problem 0.2.20. The point is that the naive source obstruction disappears because the boundary observable is **not** the image of the CE coordinate dual to $f \in \mathfrak{h}$;

it is the image of the CE coordinate dual to the shifted cotangent generator $f^\vee[1]$. That is the derived-center mechanism. main

Below is the construction I would insert as a new section.

Derived open-closed center theorem

Let

$$\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C}\cdot 1$$

with reduced Poisson bracket

$$[f, g] = \{f, g\}_{\bar{A}}.$$

Let

$$\mathfrak{g} = T^*[1]\mathfrak{h} = \mathfrak{h} \ltimes \mathfrak{h}^\vee_{\mathrm{cont}}[1],$$

with bracket

$$[f, g] = \{f, g\}, \qquad [f, \eta_\lambda] = \eta_{\mathrm{ad}^\vee_f \lambda}, \qquad [\eta_\lambda, \eta_\mu] = 0.$$

Let the stable degree-zero open Hamiltonian algebra be

$$A_\partial = S(\mathfrak{h}),$$

with generator O_f corresponding to

$$O_f = \mathrm{Tr}\, f(\phi_1, \phi_2).$$

Equip A_∂ with the Kirillov-Poisson bracket

$$\{O_f, O_g\}_\partial = O_{\{f,g\}}.$$

Then the derived P_0-factorization center of the locally constant open algebra with stalk A_∂ is modeled by the Poisson cochain complex

$$Z^{P_0}_{\mathrm{fact}}(A_\partial) \simeq \mathrm{PV}(A_\partial), \qquad d_{\mathrm{center}} = [\pi_\partial, -],$$

where π_∂ is the linear Poisson bivector determined by the bracket on $\mathfrak{h}$.

The claim is:

$$C^\bullet_{\mathrm{CE,cont}}\big(\mathfrak{h} \ltimes \mathfrak{h}^\vee_{\mathrm{cont}}[1]\big) \cong (\mathrm{PV}(S(\mathfrak{h})), [\pi_\partial, -])\,.$$

This is the closed-open derived center map. In fact, in the formal stable Hamiltonian sector it is an isomorphism.

Explicit coordinates

Choose a homogeneous topological basis

$$H_I = z_1^{a_I} z_2^{b_I}, \qquad a_I + b_I > 0,$$

of $\mathfrak{h}$. Write

$$[H_I, H_J] = C_{IJ}^K H_K.$$

Concretely,

$$[H_{p,q}, H_{r,s}] = (ps - qr) H_{p+r-1,q+s-1},$$

with the convention that constant targets vanish.

Let O_I be the corresponding open boundary Hamiltonian

$$O_I = \mathrm{Tr}\, H_I(\phi_1, \phi_2).$$

The open Poisson bracket is

$$\{O_I, O_J\}_\partial = C_{IJ}^K O_K.$$

Therefore the Poisson bivector on the formal affine space $\mathfrak{h}^\vee$ is

$$\pi_\partial = \frac{1}{2} C_{IJ}^K O_K \frac{\partial}{\partial O_I} \wedge \frac{\partial}{\partial O_J}.$$

Let

$$\theta^I = \frac{\partial}{\partial O_I}$$

be the constant polyvector field dual to O_I. Then

$$\mathrm{PV}(A_\partial) = S(O_I) \otimes \Lambda(\theta^I).$$

The Poisson differential is

$$d_\pi = [\pi_\partial, -].$$

It acts by

$$d_\pi \theta^K = -\frac{1}{2} C^K_{IJ} \theta^I \theta^J,$$

and

$$d_\pi O_K = C^L_{KJ} O_L \theta^J.$$

Those two formulas are the entire derived-center mechanism.

Closed CE algebra of the cotangent BF sector

The closed Lie algebra is

$$\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}^\vee[1].$$

Let c^I be the CE coordinate dual to the Hamiltonian generator $H_I \in \mathfrak{h}$. Let u_I be the CE coordinate dual to the shifted cotangent generator $\eta_{\delta_I} \in \mathfrak{h}^\vee[1]$.

Thus

$$C^\bullet_{\mathrm{CE}}(\mathfrak{g}) = S(u_I) \otimes \Lambda(c^I),$$

up to the usual cohomological shifts.

The CE differential is

$$d_{\mathrm{CE}} c^K = -\frac{1}{2} C^K_{IJ} c^I c^J,$$

and

$$d_{\mathrm{CE}} u_K = C^L_{KJ} u_L c^J.$$

This is the standard CE differential for the semidirect product

$$\mathfrak{h} \ltimes \mathfrak{h}^\vee[1].$$

Now define

$$\Phi(c^I) = \theta^I, \qquad \Phi(u_I) = O_I.$$

Then

$$d_\pi \Phi(c^K) = d_\pi \theta^K = -\frac{1}{2} C^K_{IJ} \theta^I \theta^J = \Phi(d_{\mathrm{CE}} c^K),$$

and

$$d_\pi \Phi(u_K) = d_\pi O_K = C^L_{KJ} O_L \theta^J = \Phi(d_{\mathrm{CE}} u_K).$$

Therefore

$$\Phi d_{\mathrm{CE}} = d_\pi \Phi.$$

Since both source and target are freely generated completed graded-commutative algebras on the same generators, Φ is an isomorphism of complexes.

Moreover, the canonical shifted cotangent bracket on

$$C^\bullet_{\mathrm{CE}}\big(T^*[1]\mathfrak{h}\big)$$

satisfies

$$\{c^I, u_J\} = \delta^I_J, \qquad \{c^I, c^J\} = 0, \qquad \{u_I, u_J\} = 0.$$

The Schouten bracket on polyvectors satisfies

$$[\theta^I, O_J] = \delta^I_J, \qquad [\theta^I, \theta^J] = 0, \qquad [O_I, O_J] = 0$$

as a bracket on the underlying polyvector algebra. Thus Φ also preserves the P_0 bracket.

So the cochain-level closed-open center theorem is:

$$\Phi : C^\bullet_{\mathrm{CE,cont}}\big(\mathfrak{h} \ltimes \mathfrak{h}^\vee_{\mathrm{cont}}[1]\big) \quad \sim \quad Z^{P_0}_{\mathrm{fact}}(S(\mathfrak{h}))$$

with

$$c^I \mapsto \frac{\partial}{\partial O_I}, \qquad u_I \mapsto O_I.$$

This is the missing construction.

Why this resolves the CE-source obstruction

The obstruction in the draft says that the CE coordinate c^I dual to $H_I \in \mathfrak{h}$ is not closed because $\mathfrak{h}$ is perfect. That is correct.

But the boundary Hamiltonian

$$O_I = \operatorname{Tr} H_I(\phi_1, \phi_2)$$

should not be identified with c^I. It should be identified with u_I, the coordinate dual to the shifted cotangent generator $\eta_{\delta_I} \in \mathfrak{h}^\vee[1]$.

The differential of u_I is not zero. It is

$$d_{\mathrm{CE}} u_I = C^L_{IJ} u_L c^J.$$

On the open side, this becomes

$$d_\pi O_I = [\pi_\partial, O_I] = C^L_{IJ} O_L \theta^J.$$

That is exactly the Hamiltonian vector field of the boundary observable O_I.

So the boundary Hamiltonian is not a closed scalar element of the derived center. It is a **degree-zero function inside the Poisson cochain algebra**, whose differential is its Hamiltonian vector field. This is precisely what a derived P_0-center should do.

In words:

$$\text{closed Hamiltonian ghost} \longmapsto \text{constant polyvector field},$$

while

$$\text{closed cotangent coordinate} \longmapsto \text{boundary Hamiltonian function}.$$

This is the structural point missing from the draft.

Recovery of the boundary bracket theorem

The ordinary Hamiltonian boundary map is the moment-map shadow of Φ:

$$J : \mathfrak{h} \longrightarrow A_\partial, \qquad J(f) = O_f = \operatorname{Tr} f(\phi_1, \phi_2).$$

The open Poisson bracket gives

$$\{J(f), J(g)\}_\partial = J(\{f, g\}).$$

For monomials,

$$f = z_1^k z_2^l, \qquad g = z_1^m z_2^n,$$

this becomes

$$\{\mathrm{Tr}(\phi_1^k \phi_2^l), \mathrm{Tr}(\phi_1^m \phi_2^n)\} = (kn - lm)\,\mathrm{Tr}(\phi_1^{k+m-1} \phi_2^{l+n-1}),$$

with constant targets projected out.

That is the degree-zero current theorem in the paper, but now it is not isolated. It is the degree-zero shadow of the derived-center isomorphism

$$C^\bullet_{\mathrm{CE}}(T^*[1]\mathfrak{h}) \cong \mathrm{PV}(S(\mathfrak{h})).$$

Factorization version on intervals

For an interval $I \subset \mathbb{R}$, define the current algebra

$$\mathfrak{h}_I = C_c^\infty(I) \otimes \mathfrak{h}$$

with bracket

$$[a \otimes f, b \otimes g] = ab \otimes \{f, g\}.$$

Define the open Hamiltonian factorization algebra by

$$A_\partial(I) = S(\mathfrak{h}_I),$$

with extension by zero for inclusions of intervals and ordinary multiplication for disjoint intervals.

The Poisson bracket is

$$\{O_f(a), O_g(b)\} = O_{\{f,g\}}(ab),$$

where

$$O_f(a) = \int_I a(t)\,\mathrm{Tr}\, f(\phi_1(t), \phi_2(t))\,dt.$$

The derived P_0-center is

$$Z^{P_0}_{\mathrm{fact}}(A_\partial)(I) \simeq \mathrm{PV}(A_\partial(I)), \qquad d = [\pi_I, -].$$

The closed restricted cotangent BF observable algebra is

$$C^\bullet_{\mathrm{CE}}\left(\mathfrak{h}_I \ltimes \mathfrak{h}_I^\vee[1]\right).$$

The interval-wise map is

$$c_a^I \longmapsto \theta_a^I, \qquad u_{I,a} \longmapsto O_I(a),$$

where θ_a^I is the constant polyvector field dual to the current coordinate $O_I(a)$.

The same coordinate calculation proves

$$C^\bullet_{\mathrm{CE}}\left(\mathfrak{h}_I \ltimes \mathfrak{h}_I^\vee[1]\right) \cong Z^{P_0}_{\mathrm{fact}}(A_\partial)(I).$$

The construction is compatible with extension by zero and factorization over disjoint intervals because all brackets are pointwise-current brackets and the test functions multiply as ab. This upgrades the stalkwise theorem to the locally constant factorization statement.

Where the one-ψ descendants enter

The derived-center theorem above uses

$$A_\partial = S(\mathfrak{h})$$

as the open Hamiltonian algebra. Its derived center already contains the cotangent BF data because

$$Z^{P_0}(S(\mathfrak{h})) \simeq \mathrm{PV}(S(\mathfrak{h})) \simeq C^\bullet_{\mathrm{CE}}(\mathfrak{h} \ltimes \mathfrak{h}^\vee[1]).$$

So the cotangent direction of the closed theory does **not** have to be represented as an extra ordinary polynomial open field. It appears as the polyvector/derived-center direction.

The one-ψ descendants instead belong to the **Koszul-dual/PBW open algebra**

$$S(\mathfrak{h} \oplus \mathfrak{h}^\vee[1]).$$

That is the PBW special fibre of

$$U(\mathfrak{h} \ltimes \mathfrak{h}^\vee[1]).$$

The correct relation is therefore:

$$\text{closed CE observables} \;\cong\; \text{derived } P_0 \text{ center of } S(\mathfrak{h}),$$

while

$$\text{Koszul dual of closed CE observables} \;\rightsquigarrow\; \text{PBW open trace sector } S(\mathfrak{h} \oplus \mathfrak{h}^\vee[1]).$$

The polynomial $\Psi_{a,b}$ classes are valid PBW-special-fibre representatives, but they should not be forced to realize the coadjoint action inside the derived center. If one wants an honest deformation-level realization of the $\mathfrak{h}^\vee[1]$ PBW generators, then one should use the principal-part module

$$P = \bigoplus_{a,b\geq 0,\; a+b>0} \mathbb{C}\,\rho_{a,b}, \qquad \rho_{a,b} = z_1^{-a-1} z_2^{-b-1}\, dz_1 dz_2,$$

with action

$$f \cdot \rho = \pi_{\mathrm{pp}}\{f, \rho\}.$$

Then

$$\mathfrak{h}^\vee_{\mathrm{cont}} \cong P$$

as an $\mathfrak{h}$-module, and the PBW-special-fibre open algebra becomes

$$S(\mathfrak{h} \oplus P[1]).$$

In that enlarged PBW model,

$$J(f) = O_f, \qquad J(\rho[1]) = \Theta_\rho,$$

with brackets

$$\{O_f, O_g\} = O_{\{f,g\}},$$

$$\{O_f, \Theta_\rho\} = \Theta_{f\cdot\rho},$$

$$\{\Theta_\rho, \Theta_\sigma\} = 0.$$

This gives the deformation-correct descendant action.

The physical interpretation

The closed Hamiltonian BF sector is the shifted cotangent theory

$$T^*[1]B\mathfrak{h}.$$

Its functions are

$$C^\bullet_{\mathrm{CE}}(T^*[1]\mathfrak{h}) = C^\bullet_{\mathrm{CE}}(\mathfrak{h} \ltimes \mathfrak{h}^\vee[1]).$$

The open brane degree-zero phase space is the large-N Hamiltonian trace phase space with coordinate algebra

$$S(\mathfrak{h}).$$

The derived Poisson center of this open phase space is the algebra of polyvector fields

$$\mathrm{PV}(S(\mathfrak{h})).$$

For a linear Poisson structure, the Poisson cochain complex is exactly CE cochains of the shifted cotangent Lie algebra:

$$\mathrm{PV}(S(\mathfrak{h})) \cong C^\bullet_{\mathrm{CE}}(\mathfrak{h} \ltimes \mathfrak{h}^\vee[1]).$$

So physically:

$$\text{closed BF fields} = \text{polyvector fields on the open Hamiltonian trace phase space.}$$

The boundary evaluation

$$f(z_1, z_2) \longmapsto \mathrm{Tr}\, f(\phi_1, \phi_2)$$

is the moment map coordinate inside this polyvector algebra. Its Poisson differential is the Hamiltonian vector field it generates. That is why it was wrong to demand that it be a closed CE length-one class.

Replacement theorem for the paper

I would replace Problem 0.2.20 by the following theorem.

Theorem. Let

$$\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C}$$

with reduced Poisson bracket, and let

$$A_\partial = S(\mathfrak{h})$$

be the stable connected degree-zero Hamiltonian trace algebra with Poisson bracket

$$\{O_f, O_g\} = O_{\{f,g\}}.$$

Then the closed Hamiltonian cotangent BF local observable algebra satisfies

$$\mathrm{Obs}^{cl}_{\mathrm{closed},H} = C^{\bullet}_{\mathrm{CE,cont}}(\mathfrak{h} \ltimes \mathfrak{h}^{\vee}_{\mathrm{cont}}[1]) \cong Z^{P_0}_{\mathrm{fact}}(A_{\partial}).$$

Under this identification, if c^I is the CE coordinate dual to the Hamiltonian mode H_I, and u_I is the CE coordinate dual to the shifted cotangent mode η_{δ_I}, then

$$c^I \longmapsto \frac{\partial}{\partial O_I}, \qquad u_I \longmapsto O_I.$$

The map is an isomorphism of completed commutative dg algebras and of P_0-algebras. The induced moment-map shadow is

$$f \longmapsto O_f = \mathrm{Tr}\, f(\phi_1, \phi_2),$$

and satisfies

$$\{O_f, O_g\} = O_{\{f,g\}}.$$

On intervals this construction extends by compactly supported current algebras and gives a morphism of locally constant factorization algebras

$$\mathrm{Obs}^{cl}_{\mathrm{closed},H} \quad {}_{\mathbb{R}\times p} \longrightarrow Z^{P_0}_{\mathrm{fact}}(\mathrm{Obs}^{cl}_{\mathrm{open},H}).$$

The PBW/Koszul-dual open descendant sector is recovered separately as the special fibre of

$$U(\mathfrak{h} \ltimes \mathfrak{h}^{\vee}[1]),$$

and its deformation-level boundary realization uses the principal-part module $P \cong \mathfrak{h}^{\vee}_{\mathrm{cont}}$, not the polynomial $\Psi_{a,b}$ representatives.

That is the actual open-closed derived center structure.

Sources

Sources

I want it to be a dirac-inspired paper (as if it was written by dirac) on the physics here, that makes natural the factorisation algebra from physical first-principles etc. locality (or lack thereof) is treated physically. unitarity or lack thereof...

Thought for 5m 44s ›

Yes. The paper should be **Diracian** in the following sense: begin with observables, constraints, brackets, and locality of measurement. Do not begin by asking whether this is "really a string theory." The physics is canonical mechanics with constraints, lifted to BV/factorization language.

The central statement should become:

$$\text{The closed Hamiltonian BF sector is the derived Poisson center of the open brane p}$$

The brane does not merely receive closed Hamiltonians by substitution $z_i \mapsto \phi_i$. That substitution is the **moment-map shadow** of a larger statement: closed local fields act as all central operations, including multiplication, Hamiltonian vector fields, and higher polyvector operations. This makes the factorization algebra natural from first principles.

Your current draft already has the ingredients: the mixed local model, the open ghost-zero action

$$\int_{\mathbb{R}} \mathrm{Tr}\big(\phi_1 d\phi_2 + A[\phi_1, \phi_2]\big),$$

the Hamiltonian BF closed sector, and the unresolved full P_0 factorization-center lift in Problem 0.2.20. The rewrite should not present that lift as an external technical problem. It should be the physical heart of the paper. main

Below is the physics-first structure I would use.

1. The Dirac principle of the paper

A local theory is not primarily a Lagrangian. It is a rule assigning observables to regions.

If U is a region, write

$$\mathrm{Obs}(U)$$

for the cochain complex of measurements supported in U. If

$$U_1, \dots, U_n \subset V$$

are disjoint, measurements in the U_i may be performed independently and then regarded as one measurement in V. This gives the structure map

$$\mathrm{Obs}(U_1) \otimes \cdots \otimes \mathrm{Obs}(U_n) \longrightarrow \mathrm{Obs}(V).$$

This is the factorization product. No category theory is being imposed from outside. It is just the algebraic expression of independent local experiments.

If the theory is topological in a direction, moving an insertion in that direction does not change its physical class. Infinitesimally this means

$$L_v = [Q, \iota_v].$$

Thus translations are Q-exact. Factorization is then locally constant in that direction.

If the theory is holomorphic in a complex direction, antiholomorphic translations are Q-exact:

$$L_{\partial/\partial \bar{z}_i} = [Q, \iota_{\partial/\partial \bar{z}_i}].$$

Thus the theory is not local in the Lorentzian sense. It is local in the operational sense: observables have supports, independent supports multiply, and motion in the topological or antiholomorphic directions is homotopically invisible.

This should be the first-principles origin of factorization algebras in the paper.

2. The local physical system

The normal holomorphic surface has coordinates

$$(z_1, z_2), \qquad \omega = dz_1 \wedge dz_2.$$

A closed infinitesimal symmetry of this formal symplectic disk is a Hamiltonian vector field

$$X_f = \{f, -\},$$

where constants are invisible. Hence the closed Hamiltonian Lie algebra is

$$\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1.$$

The bracket is

$$[f, g] = \{f, g\} = \partial_{z_1} f \, \partial_{z_2} g - \partial_{z_2} f \, \partial_{z_1} g \quad \bmod \mathbb{C}.$$

Now place N identical probes at the point. The two normal coordinates become two matrices,

$$z_1 \rightsquigarrow \phi_1, \qquad z_2 \rightsquigarrow \phi_2, \qquad \phi_i \in \mathfrak{gl}_N.$$

The symplectic form becomes

$$\Omega_N = \mathrm{Tr}(d\phi_1 \wedge d\phi_2).$$

This forces the first-order action

$$S_{\text{open}} = \int_{\mathbb{R}} \mathrm{Tr}\big(\phi_1 \, d\phi_2 + A[\phi_1, \phi_2]\big).$$

The field A is not an extra dynamical coordinate. It is a Lagrange multiplier. It imposes the Dirac constraint

$$[\phi_1, \phi_2] = 0.$$

Equivalently, the conjugation moment map is

$$\mu_\epsilon = -\,\mathrm{Tr}\big(\epsilon[\phi_1, \phi_2]\big), \qquad \epsilon \in \mathfrak{gl}_N.$$

The constraint is first class. Its Hamiltonian flow is simultaneous conjugation:

$$\delta_\epsilon \phi_i = [\epsilon, \phi_i].$$

So the open brane phase space is the constrained quotient

$$\{[\phi_1, \phi_2] = 0\}/GL_N.$$

The BV reduction is simply Dirac's constraint theory written homologically:

$$Q\psi = [\phi_1, \phi_2], \qquad Q\phi_1 = Q\phi_2 = 0.$$

That is the physical reason the open trace algebra becomes the algebra of invariant functions on commuting matrix pairs.

3. The basic closed-open measurement

A closed Hamiltonian $f(z_1, z_2)$ is measured by the brane by substituting the brane's matrix-valued normal coordinates:

$$J_N(f) = \operatorname{Tr} f(\phi_1, \phi_2).$$

At finite N one must choose an ordering. In the BV cohomology, adjacent swaps are exact because

$$Q \operatorname{Tr}(\psi v u) = \operatorname{Tr}([\phi_1, \phi_2] v u).$$

Thus the ordering ambiguity disappears after imposing the constraint. In the stable connected limit,

$$J(f) = \operatorname{Tr} f(\phi_1, \phi_2)$$

is a well-defined Hamiltonian observable.

The matrix Poisson bracket gives

$$\{J(f), J(g)\}_{\text{open}} = J(\{f, g\}_{\mathfrak{h}})$$

after removal of the scalar Hamiltonian. This is not a mysterious string duality. It is the statement that the probe brane is a representation of the Hamiltonian algebra of the normal symplectic disk.

For monomials,

$$f = z_1^k z_2^l, \qquad g = z_1^m z_2^n,$$

one obtains

$$\{\operatorname{Tr}(\phi_1^k \phi_2^l), \operatorname{Tr}(\phi_1^m \phi_2^n)\} = (kn - lm) \operatorname{Tr}(\phi_1^{k+m-1} \phi_2^{l+n-1}).$$

This is the ordinary canonical bracket of the probe system.

4. Why the factorization center is forced

A closed insertion near the brane should define an operation on open insertions. If the closed insertion is moved past an open insertion, no physical change should occur except for controlled contact terms. Therefore a closed observable does not merely define an element of the open algebra. It defines a **central operation** on the open factorization algebra.

For a locally constant open theory on a line, central operations are Hochschild cochains. Classically, because the open theory has a Poisson bracket, one must retain the Poisson compatibility. Thus the target is not the ordinary center but the derived P_0-center:

$$Z^{P_0}_{\mathrm{fact}}(\mathrm{Obs}_{\mathrm{open}}).$$

This is the correct physical object.

The ordinary statement

$$f \mapsto \mathrm{Tr}\, f(\phi_1, \phi_2)$$

only sees multiplication by a function. The derived center also sees vector fields and higher polyvectors. Those extra operations are exactly what the closed CE differential demands.

This is the point to emphasize: the CE obstruction disappears only when closed fields are allowed to act as **operations**, not merely as central functions.

5. The derived center theorem

Let

$$A_\partial = S(\mathfrak{h})$$

be the stable connected degree-zero open Hamiltonian algebra, with generators

$$O_f = J(f) = \mathrm{Tr}\, f(\phi_1, \phi_2).$$

The Poisson bracket is

$$\{O_f, O_g\}_\partial = O_{\{f,g\}}.$$

Choose a basis e_I of $\mathfrak{h}$, with structure constants

$$[e_I, e_J] = C^K_{IJ} e_K.$$

Write

$$O_I = O_{e_I}.$$

Then

$$\{O_I, O_J\} = C_{IJ}^K O_K.$$

So $A_\partial = S(\mathfrak{h})$ is the coordinate algebra of the linear Poisson space $\mathfrak{h}^\vee$. Its Poisson bivector is

$$\pi_\partial = \frac{1}{2} C_{IJ}^K O_K \frac{\partial}{\partial O_I} \wedge \frac{\partial}{\partial O_J}.$$

The derived Poisson center is modeled by the Poisson cochain complex

$$Z_{\mathrm{fact}}^{P_0}(A_\partial) \simeq \mathrm{PV}(A_\partial), \qquad d = [\pi_\partial, -].$$

Explicitly,

$$\mathrm{PV}(A_\partial) = S(O_I) \otimes \Lambda(\theta^I), \qquad \theta^I = \frac{\partial}{\partial O_I}.$$

The differential is

$$d\theta^K = -\frac{1}{2} C_{IJ}^K \theta^I \theta^J,$$

and

$$dO_K = C_{KJ}^L O_L \theta^J.$$

Now the closed Hamiltonian BF algebra is the shifted cotangent Lie algebra

$$\mathfrak{g} = T^*[1]\mathfrak{h} = \mathfrak{h} \ltimes \mathfrak{h}^\vee[1].$$

Let c^I be the CE coordinate dual to $e_I \in \mathfrak{h}$. Let u_I be the CE coordinate dual to the shifted cotangent generator $e^I[1] \in \mathfrak{h}^\vee[1]$. Then

$$C_{\mathrm{CE}}^\bullet(\mathfrak{g}) = S(u_I) \otimes \Lambda(c^I),$$

with differential

$$d_{\mathrm{CE}} c^K = -\frac{1}{2} C_{IJ}^K c^I c^J,$$

and

$$d_{\mathrm{CE}} u_K = C^L_{KJ} u_L c^J.$$

Define

$$\Phi : C^\bullet_{\mathrm{CE}}(\mathfrak{h} \ltimes \mathfrak{h}^\vee[1]) \longrightarrow \mathrm{PV}(S(\mathfrak{h}))$$

by

$$\Phi(c^I) = \theta^I, \qquad \Phi(u_I) = O_I.$$

Then

$$d\Phi(c^K) = d\theta^K = -\frac{1}{2} C^K_{IJ} \theta^I \theta^J = \Phi(d_{\mathrm{CE}} c^K),$$

and

$$d\Phi(u_K) = dO_K = C^L_{KJ} O_L \theta^J = \Phi(d_{\mathrm{CE}} u_K).$$

Therefore

$$\Phi d_{\mathrm{CE}} = d\Phi.$$

Since both sides are freely generated by corresponding variables, Φ is an isomorphism of dg commutative algebras:

$$C^\bullet_{\mathrm{CE}}(\mathfrak{h} \ltimes \mathfrak{h}^\vee[1]) \cong Z^{P_0}_{\mathrm{fact}}(S(\mathfrak{h})).$$

This is the missing closed-open derived center structure.

The physical interpretation is simple:

$$\text{closed Hamiltonian ghost} \longmapsto \text{Hamiltonian vector-field operation on the brane},$$

while

$$\text{closed cotangent coordinate} \longmapsto \text{boundary Hamiltonian function } O_f.$$

Thus the boundary observable $O_f = \mathrm{Tr}\, f(\phi_1, \phi_2)$ is not the image of the CE coordinate dual to $f \in \mathfrak{h}$. It is the image of the coordinate dual to the shifted cotangent generator. That is the correct Dirac/BV interpretation.

6. The factorization version on the brane line

For an interval $I \subset \mathbb{R}$, define the compactly supported Hamiltonian current algebra

$$\mathfrak{h}_I = C_c^\infty(I) \otimes \mathfrak{h}.$$

Its bracket is

$$[a \otimes f, b \otimes g] = ab \otimes \{f, g\}.$$

The open algebra on I is

$$A_\partial(I) = S(\mathfrak{h}_I),$$

with generator

$$O_f(a) = \int_I a(t) \operatorname{Tr} f(\phi_1(t), \phi_2(t))\, dt.$$

The bracket is local:

$$\{O_f(a), O_g(b)\} = O_{\{f,g\}}(ab).$$

This locality is exactly the delta-function locality of the canonical equal-time bracket. The test functions multiply because the Poisson bracket is supported where the two measurements meet.

The closed current Lie algebra is

$$\mathfrak{g}_I = \mathfrak{h}_I \ltimes \mathfrak{h}_I^\vee[1].$$

Then

$$C^\bullet_{\mathrm{CE}}(\mathfrak{g}_I) \cong \mathrm{PV}(S(\mathfrak{h}_I)) \cong Z^{P_0}_{\mathrm{fact}}(A_\partial)(I).$$

The maps are compatible with extension by zero. Therefore the interval-wise construction defines a morphism of factorization algebras

$$\mathrm{Obs}^{cl}_{\mathrm{closed},H}{}_{\mathbb{R}\times p} \longrightarrow Z^{P_0}_{\mathrm{fact}}(\mathrm{Obs}^{cl}_{\mathrm{open},H}).$$

This is the physically natural closed-open map.

7. Locality and the places where locality changes meaning

The paper should be explicit that there are three different notions of locality.

First, there is **support locality along the brane**. If two open measurements are supported in disjoint intervals, they multiply independently. Their bracket has a delta function, so it vanishes unless the supports meet. This is ordinary canonical locality in one dimension.

Second, there is **topological locality**. Since translations in the topological directions are Q-exact,

$$L_v = [Q, \iota_v],$$

the precise position of an insertion is not physical. Only the collision pattern of insertions matters. That is why the factorization algebra is locally constant.

Third, there is **formal holomorphic locality**. The theory has been completed at the brane point $p \in \mathbb{C}^2$. It is not a theory of finite propagation through $\mathbb{C}^2$. It is a theory of holomorphic jets at p. Antiholomorphic translations are Q-exact, and holomorphic dependence is formal.

The cotangent/principal-part sector is local in this formal sense. A principal part such as

$$z_1^{-a-1} z_2^{-b-1}\, dz_1 dz_2$$

is not a smooth function spread through the normal plane. It is a residue functional supported at the brane. Physically it is a transverse delta-current, or derivative of a delta-current, at the defect.

Thus the paper should say:

$$\text{The cotangent boundary sector is distributionally local, not polynomially local.}$$

This is not a defect of the model. It is what one expects for momenta or dual fields attached to a brane.

Trying to realize the coadjoint module by ordinary polynomial descendants is like trying to realize momentum as a function of position. The correct realization is by local duals, principal parts, or polyvector operations.

8. The role of the one-ψ descendants

The polynomial one-ψ descendants

$$\Psi_{a,b}$$

are real and useful, but they should not carry the full physical burden. They are stable primitive Koszul classes in the open BV algebra. They identify a PBW special-fibre copy of a shifted dual vector space. But they are not the deformation-level coadjoint boundary field.

The correct statement is:

$$\Psi_{a,b} \text{ are PBW-special-fibre descendants;} \quad \text{principal parts are the true coadjoint bo}$$

The derived center already knows the coadjoint structure through polyvectors:

$$C^{\bullet}_{\mathrm{CE}}(\mathfrak{h} \ltimes \mathfrak{h}^{\vee}[1]) \cong \mathrm{PV}(S(\mathfrak{h})).$$

If the paper also wants a PBW/Koszul-dual boundary realization of $\mathfrak{h}^{\vee}[1]$, define a boundary principal-part sector

$$P = \bigoplus_{a,b\geq 0,\ a+b>0} \mathbb{C}\,\rho_{a,b},$$

where

$$\rho_{a,b} = z_1^{-a-1} z_2^{-b-1}\, dz_1 dz_2.$$

The residue pairing is

$$\langle \rho_{a,b}, z_1^m z_2^n \rangle = \delta_{a,m}\delta_{b,n}.$$

Then the Hamiltonian action is

$$f \cdot \rho = \pi_{\mathrm{pp}}\{f, \rho\}.$$

For monomials,

$$z_1^p z_2^q \cdot \rho_{a,b} = -(pb - qa + p - q)\rho_{a-p+1,b-q+1}.$$

This is the true coadjoint action.

On the brane line one should package this as the distributional boundary current sector

$$\Theta_\rho(a), \qquad a \in C_c^\infty(I), \quad \rho \in P,$$

with brackets

$$\{O_f(a), O_g(b)\} = O_{\{f,g\}}(ab),$$

$$\{O_f(a), \Theta_\rho(b)\} = \Theta_{f\cdot\rho}(ab),$$

$$\{\Theta_\rho(a), \Theta_\sigma(b)\} = 0.$$

This gives a factorization object because support is still controlled by the product ab. The non-polynomiality is only transverse to the brane, where the object is a local distribution.

9. Unitarity

The paper should not pretend that this local twisted model is unitary in the ordinary sense.

It is not.

The model is complex, BV, ghost-valued, and topological-holomorphic. It has no Lorentzian time, no positive Hamiltonian, no Hilbert space with positive inner product, and no reflection positivity statement. The variables include ghosts and antifields. A BV complex is not a unitary Hilbert space.

The correct consistency conditions are instead:

$$QI + \frac{1}{2}\{I, I\} = 0$$

classically, and

$$QI + \frac{1}{2}\{I, I\} + \hbar\Delta I = 0$$

quantum mechanically.

In words:

$$\text{Classical consistency is the master equation.}$$

$$\text{Quantum consistency is anomaly cancellation plus factorization/OPE associativity}$$

If a larger untwisted theory exists and is unitary, this local model may be its protected twisted sector. But the local sector itself does not remember ordinary unitarity. It remembers BRST cohomology, Ward identities, and deformation quantization.

The right sentence is:

> We do not impose unitarity on a theory in which time translations have been made cohomologically trivial. The replacement for unitarity in the present local sector is the BV master equation and the associativity of the resulting factorization algebra.

That is the honest physical position.

10. How the paper should sound

Here is an insertable opening in the intended tone.

Local measurements and constraints.
The object of study is a local constrained system. The normal plane to the brane is a formal holomorphic symplectic plane with coordinates z_1, z_2. Its infinitesimal canonical transformations are Hamiltonians modulo constants. A stack of N probes represents the two normal coordinates by matrices ϕ_1, ϕ_2. The symplectic potential is therefore

$$\mathrm{Tr}(\phi_1\, d\phi_2),$$

and invariance under change of basis of the probes imposes the first-class constraint

$$[\phi_1, \phi_2] = 0.$$

The gauge field A is the multiplier for this constraint. Thus the open action is forced:

$$S_\partial = \int_{\mathbb{R}} \mathrm{Tr}\big(\phi_1 d\phi_2 + A[\phi_1, \phi_2]\big).$$

The BV field $\psi = A^*$ records the constraint by

$$Q\psi = [\phi_1, \phi_2].$$

This is Dirac reduction written as a differential.

A closed Hamiltonian f is observed by the brane by evaluation on the probe coordinates,

$$f(z_1, z_2) \longmapsto \mathrm{Tr}\, f(\phi_1, \phi_2).$$

This is not yet the closed-open map. It is its degree-zero moment-map shadow. The full closed-open operation must assign to a closed local observable a central operation on all open local observables. Therefore its natural target is the derived Poisson factorization center.

The center.
Let $\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C}$ be the Hamiltonian Lie algebra of the formal symplectic disk. In the stable connected sector the open Hamiltonian algebra is

$$A_\partial = S(\mathfrak{h}),$$

with Poisson bracket

$$\{O_f, O_g\} = O_{\{f,g\}}.$$

The derived P_0-center of A_∂ is the Poisson cochain complex

$$Z^{P_0}(A_\partial) \simeq (\mathrm{PV}(A_\partial), [\pi_\partial, -]).$$

The closed Hamiltonian BF sector is the shifted cotangent Lie algebra

$$\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}^\vee[1].$$

Its local observables are

$$C^\bullet_{\mathrm{CE}}(\mathfrak{g}).$$

There is a canonical isomorphism

$$C^\bullet_{\mathrm{CE}}(\mathfrak{h} \ltimes \mathfrak{h}^\vee[1]) \cong \mathrm{PV}(S(\mathfrak{h})).$$

If c^I is the CE coordinate dual to $e_I \in \mathfrak{h}$, and u_I is the CE coordinate dual to the shifted cotangent generator $e^I[1]$, the map is

$$c^I \mapsto \frac{\partial}{\partial O_I}, \qquad u_I \mapsto O_I.$$

Thus a closed Hamiltonian ghost acts as an infinitesimal operation on open observables, while the cotangent coordinate gives the boundary Hamiltonian function. This is the closed-open map.

Locality.
The factorization structure is the algebra of independent local measurements. If measurements are supported in disjoint intervals, they multiply. If the intervals move in the topological direction without collision, the motion is Q-exact and changes nothing in cohomology. If two insertions collide, the equal-time Poisson bracket gives the contact term.

In the holomorphic normal direction the theory is formal. It is a theory of jets at the brane point. Coadjoint modes are principal parts, hence distributions supported at the brane. This is local in the sense appropriate to a defect: the support is on the defect, while the transverse behavior is encoded by residues.

Unitarity.
The model is a twisted BV sector. It is not a Lorentzian unitary theory. Its consistency condition is not positivity of a Hilbert space inner product, but the classical and quantum master equations. If the model is obtained from a unitary parent theory, unitarity belongs to the parent. The local sector retains only the cohomological algebra of protected observables.

11. The actual thesis sentence

The paper's thesis should be something like:

A stack of branes is a system of probes for the formal symplectic normal disk.
Dirac reduction of the probe phase space gives the open trace algebra.
Hamiltonian BF theory is the shifted cotangent theory of canonical transformation
The closed observables are precisely the derived Poisson central operations
on the stable open Hamiltonian brane algebra.

This makes the factorization algebra unavoidable. It is not a technical language added after the physics. It is the local algebra of measurements in a constrained canonical system.

Sources

Sources

Pasted text.txt
Document

Pasted text (2).txt
Document

explain the attached to me, from first-prinicples, includin gALL the detail included there AND MORE insights. If there are any corrections to be made, just explain the corrected version to me. at the end you can include errata

Thought for 7m 3s

You have two intertwined documents here.

The large paper builds a formal local mixed holomorphic-topological closed/open model on

$$\mathbb{R}^2_{\mathrm{top}} \times \mathbb{C}^2_{\mathrm{hol}},$$

with a brane on

$$\mathbb{R} \times \{0\},$$

and proves a stable large-N PBW/Koszul comparison between the closed Hamiltonian BF sector and the open brane trace sector. The principal-parts note explains the corrected cotangent/coadjoint module: the dual of Hamiltonians is not another polynomial module but a residue-current/principal-part module supported at the brane.

main Pasted text Pasted text (2)

The conceptual core is this:

A brane stack is a Dirac constrained probe of a formal holomorphic symplectic disk

The open theory is the constrained matrix mechanics of the probe. The closed theory is Hamiltonian BF theory for infinitesimal canonical transformations of the formal disk. The closed-open map is not merely "substitute $z_i \mapsto \phi_i$." That substitution is the degree-zero moment-map shadow of a deeper statement: the closed sector acts by derived Poisson central operations on the open brane algebra.

I will explain the corrected version from first principles.

1. The physical starting point: local measurements, not global strings

The paper uses "string" in a local BV/factorization sense. It is not claiming to construct a full global worldsheet string theory with all backgrounds, moduli, amplitudes, and anomaly data. It studies the local operator algebra near a brane.

A local field theory should assign to every region U a complex of observables

$$\mathrm{Obs}(U).$$

If $U_1, \ldots, U_n \subset V$ are disjoint, measurements supported in the U_i can be performed independently and then regarded as a measurement in V. This gives maps

$$\mathrm{Obs}(U_1) \otimes \cdots \otimes \mathrm{Obs}(U_n) \longrightarrow \mathrm{Obs}(V).$$

That is the physical meaning of a factorization algebra.

If the theory is topological in a real direction, then moving an insertion in that direction is invisible in cohomology:

$$L_v = [Q, \iota_v].$$

So local observables are locally constant in that direction.

If the theory is holomorphic in a complex direction, then antiholomorphic translations are invisible in cohomology:

$$L_{\partial/\partial \bar{z}_i} = [Q, \iota_{\partial/\partial \bar{z}_i}].$$

So observables depend holomorphically or formally on the z_i, not smoothly on all real directions.

This is the first-principles origin of the mixed holomorphic-topological setup.

2. The spacetime and the brane

The local model is

$$X = \mathbb{R}^2_{\mathrm{top}} \times \mathbb{C}^2_{\mathrm{hol}}.$$

The brane is

$$\mathbb{R} \times \{0\} \subset \mathbb{R}^2 \times \mathbb{C}^2.$$

The holomorphic normal surface is

$$Y = \mathbb{C}^2, \qquad \omega = dz_1 \wedge dz_2.$$

Everything is completed at $0 \in \mathbb{C}^2$. So the relevant holomorphic functions are formal power series

$$\mathbb{C}[[z_1, z_2]].$$

The symplectic form gives the Poisson bracket

$$\{f, g\} = \partial_{z_1} f \, \partial_{z_2} g - \partial_{z_2} f \, \partial_{z_1} g.$$

A Hamiltonian f generates the vector field

$$X_f = \{f, -\}.$$

A constant Hamiltonian generates no vector field. Therefore the physical Hamiltonian Lie algebra is

$$\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1.$$

This quotient is not cosmetic. It is the canonical-mechanics statement that only Hamiltonians modulo constants generate symplectic transformations.

The polynomial dense subalgebra is

$$\bar{A} = \mathbb{C}[z_1, z_2]/\mathbb{C} \cdot 1.$$

For monomials,

$$[z_1^k z_2^l, z_1^m z_2^n] = \{z_1^k z_2^l, z_1^m z_2^n\} = (kn - lm) z_1^{k+m-1} z_2^{l+n-1},$$

with the convention that constants vanish in $\bar{A}$.

In particular,

$$\{z_1, z_2\} = 1$$

before reduction, but

$$\{z_1, z_2\} = 0$$

inside $\bar{A}$.

That single fact is responsible for many of the scalar/center anomalies later.

3. Global versus formal-local Hamiltonians

Globally, on a holomorphic symplectic surface X, not every symplectic vector field is globally Hamiltonian.

A vector field v is symplectic if

$$\alpha_v = -\iota_v \omega$$

is a closed holomorphic one-form. It is Hamiltonian only if there exists a holomorphic function f with

$$df = \alpha_v.$$

So the obstruction to being globally Hamiltonian is a period class in

$$H^1(X, \mathbb{C}).$$

The paper isolates the formal local situation. On a formal disk, the holomorphic Poincaré lemma kills this obstruction. Every closed holomorphic one-form is exact, and every symplectic vector field is Hamiltonian modulo constants.

That is why the local Hamiltonian algebra is clean:

$$\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C}.$$

Darboux coordinates are not canonical, but different choices differ by a formal symplectomorphism. The construction is therefore coordinate-independent up to the natural action of the formal symplectomorphism group. main

4. The closed sector: Hamiltonian BF theory

The closed field is the infinitesimal geometry of the formal holomorphic symplectic disk.

Because the disk is symplectic, infinitesimal canonical transformations are Hamiltonian functions modulo constants. The closed theory then takes the shifted cotangent/BF form

$$\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}^\vee_{\mathrm{cont}}[1].$$

Here

$$\mathfrak{h}^\vee_{\mathrm{cont}}$$

is the continuous Taylor dual of $\mathfrak{h}$. The shift $[1]$ makes the cotangent generators odd in the relevant graded Lie algebra.

The bracket is

$$[f, g] = \{f, g\},$$

$$[f, \eta_\lambda] = \eta_{\mathrm{ad}^\vee_f \lambda},$$

$$[\eta_\lambda, \eta_\mu] = 0,$$

where

$$(\mathrm{ad}^\vee_f \lambda)(g) = -\lambda(\{f, g\}).$$

Thus the cotangent part is genuinely the coadjoint module.

The closed fields are

$$\alpha \in \Omega^\bullet(\mathbb{R}^2) \otimes \Omega^{0,\bullet}(\mathbb{C}^2) \otimes \mathfrak{h}[1],$$

$$\beta \in \Omega^\bullet(\mathbb{R}^2) \otimes \Omega^{0,\bullet}(\mathbb{C}^2) \otimes \mathfrak{h}^\vee_{\mathrm{cont}}[2].$$

The differential is

$$D = d_{\mathbb{R}^2} + \bar{\partial}_{\mathbb{C}^2}.$$

The BF action is

$$S_{\mathrm{closed}} = \int_{\mathbb{R}^2 \times \mathbb{C}^2} \left\langle \beta,\, D\alpha + \frac{1}{2}\{\alpha, \alpha\} \right\rangle dz_1\, dz_2.$$

The equation of motion from varying β is

$$D\alpha + \frac{1}{2}\{\alpha, \alpha\} = 0.$$

So α is a flat Hamiltonian connection/deformation field. The β-field is the cotangent multiplier imposing flatness.

The local observables of this closed BF theory are continuous Chevalley-Eilenberg cochains:

$$\mathrm{Obs}^{cl}_{\mathrm{closed},H} \simeq C^{\bullet}_{\mathrm{CE,cont}}(\mathfrak{g}) = S\big(\mathfrak{g}^{\vee}_{\mathrm{cont}}[-1]\big).$$

This is one of the central closed-sector claims of the paper. main

5. Why divergence-free polyvectors reduce to Hamiltonians

The paper motivates the Hamiltonian sector as a holomorphic-symplectic analogue of Kodaira-Spencer/BCOV theory.

On $\mathbb{C}^2$, polyvectors are

$$PV^{\bullet,\bullet}(\mathbb{C}^2) \simeq \Omega^{0,\bullet}(\mathbb{C}^2)[\partial_{z_1}, \partial_{z_2}].$$

The holomorphic volume form

$$\Omega = dz_1 \wedge dz_2$$

defines a divergence operator

$$\partial_\Omega : PV^i \to PV^{i-1}.$$

For a vector field

$$X = a_1\partial_{z_1} + a_2\partial_{z_2},$$

one has

$$\partial_\Omega X = \partial_{z_1} a_1 + \partial_{z_2} a_2.$$

Contracting with the symplectic form gives

$$\iota_X \omega = a_1\, dz_2 - a_2\, dz_1.$$

The equation

$$\partial_\Omega X = 0$$

is exactly

$$d(\iota_X \omega) = 0.$$

On a formal disk every closed holomorphic one-form is exact, so

$$\iota_X \omega = -df$$

for some f, unique modulo constants. Therefore

$$X = X_f.$$

So divergence-free vector fields are Hamiltonian vector fields, and Hamiltonians are defined modulo constants. This justifies replacing the divergence-free polyvector-one sector by

$$\bar{A} = \mathbb{C}[z_1, z_2]/\mathbb{C}.$$

6. The open brane sector from first principles

The brane is a product of two simple local objects:

1. an exact topological A-brane along $\mathbb{R} \subset T^*\mathbb{R}$;
2. a B-brane skyscraper sheaf $\mathcal{O}_0$ at $0 \in \mathbb{C}^2$.

The open string states are

$$\Omega^\bullet(\mathbb{R}) \otimes \mathrm{Ext}^\bullet_{\mathbb{C}[z_1,z_2]}(\mathcal{O}_0, \mathcal{O}_0) \otimes \mathfrak{gl}_N[1].$$

The Ext algebra is computed from the Koszul resolution of

$$k = \mathbb{C}[z_1, z_2]/(z_1, z_2).$$

The resolution is

$$0 \to Re_{12} \to Re_1 \oplus Re_2 \to R \to k \to 0,$$

with

$$d(e_1) = z_1, \qquad d(e_2) = z_2,$$

$$d(e_{12}) = z_2 e_1 - z_1 e_2.$$

Applying $\mathrm{Hom}_R(-,k)$ kills the differentials because z_i act by zero on k. Hence

$$\mathrm{Ext}^\bullet_R(k,k) \simeq \Lambda(\varepsilon_1,\varepsilon_2),$$

with

$$|\varepsilon_i| = 1.$$

This is the exterior algebra on the two derived normal directions.

The open BV field is

$$\alpha \in \Omega^\bullet(\mathbb{R}) \otimes \Lambda(\varepsilon_1,\varepsilon_2) \otimes \mathfrak{gl}_N[1].$$

The formal Chern-Simons action is

$$S_{\mathrm{open}} = \int_{\mathbb{R}\times\mathbb{C}^{0|2}} \frac{1}{2}\,\mathrm{Tr}(\alpha\, d\alpha) + \frac{1}{3}\,\mathrm{Tr}(\alpha^3).$$

The Berezin integral over $\mathbb{C}^{0|2}$ picks out the coefficient of

$$\varepsilon_1\varepsilon_2.$$

The ghost-zero part of the open field is

$$\alpha_0 = A + \phi_1\varepsilon_1 + \phi_2\varepsilon_2,$$

where

$$A \in \Omega^1(\mathbb{R}) \otimes \mathfrak{gl}_N,$$

$$\phi_1,\phi_2 \in \Omega^0(\mathbb{R}) \otimes \mathfrak{gl}_N.$$

The quadratic term gives, up to a boundary term,

$$\int_{\mathbb{R}} \mathrm{Tr}(\phi_1\, d\phi_2).$$

The cubic term gives

$$\int_{\mathbb{R}} \mathrm{Tr}\big(A[\phi_1,\phi_2]\big).$$

Therefore the ghost-zero open action is

$$S_{\text{open},0} = \int_{\mathbb{R}} \operatorname{Tr}\big(\phi_1\, d\phi_2 + A[\phi_1, \phi_2]\big).$$

This is a first-order constrained matrix mechanics. main

7. The Dirac interpretation of the open action

The term

$$\int \operatorname{Tr}(\phi_1\, d\phi_2)$$

means that ϕ_1 and ϕ_2 are canonically conjugate matrix coordinates. The symplectic form is

$$\Omega_N = \operatorname{Tr}(d\phi_1 \wedge d\phi_2).$$

Thus

$$\{(\phi_1)^i_j, (\phi_2)^k_l\} = \delta^i_l \delta^k_j,$$

and

$$\{(\phi_1)^i_j, (\phi_1)^k_l\} = 0, \qquad \{(\phi_2)^i_j, (\phi_2)^k_l\} = 0.$$

The field A is a Lagrange multiplier. Its equation of motion is

$$[\phi_1, \phi_2] = 0.$$

This is the moment-map equation for simultaneous conjugation by GL_N.

For $\epsilon \in \mathfrak{gl}_N$, the moment map component is

$$\mu_\epsilon = -\operatorname{Tr}\big(\epsilon[\phi_1, \phi_2]\big).$$

Its Hamiltonian flow is

$$\delta_\epsilon \phi_i = [\epsilon, \phi_i].$$

So the open phase space is the Dirac reduction

$$\{[\phi_1, \phi_2] = 0\}/GL_N.$$

In BV language, the constraint is imposed by introducing the antifield

$$\psi = A^*$$

with differential

$$Q\psi = [\phi_1, \phi_2],$$

$$Q\phi_1 = Q\phi_2 = 0.$$

So the reduced pointwise open BV algebra is

$$R_N = \mathbb{C}[(\phi_1)^i_j, (\phi_2)^i_j] \otimes \Lambda((\psi)^i_j),$$

with

$$Q\psi = [\phi_1, \phi_2].$$

This is exactly Dirac reduction written homologically.

8. Local operators and the de Rham jet contraction

Because the brane direction is topological, positive jets along the brane line are killed in cohomology.

On the jet algebra, for a coordinate t along the brane,

$$[d_{\mathbb{R}}, \iota_{\partial_t}] = \partial_t.$$

This gives an Euler-type contraction of positive jet degree. Therefore every local cohomology class has a representative depending only on field values at the insertion point:

$$\phi_1(t_0), \quad \phi_2(t_0), \quad \psi(t_0),$$

not on derivatives.

That is why the local open operator calculation reduces to the finite-dimensional invariant BV algebra

$$(R_N^{GL_N}, Q).$$

9. Stable large-N trace invariants

At finite N, invariant functions of two matrices are generated by traces of words

$$\mathrm{Tr}(w(\phi_1, \phi_2)),$$

but there are finite-N trace identities.

In the degreewise stable limit

$$N \to \infty,$$

Procesi-Razmyslov invariant theory says the invariant algebra becomes freely generated by cyclic trace words.

Let

$$F = \mathbb{C}\langle x, y\rangle,$$

and

$$\mathrm{Cyc}(F) = F/[F, F]$$

be cyclic words. Then the stable invariant algebra is

$$\mathbb{C}[\mathfrak{gl}_\infty^2]_{\mathrm{st}}^{GL_\infty} \simeq S\big(\mathrm{Cyc}(F)_{\mathrm{nonempty}}\big).$$

The empty trace

$$\mathrm{Tr}(1) = N$$

is not a primitive stable generator.

When the odd matrix ψ is included, marked cyclic words obey graded cyclicity. Moving one ψ around the trace gives no sign if it passes even letters, but two ψ's anticommute.

10. Adjacent swaps become Q-exact

The differential is

$$Q\psi = [x, y] = xy - yx.$$

Therefore

$$Q\ \mathrm{Tr}(\psi vu) = \mathrm{Tr}((xy - yx)vu).$$

By cyclicity,

$$Q\ \mathrm{Tr}(\psi vu) = \mathrm{Tr}(uxyv) - \mathrm{Tr}(uyxv).$$

So

$$\mathrm{Tr}(uxyv) - \mathrm{Tr}(uyxv)$$

is Q-exact.

Adjacent swaps of x and y are exact. Hence in cohomology, any trace word with k copies of x and l copies of y becomes the same class as

$$\mathrm{Tr}(x^k y^l).$$

Thus the ghost-zero connected stable trace classes are labelled by nonconstant commutative monomials

$$z_1^k z_2^l, \qquad k + l > 0.$$

The boundary Hamiltonian corresponding to $f \in \bar{A}$ is

$$O_f = \mathrm{Tr}\, f(\phi_1, \phi_2).$$

This is well-defined in cohomology, independent of ordering.

11. The one-ψ descendants

For each pair

$$k, l \geq 0,$$

define the first descendant

$$\Psi_{k,l} = \sum_{w \in W_{k,l}} \mathrm{Tr}(\psi\, w(x, y)),$$

where $W_{k,l}$ is the set of words with k x's and l y's.

The paper proves

$$Q\Psi_{k,l} = 0.$$

Indeed,

$$Q\Psi_{k,l} = \sum_{w \in W_{k,l}} \mathrm{Tr}((xy - yx)w).$$

Grouping by cyclic words of multidegree $(k+1, l+1)$, the number of cyclic xy-transitions equals the number of cyclic yx-transitions. Hence the sum cancels.

The primitive one-ψ homology in bidegree (k, l) is one-dimensional, generated by $\Psi_{k,l}$. The proof uses a finite combinatorial complex:

$$C_2(k,l) \to C_1(k,l) \to C_0(k,l).$$

Here

$$C_1(k,l) = \mathbb{C}\{W_{k,l}\}$$

is spanned by one-ψ words.

The target

$$C_0(k,l)$$

is spanned by cyclic words with $k+1$ x's and $l+1$ y's.

The boundary sends

$$w \mapsto [xyw]_{\mathrm{cyc}} - [yxw]_{\mathrm{cyc}}.$$

The two-ψ source supplies square relations. The resulting cell complex is interpreted as a chip-moving state complex on a circle. The Abel map sends a chip configuration to the sum of chip positions on a circle. Its fibers are convex polytopes, so the complex has the homology of S^1. Therefore the first homology is one-dimensional.

The nonconstant one-ψ descendants are

$$\Psi_{k,l}, \qquad k + l > 0.$$

The scalar descendant

$$\mathrm{Tr}(\psi) = \Psi_{0,0}$$

exists but is excluded from the Hamiltonian comparison because constants were removed from $\bar{A}$.

Thus the connected stable Hamiltonian open sector is

$$\mathrm{Obs}^{cl}_{\mathrm{brane},H} \simeq S\big(\bar{A} \oplus \bar{A}_{\mathrm{desc}}[1]\big).$$

The word "sector" matters. Higher-ψ primitive homology is not included in this Hamiltonian sector. main

12. The closed Koszul dual and the PBW special fibre

The closed observables are

$$B = C^{\bullet}_{\mathrm{CE,cont}}(\mathfrak{g}).$$

The Koszul dual is

$$B^{!} = R\,\mathrm{Hom}_B(\mathbb{C},\mathbb{C}) \simeq U(\mathfrak{g}).$$

The completed enveloping algebra has relations

$$fg - gf = \{f,g\},$$

$$f\eta_\lambda - \eta_\lambda f = \eta_{\mathrm{ad}^\vee_f\,\lambda},$$

$$\eta_\lambda\eta_\mu + \eta_\mu\eta_\lambda = 0.$$

Introduce the PBW/Rees algebra

$$U_\hbar(\mathfrak{g}) = T(\mathfrak{g})[[\hbar]] \Big/ \Big(xy - (-1)^{|x||y|}yx - \hbar[x,y]\Big).$$

At $\hbar = 0$,

$$U_\hbar(\mathfrak{g})/(\hbar) \simeq S(\mathfrak{g}) = S(\mathfrak{h} \oplus \mathfrak{h}^\vee[1]).$$

Choosing the monomial-dual basis identifies the degreewise polynomial part

$$S(\bar{A} \oplus \mathfrak{h}^\vee[1])$$

with

$$S(\bar{A} \oplus \bar{A}_{\mathrm{desc}}[1]).$$

This gives the paper's PBW/Koszul comparison:

$$\text{stable open Hamiltonian sector} \;\cong\; \text{PBW special fibre of the Koszul dual closed alg}$$

But this identification is only a filtered graded-commutative product identification. It does not identify the polynomial descendants with the coadjoint module at the deformation level. That failure is one of the main reasons the principal-parts correction is necessary.

13. The degree-zero boundary map

The physical bulk-to-boundary prescription is

$$z_i \mapsto \phi_i,$$

$$f(z_1, z_2) \mapsto \operatorname{Tr} f(\phi_1, \phi_2).$$

Thus

$$\partial_{\mathrm{bb}}(f) = O_f = \operatorname{Tr} f(\phi_1, \phi_2).$$

For monomials,

$$z_1^k z_2^l \mapsto \operatorname{Tr}(\phi_1^k \phi_2^l).$$

The open matrix Poisson bracket gives

$$\{\operatorname{Tr}(\phi_1^k \phi_2^l), \operatorname{Tr}(\phi_1^m \phi_2^n)\} = (kn - lm) \operatorname{Tr}(\phi_1^{k+m-1} \phi_2^{l+n-1}),$$

after passing to the stable connected Hamiltonian quotient.

This matches

$$\{z_1^k z_2^l, z_1^m z_2^n\} = (kn - lm) z_1^{k+m-1} z_2^{l+n-1}.$$

Therefore

$$\{O_f, O_g\}_{\mathrm{open}} = O_{\{f,g\}_{\bar{A}}}.$$

With compactly supported test functions a, b on the brane line,

$$I_\partial(a, f) = \int_{\mathbb{R}} a(t) \operatorname{Tr} f(\phi_1(t), \phi_2(t))\, dt,$$

and locality of the equal-time bracket gives

$$\{I_\partial(a, f), I_\partial(b, g)\} = I_\partial(ab, \{f, g\}_{\bar{A}}),$$

again after projection to the stable connected Hamiltonian quotient. The diagram on page 41 depicts the single canonical contraction responsible for the coefficient $kn - lm$; it is explicitly labelled as schematic, not a separate Feynman-integral proof.

main

14. The scalar central subtlety

Before quotienting constants, the simplest bracket is

$$\{\mathrm{Tr}\,\phi_1, \mathrm{Tr}\,\phi_2\} = N.$$

But in the reduced Hamiltonian algebra,

$$\{z_1, z_2\} = 1 \equiv 0 \mod \mathbb{C}.$$

So the formula

$$\{O_f, O_g\} = O_{\{f,g\}_{\bar{A}}}$$

is true only after projecting out the scalar trace class.

This is not a minor convention. It is the scalar Hamiltonian anomaly or center-of-mass central term. It reappears quantum mechanically as a Capelli/normal-ordering shift.

So the corrected statement is:

$$\{O_f, O_g\}_{\text{open}} = O_{\{f,g\}_{\bar{A}}} \quad \text{in the connected Hamiltonian quotient by scalar trace clas}$$

Without that quotient, the formula fails for $f = z_1$, $g = z_2$.

15. The CE-source obstruction

The paper observes that the map

$$f \mapsto O_f$$

is not a map from closed CE cochains in the naive way.

The closed observable algebra is

$$C^\bullet_{\mathrm{CE}}(\mathfrak{g}).$$

A Hamiltonian mode

$$f \in \mathfrak{h}$$

is not itself a CE cochain. The CE coordinate dual to f is a length-one cochain, but it is not closed.

Indeed, $\bar{A}$ is perfect as a Lie algebra:

$$\bar{A} = \{\bar{A}, \bar{A}\}.$$

For every monomial,

$$z_1^a z_2^b = \left\{ z_1, \frac{1}{b+1} z_1^a z_2^{b+1} \right\}$$

before quotienting constants. Hence every nonconstant Hamiltonian is a bracket.

If $\xi_{a,b}$ is the CE coordinate dual to $z_1^a z_2^b$, then

$$d_{\mathrm{CE}} \xi_{a,b} \neq 0.$$

In fact, it evaluates nontrivially on the pair whose bracket gives $z_1^a z_2^b$. No higher CE-length term can cancel the length-two component of $d\xi_{a,b}$. Therefore no CE cocycle has linear Hamiltonian part $\xi_{a,b}$.

So the naive idea

$$\text{closed CE coordinate dual to } f \longmapsto \mathrm{Tr}\, f(\phi_1, \phi_2)$$

is wrong.

The correction is the derived center construction.

16. The corrected closed-open derived center theorem

The right target of the closed-open map is not the ordinary open algebra. It is the derived Poisson factorization center.

At the stable degree-zero level, the open Hamiltonian algebra is

$$A_\partial = S(\mathfrak{h}),$$

with generators

$$O_I = O_{e_I}$$

for a basis e_I of $\mathfrak{h}$.

If

$$[e_I, e_J] = C_{IJ}^K e_K,$$

then

$$\{O_I, O_J\} = C_{IJ}^K O_K.$$

So A_∂ is the coordinate algebra of the linear Poisson space $\mathfrak{h}^\vee$.

The Poisson bivector is

$$\pi_\partial = \frac{1}{2} C_{IJ}^K O_K \frac{\partial}{\partial O_I} \wedge \frac{\partial}{\partial O_J}.$$

The derived P_0-center is modeled by Poisson cochains:

$$Z_{\mathrm{fact}}^{P_0}(A_\partial) \simeq \mathrm{PV}(A_\partial), \qquad d = [\pi_\partial, -].$$

Write

$$\theta^I = \frac{\partial}{\partial O_I}.$$

Then

$$\mathrm{PV}(A_\partial) = S(O_I) \otimes \Lambda(\theta^I).$$

The differential is

$$d\theta^K = -\frac{1}{2} C_{IJ}^K \theta^I \theta^J,$$

$$dO_K = C_{KJ}^L O_L \theta^J,$$

up to the global sign convention chosen for the Schouten bracket.

Now compare with the closed CE algebra of

$$\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}^\vee[1].$$

Let

$$c^I$$

be the CE coordinate dual to $e_I \in \mathfrak{h}$, and let

$$u_I$$

be the CE coordinate dual to the shifted cotangent generator $e^I[1] \in \mathfrak{h}^\vee[1]$.

Then

$$C^\bullet_{\mathrm{CE}}(\mathfrak{g}) = S(u_I) \otimes \Lambda(c^I),$$

with

$$d_{\mathrm{CE}} c^K = -\frac{1}{2} C^K_{IJ} c^I c^J,$$

$$d_{\mathrm{CE}} u_K = C^L_{KJ} u_L c^J.$$

Define

$$\Phi(c^I) = \theta^I, \qquad \Phi(u_I) = O_I.$$

Then

$$d\Phi(c^K) = \Phi(d_{\mathrm{CE}} c^K),$$

and

$$d\Phi(u_K) = \Phi(d_{\mathrm{CE}} u_K).$$

Therefore

$$C^\bullet_{\mathrm{CE}}(\mathfrak{h} \ltimes \mathfrak{h}^\vee[1]) \cong \mathrm{PV}(S(\mathfrak{h}), \pi_\partial) \simeq Z^{P_0}_{\mathrm{fact}}(S(\mathfrak{h})).$$

This is the missing cochain-level closed-open center theorem.

The crucial correction is:

$$\text{The boundary Hamiltonian } O_f \text{ is not the image of the CE coordinate dual to } f \in \mathfrak{h}$$

Instead,

$$O_f \text{ is the image of the CE coordinate dual to the shifted cotangent generator } f^\vee[1].$$

Meanwhile the CE coordinate dual to $f \in \mathfrak{h}$ maps to a vector-field operation:

$$c_f \mapsto \frac{\partial}{\partial O_f}.$$

So:

$$\text{closed Hamiltonian generator} \longmapsto \text{Hamiltonian/vector-field operation on the bran}$$

while

$$\text{closed cotangent coordinate} \longmapsto \text{boundary Hamiltonian function.}$$

This resolves the CE-source obstruction.

17. Why principal parts are necessary

Now we come to the principal-parts note.

The continuous dual of

$$\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C}$$

is not best thought of as abstract symbols $\delta_{a,b}$. Geometrically, it is realized by principal parts of meromorphic top forms.

Define

$$\rho_{a,b} = z_1^{-a-1} z_2^{-b-1}\, dz_1 \wedge dz_2, \qquad a, b \geq 0, \quad a + b > 0.$$

Let

$$P = \bigoplus_{\substack{a,b \geq 0 \\ a+b>0}} \mathbb{C}\, \rho_{a,b}.$$

The residue pairing is

$$\langle \rho_{a,b}, z_1^m z_2^n \rangle = \mathrm{Res}_0 \left(z_1^{m-a-1} z_2^{n-b-1}\, dz_1\, dz_2 \right) = \delta_{a,m} \delta_{b,n}.$$

Equivalently, integrating over a small torus $|z_1| = \epsilon_1$, $|z_2| = \epsilon_2$,

$$\frac{1}{(2\pi i)^2} \iint \rho_{a,b}\, z_1^m z_2^n = \delta_{a,m} \delta_{b,n}.$$

Thus P identifies with

$$\mathfrak{h}_{\mathrm{cont}}^{\vee}.$$

The exclusion

$$a + b > 0$$

removes

$$\rho_{0,0} = z_1^{-1} z_2^{-1}\, dz_1 dz_2,$$

which pairs with the constant 1. Since constants were removed from $\mathfrak{h}$, their dual must also be removed.

So:

$$P \simeq \mathfrak{h}_{\mathrm{cont}}^{\vee}$$

as the correct geometric dual, realized by brane-localized residue currents.

Pasted text Pasted text (2)

18. Principal parts carry the coadjoint action

The Hamiltonian action on principal parts is

$$f \cdot \rho = \pi_{\mathrm{pp}}\{f, \rho\},$$

where π_{pp} projects a Laurent two-form to its principal-part component.

For

$$f = z_1^p z_2^q,$$

and

$$\rho_{a,b} = z_1^{-a-1} z_2^{-b-1}\, dz_1 dz_2,$$

one computes

$$z_1^p z_2^q \cdot \rho_{a,b} = -\big(pb - qa + p - q\big)\rho_{a-p+1,b-q+1},$$

with negative-index targets set to zero.

This is exactly the coadjoint action because residue integration by parts gives

$$\langle f \cdot \rho, g \rangle = -\langle \rho, \{f, g\} \rangle.$$

That is the defining property of the coadjoint representation.

For the linear Hamiltonians, the corrected formulas are

$$z_1 \cdot \rho_{a,b} = -(b+1)\rho_{a,b+1},$$

$$z_2 \cdot \rho_{a,b} = (a+1)\rho_{a+1,b}.$$

So linear Hamiltonians act as raising operators on principal parts.

This is important because the copied principal-parts note contains a sentence saying z_1 lowers indices on principal parts. That sentence is incorrect. The formula in the paper is the correct one.

19. Why polynomial descendants are not the coadjoint module

The open one-ψ descendants $\Psi_{a,b}$ have the stable primitive action

$$\{O_{p,q}, \Psi_{a,b}\} = (pb - qa)\, \Psi_{a+p-1,b+q-1},$$

where $\Psi_{a,b}$ is the averaged version of $\Psi_{a,b}$.

The closed coadjoint action is

$$\mathrm{ad}^{\vee}_{z_1^p z_2^q}\, \delta_{a,b} = -\big(pb - qa + p - q\big)\delta_{a-p+1,b-q+1}.$$

These are not the same.

For example,

$$O_{1,0}$$

acts on polynomial descendants by

$$\Psi_{a,b} \mapsto b\, \Psi_{a,b-1}.$$

But z_1 acts coadjointly on principal parts by

$$\rho_{a,b} \mapsto -(b+1)\rho_{a,b+1}.$$

The direction is opposite.

Therefore:

$$\Psi_{a,b} \text{ identify with } \mathfrak{h}^\vee \text{ only as graded vector-space labels, not as the coadjoint modul}$$

The polynomial descendants are PBW-special-fibre representatives. They are not the deformation-level realization of the cotangent BF module.

The deformation-level realization is the principal-part module P.

20. No ordinary polynomial boundary realization of principal parts

The paper gives a simple obstruction.

The linear Hamiltonians

$$O_{1,0} = \operatorname{Tr} \phi_1, \qquad O_{0,1} = \operatorname{Tr} \phi_2$$

act on ordinary polynomial boundary observables by constant derivations:

$$\partial_{\phi_2}, \qquad -\partial_{\phi_1}.$$

On any ordinary polynomial, such derivations are locally nilpotent. Apply them enough times and the polynomial becomes zero.

But on principal parts,

$$O_{1,0} \cdot \rho_{a,b} = -(b+1)\rho_{a,b+1},$$

$$O_{0,1} \cdot \rho_{a,b} = (a+1)\rho_{a+1,b}.$$

These actions are not locally nilpotent. They raise the pole order forever.

Local nilpotence is preserved under module isomorphism. Therefore no collection of ordinary polynomial boundary observables can realize the principal-part coadjoint module.

That is the core mathematical reason the polynomial $\Psi_{a,b}$ cannot be the true cotangent boundary field.

21. The local-dual boundary sector

To realize the coadjoint module on the boundary, the paper introduces a formal local-dual sector.

Let

$$M_\partial^{pp}$$

be the reduced continuous dual of the stable connected Hamiltonian trace sector, with basis

$$\Theta_{a,b}$$

dual to

$$O_{m,n} = \mathrm{Tr}(\phi_1^m \phi_2^n).$$

Thus

$$\langle \Theta_{a,b}, O_{m,n} \rangle = \delta_{a,m}\delta_{b,n}.$$

Equivalently,

$$M_\partial^{pp} \simeq H^2_{(z_1,z_2)}\big(\mathbb{C}[z_1,z_2]\big)\, dz_1 dz_2 \big/ \mathbb{C}\, z_1^{-1} z_2^{-1} dz_1 dz_2.$$

Under

$$\Theta_{a,b} \leftrightarrow \rho_{a,b},$$

the Hamiltonian action is defined by negative transpose:

$$\langle \{O_f, \Theta\}, O_g \rangle = -\langle \Theta, \{O_f, O_g\} \rangle.$$

This gives

$$\{O_{p,q}, \Theta_{a,b}\} = -\big(pb - qa + p - q\big)\Theta_{a-p+1,b-q+1}.$$

Declare the odd local-dual ideal abelian:

$$\{\Theta_{a,b}, \Theta_{c,d}\} = 0.$$

Then the stalkwise assignment

$$z_1^p z_2^q \mapsto O_{p,q},$$

$$\rho_{a,b}[1] \mapsto \Theta_{a,b}$$

realizes

$$\bar{A} \ltimes P[1]$$

on the boundary.

The remaining problem is to promote this stalkwise local-dual sector to a genuine factorization object with compact-support functoriality, BV differential, pairings, and P_0-center homotopies.

The derived-center theorem explains how this should work conceptually: principal parts/local duals are the function-type cotangent coordinates, while Hamiltonian modes act as polyvector operations.

22. The Dirac interpretation of principal parts

This is the most important physical idea in the principal-parts note.

In ordinary mechanics, position and momentum are not the same kind of object.

Position can be multiplication by x. Momentum is

$$-i\hbar \frac{d}{dx}.$$

It is a derivation. It naturally acts on distributions; for example,

$$\frac{d}{dx}\delta(x - x_0)$$

is another distribution.

Trying to realize momentum as an ordinary function of position is the wrong category.

The same happens here.

Hamiltonians

$$f(z_1, z_2)$$

are holomorphic functions. Their duals are not another family of holomorphic functions. Their duals are residue currents:

$$\rho_{a,b} = z_1^{-a-1} z_2^{-b-1}\, dz_1 dz_2.$$

They are holomorphic analogues of derivatives of delta-functions supported at the brane.

That is why the brane cotangent sector must be distributional in the transverse holomorphic directions. A brane is a defect; defect-localized dual fields are naturally delta-currents normal to the defect.

So the corrected physical slogan is:

$$\text{Hamiltonian modes are functions; cotangent modes are brane-supported residue cu}$$

23. The Witten/open-closed interpretation

A bulk insertion brought to a boundary does not merely become a boundary function. It becomes a central operation on the whole open algebra.

The degree-zero shadow is

$$f \mapsto O_f = \operatorname{Tr} f(\phi_1, \phi_2).$$

But the full closed-open map should land in

$$Z^{P_0}_{\text{fact}}(\mathrm{Obs}_{\text{open}}),$$

the derived Poisson factorization center.

For the stable Hamiltonian algebra

$$A_\partial = S(\mathfrak{h}),$$

the derived center is

$$\mathrm{PV}(S(\mathfrak{h})).$$

This contains:

1. functions O_f;
2. vector fields $\partial/\partial O_f$;
3. higher polyvector operations.

The closed CE algebra

$$C^\bullet_{\text{CE}}(\mathfrak{h} \ltimes \mathfrak{h}^\vee[1])$$

is precisely this polyvector algebra.

Thus:

$$\text{closed Hamiltonian BF fields} = \text{polyvector central operations on the open brane pl}$$

The boundary Hamiltonian function O_f corresponds to the cotangent coordinate. The Hamiltonian generator f corresponds to a derivation/vector-field operation.

This “swap” is the key to the correct closed-open interpretation.

24. Three notions of locality

The documents distinguish three physical meanings of locality.

24.1 Support locality along the brane

On the brane line, observables have compact support.

If two observables are supported in disjoint intervals, their product is independent. The equal-time bracket has a delta function:

$$\{O_f(a), O_g(b)\} = O_{\{f,g\}}(ab).$$

The test functions multiply because the bracket is supported where the insertions collide.

This is ordinary canonical locality.

24.2 Topological locality

Along the topological directions,

$$L_v = [Q, \iota_v].$$

Moving insertions does not change cohomology unless insertions collide. Therefore the factorization algebra is locally constant in those directions.

This is why the brane line behaves like an E_1-algebra after passing to cohomology.

24.3 Formal holomorphic locality

In the holomorphic normal directions, the model is completed at the brane point. It is a theory of formal holomorphic jets, not a theory of finite-speed propagation through $\mathbb{C}^2$.

Antiholomorphic translations are Q-exact:

$$L_{\partial/\partial\bar{z}_i} = [Q, \iota_{\partial/\partial\bar{z}_i}].$$

The principal parts are local in this formal/distributional sense. They are residue currents supported at the brane, not smooth fields spread through $\mathbb{C}^2$.

So the cotangent sector is:

$$\text{distributionally local transverse to the brane.}$$

25. Unitarity

This model is not a Lorentzian unitary quantum field theory.

It is complex, BV, ghost-valued, topological-holomorphic, and local/formal. It has no positive-definite Hilbert space built into it, no ordinary time-evolution unitarity, and no reflection positivity statement.

The correct consistency condition is the classical master equation

$$QI + \frac{1}{2}\{I, I\} = 0,$$

and quantum mechanically the quantum master equation

$$QI + \frac{1}{2}\{I, I\} + \hbar\Delta I = 0.$$

So in this paper, “physical consistency” means:

$$\text{BV master equation} + \text{factorization/OPE associativity} + \text{anomaly cancellation if c}$$

If this model is a protected twisted sector of a larger unitary theory, unitarity belongs to the parent theory, not to the local twisted sector by itself.

26. The quantum deformation target

The degree-zero Hamiltonian sector has a canonical deformation quantization: the Weyl/Moyal algebra of the formal symplectic disk.

The Moyal product is

$$f * g = m \circ \exp\left(\frac{\hbar}{2}\left(\partial_{z_1} \otimes \partial_{z_2} - \partial_{z_2} \otimes \partial_{z_1}\right)\right)(f \otimes g).$$

Let

$$P = \partial_{z_1} \otimes \partial_{z_2} - \partial_{z_2} \otimes \partial_{z_1}.$$

Then

$$[f, g]_* = f * g - g * f = \hbar P(f, g) + \frac{\hbar^3}{24} P^3(f, g) + O(\hbar^5).$$

The normalized quantum Hamiltonian bracket is

$$\{f, g\}_\hbar = \hbar^{-1}[f, g]_*.$$

For monomials

$$f = z_1^k z_2^l, \qquad g = z_1^m z_2^n,$$

the r-fold bidifferential is

$$P^r(f, g) = \sum_{a=0}^{r} (-1)^a \binom{r}{a} (k)_{r-a} (l)_a (m)_a (n)_{r-a} z_1^{k+m-r} z_2^{l+n-r},$$

where

$$(s)_q = s(s-1)\cdots(s-q+1).$$

Only odd r contribute to the commutator.

Thus

$$[f, g]_* = \sum_{\substack{r \geq 1 \\ r \text{ odd}}} \frac{2}{2^r r!} \hbar^r P^r(f, g).$$

The first coefficients are

$$\hbar P(f, g),$$

$$\frac{\hbar^3}{24} P^3(f, g),$$

$$\frac{\hbar^5}{1920} P^5(f,g),$$

$$\frac{\hbar^7}{322560} P^7(f,g).$$

The diagrams on pages 50 and 51 depict the order-$\hbar^3$ boundary contraction types whose tensor target is $P^3(f,g)$. They are schematic; the paper explicitly says the analytic Costello graph weights, automorphism factors, and counterterms remain to be identified. main

27. Quantum matrix reduction and the Capelli obstruction

On the open side, quantizing

$$\int \mathrm{Tr}(\phi_1 d\phi_2)$$

gives the Weyl algebra generated by matrix entries

$$X = \phi_1, \qquad Y = \phi_2,$$

with

$$[X^i_j, Y^k_l] = \hbar\, \delta^i_l \delta^k_j.$$

The quantum comoment for conjugation is normal ordered:

$$\mu_\epsilon = -\,\mathrm{Tr}([\epsilon, X]Y).$$

It satisfies

$$\hbar^{-1}[\mu_\epsilon, a] = \epsilon \cdot a.$$

The quantum brane algebra is the BRST quantum Hamiltonian reduction

$$\mathrm{Obs}^q_{\mathrm{brane},N} = C^\bullet_{\mathrm{BRST}}(\mathfrak{gl}_N, W_N, \mu).$$

Weyl-ordered traces

$$J_N(f) = \mathrm{Tr}\,\mathrm{Op}_W(f)(X,Y)$$

are invariant and descend through quantum Hamiltonian reduction.

But this does not prove the full quantum Moyal identity. The obstruction is the normal-ordering shift:

$$\mu_\epsilon = \mathrm{Tr}\big(\epsilon(YX - XY + \hbar N I)\big).$$

So the quantum moment relation is

$$YX - XY + \hbar N I = 0.$$

Therefore

$$\mathrm{Tr}(AXY) - \mathrm{Tr}(AYX) = \hbar N \,\mathrm{Tr}(A)$$

modulo the quantum moment ideal.

This is a Capelli-type shift. It can feed lower-degree traces into the connected sector. A full quantum theorem must either cancel, renormalize, or correctly account for this shift.

So the quantum part of the paper gives the algebraic Weyl/Moyal target, not a completed quantum closed-open theorem.

28. Costello graph problem

Costello's renormalized BV formalism supplies a general graph/RG framework. But the paper does not yet identify this Hamiltonian BF brane defect with a fully controlled Costello graph construction.

The missing analytic data include:

1. a precise elliptic closed-open parent theory or reduction;
2. boundary/defect conditions for $\mathbb{R} \times \{0\}$;
3. gauge fixing and propagator;
4. anomaly cancellation;
5. the bulk-to-defect kernel;
6. the relation between Costello's propagator and the Weyl contraction $P/2$;
7. first-order graph normalization;
8. third-order graph normalization giving $1/24$;
9. connected trace extraction and treatment of the Capelli shift.

The formal algebra says the boundary product should be

$$m \circ \exp\left(\frac{\hbar}{2}P\right).$$

The graph problem is to prove that the actual renormalized BV observable product in this model gives exactly that product.

29. The corrected global thesis

The corrected paper should have two main theorems, not one overloaded theorem.

Theorem A: PBW stable trace theorem

The stable connected Hamiltonian open trace sector is

$$\mathrm{Obs}^{cl}_{\mathrm{brane},H} \simeq S(\bar{A} \oplus \bar{A}_{\mathrm{desc}}[1]).$$

The closed Hamiltonian BF observables are

$$C^{\bullet}_{\mathrm{CE,cont}}(\mathfrak{h} \ltimes \mathfrak{h}^{\vee}[1]).$$

Their Koszul dual is

$$U(\mathfrak{h} \ltimes \mathfrak{h}^{\vee}[1]).$$

The PBW special fibre is

$$S(\mathfrak{h} \oplus \mathfrak{h}^{\vee}[1]).$$

Its degreewise polynomial part matches the stable open Hamiltonian sector after identifying $\mathfrak{h}^{\vee}$ with the one-ψ descendant label set.

But the polynomial descendants do not realize the coadjoint action.

Theorem B: derived closed-open center theorem

The closed CE algebra itself is the derived Poisson center of the stable degree-zero open Hamiltonian algebra:

$$C^{\bullet}_{\mathrm{CE}}(\mathfrak{h} \ltimes \mathfrak{h}^{\vee}[1]) \cong Z^{P_0}_{\mathrm{fact}}(S(\mathfrak{h})).$$

Under this isomorphism,

$$c^I \mapsto \frac{\partial}{\partial O_I},$$

$$u_I \mapsto O_I.$$

This is the cochain-level closed-open structure.

Theorem C: principal-parts cotangent realization

The correct deformation-level boundary model for $\mathfrak{h}^\vee$ is the principal-part module

$$P = \bigoplus_{a+b>0} \mathbb{C}\,\rho_{a,b},$$

with

$$\rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2.$$

The action is

$$z_1^p z_2^q \cdot \rho_{a,b} = -\big(pb - qa + p - q\big)\rho_{a-p+1,b-q+1}.$$

This is the coadjoint action. It cannot be realized by ordinary polynomial boundary observables.

Together, these three theorems make the physics natural:

$$\begin{gathered}\text{The brane is a constrained matrix probe.}\\ \text{The closed Hamiltonian BF theory is the cotangent theory of canonical transforma}\\ \text{Closed observables are derived Poisson central operations on the open probe alge}\\ \text{Cotangent modes are transverse residue currents localized at the brane.}\end{gathered}$$

30. Errata and corrections

Here are the corrections I would make explicitly.

1. Principal-part linear action

The principal-parts note says, in one place, that acting by z_1 lowers indices on principal parts. That is wrong.

The correct formula is

$$z_1 \cdot \rho_{a,b} = -(b+1)\rho_{a,b+1},$$

$$z_2 \cdot \rho_{a,b} = (a+1)\rho_{a+1,b}.$$

So linear Hamiltonians raise pole order on principal parts.

2. Polynomial descendants do not simply “raise indices”

The polynomial descendant action is

$$O_{p,q} : \Psi_{a,b} \mapsto (pb - qa)\Psi_{a+p-1,b+q-1}.$$

Whether this raises, lowers, or preserves total degree depends on $p + q$:

$$\deg_{\text{target}} = a + b + p + q - 2.$$

Linear Hamiltonians lower total degree by one, quadratic Hamiltonians preserve total degree, and higher Hamiltonians raise it.

The real point is not “raising versus lowering” in general. The real point is that the action is not the coadjoint action.

3. The coadjoint action formula is correct in the paper

The correct coadjoint/principal-part formula is

$$z_1^p z_2^q \cdot \rho_{a,b} = -\big(pb - qa + p - q\big)\rho_{a-p+1,b-q+1}.$$

This should be treated as the reference formula.

4. The smeared bracket formula needs the scalar quotient

The formula

$$\{I_\partial(a, f), I_\partial(b, g)\} = I_\partial(ab, \{f, g\}_{\bar{A}})$$

must be read in the connected Hamiltonian quotient by scalar trace classes.

Otherwise, for $f = z_1$, $g = z_2$,

$$\{I_\partial(a, z_1), I_\partial(b, z_2)\} = N \int a(t)b(t)\, dt,$$

while

$$\{z_1, z_2\}_{\bar{A}} = 0.$$

So the scalar central term must be projected out.

5. “Removing constants” is not literally the same as removing all $U(1)$ center-of-mass modes

The constant Hamiltonian corresponds to the scalar empty trace

$$\mathrm{Tr}(1) = N.$$

The center-of-mass pair

$$\mathrm{Tr}\,\phi_1, \qquad \mathrm{Tr}\,\phi_2$$

still exists as linear observables, but their bracket produces the scalar N. So the precise statement is:

$$\text{we project out the scalar Hamiltonian/empty-trace central term,}$$

not that we remove the entire $U(1)$ center-of-mass sector.

6. The “full closed-open map remains a problem” can now be sharpened

The paper currently lists the full CE/cotangent P_0 factorization-center lift as a problem. The corrected mathematical construction is:

$$C^\bullet_{\mathrm{CE}}(\mathfrak{h} \ltimes \mathfrak{h}^\vee[1]) \cong \mathrm{PV}(S(\mathfrak{h})) \simeq Z^{P_0}_{\mathrm{fact}}(S(\mathfrak{h})).$$

The remaining work is not the abstract stalkwise derived-center isomorphism. The remaining work is to implement it inside the full unfrozen BV/factorization model, with compact supports, de Rham/Dolbeault fields, defect kernels, and the principal-part boundary realization.

7. References and labels need cleanup

The PDF contains malformed reference labels such as “[13]razmyslov” and “[2]tsygan,” and a PBW citation appears to point to the wrong bibliography number. These should be cleaned before circulation.

8. The figures should either be upgraded or demoted

The diagrams on pages 41, 50, and 51 are useful only as schematic reminders. They should not look like proofs. Either label them with propagators, boundary vertices, automorphism factors, and orientation conventions, or keep them explicitly as informal pictures.

Sources

Sources

main(4).pdf
PDF

here's the latest version of my WIP whitepaper. Attack it from all angles, and as before, suggest exactly how to level it up into the strongest version of this whitepaper conceivable

I'll treat this like a hostile referee plus a constructive coauthor: first identifying what now works, then isolating fatal vulnerabilities, then giving a concrete upgrade path.

we care about strong physics
we care about groundbreaking mathematics
we care about their synthesis
we care about advancing human knowledge

The draft's core has advanced; I'll now focus on overclaimed upgrades: factorization source conventions, principal-part interpolation, finite-N quantum radial parts, and weighted Tate/Costello foundations.

I'll raise the standard from safety to inevitability: the physics principle and theorem architecture must force the ambition, not merely gesture at it.

Thought for 8m 20s

According to the document from 2026-04-28, the latest draft is **much closer to the real paper**: it now has the Dirac probe principle, the CE/PV derived-center theorem, the principal-part correction, the scalar anomaly, and a careful analytic firewall around Costello/QME issues. The strongest version is not a smaller paper. It is a **more sharply stratified paper** where the true theorem, the deformation bridge, and the analytic quantum problem are separated so that each can be attacked and proved at maximal strength. main

My hostile verdict is this:

$$\text{The physics thesis is now excellent. The central CE/PV mathematics is right.}$$

But there are still several places where the draft **reintroduces the old obstruction under new names**. The most serious problem is that parts of the factorization and primitive-projection story still send the **wrong CE generator** or the **wrong open descendant module** to the principal-part/coadjoint sector. If you fix that, the paper becomes dramatically stronger.

1. What the paper is really trying to prove

The strongest thesis is:

$$\text{A Dirac-reduced brane probe of a formal symplectic disk has closed Hamiltonian B}$$

More explicitly:

$$\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C} \cdot 1$$

is the Lie algebra of infinitesimal canonical transformations of the formal symplectic disk. The closed field theory is the shifted cotangent BF theory

$$\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}^\vee_{\mathrm{cont}}[1].$$

The degree-zero open brane phase algebra is

$$A_\partial = S(\mathfrak{h}),$$

with Kirillov-Poisson bracket

$$\{O_f, O_g\} = O_{\{f,g\}}.$$

Then the real theorem is

$$C^\bullet_{\mathrm{CE,cont}}(\mathfrak{h} \ltimes \mathfrak{h}^\vee_{\mathrm{cont}}[1]) \cong \mathrm{PV}(S(\mathfrak{h})) \simeq Z^{P_0}_{\mathrm{fact}}(S(\mathfrak{h})).$$

The draft now states this correctly at the cochain level: the CE coordinate c^I dual to a Hamiltonian generator maps to the constant polyvector $\theta^I = \partial/\partial O_I$, while the CE coordinate u_I dual to the shifted cotangent generator maps to the boundary Hamiltonian function O_I. This is the key breakthrough. main

Physically, this says:

$$\text{closed Hamiltonian mode} \longmapsto \text{Hamiltonian vector-field operation on the brane},$$

while

$$\text{closed cotangent coordinate} \longmapsto \text{boundary trace observable}.$$

That is the Dirac-Witten synthesis. The brane does not merely receive closed functions. Closed fields become **central operations** on open measurements.

2. The central mathematical inconsistency still present

The draft correctly says that the boundary observable

$$O_f = \operatorname{Tr} f(\phi_1, \phi_2)$$

is **not** the image of the CE coordinate dual to $f \in \mathfrak{h}$. It is the image of the CE coordinate dual to the shifted cotangent generator. The text even emphasizes that $O_I = \Phi(u_I)$, not $\Phi(c^I)$, and that $d_{\mathrm{CE}} u_K$ maps to the Hamiltonian vector field $d_\pi O_K$. main

But the interval-wise factorization theorem still appears to map a length-one CE coordinate

$$(a \otimes f)^\vee[-1]$$

to the boundary current

$$B_f(a) = \int_I a(t) \operatorname{Tr} f(\phi_1(t), \phi_2(t))\, dt.$$

That is the old mistake in factorization form. The source for $B_f(a)$ cannot be the CE coordinate dual to the Hamiltonian current $a \otimes f \in \mathfrak{h}_I$. It must be the CE coordinate dual to the **shifted cotangent current** paired with $a \otimes f$. The umbrella theorem's factorization statement currently phrases the Hamiltonian factor as though the boundary function is supplied by the Hamiltonian CE coordinate, even though the cochain theorem says the opposite. main

The corrected interval theorem should be:

$$\mathfrak{h}_I = \Omega_c^\bullet(I) \otimes \mathfrak{h}, \qquad \mathfrak{g}_I = \mathfrak{h}_I \ltimes \mathfrak{h}_I^\vee[1].$$

Then

$$C^{\bullet}_{\mathrm{CE}}(\mathfrak{g}_I) \cong \mathrm{PV}(S(\mathfrak{h}_I)).$$

For $x = a \otimes f \in \mathfrak{h}_I$, define a coordinate u_x dual to the shifted cotangent generator corresponding to x. Then the current map is

$$u_{a\otimes f} \longmapsto B_f(a).$$

For $\lambda \in \mathfrak{h}_I^\vee$, the Hamiltonian CE coordinate should map to the constant polyvector

$$c_\lambda \longmapsto \theta_\lambda.$$

This one correction makes the factorization theorem consistent with your own CE-source obstruction.

3. The second serious inconsistency: $\Psi_{a,b} \to \rho_{a,b}$ cannot be a strict P_0-morphism

The draft correctly distinguishes polynomial one-ψ descendants from principal parts. It says the polynomial descendants carry the symmetric action

$$F_{p,q} : \Psi_{a,b} \mapsto (pb - qa)\Psi_{a+p-1,b+q-1},$$

whereas the principal-part/coadjoint action is

$$z_1^p z_2^q \cdot \rho_{a,b} = -(pb - qa + p - q)\rho_{a-p+1,b-q+1}.$$

The draft even states that the two actions agree only in the trivial Hamiltonian case and have opposite index-shift behavior. main

But later the chain-level primitive projection sends

$$\Psi_{a,b} \mapsto \rho_{a,b}[1]$$

and claims strict P_0-bracket compatibility with the principal-part coadjoint bracket. main

That cannot be true as stated.

Take the simplest test:

$$f = z_1.$$

On polynomial descendants,

$$\{O_{1,0}, \Psi_{a,b}\} = b\,\Psi_{a,b-1}.$$

So after the proposed projection,

$$\pi^{\mathrm{cot}}_{\mathrm{prim}}\{O_{1,0}, \Psi_{a,b}\} = b\,\rho_{a,b-1}.$$

But on principal parts,

$$z_1 \cdot \rho_{a,b} = -(b+1)\rho_{a,b+1}.$$

So

$$\{O_{1,0}, \pi^{\mathrm{cot}}_{\mathrm{prim}}(\Psi_{a,b})\} = -(b+1)\rho_{a,b+1}.$$

These are not equal:

$$b\,\rho_{a,b-1} \neq -(b+1)\rho_{a,b+1}.$$

So the map

$$\Psi_{a,b} \mapsto \rho_{a,b}$$

cannot be a strict P_0-morphism. This is not a small sign error. It is the exact structural mismatch that your own principal-part section correctly identifies.

The fix is decisive:

$$\Psi_{a,b} \text{ belongs to the PBW special-fibre open descendant sector.}$$

$$\rho_{a,b} \text{ belongs to a new distributional/principal-part boundary sector.}$$

Do **not** define the principal-part sector as a strict P_0-quotient of the polynomial Ψ-sector unless you construct a genuinely nontrivial intertwiner. The naive map $\Psi_{a,b} \mapsto \rho_{a,b}$ is not one.

The right statement is:

$$S(\bar{A} \oplus \bar{A}_{\mathrm{desc}}[1]) \cong \mathrm{gr}_{\mathrm{PBW}}\, U(\mathfrak{h} \ltimes \mathfrak{h}^\vee[1])$$

as a graded-commutative PBW special-fibre algebra, while the deformation-level cotangent boundary realization is

$$S(\bar{A} \oplus P[1]),$$

with independent generators

$$\Theta_\rho.$$

The boundary brackets are then defined by

$$\{B_f, \Theta_\rho\} = \Theta_{f\cdot\rho}, \qquad \{\Theta_\rho, \Theta_\sigma\} = 0.$$

That gives the correct coadjoint module. But Θ_ρ is not an ordinary polynomial Ψ-class.

4. The “interpolation” theorem is not yet a theorem

The draft says the polynomial $\Psi_{a,b}$ basis and the principal-part $\rho_{a,b}$ basis are “two different bases of the same $\mathfrak{g}$-module,” interpolated by the Rees/Moyal parameter $\hbar$. But the same section also says the PBW special-fibre map is not an $\mathfrak{h}$-module isomorphism and that the actions diverge explicitly. main

Those two statements cannot both be literally true.

If two bases are bases of the **same** $\mathfrak{h}$-module, the matrices of the action must be conjugate. But the polynomial action is locally nilpotent under linear Hamiltonians, while the principal-part action is not. Your own no-go theorem says ordinary polynomial boundary observables cannot realize the principal-part module precisely because local nilpotence is preserved under module isomorphism. main

So the corrected statement should be:

$$\Psi_{a,b} \text{ and } \rho_{a,b} \text{ are not two bases of the same } \mathfrak{h}\text{-module.}$$

They are two different realizations of the same **label set** or the same **PBW/cotangent formal symbol set**:

$$\bar{A}_{\mathrm{desc}} \cong \mathfrak{h}^\vee_{\mathrm{cont}}$$

as graded vector spaces, but not as $\mathfrak{h}$-modules.

To make the interpolation claim real, you need to construct an actual $\hbar$-dependent module

$$\mathcal{M}_\hbar$$

over a Rees/Weyl algebra, together with two specializations:

$$\mathcal{M}_\hbar/(\hbar) \cong \text{polynomial PBW descendant symbols},$$

and

$$\mathcal{M}_\hbar[\hbar^{-1}] \cong \text{principal-part coadjoint module}.$$

This is not in the draft yet. The natural candidate is a **Fourier-Borel/Rees $D_\hbar$-module** at the brane:

$$D_\hbar = \mathbb{C}\langle z_1, z_2, \hbar\partial_{z_1}, \hbar\partial_{z_2}\rangle, \qquad [\hbar\partial_{z_i}, z_j] = \hbar\delta_{ij}.$$

Then take a delta-module or local-cohomology module

$$\mathcal{M}_\hbar = D_\hbar / D_\hbar(z_1, z_2),$$

or its holomorphic local-cohomology realization

$$H^2_{(z_1,z_2)}(\mathbb{C}[z_1, z_2])\, dz_1 dz_2.$$

The Hamiltonian vector fields act by Lie derivative. The principal parts are the $\hbar \neq 0$ residue-current realization. The associated graded is a cotangent-fiber symbol module. If you can identify the polynomial Ψ-action as the Fourier/Borel transform of that associated graded, you will have a genuine interpolation theorem.

Without that construction, replace "interpolated by $\hbar$" with:

> The $\Psi_{a,b}$ and $\rho_{a,b}$ bases are two distinct realizations of the same PBW label set. A genuine Rees/Fourier interpolation between them is an additional theorem, formulated below.

That would be mathematically honest and far stronger.

5. The factorization algebra must use de Rham compact supports, not just $C_c^\infty(I)$

The paper wants locality along the brane line to be a factorization algebra statement. That is right. But several interval-wise definitions use

$$\mathfrak{h}_I = C_c^\infty(I) \otimes \mathfrak{h}$$

as if this were locally constant under inclusions of intervals. It is not. There is no differential in $C_c^\infty(I)$, so inclusions of intervals are not quasi-isomorphisms.

The correct object is

$$\mathfrak{h}_I = \Omega_c^\bullet(I) \otimes \mathfrak{h},$$

with differential d_{dR}, and bracket

$$[\alpha \otimes f, \beta \otimes g] = (\alpha \wedge \beta) \otimes \{f, g\}.$$

Then compactly supported de Rham cohomology of an interval is one-dimensional in top degree, and extension by zero is the correct cosheaf operation. The degree-zero current formula

$$B_f(a) = \int_I a(t) \operatorname{Tr} f(\phi_1(t), \phi_2(t))\, dt$$

is then the shadow of a top-degree compactly supported form

$$a(t)\, dt \otimes f.$$

This resolves the physical locality correctly:

$$\text{support locality} + \text{topological de Rham contraction} = \text{locally constant factorizatior}$$

The draft already knows the de Rham contraction is essential, but the theorem statements should bake it into the definitions rather than add it afterward. main

6. The quantum radial-parts theorem is potentially huge, but currently underproved

The draft now claims an all-N reduced Moyal commutator theorem using a Harish-Chandra radial-parts identity:

$$\mathrm{Rad}_0(J_N(f)) = \Delta^{-1} S_N(f) \Delta,$$

and concludes that

$$D_N(f, g) = \hbar^{-1}[J_N(f), J_N(g)] - J_N(\{f, g\}_\hbar)$$

lies in the quantum moment-map ideal. main

If true, this is one of the most powerful results in the paper. It would upgrade the quantum section from “formal Moyal target” to a genuine quantum Hamiltonian reduction theorem.

But this proof needs much more armor. A hostile reader will ask:

1. Which exact radial-parts theorem is being invoked?
2. Does it apply to all Weyl-ordered mixed traces $\mathrm{Tr}\,\mathrm{Op}_W(f)(X,Y)$, not just invariant functions or constant-coefficient invariant differential operators?
3. Are there Calogero-Moser/Dunkl correction terms after conjugation by Δ?
4. Does the radial map remain injective on the precise quantum Hamiltonian reduction quotient you use?
5. Is the normalizer quotient by the moment ideal being taken before or after GL_N-invariants?
6. Does the connected single-trace projection commute with the radial-parts identification?
7. Does the stable $N \to \infty$ cumulant extraction preserve the bracket identity, or can triangular Capelli terms feed lower connected traces?

The proof sketch by “pure powers and Hamiltonian shears generate everything” is suggestive, but not yet theorem-level. main

To level this up, create a standalone theorem:

Theorem: Radial-Moyal theorem for Weyl traces.
Let $W_N = D_\hbar(\mathfrak{gl}_N)$, let I_N be the quantum moment ideal for conjugation, and define

$$J_N(f) = \mathrm{Tr}\,\mathrm{Op}_W(f)(X,Y).$$

Then in

$$N_{W_N}(I_N)/I_N$$

one has

$$\hbar^{-1}[J_N(f), J_N(g)] = J_N(\{f,g\}_\hbar)$$

for all $f, g \in \mathbb{C}[z_1, z_2][[\hbar]]$, modulo the scalar Capelli class.

The proof should proceed by identifying the reduced algebra with a spherical/Dunkl/radial algebra and proving that the images of $J_N(f)$ are precisely sums

of one-particle Weyl operators. If correction terms appear, compute them and show they vanish in the connected scalar-reduced quotient.

This theorem is worth an appendix by itself. Do not bury it in the quantum section.

7. The weighted Tate completion is a regulator, not the original dual

The weighted Tate section is intelligent, but the paper must state more bluntly what it does. The draft defines

$$\mathfrak{h}_w = \lim_{w_0} \prod_{d \le w_0} H_d^{(w)}$$

and

$$\mathfrak{h}_{\mathrm{cont}}^{\vee,w} = \lim_{w_0} \prod_{d \le w_0} (H_d^{\vee})^{(w^{-1})}.$$

It explicitly says the second factor is a product coefficient module, **not** the strict continuous dual of the first product Tate space. main

That is exactly right, but it has a consequence:

$$\text{The weighted theory is a regulated coefficient theory, not literally the original stric}$$

The weighted Casimir convergence theorem is therefore not “we fixed the original Tate dual.” It is:

$$\text{we changed the coefficient category so the BV coevaluation exists.}$$

That is acceptable, and physically natural as a regulator. But then the paper needs a regulator-independence statement or a clear claim hierarchy:

1. **Strict algebraic closed theory:** $\mathfrak{h} \ltimes \mathfrak{h}_{\mathrm{cont}}^{\vee}[1]$, with $\mathfrak{h}_{\mathrm{cont}}^{\vee}$ the direct-sum principal-part module.
2. **Weighted analytic regulator:** $\mathfrak{h}_w \ltimes \mathfrak{h}_{\mathrm{cont}}^{\vee,w}[1]$, where both sides are product-type Mittag-Leffler modules.
3. **Comparison problem:** show observables or amplitudes of interest are independent of w, or have a controlled $q \to 1^+$ limit.

Without that third step, weighted Tate completion is not a completion of the original object. It is a new analytic theory with a canonical embedding of the polynomial/principal-part sector.

The draft already admits the QME obstruction remains after the Hadamard/Mittag-Leffler analysis. Keep that honesty. The Hadamard theorem reduces the wavefront and coefficient-topology problem, but does not prove the mixed brane-defect QME obstruction vanishes. main

8. The principal-part module needs a D-module proof, not "Schur rigidity"

The principal-part correction is one of the most beautiful parts of the paper:

$$P = \bigoplus_{a+b>0} \mathbb{C}\,\rho_{a,b}, \qquad \rho_{a,b} = z_1^{-a-1} z_2^{-b-1}\, dz_1 dz_2.$$

The residue pairing identifies P with the continuous dual of $\mathfrak{h}$, and the coadjoint action is

$$z_1^p z_2^q \cdot \rho_{a,b} = -(pb - qa + p - q)\rho_{a-p+1,b-q+1}.$$

This is the correct deformation-level cotangent module. main

But the uniqueness proof via "Schur rigidity" and dense orbit is too hand-wavy. Infinite-dimensional modules over infinite-dimensional Lie algebras are treacherous. Replace it with a D-module theorem:

$$P \cong H^2_{(z_1,z_2)}(\mathbb{C}[z_1, z_2])\, dz_1 dz_2 / \mathbb{C}\rho_{0,0}.$$

Then prove:

$$\mathrm{End}_{\mathfrak{h}}(P) = \mathbb{C}$$

or prove the weaker statement actually needed:

> The residue pairing is the unique continuous degree-preserving pairing compatible with Hamiltonian integration by parts, after fixing $\langle \rho_{a,b}, z_1^a z_2^b \rangle = 1$.

The proof should be residue calculus:

$$\langle f \cdot \rho, g \rangle = -\langle \rho, \{f, g\} \rangle.$$

That is enough. You do not need “Schur rigidity” unless you really prove irreducibility.

9. The scalar anomaly section is strong, but phrase it as a distinguished line

The scalar anomaly is now one of the best physical pieces. The draft identifies the cocycle

$$\omega(z_1^k z_2^l, z_1^m z_2^n) = (kn - lm)\,\mathbf{1}_{(k+m,l+n)=(1,1)}$$

with the open central bracket

$$\{\mathrm{Tr}\,\phi_1, \mathrm{Tr}\,\phi_2\} = N$$

and with the quantum Capelli shift

$$YX - XY + \hbar N I \equiv 0.$$

That is the right physical story: the scalar Hamiltonian is the $U(1)$ center-of-mass anomaly. main

The only wording to avoid is “the one-dimensional sub-line of $H^2_{\mathrm{Lie}}(\bar{A}, \mathbb{C})$” if it sounds like H^2 itself is one-dimensional. Unless you compute all of $H^2_{\mathrm{Lie}}(\bar{A}, \mathbb{C})$, say:

$$[\bar{c}] \text{ is the distinguished one-dimensional anomaly line in } H^2_{\mathrm{Lie}}(\bar{A}, \mathbb{C}).$$

That is safer and physically sharper.

10. The chain-level primitive projection should be renamed

The theorem currently says the primitive projection

$$\pi^{\mathrm{cot}}_{\mathrm{prim}} : \mathrm{Obs}^{cl,pp}_{\partial,N} \twoheadrightarrow A^{pp}_{\partial}$$

is a surjection to an “indecomposable P_0 quotient” and explicitly says multiplication of two positive primitive generators is zero after projection. main

Then it should **not** be presented as a factorization algebra equivalence or as a morphism to A^{pp}_{∂} with its ordinary commutative multiplication.

The right name is:

$$\text{primitive Lie}/P_0 \text{ quotient projection}$$

not

$$\text{cotangent factorization-center lift.}$$

A factorization algebra remembers products. If your quotient kills products of positive primitives, it is not the same algebraic object as the factorization algebra generated by those primitives. It is the **indecomposable Lie shadow** of that algebra.

This is still useful. But call it what it is.

11. The physics can be made much stronger

The paper should not sound like it is borrowing physics decorations from string theory. It should sound like it is deriving the structure from three physical principles.

Principle 1: Dirac constrained probes

The brane stack probes the formal symplectic disk. The normal coordinates become matrices:

$$z_i \rightsquigarrow \phi_i.$$

The symplectic potential is forced:

$$\theta_N = \operatorname{Tr}(\phi_1\, d\phi_2).$$

Conjugation invariance gives the moment map

$$\mu_\epsilon = -\operatorname{Tr}(\epsilon[\phi_1, \phi_2]).$$

The first-order action is therefore

$$S_\partial = \int_{\mathbb{R}} \operatorname{Tr}\big(\phi_1 d\phi_2 + A[\phi_1, \phi_2]\big).$$

The BV variable

$$\psi = A^*$$

records the constraint:

$$Q\psi = [\phi_1, \phi_2].$$

This is not an analogy. It is Dirac reduction in BV form.

Principle 2: Witten closed-open centrality

A closed insertion near a brane is not merely a boundary value. It is a central operation on all open measurements. Therefore the target is not the open algebra but

$$Z^{P_0}_{\mathrm{fact}}(\mathrm{Obs}_{\mathrm{open}}).$$

The trace

$$f \mapsto \mathrm{Tr}\, f(\phi_1, \phi_2)$$

is only the moment-map shadow.

Principle 3: Cotangent fields are transverse currents

The dual of a holomorphic Hamiltonian is not another holomorphic Hamiltonian. It is a residue current:

$$\rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2.$$

This is the holomorphic analogue of the fact that momentum is not multiplication by a function but a distributional/derivative operation. The brane is a defect, so cotangent boundary modes are naturally transverse delta-currents.

These three principles should be the first section. They make factorization algebra feel physically inevitable.

12. The strongest possible theorem package

I would reorganize the paper around five main theorems.

Theorem A: Dirac brane probe theorem

For the mixed brane

$$\mathbb{R} \times \{0\} \subset \mathbb{R}^2 \times \mathbb{C}^2,$$

the ghost-zero open BV action is the Dirac first-order constrained system

$$S_\partial = \int_{\mathbb{R}} \mathrm{Tr}(\phi_1 d\phi_2 + A[\phi_1, \phi_2]),$$

and the reduced local BV algebra is

$$R_N = \mathbb{C}[\phi_1, \phi_2] \otimes \Lambda(\psi), \qquad Q\psi = [\phi_1, \phi_2].$$

Theorem B: Derived center theorem

Let

$$A_\partial = S(\mathfrak{h}), \qquad \mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}^\vee_{\mathrm{cont}}[1].$$

Then

$$C^\bullet_{\mathrm{CE,cont}}(\mathfrak{g}) \cong \mathrm{PV}(A_\partial) \simeq Z^{P_0}_{\mathrm{fact}}(A_\partial).$$

On generators:

$$c_\lambda \mapsto \theta_\lambda, \qquad u_x \mapsto O_x.$$

This theorem should be the central mathematical result.

Theorem C: Defect current factorization theorem

For intervals $I \subset \mathbb{R}$,

$$\mathfrak{h}_I = \Omega^\bullet_c(I) \otimes \mathfrak{h}, \qquad \mathfrak{g}_I = \mathfrak{h}_I \ltimes \mathfrak{h}_I^\vee[1],$$

and

$$C^\bullet_{\mathrm{CE}}(\mathfrak{g}_I) \cong Z^{P_0}_{\mathrm{fact}}(S(\mathfrak{h}_I)).$$

The current

$$\int_I a(t)\, \mathrm{Tr}\, f(\phi(t))\, dt$$

is the image of $u_{a(t)dt \otimes f}$, not of a Hamiltonian CE coordinate.

Theorem D: Principal-part cotangent theorem

The continuous dual is

$$\mathfrak{h}^\vee_{\mathrm{cont}} \cong P = H^2_{(z_1,z_2)}(\mathbb{C}[z_1, z_2])\, dz_1 dz_2 / \mathbb{C}\rho_{0,0}.$$

The coadjoint action is

$$z_1^p z_2^q \cdot \rho_{a,b} = -(pb - qa + p - q)\rho_{a-p+1,b-q+1}.$$

The boundary realization uses independent principal-part currents

$$\Theta_\rho,$$

not ordinary polynomial Ψ-classes.

Theorem E: PBW descendant shadow

The polynomial one-ψ descendants give

$$S(\bar{A} \oplus \bar{A}_{\mathrm{desc}}[1])$$

as the PBW special-fibre stable trace sector. The identification

$$\bar{A}_{\mathrm{desc}} \cong \mathfrak{h}_{\mathrm{cont}}^\vee$$

is a graded vector-space identification only, unless a Rees/Fourier intertwiner is separately constructed.

This theorem preserves the real open descendant calculation without making it do impossible work.

Theorem F: Quantum radial-Moyal theorem

If you can fully prove the all-N radial-parts statement, make it a main theorem:

$$\hbar^{-1}[J_N(f), J_N(g)] = J_N(\{f, g\}_\hbar) \quad \text{in quantum Hamiltonian reduction, modulo } [\bar{c}].$$

This would be genuinely groundbreaking. It would say that the brane quantum Hamiltonian reduction realizes the formal Moyal algebra exactly on stable connected traces.

13. What to remove or move

The “cross-volume horizon” material should not be in the main paper. It currently dilutes the core theorem. The local paper is already ambitious. References to Vol III, Igusa, BKM, convention transfers, and downstream comparison contracts should be moved to a private convention appendix or a separate “programmatic outlook” note. The draft itself says these transfers are local-only and non-transferable without matched-convention proofs. main

A paper this mathematically delicate needs to feel surgically self-contained.

14. Replacement abstract

Here is the abstract I would use:

> We construct a formal local closed-open model for a mixed holomorphic-topological brane on
>
> $$\mathbb{R}^2 \times \mathbb{C}^2$$
>
> near a brane line $\mathbb{R} \times \{0\}$. The open ghost-zero theory is the Dirac constrained matrix system
>
> $$\int_{\mathbb{R}} \mathrm{Tr}(\phi_1 d\phi_2 + A[\phi_1, \phi_2]),$$
>
> whose BV reduction is $Q\psi = [\phi_1, \phi_2]$. The closed Hamiltonian sector is shifted cotangent BF theory for the reduced Hamiltonian Lie algebra
>
> $$\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C}.$$
>
> We prove that the closed CE algebra
>
> $$C^\bullet_{\mathrm{CE}}(\mathfrak{h} \ltimes \mathfrak{h}^\vee_{\mathrm{cont}}[1])$$
>
> is canonically isomorphic to the derived Poisson factorization center of the stable open Hamiltonian algebra $S(\mathfrak{h})$. Under this isomorphism, the CE coordinate dual to a Hamiltonian mode maps to a constant polyvector operation, while the CE coordinate dual to the shifted cotangent mode maps to the boundary trace Hamiltonian $\mathrm{Tr}\, f(\phi_1, \phi_2)$. The continuous dual $\mathfrak{h}^\vee_{\mathrm{cont}}$ is identified with the principal-part residue module supported at the brane, making the cotangent sector a distributional defect-local current. We separately compute the PBW special-fibre stable open descendant sector generated by the one-ψ Koszul classes and show why it is a graded shadow, not the deformation-level coadjoint module. Finally, we isolate the scalar $U(1)$ center anomaly and formulate the all-order Weyl/Moyal quantum Hamiltonian reduction problem.

This abstract is strong, not modest. It also avoids false equivalences.

15. Concrete errata

Here are the exact edits I would make.

Erratum 1: Fix the interval source

Replace every statement of the form

$$(a \otimes f)^\vee[-1] \mapsto B_f(a)$$

when $(a \otimes f)^\vee$ is the CE coordinate dual to a Hamiltonian current.

Correct version:

$$u_{a\otimes f} \mapsto B_f(a),$$

where $u_{a\otimes f}$ is the CE coordinate dual to the shifted cotangent generator corresponding, by reflexive pairing, to $a \otimes f \in \mathfrak{h}_I$.

Erratum 2: Replace $C_c^\infty(I)$ by $\Omega_c^\bullet(I)$

For locally constant factorization, use

$$\mathfrak{h}_I = \Omega_c^\bullet(I) \otimes \mathfrak{h}.$$

The degree-zero current formula is the top-degree component.

Erratum 3: Remove "Ψ and ρ are different bases of the same $\mathfrak{g}$-module"

Replace with:

$$\Psi_{a,b} \text{ and } \rho_{a,b} \text{ are two distinct module realizations of the same PBW label set.}$$

A genuine Rees/Fourier interpolation is a further theorem.

Erratum 4: Delete strict P_0-compatibility of $\Psi_{a,b} \mapsto \rho_{a,b}$

The map

$$\Psi_{a,b} \mapsto \rho_{a,b}$$

does not preserve the Hamiltonian action. The z_1 test already breaks it.

Erratum 5: Make principal-part currents independent boundary generators

Define

$$A_{\partial}^{pp}(I) = S(\mathfrak{h}_I) \otimes S(P_I[1])$$

with independent generators Θ_ρ, and brackets

$$\{B_f, \Theta_\rho\} = \Theta_{f\cdot\rho}, \qquad \{\Theta_\rho, \Theta_\sigma\} = 0.$$

Do not claim this is obtained from ordinary polynomial open observables without an added distributional boundary sector.

Erratum 6: Rename the primitive projection

Call it the **primitive indecomposable P_0-shadow**, not a factorization algebra equivalence.

Erratum 7: Strengthen or condition the radial-parts theorem

Either prove the all-N radial-Moyal theorem as a standalone appendix, or phrase it as conditional on the exact radial-parts identity for mixed Weyl traces.

Erratum 8: Treat weighted Tate as a regulator

Say:

$$(\mathfrak{h}_w, \mathfrak{h}_{\mathrm{cont}}^{\vee,w})$$

is a weighted analytic coefficient replacement, not the strict original cotangent pair. Add a regulator-independence problem or theorem.

Erratum 9: Replace “Schur rigidity”

Use a residue/D-module uniqueness theorem.

Erratum 10: Move cross-volume material out of the main body

Keep the local paper self-contained.

Bottom line

The paper is now aiming at the right groundbreaking synthesis:

$$\text{Dirac constrained brane mechanics} + \text{Witten closed-open centrality} + \text{Grothendiec}$$

That is a serious idea.

The one thing that must be fixed is the temptation to identify all cotangent-looking things:

$$\Psi_{a,b} \neq \rho_{a,b} \quad \text{as } \mathfrak{h}\text{-modules.}$$

And

$$B_f \neq \Phi(c_f).$$

The corrected identifications are:

$$B_f = \Phi(u_f),$$

and

$$\rho_{a,b} \text{ is a distributional principal-part boundary mode, not a polynomial } \Psi_{a,b} \text{ class.}$$

Once those two corrections propagate through the factorization and primitive-projection sections, the paper becomes much harder to attack and much more original. It will no longer read as a paper trying to force a string slogan onto matrix traces. It will read as a new local theorem about **how closed Hamiltonian BF theory emerges as the derived Poisson center of a Dirac-reduced brane probe**, with the cotangent sector realized by holomorphic residue currents at the defect.

Sources

Sources

generate an EXHAUSTIVE set of recommendations for technical fixes, modifications, new insertions, rewrites etc.

I'll treat this as a referee-plus-coauthor overhaul: isolate technical failure points and structural contradictions, then turn them into concrete theorem replacements, rewrites, appendices, and proof obligations.

The key pattern is layer separation: CE observables, PBW shadows, polynomial descendants, principal parts, interval factorization, and quantum analytic claims must

each get distinct theorem status.

Thought for 12m 18s ›

According to a document from 2026-04-28, the current WIP has crossed an important threshold: it now contains the right central thesis—closed Hamiltonian BF observables as the derived Poisson center of the Dirac-reduced brane probe algebra—but it still needs a hard technical cleanup so that the theorem statements, source conventions, factorization language, cotangent module, quantum claims, and analytic claims all pull in the same direction. The draft's own abstract already contains the key ingredients: the CE/PV isomorphism, principal-part residue module, factorization upgrade, Moyal/Capelli sector, U(1) anomaly, weighted Tate regulator, nilpotent truncation, and QME caveats. The issue is not ambition; the issue is that several layers are currently fused together when they should be separated and then recomposed with explicit maps. main main

Below is an exhaustive technical punch list.

0. Executive diagnosis

The strongest version of the paper should be organized around this spine:

$$\text{Dirac brane probe} \quad \Longrightarrow \quad \text{stable open Hamiltonian trace algebra} \quad \Longrightarrow \quad \text{closed}$$

with the theorem

$$C^{\bullet}_{\mathrm{CE,cont}}\big(\mathfrak{h} \ltimes \mathfrak{h}^{\vee}_{\mathrm{cont}}[1]\big) \cong \mathrm{PV}(S(\mathfrak{h})) \simeq Z^{P_0}_{\mathrm{fact}}(S(\mathfrak{h})).$$

The draft correctly says that the CE coordinate c^I dual to a Hamiltonian generator maps to the constant polyvector $\theta^I = \partial/\partial O_I$, while the CE coordinate u_I dual to the shifted cotangent generator maps to the boundary Hamiltonian function $O_I = \mathrm{Tr}\, H_I(\phi_1, \phi_2)$. This is the central mathematical insight. main

But three old confusions still leak into the current draft:

$$B_f \neq \Phi(c_f), \qquad B_f = \Phi(u_f).$$

$$\Psi_{a,b} \not\simeq \rho_{a,b} \quad \text{as } \mathfrak{h}\text{-modules.}$$

$$C_c^\infty(I) \text{ is not enough for locally constant factorization; use } \Omega_c^\bullet(I).$$

Fixing those three points will remove the main mathematical vulnerabilities.

1. Reorganize the paper around six clean theorem layers

The current "umbrella theorem" tries to state too many things at once. It is good as a programmatic summary, but bad as the first serious theorem because it mixes proved algebra, reduced models, conditional analytic completions, quantum statements, and future graph problems. The current local theorem has six interlocking statements: CE/PV isomorphism, principal-part coadjoint identification, factorization upgrade, cotangent factor, all-order quantum deformation, and U(1) anomaly. That is the right content, but it should be split into independent theorem blocks. main

Use this architecture:

Theorem A: Dirac brane probe theorem

State and prove:

$$S_\partial = \int_{\mathbb{R}} \mathrm{Tr}\big(\phi_1 d\phi_2 + A[\phi_1, \phi_2]\big),$$

$$Q\psi = [\phi_1, \phi_2], \qquad Q\phi_1 = Q\phi_2 = 0.$$

This theorem should establish the open model as a first-class Dirac constrained system, not as a stringy ansatz.

Theorem B: Stable Hamiltonian trace theorem

State:

$$\mathrm{Obs}^{cl}_{\partial,H} \simeq S(\bar{A} \oplus \bar{A}_{\mathrm{desc}}[1])$$

as the **stable connected PBW-special-fibre Hamiltonian sector**, with $\Psi_{a,b}$ included only as the primitive one-ψ stable Koszul classes.

Theorem C: CE/PV derived center theorem

This is the central theorem:

$$C^\bullet_{\mathrm{CE,cont}}(\mathfrak{h} \ltimes \mathfrak{h}^\vee_{\mathrm{cont}}[1]) \cong \mathrm{PV}(S(\mathfrak{h})).$$

On generators:

$$c^I \mapsto \theta^I, \qquad u_I \mapsto O_I.$$

This theorem must be given before any PBW/Koszul-dual comparison.

Theorem D: Principal-part cotangent theorem

State:

$$\mathfrak{h}^\vee_{\mathrm{cont}} \cong P = \bigoplus_{a+b>0} \mathbb{C}\rho_{a,b}, \qquad \rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2,$$

with residue pairing

$$\langle \rho_{a,b}, z_1^m z_2^n \rangle = \delta_{a,m}\delta_{b,n}.$$

Then prove the coadjoint action:

$$z_1^p z_2^q \cdot \rho_{a,b} = -(pb - qa + p - q)\rho_{a-p+1,b-q+1}.$$

Do not rely on “Schur rigidity” unless you actually prove it.

Theorem E: Factorization-current theorem

Use compactly supported de Rham forms:

$$\mathfrak{h}_I = \Omega^\bullet_c(I) \otimes \mathfrak{h},$$

not just $C^\infty_c(I) \otimes \mathfrak{h}$. Then prove the derived-center statement intervalwise.

Theorem F: Quantum Hamiltonian/Moyal theorem

Separate the unconditional algebraic Moyal/Weyl target from the conditional Costello graph realization. The draft currently says the degree-zero quantum sector is proved through Capelli renormalization and radial-parts identities, while the full descendant/cotangent quantum lift remains conditional on Tate/QME problems. Keep that separation explicit. main

2. Fix the central source-convention error in the factorization theorem

The draft correctly states in the abstract and cochain theorem that the boundary trace observable is the image of the shifted cotangent CE coordinate, not the unclosed

Hamiltonian CE coordinate. main

But Theorem 0.1.20(iii) and Theorem 0.2.21 still say, in effect,

$$(a \otimes f)^\vee[-1] \longmapsto B_f(a) = \int_I a(t) \operatorname{Tr} f(\phi(t))\, dt.$$

That is the old source mistake in interval language. The source of $B_f(a)$ must be the CE coordinate dual to the **shifted cotangent generator** paired with $a \otimes f$, not the CE coordinate dual to the Hamiltonian current itself. The retrieved theorem statement still displays precisely the problematic assignment. main

Replace every such line by:

$$u_{a\otimes f} \longmapsto B_f(a) = \int_I a(t) \operatorname{Tr} f(\phi_1(t), \phi_2(t))\, dt.$$

Then define:

$$c_\lambda \longmapsto \theta_\lambda, \qquad \lambda \in \mathfrak{h}_I^\vee.$$

The corrected interval theorem should read:

$$\mathfrak{g}_I = \mathfrak{h}_I \ltimes \mathfrak{h}_I^\vee[1],$$

$$C^\bullet_{\mathrm{CE}}(\mathfrak{g}_I) \cong \mathrm{PV}(S(\mathfrak{h}_I)),$$

$$c_\lambda \mapsto \theta_\lambda, \qquad u_x \mapsto O_x, \qquad x \in \mathfrak{h}_I.$$

For the top-degree current $x = a(t)dt \otimes f$,

$$u_x \mapsto B_f(a).$$

This one correction should propagate through:

$$0.1.14(iii), \quad 0.2.21, \quad 0.2.23, \quad \text{the abstract}, \quad \text{the proof of factorization compatibi}$$

3. Replace $C_c^\infty(I)$ by $\Omega_c^\bullet(I)$ in all locally constant factorization statements

The current draft often defines

$$\mathfrak{h}_I = C_c^\infty(I) \otimes \mathfrak{h}.$$

That is fine for the degree-zero current bracket, but it is not the right object for locally constant factorization. For local constancy on intervals, use the compactly supported de Rham cosheaf:

$$\mathfrak{h}_I = \Omega_c^\bullet(I) \otimes \mathfrak{h}.$$

The bracket is

$$[\alpha \otimes f, \beta \otimes g] = (\alpha \wedge \beta) \otimes \{f, g\}.$$

The degree-zero current formula is recovered by taking top-degree compactly supported forms:

$$\alpha = a(t)dt.$$

Then

$$B_f(a) = \int_I a(t) \operatorname{Tr} f(\phi_1(t), \phi_2(t))\, dt.$$

This change matters because $C_c^\infty(I)$ alone has no de Rham differential and does not make interval inclusions quasi-isomorphisms. The draft already knows the de Rham contraction is the reason positive jets vanish and factorization becomes locally constant; the theorem definitions need to reflect that. The current text uses $C_c^\infty(I)$ in the support-locality section and in the factorization theorem. main

Recommended replacement:

> For an interval I, define the compactly supported Hamiltonian current dg Lie algebra
>
> $$\mathfrak{h}_I = \Omega_c^\bullet(I) \otimes \mathfrak{h}.$$
>
> The locally constant factorization algebra is obtained by taking Chevalley-Eilenberg cochains on $\mathfrak{h}_I$, with extension by zero. The displayed current formula $B_f(a)$ is the top-degree component $a(t)dt \otimes f$.

4. Remove the claim that $\Psi_{a,b}$ and $\rho_{a,b}$ are “different bases of the same $\mathfrak{g}$-module”

This is the most dangerous mathematical sentence in the draft.

The current theorem says that the polynomial descendants $\Psi_{a,b}$ act by symmetric raising, the principal parts $\rho_{a,b}$ act by coadjoint lowering, and that "the two are not in conflict" because they are "different bases of the same $\mathfrak{g}$-module" interpolated by $\hbar$.

main

That cannot be right as written.

The draft later contains the correct no-go theorem: ordinary polynomial boundary observables cannot realize the principal-part coadjoint module, because linear Hamiltonians act locally nilpotently on polynomials but not on principal parts. main

So the correct statement is:

$$\Psi_{a,b} \text{ and } \rho_{a,b} \text{ are not two bases of the same } \mathfrak{h}\text{-module.}$$

They are two realizations of the same **label set** or **PBW cotangent symbol set**:

$$\bar{A}_{\mathrm{desc}} \cong \mathfrak{h}^{\vee}_{\mathrm{cont}}$$

as graded vector spaces, not as $\mathfrak{h}$-modules.

Replace:

> The two are not in conflict: they are different bases of the same $\mathfrak{g}$-module, interpolated by the Rees parameter $\hbar$.

with:

> The two are not the same $\mathfrak{h}$-module. The $\Psi_{a,b}$ form the PBW-special-fibre polynomial descendant realization; the $\rho_{a,b}$ form the deformation-correct coadjoint realization. They carry the same bidegree labels, but the actions are not conjugate. A genuine Rees/Fourier interpolation between them would require an explicit $\hbar$-dependent $D_\hbar$-module or local-cohomology module and is not implied by PBW alone.

Then add a named problem:

Problem: Rees/Fourier interpolation of the descendant and principal-part modules. Construct an $\hbar$-flat module $\mathcal{M}_\hbar$ over the Rees Weyl algebra

$$D_\hbar = \mathbb{C}\langle z_1, z_2, \hbar\partial_{z_1}, \hbar\partial_{z_2}\rangle / ([\hbar\partial_{z_i}, z_j] = \hbar\delta_{ij})$$

such that

$$\mathcal{M}_\hbar/(\hbar) \cong \bar{A}_{\mathrm{desc}}$$

as PBW symbol module, while

$$\mathcal{M}_\hbar[\hbar^{-1}]$$

identifies with the principal-part/local-cohomology coadjoint module.

5. Correct the word “raising/lowering” for principal parts

The current text sometimes says principal parts carry a “lowering” action. The formula is:

$$z_1^p z_2^q \cdot \rho_{a,b} = -(pb - qa + p - q)\rho_{a-p+1,b-q+1}.$$

Depending on p, q, this may raise one index, lower another, or kill the term. For example:

$$z_1 \cdot \rho_{a,b} = (a-1)\rho_{a,b+1}$$

using the displayed formula with $p = 1, q = 0$, if the indexing convention is exactly as written. But in earlier discussions and principal-part notes, the expected coadjoint action under z_1 was also described as increasing the b-pole order. The important fact is not universally “lowering”; the important fact is that the action is the negative transpose/coadjoint action and is not locally nilpotent in the same way as the polynomial action.

Recommendation: replace “lowering action” with:

$$\text{coadjoint principal-part action}$$

or:

$$\text{negative-transpose residue action}$$

Then add a small table:

f	$f \cdot \rho_{a,b}$	effect
z_1	compute explicitly from convention	raises/lowers pole index depending on
z_2	compute explicitly from convention	raises/lowers pole index depending on
$z_1 z_2$	diagonal weight action	Euler-type action

This will prevent a referee from attacking a misleading “raising/lowering” slogan.

6. Re-prove principal-part uniqueness using local cohomology, not Schur rigidity

The draft says the residue pairing is unique up to scalar "by Schur rigidity." main

That is risky. Infinite-dimensional $\mathfrak{h}$-modules are subtle, and an unproved Schur lemma in this context will attract criticism.

Replace the uniqueness statement with a concrete local-cohomology theorem:

$$P \cong H^2_{(z_1,z_2)}(\mathbb{C}[z_1,z_2])\, dz_1 dz_2 / \mathbb{C}\rho_{0,0}.$$

Then prove the residue pairing directly:

$$\langle \rho_{a,b}, z_1^m z_2^n \rangle = \mathrm{Res}_0\, z_1^{m-a-1} z_2^{n-b-1} dz_1 dz_2.$$

Prove equivariance by integration by parts:

$$\langle f \cdot \rho, g \rangle = -\langle \rho, \{f, g\} \rangle.$$

Then state the uniqueness you actually need:

> The residue pairing is the unique continuous degreewise pairing satisfying coefficient extraction and Hamiltonian integration by parts, after normalizing
>
> $$\langle \rho_{a,b}, z_1^a z_2^b \rangle = 1.$$

This is stronger than a vague Schur claim because it is checkable.

7. Separate the three cotangent objects

Right now the draft has three different "cotangent" objects that sometimes blur:

1. the CE cotangent coordinate u_I;
2. the PBW-special-fibre open descendant $\Psi_{a,b}$;
3. the principal-part boundary mode $\Theta_{\rho_{a,b}}$.

These must never be identified without specifying the level.

Add a "Cotangent dictionary" table:

Object	Lives where	Action	
u_I	$C^\bullet_{\mathrm{CE}}(\mathfrak{h} \ltimes \mathfrak{h}^\vee[1])$	$d_{\mathrm{CE}} u_K = C^L_{KJ} u_L c^J$	closed CE c
O_I	$\mathrm{PV}(S(\mathfrak{h}))$	$d_\pi O_K = C^L_{KJ} O_L \theta^J$	boundary H
$\Psi_{a,b}$	stable open Koszul homology	symmetric PBW action	PBW speci
$\rho_{a,b}$	$\mathfrak{h}^\vee_{\mathrm{cont}}$	$\mathrm{ad}^\vee$	closed
Θ_ρ	A^{pp}_∂	$\{B_f, \Theta_\rho\} = \Theta_{f\cdot\rho}$	reduced distrib

Then add this warning:

$$\Psi_{a,b} \mapsto \rho_{a,b} \text{ is not a } P_0\text{-morphism.}$$

8. Rename the “primitive projection” theorem

The current abstract says the chain-level primitive projection

$$\pi_{\mathrm{prim}} : \mathrm{Obs}^{cl}_{open,N}(I) \to \Omega^\bullet_c(I) \otimes \bar{A}$$

is constructed from Procesi-Razmyslov cycle merging and de Rham contraction, with P_0-bracket homotopy identically zero. main

If this projection kills products of positive primitive generators, it should not be sold as a factorization algebra equivalence. It is an indecomposable or primitive Lie shadow.

Rename it:

$$\text{primitive indecomposable } P_0\text{-shadow}$$

not:

$$\text{factorization algebra lift.}$$

A factorization algebra remembers products. A primitive quotient that kills products is a useful Lie-level shadow, but it is not the same object.

Recommended theorem title:

> **Theorem. Primitive indecomposable P_0-projection of the open trace factorization algebra.**

Then state explicitly:

$$\pi_{\mathrm{prim}}(XY) = 0 \quad \text{for positive primitive } X, Y.$$

This will save you from the objection that the map cannot preserve the full factorization product.

9. Recast the principal-part boundary sector as an added reduced sector, not as derived from polynomial observables

The draft correctly says the principal-part current algebra

$$A_{\partial}^{pp}(I) = S(\mathfrak{h}_I) \otimes S(P_I[1])$$

is a reduced post-contraction P_0-factorization algebra, and that $Q\Theta_\rho = 0$ is part of the added local-dual factor rather than derived from the unreduced BV model. main

Make this even more explicit.

Use the language:

$$A_{\partial}^{pp} \text{ is an enlarged reduced defect-current model.}$$

Do not call it “the open observable algebra” without qualifiers. It is the minimal reduced boundary object that realizes the coadjoint sector. The actual unreduced BV origin remains a separate problem.

Add this paragraph:

> The principal-part sector is not obtained from ordinary polynomial matrix observables. It is adjoined as the reduced local-dual defect-current module forced by the coadjoint action. Its unreduced BV realization would require a distributional current complex on the brane and homotopies expressing centrality against arbitrary open observables.

That matches the draft’s own status remark. main

10. Strengthen the derived-center proof by adding a coordinate-free version

The current proof is coordinate-heavy:

$$[e_I, e_J] = C_{IJ}^K e_K,$$

$$d_{\mathrm{CE}} c^K = -\frac{1}{2} C_{IJ}^K c^I c^J, \qquad d_{\mathrm{CE}} u_K = C_{KJ}^L u_L c^J,$$

$$d_\pi \theta^K = -\frac{1}{2} C_{IJ}^K \theta^I \theta^J, \qquad d_\pi O_K = C_{KJ}^L O_L \theta^J.$$

That is good, but add a coordinate-free theorem immediately before or after it:

Let $A = S(\mathfrak{h})$ be the coordinate algebra of $\mathfrak{h}^\vee$. The linear Poisson structure on $\mathfrak{h}^\vee$ gives

$$\mathrm{PV}(A) = S(\mathfrak{h}) \otimes \Lambda(\mathfrak{h}^\vee)$$

with Lichnerowicz differential. Then

$$\mathrm{PV}(S(\mathfrak{h}), \pi_{\mathfrak{h}}) \cong C^\bullet_{\mathrm{CE}}(\mathfrak{h}; S(\mathfrak{h}))$$

where $S(\mathfrak{h})$ is the coadjoint representation. But

$$C^\bullet_{\mathrm{CE}}(\mathfrak{h}; S(\mathfrak{h})) \cong C^\bullet_{\mathrm{CE}}(\mathfrak{h} \ltimes \mathfrak{h}^\vee[1]).$$

This gives the conceptual proof:

$$\text{Poisson cochains on } \mathfrak{h}^\vee = \text{CE cochains of the shifted cotangent Lie algebra.}$$

Then the coordinate proof becomes a verification of signs.

11. Add a sign-convention appendix

The paper has too many shifts and signs for the main body to carry all convention risk. Add an appendix:

Appendix A: Shift, CE, Schouten, and BV sign conventions

Include:

$$V[1]^n = V^{n+1} \quad \text{or} \quad V[1]^n = V^{n-1}$$

depending on your convention. The draft says an element of $V[1]$ has parity reversed and cohomological degree one lower; make that globally consistent.

Define:

$$d_{\mathrm{CE}} c^K = -\frac{1}{2} C^K_{IJ} c^I c^J .$$

Define the semidirect differential:

$$d_{\mathrm{CE}} u_K = C^L_{KJ} u_L c^J$$

and prove $d^2_{\mathrm{CE}} u_K = 0$.

Define the Schouten bracket convention:

$$[\theta^I, O_J] = \delta^I_J$$

not the negative.

Define:

$$d_\pi = [\pi, -]$$

and compute:

$$d_\pi O_K, \qquad d_\pi \theta^K .$$

Then include a sign checklist:

$$\Phi(d_{\mathrm{CE}} c^K) = d_\pi \Phi(c^K),$$

$$\Phi(d_{\mathrm{CE}} u_K) = d_\pi \Phi(u_K).$$

This appendix will make the central theorem reproducible.

12. Add a “claim-strength ledger” table near the abstract

The current abstract is too dense and reads like a dozen theorems at once. It includes unconditional statements, conditional statements, reduced statements, post-contraction statements, weighted regulator statements, graphwise statements, and open problems in one paragraph. main

Add a table:

Statement	Status
Dirac open BV model	proved
Stable trace sector	proved
CE/PV derived center	proved
Principal-part coadjoint module	proved
Interval factorization	proved after de Rham current corr
Reduced principal-part boundary sector	proved as enlarged reduced mo
Unreduced BV cotangent lift	open/conditional
Quantum degree-zero Moyal	proved/conditional depending on radial
Quantum descendant lift	conditional
Costello graph realization	open

This will make the paper look stronger, not weaker, because it shows exactly which mathematical claims are proved.

13. Rewrite the abstract

The abstract currently contains too many technical claims before the reader has the dictionary. Use two paragraphs: one physics paragraph, one theorem paragraph.

Recommended replacement:

> This paper studies the formal local closed-open Hamiltonian sector of a mixed holomorphic-topological BV model near a brane line
>
> $$\mathbb{R} \times \{0\} \subset \mathbb{R}^2 \times \mathbb{C}^2.$$
>
> The holomorphic normal disk has coordinates z_1, z_2 and symplectic form $dz_1 \wedge dz_2$. A stack of N probes replaces z_i by matrices ϕ_i, and Dirac reduction of the first-order action
>
> $$\int_{\mathbb{R}} \mathrm{Tr}(\phi_1 d\phi_2 + A[\phi_1, \phi_2])$$
>
> imposes the commuting-pair constraint $Q\psi = [\phi_1, \phi_2]$.
>
> The closed Hamiltonian sector is shifted cotangent BF theory for
>
> $$\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C}, \qquad \mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}^\vee_{\mathrm{cont}}[1].$$
>
> The main theorem identifies closed CE observables with the derived Poisson factorization center of the stable open Hamiltonian algebra:

$$C^{\bullet}_{\mathrm{CE,cont}}(\mathfrak{g}) \cong \mathrm{PV}(S(\mathfrak{h})) \simeq Z^{P_0}_{\mathrm{fact}}(S(\mathfrak{h})).$$

Under this isomorphism, the CE coordinate dual to a Hamiltonian generator maps to a constant polyvector operation, while the CE coordinate dual to the shifted cotangent generator maps to the boundary trace Hamiltonian $\mathrm{Tr}\, f(\phi_1, \phi_2)$. The continuous dual $\mathfrak{h}^{\vee}_{\mathrm{cont}}$ is realized by principal-part residue currents at the brane. Polynomial one-ψ descendants form the PBW-special-fibre open shadow; they are not the deformation-level coadjoint module. The scalar $U(1)$ center anomaly and the quantum Weyl/Moyal deformation are treated separately, with the full brane-defect Costello graph realization isolated as an analytic problem.

This abstract says exactly what is new.

14. Rewrite the “thesis” section as a theorem-corollary

The current thesis sentence is excellent:

A stack of branes is a system of probes for the formal symplectic normal disk. Dirac reduction of the probe phase space gives the open trace algebra. Hamiltonian BF theory is the shifted cotangent theory of canonical transformations. The closed observables are precisely the derived Poisson central operations on the stable open Hamiltonian brane algebra.

Keep it, but make it a corollary after the theorem, not just a slogan. The draft already attempts this in Remark 0.1.22. main

Recommended formulation:

Corollary. Dirac-Witten closed-open principle.
For the formal symplectic disk and the point brane stack, the Witten closed-open map is the derived center map

$$\mathrm{Obs}^{cl}_{closed,H} \to Z^{P_0}_{fact}(\mathrm{Obs}^{cl}_{open,H}),$$

and under the CE/PV identification it is the identity map on polyvector operations.

This will make the physics synthesis feel inevitable.

15. Add a “Dirac first-principles” derivation box

Insert near the beginning:

$$\Omega_N = \mathrm{Tr}(d\phi_1 \wedge d\phi_2),$$

$$\mu_\epsilon = -\mathrm{Tr}(\epsilon[\phi_1, \phi_2]),$$

$$\iota_{X_{\mu_\epsilon}} \Omega_N = -d\mu_\epsilon,$$

$$X_{\mu_\epsilon}(\phi_i) = [\epsilon, \phi_i].$$

Then:

$$S_\partial = \int \mathrm{Tr}(\phi_1 d\phi_2) - \mu_A = \int \mathrm{Tr}(\phi_1 d\phi_2 + A[\phi_1, \phi_2]).$$

This should appear before any string terminology. It grounds the model in canonical mechanics.

16. Add a “Witten centrality” derivation box

After the Dirac box, add:

A closed insertion approaching a brane defines not merely an open element but a central operation:

$$\mathrm{Obs}_{closed} \to Z_{fact}(\mathrm{Obs}_{open}).$$

Classically, because the open algebra has a Poisson bracket, the target is

$$Z^{P_0}_{fact}(\mathrm{Obs}_{open}).$$

For a linear Poisson algebra $S(\mathfrak{h})$,

$$Z^{P_0}_{fact}(S(\mathfrak{h})) \simeq \mathrm{PV}(S(\mathfrak{h})).$$

Then the closed BF sector appears as this center.

This is the physical reason for factorization. It should be front-loaded.

17. Fix the “closed Hamiltonian factor before adjoining cotangent generators” theorem

Theorem 0.2.21 currently seems to define the closed Hamiltonian-mode subfactorization algebra before adjoining cotangent generators and maps its CE cochains into central operations. But the full derived-center theorem requires the shifted cotangent extension. A CE algebra of $\mathfrak{h}_I$ alone gives only the exterior/vector-field part, not the functions O_f.

Correct statement:

$$C^\bullet_{\mathrm{CE}}(\mathfrak{h}_I) \cong \Lambda(\mathfrak{h}_I^\vee)$$

maps to constant polyvectors only:

$$c_\lambda \mapsto \theta_\lambda.$$

It does **not** contain $B_f(a)$ as a function.

The boundary Hamiltonian current appears only after extending to

$$\mathfrak{g}_I = \mathfrak{h}_I \ltimes \mathfrak{h}_I^\vee[1].$$

Then

$$u_{a\otimes f} \mapsto B_f(a).$$

So rewrite Theorem 0.2.21 into two parts:

Part 1: Hamiltonian ghosts only

$$C^\bullet_{\mathrm{CE}}(\mathfrak{h}_I) \to \mathrm{PV}^{\mathrm{const}}(S(\mathfrak{h}_I)), \qquad c_\lambda \mapsto \theta_\lambda.$$

Part 2: shifted cotangent extension

$$C^\bullet_{\mathrm{CE}}(\mathfrak{h}_I \ltimes \mathfrak{h}_I^\vee[1]) \cong \mathrm{PV}(S(\mathfrak{h}_I)), \qquad u_x \mapsto O_x.$$

That removes the source mismatch entirely.

18. Recast the PBW/Koszul-dual theorem as a separate “shadow” theorem

The draft’s older theorem said the stable open Hamiltonian sector matches the PBW special fibre of the Koszul dual closed algebra. That remains useful, but it should not compete with the CE/PV theorem.

State:

$$(C^\bullet_{\mathrm{CE}}(\mathfrak{g}))^! \simeq U(\mathfrak{g}).$$

PBW Rees:

$$U_\hbar(\mathfrak{g})/(\hbar) \simeq S(\mathfrak{h} \oplus \mathfrak{h}^\vee[1]).$$

Then:

$$S(\bar{A} \oplus \bar{A}_{\mathrm{desc}}[1])$$

is a degreewise polynomial PBW-special-fibre shadow of this associated graded.

Add a warning:

$$\text{PBW special fibre does not encode the coadjoint action.}$$

This lets you keep the beautiful one-ψ computation while preventing it from corrupting the deformation-level story.

19. Add a proof that $\check{\mathfrak{h}}$ is perfect in the formal completed setting

You already prove perfectness for polynomials:

$$z_1^a z_2^b = \left\{ z_1, \frac{1}{b+1} z_1^a z_2^{b+1} \right\}.$$

For formal power series, include the continuity argument:

Let

$$f = \sum_{a+b>0} f_{a,b} z_1^a z_2^b.$$

Then

$$f = \left\{ z_1, \sum_{a,b\geq 0} \frac{f_{a,b}}{b+1} z_1^a z_2^{b+1} \right\}$$

in the completed topology, modulo constants.

This proves:

$$[\mathfrak{h}, \mathfrak{h}] = \mathfrak{h}.$$

Then the CE-source obstruction follows in the exact topological Lie algebra used by the main theorem.

20. Add a local-cohomology explanation of the $a + b > 0$ cutoff

The principal-parts note explains that $\rho_{0,0}$ pairs with the constant 1, which is removed from $\mathfrak{h}$. Pasted text

Insert this in the paper:

$$\rho_{0,0} = z_1^{-1} z_2^{-1} dz_1 dz_2, \qquad \langle \rho_{0,0}, 1 \rangle = 1.$$

Since

$$\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C},$$

the dual is

$$P = H^2_{(z_1,z_2)}(\mathbb{C}[z_1, z_2]) dz_1 dz_2 / \mathbb{C}\rho_{0,0}.$$

Then tie it to the scalar anomaly:

$$\rho_{0,0} \leftrightarrow 1 \leftrightarrow \mathrm{Tr}(1) = N.$$

This makes the residue cutoff physically meaningful.

21. Tighten the U(1) anomaly statement

The scalar anomaly section is strong. The draft identifies the classical cocycle

$$\omega(z_1^k z_2^l, z_1^m z_2^n) = (kn - lm) \mathbf{1}_{(k+m, l+n) = (1,1)}$$

with the brane bracket

$$\{\mathrm{Tr}\, \phi_1, \mathrm{Tr}\, \phi_2\} = N$$

and with the quantum Capelli shift

$$YX - XY + \hbar NI \equiv 0.$$

This is a major asset. Keep it central. main

But avoid claiming that $[\bar{c}]$ spans all of $H^2_{\mathrm{Lie}}(\bar{A}, \mathbb{C})$ unless you compute the whole cohomology.

Use:

$$[\bar{c}] \text{ is the distinguished scalar-extension line in } H^2_{\mathrm{Lie}}(\bar{A}, \mathbb{C}).$$

not:

$$H^2_{\mathrm{Lie}}(\bar{A}, \mathbb{C}) \cong \mathbb{C}[\bar{c}].$$

unless you prove the latter.

22. Clarify "scalar reduction" versus "removing $U(1)$ center-of-mass"

Do not say the whole $U(1)$ center-of-mass sector is removed. The linear traces

$$\mathrm{Tr}\,\phi_1, \qquad \mathrm{Tr}\,\phi_2$$

remain as observables, but their bracket produces the scalar

$$N.$$

The precise operation is:

$$\text{remove the scalar empty-trace primitive } \mathrm{Tr}(1) = N.$$

That kills the central value of the bracket, not necessarily the linear trace variables.

Add this explicit warning after every formula where

$$\{z_1, z_2\} = 0$$

in $\bar{A}$.

23. Upgrade the finite-N versus stable-N discussion

The stable large-N sector currently uses degreewise stabilization under block restriction. Keep that, but add a separate subsection:

Three large-N operations

1. **Degreewise invariant stabilization**:
   finite word degree d, take $N \geq d$.
2. **Primitive connected trace extraction**:
   pass from the symmetric multi-trace algebra to single-trace primitives.
3. **Scalar Hamiltonian reduction**:
   omit the empty trace as primitive and quotient scalar central brackets.

These are not the same operation.

Add:

$$\mathrm{Prim}_{st} \neq \mathrm{Conn} \neq (-)/\mathbb{C}\,\mathrm{Tr}(1).$$

Then define each.

This will prevent a reader from thinking the large-N limit is hiding scalar anomalies or trace identities.

24. Strengthen the one-ψ homology proof

The one-ψ primitive homology theorem is one of the paper's most original algebraic calculations. It currently relies on the chip-moving state complex and Abel-map argument. It is plausible but still vulnerable.

Add:

1. A fully formal definition of $C_2(k,l)$, $C_1(k,l)$, $C_0(k,l)$.
2. A sign lemma proving the two-ψ boundary formula.
3. A cellular-chain lemma identifying that complex with the two-skeleton of $Y_{k,l}$.
4. A proof that attaching cells of dimension ≥ 3 does not alter H_1.
5. A proof that the Abel map is well-defined under cyclic relabeling.
6. A lemma on stabilizers: quotient of a convex fiber by a finite affine stabilizer is contractible.
7. A final lemma identifying $\Psi_{k,l}$ with the fundamental S^1-cycle.

Then move the script verification to an appendix. The script is useful, but the theorem must stand independently.

25. Put all computer checks into a reproducibility appendix

The draft mentions scripts like `scripts/check_one_psi_homology.py` and Moyal coefficient scripts. main

Add an appendix:

Appendix B: Finite checks and reproducibility

Include:

Script	Checks	Range
`check_one_psi_homology.py`	$\dim H_1 = 1,$ descendant action	$0 \le k, l \le 5$
`check_moyal_coefficients.py`	P^r coefficients	exponents $\le$

Then say clearly:

> These checks verify finite instances and guard against sign errors. They are not used as proof except where explicitly stated as computational evidence.

26. Rework the weighted Tate section as a regulator theorem

The weighted Tate completion currently says the weighted pair makes the formal Casimir converge, extends the finite-scale heat-kernel propagator, and gives graphwise RG control. It also says the second factor is a product coefficient module, not the strict continuous dual of the first product Tate space. main

That means it is not simply "the original cotangent theory completed correctly." It is a **regulated coefficient replacement**.

Rename:

$$(\mathfrak{h}_w, \mathfrak{h}_{cont}^{\vee,w})$$

as:

$$\text{weighted coefficient regulator}$$

not:

$$\text{the continuous dual of } \mathfrak{h}_w.$$

Add:

> The weighted theory is a regulator category in which the BV coevaluation kernel exists. It contains the polynomial/direct-sum principal-part sector densely, but it is not identical to the strict Tate cotangent theory. Regulator independence is a separate question.

Then add a problem:

Problem: Regulator independence.
Show that the derived-center map and all finite-degree observables stabilize as $q \to 1^+$, or prove that any two exponential weights $q, q' > 1$ give canonically equivalent finite-degree local observables.

27. Remove overstrong claims in weighted quantum extension

The current weighted theorem states that the all-order quantum derived-center isomorphism extends to the weighted cotangent direction provided the QME compatibility identity holds. That is fine, but phrase it as conditional, not as part of the same theorem statement as the proved Casimir/propagator claims. main

Split:

Theorem: Weighted coefficient kernel

Unconditional:

$$C_{\mathfrak{h},w} \in \mathfrak{h}_w \otimes \mathfrak{h}_{cont}^{\vee,w}.$$

Theorem: Graphwise RG boundedness

Unconditional for each fixed graph.

Conditional theorem: Weighted quantum cotangent lift

Conditional on QME/locality obstruction vanishing.

This separation will make the analytic claims credible.

28. Add a “Hadamard/Mittag-Leffler does not imply QME” warning

The draft correctly says the Hadamard/Mittag-Leffler reduction controls wavefront and finite-window locality but does not compute the obstruction class. main

Make this a boxed warning:

$$\text{Hadamard control is not anomaly cancellation.}$$

Then:

> The Hadamard theorem controls distributional wavefront sets and passage to the Mittag-Leffler coefficient limit. It does not prove that the brane-defect counterterm system solves the quantum master equation.

This protects the paper from analytic overclaiming.

29. Reassess the all-N radial-parts theorem

The latest draft seems to claim the degree-zero quantum sector is now fully proved using Capelli renormalization, finite-N radial-parts identity, and stable connected limit, while still separating the cotangent and Costello graph obligations. main

This is potentially groundbreaking but must be armored.

Add a standalone appendix:

Appendix C: Radial-parts and Capelli-renormalized Weyl traces

It must answer:

1. What algebra is the quantum Hamiltonian reduction?

$$\mathcal{R}_N = N_{W_N}(W_N\mu(\mathfrak{gl}_N))/W_N\mu(\mathfrak{gl}_N).$$

2. What is the exact definition of

$$J_N(f) = \operatorname{Tr} \operatorname{Op}_W(f)(X, Y)?$$

3. What is the Capelli-renormalized generator?

$$J_N(f) = J_N(f) + \sum_{r \geq 1} \hbar^r C_r(f, N).$$

4. What radial map is used?

$$\operatorname{Rad}_0 : \mathcal{R}_N \to \text{Dunkl/symmetric Weyl algebra.}$$

5. Is the radial map injective on the relevant subspace?
6. Does the image of $J_N(f)$ become a one-particle Weyl operator sum?
7. Do Dunkl/Calogero correction terms vanish, cancel, or become scalar?
8. Does connected single-trace extraction commute with the bracket?

Without these details, the radial-parts theorem will look like a magic black box.

30. Clarify the relation between quantum Moyal and Costello graph normalization

The draft correctly says that the Weyl/Moyal computation fixes the formal target coefficients and does not reconstruct the elliptic BV source. main

Make this even clearer:

$$\text{Moyal algebra theorem} \neq \text{Costello graph theorem.}$$

State:

1. The reduced open-line Weyl model gives

$$f \star g = m \circ \exp\left(\frac{\hbar}{2} P\right)(f \otimes g).$$

2. This proves the algebraic target.
3. A Costello graph theorem must still identify:
   - elliptic complex,
   - gauge fixing,
   - propagator,
   - brane-defect kernel,
   - counterterms,

- anomaly cancellation,
- connected trace extraction.

The current draft says this in places; make it an explicit theorem/problem boundary.

31. Add a full "source-target-status" diagram

Insert a diagram like:

$$\begin{array}{ccc} C^\bullet_{\mathrm{CE}}(\mathfrak{h} \ltimes \mathfrak{h}^\vee[1]) & \overset{\Phi}{} & \mathrm{PV}(S(\mathfrak{h})) \\ \downarrow \text{Koszul dual} & & \downarrow \text{functions} \\ U(\mathfrak{h} \ltimes \mathfrak{h}^\vee[1]) & \overset{\mathrm{gr}_{PBW}}{} & S(\mathfrak{h} \oplus \mathfrak{h}^\vee[1]) \\ & & \downarrow \text{PBW label identification} \\ & & S(\bar{A} \oplus \bar{A}_{\mathrm{desc}}[1]). \end{array}$$

Then annotate:

$$u_f \mapsto O_f, \qquad c_f \mapsto \partial_{O_f},$$

$$\delta_{a,b}[1] \mapsto \Psi_{a,b} \quad \text{only in PBW special fibre,}$$

$$\delta_{a,b} \mapsto \rho_{a,b} \quad \text{as coadjoint module.}$$

This diagram will prevent misreadings.

32. Replace the "Hamiltonian factor" terminology

The phrase "Hamiltonian factor" is ambiguous. Sometimes it means:

1. degree-zero open trace functions O_f;
2. CE coordinates dual to Hamiltonian modes c_f;
3. PBW Hamiltonian generators f;
4. closed Hamiltonian field α.

Use a three-level notation:

$$\mathfrak{h} \quad \text{Hamiltonian Lie algebra}$$

$$O_f \quad \text{boundary Hamiltonian function}$$

$$c_f \quad \text{CE Hamiltonian ghost coordinate}$$

$$u_f \quad \text{CE cotangent coordinate paired with } f.$$

Then never write "$f \mapsto O_f$" without saying whether f is a PBW generator, a moment-map shadow, or a cotangent-coordinate label.

33. Rename $A_\partial = S(\mathfrak{h})$ more carefully

The open algebra $S(\mathfrak{h})$ is not literally the functions on $\mathfrak{h}$; it is the coordinate algebra on $\mathfrak{h}^\vee$, with linear functions labelled by $\mathfrak{h}$. Say:

$$A_\partial = S(\mathfrak{h}) = \mathcal{O}_{\mathrm{poly}}(\mathfrak{h}^\vee).$$

Then O_f is the linear function on $\mathfrak{h}^\vee$ corresponding to $f \in \mathfrak{h}$.

This will make the CE/PV identification conceptually correct.

34. Define the topology on $S(\mathfrak{h})$ once and enforce it

The paper uses S_{Tate}, S, S, degreewise polynomial subalgebras, completed CE cochains, and weighted completions. Add a notation table:

$$\begin{array}{cc} S(V) & \text{ordinary graded-commutative algebra} \\ S(V) & \text{completion in polynomial/CE length or Taylor degree} \\ S_{\mathrm{Tate}}(V) & \text{Tate-completed symmetric algebra} \\ S_{\mathrm{poly}}(\bar{A}) & \text{degreewise polynomial dense subalgebra} \end{array}$$

Then specify exactly which completion is used in each theorem.

35. Add "finite-degree support" lemmas for all infinite sums

Whenever you use structure constants

$$[H_{a,b}, H_{c,d}] = (ad - bc) H_{a+c-1,b+d-1},$$

state why all sums are continuous in the chosen topology.

For example, in $d_{\mathrm{CE}} u_K$, the sum over J, L satisfying

$$[H_K, H_J] = \sum_L C^L_{KJ} H_L$$

may be infinite in formal degree. Prove that for each finite output degree/window, only finitely many terms contribute. This will protect the completed CE differential.

36. Move the cross-volume horizon out of the main paper

The current table of contents includes a “Cross-volume horizon” and convention transfers to other projects. main

Move it to a separate private note or an appendix titled:

> Programmatic convention notes, not part of the theorem.

The main paper is already ambitious. Cross-volume material weakens the perceived seriousness by making the paper look like a component of a larger speculative ecosystem.

Keep only one short paragraph:

> No compact CY3, Igusa, Borcherds, or BKM claim is used in this paper. Any such transfer requires a separate matched-conventions proof.

37. Remove “twisted M-theory universality” from the main theorem path

The current draft includes an admissible protected Hamiltonian restriction and a twisted-M-theory universality theorem. This is a good outlook, but dangerous in the main proof unless it is clearly conditional. main

Recommendation:

- Move the universality theorem to an “Outlook and physical origin” section.
- Rename it “Conditional protected-sector realization.”
- Do not call it “universality” unless you prove admissibility for a class of Costello backgrounds.

Use:

$$\text{Conditional realization in admissible protected twisted-M sectors}$$

not:

$$\text{Twisted-M-theory universality}$$

unless the proof supplies the admissibility theorem.

38. Rephrase “string” carefully

The draft’s local-string definition is good: “local string theory” means the portion visible to local operators, BV fields, factorization observables, and brane open theory—not a global worldsheet theory. main

Keep that, but make it a disclaimer in the first two pages:

> The word “string” is used in the Costello-Witten string-field-theoretic local-operator sense. We do not construct global worldsheet moduli, scattering amplitudes, or a unitary Lorentzian theory.

Then avoid “M-theory” in theorem titles. Use “Hamiltonian BF brane model” in theorem titles.

39. Strengthen the unitarity section

The draft already says the model is not Lorentzian unitary and replaces unitarity with CME/QME plus OPE associativity. Keep that. Add:

$$\text{No reflection positivity claim is made.}$$

$$\text{No Hilbert-space completion is part of the local theorem.}$$

$$\text{If a unitary parent exists, this paper studies only a protected cohomological sector.}$$

This protects the physics.

40. Add a “physical dictionary” after the theorem

Insert:

Physics	Mathematics
normal coordinates of brane	$\phi_1, \phi_2 \in \mathfrak{gl}_N$
canonical symplectic form	$\mathrm{Tr}(d\phi_1 \wedge d\phi_2)$
gauge constraint	$[\phi_1, \phi_2] = 0$
BV Koszul variable	$Q\psi = [\phi_1, \phi_2]$
closed canonical transformations	$\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C}$
closed cotangent field	$\mathfrak{h}^\vee_{\mathrm{cont}}[1]$
brane measurement of f	$O_f = \mathrm{Tr}\, f(\phi_1, \phi_2)$
closed central operation	$\mathrm{PV}(S(\mathfrak{h}))$
cotangent defect current	$\rho_{a,b}$

This will make the synthesis readable.

41. Insert a “why principal parts are forced” subsection

Use the Dirac/Witten explanation from the principal-parts note. The attached note says principal parts are brane-localized currents, not functions, and explains them as the holomorphic analogue of delta-functions/momentum distributions. Pasted text

Pasted text (2)

Add this section:

Principal parts are transverse delta-currents

A Hamiltonian f is a holomorphic function. Its dual is not another holomorphic function. It is a current supported at the brane:

$$\rho_{a,b} = z_1^{-a-1} z_2^{-b-1} dz_1 dz_2.$$

The residue pairing is the holomorphic analogue of pairing a test function with a derivative of a delta distribution. The cotangent field is therefore distributionally local transverse to the brane.

This will make the physics powerful and intuitive.

42. Add an explicit no-go theorem for $\Psi \to \rho$

Make this a theorem, not just a remark.

Theorem. No polynomial descendant realization of the coadjoint module. There is no $\mathfrak{h}$-module isomorphism from the ordinary polynomial one-ψ descendant module to the principal-part coadjoint module.

Proof idea:

- $O_{1,0}$ and $O_{0,1}$ act by constant derivations on polynomial observables.
- Constant derivations are locally nilpotent on polynomials.
- Principal-part action has infinite pole-order chains.
- Local nilpotence is preserved by module isomorphism.
- Contradiction.

This theorem is already sketched in the draft; elevate it and use it to discipline all later claims. main

43. Rephrase the "Rees interpolation" as a conjectural $D_\hbar$-module theorem

Do not say the Rees parameter $\hbar$ automatically interpolates Ψ and ρ. Say:

> The PBW/Rees deformation suggests a Fourier-Borel $D_\hbar$-module interpolation between the polynomial descendant symbols and the principal-part coadjoint module.

Then state:

Conjecture. There exists an $\hbar$-flat holonomic $D_\hbar$-module $\mathcal{M}_\hbar$ supported at the brane such that:

$$\mathrm{gr}_\hbar \mathcal{M}_\hbar \cong \bar{A}_{\mathrm{desc}},$$

$$\mathcal{M}_\hbar[\hbar^{-1}] \cong P.$$

This makes the idea groundbreaking rather than false.

44. Clarify the CE/PV theorem's relationship to HKR

The draft uses HKR in the factorization-center proof. That can be risky in Tate-completed infinite-dimensional settings.

Split the argument:

1. **Algebraic identity**:

$$C^\bullet_{\mathrm{CE}}(\mathfrak{h} \ltimes \mathfrak{h}^\vee[1]) \cong \mathrm{PV}(S(\mathfrak{h})).$$

   This is direct, no HKR needed.
2. **Center identification**:

$$\mathrm{PV}(S(\mathfrak{h})) \simeq Z^{P_0}_{fact}(S(\mathfrak{h})).$$

   This uses the P_0-center/Poisson cochains formalism.
3. **HKR** only if you discuss ordinary E_1 center:

$$HH^\bullet(S(V)) \simeq \mathrm{PV}(S(V)).$$

Make clear which theorem uses which identification.

45. Add a finite-dimensional approximation proof for Tate CE/PV

For the infinite formal $\mathfrak{h}$, prove CE/PV by finite quotients:

Let

$$\mathfrak{h}_{\leq d} = \mathfrak{h}/\mathfrak{m}^{d+1}$$

where defined appropriately, or use finite Taylor windows.

For each finite window:

$$C^\bullet_{\mathrm{CE}}(\mathfrak{h}_{\leq d} \ltimes \mathfrak{h}^\vee_{\leq d}[1]) \cong \mathrm{PV}(S(\mathfrak{h}_{\leq d})).$$

Then pass to inverse limits, checking Mittag-Leffler exactness. This is more robust than asserting the Tate version directly.

46. Clean up the Tate Quillen-equivalence section

The current Tate Quillen section is very ambitious: admissible model envelopes, Hinich transfer, Berger-Moerdijk, Lefèvre-Hasegawa, Lurie, P_0-algebra categorical promotion. This can be valuable, but it risks overwhelming the main theorem.

Recommendations:

1. Move the full model-category construction to an appendix.
2. In the main text, state only:

   In the admissible filtered Tate envelope, the strict cochain isomorphism promot
3. Add a disclaimer:

   The central theorem does not depend on the raw unbounded Tate category admitting a global model structure.

4. Do not claim more than the envelope supplies.

The draft itself already says the model structure is in a fixed admissible envelope, not the raw category. Keep that firewall.

47. Make the nilpotent truncation section less central

The nilpotent truncation

$$\mathfrak{m}^3/\mathfrak{m}^{N+1}$$

is mathematically useful but physically secondary. It removes linear Hamiltonians, the $\mathfrak{sl}_2$ quadratic sector, constants, and the U(1) anomaly. The draft correctly lists lost sectors. main

Move it to:

> Appendix: finite nilpotent test sectors.

Do not make it part of the main theorem unless the reader needs it to believe the full result. It is a finite check/regulator, not the physical model.

48. Strengthen the “naive truncation fails” proof

The statement

$$\mathfrak{m}^2/\mathfrak{m}^{N+1}$$

fails because degree-two Hamiltonians form $\mathfrak{sl}_2$ is good. Show explicitly:

$$\{z_1^2, z_2^2\} = 4z_1z_2,$$

$$\{z_1z_2, z_1^2\} = -2z_1^2,$$

$$\{z_1z_2, z_2^2\} = 2z_2^2.$$

Then identify the standard $\mathfrak{sl}_2$ basis. This is a nice mathematical point.

49. Rewrite the quantum m^3/m^{N+1} obstruction cleanly

The draft says

$$P^3(z_1^4, z_2^4) = 576z_1z_2.$$

Then the Moyal bracket contains a degree-two term, so $\mathfrak{m}^3$ is not closed quantum mechanically. main

Turn this into a lemma:

Lemma. The nilpotent classical truncation is not Moyal-flat.

Proof:

$$P^3(z_1^4, z_2^4) = 576z_1z_2,$$

so

$$\{z_1^4, z_2^4\}_\hbar = 16z_1^3z_2^3 + 24\hbar^2z_1z_2 + \cdots .$$

The $\hbar^2z_1z_2$ term exits $\mathfrak{m}^3$.

This is a strong result and should be easy to find.

50. Add an “unreduced versus reduced” notation convention

Use superscripts consistently:

$$A_\partial^{red}, \qquad A_\partial^{unred}, \qquad A_\partial^{pp}, \qquad A_\partial^{Ham}.$$

Define:

$$A_\partial^{Ham} = S(\mathfrak{h})$$

after scalar reduction.

$$A_\partial^{pp} = S(\mathfrak{h}) \otimes S(P[1])$$

as reduced principal-part extension.

$$\mathrm{Obs}^{BV}_{open,N}$$

for the unreduced BV factorization algebra.

This will stop “reduced post-contraction” from being repeated in prose without clear notation.

51. Clean up theorem numbering and cross-references

The latest draft references many theorem numbers: 0.2.26, 0.2.126, 0.2.124, 0.2.144, 0.2.148, 0.2.173, 0.2.175, 0.2.170. In a 130-page draft, this is hard to track.

Add a “map of dependencies” after the theorem:

Result	Depends on
CE/PV theorem	Lie structure constants + CE/PV differential
principal parts	Grothendieck residue
stable traces	Procesi-Razmyslov + Koszul differential
quantum Moyal	Weyl algebra + Capelli radial parts
weighted Tate	Mittag-Leffler finite windows
Costello graph realization	open

This will make the paper navigable.

52. Add a “notation index”

The paper has many symbols. Add an index:

$\bar{A}$	$\mathbb{C}[z_1, z_2]/\mathbb{C}\cdot 1$
$\mathfrak{h}$	$\mathbb{C}[[z_1, z_2]]/\mathbb{C}\cdot 1$
P	principal-part residue module
O_f	boundary Hamiltonian function
$B_f(a)$	smeared boundary current
c^I	CE coordinate dual to Hamiltonian generator
u_I	CE coordinate dual to shifted cotangent generator
θ^I	$\partial/\partial O_I$
$\Psi_{a,b}$	polynomial one-ψ descendant
Θ_ρ	principal-part boundary current
$[\bar{c}]$	scalar anomaly class

This is essential.

53. Make P_I a density-dual object, not arbitrary distributions

The draft defines

$$P_I = D'_c(I)\otimes P.$$

If you replace $\mathfrak{h}_I$ by $\Omega^\bullet_c(I)\otimes\mathfrak{h}$, then its continuous dual is something like currents/densities on I tensored with P, with degree shifts.

Define carefully:

$$\mathfrak{h}_I^\vee = (\Omega^\bullet_c(I))^\vee\otimes P = \Omega^{-\bullet}(I)\otimes\mathcal{D}'(I)\otimes P$$

depending on your convention. At the top-degree current level, $a(t)dt$ pairs with a distribution $B(t)$.

Do not multiply two arbitrary distributions. The bracket

$$aB$$

is legal when a is smooth and B is a distribution. But a phrase like “multiplication of smooth functions and distributions is associative and commutative” needs precision: distributions do not multiply generally. In your bracket, you only multiply a smooth test function by a distribution. Say that.

In the principal-part sector:

$$\{B_f(a), \Theta_\rho(B)\} = \Theta_{f\cdot\rho}(aB)$$

is valid because $a \in C_c^\infty(I)$ acts on $B \in D_c'(I)$. Do not imply $B \cdot C$ is used.

54. Add a distribution-locality subsection

The principal-part sector is distributionally local transverse to the brane. The brane-line part uses compactly supported currents. Add:

> The product aB is multiplication of a distribution by a smooth function. No product of two arbitrary distributions is required. The bracket of two Θ-currents is declared zero, so no BC product appears.

This will avoid a serious analytic objection.

55. Fix the phrase "strict P_0 structure follows from delta-function locality"

The delta-function equal-time bracket gives a local Poisson bracket, but strictness also depends on working in the reduced post-contraction model and ignoring contact-term renormalization. Phrase:

> In the classical reduced current model, the equal-time delta bracket gives a strict P_0 bracket. In the unreduced BV/factorization model, the corresponding statement is represented by homotopies and contact-term choices.

The current theorem says strict non-homotopical P_0 follows from delta-function locality.

main Add the reduced-model caveat.

56. Strengthen the global-local section

The global local-Hamiltonian restriction is good. Keep it. But add a one-line theorem:

$$0 \to \mathbb{C}_X \to \mathcal{O}_X \to \mathcal{O}_X/\mathbb{C}_X \to 0$$

gives

$$H^0(X, \mathcal{O}_X/\mathbb{C}_X) \to H^1(X, \mathbb{C}_X)$$

as the period obstruction.

Then say:

> The main theorem is formal local and depends only on the completed Darboux disk. Globalization requires descent for local Hamiltonian sheaves and anomaly choices.

This avoids overglobalization.

57. Clarify the relationship to BCOV/Kodaira-Spencer

The draft already says the surface model is not a truncation of CY3 BCOV. Keep that. Add:

$$\text{Hamiltonian BF on a holomorphic symplectic surface} \neq \text{BCOV on a Calabi-Yau th}$$

Use “Hamiltonian BF” as primary terminology. Use “Kodaira-Spencer analogue” only as motivation.

58. Clarify “closed mixed Hamiltonian sector” fields

The closed fields are:

$$\alpha \in \Omega^\bullet(\mathbb{R}^2) \otimes \Omega^{0,\bullet}(\mathbb{C}^2) \otimes \mathfrak{h}[1],$$

$$\beta \in \Omega^\bullet(\mathbb{R}^2) \otimes \Omega^{0,\bullet}(\mathbb{C}^2) \otimes \mathfrak{h}^\vee_{\mathrm{cont}}[2].$$

Add a table of ghost-zero components and their degrees. The draft gives component descriptions; make them easy to check.

59. Fix or justify the BF pairing degree

The draft says the BV pairing has degree -1 and gives degree calculations. Add a small derivation:

If α has shift $[1]$ and β has shift $[2]$, then

$$|\alpha| = p - 1, \qquad |\beta| = q - 2.$$

For top form $p+q=4$,

$$|\alpha|+|\beta|=p+q-3=1.$$

Then integration has degree? Make sure this matches your claimed degree -1. If your convention includes density degree or BV dual shift, spell it out.

A sign/degree mismatch here would be an easy target.

60. Make the open ghost-zero truncation proof more physical

The current proof says setting nonzero ghost-number components to zero is consistent because ghost-number preservation prevents discarded equations from surviving. That is good. Add:

> This is a classical truncation, not a statement about the full BV quantum measure. Quantum corrections may reintroduce the discarded fields unless protected by symmetry.

This protects the physics.

61. Add a finite-N statement for the commuting-pair quotient

At finite N, the Koszul complex

$$Q\psi=[\phi_1,\phi_2]$$

computes functions on the derived commuting scheme modulo GL_N-invariants. But the commuting scheme is not always a complete intersection for all N in naive form. Avoid saying the Koszul complex is exact beyond what is proven.

Phrase:

> The BV complex is the Koszul presentation of the derived zero locus of the moment map. Its H^0 is the classical invariant ring of the underived commuting-pair quotient

when the derived structure is ignored; higher cohomology records derived intersection information.

This is safer and more modern.

62. Clarify gauge BRST versus invariant reduction

The draft says GL_N-invariants are taken in characteristic zero, so the Chevalley part acts trivially on trace functions. Add:

We compute invariant functions on the derived moment-map zero locus, not the fu

The full BRST quotient includes ghosts. If you suppress them by taking invariants, say the group is linearly reductive and invariants are exact.

63. Rework the finite stable invariant theorem citation and malformed labels

The current older text had malformed references like `[13]razmyslov` and `[2]tsygan`; the new draft may have fixed some but search still found malformed labels in earlier versions. main

Do a full search-replace audit:

- `[13]razmyslov` → `\cite{razmyslov}` or `\cite[...]{razmyslov}`
- `[2]tsygan` → `\cite{tsygan}`
- Ensure Loday-Vallette, Priddy, Hinich, Berger-Moerdijk, Lefèvre-Hasegawa numbers align.

Add bibliography entries for:

- Procesi invariant theory
- Razmyslov trace identities
- Loday-Quillen
- Tsygan
- Loday-Vallette
- Costello-Gwilliam
- Costello renormalization
- Costello-Li BCOV

- Gwilliam-Williams holomorphic string
- Beilinson-Drinfeld if weighted Tate/chiral analogy remains

64. Add a theorem-proof boundary for LQT

The draft uses LQT as a template, not a direct proof. Keep this explicit. The latest version says LQT supplies the stable-trace analogy and the reduced computation uses Procesi-Razmyslov plus the explicit Koszul differential. main

Add:

$$\text{LQT is motivational here, not the proof of the Hamiltonian sector.}$$

This avoids a reviewer saying the constrained commuting-pair Koszul complex is not directly LQT.

65. Add a “sector exclusions” theorem

State exactly what is excluded:

$$\mathrm{Tr}(1), \qquad \mathrm{Tr}(\psi), \qquad \text{higher-}\psi\text{ primitive homology}, \qquad \text{finite-}N\text{ trace identities},$$

Then state why each is excluded.

This will prevent “sector” from looking ad hoc.

66. Add an “included/excluded” table for the open algebra

Class	Included?	Reason
$\mathrm{Tr}(\phi_1^k\phi_2^l), k+l>0$	yes	$\bar{A}$
$\mathrm{Tr}(1)=N$	unit yes, primitive no	scalar reduction
$\Psi_{k,l}, k+l>0$	yes in PBW shadow	$\bar{A}_{\mathrm{desc}}$
$\mathrm{Tr}(\psi)$	no	dual to removed constant
two-ψ primitives	no	outside Hamiltonian sector
Θ_ρ	yes only in enlarged pp model	distributional cotangent

This will make the sector transparent.

67. Strengthen the scalar-central formula in every bracket theorem

Every time you state:

$$\{B_f(a), B_g(b)\} = B_{\{f,g\}}(ab),$$

add:

$$\text{in the scalar-reduced Hamiltonian quotient.}$$

Then include unreduced formula:

$$\{B_f(a), B_g(b)\} = B_{\{f,g\}_{\bar{A}}}(ab) + N\,\omega(f,g)\int_I a(t)b(t)\,dt.$$

The current theorem includes this central piece in part (vi). Make sure every earlier bracket theorem references it. main

68. Add a low-degree sanity-check subsection

Include:

$$\{z_1, z_2\} = 1 \equiv 0 \quad \text{in } \bar{A},$$

but

$$\{\operatorname{Tr}\phi_1, \operatorname{Tr}\phi_2\} = N.$$

Then:

$$\{z_1^2, z_2^2\} = 4z_1z_2,$$

$$\{\operatorname{Tr}\phi_1^2, \operatorname{Tr}\phi_2^2\} = 4\operatorname{Tr}(\phi_1\phi_2)$$

after connected reduction.

This makes conventions checkable.

69. Add a “finite rank warning”

At finite N, trace identities remain. The stable theorem holds degreewise. Put this warning before every use of freely generated trace algebras:

$$\text{Free trace generation is degreewise stable, not finite-}N.$$

The draft says this; make it a recurring flag in theorem statements. main

70. Recast the quantum theorem's source notation

The current quantum statement writes something like:

$$\Phi_\hbar^{(0)}(f) = [J_N(f)]_{N\to\infty}.$$

But in the classical derived-center theorem, functions O_f correspond to u_f, not f as a CE Hamiltonian coordinate.

So for quantum derived-center notation, use:

$$\Phi_\hbar^{(0)}(u_f) = J_\infty(f),$$

and

$$\Phi_\hbar^{(0)}(c_f) = \text{quantum derivation / commutator operation.}$$

If the degree-zero quantum theorem only treats trace functions, say it treats the u_f-coordinate/function part of the derived center.

71. Define the quantum $P_0, \hbar$-center carefully

The notation

$$Z_{fact}^{P_0,\hbar}$$

needs definition. At quantum level the bracket is no longer a classical P_0 bracket; it is a deformation, often an E_1/associative center or a Hochschild complex with $\hbar^{-1}$ commutator.

Define:

$$d_\hbar = [\pi_\hbar, -]$$

if you mean quantum Poisson cochains, or:

$$Z^{E_1}_{fact}(\mathrm{Obs}^{\hbar}_{open}) \simeq HH^{\bullet}(\mathrm{Obs}^{\hbar}_{open})$$

if you mean associative central operations.

Do not blur P_0 and quantum E_1.

72. Separate Weyl/Moyal deformation of the disk from Weyl algebra of matrices

There are two Weyl algebras:

1. formal disk Weyl algebra:

$$(\mathbb{C}[[z_1, z_2]][[\hbar]], \star)$$

2. matrix Weyl algebra:

$$W_N = D_\hbar(\mathfrak{gl}_N)$$

The trace map

$$J_N(f) = \mathrm{Tr}\,\mathrm{Op}_W(f)(X, Y)$$

connects them. Define both before using either.

73. Write the Capelli triangular renormalization explicitly

If the quantum theorem depends on Capelli-renormalized stable traces, give an explicit triangular formula for small degrees:

$$J_N(z_1 z_2) = \mathrm{Tr}\,\mathrm{Sym}(XY) + c_1 \hbar N,$$

$$J_N(z_1^2 z_2^2) = \mathrm{Tr}\,\mathrm{Sym}(X^2 Y^2) + c_1 \hbar N\,\mathrm{Tr}\,\mathrm{Sym}(XY) + c_2 \hbar^2 N^2.$$

Even if coefficients depend on convention, show the pattern.

This will make the Capelli renormalization tangible.

74. Put the radial-parts proof in "finite N, then stable N" order

Do not jump to stable connected $N \to \infty$. Prove:

1. finite-N Weyl trace descends to reduction;
2. finite-N radial-parts identity;
3. finite-N commutator identity modulo scalar/Capelli class;
4. stable connected extraction.

Each step should be a lemma.

75. Add a "quantum scalar anomaly" theorem with Rees specialization

You already have the idea:

$$\text{classical } \omega \leftrightarrow \text{quantum } \hbar N.$$

Make it a theorem independent of the full Moyal theorem:

$$U_\hbar(\bar{A}[\bar{c}])/(\hbar) \cong S(\bar{A}[\bar{c}]).$$

Then:

$$[X, Y] = \hbar \bar{c}.$$

Boundary:

$$[\operatorname{Tr} X, \operatorname{Tr} Y] = \hbar N.$$

This is elegant and physically meaningful.

76. Remove or rewrite "single nontrivial class" wording

Use:

$$\text{distinguished nontrivial scalar-extension class}$$

rather than:

$$\text{single nontrivial class}$$

unless you have classified all H^2.

77. Add a “no global unitarity” theorem-style statement

The unitarity section can be short but precise:

> This is a cohomological protected sector. The relevant consistency equations are CME/QME and factorization associativity. No positive Hilbert-space inner product is constructed or required.

This is already in the draft; keep it near the start, not buried. main

78. Add examples of physical observables

To make physics stronger, include examples:

- $O_{1,0} = \mathrm{Tr}\,\phi_1$: center-of-mass normal coordinate.
- $O_{0,1} = \mathrm{Tr}\,\phi_2$: conjugate center-of-mass coordinate.
- $O_{2,0}, O_{1,1}, O_{0,2}$: quadratic deformations generating $\mathfrak{sl}_2$ linear symplectic transformations.
- $\rho_{1,0}, \rho_{0,1}$: first derivative delta-currents transverse to the brane.
- $u_{1,0}$: CE cotangent coordinate mapping to $O_{1,0}$.

This will help readers.

79. Make the $\mathfrak{sl}_2$ quadratic sector physically explicit

The quadratic Hamiltonians

$$z_1^2,\ z_1 z_2,\ z_2^2$$

generate linear symplectic transformations of the disk. They are not nilpotent. Explain physically:

$$H_2 \cong \mathfrak{sp}_2 \cong \mathfrak{sl}_2.$$

This makes the nilpotent truncation’s loss obvious.

80. Add a “what would prove the full Costello theorem” checklist

The draft’s graph-realization problem is good. Make it a formal checklist:

To prove the full Costello realization, one must construct:

1. an elliptic mixed bulk BV complex extending the formal jet model;
2. a brane-defect boundary condition for $\mathbb{R} \times \{0\}$;
3. a gauge fixing Q^{GF};
4. a heat kernel K_t;
5. a propagator $P_{\epsilon,L}$;
6. a bulk-to-boundary kernel;
7. a renormalization scheme;
8. counterterms solving QME;
9. scalar anomaly renormalization;
10. connected trace normalization;
11. first graph coefficient 1;
12. third graph coefficient $1/24$;
13. proof that all odd coefficients match Moyal.

The current draft already lists many of these; consolidate them into one boxed problem.

81. Treat figures as either real graph notation or remove them

If the paper has schematic graph figures, upgrade them:

Each graph should display:

$$\Gamma, \quad |\operatorname{Aut}\Gamma|, \quad P_\partial, \quad \text{boundary order}, \quad \text{kernel orientation}, \quad \text{coefficient}.$$

If they are purely schematic, label them “schematic only; not used in proof.”

82. Add a proof that the closed BF action satisfies CME

The draft says Jacobi and D derivation imply CME. Add the explicit BF AKSZ calculation:

$$S = \int \langle \beta, D\alpha + \frac{1}{2}[\alpha, \alpha] \rangle.$$

Then:

$$\{S, S\} = 2 \int \left\langle D\alpha + \frac{1}{2}[\alpha, \alpha], D\alpha + \frac{1}{2}[\alpha, \alpha] \right\rangle_{\text{appropriate}} + \text{Jacobi terms}$$

or use the standard BF master-equation lemma. State the invariant pairing and coadjoint action are necessary.

83. Ensure the semidirect bracket degree signs are correct

For

$$\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}^\vee[1],$$

$\mathfrak{h}^\vee[1]$ is odd. Check:

$$[f, \eta_\lambda] = \eta_{\mathrm{ad}_f^\vee \lambda}$$

with signs determined by the shift.

$$[\eta_\lambda, \eta_\mu] = 0$$

because the shifted cotangent part is abelian.

Add the graded Jacobi proof.

84. Add an explicit proof that $d_{\mathrm{CE}}^2 = 0$ for u_K

Compute:

$$d^2 u_K = d(C_{KJ}^L u_L c^J) = C_{KJ}^L C_{LR}^M u_M c^R c^J - \frac{1}{2} C_{KJ}^L C_{RS}^J u_L c^R c^S.$$

Then show it vanishes by representation property of the coadjoint action / Jacobi.

This will make the central theorem complete.

85. Fix all “basis dependence” language

The CE/PV isomorphism is basis-independent, but the formula is basis-dependent. Say:

> The formulas below are written in a continuous homogeneous basis. The map is independent of this choice because it is the canonical identification between functions and polyvectors on the linear Poisson space $\mathfrak{h}^\vee$.

Similarly, the residue basis $\rho_{a,b}$ is coordinate-dependent under formal Darboux transformations, but the local-cohomology residue pairing is coordinate-invariant up to the symplectic volume form. Add this.

86. Add formal symplectomorphism covariance

You already mention Darboux coordinate independence. Add explicit statement:

For $\varphi \in \mathrm{Symp}(\mathbb{C}^2, 0)$,

$$f \mapsto f \circ \varphi^{-1},$$

$$\rho \mapsto \varphi_* \rho,$$

$$O_f \mapsto O_{f \circ \varphi^{-1}}$$

and the CE/PV map commutes.

This gives coordinate-free strength.

87. Clarify the closed β-field as cotangent field, not “anti-field” confusion

Physicists may confuse β with antifield. Say:

> β is the BF cotangent field; in BV it participates in the shifted symplectic pairing, but it is not merely the antifield of α in a gauge-fixed presentation.

This will help.

88. Add a glossary of “Hamiltonian,” “cotangent,” “descendant,” “principal part”

These terms are overloaded. Define them in one place.

89. Clean the title

Current title “Remarks on Mixed Holomorphic-Topological Strings” undersells and overgeneralizes. Suggested titles:

1. **Dirac Brane Probes and the Derived Poisson Center of Hamiltonian BF Theory**
2. **Closed Hamiltonian BF Theory as the Derived Center of a Dirac-Reduced Brane Probe**
3. **Principal Parts, Matrix Probes, and the Local Closed-Open Center**
4. **A Local Dirac-Witten Model for Mixed Holomorphic-Topological Branes**

Best title:

$$Closed\ Hamiltonian\ BF\ Theory\ as\ the\ Derived\ Poisson\ Center\ of\ a\ Dirac\ Brane\ Pr$$

Subtitle:

A formal local mixed holomorphic-topological model on $\mathbb{R}^2 \times \mathbb{C}^2$

90. Add a “main contributions” list

After the abstract:

1. Dirac derivation of the open matrix brane action.
2. Stable trace/Koszul computation of the open Hamiltonian sector.
3. CE/PV derived-center theorem.
4. Principal-part residue realization of $\mathfrak{h}^\vee$.
5. U(1) scalar anomaly class and Capelli shift.
6. Quantum Weyl/Moyal target and radial-parts theorem/status.
7. Weighted Tate regulator and analytic QME problem.

This will make the novelty unmistakable.

91. Replace “groundbreaking” claims with theorem names

Do not say the paper advances human knowledge in prose. Let the theorem names do it:

- **Derived center theorem**
- **Principal-part cotangent theorem**
- **Dirac probe theorem**
- **Scalar anomaly theorem**
- **Radial-Moyal theorem**

Strong theorem names make the ambition legible without rhetoric.

92. Add “locality from first principles” section

The draft already has three notions of locality: support locality, topological locality, formal holomorphic locality. main

Rewrite them as a derivation:

1. Canonical equal-time brackets give support-local contact terms:

$$\delta(t-s).$$

2. Q-exact translations give topological locality:

$$L_{\partial_t} = [Q, \iota_{\partial_t}].$$

3. Dolbeault Q-exact antiholomorphic translations give holomorphic locality:

$$L_{\partial/\partial\bar{z}_i} = [Q, \iota_{\partial/\partial\bar{z}_i}].$$

4. Principal parts are local cohomology classes supported at p.

Then conclude:

$$\text{Factorization is the algebra of local measurements, not added categorical decoratic}$$

93. Add “lack of locality” where appropriate

The model is not local in the Lorentzian finite-propagation sense. Say:

> The formal holomorphic direction is not a finite-propagation spacetime direction. It is a jet direction. Locality there means holomorphic factorization and formal support at p, not causal propagation.

This addresses physics concerns.

94. Add a “what is not proved” section before conclusion

Not as modesty—rather as precision.

Not proved:

1. full global twisted-M-theory reduction;
2. unreduced BV principal-part current realization;
3. full quantum descendant/cotangent lift;
4. Costello graph realization;
5. QME anomaly cancellation for the mixed brane-defect system;
6. regulator independence of weighted Tate;
7. Rees/Fourier interpolation between Ψ and ρ.

Then say which problems are now precisely formulated.

95. Add a conclusion that states the synthesis

End with:

> Closed fields are not merely functions evaluated on branes; they are derived centra

Then:

> The cotangent closed field is not a polynomial descendant; it is a transverse residue

That is the memorable physics/mathematics synthesis.

96. Fix theorem status of “full closed CE/cotangent lift”

The older draft had the full lift as Problem 0.2.20; the latest draft says the CE/PV theorem resolves the reduced cochain-level lift but still leaves unreduced BV/factorization and quantum descendants. main

Update all old problem statements:

- The **abstract CE/PV cochain-level derived center** is solved.
- The **reduced post-contraction principal-part boundary model** is solved.
- The **unreduced BV/factorization principal-part lift** remains open/conditional.
- The **quantum cotangent descendant lift** remains conditional.

This distinction must be globally consistent.

97. Add "old problem resolved" notation

If the draft still references "Problem 0.2.46" as the full closed CE/cotangent factorization-center lift, specify which part is resolved:

$$\text{Problem 0.2.46(a): abstract CE/PV source convention} \quad \text{resolved by Theorem 0.2.2}$$

$$\text{Problem 0.2.46(b): unreduced brane-defect BV current realization} \quad \text{open.}$$

This prevents contradiction.

98. Write a replacement for Theorem 0.1.20

Use this structure:

Theorem. Local Hamiltonian closed-open center theorem.
Let

$$\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C}$$

and

$$\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}^\vee_{\mathrm{cont}}[1].$$

Let

$$A_\partial = S(\mathfrak{h})$$

with Kirillov-Poisson bracket.

Then:

1. **Derived center:**

$$C^\bullet_{\mathrm{CE,cont}}(\mathfrak{g}) \cong \mathrm{PV}(A_\partial) \simeq Z^{P_0}_{fact}(A_\partial).$$

2. **Generator dictionary:**

$$c^I \mapsto \theta^I, \qquad u_I \mapsto O_I.$$

3. **Moment-map shadow:**

$$u_f \mapsto O_f = \operatorname{Tr} f(\phi_1, \phi_2).$$

4. **Principal-part dual:**

$$\mathfrak{h}_{\mathrm{cont}}^{\vee} \cong P$$

   by residue.

5. **Stable PBW open shadow:**

$$\operatorname{gr}_{PBW} U(\mathfrak{g})$$

   degreewise matches

$$S(\bar{A} \oplus \bar{A}_{\mathrm{desc}}[1]).$$

6. **Scalar anomaly:**
   the unreduced lift is obstructed by $[\bar{c}]$.
7. **Quantum status:**
   degree-zero Weyl/Moyal theorem as separately proven; cotangent/Costello graph lift conditional.

This theorem is both strong and precise.

99. Add precise language for "moment-map shadow"

Define:

$$J : \mathfrak{h} \to A_\partial, \qquad J(f) = O_f.$$

Then in the derived center theorem, $J(f)$ is:

$$J(f) = \Phi(u_f).$$

The phrase "moment-map shadow" should always refer to the composite:

$$\mathfrak{h} \xrightarrow{f \mapsto u_f} C^\bullet_{\mathrm{CE}}(\mathfrak{g}) \xrightarrow{\Phi} A_\partial.$$

Not to $f \mapsto c_f$.

100. Add an “ordinary center is not enough” proof

Because A_∂ is graded-commutative, multiplication by O_f is ordinarily central. That is not the theorem. The theorem is Poisson centrality/derived center.

Add:

> Ordinary Hochschild centrality of O_f is tautological in the reduced graded-commutative model. The nontrivial statement is that O_f has Lichnerowicz differential equal to its Hamiltonian vector-field operation:
>
> $$d_\pi O_f = [\pi, O_f] = X_{O_f}.$$
>
> This is exactly the CE differential of u_f.

This sharpens the claim.

101. Add explicit formula for $d_\pi O_f$

For general f:

$$d_\pi O_f = \sum_i O_{\{f,e_i\}}\theta^i$$

or with basis:

$$d_\pi O_K = C^L_{KJ} O_L \theta^J.$$

Then say:

> Thus O_f is not closed as a Poisson cochain unless f is central; it is a degree-zero function whose derived-center differential is the Hamiltonian vector field it generates.

This is conceptually central.

102. Add a “physical reading of $d_\pi O_f$” paragraph

In physics:

$$d_\pi O_f = X_{O_f}.$$

So a boundary Hamiltonian function determines an infinitesimal canonical transformation of all open observables. This is exactly the central operation produced by moving the closed insertion through the brane OPE.

This ties CE differential to contact terms.

103. Make the P_0-center definition explicit

Define a P_0-algebra as a commutative dg algebra with degree 1 Poisson bracket or degree? Depending on convention. Then define its center:

$$Z^{P_0}(A) = (\mathrm{PV}(A), [\pi, -])$$

for the open Poisson algebra A. The bracket on the center is the Schouten bracket.

Without this definition, readers may object that "center" is being used informally.

104. Add a "why CE of $T^*[1]\mathfrak{h}$ equals functions on derived center" conceptual proof

Use shifted symplectic language:

$$T^*[1]B\mathfrak{h}$$

has functions:

$$\mathcal{O}(T^*[1]B\mathfrak{h}) = C^\bullet_{\mathrm{CE}}(\mathfrak{h} \ltimes \mathfrak{h}^\vee[1]).$$

The open phase space is $\mathfrak{h}^\vee$ with linear Poisson structure. Polyvector fields on $\mathfrak{h}^\vee$ are functions on $T^*[1]\mathfrak{h}^\vee$. Under linear duality:

$$T^*[1]\mathfrak{h}^\vee \sim \mathfrak{h}^\vee \oplus \mathfrak{h}[1].$$

This gives the same coordinates:

$$O_I, \theta^I.$$

This geometric picture is powerful.

105. Add a “Dirac bracket versus Poisson bracket” clarification

The open phase space is constrained:

$$[\phi_1, \phi_2] = 0.$$

You use the original canonical Poisson bracket on gauge-invariant functions and pass to Hamiltonian reduction. Explain:

- The constraints are first-class.
- The Poisson bracket of gauge-invariant functions descends to the reduced phase space.
- The bracket on trace functions is computed upstairs and then reduced.

Do not call it a Dirac bracket unless you mean gauge-fixed second-class Dirac bracket. Use “Dirac reduction” for constraints.

106. Make $\mathrm{Tr}\, f(\phi_1, \phi_2)$ ordering conventions precise

At finite N, noncommuting matrices require ordering. The draft chooses ordered lift and proves adjacent swaps are Q-exact. Add:

$$\mathrm{Tr}\, f(\phi_1, \phi_2)$$

means the class in $H^0(Q)$ independent of ordered lift.

At quantum level, it means Weyl ordering:

$$\mathrm{Tr}\, \mathrm{Op}_W(f)(X, Y).$$

Keep these separate.

107. Add a “classical ordering independence” theorem

The theorem should read:

For any two words w, w' with same multidegree,

$$\mathrm{Tr}\, w(\phi_1, \phi_2) - \mathrm{Tr}\, w'(\phi_1, \phi_2) = Q(\cdots).$$

Proof via adjacent swaps. Then the boundary map from commutative f is well-defined.

108. Add "quantum ordering dependence is not Q-exact automatically"

At quantum level, Weyl ordering is a choice, and normal-ordering shifts matter. State:

> The classical Q-exact ordering independence is replaced quantum mechanically by a Weyl-ordering convention plus Capelli renormalization.

This protects against conflating classical and quantum.

109. Add "finite N trace identities are not killed by Q"

Adjacent swaps become exact, but finite N trace identities remain. Stable degreewise limit removes them. State:

$$H^0(R_N^{GL_N}, Q)$$

still satisfies finite-N identities. Only after stable $N \to \infty$ do primitives become freely generated.

110. Treat $\mathrm{Tr}(\psi)$ carefully

$\mathrm{Tr}\,\psi$ is a genuine cycle. It is excluded because it is dual to the removed constant. Add:

$$Q\,\mathrm{Tr}\,\psi = \mathrm{Tr}[\phi_1, \phi_2] = 0.$$

Then:

$$\mathrm{Tr}\,\psi \leftrightarrow \rho_{0,0} \leftrightarrow 1$$

and removed by scalar reduction.

This is a nice consistency check.

111. Add a "full higher-ψ problem" section

The draft has Problem 0.2.36 in older version. Keep a problem:

> Determine the full primitive stable Koszul homology of
>
> $$\mathbb{C}\langle x, y, \psi\rangle_{\mathrm{cyc}}, \quad Q\psi = [x, y].$$
>
> The present paper only uses the no-ψ and one-ψ Hamiltonian sector.

This will be valuable for future research.

112. Add a "relation to cyclic homology" appendix

LQT is a template. You can add a speculative/conjectural relation:

The complex with $Q\psi = [x, y]$ resembles cyclic homology of the derived commutative algebra of the plane or Koszul resolution of commuting matrices. Explain but do not overclaim.

113. Fix notation A, $\bar{A}$, A_∂, $A_\hbar$

The draft sometimes uses A, $\bar{A}$, A_∂, A_∂^{Ham}, $A_\hbar$. Add:

$$A_{\mathrm{poly}} = \mathbb{C}[z_1, z_2],$$

$$\bar{A} = A_{\mathrm{poly}}/\mathbb{C},$$

$$\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C},$$

$$A_\partial = S(\mathfrak{h}),$$

$$A_{\partial,\mathrm{poly}} = S(\bar{A}).$$

Use these consistently.

114. Define "continuous dual" in two cases

Strict continuous dual:

$$(\prod_{d\geq 1} H_d)^\vee_{\mathrm{cont}} = \bigoplus_{d\geq 1} H_d^\vee.$$

Weighted coefficient dual:

$$\mathfrak{h}^{\vee,w}_{cont} = \prod_{d\geq 1} w(d)^{-1} H_d^\vee$$

is not the strict continuous dual.

Put this in a boxed note.

115. Add a proof that the unweighted Casimir does not converge

The draft has it. Keep it short and visible:

$$C_{\mathfrak{h}} = \sum_{d,i} e_{d,i} \otimes e_d^i.$$

Second factor has nonzero component in every $H_d^\vee$, so it lies in

$$\prod_d H_d^\vee,$$

not

$$\bigoplus_d H_d^\vee.$$

Therefore

$$C_{\mathfrak{h}} \notin \mathfrak{h} \otimes \mathfrak{h}^\vee_{\mathrm{cont}}.$$

This is an important analytic obstruction.

116. Clarify that the CE/PV theorem does not require the Casimir

The draft says the Quillen equivalence and CE/PV map are independent of Casimir convergence. Keep and emphasize:

$$\text{Cochain algebraic derived center does not require a BV propagator.}$$

$$\text{Costello graph realization does require a coefficient kernel.}$$

This is a strong conceptual separation.

117. Reassess “universal bar-cobar/Koszul template”

The current theorem says the construction is universal for any pro-finite-dimensional Lie algebra satisfying Tate-conilpotency, but the full Hamiltonian algebra with linear and H_2 modes is not lower-central conilpotent. main

Make sure the theorem does not imply the full $\mathfrak{h}$ uses the conilpotent template. Say:

> The CE/PV isomorphism for the formal symplectic disk is proved directly by generator calculation. The conilpotent bar-cobar theorem is a separate template valid in its own range.

This avoids a logical gap.

118. Recast the Tate Quillen section as “optional categorical enhancement”

Label it:

> Optional categorical enhancement: admissible Tate bar-cobar envelope.

This prevents it from being seen as necessary for the main theorem.

119. Add exact external theorem citations for the factorization center

The paper should cite precise propositions in Costello-Gwilliam/Lurie/Calaque-Pantev-Toën-Vaquié-Vezzosi/Cattaneo-Felder if using Poisson center = Poisson cochains. Since I cannot verify the exact bibliographic best choice from the file alone, add a TODO:

> Insert precise reference for P_0-center of a Poisson algebra as Poisson cochains.

Do not leave it as folklore.

120. Add a “comparison with Gwilliam-Williams” compact paragraph

The draft does this. Keep it short:

- GW: holomorphic bosonic string on Riemann surface, BRST ghosts, critical dimension.
- This paper: local target-space Hamiltonian BF sector on $\mathbb{R}^2 \times \mathbb{C}^2$.

No imported critical dimension.

121. Remove apologetic phrasing

Avoid:

- “not a theorem”
- “only a shadow”
- “not yet”

unless in claim ledger.

Instead use:

- “The theorem is the reduced formal local theorem.”
- “The analytic graph realization is a separate problem.”
- “The PBW descendant sector is a special-fibre theorem.”

This sounds stronger.

122. Add “proof skeleton” before long proofs

For every long theorem, add:

> Proof strategy:
>
> 1. “identify generators;”
> 2. “compute differential on generators;”

3. “verify bracket compatibility;”
4. “extend by Leibniz and completion.”

This improves readability.

123. Shorten the introduction by moving technical machinery later

The current introduction contains the umbrella theorem, weighted Tate, nilpotent truncation, Quillen adjunction, Hadamard parametrix, cross-volume convention contracts. That is too much.

Recommended introduction structure:

1. Dirac brane probe.
2. Closed Hamiltonian BF.
3. Derived center theorem.
4. Principal parts.
5. Stable PBW shadow.
6. Quantum/anomaly status.
7. Theorem list.

Move weighted Tate, Quillen, nilpotent truncation, cross-volume to later sections/appendices.

124. Add a “minimal reading path”

At the end of introduction:

> The core classical theorem is contained in §§0.2.1–0.2.6 and §§principal-parts. The quantum and analytic extensions are in later sections. The categorical Tate and weighted regulator appendices may be skipped on first reading.

This helps.

125. Add exact replacement for the problematic part (ii)

Replace current part (ii):

The two are not in conflict: they are different bases of the same $\mathfrak{g}$-module, interpolated by the Rees parameter $\hbar$.

with:

The two are not in conflict because they answer different questions. The $\Psi_{a,b}$ are PBW-special-fibre polynomial descendant symbols in the stable open Koszul homology. The $\rho_{a,b}$ are the deformation-correct coadjoint modes of $\mathfrak{h}^\vee_{\mathrm{cont}}$, realized as residue currents. They carry the same bidegree labels but not the same $\mathfrak{h}$-module structure. A Rees/Fourier $D_\hbar$-module interpolating these realizations is a further construction, not a consequence of the present PBW identification.

126. Add exact replacement for problematic part (iii)

Replace:

$$(a \otimes f)^\vee[-1] \mapsto B_f(a)$$

with:

$$u_{a(t)dt\otimes f} \mapsto B_f(a) = \int_I a(t)\,\mathrm{Tr}\, f(\phi_1(t), \phi_2(t))\, dt,$$

and:

$$c_\lambda \mapsto \theta_\lambda.$$

Then add:

Here u_x is the CE coordinate dual to the shifted cotangent generator paired with $x \in \mathfrak{h}_I$. The CE coordinate dual to $x \in \mathfrak{h}_I$ maps to the corresponding constant polyvector operation, not to the boundary function.

127. Add exact replacement for the cotangent reduced theorem

State:

$$A^{pp}_\partial(I) = S(\mathfrak{h}_I) \otimes S(P_I[1])$$

is an **enlarged reduced principal-part defect-current model**.

Then:

$$\partial_{bb}^{pp}(f) = B_f, \qquad \partial_{bb}^{pp}(\rho[1]) = \Theta_\rho.$$

Bracket:

$$\{B_f, \Theta_\rho\} = \Theta_{f\cdot\rho}.$$

Status:

> This is a reduced P_0-model. Its unreduced BV/factorization realization is not automatic.

128. Add exact replacement for quantum theorem status

Use:

> The degree-zero quantum theorem concerns the u_f-function part of the derived center. It proves the Moyal commutator for Capelli-renormalized stable Weyl traces. It does not prove the quantum cotangent/principal-part descendant sector, which remains conditional on weighted Tate/QME input.

This aligns with Corollary 0.2.174. main

129. Add a theorem about open-line Weyl product independent of Costello

The draft has Proposition 0.2.177 saying the reduced open-line Weyl model gives the Moyal product and uses no Costello specialization. main

Make this a standalone theorem:

$$F \star G = m \circ \exp\left(\frac{\hbar}{2} P_\partial\right)(F \otimes G).$$

Then:

> This theorem is internal to the reduced first-order open-line Weyl model. It is not the

Costello graph realization.

This is clear and strong.

130. Add a “normalization conventions” appendix for Moyal coefficients

Include:

$$f \star g = m \circ e^{\hbar P/2}(f \otimes g).$$

Then:

$$[f,g]_\star = 2 \sum_{r \text{ odd}} \frac{\hbar^r}{2^r r!} P^r(f,g).$$

So:

$$r = 1: \quad \hbar P(f,g),$$

$$r = 3: \quad \frac{\hbar^3}{24} P^3(f,g).$$

This avoids coefficient confusion.

131. Add exact formula for P^r on monomials

Use:

$$P^r(z_1^k z_2^l, z_1^m z_2^n) = \sum_{a=0}^{r} (-1)^a \binom{r}{a} (k)_{r-a} (l)_a (m)_a (n)_{r-a} z_1^{k+m-r} z_2^{l+n-r}.$$

Then state only odd r enter commutator.

This is useful and concrete.

132. Check $P^3(z_1^4, z_2^4)$ coefficient consistency

Using the formula:

For $f = z_1^4$, $g = z_2^4$, only $a = 0$ contributes if $P = \partial_{z_1} \otimes \partial_{z_2} - \partial_{z_2} \otimes \partial_{z_1}$:

$$P^3(f, g) = (4)_3(4)_3 z_1^1 z_2^1 = 24 \cdot 24 z_1 z_2 = 576 z_1 z_2.$$

Then in the normalized bracket, coefficient is

$$\frac{\hbar^2}{24} \cdot 576 = 24\hbar^2 z_1 z_2.$$

The draft has this; keep it.

133. Add "finite N versus N in $\mathfrak{m}^N$" notation warning

You use N both for matrix size and truncation order. Avoid this. Use:

$$R$$

for matrix rank, and

$$M$$

or

$$D$$

for truncation degree.

Example:

$$\mathfrak{h}^{(3)}_{\leq D} = \mathfrak{m}^3 / \mathfrak{m}^{D+1}.$$

This avoids confusion.

134. Add "rank N" and "truncation D" global convention

At the start:

$$N = \text{brane rank}, \qquad D = \text{Taylor truncation degree}.$$

Then update all nilpotent truncation statements.

135. Fix "dimension $1+4$" phrasing

The draft says $1+4$-real-dimensional spacetime $\mathbb{R}^2 \times \mathbb{C}^2$ around a line. But $\mathbb{R}^2 \times \mathbb{C}^2$ is $2+4=6$ real dimensions. The brane is one real dimension. If you say "$1+4$," explain that one of the $\mathbb{R}^2$ directions is the brane and one is transverse topological. Better:

$$\mathbb{R}_{\text{brane}} \times \mathbb{R}_{\text{top,normal}} \times \mathbb{C}^2_{\text{hol}}.$$

Total real dimension 6, brane dimension 1.

136. Add notation for x_1, x_2 topological coordinates

The brane is $\mathbb{R}_{x_1} \times \{x_2 = 0\} \times \{p\}$. Use this consistently.

137. Avoid "boundary" when it is a defect

The brane is a line defect, not a boundary of spacetime. The draft says this in Construction 0.2.16. Keep it globally.

Use:

$$\text{bulk-to-defect}$$

or

$$\text{bulk-to-brane}$$

instead of "boundary" when speaking geometrically. "Boundary operator" is okay in open-string language, but define it.

138. Add "bulk-to-boundary" versus "bulk-to-defect" convention

At the start:

> We use "boundary" in the open-string/operator sense and "defect" in the spacetime sense. The brane is a defect, not a boundary component of $\mathbb{R}^2 \times \mathbb{C}^2$.

139. Add exact proof of Darboux equivariance of substitution

Under formal symplectomorphism φ:

$$f \mapsto f \circ \varphi,$$

$$(\phi_1, \phi_2) \mapsto \varphi(\phi_1, \phi_2)$$

is not literal for noncommuting matrices at finite N, but after Q-cohomology/stable commutative reduction it is well-defined. State this caveat.

140. Treat noncommutative formal symplectomorphisms carefully

At finite N, substituting a formal power series symplectomorphism into matrices requires ordering. Avoid strong equivariance claims before the commuting-pair reduction. Say:

> After passage to $H^0(Q)$, ordering ambiguities are exact, so the formal Darboux change acts on the commutative Hamiltonian labels.

141. Add an “index conventions for monomials” note

Use:

$$H_{a,b} = z_1^a z_2^b, \qquad a, b \geq 0, \quad a + b > 0.$$

Set:

$$H_{m,n} = 0$$

if $m < 0$, $n < 0$, or $m + n = 0$ after scalar reduction.

This should appear before structure constants.

142. Check the sign of the Hamiltonian vector field

You use:

$$X_f = (\partial_{z_1} f)\partial_{z_2} - (\partial_{z_2} f)\partial_{z_1}, \qquad \iota_{X_f}\omega = -df.$$

Then $X_f(g) = ?$

Compute:

$$X_f(g) = (\partial_1 f)(\partial_2 g) - (\partial_2 f)(\partial_1 g) = \{f, g\}.$$

Good. Keep this convention globally.

143. Add a "sign sanity check" for z_1, z_2

$$X_{z_1} = \partial_{z_2}, \qquad X_{z_2} = -\partial_{z_1}.$$

$$\{z_1, z_2\} = 1.$$

This will fix signs.

144. Verify principal-part action signs against the convention

Given

$$f \cdot \rho$$

defined by

$$\langle f \cdot \rho, g\rangle = -\langle \rho, \{f, g\}\rangle,$$

derive the monomial formula explicitly in the paper.

Let $f = z_1^p z_2^q$, $g = z_1^m z_2^n$. Then:

$$\{f, g\} = (pn - qm) z_1^{p+m-1} z_2^{q+n-1}.$$

Pair with $\rho_{a,b}$:

$$-\langle \rho_{a,b}, \{f, g\}\rangle = -(pn - qm)\delta_{a,p+m-1}\delta_{b,q+n-1}.$$

So

$$m = a - p + 1, \qquad n = b - q + 1.$$

Coefficient:

$$-(p(b-q+1) - q(a-p+1)) = -(pb - pq + p - qa + pq - q) = -(pb - qa + p -$$

Thus:

$$f \cdot \rho_{a,b} = -(pb - qa + p - q)\rho_{a-p+1,b-q+1}.$$

Good. Include this derivation.

145. Clarify "negative-index targets zero"

If $a - p + 1 < 0$ or $b - q + 1 < 0$, the target basis vector is absent and the action is zero. State this.

146. Add principal-part examples computed from the formula

$$z_1 \cdot \rho_{a,b} = -(b+1)\rho_{a,b+1}$$

or check carefully from formula:

Set $p = 1, q = 0$:

$$-(pb - qa + p - q) = -(b+1).$$

Target:

$$\rho_{a-1+1,b-0+1} = \rho_{a,b+1}.$$

So:

$$z_1 \cdot \rho_{a,b} = -(b+1)\rho_{a,b+1}.$$

Set $p = 0, q = 1$:

$$-(0 - a + 0 - 1) = a + 1.$$

Target:

$$\rho_{a+1,b}.$$

So:

$$z_2 \cdot \rho_{a,b} = (a+1)\rho_{a+1,b}.$$

These are the correct examples. Put them in the paper. They are clearer than “lowering.”

147. Fix any place saying z_1 lowers principal-part indices

If any note or draft sentence says z_1 lowers ρ, correct it. It raises b-pole order under the formula above.

148. Compare with polynomial descendant action explicitly

Polynomial descendant action:

$$O_{p,q} : \Psi_{a,b} \mapsto (pb - qa)\Psi_{a+p-1,b+q-1}.$$

For z_1:

$$p = 1, q = 0 : \quad \Psi_{a,b} \mapsto b\Psi_{a,b-1}.$$

For principal parts:

$$\rho_{a,b} \mapsto -(b+1)\rho_{a,b+1}.$$

Then state:

$$\text{opposite shift, different coefficient, different module.}$$

This is the cleanest demonstration.

149. Do not call the polynomial action simply “raising”

For z_1, it lowers b. For higher p, q, it may raise total degree by $p + q - 2$. Say:

$$\text{symmetric PBW action}$$

instead of raising.

150. Add a “module mismatch lemma”

Lemma. The map $\Psi_{a,b} \mapsto \rho_{a,b}$ does not intertwine z_1.

Proof:

$$\Psi_{a,b} \mapsto b\Psi_{a,b-1} \mapsto b\rho_{a,b-1},$$

but

$$\rho_{a,b} \mapsto -(b+1)\rho_{a,b+1}.$$

Not equal.

Then:

$$\Psi \mapsto \rho \text{ is only a label identification.}$$

151. Add “the Rees interpolation must change category”

If you want an interpolation, say it cannot be inside ordinary polynomial modules. It must pass through distributions or $D_\hbar$-modules.

152. Rework principal-parts note into paper section

The attached principal-parts explanation is excellent but has a few incorrect or imprecise raising/lowering sentences. Bring in its conceptual content but correct the formulas. Cite or integrate it. The note emphasizes that principal parts are brane-localized currents and polynomial Ψ's are PBW special-fibre representatives rather than deformation-level coadjoint modes. Pasted text Pasted text (2)

153. Make the “position vs momentum” analogy mathematically precise

Add:

$$\text{functions} \leftrightarrow \text{coordinates on } \mathfrak{h}^\vee,$$

$$\text{polyvector fields} \leftrightarrow \text{differential operators/derivations},$$

$$\text{principal parts} \leftrightarrow \text{distributions supported at the brane.}$$

Avoid overextending the quantum momentum analogy beyond motivation.

154. Add "defect current" notation for $\Theta_\rho(B)$

Use:

$$\Theta_\rho(B)$$

where $B \in D'_c(I)$. For ordinary smeared principal-part mode with smooth density $b(t)dt$, write:

$$\Theta_\rho(b) = \Theta_\rho(b(t)dt)$$

if appropriate. Define the pairing with $\Omega^\bullet_c(I)$.

155. Clarify Θ_ρ parity

Since $P[1]$ is shifted, Θ_ρ is odd/even depending on convention. State:

$$|\Theta_\rho| = ?$$

Then verify:

$$\{\Theta_\rho, \Theta_\sigma\} = 0$$

is compatible with graded antisymmetry.

156. Add a finite-dimensional toy model

Before $\mathbb{C}[[z_1, z_2]]$, give a finite-dimensional Lie algebra $\mathfrak{l}$. Show:

$$C^\bullet_{\mathrm{CE}}(\mathfrak{l} \ltimes \mathfrak{l}^\vee[1]) \cong \mathrm{PV}(S(\mathfrak{l})).$$

This will make the infinite Tate case intuitive.

157. Add the exact toy proof

Let $\mathfrak{l}$ have basis e_i. Then:

$$C^\bullet_{\mathrm{CE}}(\mathfrak{l} \ltimes \mathfrak{l}^\vee[1]) = S(u_i) \otimes \Lambda(c^i).$$

$$\mathrm{PV}(S(\mathfrak{l})) = S(O_i) \otimes \Lambda(\theta^i).$$

Map:

$$u_i \mapsto O_i, \quad c^i \mapsto \theta^i.$$

This becomes the main theorem by completion.

158. Add "what is $Z^{P_0}_{fact}$ on an interval?"

For a locally constant factorization algebra from A, the center stalk is Hochschild cochains. For Poisson compatibility, the P_0-center is Poisson cochains. On intervals:

$$Z^{P_0}_{fact}(A_\partial)(I) \simeq \mathrm{PV}(A_\partial(I)).$$

State how extension by zero acts on polyvectors.

159. Fix factorization product direction arrows

Factorization algebras are covariant for opens with extension by zero in the observables-as-compact-support convention. CE cochains are contravariant unless using chains/cosheaves. The draft has a diagram with vertical reversal. Make sure all arrows are correct.

Recommendation: define every object as a **factorization algebra of compactly supported observables** so maps are covariant.

For CE cochains on currents, be careful:

$$I \mapsto \mathfrak{h}_I$$

is a cosheaf. Then

$$I \mapsto C^\bullet_{\mathrm{CE}}(\mathfrak{h}_I)$$

is contravariant if CE cochains use duals. To get a factorization algebra, one often uses CE chains or observables as functions on fields with compact support. Clarify this.

This is a serious technical point. If you cannot make CE cochains covariant cleanly, define interval objects through polyvectors on $S(\mathfrak{h}_I)$ and use extension by zero there.

160. Consider using factorization coalgebras for CE cochains

If $I \mapsto \mathfrak{h}_I$ is a cosheaf, then CE chains are naturally a factorization coalgebra and CE cochains are dual. In locally constant finite-dimensional settings this is harmless; in Tate settings it matters.

Add a paragraph:

> We use compactly supported observables/density-duals so that the completed CE cochain assignment is covariant under extension by zero. Equivalently, one may formulate the source as CE chains and dualize in the admissible Tate envelope.

Then prove the convention.

161. Add exact treatment of compact support and distributions

The source $\mathfrak{h}_I = \Omega_c^\bullet(I) \otimes \mathfrak{h}$ has compact support. Its dual involves distributions on I. The function $B_f(a)$ uses a compactly supported density, not a distribution. Keep source and dual separate.

162. Add a “density versus function” convention

The current $a(t)$ under $\int a(t)dt$ is really a density. Write:

$$a(t)dt \in \Omega_c^1(I).$$

Then avoid ambiguity in orientation and coordinate changes.

163. Add orientation convention for the brane line

Fix dt and orientation of $\mathbb{R}$. The symplectic potential depends on orientation.

164. Clarify boundary terms in the open action

When reducing:

$$\frac{1}{2}\int \mathrm{Tr}(\phi_1 d\phi_2 - \phi_2 d\phi_1)$$

to

$$\int \mathrm{Tr}\, \phi_1 d\phi_2$$

you discard a boundary term. State:

- compactly supported variations;
- local operators in the interior;
- or intervals with fixed boundary values.

The draft does this; keep it.

165. Add proof of gauge invariance of open action

The draft includes it. Keep it but move into Dirac theorem.

166. Clarify the role of A as a 1-form gauge field

A is both Lagrange multiplier and connection. In 1D, every connection has no curvature but imposes moment map. Say this.

167. Add a BRST/BV field table

For open fields:

field	form/Ext degree	ghost number
A	$\Omega^1 \otimes 1$	0
ϕ_1	$\Omega^0 \otimes \epsilon_1$	0
ϕ_2	$\Omega^0 \otimes \epsilon_2$	0
$\psi = A^*$	$?$	1 or -1

Make signs/degrees consistent.

168. Define ψ degree consistently

The draft calls $\psi = A^* \in \mathfrak{gl}_N[1]$ odd. Depending on cohomological convention, $Q\psi = [\phi_1, \phi_2]$ means $|Q| = 1$, so if ϕ degree 0, then ψ degree -1 or 1 depending chain/cochain convention. State your convention once.

169. Revisit "$S(V)$ means free graded-commutative algebra" with shifts

If $V[1]$ has parity reversed, then $S(V[1])$ is exterior on even V. Make sure:

$$S(\bar{A}_{\mathrm{desc}}[1])$$

is exterior if descendants are odd. The draft says $S(V)$ free graded-commutative. Good.

170. Add "odd ideal abelian" sign check

If $P[1]$ is odd, then in graded Lie algebra:

$$[\rho[1], \sigma[1]] = 0$$

is symmetric or antisymmetric depending bracket degree. Verify with your P_0 convention.

171. Avoid calling $P[1]$ an "odd local-dual ideal" without degree convention

Say “shifted local-dual ideal” and give degree.

172. Insert a proof of Jacobi for A_{∂}^{pp}

The draft says Jacobi follows from coadjoint representation and smearing. Keep but refine distribution multiplication. Use:

$$[f,[g,\rho]]-[g,[f,\rho]]=[[f,g],\rho]$$

by coadjoint representation. Smearing:

$$a(bB)-b(aB)=abB$$

for smooth a,b and distribution B.

173. Add “do not multiply distributions” note in Jacobi proof

For the bracket involving two Θ's, zero avoids distribution products.

174. Add exact definition of $f\cdot\rho=\pi_{pp}\{f,\rho\}$

Since ρ is a top form, $\{f,\rho\}$ needs definition. Define it as Lie derivative:

$$f\cdot\rho=-\mathcal{L}_{X_f}\rho$$

or whichever sign matches coadjoint action. Then show it equals projection of the Poisson bracket expression.

This is more geometrically correct than “Poisson bracket of a function with a form” unless you define it.

Recommended:

$$f\cdot\rho=-\mathcal{L}_{X_f}\rho$$

because coadjoint action on densities is negative Lie derivative. Verify sign against formula.

175. Use “residue currents” instead of “Laurent forms” when emphasizing physics

Mathematically P is represented by Laurent top forms modulo holomorphic parts. Physically they are currents. Use both.

176. Add a Grothendieck residue citation

If not already in bibliography, cite a standard source for local residues/local cohomology.

177. Re-evaluate “unique $\mathfrak{h}$-equivariant continuous pairing”

If the pairing is between P and $\mathfrak{h}$, equivariance follows. But “unique $\mathfrak{h}$-equivariant continuous pairing of total degree zero” might require proof. Either prove or replace with “canonical.”

178. Add a local cohomology exact sequence

For $R = \mathbb{C}[z_1, z_2]$,

$$H^2_{\mathfrak{m}}(R) \cong \mathbb{C}[z_1^{-1}, z_2^{-1}]/(\mathbb{C}[z_1, z_1^{-1}] + \mathbb{C}[z_2, z_2^{-1}])$$

or the standard Čech model. This grounds P.

179. Add the Čech representative

$$H^2_{\mathfrak{m}}(R) = R[z_1^{-1}, z_2^{-1}]/(R[z_1^{-1}] + R[z_2^{-1}]).$$

Basis:

$$z_1^{-a-1} z_2^{-b-1}.$$

Then multiply by $dz_1 dz_2$.

180. Tie principal parts to Verdier/local duality

Mention:

$$H^2_{\mathfrak{m}}(R)\, dz_1 dz_2$$

is the Matlis dual of the completed local ring $R^\wedge_{\mathfrak{m}}$. This is exactly the continuous dual.

This is mathematically powerful.

181. Add “Matlis dual” phrase

$$P \simeq \mathrm{Hom}_{cont}(\mathbb{C}[[z_1, z_2]]/\mathbb{C}, \mathbb{C})$$

is the reduced Matlis dual.

182. Recast principal-part theorem in Matlis dual language

Theorem. The reduced Matlis dual of $\mathfrak{h}$ is P, and the Hamiltonian Lie derivative realizes the coadjoint action.

This is much more standard than “principal-part residue module” alone.

183. Add proof of continuity of residue pairing

A continuous functional on $\mathbb{C}[[z_1, z_2]]$ with $\mathfrak{m}$-adic topology depends on finitely many coefficients. But P as direct sum of $\rho_{a,b}$ exactly gives finite coefficient-extraction sums. State this.

Weighted product P_w is a different completion.

184. Clarify completed $\mathfrak{h}^\vee$ in closed BF action

If closed BF uses $\mathfrak{h}^\vee_{cont}$ as direct sum, then β-fields have finite Taylor-dual support. If weighted completion uses product, then β has infinite dual support. State which model each theorem uses.

185. Add “strict model” versus “weighted model” notation

$$\mathfrak{g}_{\mathrm{str}} = \mathfrak{h} \ltimes P[1],$$

$$\mathfrak{g}_w = \mathfrak{h}_w \ltimes P_w[1].$$

Do not use the same $\mathfrak{g}$ for both.

186. Add “dense inclusion” theorem

$$P \hookrightarrow P_w$$

for finite-support principal parts. Then the strict polynomial sector embeds into weighted regulator.

187. Add exact statement of graphwise RG limitation

The weighted theorem says each fixed graph is controlled. It does not prove convergence of sum over graphs. Add:

> Graphwise boundedness is not perturbative renormalizability by itself. The full effective interaction is a formal sum over graphs and loop order; locality and QME are separate.

188. Move Costello-Li failure analysis to appendix or keep concise

The draft’s Costello-Li hypothesis failure section is detailed. It is useful but may distract. In the main text, summarize:

- noncompact;
- mixed topological-holomorphic;
- brane defect;
- $\mathfrak{gl}_N$ not $\mathfrak{gl}(N|N)$;
- coefficient Tate direction.

Then put details in appendix.

189. Avoid sounding like you refute Costello-Li

Say:

> Costello-Li's theorem is not applicable to this local mixed defect model without additional work.

not:

> Costello-Li route fails

unless precisely framed.

190. Add a "why ordinary $\mathfrak{gl}_N$ retains U(1) anomaly" paragraph

Costello-Li supertrace cancellation uses $\mathfrak{gl}(N|N)$. Here ordinary $\mathfrak{gl}_N$ has trace N. Tie it to $[\bar{c}]$.

191. Clarify "Capelli shift as U(1) charge"

The draft says $\hbar N$ is charge of central direction. Add:

$$\mathrm{Tr}(I) = N.$$

Thus the central moment relation has scalar N.

192. Add "background independence" caveat

The formal local model depends only on the formal Darboux disk, but global backgrounds require choices. State clearly.

193. Add a "minimal theorem-only version" appendix

A reader should be able to extract a pure math theorem:

Theorem. For any Lie algebra $\mathfrak{h}$, CE cochains of $T^*[1]\mathfrak{h}$ are Poisson cochains of $S(\mathfrak{h})$.

Theorem. For $\mathfrak{h} = \mathbb{C}[[z_1, z_2]]/\mathbb{C}$, $\mathfrak{h}^\vee_{cont}$ is the reduced local cohomology module P.

Theorem. Stable trace/Koszul computation yields $S(\bar{A} \oplus \bar{A}_{desc}[1])$.

This makes the mathematical contribution independent of physics.

194. Add a “physics-only summary” appendix

Conversely, add a 2-page nontechnical physics summary.

This helps interdisciplinary readership.

195. Check all theorem names for consistency

Avoid:

- “Koszul duality” for CE/PV center theorem.
- “factorization equivalence” when it is reduced post-contraction.
- “universal” when conditional.
- “closed string” when formal local closed sector.

Use:

- “derived-center”
- “PBW special-fibre”
- “reduced principal-part boundary model”
- “weighted regulator”
- “conditional graph realization”

196. Rewrite the table of contents

Suggested:

1. Dirac probes and local measurements
2. The local mixed BV model
3. Open brane reduction and stable traces
4. Closed Hamiltonian BF theory
5. The CE/PV derived-center theorem
6. Principal parts and defect-local cotangent currents

7. Factorization on the brane line
8. PBW special-fibre descendants
9. Scalar anomaly and Capelli shift
10. Quantum Weyl/Moyal degree-zero theorem
11. Weighted Tate regulator and analytic QME problem
12. Appendices: signs, local cohomology, one-ψ homology, radial parts, finite checks

197. Move “Tate Quillen equivalence” after CE/PV theorem

Do not introduce model category machinery before the reader understands the central theorem. Place it after, as enhancement.

198. Move “Cross-volume convention contract” out entirely

This is not part of the whitepaper. Keep as a private doc.

199. Add a final “research program” list

Future theorems:

1. Rees/Fourier interpolation $\Psi \leftrightarrow P$.
2. Unreduced BV realization of principal-part currents.
3. Regulator independence of weighted Tate.
4. Full quantum descendant lift.
5. Costello graph realization of Moyal coefficients.
6. Global descent over holomorphic symplectic surfaces.
7. Full higher-ψ primitive Koszul homology.

This turns open issues into a coherent program.

200. Highest-priority edits to make first

Do these before anything else:

1. Replace every interval map

$$(a \otimes f)^\vee[-1] \mapsto B_f(a)$$

by

$$u_{a(t)dt \otimes f} \mapsto B_f(a).$$

2. Replace $C_c^\infty(I)$ by $\Omega_c^\bullet(I)$ in locally constant factorization theorems.
3. Delete "Ψ and ρ are different bases of the same $\mathfrak{g}$-module."
4. Add the no-go lemma showing $\Psi_{a,b} \mapsto \rho_{a,b}$ is not an $\mathfrak{h}$-module map.
5. Replace "Schur rigidity" with a local-cohomology/Matlis-dual proof.
6. Rename principal-part sector as an enlarged reduced defect-current model.
7. Split the umbrella theorem into theorem blocks.
8. Add a claim-strength ledger.
9. Move cross-volume material out of the main paper.
10. Add a sign-convention appendix.

201. One-paragraph strongest-version vision

The strongest conceivable version of this whitepaper is not a speculative string note. It is a theorem-driven Dirac-Witten local model: a brane stack is a constrained matrix probe of a formal symplectic disk; its stable Hamiltonian trace algebra is the open measurement algebra; the closed Hamiltonian BF theory is the shifted cotangent theory of canonical transformations; and the closed CE observables are exactly the derived Poisson central operations on the open algebra. The polynomial one-ψ descendants are the PBW-special-fibre shadow, while the true cotangent modes are principal-part residue currents supported at the defect. The scalar $U(1)$ anomaly is the same class classically and quantum mechanically, and the degree-zero quantum sector is governed by the Weyl/Moyal deformation, with the full Costello graph realization isolated as the remaining analytic theorem.

Sources

Sources