

**CHEMISTRY
HIGHER LEVEL
PAPER 1**

Monday 18 November 2002 (afternoon)

1 hour

INSTRUCTIONS TO CANDIDATES

- Do not open this examination paper until instructed to do so.
- Answer all the questions.
- For each question, choose the answer you consider to be the best and indicate your choice on the answer sheet provided.


Periodic Table

1 H 1.01	<div>Atomic Number</div> <div>Atomic Mass</div>																2 He 4.00
3 Li 6.94	4 Be 9.01											5 B 10.81	6 C 12.01	7 N 14.01	8 O 16.00	9 F 19.00	10 Ne 20.18
11 Na 22.99	12 Mg 24.31											13 Al 26.98	14 Si 28.09	15 P 30.97	16 S 32.06	17 Cl 35.45	18 Ar 39.95
19 K 39.10	20 Ca 40.08	21 Sc 44.96	22 Ti 47.90	23 V 50.94	24 Cr 52.00	25 Mn 54.94	26 Fe 55.85	27 Co 58.93	28 Ni 58.71	29 Cu 63.55	30 Zn 65.37	31 Ga 69.72	32 Ge 72.59	33 As 74.92	34 Se 78.96	35 Br 79.90	36 Kr 83.80
37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 92.91	42 Mo 95.94	43 Tc 98.91	44 Ru 101.07	45 Rh 102.91	46 Pd 106.42	47 Ag 107.87	48 Cd 112.40	49 In 114.82	50 Sn 118.69	51 Sb 121.75	52 Te 127.60	53 I 126.90	54 Xe 131.30
55 Cs 132.91	56 Ba 137.34	57 † La 138.91	72 Hf 178.49	73 Ta 180.95	74 W 183.85	75 Re 186.21	76 Os 190.21	77 Ir 192.22	78 Pt 195.09	79 Au 196.97	80 Hg 200.59	81 Tl 204.37	82 Pb 207.19	83 Bi 208.98	84 Po (210)	85 At (210)	86 Rn (222)
87 Fr (223)	88 Ra (226)	89 ‡ Ac (227)	104 Rf (261)	105 Db (262)	106 Sg (263)	107 Bh (262)	108 Hs	109 Mt									

†	58 Ce 140.12	59 Pr 140.91	60 Nd 144.24	61 Pm 146.92	62 Sm 150.35	63 Eu 151.96	64 Gd 157.25	65 Tb 158.92	66 Dy 162.50	67 Ho 164.93	68 Er 167.26	69 Tm 168.93	70 Yb 173.04	71 Lu 174.97
---	---------------------------	---------------------------	---------------------------	---------------------------	---------------------------	---------------------------	---------------------------	---------------------------	---------------------------	---------------------------	---------------------------	---------------------------	---------------------------	---------------------------

‡	90 Th 232.04	91 Pa 231.04	92 U 238.03	93 Np (237)	94 Pu (242)	95 Am (243)	96 Cm (247)	97 Bk (247)	98 Cf (251)	99 Es (254)	100 Fm (257)	101 Md (258)	102 No (259)	103 Lr (260)
---	---------------------------	---------------------------	--------------------------	--------------------------	--------------------------	--------------------------	--------------------------	--------------------------	--------------------------	--------------------------	---------------------------	---------------------------	---------------------------	---------------------------



1. Consider the following reaction:



2.0 dm³ of 0.50 mol dm⁻³ CaCl₂(aq) is mixed with 1.0 dm³ of 2.0 mol dm⁻³ AgNO₃(aq). What are the concentrations of Ca²⁺(aq) and NO₃⁻(aq) after mixing?


	[Ca ²⁺] / mol dm ⁻³	[NO ₃ ⁻] / mol dm ⁻³
A.	0.66	0.33
B.	0.33	0.66
C.	1.0	2.0
D.	3.0	1.5

2. Formation of polyethene from calcium carbide, CaC₂, can take place as follows:



What mass of polyethene is obtained from 64 kg of CaC₂ ?

- A. 7 kg
B. 14 kg
C. 21 kg
D. 28 kg
3. Ammonia is manufactured by the synthesis of nitrogen and hydrogen as follows



56.0 g of N₂ produces 34.0 g of NH₃.

What is the percentage yield of ammonia?

- A. 50
B. 68
C. 74
D. 100

4. Isotopes are elements with
- A. the same atomic number and the same number of neutrons.
 - B. the same mass number but a different number of neutrons.
 - C. the same atomic number but a different number of neutrons.
 - D. different atomic and mass numbers but the same number of neutrons.
5. A transition metal ion X^{3+} has the electronic configuration $[\text{Ar}] 3d^4$. What is the atomic number of element X?
- A. 22
 - B. 24
 - C. 25
 - D. 27
6. Which of the following electronic configurations gives rise to the largest increase between the second and third ionisation energies?
- A. $1s^2 2s^2$
 - B. $1s^2 2s^2 2p^2$
 - C. $1s^2 2s^2 2p^6 3s^2$
 - D. $1s^2 2s^2 2p^6 3s^1$

7. Which of the following displacement reactions is possible?
- A. $\text{Br}_2(\text{aq}) + 2\text{Cl}^-(\text{aq}) \rightarrow 2\text{Br}^-(\text{aq}) + \text{Cl}_2(\text{aq})$
 - B. $\text{I}_2(\text{aq}) + 2\text{Cl}^-(\text{aq}) \rightarrow 2\text{I}^-(\text{aq}) + \text{Cl}_2(\text{aq})$
 - C. $\text{Cl}_2(\text{aq}) + 2\text{I}^-(\text{aq}) \rightarrow 2\text{Cl}^-(\text{aq}) + \text{I}_2(\text{aq})$
 - D. $\text{I}_2(\text{aq}) + 2\text{Br}^-(\text{aq}) \rightarrow 2\text{I}^-(\text{aq}) + \text{Br}_2(\text{aq})$
8. An element E of mass number 40 has the electronic configuration 2. 8. 8. 2. Which statement regarding this element is **not** correct?
- A. It belongs to group 2 of the periodic table.
 - B. It has 20 neutrons.
 - C. It belongs to period 4 of the periodic table.
 - D. The formula of its oxide is EO_2 .
9. Which ions are listed in order of **decreasing** ionic radius (highest first)?
- A. $\text{Mg}^{2+}, \text{Na}^+, \text{F}^-, \text{O}^{2-}$
 - B. $\text{O}^{2-}, \text{F}^-, \text{Na}^+, \text{Mg}^{2+}$
 - C. $\text{F}^-, \text{O}^{2-}, \text{Na}^+, \text{Mg}^{2+}$
 - D. $\text{Mg}^{2+}, \text{Na}^+, \text{O}^{2-}, \text{F}^-$

10. Consider the following coordination compounds



What are the charges on the complex ions?

	I	II	III
A.	+2	+1	0
B.	−2	−1	0
C.	0	+1	+2
D.	0	−1	−2



11. Which intermolecular forces exist in dry ice, $\text{CO}_2(\text{s})$?

- A. Covalent bonds
- B. Dipole-dipole attractions
- C. Van der Waal's forces
- D. Hydrogen bonds

12. When the species NH_2^- , NH_3 and NH_4^+ are arranged in **increasing** order of H–N–H bond angle, the correct order is


- A. NH_2^- , NH_3 , NH_4^+
- B. NH_4^+ , NH_3 , NH_2^-
- C. NH_3 , NH_4^+ , NH_2^-
- D. NH_3 , NH_2^- , NH_4^+

13. The elements X and Y have the following electronic configurations:



What is the formula of the compound formed between X and Y?

- A. XY_2
 - B. X_5Y_2
 - C. X_2Y_5
 - D. XY_5
14. Which statements about the following molecule are correct?



- I. Three carbon atoms are sp^3 hybridized.
 - II. Three carbon atoms are sp^2 hybridized.
 - III. Two carbon atoms are sp hybridized.
- A. I and II only
 - B. I, II and III
 - C. II and III only
 - D. I and III only

15. Under what conditions would a given mass of oxygen gas occupy the greatest volume?
- High temperature and high pressure
 - High temperature and low pressure
 - Low temperature and low pressure
 - Low temperature and high pressure
16. The volume of a gas measured at 27 °C and 101.3 kPa is 20.0 dm³. What final temperature would be required to increase the volume to 40.0 dm³ at 101.3 kPa?
- 54 °C
 - 300 °C
 - 327 °C
 - 600 °C

17. Consider the following reaction:



Bond enthalpies (in kJ mol⁻¹) involved in the reaction are

N ≡ N	x
H–H	y
N–H	z


Which calculation will give the value of ΔH^\ominus ?

- $x + 3y - 6z$
- $6z - x + 3y$
- $x - 3y + 6z$
- $x + 3y - 2z$

18. If 3600 J of heat is added to 180 g of $\text{C}_2\text{H}_5\text{OH}(\text{l})$, its temperature increases from 18.5°C to 28.5°C . What is the specific heat capacity of $\text{C}_2\text{H}_5\text{OH}(\text{l})$?

- A. $0.500 \text{ J g}^{-1} ^\circ\text{C}^{-1}$
 B. $2.00 \text{ J g}^{-1} ^\circ\text{C}^{-1}$
 C. $20.0 \text{ J g}^{-1} ^\circ\text{C}^{-1}$
 D. $200 \text{ J g}^{-1} ^\circ\text{C}^{-1}$



19. The following reaction takes place in an internal combustion engine:



What are the signs for ΔH^\ominus , ΔS^\ominus and ΔG^\ominus for this reaction?

	ΔH^\ominus	ΔS^\ominus	ΔG^\ominus
A.	–	+	+
B.	–	+	–
C.	–	–	–
D.	+	–	–

20. Consider the following equations:



What is the standard enthalpy change of formation (ΔH^\ominus_f) for $\text{H}_2\text{SO}_4(\text{l})$?

- A. -812 kJ
 B. $+812 \text{ kJ}$
 C. -526 kJ
 D. $+526 \text{ kJ}$

21. In general, the rate of a reaction can be increased by all of the following **except**

- A. increasing the temperature.
- B. increasing the activation energy.
- C. increasing the concentration of reactants.
- D. increasing the surface area of the reactants.

22. The following experimental data was obtained for the reaction $X + Y \rightarrow \text{products}$.

$[X] / \text{mol dm}^{-3}$	$[Y] / \text{mol dm}^{-3}$	Initial rate $/ \text{mol dm}^{-3} \text{sec}^{-1}$
0.10	0.10	4.0×10^{-4}
0.20	0.20	1.6×10^{-3}
0.50	0.10	1.0×10^{-2}
0.50	0.50	1.0×10^{-2}


What is the order of reaction with respect to X and the order of reaction with respect to Y?

- A. 2 and 0
- B. 0 and 2
- C. 2 and 1
- D. 1 and 0

23. The rate of a gaseous reaction is given by the expression $\text{rate} = k [P][Q]$. If the volume of the reaction vessel is reduced to $\frac{1}{4}$ of the initial volume, what will be the ratio of the new rate to the original rate?


- A. 1 : 4
- B. 1 : 16
- C. 4 : 1
- D. 16 : 1

24. The volume of the reaction vessel containing the following equilibrium mixture




is increased. When equilibrium is re-established, which of the following will occur?

- A. The amount of $\text{SO}_2\text{Cl}_2(\text{g})$ will increase.
 - B. The amount of $\text{SO}_2\text{Cl}_2(\text{g})$ will decrease.
 - C. The amount of $\text{Cl}_2(\text{g})$ will remain unchanged.
 - D. The amount of $\text{Cl}_2(\text{g})$ will decrease.
25. A 1.0 dm^3 reaction vessel contains initially 1.0 mol of $\text{NO}_2(\text{g})$ and 1.0 mol of $\text{N}_2\text{O}_4(\text{g})$. At equilibrium, 0.75 mol of $\text{N}_2\text{O}_4(\text{g})$ are present. What is the value of K_c ?



- A. 0.33
 - B. 0.50
 - C. 2.0
 - D. 3.0
26. What affects the amount of $\text{X}_3\text{Y}(\text{g})$ at equilibrium in the following exothermic reaction?




- A. Temperature, pressure and a catalyst
- B. Temperature and pressure
- C. Temperature only
- D. Pressure only

27. When the following 0.10 mol dm^{-3} solutions are arranged in order of **increasing** pH (lowest first), what is the correct order?



- A. $\text{NaOH}, \text{NH}_3, \text{CH}_3\text{COOH}, \text{HCl}$
- B. $\text{HCl}, \text{CH}_3\text{COOH}, \text{NH}_3, \text{NaOH}$
- C. $\text{HCl}, \text{CH}_3\text{COOH}, \text{NaOH}, \text{NH}_3$
- D. $\text{NaOH}, \text{NH}_3, \text{HCl}, \text{CH}_3\text{COOH}$

28. Consider a weak acid HA dissolved in water.



Which statements are correct?

- I. $\text{A}^-(\text{aq})$ is a much stronger base than $\text{H}_2\text{O}(\text{l})$.
- II. HA dissociates only to a very small extent in aqueous solution.
- III. The concentration of $\text{H}_3\text{O}^+(\text{aq})$ is much greater than the concentration of $\text{HA}(\text{aq})$.

- A. I, II and III
- B. II and III only
- C. I and II only
- D. I and III only

29. When the following aqueous solutions are arranged in order of **increasing** electrical conductivity (lowest first), what is the correct order?

I. $0.10 \text{ mol dm}^{-3} \text{ CH}_3\text{COOH}$

II. $0.10 \text{ mol dm}^{-3} \text{ CH}_3\text{CH}_2\text{OH}$

III. $0.10 \text{ mol dm}^{-3} \text{ CH}_3\text{COONa}$

A. I, II, III

B. III, II, I

C. I, III, II

D. II, I, III

30. A certain buffer solution contains equal concentrations of $\text{X}^-(\text{aq})$ and $\text{HX}(\text{aq})$. The K_b value for $\text{X}^-(\text{aq})$ is 1.0×10^{-10} . What is the pH of the buffer?


A. 1

B. 4

C. 5

D. 10

31. In the reaction



A. Br_2 is only oxidised.

B. Br_2 is only reduced.

C. Br_2 is neither oxidised nor reduced.

D. Br_2 is both oxidised and reduced.

32. Consider the following statements regarding electrolysis of molten lead(II) bromide.

- I. Oxidation takes place at the anode where lead ions gain electrons.
- II. Reduction takes place at the cathode where lead ions gain electrons.
- III. Oxidation takes place at the anode where bromide ions lose electrons.
- IV. Reduction takes place at the cathode where bromide ions lose electrons.

Which of the above statements are correct?

- A. I and II only
- B. I and IV only
- C. II and III only
- D. II and IV only

33. The standard electrode potentials of three elements are as follows:

X	+1.09 V
Y	+0.54 V
Z	+1.36 V

Which statement is correct?

- A. Z will oxidise $Y^-(aq)$ and $X^-(aq)$
- B. Y will oxidise $X^-(aq)$ and $Z^-(aq)$
- C. X will oxidise $Y^-(aq)$ and $Z^-(aq)$
- D. Z will oxidise $Y^-(aq)$ but not $X^-(aq)$

34. One Faraday of electricity was passed through the electrolytic cells placed in series containing solutions of $Ag^+(aq)$, $Ni^{2+}(aq)$ and $Cr^{3+}(aq)$. What mass of Ag, Ni and Cr respectively will be deposited?

[A_r values: Ag = 108, Ni = 59, Cr = 52]

- A. 36 g, 29.5 g and 52 g
- B. 108 g, 59 g and 52 g
- C. 108 g, 29.5 g and 17.3 g
- D. 108 g, 118 g and 156 g




35. Consider the following reaction:



What will be the final product if aminoethane (ethylamine) is used instead of NH_3 ?

- A. $\text{CH}_3\text{CONHCH}_2\text{CH}_3$
- B. $\text{CH}_3\text{CONHCH}_3$
- C. CH_3CONH_2
- D. $\text{CH}_3\text{CONH}_2\text{CH}_2\text{CH}_3$


36. Which of the following compounds is optically active?



37. How many different environments for hydrogen atoms are present in the ^1H NMR spectrum of the following compound?



- A. 3
B. 4
C. 5
D. 9
38. Consider the following reactions:



What are reagents I and II respectively?

- A. $\text{H}^+ / \text{Cr}_2\text{O}_7^{2-}(\text{aq})$ LiAlH_4
B. H_2/Ni LiAlH_4
C. LiAlH_4 $\text{H}^+ / \text{Cr}_2\text{O}_7^{2-}(\text{aq})$
D. $\text{H}^+/\text{MnO}_4^-(\text{aq})$ $\text{H}^+ / \text{Cr}_2\text{O}_7^{2-}(\text{aq})$
39. An organic liquid L has a relative molecular mass of 46. On heating with concentrated H_2SO_4 at 170°C , a colourless gas is evolved which decolourises $\text{Br}_2(\text{aq})$. What is the organic liquid L?
- A. $\text{CH}_3\text{CH}_2\text{OH}$
B. CH_3OCH_3
C. $\text{CH}_3\text{CH}=\text{CH}_2$
D. CH_3OH

40. The alkaline hydrolysis of primary halogenoalkanes usually follows an S_N2 mechanism. For which compound would the rate of hydrolysis be fastest?

- A. $\text{CH}_3\text{CH}_2\text{CH}_2\text{F}$
 - B. $\text{CH}_3\text{CH}_2\text{CH}_2\text{Cl}$
 - C. $\text{CH}_3\text{CH}_2\text{CH}_2\text{Br}$
 - D. $\text{CH}_3\text{CH}_2\text{CH}_2\text{I}$
-