

INTERNATIONAL
BACCALAURÉAT
BACHILLERATO



BACCALAUREATE
INTERNATIONAL
INTERNACIONAL

M98/510/H(1)M

MARKSCHEME

May 1998

MATHEMATICS

Higher Level

Paper 1

1. Since, $\sin \theta < 0$, $\cos \theta = \frac{2}{5}$, $\sin \theta = -\sqrt{1 - \cos^2 \theta} = -\sqrt{1 - \frac{4}{25}} = -\frac{\sqrt{21}}{5}$ (M1)(A1)

Hence, $\tan \theta = \frac{\sin \theta}{\cos \theta} = -\frac{\sqrt{21}}{2}$ and $\sec \theta = \frac{1}{\cos \theta} = \frac{5}{2}$ (A1)(A1)

Answers: $\sin \theta = -\frac{\sqrt{21}}{5}$, $\tan \theta = -\frac{\sqrt{21}}{2}$, $\sec \theta = \frac{5}{2}$ (C2)(C1)(C1)

2. (a) $\frac{1}{8} + 3k + \frac{1}{6}k + \frac{1}{4} + \frac{1}{6}k = 1$ (M1)

Thus, $\frac{10k}{3} = \frac{5}{8}$ and $k = \frac{3}{16}$ (A1)

(b)

x	0	1	2	3	4
$p(X=x)$	$\frac{1}{8}$	$\frac{9}{16}$	$\frac{1}{32}$	$\frac{1}{4}$	$\frac{1}{32}$

$p(0 < X < 4) = \frac{9}{16} + \frac{1}{32} + \frac{1}{4} = \frac{27}{32}$ (M1)(A1)

Answers: (a) $k = \frac{3}{16}$ (C2)

(b) $p(0 < X < 4) = \frac{27}{32}$ (C2)

3. $(\sqrt{3})^{126} = 3^{63}$ (M1)

Hence, $3^{x^2-1} = 3^{63}$ (A1)

Therefore, $x^2 - 1 = 63$ or $x = \pm 8$ (M1)(A1)

Answers: $x = \pm 8$ (C4)

4.

$$\text{Let } -2 + i2\sqrt{3} = r(\cos\theta + i\sin\theta)$$

$$\text{Then } r = |-2 + i2\sqrt{3}| = \sqrt{4 + 12} = 4$$

$$\text{and } \tan\theta = \frac{-2}{2\sqrt{3}} = -\frac{1}{\sqrt{3}}$$

$$\text{Hence } \theta = \frac{3}{2}\pi \quad (0 \leq \theta < 2\pi)$$

$$\text{Thus } z = 4e^{i(3\pi/2)} = 4e^{i(3\pi/2 + 2k\pi)}, \quad k = 0 \pm 1, \pm 2, \dots$$

(R1)

(C4)

Note: Award (C4) for $z = 4e^{i(2/3\pi)}$

5.

$$P(A \cap B) = P(A)P(B) = \left(\frac{1}{4}\right)\left(\frac{1}{8}\right) = \frac{1}{32}$$

$$P(B) = \frac{P(A \cap B)}{P(A)} = \frac{1/32}{1/4} = \frac{1}{8}$$

$$P(A) = \frac{1/32}{1/8} = \frac{1}{4}$$

$$\text{Answers: } P(A) = \frac{1}{4}, \quad P(B) = \frac{1}{8}$$

(C2)(C2)

6.

Let X be the mean test score

$$P(X > 80) = P\left(Z > \frac{80 - 60}{\frac{10}{\sqrt{2}}}\right) = P(Z > 2)$$

$$= 1 - 0.9773 = 0.0227$$

(M1)(A1)

(C4)

$$\text{Answer: } P(X > 80) = 0.0227$$

(Also accept 0.0228 which is obtainable through calculator)

Note: Some candidates may use a continuity correction as follows:

$$X \sim N(60, 10^2)$$

$$\text{Hence } P(X > 80) = P\left(Z > \frac{80.5 - 60}{10}\right) = P(Z > 2.05)$$

$$= 1 - 0.9798 = 0.0202$$

(M1)(A1)

(C4)

$$\text{Answer: } P(X > 80) = 0.0202$$

7. (a) $2 + 4(n-1) = 58$ or $4n - 2 = 58 \Rightarrow n = 15$ (M1)(A1)

(b) Sum of 15 terms of a geometric sequence with first term 2
and common ratio $\frac{1}{2}$ is $2 \left(\frac{1 - (1/2)^{15}}{1 - 1/2} \right) = 4 \left(1 - \frac{1}{2^{15}} \right)$ (M1)(A1)

Answers: (a) $n = 15$ (C2)

(b) $4 \left(1 - \frac{1}{2^{15}} \right)$ or $\frac{32767}{8192}$ (C2)

8. $E(X) = (1)\frac{2}{9} + 2\left(\frac{1}{9}\right) + 3\left(\frac{2}{9}\right) + 4\left(\frac{1}{9}\right) + 5\left(\frac{2}{9}\right) + (6)\left(\frac{1}{9}\right)$ (M1)

$= \frac{2}{9} + \frac{2}{9} + \frac{6}{9} + \frac{4}{9} + \frac{10}{9} + \frac{6}{9} = \frac{30}{9} = 3\frac{2}{3} = 3\frac{1}{3}$ (A1)

$E(X^2) = (1)^2\frac{2}{9} + (2)^2\frac{1}{9} + (3)^2\frac{2}{9} + (4)^2\frac{1}{9} + (5)^2\frac{2}{9} + (6)^2\frac{1}{9}$
 $= \frac{2}{9} + \frac{4}{9} + \frac{18}{9} + \frac{16}{9} + \frac{50}{9} + \frac{36}{9} = \frac{126}{9} = 14$

$\text{Var}(X) = E(X^2) - (E(X))^2 = 14 - \left(\frac{10}{3}\right)^2$ (M1)

$= 14 - \frac{100}{9} = \frac{126 - 100}{9} = \frac{26}{9}$ (A1)

Answers: $E(X) = \frac{10}{3}$, $\text{Var}(X) = \frac{26}{9}$ (C2)(C2)

9. $\sin x \tan x = \sin x \Rightarrow \sin x (\tan x - 1) = 0$ (M1)

$\sin x = 0$ when $x = 0$, $x = \pi$, or $x = 2\pi$ (A1)

$\tan x - 1 = 0$ when $x = \frac{\pi}{4}$ or $x = \frac{5\pi}{4}$ (M1)(A1)

The solutions are $x = 0, \frac{\pi}{4}, \pi, \frac{5\pi}{4}, 2\pi$

Answers: $x = 0, \frac{\pi}{4}, \pi, \frac{5\pi}{4}, 2\pi$ (C4)

10. The normal to the planes are $\vec{n}_1 = 2\vec{i} + 3\vec{j} - \vec{k}$ and $\vec{n}_2 = 7\vec{i} - \vec{j} + 3\vec{k}$ (A1)(A1)

Angle between the two planes is given by

$$\arccos \frac{|\vec{n}_1 \cdot \vec{n}_2|}{|\vec{n}_1| |\vec{n}_2|} = \arccos \frac{14 - 3 - 3}{\sqrt{14} \sqrt{59}} = \arccos \frac{\sqrt{826}}{8}$$

$$= 73.8^\circ$$

Answer: 73.8° (C4)

11. $f'(x) = \frac{\ln x}{\arcsin x} - \frac{\sqrt{1-x^2}}{x} (\ln x)^2$ (M1)(M1)(M1)(A1)

$$= \frac{x\sqrt{1-x^2}(\ln x)^2}{x \ln x - \sqrt{1-x^2} \arcsin x}$$

Answer: $f'(x) = \frac{x\sqrt{1-x^2}(\ln x)^2}{x \ln x - \sqrt{1-x^2} \arcsin x}$ (C4)


or any equivalent form. (Simplification of the final answer is not required.)

12. Area $= 4 \int_1^0 y \, dx = 4 \int_1^0 \sqrt{x^2 - x^4} \, dx = 4 \int_1^0 x\sqrt{1-x^2} \, dx$ (M1)(M1)

$$= \left[\left(-\frac{2}{4} \right) \left(\frac{3}{2} \right) (1-x^2)^{3/2} \right]_1^0 = \left(-\frac{3}{4} \right) (-1) = \frac{3}{4}$$


Answer: Area $= \frac{3}{4}$ (C4)

13.



$|f(x)|$

(C1)



$\frac{1}{f(x)}$

Asymptotes
(C1).
Curves (C2).
Deduct 1
mark for
each
mistake.

14.

$$\int \frac{dy}{y} = \int \cos x \, dx, \quad 0 < x < \infty \Rightarrow \ln|y| = \sin x + C$$

Since $y > 0$, $y = Ae^{\sin x}$, A being a constant

(M1)(A1)

Since, $y = 1$ when $x = \frac{\pi}{2}$, we get,

$$Ae^{\sin \pi/2} = 1 \text{ or } A = \frac{1}{e}$$

(M1)

$$\text{Hence, } y = \left(\frac{1}{e}\right) e^{\sin x} = e^{\sin x - 1}$$

(A1)

$$\text{Answer: } y = e^{\sin x - 1}$$

(C4)

Note: Some students may solve the problem by using integrating factor.
For $e^{-\int \cos x \, dx} = e^{-\sin x}$ as the integrating factor award (C1) and proceed according to the markscheme above.

15. (a)

$$6 \int_0^k (x^2 + x) \, dx = 6 \left(\frac{k^3}{3} + \frac{k^2}{2} \right) = 2k^3 + 3k^2 = 1$$

(M1)

$$\Rightarrow 2k^3 + 3k^2 - 1 = 0 \Rightarrow (k+1)(2k^2 + k - 1) = 0$$

Therefore, $k = -1$ or $k = \frac{1}{2}$

Since $k > 0$, $k = \frac{1}{2}$

(A1)

$$(b) \quad E(X) = 6 \int_{1/2}^0 (x^2 + x) x \, dx = 6 \left[\frac{x^4}{4} + \frac{x^3}{3} \right]_{1/2}^0$$

(M1)

$$= 6 \left[\frac{1}{64} + \frac{1}{24} \right] = \frac{32}{11}$$

(A1)

Answers: (a) $k = \frac{1}{2}$

(C2)

$$(b) \quad E(X) = \frac{11}{32}$$

(C2)

16. Differentiating $x^3 + y^3 = 6xy$ implicitly with respect to x , we get

$$3x^2 + 3y^2 y' = 6y + 6xy' \Rightarrow y' = \frac{2y - x^2}{y^2 - 2x} \quad (M1)(A1)$$

Slope at $(3, 3)$ is $(y')_{(3,3)} = -1 \quad (A1)$

Tangent has equation $y - 3 = (-1)(x - 3)$ i.e. $x + y = 6 \quad (A1)$

Answer: $x + y = 6 \quad (C4)$

17. $\int \arctan x \, dx = x \arctan x - \int \frac{x}{1+x^2} \, dx \quad (M1)(A1)$

$$= x \arctan x - \frac{1}{2} \int \frac{2x}{1+x^2} \, dx$$

$$= x \arctan x - \frac{1}{2} \ln(1+x^2) + C \quad (M1)(A1)$$

Answer: $x \arctan x - \frac{1}{2} \ln(1+x^2) + C \quad (C4)$

18. There is a non-zero solution if and only if

$$\begin{vmatrix} 2 & -2 & k \\ 1 & 0 & 4 \\ k & 1 & 1 \end{vmatrix} = 0 \quad (R1)$$

$$\Rightarrow 2(-4) + 2(1-4k) + k = 0 \quad (M1)(A1)$$

$$\Rightarrow -7k = 6 \text{ or } k = -\frac{6}{7} \quad (A1)$$

Answer: $k = -\frac{6}{7} \quad (C4)$

19. $f(x)$ is defined so long as $x^2 - 4 \geq 0$

But $x^2 - 4 \geq 0$ if and only if $|x| \geq 2 \Rightarrow x \leq -2$ or $x \geq 2$

So the domain is $\{x \in \mathbb{R} | x \leq -2 \text{ or } x \geq 2\}$

Since, $f(x) = e^{3x^2} + \sqrt{x^2 - 4}$, we find that $f(-2) = f(2) = e^{12}$

Further, we observe that e^{3x^2} and $\sqrt{x^2 - 4}$ increase as $x \geq 2$ or $x \leq -2$

Also $\lim_{x \rightarrow \infty} f(x) = \infty$ and $\lim_{x \rightarrow -\infty} f(x) = \infty$

So the range of f is $\{x \in \mathbb{R} | e^{12} \leq x\}$

Answer: Domain: $\{x \in \mathbb{R} | x \leq -2 \text{ or } x \geq 2\}$

Range: $\{y \in \mathbb{R} | e^{12} \leq y\}$

20. (a) Since $z = x + iy$ and $z^* = x - iy$,

$|z - 2 - i\sqrt{3}| = (\sqrt{2})|z^* - 1 + i\sqrt{3}|$ is equivalent to

$|(x - 2) + i(y - \sqrt{3})| = (\sqrt{2})|(x - 1) - i(y - \sqrt{3})|$

Thus, we get $\{(x - 2)^2 + (y - \sqrt{3})^2\}^{1/2} = (\sqrt{2})\{(x - 1)^2 + (y - \sqrt{3})^2\}^{1/2}$

On squaring both sides, we obtain,

$$(x - 2)^2 + (y - \sqrt{3})^2 = 2(x - 1)^2 + 2(y - \sqrt{3})^2$$

$$\Rightarrow x^2 - 4x + 4 = 2x^2 - 4x + 2 + (y - \sqrt{3})^2$$

$$\Rightarrow x^2 + (y - \sqrt{3})^2 = 2 \text{ or } x^2 + y^2 - 2\sqrt{3}y + 1 = 0$$

(b) This is a circle of radius $\sqrt{2}$ with its centre at $(0, \sqrt{3})$.

Answers: (a) Equation of the circle is $x^2 + (y - \sqrt{3})^2 = 2$

$$\text{or } x^2 + y^2 - 2\sqrt{3}y + 1 = 0$$

(b) Centre of the circle is $(0, \sqrt{3})$, radius is $\sqrt{2}$