

ARLAB

ME/CprE/ComS 557

Computer Graphics and Geometric Modeling

Quaternions


October 27th, 2015

Rafael Radkowski

IOWA STATE UNIVERSITY
OF SCIENCE AND TECHNOLOGY

Book

ARLAB




Vince, J (2011). Quaternions for Computer Graphics. 1st. ed. London: Springer

Why Quaternions

ARLAB

Describing a path / locomotion as a sequence of matrices can result in an unexpected outcome due to the link of the rotation axis to each other in Cartesian space with Euler angles.




- Easier to "add" a new position / orientation to the current position / orientation
- Interpolation between points is possible.

Gimble Lock Problem



The sequence of transformations (rotations in our case) can be considered as hierarchy of transformations.



$$T = R_y \cdot R_x \cdot R_z$$

$$T = \begin{bmatrix} \cos(\beta) & 0 & -\sin(\beta) & 0 \\ 0 & 1 & 0 & 0 \\ \sin(\beta) & 0 & \cos(\beta) & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 & 0 & 0 \\ \cos(\alpha) & -\sin(\alpha) & 0 & 0 \\ \sin(\alpha) & \cos(\alpha) & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} \cos(\gamma) & \sin(\gamma) & 0 & 0 \\ -\sin(\gamma) & \cos(\gamma) & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$


The transformations are linked: the axis y, and x also rotates when an object is rotated around the z-axis.

Gimble Lock Problem



Hierarchy of transformations

z
└ x
 └ y




Gimble Lock Problem




Hierarchy of transformations

z
└ x
 └ y



History

ARLAB



$$i^2 = j^2 = k^2 = -1$$

Quaternions were discovered by Sir William Rowan Hamilton on Monday October 16th 1843 in Dublin, Ireland.

Complex Numbers

ARLAB

Complex numbers introduce imaginary numbers to solve certain equations such as:

$$x + 1 = 0$$

$$x^2 = -1$$

$$x = \sqrt{-1}$$

which we know is not possible to solve with real numbers because the square of every number is always positive.

The **imaginary number** has been founded.

$$i^2 = -1$$

The set of complex numbers is the sum of a real number and an imaginary number

$$z = a + bi \quad a, b \in \mathbb{R}, i^2 = -1$$

- all real numbers are complex numbers with $b = 0$
- all imaginary numbers are complex numbers with $a = 0$

Some Rules

ARLAB

Adding and Subtracting Complex Numbers

Addition:

$$(a_1 + b_1 i) + (a_2 + b_2 i) = (a_1 + a_2) + (b_1 + b_2)i$$

Subtraction:

$$(a_1 + b_1 i) - (a_2 + b_2 i) = (a_1 - a_2) + (b_1 - b_2)i$$

Multiply a Complex Number by a Scalar:

$$\lambda(a + bi) = \lambda a + \lambda bi$$

Some Rules

ARLAB

Product of Complex Numbers:

$$z_1 = (a_1 + b_1 i)$$

$$z_2 = (a_2 + b_2 i)$$

$$z_1 z_2 = (a_1 + b_1 i)(a_2 + b_2 i)$$

$$= a_1 a_2 + a_1 b_2 i + b_1 a_2 i + b_1 b_2 i$$

$$= (a_1 a_2 - b_1 b_2) + (a_1 b_2 + b_1 a_2) i$$

Square of Complex Numbers:

$$z^2 = (a + bi)(a + bi)$$

$$= (a^2 - b^2) + 2abi$$

Complex Conjugate:

$$z = (a + bi)$$

$$|z| = \sqrt{zz^*}$$

$$= \sqrt{(a + bi)(a - bi)}$$

$$= \sqrt{a^2 + b^2}$$

The conjugate of a complex number is a complex number with the imaginary part negated



IOWA STATE UNIVERSITY
OF SCIENCE AND TECHNOLOGY

Some Rules

ARLAB

Absolute Value of a Complex Number:

$$z = (a + bi)$$

$$|z| = \sqrt{zz^*}$$

$$= \sqrt{(a + bi)(a - bi)}$$

$$= \sqrt{a^2 + b^2}$$

We can use the conjugate of a complex number to compute the absolute value (or norm, or magnitude) of a complex number.

Power of i

$$i^0 = 1$$

$$i^1 = i$$


$$i^2 = -1$$

$$i^3 = ii^2 = -i$$

$$i^4 = i^2i^2 = 1$$

$$i^5 = ii^4 = i$$

$$i^6 = ii^5 = -1$$



This is a sequence $1, i, -1, -i, 1, i, -1, -i, 1, \dots$

The Complex Plane

ARLAB

$$i^0 = 1$$

$$i^1 = i$$


$$i^2 = -1$$

$$i^3 = ii^2 = -i$$

$$i^4 = i^2i^2 = 1$$

$$i^5 = ii^4 = i$$

$$i^6 = ii^5 = -1$$




This is a sequence 1, i, -1, -i, 1, i, -1, -i, 1,

We can say that if we multiply a complex number by i, we can rotate the complex number through the complex plane at 90 degree increments.

Rotation with Complex Numbers

ARLAB



Take an arbitrary point p in the complex plane:


○ $p = 2 + i$

Multiply this point with i .

○
$$\begin{aligned} q &= p \cdot i \\ &= (2 + i)i \\ &= 2i + i^2 \\ &= -1 + 2i \end{aligned}$$

Rotation with Complex Numbers

ARLAB



Take an arbitrary point p in the complex plane:

○ $p = 2 + i$

Multiply point q with i .

○ $q = -1 + 2i$

○ $r = q \cdot i$


$$= (-1 + 2i)i$$

$$= -i + 2i^2$$

$$= -2 - i$$

Rotation with Complex Numbers

ARLAB



Take an arbitrary point p in the complex plane:

○ $p = 2 + i$

Multiply point r with i .

○ $r = -2 - i$

○ $s = r \cdot i$

$$= (-2 - i)i$$

$$= -2i - i^2$$

$$= 1 - 2i$$

Rotation with Complex Numbers

ARLAB

We can also perform arbitrary rotations in the complex plane by defining a complex number of the form:

$$q = \cos \theta + i \sin \theta$$

We obtain the general formula when multiplying a point p with q :

$$p = a + b i$$

$$q = \cos \theta + i \sin \theta$$

$$pq = (a + bi)(\cos \theta + i \sin \theta)$$


$$a' + b'i = a \cos \theta - b \sin \theta + (a \sin \theta + b \cos \theta)i$$

(pq rewritten)

or in matrix form:

$$\begin{bmatrix} a' & -b' \\ b' & a' \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$$

to rotate an arbitrary point counter-clockwise.



This was in 2D / one plane out or three (rotation of a point around one axis in a plane).

How do we extend this to 3D?



Quaternions

Quaternions

ARLAB

We extend this to 3-dimensional space by adding two imaginary numbers to our number system in addition to i . One number for each additional plane.

The general form to express quaternions is:

$$q = s + xi + yj + zk \quad s, x, y, z \in \mathbb{R}, i, j, z \in \mathbb{I}$$

scalar value s

vector of complex numbers

with

$$i^2 + j^2 + k^2 = ijk = -1$$

and

$$\begin{array}{lll} ij = k & jk = i & ki = j \\ ji = -k & kj = -i & ik = -j \end{array}$$

Quaternions


ARLAB

When mapping these quaternion rules to a Cartesian coordinate system,

$$\begin{array}{lll} ij = k & jk = i & ki = j \\ ji = -k & kj = -i & ik = -j \end{array}$$

they behave similar to cross products.

$$\begin{array}{lll} \mathbf{x} \times \mathbf{y} = \mathbf{z}, & \mathbf{y} \times \mathbf{z} = \mathbf{x}, & \mathbf{z} \times \mathbf{x} = \mathbf{y} \\ \mathbf{y} \times \mathbf{x} = -\mathbf{z}, & \mathbf{z} \times \mathbf{y} = -\mathbf{x}, & \mathbf{x} \times \mathbf{z} = -\mathbf{y} \end{array}$$



Quaternions

ARLAB

Hamilton recognized that the imaginary numbers i , j , and k can be used to represent three cartesian unit vectors i , j , and k with the same properties of imaginary numbers, such that

$$i^2 + j^2 + k^2 = ijk = -1$$

Quaternions can be represented as an **ordered pair**:

$$q = [s, \mathbf{v}] \quad s \in \mathbb{R}, \mathbf{v} \in \mathbb{R}^3$$

or with individual components

$$q = [s, x\mathbf{i} + y\mathbf{j} + z\mathbf{k}] \quad s, x, y, z \in \mathbb{R}$$

which allows us to show similarities to complex numbers.

Substitute to swap between notations.

Mathematical Rules

ARLAB

Adding and subtracting quaternions:

$$q_a = [s_a, \mathbf{a}] \quad q_b = [s_b, \mathbf{b}]$$

with $\mathbf{a} = [x_a \mathbf{i} + y_a \mathbf{j} + z_a \mathbf{k}]$
similar for \mathbf{b}

$$q_a + q_b = [s_a + s_b, \mathbf{a} + \mathbf{b}]$$

$$q_a - q_b = [s_a - s_b, \mathbf{a} - \mathbf{b}]$$

but keep in mind:

$$q_a + q_b = (s_a + s_b) + (x_a + x_b)\mathbf{i} + (y_a + y_b)\mathbf{j} + (z_a + z_b)\mathbf{k}$$

$$q_a - q_b = (s_a - s_b) + (x_a - x_b)\mathbf{i} + (y_a - y_b)\mathbf{j} + (z_a - z_b)\mathbf{k}$$

Substitute to swap between notations.

Product of Quaternions

ARLAB

Multiplying two quaternions takes more time, but the math is straight forward:

Use the quaternions q_a and q_b as starting point:

$$q_a = [s_a, \mathbf{a}] \quad q_b = [s_b, \mathbf{b}]$$

multiplying:

$$\begin{aligned} q_a q_b &= [s_a, \mathbf{a}] [s_b, \mathbf{b}] \\ &= (s_a + x_a i + y_a j + z_a k)(s_b + x_b i + y_b j + z_b k) \\ &= (s_a s_b - x_a x_b - y_a y_b - z_a z_b) \\ &\quad + (s_a x_b + s_b x_a + y_a z_b - y_b z_a)i \\ &\quad + (s_a y_b + s_b y_a + z_a x_b - z_b x_a)j \\ &\quad + (s_a z_b + s_b z_a + x_a y_b - x_b y_a)k \end{aligned}$$

keep in mind

$$i^2 = -1$$

$$i^2 + j^2 + k^2 = ijk = -1$$

$$\begin{array}{lll} ij = k & jk = i & ki = j \\ ji = -k & kj = -i & ik = -j \end{array}$$

Product of Quaternions

ARLAB

Multiplying two quaternions takes more time, but the math is straight forward:

Use the quaternions q_a and q_b as starting point:

$$q_a = [s_a, \mathbf{a}] \quad q_b = [s_b, \mathbf{b}]$$

multiplying:

$$\begin{aligned} q_a q_b &= [s_a, \mathbf{a}] [s_b, \mathbf{b}] \\ &= (s_a + x_a i + y_a j + z_a k)(s_b + x_b i + y_b j + z_b k) \\ &= (s_a s_b - x_a x_b - y_a y_b - z_a z_b) \\ &\quad + (s_a x_b + s_b x_a + y_a z_b - y_b z_a)i \\ &\quad + (s_a y_b + s_b y_a + z_a x_b - z_b x_a)j \\ &\quad + (s_a z_b + s_b z_a + x_a y_b - x_b y_a)k \end{aligned}$$

note, change the complex number:

$$y_a j \cdot z_b k = (y_a z_b) i \rightarrow jk = i$$

keep in mind

$$i^2 = -1$$

$$i^2 + j^2 + k^2 = ijk = -1$$

$$\begin{array}{lll} ij = k & jk = i & ki = j \\ ji = -k & kj = -i & ik = -j \end{array}$$

Product of Quaternions



Now let's replace the imaginary number by ordered pairs

$$i = [0, \mathbf{i}] \ j = [0, \mathbf{j}] \ k = [0, \mathbf{k}] \text{ and } [1, \mathbf{0}] = 1$$

Pure quaternion

Real quaternion

gives

$$\begin{aligned} [s_a, \mathbf{a}][s_b, \mathbf{b}] &= (s_a s_b - x_a x_b - y_a y_b - z_a z_b)[1, \mathbf{0}] \\ &\quad + (s_a x_b + s_b x_a + y_a z_b - y_b z_a)[0, \mathbf{i}] \\ &\quad + (s_a y_b + s_b y_a + z_a x_b - z_b x_a)[0, \mathbf{j}] \\ &\quad + (s_a z_b + s_b z_a + x_a y_b - x_b y_a)[0, \mathbf{k}] \end{aligned}$$

rewriting as ordered pairs gives

$$\begin{aligned} [s_a, \mathbf{a}][s_b, \mathbf{b}] &= [s_a s_b - x_a x_b - y_a y_b - z_a z_b, \mathbf{0}] \\ &\quad + [0, (s_a x_b + s_b x_a + y_a z_b - y_b z_a) \mathbf{i}] \\ &\quad + [0, (s_a y_b + s_b y_a + z_a x_b - z_b x_a) \mathbf{j}] \\ &\quad + [0, (s_a z_b + s_b z_a + x_a y_b - x_b y_a) \mathbf{k}] \end{aligned}$$

Product of Quaternions

ARLAB

We can extract the common components and further rewrite as:

$$\begin{aligned}[s_a, \mathbf{a}][s_b, \mathbf{b}] &= [s_a s_b - x_a x_b - y_a y_b - z_a z_b, \mathbf{0}] \\ &\quad + [0, s_a(x_b \mathbf{i} + y_b \mathbf{j} + z_b \mathbf{k}) + s_b(x_a \mathbf{i} + y_a \mathbf{j} + z_a \mathbf{k}) \\ &\quad + (y_z z_b - y_b z_a) \mathbf{i} + (z_a x_b - z_b x_a) \mathbf{j} + (x_a y_b - y_b x_a) \mathbf{k}]\end{aligned}$$

Product of Quaternions

ARLAB

We can extract the common components and further rewrite as:

$$[s_a, \mathbf{a}][s_b, \mathbf{b}] = [s_a s_b - x_a x_b - y_a y_b - z_a z_b, \mathbf{0}] + [0, s_a(x_b \mathbf{i} + y_b \mathbf{j} + z_b \mathbf{k}) + s_b(x_a \mathbf{i} + y_a \mathbf{j} + z_a \mathbf{k}) + (y_z z_b - y_b z_a) \mathbf{i} + (z_a x_b - z_b x_z) \mathbf{j} + (x_a y_b - y_b x_a) \mathbf{k}]$$

Real quaternion

Pure quaternion

We combine the two ordered pairs into a single ordered pair, because one component is always 0.

$$[s_a, \mathbf{a}][s_b, \mathbf{b}] = [s_a s_b - x_a x_b - y_a y_b - z_a z_b, s_a(x_b \mathbf{i} + y_b \mathbf{j} + z_b \mathbf{k}) + s_b(x_a \mathbf{i} + y_a \mathbf{j} + z_a \mathbf{k}) + (y_z z_b - y_b z_a) \mathbf{i} + (z_a x_b - z_b x_z) \mathbf{j} + (x_a y_b - y_b x_a) \mathbf{k}]$$

Product of Quaternions

ARLAB

$$[s_a, \mathbf{a}][s_b, \mathbf{b}] = [s_a s_b - x_a x_b - y_a y_b - z_a z_b, \\ s_a(x_b \mathbf{i} + y_b \mathbf{j} + z_b \mathbf{k}) + s_b(x_a \mathbf{i} + y_a \mathbf{j} + z_a \mathbf{k}) \\ + (y_z z_b - y_b z_a) \mathbf{i} + (z_a x_b - z_b x_a) \mathbf{j} + (x_a y_b - y_b x_a) \mathbf{k}]$$

Product of Quaternions

ARLAB

$$[s_a, \mathbf{a}][s_b, \mathbf{b}] = [s_a s_b - x_a x_b - y_a y_b - z_a z_b, \\ s_a(x_b \mathbf{i} + y_b \mathbf{j} + z_b \mathbf{k}) + s_b(x_a \mathbf{i} + y_a \mathbf{j} + z_a \mathbf{k}) \\ + (y_z z_b - y_b z_a) \mathbf{i} + (z_a x_b - z_b x_a) \mathbf{j} + (x_a y_b - y_b x_a) \mathbf{k}]$$

Dot product

pure quaternion

Cross product

with:

$$\mathbf{a} = x_a \mathbf{i} + y_a \mathbf{j} + z_a \mathbf{k}$$

$$\mathbf{b} = x_b \mathbf{i} + y_b \mathbf{j} + z_b \mathbf{k}$$

$$\mathbf{a} \cdot \mathbf{b} = x_a x_b \mathbf{i}^2 + y_a y_b \mathbf{j}^2 + z_a z_b \mathbf{k}^2$$

$$\mathbf{a} \times \mathbf{b} = (y_a z_b - y_b z_a) \mathbf{i} (z_a x_b - x_b z_a) \mathbf{j} + (x_a y_b - x_b y_a) \mathbf{k}$$

$$i^2 = j^2 = k^2 = -1$$

Product of Quaternions

ARLAB

Substituting:

$$[s_a, \mathbf{a}][s_b, \mathbf{b}] = [s_a s_b - \mathbf{a} \cdot \mathbf{b}, s_a \mathbf{b} + s_b \mathbf{a} + \mathbf{a} \times \mathbf{b}]$$

Results in an equation for the product of two quaternions.

Real and Pure Quaternions

ARLAB

A Real Quaternion is a quaternion with a vector term of 0:

$$q = [s, \mathbf{0}]$$

the Pure Quaternion as a quaternion that has a zero scalar term

$$q = [0, \mathbf{v}]$$

addition of both:

$$\begin{aligned} q &= [s, \mathbf{v}] \\ &= [s, \mathbf{0}] + [0, \mathbf{v}] \end{aligned}$$

Unit and Norm



Unit Quaternion

Given a vector \mathbf{v} , we can express this vector with its scalar magnitude and its direction:

$$\mathbf{v} = v\hat{\mathbf{v}} \quad \text{where } v = |\mathbf{v}| \quad \text{and } |\hat{\mathbf{v}}| = 1$$

Thus, we obtain a quaternion with length of the complex vector = 1.

Quaternion Norm

$$q = [s, \mathbf{v}]$$

$$|q| = \sqrt{s^2 + \mathbf{v}^2}$$

Normalization

ARLAB

We can also normalize a quaternion:

$$q' = \frac{q}{|a|} = \frac{q}{\sqrt{s^2 + \mathbf{v}^2}}$$

Example:

$$q = [1, 4\mathbf{i} + 4\mathbf{j} - 4\mathbf{k}]$$

Calculate the length:

$$|q| = \sqrt{1^2 + 4^2 + 4^2 + (-4)^2} = \sqrt{49} = 7$$

Divide the quaternion by the length (norm)

$$q' = \frac{q}{|q|} = \frac{1 + 4\mathbf{i} + 4\mathbf{j} - 4\mathbf{k}}{7} = \frac{1}{7} + \frac{4}{7}\mathbf{i} + \frac{4}{7}\mathbf{j} - \frac{4}{7}\mathbf{k}$$

Quaternion Conjugate and Inverse

ARLAB

The quaternion conjugate can be computed by negating the imaginary part of the quaternion:

$$q = [s, \mathbf{v}]$$

$$q^* = [s, -\mathbf{v}]$$

The inverse of a quaternion can be computed as:

$$q^{-1} = \frac{q^*}{|q|^2}$$

Quaternion Dot Product

ARLAB

Similar to the vector dot product, we can also define a quaternion dot product by multiplying the scalar parts:

$$q_a = [s_a, x_a\mathbf{i} + y_a\mathbf{j} + z_a\mathbf{k}]$$

$$q_b = [s_b, x_b\mathbf{i} + y_b\mathbf{j} + z_b\mathbf{k}]$$

$$q_a \cdot q_b = s_a s_b + x_a x_b + y_a y_b + z_a z_b$$

Which also represents the angle between two quaternions:

$$\cos \theta = \frac{s_a s_b + x_a x_b + y_a y_b + z_a z_b}{|q_1||q_2|}$$

or with a unit-quaternion, length = 1 (similar to normalized vectors)

$$\cos \theta = s_a s_b + x_a x_b + y_a y_b + z_a z_b$$



Rotation with Quaternions

Rotation with Quaternions


ARLAB

Remember the Rotor we defined for complex numbers:

$$q = \cos \theta + i \sin \theta$$

Knowing the similarity to complex numbers, we can express a general Rotor for 3D space such as:

$$q = [\cos \theta, \sin \theta \mathbf{v}]$$



Let's test

Let's consider a vector p that we want to rotate,
as pure quaternion:


$$p = [0, 2\mathbf{i}]$$

we want to rotate $\theta = 45^\circ$ around \mathbf{k}

$$q = [\cos \theta, \sin \theta \mathbf{k}]$$

(I removed the components \mathbf{i}, \mathbf{j})

$$q = \left[\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2} \mathbf{k} \right]$$




(Quaternions work in 4D,
drawing 3D is actually not correct.)

Let's test

and rotate:

$$\begin{aligned} p' &= qp \\ &= \left[\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}\mathbf{k} \right] [0, 2\mathbf{i}] \\ &= \left[0, 2\frac{\sqrt{2}}{2}\mathbf{i} + 2\frac{\sqrt{2}}{2}\mathbf{k} \times \mathbf{i} \right] \\ &= [0, \sqrt{2}\mathbf{i} + \sqrt{2}\mathbf{j}] \end{aligned}$$



(Quaternions work in 4D,
drawing 3D is actually not correct.)

Let's test

and rotate:


$$p' = qp$$

$$= \left[\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2} \mathbf{k} \right] [0, 2\mathbf{i}]$$

Moving the scalar term to the front.

$$= \left[0, 2 \frac{\sqrt{2}}{2} \mathbf{i} + 2 \frac{\sqrt{2}}{2} \mathbf{k} \times \mathbf{i} \right]$$

$$= [0, \sqrt{2}\mathbf{i} + \sqrt{2}\mathbf{j}]$$



Multiplication


$$[s_a, \mathbf{a}][s_b, \mathbf{b}] = [s_a s_b - \mathbf{a} \cdot \mathbf{b}, s_a \mathbf{b} + s_b \mathbf{a} + \mathbf{a} \times \mathbf{b}]$$

(Quaternions work in 4D,
drawing 3D is actually not correct.)

Let's test

and rotate:

$$\begin{aligned} p' &= qp \\ &= \left[\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}\mathbf{k} \right] [0, 2\mathbf{i}] \\ &= \left[0, 2\frac{\sqrt{2}}{2}\mathbf{i} + 2\frac{\sqrt{2}}{2}\mathbf{k} \times \mathbf{i} \right] \\ &= [0, \sqrt{2}\mathbf{i} + \sqrt{2}\mathbf{j}] \end{aligned}$$



We should maintain the magnitude

$$|p'| = \sqrt{\sqrt{2}^2 + \sqrt{2}^2} = 2$$

(Quaternions work in 4D,
drawing 3D is actually not correct.)

Arbitrary Rotation Vector

ARLAB

Let's consider a vector p that we want to rotate, an pure quaternion:

$$p = [0, 2\mathbf{i}]$$


$$\mathbf{p} = 2\mathbf{i}$$

we want to rotate $\theta = 45^\circ$ around $\hat{\mathbf{v}}$

$$\hat{\mathbf{v}} = \frac{\sqrt{2}}{2}\mathbf{i} + \frac{\sqrt{2}}{2}\mathbf{k}$$

with

$$q = [\cos \theta, \sin \theta \hat{\mathbf{v}}]$$



Arbitrary Rotation Vector


ARLAB

Calculate

$$\begin{aligned} p' &= qp \\ &= [\cos \theta, \sin \theta \hat{\mathbf{v}}][0, \mathbf{p}] \\ &= [-\sin \theta \hat{\mathbf{v}} \cdot \mathbf{p}, \cos \theta \mathbf{p} + \sin \theta \hat{\mathbf{v}} \times \mathbf{p}] \end{aligned}$$

Multiplication

$$[s_a, \mathbf{a}][s_b, \mathbf{b}] = [s_a s_b - \mathbf{a} \cdot \mathbf{b}, s_a \mathbf{b} + s_b \mathbf{a} + \mathbf{a} \times \mathbf{b}]$$



Substituting θ and \mathbf{p} and $\hat{\mathbf{v}}$

$$\begin{aligned} p' &= \left[-\frac{\sqrt{2}}{2} \left(\frac{\sqrt{2}}{2} \mathbf{i} + \frac{\sqrt{2}}{2} \mathbf{k} \right) \cdot (2\mathbf{i}), \frac{\sqrt{2}}{2} 2\mathbf{i} + \frac{\sqrt{2}}{2} \left(\frac{\sqrt{2}}{2} \mathbf{i} + \frac{\sqrt{2}}{2} \mathbf{k} \right) \times 2\mathbf{i} \right] \\ &= [-1, \sqrt{2}\mathbf{i} + \mathbf{j}] \end{aligned}$$

Arbitrary Rotation Vector

ARLAB


This does not look right to me.

$$p' = [-1, \sqrt{2}\mathbf{i} + \mathbf{j}]$$

The length of p' is also $\sqrt{3}$ instead of 2

now?

we have to get it back to 2.



Arbitrary Rotation Vector

ARLAB

We post-multiply the solution with the inverse of q

$$q = \left[\cos \theta, \sin \theta \left(\frac{\sqrt{2}}{2} \mathbf{i} + \frac{\sqrt{2}}{2} \mathbf{k} \right) \right]$$


Let's compute the inverse

$$q^{-1} = \left[\cos \theta, -\sin \theta \left(\frac{\sqrt{2}}{2} \mathbf{i} + \frac{\sqrt{2}}{2} \mathbf{k} \right) \right]$$

for 45°

$$q^{-1} = \left[\frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2} \left(\frac{\sqrt{2}}{2} \mathbf{i} + \frac{\sqrt{2}}{2} \mathbf{k} \right) \right]$$

$$= \frac{1}{2} [\sqrt{2}, -\mathbf{i} - \mathbf{k}]$$



Arbitrary Rotation Vector

ARLAB

Now we compute


$$qp = [-1, \sqrt{2}\mathbf{i} + \mathbf{j}]$$

$$qpq^{-1} = [-1, \sqrt{2}\mathbf{i} + \mathbf{j}] \frac{1}{2} [\sqrt{2}, -\mathbf{i} - \mathbf{k}]$$

= ...

$$= [0, \mathbf{i} + \sqrt{2}\mathbf{j} + \mathbf{k}]$$

The length is 2 again




Quaternions and Rotations

ARLAB

A quaternion can represent a rotation by an angle θ around a unit axis \mathbf{a} :

$$q = \left[\cos \frac{\theta}{2}, \sin \frac{\theta}{2} \hat{\mathbf{v}} \right] = \left[\cos \frac{\theta}{2}, \underline{\sin \frac{\theta}{2} i + \sin \frac{\theta}{2} j + \sin \frac{\theta}{2} k} \right]$$

The “projections” of a to x , y , and z



- the three complex numbers represent a vector in 3d space. A quaternion in \mathbb{R}^4 is a vector in \mathbb{R}^3 , if the scalar value is 0.
- the scalar value theta represents the rotation about this vector a.

Thank you!

Questions

Rafael Radkowski, Ph.D.
Iowa State University
Virtual Reality Applications Center
1620 Howe Hall
Ames, Iowa 50011, USA
+1 515.294.5580
+1 515.294.5530(fax)
rafael@iastate.edu
<http://arlabs.me.iastate.edu>



IOWA STATE UNIVERSITY
OF SCIENCE AND TECHNOLOGY