

Zero-Knowledge Proofs for Committed Symmetric Boolean Functions

SAN LING, KHOA NGUYEN, DUONG HIEU PHAN, HANH TANG AND HUAXIONG WANG

NANYANG TECHNOLOGICAL UNIVERSITY, SINGAPORE

LTCI, TELECOM PARIS, INSTITUT POLYTECHNIQUE DE PARIS, FRANCE

Overview

- Introduction
- Backgrounds
- Technique for Evaluating Symmetric Boolean Functions in Zero-Knowledge
- Zero-Knowledge Proof for Symmetric Boolean Functions



Introduction

Previous Works and Motivation

- **Previous works.** Most existing works focus on ZKP for correct evaluation of **private input** (encrypted or committed) from a **publicly known function**.
- **Our setting.** ZKP for correct evaluation of **private symmetric Boolean functions** on **private inputs**.
- **Possible applications.** Policy-based anonymous authentication, privacy-preserving access controls for encrypted databases, accountable function evaluations, ...

Symmetric Boolean Functions

- An n -ary symmetric Boolean function $f: \{0,1\}^n \rightarrow \{0,1\}$ is represented by

$$\mathbf{v} = \mathbf{v}(f) = (v_0, v_1, \dots, v_n) \in \{0,1\}^{n+1}.$$

- On input $\mathbf{x} \in \{0,1\}^n$, $f(\mathbf{x})$ returns v_w where $w = \text{weight}(\mathbf{x})$.

- 2^{n+1} different symmetric Boolean functions.

- Examples.

1. **Threshold functions:** $T_k(\mathbf{x}) = 1 \Leftrightarrow \text{weight}(\mathbf{x}) \geq k$.
2. **Parity functions:** $\text{PAR}(\mathbf{x}) = 1 \Leftrightarrow \text{weight}(\mathbf{x})$ is odd.
3. **Sorting functions:** $\text{SORT}(\mathbf{x}) = (T_1(\mathbf{x}), T_2(\mathbf{x}), \dots, T_n(\mathbf{x}))$.

Problem Statement

- Given a public bit b and commitments to $\mathbf{x} \in \{0,1\}^n$ and f as follows:

$$\mathbf{c}_x = \text{Com}_{ck}(\mathbf{x}; \rho_x) \text{ and } \mathbf{c}_f = \text{Com}_{ck}(\mathbf{v}(f); \rho_f).$$

- Construct ZK proof for knowledge of \mathbf{x} and $\mathbf{v}(f)$ such that $f(\mathbf{x}) = b$.
- **Common inputs.** $ck, \mathbf{c}_x, \mathbf{c}_f$ and b .
- **Prover's inputs.** $\mathbf{x}, \mathbf{v}(f)$, commitment randomness ρ_x, ρ_f .
- **Relation**

$$R_{\text{sym}} = \{(ck, \mathbf{c}_x, \mathbf{c}_f, b); \mathbf{x}, \mathbf{v}(f), \rho_x, \rho_f : \mathbf{c}_x = \text{Com}_{ck}(\mathbf{x}; \rho_x), \mathbf{c}_f = \text{Com}_{ck}(\mathbf{v}(f); \rho_f), f(\mathbf{x}) = b\}.$$



Backgrounds

LPN-Based Commitments [JKPT12]

- n : the bit-length of message.
- Commitment key $(A_{1,x}, A_2) \in \{0,1\}^{k \times n} \times \{0,1\}^{k \times s}$.
- To commit n -bit message x , compute

$$c_x = A_{1,x} \cdot x \oplus A_2 \cdot s_x \oplus e_x$$

where $s_x \stackrel{\$}{\leftarrow} \{0,1\}^s$ and e_x is sampled from appropriate Bernoulli distribution.

- Similarly, to commit $(n + 1)$ -bit vector $v = v(f)$, use commitment key $(A_{1,f}, A_2)$ and compute

$$c_f = A_{1,f} \cdot v \oplus A_2 \cdot s_f \oplus e_f.$$

Stern-Like Σ -Protocol [LLMNW16]

- Stern-like Σ -protocol aims to show the knowledge of secret vector $\mathbf{w} = (\mathbf{w}_1 \parallel \mathbf{w}_2)$ satisfying

$$\mathbf{M}_1 \cdot \mathbf{w}_1 \oplus \mathbf{M}_2 \cdot \mathbf{w}_2 = \mathbf{u} \quad \text{and} \quad \mathbf{w}_1 \in \text{VALID}$$

for some public matrices $\mathbf{M}_1, \mathbf{M}_2$, public vector \mathbf{u} and set VALID containing \mathbf{w}_1 .

- **Relation**

$$R_{\text{abstract}} = \{(\mathbf{M}_1, \mathbf{M}_2, \mathbf{u}); \mathbf{w}_1, \mathbf{w}_2 : \mathbf{M}_1 \cdot \mathbf{w}_1 \oplus \mathbf{M}_2 \cdot \mathbf{w}_2 = \mathbf{u} \wedge \mathbf{w}_1 \in \text{VALID}\}$$

- Stern-like ZK Proof of Knowledge is constructable if there exists a set of permutations S satisfying

$$\left\{ \begin{array}{l} \forall \varphi \in S: \mathbf{w} \in \text{VALID} \Leftrightarrow \varphi(\mathbf{w}) \in \text{VALID} \\ \text{If } \mathbf{w} \in \text{VALID} \text{ and } \varphi \text{ is uniform in } S, \text{ then } \varphi(\mathbf{w}) \text{ is uniform in VALID.} \end{array} \right.$$


- **Purpose.** Reduce R_{sym} to R_{abstract} .

Recall. Stern-Like Technique for Valid Openings of LPN-Based Commitments

- **Recall.** $\mathbf{c}_x = \mathbf{A}_{1,x} \cdot \mathbf{x} \oplus \mathbf{A}_2 \cdot \mathbf{s}_x \oplus \mathbf{e}_x$ where $\mathbf{s}_x \xleftarrow{\$} \{0,1\}^s$ and $\mathbf{e}_x \in \{0,1\}^\kappa$ is sampled from appropriate Bernoulli distribution.
- With overwhelming probability, $\text{weight}(\mathbf{e}_x) \leq t$ for some t .
- Extend \mathbf{e}_x to $(\mathbf{e}_x \parallel \mathbf{e}'_x) \in \{0,1\}^{\kappa+t}$, where $\mathbf{e}'_x \in \{0,1\}^t$ such that $\text{weight}(\mathbf{e}_x \parallel \mathbf{e}'_x) = t$.
- **Fact.** $\text{weight}(\mathbf{e}_x) \leq t \Leftrightarrow \exists \mathbf{e}'_x \in \{0,1\}^t$ s.t $\text{weight}(\mathbf{e}_x \parallel \mathbf{e}'_x) = t$.

$$\mathbf{c}_x = [\mathbf{I}_\kappa | \mathbf{0}^{\kappa \times t}] \cdot (\mathbf{e}_x \parallel \mathbf{e}'_x) \oplus [\mathbf{A}_{1,x} | \mathbf{A}_2] \cdot (\mathbf{x} \parallel \mathbf{s}_x) \text{ and } \text{weight}(\mathbf{e}_x \parallel \mathbf{e}'_x) = t.$$

- Define $\text{VALID}_{\text{LPN}} = \{\mathbf{w} \in \{0,1\}^{\kappa+t} : \text{weight}(\mathbf{w}) = t\}$.
- Reducible to R_{abstract} by defining set of permutations to be symmetric group over $\kappa + t$ elements.



Technique for Evaluating Symmetric Boolean Functions in ZK

ZKP for Symmetric Boolean Functions

- Prover needs to convince verifier that

$$\begin{cases} \mathbf{c}_x = \mathbf{A}_{1,x} \cdot \mathbf{x} \oplus \mathbf{A}_2 \cdot \mathbf{s}_x \oplus \mathbf{e}_x, \\ \mathbf{c}_f = \mathbf{A}_{1,f} \cdot \mathbf{v} \oplus \mathbf{A}_2 \cdot \mathbf{s}_f \oplus \mathbf{e}_f, \\ f(\mathbf{x}) = b \end{cases}$$

where $\mathbf{v} = \mathbf{v}(f)$.

- ZKP techniques for valid openings of commitments are available.
- How to simultaneously show that $f(\mathbf{x}) = b$?

Technique for Handling $f(\mathbf{x}) = b$

- **Recall.** $\mathbf{v} = \mathbf{v}(f) = (v_0, \dots, v_n)$ as f is a symmetric Boolean function.
- Define $w = \text{weight}(\mathbf{x})$.

- Hence,

$$f(\mathbf{x}) = b \Leftrightarrow v_w = b.$$

- To extract v_w from \mathbf{v} , define $\mathbf{y} = U(w) = (y_0, \dots, y_n) = (0, \dots, 0, 1, 0, \dots, 0)$ the w^{th} basis vector.
- Hence,

$$f(\mathbf{x}) = b \Leftrightarrow v_w = b \Leftrightarrow \langle \mathbf{v}, \mathbf{y} \rangle = b.$$

1. How to construct \mathbf{y} ?
2. How to show that $\langle \mathbf{v}, \mathbf{y} \rangle = b$?

Constructing $\mathbf{y} = U(w)$

- **Recall.** $\mathbf{x} \in \{0,1\}^n$, $w = \text{weight}(\mathbf{x})$, $\mathbf{y} = U(w) \in \{0,1\}^{n+1}$.
- **Observation.** Number of 0's in the right of 1 in \mathbf{y} is equal to $n - w$.
- **Example.** $\mathbf{x} = (1,0,1,0,0,1,1) \in \{0,1\}^7 \Rightarrow \mathbf{y} = (0,0,0,0,1,0,0,0) \in \{0,1\}^8$.
- To show that \mathbf{y} is well-formed, construct $\mathbf{z} \in \{0,1\}^{n+1}$ by inverting all 0's in the right of 1 in \mathbf{y} .
- **Example.** $\mathbf{y} = (0,0,0,0,1,0,0,0) \in \{0,1\}^8 \Rightarrow \mathbf{z} = (0,0,0,0,1,1,1,1) \in \{0,1\}^8$.
- **Facts.** By setting $\mathbf{z} = (z_0, \dots, z_n) \in \{0,1\}^8$, then

$$\mathbf{y} = U(w) \Leftrightarrow \begin{cases} z_0 = y_0, \\ z_i = y_i \oplus z_{i-1} \quad \forall i \in \{1, \dots, n\}, \\ \text{weight}(\mathbf{y}) = 1 \\ \text{weight}(\mathbf{z}) + \text{weight}(\mathbf{x}) = n + 1. \end{cases}$$

Showing $\langle \mathbf{v}, \mathbf{y} \rangle = b$

- **Recall.** $\mathbf{y} = (y_0, y_1, \dots, y_n) = U(j) \in \{0,1\}^{n+1}$ and $\mathbf{v} = \mathbf{v}(f) = (v_0, v_1, \dots, v_n) \in \{0,1\}^{n+1}$.
- $\text{ext}(\mathbf{y}) = (\mathbf{y}_0 \| \mathbf{y}_1 \| \dots \| \mathbf{y}_n) = (y_0, 0, y_1, 0, \dots, y_n, 0) \in \{0,1\}^{2n+2}$ where $\mathbf{y}_i = (y_i, 0)$.
- **Observation.** $\text{weight}(\mathbf{y}_j) = 1 \bmod 2$ and $\text{weight}(\mathbf{y}_i) = 0 \bmod 2 \ \forall i \neq j$.
- $\text{enc}(\mathbf{v}) = (\mathbf{v}_0 \| \mathbf{v}_1 \| \dots \| \mathbf{v}_n) = (\bar{v}_0, v_0, \bar{v}_1, v_1, \dots, \bar{v}_n, v_n) \in \{0,1\}^{2n+2}$ where $\mathbf{v}_i = (\bar{v}_i, v_i)$.
- **Observation.** $\text{weight}(\mathbf{v}_i) = 1 \bmod 2 \ \forall i$.
- Define $\mathbf{b} = (\mathbf{b}_0 \| \mathbf{b}_1 \| \dots \| \mathbf{b}_n) = \text{ext}(\mathbf{y}) \oplus \text{enc}(\mathbf{v})$ where $\mathbf{b}_i = (b_i \oplus \bar{v}_i, v_i)$ s.t

$$\begin{cases} \text{weight}(\mathbf{b}_j) = 0 \bmod 2, \\ \text{weight}(\mathbf{b}_i) = 1 \bmod 2 \ \forall i \neq j. \end{cases}$$

Showing $\langle \mathbf{v}, \mathbf{y} \rangle = b$ (continued)

- Define $\text{good}(b) = \{(\mathbf{b}'_0 \| \mathbf{b}'_1 \| \dots \| \mathbf{b}'_n) : \text{exists unique } j \text{ s.t. } \mathbf{b}'_j = (b, b) \text{ and } \text{weight}(\mathbf{b}'_i) = 1 \ \forall i \neq j\}$.

$$\text{ext}(\mathbf{y}) \oplus \text{enc}(\mathbf{v}) \in \text{good}(b) \Leftrightarrow \langle \mathbf{v}, \mathbf{y} \rangle = b.$$

- Now, assume that $\text{weight}(\mathbf{y})$ is unknown and $\mathbf{b} = \text{ext}(\mathbf{y}) \oplus \text{enc}(\mathbf{v}) \in \text{good}(b)$.
- $\mathbf{b} = (\mathbf{b}_0 \| \mathbf{b}_1 \| \dots \| \mathbf{b}_n) \in \text{good}(b) \Rightarrow \mathbf{b}_j = (b, b)$ for some unique j and $\text{weight}(\mathbf{b}_i) = 1 \ \forall i \neq j$.
- $\text{enc}(\mathbf{v}) = (\mathbf{v}_0 \| \mathbf{v}_1 \| \dots \| \mathbf{v}_n) \Rightarrow \text{weight}(\mathbf{v}_i) = 1 \bmod 2 \ \forall i$.
- $\text{ext}(\mathbf{y}) = (\mathbf{y}_0 \| \mathbf{y}_1 \| \dots \| \mathbf{y}_n) = \mathbf{b} \oplus \text{enc}(\mathbf{v}) \Rightarrow \text{weight}(\mathbf{y}_j) = 1$ and $\text{weight}(\mathbf{y}_i) = 0 \bmod 2 \ \forall i \neq j$.

$$\left. \begin{array}{l} \mathbf{y}_j = (y_j, 0) \Rightarrow y_j = 1 \\ \forall i \neq j, \mathbf{y}_i = (y_i, 0) \Rightarrow y_i = 0 \end{array} \right\} \Rightarrow \mathbf{y} = (y_0, \dots, y_n) \text{ is a unit vector.}$$

- In summary.

$$\mathbf{b} = \text{ext}(\mathbf{y}) \oplus \text{enc}(\mathbf{v}) \in \text{good}(b) \Leftrightarrow \langle \mathbf{v}, \mathbf{y} \rangle = b \text{ and } \mathbf{y} \text{ is unit vector}$$

Putting pieces together

Theorem. $x \in \{0,1\}^n$, $v = v(f) = (v_0, v_1, \dots, v_n)$, $b \in \{0,1\}$, the following statements are equivalent:

- i. $f(x) = b$.
- ii. There exists $b_0, b_1, \dots, b_n \in \{0,1\}$ and $z = (z_0, z_1, \dots, z_n) \in \{0,1\}^{n+1}$ satisfying

$$\left\{ \begin{array}{l} b_0 \oplus v_0 \oplus z_0 = 1, \\ b_i \oplus v_i \oplus z_i \oplus z_{i-1} = 1 \quad \forall i \in \{1, \dots, n\}, \\ \text{weight}(x) + \text{weight}(z) = n + 1, \\ (b_0, v_0, \dots, b_n, v_n) \in \text{good}(b). \end{array} \right.$$



ZKP for Symmetric Boolean Functions

ZKP for Symmetric Boolean Functions

Prover shows that $\exists \mathbf{e}'_x \in \{0,1\}^t, \mathbf{e}'_f \in \{0,1\}^t, b_0, b_1, \dots, b_n \in \{0,1\}, \mathbf{z} = (z_0, z_1, \dots, z_n) \in \{0,1\}^{n+1}$ s.t

$$\left\{ \begin{array}{l} \mathbf{c}_x = \mathbf{A}_{1,x} \cdot \mathbf{x} \oplus \mathbf{A}_2 \cdot \mathbf{s}_x \oplus [\mathbf{I}_\kappa | \mathbf{0}^{\kappa \times t}] \cdot (\mathbf{e}_x \| \mathbf{e}'_x), \\ \mathbf{c}_f = \mathbf{A}_{1,f} \cdot \mathbf{v} \oplus \mathbf{A}_2 \cdot \mathbf{s}_f \oplus [\mathbf{I}_\kappa | \mathbf{0}^{\kappa \times t}] \cdot (\mathbf{e}_f \| \mathbf{e}'_f), \\ b_0 \oplus v_0 \oplus z_0 = 1, \\ b_i \oplus v_i \oplus z_i \oplus z_{i-1} = 1 \quad \forall i \in \{1, \dots, n\}, \\ \text{weight}(\mathbf{e}_x \| \mathbf{e}'_x) = \text{weight}(\mathbf{e}_f \| \mathbf{e}'_f) = t, \\ \text{weight}(\mathbf{x}) + \text{weight}(\mathbf{z}) = n + 1, \\ (b_0, v_0, \dots, b_n, v_n) \in \text{good}(b). \end{array} \right. \Leftrightarrow \left\{ \begin{array}{l} \mathbf{c}_x = \mathbf{A}_{1,x} \cdot \mathbf{x} \oplus \mathbf{A}_2 \cdot \mathbf{s}_x \oplus [\mathbf{I}_\kappa | \mathbf{0}^{\kappa \times t}] \cdot (\mathbf{e}_x \| \mathbf{e}'_x), \\ \mathbf{c}_f = \mathbf{A}_{1,f} \cdot \mathbf{v} \oplus \mathbf{A}_2 \cdot \mathbf{s}_f \oplus [\mathbf{I}_\kappa | \mathbf{0}^{\kappa \times t}] \cdot (\mathbf{e}_f \| \mathbf{e}'_f), \\ b_0 \oplus v_0 \oplus z_0 = 1, \\ b_i \oplus v_i \oplus z_i \oplus z_{i-1} = 1 \quad \forall i \in \{1, \dots, n\}, \end{array} \right.$$

and

$$\left\{ \begin{array}{l} \text{weight}(\mathbf{e}_x \| \mathbf{e}'_x) = \text{weight}(\mathbf{e}_f \| \mathbf{e}'_f) = t, \\ \text{weight}(\mathbf{x}) + \text{weight}(\mathbf{z}) = n + 1, \\ (b_0, v_0, \dots, b_n, v_n) \in \text{good}(b). \end{array} \right.$$

ZKP for Symmetric Boolean Functions

Prover shows that $\exists \mathbf{e}'_x \in \{0,1\}^t, \mathbf{e}'_f \in \{0,1\}^t, b_0, b_1, \dots, b_n \in \{0,1\}, \mathbf{z} = (z_0, z_1, \dots, z_n) \in \{0,1\}^{n+1}$ s.t

$$\begin{cases} \mathbf{c}_x = \mathbf{A}_{1,x} \cdot \mathbf{x} \oplus \mathbf{A}_2 \cdot \mathbf{s}_x \oplus [\mathbf{I}_\kappa | \mathbf{0}^{\kappa \times t}] \cdot (\mathbf{e}_x \| \mathbf{e}'_x), \\ \mathbf{c}_f = \mathbf{A}_{1,f} \cdot \mathbf{v} \oplus \mathbf{A}_2 \cdot \mathbf{s}_f \oplus [\mathbf{I}_\kappa | \mathbf{0}^{\kappa \times t}] \cdot (\mathbf{e}_f \| \mathbf{e}'_f), \\ b_0 \oplus v_0 \oplus z_0 = 1, \\ b_i \oplus v_i \oplus z_i \oplus z_{i-1} = 1 \quad \forall i \in \{1, \dots, n\}, \end{cases} \Rightarrow$$

and

$$\begin{cases} \text{weight}(\mathbf{e}_x \| \mathbf{e}'_x) = \text{weight}(\mathbf{e}_f \| \mathbf{e}'_f) = t, \\ \text{weight}(\mathbf{x}) + \text{weight}(\mathbf{z}) = n + 1, \\ (b_0, v_0, \dots, b_n, v_n) \in \text{good}(b). \end{cases}$$

$$\text{VALID}_{\text{SYM}} = B(\kappa + t, t) \| B(\kappa + t, t) \| B(2n + 1, n + 1) \| \text{good}(b)$$

Possible to construct set of permutations S_{SYM}

Secret $\mathbf{w}_1 = (\mathbf{e}_x \| \mathbf{e}'_x \| \mathbf{e}_f \| \mathbf{e}'_f \| \mathbf{x} \| \mathbf{z} \| \mathbf{b}) \in \text{VALID}_{\text{SYM}},$


Secret $\mathbf{w}_2 = (\mathbf{s}_x \| \mathbf{s}_f),$

By linear algebra, define public $\mathbf{M}_1, \mathbf{M}_2, \mathbf{u}$ satisfying

$$\mathbf{M}_1 \cdot \mathbf{w}_1 \oplus \mathbf{M}_2 \cdot \mathbf{w}_2 = \mathbf{u}$$

where $\mathbf{u} = (\mathbf{c}_x \| \mathbf{c}_f \| \mathbf{1}^{n+1}).$

\Rightarrow Reducible to $R_{\text{abstract}}.$



Thank you!
Q&A

References

[JKPT12] A. Jain, S. Krenn, K. Pietrzak, and A. Tentes. Commitments and efficient zero-knowledge proofs from learning parity with noise. In *ASIACRYPT 2012*, volume 7658 of *LNCS*, pages 663–680. Springer, 2012.

[LLMNW16] B. Libert, S. Ling, F. Mouhartem, K. Nguyen, and H. Wang. Signature schemes with efficient protocols and dynamic group signatures from lattice assumptions. In *ASIACRYPT 2016*, volume 10032 of *LNCS*, pages 373–403, 2016.