

Quantum measurements on relativistic quantum field theory

Rafael Córdoba López

Universidad de los Andes

March 1, 2021

Motivation

1. Measurements on bounded regions in space time → compact region.

Motivation

1. Measurements on bounded regions in space time \rightarrow compact region.
2. Lorentz invariance (not preferred frames).

Motivation

1. Measurements on bounded regions in space time \rightarrow compact region.
2. Lorentz invariance (not preferred frames).
3. Causally disjoint sets \rightarrow no particular order of measure.

Motivation

1. Measurements on bounded regions in space time \rightarrow compact region.
2. Lorentz invariance (not preferred frames).
3. Causally disjoint sets \rightarrow no particular order of measure.
4. ¿Sorkin's protocol?




Figure: Sorkin's protocol[?]

Definitions and notation I

Let $x \in M, S \subseteq M$

Axioms/Assumptions on AQFT \mathcal{A}

Axioms/Assumptions on AQFT \mathcal{A}

1. Isotony: $N_1 \subset N_2 \subset M \implies \mathcal{A}(M; N_1) \subset \mathcal{A}(M; N_2)$

Axioms/Assumptions on AQFT \mathcal{A}

1. Isotony: $N_1 \subset N_2 \subset M \implies \mathcal{A}(M; N_1) \subset \mathcal{A}(M; N_2)$
2. Einstein causality: $N_1 \subset N_2^\perp \implies [\mathcal{A}(M; N_1), \mathcal{A}(M; N_2)] = 0$

Axioms/Assumptions on AQFT \mathcal{A}

1. Isotony: $N_1 \subset N_2 \subset M \implies \mathcal{A}(M; N_1) \subset \mathcal{A}(M; N_2)$
2. Einstein causality: $N_1 \subset N_2^\perp \implies [\mathcal{A}(M; N_1), \mathcal{A}(M; N_2)] = 0$
3. Compatibility: $N \subset M, N = ch(N)$

then, exist a (Compatibility map) $\alpha : \mathcal{A}(M|_N) \rightarrow \mathcal{A}(M)$ with

$$\alpha_{M;N}(\mathcal{A}(M|_N)) = \mathcal{A}(M; N)$$

$$M_3 \subset M_2 \subset M_1$$

$$\alpha_{M_1;M_2} \circ \alpha_{M_2;M_3} = \alpha_{M_1;M_3}$$

Axioms/Assumptions on AQFT \mathcal{A}

1. Isotony: $N_1 \subset N_2 \subset M \implies \mathcal{A}(M; N_1) \subset \mathcal{A}(M; N_2)$
2. Einstein causality: $N_1 \subset N_2^\perp \implies [\mathcal{A}(M; N_1), \mathcal{A}(M; N_2)] = 0$
3. Compatibility: $N \subset M, N = ch(N)$


then, exist a (Compatibility map) $\alpha : \mathcal{A}(M|_N) \rightarrow \mathcal{A}(M)$ with

$$\alpha_{M;N}(\mathcal{A}(M|_N)) = \mathcal{A}(M; N)$$

$$M_3 \subset M_2 \subset M_1$$

$$\alpha_{M_1;M_2} \circ \alpha_{M_2;M_3} = \alpha_{M_1;M_3}$$

4. Time-slice property (existence of dynamics)



Axioms/Assumptions on AQFT \mathcal{A}

1. Isotony: $N_1 \subset N_2 \subset M \implies \mathcal{A}(M; N_1) \subset \mathcal{A}(M; N_2)$
2. Einstein causality: $N_1 \subset N_2^\perp \implies [\mathcal{A}(M; N_1), \mathcal{A}(M; N_2)] = 0$
3. Compatibility: $N \subset M, N = ch(N)$


then, exist a (Compatibility map) $\alpha : \mathcal{A}(M|_N) \rightarrow \mathcal{A}(M)$ with

$$\alpha_{M;N}(\mathcal{A}(M|_N)) = \mathcal{A}(M; N)$$

$$M_3 \subset M_2 \subset M_1$$

$$\alpha_{M_1;M_2} \circ \alpha_{M_2;M_3} = \alpha_{M_1;M_3}$$

4. Time-slice property (existence of dynamics)



The measurement and the probe II




Figure: In and out algebras

On \mathbf{M} the algebra of observables $\mathcal{U}(\mathbf{M}) = \mathcal{A}(\mathbf{M}) \otimes \mathcal{B}(\mathbf{M})$ with
Compatibility map $\alpha_{M;N} \otimes \alpha_{M;N}$

The measurement and the probe III




Figure: Compatibility

$L \subset L'$ open the following diagram commutes:

$$\begin{array}{ccc}
 \mathcal{A}(L') \otimes \mathcal{B}(L') & \xrightarrow{\alpha_{L;L'} \otimes \beta_{L;L'}} & \mathcal{A}(L) \otimes \mathcal{B}(L) \\
 \chi_{L'} \downarrow & & \downarrow \chi_L \\
 \mathcal{C}(L') & \xrightarrow{\gamma_{L;L'}} & \mathcal{C}(L)
 \end{array}$$

The regions M^\pm are big enough to have Cauchy surfaces
time-slice ppty \rightarrow The morphisms α, β, γ and χ are isomorphism on M^\pm .

$$\tau^\pm : \mathcal{U}(M) \rightarrow \mathcal{C}(M)$$

$$\tau^\pm = \kappa^\pm \circ (\alpha^\pm \circ \beta^\pm)^{-1} = \gamma^\pm \circ \chi^\pm \circ (\alpha^\pm \circ \beta^\pm)^{-1}$$

τ identifies the uncoupled system with the coupled at early (-) or late (+) times. Which makes the **scattering morphism**

$$\Theta = (\tau^-)^{-1} \circ \tau^+$$

an automorphism.

Properties of the scattering morphism

Properties of the scattering morphism

- $K \subset K'$ with K' compact then the scattering morphism constructed in K' is the same as the one in K .

Measurement scheme III

Then,

$$A = \varepsilon_\sigma(B) := \eta_\sigma(\Theta(1 \otimes B))$$

where $\eta_\sigma(B)$ is called the **induced system observable**

$$\omega(A) = \omega(\eta_\sigma(\Theta(1 \otimes B))) = (\omega \otimes \sigma)(\Theta(1 \otimes B)) = \omega_\sigma(\tilde{B})$$

The system $(\mathcal{B}, \mathcal{C}, \chi(\tau^\pm), \sigma)$ is called the **measurment scheme** of the induced system observable.

Properties of the induced system observable

Let σ be a preparation state of \mathcal{B} , $A = \varepsilon_\sigma(B)$ is the unique solution to

$$\omega(A) = \omega_\sigma(\tilde{B})$$

Furthermore, ε_σ is a completely positive linear map and

- $\varepsilon_\sigma(1) = 1$
- $\varepsilon_\sigma(B^*) = \varepsilon_\sigma(B)^* \implies \varepsilon_\sigma(B^*) = \varepsilon_\sigma(B)$
- $\varepsilon_\sigma(B^*B) - \varepsilon_\sigma(B)^*\varepsilon_\sigma(B)$ is a positive operator

Properties of the induced system observable

Let σ be a preparation state of \mathcal{B} , $A = \varepsilon_\sigma(B)$ is the unique solution to

$$\omega(A) = \omega_\sigma(\tilde{B})$$

Furthermore, ε_σ is a completely positive linear map and

- $\varepsilon_\sigma(1) = 1$
- $\varepsilon_\sigma(B^*) = \varepsilon_\sigma(B)^* \implies \varepsilon_\sigma(B^*) = \varepsilon_\sigma(B)$
- $\varepsilon_\sigma(B^*B) - \varepsilon_\sigma(B)^*\varepsilon_\sigma(B)$ is a positive operator
-

$$\begin{aligned}\text{Var}(\tilde{B}; \omega_\sigma) &= \omega_\sigma(\tilde{B}^2) - \omega_\sigma(\tilde{B})^2 = \omega(\varepsilon_\sigma(B^2)) - \omega(\varepsilon_\sigma(B))^2 \\ &\geq \omega(A^2) - \omega(A)^2 = \text{Var}(A; \omega)\end{aligned}$$

post-selected state conditioned on B, is

$$\omega' = \frac{\mathcal{I}_\sigma(B)(\omega)}{\mathcal{I}_\sigma(B)(\omega)(1)}$$

A non-selective measurement results in

$\omega_B = \mathcal{I}_\sigma(B)(\omega) + \mathcal{I}_\sigma(1 - B)(\omega) = \mathcal{I}_\sigma(1)(\omega)$ which is independent of B.

Theorem

For a localizable in K^\perp , $\omega'(A) = \frac{\omega(A\varepsilon_\sigma(B))}{\omega(\varepsilon_\sigma(B))}$

Theorem

For a localizable in K^\perp , $\omega'(A) = \frac{\omega(A\varepsilon_\sigma(B))}{\omega(\varepsilon_\sigma(B))}$

Corollary

$\omega'(A) = \omega(A)$ iff A is uncorrelated with $\varepsilon_\sigma(B)$ in ω $\omega'(A) = \omega(A)$ for nonselective measurement of B .

If ω has a Reeh-Schlieder property (e.g. Minkowski vacuum state)

$$\omega'(A) = \omega(A) \iff \varepsilon_\sigma(B) = \omega(\varepsilon_\sigma(B))1$$

for observables localizable in k^\perp Post selection on any nontrivial measurements alters expectation values on K^\perp This is due the correlations.

Theorem

For a localizable in K^\perp , $\omega'(A) = \frac{\omega(A\varepsilon_\sigma(B))}{\omega(\varepsilon_\sigma(B))}$

Corollary

$\omega'(A) = \omega(A)$ iff A is uncorrelated with $\varepsilon_\sigma(B)$ in ω $\omega'(A) = \omega(A)$ for nonselective measurement of B .

If ω has a Reeh-Schlieder property (e.g. Minkowski vacuum state)

$$\omega'(A) = \omega(A) \iff \varepsilon_\sigma(B) = \omega(\varepsilon_\sigma(B))1$$

for observables localizable in k^\perp Post selection on any nontrivial measurements alters expectation values on K^\perp This is due the correlations. We have not assumed any rule that ω changes ω' across a surface in M . nor have we found any indication that such a rule is desirable.



Sorkin's protocol I

Model A and B using probes.




Figure: Sorkin's protocol[?]

Sorkin's protocol II

The scattering morphism gives $\Theta_2 C \otimes_2 Id \in \mathcal{U}(M; N)$ for $N \subset K_A^\perp \cap M_B^-$

Hence, C cannot determine what A has measured:

$$\begin{aligned}\omega_{AB}(C) &= (\omega \otimes \sigma_1 \otimes \sigma_2)((\Theta_1 \otimes_3 Id) \circ (\Theta_2 \otimes_2 Id)(C \otimes_2 1)) \\ &= (\omega \otimes \sigma_1 \otimes \sigma_2)(\Theta_2(C \otimes 1 \otimes 1)) = \omega_B(C)\end{aligned}$$

A Specific Probe Model I

Let the probe and the system both free scalar fields with a linear coupling on a bounded region.

$$\mathcal{L}_0 = -(\nabla_a \phi)(\nabla^a \phi) + m_\phi^2 \phi^2 - (\nabla_a \psi)(\nabla^a \psi) + m_\psi^2 \psi^2 := P\phi + Q\psi$$

$$\mathcal{L}_{\text{int}} = \rho \phi \psi, \quad \rho \in C_0^\infty(M), \quad K = \text{supp } \rho$$

$$E_p^\pm : C_0^\infty(M)(M) \rightarrow C_0^\infty(M)(M)$$

$$E_P^\pm P f = f, P E_P^\pm f = f, E_P^\pm f \subset J^\pm(\text{supp} f)$$

$E_P = E_P^- - E_P^+$ then every ϕ can be written as $\phi = E_P f$.

$$\underbrace{\begin{pmatrix} P & R \\ R & Q \end{pmatrix}}_T \underbrace{\begin{pmatrix} \phi \\ \psi \end{pmatrix}}_\zeta = 0$$

Quantization I

Let f, h be a (smooth) compactly supported functions on M

- $f \rightarrow \psi(f)$
- $\psi(f) = \psi(f)^*$,
- $\psi(Pf) = 0$
- $[\psi(f), \psi(h)] = iE_p[f, h]1$

$$E_p(f, h) := \int_M d\text{vol } f E_p h$$

Quantization II

If the region $L \subset K^\perp$ the algebras $\mathcal{A}(L) \otimes \mathcal{B}(L)$ and $\mathcal{C}(L)$ coincide as we saw earlier (commutative diagram).

By the time-slice property it is enough to consider the action of the scattering map on a generator $\zeta_0(F)$ of $\mathcal{A}(L) \otimes \mathcal{B}(L)$ with F (compactly) supported on M^+ Giving

$$\Theta \zeta_0(F) = \zeta_0(F - \tilde{R}E_T F) = \zeta_0(F - \tilde{R}E_T^- F)$$

where $\tilde{R} = T - P \oplus Q$

A Specific Probe Model I

$$\Theta(1 \otimes \psi(h)) = \phi(f^-) \otimes 1 + 1 \otimes \psi(h^-)$$

$$\begin{pmatrix} f^- \\ h^- \end{pmatrix} = \begin{pmatrix} 0 \\ h \end{pmatrix} - \begin{pmatrix} 0 & R \\ R & 0 \end{pmatrix} E_T^- \begin{pmatrix} 0 \\ h \end{pmatrix}$$

f^- supported in K and h^- in $M^+ \cup K \implies$

$$\Theta(1 \otimes e^{i\psi(h)}) = e^{i\phi(f^-)} \otimes e^{i\psi(h^-)}$$

Induced observables I

$$\varepsilon_\sigma(\psi(h)) = \eta_\sigma(\Theta(1 \otimes \psi(h))) = \phi(f^-) + \sigma(\psi(h^-))1$$

\Rightarrow

$$\varepsilon_\sigma(e^{i\psi(h)}) = \eta_\sigma(\Theta(1 \otimes e^{i\psi(h)})) = \sigma(e^{i\psi(h^-)})e^{i\phi(f^-)}$$

Sharpness of the observable $\tilde{\psi}(h)$ on the state σ

$$\begin{aligned} \text{Var}(\tilde{\psi}(h); \omega_\sigma) &= \omega_\sigma(\tilde{\psi}^2(h)) - \omega_\sigma(\tilde{\psi}(h))^2 \\ &= \omega(\varepsilon_\sigma(\psi(h)^2)) - \omega(\varepsilon_\sigma(\psi(h)))^2 \\ &= \omega(\phi(f^-)^2) + 2\omega(\phi(f^-))\sigma(\psi(h^-)) + \sigma(\psi(h^-))^2 \\ &\quad - (\omega(\phi(f^-))^2 + 2\omega(\phi(f^-))\sigma(\psi(h^-)) + \sigma(\psi(h^-))^2) \\ &= \text{Var}(\phi(f^-); \omega) + \text{Var}(\tilde{\psi}(h^-); \sigma) \end{aligned}$$



References I



Christopher J. Fewster and Rainer Verch
Quantum Fields and Local Measurements
Commun. Math. Phys. 378, 851–889 (2020)



Henning Bostelmann, Christopher J. Fewster and Maximilian H. Riep
Impossible measurements require impossible apparatus