

Lecture 6.1: Last time: more about spinors with/without real forms
 - related table of Clifford modules to
 Grauert/Localization spin groups

- consistently the Dirac spinor over \mathbb{C} using a polarization ($L \subseteq V_{\mathbb{C}}$)
 "old canonical quantization" $\max_{\vec{x}} \uparrow$ isotropic

- Gamma matrices from Clifford multiplication
 - built a pairing on the Dirac spinor: $S = \Lambda(L^\vee) = \text{functions on } \Pi L$
 (fermionic integration is strange: $(s, t) \rightarrow (s, t)_{\text{top}}$)

$$\begin{cases} \int d\theta \, 1 = 0 \\ \int d\theta \, \theta = 1 \end{cases}$$

$$S(\Lambda(s), \Lambda(t)) = (s, t)_{\text{Clifford action}}$$

this time: Finally write down all super~~transitions~~ algebras! $N = \mathbb{C}^2 \otimes S_{\mathbb{C}}$

V acts on itself by translations \Rightarrow abelian Lie algebra

V is a rep. of $\text{spin}(V) \Rightarrow$ "supercharges" should be as well

By spin and statistics, supercharges sit in (some # of copies of) the spin rep.
 (space of supercharges $= \Pi N$). The bracket $\Pi N \times \Pi N \rightarrow V$ should
 be symmetric, $\text{spin}(V)$ -equivariant, built from Π .

Dim (1): $\text{spin}(V) = \mathbb{Z}_2$, acts by parity.

Simplest example: $V = \mathbb{R}$, $N = \mathbb{R}$, bracket = tensor product ($Q^2 = p$)

Extend this $\leadsto N = \mathbb{R}^n = S^0 U_n$, $V = \mathbb{R}$, bracket = ~~tensor product~~ $\{Q_i, Q_j\} = p \delta_{ij}$
 \uparrow has a symmetric pairing

Dim 2: Lorentzian signature \Rightarrow New auto morphisms: $SO(U)$

$\text{Spin}(1,1) \cong M_2(\mathbb{R}) \cong M_2(\mathbb{C}) \cong M_2(\mathbb{R})$
 $\text{Spin}(1,1)^+ \cong \text{Spin}(1,0) = \mathbb{R} \oplus \mathbb{R} \Rightarrow$ Two real spin rep.

To construct them $Q = t \pm x \in V$. $\Lambda^0(L^\vee), \Lambda^1(L^\vee)$ are our two spin reps

Clifford multiplication $\leadsto \Lambda^0(L^\vee) = \mathbb{C} \xrightarrow{e_+} \Lambda^1(L^\vee) = \mathbb{C}$

$$\langle \Gamma(s_{\pm}, s_{\pm}), e_{+, -} \rangle = \begin{cases} (s_+, e_+ s_+) = (s_+, s_-) = I \Rightarrow \Gamma(s_+, s_+) = e_- \\ (s_+, e_- s_+) = 0 \\ (s_+, e_{\pm} s_-) = (s_+, s_+) = 0 \end{cases}$$

we can add just s_+ to the algebra: $(s_-, e_- s_-) = (s_-, s_+) = I \Rightarrow \Gamma(s_-, s_-) = e_+$

$$N = S_+^{\otimes 2}, V = V_+ \oplus V_- \quad \left(\text{we get } 1+1 \leadsto -\frac{\partial^2}{\partial t^2} + \frac{\partial^2}{\partial x^2} = 0 \right)$$

$$S_+^{\otimes 2} = V_- \quad \text{in coords: } Q_-, P_-, P_- \quad \left(-\frac{\partial^2}{\partial t^2} + \frac{\partial^2}{\partial x^2} \right) \left(\frac{\partial^2}{\partial t^2} + \frac{\partial^2}{\partial x^2} \right) = 0$$

the $(1,1)$ looks like 2 indep. copies of SQM $P^2 = P_+ P_-$

$N = (N_+, N_-)$ "amount of extended susy"

→ unique 2-dim real spin rep.
 $E_1, X_1, X_2 \rightsquigarrow E_T = E \pm X_1, X_2$
 generated by e_+ as before



$\rho(E) = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \rho(E_1) = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \rho(E_2) = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$
 $\rho(E_1^2) = \rho(E_2^2) = \rho(E_1 E_2) = \rho(E_2 E_1) = \rho(E_1^2 + E_2^2) = \rho(2E) = 2\rho(E)$

Table: Constructing supersymmetric algebras in various dimensions

dim (bosons)	dim (fermions)	minimal # of supersymmetries	algebra
0 (HW)	1	1	$GL(N)$
1 (any on)	1	1	$SO(N)$
2	2 (Weyl)	1	$SO(N) \ltimes SO(N)$
3	2	1	$SO(N)$
4	4 (Majorana)	1	$U(N) \ltimes Sp(N)$
6	8 (Majorana-Weyl)	1	$U(N) \ltimes Sp(N)$

Unique spinor: $S = \mathbb{R}^2$
 $Spin(1,2) \cong SL(2, \mathbb{R})$
 $S \otimes S = Sym^2(S) \oplus \wedge^2(S)$
 $S \otimes S \cong S \otimes S \cong \mathbb{R}^2$
 $SO(N) \ltimes Spin(N)$
 To extend $SO(N) \ltimes Spin(N)$ to $SO(N) \ltimes Spin(N) \ltimes Spin(N)$
 $Spin(1,2) \cong SL(2, \mathbb{R})$
 $Spin(1,2) \cong SL(2, \mathbb{R})$
 $Spin(1,2) \cong SL(2, \mathbb{R})$

$S = \mathbb{C}^2, \bar{S} = \mathbb{C}^2 = S^V \leadsto S \otimes \bar{S} = \text{Hom}(S, S) = M_2(\mathbb{C})$
 $(S \otimes \bar{S})_0 \cong \text{traceless } 2 \times 2 \text{ complex matrices}$
 $\begin{bmatrix} a \\ b \end{bmatrix} \xleftrightarrow{S} \begin{bmatrix} a & b \end{bmatrix} \xrightarrow{S} \begin{bmatrix} -b \\ a \end{bmatrix}$

$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \otimes \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix}$
 $i \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \otimes \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$
 $SL(2, \mathbb{C}) \subset \text{hermitian } 2 \times 2 \text{ det. } \mathbb{R}^3$

$[S \otimes \bar{S}] = \text{a rep of } \mathfrak{spin}(3) \leadsto \Lambda^2(S \otimes \bar{S}) = \Lambda^2(S) \oplus \Lambda^2(\bar{S})$
 $(S \otimes \bar{S})_{\mathbb{R}}$ is a four real dimension (majorana)
 $\text{normally } \leadsto \{Q_\alpha, \bar{Q}_\beta\} = P_{\alpha\beta} = \gamma^\mu_{\alpha\beta} P_\mu$
 $Q_\alpha^\dagger = \bar{Q}_\alpha$

Extended susy in 4d:
 $S \otimes_{\mathbb{C}} U, U = \mathbb{C}^N \quad (S \otimes_{\mathbb{C}} U \oplus \bar{S} \otimes_{\mathbb{C}} \bar{U})_{\mathbb{R}} \quad U \otimes_{\mathbb{R}} \bar{U} \rightarrow \mathbb{C}$
 Hermitian inner product

Dim 6 $Cl(5)^+ \cong Cl(1,4) \cong M_2(Cl(0,3)) \cong M_2(\mathbb{H}) \oplus M_2(\mathbb{H})$

$(Spin(5))_0 \cong SL(2, \mathbb{H})$
 $Cl(5S) \leadsto 2 \text{ chiral spinors}$
 $S^+ \cong \mathbb{H}^2 \cong \mathbb{C}^4$
 $S^- \cong \mathbb{H}^2 \cong \mathbb{C}^4$

Dirac construction: $L \subseteq \mathbb{C}^6$
 $S_+ = \Lambda^{\text{even}}(L^V) \cong \Lambda^0 \oplus \Lambda^2 \cong \mathbb{C} \oplus \Lambda^2 L^V$
 $S_- = \Lambda^{\text{odd}}(L^V) \cong \Lambda^1 \oplus \Lambda^3 \cong L^V \oplus \Lambda^3 L^V$
 $\Gamma: S_+ \otimes S_+ \rightarrow V$
 antisymmetrically!
 $\dim(\Lambda^2 S_+) = 6 = \dim(V)$
 \hookrightarrow can be diagonalized

$\dim^2(S \otimes U) = \dim^2(S) \oplus \dim^2(U) \oplus \Lambda^2(S) \otimes \Lambda^2(U)$
 $(S_+ \otimes_{\mathbb{C}} U)_{\mathbb{R}} \cong \mathbb{R}^8$ "symplectic - quaternionic way"
 $\Lambda^2 S_+ \cong V, \Lambda^2 U \cong \mathbb{R}$
 $\mathbb{R} \xrightarrow{V} \mathbb{R}$ symplectic form.

Brief recap: Super symmetry \leftarrow Bosons exchange statistics
 Motivation: The Super-Poincare Hilbert Space is a super vector space

- constraints of susy v BPS bound (only is positive)

boson-fermion pairs degenerate

They are exactly solvable

- identification of fermionic degrees of freedom (fermionic phase spaces are inner product spaces)

- Clifford algebra (canonical anticommutative relations)

- classified their rep. in 1972, depending on

- spinors, map $(R^{0,1} \rightarrow R^{1,0})$ is isomorphic

odd degrees of freedom arise as quantum or spinor fields

There are Weyl, Majorana, symplectic Majorana, Dirac, and spinors

- supersymmetric algebra for $d=2,3,4,6$ minus $N \geq 1$

real forms ... (figure out how 8 matrices work out in general dim)

Goal: understand susy field theories and interactions in higher spacetime dim

A particularly nice way "5-spaces"

Also: There is a "no-go" theorem by Coleman-Mandula, is a

relativistic QFT has an algebra of symmetries \mathfrak{g} that includes

fermionic symmetries $p(a, p \in \mathfrak{g})$ then as (assumptions) $\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{h}$

and \mathfrak{h} acts internally

Don't apply to super Lie algebras!

Fixed up by Haag-Lopuszanski-Schüster: only the "exotic" super Lie

algebras are allowed

General structure of super-Poincaré algebras

- choose the space of supercharges $W = \mathfrak{g} \otimes U^1, \{u, u^3\} \rightarrow 1$

$0 \rightarrow T \rightarrow \text{super } T \rightarrow \pi W \rightarrow 0$

$\mathfrak{p}^+ \oplus \mathfrak{p}^- \oplus \mathfrak{p}^0 \oplus \mathfrak{p}^1 \oplus \mathfrak{p}^2 \oplus \mathfrak{p}^3 \oplus \mathfrak{p}^4 \oplus \mathfrak{p}^5 \oplus \mathfrak{p}^6 \oplus \mathfrak{p}^7 \oplus \mathfrak{p}^8 \oplus \mathfrak{p}^9 \oplus \mathfrak{p}^{10} \oplus \mathfrak{p}^{11} \oplus \mathfrak{p}^{12} \oplus \mathfrak{p}^{13} \oplus \mathfrak{p}^{14} \oplus \mathfrak{p}^{15} \oplus \mathfrak{p}^{16} \oplus \mathfrak{p}^{17} \oplus \mathfrak{p}^{18} \oplus \mathfrak{p}^{19} \oplus \mathfrak{p}^{20} \oplus \mathfrak{p}^{21} \oplus \mathfrak{p}^{22} \oplus \mathfrak{p}^{23} \oplus \mathfrak{p}^{24} \oplus \mathfrak{p}^{25} \oplus \mathfrak{p}^{26} \oplus \mathfrak{p}^{27} \oplus \mathfrak{p}^{28} \oplus \mathfrak{p}^{29} \oplus \mathfrak{p}^{30} \oplus \mathfrak{p}^{31} \oplus \mathfrak{p}^{32} \oplus \mathfrak{p}^{33} \oplus \mathfrak{p}^{34} \oplus \mathfrak{p}^{35} \oplus \mathfrak{p}^{36} \oplus \mathfrak{p}^{37} \oplus \mathfrak{p}^{38} \oplus \mathfrak{p}^{39} \oplus \mathfrak{p}^{40} \oplus \mathfrak{p}^{41} \oplus \mathfrak{p}^{42} \oplus \mathfrak{p}^{43} \oplus \mathfrak{p}^{44} \oplus \mathfrak{p}^{45} \oplus \mathfrak{p}^{46} \oplus \mathfrak{p}^{47} \oplus \mathfrak{p}^{48} \oplus \mathfrak{p}^{49} \oplus \mathfrak{p}^{50} \oplus \mathfrak{p}^{51} \oplus \mathfrak{p}^{52} \oplus \mathfrak{p}^{53} \oplus \mathfrak{p}^{54} \oplus \mathfrak{p}^{55} \oplus \mathfrak{p}^{56} \oplus \mathfrak{p}^{57} \oplus \mathfrak{p}^{58} \oplus \mathfrak{p}^{59} \oplus \mathfrak{p}^{60} \oplus \mathfrak{p}^{61} \oplus \mathfrak{p}^{62} \oplus \mathfrak{p}^{63} \oplus \mathfrak{p}^{64} \oplus \mathfrak{p}^{65} \oplus \mathfrak{p}^{66} \oplus \mathfrak{p}^{67} \oplus \mathfrak{p}^{68} \oplus \mathfrak{p}^{69} \oplus \mathfrak{p}^{70} \oplus \mathfrak{p}^{71} \oplus \mathfrak{p}^{72} \oplus \mathfrak{p}^{73} \oplus \mathfrak{p}^{74} \oplus \mathfrak{p}^{75} \oplus \mathfrak{p}^{76} \oplus \mathfrak{p}^{77} \oplus \mathfrak{p}^{78} \oplus \mathfrak{p}^{79} \oplus \mathfrak{p}^{80} \oplus \mathfrak{p}^{81} \oplus \mathfrak{p}^{82} \oplus \mathfrak{p}^{83} \oplus \mathfrak{p}^{84} \oplus \mathfrak{p}^{85} \oplus \mathfrak{p}^{86} \oplus \mathfrak{p}^{87} \oplus \mathfrak{p}^{88} \oplus \mathfrak{p}^{89} \oplus \mathfrak{p}^{90} \oplus \mathfrak{p}^{91} \oplus \mathfrak{p}^{92} \oplus \mathfrak{p}^{93} \oplus \mathfrak{p}^{94} \oplus \mathfrak{p}^{95} \oplus \mathfrak{p}^{96} \oplus \mathfrak{p}^{97} \oplus \mathfrak{p}^{98} \oplus \mathfrak{p}^{99}$

- fermion is $\mathfrak{p}^+ \oplus \mathfrak{p}^- \oplus \mathfrak{p}^0 \oplus \mathfrak{p}^1 \oplus \mathfrak{p}^2 \oplus \mathfrak{p}^3 \oplus \mathfrak{p}^4 \oplus \mathfrak{p}^5 \oplus \mathfrak{p}^6 \oplus \mathfrak{p}^7 \oplus \mathfrak{p}^8 \oplus \mathfrak{p}^9 \oplus \mathfrak{p}^{10} \oplus \mathfrak{p}^{11} \oplus \mathfrak{p}^{12} \oplus \mathfrak{p}^{13} \oplus \mathfrak{p}^{14} \oplus \mathfrak{p}^{15} \oplus \mathfrak{p}^{16} \oplus \mathfrak{p}^{17} \oplus \mathfrak{p}^{18} \oplus \mathfrak{p}^{19} \oplus \mathfrak{p}^{20} \oplus \mathfrak{p}^{21} \oplus \mathfrak{p}^{22} \oplus \mathfrak{p}^{23} \oplus \mathfrak{p}^{24} \oplus \mathfrak{p}^{25} \oplus \mathfrak{p}^{26} \oplus \mathfrak{p}^{27} \oplus \mathfrak{p}^{28} \oplus \mathfrak{p}^{29} \oplus \mathfrak{p}^{30} \oplus \mathfrak{p}^{31} \oplus \mathfrak{p}^{32} \oplus \mathfrak{p}^{33} \oplus \mathfrak{p}^{34} \oplus \mathfrak{p}^{35} \oplus \mathfrak{p}^{36} \oplus \mathfrak{p}^{37} \oplus \mathfrak{p}^{38} \oplus \mathfrak{p}^{39} \oplus \mathfrak{p}^{40} \oplus \mathfrak{p}^{41} \oplus \mathfrak{p}^{42} \oplus \mathfrak{p}^{43} \oplus \mathfrak{p}^{44} \oplus \mathfrak{p}^{45} \oplus \mathfrak{p}^{46} \oplus \mathfrak{p}^{47} \oplus \mathfrak{p}^{48} \oplus \mathfrak{p}^{49} \oplus \mathfrak{p}^{50} \oplus \mathfrak{p}^{51} \oplus \mathfrak{p}^{52} \oplus \mathfrak{p}^{53} \oplus \mathfrak{p}^{54} \oplus \mathfrak{p}^{55} \oplus \mathfrak{p}^{56} \oplus \mathfrak{p}^{57} \oplus \mathfrak{p}^{58} \oplus \mathfrak{p}^{59} \oplus \mathfrak{p}^{60} \oplus \mathfrak{p}^{61} \oplus \mathfrak{p}^{62} \oplus \mathfrak{p}^{63} \oplus \mathfrak{p}^{64} \oplus \mathfrak{p}^{65} \oplus \mathfrak{p}^{66} \oplus \mathfrak{p}^{67} \oplus \mathfrak{p}^{68} \oplus \mathfrak{p}^{69} \oplus \mathfrak{p}^{70} \oplus \mathfrak{p}^{71} \oplus \mathfrak{p}^{72} \oplus \mathfrak{p}^{73} \oplus \mathfrak{p}^{74} \oplus \mathfrak{p}^{75} \oplus \mathfrak{p}^{76} \oplus \mathfrak{p}^{77} \oplus \mathfrak{p}^{78} \oplus \mathfrak{p}^{79} \oplus \mathfrak{p}^{80} \oplus \mathfrak{p}^{81} \oplus \mathfrak{p}^{82} \oplus \mathfrak{p}^{83} \oplus \mathfrak{p}^{84} \oplus \mathfrak{p}^{85} \oplus \mathfrak{p}^{86} \oplus \mathfrak{p}^{87} \oplus \mathfrak{p}^{88} \oplus \mathfrak{p}^{89} \oplus \mathfrak{p}^{90} \oplus \mathfrak{p}^{91} \oplus \mathfrak{p}^{92} \oplus \mathfrak{p}^{93} \oplus \mathfrak{p}^{94} \oplus \mathfrak{p}^{95} \oplus \mathfrak{p}^{96} \oplus \mathfrak{p}^{97} \oplus \mathfrak{p}^{98} \oplus \mathfrak{p}^{99}$

- super-Poincaré is super $T \times \text{spin}(d-1)$

Two ways of building supersymmetric field theories

- components

if I think about Kaluza-Klein reduction, e.g. from 5d to 4d

Let's say I have a gauge field $A_\mu(x) \in \mathfrak{so}(3,1)$ $R^{3,1} \times S^1$

$A = A_\mu dx^\mu = A_\mu(x, \theta) dx^\mu + A_5(x, \theta) d\theta$

$A_5 \sim \sum A_n^{(1)}(y) e^{in\theta}$ little small: very massive for $n \neq 0$

$\int_{S^1} A_n^{(1)}(y) dy + A_n^{(0)}(y) d\theta$

If I have an odd collection: $\mathbb{R}^{1|1}$ Scalar Superfield: $\mathcal{F}(y^i, \epsilon)$
 A super space is like a matrix algebra but its exact inverse is
 that it doesn't have to be small to be invertible $\xrightarrow{\text{bosonic scalar}} \mathcal{F}(y) + \epsilon \mathcal{F}_1(y)$
 $\epsilon^2 = 0$ \uparrow fermionic scalar

Exponentiating Supertranslations, $E \in d=1, N=2$

Translations generated by P , Symmetries Q_1, Q_2 with $\{Q_i, Q_j\} = 2\delta_{ij}P$
 only no zero bracket. Consider a finite translation: $T_a = \exp(a \frac{\partial}{\partial x})$, $P = \frac{\partial}{\partial x}$
 (use convention that P is anti hermitian) hence $T_a: \mathcal{F}(x) \rightarrow \mathcal{F}(x+a)$
could try (let's see Q_i translations in odd directions)

$$\exp(\epsilon \frac{\partial}{\partial \epsilon}) \rightarrow \mathcal{F}(\epsilon) = \mathcal{F}_0 + \epsilon \mathcal{F}_1$$

$$\epsilon \rightarrow \epsilon + \epsilon_1 = (\mathcal{F}_0 + \epsilon \mathcal{F}_1) + \epsilon_1 \mathcal{F}_1$$

$$\text{then } 1 + \epsilon \frac{\partial}{\partial \epsilon}$$

BCH formula $\rightarrow e^A e^B = e^{A+B + \frac{1}{2}[A,B] + \dots}$ For super translations
 algebras

$$\Rightarrow e^{\epsilon Q_1} e^{\epsilon_1 Q_1} = e^{\epsilon Q_1 + \epsilon_1 Q_1 + \epsilon \epsilon_1 P} \sim Q_1 = \frac{\partial}{\partial \epsilon} + \epsilon \frac{\partial}{\partial x}$$

$$Q_1 \mathcal{F} = (\frac{\partial}{\partial \epsilon} + \epsilon \frac{\partial}{\partial x})(\mathcal{F}_0 + \epsilon \mathcal{F}_1) = \mathcal{F}_1 + \epsilon \mathcal{F}_0'$$

Finite super translations: $\exp(\epsilon Q_1) = 1 + \epsilon Q_1 = 1 + \epsilon \frac{\partial}{\partial \epsilon} + \epsilon \epsilon_1 \frac{\partial}{\partial x}$
 $\mathcal{F}_0 + \epsilon \mathcal{F}_1 \rightarrow \mathcal{F}_0 + \epsilon_1 \mathcal{F}_1 + \epsilon \epsilon_1 \mathcal{F}_0'$


i.e.

$$\mathcal{F}(x, \epsilon) \xrightarrow{T_{\epsilon_1}} \mathcal{F}(x + \epsilon_1 \epsilon, \epsilon + \epsilon_1) \xrightarrow{T_{\epsilon_1}} \mathcal{F}(x + \epsilon_1 \epsilon + (\epsilon + \epsilon_1) \epsilon_1, \epsilon + \epsilon_1 + \epsilon_1^2) \\ = \mathcal{F}(x + (\epsilon_1 + \epsilon_1^2) \epsilon + \epsilon_1 \epsilon_1, \epsilon + (\epsilon_1 + \epsilon_1^2))$$

Last time Started discussing Superfields
 (Functions of fields of superspace)

Exam review: wed 3.8, 14:00
 Exam: Fri 5.8, 16:00
 Essay: Sat 9.8, 13:00

Superspace



\sim Super spacetime is modeled on finite supertranslations

BCH Formula

$$e^A e^B = e^{A+B + \frac{1}{2}[A,B] + \dots}$$

(b-a) finite translations

Simple example of a superfield: $N=2$ SQM ~ 2 symmetries $\sim \{Q_i, Q_j\} = \delta_{ij}P$, 5 fields

Corresponding superspace: $T = \exp(t) \simeq \mathbb{R}^{1|2}$

$\uparrow t = \mathbb{R}^{1|2}$

then T is equipped with one even and 2 odd coordinates

2 bosonic fields

A general function looks like $\mathcal{F}(x, \theta_1, \theta_2) = \mathcal{F}_0(x) + \mathcal{F}_1(x) \theta_1 + \mathcal{F}_2(x) \theta_2$

+ $\mathcal{F}_3(x) \theta_1 \theta_2$ 2 fermionic fields

How super symmetry transf act?

$$\exp(a \cdot P) \sim \exp(a \frac{\partial}{\partial x}) : \mathcal{F}(x, \theta_1, \theta_2)$$

$$\text{on odd: } \exp(\epsilon_1 Q_1 + \epsilon_2 Q_2) = \exp(\epsilon_1 Q_1) \exp(\epsilon_2 Q_2) \exp(-\epsilon_1 \epsilon_2 P) \rightarrow \mathcal{F}(x + \epsilon_1 \epsilon_2, \theta_1, \theta_2)$$

$$\int \partial f = a \int \partial f$$

$$\frac{\partial}{\partial t} = \frac{\partial}{\partial t} + \frac{\partial}{\partial x} \frac{dx}{dt} = \frac{\partial}{\partial t} + v \frac{\partial}{\partial x}$$

$$\int \frac{\partial}{\partial x} f(x) dx = f(x) \Big|_a^b$$

$$\int \frac{\partial}{\partial x} f(x) dx = f(x) \Big|_a^b$$

$$\int \frac{\partial}{\partial x} f(x) dx = f(x) \Big|_a^b$$

$$\int \frac{\partial}{\partial x} f(x) dx = f(x) \Big|_a^b$$

$$\int \frac{\partial}{\partial x} f(x) dx = f(x) \Big|_a^b$$

$$\int \frac{\partial}{\partial x} f(x) dx = f(x) \Big|_a^b$$

$$\int \frac{\partial}{\partial x} f(x) dx = f(x) \Big|_a^b$$

$$\int \frac{\partial}{\partial x} f(x) dx = f(x) \Big|_a^b$$

$$\int \frac{\partial}{\partial x} f(x) dx = f(x) \Big|_a^b$$

The functions on a group are a rep of G by translating. In fact, in two ways: left and right translations. Right translations give other rep. of symmetry.

$$\frac{\partial}{\partial x} f(x) = f(x) \frac{\partial}{\partial x}$$

Note that f is a superfield but ∂f is not a superfield!

② Integrate out extra boson f_2 ("auxiliary field")

① what do action functionals look like?

$$= (f_1 + f_2 \partial_2 + \partial_2 f_1 + \partial_2^2 f_1) (f_1 + f_2 \partial_2 + \partial_2 f_1 + \partial_2^2 f_1)$$

$$= (f_1 + f_2 \partial_2 + \partial_2 f_1 + \partial_2^2 f_1) (f_1 + f_2 \partial_2 + \partial_2 f_1 + \partial_2^2 f_1)$$

$$\left\{ \frac{\partial}{\partial x} + \partial_2 \frac{\partial}{\partial x} + \partial_2^2 \frac{\partial}{\partial x} \right\} = 2 \frac{\partial}{\partial x}$$

$$\exp(\epsilon_1 \partial_1) = \exp(\epsilon_1 (\frac{\partial}{\partial x} + \partial_2 \frac{\partial}{\partial x}))$$

$$\left\{ \begin{array}{l} \partial f_1 = \epsilon_1 f_1 \\ \partial f_2 = \epsilon_1 f_2 \\ \partial f_3 = \epsilon_1 f_3 \end{array} \right\}$$

$$\int d\theta = \frac{\partial}{\partial \theta} \leadsto \int d\theta f(\theta) = \int d\theta (f_0 + \frac{f_1}{a} \theta) = \frac{f_1}{a} \Rightarrow$$

$$d\theta = \frac{d\phi}{a} = \left(\frac{\partial \phi}{\partial \theta}\right)^{-1} d\phi \leftarrow \text{recall } \det\left(\frac{A}{B}\right) = \frac{\det A}{\det B}$$

Try a kinetic potential for the super field F .

$$\int_{\mathbb{R}^{1|2}} dx d\theta_1 d\theta_2 D_1 F D_2 F$$

$$\begin{aligned} D_1 F &= -f_1 - f_{12} \theta_2 + \frac{\partial f_0}{\partial x} \theta_1 + \frac{\partial f_2}{\partial x} \theta_1 \theta_2 \\ D_2 F &= -f_2 + f_{12} \theta_1 + \frac{\partial f_0}{\partial x} \theta_2 - \frac{\partial f_1}{\partial x} \theta_1 \theta_2 \\ &= \left[f_1 \frac{\partial f_1}{\partial x} - f_2 \frac{\partial f_2}{\partial x} + \left(\frac{\partial f_0}{\partial x}\right)^2 - f_{12}^2 \right] \theta_1 \theta_2 + \dots \end{aligned}$$

$\underbrace{\quad}_{\text{2-fermion kinetic terms}} \quad \underbrace{\quad}_{\text{boson kinetic term}} \quad \underbrace{\quad}_{\text{no derivatives} \leadsto \text{eq. of motion} \rightarrow f_{12}=0}$

Last time: 2 possible approaches to susy field theories.
 (1) "components" write down an ordinary theory ($S = \int \mathcal{L}(\phi, \psi)$)
 \leadsto come up with some susy transformations
 $(\delta \phi = \epsilon \psi, \delta \psi = \epsilon \not{\partial} \phi)$ check by hand that $\delta S = 0$ i.e. there is

e.g. 1dim $N=2$ susy (a long time ago)

(2) use super fields i.e. functions on the superspace and write $S = \int dx d\theta \mathcal{L}(\Phi)$
 $\mathbb{R}^{1|2}$ \uparrow number of super charges depends on the dim of the spin rep.

- susy trans are automatic: $Q = \frac{\partial}{\partial \theta} + \theta \frac{\partial}{\partial x}$
 If I use N for spacetime and κ for index on space w of susys.

$$Q_\alpha = \frac{\partial}{\partial \theta^\alpha} + \theta^\beta \gamma_{\alpha\beta}^\mu \frac{\partial}{\partial x^\mu}, \quad \{Q_\alpha, Q_\beta\} = 2\gamma_{\alpha\beta}^\mu P_\mu$$

A general element $Q \in W \leadsto Q = \epsilon^\alpha Q_\alpha$

$$\begin{aligned} \{Q_\alpha, Q_\beta\} &= \left\{ \frac{\partial}{\partial \theta^\alpha} + \theta^\lambda \gamma_{\alpha\lambda}^\mu \frac{\partial}{\partial x^\mu}, \frac{\partial}{\partial \theta^\beta} + \theta^\nu \gamma_{\beta\nu}^\mu \frac{\partial}{\partial x^\mu} \right\} = \delta_{\alpha\beta}^\mu \gamma_{\mu\lambda}^\nu \frac{\partial}{\partial x^\nu} \\ &= \gamma_{\alpha\beta}^\nu \frac{\partial}{\partial x^\nu} + \gamma_{\alpha\beta}^\mu \frac{\partial}{\partial x^\mu} = 2\gamma_{\alpha\beta}^\mu \frac{\partial}{\partial x^\mu} \end{aligned}$$

There are also "covariant derivatives"

$$D_\alpha = -\frac{\partial}{\partial \theta^\alpha} + \theta^\beta \gamma_{\alpha\beta}^\mu \frac{\partial}{\partial x^\mu} \leadsto \{Q_\alpha, P_\beta\} = 0$$

$$\{D_\alpha, D_\beta\} = 0$$

Note that

$$Q \mathcal{L}(\Phi) \leadsto S = \int d\theta dx Q \mathcal{L}$$

\uparrow has $\left(\frac{\partial}{\partial \theta} + \theta \frac{\partial}{\partial x}\right)$

optimal $\{a, a^p\} = a^p$ basis

$$\Phi = \Phi(x, \theta_1, \theta_2) = f_1(x) + f_2(x) \theta_1 + f_3(x) \theta_2 + f_4(x) \theta_2^2$$

I need a form of the form $(f_1^2 \theta_1^2 + f_2^2 \theta_1 \theta_2 + f_3^2 \theta_2^2)$ guess $\sim \sqrt{D_1 \Phi D_2 \Phi} \cdot (D_1 \Phi) = (-\frac{\partial}{\partial x} + \theta_1 \frac{\partial}{\partial \theta_1}) \Phi = (-f_1 - f_2 \theta_1 + \theta_1 \frac{\partial}{\partial \theta_1}) \Phi$

$$(D_2 \Phi) = (-f_2 + f_2 \theta_1 + \theta_2 \frac{\partial}{\partial \theta_1} - \frac{\partial}{\partial \theta_2}) \Phi = (-\frac{\partial}{\partial \theta_2} + \theta_2 \frac{\partial}{\partial \theta_2} + \theta_1 \frac{\partial}{\partial \theta_1}) \Phi$$

$$\int d\theta_1 d\theta_2 \Phi = f_1 \frac{\partial}{\partial \theta_1} - f_2 \frac{\partial}{\partial \theta_2} + \left(\frac{\partial f_1}{\partial \theta_1} - f_2 \right) \Phi$$

Dirac language
kheic
Hess

We constructed

$N=2$ SQM using a "superpotential" $W(\phi)$ mass term? Φ^2 is reasonable.

$$\int d\theta^2 \Phi^2 = 2(f_1 f_2 + f_2 f_3 + f_3 f_4) \sim \int d\theta^2 \Phi (D_1 \Phi) (D_2 \Phi) + \Phi^2$$

$$= \left(\frac{\partial f_1}{\partial \theta_1} \right)^2 + f_1 \frac{\partial f_2}{\partial \theta_1} - f_2 \frac{\partial f_1}{\partial \theta_2} + \frac{\partial f_2}{\partial \theta_2} + m f_2^2 + m f_3 f_4$$

LEM: $\frac{\partial f_2}{\partial \theta_2} = 0 = -2f_2 + m f_3$

Dirac mass
(1) a magnetic basis

More generally, circle with $W(\Phi)$ for some W

If $W(\Phi) = \Phi^2 \sim \int d\theta^2 \Phi^2 = \int d\theta^2 f_1 \theta_1^2 + f_2 \theta_1 \theta_2 + f_3 \theta_2^2 = \int d\theta^2 (f_1 \theta_1^2 + f_2 \theta_1 \theta_2 + f_3 \theta_2^2)$

ECM for $f_{12} \sim 2f_{12} = 2f_{12} = w'(f_2)$

How does one see the component is easy to find?

$$(D_1 = \frac{\partial}{\partial \theta_1} + \theta_1 \frac{\partial}{\partial x}) \Phi = \frac{\partial \Phi}{\partial \theta_1} + \theta_1 \frac{\partial \Phi}{\partial x} = f_1 + f_2 \theta_1 + \theta_1 \frac{\partial f_1}{\partial \theta_1} + \theta_1 \frac{\partial f_2}{\partial \theta_1} \theta_2$$

$$\Rightarrow \delta \phi_1 = f_1, \delta \phi_2 = f_2, \delta \phi_3 = f_3, \delta \phi_4 = f_4$$

$$\begin{aligned} dQ_2 f_0 &= f_2 & \int_Q f_0 &= F_1 \\ dQ_2 f_1 &= -\frac{1}{2} \omega'(f_0) & dQ_1 f_1 &= \frac{\partial f_0}{\partial x} \\ dQ_2 f_2 &= \frac{\partial f_0}{\partial x} & dQ_1 f_2 &= \frac{1}{2} \omega'(f_0) \end{aligned}$$

	how many fields (off shell)	how many degrees of freedom (on shell)
Scalar	1	1
Gauge fields	d	$d-2$
Dirac fermion (massive)	$2 \lfloor d/2 \rfloor$	$2 \lfloor d/2 \rfloor - 1$
Auxiliary scalar	1	0

	on shell	off shell
scalar	1	1 $\in \mathbb{R}$
Fermi	1	2 $\in \mathbb{C}$
gauge	0	1 $\in \mathbb{R}$

In 4-dim minimal ($N=1$) susy what might be expected ^{(2) 4} super p
Minimal fermions have 4 real components (where Majorana or Weyl)
 \Rightarrow on-shell 2-degrees of freedom

	on shell	off shell
fermi	2	4 = 2cpx
scalars	2	2 = 1cpx
auxiliary	0	2 = 1cpx

	on shell	off shell
fermi	2	4
vector	2	4

Chiral superfield

(components of Φ)

$\langle \theta \rangle$	1	+	vector	←	one scalar
$\langle \theta \rangle$	4	-	quadr	←	one Dirac spinor
$\langle \theta \rangle^2$	6	+		←	
$\langle \theta \rangle^3$	4	-		←	
$\langle \theta \rangle^4$	1	+		←	

$\theta^\alpha \theta^\beta$
 \uparrow
 1 scalar
 one Dirac spinor

$\theta^\alpha \bar{\theta}^{\dot{\beta}}$
 \uparrow
 1 vector
 "vectors as bispinors"

$\bar{\theta}^{\dot{\alpha}} \bar{\theta}^{\dot{\beta}}$
 \uparrow
 1 scalar

$[S_+, \bar{S}_-] \approx V$
 "vectors as bispinors"

Real superfield

1 real scalar

1 Majorana

{ 1 Real vector
1 complex (two real) scalars }

1 Majorana

1 Real scalar

To pick out subrepresentations of the free superfields
need to impose constraints compatible w/ supersymmetry
can act w/ D operator for example
Chiral superfield: $\bar{D}_\alpha \Phi = 0$. If $D_\mu \Phi = \bar{D}_2 \Phi = 0 \Rightarrow \{D_1, \bar{D}_2\} \Phi = 0$
 \downarrow
 $\left(-\frac{2}{2\lambda^2} + \theta^\mu \sigma_{\mu\alpha}^\nu \frac{\partial}{\partial \lambda^\nu} \right) \Phi = 0$ If $D_2 \Phi = 0, \bar{D}_2 \Phi = 0 \Rightarrow \rho \Phi = 2\lambda_2 \Phi$
for massive case

1st try $\gamma^\mu = \gamma^\mu + \theta^\alpha \frac{\sigma^\mu}{\alpha \beta} \bar{\theta}^\beta \rightarrow \begin{cases} D_\alpha = -\frac{2}{2\theta^\alpha} \\ D_\alpha = -\frac{2}{2\theta^\alpha} + 2\bar{\theta}^\alpha \frac{2}{2\gamma} \end{cases}$ $\frac{\partial \phi}{\partial x^2} = 0 \rightarrow$ no degrees of freedom

2 complex
1 auxiliary

$\Rightarrow \underline{\Phi} = \Phi(X, \theta)$ In the literature, a single chiral multiplet

$$① \int_{\mathbb{R}^{1,3}} d^4x d^2\theta d^2\bar{\theta} K(\Phi, \bar{\Phi})$$

\uparrow
 $K=K$
covariant to be built

The condition $\bar{D}\Phi=0$ gives other possibilities

$$② \int d^4x d^2\theta W(\Phi) + h.c.$$

holomorphic

"D-terms"

argument of SUSY

$$\int d^2\theta Q(\dots)$$

$$\int d^2\theta \left(\frac{\partial}{\partial \theta} + \bar{\theta} \frac{\partial}{\partial \chi} \right) (\dots)$$

"F-terms"

↑
varies
a θ

↑
total
derivative

$$\int d^2\theta \left(\frac{\partial}{\partial \theta} + \bar{\theta} \frac{\partial}{\partial \chi} \right)$$

problem, but since we are integrating on Φ chiral events

$$\frac{\partial}{\partial \theta} = \theta \frac{\partial}{\partial \chi} \rightarrow \text{will be a total derivative} \checkmark$$

Guess for kinetic term for chiral multiplets

$$\int d^4x d^2\theta d^2\bar{\theta} (\bar{\Phi} \Phi)$$

$$\Phi = \phi(\gamma) + \chi_\alpha(\gamma) \theta^\alpha + F(\gamma) \theta^2, \quad \gamma = x + \theta \sigma \bar{\theta} \rightarrow \Phi \in \mathbb{C}$$

$$\Rightarrow \phi(\gamma) = \phi(x) + \frac{\partial \phi}{\partial x^\mu} \theta \sigma^\mu \bar{\theta} + \frac{\partial^2 \phi}{\partial x^\mu \partial x^\nu} (\theta \sigma^\mu \bar{\theta}) (\theta \sigma^\nu \bar{\theta})$$

$$\Rightarrow \theta^\alpha \chi_\alpha(\gamma) = \theta^\alpha \chi_\alpha(x) + \frac{\partial \chi_\alpha}{\partial x^\mu} \theta^\alpha \theta^\beta \sigma^\mu_{\beta\dot{\alpha}} \bar{\theta}^{\dot{\alpha}} + \frac{\partial^2 \chi_\alpha}{\partial x^\mu \partial x^\nu} \theta^\alpha \theta^\beta \theta^\gamma \sigma^\mu_{\beta\dot{\alpha}} \sigma^\nu_{\gamma\dot{\beta}} \bar{\theta}^{\dot{\alpha}} \bar{\theta}^{\dot{\beta}}$$

only one way to contract

$$\theta^2 \partial^\mu \chi^\alpha \sigma^\mu_{\alpha\dot{\beta}} \bar{\theta}^{\dot{\beta}}$$

$$\Rightarrow F(\gamma) \theta^2 = F(x) \theta^2$$

Last time: Superfields, supermultiplets, and interactions in 4d $\mathcal{N}=1$ SUSY. Super space: $\mathbb{R}^{1,3} \oplus \Pi S_R = \mathbb{R}^{1,3|4}$

Two ways of getting irreducible supermultiplets

- Impose a chirality constraint $\bar{D}\Phi = 0$

\Rightarrow a matter multiplet:

$$\bar{D}_\alpha \Phi = 0$$

1 complex scalar

1 Weyl fermion

1 complex auxiliary field

$$\Phi(\gamma, \theta) = \phi(\gamma) + \theta^\alpha \chi_\alpha(\gamma) + F(\gamma) \theta^2$$

- wait to also get a multiplet w/ one vector, one Majorana fermion

$$Q_\alpha = \frac{\partial}{\partial \theta^\alpha} + \theta^\beta \sigma^\mu_{\alpha\dot{\beta}} \frac{\partial}{\partial x^\mu}$$

$$(D_\alpha = -\frac{\partial}{\partial \theta^\alpha} + \theta^\beta \sigma^\mu_{\alpha\dot{\beta}} \frac{\partial}{\partial x^\mu})$$

Inside the real superfield:

power of θ

content

0

\mathbb{R} scalar

1

\mathbb{R} fermion (Majorana)

2

\mathbb{R} vector fields or complex scalar

3

Majorana fermion

4

\mathbb{R} scalar

"W-Z gauge"

Φ chiral $\sim i(\bar{\Phi} - \Phi)$

is not chiral.

$\begin{cases} \text{Im } \Phi \\ \Phi + \bar{\Phi} \\ \text{Re } \Phi \end{cases}$

Expansion of the chiral

superfield: Take Taylor series of $\Phi(y) \rightarrow \Phi(x)$

$$\begin{aligned} & \Phi(x) \\ & \theta^2 F(x) + \theta^\alpha \bar{\theta}^{\dot{\beta}} \partial_{\alpha\dot{\beta}} \Phi(x) + \frac{1}{2} \theta^2 \bar{\theta}^2 \partial^2 \Phi(x) \\ & \text{and supercovariates} \end{aligned}$$

$$\phi(x) + \bar{\phi}(x) \in \text{real}$$

$$\begin{aligned} & \theta^4 \quad \bar{\theta}^4 \\ & \theta^2 F \quad \theta^\alpha \bar{\theta}^{\dot{\beta}} (\partial_{\alpha\dot{\beta}} (\phi - \bar{\phi})) \quad \bar{\theta}^2 F \\ & \theta^2 \bar{\theta}^2 \partial^2 \phi \quad \theta^2 \bar{\theta}^2 \partial^2 \bar{\phi} \\ & \theta^2 \bar{\theta}^2 (\partial^2 (\phi + \bar{\phi})) \end{aligned}$$

Wess-Zumino

due to transform this

then

$$V \sim V + (\Phi + \bar{\Phi})$$

real superfield $V: \mathbb{R}^{3|4} \rightarrow \mathbb{R}$

Discussion:

In any gauge theory, there's a current

$$J = \frac{\delta Z}{\delta A}$$

$A \leftarrow$ gauge field
minimal coupling

what's the super version? $\sim A \rightarrow V$ real superfields \in

$\nabla^2 \phi = \bar{\nabla}^2 \bar{\phi} = 0$ give "correct type" multiplets

$$\mathcal{L} \sim A_{\mu} J^{\mu}$$

$$\partial_{\mu} J^{\mu} = 0$$

The simplest interacting theory is the Wess-Zumino model: chiral superfields + superpotential interaction

$$D\text{-term: } \int d^4x d^2\theta d^2\bar{\theta} K(\Phi, \bar{\Phi}), \text{ simplest } K = \bar{\Phi}\Phi$$

vector multiplet

on shell | off shell

A_{μ}	2	4
λ	2	4
D	0	1
Imp	0	-1

!

$$\text{Use } \sim \theta^2 \bar{\theta}^{\dot{\beta}} \partial_{\alpha\dot{\beta}} \psi_{\alpha}(x) (\sigma^{\mu} \bar{\varphi}_{\dot{\alpha}}(x)) \leftarrow \text{Dirac terms}$$

$$\sim \int d^4x (\bar{\phi} \partial^2 \phi + \bar{\psi} \not{\partial} \psi + F \bar{F})$$

Mass is not here \nearrow not required by SUSY. To get mass, get F-term

$$F\text{-term: } \int d^4x d^2\theta W(\Phi) + \text{h.c.}, \text{ simplest } W = \Phi^2$$

$\left. \begin{matrix} \text{chiral} \\ \text{chiral} \end{matrix} \right\}$ need

$$\text{look } \sim \int d^4x d^2\theta \psi^2 + F\phi + \text{h.c.}$$

$$\mathcal{L} \sim (F\bar{F} + F\phi + \bar{F}\bar{\phi})$$

\downarrow mass for F is auxiliary. EOM $\sim F = -\bar{\phi}$

\leadsto generate the mass term. $\phi\bar{\phi}$ after integrating out F.

using super fields

Two types of superfields: - chiral (cpx w/a constraint)

- Vector (real w/a gauge invariance)

Two types of interactions terms: + F-terms (integrate w/a chiral over

$$\int d^3\theta + h.c.$$

- D-terms (integrate $\int d^2\theta d^2\bar{\theta}$)

$$\Phi^i \rightarrow \Phi^i + \delta\Phi^i$$

$$R: \Phi \rightarrow R \Phi$$

$$W: \Phi \rightarrow W \Phi$$

$$\Phi \rightarrow \Phi + \delta\Phi$$

$$\Phi \rightarrow \Phi + \delta\Phi$$

$$\Phi \rightarrow \Phi + \delta\Phi$$

$$\Phi \rightarrow \Phi + \delta\Phi$$

$$\Phi \rightarrow \Phi + \delta\Phi$$

$$\Phi \rightarrow \Phi + \delta\Phi$$

$$\Phi \rightarrow \Phi + \delta\Phi$$

$$\Phi \rightarrow \Phi + \delta\Phi$$

$$\Phi \rightarrow \Phi + \delta\Phi$$

$$\Phi \rightarrow \Phi + \delta\Phi$$

$$\Phi \rightarrow \Phi + \delta\Phi$$

$$\Phi \rightarrow \Phi + \delta\Phi$$

$$\Phi \rightarrow \Phi + \delta\Phi$$

$$\Phi \rightarrow \Phi + \delta\Phi$$

$$\Phi \rightarrow \Phi + \delta\Phi$$

$$\Phi \rightarrow \Phi + \delta\Phi$$

$$\Phi \rightarrow \Phi + \delta\Phi$$

$$\Phi \rightarrow \Phi + \delta\Phi$$

$$\Phi \rightarrow \Phi + \delta\Phi$$

$$\Phi \rightarrow \Phi + \delta\Phi$$

$$\Phi \rightarrow \Phi + \delta\Phi$$

$$\Phi \rightarrow \Phi + \delta\Phi$$

$$\Phi \rightarrow \Phi + \delta\Phi$$

$$Z = K(\Phi, \bar{\Phi})|_D + W(\Phi)|_F$$

$$K = \Phi^\dagger \Phi = \text{quadratic}$$

$$W = \Phi^\dagger \Phi = \text{quadratic}$$

$$W = \Phi^\dagger \Phi = \text{quadratic}$$

$$W = \Phi^\dagger \Phi = \text{quadratic}$$

$$W = \Phi^\dagger \Phi = \text{quadratic}$$

$$W = \Phi^\dagger \Phi = \text{quadratic}$$

$$W = \Phi^\dagger \Phi = \text{quadratic}$$

$$W = \Phi^\dagger \Phi = \text{quadratic}$$

$$W = \Phi^\dagger \Phi = \text{quadratic}$$

$$W = \Phi^\dagger \Phi = \text{quadratic}$$

$$W = \Phi^\dagger \Phi = \text{quadratic}$$

$$W = \Phi^\dagger \Phi = \text{quadratic}$$

$$W = \Phi^\dagger \Phi = \text{quadratic}$$

$$W = \Phi^\dagger \Phi = \text{quadratic}$$

$$W = \Phi^\dagger \Phi = \text{quadratic}$$

$$W = \Phi^\dagger \Phi = \text{quadratic}$$

$$W = \Phi^\dagger \Phi = \text{quadratic}$$

$$W = \Phi^\dagger \Phi = \text{quadratic}$$

$$W = \Phi^\dagger \Phi = \text{quadratic}$$

$$W = \Phi^\dagger \Phi = \text{quadratic}$$

$$W = \Phi^\dagger \Phi = \text{quadratic}$$

$$W = \Phi^\dagger \Phi = \text{quadratic}$$

$$W = \Phi^\dagger \Phi = \text{quadratic}$$

over

chiral

over

chiral

over

chiral

over

chiral

over

chiral

over

chiral

over

chiral

over

chiral

over

chiral

over

chiral

over

chiral

over

chiral

over

chiral

over

chiral

over

chiral

over

chiral

over

EOM: $\frac{\delta \mathcal{L}}{\delta \bar{\Phi}^i} = g_{i\bar{j}} \bar{F}^{\bar{j}} + \frac{\partial g_{i\bar{k}}}{\partial \bar{\Phi}^{\bar{k}}} \bar{\Phi}^{\bar{k}} \bar{F}^{\bar{j}} = 0 \Rightarrow \bar{F}^{\bar{j}} = -g^{\bar{j}\bar{k}} \frac{\partial g_{i\bar{k}}}{\partial \bar{\Phi}^{\bar{k}}} \bar{\Phi}^{\bar{k}} \bar{F}^{\bar{j}}$

$\frac{\partial^2 \mathcal{L}}{\partial \bar{\Phi}^i \partial \bar{\Phi}^{\bar{j}}} \bar{\Phi}^{\bar{j}} = \frac{\partial^2 g_{i\bar{k}}}{\partial \bar{\Phi}^{\bar{k}} \partial \bar{\Phi}^{\bar{j}}} \bar{\Phi}^{\bar{j}}$

$\frac{\partial g_{i\bar{k}}}{\partial \bar{\Phi}^{\bar{k}}} = \frac{\partial^2 \mathcal{L}}{\partial \bar{\Phi}^{\bar{k}} \partial \bar{\Phi}^{\bar{j}}} \bar{\Phi}^{\bar{j}} = \frac{\partial g_{k\bar{j}}}{\partial \bar{\Phi}^{\bar{j}}}$ (Kähler identity)

Also get terms of the form: $\left(\frac{\partial^2 \mathcal{L}}{\partial \bar{\Phi}^{\bar{k}} \partial \bar{\Phi}^{\bar{j}}} \bar{\Phi}^{\bar{k}} \bar{\Phi}^{\bar{j}} \right) \left(\Gamma_{mn}^i \psi^m \psi^n \right) g_{i\bar{j}}$

Combine to produce 4-fermi interactions of the form

$R_{i\bar{j}k\bar{l}} \psi^i \psi^k \bar{\psi}^{\bar{j}} \bar{\psi}^{\bar{l}}$

R-symmetry $\mathcal{L} = \frac{1}{2} (\dot{\phi}^2 - \phi^2)$

what about going to gauge fields?

$V: \mathbb{R}^{1,3,4} \rightarrow \mathbb{R}$

$V \sim V + (\Psi + \bar{\Psi})$

Chiral: $\Psi \xrightarrow{e} \Psi$
 $\bar{\Psi} \xrightarrow{e} \bar{\Psi}$
 $\mathcal{D}_\mu \Psi = 0$

Ansatz: $\Phi \rightarrow e^{i\Psi} \bar{\Phi}$

Problem:

$K: \bar{\Phi} \Phi \rightarrow \bar{\Phi} e^{i\Psi} e^{i\Psi} \Phi$

Solution:

$\bar{\Phi} e^{-i\Psi} \Phi = \bar{\Phi} \Phi - i \bar{\Phi} \Psi \Phi + \frac{1}{2} \bar{\Phi} \Psi^2 \Phi$

in WZ gauge stops at V^2

More on R-symmetry, holomorphy and non-renormalization theorems

- ↳ A symmetry that acts nontrivially on the supercharges (acts on the vector space)
- Doesn't commute with supersymmetry
- but is an automorphism of super transformations.
- (similar to Lorentz group but it doesn't act in the bosonic spacetime)

In the context of 4-d, $\mathcal{N}=1$ theories, R-symmetry is $U(1)$

$Q_\alpha, \bar{Q}_{\dot{\alpha}}, P_{\alpha\dot{\alpha}} \quad \{Q_\alpha, \bar{Q}_{\dot{\alpha}}\} = P_{\alpha\dot{\alpha}}$

Charge R-charge

$+1 \quad -1 \quad 0$
 $\downarrow \quad \downarrow \quad \downarrow$
 $e^{iQ_\alpha} \quad e^{-i\bar{Q}_{\dot{\alpha}}} \quad e^{iP_{\alpha\dot{\alpha}}}$

without breaking susy

- R-symmetry may or not may be broken in any given theory.
- Determines R-charges of components but not homogeneously.

Ex: Chiral Superfield $\Phi(x, \theta, \bar{\theta}) = \phi(x) + \theta^\alpha \psi_\alpha(x) + \theta^2 F(x) + \dots$

3-1-1

R-change
Flavour-change

Imaging promoting

Another way to see this case is by symmetry.


—probably with a pipe.

$$\frac{K}{w} = 544 \text{ m}$$

$$\Phi \sim K \Phi$$

The 10-10-10 rule

Wave function renormalization (Kähler Potential)



That's one way to organize these...

Not homogeneous \Rightarrow a \neq b symmetric

Recalling that the HW is considered as

In a more general theory $w(\vec{r}, \vec{r}') = e^{i\vec{k} \cdot (\vec{r} - \vec{r}')} = e^{i\vec{k} \cdot \vec{r}} e^{-i\vec{k} \cdot \vec{r}'}$ for some change of variables $\vec{r} \rightarrow \vec{r}'$ w/ unitary symmetry.

Using Eqn of F_y $F = \frac{\partial W}{\partial \delta}$

\downarrow
 $r = -2 \Rightarrow r = -\frac{2}{2}$
 \downarrow impossible! \rightarrow sum of other

$\exists M(\Phi) \in$ must be a monomial (If R-symmetry is broken)


In a theory of 16mm s.p. field

[illegible]

	R charge	Flavour charge	
m	$-2-2r$	-2	
$(W, \frac{1}{2})$	-2	0	$\rightarrow W_{eff} = \frac{1}{2} m \Phi^2 \delta(\frac{\lambda \Phi}{m})$
λ	$-2-3r$	-3	
$\lambda \Phi / m$	0	0	$= W!$

holomorphic function.

One major consequence of not renormalizing for W is the existence of moduli spaces of vacuum in supersymmetric theories.



In S-Sy it doesn't happen: $W(\Phi_i) \rightarrow V(\phi_i) = \sum_i \left| \frac{\partial W}{\partial \phi_i} \right|^2$
 zero iff $\frac{\partial W}{\partial \phi_i} = 0 \Rightarrow$ sits at a critical point in fields space. $\hookrightarrow \text{vacua} = \text{crit}(W)$

$$W(\Phi_1, \Phi_2) = \frac{1}{2} m \Phi_2^2 + \frac{1}{2} \lambda \Phi_1^2 \Phi_2^2$$

\uparrow
 $2r_2 = -2$
 $\boxed{r_2 = -1}$

\uparrow
 $2r_1 + r_2 = -2$
 $2r_1 - 1 = -2 \rightarrow \boxed{r_1 = -\frac{1}{2}}$
 $\hookrightarrow \text{good } \checkmark$

$\hookrightarrow \text{crit?}$
 $\frac{\partial W}{\partial \Phi_2} = m \Phi_2 + \lambda \Phi_1^2 = 0$
 $\frac{\partial W}{\partial \Phi_1} = \lambda \Phi_1 \Phi_2^2 = 0$
 \downarrow
 only at $\Phi_1 = \Phi_2 = 0$


but also if $m \rightarrow 0$, $\begin{cases} \Phi_1 = 0 \\ \Phi_2 = \text{anything} \end{cases}$

Lecture 11.2


Recall Z_m and Z_1 are not indep. \hookrightarrow continuity may vacuum!

$\hookrightarrow \mathcal{L} \sim \bar{\Phi} \Phi|_D + W(\Phi)|_F \sim \Phi \rightarrow \tilde{z}^{\frac{1}{2}} \Phi$
 $\left\{ \begin{array}{l} z \bar{\Phi} \Phi|_D \\ z^{\frac{1}{2}} \Phi|_F \end{array} \right. \sim z^{\frac{1}{2}} \Phi$

For m depends on Φ and Φ are not independent



non perturbative corrections can (and do) occur
Ex: SQM (N=2) w/ target space $\mathbb{C}^2 \setminus \{0\}$
 $W = \phi^3 \rightarrow V \sim \left| \frac{\partial W}{\partial \phi} \right|^2 \sim \phi^4$



$$\hookrightarrow \phi^4 - m^2 \phi^2 + c$$

$$\left(\phi^2 - \frac{m^2}{2} \right)^2$$

$$\left(\frac{\partial W}{\partial \phi} \right)^2$$

Action for Super QED

Remember, V is a gauge field $\rightarrow V_\mu \Phi \sim e^{i\alpha} \Phi$ then $\bar{\Phi} \Phi|_D$ is not gauge invariant

transformations

then our naive guess

remember

$$W_2 = -\frac{1}{2} \bar{D}^2 D^2 V$$

W is: supergauge inv

- chiral: $D^2 W_2 = 0$

- fermion

$\bar{D}^2 D^2 V \rightarrow W$

fermion and W

want to write: $W^a W_a + h.c.$

V is $W=2$ gauge

$$V = \partial^a \bar{\partial}_a A(x)(y) + \partial^2 \bar{\partial}^2 \lambda(x)(y) + \partial^2 \bar{\partial}^2 \lambda(x)(y)$$

$$D_x = \frac{\partial}{\partial x} - 2 \bar{\partial}^2 \partial$$

$$W(x)(y) = \lambda(x)(y) + \frac{1}{2} \partial^2 D + \partial^2 F + \frac{1}{2} \partial^2 \bar{\partial}^2 \lambda(x)(y)$$

$$F \rightarrow (F_{\alpha\beta}, F_{\dot{\alpha}\dot{\beta}})$$

$$F_{\alpha\beta} F^{\alpha\beta} = F_{\alpha\beta} F^{\alpha\beta} + \partial_{\alpha\beta} A^{\alpha\beta} + \partial_{\dot{\alpha}\dot{\beta}} A^{\dot{\alpha}\dot{\beta}}$$

where action $F_{\alpha\beta} F^{\alpha\beta} = F_{\alpha\beta} F^{\alpha\beta}$

$F_{\alpha\beta} = \partial_{\alpha\beta} A^{\alpha\beta}$

$F_{\alpha\beta} = \partial_{\alpha\beta} A^{\alpha\beta}$

$F_{\alpha\beta} = \partial_{\alpha\beta} A^{\alpha\beta}$

$F_{\alpha\beta} = \partial_{\alpha\beta} A^{\alpha\beta}$

$F_{\alpha\beta} = \partial_{\alpha\beta} A^{\alpha\beta}$

$F_{\alpha\beta} = \partial_{\alpha\beta} A^{\alpha\beta}$

$F_{\alpha\beta} = \partial_{\alpha\beta} A^{\alpha\beta}$

$F_{\alpha\beta} = \partial_{\alpha\beta} A^{\alpha\beta}$

$F_{\alpha\beta} = \partial_{\alpha\beta} A^{\alpha\beta}$

$F_{\alpha\beta} = \partial_{\alpha\beta} A^{\alpha\beta}$

$F_{\alpha\beta} = \partial_{\alpha\beta} A^{\alpha\beta}$

$F_{\alpha\beta} = \partial_{\alpha\beta} A^{\alpha\beta}$

$F_{\alpha\beta} = \partial_{\alpha\beta} A^{\alpha\beta}$

$F_{\alpha\beta} = \partial_{\alpha\beta} A^{\alpha\beta}$

$F_{\alpha\beta} = \partial_{\alpha\beta} A^{\alpha\beta}$

$F_{\alpha\beta} = \partial_{\alpha\beta} A^{\alpha\beta}$

$F_{\alpha\beta} = \partial_{\alpha\beta} A^{\alpha\beta}$

Terms involving D : $\frac{1}{2} D^2 + \sum D \Phi_i \bar{\Phi}_i + \xi D =$

now $D = \left(\sum q_i |\Phi_i|^2 + \xi \right)$

brings

$\begin{aligned} \text{Eqn} \\ \frac{1}{2} D^2 + cD \\ \Rightarrow D = -c \\ \Rightarrow \frac{1}{2} D^2 + cD = -\frac{1}{2} c^2 \end{aligned}$

Lecture 2.1

Supersymmetric gauge theories:

- Last time \leadsto Abelian susy gauge theory

V : real superfield (vector multiplet) $\leadsto A, (\lambda, \bar{\lambda}), D$

$\{\bar{\Phi}_i, \Phi_i\}$ charged chiral multiplets $\leadsto \varphi_i, \psi_i, F_i$

Weyl fermions

g : electro magnetic coupling constant

J : Fayet-Iliopoulos parameter

$W(\Phi_i)$: gauge inv. holomorphic superpotential

$$\mathcal{L} = \sum_i \underbrace{\bar{\Phi}_i e^{2q_i V} \Phi_i}_{\text{Scalar kinetic terms}} + \underbrace{J V}_{\text{Fayet-Iliopoulos terms}} + \underbrace{W^X W_X}_{\text{"Super Maxwell" term}} + \text{h.c.} + \underbrace{W(\Phi_i)}_{\text{super potential interactions}}$$

Since we have chiral fermions coupled to gauge fields, we


should also check that the theory is free of gauge anomalies

anomalies $\Rightarrow \partial_\mu J^\mu \neq 0$ in the QFT theory.

What sort of diagram would I imagine I have to compute?

Generally for the operator J_μ : $\mathcal{L} = A_\mu J^\mu + \dots$, so looking at

$$\text{"problematic terms"} \quad J = \sum_i q_i \bar{\psi}_i \gamma_\mu \psi_i + \dots \quad (\text{minimally coupled})$$

\Rightarrow 

$\langle \partial_\mu J^\mu \rangle = \langle \text{tr} F \wedge F \rangle \sim q_i \text{tr} F \wedge F$

$J = \frac{\delta \mathcal{L}}{\delta A} = 3\text{-form}$

$\Rightarrow dJ = 4\text{-form}$

There is also a "gravitational anomaly" $\frac{\delta \mathcal{L}}{\delta g_{\mu\nu}} = T_{\mu\nu}$

the gravitational anomaly only vanishes iff $\sum q_i = 0$

An obvious nontrivial solution to both anomalies:

chiral fields in pairs (Φ_i^+, Φ_i^-) with charge $\pm q_i$ \leadsto "super QED"

What about non-abelian gauge theories? Pick a gauge group

$\phi \pm = \phi \pm + \frac{1}{2} \int$
 $\phi \pm = \phi \pm + \frac{1}{2} \int$
 parameter
 we can think of this as a gauge inv.

To fix gauge 1 cond imposed
 $(r, \theta + \chi, \theta - \chi)$
 $(r, \theta + \chi, \theta - \chi)$
 different charges

$\nabla(\Phi) = \frac{1}{2} D^2 = \frac{1}{2} (|\phi_+|^2 - |\phi_-|^2 + \int)$
 $\nabla(\Phi) = \frac{1}{2} D^2 = \frac{1}{2} (|\phi_+|^2 - |\phi_-|^2 + \int)$
 $\nabla(\Phi) = \frac{1}{2} D^2 = \frac{1}{2} (|\phi_+|^2 - |\phi_-|^2 + \int)$

Let's consider $S(\Phi \pm)$ with fields $\Phi \pm$ of charge ± 1
 $\frac{1}{2} D^2 + \int D + \int \phi_+ - \int \phi_-$
 EOM for D : $D = -(\int + |\phi_+|^2 - |\phi_-|^2)$

do I got a gauge inv. term from (4) like that?
 (Just trying stuff with work)
 we go to $w=2$ gauge, as before.

$e = e^{V - (X + \frac{1}{2}) - \frac{1}{2} [V, X] - \frac{1}{2} [X, V] + \dots}$

Note: we have commutators in the group algebras, so this time
 commutators don't vanish, we just choose to work in linear order

to find the form of Φ we use BCH
 $\Phi = e^{V - X - \frac{1}{2} [V, X] - \frac{1}{2} [X, V] + \dots}$
 $\Phi = e^{V - X - \frac{1}{2} [V, X] - \frac{1}{2} [X, V] + \dots}$
 $\Phi = e^{V - X - \frac{1}{2} [V, X] - \frac{1}{2} [X, V] + \dots}$

we want the covariant kinetic terms
 to make sense.
 Generally: $A_\mu \in \mathfrak{g}$
 infinitesimally: $A \rightarrow A - d\lambda + [A, \lambda]$
 finitely: $A \rightarrow U^{-1} A U + U^{-1} dU$
 algebraically: $0 + e^{\lambda} d\lambda e^{\lambda}$

Lecture 12 Moduli spaces of vacua (mostly in gauge theory)

Example: $N=1$ SQED with one flavour
(abelian gauge theory) (Φ_{\pm} chiral multiplet $q_{\pm} = \pm 1$)


$$\text{FI parameter } \sum q_i = 0$$

$$\sum q_i^3 = 0$$


If $W=0 \Rightarrow$ potential for scalars comes from D-terms. $\mathcal{L} \sim \frac{1}{2} D^2 + D \Phi_+ \phi_+ - D \bar{\Phi}_- \phi_- + \dots \Rightarrow D = \bar{\Phi}_- \phi_- - \bar{\Phi}_+ \phi_+ + \dots$

Before taking quotient by gauge symmetries, (ϕ_+, ϕ_-)
 $\mathcal{F}^2 \geq \{D=0\}$

$$\mathcal{M} = \{D=0\} / U(1) = \mathbb{C}^2 / \mathbb{Z}_2 =$$



if $S \neq 0$
 $\Rightarrow \phi_+ = \phi_- = 0$
is not a vacuum
 \Rightarrow gauge symmetry is broken



Recall: $\{D=0\} = \{(\phi_+, \phi_-) \mid |\phi_+|^2 = |\phi_-|^2\}$, parametrize $\phi_+ = r e^{i\theta_+}$
 $\phi_- = r e^{i\theta_-}$

photon mass

$$\bar{\Phi}_+ e^V \Phi_+ + \bar{\Phi}_- e^{-V} \Phi_- \supset (\bar{\Phi}_+ \phi_+ + \bar{\Phi}_- \phi_-) A^\mu A_\mu$$

$$\frac{\partial_+ + \partial_-}{2}$$

$$= 2|\phi_+|^2 + \dots \Rightarrow m_\gamma \sim \sqrt{V|\phi_+|^2 + \dots}$$

(potentially singular)

\Rightarrow The moduli space is always a \mathbb{C}^n manifold (parametrized by chiral superfields)

\Rightarrow In fact, it always has a Kähler metric! (D-terms of the massless scalars are the Kähler potential)

$$K = \bar{\Phi}_+ \Phi_+ + \bar{\Phi}_- \Phi_- = 2|\phi_+|^2 \quad (\text{if D-term condition is satisfied})$$

on the space $\{D=0\}$

Rewrite in terms of gauge inv. observables.

$$\phi_+ \phi_- = M \Rightarrow M \bar{M} = \bar{\Phi}_+ \phi_- \phi_+ \phi_- = \left(\frac{K}{2}\right)^2 \Rightarrow K = 2\sqrt{M \bar{M}}$$

$$g_{M \bar{M}} = \frac{\partial^2 K}{\partial M \partial \bar{M}} = \frac{\partial}{\partial M} \frac{2\sqrt{M \bar{M}}}{\partial \bar{M}} = \frac{1}{2} \frac{M \bar{M}}{(M \bar{M})^{3/2}} + \left(\frac{1}{M \bar{M}}\right)^{1/2} = \frac{1}{2} \frac{1}{(M \bar{M})^{1/2}}$$

In the low energy limit "deep IR", only massless d.o.f. are relevant

Examination 14

a) $M = \{v \in V : D(v) = 0\} / u(1) = V / \Phi^X$

$$\underbrace{|\psi_+\rangle}_{v_1}^2 - \underbrace{|\psi_-\rangle}_{v_2}^2 = r$$

D-term

$$M = \{v \in \Phi^2 \mid |v_1|^2 - |v_2|^2 = r\} / u(1)$$

$r \in \mathbb{R}$

show that $M \cong (\Phi^2 \setminus \Delta_r) / \Phi^X$

GIT

for $r > 0$ $\Delta_r \subset \Phi^2$ set of elements not containing solution to D-term eqn

$$\Delta_r = \{(0, v_2) \in \Phi^2\}$$

let $v \in \Phi^2 \setminus \Delta_r$ arbitrary. Claim: $\Phi^X v$ contains exactly one solution to D-term eqn.

consider cv , $c \in \Phi^X$, plug into D-term eqn.

$$|cv_1|^2 - |c^{-1}v_2|^2 = |c|^2 |v_1|^2 - |c|^{-2} |v_2|^2 = r$$

$$\leadsto |c|^4 |v_1|^2 - r |c|^2 - |v_2|^2 = 0$$

$$|c|^2 = \frac{r}{|v_1|^2} \pm \sqrt{\frac{r^2}{4|v_1|^2} + \frac{|v_2|^2}{|v_1|^2}}$$

only take \pm s.t. $|c|^2 > 0$

Always has a real solution for $v \in \Phi^2 \setminus \Delta_r$ which fixes $|c|^2$
 $\Rightarrow c$ up to a phase $\Rightarrow M \cong (\Phi^2 \setminus \Delta_r) / \Phi^X$

its analogous for $r < 0$ $|c|^2 = \frac{-r}{|v_1|^2} \pm \sqrt{\frac{r^2}{4|v_1|^2} + \frac{|v_2|^2}{|v_1|^2}}$

for $r = 0$ $\{v \in \Phi^2 \mid |v_1|^2 - |v_2|^2 = 0\} / u(1) = \Phi^2 / \Phi^X$

$$\Phi^2 / \Phi^X \ni [v] = \{cv = (cv_1, c^{-1}v_2) \mid c \in \Phi^X\}$$

\downarrow
 $\Phi^2 / \Phi^X \rightarrow \mathbb{C}^* / \mathbb{Z}_2$ look for $(c, \tilde{v}) \in \Phi^X \times \Phi$ s.t. $\left. \begin{aligned} cv_1 &= \tilde{v} \\ c^{-1}v_2 &= \tilde{v} \end{aligned} \right\} \frac{v_2}{v_1} = c^2$

$$\Rightarrow \tilde{v} = \pm \sqrt{\frac{v_2}{v_1}} v_1 \leadsto \text{act } [v] \text{ rotates elements } (\pm \tilde{v}, \pm \tilde{v})$$

$\Rightarrow \Phi^2 / \Phi^X \xrightarrow{\sim} \mathbb{C}^* / \mathbb{Z}_2$, $[v]_{\mathbb{C}^*} \rightarrow [\tilde{v}]_{\mathbb{Z}_2}$ (group $(\mathbb{C}^*) / \mathbb{Z}_2$ is not well defined)

b) $D = |x_+|^2 - |x_-|^2 + 2|x_+|^2 - 2|y_-|^2 = 0$

look for LC_2 above the size of LC_1

c) Extrakt of invariant motoric IC :

$$A_1 = X_+ X_- , A_2 = Y_+ Y_- , B_1 = Y_+ X_- , B_2 = Y_- X_+ X_+ \\ \text{next step } (A_2 A_1 = B_1 B_2) \Rightarrow \dim M = \# \text{vars} - \# \text{constraints} = 3$$

d) Group linear Sigma models:

[illegible]

stands by

$$(A \cdot B) \cdot C = (A \cdot (B \cdot C)) \quad \text{Associativity}$$

Choose basis $e_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, e_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ of \mathbb{C}^2 or \mathbb{R}^2 so $V \oplus W = \mathbb{C}^2$ or \mathbb{R}^2 induces a basis of $\{e_1, e_2\} \in \mathbb{C}^2$ or \mathbb{R}^2 the complexified space then $V \oplus W = \mathbb{C}^2$ or \mathbb{R}^2

$$(h\phi)_\lambda = (h'(\phi_\lambda))_\lambda \quad \begin{cases} \wedge^1 \phi + \phi = \phi \\ \wedge^1 h + h = h \end{cases} \quad \phi \wedge \phi = \phi \quad h \wedge h = h$$

$$(k, \phi) = \begin{matrix} \text{state } 1 \\ \text{state } 2 \\ \text{state } 3 \end{matrix} \quad (k, \phi) = \begin{matrix} \text{state } 1 \\ \text{state } 2 \\ \text{state } 3 \end{matrix} \quad (k, \phi) = \begin{matrix} \text{state } 1 \\ \text{state } 2 \\ \text{state } 3 \end{matrix}$$

$$h(\phi) = \phi^2 \quad \text{and} \quad h(\phi^2) = \phi^4$$

$$(\phi f)_\omega = \int_{\mathbb{R}^n} \phi(x) f(x) dx = \int_{\mathbb{R}^n} \phi(x) \chi_{[0,1]}(x) dx = \int_0^1 \phi(x) dx = S$$



Wahl $N = (3, 2)$: $S = \phi \oplus (R_2^A, \delta) \oplus (R_2^V, \delta)$

✓ R-symmetry $SO(B)^A \times SO(C)^A \simeq U(1)^A \times U(1)^A$


$$\begin{aligned} \wedge^2 \mathcal{O}_X &= \mathcal{O}_X \\ \wedge^1 \mathcal{O}_X &= \mathcal{O}_X \\ \wedge^0 \mathcal{O}_X &= \mathcal{O}_X \end{aligned}$$

Lecture: One loop corrections to the Kähler potential
 (D-term for chiral superfields = metric on moduli spaces)


$$W = \frac{1}{2} X^2 Y$$




$\bar{Y}Y|_D = K(\bar{Y}, Y) \rightarrow \frac{1}{2} X^2 Y + \frac{1}{2} X^2 Y_0$
 $\sim \int \frac{d^3 p}{(p^2 + m_X^2)^2} \sim \log(\frac{\Lambda}{Y_0})$
 $\frac{\partial W}{\partial X} \sim \frac{1}{2} Y_0 X^2$
 $(X=0, Y=Y_0)$



Singularity



Singularities add a degree of freedom.



Moduli spaces in QCD: (nonabelian gauge theory)

$$\frac{\{D=0\}}{G} = \frac{\text{no D-term}}{G_\Phi}$$

Ex: $G = U(1), G_\Phi = \mathbb{C}^X$

$$\mathbb{C}^2 / \mathbb{C}^X = \frac{\{D=0\}}{U(1)}$$

- Pick a gauge group $G = SU(N_c), N_c = \text{"number of colours"}$
 $g = \text{adj rep}, \dim(g) = N_c^2 - 1$
 $G = \mathbb{C}^{N_c} \sim G \& G C$

Vector multiplet valued in g : $(A_\mu^a, \lambda^a, D^a)$


- Matter: $V = \text{some other rep of } G$. Chiral multiplets valued in V .
 (Ex: $V = \mathbb{C}$ fundamental rep.)

- To guarantee anomaly cancellation, take reps + conjugate reps (e.g. $\mathbb{C} \oplus \bar{\mathbb{C}}$). sufficient but not necessary.

SQCD in the literature means " N_f fundamental flavours"
 number of flavours

Flavor space $F = \mathbb{C}^{N_f}$

\Rightarrow matter fields are: $\int \Phi_+ \in F \otimes \mathbb{C}$
 $\int \bar{\Phi}_- \in \bar{F} \otimes \bar{\mathbb{C}}$



$$\begin{array}{c} \bar{\Phi}_+ \bar{e}^{-\nu} \Phi_+ + \dots \\ \downarrow \quad \downarrow \quad \downarrow \\ \bar{F} \otimes \bar{C} \oplus F \otimes C \end{array}$$

$$\begin{array}{l} \exp \quad \Phi \rightarrow e^X \\ R \xrightarrow{\exp} R^* \text{ (non compact)} \\ i(R \xrightarrow{\exp} U(1)) \text{ (compact)} \end{array} \quad \begin{array}{l} \alpha \rightarrow \exp(\alpha) = e^\alpha \\ G = \Phi^X \end{array}$$

$$\begin{array}{l} \Phi^X = GL(N, \mathbb{C}) \supseteq U(1) \\ \hookrightarrow GL(N, \mathbb{R})_+ \\ GL(N, \mathbb{C}) \xrightarrow{U(1)} GL(N, \mathbb{R}) \end{array}$$

Re take vevs, $2N_c N_f \rightarrow (F \otimes C) \oplus (\bar{F} \otimes \bar{C}) / SLCC$
 In general I break $SU(N_c)$ to $SU(N_c - N_f)$

And (more) on moduli spaces is "super-QCD"

$V = g$ -gauge field
 $G = SU(N_c)$, $C = \Phi^{N_c}$, $N_f = \text{"Flavours"}$, $F = \Phi^{N_f}$

$$\begin{cases} \bar{\Phi}_+ \in C \\ \bar{\Phi}_- \in \bar{C} \end{cases}$$

$$\leadsto \begin{cases} \Phi_+ \in C \otimes F \\ \Phi_- \in \bar{C} \otimes \bar{F} \end{cases}$$

To compute the moduli space: space of possible VEVs = $C^{2N_c N_f} = (C \otimes F) \oplus (\bar{C} \otimes \bar{F})$
 $\begin{cases} \text{impose D-term conditions} \\ \text{divide out the action of the gauge group } SU(N_c) \end{cases}$
 or, divide out by $G_C = SL(N_c)$ (A)

The phase space of a (classical) dynamical system is a symplectic manifold
 symplectics act by $G \rightarrow \text{Symp}(\mathbb{R}^{2n}, \omega)$

different form $\leadsto g \rightarrow \text{Vect}^{\text{symp}}(\mathbb{R}^{2n}, \omega)$
 \hookrightarrow symplectic vector fields $\left(\begin{array}{l} X \in \text{Vect}(\mathbb{R}^{2n}) \\ \text{is symplectic when} \\ \mathcal{L}_X \omega = 0 \end{array} \right)$

\mathbb{R}^n $i_X \omega$ is closed $\iff X$ is symplectic
 If $i_X \omega = \text{exact} = d\lambda \leadsto X$ is "Hamiltonian" $d i_X \omega + i_X d\omega = 0$ exact is ~~not~~ symplectic

here $g \rightarrow C^\infty(\mathbb{R}^{2n}, \omega)$ is the moment map assign the generator to the observable i.e.,
 (Poisson algebra) Generator of translations \rightarrow momentum

moment map allows us to kill things
 in both ways, symplectic and observables

$\begin{array}{c} T^*S^1 \\ \hookrightarrow (S^1, \theta) \end{array}$ $p \in C^\infty(\mathbb{R})$
 $\omega = d\theta \otimes dp_0$
 $X = \frac{\partial}{\partial p_0} \leadsto i_X \omega = d\theta$
 $\leadsto d\theta = d\lambda$
 Example of symplectic but not hamiltonian

gauge symmetry

is a thing or symplectic space \Rightarrow define a moment map for action of P -terms valued in \mathfrak{g}^*


$$(L^2 \mathfrak{g}^{\text{ad}}(X)) \cong \mathfrak{g}$$

$$P \times \frac{\partial}{\partial t} \Rightarrow \text{moment map is } P$$


$$R^2/r \approx c/c$$

it's nice to think of $(\phi_t, \bar{\phi})$ as a pair of linear maps

$$\phi_t \in \text{Hom}(\bar{F}, C) \quad \phi \in \bar{C} \otimes F \cong \text{Hom}(C, \bar{F})$$





there are N_F^2 maps $F \rightarrow \bar{F}$ (if $N_C > N_F$)
 If $N_C < N_F$ and $N_F \times N_F$ matrix of rank $\leq N_C$
 on the symplectic side $\dim \Phi(N) = 2N_C N_F - \# \text{ beta gauge generators}$
 $N_F < N_C$: break $\text{SU}(N_C)$ and $\text{SU}(N_C - N_F)$
 $\Rightarrow \dim \mathfrak{g}(N) = N_F^2 = N_C^2 - (N_C - N_F)^2 - 1$
 $\Rightarrow \dim \mathfrak{g}(N) = 2N_C N_F - N_C^2 + 1$
 is $N_F \geq N_C$: break $\text{SU}(N_C)$ completely (generically)
 $\Rightarrow \dim \mathfrak{g}(N) = 2N_C N_F - N_C^2 + 1$



If $N_F = N_C$ are square matrices $\Rightarrow \det C(\phi_t) = \det \bar{C} = 1$ is a gauge inv. op.
 $N_C^2 \quad \phi_t \phi = M_1 \quad \det M = 1 \quad \text{local}$
 $N_C^2 \quad \phi_t \phi = M_1 \quad \det M = 1 \quad \text{local}$
 (bracketing paths) 3 quarks \rightarrow 3 color

$\mathbb{P}^{odd} = \mathbb{S}$
 $\mathbb{S} = \mathbb{V}^{even}$
 $\mathbb{V}^{odd} = \mathbb{S}$
 $\mathbb{V}^{even} = \mathbb{S}$

as tensor products of spinors of spin(k) and spin(d-k)



To dimensionally reduce, I will set some number of fermions to 0 (dimensional reduction)

$\mathbb{D}^2 (x_1, x_2, x_3, x_4)$
 $w_1 = dx_1 dx_2 + dx_3 dx_4$
 $w_2 = dx_1 dx_3 + dx_2 dx_4$
 $w_3 = dx_1 dx_4 + dx_2 dx_3$

Killing spinors: $\mathbb{D}^2 (2, 2) \sim \mathbb{D}^2 dx + \mathbb{D}^2 dx + \mathbb{D}^2 dx + \mathbb{D}^2 dx = 2 \cdot (\mathbb{D}^2 dx)$

multiplicity	
vector	hypermultiplet
$N=2$	$N=1$
$N=1$	$N=1$

$N=2$ multiplets in 4d:

$\mathbb{N}^2 = \mathbb{N}^2$
 $\mathbb{N}^2 = \mathbb{N}^2$

$$\Rightarrow \Lambda^*(\mathbb{C}^3)^V \cong \Lambda^*(\mathbb{C}^2)^V \oplus \Lambda^*(\mathbb{C})^V$$

$$\Lambda^{\text{even}}(\mathbb{C}^3)^V \cong \Lambda^{\text{even}}(\mathbb{C}^2)^V \oplus \Lambda^{\text{even}}(\mathbb{C})^V + \Lambda^{\text{odd}}(\mathbb{C}^2)^V \oplus \Lambda^{\text{odd}}(\mathbb{C})^V$$

$$\Rightarrow S_+^{6d} = S_+^{4d} \oplus S_+^{(2d)} \oplus S_-^{(2d)} \oplus S_-^{(2d)}$$

$$\text{total spins} \quad S_+^{(4d)} \oplus \mathbb{C}^2 \otimes S_+^{(2d)} \oplus \underbrace{S_-^{(4d)} \oplus \mathbb{C}^2 \otimes S_-^{(2d)}}_{\text{CPX conjugate}}$$

R-symmetry:

$$6d \quad Sp(1) \cong SU(2)$$

$$4d \quad U(2) \cong SU(2) \times U(1) / \mathbb{Z}_2$$

The 4d R-symmetry consists of 6d R-symmetry together with fermion rotations.

Given this, I might expect 4d $N=2$ multiplets to arise by dimensionally reducing 6d $N=1$ multiplets.

In the example of the vector multiplet:

6d Gauge Field		chiral fermions		auxiliary field	
on shell	off shell	off	on	on shell	off shell
4	6	8	4	0	2 + (1)

when I reduce, I have to decompose fields as reps. of $Spin(1,3) \times Spin(2) \subseteq Spin(1,5)$. vector \rightarrow vect + 2 real scalars (w/ $Spin(2)$ charge)

After reduction, I effectively have

- one gauge field
- 2 physical scalars (R)
- eight fermionic real d.o.f.

counting works out too: $(N=1)$ 1 vector + 1 $N=1$ chiral

4d
 $E_{N=1}$

\downarrow

2d
 $E_{N=2}$

6d
 $E_{N=1,4}$

\downarrow

4d
 $E_{N=2}$

How could I get more? SUSY?


4d $N=4$? (1 $N=1$ vector + 3 chiral)

(1 $N=2$ vector + 1 $N=2$ hyper)

16

16

the R-symmetry or almost all of it arises by dimensional reduction. Is the additional $u(1)$ important? At any it doesn't act on R-symmetry or almost all of it arises by dimensional reduction. For practical purposes we only discussed $su(4)$ -R-symmetry. Could write down a (Lorentz) action for it and by looking



$$\text{spin}(6) \cong \text{so}(6)$$

we are seeing the structure

$$s_{\text{red}} = s_{\text{red}} + s_{\text{red}} + s_{\text{red}}$$

reduction: $\text{spin}(6) \rightarrow \text{so}(6)$

to the R-symmetry: $\text{so}(6) \rightarrow \text{spin}(6)$


vector of $\text{spin}(6)$

Label degrees of freedom: $\text{so}(6) \rightarrow \text{spin}(6) \rightarrow \text{so}(6)$

$$\begin{array}{c} 16 \\ \hline 8 \end{array} \quad \begin{array}{c} 10 \\ \hline 8 \end{array} \quad \begin{array}{c} 8 \\ \hline 8 \end{array}$$

vector

Let d.c.f. be multiplied



10 d guess:

$$\mathcal{L} = \text{Tr}(F \wedge \star F) + \text{tr}(\lambda \star \lambda)$$

↓ reduce:

$$A \rightarrow (A, \phi)$$

$$\left\{ \begin{aligned} & dA \wedge \star A + \star dA \wedge [A, A] + |[A, A]|^2 \\ & \downarrow \\ & d\phi \wedge \star \phi + d\phi \wedge A \phi + A \phi A \phi \end{aligned} \right.$$

combine into $D\phi \wedge \star D\phi$

$$\lambda \star \lambda + \lambda A \lambda + \underbrace{\lambda \phi \lambda}_{\text{Yukawa couplings}} + |[A, \phi]|^2 \leftarrow \text{scalar potential}$$

Lesson: 4d $N=4$ is not 4d $N=1$ gauge theory minimally coupled to 3 adjoint chiral fields.

need to add the superpotential $W =$