

Analysis Note of the Experiment

Measurement of diffractive Central Exclusive Production of h^+h^- pairs ($h = \pi, K, p$) in proton-proton collisions at $\sqrt{s} = 200$ GeV with forward proton reconstruction in Roman Pot detectors

Leszek Adamczyk , Łukasz Fulek , Mariusz Przybycień , and Rafał Sikora

AGH University of Science and Technology, FPACS, Kraków, Poland

September 13, 2019

Abstract

In this note we present analysis of diffractive Central Exclusive Production using 2015 data from proton-proton collisions at $\sqrt{s} = 200$ GeV. This dataset was collected with newly installed Roman Pot detectors in Phase II* configuration which ensured efficient triggering and measuring diffractively scattered protons. We describe intermediate stages of analysis involving choice of selection cuts, comparison of data with Monte Carlo models folded into detector acceptance, and study of systematic uncertainties specific to the analysis. Finally, we show the physics outcome of the analysis. Parts of the analysis which are of more technical nature (calculation of efficiencies, derivation of corrections to efficiencies, adjustment of the STAR simulation, systematic uncertainty of efficiencies) are described in a supplementary analysis note [1].

ver. 0.95

Contents

List of contributions	4
Change log	4
1 Introduction	5
1.1 Central Exclusive Production	5
1.2 Double IPomeron Exchange	5
1.3 Physics motivation for the measurement	6
1.3.1 DIPE differential cross-sections, mass spectrum	6
1.3.2 Absorption effects	6
1.3.3 Size of interaction region	6
2 Data set	7
2.1 Trigger	7
2.2 Reconstruction software	8
2.3 Data format	8
3 Event selection	9
3.1 List of cuts	9
3.2 Description of cuts	10
3.2.1 (C1,C2) Primary vertex and its z -position	10
3.2.2 (C3) TPC tracks	10
3.2.3 (C4) RP tracks	14
3.2.4 (C5) TPC-RP z -vertex matching	15
3.2.5 (C6) BBC-large signal veto	16
3.2.6 (C7) TOF clusters limit	18
3.2.7 (C8) Particle identification	19
3.2.8 (C9) Exclusivity cut (missing p_T cut)	23
4 Backgrounds	27
4.1 Sources of background	27
4.1.1 Non-exclusive background	27
4.1.2 Exclusive background (particle misidentification)	28
4.2 Non-exclusive background determination	28
4.3 Normalization of signal and background models	28
5 Corrections	34
5.1 Method of corrections application	34
5.2 Acceptances and efficiencies	34
5.2.1 Trigger efficiency	34
5.2.1.1 Online TOF efficiency	34
5.2.1.2 Online veto (BBC-small and ZDC veto)	35
5.2.1.3 RP triggering efficiency	36
5.2.1.4 Up and Down RP combination veto (due to dead material)	36
5.2.2 Reconstruction and selection efficiency	37
5.2.2.1 TPC z -vertex cut (C2)	37
5.2.2.2 TPC-RP z -vertex matching (C5)	38
5.2.2.3 Primary vertices limit (C1), BBC-large veto (C6), TOF clusters limit (C7) and Up and Down RP combination veto (due to pile-up)	38

5.2.2.4	Missing p_T cut (C9)	39
5.2.2.5	Particle identification (C8)	40
5.2.3	RP track acceptance and reconstruction efficiency	43
5.2.4	TPC vertex reconstruction efficiency	44
5.3	Particle energy loss	45
5.4	Fake tracks and migrations into and out of fiducial region	46
5.5	Background subtraction	49
5.6	Closure tests	49
5.6.1	Data pull	49
5.6.2	Monte Carlo closure tests	50
5.6.2.1	TPC and TOF efficiency correction validation	50
5.6.2.2	RP efficiency correction validation	50
5.6.2.3	Total correction validation	50
6	Systematic uncertainties	52
6.1	Trigger veto effect (due to dead material)	52
6.2	Discussion of systematic effects	53
6.3	Graphical representation of systematic uncertainties	54
7	Physics results	59
Appendix A	BBC response	66
Appendix B	Formulation of total RP efficiency	75
Appendix C	Reconstruction of m_{TOF}^2	77
Appendix D	RP efficiency	78
D.1	RP track acceptance, reconstruction and selection efficiency	78
D.2	RP trigger veto probability related to dead material	84
Appendix E	Particle identification efficiency	90
List of Figures		94
List of Tables		94
References		96

List of contributions

Leszek Adamczyk	Analysis coordination/supervision, production of picoDST, production of embedded MC samples
Lukasz Fulek	Analysis support
Mariusz Przybycień	Analysis supervision
Rafał Sikora*	Main analyzer, write-up author

* - contact editor

Change log

30 May 2019 ver. 1.0 Initial revision

1. Introduction

1.1 Central Exclusive Production

The Central Exclusive Production (CEP) takes place when interacting particles form a state in the mid-rapidity region (“central production”) whose all constituents/decay products are measured in the detector (“exclusive”). The initial state particles can either dissociate, excite or stay intact. The latter case of CEP in proton-proton collisions can be written as

$$p + p \rightarrow p + X + p \quad (1.1)$$

and depicted as in Fig. 1.2. Mass and rapidity of state X is given by

$$M_X = \sqrt{s \left(\xi_1 \xi_2 \sin^2(\alpha/2) - (1 - \xi_1 - \xi_2) \cos^2(\alpha/2) \right)} \stackrel{\alpha=\pi}{=} \sqrt{s \xi_1 \xi_2}, \quad (1.2)$$

$$y_X = \frac{1}{2} \ln \frac{\xi_1}{\xi_2}, \quad (1.3)$$

where α is angle between scattered protons and $\xi = (p_0 - p)/p_0$ is the fractional momentum loss of proton.

1.2 Double IPomeron Exchange

Reaction from Eq. (1.1) can exhibit purely electromagnetic ($\gamma\gamma$ interaction), mixed ($\gamma\mathcal{O}$ interaction) or purely strong nature ($\mathcal{O}\mathcal{O}$ interaction). The last type is dominant at RHIC energies. It is characterized by the lack of hard scale (if protons are scattered at small angles), therefore perturbative QCD cannot be applied and Regge theory [2] is used instead. An object \mathcal{O} does not have unequivocal QCD representation - in Regge formalism it is the so-called “trajectory” (IReggeon, IR). IReggeon with quantum numbers of vacuum is called “IPomeron” (IP) and IP-IP reaction (Fig. 1.1) is called “Double IPomeron Exchange“.

Processes involving IPomeron exchange are referred as diffraction due to cross-section in scattering angle resembling similar shape to intensity pattern of diffracted light. For low values of Mandelstam t (small scattering angles) cross-section takes exponential form

$$\frac{d\sigma}{dt} \propto e^{-B|t|}, \quad (1.4)$$

where the slope parameter B reflects the size of target at which IPomerons scatter.

Diffractive events have specific property of the ”rapidity gap“ which is an angular region free of hadrons. In DIPPE two such gaps are present, marked in Fig. 1.2 as $\Delta\eta_1$ and $\Delta\eta_2$. Figure 1.3 shows the topology of the DIPPE event on top of the STAR detector, with centrally produced particles marked with green arrows and two forward protons escaping the interaction point inside the beam pipe drawn with red arrows.




Figure 1.2: CEP represented in $\eta\phi$ space.




Figure 1.3: Central Production event at STAR.

DIPE is a spin-parity filter - the fact that scattered particles have all quantum numbers unchanged after the interaction, production of central states satisfying Eq. (1.5) is enhanced

$$I^G J^{PC} = 0^+ \text{even}^{++}. \quad (1.5)$$

The lowest order QCD picture of the IPomeron is a pair of oppositely colored gluons (colour singlet). This fact makes the DIPE recognized as the gluon-rich environment process in which bound states of gluons ("glueballs") or hybrid mesons could be preferably produced.

For detailed introduction to the topic of diffraction see Refs. [3, 4].

1.3 Physics motivation for the measurement

STAR collected in 2015 large dataset dedicated for measurement of the Central Diffraction (DIPE in particular). Since that year the experiment was enriched with Roman Pot Phase II* subsystem and thus gained possibility of detection of forward protons. It enabled studies of properties of the central state with respect to observables related to exchanged IPomerons. No such measurement was performed before at that high c.m.s. energy ($\sqrt{s} = 200$ GeV, contamination from Reggeon exchanges is small) which makes it particularly attractive. A brief list of physics issues that can be covered with the study described in this note is briefly introduced below.

1.3.1 DIPE differential cross-sections, mass spectrum

As stated in Sec. 1.2 DIPE is a soft process whose theoretical description is done mainly using phenomenological tools, thus measurement of differential cross-sections is needed to verify various production models.

The main focus is put on the simplest state (and most numerous) produced in DIPE, namely a pair of oppositely charged pions, $\pi^+\pi^-$. It can be formed either in a non-resonant or resonant mechanism. In the first case the $\pi^+\pi^-$ continuum is formed by the exchange of the off-shell pion between IPomerons. Currently there are two models of this reaction on the market [5, 6], [7]. In the second case the IPomerons directly couple into resonance (e.g. $f_2(1270)$), which then decays to $\pi^+\pi^-$. Attempts to calculate cross-section for this production mechanism are presented in Ref. [6] and [8].

Understanding of the mass spectrum in $\pi^+\pi^-$ channel is important to learn about relative contribution from continuum and resonant production, as well as relative production of resonances. Recognition of resonant states may indicate candidates for low-mass glueballs of $J^{PC} = 0^{++}$, however presence of underlaying scalar $q\bar{q}$ states makes this task challenging.

Other channels, like K^+K^- , are also of great interest. Comparison of the cross-sections for production of $\pi^+\pi^-$ and K^+K^- gives information about strength of the IPomeron coupling to different quark flavors. Also, structures in $d\sigma/dm$ can be easier attributed to resonances by measuring more than one channel and known branching ratios thereof.

Detection of intact protons scattered at very small angle with respect to the beamline enables determination of the reaction plane which makes the Partial Wave Analysis (PWA) possible. It also allows to look at the cross-sections more differentially, especially with respect to properties of exchanged IPomerons, like carried squared four-momentum t , azimuthal separation of IPomerons in the transverse plane $\Delta\varphi$ or relative momentum of IPomerons Δp_T . The last quantity was proposed to distinguish pure $q\bar{q}$ states from these with gluonic content [9].

1.3.2 Absorption effects

One can imagine in diagram in Fig. 1.1 additional soft lines e.g. between protons in the initial state or one of IPomerons and final state proton. These so-called rescattering effects (or absorption effects) lead to production of hadrons other than these belonging to central state X hence the diffractive signature of an event in form of rapidity gap is no longer present. Measurement of the probability that the state X will remain exclusive and forward protons will remain intact, in other words the rapidity gap survival probability S^2 , would be valuable ingredient for development of absorption models.

1.3.3 Size of interaction region

From the measurement of protons in Roman Pots one is able to reconstruct squared four-momenta transferred in proton-IPomeron vertices and determine the differential cross-section $d\sigma/d|t|$. Fit of exponent allows to extract the slope parameter B , which may depend on the IPomeron-IPomeron c.m.s. energy, or in other words on the mass of diffractive system X . Knowledge on the slope parameter gives insight to the form factor of the object at which IPomeron scatters.

2. Data set

2.1 Trigger

Presented analysis made use of data taken in 2015 during $p + p$ collisions at $\sqrt{s} = 200$ GeV. The main trigger designed for studies of Central Diffraction in run 15 was RP_CPT2. It was formed of the following conditions combined by logical AND ($\&\&$):

1. **(ET $\&\&$!IT) || (!ET $\&\&$ IT)** = signal in at least one RP on each side of the STAR central detector - to ensure presence of two forward-scattered protons; a veto was imposed on simultaneous signal in RPs above and below the beamline, which might have originated either from proton dissociation, or pile-up event, or beam halo proton etc.,
2. **!BBC-E $\&\&$!BBC-W $\&\&$!ZDC-E $\&\&$!ZDC-W** = veto on any signal in small BBC tiles or ZDCs on any side of STAR central detector - such requirement is in accordance with the double-gap topology of CEP events, it mostly filtered out CEP events with parallel pile-up event(s),
3. **TOF ≥ 2** = at least 2 hits in TOF - aim of this condition was to ensure activity in the mid-rapidity; since the lowest multiplicity allowed in CEP is 2, that was the lower threshold of L0 TOF multiplicity.

This trigger was running with an average prescale of 5 and average DAQ rate of 250 Hz, which allowed to collect in total about 560 M events corresponding to 16.5 pb^{-1} of integrated luminosity. More information about number of events per run, rates etc. can be found under link provided in Ref. [10], which contains selected data from STAR run log [11]. Luminosity data used in this analysis comes from Ref. [12].

All RP triggers which were intended for usage in diffractive physics analyses or efficiency studies are listed in Tab. 2.1. Components used in definitions of these triggers are outlined in Fig. 2.1. Detailed explanation of all trigger bits can be found in Refs. [13, 14]. Explanation of naming convention in Roman Pot system can be found in Ref. [15].



Figure 2.1: Sketch of the trigger components used in definitions of diffractive triggers in run 15.

Trigger name	Definition	Events [M]	Comment
RP_CPT	EOR $\&\&$ WOR	73.3	Loose trigger (mostly elastic events) designed for monitoring/trigger efficiency study
RP_CPT	EOR $\&\&$ WOR $\&\&$!BBC-E $\&\&$!BBC-W $\&\&$!ZDC-E $\&\&$!ZDC-W $\&\&$ TOF ≥ 1	38.9	Intended to be main CEP trigger (later switched to RP_CPT2 due to large prescale)
RP_CPT2	(ET $\&\&$!IT) (!ET $\&\&$ IT) $\&\&$!BBC-E $\&\&$!BBC-W $\&\&$!ZDC-E $\&\&$!ZDC-W $\&\&$ TOF ≥ 2	556.5	Main CEP trigger Note: On Apr 14 added upper TOF limit (10)
RP_CPTX	IT $\&\&$!BBC-E $\&\&$!BBC-W $\&\&$!ZDC-E $\&\&$!ZDC-W $\&\&$ TOF ≥ 2	40.1	The same as RP_CPT2 but only IT configuration
RP_CPEI	ET $\&\&$ IT $\&\&$!BBC-E $\&\&$!BBC-W $\&\&$!ZDC-E $\&\&$!ZDC-W $\&\&$ TOF ≥ 2	15.6	Control trigger for CPT2 to estimate effect of !(ET $\&\&$ IT) veto

Table 2.1: Central Diffraction physics triggers and control triggers involving Roman Pot detectors in run 15.

2.2 Reconstruction software

Raw data was processed with STAR libraries in versions SL17f. All four trigger datasets were processed: production_pp200trans_2015, production_pp200long2_2015, production_pp200long3_2015 and production_pp200long_2015 (see [16]).

The following BFC options were used in the reconstruction:

```
DbV20160418,pp2015c,btof,mtd,mtdCalib,pp2pp,-beamline,beamline3D,useBTOFmatchOnly,VFStoreX,  
fmsDat,fmsPoint,fpsDat,BEmcChkStat,-evout,CorrX,OSpaceZ2,OGridLeak3D,-hitfilt
```

Main attention should be put on option **useBTOFmatchOnly** which forced vertexing algorithm to form vertices only from the global TPC tracks which are matched with hits in the TOF system. This solution was found to yield significantly larger signal reconstruction efficiency (vertexing efficiency) and better resolutions. The study which lead to above conclusions, presented in Ref. [17], was performed on the same dataset processed with older libraries SL15k (without useBTOFmatchOnly option).

2.3 Data format

The analyzed data was stored in ROOT files in the picoDST format which was in large part a skimmed MuDST (standard STAR format). The picoDST format was introduced in Ref. [18]. PicoDST description files (C++ headers etc.) can be found in the analysis code repository [19].

3. Event selection

Complete list of analysis cuts used for signal extraction is presented in Sec. 3.1. Detailed description of each cut can be found in Sec. 3.2.

3.1 List of cuts¹

C1. Exactly 1 primary vertex with TPC track(s) matched with hits in TOF.

C2. TPC vertex from C1 is placed within $|z_{\text{vx}}| < 80$ cm.

C3. Exactly 2 opposite-sign primary TPC tracks (C3.2) of good quality (C3.4) matched with hits in TOF (C3.1), reconstructed within kinematic region of high TPC acceptance (C3.3), with small distance of closest approach (DCA) to the vertex (C3.5) and high proximity to each other at the beamline (C3.6).

C3.1. Exactly 2 TOF-matched (match flag > 0) primary tracks and no additional primary tracks matched with BEMC clusters,

C3.2. Tracks are of opposite signs,

C3.3. Both tracks are contained within the kinematic range: $|\eta| < 0.7$, $p_T > 0.2$ GeV/ c ,

C3.4. Both tracks satisfy quality criteria: $N_{\text{hits}}^{\text{fit}} \geq 25$, $N_{\text{hits}}^{\text{dE/dx}} \geq 15$, $|d_0| < 1.5$ cm,

C3.5. Both tracks match well to the primary vertex: $\text{DCA}(R) < 1.5$ cm, $|\text{DCA}(z)| < 1$ cm,

C3.6. Tracks are close to each other at the beamline: $|\Delta z_0| < 2$ cm.

C4. Exactly 1 RP track on each side of STAR central detector (C4.3) of good quality (C4.1), with local angles consistent with the IP being the track origin (C4.2), lying within fiducial region of high geometrical acceptance (C4.4).

C4.1. RP tracks contain only track-points with at least 3 (out of 4) planes used in reconstruction,

C4.2. Local angles (θ_x^{RP} , θ_y^{RP}) consistent with expectation for protons originating from the IP

$$-2 \text{ mrad} < \theta_x^{\text{RP}} - x^{\text{RP}}/|z^{\text{RP}}| < 4 \text{ mrad}, \quad -2 \text{ mrad} < \theta_y^{\text{RP}} - y^{\text{RP}}/|z^{\text{RP}}| < 2 \text{ mrad},$$

C4.3. Exactly 1 track passing cuts C4.1-C4.2 per side,

C4.4. Tracks passing cut C4.3 lie within the fiducial (p_x , p_y) region defined as:

$$0.2 < |p_y| < 0.4, \quad -0.2 < p_x, \quad (p_x + 0.3)^2 + p_y^2 < 0.5^2 \quad (\text{all in GeV}).$$

C5. Vertex z -positions measured in TPC and reconstructed from the difference of proton detection time in west and east RPs are consistent with each other within the resolution (at $3.5\sigma_{\Delta z_{\text{vtx}}}$ level):

$$|\Delta z_{\text{vtx}}| = |z_{\text{vx}}^{\text{TPC}} - z_{\text{vx}}^{\text{RP}}| < 36 \text{ cm}.$$

C6. No signal in any tile of BBC-large (east or west) with $\text{ADC} > \text{ADC}_{\text{thr}}$ and $100 < \text{TDC} < 2400$, where ADC_{thr} is specific for each channel.

C7. Maximally 3 reconstructed TOF clusters $N_{\text{clstrs}}^{\text{TOF}} \leq 3$.

C8. Particle (pair) identification:

C8.1. Identification of particle pairs based on χ^2 or $n\sigma$ (from dE/dx) and m_{TOF}^2 :

```
if  $\chi^2(\pi\pi) > 9$  and  $\chi^2(KK) > 9$  and  $\chi^2(pp) < 9$  and  $m_{\text{TOF}}^2 > 0.6$  GeV/ $c^2$  →  $p\bar{p}$ 
elif  $\chi^2(\pi\pi) > 9$  and  $\chi^2(KK) < 9$  and  $\chi^2(pp) > 9$  and  $m_{\text{TOF}}^2 > 0.15$  GeV/ $c^2$  →  $K^+K^-$ 
elif  $|n\sigma_{\text{pion}}^{\text{trk1}}| < 3$  and  $|n\sigma_{\text{pion}}^{\text{trk2}}| < 3$  →  $\pi^+\pi^-$ .
```

C8.2. Restricting fiducial cuts on K^+K^- and $p\bar{p}$:

```
if  $p\bar{p}$ :  $p_T > 0.4$  GeV/ $c$ ,  $\min(p_T^+, p_T^-) < 1.1$  GeV/ $c$ 
if  $K^+K^-$ :  $p_T > 0.3$  GeV/ $c$ ,  $\min(p_T^+, p_T^-) < 0.7$  GeV/ $c$ 
```

C9. Missing (total) momentum of TPC tracks and RP tracks $p_T^{\text{miss}} < 75$ MeV/ c .

¹Some cuts (e.g. C3) are decomposed to constituent sub-cuts. Cut is formed by the logical AND of all its sub-cuts. Events must pass all cuts to be identified as a signal.

3.2 Description of cuts

3.2.1 (C1,C2) Primary vertex and its z -position

As it was designed in the trigger logic, we aim to perform CEP analysis in a clean, pile-up-free environment, therefore we cut on primary vertex multiplicity (Fig. 3.1) to reject events with more than one interaction per bunch crossing. We require exactly one primary vertex containing TPC tracks matched with hits in TOF (matching of the track with hit in TOF is identified with the TOF match flag being different from 0). Later in the text we refer to such events as a single “TOF vertex” events.



Figure 3.1: Primary vertex multiplicity. Red arrow marks bin with events with exactly one primary vertex (with track(s) matched with hit in TOF), which are used in physics analysis.

The single TOF vertex is required to be placed within a range (-80 cm, 80 cm) along the z -axis (Fig. 3.2). Events with vertices away from the nominal IP have low acceptance both for the central tracks and the forward protons (comparing to events with vertices close to nominal IP), therefore we reject them as their inclusion to analysis would naturally introduce large systematic uncertainties. See Sec. 3.2.3 in Ref. [1]

3.2.2 (C3) TPC tracks

The TPC track selection starts from the selection of events with exactly two primary tracks matched with hit in TOF (Fig. 3.3). Matching with TOF guarantee that analyzed tracks originate from the triggered bunch crossing (ensures that tracks are “in-time”). It is in accordance with the trigger logic which required at least 2 L0 TOF hits, as well as it enables more accurate particle identification with merged time-of-flight and dE/dx method, comparing to sole usage of dE/dx . Primary tracks not matched with hit in TOF, whose average multiplicity in single TOF vertex is ~ 8 , are hardly distinguished between real and fake (off-time) tracks, which is an additional reason for not analyzing events with only one TOF-matched primary TPC track (the other track might be unmatched due to TOF inefficiency).

Primary TPC tracks from the single TOF vertex which are matched with TOF are allowed to be also matched with BEMC clusters. Matching with BEMC cluster is claimed if the distance in $\eta - \phi$ space between the BEMC cluster position (η_{clus} , ϕ_{clus}) and projected position of the track in BEMC (η_{proj} , ϕ_{proj}), defined as

$$R = \sqrt{(\eta_{\text{clus}} - \eta_{\text{proj}})^2 + (\phi_{\text{clus}} - \phi_{\text{proj}})^2}, \quad (3.1)$$

is less than $R_{\text{max}}^{\text{match}} = 0.05$. Distribution of the distance between the primary TPC track and the closest BEMC cluster is shown in Fig. 3.4.

However, if there are any primary TPC tracks matched with BEMC cluster and not matched with TOF in the single TOF vertex with two TOF-matched tracks, an event is rejected. Such configuration implies higher-than-2 multiplicity of the real tracks in the vertex, hence an event is unlikely a Central Exclusive Production of two particles.



Figure 3.3: Multiplicity of primary TPC tracks matched with hit in TOF for single TOF vertex events. Red arrow marks bin with events with exactly two primary tracks matched with hit in TOF, which are used in physics analysis.




Figure 3.4: Distribution of a distance in $\eta - \phi$ space between the BEMC cluster closest to primary TPC track matched (filled circle) or not matched (open circle) with hit in TOF, for single TOF vertex events. Red dashed line indicate matching threshold $R_{\max}^{\text{match}} = 0.05$.



(a)




(b)




(c)

Figure 3.5: Comparison of distribution of the number of hits used in TPC track reconstruction $N_{\text{hits}}^{\text{fit}}$ (3.5a), number of hits used in specific energy loss reconstruction $N_{\text{hits}}^{\text{dE/dx}}$ (3.5b) and fraction of number of hits potentially generated by the track and finally used in the reconstruction $N_{\text{hits}}^{\text{fit}}/N_{\text{hits}}^{\text{poss}}$ (3.5c) in the data and embedded MC. Normalizations of the signal and backgrounds were established from the comparison of p_T^{miss} and $\Delta\theta$ distributions after full selection (without cut on the presented quantity and without exclusivity cut), as described in Sec. 4.3. Red dashed line and red arrow indicate the range of each quantity which is accepted in analysis.



(a)



(b)

Figure 3.6: Comparison of the track pseudorapidity η (3.6a) and the track azimuthal angle ϕ (3.6b) in the data and embedded MC. Normalizations of the signal and backgrounds were established from the comparison of p_T^{miss} and $\Delta\theta$ distributions after full selection (without cut on the presented quantity and without exclusivity cut), as described in Sec. 4.3. Red dashed lines and red arrows indicate the range of each quantity which is accepted in analysis.

3.2.3 (C4) RP tracks




Figure 3.7: Typical correlation between local angle (y -axis) and position (x -axis) of RP tracks matched with true level primary protons for x - (a) and y -coordinate (b), here shown for branch WU. The same events are contained in c and d for x - and y -coordinate respectively, where difference between reconstructed local angle and local angle expected from the elastic track is histogrammed. Red lines and arrows visualize cuts imposed on RP tracks for final selection (cuts C4.2).




Figure 3.8: (left) Merged distributions of diffractively scattered protons momenta p_y vs. p_x in exclusive h^+h^- events reconstructed with the East and West RP stations, together with the kinematic region used in the measurement marked with the black line. (right) Distributions of measured four momenta transfers at the proton vertices for exclusive h^+h^- events with all particles in the fiducial phase space are shown for East and West stations with yellow and blue color, respectively.

3.2.4 (C5) TPC-RP z -vertex matching

In CEP tracks in the central detector and tracks in Roman Pots originate from the same interaction vertex. Measurement of the time of detection of forward protons in RPs gives access to reconstruction of the position of the vertex

$$z_{\text{vtx}}^{\text{RP}} = c \cdot \frac{t_{\text{W}}^{\text{RP}} - t_{\text{E}}^{\text{RP}}}{2} \quad (3.2)$$

independently from TPC, which allows their comparison and rejection of the background if the two values disagree. Time of detection of proton in RP is provided in StMuRpsTrack object - it is an average of all TAC values from PMTs in RPs used to form a track, corrected for the slewing effect and adjusted to have the best correlation with the z -position of the vertex measured in TPC, translated to unit of time (all these steps are done at the level of raw data reconstruction). In Fig. 3.9 the comparisons of the $z_{\text{vtx}}^{\text{RP}}$ and $z_{\text{vtx}}^{\text{TPC}}$ are shown with some preselection cuts applied. A clear signal from the Central Diffraction (and thus CEP) process is manifesting in high correlation of the two values (diagonal in Fig. 3.9a) or significant and relatively narrow peak centered at 0 for the difference of two values (Fig. 3.9b). The sum of two Gaussian distributions was fitted to data in Fig. 3.9b yielding good description of the distribution of Δz_{vtx} with the width parameters equal 10.3 cm (CD signal) and 73.9 cm (pile-up). The first parameter reflects the time resolution of RPs (the $z_{\text{vtx}}^{\text{RP}}$ measurement), as the TPC resolution is much better (~ 1 cm). Value of the second parameter, consistent with $\sqrt{2}\sigma_{z_{\text{vtx}}} \approx \sqrt{2} \cdot 52 \text{ cm} \approx 73.5 \text{ cm}$, confirms that the wide distribution under the narrow signal peak is uncorrelated background, in other words forward protons originating from a different vertex than the central tracks. To reject this background without significant loss of the signal, we introduce $3.5\sigma_{\Delta z_{\text{vtx}}}$ cut on Δz_{vtx} .




Figure 3.9: Correlation (Fig. 3.9a) and difference (Fig. 3.9b) of z -vertex position measured in Roman Pots and TPC in RP_CPT2 triggers, after preselection described in the plots.

3.2.5 (C6) BBC-large signal veto

At the trigger level a veto on signal in small BBC detectors was used. During offline analysis we found that the non-exclusive background can be efficiently rejected if an additional veto on signal in large BBC detectors is added. It is connected with the fact that vast majority of selected RP_CPT2 triggers were from the central diffraction process to which CEP belongs. Many of central diffraction events have particles produced in the rapidity region outside the TPC and TOF acceptance, some hitting large BBC tiles. Presence of signal in large BBC is therefore a signature of background or a pile-up interaction.


The response of large BBC tiles is different from that of small BBC tiles, as shown in sample plots in Fig. ?? (similar distributions for all channels can be found in Appendix A). Typically in small BBC tiles a peak visible in ADC distribution around $100 - 150$ (Figs. 3.10a,3.10c), a signature of good separation of the electronics noise and signal from the ionizing particle. No such feature is observed in corresponding distribution for large BBC tile (Figs. 3.10b,3.10d), which can be explained by the difference in geometry (in size) of small and large tiles. In large BBC tiles the path that scintillation light must travel to reach PMT is much longer in comparison to small BBC tiles (multiple reflections on the main tile surface due to small thickness of the tile) therefore it is highly attenuated and extended in time. This is possible reason of lack of signal peak in the ADC distribution in large BBC tile spectrum (Fig. 3.10d), as well as the late-TAC (TAC $< \sim 600$, ADC < 100) tail in the ADC vs. TAC spectrum (slewing effect, Fig. 3.10b). Nevertheless, the above features of BBC-large response does not disqualify this detector from being used as a veto detector, as in this case lower efficiency of the detector only reduce the background rejection power.

Each channel of the BBC-large has different response to signal from ionizing particle, as well as different level of noise. We decided to set up a signal threshold for each channel based on a study of the noise in abort gaps (in zero-bias data). This noise, in principle, should be solely the electronics noise. We checked for each channel the probability to detect a signal with ADC above certain threshold and with TAC contained within 100 and 2400 (the same window is default for small BBC). The result is shown in Fig. 3.11. Next, we established final ADC thresholds in each BBC-large channel by requiring that the noise in BBC-large would cause a veto in maximally 3.5% of events. To transform it to ADC_{thr} we first assumed that the noise is uncorrelated between the channels. With this assumption one can connect the probability of the veto in whole BBC-large detector (east and west) caused by noise $\mathcal{P}_{\text{veto}}^{\text{noise}}$ with the probability of the signal induced by noise in single BBC-large channel $\mathcal{P}_{i,\text{sig}}^{\text{noise}}$:


$$\mathcal{P}_{\text{veto}}^{\text{noise}} = 1 - \mathcal{P}_{i,\text{veto}}^{\text{noise}} = 1 - (1 - \mathcal{P}_{i,\text{sig}}^{\text{noise}})^{N_{\text{ch}}^{\text{BBC}}}. \quad (3.3)$$

In the equation above $N_{\text{ch}}^{\text{BBC}}$ denotes number of active channels in BBC-large. From plots contained in Appendix A one can read that there were 14 active channels in BBC-large. 2 dead channels were found on the west side (40 and 42). By transforming Eq. 3.3 to the form presented below we can calculate the threshold probability for a single BBC-large channel:


$$\mathcal{P}_{i,\text{sig}}^{\text{noise}} = 1 - \sqrt[N_{\text{ch}}^{\text{BBC}}]{1 - \mathcal{P}_{\text{veto}}^{\text{noise}}} = 1 - \sqrt[14]{1 - 0.035} \approx 0.0025. \quad (3.4)$$




(a)



(b)



(c)



(d)

Figure 3.10: Sample BBC-small (left column) and BBC-large (right column) response in zero-bias data. Top row shows TAC vs. ADC distributions, bottom row shows projection of the corresponding two-dimensional ditribution on x -axis (ADC) in the TAC range quoted in the legend, for both abort gaps and colliding bunches. Red lines and arrows indicate thresholds for a signal in presented channels.

In the last step we translated this number to ADC threshold for each channel of BBC-large. For this purpose we used Fig. 3.11. The x -axis projection of the crossing point of each color line with the y -axis value of 0.0025 defines ADC_{thr} for each particular channel. These numbers are listed in Tab. 3.1. The event was dropped from analysis if any of the BBC-large channels registered signal of strength $\text{ADC}_i > \text{ADC}_{i,thr}$ and $100 < \text{TAC}_i < 2400$.

Observation of high purification of CEP sample with described BBC-large veto in the data from run 15 was helpful to improve the CEP trigger for run 17. The improved trigger called RP_CPT2noBBCL was similar to RP_CPT2 with an addition of BBC-large veto using ADC threshold of 50.




Figure 3.11: Percentage of events in abort gaps from zero-bias triggers with the ADC counts larger than the ADC threshold given in the x -axis, for each BBC-large channel. Measured points with statistical uncertainties are connected with a smooth line of corresponding color for better visualization.

3.2.6 (C7) TOF clusters limit

The TOF is mainly used to distinguish real TPC tracks from the fakes, as well as it helps to identify particles. However, we also used it to reject non-CEP events in which the TPC tracks were not reconstructed or were not successfully matched to TOF hit. For this we introduced a concept of a TOF cluster - a group of offline TOF hits close in space and time. We expect that such cluster of hits is induced by the single primary particle, eventually associated with the secondaries (e.g. delta rays).

We define a TOF cluster as a group of reconstructed TOF hits with the (ϕ, η) space distance R to neighbouring hit (defined similarly to Eq. (3.1) not larger than 0.1 and with the time distance to the same hit Δt not larger than 1.5 ns. In other words, TOF clusters are formed by the offline hits that form at least one pair with the other hit in the cluster satisfying

$$R < 0.1, \quad \Delta t < 1.5 \text{ ns.} \quad (3.5)$$

Per event no more than 1 additional TOF cluster was allowed, thus in total the number of reconstructed TOF clusters $N_{\text{clstrs}}^{\text{TOF}}$ could not exceed 3.

i	East		West	
	ADC_{thr}	i	ADC_{thr}	i
16	27	40	0	
17	30	41	31	
18	26	42	0	
19	37	43	14	
20	25	44	29	
21	55	45	30	
22	43	46	33	
23	27	47	22	

Table 3.1: Offline ADC thresholds in BBC-large.




Figure 3.12: NTofClusters.

3.2.7 (C8) Particle identification

Particles were identified using combined information from the TPC (dE/dx) and TOF (time of hit detection in the TOF subsystem). Merging informations from two sources led to reduction of misidentifications, as well as gave access to higher kaon and proton momentum range where dE/dx of different species overlap.

Compatibility of track dE/dx with that expected from particle of type X was determined using the quantity $n\sigma_X$ widely used at STAR, defined as

$$n\sigma_X = \ln \left[(dE/dx)^{\text{measured}} / (dE/dx)_X^{\text{theory}} \right] / \sigma_{dE/dx}, \quad (3.6)$$

where $(dE/dx)^{\text{measured}}$ is the ionization energy loss of the TPC track, $(dE/dx)_X^{\text{theory}}$ is the Bethe-Bloch [20] expectation for the given particle type ($X = \pi, K, p$) at reconstructed track momentum, and $\sigma_{dE/dx}$ is the statistical uncertainty of $\ln(dE/dx)^{\text{measured}}$. Quantity $n\sigma_X$ is in fact a pull: $(dE/dx)^{\text{measured}}$ is (in first order) an average over Landau \otimes normal-distributed dE/dx of single TPC hits forming the track, hence according to the central limit theorem² the $(dE/dx)^{\text{measured}}$ is distributed log-normally and $\ln(dE/dx)^{\text{measured}}$ - normally. From $n\sigma_X$ of the two tracks the χ^2 statistic for a XX pair hypothesis was calculated:

$$\chi^2_{dE/dx}(XX) = (n\sigma_X^{\text{trk1}})^2 + (n\sigma_X^{\text{trk2}})^2. \quad (3.7)$$

Sometimes we also quote $n\sigma_X^{\text{pair}}$ quantity (which is no longer a Gaussian pull) connected with χ^2 through relation

$$n\sigma_X^{\text{pair}} = \sqrt{\chi^2_{dE/dx}(XX)} = \sqrt{(n\sigma_X^{\text{trk1}})^2 + (n\sigma_X^{\text{trk2}})^2}. \quad (3.8)$$

The time of detection of particle in the TOF system was used to reconstruct its squared mass m_{TOF}^2 . For this purpose the time of primary interaction is typically used ("start time"), reconstructed by detecting fragments of dissociated beam particles in VPD detectors on both sides of the interaction point³. However, it is not accessible in CEP as the initial protons survive the interaction intact. We therefore assumed that both central tracks are of the same type which is natural consequence of quantum number conservation. With this assumption the time difference between TOF hits and measured tracks' momenta and lengths of helical paths between the primary vertex and TOF then allow to calculate m_{TOF}^2 . The derivation of formula used to obtain m_{TOF}^2 is presented in Appendix C.

Particle identification involved a few steps. First, the pp hypothesis was verified:

$$\underbrace{\chi^2_{dE/dx}(pp) < 9}_{\text{likely } pp} \quad \& \quad \underbrace{m_{\text{TOF}}^2 > 0.6 \text{ GeV}^2}_{\text{unlikely } \pi\pi \text{ or } KK} \quad \& \quad \underbrace{\chi^2_{dE/dx}(\pi\pi) > 9 \quad \& \quad \chi^2_{dE/dx}(KK) > 9}_{\text{unlikely } \pi\pi \text{ or } KK}. \quad (3.9)$$

²Keeping in mind that it assumes finite mean and variance of the distribution that summed components follow.

³Time measured from protons in the RP detectors cannot be used because RP readout runs on independent clock from that used by VPD and TOF.

If any of above was not satisfied, the pair was checked for compatibility with KK hypothesis:

$$\overbrace{\chi_{dE/dx}^2(KK) < 9 \text{ } \& \text{ } m_{\text{TOF}}^2 > 0.15 \text{ GeV}^2}^{\text{likely } KK} \text{ } \& \text{ } \underbrace{\chi_{dE/dx}^2(\pi\pi) > 9}_{\text{unlikely } \pi\pi} \text{ } \& \text{ } \underbrace{\chi_{dE/dx}^2(pp) > 9}_{\text{unlikely } pp}. \quad (3.10)$$

In case the pair was neither recognized as $p\bar{p}$ or K^+K^- , it was assumed to be a $\pi^+\pi^-$ pair if the dE/dx of positive and negative charge track was consistent with pion hypothesis at 3σ level:

$$|n\sigma_\pi^{\text{trk}1}| < 3 \text{ } \& \text{ } |n\sigma_\pi^{\text{trk}2}| < 3. \quad (3.11)$$




Figure 3.13: Two-dimensional distributions of $n\sigma_{\pi}^{\text{pair}}$ vs. $n\sigma_K^{\text{pair}}$ (a), $n\sigma_{\pi}^{\text{pair}}$ vs. $n\sigma_p^{\text{pair}}$ (b) and $n\sigma_K^{\text{pair}}$ vs. $n\sigma_p^{\text{pair}}$ (c) for exclusive event candidates after full event selection except PID cuts (except cuts C8). Dashed lines indicate the value of $n\sigma_{\text{X}}^{\text{pair}}$ which is used in pair identification C8 ($n\sigma_{\text{X}}^{\text{pair}} = 9$ which is equivalent to $\chi^2(XX) = 9$).

In Fig. 3.13 we present two-dimensional distributions of $n\sigma^{\text{pair}}$ variables which help better understand the behaviour and aim of $n\sigma^{\text{pair}}$ (χ^2) cuts in Eqs. (3.9), (3.10). Regions of enriched population of specific pair species are appropriately labeled. Similar connections between $n\sigma^{\text{pair}}$ and m_{TOF}^2 are shown in Fig. 3.14.




Figure 3.14: Two-dimensional distributions of $n\sigma_{\pi}^{\text{pair}}$ (top row), $n\sigma_K^{\text{pair}}$ (middle row) and $n\sigma_p^{\text{pair}}$ (bottom row) vs. m_{TOF}^2 . The left column contains all clean BBC-large events with single TOF vertex and two opposite sign TOF-matched tracks (passing cuts C1, C3.1, C3.2 and C6), which provides excellent statistics to see the signatures or pairs of specific ID. The right column contains exclusive event candidates after full event selection except PID cuts (except cuts C8). Dashed red line and arrow indicate the cut imposed on plotted quantities which are used to select exclusive pairs of given particle species (keep in mind that these are not the only cuts).




Figure 3.15: Raw distributions of $\chi^2_{dE/dx}$ (left column) and m_{TOF}^2 (right column) for exclusive $\pi^+\pi^-$ (top row), K^+K^- (middle row) and $p\bar{p}$ (bottom row) candidates after full event selection. Dashed red line and arrow indicate the value of cut imposed on plotted quantity to select exclusive pairs of given particle species. Presented distributions were obtained after all the cuts were applied, except the cut on presented quantity in the last step in PID algorithm used to select pairs of given species.

3.2.8 (C9) Exclusivity cut (missing p_T cut)

The most important cut which is used in this analysis to select events of exclusively produced pairs of particles is the missing transverse momentum, or the total transverse momentum cut. It benefits from detection and reconstruction of the forward proton in RP detectors - a rare capability among high energy physics experiments which STAR provides. The observable p_T^{miss} used to select exclusive event is defined as

$$p_T^{\text{miss}} = \left(\vec{p}_{p'}^E + \vec{p}_{h^+} + \vec{p}_{h^-} + \vec{p}_{p'}^W \right)_T = \sqrt{\left(p_x^{\text{miss}} \right)^2 + \left(p_y^{\text{miss}} \right)^2}, \quad (3.12)$$

with the other total momentum components defined analogously:

$$p_x^{\text{miss}} = \left(\vec{p}_{p'}^E + \vec{p}_{h^+} + \vec{p}_{h^-} + \vec{p}_{p'}^W \right)_x, \quad (3.13) \quad p_y^{\text{miss}} = \left(\vec{p}_{p'}^E + \vec{p}_{h^+} + \vec{p}_{h^-} + \vec{p}_{p'}^W \right)_y. \quad (3.14)$$

Figure 3.16 visualize the (anti-)correlation between the momentum components of the forward system (sum of two forward protons momenta) and the central system (sum of two central tracks momenta). The enhanced band at anti-diagonal restricted by dashed lines contains events balanced in momentum, a signature of exclusivity. Events outside this band are the non exclusive backgrounds, in most cases Central Diffraction events with some particles undected (due to detector inefficiency or produced outside acceptance). Slight horizontal enhancement in all distributions around $[\vec{p}_{p'}^W + \vec{p}_{p'}^E]_x = [\vec{p}_{p'}^W + \vec{p}_{p'}^E]_y = 0$ is a signature of the elastic proton-proton scattering background with some non-elastic pile-up interaction which mimics the CEP event. All these backgrounds are reasonably low after the exclusivity cut, as described in Sec. 4.1.1.

The momentum balance is shown one-dimensionally in Fig. 3.17, with the sum of x - and y -components of momentum shown respectively in the left and right column for each analyzed particle species. The sum of signal and background (both assumed to be described a Gaussian) was fitted to p_x^{miss} and p_y^{miss} distributions. Results of the fit are given in each subfigure. One can notice that the widths of Gaussian functions representing the exclusive signal are consistent among species and amount $\sigma_{p_x^{\text{miss}}} = 27.4$ MeV for the x -component of total momentum, and $\sigma_{p_y^{\text{miss}}} = 28.1$ MeV for the y -component of total momentum, taking the values of the lowest statistical uncertainty - for $\pi^+\pi^-$. These values are measures of the total momentum resolution respectively for p_x^{miss} and p_y^{miss} . Having these number it is possible to form an elliptical on the missing momentum:

$$\left(\frac{p_x^{\text{miss}}}{\sigma_{p_x^{\text{miss}}}} \right)^2 + \left(\frac{p_y^{\text{miss}}}{\sigma_{p_y^{\text{miss}}}} \right)^2 < n_{\text{cut}}^2 \quad (3.15)$$

where n_{cut} is the parameter denoting radius of limiting ellipsis in units of standard deviations of distributions of total momentum components (resolutions). Since these resolutions are nearly identical ($\sigma_{p_x^{\text{miss}}} = \sigma_{p_y^{\text{miss}}} = \sigma_{p_{x,y}^{\text{miss}}}$) such cut can be reduced (multiplying Ineq. 3.15 by $\sigma_{p_{x,y}^{\text{miss}}}^2$) to one-dimensional cut on a single quantity:

$$(p_x^{\text{miss}})^2 + (p_y^{\text{miss}})^2 < \left(n_{\text{cut}} \cdot \sigma_{p_{x,y}^{\text{miss}}} \right)^2 \quad \xrightarrow{\sqrt{}} \quad p_T^{\text{miss}} < n_{\text{cut}} \cdot \sigma_{p_{x,y}^{\text{miss}}} \quad (3.16)$$

In current analysis the n_{cut} was set to 2.5, which translates to threshold value 2.5×30 MeV = 75 MeV.




Figure 3.16: Two-dimensional distributions of sum of forward protons momenta (x -axis) and sum of central tracks momenta (y -axis) for exclusive $\pi^+\pi^-$ (top row), K^+K^- (middle row) and $p\bar{p}$ (bottom row) event candidates after full event selection, except the exclusivity cut C9. Left and right column shows correlation of respectively x - and y -component of tracks' momenta. Anti-diagonal representing perfect momentum balance of the central and forward system is limited with dashed lines extending by $\pm 2.5\sigma$ ($\sigma \approx 30$ MeV) around the anti-diagonal. Three distinct horizontal regions in plots on the right hand side correspond to different forward proton configurations: elastic-like (protons in branches EU&WD or ED&WU, $|[\vec{p}_p^W + \vec{p}_p^E]_y| < 0.2$ GeV) and anti-elastic configuration (protons in branches ED&WD or EU&WU, $|[\vec{p}_p^W + \vec{p}_p^E]_y| > 0.4$ GeV).




Figure 3.17: Raw distributions of $p_{\text{miss}}^{\text{x}}$ (left column) and $p_{\text{miss}}^{\text{y}}$ (right column) for exclusive $\pi^+\pi^-$ (top row), K^+K^- (middle row) and $p\bar{p}$ (bottom row) candidates after full event selection, except exclusivity cut C9. Solid red line represents the fit of sum of two Gaussian functions representing the exclusive event signal (orange) and non-exclusive background (violet). Parameters of the total momentum resolution for signal events obtained from the fit (given in the plots) roughly agree between all species.




Figure 3.18: Uncorrected distributions of the CEP event candidates for missing transverse momentum p_T^{miss} for $\pi^+\pi^-$ (top), K^+K^- (middle) and $p\bar{p}$ (bottom) pairs. Distributions for opposite-sign and same-sign particle pairs are shown as black and red symbols, respectively. The vertical error bars represent statistical uncertainties. The horizontal bars represent bin sizes. Distribution for $\pi^+\pi^-$ channel with MC predictions for both signal and background can be found in Fig. 4.7a.

4. Backgrounds

4.1 Sources of background

4.1.1 Non-exclusive background

The main background present in the final exclusive $\pi^+\pi^-/K^+K^-/p\bar{p}$ sample is the non-exclusive background. There are several classes of events which mimic topology of h^+h^- CEP: two forward protons, two opposite sign central tracks and rapidity gaps. Below we list the most probable cases:

- Single physics processes:
 - Central Diffraction (Fig. 4.1b) - this process differs from CEP of h^+h^- only by the number of particles produced in the mid-rapidity; protons originate from the same vertex as the central tracks, hence correlation of reconstructed vertex position from RPs and TPC is still observed.
- Coincidences (pile-up):
 - inelastic + elastic interaction (Fig. 4.1c) - there may be overlap of protons from elastic scattering interaction and activity in the central detector from another (inelastic) interaction; it should be suppressed by the rapidity gap veto in BBC-small (online) and BBC-large (offline); easy to identify through protons collinearity and lack of correlation of z -vertex from RPs and TPC.
 - Single Diffraction + beam halo - there may be overlap of proton from SD on one side and beam halo proton on the opposite side, and activity in the central detector from diffractive state; it should be suppressed by the rapidity gap vetos and (low) beam halo rate;
 - 2×beam halo + inelastic interaction.

} negligibly low

These backgrounds are graphically presented in Fig. 4.1.




Figure 4.1: Sketches of main processes exhibiting h^+h^- CEP event topology: the exclusive h^+h^- signal itself (a), central diffraction event with some particles not detected (b) and elastic proton-proton scattering event with pile-up inelastic interaction in the central region (c). Particles represented by arrows are: forward scattered protons (blue), detected mid-rapidity particles (green) and undetected particles (dashed gray). Black dots mark primary interaction vertices.

4.1.2 Exclusive background (particle misidentification)

Another source of background which is connected with finite particle identification power is the exclusive background from the particle species other than species under study.




Figure 4.2: Graph illustrating the misidentification problem - the origin of exclusive background in selected samples. Gray arrows represent event rejection due to failed PID selection (C8). Magenta arrows indicate non-exclusive backgrounds described in Sec. 4.1.1. Solid black arrows represent successful identification, whereas dashed black lines show misidentification paths.

$$N_R^{\pi\pi} = \underbrace{\epsilon^{\pi\pi} \cdot N_T^{\pi\pi}}_{\text{true pion pairs}} + \underbrace{\lambda^{KK \rightarrow \pi\pi} \cdot N_T^{KK}}_{\text{kaon pairs reconstructed as pion pairs}} + \underbrace{\lambda^{p\bar{p} \rightarrow \pi\pi} \cdot N_T^{p\bar{p}}}_{\text{proton pairs reconstructed as pion pairs}} + \textcolor{magenta}{N_{bkgd}^{\pi\pi}} \quad (4.1a)$$

$$N_R^{KK} = \underbrace{\lambda^{\pi\pi \rightarrow KK} \cdot N_T^{\pi\pi}}_{\text{pion pairs reconstructed as kaon pairs}} + \underbrace{\epsilon^{KK} \cdot N_T^{KK}}_{\text{true kaon pairs}} + \underbrace{\lambda^{p\bar{p} \rightarrow KK} \cdot N_T^{p\bar{p}}}_{\text{proton pairs reconstructed as kaon pairs}} + \textcolor{magenta}{N_{bkgd}^{KK}} \quad (4.1b)$$

$$N_R^{p\bar{p}} = \underbrace{\lambda^{\pi\pi \rightarrow p\bar{p}} \cdot N_T^{\pi\pi}}_{\text{pion pairs reconstructed as proton pairs}} + \underbrace{\lambda^{KK \rightarrow p\bar{p}} \cdot N_T^{KK}}_{\text{kaon pairs reconstructed as proton pairs}} + \underbrace{\epsilon^{p\bar{p}} \cdot N_T^{p\bar{p}}}_{\text{true proton pairs}} + \textcolor{magenta}{N_{bkgd}^{p\bar{p}}} \quad (4.1c)$$

Eqs. (4.1) can be written in the matrix form, as shown in Eq. (4.2), from which it is straightforward to obtain final formula for unfolded number of events of given ID, Eq. (4.3):

$$\begin{bmatrix} N_R^{\pi\pi} - \textcolor{magenta}{N_{bkgd}^{\pi\pi}} \\ N_R^{KK} - \textcolor{magenta}{N_{bkgd}^{KK}} \\ N_R^{p\bar{p}} - \textcolor{magenta}{N_{bkgd}^{p\bar{p}}} \end{bmatrix} = \underbrace{\begin{bmatrix} \epsilon^{\pi\pi} & \lambda^{KK \rightarrow \pi\pi} & \lambda^{p\bar{p} \rightarrow \pi\pi} \\ \lambda^{\pi\pi \rightarrow KK} & \epsilon^{KK} & \lambda^{p\bar{p} \rightarrow KK} \\ \lambda^{\pi\pi \rightarrow p\bar{p}} & \lambda^{KK \rightarrow p\bar{p}} & \epsilon^{p\bar{p}} \end{bmatrix}}_{\text{"mixing matrix" } \Lambda} \begin{bmatrix} N_T^{\pi\pi} \\ N_T^{KK} \\ N_T^{p\bar{p}} \end{bmatrix} \rightarrow \begin{bmatrix} N_T^{\pi\pi} \\ N_T^{KK} \\ N_T^{p\bar{p}} \end{bmatrix} = \Lambda^{-1} \begin{bmatrix} N_R^{\pi\pi} - \textcolor{magenta}{N_{bkgd}^{\pi\pi}} \\ N_R^{KK} - \textcolor{magenta}{N_{bkgd}^{KK}} \\ N_R^{p\bar{p}} - \textcolor{magenta}{N_{bkgd}^{p\bar{p}}} \end{bmatrix} \quad (4.2)$$

4.2 Non-exclusive background determination

Determination of non-exclusive background makes use of the missing transverse momentum which is distributed differently for the signal and for aforementioned background. High statistics of the data allows to apply a data-driven method. What is more, not only integrated over observables but even differentially as a function of these.

It has already been demonstrated in Sec. 3.2.8 that p_T^{miss} from exclusive events is much narrower compared to background, visible as a peak near the axis origin. Additional feature of p_T^{miss} - probability density approaching to zero with p_T^{miss} moving towards the axis origin, enables performing a polynomial fit to p_T^{miss} distribution in the background-dominated range and extrapolation of this polynomial down to $p_T^{\text{miss}} = 0$ under the peak of CEP signal. The procedure used to determine non-exclusive background content in the signal region is described below.

4.3 Normalization of signal and background models

Consistency between data and MC, valuable to demonstrate good understanding of the backgrounds and data themselves, has been tested for exclusive $\pi^+\pi^-$ channel¹. We considered here only non-exclusive backgrounds because of negligible contribution from misidentifications.

The following MC samples were used in this study:

¹Other channels - K^+K^- and $p\bar{p}$ - were not subjected to similar study because of poor MC statistics. However, structure of backgrounds and level of agreement with MC is expected to be very similar to that presented for $\pi^+\pi^-$.

- Exclusive $\pi^+\pi^-$ (signal) - events from GenEx [21] generator passed through Geant3 simulation of the STAR detector (STARsim) and Geant4 simulation of the RP Phase II* detectors, fully embedded into zero-bias data,
- CD (background) - Central Diffraction events from Pythia 8.1 generator, filtered at generation to ensure lack of signal in BBC-large, passed through Geant3 simulation of the STAR detector (STARsim) and Geant4 simulation of the RP Phase II* detectors, partially embedded into zero-bias data (only the simulated RP response embedded),
- MB+elastic, (background) - Minimum Bias events from Pythia 8.1 generator filtered at generation to ensure lack of signal in BBC-large, passed through Geant3 simulation of the STAR detector (STARsim) and Geant4 simulation of the RP Phase II* detectors, partially embedded into elastic trigger (RP_ET) data (only the simulated RP response embedded).

Listed background samples were not fully embedded since an enormous CPU time would be required to obtain satisfactory statistics. It was also found unnecessary to embed TPC tracks into zero-bias data to obtain reliable agreement between distributions of desired quantities presented below.

MC samples from Pythia generator were additionally filtered before passing through Geant to increase generation efficiency, as well as overcome difficulty arising from missing simulation of the BBC-large in STARsim. For each event, all charged particles were analytically propagated through the magnetic field of TPC with the helical paths resulting from their hadron-level momenta. If any of these particles crossed the volume of BBC-large detector, event was dropped from generation.

Normalization of backgrounds was done separately for two ranges of $\Delta\varphi$. First, MB+elastic background was normalized. By definition this was done only for $\Delta\varphi$ bin representing elastic-like configuration of forward protons ($\Delta\varphi > 90^\circ$). The MB+elastic MC was scaled to have the same integral as the data in range $|\Delta z_{\text{vtx}}| > 100$ cm. In this range we assumed sole presence of this type of background, which is characterized by very wide distribution of Δz_{vtx} because of TPC and RP vertices being independent. Comparison plots are contained in Figs. 4.3 and 4.4. An important cross-check for correctness of this assumption is shown in Fig. 4.5, where the data vs. MC collinearity $\Delta\theta$ is presented, defined as

$$\Delta\theta = \sqrt{(\Delta\theta_x)^2 + (\Delta\theta_y)^2} = \sqrt{(\theta_x^W + \theta_x^E)^2 + (\theta_y^W + \theta_y^E)^2}. \quad (4.4)$$

One can notice part of distribution close to 0, with nearly perfectly collinear protons. The data is well described by MC, which would unlikely be the case without contribution from the red histogram representing MB+elastic background. An interesting observation related to this background contribution is that almost all MB+elastic events in the final plots originate from the Central Diffraction process, with the forward protons outside of RP acceptance - non-diffractive events do not pass tight CEP event selection.

In the second step the CD MC was normalized. It was scaled to have the same integral as the data (minus MB+elastic MC in $\Delta\varphi > 90^\circ$ sub-sample) in range $p_T^{\text{miss}} > 150$ MeV, where no exclusive signal is expected.

In the last step the exclusive $\pi^+\pi^-$ MC was normalized. It was scaled to have the same integral as the data (minus all considered non-exclusive backgrounds) in range $p_T^{\text{miss}} < 75$ MeV, where exclusive signal is dominant. The result of this procedure for the distribution of p_T^{miss} is given in Figs. 4.6 and 4.7. Joint distribution for each quantity - without differentiation with respect to $\Delta\varphi$ - was obtained by adding corresponding event counts from two $\Delta\varphi$ ranges.

As can be observed in the comparison plots, presented data are generally well described by MC. In case of Δz_{vtx} some imperfection in the position and width of the simulated signal peak can be noticed, most probably arising from slightly underestimated timing resolution of the RP trigger counters in the simulation. Distribution of p_T^{miss} is quite well described, for both signal and control channel. The ratio of number of opposite-sign pairs to same-sign pairs is compatible between data and MC, which was possible to achieve by rejecting in Pythia contributions from events with the central state consisting from two opposite-sign pions and at least one neutral particle. These events should be suppressed by the DIPE condition (1.5), which seem to be not taken into account in Pythia at the hadronization level. We demonstrate in Fig. 4.7c that if these events are preserved, Pythia MC cannot describe data in the background-dominating region (large p_T^{miss}).






Figure 4.3: Comparison of Δz_{vtx} for CEP $\pi^+\pi^-$ events in two ranges of $\Delta\varphi$ (left: $\Delta\varphi < 90^\circ$, right: $\Delta\varphi > 90^\circ$) between data and embedded MC after full selection (except cut on presented quantity). Plots in top and bottom row differ only in the y -axis (top: logarithmic, bottom: linear). Data are represented by black points, while stacked MC predictions are drawn as histograms of different colors. Histogram from each MC process has been normalized according to prescription in the text. Vertical error bars represent statistical uncertainties, horizontal bars represent bin sizes.



(a)



(b)

Figure 4.4: Comparison of Δz_{vtx} for CEP $\pi^+ \pi^-$ events between data and embedded MC after full selection (except cut on presented quantity). Left and right plot differ only in the y-axis (left: logarithmic, right: linear). Data are represented by black points, while stacked MC predictions are drawn as histograms of different colors. Histogram from each MC process has been normalized according to prescription in the text. Vertical error bars represent statistical uncertainties, horizontal bars represent bin sizes.




Figure 4.5: Comparison of colinearity $\Delta\theta$ for CEP $\pi^+ \pi^-$ events with $\Delta\varphi > 90^\circ$ between data and embedded MC after full selection. Data are represented by black points, while stacked MC predictions are drawn as histograms of different colors. Histogram from each MC process has been normalized according to prescription in the text. Vertical error bars represent statistical uncertainties, horizontal bars represent bin sizes.




Figure 4.6: Comparison of p_T^{miss} for CEP $\pi^+\pi^-$ events in two ranges of $\Delta\varphi$ (left: $\Delta\varphi < 90^\circ$, right: $\Delta\varphi > 90^\circ$) between data and embedded MC after full selection (except cut on presented quantity). Plots in top and bottom row differ only in the y -axis (top: linear, bottom: logarithmic). In addition to signal channel (opposite-sign particles) also control background channel (same-sign particles) is contained in the plots. Data are represented by black (opposite-sign) or red (same-sign) points, while stacked MC predictions are drawn as filled (opposite-sign) or hatched (same-sign) histograms of different colors. Histogram from each MC process has been normalized according to prescription in the text. Vertical error bars represent statistical uncertainties, horizontal bars represent bin sizes.






Figure 4.7: Comparison of p_T^{miss} for CEP $\pi^+\pi^-$ events between data and embedded MC after full selection (except cut on presented quantity). Top left and top right plot differ only in the y -axis (left: linear, right: logarithmic). In addition to signal channel (opposite-sign particles) also control background channel (same-sign particles) is contained in the plots. Data are represented by black (opposite-sign) or red (same-sign) points, while stacked MC predictions are drawn as filled (opposite-sign) or hatched (same-sign) histograms of different colors. Histogram from each MC process has been normalized according to prescription in the text. Vertical error bars represent statistical uncertainties, horizontal bars represent bin sizes.

Left bottom plot differ from the top plots in the content of Pythia MCs - in the bottom plot events with $\pi^\pm\pi^\mp + \text{neutrals}$ in the central state were preserved, demonstrating significant inconsistency between data and MC in the ratio of opposite-sign to same-sign events if such events are not rejected.

5. Corrections

5.1 Method of corrections application

$$\frac{d\sigma}{dq} = \frac{1}{\Delta q} \times \frac{1}{\varepsilon} \times \frac{N^w - N_{\text{bkgd}}^w}{L_{\text{int}}^{\text{eff}}} \quad (5.1)$$

$$L_{\text{int}}^{\text{eff}} = \sum_{\text{run}} L_{\text{int}}^{\text{run}} \times \epsilon_{\text{veto}}(L^{\text{run}}) \quad (5.2)$$

$$\varepsilon = \epsilon_{\text{ET/IT}} \times \epsilon_{\text{vrtx}}(q) \times \epsilon_{C2} \times \epsilon_{C5} \times \epsilon_{C9} \times \epsilon_{\text{PID}}(q) \quad (5.3)$$

$$N^w = \sum_{\text{event}} w_{\text{event}} \quad (5.4)$$

$$w = \left[\prod_{\text{sign}} \epsilon_{\text{TOF}}(\text{sign}, \text{PID}, p_T, z_{vx}, \eta) \times \prod_{\text{sign}} \epsilon_{\text{TPC}}(\text{sign}, \text{PID}, p_T, z_{vx}, \eta) \times \prod_{\text{side}} \epsilon_{\text{RP}}^{\text{side}}(p_x, p_y) \right]^{-1}, \quad (5.5)$$

sign $\in \{+, -\}$, side $\in \{E, W\}$




Figure 5.1: Distribution of weights assigned to selected CEP events in $\pi^+\pi^-$ (black circle), K^+K^- (red square) and $p\bar{p}$ (blue triangle) channel.

5.2 Acceptances and efficiencies

In this section we present calculation of all efficiencies except TPC track reconstruction and TOF hit reconstruction and matching efficiency, which were discussed and presented in Ref. [1].

5.2.1 Trigger efficiency

5.2.1.1 Online TOF efficiency

The efficiency of the part of the trigger related to TOF subsystem $\mathcal{E}_{\text{TOF}}^{\text{trig}}$ was calculated using the zero-bias data. Events with exactly 1 TOF vertex (cut C1), exactly 2 primary good quality TOF-matched TPC tracks (cuts C3.1, C3.4) and maximally 3 TOF clusters (cut C7) were used to calculate this efficiency defined as a probability of having at least two TOF hits on the trigger level in selected set of events:

$$\mathcal{E}_{\text{TOF}}^{\text{trig}} = \frac{\#\text{events with } \geq 2 \text{ L0 TOF multiplicity and } 1 \text{ TOF vtx and } 2 \text{ TOF trks and } \leq 3 \text{ TOF cltrs}}{\#\text{events with } 1 \text{ TOF vtx and } 2 \text{ TOF trks and } \leq 3 \text{ TOF cltrs}} \quad (5.6)$$




Figure 5.2: TOF trigger efficiency as a function of lower p_T of two TOF-matched tracks. Black points denote efficiency of requirement of at least 2 online TOF hits, while open blue circles represent efficiency with added upper multiplicity limit (10). These efficiencies are identical therefore the same efficiency (98.7%, solid red line) is used as a correction factor for all analyzed events.

It has been verified that described efficiency does not depend on the transverse momentum of two tracks. In Fig. 5.2 we show dependence of the TOF trigger efficiency in one dimension, as a function of lower p_T of two TOF-matched tracks. Data points are consistent with constant value, herefore a single number for this efficiency is used, equal to 98.7%. It is important to account for the fact that at some point of data-taking (see Tab. 2.1) an upper limit (< 10) on L0 TOF multiplicity was imposed. Study shows that with the cut on offline TOF cluster multiplicity (cut C7) the efficiency of the trigger with and without upper limit of online TOF multiplicity remains unchanged, so the same value of the efficiency is used for the entire dataset.

5.2.1.2 Online veto (BBC-small and ZDC veto)

Vetoeing signal in BBC-small and ZDC detectors on both sides of STAR was implemented in the logic of RP_CPT2 trigger. Common correction of the online and offline vetoes which is used in the correction procedure explained in Sec. 5.1 is presented in Sec. 5.2.2.3. However, to help quantify effect of just the online vetoes in BBC-small and ZDCs we show the Fig. 5.3 with the efficiency of the joint BBC-small and ZDC veto as a function of the instantaneous luminosity calculated from the zero-bias data. Details of the way the efficiency was calculated as well as description of the data in the Figure is the same as explained in Sec. 5.2.2.3.




Figure 5.3: Overall efficiency of the online BBC-small and ZDC veto as a function of instantaneous luminosity.

5.2.1.3 RP triggering efficiency

Based on preliminary studies preformed during data taking using fast offline data (MuDst) we concluded that the RP triggering efficiency is very close to 100 % and eventual correction (which is going to be evaluated) is much smaller than overall systematic uncertainty. For the time being we assume that

$$\mathcal{E}(\text{TR}^E \wedge \text{TR}^W) = \mathcal{E}(\text{TR}^W) \times \mathcal{E}(\text{TR}^E) = 1. \quad (5.7)$$

5.2.1.4 Up and Down RP combination veto (due to dead material)

Probability that secondaries induced by proton with successfully reconstructed and selected RP track generate a trigger signal in the other RP branch on the same side was calculated using the embedded MC (see Sec. 6.3 in Ref. [1] for details of RP simulation in Geant4). Forward protons from CEP process provided by GenEx [21] were simultaneously generated from the interaction point spatially distributed the same as in the data. MC samples for all runs with RP_CPT2 triggers were produced to account for non-constant positions of the RP detectors throughout the run 15. Number of simulated events for each run was proportional to number of RP_CPT2 triggers in given run. The angular divergence of the beams was also simulated.

The discussed probability, $\mathcal{P}_{\text{DM veto}}^{\text{side}}$, was calculated as a probability that a MC-trigger signal is present in the branch on given side other than east and west branches where primary forward protons are expected from their initial momenta, under condition that these east and west branches detect a MC-trigger signal and there is no veto due to simultaneous ET&IT trigger bits in the overlayed data (no pile-up veto). By MC-trigger we understand the trigger signal reconstructed solely from the simulated data (not from the data embedded into).

Technically the $\mathcal{P}_{\text{DM veto}}^{\text{side}}$ was obtained in the following procedure:

1. No simultaneous ET&IT trigger bits were allowed in the data of an event that simulated signal was embedded into.
2. It was verified if there are MC-trigger signals in east and west branches that the primary forward protons were expected to reach based on their p_y ($p_y > 0$ - branch UP, $p_y < 0$ - branch DOWN). These events formed set *A*.
3. Events with the MC-trigger signal in RP branch other than the branch with MC-trigger signal expected from proton p_y on given side formed set *B*.
4. The probability was determined by the ratio of histograms from set *B* and set *A*:

$$\begin{aligned} \mathcal{P}_{\text{DM veto}}^{\text{side}}(p_x, p_y, z_{\text{vtx}}) &= \mathcal{E}^{\text{side}}(\text{Veto}^{\text{DM}} | \text{!Veto}^{\text{PU}} \wedge \text{TR}^E \wedge \text{TR}^W) = \\ &= \frac{(p_x, p_y, z_{\text{vtx}}) \text{ histogram for protons from set } B}{(p_x, p_y, z_{\text{vtx}}) \text{ histogram for protons from set } A} \end{aligned} \quad (5.8)$$

It should be noted that the momentum components (p_x, p_y) were taken from the proton with accounted effect of the beam divergence (after the original initial momentum smearing). Sample probability of a dead-material-induced veto is shown in Fig. 5.4, with all the remaining results contained in Appendix D.

The efficiency of the discussed veto which is finally used to correct the data is opposite of the veto probability, namely

$$\epsilon_{\text{DM veto}}^{\text{side}} = 1 - \mathcal{P}_{\text{DM veto}}^{\text{side}}. \quad (5.9)$$




Figure 5.4: Sample probability of ET&IT trigger veto due to forward proton interaction with dead material. Results were obtained from forward proton MC simulation embedded into zero-bias data.

5.2.2 Reconstruction and selection efficiency

5.2.2.1 TPC z -vertex cut (C2)

Removing from analysis specific range of z -positions of primary vertices effectively reduces accepted luminosity with respect to that delivered by the collider. This loss of luminosity has to be accounted when calculating the cross sections, which is the goal of presented analysis. Assuming that the distribution of z_{vtx} of all primary interactions that take place when east and west beams overlap follows a normal distribution (\mathcal{N}), then the formula describing efficiency of the cut C2 has the following form:

$$\epsilon_{z_{\text{vtx}}} = \int_{z_{\text{vtx}}^{\min}}^{z_{\text{vtx}}^{\max}} \mathcal{N}(z_{\text{vtx}}; \mu, \sigma) dz_{\text{vtx}} = \frac{1}{2} \left[\text{Erf} \left(\frac{z_{\text{vtx}}^{\max} - \mu}{\sqrt{2}\sigma} \right) - \text{Erf} \left(\frac{z_{\text{vtx}}^{\min} - \mu}{\sqrt{2}\sigma} \right) \right], \quad (5.10)$$

where z_{vtx}^{\min} and z_{vtx}^{\max} are respectively minimum and maximum value of the longitudinal position of the vertex accepted in analysis (in our case these are -80 cm and 80 cm), and parameters μ and σ are respectively the mean and standard deviation of the normal distribution:

$$\mu = \langle z_{\text{vtx}} \rangle, \quad \sigma = \sigma(z_{\text{vtx}}).$$

Real parameters of z_{vtx} distribution were studied separately for every single fill of the collider. It is motivated by the fact that each fill of the machine is nearly independent from the previous thus the shape and position of z_{vtx} may vary on fill by fill basis. We neglect possible changes within the fill (e.g. widening of the distribution due to intrabeam scattering etc.) arguing that this effect if expected to be smaller than the systematic uncertainties related to determination of position and width of z_{vtx} distribution.

For every fill of RHIC the distribution of z_{vtx} of single TOF vertices was prepared, as shown in Fig. 5.5. This distribution was fitted with the Gaussian function in a range $z_{\text{vtx}} \in [-120 \text{ cm}, 120 \text{ cm}]$. The output parameters of all fits were plotted as a function of the fill number, as shown in Fig. 5.6. The efficiency used in the correction procedure was calculated independently for each fill using presented values of $\langle z_{\text{vtx}} \rangle$ and $\sigma(z_{\text{vtx}})$. Typical numerical value of this efficiency equals $\sim 88\%$.




Figure 5.5: Sample distribution of z_{vtx} of single TOF vertices together with the fit of normal distribution (dashed blue) extended outside the range of the fit (solid red). Hashed red area represents part of the distribution rejected by cut C2 with the cut value marked with dashed red vertical lines and arrows.




Figure 5.6: Mean (top panel) and width (bottom panel) parameters of the normal distribution obtained from a fit of normal distribution to z_{vtx} distribution in a range $[-120 \text{ cm}, 120 \text{ cm}]$ for each RHIC fill as a function of the fill number.

5.2.2.2 TPC-RP z -vertex matching (C5)

5.2.2.3 Primary vertices limit (C1), BBC-large veto (C6), TOF clusters limit (C7) and Up and Down RP combination veto (due to pile-up)

Combined efficiency of the online veto in BBC-small and ZDC (Sec. 5.2.1.2) and offline cuts (vetoes) on extra TPC-TOF vertices, extra TOF clusters, signal in BBC-large and simultaneous signal in Up and Down RPs, was calculated using the zero-bias data. For each run a fraction of events (for colliding bunches) was calculated in which all mentioned cuts would be satisfied in case of the CEP $\pi^+\pi^-/K^+K^-/p\bar{p}$ event (event would not be vetoed). One can transform this prescription to simple formula below:

$$\epsilon_{b_E b_W}^{\text{veto}} = \frac{\# \text{events in the run without TOF vertices, without signal in BBC-S, BBC-L, ZDC,}}{\text{RP branches other than } b_E, b_W, \text{ and with no more than 1 reconstructed TOF cluster}} \quad (5.11)$$

In Fig. 5.7 this efficiency is presented as a function instantaneous luminosity delivered by the machine, for each combination of east and west RP branches. Result for each combination is nearly identical as the effect of ET&IT trigger veto in RPs is not dominant, as well as trigger in all branches had similar acceptance. The data points were fitted with the exponential function (of the form contained in the figure) which reflects the fact that this efficiency should behave similar to the probability of lack of any interaction in the bunch crossing given by the Poisson distribution:

$$\text{Pois}(0; \mu) = \frac{\mu^0}{0!} \times e^{-\mu} = e^{-\mu}. \quad (5.12)$$

Comparison of the μ in Eq. (5.12) with the fit parameters in Fig. 5.7 leads to approximate determination of the average interaction probability per bunch crossing equal $0.2 - 0.9$. The result of the fit, $\epsilon_{b_E b_W}^{\text{veto}}(\mathcal{L})$, is finally used to correct measured data as described in Sec. 5.1.

Comparison of efficiencies in Fig. 5.7 with similar efficiency in Fig. 5.3 demonstrates that offline selection has much smaller impact on the loss of signal events than online selection. It has to be underlined that online vetoes were necessary to set trigger purity to satisfactory level, as well as reduce prescale of the trigger.




Figure 5.7: Overall efficiency of the online BBC-small, ZDC and ET&IT trigger veto, primary vertices limit (C1), BBC-large veto (C6) and TOF clusters limit (C7) as a function of instantaneous luminosity for all possible combinations of east and west RP branches. Red and blue points represent runs lasting for less and more than 20 minutes, respectively. Black dotted lines represent fits of exponential functions to blue points.

5.2.2.4 Missing p_T cut (C9)

The resolution of the total transverse momentum is in general determined by the angular divergence of the proton beams. This angular divergence is connected with imperfect collimation of the beams - protons do not circulate in RHIC along a single closed line, they rather move within certain volume¹. Based on known parameters of the beams in run 15 (see Ref. [22]) expected divergence for a single beam (and single transverse spatial coordinate) amounts 180 μrad. In forward proton track reconstruction the direction of the incoming proton is assumed to be the same in each event (aligned with nominal z -axis), hence effectively the transverse components of reconstructed forward protons momenta are smeared by this amount. One can check the comparison of the collinearities of elastically scattered protons in the data and embedded MC (Fig. 10.19 of Ref. [1]) to see the satisfactory agreement between the two with assumed MC divergence equal to 180 μrad. For 100 GeV beam such divergence results in smearing of the transverse momentum components for (elastically) scattered protons equal $100 \text{ GeV} \times 180 \cdot 10^{-6} = 18 \text{ MeV}$. Using this value to calculate the resolution of total $p_x(p_y)$ momentum of two forward protons we get $\sqrt{2} \times 18 \text{ MeV} = 25.5 \text{ MeV}$. One can compare it with the output of fits to distributions of $p_x^{\text{miss}}(p_y^{\text{miss}})$ and find that indeed the angular beam divergence dominates the resolution of the total transverse momentum in CEP events.

There is, however, another ingredient to the total transverse momentum resolution, namely the momentum resolution of the central tracks, whose significance rises up together with increasing track p_T . Most of CEP events is characterized by low invariant mass of the central tracks pair ($\lesssim 1.5 \text{ GeV}$) which is inextricably linked with low momentum of tracks ($\lesssim 0.7 \text{ GeV}$), therefore an effect of p_T^{miss} widening is barely visible in the missing momentum distribution integrated over mass (e.g. Fig. 3.17). Nevertheless one can calculate the efficiency of the exclusivity cut as a function of the central tracks momenta to directly see this effect.

Physics model of the CEP process from GenEx was used in embedded signal MC, therefore there was no

¹In other words, their phase-space during orbitation is not point-like, it is elliptic. The area of this elliptic phase space is called emittance. The lower the emittance is, the better collimation of the beam is and lower transverse size of the beam is.

sufficient statistics (too low population of high- p_T tracks) to calculate the efficiency of p_T^{miss} cut as a function of central tracks momenta. In such case the efficiency of this cut was calculated using simplified MC method described below.

A simple CEP event generator was used to produce large sample of CEP events (kinematics of protons and pions) in the fiducial phase space of the measurement. For each event the hadronic level momenta of positive and negative central tracks were smeared according to resolution function obtained from the embedded MC (6 MeV if $p_T < 0.3$ GeV, $2.4 \text{ MeV} + 1.2\% \times p_T$ if $p_T > 0.3$ GeV). Next, the p_x^{miss} and p_y^{miss} were calculated from these smeared pions momenta and hadronic level forward protons momenta. Both p_x^{miss} and p_y^{miss} were added the Gaussian component corresponding to the experimental resolution extracted in Figs. 3.17a and 3.17b (means and standard deviations printed in the plots were used). At the end, p_T^{miss} was calculated with the use of Eq. (3.12). The efficiency of the exclusivity cut was defined as a following ratio:

$$\epsilon_{p_T^{\text{miss}}}(p^{\max}, p^{\min}) = \frac{N_{p_T^{\text{miss}}}(p^{\max}, p^{\min})}{N_{\text{all}}(p^{\max}, p^{\min})}, \quad (5.13)$$

where N_{all} is a number of all generated CEP events and $N_{p_T^{\text{miss}}}$ is number of events passing exclusivity cut C9. The result is shown in Fig. ???. The cut efficiency for very low tracks momenta (low pair invariant mass) obtained from simple MC (96.8%) agrees perfectly with the efficiency calculated using embedded MC sample for the same momenta range (96.4%), which validates presented efficiency. Clearly, the deterioration of the central track momentum resolution starts to play significant role in efficiency of p_T^{miss} cut at about 1 GeV track momentum.




Figure 5.8: Efficiency of p_T^{miss} cut as a function of higher (x -axis) and lower (y -axis) momentum of the central tracks, calculated with MC method described in the text of this Section. Efficiency in a few points of (p^{\min}, p^{\max}) space is printed on top of the plot. Lower right corner has no entries/low statistics due to kinematic constraints in the fiducial phase space; this region is not populated with events in the data.

5.2.2.5 Particle identification (C8)

Correction reflecting efficiency of identification requires good modeling of detector response in terms of dE/dx and TOF time ($\rightarrow m_{\text{TOF}}^2$) measurement which were used for this purpose as described in Sec. 3.2.7. In addition to this, significant number of simulated events is needed to reduce statistical uncertainties of efficiency. The former was provided by adjusting dE/dx spectra from embedded MC to match the data, as elaborated in Chap. 7 of Ref. [1]. The latter, however, was not easy to achieve for exclusive K^+K^- and $p\bar{p}$ whose identification is most challenging and information about identification efficiency is the most needed among studied species. Specially for study of particle (exclusive pair) identification a dedicated MC simulation was prepared.

This dedicated MC simulation was designed to work as follows (simulation of single CEP event of predefined pair ID is described):

1. The position of z_{vtx} was drawn from predefined distribution.
2. Kinematics of central state particles was set: momentum (magnitude) p , pseudorapidity η and azimuthal angle ϕ of positive and negative charge particles were drawn from predefined distributions.
3. Both particles were tested if doubled radius of curvature $2R$ of associated track in the magnetic field of the TPC ($B = 0.5$ T, $R \propto p_T/B$) is smaller than the radius of TOF detector barrel (assumed 212 cm). If not then event was skipped and procedure was restarted (back to 1.).




Figure 5.9: Correlation between uncertainty of the natural logarithm of $dE/dx/(1 \text{ keV}/\text{cm})$ and track TOF path length (5.9a) and correlation between base 2 logarithm of dx and track TOF path length (5.9b). The distributions were obtained for the exclusive event candidates after full selection, with all three types of particle pairs combined.

4. The particles were propagated from the vertex at $(0, 0, z_{\text{vtx}})$ through the magnetic field of TPC using Newton's method with the time step (in the laboratory) equal 100 ps, corresponding to space step $< 3 \text{ cm}$.
5. After step 4. the position of the TOF cell was known allowing to calculate the TOF path length L between the vertex and position of the TOF hit. Also the TOF hit time t was then known, further smeared by adding random number from normal distribution with mean at 0 and standard deviation $\sigma_{\text{TOF}} = 100 \text{ ps}$ to account for the finite TOF time resolution. In addition to this, reconstructed tracks' (transverse) momenta were defined as the true momenta smeared by 6 MeV if $p_T < 0.3 \text{ GeV}$ or by $2.4 \text{ MeV} + 1.2\% \times p_T$ if $p_T > 0.3 \text{ GeV}$, to account for finite TPC momentum resolution. At this stage it was possible to calculate m_{TOF}^2 using Eq. (C.8).
6. The dE/dx measurement was simulated. For each particle a dE/dx was drawn from the distribution of the form given by Eq. (7.6) and of parameters (for given particle ID and momentum) which were extracted from the data and tabulated in Tab. 7.1, all contained in Chap. 7 of Ref. [1]. This assured that the simulated dE/dx exactly matched the data. Once dE/dx for both particles (tracks) was obtained, value of the dE/dx error (more strictly: uncertainty of $\ln(dE/dx [\text{keV}/\text{cm}])$) and $\log_2(dx)$ was also set up. These quantities depend on the number of TPC hit points used in the reconstruction of dE/dx (the more hits in tracks, the better resolution of dE/dx and higher dx), which obviously is not accessible without full STAR simulation in Geant. This problem was solved by extracting dependence of $\sigma(\ln(dE/dx))$ and $\log_2(dx)$ on the TOF path length from the data (from CEP events, Fig. 5.9). Since the length of the TOF path is very strongly correlated with the number of hits forming the track and thus number of hits used to reconstruct dE/dx , one is allowed to draw $\sigma(\ln(dE/dx))$ and $\log_2(dx)$ from their distributions for particular TOF path lengths calculated in 5. and use as measured ones. In this way the simulation preserves relevant connections between dE/dx -related quantities. After these steps are taken the $n\sigma_X$ ($X = \pi, K, p$) variables are calculated for each track using the definition, in exactly the same way as it is done during standard data reconstruction.
7. Event information needed to study pair identification was stored in the ROOT tree: ID of particles forming a pair, their three-momenta, m_{TOF}^2 , $n\sigma_\pi$, $n\sigma_K$ and $n\sigma_p$.

For the purpose of determination of pair identification efficiency in CEP analysis described in this note, parameters of vertex distribution were set to match the data: $\langle z_{\text{vtx}} \rangle = 0$ and $\sigma(z_{\text{vtx}}) = 50 \text{ cm}$, as well as z_{vtx} was required to lie within the analysis limits (cut C2). Distribution of particle η was set to flat and limited to analyzed range $|\eta| < 0.7$, while particle ϕ was defined as uniformly distributed in full azimuth ($2\pi \text{ rad}$), both fairly agreeing with expectations from physics models and observations in data.




Figure 5.10: Pair identification efficiency (diagonal) and misidentification probability (off-diagonal) as a function of tracks' p_T for $\pi^+\pi^-$, K^+K^- and $p\bar{p}$ pairs. The results were obtained from the dedicated MC simulation described in the text. Blue lines and arrows mark the cut value on lower of the track p_T 's for kaons and protons.

The identification efficiency/misidentification probability was studied in two dimensions - as a function of particles' momenta. Because of that the shape of momentum distribution for PID efficiency study was not required to match the data and was defined flat, spanning between 0.2 GeV and 3 GeV, to provide low statistical uncertainties over full momentum space. We considered particles' momenta as the best quantities to study pair identification because $n\sigma_X$ and m_{TOF}^2 and their resolutions depend nearly solely on magnitude of momentum.

The identification efficiency ϵ^X and misidentification probability $\lambda^{X \rightarrow Y}$ were calculated for each species using the pair selection cuts C8.1 described in more detail in Sec. 3.2.7, following the definitions provided in Eqs. (5.14) and (5.15):

$$\epsilon^X(p_{\text{max}}^{\text{max}}, p_{\text{min}}^{\text{min}}) = \frac{N_{\text{true}X}^{\text{reco}X}(p_{\text{max}}^{\text{max}}, p_{\text{min}}^{\text{min}})}{N_{\text{true}X}(p_{\text{max}}^{\text{max}}, p_{\text{min}}^{\text{min}})}, \quad (5.14) \quad \lambda^{X \rightarrow Y}(p_{\text{max}}^{\text{max}}, p_{\text{min}}^{\text{min}}) = \frac{N_{\text{true}X}^{\text{reco}Y}(p_{\text{max}}^{\text{max}}, p_{\text{min}}^{\text{min}})}{N_{\text{true}X}(p_{\text{max}}^{\text{max}}, p_{\text{min}}^{\text{min}})}, \quad (5.15)$$

where $N_{\text{true}X}$ is number of XX pairs, $N_{\text{true}X}^{\text{reco}X}$ is number of XX pairs correctly recognized as XX , $N_{\text{true}X}^{\text{reco}Y}$ is number of XX pairs misidentified as YY and $p_{\text{max}}^{\text{max}}$ and $p_{\text{min}}^{\text{min}}$ are respectively higher and lower momentum of particles in the pair. The results as a function of $(p_{\text{max}}^{\text{max}}, p_{\text{min}}^{\text{min}})$ are presented in Fig. E.1 (Appendix E). Here in Fig. 5.10 we show the result presented as a function of higher and lower transverse momentum of particles in

the pair (p_T^{\max} , p_T^{\min}). This result clearly indicated necessity of adding a cut on the lower transverse momentum of particle in the pair for K^+K^- and $p\bar{p}$, as above certain p_T^{\min} the identification efficiency drops to a very low level (< 10%). The cuts (C8.2) are $p_T^{\min} > 0.7$ GeV for kaons and $p_T^{\min} > 1.1$ GeV for protons, indicated with dashed blue lines and arrows.

5.2.3 RP track acceptance and reconstruction efficiency

To calculate RP acceptance and track reconstruction efficiency the embedded MC technique was used. The same sample was used as that described in Sec. 5.2.1.4, used for calculation of the dead-material related trigger veto.

The joint RP acceptance and track reconstruction efficiency for a given STAR side, $\epsilon_{RP}^{\text{side}}$, was calculated as a probability that a single good quality RP track (satisfying cuts C4.1-C4.2) matched with true-level primary forward proton is reconstructed on given side in the branch expected based on sign of p_y of the proton, under condition that there is a trigger signal in that branch and there is no trigger signal in the other branch on the same side.

Technically the $\epsilon_{RP}^{\text{side}}$ was obtained in the following procedure:

1. It was verified if there is a trigger signal in the branch that the primary forward proton is expected to reach based on its p_y ($p_y > 0$ - branch UP, $p_y < 0$ - branch DOWN). Additionally required lack of trigger signal in the other branch on the same side. These events formed *set A*.
2. The nominal RP track selection algorithm was used to find a single good quality track (cuts C4.1-C4.2) on given side. If exactly one such track was found, it was additionally checked if it is matched with true-level primary proton. These events formed *set B*.
3. The efficiency was determined by the ratio of histograms from *set B* and *set A*:

$$\epsilon_{RP}^{\text{side}}(p_x, p_y, z_{\text{vtx}}) = \mathcal{E} \left(\text{RP}^{\text{side}} \mid \text{TR}^{\text{side}} \wedge \neg \text{TR}^{\text{side}} \right) = \frac{(p_x, p_y, z_{\text{vtx}}) \text{ histogram for protons from set } B}{(p_x, p_y, z_{\text{vtx}}) \text{ histogram for protons from set } A} \quad (5.16)$$

It should be noted that the momentum components (p_x, p_y) were taken from the proton with accounted effect of the beam divergence (after the original initial momentum smearing).




Figure 5.11: Sample RP track reconstruction efficiency in a single z -vertex bin on the east STAR side. The efficiency was calculated using forward proton MC simulation embedded into zero-bias data. Green envelopes mark the fiducial region of the measurement, while dashed yellow lines mark the part of the fiducial region with a data-driven efficiency correction needed, as explained in Sec. 10.3.1 of Ref. [1].




Figure 5.12: Probability of simultaneous inefficiency of the RP track reconstruction calculated from the CEP MC embedded into zero-bias data.

The efficiency calculated in the way described above is, by design, the acceptance, reconstruction and selection efficiency for a single proton on one side of the IP. In CEP event there are two independent forward protons which may be simultaneously not reconstructed or rejected by the selection algorithm due to e.g. elastic pile-up interaction providing additional good quality proton tracks on both sides of IP. One could, in principle, calculate 5-dimensional efficiency for both forward protons (in variables $p_x^E, p_y^E, p_x^W, p_y^W$ and z_{vtx}) which would ultimately account for the simultaneous east and west inefficiency, however this would require orders of magnitude larger statistics of MC to provide reasonably low statistical uncertainty of the efficiency. Instead, on top of the 3-dimensional reconstruction and selection efficiencies for east and west RPs we calculate (from embedded MC) probability that the proton tracks are simultaneously not reconstructed or selected on the east and west side, despite the trigger signal solely in expected RP branches (no trigger veto) and true-level (p_x, p_y) of both forward protons contained in the fiducial region. This probability, denoted as

$$\mathcal{P}_{\text{loss}}^{\text{E\&W}} = \mathcal{E} \left(!\text{RP}^E \wedge !\text{RP}^W \mid \text{TR}^E \wedge \text{TR}^W \wedge !\text{Veto} \right), \quad (5.17)$$

is presented in Fig. 5.12 for all four combinations of east and west RP branches.

5.2.4 TPC vertex reconstruction efficiency

The definition of vertex reconstruction efficiency (ϵ_{vtx}) established in this analysis is the probability that two global tracks, both associated with true-level primary particles from the kinematic region of the measurement, both satisfying kinematic and quality criteria (cuts C3.3 and C3.4) and both matched with hits in TOF, form a vertex listed in the collection of reconstructed primary vertices and DCA(R) and DCA(z) of both global tracks calculated w.r.t. this vertex is contained within the limits of cut C3.5.

We calculated this efficiency as a function of the longitudinal separation between two tracks (global helices) Δz_0 . Illustration of this quantity is given in Fig. 5.13. We consider this a natural quantity to present the vertexing efficiency - the closer to each other the helices are on the beamline, the more probable it is that two tracks will form a common primary vertex. It is in accordance with the way the vertexing algorithm works.

The vertexing efficiency was calculated from the data in the following way:

1. Data from RP_CPT2 trigger were used. Events were selected with nearly the same cuts as in nominal CEP analysis (Sec. 3.1). The requirement of exactly one primary vertex with exactly two primary TOF tracks was dropped. Instead, analysis utilized only global TOF tracks - exactly two global TOF tracks were required (cut C3.1 without primary track requirement), passing also cuts C3.2-C3.4. In this case the position of the vertex was reconstructed as

$$z_{\text{vtx}} = \frac{z_0^+ + z_0^-}{2}, \quad (5.18)$$

where z_0^+ and z_0^- are longitudinal impact parameters (z -coordinates of points of closest approach to the beamline) of positive and negative charge particle tracks, respectively. The vertex position was normally required to satisfy cut C2. Events after full selection, classified as exclusive $\pi^+\pi^-$ candidates, formed set A.

2. The two global TOF tracks were checked if they have associated primary tracks, and the two primary tracks are assigned to the same primary vertex. If yes, the tracks were additionally subjected to cut C3.5. Events passing described selection formed set B.
3. The efficiency was determined by the ratio of histograms from set B and set A:

$$\epsilon_{\text{vtx}}(|\Delta z_0|) = \frac{|\Delta z_0| \text{ histogram for events from set } B}{|\Delta z_0| \text{ histogram for events from set } A}. \quad (5.19)$$

Distribution of $|\Delta z_0|$ between two CEP global track candidates after full selection (setA) is presented in Fig. 5.14. The vertexing efficiency obtained with described method is shown in Fig. 5.15. Solid green points represent efficiency calculated with the non-exclusive background preserved, while open black points represent efficiency with this background subtracted (using method described in Sec. 5.5). Since the vertexing efficiency does not depend on the physics process and background is purely of physics origin the black and green points




Figure 5.13: Sketch with definition of the longitudinal separation of two tracks (helices) Δz_0 .

should overlap. Such picture emerges from presented comparison. We have calculated the same efficiency using CEP MC embedded into zero-bias data, the result is shown in Fig. 5.15 with red points. There is very good agreement between vertexing efficiency in the data and embedded MC for $|\Delta z_0| < 1$ cm, where most ($\sim 80\%$) of the signal is present. The differences in high- $|\Delta z_0|$ tail are understood as a result of the imperfect description of the pointing resolution (here: the transverse resolution) of TPC tracks in STARsim. Although the pointing resolution is adjusted to gain more accurate description of the data by MC simulation as described in Chapter 8 of Ref. [1], this does not help with the vertexing which is performed at the level of raw data (MC) processing to MuDst. Another reason could be different p_T (thus also d_0) spectrum of CEP tracks in the data and MC (GenEx).




Figure 5.14: Distribution of Δz_0 between two CEP-candidate global tracks (green and black points) together with fraction of non-exclusive background in black distribution as a function of Δz_0 (violet points).

Based on the width of $|\Delta z_0|$ distribution, the background content as a function of $|\Delta z_0|$, and the value of ϵ_{vtx} as a function of $|\Delta z_0|$, we decided to accept in analysis only tracks which satisfy $|\Delta z_0| < 2$ cm. This assures that the vertexing efficiency does not drop below $\approx 30\%$, as well as it coincides with the primary tracks requirement of $|\text{DCA}(z)| < 1$ cm. In the correction procedure we nominally use the vertexing efficiency represented by open black points in Fig. 5.15 (to correct the MC e.g. in closure tests we use red points instead). The additional correction factor connected with the cut on maximum $|\Delta z_0|$ (2 cm) was calculated from the data and equals 2.4% (Fig. 5.14). This correction is used as an additional normalization correction factor, which is different from $\epsilon_{\text{vtx}}(|\Delta z_0|)$ applied in form of a weight to each selected CEP event.

5.3 Particle energy loss

Energy loss correction as a function of reconstructed particle p_T in bins of z -position of reconstructed vertex has been calculated and presented in Chapter 5 of Ref. [1] for all analyzed particle species and both positive and negative charges. The correction was applied independently for each particle in the following procedure:

1. After central particles were identified (cut C8) an absolute value of the particle transverse momentum correction ($-\Delta p_T = p_T^{\text{meas}} - p_T^{\text{true}}$) was read from the histogram corresponding to reconstructed z_{vtx} and to assigned particle ID (Appendix C of Ref. [1]).
2. The momentum correction factor f_p^{corr} was calculated:

$$f_p^{\text{corr}} = \frac{p_T^{\text{meas}} + \Delta p_T}{p_T^{\text{meas}}}. \quad (5.20)$$

3. New, corrected momentum \vec{p}^{corr} was assigned to the particle:

$$\vec{p}^{\text{corr}} = f_p^{\text{corr}} \cdot \vec{p}^{\text{meas}}. \quad (5.21)$$




Figure 5.15: Vertexing efficiency calculated from the data (open black and full green points) and from embedded MC (red points) as a function of Δz_0 between two CEP-candidate global tracks.

In this way all three components of particle momentum are corrected so that the pseudorapidity of a particle remains unchanged.

This new momentum was further used in determination of total transverse momentum of all reconstructed particles, p_T^{miss} , as well as in applying TPC and TOF efficiency corrections and preparing histograms (cross sections) of physics quantities (e.g. invariant mass, rapidity of a pair of central tracks).

5.4 Fake tracks and migrations into and out of fiducial region

Finite resolutions play non-negligible role in presented analysis. Smearing of reconstructed quantities with respect to their true value at the hadron level makes some tracks from outside of the fiducial phase space migrating/leaking into it, as well as some tracks from inside the fiducial region escaping from it.

Correction related to migrations and fake tracks was calculated using quantities described in the listing below. It applies to both TPC and RP tracks.

- N_{reco} - number of true level-matched tracks reconstructed inside fiducial region,
- $N_{\text{migr.}}^{\text{in}}$ - number of true level-matched tracks from outside of fiducial region (at the hadron level), reconstructed inside fiducial regions,
- $N_{\text{migr.}}^{\text{out}}$ - number of true level-matched tracks from inside of fiducial region (at the hadron level), reconstructed outside fiducial region,
- N_{fakes} - number of tracks not matched with true level primary but consistent with it, reconstructed inside fiducial region.

All quantities above are represented as a function of reconstructed values (TPC tracks: p_T and η , RP tracks: p_x and p_y), except $N_{\text{migr.}}^{\text{out}}$, for which true (hadron) level values are used. Definitions of fraction of migrations into and out of fiducial region, as well as fraction of fake tracks inside the fiducial region, are provided in Eqs. (5.22)-(5.24) in the same order:

$$f_{\text{migr.}}^{\text{in}} = \frac{N_{\text{migr.}}^{\text{in}}}{N_{\text{reco}}}, \quad (5.22) \quad f_{\text{migr.}}^{\text{out}} = \frac{N_{\text{migr.}}^{\text{out}}}{N_{\text{reco}}}, \quad (5.23) \quad f_{\text{fakes}} = \frac{N_{\text{fakes}}}{N_{\text{reco}}}. \quad (5.24)$$

Definition of number of fake tracks requires some elaboration. The necessary condition to claim a track being fake is that it is not matched with a true level primary particle. However, such tracks should typically have momentum significantly different from that of unmatched primary, therefore total momentum cut C9 would not be satisfied and thus these tracks should not be subtracted. For this reason, an additional condition to assign a track as fake is that its reconstructed momentum is consistent with that of primary particle. In case of TPC tracks we use cut on the distance δ between true and reconstructed (η, ϕ) , $\delta < 0.15$, introduced in Sec. 3.2.2 of Ref. [1]. For the RP tracks we use cut on the difference between true and reconstructed transverse momentum not greater than 70 MeV. Such value, somewhat arbitrary, is driven by the value used in cut C9. We do not consider any systematic effect connected with the fake RP track definition as this effect is believed to be much smaller than most of other systematic uncertainties. In Figs. 5.16 and 5.17 we present distributions of quantities for matched and not matched tracks with magenta circles drawn on top marking the conditions used in fake track definitions.

Fraction of tracks, for both TPC and RP detectors, migrating from/to fiducial region and being fake, is shown in Figs. 5.18 and 5.19. Clearly, all mentioned effects are dominant for RP tracks.

Multiplicative correction factor (per track) that takes into account all migrations and fakes can be formed as

$$C_{\text{m,f}} = \frac{N_{\text{reco}} + N_{\text{migr.}}^{\text{out}} - N_{\text{migr.}}^{\text{in}} - N_{\text{fakes}}}{N_{\text{reco}}} = 1 + f_{\text{migr.}}^{\text{out}} - f_{\text{migr.}}^{\text{in}} - f_{\text{fakes}}. \quad (5.25)$$

This correction is graphically represented in Fig. 5.20 for both TPC (5.20a) and RP (5.20b) tracks. One can easily notice that most important role plays correction related to migrations at the edge of fiducial region defining accepted forward scattered protons, reaching as much as 25% correction for tracks at the corners of allowed (p_x, p_y) . This is a direct consequence of sizeable angular divergence of proton beams.




Figure 5.16: Difference between reconstructed and true parameters of the TPC tracks. Magenta circle of radius 0.15 drawn on top of distributions represent cut on maximum difference between true and reconstructed η and φ of tracks used for determination of fake tracks fraction, as explained in the text. The same z -axis (color) scale has been set up for both plots to better visualize relative content of true and fake tracks.




Figure 5.17: Difference between reconstructed and true momentum components of RP tracks matched (a) and not matched (b) with true level primary forward protons. Magenta circle of radius 70 MeV drawn on top of distributions represent cut on maximum difference between true and reconstructed transverse momentum used for determination of fake tracks fraction, as explained in the text. The same z -axis (color) scale has been set up for both plots to better visualize relative content of true and fake tracks.




Figure 5.18: Fraction of TOF-matched TPC tracks migrating into the fiducial central tracks phase space from outside of it (a), migrating outside the fiducial phase space from inside of it (b), and fraction of fake TOF matched TPC tracks reconstructed in the fiducial phase space (c). In case of reconstructed (p_T , η) escaping the fiducial phase space (b) the true value of p_T and η is used; otherwise reconstructed values are plotted.




Figure 5.19: Fraction of RP tracks migrating into the fiducial forward proton momentum area from outside of it (a), migrating outside the fiducial area from inside of it (b), and fraction of fake RP tracks reconstructed in the fiducial area (c). In case of reconstructed tracks momentum escaping the fiducial area (b) the true value of p_x and p_y is used; otherwise reconstructed momentum components are plotted.




Figure 5.20: Joint correction factors for migrations and fake tracks for TOF-matched TPC tracks (5.20a) and RP tracks (5.20b).

5.5 Background subtraction

5.6 Closure tests

5.6.1 Data pull

Sanity of the data after applied corrections was verified by preparing a pull histogram of the number of reconstructed and selected CEP events per unit of integrated luminosity (in other words - an integrated cross section). The pull quantity was calculated per run, defined as

$$\text{Pull}_i = \begin{cases} \frac{\sigma_i - \langle \sigma \rangle}{u^-(\sigma_i)}, & \text{if } \sigma_i - \langle \sigma \rangle \geq 0, \\ \frac{\sigma_i - \langle \sigma \rangle}{u^+(\sigma_i)}, & \text{if } \sigma_i - \langle \sigma \rangle < 0, \end{cases} \quad (5.26)$$

where σ_i is an integrated cross section calculated for i -th run with statistical uncertainty equal to $u^-(\sigma_i)$ or $u^+(\sigma_i)$ (lower or upper 68% confidence level interval), and $\langle \sigma \rangle$ is an expectation value of integrated cross section. For each run the integrated cross section was calculated as

$$\sigma_i = \frac{N_i}{L_i}. \quad (5.27)$$

where N_i is a number of CEP events in i -th run and L_i is a luminosity collected during i -th run. The expectation value $\langle \sigma \rangle$ was calculated using an estimator from Eq. (5.28):

$$\langle \sigma \rangle = \frac{\sum_i \sigma_i \times u^{-2}(\sigma_i)}{\sum_i u^{-2}(\sigma_i)} \quad (5.28)$$

which is simply an average integrated cross section calculated with full dataset, weighted by the uncertainties (by reciprocal of variance).

Final pull histogram is shown in Fig. 5.21 for raw event counts (open circles) and efficiency-corrected events weighted according to prescription from Eq. (5.5) (filled circles). Parameters of the Gaussian fits to data points show agreement of the mean value with 0, as expected from the pull, and significant improvement of the width of distribution after appliance of corrections, which should be consistent with 1.




Figure 5.21: Pull histogram of number of events per unit of integrated luminosity. Open circles fitted with dashed blue Gaussian mark uncorrected data, whereas black dots fitted with red curve denote corrected (weighted) data. Parameters obtained from the fit are provided in the plot.

5.6.2 Monte Carlo closure tests

5.6.2.1 TPC and TOF efficiency correction validation

5.6.2.2 RP efficiency correction validation

5.6.2.3 Total correction validation




Figure 5.22: Full closure test.

6. Systematic uncertainties

6.1 Trigger veto effect (due to dead material)

Systematic uncertainty related to the trigger veto correction (Sec. 5.2.1.4) has been studied with elastic scattering events, in a way similar to analyses presented in Sec. 10.3 (and following) of Ref. [1].

Triggers dedicated for elastic scattering process (RP_ET triggers) have been studied. The trigger required signal in at least one PMT (out of four) in two RP branches opposite to each other with respect to interaction region. In addition to trigger selection, vetoes were imposed offline on any activity in other STAR detectors, such as BBC (small and large), ZDC, TOF and VPD - it reduced probability of a non-elastic pile-up interaction. It has been demonstrated in Sec. 10.3.1 of Ref. [1] that once the single good quality RP track is required on one side, such sample consists only of elastic proton-proton scattering events.

Each side (branch) was analyzed independently; when single good quality RP track, in addition of $|\xi| < 0.01$, was found on the east side, the systematics for west side was investigated (and vice versa). Two histograms were filled per event. The first histogram was filled with all selected events. The second histogram was filled only if there was no simultaneous signal in upper and lower RP on studied side (no ET and IT trigger bits fired at the same time) - just the same, as it was implemented in RP_CPT2 trigger. However, the second histogram was filled with weight equal to inverse efficiency of the veto, $1/\epsilon_{\text{DM veto}}^{\text{side}}$. The ratio of the second to first histogram,

$$R_{\text{DM veto}} = \frac{\text{histogram of events with satisfied veto, filled with weight } 1/\epsilon_{\text{DM veto}}^{\text{side}}}{\text{histogram of all events, filled with unit weight}}, \quad (6.1)$$

has been presented in Fig. 6.1 as a function of p_x and p_y of elastically scattered proton on studied side. If single good quality proton track was reconstructed in studied branch, parameters of that track were histogrammed. Otherwise, transverse components of momentum of unreconstructed elastic proton (e.g. due to induced shower) were estimated as $-p_x$ and $-p_y$ of elastic proton track on the opposite side.




Figure 6.1: Ratio $R_{\text{DM veto}}$ illustrating systematic uncertainty of the dead material trigger veto correction in east (left) and west (right) Roman Pots.

Results presented in Fig. 6.1 indicate imperfect description of the dead material of the elements surrounding RPs (DX-D0 chamber, RF shield), which has been also overed in studies of systematic uncertainties related to RP track reconstruction efficiency, presented in Ref. [1]. Based on current study, the correction and systematic uncertainty of the dead material veto efficiency $\epsilon_{\text{DM veto}}^{\text{side}}$ are assumed to have the following form:

- multiplicative correction is applied to $\epsilon_{\text{DM veto}}^{\text{side}}$, equal to $1 + \frac{1}{2}(R_{\text{DM veto}} - 1)$,
- systematic uncertainty is assumed to be equal to $1 + \frac{1}{2}|R_{\text{DM veto}} - 1|$, therefore propagation of this systematic effect is done by variating the efficiency $\epsilon_{\text{DM veto}}^{\text{side}}$ by the multiplicative factor equal to $\pm [1 + \frac{1}{2}|R_{\text{DM veto}} - 1|]$.

6.2 Discussion of systematic effects

The following contributions to the overall systematic uncertainty have been studied. Influence of each systematic effect on measured cross sections has been tested by changing amount of the quantity that the systematic effect refers to and comparing the result with that obtained using nominal values.

1. Representativeness of the embedding sample ($\Delta\epsilon_{\text{TPC}}(\text{embed. stat.})$).

Zero-bias data available for the MC embedding is only a fraction of all zero-bias triggers, therefore physics data may be not fully represented in MC events used for determination of the TPC track reconstruction efficiency. This effect was studied by comparing estimated average levels of pile-up in the data and embedded MC. The difference was found to be of the order of 1%, which we establish as a symmetric systematic uncertainty on the TPC track reconstruction efficiency (per track). See the last paragraph of Sec. 10.1.1 of Ref. [1] for more details.

2. Embedding procedure/off-time pile-up effect ($\Delta\epsilon_{\text{TPC}}(\text{pile-up})$).

Reliability and precision of the embedding technique was verified and quantitatively estimated in the procedure described in Sec. 10.1.1 of Ref. [1]. Embedded MC samples were divided into sub-samples representing different levels of off-time pile-up/density of hit points in TPC. With dedicated analysis it was possible to verify if the TPC track reconstruction efficiency is compatible between all sub-samples when the effect of pile-up (changing number of hits forming a track) is reduced. The average systematic uncertainty related to the embedding procedure is < 1% (per track, Fig. 10.5 of Ref. [1]).

3. Modelling of the dead material in front of the TPC ($\Delta\epsilon_{\text{TPC}}(\text{dead mat.})$).

Not all detector elements are fully modeled in the MC simulation, quite often some simplifications are used. This leads to inaccuracies in efficiencies derived from the simulation. We estimated systematic uncertainty related to amount of simulated material between the primary vertex and STAR TPC to be 25% which translates to $\approx 0.5\%$ of uncertainty of the TPC track reconstruction efficiency. See Chap. 9 of Ref. [1] for details.

4. Vertexing ($\Delta\epsilon_{\text{vtx}}$).

Vertexing efficiency has been obtained using data-driven method presented in Sec. 5.2.4, thus systematic uncertainty related to this efficiency has been significantly reduced. Systematic uncertainty has been estimated as a difference between efficiency with and without subtracted background.

5. Modelling of the TOF system and validity of derived efficiency corrections ($\Delta\epsilon_{\text{TOF}}$).

The efficiency of matching TOF hits with the TPC tracks has been extracted from embedded MC sample. It has been confronted with TOF efficiency extracted from the data using two independent techniques: tag&probe (Sec. 4.1 of Ref. [1]) and HFT-tagging (Sec. 10.2.2 of Ref. [1]). Average difference between the data and MC efficiency has been used as a correction to MC efficiency, while half of the difference between data-extracted efficiencies has been treated as a systematic uncertainty (Fig. 10.16, Ref. [1]). This amounts 1%-3% (per track), depending on particle species.

6. Modelling of the RP system and validity of derived efficiency corrections ($\Delta\epsilon_{\text{RP}}$, $\Delta\epsilon_{\text{RP}}^{\text{trig.}}$ and $\Delta\epsilon_{\text{RP}}^{\text{DM veto}}$).

Reliability of RP simulation which was used to extract efficiency corrections with MC embedded into zero-bias data, has been verified and quantitatively estimated in the procedure described in Sec. 10.3 of Ref. [1]. For this purpose elastic proton-proton scattering events have been used. The same analysis has been performed on embedded elastic scattering MC and the data, leading to estimates of the RP acceptance and track reconstruction efficiency. The differences between two results has been considered as a measure of the systematic uncertainty that covers RP track reconstruction efficiency itself, detectors alignment, embedding technique. Similar studies have been performed to determine systematic uncertainty related to the trigger veto efficiency correction, as presented in Sec. 6.1.

7. Pile-up veto correction ($\Delta\epsilon_{\text{veto}}$).

Luminosity-dependent correction related to veto of pile-up interactions is derived from the zero-bias data on run by run basis. Residual systematic uncertainty has been estimated as a difference between the correction factor calculated for particular run, and correction factor obtained from the exponential fit to all points representing correction factors as a function of instantaneous luminosity.

8. Longitudinal shape and position of the primary vertex distribution ($\Delta\langle z_{\text{vtx}} \rangle$ and $\Delta\sigma(z_{\text{vtx}})$).

Comparison of the z_{vtx} distribution as seen in the TPC and in RPs leads to conservative estimate of the uncertainty of central position of the vertex equal 2 cm, and the spread (standard deviation) of vertex equal 3 cm.

9. Luminosity determination ($\Delta\mathcal{L}$).

Uncertainty of the integrated luminosity has been estimated to 7%, which is still subject to change.

6.3 Graphical representation of systematic uncertainties

In the remaining part of this chapter we present relative contributions of effects listed in Sec. 6.2 to the total systematic uncertainties on differential fiducial cross sections presented in Chap. 7. Numbering of figures is preserved with respect to corresponding cross section results in the next chapter. The color code is the same in all figures, with the legend explaining the meaning of each color attached at the bottom of Fig. 6.2.




Figure 6.2: Systematic uncertainties of the differential cross sections for CEP of charged particle pairs $\pi^+\pi^-$ as a function of the invariant mass of the pair in the fiducial region explained on the plots.




Figure 6.3: Systematic uncertainties of the differential cross sections for CEP of charged particle pairs K^+K^- (left) and $p\bar{p}$ (right) as a function of the invariant mass of the pair in the fiducial region explained on the plots.




Figure 6.4: Systematic uncertainties of the differential cross sections for CEP of charged particle pairs $\pi^+\pi^-$ (left), K^+K^- (middle) and $p\bar{p}$ (right) as a function of the pair rapidity measured in the fiducial region explained on the plots.




Figure 6.5: Systematic uncertainties of the differential cross sections for CEP of charged particle pairs $\pi^+\pi^-$ (left column), K^+K^- (middle column) and $p\bar{p}$ (right column) as a function of the difference of azimuthal angles of the forward scattered protons (top) and of the sum of the squares of the four-momenta losses in the proton vertices (bottom) measured in the fiducial region explained on the plots.




Figure 6.6: Systematic uncertainties of the differential cross sections for CEP of charged particle pairs $\pi^+\pi^-$ (top), K^+K^- (middle) and $p\bar{p}$ (bottom) as a function of the invariant mass of the pair in two $\Delta\phi$ regions: $\Delta\phi < 90^\circ$ (left column) and $\Delta\phi > 90^\circ$ (right column) measured in the fiducial region explained on the plots.




Figure 6.7: Systematic uncertainties of the differential cross sections for CEP of $\pi^+\pi^-$ pairs as a function of the rapidity of the pair (left column) difference of azimuthal angles of the forward scattered protons (middle column) and of the sum of the squares of the four-momentum losses in the proton vertices (right column) measured in the fiducial region explained on the plots, separately for three ranges of the $\pi^+\pi^-$ pair invariant mass: $m < 1$ GeV (top), $1 < m < 1.5$ GeV (middle) and $m > 1.5$ GeV (bottom).




Figure 6.8: Systematic uncertainties of the differential cross sections $d\sigma/dm(\pi^+\pi^-)$ for CEP of $\pi^+\pi^-$ pairs in two $|\vec{p}'_{1,T} - \vec{p}'_{2,T}|$ regions: $|\vec{p}'_{1,T} - \vec{p}'_{2,T}| < 0.12$ GeV (left) and $|\vec{p}'_{1,T} - \vec{p}'_{2,T}| > 0.12$ GeV (right) in the fiducial region and $\Delta\phi < 90$ degree.




Figure 6.9: Systematic uncertainties of the differential cross sections for CEP of $\pi^+\pi^-$ pairs as a function of $\cos\theta^{\text{CS}}$ (top) and of ϕ^{CS} (bottom) measured in the fiducial region explained on the plots, separately for three ranges of the $\pi^+\pi^-$ pair invariant mass: $m < 1$ GeV (left column), $1 < m < 1.5$ GeV (middle column) and $m > 1.5$ GeV (right column).

7. Physics results




Figure 7.1: Differential cross sections for CEP of charged particle pairs $\pi^+\pi^-$ as a function of the invariant mass of the pair in the fiducial region explained on the plots. Data are shown as solid points with error bars representing the statistical uncertainties. The typical systematic uncertainties are shown as gray boxes for only few data points as they are almost fully correlated between neighboring bins. Predictions from MC models GenEx, DiMe and MBR are shown as histograms. In the lower panels in the bottom plots the ratios of the MC predictions scaled to data and the data are shown.




Figure 7.2: Differential cross sections for CEP of charged particle pairs K^+K^- (left) and $p\bar{p}$ (right) as a function of the invariant mass of the pair in the fiducial region explained on the plots. Data are shown as solid points with error bars representing the statistical uncertainties. The typical systematic uncertainties are shown as gray boxes for only few data points as they are almost fully correlated between neighboring bins. Predictions from MC models GenEx, DiMe and MBR are shown as histograms. In the lower panels in the bottom plots the ratios of the MC predictions scaled to data and the data are shown.




Figure 7.3: Differential cross sections for CEP of charged particle pairs $\pi^+\pi^-$ (left), K^+K^- (middle) and $p\bar{p}$ (right) as a function of the pair rapidity measured in the fiducial region explained on the plots. Data are shown as solid points with error bars representing the statistical uncertainties. The typical systematic uncertainties are shown as gray boxes for only few data points as they are almost fully correlated between neighboring bins. Predictions from MC models GenEx, DiMe and MBR are shown as histograms. In the lower panels in the bottom plots the ratios of the MC predictions scaled to data and the data are shown.




Figure 7.4: Differential cross sections for CEP of charged particle pairs $\pi^+\pi^-$ (left column), K^+K^- (middle column) and $p\bar{p}$ (right column) as a function of the difference of azimuthal angles of the forward scattered protons (top) and of the sum of the squares of the four-momenta losses in the proton vertices (bottom) measured in the fiducial region explained on the plots. Data are shown as solid points with error bars representing the statistical uncertainties. The typical systematic uncertainties are shown as gray boxes for only few data points as they are almost fully correlated between neighboring bins. Predictions from MC models GenEx, DiMe and MBR are shown as histograms. In the lower panels the ratios of the MC predictions scaled to data and the data are shown.




Figure 7.5: Differential cross sections for CEP of charged particle pairs $\pi^+\pi^-$ (top), K^+K^- (middle) and $p\bar{p}$ (bottom) as a function of the invariant mass of the pair in two $\Delta\phi$ regions: $\Delta\phi < 90$ degree (left column) and $\Delta\phi > 90$ degree (right column) measured in the fiducial region explained on the plots. Data are shown as solid points with error bars representing the statistical uncertainties. The typical systematic uncertainties are shown as gray boxes for only few data points as they are almost fully correlated between neighboring bins. Predictions from MC models GenEx, DiMe and MBR are shown as histograms.




Figure 7.6: Differential cross sections for CEP of $\pi^+\pi^-$ pairs as a function of the rapidity of the pair (left column) difference of azimuthal angles of the forward scattered protons (middle column) and of the sum of the squares of the four-momenta losses in the proton vertices (right column) measured in the fiducial region explained on the plots, separately for three ranges of the $\pi^+\pi^-$ pair invariant mass: $m < 1$ GeV (top), $1 < m < 1.5$ GeV (middle) and $m > 1.5$ GeV (bottom). Data are shown as solid points with error bars representing the statistical uncertainties. The typical systematic uncertainties are shown as gray boxes for only few data points as they are almost fully correlated between neighboring bins. Predictions from MC models GenEx, DiMe and MBR are shown as histograms. In the lower panels the ratios of the MC predictions scaled to data and the data are shown.




Figure 7.7: Differential cross sections $d\sigma/dm(\pi^+\pi^-)$ for CEP of $\pi^+\pi^-$ pairs in two $|\vec{p}'_{1,T} - \vec{p}'_{2,T}|$ regions: $|\vec{p}'_{1,T} - \vec{p}'_{2,T}| < 0.12$ GeV (left) and $|\vec{p}'_{1,T} - \vec{p}'_{2,T}| > 0.12$ GeV (right) in the fiducial region and $\Delta\phi < 90$ degree. There is no difference for two $|\vec{p}'_{1,T} - \vec{p}'_{2,T}|$ regions. Data are shown as solid points with error bars representing the statistical uncertainties. The typical systematic uncertainties are shown as gray boxes for only few data points as they are almost fully correlated between neighboring bins. Predictions from MC models GenEx, DiMe and MBR are shown as histograms.







Figure 7.8: Differential cross sections for CEP of $\pi^+\pi^-$ pairs as a function of $\cos\theta^{\text{CS}}$ (top) and of ϕ^{CS} (bottom) measured in the fiducial region explained on the plots, separately for three ranges of the $\pi^+\pi^-$ pair invariant mass: $m < 1 \text{ GeV}$ (left column), $1 < m < 1.5 \text{ GeV}$ (middle column) and $m > 1.5 \text{ GeV}$ (right column). Data are shown as solid points with error bars representing the statistical uncertainties. The typical systematic uncertainties are shown as gray boxes for only few data points as they are almost fully correlated between neighboring bins. Predictions from MC models GenEx, DiMe and MBR are shown as histograms. In the lower panels the ratios of the MC predictions scaled to data and the data are shown.


A. BBC response


Figure A.1: Two-dimensional distribution of ADC vs. TAC counts per BBC-small channel in abort gaps (left) and colliding bunches (middle), and one-dimensional projection on x -axis (ADC) for selected ranges of TAC for colliding bunches and abort gaps (right). Each row represents single channel (small BBC tile). Red lines and arrows indicate thresholds for a signal in given channel.

















Figure A.2: Two-dimensional distribution of ADC vs. TAC counts per BBC-large channel in abort gaps (left) and colliding bunches (middle), and one-dimensional projection on x -axis (ADC) for selected ranges of TAC for colliding bunches and abort gaps (right). Each row represents single channel (large BBC tile). Red lines and arrows indicate thresholds for a signal in given channel.







B. Formulation of total RP efficiency

Definitions:

- RP^E - single good quality track (satisfying cuts C4.1-C4.2) on the east side,
- RP^W - single good quality track (satisfying cuts C4.1-C4.2) on the west side,
- TR^E - trigger signal in the RP branch with single good track on the east side,
- $TR^{\neq E}$ - trigger signal in the RP branch other than branch with single good track on the east side,
- TR^W - trigger signal in the RP branch with single good track on the west side,
- $TR^{\neq W}$ - trigger signal in the RP branch other than branch with single good track on the west side,
- Veto - trigger veto on the simultaneous trigger signal in Up and Down RPs (ET&IT),
- $Veto^{PU}$ - trigger veto on ET&IT (Veto) due to pile-up interactions,
- $Veto^{DM}$ - trigger veto on ET&IT (Veto) due to forward proton interaction with dead material.

The total efficiency related to both east and west forward protons in CEP event has the following form:

$$\mathcal{E}(RP^E \wedge RP^W \wedge TR^E \wedge TR^W \wedge !Veto) = \mathcal{E}(RP^E \wedge RP^W | TR^E \wedge TR^W \wedge !Veto) \times \mathcal{E}(TR^E \wedge TR^W \wedge !Veto), \quad (B.1)$$

where the r.h.s. part of the equation is factorized using the rules of conditional probability to two components describing reconstruction and selection efficiency (first) and trigger efficiency (second).

The reconstruction and selection efficiency part can be represented as a product of single-proton reconstruction and selection efficiencies described in Sec. 5.2.3 with an additional scaling factor in the denominator:

$$\mathcal{E}(RP^E \wedge RP^W | TR^E \wedge TR^W \wedge !Veto) = \frac{\mathcal{E}(RP^E | TR^E \wedge !TR^{\neq E}) \times \mathcal{E}(RP^W | TR^W \wedge !TR^{\neq W})}{1 - \mathcal{E}(!RP^E \wedge !RP^W | TR^E \wedge TR^W \wedge !Veto)} \quad (B.2)$$

This denominator accounts for the fact that the efficiencies of the east and west RPs are correlated. To be more precise, unsuccessful reconstruction/selection of RP track on the east and west side in the same event can be a result of a pile-up interaction, typically of elastic proton-proton scattering, producing additional tracks/showers simultaneously in east and west RPs and thus introducing simultaneous east and west RP inefficiency. Without this factor the sole product of east and west efficiencies would account for discussed inefficiency twice - the total RP efficiency would be then underestimated. The effect is small (as can be read from Fig. 5.12), but we correct for it for completeness.

In the component of RP efficiency related to the trigger we can use again the conditional probability and factorize it to part connected with the trigger veto (first) and the efficiency of detecting a signal of both forward protons by the trigger system (second):

$$\mathcal{E}(TR^E \wedge TR^W \wedge !Veto) = \mathcal{E}(!Veto | TR^E \wedge TR^W) \times \mathcal{E}(TR^E \wedge TR^W). \quad (B.3)$$

Efficiency of the triggering $\mathcal{E}(TR^E \wedge TR^W)$ is basically 1. Efficiency of the (lack of) veto if forward protons are triggering in east and west RPs can be decomposed to efficiency of the veto induced by the pile-up interaction in the same bunch crossing ($Veto^{PU}$) and efficiency of the veto induced by the interaction of the CEP protons with the material of the accelerator and detectors ($Veto^{DM}$):

$$\begin{aligned} \mathcal{E}(!Veto | TR^E \wedge TR^W) &= \left| Veto = Veto^{PU} \vee Veto^{DM} \right| = \mathcal{E}(!Veto^{PU} \wedge !Veto^{DM} | TR^E \wedge TR^W) = \\ &= \mathcal{E}(!Veto^{DM} | !Veto^{PU} \wedge TR^E \wedge TR^W) \times \mathcal{E}(!Veto^{PU} | TR^E \wedge TR^W). \end{aligned} \quad (B.4)$$

The first term of the last part of Eq. (B.4) described in Sec. 5.2.1.3 can be safely factorized as the probability of veto induced by the primary CEP proton on the east side is totally independent from the similar probability on the west side:

$$\begin{aligned} \mathcal{E}(!Veto^{DM} | !Veto^{PU} \wedge TR^E \wedge TR^W) &= \\ &= \mathcal{E}^E(!Veto^{DM} | !Veto^{PU} \wedge TR^E \wedge TR^W) \times \mathcal{E}^W(!Veto^{DM} | !Veto^{PU} \wedge TR^E \wedge TR^W) \end{aligned} \quad (B.5)$$

The second term of the last part of Eq. (B.4) related with the pile-up is incorporated to overall efficiency of online and offline vetoes described in Sec. 5.2.2.3 - this is required by the correlation of vetoes, the possibility that vetoes in independent subdetectors take place in the same bunch crossing. This could happen if e.g. single diffraction event occurs on top of the CEP event, yielding a BBC signal and RP signal, both vetoing RP_CPT2 trigger.

C. Reconstruction of m_{TOF}^2

Definitions:

- t_0 - time of the primary pp interaction
- $t_{1,2}$ - time of detection of the hit in TOF by particle 1(2)
- $L_{1,2}$ - helical path of the particle 1(2) from the interaction vertex to the TOF cell with reconstructed hit,
- $p_{1,2}$ - magnitude of momentum of particle 1(2),
- $m_{1,2}$ - mass of particle 1(2),




Figure C.1: Scheme of two central tracks of lengths L_1 and L_2 , produced in common vertex in moment t_0 , hitting cells in TOF detector in moments t_1 and t_2 .

From the simple algebra below which describes relation between track lengths, momenta and times of hit detection one can derive formula for the squared mass of two particles, assuming that their masses are equal (particles are of the same type).

Below we assume $c = 1$. We can write a set of two equations connecting the time that it takes for each particle to reach the TOF, starting from the interaction vertex:

$$\begin{cases} t_1 - t_0 = L_1 \sqrt{1 + \frac{m_1^2}{p_1^2}}, \\ t_2 - t_0 = L_2 \sqrt{1 + \frac{m_2^2}{p_2^2}}. \end{cases} \quad (\text{C.1})$$

By adding the two equations above we get

$$\Delta t = t_1 - t_2 = L_1 \sqrt{1 + \frac{m_1^2}{p_1^2}} - L_2 \sqrt{1 + \frac{m_2^2}{p_2^2}}. \quad (\text{C.2})$$

In CEP of two opposite-sign particles always the same species of particles are produced, therefore

$$m_1 = m_2 = m. \quad (\text{C.3})$$

If we substitute m_1 and m_2 with m in Eq. (C.2) and transform the equation to remove the square roots we get a quadratic equation of the form

$$\mathcal{A} \times (m_{\text{TOF}}^2)^2 + \mathcal{B} \times m_{\text{TOF}}^2 + \mathcal{C} = 0. \quad (\text{C.4})$$

Parameters of the Eq. (C.4) are given below:

$$\mathcal{A} = -2 \frac{L_1^2 L_2^2}{p_1^2 p_2^2} + \frac{L_1^4}{p_1^4} + \frac{L_2^4}{p_2^4}, \quad (\text{C.5})$$

$$\mathcal{B} = -2 L_1^2 L_2^2 \left(\frac{1}{p_1^2} + \frac{1}{p_2^2} \right) + \frac{2 L_1^4}{p_1^2} + \frac{2 L_2^4}{p_2^2} - 2 (\Delta t)^2 \left(\frac{L_1^2}{p_1^2} + \frac{L_2^2}{p_2^2} \right), \quad (\text{C.6})$$

$$\mathcal{C} = (\Delta t)^4 - 2 (\Delta t)^2 (L_1^2 + L_2^2) + L_1^4 + L_2^4 - 2 L_1^2 L_2^2, \quad (\text{C.7})$$


together with the final formula for a physical root of the quadratic equation which is used in the m_{TOF}^2 reconstruction:


$$m_{\text{TOF}}^2 = \frac{-\mathcal{B} + \sqrt{\mathcal{B}^2 - 4\mathcal{A}\mathcal{C}}}{2\mathcal{A}}. \quad (\text{C.8})$$

D. RP efficiency

D.1 RP track acceptance, reconstruction and selection efficiency

Figure D.1: RP track acceptance, reconstruction and selection efficiency on the east side obtained from MC simulation embedded into zero-bias data. Each plot corresponds to z -vertex range given in the plot.











Figure D.2: RP track acceptance, reconstruction and selection efficiency on the west side obtained from MC simulation embedded into zero-bias data. Each plot corresponds to z -vertex range given in the plot.









D.2 RP trigger veto probability related to dead material

Figure D.3: Probability of ET&IT trigger veto due to forward proton interaction with dead material on the east side. Results were obtained from forward proton MC simulation embedded into zero-bias data. Each plot corresponds to z -vertex range given in the plot.











Figure D.4: Probability of ET&IT trigger veto due to forward proton interaction with dead material on the west side. Results were obtained from forward proton MC simulation embedded into zero-bias data. Each plot corresponds to z -vertex range given in the plot.







E. Particle identification efficiency




Figure E.1: Pair identification efficiency (diagonal) and misidentification probability (off-diagonal) as a function of tracks' momentum (higher, x -axis, and lower, y -axis) for $\pi^+\pi^-$, K^+K^- and $p\bar{p}$ pairs. The results were obtained from the dedicated MC simulation described in Sec. 5.2.2.5.

List of Figures

1.1	Diagram of DIPE process.	5
1.2	CEP represented in η - ϕ space.	5
1.3	Central Production event at STAR.	5
2.1	Sketch of the trigger components used in definitions of diffractive triggers in run 15.	7
3.1	Primary vertex multiplicity. Red arrow marks bin with events with exactly one primary vertex (with track(s) matched with hit in TOF), which are used in physics analysis.	10
3.2	z -position of the primary vertex in single TOF vertex events. Red dashed line indicate range of longitudinal vertex position accepted in analysis.	10
3.3	Multiplicity of primary TPC tracks matched with hit in TOF for single TOF vertex events	11
3.4	Distribution of a distance in $\eta - \phi$ space between the BEMC cluster closest to primary TPC track (R_{\min})	11
3.5	Comparison of distribution of $N_{\text{hits}}^{\text{fit}}$, $N_{\text{hits}}^{\text{dE/dx}}$ and $N_{\text{hits}}^{\text{fit}}/N_{\text{hits}}^{\text{poss}}$ in the data and embedded MC	12
3.6	Comparison of distribution of track η and ϕ in the data and embedded MC	13
3.7	Local angle vs. position of RP tracks matched with true level primary protons.	14
3.8	(left) Merged distributions of diffractively scattered protons momenta p_y vs. p_x in exclusive h^+h^- events reconstructed with the East and West RP stations, together with the kinematic region used in the measurement marked with the black line. (right) Distributions of measured four momenta transfers at the proton vertices for exclusive h^+h^- events with all particles in the fiducial phase space are shown for East and West stations with yellow and blue color, respectively.	15
3.9	Correlation and difference of z -vertex position measured in Roman Pots and TPC.	16
3.10	Sample BBC-small and BBC-large response in zero-bias triggers.	17
3.11	Probability of false BBC-large signal (noise-induced).	18
3.12	NTofClusters.	19
3.13	$n\sigma_X^{\text{pair}}$ vs. $n\sigma_Y^{\text{pair}}$	20
3.14	$n\sigma_X^{\text{pair}}$ vs. m_{TOF}^2	21
3.15	$\chi^2_{dE/dx}$ and m_{TOF}^2 for exclusive $\pi^+\pi^-$, K^+K^- and $p\bar{p}$ candidates.	22
3.16	Two-dimensional distributions of sum of forward protons momenta and sum of central tracks momenta for exclusive $\pi^+\pi^-$ (top row), K^+K^- (middle row) and $p\bar{p}$ (bottom row) event candidates.	24
3.17	Raw distributions of p_x^{miss} and p_y^{miss} for exclusive $\pi^+\pi^-$, K^+K^- and $p\bar{p}$ candidates.	25
3.18	26
4.1	Sketches of main processes with CEP event topology.	27
4.2	Graph illustrating the misidentification problem.	28
4.3	Comparison of Δz_{vtx} for CEP $\pi^+\pi^-$ events in two ranges of $\Delta\varphi$ between data and embedded MC.	30
4.4	Comparison of Δz_{vtx} for CEP $\pi^+\pi^-$ events between data and embedded MC.	31
4.5	Comparison of collinearity $\Delta\theta$ for CEP $\pi^+\pi^-$ events with $\Delta\varphi > 90^\circ$, between data and embedded MC.	31
4.6	Comparison of p_T^{miss} for CEP $\pi^+\pi^-$ events in two ranges of $\Delta\varphi$ between data and embedded MC.	32
4.7	Comparison of p_T^{miss} for CEP $\pi^+\pi^-$ events between data and embedded MC.	33
5.1	Distribution of weights assigned to selected CEP events.	34
5.2	TOF trigger efficiency as a function of lower p_T of two TOF-matched tracks.	35
5.3	Overall efficiency of the online BBC-small and ZDC veto as a function of instantaneous luminosity.	35
5.4	Sample probability of ET&IT trigger veto due to forward proton interaction with dead material.	37
5.5	Sample distribution of z_{vtx} with fitted normal distribution.	38
5.6	Mean and width of the z_{vtx} distribution as a function of the RHIC fill number.	38

5.7	Overall efficiency of online and offline cuts as a function of instantaneous luminosity.	39
5.8	Efficiency of p_T^{miss} cut as a function of central tracks' momenta.	40
5.9	dE/dx error vs. TOF path length and $\log_2(dx)$ vs. TOF path length for exclusive event candidates.	41
5.10	Pair identification efficiency and misidentification probability as a function of tracks' p_T	42
5.11	Sample RP track reconstruction efficiency in a single z -vertex bin.	43
5.12	Probability of simultaneous inefficiency of the RP track reconstruction calculated from the CEP MC embedded into zero-bias data.	
	43	
5.13	Sketch with definition of Δz_0	44
5.14	Distribution of Δz_0 together with fraction of non-exclusive background.	45
5.15	Vertexing efficiency.	45
5.16	Difference between reconstructed and true parameters of the TPC tracks.	47
5.17	Difference between reconstructed and true momentum components of RP tracks.	47
5.18	Fractions of migrating and fake TPC tracks matched with TOF.	48
5.19	Fractions of migrating and fake RP tracks.	48
5.20	Joint corrections for migrations and fake tracks for TPC and RP tracks.	49
5.21	Pull histogram of number of events per unit of integrated luminosity.	50
5.22	Full closure test.	51
6.1	Estimated systematic uncertainty related to trigger veto induced by interaction with dead material.	52
6.2	Systematic uncertainties of the differential cross sections for CEP of charged particle pairs $\pi^+\pi^-$ as a function of the invariant mass of the pair in the fiducial region explained on the plots.	54
6.3	Systematic uncertainties of the differential cross sections for CEP of charged particle pairs K^+K^- (left) and $p\bar{p}$ (right) as a function of the invariant mass of the pair in the fiducial region explained on the plots.	54
6.4	Systematic uncertainties of the differential cross sections for CEP of charged particle pairs $\pi^+\pi^-$ (let), K^+K^- (middle) and $p\bar{p}$ (right) as a function of the pair rapidity measured in the fiducial region explained on the plots.	55
6.5	Systematic uncertainties of the differential cross sections for CEP of charged particle pairs $\pi^+\pi^-$ (left column), K^+K^- (middle column) and $p\bar{p}$ (right column) as a function of the difference of azimuthal angles of the forward scattered protons (top) and of the sum of the squares of the four-momenta losses in the proton vertices (bottom) measured in the fiducial region explained on the plots.	55
6.6	Systematic uncertainties of the differential cross sections for CEP of charged particle pairs $\pi^+\pi^-$ (top), K^+K^- (middle) and $p\bar{p}$ (bottom) as a function of the invariant mass of the pair in two $\Delta\phi$ regions: $\Delta\phi < 90$ degree (left column) and $\Delta\phi > 90$ degree (right column) measured in the fiducial region explained on the plots.	56
6.7	Systematic uncertainties of the differential cross sections for CEP of $\pi^+\pi^-$ pairs as a function of the rapidity of the pair (left column) difference of azimuthal angles of the forward scattered protons (middle column) and of the sum of the squares of the four-momenta losses in the proton vertices (right column) measured in the fiducial region explained on the plots, separately for three ranges of the $\pi^+\pi^-$ pair invariant mass: $m < 1$ GeV (top), $1 < m < 1.5$ GeV (middle) and $m > 1.5$ GeV (bottom).	57
6.8	Systematic uncertainties of the differential cross sections $d\sigma/dm(\pi^+\pi^-)$ for CEP of $\pi^+\pi^-$ pairs in two $ \vec{p}'_{1,T} - \vec{p}'_{2,T} $ regions: $ \vec{p}'_{1,T} - \vec{p}'_{2,T} < 0.12$ GeV (left) and $ \vec{p}'_{1,T} - \vec{p}'_{2,T} > 0.12$ GeV (right) in the fiducial region and $\Delta\phi < 90$ degree.	57
6.9	Systematic uncertainties of the differential cross sections for CEP of $\pi^+\pi^-$ pairs as a function of $\cos\theta^{\text{CS}}$ (top) and of ϕ^{CS} (bottom) measured in the fiducial region explained on the plots, separately for three ranges of the $\pi^+\pi^-$ pair invariant mass: $m < 1$ GeV (left column), $1 < m < 1.5$ GeV (middle column) and $m > 1.5$ GeV (right column).	58

7.1	Differential cross sections for CEP of charged particle pairs $\pi^+\pi^-$ as a function of the invariant mass of the pair in the fiducial region explained on the plots. Data are shown as solid points with error bars representing the statistical uncertainties. The typical systematic uncertainties are shown as gray boxes for only few data points as they are almost fully correlated between neighboring bins. Predictions from MC models GenEx, DiMe and MBR are shown as histograms. In the lower panels in the bottom plots the ratios of the MC predictions scaled to data and the data are shown.	59
7.2	Differential cross sections for CEP of charged particle pairs K^+K^- (left) and $p\bar{p}$ (right) as a function of the invariant mass of the pair in the fiducial region explained on the plots. Data are shown as solid points with error bars representing the statistical uncertainties. The typical systematic uncertainties are shown as gray boxes for only few data points as they are almost fully correlated between neighboring bins. Predictions from MC models GenEx, DiMe and MBR are shown as histograms. In the lower panels in the bottom plots the ratios of the MC predictions scaled to data and the data are shown.	60
7.3	Differential cross sections for CEP of charged particle pairs $\pi^+\pi^-$ (left), K^+K^- (middle) and $p\bar{p}$ (right) as a function of the pair rapidity measured in the fiducial region explained on the plots. Data are shown as solid points with error bars representing the statistical uncertainties. The typical systematic uncertainties are shown as gray boxes for only few data points as they are almost fully correlated between neighboring bins. Predictions from MC models GenEx, DiMe and MBR are shown as histograms. In the lower panels in the bottom plots the ratios of the MC predictions scaled to data and the data are shown.	60
7.4	Differential cross sections for CEP of charged particle pairs $\pi^+\pi^-$ (left column), K^+K^- (middle column) and $p\bar{p}$ (right column) as a function of the difference of azimuthal angles of the forward scattered protons (top) and of the sum of the squares of the four-momenta losses in the proton vertices (bottom) measured in the fiducial region explained on the plots. Data are shown as solid points with error bars representing the statistical uncertainties. The typical systematic uncertainties are shown as gray boxes for only few data points as they are almost fully correlated between neighboring bins. Predictions from MC models GenEx, DiMe and MBR are shown as histograms. In the lower panels the ratios of the MC predictions scaled to data and the data are shown.	61
7.5	Differential cross sections for CEP of charged particle pairs $\pi^+\pi^-$ (top), K^+K^- (middle) and $p\bar{p}$ (bottom) as a function of the invariant mass of the pair in two $\Delta\phi$ regions: $\Delta\phi < 90$ degree (left column) and $\Delta\phi > 90$ degree (right column) measured in the fiducial region explained on the plots. Data are shown as solid points with error bars representing the statistical uncertainties. The typical systematic uncertainties are shown as gray boxes for only few data points as they are almost fully correlated between neighboring bins. Predictions from MC models GenEx, DiMe and MBR are shown as histograms.	62
7.6	Differential cross sections for CEP of $\pi^+\pi^-$ pairs as a function of the rapidity of the pair (left column) difference of azimuthal angles of the forward scattered protons (middle column) and of the sum of the squares of the four-momenta losses in the proton vertices (right column) measured in the fiducial region explained on the plots, separately for three ranges of the $\pi^+\pi^-$ pair invariant mass: $m < 1$ GeV (top), $1 < m < 1.5$ GeV (middle) and $m > 1.5$ GeV (bottom). Data are shown as solid points with error bars representing the statistical uncertainties. The typical systematic uncertainties are shown as gray boxes for only few data points as they are almost fully correlated between neighboring bins. Predictions from MC models GenEx, DiMe and MBR are shown as histograms. In the lower panels the ratios of the MC predictions scaled to data and the data are shown.	63
7.7	Differential cross sections $d\sigma/dm(\pi^+\pi^-)$ for CEP of $\pi^+\pi^-$ pairs in two $ \vec{p}'_{1,T} - \vec{p}'_{2,T} $ regions: $ \vec{p}'_{1,T} - \vec{p}'_{2,T} < 0.12$ GeV (left) and $ \vec{p}'_{1,T} - \vec{p}'_{2,T} > 0.12$ GeV (right) in the fiducial region and $\Delta\phi < 90$ degree. There is no difference for two $ \vec{p}'_{1,T} - \vec{p}'_{2,T} $ regions. Data are shown as solid points with error bars representing the statistical uncertainties. The typical systematic uncertainties are shown as gray boxes for only few data points as they are almost fully correlated between neighboring bins. Predictions from MC models GenEx, DiMe and MBR are shown as histograms.	64

7.8	Differential cross sections for CEP of $\pi^+\pi^-$ pairs as a function of $\cos\theta^{\text{CS}}$ (top) and of ϕ^{CS} (bottom) measured in the fiducial region explained on the plots, separately for three ranges of the $\pi^+\pi^-$ pair invariant mass: $m < 1 \text{ GeV}$ (left column), $1 < m < 1.5 \text{ GeV}$ (middle column) and $m > 1.5 \text{ GeV}$ (right column). Data are shown as solid points with error bars representing the statistical uncertainties. The typical systematic uncertainties are shown as gray boxes for only few data points as they are almost fully correlated between neighboring bins. Predictions from MC models GenEx, DiMe and MBR are shown as histograms. In the lower panels the ratios of the MC predictions scaled to data and the data are shown.	65
A.1	Distribution of ADC vs. TAC counts (2D) and ADC conunts (1D) per BBC-small channel in abort gaps and colliding bunches.	66
A.2	Distribution of ADC vs. TAC counts (2D) and ADC conunts (1D) per BBC-large channel in abort gaps and colliding bunches.	72
C.1	Scheme of two central tracks with common vertex, hitting cells in TOF detector.	77
D.1	RP track acceptance, reconstruction and selection efficiency on the east (MC embedded into zero-bias data).	78
D.2	RP track acceptance, reconstruction and selection efficiency on the west (MC embedded into zero-bias data).	81
D.3	Probability of ET&IT trigger veto due to forward proton interaction with dead material on the east.	84
D.4	Probability of ET&IT trigger veto due to forward proton interaction with dead material on the west.	87
E.1	Pair identification efficiency and misidentification probability as a function of tracks' momentum.	90

List of Tables

2.1	Central Diffraction physics triggers and control triggers involving Roman Pot detectors in run 15.	7
3.1	Offline ADC thresholds in BBC-large.	18

References

- [1] L. Adamczyk, L. Fulek, and R. Sikora, “Supplementary note on diffractive analyses of 2015 proton-proton data.” <https://drupal.star.bnl.gov/STAR/blog/rafals/supplementary-note-diffractive-analyses-2015-proton-proton-data>, January, 2019.
- [2] P. D. B. Collins, *An Introduction to Regge Theory and High-Energy Physics*. Cambridge Monographs on Mathematical Physics. Cambridge Univ. Press, Cambridge, UK, 2009.
<http://www-spires.fnal.gov/spires/find/books/www?cl=QC793.3.R4C695>.
- [3] S. Donnachie, H. G. Dosch, O. Nachtmann, and P. Landshoff, *Pomeron physics and QCD*, vol. 19. 2002.
- [4] V. Barone and E. Predazzi, *High-Energy Particle Diffraction*, vol. v.565 of *Texts and Monographs in Physics*. Springer-Verlag, Berlin Heidelberg, 2002.
<http://www-spires.fnal.gov/spires/find/books/www?cl=QC794.6.C6B37::2002>.
- [5] P. Lebiedowicz and A. Szczurek, “Exclusive $pp \rightarrow pp\pi^+\pi^-$ reaction: From the threshold to LHC,” *Phys. Rev.* **D81** (2010) 036003, [arXiv:0912.0190 \[hep-ph\]](https://arxiv.org/abs/0912.0190).
- [6] P. Lebiedowicz, O. Nachtmann, and A. Szczurek, “Central exclusive diffractive production of $\pi^+\pi^-$ continuum, scalar and tensor resonances in pp and $p\bar{p}$ scattering within tensor pomeron approach,” *Phys. Rev.* **D93** no. 5, (2016) 054015, [arXiv:1601.04537 \[hep-ph\]](https://arxiv.org/abs/1601.04537).
- [7] L. A. Harland-Lang, V. A. Khoze, and M. G. Ryskin, “Modelling exclusive meson pair production at hadron colliders,” *Eur. Phys. J.* **C74** (2014) 2848, [arXiv:1312.4553 \[hep-ph\]](https://arxiv.org/abs/1312.4553).
- [8] R. Fiore, L. Jenkovszky, and R. Schicker, “Resonance production in Pomeron–Pomeron collisions at the LHC,” *Eur. Phys. J.* **C76** no. 1, (2016) 38, [arXiv:1512.04977 \[hep-ph\]](https://arxiv.org/abs/1512.04977).
- [9] F. E. Close and A. Kirk, “A Glueball - q anti-q filter in central hadron production,” *Phys. Lett.* **B397** (1997) 333–338, [arXiv:hep-ph/9701222 \[hep-ph\]](https://arxiv.org/abs/hep-ph/9701222).
- [10] <https://online.star.bnl.gov/rp/pp200/>.
- [11] <https://online.star.bnl.gov/RunLogRun15/>.
- [12] <http://www.star.bnl.gov/protected/common/common2015/trigger2015/lumipp200GeV/>.
- [13] I. Alekseev and J. H. Lee, “Trigger Definitions for RP phase-II* for Run15.”
https://drupal.star.bnl.gov/STAR/system/files/trigger_updated_run15_v2.doc, December, 2014.
- [14] I. Alekseev, “Trigger with Roman Pots Phase II*.”
https://drupal.star.bnl.gov/STAR/system/files/trigger4RP2_v2.pdf, December, 2014.
- [15] R. Sikora, “Naming convention and layout of the Roman Pot Phase II* subsystem in the STAR experiment.” https://drupal.star.bnl.gov/STAR/system/files/RomanPotNomenclature_0.pdf, October, 2014.
- [16] <http://www.star.bnl.gov/public/comp/prod/DataSummary.html>.
- [17] <https://drupal.star.bnl.gov/STAR/event/2017/09/27/software-and-computing-meeting/revertexing-2015-rp-stream>.
- [18] L. Adamczyk. https://drupal.star.bnl.gov/STAR/system/files/talk_42.pdf, March, 2016.
- [19] <https://github.com/rafalsikora/CEP-STAR>.

- [20] H. Bichsel, “A method to improve tracking and particle identification in TPCs and silicon detectors,” *Nucl. Instrum. Meth.* **A562** (2006) 154–197.
- [21] R. A. Kycia, J. Chwastowski, R. Staszewski, and J. Turnau, “GenEx: A simple generator structure for exclusive processes in high energy collisions,” *Commun. Comput. Phys.* **24** no. 3, (2018) 860–884, arXiv:1411.6035 [hep-ph].
- [22] “RHIC Run 15 Overview.” <http://www.agsrhichome.bnl.gov/RHIC/Runs/index.html#Run-15>.