

SVM

	Przewidziana klasa							
	Czerniak	Rak kolczystokomórkowy	Rak podstawnokomórkowy	Rogowacenie słoneczne	Włókniak	Zmiana naczyniowa	Znamię barwnikowe	Łagodna zmiana
Czerniak	260 (57.5%)	3 (0.7%)	17 (3.8%)	3 (0.7%)	0 (0.0%)	1 (0.2%)	146 (32.3%)	22 (4.9%)
Rak kolczystokomórkowy	4 (6.3%)	23 (36.5%)	15 (23.8%)	11 (17.5%)	0 (0.0%)	0 (0.0%)	6 (9.5%)	4 (6.3%)
Rak podstawnokomórkowy	7 (2.1%)	2 (0.6%)	287 (86.2%)	12 (3.6%)	0 (0.0%)	0 (0.0%)	21 (6.3%)	4 (1.2%)
Rogowacenie słoneczne	10 (11.6%)	2 (2.3%)	17 (19.8%)	40 (46.5%)	0 (0.0%)	0 (0.0%)	3 (3.5%)	14 (16.3%)
Włókniak	0 (0.0%)	0 (0.0%)	2 (8.3%)	1 (4.2%)	3 (12.5%)	0 (0.0%)	17 (70.8%)	1 (4.2%)
Zmiana naczyniowa	0 (0.0%)	0 (0.0%)	5 (20.0%)	0 (0.0%)	0 (0.0%)	17 (68.0%)	3 (12.0%)	0 (0.0%)
Znamię barwnikowe	44 (3.4%)	2 (0.2%)	19 (1.5%)	1 (0.1%)	2 (0.2%)	0 (0.0%)	1202 (93.4%)	17 (1.3%)
Łagodna zmiana	28 (10.6%)	6 (2.3%)	17 (6.5%)	16 (6.1%)	0 (0.0%)	0 (0.0%)	54 (20.5%)	142 (54.0%)