

#### Probability and Statistics

UNIT 4.—

# Statistical Hypothesis Testing

#### Unit 4. Statistical Hypothesis Testing

- What this unit is about:
  - ► You will learn about the principles of hypothesis testing.
  - ► You will learn about the t-test for one sample.
  - ► You will learn about the t-test for independent samples and paired samples.
  - ▶ You will learn about the different kinds of Chi-squared test.
  - ▶ You will learn about the F-test.
- Expected outcome:
  - ► Ability to apply the different kinds of hypothesis testing methods.
- How to check your progress:
  - ► Coding Exercises.
  - Quiz.



#### **Probability and Statistics**

#### UNIT 1. Understanding of Probability

- 1.1. Probability Theory.
- 1.2. Probability Rules.
- 1.3. Random Variable.
- 1.4. Discrete Probability Distribution.

#### Unit 2. Understanding of Statistics I

- 2.1. Continuous Probability Density.
- 2.2. Conjoint Probability.

#### Unit 3. Understanding of Statistics II

- 3.1. Descriptive Statistics.
- 3.2. Central Limit Theorem.
- 3.3. Estimation Theory.

#### **Unit 4. Statistical Hypothesis Testing**

4.1. Principles of Hypothesis Testing.

4.2. Hypothesis Testing in Action.

# Together for Tomorrow! Enabling People Education for Eutrus Generations

#### **4.1.** Principles of Hypothesis Testing.

#### Hypothesis Testing (1/8)

- Principles of the hypothesis testing in statistics:
  - ▶ Tests a hypothesis about the population parameters using the sample statistics.
    - Ex) "The average sleep time of adults in USA is 7 hours."  $\Leftarrow$  True or False?
    - Ex) "The average annual wage in Islamabad is Rs. 85,000."  $\Leftarrow$  True or False?

#### **4.1.** Principles of Hypothesis Testing.




### Hypothesis Testing (2/8)

Principles of the hypothesis testing in statistics:

#### Judge







### Hypothesis Testing (3/8)

- Principles of the hypothesis testing in statistics:
  - ▶ The defendant is assumed innocent until proven otherwise.
  - ► The defendant doesn't need to prove his/her innocence.
  - ▶ The main objective of the trial is to prove the guiltiness of the defendant.
  - ► The judge delivers the verdict based on the presented evidence.

#### **4.1.** Principles of Hypothesis Testing.



### Hypothesis Testing (4/8)

- Principles of the hypothesis testing in statistics:
  - ▶ Null hypothesis *H*0: what is assumed until proven otherwise.
  - ▶ Alternative hypothesis *H*1: what needs to be proven.
  - ► Test statistic: a quantity used as the evidence; calculated from the sample.
  - p-value: probability of observing the current test statistic or more extreme one assuming that the null hypothesis is true.
    - a) If the p-value is small: weakens the assumption of the null hypothesis.
    - b) If the p-value is large: strengthens the assumption of the null hypothesis.

#### 4.1. Principles of Hypothesis Testing.




### Hypothesis Testing (5/8)

- Principles of the hypothesis testing in statistics:
  - Significance level  $\alpha$ :
    - a) It's the reference probability used when deciding whether the p-value is small enough or not.
    - b) If p-value  $\geq \alpha$ , the  $H_0$  is maintained. If p-value  $< \alpha$ , the  $H_0$  is rejected in favor of the  $H_1$ .
    - c) This can also be interpreted as the maximum probability of rejecting the  $H_0$  even when it's true.
    - d)  $1-\alpha$  is the probability that the true  $H_0$  is retained.
  - ▶ Statistical power: probability of rejecting the false  $H_0$  in favor of  $H_1$ .



### Hypothesis Testing (6/8)

- Principles of the hypothesis testing in statistics:
  - ► Procedure using the test statistic:




#### **4.1.** Principles of Hypothesis Testing.



### Hypothesis Testing (7/8)

- Principles of the hypothesis testing in statistics:
  - ► Procedure using the p-value:



#### **4.1.** Principles of Hypothesis Testing.



# Hypothesis Testing (8/8)

Principles of the hypothesis testing in statistics:

ACTUALLY TEST RESULT	$m{H_0}$ is true	$H_0$ IS FALSE
$H_0$ RETAINED	Correct Decision. Probability $=1-lpha$	<b>Type 2 error.</b> Probability= $eta$
$H_0$ REJECTED & $H_1$ ADOPTED	<b>Type 1 error.</b> Probability $= lpha$	<b>Correct Decision.</b> Probability $=1-eta$



#### **Probability and Statistics**

#### UNIT 1. Understanding of Probability

- 1.1. Probability Theory.
- 1.2. Probability Rules.
- 1.3. Random Variable.
- 1.4. Discrete Probability Distribution.

#### Unit 2. Understanding of Statistics I

- 2.1. Continuous Probability Density.
- 2.2. Conjoint Probability.

#### Unit 3. Understanding of Statistics II

- 3.1. Descriptive Statistics.
- 3.2. Central Limit Theorem.
- 3.3. Estimation Theory.


#### **Unit 4. Statistical Hypothesis Testing**

- 4.1. Principles of Hypothesis Testing.
- 4.2. Hypothesis Testing in Action.



### Hypothesis Test of the Means (1/8)

- One sample t-test:
  - ► There is one population and one sample.



► Student-t distribution is used to interpret the test statistic calculated as following:


$$t = \frac{\overline{x} - \mu_0}{S / \sqrt{n}}$$





### Hypothesis Test of the Means (2/8)

- One sample t-test:
  - ► Right tail test:

$$H_0: \mu \leq \mu_0$$








Position of the test statistic



### Hypothesis Test of the Means (3/8)

- One sample t-test:
  - ► Left tail test:





### Hypothesis Test of the Means (4/8)

- One sample t-test:
  - ► Two tail test:


 $H_0: \mu = \mu_0$   $H_1: \mu \neq \mu_0$  "different"

• p-value =  $2 \times P(X > |test \ statistic|)$ Reject  $H_0$   $\alpha/2$ Position of the test statistic.



### Hypothesis Test of the Means (5/8)

- Independent two sample t-test:
  - ► There are two populations and two samples.





# Hypothesis Test of the Means (6/8)

#### Independent two sample t-test:

► Right tail test:

$$H_0: \mu_1 - \mu_2 \leq 0 \iff \mu_1 \leq \mu_2$$

$$H_1: \mu_1 - \mu_2 > 0 \iff \mu_1 > \mu_2$$

Left tail test:

$$H_0: \mu_1 - \mu_2 \ge 0 \iff \mu_1 \ge \mu_2$$

$$H_1: \mu_1 - \mu_2 < 0 \iff \mu_1 < \mu_2$$

► Two tail test:

$$H_0: \mu_1 - \mu_2 = 0 \iff \mu_1 = \mu_2$$

$$H_1: \mu_1 - \mu_2 \neq 0 \iff \mu_1 \neq \mu_2$$


You should consider two cases:

- 1) Equal variances.
- 2) Unequal variances.



### Hypothesis Test of the Means (7/8)

- Paired two sample t-test:
  - ▶ There are two populations and two samples. There is "one on one" pairing.



Ex) Change in the blood pressure of the same test subjects before and after taking a new drug.



### Hypothesis Test of the Means (8/8)

- Analysis of Variance (ANOVA):
  - ▶ So far with the t-test, we had one or two groups (samples).
  - ► ANOVA can detect differences in the means of two or more groups.

Null hypothesis

$$H_0$$
:  $\mu 1 = \mu 2 = \mu 3 = ...$ 

Alternative hypothesis  $H_1$ : There is at least one case where  $\mu i \neq \mu j$ 

- ► Assumptions: 1) The distribution of the data is Normal.
  - 2) The group variances are the same.
  - 3) The groups are independent from each other.
- ► F distribution is used to calculate the p-value.



### Coding Exercise #0307

Follow practice steps on 'ex\_0307.ipynb' file.



### Hypothesis Test of the Frequencies (1/4)

- Chi-squared test for one way table:
  - ▶ One way table or "frequency table" summarizes a categorical variable.
  - ► Compares the observed frequencies with a given model (expected frequencies).

Null hypothesis  $H_0$ : The observed frequency table and the expected model agree.

Alternative hypothesis  $H_1$ : The observed frequency table and the expected model are different.

► Also called the "Goodness of fit test".



### Hypothesis Test of the Frequencies (2/4)

- Chi-squared test for one way table:
  - ► The test statistic is:

$$test \ statistic = \sum_{i=1}^{k} \frac{(O_i - E_i)^2}{E_i}$$

- $\triangleright$   $E_i$  are the expected frequencies and  $O_i$  are the observed frequencies.
- $\triangleright$  k is the number of categories or types.
- ▶ The test statistic follows the Chi-square distribution of degree of freedom = k-1.



### Hypothesis Test of the Frequencies (3/4)

- Chi-squared test for two way table:
  - ► A contingency table summarizes two categorical variables.
    - Ex) Confusion matrix (machine learning).
  - ▶ Uses the frequencies to test the existence of relationship between two categorical variables.

Null hypothesis  $H_0$ : The categorical variables are independent.

Alternative hypothesis  $H_1$ : The categorical variables are not independent.

► Also called "Independence test".



### Hypothesis Test of the Frequencies (4/4)

- Chi-squared test for two way table:
  - ► The test statistic is:

$$test \ statistic = \sum_{i=1}^{r} \sum_{j=1}^{c} \frac{\left(O_{ij} - E_{ij}\right)^{2}}{E_{ij}}$$

- $\triangleright$   $E_i$  are the expected frequencies and  $O_i$  are the observed frequencies.
- ightharpoonup r is the number of rows and c is the number of columns in the two way table.
- ▶ The test statistic follows the Chi-square distribution of degree of freedom =  $(r-1) \times (c-1)$ .



### Coding Exercise #0308

Follow practice steps on 'ex\_0308.ipynb' file.




### Coding Exercise #0309

Follow practice steps on 'ex\_0309.ipynb' file.



### Hypothesis Test of the Variances (1/4)

- One sample t-test:
  - There is one population and one sample.





### Hypothesis Test of the Variances (2/4)

- Chi-squared test of variance:
  - ► There are left tail test, right tail test and two tail test. ← Just like in t-test.
  - ightharpoonup The test statistic is calculated as: (n= sample size)

$$test \ statistic = \frac{(n-1)S^2}{\sigma_0^2}$$

► The test statistic follows the Chi-square distribution of degree of freedom = n-1.




### Coding Exercise #0310

Follow practice steps on 'ex\_0310.ipynb' file.



### Hypothesis Test of the Variances (3/4)

- F-test of variance ratio:
  - There are two populations and two samples.





### Hypothesis Test of the Variances (4/4)

- F-test of variance ratio:
  - ▶ There are left tail test, right tail test and two tail test.
  - ▶ The test statistic is calculated as a ratio of the sample variances:

$$test \ statistic = \frac{S_1^2}{S_2^2}$$

▶ The test statistic follows the F distribution  $F(n_1 - 1, n_2 - 1)$ . Here  $n_1$  and  $n_2$  are the sample sizes.



### Hypothesis Test Summary

Here, this summarizes the hypothesis tests we have covered so far:

HYPOTHESIS TEST	PROBABILITY DENSITY DISTRIBUTION
One sample t-test, Independent two sample t-test, Paired sample t-test.	Student-t
ANOVA	F
Chi-squared test of one way table, Chi-squared test of two way table.	Chi-square
Chi-squared test of variance.	Chi-square
F-test of variance ratio.	F




### Coding Exercise #0311





### Coding Exercise #0312







#### SAMSUNG

# Together for Tomorrow! Enabling People

**Education for Future Generations** 

©2019 SAMSUNG. All rights reserved.

Samsung Electronics Corporate Citizenship Office holds the copyright of book.

This book is a literary property protected by copyright law so reprint and reproduction without permission are prohibited.

To use this book other than the curriculum of Samsung innovation Campus or to use the entire or part of this book, you must receive written consent from copyright holder.