

Dedicated to Prof. Wai Chee Shiu
on the occasion of his 66th birthday

COMPLETE CHARACTERIZATION OF GRAPHS WITH LOCAL TOTAL ANTIMAGIC CHROMATIC NUMBER 3

Gee-Choon Lau

Communicated by Ingo Schiermeyer

Abstract. A total labeling of a graph $G = (V, E)$ is said to be local total antimagic if it is a bijection $f : V \cup E \rightarrow \{1, \dots, |V| + |E|\}$ such that adjacent vertices, adjacent edges, and pairs of an incident vertex and edge have distinct induced weights where the induced weight of a vertex v is $w_f(v) = \sum f(e)$ with e ranging over all the edges incident to v , and the induced weight of an edge uv is $w_f(uv) = f(u) + f(v)$. The local total antimagic chromatic number of G , denoted by $\chi_{lt}(G)$, is the minimum number of distinct induced vertex and edge weights over all local total antimagic labelings of G . In this paper, we first obtain general lower and upper bounds for $\chi_{lt}(G)$ and sufficient conditions to construct a graph H with k pendant edges and $\chi_{lt}(H) \in \{\Delta(H) + 1, k + 1\}$. We then completely characterize graphs G with $\chi_{lt}(G) = 3$. Many families of (disconnected) graphs H with k pendant edges and $\chi_{lt}(H) \in \{\Delta(H) + 1, k + 1\}$ are also obtained.

Keywords: local total antimagic, local total antimagic chromatic number.

Mathematics Subject Classification: 05C78, 05C69.

1. INTRODUCTION

Let $G = (V, E)$ be a simple and loopless graph of order p and size q . For integers $a < b$, let $[a, b] = \{n \in \mathbb{Z} \mid a \leq n \leq b\}$. If no adjacent vertices of G are assigned the same color, then G has a proper (vertex) coloring. The smallest required number of colors is the chromatic number of G , denoted $\chi(G)$. The most famous problem on the chromatic number is the 4-Color Conjecture which states that every planar graph G has $\chi(G) = 4$. Interested readers may refer to [2] for the proof by Appel and Haken. If no two adjacent edges of G are assigned the same color, then G has a proper edge coloring. The smallest required number of colors is the chromatic index of G , denoted $\chi'(G)$. Interested readers may refer to [16] for an interesting brief history. A total coloring of G is a coloring of the vertices and edges of G such that x and y are assigned distinct colors whenever x and y are adjacent vertices or edges, or incident vertex and

edge. The smallest number of colors required for a total coloring is the total chromatic number of G , denoted $\chi_t(G)$. The conjecture that

$$\Delta(G) + 1 \leq \chi_t(G) \leq \Delta(G) + 2$$

remained unsolved (see [4, 6]) where $\Delta(G)$ is the maximum degree of G .

A bijection $f : E \rightarrow [1, q]$ is a local antimagic labeling if $f^+(u) \neq f^+(v)$ for every edge uv of G where $f^+(u)$ is the sum of all the incident edge label(s) of u under f . Let $f^+(u)$ be the color of u . Clearly, f^+ is a proper vertex coloring of G induced by f . The smallest number of vertex colors taken over all local antimagic labeling f is called the local antimagic chromatic number of G , denoted $\chi_{la}(G)$ (see [3, 7–9, 11, 12] for the many results available).

A bijection $g : V \rightarrow [1, p]$ is a local edge antimagic labeling if $g_{le}(e_1) \neq g_{le}(e_2)$ for every two adjacent edges e_1 and e_2 of G where $g_{le}(uv) = g(u) + g(v)$. Let $g_{le}(e)$ be the induced edge color of e under g . Clearly, g_{le} is a proper edge coloring of G induced by g . The smallest number of edge color(s) over all local edge antimagic labeling g is called the local edge antimagic chromatic number of G , denoted $\chi_{lea}(G)$ [1].

A bijection $f : V \cup E \rightarrow [1, p+q]$ is a local antimagic total labeling if $w^+(u) \neq w^+(v)$ for every two adjacent vertices u and v where $w^+(u) = f(u) + \sum f(e)$ over all edge(s) e incident to u . Let $w^+(u)$ be the induced vertex color of u under f . Clearly, w^+ is a proper vertex coloring of G induced by f . The smallest number of vertex colors over all local antimagic total labeling f is called the local antimagic total chromatic number of G , denoted $\chi_{lat}(G)$. In [10, 13], the authors proved that every graph is local antimagic total and obtained the exact $\chi_{lat}(G)$ for many families of graphs G .

Motivated by the concepts of local (edge) antimagic labeling and total labeling of G , Sandhiya and Nalliah [14] defined the concept of local total antimagic labeling. A bijection $f : V \cup E \rightarrow [1, p+q]$ such that the weight of a vertex u is $w_f(u) = \sum f(e)$ over all edge(s) e incident to u , and the weight of an edge $e = uv$ is $w_f(e) = f(u) + f(v)$, is called a *local total antimagic labeling* if (i) every two adjacent vertices and adjacent edges have distinct weights, and (ii) every vertex and its incident edge(s) have distinct weights. The mapping w_f (or w if no ambiguity) is called a *local total antimagic labeling of G induced by f* , and the weights assigned to vertices and edges are called *induced total colors* under f .

The *local total color number* of a local total antimagic labeling f is the number of distinct induced total colors under f , denoted by $w(f)$. Moreover, f is called a *local total antimagic $w(f)$ -coloring* and G is *local total antimagic $w(f)$ -colorable*. The *local total antimagic chromatic number* $\chi_{lt}(G)$ is defined to be the minimum number of colors taken over all local total antimagic colorings of G induced by local total antimagic labelings of G . Let $G + H$ and mG denote the disjoint union of graphs G and H , and m copies of G , respectively. By definition, any graph with an isolated vertex or a K_2 component does not admit a local total antimagic labeling. In this paper, we only consider graph G of order $p \geq 3$ and size $q \geq 2$. Thus, G contains neither isolated vertex nor K_2 component.

In Section 2, we obtained various sufficient conditions on bounds of $\chi_{lt}(G)$. In Section 3, we give complete characterization of graphs G with $\chi_{lt}(G) = 3$. In Section 4, we first obtained various families of disconnected graph with k pendant edges and

$\chi_{lt}(G) = k + 1$. Using the results in Section 3, we then obtained various new families of graphs H , constructed from G , having s pendant edges and $\chi_{lt}(G) = s + 1$. Open problems and conjectures are given in Section 5.

2. SUFFICIENT CONDITIONS

In [14], the authors proved that a graph G with $k \geq 1$ pendant edge(s) has $\chi_{lt}(G) \geq k + 1$. We now give a necessary condition and a proof for equality to hold.

Lemma 2.1. *Suppose G has $k \geq 1$ pendant edges, then $\chi_{lt}(G) \geq k + 1$. If $\chi_{lt}(G) = k + 1$, then $p + q$ is assigned to a pendant vertex or a pendant edge.*

Proof. For completeness, we only need to prove the second statement. Assume that $\chi_{lt}(G) = k + 1$. Suppose $p + q$ is assigned to a non-pendant vertex, say x . Now, all the k pendant vertices have distinct weights at most $p + q - 1$. Moreover, x is incident to two edges, say e and e' . Clearly, $w(e), w(e')$ are distinct weights at least $p + q + 1$. Suppose $p + q$ is assigned to a non-pendant edge, say xy . Now, all the k pendant vertices have distinct weights at most $p + q - 1$ and $w(x), w(y)$ are distinct weights at least $p + q + 1$. In both cases, there are at least $k + 2$ distinct weight so that $\chi_{lt}(G) \geq k + 2$. This contradicts $\chi_{lt}(G) = k + 1$. \square

Corollary 2.2. *Suppose G is a graph with $k \geq 1$ pendant edges, then*

$$\chi_{lt}(G) \geq \max\{\Delta(G) + 1, k + 1\} \geq 3.$$

Proof. By definition and Lemma 2.1, $\chi_{lt}(G) \geq \max\{\Delta(G) + 1, k + 1\}$. Thus, if $\chi_{lt}(G) = 2$, then $\Delta(G) = 1$ or $k = 1$. If the former holds, G is a 1-regular graph that does not admit a local total antimagic labeling. If the latter holds, $\Delta(G) \geq 3$ so that $\chi_{lt}(G) \geq \chi_t(G) \geq 4$, a contradiction. Consequently, $\chi_{lt}(G) \geq 3$ if G admits a local total antimagic labeling. \square

In [14], the authors also proved that if G is an r -regular graph, $r \geq 2$, that admits a $\chi_{la}(G)$ -labeling and a $\chi_{lea}(G)$ -labeling, then $\chi_{lt}(G) \leq \chi_{la}(G) + \chi_{lea}(G)$. We now give a more general result.

Theorem 2.3. *Suppose G admits a $\chi_{lea}(G)$ -labeling. If G has (i) $\delta(G) \geq 2$; or else (ii) $\delta(G) = 1$ so that G has $k \geq 1$ pendant vertices, v_p non-pendant vertices and e_p non-pendant edges with $e_p > v_p + k - 2$ that admits a $\chi_{la}(G)$ -labeling that assigns the non-pendant edges by integers in $[1, e_p]$, then G admits a local total antimagic labeling. Moreover, $\chi_{lt}(G) \leq \chi_{la}(G) + \chi_{lea}(G)$.*

Proof. Suppose G has order p and size q . By definition, G has no K_2 components. Let $f : E \rightarrow [1, q]$ (and $h : V \rightarrow [1, p]$) be a $\chi_{la}(G)$ - (and $\chi_{lea}(G)$ -) labeling of G . Define a bijective total labeling $g : V \cup E \rightarrow [1, p + q]$ such that $g(u) = h(u)$ for each vertex u of G , and $g(e) = f(e) + p$ for each edge e of G . Therefore, $\{g(u)\} = [1, p]$ and $\{g(e)\} = [p + 1, p + q]$. Clearly, $w_g(u) = w_g(v)$ if and only if $f^+(u) = f^+(v)$, and $w_g(e_1) = w_g(e_2)$ if and only if $h_{le}(e_1) = h_{le}(e_2)$ for all $u, v \in V(G)$ and $e_1, e_2 \in E(G)$.

(i) $\delta \geq 2$. In this case, G has no pendant edges. Thus, every edge weight under g is at most $2p - 1$ and every vertex weight under g is at least $2p + 3$. Therefore, every edge weight is less than every vertex weight. Consequently, g is a local total antimagic labeling that induces $\chi_{la}(G) + \chi_{lea}(G)$ distinct weights.

(ii) $\delta = 1$. Thus, G has order $p = v_p + k$ and size $q = e_p + k$. By the given assumption, the $\chi_{la}(G)$ -labeling f of G assigns each non-pendant edge with an integer in $[1, e_p]$. In this case, under g , every vertex is assigned an integer in $[1, v_p + k]$, every non-pendant edge is assigned an integer in $[v_p + k + 1, v_p + k + e_p]$ while every pendant edge is assigned an integer in $[v_p + k + e_p + 1, v_p + 2k + e_p]$. Now, every edge weight is at most $2(v_p + k) - 1$. Every pendant vertex weight is at least $v_p + k + e_p + 1 > 2(v_p + k) - 1$ and every non pendant vertex weight is at least $2(v_p + k) + 3$. Therefore, every edge weight is less than every vertex weight. Consequently, g is a local total antimagic labeling that induces $\chi_{la}(G) + \chi_{lea}(G)$ distinct weights. \square

Suppose G is a graph with $\chi_{lt}(G) = \Delta(G) + 1$. We now give sufficient condition to construct new graph H with $k \geq 2$ pendants edges from G to have $\chi_{lt}(H) = \Delta(H) + 1$.

Theorem 2.4. *For $k, s \geq 1, ks \geq 2$, let G be a graph of order p and size q with $d \geq 0$ pendant edges such that*

- (a) $\chi_{lt}(G) = t = \Delta(G) + 1$ and the corresponding local total antimagic labeling f assigns k to a maximum degree vertex v of G ;
- (b) for $k \geq 1, s \geq 1$,

$$w_f(v) + \sum_{j=1}^s \sum_{i=1}^k (p + q + 2jk + 1 - i) \neq w_f(x),$$

for each vertex x adjacent to v ;

- (c) v is the only element with weight $w_f(v)$, or else there is a non-pendant vertex element in $V(G) \cup E(G)$ with weight in $\{p + q + (2j - 1)k + i \mid 1 \leq j \leq s, 1 \leq i \leq k\}$.

Suppose $G_v(k, s)$ is obtained from G by attaching $ks \geq 2$ pendant edges to v . If Conditions (a), (b) are satisfied, then

$$\Delta(G_v(k, s)) + 1 \leq \chi_{lt}(G_v(k, s)) \leq \Delta(G_v(k, s)) + 2.$$

The lower bound holds if Condition (c) is also satisfied.

Proof. Let G be a graph satisfying Condition (a). Clearly, v is not a pendant vertex. Lemma 2.1 implies that $\Delta(G) \geq d$ if $d > 0$. Note that $G_v(k, s)$ has $ks + d$ pendant edges. So,

$$\Delta(G_v(k, s)) = \Delta(G) + ks \geq ks + d.$$

By definition,

$$\chi_{lt}(G_v(k, s)) \geq \Delta(G_v(k, s)) + 1 = \Delta(G) + ks + 1.$$

Let the $ks \geq 2$ pendant edges added to G to get H be $e_{j,i} = vx_{j,i}$ for $1 \leq j \leq s$, $1 \leq i \leq k$. For simplicity, if $k = 1$, let $e_{j,1} = e_j$ and $x_{j,1} = x_j$. Define a total

labeling $g : V(G_v(k, s)) \cup E(G_v(k, s)) \rightarrow [1, p + q + 2ks]$ such that $g(x) = f(x)$ for each $x \in V(G) \cup E(G)$. For the remaining vertices and edges, we do as follows.

- (i) For $k = 1$, let $g(x_j) = p + q + 2j - 1$ and $g(e_j) = p + q + 2(s - j + 1)$. If $s \geq 3$ is odd, then swap the edge labels of $e_{(s+1)/2}$ and $e_{(s-1)/2}$.
- (ii) For $k \geq 2$, let $g(x_{j,i}) = p + q + 2(j - 1)k + i$, and $g(e_{j,i}) = p + q + 2jk + 1 - i$ for $1 \leq j \leq s, 1 \leq i \leq k$. If $k \geq 3$ is odd, then swap the edge labels of $e_{j,(k+1)/2}$ and $e_{j,(k-1)/2}$ for each $1 \leq j \leq s$.

Clearly, for distinct $x, y \in (V(G) \cup E(G)) \setminus \{v\}$, $w_g(x) = w_g(y)$ if and only if $w_f(x) = w_f(y)$. Observe that

$$\begin{aligned} \{g(e_{j,i}) \mid 1 \leq j \leq s, 1 \leq i \leq k\} &= \bigcup_{j=1}^s [p + q + (2j - 1)k + 1, p + q + 2jk] \\ &= \{w_g(e_{j,i}) \mid 1 \leq j \leq s, 1 \leq i \leq k\} \\ &= \{w_g(x_{j,i}) \mid 1 \leq j \leq s, 1 \leq i \leq k\}, \end{aligned}$$

denoted W . Clearly, all the elements in W are distinct. Moreover, every edge e of G that is incident to v has weight

$$\begin{aligned} w_g(e) = w_f(e) &\leq 2p + 2q - 1 \\ &< w_f(v) + \sum_{j=1}^s \sum_{i=1}^k (p + q + (2j - 1)k + i) = w_g(v). \end{aligned}$$

So, Condition (b) implies that g is a local total antimagic labeling of H with weights set $W_g = W \cup (W_f \setminus \{w_f(v)\}) \cup \{w_g(v)\}$ that has size at most

$$ks + t + 1 = \Delta(G) + ks + 2 = \Delta(G_v(k, s)) + 2.$$

Thus,

$$\Delta(G_v(k, s)) + 1 \leq \chi_{lt}(G_v(k, s)) \leq \Delta(G_v(k, s)) + 2.$$

Consider Condition (c). Suppose v is the only element with weight $w_f(v)$. Now g induces $t - 1$ distinct weights among the elements in $(V(G) \cup E(G)) \setminus \{v\}$. Therefore, W_g has size

$$ks + (t - 1) + 1 = ks + t = \Delta(G_v(k, s)) + 1.$$

Otherwise, since there is a non-pendant vertex element in $(V(G) \cup E(G)) \setminus \{v\}$ with weight in $\{p + q + (2j - 1)k + i \mid 1 \leq j \leq s, 1 \leq i \leq k\}$, we also have W_g has size $\Delta(G_v(k, s)) + 1$. This completes the proof. \square

Suppose G has $d_G \geq 1$ pendant edges and $\chi_{lt}(G) = d_G + 1 \geq \Delta(G) + 1$. Let H be obtained from G by attaching pendant edges to a vertex of G such that H has d_H pendant edges. We now give sufficient condition for $\chi_{lt}(H) = d_H + 1$. Let $\deg_G(v)$ be the degree of vertex v in G .

Theorem 2.5. For $k, s \geq 1$, $ks \geq 2$, let G be a graph of order p and size q with d_G pendant edges such that

- (i) $\chi_{lt}(G) = d_G + 1 \geq \Delta(G) + 1 \geq 3$ and the corresponding local total antimagic labeling f assigns k to a vertex v of G ,
- (ii) if x is a vertex adjacent to v , then

$$w_f(x) \neq w_f(v) + \sum_{j=1}^s \sum_{i=1}^k (p + q + (2j - 1)k + i),$$

- (iii) if v is not the only element with weight $w_f(v)$, then w or else $w_f(v) \in \{g(e_{j,i}) \mid 1 \leq j \leq s, 1 \leq i \leq k\}$, where w is the only weight under f which is not a pendant edge label.

Suppose $G_v(k, s)$ is obtained from G by attaching ks pendant edges to v .

- (a) If v is a pendant vertex, then

$$d_H + 1 = ks + d_G \leq \chi_{lt}(G_v(k, s)) \leq ks + d_G + 1 = d_H + 2.$$

The lower bound holds if (i) there is a weight of G under f equal to $p+q+(2j-1)k+i$ for $1 \leq j \leq s, 1 \leq i \leq k$; or else (ii) v is the only element with weight $w_f(v)$ under f .

- (b) If v is not a pendant vertex, then $\chi_{lt}(G_v(k, s)) = ks + d_G + 1$.

Proof. Let the ks vertices added to G to get $H = G_v(k, s)$ be $x_{j,i}$ for $1 \leq j \leq s, 1 \leq i \leq k$ and let $vx_{j,i}$ be $e_{j,i}$. Since H has order $p + ks$ and size $q + ks$, we define a total labeling $g : V(H) \cup E(H) \rightarrow [1, p + q + 2ks]$ such that g is as defined in the proof of Theorem 2.4. Obviously, every edge incident to v has weight less than $w_g(e_{j,i})$. Moreover, all but one of the weights of G under f , say w , must be an edge label of G under f . Moreover,

$$w_g(v) = w_f(v) + \sum_{j=1}^s \sum_{i=1}^k (p + q + (2j - 1)k + i).$$

By definition and Lemma 2.1, we know $d_G \geq \Delta(G) \geq 2$.

(a) If v is a pendant vertex, then H has $d_H = ks + d_G - 1 \geq ks + 1$ pendant edges with $\deg_H(v) = ks + 1$. Since $ks \geq 2$, we also have $ks + d_G - 1 \geq d_G + 1 > \Delta(G)$ so that $ks + d_G - 1 \geq \max\{ks + 1, \Delta(G)\} = \Delta(H)$. By Lemma 2.1, $\chi_{lt}(H) \geq ks + d_G = d_H + 1$. Note that $w_g(x) = w_g(y)$ if and only if $w_f(x) = w_f(y)$ for each $x, y \in (V(G) \cup E(G)) \setminus \{v\}$. Clearly, $w_f(v)$ is unique. If w does not equal to $p + q + (2j - 1)k + i$ for $1 \leq j \leq s, 1 \leq i \leq k$, then Condition (ii) implies that g is a local total antimagic labeling that induces $d_G + ks + 1$ distinct weights so that $d_G + ks \leq \chi_{lt}(H) \leq d_G + ks + 1$. Otherwise, g induces $d_G + ks$ distinct weights and the lower bound holds.

(b) If v is not a pendant vertex, then $\Delta(G) \geq \deg_G(v) = r \geq 2$ and H has $ks + d_G$ pendant edges with $\deg_H(v) = ks + r$ so that $ks + d_G \geq \max\{ks + r, \Delta(G)\} = \Delta(H)$.

By Lemma 2.1, $\chi_{lt}(H) \geq ks + d_G + 1$. Note that $w_g(x) = w_g(y)$ if and only if $w_f(x) = w_f(y)$ for each $x, y \in (V(G) \cup E(G)) \setminus \{v\}$. If v is the only element with weight $w_f(v)$, then Condition (ii) implies that g is a local total antimagic labeling that induces $ks + d_G + 1$ distinct weights. Otherwise, Condition (iii) implies that g is a local total antimagic chromatic labeling that induces $ks + d_G + 1$ distinct weights. Thus, $\chi_{lt}(H) = ks + d_G + 1$.

This completes the proof. \square

Theorem 2.6. *Suppose G is a graph of order p and size q with exactly one vertex of maximum degree $\Delta \geq 3$ which is not adjacent to any pendant vertex and all other vertices of G has degree at most $m < \Delta$. Moreover, G has $k \geq \Delta \geq 2$ pendant edges such that*

$$\Delta(\Delta + 1) > \max\{m[2(p + q) - m + 1], 4(p + q) - 2\}.$$

If G admits a local total antimagic labeling, then $\chi_{lt}(G) \geq k + 2$.

Proof. Let f be a local total antimagic labeling of G . If $p + q$ is assigned to a non-pendant vertex or edge, by Lemma 2.1, $\chi_{lt}(G) \geq k + 2$. Suppose $p + q$ is assigned to a pendant edge that has a non-pendant end-vertex x . Now, all the k induced pendant vertex labels are distinct and at most $p + q$. Suppose u is the vertex of maximum degree Δ , then $w(u) \geq \Delta(\Delta + 1)/2$ and $p + q + 1 \leq w(x) \leq m[2(p + q) - m + 1]/2$. By the given hypothesis, $w(u) > w(x) > w(y)$ for every pendant vertex y . Thus, f induces at least $k + 2$ distinct vertex weights.

Suppose $p + q$ is assigned to a pendant vertex, then the adjacent pendant edge, say e , has $p + q + 1 \leq w(e) \leq 2(p + q) - 1$. Moreover, all the k induced pendant vertex labels are distinct and at most $p + q - 1$. By the given hypothesis, $w(u) \geq \Delta(\Delta + 1)/2 > 2(p + q) - 1 \geq w(e)$. Thus, f induces at least $k + 2$ distinct vertex weights. \square

Let f_n be the friendship graph obtained from $n \geq 2$ copies of K_3 with a common vertex c . Let $f_n(k)$, $k \leq 2n - 3$, be obtained from f_n by attaching exactly k pendant edges to every degree 2 vertex of f_n . Now, $f_n(k)$ has order $p = n(2k + 2) + 1$ and size $q = n(2k + 3)$ with exactly one vertex c with maximum degree $\Delta = 2n$ with is not adjacent to any pendant vertex and all other vertices has degree at most $m = k + 2 \leq 2n - 1 < \Delta$. Thus,

$$\begin{aligned} \max\{m[(2(p + q) - m + 1)], 4(p + q) - 2\} &= m[2(p + q) - m + 1] \\ &= 2n(k + 2)(4k + 5) - (k + 2)(k - 1). \end{aligned}$$

Thus,

$$\Delta(\Delta + 1) > m[2(p + q) - m + 1]$$

implies that

$$2n(2n + 1) - 2n(k + 2)(4k + 5) + (k + 2)(k - 1) > 0.$$

By Lemma 2.1 and Theorem 2.6, we have the following.

Corollary 2.7. *For $2n \geq k+3 \geq 4$, and $2n(2n+1)-2n(k+2)(4k+5)+(k+2)(k-1) > 0$, $\chi_{lt}(f_n(k)) \geq 2nk + 2$.*

Note that the above corollary is always attainable. For example, take $k = 1$, we have $n > 13$.

3. GRAPHS WITH $\chi_{lt} = 3$

By Corollary 2.2, $\chi_{lt}(G) = 3$ only if $\Delta(G) = 2$ possibly with exactly one path component of order at least 3. Let $P_n = u_1u_2 \dots u_n$ be the path of order $n \geq 3$ with $e_i = u_iu_{i+1}$ for $1 \leq i \leq n-1$. The authors in [14] also obtained $\chi_{lt}(P_n) = 3$ for $n = 3, 6$ and $\chi_{lt}(P_n) = 4$ for $n = 4$ and odd $n \geq 5$. Moreover, $3 \leq \chi_{lt}(P_n) \leq 5$ for even $n \geq 8$. We first improve the lower bound of the last statement in the following lemma.

Lemma 3.1. *For $n = 3, 6$, $\chi_{lt}(P_n) = 3$. If $n = 4$ or $n \geq 5$ is odd, $\chi_{lt}(P_n) = 4$. Otherwise, $4 \leq \chi_{lt}(P_n) \leq 5$ for even $n \geq 8$.*

Proof. From the proofs in [14, Theorems 2.3 and 2.5], we only need to show that $\chi_{lt}(P_n) \geq 4$ for $n \geq 7$. Suppose $\chi_{lt}(P_n) = 3$ and f is a corresponding local total antimagic 3-coloring. Without loss of generality, we may assume the 3 distinct weights are a, b, c such that

$$\begin{aligned} a &= w(u_1) = w(e_2) = w(u_4) = w(e_5), \\ b &= w(e_1) = w(u_3) = w(e_4) = w(u_6) \\ c &= w(u_2) = w(e_3) = w(u_5) = w(e_6). \end{aligned}$$

Thus, we have

$$\begin{aligned} a &= f(e_1) = f(e_3) + f(e_4) \\ b &= f(e_2) + f(e_3) = f(e_5) + f(e_6) \\ c &= f(e_1) + f(e_2) = f(e_4) + f(e_5). \end{aligned}$$

Thus, from $a + b = c$, we get $f(e_3) = f(e_3) + f(e_6)$, a contradiction. Therefore, $\chi_{lt}(P_n) \geq 4$. The lemma holds. \square

By Corollary 2.2, the proofs in [14, Theorems 2.3 and 2.5] and the argument of Lemma 3.1, we have the following.

Corollary 3.2. *If $\chi_{lt}(G) = 3$, then G is a 2-regular graph or a path $P_n, n = 3, 6$, or $H + P_3$ or $H + P_6$, where H is a 2-regular graph.*

In what follows, if a 2-regular graph has i -th component of order $n \geq 3$, then the consecutive vertices and edges are $u_{i,1}, e_{i,1}, u_{i,2}, e_{i,2}, \dots, u_{i,n}, e_{i,n}$. In particular, a cycle of order n has consecutive vertices and edges $u_1, e_1, u_2, e_2, \dots, u_n, e_n$. We first give a family of 2-regular graphs G with $\chi_{lt}(G) = 3$.

Theorem 3.3. For $m \geq 1$, $\chi_{lt}(mC_6) = 3$.

Proof. Since mC_6 has $6m$ vertices and $6m$ edges, we define a bijection $f : V(mC_6) \cup E(mC_6) \rightarrow [1, 12m]$ such that for $1 \leq i \leq m$,

- (i) $f(u_{i,1}) = 3i - 2$, $f(u_{i,3}) = 3i - 1$, $f(u_{i,5}) = 3i$,
- (ii) $f(u_{i,2}) = 12m + 3 - 3i$, $f(u_{i,4}) = 12m + 1 - 3i$, $f(u_{i,6}) = 12m + 2 - 3i$,
- (iii) $f(e_{i,1}) = 6m - 1 + 3i$, $f(e_{i,3}) = 6m + 3i$, $f(e_{i,5}) = 6m - 2 + 3i$,
- (iv) $f(e_{i,2}) = 6m + 1 - 3i$, $f(e_{i,4}) = 6m + 2 - 3i$, $f(e_{i,6}) = 6m + 3 - 3i$.

Clearly, the weights of $u_{i,1}$, $e_{i,1}$, $u_{i,2}$, $e_{i,2}$, $u_{i,3}$, $e_{i,3}$, $u_{i,4}$, $e_{i,4}$, $u_{i,5}$, $e_{i,5}$, $u_{i,6}$, $e_{i,6}$ are $12m + 2$, $12m + 1$, $12m$ repeatedly for $1 \leq i \leq m$. Thus, $\chi_{lt}(mC_6) \leq 3$. Since $\chi_{lt}(mC_6) \geq \chi_t(mC_6) = 3$, the theorem holds. \square

Example 3.4. Figure 1 gives the local total antimagic 3-coloring of $2C_6$ as defined above with induced weights 24, 25, 26.




Fig. 1. $2C_6$ with local total antimagic 3-coloring

Theorem 3.5. Suppose G is a 2-regular graph or else the disjoint union of a 2-regular graph and a path of order at least 3. If G has a C_n component, $n \neq 6$, then $\chi_{lt}(G) \geq 4$. Moreover, for $m \geq 1$, $\chi_{lt}(C_3) = \chi_{lt}(mC_4) = \chi_{lt}(C_5) = 4$, and $\chi_{lt}(C_8) \leq 5$.

Proof. By definition, $\chi_{lt}(G) \geq 3$. If $n \equiv 1, 2 \pmod{3}$, then $\chi_{lt}(G) \geq \chi_t(G) \geq 4$. Suppose $n \equiv 0 \pmod{3}$ and that $\chi_{lt}(G) = \chi_t(G) = 3$. Let f be a required local total antimagic 3-coloring of G . Suppose $n = 3$. Without loss of generality, we may assume the 3 distinct weights are a, b, c such that $a = w(u_1) = w(e_2)$, $b = w(e_1) = w(u_3)$ and $c = w(u_2) = w(e_3)$. Thus, we have

$$\begin{aligned} a &= f(e_1) + f(e_3) = f(u_2) + f(u_3) \\ b &= f(e_2) + f(e_3) = f(u_1) + f(u_2) \\ c &= f(e_1) + f(e_2) = f(u_1) + f(u_3). \end{aligned}$$

Now, $a + b - c$ gives $f(e_3) = f(u_2)$, a contradiction.

Suppose $n \geq 9$. Similarly, we may assume $a = w(u_j)$, $j \equiv 1 \pmod{3}$, $b = w(u_j)$, $j \equiv 0 \pmod{3}$, $c = w(u_j)$, $j \equiv 2 \pmod{3}$. Thus, we have

$$\begin{aligned} a &= f(e_3) + f(e_4) = f(e_6) + f(e_7), \\ b &= f(e_2) + f(e_3) = f(e_5) + f(e_6), \\ c &= f(e_1) + f(e_2) = f(e_4) + f(e_5). \end{aligned}$$

From $a-b+c$, we get $f(e_4)+f(e_1) = f(e_7)+f(e_4)$ so that $f(e_1) = f(e_7)$, a contradiction. Thus, $\chi_{lt}(G) \geq 4$ if G has a component of order $n \neq 6$.

Consider C_3 , label the vertices and edges $u_{1,1}, e_{1,1}, u_{1,2}, e_{1,2}, u_{1,3}, e_{1,3}$ by 1, 3, 5, 4, 6, 2 bijectively so that the corresponding weights are 5, 6, 7, 11, 6, 7. Thus, $\chi_{lt}(C_3) = 4$.

Consider mC_4 . Since mC_4 has $4m$ vertices and $4m$ edges, we define a bijection $f : V(mC_4) \cup E(mC_4) \rightarrow [1, 8m]$ such that for $1 \leq i \leq m$,

- (i) $f(u_{i,1}) = i, f(u_{i,2}) = 7m + 1 - i, f(u_{i,3}) = 3m + i, f(u_{i,4}) = 6m + 1 - i,$
- (ii) $f(e_{i,1}) = m + i, f(e_{i,2}) = 5m + 1 - i, f(e_{i,3}) = 2m + i, f(e_{i,4}) = 8m + 1 - i.$

Clearly, the weights of $u_{i,1}, e_{i,1}, u_{i,2}, e_{i,2}, u_{i,3}, e_{i,3}, u_{i,4}, e_{i,4}$ are $9m + 1, 7m + 1, 6m + 1, 10m + 1, 7m + 1, 9m + 1, 10m + 1, 6m + 1$ respectively. Thus, $\chi_{lt}(mC_4) = 4$.

For C_5 , label the vertices and edges $u_{1,1}, e_{1,1}, u_{1,2}, e_{1,2}, \dots, u_{1,5}, e_{1,5}$ by 1, 2, 7, 8, 5, 6, 3, 4, 9, 10 bijectively so that the corresponding weights are 12, 8, 10, 12, 14, 8, 10, 12, 14, 10. Thus, $\chi_{lt}(C_n) = 4$ for $n = 4, 5$. For C_8 , label the vertices and edges $u_{1,1}, e_{1,1}, u_{1,2}, e_{1,2}, \dots, u_{1,8}, e_{1,8}$ by 1, 10, 16, 5, 2, 11, 13, 6, 3, 12, 14, 7, 4, 9, 15, 8 bijectively so that the corresponding weights are 18, 17, 15, 18, 16, 15, 19, 16, 18, 17, 19, 18, 16, 19, 17, 16. Thus, $\chi_{lt}(C_8) \leq 5$. \square

Theorem 3.6. *For $m \geq 1$, $\chi_{lt}(G) = 3$ if and only if $G = P_n, n = 3, 6$ or $G = mC_6$ or $G = mC_6 + P_6, (m \geq 0)$.*

Proof. Necessity: Suppose $\chi_{lt}(G) = 3$. By Corollary 3.2, Theorem 3.3 and the proof of Theorem 3.5, we know $G = P_n, n = 3, 6$ or $G = mC_6, G = mC_6 + P_3$ or $G = mC_6 + P_6$ for $m \geq 1$. We shall prove that $\chi_{lt}(mC_6 + P_3) \neq 3$. Suppose equality holds and the P_3 component has vertices and edges v_1, h_1, v_2, h_2, v_3 consecutively. Let f be a local total antimagic 3-coloring of $mC_6 + P_3$ with induced weights a, b, c . Without loss of generality, we may assume the weights of $u_{i,1}, e_{i,1}, \dots, u_{i,6}, e_{i,6}$ of the i -th C_6 are a, b, c repeatedly, and the weights of v_1, h_1, v_2, h_2, v_3 are a, b, c, a, b respectively. Thus, we have

$$\begin{aligned} a &= f(e_{i,6}) + f(e_{i,1}) = f(h_1), \\ b &= f(e_{i,2}) + f(e_{i,3}) = f(h_2), \\ c &= f(e_{i,1}) + f(e_{i,2}) = f(h_1) + f(h_2). \end{aligned}$$

The right hand side above gives $a+b-c = 0$ so that $f(e_{i,6})+f(e_{i,3}) = 0$, a contradiction. Thus, $\chi_{lt}(mC_6 + P_3) \geq 4$. Therefore, $G = P_n, n = 3, 6$ or $G = mC_6$ or $G = mC_6 + P_6$ for $m \geq 1$.

Sufficiency: By Lemma 3.1 and Theorem 3.3, we only need to show that $\chi_{lt}(mC_6 + P_6) = 3$ for $m \geq 1$. Suppose the P_6 component has vertices and edges $v_1, h_1, \dots, v_5, h_5, v_6$ consecutively. Since G has $6m + 6$ vertices and $6m + 5$ edges, we define a bijection $f : V(G) \cup E(G) \rightarrow [1, 12m + 11]$ such that for $1 \leq i \leq m$,

- (i) $f(v_1) = 6m + 7, f(v_2) = 6m + 3, f(v_3) = 6m + 8, f(v_4) = 6m + 4, f(v_5) = 6m + 6,$
 $f(v_6) = 6m + 5,$
- (ii) $f(h_1) = 12m + 11, f(h_2) = 1, f(h_3) = 12m + 9, f(h_4) = 2, f(h_5) = 12m + 10,$

- (iii) $f(u_{i,1}) = 3i, f(u_{i,3}) = 3i + 1, f(u_{i,5}) = 3i + 2,$
- (iv) $f(u_{i,2}) = 12m + 11 - 3i, f(u_{i,4}) = 12m + 9 - 3i, f(u_{i,6}) = 12m + 10 - 3i,$
- (v) $f(e_{i,1}) = 6m + 7 + 3i, f(e_{i,3}) = 6m + 8 + 3i, f(e_{i,5}) = 6m + 6 + 3i,$
- (vi) $f(e_{i,2}) = 6m + 3 - 3i, f(e_{i,4}) = 6m + 4 - 3i, f(e_{i,6}) = 6m + 5 - 3i.$

Clearly, the weights of $u_{i,1}, e_{i,1}, \dots, u_{i,6}, e_{i,6}$ are $12m + 12, 12m + 11, 12m + 10$ repeatedly for $1 \leq i \leq m$ whereas the weights of $v_1, h_1, \dots, v_5, h_5, v_6$ are $12m + 11, 12m + 10, 12m + 12, 12m + 11, 12m + 10, 12m + 12, 12m + 11, 12m + 10,$ respectively. Thus, f is a local total antimagic labeling and $\chi_{lt}(G) = 3$. \square

Example 3.7. Figure 2 gives the local total antimagic 3-coloring of $2C_6 + P_6$ as defined above with induced weights 34, 35, 36.




Fig. 2. $2C_6 + P_6$ with local total antimagic 3-coloring

Corollary 3.8. Suppose $m \geq 1$ is odd and $s \geq \frac{m+1}{2}$. If $G_v(s)$ is obtained from mC_4 by attaching $s \geq 2$ pendant edges to a single vertex of mC_4 , then

$$\Delta(G_v(s)) + 1 = s + 3 \leq \chi_{lt}(G_v(s)) \leq s + 4 = \Delta(G_v(s)) + 2.$$

Proof. Let f be the local total antimagic labeling of mC_4 as defined in the proof of Theorem 3.5. Without loss of generality, assume the s edges are attached to $u_{1,1}$. Define a labeling g of $G_v(s)$ such that $g(x) = f(x)$ if $x \in V(G) \cup E(G)$, and label the vertices and edges of the s added pendant edges as in the proof of Theorem 2.4 for $k = 1$. Now, g is a local total antimagic labeling that induces weights $9m + 1, 7m + 1, 6m + 1, 10m + 1, 8m + 2i, 1 \leq i \leq s, \sum_{i=1}^s (8m + 2i) = 8ms + s(s + 1)$. Since $s \geq \frac{m+1}{2}$, $9m + 1 \in \{8m + 2i \mid 1 \leq i \leq s\}$, there are exactly $s + 4$ distinct weight so that $\chi_{lt}(G_v(s)) \leq s + 4$. Since $\Delta(G_v(s)) = s + 2$ and $G_v(s)$ has s pendant edges, by Corollary 2.2, $\chi_{lt}(G_v(s)) \geq s + 3$. This completes the proof. \square

4. χ_{lt} = NUMBER OF PENDANT EDGES + 1

We first note that applying Theorem 2.4 to the vertex of $mC_6, m \geq 1$ with label 1 as in Theorem 3.3, we get the following corollary.

Corollary 4.1. Suppose $m \geq 1$. If $G_v(s)$ is obtained from mC_6 by attaching $s \geq 2$ pendant edges to a single vertex of mC_6 , then $\chi_{lt}(G_v(s)) = \Delta(G_v(s)) + 1 = s + 3$.

In [15], the authors proved that $\chi_{lt}(nP_3) = 2n + 1$ for $n \geq 1$. We can now extend the obtained labeling to the following theorem.

Theorem 4.2. For $m \geq 1, n \geq 2$, $\chi_{lt}(mC_6 + P_3) = 4$ and $\chi_{lt}(mC_6 + nP_3) = 2n + 1$.

Proof. Consider $G = mC_6 + P_3$. From the proof of Theorem 3.6, we know that $\chi_{lt}(G) \geq 4$. Since G has $6m + 3$ vertices and $6m + 2$ edges, we define a bijection $f : V(G) \cup E(G) \rightarrow [1, 12m + 5]$ such that for $1 \leq i \leq m$,

- (i) $f(v_1) = 1, f(h_1) = 12m + 5, f(v_2) = 12m + 3, f(h_2) = 12m + 4, f(v_3) = 2,$
- (ii) $f(u_{i,1}) = 3i, f(u_{i,3}) = 3i + 1, f(u_{i,5}) = 3i + 2,$
- (iii) $f(u_{i,2}) = 12m + 5 - 3i, f(u_{i,4}) = 12m + 3 - 3i, f(u_{i,6}) = 12m + 4 - 3i,$
- (iv) $f(e_{i,1}) = 6m + 1 + 3i, f(e_{i,3}) = 6m + 2 + 3i, f(e_{i,5}) = 6m + 3i,$
- (v) $f(e_{i,2}) = 6m + 3 - 3i, f(e_{i,4}) = 6m + 4 - 3i, f(e_{i,6}) = 6m + 5 - 3i.$

Clearly, the weights of $u_{i,1}, e_{i,1}, u_{i,2}, e_{i,2}, u_{i,3}, e_{i,3}, u_{i,4}, e_{i,4}, u_{i,5}, e_{i,5}, u_{i,6}, e_{i,6}$ are $12m + 6, 12m + 5, 12m + 4$ repeatedly for $1 \leq i \leq m$ whereas the weights of v_1, h_1, v_2, h_2, v_3 are $12m + 5, 12m + 4, 24m + 9, 12m + 5, 12m + 4$ respectively. Thus, $\chi_{lt}(G) \leq 4$. Consequently, $\chi_{lt}(G) = 4$.

Consider $n \geq 2$. Now, $G = mC_6 + nP_3$ has $2n$ pendant vertices and maximum degree 3. By Corollary 2.2, we have $\chi_{lt}(G) \geq 2n + 1$. Suppose the nP_3 has vertex set $\{v_{j,1}, v_{j,2}, v_{j,3}\}$ and edge set $\{v_{j,1}v_{j,2}, v_{j,2}v_{j,3}\}$ for $1 \leq j \leq n$. Define a total labeling $f : V(G) \cup E(G) \rightarrow [1, 12m + 5n]$ such that for $1 \leq i \leq m, 1 \leq j \leq n$,

- (a) $f(v_{j,1}) = j, f(v_{j,1}v_{j,2}) = 5n + 12m + 1 - j, f(v_{j,2}) = 3n + 12m + 1 - j,$
 $f(v_{j,2}v_{j,3}) = 3n + 12m + j, f(v_{j,3}) = n + j,$
- (b) $f(u_{i,1}) = 2n + 3i - 2, f(u_{i,3}) = 2n + 3i - 1, f(u_{i,5}) = 2n + 3i,$
- (c) $f(u_{i,2}) = 2n + 12m + 3 - 3i, f(u_{i,4}) = 2n + 12m + 1 - 3i, f(u_{i,6}) = 2n + 12m + 2 - 3i,$
- (d) $f(e_{i,1}) = 2n + 6m - 1 + 3i, f(e_{i,3}) = 2n + 6m + 3i, f(e_{i,5}) = 2n + 6m - 2 + 3i,$
- (e) $f(e_{i,2}) = 2n + 6m + 1 - 3i, f(e_{i,4}) = 2n + 6m + 2 - 3i, f(e_{i,6}) = 2n + 6m + 3 - 3i.$

We now have $w(v_{j,1}) = 5n + 12m + 1 - j, w(v_{j,3}) = 3n + 12m + j$ for $1 \leq j \leq n$, $w(v_{j,2}) = 8n + 24m + 1, w(v_{j,1}v_{j,2}) = 3n + 12m + 1, w(v_{j,2}v_{j,3}) = 4n + 12m + 1$ so that the nP_3 components have $2n + 1$ distinct weights. Moreover, the weights of $u_{i,1}, e_{i,1}, u_{i,2}, e_{i,2}, u_{i,3}, e_{i,3}, u_{i,4}, e_{i,4}, u_{i,5}, e_{i,5}, u_{i,6}, e_{i,6}$ are $4n + 12m + 2, 4n + 12m + 1, 4n + 12m$ (that are the $w(v_{n-1,1}), w(v_{n,1}), w(v_{n,3})$) repeatedly for $1 \leq i \leq m$. Thus, f is a local total antimagic labeling that induces $2n + 1$ distinct weights so that $\chi_{lt}(G) \leq 2n + 1$. Consequently, $\chi_{lt}(G) = 2n + 1$. \square

Example 4.3. Figure 3 gives the local total antimagic 5-coloring of $2C_6 + 2P_3$ with induced weights 31, 32, 33, 34, 65 as defined above.




Fig. 3. $2C_6 + 2P_3$ with local total antimagic 5-coloring

In [14], the authors also proved that

$$\chi_{lt}(mP_6) = 2m + 1$$

for $m \geq 2$, and that

$$\chi_{lt}(mP_6 + nP_3) = 2m + 2n + 1$$

for $m \geq 1, n \geq 2$. We can now prove the following theorems.

Theorem 4.4. For $m, n \geq 1$, $\chi_{lt}(mC_6 + nP_6) = 2n + 1$.

Proof. Let $G = mC_6 + nP_6$. By Theorem 3.6, we only need to consider $m \geq 1, n \geq 2$. Since G has $2n$ pendant edges and $\Delta(G) = 2$, by Corollary 2.2, $\chi_{lt}(G) \geq 2n + 1$. We shall show that $\chi_{lt}(G) \leq 2n + 1$. Keep the notations for the mC_6 , $m \geq 1$. For $1 \leq t \leq n$, let the vertex and edge sets of the t -th component of the nP_6 be $\{y_{t,a} \mid 1 \leq a \leq 6\}$ and $\{z_{t,a} = y_{t,a}y_{t,a+1} \mid 1 \leq a \leq 5\}$ respectively. Since G has order $6m + 6n$ and size $6m + 5n$, we define a bijection $f : V(G) \cup E(G) \rightarrow [1, 12m + 11n]$ such that for $1 \leq i \leq m, 1 \leq t \leq n$,

- (a) $f(u_{i,1}) = 2n + 3i - 2, f(u_{i,3}) = 2n + 3i - 1, f(u_{i,5}) = 2n + 3i,$
- (b) $f(u_{i,2}) = 8n + 12m - 3i + 3, f(u_{i,4}) = 8n + 12m - 3i + 1, f(u_{i,6}) = 8n + 12m - 3i + 2,$
- (c) $f(e_{i,1}) = 8n + 6m + 3i - 1, f(e_{i,3}) = 8n + 6m + 3i, f(e_{i,5}) = 8n + 6m + 3i + 1,$
- (d) $f(e_{i,2}) = 2n + 6m - 3i + 1, f(e_{i,4}) = 2n + 6m - 3i + 2, f(e_{i,6}) = 2n + 6m - 3i + 3,$
- (e) $f(y_{t,1}) = 8n + 6m - 3t + 2, f(y_{t,3}) = 8n + 6m - 3t + 3, f(y_{t,5}) = 8n + 6m - 3t + 1,$
- (f) $f(y_{t,2}) = 2n + 6m + 3t - 2, f(y_{t,4}) = 2n + 6m + 3t - 1, f(y_{t,6}) = 2n + 6m + 3t,$
- (g) $f(z_{t,1}) = 11n + 12m + 1 - t, f(z_{t,3}) = 9n + 12m + 1 - t, f(z_{t,5}) = 10n + 12m + 1 - t,$
- (h) $f(z_{t,2}) = t, f(z_{t,4}) = n + t.$

One can check that the induced weights of each component of the mC_6 , starting from $w(u_{i,1})$ and $w(e_{i,1})$, are $10n + 12m + 2, 10n + 12m + 1, 10n + 12m$ repeatedly and the induced non-pendant vertex weights of each component of the nP_6 , starting from $w(z_{t,1})$, are $10n + 12m, 11n + 12m + 1, 10n + 12m + 1, 9n + 12m + 1, 10n + 12m + 2, 10n + 12m + 1, 10n + 12m, 11n + 12m + 1, 10n + 12m + 1$ consecutively, while $w(y_{t,1}) = 11n + 12m + 1 - t$ and $w(y_{t,6}) = 10n + 12m + 1 - t$ for $1 \leq t \leq n$. Thus, f is a local total antimagic labeling that induces $2n + 1$ distinct weights so that $\chi_{lt}(G) \leq 2n + 1$. This completes the proof. \square

Example 4.5. Figure 4 gives the local total antimagic 5-coloring of $2C_6 + 2P_6$ as defined above with induced weights 43, 44, 45, 46, 47.




Fig. 4. $2C_6 + 2P_6$ with local total antimagic 5-coloring

Theorem 4.6. For $m, n \geq 1, a \geq 2$, if $n \geq 2a + 2$ or else $a \geq 2n$, then

$$\chi_{lt}(mC_6 + nP_6 + aP_3) = 2n + 2a + 1.$$

Otherwise,

(i) if $n = 2a + 1$ or else $s \in [n + 1, 2n - 1]$, then

$$2n + 2a + 1 \leq \chi_{lt}(mC_6 + nP_6 + aP_3) \leq 2n + 2a + 2.$$

(ii) if $n = 2a$ or else $a = n$, then

$$2n + 2a + 1 \leq \chi_{lt}(mC_6 + nP_6 + aP_3) \leq 2n + 2a + 3.$$

(iii) if $n \in [a + 1, 2a - 1]$ or else $a = n - 1$, then

$$2n + 2a + 1 \leq \chi_{lt}(mC_6 + nP_6 + aP_3) \leq 2n + 2a + 4.$$

(iv) if $n \leq a$ or else $a \leq n - 2$, then

$$2n + 2a + 1 \leq \chi_{lt}(mC_6 + nP_6 + aP_3) \leq 2n + 2a + 5.$$

Proof. Let $G = mC_6 + nP_6 + aP_3$ for $m, n \geq 1, a \geq 2$. Since G has $2n + 2a$ pendant edges and $\Delta(G) = 2$, by Corollary 2.2, $\chi_{lt}(G) \geq 2n + 2a + 1$. We shall show that $\chi_{lt}(G)$ has the given upper bound under the given conditions. Keep the notations of $mC_6 + nP_6$ as defined in the proof of Theorem 4.4. Let f be the total labeling as defined in the proof of Theorem 4.4. For $1 \leq r \leq a$, let the vertex and edge sets of the r -th component of the aP_3 be $\{v_{r,b} \mid 1 \leq b \leq 3\}$ and $\{h_{r,b} = v_{r,b}v_{r,b+1} \mid 1 \leq b \leq 2\}$ respectively.

Since G has order $6m + 6n + 3a$ and size $6m + 5n + 2a$, we define a bijection $g : V(G) \cup E(G) \rightarrow [1, 12m + 11n + 5a]$ such that for $1 \leq i \leq m, 1 \leq t \leq n, 1 \leq r \leq a$, $g(x) = f(x) + 2a$ for $x \in V(mC_6 + nP_6) \cup E(mC_6 + nP_6)$. Moreover,

$$\begin{aligned} g(v_{r,1}) &= r, \\ g(h_{r,1}) &= 12m + 11n + 5a + 1 - r, \\ g(v_{r,2}) &= 12m + 11n + 3a + 1 - r, \\ g(h_{r,2}) &= 12m + 11n + 3a + r, \\ g(v_{r,3}) &= a + r. \end{aligned}$$

Similar to the proof of Theorem 4.4, one can check that the induced weights of each component of the mC_6 are $10n + 12m + 4a + 2$, $10n + 12m + 4a + 1$, $10n + 12m + 4a$ repeatedly, the induced non-pendant vertex weights of each component of the nP_6 are $10n + 12m + 4a$, $11n + 12m + 4a + 1$, $10n + 12m + 4a + 1$, $9n + 12m + 4a + 1$,

$10n + 12m + 4a + 2$, $10n + 12m + 4a + 1$, $10n + 12m + 4a$, $11n + 12m + 4a + 1$, $10n + 12m + 4a + 1$, while

$$\begin{aligned} w_g(y_{t,1}) &= 11n + 12m + 2a + 1 - t, \\ w_g(y_{t,5}) &= 10n + 12m + 2a + 1 - t, \\ w_g(v_{r,1}) &= 11n + 12m + 5a + 1 - r, \\ w_g(h_{r,1}) &= 11n + 12m + 3a + 1, \\ w_g(v_{r,2}) &= 22n + 24m + 8a + 1, \\ w_g(h_{r,2}) &= 11n + 12m + 4a + 1, \\ w_g(v_{r,3}) &= 11n + 12m + 3a + r \end{aligned}$$

for $1 \leq t \leq n$, $1 \leq r \leq a$. Clearly, $w_g(v_{r,2}) = 22n + 24m + 8a + 1$ is not in the pendant vertex weights set

$$W = [9n + 12m + 2a + 1, 11n + 12m + 2a] \cup [11n + 12m + 3a + 1, 11n + 12m + 5a].$$

Note that this labeling is local total antimagic. Moreover, there are exactly $2n + 2a + 1$ distinct weights if and only if all the induced weights set of the non-pendant vertices of $mC_6 + nP_6$ components, namely $\{9n + 12m + 4a + 1, 10n + 12m + 4a, 10n + 12m + 4a + 1, 10n + 12m + 4a + 2, 11n + 12m + 4a + 1\}$ is a subset of W . This implies that

- (a) $2n \geq 2a + 1$, or else $a \geq 2n$ for $9n + 12m + 4a + 1 \in W$,
- (b) $n \geq 2a$, or else $a \geq n + 1$ for $10n + 12m + 4a \in W$,
- (c) $n \geq 2a + 1$, or else $a \geq n$ for $10n + 12m + 4a + 1 \in W$,
- (d) $n \geq 2a + 2$, or else $a \geq n - 1$ for $10n + 12m + 4a + 2 \in W$.

Therefore, if $n \geq 2a + 2$ or else $a \geq 2n$, then

$$\chi_{lt}(mC_6 + nP_6 + aP_3) \leq 2n + 2a + 1.$$

Otherwise,

- (i) if $n = 2a + 1$ or else $a \in [n + 1, 2n - 1]$, then there are $2n + 2a + 2$ distinct weights so that

$$2n + 2a + 1 \leq \chi_{lt}(mC_6 + nP_6 + aP_3) \leq 2n + 2a + 2.$$

- (ii) if $n = 2a$ or else $a = n$, then there are $2n + 2a + 3$ distinct weights so that

$$2n + 2a + 1 \leq \chi_{lt}(mC_6 + nP_6 + sP_3) \leq 2n + 2a + 3.$$

- (iii) if $n \in [a + 1, 2a - 1]$ or else $a = n - 1$, then there are $2n + 2a + 4$ distinct weights so that

$$2n + 2a + 1 \leq \chi_{lt}(mC_6 + nP_6 + aP_3) \leq 2n + 2s + 4.$$

- (iv) if $n \leq a$ or else $a \leq n - 2$, then there are $2n + 2a + 5$ distinct weights so that

$$2n + 2a + 1 \leq \chi_{lt}(mC_6 + nP_6 + aP_3) \leq 2n + 2a + 5.$$

This completes the proof. \square

Example 4.7. Figures 5 and 6 give the local total antimagic 11-coloring of $C_6 + 4P_6 + P_3$ with $n = 4$, $a = 1$ and induced weights $51, 52, \dots, 58, 60, 61, 121$, as well as local total antimagic 9-coloring of $C_6 + P_6 + 3P_3$ with $n = 1$, $a = 3$ and induced weights $28, 29, 33, 34, \dots, 38, 71$ as defined above.




Fig. 5. $C_6 + 4P_6 + P_3$ with local total antimagic 13-coloring




Fig. 6. $C_6 + P_6 + 3P_3$ with local total antimagic 9-coloring

Theorem 4.8. For $m \geq 1, n \geq 2$, let $G = mC_6 + nP_3$ with a local total antimagic labeling as defined in the proof of Theorem 4.2. Suppose $s \geq 1$, $1 \leq j' \leq s$, $1 \leq i' \leq k$, $1 \leq i \leq m$, $1 \leq j \leq n$, $ks \geq 2$ and the following ten conditions:

- (a) $v = v_{1,1}$, $k = 1$, n is odd and $s \geq (3n + 24m + 1)/2$,
- (b) $v = v_{j,1}$ for $j \in [2, n]$, $k = j$ and $2(j' - 1)j + 1 \leq 3n + 12m + 1 \leq 2j'j$,
- (c) $v = v_{j,3}$ for $j \in [1, n]$, $k = n + j$ and

$$(2j' - 1)(n + j) + 1 \leq 3n + 12m + 1 \leq 2j'(n + j),$$

- (d) $v = v_{j,2}$, $k = 3n + 12m + 1 - j$, $s \geq 1$,
- (e) $v = u_{i,1}$, $k = 2n + 3i - 2$, and

$$8n + 24m + 1 \in \bigcup_{j'=1}^s [5n + 12m + (2j' - 1)(2n + 3i - 2) + 1, 5n + 12m + 2j'(2n + 3i - 2)],$$

- (f) $v = u_{i,3}$, $k = 2n + 3i - 1$, and

$$8n + 24m + 1 \in \bigcup_{j'=1}^s [5n + 12m + (2j' - 1)(2n + 3i - 1) + 1, 5n + 12m + 2j'(2n + 3i - 1)],$$

- (g) $v = u_{i,5}$, $k = 2n + 3i$, and

$$8n + 24m + 1 \in \bigcup_{j'=1}^s [5n + 12m + (2j' - 1)(2n + 3i) + 1, 5n + 12m + 2j'(2n + 3i)],$$

- (h) $v = u_{i,2}$, $k = 2n + 12m + 3 - 3i$, $s \geq 1$,
- (i) $v = u_{i,4}$, $k = 2n + 12m + 1 - 3i$, $s \geq 1$,
- (j) $v = u_{i,6}$, $k = 2n + 12m + 2 - 3i$, $s \geq 1$.

Each of Conditions (a) to (c) implies that $\chi_{lt}(G_v(k, s)) = ks + 2n$, whereas each of Conditions (d) to (j) implies that $\chi_{lt}(G_v(k, s)) = ks + 2n + 1$.

Proof. Let f be the local total antimagic labeling of G as defined in the proof of Theorem 4.2. Clearly, if v is a pendant vertex, then $v \in \{v_{j,1}, v_{j,3} \mid 1 \leq j \leq n\}$ and $f(v) = k = j$ for $v = v_{j,1}$ and $k = n + j$ for $v = v_{j,3}$. Moreover, $\Delta(G_v(k, s)) = ks + 1$ and $G_v(k, s)$ has $ks + 2n - 1$ pendant edges. By Corollary 2.2, $\chi_{lt}(G_v(k, s)) \geq ks + 2n$. If v is not a pendant vertex, then

$$v \in \{u_{i,1}, \dots, u_{i,6}, v_{j,2} \mid 1 \leq i \leq m, 1 \leq j \leq n\}$$

and

$$f(v) = k \in \{2n + 3i - 2, 2n + 3i - 1, 2n + 3i, 2n + 12m + 3 - 3i, 2n + 12m + 2 - 3i, 2n + 12m + 1 - 3i, 3n + 12m + 1 - j \mid 1 \leq i \leq m, 1 \leq j \leq n\}.$$

Moreover, $\Delta(G_v(k, s)) = ks + 2$ and $G_v(k, s)$ has $ks + 2n$ pendant edges. Suppose the added ks pendant edges incident to v are $e_{i',j'}$ and the corresponding pendant vertices are $x_{i',j'}$ for $1 \leq j' \leq s$, $1 \leq i' \leq k$. By Corollary 2.2, $\chi_{lt}(G_v(k, s)) \geq ks + 2n + 1$.

Since $G_v(k, s)$ has $3n + 6m + ks$ vertices and $2n + 6m + ks$ edges, we define a total labeling $g : V(G_v(k, s)) \cup E(G_v(k, s)) \rightarrow [1, 5n + 12m + 2ks]$ such that $g(z) = f(z)$ for $z \in V(G) \cup E(G)$. Otherwise, $g(z)$ is as defined in the proof of Theorems 2.4 or 2.5. One can check that

$$w_g(v) = w_f(v) + \sum_{j'=1}^s \sum_{i'=1}^k (12m + 5n + (2j' - 1)k + i'),$$

and

$$w(z) \in [3n + 12m + 1, 5n + 12m] \cup \{8n + 24m + 1\} \cup \{5n + 12m + (2j' - 1)k + i' \mid 1 \leq j' \leq s, 1 \leq i' \leq k\}$$

for $z \in (V(G_v(k, s)) \cup E(G_v(k, s))) \setminus \{v\}$. Moreover, g is a local total antimagic labeling. Thus, $\chi_{lt}(G_v(k, s)) \leq ks + 2n + 2$ if all the weights are distinct. We shall need to check the conditions under Theorem 2.5 in the following 10 cases.

- (a) Suppose $v = v_{1,1}$ so that $k = 1$. Thus,

$$\{w_g(e_{j',1})\} = \{5n + 12m + 2j' \mid 1 \leq j' \leq s, s \geq 2\}.$$

Therefore, $8n + 24m + 1 \in \{w_g(e_{j',1})\}$ so that n is odd and

$$s \geq (3n + 24m + 1)/2.$$

(b) Suppose $v = v_{j,1}$, $2 \leq j \leq n$ so that $k = j \geq 2$. Therefore,

$$\begin{aligned} 8n + 24m + 1 &\in \{w_g(e_{j',i'})\} \\ &= \bigcup_{j'=1}^s [5n + 12m + (2j' - 1)j + 1, 5n + 12m + 2j'j], \end{aligned}$$

$2 \leq j \leq n$, so that

$$2(j' - 1)j + 1 \leq 3n + 12m + 1 \leq 2j'j.$$

As an example, take $n = 3, m = 1$, we can choose $j = 3, s \geq j' = 4$ to get

$$8n + 24m + 1 = 49 \in [49, 51] = [5n + 12m + (2j' - 1)j + 1, 5n + 12m + 2j'j]$$

as required.

(c) Suppose $v = v_{j,3}$, $1 \leq j \leq n$ so that $k = n + j$. Therefore,

$$\begin{aligned} 8n + 24m + 1 &\in \{w_g(e_{j',i'})\} \\ &= \bigcup_{j'=1}^s [5n + 12m + (2j' - 1)(n + j) + 1, 5n + 12m + 2j'(n + j)], \end{aligned}$$

$1 \leq j \leq n$, so that

$$(2j' - 1)(n + j) + 1 \leq 3n + 12m + 1 \leq 2j'(n + j).$$

As an example, take $n = m = 3$, we can choose $j = 1, s \geq j' = 6$ to get

$$\begin{aligned} 8n + 24m + 1 &= 97 \in [96, 99] \\ &= [5n + 12m + (2j' - 1)(n + j) + 1, 5n + 12m + 2j'(n + j)] \end{aligned}$$

as required.

(d) Suppose $v = v_{j,2}$ so that $k = 3n + 12m + 1 - j$. Thus,

$$\begin{aligned} &\{w_g(e_{j',i'})\} \\ &= \{5n + 12m + (2j' - 1)(3n + 12m + 1 - j) + i' \mid 1 \leq i' \leq k, 1 \leq j' \leq s\} \\ &= \bigcup_{j'=1}^s [5n + 12m + (2j' - 1)(3n + 12m + 1 - j) + 1, \\ &\quad 5n + 12m + 2j'(3n + 12m + 1 - j)]. \end{aligned}$$

If $j' = 1$, then

$$8n + 24m + 1 \in [8n + 24m + 2 - j, 11n + 36m + 2 - 2j]$$

since $j \leq n$. Thus, g induces $ks + 2n + 1$ distinct weights. Therefore, if $G = mC_6 + nP_3$, then

$$\chi_{lt}(G_{v_{j,2}}(3n + 12m + 1 - j, s)) = (3n + 12m + 1 - j)s + 2n + 1$$

for $s \geq 1$.

(e) Suppose $v = u_{i,1}$ so that $k = 2n + 3i - 2$. Thus,

$$\begin{aligned} \{w_g(e_{j',i'})\} &= \{5n + 12m + (2j' - 1)k + i' \mid 1 \leq i' \leq k, 1 \leq j' \leq s\} \\ &= \bigcup_{j'=1}^s [5n + 12m + (2j' - 1)(2n + 3i - 2) + 1, 5n + 12m + 2j'(2n + 3i - 2)], \end{aligned}$$

denoted U , for $1 \leq i \leq m, 1 \leq j' \leq s$. Thus, if $8n + 24m + 1 \in U$, then g induces $ks + 2n + 1$ distinct weights. As an example, take $m = 1, n = 3, i = 1, k = 7, j' = 2$, we get $8n + 24m + 1 = 49$ and

$$[5n + 12m + (2j' - 1)(2n + 3i - 2) + 1, 5n + 12m + 2j'(2n + 3i - 2)] = [49, 55].$$

Thus, for $G = C_6 + 3P_3$,

$$\chi_{lt}(G_{u_{1,1}}(7, s)) = 7s + 7$$

for $s \geq 2$.

(f) Suppose $v = u_{i,3}$ so that $k = 2n + 3i - 1$. By a similar argument, we get g induces $ks + 2n + 1$ distinct weights if

$$\begin{aligned} 8n + 24m + 1 &\in \\ &\bigcup_{j'=1}^s [5n + 12m + (2j' - 1)(2n + 3i - 1) + 1, 5n + 12m + 2j'(2n + 3i - 1)] \end{aligned}$$

for $1 \leq i \leq m, 1 \leq j' \leq s$. As an example: take $m = 2, n = 3, i = 2, k = 11, j' = 2$, we get $8n + 24m + 1 = 73$ and

$$[5n + 12m + (2j' - 1)(2n + 3i - 1) + 1, 5n + 12m + 2j'(2n + 3i - 1)] = [73, 83].$$

Thus, for $G = 2C_6 + 3P_3$,

$$\chi_{lt}(G_{u_{2,3}}(11, s)) = 11s + 7$$

for $s \geq 2$.

(g) Suppose $v = u_{i,5}$ so that $k = 2n + 3i$. By a similar argument, we get g induces $ks + 2n + 1$ distinct weights if

$$8n + 24m + 1 \in \bigcup_{j'=1}^s [5n + 12m + (2j' - 1)(2n + 3i) + 1, 5n + 12m + 2j'(2n + 3i)].$$

As an example: take $m = n = i = 3, k = 15, j' = 2$, we get $8n + 24m + 1 = 97$ and

$$[5n + 12m + (2j' - 1)(2n + 3i) + 1, 5n + 12m + 2j'(2n + 3i)] = [97, 111].$$

Thus, for $G = 3C_6 + 3P_3$,

$$\chi_{lt}(G_{u_{3,5}}(15, s)) = 15s + 7$$

for $s \geq 2$.

- (h) Suppose $v = u_{i,2}$ so that $k = 2n + 12m + 3 - 3i$. We get g induces $ks + 2n + 1$ distinct weights if

$$8n + 24m + 1 \in \bigcup_{j'=1}^s [5n + 12m + (2j' - 1)(2n + 12m + 3 - 3i) + 1, \\ 5n + 12m + 2j'(2n + 12m + 3 - 3i)].$$

For $j' = 1$, we get

$$8n + 24m + 1 \in [7n + 12m + 4 - 3i, 9n + 36m + 6 - 6i]$$

since $i \leq m$. Thus, for $G = mC_6 + nP_3$,

$$\chi_{lt}(G_{u_{i,2}}(2n + 12m + 3 - 3i, s)) = (2n + 12m + 3 - 3i)s + 2n + 1$$

for $s \geq 1$.

- (i) Suppose $v = u_{i,4}$ so that $k = 2n + 12m + 1 - 3i$. We get g induces $ks + 2n + 1$ distinct weights if

$$8n + 24m + 1 \in \bigcup_{j'=1}^s [5n + 12m + (2j' - 1)(2n + 12m + 1 - 3i) + 1, \\ 5n + 12m + 2j'(2n + 12m + 1 - 3i)].$$

For $j' = 1$, we get

$$8n + 24m + 1 \in [7n + 24m + 2 - 3i, 9n + 36m + 3 - 6i]$$

since $i \leq m$. Thus, for $G = mC_6 + nP_3$,

$$\chi_{lt}(G_{u_{i,4}}(2n + 12m + 1 - 3i, s)) = (2n + 12m + 1 - 3i)s + 2n + 1$$

for $s \geq 1$.

- (j) Suppose $v = u_{i,6}$ so that $k = 2n + 12m + 2 - 3i$. We get g induces $ks + 2n + 1$ distinct weights if

$$8n + 24m + 1 \in \bigcup_{j'=1}^s [5n + 12m + (2j' - 1)(2n + 12m + 2 - 3i) + 1, \\ 5n + 12m + 2j'(2n + 12m + 2 - 3i)].$$

For $j' = 1$, we get

$$8n + 24m + 1 \in [7n + 24m + 3 - 3i, 9n + 36m + 4 - 6i]$$

since $i \leq m$. Thus, for $G = mC_6 + nP_3$,

$$\chi_{lt}(G_{u_{i,6}}(2n + 12m + 2 - 3i, s)) = (2n + 12m + 2 - 3i)s + 2n + 1$$

for $s \geq 1$.

For (a) to (c), Theorem 2.5(a)(i) implies that $\chi_{lt}(G_v(k, s)) = ks + 2n$. For (d) to (j), Theorem 2.5(b) implies that $\chi_{lt}(G_v(k, s)) = ks + 2n + 1$. This completes the proof. \square

Example 4.9. Figure 7 gives the local total antimagic 65-coloring of $G_v(k, s)$ for $G = 2C_6 + 2P_3$, $v = v_{1,2}$, $k = 30$, $s = 2$ with induced weights 31, 32, 33, 34, 65, 66, ..., 94, 125, 126, ..., 154 and 6635 being the weight of $v_{1,2}$ in $G_v(k, s)$ as defined above.




Fig. 7. $G_v(k, s)$, $v = v_{1,2}$, $k = 30$, $s = 2$ with local total antimagic 65-coloring

Note that Theorem 2.5 is not applicable to $mC_6 + nP_6$ for $m, n \geq 1$. Let $G = mC_6 + nP_6 + aP_3$ with vertex and edge sets as defined in the proof of Theorem 4.6. We now have the following theorem.

Theorem 4.10. For $m, n \geq 1$, $a \geq 2$ and $G = mC_6 + nP_6 + aP_3$ such that $n \geq 2a + 2$ or else $a \geq 2n$ with the following 15 conditions:

- (1) $v = y_{t,1}$, $k = 8n + 6m - 3t + 2a + 2$, $t \leq (5n + 3)/6$,
- (2) $v = y_{t,6}$, $k = 2n + 6m + 3t + 2a$ and
 - (a) $s \geq 1$, $t \in [1, n]$, $a \geq 2n$ only, or else
 - (b) $s \geq 2$, $(3n - 12m - 5a + 1)/12 \leq t \leq (5n - 6m - 3a + 1)/9$, or else
 - (c) $s \geq 3$, $t \leq (n - 18m - 7a + 1)/15$.
- (3) $v = v_{r,1}$, $r \in [1, a]$, $k = r$, $(2j' - 1)r \leq 11n + 12m + 3a + 1 \leq 2j'r$,
- (4) $v = v_{r,3}$, $r \in [1, a]$, $k = a + r$,

$$(2j' - 1)(a + r) \leq 11n + 12m + 3a + 1 \leq 2j'(a + r),$$

- (5) $v = u_{i,1}$, $i \in [1, m]$, $k = 2n + 2a + 3i - 2$,

$$(2j' - 1)(2n + 2a + 3i - 2) \leq 11n + 12m + 3a + 1 \leq 2j'(2n + 2a + 3i - 2),$$

- (6) $v = u_{i,3}$, $i \in [1, m]$, $k = 2n + 2a + 3i - 1$,

$$(2j' - 1)(2n + 2a + 3i - 1) \leq 11n + 12m + 3a + 1 \leq 2j'(2n + 2a + 3i - 1),$$

- (7) $v = u_{i,5}$, $i \in [1, m]$, $k = 2n + 2a + 3i$,

$$(2j' - 1)(2n + 2a + 3i) \leq 11n + 12m + 3a + 1 \leq 2j'(2n + 2a + 3i),$$

- (8) $v = u_{i,2}$, $i \in [1, m]$, $k = 8n + 12m + 2a - 3i + 3$, $s \geq 1$,
- (9) $v = u_{i,4}$, $i \in [1, m]$, $k = 8n + 12m + 2a - 3i + 1$, $s \geq 1$,
- (10) $v = u_{i,6}$, $i \in [1, m]$, $k = 8n + 12m + 2a - 3i + 2$, $s \geq 1$,

- (11) $v = y_{t,3}, k = 8n + 6m + 2a - 3t + 3, t \leq (5n + a + 5)/6, n \geq a + 5$ for $a = 1, 2$ and $n \geq 2a + 2$ otherwise,
 (12) $v = y_{t,5}, k = 8n + 6m + 2a - 3t + 1, t \leq (5n + a + 1)/6$ and $n \geq 2a + 2$ only,
 (13) $v = y_{t,2}, k = 2n + 6m + 2a + 3t - 2$ and
 (a) $s \geq 1, t \in [1, n], a \geq n + 5$ for $a = 1, 2, 3, 4$ and $a \geq 2n$ otherwise, or else
 (b) $s \geq 2, (3n - 12m - 5a + 9)/12 \leq t \leq (5n - 6m - a + 7)/3$, or else
 (c) $s \geq 3, t \leq (n - 18m - 7a + 13)/15$,
 (14) $v = y_{t,4}, k = 2n + 6m + 2a + 3t - 1$, and
 (a) $s \geq 1, t \in [1, n], a \geq n + 3$ for $a = 1, 2$ and $a \geq 2n$ otherwise, or else
 (b) $s \geq 2, (3n - 12m - 5a + 5)/12 \leq t \leq (5n - 6m - 3a + 4)/9$, or else
 (c) $s \geq 3, t \leq (n - 18m - 7a + 6)/15$.
 (15) $v = v_{r,2}, r \in [1, a], k = 12m + 11n + 3a + 1 - r, s \geq 1$.

Each of Conditions (1) to (4) implies that $\chi_{lt}(G_v(k, s)) = 2n + 2a + ks$, whereas each of Conditions (5) to (15) implies that $\chi_{lt}(G_v(k, s)) = 2n + 2a + ks + 1$.

Proof. Let g be the local total antimagic labeling of G as defined in the proof of Theorem 4.6. By an argument similar to the proof of Theorem 4.8 and Theorem 2.5(iii), we shall show that the only weight $w_g(v_{r,2}) = 22n + 24m + 8a + 1$ which is not a pendant edge label under g must be in

$$\{w_g(e_{j',i'})\} = \bigcup_{j'=1}^s [11n + 12m + 5a + (2j' - 1)k + 1, 11n + 12m + 5a + 2j'k].$$

We also check the conditions under Theorem 2.5. We have the following first four cases for v being a pendant vertex and next 11 cases for v not a pendant vertex.

- (1) Suppose $v = y_{t,1}, t \in [1, n]$ with $k = 8n + 6m - 3t + 2a + 2$. Therefore, $22n + 24m + 8a + 1 \in \{w_g(e_{j',i'})\}$ means

$$(2j' - 1)(8n + 6m - 3t + 2a + 2) + 1 \leq 11n + 12m + 3a + 1 \leq 2j'(8n + 6m - 3t + 2a + 2).$$

Thus, $j' = 1$ so that

$$8n + 6m - 3t + 2a + 3 \leq 11n + 12m + 3a + 1 \leq 16n + 12m - 6t + 4a + 4.$$

Consequently, $t \leq (5n + 3)/6$.

- (2) Suppose $v = y_{t,6}, t \in [1, n]$ with $k = 2n + 6m + 3t + 2a$. Therefore, we must have

$$(2j' - 1)(2n + 6m + 3t + 2a) \leq 11n + 12m + 3a + 1 \leq 2j'(2n + 6m + 3t + 2a).$$

Thus, $j' \leq 3$. If $j' = 1$, then

$$2n + 6m + 3t + 2a \leq 11n + 12m + 3a + 1 \leq 4n + 12m + 6t + 4a$$

so that

$$(7n - a + 1)/6 \leq t \leq (9n + 6m + a + 1)/3.$$

Since $t \leq n$, we then have $a \geq n + 1$. The conditions $n \geq 2a + 2$ or else $a \geq 2n$ further implies that we must have $a \geq 2n$ for this case. If $s \geq j' = 2$, then

$$6n + 18m + 9t + 6a \leq 11n + 12m + 3a + 1 \leq 8n + 24m + 12t + 8a$$

so that

$$(3n - 12m - 5a + 1)/12 \leq t \leq (5n - 6m - 3a + 1)/9.$$

If $s \geq j' = 3$, then

$$10n + 30m + 15t + 10a \leq 11n + 12m + 3a + 1 \leq 12n + 36m + 18t + 12a$$

so that

$$t \leq (n - 18m - 7a + 1)/15.$$

- (3) Suppose $v = v_{r,1}$, $r \in [1, a]$ with $k = r$. Therefore, we must have

$$(2j' - 1)r \leq 11n + 12m + 3a + 1 \leq 2j'r.$$

Note that $s \geq 2$ if $r = 1$, and $s \geq 1$ otherwise.

- (4) Suppose $v = v_{r,3}$ with $k = a + r$. Therefore, we must have

$$(2j' - 1)(a + r) \leq 11n + 12m + 3a + 1 \leq 2j'(a + r).$$

- (5) Suppose $v = u_{i,1}$, $i \in [1, m]$ with $k = 2n + 2a + 3i - 2$. Therefore, we must have

$$(2j' - 1)(2n + 2a + 3i - 2) \leq 11n + 12m + 3a + 1 \leq 2j'(2n + 2a + 3i - 2).$$

- (6) Suppose $v = u_{i,3}$ with $k = 2n + 2a + 3i - 1$. Therefore, we must have

$$(2j' - 1)(2n + 2a + 3i - 1) \leq 11n + 12m + 3a + 1 \leq 2j'(2n + 2a + 3i - 1).$$

- (7) Suppose $v = u_{i,5}$ with $k = 2n + 2a + 3i$. Therefore, we must have

$$(2j' - 1)(2n + 2a + 3i) \leq 11n + 12m + 3a + 1 \leq 2j'(2n + 2a + 3i).$$

- (8) Suppose $v = u_{i,2}$ with $k = 8n + 12m + 2a - 3i + 3$. Therefore, we must have

$$(2j' - 1)(8n + 12m + 2a - 3i + 3) \leq 11n + 12m + 3a + 1 \leq 2j'(8n + 12m + 2a - 3i + 3).$$

Since $i \leq m$, we must have $j' = 1$ and the equality always holds.

- (9) Suppose $v = u_{i,4}$ with $k = 8n + 12m + 2a - 3i + 1$. Therefore, we must have

$$(2j' - 1)(8n + 12m + 2a - 3i + 1) \leq 11n + 12m + 3a + 1 \leq 2j'(8n + 12m + 2a - 3i + 1).$$

Similar to (8), the equality always hold.

- (10) Suppose $v = u_{i,6}$ with $k = 8n + 12m + 2a - 3i + 2$. Therefore, we must have

$$(2j' - 1)(8n + 12m + 2a - 3i + 2) \leq 11n + 12m + 3a + 1 \leq 2j'(8n + 12m + 2a - 3i + 2).$$

Similar to (8), the equality always hold.

- (11) Suppose $v = y_{t,3}, t \in [1, n]$ with $k = 8n + 6m + 2a - 3t + 3$. Therefore, we must have

$$(2j' - 1)(8n + 6m + 2a - 3t + 3) \leq 11n + 12m + 3a + 1 \leq 2j'(8n + 6m + 2a - 3t + 3).$$

Thus, $j' = 1$ so that $t \leq (5n + a + 5)/6$. Since $t \leq n$, we have $a + 5 \leq n$. The conditions $n \geq 2a + 2$ or else $a \geq 2n$ further implies that we must have $n \geq a + 5$ for $a = 1, 2$, and $n \geq 2a + 2$ otherwise.

- (12) Suppose $v = y_{t,5}$ with $k = 8n + 6m + 2a - 3t + 1$. Therefore, we must have

$$(2j' - 1)(8n + 6m + 2a - 3t + 1) \leq 11n + 12m + 3a + 1 \leq 2j'(8n + 6m + 2a - 3t + 1).$$

Thus, $j' = 1$ so that $t \leq (5n + a + 1)/6$. Since $t \leq n$, we must have $a + 1 \leq n$. Similar to (11), we must have $n \geq 2a + 2$ for this case.

- (13) Suppose $v = y_{t,2}$ with $k = 2n + 6m + 2a + 3t - 2$. Therefore, we must have

$$(2j' - 1)(2n + 6m + 2a + 3t - 2) \leq 11n + 12m + 3a + 1 \leq 2j'(2n + 6m + 2a + 3t - 2).$$

Thus, $j' \leq 3$. If $j' = 1$, then

$$2n + 6m + 2a + 3t - 2 \leq 11n + 12m + 3a + 1 \leq 4n + 12m + 4a + 6t - 4$$

so that

$$(7n - a + 5)/6 \leq t \leq (9n + 6m + a + 3)/3.$$

Since $t \leq n$, we then have $a \geq n + 5$. The conditions $n \geq 2a + 2$ or else $a \geq 2n$ further implies that we must have $a \geq n + 5$ for $a = 1, 2, 3, 4$ and $a \geq 2n$ otherwise. If $s \geq j' = 2$, then

$$6n + 18m + 6a + 9t - 6 \leq 11n + 12m + 3a + 1 \leq 8n + 24m + 8a + 12t - 8$$

so that

$$(3n - 12m - 5a + 9)/12 \leq t \leq (5n - 6m - 3a + 7)/9.$$

If $s \geq j' = 3$, then

$$10n + 30m + 10a + 15t - 10 \leq 11n + 12m + 3a + 1 \leq 12n + 36m + 12a + 18t - 12$$

so that

$$t \leq (n - 18m - 7a + 13)/15.$$

- (14) Suppose $v = y_{t,4}$ with $k = 2n + 6m + 2a + 3t - 1$. Therefore, we must have

$$(2j' - 1)(2n + 6m + 2a + 3t - 1) \leq 11n + 12m + 3a + 1 \leq 2j'(2n + 6m + 2a + 3t - 1).$$

Thus, $j' \leq 3$. If $j' = 1$, then

$$2n + 6m + 2a + 3t - 1 \leq 11n + 12m + 3a + 1 \leq 4n + 12m + 4a + 6t - 2$$

so that

$$(7n - a + 3)/6 \leq t \leq (9n + 6m + a + 2)/3.$$

Since $t \leq n$, we then have $a \geq n + 3$. The conditions $n \geq 2a + 2$ or else $a \geq 2n$ further implies that we must have $a \geq n + 3$ for $a = 1, 2$ and $a \geq 2n$ otherwise. If $s \geq j' = 2$, then

$$6n + 18m + 6a + 9t - 3 \leq 11n + 12m + 3a + 1 \leq 8n + 24m + 8a + 12t - 4$$

so that

$$(3n - 12m - 5a + 5)/12 \leq t \leq (5n - 6m - 3a + 4)/9.$$

If $s \geq j' = 3$, then

$$10n + 30m + 10a + 15t - 5 \leq 11n + 12m + 3a + 1 \leq 12n + 36m + 12a + 18t - 6$$

so that

$$t \leq (n - 18m - 7a + 6)/15.$$

- (15) Suppose $v = v_{r,2}$, $r \in [1, a]$ with $k = 12m + 11n + 3a + 1 - r$. Therefore, we must have

$$(2j' - 1)(12m + 11n + 3a + 1 - r) \leq 11n + 12m + 3a + 1 \leq 2j'(12m + 11n + 3a + 1 - r).$$

Since $r \leq a$, we must have $j' = 1$ and the equality always holds.

For (1) to (4), Theorem 2.5(a)(i) implies that $\chi_{lt}(G_v(k, s)) = 2n + 2a + ks$. For (5) to (15), By Theorem 2.5(b), we have $\chi_{lt}(G_v(k, s)) = 2n + 2a + ks + 1$. This completes the proof. \square

Example 4.11. Figure 8 gives the local total antimagic 39-coloring of $G_v(k, s)$ for $G = C_6 + P_6 + 3P_3$, $v = v_{3,2}$, $k = 30$, $s = 1$ with induced weights 28, 29, 33, 34, 35, 36, 37, 38, 69, 70, ..., 98 and 2576 being the weight of $v_{3,2}$ in $G_v(k, s)$ as defined above.




Fig. 8. $G_v(k, s)$, $v = v_{3,2}$, $k = 30$, $s = 1$ with local total antimagic 39-coloring

5. CONCLUSIONS

In this paper, we successfully characterize graphs with $\chi_{lt}(G) = 3$. Moreover, we obtain a sufficient condition for a graph G with $k \geq 2$ pendant edges such that $\chi_{lt}(G) \geq k + 2$.

A family of graphs G with $\chi_{lt}(G) \geq k+2$ is given. We then obtain several families of graphs with (i) $\Delta(G) = k+2$ and $\chi_{lt}(G) = k+3$, or (ii) $k \geq \Delta(G)$ and $\chi_{lt}(G) = k+1$. The following problems arise naturally.

Problem 5.1. For $2n \geq k+3 \geq 4$ and $2n(2n+1) - 2n(k+2)(5k+5) + (k+2)(k-1) > 0$, show that $\chi_{lt}(f_n(k)) = 2nk + 2$. Otherwise, $\chi_{lt}(f_n(k)) = 2nk + 1$.

Problem 5.2. Determine $\chi_{lt}(G)$ for $G \cong P_n, C_n$ for $n \geq 3$ completely.

Problem 5.3. Determine $\chi_{lt}(G_v(k, s))$ for $G = mC_6 + nP_6$, $m \geq 1, n \geq 1$.

Problem 5.4. Determine $\chi_{lt}(mC_6 + nP_6 + aP_3)$ for $m, a \geq 1, (a+1)/2 \leq n \leq 2a+1$.

We note that $P_n, n \geq 3$ is the only graph with $\Delta(P_n) = 2$ = the number of pendant edges. Thus, we also pose the following problem.

Problem 5.5. Determine $\chi_{lt}(G)$ for G with $k \geq 3$ pendant edges and $\Delta(G) = k$.

In all the known results on graphs G with $k \geq \Delta(G)$ pendant edges, we have $k+1 \leq \chi_{lt}(G) \leq k+2$. Similar to the conjecture of $\Delta(G)+1 \leq \chi_t(G) \leq \Delta(G)+2$, we end the paper with the following conjectures.

Conjecture 5.6. If G has $k \leq \Delta(G)$ pendant edges, then

$$\Delta(G) + 1 \leq \chi_{lt}(G) \leq \Delta(G) + 2.$$

Conjecture 5.7. If G has $k \geq \Delta(G)$ pendant edges, then

$$k + 1 \leq \chi_{lt}(G) \leq k + 2.$$

Acknowledgements

The author thanks the anonymous referee for the many suggestions that helped improve the paper a great deal. Most results of this paper were obtained while the author was affiliated to Universiti Teknologi MARA (Segamat Campus). He is grateful to the university for all the supports given.


REFERENCES

- [1] I.H. Agustin, M. Hasan, Dafik, R. Alfarisi, R.M. Prihandini, *Local edge antimagic coloring of graphs*, Far East J. Math. Sci. **102** (2017), 1925–1941.
- [2] K. Appel, W. Haken, *Every map is four colourable*, Bull. Am. Math. Soc. **82** (1976), 711–712.
- [3] S. Arumugam, K. Premalatha, M. Bača, A. Semaničová-Feňovčíková, *Local antimagic vertex coloring of a graph*, Graphs Combin. **33** (2017), 275–285.
- [4] M. Behzad, *Graphs and their chromatic numbers*, PhD Thesis, Michigan State University, 1965.
- [5] J. Bensmail, M. Senhaji, K. Szabo Lyngsie, *On a combination of the 1–2–3 conjecture and the antimagic labelling conjecture*, Discrete Math. Theoret. Comput. Sci. **19** (2017), no. 1, #22.

- [6] J. Geetha, N. Narayanan, K. Somasundaram, *Total colorings-a survey*, AKCE Int. J. Graphs Comb. **20** (2023), no. 3.
- [7] J. Haslegrave, *Proof of a local antimagic conjecture*, Discret. Math. Theor. Comput. Sci. **20** (2018), no. 1.
- [8] G.C. Lau, W.C. Shiu, *On join product and local antimagic chromatic number of regular graphs*, Acta Math. Hungar. **169** (2023), no. 1, 108–133.
- [9] G.C. Lau, W.C. Shiu, *On local antimagic chromatic number of lexicographic product graphs*, Acta Math. Hungar. **169** (2023), no. 1, 158–170.
- [10] G.C. Lau, W.C. Shiu, *On local antimagic total labeling of complete graphs amalgamation*, Opuscula Math. **43** (2023), no. 3, 429–453.
- [11] G.C. Lau, H.K. Ng, W.C. Shiu, *Affirmative solutions on local antimagic chromatic number*, Graphs Combin. **36** (2020), 1337–1354.
- [12] G.C. Lau, W.C. Shiu, H.K. Ng, *On local antimagic chromatic number of cycle-related join graphs*, Discuss. Math. Graph Theory **4** (2021), no. 1.
- [13] G.C. Lau, K. Schaffer, W.C. Shiu, *Every graph is local antimagic total and its applications*, Opuscula Math. **43** (2023), no. 6, 841–864.
- [14] V. Sandhiya, M. Nalliah, *Local total antimagic chromatic number of graphs*, Heliyon **9** (2023), e17761.
- [15] V. Sandhiya, M. Nalliah, private communication.
- [16] B. Toft, R. Wilson, *A brief history of edge-colorings – with personal reminiscences*, Discrete Math. Lett. **6** (2021), 38–46.

Gee-Choon Lau

geeclau@yahoo.com

 <https://orcid.org/0000-0002-9777-6571>

77D, Jalan Subuh, 85000, Johor, Malaysia

Received: January 30, 2024.

Revised: February 4, 2025.

Accepted: February 14, 2025.