
Figure 6 of arXiv:1708.02243: contours of 3 (exclusion) and 5 (discovery) signal events at 300 fb^{-1}, no background, as in the paper
left: signal SCT z = 1.16 mm (literal text); right: signal without SCT z smearing (what the paper's results require)



Normalisation: σ = MadGraph5 3.7.3 LO (0 + 1 jet, MLM-matched with Pythia8) for one quirk species, no K-factor; the paper gives no σ. The "N_Q = 2" of the paper's panel titles is NOT applied (no factor 2 on σ). Colored: a factor 1.5–1.7 on σ brings our maxima close to the paper's; EW: our maxima are already at the paper's (+2–4%), a factor > 1 would overshoot (sensitivity, not a fit; F6_kfactor).
Trigger (both channels, the paper's described method): ε_trig = N(p_T($Q\bar{Q}$) > 200 GeV) / N(matched) of the same matched + shower sample that gives ε_fid and ε_reco (EW 0.148 at 800 GeV). Alternative, not stated in the paper: the EW LO ratio σ($Q\bar{Q}$ + j, p_T(j) > 200) / σ($Q\bar{Q}$) (0.096 at 800 GeV, = Table I's 0.10) is lower by a factor 0.54–0.75 and gives EW 712 / 643 GeV (exclusion / discovery, no SCT z smearing). Colored: the both-charged factor (~0.33) is included (thin orange: without it).
Red ×: at low Λ for colored 600–1200 GeV and EW 100–200 GeV our samples (40–70 events per mass) are too small; the paper's lower-left edge passes there and is not tested here (larger samples needed). Λ = 15–25 keV: computed (the integrator's 20-million-step ceiling removes ≤ 13% of the events fiducial at 10 keV there); Λ > 25 keV: not computed, because that ceiling removes 5–56% of them at 30–35 keV (needs a larger ceiling and more memory). The maxima are at 3–5 keV and are unaffected.
Annihilation is not simulated (as in the paper). Worst case (α = 0.05, annihilation chance at every second meeting) it lowers the signal near the maxima by ~17–24% (EW 4–5 keV) and ~17% (colored 5 keV), moving the maxima inward by ~30 GeV (colored) and ~25–40 GeV (EW).