

DISCHARGE SUMMARY - SYNTHETIC PATIENT DATA (NOT REAL)

Patient: Mary Johnson

Date of Birth: 03/22/1950

Discharge Date: 2026-06-12

Admit Date: 2026-06-10

Hospital: Northside Orthopedic Center

Attending Physician: Dr. Angela Reyes

Primary Specialty: Orthopedic Surgery

DIAGNOSIS

Primary Diagnosis: Total Right Knee Arthroplasty

ICD-10 Code: Z96.651

MEDICATIONS AT DISCHARGE

1. Oxycodone/Acetaminophen 5/325mg - Every 6 hours as needed for pain (do not drive)
2. Aspirin 325mg - Once daily with food for blood clot prevention
3. Celecoxib 200mg - Twice daily with food for swelling and pain
4. Enoxaparin (Lovenox) injection 40mg - Once daily under skin for 10 days

FOLLOW-UP APPOINTMENTS

- Orthopedic surgeon Dr. Reyes - 10-14 days post discharge for wound check
- Physical therapy - begin 2 days after discharge, 3 times per week for 6 weeks

ACTIVITY RESTRICTIONS

- Walk with walker for full weight bearing as tolerated
- Do not kneel on the operated knee for 8 weeks
- No driving until cleared by surgeon (typically 4-6 weeks)
- Elevate leg above heart level when resting to reduce swelling
- Perform ankle pumps every hour while awake
- Ice knee for 15-20 minutes several times per day

DIETARY RESTRICTIONS

- No specific dietary restrictions
- Stay well hydrated - drink 6-8 glasses of water daily
- Constipation is common with opioid pain medication - increase fiber intake

GO TO THE EMERGENCY ROOM IMMEDIATELY IF YOU EXPERIENCE

- Sudden chest pain or difficulty breathing (possible blood clot in lungs)
- Severe calf pain or swelling (possible blood clot in leg)
- Signs of infection: fever above 101.5F, wound drainage, increasing redness
- Sudden severe pain not controlled by medication

CALL YOUR DOCTOR WITHIN 24 HOURS IF YOU NOTICE

- Wound is oozing or has increased redness or warmth

- Knee swelling is getting significantly worse
- Numbness or tingling in foot or leg
- Fever between 100-101.5 degrees F