

DISCHARGE SUMMARY - SYNTHETIC PATIENT DATA (NOT REAL)

Patient: Helen Rodriguez

Date of Birth: 06/10/1940

Discharge Date: 2026-06-13

Admit Date: 2026-06-08

Hospital: St. Mary Orthopedic Hospital

Attending Physician: Dr. William Nguyen

Primary Specialty: Orthopedic Surgery

DIAGNOSIS

Primary Diagnosis: Right Hip Fracture - Open Reduction Internal Fixation (ORIF)

ICD-10 Code: S72.001A

Secondary: Osteoporosis, Hypertension

MEDICATIONS AT DISCHARGE

1. Enoxaparin (Lovenox) 40mg - Once daily subcutaneous injection for 28 days
2. Acetaminophen 650mg - Every 6 hours scheduled (do not exceed 3g per day total)
3. Oxycodone 5mg - Every 4-6 hours as needed for breakthrough pain
4. Calcium Carbonate 600mg + Vitamin D3 800IU - Twice daily with food
5. Alendronate (Fosamax) 70mg - Once weekly on same day, on empty stomach with full glass of water
6. Amlodipine 5mg - Once daily for blood pressure

FOLLOW-UP APPOINTMENTS

- Orthopedic surgeon Dr. Nguyen - 10-14 days for wound check and X-ray
- Physical therapy - daily or home PT 3x weekly
- Occupational therapy - home safety evaluation within 5 days
- Primary care - 2 weeks for blood pressure and medication review
- Bone density scan (DEXA) - schedule within 3 months

ACTIVITY RESTRICTIONS

- Weight bearing as tolerated on right leg - use walker at all times
- Do not cross legs or bend hip past 90 degrees
- Do not pivot on the operated leg
- Sleep with pillow between knees to keep hip in proper position
- Fall prevention is critical - install grab bars, remove hazards from home
- No driving for minimum 6-8 weeks and until cleared by surgeon

DIETARY RESTRICTIONS

- High protein diet to support healing - eggs, fish, chicken, legumes
- Adequate calcium and vitamin D intake (supplements provided)
- Stay well hydrated to reduce blood clot risk
- Take Alendronate on empty stomach, stay upright 30 minutes after

GO TO THE EMERGENCY ROOM IMMEDIATELY IF YOU EXPERIENCE

- Sudden chest pain or difficulty breathing (signs of pulmonary embolism)
- Severe swelling, redness, or warmth in calf (signs of DVT)
- Signs of hip dislocation: sudden severe pain, leg appears shorter or rotated
- Surgical site infection: increasing redness, fever above 101F, wound drainage
- Loss of feeling in leg or foot

CALL YOUR DOCTOR WITHIN 24 HOURS IF YOU NOTICE

- Mild calf pain or swelling in the leg
- Wound drainage or increased wound redness
- Fever 100-101 degrees F
- Nausea preventing taking medications
- Falling at home - even if no apparent injury
- Blood pressure readings above 160/100