


Name: Ragib Ahsan
Eco602: Week10_Using_Models1

Q1 Answer:



Q2 Answer:

p-value = 0.04097

#R Code- `shapiro.test(catrate$cat.rate)`

Q3 Answer:

Hypothesis- Collected data was from distributed data set.

Q4 Answer:

Lower p-value indicates that the data is from non-normally distributed population.

Q5 Answer:

#R Code- `t.test(catrate$cat.rate, mu = 0.2857143)`

Q6 Answer:

Hypothesis- salamander catastrophic reproduction rate is the same as the pond late filling rate.

Q7 Answer:

Because of the main interest in non-directionality of the data, it's a two-tailed test.

Q8 Answer:

p-value = 0.01193, which is below the range of 0.05. Because of that a notable difference is expected between the two data sets (salamander reproductive catastrophic rate and pond late filling rate), which could be the false positive rate of 1.20%, suggests no difference in the groups of 1.20% of the time.

Q9 Answer:

Confidence interval= 0.3526250 to 0.7261295, without zero.

Q10 Answer:

There could be a strong evidence to reject the null-hypothesis, because the p-value is lower.

Q11 Answer:

```
#R Code- wilcox.test(x= catrate$cat.rat, mu = 0.2857143)
```

Q12 Answer:

P-value from the Wilcoxon test is 0.006275, which is lower compared to the previous p-value- 0.01193.

Q13 Answer:

Yes, Wilcoxon test p-value could lead to reject the null hypothesis and suggests about the difference between the two datasets (reproductive catastrophic rate and pond late-filling rate). So, the rejection of null-hypothesis is possible based on this.

Q14 Answer:

Both tests indicate the lower p-values, which discourage the accept the null-hypothesis. Specifically, the p-value from Wilcoxon test, suggests to reject the null-hypothesis.

Q15 Answer:

I would choose the Wilcoxon test because, the data distribution type is more appropriate for this Wilcoxon test here.

Q16 Answer:

```
#R Code- shapiro.test(dat_adelie$flipper_length_mm)
```

Q17 Answer:

Both data sets are normally distributed. Because, their p-value is high enough to be the normally distributed data (for both Adelie and Chinstrap penguins- 0.72, and 0.8106). Here, these p-values supports the null-hypothesis.

Q18 Answer:

I had an issue, to get a combined image file with specific width and length.

Q19 Answer:

The alternative hypothesis- the flipper lengths are different in the Adelie penguins and Chinstrap penguins.

Q20 Answer:

```
t.test(flipper_length_mm ~ species, data = penguin_dat)
```