

BEYOND STANDARD MODEL

Problems of the Standard Model. Search Strategy. Indirect search. Supersymmetry

Lecture 18

DIPARTIMENTO DI FISICA



SAPIENZA
UNIVERSITÀ DI ROMA

Shahram Rahatlou

Fisica delle Particelle Elementari, Anno Accademico 2015-16

<http://www.roma1.infn.it/people/rahatlou/particelle/>

SUCCESS OF STANDARD MODEL


- LEP Precisions measurements have proved SM to be reliable and correct in describing almost all phenomenology so far
- CP violation well described and verified experimentally
- Higgs discovered with mass ~ 125 GeV compatible with precision tests at LEP
- Neutrinos also integrated well within the Standard Model
- *Shortcomings of the Standard Model*
 - cannot incorporate gravity
 - cannot survive any attempt at unification of all forces
 - cannot explain why all physics below TeV scale!
 - cannot explain why Higgs mass is so fine-tuned
 - cannot explain what happens up to Planck scale

HINTS OF NEW PHYSICS

- Neutrinos have very small but non-zero mass
 - so far we have always assumed them to be massless
- Astrophysical proof of existence of cold dark matter and we also need a large amount of yet-to-be-understood dark energy


COMPOSITION OF THE COSMOS

- Still no idea about origin of masses and mixing structure




CORRECTIONS To HIGGS MASS


- Propagator of Higgs mass affected by higher order corrections



$$\delta m_h^2 \sim -\frac{3}{8\pi^2} \lambda_t^2 \Lambda^2$$




$$\delta m_h^2 \sim \frac{9}{64\pi^2} g^2 \Lambda^2$$



$$\delta m_h^2 \sim \frac{1}{16\pi^2} \lambda^2 \Lambda^2$$

- Such terms change the bare Higgs mass. Regularization is needed to keep corrections finite

HIGGS PROPAGATOR




a)

$$\begin{aligned} &= \int \frac{d^4 p}{(2\pi)^4} (-) N_c \text{Tr} \left[i\lambda_t \frac{i}{p} i\lambda_t \frac{i}{p} \right] \\ &= -N_c \lambda_t^2 \int \frac{d^4 p}{(2\pi)^4} \text{Tr} \left[\frac{1}{p^2} \right] \quad \text{Tr}[1] = 4 \\ &= -\frac{4N_c \lambda_t^2}{(2\pi)^4} \int \frac{d^4 p}{p^2} \end{aligned}$$

- quadratic divergence with cutoff M_{UV}


HIGGS MASS CORRECTIONS

- High energy (UltraViolet) cutoff M_{UV} to regularize corrections
 - M_{UV} is highest energy accessible by known particles
 - For $E > M_{UV}$ new particles enter the loop
- Fermionic corrections depend on fermion mass and coupling to Higgs
 - top contribution by far most dominant




$$\Delta m_H^2 = \frac{\lambda_f^2}{16\pi^2} [-2M_{UV}^2 + 6m_f^2 \ln(M_{UV}/m_f) + \dots]$$

- Higgs mass is sensitive to highest energy scale in the theory
- Divergences in Higgs mass if M_{UV} too large
- How can we determine the energy scale?




SCALE FOR NEW PHYSICS?

- Since birth of particle physics experiments have explored many orders of magnitude in energy
- Different phenomena have appeared at different scales
- Standard Model and EW breaking occurs up to TeV scale
- How to determine scale of new physics beyond SM ?



GRAND UNIFICATION



- Running of coupling constants leads to unification of different forces at very high energy scale
- Gravity also can be included at Planck scale

PLANCK SCALE

- Mass with same Compton wavelength and Schwarzschild radius
- Compton wavelength: defines length scale where quantum mechanics must be used
 - decreases with larger mass
 - e.g. for a photon the corpuscular nature of light becomes relevant
- Schwarzschild radius is the radius in which a confined mass object becomes a black hole
 - classically radius such that escape velocity equal to speed of light
 - increases for larger mass
- Planck mass or scale defines the scale at which quantum and gravitational effects are both relevant and comparable

$$E = mc^2 = \hbar\nu \rightarrow mc^2 = \frac{\hbar c}{\lambda_c} \rightarrow \lambda_c = \frac{\hbar}{mc^2}$$

$$\frac{1}{2} M v^2 = \frac{GMm}{r} \rightarrow r_s = \frac{2Gm}{c^2}$$

$$m_P = \sqrt{\frac{c\hbar}{G}} = 1.2 \cdot 10^{19} \text{ GeV}/c^2$$

$$\ell_P = \frac{\hbar}{m_P c} = \sqrt{\frac{\hbar G}{c^3}} = 1.6 \cdot 10^{-35} \text{ m}$$

$$t_P = \ell_P / c = \sqrt{G\hbar/c^5} = 5.4 \cdot 10^{-44} \text{ s}$$

NATURALNESS OR FINE-TUNING PROBLEM

$$m_h^2 \sim m_{h0}^2 + \delta m_h^2$$

- We know that $m_h = 125$ GeV
- Nature sets m_{h0} at Planck scale and we observe the physical mass after all higher order correction terms
- For Higgs mass to be finite at EW scale, corrections must balance the bare mass over 16 order of magnitude
- This is not a consistency problem for the theory but requires **incredible fine-tuning of parameters** to achieve such precise cancellation
- Such accidental features although possible are extremely unlikely
- Nature generally prefers rules and symmetries to accidents

WHAT IS A NATURAL SCALE?

- We could afford corrections of the order of the Higgs mass
 - fermion mass corrections are proportional to their mass
 - ▶ approximate chiral symmetry
- If $M_{UV} = 1 \text{ TeV}$ the Higgs mass fine tuning would be natural
- This implies that new particles and processes to be discovered at LHC!

YET ANOTHER HEADACHE: HIERARCHY PROBLEM

- Plank scale occurs 16 orders of magnitude higher than Standard Model
 - $\Lambda_{\text{pl}} \gg \text{EW scale}$
- Almost all known phenomena so far occur within TeV scale
- Past experience tells us that going up in energy scale we find new phenomena
- Such a desert in energy scale is very unlikely and hard to explain
- This is known as the hierarchy problem
 - Huge energy gap between new physics scale and electroweak scale
- LHC will explore the gap by searching for signatures and phenomena not predicted by Standard Model at TeV scale

SEARCH STRATEGIES

- Indirect Searches
 - New physics by observing deviations from predictions
 - Enhanced decay and production rates due to new particles in loops
 - ▶ $B_s \rightarrow \mu\mu$ branching ratio
- Supersymmetry
 - complete theory with **few** free parameters
 - scan parameter space
 - Rich and well defined phenomenology
 - Well defined Higgs sector
- Direct search for exotic particles
 - kinematic observables or mass of new particles
 - Many models and many signatures
 - No comprehensive theory
 - very good signal to noise

$B_s \rightarrow \mu\mu$

B PRODUCTION AT HADRON COLLIDERS


- At B factor \times sec ~ 1 nb

$$\sigma = N_c Q_f^2 \frac{4\pi\alpha^2}{3s} = N_c Q_f^2 \frac{86.8 \text{ nb}}{s(\text{GeV}^2)}$$


- At LHC @ 7 TeV \times sec ~ 0.3 mb

$$\frac{d\sigma}{d\Omega}(q\bar{q}, gg \rightarrow q\bar{q}) \sim \frac{\alpha_s^2}{s}$$



Lowest order




Flavor excitation



Gluon splitting




B PRODUCTION TOPOLOGY




- Events of interest almost exclusively in forward region
 - Need excellent coverage at high η
- All flavors of b hadrons produced
 - Relative production ratio: $B_d/B_u/B_s/B_c/b\text{-baryons}$ 4:4:1:0.01:1

LHCb DETECTOR



- Dedicated detector covering $2 < \eta < 5$ region
 - Measurements also with ATLAS and CMS
- Extremely competitive for states with charged tracks but no access to neutrino
 - no semileptonic decays


FLIGHT LENGTH



- Large impact parameters and very displaced secondary vertex
 - Large boost and flight length of ~ 1 cm
 - ▶ compare to 250 microns at B factory

$B_s \rightarrow \mu\mu$

- Rare B decay in Standard Model




$$B_s \rightarrow \mu^+ \mu^- = (3.2 \pm 0.2) \times 10^{-9}$$

$$B_d \rightarrow \mu^+ \mu^- = (1.0 \pm 0.1) \times 10^{-10}$$

BACKGROUNDS



$$\begin{aligned} B^0 &\rightarrow \pi^- \mu^+ \nu \\ B_s &\rightarrow K^- \mu^+ \nu \\ \Lambda_b &\rightarrow p \mu \nu \end{aligned}$$

$$\begin{aligned} B^{0/+} &\rightarrow \pi^{0/+} \mu \mu \\ B_c &\rightarrow J/\psi(\mu \mu) \mu \nu \end{aligned}$$





- 2 real muons
- 1 real muon and 1 fake muon
- But also two fake muons: very small BF for signal!

BACKGROUND REJECTION





- Combination of two muons from different B mesons
- Multivariate tools to combine information and maximize background rejection
 - total of 12 B and muon variables
 - ▶ impact parameter of muon tracks
 - ▶ muon isolation
 - ▶ p_T of B
- Use each bin as a separate channel since different S/B
 - maximize power of each bin







EVIDENCE AT LHCb





	90 % CL	95 % CL
Exp. bkg	3.5×10^{-10}	4.4×10^{-10}
Exp. bkg+SM	4.5×10^{-10}	5.4×10^{-10}
Observed	6.3×10^{-10}	7.4×10^{-10}

EVIDENCE AT CMS



ANOTHER TEST OF STANDARD MODEL




- Evidence at LHCb
 - <http://arxiv.org/abs/1211.2674>
- Evidence at CMS
 - <http://arxiv.org/abs/1307.5025>
- Combination of CMS and LHCb rare decays
 - <http://cds.cern.ch/record/1564324>

SUPERSYMMETRY

A SOLUTION TO HIGGS MASS DIVERGENCE

- scalar particles contribution to Higgs mass also quadratically divergent with MUV
- Contribution with opposite sign compared to fermions



$$\Delta m_H^2 = \frac{\lambda_S}{16\pi^2} [M_{\text{UV}}^2 - 2m_S^2 \ln(M_{\text{UV}}/m_S) + \dots]$$

$$\Delta m_H^2 = \frac{\lambda_f^2}{16\pi^2} [-2M_{\text{UV}}^2 + 6m_f^2 \ln(M_{\text{UV}}/m_f) + \dots]$$

- If such scalar particles existed, fermion and scalar contributions could cancel each other **exactly and naturally** without fine tuning
- Such conspiracy is generally known as a **symmetry** of the theory!

ORIGIN OF SUSY

- ❖ Before discovery of antimatter, electromagnetism had a problem

$$(m_e)^2 = (m_{e,0})^2 + \Delta E_{\text{Coulomb}}$$


$= 0.5 \text{ MeV}$

$$\Delta E_{\text{Coulomb}} = \frac{1}{4\pi\epsilon_0} \frac{e^2}{r_e}$$

$r_e < 10^{-17} \text{ cm} \rightarrow \underline{\Delta E > 10 \text{ GeV}}$

- ❖ **Solution:** for every fermion, add an antifermion

- Additional diagram that **cancels ΔE !**
- Also **doubles** degrees of freedom of theory



SUPERSYMMETRY (SUSY)

- New symmetry of Lagrangian to cancel quadratic divergences of Higgs mass
- Create super partners for each fermion and boson of SM

$$Q^\dagger |\text{Boson}\rangle = |\text{Fermion}\rangle; \quad Q^\dagger |\text{Fermion}\rangle = |\text{Boson}\rangle$$

$$Q |\text{Boson}\rangle = |\text{Fermion}\rangle; \quad Q |\text{Fermion}\rangle = |\text{Boson}\rangle$$

- Q operator is a spinor: spin 1/2 and defined helicity
- Algebra of Q operator constrained when acting on chiral fermions

$$\{Q, Q^\dagger\} = P^\mu$$

$$\{Q, Q\} = \{Q^\dagger, Q^\dagger\} = 0$$

$$[P^\mu, Q] = [P^\mu, Q^\dagger] = 0$$

- $P^\mu = (H, \vec{P})$ is the momentum operator
- T^a are the gauge operators

WHERE ARE SUSY PARTICLES?

- Many SUSY particles have already been observed
 - leptons, quarks, W, and Z
 - same particles included in SM
- But no SUSY partner of SM observed yet!
- If SUSY is an exact symmetry, we should have seen SUSY partners of known particles with the same mass
- SUSY is certainly broken!
 - Spontaneous SUSY breaking must be added by hand and still avoid divergences in Higgs mass corrections
 - Different symmetry breaking mechanisms on the market
 - we'll see a short summary later

SUSY BREAKING MECHANISM

- Soft breaking means SUSY is not exact but still addresses the naturalness/hierarchy problem

$$\Delta m_H^2 = \frac{1}{16\pi^2} (\lambda_S - \lambda_F^2) M_{\text{UV}}^2 + \dots$$

- Add SUSY breaking term in Lagrangian $\mathcal{L} = \mathcal{L}_{\text{SUSY}} + \mathcal{L}_{\text{soft}}$

$\mathcal{L}_{\text{SUSY}}$ contains all of the gauge, Yukawa, and dimensionless scalar couplings, and preserves exact supersymmetry


$\mathcal{L}_{\text{soft}}$ violates supersymmetry, and contains only mass terms and couplings with *positive* mass dimension.

- Couplings of fermions and scalars are not identical but only slightly different so that corrections are logarithmically divergent

$$\Delta m_H^2 = m_{\text{soft}}^2 \left[\frac{\lambda}{16\pi^2} \ln(M_{\text{UV}}/m_{\text{soft}}) + \dots \right]$$

- Sufficient to have $m_{\text{soft}} \sim \text{TeV}$
- Such breaking saves the Higgs mass and opens up possibility of new phenomena at TeV scale to be tested now at LHC

MASS SCALE OF SUSY



- m_{soft} determines the scale of SUSY particles
- SUSY particles cannot be light or lighter than already known particles

SOFT SUSY BREAKING IN MSSM

$$\begin{aligned}
\mathcal{L}_{\text{soft}}^{\text{MSSM}} = & -\frac{1}{2} (M_3 \tilde{g}\tilde{g} + M_2 \tilde{W}\tilde{W} + M_1 \tilde{B}\tilde{B}) + \text{c.c.} \\
& - (\tilde{u} \mathbf{a_u} \tilde{Q} H_u - \tilde{d} \mathbf{a_d} \tilde{Q} H_d - \tilde{e} \mathbf{a_e} \tilde{L} H_d) + \text{c.c.} \\
& - \tilde{Q}^\dagger \mathbf{m_Q^2} \tilde{Q} - \tilde{L}^\dagger \mathbf{m_L^2} \tilde{L} - \tilde{u} \mathbf{m_{\tilde{u}}^2} \tilde{u}^\dagger - \tilde{d} \mathbf{m_{\tilde{d}}^2} \tilde{d}^\dagger - \tilde{e} \mathbf{m_{\tilde{e}}^2} \tilde{e}^\dagger \\
& - m_{H_u}^2 H_u^* H_u - m_{H_d}^2 H_d^* H_d - (b H_u H_d + \text{c.c.})
\end{aligned}$$

first line gives masses to the MSSM gauginos (gluino \tilde{g} , winos \tilde{W} , bino \tilde{B})

second line consists of (scalar)³ interactions.


third line is (mass)² terms for the squarks and sleptons.

last line is Higgs (mass)² terms.

$$\begin{aligned}
M_1, M_2, M_3, \mathbf{a_u}, \mathbf{a_d}, \mathbf{a_e} &\sim m_{\text{soft}}; \\
\mathbf{m_{\tilde{Q}}^2}, \mathbf{m_{\tilde{L}}^2}, \mathbf{m_{\tilde{u}}^2}, \mathbf{m_{\tilde{d}}^2}, \mathbf{m_{\tilde{e}}^2}, m_{H_u}^2, m_{H_d}^2, b &\sim m_{\text{soft}}^2
\end{aligned}$$

- **Total of > 100 parameters in the Lagrangian!**
 - Different number of additional parameters for each type of SUSY breaking
 - In SM there are 27 parameters between masses, couplings, and CP violation parameters

SUSY Zoo



Need to extend SM Higgs sector
Two doublets → **five** Higgs bosons

h, H, A, H⁺, H⁻

CP-even, light:
'our' 125 GeV Higgs

Gauginos and higgsinos mix:

X₁[±], X₂[±] — **four charginos**

X₁⁰, X₂⁰, X₃⁰, X₄⁰ — **four neutralinos**

Lightest Susy Particle (LSP):
stable if R-Parity is conserved

Dark Matter candidate

UNDISCOVERED MSSM ZOOLOGY

Names	Spin	P_R	Mass Eigenstates	Gauge Eigenstates
Higgs bosons	0	+1	$h^0 \ H^0 \ A^0 \ H^\pm$	$H_u^0 \ H_d^0 \ H_u^+ \ H_d^-$
squarks	0	-1	$\tilde{u}_L \ \tilde{u}_R \ \tilde{d}_L \ \tilde{d}_R$ $\tilde{s}_L \ \tilde{s}_R \ \tilde{c}_L \ \tilde{c}_R$ $\tilde{t}_1 \ \tilde{t}_2 \ \tilde{b}_1 \ \tilde{b}_2$	“ ” “ ” $\tilde{t}_L \ \tilde{t}_R \ \tilde{b}_L \ \tilde{b}_R$
sleptons	0	-1	$\tilde{e}_L \ \tilde{e}_R \ \tilde{\nu}_e$ $\tilde{\mu}_L \ \tilde{\mu}_R \ \tilde{\nu}_\mu$ $\tilde{\tau}_1 \ \tilde{\tau}_2 \ \tilde{\nu}_\tau$	“ ” “ ” $\tilde{\tau}_L \ \tilde{\tau}_R \ \tilde{\nu}_\tau$
neutralinos	1/2	-1	$\tilde{N}_1 \ \tilde{N}_2 \ \tilde{N}_3 \ \tilde{N}_4$	$\tilde{B}^0 \ \tilde{W}^0 \ \tilde{H}_u^0 \ \tilde{H}_d^0$
charginos	1/2	-1	$\tilde{C}_1^\pm \ \tilde{C}_2^\pm$	$\tilde{W}^\pm \ \tilde{H}_u^+ \ \tilde{H}_d^-$
gluino	1/2	-1	\tilde{g}	“ ”

NEW SYMMETRY: R-PARITY

$$R = (-1)^{3(B-L)+2S}$$



- Add new conservation law to protect against lepton and baryon number violation
- R-parity combines spin, baryon, and lepton quantum numbers
 - particles: $R = +1$
 - SUSY particles: $R = -1$
- Important phenomenological consequences
 - Lagrangian invariance requires all terms to be quadratic in negative R-parity
 - SUSY particles can only be produced in pairs
 - $R = -1$ particles must always decay in final states with at least one $R = -1$ particle
 - ▶ lightest SUSY particle (LSP) must be stable
 - Two $R = -1$ particles can annihilate and produce ONLY $R = +1$ particles
 - ▶ important for Dark Matter searches

NEXT LIGHTEST SUPERSYMMETRIC PARTICLE (NLSP)

- Two particles play crucial role in SUSY searched
- Lightest supersymmetric particle must be stable and hence escape detection
 - missing energy in SUSY processes
 - SUSY masses are expected to be large therefore expect large missing energy
- Next to lightest supersymmetric particle (NLSP)
 - Because of R-parity conservation must always decay to LSP
 - ▶ two body decay of NLSP $\rightarrow X + \text{LSP}$
 - X will always be an ordinary $R = +1$ particle
 - distinctive kinematic signature for X helps searching for NLSP



SUSY PRODUCTION VIA QCD

Gluino pairs





etc.

Squark pairs



etc.

Squark + gluino




etc.


- QCD production dominates but given heavy mass for SUSY, cross section strongly depends on squark and gluino masses

SUSY PRODUCTION VIA ELECTROWEAK


Chargino pairs










Neutralino pairs



Chargino +
neutralino






Slepton pairs



- Smaller cross section since EW coupling is smaller compared to QCD

SUSY CROSS SECTION



SUSY vs. STANDARD MODEL

