







Lepton Universality in Weak Interactions



How do we know



$$\tau^- \rightarrow e^-/\mu^- + \bar{\nu}_e/\mu + \nu_\tau$$



$$l = e^-/\mu^-$$

$$\Gamma(\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau) = \frac{g_\tau^2}{M_W^2} \frac{g_l^2}{M_W^2} m_\tau^5 \rho_l$$

$\sim \frac{1}{G_F^2}$

phase space of $\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau$

$$\frac{\Gamma(\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau)}{\Gamma(\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau)} = \frac{g_\mu^2}{g_e^2} \frac{\rho_{\tau \rightarrow \mu}}{\rho_{\tau \rightarrow e}}$$

$$\frac{\Gamma(\tau^- \rightarrow \mu^- + X)}{\Gamma_{\text{tot}}} = \text{BR}(\tau^- \rightarrow \mu^- + X)$$

produce $\sim \tau^-$ leptons.

$$\text{BR} = \frac{\# \text{ decays int } \mu^-}{\# \text{ total } \tau^-}$$

$$m_\tau = 1.78 \text{ GeV}$$

$$m_\mu = 106 \text{ MeV} \quad m_e = 0.5 \text{ MeV}$$

| τ^- DECAY MODES | Fraction (Γ_i/Γ) | Scale factor/ Confidence level | p (MeV/c) |
|---|--------------------------------------|-----------------------------------|----------------|
| Modes with one charged particle | | | |
| particle $^- \geq 0$ neutrals $\geq 0K^0 \nu_\tau$ ("1-prong") | $(85.24 \pm 0.06) \%$ | | - |
| particle $^- \geq 0$ neutrals $\geq 0K_L^0 \nu_\tau$ | $(84.58 \pm 0.06) \%$ | | - |
| $\mu^- \bar{\nu}_\mu \nu_\tau$ | [g] $(17.39 \pm 0.04) \%$ | | 885 |
| $\mu^- \bar{\nu}_\mu \nu_\tau \gamma$ | [e] $(3.67 \pm 0.08) \times 10^{-3}$ | | 885 |
| $e^- \bar{\nu}_e \nu_\tau$ | [g] $(17.82 \pm 0.04) \%$ | | 888 |
| $e^- \bar{\nu}_e \nu_\tau \gamma$ | [e] $(1.83 \pm 0.05) \%$ | | 888 |
| $h^- \geq 0K_L^0 \nu_\tau$ | $(12.03 \pm 0.05) \%$ | | 883 |
| $h^- \nu_\tau$ | $(11.51 \pm 0.05) \%$ | | 883 |
| $\pi^- \nu_\tau$ | [g] $(10.82 \pm 0.05) \%$ | | 883 |
| $K^- \nu_\tau$ | [g] $(6.96 \pm 0.10) \times 10^{-3}$ | | 820 |
| $h^- \geq 1$ neutrals ν_τ | $(37.01 \pm 0.09) \%$ | | - |
| $h^- \geq 1\pi^0 \nu_\tau$ (ex. K^0) | $(36.51 \pm 0.09) \%$ | | - |
| $h^- \pi^0 \nu_\tau$ | $(25.93 \pm 0.09) \%$ | | 878 |
| $\pi^- \pi^0 \nu_\tau$ | [g] $(25.49 \pm 0.09) \%$ | | 878 |
| $\pi^- \pi^0 \text{non-}\rho(770) \nu_\tau$ | $(3.0 \pm 3.2) \times 10^{-3}$ | | 878 |
| $K^- \pi^0 \nu_\tau$ | [g] $(4.33 \pm 0.15) \times 10^{-3}$ | | 814 |
| $h^- \geq 2\pi^0 \nu_\tau$ | $(10.81 \pm 0.09) \%$ | | - |
| $h^- 2\pi^0 \nu_\tau$ | $(9.48 \pm 0.10) \%$ | | 862 |
| $h^- 2\pi^0 \nu_\tau$ (ex. K^0) | $(9.32 \pm 0.10) \%$ | | 862 |
| $\pi^- 2\pi^0 \nu_\tau$ (ex. K^0) | [g] $(9.26 \pm 0.10) \%$ | | 862 |
| $\pi^- 2\pi^0 \nu_\tau$ (ex. K^0) | < 9 | $\times 10^{-3} \text{ CL}=95\%$ | 862 |

compute phase space.
measure $BR(\bar{e}, \mu^-)$

\Rightarrow estimate $\frac{g_\mu^2}{g_e^2}$



$$\left(\frac{g_\mu}{g_e}\right)^2 \simeq 0.976 \rightarrow \frac{g_\mu}{g_e} = 1.001 \pm 0.002$$

$\mu \leftrightarrow e$ are the same for Weak int.

what about $\tau \leftrightarrow \mu$ universality?

BR: $\mu^- \rightarrow e^- \bar{\nu}_e \nu_\mu$

$\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau$




$BR(\mu^- \rightarrow e^- \bar{\nu}_e \nu_\mu) \simeq 100\%$




$BR(\tau^- \rightarrow e^- \bar{\nu}_e \nu_\tau) \simeq 17.8\%$

$$\Gamma_{tot}^\mu \simeq \Gamma_{\mu \rightarrow e}$$

$\tau^- \rightarrow \nu_\tau +$




$\begin{matrix} S \\ \bar{U} \end{matrix} \begin{matrix} d \\ \bar{c} \\ \bar{s} \\ \bar{b} \end{matrix}$
 $m \simeq 1.86 \text{ GeV}$
 $> m_\tau$





$\tau^- \rightarrow \nu_\tau + \pi^- + \pi^0$

$$\Gamma_{\tau \rightarrow \mu} \simeq \Gamma_{\tau \rightarrow e} \simeq \Gamma_{ud}/3$$



3 colors



$$\Gamma_{\text{tot}}^{\tau} \simeq \Gamma_{\tau \rightarrow e} + \Gamma_{\tau \rightarrow \mu} + \underbrace{\left(\Gamma_{\text{had}} \right)}_{\sim 3} \simeq 5 \Gamma_{\tau \rightarrow e}$$

$$\Gamma_{\tau \rightarrow e} \simeq \frac{1}{5} \Gamma_{\text{tot}} = \frac{1}{5} \frac{1}{\tau_{\tau}} \quad \tau_{\tau} \quad \tau \text{ lifetime.}$$

measure.

$$\frac{\Gamma(\mu^+ \rightarrow e^+ \bar{\nu}_e \nu_{\mu})}{\Gamma(\tau^- \rightarrow e^- \bar{\nu}_e \nu_{\tau})} = \underbrace{\frac{1}{\tau_{\mu}}}_{\text{measure}} \frac{\text{BR}(\mu^+ \rightarrow e^+ \bar{\nu}_e \nu_{\mu})}{\text{BR}(\tau^- \rightarrow e^- \bar{\nu}_e \nu_{\tau})}$$

$$\frac{\Gamma(\tau^- \rightarrow e^- \bar{\nu}_e \nu_{\tau})}{\Gamma_{\text{tot}}} = \text{BR}(\tau^- \rightarrow e^-) \Rightarrow$$

$$\Gamma(\tau^- \rightarrow e^-) = \Gamma_{\text{tot}}^{\tau} \times \text{BR}(\tau^- \rightarrow e^-) = \frac{1}{\tau_{\tau}} \text{BR}(\tau^- \rightarrow e^-)$$

$$\hookrightarrow \Gamma \simeq \frac{g_{\tau}^2}{M_W^2} \frac{g_e^2}{M_W^2} m_{\tau}^5 \rho_{\tau \rightarrow e}.$$

$$\Rightarrow \frac{g_{\mu}^2}{g_{\tau}^2} \underbrace{\frac{m_{\mu}^5}{m_{\tau}^5}}_{\text{measured.}} \underbrace{\frac{\rho_{\mu \rightarrow e}}{\rho_{\tau \rightarrow e}}}_{\text{computed.}}$$

$$\frac{g_{\mu}}{g_{\tau}} = 1.001 \pm 0.003 \Rightarrow \text{universality of } \tau \leftrightarrow \mu.$$

$\Rightarrow \tau^-$ is a lepton.

$$g_W^e \simeq g_W^{\mu} \simeq g_W^{\tau}$$

Universality with hadronic τ decays.


$$\frac{\Gamma_{\tau \rightarrow \mu}}{\Gamma_{\mu \rightarrow e}} = \frac{m_{\tau}^5}{m_{\mu}^5}.$$

$$\hookrightarrow \Gamma_{\mu \rightarrow e} \simeq \Gamma_{\text{tot}}^{\mu} = \frac{1}{\tau_{\mu}} \quad \Gamma_{\tau \rightarrow \mu} \simeq \frac{1}{5} \Gamma_{\text{tot}}^{\tau} = \frac{1}{5} \frac{1}{\tau_{\tau}}.$$

$$\tau_{\tau} = \frac{1}{5} \frac{m_{\mu}^5}{m_{\tau}^5} \tau_{\mu} = 3.1 \times 10^{-13} \text{ s. from PDG values.}$$



$$\tau_{\tau}^{\text{exp}} = 2.9 \times 10^{-13} \text{ s.}$$

Weak interaction in Quarks



V-A structure not so clear with hadrons.
 \Rightarrow corrections from strong interaction.

$$n \rightarrow p e^- \bar{\nu}_e$$



$$\pi^- \rightarrow \pi^0 e^- \bar{\nu}_e$$

$m \begin{matrix} 139 \\ 135 \end{matrix}$ $Q \approx 3.5 \text{ MeV.}$

π^+ DECAY MODES

π^- modes are charge conjugates of the modes below.


For decay limits to particles which are not established, see the section on Searches for Axions and Other Very Light Bosons.

| Mode | Fraction (Γ_i/Γ) | Confidence level |
|--|--|----------------------|
| $\Gamma_1 \quad \mu^+ \nu_\mu$ | [a] (99.98770 \pm 0.00004) % | |
| $\Gamma_2 \quad \mu^+ \nu_\mu \gamma$ | [b] (2.00 \pm 0.25) $\times 10^{-4}$ | |
| $\Gamma_3 \quad e^+ \nu_e$ | [a] (1.230 \pm 0.004) $\times 10^{-4}$ | |
| $\Gamma_4 \quad e^+ \nu_e \gamma$ | [b] (7.39 \pm 0.05) $\times 10^{-7}$ | |
| $\Gamma_5 \quad e^+ \nu_e \pi^0$ | (1.036 \pm 0.006) $\times 10^{-8}$ | |
| $\Gamma_6 \quad e^+ \nu_e e^+ e^-$ | (3.2 \pm 0.5) $\times 10^{-9}$ | |
| $\Gamma_7 \quad \mu^+ \nu_\mu \nu \bar{\nu}$ | < 9 | $\times 10^{-6}$ 90% |
| $\Gamma_8 \quad e^+ \nu_e \nu \bar{\nu}$ | < 1.6 | $\times 10^{-7}$ 90% |



Lepton Family number (LF) or Lepton number (L) violating modes

| | | | | |
|---------------------------------------|----|-----------|------------------|-----|
| $\Gamma_9 \quad \mu^+ \bar{\nu}_e$ | L | [c] < 1.5 | $\times 10^{-3}$ | 90% |
| $\Gamma_{10} \quad \mu^+ \nu_e$ | LF | [c] < 8.0 | $\times 10^{-3}$ | 90% |
| $\Gamma_{11} \quad \mu^- e^+ e^+ \nu$ | LF | < 1.6 | $\times 10^{-6}$ | 90% |


why no $\pi^- \rightarrow \pi^0 e^- \bar{\nu}_e$ $Q < 0$.



π decays



$$K^- \rightarrow \pi^0 e^- \bar{\nu}_e$$



$$\pi^- \rightarrow \pi^0 e^- \bar{\nu}_e$$

$\Delta S = 0$




$$\Delta S = 1$$

$$K^- \rightarrow \pi^0 e^- \bar{\nu}_e$$


$$\begin{pmatrix} u \\ d \end{pmatrix} \begin{pmatrix} s \\ s \end{pmatrix} \begin{pmatrix} t \\ b \end{pmatrix}$$

charged current W^- charges quark flavor

| K^+ DECAY MODES | Fraction (Γ_i/Γ) | Scale factor/ Confidence level (MeV/c) | p |
|--|------------------------------------|---|-----|
| Leptonic and semileptonic modes | | | |
| $e^+ \nu_e$ | $(1.582 \pm 0.007) \times 10^{-5}$ | | 247 |
| $\mu^+ \nu_\mu$ | $(63.56 \pm 0.11) \%$ | $S=1.2$ | 236 |
| $\pi^0 e^+ \nu_e$ | $(5.07 \pm 0.04) \%$ | $S=2.1$ | 228 |
| Called K_{e3}^+ | | | |
| $\pi^0 \mu^+ \nu_\mu$ | $(3.352 \pm 0.033) \%$ | $S=1.9$ | 215 |
| Called $K_{\mu3}^+$ | | | |
| $\pi^0 \pi^0 e^+ \nu_e$ | $(2.55 \pm 0.04) \times 10^{-5}$ | $S=1.1$ | 206 |
| $\pi^+ \pi^- e^+ \nu_e$ | $(4.247 \pm 0.024) \times 10^{-5}$ | | 203 |
| $\pi^+ \pi^- \mu^+ \nu_\mu$ | $(1.4 \pm 0.9) \times 10^{-5}$ | | 151 |
| $\pi^0 \pi^0 \pi^0 e^+ \nu_e$ | $< 3.5 \times 10^{-6}$ | $CL=90\%$ | 135 |
| Hadronic modes | | | |
| $\pi^+ \pi^0$ | $(20.67 \pm 0.08) \%$ | $S=1.2$ | 205 |
| $\pi^+ \pi^0 \pi^0$ | $(1.760 \pm 0.023) \%$ | $S=1.1$ | 133 |
| $\pi^+ \pi^+ \pi^-$ | $(5.583 \pm 0.024) \%$ | | 125 |



$$\propto G_F^2 \frac{f_\pi^2}{m_\pi^3} m_\mu^2 (m_\pi^2 - m_\mu^2)$$




$$\propto G_F^2 \frac{f_K^2}{m_K^3} m_\mu^2 (m_K^2 - m_\mu^2)$$

$$\frac{\Gamma(K^+ \rightarrow \mu^+ \nu_\mu)}{\Gamma(\pi^+ \rightarrow \mu^+ \nu_\mu)} \sim \frac{G_F^2}{G_F^2} \frac{f_\pi^2}{f_K^2} (\quad)$$

$$\frac{BR(K^+ \rightarrow \mu^+ \nu_\mu)}{BR(\pi^+ \rightarrow \mu^+ \nu_\mu)}$$

$$\frac{G_F^2(K^+ \rightarrow \mu^+ \nu_\mu, \Delta S=1)}{G_F^2(\pi^+ \rightarrow \mu^+ \nu_\mu, \Delta S=0)} = 0.05$$




$$n \rightarrow p e^- \bar{\nu}_e$$

$$\Gamma \propto G_F^2 \rho_{phase}$$

$\Delta S=0$

$$\frac{G_F^2(\Delta S=1)}{G_F^2(\Delta S=0)} \approx 0.05$$

$$(\Delta S=1) \ll \Gamma(\Delta S=0)$$



$$\Delta S=1$$

$$\Sigma^- \rightarrow p e^- \bar{\nu}_e$$

$$\Gamma \propto G_F^2 \rho_{phase \text{ since}}$$



n DECAY MODES

| Mode | Fraction (Γ_i/Γ) | Confidence level |
|--|------------------------------------|------------------|
| Γ_1 $pe^- \bar{\nu}_e$ | 100 % | |
| Γ_2 $pe^- \bar{\nu}_e \gamma$ | [a] $(9.2 \pm 0.7) \times 10^{-3}$ | |
| Γ_3 hydrogen-atom $\bar{\nu}_e$ | < 2.7 $\times 10^{-3}$ | 95% |

Σ^- DECAY MODES


| Mode | Fraction (Γ_i/Γ) | Confidence level |
|--------------------------------------|------------------------------------|------------------|
| Γ_1 $n\pi^-$ | $(99.848 \pm 0.005) \%$ | |
| Γ_2 $n\pi^- \gamma$ | [a] $(4.6 \pm 0.6) \times 10^{-4}$ | |
| Γ_3 $ne^- \bar{\nu}_e$ | $(1.017 \pm 0.034) \times 10^{-3}$ | |
| Γ_4 $n\mu^- \bar{\nu}_\mu$ | $(4.5 \pm 0.4) \times 10^{-4}$ | |
| Γ_5 $\Lambda e^- \bar{\nu}_e$ | $(5.73 \pm 0.27) \times 10^{-5}$ | |
| Γ_6 $\Sigma^+ X$ | < 1.2 $\times 10^{-4}$ | 90% |

Cabibbo's Solution.



$$d' = \cos\theta \cdot d + \sin\theta \cdot s \quad \text{mixture of } d, s.$$

$$s' = -\sin\theta \cdot d + \cos\theta \cdot s.$$



UNITARY SYMMETRY AND LEPTONIC DECAYS

Nicola Cabibbo

CERN, Geneva, Switzerland

(Received 29 April 1963)

We present here an analysis of leptonic decays based on the unitary symmetry for strong interactions, in the version known as "eightfold way,"¹ and the $V-A$ theory for weak interactions.^{2,3} Our basic assumptions on J_μ , the weak current of strong interacting particles, are as follows:

(1) J_μ transforms according to the eightfold representation of SU_3 . This means that we neglect currents with $\Delta S = -\Delta Q$, or $\Delta I = 3/2$, which should belong to other representations. This limits the scope of the analysis, and we are not

able to treat the complex of K^0 leptonic decays, or $\Sigma^+ \rightarrow n + e^+ + \nu$ in which $\Delta S = -\Delta Q$ currents play a role. For the other processes we make the hypothesis that the main contributions come from that part of J_μ which is in the eightfold representation.

(2) The vector part of J_μ is in the same octet as the electromagnetic current. The vector contribution can then be deduced from the electromagnetic properties of strong interacting particles. For $\Delta S = 0$, this assumption is equivalent to vector-

To determine θ , let us compare the rates for $K^+ \rightarrow \mu^+ + \nu$ and $\pi^+ \rightarrow \mu^+ + \nu$; we find

$$\Gamma(K^+ \rightarrow \mu\nu) / \Gamma(\pi^+ \rightarrow \mu\nu)$$

$$= \tan^2\theta \frac{M_K (1 - M_\mu^2/M_K^2)^2}{M_\pi (1 - M_\mu^2/M_\pi^2)^2}. \quad (3)$$

From the experimental data, we then get^{5,6}

$$\theta = 0.257. \quad (4)$$

For an independent determination of θ , let us consider $K^+ \rightarrow \pi^0 + e^+ + \nu$. The matrix element for this process can be connected to that for $\pi^+ \rightarrow \pi^0 + e^+ + \nu$, known from the conserved vector-current hypothesis (2nd assumption). From the rate⁶ for $K^+ \rightarrow \pi^0 + e^+ + \nu$, we get

$$\theta = 0.26. \quad (5)$$

The two determinations coincide within experimental errors; in the following we use $\theta = 0.26$.

We go now to the leptonic decays of the baryons, of the type $A \rightarrow B + e + \nu$. The matrix element of

$$\pi^+ \rightarrow \mu^+ \nu \propto G_F^2 \cos^2\theta \cdot m_\mu^2 (m_\pi^2 - m_\mu^2)$$


$$K^+ \rightarrow \mu^+ \nu \propto G_F^2 \sin^2\theta \cdot m_\mu^2 (m_K^2 - m_\mu^2)$$

$$u \rightarrow d \quad g \cos\theta$$

$$u \rightarrow s \quad g \sin\theta$$

$$\frac{\Gamma(K^+ \rightarrow \mu^+ \nu_\mu)}{\Gamma(K^+ \rightarrow \mu^+ \nu_\mu)} \approx \tan^2 \theta \quad \left(\leftarrow - \right)$$

$$\theta \approx 13^\circ$$



$$\cos^2 \theta = 0.97$$

$$\sin^2 \theta = 0.03$$

$$\tan^2 \theta \approx 0.05$$

$$\Gamma(\Delta S = 1) \propto \sin^2 \theta.$$




$$G_{F_K}^2 \approx \left(\frac{g_W^2}{M_W^2} \right)^2 \sin^2 \theta. \quad \Delta S = 1.$$

$$G_{F_\pi}^2 \approx \left(\frac{g_W^2}{M_W^2} \right)^2 \cos^2 \theta. \quad \Delta S = 0.$$

Flavor eigenstates \neq weak eigenstates
(d, s) (d', s').

$$\Gamma(K^+ \rightarrow \mu^+ \nu_\mu) = 64\%$$

$$\Gamma(K^0 \rightarrow \mu^+ \mu^-) = 7 \times 10^{-9}$$




W^0 : neutral current
flavor changing.

not allowed.

| K ⁺ DECAY MODES | Scale factor/ Confidence level (MeV/c) | | p |
|--|---|--------|-----|
| | Fraction (Γ _i /Γ) | | |
| Leptonic and semileptonic modes | | | |
| e ⁺ ν _e | (1.582±0.007) × 10 ^{−5} | | 247 |
| μ ⁺ ν _μ | (63.56 ±0.11) % | S=1.2 | 236 |
| π ⁰ e ⁺ ν _e | (5.07 ±0.04) % | S=2.1 | 228 |
| Called K ⁺ _{e3} . | | | |
| π ⁰ μ ⁺ ν _μ | (3.352±0.033) % | S=1.9 | 215 |
| Called K ⁺ _{μ3} . | | | |
| π ⁰ π ⁰ e ⁺ ν _e | (2.55 ±0.04) × 10 ^{−5} | S=1.1 | 206 |
| π ⁺ π [−] e ⁺ ν _e | (4.247±0.024) × 10 ^{−5} | | 203 |
| π ⁺ π [−] μ ⁺ ν _μ | (1.4 ±0.9) × 10 ^{−5} | | 151 |
| π ⁰ π ⁰ π ⁰ e ⁺ ν _e | < 3.5 × 10 ^{−6} | CL=90% | 135 |
| Hadronic modes | | | |
| π ⁺ π ⁰ | (20.67 ±0.08) % | S=1.2 | 205 |
| π ⁺ π ⁰ π ⁰ | (1.760±0.023) % | S=1.1 | 133 |
| π ⁺ π ⁺ π [−] | (5.583±0.024) % | | 125 |

if w^0 exists.



$$K^+ \rightarrow \pi^+ l^+ l^-$$