

BAAZ & UDAY JKBOSE

2026 CLASS 11TH



ANIMAL KINGDOM

LECTURE-4

ZOOLOGY


DAVOOD SIR (CEO of NG)




Topics To Be Covered

- 6 **Phylum Platyhelminthes**
- 7 **Phylum Aschelminthes**
- 8 **Previous Year NEET MCQ's**

Phylum Platyhelminthes (The Flatworms)



Phylum Platyhelminthes (The Flatworms)




Phylum Platyhelminthes (The Flatworms)

↳ Named for their dorso-ventrally flattened body (leaf like).

- ① Habitat: → Mostly Endoparasites (present in Animal body)
- ② Body structure: → Bilateral symmetry, Triploblastic, organ level of organisation & Acoelomates.
- ③ Parasitic Adaption: → Hook & Sucker aid in attachment to hosts body.
- ④ Digestive system: → those platyhelminthes having digestive system are called flukes (incomplete digestive system)
those platyhelminthes having no mouth & digestive system are called → Tapeworms.
- ⑤ Excretion: → Flame cells / protonephridia helps in excretion and osmoregulation.
- ⑥ Reproduction: → they are hermaphrodites (Bisexual), with internal fertilization.
Fertilization is indirect (Larval stage)
• Some members (Planaria) show asexual Reproduction → Regeneration.
- ⑦ Examples: → Taenia (Tapeworm), Fasciola [liver fluke].

↓
Absorbs digested food from hosts body through their body surface.

Phylum Aschelminthes (The Roundworms)



Phylum Aschelminthes (The Roundworms)

- ↳ Body is **circular** in cross section due to circular muscles → that is why we call them Roundworms.
- ① Habitat → Free living Aquatic, Terrestrial or Endoparasites of Animal or plant body.
 - ② Body Structure → Bilateral symmetry, Triploblastic, Pseudocoelomates, having organ system level of organisation.
 - ③ Digestive System → **Digestive System** is complete (Mouth and Anus) with muscular pharynx.
 - ④ Excretion → Metabolic wastes from body are removed through **Excretory pore of Excretory tube**.
 - ⑤ Reproduction → Sexes are separate (sexual dimorphism), **Male are shorter** and having copulatory setae whereas **females are longer**.
Fertilization is internal and development is direct or indirect.
 - ⑥ Examples → Ascaris (Roundworms).
Wuchereria (Filaria worm).
Ancylostoma (Hookworm).

Kaun Banayga Nabkaar: NEET-2024

Consider the following statements.

- A. Annelids are true coelomates. *T.*
- B. Poriferans are pseudocoelomates. *→ F*
- C. Aschelminths are acoelomates. *F*
- D. Platyhelminthes are pseudocoelomates. *F*

Choose the correct sequence of stages from the options given below.

- A** B Only
- B** A Only. *✓*
- C** C Only
- D** D Only



Kaun Banayga Nabkaar : NEET-2023

Select the correct statements.

- ☒ (A) Platyhelminthes are triploblastic pseudocoelomate and bilaterally symmetrical organisms.
- ☒ (B) Ctenophores reproduced only sexually and fertilisation is external.
- ☒ (C) In tapeworm, fertilisation is internal but sexes are not separate.
- ☒ (D) Ctenophores are exclusively marine diploblastic and bioluminescent organisms.
- ☒ (E) In sponges, fertilisation is external and development direct.

Choose the correct answer from the option given below.

A (A), (C) and (D) only

☒ **B** (B), (C) and (D) only.

C (A) and (E) only

D (B) and (D) only



Thank
You