

# Dynamical Many Body Localization in Periodically Driven Long-Range Spin Systems: An Eccentric Phase Crossover

Mahbub Rahaman,<sup>1</sup> Takashi Mori,<sup>2</sup> and Analabha Roy<sup>1</sup>

<sup>1</sup>*Department of Physics, The University of Burdwan, Golapbag, Bardhaman - 713 104, India*

<sup>2</sup>*RIKEN CEMS, 2-1 Hirosawa, Wako, Saitama, 351-0198, Japan*

The phenomenon of many-body localization occurs in quantum systems under specific resonance conditions, preventing thermalization and allowing the system to maintain its initial state. Quantum dynamical localization has been validated for multi-body spin systems under the influence of transverse field. The validation has been parameterized by the Inverse Participation ratio (IPR) of Floquet modes. Power-law dependence with long-range. The study applied  $J_{ij} = 1/|i-j|^\beta$  and examined the short-range Transverse Field Ising Model for  $\beta = \infty$  and the long-range Lipkin Meshkov-Glick (LMG) model for  $\beta = 0$  at varying drive frequencies. The Ising model displays complete localization at high frequencies, while localization persists at low frequencies at the localization point. At the resonance point of system localization, LMG exhibits high-frequency localization in the long-range scenario. However, at lower frequencies, localization deteriorates following an inverse system size law. Chaos and regularity alternate at different frequencies in LMG at the phase-space continuum limit. IPR for LMG is evolved by an adiabatic increase in drive frequencies; the IPR rises gradually up to unity thereby manifesting a phase crossover which proposes a future protocol for the MBL engine.

Keywords: Dynamical localization, Thermalization, Phase Crossover

Periodically driven Quantum Many Body Systems can experience Dynamical Freezing (DMF) when dynamical hysteresis stops observables from reaching their diagonal averaged values and thermalizing to infinite temperature [1–3]. Under certain resonance conditions in the drive parameters, DMF can cause the response to ‘freeze’ completely to its initial value at all times. This arises as a consequence of additional approximate symmetries that occur at resonance. DMF has been demonstrated via the Rotating Wave Approximation (RWA) in the driven TFIM with nearest neighbour interactions [4] and is shown to be protected when translational invariance is explicitly broken (say, by disorder) [5, 6].

The utilization of Floquet theory simplifies the analysis of time-periodic systems. For closed quantum systems governed by the time-dependent Schrödinger equation, the *Floquet Hamiltonian* allows for a mapping of the time-dependent dynamics into the dynamics of a time-independent effective Hamiltonian, provided the system is strobed at integer multiples of the time period of the drive. The time independent eigenstates of the effective Hamiltonian correspond to quasi-stationary *Floquet Modes* of the original Hamiltonian. The temporal progression of the system comes from phase coefficients that capture the dynamics [7, 8].

Any sufficiently complex non-integrable Many Body System is expected to thermalize according to the Eigenstate Thermalization Hypothesis (ETH) despite the fact that closed quantum dynamics preserves the memory of the initial state of the system. This arises due to the properties of the matrix elements of ob-

servables in typical states[9]. The ETH can be readily adapted to time-periodic systems using Floquet theory (the Floquet-ETH, or FETH). Nonetheless, the conditions for ETH to hold are not particularly strong, and the density matrix of the system can fail to approach one that is described by a thermal expression. In such cases, the system is said to undergo *Many Body Localization* (MBL)[10]. This phenomenon is stable against local perturbations, and constitutes an exotic state of matter with far-reaching implications in theoretical physics, as well as in practical applications[11].

The addition of disorder has been identified as a crucial component in the onset of MBL. In that case, thermalization is prevented by disorder-induced localization. An alternative approach to realizing MBL in clean many-body system involves *Floquet Engineering*, where a time-periodic drive is introduced, and the drive parameters tuned so as to introduce a clustering of quasistationary energies in a manner similar to localization[9].

In this article, we propose that additional approximate symmetries can be Floquet-engineered in quantum many body systems with lower symmetry than the TFIM, such as those with long-range interactions. This results in both DMF and MBL occurring simultaneously at resonant values of the drive parameters, and complete thermal behaviour at other values. This phenomenon is distinct from DMF in the TFIM, since clean TFIM systems, being integrable, never thermalize.

To demonstrate the onset of MBL, we investigate the driven Lipkin-Meshkov-Glick (LMG) model, a long-range system that is a special case of the more general

Curie-Weiss model, wherein the nearest-neighbour exchange in the TFIM is extended to longer ranges with a power law dependence,  $J_{ij} \sim 1/|i - j|^\beta$  [12–14]. Setting  $\beta = \infty$  recovers the TFIM, and setting  $\beta = 0$  yields the LMG model. We have recovered the onset of DMF in this system and have supported our result with numerical simulations.

In addition, we compare the degree of localization of the quasi-stationary Floquet modes in both limits of  $\beta$ . In order to do so, we look at the Inverse Participation Ratio (IPR) of the Floquet modes in the representation given by the eigenstates of the symmetry-breaking field. The IPR, closely related to the concept of quantum purity, is defined as the formal sum of the square of the density in some physically meaningful space or representation. A high IPR of a stationary state denotes low participation in most of the representation, and a low IPR distributes participation uniformly across the representation, leading to ergodic dynamics[15]. Thus, IPR [16] is a useful tool for witnessing MBL of a quantum system. For an MBL system, the IPR is unity, and it scales inversely with the system size when it is thermally distributed [17].

In the first section of this paper, we present all essential theoretical frameworks. Our results for the LMG model are presented next in section II. In that section, we have used the Rotating Wave Approximation (RWA) [18], where only the slowest rotating terms in the Fourier expansion of the Hamiltonian in a frame co-rotating with the symmetry breaking drive field are retained. In addition, we have the obtained analytical expressions for the Floquet modes and their IPR. They are used to probe the system dynamics in the high and low-frequency domains at both limits of  $\beta$ . In section III we have used phase space plots to contrast the low and high frequency limits of the LMG model in the thermodynamic limit by mapping it to an equivalent classical Hamiltonian system. Finally, in section IV, we have looked at numerical computations of the IPR of the Floquet modes for different values of the drive parameters, well beyond those that allow for the RWA. We observed that, if the system is driven by an adiabatically increasing drive frequency from low to high limit while remaining in the resonance region, a sharp crossover from a thermal to an MBL phase occurs. We conclude with discussions and outlook.

## I. BACKGROUND

The Eigenstate Thermalization Hypothesis (ETH) is a series of conjectures that allows for the thermalization of an isolated quantum many body system. The state of the system,  $|\psi(t)\rangle$ , evolves according to the Schrödinger equation  $\hat{H}|\psi(t)\rangle = i\frac{\partial}{\partial t}|\psi\rangle$ . The Hamiltonian  $\hat{H}$  is assumed to be *non-integrable*, in that it lacks an extensive number of *local* additive conserved quantities, that is to say, there are no set of observables  $\hat{O}_s$  such that  $\hat{H} = \sum_s \hat{O}_s$  for any extensive index  $s$ . Here, the  $\hat{O}_s$  constitute an arbitrary CSCO (complete set of commuting observables) that are *local*, having sub-extensive support in the system size. In addition, we postulate the existence of an equivalent Hamiltonian  $\hat{H}_{eq}$  for every Hamiltonian  $\hat{H}$  as well as an "equilibrium" value  $A_{eq}$  for every observable  $\hat{A}$ , such that

$$A_{eq}(E) \equiv \frac{\text{Tr}(\hat{A}e^{-\beta\hat{H}_{eq}})}{\text{Tr}(e^{-\beta\hat{H}_{eq}})}. \quad (1)$$

where  $E = \langle\psi(t)|\hat{H}|\psi(t)\rangle$  is the conserved energy of the system, and  $\beta = 1/(k_B T)$  is the inverse temperature,  $H_{eq}$  is an effective Hamiltonian that captures the long-time average dynamics of the system, and  $k_B$  is the Boltzmann constant.

To put it simply, ETH proposes that this many-body Hamiltonian undergoes thermalization as seen in the *long-time averages* of observables, with the eigenstates bearing resemblance to thermal states. The aforementioned hypothesis serves as a valuable instrument for comprehending the conduct of stimulated quantum systems and their correlation with thermal equilibrium. This assertion can be justified by examining the expectation value of an observable  $\hat{A}$  as it evolves under the Schrödinger equation. To see this, we first expand the state of the system  $|\psi(t)\rangle$  as:

$$|\psi(t)\rangle = \sum_m c_m(t) |m(0)\rangle,$$

where  $|m(0)\rangle$  represents the eigenstates of  $\hat{H}(0)$  with energy  $E_m$ . The coefficients  $c_m(t)$  describe the time-dependent amplitude of the expansion. Plugging these expansions into the expression for the expectation value, we obtain the long-time average of the expectation value [19]:

$$\overline{\langle\hat{A}(t)\rangle} = \sum_{m,k} \overline{c_m^*(t)c_k(t)} \langle m(0)|\hat{A}|k(0)\rangle, \quad (2)$$

Had the system been integrable, the large number of conserved quantities would restrict mixing between the states during unitary evolution. In the non-integrable case, the system explores the entire Hilbert space spanned by eigenstates with eigenvalues close to  $E$  more-or-less uniformly. In that case, the matrix elements  $\langle m(0)|\hat{A}|k(0)\rangle$  are said to satisfy the Srednicki ansatz [20, 21]:

$$\langle m(0)|\hat{A}|k(0)\rangle \approx A_{eq}\delta_{mk} + e^{-\frac{1}{2}S\left(\frac{E_m+E_k}{2}\right)} f\left(\frac{E_m+E_k}{2}, E_m - E_k\right) R_{mk}. \quad (3)$$

Here,  $S$  is the thermodynamic entropy and  $R_{mk}$  are elements of a random matrix with vanishing mean and unit variance. What this means for the ensuing dynamics is that the system explores the accessible Hilbert space uniformly, and the matrix elements  $\langle m(0) | \hat{A}(t) | k(0) \rangle$  become indistinguishable for most pairs of  $m$  and  $k$ . Applying this ansatz and taking the thermodynamic limit by ignoring terms  $\mathcal{O}(e^{-S/2})$ , the expression for the expectation value becomes:

$$\overline{\langle \hat{A}(t) \rangle} \approx \sum_m \overline{|c_m(t)|^2} A_{eq} = A_{eq},$$

Therefore, in the limit of large systems the expectation value of an observable  $\hat{A}$  is approximately equal to the thermal expectation value  $A_{eq}$ . This is the essence of the ETH, which suggests that individual eigenstates of a quantum system can be described by statistical mechanics in the long-time limit.

We now generalize the ETH to non-integrable many body systems that are closed, but not isolated. In that case, it is possible to impart a periodic time-dependence on the Hamiltonian while still ensuring unitary evolution. If the time period of the drive is  $T$ , and the corresponding drive frequency  $\omega \equiv 2\pi/T$ , the Floquet theorem states that the solutions to the Schrödinger equation can be written as  $|\psi(t)\rangle = e^{-i\epsilon t/\hbar} |\phi(t)\rangle$ , where the  $|\phi(t)\rangle$  are  $T$ -periodic states called *Floquet Modes*, the corresponding  $\epsilon \in \mathbb{R}$ , are called *quasienergies*. Quasienergy values are not unique, and can be made to be bounded within a Floquet photon, viz. a range  $[-\omega/2, \omega/2]$  [22, 23]. As a consequence, the unitary evolution operator can be split into two parts as follows [24].

$$U(t) = e^{-i\hat{K}_F(t)} e^{-i\hat{H}_F t}. \quad (4)$$

Here, the micromotion operator  $\hat{K}_F(t)$  is time-periodic in  $T$ , with  $\hat{K}_F(0) = 0$ , and the Floquet Hamiltonian  $\hat{H}_F = \hat{H}(t) - i \frac{\partial}{\partial t} \Big|_{t=T}$ . Thus, if the system is strobed at integer multiples of  $T$  only, then the unitary evolution matches that of a time independent Hamiltonian  $H_F$ . This can capture a representative cross-section of the exact dynamics at large frequencies.

In such systems, the Floquet Eigenstate Thermalization Hypothesis (FETH) posits that subject to specific conditions and in the context of a system of significant size, the Floquet modes themselves exhibit thermal state-like behavior, i.e.,  $\hat{H}_{eq} \approx \hat{H}_F$  in eqn 1. However, in contrast to the isolated systems, the loss of energy conservation allows for the mixing of all Floquet modes in the ensuing dynamics, not just those with quasienergies near  $E$ . Were this to actually happen in the ensuing dynamics, it can be reconciled with ETH [25] by ensuring that  $\beta = 0$  in eq 1. In other words, the nonequilibrium steady state of the system

tends to an infinite temperature, maximum entropy density matrix.

However, drive parameters like amplitude, frequency, and duty-cycle strongly affect the structure of the Floquet modes  $|\phi\rangle$ . Thus, they can be engineered to prevent the kind of full mixing that would lead to infinite temperatures, manifesting suppression of thermalization dynamically. Thus, this type of *Floquet Engineering* can produce *Dynamical Many Body Localization* (DMBL), where the system fails to reach thermal equilibrium and remains localised, possibly near its initial state, even at large times. This paradigm seems similar to standard Many-Body Localization [26, 27], where disorder, locality, and integrability can cause athermality via breakdown in the Srednicki ansatz. However, DMBL is a purely dynamical phenomenon, and thus can occur regardless of disorder, locality of observables, or system integrability, all of which have been studied for MBL onset [28–30].

Integrable Many Body systems do not exhibit thermalization. When subjected to time-periodic drives, Floquet engineering allows for the introduction of additional approximate conserved quantities that dynamically suppress the evolution of certain observables by hysteresis. This type of *freezing* of response has been shown in integrable systems [6]. A paradigmatic example is the driven Transverse Field Ising model (TFIM) in one dimension [31]. The Hamiltonian is given by

$$\hat{H}(t) = \hat{H}_0 + h_z(t) \hat{H}_1 \quad (5)$$

$$\hat{H}_0 = -\frac{1}{2} \sum_{i=1}^N \sigma_i^x \sigma_{i+1}^x \quad (6)$$

$$\hat{H}_1 = -\frac{1}{2} \sum_{i=1}^N \sigma_i^z. \quad (7)$$

Here, the undriven Hamiltonian  $\hat{H}_0$  consists of nearest-neighbour interactions between  $N$  number of spin-1/2 particles on a one-dimensional spin network. The transverse field is denoted by  $\hat{H}_1$ , and is being varied by a time-periodic and harmonic signal  $h_z(t) = h_0 + h \cos \omega t$ , yielding a time period  $T = 2\pi/\omega$  with amplitude  $h$ , drive frequency  $\omega$ , and d.c. field  $h_0$ . This Hamiltonian can be readily transformed into a spinless pseudo-fermionic system via the Jordan-Wigner transformation [4]. When written in momentum space spanned by spinors  $\psi_k = (c_{-k}, c_k^\dagger)^T$  of fermions at momentum  $k$  created (annihilated) by operators  $c_k^\dagger$  ( $c_k$ ), the effective Hamiltonian

$$H(t) = \sum_{(k, -k)-\text{pairs}} \psi_k^\dagger \left[ \left( f_k - h_z(t) \right) \tau_z + \tau_x \Delta_k \right] \psi_k \quad (8)$$

with  $f_k = J \cos k$ ,  $\Delta_k = J \sin k$ ,  $\tau_{xyz}$  are the three Pauli Matrices, and the sum is over distinct

( $k, -k$ ) *Cooper Pairs*. A transformation to the rotated frame is achieved by means of the unitary transformation operator  $U(t) = \prod_k U_k(t)$ , where  $U_k(t) = \exp\left\{\left[\frac{ih}{\omega} \sin \omega t\right] \tau_z\right\}$ . The resulting transformed Hamiltonian is

$$H'(t) = \sum_{(k, -k)-\text{pairs}} \psi_k^\dagger \left[ \tau_z f_k + \tau_x \cos(\eta \sin \omega t) + \tau_y \cos(\eta \sin \omega t) \right] \psi_k,$$

where we defined  $\eta = 2h/\omega$ . Using the Jacobi-Anger formula [32]

$$e^{i\eta \sin \omega t} = \sum_{n=-\infty}^{\infty} J_n(\eta) e^{in\omega t}, \quad (9)$$

where  $J_n(\eta)$  are Bessel Functions, the transformed Hamiltonian simplifies to

$$H'(t) = \sum_{\substack{(k, -k) \\ \text{pairs}}} \psi_k^\dagger \left\{ \tau_z f_k + 2\tau_x \Delta_k \sum_{n \geq 0} J_{2n}(\eta) \cos(2n\omega t) - 2\tau_y \Delta_k \sum_{n \geq 0} J_{2n+1}(\eta) \sin[(2n+1)\omega t] \right\} \psi_k, \quad (10)$$

In the frequency regime  $\omega \gg f_k$ , the long-time average  $H^{RWA} \equiv \lim_{n \rightarrow \infty} \frac{1}{nT} \int_0^{nT} dt H'(t)$  can serve as a suitable approximation for  $H'(t)$ . This approximation, known as the *Rotated Wave Approximation* (RWA), eliminates the oscillating modes and results in an effective Hamiltonian that is independent of time.

$$H^{RWA} = \sum_{(k, -k)-\text{pairs}} \psi_k^\dagger \left[ f_k \tau_z + 2J_0(\eta) \Delta_k \tau_x \right] \psi_k, \quad (11)$$

It is evident that by manipulating the drive parameters, specifically the amplitude denoted by  $h$  and the frequency denoted by  $\omega$ , in a manner such that  $\eta$  is positioned on a root of  $J_0(\eta)$ , the fermion number can be conserved to a significant extent at this particular resonance. Consequently, it is feasible to exercise direct control over  $H^{RWA}$ , resulting in a comprehensive suppression of the dynamics of otherwise responsive observables.

This phenomenon is highly general in nature, and can be readily adapted to non-integrable systems. In such cases, freezing has the additional effect of inducing DMBL, suppressing thermalization to infinite temperatures. Numerical quantification of localization of a specific (quasi) stationary state in a physically significant representation can be achieved through the

computation of the *Inverse Participation Ratio* (IPR). In the position representation, the IPR for a state  $|\psi\rangle$  [33–36] is defined as

$$\phi_{IPR} \equiv \int d\mathbf{x} |\langle \mathbf{x} | \psi \rangle|^4$$

This definition can be generalized to the IPR of a state  $|\phi\rangle$  in a representation given by complete orthonormal basis  $|m\rangle$  as

$$\phi_{IPR} \equiv \sum_m |\langle m | \psi \rangle|^4. \quad (12)$$

The smallest value of the IPR corresponds to a fully delocalized state,  $\psi(x) = 1/\sqrt{N}$  for a system of size  $N$  [36, 37]. Values of the IPR close to unity correspond to localized states [38]. For a periodically driven system, we look at the IPR of the quasi-stationary Floquet modes at  $t = T$ , where  $t = 2\pi/\omega$  for drive frequency  $\omega$ . In the TFIM model, equation 11 indicates that, at resonance, when  $J_0(\eta) = 0$ , the Floquet modes are approximately given by the fermionic Fock states, which have a trivially unit IPR in the representation of the eigenmodes of the transverse field  $\hat{H}_1$  in equation 5. Here, a particular Floquet mode can be decomposed into a direct product of cooper-pair states as  $|\phi\rangle = \prod_{k, -k} |\phi_k^n\rangle$ . In the RWA limit and at resonance,  $|\phi_k^n\rangle$  has values of  $|0\rangle, |k, -k\rangle$  for two values of  $n = 0, 1$  respectively. We define the reduced IPR of  $|\phi_k^n\rangle \forall k$  to be

$$\phi_{IPR}^{(n)}(k) = |\langle 0 | \phi_k^n \rangle|^4 + |\langle +k, -k | \phi_k^n \rangle|^4, \quad (13)$$

where  $n = 0, 1$ . In the RWA limit and at resonance, this quantity is unity, indicating very low participation and the onset of freezing. Figure 1 shows results from numerically simulating the TFIM dynamics. The reduced IPR for a particular Floquet mode recovered by simulating the exact Schrödinger dynamics over a single time period of the drive, and plotted as a function of momentum  $k$  for different  $\eta$ 's. At resonance, when  $\eta$  lies at the root of the Bessel function  $J_0(\eta)$ , the IPR is exactly unity for all momenta. Outside this resonance, the IPR is unity only for some momenta because the effective Hamiltonian is perfect diagonal at  $k \in \{-\pi, 0, \pi\}$  as can be seen in the cross-sectional plots of figure 1. As we move away from the resonance point, IPR reduces from unity. However, as the TFIM is an integrable spin model, the IPR never drops to a value that is small enough to indicate thermalization. At low frequencies, RWA fails due to the unavailability of zero off-diagonal terms in the effective transformed Hamiltonian, as well as the absence of integrability breaking terms to counteract the off diagonal terms. Consequently, the IPR remains quite high ( $\sim 0.5$ ) even at the resonance point as can be seen figure ???. At low frequency, this is valid for all momentum and parameter  $\eta$ , see figure 2.




FIG. 1. Reduced IPR (defined in equation 13) for one of the two Floquet modes obtained from the exact dynamics of the TFIM for size  $N = 100$  and  $\omega = 90$  for the entire Brillouin zone (top panel, ordinate) and a few drive amplitudes (top panel, abscissa). The dashed lines indicate the roots of  $J_0(\eta)$ . The bottom panel shows cross-sections for four different chosen amplitudes.

Because the dependence of observable expectations on the eigenstates is always fairly strong for integrable systems like the TFIM, such systems will never exhibit any kind of thermal behaviour unless integrability breaking terms (such as strong disorder) are included [6]. As a result, it is not physically meaningful to refer to the unit IPR region as "*Many Body Localization*", because the parameter space lacks a thermalized region to contrast with this state. The type of Floquet Engineering described above, on the other hand, can be easily applied to a broad class of nonintegrable systems where FETH is expected to hold in certain regions. Long-range spin systems, in particular, where the exchange energies between far-off spins are taken into account in the model Hamiltonian, are good candidates because they are known to thermalize when




FIG. 2. IPR for TFIM is approximately 0.5 at lower frequency and 1 at higher frequency. The slow change in IPR from minimum to unity over a large frequency range indicates the absence of a mobility edge and, consequently, the absence of a phase transition. In addition, at the point of resonance at low frequency the system is localized yet not at fully freezing state, where at high frequency it is localized to a fully frozen. A low-frequency regime, at frequency  $\omega = 2.0$  plot for TFIM at four different density points for system size  $N = 500$ . IPR localization persists nonetheless.

driven with low frequencies [39].

## II. LIPKIN MESHKOV GLICK MODEL: LONG RANGE INTERACTION

The periodically driven Curie-Weiss model for  $N$  long-range spins is described by the Hamiltonian

$$\hat{H}(t) = \hat{H}_0 + [h \cos(\omega t) + h_0] \hat{H}_1. \quad (14)$$

Here, the undriven part  $\hat{H}_0$  and the driven part  $\hat{H}_1$  are, respectively,

$$\begin{aligned}\hat{H}_0 &= \frac{1}{2} \sum_{ij} J_{ij} \hat{\sigma}_i^z \hat{\sigma}_j^z, \\ \hat{H}_1 &= \sum_{i=1}^N \hat{\sigma}_i^x.\end{aligned}\quad (15)$$

The Heisenberg exchange energy of the bond between spins  $i$  and  $j$  is given by

$$J_{ij} = \frac{J_\alpha}{N^{1-\alpha}} \frac{1}{r_{ij}^\alpha}, \quad (16)$$

with  $r_{ij}$  representing the smallest graph distance between them. Putting  $\alpha = 0$  yields the Lipkin Meshkov Glick (LMG) model with all-to-all interaction, yielding  $J_{ij} = J_0/N$ . We choose to maintain the extensivity of the interaction energy by enforcing the condition

$$\frac{J_0}{N} \sum_{i \neq j} 1 = \frac{J_0}{N} \frac{N(N-1)}{2} = 1,$$

yielding the Kac-norm  $J_0 = 2/(N-1)$ . The Hamiltonian equation 14 commutes with  $P_{ij} \equiv \frac{1}{2}(1 + \vec{\sigma}_i \cdot \vec{\sigma}_j)$ . In addition, the conservation of total angular momentum  $S^2 = |\vec{S}|^2$ , with  $\vec{S} = S^x \hat{x} + S^y \hat{y} + S^z \hat{z} \equiv \frac{1}{2} \sum_i \vec{\sigma}_i$  is assured by the identity  $[S^2, H(t)] = 0$ . We now choose to populate the system in a state with  $S^2 = \frac{N}{2} \left( \frac{N}{2} + 1 \right)$ . In that case, the dynamics remains invariant in the  $N+1$ -dimensional space spanned by the common eigen states of  $P_{ij}$ ,  $|S|^2$  and  $S_z$ ; these are also eigenstates in the so-called TSS subspace [40]. Let the eigenvalues be  $s_n$ , and the eigen vectors be  $|s_n\rangle$ , where  $s_n = -\frac{1}{2} + \frac{n}{N}$  and the index  $n = 0(1)N$  has  $N+1$  values. The dynamics is restricted to this invariant subspace, wherein the matrix elements of the Hamiltonian are given by

$$\begin{aligned}(H_0)_{ij} &= -\frac{4}{N-1} s_i^2 \delta_{ij}, \\ (H_1)_{ij} &= \left[ \sqrt{\frac{N}{2} \left( \frac{N}{2} + 1 \right) - N s_i (N s_{i+1})} \delta_{i+1,j} \right. \\ &\quad \left. + \sqrt{\frac{N}{2} \left( \frac{N}{2} + 1 \right) - N s_i (N s_{i-1})} \delta_{i-1,j} \right] \quad (17)\end{aligned}$$

Now we transform the Hamiltonian to the Rotated frame given by the transformation

$$\hat{U}(t) = \exp \left[ i \frac{h}{\omega} \sin(\omega t) \hat{H}_1 \right] \quad (18)$$

Defining  $\tau = \frac{h}{\omega} \sin \omega t$ , we use the fact that  $\hat{H}_1 = 2S^x$ , and the following identity obtained using the Baker-Campbell-Hausdorff formula,

$$e^{i2\tau \hat{S}^x} \hat{S}^z e^{-i2\tau \hat{S}^x} = \hat{S}^z \cos(2\tau) + \hat{S}^y \sin(2\tau), \quad (19)$$

to simplify the transformed Hamiltonian  $\tilde{H} = \hat{U} \hat{H} \hat{U}^\dagger - \hat{U}^\dagger \partial_t \hat{U}$ , yielding

$$\begin{aligned}\tilde{H}(t) &= -\frac{1}{N-1} \left[ (S^z)^2 (1 + \cos 4\tau) + (S^y)^2 (1 - \cos 4\tau) \right. \\ &\quad \left. + \{S^y, S^z\} \sin 4\tau \right] - 2h_0 S^x \quad (20)\end{aligned}$$

We define  $\eta \equiv 4h/\omega$  and use the Jacobi-Anger formula in eqn 9 to simplify the expression for  $\tilde{H}(t)$ , then assume that  $\omega$  is large enough to smooth out the harmonic components present in the Hamiltonian. Ignoring constant terms, this yields

$$\tilde{H}_{\text{RWA}} = \frac{(\hat{S}^x)^2}{N-1} - 2h_0 \hat{S}^x - \frac{J_0(\eta)}{N-1} \left[ (\hat{S}^z)^2 - (\hat{S}^y)^2 \right] \quad (21)$$

Proof  
from  
here

If the drive amplitude  $h$  is adjusted such that  $\eta$  lies at a root of  $J_0(\eta)$  (the localization point), the RWA Hamiltonian is diagonal in the representation of the transverse field  $\hat{S}^x$ , yielding an IPR of unity in that representation, similar to the Ising case. Note however, that if the DC transverse field  $h_0$  is set to 0, then, at the localization point, the RWA Hamiltonian  $\tilde{H}_{\text{RWA}} \sim (\hat{S}^x)^2$ , each of whose eigenvalues (given by  $(\frac{N}{2}-m)^2$  where  $m \in 0(1)N$ , and  $(\frac{N}{2}-m)$  are the eigenvalues of  $\hat{S}^x$ ) are two-fold degenerate. This produces infinitely many (Floquet) eigenmodes in the degenerate subspace whose IPRs may not always be unity in the  $S^x$  representation. Thus, the absence of the DC field may produce delocalization in the Floquet states even at the localization points, and this necessitated the inclusion of a DC field  $h_0$  in order to break the symmetry. Finally, note that not all values of the DC field  $h_0$  remove all degeneracies in  $\tilde{H}_{\text{RWA}}$ . To see this, note that, at the localization point, the eigenvalues of  $\tilde{H}_{\text{RWA}}$  are given by

$$\text{Eigs} \left[ \tilde{H}_{\text{RWA}} \right] = \frac{\left( \frac{N}{2} - m \right)^2}{N-1} - 2h_0 \left( \frac{N}{2} - m \right) \quad (22)$$

In order to ensure that no degeneracies occur, we have to adjust  $h_0$  to ensure that for any two integers  $m_1, m_2 \leq N$  the condition following is always met,

$$\frac{\left( \frac{N}{2} - m_1 \right)^2}{N-1} - 2h_0 \left( \frac{N}{2} - m_1 \right) \neq \frac{\left( \frac{N}{2} - m_2 \right)^2}{N-1} - 2h_0 \left( \frac{N}{2} - m_2 \right) \quad (23)$$

If  $N \gg 1$  (substantially large), then this condition can be met by assuring that  $(1 - 2h_0)^{-1}$  is never an integer that is divisible by  $N$ . To ensure this in our numerical simulations we have kept  $h_0$  at a small irrational value.

This result is supported by exact numerical results, as can be seen in the plots below in FIG. 3. There, we show plots of the IPR of the Floquet mode  $|\phi^n\rangle$  for all  $n$  corresponding to eigenvalues of  $S^x$  for a fixed eigenvalue of  $S^2 = N/2(N/2 + 1)$ . The IPR is thus

$$\phi_{IPR}(n) = \sum_m |\langle m | \phi^n \rangle|^4 \quad (24)$$

where  $|m\rangle$  is the  $m^{th}$  eigenstate of  $\hat{S}^x$ .

Now, we focus at numerical simulations for  $H(t)$  via the IPR of the Floquet state in the representation of the transverse field i.e. the eigenstates of  $S^x$ . We kept frequency  $\omega = 90$  as a large enough value for RWA to hold, and  $N = \mathcal{O}(10^2)$  which our standard computational resources will allow.

The uppermost section displays a density plot that depicts the Inverse Participation Ratio (IPR) of the Floquet states in FIG3. The abscissa represents  $\eta = 4h/\omega$ , while the ordinate represents the spin index  $m/(2N+1)$ . The vertical lines that are dashed and colored blue correspond to the roots of the Bessel function of the first kind with order zero, denoted as  $J_0(\eta)$ . Evidently, the Inverse Participation Ratio (IPR) approaches a value of one for high values of the root of the Bessel function  $J_0(\eta)$ , which suggests a state of full localization of the Many Body system. Nevertheless, a deviation from unity occurs at the smallest root of  $J_0(\eta)$ . The observed phenomenon can be attributed to the fact that when  $\eta$  approaches the smallest root of  $J_0(\eta)$ , which is approximately equal to 2.405, the amplitudes of the higher order terms in the RW expansion, denoted by  $\mathcal{O}(J_n(\eta))$ , become significant enough to contribute to the process of delocalization. Thus, a greater magnitude of the root is favored for the purpose of investigating the phenomena.

The bottom panel contains cross sections of the full IPR plot for selected values of  $\eta$  as indicated in the legend. When the drive amplitude  $h$  is adjusted to make  $J_0(\eta) \neq 0$ , the Floquet States are mixed, but not entirely thermal, since the IPR does not fall to  $\mathcal{O}(N^{-1})$ , indicating that localization persists to some extent always. So, as long as there is an appropriate DC field,  $\hat{S}^x$  is mostly conserved and  $\hat{H}_F$  is mostly diagonal in the  $\hat{S}^x$  representation at the freezing point. The small

deviations from this conservation occur due to the role of higher order terms in the Fourier expansion of the Hamiltonian on the rotating basis that contribute additional time-periodic terms to the RWA Hamiltonian, as can be seen in FIG.II.




FIG. 3. The IPR for exact dynamics for spin model size  $N \sim 30$ . The upper panel shows a density plot of the IPR of Floquet modes for different parameter points corresponding to the ratio between strength and the frequency of the symmetry-breaking field which is the lowest root among Bessel's roots of the first kind and zeroth order *i.e.*  $\frac{4h}{\omega}$ . The constant dc part of the symmetry-breaking field is kept at a small irrational number to avoid the symmetry arising from degeneracy in Flqouet states. The IPR density is high at Bessel's first order roots,  $J_0\left(\frac{4h}{\omega}\right) = 0$  for all the available floquet states. The lower panel shows the crossectional plot of IPR for four different floquet IPR density points. The point corresponding  $J_0\left(\frac{4h}{\omega}\right) = 0$  has unity value for all states where points away from the roots have lower and high fluctuations.

The full Hamiltonian for the LMG model on the rotated basis is

$$\tilde{H}^{RWA}(t) \sim \frac{(\hat{S}^x)^2}{N-1} - 2h_0\hat{S}^x - \frac{J_0(\eta)}{N-1} \left[ (\hat{S}^z)^2 - (\hat{S}^y)^2 \right] - \frac{2}{N-1} \sum_{n=1}^{\infty} J_{2n}(\eta) \left[ (\hat{S}^z)^2 - (\hat{S}^y)^2 \right] \cos(2n\omega t) - \frac{2}{N-1} \sum_{n=1}^{\infty} J_{2n-1}(\eta) \{ \hat{S}^y, \hat{S}^z \} \sin[(2n-1)\omega t] \quad (25)$$




FIG. 4. The comparison between IPR for exact dynamics and RWA with corresponding correction orders. IPR is calculated for four different  $\eta$ 's and corresponding  $J_0(\eta) \neq 0$  values for colors, Blue:  $\eta = 2.40, J_0(\eta) = 0.0$ , dashed orange:  $\eta = 18.07, J_0(\eta) = 0.0$ , Green:  $\eta = 13.03, J_0(\eta) = 0.21$ , Red:  $\eta = 1.62, J_0(\eta) = 0.44$ . IPR plots for RWA with zeroth order aren't enough to describe the exact dynamics, but plots for RWA with first-order correction are similar to the exact dynamics.

The IPR plots pertaining to exact dynamics indicate that the freezing point, represented by the initial blue curve corresponding to Bessel's root of zeroth order

of first kind, is insufficient for achieving many-body localization, as depicted in the top panel. The reason for this phenomenon is attributed to the greater magnitude of other Bessel roots of higher order, which in turn leads to delocalization. However, it is worth noting that at elevated localization points (as depicted by the dashed orange curve), localization is particularly conspicuous. In terms of system distribution, a range of curves, denoted by both green and red, illustrates the degree of distribution. The zeroth order RWA in the central panel exhibits a consistent pattern at locations distant from the resonance point, as evidenced by the congruent green and red curves. However, the curves for both higher and lower localization points display complete localization, which contradicts the precise findings presented in the top panel. This necessitates the incorporation of higher-order corrections into the Rotating Wave Approximation (RWA). The application of the first-order correction to RWA in the lower panel results in a curve structure that is recognizable and consistent with the exact dynamics.

### III. CLASSICAL LIPKIN DYNAMICS

In the context of the classical (continuum) limit, where  $N$  approaches infinity, the disparity between neighboring values of  $s_i$  in equation 17 can be disregarded, resulting in a Hamiltonian per particle of  $h(t) \equiv \frac{1}{N}$  and the Hamiltonian per particle becomes

$$h(t) \equiv \frac{1}{N} H(t) = h + h_0 \cos(\omega t) h_1, \quad (26)$$

where,

$$\begin{aligned} (h)_{ij} &\approx -2s_i^2 \delta_{ij}, \\ H_0 &\rightarrow -2s^2 \\ (h_1)_{ij} &\approx \sqrt{1 - 4s_i^2} [\delta_{i+1,j} + \delta_{i-1,j}] \\ H_1 &\rightarrow \sqrt{1 - 4s_i^2} \cos p, \end{aligned} \quad (27)$$

In the continuum limit, the Lipkin system can be described by  $p, q$  with corresponding Hamiltonian [41]:

$$H = -2q^2 - h(t) \sqrt{1 - 4q^2} \cos p, \quad (28)$$

which yields the Hamiltonian dynamical system

$$\begin{aligned}\frac{dq}{dt} &= h(t)\sqrt{1-4q^2}\sin p \\ \frac{dp}{dt} &= 4q\left[1 - \frac{h(t)\cos p}{\sqrt{1-4q^2}}\right]\end{aligned}\quad (29)$$

We recovered Hamiltonian numerically 28 using equations 29 to obtain position and momentum coordinates  $p, q$  and plotted in the *Poincaré surface of section* (PSOS) strobbed at integer multiple of time period. A chaotic Poincaré pattern is found for all small drive amplitude s.t.  $A/J < 0.5$  and also a regular pattern emerges at for ratios  $A/J \geq 0.5$  both at small drive frequency  $\omega \sim 2.0$  [39].

However, under a regime of sufficiently high frequency, the system exhibits distinct behavior. The study involves a comparison of the Poincaré sections of the ensuing dynamics under the influence of  $h(t) = h \cos \omega t$ , where two distinct scenarios have been considered. The first scenario pertains to the case where frequency  $\omega = 2.5$ , corresponds to a lower frequency and, consequently, a smaller amplitude. The second scenario pertains to the case where frequency  $\omega = 90.0$ , corresponds to a higher frequency and, correspondingly, a higher amplitude. Both scenarios have been evaluated at  $J_0(4h/\omega) = 0$ . The Husimi Q-functions of the acquired Floquet States are compared with these. The quantum phase space is characterized by the Spectral Average of the Husimi functions of all the Floquet modes  $|\phi^n\rangle$  for a given value of  $S^2$ . Specifically, for a coherent state  $|q, p\rangle$ , the corresponding plot is presented in FIG.5.

$$H(q, p) \equiv \frac{1}{(2N+1)\pi} \sum_n \langle q, p | \phi^n \rangle \langle \phi^n | q, p \rangle$$

The classical plots are dominated by chaos at low values of  $\omega$ , as noted in Kidd's work [42]. Conversely, at high values of  $\omega$ , the Poincaré plot exhibits a distinct regular pattern, indicating localization.

#### IV. THERMALITY TO ATHERMALITY: A PHASE CROSSOVER

The analysis of the Lipkin-Glick-Meshkov model reveals two distinct scenarios at low and high external drive frequencies. As a result, it is hypothesized that a change in phase may occur due to the influence of frequency. IPR of the Floquet modes is computed numerically and plotted in FIG.6 for numerous frequencies in ascending order from frequency  $\omega = 1.0$  upto  $\omega = 50.0$  so that the system can vary adiabatically, along with the associated drive amplitude  $h$  for the localization resonant point, which is  $J_0(\frac{4h}{\omega})$  (Top panel). In the




FIG. 5. The above panel describes the phase-space Poincaré distribution symmetry breaking smaller drive frequency  $\omega = 2.5$  (left) and higher frequency  $\omega = 90.0$  (right) for system size  $N = 500$  with 100 realisation numbers. At smaller frequencies, the Poincaré picture contains chaotic behaviour (top left panel) whereas at the higher frequencies, it is a normal Poincaré picture which represents discrete freezing behaviour (top right panel). The bottom panel is Husimi Q-function average plot for a smaller frequency (bottom left) and has a uniform distribution with less contrast in colour. This means a Q-function distribution in chaotic behaviour. At the right bottom, the Husimi plot has distinct colour contrast in the Q-function average value which represents a regular dynamics pattern in the system.

low-frequency range spanning from  $\omega = 1.0$  to approximately  $\omega \approx 9.0$ , IPR exhibits values well below unity. Moreover, the IPR gradually diminishes with increasing system size, following a system size inverse proportional trend, which confirms the participation distribution (as shown in the bottom panel). As the limit of  $N$  approaches infinity, the inverse participation ratio (IPR) tends towards zero, indicating a fully delocalized state. The study conducted by IPR revealed a gradual increase in the unity of IPR over a certain frequency range, specifically at approximately  $\omega = 5.0$ . The increase in IPR is not characterized by a sudden surge, but rather a gradual increase that exhibits a phase crossover[27, 43]. As the size of the system increases, the crossover region becomes smoother.

The numerical evolution of the system is carried out using the exact Hamiltonian for a duration of 500 time periods. The average of the Hamiltonian, denoted




FIG. 6. The plotting of the Inverse Participation Ratio (IPR) at Bessel's first root with first kind  $J_0\left(\frac{4h}{\omega}\right)$  for varying drive amplitudes and frequencies in a spin array of LMG type, for different spin sizes  $N = 10, 30, 50, 100$ . The adiabatic increase of the drive frequency is observed in the range of  $O(1)$  to  $O(50)$  for varying system sizes of  $N = 10, 30, 50$ , and 100. The study reveals that the IPR exhibits a decrease in magnitude at low frequencies, followed by an increase to unity at high frequencies, across various system sizes (top panel). The findings indicate that the system's inverse participation ratio (IPR) diminishes in proportion to  $N$  in low-frequency regions, thereby validating the distribution of the system's state participation, as demonstrated in the lower panel. The increase in IPR with respect to system frequency exhibits a consistently gradual trend until it reaches a high frequency where IPR is unity, which is contingent upon system parameters. The lower panel indicates that the IPR decreases proportionally with the system size, which can be attributed to the distribution of states at lower frequencies.

as  $\langle \hat{H} \rangle$ , is then computed to determine the energy spreading through the standard deviation. The thermal conditions necessitate minimal fluctuations in energy distribution, as the spread of states leads to a limited standard deviation[44]. As a result, finite standard deviation arises in athermal regions. A graph depicting the standard deviation of the energy expectation value is presented for a system Hamiltonian adiabatically varied with drive frequency. The standard deviation of the Hamiltonian, denoted as  $\langle H \rangle_{std}$ , exhibits a small value for frequencies up to approximately  $\omega \sim 5.0$ . Subsequently, there is an increase in that follows a linear pattern as the frequency in-

creases further  $\langle H \rangle_{std}$  in FIG7. A thermal region is observed at a lower frequency, while an athermal region is detected at a higher frequency.




FIG. 7. The standard deviation of the mean value of  $H$ , denoted by  $\langle H \rangle_{std}$ , has been calculated over a span of 500 time periods. A singularity has been identified at approximately  $\omega = 5.0$ . The standard expectation value of the Hamiltonian, denoted as  $\langle H \rangle_{std}$ , exhibits a diminutive magnitude below the singularity, while a linear and gradual augmentation is observed at higher frequencies. A small standard deviation indicates the presence of a thermal region, whereas a larger finite standard deviation suggests the existence of a regular athermal region.

## V. OBSERVATIONS AND DISCUSSION

The quantum many-body localization in the Ising model is observed at high drive frequency and corresponding high drive amplitude, where the localization points are determined by setting  $J_0(2h/\omega) = 0$  for both exact and Rotated wave simulations. The delocalization away from the localization points is relatively weak. The integrability of the Ising model allows for localization to be observed at low values of  $\omega$ , despite the breakdown of the rotating wave approximation. Analytical methods beyond the adiabatic limit are intricate. The LMG model exhibits distinct localization at localization points, which are determined by the condition  $J_0(4h/\omega) = 0$ , in both the exact and Rotated Wave simulations. However, there is some degree of delocalization observed beyond these points, albeit weak. The non-integrability of the LMG model and the established onset of chaos in the thermodynamic limit at low  $\omega$  lead to a near-complete delocalization in the Inverse Participation Ratio (IPR) of the Floquet states at small  $\omega$ .

Consequently, we adiabatically varied  $\omega, h$  in the LMG model, under the constraint that  $\eta = 4h/\omega$

was held at a root of  $J_0(\eta)$ . There, we observed a crossover or phase change from nearly fully thermal to completely localised behaviour (see FIG.6). Even if the limitation is relaxed, the macroscopic behaviour changes from thermal to athermal. At low frequencies, IPR is observed to decrease with increasing system size  $N$ , and when  $N$  is big, i.e.,  $\rightarrow \infty$ , IPR appears to evaporate, resulting in a fully thermalized state at  $\eta = 4h/\omega$ . This is in contrast to the Ising model, which lacks such a change in phase. Thus, the incorporation of long-range interactions appears to trigger a change in phase from the thermal phase to the localised phase, a property that will prove useful in the design of MBL engines.

## VI. CONCLUSION AND OUTLOOK

The study delved into the onset of Dynamical Many-Body Localization in the Lipkin-Glick-Meshkov spin model, specifically in regularly driven long-range spins, serving as a paradigmatic illustration. The parametrization of many-body localization is based on the Inverse Participation Ratio of the Floquet eigenstates. The present study conducted a numerical comparison between the inverse participation ratio (IPR) of the LMG model and the transverse field Ising model (TFIM) at both low and high drive frequencies. The present study involved an examination of the phase space dynamics of the LMG model. Specifically, we sought to analyze the emergence of thermal behavior at low frequencies and localization at high frequencies, as well as the appearance of additional approximately conserved quantities in the high-frequency regime for both models.

*Conclusion:* Long-range spins exhibit strong localization in spin-coordinate space for the LMG model when the drive frequency is  $\omega \gg J$ , where  $J$  represents the spin exchange energy. The localization of the LMG model occurs at specific resonances of the drive frequency  $\omega$  and amplitude  $h$ , where  $J_0(4h/\omega) = 0$ . The TFIM case has shown comparable localization (in momentum space) at resonances determined by  $J_0(2h/\omega) = 0$ . However, the mechanism differs in long-range systems as a distinct observable ( $S_x$ )<sup>2</sup> is conserved in the LMG case, where  $S$  represents the total spin. Upon the removal of accidental degeneracy through the application of a DC transverse field in the form of  $\sim \hat{S}_x$ , the eigenstates can be mapped into a coordinate representation, leading to a robust spa-

tial localization. The periodically driven LMG model exhibits a pronounced mobility edge when the frequency is gradually increased from  $\omega \sim J$  to  $\omega \gg J$  in an adiabatic manner. In the first region where frequency  $\omega$  is small, the quantum system is expected to undergo thermalization at exceedingly high temperatures due to the attainment of dynamical chaos in classical dynamics. However, in the latter high frequency regime, the perpetual postponement occurs due to dynamical localization. The point of transition in mobility between these two regimes exhibits anomalous characteristics when analyzed in the context of thermodynamic principles, indicating a quantum change in phase which detected as *phase – crossover* between them. The absence of thermal behavior in the low-frequency limit of the short-range transverse field Ising model (TFIM) can be attributed to the significant magnitude of the inverse participation ratio (IPR). The induction of Thermal and Localized states in long-range systems can be achieved through Floquet engineering, without the need for disorder.

*Outlook:* A system characterized by high symmetry has been thoroughly examined in a pristine state. Thermalization takes place in all scenarios, alongside integrability-disrupting factors, such as disorder. However, akin to TFIM, the onset of thermalization in LMG can be postponed when the condition  $\omega \gg J$  and  $J_0(4h/\omega) = 0$  is satisfied. In systems possessing a Hamiltonian of the form given by equations 15 and 16, the time-independent Schrödinger equation can be solved analytically for certain potentials. The intermediate spin-spin interaction power law limits have received less attention compared to the infinite and long-range limit. Further research can be conducted on this topic. The LMG spin configuration undergoes a phase cross-over due to the adiabatic increase in drive frequency. This suggests the possibility of a future MBL engine that can operate between the thermal and localized regimes, with a thermodynamic cycle. Additionally, there exist prospects for diabatic corrections.

*Acknowledgement:* One of the authors, MR would like to acknowledge The University of Burdwan for support via a state-funded fellowship, as well as for providing computational resources. AR acknowledges support from the University Grants Commission (UGC) of India, via BSR Startup Grant No. F.30-425/2018(BSR), as well as from the Science and Engineering Research Board (SERB) Core Research Grant No. CRG/2018/004002.

---

[1] P. Bordia, H. Lüschen, U. Schneider, M. Knap, and I. Bloch, Periodically driving a many-body localized

quantum system, **13**, 460.  
[2] S. Sahoo, I. Schneider, and S. Eggert, Periodically

- driven many-body systems: A floquet density matrix renormalization group study, .
- [3] A. Das, Exotic freezing of response in a quantum many-body system, **82**, 172402.
- [4] G. B. Mbeng, A. Russomanno, and G. E. Santoro, The quantum ising chain for beginners, 2009.09208 [cond-mat, physics:quant-ph].
- [5] H. S. Yamada and K. S. Ikeda, Localization and delocalization properties in quasi-periodically-driven one-dimensional disordered systems, **105**, 054201.
- [6] A. Roy and A. Das, Fate of dynamical many-body localization in the presence of disorder, **91**, 121106 ()�.
- [7] H. Li, B. Shapiro, and T. Kottos, Floquet scattering theory based on effective Hamiltonians of driven systems, Phys. Rev. B **98**, 121101 (2018), publisher: American Physical Society.
- [8] A. Eckardt and E. Anisimovas, High-frequency approximation for periodically driven quantum systems from a Floquet-space perspective, New J. Phys. **17**, 093039 (2015), publisher: IOP Publishing.
- [9] L. Zhang, V. Khemani, and D. A. Huse, A floquet model for the many-body localization transition, Phys. Rev. B **94**, 224202 (2016), publisher: American Physical Society.
- [10] V. Khemani, A. Lazarides, R. Moessner, and S. Sondhi, Phase structure of driven quantum systems, **116**, 250401, publisher: American Physical Society.
- [11] N. Yunger Halpern, C. D. White, S. Gopalakrishnan, and G. Refael, Quantum engine based on many-body localization, **99**, 024203.
- [12] A. Campa, T. Dauxois, and S. Ruffo, Statistical mechanics and dynamics of solvable models with long-range interactions, **480**, 57.
- [13] T. Eisele and R. S. Ellis, Multiple phase transitions in the generalized curie-weiss model, **52**, 161.
- [14] A. Canning, A class of long range ising spin models described by curie-weiss mean field theory, **185**, 254.
- [15] D. Vu, K. Huang, X. Li, and S. Das Sarma, Fermionic manybody localization for random and quasiperiodic systems in the presence of short- and long-range interaction, Phys. Rev. Lett. **128**, 146601 (2022), publisher: American Physical Society.
- [16] G. Misguich, V. Pasquier, and J.-M. Luck, Inverse participation ratios in the xxz spin chain, Phys. Rev. B **94**, 155110 (2016).
- [17] M. Calixto and E. Romera, Inverse participation ratio and localization in topological insulator phase transitions, **2015**, P06029.
- [18] K. Fujii, Introduction to the rotating wave approximation (RWA): Two coherent oscillations, **8**, 2042.
- [19] D. A. Abanin, E. Altman, I. Bloch, and M. Serbyn, Colloquium: Many-body localization, thermalization, and entanglement, Rev. Mod. Phys. **91**, 021001 (2019), publisher: American Physical Society.
- [20] M. Srednicki, Chaos and quantum thermalization, **50**, 888.
- [21] M. Srednicki, The approach to thermal equilibrium in quantized chaotic systems, Journal of Physics A: Mathematical and General **32**, 1163 (1999).
- [22] M. Holthaus, Floquet engineering with quasienergy bands of periodically driven optical lattices, J. Phys. B: At. Mol. Opt. Phys. **49**, 013001 (2016).
- [23] M. Vogl, M. Rodriguez-Vega, and G. A. Fiete, Effective Floquet Hamiltonian in the low-frequency regime, Phys. Rev. B **101**, 024303 (2020), publisher: American Physical Society.
- [24] M. Bukov, L. D'Alessio, and A. Polkovnikov, Universal High-Frequency Behavior of Periodically Driven Systems: from Dynamical Stabilization to Floquet Engineering, ArXiv: 1407.4803v2.
- [25] L. D'Alessio and M. Rigol, Long-time behavior of isolated periodically driven interacting lattice systems, Phys. Rev. X **4**, 041048 (2014).
- [26] R. Yousefjani, S. Bose, and A. Bayat, Floquet-induced localization in long-range many-body systems, Phys. Rev. Res. **5**, 013094 (2023).
- [27] P. Sierant, M. Lewenstein, A. Scardicchio, and J. Zakrzewski, Stability of many-body localization in floquet systems, Phys. Rev. B **107**, 115132 (2023).
- [28] S. Rozhin, Abolfazl, Floquet-induced localization in long-range many-body systems 10.1103/PhysRevResearch.5.013094.
- [29] L. N. Alet Fabien, Many-body localization: An introduction and selected topics **19**, 10.1016/j.crhy.2018.03.003.
- [30] S. J. Garratt and S. Roy, Resonant energy scales and local observables in the many-body localized phase, **106**, 054309, publisher: American Physical Society.
- [31] R. B. Stinchcombe, Ising model in a transverse field. i. basic theory, **6**, 2459.
- [32] F. E. H. George Arfken, Hans Weber, *Mathematical Methods for Physicists*.
- [33] S. Mukherjee, A. Spracklen, D. Choudhury, N. Goldman, P. Öhberg, E. Andersson, and R. R. Thomson, Modulation-assisted tunneling in laser-fabricated photonic wannier-stark ladders, **17**, 115002, publisher: IOP Publishing.
- [34] S.-H. Lin, B. Sbierski, F. Dorfner, C. Karrasch, and F. Heidrich-Meisner, Many-body localization of spinless fermions with attractive interactions in one dimension, **4**, 002.
- [35] N. C. Murphy, R. Wortis, and W. A. Atkinson, Generalized inverse participation ratio as a possible measure of localization for interacting systems, **83**, 184206, publisher: American Physical Society.
- [36] E. J. Torres-Herrera, I. Vallejo-Fabila, A. J. Martínez-Mendoza, and L. F. Santos, Self-averaging in many-body quantum systems out of equilibrium: Time dependence of distributions, **102**, 062126, publisher: American Physical Society.
- [37] N. Trivedi and D. Heidarian, Can disorder drive a mott insulator into a metal in 2d?, **160**, 296.
- [38] G. Misguich, V. Pasquier, and J.-M. Luck, Inverse participation ratios in the XXZ spin chain, **94**, 155110, publisher: American Physical Society.
- [39] A. Russomanno, R. Fazio, and G. E. Santoro, Thermalization in a periodically driven fully connected quantum ising ferromagnet, **110**, 37005.
- [40] T. Mori, Prethermalization in the transverse-field ising chain with long-range interactions, **52**, 054001, publisher: IOP Publishing.
- [41] B. Sciolla and G. Biroli, Quantum quenches and

- off-equilibrium dynamical transition in the infinite-dimensional bose-hubbard model, **105**, 220401.
- [42] R. A. Kidd, M. K. Olsen, and J. F. Corney, Quantum chaos in a bose-hubbard dimer with modulated tunneling, Phys. Rev. A **100**, 013625 (2019).
- [43] S. Sachdev, *Quantum Phase Transitions*, 2nd ed. (Cambridge University Press, Cambridge, 2011).
- [44] P. Reimann, Symmetry-prohibited thermalization after a quantum quench, **2021**, 103106, publisher: IOP Publishing and SISSA.
- [45] L. Reichl, *The Transition to Chaos: Conservative Classical and Quantum Systems*, Fundamental Theories of Physics, Vol. 200 (Springer International Publishing).
- [46] A. Roy and A. Das, Fate of dynamical many-body localization in the presence of disorder, **91**, 121106 () .
- [47] N. Srivatsa, R. Moessner, and A. E. Nielsen, Many-body delocalization via emergent symmetry, **125**, 240401.
- [48] Quantum ising phase transition.
- [49] B. Marcos, A. Gabrielli, and M. Joyce, Relaxation of quasi-stationary states in long range interacting systems and a classification of the range of pair interactions, **10**, 676.
- [50] A. Haldar and A. Das, Dynamical many-body localization and delocalization in periodically driven closed quantum systems, **529**, 1600333.
- [51] J. M. Deutsch, Quantum statistical mechanics in a closed system, **43**, 2046 () .
- [52] E. W. Hobson, On the Second Mean-Value Theorem of the Integral Calculus (1909).
- [53] M. Rigol and M. Srednicki, Alternatives to eigenstate thermalization, Physical Review Letters **108**, 10.1103/physrevlett.108.110601 (2012).
- [54] J. M. Deutsch, Eigenstate thermalization hypothesis, **81**, 296 () .
- [55] R. Nandkishore and D. A. Huse, Many-body localization and thermalization in quantum statistical mechanics, Annual Review of Condensed Matter Physics **6**, 15 (2015), <https://doi.org/10.1146/annurev-conmatphys-031214-014726>.