## TABLE 2. SPECIES RANKED IN DESCENDING ORDER OF TOTAL FY 2016 REPORTED EXPENDITURES, NOT INCLUDING LAND ACQUISITION COSTS

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 1    | Steelhead (Oncorhynchus (=Salmo) mykiss) - Middle Columbia River DPS  | Т      | \$64,365,536  | \$64,365,536       |
| 2    | Salmon, Chinook (Oncorhynchus (=Salmo) tshawytscha) -<br>Snake River spring/summer-run ESU                  | Т      | \$61,231,560  | \$125,597,096      |
| 3    | Steelhead (Oncorhynchus (=Salmo) mykiss) - Snake River Basin DPS  | Т      | \$60,270,758  | \$185,867,854      |
| 4    | Salmon, Chinook (Oncorhynchus (=Salmo) tshawytscha) - Upper Columbia spring-run ESU                         | Е      | \$44,865,295  | \$230,733,149      |
| 5    |   | E      | \$43,237,405  | \$273,970,554      |
| 6    | Trout, bull (Salvelinus confluentus) - U.S.A., conterminous, lower 48 states                                | Т      | \$40,233,890  | \$314,204,444      |
| 7    | Salmon, Chinook (Oncorhynchus (=Salmo) tshawytscha) - Snake River fall-run ESU                              | Т      | \$39,689,609  | \$353,894,053      |
| 8    | Salmon, Chinook (Oncorhynchus (=Salmo) tshawytscha) -<br>Lower Columbia River ESU                           | Т      | \$33,557,518  | \$387,451,571      |
| 9    | Salmon, Chinook (Oncorhynchus (=Salmo) tshawytscha) - Puget Sound ESU                                       | Т      | \$29,187,297  | \$416,638,868      |
| 10   | Steelhead (Oncorhynchus (=Salmo) mykiss) - Upper Columbia River DPS   | Т      | \$28,642,935  | \$445,281,803      |
| 11   | Salmon, sockeye (Oncorhynchus (=Salmo) nerka) - Snake<br>River ESU  | Е      | \$28,114,109  | \$473,395,912      |
| 12   | Woodpecker, red-cockaded (Picoides borealis) - Wherever found   | Е      | \$25,032,102  | \$498,428,014      |
| 13   | Sea lion, Steller (Eumetopias jubatus) - Western DPS  | E      | \$22,127,195  | \$520,555,209      |
| 14   | Salmon, coho (Oncorhynchus (=Salmo) kisutch) - Lower Columbia River ESU                                     | Т      | \$19,639,995  | \$540,195,204      |
| 15   | Tortoise, desert (Gopherus agassizii) - Wherever found, except AZ south and east of Colorado R., and Mexico | Т      | \$18,554,604  | \$558,749,808      |
| 16   | Salmon, coho (Oncorhynchus (=Salmo) kisutch) - Southern Oregon - Northern California Coast ESU              | Т      | \$18,498,254  | \$577,248,062      |
| 17   | Salmon, Chinook (Oncorhynchus (=Salmo) tshawytscha) - Upper Willamette River ESU                            | Т      | \$18,271,433  | \$595,519,495      |
| 18   | Steelhead (Oncorhynchus (=Salmo) mykiss) - Lower Columbia River DPS   | Т      | \$18,186,684  | \$613,706,179      |
| 19   | Sturgeon, white (Acipenser transmontanus) - U.S.A. (ID, MT), Canada (B.C.), Kootenai R. system              | Е      | \$16,013,924  | \$629,720,103      |
| 20   | Owl, northern spotted (Strix occidentalis caurina) - Wherever found   | Т      | \$15,624,955  | \$645,345,058      |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 21   | Salmon, chum (Oncorhynchus keta) - Columbia River ESU   | Т      | \$14,811,627  | \$660,156,685      |
| 22   | Bat, Indiana (Myotis sodalis) - Wherever found  | E      | \$14,035,548  | \$674,192,233      |
| 23   | Sucker, razorback (Xyrauchen texanus) - Wherever found  | E      | \$13,945,914  | \$688,138,147      |
| 24   | Bat, Northern long-eared (Myotis septentrionalis) - Wherever found  | Т      | \$12,743,822  | \$700,881,969      |
| 25   | Smelt, delta (Hypomesus transpacificus) - Wherever found  | Т      | \$12,019,452  | \$712,901,421      |
| 26   | Steelhead (Oncorhynchus (=Salmo) mykiss) - Puget Sound DPS  | Т      | \$11,901,567  | \$724,802,988      |
| 27   | Bear, grizzly (Ursus arctos horribilis) - Bitterroot Ecosystem Recovery Zone Population   | EXPN   | \$11,724,619  | \$736,527,607      |
| 28   | Plover, piping (Charadrius melodus) - [Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered. | Т      | \$11,306,784  | \$747,834,391      |
| 29   | Chub, bonytail (Gila elegans) - Wherever found  | E      | \$10,909,990  | \$758,744,381      |
| 30   | Steelhead (Oncorhynchus (=Salmo) mykiss) - Upper Willamette River DPS   | Т      | \$10,828,635  | \$769,573,016      |
| 31   | Whale, North Atlantic Right (Eubalaena glacialis) - Wherever found  | Е      | \$10,308,042  | \$779,881,058      |
| 32   | Salmon, Chinook (Oncorhynchus (=Salmo) tshawytscha) - Sacramento River winter-run ESU   | E      | \$10,256,141  | \$790,137,199      |
| 33   | Salmon, Atlantic (Salmo salar) - Gulf of Maine DPS  | E      | \$10,013,881  | \$800,151,080      |
| 34   | Salmon, coho (Oncorhynchus (=Salmo) kisutch) - Oregon Coast ESU   | Т      | \$9,784,535   | \$809,935,615      |
| 35   | Chub, humpback (Gila cypha) - Wherever found  | E      | \$9,311,369   | \$819,246,984      |
| 36   | Flycatcher, southwestern willow (Empidonax traillii extimus) - Wherever found   | E      | \$8,946,102   | \$828,193,086      |
| 37   | Minnow, Rio Grande Silvery (Hybognathus amarus) -<br>Wherever found, except where listed as an experimental<br>population                                     | Е      | \$8,544,342   | \$836,737,428      |
| 38   | Pikeminnow (=squawfish), Colorado (Ptychocheilus lucius) - Wherever found, except where listed as an experimental population                                  | Е      | \$8,519,374   | \$845,256,802      |
| 39   | Elephant, African (Loxodonta africana) - Wherever found   | Т      | \$8,029,080   | \$853,285,882      |
| 40   | Tern, least (Sterna antillarum) - interior pop.   | E      | \$7,892,269   | \$861,178,151      |
| 41   | Ocelot (Leopardus (=Felis) pardalis) - wherever found   | E      | \$7,183,922   | \$868,362,073      |
| 42   | Whale, humpback (Megaptera novaeangliae) - Wherever found   | Е      | \$7,070,114   | \$875,432,187      |
| 43   | Salmon, chum (Oncorhynchus keta) - Hood Canal summer-run ESU  | Т      | \$6,771,036   | \$882,203,223      |
| 44   | Cuckoo, yellow-billed (Coccyzus americanus) - Western U.S. DPS  | Т      | \$6,343,860   | \$888,547,083      |
| 45   | Steelhead (Oncorhynchus (=Salmo) mykiss) - Central California Coast DPS   | Т      | \$6,222,951   | \$894,770,034      |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 46   | Salmon, sockeye (Oncorhynchus (=Salmo) nerka) - Ozette Lake ESU   | Т      | \$5,518,535   | \$900,288,569      |
| 47   | Manatee, West Indian (Trichechus manatus) - Wherever found  | Е      | \$5,430,465   | \$905,719,034      |
| 48   | Steelhead (Oncorhynchus (=Salmo) mykiss) - California<br>Central Valley DPS   | Т      | \$5,428,021   | \$911,147,055      |
| 49   | Sea turtle, loggerhead (Caretta caretta) - Northwest Atlantic Ocean DPS   | Т      | \$5,221,940   | \$916,368,995      |
| 50   | Frog, California red-legged (Rana draytonii) - Wherever found   | Т      | \$5,166,801   | \$921,535,796      |
| 51   | Bear, grizzly (Ursus arctos horribilis) - U.S.A., conterminous (lower 48) States, except where listed as an experimental population or delisted       | Т      | \$5,109,765   | \$926,645,561      |
| 52   | Salmon, coho (Oncorhynchus (=Salmo) kisutch) - Central California Coast ESU   | E      | \$5,037,464   | \$931,683,025      |
| 53   | Sea turtle, leatherback (Dermochelys coriacea) - Wherever found   | E      | \$4,887,129   | \$936,570,154      |
| 54   | Plover, western snowy (Charadrius alexandrinus nivosus) - Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast) | Т      | \$4,878,512   | \$941,448,666      |
| 55   | Salmon, Chinook (Oncorhynchus (=Salmo) tshawytscha) - California Coastal ESU  | Т      | \$4,852,009   | \$946,300,675      |
| 56   | Murrelet, marbled (Brachyramphus marmoratus) - U.S.A. (CA, OR, WA)  | Т      | \$4,721,414   | \$951,022,089      |
| 57   | Gorilla (Gorilla gorilla) - Wherever found  | E      | \$4,588,000   | \$955,610,089      |
| 58   | Steelhead (Oncorhynchus (=Salmo) mykiss) - Northern California DPS  | Т      | \$4,527,260   | \$960,137,349      |
| 59   | Combshell, Cumberlandian (Epioblasma brevidens) -<br>Wherever found; Except where listed as Experimental<br>Populations                               | Е      | \$4,445,161   | \$964,582,510      |
| 60   | Sea turtle, Kemp's ridley (Lepidochelys kempii) - Wherever found  | Е      | \$4,269,587   | \$968,852,097      |
| 61   | Steelhead (Oncorhynchus (=Salmo) mykiss) - Southern California DPS  | Е      | \$4,085,228   | \$972,937,325      |
| 62   | Wolf, gray (Canis lupus) - WY, EXPN population  | EXPN   | \$4,083,384   | \$977,020,709      |
| 63   | Owl, Mexican spotted (Strix occidentalis lucida) - Wherever found   | Т      | \$3,940,575   | \$980,961,284      |
| 64   | Trout, Lahontan cutthroat (Oncorhynchus clarkii henshawi) - Wherever found  | Т      | \$3,693,716   | \$984,655,000      |
| 65   | Seal, Hawaiian monk (Monachus schauinslandi) - Wherever found   | Е      | \$3,683,303   | \$988,338,303      |
| 66   | Tortoise, gopher (Gopherus polyphemus) - West of Mobile and Tombigbee Rivers  | Т      | \$3,677,412   | \$992,015,715      |
| 67   | Steelhead (Oncorhynchus (=Salmo) mykiss) - South-<br>Central California Coast DPS   | Т      | \$3,587,667   | \$995,603,382      |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 68   | Panther, Florida (Puma (=Felis) concolor coryi) - Wherever found  | Е      | \$3,581,925   | \$999,185,307      |
| 69   | Sea turtle, hawksbill (Eretmochelys imbricata) - Wherever found   | Е      | \$3,551,590   | \$1,002,736,897    |
| 70   | Bat, Hawaiian hoary (Lasiurus cinereus semotus) - Wherever found  | Е      | \$3,424,744   | \$1,006,161,641    |
| 71   | Wolf, gray (Canis lupus) - MN   | Т      | \$3,358,824   | \$1,009,520,465    |
| 72   | Sage-grouse, Gunnison (Centrocercus minimus) - Wherever found   | Т      | \$3,266,168   | \$1,012,786,633    |
| 73   | Whale, killer (Orcinus orca) - Southern Resident DPS  | E      | \$3,234,532   | \$1,016,021,165    |
| 74   | Trout, bull (Salvelinus confluentus) - Clackamas River subbasin experimental population   | EXPN   | \$3,233,099   | \$1,019,254,264    |
| 75   | Sturgeon (Gulf subspecies), Atlantic (Acipenser oxyrinchus (=oxyrhynchus) desotoi) - Wherever found   | Т      | \$3,193,864   | \$1,022,448,128    |
| 76   | Bear, polar (Ursus maritimus) - Wherever found  | Т      | \$3,139,348   | \$1,025,587,476    |
| 77   | Salmon, Chinook (Oncorhynchus (=Salmo) tshawytscha) - Central Valley spring-run ESU   | Т      | \$3,101,029   | \$1,028,688,505    |
| 78   | Stork, wood (Mycteria americana) - AL, FL, GA, MS, NC, SC   | Т      | \$3,071,456   | \$1,031,759,961    |
| 79   | Chub, Borax Lake (Gila boraxobius) - Wherever found   | E      | \$3,059,410   | \$1,034,819,371    |
| 80   | Whale, finback (Balaenoptera physalus) - Wherever found   | E      | \$2,990,369   | \$1,037,809,740    |
| 81   | Fox, San Joaquin kit (Vulpes macrotis mutica) - wherever found  | Е      | \$2,990,031   | \$1,040,799,771    |
| 82   | Whale, blue (Balaenoptera musculus) - Wherever found  | E      | \$2,958,689   | \$1,043,758,460    |
| 83   | Wolf, Mexican (Canis lupus baileyi) - U.S.A. (portions of AZ and NM)see 17.84(k)  | EXPN   | \$2,905,023   | \$1,046,663,483    |
| 84   | Tern, California least (Sterna antillarum browni) - Wherever found  | E      | \$2,856,294   | \$1,049,519,777    |
| 85   | Sturgeon, green (Acipenser medirostris) - Southern DPS  | Т      | \$2,755,720   | \$1,052,275,497    |
| 86   | Salamander, California tiger (Ambystoma californiense) - U.S.A. (CA - Santa Barbara County)   | E      | \$2,742,174   | \$1,055,017,671    |
| 87   | Wolf, gray (Canis lupus) - U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico. | E      | \$2,721,544   | \$1,057,739,215    |
| 88   | Sucker, Lost River (Deltistes luxatus) - Wherever found   | E      | \$2,694,599   | \$1,060,433,814    |
| 89   | Sucker, shortnose (Chasmistes brevirostris) - Wherever found  | Е      | \$2,687,499   | \$1,063,121,313    |
| 90   | Rail, Yuma clapper (Rallus longirostris yumanensis) - Wherever found  | Е      | \$2,529,587   | \$1,065,650,900    |
| 91   | Ferret, black-footed (Mustela nigripes) - U.S.A. (WY and specified portions of AZ, CO, MT, SD, and UT, see 17.84(g)(9))   | EXPN   | \$2,479,747   | \$1,068,130,647    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 92   | Ferret, black-footed (Mustela nigripes) - Wherever found, except where listed as an experimental population | E      | \$2,446,190   | \$1,070,576,837    |
| 93   | Caracara, Audubon's crested (Polyborus plancus audubonii) - FL pop.   | Т      | \$2,408,930   | \$1,072,985,767    |
| 94   | Manatee, West African (Trichechus senegalensis) - Wherever found  | Т      | \$2,375,240   | \$1,075,361,007    |
| 95   | Rabbit, Columbia Basin Pygmy (Brachylagus idahoensis) - Columbia Basin DPS                                  | E      | \$2,372,632   | \$1,077,733,639    |
| 96   | Gnatcatcher, coastal California (Polioptila californica californica) - Wherever found                       | Т      | \$2,305,975   | \$1,080,039,614    |
| 97   | Mouse, salt marsh harvest (Reithrodontomys raviventris) - wherever found                                    | E      | \$2,292,276   | \$1,082,331,890    |
| 98   | Vireo, least Bell's (Vireo bellii pusillus) - Wherever found  | E      | \$2,289,658   | \$1,084,621,548    |
| 99   | Toad, arroyo (=arroyo southwestern) (Anaxyrus californicus) - Wherever found                                | E      | \$2,282,777   | \$1,086,904,325    |
| 100  | Sea turtle, green (Chelonia mydas) - North Atlantic DPS   | Т      | \$2,249,167   | \$1,089,153,492    |
| 101  | Frog, Oregon spotted (Rana pretiosa) - Wherever found   | Т      | \$2,207,445   | \$1,091,360,937    |
| 102  | Tiger (Panthera tigris) - Wherever found  | E      | \$2,089,700   | \$1,093,450,637    |
| 103  | Whale, sperm (Physeter catodon (=macrocephalus)) - Wherever found   | E      | \$2,080,368   | \$1,095,531,005    |
| 104  | Lynx, Canada (Lynx canadensis) - Contiguous U.S. DPS  | Т      | \$2,066,279   | \$1,097,597,284    |
| 105  | Snake, eastern indigo (Drymarchon corais couperi) - Wherever found  | Т      | \$2,023,270   | \$1,099,620,554    |
| 106  | Elephant, Asian (Elephas maximus) - Wherever found  | E      | \$2,007,600   | \$1,101,628,154    |
| 107  | Shrike, San Clemente loggerhead (Lanius ludovicianus mearnsi) - Wherever found                              | E      | \$1,998,364   | \$1,103,626,518    |
| 108  | Darter, duskytail (Etheostoma percnurum) - Wherever found   | E      | \$1,987,710   | \$1,105,614,228    |
| 109  | Bat, gray (Myotis grisescens) - Wherever found  | E      | \$1,961,039   | \$1,107,575,267    |
| 110  | Condor, California (Gymnogyps californianus) - U.S.A. (specific portions of Arizona, Nevada, and Utah)      | EXPN   | \$1,935,132   | \$1,109,510,399    |
| 111  | Fairy shrimp, vernal pool (Branchinecta lynchi) - Wherever found  | Т      | \$1,894,898   | \$1,111,405,297    |
| 112  | Sea turtle, loggerhead (Caretta caretta) - North Pacific Ocean DPS  | E      | \$1,830,084   | \$1,113,235,381    |
| 113  | Snake, giant garter (Thamnophis gigas) - Wherever found   | Т      | \$1,807,870   | \$1,115,043,251    |
| 114  | Whale, beluga (Delphinapterus leucas) - Cook Inlet DPS  | E      | \$1,772,204   | \$1,116,815,455    |
| 115  | Parrot, Puerto Rican (Amazona vittata) - Wherever found   | E      | \$1,738,365   | \$1,118,553,820    |
| 116  | Eulachon (Thaleichthys pacificus) - Southern DPS  | Т      | \$1,720,738   | \$1,120,274,558    |
| 117  | Whale, Sei (Balaenoptera borealis) - Wherever found   | Е      | \$1,695,648   | \$1,121,970,206    |
| 118  | Sea turtle, green (Chelonia mydas) - Central West Pacific DPS   | Е      | \$1,667,408   | \$1,123,637,614    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 119  | Shiner, Pecos bluntnose (Notropis simus pecosensis) - Wherever found   | Т      | \$1,648,550   | \$1,125,286,164    |
| 120  | Abalone, White (Haliotis sorenseni) - North America (West Coast from Point Conception, CA, U.S.A., to Punta Abreojos, Baja California, Mexico)               | E      | \$1,636,960   | \$1,126,923,124    |
| 121  | Scrub-jay, Florida (Aphelocoma coerulescens) - Wherever found  | Т      | \$1,541,169   | \$1,128,464,293    |
| 122  | Shiner, Topeka (Notropis topeka (=tristis)) - Wherever found, except where listed as an experimental population  | Е      | \$1,519,345   | \$1,129,983,638    |
| 123  | Plover, piping (Charadrius melodus) - [Great Lakes watershed DPS] - Great Lakes, watershed in States of IL, IN, MI, MN, NY, OH, PA, and WI and Canada (Ont.) | Е      | \$1,512,833   | \$1,131,496,471    |
| 124  | Horned lark, streaked (Eremophila alpestris strigata) - Wherever found   | Т      | \$1,476,867   | \$1,132,973,338    |
| 125  | Sturgeon, shortnose (Acipenser brevirostrum) - Wherever found  | E      | \$1,400,005   | \$1,134,373,343    |
| 126  | Checkerspot, Taylor's (=whulge) (Euphydryas editha taylori) - Wherever found   | E      | \$1,398,297   | \$1,135,771,640    |
| 127  | Butterfly, Karner blue (Lycaeides melissa samuelis) -<br>Wherever found  | E      | \$1,346,957   | \$1,137,118,597    |
| 128  | Eider, Steller's (Polysticta stelleri) - AK breeding pop.  | Т      | \$1,313,186   | \$1,138,431,783    |
| 129  | Topminnow, Gila (incl. Yaqui) (Poeciliopsis occidentalis) - Wherever found   | Е      | \$1,305,800   | \$1,139,737,583    |
| 130  | Sea turtle, green (Chelonia mydas) - East Pacific DPS  | Т      | \$1,278,531   | \$1,141,016,114    |
| 131  | Chub, Gila (Gila intermedia) - Wherever found  | E      | \$1,237,727   | \$1,142,253,841    |
| 132  | Coral, staghorn (Acropora cervicornis) - Wherever found  | Т      | \$1,234,097   | \$1,143,487,938    |
| 133  | Beetle, American burying (Nicrophorus americanus) -<br>Wherever found, except where listed as an experimental<br>population                                  | Е      | \$1,225,274   | \$1,144,713,212    |
| 134  | Steelhead (Oncorhynchus (=Salmo) mykiss) - Middle Columbia River DPS   | EXPN   | \$1,213,970   | \$1,145,927,182    |
| 135  | Sturgeon, Atlantic (Acipenser oxyrinchus oxyrinchus) - Gulf of Maine DPS   | Т      | \$1,192,663   | \$1,147,119,845    |
| 136  | Turtle, bog (Clemmys muhlenbergii) - Wherever found, except GA, NC, SC, TN, VA   | Т      | \$1,183,444   | \$1,148,303,289    |
| 137  | Sturgeon, Atlantic (Acipenser oxyrinchus oxyrinchus) - Chesapeake Bay DPS  | Е      | \$1,151,249   | \$1,149,454,538    |
| 138  | Kite, Everglade snail (Rostrhamus sociabilis plumbeus) - Wherever found  | Е      | \$1,136,523   | \$1,150,591,061    |
| 139  | Prairie-chicken, lesser (Tympanuchus pallidicinctus) - Wherever found  | Т      | \$1,134,526   | \$1,151,725,587    |
| 140  | Abalone, Black (Haliotis cracherodii) - Wherever found   | E      | \$1,127,055   | \$1,152,852,642    |
| 141  | Tadpole shrimp, vernal pool (Lepidurus packardi) - Wherever found  | Е      | \$1,116,870   | \$1,153,969,512    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 142  | Crow, Hawaiian (='alala) (Corvus hawaiiensis) - Wherever found                                      | Е      | \$1,113,181   | \$1,155,082,693    |
| 143  | Shearwater, Newell's Townsend's (Puffinus auricularis newelli) - Wherever found                     | Т      | \$1,106,222   | \$1,156,188,915    |
| 144  | Sparrow, Cape Sable seaside (Ammodramus maritimus mirabilis) - Wherever found                       | E      | \$1,105,627   | \$1,157,294,542    |
| 145  | Rhinoceros, black (Diceros bicornis) - Africa   | E      | \$1,087,500   | \$1,158,382,042    |
| 146  | Frog, Sierra Nevada Yellow-legged (Rana sierrae) - Wherever found                                   | E      | \$1,076,657   | \$1,159,458,699    |
| 147  | Whale, false killer (Pseudorca crassidens) - Main Hawaiian Islands Insular DPS                      | E      | \$1,073,275   | \$1,160,531,974    |
| 148  | Sturgeon, Atlantic (Acipenser oxyrinchus oxyrinchus) - South Atlantic DPS                           | E      | \$1,067,114   | \$1,161,599,088    |
| 149  | Knot, red (Calidris canutus rufa) - Wherever found  | Т      | \$1,051,480   | \$1,162,650,568    |
| 150  | Warbler (=wood), golden-cheeked (Dendroica chrysoparia) - Wherever found                            | E      | \$1,050,715   | \$1,163,701,283    |
| 151  | Sea turtle, green (Chelonia mydas) - South Atlantic DPS   | Т      | \$1,040,944   | \$1,164,742,227    |
| 152  | Petrel, Hawaiian (Pterodroma sandwichensis) - Wherever found  | E      | \$1,037,195   | \$1,165,779,422    |
| 153  | Sea turtle, green (Chelonia mydas) - Central North Pacific DPS                                      | Т      | \$1,036,007   | \$1,166,815,429    |
| 154  | Lizard, blunt-nosed leopard (Gambelia silus) - Wherever found                                       | E      | \$1,027,064   | \$1,167,842,493    |
| 155  | Gibbons (Hylobates spp. (including Nomascus)) - Wherever found                                      | E      | \$1,018,400   | \$1,168,860,893    |
| 156  | Wedgemussel, dwarf (Alasmidonta heterodon) - Wherever found   | E      | \$1,016,736   | \$1,169,877,629    |
| 157  | Crow, Mariana (=aga) (Corvus kubaryi) - Wherever found  | E      | \$990,338     | \$1,170,867,967    |
| 158  | Rail, California clapper (Rallus longirostris obsoletus) - Wherever found                           | E      | \$983,279     | \$1,171,851,246    |
| 159  | Wolf, red (Canis rufus) - U.S.A. (portions of NC and TN)  | EXPN   | \$973,097     | \$1,172,824,343    |
| 160  | Whale, bowhead (Balaena mysticetus) - Wherever found  | E      | \$972,511     | \$1,173,796,854    |
| 161  | Spikedace (Meda fulgida) - Wherever found   | E      | \$967,603     | \$1,174,764,457    |
| 162  | Bear, Louisiana black (Ursus americanus luteolus) - Wherever found                                  | Т      | \$962,617     | \$1,175,727,074    |
| 163  | Minnow, loach (Tiaroga cobitis) - Wherever found  | E      | \$956,341     | \$1,176,683,415    |
| 164  | Sawfish, smalltooth (Pristis pectinata) - US DPS - Smalltooth sawfish originating from U.S. waters. | Е      | \$946,959     | \$1,177,630,374    |
| 165  | Sucker, June (Chasmistes liorus) - Wherever found   | E      | \$934,900     | \$1,178,565,274    |
| 166  | Pupfish, Devils Hole (Cyprinodon diabolis) - Wherever found   | Е      | \$932,403     | \$1,179,497,677    |
| 167  | Sucker, Santa Ana (Catostomus santaanae) - 3 CA river basins  | Т      | \$922,402     | \$1,180,420,079    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 168  | Goose, Hawaiian (Branta (=Nesochen) sandvicensis) - Wherever found   | Е      | \$909,535     | \$1,181,329,614    |
| 169  | Pocket gopher, Tenino (Thomomys mazama tumuli) - Wherever found  | Т      | \$908,775     | \$1,182,238,389    |
| 170  | Beetle, valley elderberry longhorn (Desmocerus californicus dimorphus) - Wherever found                                  | Т      | \$907,854     | \$1,183,146,243    |
| 171  | Stilt, Hawaiian (Himantopus mexicanus knudseni) - Wherever found   | E      | \$898,229     | \$1,184,044,472    |
| 172  | Frog, Chiricahua leopard (Rana chiricahuensis) - Wherever found  | Т      | \$898,059     | \$1,184,942,531    |
| 173  | Skink, sand (Neoseps reynoldsi) - Wherever found   | Т      | \$894,285     | \$1,185,836,816    |
| 174  | Crane, whooping (Grus americana) - Wherever found, except where listed as an experimental population                     | E      | \$882,113     | \$1,186,718,929    |
| 175  | Higgins eye (pearlymussel) (Lampsilis higginsii) - Wherever found  | E      | \$881,978     | \$1,187,600,907    |
| 176  | Sturgeon, Atlantic (Acipenser oxyrinchus oxyrinchus) - New York Bight DPS  | E      | \$858,655     | \$1,188,459,562    |
| 177  | Otter, southern sea (Enhydra lutris nereis) - Wherever found   | Т      | \$836,029     | \$1,189,295,591    |
| 178  | Alligator, American (Alligator mississippiensis) - Wherever found  | SAT    | \$835,957     | \$1,190,131,548    |
| 179  | Jaguar (Panthera onca) - Wherever found  | E      | \$829,353     | \$1,190,960,901    |
| 180  | Mouse, Pacific pocket (Perognathus longimembris pacificus) - Wherever found  | E      | \$806,257     | \$1,191,767,158    |
| 181  | Butterfly, bay checkerspot (Euphydryas editha bayensis) - Wherever found   | Т      | \$790,057     | \$1,192,557,215    |
| 182  | Sea turtle, green (Chelonia mydas) - Central South Pacific DPS   | E      | \$782,880     | \$1,193,340,095    |
| 183  | Rockfish, yelloweye (Sebastes ruberrimus) - Puget Sound - Georgia Basin DPS  | Т      | \$727,095     | \$1,194,067,190    |
| 184  | Mussel, snuffbox (Epioblasma triquetra) - Wherever found   | E      | \$723,285     | \$1,194,790,475    |
| 185  | Rockfish, canary (Sebastes pinniger) - Puget Sound - Georgia Basin DPS   | Т      | \$721,521     | \$1,195,511,996    |
| 186  | Sea turtle, olive ridley (Lepidochelys olivacea) - Wherever found, except when listed as endangered under 50 CFR 224.101 | Т      | \$721,272     | \$1,196,233,268    |
| 187  | Warbler, Kirtland's (Setophaga kirtlandii (= Dendroica kirtlandii)) - Wherever found                                     | Е      | \$684,806     | \$1,196,918,074    |
| 188  | Bocaccio (Sebastes paucispinis) - Puget Sound - Georgia Basin DPS  | Е      | \$681,045     | \$1,197,599,119    |
| 189  | Deer, Columbian white-tailed (Odocoileus virginianus leucurus) - Columbia River DPS                                      | Е      | \$665,268     | \$1,198,264,387    |
| 190  | Coot, Hawaiian (Fulica americana alai) - Wherever found  | Е      | \$657,706     | \$1,198,922,093    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 191  | Bat, Virginia big-eared (Corynorhinus (=Plecotus) townsendii virginianus) - Wherever found                    | Е      | \$654,657     | \$1,199,576,750    |
| 192  | Condor, California (Gymnogyps californianus) - U.S.A. only, except where listed as an experimental population | E      | \$637,125     | \$1,200,213,875    |
| 193  | Butterfly, Fender's blue (Icaricia icarioides fenderi) - Wherever found                                       | Е      | \$630,579     | \$1,200,844,454    |
| 194  | Sparrow, San Clemente sage (Amphispiza belli clementeae) - Wherever found                                     | Т      | \$622,397     | \$1,201,466,851    |
| 195  | Elktoe, Appalachian (Alasmidonta raveneliana) - Wherever found  | E      | \$617,962     | \$1,202,084,813    |
| 196  | Salamander, frosted flatwoods (Ambystoma cingulatum) - Wherever found   | Т      | \$615,465     | \$1,202,700,278    |
| 197  | Coral, Mountainous Star (Orbicella faveolata) - Wherever found  | Т      | \$602,894     | \$1,203,303,172    |
| 198  | Eider, spectacled (Somateria fischeri) - Wherever found   | Т      | \$595,886     | \$1,203,899,058    |
| 199  | Gartersnake, northern Mexican (Thamnophis eques megalops) - Wherever found                                    | Т      | \$589,474     | \$1,204,488,532    |
| 200  | Rhinoceros, Sumatran (Dicerorhinus sumatrensis) - Wherever found  | E      | \$588,000     | \$1,205,076,532    |
| 201  | Pogonia, small whorled (Isotria medeoloides) - Wherever found   | Т      | \$578,963     | \$1,205,655,495    |
| 202  | Coral, elkhorn (Acropora palmata) - Wherever found  | Т      | \$577,494     | \$1,206,232,989    |
| 203  | Crane, Mississippi sandhill (Grus canadensis pulla) - Wherever found  | E      | \$575,239     | \$1,206,808,228    |
| 204  | Darter, Cherokee (Etheostoma scotti) - Wherever found   | Т      | \$565,120     | \$1,207,373,348    |
| 205  | Fritillary, Gentner's (Fritillaria gentneri) - Wherever found   | E      | \$564,567     | \$1,207,937,915    |
| 206  | Chaffseed, American (Schwalbea americana) - Wherever found  | E      | \$561,368     | \$1,208,499,283    |
| 207  | Paintbrush, golden (Castilleja levisecta) - Wherever found  | Т      | \$561,097     | \$1,209,060,380    |
| 208  | Darter, Etowah (Etheostoma etowahae) - Wherever found   | Е      | \$554,620     | \$1,209,615,000    |
| 209  | Lupine, Kincaid's (Lupinus sulphureus ssp. kincaidii) - Wherever found  | Т      | \$553,207     | \$1,210,168,207    |
| 210  | Duck, Hawaiian (=koloa) (Anas wyvilliana) - Wherever found  | E      | \$553,014     | \$1,210,721,221    |
| 211  | Woundfin (Plagopterus argentissimus) - Wherever found, except where listed as an experimental population      | E      | \$551,800     | \$1,211,273,021    |
| 212  | Bean, rayed (Villosa fabalis) - Wherever found  | E      | \$534,947     | \$1,211,807,968    |
| 213  | Pronghorn, Sonoran (Antilocapra americana sonoriensis) - U.S.A. (AZ), Mexico                                  | EXPN   | \$530,979     | \$1,212,338,947    |
| 214  | Bat, lesser long-nosed (Leptonycteris curasoae yerbabuenae) - Wherever found                                  | E      | \$523,261     | \$1,212,862,208    |
| 215  | Prairie-chicken, Attwater's greater (Tympanuchus cupido attwateri) - Wherever found                           | Е      | \$520,400     | \$1,213,382,608    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 216  | Pupfish, desert (Cyprinodon macularius) - Wherever found  | E      | \$517,456     | \$1,213,900,064    |
| 217  | Chub, Virgin River (Gila seminuda (=robusta)) - Wherever found                                  | Е      | \$516,900     | \$1,214,416,964    |
| 218  | Skipperling, Poweshiek (Oarisma poweshiek) - Wherever found                                     | E      | \$516,464     | \$1,214,933,428    |
| 219  | Whale, North Pacific Right (Eubalaena japonica) - Wherever found                                | E      | \$510,039     | \$1,215,443,467    |
| 220  | Goby, tidewater (Eucyclogobius newberryi) - Wherever found                                      | E      | \$509,665     | \$1,215,953,132    |
| 221  | Coral, Lobed Star (Orbicella annularis) - Wherever found  | Т      | \$507,164     | \$1,216,460,296    |
| 222  | Ladies'-tresses, Ute (Spiranthes diluvialis) - Wherever found                                   | Т      | \$504,649     | \$1,216,964,945    |
| 223  | Coral, Pillar (Dendrogyra cylindricus) - Wherever found   | T      | \$503,520     | \$1,217,468,465    |
| 224  | Coral, Rough Cactus (Mycetophyllia ferox) - Wherever found                                      | Т      | \$503,364     | \$1,217,971,829    |
| 225  | Vireo, Black-capped (Vireo atricapilla) - Wherever found  | E      | \$483,186     | \$1,218,455,015    |
| 226  | Crane, whooping (Grus americana) - U.S.A. (CO, ID, FL, NM, UT, and the western half of Wyoming) | EXPN   | \$477,059     | \$1,218,932,074    |
| 227  | Tern, roseate (Sterna dougallii dougallii) - northeast U.S. nesting pop.                        | E      | \$470,225     | \$1,219,402,299    |
| 228  | Panda, giant (Ailuropoda melanoleuca) - Wherever found  | E      | \$469,200     | \$1,219,871,499    |
| 229  | Bat, Florida bonneted (Eumops floridanus) - Wherever found                                      | E      | \$465,300     | \$1,220,336,799    |
| 230  | Coral, Boulder star (Orbicella franksi) - Wherever found  | Т      | \$461,318     | \$1,220,798,117    |
| 231  | Rail, light-footed clapper (Rallus longirostris levipes) - Wherever found                       | E      | \$459,816     | \$1,221,257,933    |
| 232  | Sturgeon, Atlantic (Acipenser oxyrinchus oxyrinchus) - Carolina DPS                             | E      | \$459,137     | \$1,221,717,070    |
| 233  | Butterfly, Oregon silverspot (Speyeria zerene hippolyta) - Wherever found                       | Т      | \$458,095     | \$1,222,175,165    |
| 234  | Salamander, California tiger (Ambystoma californiense) - U.S.A. (Central CA DPS)                | Т      | \$457,779     | \$1,222,632,944    |
| 235  | Heelsplitter, Carolina (Lasmigona decorata) - Wherever found                                    | E      | \$454,057     | \$1,223,087,001    |
| 236  | Kangaroo rat, Stephens' (Dipodomys stephensi (incl. D. cascus)) - Wherever found                | E      | \$447,930     | \$1,223,534,931    |
| 237  | Crocodile, American (Crocodylus acutus) - U.S.A. (FL)   | T      | \$447,877     | \$1,223,982,808    |
| 238  | Caribou, Woodland (Rangifer tarandus caribou) - Selkirk Mountain population                     | E      | \$446,061     | \$1,224,428,869    |
| 239  | Mouse, Perdido Key beach (Peromyscus polionotus trissyllepsis) - Wherever found                 | E      | \$445,399     | \$1,224,874,268    |
| 240  | Salamander, Reticulated flatwoods (Ambystoma bishopi) - Wherever found                          | Е      | \$444,373     | \$1,225,318,641    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 241  | Butterfly, Quino checkerspot (Euphydryas editha quino (=E. e. wrighti)) - Wherever found                                   | Е      | \$438,641     | \$1,225,757,282    |
| 242  | Falcon, northern aplomado (Falco femoralis septentrionalis) - U.S.A (AZ, NM)   | EXPN   | \$432,253     | \$1,226,189,535    |
| 243  | Mucket, pink (pearlymussel) (Lampsilis abrupta) - Wherever found   | E      | \$432,186     | \$1,226,621,721    |
| 244  | Sumac, Michaux's (Rhus michauxii) - Wherever found   | E      | \$424,635     | \$1,227,046,356    |
| 245  | Trout, Greenback Cutthroat (Oncorhynchus clarki stomias) - Wherever found  | Т      | \$423,077     | \$1,227,469,433    |
| 246  | Whale, gray (Eschrichtius robustus) - Western North Pacific DPS  | E      | \$421,797     | \$1,227,891,230    |
| 247  | Peppergrass, Slickspot (Lepidium papilliferum) -   | Т      | \$417,702     | \$1,228,308,932    |
| 248  | Skink, bluetail mole (Eumeces egregius lividus) - Wherever found   | Т      | \$411,300     | \$1,228,720,232    |
| 249  | Catchfly, Spalding's (Silene spaldingii) - Wherever found  | Т      | \$410,926     | \$1,229,131,158    |
| 250  | Frog, dusky gopher (Rana sevosa) - Wherever found  | E      | \$410,519     | \$1,229,541,677    |
| 251  | Trout, Gila (Oncorhynchus gilae) - Wherever found  | Т      | \$410,501     | \$1,229,952,178    |
| 252  | Otter, Northern Sea (Enhydra lutris kenyoni) - Southwest Alaska DPS  | Т      | \$402,909     | \$1,230,355,087    |
| 253  | Rail, Guam (Rallus owstoni) - Wherever found, except where listed as an experimental population                            | Е      | \$396,574     | \$1,230,751,661    |
| 254  | Pronghorn, Sonoran (Antilocapra americana sonoriensis) - Wherever found, except where listed as an experimental population | Е      | \$394,068     | \$1,231,145,729    |
| 255  | Snails, Oahu tree (Achatinella spp.) - Wherever found  | E      | \$390,715     | \$1,231,536,444    |
| 256  | Water-umbel, Huachuca (Lilaeopsis schaffneriana var. recurva) - Wherever found   | E      | \$390,200     | \$1,231,926,644    |
| 257  | Mouse, Preble's meadow jumping (Zapus hudsonius preblei) - wherever found  | Т      | \$390,108     | \$1,232,316,752    |
| 258  | Akikiki (Oreomystis bairdi) - Wherever found   | E      | \$385,946     | \$1,232,702,698    |
| 259  | Chimpanzee (Pan troglodytes) - Wherever found  | E      | \$378,000     | \$1,233,080,698    |
| 260  | Bush-mallow, San Clemente Island (Malacothamnus clementinus) - Wherever found  | Е      | \$376,610     | \$1,233,457,308    |
| 261  | Fairy shrimp, longhorn (Branchinecta longiantenna) - Wherever found  | Е      | \$376,067     | \$1,233,833,375    |
| 262  | Amaranth, seabeach (Amaranthus pumilus) - Wherever found   | Т      | \$374,370     | \$1,234,207,745    |
| 263  | Akekee (Loxops caeruleirostris) - Wherever found   | E      | \$373,849     | \$1,234,581,594    |
| 264  | Dragonfly, Hine's emerald (Somatochlora hineana) - Wherever found  | Е      | \$373,758     | \$1,234,955,352    |
| 265  | Clover, running buffalo (Trifolium stoloniferum) - Wherever found  | Е      | \$373,442     | \$1,235,328,794    |
| 266  | Skipper, Dakota (Hesperia dacotae) - Wherever found  | Т      | \$373,074     | \$1,235,701,868    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 267  | Mouse, Alabama beach (Peromyscus polionotus ammobates) - Wherever found          | Е      | \$371,090     | \$1,236,072,958    |
| 268  | Kingfisher, Guam (Todiramphus cinnamominus) -<br>Wherever found                  | Е      | \$357,189     | \$1,236,430,147    |
| 269  | Toad, Houston (Bufo houstonensis) - Wherever found                               | E      | \$355,000     | \$1,236,785,147    |
| 270  | No Common Name (Acropora globiceps) - Wherever found                             | Т      | \$349,425     | \$1,237,134,572    |
| 271  | Howellia, water (Howellia aquatilis) - Wherever found                            | Т      | \$348,486     | \$1,237,483,058    |
| 272  | Salamander, Sonora tiger (Ambystoma tigrinum stebbinsi) - Wherever found         | Е      | \$347,032     | \$1,237,830,090    |
| 273  | Mouse, New Mexico meadow jumping (Zapus hudsonius luteus) - Wherever found       | Е      | \$345,292     | \$1,238,175,382    |
| 274  | No Common Name (Acropora jacquelinae) - Wherever found                           | Т      | \$345,100     | \$1,238,520,482    |
| 274  | No Common Name (Euphyllia paradivisa) - Wherever found                           | Т      | \$345,100     | \$1,238,865,582    |
| 274  | No Common Name (Acropora speciosa) - Wherever found                              | Т      | \$345,100     | \$1,239,210,682    |
| 274  | No Common Name (Isopora crateriformis) - Wherever found                          | Т      | \$345,100     | \$1,239,555,782    |
| 275  | No Common Name (Acropora rudis) - Wherever found                                 | Т      | \$345,000     | \$1,239,900,782    |
| 276  | Pocket gopher, Roy Prairie (Thomomys mazama glacialis) - Wherever found          | Т      | \$340,833     | \$1,240,241,615    |
| 277  | Frog, mountain yellow-legged (Rana muscosa) - Northern California DPS            | E      | \$339,152     | \$1,240,580,767    |
| 278  | No Common Name (Seriatopora aculeata) - Wherever found                           | Т      | \$339,125     | \$1,240,919,892    |
| 279  | Bobwhite, masked (quail) (Colinus virginianus ridgwayi) - Wherever found         | E      | \$337,212     | \$1,241,257,104    |
| 280  | Buckwheat, scrub (Eriogonum longifolium var. gnaphalifolium) - Wherever found    | Т      | \$336,271     | \$1,241,593,375    |
| 281  | No Common Name (Pavona diffluens) - Wherever found                               | Т      | \$335,424     | \$1,241,928,799    |
| 282  | No Common Name (Acropora lokani) - Wherever found                                | T      | \$335,000     | \$1,242,263,799    |
| 282  | No Common Name (Acropora pharaonis) - Wherever found                             | Т      | \$335,000     | \$1,242,598,799    |
| 282  | No Common Name (Porites napopora) - Wherever found                               | Т      | \$335,000     | \$1,242,933,799    |
| 283  | Sucker, Warner (Catostomus warnerensis) - Wherever found                         | Т      | \$327,204     | \$1,243,261,003    |
| 284  | Bonamia, Florida (Bonamia grandiflora) - Wherever found                          | Т      | \$323,908     | \$1,243,584,911    |
| 285  | Pocketbook, fat (Potamilus capax) - Wherever found                               | E      | \$320,884     | \$1,243,905,795    |
| 286  | Kangaroo rat, giant (Dipodomys ingens) - Wherever found                          | E      | \$320,683     | \$1,244,226,478    |
| 287  | Prairie dog, Utah (Cynomys parvidens) - Wherever found                           | Т      | \$319,085     | \$1,244,545,563    |
| 288  | Jewelflower, Metcalf Canyon (Streptanthus albidus ssp. albidus) - Wherever found | Е      | \$315,370     | \$1,244,860,933    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 289  | Bat, Ozark big-eared (Corynorhinus (=Plecotus) townsendii ingens) - Wherever found   | Е      | \$314,682     | \$1,245,175,615    |
| 290  | Snail, Snake River physa (Physa natricina) - Wherever found  | Е      | \$313,592     | \$1,245,489,207    |
| 291  | Pigeon wings (Clitoria fragrans) - Wherever found  | Т      | \$311,008     | \$1,245,800,215    |
| 292  | Sea turtle, loggerhead (Caretta caretta) - Northeast Atlantic Ocean DPS  | E      | \$310,010     | \$1,246,110,225    |
| 293  | Squirrel, Northern Idaho Ground (Urocitellus brunneus) - Wherever found  | Т      | \$308,283     | \$1,246,418,508    |
| 294  | Toad, Wyoming (Bufo hemiophrys baxteri) - Wherever found   | Е      | \$305,808     | \$1,246,724,316    |
| 295  | Townsendia, Last Chance (Townsendia aprica) - Wherever found   | Т      | \$305,252     | \$1,247,029,568    |
| 296  | Mussel, sheepnose (Plethobasus cyphyus) - Wherever found   | Е      | \$305,081     | \$1,247,334,649    |
| 297  | Spectaclecase (mussel) (Cumberlandia monodonta) - Wherever found   | Е      | \$302,089     | \$1,247,636,738    |
| 298  | Orangutan (Pongo pygmaeus) - Wherever found  | E      | \$291,200     | \$1,247,927,938    |
| 299  | Tortoise, desert (Gopherus agassizii) - AZ south and east of Colorado R., and Mexico, when found outside of Mexico or said range in AZ | SAT    | \$291,113     | \$1,248,219,051    |
| 300  | Milk-vetch, Peirson's (Astragalus magdalenae var. peirsonii) - Wherever found  | Т      | \$287,872     | \$1,248,506,923    |
| 301  | Spinymussel, Tar River (Elliptio steinstansana) - Wherever found   | E      | \$286,942     | \$1,248,793,865    |
| 302  | Pink, swamp (Helonias bullata) - Wherever found  | Т      | \$283,214     | \$1,249,077,079    |
| 303  | Desert-parsley, Bradshaw's (Lomatium bradshawii) - Wherever found  | Е      | \$279,503     | \$1,249,356,582    |
| 304  | Vole, Amargosa (Microtus californicus scirpensis) - Wherever found   | Е      | \$279,419     | \$1,249,636,001    |
| 305  | Sheep, Sierra Nevada bighorn (Ovis canadensis sierrae) - Sierra Nevada   | E      | \$277,678     | \$1,249,913,679    |
| 306  | Rhinoceros, great Indian (Rhinoceros unicornis) - Nepal, India   | Е      | \$277,200     | \$1,250,190,879    |
| 307  | Elepaio, Oahu (Chasiempis ibidis) - Wherever found   | Е      | \$275,254     | \$1,250,466,133    |
| 308  | Mapleleaf, winged (Quadrula fragosa) - Wherever found, except where listed as an experimental population                               | Е      | \$274,016     | \$1,250,740,149    |
| 309  | Sea turtle, loggerhead (Caretta caretta) - South Atlantic Ocean DPS  | Т      | \$273,258     | \$1,251,013,407    |
| 310  | Fiddleneck, large-flowered (Amsinckia grandiflora) - Wherever found  | Е      | \$273,050     | \$1,251,286,457    |
| 311  | Goldfields, Contra Costa (Lasthenia conjugens) - Wherever found  | E      | \$263,197     | \$1,251,549,654    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 312  | Swiftlet, Mariana gray (Aerodramus vanikorensis bartschi) - Wherever found   | Е      | \$262,043     | \$1,251,811,697    |
| 313  | Butterfly, Mitchell's satyr (Neonympha mitchellii mitchellii) - Wherever found   | E      | \$261,140     | \$1,252,072,837    |
| 314  | Beargrass, Britton's (Nolina brittoniana) - Wherever found   | E      | \$259,700     | \$1,252,332,537    |
| 315  | Wolf, maned (Chrysocyon brachyurus) - Wherever found   | E      | \$259,000     | \$1,252,591,537    |
| 316  | Shiner, Arkansas River (Notropis girardi) - Arkansas River Basin (AR, KS, NM, OK, TX)  | Т      | \$258,778     | \$1,252,850,315    |
| 317  | Crane, whooping (Grus americana) - U.S.A. (AL, AR, CO, FL, GA, ID, IL, IN, IA, KY, LA, MI, MN, MS, MO, NC, NM, OH, SC, TN, UT, VA, WI, WV, western half of WY) | EXPN   | \$258,404     | \$1,253,108,719    |
| 318  | Pimpleback, orangefoot (pearlymussel) (Plethobasus cooperianus) - Wherever found   | E      | \$257,820     | \$1,253,366,539    |
| 319  | Snake, San Francisco garter (Thamnophis sirtalis tetrataenia) - Wherever found   | E      | \$255,681     | \$1,253,622,220    |
| 320  | Cavefish, Ozark (Amblyopsis rosae) - Wherever found  | Т      | \$255,566     | \$1,253,877,786    |
| 321  | No Common Name (Hesperomannia arborescens) - Wherever found  | E      | \$254,541     | \$1,254,132,327    |
| 322  | Yerba santa, Lompoc (Eriodictyon capitatum) - Wherever found   | E      | \$253,894     | \$1,254,386,221    |
| 323  | Ceanothus, coyote (Ceanothus ferrisae) - Wherever found  | Е      | \$252,920     | \$1,254,639,141    |
| 324  | Rhinoceros, northern white (Ceratotherium simum cottoni) - Zaire, Uganda, Sudan, Central African Republic  | E      | \$251,000     | \$1,254,890,141    |
| 325  | Pocket gopher, Yelm (Thomomys mazama yelmensis) - Wherever found   | Т      | \$250,505     | \$1,255,140,646    |
| 326  | Four-o'clock, MacFarlane's (Mirabilis macfarlanei) - Wherever found  | Т      | \$250,126     | \$1,255,390,772    |
| 327  | Toad, Yosemite (Anaxyrus canorus) - Wherever found   | Т      | \$249,034     | \$1,255,639,806    |
| 328  | Butterfly, Saint Francis' satyr (Neonympha mitchellii francisci) - Wherever found  | E      | \$248,939     | \$1,255,888,745    |
| 329  | Shiner, Cape Fear (Notropis mekistocholas) - Wherever found  | E      | \$247,380     | \$1,256,136,125    |
| 330  | Wolf, red (Canis rufus) - Wherever found, except where listed as an experimental population  | E      | \$246,268     | \$1,256,382,393    |
| 331  | Wooly-threads, San Joaquin (Monolopia (=Lembertia) congdonii) - Wherever found   | E      | \$245,580     | \$1,256,627,973    |
| 332  | Nanu (Gardenia mannii) - Wherever found  | E      | \$245,279     | \$1,256,873,252    |
| 333  | Gallinule, Hawaiian common (Gallinula galeata sandvicensis) - Wherever found   | Е      | \$244,074     | \$1,257,117,326    |
| 334  | No Common Name (Abutilon sandwicense) - Wherever found   | Е      | \$243,051     | \$1,257,360,377    |
| 335  | Ha`iwale (Cyrtandra dentata) - Wherever found  | E      | \$241,982     | \$1,257,602,359    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 336  | Woodland-star, San Clemente Island (Lithophragma maximum) - Wherever found   | Е      | \$240,208     | \$1,257,842,567    |
| 337  | Pangolin, Temnick's ground (Manis temminckii) - Wherever found   | Е      | \$238,500     | \$1,258,081,067    |
| 338  | lagu, Hayun (=(Guam), Tronkon guafi (Rota)) (Serianthes nelsonii) - Wherever found                                   | Е      | \$234,647     | \$1,258,315,714    |
| 339  | Loosestrife, rough-leaved (Lysimachia asperulaefolia) - Wherever found   | Е      | \$232,167     | \$1,258,547,881    |
| 340  | Sparrow, Florida grasshopper (Ammodramus savannarum floridanus) - Wherever found                                     | Е      | \$231,330     | \$1,258,779,211    |
| 341  | Crane, whooping (Grus americana) - U.S.A (Southwestern Louisiana)  | EXPN   | \$228,260     | \$1,259,007,471    |
| 342  | Bat, Mariana fruit (=Mariana flying fox) (Pteropus mariannus mariannus) - Wherever found                             | Т      | \$225,901     | \$1,259,233,372    |
| 343  | Rhinoceros, Javan (Rhinoceros sondaicus) - Sikkim,<br>Thailand, Malaysia, Indonesia, Indochina, Burma,<br>Bangladesh | Е      | \$225,000     | \$1,259,458,372    |
| 344  | Logperch, Roanoke (Percina rex) - Wherever found   | E      | \$224,203     | \$1,259,682,575    |
| 345  | No Common Name (Tuberolabium guamense) - Wherever found  | Т      | \$222,090     | \$1,259,904,665    |
| 346  | Darter, Okaloosa (Etheostoma okaloosae) - Wherever found   | Т      | \$222,072     | \$1,260,126,737    |
| 347  | Daisy, Willamette (Erigeron decumbens) - Wherever found  | E      | \$220,188     | \$1,260,346,925    |
| 348  | Seal, Guadalupe fur (Arctocephalus townsendi) - Wherever found   | Т      | \$218,225     | \$1,260,565,150    |
| 349  | Ho`awa (Pittosporum napaliense) - Wherever found   | E      | \$217,857     | \$1,260,783,007    |
| 350  | Hawk, Hawaiian (='lo) (Buteo solitarius) - Wherever found  | E      | \$216,759     | \$1,260,999,766    |
| 351  | Tui chub, Mohave (Gila bicolor ssp. mohavensis) -<br>Wherever found  | Е      | \$216,558     | \$1,261,216,324    |
| 352  | Meadowfoam, large-flowered woolly (Limnanthes pumila grandiflora) - Wherever found                                   | Е      | \$215,907     | \$1,261,432,231    |
| 353  | Kangaroo rat, Fresno (Dipodomys nitratoides exilis) - Wherever found   | Е      | \$214,381     | \$1,261,646,612    |
| 354  | Butterfly, Uncompangre fritillary (Boloria acrocnema) - Wherever found   | Е      | \$213,215     | \$1,261,859,827    |
| 355  | Gartersnake, narrow-headed (Thamnophis rufipunctatus) - Wherever found   | Т      | \$213,200     | \$1,262,073,027    |
| 356  | Cheetah (Acinonyx jubatus) - Wherever found  | E      | \$212,100     | \$1,262,285,127    |
| 357  | Moorhen, Mariana common (Gallinula chloropus guami) - Wherever found   | E      | \$209,257     | \$1,262,494,384    |
| 358  | Snail, Bliss Rapids (Taylorconcha serpenticola) - Wherever found   | Т      | \$208,983     | \$1,262,703,367    |
| 359  | Dace, Moapa (Moapa coriacea) - Wherever found  | E      | \$204,404     | \$1,262,907,771    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 360  | Heartleaf, dwarf-flowered (Hexastylis naniflora) - Wherever found                          | Т      | \$202,876     | \$1,263,110,647    |
| 361  | Pikeminnow (=squawfish), Colorado (Ptychocheilus lucius) - Salt and Verde R. drainages, AZ | EXPN   | \$202,689     | \$1,263,313,336    |
| 362  | Falcon, Eurasian peregrine (Falco peregrinus peregrinus) - Wherever found                  | E      | \$202,655     | \$1,263,515,991    |
| 363  | Totoaba (seatrout or weakfish) (Cynoscion macdonaldi) - Wherever found                     | E      | \$200,000     | \$1,263,715,991    |
| 364  | Lion (Panthera leo ssp. leo) - Wherever found  | E      | \$199,900     | \$1,263,915,891    |
| 365  | Darter, Niangua (Etheostoma nianguae) - Wherever found                                     | Т      | \$197,826     | \$1,264,113,717    |
| 366  | Kuahiwi laukahi (Plantago princeps) - Wherever found                                       | E      | \$195,991     | \$1,264,309,708    |
| 367  | Spiraea, Virginia (Spiraea virginiana) - Wherever found                                    | Т      | \$195,197     | \$1,264,504,905    |
| 368  | Piperia, Yadon's (Piperia yadonii) - Wherever found  | Е      | \$194,021     | \$1,264,698,926    |
| 369  | Thistle, Pitcher's (Cirsium pitcheri) - Wherever found                                     | Т      | \$193,204     | \$1,264,892,130    |
| 370  | Towhee, Inyo California (Pipilo crissalis eremophilus) - Wherever found                    | Т      | \$192,406     | \$1,265,084,536    |
| 371  | Rabbitsfoot (Quadrula cylindrica cylindrica) - Wherever found                              | Т      | \$189,057     | \$1,265,273,593    |
| 372  | Cactus, Colorado hookless (Sclerocactus glaucus) - Wherever found                          | Т      | \$188,088     | \$1,265,461,681    |
| 373  | Salamander, Austin blind (Eurycea waterlooensis) - Wherever found                          | E      | \$188,000     | \$1,265,649,681    |
| 374  | Salamander, Jollyville Plateau (Eurycea tonkawae) -<br>Wherever found                      | Т      | \$185,000     | \$1,265,834,681    |
| 375  | Ohai (Sesbania tomentosa) - Wherever found   | Е      | \$176,279     | \$1,266,010,960    |
| 376  | Spinymussel, James (Pleurobema collina) - Wherever found                                   | E      | \$173,483     | \$1,266,184,443    |
| 377  | Stickleback, unarmored threespine (Gasterosteus aculeatus williamsoni) - Wherever found    | E      | \$171,971     | \$1,266,356,414    |
| 378  | Catfish, Yaqui (Ictalurus pricei) - Wherever found   | Т      | \$170,900     | \$1,266,527,314    |
| 379  | Orchid, western prairie fringed (Platanthera praeclara) - Wherever found                   | Т      | \$170,320     | \$1,266,697,634    |
| 380  | Dace, desert (Eremichthys acros) - Wherever found  | Т      | \$169,262     | \$1,266,866,896    |
| 381  | Springsnail, Bruneau Hot (Pyrgulopsis bruneauensis) - Wherever found                       | E      | \$167,600     | \$1,267,034,496    |
| 382  | Duck, Laysan (Anas laysanensis) - Wherever found   | Е      | \$166,648     | \$1,267,201,144    |
| 383  | Pondberry (Lindera melissifolia) - Wherever found  | E      | \$166,458     | \$1,267,367,602    |
| 384  | Popcornflower, rough (Plagiobothrys hirtus) - Wherever found                               | E      | \$166,000     | \$1,267,533,602    |
| 385  | Fairy shrimp, San Diego (Branchinecta sandiegonensis) - Wherever found                     | E      | \$163,618     | \$1,267,697,220    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 386  | Skipper, Carson wandering (Pseudocopaeodes eunus obscurus) - Wherever found   | Е      | \$162,718     | \$1,267,859,938    |
| 387  | Kangaroo rat, Tipton (Dipodomys nitratoides nitratoides) - Wherever found   | Е      | \$161,924     | \$1,268,021,862    |
| 388  | Spineflower, Orcutt's (Chorizanthe orcuttiana) - Wherever found   | Е      | \$160,302     | \$1,268,182,164    |
| 389  | Thrush, small Kauai (=puaiohi) (Myadestes palmeri) -<br>Wherever found  | Е      | \$159,325     | \$1,268,341,489    |
| 390  | Fox, San Miguel Island (Urocyon littoralis littoralis) - wherever found   | Е      | \$157,712     | \$1,268,499,201    |
| 391  | Riffleshell, northern (Epioblasma torulosa rangiana) - Wherever found   | Е      | \$157,146     | \$1,268,656,347    |
| 392  | Mahoe (Alectryon macrococcus) - Wherever found  | E      | \$156,917     | \$1,268,813,264    |
| 393  | Ma`o hau hele, (=native yellow hibiscus) (Hibiscus brackenridgei) - Wherever found  | Е      | \$156,472     | \$1,268,969,736    |
| 394  | Sea turtle, green (Chelonia mydas) - East Indian-West Pacific DPS   | Т      | \$155,000     | \$1,269,124,736    |
| 395  | Clubshell (Pleurobema clava) - Wherever found; Except where listed as Experimental Populations  | Е      | \$154,519     | \$1,269,279,255    |
| 396  | Dace, Foskett speckled (Rhinichthys osculus ssp.) - Wherever found  | Т      | \$153,490     | \$1,269,432,745    |
| 397  | Madtom, Neosho (Noturus placidus) - Wherever found  | Т      | \$147,942     | \$1,269,580,687    |
| 398  | Quillwort, Louisiana (Isoetes Iouisianensis) - Wherever found   | Е      | \$147,932     | \$1,269,728,619    |
| 399  | Rabbit, riparian brush (Sylvilagus bachmani riparius) - Wherever found  | Е      | \$146,666     | \$1,269,875,285    |
| 400  | Kangaroo rat, San Bernardino Merriam's (Dipodomys merriami parvus) - Wherever found   | Е      | \$143,970     | \$1,270,019,255    |
| 401  | Fanshell (Cyprogenia stegaria) - Wherever found   | E      | \$142,717     | \$1,270,161,972    |
| 402  | Harvestman, Bone Cave (Texella reyesi) - Wherever found   | E      | \$142,500     | \$1,270,304,472    |
| 403  | Sunflower, Schweinitz's (Helianthus schweinitzii) -<br>Wherever found   | Е      | \$141,720     | \$1,270,446,192    |
| 404  | Cui-ui (Chasmistes cujus) - Wherever found  | E      | \$141,644     | \$1,270,587,836    |
| 405  | Lomatium, Cook's (Lomatium cookii) - Wherever found   | E      | \$140,980     | \$1,270,728,816    |
| 406  | Amole, purple (Chlorogalum purpureum) - Wherever found  | Т      | \$140,914     | \$1,270,869,730    |
| 407  | Pearlymussel, cracking (Hemistena lata) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL | EXPN   | \$140,460     | \$1,271,010,190    |
| 408  | Mehamehame (Flueggea neowawraea) - Wherever found   | E      | \$140,045     | \$1,271,150,235    |
| 409  | No Common Name (Phyllostegia hirsuta) - Wherever found  | E      | \$139,777     | \$1,271,290,012    |
| 410  | Dace, blackside (Phoxinus cumberlandensis) - Wherever found   | Т      | \$137,281     | \$1,271,427,293    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 411  | Pearlshell, Louisiana (Margaritifera hembeli) - Wherever found                           | Т      | \$137,260     | \$1,271,564,553    |
| 412  | Haha (Cyanea koolauensis) - Wherever found   | E      | \$135,895     | \$1,271,700,448    |
| 413  | Parrotbill, Maui (Kiwikiu) (Pseudonestor xanthophrys) - Wherever found                   | Е      | \$134,589     | \$1,271,835,037    |
| 414  | Penstemon, blowout (Penstemon haydenii) - Wherever found                                 | E      | \$133,835     | \$1,271,968,872    |
| 415  | Ivesia, Webber's (Ivesia webberi) - Wherever found                                       | Т      | \$133,278     | \$1,272,102,150    |
| 416  | Kamakahala (Labordia cyrtandrae) - Wherever found  | E      | \$133,083     | \$1,272,235,233    |
| 417  | Butterfly, El Segundo blue (Euphilotes battoides allyni) - Wherever found                | E      | \$132,989     | \$1,272,368,222    |
| 418  | Whipsnake (=striped racer), Alameda (Masticophis lateralis euryxanthus) - Wherever found |        | \$132,240     | \$1,272,500,462    |
| 419  | Haha (Cyanea superba) - Wherever found   | E      | \$131,924     | \$1,272,632,386    |
| 420  | No Common Name (Neraudia angulata) - Wherever found                                      | E      | \$131,878     | \$1,272,764,264    |
| 421  | No Common Name (Neraudia ovata) - Wherever found   | E      | \$131,788     | \$1,272,896,052    |
| 422  | Bankclimber, purple (mussel) (Elliptoideus sloatianus) - Wherever found                  | Т      | \$131,570     | \$1,273,027,622    |
| 423  | Kulu`i (Nototrichium humile) - Wherever found  | E      | \$131,299     | \$1,273,158,921    |
| 424  | `Ahinahina (Argyroxiphium sandwicense ssp. macrocephalum) - Wherever found               | Т      | \$131,167     | \$1,273,290,088    |
| 425  | Threeridge, fat (mussel) (Amblema neislerii) - Wherever found                            | E      | \$131,056     | \$1,273,421,144    |
| 426  | No Common Name (Schiedea kaalae) - Wherever found  | E      | \$130,959     | \$1,273,552,103    |
| 427  | No Common Name (Phyllostegia mollis) - Wherever found                                    | E      | \$129,320     | \$1,273,681,423    |
| 428  | No Common Name (Kadua degeneri) - Wherever found   | E      | \$128,798     | \$1,273,810,221    |
| 429  | Pupfish, Warm Springs (Cyprinodon nevadensis pectoralis) - Wherever found                | E      | \$128,526     | \$1,273,938,747    |
| 430  | No Common Name (Schiedea obovata) - Wherever found                                       | E      | \$128,451     | \$1,274,067,198    |
| 431  | No Common Name (Stenogyne kanehoana) - Wherever found                                    | E      | \$128,205     | \$1,274,195,403    |
| 432  | Haha (Cyanea grimesiana ssp. obatae) - Wherever found                                    | E      | \$128,147     | \$1,274,323,550    |
| 433  | No Common Name (Schiedea nuttallii) - Wherever found                                     | E      | \$128,013     | \$1,274,451,563    |
| 434  | No Common Name (Phyllostegia kaalaensis) - Wherever found                                | Е      | \$127,853     | \$1,274,579,416    |
| 435  | `Akoko (Euphorbia herbstii) - Wherever found   | E      | \$127,360     | \$1,274,706,776    |
| 436  | Oha (Delissea subcordata) - Wherever found   | E      | \$126,706     | \$1,274,833,482    |
| 437  | Palila (honeycreeper) (Loxioides bailleui) - Wherever found                              | E      | \$126,442     | \$1,274,959,924    |
| 438  | Nioi (Eugenia koolauensis) - Wherever found  | E      | \$126,323     | \$1,275,086,247    |
| 439  | No Common Name (Kadua parvula) - Wherever found  | E      | \$125,851     | \$1,275,212,098    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 440  | Kamanomano (Cenchrus agrimonioides) - Wherever found  | E      | \$125,667     | \$1,275,337,765    |
| 441  | No Common Name (Tetramolopium filiforme) - Wherever found   | Е      | \$125,176     | \$1,275,462,941    |
| 442  | Na`ena`e (Dubautia herbstobatae) - Wherever found   | E      | \$124,525     | \$1,275,587,466    |
| 442  | Lo`ulu (Pritchardia kaalae) - Wherever found  | E      | \$124,525     | \$1,275,711,991    |
| 443  | Rhinoceros, Southern white (Ceratotherium simum ssp. simum) - Wherever found  | SAT    | \$124,500     | \$1,275,836,491    |
| 444  | No Common Name (Schiedea trinervis) - Wherever found  | E      | \$124,380     | \$1,275,960,871    |
| 445  | Haha (Cyanea longiflora) - Wherever found   | E      | \$124,221     | \$1,276,085,092    |
| 446  | No Common Name (Sanicula mariversa) - Wherever found  | E      | \$124,076     | \$1,276,209,168    |
| 447  | Lau `ehu (Panicum niihauense) - Wherever found  | E      | \$123,588     | \$1,276,332,756    |
| 448  | `Akoko (Euphorbia celastroides var. kaenana) - Wherever found   | Е      | \$123,434     | \$1,276,456,190    |
| 449  | Lampmussel, Alabama (Lampsilis virescens) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL | EXPN   | \$123,340     | \$1,276,579,530    |
| 450  | Leopard (Panthera pardus) - except where threatened   | E      | \$122,600     | \$1,276,702,130    |
| 451  | No Common Name (Spermolepis hawaiiensis) - Wherever found   | Е      | \$122,499     | \$1,276,824,629    |
| 452  | Picture-wing fly, Hawaiian (Drosophila montgomeryi) - Wherever found  | Е      | \$121,759     | \$1,276,946,388    |
| 452  | Fly, Hawaiian picture-wing (Drosophila substenoptera) - Wherever found  | Е      | \$121,759     | \$1,277,068,147    |
| 453  | Po`e (Portulaca sclerocarpa) - Wherever found   | E      | \$121,607     | \$1,277,189,754    |
| 454  | Shrimp, California freshwater (Syncaris pacifica) - Wherever found  | Е      | \$120,984     | \$1,277,310,738    |
| 455  | Moth, Blackburn's sphinx (Manduca blackburni) - Wherever found  | Е      | \$120,513     | \$1,277,431,251    |
| 456  | Sturgeon, Shovelnose (Scaphirhynchus platorynchus) - Wherever found   | SAT    | \$119,217     | \$1,277,550,468    |
| 457  | Ma`oli`oli (Schiedea hawaiiensis) - Wherever found  | E      | \$118,908     | \$1,277,669,376    |
| 458  | Pocket gopher, Olympia (Thomomys mazama pugetensis) - Wherever found  | Т      | \$117,630     | \$1,277,787,006    |
| 459  | Sea turtle, loggerhead (Caretta caretta) - North Indian Ocean DPS   | Е      | \$117,037     | \$1,277,904,043    |
| 460  | Albatross, short-tailed (Phoebastria (=Diomedea) albatrus) - Wherever found   | Е      | \$116,783     | \$1,278,020,826    |
| 461  | No Common Name (Silene lanceolata) - Wherever found   | E      | \$116,296     | \$1,278,137,122    |
| 462  | A`e (Zanthoxylum hawaiiense) - Wherever found   | E      | \$116,012     | \$1,278,253,134    |
| 463  | Popolo ku mai (Solanum incompletum) - Wherever found  | E      | \$115,998     | \$1,278,369,132    |
| 464  | Tiger beetle, Salt Creek (Cicindela nevadica lincolniana) - Wherever found  | Е      | \$114,073     | \$1,278,483,205    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 465  | Aster, decurrent false (Boltonia decurrens) - Wherever found  | Т      | \$113,492     | \$1,278,596,697    |
| 466  | No Common Name (Stenogyne angustifolia angustifolia) - Wherever found   | Е      | \$113,324     | \$1,278,710,021    |
| 467  | Rabbit, Lower Keys marsh (Sylvilagus palustris hefneri) - Wherever found  | E      | \$113,224     | \$1,278,823,245    |
| 468  | Woolly-star, Santa Ana River (Eriastrum densifolium ssp. sanctorum) - Wherever found  | E      | \$112,626     | \$1,278,935,871    |
| 469  | Coneflower, smooth (Echinacea laevigata) - Wherever found   | Е      | \$112,527     | \$1,279,048,398    |
| 470  | No Common Name (Asplenium peruvianum var. insulare) - Wherever found  | Е      | \$112,103     | \$1,279,160,501    |
| 471  | No Common Name (Schiedea haleakalensis) - Wherever found  | Е      | \$111,691     | \$1,279,272,192    |
| 472  | No Common Name (Silene hawaiiensis) - Wherever found  | Т      | \$111,670     | \$1,279,383,862    |
| 473  | Honohono (Haplostachys haplostachya) - Wherever found   | E      | \$111,212     | \$1,279,495,074    |
| 474  | Chub, spotfin (Erimonax monachus) - Wherever found, except where listed as an experimental population                           | Т      | \$110,969     | \$1,279,606,043    |
| 475  | Kio`ele (Kadua coriacea) - Wherever found   | E      | \$109,346     | \$1,279,715,389    |
| 475  | No Common Name (Vigna o-wahuensis) - Wherever found   | E      | \$109,346     | \$1,279,824,735    |
| 476  | Falcon, northern aplomado (Falco femoralis septentrionalis) - Wherever found, except where listed as an experimental population | Е      | \$108,459     | \$1,279,933,194    |
| 477  | Turtle, bog (Clemmys muhlenbergii) - U.S.A. (GA, NC, SC, TN, VA)  | SAT    | \$107,565     | \$1,280,040,759    |
| 478  | Manzanita, Franciscan (Arctostaphylos franciscana) - Wherever found   | Е      | \$107,500     | \$1,280,148,259    |
| 479  | Chub, Pahranagat roundtail (Gila robusta jordani) - Wherever found  | Е      | \$107,289     | \$1,280,255,548    |
| 480  | Aupaka (Isodendrion hosakae) - Wherever found   | E      | \$107,110     | \$1,280,362,658    |
| 481  | Larkspur, San Clemente Island (Delphinium variegatum ssp. kinkiense) - Wherever found   | E      | \$106,934     | \$1,280,469,592    |
| 482  | Haha (Cyanea asplenifolia) - Wherever found   | E      | \$106,802     | \$1,280,576,394    |
| 483  | No Common Name (Tetramolopium arenarium) - Wherever found   | E      | \$106,685     | \$1,280,683,079    |
| 484  | Fairy shrimp, Riverside (Streptocephalus woottoni) - Wherever found   | Е      | \$106,600     | \$1,280,789,679    |
| 485  | Milk-vetch, Sentry (Astragalus cremnophylax var. cremnophylax) - Wherever found   | Е      | \$106,272     | \$1,280,895,951    |
| 486  | Buckwheat, steamboat (Eriogonum ovalifolium var. williamsiae) - Wherever found  | Е      | \$106,041     | \$1,281,001,992    |
| 487  | No Common Name (Lipochaeta venosa) - Wherever found   | E      | \$105,795     | \$1,281,107,787    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 488  | Indian paintbrush, San Clemente Island (Castilleja grisea) - Wherever found   | Т      | \$105,422     | \$1,281,213,209    |
| 489  | Deer, musk (Moschus spp. (all species)) - Afghanistan,<br>Bhutan, Burma, China [Tibet, Yunnan], India, Nepal,<br>Pakistan, Sikkim   | E      | \$105,135     | \$1,281,318,344    |
| 490  | Bear, brown (Ursus arctos pruinosus) - Wherever found   | E      | \$104,800     | \$1,281,423,144    |
| 491  | Harperella (Ptilimnium nodosum) - Wherever found  | E      | \$104,412     | \$1,281,527,556    |
| 492  | Lotus, San Clemente Island (Acmispon dendroideus var. traskiae (=Lotus d. ssp. traskiae)) - Wherever found  | Т      | \$104,326     | \$1,281,631,882    |
| 493  | Darter, leopard (Percina pantherina) - Wherever found   | Т      | \$103,823     | \$1,281,735,705    |
| 494  | Pupfish, Ash Meadows Amargosa (Cyprinodon nevadensis mionectes) - Wherever found  | E      | \$103,103     | \$1,281,838,808    |
| 495  | Rockcress, Santa Cruz Island (Sibara filifolia) - Wherever found  | E      | \$102,338     | \$1,281,941,146    |
| 496  | Squirrel, Carolina northern flying (Glaucomys sabrinus coloratus) - Wherever found  | Е      | \$101,396     | \$1,282,042,542    |
| 497  | Limpet, Banbury Springs (Lanx sp.) - Wherever found   | E      | \$101,364     | \$1,282,143,906    |
| 498  | Trout, Apache (Oncorhynchus apache) - Wherever found  | Т      | \$101,113     | \$1,282,245,019    |
| 499  | Tiger beetle, Puritan (Cicindela puritana) - Wherever found   | Т      | \$100,829     | \$1,282,345,848    |
| 500  | Mussel, scaleshell (Leptodea leptodon) - Wherever found   | E      | \$100,752     | \$1,282,446,600    |
| 501  | Haiwale (Cyrtandra ferripilosa) - Wherever found  | E      | \$100,728     | \$1,282,547,328    |
| 501  | No Common Name (Phyllostegia haliakalae) - Wherever found   | Е      | \$100,728     | \$1,282,648,056    |
| 502  | Butterfly, Behren's silverspot (Speyeria zerene behrensii) - Wherever found   | E      | \$100,572     | \$1,282,748,628    |
| 503  | Pupfish, Owens (Cyprinodon radiosus) - Wherever found   | E      | \$100,476     | \$1,282,849,104    |
| 504  | `Oha wai (Clermontia samuelii) - Wherever found   | E      | \$100,306     | \$1,282,949,410    |
| 505  | Seagrass, Johnson's (Halophila johnsonii) - Wherever found  | Т      | \$99,100      | \$1,283,048,510    |
| 506  | Bear-poppy, Dwarf (Arctomecon humilis) - Wherever found   | E      | \$98,350      | \$1,283,146,860    |
| 507  | Bladderpod, Short's (Physaria globosa) - Wherever found   | E      | \$98,310      | \$1,283,245,170    |
| 508  | Baccharis, Encinitas (Baccharis vanessae) - Wherever found  | Т      | \$98,204      | \$1,283,343,374    |
| 509  | Shiner, Topeka (Notropis topeka (=tristis)) - U.S.A. (MO-specified portions of Little Creek, Big Muddy Creek, and Spring Creek watersheds in Adair, Gentry, Harrison, Putnam, Sullivan, and Worth Counties; see 17.84(d)(1)(i)) | EXPN   | \$98,100      | \$1,283,441,474    |
| 510  | Polygala, Lewton's (Polygala lewtonii) - Wherever found   | Е      | \$97,842      | \$1,283,539,316    |
| 511  | Manzanita, Del Mar (Arctostaphylos glandulosa ssp. crassifolia) - Wherever found  | Е      | \$97,000      | \$1,283,636,316    |
| 512  | Fadang (Cycas micronesica) - Wherever found   | Т      | \$96,332      | \$1,283,732,648    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 513  | Mountain beaver, Point Arena (Aplodontia rufa nigra) - Wherever found                  | Е      | \$96,077      | \$1,283,828,725    |
| 514  | Goldfields, Burke's (Lasthenia burkei) - Wherever found                                | E      | \$95,697      | \$1,283,924,422    |
| 515  | Frog, mountain yellow-legged (Rana muscosa) - Southern California DPS                  | Е      | \$95,650      | \$1,284,020,072    |
| 516  | Lizard, Coachella Valley fringe-toed (Uma inornata) - Wherever found                   | Т      | \$95,443      | \$1,284,115,515    |
| 517  | Ring pink (mussel) (Obovaria retusa) - Wherever found                                  | E      | \$94,820      | \$1,284,210,335    |
| 518  | Harvestman, Bee Creek Cave (Texella reddelli) - Wherever found                         | E      | \$94,500      | \$1,284,304,835    |
| 518  | Pseudoscorpion, Tooth Cave (Tartarocreagris texana) - Wherever found                   | E      | \$94,500      | \$1,284,399,335    |
| 518  | Spider, Tooth Cave (Neoleptoneta myopica) - Wherever found                             | Е      | \$94,500      | \$1,284,493,835    |
| 519  | Boa, Puerto Rican (Epicrates inornatus) - Wherever found                               | E      | \$94,130      | \$1,284,587,965    |
| 520  | Ambrosia, San Diego (Ambrosia pumila) - Wherever found                                 | E      | \$93,624      | \$1,284,681,589    |
| 521  | Creeper, Hawaii (Oreomystis mana) - Wherever found                                     | E      | \$92,540      | \$1,284,774,129    |
| 522  | Bird's-beak, salt marsh (Cordylanthus maritimus ssp. maritimus) - Wherever found       | Е      | \$92,498      | \$1,284,866,627    |
| 523  | Blackbird, yellow-shouldered (Agelaius xanthomus) - Wherever found                     | Е      | \$92,100      | \$1,284,958,727    |
| 524  | Squirrel, Mount Graham red (Tamiasciurus hudsonicus grahamensis) - Wherever found      | E      | \$91,300      | \$1,285,050,027    |
| 525  | Butterfly, Lange's metalmark (Apodemia mormo langei) - Wherever found                  | Е      | \$91,104      | \$1,285,141,131    |
| 526  | Akepa, Hawaii (Loxops coccineus) - Wherever found                                      | E      | \$90,973      | \$1,285,232,104    |
| 527  | Goldenrod, Houghton's (Solidago houghtonii) - Wherever found                           | Т      | \$89,949      | \$1,285,322,053    |
| 528  | Joint-vetch, Sensitive (Aeschynomene virginica) - Wherever found                       | Т      | \$89,484      | \$1,285,411,537    |
| 529  | Skipper, Laguna Mountains (Pyrgus ruralis lagunae) - Wherever found                    | Е      | \$88,979      | \$1,285,500,516    |
| 530  | Potentilla, Hickman's (Potentilla hickmanii) - Wherever found                          | Е      | \$88,894      | \$1,285,589,410    |
| 531  | Butterfly, Palos Verdes blue (Glaucopsyche lygdamus palosverdesensis) - Wherever found | E      | \$88,031      | \$1,285,677,441    |
| 532  | Checker-mallow, Nelson's (Sidalcea nelsoniana) - Wherever found                        | Т      | \$87,264      | \$1,285,764,705    |
| 533  | Slabshell, Chipola (Elliptio chipolaensis) - Wherever found                            | Т      | \$86,206      | \$1,285,850,911    |
| 534  | Glade cress, Kentucky (Leavenworthia exigua laciniata) - Wherever found                | Т      | \$85,706      | \$1,285,936,617    |
| 535  | Jewelflower, California (Caulanthus californicus) - Wherever found                     | Е      | \$85,360      | \$1,286,021,977    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 536  | Pigtoe, oval (Pleurobema pyriforme) - Wherever found                            | E      | \$85,196      | \$1,286,107,173    |
| 537  | Tern, roseate (Sterna dougallii dougallii) - Western Hemisphere except NE U.S.  | Т      | \$84,705      | \$1,286,191,878    |
| 538  | Brodiaea, thread-leaved (Brodiaea filifolia) - Wherever found                   | Т      | \$84,547      | \$1,286,276,425    |
| 539  | Darter, bluemask (Etheostoma akatulo) - Wherever found                          | E      | \$84,490      | \$1,286,360,915    |
| 540  | Geranium, Hawaiian red-flowered (Geranium arboreum) - Wherever found            | E      | \$84,373      | \$1,286,445,288    |
| 541  | Monkshood, northern wild (Aconitum noveboracense) - Wherever found              | Т      | \$84,171      | \$1,286,529,459    |
| 542  | Butterfly, Myrtle's silverspot (Speyeria zerene myrtleae) - Wherever found      | E      | \$83,971      | \$1,286,613,430    |
| 543  | Warea, wide-leaf (Warea amplexifolia) - Wherever found                          | E      | \$82,900      | \$1,286,696,330    |
| 544  | No Common Name (Phyllostegia bracteata) - Wherever found                        | Е      | \$81,822      | \$1,286,778,152    |
| 545  | Pigtoe, rough (Pleurobema plenum) - Wherever found                              | E      | \$81,720      | \$1,286,859,872    |
| 546  | Heelsplitter, Alabama (=inflated) (Potamilus inflatus) - Wherever found         | Т      | \$81,633      | \$1,286,941,505    |
| 547  | Dropwort, Canby's (Oxypolis canbyi) - Wherever found                            | E      | \$81,313      | \$1,287,022,818    |
| 548  | Hawk, Puerto Rican sharp-shinned (Accipiter striatus venator) - Wherever found  | E      | \$81,103      | \$1,287,103,921    |
| 549  | Mouse, southeastern beach (Peromyscus polionotus niveiventris) - wherever found | Т      | \$80,800      | \$1,287,184,721    |
| 550  | Darter, relict (Etheostoma chienense) - Wherever found                          | E      | \$80,187      | \$1,287,264,908    |
| 551  | Mouse, Choctawhatchee beach (Peromyscus polionotus allophrys) - Wherever found  | Е      | \$79,838      | \$1,287,344,746    |
| 552  | Darter, boulder (Etheostoma wapiti) - Wherever found                            | E      | \$79,660      | \$1,287,424,406    |
| 553  | Layia, beach (Layia carnosa) - Wherever found                                   | E      | \$79,133      | \$1,287,503,539    |
| 554  | Ziziphus, Florida (Ziziphus celata) - Wherever found                            | E      | \$79,000      | \$1,287,582,539    |
| 555  | Ko`oloa`ula (Abutilon menziesii) - Wherever found                               | E      | \$78,670      | \$1,287,661,209    |
| 556  | Ischaemum, Hilo (Ischaemum byrone) - Wherever found                             | E      | \$78,166      | \$1,287,739,375    |
| 557  | Crocodile, African slender-snouted (Crocodylus cataphractus) - Wherever found   | E      | \$78,000      | \$1,287,817,375    |
| 558  | Chub, Owens Tui (Gila bicolor ssp. snyderi) - Wherever found                    | Е      | \$77,728      | \$1,287,895,103    |
| 559  | Pocketbook, Ouachita rock (Arkansia wheeleri) - Wherever found                  | Е      | \$76,880      | \$1,287,971,983    |
| 560  | Mallow, Kern (Eremalche kernensis) - Wherever found                             | E      | \$76,710      | \$1,288,048,693    |
| 561  | Ko`oko`olau (Bidens micrantha ssp. kalealaha) - Wherever found                  | E      | \$76,167      | \$1,288,124,860    |
| 562  | Beetle, Casey's June (Dinacoma caseyi) - Wherever found                         | E      | \$75,859      | \$1,288,200,719    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 563  | Meshweaver, Madla's Cave (Cicurina madla) - Wherever found   | Е      | \$75,533      | \$1,288,276,252    |
| 564  | Shrew, Buena Vista Lake ornate (Sorex ornatus relictus) - Wherever found                                   | Е      | \$75,504      | \$1,288,351,756    |
| 565  | Milk-vetch, Lane Mountain (Astragalus jaegerianus) - Wherever found  | Е      | \$75,429      | \$1,288,427,185    |
| 566  | Beetle, [no common name] (Rhadine exilis) - Wherever found   | Е      | \$75,333      | \$1,288,502,518    |
| 567  | Button-celery, San Diego (Eryngium aristulatum var. parishii) - Wherever found                             | Е      | \$75,283      | \$1,288,577,801    |
| 568  | Akiapolaau (Hemignathus wilsoni) - Wherever found  | E      | \$75,115      | \$1,288,652,916    |
| 569  | Pitcher-plant, mountain sweet (Sarracenia rubra ssp. jonesii) - Wherever found                             | Е      | \$74,998      | \$1,288,727,914    |
| 570  | Pocketbook, shinyrayed (Lampsilis subangulata) - Wherever found  | E      | \$74,290      | \$1,288,802,204    |
| 571  | Bison, wood (Bison bison athabascae) - Wherever found  | Т      | \$73,000      | \$1,288,875,204    |
| 572  | Tiger beetle, Northeastern beach (Cicindela dorsalis dorsalis) - Wherever found                            | Т      | \$72,944      | \$1,288,948,148    |
| 573  | Bat, Mexican long-nosed (Leptonycteris nivalis) - Wherever found   | E      | \$72,850      | \$1,289,020,998    |
| 574  | Orchid, eastern prairie fringed (Platanthera leucophaea) - Wherever found                                  | Т      | \$72,474      | \$1,289,093,472    |
| 575  | Butterfly, Smith's blue (Euphilotes enoptes smithi) - Wherever found                                       | Е      | \$71,411      | \$1,289,164,883    |
| 576  | Hellbender, Ozark (Cryptobranchus alleganiensis bishopi) - Wherever found                                  | Е      | \$71,200      | \$1,289,236,083    |
| 577  | Butterfly, Mariana eight-spot (Hypolimnas octocula marianensis) - Wherever found                           | Е      | \$71,160      | \$1,289,307,243    |
| 578  | Dace, Ash Meadows speckled (Rhinichthys osculus nevadensis) - Wherever found                               | Е      | \$70,364      | \$1,289,377,607    |
| 579  | Beetle, [no common name] (Rhadine infernalis) - Wherever found   | Е      | \$70,333      | \$1,289,447,940    |
| 580  | No Common Name (Osmoxylon mariannense) - Wherever found  | Е      | \$70,328      | \$1,289,518,268    |
| 581  | Milk-vetch, Ash meadows (Astragalus phoenix) - Wherever found  | Т      | \$70,044      | \$1,289,588,312    |
| 582  | Warbler, elfin-woods (Setophaga angelae) - Wherever found  | Т      | \$70,000      | \$1,289,658,312    |
| 583  | Fly, Delhi Sands flower-loving (Rhaphiomidas terminatus abdominalis) - Wherever found                      | Е      | \$69,748      | \$1,289,728,060    |
| 584  | Mussel, oyster (Epioblasma capsaeformis) - Wherever found; Except where listed as Experimental Populations | Е      | \$69,460      | \$1,289,797,520    |
| 585  | Beetle, American burying (Nicrophorus americanus) - Ex Pop, SW Missouri                                    | EXPN   | \$69,200      | \$1,289,866,720    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 586  | Cactus, Mesa Verde (Sclerocactus mesae-verdae) - Wherever found                          | Т      | \$68,800      | \$1,289,935,520    |
| 587  | Butterfly plant, Colorado (Gaura neomexicana var. coloradensis) - Wherever found         | Т      | \$68,679      | \$1,290,004,199    |
| 588  | No Common Name (Nesogenes rotensis) - Wherever found                                     | E      | \$67,813      | \$1,290,072,012    |
| 589  | Butterfly, callippe silverspot (Speyeria callippe callippe) - Wherever found             | Е      | \$67,745      | \$1,290,139,757    |
| 590  | Potato-bean, Price's (Apios priceana) - Wherever found                                   | Т      | \$67,037      | \$1,290,206,794    |
| 591  | Rattlesnake, New Mexican ridge-nosed (Crotalus willardi obscurus) - Wherever found       | Т      | \$66,900      | \$1,290,273,694    |
| 592  | `Aiea (Nothocestrum breviflorum) - Wherever found  | E      | \$66,884      | \$1,290,340,578    |
| 593  | Beauty, Harper's (Harperocallis flava) - Wherever found                                  | E      | \$66,495      | \$1,290,407,073    |
| 594  | Bean, purple (Villosa perpurpurea) - Wherever found                                      | E      | \$66,480      | \$1,290,473,553    |
| 595  | Cactus, Wright fishhook (Sclerocactus wrightiae) - Wherever found                        | Е      | \$66,400      | \$1,290,539,953    |
| 596  | Wallflower, Menzies' (Erysimum menziesii) - Wherever found                               | Е      | \$66,052      | \$1,290,606,005    |
| 597  | Ayenia, Texas (Ayenia limitaris) - Wherever found  | E      | \$65,800      | \$1,290,671,805    |
| 598  | Spinedace, White River (Lepidomeda albivallis) - Wherever found                          | Е      | \$65,738      | \$1,290,737,543    |
| 599  | Cactus, Arizona hedgehog (Echinocereus triglochidiatus var. arizonicus) - Wherever found | Е      | \$65,197      | \$1,290,802,740    |
| 600  | Milk-vetch, Fish Slough (Astragalus lentiginosus var. piscinensis) - Wherever found      | Т      | \$64,440      | \$1,290,867,180    |
| 601  | Cactus, Bakersfield (Opuntia treleasei) - Wherever found                                 | E      | \$63,874      | \$1,290,931,054    |
| 602  | Darter, slackwater (Etheostoma boschungi) - Wherever found                               | Т      | \$63,819      | \$1,290,994,873    |
| 603  | Kamakahala (Labordia triflora) - Wherever found  | E      | \$63,529      | \$1,291,058,402    |
| 604  | No Common Name (Neraudia sericea) - Wherever found                                       | E      | \$63,478      | \$1,291,121,880    |
| 605  | Milkweed, Mead's (Asclepias meadii) - Wherever found                                     | Т      | \$63,310      | \$1,291,185,190    |
| 606  | Skullcap, large-flowered (Scutellaria montana) - Wherever found                          | Т      | \$63,062      | \$1,291,248,252    |
| 607  | Popolo (Cyanea solanacea) - Wherever found   | E      | \$62,464      | \$1,291,310,716    |
| 607  | Ha`iwale (Cyrtandra filipes) - Wherever found  | E      | \$62,464      | \$1,291,373,180    |
| 607  | No Common Name (Cyanea profuga) - Wherever found   | E      | \$62,464      | \$1,291,435,644    |
| 608  | No Common Name (Stenogyne bifida) - Wherever found                                       | E      | \$61,941      | \$1,291,497,585    |
| 609  | Nohoanu (Geranium hanaense) - Wherever found   | E      | \$61,802      | \$1,291,559,387    |
| 610  | Fairy shrimp, Conservancy (Branchinecta conservatio) - Wherever found                    | Е      | \$61,386      | \$1,291,620,773    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 611  | Honeycreeper, (Akohekohe) crested (Palmeria dolei) - Wherever found  | Е      | \$61,120      | \$1,291,681,893    |
| 612  | Hawk, Puerto Rican broad-winged (Buteo platypterus brunnescens) - Wherever found   | Е      | \$61,103      | \$1,291,742,996    |
| 613  | Lo`ulu (Pritchardia munroi) - Wherever found   | E      | \$61,013      | \$1,291,804,009    |
| 614  | Moccasinshell, Gulf (Medionidus penicillatus) - Wherever found   | E      | \$60,966      | \$1,291,864,975    |
| 615  | Mucket, Neosho (Lampsilis rafinesqueana) - Wherever found  | E      | \$60,613      | \$1,291,925,588    |
| 616  | Darter, snail (Percina tanasi) - Wherever found  | Т      | \$60,610      | \$1,291,986,198    |
| 617  | Snail, Guam tree (Partula radiolata) - Wherever found  | E      | \$60,483      | \$1,292,046,681    |
| 618  | Shiner, blue (Cyprinella caerulea) - Wherever found  | Т      | \$60,130      | \$1,292,106,811    |
| 619  | Mesa-mint, San Diego (Pogogyne abramsii) - Wherever found  | Е      | \$59,733      | \$1,292,166,544    |
| 620  | Pennyroyal, Todsen's (Hedeoma todsenii) - Wherever found   | Е      | \$59,000      | \$1,292,225,544    |
| 621  | Milk-vetch, Applegate's (Astragalus applegatei) - Wherever found   | Е      | \$58,900      | \$1,292,284,444    |
| 622  | Salamander, Santa Cruz long-toed (Ambystoma macrodactylum croceum) - Wherever found  | Е      | \$58,809      | \$1,292,343,253    |
| 623  | Mouse, St. Andrew beach (Peromyscus polionotus peninsularis) - Wherever found  | Е      | \$58,800      | \$1,292,402,053    |
| 624  | Snake, black pine (Pituophis melanoleucus lodingi) - Wherever found  | Т      | \$57,925      | \$1,292,459,978    |
| 625  | Brodiaea, Chinese Camp (Brodiaea pallida) - Wherever found   | Т      | \$57,810      | \$1,292,517,788    |
| 626  | Leopard, clouded (Neofelis nebulosa) - Wherever found  | E      | \$57,800      | \$1,292,575,588    |
| 627  | Alani (Melicope ovalis) - Wherever found   | E      | \$57,714      | \$1,292,633,302    |
| 628  | Cactus, Winkler (Pediocactus winkleri) - Wherever found  | Т      | \$57,600      | \$1,292,690,902    |
| 629  | Bean, Cumberland (pearlymussel) (Villosa trabalis) -<br>Wherever found; Except where listed as Experimental<br>Populations | Е      | \$57,560      | \$1,292,748,462    |
| 630  | Pearlymussel, slabside (Pleuronaia dolabelloides) -<br>Wherever found  | E      | \$57,410      | \$1,292,805,872    |
| 631  | Ko`oko`olau (Bidens wiebkei) - Wherever found  | E      | \$57,204      | \$1,292,863,076    |
| 632  | Akepa, Maui (Loxops ochraceus) - Wherever found  | E      | \$57,133      | \$1,292,920,209    |
| 633  | Lily, Western (Lilium occidentale) - Wherever found  | E      | \$57,000      | \$1,292,977,209    |
| 634  | Alani (Melicope reflexa) - Wherever found  | E      | \$56,950      | \$1,293,034,159    |
| 634  | Haha (Cyanea mannii) - Wherever found  | E      | \$56,950      | \$1,293,091,109    |
| 635  | Ladies'-tresses, Navasota (Spiranthes parksii) - Wherever found  | Е      | \$56,900      | \$1,293,148,009    |
| 636  | Wild-rice, Texas (Zizania texana) - Wherever found   | E      | \$56,600      | \$1,293,204,609    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 637  | Squirrel, Delmarva Peninsula fox (Sciurus niger cinereus) - Wherever found, except where listed as an experimental population | Е      | \$56,060      | \$1,293,260,669    |
| 638  | Turtle, Plymouth Redbelly (Pseudemys rubriventris bangsi) - Wherever found  | E      | \$55,920      | \$1,293,316,589    |
| 639  | Bladderpod, Missouri (Physaria filiformis) - Wherever found   | Т      | \$55,373      | \$1,293,371,962    |
| 640  | Haha (Cyanea maritae) - Wherever found  | E      | \$55,037      | \$1,293,426,999    |
| 641  | Kula wahine noho (Isodendrion pyrifolium) - Wherever found  | Е      | \$54,969      | \$1,293,481,968    |
| 642  | Riffleshell, tan (Epioblasma florentina walkeri (=E. walkeri)) - Wherever found   | E      | \$54,960      | \$1,293,536,928    |
| 643  | No Common Name (Wikstroemia villosa) - Wherever found   | E      | \$54,872      | \$1,293,591,800    |
| 644  | Thelypody, Howell's spectacular (Thelypodium howellii spectabilis) - Wherever found   | Т      | \$54,615      | \$1,293,646,415    |
| 645  | Butterweed, Layne's (Senecio layneae) - Wherever found  | Т      | \$54,020      | \$1,293,700,435    |
| 646  | Nohoanu (Geranium multiflorum) - Wherever found   | E      | \$53,808      | \$1,293,754,243    |
| 647  | `Akoko, Ewa Plains (Euphorbia skottsbergii var. skottsbergii) - Wherever found  | Е      | \$53,793      | \$1,293,808,036    |
| 648  | Pearlymussel, littlewing (Pegias fabula) - Wherever found   | E      | \$53,660      | \$1,293,861,696    |
| 649  | Nui, haha (Cyanea horrida) - Wherever found   | E      | \$52,885      | \$1,293,914,581    |
| 650  | Arrowhead, bunched (Sagittaria fasciculata) - Wherever found  | E      | \$52,359      | \$1,293,966,940    |
| 651  | Horse, Przewalski's (Equus przewalskii) - Wherever found  | E      | \$52,000      | \$1,294,018,940    |
| 652  | Wawae`iole (Huperzia mannii) - Wherever found   | E      | \$51,857      | \$1,294,070,797    |
| 653  | Bulrush, Northeastern (Scirpus ancistrochaetus) - Wherever found  | E      | \$51,825      | \$1,294,122,622    |
| 654  | Boa, Jamaican (Epicrates subflavus) - Wherever found  | E      | \$51,667      | \$1,294,174,289    |
| 655  | Haha (Cyanea kunthiana) - Wherever found  | E      | \$51,290      | \$1,294,225,579    |
| 656  | Sheep, Peninsular bighorn (Ovis canadensis nelsoni) - Peninsular CA pop.  | E      | \$51,257      | \$1,294,276,836    |
| 657  | Ko`oko`olau (Bidens campylotheca pentamera) - Wherever found  | E      | \$51,246      | \$1,294,328,082    |
| 658  | Woodrat, Key Largo (Neotoma floridana smalli) - Wherever found  | E      | \$51,210      | \$1,294,379,292    |
| 659  | Haha (Cyanea copelandii ssp. haleakalaensis) - Wherever found   | Е      | \$51,167      | \$1,294,430,459    |
| 660  | Haha (Cyanea hamatiflora ssp. hamatiflora) - Wherever found   | Е      | \$50,611      | \$1,294,481,070    |
| 661  | Grass, Eureka Dune (Swallenia alexandrae) -   | E      | \$50,564      | \$1,294,531,634    |
| 662  | Evening-primrose, Eureka Valley (Oenothera avita ssp. eurekensis) -   | Е      | \$50,464      | \$1,294,582,098    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 663  | Yellowhead, desert (Yermo xanthocephalus) - Wherever found  | Т      | \$50,279      | \$1,294,632,377    |
| 664  | Trillium, relict (Trillium reliquum) - Wherever found   | E      | \$50,169      | \$1,294,682,546    |
| 665  | Ko`oko`olau (Bidens campylotheca waihoiensis) - Wherever found  | Е      | \$50,152      | \$1,294,732,698    |
| 666  | Deer, Eld's brow-antlered (Cervus eldi) - Wherever found  | E      | \$50,100      | \$1,294,782,798    |
| 666  | Cactus, Acuna (Echinomastus erectocentrus var. acunensis) - Wherever found                                  | E      | \$50,100      | \$1,294,832,898    |
| 667  | Crocodile, Nile (Crocodylus niloticus) - Wherever found   | Т      | \$50,000      | \$1,294,882,898    |
| 667  | Crocodile, saltwater (Crocodylus porosus) - Australia   | Т      | \$50,000      | \$1,294,932,898    |
| 667  | Sturgeon, Beluga (Huso huso) - Wherever found   | Т      | \$50,000      | \$1,294,982,898    |
| 668  | Madtom, yellowfin (Noturus flavipinnis) - Wherever found, except where listed as an experimental population | Т      | \$49,980      | \$1,295,032,878    |
| 669  | Moccasinshell, Ochlockonee (Medionidus simpsonianus) - Wherever found                                       | E      | \$49,776      | \$1,295,082,654    |
| 670  | No Common Name (Dendrobium guamense) - Wherever found   | Т      | \$49,639      | \$1,295,132,293    |
| 671  | Damselfly, Pacific Hawaiian (Megalagrion pacificum) - Wherever found  | Е      | \$49,123      | \$1,295,181,416    |
| 672  | Milkvetch, Deseret (Astragalus desereticus) - Wherever found  | Т      | \$48,514      | \$1,295,229,930    |
| 673  | Snail, Pecos assiminea (Assiminea pecos) - Wherever found   | E      | \$48,275      | \$1,295,278,205    |
| 674  | Kidneyshell, fluted (Ptychobranchus subtentum) - Wherever found   | E      | \$48,260      | \$1,295,326,465    |
| 675  | Pigtoe, fuzzy (Pleurobema strodeanum) - Wherever found  | Т      | \$48,203      | \$1,295,374,668    |
| 676  | Thistle, Chorro Creek bog (Cirsium fontinale var. obispoense) - Wherever found                              | Е      | \$48,088      | \$1,295,422,756    |
| 677  | Rock cress, Shale barren (Arabis serotina) - Wherever found   | Е      | \$48,047      | \$1,295,470,803    |
| 678  | Butterwort, Godfrey's (Pinguicula ionantha) - Wherever found  | Т      | \$47,569      | \$1,295,518,372    |
| 679  | Fern, American hart's-tongue (Asplenium scolopendrium var. americanum) - Wherever found                     | Т      | \$47,079      | \$1,295,565,451    |
| 680  | Cactus, Pima pineapple (Coryphantha scheeri var. robustispina) - Wherever found                             | Е      | \$47,014      | \$1,295,612,465    |
| 681  | Wallflower, Contra Costa (Erysimum capitatum var. angustatum) - Wherever found                              | Е      | \$46,627      | \$1,295,659,092    |
| 682  | Spinedace, Big Spring (Lepidomeda mollispinis pratensis) - Wherever found                                   | Т      | \$46,317      | \$1,295,705,409    |
| 683  | Bean, Choctaw (Villosa choctawensis) - Wherever found   | E      | \$46,045      | \$1,295,751,454    |
| 684  | Amphipod, Noel's (Gammarus desperatus) - Wherever found   | Е      | \$45,915      | \$1,295,797,369    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 685  | Cactus, Knowlton's (Pediocactus knowltonii) - Wherever found   | E      | \$45,855      | \$1,295,843,224    |
| 686  | Butterfly, mission blue (Icaricia icarioides missionensis) - Wherever found  | E      | \$45,127      | \$1,295,888,351    |
| 687  | Wolf, Mexican (Canis lupus baileyi) - Wherever found, except where listed as an experimental population  | Е      | \$45,000      | \$1,295,933,351    |
| 687  | Cat, Andean (Felis jacobita) - Wherever found  | E      | \$45,000      | \$1,295,978,351    |
| 688  | Wartyback, white (pearlymussel) (Plethobasus cicatricosus) - Wherever found  | E      | \$44,860      | \$1,296,023,211    |
| 689  | Avens, spreading (Geum radiatum) - Wherever found  | E      | \$44,315      | \$1,296,067,526    |
| 690  | Aplokating-palaoan (Psychotria malaspinae) - Wherever found  | E      | \$44,100      | \$1,296,111,626    |
| 691  | Sturgeon, Alabama (Scaphirhynchus suttkusi) - Wherever found   | E      | \$43,957      | \$1,296,155,583    |
| 692  | Spineflower, Monterey (Chorizanthe pungens var. pungens) - Wherever found  | Т      | \$43,931      | \$1,296,199,514    |
| 693  | Cactus, black lace (Echinocereus reichenbachii var. albertii) - Wherever found   | E      | \$43,917      | \$1,296,243,431    |
| 694  | Rail, Guam (Rallus owstoni) - Rota   | EXPN   | \$43,913      | \$1,296,287,344    |
| 695  | Fern, Florida bristle (Trichomanes punctatum ssp. floridanum) - Wherever found   | Е      | \$43,600      | \$1,296,330,944    |
| 696  | Halumtano, Cebello (Bulbophyllum guamense) - Wherever found  | Т      | \$43,378      | \$1,296,374,322    |
| 697  | Snail, Morro shoulderband (=Banded dune)<br>(Helminthoglypta walkeriana) - Wherever found  | E      | \$43,347      | \$1,296,417,669    |
| 697  | Gerardia, sandplain (Agalinis acuta) - Wherever found  | E      | \$43,347      | \$1,296,461,016    |
| 698  | Pigtoe, narrow (Fusconaia escambia) - Wherever found   | Т      | \$43,259      | \$1,296,504,275    |
| 699  | Jaguarundi, Gulf Coast (Herpailurus (=Felis) yagouaroundi cacomitli) - Wherever found  | Е      | \$43,122      | \$1,296,547,397    |
| 700  | Blazingstar, Heller's (Liatris helleri) - Wherever found   | Т      | \$43,095      | \$1,296,590,492    |
| 701  | Gumplant, Ash Meadows (Grindelia fraxinipratensis) - Wherever found  | Т      | \$43,032      | \$1,296,633,524    |
| 702  | Sandshell, Southern (Hamiota australis) - Wherever found   | Т      | \$42,939      | \$1,296,676,463    |
| 703  | Purple Cat's paw (=Purple Cat's paw pearlymussel) (Epioblasma obliquata obliquata) - Wherever found; Except where listed as Experimental Populations | Е      | \$42,833      | \$1,296,719,296    |
| 704  | Snail, fragile tree (Samoana fragilis) - Wherever found  | E      | \$42,467      | \$1,296,761,763    |
| 705  | Millerbird, Nihoa (old world warbler) (Acrocephalus familiaris kingi) - Wherever found   | E      | \$42,271      | \$1,296,804,034    |
| 706  | Snail, Humped tree (Partula gibba) - Wherever found  | E      | \$42,142      | \$1,296,846,176    |
| 707  | Heather, mountain golden (Hudsonia montana) - Wherever found   | Т      | \$41,993      | \$1,296,888,169    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 708  | Cactus, Tobusch fishhook (Sclerocactus brevihamatus ssp. tobuschii) - Wherever found                                      | Е      | \$41,600      | \$1,296,929,769    |
| 709  | Irisette, white (Sisyrinchium dichotomum) - Wherever found  | E      | \$41,540      | \$1,296,971,309    |
| 710  | Salamander, Cheat Mountain (Plethodon nettingi) - Wherever found  | Т      | \$41,366      | \$1,297,012,675    |
| 711  | Spurge, telephus (Euphorbia telephioides) - Wherever found  | Т      | \$41,310      | \$1,297,053,985    |
| 712  | Milk-vetch, Holmgren (Astragalus holmgreniorum) - Wherever found  | Е      | \$41,300      | \$1,297,095,285    |
| 713  | Meadowfoam, Butte County (Limnanthes floccosa ssp. californica) - Wherever found  | Е      | \$40,920      | \$1,297,136,205    |
| 714  | Fanshell (Cyprogenia stegaria) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1)) | EXPN   | \$40,416      | \$1,297,176,621    |
| 715  | Prairie-clover, leafy (Dalea foliosa) - Wherever found  | E      | \$40,292      | \$1,297,216,913    |
| 716  | Kidneyshell, southern (Ptychobranchus jonesi) - Wherever found  | Е      | \$40,105      | \$1,297,257,018    |
| 717  | Shrimp, Alabama cave (Palaemonias alabamae) - Wherever found  | Е      | \$40,100      | \$1,297,297,118    |
| 718  | Bonytongue, Asian (Scleropages formosus) - Wherever found   | Е      | \$40,000      | \$1,297,337,118    |
| 719  | Cycladenia, Jones (Cycladenia humilis var. jonesii) - Wherever found  | Т      | \$39,640      | \$1,297,376,758    |
| 720  | Bluegrass, San Bernardino (Poa atropurpurea) - Wherever found   | Е      | \$39,483      | \$1,297,416,241    |
| 721  | Reed-mustard, shrubby (Schoenocrambe suffrutescens) - Wherever found  | Е      | \$39,424      | \$1,297,455,665    |
| 722  | Moccasinshell, Alabama (Medionidus acutissimus) - Wherever found  | Т      | \$39,398      | \$1,297,495,063    |
| 723  | Isopod, Madison Cave (Antrolana lira) - Wherever found  | Т      | \$38,903      | \$1,297,533,966    |
| 724  | Milk-vetch, Mancos (Astragalus humillimus) - Wherever found   | E      | \$38,900      | \$1,297,572,866    |
| 725  | Pitcher-plant, green (Sarracenia oreophila) - Wherever found  | Е      | \$38,603      | \$1,297,611,469    |
| 726  | Orcutt grass, San Joaquin (Orcuttia inaequalis) - Wherever found  | Т      | \$38,591      | \$1,297,650,060    |
| 727  | Lichen, rock gnome (Gymnoderma lineare) - Wherever found  | Е      | \$38,421      | \$1,297,688,481    |
| 728  | Bittercress, small-anthered (Cardamine micranthera) - Wherever found  | Е      | \$38,395      | \$1,297,726,876    |
| 729  | Salamander, California tiger (Ambystoma californiense) - U.S.A. (CA - Sonoma County)                                      | Е      | \$38,150      | \$1,297,765,026    |
| 730  | Cactus, Fickeisen plains (Pediocactus peeblesianus fickeiseniae) - Wherever found   | Е      | \$38,022      | \$1,297,803,048    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 731  | Ipomopsis, Holy Ghost (Ipomopsis sancti-spiritus) - Wherever found                   | Е      | \$37,800      | \$1,297,840,848    |
| 732  | Pupfish, Comanche Springs (Cyprinodon elegans) - Wherever found                      | Е      | \$37,709      | \$1,297,878,557    |
| 733  | No Common Name (Tabernaemontana rotensis) - Wherever found                           | Т      | \$37,667      | \$1,297,916,224    |
| 734  | Blazingstar, Ash Meadows (Mentzelia leucophylla) - Wherever found                    | Т      | \$37,663      | \$1,297,953,887    |
| 735  | Ground-plum, Guthrie's (=Pyne's) (Astragalus bibullatus) - Wherever found            | E      | \$37,600      | \$1,297,991,487    |
| 736  | Deer, key (Odocoileus virginianus clavium) - Wherever found                          | E      | \$37,400      | \$1,298,028,887    |
| 737  | Finch, Nihoa (honeycreeper) (Telespyza ultima) - Wherever found                      | E      | \$37,160      | \$1,298,066,047    |
| 738  | Oryx, scimitar-horned (Oryx dammah) - Wherever found                                 | E      | \$37,000      | \$1,298,103,047    |
| 738  | No Common Name (Lepanthes eltoroensis) - Wherever found                              | Е      | \$37,000      | \$1,298,140,047    |
| 739  | Shiner, smalleye (Notropis buccula) - Wherever found                                 | E      | \$36,900      | \$1,298,176,947    |
| 739  | Gooseberry, Miccosukee (Ribes echinellum) - Wherever found                           | Т      | \$36,900      | \$1,298,213,847    |
| 740  | Stickseed, showy (Hackelia venusta) - Wherever found                                 | E      | \$36,715      | \$1,298,250,562    |
| 741  | Evening-primrose, Antioch Dunes (Oenothera deltoides ssp. howellii) - Wherever found | Е      | \$36,686      | \$1,298,287,248    |
| 742  | Elktoe, Cumberland (Alasmidonta atropurpurea) - Wherever found                       | Е      | \$36,660      | \$1,298,323,908    |
| 743  | Cat's-eye, Terlingua Creek (Cryptantha crassipes) - Wherever found                   | Е      | \$36,648      | \$1,298,360,556    |
| 744  | Minnow, Devils River (Dionda diaboli) - Wherever found                               | Т      | \$36,530      | \$1,298,397,086    |
| 745  | Dace, Laurel (Chrosomus saylori) - Wherever found                                    | E      | \$36,500      | \$1,298,433,586    |
| 746  | Bluet, Roan Mountain (Hedyotis purpurea var. montana) - Wherever found               | Е      | \$36,287      | \$1,298,469,873    |
| 747  | Chaff-flower, round-leaved (Achyranthes splendens var. rotundata) - Wherever found   | Е      | \$36,228      | \$1,298,506,101    |
| 748  | Rosemary, Apalachicola (Conradina glabra) - Wherever found                           | E      | \$36,000      | \$1,298,542,101    |
| 749  | Ufa-halomtano (Heritiera longipetiolata) - Wherever found                            | E      | \$35,987      | \$1,298,578,088    |
| 750  | Uhi uhi (Mezoneuron kavaiense) - Wherever found                                      | E      | \$35,966      | \$1,298,614,054    |
| 751  | Niterwort, Amargosa (Nitrophila mohavensis) - Wherever found                         | Е      | \$35,935      | \$1,298,649,989    |
| 752  | Palo de rosa (Ottoschulzia rhodoxylon) - Wherever found                              | E      | \$35,700      | \$1,298,685,689    |
| 753  | Sunshine, Sonoma (Blennosperma bakeri) - Wherever found                              | Е      | \$35,639      | \$1,298,721,328    |
| 754  | No Common Name (Eugenia bryanii) - Wherever found                                    | E      | \$35,348      | \$1,298,756,676    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 755  | Shiner, sharpnose (Notropis oxyrhynchus) - Wherever found   | Е      | \$35,300      | \$1,298,791,976    |
| 755  | Iris, dwarf lake (Iris lacustris) - Wherever found  | Т      | \$35,300      | \$1,298,827,276    |
| 756  | Bear, American black (Ursus americanus) - County range of LA b.bear                                       | SAT    | \$35,171      | \$1,298,862,447    |
| 757  | Mountainbalm, Indian Knob (Eriodictyon altissimum) -<br>Wherever found                                    | Е      | \$35,102      | \$1,298,897,549    |
| 758  | Crayfish, Big Sandy (Cambarus callainus) - Wherever found   | Т      | \$34,253      | \$1,298,931,802    |
| 759  | Butterfly, Miami Blue (Cyclargus (=Hemiargus) thomasi bethunebakeri) - Wherever found                     | Е      | \$34,200      | \$1,298,966,002    |
| 760  | Ambrosia, south Texas (Ambrosia cheiranthifolia) - Wherever found   | Е      | \$34,077      | \$1,299,000,079    |
| 761  | Sedge, golden (Carex lutea) - Wherever found  | E      | \$33,849      | \$1,299,033,928    |
| 762  | Phacelia, DeBeque (Phacelia submutica) - Wherever found   | Т      | \$33,613      | \$1,299,067,541    |
| 763  | `aku (Cyanea tritomantha) - Wherever found  | E      | \$33,606      | \$1,299,101,147    |
| 764  | Manzanita, Ione (Arctostaphylos myrtifolia) - Wherever found  | Т      | \$33,521      | \$1,299,134,668    |
| 765  | Butterfly, Bartram's hairstreak (Strymon acis bartrami) - Wherever found                                  | Е      | \$33,500      | \$1,299,168,168    |
| 766  | Ebonyshell, round (Fusconaia rotulata) - Wherever found   | E      | \$33,019      | \$1,299,201,187    |
| 767  | Condor, Andean (Vultur gryphus) - Wherever found  | E      | \$33,000      | \$1,299,234,187    |
| 768  | Darter, Cumberland (Etheostoma susanae) - Wherever found  | E      | \$32,720      | \$1,299,266,907    |
| 769  | Sunflower, Pecos (=puzzle, =paradox) (Helianthus paradoxus) - Wherever found                              | Т      | \$32,700      | \$1,299,299,607    |
| 770  | Meadowrue, Cooley's (Thalictrum cooleyi) - Wherever found   | E      | \$32,687      | \$1,299,332,294    |
| 771  | Sedge, Navajo (Carex specuicola) - Wherever found   | Т      | \$32,460      | \$1,299,364,754    |
| 772  | Ko`oko`olau (Bidens micrantha ctenophylla) - Wherever found   | Е      | \$32,359      | \$1,299,397,113    |
| 773  | Goldenrod, white-haired (Solidago albopilosa) - Wherever found  | Т      | \$32,200      | \$1,299,429,313    |
| 773  | Cobana negra (Stahlia monosperma) - Wherever found  | Т      | \$32,200      | \$1,299,461,513    |
| 774  | Darter, amber (Percina antesella) - Wherever found  | E      | \$32,110      | \$1,299,493,623    |
| 775  | Pigtoe, finerayed (Fusconaia cuneolus) - Wherever found; Except where listed as Experimental Populations  | E      | \$32,000      | \$1,299,525,623    |
| 776  | Poolfish, Pahrump (Empetrichthys latos) - Wherever found  | Е      | \$31,947      | \$1,299,557,570    |
| 777  | Haha (Cyanea shipmanii) - Wherever found  | E      | \$31,751      | \$1,299,589,321    |
| 778  | Pearlymussel, cracking (Hemistena lata) - Wherever found; Except where listed as Experimental Populations | Е      | \$31,560      | \$1,299,620,881    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 779  | Frog, Panamanian golden (Atelopus varius zeteki) - Wherever found  | E      | \$31,500      | \$1,299,652,381    |
| 780  | Goldenrod, Blue Ridge (Solidago spithamaea) - Wherever found   | Т      | \$31,499      | \$1,299,683,880    |
| 781  | Skullcap, Florida (Scutellaria floridana) - Wherever found   | Т      | \$31,355      | \$1,299,715,235    |
| 782  | Owl's-clover, fleshy (Castilleja campestris ssp. succulenta) - Wherever found  | Т      | \$31,334      | \$1,299,746,569    |
| 783  | Gambusia, Pecos (Gambusia nobilis) - Wherever found  | E      | \$31,010      | \$1,299,777,579    |
| 784  | Milk-vetch, Jesup's (Astragalus robbinsii var. jesupi) - Wherever found  | Е      | \$30,923      | \$1,299,808,502    |
| 785  | Salamander, Shenandoah (Plethodon shenandoah) - Wherever found   | E      | \$30,800      | \$1,299,839,302    |
| 786  | Lo`ulu (Pritchardia lanigera) - Wherever found   | E      | \$30,786      | \$1,299,870,088    |
| 787  | Spinedace, Little Colorado (Lepidomeda vittata) - Wherever found   | Т      | \$30,602      | \$1,299,900,690    |
| 788  | Meadowfoam, Sebastopol (Limnanthes vinculans) - Wherever found   | E      | \$30,039      | \$1,299,930,729    |
| 789  | Darter, fountain (Etheostoma fonticola) - Wherever found   | E      | \$30,000      | \$1,299,960,729    |
| 790  | Lupine, scrub (Lupinus aridorum) - Wherever found  | E      | \$29,700      | \$1,299,990,429    |
| 791  | Plum, scrub (Prunus geniculata) - Wherever found   | E      | \$29,600      | \$1,300,020,029    |
| 792  | Rice rat (Oryzomys palustris natator) - lower FL Keys  | E      | \$29,510      | \$1,300,049,539    |
| 793  | Clubshell (Pleurobema clava) - AL: Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL | EXPN   | \$29,480      | \$1,300,079,019    |
| 794  | Clubshell, southern (Pleurobema decisum) - Wherever found  | E      | \$29,374      | \$1,300,108,393    |
| 795  | Birds-in-a-nest, white (Macbridea alba) - Wherever found   | Т      | \$29,334      | \$1,300,137,727    |
| 796  | Gilia, Monterey (Gilia tenuiflora ssp. arenaria) - Wherever found  | E      | \$29,162      | \$1,300,166,889    |
| 797  | Iguana, Cuban ground (Cyclura nubila nubila) - (Cyclura nubila nubila)   | Т      | \$29,148      | \$1,300,196,037    |
| 798  | Torreya, Florida (Torreya taxifolia) - Wherever found  | E      | \$28,411      | \$1,300,224,448    |
| 799  | Navarretia, spreading (Navarretia fossalis) - Wherever found   | Т      | \$28,382      | \$1,300,252,830    |
| 800  | Bird's beak, palmate-bracted (Cordylanthus palmatus) - Wherever found  | E      | \$28,050      | \$1,300,280,880    |
| 801  | Kopa (Kadua cordata remyi) - Wherever found  | E      | \$28,018      | \$1,300,308,898    |
| 802  | Campion, fringed (Silene polypetala) - Wherever found  | E      | \$27,746      | \$1,300,336,644    |
| 803  | Cactus, Uinta Basin hookless (Sclerocactus wetlandicus) - Wherever found   | Т      | \$27,718      | \$1,300,364,362    |
| 804  | Hala pepe (Pleomele hawaiiensis) - Wherever found  | E      | \$27,704      | \$1,300,392,066    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 805  | Milk-vetch, Coachella Valley (Astragalus lentiginosus var. coachellae) - Wherever found         | Е      | \$27,584      | \$1,300,419,650    |
| 806  | Snail, Iowa Pleistocene (Discus macclintocki) - Wherever found                                  | E      | \$27,500      | \$1,300,447,150    |
| 806  | Cactus, Kuenzler hedgehog (Echinocereus fendleri var. kuenzleri) - Wherever found               | Е      | \$27,500      | \$1,300,474,650    |
| 807  | Boa, Virgin Islands tree (Epicrates monensis granti) - Wherever found                           | Е      | \$26,800      | \$1,300,501,450    |
| 808  | Nehe (Melanthera tenuifolia) - Wherever found   | E      | \$26,740      | \$1,300,528,190    |
| 809  | Spinymussel, Altamaha (Elliptio spinosa) - Wherever found                                       | E      | \$26,606      | \$1,300,554,796    |
| 810  | Pigtoe, shiny (Fusconaia cor) - Wherever found; Except where listed as Experimental Populations | Е      | \$26,600      | \$1,300,581,396    |
| 811  | Milkweed, Welsh's (Asclepias welshii) - Wherever found  | Т      | \$26,500      | \$1,300,607,896    |
| 812  | Orcutt grass, California (Orcuttia californica) - Wherever found                                | Е      | \$26,452      | \$1,300,634,348    |
| 813  | Dudleya, Santa Clara Valley (Dudleya setchellii) -<br>Wherever found                            | Е      | \$26,370      | \$1,300,660,718    |
| 814  | Mallow, Gierisch (Sphaeralcea gierischii) - Wherever found                                      | E      | \$26,300      | \$1,300,687,018    |
| 815  | Skyrocket, Pagosa (Ipomopsis polyantha) - Wherever found  | Е      | \$26,292      | \$1,300,713,310    |
| 816  | Butterfly, Florida leafwing (Anaea troglodyta floridalis) - Wherever found                      | Е      | \$26,000      | \$1,300,739,310    |
| 817  | Kiponapona (Phyllostegia racemosa) - Wherever found   | E      | \$25,986      | \$1,300,765,296    |
| 818  | Orcutt grass, hairy (Orcuttia pilosa) - Wherever found  | E      | \$25,760      | \$1,300,791,056    |
| 819  | Dace, Independence Valley speckled (Rhinichthys osculus lethoporus) - Wherever found            | E      | \$25,625      | \$1,300,816,681    |
| 820  | Coqui, golden (Eleutherodactylus jasperi) - Wherever found                                      | Т      | \$25,500      | \$1,300,842,181    |
| 821  | Pigtoe, tapered (Fusconaia burkei) - Wherever found   | Т      | \$25,425      | \$1,300,867,606    |
| 822  | Mustard, Penland alpine fen (Eutrema penlandii) -<br>Wherever found                             | Т      | \$25,355      | \$1,300,892,961    |
| 823  | Vetch, Hawaiian (Vicia menziesii) - Wherever found  | E      | \$25,210      | \$1,300,918,171    |
| 824  | Thistle, La Graciosa (Cirsium Ioncholepis) - Wherever found                                     | Е      | \$25,010      | \$1,300,943,181    |
| 825  | Rocksnail, round (Leptoxis ampla) - Wherever found  | Т      | \$25,000      | \$1,300,968,181    |
| 826  | Springfish, Railroad Valley (Crenichthys nevadae) - Wherever found                              | Т      | \$24,774      | \$1,300,992,955    |
| 827  | Butterfly, Mariana wandering (Vagrans egistina) - Wherever found                                | Е      | \$24,732      | \$1,301,017,687    |
| 828  | No Common Name (Tetramolopium rockii) - Wherever found  | Т      | \$24,655      | \$1,301,042,342    |
| 829  | Clubshell, black (Pleurobema curtum) - Wherever found   | E      | \$24,500      | \$1,301,066,842    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 830  | `Aiakeakua, popolo (Solanum sandwicense) - Wherever found   | Е      | \$24,333      | \$1,301,091,175    |
| 831  | Aster, Ruth's golden (Pityopsis ruthii) - Wherever found  | E      | \$24,320      | \$1,301,115,495    |
| 832  | Crayfish, Guyandotte River (Cambarus veteranus) - Wherever found  | Е      | \$24,253      | \$1,301,139,748    |
| 833  | Warbler, nightingale reed (old world warbler)<br>(Acrocephalus luscinia) - Wherever found                 | Е      | \$24,165      | \$1,301,163,913    |
| 834  | Fern, Aleutian shield (Polystichum aleuticum) - Wherever found  | Е      | \$24,000      | \$1,301,187,913    |
| 835  | Mesa-mint, Otay (Pogogyne nudiuscula) - Wherever found  | E      | \$23,847      | \$1,301,211,760    |
| 836  | Alopecurus, Sonoma (Alopecurus aequalis var. sonomensis) - Wherever found                                 | E      | \$23,839      | \$1,301,235,599    |
| 837  | Naucorid, Ash Meadows (Ambrysus amargosus) - Wherever found   | Т      | \$23,717      | \$1,301,259,316    |
| 838  | Dace, Clover Valley speckled (Rhinichthys osculus oligoporus) - Wherever found                            | Е      | \$23,712      | \$1,301,283,028    |
| 839  | Sea turtle, olive ridley (Lepidochelys olivacea) - Breeding colony populations on Pacific coast of Mexico | Е      | \$23,555      | \$1,301,306,583    |
| 840  | Sunray, Ash Meadows (Enceliopsis nudicaulis var. corrugata) - Wherever found                              | Т      | \$23,539      | \$1,301,330,122    |
| 841  | Flannelbush, Mexican (Fremontodendron mexicanum) - Wherever found   | Е      | \$23,410      | \$1,301,353,532    |
| 842  | Kangaroo rat, Morro Bay (Dipodomys heermanni morroensis) - Wherever found                                 | Е      | \$23,256      | \$1,301,376,788    |
| 843  | A`e (Zanthoxylum dipetalum var. tomentosum) - Wherever found  | Е      | \$23,217      | \$1,301,400,005    |
| 844  | Sea turtle, green (Chelonia mydas) - North Indian DPS   | Т      | \$23,000      | \$1,301,423,005    |
| 844  | Gourd, Okeechobee (Cucurbita okeechobeensis ssp. okeechobeensis) - Wherever found                         | Е      | \$23,000      | \$1,301,446,005    |
| 844  | Beardtongue, Penland (Penstemon penlandii) - Wherever found   | Е      | \$23,000      | \$1,301,469,005    |
| 845  | Pearlymussel, dromedary (Dromus dromas) - Wherever found; Except where listed as Experimental Populations | Е      | \$22,960      | \$1,301,491,965    |
| 846  | Snail, Lanai tree (Partulina semicarinata) - Wherever found   | E      | \$22,937      | \$1,301,514,902    |
| 846  | Snail, Lanai tree (Partulina variabilis) - Wherever found   | Е      | \$22,937      | \$1,301,537,839    |
| 847  | Butterfly, Schaus swallowtail (Heraclides aristodemus ponceanus) - Wherever found                         | Е      | \$22,830      | \$1,301,560,669    |
| 848  | Haha (Cyanea crispa) - Wherever found   | Е      | \$22,817      | \$1,301,583,486    |
| 849  | Vole, Hualapai Mexican (Microtus mexicanus hualpaiensis) - Wherever found                                 | E      | \$22,700      | \$1,301,606,186    |
| 850  | Moth, Kern primrose sphinx (Euproserpinus euterpe) - Wherever found                                       | Т      | \$22,690      | \$1,301,628,876    |
| 851  | Haha (Cyanea stictophylla) - Wherever found   | E      | \$22,532      | \$1,301,651,408    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 852  | Springsnail, Three Forks (Pyrgulopsis trivialis) - Wherever found                | Е      | \$22,400      | \$1,301,673,808    |
| 853  | Checker-mallow, Keck's (Sidalcea keckii) - Wherever found                        | E      | \$22,322      | \$1,301,696,130    |
| 854  | Crayfish, Shasta (Pacifastacus fortis) - Wherever found                          | E      | \$22,200      | \$1,301,718,330    |
| 855  | Sandlace (Polygonella myriophylla) - Wherever found                              | E      | \$22,000      | \$1,301,740,330    |
| 856  | Wawae`iole (Huperzia nutans) - Wherever found                                    | E      | \$21,880      | \$1,301,762,210    |
| 857  | Alani (Melicope zahlbruckneri) - Wherever found                                  | E      | \$21,866      | \$1,301,784,076    |
| 858  | Salamander, Jemez Mountains (Plethodon neomexicanus) - Wherever found            | Е      | \$21,800      | \$1,301,805,876    |
| 858  | Sucker, Modoc (Catostomus microps) - Wherever found                              | E      | \$21,800      | \$1,301,827,676    |
| 859  | Sandwort, Marsh (Arenaria paludicola) - Wherever found                           | E      | \$21,725      | \$1,301,849,401    |
| 860  | Mint, longspurred (Dicerandra cornutissima) - Wherever found                     | Е      | \$21,700      | \$1,301,871,101    |
| 861  | Lousewort, Furbish (Pedicularis furbishiae) - Wherever found                     | E      | \$21,574      | \$1,301,892,675    |
| 862  | Barberry, island (Berberis pinnata ssp. insularis) - Wherever found              | Е      | \$21,561      | \$1,301,914,236    |
| 863  | Checkermallow, Wenatchee Mountains (Sidalcea oregana var. calva) - Local Endemic | Е      | \$21,520      | \$1,301,935,756    |
| 864  | Pigtoe, southern (Pleurobema georgianum) - Wherever found                        | Е      | \$21,440      | \$1,301,957,196    |
| 865  | Butterfly, San Bruno elfin (Callophrys mossii bayensis) - Wherever found         | Е      | \$21,177      | \$1,301,978,373    |
| 866  | Silverside, Waccamaw (Menidia extensa) - Wherever found                          | Т      | \$21,112      | \$1,301,999,485    |
| 867  | Mouse, Anastasia Island beach (Peromyscus polionotus phasma) - Wherever found    | Е      | \$21,100      | \$1,302,020,585    |
| 868  | Springfish, White River (Crenichthys baileyi baileyi) - Wherever found           | Е      | \$21,068      | \$1,302,041,653    |
| 869  | Trillium, persistent (Trillium persistens) - Wherever found                      | E      | \$21,000      | \$1,302,062,653    |
| 870  | Milk-vetch, Braunton's (Astragalus brauntonii) - Wherever found                  | Е      | \$20,960      | \$1,302,083,613    |
| 871  | Manioc, Walker's (Manihot walkerae) - Wherever found                             | Е      | \$20,800      | \$1,302,104,413    |
| 871  | Lead-plant, Crenulate (Amorpha crenulata) - Wherever found                       | Е      | \$20,800      | \$1,302,125,213    |
| 872  | Logperch, Conasauga (Percina jenkinsi) - Wherever found                          | E      | \$20,740      | \$1,302,145,953    |
| 873  | Mouse, Key Largo cotton (Peromyscus gossypinus allapaticola) - Wherever found    | Е      | \$20,710      | \$1,302,166,663    |
| 874  | Bladderpod, White Bluffs (Physaria douglasii ssp. tuplashensis) - Wherever found | Т      | \$20,675      | \$1,302,187,338    |
| 875  | `Oha wai (Clermontia peleana) - Wherever found                                   | E      | \$20,673      | \$1,302,208,011    |
| 876  | Pocketbook, finelined (Lampsilis altilis) - Wherever found                       | Т      | \$20,656      | \$1,302,228,667    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 877  | Pa`iniu (Astelia waialealae) - Wherever found   | E      | \$20,598      | \$1,302,249,265    |
| 878  | Salamander, Barton Springs (Eurycea sosorum) - Wherever found   | Е      | \$20,500      | \$1,302,269,765    |
| 878  | Orcutt grass, slender (Orcuttia tenuis) - Wherever found  | Т      | \$20,500      | \$1,302,290,265    |
| 879  | Koki`o (Kokia drynarioides) - Wherever found  | Е      | \$20,404      | \$1,302,310,669    |
| 880  | Aster, Florida golden (Chrysopsis floridana) - Wherever found   | E      | \$20,400      | \$1,302,331,069    |
| 881  | Naupaka, dwarf (Scaevola coriacea) - Wherever found   | E      | \$20,398      | \$1,302,351,467    |
| 882  | Pigeon, Puerto Rican plain (Columba inornata wetmorei) - Wherever found   | E      | \$20,340      | \$1,302,371,807    |
| 883  | Iguana, Fiji banded (Brachylophus fasciatus) - Wherever found   | E      | \$20,000      | \$1,302,391,807    |
| 884  | `Oha wai (Clermontia lindseyana) - Wherever found   | E      | \$19,779      | \$1,302,411,586    |
| 885  | No Common Name (Phyllostegia parviflora) - Wherever found   | Е      | \$19,755      | \$1,302,431,341    |
| 886  | No Common Name (Phyllostegia floribunda) - Wherever found   | E      | \$19,717      | \$1,302,451,058    |
| 887  | Petrel, Bermuda (Pterodroma cahow) - Wherever found   | Е      | \$19,575      | \$1,302,470,633    |
| 888  | Salamander, Texas blind (Typhlomolge rathbuni) - Wherever found   | E      | \$19,500      | \$1,302,490,133    |
| 889  | Chub, Hutton tui (Gila bicolor ssp.) - Wherever found   | Т      | \$19,480      | \$1,302,509,613    |
| 890  | Haha (Cyanea grimesiana ssp. grimesiana) - Wherever found   | E      | \$19,277      | \$1,302,528,890    |
| 891  | Cactus, San Rafael (Pediocactus despainii) - Wherever found   | E      | \$19,236      | \$1,302,548,126    |
| 892  | Rattleweed, hairy (Baptisia arachnifera) - Wherever found   | Е      | \$19,200      | \$1,302,567,326    |
| 893  | Trout, Paiute cutthroat (Oncorhynchus clarkii seleniris) - Wherever found   | Т      | \$19,137      | \$1,302,586,463    |
| 894  | Lilliput, pale (pearlymussel) (Toxolasma cylindrellus) - Wherever found   | E      | \$19,120      | \$1,302,605,583    |
| 895  | Fly, Hawaiian picture-wing (Drosophila sharpi) - Wherever found   | E      | \$19,067      | \$1,302,624,650    |
| 896  | Monkeyface, Cumberland (pearlymussel) (Quadrula intermedia) - Wherever found; Except where listed as Experimental Populations | Е      | \$19,000      | \$1,302,643,650    |
| 896  | Isopod, Lee County cave (Lirceus usdagalun) - Wherever found  | E      | \$19,000      | \$1,302,662,650    |
| 897  | Tree snail, Newcomb's (Newcombia cumingi) - Wherever found  | Е      | \$18,824      | \$1,302,681,474    |
| 898  | Pigtoe, Georgia (Pleurobema hanleyianum) - Wherever found   | E      | \$18,800      | \$1,302,700,274    |
| 899  | Ihi`ihi (Marsilea villosa) - Wherever found   | E      | \$18,786      | \$1,302,719,060    |
| 900  | Turtle, ringed map (Graptemys oculifera) - Wherever found   | Т      | \$18,765      | \$1,302,737,825    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 901  | Fly, Hawaiian picture-wing (Drosophila musaphilia) - Wherever found   | Е      | \$18,733      | \$1,302,756,558    |
| 902  | Vole, Florida salt marsh (Microtus pennsylvanicus dukecampbelli) - Wherever found                           | E      | \$18,700      | \$1,302,775,258    |
| 903  | Fern, pendant kihi (Adenophorus periens) - Wherever found   | E      | \$18,698      | \$1,302,793,956    |
| 904  | No Common Name (Asplenium dielmannii) - Wherever found  | E      | \$18,595      | \$1,302,812,551    |
| 905  | Alani (Melicope lydgatei) - Wherever found  | E      | \$18,574      | \$1,302,831,125    |
| 906  | Megapode, Micronesian (Megapodius laperouse) - Wherever found   | E      | \$18,527      | \$1,302,849,652    |
| 907  | Fox, Santa Catalina Island (Urocyon littoralis catalinae) - Wherever found                                  | Т      | \$18,505      | \$1,302,868,157    |
| 908  | Kauila (Colubrina oppositifolia) - Wherever found   | E      | \$18,419      | \$1,302,886,576    |
| 909  | Wild-buckwheat, gypsum (Eriogonum gypsophilum) - Wherever found   | Т      | \$18,200      | \$1,302,904,776    |
| 910  | No Common Name (Platanthera holochila) - Wherever found   | Е      | \$18,094      | \$1,302,922,870    |
| 911  | Haha (Cyanea humboldtiana) - Wherever found   | E      | \$18,090      | \$1,302,940,960    |
| 912  | Centaury, spring-loving (Centaurium namophilum) - Wherever found  | Т      | \$17,987      | \$1,302,958,947    |
| 913  | Amphipod, Hay's Spring (Stygobromus hayi) - Wherever found  | E      | \$17,976      | \$1,302,976,923    |
| 914  | `Awikiwiki (Canavalia molokaiensis) - Wherever found  | E      | \$17,964      | \$1,302,994,887    |
| 915  | Wild buckwheat, clay-loving (Eriogonum pelinophilum) - Wherever found                                       | E      | \$17,905      | \$1,303,012,792    |
| 916  | Toad, Puerto Rican crested (Peltophryne lemur) - Wherever found   | Т      | \$17,900      | \$1,303,030,692    |
| 916  | Wireweed (Polygonella basiramia) - Wherever found   | E      | \$17,900      | \$1,303,048,592    |
| 917  | ,   | E      | \$17,806      | \$1,303,066,398    |
| 918  | Rock-cress, Braun's (Arabis perstellata) - Wherever found   | E      | \$17,720      | \$1,303,084,118    |
| 919  | Rosemary, Cumberland (Conradina verticillata) - Wherever found  | Т      | \$17,680      | \$1,303,101,798    |
| 920  | Ceanothus, Pine Hill (Ceanothus roderickii) - Wherever found  | Е      | \$17,649      | \$1,303,119,447    |
| 921  | Sunburst, San Joaquin adobe (Pseudobahia peirsonii) - Wherever found  | Т      | \$17,628      | \$1,303,137,075    |
| 922  | Lampmussel, Alabama (Lampsilis virescens) - Wherever found; Except where listed as Experimental Populations | Е      | \$17,600      | \$1,303,154,675    |
| 922  | Amphipod, Peck's cave (Stygobromus (=Stygonectes) pecki) - Wherever found                                   | Е      | \$17,600      | \$1,303,172,275    |
| 922  | Dawn-flower, Texas prairie (Hymenoxys texana) -<br>Wherever found   | E      | \$17,600      | \$1,303,189,875    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 923  | Springfish, Hiko White River (Crenichthys baileyi grandis) - Wherever found             | Е      | \$17,568      | \$1,303,207,443    |
| 924  | Pocketbook, speckled (Lampsilis streckeri) - Wherever found                             | E      | \$17,500      | \$1,303,224,943    |
| 925  | Darter, yellowcheek (Etheostoma moorei) - Wherever found                                | E      | \$17,400      | \$1,303,242,343    |
| 926  | `Ohe`ohe (Polyscias gymnocarpa) - Wherever found  | E      | \$17,303      | \$1,303,259,646    |
| 927  | Wild coffee, Oahu (=kopiko) (Psychotria hexandra ssp. oahuensis) - Wherever found       | E      | \$17,280      | \$1,303,276,926    |
| 928  | No Common Name (Polyscias racemosa) - Wherever found                                    | E      | \$17,223      | \$1,303,294,149    |
| 929  | No Common Name (Tetramolopium remyi) - Wherever found                                   | E      | \$17,209      | \$1,303,311,358    |
| 930  | Goetzea, beautiful (Goetzea elegans) - Wherever found                                   | E      | \$17,100      | \$1,303,328,458    |
| 931  | Haiwale (Cyrtandra paliku) - Wherever found   | E      | \$17,067      | \$1,303,345,525    |
| 932  | Haha (Cyanea kolekoleensis) - Wherever found  | E      | \$17,023      | \$1,303,362,548    |
| 933  | Snakeroot (Eryngium cuneifolium) - Wherever found                                       | E      | \$17,000      | \$1,303,379,548    |
| 934  | Koki`o (Kokia kauaiensis) - Wherever found  | E      | \$16,913      | \$1,303,396,461    |
| 935  | Fatmucket, Arkansas (Lampsilis powellii) - Wherever found                               | Т      | \$16,901      | \$1,303,413,362    |
| 936  | Heau (Exocarpos luteolus) - Wherever found  | E      | \$16,819      | \$1,303,430,181    |
| 937  | Sculpin, Grotto (Cottus specus) - Wherever found  | E      | \$16,818      | \$1,303,446,999    |
| 938  | No Common Name (Schiedea membranacea) - Wherever found                                  | E      | \$16,792      | \$1,303,463,791    |
| 939  | Crayfish, Nashville (Orconectes shoupi) - Wherever found                                | E      | \$16,700      | \$1,303,480,491    |
| 940  | Cactus, Siler pincushion (Pediocactus (=Echinocactus,=Utahia) sileri) - Wherever found  | Т      | \$16,600      | \$1,303,497,091    |
| 941  | Salamander, San Marcos (Eurycea nana) - Wherever found                                  | Т      | \$16,500      | \$1,303,513,591    |
| 942  | Ambersnail, Kanab (Oxyloma haydeni kanabensis) -<br>Wherever found                      | E      | \$16,300      | \$1,303,529,891    |
| 943  | Damselfly, blackline Hawaiian (Megalagrion nigrohamatum nigrolineatum) - Wherever found | E      | \$16,296      | \$1,303,546,187    |
| 944  | Kuahiwi laukahi (Plantago hawaiensis) - Wherever found                                  | E      | \$16,252      | \$1,303,562,439    |
| 945  | No Common Name (Pteris lidgatei) - Wherever found                                       | Е      | \$16,200      | \$1,303,578,639    |
| 946  | Haha (Cyanea rivularis) - Wherever found  | E      | \$16,155      | \$1,303,594,794    |
| 947  | Iliau, dwarf (Wilkesia hobdyi) - Wherever found   | E      | \$16,110      | \$1,303,610,904    |
| 948  | Sucker, Zuni bluehead (Catostomus discobolus yarrowi) - Wherever found                  | Е      | \$15,983      | \$1,303,626,887    |
| 949  | No Common Name (Maesa walkeri) - Wherever found   | Т      | \$15,932      | \$1,303,642,819    |
| 950  | No Common Name (Amaranthus brownii) - Wherever found                                    | Е      | \$15,705      | \$1,303,658,524    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 951  | Rabbitsfoot, rough (Quadrula cylindrica strigillata) - Wherever found   | Е      | \$15,700      | \$1,303,674,224    |
| 951  | Cladonia, Florida perforate (Cladonia perforata) - Wherever found       | Е      | \$15,700      | \$1,303,689,924    |
| 952  | Mint, Lakela's (Dicerandra immaculata) - Wherever found                 | E      | \$15,500      | \$1,303,705,424    |
| 952  | No Common Name (Schoepfia arenaria) - Wherever found                    | Т      | \$15,500      | \$1,303,720,924    |
| 953  | `Oha wai (Clermontia pyrularia) - Wherever found                        | E      | \$15,463      | \$1,303,736,387    |
| 954  | Haha (Cyanea kuhihewa) - Wherever found                                 | E      | \$15,428      | \$1,303,751,815    |
| 954  | Haha (Cyanea eleeleensis) - Wherever found                              | E      | \$15,428      | \$1,303,767,243    |
| 954  | Haha (Cyanea dolichopoda) - Wherever found                              | E      | \$15,428      | \$1,303,782,671    |
| 955  | No Common Name (Platydesma remyi) - Wherever found                      | E      | \$15,408      | \$1,303,798,079    |
| 955  | No Common Name (Cyrtandra wagneri) - Wherever found                     | E      | \$15,408      | \$1,303,813,487    |
| 956  | Kaulu (Pteralyxia macrocarpa) - Wherever found                          | E      | \$15,369      | \$1,303,828,856    |
| 957  | Haha (Cyanea stjohnii) - Wherever found                                 | E      | \$15,322      | \$1,303,844,178    |
| 958  | Sandwort, Cumberland (Arenaria cumberlandensis) - Wherever found        | Е      | \$15,220      | \$1,303,859,398    |
| 959  | Leopard, snow (Uncia (=Panthera) uncia) - Wherever found                | Е      | \$15,200      | \$1,303,874,598    |
| 959  | Snake, Atlantic salt marsh (Nerodia clarkii taeniata) - Wherever found  | Т      | \$15,200      | \$1,303,889,798    |
| 960  | A`e (Zanthoxylum oahuense) - Wherever found                             | E      | \$15,135      | \$1,303,904,933    |
| 961  | Pinkroot, gentian (Spigelia gentianoides) - Wherever found              | E      | \$15,058      | \$1,303,919,991    |
| 962  | `Anunu (Sicyos albus) - Wherever found                                  | E      | \$15,019      | \$1,303,935,010    |
| 963  | Madtom, chucky (Noturus crypticus) - Wherever found                     | E      | \$15,000      | \$1,303,950,010    |
| 963  | Beetle, Coffin Cave mold (Batrisodes texanus) - Wherever found          | Е      | \$15,000      | \$1,303,965,010    |
| 963  | No Common Name (Eugenia woodburyana) - Wherever found                   | Е      | \$15,000      | \$1,303,980,010    |
| 963  | Quillwort, black spored (Isoetes melanospora) - Wherever found          | Е      | \$15,000      | \$1,303,995,010    |
| 964  | Turtle, yellow-blotched map (Graptemys flavimaculata) - Wherever found  | Т      | \$14,893      | \$1,304,009,903    |
| 965  | No Common Name (Pittosporum hawaiiense) - Wherever found                | Е      | \$14,820      | \$1,304,024,723    |
| 966  | Bedstraw, El Dorado (Galium californicum ssp. sierrae) - Wherever found | Е      | \$14,810      | \$1,304,039,533    |
| 967  | Beaked-rush, Knieskern's (Rhynchospora knieskernii) - Wherever found    | Т      | \$14,806      | \$1,304,054,339    |
| 968  | Pua `ala (Brighamia rockii) - Wherever found                            | E      | \$14,769      | \$1,304,069,108    |
| 969  | `Aiea (Nothocestrum peltatum) - Wherever found                          | E      | \$14,764      | \$1,304,083,872    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 969  | No Common Name (Remya montgomeryi) - Wherever found                       | Е      | \$14,764      | \$1,304,098,636    |
| 970  | No Common Name (Lobelia oahuensis) - Wherever found                       | E      | \$14,744      | \$1,304,113,380    |
| 971  | Cactus, Pariette (Sclerocactus brevispinus) - Wherever found              | Т      | \$14,726      | \$1,304,128,106    |
| 972  | Haha (Cyanea hamatiflora ssp. carlsonii) - Wherever found                 | E      | \$14,672      | \$1,304,142,778    |
| 973  | Grass, Tennessee yellow-eyed (Xyris tennesseensis) - Wherever found       | Е      | \$14,653      | \$1,304,157,431    |
| 974  | Tuctoria, Greene's (Tuctoria greenei) - Wherever found                    | E      | \$14,567      | \$1,304,171,998    |
| 975  | Alani (Melicope adscendens) - Wherever found                              | E      | \$14,543      | \$1,304,186,541    |
| 976  | No Common Name (Geocarpon minimum) - Wherever found                       | Т      | \$14,503      | \$1,304,201,044    |
| 977  | Iguana, Mona ground (Cyclura stejnegeri) - Wherever found                 | Т      | \$14,500      | \$1,304,215,544    |
| 977  | Monkeyface, Appalachian (pearlymussel) (Quadrula sparsa) - Wherever found | E      | \$14,500      | \$1,304,230,044    |
| 978  | Ha`iwale (Cyrtandra giffardii) - Wherever found                           | E      | \$14,450      | \$1,304,244,494    |
| 979  | Silversword, Mauna Loa (=Ka'u) (Argyroxiphium kauense) - Wherever found   | Е      | \$14,420      | \$1,304,258,914    |
| 980  | Fleabane, Zuni (Erigeron rhizomatus) - Wherever found                     | Т      | \$14,404      | \$1,304,273,318    |
| 981  | Haha (Cyanea lanceolata) - Wherever found                                 | E      | \$14,355      | \$1,304,287,673    |
| 982  | Tarplant, Santa Cruz (Holocarpha macradenia) - Wherever found             | Т      | \$14,303      | \$1,304,301,976    |
| 983  | Cactus, Cochise pincushion (Coryphantha robbinsiorum) - Wherever found    | Т      | \$14,300      | \$1,304,316,276    |
| 984  | Watercress, Gambel's (Rorippa gambellii) - Wherever found                 | Е      | \$14,277      | \$1,304,330,553    |
| 985  | No Common Name (Phyllostegia velutina) - Wherever found                   | Е      | \$14,192      | \$1,304,344,745    |
| 986  | No Common Name (Acropora retusa) - Wherever found                         | Т      | \$14,122      | \$1,304,358,867    |
| 987  | Erubia (Solanum drymophilum) - Wherever found                             | E      | \$14,100      | \$1,304,372,967    |
| 988  | Grass, Colusa (Neostapfia colusana) - Wherever found                      | Т      | \$14,089      | \$1,304,387,056    |
| 989  | Lizard, St. Croix ground (Ameiva polops) - Wherever found                 | E      | \$14,000      | \$1,304,401,056    |
| 990  | No Common Name (Schiedea kauaiensis) - Wherever found                     | Е      | \$13,922      | \$1,304,414,978    |
| 991  | No Common Name (Polyscias bisattenuata) - Wherever found                  | Е      | \$13,904      | \$1,304,428,882    |
| 992  | Snail, painted snake coiled forest (Anguispira picta) - Wherever found    | Т      | \$13,900      | \$1,304,442,782    |
| 993  | Awiwi (Schenkia sebaeoides) - Wherever found                              | E      | \$13,879      | \$1,304,456,661    |
| 994  | No Common Name (Gouania vitifolia) - Wherever found                       | E      | \$13,877      | \$1,304,470,538    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 995  | Lo`ulu (Pritchardia schattaueri) - Wherever found  | E      | \$13,820      | \$1,304,484,358    |
| 996  | Daisy, Lakeside (Hymenoxys herbacea) - Wherever found  | Т      | \$13,809      | \$1,304,498,167    |
| 997  | Kopiko (Psychotria grandiflora) - Wherever found   | E      | \$13,772      | \$1,304,511,939    |
| 998  | No Common Name (Nervilia jacksoniae) - Wherever found  | Т      | \$13,764      | \$1,304,525,703    |
| 999  | Nightjar, Puerto Rican (Caprimulgus noctitherus) - Wherever found                              | Е      | \$13,700      | \$1,304,539,403    |
| 999  | Wire-lettuce, Malheur (Stephanomeria malheurensis) - Wherever found                            | Е      | \$13,700      | \$1,304,553,103    |
| 1000 | Alani (Melicope knudsenii) - Wherever found  | E      | \$13,686      | \$1,304,566,789    |
| 1001 | Ha`iwale (Cyrtandra viridiflora) - Wherever found  | E      | \$13,683      | \$1,304,580,472    |
| 1002 | Butterfly, Mount Charleston blue (Icaricia (Plebejus) shasta charlestonensis) - Wherever found | Е      | \$13,678      | \$1,304,594,150    |
| 1003 | Spurge, Hoover's (Chamaesyce hooveri) - Wherever found   | Т      | \$13,662      | \$1,304,607,812    |
| 1004 | Shrimp, Kentucky cave (Palaemonias ganteri) - Wherever found                                   | Е      | \$13,650      | \$1,304,621,462    |
| 1005 | No Common Name (Silene perlmanii) - Wherever found   | E      | \$13,646      | \$1,304,635,108    |
| 1006 | No Common Name (Phyllostegia pilosa) - Wherever found  | E      | \$13,609      | \$1,304,648,717    |
| 1007 | Orcutt grass, Sacramento (Orcuttia viscida) - Wherever found                                   | Е      | \$13,530      | \$1,304,662,247    |
| 1008 | No Common Name (Silene alexandri) - Wherever found   | E      | \$13,529      | \$1,304,675,776    |
| 1008 | Haha (Cyanea procera) - Wherever found   | E      | \$13,529      | \$1,304,689,305    |
| 1008 | No Common Name (Lysimachia maxima) - Wherever found  | E      | \$13,529      | \$1,304,702,834    |
| 1008 | No Common Name (Phyllostegia mannii) - Wherever found  | E      | \$13,529      | \$1,304,716,363    |
| 1008 | No Common Name (Phyllostegia hispida) - Wherever found   | Е      | \$13,529      | \$1,304,729,892    |
| 1009 | Frankenia, Johnston's (Frankenia johnstonii) - Wherever found                                  | Е      | \$13,500      | \$1,304,743,392    |
| 1010 | Koki`o, Cooke's (Kokia cookei) - Wherever found  | E      | \$13,478      | \$1,304,756,870    |
| 1010 | Alani (Melicope mucronulata) - Wherever found  | E      | \$13,478      | \$1,304,770,348    |
| 1011 | Alani (Melicope degeneri) - Wherever found   | E      | \$13,348      | \$1,304,783,696    |
| 1011 | Kamakahala (Labordia lydgatei) - Wherever found  | E      | \$13,348      | \$1,304,797,044    |
| 1012 | Rose-mallow, Neches River (Hibiscus dasycalyx) - Wherever found                                | Т      | \$13,343      | \$1,304,810,387    |
| 1013 | No Common Name (Phyllostegia warshaueri) - Wherever found                                      | Е      | \$13,311      | \$1,304,823,698    |
| 1014 | Mint, scrub (Dicerandra frutescens) - Wherever found   | E      | \$13,300      | \$1,304,836,998    |
| 1015 | Buckwheat, Umtanum Desert (Eriogonum codium) - Wherever found                                  | Т      | \$13,275      | \$1,304,850,273    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 1016 | Acornshell, southern (Epioblasma othcaloogensis) - Wherever found  | Е      | \$13,201      | \$1,304,863,474    |
| 1017 | Hau kuahiwi (Hibiscadelphus giffardianus) - Wherever found   | Е      | \$13,151      | \$1,304,876,625    |
| 1018 | Hypericum, highlands scrub (Hypericum cumulicola) - Wherever found   | Е      | \$13,100      | \$1,304,889,725    |
| 1018 | Chumbo, Higo (Harrisia portoricensis) - Wherever found   | Т      | \$13,100      | \$1,304,902,825    |
| 1018 | No Common Name (Vernonia proctorii) - Wherever found   | E      | \$13,100      | \$1,304,915,925    |
| 1019 | Darter, goldline (Percina aurolineata) - Wherever found  | Т      | \$13,070      | \$1,304,928,995    |
| 1020 | `Ahinahina (Argyroxiphium sandwicense ssp. sandwicense) - Wherever found                                     | Е      | \$13,033      | \$1,304,942,028    |
| 1021 | Damselfly, oceanic Hawaiian (Megalagrion oceanicum) - Wherever found   | Е      | \$13,014      | \$1,304,955,042    |
| 1022 | Nohoanu (Geranium kauaiense) - Wherever found  | E      | \$13,005      | \$1,304,968,047    |
| 1023 | Pearlymussel, birdwing (Lemiox rimosus) - Wherever found; Except where listed as Experimental Populations    | Е      | \$13,000      | \$1,304,981,047    |
| 1023 | Manaca, palma de (Calyptronoma rivalis) - Wherever found   | Т      | \$13,000      | \$1,304,994,047    |
| 1023 | No Common Name (Myrcia paganii) - Wherever found   | E      | \$13,000      | \$1,305,007,047    |
| 1024 | No Common Name (Pittosporum halophilum) - Wherever found   | Е      | \$12,920      | \$1,305,019,967    |
| 1025 | Pawpaw, four-petal (Asimina tetramera) - Wherever found  | E      | \$12,900      | \$1,305,032,867    |
| 1026 | Gardenia (=Na`u), Hawaiian (Gardenia brighamii) -<br>Wherever found  | Е      | \$12,894      | \$1,305,045,761    |
| 1027 | Phacelia, North Park (Phacelia formosula) - Wherever found   | Е      | \$12,707      | \$1,305,058,468    |
| 1028 | Na`ena`e (Dubautia plantaginea ssp. humilis) - Wherever found  | Е      | \$12,691      | \$1,305,071,159    |
| 1029 | Sculpin, pygmy (Cottus paulus (=pygmaeus)) - Wherever found  | Т      | \$12,667      | \$1,305,083,826    |
| 1030 | `Awikiwiki (Canavalia napaliensis) - Wherever found  | E      | \$12,666      | \$1,305,096,492    |
| 1031 | `aku`aku (Cyanea platyphylla) - Wherever found   | E      | \$12,639      | \$1,305,109,131    |
| 1032 | Mapleleaf, winged (Quadrula fragosa) - U.S.A. (ALspecified portions of the Tennessee River; see 17.85(a)(1)) | EXPN   | \$12,500      | \$1,305,121,631    |
| 1032 | Sunflower, whorled (Helianthus verticillatus) - Wherever found   | Е      | \$12,500      | \$1,305,134,131    |
| 1033 | Haha (Cyanea obtusa) - Wherever found  | E      | \$12,464      | \$1,305,146,595    |
| 1033 | No Common Name (Stenogyne kauaulaensis) - Wherever found   | Е      | \$12,464      | \$1,305,159,059    |
| 1034 | Milkvetch, Osterhout (Astragalus osterhoutii) - Wherever found   | Е      | \$12,441      | \$1,305,171,500    |
| 1035 | Bedstraw, island (Galium buxifolium) - Wherever found  | E      | \$12,404      | \$1,305,183,904    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 1036 | Beetle, Comal Springs riffle (Heterelmis comalensis) - Wherever found      | Е      | \$12,400      | \$1,305,196,304    |
| 1036 | Beetle, Comal Springs dryopid (Stygoparnus comalensis) - Wherever found    | E      | \$12,400      | \$1,305,208,704    |
| 1036 | Rhododendron, Chapman (Rhododendron chapmanii) - Wherever found            | Е      | \$12,400      | \$1,305,221,104    |
| 1037 | Kidneyshell, triangular (Ptychobranchus greenii) -<br>Wherever found       | Е      | \$12,359      | \$1,305,233,463    |
| 1038 | No Common Name (Tetramolopium lepidotum ssp. lepidotum) - Wherever found   | E      | \$12,251      | \$1,305,245,714    |
| 1039 | Meshweaver, Braken Bat Cave (Cicurina venii) - Wherever found              | Е      | \$12,200      | \$1,305,257,914    |
| 1040 | Haha (Cyanea lobata) - Wherever found                                      | E      | \$12,195      | \$1,305,270,109    |
| 1041 | No Common Name (Diplazium molokaiense) - Wherever found                    | Е      | \$12,172      | \$1,305,282,281    |
| 1042 | Springsnail, Koster's (Juturnia kosteri) - Wherever found                  | E      | \$12,150      | \$1,305,294,431    |
| 1043 | Pelos del diablo (Aristida portoricensis) - Wherever found                 | E      | \$12,100      | \$1,305,306,531    |
| 1043 | Palo de ramon (Banara vanderbiltii) - Wherever found                       | E      | \$12,100      | \$1,305,318,631    |
| 1044 | Beetle, Tooth Cave ground (Rhadine persephone) - Wherever found            | Е      | \$12,000      | \$1,305,330,631    |
| 1044 | Beetle, Helotes mold (Batrisodes venyivi) - Wherever found                 | E      | \$12,000      | \$1,305,342,631    |
| 1044 | Spider, Government Canyon Bat Cave (Neoleptoneta microps) - Wherever found | Е      | \$12,000      | \$1,305,354,631    |
| 1044 | Meshweaver, Government Canyon Bat Cave (Cicurina vespera) - Wherever found | Е      | \$12,000      | \$1,305,366,631    |
| 1045 | Haha (Cyanea truncata) - Wherever found                                    | E      | \$11,981      | \$1,305,378,612    |
| 1046 | Rock-cress, Hoffmann's (Arabis hoffmannii) - Wherever found                | Е      | \$11,976      | \$1,305,390,588    |
| 1047 | No Common Name (Delissea undulata) - Wherever found                        | E      | \$11,958      | \$1,305,402,546    |
| 1048 | Lo`ulu (Pritchardia maideniana) - Wherever found                           | E      | \$11,948      | \$1,305,414,494    |
| 1049 | No Common Name (Bonamia menziesii) - Wherever found                        | E      | \$11,947      | \$1,305,426,441    |
| 1050 | `Oha wai (Clermontia oblongifolia ssp. brevipes) - Wherever found          | E      | \$11,931      | \$1,305,438,372    |
| 1051 | `Ala `ala wai nui (Peperomia subpetiolata) - Wherever found                | Е      | \$11,926      | \$1,305,450,298    |
| 1052 | Pamakani (Tetramolopium capillare) - Wherever found                        | E      | \$11,890      | \$1,305,462,188    |
| 1052 | Kamakahala (Labordia tinifolia var. lanaiensis) - Wherever found           | E      | \$11,890      | \$1,305,474,078    |
| 1052 | No Common Name (Lysimachia lydgatei) - Wherever found                      | E      | \$11,890      | \$1,305,485,968    |
| 1052 | Kohe malama malama o kanaloa (Kanaloa kahoolawensis) - Wherever found      | Е      | \$11,890      | \$1,305,497,858    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 1053 | Rush-rose, island (Helianthemum greenei) - Wherever found                                    | Т      | \$11,880      | \$1,305,509,738    |
| 1053 | Schiedea, Diamond Head (Schiedea adamantis) - Wherever found                                 | E      | \$11,880      | \$1,305,521,618    |
| 1053 | Opuhe (Urera kaalae) - Wherever found  | E      | \$11,880      | \$1,305,533,498    |
| 1054 | Bush-mallow, Santa Cruz Island (Malacothamnus fasciculatus var. nesioticus) - Wherever found | Е      | \$11,876      | \$1,305,545,374    |
| 1054 | Phacelia, island (Phacelia insularis ssp. insularis) - Wherever found                        | E      | \$11,876      | \$1,305,557,250    |
| 1055 | Hau kuahiwi (Hibiscadelphus hualalaiensis) - Wherever found                                  | E      | \$11,875      | \$1,305,569,125    |
| 1056 | Goldenrod, Short's (Solidago shortii) - Wherever found                                       | E      | \$11,860      | \$1,305,580,985    |
| 1057 | Olulu (Brighamia insignis) - Wherever found  | E      | \$11,843      | \$1,305,592,828    |
| 1058 | Rockcress, Georgia (Arabis georgiana) - Wherever found                                       | Т      | \$11,808      | \$1,305,604,636    |
| 1059 | No Common Name (Schiedea diffusa ssp. macraei) - Wherever found                              | Е      | \$11,764      | \$1,305,616,400    |
| 1060 | Alani (Melicope haupuensis) - Wherever found   | E      | \$11,744      | \$1,305,628,144    |
| 1060 | No Common Name (Viola helenae) - Wherever found  | E      | \$11,744      | \$1,305,639,888    |
| 1060 | Nani wai`ale`ale (Viola kauaiensis var. wahiawaensis) -<br>Wherever found                    | Е      | \$11,744      | \$1,305,651,632    |
| 1060 | Kolea (Myrsine mezii) - Wherever found   | E      | \$11,744      | \$1,305,663,376    |
| 1060 | Haha (Cyanea undulata) - Wherever found  | E      | \$11,744      | \$1,305,675,120    |
| 1060 | Na`ena`e (Dubautia pauciflorula) - Wherever found  | E      | \$11,744      | \$1,305,686,864    |
| 1060 | No Common Name (Phyllostegia waimeae) - Wherever found                                       | Е      | \$11,744      | \$1,305,698,608    |
| 1060 | Lo`ulu (Pritchardia viscosa) - Wherever found  | E      | \$11,744      | \$1,305,710,352    |
| 1060 | Kolea (Myrsine knudsenii) - Wherever found   | E      | \$11,744      | \$1,305,722,096    |
| 1061 | Combshell, southern (Epioblasma penita) - Wherever found                                     | Е      | \$11,734      | \$1,305,733,830    |
| 1062 | No Common Name (Lysimachia pendens) - Wherever found   | Е      | \$11,700      | \$1,305,745,530    |
| 1063 | No Common Name (Delissea rhytidosperma) - Wherever found                                     | E      | \$11,693      | \$1,305,757,223    |
| 1063 | No Common Name (Stenogyne campanulata) - Wherever found                                      | E      | \$11,693      | \$1,305,768,916    |
| 1063 | No Common Name (Stenogyne kealiae) - Wherever found  | E      | \$11,693      | \$1,305,780,609    |
| 1064 | Beetle, Mount Hermon June (Polyphylla barbata) -<br>Wherever found                           | Е      | \$11,681      | \$1,305,792,290    |
| 1064 | Pentachaeta, Lyon's (Pentachaeta Iyonii) - Wherever found                                    | E      | \$11,681      | \$1,305,803,971    |
| 1064 | Tarplant, Gaviota (Deinandra increscens ssp. villosa) - Wherever found                       | Е      | \$11,681      | \$1,305,815,652    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 1064 | Monkeyflower, Vandenberg (Diplacus vandenbergensis) - Wherever found                | Е      | \$11,681      | \$1,305,827,333    |
| 1065 | Poppy, Sacramento prickly (Argemone pleiacantha ssp. pinnatisecta) - Wherever found | E      | \$11,660      | \$1,305,838,993    |
| 1066 | Pussypaws, Mariposa (Calyptridium pulchellum) - Wherever found                      | Т      | \$11,620      | \$1,305,850,613    |
| 1066 | Morning-glory, Stebbins' (Calystegia stebbinsii) - Wherever found                   | Е      | \$11,620      | \$1,305,862,233    |
| 1067 | No Common Name (Cyanea purpurellifolia) - Wherever found                            | Е      | \$11,587      | \$1,305,873,820    |
| 1068 | Haha (Cyanea asarifolia) - Wherever found   | E      | \$11,553      | \$1,305,885,373    |
| 1069 | Ha`iwale (Cyrtandra kaulantha) - Wherever found                                     | E      | \$11,536      | \$1,305,896,909    |
| 1069 | No Common Name (Cyrtandra gracilis) - Wherever found                                | E      | \$11,536      | \$1,305,908,445    |
| 1069 | Alani (Melicope hiiakae) - Wherever found   | E      | \$11,536      | \$1,305,919,981    |
| 1069 | Haha (Cyanea remyi) - Wherever found  | E      | \$11,536      | \$1,305,931,517    |
| 1070 | No Common Name (Lobelia monostachya) - Wherever found                               | E      | \$11,503      | \$1,305,943,020    |
| 1071 | Ha`iwale (Cyrtandra oenobarba) - Wherever found                                     | E      | \$11,502      | \$1,305,954,522    |
| 1072 | Sea turtle, green (Chelonia mydas) - Mediterranean DPS                              | E      | \$11,500      | \$1,305,966,022    |
| 1072 | Sea turtle, loggerhead (Caretta caretta) - Mediterranean<br>Sea DPS                 | E      | \$11,500      | \$1,305,977,522    |
| 1072 | No Common Name (Cranichis ricartii) - Wherever found                                | E      | \$11,500      | \$1,305,989,022    |
| 1073 | `Akoko (Euphorbia haeleeleana) - Wherever found                                     | E      | \$11,472      | \$1,306,000,494    |
| 1074 | Pu`uka`a (Cyperus trachysanthos) - Wherever found                                   | E      | \$11,468      | \$1,306,011,962    |
| 1075 | Mapele (Cyrtandra cyaneoides) - Wherever found                                      | E      | \$11,377      | \$1,306,023,339    |
| 1076 | No Common Name (Schiedea laui) - Wherever found                                     | E      | \$11,366      | \$1,306,034,705    |
| 1077 | Madtom, pygmy (Noturus stanauli) - Wherever found                                   | E      | \$11,360      | \$1,306,046,065    |
| 1078 | Cactus, Key tree (Pilosocereus robinii) - Wherever found                            | E      | \$11,318      | \$1,306,057,383    |
| 1079 | Bush-clover, prairie (Lespedeza leptostachya) - Wherever found                      | Т      | \$11,260      | \$1,306,068,643    |
| 1080 | Kolea (Myrsine linearifolia) - Wherever found                                       | Т      | \$11,227      | \$1,306,079,870    |
| 1081 | Lupine, clover (Lupinus tidestromii) - Wherever found                               | E      | \$11,194      | \$1,306,091,064    |
| 1082 | No Common Name (Abutilon eremitopetalum) - Wherever found                           | E      | \$11,181      | \$1,306,102,245    |
| 1082 | Haha (Cyanea magnicalyx) - Wherever found   | E      | \$11,181      | \$1,306,113,426    |
| 1083 | No Common Name (Achyranthes mutica) - Wherever found                                | E      | \$11,120      | \$1,306,124,546    |
| 1084 | No Common Name (Aristida chaseae) - Wherever found                                  | E      | \$11,100      | \$1,306,135,646    |
| 1084 | No Common Name (Lyonia truncata var. proctorii) - Wherever found                    | Е      | \$11,100      | \$1,306,146,746    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 1085 | Sunburst, Hartweg's golden (Pseudobahia bahiifolia) - Wherever found           | Е      | \$11,090      | \$1,306,157,836    |
| 1086 | Haha (Cyanea calycina) - Wherever found  | E      | \$11,077      | \$1,306,168,913    |
| 1087 | Chub, Chihuahua (Gila nigrescens) - Wherever found                             | Т      | \$11,076      | \$1,306,179,989    |
| 1088 | Manzanita, pallid (Arctostaphylos pallida) - Wherever found                    | Т      | \$11,000      | \$1,306,190,989    |
| 1088 | Larkspur, Baker's (Delphinium bakeri) - Wherever found                         | E      | \$11,000      | \$1,306,201,989    |
| 1088 | Blazingstar, scrub (Liatris ohlingerae) - Wherever found                       | E      | \$11,000      | \$1,306,212,989    |
| 1088 | No Common Name (Cordia bellonis) - Wherever found                              | E      | \$11,000      | \$1,306,223,989    |
| 1088 | Palo de nigua (Cornutia obovata) - Wherever found                              | E      | \$11,000      | \$1,306,234,989    |
| 1088 | Uvillo (Eugenia haematocarpa) - Wherever found                                 | E      | \$11,000      | \$1,306,245,989    |
| 1088 | No Common Name (Ilex sintenisii) - Wherever found                              | E      | \$11,000      | \$1,306,256,989    |
| 1089 | Lehua makanoe (Lysimachia daphnoides) - Wherever found                         | Е      | \$10,977      | \$1,306,267,966    |
| 1089 | Alani (Melicope puberula) - Wherever found                                     | E      | \$10,977      | \$1,306,278,943    |
| 1090 | No Common Name (Polyscias flynnii) - Wherever found                            | E      | \$10,913      | \$1,306,289,856    |
| 1091 | Springsnail, Alamosa (Tryonia alamosae) - Wherever found                       | Е      | \$10,905      | \$1,306,300,761    |
| 1092 | Haha (Cyanea dunbariae) - Wherever found                                       | E      | \$10,904      | \$1,306,311,665    |
| 1093 | Finch, Laysan (honeycreeper) (Telespyza cantans) - Wherever found              | Е      | \$10,898      | \$1,306,322,563    |
| 1094 | `Awikiwiki (Canavalia pubescens) - Wherever found                              | E      | \$10,876      | \$1,306,333,439    |
| 1094 | No Common Name (Schiedea jacobii) - Wherever found                             | E      | \$10,876      | \$1,306,344,315    |
| 1095 | Milkpea, Small's (Galactia smallii) - Wherever found                           | E      | \$10,800      | \$1,306,355,115    |
| 1096 | Fox, Santa Cruz Island (Urocyon littoralis santacruzae) - wherever found       | Е      | \$10,773      | \$1,306,365,888    |
| 1097 | Bat, Pacific sheath-tailed (Emballonura semicaudata rotensis) - Wherever found | Е      | \$10,760      | \$1,306,376,648    |
| 1098 | No Common Name (Lobelia niihauensis) - Wherever found                          | Е      | \$10,749      | \$1,306,387,397    |
| 1099 | Chub, slender (Erimystax cahni) - Wherever found                               | Т      | \$10,740      | \$1,306,398,137    |
| 1100 | Jacquemontia, beach (Jacquemontia reclinata) - Wherever found                  | Е      | \$10,700      | \$1,306,408,837    |
| 1101 | No Common Name (Sanicula purpurea) - Wherever found                            | E      | \$10,686      | \$1,306,419,523    |
| 1102 | Kamakahala (Labordia tinifolia var. wahiawaensis) -<br>Wherever found          | Е      | \$10,666      | \$1,306,430,189    |
| 1103 | Thornmint, San Diego (Acanthomintha ilicifolia) - Wherever found               |        | \$10,620      | \$1,306,440,809    |
| 1104 | No Common Name (Polyscias lydgatei) - Wherever found                           | E      | \$10,535      | \$1,306,451,344    |
| 1105 | Aupaka (Isodendrion laurifolium) - Wherever found                              | E      | \$10,528      | \$1,306,461,872    |
| 1105 | Dudleya, Verity's (Dudleya verityi) - Wherever found                           | Т      | \$10,528      | \$1,306,472,400    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 1106 | Cactus, star (Astrophytum asterias) - Wherever found  | E      | \$10,500      | \$1,306,482,900    |
| 1106 | Boxwood, Vahl's (Buxus vahlii) - Wherever found   | E      | \$10,500      | \$1,306,493,400    |
| 1106 | Quillwort, mat-forming (Isoetes tegetiformans) - Wherever found   | Е      | \$10,500      | \$1,306,503,900    |
| 1107 | Tarplant, Otay (Deinandra (=Hemizonia) conjugens) - Wherever found  | Т      | \$10,438      | \$1,306,514,338    |
| 1108 | Dogweed, ashy (Thymophylla tephroleuca) - Wherever found  | Е      | \$10,400      | \$1,306,524,738    |
| 1109 | Turtle, flattened musk (Sternotherus depressus) - Black Warrior R. system upstream from Bankhead Dam  | Т      | \$10,313      | \$1,306,535,051    |
| 1110 | Haha (Cyanea munroi) - Wherever found   | E      | \$10,299      | \$1,306,545,350    |
| 1111 | Ha`iwale (Cyrtandra polyantha) - Wherever found   | E      | \$10,292      | \$1,306,555,642    |
| 1111 | Haha (Cyanea pinnatifida) - Wherever found  | E      | \$10,292      | \$1,306,565,934    |
| 1112 | Sneezeweed, Virginia (Helenium virginicum) - Wherever found   | Т      | \$10,229      | \$1,306,576,163    |
| 1113 | No Common Name (Gouania meyenii) - Wherever found   | E      | \$10,207      | \$1,306,586,370    |
| 1113 | Bladderpod, kodachrome (Lesquerella tumulosa) - Wherever found  | Е      | \$10,207      | \$1,306,596,577    |
| 1114 | Mussel, oyster (Epioblasma capsaeformis) - AL; Free-<br>Flowing Reach of the Tennessee River below the Wilson<br>Dam, Colbert and Lauderdale Counties, AL | EXPN   | \$10,200      | \$1,306,606,777    |
| 1115 | Naenae (Dubautia kalalauensis) - Wherever found   | E      | \$10,156      | \$1,306,616,933    |
| 1115 | No Common Name (Lysimachia iniki) - Wherever found  | E      | \$10,156      | \$1,306,627,089    |
| 1116 | Clarkia, Vine Hill (Clarkia imbricata) - Wherever found   | E      | \$10,139      | \$1,306,637,228    |
| 1117 | Combshell, upland (Epioblasma metastriata) - Wherever found   | Е      | \$10,121      | \$1,306,647,349    |
| 1118 | No Common Name (Phyllostegia knudsenii) - Wherever found  | Е      | \$10,105      | \$1,306,657,454    |
| 1118 | Kauai hau kuahiwi (Hibiscadelphus distans) - Wherever found   | E      | \$10,105      | \$1,306,667,559    |
| 1118 | Alani (Melicope quadrangularis) - Wherever found  | E      | \$10,105      | \$1,306,677,664    |
| 1118 | No Common Name (Phyllostegia wawrana) - Wherever found  | Е      | \$10,105      | \$1,306,687,769    |
| 1118 | Hau kuahiwi (Hibiscadelphus woodii) - Wherever found  | E      | \$10,105      | \$1,306,697,874    |
| 1118 | No Common Name (Lysimachia venosa) - Wherever found   | E      | \$10,105      | \$1,306,707,979    |
| 1118 | No Common Name (Schiedea attenuata) - Wherever found  | Е      | \$10,105      | \$1,306,718,084    |
| 1118 | Naenae (Dubautia kenwoodii) - Wherever found  | Е      | \$10,105      | \$1,306,728,189    |
| 1119 | Pigtoe, Cumberland (Pleurobema gibberum) - Wherever found   | Е      | \$10,100      | \$1,306,738,289    |
| 1119 | No Common Name (Varronia rupicola) - Wherever found   | Т      | \$10,100      | \$1,306,748,389    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 1119 | Cactus, Peebles Navajo (Pediocactus peeblesianus var. peeblesianus) - Wherever found | Е      | \$10,100      | \$1,306,758,489    |
| 1119 | Higuero de sierra (Crescentia portoricensis) - Wherever found                        | E      | \$10,100      | \$1,306,768,589    |
| 1120 | Paintbrush, soft-leaved (Castilleja mollis) - Wherever found                         | E      | \$10,072      | \$1,306,778,661    |
| 1121 | No Common Name (Lysimachia scopulensis) - Wherever found                             | E      | \$10,061      | \$1,306,788,722    |
| 1122 | Diellia, Asplenium-leaved (Asplenium dielerectum) - Wherever found                   | E      | \$10,046      | \$1,306,798,768    |
| 1123 | Beardtongue, Parachute (Penstemon debilis) - Wherever found                          | Т      | \$10,007      | \$1,306,808,775    |
| 1124 | Gecko, Monito (Sphaerodactylus micropithecus) - Wherever found                       | E      | \$10,000      | \$1,306,818,775    |
| 1124 | Salamander, Georgetown (Eurycea naufragia) - Wherever found                          | Т      | \$10,000      | \$1,306,828,775    |
| 1124 | Salamander, Salado (Eurycea chisholmensis) - Wherever found                          | Т      | \$10,000      | \$1,306,838,775    |
| 1124 | No Common Name (Daphnopsis hellerana) - Wherever found                               | Е      | \$10,000      | \$1,306,848,775    |
| 1124 | Vervain, Red Hills (Verbena californica) - Wherever found                            | Т      | \$10,000      | \$1,306,858,775    |
| 1125 | No Common Name (Platydesma cornuta cornuta) - Wherever found                         | E      | \$9,955       | \$1,306,868,730    |
| 1126 | No Common Name (Cyrtandra waiolani) - Wherever found                                 | E      | \$9,948       | \$1,306,878,678    |
| 1126 | No Common Name (Stenogyne cranwelliae) - Wherever found                              | E      | \$9,948       | \$1,306,888,626    |
| 1127 | Haha (Cyanea marksii) - Wherever found   | E      | \$9,938       | \$1,306,898,564    |
| 1128 | Phlox, Yreka (Phlox hirsuta) - Wherever found  | E      | \$9,900       | \$1,306,908,464    |
| 1129 | Ha`iwale (Cyrtandra tintinnabula) - Wherever found                                   | E      | \$9,894       | \$1,306,918,358    |
| 1130 | No Common Name (Viola oahuensis) - Wherever found                                    | E      | \$9,757       | \$1,306,928,115    |
| 1130 | Kolea (Myrsine juddii) - Wherever found  | E      | \$9,757       | \$1,306,937,872    |
| 1130 | Haha (Cyanea acuminata) - Wherever found   | E      | \$9,757       | \$1,306,947,629    |
| 1131 | Shrimp, anchialine pool (Vetericaris chaceorum) - Wherever found                     | Е      | \$9,683       | \$1,306,957,312    |
| 1131 | Cypress, Santa Cruz (Cupressus abramsiana) - Wherever found                          | Т      | \$9,683       | \$1,306,966,995    |
| 1132 | Love grass, Fosberg's (Eragrostis fosbergii) - Wherever found                        | Е      | \$9,653       | \$1,306,976,648    |
| 1133 | Bird's-beak, soft (Cordylanthus mollis ssp. mollis) - Wherever found                 | Е      | \$9,600       | \$1,306,986,248    |
| 1133 | Bariaco (Trichilia triacantha) - Wherever found                                      | E      | \$9,600       | \$1,306,995,848    |
| 1134 | No Common Name (Asplenium dielpallidum) - Wherever found                             | Е      | \$9,540       | \$1,307,005,388    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 1135 | Phlox, Texas trailing (Phlox nivalis ssp. texensis) - Wherever found        | Е      | \$9,500       | \$1,307,014,888    |
| 1136 | Haha (Cyanea copelandii ssp. copelandii) - Wherever found                   | E      | \$9,414       | \$1,307,024,302    |
| 1137 | Alani (Melicope paniculata) - Wherever found                                | E      | \$9,371       | \$1,307,033,673    |
| 1138 | Halomtano, Berenghenas (Solanum guamense) - Wherever found                  | E      | \$9,338       | \$1,307,043,011    |
| 1139 | No Common Name (Tinospora homosepala) - Wherever found                      | Е      | \$9,308       | \$1,307,052,319    |
| 1140 | Isopod, Socorro (Thermosphaeroma thermophilus) -<br>Wherever found          | Е      | \$9,298       | \$1,307,061,617    |
| 1141 | Snail, Chittenango ovate amber (Succinea chittenangoensis) - Wherever found | Т      | \$9,279       | \$1,307,070,896    |
| 1142 | No Common Name (Cyrtandra nanawaleensis) - Wherever found                   | Е      | \$9,250       | \$1,307,080,146    |
| 1143 | No Common Name (Remya kauaiensis) - Wherever found                          | E      | \$9,221       | \$1,307,089,367    |
| 1144 | Flax, Carter's small-flowered (Linum carteri carteri) - Wherever found      | Е      | \$9,200       | \$1,307,098,567    |
| 1145 | Bluegrass, Hawaiian (Poa sandvicensis) - Wherever found                     | E      | \$9,199       | \$1,307,107,766    |
| 1145 | No Common Name (Poa siphonoglossa) - Wherever found                         | E      | \$9,199       | \$1,307,116,965    |
| 1145 | Haha (Cyanea recta) - Wherever found  | Т      | \$9,199       | \$1,307,126,164    |
| 1146 | White-eye, Rota bridled (Zosterops rotensis) - Wherever found               | E      | \$9,178       | \$1,307,135,342    |
| 1147 | Kuawawaenohu (Schiedea lychnoides) - Wherever found                         | E      | \$9,176       | \$1,307,144,518    |
| 1148 | No Common Name (Phyllanthus saffordii) - Wherever found                     | E      | \$9,169       | \$1,307,153,687    |
| 1149 | `Akoko (Euphorbia eleanoriae) - Wherever found                              | E      | \$9,161       | \$1,307,162,848    |
| 1150 | Beetle, delta green ground (Elaphrus viridis) - Wherever found              | Т      | \$9,160       | \$1,307,172,008    |
| 1151 | Curassow, blue-billed (Crax alberti) - Wherever found                       | E      | \$9,100       | \$1,307,181,108    |
| 1152 | Palo de jazmin (Styrax portoricensis) - Wherever found                      | E      | \$9,000       | \$1,307,190,108    |
| 1153 | Coqui, Llanero (Eleutherodactylus juanariveroi) - Wherever found            | E      | \$8,990       | \$1,307,199,098    |
| 1154 | `Akoko (Euphorbia rockii) - Wherever found                                  | E      | \$8,970       | \$1,307,208,068    |
| 1155 | Damselfly, crimson Hawaiian (Megalagrion leptodemas) - Wherever found       | Е      | \$8,798       | \$1,307,216,866    |
| 1156 | Creeper, Oahu (Paroreomyza maculata) - Wherever found                       | E      | \$8,760       | \$1,307,225,626    |
| 1157 | Fringe-tree, pygmy (Chionanthus pygmaeus) - Wherever found                  | Е      | \$8,700       | \$1,307,234,326    |
| 1157 | Milk-vetch, Shivwits (Astragalus ampullarioides) - Wherever found           | E      | \$8,700       | \$1,307,243,026    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 1158 | Spineflower, slender-horned (Dodecahema leptoceras) - Wherever found   | E      | \$8,680       | \$1,307,251,706    |
| 1159 | Twinpod, Dudley Bluffs (Physaria obcordata) - Wherever found   | Т      | \$8,648       | \$1,307,260,354    |
| 1160 | Bladderpod, Dudley Bluffs (Lesquerella congesta) -<br>Wherever found   | Т      | \$8,600       | \$1,307,268,954    |
| 1161 | `Akoko (Euphorbia remyi var. remyi) - Wherever found   | E      | \$8,584       | \$1,307,277,538    |
| 1161 | Koki`o ke`oke`o (Hibiscus waimeae ssp. hannerae) -<br>Wherever found   | Е      | \$8,584       | \$1,307,286,122    |
| 1162 | Spider, Kauai cave wolf or pe'e pe'e maka 'ole (Adelocosa anops) - Wherever found  | Е      | \$8,558       | \$1,307,294,680    |
| 1162 | Amphipod, Kauai cave (Spelaeorchestia koloana) - Wherever found  | Е      | \$8,558       | \$1,307,303,238    |
| 1163 | No Common Name (Schiedea viscosa) - Wherever found   | E      | \$8,485       | \$1,307,311,723    |
| 1164 | Na`ena`e (Dubautia waialealae) - Wherever found  | E      | \$8,460       | \$1,307,320,183    |
| 1165 | Gilia, Hoffmann's slender-flowered (Gilia tenuiflora ssp. hoffmannii) - Wherever found   | Е      | \$8,451       | \$1,307,328,634    |
| 1166 | No Common Name (Lobelia koolauensis) - Wherever found  | Е      | \$8,443       | \$1,307,337,077    |
| 1167 | Kaulu (Pteralyxia kauaiensis) - Wherever found   | E      | \$8,357       | \$1,307,345,434    |
| 1168 | Mustard, Carter's (Warea carteri) - Wherever found   | E      | \$8,300       | \$1,307,353,734    |
| 1169 | Alani (Melicope balloui) - Wherever found  | E      | \$8,298       | \$1,307,362,032    |
| 1170 | Madtom, smoky (Noturus baileyi) - Wherever found   | E      | \$8,280       | \$1,307,370,312    |
| 1171 | Pamakani (Viola chamissoniana ssp. chamissoniana) - Wherever found   | E      | \$8,277       | \$1,307,378,589    |
| 1172 | Cactus, Lee pincushion (Coryphantha sneedii var. leei) - Wherever found  | Т      | \$8,270       | \$1,307,386,859    |
| 1173 | Thistle, Sacramento Mountains (Cirsium vinaceum) - Wherever found  | Т      | \$8,250       | \$1,307,395,109    |
| 1174 | Pearlymussel, dromedary (Dromus dromas) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL                  | EXPN   | \$8,200       | \$1,307,403,309    |
| 1175 | Haha (Cyanea duvalliorum) - Wherever found   | E      | \$8,146       | \$1,307,411,455    |
| 1176 | Chub, spotfin (Erimonax monachus) - U.S.A. (TN-specified portions of the Tellico River; see 17.84(m)(1)(i))  | EXPN   | \$8,100       | \$1,307,419,555    |
| 1176 | Bean, Cumberland (pearlymussel) (Villosa trabalis) - AL;<br>Free-Flowing Reach of the Tennessee River below the<br>Wilson Dam, Colbert and Lauderdale Counties, AL | EXPN   | \$8,100       | \$1,307,427,655    |
| 1176 | Combshell, Cumberlandian (Epioblasma brevidens) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL          | EXPN   | \$8,100       | \$1,307,435,755    |
| 1176 | No Common Name (Catesbaea melanocarpa) - Wherever found  | Е      | \$8,100       | \$1,307,443,855    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 1177 | `Oha wai (Clermontia drepanomorpha) - Wherever found                                | E      | \$8,085       | \$1,307,451,940    |
| 1178 | No Common Name (Schiedea hookeri) - Wherever found                                  | E      | \$8,077       | \$1,307,460,017    |
| 1179 | Rush-pea, slender (Hoffmannseggia tenella) - Wherever found                         | Е      | \$8,076       | \$1,307,468,093    |
| 1180 | Holei (Ochrosia kilaueaensis) - Wherever found                                      | E      | \$8,051       | \$1,307,476,144    |
| 1181 | No Common Name (Viola lanaiensis) - Wherever found                                  | E      | \$8,015       | \$1,307,484,159    |
| 1182 | No Common Name (Platydesma cornuta decurrens) - Wherever found                      | E      | \$8,011       | \$1,307,492,170    |
| 1183 | Trout, Little Kern golden (Oncorhynchus aguabonita whitei) - Wherever found         | Т      | \$8,009       | \$1,307,500,179    |
| 1184 | Capa rosa (Callicarpa ampla) - Wherever found                                       | E      | \$8,000       | \$1,307,508,179    |
| 1184 | No Common Name (Chamaecrista glandulosa var. mirabilis) - Wherever found            | E      | \$8,000       | \$1,307,516,179    |
| 1184 | No Common Name (Mitracarpus polycladus) - Wherever found                            | E      | \$8,000       | \$1,307,524,179    |
| 1184 | Chupacallos (Pleodendron macranthum) - Wherever found                               | E      | \$8,000       | \$1,307,532,179    |
| 1184 | Palo colorado (Ternstroemia luquillensis) - Wherever found                          | E      | \$8,000       | \$1,307,540,179    |
| 1184 | No Common Name (Gesneria pauciflora) - Wherever found                               | Т      | \$8,000       | \$1,307,548,179    |
| 1184 | Fern, Elfin tree (Cyathea dryopteroides) - Wherever found                           | E      | \$8,000       | \$1,307,556,179    |
| 1184 | No Common Name (Thelypteris inabonensis) - Wherever found                           | E      | \$8,000       | \$1,307,564,179    |
| 1185 | Manzanita, Santa Rosa Island (Arctostaphylos confertiflora) - Wherever found        | E      | \$7,986       | \$1,307,572,165    |
| 1186 | No Common Name (Gouania hillebrandii) - Wherever found                              | Е      | \$7,964       | \$1,307,580,129    |
| 1186 | Alani (Melicope munroi) - Wherever found  | E      | \$7,964       | \$1,307,588,093    |
| 1186 | Ha`iwale (Cyrtandra munroi) - Wherever found  | E      | \$7,964       | \$1,307,596,057    |
| 1186 | No Common Name (Schiedea lydgatei) - Wherever found                                 | E      | \$7,964       | \$1,307,604,021    |
| 1187 | No Common Name (Hesperomannia arbuscula) - Wherever found                           | Е      | \$7,950       | \$1,307,611,971    |
| 1188 | Makou (Peucedanum sandwicense) - Wherever found                                     | Т      | \$7,901       | \$1,307,619,872    |
| 1189 | Turtle, Alabama red-bellied (Pseudemys alabamensis) - Wherever found                | Е      | \$7,870       | \$1,307,627,742    |
| 1190 | Woodrat, riparian (=San Joaquin Valley) (Neotoma fuscipes riparia) - Wherever found | Е      | \$7,866       | \$1,307,635,608    |
| 1191 | Snail, Newcomb's (Erinna newcombi) - Wherever found                                 | Т      | \$7,848       | \$1,307,643,456    |
| 1192 | Chub, Yaqui (Gila purpurea) - Wherever found  | E      | \$7,800       | \$1,307,651,256    |
| 1193 | Dudleya, Santa Monica Mountains (Dudleya cymosa ssp. ovatifolia) - Wherever found   | Т      | \$7,787       | \$1,307,659,043    |
| 1194 | Daisy, Parish's (Erigeron parishii) - Wherever found                                | Т      | \$7,759       | \$1,307,666,802    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 1195 | Pilo (Kadua laxiflora) - Wherever found                                  | E      | \$7,738       | \$1,307,674,540    |
| 1196 | Darter, diamond (Crystallaria cincotta) - Wherever found                 | E      | \$7,735       | \$1,307,682,275    |
| 1197 | Koki`o ke`oke`o (Hibiscus arnottianus ssp. immaculatus) - Wherever found | Е      | \$7,677       | \$1,307,689,952    |
| 1198 | `Anaunau (Lepidium arbuscula) - Wherever found                           | E      | \$7,636       | \$1,307,697,588    |
| 1198 | Alani (Melicope saint-johnii) - Wherever found                           | E      | \$7,636       | \$1,307,705,224    |
| 1199 | Hala pepe (Pleomele forbesii) - Wherever found                           | E      | \$7,635       | \$1,307,712,859    |
| 1200 | Springsnail, Roswell (Pyrgulopsis roswellensis) - Wherever found         |        | \$7,600       | \$1,307,720,459    |
| 1200 | Cavesnail, Tumbling Creek (Antrobia culveri) - Wherever found            | E      | \$7,600       | \$1,307,728,059    |
| 1200 | Rosemary, short-leaved (Conradina brevifolia) - Wherever found           | E      | \$7,600       | \$1,307,735,659    |
| 1201 | Nehe (Lipochaeta lobata var. leptophylla) - Wherever found               | E      | \$7,550       | \$1,307,743,209    |
| 1202 | Boa, Mona (Epicrates monensis monensis) - Wherever found                 | Т      | \$7,500       | \$1,307,750,709    |
| 1202 | Guajon (Eleutherodactylus cooki) - Wherever found                        | Т      | \$7,500       | \$1,307,758,209    |
| 1202 | Blindcat, Mexican (catfish) (Prietella phreatophila) - Wherever found    | Е      | \$7,500       | \$1,307,765,709    |
| 1202 | Prickly-ash, St. Thomas (Zanthoxylum thomasianum) - Wherever found       | E      | \$7,500       | \$1,307,773,209    |
| 1202 | No Common Name (Auerodendron pauciflorum) - Wherever found               | E      | \$7,500       | \$1,307,780,709    |
| 1202 | No Common Name (Tectaria estremerana) - Wherever found                   | E      | \$7,500       | \$1,307,788,209    |
| 1203 | Papala (Charpentiera densiflora) - Wherever found                        | E      | \$7,343       | \$1,307,795,552    |
| 1203 | Alani (Melicope christophersenii) - Wherever found                       | E      | \$7,343       | \$1,307,802,895    |
| 1203 | No Common Name (Doryopteris angelica) - Wherever found                   | E      | \$7,343       | \$1,307,810,238    |
| 1204 | Haha (Cyanea glabra) - Wherever found                                    | E      | \$7,305       | \$1,307,817,543    |
| 1205 | No Common Name (Asplenium unisorum) - Wherever found                     | E      | \$7,290       | \$1,307,824,833    |
| 1206 | Damselfly, Rota blue (Ischnura luta) - Wherever found                    | E      | \$7,278       | \$1,307,832,111    |
| 1207 | found  | Е      | \$7,250       | \$1,307,839,361    |
| 1208 | ,  | Е      | \$7,200       | \$1,307,846,561    |
| 1208 | Water-willow, Cooley's (Justicia cooleyi) - Wherever found               | Е      | \$7,200       | \$1,307,853,761    |
| 1209 | Kamakahala (Labordia pumila) - Wherever found                            | Е      | \$7,193       | \$1,307,860,954    |
| 1209 | Na`ena`e (Dubautia plantaginea magnifolia) - Wherever found              | Е      | \$7,193       | \$1,307,868,147    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 1210 | Sandalwood, Lanai (=`iliahi) (Santalum haleakalae var. lanaiense) - Wherever found | E      | \$7,177       | \$1,307,875,324    |
| 1211 | Milk-vetch, triple-ribbed (Astragalus tricarinatus) - Wherever found               | E      | \$7,160       | \$1,307,882,484    |
| 1212 | Springsnail, Phantom (Pyrgulopsis texana) - Wherever found                         | E      | \$7,100       | \$1,307,889,584    |
| 1212 | Leather flower, Alabama (Clematis socialis) - Wherever found                       | E      | \$7,100       | \$1,307,896,684    |
| 1212 | Ladies-tresses, Canelo Hills (Spiranthes delitescens) - Wherever found             | E      | \$7,100       | \$1,307,903,784    |
| 1213 | Skipper, Pawnee montane (Hesperia leonardus montana) - Wherever found              | Т      | \$7,071       | \$1,307,910,855    |
| 1214 | Remya, Maui (Remya mauiensis) - Wherever found                                     | E      | \$7,050       | \$1,307,917,905    |
| 1215 | No Common Name (Schiedea verticillata) - Wherever found                            | E      | \$7,036       | \$1,307,924,941    |
| 1215 | No Common Name (Phyllostegia glabra var. lanaiensis) - Wherever found              | E      | \$7,036       | \$1,307,931,977    |
| 1216 | Cat, tiger (Leopardus (=Felis) tigrinus) - Wherever found                          | E      | \$7,000       | \$1,307,938,977    |
| 1216 | Beetle, Kretschmarr Cave mold (Texamaurops reddelli) - Wherever found              | E      | \$7,000       | \$1,307,945,977    |
| 1216 | Meshweaver, Robber Baron Cave (Cicurina baronia) - Wherever found                  | E      | \$7,000       | \$1,307,952,977    |
| 1216 | Holly, Cook's (Ilex cookii) - Wherever found                                       | Е      | \$7,000       | \$1,307,959,977    |
| 1216 | No Common Name (Ternstroemia subsessilis) - Wherever found                         | E      | \$7,000       | \$1,307,966,977    |
| 1216 | No Common Name (Elaphoglossum serpens) - Wherever found                            | E      | \$7,000       | \$1,307,973,977    |
| 1216 | No Common Name (Thelypteris verecunda) - Wherever found                            | E      | \$7,000       | \$1,307,980,977    |
| 1217 | Haha (Cyanea gibsonii) - Wherever found  | Е      | \$6,978       | \$1,307,987,955    |
| 1218 | Pilo kea lau li`i (Platydesma rostrata) - Wherever found                           | Е      | \$6,973       | \$1,307,994,928    |
| 1218 | Lo`ulu (Pritchardia napaliensis) - Wherever found                                  | Е      | \$6,973       | \$1,308,001,901    |
| 1218 | Kopiko (Psychotria hobdyi) - Wherever found  | Е      | \$6,973       | \$1,308,008,874    |
| 1219 | Ko`oko`olau (Bidens conjuncta) - Wherever found                                    | E      | \$6,950       | \$1,308,015,824    |
| 1219 | No Common Name (Schiedea sarmentosa) - Wherever found                              | E      | \$6,950       | \$1,308,022,774    |
| 1219 | No Common Name (Schiedea salicaria) - Wherever found                               | E      | \$6,950       | \$1,308,029,724    |
| 1219 | Ha`iwale (Cyrtandra oxybapha) - Wherever found                                     | E      | \$6,950       | \$1,308,036,674    |
| 1219 | Kolea (Myrsine vaccinioides) - Wherever found                                      | E      | \$6,950       | \$1,308,043,624    |
| 1219 | Nohoanu (Geranium hillebrandii) - Wherever found                                   | E      | \$6,950       | \$1,308,050,574    |
| 1219 | Hala pepe (Pleomele fernaldii) - Wherever found                                    | E      | \$6,950       | \$1,308,057,524    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 1220 | No Common Name (Festuca molokaiensis) - Wherever found   | Е      | \$6,910       | \$1,308,064,434    |
| 1221 | Whitlow-wort, papery (Paronychia chartacea) - Wherever found   | Т      | \$6,900       | \$1,308,071,334    |
| 1222 | Amphianthus, little (Amphianthus pusillus) - Wherever found  | Т      | \$6,850       | \$1,308,078,184    |
| 1222 | Barbara's Buttons, Mohr's (Marshallia mohrii) - Wherever found   | Т      | \$6,850       | \$1,308,085,034    |
| 1223 | `Akoko (Euphorbia kuwaleana) - Wherever found  | E      | \$6,849       | \$1,308,091,883    |
| 1224 | Fox, Santa Rosa Island (Urocyon littoralis santarosae) - wherever found  | Е      | \$6,839       | \$1,308,098,722    |
| 1224 | Butterfly, lotis blue (Lycaeides argyrognomon lotis) - Wherever found  | Е      | \$6,839       | \$1,308,105,561    |
| 1225 | No Common Name (Cyperus pennatiformis) - Wherever found  | Е      | \$6,827       | \$1,308,112,388    |
| 1226 | Panicgrass, Carter's (Panicum fauriei var. carteri) - Wherever found   | Е      | \$6,815       | \$1,308,119,203    |
| 1227 | Blossom, tubercled (pearlymussel) (Epioblasma torulosa torulosa) - Wherever found; Except where listed as Experimental Populations | Е      | \$6,764       | \$1,308,125,967    |
| 1228 | Reedgrass, Hillegrand's (Calamagrostis hillebrandii) - Wherever found  | Е      | \$6,737       | \$1,308,132,704    |
| 1228 | Pauoa (Ctenitis squamigera) - Wherever found   | E      | \$6,737       | \$1,308,139,441    |
| 1229 | Crownscale, San Jacinto Valley (Atriplex coronata var. notatior) - Wherever found  | Е      | \$6,728       | \$1,308,146,169    |
| 1230 | Cactus, Brady pincushion (Pediocactus bradyi) - Wherever found   | E      | \$6,650       | \$1,308,152,819    |
| 1230 | Nehe (Melanthera kamolensis) - Wherever found  | Е      | \$6,650       | \$1,308,159,469    |
| 1231 | No Common Name (Gonocalyx concolor) - Wherever found   | E      | \$6,600       | \$1,308,166,069    |
| 1232 | Haha (Cyanea mceldowneyi) - Wherever found   | E      | \$6,575       | \$1,308,172,644    |
| 1233 | Liliwai (Acaena exigua) - Wherever found   | E      | \$6,516       | \$1,308,179,160    |
| 1234 | Milk-vetch, heliotrope (Astragalus montii) - Wherever found  | Т      | \$6,503       | \$1,308,185,663    |
| 1235 | Anole, Culebra Island giant (Anolis roosevelti) - Wherever found   | Е      | \$6,500       | \$1,308,192,163    |
| 1235 | Shiner, palezone (Notropis albizonatus) - Wherever found   | E      | \$6,500       | \$1,308,198,663    |
| 1235 | No Common Name (Agave eggersiana) - Wherever found   | E      | \$6,500       | \$1,308,205,163    |
| 1235 | No Common Name (Mitracarpus maxwelliae) - Wherever found   | Е      | \$6,500       | \$1,308,211,663    |
| 1235 | Peperomia, Wheeler's (Peperomia wheeleri) - Wherever found   | Е      | \$6,500       | \$1,308,218,163    |
| 1235 | No Common Name (Adiantum vivesii) - Wherever found   | E      | \$6,500       | \$1,308,224,663    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 1235 | No Common Name (Polystichum calderonense) - Wherever found   | Е      | \$6,500       | \$1,308,231,163    |
| 1235 | No Common Name (Thelypteris yaucoensis) - Wherever found   | E      | \$6,500       | \$1,308,237,663    |
| 1236 | Bladderpod, Spring Creek (Lesquerella perforata) - Wherever found  | E      | \$6,490       | \$1,308,244,153    |
| 1237 | Aupaka (Isodendrion longifolium) - Wherever found  | Т      | \$6,472       | \$1,308,250,625    |
| 1237 | Alani (Melicope pallida) - Wherever found  | E      | \$6,472       | \$1,308,257,097    |
| 1237 | No Common Name (Lysimachia filifolia) - Wherever found   | E      | \$6,472       | \$1,308,263,569    |
| 1238 | Oxytheca, Cushenbury (Oxytheca parishii var. goodmaniana) - Wherever found   | E      | \$6,443       | \$1,308,270,012    |
| 1239 | `Oha wai (Clermontia oblongifolia ssp. mauiensis) - Wherever found   | E      | \$6,417       | \$1,308,276,429    |
| 1240 | Polygonum, Scotts Valley (Polygonum hickmanii) - Wherever found  | E      | \$6,411       | \$1,308,282,840    |
| 1241 | No Common Name (Keysseria (=Lagenifera) erici) - Wherever found  | E      | \$6,406       | \$1,308,289,246    |
| 1241 | No Common Name (Keysseria (=Lagenifera) helenae) - Wherever found  | E      | \$6,406       | \$1,308,295,652    |
| 1242 | Springsnail, San Bernardino (Pyrgulopsis bernardina) - Wherever found  | Т      | \$6,400       | \$1,308,302,052    |
| 1243 | Ma`oli`oli (Schiedea kealiae) - Wherever found   | Е      | \$6,359       | \$1,308,308,411    |
| 1244 | Walnut (=Nogal), West Indian (Juglans jamaicensis) - Wherever found  | E      | \$6,348       | \$1,308,314,759    |
| 1245 | Haha (Cyanea mauiensis) - Wherever found   | E      | \$6,322       | \$1,308,321,081    |
| 1246 | Ha`iwale (Cyrtandra subumbellata) - Wherever found   | E      | \$6,315       | \$1,308,327,396    |
| 1246 | Ha`iwale (Cyrtandra crenata) - Wherever found  | E      | \$6,315       | \$1,308,333,711    |
| 1246 | No Common Name (Trematolobelia singularis) - Wherever found  | E      | \$6,315       | \$1,308,340,026    |
| 1247 | Snail, Langford's tree (Partula langfordi) - Wherever found  | E      | \$6,314       | \$1,308,346,340    |
| 1248 | Moccasinshell, Coosa (Medionidus parvulus) - Wherever found  | E      | \$6,300       | \$1,308,352,640    |
| 1249 | No Common Name (Bidens hillebrandiana ssp. hillebrandiana) - Wherever found  | E      | \$6,250       | \$1,308,358,890    |
| 1250 | Milk-vetch, Cushenbury (Astragalus albens) - Wherever found  | E      | \$6,243       | \$1,308,365,133    |
| 1251 | Aumakua, Palapalai (Dryopteris crinalis var. podosorus) - Wherever found   | Е      | \$6,223       | \$1,308,371,356    |
| 1252 | Snake, copperbelly water (Nerodia erythrogaster neglecta) - Indiana north of 40 degrees north latitude, Michigan, Ohio | Т      | \$6,192       | \$1,308,377,548    |
| 1253 | Phacelia, clay (Phacelia argillacea) - Wherever found  | E      | \$6,180       | \$1,308,383,728    |
| 1254 | Koholapehu (Dubautia latifolia) - Wherever found   | E      | \$6,179       | \$1,308,389,907    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 1254 | No Common Name (Hesperomannia lydgatei) - Wherever found                         | Е      | \$6,179       | \$1,308,396,086    |
| 1254 | Hibiscus, Clay's (Hibiscus clayi) - Wherever found                               | E      | \$6,179       | \$1,308,402,265    |
| 1254 | Bluegrass, Mann's (Poa mannii) - Wherever found                                  | E      | \$6,179       | \$1,308,408,444    |
| 1254 | Ma`oli`oli (Schiedea apokremnos) - Wherever found                                | E      | \$6,179       | \$1,308,414,623    |
| 1254 | No Common Name (Schiedea helleri) - Wherever found                               | E      | \$6,179       | \$1,308,420,802    |
| 1254 | No Common Name (Schiedea spergulina var. leiopoda) - Wherever found              | Е      | \$6,179       | \$1,308,426,981    |
| 1254 | Laulihilihi (Schiedea stellarioides) - Wherever found                            | E      | \$6,179       | \$1,308,433,160    |
| 1254 | Na`ena`e (Dubautia imbricata imbricata) - Wherever found                         | E      | \$6,179       | \$1,308,439,339    |
| 1254 | Kamakahala (Labordia helleri) - Wherever found                                   | E      | \$6,179       | \$1,308,445,518    |
| 1254 | Lo`ulu, (=Na`ena`e) (Pritchardia hardyi) - Wherever found                        | E      | \$6,179       | \$1,308,451,697    |
| 1254 | No Common Name (Phyllostegia renovans) - Wherever found                          | Е      | \$6,179       | \$1,308,457,876    |
| 1254 | No Common Name (Doryopteris takeuchii) - Wherever found                          | Е      | \$6,179       | \$1,308,464,055    |
| 1255 | No Common Name (Cyperus fauriei) - Wherever found                                | E      | \$6,163       | \$1,308,470,218    |
| 1256 | Skink, Slevin's (Emoia slevini) - Wherever found                                 | E      | \$6,145       | \$1,308,476,363    |
| 1257 | Catspaw, white (pearlymussel) (Epioblasma obliquata perobliqua) - Wherever found | Е      | \$6,110       | \$1,308,482,473    |
| 1258 | Shiner, beautiful (Cyprinella formosa) - Wherever found                          | Т      | \$6,100       | \$1,308,488,573    |
| 1259 | Fly, Hawaiian picture-wing (Drosophila obatai) - Wherever found                  | Е      | \$6,026       | \$1,308,494,599    |
| 1260 | Lo`ulu (Pritchardia remota) - Wherever found                                     | E      | \$6,022       | \$1,308,500,621    |
| 1260 | Alani (Melicope makahae) - Wherever found  | E      | \$6,022       | \$1,308,506,643    |
| 1260 | Ha`iwale (Cyrtandra sessilis) - Wherever found                                   | E      | \$6,022       | \$1,308,512,665    |
| 1260 | Hulumoa (Korthalsella degeneri) - Wherever found                                 | E      | \$6,022       | \$1,308,518,687    |
| 1260 | Ko`oko`olau (Bidens amplectens) - Wherever found                                 | E      | \$6,022       | \$1,308,524,709    |
| 1261 | Crow, white-necked (Corvus leucognaphalus) - Wherever found                      | Е      | \$6,000       | \$1,308,530,709    |
| 1261 | Gambusia, Big Bend (Gambusia gaigei) - Wherever found                            | E      | \$6,000       | \$1,308,536,709    |
| 1261 | Darter, watercress (Etheostoma nuchale) - Wherever found                         | E      | \$6,000       | \$1,308,542,709    |
| 1261 | No Common Name (Calyptranthes thomasiana) - Wherever found                       | Е      | \$6,000       | \$1,308,548,709    |
| 1261 | No Common Name (Leptocereus grantianus) - Wherever found                         | Е      | \$6,000       | \$1,308,554,709    |
| 1261 | Thoroughwort, Cape Sable (Chromolaena frustrata) - Wherever found                | Е      | \$6,000       | \$1,308,560,709    |
| 1262 | Springsnail, Chupadera (Pyrgulopsis chupaderae) - Wherever found                 | Е      | \$5,980       | \$1,308,566,689    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 1263 | Snail, Stock Island tree (Orthalicus reses (not incl. nesodryas)) - Wherever found  | Т      | \$5,910       | \$1,308,572,599    |
| 1264 | Bluecurls, Hidden Lake (Trichostema austromontanum ssp. compactum) - Wherever found   | Т      | \$5,800       | \$1,308,578,399    |
| 1265 | Nehe (Lipochaeta micrantha) - Wherever found  | E      | \$5,745       | \$1,308,584,144    |
| 1266 | `Akoko (Euphorbia deppeana) - Wherever found  | E      | \$5,528       | \$1,308,589,672    |
| 1266 | No Common Name (Asplenium dielfalcatum) - Wherever found  | E      | \$5,528       | \$1,308,595,200    |
| 1267 | Seablite, California (Suaeda californica) -   | E      | \$5,494       | \$1,308,600,694    |
| 1268 | No Common Name (Kadua stjohnii) - Wherever found  | E      | \$5,436       | \$1,308,606,130    |
| 1269 | 'Akoko (Euphorbia halemanui) - Wherever found   | E      | \$5,392       | \$1,308,611,522    |
| 1269 | 'Awiwi (Kadua cookiana) - Wherever found  | E      | \$5,392       | \$1,308,616,914    |
| 1269 | Ha`iwale (Cyrtandra limahuliensis) - Wherever found   | Т      | \$5,392       | \$1,308,622,306    |
| 1270 | Nehe (Lipochaeta waimeaensis) - Wherever found  | E      | \$5,374       | \$1,308,627,680    |
| 1271 | Bean, sea (Mucuna sloanei persericea) - Wherever found  | E      | \$5,362       | \$1,308,633,042    |
| 1272 | Clubshell, ovate (Pleurobema perovatum) - Wherever found  | Е      | \$5,354       | \$1,308,638,396    |
| 1273 | Lessingia, San Francisco (Lessingia germanorum (=L.g. var. germanorum)) - Wherever found  | Е      | \$5,300       | \$1,308,643,696    |
| 1274 | Pigtoe, rough (Pleurobema plenum) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))        | EXPN   | \$5,260       | \$1,308,648,956    |
| 1275 | `Akoko (Euphorbia remyi var. kauaiensis) - Wherever found   | E      | \$5,235       | \$1,308,654,191    |
| 1276 | Bellflower, Brooksville (Campanula robinsiae) - Wherever found  | E      | \$5,200       | \$1,308,659,391    |
| 1276 | Spurge, deltoid (Chamaesyce deltoidea ssp. deltoidea) - Wherever found  | Е      | \$5,200       | \$1,308,664,591    |
| 1277 | No Common Name (Schiedea spergulina var. spergulina) - Wherever found   | Т      | \$5,101       | \$1,308,669,692    |
| 1278 | Sunfish, spring pygmy (Elassoma alabamae) - Wherever found  | Т      | \$5,100       | \$1,308,674,792    |
| 1279 | Fly, Hawaiian picture-wing (Drosophila aglaia) - Wherever found   | E      | \$5,012       | \$1,308,679,804    |
| 1280 | Damselfly, flying earwig Hawaiian (Megalagrion nesiotes) - Wherever found   | E      | \$5,009       | \$1,308,684,813    |
| 1281 | Paudedo (Hedyotis megalantha) - Wherever found  | E      | \$5,005       | \$1,308,689,818    |
| 1282 | Madtom, yellowfin (Noturus flavipinnis) - U.S.A. (TN, VA-specified portions of the Holston River and watershed; see 17.84(e)(1)(i)) | EXPN   | \$5,000       | \$1,308,694,818    |
| 1282 | Snail, noonday (Mesodon clarki nantahala) - Wherever found  | Т      | \$5,000       | \$1,308,699,818    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 1282 | Harvestman, Cokendolpher Cave (Texella cokendolpheri) - Wherever found  | Е      | \$5,000       | \$1,308,704,818    |
| 1282 | Mint, Garrett's (Dicerandra christmanii) - Wherever found   | E      | \$5,000       | \$1,308,709,818    |
| 1282 | ,   | E      | \$5,000       | \$1,308,714,818    |
| 1283 | Wherever found  | Т      | \$4,992       | \$1,308,719,810    |
| 1284 | Blossom, turgid (pearlymussel) (Epioblasma turgidula) -<br>AL; Free-Flowing Reach of the Tennessee River below the<br>Wilson Dam, Colbert and Lauderdale Counties, AL | EXPN   | \$4,960       | \$1,308,724,770    |
| 1285 | Wahane (Pritchardia aylmer-robinsonii) - Wherever found   | E      | \$4,944       | \$1,308,729,714    |
| 1286 | Ridge-cress, Barneby (Lepidium barnebyanum) - Wherever found  | Е      | \$4,871       | \$1,308,734,585    |
| 1287 | Nehe (Lipochaeta fauriei) - Wherever found  | Е      | \$4,864       | \$1,308,739,449    |
| 1288 | Pigtoe, finerayed (Fusconaia cuneolus) - AL: Free-Flowing<br>Reach of the Tennessee River below the Wilson Dam,<br>Colbert and Lauderdale Counties, AL                | EXPN   | \$4,860       | \$1,308,744,309    |
| 1288 | Pigtoe, shiny (Fusconaia cor) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL                               | EXPN   | \$4,860       | \$1,308,749,169    |
| 1289 | Flannelbush, Pine Hill (Fremontodendron californicum ssp. decumbens) - Wherever found   | Е      | \$4,850       | \$1,308,754,019    |
| 1290 | Pearlymussel, birdwing (Lemiox rimosus) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert, and Lauderdale Counties, AL                    | EXPN   | \$4,780       | \$1,308,758,799    |
| 1291 | Riversnail, Anthony's (Athearnia anthonyi) - Wherever found; Except where listed as Experimental Populations  | Е      | \$4,700       | \$1,308,763,499    |
| 1291 | found   | E      | \$4,700       | \$1,308,768,199    |
| 1291 | Wherever found ` ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' ' '  | Е      | \$4,700       | \$1,308,772,899    |
| 1291 | Monkey-flower, Michigan (Mimulus michiganensis) - Wherever found  | Е      | \$4,700       | \$1,308,777,599    |
| 1292 | found   | Е      | \$4,630       | \$1,308,782,229    |
| 1293 | Cliffrose, Arizona (Purshia (=Cowania) subintegra) - Wherever found   | Е      | \$4,600       | \$1,308,786,829    |
| 1294 | Clarkia, Pismo (Clarkia speciosa ssp. immaculata) - Wherever found  | Е      | \$4,594       | \$1,308,791,423    |
| 1295 | Sea turtle, loggerhead (Caretta caretta) - South Pacific Ocean DPS  | Е      | \$4,537       | \$1,308,795,960    |
| 1296 | Puma (=cougar), eastern (Puma (=Felis) concolor couguar) - Wherever found   | E      | \$4,413       | \$1,308,800,373    |
| 1297 | 1 / 7 ( 1 /   | E      | \$4,400       | \$1,308,804,773    |
| 1297 | Cactus, Florida semaphore (Consolea corallicola) - Wherever found   | Е      | \$4,400       | \$1,308,809,173    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 1298 | Nukupuu, Maui (Hemignathus affinis) - Wherever found                                  | Е      | \$4,390       | \$1,308,813,563    |
| 1299 | Darter, rush (Etheostoma phytophilum) - Wherever found                                | Е      | \$4,284       | \$1,308,817,847    |
| 1300 | Thrush, Molokai (Myadestes lanaiensis rutha) - Wherever found                         | Е      | \$4,248       | \$1,308,822,095    |
| 1301 | Fly, Hawaiian picture-wing (Drosophila digressa) - Wherever found                     | E      | \$4,197       | \$1,308,826,292    |
| 1301 | Fly, Hawaiian picture-wing (Drosophila differens) - Wherever found                    | E      | \$4,197       | \$1,308,830,489    |
| 1302 | Gazelle, dama (Gazella dama) - Wherever found   | E      | \$4,100       | \$1,308,834,589    |
| 1302 | Buttercup, autumn (Ranunculus aestivalis (=acriformis)) - Wherever found              | E      | \$4,100       | \$1,308,838,689    |
| 1302 | Sand-verbena, large-fruited (Abronia macrocarpa) - Wherever found                     | Е      | \$4,100       | \$1,308,842,789    |
| 1302 | Pitcher-plant, Alabama canebrake (Sarracenia rubra ssp. alabamensis) - Wherever found | E      | \$4,100       | \$1,308,846,889    |
| 1303 | Milk-vetch, coastal dunes (Astragalus tener var. titi) - Wherever found               | E      | \$4,084       | \$1,308,850,973    |
| 1304 | Paintbrush, Tiburon (Castilleja affinis ssp. neglecta) - Wherever found               | E      | \$4,060       | \$1,308,855,033    |
| 1305 | Darter, vermilion (Etheostoma chermocki) - Wherever found                             | Е      | \$4,000       | \$1,308,859,033    |
| 1305 | Rocksnail, interrupted (=Georgia) (Leptoxis foremani) - Wherever found                | Е      | \$4,000       | \$1,308,863,033    |
| 1305 | Thistle, Suisun (Cirsium hydrophilum var. hydrophilum) - Wherever found               | E      | \$4,000       | \$1,308,867,033    |
| 1306 | Fly, Hawaiian picture-wing (Drosophila heteroneura) - Wherever found                  | E      | \$3,998       | \$1,308,871,031    |
| 1306 | Fly, Hawaiian picture-wing (Drosophila mulli) - Wherever found                        | Т      | \$3,998       | \$1,308,875,029    |
| 1306 | Picture-wing fly, Hawaiian (Drosophila neoclavisetae) - Wherever found                | Е      | \$3,998       | \$1,308,879,027    |
| 1306 | Fly, Hawaiian picture-wing (Drosophila tarphytrichia) - Wherever found                | Е      | \$3,998       | \$1,308,883,025    |
| 1306 | Fly, Hawaiian picture-wing (Drosophila hemipeza) - Wherever found                     | Е      | \$3,998       | \$1,308,887,023    |
| 1306 | Fly, Hawaiian picture-wing (Drosophila ochrobasis) - Wherever found                   | E      | \$3,998       | \$1,308,891,021    |
| 1307 | Clover, Monterey (Trifolium trichocalyx) - Wherever found                             | E      | \$3,994       | \$1,308,895,015    |
| 1308 | Shark, Scalloped Hammerhead (Sphyrna lewini) - Eastern Pacific DPS                    | E      | \$3,944       | \$1,308,898,959    |
| 1309 | Reed-mustard, clay (Schoenocrambe argillacea) - Wherever found                        | Т      | \$3,942       | \$1,308,902,901    |
| 1310 | Dudleya, Conejo (Dudleya abramsii ssp. parva) - Wherever found                        | Т      | \$3,894       | \$1,308,906,795    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 1310 | Dudleya, marcescent (Dudleya cymosa ssp. marcescens) - Wherever found  | Т      | \$3,894       | \$1,308,910,689    |
| 1310 | Cypress, Gowen (Cupressus goveniana ssp. goveniana) - Wherever found   | Т      | \$3,894       | \$1,308,914,583    |
| 1311 | Malacothrix, Santa Cruz Island (Malacothrix indecora) - Wherever found   | Е      | \$3,891       | \$1,308,918,474    |
| 1312 | Pawpaw, Rugel's (Deeringothamnus rugelii) - Wherever found   | Е      | \$3,800       | \$1,308,922,274    |
| 1313 | Babirusa (Babyrousa babyrussa) - Wherever found  | E      | \$3,650       | \$1,308,925,924    |
| 1313 | Dwarf-flax, Marin (Hesperolinon congestum) - Wherever found  | Т      | \$3,650       | \$1,308,929,574    |
| 1314 | Buckwheat, Ione (incl. Irish Hill) (Eriogonum apricum (incl. var. prostratum)) - Wherever found  | Е      | \$3,600       | \$1,308,933,174    |
| 1315 | Spineflower, Sonoma (Chorizanthe valida) - Wherever found  | Е      | \$3,550       | \$1,308,936,724    |
| 1316 | Spurge, Garber's (Chamaesyce garberi) - Wherever found   | Т      | \$3,515       | \$1,308,940,239    |
| 1317 | Harebells, Avon Park (Crotalaria avonensis) - Wherever found   | Е      | \$3,500       | \$1,308,943,739    |
| 1318 | Milk-vetch, Ventura Marsh (Astragalus pycnostachyus var. lanosissimus) - Wherever found  | Е      | \$3,461       | \$1,308,947,200    |
| 1319 | Dugong (Dugong dugon) - Wherever found   | E      | \$3,450       | \$1,308,950,650    |
| 1320 | Po`ouli (honeycreeper) (Melamprosops phaeosoma) - Wherever found   | Е      | \$3,444       | \$1,308,954,094    |
| 1321 | Bladderpod, Zapata (Lesquerella thamnophila) - Wherever found  | Е      | \$3,400       | \$1,308,957,494    |
| 1322 | Shiner, Cahaba (Notropis cahabae) - Wherever found   | E      | \$3,363       | \$1,308,960,857    |
| 1323 | Seal, Saimaa (Phoca hispida saimensis) - Wherever found  | E      | \$3,332       | \$1,308,964,189    |
| 1324 | Clarkia, Presidio (Clarkia franciscana) - Wherever found   | E      | \$3,320       | \$1,308,967,509    |
| 1325 | Clover, showy Indian (Trifolium amoenum) - Wherever found  | E      | \$3,300       | \$1,308,970,809    |
| 1326 | Blossom, green (pearlymussel) (Epioblasma torulosa gubernaculum) - Wherever found  | Е      | \$3,240       | \$1,308,974,049    |
| 1326 | Monkeyface, Appalachian (pearlymussel) (Quadrula sparsa) - USA (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1)) | EXPN   | \$3,240       | \$1,308,977,289    |
| 1327 | Oryx, Arabian (Oryx leucoryx) - Wherever found   | E      | \$3,200       | \$1,308,980,489    |
| 1327 | Lemurs (Lemuridae (incl. genera Lemur, Phaner,<br>Hapalemur, Lepilemur, Microcebus, Allocebus,<br>Cheirogaleus, Varecia)) - Wherever found       | Е      | \$3,200       | \$1,308,983,689    |
| 1328 | Manzanita, Presidio (Arctostaphylos hookeri var. ravenii) - Wherever found   | Е      | \$3,100       | \$1,308,986,789    |
| 1329 | Mucket, orangenacre (Lampsilis perovalis) - Wherever found   | Т      | \$3,046       | \$1,308,989,835    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 1330 | Marstonia, royal (snail) (Pyrgulopsis ogmorhaphe) - Wherever found   | Е      | \$3,000       | \$1,308,992,835    |
| 1330 | Amphipod, Illinois cave (Gammarus acherondytes) - Wherever found   | Е      | \$3,000       | \$1,308,995,835    |
| 1330 | Amphipod, Pecos (Gammarus pecos) - Wherever found  | E      | \$3,000       | \$1,308,998,835    |
| 1331 | Cactus, Nichol's Turk's head (Echinocactus horizonthalonius var. nicholii) - Wherever found  | Е      | \$2,900       | \$1,309,001,735    |
| 1331 | Polygala, tiny (Polygala smallii) - Wherever found   | E      | \$2,900       | \$1,309,004,635    |
| 1332 | Spider, spruce-fir moss (Microhexura montivaga) - Wherever found   | E      | \$2,800       | \$1,309,007,435    |
| 1333 | Amphipod, diminutive (Gammarus hyalleloides) - Wherever found  | E      | \$2,600       | \$1,309,010,035    |
| 1333 | Locoweed, Fassett's (Oxytropis campestris var. chartacea) - Wherever found   | Т      | \$2,600       | \$1,309,012,635    |
| 1333 | Bladderpod, white (Lesquerella pallida) - Wherever found   | E      | \$2,600       | \$1,309,015,235    |
| 1333 | Gladecress, Texas golden (Leavenworthia texana) - Wherever found   | Е      | \$2,600       | \$1,309,017,835    |
| 1334 | Minnow, Rio Grande Silvery (Hybognathus amarus) - Rio Grande, from Little Box Canyon to Amistad Dam                                | EXPN   | \$2,500       | \$1,309,020,335    |
| 1334 | Pearlymussel, birdwing (Lemiox rimosus) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1)) | EXPN   | \$2,500       | \$1,309,022,835    |
| 1334 | Tiger beetle, Ohlone (Cicindela ohlone) - Wherever found   | E      | \$2,500       | \$1,309,025,335    |
| 1334 | Clarkia, Springville (Clarkia springvillensis) - Wherever found  | Т      | \$2,500       | \$1,309,027,835    |
| 1334 | Leather flower, Morefield's (Clematis morefieldii) - Wherever found  | E      | \$2,500       | \$1,309,030,335    |
| 1335 | Chub, Sonora (Gila ditaenia) - Wherever found  | Т      | \$2,400       | \$1,309,032,735    |
| 1336 | Woodpecker, ivory-billed (Campephilus principalis) - Wherever found  | E      | \$2,347       | \$1,309,035,082    |
| 1337 | Roseroot, Leedy's (Rhodiola integrifolia ssp. leedyi) - Wherever found   | Т      | \$2,335       | \$1,309,037,417    |
| 1338 | Ceanothus, Vail Lake (Ceanothus ophiochilus) - Wherever found  | Т      | \$2,204       | \$1,309,039,621    |
| 1339 | Gladecress, Fleshy-fruit (Leavenworthia crassa) - Wherever found   | Е      | \$2,200       | \$1,309,041,821    |
| 1340 | Snail, tulotoma (Tulotoma magnifica) - Wherever found  | Т      | \$2,175       | \$1,309,043,996    |
| 1341 | Paintbrush, ash-grey (Castilleja cinerea) - Wherever found   | Т      | \$2,087       | \$1,309,046,083    |
| 1342 | Parrot, thick-billed (Rhynchopsitta pachyrhyncha) - Wherever found   | Е      | \$2,000       | \$1,309,048,083    |
| 1343 | Buckwheat, cushenbury (Eriogonum ovalifolium var. vineum) - Wherever found   | Е      | \$1,943       | \$1,309,050,026    |

| Rank | Species (50 CFR Part 17)   | Status | Species Total | Cumulatative Total |
|------|--|--------|---------------|--------------------|
| 1344 | Ivesia, Ash Meadows (Ivesia kingii var. eremica) - Wherever found  | Т      | \$1,914       | \$1,309,051,940    |
| 1345 | Woundfin (Plagopterus argentissimus) - Gila R. drainage, AZ, NM  | EXPN   | \$1,900       | \$1,309,053,840    |
| 1345 | Lupine, Nipomo Mesa (Lupinus nipomensis) - Wherever found  | E      | \$1,900       | \$1,309,055,740    |
| 1346 | Madtom, Scioto (Noturus trautmani) - Wherever found  | Е      | \$1,881       | \$1,309,057,621    |
| 1347 | Cactus, Lloyd's Mariposa (Echinomastus mariposensis) - Wherever found  | Т      | \$1,877       | \$1,309,059,498    |
| 1348 | Salamander, Red Hills (Phaeognathus hubrichti) - Wherever found  | Т      | \$1,839       | \$1,309,061,337    |
| 1349 | Gambusia, San Marcos (Gambusia georgei) - Wherever found   | E      | \$1,800       | \$1,309,063,137    |
| 1349 | Snail, Virginia fringed mountain (Polygyriscus virginianus) - Wherever found   | E      | \$1,800       | \$1,309,064,937    |
| 1350 | Taraxacum, California (Taraxacum californicum) -<br>Wherever found   | E      | \$1,767       | \$1,309,066,704    |
| 1351 | Jewelflower, Tiburon (Streptanthus niger) - Wherever found   | E      | \$1,750       | \$1,309,068,454    |
| 1352 | Cactus, Chisos Mountain hedgehog (Echinocereus chisoensis var. chisoensis) - Wherever found  | Т      | \$1,736       | \$1,309,070,190    |
| 1353 | Cat, leopard (Prionailurus (=Felis) bengalensis bengalensis) - Wherever found  | E      | \$1,730       | \$1,309,071,920    |
| 1354 | Purple cat's paw (=purple cat's paw pearlymussel) (Epioblasma obliquata obliquata) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL | EXPN   | \$1,720       | \$1,309,073,640    |
| 1355 | Prickly-apple, fragrant (Cereus eriophorus var. fragrans) - Wherever found   | E      | \$1,700       | \$1,309,075,340    |
| 1355 | Pawpaw, beautiful (Deeringothamnus pulchellus) - Wherever found  | E      | \$1,700       | \$1,309,077,040    |
| 1356 | Fern, Alabama streak-sorus (Thelypteris pilosa var. alabamensis) - Wherever found  | Т      | \$1,600       | \$1,309,078,640    |
| 1357 | 'O'u (honeycreeper) (Psittirostra psittacea) - Wherever found  | E      | \$1,507       | \$1,309,080,147    |
| 1358 | Madtom, pygmy (Noturus stanauli) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.84(t)(1)(i))   | EXPN   | \$1,500       | \$1,309,081,647    |
| 1358 | Springsnail, Socorro (Pyrgulopsis neomexicana) - Wherever found  | E      | \$1,500       | \$1,309,083,147    |
| 1358 | Cory cactus, bunched (Coryphantha ramillosa) - Wherever found  | Т      | \$1,500       | \$1,309,084,647    |
| 1358 | Water-plantain, Kral's (Sagittaria secundifolia) - Wherever found  | Т      | \$1,500       | \$1,309,086,147    |
| 1359 | Thornmint, San Mateo (Acanthomintha obovata ssp. duttonii) - Wherever found  | Е      | \$1,413       | \$1,309,087,560    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 1360 | Warbler (=wood), Bachman's (Vermivora bachmanii) - Wherever found                           | E      | \$1,397       | \$1,309,088,957    |
| 1361 | Poppy-mallow, Texas (Callirhoe scabriuscula) - Wherever found                               | E      | \$1,300       | \$1,309,090,257    |
| 1362 | Curlew, Eskimo (Numenius borealis) - Wherever found   | E      | \$1,294       | \$1,309,091,551    |
| 1363 | Grass, Solano (Tuctoria mucronata) - Wherever found   | E      | \$1,200       | \$1,309,092,751    |
| 1363 | Manzanita, Morro (Arctostaphylos morroensis) - Wherever found                               | Т      | \$1,200       | \$1,309,093,951    |
| 1364 | Dace, Kendall Warm Springs (Rhinichthys osculus thermalis) - Wherever found                 | E      | \$1,131       | \$1,309,095,082    |
| 1365 | Wild-buckwheat, southern mountain (Eriogonum kennedyi var. austromontanum) - Wherever found | Т      | \$1,103       | \$1,309,096,185    |
| 1366 | Cactus, Sneed pincushion (Coryphantha sneedii var. sneedii) - Wherever found                | E      | \$1,100       | \$1,309,097,285    |
| 1367 | Pigtoe, heavy (Pleurobema taitianum) - Wherever found                                       | E      | \$1,016       | \$1,309,098,301    |
| 1368 | Akialoa, Kauai (honeycreeper) (Akialoa stejnegeri) - Wherever found                         | E      | \$1,004       | \$1,309,099,305    |
| 1369 | Gambusia, Clear Creek (Gambusia heterochir) - Wherever found                                | E      | \$1,000       | \$1,309,100,305    |
| 1369 | Beetle, Hungerford's crawling water (Brychius hungerfordi) - Wherever found                 | E      | \$1,000       | \$1,309,101,305    |
| 1370 | 'O'o, Kauai (honeyeater) (Moho braccatus) - Wherever found                                  | E      | \$990         | \$1,309,102,295    |
| 1371 | Darter, bayou (Etheostoma rubrum) - Wherever found  | Т      | \$963         | \$1,309,103,258    |
| 1372 | Ragwort, San Francisco Peaks (Packera franciscana) - Wherever found                         | Т      | \$915         | \$1,309,104,173    |
| 1373 | Sandwort, Bear Valley (Arenaria ursina) - Wherever found                                    | Т      | \$803         | \$1,309,104,976    |
| 1374 | Thrush, large Kauai (=kamao) (Myadestes myadestinus) - Wherever found                       | E      | \$801         | \$1,309,105,777    |
| 1374 | Creeper, Molokai (Paroreomyza flammea) - Wherever found                                     | E      | \$801         | \$1,309,106,578    |
| 1375 | Snowbells, Texas (Styrax texanus) - Wherever found  | Е      | \$800         | \$1,309,107,378    |
| 1376 | White-eye, bridled (Zosterops conspicillatus conspicillatus) - Wherever found               | E      | \$761         | \$1,309,108,139    |
| 1377 | Prickly-apple, aboriginal (Harrisia (=Cereus) aboriginum (=gracilis)) - Wherever found      | E      | \$700         | \$1,309,108,839    |
| 1378 | Tamarin, cotton-top (Saguinus oedipus) - Wherever found                                     | E      | \$600         | \$1,309,109,439    |
| 1378 | Snail, armored (Pyrgulopsis (=Marstonia) pachyta) - Wherever found                          | Е      | \$600         | \$1,309,110,039    |
| 1378 | Lioplax, cylindrical (snail) (Lioplax cyclostomaformis) - Wherever found                    | E      | \$600         | \$1,309,110,639    |
| 1378 | Campeloma, slender (Campeloma decampi) - Wherever found                                     | E      | \$600         | \$1,309,111,239    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 1378 | Crayfish, cave (Cambarus aculabrum) - Wherever found                              | E      | \$600         | \$1,309,111,839    |
| 1378 | Bladderpod, lyrate (Lesquerella lyrata) - Wherever found                          | Т      | \$600         | \$1,309,112,439    |
| 1378 | Oak, Hinckley (Quercus hinckleyi) - Wherever found                                | Т      | \$600         | \$1,309,113,039    |
| 1379 | ssp. bernardina) - Wherever found   | E      | \$583         | \$1,309,113,622    |
| 1379 | Mustard, slender-petaled (Thelypodium stenopetalum) - Wherever found              | Е      | \$583         | \$1,309,114,205    |
| 1380 | Salamander, desert slender (Batrachoseps aridus) - Wherever found                 | Е      | \$580         | \$1,309,114,785    |
| 1381 | Pigtoe, flat (Pleurobema marshalli) - Wherever found                              | E      | \$550         | \$1,309,115,335    |
| 1381 | 1 ,   | E      | \$550         | \$1,309,115,885    |
| 1381 | infantilis) - Wherever found  | Е      | \$550         | \$1,309,116,435    |
| 1381 | Mariposa lily, Tiburon (Calochortus tiburonensis) - Wherever found                | Т      | \$550         | \$1,309,116,985    |
| 1382 | found   | Е      | \$500         | \$1,309,117,485    |
| 1382 | Liveforever, Santa Barbara Island (Dudleya traskiae) - Wherever found             | Е      | \$500         | \$1,309,117,985    |
| 1382 | Pitaya, Davis' green (Echinocereus viridiflorus var. davisii) - Wherever found    | Е      | \$500         | \$1,309,118,485    |
| 1383 | Checker-mallow, pedate (Sidalcea pedata) - Wherever found                         | Е      | \$483         | \$1,309,118,968    |
| 1384 | Solenodon, Cuban (Solenodon cubanus) - Wherever found                             | E      | \$348         | \$1,309,119,316    |
| 1384 | Hutia, Cabrera's (Capromys angelcabrerai) - Wherever found                        | Е      | \$348         | \$1,309,119,664    |
| 1384 | Hutia, dwarf (Capromys nana) - Wherever found                                     | E      | \$348         | \$1,309,120,012    |
| 1384 | Hutia, large-eared (Capromys auritus) - Wherever found                            | E      | \$348         | \$1,309,120,360    |
| 1384 | Crocodile, Cuban (Crocodylus rhombifer) - Wherever found                          | Е      | \$348         | \$1,309,120,708    |
| 1384 | ,   | E      | \$348         | \$1,309,121,056    |
| 1385 | Kite, Cuba hook-billed (Chondrohierax uncinatus wilsonii) - Wherever found        | Е      | \$347         | \$1,309,121,403    |
| 1385 | Parrot, Bahaman or Cuban (Amazona leucocephala) - Wherever found                  | Е      | \$347         | \$1,309,121,750    |
| 1386 | Shark, Scalloped Hammerhead (Sphyrna lewini) - Indo-<br>West Pacific DPS          | Т      | \$335         | \$1,309,122,085    |
| 1387 | Serow (Naemorhedus (=Capricornis) sumatraensis) - Wherever found                  | Е      | \$320         | \$1,309,122,405    |
| 1388 | Shark, Scalloped Hammerhead (Sphyrna lewini) - Central and Southwest Atlantic DPS | Т      | \$316         | \$1,309,122,721    |
| 1389 | Lechwe, red (Kobus leche) - Wherever found  | Т      | \$300         | \$1,309,123,021    |
| 1389 | Mandrill (Mandrillus (=Papio) sphinx) - Wherever found                            | Е      | \$300         | \$1,309,123,321    |

| Rank | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|------|---|--------|---------------|--------------------|
| 1389 | Zebra, Grevy's (Equus grevyi) - Wherever found  | Т      | \$300         | \$1,309,123,621    |
| 1389 | Spineflower, Howell's (Chorizanthe howellii) - Wherever found   | Е      | \$300         | \$1,309,123,921    |
| 1389 | Reed-mustard, Barneby (Schoenocrambe barnebyi) - Wherever found   | Е      | \$300         | \$1,309,124,221    |
| 1389 | Spineflower, Ben Lomond (Chorizanthe pungens var. hartwegiana) - Wherever found   | Е      | \$300         | \$1,309,124,521    |
| 1389 | Spineflower, Scotts Valley (Chorizanthe robusta var. hartwegii) - Wherever found  | Е      | \$300         | \$1,309,124,821    |
| 1389 | Spineflower, Robust (Chorizanthe robusta var. robusta) - Wherever found   | E      | \$300         | \$1,309,125,121    |
| 1390 | Pigtoe, dark (Pleurobema furvum) - Wherever found   | E      | \$284         | \$1,309,125,405    |
| 1391 | Pearlymussel, Curtis (Epioblasma florentina curtisii) - Wherever found  | Е      | \$278         | \$1,309,125,683    |
| 1392 | Bat, little Mariana fruit (Pteropus tokudae) - Wherever found   | E      | \$254         | \$1,309,125,937    |
| 1393 | Pondweed, Little Aguja (=Creek) (Potamogeton clystocarpus) - Wherever found   | E      | \$250         | \$1,309,126,187    |
| 1394 | Macaw, military (Ara militaris) - Wherever found  | E      | \$240         | \$1,309,126,427    |
| 1395 | Monardella, willowy (Monardella viminea) - Wherever found   | E      | \$232         | \$1,309,126,659    |
| 1396 | Gazelle, slender-horned (Gazella leptoceros) - Wherever found   | Е      | \$200         | \$1,309,126,859    |
| 1396 | Margay (Leopardus (=Felis) wiedii) - Wherever found   | E      | \$200         | \$1,309,127,059    |
| 1396 | Bontebok (Damaliscus pygarus (=dorcas) dorcas) -<br>Wherever found  | E      | \$200         | \$1,309,127,259    |
| 1396 | Baboon, gelada (Theropithecus gelada) - Wherever found  | Т      | \$200         | \$1,309,127,459    |
| 1396 | Monkey, Diana (Cercopithecus diana) - Wherever found  | E      | \$200         | \$1,309,127,659    |
| 1396 | Zebra, Hartmann's mountain (Equus zebra hartmannae) - Wherever found  | Т      | \$200         | \$1,309,127,859    |
| 1396 | Addax (Addax nasomaculatus) - Wherever found  | E      | \$200         | \$1,309,128,059    |
| 1396 | Koala (Phascolarctos cinereus) - Australia  | Т      | \$200         | \$1,309,128,259    |
| 1396 | Cavefish, Alabama (Speoplatyrhinus poulsoni) - Wherever found   | Е      | \$200         | \$1,309,128,459    |
| 1396 | Blossom, yellow (pearlymussel) (Epioblasma florentina florentina) - Wherever found; Except where listed as Experimental Populations                                   | Е      | \$200         | \$1,309,128,659    |
| 1396 | Monkeyface, Cumberland (pearlymussel) (Quadrula intermedia) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL | EXPN   | \$200         | \$1,309,128,859    |
| 1397 | Rocksnail, plicate (Leptoxis plicata) - Wherever found  | E      | \$149         | \$1,309,129,008    |
| 1398 | Onion, Munz's (Allium munzii) - Wherever found  | E      | \$116         | \$1,309,129,124    |

| Rank  | Species (50 CFR Part 17)  | Status | Species Total | Cumulatative Total |
|-------|---|--------|---------------|--------------------|
| 1399  | Jaguarundi, Sinaloan (Herpailurus (=Felis) yagouaroundi tolteca) - Wherever found   | Е      | \$100         | \$1,309,129,224    |
| 1399  | Darter, duskytail (Etheostoma percnurum) - Tellico River, TN  | EXPN   | \$100         | \$1,309,129,324    |
| 1399  | Sawfish, Dwarf (Pristis clavata) - Wherever found   | E      | \$100         | \$1,309,129,424    |
| 1399  | Blossom, tubercled (pearlymussel) (Epioblasma torulosa torulosa) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL  | EXPN   | \$100         | \$1,309,129,524    |
| 1399  | Blossom, yellow (pearlymussel) (Epioblasma florentina florentina) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL | EXPN   | \$100         | \$1,309,129,624    |
| 1399  | Pearlshell, Alabama (Margaritifera marrianae) - Wherever found  | E      | \$100         | \$1,309,129,724    |
| 1399  | Tryonia, Phantom (Tryonia cheatumi) - Wherever found  | E      | \$100         | \$1,309,129,824    |
| 1399  | Hornsnail, rough (Pleurocera foremani) - Wherever found   | E      | \$100         | \$1,309,129,924    |
| 1399  | Crayfish, cave (Cambarus zophonastes) - Wherever found  | E      | \$100         | \$1,309,130,024    |
| 1399  | Milk-vetch, Clara Hunt's (Astragalus clarianus) - Wherever found  | E      | \$100         | \$1,309,130,124    |
| 1399  | Rock-cress, McDonald's (Arabis macdonaldiana) - Wherever found  | E      | \$100         | \$1,309,130,224    |
| 1399  | Wallflower, Ben Lomond (Erysimum teretifolium) - Wherever found   | E      | \$100         | \$1,309,130,324    |
| 1399  | Primrose, Maguire (Primula maguirei) - Wherever found   | Т      | \$100         | \$1,309,130,424    |
| 1399  | Penny-cress, Kneeland Prairie (Thlaspi californicum) - Wherever found   | Е      | \$100         | \$1,309,130,524    |
| 1399  | Fringepod, Santa Cruz Island (Thysanocarpus conchuliferus) - Wherever found   | Е      | \$100         | \$1,309,130,624    |
| 1400  | Crane, hooded (Grus monacha) - Wherever found   | Е      | \$80          | \$1,309,130,704    |
| 1401  | Crane, Japanese (Grus japonensis) - Wherever found  | Е      | \$20          | \$1,309,130,724    |
| Total |   |        |               | \$1,309,130,724    |