IDA REPORT - ASSIGNMENT 2

Name: Katinni Rahul

Roll No.: S20200010091

Question 1:

```
#Given Data
mean <- 5818
n <- 24
52 <- 7273809
mu <- 5800

# Test Statistic
t <- (mean - mu) / (S2 / n)^(1/2)

# DOF (degrees of freedom)
df <- (n - 1)

# Critical-t-value from table
tc <- abs(qt(0.1, df))

# Results
cat("Test Statistic = ", t)
cat("Degrees of Freedom = ", df)
cat("Critical-t-value = ", tc)

if[t < tc]{
    cat("Failed to Reject the null Hypothesis meaning the mean is less than 5,800 tons")
}else {
    cat("Rejected the null hypothesis that the mean is less than 5,800 tons")
}</pre>
```

Output:

Failed to Reject the null Hypothesis meaning the mean is less than 5,800 tons

Question 2:

```
#question 2
# Given data
tar = c(14.5, 14.2, 14.4, 14.3, 14.6)
# Sample size
n = length(tar)
# Sample mean
xbar = mean(tar)
mu = 14.0
# Sample standard deviation
sd = sd(tar)
df = n - 1
# Test statistic
t = (xbar - mu) / (sd/sqrt(n))
tc <- abs(qt(0.025, df))
# Report
cat("Sample mean (xbar): ", xbar)
cat("Claimed population mean (mu): ", mu)
cat("t-stat: ", t)
cat("Critical-t-value = ", tc)
if(t < tc){}
 cat("Failed to Reject null hypothesis that population mean is 14.0")
} else {
 cat("Reject null hypothesis that population mean is 14.0")
```

Output:

```
Reject null hypothesis that population mean is 14.0 >
```

Question 3b:

```
#Question 3b
# Paired t-test
before <- c(45, 73, 46, 124, 33, 57, 83, 34, 26, 17)
# Sample size
n = length(before)
# Degrees of freedom
df = n - 1
# After data
after <- c(36, 60, 44, 119, 35, 51, 77, 29, 24, 11)
t_test <- t.test(before, after, paired = TRUE)</pre>
# Critical-t-value from table for two-tailed take given alpha/2
tc <- abs(qt(0.025, df))
# Test statistic
t = t_test$statistic
# Report test at 0.05 level
cat("t-statistic = ", t)
cat("Critical-t-value = ", tc)
if(t < tc){
 cat("Failed to reject null hypothesis that safety program had no effect")
} else {
  cat("Rejected null hypothesis that safety program had no effect")
```

Output:

Rejected null hypothesis that safety program had no effect

Question 3c:

```
#question 3c
# Given Data
bfr <- c(45, 73, 46, 124, 33, 57, 83, 34, 26, 17)
afr <- c(36, 60, 44, 119, 35, 51, 77, 29, 24, 11)

# Sign test
sign_test <- wilcox.test(bfr, afr, paired = TRUE, exact = FALSE)

# Extract p-value
p_value <- sign_test$p.value

# Report test at 0.05 level
cat("Number of negative differences =", sign_test$statistic)
cat("p-value =", p_value)

if(p_value < 0.05){
    cat("Rejected null hypothesis that safety program had no effect")
} else {
    cat("Failed to reject null hypothesis that safety program had no effect")
}</pre>
```

Output:

Rejected null hypothesis that safety program had no effect

Question 4:

```
#question 4
# Given Data
data <- c(0.265, 0.263, 0.266, 0.267, 0.267, 0.265, 0.267, 0.265, 0.268, 0.268, 0.268)
# Claimed mean
mu <- 0.265
# Wilcoxon signed-rank test
wilcox_test <- wilcox.test(data, mu, alternative = "two.sided", exact = FALSE)
# Test statistic
t = wilcox_test$statistic
# Extract p-value
p_value <- wilcox_test$p.value
# Output
cat("Test statistic = ", t)
cat("p-value = ", p_value)

if(p_value < 0.05){
    cat("Rejected the claim that the mean diameter is 0.265 mm")
} else {
    cat("Failed to reject the claim that the mean diameter is 0.265 mm")
}</pre>
```

Output:

Failed to reject the claim that the mean diameter is 0.265 mm