


Orthogonal transformations and rotations



What are orthogonal transformations

Let V be an inner product space and T be a linear transformation from V to V . T is said to be **orthogonal transformation** if

$$\langle T\mathbf{v}, T\mathbf{w} \rangle = \langle \mathbf{v}, \mathbf{w} \rangle \quad \forall \mathbf{v}, \mathbf{w} \in V.$$



What are orthogonal transformations

Let V be an inner product space and T be a linear transformation from V to V . T is said to be **orthogonal transformation** if

$$\langle Tv, Tw \rangle = \langle v, w \rangle \quad \forall v, w \in V.$$

When $V = \mathbb{R}^n$ with the usual inner product, a linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is orthogonal if it **preserves angles and lengths**.

What are orthogonal transformations

Let V be an inner product space and T be a linear transformation from V to V . T is said to be **orthogonal transformation** if

$$\langle Tv, Tw \rangle = \langle v, w \rangle \quad \forall v, w \in V.$$

When $V = \mathbb{R}^n$ with the usual inner product, a linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is orthogonal if it **preserves angles and lengths**.

Fact : It is enough to demand that the linear transformation preserves lengths. In that case, angles automatically get preserved (think of triangle congruences).




Example




Finding the rotation matrix in \mathbb{R}^2

Consider the standard basis $\{(1, 0), (0, 1)\}$ of \mathbb{R}^2 .




Finding the rotation matrix in \mathbb{R}^2

Consider the standard basis $\{(1, 0), (0, 1)\}$ of \mathbb{R}^2 . Rotate the plane by an angle θ . The vectors obtained after rotation tell us the matrix corresponding to this linear transformation.



Finding the rotation matrix in \mathbb{R}^2


Let T_θ be the corresponding linear transformation. Then



Finding the rotation matrix in \mathbb{R}^2

Let T_θ be the corresponding linear transformation. Then

$$T_\theta(1, 0) = (\cos(\theta), \sin(\theta)) \quad \text{and} \quad T_\theta(0, 1) = (-\sin(\theta), \cos(\theta)).$$



Finding the rotation matrix in \mathbb{R}^2

Let T_θ be the corresponding linear transformation. Then

$$T_\theta(1, 0) = (\cos(\theta), \underline{\sin(\theta)}) \quad \text{and} \quad T_\theta(0, 1) = (-\underline{\sin(\theta)}, \cos(\theta)).$$

Thus the matrix corresponding to this linear transformation is

$$R_\theta = \begin{pmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{pmatrix}.$$

Finding the rotation matrix in \mathbb{R}^2

Let T_θ be the corresponding linear transformation. Then

$$T_\theta(1, 0) = (\cos(\theta), \sin(\theta)) \quad \text{and} \quad T_\theta(0, 1) = (-\sin(\theta), \cos(\theta)).$$

Thus the matrix corresponding to this linear transformation is

$$R_\theta = \begin{pmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{pmatrix}.$$

Note that $R_\theta^T = R_{-\theta}$ and $R_\theta^T R_\theta = R_\theta R_\theta^T = I$.

Finding the rotation matrix in \mathbb{R}^2

Let T_θ be the corresponding linear transformation. Then

$$T_\theta(1, 0) = (\cos(\theta), \sin(\theta)) \quad \text{and} \quad T_\theta(0, 1) = (-\sin(\theta), \cos(\theta)).$$

Thus the matrix corresponding to this linear transformation is

$$R_\theta = \begin{pmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{pmatrix}.$$


$$\begin{aligned} R_{-\theta} &= \begin{pmatrix} \cos(-\theta) & \sin(-\theta) \\ -\sin(-\theta) & \cos(-\theta) \end{pmatrix} \\ &= R_\theta^T. \end{aligned}$$

Note that $R_\theta^T = R_{-\theta}$ and $R_\theta^T R_\theta = R_\theta R_\theta^T = I$.

Further note that since angles and lengths are preserved and the standard basis is orthonormal, the rotated vectors are also orthonormal and therefore yield an orthonormal basis of \mathbb{R}^2 .

Rotations in \mathbb{R}^3

Consider the rotations about the axes in \mathbb{R}^3 .



Rotations in \mathbb{R}^3


Consider the rotations about the axes in \mathbb{R}^3 . Since these clearly preserve angles and distances and are linear transformations, they are orthogonal transformations.



Rotations in \mathbb{R}^3

Consider the rotations about the axes in \mathbb{R}^3 . Since these clearly preserve angles and distances and are linear transformations, they are orthogonal transformations.

Rotations about the axes can be described by considering its effect on the standard basis $\{e_1, e_2, e_3\}$.



Rotations in \mathbb{R}^3

Consider the rotations about the axes in \mathbb{R}^3 . Since these clearly preserve angles and distances and are linear transformations, they are orthogonal transformations.

Rotations about the axes can be described by considering its effect on the standard basis $\{e_1, e_2, e_3\}$.

When considering the rotation about the Z-axis, e_3 remains unchanged and the XY-plane gets rotated exactly as in the previous case of \mathbb{R}^2 .

Rotations in \mathbb{R}^3

Consider the rotations about the axes in \mathbb{R}^3 . Since these clearly preserve angles and distances and are linear transformations, they are orthogonal transformations.

Rotations about the axes can be described by considering its effect on the standard basis $\{e_1, e_2, e_3\}$.


When considering the rotation about the Z-axis, e_3 remains unchanged and the XY-plane gets rotated exactly as in the previous case of \mathbb{R}^2 . Therefore its matrix is

$$T_3(\theta) = \begin{pmatrix} \cos(\theta) & -\sin(\theta) & 0 \\ \sin(\theta) & \cos(\theta) & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Rotations in \mathbb{R}^3 (contd.)

Similarly, the matrix corresponding to rotation about the X-axis is

$$T_1(\theta) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos(\theta) & -\sin(\theta) \\ 0 & \sin(\theta) & \cos(\theta) \end{pmatrix}$$



Rotations in \mathbb{R}^3 (contd.)

Similarly, the matrix corresponding to rotation about the X-axis is

$$T_1(\theta) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos(\theta) & -\sin(\theta) \\ 0 & \sin(\theta) & \cos(\theta) \end{pmatrix}$$

and the matrix corresponding to rotation about the Y-axis is

$$T_2(\theta) = \begin{pmatrix} \cos(\theta) & 0 & -\sin(\theta) \\ 0 & 1 & 0 \\ \sin(\theta) & 0 & \cos(\theta) \end{pmatrix}.$$

Rotations in \mathbb{R}^3 (contd.)

Similarly, the matrix corresponding to rotation about the X-axis is

$$T_1(\theta) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos(\theta) & -\sin(\theta) \\ 0 & \sin(\theta) & \cos(\theta) \end{pmatrix}$$

and the matrix corresponding to rotation about the Y-axis is


$$T_2(\theta) = \begin{pmatrix} \cos(\theta) & 0 & -\sin(\theta) \\ 0 & 1 & 0 \\ \sin(\theta) & 0 & \cos(\theta) \end{pmatrix}.$$

Notice : $T_i(\theta)^T = T_i(-\theta)$ and $T_i(\theta)^T T_i(\theta) = T_i(\theta) T_i(\theta)^T = I$.

Another example of an orthogonal transformation

Let us define a linear transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$, where

$$T(x_1, x_2, x_3) = \frac{1}{3}(x_1 - 2x_2 + 2x_3, 2x_1 - x_2 - 2x_3, 2x_1 + 2x_2 + x_3).$$




Another example of an orthogonal transformation

Let us define a linear transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$, where

$$T(x_1, x_2, x_3) = \frac{1}{3}(x_1 - 2x_2 + 2x_3, 2x_1 - x_2 - 2x_3, 2x_1 + 2x_2 + x_3).$$

Then evaluating T on the standard basis $\{e_1, e_2, e_3\}$ yields :



Another example of an orthogonal transformation

Let us define a linear transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$, where

$$T(x_1, x_2, x_3) = \frac{1}{3}(x_1 - 2x_2 + 2x_3, 2x_1 - x_2 - 2x_3, 2x_1 + 2x_2 + x_3).$$

Then evaluating T on the standard basis $\{e_1, e_2, e_3\}$ yields :

$$T(e_1) = v_1 = \left(\frac{1}{3}, \frac{2}{3}, \frac{2}{3} \right)$$

$$T(e_2) = v_2 = \left(-\frac{2}{3}, -\frac{1}{3}, \frac{2}{3} \right)$$

$$T(e_3) = v_3 = \left(\frac{2}{3}, -\frac{2}{3}, \frac{1}{3} \right) .$$

Another example of an orthogonal transformation

Let us define a linear transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$, where

$$T(x_1, x_2, x_3) = \frac{1}{3}(x_1 - 2x_2 + 2x_3, 2x_1 - x_2 - 2x_3, 2x_1 + 2x_2 + x_3).$$

Then evaluating T on the standard basis $\{e_1, e_2, e_3\}$ yields :

$$T(e_1) = v_1 = \left(\frac{1}{3}, \frac{2}{3}, \frac{2}{3} \right) \quad \leftarrow$$

$$T(e_2) = v_2 = \left(-\frac{2}{3}, -\frac{1}{3}, \frac{2}{3} \right) \quad \leftarrow$$

$$T(e_3) = v_3 = \left(\frac{2}{3}, -\frac{2}{3}, \frac{1}{3} \right) \quad . \quad \leftarrow$$

Thus, the matrix corresponding to T is $A = \frac{1}{3} \begin{pmatrix} 1 & -2 & 2 \\ 2 & -1 & -2 \\ 2 & 2 & 1 \end{pmatrix}$.

Orthogonal matrices


As $\{v_1, v_2, v_3\}$ is an orthogonal set, the linear transformation T is an orthogonal transformation.



Orthogonal matrices

As $\{v_1, v_2, v_3\}$ is an orthogonal set, the linear transformation T is ^{normal} an orthogonal transformation.

Observe that the $AA^T = A^T A = I_3$.



Orthogonal matrices

As $\{v_1, v_2, v_3\}$ is an orthogonal set, the linear transformation T is an orthogonal transformation.

Observe that the $AA^T = A^T A = I_3$.

A square matrix A is called an **orthogonal matrix** if $AA^T = A^T A = I$


$$\|T\mathbf{x}\| = \sqrt{\langle T\mathbf{x}, T\mathbf{x} \rangle} = \sqrt{(T\mathbf{x}) \cdot (T\mathbf{x})} = \sqrt{(A\mathbf{x})^T A\mathbf{x}} = \sqrt{\mathbf{x}^T A^T A \mathbf{x}} = \sqrt{\mathbf{x}^T \mathbf{x}} = \|\mathbf{x}\|.$$

Orthogonal matrices

As $\{v_1, v_2, v_3\}$ is an orthogonal set, the linear transformation T is an orthogonal transformation.

Observe that the $AA^T = A^T A = I_3$.

A square matrix A is called an **orthogonal matrix** if $AA^T = A^T A = I$



Thank you

