


Lengths and angles

Sarang S. Sane



The dot product of two vectors in \mathbb{R}^2


Consider the two vectors $(3, 4)$ and $(2, 7)$ in \mathbb{R}^2 .



The dot product of two vectors in \mathbb{R}^2

Consider the two vectors $(3, 4)$ and $(2, 7)$ in \mathbb{R}^2 . The **dot product** of these two vectors gives us a scalar as follows:

$$(3, 4) \cdot (2, 7) = 3 \times 2 + 4 \times 7 = 6 + 28 = 34$$



The dot product of two vectors in \mathbb{R}^2

Consider the two vectors $(3, 4)$ and $(2, 7)$ in \mathbb{R}^2 . The **dot product** of these two vectors gives us a scalar as follows:

$$(3, 4) \cdot (2, 7) = 3 \times 2 + 4 \times 7 = 6 + 28 = 34$$

For two general vectors (x_1, y_1) and (x_2, y_2) in \mathbb{R}^2 , the **dot product** of these two vectors is the scalar computed as follows :

The dot product of two vectors in \mathbb{R}^2

Consider the two vectors $(3, 4)$ and $(2, 7)$ in \mathbb{R}^2 . The **dot product** of these two vectors gives us a scalar as follows:

$$(3, 4) \cdot (2, 7) = 3 \times 2 + 4 \times 7 = 6 + 28 = 34$$

For two general vectors (x_1, y_1) and (x_2, y_2) in \mathbb{R}^2 , the **dot product** of these two vectors is the scalar computed as follows :

$$(x_1, y_1) \cdot (x_2, y_2) = x_1 x_2 + y_1 y_2.$$


The length of a vector in \mathbb{R}^2

Let us find the length of the vector $(3, 4)$ in \mathbb{R}^2 .




The length of a vector in \mathbb{R}^2

Let us find the length of the vector $(3, 4)$ in \mathbb{R}^2 .



The length of a vector in \mathbb{R}^2


Let us find the length of the vector $(3, 4)$ in \mathbb{R}^2 .



Using Pythagoras' theorem, the length of the vector $(3, 4)$ is $\sqrt{3^2 + 4^2} = 5$ units.

The relation between length and dot product in \mathbb{R}^2

Observe that $(3, 4) \cdot (3, 4) = 3^2 + 4^2$,



The relation between length and dot product in \mathbb{R}^2

Observe that $(3, 4) \cdot (3, 4) = 3^2 + 4^2$, and hence the length of $(3, 4)$ is the square root of the dot product of the vector with itself.

$$\text{Length of the vector } (3, 4) = \sqrt{(3, 4) \cdot (3, 4)} = \sqrt{3^2 + 4^2} = \sqrt{\cancel{3}^{\times} \cancel{4}^{\times}} = \sqrt{25} = 5.$$

The relation between length and dot product in \mathbb{R}^2

Observe that $(3, 4) \cdot (3, 4) = 3^2 + 4^2$, and hence the length of $(3, 4)$ is the square root of the dot product of the vector with itself.


$$\text{Length of the vector } (3, 4) = \sqrt{(3, 4) \cdot (3, 4)} = \sqrt{3^2 + 4^2} = \sqrt{\cancel{9} + \cancel{16}} = \cancel{\sqrt{25}} = 5.$$

More generally, the length of the vector $(x, y) \in \mathbb{R}^2$ is $\sqrt{x^2 + y^2} = \sqrt{(x, y) \cdot (x, y)}$.

$$\begin{aligned} & \sqrt{x^2 + y^2} \\ &= \sqrt{(x, y) \cdot (x, y)} \end{aligned}$$


The angle between two vectors in \mathbb{R}^2

- The angle between the vectors u and v ~~and~~ measures how far the direction is of v from u (or vice versa).




The angle between two vectors in \mathbb{R}^2

- ▶ The angle between the vectors u and v and measures how far the direction is of v from u (or vice versa). e.g. θ is the angle between $u = (3, 4)$ and $v = (1, 5)$.



The angle between two vectors in \mathbb{R}^2


- ▶ The angle between the vectors u and v and measures how far the direction is of v from u (or vice versa). e.g. θ is the angle between $u = (3, 4)$ and $v = (1, 5)$.



- ▶ It is measured in degrees (between 0 and 360) or radians (between 0 and 2π).

The angle between two vectors in \mathbb{R}^2

- ▶ The angle between the vectors u and v and measures how far the direction is of v from u (or vice versa). e.g. θ is the angle between $u = (3, 4)$ and $v = (1, 5)$.




- ▶ It is measured in degrees (between 0 and 360) or radians (between 0 and 2π). *on measured*
- ▶ The angle is often described by computing its trigonometric functions (e.g. \sin , \cos , \tan).

The dot product and the angle between two vectors in \mathbb{R}^2

Let u and v be two vectors in \mathbb{R}^2 . Then we can compute the angle θ between the vectors u and v using the dot products as :

$$\cos(\theta) = \frac{u \cdot v}{\sqrt{(v \cdot v) \times (u \cdot u)}} \quad \text{i.e.} \quad \theta = \cos^{-1} \left(\frac{u \cdot v}{\sqrt{(v \cdot v) \times (u \cdot u)}} \right).$$




$$\cos(\theta) = \frac{\text{length of adj. side}}{\text{length of hypotenuse}}$$

$$\sin(\theta) = \frac{\text{length of the opp. side}}{\text{length of hypotenuse}}$$

$$\tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)} = \frac{\text{length(opp.)}}{\text{length(adj.)}}$$

The dot product of two vectors in \mathbb{R}^3


Consider the two vectors $(1, 2, 3)$ and $(2, 0, 1)$ in \mathbb{R}^3 .



The dot product of two vectors in \mathbb{R}^3

Consider the two vectors $(1, 2, 3)$ and $(2, 0, 1)$ in \mathbb{R}^3 . The **dot product** of these two vectors gives us a scalar as follows:

$$(1, 2, 3) \cdot (2, 0, 1) = 1 \times 2 + 2 \times 0 + 3 \times 1 = 2 + 0 + 3 = 5$$



The dot product of two vectors in \mathbb{R}^3

Consider the two vectors $(1, 2, 3)$ and $(2, 0, 1)$ in \mathbb{R}^3 . The **dot product** of these two vectors gives us a scalar as follows:

$$(1, 2, 3) \cdot (2, 0, 1) = 1 \times 2 + 2 \times 0 + 3 \times 1 = 2 + 0 + 3 = 5$$

For two general vectors (x_1, y_1, z_1) and (x_2, y_2, z_2) in \mathbb{R}^3 , the **dot product** of these two vectors is the scalar computed as follows :

The dot product of two vectors in \mathbb{R}^3

Consider the two vectors $(1, 2, 3)$ and $(2, 0, 1)$ in \mathbb{R}^3 . The **dot product** of these two vectors gives us a scalar as follows:

$$(1, 2, 3) \cdot (2, 0, 1) = 1 \times 2 + 2 \times 0 + 3 \times 1 = 2 + 0 + 3 = 5$$

For two general vectors (x_1, y_1, z_1) and (x_2, y_2, z_2) in \mathbb{R}^3 , the **dot product** of these two vectors is the scalar computed as follows :

$$(x_1, y_1, z_1) \cdot (x_2, y_2, z_2) = x_1 x_2 + y_1 y_2 + z_1 z_2.$$


Length of a vector in \mathbb{R}^3

Let us find the length of the vector $(4, 3, 3)$ in \mathbb{R}^3 .




Length of a vector in \mathbb{R}^3

Let us find the length of the vector $(4, 3, 3)$ in \mathbb{R}^3 .



Length of a vector in \mathbb{R}^3


Let us find the length of the vector $(4, 3, 3)$ in \mathbb{R}^3 .



By using Pythagoras' theorem, the length of $(4, 3, 3)$ is $\sqrt{4^2 + 3^2 + 3^2} = \sqrt{34}$ units.

The length and dot product in \mathbb{R}^3

Observe that $(4, 3, 3) \cdot (4, 3, 3) = 4^2 + 3^2 + 3^2$ and hence the length of $(4, 3, 3)$ can be expressed as the square root of dot product of the vector with itself.



The length and dot product in \mathbb{R}^3

Observe that $(4, 3, 3) \cdot (4, 3, 3) = 4^2 + 3^2 + 3^2$ and hence the length of $(4, 3, 3)$ can be expressed as the square root of dot product of the vector with itself.

Length of the vector $(4, 3, 3)$

$$\begin{aligned}&= \sqrt{(4, 3, 3) \cdot (4, 3, 3)} \\&= \sqrt{4^2 + 3^2 + 3^2} \\&= \sqrt{34} \text{ units}\end{aligned}$$

The length and dot product in \mathbb{R}^3

Observe that $(4, 3, 3) \cdot (4, 3, 3) = 4^2 + 3^2 + 3^2$ and hence the length of $(4, 3, 3)$ can be expressed as the square root of dot product of the vector with itself.

Length of the vector $(4, 3, 3)$

$$\begin{aligned}&= \sqrt{(4, 3, 3) \cdot (4, 3, 3)} \\&= \sqrt{4^2 + 3^2 + 3^2} \\&= \sqrt{34} \text{ units}\end{aligned}$$

More generally, the length of the vector $(x, y, z) \in \mathbb{R}^3$ is $\sqrt{x^2 + y^2 + z^2} = \sqrt{(x, y, z) \cdot (x, y, z)}$.

The angle between two vectors in \mathbb{R}^3 and the dot product


The angle between the vectors u and v in \mathbb{R}^3 is the angle between them computed by passing a plane through them.



The angle between two vectors in \mathbb{R}^3 and the dot product

The angle between the vectors u and v in \mathbb{R}^3 is the angle between them computed by passing a plane through them.

It measures how far the direction is of v from u (or vice versa) on that plane.



The angle between two vectors in \mathbb{R}^3 and the dot product

The angle between the vectors u and v in \mathbb{R}^3 is the angle between them computed by passing a plane through them.

It measures how far the direction is of v from u (or vice versa) on that plane.

Let u and v be two vectors in \mathbb{R}^3 . Then we can compute the angle θ between the vectors u and v using the dot product as :

$$\cos(\theta) = \frac{u \cdot v}{\sqrt{(v \cdot v) \times (u \cdot u)}} \quad i.e. \quad \theta = \cos^{-1} \left(\frac{u \cdot v}{\sqrt{(v \cdot v) \times (u \cdot u)}} \right).$$

The angle between two vectors in \mathbb{R}^3 and the dot product

The angle between the vectors u and v in \mathbb{R}^3 is the angle between them computed by passing a plane through them.

It measures how far the direction is of v from u (or vice versa) on that plane.

Let u and v be two vectors in \mathbb{R}^3 . Then we can compute the angle θ between the vectors u and v using the dot product as :

$$\cos(\theta) = \frac{u \cdot v}{\sqrt{(v \cdot v) \times (u \cdot u)}} \quad \text{i.e.} \quad \theta = \cos^{-1} \left(\frac{u \cdot v}{\sqrt{(v \cdot v) \times (u \cdot u)}} \right).$$



Examples of computing angles in \mathbb{R}^3


Let us compute the angle θ between $(1, 0, 0)$ and $(1, 0, 1)$.



Examples of computing angles in \mathbb{R}^3

Let us compute the angle θ between $(1, 0, 0)$ and $(1, 0, 1)$.

$$(1, 0, 0) \cdot (1, 0, 1) = 1, (1, 0, 1) \cdot (1, 0, 1) = 2, (1, 0, 0) \cdot (1, 0, 0) = 1.$$




Examples of computing angles in \mathbb{R}^3

Let us compute the angle θ between $(1, 0, 0)$ and $(1, 0, 1)$.

$$(1, 0, 0) \cdot (1, 0, 1) = 1, (1, 0, 1) \cdot (1, 0, 1) = 2, (1, 0, 0) \cdot (1, 0, 0) = 1.$$

$$\text{Hence, } \theta = \cos^{-1} \left(\frac{1}{\sqrt{2}} \right) = \frac{\pi}{4} \text{ radians or } 45^\circ.$$



Examples of computing angles in \mathbb{R}^3

Let us compute the angle θ between $(1, 0, 0)$ and $(1, 0, 1)$.

$$(1, 0, 0) \cdot (1, 0, 1) = 1, (1, 0, 1) \cdot (1, 0, 1) = 2, (1, 0, 0) \cdot (1, 0, 0) = 1.$$

$$\text{Hence, } \theta = \cos^{-1} \left(\frac{1}{\sqrt{2}} \right) = \frac{\pi}{4} \text{ radians or } 45^\circ.$$

Similarly, the angle between between $(1, 0, 0)$ and $(1, 1, 1)$ is


$$\cos^{-1} \left(\frac{1}{\sqrt{3}} \right).$$

$$(1, 0, 0) \cdot (1, 1, 1) = \frac{1}{\sqrt{3}}, (1, 1, 1) \cdot (1, 1, 1) = 3, \\ (1, 0, 0) \cdot (1, 1, 1) = \frac{1}{\sqrt{3}} \cdot \frac{1}{\sqrt{3}} \cdot (1, 1, 1) \cdot (1, 1, 1) = 1.$$

$$\theta = \cos^{-1} \left(\frac{1}{\sqrt{3}} \right) = \omega \approx \left(\frac{1}{\sqrt{3}} \right) .$$

Dot products in \mathbb{R}^n : length and angle

Let $u = (u_1, u_2, \dots, u_n)$ and $v = (v_1, v_2, \dots, v_n)$ be vectors in \mathbb{R}^n .




Dot products in \mathbb{R}^n : length and angle

Let $u = (u_1, u_2, \dots, u_n)$ and $v = (v_1, v_2, \dots, v_n)$ be vectors in \mathbb{R}^n .

The dot product of the two vectors u and v is defined as

$$u \cdot v = u_1 v_1 + u_2 v_2 + \dots + u_n v_n.$$



Dot products in \mathbb{R}^n : length and angle


Let $u = (u_1, u_2, \dots, u_n)$ and $v = (v_1, v_2, \dots, v_n)$ be vectors in \mathbb{R}^n .

The dot product of the two vectors u and v is defined as

$$u \cdot v = u_1 v_1 + u_2 v_2 + \dots + u_n v_n.$$

The length of the vector u is denoted by $\|u\|$ and defined by

$$\|u\| = \sqrt{u \cdot u}.$$



Dot products in \mathbb{R}^n : length and angle

Let $u = (u_1, u_2, \dots, u_n)$ and $v = (v_1, v_2, \dots, v_n)$ be vectors in \mathbb{R}^n .

The dot product of the two vectors u and v is defined as

$$u \cdot v = u_1 v_1 + u_2 v_2 + \dots + u_n v_n.$$

The length of the vector u is denoted by $\|u\|$ and defined by

$$\|u\| = \sqrt{u \cdot u}.$$

The angle θ between the two vectors u and v is measured on the 2-dimensional plane spanned by u and v and can be computed as :

Dot products in \mathbb{R}^n : length and angle

Let $u = (u_1, u_2, \dots, u_n)$ and $v = (v_1, v_2, \dots, v_n)$ be vectors in \mathbb{R}^n .

The dot product of the two vectors u and v is defined as

$$u \cdot v = u_1 v_1 + u_2 v_2 + \dots + u_n v_n.$$

The length of the vector u is denoted by $\|u\|$ and defined by

$$\|u\| = \sqrt{u \cdot u}.$$

The angle θ between the two vectors u and v is measured on the 2-dimensional plane spanned by u and v and can be computed as :

$$\cos(\theta) = \frac{u \cdot v}{\|u\| \times \|v\|} \quad i.e. \quad \theta = \cos^{-1} \left(\frac{u \cdot v}{\sqrt{(v \cdot v) \times (u \cdot u)}} \right).$$

Thank you

