

Determinants (Part 2)


Sarang S. Sane

Recall from part 1 :



Recall from part 1 :


- ▶ $A = [a]$ $\det(A) = a$



Recall from part 1 :


$$\blacktriangleright A = [a] \quad \det(A) = a$$

$$\blacktriangleright A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$




Recall from part 1 :

- ▶ $A = [a] \quad \det(A) = a$
- ▶ $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \quad \det(A) = ad - bc$



Recall from part 1 :

- ▶ $A = [a] \quad \det(A) = a$
- ▶ $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \quad \det(A) = ad - bc$
- ▶ $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$



Recall from part 1 :

► $A = [a] \quad \det(A) = a$

► $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \quad \det(A) = ad - bc$

► $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$

Expanding with respect to the 1st row :

$$\det(A) = a_{11} \times \det \begin{bmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{bmatrix} - a_{12} \times \det \begin{bmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{bmatrix} + a_{13} \times \det \begin{bmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix}$$

Recall from part 1 :

► $A = [a] \quad \det(A) = a$

► $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \quad \det(A) = ad - bc$


► $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$

Expanding with respect to the 1st row :

$$\begin{aligned}\det(A) &= a_{11} \times \det \begin{bmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{bmatrix} - a_{12} \times \det \begin{bmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{bmatrix} + a_{13} \times \det \begin{bmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix} \\ &= a_{11}a_{22}a_{33} - a_{11}a_{23}a_{32} - a_{12}a_{21}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32} - a_{13}a_{22}a_{31}\end{aligned}$$

An example

$$A = \begin{bmatrix} 2 & 4 & 3 \\ 0 & 8 & 7 \\ 0 & 0 & 9 \end{bmatrix}$$



An example

$$A = \begin{bmatrix} 2 & 4 & 3 \\ 0 & 8 & 7 \\ 0 & 0 & 9 \end{bmatrix}$$

$$\det(A) = 2 \times \det \begin{bmatrix} 8 & 7 \\ 0 & 9 \end{bmatrix} - 4 \times \det \begin{bmatrix} 0 & 7 \\ 0 & 9 \end{bmatrix} + 3 \times \det \begin{bmatrix} 0 & 8 \\ 0 & 0 \end{bmatrix}$$

An example

$$A = \begin{bmatrix} 2 & 4 & 3 \\ 0 & 8 & 7 \\ 0 & 0 & 9 \end{bmatrix}$$

$$\begin{aligned} \det(A) &= 2 \times \det \begin{bmatrix} 8 & 7 \\ 0 & 9 \end{bmatrix} - 4 \times \det \begin{bmatrix} 0 & 7 \\ 0 & 9 \end{bmatrix} + 3 \times \det \begin{bmatrix} 0 & 8 \\ 0 & 0 \end{bmatrix} \\ &= 2(72 - 0) - 4(0 - 0) + 3(0 - 0) \end{aligned}$$

An example

$$A = \begin{bmatrix} 2 & 4 & 3 \\ 0 & 8 & 7 \\ 0 & 0 & 9 \end{bmatrix}$$

$$\begin{aligned} \det(A) &= 2 \times \det \begin{bmatrix} 8 & 7 \\ 0 & 9 \end{bmatrix} - 4 \times \det \begin{bmatrix} 0 & 7 \\ 0 & 9 \end{bmatrix} + 3 \times \det \begin{bmatrix} 0 & 8 \\ 0 & 0 \end{bmatrix} \\ &= 2(72 - 0) - 4(0 - 0) + 3(0 - 0) \\ &= 2(72) - 4(0) + 3(0) \end{aligned}$$

An example


$$A = \begin{bmatrix} 2 & 4 & 3 \\ 0 & 8 & 7 \\ 0 & 0 & 9 \end{bmatrix}$$

$$\begin{aligned} \det(A) &= 2 \times \det \begin{bmatrix} 8 & 7 \\ 0 & 9 \end{bmatrix} - 4 \times \det \begin{bmatrix} 0 & 7 \\ 0 & 9 \end{bmatrix} + 3 \times \det \begin{bmatrix} 0 & 8 \\ 0 & 0 \end{bmatrix} \\ &= 2(72 - 0) - 4(0 - 0) + 3(0 - 0) \\ &= 2(72) - 4(0) + 3(0) \\ &= 144 \end{aligned}$$

$$A = \begin{bmatrix} 2 & 4 & 3 \\ 0 & 8 & 7 \\ 0 & 0 & 9 \end{bmatrix}$$

$$\begin{aligned} \det(A) &= 2 \times \det \begin{bmatrix} 8 & 7 \\ 0 & 9 \end{bmatrix} - 4 \times \det \begin{bmatrix} 0 & 7 \\ 0 & 9 \end{bmatrix} + 3 \times \det \begin{bmatrix} 0 & 8 \\ 0 & 0 \end{bmatrix} \\ &= 2(72 - 0) - 4(0 - 0) + 3(0 - 0) \\ &= 2(72) - 4(0) + 3(0) \\ &= 144 \\ &= 2 \times 8 \times 9 \end{aligned}$$

$$A = \begin{bmatrix} 2 & 4 & 3 \\ 0 & 8 & 7 \\ 0 & 0 & 9 \end{bmatrix}$$




$$\begin{aligned} \det(A) &= 2 \times \det \begin{bmatrix} 8 & 7 \\ 0 & 9 \end{bmatrix} - 4 \times \det \begin{bmatrix} 0 & 7 \\ 0 & 9 \end{bmatrix} + 3 \times \det \begin{bmatrix} 0 & 8 \\ 0 & 0 \end{bmatrix} \\ &= 2(72 - 0) - 4(0 - 0) + 3(0 - 0) \\ &= 2(72) - 4(0) + 3(0) \\ &= 144 \\ &= 2 \times 8 \times 9 \end{aligned}$$



This is an upper triangular matrix. For such matrices, the determinant is the product of the diagonal elements.

The transpose of a matrix and its determinant



The transpose of a matrix and its determinant

The transpose of $A_{m \times n}$ is the $n \times m$ matrix with (i, j) -th entry A_{ji} .




The transpose of a matrix and its determinant

The transpose of $A_{m \times n}$ is the $n \times m$ matrix with (i, j) -th entry A_{ji} .

Notation : A^T

Definition : $(A^T)_{ij} = A_{ji}$




The transpose of $A_{m \times n}$ is the $n \times m$ matrix with (i, j) -th entry A_{ji} .

Notation : A^T

Definition : $(A^T)_{ij} = A_{ji}$

Let $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$ then $A^T = \begin{bmatrix} a_{11} & a_{21} & a_{31} \\ a_{12} & a_{22} & a_{32} \\ a_{13} & a_{23} & a_{33} \end{bmatrix}$



The transpose of a matrix and its determinant

The transpose of $A_{m \times n}$ is the $n \times m$ matrix with (i, j) -th entry A_{ji} .

Notation : A^T


Definition : $(A^T)_{ij} = A_{ji}$

Let $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$ then $A^T = \begin{bmatrix} a_{11} & a_{21} & a_{31} \\ a_{12} & a_{22} & a_{32} \\ a_{13} & a_{23} & a_{33} \end{bmatrix}$

$$\begin{aligned} \det(A^T) &= a_{11} \times \det \begin{bmatrix} a_{22} & a_{32} \\ a_{23} & a_{33} \end{bmatrix} - a_{21} \times \det \begin{bmatrix} a_{12} & a_{32} \\ a_{13} & a_{33} \end{bmatrix} + a_{31} \times \det \begin{bmatrix} a_{12} & a_{22} \\ a_{13} & a_{23} \end{bmatrix} \\ &= a_{11}(a_{22}a_{33} - a_{32}a_{23}) - a_{21}(a_{12}a_{33} - a_{32}a_{13}) + a_{31}(a_{12}a_{23} - a_{22}a_{13}) \\ &= a_{11}a_{22}a_{33} - a_{11}a_{32}a_{23} - a_{21}a_{12}a_{33} + a_{21}a_{32}a_{13} + a_{31}a_{12}a_{23} - a_{31}a_{22}a_{13} \\ &= \det(A) \end{aligned}$$

Minors and Cofactors

If A is an $n \times n$ square matrix with $n \leq 4$. Then the minor of the entry in the i -th row and j -th column is the determinant of the submatrix formed by deleting the i -th row and j -th column.




Minors and Cofactors

If A is an $n \times n$ square matrix with $n \leq 4$. Then the minor of the entry in the i -th row and j -th column is the determinant of the submatrix formed by deleting the i -th row and j -th column.

Name : the (i,j) -th minor

Notation : M_{ij}



Minors and Cofactors

If A is an $n \times n$ square matrix with $n \leq 4$. Then the minor of the entry in the i -th row and j -th column is the determinant of the submatrix formed by deleting the i -th row and j -th column.

Name : the (i,j) -th minor Notation : M_{ij}

Example : $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$ $M_{11} = \det \begin{bmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{bmatrix}.$

Minors and Cofactors

If A is an $n \times n$ square matrix with $n \leq 4$. Then the minor of the entry in the i -th row and j -th column is the determinant of the submatrix formed by deleting the i -th row and j -th column.

Name : the (i,j) -th minor Notation : M_{ij}

Example : $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$ $M_{11} = \det \begin{bmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{bmatrix}.$

The (i,j) -th cofactor $C_{ij} = (-1)^{i+j} M_{ij}.$

If A is an $n \times n$ square matrix with $n \leq 4$. Then the minor of the entry in the i -th row and j -th column is the determinant of the submatrix formed by deleting the i -th row and j -th column.

Name : the (i,j) -th minor Notation : M_{ij}


Example : $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$ $M_{11} = \det \begin{bmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{bmatrix}.$

The (i,j) -th cofactor $C_{ij} = (-1)^{i+j} M_{ij}.$

Above example : $C_{11} = (-1)^{1+1} M_{11} = M_{11}$

$$C_{23} = (-1)^{2+3} M_{23} = -M_{23}$$


Determinant in terms of minors and cofactors



Determinant in terms of minors and cofactors

Observe that : For $A_{3 \times 3}$


$$\det(A) = a_{11} \times \det \begin{bmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{bmatrix} - a_{12} \times \det \begin{bmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{bmatrix} + a_{13} \times \det \begin{bmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix}$$



Determinant in terms of minors and cofactors

Observe that : For $A_{3 \times 3}$

$$\begin{aligned} \det(A) &= a_{11} \times \det \begin{bmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{bmatrix} - a_{12} \times \det \begin{bmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{bmatrix} + a_{13} \times \det \begin{bmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix} \\ &= a_{11} \times M_{11} - a_{12} \times M_{12} + a_{13} \times M_{13} \end{aligned}$$



Observe that : For $A_{3 \times 3}$

$$\begin{aligned} \det(A) &= a_{11} \times \det \begin{bmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{bmatrix} - a_{12} \times \det \begin{bmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{bmatrix} + a_{13} \times \det \begin{bmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix} \\ &= a_{11} \times M_{11} - a_{12} \times M_{12} + a_{13} \times M_{13} \\ &= a_{11} \times C_{11} + a_{12} \times C_{12} + a_{13} \times C_{13} \end{aligned}$$

Determinant in terms of minors and cofactors

Observe that : For $A_{3 \times 3}$

$$\begin{aligned} \det(A) &= a_{11} \times \det \begin{bmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{bmatrix} - a_{12} \times \det \begin{bmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{bmatrix} + a_{13} \times \det \begin{bmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix} \\ &= a_{11} \times M_{11} - a_{12} \times M_{12} + a_{13} \times M_{13} \\ &= a_{11} \times C_{11} + a_{12} \times C_{12} + a_{13} \times C_{13} \end{aligned}$$

This formula holds for $A_{2 \times 2}$.

Determinant in terms of minors and cofactors

Observe that : For $A_{3 \times 3}$

$$\begin{aligned} \det(A) &= a_{11} \times \det \begin{bmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{bmatrix} - a_{12} \times \det \begin{bmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{bmatrix} + a_{13} \times \det \begin{bmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix} \\ &= a_{11} \times M_{11} - a_{12} \times M_{12} + a_{13} \times M_{13} \\ &= a_{11} \times C_{11} + a_{12} \times C_{12} + a_{13} \times C_{13} \end{aligned}$$

This formula holds for $A_{2 \times 2}$. We use it to generalize the determinant beyond $n = 3$. Generalization to $A_{4 \times 4}$:

Observe that : For $A_{3 \times 3}$

$$\begin{aligned} \det(A) &= a_{11} \times \det \begin{bmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{bmatrix} - a_{12} \times \det \begin{bmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{bmatrix} + a_{13} \times \det \begin{bmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix} \\ &= \underline{a_{11} \times M_{11} - a_{12} \times M_{12} + a_{13} \times M_{13}} \\ &= a_{11} \times C_{11} + a_{12} \times C_{12} + a_{13} \times C_{13} \end{aligned}$$

This formula holds for $A_{2 \times 2}$. We use it to generalize the determinant beyond $n = 3$. Generalization to $A_{4 \times 4}$:

Definition

$$\det(A) = \sum_{j=1}^4 (-1)^{1+j} a_{1j} M_{1j} = \sum_{i=1}^4 a_{1j} C_{1j}$$

$$\begin{aligned} &= a_{11} \times \det \begin{bmatrix} a_{22} & a_{23} & a_{24} \\ a_{32} & a_{33} & a_{34} \\ a_{42} & a_{43} & a_{44} \end{bmatrix} - a_{12} \times \det \begin{bmatrix} a_{21} & a_{23} & a_{24} \\ a_{31} & a_{33} & a_{34} \\ a_{41} & a_{43} & a_{44} \end{bmatrix} + a_{13} \times \det \begin{bmatrix} a_{21} & a_{22} & a_{24} \\ a_{31} & a_{32} & a_{34} \\ a_{41} & a_{42} & a_{44} \end{bmatrix} \\ &\quad - a_{14} \times \det \begin{bmatrix} a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{bmatrix} \end{aligned}$$


Inductive definition of the determinant

Suppose $A_{n \times n}$ is given and we know how to define determinants for $n - 1 \times n - 1$ matrices.



Inductive definition of the determinant


Suppose $A_{n \times n}$ is given and we know how to define determinants for $n - 1 \times n - 1$ matrices. Define minors and cofactors as before. :



Inductive definition of the determinant

Suppose $A_{n \times n}$ is given and we know how to define determinants for $n - 1 \times n - 1$ matrices. Define minors and cofactors as before. :

The (i, j) -th minor is the determinant of the submatrix formed by deleting the i -th row and j -th column.




Inductive definition of the determinant

Suppose $A_{n \times n}$ is given and we know how to define determinants for $n - 1 \times n - 1$ matrices. Define minors and cofactors as before. :

The (i, j) -th minor is the determinant of the submatrix formed by deleting the i -th row and j -th column.

The (i, j) -th cofactor $C_{ij} = (-1)^{i+j} M_{ij}$.



Suppose $A_{n \times n}$ is given and we know how to define determinants for $n-1 \times n-1$ matrices. Define minors and cofactors as before. :

The (i, j) -th minor is the determinant of the submatrix formed by deleting the i -th row and j -th column.

The (i, j) -th cofactor $C_{ij} = (-1)^{i+j} M_{ij}$.

$$\begin{aligned}\det(I_n) &= \det(I_{n-1}) \\ &= \det(I_{n-2}) \\ &= \dots = \det(I_3) \\ &= \det(I_2) \\ &= 1.\end{aligned}$$

Definition

$$\det(A) = \sum_{j=1}^n (-1)^{1+j} a_{1j} M_{1j} = \sum_{i=1}^n a_{1j} C_{1j}$$

$$\det(A) = a_{11} \times \det \begin{bmatrix} a_{22} & a_{23} & \dots & a_{2n} \\ a_{32} & a_{33} & \dots & a_{3n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n2} & a_{n3} & \dots & a_{nn} \end{bmatrix} - a_{12} \times \det \begin{bmatrix} a_{21} & a_{23} & \dots & a_{2n} \\ a_{31} & a_{33} & \dots & a_{3n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n3} & \dots & a_{nn} \end{bmatrix} + a_{13} \times \det \begin{bmatrix} a_{21} & a_{22} & \dots & a_{2n} \\ a_{31} & a_{32} & \dots & a_{3n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix} - a_{14} \times \det \begin{bmatrix} a_{21} & a_{22} & \dots & a_{2n} \\ a_{31} & a_{32} & \dots & a_{3n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$$

$$\begin{aligned}\det(I_{n \times n}) &= \det \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{pmatrix} \\ &= 1 \times \det \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{pmatrix} - 0 \times \det \begin{pmatrix} 0 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 \end{pmatrix} + 0 \times \det \begin{pmatrix} 0 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 \end{pmatrix} \\ &= 1 \times \det(I_{n-1})\end{aligned}$$

Expansion along any row or column



Expansion along any row or column

$$\det(A) = \sum_{j=1}^n (-1)^{i+j} a_{ij} M_{ij} \quad \text{for a fixed } i$$



$$\det(A) = \sum_{j=1}^n (-1)^{i+j} a_{ij} M_{ij} \quad \text{for a fixed } i$$

\leftarrow expansion along the i -th row

$$= \sum_{i=1}^n (-1)^{i+j} a_{ij} M_{ij}$$

\leftarrow for a fixed j

\leftarrow expansion along the j -th column

$\det(A_{2+3}) =$

$$= -a_{21} \times \det \begin{bmatrix} a_{12} & a_{13} \\ a_{32} & a_{33} \end{bmatrix} + a_{22} \times \det \begin{bmatrix} a_{11} & a_{13} \\ a_{31} & a_{33} \end{bmatrix} + (-1)^{2+3} a_{23} \times \det \begin{bmatrix} a_{11} & a_{12} \\ a_{31} & a_{32} \end{bmatrix}$$

$$= (-1)^{1+2} a_{12} \times \det \begin{bmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{bmatrix} + (-1)^{2+2} a_{22} \times \det \begin{bmatrix}] \\] \end{bmatrix} + (-1)^{2+3} a_{32} \times \det \begin{bmatrix}] \\] \end{bmatrix}$$

Important properties and identities

Property 1 : Determinant of a product is product of the determinants.



Property 1 : Determinant of a product is product of the determinants. Related identity : $\det(AB) = \det(A)\det(B)$

$$\det(A^n) = \det(A)^n$$

$$\det(A^{-1}) = \frac{1}{\det(A)} = \det(A)^{-1}$$

$$\det(P^T A P) = \det(A)$$

$$\det(AB) = \det(BA)$$

$$\det(AB) = \det(A)^2$$

$$\det(A^T A) = \det(A)^2$$

$\det(A^T)$
 || expand
 along 1st
 column
 + induction
 $\det(A)$

Property 2 : Switching two rows or columns changes the sign.

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

$$\tilde{A} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

$\leftarrow i^{\text{th}}$
 $\leftarrow j^{\text{th}}$
 $\leftarrow m^{\text{th}}$
 $\leftarrow n^{\text{th}}$

$$\det(\tilde{A}) = - \det(A)$$

↑
↑ Expand along i^{th} row
& use induction

Property 3 : Adding multiples of a row to another row leaves the determinant unchanged.

A

$$\det(\tilde{A}) = \det(A)$$

$$\tilde{A} = \begin{bmatrix} a_{11} + t_{1j_1} & a_{12} + t_{1j_2} & \dots & a_{1n} + t_{1j_n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$$

$$\tilde{A} = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} + t_{2j_1} & a_{22} + t_{2j_2} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} + t_{nj_1} & a_{n2} + t_{nj_2} & \dots & a_{nn} \end{bmatrix}$$

\downarrow
 k^{th} column

Important properties and identities

Property 3 : Adding multiples of a row to another row leaves the determinant unchanged.



Property 3' : Adding multiples of a column to another column leaves the determinant unchanged.

Important properties and identities

Property 4 : Scalar multiplication of a row by a constant t multiplies the determinant by t .



Property 4 : Scalar multiplication of a row by a constant t multiplies the determinant by t .

(A)

$$\det(\tilde{A}) = \sum_{j=1}^n (-1)^{i+j} \tilde{a}_{ij} \tilde{M}_{ij}$$

$$\tilde{A} = \begin{bmatrix} t a_{11} & t a_{12} & \dots & t a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$$


$$= t \left(\sum_{j=1}^n (-1)^{i+j} a_{ij} M_{ij} \right) = t \det(A).$$

Warning: $\det(tA_{mn}) = t^n \det(A)$

Property 4' : Scalar multiplication of a column by a constant t multiplies the determinant by t .


Useful computational tips

- 1) The determinant of a matrix with a row or column of zeros is 0.




Useful computational tips

- 1) The determinant of a matrix with a row or column of zeros is 0.
- 2) The determinant of a matrix in which one row (or column) is a linear combination of other rows (resp. columns) is 0.



Useful computational tips

- 1) The determinant of a matrix with a row or column of zeros is 0.
- 2) The determinant of a matrix in which one row (or column) is a linear combination of other rows (resp. columns) is 0.
- 3) Scalar multiplication of a row by a constant t multiplies the determinant by t .

- 
- 1) The determinant of a matrix with a row or column of zeros is 0.
 - 2) The determinant of a matrix in which one row (or column) is a linear combination of other rows (resp. columns) is 0.
 - 3) Scalar multiplication of a row by a constant t multiplies the determinant by t .
 - 4) While computing the determinant, you can choose to compute it using expansion along a suitable row or column.

Thank you

