



Tangents for scalar-valued multivariable functions



Recall : tangent lines to curves



Recall : tangent lines to curves


A **tangent line** to a curve C at a point p (on C) is a line which represents the *instantaneous* direction in which the curve C moves at the point p .



Recall : tangent lines to curves

A **tangent line** to a curve C at a point p (on C) is a line which represents the *instantaneous* direction in which the curve C moves at the point p .


Traditionally, it was thought of as a line which *just touches* the curve at that point.



Recall : tangent lines to curves

A **tangent line** to a curve C at a point p (on C) is a line which represents the *instantaneous* direction in which the curve C moves at the point p .


Traditionally, it was thought of as a line which *just touches* the curve at that point.



Recall : tangent lines to curves

A **tangent line** to a curve C at a point p (on C) is a line which represents the *instantaneous* direction in which the curve C moves at the point p .

Traditionally, it was thought of as a line which *just touches* the curve at that point.



Recall : tangent lines to curves


A **tangent line** to a curve C at a point p (on C) is a line which represents the *instantaneous* direction in which the curve C moves at the point p .

Traditionally, it was thought of as a line which *just touches* the curve at that point.




The tangent line to a function of one variable

Let $f : D \rightarrow \mathbb{R}$ be a function where D is a subset of \mathbb{R} .



The tangent line to a function of one variable

Let $f : D \rightarrow \mathbb{R}$ be a function where D is a subset of \mathbb{R} . Assume that $\Gamma(f)$, the graph of f is a curve. Let $x \in D$.



The tangent line to a function of one variable

Let $f : D \rightarrow \mathbb{R}$ be a function where D is a subset of \mathbb{R} . Assume that $\Gamma(f)$, the graph of f is a curve. Let $x \in D$.

Then a **tangent (line) to f at x** is a tangent (line) to $\Gamma(f)$ at the point $(x, f(x))$.



The tangent line to a function of one variable

Let $f : D \rightarrow \mathbb{R}$ be a function where D is a subset of \mathbb{R} . Assume that $\Gamma(f)$, the graph of f is a curve. Let $x \in D$.

Then a **tangent (line) to f at x** is a tangent (line) to $\Gamma(f)$ at the point $(x, f(x))$.




Figure: Tangent lines for $y = x^3$

Tangents for $f(x, y)$


Let $f(x, y)$ be a function defined on a domain D in \mathbb{R}^2 containing some open ball around the point \tilde{a} .



Tangents for $f(x, y)$

Let $f(x, y)$ be a function defined on a domain D in \mathbb{R}^2 containing some open ball around the point \tilde{a} .

Consider a line L in D passing through \tilde{a} and restrict f to L .



Tangents for $f(x, y)$

Let $f(x, y)$ be a function defined on a domain D in \mathbb{R}^2 containing some open ball around the point \tilde{a} .

Consider a line L in D passing through \tilde{a} and restrict f to L .

Since it is now a function of one variable, we can consider the tangent to f at \tilde{a} over L as in the previous slide.

Tangents for $f(x, y)$

Let $f(x, y)$ be a function defined on a domain D in \mathbb{R}^2 containing some open ball around the point \tilde{a} .

Consider a line L in D passing through \tilde{a} and restrict f to L .

Since it is now a function of one variable, we can consider the tangent to f at \tilde{a} over L as in the previous slide.

If the line L is in the direction of the unit vector u , then the tangent (if it exists) will be the line with slope $f_u(\tilde{a})$ and passing through the point $(\tilde{a}, f(\tilde{a}))$.

Tangents for $f(x, y)$

Let $f(x, y)$ be a function defined on a domain D in \mathbb{R}^2 containing some open ball around the point \tilde{a} .

Consider a line L in D passing through \tilde{a} and restrict f to L .

Since it is now a function of one variable, we can consider the tangent to f at \tilde{a} over L as in the previous slide.

If the line L is in the direction of the unit vector u , then the tangent (if it exists) will be the line with slope $f_u(\tilde{a})$ and passing through the point $(\tilde{a}, f(\tilde{a}))$.

The equations of the tangent line

$$\left. \begin{aligned} x(t) &= a + tu_1 \\ y(t) &= b + tu_2 \\ z(t) &= f(a, b) + t f_u(a, b). \end{aligned} \right\} \text{Parametric eqns.}$$

$$\frac{x-a}{u_1} = \frac{y-b}{u_2} = \frac{z-f(a,b)}{f_u(a,b)} \left. \right\} \text{symm. eqns.}$$

$$\begin{aligned} (x(t), y(t), z(t)) \\ = (a, b, f(a, b)) + t(u_1, u_2, f_u(a, b)) \end{aligned}$$

vector form.

$u = (u_1, u_2)$
 unit vector on L.
 $a = (a, b)$ is the point.
 $f_u(a, b)$ is the directional derivative at (a, b) .

$$\begin{aligned} L: z &= 0, \\ u_1(y-b) &= u_2(x-a). \\ P: u_1(y-b) &= u_2(x-a). \end{aligned}$$

$$\begin{aligned} x(t) &= a + t u_1 \\ y(t) &= b + t u_2 \\ z(t) &= 0. \\ (x(t), y(t), z(t)) \\ = (a, b, 0) + t(u_1, u_2, 0) \end{aligned}$$

Examples

$f(x, y) = x + y$; tangent at $(1, 1)$ in the direction of $(1, 0)$

$$f_u(1,1) = \frac{\partial f}{\partial x}(x,y) = 1.$$
$$(x(t), y(t), z(t)) = (1, 1, 2) + t(1, 0, 1).$$
$$x(t) = 1+t, y(t) = 1, z(t) = 2+t.$$

$f(x, y) = xy$; tangent at $(1, 1)$ in the direction of $(3, 4)$

$$u = \left(\frac{3}{5}, \frac{4}{5}\right).$$
$$f_u(1,1) = 1 \times \frac{3}{5} + 1 \times \frac{4}{5} = \frac{7}{5}.$$
$$\nabla f(x,y) = (y, x)$$
$$\nabla f(1,1) = (1, 1).$$
$$(x(t), y(t), z(t)) = (1, 1, 1) + t\left(\frac{3}{5}, \frac{4}{5}, \frac{7}{5}\right)$$
$$= \left(1 + \frac{3t}{5}, 1 + \frac{4t}{5}, 1 + \frac{7t}{5}\right).$$

$f(x, y) = \sin(xy)$; tangent at $(\pi, 1)$ in the direction of $(1, 2)$

$$u = \frac{1}{\sqrt{5}}(1, 2).$$
$$\nabla f(x,y) = (y \cos(xy), x \cos(xy)). \therefore \nabla f(\pi, 1) = (-1, -\pi).$$
$$f_u(\pi, 1) = (-1) \times \frac{1}{\sqrt{5}} + (-\pi) \times \frac{2}{\sqrt{5}} = \frac{-2\pi - 1}{\sqrt{5}}.$$

$$(x(t), y(t), z(t)) = (\pi, 1, 0) + t\left(\frac{1}{\sqrt{5}}, \frac{2}{\sqrt{5}}, -\frac{2\pi + 1}{\sqrt{5}}\right).$$

Tangents for scalar-valued multivariable functions


Let $\tilde{f}(\tilde{x})$ be a function defined on a domain D in \mathbb{R}^n containing some open ball around the point \tilde{a} .



Tangents for scalar-valued multivariable functions

Let $\tilde{f}(\tilde{x})$ be a function defined on a domain D in \mathbb{R}^n containing some open ball around the point \tilde{a} .

Consider a line L in D passing through \tilde{a} and restrict f to L .



Tangents for scalar-valued multivariable functions

Let $\tilde{f}(\tilde{x})$ be a function defined on a domain D in \mathbb{R}^n containing some open ball around the point \tilde{a} .

Consider a line L in D passing through \tilde{a} and restrict f to L .

Since it is now a function of one variable, we can consider the tangent to f at \tilde{a} over L as before.

Tangents for scalar-valued multivariable functions

Let $f(\tilde{x})$ be a function defined on a domain D in \mathbb{R}^n containing some open ball around the point \tilde{a} .

Consider a line L in D passing through \tilde{a} and restrict f to L .

Since it is now a function of one variable, we can consider the tangent to f at \tilde{a} over L as before.

If the line L is in the direction of the unit vector u , then the tangent (if it exists) will be the line with slope $f_u(\tilde{a})$ and passing through the point $(\tilde{a}, f(\tilde{a}))$.

Parametric equations and an example

Similar to the two-variable case, we can deduce the equations of the tangent line as :

$\tilde{x} = (x_1, x_2, \dots, x_n)$, \tilde{z} is the variable in which we are measuring the fn.

$\tilde{a} = (a_1, \dots, a_n)$, $u = (u_1, \dots, u_n)$.
Line through \tilde{a} in the direction of u is $x_i(t) = a_i + t u_i$; $\tilde{z} = 0$.
 $(\tilde{x}(t), \tilde{z}(t)) = (\tilde{a}, 0) + t(u, 0)$.

\therefore The tangent line to f at \tilde{a} above 2 is $(\tilde{x}(t), \tilde{z}(t)) = (\tilde{a}, f(\tilde{a})) + t(u, f_u(\tilde{a}))$.
 $x_i(t) = a_i + t u_i$, $\tilde{z}(t) = f(\tilde{a}) + t f_u(\tilde{a})$.

Example : $f(x, y) = xy + yz + zx$; tangent at $(1, 1, 1)$ in the direction $(-1, -2, 2)$

$$u = \frac{1}{3}(-1, -2, 2). \quad \nabla f(x, y, z) = (y+z, z+x, x+y); \quad \nabla f(1, 1, 1) = (2, 2, 2).$$

$$f_u(1, 1, 1) = -2/3$$

$$(\tilde{x}(t), \tilde{y}(t), \tilde{z}(t), u(t)) = (1, 1, 1, 3) + t\left(\frac{-1}{3}, \frac{-2}{3}, \frac{2}{3}, -\frac{2}{3}\right)$$

$$\tilde{x}(t) = 1 - t/3, \quad \tilde{y}(t) = 1 - 2t/3, \quad \tilde{z}(t) = 1 + 2t/3, \quad u(t) = 3^{-2+t}/5.$$

Caution : tangents need not always exist.

$$f(x, y) = \begin{cases} \frac{xy}{x^2+y^2} & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0) \end{cases}$$

$$f_u(0, 0) = \begin{cases} 0 & \text{if } (u_1, u_2) = \pm e_1 \text{ or } \pm e_2 \\ \text{DNE} & \text{o.w.} \end{cases}$$


The tangent lines in all directions other than along the x or y-axes at $(0, 0)$ DNE.

$$f(x, y) = |x| + |y|$$

For many directions at many points, the tangent line will not exist.

When do all the tangents exist?


This is equivalent to asking when do all the directional derivatives exist.



When do all the tangents exist?

This is equivalent to asking when do all the directional derivatives exist.

Let $f(x_1, x_2, \dots, x_n)$ be a function defined on a domain D in \mathbb{R}^n containing some open ball around the point a .



When do all the tangents exist?

This is equivalent to asking when do all the directional derivatives exist.

Let $f(x_1, x_2, \dots, x_n)$ be a function defined on a domain D in \mathbb{R}^n containing some open ball around the point a .

Theorem

Suppose ∇f exists and is continuous on some open ball around the point a .

When do all the tangents exist?

This is equivalent to asking when do all the directional derivatives exist.

Let $f(x_1, x_2, \dots, x_n)$ be a function defined on a domain D in \mathbb{R}^n containing some open ball around the point a .

Theorem

Suppose ∇f exists and is continuous on some open ball around the point \tilde{a} . Then for every unit vector u , the directional derivative $f_u(\tilde{a})$ exists and equals $\nabla f(\tilde{a}) \cdot u$.

When do all the tangents exist?

This is equivalent to asking when do all the directional derivatives exist.

Let $f(x_1, x_2, \dots, x_n)$ be a function defined on a domain D in \mathbb{R}^n containing some open ball around the point a .

Theorem

Suppose ∇f exists and is continuous on some open ball around the point \tilde{a} . Then for every unit vector u , the directional derivative $f_u(\tilde{a})$ exists and equals $\nabla f(\tilde{a}) \cdot u$.

Conclusion : All the tangents at a point \tilde{a} exist when ∇f exists and is continuous on some open ball around the point \tilde{a} .

Thank you

