

The Gaussian Elimination Method

Sarang S. Sane

Recall

We have seen the following methods to find the solutions to a system of linear equations $Ax = b$:

Recall

We have seen the following methods to find the solutions to a system of linear equations $Ax = b$:

- ▶ If A is invertible, then the solution is unique and is given by $A^{-1}b$. The solution can be found by using :

Recall

We have seen the following methods to find the solutions to a system of linear equations $Ax = b$:

- ▶ If A is invertible, then the solution is unique and is given by $A^{-1}b$. The solution can be found by using :
 1. Cramer's rule.

Recall

We have seen the following methods to find the solutions to a system of linear equations $Ax = b$:

- ▶ If A is invertible, then the solution is unique and is given by $A^{-1}b$. The solution can be found by using :
 1. Cramer's rule.
 2. the adjugate matrix to calculate A^{-1} .

Recall

We have seen the following methods to find the solutions to a system of linear equations $Ax = b$:

- ▶ If A is invertible, then the solution is unique and is given by $A^{-1}b$. The solution can be found by using :
 1. Cramer's rule.
 2. the adjugate matrix to calculate A^{-1} .

- ▶ If A is in (reduced) row echelon form, we can find all the solutions as follows :
 1. Find the dependent variables (corr. to columns with leading entries) and independent variables (corr. to other columns).

Recall

We have seen the following methods to find the solutions to a system of linear equations $Ax = b$:

- ▶ If A is invertible, then the solution is unique and is given by $A^{-1}b$. The solution can be found by using :
 1. Cramer's rule.
 2. the adjugate matrix to calculate A^{-1} .

- ▶ If A is in (reduced) row echelon form, we can find all the solutions as follows :
 1. Find the dependent variables (corr. to columns with leading entries) and independent variables (corr. to other columns).
 2. Assign a value to each independent variable. Calculate the values of each dependent variable using the unique equation in which it occurs.

Contents

Contents

- ▶ The augmented matrix for a system of linear equations.

Contents

- ▶ The augmented matrix for a system of linear equations.
- ▶ The Gaussian elimination method to determine all solutions of a system of linear equations.

Contents

- ▶ The augmented matrix for a system of linear equations.
- ▶ The Gaussian elimination method to determine all solutions of a system of linear equations.
- ▶ Computing the inverse using Gaussian elimination.

The augmented matrix

Let $Ax = b$ be a system of linear equations where A is an $m \times n$ matrix and b is a $m \times 1$ column vector.

The augmented matrix

Let $Ax = b$ be a system of linear equations where A is an $m \times n$ matrix and b is a $m \times 1$ column vector.

The augmented matrix of this system is defined as the matrix of size $m \times (n + 1)$ whose first n columns are the columns of A and the last column is b .

The augmented matrix

Let $Ax = b$ be a system of linear equations where A is an $m \times n$ matrix and b is a $m \times 1$ column vector.

The augmented matrix of this system is defined as the matrix of size $m \times (n + 1)$ whose first n columns are the columns of A and the last column is b .

We denote the augmented matrix by $[A|b]$ and put a vertical line between the first n columns and the last column b while writing it.

$$\left[\begin{array}{cccc} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{array} \right] \left| \begin{array}{c} b_1 \\ b_2 \\ \vdots \\ b_m \end{array} \right]$$

A

The augmented matrix

Let $Ax = b$ be a system of linear equations where A is an $m \times n$ matrix and b is a $m \times 1$ column vector.

The augmented matrix of this system is defined as the matrix of size $m \times n + 1$ whose first n columns are the columns of A and the last column is b .

We denote the augmented matrix by $[A|b]$ and put a vertical line between the first n columns and the last column b while writing it.

Example

$$3x_1 + 2x_2 + x_3 + x_4 = 6$$

$$x_1 + x_2 = 2$$

$$7x_2 + x_3 + x_4 = 8$$

Example

$$3x_1 + 2x_2 + x_3 + x_4 = 6$$

$$x_1 + x_2 = 2$$

$$7x_2 + x_3 + x_4 = 8$$

where $A = \begin{bmatrix} 3 & 2 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 7 & 1 & 1 \end{bmatrix}$ and $b = \begin{bmatrix} 6 \\ 2 \\ 8 \end{bmatrix}$.

Example

$$3x_1 + 2x_2 + x_3 + x_4 = 6$$

$$x_1 + x_2 = 2$$

$$7x_2 + x_3 + x_4 = 8$$

$$\text{where } A = \begin{bmatrix} 3 & 2 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 7 & 1 & 1 \end{bmatrix} \quad \text{and} \quad b = \begin{bmatrix} 6 \\ 2 \\ 8 \end{bmatrix}.$$

$$\text{The augmented matrix is } [A|b] = \left[\begin{array}{cccc|c} 3 & 2 & 1 & 1 & 6 \\ 1 & 1 & 0 & 0 & 2 \\ 0 & 7 & 1 & 1 & 8 \end{array} \right].$$

The Gaussian elimination method

Consider the system of linear equations $Ax = b$.

The Gaussian elimination method

Consider the system of linear equations $Ax = b$.

1. Form the augmented matrix of the system $[A|b]$.

The Gaussian elimination method

Consider the system of linear equations $Ax = b$.

1. Form the augmented matrix of the system $[A|b]$.
2. Perform the same operations on $[A|b]$ that were used to bring A into reduced row echelon form.

The Gaussian elimination method

Consider the system of linear equations $Ax = b$.

1. Form the augmented matrix of the system $[A|b]$.
2. Perform the same operations on $[A|b]$ that were used to bring A into reduced row echelon form.
3. Let R be the submatrix of the obtained matrix of the first n columns and c be the submatrix of the obtained matrix consisting of the last column.

The Gaussian elimination method

Consider the system of linear equations $Ax = b$.

1. Form the augmented matrix of the system $[A|b]$.
2. Perform the same operations on $[A|b]$ that were used to bring A into reduced row echelon form.
3. Let R be the submatrix of the obtained matrix of the first n columns and c be the submatrix of the obtained matrix consisting of the last column.

We write the obtained matrix as $[R|c]$. Notice that R is the reduced row echelon matrix obtained by row reducing A .

The Gaussian elimination method

Consider the system of linear equations $Ax = b$.

1. Form the augmented matrix of the system $[A|b]$.
2. Perform the same operations on $[A|b]$ that were used to bring A into reduced row echelon form.
3. Let R be the submatrix of the obtained matrix of the first n columns and c be the submatrix of the obtained matrix consisting of the last column.

We write the obtained matrix as $[R|c]$. Notice that R is the reduced row echelon matrix obtained by row reducing A .

The solutions of $Ax = b$ are precisely the solutions of $Rx = c$.

The Gaussian elimination method

4. Form the corresponding system of linear equations $Rx = c$.

The Gaussian elimination method

4. Form the corresponding system of linear equations $Rx = c$.
5. Find ALL the solutions of $Rx = c$ and hence of $Ax = b$.

The Gaussian elimination method

4. Form the corresponding system of linear equations $Rx = c$.
5. Find ALL the solutions of $Rx = c$ and hence of $Ax = b$.

Since R is in reduced row echelon form, we can find ALL its solutions (as described earlier).

Example

$$\left[\begin{array}{cccc|c} 3 & 2 & 1 & 1 & 6 \\ 1 & 1 & 0 & 0 & 2 \\ 0 & 7 & 1 & 1 & 8 \end{array} \right]$$

Example

$$\left[\begin{array}{cccc|c} 3 & 2 & 1 & 1 & 6 \\ 1 & 1 & 0 & 0 & 2 \\ 0 & 7 & 1 & 1 & 8 \end{array} \right] \xrightarrow[\sim]{R_1/3}$$

$$\left[\begin{array}{cccc|c} 1 & 2/3 & 1/3 & 1/3 & 2 \\ 1 & 1 & 0 & 0 & 2 \\ 0 & 7 & 1 & 1 & 8 \end{array} \right]$$

Example

$$\left[\begin{array}{cccc|c} 3 & 2 & 1 & 1 & 6 \\ 1 & 1 & 0 & 0 & 2 \\ 0 & 7 & 1 & 1 & 8 \end{array} \right] \xrightarrow{R_1/3}$$

$$\left[\begin{array}{cccc|c} 1 & 2/3 & 1/3 & 1/3 & 2 \\ 1 & 1 & 0 & 0 & 2 \\ 0 & 7 & 1 & 1 & 8 \end{array} \right]$$

$$\left. \vphantom{\left[\begin{array}{cccc|c} 1 & 2/3 & 1/3 & 1/3 & 2 \\ 1 & 1 & 0 & 0 & 2 \\ 0 & 7 & 1 & 1 & 8 \end{array} \right]} \right\} R_2 - R_1$$

$$\left[\begin{array}{cccc|c} 1 & 2/3 & 1/3 & 1/3 & 2 \\ 0 & 1/3 & -1/3 & -1/3 & 0 \\ 0 & 7 & 1 & 1 & 8 \end{array} \right]$$

Example

$$\left[\begin{array}{cccc|c} 3 & 2 & 1 & 1 & 6 \\ 1 & 1 & 0 & 0 & 2 \\ 0 & 7 & 1 & 1 & 8 \end{array} \right] \xrightarrow{R_1/3}$$

$$\left[\begin{array}{cccc|c} 1 & 2/3 & 1/3 & 1/3 & 2 \\ 1 & 1 & 0 & 0 & 2 \\ 0 & 7 & 1 & 1 & 8 \end{array} \right]$$

$$\xrightarrow{R_2 - R_1}$$

$$\left[\begin{array}{cccc|c} 1 & 2/3 & 1/3 & 1/3 & 2 \\ 0 & 1 & -1 & -1 & 0 \\ 0 & 7 & 1 & 1 & 8 \end{array} \right] \xleftarrow{3R_2}$$

$$\left[\begin{array}{cccc|c} 1 & 2/3 & 1/3 & 1/3 & 2 \\ 0 & 1/3 & -1/3 & -1/3 & 0 \\ 0 & 7 & 1 & 1 & 8 \end{array} \right]$$

Example

$$\left[\begin{array}{cccc|c} 3 & 2 & 1 & 1 & 6 \\ 1 & 1 & 0 & 0 & 2 \\ 0 & 7 & 1 & 1 & 8 \end{array} \right] \xrightarrow{R_1/3}$$

$$\left[\begin{array}{cccc|c} 1 & 2/3 & 1/3 & 1/3 & 2 \\ 1 & 1 & 0 & 0 & 2 \\ 0 & 7 & 1 & 1 & 8 \end{array} \right]$$

$$\xrightarrow{R_2 - R_1}$$

$$\left[\begin{array}{cccc|c} 1 & 2/3 & 1/3 & 1/3 & 2 \\ 0 & 1 & -1 & -1 & 0 \\ 0 & 7 & 1 & 1 & 8 \end{array} \right] \xleftarrow{3R_2}$$

$$\left[\begin{array}{cccc|c} 1 & 2/3 & 1/3 & 1/3 & 2 \\ 0 & 1 & -1/3 & -1/3 & 0 \\ 0 & 7 & 1 & 1 & 8 \end{array} \right]$$

$$\xrightarrow{R_3 - 7R_2}$$

$$\left[\begin{array}{cccc|c} 1 & 2/3 & 1/3 & 1/3 & 2 \\ 0 & 1 & -1 & -1 & 0 \\ 0 & 0 & 8 & 8 & 8 \end{array} \right]$$

Example

$$\left[\begin{array}{cccc|c} 3 & 2 & 1 & 1 & 6 \\ 1 & 1 & 0 & 0 & 2 \\ 0 & 7 & 1 & 1 & 8 \end{array} \right] \xrightarrow{R_1/3}$$

$$\left[\begin{array}{cccc|c} 1 & 2/3 & 1/3 & 1/3 & 2 \\ 1 & 1 & 0 & 0 & 2 \\ 0 & 7 & 1 & 1 & 8 \end{array} \right]$$

$$\left. \vphantom{\begin{array}{c} \\ \\ \end{array}} \right\} R_2 - R_1$$

$$\left[\begin{array}{cccc|c} 1 & 2/3 & 1/3 & 1/3 & 2 \\ 0 & 1 & -1 & -1 & 0 \\ 0 & 7 & 1 & 1 & 8 \end{array} \right] \xleftarrow{3R_2}$$

$$\left[\begin{array}{cccc|c} 1 & 2/3 & 1/3 & 1/3 & 2 \\ 0 & 1 & -1/3 & -1/3 & 0 \\ 0 & 7 & 1 & 1 & 8 \end{array} \right]$$

$$\left. \vphantom{\begin{array}{c} \\ \\ \end{array}} \right\} R_3 - 7R_2$$

$$\left[\begin{array}{cccc|c} 1 & 2/3 & 1/3 & 1/3 & 2 \\ 0 & 1 & -1 & -1 & 0 \\ 0 & 0 & 8 & 8 & 8 \end{array} \right] \xrightarrow{R_3/8}$$

$$\left[\begin{array}{cccc|c} 1 & 2/3 & 1/3 & 1/3 & 2 \\ 0 & 1 & -1 & -1 & 0 \\ 0 & 0 & 1 & 1 & 1 \end{array} \right]$$

Example (contd.)

$$\left[\begin{array}{cccc|c} 1 & 2/3 & 1/3 & 1/3 & 2 \\ 0 & 1 & -1 & -1 & 0 \\ 0 & 0 & 1 & 1 & 1 \end{array} \right]$$

$$R_2 + R_3$$

$$R_1 - \frac{1}{3}R_3$$

$$\left[\begin{array}{cccc|c} 1 & 2/3 & 0 & 0 & 5/3 \\ 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 & 1 \end{array} \right]$$

$$\left. \vphantom{\left[\begin{array}{cccc|c} 1 & 2/3 & 0 & 0 & 5/3 \\ 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 & 1 \end{array} \right]} \right\} R_1 - \frac{2}{3}R_2$$

$$Rx = c$$

$x_1, x_2, x_3 \rightarrow$ dependent

$x_4 \rightarrow$ independent

$$x_4 = c$$

$$x_1 = 1$$

$$x_2 = 1$$

$$x_3 + x_4 = 1$$

$$\Rightarrow x_3 = 1 - c$$

$$\left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 & 1 \end{array} \right]$$

Set of solns. of $Rx = c$ & hence $Ax = b$

$$\left\{ \begin{array}{l} (x_1=1, x_2=1, \\ x_3=1-c, x_4=c) \end{array} \mid c \in \mathbb{R} \right\}$$

$$= \left\{ \begin{bmatrix} 1 \\ 1 \\ 1-c \\ c \end{bmatrix} \mid c \in \mathbb{R} \right\}$$

Another example

$$x_1 + x_2 + x_3 = 2$$

$$x_2 - 3x_3 = 1$$

$$2x_1 + x_2 + 5x_3 = 0$$

Another example

$$x_1 + x_2 + x_3 = 2$$

$$x_2 - 3x_3 = 1$$

$$2x_1 + x_2 + 5x_3 = 0$$

The matrix representation of this system of linear equations is:

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & -3 \\ 2 & 1 & 5 \end{bmatrix} \quad x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \quad b = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}$$

Another example

$$x_1 + x_2 + x_3 = 2$$

$$x_2 - 3x_3 = 1$$

$$2x_1 + x_2 + 5x_3 = 0$$

The matrix representation of this system of linear equations is:

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & -3 \\ 2 & 1 & 5 \end{bmatrix} \quad x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \quad b = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}$$

The augmented matrix is $\left[\begin{array}{ccc|c} 1 & 1 & 1 & 2 \\ 0 & 1 & -3 & 1 \\ 2 & 1 & 5 & 0 \end{array} \right]$

Another example (contd.)

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 2 \\ 0 & 1 & -3 & 1 \\ 2 & 1 & 5 & 0 \end{array} \right]$$

Another example (contd.)

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 2 \\ 0 & 1 & -3 & 1 \\ 2 & 1 & 5 & 0 \end{array} \right] \xrightarrow{\widetilde{R_3 - 2R_1}} \left[\begin{array}{ccc|c} 1 & 1 & 1 & 2 \\ 0 & 1 & -3 & 1 \\ 0 & -1 & 3 & -4 \end{array} \right]$$

Another example (contd.)

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 2 \\ 0 & 1 & 1 & 1 \\ 2 & 1 & 5 & 0 \end{array} \right]$$

$\xrightarrow{R_3 - 2R_1}$

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 2 \\ 0 & 1 & -3 & 1 \\ 0 & -1 & 3 & -4 \end{array} \right]$$

$\xrightarrow{R_3 + R_2}$

$$\left[\begin{array}{ccc|c} 1 & \cancel{1} & 1 & 1 \\ 0 & 1 & -3 & 1 \\ 0 & 0 & 0 & -3 \end{array} \right]$$

$\downarrow R_1 - R_2$

$$\left[\begin{array}{ccc|c} 1 & 0 & 4 & 0 \\ 0 & 1 & -3 & 1 \\ 0 & 0 & 0 & -3 \end{array} \right]$$

$\underbrace{\hspace{10em}}_R \quad \underbrace{\hspace{10em}}_c$

$Rx = c.$

This system does not have solutions.
 $\therefore Ax = b$ does not have solutions

Homogeneous system of linear equations

Homogeneous system of linear equations

0 is always a solution of a homogeneous system of linear equations $Ax = 0$. This solution is called the *trivial solution*.

Homogeneous system of linear equations

0 is always a solution of a homogeneous system of linear equations $Ax = 0$. This solution is called the *trivial solution*.

For a homogeneous system, there are only two different possibilities :

Homogeneous system of linear equations

0 is always a solution of a homogeneous system of linear equations $Ax = 0$. This solution is called the *trivial solution*.

For a homogeneous system, there are only two different possibilities :

- ▶ 0 is the unique solution.

Homogeneous system of linear equations

0 is always a solution of a homogeneous system of linear equations $Ax = 0$. This solution is called the *trivial solution*.

For a homogeneous system, there are only two different possibilities :

- ▶ 0 is the unique solution.
- ▶ there are infinitely many solutions other than 0.

$$\begin{aligned} Ax &= 0 \\ x_1 &= w_1, x_2 = w_2, \\ &\dots, x_n = w_n \\ &\text{is a soln.} \\ \text{then } & \text{so is} \\ x_1 &= tw_1, x_2 = tw_2, \\ &\dots, x_n = tw_n. \end{aligned}$$


Homogeneous system of linear equations

0 is always a solution of a homogeneous system of linear equations $Ax = 0$. This solution is called the *trivial solution*.

For a homogeneous system, there are only two different possibilities :

- ▶ 0 is the unique solution
- ▶ there are infinitely many solutions other than 0.

In a homogeneous system of equations, if there are more variables than equations, then it is guaranteed to have nontrivial solutions.



Computing the inverse

Computing the inverse of an invertible matrix A is equivalent to :

Computing the inverse

Computing the inverse of an invertible matrix A is equivalent to :

Finding solutions of $Ax = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$, $Ay = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$ and $Az = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$.

$$\left[A \mid \begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right]$$

$$[A|I]$$

reduction to reduced row echelon form

$$[I \mid A^{-1}]$$

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 2 & 4 & 8 & 0 & 1 & 0 \\ 3 & 9 & 27 & 0 & 0 & 1 \end{array} \right]$$

$R_2 - 2R_1$
 $R_3 - 3R_1$

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 2 & 6 & -2 & 1 & 0 \\ 0 & 6 & 24 & -3 & 0 & 1 \end{array} \right]$$

$R_2/2$

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 3 & -1 & 1/2 & 0 \\ 0 & 6 & 24 & -3 & 0 & 1 \end{array} \right]$$

$R_3 - 6R_2$

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 3 & -1 & 1/2 & 0 \\ 0 & 0 & 6 & 3 & -3 & 1 \end{array} \right]$$

$R_3/6$

$$\left[\begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 3 & -1 & 1/2 & 0 \\ 0 & 0 & 1 & 1/2 & -1/2 & 1/6 \end{array} \right]$$

$$A \begin{bmatrix} x' & y' & z' \end{bmatrix}$$

$$\begin{bmatrix} Ax' & Ay' & Az' \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I$$

Thank you