


What is the rank/dimension of a vector space?



# Basis for a vector space (recall)



## Basis for a vector space (recall)

A basis  $B$  of a vector space  $V$  is a linearly independent subset of  $V$  that spans  $V$ .



## Basis for a vector space (recall)

A basis  $B$  of a vector space  $V$  is a linearly independent subset of  $V$  that spans  $V$ .

Equivalently : A basis  $B$  of a vector space  $V$  is a subset  $B \subseteq V$  such that every element of  $V$  can be **uniquely** written as a linear combination of elements of  $B$ .

## Basis for a vector space (recall)

A basis  $B$  of a vector space  $V$  is a linearly independent subset of  $V$  that spans  $V$ .

Equivalently : A basis  $B$  of a vector space  $V$  is a subset  $B \subseteq V$  such that every element of  $V$  can be uniquely written as a linear combination of elements of  $B$ .

i.e. if  $B = \{v_1, v_2, \dots, v_n\}$  then for every  $v \in V$ , there is a unique set of scalars  $a_1, a_2, \dots, a_n \in \mathbb{R}$  such that  $v = \sum_{i=1}^n a_i v_i$ .

# What is the rank/dimension of a vector space



# What is the rank/dimension of a vector space


The dimension (or rank) of a vector space is the **size (or cardinality) of a basis of the vector space.**



# What is the rank/dimension of a vector space

The dimension (or rank) of a vector space is the **size (or cardinality) of a basis of the vector space.**

for this course : if  $B$  is a basis of  $V$ , then the rank is the number of elements in  $B$ .




# What is the rank/dimension of a vector space

The dimension (or rank) of a vector space is the **size (or cardinality) of a basis of the vector space.**

for this course : if  $B$  is a basis of  $V$ , then the rank is the number of elements in  $B$ .

For every vector space there exists a basis, and all bases of a vector space have the same number of elements (or cardinality) ;



# What is the rank/dimension of a vector space

The dimension (or rank) of a vector space is the **size (or cardinality) of a basis of the vector space.**


for this course : if  $B$  is a basis of  $V$ , then the rank is the number of elements in  $B$ .

For every vector space there exists a basis, and all bases of a vector space have the same number of elements (or cardinality) ; hence, the dimension (or rank) of a vector space (say  $V$ ) is uniquely defined and denoted by  $\dim(V)$  (or  $\text{rank}(V)$ ) respectively.

# Dimension of $\mathbb{R}^n$

Recall the  $i^{th}$  **standard basis vector** in  $\mathbb{R}^n$ .

$$e_i = (0, 0, \dots, 0, 1, 0, \dots, 0)$$




# Dimension of $\mathbb{R}^n$

Recall the  $i^{th}$  **standard basis vector** in  $\mathbb{R}^n$ .

$$e_i = (0, 0, \dots, 0, 1, 0, \dots, 0)$$

i.e. the  $i$ -th co-ordinate is 1 and 0 elsewhere.



# Dimension of $\mathbb{R}^n$

Recall the  $i^{th}$  **standard basis vector** in  $\mathbb{R}^n$ .

$$e_i = (0, 0, \dots, 0, 1, 0, \dots, 0)$$

i.e. the  $i$ -th co-ordinate is 1 and 0 elsewhere.

Recall that the set  $\{e_1, e_2, \dots, e_n\}$  is a basis of  $\mathbb{R}^n$  called the **standard basis**.

# Dimension of $\mathbb{R}^n$

Recall the  $i^{th}$  **standard basis vector** in  $\mathbb{R}^n$ .

$$e_i = (0, 0, \dots, 0, 1, 0, \dots, 0)$$

i.e. the  $i$ -th co-ordinate is 1 and 0 elsewhere.

Recall that the set  $\{e_1, e_2, \dots, e_n\}$  is a basis of  $\mathbb{R}^n$  called the **standard basis**.

Hence the dimension of  $\mathbb{R}^n$  is  $n$ .

## Example


Let us calculate the dimension of the subspace  $W$  of  $\mathbb{R}^3$  spanned by  $\{(1, 0, 0), (0, 1, 0), (3, 5, 0)\}$ .



## Example

Let us calculate the dimension of the subspace  $W$  of  $\mathbb{R}^3$  spanned by  $\{(1, 0, 0), (0, 1, 0), (3, 5, 0)\}$ .

Observe that,  $3(1, 0, 0) + 5(0, 1, 0) = (3, 5, 0)$ .




## Example

Let us calculate the dimension of the subspace  $W$  of  $\mathbb{R}^3$  spanned by  $\{(1, 0, 0), (0, 1, 0), (3, 5, 0)\}$ .

Observe that,  $3(1, 0, 0) + 5(0, 1, 0) = (3, 5, 0)$ .

Hence the set is not linearly independent.



## Example

Let us calculate the dimension of the subspace  $W$  of  $\mathbb{R}^3$  spanned by  $\{(1, 0, 0), (0, 1, 0), (3, 5, 0)\}$ .

Observe that,  $3(1, 0, 0) + 5(0, 1, 0) = (3, 5, 0)$ .

Hence the set is not linearly independent.

Hence we delete the vector  $(3, 5, 0)$  from this set.

## Example

Let us calculate the dimension of the subspace  $W$  of  $\mathbb{R}^3$  spanned by  $\{(1, 0, 0), (0, 1, 0), (3, 5, 0)\}$ .

Observe that,  $3(1, 0, 0) + 5(0, 1, 0) = (3, 5, 0)$ .

Hence the set is not linearly independent.

Hence we delete the vector  $(3, 5, 0)$  from this set.

The remaining two vectors form a linearly independent set.

## Example

Let us calculate the dimension of the subspace  $W$  of  $\mathbb{R}^3$  spanned by  $\{(1, 0, 0), (0, 1, 0), (3, 5, 0)\}$ .

Observe that,  $3(1, 0, 0) + 5(0, 1, 0) = (3, 5, 0)$ .

Hence the set is not linearly independent.

Hence we delete the vector  $(3, 5, 0)$  from this set.

The remaining two vectors form a linearly independent set.

Hence the set  $\{(1, 0, 0), (0, 1, 0)\}$  forms a basis of the subspace  $W$  spanned by  $\{(1, 0, 0), (0, 1, 0), (3, 5, 0)\}$ .

## Example

Let us calculate the dimension of the subspace  $W$  of  $\mathbb{R}^3$  spanned by  $\{(1, 0, 0), (0, 1, 0), (3, 5, 0)\}$ .

Observe that,  $3(1, 0, 0) + 5(0, 1, 0) = (3, 5, 0)$ .

Hence the set is not linearly independent.

Hence we delete the vector  $(3, 5, 0)$  from this set.


The remaining two vectors form a linearly independent set.

Hence the set  $\{(1, 0, 0), (0, 1, 0)\}$  forms a basis of the subspace  $W$  spanned by  $\{(1, 0, 0), (0, 1, 0), (3, 5, 0)\}$ .


Hence dimension of the subspace  $W$  is 2.

## Example contd.

Geometrically the subspace  $W$  is the  $XY$ -plane.



## Example : in terms of matrices



## Example : in terms of matrices


Write the vectors which span (or generate)  $W$  as rows of a matrix :



## Example : in terms of matrices

Write the vectors which span (or generate)  $W$  as rows of a matrix :

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 5 & 0 \end{bmatrix}$$



## Example : in terms of matrices

Write the vectors which span (or generate)  $W$  as rows of a matrix :

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 5 & 0 \end{bmatrix}$$

Apply row reduction to this matrix.

## Example : in terms of matrices

Write the vectors which span (or generate)  $W$  as rows of a matrix :

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 5 & 0 \end{bmatrix}.$$

Apply row reduction to this matrix.

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 5 & 0 \end{bmatrix} \xrightarrow{R_3 - 3R_1} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 5 & 0 \end{bmatrix} \xrightarrow{R_3 - 5R_2} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

## Example : in terms of matrices

Write the vectors which span (or generate)  $W$  as rows of a matrix :

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 5 & 0 \end{bmatrix}$$

Apply row reduction to this matrix.

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 5 & 0 \end{bmatrix} \xrightarrow{R_3 - 3R_1} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 5 & 0 \end{bmatrix} \xrightarrow{R_3 - 5R_2} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

The final matrix is the row echelon form of the original matrix and its rows form a basis of the subspace  $W$ .

## Example : in terms of matrices

Write the vectors which span (or generate)  $W$  as rows of a matrix :

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 5 & 0 \end{bmatrix}.$$

Apply row reduction to this matrix.

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 5 & 0 \end{bmatrix}$$

$$R_3 - 3R_1$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 5 & 0 \end{bmatrix}$$

$$R_3 - 5R_2$$


$$\boxed{\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}}$$

The final matrix is the row echelon form of the original matrix and its rows form a basis of the subspace  $W$ .

In particular, the number of non-zero rows is  $2 = \dim(W)$ .

# Rank of a matrix


Let  $A$  be an  $m \times n$  matrix.



# Rank of a matrix

Let  $A$  be an  $m \times n$  matrix.


- ▶ The **column space** of  $A$  is the subspace of  $\mathbb{R}^m$  spanned by the column vectors of  $A$ .



# Rank of a matrix

Let  $A$  be an  $m \times n$  matrix.

- ▶ The **column space** of  $A$  is the subspace of  $\mathbb{R}^m$  spanned by the column vectors of  $A$ .
- ▶ The **row space** of  $A$  is the subspace of  $\mathbb{R}^n$  spanned by the row vectors of  $A$ .



# Rank of a matrix

Let  $A$  be an  $m \times n$  matrix.

- ▶ The **column space** of  $A$  is the subspace of  $\mathbb{R}^m$  spanned by the column vectors of  $A$ .
- ▶ The **row space** of  $A$  is the subspace of  $\mathbb{R}^n$  spanned by the row vectors of  $A$ .
- ▶ The dimension of the column space of  $A$  is defined as the **column rank** of  $A$ .

# Rank of a matrix

Let  $A$  be an  $m \times n$  matrix.

- ▶ The **column space** of  $A$  is the subspace of  $\mathbb{R}^m$  spanned by the column vectors of  $A$ .
- ▶ The **row space** of  $A$  is the subspace of  $\mathbb{R}^n$  spanned by the row vectors of  $A$ .
- ▶ The dimension of the column space of  $A$  is defined as the **column rank** of  $A$ .
- ▶ The dimension of row space of  $A$  is defined as the **row rank** of  $A$ .

# Rank of a matrix

Let  $A$  be an  $m \times n$  matrix.

- ▶ The **column space** of  $A$  is the subspace of  $\mathbb{R}^m$  spanned by the column vectors of  $A$ .
- ▶ The **row space** of  $A$  is the subspace of  $\mathbb{R}^n$  spanned by the row vectors of  $A$ .
- ▶ The dimension of the column space of  $A$  is defined as the **column rank** of  $A$ .
- ▶ The dimension of row space of  $A$  is defined as the **row rank** of  $A$ .

Fact : **Column rank = Row rank** and this number is called the **rank** of  $A$ .

## Example

Let us find the rank of the matrix  $A = \begin{bmatrix} 1 & 0 & 1 \\ -2 & -3 & 1 \\ 3 & 3 & 0 \end{bmatrix}$ .



## Example

Let us find the rank of the matrix  $A = \begin{bmatrix} 1 & 0 & 1 \\ -2 & -3 & 1 \\ 3 & 3 & 0 \end{bmatrix}$ .

Reduce it to row echelon form:



## Example

Let us find the rank of the matrix  $A = \begin{bmatrix} 1 & 0 & 1 \\ -2 & -3 & 1 \\ 3 & 3 & 0 \end{bmatrix}$ .  $\stackrel{R_2 \leftarrow R_2 + 2R_1}{=} \text{Row space}(A)$

Reduce it to row echelon form:

$$\begin{bmatrix} 1 & 0 & 1 \\ -2 & -3 & 1 \\ 3 & 3 & 0 \end{bmatrix}$$

$$R_2 \leftarrow R_2 + 2R_1$$

$$\begin{bmatrix} 1 & 0 & 1 \\ 0 & -3 & 3 \\ 3 & 3 & 0 \end{bmatrix}$$

$$R_3 \leftarrow 3R_1$$

$$\begin{bmatrix} 1 & 0 & 1 \\ 0 & -3 & 3 \\ 0 & 3 & -3 \end{bmatrix}$$

$$A \xrightarrow{R}$$

via row operations  
then rank is unchanged.

$$\begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{bmatrix}$$

$$R_3 \leftarrow 3R_2$$

$$\begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & -1 \\ 0 & 3 & -3 \end{bmatrix}$$

$$\left\{ \begin{array}{l} -R_2/3 \\ \downarrow \end{array} \right.$$

There are two non-zero rows. Hence  $\text{rank}(A) = 2$ .

## Example

Let us find the rank of the matrix  $A = \begin{bmatrix} 1 & 0 & 1 \\ -2 & -3 & 1 \\ 3 & 3 & 0 \end{bmatrix}$ .

Reduce it to row echelon form:

$$\begin{bmatrix} 1 & 0 & 1 \\ -2 & -3 & 1 \\ 3 & 3 & 0 \end{bmatrix}$$

$$R_2 + 2R_1$$

$$\begin{bmatrix} 1 & 0 & 1 \\ 0 & -3 & 3 \\ 3 & 3 & 0 \end{bmatrix}$$

$$R_3 - 3R_1$$

$$\begin{bmatrix} 1 & 0 & 1 \\ 0 & -3 & 3 \\ 0 & 3 & -3 \end{bmatrix}$$

$$\left\{ \begin{array}{l} -R_2/3 \\ \downarrow \end{array} \right.$$

$$\begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{bmatrix}$$

$$R_3 - 3R_2$$

$$\begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & -1 \\ 0 & 3 & -3 \end{bmatrix}$$

There are two non-zero rows. Hence  $\text{rank}(A) = 2$ .

# Thank you

