


Critical points for multivariable functions



Recall : Critical points for functions of one variable


A point a is called a **critical point** of a function $f(x)$ if either f is not differentiable at a or $f'(a) = 0$.



Recall : Critical points for functions of one variable

A point a is called a **critical point** of a function $f(x)$ if either f is not differentiable at a or $f'(a) = 0$.

If f is differentiable at a point a of **local extremum**, it satisfies $f'(a) = 0$ and so every point of local extremum is a critical point.



Recall : Critical points for functions of one variable

A point a is called a **critical point** of a function $f(x)$ if either f is not differentiable at a or $f'(a) = 0$.

If f is differentiable at a point a of **local extremum**, it satisfies $f'(a) = 0$ and so every point of local extremum is a critical point.




Figure: $f(x) = (x^2 - 4x + 3.8)(x + 2)x^3$

Recall : Critical points for functions of one variable

A point a is called a **critical point** of a function $f(x)$ if either f is not differentiable at a or $f'(a) = 0$.

If f is differentiable at a point a of **local extremum**, it satisfies $f'(a) = 0$ and so every point of local extremum is a critical point.




Figure: $f(x) = (x^2 - 4x + 3.8)(x + 2)x^3$

Not every critical point is a point of local extremum.

Recall : Critical points for functions of one variable

A point a is called a **critical point** of a function $f(x)$ if either f is not differentiable at a or $f'(a) = 0$.

If f is differentiable at a point a of **local extremum**, it satisfies $f'(a) = 0$ and so every point of local extremum is a critical point.





Figure: $f(x) = (x^2 - 4x + 3.8)(x + 2)x^3$

Not every critical point is a point of local extremum. A **saddle point** is a critical point which is not a point of local extremum.

Points of local extrema for multivariable functions


Let $\tilde{f}(\tilde{x})$ be a function defined on a domain D in \mathbb{R}^n and suppose $\tilde{a} \in D$.



Points of local extrema for multivariable functions

Let $f(\tilde{x})$ be a function defined on a domain D in \mathbb{R}^n and suppose $\tilde{a} \in D$.

The point \tilde{a} is a **local maximum** (or point of local maximum) of f if for some open ball B containing \tilde{a} , $f(\tilde{x}) \leq f(\tilde{a})$ whenever $\tilde{x} \in B \cap D$.



Points of local extrema for multivariable functions

Let $\tilde{f}(\tilde{x})$ be a function defined on a domain D in \mathbb{R}^n and suppose $\tilde{a} \in D$.

The point \tilde{a} is a **local maximum** (or point of local maximum) of f if for some open ball B containing \tilde{a} , $\tilde{f}(\tilde{x}) \leq \tilde{f}(\tilde{a})$ whenever $\tilde{x} \in B \cap D$.

The point \tilde{a} is a **local minimum** (or point of local minimum) of f if for some open ball B containing \tilde{a} , $\tilde{f}(a) \leq \tilde{f}(\tilde{x})$ whenever $\tilde{x} \in B \cap D$.

Points of local extrema for multivariable functions

Let $\tilde{f}(\tilde{x})$ be a function defined on a domain D in \mathbb{R}^n and suppose $\tilde{a} \in D$.

The point \tilde{a} is a **local maximum** (or point of local maximum) of f if for some open ball B containing \tilde{a} , $\tilde{f}(\tilde{x}) \leq \tilde{f}(\tilde{a})$ whenever $\tilde{x} \in B \cap D$.

The point \tilde{a} is a **local minimum** (or point of local minimum) of f if for some open ball B containing \tilde{a} , $\tilde{f}(a) \leq \tilde{f}(\tilde{x})$ whenever $\tilde{x} \in B \cap D$.

A **local extremum** (or point of local extremum) of f is either a local maximum or a local minimum of f .

The gradient vector at points of local extrema



The gradient vector at points of local extrema


Let $f(\tilde{x})$ be a function defined on a domain D in \mathbb{R}^n containing some open ball around a point a of local extremum.



The gradient vector at points of local extrema

Let $f(\tilde{x})$ be a function defined on a domain D in \mathbb{R}^n containing some open ball around a point a of local extremum.

Restrict f to a line L passing through \tilde{a} and view it as a function of one variable on L .



The gradient vector at points of local extrema

Let $f(\tilde{x})$ be a function defined on a domain D in \mathbb{R}^n containing some open ball around a point \tilde{a} of local extremum.

Restrict f to a line L passing through \tilde{a} and view it as a function of one variable on L .

Then \tilde{a} is a local extremum for the restricted function on L and hence the directional directive of f in the direction of the line L (if it exists) at \tilde{a} is 0.

The gradient vector at points of local extrema

Let $f(\tilde{x})$ be a function defined on a domain D in \mathbb{R}^n containing some open ball around a point a of local extremum.

Restrict f to a line L passing through \tilde{a} and view it as a function of one variable on L .

Then \tilde{a} is a local extremum for the restricted function on L and hence the directional directive of f in the direction of the line L (if it exists) at \tilde{a} is 0.

In particular, those partial derivatives which exist at \tilde{a} must be 0.

The gradient vector at points of local extrema

Let $f(\tilde{x})$ be a function defined on a domain D in \mathbb{R}^n containing some open ball around a point \tilde{a} of local extremum.


Restrict f to a line L passing through \tilde{a} and view it as a function of one variable on L .

Then \tilde{a} is a local extremum for the restricted function on L and hence the directional directive of f in the direction of the line L (if it exists) at \tilde{a} is 0.

In particular, those partial derivatives which exist at \tilde{a} must be 0.


If $\nabla f(\tilde{a})$ exists for a local extremum \tilde{a} , then $\nabla f(\tilde{a}) = 0$. \leftarrow ^{vector}

Critical points



Critical points

A point \tilde{a} is called a **critical point** of a function $f(\tilde{x})$ if either $\nabla f(\tilde{a})$ does not exist or $\nabla f(\tilde{a})$ exists and $\nabla f(\tilde{a}) = 0$.



Critical points

A point \tilde{a} is called a **critical point** of a function $f(\tilde{x})$ if either $\nabla f(\tilde{a})$ does not exist or $\nabla f(\tilde{a})$ exists and $\nabla f(\tilde{a}) = 0$.

Example : Critical points of $f(x, y) = x^2 + 6xy + 4y^2 + 2x - 4y$.

$$\frac{\partial f}{\partial x} = 2x + 6y + 2 \quad \frac{\partial f}{\partial y} = 6x + 8y - 4.$$

$$\nabla f(x, y) = (2x + 6y + 2, 6x + 8y - 4) = (0, 0).$$

Set $\nabla f = 0$

i.e. $\begin{cases} 2x + 6y + 2 = 0 \\ 6x + 8y - 4 = 0 \end{cases}$

$$\left[\begin{array}{cc|c} 2 & 6 & -2 \\ 6 & 8 & 4 \end{array} \right]$$

$\xrightarrow{R_1/2}$

$$\left[\begin{array}{cc|c} 1 & 3 & -1 \\ 0 & 1 & -1 \end{array} \right]$$

$$\xrightarrow{R_2 - 10R_1}$$

$$\left[\begin{array}{cc|c} 1 & 3 & -1 \\ 0 & -10 & 10 \end{array} \right]$$


$$\xrightarrow{R_2 - 6R_1}$$

$$\left[\begin{array}{cc|c} 1 & 3 & -1 \\ 0 & 8 & 4 \end{array} \right]$$

$\xrightarrow{R_1 - 3R_2}$

$$\left[\begin{array}{cc|c} 1 & 0 & 2 \\ 0 & 1 & -1 \end{array} \right] \Rightarrow \begin{cases} x = 2, y = -1 \\ \therefore \text{Critical pt. of } f \text{ is } (2, -1) \end{cases}$$

Saddle points




Saddle points

Every local extremum is a critical point.



Saddle points


Every local extremum is a critical point. Unfortunately, not all critical points are local extrema.



Saddle points

Every local extremum is a critical point. Unfortunately, not all critical points are local extrema.

Example : $f(x, y) = x^3$.




Saddle points

Every local extremum is a critical point. Unfortunately, not all critical points are local extrema.

Example : $f(x, y) = x^3$.

A **saddle point** is a critical point \tilde{a} such that $\tilde{\nabla f(a)}$ exists and $\tilde{\nabla f(a)} = 0$ but \tilde{a} is not a local extremum.




Saddle points

Every local extremum is a critical point. Unfortunately, not all critical points are local extrema.

Example : $f(x, y) = x^3$.


A **saddle point** is a critical point \tilde{a} such that $\nabla f(\tilde{a})$ exists and $\nabla f(\tilde{a}) = 0$ but \tilde{a} is not a local extremum.

$$f(x, y) = x^2 - y^2$$



Absolute (or global) extrema


Let $\tilde{f}(\tilde{x})$ be a function defined on a domain D in \mathbb{R}^n and suppose $\tilde{a} \in D$.



Absolute (or global) extrema

Let $\tilde{f}(\tilde{x})$ be a function defined on a domain D in \mathbb{R}^n and suppose $\tilde{a} \in D$.

The point \tilde{a} is an **absolute maximum** (or global maximum) of f if $\tilde{f}(\tilde{x}) \leq \tilde{f}(\tilde{a})$ for all $\tilde{x} \in D$.




Absolute (or global) extrema

Let $f(\tilde{x})$ be a function defined on a domain D in \mathbb{R}^n and suppose $a \in D$.

The point \tilde{a} is an **absolute maximum** (or global maximum) of f if $f(\tilde{x}) \leq f(\tilde{a})$ for all $\tilde{x} \in D$.

The point \tilde{a} is an **absolute minimum** (or global minimum) of f if $f(\tilde{a}) \leq f(\tilde{x})$ for all $\tilde{x} \in D$.





Absolute (or global) extrema


Let $f(\tilde{x})$ be a function defined on a domain D in \mathbb{R}^n and suppose $a \in D$.

The point \tilde{a} is an **absolute maximum** (or global maximum) of f if $f(\tilde{x}) \leq f(\tilde{a})$ for all $\tilde{x} \in D$.

The point \tilde{a} is an **absolute minimum** (or global minimum) of f if $f(\tilde{a}) \leq f(\tilde{x})$ for all $\tilde{x} \in D$.




Existence of absolute maximum/minimum



Existence of absolute maximum/minimum

A domain D in \mathbb{R}^n is called **closed** if it contains all its boundary points.



Existence of absolute maximum/minimum


A domain D in \mathbb{R}^n is called **closed** if it contains all its boundary points. A domain D in \mathbb{R}^n is called **bounded** if it is contained inside a ball around 0 with finite radius.



Existence of absolute maximum/minimum

A domain D in \mathbb{R}^n is called **closed** if it contains all its boundary points. A domain D in \mathbb{R}^n is called **bounded** if it is contained inside a ball around 0 with finite radius.

Fact : If the domain D is closed and bounded and f is continuous on D , then the global maximum and minimum must exist.




Existence of absolute maximum/minimum

A domain D in \mathbb{R}^n is called **closed** if it contains all its boundary points. A domain D in \mathbb{R}^n is called **bounded** if it is contained inside a ball around 0 with finite radius.

Fact : If the domain D is closed and bounded and f is continuous on D , then the global maximum and minimum must exist.

Note that the global maximum and minimum are in particular local maxima or local minima



Existence of absolute maximum/minimum

A domain D in \mathbb{R}^n is called **closed** if it contains all its boundary points. A domain D in \mathbb{R}^n is called **bounded** if it is contained inside a ball around 0 with finite radius.

Fact : If the domain D is closed and bounded and f is continuous on D , then the global maximum and minimum must exist.

Note that the global maximum and minimum are in particular local maxima or local minima unless they are on **boundary points**.

Existence of absolute maximum/minimum

A domain D in \mathbb{R}^n is called **closed** if it contains all its boundary points. A domain D in \mathbb{R}^n is called **bounded** if it is contained inside a ball around 0 with finite radius.

Fact : If the domain D is closed and bounded and f is continuous on D , then the global maximum and minimum must exist.

Note that the global maximum and minimum are in particular local maxima or local minima unless they are on **boundary points**.

Thus to find the global maximum and minimum, we find the critical points

- ▶ inside the domain D
 - ▶ on the boundary of D
 - ▶ on the boundary of the boundary of D
- ... and check the value of f on all of them.

Example

Find the absolute maximum and minimum of the function

$f(x, y) = x^3 + y^3 - 3x - 3y^2 + 1$ over the square with vertices $(0, 0), (2, 0), (2, 2), (0, 2)$.

$$\nabla f = (3x^2 - 3, 3y^2 - 6y) \cdot$$

$$3x^2 - 3 = 0, \quad 3y^2 - 6y = 0.$$

Sect to 0.

$$x^2 = 1, \quad y(y-2) = 0.$$

$$x = 1, \quad y = 0 \text{ or } 2.$$

(1, 0)


$$f(x, 0) = x^3 - 3x + 1$$

$$f'(x) = 0 \Rightarrow 3x^2 - 3 = 0 \Rightarrow 3x^2 - 3x + 1 = x^3 - 3x - 3.$$

$$f'(x_2) = 0 \Rightarrow x_2$$

$$f'(x) = 0 \Rightarrow x \text{ occurs at } (2,0) \\ (1,2)$$

Abc. max.
min.



$$f^{(0,y)} = y^3 - 3y^2 + 1.$$

$$f'(y) = 0$$

$$3y^2 - 6y = 0$$

$$(0,0), (0,2).$$

$$f(2,y)$$

$$= y^3 - 3y^2 + c.$$

Thank you

