

The direction of steepest ascent/descent




Tracing water flowing down a hill



Tracing water flowing down a hill

For an electrification project, Mohan Bhargava is trying to understand how water flows down a particular hill.



Tracing water flowing down a hill

For an electrification project, Mohan Bhargava is trying to understand how water flows down a particular hill.


How does water flow downhill?



Tracing water flowing down a hill

For an electrification project, Mohan Bhargava is trying to understand how water flows down a particular hill.


How does water flow downhill? It will move in the direction where the altitude decreases most rapidly.




Tracing water flowing down a hill

For an electrification project, Mohan Bhargava is trying to understand how water flows down a particular hill.

How does water flow downhill? It will move in the direction where the altitude decreases most rapidly.




Tracing water flowing down a hill (contd.)



Tracing water flowing down a hill (contd.)


Mohan models the hill as in the previous picture using the graph of a function $h(x)$ where h is the altitude of a point.



Tracing water flowing down a hill (contd.)

Mohan models the hill as in the previous picture using the graph of a function $h(x)$ where h is the altitude of a point.

He then calculates the derivative $h'(x)$ and computes at which points it is negative, positive and 0.



Tracing water flowing down a hill (contd.)

Mohan models the hill as in the previous picture using the graph of a function $h(x)$ where h is the altitude of a point.

He then calculates the derivative $h'(x)$ and computes at which points it is negative, positive and 0.

- ▶ If the derivative is negative at a point, water flows to the right from that point.

Tracing water flowing down a hill (contd.)

Mohan models the hill as in the previous picture using the graph of a function $h(x)$ where h is the altitude of a point.

He then calculates the derivative $h'(x)$ and computes at which points it is negative, positive and 0.

- ▶ If the derivative is negative at a point, water flows to the right from that point.
- ▶ If the derivative is positive at a point, water flows to the right from that point.


Tracing water flowing down a hill (contd.)

Mohan models the hill as in the previous picture using the graph of a function $h(x)$ where h is the altitude of a point.

He then calculates the derivative $h'(x)$ and computes at which points it is negative, positive and 0.


- ▶ If the derivative is negative at a point, water flows to the right from that point.
- ▶ If the derivative is positive at a point, water flows to the right from that point.
- ▶ If the derivative is 0 at a point, water will remain stationary at that point.

Tracing water flowing down a hill (contd.)



Tracing water flowing down a hill (contd.)


Mohan's friend Gita points out that a one dimensional model (i.e. a cross section) will not be useful and that a two dimensional model will be more useful.



Tracing water flowing down a hill (contd.)

Mohan's friend Gita points out that a one dimensional model (i.e. a cross section) will not be useful and that a two dimensional model will be more useful.


She uses a 2-dimensional function $h(x, y)$ to model the altitude h .



Tracing water flowing down a hill (contd.)

Mohan's friend Gita points out that a one dimensional model (i.e. a cross section) will not be useful and that a two dimensional model will be more useful.

She uses a 2-dimensional function $h(x, y)$ to model the altitude h .




Tracing water flowing down a hill (contd.)



Tracing water flowing down a hill (contd.)


Mohan asks her how she will find out the direction in which the water will flow since there are now more than 2 directions to contend with.



Tracing water flowing down a hill (contd.)

Mohan asks her how she will find out the direction in which the water will flow since there are now more than 2 directions to contend with.

Gita tells Mohan that water will flow in the direction in which the altitude decreases fastest, i.e. along the steepest slope downward.



Tracing water flowing down a hill (contd.)

Mohan asks her how she will find out the direction in which the water will flow since there are now more than 2 directions to contend with.

Gita tells Mohan that water will flow in the direction in which the altitude decreases fastest, i.e. along the steepest slope downward.


She says that to compute it, we have to find the direction in which the function h decreases fastest

Tracing water flowing down a hill (contd.)

Mohan asks her how she will find out the direction in which the water will flow since there are now more than 2 directions to contend with.

Gita tells Mohan that water will flow in the direction in which the altitude decreases fastest, i.e. along the steepest slope downward.

She says that to compute it, we have to find the direction in which the function h decreases fastest or equivalently the rate of decrease of h is fastest.



Tracing water flowing down a hill (contd.)

Mohan asks her how she will find out the direction in which the water will flow since there are now more than 2 directions to contend with.

Gita tells Mohan that water will flow in the direction in which the altitude decreases fastest, i.e. along the steepest slope downward.

She says that to compute it, we have to find the direction in which the function h decreases fastest or equivalently the rate of decrease of h is fastest.

This is same as finding the vector u in which the directional derivative h_u is largest in absolute value amongst those for which it has negative sign

Tracing water flowing down a hill (contd.)

Mohan asks her how she will find out the direction in which the water will flow since there are now more than 2 directions to contend with.

Gita tells Mohan that water will flow in the direction in which the altitude decreases fastest, i.e. along the steepest slope downward.

She says that to compute it, we have to find the direction in which the function h decreases fastest or equivalently the rate of decrease of h is fastest.

This is same as finding the vector u in which the directional derivative h_u is largest in absolute value amongst those for which it has negative sign i.e. u such that

1. $h_u \leq h_v$ for all $v \in \mathbb{R}^2$ and
2. $h_u < 0$.

In what direction is the directional derivative minimized?

Let $f(x_1, x_2, \dots, x_n)$ be a function defined on a domain D in \mathbb{R}^n containing some open ball around the point a .



In what direction is the directional derivative minimized?

Let $f(x_1, x_2, \dots, x_n)$ be a function defined on a domain D in \mathbb{R}^n containing some open ball around the point a .

Suppose ∇f exists and is continuous on some open ball around the point a .



In what direction is the directional derivative minimized?

Let $f(x_1, x_2, \dots, x_n)$ be a function defined on a domain D in \mathbb{R}^n containing some open ball around the point \tilde{a} .

Suppose ∇f exists and is continuous on some open ball around the point \tilde{a} .

$$\begin{aligned}f_u &= \nabla f(\tilde{a}) \cdot u \\&= \|\nabla f(\tilde{a})\| \|u\| \cos(\theta) \\&= \|\nabla f(\tilde{a})\| \omega_s(\theta)\end{aligned}$$

where θ is
the angle
between $\nabla f(\tilde{a})$
and u .

$\omega_s(\theta)$ is minimized when $\theta = \pi$
i.e. u is pointing in the direction
opposite to $\nabla f(\tilde{a})$.

∴ The minimum value of f_u is attained when
 $u = -\nabla f(\tilde{a}) / \|\nabla f(\tilde{a})\|$ & is equal to $-\|\nabla f(\tilde{a})\|$.

Directions in which the directional derivative is maximized or remains unchanged


Assume the same hypothesis as the previous slide.

$$f_u = \|\nabla f(a)\| \|u\| \cos(\theta) = \|\nabla f(a)\| \cos(\theta)$$

It is maximized when $\theta = 0$, i.e. u is in the same direction as $\nabla f(a)$ i.e. $u = \frac{\nabla f(a)}{\|\nabla f(a)\|}$.

It remains unchanged when $f_u = 0$ i.e. $\theta = \pi/2$ i.e. u is orthogonal / perpendicular to $\nabla f(a)$.

Directions : steepest ascent, steepest descent, no change



Directions : steepest ascent, steepest descent, no change

Assume the same hypothesis as the previous slide.



Directions : steepest ascent, steepest descent, no change

Assume the same hypothesis as the previous slide.

Property	In terms of directional derivatives	Direction
Steepest ascent	f_u is positive and maximum	$u = \nabla f / \nabla f $
Steepest descent	f_u is negative and minimum	$u = -\nabla f / \nabla f $
No change	$f_u = 0$	u is orthogonal to ∇f

Examples

1. $f(x, y) = \sin(xy)$

$$\nabla f(x, y) = (y \cos(xy), x \cos(xy))$$

At $(\pi, 1)$ what is the direction of steepest descent
on the graph of this function?

$$\nabla f(\pi, 1) = (\pi \cos(\pi), \pi \times \cos(\pi)) = (-1, -\pi).$$

$$u = \frac{-\nabla f(\pi, 1)}{\|\nabla f(\pi, 1)\|} = \frac{(-1, -\pi)}{\sqrt{1+\pi^2}}.$$

2. $f(x, y, z) = x^2 + y^2 + z^2$

$$\nabla f(x, y, z) = (2x, 2y, 2z)$$

At $(1, 1, 1)$ what is the direction in which the fn.

$$\nabla f(1, 1, 1) = (2, 2, 2) \text{ fastest increase? In the direction } u = \frac{(2, 2, 2)}{\sqrt{2^2+2^2+2^2}} = \frac{1}{\sqrt{3}}(1, 1, 1).$$

$\nabla f(1, 1, 1) = (2, 2, 2)$. In the direction $u = \frac{(2, 2, 2)}{\sqrt{2^2+2^2+2^2}} = \frac{1}{\sqrt{3}}(1, 1, 1)$, does the fn. remain constant?


In which direction does the fn. remain constant?
e.g. $(1, -1, 0) \frac{1}{\sqrt{2}}$, $\frac{1}{\sqrt{2}}(1, 0, -1)$.

A cautionary tale

$$f(x, y) = \begin{cases} \frac{xy}{x^2+y^2} & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{if } (x, y) = (0, 0) \end{cases}$$

$$\nabla f(0, 0) = (0, 0).$$

$f_u(0, 0)$ DNE unless $u = e_1$ or e_2 .



Thank you

