

What are vector spaces?

Sarang S. Sane

Recall : vectors in \mathbb{R}^n

Recall : vectors in \mathbb{R}^n

Consider vectors $(x_1, x_2, \dots, x_n), (y_1, y_2, \dots, y_n)$ in \mathbb{R}^n and $c \in \mathbb{R}$.

Recall : vectors in \mathbb{R}^n

Consider vectors $(x_1, x_2, \dots, x_n), (y_1, y_2, \dots, y_n)$ in \mathbb{R}^n and $c \in \mathbb{R}$.

- ▶ Recall **addition** of these vectors is defined as :

Recall : vectors in \mathbb{R}^n

Consider vectors $(x_1, x_2, \dots, x_n), (y_1, y_2, \dots, y_n)$ in \mathbb{R}^n and $c \in \mathbb{R}$.

- ▶ Recall **addition** of these vectors is defined as :

$$(x_1, x_2, \dots, x_n) + (y_1, y_2, \dots, y_n) = (x_1 + y_1, x_2 + y_2, \dots, x_n + y_n)$$

Recall : vectors in \mathbb{R}^n

Consider vectors $(x_1, x_2, \dots, x_n), (y_1, y_2, \dots, y_n)$ in \mathbb{R}^n and $c \in \mathbb{R}$.

- ▶ Recall **addition** of these vectors is defined as :

$$(x_1, x_2, \dots, x_n) + (y_1, y_2, \dots, y_n) = (x_1 + y_1, x_2 + y_2, \dots, x_n + y_n)$$

- ▶ and **scalar multiplication** is defined as :

Recall : vectors in \mathbb{R}^n

Consider vectors $(x_1, x_2, \dots, x_n), (y_1, y_2, \dots, y_n)$ in \mathbb{R}^n and $c \in \mathbb{R}$.

- ▶ Recall **addition** of these vectors is defined as :

$$(x_1, x_2, \dots, x_n) + (y_1, y_2, \dots, y_n) = (x_1 + y_1, x_2 + y_2, \dots, x_n + y_n)$$

- ▶ and **scalar multiplication** is defined as :

$$c(x_1, x_2, \dots, x_n) = (cx_1, cx_2, \dots, cx_n)$$

Properties of addition and scalar multiplication

Let v, w and v' be vectors in \mathbb{R}^n and $a, b \in \mathbb{R}$.

Properties of addition and scalar multiplication

Let v, w and v' be vectors in \mathbb{R}^n and $a, b \in \mathbb{R}$.

i) $v + w = w + v.$

Properties of addition and scalar multiplication

Let v, w and v' be vectors in \mathbb{R}^n and $a, b \in \mathbb{R}$.

- i) $v + w = w + v.$
- ii) $(v + w) + v' = v + (w + v').$

Properties of addition and scalar multiplication

Let v, w and v' be vectors in \mathbb{R}^n and $a, b \in \mathbb{R}$.

- i) $v + w = w + v$.
- ii) $(v + w) + v' = v + (w + v')$.
- iii) The 0 vector satisfies that $v + 0 = 0 + v = v$.

Properties of addition and scalar multiplication

Let v, w and v' be vectors in \mathbb{R}^n and $a, b \in \mathbb{R}$.

- i) $v + w = w + v$.
- ii) $(v + w) + v' = v + (w + v')$.
- iii) The 0 vector satisfies that $v + 0 = 0 + v = v$.
- iv) The vector $-v$ satisfies that $v + (-v) = 0$.

Properties of addition and scalar multiplication

Let v, w and v' be vectors in \mathbb{R}^n and $a, b \in \mathbb{R}$.

- i) $v + w = w + v$.
- ii) $(v + w) + v' = v + (w + v')$.
- iii) The 0 vector satisfies that $v + 0 = 0 + v = v$.
- iv) The vector $-v$ satisfies that $v + (-v) = 0$.
- v) $1v = v$.

Properties of addition and scalar multiplication

Let v, w and v' be vectors in \mathbb{R}^n and $a, b \in \mathbb{R}$.

- i) $v + w = w + v$.
- ii) $(v + w) + v' = v + (w + v')$.
- iii) The 0 vector satisfies that $v + 0 = 0 + v = v$.
- iv) The vector $-v$ satisfies that $v + (-v) = 0$.
- v) $1v = v$.
- vi) $(ab)v = a(bv)$.

Properties of addition and scalar multiplication

Let v, w and v' be vectors in \mathbb{R}^n and $a, b \in \mathbb{R}$.

- i) $v + w = w + v$.
- ii) $(v + w) + v' = v + (w + v')$.
- iii) The 0 vector satisfies that $v + 0 = 0 + v = v$.
- iv) The vector $-v$ satisfies that $v + (-v) = 0$.
- v) $1v = v$.
- vi) $(ab)v = a(bv)$.
- vii) $a(v + w) = av + aw$.

Properties of addition and scalar multiplication

Let v, w and v' be vectors in \mathbb{R}^n and $a, b \in \mathbb{R}$.

- i) $v + w = w + v.$
- ii) $(v + w) + v' = v + (w + v').$
- iii) The 0 vector satisfies that $v + 0 = 0 + v = v.$
- iv) The vector $-v$ satisfies that $v + (-v) = 0.$
- v) $1v = v.$
- vi) $(ab)v = a(bv).$
- vii) $a(v + w) = av + aw.$
- viii) $(a + b)v = av + bv$

Properties of addition and scalar multiplication

Let v, w and v' be vectors in \mathbb{R}^n and $a, b \in \mathbb{R}$.

- i) $v + w = w + v$.
- ii) $(v + w) + v' = v + (w + v')$.
- iii) The 0 vector satisfies that $v + 0 = 0 + v = v$.
- iv) The vector $-v$ satisfies that $v + (-v) = 0$.
- v) $1v = v$.
- vi) $(ab)v = a(bv)$.
- vii) $a(v + w) = av + aw$.
- viii) $(a + b)v = av + bv$

A **vector space** is a set with two operations (called **addition** and **scalar multiplication** with the above properties (i)-(viii).

Defintion of a vector space

A vector space V over \mathbb{R} is a set along with two functions

$$+ : V \times V \rightarrow V \quad \text{and} \quad \cdot : \mathbb{R} \times V \rightarrow V$$

Defintion of a vector space

A vector space V over \mathbb{R} is a set along with two functions

$$+ : V \times V \rightarrow V \quad \text{and} \quad \cdot : \mathbb{R} \times V \rightarrow V$$

(i.e. for each pair of elements v_1 and v_2 in V , there is a unique element $v_1 + v_2$ in V , and for each $c \in \mathbb{R}$ and $v \in V$ there is a unique element $c \cdot v$ in V)

Defintion of a vector space

A vector space V over \mathbb{R} is a set along with two functions

$$+ : V \times V \rightarrow V \quad \text{and} \quad \cdot : \mathbb{R} \times V \rightarrow V$$

(i.e. for each pair of elements v_1 and v_2 in V , there is a unique element $v_1 + v_2$ in V , and for each $c \in \mathbb{R}$ and $v \in V$ there is a unique element $c \cdot v$ in V)

It is standard to suppress the \cdot and only write cv instead of $c.v$.

Defintion of a vector space

A vector space V over \mathbb{R} is a set along with two functions

$$+ : V \times V \rightarrow V \quad \text{and} \quad \cdot : \mathbb{R} \times V \rightarrow V$$

(i.e. for each pair of elements v_1 and v_2 in V , there is a unique element $v_1 + v_2$ in V , and for each $c \in \mathbb{R}$ and $v \in V$ there is a unique element $c \cdot v$ in V)

It is standard to suppress the \cdot and only write cv instead of $c.v$.

The functions $+$ and \cdot are required to satisfy the following rules :

Formal definition (Contd.)

i) $v_1 + v_2 = v_2 + v_1$ for all $v_1, v_2 \in V$

Formal definition (Contd.)

- i) $v_1 + v_2 = v_2 + v_1$ for all $v_1, v_2 \in V$
- ii) $(v_1 + v_2) + v_3 = v_1 + (v_2 + v_3)$ for all $v_1, v_2, v_3 \in V$

Formal definition (Contd.)

- i) $v_1 + v_2 = v_2 + v_1$ for all $v_1, v_2 \in V$
- ii) $(v_1 + v_2) + v_3 = v_1 + (v_2 + v_3)$ for all $v_1, v_2, v_3 \in V$
- iii) There exists an element in V denoted by 0 such that
 $v + 0 = v$ for all $v \in V$

Formal definition (Contd.)

- i) $v_1 + v_2 = v_2 + v_1$ for all $v_1, v_2 \in V$
- ii) $(v_1 + v_2) + v_3 = v_1 + (v_2 + v_3)$ for all $v_1, v_2, v_3 \in V$
- iii) There exists an element in V denoted by 0 such that
 $v + 0 = v$ for all $v \in V$
- iv) For each element $v \in V$ there exists an element $v' \in V$ such
that $v + v' = 0$

Formal definition (Contd.)

- i) $v_1 + v_2 = v_2 + v_1$ for all $v_1, v_2 \in V$
- ii) $(v_1 + v_2) + v_3 = v_1 + (v_2 + v_3)$ for all $v_1, v_2, v_3 \in V$
- iii) There exists an element in V denoted by 0 such that
 $v + 0 = v$ for all $v \in V$
- iv) For each element $v \in V$ there exists an element $v' \in V$ such
that $v + v' = 0$
- v) For each element $v \in V$, $1v = v$

Formal definition (Contd.)

- i) $v_1 + v_2 = v_2 + v_1$ for all $v_1, v_2 \in V$
- ii) $(v_1 + v_2) + v_3 = v_1 + (v_2 + v_3)$ for all $v_1, v_2, v_3 \in V$
- iii) There exists an element in V denoted by 0 such that
 $v + 0 = v$ for all $v \in V$
- iv) For each element $v \in V$ there exists an element $v' \in V$ such
that $v + v' = 0$
- v) For each element $v \in V$, $1v = v$
- vi) For each pair of elements $a, b \in \mathbb{R}$ and each element $v \in V$,
 $(ab)v = a(bv)$

Formal definition (Contd.)

- i) $v_1 + v_2 = v_2 + v_1$ for all $v_1, v_2 \in V$
- ii) $(v_1 + v_2) + v_3 = v_1 + (v_2 + v_3)$ for all $v_1, v_2, v_3 \in V$
- iii) There exists an element in V denoted by 0 such that
 $v + 0 = v$ for all $v \in V$
- iv) For each element $v \in V$ there exists an element $v' \in V$ such that $v + v' = 0$
- v) For each element $v \in V$, $1v = v$
- vi) For each pair of elements $a, b \in \mathbb{R}$ and each element $v \in V$,
 $(ab)v = a(bv)$
- vii) For each element $a \in \mathbb{R}$ and each pair of elements v_1 and v_2 ,
 $a(v_1 + v_2) = av_1 + av_2$

Formal definition (Contd.)

- i) $v_1 + v_2 = v_2 + v_1$ for all $v_1, v_2 \in V$
- ii) $(v_1 + v_2) + v_3 = v_1 + (v_2 + v_3)$ for all $v_1, v_2, v_3 \in V$
- iii) There exists an element in V denoted by 0 such that
 $v + 0 = v$ for all $v \in V$
- iv) For each element $v \in V$ there exists an element $v' \in V$ such that $v + v' = 0$
- v) For each element $v \in V$, $1v = v$
- vi) For each pair of elements $a, b \in \mathbb{R}$ and each element $v \in V$,
 $(ab)v = a(bv)$
- vii) For each element $a \in \mathbb{R}$ and each pair of elements v_1 and v_2 ,
 $a(v_1 + v_2) = av_1 + av_2$
- viii) For each pair of elements $a, b \in \mathbb{R}$ and each element $v \in V$,
 $(a + b)v = av + bv$

Example : Matrices

Let $M_{m \times n}(\mathbb{R})$ be the set of all $m \times n$ matrices with real numbers.

Example : Matrices

Let $M_{m \times n}(\mathbb{R})$ be the set of all $m \times n$ matrices with real numbers.

Recall that we have defined addition and scalar multiplication on $M_{m \times n}(\mathbb{R})$ as follows :

Example : Matrices

Let $M_{m \times n}(\mathbb{R})$ be the set of all $m \times n$ matrices with real numbers.

Recall that we have defined addition and scalar multiplication on $M_{m \times n}(\mathbb{R})$ as follows :

- ▶ $(A + B)_{ij} = A_{ij} + B_{ij}$

Example : Matrices

Let $M_{m \times n}(\mathbb{R})$ be the set of all $m \times n$ matrices with real numbers.

Recall that we have defined addition and scalar multiplication on $M_{m \times n}(\mathbb{R})$ as follows :

- ▶ $(A + B)_{ij} = A_{ij} + B_{ij}$
- ▶ $(cA)_{ij} = c(A)_{ij}$

Example : Matrices

Let $M_{m \times n}(\mathbb{R})$ be the set of all $m \times n$ matrices with real numbers.

Recall that we have defined addition and scalar multiplication on $M_{m \times n}(\mathbb{R})$ as follows :

- ▶ $(A + B)_{ij} = A_{ij} + B_{ij}$
- ▶ $(cA)_{ij} = c(A)_{ij}$

where $A, B \in M_{m \times n}(\mathbb{R})$ and $c \in \mathbb{R}$.

Example : Matrices

Let $M_{m \times n}(\mathbb{R})$ be the set of all $m \times n$ matrices with real numbers.

Recall that we have defined addition and scalar multiplication on $M_{m \times n}(\mathbb{R})$ as follows :

- ▶ $(A + B)_{ij} = A_{ij} + B_{ij}$
- ▶ $(cA)_{ij} = c(A)_{ij}$

where $A, B \in M_{m \times n}(\mathbb{R})$ and $c \in \mathbb{R}$.

Then $M_{m \times n}(\mathbb{R})$ along with addition and scalar multiplication forms a vector space.

Example : Solutions of a homogeneous system

Consider the set of solutions V of a homogeneous system $Ax = 0$ where $A \in M_{m \times n}(\mathbb{R})$ (i.e. this is a homogeneous system of m linear equations in n variables).

Example : Solutions of a homogeneous system

Consider the set of solutions V of a homogeneous system $Ax = 0$ where $A \in M_{m \times n}(\mathbb{R})$ (i.e. this is a homogeneous system of m linear equations in n variables).

Note that if $v, w \in V$ then

$$A(v + w) = Av + Aw = 0 + 0 = 0.$$

Example : Solutions of a homogeneous system

Consider the set of solutions V of a homogeneous system $Ax = 0$ where $A \in M_{m \times n}(\mathbb{R})$ (i.e. this is a homogeneous system of m linear equations in n variables).

Note that if $v, w \in V$ then

$$A(v + w) = Av + Aw = 0 + 0 = 0.$$

and if $c \in \mathbb{R}$ then

$$A(cv) = c(Av) = c(0) = 0.$$

Example : Solutions of a homogeneous system

Consider the set of solutions V of a homogeneous system $Ax = 0$ where $A \in M_{m \times n}(\mathbb{R})$ (i.e. this is a homogeneous system of m linear equations in n variables).

Note that if $v, w \in V$ then

$$A(v + w) = Av + Aw = 0 + 0 = 0. \Rightarrow v + w \in V$$

and if $c \in \mathbb{R}$ then

$$A(cv) = c(Av) = c(0) = 0. \Rightarrow cv \in V$$

So addition and scalar multiplication on \mathbb{R}^n restricts to the solution set. Hence it is a vector space.

Example : Solutions of a homogeneous system

Consider the set of solutions V of a homogeneous system $Ax = 0$ where $A \in M_{m \times n}(\mathbb{R})$ (i.e. this is a homogeneous system of m linear equations in n variables).

Note that if $v, w \in V$ then

$$A(v + w) = Av + Aw = 0 + 0 = 0. \quad \Rightarrow v+w \in V$$

and if $c \in \mathbb{R}$ then

$$A(cv) = c(Av) = c(0) = 0. \quad \Rightarrow cv \in V$$

So addition and scalar multiplication on \mathbb{R}^n restricts to the solution set. Hence it is a vector space.

This is an example of a subspace of a vector space.

Non-example

Let us define addition and scalar multiplication in \mathbb{R}^2 as follows:

- ▶ $(x_1, x_2) + (y_1, y_2) = (x_1 + y_1, x_2 - y_2)$
- ▶ $c(x_1, x_2) = (cx_1, cx_2)$

Non-example

Let us define addition and scalar multiplication in \mathbb{R}^2 as follows:

- $(x_1, x_2) + (y_1, y_2) = (x_1 + y_1, x_2 - y_2)$
- $c(x_1, x_2) = (cx_1, cx_2)$

Check that (i), (ii) and (viii) fail to hold.

(i) fails :

$$(x_1, x_2) + (y_1, y_2) = (x_1 + y_1, x_2 - y_2)$$
$$(y_1, y_2) + (x_1, x_2) = (y_1 + x_1, y_2 - x_2)$$
$$(0, 0) + (1, 1) = (1, -1)$$
$$(1, 1) + (0, 0) = (1, 1)$$

Non-example

Let us define addition and scalar multiplication in \mathbb{R}^2 as follows:

- ▶ $(x_1, x_2) + (y_1, y_2) = (x_1 + y_1, 0)$
- ▶ $c(x_1, x_2) = (cx_1, 0)$

Non-example

Let us define addition and scalar multiplication in \mathbb{R}^2 as follows:

- ▶ $(x_1, x_2) + (y_1, y_2) = (x_1 + y_1, 0)$
- ▶ $c(x_1, x_2) = (cx_1, 0)$

Check that (iii), (iv) and (v) fail to hold.

Cancellation law of vector addition

If $v_1, v_2, v_3 \in V$ such that $v_1 + v_3 = v_2 + v_3$, then $v_1 = v_2$.

$$(x_1, x_2, x_3) + (y_1, y_2, y_3) = (z_1, z_2, z_3) + (y_1, y_2, y_3)$$

in \mathbb{R}^3

$$\begin{aligned} & \Rightarrow x_i + y_i = z_i + y_i \quad \forall i=1, 2, 3 \\ & \text{Subtract } y_i \text{ from both sides} \Rightarrow x_i = z_i \quad \forall i=1, 2, 3. \end{aligned}$$

on gen.
in \mathbb{R}^n

$$v_1 + v_3 = v_2 + v_3 \Rightarrow (v_1 + v_3) + v_3' = (v_2 + v_3) + v_3'$$
$$\begin{aligned} & \Rightarrow v_1 + (v_3 + v_3') = v_2 + (v_3 + v_3') \\ & \Rightarrow v_1 + 0 = v_2 + 0 \\ & \Rightarrow v_1 = v_2. \end{aligned}$$
$$v_3 + v_3' = 0$$

Cancellation law of vector addition

If $v_1, v_2, v_3 \in V$ such that $v_1 + v_3 = v_2 + v_3$, then $v_1 = v_2$.

Corollaries:

- ▶ The vector 0 described in (iii) is unique.

Cancellation law of vector addition

If $v_1, v_2, v_3 \in V$ such that $v_1 + v_3 = v_2 + v_3$, then $v_1 = v_2$.

Corollaries:

- The vector 0 described in (iii) is unique.

Suppose $\exists w \in V$ s.t. $v + w = v \neq v \in V$.

$$v + w = v + 0 \Rightarrow w = 0.$$

- The vector v' described in (iv) is unique and it is standard to refer to it as $-v$.

Suppose v'' also satisfies this.

$$v + v' = 0$$
$$v + v' = 0 = v + v'' \therefore v' = v''.$$

Then $v + v' = 0 = v + v''$ Cancel v

Some more important properties

In any vector space V the following statements are true.

Some more important properties

In any vector space V the following statements are true.

- ▶ $0v = 0$ for each $v \in V$.

Some more important properties

In any vector space V the following statements are true.

- ▶ $0v = 0$ for each $v \in V$.

- ▶ $(-c)v = -(cv) = c(-v)$ for each $c \in \mathbb{R}$ and for each $v \in V$.

Some more important properties

In any vector space V the following statements are true.

- ▶ $0v = 0$ for each $v \in V$.

$$\begin{array}{lcl} (0+0)v & = & 0v + 0v \\ \text{" } 0v & & \end{array} \quad \left. \begin{array}{l} 0v = 0v + 0v \\ \Rightarrow 0v + 0 = 0v + 0v \\ \text{Cancel } 0v \Rightarrow 0 = 0v. \end{array} \right\}$$

- ▶ $(-c)v = -(cv) = c(-v)$ for each $c \in \mathbb{R}$ and for each $v \in V$.

$$\begin{array}{lcl} (c + (-c))v & = & cv + (-c)v \\ \text{" } 0v = 0 & & \end{array} \quad \begin{array}{l} \Rightarrow cv + (-c)v = 0 \\ \Rightarrow (-c)v = -cv. \end{array}$$

- ▶ $c0 = 0$ for each $c \in \mathbb{R}$.

Check this!

Earlier Example : Stock taking

Stock taking in a grocery shop :

Items	In stock	Buyer A	Buyer B	Buyer C	New stock
Rice in kg	150	8	12	3	100
Dal in kg	50	8	5	2	75
Oil in Litres	35	4	7	5	30
Biscuits in packets	70	10	10	5	80
Soap Bars	25	4	2	1	30

Vector space : { (Quantity of rice in kg, quantity of dal in kg, the no. of biscuit packets,
quantity of oil in litres, the no. of soap bars) }

IS like \mathbb{R}^5 .

-ve components to demand
+ve " to supply.


Example : Affine flats

Let V be a plane parallel to the XY -plane. We will define an "addition" and "scalar multiplication" of points on V .

Example : Affine flats

Let V be a plane parallel to the XY -plane. We will define an "addition" and "scalar multiplication" of points on V .

Scalar multiplication : Let $Q \in V$ and $c \in \mathbb{R}$. Project Q onto the XY -plane, scale the resulting vector by c and project the result back to V . Define cQ to be the tip of the obtained arrow.



Example : Affine flats (contd.)

Addition :

Example : Affine flats (contd.)

Addition :

Check that this is a vector space both geometrically (visualize!) and by writing down the algebra.

Thank you