


Projections using inner products

Sarang S. Sane


Shortest distances in \mathbb{R}^2

A and B are points in the plane \mathbb{R}^2 and we want to find the nearest point from B on the line passing through A and the origin.




Shortest distances in \mathbb{R}^2

A and B are points in the plane \mathbb{R}^2 and we want to find the nearest point from B on the line passing through A and the origin. Drop a perpendicular from B on to the line.




Shortest distances in \mathbb{R}^2

A and B are points in the plane \mathbb{R}^2 and we want to find the nearest point from B on the line passing through A and the origin. Drop a perpendicular from B on to the line. Let a and b be the vectors corresponding to the points A and B respectively.




Shortest distances in \mathbb{R}^3

Suppose we have a vector b in \mathbb{R}^3 , and we want to find out the nearest point ~~of~~ ^{to} b on the two dimensional plane generated by the vectors $(1, 0, 0)$ and $(0, 1, 0)$.



The projection of a vector to a subspace



The projection of a vector to a subspace

Let V be an inner product space , $v \in V$ and $W \subseteq V$ be a subspace.



The projection of a vector to a subspace


Let V be an inner product space , $v \in V$ and $W \subseteq V$ be a subspace. Then the **projection of v onto W** is the vector in W , denoted by $\text{proj}_W(v)$, computed as follows :



The projection of a vector to a subspace

Let V be an inner product space , $v \in V$ and $W \subseteq V$ be a subspace. Then the **projection of v onto W** is the vector in W , denoted by $\text{proj}_W(v)$, computed as follows :

Find an orthonormal basis $\{v_1, v_2, \dots, v_n\}$ for V .



The projection of a vector to a subspace

Let V be an inner product space , $v \in V$ and $W \subseteq V$ be a subspace. Then the **projection of v onto W** is the vector in W , denoted by $\text{proj}_W(v)$, computed as follows :

Find an orthonormal basis $\{v_1, v_2, \dots, v_n\}$ for V .

Define $\text{proj}_W(v) = \sum_{i=1}^n \langle v, v_i \rangle v_i.$

The projection of a vector to a subspace

Let V be an inner product space, $v \in V$ and $W \subseteq V$ be a subspace. Then the **projection of v onto W** is the vector in W , denoted by $\text{proj}_W(v)$, computed as follows :

Find an orthonormal basis $\{v_1, v_2, \dots, v_n\}$ for ~~V~~ W .

Define

$$\text{proj}_W(v) = \boxed{\sum_{i=1}^n \langle v, v_i \rangle v_i.}$$

Fact : The definition is independent of the chosen orthonormal basis (i.e. the expression on the RHS does not change even if you choose a different orthonormal basis).

The projection of a vector to a subspace

Let V be an inner product space , $v \in V$ and $W \subseteq V$ be a subspace. Then the **projection of v onto W** is the vector in W , denoted by $\text{proj}_W(v)$, computed as follows :

Find an orthonormal basis $\{v_1, v_2, \dots, v_n\}$ for W .

Define $\text{proj}_W(v) = \sum_{i=1}^n \langle v, v_i \rangle v_i.$

Fact : The definition is independent of the chosen orthonormal basis (i.e. the expression on the RHS does not change even if you choose a different orthonormal basis).

The projection of v onto W is the vector in W closest to v .

The projection of a vector to a subspace

Let V be an inner product space, $v \in V$ and $W \subseteq V$ be a subspace. Then the **projection of v onto W** is the vector in W , denoted by $\text{proj}_W(v)$, computed as follows :

Find an orthonormal basis $\{v_1, v_2, \dots, v_n\}$ for ~~W~~ .

Define $\text{proj}_W(v) = \sum_{i=1}^n \langle v, v_i \rangle v_i.$


Fact : The definition is independent of the chosen orthonormal basis (i.e. the expression on the RHS does not change even if you choose a different orthonormal basis).

*Find $w \in W$ s.t. $\|v - w\|$ is smallest.
Ans. $w = \text{proj}_W(v)$.*

The projection of v onto W is the vector in W closest to v . Note that "closest" is in terms of the distance based on the norm induced by the inner product.

Previous examples

$V = \mathbb{R}^2$, $W = \langle(3, 1)\rangle$, $v = (1, 3)$. Then $\text{proj}_W(v) = (1.8, 0.6)$.



Previous examples

$V = \mathbb{R}^2$, $W = \langle(3, 1)\rangle$, $v = (1, 3)$. Then $\text{proj}_W(v) = (1.8, 0.6)$. ✓

$\frac{1}{\sqrt{10}}(3, 1)$ is an o.n. basis for W .

$$\begin{aligned}\text{proj}_W(v) &= \left\langle v, \frac{1}{\sqrt{10}}(3, 1) \right\rangle \frac{1}{\sqrt{10}}(3, 1) \\ &= \frac{\langle (1, 3), (3, 1) \rangle}{\sqrt{10}} (3, 1) = \frac{\frac{1}{10}(3, 1)}{\sqrt{10}} (3, 1) \\ &= \frac{6}{10} (3, 1) = (1.8, 0.6).\end{aligned}$$

$V = \mathbb{R}^3$, $W = \langle(1, 0, 0), (0, 1, 0)\rangle$, $v = (2, 3, 5)$.

Then $\text{proj}_W(v) = (2, 3, 0)$.

o.n. basis $\langle(1, 0, 0), (0, 1, 0)\rangle$.

$$\begin{aligned}\text{proj}_W(v) &= \left\langle (2, 3, 5), \frac{(1, 0, 0)}{\sqrt{1+0+0}} \right\rangle (1, 0, 0) + \left\langle (2, 3, 5), \frac{(0, 1, 0)}{\sqrt{0+1+0}} \right\rangle (0, 1, 0) \\ &= 2(1, 0, 0) + 3(0, 1, 0) = (2, 3, 0).\end{aligned}$$

Projection on a vector and orthogonal bases


Let V be an inner product space and $v, w \in V$.



Projection on a vector and orthogonal bases

Let V be an inner product space and $v, w \in V$. Define

$$\text{proj}_w(v) = \text{proj}_{\langle w \rangle}(v).$$




Projection on a vector and orthogonal bases

Let V be an inner product space and $v, w \in V$. Define

$$\text{proj}_w(v) = \text{proj}_{\langle w \rangle}(v).$$

Note that an orthonormal basis for $\langle w \rangle$ is $\frac{w}{\|w\|}$



Projection on a vector and orthogonal bases

Let V be an inner product space and $v, w \in V$. Define

$$\text{proj}_w(v) = \text{proj}_{\langle w \rangle}(v).$$

Note that an orthonormal basis for $\langle w \rangle$ is $\frac{w}{\|w\|}$ and hence

$$\text{proj}_w(v) = \left\langle v, \frac{w}{\|w\|} \right\rangle \frac{w}{\|w\|} = \frac{\langle v, w \rangle}{\|w\|^2} w = \frac{\langle v, w \rangle}{\langle w, w \rangle} w.$$

Projection on a vector and orthogonal bases

Let V be an inner product space and $v, w \in V$. Define

$$\text{proj}_w(v) = \text{proj}_{\langle w \rangle}(v).$$

Note that an orthonormal basis for $\langle w \rangle$ is $\frac{w}{\|w\|}$ and hence

$$\text{proj}_w(v) = \left\langle v, \frac{w}{\|w\|} \right\rangle \frac{w}{\|w\|} = \frac{\langle v, w \rangle}{\|w\|^2} w = \frac{\langle v, w \rangle}{\langle w, w \rangle} w.$$

Similarly, if $\{v_1, v_2, \dots, v_n\}$ is an orthogonal basis for a subspace W , then $\left\{ \frac{v_1}{\|v_1\|}, \frac{v_2}{\|v_2\|}, \dots, \frac{v_n}{\|v_n\|} \right\}$ is an orthonormal basis for W

Projection on a vector and orthogonal bases

Let V be an inner product space and $v, w \in V$. Define

$$\text{proj}_w(v) = \text{proj}_{\langle w \rangle}(v).$$

Note that an orthonormal basis for $\langle w \rangle$ is $\frac{w}{\|w\|}$ and hence

$$\text{proj}_w(v) = \left\langle v, \frac{w}{\|w\|} \right\rangle \frac{w}{\|w\|} = \frac{\langle v, w \rangle}{\|w\|^2} w = \frac{\langle v, w \rangle}{\langle w, w \rangle} w.$$

Similarly, if $\{v_1, v_2, \dots, v_n\}$ is an orthogonal basis for a subspace W , then $\left\{ \frac{v_1}{\|v_1\|}, \frac{v_2}{\|v_2\|}, \dots, \frac{v_n}{\|v_n\|} \right\}$ is an orthonormal basis for W and hence

$$\text{proj}_W(v) = \sum_{i=1}^n \left\langle v, \frac{v_i}{\|v_i\|} \right\rangle \frac{v_i}{\|v_i\|} = \sum_{i=1}^n \underbrace{\frac{\langle v, v_i \rangle}{\langle v_i, v_i \rangle} v_i}_{= \text{proj}_{v_i}(v)} = \sum_{i=1}^n \text{proj}_{v_i}(v).$$


Example

Let W be the 2-dimensional subspace of $V = \mathbb{R}^3$ spanned by the orthogonal vectors $v_1 = (1, 2, 1)$ and $v_2 = (1, -1, 1)$.



Example


Let W be the 2-dimensional subspace of $V = \mathbb{R}^3$ spanned by the orthogonal vectors $v_1 = (1, 2, 1)$ and $v_2 = (1, -1, 1)$. What is the projection of $v = (-2, 2, 2)$ on W ?



Example

Let W be the 2-dimensional subspace of $V = \mathbb{R}^3$ spanned by the orthogonal vectors $v_1 = (1, 2, 1)$ and $v_2 = (1, -1, 1)$. What is the projection of $v = (-2, 2, 2)$ on W ?

$$\text{proj}_{v_1} v = \frac{\langle v, v_1 \rangle}{\langle v_1, v_1 \rangle} v_1 = \frac{4}{6}(1, 2, 1) = \frac{2}{3}(1, 2, 1).$$



Example

Let W be the 2-dimensional subspace of $V = \mathbb{R}^3$ spanned by the orthogonal vectors $v_1 = (1, 2, 1)$ and $v_2 = (1, -1, 1)$. What is the projection of $v = (-2, 2, 2)$ on W ?

$$\text{proj}_{v_1} v = \frac{\langle v, v_1 \rangle}{\langle v_1, v_1 \rangle} v_1 = \frac{4}{6}(1, 2, 1) = \frac{2}{3}(1, 2, 1).$$

$$\text{proj}_{v_2} v = \frac{\langle v, v_2 \rangle}{\langle v_2, v_2 \rangle} v_2 = -\frac{2}{3}(1, -1, 1).$$

Example

Let W be the 2-dimensional subspace of $V = \mathbb{R}^3$ spanned by the orthogonal vectors $v_1 = (1, 2, 1)$ and $v_2 = (1, -1, 1)$. What is the projection of $v = (-2, 2, 2)$ on W ?

$$\text{proj}_{v_1} v = \frac{\langle v, v_1 \rangle}{\langle v_1, v_1 \rangle} v_1 = \frac{4}{6}(1, 2, 1) = \frac{2}{3}(1, 2, 1).$$

$$\text{proj}_{v_2} v = \frac{\langle v, v_2 \rangle}{\langle v_2, v_2 \rangle} v_2 = -\frac{2}{3}(1, -1, 1).$$

$$\begin{aligned}\text{Hence } \text{proj}_W(v) &= \text{Proj}_{v_1}(v) + \text{Proj}_{v_2}(v) \\ &= \frac{2}{3}(1, 2, 1) - \frac{2}{3}(1, -1, 1) \\ &= (0, 2, 0).\end{aligned}$$

Projection as a linear transformation


Let V be an inner product space and W be a subspace.



Projection as a linear transformation

Let V be an inner product space and W be a subspace.

Then the projection of vectors in V to W is a linear transformation from V to V with image W .



Projection as a linear transformation

Let V be an inner product space and W be a subspace.

Then the projection of vectors in V to W is a linear transformation from V to V with image W .

Choose o.n. basis $\{w_1, \dots, w_n\}$ for W .

$$\begin{aligned} P_W(v) &= \text{Proj}_W(v) \\ P_W(v_1 + v_2) &= P_W(v_1) + P_W(v_2) \\ \text{Proj}_W(v_1 + v_2) &= \sum_{i=1}^n \langle v_1 + v_2, w_i \rangle w_i \\ &= \sum_{i=1}^n (\langle v_1, w_i \rangle + \langle v_2, w_i \rangle) w_i \\ &= \sum_{i=1}^n \langle v_1, w_i \rangle w_i + \sum_{i=1}^n \langle v_2, w_i \rangle w_i \\ &= c \sum_{i=1}^n \langle v, w_i \rangle w_i \\ P_W(cv) &= c P_W(v) \end{aligned}$$

Denote this linear transformation as P_W .

Some properties of the projection P_W


The linear transformation P_W has some interesting properties (some of which actually characterize it) :



Some properties of the projection P_W

The linear transformation P_W has some interesting properties (some of which actually characterize it) :


- i) $P_W(v) = v$, for all $v \in W$.



Some properties of the projection P_W

The linear transformation P_W has some interesting properties (some of which actually characterize it) :

- i) $P_W(v) = v$, for all $v \in W$.
- ii) $\overline{\text{Range}}(P_W) = W$.



Some properties of the projection P_W

The linear transformation P_W has some interesting properties (some of which actually characterize it) :

- i) $P_W(v) = v$, for all $v \in W$.
- ii) $\text{Range}(P_W) = W$.
- iii) $W^\perp = \{v \mid v \in V, \text{ such that } \langle v, w \rangle = 0 \forall w \in W\}$ is the null space of P_W .

Some properties of the projection P_W


The linear transformation P_W has some interesting properties (some of which actually characterize it) :

- i) $P_W(v) = v$, for all $v \in W$.
- ii) $\text{Range}(P_W) = W$.
- iii) $W^\perp = \{v \mid v \in V, \text{ such that } \langle v, w \rangle = 0 \forall w \in W\}$ is the null space of P_W .
- iv) $P_W^2 = P_W$.

Some properties of the projection P_W

The linear transformation P_W has some interesting properties (some of which actually characterize it) :

- i) $P_W(v) = v$, for all $v \in W$.

- ii) Image(P_W) = W .
- iii) $W^\perp = \{v \mid v \in V, \text{ such that } \langle v, w \rangle = 0 \forall w \in W\}$ is the null space of P_W .
- iv) $P_W^2 = P_W$.

- v) $\|P_W(v)\| \leq \|v\|$.

Thank you

