

What is a basis for a vector space?

Sarang S. Sane

Linear dependence and independence (recall)

Let v_1, v_2, \dots, v_n be a set of vectors in the vector space V .

Linear dependence and independence (recall)

Let v_1, v_2, \dots, v_n be a set of vectors in the vector space V .

The set v_1, v_2, \dots, v_n is said to be **linearly dependent**, if there exist scalars a_1, a_2, \dots, a_n , not all zero, such that

$$a_1 v_1 + a_2 v_2 + \dots + a_n v_n = 0$$

Linear dependence and independence (recall)

Let v_1, v_2, \dots, v_n be a set of vectors in the vector space V .

The set v_1, v_2, \dots, v_n is said to be **linearly dependent**, if there exist scalars a_1, a_2, \dots, a_n , not all zero, such that

$$a_1 v_1 + a_2 v_2 + \dots + a_n v_n = 0$$

The set v_1, v_2, \dots, v_n is said to be **linearly independent**, if the only choice of scalars a_1, a_2, \dots, a_n such that $a_1 v_1 + a_2 v_2 + \dots + a_n v_n = 0$ is with $a_i = 0$ for all i .

Span of a set of vectors

The span of a set S (of vectors) is defined as the set of all finite linear combinations of elements(vectors) of S , and denoted by $\text{Span}(S)$.

Span of a set of vectors

The span of a set S (of vectors) is defined as the set of all finite linear combinations of elements(vectors) of S , and denoted by $Span(S)$.

$$\text{i.e. } Span(S) = \{\sum_{i=1}^n a_i v_i \in V \mid a_1, a_2, \dots, a_n \in \mathbb{R}\}$$

Span of a set of vectors

The span of a set S (of vectors) is defined as the set of all finite linear combinations of elements(vectors) of S , and denoted by $Span(S)$.

$$\text{i.e. } Span(S) = \{\sum_{i=1}^n a_i v_i \in V \mid a_1, a_2, \dots, a_n \in \mathbb{R}\}$$

Example

Let $S = \{(1, 0)\} \subset \mathbb{R}^2$. Then

$$Span(S) = \{a(1, 0) \mid a \in \mathbb{R}\} = \{(a, 0) \mid a \in \mathbb{R}\}$$

Span of a set of vectors

The span of a set S (of vectors) is defined as the set of all finite linear combinations of elements(vectors) of S , and denoted by $Span(S)$.

$$\text{i.e. } Span(S) = \{\sum_{i=1}^n a_i v_i \in V \mid a_1, a_2, \dots, a_n \in \mathbb{R}\}$$

Example

Let $S = \{(1, 0)\} \subset \mathbb{R}^2$. Then

$$Span(S) = \{a(1, 0) \mid a \in \mathbb{R}\} = \{(a, 0) \mid a \in \mathbb{R}\}$$

Thus, $Span(S)$ is the X -axis in \mathbb{R}^2 .

More examples : in \mathbb{R}^2

Let $S = \{(1, 1)\} \subset \mathbb{R}^2$.

More examples : in \mathbb{R}^2


Let $S = \{(1, 1)\} \subset \mathbb{R}^2$.

Then $\text{Span}(S) = \{a(1, 1) | a \in \mathbb{R}\} = \{(a, a) | a \in \mathbb{R}\}$.

More examples : in \mathbb{R}^2

Let $S = \{(1, 1)\} \subset \mathbb{R}^2$.

Then $\text{Span}(S) = \{a(1, 1) | a \in \mathbb{R}\} = \{(a, a) | a \in \mathbb{R}\}$.



More examples : in \mathbb{R}^3

More examples : in \mathbb{R}^3

Let $S = \{(1, 0, 0), (0, 1, 0)\} \subset \mathbb{R}^3$.

More examples : in \mathbb{R}^3

Let $S = \{(1, 0, 0), (0, 1, 0)\} \subset \mathbb{R}^3$. Then

$$\text{Span}(S) = \{a(1, 0, 0) + b(0, 1, 0) \mid a, b \in \mathbb{R}\} = \{(a, b, 0) \mid a, b \in \mathbb{R}\}.$$

More examples : in \mathbb{R}^3

Let $S = \{(1, 0, 0), (0, 1, 0)\} \subset \mathbb{R}^3$. Then

$$\text{Span}(S) = \{a(1, 0, 0) + b(0, 1, 0) \mid a, b \in \mathbb{R}\} = \{(a, b, 0) \mid a, b \in \mathbb{R}\}.$$




Figure: $\text{Span}(S)$ is the XY-plane

Spanning set for a vector space

Let V be a vector space. A set $S \subseteq V$ is a **spanning set** for V if $\text{Span}(S) = V$.

Spanning set for a vector space

Let V be a vector space. A set $S \subseteq V$ is a **spanning set** for V if $\text{Span}(S) = V$.

Example

► If $S = \{(1, 0), (0, 1)\}$ then $\text{Span}(S) = \mathbb{R}^2$

Spanning set for a vector space

Let V be a vector space. A set $S \subseteq V$ is a **spanning set** for V if $\text{Span}(S) = V$.

Example

- ▶ If $S = \{(1, 0), (0, 1)\}$ then $\text{Span}(S) = \mathbb{R}^2$
- ▶ If $S = \{(1, 0), (0, 1), (1, 2)\}$ then $\text{Span}(S) = \mathbb{R}^2$

Spanning set for a vector space

Let V be a vector space. A set $S \subseteq V$ is a **spanning set** for V if $\text{Span}(S) = V$.

Example

- ▶ If $S = \{(1, 0), (0, 1)\}$ then $\text{Span}(S) = \mathbb{R}^2$
- ▶ If $S = \{(1, 0), (0, 1), (1, 2)\}$ then $\text{Span}(S) = \mathbb{R}^2$
- ▶ If $S = \{(1, 1), (0, 1)\}$ then $\text{Span}(S) = \mathbb{R}^2$

Spanning set for a vector space

Let V be a vector space. A set $S \subseteq V$ is a **spanning set** for V if $\text{Span}(S) = V$.

Example

- ▶ If $S = \{(1, 0), (0, 1)\}$ then $\text{Span}(S) = \mathbb{R}^2$
- ▶ If $S = \{(1, 0), (0, 1), \boxed{(1, 2)}\}$ then $\text{Span}(S) = \mathbb{R}^2$
- ▶ If $S = \{(1, 1), (0, 1)\}$ then $\text{Span}(S) = \mathbb{R}^2$
- ▶ If $S = \{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}$ then $\text{Span}(S) = \mathbb{R}^3$

$$\begin{aligned} T &\subseteq S \\ \text{Span}(T) &\subseteq \text{Span}(S) \\ T &\subseteq \text{Span}(S) \\ \text{Span}(T) &\subseteq \text{Span}(S) \end{aligned}$$

$$(x, y) \in \mathbb{R}^2 \quad (x, y) = x(1, 0) + y(0, 1).$$

$$(x, y, z) \in \mathbb{R}^3 \quad (x, y, z) = x(1, 0, 0) + y(0, 1, 0) + z(0, 0, 1).$$

$$\begin{aligned} (1, 0) &= (1, 1) - (0, 1) \therefore (1, 0) \in \text{Span}(\{(1, 1), (0, 1)\}) \\ &\therefore \{(1, 1), (0, 1)\} \subseteq \text{Span} \end{aligned}$$

Example : Adding vectors to obtain a spanning set for \mathbb{R}^3

Example : Adding vectors to obtain a spanning set for \mathbb{R}^3

We will try to "build" a spanning set for the vector space \mathbb{R}^3 .

Example : Adding vectors to obtain a spanning set for \mathbb{R}^3

We will try to "build" a spanning set for the vector space \mathbb{R}^3 .

Start with S_0 to be the empty set \emptyset . Then $\text{Span}(S_0) = \text{Span}(\emptyset) = \{(0, 0, 0)\}$.

Example : Adding vectors to obtain a spanning set for \mathbb{R}^3

We will try to "build" a spanning set for the vector space \mathbb{R}^3 .

Start with S_0 to be the empty set \emptyset . Then $\text{Span}(S_0) = \text{Span}(\emptyset) = \{(0, 0, 0)\}$.

Since this is not the full vector space, append a vector outside $\text{Span}(S_0)$ in \mathbb{R}^3 e.g. $(0, 2, 1)$ to S_0 and call the new set S_1 .

Example : Adding vectors to obtain a spanning set for \mathbb{R}^3

We will try to "build" a spanning set for the vector space \mathbb{R}^3 .

Start with S_0 to be the empty set \emptyset . Then $\text{Span}(S_0) = \text{Span}(\emptyset) = \{(0, 0, 0)\}$.


Since this is not the full vector space, append a vector outside $\text{Span}(S_0)$ in \mathbb{R}^3 e.g. $(0, 2, 1)$ to S_0 and call the new set S_1 .

So $S_1 = S_0 \cup \{(0, 2, 1)\}$.

Example (Contd.)

Example (Contd.)

$\text{Span}(S_1)$ is the line shown in the picture below.



Choose a vector outside $\text{Span}(S_1)$ e.g. $(2, 2, 0)$, append it to S_1 and call the new set S_2 .

Choose a vector outside $\text{Span}(S_1)$ e.g. $(2, 2, 0)$, append it to S_1 and call the new set S_2 .

So $S_2 = S_1 \cup \{(2, 2, 0)\}$.

Choose a vector outside $\text{Span}(S_1)$ e.g. $(2, 2, 0)$, append it to S_1 and call the new set S_2 .


So $S_2 = S_1 \cup \{(2, 2, 0)\}$.

$\text{Span}(S_2)$ is the plane shown in the picture.

Choose a vector outside $\text{Span}(S_1)$ e.g. $(2, 2, 0)$, append it to S_1 and call the new set S_2 .

So $S_2 = S_1 \cup \{(2, 2, 0)\}$.

$\text{Span}(S_2)$ is the plane shown in the picture.



Again choose a vector outside $\text{Span}(S_2)$, e.g. $(0, 0, 5)$, append it to S_2 and call the new set S_3 .

Again choose a vector outside $\text{Span}(S_2)$, e.g. $(0, 0, 5)$, append it to S_2 and call the new set S_3 .

So $S_3 = S_2 \cup \{(0, 0, 5)\}$.

Again choose a vector outside $\text{Span}(S_2)$, e.g. $(0, 0, 5)$, append it to S_2 and call the new set S_3 .

So $S_3 = S_2 \cup \{(0, 0, 5)\}$.

Any arbitrary vector $(x, y, z) \in \mathbb{R}^3$ can be written as follows:

Again choose a vector outside $\text{Span}(S_2)$, e.g. $(0, 0, 5)$, append it to S_2 and call the new set S_3 .

So $S_3 = S_2 \cup \{(0, 0, 5)\}$.

Any arbitrary vector $(x, y, z) \in \mathbb{R}^3$ can be written as follows:

$$(x, y, z) = \frac{y-x}{2}(0, 2, 1) + \frac{x}{2}(2, 2, 0) + \frac{x-y+2z}{10}(0, 0, 5)$$

Again choose a vector outside $\text{Span}(S_2)$, e.g. $(0, 0, 5)$, append it to S_2 and call the new set S_3 .

So $S_3 = S_2 \cup \{(0, 0, 5)\}$.

Any arbitrary vector $(x, y, z) \in \mathbb{R}^3$ can be written as follows:

$$(x, y, z) = \frac{y-x}{2}(0, 2, 1) + \frac{x}{2}(2, 2, 0) + \frac{x-y+2z}{10}(0, 0, 5)$$

Hence

$$\text{Span}(S_3) = \mathbb{R}^3$$

Another example

Another example

Start with S_0 to be the empty set \emptyset as before.

Another example

Start with S_0 to be the empty set \emptyset as before.

Thus $S_0 = \emptyset$ and hence $\text{Span}(S_0) = \text{Span}(\emptyset) = \{(0, 0, 0)\}$.

Another example

Start with S_0 to be the empty set \emptyset as before.

Thus $S_0 = \emptyset$ and hence $\text{Span}(S_0) = \text{Span}(\emptyset) = \{(0, 0, 0)\}$.

Append any vector not in $\text{Span}(S_0)$ e.g. $(3, 0, 0)$ to S_0 and call the new set S_1 .

Another example

Start with S_0 to be the empty set \emptyset as before.


Thus $S_0 = \emptyset$ and hence $\text{Span}(S_0) = \text{Span}(\emptyset) = \{(0, 0, 0)\}$.

Append any vector not in $\text{Span}(S_0)$ e.g. $(3, 0, 0)$ to S_0 and call the new set S_1 .

Hence $S_1 = S_0 \cup \{(3, 0, 0)\}$.

Example (contd.)

$\text{Span}(S_1)$ is the X -axis, as shown below.



Example (contd.)

Choose a vector outside $\text{Span}(S_1)$ e.g. $(2, 2, 1)$ and append it to S_1 and call the new set S_2 .

Example (contd.)


Choose a vector outside $\text{Span}(S_1)$ e.g. $(2, 2, 1)$ and append it to S_1 and call the new set S_2 .

Then $S_2 = S_1 \cup \{(2, 2, 1)\}$ and $\text{Span}(S_2)$ is the plane shown below.

Example (contd.)

Choose a vector outside $\text{Span}(S_1)$ e.g. $(2, 2, 1)$ and append it to S_1 and call the new set S_2 .

Then $S_2 = S_1 \cup \{(2, 2, 1)\}$ and $\text{Span}(S_2)$ is the plane shown below.



Again choose a vector outside $\text{Span}(S_2)$ e.g. $(1, 3, 3)$, append it to S_2 and call the new set S_3 .

Again choose a vector outside $\text{Span}(S_2)$ e.g. $(1, 3, 3)$, append it to S_2 and call the new set S_3 .

Then $S_3 = S_2 \cup \{(1, 3, 3)\}$.

Again choose a vector outside $\text{Span}(S_2)$ e.g. $(1, 3, 3)$, append it to S_2 and call the new set S_3 .

Then $S_3 = S_2 \cup \{(1, 3, 3)\}$.

Any arbitrary vector $(x, y, z) \in \mathbb{R}^3$ can be written as follows:

$$(x, y, z) = \frac{3x-5y+4z}{9}(3, 0, 0) + (y-z)(2, 2, 1) + \frac{2z-y}{3}(1, 3, 3)$$

Again choose a vector outside $\text{Span}(S_2)$ e.g. $(1, 3, 3)$, append it to S_2 and call the new set S_3 .

Then $S_3 = S_2 \cup \{(1, 3, 3)\}$.

Any arbitrary vector $(x, y, z) \in \mathbb{R}^3$ can be written as follows:

$$(x, y, z) = \frac{3x-5y+4z}{9}(3, 0, 0) + (y-z)(2, 2, 1) + \frac{2z-y}{3}(1, 3, 3)$$

Hence

$$\text{Span}(S_3) = \mathbb{R}^3$$

What is a basis?

A basis B of a vector space V is a linearly independent subset of V that spans V .

What is a basis?

A basis B of a vector space V is a linearly independent subset of V that spans V .

Example

Let $e_i \in \mathbb{R}^n$ be the vector with i^{th} coordinate 1 and all other coordinates 0

What is a basis?

A basis B of a vector space V is a linearly independent subset of V that spans V .

Example

Let $e_i \in \mathbb{R}^n$ be the vector with i^{th} coordinate 1 and all other coordinates 0 e.g. $e_1 = (1, 0, 0, \dots, 0)$.

What is a basis?

A basis B of a vector space V is a linearly independent subset of V that spans V .

Example

Let $e_i \in \mathbb{R}^n$ be the vector with i^{th} coordinate 1 and all other coordinates 0 e.g. $e_1 = (1, 0, 0, \dots, 0)$.

The set $\varepsilon = \{e_1, e_2, \dots, e_n\} \subseteq \mathbb{R}^n$ is a basis for \mathbb{R}^n . ~~consisting of~~

$$(x_1, x_2, \dots, x_n) \\ = x_1 e_1 + x_2 e_2 + \dots + x_n e_n$$

$$\therefore \text{Span}(\varepsilon) = \mathbb{R}^n.$$

$$\sum_{i=1}^n a_i e_i = 0 \Rightarrow \begin{array}{l} i^{\text{th}} \text{ coordinate of LHS is } a_i \\ \Rightarrow a_j = 0 \quad \forall j. \end{array}$$

$\therefore \varepsilon$ is lin. indep.

Equivalent conditions for B to be a basis

The following conditions are equivalent to a subset $B \subseteq V$ being a basis :

Equivalent conditions for B to be a basis

The following conditions are equivalent to a subset $B \subseteq V$ being a basis :

- i) B is linearly independent and $\text{Span}(B) = V$.

Equivalent conditions for B to be a basis

The following conditions are equivalent to a subset $B \subseteq V$ being a basis :

- i) B is linearly independent and $\text{Span}(B) = V$.
- ii) B is a maximal linearly independent set.

Equivalent conditions for B to be a basis

The following conditions are equivalent to a subset $B \subseteq V$ being a basis :

- i) B is linearly independent and $\text{Span}(B) = V$.
- ii) B is a maximal linearly independent set.
- iii) B is a minimal spanning set.

Suppose B is a basis.
 $\therefore B$ is lin. indept.
Suppose $B' = B \cup \{v\}$.
 $\therefore v = \sum_{i=1}^n a_i v_i$ where $v_1, \dots, v_n \in B$.
 $\therefore B'$ is a lin. dep. set.

maximal lin.
indept. means

① it is lin. indept.

② appending any
vector makes
it lin. dep.

minimal spanning
means

① it is spanning

② it is no longer
spanning if
we delete any
vector

How do we find a basis?

How do we find a basis?

We can find a basis by any one of the methods described below :

How do we find a basis?

We can find a basis by any one of the methods described below :

- i) Start with the \emptyset and keep appending vectors which are not in the span of the set thus far obtained, until we obtain a spanning set.

How do we find a basis?

We can find a basis by any one of the methods described below :

- i) Start with the \emptyset and keep appending vectors which are not in the span of the set thus far obtained, until we obtain a spanning set.

Examples 1 & 2

- ii) Take a spanning set and keep deleting vectors which are linear combinations of the other vectors, until the remaining vectors satisfy that they are not a linear combination of the other remaining ones.

Example : Method 1 : $V = \mathbb{R}^2$

Let us start with the empty set and append a non-zero vector e.g. $(1, 2)$.

Example : Method 1 : $V = \mathbb{R}^2$

Let us start with the empty set and append a non-zero vector e.g. $(1, 2)$.

Now choose another vector which is not in the span of the earlier vector e.g. $(2, 3)$.

Example : Method 1 : $V = \mathbb{R}^2$

Let us start with the empty set and append a non-zero vector e.g. $(1, 2)$.

Now choose another vector which is not in the span of the earlier vector e.g. $(2, 3)$.

$$\text{Span}(\{(1, 2), (2, 3)\}) = \mathbb{R}^2.$$

Example : Method 1 : $V = \mathbb{R}^2$

Let us start with the empty set and append a non-zero vector e.g. $(1, 2)$.

Now choose another vector which is not in the span of the earlier vector e.g. $(2, 3)$.

$$\text{Span}(\{(1, 2), (2, 3)\}) = \mathbb{R}^2.$$

Hence this set forms a basis for \mathbb{R}^2 .

Example : Method 2 : $V = \mathbb{R}^3$

Let us start with the set

$$S = \{(1, 0, 0), (1, 2, 0), (1, 0, 3), (0, 2, 3), (0, 4, 2)\}$$

Example : Method 2 : $V = \mathbb{R}^3$

Let us start with the set

$$S = \{(1, 0, 0), (1, 2, 0), (1, 0, 3), (0, 2, 3), (0, 4, 2)\}$$

Check that $\text{Span}(S) = \mathbb{R}^3$.

Example : Method 2 : $V = \mathbb{R}^3$

Let us start with the set

$$S = \{(1, 0, 0), (1, 2, 0), (1, 0, 3), (0, 2, 3), (0, 4, 2)\}$$

Check that $\text{Span}(S) = \mathbb{R}^3$.

Now observe that, $(0, 4, 2) = 2(1, 2, 0) + \frac{2}{3}(1, 0, 3) - \frac{8}{3}(1, 0, 0)$.

Example : Method 2 : $V = \mathbb{R}^3$

Let us start with the set

$$S = \{(1, 0, 0), (1, 2, 0), (1, 0, 3), (0, 2, 3), (0, 4, 2)\}$$

Check that $\text{Span}(S) = \mathbb{R}^3$.

Now observe that, $(0, 4, 2) = 2(1, 2, 0) + \frac{2}{3}(1, 0, 3) - \frac{8}{3}(1, 0, 0)$.

So delete $(0, 4, 2)$.

Example : Method 2 : $V = \mathbb{R}^3$

Let us start with the set

$$S = \{(1, 0, 0), (1, 2, 0), (1, 0, 3), (0, 2, 3), (0, 4, 2)\}$$

Check that $\text{Span}(S) = \mathbb{R}^3$.

Now observe that, $(0, 4, 2) = 2(1, 2, 0) + \frac{2}{3}(1, 0, 3) - \frac{8}{3}(1, 0, 0)$.

So delete $(0, 4, 2)$.

Hence our new set of vectors is

$$S_1 = \{(1, 0, 0), (1, 2, 0), (1, 0, 3), (0, 2, 3)\}$$

Method 2 (contd.)

Observe that $(0, 2, 3) = (1, 2, 0) + (1, 0, 3) - 2(1, 0, 0)$.

Method 2 (contd.)

Observe that $(0, 2, 3) = (1, 2, 0) + (1, 0, 3) - 2(1, 0, 0)$.

Hence delete $(0, 2, 3)$.

Method 2 (contd.)

Observe that $(0, 2, 3) = (1, 2, 0) + (1, 0, 3) - 2(1, 0, 0)$.

Hence delete $(0, 2, 3)$.

Hence our new set of vectors is

$$S_2 = \{(1, 0, 0), (1, 1, 0), (1, 0, 1)\}$$

Method 2 (contd.)

Observe that $(0, 2, 3) = (1, 2, 0) + (1, 0, 3) - 2(1, 0, 0)$.

Hence delete $(0, 2, 3)$.

Hence our new set of vectors is

$$S_2 = \{(1, 0, 0), (1, 1, 0), (1, 0, 1)\}$$

None of these vectors is a linear combination of the other two vectors.

Method 2 (contd.)

Observe that $(0, 2, 3) = (1, 2, 0) + (1, 0, 3) - 2(1, 0, 0)$.

Hence delete $(0, 2, 3)$.

Hence our new set of vectors is

$$S_2 = \{(1, 0, 0), (1, 1, 0), (1, 0, 1)\}$$

None of these vectors is a linear combination of the other two vectors.

Hence S_2 forms a basis of R^3 .

Thank you