

Linear independence

Sarang S. Sane

Linear dependence (recall)

A set of vectors v_1, v_2, \dots, v_n from a vector space V is said to be **linearly dependent** if there exists scalars a_1, a_2, \dots, a_n , not all zero, such that

$$a_1 v_1 + a_2 v_2 + \dots + a_n v_n = 0$$

Linear dependence (recall)

A set of vectors v_1, v_2, \dots, v_n from a vector space V is said to be **linearly dependent** if there exists scalars a_1, a_2, \dots, a_n , not all zero, such that

$$a_1 v_1 + a_2 v_2 + \dots + a_n v_n = 0$$

Equivalently : v_1, v_2, \dots, v_n are **linearly dependent** if the 0 vector can be expressed as a linear combination of v_1, v_2, \dots, v_n with non-zero coefficients (i.e. at least one coefficient is non-zero).

Definition of linear independence

A set of vectors v_1, v_2, \dots, v_n from a vector space V is said to be **linearly independent** if v_1, v_2, \dots, v_n are not linearly dependent.

Definition of linear independence

A set of vectors v_1, v_2, \dots, v_n from a vector space V is said to be **linearly independent** if v_1, v_2, \dots, v_n are not linearly dependent.

Equivalently : A set of vectors v_1, v_2, \dots, v_n from a vector space V is said to be **linearly independent**, if the equation

$$a_1 v_1 + a_2 v_2 + \dots + a_n v_n = 0$$

can only be satisfied when $a_i = 0$ for all $i = 1, 2, \dots, n$.

Definition of linear independence

A set of vectors v_1, v_2, \dots, v_n from a vector space V is said to be **linearly independent** if v_1, v_2, \dots, v_n are not linearly dependent.

Equivalently : A set of vectors v_1, v_2, \dots, v_n from a vector space V is said to be **linearly independent**, if the equation

$$a_1 v_1 + a_2 v_2 + \dots + a_n v_n = 0$$

can only be satisfied when $a_i = 0$ for all $i = 1, 2, \dots, n$.

Equivalently : A set of vectors v_1, v_2, \dots, v_n from a vector space V is said to be **linearly independent** if the only linear combination of v_1, v_2, \dots, v_n which equals 0 is the linear combination with all coefficients 0.

Example in \mathbb{R}^2

Consider the two vectors $(-1, 3)$ and $(2, 0)$ in \mathbb{R}^2 .

Example in \mathbb{R}^2

Consider the two vectors $(-1, 3)$ and $(2, 0)$ in \mathbb{R}^2 .

Consider the following equation :

$$a(-1, 3) + b(2, 0) = (0, 0)$$

Example in \mathbb{R}^2

Consider the two vectors $(-1, 3)$ and $(2, 0)$ in \mathbb{R}^2 .

Consider the following equation :

$$a(-1, 3) + b(2, 0) = (0, 0)$$

Hence we have the following system of linear equations:

$$-a + 2b = 0 \text{ and } 3a = 0.$$

Example in \mathbb{R}^2

Consider the two vectors $(-1, 3)$ and $(2, 0)$ in \mathbb{R}^2 .

Consider the following equation :

$$a(-1, 3) + b(2, 0) = (0, 0)$$

Hence we have the following system of linear equations:

$$-a + 2b = 0 \text{ and } 3a = 0.$$

Hence $a = 0$, $b = 0$ is the unique solution of the system of linear equations, which implies that the vectors $(-1, 3)$ and $(2, 0)$ are linearly independent.

The 0 vector

Let v_1, v_2, \dots, v_n be a set of vectors containing the 0 vector.

The 0 vector

Let v_1, v_2, \dots, v_n be a set of vectors containing the 0 vector.

Suppose $v_i = 0$. Then we can choose $a_i = 1$ and $a_j = 0$ for $j \neq i$.

The 0 vector

Let v_1, v_2, \dots, v_n be a set of vectors containing the 0 vector.

Suppose $v_i = 0$. Then we can choose $a_i = 1$ and $a_j = 0$ for $j \neq i$.

Then the linear combination $a_1 v_1 + a_2 v_2 + \dots + a_n v_n$ is 0 but not all coefficients are 0.

The 0 vector

Let v_1, v_2, \dots, v_n be a set of vectors containing the 0 vector.

Suppose $v_i = 0$. Then we can choose $a_i = 1$ and $a_j = 0$ for $j \neq i$.

Then the linear combination $a_1 v_1 + a_2 v_2 + \dots + a_n v_n$ is 0 but not all coefficients are 0.

Hence, a set of vectors v_1, v_2, \dots, v_n containing the 0 vector is always a linearly dependent set.

When are two non-zero vectors linearly independent?

Let v_1 and v_2 be two non-zero vectors.

When are two non-zero vectors linearly independent?

Let v_1 and v_2 be two non-zero vectors.

Suppose that v_1 and v_2 are linearly dependent.

When are two non-zero vectors linearly independent?

Let v_1 and v_2 be two non-zero vectors.

Suppose that v_1 and v_2 are linearly dependent.

Then $a_1 v_1 + a_2 v_2 = 0$ for some coefficients a_1 and a_2 .

at least
one of
 a_1 or a_2 is
not zero

When are two non-zero vectors linearly independent?

Let v_1 and v_2 be two non-zero vectors.

Suppose that v_1 and v_2 are linearly dependent.

Then $a_1 v_1 + a_2 v_2 = 0$ for some coefficients a_1 and a_2 .

Note that since the vectors are non-zero, both a_1 and a_2 must be non-zero.

When are two non-zero vectors linearly independent?

Let v_1 and v_2 be two non-zero vectors.

Suppose that v_1 and v_2 are linearly dependent.

Then $a_1 v_1 + a_2 v_2 = 0$ for some coefficients a_1 and a_2 .

Note that since the vectors are non-zero, both a_1 and a_2 must be non-zero.

Dividing by a_1 and putting $c = -a_2/a_1$, we get that $v_1 = cv_2$.

When are two non-zero vectors linearly independent?

Let v_1 and v_2 be two non-zero vectors.

Suppose that v_1 and v_2 are linearly dependent.

Then $a_1 v_1 + a_2 v_2 = 0$ for some coefficients a_1 and a_2 .

Note that since the vectors are non-zero, both a_1 and a_2 must be non-zero.

Dividing by a_1 and putting $c = -a_2/a_1$, we get that $v_1 = cv_2$.

Hence v_1 and v_2 are multiples of each other.

When are two non-zero vectors linearly independent?

Let v_1 and v_2 be two non-zero vectors.

Suppose that v_1 and v_2 are linearly dependent.

Then $a_1 v_1 + a_2 v_2 = 0$ for some coefficients a_1 and a_2 .

Note that since the vectors are non-zero, both a_1 and a_2 must be non-zero.

Dividing by a_1 and putting $c = -a_2/a_1$, we get that $v_1 = cv_2$.

Hence v_1 and v_2 are multiples of each other.

We can reverse the implications above and conclude that if v_1 and v_2 are multiples of each other then they are linearly dependent.

When are two non-zero vectors linearly independent?

Let v_1 and v_2 be two non-zero vectors.

Suppose that v_1 and v_2 are linearly dependent.

Then $a_1 v_1 + a_2 v_2 = 0$ for some coefficients a_1 and a_2 . *(where at least one of a_1 or a_2 is not 0)*

Note that since the vectors are non-zero, both a_1 and a_2 must be non-zero.

Dividing by a_1 and putting $c = -a_2/a_1$, we get that $v_1 = cv_2$.

Hence v_1 and v_2 are multiples of each other.

We can reverse the implications above and conclude that if v_1 and v_2 are multiples of each other then they are linearly dependent.

Conclusion : Two non-zero vectors are **linearly independent** precisely when they are **not multiples of each other** .

Linear independence of three vectors

Suppose v_1, v_2 and v_3 are linearly dependent.

Linear independence of three vectors

Suppose v_1, v_2 and v_3 are linearly dependent.

Then $a_1 v_1 + a_2 v_2 + a_3 v_3 = 0$ for some coefficients a_1, a_2, a_3 where at least one of the coefficients is non-zero.

Linear independence of three vectors

Suppose v_1, v_2 and v_3 are linearly dependent.

Then $a_1 v_1 + a_2 v_2 + a_3 v_3 = 0$ for some coefficients a_1, a_2, a_3 where at least one of the coefficients is non-zero.

If $a_1 \neq 0$, then we can write $v_1 = b_2 v_2 + b_3 v_3$ where $b_2 = -a_2/a_1$ and $b_3 = -a_3/a_1$.

Linear independence of three vectors

Suppose v_1, v_2 and v_3 are linearly dependent.

Then $a_1 v_1 + a_2 v_2 + a_3 v_3 = 0$ for some coefficients a_1, a_2, a_3 where at least one of the coefficients is non-zero.

If $a_1 \neq 0$, then we can write $v_1 = b_2 v_2 + b_3 v_3$ where $b_2 = -a_2/a_1$ and $b_3 = -a_3/a_1$. Hence, v_1 is a linear combination of the other two vectors.

Linear independence of three vectors

Suppose v_1, v_2 and v_3 are linearly dependent.

Then $a_1 v_1 + a_2 v_2 + a_3 v_3 = 0$ for some coefficients a_1, a_2, a_3 where at least one of the coefficients is non-zero.

If $a_1 \neq 0$, then we can write $v_1 = b_2 v_2 + b_3 v_3$ where $b_2 = -a_2/a_1$ and $b_3 = -a_3/a_1$. Hence, v_1 is a linear combination of the other two vectors.

Similarly if $a_2 \neq 0$, v_2 is a linear combination of the other two vectors and if $a_3 \neq 0$, v_3 is a linear combination of the other two vectors.

Linear independence of three vectors

Suppose v_1, v_2 and v_3 are linearly dependent.

Then $a_1 v_1 + a_2 v_2 + a_3 v_3 = 0$ for some coefficients a_1, a_2, a_3 where at least one of the coefficients is non-zero.

If $a_1 \neq 0$, then we can write $v_1 = b_2 v_2 + b_3 v_3$ where $b_2 = -a_2/a_1$ and $b_3 = -a_3/a_1$. Hence, v_1 is a linear combination of the other two vectors.

Similarly if $a_2 \neq 0$, v_2 is a linear combination of the other two vectors and if $a_3 \neq 0$, v_3 is a linear combination of the other two vectors.

Since the implications are reversible, we obtain that v_1, v_2 and v_3 are linearly dependent exactly when one of the vectors is a linear combination of the others.

Linear independence of three vectors

Suppose v_1, v_2 and v_3 are linearly dependent.

Then $a_1 v_1 + a_2 v_2 + a_3 v_3 = 0$ for some coefficients a_1, a_2, a_3 where at least one of the coefficients is non-zero.

If $a_1 \neq 0$, then we can write $v_1 = b_2 v_2 + b_3 v_3$ where $b_2 = -a_2/a_1$ and $b_3 = -a_3/a_1$. Hence, v_1 is a linear combination of the other two vectors.

Similarly if $a_2 \neq 0$, v_2 is a linear combination of the other two vectors and if $a_3 \neq 0$, v_3 is a linear combination of the other two vectors.

Since the implications are reversible, we obtain that v_1, v_2 and v_3 are linearly dependent exactly when one of the vectors is a linear combination of the others.

Conclusion : If three vectors are linearly independent then none of these vectors is a linear combination of the other two.

Example in \mathbb{R}^3

Let us consider three vectors $(1, 1, 2)$, $(1, 2, 0)$ and $(0, 2, 1)$ in \mathbb{R}^3 and also consider the following equation:

$$a(1, 1, 2) + b(1, 2, 0) + c(0, 2, 1) = (0, 0, 0)$$

Example in \mathbb{R}^3

Let us consider three vectors $(1, 1, 2)$, $(1, 2, 0)$ and $(0, 2, 1)$ in \mathbb{R}^3 and also consider the following equation:

$$a(1, 1, 2) + b(1, 2, 0) + c(0, 2, 1) = (0, 0, 0)$$

Hence we have the following system of linear equations:

$$a + b = 0 \quad a + 2b + 2c = 0 \quad 2a + c = 0.$$

Example in \mathbb{R}^3

Let us consider three vectors $(1, 1, 2)$, $(1, 2, 0)$ and $(0, 2, 1)$ in \mathbb{R}^3 and also consider the following equation:

$$a(1, 1, 2) + b(1, 2, 0) + c(0, 2, 1) = (0, 0, 0)$$

Hence we have the following system of linear equations:

$$a + b = 0 \quad a + 2b + 2c = 0 \quad 2a + c = 0.$$

Substituting $b = -a$ and $c = -2a$ in the middle equation yields that $a = 0$, $b = 0$, $c = 0$ is the unique solution of this system. Hence the vectors $(1, 1, 2)$, $(1, 2, 0)$ and $(0, 2, 1)$ are linearly independent.

How to check linear independence in \mathbb{R}^m

How do we check if $v_1, v_2, \dots, v_n \in \mathbb{R}^m$ are linearly independent?

How to check linear independence in \mathbb{R}^m

How do we check if $v_1, v_2, \dots, v_n \in \mathbb{R}^m$ are linearly independent?

In terms of coordinates, let $v_j = (v_{1j}, v_{2j}, \dots, v_{mj}) ; j = 1, 2, \dots, n$.

How to check linear independence in \mathbb{R}^m

How do we check if $v_1, v_2, \dots, v_n \in \mathbb{R}^m$ are linearly independent?

In terms of coordinates, let $v_j = (v_{1j}, v_{2j}, \dots, v_{mj}) ; j = 1, 2, \dots, n$.

Let us write the linear combination of these vectors with *arbitrary* coefficients a_1, a_2, \dots, a_n and equate it to 0 :

$$a_1 v_1 + a_2 v_2 + \dots + a_n v_n = 0.$$

How to check linear independence in \mathbb{R}^m

How do we check if $v_1, v_2, \dots, v_n \in \mathbb{R}^m$ are linearly independent?

In terms of coordinates, let $v_j = (v_{1j}, v_{2j}, \dots, v_{mj}) ; j = 1, 2, \dots, n$.

Let us write the linear combination of these vectors with *arbitrary* coefficients a_1, a_2, \dots, a_n and equate it to 0 :

$$a_1 v_1 + a_2 v_2 + \dots + a_n v_n = 0.$$

Considering each coordinate, we have the following identities :

$$v_{11}a_1 + v_{12}a_2 + \dots + v_{1n}a_n = 0$$

$$v_{21}a_1 + v_{22}a_2 + \dots + v_{2n}a_n = 0$$

$$\vdots \quad \quad \quad \vdots \quad \quad \quad \ddots \quad \quad \quad \vdots \quad \quad \quad \vdots$$

$$v_{m1}a_1 + v_{m2}a_2 + \dots + v_{mn}a_n = 0$$

Since the a_i are arbitrary (unknown), we can treat this like a homogeneous system of linear equations with coefficients v_{ij} and unknowns a_i .

Since the a_i are arbitrary (unknown), we can treat this like a homogeneous system of linear equations with coefficients v_{ij} and unknowns a_i .

For linear independence, we have to check if the only choice of a_i 's satisfying the above identities is $a_i = 0$.

Since the a_i are arbitrary (unknown), we can treat this like a homogeneous system of linear equations with coefficients v_{ij} and unknowns a_i .

For linear independence, we have to check if the only choice of a_i 's satisfying the above identities is $a_i = 0$.

Equivalently, in terms of the homogeneous system of linear equations, we have to check that its only solution is the 0 solution.

Since the a_i are arbitrary (unknown), we can treat this like a homogeneous system of linear equations with coefficients v_{ij} and unknowns a_i .

For linear independence, we have to check if the only choice of a_i 's satisfying the above identities is $a_i = 0$.

Equivalently, in terms of the homogeneous system of linear equations, we have to check that its only solution is the 0 solution.

Conclusion : To check $v_1, v_2, \dots, v_n \in \mathbb{R}^m$ are linearly independent, we have to check that the homogeneous system of linear equations $Vx = 0$ has only the trivial solution, where the j^{th} column of V is v_j .

Example : 2×2

Consider the two vectors $(5, 2)$ and $(1, 3)$ in \mathbb{R}^2 .

Example : 2×2

Consider the two vectors $(5, 2)$ and $(1, 3)$ in \mathbb{R}^2 . Write the linear combination of these two vectors with *unknown* coefficients x_1 and x_2 and equate it to 0 : $x_1(5, 2) + x_2(1, 3) = (0, 0)$.

Example : 2×2

Consider the two vectors $(5, 2)$ and $(1, 3)$ in \mathbb{R}^2 . Write the linear combination of these two vectors with *unknown* coefficients x_1 and x_2 and equate it to 0 : $x_1(5, 2) + x_2(1, 3) = (0, 0)$.

Hence we have the system of linear equations:

$$5x_1 + x_2 = 0$$

$$2x_1 + 3x_2 = 0$$

Example : 2×2

Consider the two vectors $(5, 2)$ and $(1, 3)$ in \mathbb{R}^2 . Write the linear combination of these two vectors with *unknown* coefficients x_1 and x_2 and equate it to 0 : $x_1(5, 2) + x_2(1, 3) = (0, 0)$.

Hence we have the system of linear equations:

$$5x_1 + x_2 = 0$$

$$2x_1 + 3x_2 = 0$$

Since the corresponding matrix $\begin{bmatrix} 5 & 1 \\ 2 & 3 \end{bmatrix}$ is invertible, the system of linear equations has a unique solution $x_1 = x_2 = 0$.

Example : 2×2

Consider the two vectors $(5, 2)$ and $(1, 3)$ in \mathbb{R}^2 . Write the linear combination of these two vectors with *unknown* coefficients x_1 and x_2 and equate it to 0 : $x_1(5, 2) + x_2(1, 3) = (0, 0)$.

Hence we have the system of linear equations:

$$5x_1 + x_2 = 0$$

$$2x_1 + 3x_2 = 0$$

Since the corresponding matrix $\begin{bmatrix} 5 & 1 \\ 2 & 3 \end{bmatrix}$ is invertible, the system of linear equations has a unique solution $x_1 = x_2 = 0$.

Hence the vectors $(5, 2)$ and $(1, 3)$ are linearly independent.

Example : 3×2

Consider the two vectors $(1, 2, 0)$ and $(3, 3, 5)$ in \mathbb{R}^3 .

Example : 3×2

Consider the two vectors $(1, 2, 0)$ and $(3, 3, 5)$ in \mathbb{R}^3 . Write the linear combination of these two vectors with *unknown* coefficients x_1 and x_2 and equate it to 0 : $x_1(1, 2, 0) + x_2(3, 3, 5) = (0, 0, 0)$.

Example : 3×2

Consider the two vectors $(1, 2, 0)$ and $(3, 3, 5)$ in \mathbb{R}^3 . Write the linear combination of these two vectors with *unknown* coefficients x_1 and x_2 and equate it to 0 : $x_1(1, 2, 0) + x_2(3, 3, 5) = (0, 0, 0)$.

Hence we have the system of linear equations:

$$x_1 + 3x_2 = 0$$

$$2x_1 + 3x_2 = 0$$

$$0x_1 + 5x_2 = 0$$

Example : 3×2

Consider the two vectors $(1, 2, 0)$ and $(3, 3, 5)$ in \mathbb{R}^3 . Write the linear combination of these two vectors with *unknown* coefficients x_1 and x_2 and equate it to 0 : $x_1(1, 2, 0) + x_2(3, 3, 5) = (0, 0, 0)$.

Hence we have the system of linear equations:

$$x_1 + 3x_2 = 0$$

$$2x_1 + 3x_2 = 0$$

$$0x_1 + 5x_2 = 0$$

It is easy to check that the system of linear equations has a unique solution $x_1 = x_2 = 0$ (or we can use Gaussian elimination and check that the only solution is the trivial one).

Example : 3×2

Consider the two vectors $(1, 2, 0)$ and $(3, 3, 5)$ in \mathbb{R}^3 . Write the linear combination of these two vectors with *unknown* coefficients x_1 and x_2 and equate it to 0 : $x_1(1, 2, 0) + x_2(3, 3, 5) = (0, 0, 0)$.

Hence we have the system of linear equations:

$$x_1 + 3x_2 = 0$$

$$2x_1 + 3x_2 = 0$$

$$0x_1 + 5x_2 = 0$$

It is easy to check that the system of linear equations has a unique solution $x_1 = x_2 = 0$ (or we can use Gaussian elimination and check that the only solution is the trivial one).

Hence the vectors $(1, 2, 0)$ and $(3, 3, 5)$ are linearly independent.

Example : 2×3

Consider the three vectors $(1, 2)$, $(1, 3)$ and $(3, 4)$ in \mathbb{R}^2 .

Example : 2×3

Consider the three vectors $(1, 2)$, $(1, 3)$ and $(3, 4)$ in \mathbb{R}^2 . Equate the linear combination of these three vectors with *unknown* coefficients x_1, x_2 and x_3 to 0 : $x_1(1, 2) + x_2(1, 3) + x_3(3, 4) = (0, 0)$.

Example : 2×3

Consider the three vectors $(1, 2)$, $(1, 3)$ and $(3, 4)$ in \mathbb{R}^2 . Equate the linear combination of these three vectors with *unknown* coefficients x_1, x_2 and x_3 to 0 : $x_1(1, 2) + x_2(1, 3) + x_3(3, 4) = (0, 0)$.

Hence we have the system of linear equations:

$$1x_1 + 1x_2 + 3x_3 = 0$$

$$2x_1 + 3x_2 + 4x_3 = 0$$

Example : 2×3

Consider the three vectors $(1, 2)$, $(1, 3)$ and $(3, 4)$ in \mathbb{R}^2 . Equate the linear combination of these three vectors with *unknown* coefficients x_1, x_2 and x_3 to 0 : $x_1(1, 2) + x_2(1, 3) + x_3(3, 4) = (0, 0)$.

Hence we have the system of linear equations:

$$1x_1 + 1x_2 + 3x_3 = 0$$

$$2x_1 + 3x_2 + 4x_3 = 0$$

The augmented matrix for this system is $\left[\begin{array}{ccc|c} 1 & 1 & 3 & 0 \\ 2 & 3 & 4 & 0 \end{array} \right]$.

Example : 2×3

Consider the three vectors $(1, 2)$, $(1, 3)$ and $(3, 4)$ in \mathbb{R}^2 . Equate the linear combination of these three vectors with *unknown* coefficients x_1, x_2 and x_3 to 0 : $x_1(1, 2) + x_2(1, 3) + x_3(3, 4) = (0, 0)$.

Hence we have the system of linear equations:

$$1x_1 + 1x_2 + 3x_3 = 0$$

$$2x_1 + 3x_2 + 4x_3 = 0$$

The augmented matrix for this system is $\left[\begin{array}{ccc|c} 1 & 1 & 3 & 0 \\ 2 & 3 & 4 & 0 \end{array} \right]$.

Gaussian elimination yields infinitely many solutions for this system of the form $x_1 = -5c, x_2 = 2c, x_3 = c$ where $c \in \mathbb{R}$.

Example : 2×3

Consider the three vectors $(1, 2)$, $(1, 3)$ and $(3, 4)$ in \mathbb{R}^2 . Equate the linear combination of these three vectors with *unknown* coefficients x_1, x_2 and x_3 to 0 : $x_1(1, 2) + x_2(1, 3) + x_3(3, 4) = (0, 0)$.

Hence we have the system of linear equations:

$$1x_1 + 1x_2 + 3x_3 = 0$$

$$2x_1 + 3x_2 + 4x_3 = 0$$

The augmented matrix for this system is $\left[\begin{array}{ccc|c} 1 & 1 & 3 & 0 \\ 2 & 3 & 4 & 0 \end{array} \right]$.

Gaussian elimination yields infinitely many solutions for this system of the form $x_1 = -5c, x_2 = 2c, x_3 = c$ where $c \in \mathbb{R}$. **Hence the vectors $(1, 2)$, $(1, 3)$ and $(3, 4)$ are linearly dependent.**

Example : 3×3

Consider the three vectors $(1, 2, 0)$, $(0, 2, 4)$ and $(3, 0, 0)$ in \mathbb{R}^3 .

Example : 3×3

Consider the three vectors $(1, 2, 0)$, $(0, 2, 4)$ and $(3, 0, 0)$ in \mathbb{R}^3 .
Equate the linear combination of these three vectors with *unknown* coefficients x_1, x_2 and x_3 to 0 :

$$x_1(1, 2, 0) + x_2(0, 2, 4) + x_3(3, 0, 0) = (0, 0, 0).$$

Example : 3×3

Consider the three vectors $(1, 2, 0)$, $(0, 2, 4)$ and $(3, 0, 0)$ in \mathbb{R}^3 .
Equate the linear combination of these three vectors with *unknown* coefficients x_1, x_2 and x_3 to 0 :

$$x_1(1, 2, 0) + x_2(0, 2, 4) + x_3(3, 0, 0) = (0, 0, 0).$$

Hence we have the system of linear equations:

$$x_1 + 0x_2 + 3x_3 = 0$$

$$2x_1 + 2x_2 + 0x_3 = 0$$

$$0x_1 + 4x_2 + 0x_3 = 0$$

Example : 3×3

Consider the three vectors $(1, 2, 0)$, $(0, 2, 4)$ and $(3, 0, 0)$ in \mathbb{R}^3 .
Equate the linear combination of these three vectors with *unknown* coefficients x_1, x_2 and x_3 to 0 :

$$x_1(1, 2, 0) + x_2(0, 2, 4) + x_3(3, 0, 0) = (0, 0, 0).$$

Hence we have the system of linear equations:

$$x_1 + 0x_2 + 3x_3 = 0$$

$$2x_1 + 2x_2 + 0x_3 = 0$$

$$0x_1 + 4x_2 + 0x_3 = 0$$

Since the matrix $\begin{bmatrix} 1 & 0 & 3 \\ 2 & 2 & 0 \\ 0 & 4 & 0 \end{bmatrix}$ is invertible, the system of linear equations has a unique solution $x_1 = x_2 = x_3 = 0$.

Example : 3×3

Consider the three vectors $(1, 2, 0)$, $(0, 2, 4)$ and $(3, 0, 0)$ in \mathbb{R}^3 .
Equate the linear combination of these three vectors with *unknown* coefficients x_1, x_2 and x_3 to 0 :

$$x_1(1, 2, 0) + x_2(0, 2, 4) + x_3(3, 0, 0) = (0, 0, 0).$$

Hence we have the system of linear equations:

$$x_1 + 0x_2 + 3x_3 = 0$$

$$2x_1 + 2x_2 + 0x_3 = 0$$

$$0x_1 + 4x_2 + 0x_3 = 0$$

Since the matrix $\begin{bmatrix} 1 & 0 & 3 \\ 2 & 2 & 0 \\ 0 & 4 & 0 \end{bmatrix}$ is invertible, the system of linear equations has a unique solution $x_1 = x_2 = x_3 = 0$. **Hence the vectors $(1, 2, 0)$, $(0, 2, 4)$ and $(3, 0, 0)$ are linearly independent.**

More than 2 vectors in \mathbb{R}^2

Suppose we have n vectors in \mathbb{R}^2 where $n \geq 3$.

More than 2 vectors in \mathbb{R}^2

Suppose we have n vectors in \mathbb{R}^2 where $n \geq 3$. To check linear independence, we have to check whether the corresponding homogeneous linear system $Vx = 0$ has the unique solution $x = 0$.

More than 2 vectors in \mathbb{R}^2

Suppose we have n vectors in \mathbb{R}^2 where $n \geq 3$. To check linear independence, we have to check whether the corresponding homogeneous linear system $Vx = 0$ has the unique solution $x = 0$.

Since $n \geq 3 > 2$, this is a homogeneous system with more unknowns (n) than equations (2).

More than 2 vectors in \mathbb{R}^2

Suppose we have n vectors in \mathbb{R}^2 where $n \geq 3$. To check linear independence, we have to check whether the corresponding homogeneous linear system $Vx = 0$ has the unique solution $x = 0$.

Since $n \geq 3 > 2$, this is a homogeneous system with more unknowns (n) than equations (2).

We have seen in the previous week that Gaussian elimination will yield infinitely many solutions.

More than 2 vectors in \mathbb{R}^2

Suppose we have n vectors in \mathbb{R}^2 where $n \geq 3$. To check linear independence, we have to check whether the corresponding homogeneous linear system $Vx = 0$ has the unique solution $x = 0$.

Since $n \geq 3 > 2$, this is a homogeneous system with more unknowns (n) than equations (2).

We have seen in the previous week that Gaussian elimination will yield infinitely many solutions.

Hence, any set of n vectors in \mathbb{R}^2 with $n \geq 3$ are linearly dependent.

More than n vectors in \mathbb{R}^n

The same argument as for \mathbb{R}^2 in the previous slide yields :

More than n vectors in \mathbb{R}^n

The same argument as for \mathbb{R}^2 in the previous slide yields :

Hence, any set of r vectors in \mathbb{R}^n with $r > n$ are linearly dependent.

Example in \mathbb{R}^3

Consider the four vectors $(1, 2, 0)$, $(0, 2, 4)$, $(3, 0, 0)$ and $(1, 2, 3)$ in \mathbb{R}^3 .

Example in \mathbb{R}^3

Consider the four vectors $(1, 2, 0)$, $(0, 2, 4)$, $(3, 0, 0)$ and $(1, 2, 3)$ in \mathbb{R}^3 . To check linear independence, we write the corresponding system of linear equations :

$$x_1 + 0x_2 + 3x_3 + x_4 = 0$$

$$2x_1 + 2x_2 + 0x_3 + 2x_4 = 0$$

$$0x_1 + 4x_2 + 0x_3 + 3x_4 = 0$$

Example in \mathbb{R}^3

Consider the four vectors $(1, 2, 0)$, $(0, 2, 4)$, $(3, 0, 0)$ and $(1, 2, 3)$ in \mathbb{R}^3 . To check linear independence, we write the corresponding system of linear equations :

$$x_1 + 0x_2 + 3x_3 + x_4 = 0$$

$$2x_1 + 2x_2 + 0x_3 + 2x_4 = 0$$

$$0x_1 + 4x_2 + 0x_3 + 3x_4 = 0$$

To solve this system, we consider the augmented matrix

$$\left[\begin{array}{cccc|c} 1 & 0 & 3 & 1 & 0 \\ 2 & 2 & 0 & 2 & 0 \\ 0 & 4 & 0 & 3 & 0 \end{array} \right] \text{ and apply Gaussian elimination.}$$

Row reduction results in the augmented matrix $\left[\begin{array}{cccc|c} 1 & 0 & 0 & 1/4 & 0 \\ 0 & 1 & 0 & 3/4 & 0 \\ 0 & 0 & 1 & 1/4 & 0 \end{array} \right]$

Row reduction results in the augmented matrix $\left[\begin{array}{cccc|c} 1 & 0 & 0 & 1/4 & 0 \\ 0 & 1 & 0 & 3/4 & 0 \\ 0 & 0 & 1 & 1/4 & 0 \end{array} \right]$

Thus we obtain solutions : $x_1 = -\frac{c}{4}, x_2 = -\frac{3c}{4}, x_3 = -\frac{c}{4}, x_4 = c$
where $c \in \mathbb{R}$.

Row reduction results in the augmented matrix $\left[\begin{array}{cccc|c} 1 & 0 & 0 & 1/4 & 0 \\ 0 & 1 & 0 & 3/4 & 0 \\ 0 & 0 & 1 & 1/4 & 0 \end{array} \right]$

Thus we obtain solutions : $x_1 = -\frac{c}{4}$, $x_2 = -\frac{3c}{4}$, $x_3 = -\frac{c}{4}$, $x_4 = c$ where $c \in \mathbb{R}$.

So we can write

$$-\frac{c}{4}(1, 2, 0) - \frac{3c}{4}(0, 2, 4) - \frac{c}{4}(3, 0, 0) + c(1, 2, 3) = 0 \text{ for } c \in \mathbb{R}.$$

Row reduction results in the augmented matrix $\left[\begin{array}{cccc|c} 1 & 0 & 0 & 1/4 & 0 \\ 0 & 1 & 0 & 3/4 & 0 \\ 0 & 0 & 1 & 1/4 & 0 \end{array} \right]$

Thus we obtain solutions : $x_1 = -\frac{c}{4}, x_2 = -\frac{3c}{4}, x_3 = -\frac{c}{4}, x_4 = c$ where $c \in \mathbb{R}$.

So we can write

$$-\frac{c}{4}(1, 2, 0) - \frac{3c}{4}(0, 2, 4) - \frac{c}{4}(3, 0, 0) + c(1, 2, 3) = 0 \text{ for } c \in \mathbb{R}.$$

In particular with $c = 4$

$$-1(1, 2, 0) - 3(0, 2, 4) - 1(3, 0, 0) + 4(1, 2, 3) = 0$$

Row reduction results in the augmented matrix $\left[\begin{array}{cccc|c} 1 & 0 & 0 & 1/4 & 0 \\ 0 & 1 & 0 & 3/4 & 0 \\ 0 & 0 & 1 & 1/4 & 0 \end{array} \right]$

Thus we obtain solutions : $x_1 = -\frac{c}{4}, x_2 = -\frac{3c}{4}, x_3 = -\frac{c}{4}, x_4 = c$ where $c \in \mathbb{R}$.

So we can write

$$-\frac{c}{4}(1, 2, 0) - \frac{3c}{4}(0, 2, 4) - \frac{c}{4}(3, 0, 0) + c(1, 2, 3) = 0 \text{ for } c \in \mathbb{R}.$$

In particular with $c = 4$

$$-1(1, 2, 0) - 3(0, 2, 4) - 1(3, 0, 0) + 4(1, 2, 3) = 0$$

Hence the vectors $(1, 2, 0)$, $(0, 2, 4)$, $(3, 0, 0)$ and $(1, 2, 3)$ are linearly dependent.

Relationship with determinant

To check whether a set of n vectors in \mathbb{R}^n are linearly independent, we have to find the solutions of the homogeneous system $Vx = 0$ where V is an $n \times n$ matrix obtained by arranging the vectors in columns.

Relationship with determinant

To check whether a set of n vectors in \mathbb{R}^n are linearly independent, we have to find the solutions of the homogeneous system $Vx = 0$ where V is an $n \times n$ matrix obtained by arranging the vectors in columns.

Since V is a square matrix, it has unique solution $x = 0$ if and only if \checkmark ~~A~~ is invertible if and only if \checkmark $\det(\checkmark \text{ ~~A~~ }) \neq 0$.

Relationship with determinant

To check whether a set of n vectors in \mathbb{R}^n are linearly independent, we have to find the solutions of the homogeneous system $Vx = 0$ where V is an $n \times n$ matrix obtained by arranging the vectors in columns.

Since V is a square matrix, it has unique solution $x = 0$ if and only if $\checkmark A$ is invertible if and only if $\checkmark \det(A) \neq 0$.

- ▶ If A is invertible then there exists A^{-1} such that $AA^{-1} = \underline{1} = A^{-1}A$. Hence $\det(A) \cdot \det(A^{-1}) = 1$ which implies $\det(A) \neq 0$.

Relationship with determinant

To check whether a set of n vectors in \mathbb{R}^n are linearly independent, we have to find the solutions of the homogeneous system $Vx = 0$ where V is an $n \times n$ matrix obtained by arranging the vectors in columns.

Since V is a square matrix, it has unique solution $x = 0$ if and only if A is invertible if and only if $\det(A) \neq 0$.



- ▶ If A is invertible then there exists A^{-1} such that $AA^{-1} = \underline{1} = A^{-1}A$. Hence $\det(A) \cdot \det(A^{-1}) = 1$ which implies $\det(A) \neq 0$.
- ▶ Now if $\det(A) \neq 0$ then $A^{-1} = \frac{1}{\det(A)} \text{adj}(A)$ exists.

Example

Let us consider the vectors $(1, 4, 2)$, $(0, 4, 3)$ and $(1, 1, 0)$ in \mathbb{R}^3 .

Example

Let us consider the vectors $(1, 4, 2)$, $(0, 4, 3)$ and $(1, 1, 0)$ in \mathbb{R}^3 .

We obtain the matrix

$$V = \begin{bmatrix} 1 & 0 & 1 \\ 4 & 4 & 1 \\ 2 & 3 & 0 \end{bmatrix}.$$

Example

Let us consider the vectors $(1, 4, 2)$, $(0, 4, 3)$ and $(1, 1, 0)$ in \mathbb{R}^3 .

We obtain the matrix

$$V = \begin{bmatrix} 1 & 0 & 1 \\ 4 & 4 & 1 \\ 2 & 3 & 0 \end{bmatrix}.$$

Since $\det(V) = 1 \neq 0$, the matrix V is invertible and hence **the vectors $(1, 4, 2)$, $(0, 4, 3)$ and $(1, 1, 0)$ are linearly independent.**

Thank you

/