

Rahul Bhadani

Signals and Systems for Computer Engineers

WORKSHEET

*Department of Electrical & Computer Engineering,
The University of Alabama in Huntsville*

Content

1	Mathematical Preliminaries	3
1.1	Complex Numbers	3
1.2	Trigonometry	7


1. Mathematical Preliminaries

1.1 Complex Numbers

1. **4pts** Plot the following complex numbers on the complex plane:

- $2 + i3$
- $3 - i2$
- $-2 - i2$
- $-4 + j2$

Solution:



2. **4pts** Express $\frac{-1+3i}{2+5i}$ in the form $a + ib$.

Solution:

$$\begin{aligned}
\frac{-1+3i}{2+5i} &= \frac{-1+3i}{2+5i} \times \frac{2-5i}{2-5i} \\
&= \frac{-2+5i+6i-15i^2}{2^2+5^2} \\
&= \frac{-2+11i+15}{4+25} \quad (1.1) \\
&= \frac{13+11i}{29} \\
&= \frac{13}{29} + \frac{11}{29}i
\end{aligned}$$

3. **2pts** If $Z = 3 + i5$ is a complex number, what is the value of the modulus $|Z|$?

Solution:

$$|Z| = \sqrt{3^2 + 5^2} = \sqrt{9 + 25} = \sqrt{34}$$

4. **4pts** Find the roots of the equation $x^2 + x + 1 = 0$.

Solution:

For a quadratic equation $ax^2 + bx + c = 0$, the roots are given by $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$. In this case, $a = 1$, $b = 1$, $c = 1$.

$$x = \frac{-1 \pm \sqrt{1^2 - 4(1)(1)}}{2(1)} = \frac{-1 \pm \sqrt{1-4}}{2} = \frac{-1 \pm \sqrt{-3}}{2} = \frac{-1 \pm i\sqrt{3}}{2}$$

5. **2pts** Write the following complex numbers in the polar form:

(a) $z = 1 + i$

(b) $w = \sqrt{3} - i$

Solution for (a): $|z| = \sqrt{1^2 + 1^2} = \sqrt{2}$ $\tan \theta = \frac{1}{1} = 1$, so $\theta = \frac{\pi}{4}$. In polar form: $z = \sqrt{2}(\cos \frac{\pi}{4} + i \sin \frac{\pi}{4})$.

Solution for (b): $|w| = \sqrt{(\sqrt{3})^2 + (-1)^2} = \sqrt{3+1} = \sqrt{4} = 2$
 $\tan \theta = \frac{-1}{\sqrt{3}}$, so $\theta = -\frac{\pi}{6}$ (since the number is in the 4th quadrant).
 In polar form: $w = 2(\cos(-\frac{\pi}{6}) + i \sin(-\frac{\pi}{6}))$.

6. **4pts** Find the product of the complex numbers $1 + i$ and $\sqrt{3} - i$ in the polar form.

Solution:

From the previous problem, we have: $z_1 = 1 + i = \sqrt{2}e^{i\pi/4}$

$$z_2 = \sqrt{3} - i = 2e^{-i\pi/6}$$

$$z_1 z_2 = (\sqrt{2}e^{i\pi/4})(2e^{-i\pi/6}) = 2\sqrt{2}e^{i(\pi/4 - \pi/6)} = 2\sqrt{2}e^{i(\pi/12)}$$

In standard polar form: $2\sqrt{2}(\cos \frac{\pi}{12} + i \sin \frac{\pi}{12})$.

7. **2pts** Find $(\frac{1}{2} + \frac{1}{2}i)^{10}$.

Solution:

First, convert to polar form:

$$\frac{1}{2} + \frac{1}{2}i = \frac{\sqrt{2}}{2}(\cos \frac{\pi}{4} + i \sin \frac{\pi}{4})$$

Using De Moivre's Theorem:

$$\begin{aligned} (\frac{1}{2} + \frac{1}{2}i)^{10} &= (\frac{\sqrt{2}}{2})^{10}(\cos(10 \cdot \frac{\pi}{4}) + i \sin(10 \cdot \frac{\pi}{4})) \\ &= (\frac{2^{1/2}}{2})^{10}(\cos(\frac{5\pi}{2}) + i \sin(\frac{5\pi}{2})) \\ &= (\frac{1}{2^{1/2}})^{10}(\cos(\frac{\pi}{2} + 2\pi) + i \sin(\frac{\pi}{2} + 2\pi)) \\ &= (\frac{1}{2^5})(0 + i \cdot 1) = \frac{i}{32} \end{aligned}$$

8. **2pts** Evaluate or Simplify:

(a) $e^{i\pi}$

(b) $e^{-1+i\pi/2}$

Solution for (a): $e^{i\pi} = \cos \pi + i \sin \pi = -1 + i(0) = -1$.

Solution for (b): $e^{-1+i\pi/2} = e^{-1}e^{i\pi/2} = e^{-1}\left(\cos \frac{\pi}{2} + i \sin \frac{\pi}{2}\right) = \frac{1}{e}\left(0 + i(1)\right) = \frac{i}{e}$

9. **2pts** Evaluate the expression below and write your answer in the form $a + ib$.

(a) $(5 - i6) + (3 + i2)$

(b) $\frac{3}{4-i3}$

Solution for (a): $(5 - i6) + (3 + i2) = (5 + 3) + (-6 + 2)i = 8 - 4i$.

Solution for (b): $\frac{3}{4-i3} = \frac{3}{4-i3} \times \frac{4+i3}{4+i3} = \frac{12+i9}{4^2+3^2} = \frac{12+i9}{16+9} = \frac{12+i9}{25} = \frac{12}{25} + \frac{9}{25}i$.

10. **2pts** Find the complex conjugate and modulus of the number:

(a) $12 + i5$

(b) $-1 + 2\sqrt{2}i$

Solution for (a):

Complex conjugate: $12 - i5$.

Modulus: $|12 + i5| = \sqrt{12^2 + 5^2} = \sqrt{144 + 25} = \sqrt{169} = 13$.

Solution for (b):

Complex conjugate: $-1 - 2\sqrt{2}i$.

Modulus: $|-1 + 2\sqrt{2}i| = \sqrt{(-1)^2 + (2\sqrt{2})^2} = \sqrt{1 + 8} = \sqrt{9} = 3$.

11. **2pts** Apply De Moivre's Theorem to simplify:

(a) $(1 + i)^{20}$

(b) $(1 - \sqrt{3}i)^5$

(c) $(1 - i)^8$

Solution for (a):

Convert $1 + i$ to polar form: $r = \sqrt{1^2 + 1^2} = \sqrt{2}$,

$$\theta = \tan^{-1}\left(\frac{1}{1}\right) = \frac{\pi}{4}.$$

$$1 + i = \sqrt{2}\left(\cos \frac{\pi}{4} + i \sin \frac{\pi}{4}\right).$$

$$\begin{aligned}(1 + i)^{20} &= (\sqrt{2})^{20}(\cos(20 \cdot \frac{\pi}{4}) + i \sin(20 \cdot \frac{\pi}{4})) \\ &= 2^{10}(\cos(5\pi) + i \sin(5\pi)) = 1024(-1 + 0i) = -1024.\end{aligned}$$

Solution for (b):

Convert $1 - \sqrt{3}i$ to polar form: $r = \sqrt{1^2 + (-\sqrt{3})^2} = 2$, $\theta = \tan^{-1}\left(\frac{-\sqrt{3}}{1}\right) = -\frac{\pi}{3}$.

$$1 - \sqrt{3}i = 2\left(\cos\left(-\frac{\pi}{3}\right) + i \sin\left(-\frac{\pi}{3}\right)\right).$$

$$\begin{aligned}(1 - \sqrt{3}i)^5 &= 2^5(\cos(5 \cdot (-\frac{\pi}{3})) + i \sin(5 \cdot (-\frac{\pi}{3}))) \\ &= 32(\cos(-\frac{5\pi}{3}) + i \sin(-\frac{5\pi}{3})) = 32(\cos(\frac{\pi}{3}) + i \sin(\frac{\pi}{3})) \\ &= 32(\frac{1}{2} + i\frac{\sqrt{3}}{2}) = 16(1 + i\sqrt{3}).\end{aligned}$$

Solution for (c):

Convert $1 - i$ to polar form: $r = \sqrt{1^2 + (-1)^2} = \sqrt{2}$, $\theta = \tan^{-1}\left(\frac{-1}{1}\right) = -\frac{\pi}{4}$.

$$1 - i = \sqrt{2}\left(\cos\left(-\frac{\pi}{4}\right) + i \sin\left(-\frac{\pi}{4}\right)\right).$$

$$\begin{aligned}(1 - i)^8 &= (\sqrt{2})^8(\cos(8 \cdot (-\frac{\pi}{4})) + i \sin(8 \cdot (-\frac{\pi}{4}))) \\ &= 2^4(\cos(-2\pi) + i \sin(-2\pi)) = 16(1 + 0i) = 16.\end{aligned}$$

12. **3pts** Use Euler's formula to prove the following formulas for $\cos x$ and $\sin x$:

(a) $\cos x = \frac{e^{ix} + e^{-ix}}{2}$

(b) $\sin x = \frac{e^{ix} - e^{-ix}}{2i}$

Solution: Euler's formula states: $e^{ix} = \cos x + i \sin x$ $e^{-ix} = \cos x - i \sin x$


To prove (a), add the two equations: $e^{ix} + e^{-ix} = (\cos x + i \sin x) + (\cos x - i \sin x) = 2 \cos x$. Therefore, $\cos x = \frac{e^{ix} + e^{-ix}}{2}$.

To prove (b), subtract the second equation from the first: $e^{ix} - e^{-ix} = (\cos x + i \sin x) - (\cos x - i \sin x) = 2i \sin x$. Therefore, $\sin x = \frac{e^{ix} - e^{-ix}}{2i}$.

1.2 Trigonometry

1. **4 pts** Draw a circle and mark $\frac{\pi}{6}$, $\frac{\pi}{3}$, $\frac{\pi}{4}$, and $\frac{3\pi}{4}$.

Solution:



2. **1 pts** Convert $\frac{3\pi}{2}$ radians into degrees.

Solution:

$$\frac{3\pi}{2} \times \frac{180^\circ}{\pi} = 270^\circ \quad (1.2)$$

3. **3 pts** Find the value of θ : $4 \sin^2 \theta = 3$.

Solution:

$$\begin{aligned} 4 \sin^2 \theta &= 3 \\ \sin^2 \theta &= \frac{3}{4} \\ \sin \theta &= \pm \frac{\sqrt{3}}{2} \end{aligned} \quad (1.3)$$

If $\sin \theta = \frac{\sqrt{3}}{2}$, then $\theta = 60^\circ$ or $\theta = 120^\circ$ (Quadrant I and II).

If $\sin \theta = -\frac{\sqrt{3}}{2}$, then $\theta = 240^\circ$ or $\theta = 300^\circ$ (Quadrant III and IV).

4. **3 pts** Find the value of x : $2 \sin^2 x - 3 \sin x + 1 = 0$.

Solution:

$$\begin{aligned}
2 \sin^2 x - 3 \sin x + 1 &= 0 \\
2 \sin^2 x - 2 \sin x - \sin x + 1 &= 0 \\
2 \sin x (\sin x - 1) - 1 (\sin x - 1) &= 0 \\
(2 \sin x - 1)(\sin x - 1) &= 0
\end{aligned} \tag{1.4}$$

This gives two possible solutions:

$$2 \sin x - 1 = 0 \implies \sin x = \frac{1}{2}.$$

$$\sin x - 1 = 0 \implies \sin x = 1.$$

$$\text{For } \sin x = \frac{1}{2}, x = \frac{\pi}{6} \text{ or } x = \frac{5\pi}{6}.$$

$$\text{For } \sin x = 1, x = \frac{\pi}{2}.$$

The general solutions are $x = \frac{\pi}{6} + 2\pi K$, $x = \frac{5\pi}{6} + 2\pi K$, and $x = \frac{\pi}{2} + 2\pi K$.

5. **5 pts** Prove: $(1 - \sin^2(t))(1 + \tan^2(t)) = 1$.

Solution:

$$\begin{aligned}
(1 - \sin^2(t))(1 + \tan^2(t)) &= (\cos^2(t))(1 + \tan^2(t)) \\
&= \cos^2(t) + \cos^2(t) \tan^2(t) \\
&= \cos^2(t) + \cos^2(t) \frac{\sin^2(t)}{\cos^2(t)} \\
&= \cos^2(t) + \sin^2(t) \\
&= 1
\end{aligned} \tag{1.5}$$

6. **5 pts** Prove: $\frac{\sin^3(t) + \cos^3(t)}{\sin(t) + \cos(t)} = 1 - \sin(t) \cos(t)$.

Solution:Using the identity $a^3 + b^3 = (a + b)(a^2 - ab + b^2)$.Let $a = \sin(t)$ and $b = \cos(t)$.

$$\begin{aligned}
\frac{\sin^3(t) + \cos^3(t)}{\sin(t) + \cos(t)} &= \frac{(\sin(t) + \cos(t))(\sin^2(t) - \sin(t) \cos(t) + \cos^2(t))}{\sin(t) + \cos(t)} \\
&= \sin^2(t) + \cos^2(t) - \sin(t) \cos(t) \\
&= 1 - \sin(t) \cos(t)
\end{aligned} \tag{1.6}$$

7. **2 pts** What is the value of $\sin \theta$ and $\cos \theta$ given $\tan \theta = \frac{4}{3}$?

Solution:


$$\tan \theta = \frac{4}{3} = \frac{\text{Perpendicular}}{\text{Base}}.$$

The hypotenuse is $h = \sqrt{3^2 + 4^2} = \sqrt{9 + 16} = \sqrt{25} = 5$.

$$\sin \theta = \frac{\text{Perpendicular}}{\text{Hypotenuse}} = \frac{4}{5}.$$

$$\cos \theta = \frac{\text{Base}}{\text{Hypotenuse}} = \frac{3}{5}.$$

8. **2 pts** Find the value of $\sin \theta$ and $\cos \theta$ from the triangle.



Solution:

From the right-angled triangle with sides 1, 1 and hypotenuse $\sqrt{2}$:

$$\sin \theta = \frac{\text{Opposite}}{\text{Hypotenuse}} = \frac{1}{\sqrt{2}}.$$

$$\cos \theta = \frac{\text{Adjacent}}{\text{Hypotenuse}} = \frac{1}{\sqrt{2}}.$$

9. **3 pts** Prove $\sin^{-1}(\frac{\sqrt{2}}{2}) - \sin^{-1}(\frac{1}{2}) = \frac{\pi}{12}$.

Solution:

$$\sin^{-1}(\frac{\sqrt{2}}{2}) = \frac{\pi}{4}.$$

$$\sin^{-1}(\frac{1}{2}) = \frac{\pi}{6}.$$

$$\sin^{-1}(\frac{\sqrt{2}}{2}) - \sin^{-1}(\frac{1}{2}) = \frac{\pi}{4} - \frac{\pi}{6} = \frac{3\pi - 2\pi}{12} = \frac{\pi}{12}.$$

10. **2 pts** Find the value of x given $2 \sin x = 1$.

Solution:

$$2 \sin x = 1 \implies \sin x = \frac{1}{2}.$$

$$\text{For } \sin x = \frac{1}{2}, x = \frac{\pi}{6} \text{ or } x = \frac{5\pi}{6}.$$