

CPE 381: Fundamentals of Signals and Systems for Computer Engineers

01 Introduction and Preliminaries

Fall 2025

Rahul Bhadani

Outline

1. Course Logistics

2. Motivation

3. Mathematical Preliminaries

About Me

Rahul Bhadani

Assistant Professor

Electrical and Computer Engineering

The University of Alabama in Huntsville

Office: 217-H

Email: rahul.bhadani@uah.edu

Web: <https://rahulbhadani.github.io/>



Research Interests

Cyber-physical Systems, Intelligent Transportation, Connected-and-Autonomous Driving, Applied machine learning, Quantum Information Science

Course Logistics

Lecture:

M/W 11:20 AM - 12:40 PM

Location:

ENG 240

Prerequisites:

- ⚡ EE 213 - Electrical Circuit Analysis I
- ⚡ MA 238 – Applied Differential Equations

Office Hours:

⚡ Tuesday: 2:00 PM - 4:00 PM

⚡ Wednesday: 12:45 PM - 2:00 PM

Instructor Email: rahul.bhadani@uah.edu

Textbooks

Required: *Signals and Systems using Matlab*. Luis F. Chaparro.

Elsevier, 3rd Edition, 2019.

ISBN: 978-0-12-814204-2, eBook ISBN: 9780128142059

Suggested Reading: *Linear Systems and Signals*, B. P. Lathi, Roger Green, Oxford University Press, 2017, 3rd Edition

Other Notable Textbooks:

⚡ Schaum's Outline of Signals and Systems

⚡ *Signals and Systems*. Haykin, Simon, and Barry Van Veen. John Wiley & Sons, 2007.

⚡ Asadi, Farzin. Signals and Systems with MATLAB and Simulink. Springer, Dec 2023.

ISBN: 9783031456220, 303145622X

Grading

Homework:	25%
Quizzes:	5%
Attendance/In-Class Participation:	10%
Mid-term Exam 1:	15%
Mid-term Exam 2:	15%
Final Exam:	30%

Grading Scale	
Percentage	Grade
90% - 100%	A
75% - 89%	B
60% - 74%	C
45% - 59%	D
0% - 44%	F

Exam Schedule

- ⚡ **Mid Term 1:** September 30, Monday
- ⚡ **Mid Term 2:** November 06, Wednesday
- ⚡ **Final Exam:** December 13, Friday

Tentative Topics

- ⚡ Introduction, Mathematical Preliminaries
- ⚡ Continuous and Discrete Signals
- ⚡ Linear-time Invariant (LTI) Systems
- ⚡ Laplace Transform
- ⚡ Fourier Series for Frequency Analysis
- ⚡ Fourier Transform
- ⚡ Sampling Theory
- ⚡ Discrete-Time Signals and Systems
- ⚡ Z-transform
- ⚡ Discrete Fourier Analysis

Refer to the syllabus for the detailed information on the syllabus.

In-Class Activity


Introduce Yourself

- ⚡ Why Computer Engineering?
- ⚡ Why do you want to take this course?
- ⚡ One fun fact about you that's not on social media





Motivation


Why Study Signals and Systems



We are in digital era




Sensors and Mobile Devices



Cyber-physical Systems




Connected and Autonomous Vehicles




Microgrid

Signals in Nature

Everything in nature is analog and continuous. We use hardware/software interfaces to digitize them, extract information, make inferences, and send them back to the user.



Sampling continuous time signals



$x[n] = x(nT_s)$, T_s = sample time.
Code for the figure: https://github.com/rahulbhadani/CPE381_FA25/blob/master/Code/sampled_sine_wave.m

Inherent Discrete Time Signals

Nature may be continuous but human activities naturally lead to some discrete time signals.

Examples:

- ⚡ Stock market closing data
- ⚡ Adaptive cruise control states
- ⚡ Thermostat setpoints



Mathematical Preliminaries

Trigonometry

⚡ Sine, Cosine, Tangent:


$$-\sin(\theta) = \frac{p}{h}$$

$$-\cos(\theta) = \frac{b}{h}$$

$$-\tan(\theta) = \frac{p}{b}$$

⚡ Pythagorean Identity:


$$\sin^2(\theta) + \cos^2(\theta) = 1$$



⚡ Unit Circle:

Trigonometric Identities

What is the value of h based on the diagram?



⚡ $\cos \theta =$

⚡ $\sin \theta =$

⚡ $\sin(\alpha + \beta) =$

⚡ $\cos(\alpha + \beta) =$

⚡ $\sin^2(\theta) =$ in terms of $\cos(2\theta)$

⚡ $\cos^2(\theta) =$ in terms of $\cos(2\theta)$

⚡ $\sin 2\theta =$ in terms of θ


⚡ $\cos 2\theta =$ in terms of θ

⚡ $\sin(-\theta) =$

⚡ $\cos(-\theta) =$

Trigonometric Identities

What is the value of h based on the diagram?




- ⚡ $\tan(\alpha + \beta) =$
- ⚡ $\tan(\alpha - \beta) =$
- ⚡ $\sin 30^\circ =$
- ⚡ $\sin 45^\circ =$
- ⚡ $\sin 60^\circ =$

- ⚡ $\cos 30^\circ =$
- ⚡ $\cos 45^\circ =$
- ⚡ $\cos 60^\circ =$
- ⚡ $\tan 30^\circ =$
- ⚡ $\tan 45^\circ =$
- ⚡ $\tan 60^\circ =$

What's the range of $\sin \theta$ and $\cos \theta$?

Trigonometric Functions in Different Quadrant

Four Quadrants:



Considering a unit circle, in four different quadrants (x, y) signs are different, Hence the trigonometric ratios are also different in signs.


Quadrant	sin	cos	tan
I	+	+	+
II	+	-	-
III	-	-	+
IV	-	+	-

Table: Signs of sin, cos, and tan in all four quadrants

Complex Number

$$z = x + iy$$

$$i^2 = -1$$



Two representations of complex numbers:

$$z = re^{i\theta}$$

$$\text{Conjugate: } \bar{z} = re^{-i\theta}$$

$$\blacksquare r = \sqrt{x^2 + y^2}$$

$$\blacksquare \theta = \tan^{-1} \frac{y}{x}$$

Euler's Identity:

$$\blacksquare e^{i\theta} = \cos \theta + i \sin \theta$$

$$\blacksquare e^{-i\theta} =$$

$$\blacksquare \cos \theta =$$

(in terms of exponential)

$$\blacksquare \sin \theta =$$

(in terms of exponential)

Complex Number

Properties of Conjugates

- ⚡ $\overline{z+w} = \bar{z} + \bar{w}$
- ⚡ $\overline{zw} = \bar{z}\bar{w}$
- ⚡ $\overline{z^n} = \bar{z}^n$
- ⚡ $z\bar{z} = |z|^2$ **Prove it!**

De Moivre's Theorem If $z = r(\cos \theta + i \sin \theta)$ and n is a positive integer, then

$$z^n = [r(\cos \theta + i \sin \theta)^n] = r^n(\cos n\theta + i \sin n\theta)$$

Roots of a complex number $z = 0$ has n distinct roots:

$$w_k = r^{1/n} \left[\cos \left(\frac{\theta + 2k\pi}{n} \right) + i \sin \left(\frac{\theta + 2k\pi}{n} \right) \right]$$

Complex Exponentials

We also need to give a meaning to the expression e^z when $z = x + iy$ is a complex number.

Based on the infinite series using Taylor's series expansion for e^x , we have

$$e^z = \sum_{n=0}^{\infty} \frac{z^n}{n!} = 1 + z + \frac{z^2}{2!} + \frac{z^3}{3!} + \dots$$

Can you find out, what should be the value of e^{iy} where y is the real number?

Note: $\cos y = 1 - \frac{y^2}{2!} + \frac{y^4}{4!} - \frac{y^6}{6!} + \dots$ and $\sin y = y - \frac{y^3}{3!} + \frac{y^5}{5!} - \dots$

Derivatives

Rate of change of a quantity with respect to another quantity.

In signals and systems, we are usually interested in the rate of change with respect to time.

$$f'(t) = \frac{df(t)}{dt} = \lim_{h \rightarrow 0} \frac{f(t + h) - f(t)}{h}$$

In computer implementation, h is Δt . Δt is the difference of timestamps between two consecutive samples of a signal.

Derivative Review



$$\begin{array}{l} \textcolor{red}{\checkmark} f'(x) = \frac{df(x)}{dx} \\ \textcolor{red}{\checkmark} f'(t) = \end{array}$$

$$\begin{array}{l} \textcolor{red}{\checkmark} \frac{d}{dx}(x^n) = nx^{n-1} \\ \textcolor{red}{\checkmark} \frac{d}{dx}(\ln(x)) = \frac{1}{x} \\ \textcolor{red}{\checkmark} \frac{d}{dx}(e^x) = e^x \\ \textcolor{red}{\checkmark} \frac{d}{dx}(\sin(x)) = \cos(x) \\ \textcolor{red}{\checkmark} \frac{d}{dx}(\cos(x)) = -\sin(x) \\ \textcolor{red}{\checkmark} \frac{d}{dx}(\sin^{-1}(x)) = \frac{1}{\sqrt{1-x^2}} \\ \textcolor{red}{\checkmark} \frac{d}{dx}(\tan(x)) = \sec^2(x) \\ \textcolor{red}{\checkmark} \frac{d}{dx}(\cot(x)) = -\csc^2(x) \end{array}$$

$$\begin{array}{l} \textcolor{red}{\checkmark} \frac{d}{dx}(\sec^{-1}(x)) = \frac{1}{x\sqrt{x^2-1}} \\ \textcolor{red}{\checkmark} \frac{d}{dx}(\sec(x)) = \sec(x)\tan(x) \\ \textcolor{red}{\checkmark} \frac{d}{dx}(\csc(x)) = -\csc(x)\cot(x) \\ \textcolor{red}{\checkmark} \frac{d}{dx}(\tan^{-1}(x)) = \frac{1}{x^2+1} \\ \textcolor{red}{\checkmark} (F \cdot S)' = F \cdot S' + F' \cdot S \\ \textcolor{red}{\checkmark} \left(\frac{N}{D}\right)' = \frac{D \cdot N' - N \cdot D'}{D^2} \\ \textcolor{red}{\checkmark} [f(g(x))]' = f'(g(x)) \cdot g'(x) \end{array}$$

Geometrical Interpretation of Derivatives

The derivative $f'(x)$ can be interpreted as the slope of the tangent line at the point (x, y) on the graph of the function $y = f(x)$.



Geometrical Interpretation of Derivatives

We can use the geometric interpretation to help study the behavior and the graph of $f(x)$.

- ⚡ $f'(x) > 0$ exactly when $f(x)$ is increasing.
 - ⚡ $f'(x) < 0$ exactly when $f(x)$ is decreasing.
 - ⚡ $f'(x) = 0$ exactly when $f(x)$ has a horizontal tangent.
-

In many applications, we want to find local maximum and local minimum values. In these applications, it makes sense to look at the locations on a graph where the slope is 0. In other words, we often use the following method to find all locations where the slope of the graph is 0.

- ⚡ Find the formula for the derivative: $f'(x)$.
- ⚡ Set this formula equal to 0 and solve for x .


Integration

**Integration is antiderivative but you can also understand it as an area under the curve.
In other words, integration is the summation.**

If $\frac{d}{dx}F(x) = f(x)$ then $\int f(x)dx = F(x).$

Geometric Interpretation of Integral

Integral gives the area under a curve



$$\int_0^1 x^2 dx$$

Table of Integrals: Basic Form

$$\int x^n dx = \frac{1}{n+1}x^{n+1}$$

$$\int \frac{1}{x} dx = \ln x$$

$$\int u dv = uv - \int v du$$

$$\int u(x)v'(x) dx = u(x)v(x) - \int v(x)u'(x) dx$$

Table of Integrals: More Functions

$$\int \frac{1}{ax+b} dx = \frac{1}{a} \ln(ax + b)$$

$$\int \frac{1}{(x+a)^2} dx = -\frac{1}{x+a}$$

$$\int (x+a)^n dx = \frac{(x+a)^{n+1}}{n+1}, \quad n \neq -1$$

$$\int x(x+a)^n dx = \frac{(x+a)^{n+1}(nx+x-a)}{(n+2)(n+1)}$$

$$\int \frac{dx}{1+x^2} = \tan^{-1} x$$

$$\int \frac{dx}{a^2+x^2} = \frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right)$$

$$\int \frac{x dx}{a^2+x^2} = \frac{1}{2} \ln(a^2 + x^2)$$

$$\int \frac{x^2 dx}{a^2+x^2} = x - a \tan^{-1} \left(\frac{x}{a} \right)$$

$$\int \frac{x^3 dx}{a^2+x^2} = \frac{1}{2}x^2 - \frac{1}{2}a^2 \ln(a^2 + x^2)$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}$$

$$\int \sin x dx = -\cos x$$

$$\int \cos x dx = \sin x$$

$$\int \tan x dx = -\ln \cos x$$

Differential Equations

Differential equations are simply equations that have derivatives in them.

Example:

⚡ $\frac{dy}{dx} + 5y = 0$

⚡ $y'' + 2y' + 10y = 0$

⚡ $y'' + 3y' + 2y = 3 \cos t$

The order of a differential equation is the order of the highest derivative in the equation. They are all examples of 'ordinary differential equations'.

Solving Differential Equations

Consider

$$y' = 2y + 3$$

The Fundamental Theorem of Calculus implies $y(t) = \int y'(t)dt$, so we get

$$y(t) = 2 \int y(t)dt + 3t + C$$

This is not the closed form and doesn't entirely solve to find a solution y . We look at some formula (and their derivation) using the rules of differentiation on how to get a closed-form solution.

Solving Differential Equations

The linear differential equation

$$y' = ay + b$$

with $a \neq 0, b$ has infinitely many solutions

$$y(t) = ce^{at} - \frac{b}{a}$$

Proof discussed in class.

Up Next

- ⚡ Representation of Signals: Continuous and Discrete
- ⚡ Continuous Signals
- ⚡ Using MATLAB