

Homework 1: Pre-requisites and Numerical Methods for Signals

CPE 381

Instructor: Rahul Bhadani

Due: Sept 11, 2026, 11:59 PM
100 points

You are allowed to use a generative model-based AI tool for your assignment. However, you must submit an accompanying reflection report detailing how you used the AI tool, the specific query you made, and how it improved your understanding of the subject. You are also required to submit screenshots of your conversation with any large language model (LLM) or equivalent conversational AI, clearly showing the prompts and your login avatar. Some conversational AIs provide a way to share a conversation link, and such a link is desirable for authenticity. Failure to do so may result in actions taken in compliance with the plagiarism policy.

Additionally, you must include your thoughts on how you would approach the assignment if such a tool were not available. Failure to provide a reflection report for every assignment where an AI tool is used may result in a penalty, and subsequent actions will be taken in line with the plagiarism policy.

Submission instruction:

Upload a .pdf on Canvas with the format CPE381-LastFirst-HW-XX. For example, if your name is Sam Wells, your file name should be CPE381-WellsSam-HW-XX.pdf. If there is a programming assignment, then you should include your source code along with your PDF files in a zip file CPE381-LastFirst-HW-XX.zip. Your submission must contain your name and UAH Charger ID or your UAH email address. Please number your pages as well.

Theory

1 Let's face the truth. (10 points)

Prove:

1. $1 + e^{j\pi} = 0.$

2. $\sin(-\theta) = -\sin \theta$ using Euler's identity.

Answer

1. Using Euler's Identity, we can write $e^{j\pi}$ as

$$e^{j\pi} = \cos \pi + j \sin \pi$$

but $\cos \pi = -1$, and $\sin \pi = 0$, hence $e^{j\pi} = -1$. Thus, $1 + e^{j\pi} = 0$.

2. Using Euler's identity,

$$\sin \theta = \frac{e^{j\theta} - e^{-j\theta}}{2j}$$

Replace θ by $-\theta$,

$$\Rightarrow \sin(-\theta) = \frac{e^{-j\theta} - e^{j\theta}}{2j}$$

Take the minus sign out of the numerator from Right-hand side of the above equation

$$\Rightarrow \sin(-\theta) = -\frac{e^{j\theta} - e^{-j\theta}}{2j} = -\sin \theta$$

(1)

2 Tigonometry Fun (20 points)

1. Verify the following identities: $1 - \frac{\cos^2 \theta}{1 + \sin \theta} = \sin \theta$

2. Solve $\cos 2x - 3 \sin x + 1 = 0$ for x .

Answer

1. We know, $\sin^2 \theta + \cos^2 \theta = 1$

$$1 - \frac{1 - \sin^2 \theta}{1 + \sin \theta} = 1 - \frac{(1 - \sin \theta)(1 + \sin \theta)}{1 + \sin \theta} = 1 - (1 - \sin \theta) = \sin \theta \quad (2)$$

2.

$$\begin{aligned}
 \cos 2x - 3 \sin x + 1 &= 0 \\
 (1 - 2 \sin^2 x) - 3 \sin x + 1 &= 0 \\
 -2 \sin^2 x - 3 \sin x + 2 &= 0 \\
 2 \sin^2 x + 3 \sin x - 2 &= 0 \\
 (2 \sin x - 1)(\sin x + 2) &= 0 \\
 \Rightarrow \sin x &= \frac{1}{2} \text{ and } \sin x = -2
 \end{aligned} \tag{3}$$

For $\sin x = \frac{1}{2}$, $x = \frac{\pi}{6}$ and $\frac{5\pi}{6}$. For $\sin x = -2$, there is no solution as $\sin x \in [-1, 1]$.

3 It's Complicated (20 points)

- Using Euler's formula evaluate the following complex numbers: (a) $z = e^{j2\pi}$ (b) $z = e^{j\pi}$
- Using Euler's formula derive the following: $\cos^2 \theta = \frac{1}{2}(1 + \cos(2\theta))$

Answer

- $z = e^{j2\pi} = \cos(2\pi) + j \sin(2\pi) = 1 + 0 = 1$
 - $z = e^{j\pi} = \cos(\pi) + j \sin(\pi) = -1 + 0 = -1$
- $$\begin{aligned}
 \cos^2 \theta &= \left(\frac{e^{j\theta} + e^{-j\theta}}{2} \right)^2 = \frac{e^{2j\theta} + e^{-2j\theta} + 2e^{j\theta}e^{-j\theta}}{4} \\
 &= \frac{e^{2j\theta} + e^{-2j\theta}}{4} + \frac{2}{4} = \frac{\cos(2\theta)}{2} + \frac{1}{2} = \frac{1}{2}(1 + \cos(2\theta))
 \end{aligned}$$

4 It's all Derivatives (10 points)

- Taylor's Formula is written as

$$f(x) \approx g_n(x) = f(c) + f'(c)(x - c) + \frac{f''(c)}{2!}(x - c)^2 + \cdots + \frac{f^{(n)}(c)}{n!}(x - c)^n$$

If $f(x) = e^{\sin x}$, what is the Taylor's second degree ($n = 2$) polynomial at around point $c = 0$?

2. Differentiate: (a) $f(x) = x(\ln x - 1)$, (b) $f(x) = \ln(\tan x)$.

Answer

$$1. \quad g_2(x) = f(0) + f'(0)x + \frac{f''(0)}{2}x^2$$

$$f(x) = e^{\sin x} \Rightarrow f(0) = 1$$

$$f'(x) = e^{\sin x} \cos x \Rightarrow f'(0) = 1$$

$$f''(x) = e^{\sin x} \cos^2 x - e^{\sin x} \sin x \Rightarrow f''(0) = 1$$

$$\text{Hence, } g_2(x) = 1 + x + \frac{x^2}{2}$$

Coding

Note: You will need to provide all relevant MATLAB code in a zip file along with the PDF contain your solution/responses.

5 Geometric Progression (GP) and Convergence (10 Points)

Consider a GP with the first term as $a = 2.0$ and the common ratio as $r = \frac{1}{3}$. The formula for sum of a GP is

$$S_n = \frac{a(r^n - 1)}{(r - 1)}$$

Write a MATLAB code to plot the sum S_n as a function of n to demonstrate that the sum indeed converge as n approaches a very large value. You may choose n upto 10000. Label axes properly in your plot.

Answer

We may not need n upto 10000 as convergence happens much faster. We will just choose n upto 100.

```

Sn = @(a, r, n) (a.*(r.^n - 1))./(r-1);
a = 2.0;
r = 1.0/3.0;
% Varies n from 1 to 100 spaced by 1
N = 1:1:100;
SUM = zeros(1, length(N));
for i = 1:length(N)
    SUM(i) = Sn(a, r, N(i));
    SUM(i)
end
f = figure;
plot(N, SUM, 'LineWidth',2, 'Color','#432423');
grid on;
grid minor;
set(gca, 'XColor', [0, 0, 0], 'YColor', [0, 0, 0], 'TickDir', 'out');
xaxis = get(gca, 'XAxis');
xaxis.TickLabelInterpreter = 'latex';
yaxis = get(gca, 'YAxis');
yaxis.TickLabelInterpreter = 'latex';
set(gca, 'FontSize', 18);
xlabel('n', 'Interpreter', 'latex');
ylabel('$S_n$', 'Interpreter', 'latex');
ylim([1.9, 3.1]);
exportgraphics(f, 'figures/HW01_Geometric_Progression.pdf', 'BackgroundColor', 'none');

```




Figure 1: Convergence of a Geometric Progression.

6 Discrete Derivative and Sampling Time (20 Points)

Consider a quadratic signal $f(t) = 3t^2 + 4t + 5$. You are required to compute the discrete derivative of $f(t)$. Compute discrete derivative using **Backward difference formula** as shown below:

$$\Delta f(t_k) = \frac{f(t_k) - f(t_{k-1})}{t_k - t_{k-1}}$$

where k is the index for k -th time-points.

Consider three scenarios:

1. Compute discrete derivative with the above formula while consider uniformly sampling time of $\Delta t = 1.0$ seconds.
2. Compute discrete derivative with the above formula while consider uniformly sampling time of $\Delta t = 0.5$ seconds.
3. Compute discrete derivative with the above formula while consider uniformly sampling time of $\Delta t = 0.05$ seconds.

Assume the final time to be $t = 10$ seconds.

Compare them with the analytically solved derivative $f'(t)$. Plot all four versions of derivatives. Label your axes properly, and choose appropriate DisplayName and legend for visualization.

Answer

We may not need n upto 10000 as convergence happens much faster. We will just choose n upto 100.

```

%% Discrete Derivative and Sampling Time

f = @(t) 3.*t.*t + 4.*t + 5;
true_diff_f = @(t) 6.*t + 4;
t1 = 0:1:10;
x1 = f(t1);
t2 = 0:0.5:10;
x2 = f(t2);
t3 = 0:0.05:10;
x3 = f(t3);
y1 = delta_f(t1, x1);
y2 = delta_f(t2, x2);
y3 = delta_f(t3, x3);
y_true = true_diff_f(t3);
f = figure;
plot(t1, y1, 'LineWidth',2, 'Color','#432423', 'DisplayName','\Delta t = 1');
hold on;
plot(t2, y2, 'LineWidth',2, 'Color','#A42368', 'DisplayName','\Delta t = 0.5');
plot(t3, y3, 'LineWidth',2, 'Color','#989233', 'DisplayName','\Delta t = 0.05');
plot(t3, y_true, 'LineWidth',2, 'Color','#F14056', ...
     'LineStyle','--', 'DisplayName','True Derivative');
grid on;
grid minor;
set(gca, 'XColor', [0, 0, 0], 'YColor', [0, 0, 0], 'TickDir', 'out');
xaxis = get(gca, 'XAxis');
xaxis.TickLabelInterpreter = 'latex';
yaxis = get(gca, 'YAxis');
yaxis.TickLabelInterpreter = 'latex';
set(gca, 'FontSize', 18);
xlabel('t','Interpreter', 'latex');
ylabel('Derivative of f','Interpreter', 'latex');
legend('Location', 'northwest');
exportgraphics(f, 'figures/HW01_Discrete_Derivative_Sampling_Time.pdf', ...
    'BackgroundColor', 'none');
function y = delta_f(t, x)
    y = zeros(1, length(t));
    y(1) = 0.0;
    for i = 2:length(t)
        y(i) = (x(i) - x(i-1))./(t(i) - t(i-1));
    end
end
end

```




Figure 2: Discrete Derivative and Sampling Time.

7 Area Under Curve (10 Points)

Compute the integration of $f(t) = t^2$ as an area under the curve numerically. Consider the lower bound as $t = 0$ and upper bound as $t = 5$. For this part of the assignment, you will write your own function. Create time bins with different sampling time: (i) 0.1 seconds; (ii) 0.5 seconds; (iii) and 1 second. Approximate area as sum of rectangles with height of each rectangle being the left timestamp value. Compute area under all three conditions. How much they differ from the true solution?

Answer

Note to grader: If students used some other final time to compute the integration, please give them full marks appropriately.

Consider the initial time to be $t = 0.0$. Further, consider the final time to be $t = 5$.

In this case, the true area under the curve is

$$\int_0^5 t^2 dt = \left. \frac{t^3}{3} \right|_0^5 = \frac{5^3}{3} - 0 = \frac{125}{3} = 41.667 \quad (4)$$

```

%% Area Under Curve
f = @(t) t.^2;
t1 = 0:0.1:5;
t2 = 0:0.5:5;
t3 = 0:1:5;
x1 = f(t1);
x2 = f(t2);
x3 = f(t3);
area_1 = area_under_curve(t1, x1);
area_2 = area_under_curve(t2, x2);
area_3 = area_under_curve(t3, x3);
true_area = 1000.0/3.0;
fprintf("Considering time to be between 0 to 10,\n")
fprintf("True area is %f\n", true_area)
fprintf("Area with Delta t = 0.1 is %f\n", area_1)
fprintf("Area with Delta t = 0.5 is %f\n", area_2)
fprintf("Area with Delta t = 1.0 is %f\n", area_3)

fprintf("Area Difference with Delta t = 0.1 is %f\n", true_area - area_1)
fprintf("Area Difference with Delta t = 0.5 is %f\n", true_area - area_2)
fprintf("Area Difference with Delta t = 1.0 is %f\n", true_area - area_3)

function SUM = area_under_curve(t, x)
    SUM = 0.0;
    for i = 1:length(t)-1
        sum = x(i).*( t(i+1) - t(i) );
        SUM = SUM + sum;
    end
end

```

Output:

```

Considering time to be between 0 to 5,
True area is 41.666667
Area with Delta t = 0.1 is 40.425000
Area with Delta t = 0.5 is 35.625000
Area with Delta t = 1.0 is 30.000000
Area Difference with Delta t = 0.1 is 1.241667
Area Difference with Delta t = 0.5 is 6.041667
Area Difference with Delta t = 1.0 is 11.666667

```

§- §- §