

Context Free Languages

$$G_{pal} = (\{P\}, \{0, 1\}, A, P)$$

1. $P \rightarrow \epsilon$
2. $P \rightarrow 0$
3. $P \rightarrow 1$
4. $P \rightarrow 0P0$
5. $P \rightarrow 1P1$

A context-free grammar for palindromes

Prove that $L(G_{pal})$ is the set of palindromes over the given alphabet.

$$G_{pal} = (\{P\}, \{0, 1\}, A, P)$$

1. $P \rightarrow \epsilon$
2. $P \rightarrow 0$
3. $P \rightarrow 1$
4. $P \rightarrow 0P0$
5. $P \rightarrow 1P1$

A context-free grammar for palindromes

Prove that $L(G_{pal})$ is the set of palindromes over the given alphabet.

- This proof has two parts (\Rightarrow and \Leftarrow)

$$G_{pal} = (\{P\}, \{0, 1\}, A, P)$$

1. $P \rightarrow \epsilon$
2. $P \rightarrow 0$
3. $P \rightarrow 1$
4. $P \rightarrow 0P0$
5. $P \rightarrow 1P1$

A context-free grammar for palindromes

- This proof has two parts (\Rightarrow and \Leftarrow)
 - 1) $(w = w^R) \Rightarrow w \in L(G_{pal})$
 - 2) $w \in L(G_{pal}) \Rightarrow (w = w^R)$

- This proof has two parts (\Rightarrow and \Leftarrow)

$$1) (w = w^R) \Rightarrow w \in L(G_{pal})$$

$$2) w \in L(G_{pal}) \Rightarrow (w = w^R)$$

$$(w = w^R) \Rightarrow w \in L(G_{pal})$$

1. $P \rightarrow \epsilon$
2. $P \rightarrow 0$
3. $P \rightarrow 1$
4. $P \rightarrow 0P0$
5. $P \rightarrow 1P1$

A context-free grammar for palindromes

- Proof [by induction on $|w|$]:

BASIS: We use lengths 0 and 1 as the basis.

If $|w| = 0$ or $|w| = 1$, then w is ϵ , 0, or 1.

Since there are productions $P \rightarrow \epsilon$, $P \rightarrow 0$, and $P \rightarrow 1$, we conclude that $P \xRightarrow{*} w$ in any of these basis cases.

- Note, $w \in L(G_{pal})$ is same $P \xRightarrow{*} w$
- Note, $(w = w^R)$ means w begins and ends with the same character.

BASIS: We use lengths 0 and 1 as the basis.

If $|w| = 0$ or $|w| = 1$, then w is ϵ , 0, or 1.

Since there are productions $P \rightarrow \epsilon$, $P \rightarrow 0$, and $P \rightarrow 1$, we conclude that $P \xRightarrow{*} w$ in any of these basis cases.

BASIS: We use lengths 0 and 1 as the basis.

If $|w| = 0$ or $|w| = 1$, then w is ϵ , 0, or 1.

Since there are productions $P \rightarrow \epsilon$, $P \rightarrow 0$, and $P \rightarrow 1$, we conclude that $P \xRightarrow{*} w$ in any of these basis cases.

Inductive Hypothesis: Let for $|w| \leq k$ where $(w = w^R)$, $P \xRightarrow{*} w$ is true.

Inductive Step: We need to show for $|w| = k + 1$, $P \xRightarrow{*} w$ is true.

BASIS: We use lengths 0 and 1 as the basis.

If $|w| = 0$ or $|w| = 1$, then w is ϵ , 0, or 1.

Since there are productions $P \rightarrow \epsilon$, $P \rightarrow 0$, and $P \rightarrow 1$, we conclude that $P \xRightarrow{*} w$ in any of these basis cases.

Inductive Hypothesis: Let for $|w| \leq k$ where $(w = w^R)$, $P \xRightarrow{*} w$ is true.

Inductive Step: We need to show for $|w| = k + 1$, $P \xRightarrow{*} w$ is true.

Note, $w = 0x0$ or $w = 1x1$, where $|x| = k - 1$.

Then, $P \Rightarrow 0P0 \xRightarrow{*} 0x0$ (Since $|x| \leq k$, so $P \xRightarrow{*} x$ is true).

So, $P \xRightarrow{*} w$ is true. With a similar argument, $P \Rightarrow 1P1 \xRightarrow{*} 1x1$

BASIS: We use lengths 0 and 1 as the basis.

If $|w| = 0$ or $|w| = 1$, then w is ϵ , 0, or 1.

Since there are productions $P \rightarrow \epsilon$, $P \rightarrow 0$, and $P \rightarrow 1$, we conclude that $P \xRightarrow{*} w$ in any of these basis cases.

Inductive Hypothesis: Let for $|w| \leq k$ where $(w = w^R)$, $P \xRightarrow{*} w$ is true.

Inductive Step: We need to show for $|w| = k + 1$, $P \xRightarrow{*} w$ is true.

Note, $w = 0x0$ or $w = 1x1$, where $|x| = k - 1$.

Then, $P \Rightarrow 0P0 \xRightarrow{*} 0x0$ (Since $|x| \leq k$, so $P \xRightarrow{*} x$ is true).

So, $P \xRightarrow{*} w$ is true. With a similar argument, $P \Rightarrow 1P1 \xRightarrow{*} 1x1$

This completes our proof for : $(w = w^R) \Rightarrow w \in L(G_{pal})$

$$w \in L(G_{pal}) \Rightarrow (w = w^R)$$

- Proof [by induction on number of steps in the derivation]:

BASIS: If the derivation is one step, then it must use one of the three productions that do not have P in the body. That is, the derivation is $P \Rightarrow \epsilon$, $P \Rightarrow 0$, or $P \Rightarrow 1$. Since ϵ , 0 , and 1 are all palindromes, the basis is proven.

INDUCTION:

- Assume for n steps it is true.
- Then, show for $(n+1)$ steps it must be true.

Left as an exercise.

Sentential Forms

$G = (\bar{V}, T, P, S)$ is a $\bar{\text{CFG}}$, then any string α in $(V \cup T)^*$ such that $S \xRightarrow{*} \alpha$ is a *sentential form*.

Sentential Forms

$G = (\bar{V}, T, P, S)$ is a CFG, then any

string α in $(V \cup T)^*$ such that $S \xRightarrow{*} \alpha$ is a *sentential form*.

1. $E \rightarrow I$
2. $E \rightarrow E + E$
3. $E \rightarrow E * E$
4. $E \rightarrow (E)$
5. $I \rightarrow a$
6. $I \rightarrow b$
7. $I \rightarrow Ia$
8. $I \rightarrow Ib$
9. $I \rightarrow I0$
10. $I \rightarrow I1$

A context-free grammar for simple expressions

$$E \Rightarrow E * E \Rightarrow E * (E) \Rightarrow E * (E + E) \Rightarrow E * (I + E)$$

Sentential Forms

$G = (\bar{V}, T, P, S)$ is a $\bar{\text{CFG}}$, then any string α in $(V \cup T)^*$ such that $S \xRightarrow{*} \alpha$ is a *sentential form*.

If $S \xRightarrow[lm]{*} \alpha$, then α is a *left-sentential form*,

and if $S \xRightarrow[rm]{*} \alpha$, then α is a *right-sentential form*.

Note that the language $L(G)$ is those sentential forms that are in T^* ; i.e., they consist solely of terminals.

1. $E \rightarrow I$
2. $E \rightarrow E + E$
3. $E \rightarrow E * E$
4. $E \rightarrow (E)$

5. $I \rightarrow a$
6. $I \rightarrow b$
7. $I \rightarrow Ia$
8. $I \rightarrow Ib$
9. $I \rightarrow I0$
10. $I \rightarrow I1$

A context-free grammar for simple expressions

$$E \Rightarrow E * E \Rightarrow E * (E) \Rightarrow E * (E + E) \Rightarrow E * (I + E)$$

- Is this sentential form left-sentential? Or right-sentential?

1. $E \rightarrow I$
2. $E \rightarrow E + E$
3. $E \rightarrow E * E$
4. $E \rightarrow (E)$
5. $I \rightarrow a$
6. $I \rightarrow b$
7. $I \rightarrow Ia$
8. $I \rightarrow Ib$
9. $I \rightarrow I0$
10. $I \rightarrow I1$

A context-free grammar for simple expressions

$$E \Rightarrow E * E \Rightarrow E * (E) \Rightarrow E * (E + E) \Rightarrow E * (I + E)$$

- Is this sentential form left-sentential? Or right-sentential?
- It is a sentential form. But neither left nor right.

Exercise 5.1.2: The following grammar generates the language of regular expression $0^*1(0+1)^*$:

$$\begin{aligned} S &\rightarrow A1B \\ A &\rightarrow 0A \mid \epsilon \\ B &\rightarrow 0B \mid 1B \mid \epsilon \end{aligned}$$

Give leftmost and rightmost derivations of the following strings:

* a) 00101.

b) 1001.

c) 00011.

Note, the given grammar is not a regular grammar (even-though it generates a regular language).

Can you find $L(G)$?

- $S \rightarrow aS|bS|a|b|\epsilon$

Can you find $L(G)$?

- $S \rightarrow aS|bS|a|b|\epsilon$
- Answer: All strings. Σ^*

Can you find $L(G)$?

1. $S \rightarrow S_1 S | \epsilon,$

2. $S_1 \rightarrow a S_1 b | ab$

Can you find $L(G)$?

1. $S \rightarrow S_1 S | \epsilon,$
2. $S_1 \rightarrow a S_1 b | ab$

Recall that the $S \rightarrow a S b | \epsilon$ generates $\{a^n b^n | n \geq 0\}$.

Can you find $L(G)$?

1. $S \rightarrow S_1 S | \epsilon,$

2. $S_1 \rightarrow a S_1 b | ab$

Recall that the $S \rightarrow a S b | \epsilon$ generates $\{a^n b^n | n \geq 0\}$.

Starting from S_1 we get $\{a^n b^n | n \geq 1\}$

Can you find $L(G)$?

1. $S \rightarrow S_1 S | \epsilon,$
2. $S_1 \rightarrow a S_1 b | ab$

Recall that the $S \rightarrow a S b | \epsilon$ generates $\{a^n b^n | n \geq 0\}$.

Starting from S_1 we get $\{a^n b^n | n \geq 1\}$

The answer:

$$a^{n_1} b^{n_1} a^{n_2} b^{n_2} \dots a^{n_k} b^{n_k} \in L(G)$$

$$L(G) = (\{a^n b^n | n \geq 1\})^*$$

Can you find $L(G)$?

$$S \rightarrow SS \mid [S] \mid (S) \mid [] \mid ()$$

Can you find $L(G)$?

$$S \rightarrow SS \mid [S] \mid (S) \mid [] \mid ()$$

Set of all balanced parentheses with alphabet
 $\{ (,), [,] \}$

Can you find $L(G)$?

1. $S \rightarrow aB|bA$

2. $B \rightarrow b|bS|aBB$

3. $A \rightarrow a|aS|bAA$

Can you find $L(G)$?

1. $S \rightarrow SaSbS | SbSaS | \epsilon$