

## 8. Continuous random variables

## Last time

- Dynamics of discrete stochastic processes
- Markov chains
- Stationary distribution

## Goals for today

- Continuous limit
- Continuous probability distributions
- Probability density function

## Geometric random variable

- Recall **geometric random variable**  $X$

## Geometric random variable

- Recall **geometric random variable**  $X$
- Integer time steps
- Probability to decay / recover at each time step is  $p$

## Geometric random variable

- Recall **geometric random variable**  $X$
- Integer time steps
- Probability to decay / recover at each time step is  $p$
- $\mathbb{P}(X = n) = p(1 - p)^{n-1}$  for  $n = 1, 2, \dots$

## Probability mass function

- For a **discrete(-valued) random variable**  $X$ :
- Define **probability mass function**  $f_X$  as

$$f_X(n) := \mathbb{P}[X = n]$$

i.e. the probability that  $X$  takes the value  $n$

## Probability mass function

- For a **discrete(-valued) random variable**  $X$ :
- Define **probability mass function**  $f_X$  as

$$f_X(n) := \mathbb{P}[X = n]$$

i.e. the probability that  $X$  takes the value  $n$

- For geometric random variable  $X$  we have

$$f_X(n) := p(1 - p)^{n-1}$$



## Probability mass function

- For a **discrete(-valued) random variable**  $X$ :
- Define **probability mass function**  $f_X$  as

$$f_X(n) := \mathbb{P}[X = n]$$

i.e. the probability that  $X$  takes the value  $n$

- For geometric random variable  $X$  we have

$$f_X(n) := p(1 - p)^{n-1}$$

- Satisfies  $\sum_{n=1}^{\infty} f_X(n) = 1$

## Towards continuous time

- Suppose want jumps not only at integer times
- Jump at times  $t_n := n \delta$  for  $n = 1, 2, \dots$

## Towards continuous time

- Suppose want jumps not only at integer times
- Jump at times  $t_n := n \delta$  for  $n = 1, 2, \dots$
- E.g. if  $\delta = \frac{1}{2}$ , jump twice per day

## Towards continuous time

- Suppose want jumps not only at integer times
- Jump at times  $t_n := n \delta$  for  $n = 1, 2, \dots$
- E.g. if  $\delta = \frac{1}{2}$ , jump twice per day
- If we want to *reproduce* same dynamics, how should we choose decay probability  $p$ ?

## Towards continuous time

- Suppose want jumps not only at integer times
- Jump at times  $t_n := n \delta$  for  $n = 1, 2, \dots$
- E.g. if  $\delta = \frac{1}{2}$ , jump twice per day
- If we want to *reproduce* same dynamics, how should we choose decay probability  $p$ ?
- $p$  must depend on  $\delta$ , so call it  $p(\delta)$

## Towards continuous time

- In order to compare the two different processes, can't look at instantaneous jumps

## Towards continuous time

- In order to compare the two different processes, can't look at instantaneous jumps
- Need to look at **cumulative distribution function**:

$$F_X(n) := \mathbb{P}[X \leq n]$$

## Towards continuous time

- In order to compare the two different processes, can't look at instantaneous jumps
- Need to look at **cumulative distribution function**:

$$F_X(n) := \mathbb{P}[X \leq n]$$

- “The probability that have decayed *by time*  $n$ ”



## Towards continuous time

- In order to compare the two different processes, can't look at instantaneous jumps
- Need to look at **cumulative distribution function**:

$$F_X(n) := \mathbb{P}[X \leq n]$$

- “The probability that have decayed *by time*  $n$ ”
- Actually now can talk about any *continuous* (real) time  $t$ :

$$F_X(t) := \mathbb{P}[X \leq t]$$

## Towards continuous time II

- How can we write the cumulative distribution function in terms of  $f_X$ ?

## Towards continuous time II

- How can we write the cumulative distribution function in terms of  $f_X$ ?
- We need to find the largest integer value less than  $t$

## Towards continuous time II

- How can we write the cumulative distribution function in terms of  $f_X$ ?
- We need to find the largest integer value less than  $t$
- We obtain

$$F_X(t) = \sum_{j=1}^{\lfloor t \rfloor} f_X(j)$$

## Towards continuous time II

- How can we write the cumulative distribution function in terms of  $f_X$ ?
- We need to find the largest integer value less than  $t$
- We obtain

$$F_X(t) = \sum_{j=1}^{\lfloor t \rfloor} f_X(j)$$

- $\lfloor t \rfloor$  is the **floor** function:
- $\lfloor t \rfloor :=$  largest integer  $\leq t$
- `floor(t)` in Julia

## Towards continuous time III

- Now think about jumps at times  $t_n = n\delta$
- Call resulting process  $Y_\delta$

## Towards continuous time III

- Now think about jumps at times  $t_n = n\delta$
- Call resulting process  $Y_\delta$
- Also a discrete(-valued) random variable

## Towards continuous time III

- Now think about jumps at times  $t_n = n\delta$
- Call resulting process  $Y_\delta$
- Also a discrete(-valued) random variable
  
- Its probability mass function is

$$f_{Y_\delta}(n) = p(\delta) [1 - p(\delta)]^{n-1}$$



## Towards continuous time III

- Now think about jumps at times  $t_n = n\delta$
- Call resulting process  $Y_\delta$
- Also a discrete(-valued) random variable
  
- Its probability mass function is

$$f_{Y_\delta}(n) = p(\delta) [1 - p(\delta)]^{n-1}$$

- This is the probability after  $n$  *time-steps*
- But now these jumps occur at *times*  $n\delta$

## Towards continuous time IV

- We need  $F_{Y_\delta}(t)$
- Probability that  $Y_\delta$  has decayed by *time*  $t$

## Towards continuous time IV

- We need  $F_{Y_\delta}(t)$
- Probability that  $Y_\delta$  has decayed by *time*  $t$
- We showed on Problem Set 2 that

$$F_X(n) = 1 - (1 - p)^n$$

## Towards continuous time IV

- We need  $F_{Y_\delta}(t)$
- Probability that  $Y_\delta$  has decayed by *time*  $t$
- We showed on Problem Set 2 that

$$F_X(n) = 1 - (1 - p)^n$$

- For  $Y_\delta$  we use  $t = n \delta$ , so  $n = t/\delta$

## Towards continuous time IV

- We need  $F_{Y_\delta}(t)$
- Probability that  $Y_\delta$  has decayed by *time*  $t$
- We showed on Problem Set 2 that

$$F_X(n) = 1 - (1 - p)^n$$

- For  $Y_\delta$  we use  $t = n \delta$ , so  $n = t/\delta$
- So

$$F_{Y_\delta}(t) = 1 - [1 - p(\delta)]^{\lfloor t/\delta \rfloor}$$

## Continuous limit

- We could impose that at integers  $n$  we want to have the same probability as for the original process  $X$

## Continuous limit

- We could impose that at integers  $n$  we want to have the same probability as for the original process  $X$
- Instead let's look directly at the **continuous limit**
- i.e.  $\delta \rightarrow 0$

## Continuous limit

- We could impose that at integers  $n$  we want to have the same probability as for the original process  $X$
- Instead let's look directly at the **continuous limit**
- i.e.  $\delta \rightarrow 0$
- We need to take  $p(\delta)$  in a way that “makes sense”
- I.e. where nothing goes to  $\infty$  or to 0



## Continuous limit II

- When  $\delta \rightarrow 0$  we need  $p(\delta) \rightarrow 0$ , since otherwise we will have an infinite number of finite decays in a finite time

## Continuous limit II

- When  $\delta \rightarrow 0$  we need  $p(\delta) \rightarrow 0$ , since otherwise we will have an infinite number of finite decays in a finite time
- When  $\delta \rightarrow 0$ , we can approximate

$$1 - p(\delta) \simeq \exp[-p(\delta)]$$

## Continuous limit II

- When  $\delta \rightarrow 0$  we need  $p(\delta) \rightarrow 0$ , since otherwise we will have an infinite number of finite decays in a finite time
- When  $\delta \rightarrow 0$ , we can approximate

$$1 - p(\delta) \simeq \exp[-p(\delta)]$$

- So

$$F_{Y_\delta}(t) \simeq 1 - \{\exp[-p(\delta)]\}^{t/\delta}$$

## Continuous limit II

- When  $\delta \rightarrow 0$  we need  $p(\delta) \rightarrow 0$ , since otherwise we will have an infinite number of finite decays in a finite time
- When  $\delta \rightarrow 0$ , we can approximate

$$1 - p(\delta) \simeq \exp[-p(\delta)]$$

- So

$$F_{Y_\delta}(t) \simeq 1 - \{\exp[-p(\delta)]\}^{t/\delta}$$

- Hence

$$F_{Y_\delta}(t) \simeq 1 - \{\exp[-p(\delta)]\}^{t/\delta}$$

## Continuous limit III

- We find

$$F_{Y_\delta}(t) \simeq 1 - \exp \left[ -p(\delta) \frac{t}{\delta} \right]$$

## Continuous limit III

- We find

$$F_{Y_\delta}(t) \simeq 1 - \exp \left[ -p(\delta) \frac{t}{\delta} \right]$$

- Hence we need  $p(\delta) \propto \delta$  when  $\delta \rightarrow 0$

## Continuous limit III

- We find

$$F_{Y_\delta}(t) \simeq 1 - \exp \left[ -p(\delta) \frac{t}{\delta} \right]$$

- Hence we need  $p(\delta) \propto \delta$  when  $\delta \rightarrow 0$
- Take

$$p(\delta) =: \lambda \delta$$

## Continuous limit III

- We find

$$F_{Y_\delta}(t) \simeq 1 - \exp \left[ -p(\delta) \frac{t}{\delta} \right]$$

- Hence we need  $p(\delta) \propto \delta$  when  $\delta \rightarrow 0$
- Take

$$p(\delta) =: \lambda \delta$$

- $\lambda$  is called the **rate** of the continuous process
- Decay probability *per unit time*
- Limiting **continuous random variable** with



## Exponential distribution

- A random variable  $Z$  with the cumulative distribution function

$$F(t) = 1 - \exp(-\lambda t)$$

is called an **exponential random variable**

# Exponential distribution

- A random variable  $Z$  with the cumulative distribution function

$$F(t) = 1 - \exp(-\lambda t)$$

is called an **exponential random variable**

- $F_Z(t) = \mathbb{P}(Z \leq t)$
- Can write

$$F_Z(t) = \int_0^t f_Z(s) ds$$

- $f_Z(s)$  is the **probability density function**

# Continuous random variables

- What is a continuous random variable?}
- Random procedure where outcome can take **continuous range of values**
- E.g. `rand()`: outcome any real number between 0 and 1
- So called **continuous random variable**

## Summary statistics

- **Mean** and **variance** make sense, just as for discrete random variables.
- How describe **probability distribution** of continuous random variable?
- For discrete random variable *count* number of times each value occurred
- Impossible for continuous random variables
- Uncountably infinite possible values for outcome

## We can't count

- For (many) continuous random variables  $X$  we have
$$\mathbb{P}(X = x) = 0 \quad \forall x$$
- Never expect to repeat outcomes in a simulation
- Counting is useless!
- But values still concentrate around  $\pi$  (mean / expectation) as in discrete case
- How replace counting?

## Probability density function (PDF)

- Idea: Calculate  $\mathbb{P}(a \leq X \leq b)$
- I.e. prob. that outcome *lies in certain range*
- For discrete r.v.s this is the *sum* of probabilities
- Analogous idea for continuous r.v.s: *integral*
- So “expect”

$$\mathbb{P}(a \leq X \leq b) = \int_a^b f_X(x) dx$$

for some function  $f_X$

- NB: This is *not* always true

## Probability density function II

- $f_X$  is the **probability density function** of  $X$
- $f_X(x) dx$  is prob. that  $X \in [x, x + dx]$
- $f_X$  is not a probability; it's a *density* of probability

## Calculating a PDF: histograms

- It's “easy” to calculate approximations of the PDF
- Fix **bin width**  $h$
- Bin edges  $x_n := x_0 + h n$
- *Count* points in  $[x_n, x_{n+1})$
- Do this for several such intervals to get **histogram**



# Histograms II

- Draw bar whose *area* is proportional to frequency in that bin
- Sum of areas = 1
- How choose bin width?
- Choose to give “best” result. Several interpretations
- Alternative: **kernel density estimate**: for each  $x$ , count number of points near  $x$

# Histograms in Julia

## ■ Three options:

1 Make your own!

2 `histogram(data)` function in `Plots.jl`:

- Draws histogram
- Does not allow access to data in histogram

3 `fit(Histogram, data)` in `StatsBase.jl`:

- Need `StatsPlots.jl` to plot
- Returns data

```
fit(Histogram, data)
```

```
using StatsBase
```

```
data = rand(100)
```

```
h = fit(Histogram, data, nbins=50)
```

```
using StatsPlots
```

```
plot(h)
```

## Cumulative distribution function (CDF)

- Histograms lose information: lump data together in single bin
- Cumulative distribution function does not lose information:

$$F(x) := \mathbb{P}(X \leq x)$$

- Empirical CDF: Step function that increases at each data point

# Normal distribution

- PDF of standard normal distribution:

$$f(x) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2}\right)$$

- Famous bell curve
- CDF cannot be written in terms of standard functions
- Introduce new “error function”, erf
- Quadratic on log-linear (log  $y$ -axis)

## Why is the normal distribution so ubiquitous?

- **Central limit theorem:**  
*Sum of independent random variables converges to a normal distribution*
- Limiting shape of “centre” of distribution (not tails)
- Summands (things being summed) can be *different*

## Why is the CLT true?

- Dice example (PS2): means increase linearly; standard deviations increase *slower*
- So everything concentrates around mean with zero (relative) width in limit
- CLT: centre around mean and *rescale*; obtain limiting normal shape
- Says how positive and negative deviations tend to cancel each other
- PDF does *not* always “converge”: **weak convergence**

# Does the Central Limit Theorem always hold?

- No!
- Only if mean and variance are finite
- e.g. Sample from a Pareto distribution (power-law tail)

$\alpha = 4$

```
data = [sum(rand(Pareto( $\alpha$ , 1.0), 100)) for i in 1:10000]
histogram(data) # satisfies CLT
```

$\alpha = 1.5$

```
data = [sum(rand(Pareto( $\alpha$ , 1.0), 100)) for i in 1:10000]
histogram(data) # doesn't satisfy CLT
```

- Then convergence to other distributions: Lévy stable distributions
- Long tail often corresponds to some kind of “memory



# Review

- Exact first-passage distribution and diverging (infinite) mean hitting time
- Continuous random variables
- Probability density function (PDF)
- Central Limit Theorem