

Supplemental Information

Comprehensive Analysis of Protein Glycation Reveals Its Potential Impacts on Protein Degradation and Gene Expression in Human Cells

Fangxu Sun[‡], Suttipong Suttapitugsakul[‡], Haopeng Xiao[‡] and Ronghu Wu^{‡,*}

[‡]School of Chemistry and Biochemistry and the Petit Institute for Bioengineering and Bioscience,
Georgia Institute of Technology, Atlanta, Georgia 30332, USA

Supplemental Figures:

Figure S1. The abundance distributions of proteins from the database and the glycosylated proteins identified in this work.

Figure S2. The distribution of the glycosylated sites identified on lysine (K) and arginine (R) in HEK293T, Jurkat and MCF7 cells.

Figure S3. Protein clustering based on cellular component (A) and biological process (B).

Figure S4. Normalized frequency of different types of amino acid residues around the identified lysine glycosylation sites (five residues from the center lysine).

Supplemental Tables (in Excel files):

Table S1. Glycosylation sites identified in MCF7 cells in the duplicate experiments.

Table S2. Identification of glycosylation sites in HEK293T, Jurkat and MCF7 cells.

Table S3. Identification of glycosylated proteins involved in the glycolysis in HEK293T, Jurkat and MCF7 cells.




Figure S1. The abundance distributions of proteins from the database and the glycosylated proteins identified in this work.




Figure S2. The distribution of the glycosylated sites identified on lysine (K) and arginine (R) in HEK293T, Jurkat and MCF7 cells.




Figure S3. Protein clustering based on cellular component (A) and biological process (B).




Figure S4. Normalized frequencies of different types of amino acid residues around the identified lysine glycation sites (five residues from the center lysine).