

Cartesian Products

Solutions by: Raindrops_drop

September 21, 2026

Exercises

1. Show there is a bijective correspondence of $A \times B$ and $B \times A$.

Proof. It is claimed that the function $f : A \times B \rightarrow B \times A$ defined as $f((a, b)) = (b, a)$. To show f is surjective, consider $(b, a) \in B \times A$. Then, by definition $b \in B$ and $a \in A$. As such, $(a, b) \in A \times B$, and therefore $f((a, b)) = (b, a)$. Thus, since (b, a) was an arbitrary element of $B \times A$ it has been shown that for any element in $s \in B \times A$ there exists at least one element $t \in A \times B$ such that $f(t) = s$. Hence, f is surjective.

Consider now $(a, b), (c, d) \in A \times B$ such that $f((a, b)) = f((c, d)) = (s, t)$. By definition of f , note $a = t$ and $c = t$ implying $a = c$. Similarly, it must be that $b = d$, and thus $(a, b) = (c, d)$. Therefore, f is injective and it can be thus concluded that f is bijective. $\square$

2.

- (a) Show that if $n > 1$ there is a bijective correspondence of

$$A_1 \times \cdots \times A_n \quad \text{with} \quad (A_1 \times \cdots \times A_{n-1}) \times A_n.$$

Proof. If $n = 2$, note $A_1 \times A_2 = (A_1) \times A_2$ and thus there exists a bijective correspondence between both sets, in particular this bijective map is the identity map. Thus, assume $n > 2$. We claim that $f : A_1 \times \cdots \times A_n \rightarrow (A_1 \times \cdots \times A_{n-1}) \times A_n$ defined by $(a_1, \dots, a_n) \mapsto ((a_1, \dots, a_{n-1}), a_n)$ is bijective. We begin by showing f is injective. Let $(a_1, \dots, a_n)$ and $(b_1, \dots, b_n)$ be elements in the domain set such that they both map to $((c_1, \dots, c_{n-1}), c_n)$. By construction, $a_n = c_n$ and $b_n = c_n$ implying $a_n = b_n$. Similarly, $a_1, \dots, a_{n-1}$ must all equal the corresponding element in $(c_1, \dots, c_{n-1})$, i.e., $a_1 = c_1$ and so on. In the same way, $b_i = c_i$ for $1 \leq i \leq n-1$. Therefore, $a_i = b_i$ for $1 \leq i \leq n-1$, and we thus conclude $(a_1, \dots, a_n) = (b_1, \dots, b_n)$ making f injective. To show f is surjective, consider the element $((c_1, \dots, c_{n-1}), c_n)$ in the set being mapped onto by f . By construction, $c_i \in A_i$ for $1 \leq i \leq n$. Thus, $(c_1, \dots, c_n) \in A_1 \times \cdots \times A_n$. By definition of f , $f(c_1, \dots, c_n) = ((c_1, \dots, c_{n-1}), c_n)$. Thus, we have shown f is also surjective, and therefore we conclude f is bijective. $\square$

- (b) Given the indexed family $\{A_1, A_2, \dots\}$ let $B_i = A_{2i-1} \times A_{2i}$ for each positive integer i . Show there is a bijective correspondence of $A_1 \times A_2 \times \cdots$ with $B_1 \times B_2 \times \cdots$.

Proof. Note $B_1 \times B_2 \times \cdots$ is equivalent to the set $(A_1 \times A_2) \times (A_3 \times A_4) \times \cdots$ by construction. $\square$

3. Let $A = A_1 \times A_2 \times \cdots$ and $B = B_1 \times B_2 \times \cdots$.
- (a) Show that if $B_i \subset A_i$ for all i , then $B \subset A$. (Strictly speaking, if we are given a function mapping the index set $\mathbb{Z}_+$ into the union of the sets B_i , we must change its range before it can be considered as a function mapping $\mathbb{Z}_+$ into the union of the sets A_i . We shall ignore this technicality when dealing with Cartesian products).
 - (b) Show the converse of (a) holds if B is non-empty.
 - (c) Show that if A is non-empty, each A_i is non-empty. Does the converse hold?
 - (d) What is the relation between the set $A \cup B$ and the Cartesian product of the sets $A_i \cup B_i$? What is the relation between the set $A \cap B$ and the Cartesian product of the sets $A_i \cap B_i$?
4. Let $n, m \in \mathbb{Z}_+$. Let $X \neq \emptyset$.
- (a) If $m \leq n$, find an injective map $f : X^m \rightarrow X^n$.
 - (b) Find a bijective map $g : X^m \times X^n \rightarrow X^{m+n}$.
 - (c) Find an injective map $h : X^n \rightarrow X^\omega$.
 - (d) Find a bijective map $k : X^n \times X^\omega \rightarrow X^\omega$.
 - (e) Find a bijective map $l : X^\omega \times X^\omega \rightarrow X^\omega$.
 - (f) If $A \subset B$, find an injective map $m : (A^\omega)^n \rightarrow B^\omega$.
5. Which of the following subsets of $\mathbb{R}^\omega$ can be expressed as the Cartesian product of subsets of $\mathbb{R}$?
- (a) $\{\mathbf{x} \mid x_i \text{ is an integer for all } i\}$.
 - (b) $\{\mathbf{x} \mid x_i \geq i \text{ for all } i\}$.
 - (c) $\{\mathbf{x} \mid x_i \text{ is an integer for all } i \geq 100\}$.
 - (d) $\{\mathbf{x} \mid x_2 = x_3\}$.