



American Society of Hematology  
2021 L Street NW, Suite 900,  
Washington, DC 20036  
Phone: 202-776-0544 | Fax 202-776-0545  
editorial@hematology.org

## Long-term follow-up of VIALE-C in patients with untreated AML ineligible for intensive chemotherapy

Tracking no: BLD-2022-016963R1

Andrew Wei (Peter MacCallum Cancer Centre and Royal Melbourne Hospital, Australia) Panayiotis Panayiotidis (National and Kapodistrian University of Athens, Greece) Pau Montesinos (Hospital Universitari i Politècnic La Fe, Spain) Kamel Laribi (Department of Hematology, Centre Hospitalier Le Mans, Le Mans, France, France) Vladimir Ivanov (Almazov National Medical Research Center, Russian Federation) Inho Kim (Seoul National University Hospital, Korea, Republic of) Jan Novak (University Hospital Kralovske Vinohrady and Third Faculty of Medicine, Charles University, Czech Republic) Rebecca Champion (Norton Cancer Institute, United States) Walter Fiedler (Department of Oncology, Hematology and Bone Marrow Transplantation with Section Pneumology, Hubertus Wald University Cancer Center, University Medical Center Hamburg-Eppendorf, Hamburg, Germany, Germany) Maria Pagoni (Evangelismos Hospital, Greece) Julie Bergeron (Hôpital Maisonneuve-Rosemont, Canada) Stephen Ting (Eastern Health-Monash University, Australia) Jingzhou Hou (University of Pittsburgh, United States) Achilles Anagnostopoulos (George Papanicolaou Hospital, Greece) Andrew McDonald (Albert Alberts Stem Cell Transplantation Centre, ) Vidhya Murthy (Heartlands Hospital, part of UHB, United Kingdom) Takahiro Yamauchi (Department of Hematology and Oncology, University of Fukui, Japan) Jianxiang Wang (Institute of Hematology & Blood Diseases Hospital, CAMS&PUMC, China) Qi Jiang (AbbVie, Inc., United States) Yan Sun (AbbVie, Inc, United States) Brenda Chyla (Abbvie, United States) Wellington Mendes (AbbVie, Inc., United States) Courtney DiNardo (UT MD Anderson Cancer Center, United States)

### Abstract:

**Conflict of interest:** COI declared - see note

**COI notes:** AbbVie sponsored the study (NCT03069352), contributed to its design, collection, analysis, and interpretation of the data, and participated in the writing, review, and approval of the manuscript. All authors had access to relevant data. No honoraria or payments were made for authorship. Venetoclax (ABT-199/GDC-0199) is being developed in collaboration between AbbVie and Genentech. A.H.W.: Consultant for AbbVie, Amgen, Astellas Pharma, Celgene, Janssen, MacroGenics, Novartis, Roche, and Servier; research funding from AbbVie, Celgene, Novartis, and Servier; employee of Walter and Eliza Hall Institute of Medical Research, which receives royalties related to venetoclax, and Dr Wei is entitled to a fraction of these payments. P.P.: Grant/research support from AbbVie, Genesis, Novartis, and Roche; honoraria from AbbVie, Genesis, Gilead, Janssen, Novartis, and Roche. P.M.: Grant/research support from Astellas Pharma, Celgene, Daiichi Sankyo, Janssen, Karyopharm Therapeutics, Novartis, Pfizer, and Teva; speaker/advisory role for AbbVie, Celgene, Daiichi Sankyo, Incyte, Janssen, Karyopharm Therapeutics, Novartis, Pfizer, Teva, and Tolero; consultant for Agios, Astellas Pharma, Celgene, Daiichi Sankyo, Oryzon, and Tolero. K.L.: Grant/research support from AbbVie, Novartis, Roche, Sandoz, and Takeda; personal fees from AbbVie, Astellas Pharma, BeiGene, Celgene, iQone Healthcare Switzerland, Janssen, Novartis, and Sandoz. V.I.: Investigator in AbbVie-sponsored clinical trials. I.K.: Investigator in AbbVie-sponsored clinical trials. J.N.: Consultant/advisory role for Amgen, Novartis, Pfizer, Roche, and Takeda; travel expenses from Amgen and Janssen. R.C.: Speakers bureau for Bristol Myer Squibb and AbbVie; advisory board for ADC therapeutics and Beigene. W.F.: Membership on an entity's board of directors or advisory committee for AbbVie, Amgen, ARIAD/Incyte, Celgene, Jazz Pharmaceuticals, MorphoSys AG, Novartis, Stemline, Clinigen and Pfizer; patents and royalties from Amgen; support for meeting attendance Amgen, Daiichi Sankyo, Gilead, Jazz Pharmaceuticals, and Servier; research funding from Amgen and Pfizer. M.P.: Speaker/advisory role for AbbVie, Amgen, Astellas Pharma, Genesis, Janssen, Novartis, AstraZeneca, Bristol Meyers Squibb, Gilead, Sobie and Pfizer. J.B.: Consultant for AbbVie, Servier, Amgen, Astellas Pharma, BMS, Jazz Pharmaceuticals, Novartis, and Pfizer; travel support from Amgen and Novartis. S.B.T.: Consultant for AbbVie; investigator in AbbVie-sponsored clinical trials. J.Z.H., A.A., and A.M.: Investigator in AbbVie-sponsored clinical trials. V.M.: Conference attendance support from AbbVie, Celgene, Janssen, Novartis, and Takeda; consultant for AbbVie, Celgene, Novartis, and Janssen. T.Y.: Research support/honoraria from, and advisory role for AbbVie, Astellas Pharma, Gilead, Janssen, Nippon Shinyaku, Otsuka, Pfizer, Solasia, Symbio, and Takeda. J.W.: Advisory role for AbbVie; research support from Celgene. B.C., Y.S., Q.J. and W.M.: Employees of AbbVie and may hold stock or stock options. C.D.D.: Research support from AbbVie/Genentech, Agios, BMS/Celgene, Calithera, Cleave, Daiichi Sankyo, Immune-Onc, and Loxo; consultant/advisory board member for AbbVie, Agios, Aprea, BMS/Celgene, Immune-Onc, Kura, Novartis, Takeda, and Notable Labs.

**Preprint server:** No;

**Author contributions and disclosures:** Conception and design: AbbVie, Inc. Provision, collection, and assembly of data: all authors contributed to data collection. Data analysis and interpretation done by A.H.W., Q.J., Y.S., B.C., and W.M.; all authors contributed thereafter. Manuscript writing, critical revision, and final approval: all authors.

**Non-author contributions and disclosures:** Yes; Medical writing support was provided by Susan Olalekan, PhD of AbbVie, Inc. Editorial support was provided by Angela T. Hadsell of AbbVie, Inc.

**Agreement to Share Publication-Related Data and Data Sharing Statement:** AbbVie is committed to responsible data sharing regarding the clinical trials we sponsor. This includes access to anonymized, individual, and trial-level data (analysis data sets), as well as other information (e.g., protocols and Clinical Study Reports), as long as the trials are not part of an ongoing or planned regulatory submission. This includes requests for clinical trial data for unlicensed products and indications. These clinical trial data can be requested by any qualified researchers who engage in rigorous, independent scientific research, and will be provided following review and approval of a research proposal and Statistical Analysis Plan (SAP) and execution of a Data Sharing Agreement (DSA). Data requests can be submitted at any time after approval in the US and Europe and after acceptance of this manuscript for publication. The data will be accessible for 12 months, with possible extensions considered. For more information on the process, or to submit a request, visit the following link: <https://www.abbvie.com/our-science/clinical-trials/clinical-trials-data-and-information-sharing/data-and-information-sharing-with-qualified-researchers.html>.

**Clinical trial registration information (if any):** NCT03069352, <https://beta.clinicaltrials.gov/>

**Long-term follow-up of VIALE-C in patients with untreated AML ineligible for intensive chemotherapy**

Andrew H. Wei<sup>1</sup>, Panayiotis Panayiotidis<sup>2</sup>, Pau Montesinos<sup>3</sup>, Kamel Laribi<sup>4</sup>, Vladimir Ivanov<sup>5</sup>, Inho Kim<sup>6</sup>, Jan Novak<sup>7</sup>, Rebecca Champion<sup>8</sup>, Walter Fiedler<sup>9</sup>, Maria Pagoni<sup>10</sup>, Julie Bergeron<sup>11</sup>, Stephen B. Ting<sup>12</sup>, Jing-Zhou Hou<sup>13</sup>, Achilles Anagnostopoulos<sup>14</sup>, Andrew McDonald<sup>15</sup>, Vidhya Murthy<sup>16</sup>, Takahiro Yamauchi<sup>17</sup>, Jianxiang Wang<sup>18</sup>, Qi Jiang<sup>19</sup>, Yan Sun<sup>19</sup>, Brenda Chyla<sup>19</sup>, Wellington Mendes<sup>19</sup> and Courtney D. DiNardo<sup>20</sup>

<sup>1</sup>The Peter MacCallum Cancer Centre, Royal Melbourne Hospital and Walter and Eliza Hall Institute of Medical Research, Melbourne, VIC, Australia; <sup>2</sup>National and Kapodistrian University of Athens Medical School, Laiko General Hospital, Athens, Greece; <sup>3</sup>Hospital Universitario y Politécnico La Fe, Valencia, Spain and CIBERONC, Instituto Carlos III, Madrid, Spain; <sup>4</sup>Centre Hospitalier–Le Mans, Le Mans, France; <sup>5</sup>Almazov National Medical Research Center, Saint Petersburg, Russia; <sup>6</sup>Seoul National University Hospital, Seoul, South Korea; <sup>7</sup>Department of Internal Medicine and Hematology, University Hospital Královské Vinohrady and Third Faculty of Medicine, Charles University, Prague, Czech Republic; <sup>8</sup>Norton Cancer Institute, Louisville, KY, USA; <sup>9</sup>Hubertus Wald University Cancer Center, University Medical Center Hamburg-Eppendorf, Hamburg, Germany; <sup>10</sup>Evangelismos General Hospital, Athens Greece; <sup>11</sup>CIUSSS-EMTL, Installation Maisonneuve-Rosemont, Montreal, QC, Canada; <sup>12</sup>Eastern Health and Monash University, Melbourne, VIC, Australia; <sup>13</sup>University of Pittsburgh Medical Center (UPMC) Cancer Center, Pittsburgh, PA, USA; <sup>14</sup>George Papanicolaou General Hospital, Thessalonica, Greece; <sup>15</sup>Netcare Pretoria East Hospital, Moreletapark, Pretoria, South Africa; <sup>16</sup>Heartlands Hospital, Birmingham, UK; <sup>17</sup>University of Fukui Hospital, Fukui, Japan; <sup>18</sup>Chinese Academy of Medical Sciences, Tianjin, China; <sup>19</sup>AbbVie Inc., North Chicago, IL, USA; <sup>20</sup>Department of Leukemia, The University of Texas MD Anderson Cancer Center, Houston, TX, USA

**CORRESPONDING AUTHOR:**

Andrew H. Wei

The Peter MacCallum Cancer Centre and Royal Melbourne Hospital

305 Grattan St, Melbourne, VIC 3000, Australia

Telephone: +61 (0)3 8559 7915

Fax: +61 (0)3 8559 7739

E-mail: [andrew.wei@petermac.org](mailto:andrew.wei@petermac.org)

**Target journal: Blood**

**Manuscript word count: 779**

**Abstract: no abstract**

**Tables/figures limit: 1/2**

**References: 3/25**

**Supplementary information:**

**Running title: 2-yr update of venetoclax + LDAC for untreated AML**

**Keywords:** venetoclax, BCL-2 inhibitor, low-dose cytarabine, VIALE-C, acute myeloid leukemia

Venetoclax (VEN) in combination with low dose cytarabine (LDAC) is FDA-approved for the treatment of unfit patients with newly diagnosed AML ineligible for intensive chemotherapy, based on a response rate of 54% (complete remission with or without blood count recovery [CR/CRi]) in the original phase Ib/II study.<sup>1</sup> The VIALE-C phase 3 study (NCT03069352), compared VEN vs placebo (PBO) in combination with LDAC in 210 patients with untreated AML ineligible for intensive chemotherapy.<sup>2,3</sup> The primary overall survival (OS) endpoint was event-driven and did not show a significant benefit in favor of VEN + LDAC after a median follow-up time of 12 months.<sup>2</sup> This initial analysis was associated with substantial early censoring of patients with <6 months follow-up. In a subsequent post hoc analysis with median follow-up 17.5 months (range 0.1 – 23.5), median OS was significantly longer in patients receiving VEN + LDAC (8.4 vs. 4.1 months; HR = 0.70, 95% CI 0.50, 0.99;  $P = 0.04$ ). Rates of CR/CRi were higher for patients receiving VEN + LDAC (48.3%), compared to PBO + LDAC (13.2%). In the present study, a final analysis with 2-years additional follow-up was undertaken to determine if the survival benefit of VEN +LDAC was sustained. In addition, clinical and molecular correlates of survival among patients receiving VEN + LDAC were assessed. These analyses demonstrated that survival outcome was influenced by prior exposure to hypomethylating agents, clinical response, cytogenetic risk and molecular genotype, with best outcomes observed for patients with *NPM1* mutation. This longer-term final analysis confirmed the survival improvement of VEN + LDAC in patients unfit for intensive chemotherapy.

At last follow-up on 15 February 2021 (median follow-up 34.7 months; range 0.1 – 41.3), 83.9% ( $n = 120$ ; Ven + LDAC) and 89.7% ( $n = 61$ ; PBO + LDAC) of patients had died with 7% of patients ( $n = 10$ ) still receiving Ven + LDAC. No patient was receiving LDAC + PBO at this follow-up. Two patients on Ven + LDAC were lost to follow-up, and five withdrew consent (VEN + LDAC 3 [2.1%] and PBO + LDAC 2 [2.9%]). With an additional 2 years of follow-up from the

last analysis,<sup>3</sup> improvement in median OS with VEN + LDAC vs PBO + LDAC was unchanged (8.4 vs 4.1 months) (Fig. 1A, Supplementary Table 1). Two-year OS was 21.5% for patients in the VEN + LDAC arm and 12.4% for patients receiving PBO + LDAC (number needed to treat = 11). No new adverse event signal was noted (Supplementary Table 2). We next investigated correlates of outcome in the VEN + LDAC treated arm.

Clinical response rates were similar to the previously published 6-month follow-up (supplemental table 1), with 28.7% and 19.6% achieving CR and CRi, respectively. At two years, clinical response was sustained in 34.3% and 31.6% of patients initially achieving CR or CR/CRi, respectively, in the VEN + LDAC arm, indicating a similar duration of response for both response categories (Fig. 1B). In contrast to the VIALE-A study, patients enrolled in the VIALE-C trial included 20% with a prior history of HMA treatment. Patients not receiving prior HMA treatment had a longer OS than those who received prior HMA therapy, 8.9 months (95% CI 6.6, 10.9) vs 5.6 months (95% CI 3.4, 9.6), respectively (Fig. 1C). Among the 29% of patients achieving CR, median OS was 24.3 months (95% CI 20.1, 28.3)(Fig. 1D). Of the 48% patients achieving CR/CRi, median OS was 20.7 months (95% CI 12.7, 24.5), compared to 3.4 months (95% CI 2.1, 4.1) for those not achieving remission (Fig. 1D). OS for patients with *de novo* AML was 9.2 months (95% CI 7.2, 12.7), compared to 5.6 months (95%CI 3.4, 9.8) in patients with secondary AML (Fig. 1E). Patients categorized according to NCCN classification as intermediate risk had a longer OS, 11.2 months (95% CI 7.9, 16.4) compared to 4.4 months (95%CI 3.0,6.4) in poor risk patients (Fig. 1E). We further analyzed OS according to the presence of somatic mutations in patients treated with VEN + LDAC. Median OS was 25.3 (95% CI 9.9, -), 11.2 (95% CI 3.4, 23.6), 5.9 (95% CI 1.6, 20.8) and 2.8 (95% CI 2.1, 3.6) months in patients with *NPM1*, *IDH1/2*, *FLT3* and *TP53* mutations, respectively (Fig. 1F). These outcomes should be interpreted with caution, due to the limited size of these sub-groups.

The mOS difference between patients with a *NPM1* and *IDH1/2* baseline mutations can be attributed to other prognostic factors such as secondary AML, poor cytogenetic risk and prior HMA treatment. Patients with *IDH1/2* mutation presented secondary AML (38%), poor cytogenetic risk per NCCN 2016 classification (24%) and received prior treatment with HMA (19%) more frequently when compared to *NPM1* patients (16%, 16% and 5%, respectively) (Supplemental table 3).

In conclusion, among patients with newly diagnosed AML ineligible for intensive chemotherapy, longer-term follow-up confirmed that patients receiving VEN + LDAC had longer median OS than patients receiving PBO + LDAC. CR/CRi responses in the VEN + LDAC were durable, with 31.6% remaining in remission for >2 years. Notably, for patients with *NPM1* mutation treated with VEN + LDAC, OS at 24 months was ~50%. In contrast, outcomes for patients with *TP53* mutation remained poor. This 2-year follow-up analysis confirms long-term benefit for patients treated with VEN + LDAC, with no new safety findings.

### Data sharing statement

AbbVie is committed to responsible data sharing regarding the clinical trials we sponsor. This includes access to anonymized, individual, and trial-level data (analysis data sets), as well as other information (e.g., protocols and Clinical Study Reports), as long as the trials are not part of an ongoing or planned regulatory submission. This includes requests for clinical trial data for unlicensed products and indications.

These clinical trial data can be requested by any qualified researchers who engage in rigorous, independent scientific research, and will be provided following review and approval of a research proposal and Statistical Analysis Plan (SAP) and execution of a Data Sharing Agreement (DSA). Data requests can be submitted at any time after acceptance of this manuscript for publication. The data will be accessible for 12 months, with possible extensions considered. For more information on the process, or to submit a request, visit the following link:

<https://www.abbvie.com/our-science/clinical-trials/clinical-trials-data-and-information-sharing/data-and-information-sharing-with-qualified-researchers.html>.

### **Acknowledgements**

AbbVie and the authors thank all the trial investigators and patients who participated in this clinical trial. Medical writing support was provided by Susan Olalekan, PhD of AbbVie, Inc. Editorial support was provided by Angela T. Hadsell of AbbVie, Inc.

### **Author contributions**

Conception and design: AbbVie, Inc. Provision, collection, and assembly of data: all authors contributed to data collection. Data analysis and interpretation done by A.H.W., Q.J., Y.S., B.C., and W.M.; all authors contributed thereafter. Manuscript writing, critical revision, and final approval: all authors.

### **Conflict of interest**

AbbVie sponsored the study (NCT03069352), contributed to its design, collection, analysis, and interpretation of the data, and participated in the writing, review, and approval of the manuscript.



All authors had access to relevant data. No honoraria or payments were made for authorship. Venetoclax (ABT-199/GDC-0199) is being developed in collaboration between AbbVie and Genentech.

A.H.W.: Consultant for AbbVie, Amgen, Astellas Pharma, Celgene, Janssen, MacroGenics, Novartis, Roche, and Servier; research funding from AbbVie, Celgene, Novartis, and Servier; employee of Walter and Eliza Hall Institute of Medical Research, which receives royalties related to venetoclax, and Dr Wei is entitled to a fraction of these payments. P.P.: Grant/research support from AbbVie, Genesis, Novartis, and Roche; honoraria from AbbVie, Genesis, Gilead, Janssen, Novartis, and Roche. P.M.: Grant/research support from Astellas Pharma, Celgene, Daiichi Sankyo, Janssen, Karyopharm Therapeutics, Novartis, Pfizer, and Teva; speaker/advisory role for AbbVie, Celgene, Daiichi Sankyo, Incyte, Janssen, Karyopharm Therapeutics, Novartis, Pfizer, Teva, and Tolero; consultant for Agios, Astellas Pharma, Celgene, Daiichi Sankyo, Oryzon, and Tolero. K.L.: Grant/research support from AbbVie, Novartis, Roche, Sandoz, and Takeda; personal fees from AbbVie, Astellas Pharma, BeiGene, Celgene, iQone Healthcare Switzerland, Janssen, Novartis, and Sandoz. V.I.: Investigator in AbbVie-sponsored clinical trials. I.K.: Investigator in AbbVie-sponsored clinical trials. J.N.: Consultant/advisory role for Amgen, Novartis, Pfizer, Roche, and Takeda; travel expenses from Amgen and Janssen. R.C.: Speakers bureau for Bristol Myer Squibb and AbbVie; advisory board for ADC therapeutics and Beigene. W.F.: Membership on an entity's board of directors or advisory committee for AbbVie, Amgen, ARIAD/Incyte, Celgene, Jazz Pharmaceuticals, MorphoSys AG, Novartis, Stemline, Clinigen and Pfizer; patents and royalties from Amgen; support for meeting attendance Amgen, Daiichi Sankyo, Gilead, Jazz Pharmaceuticals, and Servier; research funding from Amgen and Pfizer. M.P.: Speaker/advisory role for AbbVie, Amgen, Astellas Pharma, Genesis, Janssen, Novartis, AstraZeneca, Bristol Meyers Squibb, Gilead, Sobie and Pfizer. J.B.: Consultant for AbbVie, Servier, Amgen, Astellas Pharma, BMS,

Jazz Pharmaceuticals, Novartis, and Pfizer; travel support from Amgen and Novartis. S.B.T.: Consultant for AbbVie; investigator in AbbVie-sponsored clinical trials. J.Z.H., A.A., and A.M.: Investigator in AbbVie-sponsored clinical trials. V.M.: Conference attendance support from AbbVie, Celgene, Janssen, Novartis, and Takeda; consultant for AbbVie, Celgene, Novartis, and Janssen. T.Y.: Research support/honoraria from, and advisory role for AbbVie, Astellas Pharma, Gilead, Janssen, Nippon Shinyaku, Otsuka, Pfizer, Solasia, Symbio, and Takeda. J.W.: Advisory role for AbbVie; research support from Celgene. B.C., Y.S., Q.J. and W.M.: Employees of AbbVie and may hold stock or stock options. C.D.D.: Research support from AbbVie/Genentech, Agios, BMS/Celgene, Calithera, Cleave, Daiichi Sankyo, Immune-Onc, and Loxo; consultant/advisory board member for AbbVie, Agios, Aprea, BMS/Celgene, Immune-Onc, Kura, Novartis, Takeda, and Notable Labs.

### References (3 out of 25)

1. Wei AH, Strickland Jr SA, Hou J-Z, et al. Venetoclax Combined With Low-Dose Cytarabine for Previously Untreated Patients With Acute Myeloid Leukemia: Results From a Phase Ib/II Study. *Journal of Clinical Oncology*. 2019;JCO. 18.01600.
2. Wei AH, Montesinos P, Ivanov V, et al. Venetoclax plus LDAC for newly diagnosed AML ineligible for intensive chemotherapy: a phase 3 randomized placebo-controlled trial. *Blood*. 2020;135(24):2137-2145.
3. Wei AH, Panayiotidis P, Montesinos P, et al. 6-month follow-up of VIALE-C demonstrates improved and durable efficacy in patients with untreated AML ineligible for intensive chemotherapy. *Blood cancer journal*. 2021;11(10):1-8.

## Figure Legend

**Figure 1. Survival outcomes and response in patients treated with venetoclax compared to placebo.** A) Kaplan-Meier overall survival curves of all patients. Number of patients at risk for each time is shown below and separated by treatment arms. B) Kaplan-Meier duration of response curves in patients treated with VEN + LDAC. Number of patients at risk for each time is shown below and separated by response. C) Kaplan-Meier overall survival by prior HMA treatment curves in patients treated with VEN + LDAC. Number of patients at risk for each time is shown below and separated by prior HMA treatment. D) Kaplan-Meier overall survival by remission status curves in patients treated with VEN + LDAC. Number of patients at risk for each time is shown below and separated by response. E) Kaplan-Meier overall survival by AML type curves in patients treated with VEN + LDAC. Number of patients at risk for each time is shown below and separated by AML type. F) Kaplan-Meier overall survival by mutational status curves in patients treated with VEN + LDAC. Number of patients at risk for each time is shown below and separated by mutational status. AML, acute myeloid leukemia; CR, complete remission; CRi, complete remission with incomplete blood count recovery; DOR, duration of remission; HMA, hypomethylating agent; LDAC, low dose cytarabine; PBO, placebo; VEN, venetoclax

Figure 1

