

# Nuclear Receptors in Energy Metabolism

# 4

Alina A. Walth-Hummel, Stephan Herzig,  
and Maria Rohm

## Abstract

Nuclear receptors are master regulators of energy metabolism through the conversion of extracellular signals into gene expression signatures. The function of the respective nuclear receptor is tissue specific, signal and co-factor dependent. While normal nuclear receptor function is central to metabolic physiology, aberrant nuclear receptor signal-

ing is linked to various metabolic diseases such as type 2 diabetes mellitus, obesity, or hepatic steatosis. Thus, the tissue specific manipulation of nuclear receptors is a major field in biomedical research and represents a treatment approach for metabolic syndrome. This chapter focuses on key nuclear receptors involved in regulating the metabolic function of liver, adipose tissue, skeletal muscle, and pancreatic  $\beta$ -cells. It also addresses the importance of nuclear co-factors for fine-tuning of nuclear receptor function. The mode of action, role in energy metabolism, and therapeutic potential of prominent nuclear receptors is outlined.

A. A. Walth-Hummel · M. Rohm (✉)  
Institute for Diabetes and Cancer, Helmholtz Center  
Munich, Neuherberg, Germany

Joint Heidelberg-IDC Translational Diabetes  
Program, Inner Medicine 1, Heidelberg University  
Hospital, Heidelberg, Germany

German Center for Diabetes Research (DZD),  
Neuherberg, Germany  
e-mail: [maria.rohm@helmholtz-muenchen.de](mailto:maria.rohm@helmholtz-muenchen.de)

S. Herzig  
Institute for Diabetes and Cancer, Helmholtz Center  
Munich, Neuherberg, Germany

Joint Heidelberg-IDC Translational Diabetes  
Program, Inner Medicine 1, Heidelberg University  
Hospital, Heidelberg, Germany

German Center for Diabetes Research (DZD),  
Neuherberg, Germany

Chair Molecular Metabolic Control, Technical  
University Munich,  
Munich, Germany

## Keywords

Energy homeostasis · Glucose and lipid  
metabolism · Nuclear receptor-based thera-  
pies · Metabolic syndrome · Transcriptional  
co-factors

## 4.1 Introduction and Outline

Nuclear receptors govern multiple essential functions in metabolism. In the current chapter, we aim to introduce the most important nuclear receptor-related functions and factors in the organ-specific regulation of glucose and lipid

metabolism, as well as diseases associated with their malfunction, and novel approaches to target them. Notably, there is a multitude of additional nuclear receptors, classified either as orphan receptors including the estrogen related receptor (ERR) and the retinoic acid related receptor (ROR), or with known ligands such as the estrogen receptor (ER), the androgen receptor (AR), and the retinoic acid receptor (RAR). These receptors are also involved to some extent in the regulation of metabolism, but cannot be covered exhaustively within the scope of this chapter. Our increasing understanding of the organ- and context-specific regulation of nuclear receptors and their co-factors (Box 4.1) has already led to the development of promising therapeutics for common diseases and will likely yield novel treatment approaches for metabolic diseases in the future. We here discuss the roles of the most prominent nuclear receptors in metabolism, PPARs, LXR, FXR, and GR, in the major metabolic organs and summarize the current state of play as regards therapeutic targeting of these receptors in metabolic diseases.

## 4.2 Liver

Constant food accessibility and an overly sedentary lifestyle have led to an obesity pandemic. The imbalance of energy availability and expenditure is detrimental especially for the liver, which is one of the central organs for metabolism. Consequently, conditions such as non-alcoholic fatty liver disease (NAFLD) and non-alcoholic steatohepatitis (NASH) are on the rise. To date, there are no efficient treatments for these diseases. Increasing knowledge of the diverse functions of nuclear receptors has led to growing interest in pharmacological compounds that can manipulate their activity. Tissue- or pathway-specific manipulation of nuclear receptors could represent novel treatment possibilities.

### 4.2.1 PPAR $\alpha$ Is the Key to Liver Lipid Metabolism

Peroxisome proliferator-activated receptor alpha (PPAR $\alpha$ ) is a nuclear receptor highly abundant in liver and other tissues with high rates of fatty acid oxidation such as cardiac muscle, brown adipose tissue, and kidney [1–3]. PPAR $\alpha$  was initially identified as an activator of peroxisome proliferation induced by hepatocarcinogens [4], but was since established as a master regulator of liver lipid metabolism. PPAR $\alpha$  expression levels were found to be reduced in NAFLD patients, but increased in parallel with NAFLD histological improvements secondary to lifestyle intervention or bariatric surgery [5]. In line with this, hepatocyte specific disruption of PPAR $\alpha$  in mice resulted in steatosis and steatohepatitis indicating an essential role in lipid utilization [6]. Indeed, PPAR $\alpha$  expression is increased during suckling [7] and fasting [8], both states in which fat serves as the primary energy source. During fasting, fatty acids released from adipose tissue serve as endogenous ligands for PPAR $\alpha$  and promote the activation of the majority of pathways involved in lipid catabolism, including lipid uptake, intracellular lipid trafficking, peroxisomal and mitochondrial  $\beta$ -oxidation, and ketone body synthesis. Cellular lipid uptake, which is the first step in lipid catabolism, is facilitated by the fatty acid transporters fatty acid transporter (FAT/CD36) and fatty acid transporter protein (FATP). Both are direct target genes of PPAR $\alpha$  [9] highlighting the importance of PPAR $\alpha$  function not only for hepatic cellular metabolism but also for fatty acid clearance from the periphery. Apart from lipid uptake, PPAR $\alpha$  regulates medium-chain acyl-coenzyme A (CoA) dehydrogenase [10, 11] and acyl-CoA oxidase 1 [11] which are the rate limiting enzymes of mitochondrial and peroxisomal fatty acid  $\beta$ -oxidation (Fig. 4.1). These miscellaneous regulatory functions of PPAR $\alpha$  in lipid metabolism sparked interest in developing compounds based on PPAR $\alpha$  target gene products to counteract abnormalities and disorders associated with the metabolic syn-


#### Box 4.1: Transcriptional Co-factors as Tissue- and Context-Specific Regulators of Nuclear Receptor Function

The transcriptional control of biological processes requires tight regulation and the ability to rapidly adapt in response to metabolic changes. NRs facilitate the transcription of target genes ligand dependently, however also rely on the recruitment of co-factors. These co-factors either induce or suppress transcription and are referred to as co-activators or co-repressors, respectively. Co-factors are not exclusive to the NRs, however in general, unliganded NRs preferentially interact with co-repressors and thereby inhibit transcription while ligand binding promotes NR:co-activator interaction which facilitates transcription. Interestingly the transcriptional co-factors transducin  $\beta$ -like protein 1 (TBL1), and TBL-related 1 (TBLR1) were reported to act as so called “nuclear exchange factors”, which regulate gene repression and expression by exchanging co-repressors and co-activators [147]. Co-factors modulate NR activity by determination of cellular localization, regulation of NR stability, or posttranslational modification of the NR itself or the chromatin [148–150]. One of the better studied repressor complexes is the NR co-repressor (NCoR) and silencing mediator of retinoic acid and thyroid hormone receptors (SMRT) co-repressor complex (NCoR/SMRT complex). This repressor complex is comprised of the core subunits NCoR or SMRT as well as histone deacetylase 3 (HDAC3), G-protein pathway suppressor 2 (GPS2), TBL1, and TBLR1 [151, 152]. Repression through the NCoR/

SMRT complex is partly achieved by HDAC3 mediated deacetylation of chromatin, which disables transcription. HDAC3 is a negative regulator of white adipose tissue (WAT) browning [153], and a modulator of liver lipid metabolism [154] and pancreatic  $\beta$ -cell function [155, 156], indicating its prominent and tissue-specific function in molecular metabolic regulation. GPS2 regulates liver lipid metabolism [157] as well as adipose tissue metabolism and its endocrine function [158, 159]. Interestingly, while hepatocyte specific loss of GPS2 ameliorates NASH [157], TBL1 and TBLR1 loss of function in the liver promotes steatosis development [160], indicating that the complex core components regulate transcriptional events independently of NCoR and SMRT and/or through recruitment of additional regulatory units. In addition, TBL1 and TBLR1 were shown to directly interact with NRs and facilitate diverse tissue specific metabolic events including proliferation in pancreatic cancer cells [161, 162] and adipose tissue lipid metabolism [163]. It is currently estimated that the group of co-regulators includes around 150–400 proteins in humans [164], providing a unique tissue- and context-specific targeting opportunity in future research. As disruption of co-factor function results in various metabolic diseases, insights into the mechanistic action of these co-factors are essential. The diverse features of these co-factors and their ability to function as complexes but also independently highlight their importance for the maintenance of a normal and healthy metabolism and their potential in the development of novel therapeutic drugs.

drome. A recent prominent example which gained interest as a pharmacological target is fibroblast growth factor (FGF) 21 [12]. FGF21 administration reduced body weight, blood glucose levels, circulating plasma insulin, and hepatic gluconeogenesis in diet-induced and genetic mouse models of obesity, as well as in non-human primates [13–

16]. Additionally, while FGF21 ablation resulted in severe hepatic steatosis and inflammation [17, 18], elevation of circulating FGF21 levels reversed fatty liver and NASH [18, 19]. Due to poor pharmacokinetic properties of natural FGF21, several modified FGF21 analogues were developed, which are currently undergoing clinical trials.



**Fig. 4.1 Nuclear receptors involved in regulating liver metabolism.** Schematic representation of the main functions of nuclear receptors in the liver and their ligands, main mode of action, and key target genes. (a) Upon ligand binding, PPARα induces lipid catabolism through direct induction of genes involved in cellular lipid uptake (*FAT/CD36*, *FATP*), fatty acid β-oxidation (*ACO*, *MCAD*, *CPT1*), and ketone body synthesis. (b) Apart from cholesterol metabolism (*CYP7A1*, *ABCG5*, *ABCG8*), LXR con-

trols genes involved in carbohydrate (*ChREBP*) and triglyceride (*SREBP1c*, *APOE*, *LPL*) metabolism (c) FXR controls hepatic bile acid metabolism and is a known LXR counterplayer. (d) Glucocorticoid induced GR activity promotes gluconeogenesis (*PEPCK*, *G6Pase*) and lipid metabolism (*ANGPTL4*, *FOXO1*) in the hepatocytes. *FAT/CD36* fatty acid transporter, *FATP* fatty acid transport protein, *ACO* acyl-CoA oxidase, *MACD* medium-chain acyl-CoA dehydrogenase, *FGF21* fibroblast growth factor

Recently completed phase 2 studies assessed the efficacy and safety of FGF21 covalently conjugated to polyethylene glycol (PEGylation). PEGylated FGF21 significantly reduced hepatic fat, liver stiffness, fibrosis markers, and markers of liver damage [20, 21]. In contrast to recombinant FGF21 and other FGF21 analogues, PEGylated FGF21 did not induce bone loss or increased blood pressure in the pre-clinical setting [22–24], suggesting a low risk for chronic treatment in patients. Currently two phase 2b studies are ongoing, which evaluate the effects of PEGylated FGF21 in NASH patients with severe fibrosis ([ClinicalTrials.gov](https://clinicaltrials.gov) Identifiers: NCT03486899 and NCT03486912). Another long-acting FGF21 fusion protein (Fc-FGF21) was shown to improve glucose metabolism and plasma lipid levels across species, including humans [25, 26]. Completion of a phase 2 clinical study, which investigated the effects of Fc-FGF21 in NASH patients, reported reduction of the hepatic fat fraction and markers of liver damage, while improving glycemic control [27], indicating the potential of this compound as a novel treatment possibility for NASH and NAFLD. The current lack of approved treatments for these conditions underlines the significance of the pharmacological improvements in obesity related morbidities after FGF21 analogue administration. As a direct PPAR $\alpha$  target gene, this indicates how crucial the physiological function of PPAR $\alpha$  in the liver is, and how promising manipulations of these pathways are to develop novel treatment possibilities for the metabolic syndrome and its manifestation as NAFLD and NASH.

#### 4.2.2 LXR and FXR Are Regulators of Cholesterol Metabolism

The two liver X receptor (LXR) isoforms,  $\alpha$  and  $\beta$ , are key regulators of cholesterol, triglyceride, and carbohydrate metabolism in the liver [28–

30]. LXR $\alpha$  was initially discovered in the liver where it is highly abundant [31], whereas LXR $\beta$  is ubiquitously expressed [32]. Cholesterol, cholesterol derivatives, and cholesterol precursors were identified as natural LXR ligands indicating a central role of LXR in cholesterol metabolism [33]. Indeed, LXR is an intracellular cholesterol sensor and modulator by directly regulating genes involved in reverse cholesterol transport (RCT), conversion of cholesterol into bile acids, and intestinal excretion of cholesterol. In rodents, LXR induced cholesterol 7  $\alpha$ -hydrolase (CYP7A1) expression upon ligand binding, which is the first step and the rate-limiting enzyme for bile acid synthesis [28]. Interestingly, binding of bile acids to the farnesoid X receptor (FXR), another nuclear receptor highly abundant in the liver, downregulated CYP7A1 expression [34], identifying FXR as a LXR counterplayer. The downregulation of CYP7A1 was in part facilitated by the FXR target gene FGF15 in mice and its orthologue FGF19 in humans [35]. In addition to bile acid synthesis inhibition, FXR promoted the modification of bile acids into less toxic molecules [36] and hepatic bile acid efflux via ATP-binding cassettes ABCB11 and ABCB4 [37–39], while LXR $\alpha$  and  $\beta$  regulated cholesterol efflux from the liver into the bile via ABCG5 and ABCG8 [40] (Fig. 4.1).

Despite sharing 78% similarity in their amino acid sequence [41], LXR $\alpha$  and  $\beta$  do not possess identical functions in metabolism [42]. In mice lacking LXR $\alpha$  only, cholesterol removal from the body was severely impaired [28, 42], while LXR $\beta$  knockout (KO) mice were protected from such a phenotype [42]. Interestingly, LXRs were also identified to regulate glucose metabolism through energy utilization in brown fat [43], pancreatic insulin secretion [44], and direct up-regulation of the glucose transporter GLUT4 in adipose tissue and muscle [30, 45, 46]. Additionally, ligand activated LXR activity inhibited the gluconeogenic program through

**Fig. 4.1** (continued) 21, *CPT1* carnitine palmitoyltransferase 1, *CYP7A1* cholesterol 7  $\alpha$ -hydrolase, *ABCG5* ATP-binding cassette G 5, *SREBP1c* sterol regulatory element-binding transcription factor 1, *ChREBP* carbohydrate-responsive element-binding protein, *APOE*

apolipoprotein E, *LPL* lipoprotein lipase, *BA* bile acid, *PEPCK* phosphoenolpyruvate carboxykinase, *G6Pase* glucose-6-phosphatase, *ANGPTL4* angiopoietin-like 4, *FOXO1* forkhead box protein O1



down-regulation of peroxisome proliferator-activated receptor- $\gamma$  coactivator 1- $\alpha$  (PGC-1), phosphoenolpyruvate carboxykinase (PEPCK), and glucose-6-phosphatase (G6Pase) expression in the liver (Fig. 4.1). Suppression of gluconeogenesis was accompanied by increased glucokinase expression, which promoted glucose utilization and blood glucose clearance [30, 45, 47]. Similar to LXR, FXR also regulates glucose metabolism through fine tuning of gluconeogenesis. In the fed state FXR inhibited gluconeogenesis through repression of the key enzymes PEPCK and G6Pase [48, 49], while at fasted state FXR promoted the gluconeogenic program [49]. Accordingly, mice lacking FXR are prone to glucose intolerance and insulin resistance [50].

Due to the primarily beneficial action of LXR on glucose metabolism, LXR agonists were initially thought to be ideal therapeutic agents to treat hepatic steatosis and hyperglycemia. Indeed, LXR agonist administration suppressed the gluconeogenic program and thereby reduced blood glucose levels [45]. Some of the promising effects of LXR agonists on glucose metabolism were, however, shown to be rodent specific [51]. Additionally, synthetic LXR agonists promoted hepatic lipogenesis and steatosis, via transcriptional activation of the triglyceride master regulator sterol regulatory binding transcription factor (SREBP) 1 [29, 52]. Reduced lipogenesis was observed in LXR $\alpha$  KO mice in comparison to wild type mice [28]. However, LXR agonist administration in LXR $\alpha$  KO mice increased SREBP1 gene expression [29] indicating that both LXR $\alpha$  and LXR $\beta$  regulate lipogenesis and triglyceride synthesis in the liver. The species-specific effects as well as adverse effects of synthetic LXR agonists raised concerns as to the suitability of LXR agonists for lipid metabolism-associated disorders. In recent studies inverse LXR agonists gained interest for NASH/NAFLD treatment. Similar to agonists, inverse agonists bind to the same receptor, however they exert the opposite effect on the target cells. The inverse LXR agonist 10rr was found to inhibit lipogenesis by downregulating the expression of SREBP1, acetyl-CoA carboxylase (ACC), fatty acid synthase (FAS) and stearoyl-CoA desaturase-1 (SCD-1) [53]. Interestingly, the liver specific inverse LXR agonist

SR9238 exerts antifibrotic and anti-inflammatory effects on NASH [54, 55] and suppresses hepatic steatosis [56]. Surprisingly, in contrast to what was seen in LXR KO mice, reduced plasma cholesterol levels were observed, partly through downregulation of HMG CoA reductase (HMGCR) [56], a key regulator of cholesterol synthesis. So far, no studies were performed testing inverse LXR agonists in humans. However, administration or combination of such inverse agonists could provide novel implications and treatment possibilities in fatty liver associated diseases. In contrast, several FXR agonists already underwent clinical studies especially for the treatment of NASH. The FXR agonist obeticholic acid (OCA) reduced bile acid synthesis by repression of CYP7A1 gene expression. Moreover, OCA protected from NASH induced apoptosis of hepatocytes through suppression of p53 [57]. In NASH patients, treatment with OCA improved liver histology, however induced pruritus (itching) [58], which was also observed in treatment with other FXR agonists [59]. Currently a long term phase 3 study (NCT02548351) is ongoing to evaluate the effects of OCA on mortality, liver-related clinical outcomes, and long-term safety [60]. Cilofexor, another FXR agonist, was reported to improve hypertension and liver fibrosis in rats [61]. Additionally, in a phase 2 clinical trial NASH patients receiving cilofexor showed improved hepatic steatosis and liver transaminases [59]. In conclusion, FXR agonists represent novel and attractive candidates for NASH treatment. The current ongoing studies to determine safety and efficiency highlight the potential of such FXR agonists.

#### 4.2.3 GR – Linking Inflammation and Metabolism

The glucocorticoid receptor (GR) is a hormone-dependent nuclear receptor which regulates a wide range of metabolic processes including inflammation, lipid and glucose metabolism. More than 50 genes are under direct GR control in the liver alone [62]. GR directly regulates the expression of the gluconeogenic key enzymes PEPCK and G6Pase, suggesting an essential role

of GR in gluconeogenesis. Thus, suppression of GR activity in the liver improved hyperglycemia and dyslipidemia in genetic and inducible models of diabetes through down-regulation of PEPCK and G6Pase [63–65] (Fig. 4.1). Apart from gluconeogenesis, GR was shown to control lipid metabolism by regulating enzymes involved in lipogenesis, triglyceride (TG) uptake and fatty acid  $\beta$ -oxidation, resulting in hepatic lipid accumulation upon GR activity [66, 67]. Conversely, upon liver specific GR dysfunction, hepatic steatosis in db/db mice was ameliorated mainly through the induction of hairy enhancer of split 1 (HES1) gene expression [67]. Additionally, hepatocyte-specific GR KO impaired systemic bile acid distribution, hepatic bile acid uptake, and increased the susceptibility to develop cholesterol gallstones [68]. This indicates the importance of functional and balanced GR signaling in the liver. Previous studies identified regulatory factors such as microRNAs, transcription factors, or co-factors that directly interact with GR and thereby fine tune and balance its actions in the liver and other tissues [69–72]. Pharmacological activation of GR through cortisone or dexamethasone has immune-suppressive properties and is therefore commonly used in inflammatory or auto-immune diseases [73], however the use is overshadowed due to its severe metabolic side effects. Chronic glucocorticoid (GC) administration and the induced GR activity can result in metabolic abnormalities including hyperglycemia, insulin resistance, hepatic dyslipidemia, and hypertension [74]. Despite these severe negative effects that come with chronic GR activation, GCs are still widely prescribed. Interestingly, it was previously shown that due to alternative splicing, GR is expressed as two isoforms, GR $\alpha$  and GR $\beta$  [75]. Both isoforms are ubiquitously expressed in most tissues, with GR $\beta$  to a lesser extent than GR $\alpha$ . Additionally, in contrast to GR $\alpha$ , GR $\beta$  lacks a binding pocket for GCs [75, 76]. GR $\beta$  was initially shown to act as a GR $\alpha$  antagonist by binding to glucocorticoid responsive elements without inducing gene expression and also through heterodimerization with GR $\alpha$  [75, 77–79]. However, recent studies suggest that GR $\beta$  has, apart from the GR $\alpha$  antagonizing property, distinct regula-

tory functions on gene expression. Animal studies revealed that GR $\beta$  is the main regulator of hepatic gluconeogenesis and lipid storage. GR $\beta$  gene expression was elevated in the liver upon diet induced obesity [80]. Moreover, upon liver specific GR $\beta$  overexpression, hepatic and serum TG levels were significantly elevated. Additionally, GR $\beta$  overexpression resulted in hyperglycemia without alterations in circulating insulin levels suggesting increased gluconeogenesis or reduced hepatic insulin signaling [80]. Interestingly, short term GC administration induced lipolysis in adipose tissue through transcription of hormone-sensitive lipase (HSL) and adipose triglyceride lipase (ATGL) [81]. An explanation for this observation might be that short-term exposure to GC specifically activates GR $\alpha$ , which facilitates lipolysis [81]. GR $\alpha$  action however is antagonized by GR $\beta$  upon chronic GC administration [75, 77, 78] which in turn induces gluconeogenesis and lipogenesis [80], notably through interaction with PPAR $\alpha$  pathways, finally resulting in metabolic disruptions such as hepatic dyslipidemia and hyperglycemia. Given the antagonizing effect of GR $\beta$  on GR $\alpha$  and the direct regulation of hepatic glucose and lipid metabolism by GR $\beta$ , differentiation between the isoforms is essential in future studies. Specific regulation of either of the GR isoforms thus might provide novel treatment possibilities or reduce the severity of GC-associated side-effects.

---

### 4.3 Adipose Tissue


The white adipose tissue (WAT) is the central organ for energy storage. Excess energy is converted into TG and stored in lipid droplets in adipocytes. In nutrient deprived states, fatty acids are released and serve as energy source for other tissues such as liver or skeletal muscle. Additionally, adipose tissue was identified as an endocrine organ and releases a wide range of adipokines which regulate immune responses, control blood pressure, or modify glucose homeostasis. Modern lifestyle but also genetic predisposition account for excessive fat storage in WAT, which is accompanied by hyperglycemia

and dyslipidemia. In contrast, brown adipose tissue (BAT) is specialized to dissipate chemical energy in the form of non-shivering thermogenesis. The ability of white adipocytes to convert into brown-like or beige adipocytes, named browning and partly regulated by nuclear receptors (see also Box 4.1), represents an intensively studied field to target lifestyle induced obesity.

#### 4.3.1 PPAR $\gamma$ Is the Master Regulator of Adipose Tissue Function

The master regulator of adipose tissue function is PPAR $\gamma$ . Its expression is rapidly induced during adipogenesis [82], regulating hundreds of genes central to adipocyte function, including lipid transport (FABP4), fatty acid uptake (FATP, LPL), recycling of fatty acids (PEPCK), and lipolysis (GPR81) [83–87] (Fig. 4.2). Indeed, PPAR $\gamma$  KO in pre-adipocytes completely inhibited adipocyte formation [88], proving the vital importance of this nuclear receptor for adipogenesis. In patients, a mutation in the ligand-binding

domain of PPAR $\gamma$  led to partial lipodystrophy and insulin resistance [89], and mice lacking PPAR $\gamma$  in the adipose tissue displayed the same phenotype [90]. Aside from the effects on adipogenesis, PPAR $\gamma$  is also important for insulin sensitivity as it regulates adiponectin and resistin expression [91, 92]. Synthetic PPAR $\gamma$  agonists, in particular thiazolidinediones, ameliorate insulin resistance and are widely used in type 2 diabetes mellitus (T2DM) treatment [93]. Fat accumulation in insulin sensitive tissues such as liver and skeletal muscle has been shown to promote insulin resistance. PPAR $\gamma$  agonist-induced upregulation of genes involved in fatty acid uptake and storage in the adipose tissue promotes redistribution, and could thereby prevent ectopic fat accumulation in liver or skeletal muscle. Weight gain, liver damage, and cardiovascular events are, however, common side-effects upon chronic PPAR $\gamma$  activation [94, 95]. Interestingly, in vitro exposure of white adipocytes and in vivo exposure of mice to synthetic PPAR $\gamma$  agonist induced expression of brown fat marker genes including uncoupling protein 1 (UCP1), PR



**Fig. 4.2 Regulation of adipocyte function by PPAR $\gamma$ .** Upon ligand binding PPAR $\gamma$  induces genes involved in adipogenesis, lipid transport (FABP4), fatty acid uptake (FAT/CD36, FATP), lipid recycling (PEPCK, GLUT4), and lipolysis (LPL). *PUFA* polyunsaturated fatty acid, 12-

*HETE* 12-hydroxyeicosatetraenoic acid, *FAT/CD36* fatty acid transporter, *FATP* fatty acid transport protein, *GLUT4* glucose transporter type 4, *LPL* lipoprotein lipase, *PEPCK* phosphoenolpyruvate carboxykinase, *ACS* acetyl-CoA synthetase, *FABP4* fatty acid-binding protein 4



domain containing 16 (PRDM16), and Cell Death-Inducing DFFA-Like Effector A (CIDEA) [96–98]. The potential of increased energy expenditure by promoting browning in obesity treatment has been previously reviewed [99]. However, although PPAR $\gamma$  agonist administration induced browning of white adipocytes, PPAR $\gamma$  overexpression had no such effect [93]. Previous studies have shown that administration of dual agonists, compounds activating two targets simultaneously, surpass effects that are reached by conventional agonists [100]. Interestingly, *in vivo* simultaneous activation of PPAR $\alpha$  and PPAR $\gamma$  through dual agonists synergistically induced browning of white adipocytes [101]. Moreover, combinatorial PPAR $\alpha$  and  $\gamma$  activation reduced body weight and ameliorated insulin resistance in diet induced obesity, mainly through FGF21 signaling. Although PPAR $\alpha$  is the key regulator of FGF21 expression [12], its effects on browning rely on pharmacological PPAR $\gamma$  activation [101].

Not only white adipocyte metabolism and the white-to-brown transition of adipocytes are regulated by PPAR $\gamma$ , but also BAT relies on normal PPAR $\gamma$  function. In BAT, PPAR $\gamma$  is essential for adipogenesis, adipocyte differentiation, survival, and functionality [90, 102–106]. Similar to WAT, PPAR $\gamma$  ablation in BAT inhibited adipocyte formation [90, 102]. In addition, mature brown adipocytes lost their ability to induce non-shivering thermogenesis as PPAR $\gamma$  directly regulated the key thermogenic proteins UCP1 and PRDM16 [104, 107]. Accordingly, chronic treatment of mice with the PPAR $\gamma$  agonist rosiglitazone increased UCP1 levels in brown adipocytes, and thereby thermogenesis [103]. Interestingly, PGC1 $\alpha$ , a PPAR $\gamma$  target gene itself [108], directly interacted with PPAR $\gamma$  to enhance UCP1 gene expression [109]. Conversely, mice lacking PGC1 $\alpha$  failed to induce thermogenesis in response to cold exposure [110]. The interaction between PPAR $\gamma$  and PGC1 $\alpha$  is highly tissue- and target gene-specific. For example, PGC1 $\alpha$  was differentially expressed between BAT and WAT [109], suggesting a BAT-specific function. Moreover, PPAR $\gamma$ -controlled FABP4 expression was PGC1 $\alpha$  independent [109], indicating that PGC1 $\alpha$  selec-

tively facilitated PPAR $\gamma$  mediated thermogenesis in BAT. Apart from UCP1 expression, rosiglitazone administration upregulated triacylglyceride (TAG) synthesis [103], underlining the importance of PPAR $\gamma$  in BAT, as it regulates expression of key proteins but also the formation of substrates for non-shivering thermogenesis.

Taken together, PPAR $\gamma$  regulates many aspects of white and brown adipose tissue metabolism, which reveals PPAR $\gamma$  as highly promising target for metabolism associated abnormalities. Browning of adipocytes especially, but also increased combusting of energy through BAT, represent two intriguing possibilities to counteract obesity and its related morbidities. To date, the investigation of novel drugs for the treatment of obesity in humans by the induction of browning of white adipocytes has proven difficult.

---

## 4.4 Muscle


Skeletal muscle is the largest metabolically active organ in the human body. It is the major site of insulin dependent glucose uptake, glycogen storage, and fatty acid oxidation. Under metabolic disorders such as obesity and diabetes, severe changes occur in the skeletal muscle amongst which the switch from type 1 slow-twitch to type 2 fast-twitch fibers has profound consequences. The transition of the fiber types promotes insulin resistance, further driving the vicious cycle of the metabolic syndrome. Increase of the favorable type 1 slow-twitch muscles will ameliorate the metabolic syndrome by retrieving insulin sensitivity.

### 4.4.1 PPAR $\beta/\delta$ – Regulator of Skeletal Muscle

PPAR $\beta/\delta$  is the key transcription factor in skeletal muscle function and metabolism (Fig. 4.3). In the skeletal muscle PPAR $\beta/\delta$  regulates glucose and fatty acid metabolism [111, 112], myogenesis [113] and the transition from fast glycolytic 2b to slow/fast oxidative 1/2a fibers [114–116]. In mice and humans, PPAR $\beta/\delta$  expression is

**Fig. 4.3 PPAR $\beta/\delta$  and its main function in muscle metabolism.**

In the muscle, PPAR $\beta/\delta$  regulates genes involved in glucose and fatty acid metabolism. Moreover, ligand induced PPAR $\beta/\delta$  activation favours the muscle fiber transition from fast glycolytic to slow/fast oxidative. *PUFA* polyunsaturated fatty acid, *FA* fatty acid, *PDK4* pyruvate dehydrogenase kinase 4, *PGC1 $\alpha$*  peroxisome proliferator-activated receptor  $\gamma$  coactivator 1- $\alpha$



higher in slow/oxidative muscle types in comparison to fast/glycolytic muscles [117, 118] indicating a significant role in muscle type transition. Indeed, during endurance training, which promotes slow/fast oxidative fiber formation, PPAR $\beta/\delta$  expression was elevated [119]. The increase in the number of slow/oxidative muscle fibers upon PPAR $\beta/\delta$  induction resulted from increased muscle progenitor cell abundance, partly through antagonism of myostatin activity, a potent myokine inhibiting muscle growth [113, 120, 121]. Moreover, induction of overexpression of PPAR $\beta/\delta$  improved wheel-running performance, favored the number of slow/fast oxidative 1/2a fibers, and decreased body fat mass in rodents, partly regulated by the PPAR $\beta/\delta$  target gene PGC1 $\alpha$  [114, 116, 119]. Interestingly, as observed above PGC1 $\alpha$  was also identified as a PPAR $\beta/\delta$  coactivator [109], leading to a feed-forward loop which ensures constant PGC1 $\alpha$  expression and thereby maintenance of slow/oxidative fibers [122, 123]. Accordingly, muscle specific overexpression of PGC1 $\alpha$  phenocopies PPAR $\beta/\delta$  overexpression suggesting that both PGC1 $\alpha$  and PPAR $\beta/\delta$  facilitate skeletal muscle metabolism and function [123]. Moreover, PPAR $\beta/\delta$  modulates fatty acid metabolism through direct transcriptional control of enzymes involved in lipolysis, lipid uptake, and fatty acid

$\beta$ -oxidation [111, 115]. Regulation of  $\beta$ -oxidation in the muscle by PPAR $\beta/\delta$  is facilitated by its direct target gene pyruvate dehydrogenase kinase 4 (PDK4) [111]. PDK4 inactivates the pyruvate dehydrogenase complex (PDH), which is rate limiting for carbohydrate metabolism, leading to the up-regulation of fatty acid  $\beta$ -oxidation [124]. Accordingly, mice lacking muscle-specific PPAR $\beta/\delta$  suffer from dyslipidemia [115]. In diet-induced and genetic mouse models of obesity, PPAR $\beta/\delta$  agonist administration increased fatty acid  $\beta$ -oxidation and thereby improved dyslipidemia [111]. Moreover, PPAR $\beta/\delta$  agonist treatment improved insulin resistance, elevated proliferation of mitochondria, and reduced lipid droplets in skeletal muscle [111, 116], highlighting the therapeutic potential of PPAR $\beta/\delta$  agonists in the metabolic syndrome. Although agonist-activated PPAR $\beta/\delta$  was shown to oppose T2DM and obesity progression, and mimicked endurance training, none of the PPAR $\beta/\delta$  agonists has reached human application yet. This is largely because, apart from the overall positive effects on skeletal muscle metabolism, PPAR $\beta/\delta$  agonist administration was linked to liver fibrosis and hepatic carcinoma [125, 126]. In a recent study, novel and highly muscle-specific PPAR $\beta/\delta$  agonists were synthesized [127], yet remain to be tested for efficiency and side effects. The devel-

opment of tissue-specific PPAR $\beta/\delta$  agonists is crucial in order to bypass the severe side effects in other tissues which limit their potential to improve skeletal muscle function.

## 4.5 Pancreas


Pancreatic  $\beta$ -cells are the main regulators of glucose homeostasis as they secrete insulin in response to glucose, enabling glucose uptake into peripheral tissues. Chronic elevation of blood glucose levels results in  $\beta$ -cell hypertrophy, exhaustion, and dysfunction. Hallmarks of  $\beta$ -cell dysfunction are loss of identity, apoptosis, and insufficient insulin secretion, promoting the progression of T2DM. Counteracting  $\beta$ -cell dysfunction by nuclear receptor manipulation might prevent progression of diabetes.

### 4.5.1 The NR4A Family of Orphan Nuclear Receptors as Regulators of $\beta$ -Cell Physiology

The NR4A family of orphan nuclear receptors comprises of three members: nuclear receptor 4A1 (NR4A1), NR4A2, and NR4A3. All three were identified as important regulators of apoptosis, inflammation, and metabolism. While no function in  $\beta$ -cells for NR4A2 was reported thus far, the NR4A members NR4A1 and NR4A3 have gained substantial interest in pancreatic  $\beta$ -cell research and their roles are summarized in Fig. 4.4. NR4A1 expression was induced by glucose and fatty acids in  $\beta$ -cells [128, 129], indicating a significant role of NR4A1 in  $\beta$ -cell function and metabolism. Indeed, the NR4A1 promoter was hypomethylated in pancreatic islets from T2DM patients and mouse models of T2DM and in turn, induction of NR4A1 expression decreased blood glucose levels [130]. Moreover, NR4A1 deletion in insulin-secreting INS1 832/13 cells inhibited glucose stimulated insulin secretion through impaired mitochondrial respiration and tricarboxylic acid cycle [131, 132]. Additionally, NR4A1 was characterized as a direct NK homeobox 6.1

(NKX6.1) target and thereby induced  $\beta$ -cell proliferation in rat pancreatic islets through up-regulation of cell cycle activating genes [133]. Surprisingly, in MIN6 cells – a murine insulinoma  $\beta$ -cell line capable of insulin secretion in response to glucose stimulation [134] – fatty acid induced NR4A1 expression impaired insulin biosynthesis and insulin secretion through direct protein-protein interaction with forkhead box protein O1 (FOXO1) and down-regulation of pancreatic and duodenal homeobox 1 (PDX-1), MAF BZIP transcription factor A (MAFA), and neurogenic differentiation 1 (NEUROD1), essential transcription factors regulating  $\beta$ -cell identity and function [135]. Apart from insulin secretion, NR4A1 directly regulates endoplasmic reticulum (ER) stress induced apoptosis. ER stress, a result of sustained hyperglycemia and dyslipidemia, is a driver for T2DM progression by inducing apoptosis in pancreatic  $\beta$ -cells [136]. Interestingly, NR4A1 expression positively correlated with the induction of ER stress in vitro and ex vivo, while overexpression of NR4A1 ameliorated ER stress induced apoptosis [136, 137]. NR4A1 was identified as mitogen-activated protein kinase phosphatase 7 (MKP7) transcription factor, which counteracted c-Jun N-terminal kinase (JNK) activity and thereby apoptosis by dephosphorylation of JNK [138]. The lack of suitable and specific NR4A1 ligands has hampered detailed research on NR4A1 function in  $\beta$ -cells. Although cytosporone B (Csn-B) was identified as one of the first naturally occurring agonists for NR4A1, its effects on  $\beta$ -cells remain to be investigated [139]. Systemically, Csn-B administration resulted in increased blood glucose levels partly induced by upregulation of gluconeogenic genes in the liver. Additionally, Csn-B induced apoptosis in tumor cells to inhibit xenograft tumor growth [139], highlighting its promising properties to treat hypoglycemia and cancer.

Similar to NR4A1, findings on NR4A3 function in  $\beta$ -cells are contradictory. Initially, NR4A3 was proposed as novel candidate gene for  $\beta$ -cell function, as common genetic variations within the NR4A3 locus were associated with improved insulin secretion [140]. Glucose, fatty acids, and pro-inflammatory cytokines promote NR4A3



**Fig. 4.4 The orphan nuclear receptors NR4A1 and NR4A3 and their function in  $\beta$ -cells.** Ligands, main mode of action, and key target genes of NR4A1 and NR4A3 in the pancreatic  $\beta$ -cells. For both nuclear receptors, studies are contradictory. (a) NR4A1 expression promotes  $\beta$ -cell proliferation and ameliorates ER stress induced apoptosis through MKP7. However, NR4A1


expression was also associated with impaired  $\beta$ -cell functionality. (b) NR4A3 was described as positive regulator of  $\beta$ -cell proliferation. However, NR4A3 expression was linked to impaired  $\beta$ -cell function. The underlying conflicting results might be explained by the different models used in the respective studies. FA fatty acid, MKP7 Mitogen-activated protein kinase phosphatase 7

gene expression [141, 142]. Accordingly, NR4A3 expression was elevated in human islets from T2DM patients in comparison to healthy controls, while global NR4A3 deletion in mice resulted in increased  $\beta$ -cell mass, enhanced  $\beta$ -cell proliferation, and improved glucose tolerance [141]. Additionally, in vitro NR4A3 overexpression negatively correlated with insulin gene expression and secretion [142], suggesting NR4A3 activity impaired  $\beta$ -cell function (Fig. 4.4). Unexpectedly, NKX6.1 was also characterized as a direct regulator of NR4A3 expression, and NR4A3 overexpression promoted  $\beta$ -cell proliferation [133]. Various compounds directly regulate NR4A3 activity. 6-Mercaptopurine was identified as specific

NR4A3 agonist in skeletal muscle [143, 144]. Moreover, NR4A3 expression was induced by  $\beta$ -adrenergic receptor agonists, indicating a role in lipid metabolism [145]. Further, exendin-4 was shown to attenuate NR4A3 expression in vascular smooth muscle cells [146]. However, the effects and functionality on  $\beta$ -cells remain to be shown.

In summary, current data support a direct regulatory function of the orphan nuclear receptors NR4A1 and NR4A3 in  $\beta$ -cell functionality and metabolism. The conflicting results on NR4A1 and NR4A3 regulated insulin gene expression and secretion may be explained by the different models used in the respective studies. The conduction of in vivo studies with  $\beta$ -cell specific





**Fig. 4.5 Potential treatment strategies in metabolic diseases.** Exemplary treatment strategies for metabolic diseases by pharmacologically targeting nuclear receptors. (a) NAFLD & NASH treatment: Recombinant FGF21 proteins downregulate hepatic gluconeogenesis and promote glucose clearance and glycolysis. Inverse LXR agonists reduce lipogenesis and reduce plasma cholesterol levels though inhibition of HMG CoA reductase. FXR agonists reduce bile acid synthesis and thereby protect from NASH induced apoptosis in hepatocytes.

Specific GRα manipulation would avoid glucocorticoid induced side effects, which are likely induced by GRβ. (b) T2DM and obesity progression: PPARβ/δ activity alters fiber types and increases energy expenditure. A dual PPARα and PPARγ agonist induces browning of white adipocytes and thereby increases energy expenditure. *FGF21* fibroblast growth factor 21, *SREBP* sterol regulatory element-binding transcription factor 1, *ACC* acetyl-CoA carboxylase, *FAS* fatty acid synthase, *SCD1* stearoyl-CoA desaturase

NR4A manipulation will help to further understand the function of NR4A1 and NR4A3 in β-cells. Additionally, the continuous search for novel specific agonists is essential for clinical applications such as in the treatment of β-cell dysfunction in T2DM.

## 4.6 Conclusion

Nuclear receptors are key regulators of metabolism and their function is indispensable for metabolic health. Together with co-factors and other co-regulators, they govern a wide range of tissue- and context-specific functions influencing lipid and glucose metabolism. As a result,

manipulation and therapeutic targeting of nuclear receptor function has been intensively studied and continues to produce novel and promising drug candidates for metabolic diseases including T2DM, NAFLD, and NASH (Fig. 4.5).

**Acknowledgements** We thank Dr. Luke Harrison for figure design and editorial support. Figures were created with Biorender.com.

**Funding** A.W. and M.R. are funded through Helmholtz Association – Initiative and Networking Fund. M.R. is funded by the European Association for the Study of Diabetes (EASD), and the Helmholtz Future Topic Aging and Metabolic Programming (AMPro, ZT-0026). S.H. receives funding by the Deutsche Forschungsgemeinschaft (DFG; SFB 1123 and SFB1118) and the Else Kröner-Fresenius Foundation.



## References

1. Kliewer SA, Forman BM, Blumberg B, Ong ES, Borgmeyer U, Mangelsdorf DJ, Umesono K, Evans RM (1994) Differential expression and activation of a family of murine peroxisome proliferator-activated receptors. *Proc Natl Acad Sci U S A* 91:7355–7359. <https://doi.org/10.1073/pnas.91.15.7355>
2. Braissant O, Foulfelle F, Scotto C, Dauça M, Wahli W (1996) Differential expression of peroxisome proliferator-activated receptors (PPARs): tissue distribution of PPAR- $\alpha$ , - $\beta$ , and - $\gamma$  in the adult rat. *Endocrinology* 137:354–366. <https://doi.org/10.1210/endo.137.1.8536636>
3. Escher P, Braissant O, Basu-Modak S, Michalik L, Wahli W, Desvergne B (2001) Rat PPARs: quantitative analysis in adult rat tissues and regulation in fasting and refeeding. *Endocrinology* 142:4195–4202
4. Issemann I, Green S (1990) Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators. *Nature* 347:645–650. <https://doi.org/10.1038/347645a0>
5. Francque S, an Verrijken SC, Prawitt J, Paumelle R, Derudas B, Lefebvre P, Taskinen M-R, van Hul W, Mertens I, Hubens G, van Marck E, Michielsens P, van Gaal L, Staels B (2015) PPAR $\alpha$  gene expression correlates with severity and histological treatment response in patients with non-alcoholic steatohepatitis. *J Hepatol* 63:164–173. <https://doi.org/10.1016/j.jhep.2015.02.019>
6. Montagner A, Polizzi A, Fouché E, Ducheix S, Lippi Y, Lasserre F, Barquissau V, Régnier M, Lukowicz C, Benhamed F, Iroz A, Bertrand-Michel J, Al Saati T, Cano P, Mselli-Lakhal L, Mithieux G, Rajas F, Lagarrigue S, Pineau T, Loiseau N, Postic C, Langin D, Wahli W, Guillou H (2016) Liver PPAR $\alpha$  is crucial for whole-body fatty acid homeostasis and is protective against NAFLD. *Gut* 65:1202–1214. <https://doi.org/10.1136/gutjnl-2015-310798>
7. Panadero M, Herrera E, Bocos C (2005) Different sensitivity of PPAR $\alpha$  gene expression to nutritional changes in liver of suckling and adult rats. *Life Sci* 76:1061–1072. <https://doi.org/10.1016/j.lfs.2004.10.018>
8. Kersten S, Seydoux J, Peters JM, Gonzalez FJ, Desvergne B, Wahli W (1999) Peroxisome proliferator-activated receptor  $\alpha$  mediates the adaptive response to fasting. *J Clin Invest* 103:1489–1498. <https://doi.org/10.1172/JCI6223>
9. Motojima K, Passilly P, Peters JM, Gonzalez FJ, Latruffe N (1998) Expression of putative fatty acid transporter genes are regulated by peroxisome proliferator-activated receptor  $\alpha$  and  $\gamma$  activators in a tissue- and inducer-specific manner. *J Biol Chem* 273:16710–16714. <https://doi.org/10.1074/jbc.273.27.16710>
10. Gulick T, Cresci S, Caira T, Moore DD, Kelly DP (1994) The peroxisome proliferator-activated receptor regulates mitochondrial fatty acid oxidative enzyme gene expression. *Proc Natl Acad Sci U S A* 91:11012–11016. <https://doi.org/10.1073/pnas.91.23.11012>
11. Leone TC, Weinheimer CJ, Kelly DP (1999) A critical role for the peroxisome proliferator-activated receptor  $\alpha$  (PPAR $\alpha$ ) in the cellular fasting response: the PPAR $\alpha$ -null mouse as a model of fatty acid oxidation disorders. *Proc Natl Acad Sci U S A* 96:7473–7478. <https://doi.org/10.1073/pnas.96.13.7473>
12. Lundåsen T, Mary CH, Lisa-Mari N, Sabyasachi S, Bo A, Stefan EA, Mats R (2007) PPAR $\alpha$  is a key regulator of hepatic FGF21. *Biochem Biophys Res Commun* 360:437–440. <https://doi.org/10.1016/j.bbrc.2007.06.068>
13. Kharitonov A, Shiyanova TL, Koester A, Ford AM, Micanovic R, Galbreath EJ, Sandusky GE, Hammond LJ, Moyers JS, Owens RA, Gromada J, Brozinick JT, Hawkins ED, Wroblewski VJ, Li D-S, Mehrbod F, Jaskunas SR, Shanafelt AB (2005) FGF-21 as a novel metabolic regulator. *J Clin Invest* 115:1627–1635. <https://doi.org/10.1172/JCI23606>
14. Coskun T, Bina HA, Schneider MA, Dunbar JD, Hu CC, Chen Y, Moller DE, Kharitonov A (2008) Fibroblast growth factor 21 corrects obesity in mice. *Endocrinology* 149:6018–6027. <https://doi.org/10.1210/en.2008-0816>
15. Berglund ED, Li CY, Bina HA, Lynes SE, Michael MD, Shanafelt AB, Kharitonov A, Wasserman DH (2009) Fibroblast growth factor 21 controls glycemia via regulation of hepatic glucose flux and insulin sensitivity. *Endocrinology* 150:4084–4093. <https://doi.org/10.1210/en.2009-0221>
16. Kharitonov A, Wroblewski VJ, Koester A, Chen Y-F, Clutinger CK, Tigno XT, Hansen BC, Shanafelt AB, Etgen GJ (2007) The metabolic state of diabetic monkeys is regulated by fibroblast growth factor-21. *Endocrinology* 148:774–781
17. Tanaka N, Takahashi S, Zhang Y, Krausz KW, Smith PB, Patterson AD, Gonzalez FJ (2015) Role of fibroblast growth factor 21 in the early stage of NASH induced by methionine- and choline-deficient diet. *Biochim Biophys Acta (BBA) – Mol Basis Dis* 1852:1242–1252. <https://doi.org/10.1016/j.bbadis.2015.02.012>
18. Fisher FM, Chui PC, Nasser IA, Popov Y, Cunniff JC, Lundasen T, Kharitonov A, Schuppan D, Flier JS, Maratos-Flier E (2014) Fibroblast growth factor 21 limits lipotoxicity by promoting hepatic fatty acid activation in mice on methionine and choline-deficient diets. *Gastroenterology* 147:1073–83.e6. <https://doi.org/10.1053/j.gastro.2014.07.044>
19. Xu J, Lloyd DJ, Hale C, Stanislaus S, Chen M, Sivits G, Vonderfecht S, Hecht R, Li Y-S, Lindberg RA, Chen J-L, Jung DY, Zhang Z, Ko H-J, Kim JK, Véniant MM (2009) Fibroblast growth factor 21 reverses hepatic steatosis, increases energy expenditure, and improves insulin sensitivity in diet-induced

- obese mice. *Diabetes* 58:250–259. <https://doi.org/10.2337/db08-0392>
20. Sanyal A, Charles ED, Neuschwander-Tetri BA, Loomba R, Harrison SA, Abdelmalek MF, Lawitz EJ, Halegoua-DeMarzio D, Kundu S, Noviello S, Luo Y, Christian R (2019) Pegbelfermin (BMS-986036), a PEGylated fibroblast growth factor 21 analogue, in patients with non-alcoholic steatohepatitis: a randomised, double-blind, placebo-controlled, phase 2a trial. *Lancet* 392:2705–2717. [https://doi.org/10.1016/S0140-6736\(18\)31785-9](https://doi.org/10.1016/S0140-6736(18)31785-9)
21. Charles ED, Neuschwander-Tetri BA, Pablo Frias J, Kundu S, Luo Y, Tiruchera GS, Christian R (2019) Pegbelfermin (BMS-986036), PEGylated FGF21, in patients with obesity and type 2 diabetes: results from a randomized phase 2 study. *Obesity* (Silver Spring) 27:41–49. <https://doi.org/10.1002/oby.22344>
22. Wei W, Dutchak PA, Wang X, Ding X, Wang X, Bookout AL, Goetz R, Mohammadi M, Gerard RD, Dechow PC, Mangelsdorf DJ, Kliewer SA, Wan Y (2012) Fibroblast growth factor 21 promotes bone loss by potentiating the effects of peroxisome proliferator-activated receptor  $\gamma$ . *Proc Natl Acad Sci U S A* 109:3143–3148. <https://doi.org/10.1073/pnas.1200797109>
23. Thompson KE, Guillot M, Graziano MJ, Mangipudy RS, Chadwick KD (2021) Pegbelfermin, a PEGylated FGF21 analogue, has pharmacology without bone toxicity after 1-year dosing in skeletally-mature monkeys. *Toxicol Appl Pharmacol* 428:115673. <https://doi.org/10.1016/j.taap.2021.115673>
24. Kim AM, Somayaji VR, Dong JQ, Rolph TP, Weng Y, Chabot JR, Gropp KE, Talukdar S, Calle RA (2017) Once-weekly administration of a long-acting fibroblast growth factor 21 analogue modulates lipids, bone turnover markers, blood pressure and body weight differently in obese people with hypertriglyceridaemia and in non-human primates. *Diabetes Obes Metab* 19:1762–1772. <https://doi.org/10.1111/dom.13023>
25. Stanislaus S, Hecht R, Yie J, Hager T, Hall M, Spahr C, Wang W, Weiszmann J, Li Y, Deng L, Winters D, Smith S, Zhou L, Li Y, Véniant MM, Xu J (2017) A novel Fc-FGF21 with improved resistance to proteolysis, increased affinity toward  $\beta$ -Klotho, and enhanced efficacy in mice and cynomolgus monkeys. *Endocrinology* 158:1314–1327. <https://doi.org/10.1210/en.2016-1917>
26. Kaufman A, Abuqayyas L, Denney WS, Tillman EJ, Rolph T (2020) AKR-001, an Fc-FGF21 analog, showed sustained pharmacodynamic effects on insulin sensitivity and lipid metabolism in type 2 diabetes patients. *Cell Rep Med* 1:100057. <https://doi.org/10.1016/j.xcrim.2020.100057>
27. Harrison SA, Ruane PJ, Freilich BL, Neff G, Patil R, Behling CA, Hu C, Fong E, de Temple B, Tillman EJ, Rolph TP, Cheng A, Yale K (2021) Efruxifermin in non-alcoholic steatohepatitis: a randomized, double-blind, placebo-controlled, phase 2a trial. *Nat Med* 27:1262–1271. <https://doi.org/10.1038/s41591-021-01425-3>
28. Peet DJ, Turley SD, Ma W, Janowski BA, Lobaccaro JM, Hammer RE, Mangelsdorf DJ (1998) Cholesterol and bile acid metabolism are impaired in mice lacking the nuclear oxysterol receptor LXR  $\alpha$ . *Cell* 93:693–704. [https://doi.org/10.1016/S0092-8674\(00\)81432-4](https://doi.org/10.1016/S0092-8674(00)81432-4)
29. Repa JJ, Liang G, Ou J, Bashmakov Y, Lobaccaro JM, Shimomura I, Shan B, Brown MS, Goldstein JL, Mangelsdorf DJ (2000) Regulation of mouse sterol regulatory element-binding protein-1c gene (SREBP-1c) by oxysterol receptors, LXR $\alpha$  and LXR $\beta$ . *Genes Dev* 14:2819–2830. <https://doi.org/10.1101/gad.844900>
30. Commerford SR, Vargas L, Dorfman SE, Mitro N, Rocheford EC, Mak PA, Li X, Kennedy P, Mullarkey TL, Saez E (2007) Dissection of the insulin-sensitizing effect of liver X receptor ligands. *Mol Endocrinol* 21:3002–3012. <https://doi.org/10.1210/me.2007-0156>
31. Willy PJ, Umehono K, Ong ES, Evans RM, Heyman RA, Mangelsdorf DJ (1995) LXR, a nuclear receptor that defines a distinct retinoid response pathway. *Genes Dev* 9:1033–1045. <https://doi.org/10.1101/gad.9.9.1033>
32. Teboul M, Enmark E, Li Q, Wikström AC, Pelto-Huikko M, Gustafsson JA (1995) OR-1, a member of the nuclear receptor superfamily that interacts with the 9-cis-retinoic acid receptor. *Proc Natl Acad Sci U S A* 92:2096–2100. <https://doi.org/10.1073/pnas.92.6.2096>
33. Janowski BA, Willy PJ, Devi TR, Falck JR, Mangelsdorf DJ (1996) An oxysterol signalling pathway mediated by the nuclear receptor LXR $\alpha$ . *Nature* 383:728–731. <https://doi.org/10.1038/383728a0>
34. Chiang JY, Kimmel R, Weinberger C, Stroup D (2000) Farnesoid X receptor responds to bile acids and represses cholesterol 7 $\alpha$ hydroxylase gene (CYP7A1) transcription. *J Biol Chem* 275:10918–10924. <https://doi.org/10.1074/jbc.275.15.10918>
35. Inagaki T, Choi M, Moschetta A, Peng L, Cummins CL, McDonald JG, Luo G, Jones SA, Goodwin B, Richardson JA, Gerard RD, Repa JJ, Mangelsdorf DJ, Kliewer SA (2005) Fibroblast growth factor 15 functions as an enterohepatic signal to regulate bile acid homeostasis. *Cell Metab* 2:217–225. <https://doi.org/10.1016/j.cmet.2005.09.001>
36. Guo GL, Lambert G, Negishi M, Ward JM, Brewer HB, Kliewer SA, Gonzalez FJ, Sinal CJ (2003) Complementary roles of farnesoid X receptor, pregnane X receptor, and constitutive androstane receptor in protection against bile acid toxicity. *J Biol Chem* 278:45062–45071
37. Huang L, Zhao A, Lew J-L, Zhang T, Hrywna Y, Thompson JR, de Pedro N, Royo I, Blevins RA, Peláez F, Wright SD, Cui J (2003) Farnesoid X

- receptor activates transcription of the phospholipid pump MDR3. *J Biol Chem* 278:51085–51090. <https://doi.org/10.1074/jbc.M308321200>
38. Plass JRM, Mol O, Heegsma J, Geuken M, Faber KN, Jansen PLM, Müller M (2002) Farnesoid X receptor and bile salts are involved in transcriptional regulation of the gene encoding the human bile salt export pump. *Hepatology* 35:589–596. <https://doi.org/10.1053/jhep.2002.31724>
  39. Ananthanarayanan M, Balasubramanian N, Makishima M, Mangelsdorf DJ, Suchy FJ (2001) Human bile salt export pump promoter is transactivated by the farnesoid X receptor/bile acid receptor. *J Biol Chem* 276:28857–28865. <https://doi.org/10.1074/jbc.M011610200>
  40. Repa JJ, Berge KE, Pomajzl C, Richardson JA, Hobbs H, Mangelsdorf DJ (2002) Regulation of ATP-binding cassette sterol transporters ABCG5 and ABCG8 by the liver X receptors alpha and beta. *J Biol Chem* 277:18793–18800. <https://doi.org/10.1074/jbc.M109927200>
  41. Alberti S, Steffensen KR, Gustafsson JA (2000) Structural characterisation of the mouse nuclear oxysterol receptor genes LXRalpha and LXRbeta. *Gene* 243:93–103. [https://doi.org/10.1016/S0378-1119\(99\)00555-7](https://doi.org/10.1016/S0378-1119(99)00555-7)
  42. Alberti S, Schuster G, Parini P, Feltkamp D, Diczfalussy U, Rudling M, Angelin B, Björkhem I, Pettersson S, Gustafsson JA (2001) Hepatic cholesterol metabolism and resistance to dietary cholesterol in LXRbeta-deficient mice. *J Clin Invest* 107:565–573. <https://doi.org/10.1172/JCI9794>
  43. Korach-André M, Archer A, Barros RP, Parini P, Gustafsson J-Å (2011) Both liver-X receptor (LXR) isoforms control energy expenditure by regulating brown adipose tissue activity. *Proc Natl Acad Sci U S A* 108:403–408
  44. Ogihara T, Chuang J-C, Vestermarck GL, Garmey JC, Ketchum RJ, Huang X, Brayman KL, Thorne MO, Repa JJ, Mirmira RG, Evans-Molina C (2010) Liver X receptor agonists augment human islet function through activation of anaplerotic pathways and glycerolipid/free fatty acid cycling. *J Biol Chem* 285:5392–5404. <https://doi.org/10.1074/jbc.M109.064659>
  45. Laffitte BA, Chao LC, Li J, Walczak R, Hummasti S, Joseph SB, Castrillo A, Wilpitz DC, Mangelsdorf DJ, Collins JL, Saez E, Tontonoz P (2003) Activation of liver X receptor improves glucose tolerance through coordinate regulation of glucose metabolism in liver and adipose tissue. *Proc Natl Acad Sci U S A* 100:5419–5424. <https://doi.org/10.1073/pnas.0830671100>
  46. Baranowski M, Zabielski P, Błażnio-Zabielska AU, Harasim E, Chabowski A, Górski J (2014) Insulin-sensitizing effect of LXR agonist T0901317 in high-fat fed rats is associated with restored muscle GLUT4 expression and insulin-stimulated AS160 phosphorylation. *Cell Physiol Biochem* 33:1047–1057. <https://doi.org/10.1159/000358675>
  47. Cao G, Liang Y, Broderick CL, Oldham BA, Beyer TP, Schmidt RJ, Zhang Y, Staybrook KR, Suen C, Otto KA, Miller AR, Dai J, Foxworthy P, Gao H, Ryan TP, Jiang X-C, Burris TP, Eacho PI, Etgen GJ (2003) Antidiabetic action of a liver x receptor agonist mediated by inhibition of hepatic gluconeogenesis. *J Biol Chem* 278:1131–1136. <https://doi.org/10.1074/jbc.M210208200>
  48. Zhang Y, Lee FY, Barrera G, Lee H, Vales C, Gonzalez FJ, Willson TM, Edwards PA (2006) Activation of the nuclear receptor FXR improves hyperglycemia and hyperlipidemia in diabetic mice. *Proc Natl Acad Sci U S A* 103:1006–1011. <https://doi.org/10.1073/pnas.0506982103>
  49. Renga B, Mencarelli A, D'Amore C, Cipriani S, Baldelli F, Zampella A, Distrutti E, Fiorucci S (2012) Glucocorticoid receptor mediates the gluconeogenic activity of the farnesoid X receptor in the fasting condition. *FASEB J* 26:3021–3031. <https://doi.org/10.1096/fj.11-195701>
  50. Cariou B, van Harmelen K, Duran-Sandoval D, van Dijk TH, Grefhorst A, Abdelkarim M, Caron S, Torpier G, Fruchart J-C, Gonzalez FJ, Kuipers F, Staels B (2006) The farnesoid X receptor modulates adiposity and peripheral insulin sensitivity in mice. *J Biol Chem* 281:11039–11049. <https://doi.org/10.1074/jbc.M510258200>
  51. Goodwin B, Watson MA, Kim H, Miao J, Kemper JK, Kliewer SA (2003) Differential regulation of rat and human CYP7A1 by the nuclear oxysterol receptor liver X receptor-alpha. *Mol Endocrinol* 17:386–394. <https://doi.org/10.1210/me.2002-0246>
  52. Schultz JR, Tu H, Luk A, Repa JJ, Medina JC, Li L, Schwendner S, Wang S, Thoolen M, Mangelsdorf DJ, Lustig KD, Shan B (2000) Role of LXRs in control of lipogenesis. *Genes Dev* 14:2831–2838. <https://doi.org/10.1101/gad.850400>
  53. Chen Z, Chen H, Zhang Z, Ding P, Yan X, Li Y, Zhang S, Gu Q, Zhou H, Xu J (2020) Discovery of novel liver X receptor inverse agonists as lipogenesis inhibitors. *Eur J Med Chem* 206:112793. <https://doi.org/10.1016/j.ejmech.2020.112793>
  54. Huang P, Kaluba B, Jiang X, Chang S, Tang X, Mao L, Zhang Z, Huang F, Zhai L (2018) Liver X receptor inverse agonist SR9243 suppresses nonalcoholic steatohepatitis intrahepatic inflammation and fibrosis. *Biomed Res Int* 2018:8071093. <https://doi.org/10.1155/2018/8071093>
  55. Griffett K, Welch RD, Flaveny CA, Kolar GR, Neuschwander-Tetri BA, Burris TP (2015) The LXR inverse agonist SR9238 suppresses fibrosis in a model of non-alcoholic steatohepatitis. *Mol Metab* 4:353–357. <https://doi.org/10.1016/j.molmet.2015.01.009>
  56. Griffett K, Solt LA, El-Gendy BE-DM, Kamenecka TM, Burris TP (2013) A liver-selective LXR inverse

- agonist that suppresses hepatic steatosis. *ACS Chem Biol* 8:559–567. <https://doi.org/10.1021/cb300541g>
57. Goto T, Itoh M, Suganami T, Kanai S, Shirakawa I, Sakai T, Asakawa M, Yoneyama T, Kai T, Ogawa Y (2018) Obeticholic acid protects against hepatocyte death and liver fibrosis in a murine model of non-alcoholic steatohepatitis. *Sci Rep* 8:8157. <https://doi.org/10.1038/s41598-018-26383-8>
  58. Neuschwander-Tetri BA, Loomba R, Sanyal AJ, Lavine JE, van Natta ML, Abdelmalek MF, Chalasani N, Dasarthy S, Diehl AM, Hameed B, Kowdley KV, McCullough A, Terrault N, Clark JM, Tonascia J, Brunt EM, Kleiner DE, Doo E (2015) Farnesoid X nuclear receptor ligand obeticholic acid for non-cirrhotic, non-alcoholic steatohepatitis (FLINT): a multicentre, randomised, placebo-controlled trial. *Lancet* 385:956–965. [https://doi.org/10.1016/S0140-6736\(14\)61933-4](https://doi.org/10.1016/S0140-6736(14)61933-4)
  59. Patel K, Harrison SA, Elkhatab M, Trotter JF, Herring R, Rojter SE, Kayali Z, Wong VW-S, Greenbloom S, Jayakumar S, Shiffman ML, Freilich B, Lawitz EJ, Gane EJ, Harting E, Xu J, Billin AN, Chung C, Djedjos CS, Subramanian GM, Myers RP, Middleton MS, Rinella M, Noureddin M (2020) Cilofexor, a nonsteroidal FXR agonist, in patients with noncirrhotic NASH: a phase 2 randomized controlled trial. *Hepatology* 72:58–71. <https://doi.org/10.1002/hep.31205>
  60. Ratziu V, Sanyal AJ, Loomba R, Rinella M, Harrison S, Anstee QM, Goodman Z, Bedossa P, MacConell L, Shringarpure R, Shah A, Younossi Z (2019) REGENERATE: design of a pivotal, randomised, phase 3 study evaluating the safety and efficacy of obeticholic acid in patients with fibrosis due to nonalcoholic steatohepatitis. *Contemp Clin Trials* 84:105803. <https://doi.org/10.1016/j.cct.2019.06.017>
  61. Schwabl P, Hambruch E, Budas GR, Supper P, Burnet M, Liles JT, Birkel M, Brusilovskaya K, Königshofer P, Peck-Radosavljevic M, Watkins WJ, Trauner M, Breckenridge DG, Kremoser C, Reiberger T (2021) The non-steroidal FXR agonist cilofexor improves portal hypertension and reduces hepatic fibrosis in a rat NASH model. *Biomedicine* 9. <https://doi.org/10.3390/biomedicine9010060>
  62. Phuc Le P, Friedman JR, Schug J, Brestelli JE, Parker JB, Bochkis IM, Kaestner KH (2005) Glucocorticoid receptor-dependent gene regulatory networks. *PLoS Genet* 1:e16. <https://doi.org/10.1371/journal.pgen.0010016>
  63. Opherck C, Tronche F, Kellendonk C, Köhl Müller D, Schulze A, Schmid W, Schütz G (2004) Inactivation of the glucocorticoid receptor in hepatocytes leads to fasting hypoglycemia and ameliorates hyperglycemia in streptozotocin-induced diabetes mellitus. *Mol Endocrinol* 18:1346–1353. <https://doi.org/10.1210/me.2003-0283>
  64. Liang Y, Osborne MC, Monia BP, Bhanot S, Watts LM, She P, DeCarlo SO, Chen X, Demarest K (2005) Antisense oligonucleotides targeted against glucocorticoid receptor reduce hepatic glucose production and ameliorate hyperglycemia in diabetic mice. *Metabolism* 54:848–855. <https://doi.org/10.1016/j.metabol.2005.01.030>
  65. Watts LM, Mancham VP, Leedom TA, Rivard AL, McKay RA, Bao D, Neroladakis T, Monia BP, Bodenmiller DM, Cao JX-C, Zhang HY, Cox AL, Jacobs SJ, Michael MD, Sloop KW, Bhanot S (2005) Reduction of hepatic and adipose tissue glucocorticoid receptor expression with antisense oligonucleotides improves hyperglycemia and hyperlipidemia in diabetic rodents without causing systemic glucocorticoid antagonism. *Diabetes* 54:1846–1853. <https://doi.org/10.2337/diabetes.54.6.1846>
  66. Cole TG, Wilcox HG, Heimberg M (1982) Effects of adrenalectomy and dexamethasone on hepatic lipid metabolism. *J Lipid Res* 23:81–91. [https://doi.org/10.1016/S0022-2275\(20\)38176-1](https://doi.org/10.1016/S0022-2275(20)38176-1)
  67. Lemke U, Krones-Herzig A, Diaz MB, Narvekar P, Ziegler A, Vegiopoulos A, Cato AC, Bohl S, Klingmüller U, Sreaton RA, Müller-Decker K, Kersten S, Herzig S (2008) The glucocorticoid receptor controls hepatic dyslipidemia through Hes1. *Cell Metab* 8:212–223. <https://doi.org/10.1016/j.cmet.2008.08.001>
  68. Rose AJ, Díaz MB, Reimann A, Klement J, Walcher T, Krones-Herzig A, Strobel O, Werner J, Peters A, Kleyman A, Tuckermann JP, Vegiopoulos A, Herzig S (2011) Molecular control of systemic bile acid homeostasis by the liver glucocorticoid receptor. *Cell Metab* 14:123–130. <https://doi.org/10.1016/j.cmet.2011.04.010>
  69. de Guia RM, Rose AJ, Sommerfeld A, Seibert O, Strzoda D, Zota A, Feuchter Y, Krones-Herzig A, Sijmonsma T, Kirilov M, Sticht C, Gretz N, Dallinga-Thie G, Diederichs S, Klötting N, Blüher M, Berriel Diaz M, Herzig S (2015) microRNA-379 couples glucocorticoid hormones to dysfunctional lipid homeostasis. *EMBO J* 34:344–360. <https://doi.org/10.15252/embj.201490464>
  70. Glantschnig C, Koenen M, Gil-Lozano M, Karbiener M, Pickrahn I, Williams-Dautovich J, Patel R, Cummins CL, Giroud M, Hartleben G, Vogl E, Blüher M, Tuckermann J, Uhlenhaut H, Herzig S, Scheideler M (2019) A miR-29a-driven negative feedback loop regulates peripheral glucocorticoid receptor signaling. *FASEB J* 33:5924–5941. <https://doi.org/10.1096/fj.201801385RR>
  71. Hemmer MC, Wierer M, Schachtrup K, Downes M, Hübner N, Evans RM, Uhlenhaut NH (2019) E47 modulates hepatic glucocorticoid action. *Nat Commun* 10:306. <https://doi.org/10.1038/s41467-018-08196-5>
  72. Jones CL, Bhatla T, Blum R, Wang J, Paugh SW, Wen X, Bourgeois W, Bitterman DS, Raetz EA, Morrison



- DJ, Teachey DT, Evans WE, Garabedian MJ, Carroll WL (2014) Loss of TBL1XR1 disrupts glucocorticoid receptor recruitment to chromatin and results in glucocorticoid resistance in a B-lymphoblastic leukemia model. *J Biol Chem* 289:20502–20515. <https://doi.org/10.1074/jbc.M114.569889>
73. Greulich F, Wierer M, Mechtidou A, Gonzalez-Garcia O, Uhlenhaut NH (2021) The glucocorticoid receptor recruits the COMPASS complex to regulate inflammatory transcription at macrophage enhancers. *Cell Rep* 34:108742. <https://doi.org/10.1016/j.celrep.2021.108742>
  74. Schäcke H, Döcke WD, Asadullah K (2002) Mechanisms involved in the side effects of glucocorticoids. *Pharmacol Ther* 96:23–43. [https://doi.org/10.1016/s0163-7258\(02\)00297-8](https://doi.org/10.1016/s0163-7258(02)00297-8)
  75. Hinds TD Jr, Ramakrishnan S, Cash HA, Stechschulte LA, Heinrich G, Najjar SM, Sanchez ER (2010) Discovery of glucocorticoid receptor-beta in mice with a role in metabolism. *Mol Endocrinol* 24:1715–1727. <https://doi.org/10.1210/me.2009-0411>
  76. Oakley RH, Webster JC, Sar M, Parker CR Jr, Cidlowski JA (1997) Expression and subcellular distribution of the beta-isoform of the human glucocorticoid receptor. *Endocrinology* 138:5028–5038. <https://doi.org/10.1210/endo.138.11.5501>
  77. Bamberger GM, Bamberger AM, de Castro M, Chrousos GP (1995) Glucocorticoid receptor beta, a potential endogenous inhibitor of glucocorticoid action in humans. *J Clin Invest* 95:2435–2441. <https://doi.org/10.1172/JCI117943>
  78. Chatzopoulou A, Roy U, Meijer AH, Alia A, Spaink HP, Schaaf MJM (2015) Transcriptional and metabolic effects of glucocorticoid receptor  $\alpha$  and  $\beta$  signaling in zebrafish. *Endocrinology* 156:1757–1769. <https://doi.org/10.1210/en.2014-1941>
  79. Oakley RH, Jewell CM, Yudt MR, Bofetiado DM, Cidlowski JA (1999) The dominant negative activity of the human glucocorticoid receptor  $\beta$  isoform: specificity and mechanisms of action\*. *J Biol Chem* 274:27857–27866. <https://doi.org/10.1074/jbc.274.39.27857>
  80. Marino JS, Stechschulte LA, Stec DE, Nestor-Kalinoski A, Coleman S, Hinds TD Jr (2016) Glucocorticoid receptor  $\beta$  induces hepatic steatosis by augmenting inflammation and inhibition of the peroxisome proliferator-activated receptor (PPAR)  $\alpha$ . *J Biol Chem* 291:25776–25788. <https://doi.org/10.1074/jbc.M116.752311>
  81. Xu C, He J, Jiang H, Zu L, Zhai W, Pu S, Xu G (2009) Direct effect of glucocorticoids on lipolysis in adipocytes. *Mol Endocrinol* 23:1161–1170. <https://doi.org/10.1210/me.2008-0464>
  82. Chawla A, Schwarz EJ, Dimaculangan DD, Lazar MA (1994) Peroxisome proliferator-activated receptor (PPAR) gamma: adipose-predominant expression and induction early in adipocyte differentiation. *Endocrinology* 135:798–800. <https://doi.org/10.1210/endo.135.2.8033830>
  83. Tontonoz P, Graves RA, Budavari AI, Erdjument-Bromage H, Lui M, Hu E, Tempst P, Spiegelman BM (1994) Adipocyte-specific transcription factor ARF6 is a heterodimeric complex of two nuclear hormone receptors, PPAR gamma and RXR alpha. *Nucleic Acids Res* 22:5628–5634. <https://doi.org/10.1093/nar/22.25.5628>
  84. Frohnert BI, Hui TY, Bernlohr DA (1999) Identification of a functional peroxisome proliferator-responsive element in the murine fatty acid transport protein gene. *J Biol Chem* 274:3970–3977. <https://doi.org/10.1074/jbc.274.7.3970>
  85. Jeninga EH, Bugge A, Nielsen R, Kersten S, Hamers N, Dani C, Wabitsch M, Berger R, Stunnenberg HG, Mandrup S, Kalkhoven E (2009) Peroxisome proliferator-activated receptor gamma regulates expression of the anti-lipolytic G-protein-coupled receptor 81 (GPR81/Gpr81). *J Biol Chem* 284:26385–26393. <https://doi.org/10.1074/jbc.M109.040741>
  86. Tontonoz P, Hu E, Devine J, Beale EG, Spiegelman BM (1995) PPAR gamma 2 regulates adipose expression of the phosphoenolpyruvate carboxykinase gene. *Mol Cell Biol* 15:351–357. <https://doi.org/10.1128/MCB.15.1.351>
  87. Schoonjans K, Peinado-Onsurbe J, Lefebvre AM, Heyman RA, Briggs M, Deeb S, Staels B, Auwerx J (1996) PPARalpha and PPARgamma activators direct a distinct tissue-specific transcriptional response via a PPRE in the lipoprotein lipase gene. *EMBO J* 15:5336–5348
  88. Rosen ED, Sarraf P, Troy AE, Bradwin G, Moore K, Milstone DS, Spiegelman BM, Mortensen RM (1999) PPAR $\gamma$  is required for the differentiation of adipose tissue in vivo and in vitro. *Mol Cell* 4:611–617. [https://doi.org/10.1016/S1097-2765\(00\)80211-7](https://doi.org/10.1016/S1097-2765(00)80211-7)
  89. Savage DB, Tan GD, Acerini CL, Jebb SA, Agostini M, Gurnell M, Williams RL, Umpoleby AM, Thomas EL, Bell JD, Dixon AK, Dunne F, Boiani R, Cinti S, Vidal-Puig A, Karpe F, Chatterjee VKK, O'Rahilly S (2003) Human metabolic syndrome resulting from dominant-negative mutations in the nuclear receptor peroxisome proliferator-activated receptor-gamma. *Diabetes* 52:910–917. <https://doi.org/10.2337/diabetes.52.4.910>
  90. He W, Barak Y, Hevener A, Olson P, Liao D, Le J, Nelson M, Ong E, Olefsky JM, Evans RM (2003) Adipose-specific peroxisome proliferator-activated receptor gamma knockout causes insulin resistance in fat and liver but not in muscle. *Proc Natl Acad Sci U S A* 100:15712–15717. <https://doi.org/10.1073/pnas.2536828100>
  91. Iwaki M, Matsuda M, Maeda N, Funahashi T, Matsuzawa Y, Makishima M, Shimomura I (2003) Induction of adiponectin, a fat-derived antidiabetic



- and antiatherogenic factor, by nuclear receptors. *Diabetes* 52:1655–1663. <https://doi.org/10.2337/diabetes.52.7.1655>
92. Tomaru T, Steger DJ, Lefterova MI, Schupp M, Lazar MA (2009) Adipocyte-specific expression of murine resistin is mediated by synergism between peroxisome proliferator-activated receptor gamma and CCAAT/enhancer-binding proteins. *J Biol Chem* 284:6116–6125. <https://doi.org/10.1074/jbc.M808407200>
  93. Sugii S, Olson P, Sears DD, Saberi M, Atkins AR, Barish GD, Hong S-H, Castro GL, Yin Y-Q, Nelson MC, Hsiao G, Greaves DR, Downes M, Yu RT, Olefsky JM, Evans RM (2009) PPARgamma activation in adipocytes is sufficient for systemic insulin sensitization. *Proc Natl Acad Sci U S A* 106:22504–22509. <https://doi.org/10.1073/pnas.0912487106>
  94. Graham DJ, Ouellet-Hellstrom R, MacCurdy TE, Ali F, Sholley C, Worrall C, Kelman JA (2010) Risk of acute myocardial infarction, stroke, heart failure, and death in elderly Medicare patients treated with rosiglitazone or pioglitazone. *JAMA* 304:411–418. <https://doi.org/10.1001/jama.2010.920>
  95. Floyd JS, Barbehenn E, Lurie P, Wolfe SM (2009) Case series of liver failure associated with rosiglitazone and pioglitazone. *Pharmacoevidemiol Drug Saf* 18:1238–1243. <https://doi.org/10.1002/pds.1804>
  96. Fukui Y, Masui S, Osada S, Umesono K, Motojima K (2000) A new thiazolidinedione, NC-2100, which is a weak PPAR-gamma activator, exhibits potent antidiabetic effects and induces uncoupling protein 1 in white adipose tissue of KKAY obese mice. *Diabetes* 49:759–767. <https://doi.org/10.2337/diabetes.49.5.759>
  97. Petrovic N, Walden TB, Shabalina IG, Timmons JA, Cannon B, Nedergaard J (2010) Chronic peroxisome proliferator-activated receptor gamma (PPARgamma) activation of epididymally derived white adipocyte cultures reveals a population of thermogenically competent, UCP1-containing adipocytes molecularly distinct from classic brown adipocytes. *J Biol Chem* 285:7153–7164. <https://doi.org/10.1074/jbc.M109.053942>
  98. Ohno H, Shinoda K, Spiegelman BM, Kajimura S (2012) PPARγ agonists induce a white-to-brown fat conversion through stabilization of PRDM16 protein. *Cell Metab* 15:395–404. <https://doi.org/10.1016/j.cmet.2012.01.019>
  99. Jeremic N, Chaturvedi P, Tyagi SC (2017) Browning of white fat: novel insight into factors, mechanisms, and therapeutics. *J Cell Physiol* 232:61–68. <https://doi.org/10.1002/jcp.25450>
  100. Day JW, Ottaway N, Patterson JT, Gelfanov V, Smiley D, Gidda J, Findeisen H, Bruemmer D, Drucker DJ, Chaudhary N, Holland J, Hembree J, Abplanalp V, Grant E, Ruehl J, Wilson H, Kirchner H, Lockie SH, Hofmann S, Woods SC, Nogueiras R, Pfluger PT, Perez-Tilve D, DiMarchi R, Tschöp MH (2009) A new glucagon and GLP-1 co-agonist eliminates obesity in rodents. *Nat Chem Biol* 5:749–757. <https://doi.org/10.1038/nchembio.209>
  101. Kroon T, Harms M, Maurer S, Bonnet L, Alexandersson I, Lindblom A, Ahnmark A, Nilsson D, Gennemark P, O'Mahony G, Osinski V, McNamara C, Boucher J (2020) PPARγ and PPARα synergize to induce robust browning of white fat in vivo. *Mol Metab* 36:100964. <https://doi.org/10.1016/j.molmet.2020.02.007>
  102. Jones JR, Barrick C, Kim K-A, Lindner J, Blondeau B, Fujimoto Y, Shiota M, Kesterson RA, Kahn BB, Magnuson MA (2005) Deletion of PPARgamma in adipose tissues of mice protects against high fat diet-induced obesity and insulin resistance. *Proc Natl Acad Sci U S A* 102:6207–6212. <https://doi.org/10.1073/pnas.0306743102>
  103. Festuccia WT, Blanchard P-G, Richard D, Deshaies Y (2010) Basal adrenergic tone is required for maximal stimulation of rat brown adipose tissue UCP1 expression by chronic PPAR-gamma activation. *Am J Physiol Regul Integr Comp Phys* 299:R159–R167. <https://doi.org/10.1152/ajpregu.00821.2009>
  104. Lasar D, Rosenwald M, Kiehlmann E, Balaz M, Tall B, Opitz L, Lidell ME, Zamboni N, Krznar P, Sun W, Varga L, Stefanicka P, Ukropec J, Nuutila P, Virtanen K, Amri E-Z, Enerbäck S, Wahli W, Wolfrum C (2018) Peroxisome proliferator activated receptor gamma controls mature brown adipocyte inducibility through glycerol kinase. *Cell Rep* 22:760–773. <https://doi.org/10.1016/j.celrep.2017.12.067>
  105. Imai T, Takakuwa R, Marchand S, Dentz E, Bornert J-M, Messaddeq N, Wendling O, Mark M, Desvergne B, Wahli W, Chambon P, Metzger D (2004) Peroxisome proliferator-activated receptor gamma is required in mature white and brown adipocytes for their survival in the mouse. *Proc Natl Acad Sci U S A* 101:4543–4547. <https://doi.org/10.1073/pnas.0400356101>
  106. Tai T-AC, Jennermann C, Brown KK, Oliver BB, MacGinnitie MA, Wilkison WO, Brown HR, Lehmann JM, Kliewer SA, Morris DC, Graves RA (1996) Activation of the nuclear receptor peroxisome proliferator-activated receptor γ promotes brown adipocyte differentiation\*. *J Biol Chem* 271:29909–29914. <https://doi.org/10.1074/jbc.271.47.29909>
  107. Xiong W, Zhao X, Villacorta L, Rom O, Garcia-Barrio MT, Guo Y, Fan Y, Zhu T, Zhang J, Zeng R, Chen YE, Jiang Z, Chang L (2018) Brown adipocyte-specific PPARγ (peroxisome proliferator-activated receptor γ) deletion impairs perivascular adipose tissue development and enhances atherosclerosis in mice. *Arterioscler Thromb Vasc Biol* 38:1738–1747. <https://doi.org/10.1161/ATVBAHA.118.311367>
  108. Hondares E, Mora O, Yubero P, de la Concepción R, Marisa R, Iglesias M, Giralt FV (2006) Thiazolidinediones and retinoids induce peroxisome proliferator-activated receptor-coactivator (PGC)-

- 1alpha gene transcription: an autoregulatory loop controls PGC-1alpha expression in adipocytes via peroxisome proliferator-activated receptor-gamma coactivation. *Endocrinology* 147:2829–2838. <https://doi.org/10.1210/en.2006-0070>
109. Puigserver P, Wu Z, Park CW, Graves R, Wright M, Spiegelman BM (1998) A cold-inducible coactivator of nuclear receptors linked to adaptive thermogenesis. *Cell* 92:829–839. [https://doi.org/10.1016/S0092-8674\(00\)81410-5](https://doi.org/10.1016/S0092-8674(00)81410-5)
  110. Lin J, Wu P-H, Tarr PT, Lindenberg KS, St-Pierre J, Zhang C-Y, Mootha VK, Jäger S, Vianna CR, Reznick RM, Cui L, Manieri M, Donovan MX, Wu Z, Cooper MP, Fan MC, Rohas LM, Zavacki AM, Cinti S, Shulman GI, Lowell BB, Krainc D, Spiegelman BM (2004) Defects in adaptive energy metabolism with CNS-linked hyperactivity in PGC-1 $\alpha$  null mice. *Cell* 119:121–135. <https://doi.org/10.1016/j.cell.2004.09.013>
  111. Tanaka T, Yamamoto J, Iwasaki S, Asaba H, Hamura H, Ikeda Y, Watanabe M, Magoori K, Ioka RX, Tachibana K, Watanabe Y, Uchiyama Y, Sumi K, Iguchi H, Ito S, Doi T, Hamakubo T, Naito M, Auwerx J, Yanagisawa M, Kodama T, Sakai J (2003) Activation of peroxisome proliferator-activated receptor delta induces fatty acid beta-oxidation in skeletal muscle and attenuates metabolic syndrome. *Proc Natl Acad Sci U S A* 100:15924–15929. <https://doi.org/10.1073/pnas.0306981100>
  112. Nahlé Z, Hsieh M, Pietka T, Coburn CT, Grimaldi PA, Zhang MQ, Das D, Abumrad NA (2008) CD36-dependent regulation of muscle FoxO1 and PDK4 in the PPAR delta/beta-mediated adaptation to metabolic stress. *J Biol Chem* 283:14317–14326. <https://doi.org/10.1074/jbc.M706478200>
  113. Angione AR, Jiang C, Pan D, Wang Y-X, Kuang S (2011) PPAR $\delta$  regulates satellite cell proliferation and skeletal muscle regeneration. *Skelet Muscle* 1:33. <https://doi.org/10.1186/2044-5040-1-33>
  114. Chen W, Gao R, Xie X, Zheng Z, Li H, Li S, Dong F, Wang L (2015) A metabolomic study of the PPAR $\delta$  agonist GW501516 for enhancing running endurance in Kunming mice. *Sci Rep* 5:9884. <https://doi.org/10.1038/srep09884>
  115. Schuler M, Ali F, Chambon C, Duteil D, Bornert J-M, Tardivel A, Desvergne B, Wahli W, Chambon P, Metzger D (2006) PGC1 $\alpha$  expression is controlled in skeletal muscles by PPAR $\beta$ , whose ablation results in fiber-type switching, obesity, and type 2 diabetes. *Cell Metab* 4:407–414. <https://doi.org/10.1016/j.cmet.2006.10.003>
  116. Wang Y-X, Zhang C-L, Yu RT, Cho HK, Nelson MC, Bayuga-Ocampo CR, Ham J, Kang H, Evans RM (2004) Regulation of muscle fiber type and running endurance by PPARdelta. *PLoS Biol* 2:e294. <https://doi.org/10.1371/journal.pbio.0020294>
  117. Lunde IG, Ekmark M, Rana ZA, Buonanno A, Gundersen K (2007) PPARdelta expression is influenced by muscle activity and induces slow muscle properties in adult rat muscles after somatic gene transfer. *J Physiol* 582:1277–1287. <https://doi.org/10.1113/jphysiol.2007.133025>
  118. Krämer DK, Ahlsén M, Norrbom J, Jansson E, Hjeltnes N, Gustafsson T, Krook A (2006) Human skeletal muscle fibre type variations correlate with PPAR alpha, PPAR delta and PGC-1 alpha mRNA. *Acta Physiol* 188:207–216. <https://doi.org/10.1111/j.1748-1716.2006.01620.x>
  119. Luquet S, Lopez-Soriano J, Holst D, Fredenrich A, Melki J, Rassoulzadegan M, Grimaldi PA (2003) Peroxisome proliferator-activated receptor delta controls muscle development and oxidative capability. *FASEB J* 17:2299–2301. <https://doi.org/10.1096/fj.03-0269fje>
  120. Chandrashekar P, Manickam R, Ge X, Bonala S, McFarlane C, Sharma M, Wahli W, Kambadur R (2015) Inactivation of PPAR $\beta/\delta$  adversely affects satellite cells and reduces postnatal myogenesis. *Am J Physiol Endocrinol Metab* 309:E122–E131. <https://doi.org/10.1152/ajpendo.00586.2014>
  121. McPherron AC, Lawler AM, Lee SJ (1997) Regulation of skeletal muscle mass in mice by a new TGF-beta superfamily member. *Nature* 387:83–90. <https://doi.org/10.1038/387083a0>
  122. Wang Y-X, Lee C-H, Tiep S, Yu RT, Ham J, Kang H, Evans RM (2003) Peroxisome-proliferator-activated receptor  $\delta$  activates fat metabolism to prevent obesity. *Cell* 113:159–170. [https://doi.org/10.1016/S0092-8674\(03\)00269-1](https://doi.org/10.1016/S0092-8674(03)00269-1)
  123. Lin J, Wu H, Tarr PT, Zhang C-Y, Wu Z, Boss O, Michael LF, Puigserver P, Isotani E, Olson EN, Lowell BB, Bassel-Duby R, Spiegelman BM (2002) Transcriptional co-activator PGC-1 $\alpha$  drives the formation of slow-twitch muscle fibres. *Nature* 418:797–801. <https://doi.org/10.1038/nature00904>
  124. Pettersen IK, Tusubira D, Ashrafi H, Dyrstad SE, Hansen L, Liu X-Z, Nilsson LIH, Løvsletten NG, Berge K, Wergedahl H, Bjørndal B, Fluge Ø, Bruland O, Rustan AC, Halberg N, Røslund GV, Berge RK, Tronstad KJ (2019) Upregulated PDK4 expression is a sensitive marker of increased fatty acid oxidation. *Mitochondrion* 49:97–110. <https://doi.org/10.1016/j.mito.2019.07.009>
  125. Glinghammar B, Skogsberg J, Hamsten A, Ehrenborg E (2003) PPAR $\delta$  activation induces COX-2 gene expression and cell proliferation in human hepatocellular carcinoma cells. *Biochem Biophys Res Commun* 308:361–368. [https://doi.org/10.1016/S0006-291X\(03\)01384-6](https://doi.org/10.1016/S0006-291X(03)01384-6)
  126. Kostadinova R, Montagner A, Gouranton E, Fleury S, Guillou H, Dombrowicz D, Desreumaux P, Wahli W (2012) GW501516-activated PPAR $\beta/\delta$  promotes liver fibrosis via p38-JNK MAPK-induced hepatic stellate cell proliferation. *Cell Biosci* 2:34. <https://doi.org/10.1186/2045-3701-2-34>
  127. Da'adoosh B, Marcus D, Rayan A, King F, Che J, Goldblum A (2019) Discovering highly selective and diverse PPAR-delta agonists by ligand based machine learning and structural modeling. *Sci Rep* 9:1106. <https://doi.org/10.1038/s41598-019-38508-8>

128. Susini S, Roche E, Prentki M, Schlegel W (1998) Glucose and glucocorticoid peptides synergize to induce c-fos, c-jun, junB, zif-268, and nur-77 gene expression in pancreatic  $\beta$  (INS-1) cells. *FASEB J* 12:1173–1182
129. Roche E, Buteau J, Aniento I, Reig JA, Soria B, Prentki M (1999) Palmitate and oleate induce the immediate-early response genes c-fos and nur-77 in the pancreatic beta-cell line INS-1. *Diabetes* 48:2007–2014. <https://doi.org/10.2337/diabetes.48.10.2007>
130. Chen Y-T, Liao J-W, Tsai Y-C, Tsai F-J (2016) Inhibition of DNA methyltransferase 1 increases nuclear receptor subfamily 4 group A member 1 expression and decreases blood glucose in type 2 diabetes. *Oncotarget* 7:39162–39170. <https://doi.org/10.18632/oncotarget.10043>
131. Reynolds MS, Hancock CR, Ray JD, Kener KB, Draney C, Garland K, Hardman J, Bikman BT, Tessem JS (2016)  $\beta$ -Cell deletion of Nr4a1 and Nr4a3 nuclear receptors impedes mitochondrial respiration and insulin secretion. *Am J Physiol Endocrinol Metab* 311:E186–E201
132. Close A-F, Dadheech N, Lemieux H, Wang Q, Buteau J (2020) Disruption of beta-cell mitochondrial networks by the orphan nuclear receptor Nor1/Nr4a3. *Cell* 9:168. <https://doi.org/10.3390/cells9010168>
133. Tessem JS, Moss LG, Chao LC, Arlotto M, Lu D, Jensen MV, Stephens SB, Tontonoz P, Hohmeier HE, Newgard CB (2014) Nkx6.1 regulates islet  $\beta$ -cell proliferation via Nr4a1 and Nr4a3 nuclear receptors. *Proc Natl Acad Sci U S A* 111:5242–5247. <https://doi.org/10.1073/pnas.1320953111>
134. Ishihara H, Asano T, Tsukuda K, Katagiri H, Inukai K, Anai M, Kikuchi M, Yazaki Y, Miyazaki J-I, Oka Y (1993) Pancreatic beta cell line MIN6 exhibits characteristics of glucose metabolism and glucose-stimulated insulin secretion similar to those of normal islets. *Diabetologia* 36:1139–1145. <https://doi.org/10.1007/BF00401058>
135. Briand O, Helleboid-Chapman A, Ploton M, Hennuyer N, Carpentier R, Pattou F, Vandewalle B, Moerman E, Gmyr V, Kerr-Conte J, Eeckhoutte J, Staels B, Lefebvre P (2012) The nuclear orphan receptor Nur77 is a lipotoxicity sensor regulating glucose-induced insulin secretion in pancreatic  $\beta$ -cells. *Mol Endocrinol* 26:399–413. <https://doi.org/10.1210/me.2011-1317>
136. Yu C, Cui S, Zong C, Gao W, Xu T, Gao P, Chen J, Qin D, Guan Q, Liu Y (2015) The orphan nuclear receptor NR4A1 protects pancreatic  $\beta$ -cells from endoplasmic reticulum (ER) stress-mediated apoptosis. *J Biol Chem* 290:20687–20699
137. Pu Z, Liu D, Mouguegue L, Patherne HP, Jin C, Sadiq E, Qin D, Yu T, Zong C, Chen J, Zhao R (2020) NR4A1 counteracts JNK activation incurred by ER stress or ROS in pancreatic  $\beta$ -cells for protection. *J Cell Mol Med* 24:14171–14183
138. Pu Z, Yu T, Liu D, Jin C, Sadiq E, Qiao X, Li X, Chen Y, Zhang J, Tian M, Li S, Zhao R, Wang X (2021) NR4A1 enhances MKP7 expression to diminish JNK activation induced by ROS or ER-stress in pancreatic  $\beta$  cells for surviving. *Cell Death Dis* 7:133. <https://doi.org/10.1038/s41420-021-00521-0>
139. Zhan Y, Du X, Chen H, Liu J, Zhao B, Huang D, Li G, Xu Q, Zhang M, Weimer BC, Chen D, Cheng Z, Zhang L, Li Q, Li S, Zheng Z, Song S, Huang Y, Ye Z, Su W, Lin S-C, Shen Y, Wu Q (2008) Cytosporone B is an agonist for nuclear orphan receptor Nur77. *Nat Chem Biol* 4:548–556. <https://doi.org/10.1038/nchembio.106>
140. Weyrich P, Staiger H, Stancáková A, Schäfer SA, Kirchhoff K, Ullrich S, Ranta F, Gallwitz B, Stefan N, Machicao F, Kuusisto J, Laakso M, Fritsche A, Häring H-U (2009) Common polymorphisms within the NR4A3 locus, encoding the orphan nuclear receptor Nor-1, are associated with enhanced beta-cell function in non-diabetic subjects. *BMC Med Genet* 10:77. <https://doi.org/10.1186/1471-2350-10-77>
141. Close A-F, Dadheech N, Vilella BS, Rouillard C, Buteau J (2019) The orphan nuclear receptor Nor1/Nr4a3 is a negative regulator of  $\beta$ -cell mass. *J Biol Chem* 294:4889–4897. <https://doi.org/10.1074/jbc.RA118.005135>
142. Gao W, Fu Y, Yu C, Wang S, Zhang Y, Zong C, Xu T, Liu Y, Li X, Wang X (2014) Elevation of NR4A3 expression and its possible role in modulating insulin expression in the pancreatic beta cell. *PLoS One* 9:e91462. <https://doi.org/10.1371/journal.pone.0091462>
143. Wansa KDSA, Harris JM, Yan G, Ordentlich P, Muscat GEO (2003) The AF-1 domain of the orphan nuclear receptor NOR-1 mediates trans-activation, coactivator recruitment, and activation by the purine anti-metabolite 6-mercaptopurine. *J Biol Chem* 278:24776–24790. <https://doi.org/10.1074/jbc.M300088200>
144. Liu Q, Zhu X, Xu L, Fu Y, Garvey WT (2013) 6-Mercaptopurine augments glucose transport activity in skeletal muscle cells in part via a mechanism dependent upon orphan nuclear receptor NR4A3. *Am J Physiol Endocrinol Metab* 305:E1081–E1092. <https://doi.org/10.1152/ajpendo.00169.2013>
145. Pearen MA, Ryall JG, Maxwell MA, Ohkura N, Lynch GS, Muscat GEO (2006) The orphan nuclear receptor, NOR-1, is a target of beta-adrenergic signaling in skeletal muscle. *Endocrinology* 147:5217–5227. <https://doi.org/10.1210/en.2006-0447>
146. Takahashi H, Nomiyama T, Terawaki Y, Kawanami T, Hamaguchi Y, Tanaka T, Tanabe M, Bruemmer D, Yanase T (2019) GLP-1 receptor agonist exendin-4 attenuates NR4A orphan nuclear receptor NOR1 expression in vascular smooth muscle cells. *J Atheroscler Thromb* 26:183–197. <https://doi.org/10.5551/jat.43414>
147. Perissi V, Aggarwal A, Glass CK, Rose DW, Rosenfeld MG (2004) A corepressor/coactivator exchange complex required for transcriptional

- activation by nuclear receptors and other regulated transcription factors. *Cell* 116:511–526. [https://doi.org/10.1016/s0092-8674\(04\)00133-3](https://doi.org/10.1016/s0092-8674(04)00133-3)
148. Jeyakumar M, Liu X-F, Erdjument-Bromage H, Tempst P, Bagchi MK (2007) Phosphorylation of thyroid hormone receptor-associated nuclear receptor corepressor holocomplex by the DNA-dependent protein kinase enhances its histone deacetylase activity. *J Biol Chem* 282:9312–9322. <https://doi.org/10.1074/jbc.M609009200>
  149. Perissi V, Scafoglio C, Zhang J, Ohgi KA, Rose DW, Glass CK, Rosenfeld MG (2008) TBL1 and TBLR1 phosphorylation on regulated gene promoters overcomes dual CtBP and NCoR/SMRT transcriptional repression checkpoints. *Mol Cell* 29:755–766. <https://doi.org/10.1016/j.molcel.2008.01.020>
  150. Li J, Wang C-Y (2008) TBL1–TBLR1 and  $\beta$ -catenin recruit each other to Wnt target-gene promoter for transcription activation and oncogenesis. *Nat Cell Biol* 10:160–169. <https://doi.org/10.1038/ncb1684>
  151. Guenther MG, Lane WS, Fischle W, Verdin E, Lazar MA, Shiekhattar R (2000) A core SMRT corepressor complex containing HDAC3 and TBL1, a WD40-repeat protein linked to deafness. *Genes Dev* 14:1048–1057
  152. Yoon H-G, Chan DW, Huang Z-Q, Li J, Fondell JD, Qin J, Wong J (2003) Purification and functional characterization of the human N-CoR complex: the roles of HDAC3, TBL1 and TBLR1. *EMBO J* 22:1336–1346
  153. Ferrari A, Longo R, Fiorino E, Silva R, Mitro N, Cermenati G, Gilardi F, Desvergne B, Andolfo A, Magagnotti C, Caruso D, de Fabiani E, Hiebert SW, Crestani M (2017) HDAC3 is a molecular brake of the metabolic switch supporting white adipose tissue browning. *Nat Commun* 8:93. <https://doi.org/10.1038/s41467-017-00182-7>
  154. Sun Z, Feng D, Fang B, Mullican SE, You S-H, Lim H-W, Everett LJ, Nabel CS, Li Y, Selvakumaran V, Won K-J, Lazar MA (2013) Deacetylase-independent function of HDAC3 in transcription and metabolism requires nuclear receptor corepressor. *Mol Cell* 52:769–782. <https://doi.org/10.1016/j.molcel.2013.10.022>
  155. Chen W-B, Gao L, Wang J, Wang Y-G, Dong Z, Zhao J, Mi Q-S, Zhou L (2016) Conditional ablation of HDAC3 in islet beta cells results in glucose intolerance and enhanced susceptibility to STZ-induced diabetes. *Oncotarget* 7:57485–57497. <https://doi.org/10.18632/oncotarget.11295>
  156. Remsberg JR, Ediger BN, Ho WY, Damle M, Li Z, Teng C, Lanzillotta C, Stoffers DA, Lazar MA (2017) Deletion of histone deacetylase 3 in adult beta cells improves glucose tolerance via increased insulin secretion. *Mol Metab* 6:30–37. <https://doi.org/10.1016/j.molmet.2016.11.007>
  157. Liang N, Damdimopoulos A, Goñi S, Huang Z, Vedin L-L, Jakobsson T, Giudici M, Ahmed O, Pedrelli M, Barilla S, Alzaid F, Mendoza A, Schröder T, Kuiper R, Parini P, Hollenberg A, Lefebvre P, Francque S, van Gaal L, Staels B, Venticlef N, Treuter E, Fan R (2019) Hepatocyte-specific loss of GPS2 in mice reduces non-alcoholic steatohepatitis via activation of PPAR $\alpha$ . *Nat Commun* 10:1684. <https://doi.org/10.1038/s41467-019-09524-z>
  158. Drareni K, Ballaire R, Barilla S, Mathew MJ, Toubal A, Fan R, Liang N, Chollet C, Huang Z, Kondili M, Foulfelle F, Soprani A, Roussel R, Gautier J-F, Alzaid F, Treuter E, Venticlef N (2018) GPS2 deficiency triggers maladaptive white adipose tissue expansion in obesity via HIF1A activation. *Cell Rep* 24:2957–2971.e6. <https://doi.org/10.1016/j.celrep.2018.08.032>
  159. Drareni K, Ballaire R, Alzaid F, Goncalves A, Chollet C, Barilla S, Nguewa J-L, Dias K, Lemoine S, Riveline J-P, Roussel R, Dalmás E, Velho G, Treuter E, Gautier J-F, Venticlef N (2020) Adipocyte reprogramming by the transcriptional coregulator GPS2 impacts beta cell insulin secretion. *Cell Rep* 32:108141. <https://doi.org/10.1016/j.celrep.2020.108141>
  160. Kulozik P, Jones A, Mattijssen F, Rose AJ, Reimann A, Strzoda D, Kleinsorg S, Raupp C, Kleinschmidt J, Müller-Decker K, Wahli W, Sticht C, Gretz N, von Loeffelholz C, Stockmann M, Pfeiffer A, Stöhr S, Dallinga-Thie GM, Nawroth PP, Diaz MB, Herzig S (2011) Hepatic deficiency in transcriptional cofactor TBL1 promotes liver steatosis and hypertriglyceridemia. *Cell Metab* 13:389–400. <https://doi.org/10.1016/j.cmet.2011.02.011>
  161. Stoy C, Sundaram A, Rios Garcia M, Wang X, Seibert O, Zota A, Wendler S, Männle D, Hinz U, Sticht C, Muciek M, Gretz N, Rose AJ, Greiner V, Hofmann TG, Bauer A, Hoheisel J, Berriel Diaz M, Gaida MM, Werner J, Schafmeier T, Strobel O, Herzig S (2015) Transcriptional co-factor Transducin beta-like (TBL) 1 acts as a checkpoint in pancreatic cancer malignancy. *EMBO Mol Med* 7:1048–1062. <https://doi.org/10.15252/emmm.201404837>
  162. Gu J-F, Fu W, Qian H-X, Gu W-X, Zong Y, Chen Q, Lu L (2020) TBL1XR1 induces cell proliferation and inhibit cell apoptosis by the PI3K/AKT pathway in pancreatic ductal adenocarcinoma. *World J Gastroenterol* 26:3586–3602. <https://doi.org/10.3748/wjg.v26.i25.3586>
  163. Rohm M, Sommerfeld A, Strzoda D, Jones A, Sijmonsma TP, Rudofsky G, Wolfrum C, Sticht C, Gretz N, Zeyda M, Leitner L, Nawroth PP, Stulnig TM, Diaz MB, Vegiopoulos A, Herzig S (2013) Transcriptional cofactor TBLR1 controls lipid mobilization in white adipose tissue. *Cell Metab* 17:575–585. <https://doi.org/10.1016/j.cmet.2013.02.010>
  164. Schaefer U, Schmeier S, Bajic VB (2011) TcoF-DB: dragon database for human transcription co-factors and transcription factor interacting proteins. *Nucleic Acids Res* 39:D106–D110. <https://doi.org/10.1093/nar/gkq945>